



US 20180280394A1

(19) **United States**

(12) **Patent Application Publication**

Bates et al.

(10) **Pub. No.: US 2018/0280394 A1**

(43) **Pub. Date: Oct. 4, 2018**

(54) **METHODS OF TREATING LIVER DISEASE**

filed on Apr. 5, 2017, provisional application No. 62/477,697, filed on Mar. 28, 2017.

(71) Applicant: **Gilead Sciences, Inc.**, Foster City, CA (US)

Publication Classification

(72) Inventors: **Jamie Geier Bates**, Burlingame, CA (US); **David Gordon Clarkson Breckenridge**, San Mateo, CA (US); **John T. Liles**, San Jose, CA (US)

(51) **Int. Cl.**

A61K 31/519 (2006.01)

A61P 1/16 (2006.01)

A61K 31/4439 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 31/519** (2013.01); **A61K 31/4439** (2013.01); **A61P 1/16** (2018.01)

(21) Appl. No.: **15/935,759**

(22) Filed: **Mar. 26, 2018**

(57)

ABSTRACT

The present disclosure relates to a method of preventing and/or treating liver disease comprising administering an ACC inhibitor in combination with an FXR agonist to a patient in need thereof.

Related U.S. Application Data

Specification includes a Sequence Listing.

(60) Provisional application No. 62/586,354, filed on Nov. 15, 2017, provisional application No. 62/482,105,








FIG. 1

**FIG. 2**

**FIG. 3**

**FIG. 4**

**FIG. 5**






FIG. 6

**FIG. 7**

**FIG. 8**




FIG. 9

METHODS OF TREATING LIVER DISEASE

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of: U.S. Provisional Application Ser. No. 62/477,697, filed Mar. 28, 2017; U.S. Provisional Application Ser. No. 62/482,105, filed Apr. 5, 2017; and, U.S. Provisional Application Ser. No. 62/586,354, filed Nov. 15, 2017. The entireties of these applications are incorporated herein by reference.

SEQUENCE LISTING

[0002] The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is 1212P3C_2018-03-26_Seq_Listing_ST25.txt. The text file created on Mar. 26, 2018, is 2.32 KB in size and submitted electronically via EFS-Web.

FIELD

[0003] The present disclosure relates to methods of preventing and/or treating liver diseases.

BACKGROUND

[0004] Liver disease is generally classified as acute or chronic based upon the duration of the disease. Liver disease may be caused by infection, injury, exposure to drugs or toxic compounds, alcohol, impurities in foods, and the abnormal build-up of normal substances in the blood, an autoimmune process, a genetic defect (such as haemochromatosis), or unknown cause(s).

[0005] Liver disease is a leading cause of death world wide. In particular, it has been seen that a diet high in fat damages the liver in ways that are surprisingly similar to hepatitis. The American Liver Foundation estimates that more than 20 percent of the population has non-alcoholic fatty liver disease (NAFLD). It is suggested that obesity, unhealthy diets, and sedentary lifestyles may contribute to the high prevalence of NAFLD. When left untreated, NAFLD can progress to non-alcoholic steatohepatitis (NASH) causing serious adverse effects. Once NASH develops, it causes the liver to swell and scar (i.e. cirrhosis) over time.

[0006] Although preliminary reports suggest positive lifestyle changes could prevent or reverse liver damage, there are no effective medical treatments for NAFLD or NASH. Accordingly, there remains a need to provide new effective pharmaceutical agents to treat liver diseases.

SUMMARY

[0007] Disclosed herein are methods of treating and/or preventing liver disease in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an acetyl-CoA carboxylase (ACC) inhibitor in combination with a therapeutically effective amount of farnesoid X receptor (FXR) agonist. The liver disease can be any liver disease, including, but not limited to, chronic and/or metabolic liver diseases, nonalcoholic fatty liver disease (NAFLD), and nonalcoholic steatohepatitis (NASH).

[0008] In certain embodiments, provided herein is a method of treating and/or preventing nonalcoholic steatohepatitis (NASH) in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of a FXR agonist.

[0009] In the methods provided herein, the ACC inhibitor and the FXR agonist can be coadministered. In such embodiments, the ACC inhibitor and the FXR agonist can be administered together as a single pharmaceutical composition, or separately in more than one pharmaceutical composition. Accordingly, also provided herein is a pharmaceutical composition comprising a therapeutically effective amount of an ACC inhibitor and a therapeutically effective amount of a FXR agonist.

DESCRIPTION OF THE DRAWINGS

[0010] FIG. 1. Liver triglycerides in $\mu\text{mol/g}$ in the murine FFD model. (* $p<0.05$; ** $p<0.01$; *** $p<0.001$; **** $p<0.0001$ significantly different from vehicle by ANOVA). Graph shows mean $\pm$ SEM.

[0011] FIG. 2. ALT IU/L in the murine FFD model. (** $p<0.001$; significantly different from vehicle by ANOVA). Graph shows mean $\pm$ SEM.

[0012] FIG. 3. Hepatic expression of liver fibrosis gene Col1a1 measured by quantitative RT-PCR in the murine FFD model. (** $p<0.01$; **** $p<0.0001$ significantly different from vehicle by ANOVA; # significantly different from either single agent by t-test). Graph shows mean $\pm$ SEM.

[0013] FIG. 4. Hepatic expression of liver fibrosis gene Timp1 measured by quantitative RT-PCR in the murine FFD model. (* $p<0.05$; **** $p<0.0001$ significantly different from vehicle by ANOVA; # significantly different from either single agent by t-test). Graph shows mean $\pm$ SEM.

[0014] FIG. 5. Percent PSR positive area by quantitative image analysis in the rat CDHFD model. (** $p<0.01$, *** $p<0.001$, **** $p<0.0001$ significantly different from vehicle by t-test; & $p<0.001$ significantly different from start of treatment by t-test). Graph shows mean $\pm$ SEM.

[0015] FIG. 6. Percent α -SMA positive area by quantitative image analysis in the rat CDHFD model. (** $p<0.01$ significantly different from vehicle by t-test; & $p<0.001$ significantly different from start of treatment by t-test; # $p<0.05$ significantly different from either single agent by t-test). Graph shows mean $\pm$ SEM.

[0016] FIG. 7. Timp1 protein measured in plasma by ELISA in the rat CDHFD model. (* $p<0.05$ significantly different from vehicle by t-test; & $p<0.001$ significantly different from start of treatment by t-test). Graph shows mean $\pm$ SEM.

[0017] FIG. 8. Hyaluronic acid (HA) measured in plasma by ELISA in the rat CDHFD model. (** $p<0.01$, *** $p<0.001$, **** $p<0.0001$ significantly different from vehicle by t-test). Graph shows mean $\pm$ SEM.

[0018] FIG. 9. N-terminal propeptide of Type III Collagen (PIIINP) measured in plasma by ELISA in the rat CDHFD model. (* $p<0.05$, ** $p<0.01$, **** $p<0.0001$ significantly different from vehicle by t-test; & $p<0.001$ significantly different from start of treatment by t-test; # $p<0.05$ significantly different from either single agent by t-test). Graph shows mean $\pm$ SEM.

DETAILED DESCRIPTION

Definitions and General Parameters

[0019] As used in the present specification, the following terms and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise.

[0020] As used herein, the term “about” used in the context of quantitative measurements means the indicated amount $\pm 10\%$, or alternatively the indicated amount $\pm 5\%$ or $\pm 1\%$.

[0021] The term “pharmaceutically acceptable salt” refers to a salt of a compound disclosed herein that retains the biological effectiveness and properties of the underlying compound, and which is not biologically or otherwise undesirable. There are acid addition salts and base addition salts. Pharmaceutically acceptable acid addition salts may be prepared from inorganic and organic acids.

[0022] Acids and bases useful for reaction with an underlying compound to form pharmaceutically acceptable salts (acid addition or base addition salts respectively) are known to one of skill in the art. Similarly, methods of preparing pharmaceutically acceptable salts from an underlying compound (upon disclosure) are known to one of skill in the art and are disclosed in for example, Berge, at al. *Journal of Pharmaceutical Science*, January 1977 vol. 66, No. 1, and other sources.

[0023] As used herein, “pharmaceutically acceptable carrier” includes excipients or agents such as solvents, diluents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like that are not deleterious to the disclosed compound or use thereof. The use of such carriers and agents to prepare compositions of pharmaceutically active substances is well known in the art (see, e.g., Remington’s *Pharmaceutical Sciences*, Mace Publishing Co., Philadelphia, Pa. 17th Ed. (1985); and *Modern Pharmaceutics*, Marcel Dekker, Inc. 3rd Ed. (G. S. Banker & C. T. Rhodes, Eds.).

[0024] The terms “therapeutically effective amount” and “effective amount” are used interchangeably and refer to an amount of a compound that is sufficient to effect treatment as defined below, when administered to a patient (e.g., a human) in need of such treatment in one or more doses. The therapeutically effective amount will vary depending upon the patient, the disease being treated, the weight and/or age of the patient, the severity of the disease, or the manner of administration as determined by a qualified prescriber or care giver.

[0025] The term “treatment” or “treating” means administering a compound or pharmaceutically acceptable salt thereof for the purpose of: (i) delaying the onset of a disease, that is, causing the clinical symptoms of the disease not to develop or delaying the development thereof; (ii) inhibiting the disease, that is, arresting the development of clinical symptoms; and/or (iii) relieving the disease, that is, causing the regression of clinical symptoms or the severity thereof.

Liver Diseases

[0026] Liver diseases are acute or chronic damages to the liver based on the duration of the disease. The liver damage may be caused by infection, injury, exposure to drugs or toxic compounds such as alcohol or impurities in foods, an abnormal build-up of normal substances in the blood, an

autoimmune process, a genetic defect (such as haemochromatosis), or other unknown causes. Exemplary liver diseases include, but are not limited to, cirrhosis, liver fibrosis, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), alcoholic steatohepatitis (ASH), hepatic ischemia reperfusion injury, primary biliary cirrhosis (PBC), primary sclerosing cholangitis (PSC), and hepatitis, including both viral and alcoholic hepatitis.

[0027] Non-alcoholic fatty liver disease (NAFLD) is the build up of extra fat in liver cells that is not caused by alcohol. NAFLD may cause the liver to swell (i.e. steatohepatitis), which in turn may cause scarring (i.e. cirrhosis) over time and may lead to liver cancer or liver failure. NAFLD is characterized by the accumulation of fat in hepatocytes and is often associated with some aspects of metabolic syndrome (e.g. type 2 diabetes mellitus, insulin resistance, hyperlipidemia, hypertension). The frequency of this disease has become increasingly common due to consumption of carbohydrate-rich and high fat diets. A subset (~20%) of NAFLD patients develop nonalcoholic steatohepatitis (NASH).

[0028] NASH, a subtype of fatty liver disease, is the more severe form of NAFLD. It is characterized by macrovesicular steatosis, balloon degeneration of hepatocytes, and/or inflammation ultimately leading to hepatic scarring (i.e. fibrosis). Patients diagnosed with NASH progress to advanced stage liver fibrosis and eventually cirrhosis. The current treatment for cirrhotic NASH patients with end-stage disease is liver transplant.

[0029] Another common liver disease is primary sclerosing cholangitis (PSC). It is a chronic or long-term liver disease that slowly damages the bile ducts inside and outside the liver. In patients with PSC, bile accumulates in the liver due to blocked bile ducts, where it gradually damages liver cells and causes cirrhosis, or scarring of the liver. Currently, there is no effective treatment to cure PSC. Many patients having PSC ultimately need a liver transplant due to liver failure, typically about 10 years after being diagnosed with the disease. PSC may also lead to bile duct cancer.

[0030] Liver fibrosis is the excessive accumulation of extracellular matrix proteins, including collagen, that occurs in most types of chronic liver diseases. Advanced liver fibrosis results in cirrhosis, liver failure, and portal hypertension and often requires liver transplantation.

Methods

[0031] Disclosed herein is a method of treating and/or preventing liver disease in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of a FXR agonist. The presence of active liver disease can be detected by the existence of elevated enzyme levels in the blood. Specifically, blood levels of alanine aminotransferase (ALT) and aspartate aminotransferase (AST) above clinically accepted normal ranges are known to be indicative of on-going liver damage. Routine monitoring of liver disease patients for blood levels of ALT and AST is used clinically to measure progress of the liver disease while on medical treatment. Reduction of elevated ALT and AST to within the accepted normal range is taken as clinical evidence reflecting a reduction in the severity of the patient’s on-going liver damage.

[0032] In certain embodiments, the liver disease is a chronic liver disease. Chronic liver diseases involve the progressive destruction and regeneration of the liver parenchyma, leading to fibrosis and cirrhosis. In general, chronic liver diseases can be caused by viruses (such as hepatitis B, hepatitis C, cytomegalovirus (CMV), or Epstein Barr Virus (EBV)), toxic agents or drugs (such as alcohol, methotrexate, or nitrofurantoin), a metabolic disease (such as non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), haemochromatosis, or Wilson's Disease), an autoimmune disease (such as Autoimmune Chronic Hepatitis, Primary Biliary Cholangitis (formerly known as Primary Biliary Cirrhosis), or Primary Sclerosing Cholangitis, or other causes (such as right heart failure).

[0033] In one embodiment, provided herein is a method for reducing the level of cirrhosis. In one embodiment, cirrhosis is characterized pathologically by loss of the normal microscopic lobular architecture, with fibrosis and nodular regeneration. Methods for measuring the extent of cirrhosis are well known in the art. In one embodiment, the level of cirrhosis is reduced by about 5% to about 100%. In one embodiment, the level of cirrhosis is reduced by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% in the subject.

[0034] In certain embodiments, the liver disease is a metabolic liver disease. In one embodiment, the liver disease is non-alcoholic fatty liver disease (NAFLD). NAFLD is associated with insulin resistance and metabolic syndrome (obesity, combined hyperlipidemia, diabetes mellitus (type II) and high blood pressure). NAFLD is considered to cover a spectrum of disease activity, and begins as fatty accumulation in the liver (hepatic steatosis).

[0035] It has been shown that both obesity and insulin resistance probably play a strong role in the disease process of NAFLD. In addition to a poor diet, NAFLD has several other known causes. For example, NAFLD can be caused by certain medications, such as amiodarone, antiviral drugs (e.g., nucleoside analogues), aspirin (rarely as part of Reye's syndrome in children), corticosteroids, methotrexate, tamoxifen, or tetracycline. NAFLD has also been linked to the consumption of soft drinks through the presence of high fructose corn syrup which may cause increased deposition of fat in the abdomen, although the consumption of sucrose shows a similar effect (likely due to its breakdown into fructose). Genetics has also been known to play a role, as two genetic mutations for this susceptibility have been identified.

[0036] If left untreated, NAFLD can develop into non-alcoholic steatohepatitis (NASH), which is the most extreme form of NAFLD, a state in which steatosis is combined with inflammation and fibrosis. NASH is regarded as a major cause of cirrhosis of the liver. Accordingly, provided herein is a method of treating and/or preventing nonalcoholic steatohepatitis (NASH) in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of a FXR agonist.

[0037] Also provided herein is a method of treating and/or preventing liver fibrosis in a patient in need thereof, com-

prising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of a FXR agonist. Liver fibrosis is the excessive accumulation of extracellular matrix proteins including collagen that occurs in most types of chronic liver diseases. In certain embodiments, advanced liver fibrosis results in cirrhosis and liver failure. Methods for measuring liver histologies, such as changes in the extent of fibrosis, lobular hepatitis, and periportal bridging necrosis, are well known in the art.

[0038] In one embodiment, the level of liver fibrosis, which is the formation of fibrous tissue, fibroid or fibrous degeneration, is reduced by more than about 90%. In one embodiment, the level of fibrosis, which is the formation of fibrous tissue, fibroid or fibrous degeneration, is reduced by at least about 90%, at least about 80%, at least about 70%, at least about 60%, at least about 50%, at least about 40%, at least about 30%, at least about 20%, at least about 10%, at least about 5% or at least about 2%.

[0039] In one embodiment, the compounds provided herein reduce the level of fibrogenesis in the liver. Liver fibrogenesis is the process leading to the deposition of an excess of extracellular matrix components in the liver known as fibrosis. It is observed in a number of conditions such as chronic viral hepatitis B and C, alcoholic liver disease, drug-induced liver disease, hemochromatosis, autoimmune hepatitis, Wilson disease, Primary Biliary Cholangitis (formerly known as Primary Biliary Cirrhosis), sclerosing cholangitis, liver schistosomiasis and others. In one embodiment, the level of fibrogenesis is reduced by more than about 90%. In one embodiment, the level of fibrogenesis is reduced by at least about 90%, at least about 80%, at least about 70%, at least about 60%, at least about 50%, at least about 40%, at least about 30%, at least about 20%, at least about 10%, at least about 5% or at least 2%.

[0040] In still other embodiments, provided herein is a method of treating and/or preventing primary sclerosing cholangitis (PSC) in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of a FXR agonist.


[0041] It has been observed that patients having NASH are on average about 2.8 years older than healthy patients in epigenetic testing. Thus, in one embodiment, compounds useful for the treatment of NASH would be useful for slowing, improving or reversing epigenetic age or effects of aging due to NASH. In another embodiment, combination therapies for the treatment of NASH such as, for example, the combination of an ACC inhibitor with an FXR agonist as disclosed herein may be useful for improvement or reversal of aging effects due to NASH.

[0042] In one embodiment, the ACC inhibitor and the FXR agonist may be administered together in a combination formulation or in separate pharmaceutical compositions, where each inhibitor may be formulated in any suitable dosage form. In certain embodiments, the methods provided herein comprise administering separately a pharmaceutical composition comprising an ACC inhibitor and a pharmaceutically acceptable carrier or excipient and a pharmaceutical composition comprising a FXR agonist and a pharmaceutically acceptable carrier or excipient. Combination formulations according to the present disclosure comprise an ACC inhibitor and a FXR agonist together with one or more pharmaceutically acceptable carriers or excipients and

optionally other therapeutic agents. Combination formulations containing the active ingredient may be in any form suitable for the intended method of administration.


ACC Inhibitors

[0043] In certain embodiments of the methods and pharmaceutical compositions disclosed herein, the ACC inhibitor is a compound having the structure of Formula (I):



or a pharmaceutically acceptable salt thereof.

[0044] In certain embodiments of the methods and pharmaceutical compositions disclosed herein, the ACC inhibitor is a compound having the structure of Formula (II):




or a pharmaceutically acceptable salt thereof.

[0045] The compounds of Formula (I) and Formula (II) may be synthesized and characterized using methods known to those of skill in the art, such as those described in PCT International Application Publication No. WO 2013/071169. In one embodiment, the ACC inhibitor is the compound of Formula (I) or a pharmaceutically acceptable salt thereof. In one embodiment, the ACC inhibitor is the compound of Formula (II) or a pharmaceutically acceptable salt thereof.


FXR Agonist

[0046] In certain embodiments of the methods and pharmaceutical compositions disclosed herein, the FXR agonist is a compound having the structure of Formula (III):



or a pharmaceutically acceptable salt thereof.

[0047] In certain embodiments of the methods and pharmaceutical compositions disclosed herein, the FXR agonist is a compound having the structure of Formula (IV):



or a pharmaceutically acceptable salt thereof.

[0048] The compounds of Formula (III) and Formula (IV) may be synthesized and characterized using methods known to those of skill in the art, such as those described in U.S. Publication No. 2014/0221659.

Dosing and Administration

[0049] While it is possible for an active ingredient to be administered alone, it may be preferable to present them as pharmaceutical formulations or pharmaceutical compositions as described below. The formulations, both for veterinary and for human use, of the disclosure comprise at least one of the active ingredients, together with one or more acceptable carriers therefor and optionally other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and physiologically innocuous to the recipient thereof.

[0050] Each of the active ingredients can be formulated with conventional carriers and excipients, which will be selected in accord with ordinary practice. Tablets can contain excipients, glidants, fillers, binders and the like. Aqueous formulations are prepared in sterile form, and when intended for delivery by other than oral administration generally will be isotonic. All formulations will optionally contain excipients such as those set forth in the Handbook of Pharmaceutical Excipients (1986). Excipients include ascorbic acid and other antioxidants, chelating agents such as EDTA, carbohydrates such as dextrin, hydroxyalkylcellulose, hydroxyalkylmethylcellulose, stearic acid and the like. The pH of the formulations ranges from about 3 to about 11, but is ordinarily about 7 to 10.

[0051] The therapeutically effective amount of active ingredient can be readily determined by a skilled clinician using conventional dose escalation studies. Typically, each active ingredient will be administered in a dose from 0.01 milligrams to 1 gram. In one embodiment, the dosage will be from about 10 milligrams to 450 milligrams. In another embodiment, the dosage will be from about 25 to about 250 milligrams. In another embodiment, the dosage will be about 50 or 100 milligrams. In one embodiment, the dosage will be about 100 milligrams. In one embodiment, 20 mg of an ACC inhibitor is administered. In a specific embodiment, 20 mg of a compound of Formula (II) is administered. In one embodiment, 30 mg of an FXR agonist is administered. In a specific embodiment, 30 mg of a compound of Formula (III) is administered. It is contemplated that the active ingredients may be administered once, twice or three times a day. Also, the active ingredients may be administered once or twice a week, once every two weeks, once every three weeks, once every four weeks, once every five weeks, or once every six weeks.

[0052] The pharmaceutical composition for the active ingredient can include those suitable for the foregoing administration routes. The formulations can conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Techniques and formulations generally are found in Remington's Pharmaceutical Sciences (Mack Publishing Co., Easton, Pa.). Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

[0053] Formulations suitable for oral administration can be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be administered as a bolus, electuary or paste. In certain embodiments, the active ingredient may be administered as a subcutaneous injection.

[0054] A tablet can be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets can be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, or surface active agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered active ingredient moistened with an inert liquid diluent. The tablets may optionally be coated or scored and optionally are formulated so as to provide slow or controlled release of the active ingredient therefrom.

[0055] The active ingredient can be administered by any route appropriate to the condition. Suitable routes include oral, rectal, nasal, topical (including buccal and sublingual), vaginal and parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural), and the like. It will be appreciated that the preferred route may vary with for example the condition of the recipient. In

certain embodiments, the active ingredients are orally bio-available and can therefore be dosed orally. In one embodiment, the patient is human.

[0056] When used in combination in the methods disclosed herein, the ACC inhibitor and the FXR agonist can be administered together in a single pharmaceutical composition, e.g. a fixed dose combination, or separately (either concurrently or sequentially) in more than one pharmaceutical composition. In certain embodiments, the ACC inhibitor and the FXR agonist are administered together. In other embodiments, the ACC inhibitor and the FXR agonist are administered separately. In some aspects, the ACC inhibitor is administered prior to the FXR agonist. In some aspects, the FXR agonist is administered prior to the ACC inhibitor. When administered separately, the ACC inhibitor and the FXR agonist can be administered to the patient by the same or different routes of delivery.

Pharmaceutical Compositions

[0057] The pharmaceutical compositions of the disclosure comprise an effective amount of an ACC inhibitor selected from the group consisting of a compound of Formula (I) and a compound of Formula (II), or a pharmaceutically acceptable salt thereof, and an effective amount of a FXR agonist selected from the group consisting of a compound of Formula (III) and a compound of Formula (IV), or a pharmaceutically acceptable salt thereof.

[0058] When used for oral use for example, tablets, troches, lozenges, aqueous or oil suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents including sweetening agents, flavoring agents, coloring agents and preserving agents, in order to provide a palatable preparation. Tablets containing the active ingredient in admixture with non-toxic pharmaceutically acceptable excipient which are suitable for manufacture of tablets are acceptable. These excipients may be, for example, inert diluents, such as, for example, calcium or sodium carbonate, lactose, lactose monohydrate, croscarmellose sodium, povidone, calcium or sodium phosphate; granulating and disintegrating agents, such as, for example, maize starch, or alginic acid; binding agents, such as, for example, cellulose, microcrystalline cellulose, starch, gelatin or acacia; and lubricating agents, such as, for example, magnesium stearate, stearic acid or talc. Tablets may be uncoated or may be coated by known techniques including microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as, for example, glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

[0059] Formulations for oral use may be also presented as hard gelatin capsules where the active ingredient is mixed with an inert solid diluent, for example calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, such as, for example, peanut oil, liquid paraffin or olive oil.

[0060] Aqueous suspensions of the disclosure contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients include a suspending agent, such as, for example, sodium

carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as, for example, a naturally occurring phosphatide (e.g., lecithin), a condensation product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate), a condensation product of ethylene oxide with a long chain aliphatic alcohol (e.g., heptadecaethyleneoxycetanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan monooleate). The aqueous suspension may also contain one or more preservatives such as, for example, ethyl or n-propyl p-hydroxy-benzoate, one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as, for example, sucrose or saccharin.

[0061] Oil suspensions may be formulated by suspending the active ingredient in a vegetable oil, such as, for example, arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as, for example, liquid paraffin. The oral suspensions may contain a thickening agent, such as, for example, beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as, for example, those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as, for example, ascorbic acid.

[0062] Dispersible powders and granules of the disclosure suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent, and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those disclosed above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

[0063] The pharmaceutical compositions of the disclosure may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as, for example, olive oil or arachis oil, a mineral oil, such as, for example, liquid paraffin, or a mixture of these. Suitable emulsifying agents include naturally-occurring gums, such as, for example, gum acacia and gum tragacanth, naturally occurring phosphatides, such as, for example, soybean lecithin, esters or partial esters derived from fatty acids and hexitol anhydrides, such as, for example, sorbitan monooleate, and condensation products of these partial esters with ethylene oxide, such as, for example, polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, such as, for example, glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, a flavoring or a coloring agent.

[0064] The pharmaceutical compositions of the disclosure may be in the form of a sterile injectable preparation, such as, for example, a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a

sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as, for example, a solution in 1,3-butane-diol or prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may conventionally be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as, for example, oleic acid may likewise be used in the preparation of injectables.

[0065] The amount of active ingredient that may be combined with the carrier material to produce a single dosage form will vary depending upon the host treated and the particular mode of administration, such as oral administration or subcutaneous injection. For example, a time-release formulation intended for oral administration to humans may contain approximately 1 to 1000 mg of active material compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95% of the total compositions (weight:weight). The pharmaceutical composition can be prepared to provide easily measurable amounts for administration. For example, an aqueous solution intended for intravenous infusion may contain from about 3 to 500 μg of the active ingredient per milliliter of solution in order that infusion of a suitable volume at a rate of about 30 mL/hr can occur. When formulated for subcutaneous administration, the formulation is typically administered about twice a month over a period of from about two to about four months.

[0066] Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents.

[0067] The formulations can be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions are prepared from sterile powders, granules and tablets of the kind previously described. Preferred unit dosage formulations are those containing a daily dose or unit daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the active ingredient.

Examples

Example 1. Efficacy in a Mouse Model of NASH

[0068] The following study was conducted to evaluate the efficacy of the combination of an ACC inhibitor and an FXR agonist in a mouse model of non-alcoholic steatohepatitis (NASH), relative to the efficacy of the individual agents alone in the model. NASH was induced in male C57BL/6 mice by chronic administration of a "fast food" diet (FFD) high in saturated fats, cholesterol and sugars for a total of 6 months, whereas lean control animals were maintained on a

normal chow diet. A NASH phenotype was established in FFD mice compared to control mice after 6 months, and was characterized by macrovesicular steatosis, elevated ALT and AST, and increased levels of transcripts associated with hepatic stellate cell activation. See Charlton M, et al. Fast food diet mouse: novel small animal model of NASH with ballooning, progressive fibrosis, and high physiological fidelity to the human condition. *American Journal of Physiology. Gastrointestinal and Liver Physiology* 2011; 301 (5):G825-34.

[0069] After 5 months, FFD mice were subsequently treated with placebo (vehicle), an ACC inhibitor (Formula (I)), an FXR agonist (Formula (III)), or with the combination of Formula (I) and Formula (III) for 1 month. Control mice remained on a normal chow diet for the entire 6 month study period. Endpoint analyses included biochemical quantification of liver triglycerides, plasma ALT, and measurement of the pro-fibrotic transcripts Timp1 and Col1A1 in liver.

formulated in 0.5% sodium carboxymethylcellulose (medium viscosity), 1% w/w ethanol, 98.5% w/w 50 mM Tris Buffer, pH 8 in reverse osmosis water. The compound of Formula (I) was formulated at either 0.1 or 0.2 mg/mL and given in the dose provided in Table 1, and the compound of Formula (III) was formulated at 2 mg/mL and given in the dose provided in Table 1.

[0073] Starting seven days before PO dosing, animals in groups 1-6 were sham dosed with vehicle BID. The sham dosing was designed to acclimate animals to oral gavage dose administration. Starting at Day 1 of the study, animals in all dose groups were dosed three times daily; twice sequentially in the AM (7:00+/-1 hour), and once in the evening (19:00+/-1 hr), with the same volume of formulation containing no compound (group 1, vehicle) or the appropriate compounds as outlined below (Table 1) for 28 days (until dosing Day 29). Each group was split into two and half were sacrificed 2 hours post dose, and half were sacrificed 8 hours post dose on Day 29.

TABLE 1

Experimental Design and Dose Groups								
Group	Test Article	Dose (mg/kg)	Dose Vol (mL/kg)	Concentration (mg/mL)	Number of Animals	Dosing Frequency (x/day)	Dosing Duration (days)	Route
1	Vehicle	0	5	0	15	TID	29	PO
2	Vehicle	0	5	0	15	QD	29	PO
	Formula (I)	.5	5	0.1		BID	29	PO
3	Vehicle	0	5	0	15	BID	29	PO
	Formula (III)	10	5	2		QD		
4	Formula (I)	0.5	5	0.1	16	BID	29	PO
	Formula (III)	10	5	2		QD		PO
5	Vehicle (age-matched lean)	0	5	0	10	TID	29	PO

Methods

Animals

[0070] Male C57CL/6 mice (aged 12 weeks at study inception) were used in this study. All procedures used to study the animals were in the compliance with the U.S. Department of Agriculture's Animal Welfare Act (9 CFR Parts 1, 2, and 3); the Guide for the Care and Use of Laboratory Animals (Institute for Laboratory Animal Research, The National Academies Press, Washington, D.C.); and the National Institutes of Health, Office of Laboratory Animal Welfare.

In-Life Experimental Protocol for the FFD Mouse Model

[0071] The experimental design is shown in Table 1. Study animals were administered either a standard chow diet (Harlan Teklad Global Diets 2014, TD2014) or a commercially available high fat, high cholesterol diet (Research Diets Inc, DB12079B) (the FFD). Animals receiving the FFD were administered fructose/glucose in drinking water formulated as follows: 23.1 g fructose (Sigma, F2543) and 17.2 g of glucose (Sigma, 49158) was mixed into 1000 mL of drinking water.

[0072] The compound of Formula (I) or the compound of Formula (III) alone, or the combination of the compounds of Formula (I) and Formula (III), were administered for the final month of the study (month 5-month 6). The compound of Formula (I) and the compound of Formula (III) were

Quantification of Triglycerides from Murine Liver

[0074] Tissue Extraction:

[0075] Mouse liver tissue samples (25±10 mg, accurately weighed in frozen state) were homogenized and extracted with a water immiscible organic solvent mixture that extracts the triacylglyceride fraction as well as the free and esterified cholesterol fractions into the organic phase. After centrifugation, an aliquot of the organic upper layer, containing the triacylglycerides, cholesterol and cholesterol esters was diluted either 10-fold or 25-fold with ethanol. Two separate aliquots of this dilution were taken. One aliquot was analyzed for triacylglycerides, the second aliquot was used for the total cholesterol determination.

[0076] Triacylglyceride Determination:

[0077] For the triacylglyceride determination, one aliquot of the 25-fold dilution (or no dilution in the case of samples which have low triacylglyceride content) was evaporated under a stream of nitrogen. The dried extract was reconstituted stepwise with a 0.1% sodium dodecyl sulfate in PBS solution under ultrasonication followed by mixing with the Triacylglyceride Determination Reagent (Infinity™ Triglycerides Liquid Stable Reagent, Thermo Scientific, Product Data Sheet, Infinity™, Triglycerides Liquid Stable Reagent).

This reagent solution contained several enzymes, cofactors and the chromogenic reagent 4-aminoantipyrine. The determination of triacylglycerides (TAG) with this reagent was based on the method of Wako, Product Data Sheet, Triacyl-

glyceride-G Code No. 997-69801, Wako Pure Chemical Industries Ltd., Dallas, Tex., and the modifications by McGowan et al. (McGowan, M W, et al., Clin. Chem 1983;29:538) and Fossati et al (Fosseti, P. Prenciple L. Clin Chem. 1892;28:2077-80) as follows:

[0078] 1. Triglycerides are enzymatically hydrolyzed by lipase to free fatty acids and glycerol.

[0079] 2. The glycerol is phosphorylated by adenosine triphosphate (ATP) with glycerol kinase (GK) to produce glycerol-3-phosphate and adenosine diphosphate.

[0080] 3. Glycerol-3-phosphate is oxidized by dihydroxyacetone phosphate (DAP) by glycerol phosphate oxidase producing hydrogen peroxide (H₂O₂).

[0081] 4. In a Trinder⁵-type colour reaction catalyzed by peroxidase, the H₂O₂ reacts with 4-aminoantipyrine (4-AAP) and 3,5-dichloro-2-hydroxybenzene sulfonate (DHBS) to produce a red colored dye. The absorbance of this dye is proportional to the concentration of triglycerides present in the sample.

[0082] After incubation with the Triacylglyceride Determination Reagent for 30 min at 37° C., samples were transferred into a microtiter plate, and the absorbance is measured at 540 nm in a microplate reader (SpectraMax M2, Molecular Devices). Quantitation was performed using a

pyridoxal-5'-phosphate and analyzed on the Cobas Hitachi 6000 Chemistry System, Roche Diagnostics.

Gene Expression

[0084] An approximately 100 mg chunk of frozen left lateral lobe was sent to DC3 Therapeutics, LLC for lysing and RNA extraction. NanoString assays were carried out with all reagents and consumables contained in an nCounter master kit (NanoString) according to manufacturer instructions to measure RNA transcripts. Briefly, the color coded reporter probe targeting 110 liver fibrosis related genes and 6 control housekeeping genes (Table 2) were hybridized overnight in a pre-heated 65° C. thermocycler for 16 to 22 hours with 100 ng RNA samples in a reaction that includes a hybridization buffer and a capture probe. Following incubation, samples were placed on a prep station where excess probes were removed and the probe-transcript complexes were immobilized on a streptavidin coated cartridge. Finally, the cartridges were imaged in the nCounter Digital Analyzer (NanoString Technologies, Seattle, Wash.). All transcripts were normalized to the geometric mean of 6 housekeeping genes (B2m, Hprt, Pgk1, Rpl13a, Rpn1, and Sfrs4) with nSover 3.0 software.

TABLE 2

Nanostring Probes			
Gene Symbol	Accession Number	Target Sequence	
TIMP1	NM_011593.2	AAGCCTCTGTGGATATGCCACAAGTCCCAGAACCGCAGTGAAGAGTTTCTCATC ACGGGCCGCTAAGGAACGGAATTGTCACATCAGTGCCTGCAGC	
COL1A1	NM_007742.3	CAATGGTGAGACGTGGAAACCCGAGGTATGCTTGATCTGTATCTGCCACAATG GCACGGCTGTGTGCGATGACGTGCAATGCAATGAAGAAGTGGACTGT	
B2M	NM_009735.3	CATACGCCTGCAGAGTTAAGCATGCCAGTATGGCCGAGCCCAAGACCGTCTAC TGGGATCGAGACATGTGATCAAGCATCATGATGCTCTGAAGATTCTAT	
HPRT	NM_013556.2	TGCTGAGGCGGCGAGGGAGAGCGTTGGGCTTACCTCACTGCTTTCCGGAGCG GTAGACCTCTCCGCGGCTTCTCTCTAGACCGCTTTTGCCCGCA	
PGK1	NM_008828.2	CCGGCATTCTGCACGCTTCAAAAGCGCACGTCTGCCGCGCTGTTCTCTCTCTCTCT ATCTCCGGGCTTTTCGACCTCAGCGTGTGCAAAATGTCGCTT	
RPL13a	NM_009438.5	ATGGGATCCCTCCACCCTATGACAAGAAAAAGCGGATGGTGGTCCCTGCTGCT CTCAAGTTGTTTCGGCTGAAGCCTACCAGAAAGTTTGCTTACCTGGG	
RPN1	NM_133933.3	GGCAGCCTGACAGTGGGATCTCTCCATTCGTTCTTTAAGACCATCCTTCCTG CTGCCGCCAGGATGTCTATTACCGGATGAGATTGGTAATGTTTC	
SFRS4	NM_020587.2	GATGCTCACAAGGGACGCAAAACGAAGGAGTGATTGAATTTGTGCTTACTCT GATATGAAAAGAGCTTTGGAAAAGCTGGACGGAAGTGAAGTCAACG	

linear least squares regression analysis generated from fortified calibration standards using glyceryl trioleate (triolein) as triacylglyceride reference standard. Calibration standard samples were taken through the same extraction and incubation steps as the tissue samples. Weight corrections and concentration calculations were performed using Microsoft Excel 2013. Final tissue contents were given in μmol Triacylglyceride (TAG)/g Liver Tissue.

ALT

[0083] Serum was collected from all mice at terminal necropsy. Serum ALT was measured by Pyruvate with

Results

[0085] Example 1 demonstrates that a combined treatment with an ACC inhibitor and an FXR agonist results in greater efficacy than either agent administered alone in the mouse model of NASH. In particular, FIG. 1 shows a significant reduction in liver triglycerides with the combination of the compound of Formula (I) and the compound of Formula (III) relative to the individual agents, FIG. 2 shows a significant reduction in serum ALT with the combination of the compound of Formula (I) and the compound of Formula (III) relative to the individual agents, and FIG. 3 and FIG. 4 show a significant reduction in liver expression of Col1a1

and Timp1 with the combination of the compound of Formula (I) and the compound of Formula (III) relative to the individual agents, respectively.

Example 2. Efficacy in a Rat Model of NASH

[0086] The following study was conducted to evaluate the efficacy of the combination of an ACC inhibitor and an FXR agonist in a rodent model of non-alcoholic steatohepatitis (NASH) with fibrosis relative to the efficacy of the individual agents alone in the model. In this model, NASH with fibrosis was induced in male Wistar rats by administration of a choline-deficient high fat diet (CDHFD).

Animals

[0087] Male Wistar (CrI:Wi(Han)) rats (aged 8-9 weeks at arrival) were acquired from Charles River, Raleigh, N.C., and used in the current studies. This study complied with all applicable sections of the Final Rules of the Animal Welfare Act regulations (Code of Federal Regulations, Title 9), the Public Health Service Policy on Humane Care and Use of Laboratory Animals from the Office of Laboratory Animal Welfare, and the Guide for the Care and Use of Laboratory Animals from the National Research Council.

Vehicle Preparation

[0088] The vehicle, w/v 50 mM tris buffer, pH 8 in deionized water, was prepared prior to use and stored in a refrigerator set to maintain 2-8° C. To prepare 1 L, 800 mL of hot water (~80° C.) was added to an appropriate container and stirred vigorously until a steep vortex formed. 5.0 grams of sodium methylcellulose was slowly added to the sodium carboxymethylcellulose to the vortex. Stirring was continued until all carboxymethylcellulose was dissolved and the solution cooled down to ambient temperature. 5.12 g of Tris HCl was added to the container. 2.12 g of Tris base was added to the container. 10 g of ethanol was added to the container. The components were stirred for approximately 15 minutes, ensuring all solids have dissolved. QS water was added to 1 L with gentle mixing.

Study Design

[0089] Food was pro libitum and all animals on study were given a choline-deficient, high fat, high cholesterol diet (CDHFD; Research Diets, A16092003) on Day 1 of study except for group 1, the control chow group, which received standard diet (5CR4), as outlined in Table 3. On the day of sacrifice, liver was harvested and paraffin embedded, and plasma was collected and frozen. Animals were not dosed the day of sacrifice.

TABLE 3

Experimental Design and Dose Groups			
Group	Group name	n Diet (weeks)	Treatment (PO)
1	Control	10 Standard Diet (0-12)	N/A
2	Start of Treatment	10 CDHFD (0-6)	N/A
3	Vehicle	15 CDHFD (1-12)	N/A
4	Compound of Formula (I)	15 CDHFD (1-12)	10 mg/kg QD
5	Compound of Formula (III)	15 CDHFD (1-12)	30 mg/kg QD

TABLE 3-continued

Experimental Design and Dose Groups			
Group	Group name	n Diet (weeks)	Treatment (PO)
6	Compound of Formula (I) + Compound of Formula (III)	15 CDHFD (1-12)	10 mg/kg QD + 30 mg/kg QD

[0090] Tissues were collected by Charles River in Reno, Nev., processed and embedded in paraffin at Histo-tec in Hayward, Calif. and then shipped to Gilead Sciences in Foster City. Samples were sectioned at 5 µm and sections were mounted on glass slides for subsequent staining.

[0091] Picrosirius Red Staining:

[0092] Sections were pretreated in 0.2% Phosphomolybdic Acid (EMS, Cat#26357-01) and then subsequently incubated in 0.1% (W/V) Sirius Red 88-89-1 in saturated Picric acid solution (EMS, Cat#26357-02) for 1 hour at room temperature. This was followed by differentiation in 0.01N HCl (EMS, Cat#26357) and dehydration in graded alcohols.

[0093] Whole slide images of Picrosirius Red (PSR) stained slides were captured using a Leica AT2 scanner at 40× magnification. Digital slide images were checked for scanning quality, annotated and exported to appropriate network folders within Leica Digital Image Hub archive. Quantitative image analysis was performed on the whole slide images using Visiopharm image analysis software (Visiopharm, Hoersholm, Denmark) to determine the extent and intensity of PSR. The total PSR-stained area was measured and expressed as a percentage of total liver area stained. Results are shown in FIG. 7.

[0094] α-SMA:

[0095] Sections were deparaffinized in 3 changes of xylene for 5 minutes each, and subsequently rehydrated in 3 changes of 100% EtOH, 1 change of 95% EtOH, 1 change of 80% EtOH for 3 minutes each; followed by 2 successive rinses in distilled water. The sections were then incubated in Peroxidized 1 (Biocare Medical, Cat# PX968) endogenous peroxidase blocker for 5 minutes and rinsed in distilled water. Heat induced epitope retrieval was then performed using Reveal Decloaker (Biocare Medical, Cat# RV1000M) at 95° C. for 40 minutes with a Decloaking Chamber NxGen (Biocare Medical, Cat# DC2012), followed by gradual cooling with replacement of retrieval buffer with distilled water and placed in tris buffered saline (TBS). Immunohistochemistry was performed on prepared slides using an Intellipath autostainer (Biocare Medical, Cat# IPS0001) using the following steps:

[0096] 1. Apply 300 µL of Background Punisher (Biocare Medical, Cat# IP974G20) to slides and incubate for 10 minutes; followed by TBS wash.

[0097] 2. Apply 300 µL primary antibody of mouse monoclonal SMA, clone 1A4, (Biocare Medical, Cat# CM001) diluted 1:50 in Da Vinci Green diluent (Biocare Medical, Cat# PD900L). Incubate for 30 Minutes at room temperature; followed by TBS wash.

[0098] 3. Apply 300 µL of Mouse on Rat HRP Polymer (Biocare Medical, Cat# MRT621H) and incubate for 30 minutes; followed by TBS wash.

[0099] 4. Prepare DSB: 1 drop of DSB Chromogen/1 ml Substrate Buffer (Biocare Medical, Cat# BRI 4014C/

BRI 4013 respectfully). Apply 300 μ L Deep Space Black (DSB) Chromogen for 5 minutes; followed by distilled water wash.

[0100] 5. Counterstain with Nuclear Fast Red (Biocare Medical, Cat# STNFRLT) for 1 minute; followed by distilled water wash.

[0101] Slides were removed from the instrument and dehydrated through a series of graded histological grade alcohols to xylene and coverslipped.

[0102] Whole slide images of α -SMA stained slides were captured using a Leica AT2 scanner at 40 $\times$ magnification. Digital slide images were checked for scanning quality, annotated and exported to appropriate network folders within Leica Digital Image Hub archive. Quantitative image analysis was performed on the whole slide images using Visiopharm image analysis software (Visiopharm, Hoersholm, Denmark) to determine the extent and intensity of α -SMA. The total α -SMA-stained area was measured and expressed as a percentage of total liver area stained.

[0103] Plasma TIMP-1 ELISA:

[0104] Plasma TIMP-1 concentrations were determined in duplicate using a commercially available rat TIMP-1 specific ELISA kit (R&D Systems, Minneapolis, Minn., Cat # RTM100). TIMP-1 was assayed in plasma according to the manufacturer's specifications with minor modifications. Buffer RD1-21 (50 μ L) was added to ELISA plate wells pre-coated with mouse anti-TIMP-1. Prior to ELISA, a seven point standard curve of rat TIMP-1 (NSO-expressed recombinant TIMP-1: 2400-37.5 pg/mL) was generated and plasma samples were diluted 1:20 in buffer RD5-17. Samples and standards (50 μ L each) were added in duplicate to wells containing RD1-21 and incubated (room temperature) for 2 hours on an orbital plate shaker (300 rpm). Following antigen capture, plates were washed 5 times (350 μ L/well/wash) with Wash Buffer using an automated plate washer. Following washing, rat TIMP-1 conjugate (100 μ L) was added to each well and plates were incubated (room temperature) for 2 hours on an orbital plate shaker (300 rpm). Plates were then washed 5 times and Substrate Solution (100 μ L) was added to each well. Plates were incubated at room temperature for 30 minutes protected from light. Finally, Stop Solution (100 μ L) was added to each well. Optical Density (O.D.) absorbance was immediately determined at 450 nm on a SpectraMax 190 microplate reader (Molecular Devices, Sunnyvale Calif.). Relative O.D.s for each standard and sample were background corrected against blank samples, and standard curves for conversion of O.D.s to TIMP-1 concentration were generated using a 4 Parameter curve fit method. Unknown sample TIMP-1 concentrations were determined using SoftMax ProS software using a dilution factor of 20. Results are shown in FIG. 7.

[0105] Plasma PIIINP:

[0106] Plasma PIIINP concentrations were determined in duplicate using a commercially available rat Procollagen III N-Terminal Propeptide (PIIINP) ELISA Kit (Biomatik, Wilmington, Del., Cat# EKU06788). PIIINP was assayed in plasma diluted 50 fold in PBS according to the manufacturer's specifications with minor modifications. 7 standards (2,000 pg/mL, 1,000 pg/mL, 500 pg/mL, 250 pg/mL, 125 pg/mL, 62.5 pg/mL, 31.2 pg/mL) were prepared from standard stock which was reconstituted in Standard Diluent. 100 μ L each of standards, blank and samples were added into the appropriate wells. The plate was covered with the plate sealer and incubated for 1 hour at 37° C. After removing

liquid from each well, 100 μ L of Detection Reagent A working solution was added to each well and covered with the plate sealer then incubated for 1 hour at 37° C. The wells were washed with 350 μ L of 1 $\times$ Wash and sit for 1~2 minutes for 3 times. After the last wash, any remaining wash buffer was removed by decanting and blotting against absorbent paper. Then 100 μ L of Detection Reagent B working solution was added to each well, plate was covered with the plate sealer and incubated for 30 minutes at 37° C. The aspiration/wash process was repeated for total 5 times. 90 μ L of Substrate Solution was added to each well, plate was covered with a new plate sealer and incubated for 10-20 minutes at 37° C. protecting from light. The liquid turned blue by the addition of Substrate Solution Finally 50 μ L of Stop Solution was added to each well. The liquid then turned yellow. Mix the liquid by gently tapping the side of the plate. Optical Density (O.D.) absorbance was immediately determined at 450 nm on a SpectraMax 190 microplate reader (Molecular Devices, Sunnyvale Calif.). Relative O.D.s for each standard and sample were background corrected against blank samples, and standard curves for conversion of O.D.s to PIIINP concentration were generated using a 4 Parameter curve fit method. Unknown sample PIIINP concentrations were determined using SoftMax ProS software using a dilution factor of 50. Results are shown in FIG. 9.

[0107] Plasma Hyaluronic Acid (HA) Assay:

[0108] Plasma HA concentrations were determined in duplicate using a commercially available HA Test Kit (Corgenix, Inc., Broomfield, Colo., Cat#029-001). HA was assayed in plasma according to the manufacturer's specifications with minor modifications. Prior to assay, a seven point standard curve of HA reference solution (800-12.5 ng/mL) was generated and each reference sample and plasma sample was diluted 1 part to 10 parts Reaction Buffer (30 μ L reference/sample to 300 μ L Reaction Buffer). Samples and standards (100 μ L) were added in duplicate to microplate wells pre-coated with HA binding protein (HABP) and incubated (room temperature) for 60 minutes on an orbital plate shaker (300 rpm). Following antigen capture, plates were washed 4 times (350 μ L/well/wash) with PBS using an automated plate washer. Following washing, HRP-conjugated HABP (100 μ L) was added to each well and plates were incubated (room temperature) for 30 minutes on an orbital plate shaker (300 rpm). Plates were then washed 4 times and the one-component Substrate Solution (100 μ L) was added to each well. Plates were incubated at room temperature for 30 minutes protected from light. Finally, Stop Solution (100 μ L) was added to each well. Optical Density (O.D.) absorbance was immediately determined at 450 nm on a SpectraMax 190 microplate reader (Molecular Devices, Sunnyvale Calif.). Relative O.D.s for each standard and sample was background corrected against blank samples, and standard curves for conversion of O.D.s to HA concentration was generated using a 4 Parameter curve fit method. Undiluted unknown sample HA concentrations were determined using SoftMax ProS software. Results are shown in FIG. 8.

Results

[0109] Example 2 demonstrates that a combined treatment with an ACC inhibitor and an FXR agonist results in greater efficacy than either agent administered alone in the rat model of NASH. In particular, FIG. 5-9 shows a significant reduction markers of fibrosis including percent picrosirius posi-

tive area, percent α -SMA positive area, and three plasma markers associated with fibrosis, TIMP1, HA, and PIIINP with the combination of the compound of Formula (I) and the compound of Formula (III) relative to the vehicle. FIG.

6 and FIG. 9 show a significant reduction α -SMA and PIIINP with the combination of the compound of Formula (I) and the compound of Formula (III) relative to the individual agents, respectively.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 8

<210> SEQ ID NO 1
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 1

aagcctctgt ggatatgcc acaagtccca gaaccgcagt gaagagtctc tcatcacggg 60
ccgcctaagg aacggaaatt tgcacatcag tgccctgcagc 100

<210> SEQ ID NO 2
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 2

caatggtgag acgtggaaac ccgaggtatg cttgatctgt atctgccaca atggcacggc 60
tgtgtgcgat gacgtgcaat gcaatgaaga actggactgt 100

<210> SEQ ID NO 3
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 3

catacgcctg cagagttaag catgccagta tggccgagcc caagaccgtc tactgggac 60
gagacatgtg atcaagcatc atgatgctct gaagattcat 100

<210> SEQ ID NO 4
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 4

tgctgaggcg gcgagggaga gcgttgggct tacctcactg ctttcgggag cggtagcacc 60
tcctccgccc gcttctctct cagaccgctt ttgcccga 100

<210> SEQ ID NO 5
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 5

ccggcattct gcacgcttca aaagcgcacg tctgccgcgc tggtctctct ttcctcatct 60

-continued

 ccgggccttt cgacctcag gtgttgccaa aatgtcgctt 100

<210> SEQ ID NO 6
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 6

atgggatccc tccacctat gacaagaaaa agcggatggt ggtccctgct gctctcaagg 60

ttgttcggct gaagcctacc agaaagtgtg cttacctggg 100

<210> SEQ ID NO 7
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 7

ggcagcctga cagtgggatc tcctccattc gttcttttaa gaccatcctt cctgctgccg 60

cccaggatgt ctattaccgg gatgagattg gtaatgtttc 100


<210> SEQ ID NO 8
 <211> LENGTH: 101
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 8

gatgtcaca agggacgcaa aaacgaagga gtgattgaat ttgtgtctta ctctgatatg 60


aaaagagctt tggaaaagct ggacggaact gaagtcaacg g 101

1. A method of treating and/or preventing a liver disease in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of an FXR agonist, wherein the ACC inhibitor is a compound of Formula (I):




or a pharmaceutically acceptable salt thereof;

and the FXR agonist is a compound of Formula (III):




or a pharmaceutically acceptable salt thereof.

2. A method of treating and/or preventing a liver disease in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of an FXR agonist, wherein the ACC inhibitor is a compound of Formula (I):




or a pharmaceutically acceptable salt thereof;

and the FXR agonist is a compound of Formula (IV):




or a pharmaceutically acceptable salt thereof.

3. A method of treating and/or preventing a liver disease in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of an FXR agonist, wherein the ACC inhibitor is a compound of Formula (II):




or a pharmaceutically acceptable salt thereof;

and the FXR agonist is a compound of Formula (III):




or a pharmaceutically acceptable salt thereof.

4. A method of treating and/or preventing a liver disease in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of an ACC inhibitor in combination with a therapeutically effective amount of an FXR agonist, wherein the ACC inhibitor is a compound of Formula (II):



or a pharmaceutically acceptable salt thereof;

and the FXR agonist is a compound of Formula (IV):



or a pharmaceutically acceptable salt thereof.


5. The method of claim 1, wherein the liver disease is non-alcoholic steatohepatitis (NASH).

6. The method of claim 2, wherein the liver disease is non-alcoholic steatohepatitis (NASH).

7. The method of claim 3, wherein the liver disease is non-alcoholic steatohepatitis (NASH).


8. A pharmaceutical composition comprising a therapeutically effective amount of an ACC inhibitor, a therapeutically effective amount of an FXR agonist, and a pharmaceutically acceptable carrier.

cally effective amount of an FXR agonist, and a pharmaceutically acceptable carrier; wherein the ACC inhibitor is a compound of Formula (I):




or a pharmaceutically acceptable salt thereof;

and the FXR agonist is a compound of Formula (III):




or a pharmaceutically acceptable salt thereof.

9. A pharmaceutical composition comprising a therapeutically effective amount of an ACC inhibitor, a therapeutically effective amount of an FXR agonist, and a pharmaceutically acceptable carrier; wherein the ACC inhibitor is a compound of Formula (I):




or a pharmaceutically acceptable salt thereof;

and the FXR agonist is a compound of Formula (IV):




or a pharmaceutically acceptable salt thereof.

10. A pharmaceutical composition comprising a therapeutically effective amount of an ACC inhibitor, a therapeutically effective amount of an FXR agonist, and a pharmaceutically acceptable carrier; wherein the ACC inhibitor is a compound of Formula (II):




or a pharmaceutically acceptable salt thereof;

and the FXR agonist is a compound of Formula (III):




or a pharmaceutically acceptable salt thereof.

11. A pharmaceutical composition comprising a therapeutically effective amount of an ACC inhibitor, a therapeutically effective amount of an FXR agonist, and a pharmaceutically acceptable carrier; wherein the ACC inhibitor is a compound of Formula (II):



or a pharmaceutically acceptable salt thereof;

and the FXR agonist is a compound of Formula (IV):



or a pharmaceutically acceptable salt thereof.

12. The pharmaceutical composition of claim **8**, further comprising a pharmaceutically acceptable carrier.

13. The pharmaceutical composition of claim **9**, further comprising a pharmaceutically acceptable carrier.

14. The pharmaceutical composition of claim **10**, further comprising a pharmaceutically acceptable carrier.

15. The pharmaceutical composition of claim **11**, further comprising a pharmaceutically acceptable carrier.

16. The method of claim **4**, wherein the liver disease is non-alcoholic steatohepatitis (NASH).

* * * * *