

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 April 2009 (23.04.2009)

PCT

(10) International Publication Number
WO 2009/052231 A1

(51) International Patent Classification:
C07D 231/56 (2006.01) **A61P 35/00** (2006.01)
A61K 31/423 (2006.01)

(21) International Application Number:
PCT/US2008/080068

(22) International Filing Date: 16 October 2008 (16.10.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/981,265 19 October 2007 (19.10.2007) US

(71) Applicant (for all designated States except US): **ABBOTT LABORATORIES** [US/US]; 100 Abbott Park Road, Abbott Park, Illinois 60064 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BORCHARDT, Thomas, B.** [US/US]; 2433 55th Avenue, Kenosha, Wisconsin 53144 (US).


(74) Agents: **COLEMAN JAMES, Patricia** et al.; Abbott Laboratories, Dept. 0377, Bldg. AP6A-1A, 100 Abbott Park Road, Abbott Park, Illinois 60064-6008 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

(54) Title: CRYSTALLINE CHEMOTHERAPEUTIC



(57) Abstract: N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2, ways to make it, formulations comprising it and made with it and methods of treating patients having disease using it are disclosed.

WO 2009/052231 A1

CRYSTALLINE CHEMOTHERAPEUTIC

FIELD OF THE INVENTION

This invention pertains to N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2, ways to make it, formulations comprising it and made with it and methods of treating patients having disease using it.

BACKGROUND OF THE INVENTION

N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea (ABT-869) belongs to a family of protein tyrosine kinases (PTKs) which catalyze the phosphorylation of specific tyrosine residues in cellular proteins. Aberrant or excessive PTK activity has been observed in many disease states including benign and malignant proliferative disorders and diseases resulting from inappropriate activation of the immune system.

Crystallinity of ABT-869 may effect, among other physical and mechanical properties, their stability, solubility, dissolution rate, hardness, compressibility and melting point. Because ease of manufacture and formulation of ABT-869 is dependent on some, if not all, of these properties, there is an existing need in the chemical and therapeutic arts for identification of crystalline forms of ABT-869 and ways to reproducibly make them.

SUMMARY OF THE INVENTION

One embodiment of this invention, therefore, pertains to N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2 which, when measured at about 25°C with radiation at 1.54178 Å, is characterized by a powder diffraction pattern having respective 2θ values of about 6.7°, 10.1°, 11.8°, 13.4°, 14.4°, 15.4°, 16.6°, 16.8°, 17.8°, 20.5° and 23.4°.

Another embodiment pertains to formulations comprising an excipient and N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2 which, when measured at about 25°C with radiation at 1.54178 Å, is characterized by a powder diffraction pattern having respective 2θ values of about 6.7°, 10.1°, 11.8°, 13.4°, 14.4°, 15.4°, 16.6°, 16.8°, 17.8°, 20.5° and 23.4°.

Still another embodiment pertains to methods of treating cancer in a mammal comprising administering thereto, with or without one or more than one additional anticancer drugs, a therapeutically effective amount of N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2 which, when measured at about 25°C with radiation at 1.54178 Å, is characterized by a powder diffraction pattern having respective 2θ values of about 6.7°, 10.1°, 11.8°, 13.4°, 14.4°, 15.4°, 16.6°, 16.8°, 17.8°, 20.5° and 23.4°.

Still another embodiment pertains to a process for making N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2 comprising:

providing N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Hydrate Crystalline Form 1; and

removing water from the N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Hydrate Crystalline Form 1 to form N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2.

Still another embodiment comprises N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2 prepared by the process of the preceding embodiment.

Still another embodiment comprises ABT-869 Tolueneate for use in preparing N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2.

Still another embodiment comprises a salt of ABT-869 for use in preparing N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2.

Still another embodiment comprises the hydrochloride salt of ABT-869 for use in preparing N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2.

BRIEF DESCRIPTION of THE DRAWINGS

Figure 1 is a powder x-ray diffraction pattern for N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2.

DETAILED DESCRIPTION OF THE INVENTION

This invention pertains to discovery of N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2, ways to make it, ways to characterize it, formulations containing it and made with it, and methods of treating cancer using it. The terms “N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea” and “ABT-869” are meant to be used interchangeably.

The terms “ABT-869” and “an ABT-869” without any indicia of crystallinity or non-crystallinity associated with it, as used herein, mean amorphous ABT-869, a crystalline ABT-869, microcrystalline ABT-869, ABT-869 in solution, a semisolid, wax or oil form of ABT-869, mixtures thereof and the like.

The terms “crystalline” and “microcrystalline,” as used herein, mean having a regularly repeating arrangement of molecules which is maintained over a long range or external face planes.

Unless stated otherwise, percentages herein are weight/weight (w/w) percentages.

The term “hydrochloride salt,” as used herein, means having associated therewith one or more than one hydrochloride equivalent.

The term “solvent,” as used herein, means a liquid in which a compound is soluble or partially soluble enough at a given concentration to dissolve or partially dissolve the compound.

The term “anti-solvent,” as used herein, means a liquid in which a compound is insoluble enough at a given concentration to be effective for precipitating that compound from a solution.

Solvents and anti-solvents may be mixed with or without separation of phases.

It is meant to be understood that, because many solvents and anti-solvents contain impurities, the level of impurities in solvents and anti-solvents for the practice of this

invention, if present, are at a low enough concentration that they do not interfere with the intended use of the solvent in which they are present.

The term "acid," as used herein, means a compound having at least one acidic proton.

5 Examples of acids for the practice of this invention include, but are not limited to, hydrochloric acid, hydrobromic acid, trifluoroacetic acid, trichloroacetic acid, sulfuric acid, phosphoric acid and the like.

The term "base," as used herein, means a compound capable of accepting a proton.

10 Examples of bases for the practice of this invention include, but are not limited to, sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate, dibasic sodium phosphate (i.e. Na_2HPO_4 , K_2HPO_4 and the like), triethylamine, diisopropylethylamine and the like.

15 The term "isolating" as used herein, means separating ABT-869 Crystalline Form 2 from solvent, anti-solvent, or a mixture of solvent anti-solvent. This is typically accomplished by means such as centrifugation, filtration with or without vacuum, filtration with positive pressure, distillation, evaporation or a combination thereof.

20 Therapeutically acceptable amounts of ABT-869 Crystalline Form 2 depend on recipient of treatment, disorder being treated and severity thereof, composition containing it, time of administration, route of administration, duration of treatment, its potency, its rate of clearance and whether or not another drug is co-administered. The amount of ABT-869 Crystalline Form 2 used to make a formulation to be administered daily to a patient in a
25 single dose or in divided doses is from about 0.03 to about 200 mg/kg body weight. Single dose formulations contain these amounts or a combination of submultiples thereof.

ABT-869 Crystalline Form 2 may be administered with or without an excipient, typically with an excipient. Excipients include but are not limited to, for example,
30 encapsulating materials and additives such as absorption accelerators, antioxidants, binders, buffers, carriers, coating agents, coloring agents, diluents, disintegrating agents, emulsifiers, extenders, fillers, flavoring agents, glidants, humectants, lubricants, perfumes, preservatives, propellants, releasing agents, sterilizing agents, sweeteners, solubilizers, wetting agents, mixtures thereof and the like.

Excipients for preparation of formulations comprising or made with ABT-869 Crystalline Form 2 to be administered orally in solid dosage form include, for example, agar, alginic acid, aluminum hydroxide, benzyl alcohol, benzyl benzoate, 1,3-butylene glycol, carbomers, castor oil, cellulose, cellulose acetate, cocoa butter, copovidone, corn starch, corn oil, cottonseed oil, cross-povidone, diglycerides, ethanol, ethyl cellulose, ethyl laurate, ethyl oleate, fatty acid esters, gelatin, germ oil, glucose, glycerol, groundnut oil, hydroxypropylmethyl cellulose, isopropanol, isotonic saline, lactose, magnesium hydroxide, magnesium stearate, malt, mannitol, monoglycerides, olive oil, povidone, peanut oil, potassium phosphate salts, potato starch, povidone, propylene glycol, Ringer's solution, safflower oil, sesame oil, silicon dioxide, sodium carboxymethyl cellulose, sodium phosphate salts, sodium lauryl sulfate, sodium sorbitol, sodium stearyl fumarate, soybean oil, stearic acids, stearyl fumarate, sucrose, surfactants, talc, tragacanth, tetrahydrofurfuryl alcohol, triglycerides, vitamin E and derivatives thereof, water, mixtures thereof and the like.

Excipients for preparation of formulations comprising or made with ABT-869 Crystalline Form 2 to be administered ophthalmically or orally in liquid dosage forms include, for example, 1,3-butylene glycol, castor oil, corn oil, cottonseed oil, ethanol, fatty acid esters of sorbitan, germ oil, groundnut oil, glycerol, isopropanol, olive oil, polyethylene glycols, propylene glycol, sesame oil, water, mixtures thereof and the like.

Excipients for preparation of formulations comprising or made with ABT-869 Crystalline Form 2 to be administered osmotically include, for example, chlorofluorohydrocarbons, ethanol, water, mixtures thereof and the like.

Excipients for preparation of formulations comprising or made with ABT-869 Crystalline Form 2 to be administered parenterally include, for example, 1,3-butanediol, castor oil, corn oil, cottonseed oil, dextrose, germ oil, groundnut oil, liposomes, oleic acid, olive oil, peanut oil, Ringer's solution, safflower oil, sesame oil, soybean oil, U.S.P. or isotonic sodium chloride solution, water, mixtures thereof and the like.

Excipients for preparation of formulations comprising or made with ABT-869 Crystalline Form 2 to be administered rectally or vaginally include, but are not limited to, cocoa butter, polyethylene glycol, wax, mixtures thereof and the like.

ABT-869 Crystalline Form 2 is also useful when administered with anticancer drugs such as alkylating agents, angiogenesis inhibitors, antibodies, antimetabolites, antimitotics, antiproliferatives, aurora kinase inhibitors, Bcr-Abl kinase inhibitors, biologic response
5 modifiers, cyclin-dependent kinase inhibitors, cell cycle inhibitors, cyclooxygenase-2 inhibitors, leukemia viral oncogene homolog (ErbB2) receptor inhibitors, growth factor inhibitors, heat shock protein (HSP)-90 inhibitors, histone deacetylase (HDAC) inhibitors, hormonal therapies, immunologicals, intercalating antibiotics, other kinase inhibitors, including other PTKs, mammalian target of rapamycin inhibitors, mitogen-activated
10 extracellular signal-regulated kinase inhibitors, non-steroidal anti-inflammatory drugs (NSAIDs), platinum chemotherapeutics, polo-like kinase inhibitors, proteasome inhibitors, purine analogs, pyrimidine analogs, receptor tyrosine kinase inhibitors, retinoids/deltoids plant alkaloids, topoisomerase inhibitors and the like.

15 Alkylating agents include altretamine, AMD-473, AP-5280, apaziquone, bendamustine, brostallicin, busulfan, carboquone, carmustine (BCNU), chlorambucil, Cloretazine™ (VNP 40101M), cyclophosphamide, decarbazine, estramustine, fotemustine, glufosfamide, ifosfamide, KW-2170, lomustine (CCNU), mafosfamide, melphalan, mitobronitol, mitolactol, nimustine, nitrogen mustard N-oxide, ranimustine, temozolomide,
20 thiotepa, treosulfan, trofosfamide and the like.

Angiogenesis inhibitors include endothelial-specific receptor tyrosine kinase (Tie-2) inhibitors, epidermal growth factor receptor (EGFR) inhibitors, insulin growth factor-2 receptor (IGFR-2) inhibitors, matrix metalloproteinase-2 (MMP-2) inhibitors, matrix
25 metalloproteinase-9 (MMP-9) inhibitors, platelet-derived growth factor receptor (PDGFR) inhibitors, thrombospondin analogs vascular endothelial growth factor receptor tyrosine kinase (VEGFR) inhibitors and the like.

Aurora kinase inhibitors include AZD-1152, MLN-8054, VX-680 and the like.

30 Bcr-Abl kinase inhibitors include DASATINIB® (BMS-354825), GLEEVEC® (imatinib) and the like.

CDK inhibitors include AZD-5438, BMI-1040, BMS-032, BMS-387, CVT-2584,

flavopyridol, GPC-286199, MCS-5A, PD0332991, PHA-690509, seliciclib (CYC-202, R-roscovitine), ZK-304709 and the like.

COX-2 inhibitors include ABT-963, ARCOXIA[®] (etoricoxib), BEXTRA[®] (valdecoxib), BMS347070, CELEBREX[™] (celecoxib), COX-189 (lumiracoxib), CT-3, DERAMAXX[®] (deracoxib), JTE-522, 4-methyl-2-(3,4-dimethylphenyl)-1-(4-sulfamoylphenyl)-1H-pyrrole), MK-663 (etoricoxib), NS-398, parecoxib, RS-57067, SC-58125, SD-8381, SVT-2016, S-2474, T-614, VIOXX[®] (rofecoxib) and the like.

EGFR inhibitors include ABX-EGF, anti-EGFR immunoliposomes, EGF-vaccine, EMD-7200, ERBITUX[®] (cetuximab), HR3, IgA antibodies, IRESSA[®] (gefitinib), TARCEVA[®] (erlotinib or OSI-774), TP-38, EGFR fusion protein, TYKERB[®] (lapatinib) and the like.

ErbB2 receptor inhibitors include CP-724-714, CI-1033 (canertinib), HERCEPTIN[®] (trastuzumab), TYKERB[®] (lapatinib), OMNITARG[®] (2C4, petuzumab), TAK-165, GW-572016 (ionafarnib), GW-282974, EKB-569, PI-166, dHER2 (HER2 vaccine), APC-8024 (HER-2 vaccine), anti-HER/2neu bispecific antibody, B7.her2IgG3, AS HER2 trifunctional bispecific antibodies, mAB AR-209, mAB 2B-1 and the like.

Histone deacetylase inhibitors include depsipeptide, LAQ-824, MS-275, trapoxin, suberoylanilide hydroxamic acid (SAHA), TSA, valproic acid and the like.

HSP-90 inhibitors include 17-AAG-nab, 17-AAG, CNF-101, CNF-1010, CNF-2024, 17-DMAG, geldanamycin, IPI-504, KOS-953, MYCOGRAB[®], NCS-683664, PU24FCI, PU-3, radicicol, SNX-2112, STA-9090 VER49009 and the like.

MEK inhibitors include ARRY-142886, ARRY-438162 PD-325901, PD-98059 and the like.

mTOR inhibitors include AP-23573, CCI-779, everolimus, RAD-001, rapamycin, temsirolimus and the like.

Non-steroidal anti-inflammatory drugs include AMIGESIC[®] (salsalate), DOLOBID[®] (diflunisal), MOTRIN[®] (ibuprofen), ORUDIS[®] (ketoprofen), RELAFEN[®] (nabumetone), FELDENE[®] (piroxicam) ibuprofen cream, ALEVE[®] and NAPROSYN[®] (naproxen), VOLTAREN[®] (diclofenac), INDOCIN[®] (indomethacin), CLINORIL[®] (sulindac),
5 TOLECTIN[®] (tolmetin), LODINE[®] (etodolac), TORADOL[®] (ketorolac), DAYPRO[®] (oxaprozin) and the like.

PDGFR inhibitors include C-451, CP-673, CP-868596 and the like.

10 Platinum chemotherapeutics include cisplatin, ELOXATIN[®] (oxaliplatin), eptaplatin, lobaplatin, nedaplatin, PARAPLATIN[®] (carboplatin), satraplatin and the like.

Polo-like kinase inhibitors include BI-2536 and the like.

15 Thrombospondin analogs include ABT-510, ABT-567, ABT-898, TSP-1 and the like.

VEGFR inhibitors include AVASTIN[®] (bevacizumab), ABT-869, AEE-788, ANGIOZYME[™], axitinib (AG-13736), AZD-2171, CP-547,632, IM-862, Macugen (pegaptamib), NEXAVAR[®] (sorafenib, BAY43-9006), pazopanib (GW-786034), (PTK-787,
20 ZK-222584), SUTENT[®] (sunitinib, SU-11248), VEGF trap, vatalanib, ZACTIMA[™] (vandetanib, ZD-6474) and the like.

Antimetabolites include ALIMTA[®] (pemetrexed disodium, LY231514, MTA), 5-azacitidine, XELODA[®] (capecitabine), carmofur, LEUSTAT[®] (cladribine), clofarabine,
25 cytarabine, cytarabine ocfosfate, cytosine arabinoside, decitabine, deferoxamine, doxifluridine, eflornithine, EICAR, enocitabine, ethnylcytidine, fludarabine, hydroxyurea, 5-fluorouracil (5-FU) alone or in combination with leucovorin, GEMZAR[®] (gemcitabine), hydroxyurea, ALKERAN[®] (melphalan), mercaptopurine, 6-mercaptopurine riboside, methotrexate, mycophenolic acid, nelarabine, nolatrexed, ocfosfate, pelitrexol, pentostatin,
30 raltitrexed, Ribavirin, triapine, trimetrexate, S-1, tiazaforin, tegafur, TS-1, vidarabine, UFT and the like.

Antibiotics include intercalating antibiotics aclarubicin, actinomycin D, amrubicin, annamycin, adriamycin, BLENOXANE[®] (bleomycin), daunorubicin, CAELYX[®] or MYOCET[®] (doxorubicin), elsamitrucin, epirubicin, glarbuicin, ZAVEDOS[®] (idarubicin), mitomycin C, nemorubicin, neocarzinostatin, peplomycin, pirarubicin, rebeccamycin, 5 stimalamer, streptozocin, VALSTAR[®] (valrubicin), zinostatin and the like.

Topoisomerase inhibitors include aclarubicin, 9-aminocamptothecin, amonafide, amsacrine, becatecarin, belotecan, BN-80915, CAMPTOSAR[®] (irinotecan hydrochloride), camptothecin, CARDIOXANE[®] (dexrazoxine), diflomotecan, edotecarin, ELLENCE[®] or 10 PHARMORUBICIN[®] (epirubicin), etoposide, exatecan, 10-hydroxycamptothecin, gimatecan, lurtotecan, mitoxantrone, orathecin, pirarubicin, pixantrone, rubitecan, sobuzoxane, SN-38, tafluposide, topotecan and the like.

Antibodies include AVASTIN[®] (bevacizumab), CD40-specific antibodies, chTNT- 15 1/B, denosumab, ERBITUX[®] (cetuximab), HUMAX-CD4[®] (zanolimumab), IGF1R-specific antibodies, lintuzumab, PANOREX[®] (edrecolomab), RENCAREX[®] (WX G250), RITUXAN[®] (rituximab), ticilimumab, trastuzumab and the like.

Hormonal therapies include ARIMIDEX[®] (anastrozole), AROMASIN[®] 20 (exemestane), arzoxifene, CASODEX[®] (bicalutamide), CETROTIDE[®] (cetrorelix), degarelix, deslorelin, DESOPAN[®] (trilostane), dexamethasone, DROGENIL[®] (flutamide), EVISTA[®] (raloxifene), fadrozole, FARESTON[®] (toremifene), FASLODEX[®] (fulvestrant), FEMARA[®] (letrozole), formestane, glucocorticoids, HECTOROL[®] (doxercalciferol), RENAGEL[®] (sevelamer carbonate), lasofoxifene, leuprolide acetate, MEGACE[®] 25 (megesterol), MIFEPREX[®] (mifepristone), NILANDRON[™] (nilutamide), NOLVADEX[®] (tamoxifen citrate), PLENAXIS[™] (abarelix), prednisone, PROPECIA[®] (finasteride), rilostane, SUPREFACT[®] (buserelin), TRELSTAR[®] (luteinizing hormone releasing hormone (LHRH)), VANTAS (histrelin implant), VETORYL[®] (trilostane or modrastane), ZOLADEX[®] (fosrelin, goserelin) and the like.

Deltoids and retinoids include seocalcitrol (EB1089, CB1093), lexacalcitrol (KH1060), fenretinide, PANRETIN[®] (aliretinoin), ATRAGEN[®] (liposomal tretinoin), TARGRETIN[®] (bexarotene), LGD-1550 and the like.

5 Plant alkaloids include, but are not limited to, vincristine, vinblastine, vindesine, vinorelbine and the like.

Proteasome inhibitors include VELCADE[®] (bortezomib), MG132, NPI-0052, PR-171 and the like.

10

Examples of immunologicals include interferons and other immune-enhancing agents. Interferons include interferon alpha, interferon alpha-2a, interferon alpha-2b, interferon beta, interferon gamma-1a, ACTIMMUNE[®] (interferon gamma-1b), or interferon gamma-n1, combinations thereof and the like. Other agents include ALFAFERONE[®], BAM-002, BEROMUN[®] (tasonermin), BEXXAR[®] (tositumomab), CamPath[®] (alemtuzumab), CTLA4 (cytotoxic lymphocyte antigen 4), decarbazine, denileukin, epratuzumab, GRANOCYTE[®] (lenograstim), lentinan, leukocyte alpha interferon, imiquimod, MDX-010, melanoma vaccine, mitumomab, molgramostim, MYLOTARG[™] (gemtuzumab ozogamicin), NEUPOGEN[®] (filgrastim), OncoVAC-CL, OvaRex[®] (oregovomab), pemtumomab (Y-muHMFG1), PROVENGE[®], sargaramostim, sizofilan, teceleukin, TheraCys[®], ubenimex, VIRULIZIN[®], Z-100, WF-10, PROLEUKIN[®] (aldesleukin), ZADAXIN[®] (thymalfasin), ZENAPAX[®] (daclizumab), ZEVALIN[®] (90Y-Ibritumomab tiuxetan) and the like.

25 Biological response modifiers are agents that modify defense mechanisms of living organisms or biological responses, such as survival, growth, or differentiation of tissue cells to direct them to have anti-tumor activity and include krestin, lentinan, sizofiran, picibanil PF-3512676 (CpG-8954), ubenimex and the like.

Pyrimidine analogs include cytarabine (ara C or Arabinoside C), cytosine arabinoside, doxifluridine, FLUDARA[®] (fludarabine), 5-FU (5-fluorouracil), floxuridine, GEMZAR[®] (gemcitabine), TOMUDEX[®] (ratitrexed), TROXATYL[™] (triacytyluridine troxacitabine) and the like.

30

Purine analogs include LANVIS[®] (thioguanine) and PURI-NETHOL[®] (mercaptopurine).

Antimitotic agents include batabulin, epothilone D (KOS-862), N-(2-((4-hydroxyphenyl)amino)pyridin-3-yl)-4-methoxybenzenesulfonamide, ixabepilone (BMS 247550), paclitaxel, TAXOTERE[®] (docetaxel), PNU100940 (109881), patupilone, XRP-9881, vinflunine, ZK-EPO and the like.

- 5 Compounds of the present invention are also intended to be used as a radiosensitizer that enhances the efficacy of radiotherapy. Examples of radiotherapy include, but are not limited to, external beam radiotherapy, teletherapy, brachytherapy and sealed and unsealed source radiotherapy.
- 10 Additionally, ABT-869 Crystalline Form 2 may be combined with other chemotherapeutic agents such as ABRAXANE[™] (ABI-007), ABT-100 (farnesyl transferase inhibitor), ADVEXIN[®], ALTOCOR[®] or MEVACOR[®] (lovastatin), AMPLIGEN[®] (poly I:poly C12U, a synthetic RNA), APTOSYN[™] (exisulind), AREDIA[®] (pamidronic acid), arglabin, L-asparaginase, atamestane (1-methyl-3,17-dione-androsta-1,4-diene), AVAGE[®]
- 15 (tazarotene), AVE-8062, BEC2 (mitumomab), cachectin or cachexin (tumor necrosis factor), canvaxin (vaccine), CeaVac[™] (cancer vaccine), CELEUK[®] (celmoleukin), CEPLENE[®] (histamine dihydrochloride), CERVARIX[™] (human papillomavirus vaccine), CHOP[®] (C: CYTOXAN[®] (cyclophosphamide); H: ADRIAMYCIN[®] (hydroxydoxorubicin);
- O: Vincristine (ONCOVIN[®]); P: prednisone), CyPat[™], combrestatin A4P, DAB(389)EGF or
- 20 TransMID-107R[™] (diphtheria toxins), dacarbazine, dactinomycin, 5,6-dimethylxanthenone-4-acetic acid (DMXAA), eniluracil, EVIZON[™] (squalamine lactate), DIMERICINE[®] (T4N5 liposome lotion), discodermolide, DX-8951f (exatecan mesylate), enzastaurin, EPO906, GARDASIL[®] (quadrivalent human papillomavirus (Types 6, 11, 16, 18) recombinant vaccine), GASTRIMMUNE[™] (gastrin-diphtheria conjugate), GENASENSE[™] (oblimersen
- 25 sodium), GMK (ganglioside conjugate vaccine), GVAX[®] (prostate cancer vaccine), halofuginone, histerelin, hydroxycarbamide, ibandronic acid, IGN-101, IL-13-PE38, IL-13-PE38QQR (cintredekin besudotox), IL-13-pseudomonas exotoxin, interferon- α , interferon- γ , JUNOVAN[™] or MEPACT[™] (mifamurtide), lonafarnib, 5,10-methylenetetrahydrofolate,

miltefosine (hexadecylphosphocholine), NEOVASTAT[®] (AE-941), NEUTREXIN[®]
 (trimetrexate glucuronate), NIPENT[®] (pentostatin), ONCONASE[®] (a ribonuclease enzyme),
 ONCOPHAGE[®] (melanoma vaccine treatment), OncoVAX (IL-2 Vaccine), ORATHECIN[™]
 (rubitecan), OSIDEM[®] (antibody-based cell drug), OvaRex[®] MAb (murine monoclonal
 5 antibody), paclitaxel, PANDIMEX[™] (aglycone saponins from ginseng comprising
 20(S)protopanaxadiol (aPPD) and 20(S)protopanaxatriol (aPPT)), panitumumab,
 PANVAC[®]-VF (investigational cancer vaccine), pegaspargase, PEG Interferon A,
 phenoxodiol, procarbazine, rebimastat, REMOVAB[®] (catumaxomab), REVLIMID[®]
 (lenalidomide), RSR13 (efaproxiral), SOMATULINE[®] LA (lanreotide), SORIATANE[®]
 10 (acitretin), staurosporine (Streptomyces staurospores), talabostat (PT100), TARGRETIN[®]
 (bexarotene), Taxoprexin[®] (DHA-paclitaxel), TELCYTA[™] (TLK286), temilifene,
 TEMODAR[®] (temozolomide), tesmilifene, thalidomide, THERATOPE[®] (STn-KLH),
 thymitaq (2-amino-3,4-dihydro-6-methyl-4-oxo-5-(4-pyridylthio)quinazoline
 dihydrochloride), TNFerade[™] (adenovector: DNA carrier containing the gene for tumor
 15 necrosis factor- α), TRACLEER[®] or ZAVESCA[®] (bosentan), tretinoin (Retin-A), tetrandrine,
 TRISENOX[®] (arsenic trioxide), VIRULIZIN[®], ukrain (derivative of alkaloids from the
 greater celandine plant), vitaxin (anti- α v β 3 antibody), XCYTRIN[®] (motexafin
 gadolinium), XINLAY[™] (atrasentan), XYOTAX[™] (paclitaxel poliglumex), YONDELIS[™]
 (trabectedin), ZD-6126, ZINECARD[®] (dexrazoxane), zometa (zoledronic acid), zorubicin
 20 and the like.

It is also expected that ABT-869 Crystalline Form 2 would inhibit growth of cells
 derived from a pediatric cancer or neoplasm including embryonal rhabdomyosarcoma,
 pediatric acute lymphoblastic leukemia, pediatric acute myelogenous leukemia, pediatric
 25 alveolar rhabdomyosarcoma, pediatric anaplastic ependymoma, pediatric anaplastic large cell
 lymphoma, pediatric anaplastic medulloblastoma, pediatric atypical teratoid/rhabdoid tumor
 of the central nervous system, pediatric biphenotypic acute leukemia, pediatric Burkitts
 lymphoma, pediatric cancers of Ewing's family of tumors such as primitive neuroectodermal
 tumors, pediatric diffuse anaplastic Wilm's tumor, pediatric favorable histology Wilm's
 30 tumor, pediatric glioblastoma, pediatric medulloblastoma, pediatric neuroblastoma, pediatric
 neuroblastoma-derived myelocytomatosis, pediatric pre-B-cell cancers (such as leukemia),

pediatric psteosarcoma, pediatric rhabdoid kidney tumor, pediatric rhabdomyosarcoma, and pediatric T-cell cancers such as lymphoma and skin cancer and the like.

Preparation of ABT-869 and its utility as a PTK inhibitor is described in
5 commonly-owned United States Patent 7297709.

The following examples are presented to provide what is believed to be the most useful and readily understood description of procedures and conceptual aspects of this invention.

10 EXAMPLE 1

Preparation of ABT-869· $\frac{1}{4}$ Ethanolate Crystalline Form 1

A mixture of ABT-869 hydrochloride in ethyl acetate and ethanol, in which the ABT-869 hydrochloride was completely soluble, was mixed with dibasic sodium phosphate. The organic layer was separated, treated with decolorizing carbon, and filtered. A small
15 quantity of L-ascorbic acid was added, and the solution was concentrated. The ethyl acetate was removed by azeotropic distillation with ethanol. Additional ethanol may be added and the solution heated to dissolve any solid that forms. The solution was cooled to 25°C and diluted with water, causing ABT-869· $\frac{1}{4}$ Ethanolate Crystalline Form 1 to crystallize. The product was isolated, washed with water, and dried under reduced pressure, while
20 monitoring residual ethyl acetate and ethanol (by gas chromatography (GC)) and water (by Karl Fischer (KF)). A yield of 92% is typical.

EXAMPLE 2

Preparation of ABT-869 Hydrate Crystalline Form 1

25 Following the neutralization with dibasic sodium phosphate, decolorizing carbon treatment and the removal of ethyl acetate as described in EXAMPLE 1, the mixture of ABT-869 in ethanol was gradually mixed with water at 25°C, with vigorous agitation. ABT-869·H₂O Crystalline Form 1 was isolated, washed with water, and dried under reduced pressure while monitoring residual ethyl acetate and ethanol (by GC) and water (by KF).
30 The dried material may be delumped/milled to control particle size. A yield of 76% is typical.

EXAMPLE 3

Preparation of ABT-869 Crystalline Form 2

A mixture of EXAMPLE 2 was stored under nitrogen at 0% humidity.

Diffraction patterns were collected using an Inel G3000 diffractometer equipped with an incident beam germanium monochromator to provide Cu-K α_1 radiation. The X-ray generator was operated at a voltage of 40 kV and a current of 30 mA. The Inel G3000 is equipped with a position sensitive detector that monitors all diffraction data simultaneously. The detector was calibrated by collecting the attenuated direct beam for seven seconds in 1 degree intervals across a 90 degree two theta range. The calibration was checked against a silicon line position reference standard (NIST 640c). Powder X-ray diffraction (PXRD) analysis was conducted in the following manner. A sample of hydrate powder was placed in a thin layer on the sample holder and gently flattened with a glass microscope slide. The hydrate sample was then equilibrated under dry nitrogen purge at atmospheric temperature and pressure using the low temperature furnace for the Inel G3000 powder X-ray diffractometer. The low temperature furnace contains a Kapton® window that results in a broad diffraction peak at approximately 6 degrees two theta. The sample was equilibrated under nitrogen overnight, followed by PXRD data acquisition for two hours under continued nitrogen purge.

Figure 1 is a powder x-ray diffraction pattern for this Crystalline Form 2.

It is meant to be understood that relative intensities of peak heights and/or intensities in a PXRD pattern may vary and will be dependent on variables such as the temperature, size of crystal size or morphology, sample preparation, or sample height in the analysis well of the X-ray diffractometer.

It is also meant to be understood that peak positions may vary when measured with different radiation sources. For example, Cu-K α_1 , Mo-K α , Co-K α and Fe-K α radiation, having wavelengths of 1.54060 Å, 0.7107 Å, 1.7902 Å and 1.9373 Å, respectively, may provide peak positions which differ from those measured with Cu-K α radiation, which has a wavelength of 1.5478 Å.

The term “about” preceding a series of peak positions means that all of the peaks of the group which it precedes are reported in terms of angular positions (two theta) with an allowable variability of $\pm 0.1^\circ$ as specified by the U.S. Pharmacopeia, pages 1843-1884

(1995). The variability of $\pm 0.1^\circ$ is intended to be used when comparing two powder X-ray diffraction patterns. In practice, if a diffraction pattern peak from one pattern is assigned a range of angular positions (two theta) which is the measured peak position $\pm 0.1^\circ$ and if those ranges of peak positions overlap, then the two peaks are considered to have the same angular position. For example, if a peak from one pattern is determined to have a position of 11.0° , for comparison purposes the allowable variability allows the peak to be assigned a position in the range of 10.9° - 11.1° .

Accordingly, for example, the phrase “about 6.7° , 10.1° , 11.8° , 13.4° , 14.4° , 15.4° , 16.6° , 16.8° , 17.8° , 20.5° and 23.4° means about 6.7° , about 10.1° , about 11.8° , about 13.4° , about 14.4° , about 15.4° , about 16.6° , about 16.8° , about 17.8° , about 20.5° and about 23.4° , which, in turn, means $6.7^\circ \pm 0.1^\circ$, $10.1^\circ \pm 0.1^\circ$, $11.8^\circ \pm 0.1^\circ$, $13.4^\circ \pm 0.1^\circ$, $14.4^\circ \pm 0.1^\circ$, $15.4^\circ \pm 0.1^\circ$, $16.6^\circ \pm 0.1^\circ$, $16.8^\circ \pm 0.1^\circ$, $17.8^\circ \pm 0.1^\circ$, $20.5^\circ \pm 0.1^\circ$, and $23.4^\circ \pm 0.1^\circ$.


The term “about” preceding a temperature means the given temperature $\pm 2^\circ\text{C}$. For example, about 25°C means $25^\circ\text{C} \pm 2^\circ\text{C}$ or 23°C - 27°C .

The foregoing is meant to be illustrative of the invention and not intended to limit it to the disclosed embodiments. Variations and changes obvious to one skilled in the art are intended to be within the scope and nature of the invention as defined in the claims.

WE CLAIM:

1. N-[4-(3-Amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea Crystalline Form 2 which, when measured at about 25°C with radiation at 1.54178 Å, is characterized by a powder diffraction pattern having respective 2θ values of about 6.7°, 10.1°, 11.8°, 13.4°, 14.4°, 15.4°, 16.6°, 16.8°, 17.8°, 20.5° and 23.4°.

1/1
FIG. 1



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2008/080068

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D231/56 A61K31/423 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DAI Y ET AL: "Discovery of N-(4-(3-amino-1H-indazol-4-yl)phenyl)-N'-(2-fluoro-5-methylphenyl)urea (ABT-869), a 3-aminoindazole-based orally active multitargeted receptor tyrosine kinase" JOURNAL OF MEDICINAL CHEMISTRY,, vol. 50, 1 January 2007 (2007-01-01), pages 1584-1587, XP002507892 compound 17p -----	1



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

9 January 2009

Date of mailing of the international search report


19/01/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Diederer, Jeroen



nsité (comptages)
ux-Theta (deg)