

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

28 December 2023 (28.12.2023)



(10) International Publication Number

WO 2023/249875 A1

(51) International Patent Classification:

A61K 31/395 (2006.01) A61K 31/445 (2006.01)

A61K 31/435 (2006.01) A61K 31/33 (2006.01)

A61K 31/18 (2006.01)

DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, ME, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/US2023/025389

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(22) International Filing Date:

15 June 2023 (15.06.2023)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/353,985 21 June 2022 (21.06.2022) US

(71) Applicant: **ALKERMES, INC.** [US/US]; 900 Winter Street, Waltham, MA 02451 (US).

(72) Inventors: **HANDE, Sudhir, Mahadeo**; 28 Constitution Road, Lexington, MA 02421 (US). **HU, Yuan**; 4908 Stearns Hill Road, Waltham, MA 02451 (US). **CHOI, Younggi**; 37 Golden Drive, Stow, MA 01775 (US). **HUYNH, Hoan**; 43 Fiske Avenue, Waltham, MA 02453 (US). **AQUILA, Brian, M.**; 917 Elm Street, Marlborough, MA 01752 (US). **MUGGE, Ingo, Andreas**; 59 Fiske Street, Apt. 1, Waltham, MA 02451 (US). **RAYMER, Brian, Kenneth**; 3 Pilgrim Road, Holliston, MA 01746 (US). **SUH, Byung-Chul**; 504 Manor Terrace, Lexington, MA 02420 (US). **PENNINGTON, Lewis, D.**; 100 Columbia Road, Arlington, MA 02474 (US). **LEHMANN, Jonathan, Ward**; 12A Seven Springs Lane, Burlington, MA 01803 (US).

(74) Agent: **ZUCCHERO, Joseph, C.** et al.; Elmore Patent Law Group, P.C., 484 Groton Road, Westford, MA 01886 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CV, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, MG, MK, MN, MU, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, CV, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SC, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ,

(54) Title: SUBSTITUTED PYRROLIDINYL AND PIPERIDINYL COMPOUNDS AND RELATED METHODS OF TREATMENT

(57) Abstract: The present invention provides compounds useful for the treatment of narcolepsy or cataplexy in a subject in need thereof. Related pharmaceutical compositions and methods are also provided herein.



WO 2023/249875 A1

SUBSTITUTED PYRROLIDINYL AND PIPERIDINYL COMPOUNDS AND RELATED METHODS OF TREATMENT

RELATED APPLICATION

This application claims the benefit of U.S. Provisional patent application serial No.: 63/353,985, filed on June 21, 2022. The entire contents of the above-identified application is herein incorporated by reference.

5

TECHNICAL FIELD

The present invention relates to substituted pyrrolidinyl and piperidinyl compounds, particularly, substituted pyrrolidinyl and piperidinyl compounds having agonist activity.

BACKGROUND OF THE INVENTION

Orexin is a neuropeptide synthesized and released by a subpopulation of neurons within the lateral hypothalamus and its surrounding regions. It consists of two subtypes: orexin A and orexin B. Orexin A and orexin B bind to orexin receptors. Orexin receptors are G protein-coupled receptors expressed preferentially in the brain. There are two subtypes (type 1 and type 2) of orexin receptors (Cell, Vol. 92, 573-585, 1998). Activation of orexin receptors is known to be important for a variety of central nervous system functions, such as maintenance of wakefulness, energy homeostasis, reward processing and motivation (Saper *et al.*, TRENDS in Neuroscience 2001; Yamanaka *et al.*, Neuron 2003; Sakurai, Nature Reviews Neuroscience 2014).

Narcolepsy is a neurological disease that results in excessive daytime sleepiness, sudden bouts of muscular paralysis (cataplexy), and disrupted sleep patterns (Mahoney *et al.*, Nature Reviews Neuroscience, 2019). It is known that narcolepsy is caused by the degeneration of orexin neurons. Narcoleptic symptoms can be modeled in transgenic mice engineered to degenerate orexin neurons, and their symptoms can be reversed by intraventricular administration of orexin peptides (Proc. Natl. Acad. Sci. USA, Vol. 101, 4649-4654, 2004). Studies of orexin-2 receptor knockout mice have suggested that the orexin-2 receptor plays a preferential role in maintaining wakefulness (Cell, Vol. 98, 437-451, 1999, Neuron, Vol. 38, 715-730, 2003). As such, orexin-2 receptor agonists can be therapeutic agents for narcolepsy or other disorders exhibiting excessive daytime sleepiness, such as Parkinson's disease (CNS Drugs, Vol. 27, 83-90, 2013; Brain, Vol. 130, 2007, 1586-1595).


A compound having agonist activity at the orexin-2 receptor is hypothesized to be useful as a novel therapeutic agent for narcolepsy, idiopathic hypersomnia, hypersomnia, sleep apnea syndrome, disturbance of consciousness such as coma and the like, narcolepsy syndrome, hypersomnolence syndrome characterized by hypersomnia (e.g., in Parkinson's disease, Guillain-Barre syndrome or Kleine Levin syndrome), Alzheimer's disease, obesity, insulin resistance syndrome, cardiac failure, diseases related to bone loss, or sepsis and the like. (Cell Metabolism, Vol. 9, 64-76, 2009; Neuroscience, Vol. 121, 855-863, 2003; Respiration, Vol. 71, 575-579, 2004; Peptides, Vol. 23, 1683-1688, 2002; WO 2015/073707; Journal of the American College of Cardiology, Vol. 66, 2015, pages 2522-2533; WO 2015/048091; WO 2015/147240).

Some compounds having orexin-2 receptor agonist activity have been reported (U.S. Pat. No. 8,258,163; WO 2015/088000; WO 2014/198880; Journal of Medicinal Chemistry, Vol. 58, pages 7931-7937; US 20190040010; US 20190031611; US 20170226137). However, it is considered that these compounds are not satisfactory, for example, in terms of activity, pharmacokinetics, permeability into the brain/central nervous system or safety, and the development of an improved compound having orexin-2 receptor agonist activity is desired.

SUMMARY OF THE INVENTION

The present invention aims to provide piperidinyl and pyrrolidinyl compounds having orexin-2 receptor agonist activity.

Accordingly, in an initial aspect, the present invention provides a compound represented by Formula I-A or a pharmaceutically acceptable salt thereof:



wherein:

ring A is selected from the group consisting of phenyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, further wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen;

5 X is N or CH;

Y is S(=O)₂, C(=O), or S(=O)(=NR_e);

R_e is selected from the group consisting of H, C₁-C₃ alkyl, or C₃-C₅ cycloalkyl;

E is selected from the group consisting of NR_aR_b, C₁-C₃ alkylene-NR_aR_b, C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-
 10 membered heterocyclyl, C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-membered heteroaryl and C₁-C₃ alkylene-(5- to 7-membered heteroaryl), wherein the C₁-C₃ alkylene-NR_aR_b, C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-membered heterocyclyl, C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-
 15 membered heteroaryl or C₁-C₃ alkylene-(5- to 7-membered heteroaryl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy;

R_a and R_b are each, independently, selected from the group consisting of H, C₁-C₃ alkyl, C₃-C₅ cycloalkyl, and 4- to 7-membered heterocyclyl, wherein the C₁-C₃ alkyl, C₃-C₅
 20 cycloalkyl, or 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, C₁-C₃ alkyl, or C₁-C₃ alkoxy;

or, alternatively, R_a and R_b, together with the N atom to which they are attached, form a 4- to 7-membered heterocyclyl or 5- to 7-membered heteroaryl, wherein the 4- to 7-
 membered heterocyclyl or 5- to 7-membered heteroaryl is unsubstituted or substituted with
 25 one or more halogen, hydroxyl, NR_cR_d, C₁-C₃ alkyl, C₁-C₃ alkoxy, or C₁-C₃ alkyl substituted with 1-3 halogen;

R₁ is selected from the group consisting of C(=O)-C₁-C₄ alkyl, C(=O)-C₁-C₄ alkoxy, C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), C(=O)-(CR_cR_d)_n-(5- to 10-membered heteroaryl), C(=O)-O-
 30 (CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), (CR_cR_d)_n-(C₆-C₁₀ aryl) and (CR_cR_d)_n-(5- to 10-membered heteroaryl), wherein the C₁-C₄ alkyl, C₁-C₄ alkoxy, C₃-C₈ cycloalkyl, 4- to 7-membered heterocyclyl, C₆-C₁₀ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

R_c and R_d are each, independently, H, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

n is 0, 1, or 2;

each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is, independently, H, halogen, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

or, alternatively, R_3 and R_6 , together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;


or, alternatively, R_4 and R_5 , together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;

m is 0 or 1;

p is 0, 1, 2, 3, or 4; and

each R_9 is, independently, selected from the group consisting of deuterium, halogen, hydroxyl, and cyano.

In one embodiment, provided herein are compounds of Formula I-A having the structure of Formula I or a pharmaceutically acceptable salt thereof:



wherein:

ring A is selected from the group consisting of phenyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, further wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen;

X is N or CH;

Y is S(=O)₂, C(=O), or S(=O)(=NR_e);

R_e is selected from the group consisting of H, C₁-C₃ alkyl, or C₃-C₅ cycloalkyl;

E is selected from the group consisting of NR_aR_b , $\text{C}_1\text{-C}_3$ alkylene- NR_aR_b , $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_2\text{-C}_4$ alkenyl, $\text{C}_2\text{-C}_4$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_3\text{-C}_8$ cycloalkyl), 4- to 10-membered heterocyclyl, $\text{C}_1\text{-C}_3$ alkylene-(4- to 10-membered heterocyclyl), $\text{C}_6\text{-C}_{10}$ aryl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_6\text{-C}_{10}$ aryl), 5- to 7-membered heteroaryl and $\text{C}_1\text{-C}_3$ alkylene-(5- to 7-membered heteroaryl), wherein the $\text{C}_1\text{-C}_3$ alkylene- NR_aR_b , $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_2\text{-C}_4$ alkenyl, $\text{C}_2\text{-C}_4$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_3\text{-C}_8$ cycloalkyl), 4- to 10-membered heterocyclyl, $\text{C}_1\text{-C}_3$ alkylene-(4- to 10-membered heterocyclyl), $\text{C}_6\text{-C}_{10}$ aryl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_6\text{-C}_{10}$ aryl), 5- to 7-membered heteroaryl or $\text{C}_1\text{-C}_3$ alkylene-(5- to 7-membered heteroaryl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy;

R_a and R_b are each, independently, selected from the group consisting of H, $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_3\text{-C}_5$ cycloalkyl, and 4- to 7-membered heterocyclyl, wherein the $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_3\text{-C}_5$ cycloalkyl, or 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy;

or, alternatively, R_a and R_b , together with the N atom to which they are attached, form a 4- to 7-membered heterocyclyl or 5- to 7-membered heteroaryl, wherein the 4- to 7-membered heterocyclyl or 5- to 7-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d , $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_1\text{-C}_3$ alkoxy, or $\text{C}_1\text{-C}_3$ alkyl substituted with 1-3 halogen;

R_1 is selected from the group consisting of $\text{C}(=\text{O})\text{-C}_1\text{-C}_4$ alkyl, $\text{C}(=\text{O})\text{-C}_1\text{-C}_4$ alkoxy, $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-C}_3\text{-C}_8$ cycloalkyl, $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-(4- to 7-membered heterocyclyl)}$, $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-(C}_6\text{-C}_{10}\text{ aryl)}$, $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-(5- to 10-membered heteroaryl)}$, $\text{C}(=\text{O})\text{-O-(CR}_c\text{R}_d)_n\text{-C}_3\text{-C}_8$ cycloalkyl, $\text{C}(=\text{O})\text{-O-(CR}_c\text{R}_d)_n\text{-(4- to 7-membered heterocyclyl)}$, $\text{(CR}_c\text{R}_d)_n\text{-(C}_6\text{-C}_{10}\text{ aryl)}$ and $\text{(CR}_c\text{R}_d)_n\text{-(5- to 10-membered heteroaryl)}$, wherein the $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_3\text{-C}_8$ cycloalkyl, 4- to 7-membered heterocyclyl, $\text{C}_6\text{-C}_{10}$ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkyl substituted with one or more halogen or deuterium;

R_c and R_d are each, independently, H, unsubstituted $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkyl substituted with one or more halogen or deuterium;

n is 0, 1, or 2;

each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is, independently, H, halogen, unsubstituted $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkyl substituted with one or more halogen or deuterium;

or, alternatively, R_3 and R_6 , together form an unsubstituted $\text{C}_1\text{-C}_3$ alkylene or a $\text{C}_1\text{-C}_3$ alkylene substituted with one or more halogen;

or, alternatively, R₄ and R₅, together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;

m is 0 or 1;

p is 0, 1, 2, 3, or 4; and

5 each R₉ is, independently, selected from the group consisting of deuterium, halogen, hydroxyl, and cyano.

Also provided herein is a pharmaceutical composition comprising a compound of any of Formula I-A, or I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

10 In another aspect, provided herein is a method of treating narcolepsy in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

In another aspect, provided herein is a method of treating cataplexy in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a
15 pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION OF THE INVENTION

Provided herein are compounds, e.g., the compounds of Formula I-A, or I, or pharmaceutically acceptable salts thereof, that are useful in the treatment of narcolepsy or
20 cataplexy in a subject.

In a non-limiting aspect, these compounds may modulate the orexin-2 receptor. In a particular embodiment, the compounds provided herein are considered orexin-2 agonists. As such, in one aspect, the compounds provided herein are useful in treatment of narcolepsy in a subject by acting as an agonist of the orexin-2 receptor.

25

Definitions

Listed below are definitions of various terms used to describe this invention. These definitions apply to the terms as they are used throughout this specification and claims, unless otherwise limited in specific instances, either individually or as part of a larger group.

30 Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Generally, the nomenclature used herein and the laboratory procedures in cell culture, molecular genetics, organic chemistry, and peptide chemistry are those well-known and commonly employed in the art.

As used herein, the articles “a” and “an” refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element. Furthermore, use of the term “including” as well as other forms, such as “include,” “includes,” and “included,” is not limiting.

5 As used herein, the term “about” will be understood by persons of ordinary skill in the art and will vary to some extent on the context in which it is used. As used herein when referring to a measurable value such as an amount, a temporal duration, and the like, the term “about” is meant to encompass variations of $\pm 20\%$ or $\pm 10\%$, including $\pm 5\%$, $\pm 1\%$, and $\pm 0.1\%$ from the specified value, as such variations are appropriate to perform the disclosed
10 methods.

As used to herein, the term “EC₅₀” refers to the concentration of a compound required to achieve an effect that is 50% of the maximal observed effect of a compound.

The term “agonist,” as used herein, refers to a compound that, when contacted with a target of interest (e.g., the orexin-2 receptor), causes an increase in the magnitude of a certain
15 activity or function of the target compared to the magnitude of the activity or function observed in the absence of the agonist.

The term “treat,” “treated,” “treating,” or “treatment” includes the diminishment or alleviation of at least one symptom associated or caused by the state, disorder or disease being treated. In certain embodiments, the treatment comprises bringing into contact with the
20 orexin-2 receptor an effective amount of a compound of the invention for conditions related to narcolepsy or cataplexy.

As used herein, the term “prevent” or “prevention” means no disorder or disease development if none had occurred, or no further disorder or disease development if there had already been development of the disorder or disease. Also considered is the ability of one to
25 prevent some or all of the symptoms associated with the disorder or disease.

As used herein, the term “patient,” “individual” or “subject” refers to a human or a non-human mammal. Non-human mammals include, for example, livestock and pets, such as ovine, bovine, porcine, canine, feline and murine mammals. Preferably, the patient, subject, or individual is human.

30 As used herein, the terms “effective amount,” “pharmaceutically effective amount,” and “therapeutically effective amount” refer to a nontoxic but sufficient amount of an agent to provide the desired biological result. That result may be reduction or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system.

An appropriate therapeutic amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation.

As used herein, the term “pharmaceutically acceptable” refers to a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound, and is relatively non-toxic, i.e., the material may be administered to an individual without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

As used herein, the term “pharmaceutically acceptable salt” refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of the present invention include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. The phrase “pharmaceutically acceptable salt” is not limited to a mono, or 1:1, salt. For example, “pharmaceutically acceptable salt” also includes bis-salts, such as a bis-hydrochloride salt. Lists of suitable salts are found in Remington’s Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and Journal of Pharmaceutical Science, 66, 2 (1977), each of which is incorporated herein by reference in its entirety.

As used herein, the term “composition” or “pharmaceutical composition” refers to a mixture of at least one compound useful within the invention with a pharmaceutically acceptable carrier. The pharmaceutical composition facilitates administration of the compound to a patient or subject. Multiple techniques of administering a compound exist in the art including, but not limited to, intravenous, oral, aerosol, parenteral, ophthalmic, pulmonary, and topical administration.

As used herein, the term “pharmaceutically acceptable carrier” means a pharmaceutically acceptable material, composition or carrier, such as a liquid or solid filler, stabilizer, dispersing agent, suspending agent, diluent, excipient, thickening agent, solvent or

encapsulating material, involved in carrying or transporting a compound useful within the invention within or to the patient such that it may perform its intended function. Typically, such constructs are carried or transported from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be “acceptable” in the sense of being
5 compatible with the other ingredients of the formulation, including the compound useful within the invention, and not injurious to the patient. Some examples of materials that may serve as pharmaceutically acceptable carriers include: sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth;
10 malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; surface active agents; alginic acid;
15 pyrogen-free water; isotonic saline; Ringer’s solution; ethyl alcohol; phosphate buffer solutions; and other non-toxic compatible substances employed in pharmaceutical formulations.

As used herein, “pharmaceutically acceptable carrier” also includes any and all coatings, antibacterial and antifungal agents, and absorption delaying agents, and the like that
20 are compatible with the activity of the compound useful within the invention and are physiologically acceptable to the patient. Supplementary active compounds may also be incorporated into the compositions. The “pharmaceutically acceptable carrier” may further include a pharmaceutically acceptable salt of the compound useful within the invention. Other additional ingredients that may be included in the pharmaceutical compositions used in
25 the practice of the invention are known in the art and described, for example in Remington’s Pharmaceutical Sciences (Genaro, Ed., Mack Publishing Co., 1985, Easton, PA), which is incorporated herein by reference.

As used herein, the term “alkyl,” by itself or as part of another substituent means, unless otherwise stated, a straight or branched chain hydrocarbon having the number of carbon
30 atoms designated (*i.e.*, C₁₋₆ alkyl means an alkyl having one to six carbon atoms) and includes straight and branched chains. Examples include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *tert*-butyl, pentyl, neopentyl, and hexyl. Other examples of C₁-C₆-alkyl include ethyl, methyl, isopropyl, isobutyl, n-pentyl, and n-hexyl.

As used herein, the term “halo” or “halogen” alone or as part of another substituent means, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom, preferably, fluorine, chlorine, or bromine, more preferably, fluorine or chlorine.

As used herein, the term “alkylene” refers to divalent aliphatic hydrocarbyl groups, for example, having from 1 to 4 carbon atoms that are either straight-chained or branched. This term includes, by way of example, methylene (-CH₂-), ethylene (-CH₂CH₂-), n-propylene (-CH₂CH₂CH₂-), iso-propylene (-CH₂CH(CH₃)-), and the like.

As used herein, the term “alkenyl” denotes a monovalent group derived from a hydrocarbon moiety containing at least two carbon atoms and at least one carbon-carbon double bond. The double bond may or may not be the point of attachment to another group. Alkenyl groups (*e.g.*, C₂-C₈-alkenyl) include, but are not limited to, for example, ethenyl, propenyl, prop-1-en-2-yl, butenyl, 1-methyl-2-buten-1-yl, heptenyl, octenyl and the like.

As used herein, the term “alkynyl” denotes a monovalent group derived from a hydrocarbon moiety containing at least two carbon atoms and at least one carbon-carbon triple bond. The triple bond may or may not be the point of attachment to another group. Alkynyl groups (*e.g.*, C₂-C₈-alkynyl) include, but are not limited to, for example, ethynyl, propynyl, prop-1-yn-2-yl, butynyl, 1-methyl-2-butyn-1-yl, heptynyl, octynyl and the like.

As used herein, the term “alkoxy,” refers to the group -O-alkyl, wherein alkyl is as defined herein. Alkoxy includes, by way of example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec-butoxy, t-butoxy and the like.

As used herein, the term “cycloalkyl” means a non-aromatic carbocyclic system that is partially or fully saturated having 1, 2 or 3 rings wherein such rings may be fused. The term “fused” means that a second ring is present (*i.e.*, attached or formed) by having two adjacent atoms in common (*i.e.*, shared) with the first ring. Cycloalkyl also includes bicyclic structures that may be bridged or spirocyclic in nature with each individual ring within the bicycle varying from 3-8 atoms. The term “cycloalkyl” includes, but is not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, bicyclo[3.1.0]hexyl, spiro[3.3]heptanyl, and bicyclo[1.1.1]pentyl.

As used herein, the term “heterocyclyl” means a non-aromatic carbocyclic system containing 1, 2, 3 or 4 heteroatoms selected independently from N, O, and S and having 1, 2 or 3 rings wherein such rings may be fused, wherein fused is defined above. Heterocyclyl also includes bicyclic structures that may be bridged or spirocyclic in nature with each individual ring within the bicycle varying from 3-8 atoms, and containing 0, 1, or 2 N, O, or

S atoms. The term “heterocyclyl” includes cyclic esters (*i.e.*, lactones) and cyclic amides (*i.e.*, lactams) and also specifically includes, but is not limited to, epoxidyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl (*i.e.*, oxanyl), pyranyl, dioxanyl, aziridinyl, azetidiny, pyrrolidinyl, 2,5-dihydro-1H-pyrrolyl, oxazolidinyl, thiazolidinyl, piperidinyl, morpholinyl, piperazinyl, thiomorpholinyl, 1,3-oxazinanyl, 1,3-thiazinanyl, and the like. For example, the term “heterocyclyl” can include 4- to 10-membered heterocyclyl, 4- to 7-membered heterocyclyl, 5- to 10-membered heterocyclyl, 6- to 10-membered heterocyclyl, 4- to 6-membered heterocyclyl, 4-membered heterocyclyl, 5-membered heterocyclyl, 6-membered heterocyclyl, 7-membered heterocyclyl, 8-membered heterocyclyl, 9-membered heterocyclyl, or 10-membered heterocyclyl.

As used herein, the term “aromatic” refers to a carbocycle or heterocycle with one or more polyunsaturated rings and having aromatic character, *i.e.*, having $(4n + 2)$ delocalized π (pi) electrons, where n is an integer.

As used herein, the term “aryl” means an aromatic carbocyclic system containing 1, 2 or 3 rings, wherein such rings may be fused, wherein fused is defined above. If the rings are fused, one of the rings must be fully unsaturated and the fused ring(s) may be fully saturated, partially unsaturated or fully unsaturated. The term “aryl” includes, but is not limited to, phenyl, naphthyl, indanyl, and 1,2,3,4-tetrahydronaphthalenyl. For example, the term “aryl” can include C₆-C₁₀ aryl, C₆-C₈ aryl, or C₆ aryl (*i.e.*, phenyl).

As used herein, the term “heteroaryl” means an aromatic carbocyclic system containing 1, 2, 3, or 4 heteroatoms selected independently from N, O, and S and having 1, 2, or 3 rings wherein such rings may be fused, wherein fused is defined above. The term “heteroaryl” includes, but is not limited to, furanyl, thiophenyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl and the like. For example, the term “heteroaryl” can include 5- to 10-membered heteroaryl, 5- to 8-membered heteroaryl, 5- to 6-membered heteroaryl, 6- to 10-membered heteroaryl, 6- to 8-membered heteroaryl, 5-membered heteroaryl, 6-membered heteroaryl, 7-membered heteroaryl, 8-membered heteroaryl, 9-membered heteroaryl, or 10-membered heteroaryl.

It is to be understood that if an aryl, heteroaryl, cycloalkyl, or heterocyclyl moiety may be bonded or otherwise attached to a designated moiety through differing ring atoms (*i.e.*, shown or described without denotation of a specific point of attachment), then all possible points are intended, whether through a carbon atom or, for example, a trivalent


nitrogen atom. For example, the term “pyridinyl” means 2-, 3- or 4-pyridinyl, the term “thiophenyl” means 2- or 3-thiophenyl, and so forth.

As used herein, the term “substituted” means that an atom or group of atoms has replaced hydrogen as the substituent attached to another group.

5

Compounds of the Invention

Accordingly, in an initial aspect, the present invention provides a compound represented by Formula I-A or a pharmaceutically acceptable salt thereof:



10 wherein:

ring A is selected from the group consisting of phenyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, further wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen;

15 X is N or CH;

Y is S(=O)₂, C(=O), or S(=O)(=NR_e);

R_e is selected from the group consisting of H, C₁-C₃ alkyl, or C₃-C₅ cycloalkyl;

E is selected from the group consisting of NR_aR_b, C₁-C₃ alkylene-NR_aR_b, C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-
 20 membered heterocyclyl, C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-membered heteroaryl and C₁-C₃ alkylene-(5- to 7-membered heteroaryl), wherein the C₁-C₃ alkylene-NR_aR_b, C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-membered heterocyclyl, C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-

membered heteroaryl or C₁-C₃ alkylene-(5- to 7-membered heteroaryl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy;

R_a and R_b are each, independently, selected from the group consisting of H, C₁-C₃ alkyl, C₃-C₅ cycloalkyl, and 4- to 7-membered heterocyclyl, wherein the C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, C₁-C₃ alkyl, or C₁-C₃ alkoxy;

or, alternatively, R_a and R_b, together with the N atom to which they are attached, form a 4- to 7-membered heterocyclyl or 5- to 7-membered heteroaryl, wherein the 4- to 7-membered heterocyclyl or 5- to 7-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, C₁-C₃ alkyl, C₁-C₃ alkoxy, or C₁-C₃ alkyl substituted with 1-3 halogen;

R₁ is selected from the group consisting of C(=O)-C₁-C₄ alkyl, C(=O)-C₁-C₄ alkoxy, C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), C(=O)-(CR_cR_d)_n-(5- to 10-membered heteroaryl), C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), (CR_cR_d)_n-(C₆-C₁₀ aryl) and (CR_cR_d)_n-(5- to 10-membered heteroaryl), wherein the C₁-C₄ alkyl, C₁-C₄ alkoxy, C₃-C₈ cycloalkyl, 4- to 7-membered heterocyclyl, C₆-C₁₀ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

R_c and R_d are each, independently, H, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

n is 0, 1, or 2;

each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H, halogen, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

or, alternatively, R₃ and R₆, together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;


or, alternatively, R₄ and R₅, together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;

m is 0 or 1;

p is 0, 1, 2, 3, or 4; and

each R₉ is, independently, selected from the group consisting of deuterium, halogen, hydroxyl, and cyano.

In one embodiment, provided herein are compounds of Formula I-A having the structure of Formula I or a pharmaceutically acceptable salt thereof:



wherein:

- 5 ring A is selected from the group consisting of phenyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, further wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen;
- X is N or CH;
- 10 Y is S(=O)₂, C(=O), or S(=O)(=NR_e);
- R_e is selected from the group consisting of H, C₁-C₃ alkyl, or C₃-C₅ cycloalkyl;
- E is selected from the group consisting of NR_aR_b, C₁-C₃ alkylene-NR_aR_b, C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-membered heterocyclyl, C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-membered heteroaryl and C₁-C₃ alkylene-(5- to 7-membered heteroaryl), wherein the C₁-C₃ alkylene-NR_aR_b, C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-membered heterocyclyl, C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-membered heteroaryl or C₁-C₃ alkylene-(5- to 7-membered heteroaryl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy;
- 15 Ra and R_b are each, independently, selected from the group consisting of H, C₁-C₃ alkyl, C₃-C₅ cycloalkyl, and 4- to 7-membered heterocyclyl, wherein the C₁-C₃ alkyl, C₃-C₅
- 20

cycloalkyl, or 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, C₁-C₃ alkyl, or C₁-C₃ alkoxy;

or, alternatively, R_a and R_b, together with the N atom to which they are attached, form a 4- to 7-membered heterocyclyl or 5- to 7-membered heteroaryl, wherein the 4- to 7-membered heterocyclyl or 5- to 7-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, C₁-C₃ alkyl, C₁-C₃ alkoxy, or C₁-C₃ alkyl substituted with 1-3 halogen;

R₁ is selected from the group consisting of C(=O)-C₁-C₄ alkyl, C(=O)-C₁-C₄ alkoxy, C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), C(=O)-(CR_cR_d)_n-(5- to 10-membered heteroaryl), C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), (CR_cR_d)_n-(C₆-C₁₀ aryl) and (CR_cR_d)_n-(5- to 10-membered heteroaryl), wherein the C₁-C₄ alkyl, C₁-C₄ alkoxy, C₃-C₈ cycloalkyl, 4- to 7-membered heterocyclyl, C₆-C₁₀ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

R_c and R_d are each, independently, H, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

n is 0, 1, or 2;

each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H, halogen, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

or, alternatively, R₃ and R₆, together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;

or, alternatively, R₄ and R₅, together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;

m is 0 or 1;

p is 0, 1, 2, 3, or 4; and

each R₉ is, independently, selected from the group consisting of deuterium, halogen, hydroxyl, and cyano.

In one embodiment of Formula (I), ring A is a phenyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a phenyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a phenyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

5 In another embodiment of Formula (I), ring A is a phenyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.

In another embodiment of Formula (I), ring A is a phenyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.

In another embodiment of Formula (I), ring A is a phenyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from F or Cl.

10 In another embodiment of Formula (I), ring A is a phenyl ring, wherein the ring is substituted with 1-2 F substituents.

In another embodiment of Formula (I), ring A is 3-fluorophenyl.

In another embodiment of Formula (I), ring A is a pyridinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a pyridinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

20 In another embodiment of Formula (I), ring A is a pyridinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a pyridinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.

25 In another embodiment of Formula (I), ring A is a pyridinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.

In another embodiment of Formula (I), ring A is a pyridinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.

In another embodiment of Formula (I), ring A is a pyridinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.

30 In another embodiment of Formula (I), ring A is a pyridazinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a pyridazinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

5 In another embodiment of Formula (I), ring A is a pyridazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a pyridazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.

10 In another embodiment of Formula (I), ring A is a pyridazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.

In another embodiment of Formula (I), ring A is a pyridazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.

In another embodiment of Formula (I), ring A is a pyridazinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.

15 In another embodiment of Formula (I), ring A is a pyrimidinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

20 In another embodiment of Formula (I), ring A is a pyrimidinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a pyrimidinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

25 In another embodiment of Formula (I), ring A is a pyrimidinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.

In another embodiment of Formula (I), ring A is a pyrimidinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.

In another embodiment of Formula (I), ring A is a pyrimidinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.

30 In another embodiment of Formula (I), ring A is a pyrimidinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.

In another embodiment of Formula (I), ring A is a pyrazinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a pyrazinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

5 In another embodiment of Formula (I), ring A is a pyrazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a pyrazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.

10 In another embodiment of Formula (I), ring A is a pyrazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.

In another embodiment of Formula (I), ring A is a pyrazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.

In another embodiment of Formula (I), ring A is a pyrazinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.

15 In another embodiment of Formula (I), ring A is a triazinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

20 In another embodiment of Formula (I), ring A is a triazinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

In another embodiment of Formula (I), ring A is a triazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

25 In another embodiment of Formula (I), ring A is a triazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.

In another embodiment of Formula (I), ring A is a triazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.

In another embodiment of Formula (I), ring A is a triazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.

30 In another embodiment of Formula (I), ring A is a triazinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.

In another embodiment of Formula (I), p is 0. In another embodiment of Formula (I), p is 1. In another embodiment of Formula (I), p is 2. In another embodiment of Formula (I), p

is 3. In another embodiment of Formula (I), p is 4. In another embodiment of Formula (I), p is 0, 1 or 2. In another embodiment of Formula (I), p is 0 or 1.

In another embodiment of Formula (I), p is 1 and R₉ is deuterium. In another embodiment of Formula (I), p is 1 and R₉ is halogen. In another embodiment of Formula (I), p is 1 and R₉ is fluorine. In another embodiment of Formula (I), p is 1 and R₉ is hydroxyl. In another embodiment of Formula (I), p is 1 and R₉ is cyano. In another embodiment of Formula (I), p is 2 and each R₉ is hydroxyl. In another embodiment of Formula (I), p is 2 and each R₉ is halogen. In another embodiment of Formula (I), p is 2 and each R₉ is fluorine.

In another embodiment of Formula (I), E is NR_aR_b. In another embodiment of Formula (I), E is C₁-C₃ alkylene-NR_aR_b. In another embodiment of Formula (I), E is unsubstituted C₁-C₃ alkyl, unsubstituted C₂-C₄ alkenyl or unsubstituted C₂-C₄ alkynyl. In another embodiment of Formula (I), E is C₁-C₃ alkyl, C₂-C₄ alkenyl or C₂-C₄ alkynyl substituted with one or more halogen, hydroxyl, C₁-C₃ alkyl, or C₁-C₃ alkoxy. In another embodiment of Formula (I), E is unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), E is C₁-C₃ alkyl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy. In another embodiment of Formula (I), E is unsubstituted C₃-C₈ cycloalkyl. In another embodiment of Formula (I), E is C₃-C₈ cycloalkyl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy. In another embodiment of Formula (I), E is unsubstituted C₁-C₃ alkylene-(C₃-C₈ cycloalkyl). In another embodiment of Formula (I), E is C₁-C₃ alkylene-(C₃-C₈ cycloalkyl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy. In another embodiment of Formula (I), E is unsubstituted 4- to 10-membered heterocyclyl. In another embodiment of Formula (I), E is 4- to 10-membered heterocyclyl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy. In another embodiment of Formula (I), E is unsubstituted C₁-C₃ alkylene-(4- to 10-membered heterocyclyl). In another embodiment of Formula (I), E is C₁-C₃ alkylene-(4- to 10-membered heterocyclyl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy. In another embodiment of Formula (I), E is unsubstituted C₆-C₁₀ aryl. In another embodiment of Formula (I), E is C₆-C₁₀ aryl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy. In another embodiment of Formula (I), E is unsubstituted C₁-C₃ alkylene-(C₆-C₁₀ aryl). In another embodiment of Formula (I), E is C₁-C₃ alkylene-(C₆-C₁₀ aryl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy. In another embodiment of Formula (I), E is unsubstituted 5- to 7-membered

heteroaryl. In another embodiment of Formula (I), E is 5- to 7-membered heteroaryl substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy. In another embodiment of Formula (I), E is unsubstituted $\text{C}_1\text{-C}_3$ alkylene-(5- to 7-membered heteroaryl). In another embodiment of Formula (I), E is $\text{C}_1\text{-C}_3$ alkylene-(5- to 7-membered heteroaryl) substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

In another embodiment of Formula (I), E is $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_2\text{-C}_4$ alkenyl, $\text{C}_2\text{-C}_4$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_3\text{-C}_8$ cycloalkyl), 4- to 10-membered heterocyclyl, or $\text{C}_1\text{-C}_3$ alkylene-(4- to 10-membered heterocyclyl), wherein the $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_2\text{-C}_4$ alkenyl, $\text{C}_2\text{-C}_4$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_3\text{-C}_8$ cycloalkyl), 4- to 10-membered heterocyclyl, or $\text{C}_1\text{-C}_3$ alkylene-(4- to 10-membered heterocyclyl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

In another embodiment of Formula (I), E is $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_3\text{-C}_8$ cycloalkyl), 4- to 10-membered heterocyclyl, or $\text{C}_1\text{-C}_3$ alkylene-(4- to 10-membered heterocyclyl), wherein the $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_3\text{-C}_8$ cycloalkyl), 4- to 10-membered heterocyclyl, or $\text{C}_1\text{-C}_3$ alkylene-(4- to 10-membered heterocyclyl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

In another embodiment of Formula (I), E is $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, or $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_3\text{-C}_8$ cycloalkyl), wherein the $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, or $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_3\text{-C}_8$ cycloalkyl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

In another embodiment of Formula (I), E is methyl. In another embodiment of Formula (I), E is methyl substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy. In another embodiment of Formula (I), E is CF_3 . In another embodiment of Formula (I), E is CHF_2 . In another embodiment of Formula (I), E is CH_2F . In another embodiment of Formula (I), E is $\text{NH}(\text{CH}_3)$. In another embodiment of Formula (I), E is $\text{N}(\text{CH}_3)_2$.

In another embodiment of Formula (I), E is $\text{C}_6\text{-C}_{10}$ aryl, $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_6\text{-C}_{10}$ aryl), 5- to 7-membered heteroaryl or $\text{C}_1\text{-C}_3$ alkylene-(5- to 7-membered heteroaryl). In another embodiment of Formula (I), E is unsubstituted $\text{C}_6\text{-C}_{10}$ aryl or $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_6\text{-C}_{10}$ aryl). In another embodiment of Formula (I), E is $\text{C}_6\text{-C}_{10}$ aryl or $\text{C}_1\text{-C}_3$ alkylene-($\text{C}_6\text{-C}_{10}$ aryl)

substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

In another embodiment of Formula (I), E is 5- to 7-membered heteroaryl or $\text{C}_1\text{-C}_3$ alkylene-(5- to 7-membered heteroaryl). In another embodiment of Formula (I), E is
5 unsubstituted 5- to 7-membered heteroaryl or $\text{C}_1\text{-C}_3$ alkylene-(5- to 7-membered heteroaryl).
In another embodiment of Formula (I), E is 5- to 7-membered heteroaryl or $\text{C}_1\text{-C}_3$ alkylene-(5- to 7-membered heteroaryl) substituted with one or more halogen, hydroxyl, NR_cR_d , CF_3 , CHF_2 , CH_2F , $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

In another embodiment of Formula (I), Y is $\text{S}(=\text{O})_2$. In another embodiment of
10 Formula (I), Y is $\text{C}(=\text{O})$. In another embodiment of Formula (I), Y is $\text{S}(=\text{O})(=\text{NR}_c)$.

In another embodiment of Formula (I), X is N. In another embodiment of Formula (I), X is CH.

In another embodiment of Formula (I), R_1 is selected from the group consisting of $\text{C}(=\text{O})\text{-C}_1\text{-C}_4$ alkyl, $\text{C}(=\text{O})\text{-C}_1\text{-C}_4$ alkoxy, $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-C}_3\text{-C}_8$ cycloalkyl, $\text{C}(=\text{O})\text{-}$
15 $(\text{CR}_c\text{R}_d)_n\text{-(4- to 7-membered heterocyclyl)}$, $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-(C}_6\text{-C}_{10} \text{ aryl)}$, and $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-(5- to 10-membered heteroaryl)}$, wherein the $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_3\text{-C}_8$ cycloalkyl, 4- to 7-membered heterocyclyl, $\text{C}_6\text{-C}_{10}$ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R_1 is selected from the group consisting of $\text{C}(=\text{O})\text{-C}_1\text{-C}_4$ alkyl, $\text{C}(=\text{O})\text{-C}_1\text{-C}_4$ alkoxy, and $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-C}_3\text{-C}_8$ cycloalkyl, wherein
20 the $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, or $\text{C}_3\text{-C}_8$ cycloalkyl, is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R_1 is selected from the group consisting of $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-(4- to 7-membered heterocyclyl)}$, $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-(C}_6\text{-C}_{10} \text{ aryl)}$, and
25 $\text{C}(=\text{O})\text{-(CR}_c\text{R}_d)_n\text{-(5- to 10-membered heteroaryl)}$, wherein the 4- to 7-membered heterocyclyl, $\text{C}_6\text{-C}_{10}$ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkyl substituted with one or more
30 halogen or deuterium.

In another embodiment of Formula (I), R_1 is $\text{C}(=\text{O})\text{-C}_1\text{-C}_4$ alkyl, wherein the $\text{C}_1\text{-C}_4$ alkyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkoxy, wherein the C₁-C₄ alkoxy is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

5 In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, wherein the C₃-C₈ cycloalkyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

10 In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), wherein the 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), wherein the C₆-C₁₀ aryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

15 In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 10-membered heteroaryl), wherein the 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

20 In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heterocyclyl), wherein the 5- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

25 In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 6-membered heterocyclyl), wherein the 5- to 6-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

30 In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), wherein the 5-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), wherein the 6-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n-(C_6 \text{ aryl})$, wherein the C_6 aryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

5 In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n-(C_8 \text{ aryl})$, wherein the C_8 aryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n-(C_{10} \text{ aryl})$, wherein the C_{10} aryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

10 In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n-(5\text{- to }6\text{-membered heteroaryl})$, wherein the 5- to 6-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

15 In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n-(5\text{-membered heteroaryl})$, wherein the 5-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

20 In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n-(6\text{-membered heteroaryl})$, wherein the 6-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

25 In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n-(7\text{-membered heteroaryl})$, wherein the 7-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R_1 is $C(=O)-O-(CR_cR_d)_n-C_3-C_8$ cycloalkyl or $C(=O)-O-(CR_cR_d)_n-(4\text{- to }7\text{-membered heterocyclyl})$, wherein the C_3-C_8 cycloalkyl or 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

30 In another embodiment of Formula (I), R_1 is $C(=O)-O-(CR_cR_d)_n-C_3-C_8$ cycloalkyl, wherein the C_3-C_8 cycloalkyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium. In another embodiment of Formula (I), R_1 is $C(=O)-O-(CR_cR_d)_n-(4\text{- to }7\text{-membered heterocyclyl})$, wherein the 4- to 7-membered heterocyclyl is unsubstituted or

substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₅ cycloalkyl, wherein the C₃-C₅ cycloalkyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5- to 6-membered heterocyclyl), wherein the 5- to 6-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), wherein the 5-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), wherein the 6-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is unsubstituted. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is unsubstituted and further wherein n is 0. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is unsubstituted and further wherein n is 1. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl,

unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is unsubstituted. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is unsubstituted and further wherein n is 0. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is unsubstituted and further wherein n is 1. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is unsubstituted. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is unsubstituted and further wherein n is 0. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is unsubstituted and further wherein n is 1. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.

In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is unsubstituted. In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is unsubstituted and further wherein n is 0. In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is unsubstituted and further wherein n is 1. In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium and further wherein n is 1.

In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is unsubstituted. In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is unsubstituted and further wherein n is 0. In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is unsubstituted and further wherein n is 1. In another embodiment of Formula (I), R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is

substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl) wherein the 6-membered heteroaryl is unsubstituted. In another embodiment of Formula (I),
5 R₁ is (CR_cR_d)_n-(6-membered heteroaryl) wherein the 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium. In another embodiment of Formula (I), R₁ is pyridazinyl. In another embodiment of Formula (I), R₁ is 3-pyridazinyl.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl)
10 wherein the 6-membered heteroaryl is unsubstituted and further wherein n is 0. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl) wherein the 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl)
15 wherein the 6-membered heteroaryl is unsubstituted and further wherein n is 1. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl) wherein the 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.
20

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5-membered heteroaryl)
wherein the 5-membered heteroaryl is unsubstituted. In another embodiment of Formula (I),
R₁ is (CR_cR_d)_n-(5-membered heteroaryl) wherein the 5-membered heteroaryl is substituted
25 with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5-membered heteroaryl)
wherein the 5-membered heteroaryl is unsubstituted and further wherein n is 0. In another
embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5-membered heteroaryl) wherein the 5-
30 membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5-membered heteroaryl)
wherein the 5-membered heteroaryl is unsubstituted and further wherein n is 1. In another
embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5-membered heteroaryl) wherein the 5-

membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.

In another embodiment of Formula (I), each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H, halogen, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium. In another embodiment of Formula (I), each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H, halogen, or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H or halogen. In another embodiment of Formula (I), each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H or fluorine. In another embodiment of Formula (I), each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H or C₁-C₃ alkyl. In another embodiment of Formula (I), each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.

In another embodiment of Formula (I), R₃ and R₆, together, form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen. In another embodiment of Formula (I), R₃ and R₆, together, form an unsubstituted C₂ alkylene or a C₂ alkylene substituted with one or more halogen. In another embodiment of Formula (I), R₃ and R₆, together, form an azabicyclo[3.2.1]octanyl bridged bicyclic heterocyclyl.

In another embodiment of Formula (I), R₄ and R₅, together, form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen. In another embodiment of Formula (I), R₄ and R₅, together, form an unsubstituted C₂ alkylene or a C₂ alkylene substituted with one or more halogen. In another embodiment of Formula (I), R₄ and R₅, together, form an azabicyclo[3.2.1]octanyl bridged bicyclic heterocyclyl.

In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkoxy and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkoxy and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkoxy and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-C₁-C₄ alkoxy and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is

C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another
5 embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of
10 Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of
Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆,
15 R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment
20 of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another
25 embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of
30 Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of

Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered

5 heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula

10 (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another

15 embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(6-membered heterocyclyl), n is 1 or 2, and each of R₂,

20 R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl) and each

25 of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another

30 embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(C₆ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-(CR_cR_d)_n-(5-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another

embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (5-membered heteroaryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or halogen. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (5-membered heteroaryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or fluorine.

5 In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or unsubstituted C_1-C_3 alkyl. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or halogen. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or fluorine.

In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), n is 0, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), n is 0, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or unsubstituted C_1-C_3 alkyl. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), n is 0, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or halogen. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), n is 0, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or fluorine.

15 In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or unsubstituted C_1-C_3 alkyl. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or halogen. In another embodiment of Formula (I), R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or fluorine.

20 In another embodiment of Formula (I), R_1 is $C(=O)-O-(CR_cR_d)_n-C_3-C_8$ cycloalkyl and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H. In another embodiment of Formula (I), R_1 is $C(=O)-O-(CR_cR_d)_n-C_3-C_8$ cycloalkyl and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or unsubstituted C_1-C_3 alkyl. In another embodiment of Formula (I), R_1 is $C(=O)-O-(CR_cR_d)_n-C_3-C_8$ cycloalkyl and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or halogen. In another embodiment of Formula (I), R_1 is $C(=O)-O-(CR_cR_d)_n-C_3-C_8$ cycloalkyl and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1

or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of
5 Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered
10 heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of
15 Formula (I), R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or
20 unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 10-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 7-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 7-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or
25 unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 7-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(5- to 7-membered heteroaryl) and each of R₂, R₃,
30 R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or
unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-

membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(6-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.


In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl), n is 0, and each of R₂, R₃, R₄,

R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.


In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or halogen. In another embodiment of Formula (I), R₁ is (CR_cR_d)_n-(C₆ aryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or fluorine.

Each of the embodiments described herein with respect to compounds of Formula I also applies to compounds of Formula I-A.


According to Formula I-A or I herein, when ring A is pyridinyl, the position of the pyridinyl N atom are specified as shown below:




Further, according to Formula I-A or I herein, when ring A is pyridazinyl, the positions of the pyridazinyl N atoms are specified as shown below:








Further, according to Formula I-A or I herein, when ring A is pyrimidinyl, the positions of the pyrimidinyl N atoms are specified as shown below:







Further, according to Formula I-A or I herein, when ring A is pyrazinyl, the positions of the pyrazinyl N atoms are specified as shown below:








	<p>3</p>
	<p>4</p>
	<p>5</p>
	<p>6</p>
	<p>7</p>

	8
	9
	10
	11
	12









	<p>13</p>
	<p>14</p>
	<p>15</p>
	<p>16</p>
	<p>17</p>









	18
	19
	20
	21
	22








	23
	24
	25
	26
	27








	<p>28</p>
	<p>29</p>
	<p>30</p>
	<p>31</p>
	<p>32</p>


	<p>33</p>
	<p>34</p>
	<p>35</p>
	<p>36</p>
	<p>37</p>
	<p>38</p>

	<p>39</p>
	<p>40</p>
	<p>41</p>
	<p>42</p>
	<p>43</p>
	<p>44</p>
	<p>45</p>
	<p>46</p>

	<p>47</p>
	<p>48</p>
	<p>49</p>
	<p>50</p>
	<p>51</p>
	<p>52</p>
	<p>53</p>
	<p>54</p>

	<p>55</p>
	<p>56</p>
	<p>57</p>
	<p>58</p>
	<p>59</p>
	<p>60</p>
	<p>61</p>

	<p>62</p>
	<p>63</p>
	<p>64</p>
	<p>65</p>
	<p>66</p>
	<p>67</p>
	<p>68</p>



The disclosed compounds possess one or more stereocenters, and each stereocenter may exist independently in either the R or S configuration. In one embodiment, compounds described herein are present in optically active or racemic forms. It is to be understood that the compounds described herein encompass racemic, optically-active, regioisomeric and stereoisomeric forms, or combinations thereof that possess the therapeutically useful properties described herein.

Preparation of optically active forms is achieved in any suitable manner, including by way of non-limiting example, by resolution of the racemic form with recrystallization techniques, synthesis from optically-active starting materials, chiral synthesis, or chromatographic separation using a chiral stationary phase. In one embodiment, a mixture of two or more isomers is utilized as the disclosed compound described herein. In another embodiment, a pure isomer is utilized as the disclosed compound described herein. In another embodiment, compounds described herein contain one or more chiral centers. These compounds are prepared by any means, including stereoselective synthesis, enantioselective synthesis or separation of a mixture of enantiomers or diastereomers. Resolution of compounds and isomers thereof is achieved by any means including, by way of non-limiting example, chemical processes, enzymatic processes, fractional crystallization, distillation, and chromatography.

In one embodiment, the disclosed compounds may exist as tautomers. All tautomers are included within the scope of the compounds presented herein.

Compounds described herein also include isotopically-labeled compounds wherein one or more atoms is replaced by an atom having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes suitable for inclusion in the compounds described herein include and are not limited to ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{36}Cl , ^{18}F , ^{123}I , ^{125}I , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{32}P , and ^{35}S .

In one embodiment, isotopically-labeled compounds are useful in drug or substrate tissue distribution studies. In another embodiment, substitution with heavier isotopes such as deuterium affords greater metabolic stability (for example, increased *in vivo* half-life or reduced dosage requirements). In another embodiment, the compounds described herein
5 include a ^2H (*i.e.*, deuterium) isotope.

In yet another embodiment, substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , is useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled compounds are prepared by any suitable method or by processes using an appropriate isotopically-labeled reagent in place of the non-
10 labeled reagent otherwise employed.

The specific compounds described herein, and other compounds encompassed by one or more of the Formulas described herein having different substituents are synthesized using techniques and materials described herein and as described, for example, in Fieser and Fieser's Reagents for Organic Synthesis, Volumes 1-17 (John Wiley and Sons, 1991); Rodd's
15 Chemistry of Carbon Compounds, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989); Organic Reactions, Volumes 1-40 (John Wiley and Sons, 1991), Larock's Comprehensive Organic Transformations (VCH Publishers Inc., 1989), March, Advanced Organic Chemistry 4th Ed., (Wiley 1992); Carey and Sundberg, Advanced Organic Chemistry 4th Ed., Vols. A and B (Plenum 2000, 2001), and Green and Wuts, Protective Groups in
20 Organic Synthesis 3rd Ed., (Wiley 1999) (all of which are incorporated by reference for such disclosure). General methods for the preparation of compounds as described herein are modified by the use of appropriate reagents and conditions, for the introduction of the various moieties found in the Formulas as provided herein.

Compounds described herein are synthesized using any suitable procedures starting
25 from compounds that are available from commercial sources or are prepared using procedures described herein.

Methods of Treatment

The compounds of the invention can be used in a method of treating a disease or
30 condition in a subject, said method comprising administering to the subject a compound of the invention, or a pharmaceutical composition comprising a compound of the invention. In one embodiment of the methods described herein, the subject is human. In one aspect, the compounds provided herein are useful in treatment of a disease or condition by acting as an agonist of the orexin-2 receptor.

The compounds of the invention can be used to treat a disease or condition selected from the group consisting of narcolepsy, cataplexy, or hypersomnia in a subject in need thereof.

In one embodiment, the compounds of the invention can be used to treat narcolepsy in a subject. In one embodiment, the compounds of the invention can be used to treat cataplexy in a subject. In one embodiment, the compounds of the invention can be used to treat hypersomnia in a subject.

Orexin-2 receptors are important in a wide range of biological functions. This suggests that orexin-2 receptors play a role in diverse disease processes in humans or other species. The compound of the present invention is useful for treating, preventing, or ameliorating the risk of one or more of the following symptoms or diseases of various neurological and psychiatric diseases associated with alterations in sleep/wake function. That is, narcolepsy, narcolepsy with cataplexy, idiopathic hypersomnia, hypersomnia, sleep apnea syndrome, narcolepsy syndrome, hypersomnolence syndrome characterized by hypersomnia (e.g., in subjects with Kleine Levin syndrome, major depression with hypersomnia, Lewy body dementia, Parkinson's disease, progressive supranuclear paralysis, Prader-Willi syndrome, Mobius syndrome, hypoventilation syndrome, Niemann-Pick disease type C, brain contusion, cerebral infarction, brain tumor, muscular dystrophy, multiple sclerosis, multiple systems atrophy, acute disseminated encephalomyelitis, Guillain-Barre syndrome, Rasmussen's encephalitis, Wernicke's encephalitis, limbic encephalitis, or Hashimoto's encephalopathy), coma, loss of consciousness, obesity (e.g., malignant mastocytosis, exogenous obesity, hyperinsular obesity, hyperplasmic obesity, hypop hyseal adiposity, hypoplasmic obesity, hypothyroid obesity, hypothalamic obesity, symptomatic obesity, infantile obesity, upper body obesity, alimentary obesity, hypogonadal obesity, systemic mastocytosis, simple obesity, or central obesity), insulin resistance syndrome, Alzheimer's disease, disturbance of consciousness such as coma and the like, side effects and complications due to anesthesia, sleep disturbance, excessive daytime sleepiness, sleep problem, insomnia, intermittent sleep, nocturnal myoclonus, REM sleep interruption, jet lag, jet lag syndrome, sleep disorder of alternating worker, sleep disorder, night terror, depression, major depression, sleepwalking disease, enuresis, sleep disorder, Alzheimer's dusk, sundowning, diseases associated with circadian rhythm, fibromyalgia, condition arising from decline in the quality of sleep, overeating, obsessive compulsive eating disorder, obesity-related disease, hypertension, diabetes, elevated plasma insulin concentration and insulin resistance, hyperlipidemia, hyperlipemia, endometrial cancer, breast cancer, prostate cancer,

colorectal cancer, cancer, osteoarthritis, obstructive sleep apnea, cholelithiasis, gallstones, cardiac disease, abnormal heartbeat, arrhythmia, myocardial infarction, congestive cardiac failure, cardiac failure, coronary heart disease, cardiovascular disorder, polycysticovarian disease, craniopharingioma, Prader-Willi syndrome, Froelich's syndrome, growth hormone deficient, normal mutant short stature, Turner's syndrome, children suffering from acute lymphoblastic leukemia, syndrome X, reproductive hormone abnormality, declining fertility, infertility, male gonadal function decline, sexual and reproductive dysfunction such as female male hirsutism, fetal defects associated with pregnant women obesity, gastrointestinal motility disorders such as obesity-related gastroesophageal reflux, obesity hypoventilation syndrome (Pickwick syndrome), respiratory diseases such as dyspnea, inflammation such as systemic inflammation of the vascular system, arteriosclerosis, hypercholesterolemia, hyperuricemia, lower back pain, gall bladder disease, gout, kidney cancer, risk of secondary outcomes of obesity, such as lowering the risk of left ventricular hypertrophy, migraine pain, headache, neuropathic pain, Parkinson's disease, psychosis, autoimmune encephalitis, cancer related fatigue (such as excessive daytime sleepiness or fatigue associated with cancer and/or chemotherapy), cancer related nausea and vomiting, corticobasal degeneration, Huntington's disease, neuromyelitis optica, nociception, progressive supranuclear palsy, schizophrenia, systemic lupus erythematosus, traumatic brain injury, facial flushing, night sweats, diseases of the genital/urinary system, diseases related to sexual function or fertility, dysthymic disorder, bipolar disorder, bipolar I disorder, bipolar II disorder, cyclothymic disorder, acute stress disorder, agoraphobia, generalized anxiety disorder, obsessive disorder, panic attack, panic disorder, post-traumatic stress disorder (PTSD), separation anxiety disorder, social phobia, anxiety disorder, acute neurological and psychiatric disorders such as cardiac bypass surgery and post-transplant cerebral deficit, stroke, ischemic stroke, cerebral ischemia, spinal cord trauma, head trauma, perinatal hypoxia, cardiac arrest, hypoglycemic nerve injury, Huntington's chorea, amyotrophic lateral sclerosis, eye damage, retinopathy, cognitive impairment, muscle spasm, tremor, epilepsy, disorders associated with muscle spasticity, delirium, amnesic disorder, age-related cognitive decline, schizoaffective disorder, delusional disorder, drug addiction, dyskinesia, chronic fatigue syndrome, fatigue, medication-induced Parkinsonism syndrome, Jil-do La Tourette's syndrome, chorea, myoclonus, tic, restless legs syndrome, dystonia, dyskinesia, attention deficit hyperactivity disorder (ADHD), behavior disorder, urinary incontinence, withdrawal symptoms, trigeminal neuralgia, hearing loss, tinnitus, nerve damage, retinopathy, macular degeneration, vomiting,

cerebral edema, pain, bone pain, arthralgia, toothache, cataplexy, and traumatic brain injury (TBI).

Particularly, the compound of the present invention is useful as a therapeutic or prophylactic drug for narcolepsy, idiopathic hypersomnia, hypersomnia, sleep apnea
5 syndrome, narcolepsy syndrome, hypersomnolence syndrome characterized by hypersomnia (e.g., in Parkinson's disease, Guillain-Barre syndrome or Kleine Levin syndrome), Alzheimer's disease, obesity, insulin resistance syndrome, cardiac failure, diseases related to bone loss, sepsis, disturbance of consciousness such as coma and the like, side effects and complications due to anesthesia, and the like, or anesthetic antagonist.

10 In one embodiment, the compound of the present invention has orexin-2 receptor agonist activity and is useful as a prophylactic or therapeutic agent for narcolepsy.

In another embodiment, the compound of the present invention is useful as a prophylactic or therapeutic agent for narcolepsy type-1. In another embodiment, the compound of the present invention is useful as a prophylactic or therapeutic agent for
15 narcolepsy type-2. In another embodiment, the compound of the present invention is useful as a prophylactic or therapeutic agent for narcolepsy and excessive daytime sleepiness. In another embodiment, the compound of the present invention is useful as a prophylactic or therapeutic agent for narcolepsy, cataplexy, and excessive daytime sleepiness. In another embodiment, the compound of the present invention is useful as a prophylactic or therapeutic agent for narcolepsy and cataplexy. In another embodiment, the compound of the present
20 invention is useful as a prophylactic or therapeutic agent for excessive daytime sleepiness. In another embodiment, the compound of the present invention is useful as a prophylactic or therapeutic agent for idiopathic hypersomnia. In another embodiment, the compound of the present invention is useful as a prophylactic or therapeutic agent for obstructive sleep apnea.

25 In another embodiment, the compound of the present invention has orexin-2 receptor agonist activity and is useful as a prophylactic or therapeutic agent for hypersomnia in Parkinson's disease.

In another embodiment, the compound of the present invention has orexin-2 receptor agonist activity and is useful as a prophylactic or therapeutic agent for hypersomnia. In
30 another embodiment, the compound of the present invention has orexin-2 receptor agonist activity and is useful as a prophylactic or therapeutic agent for excessive daytime sleepiness associated with Parkinson's disease.

In another embodiment, the compound of the present invention has orexin-2 receptor agonist activity, and is useful as a prophylactic or therapeutic agent for excessive daytime sleepiness or fatigue associated with cancer and/or chemotherapy.

5 In another embodiment, the present invention provides a method of treating narcolepsy in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating narcolepsy type-1 in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

10 In another embodiment, the present invention provides a method of treating narcolepsy type-2 in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating narcolepsy and excessive daytime sleepiness in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

15 In another embodiment, the present invention provides a method of treating narcolepsy, cataplexy, and excessive daytime sleepiness in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating narcolepsy and cataplexy in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

20 In another embodiment, the present invention provides a method of treating excessive daytime sleepiness in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating idiopathic hypersomnia in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

30 In another embodiment, the present invention provides a method of treating excessive daytime sleepiness and idiopathic hypersomnia in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating obstructive sleep apnea in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

5 In another embodiment, the present invention provides a method of treating excessive daytime sleepiness and obstructive sleep apnea in a subject in need thereof comprising administering to the subject a compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof.

10 In any of the methods as described herein, the subject is administered a compound of Formula I. In any of the methods as described herein, the subject is administered a compound of Formula I-A.

Each of the embodiments described herein with respect to the use of compounds of Formula I also applies to compounds of Formula I-A.

15 In any of the compositions or methods as described herein, the compound of Formula I-A, or I, or a pharmaceutically acceptable salt thereof, is present and/or administered in a therapeutically effective amount.

Administration / Dosage / Formulations

In another aspect, provided herein is a pharmaceutical composition comprising at least one compound of the invention, together with a pharmaceutically acceptable carrier.

20 Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

25 In particular, the selected dosage level will depend upon a variety of factors including the activity of the particular compound employed, the time of administration, the rate of excretion of the compound, the duration of the treatment, other drugs, compounds or materials used in combination with the compound, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

30 A medical doctor, *e.g.*, physician or veterinarian, having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could begin administration of the pharmaceutical composition to dose the disclosed compound at levels lower than that required

in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

In particular embodiments, it is especially advantageous to formulate the compound in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the patients to be treated; each unit containing a predetermined quantity of the disclosed compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical vehicle. The dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the disclosed compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding/formulating such a disclosed compound for the treatment of narcolepsy or cataplexy in a patient.

In one embodiment, the compounds of the invention are formulated using one or more pharmaceutically acceptable excipients or carriers. In one embodiment, the pharmaceutical compositions of the invention comprise a therapeutically effective amount of a disclosed compound and a pharmaceutically acceptable carrier.

In some embodiments, the dose of a disclosed compound is from about 1 mg to about 1,000 mg. In some embodiments, a dose of a disclosed compound used in compositions described herein is less than about 1,000 mg, or less than about 800 mg, or less than about 600 mg, or less than about 500 mg, or less than about 300 mg, or less than about 200 mg, or less than about 100 mg, or less than about 50 mg, or less than about 20 mg, or less than about 10 mg. For example, a dose is about 10 mg, 20 mg, 25 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70 mg, 80 mg, 90 mg, 100 mg, 120 mg, 140 mg, 160 mg, 180 mg, 200 mg, 220 mg, 240, 260 mg, 280 mg, 300 mg, 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, or about 600 mg.

Routes of administration of any of the compositions of the invention include oral, nasal, rectal, intravaginal, parenteral, buccal, sublingual or topical. The compounds for use in the invention may be formulated for administration by any suitable route, such as for oral or parenteral, for example, transdermal, transmucosal (e.g., sublingual, lingual, (trans)buccal, (trans)urethral, vaginal (e.g., trans- and perivaginally), (intra)nasal and (trans)rectal), intravesical, intrapulmonary, intraduodenal, intragastrical, intrathecal, subcutaneous, intramuscular, intradermal, intra-arterial, intravenous, intrabronchial, inhalation, and topical administration. In one embodiment, the preferred route of administration is oral.

Suitable compositions and dosage forms include, for example, tablets, capsules, caplets, pills, gel caps, troches, dispersions, suspensions, solutions, syrups, granules, beads, transdermal patches, gels, powders, pellets, magmas, lozenges, creams, pastes, plasters,

lotions, discs, suppositories, liquid sprays for nasal or oral administration, dry powder or aerosolized formulations for inhalation, compositions and formulations for intravesical administration and the like. It should be understood that the formulations and compositions that would be useful in the present invention are not limited to the particular formulations and compositions that are described herein.

For oral application, particularly suitable are tablets, dragees, liquids, drops, suppositories, or capsules, caplets and gelcaps. The compositions intended for oral use may be prepared according to any method known in the art and such compositions may contain one or more agents selected from the group consisting of inert, non-toxic pharmaceutically excipients that are suitable for the manufacture of tablets. Such excipients include, for example an inert diluent such as lactose; granulating and disintegrating agents such as cornstarch; binding agents such as starch; and lubricating agents such as magnesium stearate. The tablets may be uncoated or they may be coated by known techniques for elegance or to delay the release of the active ingredients. Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert diluent.

For parenteral administration, the disclosed compounds may be formulated for injection or infusion, for example, intravenous, intramuscular or subcutaneous injection or infusion, or for administration in a bolus dose or continuous infusion. Suspensions, solutions or emulsions in an oily or aqueous vehicle, optionally containing other formulatory agents such as suspending, stabilizing or dispersing agents may be used.

Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures, embodiments, claims, and examples described herein. Such equivalents are considered to be within the scope of this invention and covered by the claims appended hereto. For example, it should be understood, that modifications in reaction conditions, including but not limited to reaction times, reaction size/volume, and experimental reagents, such as solvents, catalysts, pressures, atmospheric conditions, *e.g.*, nitrogen atmosphere, and reducing/oxidizing agents, with art-recognized alternatives and using no more than routine experimentation, are within the scope of the present application.

It is to be understood that wherever values and ranges are provided herein, all values and ranges encompassed by these values and ranges, are meant to be encompassed within the scope of the present invention. Moreover, all values that fall within these ranges, as well as the upper or lower limits of a range of values, are also contemplated by the present application.

The following examples further illustrate aspects of the present invention. However, they are in no way a limitation of the teachings or disclosure of the present invention as set forth herein.

Examples

5 The invention is further illustrated by the following examples, which should not be construed as further limiting. The practice of the present invention will employ, unless otherwise indicated, conventional techniques of organic synthesis, cell biology, cell culture, molecular biology, transgenic biology, microbiology and immunology, which are within the skill of the art.

10

General Procedures

Example 1: Synthesis Procedures

 Synthesis procedures for preparation of the compounds of the invention are readily available to the ordinary skilled artisan. Unless otherwise indicated, starting materials were
15 generally obtained from commercial sources. Synthetic procedures for related compounds can be found, for example, in U.S. Application No.: 17/556,295, filed December 20, 2021 and in PCT Application No.: PCT/US21/64484, filed December 21, 2021; both of which are expressly incorporated by reference herein in their entireties.

 The following abbreviations may be used in the synthetic examples below:

20 DCM = dichloromethane

MeOH = methanol

THF = tetrahydrofuran

 PPh₃ = triphenylphosphine

DIAD = diisopropyl azodicarboxylate

25 DME = 1,2-dimethoxyethane

*t*BuXphos = 2-di-*tert*-butylphosphino-2',4',6'-triisopropylbiphenyl

EA = ethyl acetate

 HATU = 1-[bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide
 hexafluorophosphate30 DIEA = *N,N*-diisopropylethylamine

NMP = 1-methyl-2-pyrrolidinone

EDCI = 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide

HOBt = 1-hydroxybenzotriazole

STAB = sodium triacetoxyborohydride

BH₃.THF = borane tetrahydrofuran

ACN = acetonitrile

PE = petroleum ether

EtOAc = ethyl acetate

5 DMF = dimethyl formamide

LiOH = lithium hydroxide

min = minutes

h = hours

NaH = sodium hydride

10 Pd₂(dba)₃ = tris(dibenzylideneacetone)dipalladium(0)

DMSO = dimethyl sulfoxide

Boc = *tert*-butyloxycarbonyl

Ms = methanesulfonyl


Bn = benzyl


15 Et = ethyl

Cbz = carboxybenzyl

TEA = triethylamine.


Scheme 1:



Intermediates


To a solution of 4-bromopyridin-3-olate (1 g, 5.747 mmol, 1.00 eq), (cis-4-phenylcyclohexyl)methanol (1.42 g, 7.462 mmol, 1.3 eq), PPh₃ (3.01 g, 11.476 mmol, 2 eq) in THF (2 mL) was added DIAD (2.28 mL, 11.501 mmol, 2 eq) at 0 degrees C. The resulting solution was stirred for 2 h at 70 degrees C. The resulting solution was concentrated and then purified by silica gel chromatography (0-30 % ethyl acetate/petroleum ether) to afford the product 4-bromo-3-[(cis-4-phenylcyclohexyl)methoxy]pyridine as a solid (1 g, 50 % yield).

10 LCMS: m/z (ES+), [M+H]⁺ = 346




To a solution of 4-bromo-3-[(cis-4-phenylcyclohexyl)methoxy]pyridine (1 g, 2.89 mmol, 1 eq) in DME (20 mL) was added methanesulfonamide (1.38 g, 14.508 mmol, 5 eq), Cs₂CO₃ (1.41 g, 4.328 mmol, 1.5 eq), Pd₂(dba)₃ (264.45 mg, 0.289 mmol, 0.1 eq) and *t*-BuXphos (245.3 mg, 0.578 mmol, 0.2 eq) under N₂ atmosphere. The resulting solution was stirred overnight at 80 degrees C. The resulting mixture was concentrated and purified by silica gel chromatography (0-20 % MeOH/DCM) to afford the product N-(3-[(cis-4-phenylcyclohexyl)methoxy]pyridin-4-yl)methanesulfonamide as a solid (977 mg, 94 % yield).

20 LCMS: m/z (ES+), [M+H]⁺ = 361




To a solution of N-(3-[[cis-4-phenylcyclohexyl]methoxy]pyridin-4-yl)methanesulfonamide (562 mg, 1.559 mmol, 1 eq) in ACN (10 mL) was added benzyl bromide (0.19 mL, 1.597 mmol, 1.02 eq) at room temperature. The resulting solution was stirred overnight at 80 degrees C. The mixture was concentrated and purified by reverse-phase chromatography to afford 1-benzyl-4-methanesulfonamido-3-[[cis-4-phenylcyclohexyl]methoxy]pyridin-1-ium-1-yl)-lambda2-bromanide (474 mg, 67.3 % yield) as a solid.

LCMS: m/z (ES+), [M]⁺ = 451



To a solution of (1-benzyl-4-methanesulfonamido-3-[[cis-4-phenylcyclohexyl]methoxy]-lambda4-pyridin-1-ium-1-yl)-lambda2-bromanide (474 mg, 0.892 mmol, 1 eq) in THF (40 mL) was added LiAlH₄ (270.8 mg, 7.134 mmol, 8 eq) at 0 degrees C. The resulting solution was stirred for 1 h at 70 degrees C. The reaction was quenched by MeOH (2 mL) at 0 degrees C, and then filtered. The filtrate was extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated to afford N-[1-benzyl-3-[[cis-4-phenylcyclohexyl]methoxy]-2H-pyridin-4-yl]methanesulfonamide as an oil (500 mg, crude).

LCMS: m/z (ES+), [M+1]⁺ = 453




A mixture of (N-[1-benzyl-3-[[cis-4-phenylcyclohexyl]methoxy]-4H-pyridin-4-yl]methanesulfonamide (500 mg, 1.10 mmol, 1.00 eq) and Pd(OH)₂/C (499.5 mg) in EA (20 mL) was stirred at room temperature for 20 h under an atmosphere of H₂. The mixture was filtered and the filtrate was concentrated and purified by reverse-phase chromatography to

afford N-[cis-3-[(cis-4-phenylcyclohexyl)methoxy]piperidin-4-yl]methanesulfonamide (200 mg, 49.4%) as an oil.

LCMS: m/z (ES+), [M+1]⁺ = 367

5 Compounds of the Invention

Compound 37




Piperidine ring stereochemistry - cis, racemic

Method 1: To a solution of N-[cis-3-[(cis-4-phenylcyclohexyl)methoxy]piperidin-4-yl]methanesulfonamide (112 mg, 0.306 mmol, 1 eq) in DMF (4 mL) was added
 10 cyclopropanecarboxylic acid (39.46 mg, 0.458 mmol, 1.5 eq), DIEA (78.99 mg, 0.611 mmol, 2 eq) and HATU (174.28 mg, 0.458 mmol, 1.5 eq) at room temperature. The resulting solution was stirred for 2 h at room temperature. After workup the crude material was purified by Prep-HPLC to afford N-[cis-1-cyclopropanecarbonyl-3-[(cis-4-phenylcyclohexyl)methoxy]piperidin-4-yl]methanesulfonamide (86 mg, 65% yield) as a
 15 mixture of isomers as a solid. This isomeric mixture we purified by preparative chiral HPLC to afford compound 37 (36.5 mg, 42 % yield) (absolute configuration not confirmed) as a solid.

LCMS: m/z (ES+), [M+1]⁺ = 435

¹H NMR (400 MHz, Methanol-*d*₄) δ 7.30 – 7.22 (m, 4H), 7.14 – 7.12 (m, 1H), 4.79 – 4.20 (m, 2H), 3.84 – 3.80 (m, 1H), 3.66 – 3.64 (m, 3H), 3.47 – 3.35 (m, 1H), 3.02 (s, 3H), 2.92 – 2.90 (m, 1H), 2.70 – 2.51 (m, 1H), 2.16 – 1.75 (m, 6H), 1.74 – 1.53 (m, 6H), 0.87 – 0.75 (m, 4H).

Compound 38



N-(cis-3-((cis-4-phenylcyclohexyl)methoxy)piperidin-4-yl)methanesulfonamide (100.00 mg, 0.273 mmol, 1.00 eq), TEA (82.82 mg, 0.819 mmol, 3 eq) and 1-chloro-2-methyl-1-oxopropan-2-yl acetate (89.81 mg, 0.546 mmol, 2 eq) in DCM (2.00 mL) were added into an 8 mL vial at 0 degrees C. The resulting mixture was stirred for 1 h at room temperature under argon atmosphere. The reaction was quenched with water. The resulting mixture was extracted with DCM (2 x 5 mL). The combined organic layers were washed with brine (5 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure to afford 1-(4-methanesulfonamido-3-[[[(1s,4s)-4-phenylcyclohexyl]methoxy]piperidin-1-yl]-2-methyl-1-oxopropan-2-yl acetate (130 mg, 96.3%) as a foam.


LCMS: m/z (ES+), [M+H]⁺ = 495

Into a 50 mL round-bottom flask were added 2-methyl-1-(cis-4-(methylsulfonamido)-3-((cis-4-phenylcyclohexyl)methoxy)piperidin-1-yl)-1-oxopropan-2-yl acetate (130.00 mg, 0.263 mmol, 1.00 eq) and LiOH.H₂O (55.14 mg, 1.314 mmol, 5.00 eq) in THF (3.00 mL) and H₂O (1.00 mL) at room temperature. The resulting mixture was stirred overnight at room temperature under argon atmosphere. The reaction was quenched with water. The resulting mixture was extracted with EtOAc (2 x 5 mL). The combined organic layers were washed with brine (5 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The crude product (130 mg) was purified by Prep-HPLC to afford crude product (70 mg, 59%) as a solid. This crude product (70 mg) was purified by CHIRAL-Prep-HPLC to afford compound 38 (17.1 mg, 24%,) as a solid.

LCMS: m/z (ES+), [M+H]⁺ = 453

¹H NMR (400 MHz, DMSO-*d*₆) δ 7.27 (d, *J* = 4.5 Hz, 4H), 7.16 (h, *J* = 4.1 Hz, 1H), 6.87 (s, 1H), 5.34 (s, 1H), 4.58 (s, 2H), 3.64 (d, *J* = 103.3 Hz, 4H), 3.26 (s, 1H), 2.95 (s, 3H), 2.80 (d, *J* = 18.4 Hz, 1H), 2.52 (s, 1H), 1.97 (s, 1H), 1.84 – 1.64 (m, 4H), 1.56 (s, 4H), 1.48 (s, 2H), 1.29 (d, *J* = 9.6 Hz, 6H).

Compound 39




Into a 8 mL vial were added N-(cis-3-((cis-4-phenylcyclohexyl)methoxy)piperidin-4-yl)methanesulfonamide (90.0 mg, 0.246 mmol, 1.00 eq), DIEA (95.21 mg, 0.737 mmol, 3 eq) and 2-chloro-1,3-oxazole (50.83 mg, 0.491 mmol, 2.00 eq) in NMP (2.00 mL) at room temperature. The resulting mixture was stirred overnight at 120 degrees C under argon atmosphere. The reaction was quenched with water. The resulting mixture was extracted with EtOAc (2 x 5 mL). The combined organic layers were washed with brine (5 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The crude product (120 mg) was purified by Prep-HPLC to afford N-(cis-1-(oxazol-2-yl)-3-((cis-4-phenylcyclohexyl)methoxy)piperidin-4-yl)methanesulfonamide (60 mg, 56.3%) as a solid. Above product (60 mg) was purified by CHIRAL-Prep-HPLC to afford compound 39 (10.6 mg, 18%) as a solid.

LCMS: m/z (ES⁺), [M+H]⁺ = 434

¹H NMR (400 MHz, DMSO-*d*₆) δ 7.46 (d, *J* = 1.0 Hz, 1H), 7.28 (dd, *J* = 8.0, 6.9 Hz, 2H), 7.25 – 7.20 (m, 2H), 7.19 – 7.13 (m, 1H), 6.93 (d, *J* = 7.6 Hz, 1H), 6.78 (d, *J* = 1.0 Hz, 1H), 4.12 – 4.00 (m, 1H), 3.83 (dd, *J* = 13.3, 5.0 Hz, 1H), 3.68 – 3.50 (m, 3H), 3.45 (dd, *J* = 9.0, 7.0 Hz, 1H), 3.21 – 3.07 (m, 2H), 2.96 (s, 3H), 2.52 (s, 1H), 1.91 – 1.77 (m, 2H), 1.71 (d, *J* = 11.0 Hz, 1H), 1.67 – 1.54 (m, 5H), 1.53 – 1.38 (m, 3H).

20

Compound 40 and 41



Method 2: To a solution of N-(cis-3-(((1s,4S)-4-phenylcyclohexyl)methoxy)piperidin-4-yl)methanesulfonamide (200 mg, 0.546 mmol, 1 eq) and TEA (165.65 mg, 1.637 mmol, 3 eq) in DCM (10 mL) was added methyl chloroformate (103.12 mg, 1.091 mmol, 2 eq) at 5 degrees C. The resulting solution was stirred for 2.5 h at room temperature. The reaction was quenched with water (20 mL). The resulting mixture was extracted with dichloromethane (3 x 10 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and concentrated. The residue obtained was purified by Prep-HPLC to afford methyl-cis-4-

10 (methanesulfonamido)-3-(((1s,4S)-4-phenylcyclohexyl)methoxy)piperidine-1-carboxylate (102 mg, 44 % yield) as a solid. The above material was purified by preparative chiral HPLC to afford compound 40 (36.0 mg (35 % yield) as a solid and compound 41 (35.6 mg, 34 % yield) as a solid.

Compound 40:

15 LCMS: m/z (ES⁺), $[M+NH_4]^+ = 442$


¹H NMR (400 MHz, Methanol-*d*₄) δ 7.30 – 7.27 (m, 4H), 7.19 – 7.09 (m, 1H), 4.41 – 4.34 (m, 1H), 4.10 – 4.05 (m, 1H), 3.90 – 3.81 (m, 1H), 3.72 – 3.62 (m, 5H), 3.52 – 3.46 (m, 1H), 3.00 (s, 5H), 2.60 (s, 1H), 2.05 (s, 1H), 1.92 – 1.87 (m, 4H), 1.74 – 1.61 (m, 6H).

Compound 41:

20 LCMS: m/z (ES⁺), $[M+NH_4]^+ = 442$

¹H NMR (400 MHz, Methanol-*d*₄) δ 7.30 – 7.27 (m, 4H), 7.19 – 7.09 (m, 1H), 4.41 – 4.34 (m, 1H), 4.10 – 4.05 (m, 1H), 3.90 – 3.81 (m, 1H), 3.72 – 3.62 (m, 5H), 3.52 – 3.46 (m, 1H), 3.00 (s, 5H), 2.60 (s, 1H), 2.05 (s, 1H), 1.92 – 1.87 (m, 4H), 1.74 – 1.61 (m, 6H).

Compound 42




Method 3: A mixture of N-[(3R,4S)-3-[[[(1S,4S)-4-phenylcyclohexyl]methoxy]piperidin-4-yl]methanesulfonamide (100.00 mg, 0.273 mmol, 1.00 eq), EDCI (78.45 mg, 0.409 mmol, 1.5 eq), HOBt (55.30 mg, 0.409 mmol, 1.5 eq), DIEA (105.79 mg, 0.819 mmol, 3 eq) and 1-methoxycyclobutane-1-carboxylic acid (53.26 mg, 0.409 mmol, 1.5 eq) in DCM (1.00 mL) was stirred for 3 h at room temperature. The residue was purified by Prep-TLC (PE/EtOAc 2:1) to afford crude product (100mg, 76.57%) as a solid. This crude product was purified by chiral HPLC to afford compound 42 (43mg) as a solid.

LCMS: m/z (ES⁺), $[M+H]^+ = 479.25$

¹H NMR (400 MHz, Methanol-d₄) δ 7.26 (dd, $J = 5.5, 3.9$ Hz, 4H), 7.15 (tq, $J = 5.0, 3.3, 2.9$ Hz, 1H), 4.83 (dt, $J = 14.1, 3.0$ Hz, 1H), 4.19 (dp, $J = 13.8, 2.8$ Hz, 1H), 3.89 (t, $J = 8.6$ Hz, 1H), 3.79 – 3.58 (m, 2H), 3.44 (dd, $J = 8.8, 6.8$ Hz, 1H), 3.16 (s, 2H), 3.21 – 3.06 (m, 2H), 3.01 (d, $J = 4.5$ Hz, 3H), 2.84 (dd, $J = 14.0, 1.4$ Hz, 1H), 2.63 (tt, $J = 12.1, 4.0$ Hz, 1H), 2.61 – 2.46 (m, 1H), 2.33 – 2.16 (m, 1H), 2.19 – 2.07 (m, 1H), 2.04 (dt, $J = 8.3, 4.4$ Hz, 1H), 1.98 – 1.84 (m, 1H), 1.85 (s, 2H), 1.82 (ddt, $J = 12.9, 7.8, 4.1$ Hz, 1H), 1.80 – 1.66 (m, 2H), 1.70 – 1.55 (m, 2H), 1.61 (s, 2H), 1.37 – 1.28 (m, 1H).

20 Compound 43 and 44



Compound 43 and compound 44 were synthesized using method 3.

Compound 43: (12.6 mg) as a solid.

LCMS72: m/z (ES+), $[M+H]^+ = 465.2$

^1H NMR (400 MHz, Methanol- d_4) δ 7.32 – 7.22 (m, 5H), 7.19 – 7.10 (m, 2H), 4.70 (d, $J =$
 5 14.0 Hz, 1H), 4.16 (d, $J = 14.1$ Hz, 1H), 3.90 (t, $J = 8.7$ Hz, 1H), 3.86 – 3.73 (m, 1H), 3.67 (s,
 2H), 3.42 (dd, $J = 8.7, 6.5$ Hz, 1H), 3.21 (d, $J = 14.4$ Hz, 1H), 3.02 (s, 1H), 3.01 (s, 4H), 2.90
 (d, $J = 13.8$ Hz, 1H), 2.74 (s, 2H), 2.59 (d, $J = 11.2$ Hz, 3H), 2.12 (dq, $J = 39.6, 9.7$ Hz, 4H),
 1.98 – 1.81 (m, 2H), 1.84 – 1.77 (m, 1H), 1.70 (s, 9H), 1.68 (s, 1H), 1.62 (s, 2H), 1.54 (dt, $J =$
 17.0, 8.7 Hz, 1H), 1.38 – 1.29 (m, 1H).


10

Compound 44: (11mg) as a solid.

LCMS: m/z (ES+), $[M+H]^+ = 465.2$

^1H NMR (400 MHz, Methanol- d_4) δ 7.32 – 7.22 (m, 5H), 7.14 (s, 1H), 4.70 (d, $J = 14.1$ Hz,
 1H), 4.16 (d, $J = 14.0$ Hz, 1H), 3.90 (t, $J = 8.7$ Hz, 1H), 3.76 (d, $J = 7.6$ Hz, 1H), 3.65 (d, $J =$
 15 16.6 Hz, 3H), 3.42 (dd, $J = 8.7, 6.4$ Hz, 1H), 3.19 (s, 1H), 3.01 (s, 3H), 2.90 (d, $J = 14.0$ Hz,
 1H), 2.74 (s, 2H), 2.59 (d, $J = 10.2$ Hz, 2H), 2.16 (dt, $J = 19.5, 9.9$ Hz, 1H), 2.07 (q, $J = 9.8,$
 8.6 Hz, 2H), 1.98 – 1.77 (m, 3H), 1.69 (s, 8H), 1.68 (s, 1H), 1.64 – 1.49 (m, 1H), 1.32 (q, $J =$
 7.7, 7.0 Hz, 1H), 0.12 (s, 1H).

20 *Compound 45 and 46*



Compound 45 and compound 46 were synthesized using method 3.

Compound 45: (90 mg, 71%) as a solid.

^1H NMR (400 MHz, Methanol- d_4) δ 7.27 (d, $J = 5.6$ Hz, 4H), 7.14 (s, 1H), 4.93 (d, $J = 5.9$
 25 Hz, 2H), 4.81 (d, $J = 14.5$ Hz, 1H), 4.34 (dd, $J = 5.9, 18.7$ Hz, 2H), 3.91 (t, $J = 8.8$ Hz, 1H),
 3.71 - 3.61 (m, 2H), 3.42 - 3.36 (m, 1H), 3.20 - 3.09 (m, 2H), 3.02 (d, $J = 4.8$ Hz, 3H), 2.82

(d, $J = 14.2$ Hz, 1H), 2.58 (s, 1H), 2.08 (s, 1H), 1.98 - 1.85 (m, 2H), 1.77 (d, $J = 13.5$ Hz, 3H), 1.72 - 1.60 (m, 8H).


LCMS: m/z (ES+), $[M+H]^+ = 465.2$

5 Compound 46: (10 mg, 7%) as a solid.

$^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 7.27 (d, $J = 4.5$ Hz, 4H), 7.16 (t, $J = 4.1$ Hz, 1H), 6.95 (dd, $J = 7.2, 19.8$ Hz, 1H), 4.76 (t, $J = 5.5$ Hz, 2H), 4.59 (d, $J = 13.8$ Hz, 1H), 4.24 (dd, $J = 5.8, 9.7$ Hz, 2H), 3.76 (t, $J = 9.0$ Hz, 1H), 3.55 (s, 2H), 3.28 (dd, $J = 5.9, 8.6$ Hz, 1H), 3.07 (t, $J = 12.8$ Hz, 1H), 2.95 (d, $J = 6.0$ Hz, 4H), 2.76 (d, $J = 13.9$ Hz, 1H), 1.99 (s, 1H), 1.79 (t, $J = 11.4$ Hz, 1H), 1.70 (t, $J = 13.8$ Hz, 3H), 1.64 - 1.54 (m, 5H), 1.50 (s, 5H).

LCMS: m/z (ES+), $[M+H]^+ = 465.2$

Compound 47 and 48



- 15 **Method 4:** To a stirred mixture of NaH (39.72 mg, 0.99 mmol, 2.80 eq, 60%) and ionic (218.84 mg, 0.993 mmol, 2.80 eq) was added THF (2.00 ml) dropwise at 0 degrees C under nitrogen atmosphere. After stirring for about 5 minutes, the ice bath was removed, and the solution was allowed to warm to room temperature over 10 minutes. An aliquot of this solution (volume = 1.2 mL, 0.60 mmol, 1.5 eq) was transferred into a dry 7 mL vial. The
- 20 resulting mixture was concentrated under reduced pressure. To the above mixture was added Pd-PEPPSI-IPentCl *o*-picoline (29.80 mg, 0.035 mmol, 0.10 eq), N-[(3R,4S)-3-[[[(1s,4s)-4-phenylcyclohexyl]methoxy]piperidin-4-yl]methanesulfonamide (130.00 mg, 0.355 mmol, 1.00 eq) and 3-bromo-1-methylpyrazole (57.11 mg, 0.355 mmol, 1.00 eq) in 1,4-dioxane (10 mL) was stirred overnight at 120 degrees
- 25 C under nitrogen atmosphere. After starting material was consumed, the residue was purified

by Prep-TLC (CH₂Cl₂ /MeOH 10 :1) to afford crude product (120mg) as a solid.

The crude product (120 mg) was purified by Prep-chiral-HPLC to afford compound 47 and compound 48.

Compound 47: (13.8 mg, 18.65%) as a solid.

5 LCMS: m/z (ES+), [M+H]⁺ = 447

¹H NMR (400 MHz, Methanol-d₄) δ 7.32 (d, J = 2.2 Hz, 1H), 7.26 (t, J = 7.4 Hz, 2H), 7.22 – 7.17 (m, 2H), 7.14 (t, J = 7.2 Hz, 1H), 5.72 (t, J = 1.9 Hz, 1H), 3.76 (q, J = 7.9, 7.2 Hz, 2H), 3.69 – 3.66 (m, 5H), 3.62 – 3.47 (m, 2H), 3.06 (m, 1H), 3.02– 2.90 (m, 5H), 2.61 – 2.52 (m, 1H), 2.08 (ddd, J = 23.4, 9.8, 3.9 Hz, 2H), 1.83 (dp, J = 17.3, 5.9, 4.1 Hz, 3H), 1.75 – 1.55 (m, 6H), 1.32 (t, J = 7.3 Hz, 3H).

10


Compound 48: (12.3mg, 16.62%) as a solid.

LCMS: m/z (ES+), [M+H]⁺ = 447.10

¹H NMR (400 MHz, Methanol-d₄) δ 7.32 (d, J = 2.2 Hz, 1H), 7.26 (t, J = 7.4 Hz, 2H), 7.22 – 7.17 (m, 2H), 7.14 (t, J = 7.2 Hz, 1H), 5.72 (t, J = 1.9 Hz, 1H), 3.76 (q, J = 7.9, 7.2 Hz, 2H), 3.69 – 3.66 (m, 5H), 3.62 – 3.47 (m, 2H), 3.06 (m, 1H), 3.02– 2.90 (m, 5H), 2.61 – 2.52 (m, 1H), 2.08 (ddd, J = 23.4, 9.8, 3.9 Hz, 2H), 1.83 (dp, J = 17.3, 5.9, 4.1 Hz, 3H), 1.75 – 1.55 (m, 6H), 1.32 (t, J = 7.3 Hz, 3H).

15

20 *Compound 49 and 50*



Compound 49 and compound 50 were synthesized using method 3.

Compound 49: (93.1 mg, 33.2%) as a solid.

25 LCMS: m/z (ES+), [M+H]⁺ = 467.0

¹H NMR (400 MHz, Methanol-d₄) δ 7.30 – 7.20 (m, 4H), 7.17 – 7.12 (m, 1H), 4.79 (d, 1H), 3.92 (d, 2H), 3.77 – 3.61 (m, 2H), 3.40 – 3.34 (m, 1H), 3.24 – 3.14 (m, 1H), 3.01 (d, 3H),

2.89 (d, 1H), 2.70 (d, 2H), 2.56 (s, 1H), 2.41 (dp, 2H), 1.92 (d, 3H), 1.81 – 1.71 (m, 3H), 1.67 (d, 3H), 1.63 – 1.49 (m, 3H).


Compound 50: (75.5 mg, 26.96%) as a solid.

5 LCMS: m/z (ES+), $[M+H]^+ = 467.0$

^1H NMR (400 MHz, Methanol- d_4) δ 7.31 – 7.23 (m, 4H), 7.14 (d, 1H), 4.79 (d, 1H), 3.96 – 3.62 (m, 4H), 3.40 – 3.35 (m, 1H), 3.20 (d, 1H), 3.01 (d, 3H), 2.89 (d, 1H), 2.70 (s, 2H), 2.56 (s, 1H), 2.41 (dt, 2H), 2.09 (d, 1H), 1.97 – 1.84 (m, 3H), 1.76 (d, 3H), 1.68 (d, 3H), 1.61 – 1.51 (m, 2H).

10

Compound 51 and 52



To a stirred solution of N-[(3R,4S)-3-[[[(1S,4S)-4-phenylcyclohexyl]methoxy]piperidin-4-yl]methanesulfonamide (200.00 mg, 0.546 mmol, 1.00 eq) in MeOH (5mL) were added

15 ZnCl_2 (0.74 mg, 0.005 mmol, 0.01 eq) and NaBH_3CN (51.44 mg, 0.819 mmol, 1.50 eq) in portions at 0 degrees C under nitrogen atmosphere. The resulting mixture was stirred for 3 h at 50 degrees C under nitrogen atmosphere. The residue was filtered through silica gel column with DCM/MeOH (20:1). The crude product (54mg) was purified by Prep-HPLC followed by chiral HPLC to afford products.

20 Compound 51: (1.5mg, 0.65%) as a solid.

LCMS: m/z (ES+), $[M+H]^+ = 423.20$

^1H -NMR (400 MHz, Methanol- d_4) δ 7.26 (d, $J = 6.0$ Hz, 3H), 7.15 (td, $J = 5.9, 2.9$ Hz, 1H), 4.75 – 4.55 (m, 5H), 3.77 – 3.69 (m, 1H), 3.69 – 3.48 (m, 3H), 3.00 (s, 3H), 2.76 (s, 1H), 2.61 (s, 2H), 2.21 (s, 2H), 2.08 (s, 1H), 2.04 – 1.94 (m, 1H), 1.90 (d, $J = 11.4$ Hz, 2H), 1.82 – 1.64 (m, 6H).


25

Compound 52: (0.6 mg, 0.26%).

LCMS: m/z (ES+), $[M+H]^+ = 423.20$

$^1\text{H-NMR}$ (E05570-026): (400 MHz, Methanol- d_4) δ 7.26 (d, $J = 6.0$ Hz, 3H), 7.15 (td, $J = 5.9, 2.9$ Hz, 1H), 4.75 – 4.55 (m, 5H), 3.77 – 3.69 (m, 1H), 3.69 – 3.48 (m, 3H), 3.00 (s, 3H),
 5 2.76 (s, 1H), 2.61 (s, 2H), 2.21 (s, 2H), 2.08 (s, 1H), 2.04 – 1.94 (m, 1H), 1.90 (d, $J = 11.4$ Hz, 2H), 1.82 – 1.64 (m, 6H).

Compound 53 and 54



10 Compound 53 and compound 54 were synthesized using method 4.

Compound 53: (6.3mg) as a solid.

LCMS: m/z (ES+), $[M+H]^+ = 447.1$

$^1\text{H NMR}$ (400 MHz, Methanol- d_4) δ 7.28 – 7.22 (m, 2H), 7.22 – 7.11 (m, 3H), 6.82 (d, $J = 1.5$ Hz, 1H), 6.71 (d, $J = 1.6$ Hz, 1H), 4.60 (s, 1H), 3.73 (t, $J = 8.7$ Hz, 2H), 3.70 – 3.64 (m,
 15 1H), 3.60 (dd, $J = 9.1, 6.8$ Hz, 1H), 3.53 (s, 3H), 3.46 (dd, $J = 12.6, 5.0$ Hz, 1H), 3.24 (d, $J = 12.3$ Hz, 1H), 3.03 (s, 3H), 2.96 (dd, $J = 20.7, 10.7$ Hz, 2H), 2.58 (s, 1H), 2.13 (ddd, $J = 23.3, 11.4, 7.7$ Hz, 2H), 1.94 – 1.79 (m, 3H), 1.65 (dt, $J = 25.9, 8.0$ Hz, 6H), 1.31 (s, 2H).


Compound 54: (8.8mg, 98%) as a solid.

20 LCMS: m/z (ES+), $[M+H]^+ = 447.3$

$^1\text{H NMR}$ (400 MHz, Methanol- d_4) δ 7.28 – 7.22 (m, 2H), 7.22 – 7.11 (m, 3H), 6.82 (d, $J = 1.5$ Hz, 1H), 6.71 (d, $J = 1.6$ Hz, 1H), 4.60 (s, 1H), 3.73 (t, $J = 8.7$ Hz, 2H), 3.70 – 3.64 (m, 1H), 3.60 (dd, $J = 9.1, 6.8$ Hz, 1H), 3.53 (s, 3H), 3.46 (dd, $J = 12.6, 5.0$ Hz, 1H), 3.24 (d, $J =$

12.3 Hz, 1H), 3.03 (s, 3H), 2.96 (dd, $J = 20.7, 10.7$ Hz, 2H), 2.58 (s, 1H), 2.13 (ddd, $J = 23.3, 11.4, 7.7$ Hz, 2H), 1.94 – 1.79 (m, 3H), 1.65 (dt, $J = 25.9, 8.0$ Hz, 6H), 1.31 (s, 2H).

Compound 55 and 56



5

To the stirred solution of N-(3-(((1s,4s)-4-phenylcyclohexyl)methoxy)piperidin-4-yl)methanesulfonamide (150.00 mg, 0.409 mmol, 1.00 eq) in DMF (10.00 mL) were added 3-chloropyridazine (70.31 mg, 0.614 mmol, 1.50 eq) and CsF (124.33 mg, 0.819 mmol, 2.00 eq). The mixture was stirred for 48 hours at 120 degrees C. The resulting mixture was diluted with water (20 mL). The aqueous layer was extracted with EtOAc (3x20 mL). The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (CH₂Cl₂/MeOH 20:1) to afford product (50mg, 27.48%) as a mixture of isomers. This mixture was purified by chiral HPLC to afford:

Compound 55: (6.3 mg, 90.00%) as a solid.

15 LCMS: m/z (ES+), $[M+H]^+ = 445.1$

¹H NMR(E05616-059): (400 MHz, Methanol-d₄) δ 8.38 (dd, $J = 4.2, 1.5$ Hz, 1H), 7.36 – 7.29 (m, 2H), 7.29 – 7.23 (m, 2H), 7.22 – 7.11 (m, 3H), 4.68 (d, $J = 14.4$ Hz, 1H), 4.34 (d, $J = 13.8$ Hz, 1H), 3.76 – 3.68 (m, 3H), 3.54 – 3.46 (m, 1H), 3.26 – 3.14 (m, 2H), 3.03 (s, 3H), 2.51 (s, 1H), 2.01 (dd, $J = 13.1, 9.1$ Hz, 1H), 1.89 (s, 1H), 1.85 – 1.78 (m, 1H), 1.74 (d, $J = 11.7$ Hz, 1H), 1.69 – 1.56 (m, 4H), 1.49 (d, $J = 6.1$ Hz, 3H), 1.31 (s, 1H).

20

Compound 56: (3 mg) as a solid.


LCMS: m/z (ES+), $[M+H]^+ = 445.1$

1H NMR(E05616-059): (400 MHz, Methanol-d₄) δ 8.38 (dd, $J = 4.2, 1.5$ Hz, 1H), 7.36 – 7.29 (m, 2H), 7.29 – 7.23 (m, 2H), 7.22 – 7.11 (m, 3H), 4.68 (d, $J = 14.4$ Hz, 1H), 4.34 (d, $J = 13.8$ Hz, 1H), 3.76 – 3.68 (m, 3H), 3.54 – 3.46 (m, 1H), 3.26 – 3.14 (m, 2H), 3.03 (s, 3H),

25

2.51 (s, 1H), 2.01 (dd, $J = 13.1, 9.1$ Hz, 1H), 1.89 (s, 1H), 1.85 – 1.78 (m, 1H), 1.74 (d, $J = 11.7$ Hz, 1H), 1.69 – 1.56 (m, 4H), 1.49 (d, $J = 6.1$ Hz, 3H), 1.31 (s, 1H).

Compound 57 and 58



5

To a stirred solution of N-(3-[[[(1s,4s)-4-phenylcyclohexyl]methoxy]piperidin-4-yl)methanesulfonamide (100.0 mg, 0.273 mmol, 1.00 eq) and oxetane-3-carbaldehyde (35.2 mg, 0.409 mmol, 1.50 eq) in MeOH (6.00 mL) was added STAB (86.7 mg, 0.409 mmol, 1.50 eq) in portions at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 2 h at room temperature under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 20:1) to afford product (60 mg, 50.37%) as a mixture of isomers as a solid. This mixture was purified by chiral HPLC to afford:

10

Compound 57: (25 mg, 42 %) as an oil.

15

LCMS: m/z (ES⁺), $[\text{M}+\text{H}]^+ = 437.25$

¹H NMR (400 MHz, Methanol-*d*⁴) δ 7.34-7.20 (m, 4H), 7.19-7.08 (m, 1H), 4.82-4.77 (m, 2H), 4.43 (t, $J = 6.2$ Hz, 2H), 3.75-3.50 (m, 4H), 3.30-3.24 (m, 1H), 2.99 (s, 3H), 2.75 (d, $J = 4.1$ Hz, 3H), 2.69-2.56 (m, 2H), 2.29 (d, $J = 9.5$ Hz, 2H), 2.11-1.92 (m, 2H), 1.90-1.80 (m, 2H), 1.70 (m, 7H).

20


Compound 58: (20 mg, 33%) as an oil.

LCMS: m/z (ES⁺), $[\text{M}+\text{H}]^+ = 437.20$

¹H NMR (400 MHz, Methanol-*d*⁴) δ 7.32-7.21 (m, 4H), 7.15 (t, $J = 2.2$ Hz, 1H), 4.81 (d, $J = 5.8$ Hz, 2H), 4.43 (t, $J = 6.2$ Hz, 2H), 3.75-3.49 (m, 4H), 3.30-3.23 (m, 1H), 2.99 (s, 3H),

2.85-2.72 (m, 3H), 2.62 (d, J = 9.5 Hz, 2H), 2.29 (d, J = 10.0 Hz, 2H), 2.12-2.01 (m, 1H),
2.01-1.91 (m, 1H), 1.91-1.80 (m, 2H), 1.78-1.64 (m, 7H).

Compound 59 and 60



Compound 59 and compound 60 were synthesized using method 4.

Compound 59: (10mg, 23%) as a solid.

LCMS : m/z (ES+), [M+H]⁺ = 447


10 ¹H NMR (400 MHz, Methanol-d₄) δ 7.35 -7.18 (m, 5H), 7.18-7.11 (m, 1H), 6.39 (d, 1H),
3.87-3.79 (m, 1H), 3.71 (dt, 1H), 3.67-3.56 (m, 6H), 3.39-3.35 (m, 1H), 3.02 (s, 3H), 2.93-
2.87 (m, 1H), 2.82 (dd, 1H), 2.61-2.52 (m, 1H), 2.10 (dt, 2H), 1.91-1.79 (m, 3H), 1.69 (dd,
6H).

Compound 60: (10mg, 23%) as a solid.

LCMS: m/z (ES+), [M+H]⁺ = 447

15 ¹H NMR (400 MHz, Methanol-d₄) δ 7.36-7.18 (m, 5H), 7.18-7.09 (m, 1H), 6.39 (d, 1H), 3.79
(dd, 1H), 3.71 (dd, 1H), 3.68-3.57 (m, 5H), 3.37 (s, 1H), 3.02 (s, 3H), 2.92-2.77 (m, 2H), 2.58
(dd, 1H), 2.10 (dt, 2H), 1.94-1.81 (m, 3H), 1.69 (dd, 6H).

Compound 61, 62, 63 and 64



To a stirred solution of N-[(3S,4R)-3-[[[(1S,4S)-4-phenylcyclohexyl]methoxy]piperidin-4-yl]methanesulfonamide (100.0 mg, 0.273 mmol, 1.00 eq) and 1-(oxetan-3-

- 5 yl)ethanone (81.95 mg, 0.819 mmol, 3.00 eq) in MeOH (3 mL) was added ZnCl₂ (37.19 mg, 0.273 mmol, 1.00 eq) and NaBH₃CN (51.44 mg, 0.819 mmol, 3.00 eq) at room temperature. The resulting mixture was stirred for additional overnight at room temperature.

The residue was purified by Prep-TLC (CH₂Cl₂/MeOH 15:1) to afford product as a mixture of isomers (100mg, 81%) as a solid.

- 10 The crude product (96 mg) was purified by Prep-chiral-HPLC to afford:

Compound 61: (30mg, 31.25%) as a solid.

LCMS: m/z (ES⁺), [M+H]⁺ = 451

- ¹H NMR (400 MHz, Methanol-d₄) δ 7.27 (s, 4H), 7.15 (t, 1H), 4.73 (dt, 2H), 4.61 (t, 1H), 4.44 (t, 1H), 3.67-3.51 (m, 4H), 3.25-3.10 (m, 1H), 3.09-2.98 (m, 1H), 2.98 (s, 3H), 2.82 (dd, 15 1H), 2.61 (s, 2H), 2.46 (dd, 1H), 2.35 (d, 1H), 1.94 (dt, 1H), 1.85 (t, 2H), 1.79-1.62 (m, 7H), 0.95 (d, 3H).

Compound 62: (20mg, 21%) as a solid.

LCMS: m/z (ES⁺), [M+H]⁺ = 451

- 20 ¹H NMR (400 MHz, Methanol-d₄) δ 7.35-7.21 (m, 4H), 7.15 (t, 1H), 4.72 (dd, 2H), 4.61 (t, 1H), 4.44 (t, 1H), 3.69 (dd, 1H), 3.62-3.47 (m, 3H), 3.20 (dd, 1H), 3.11-2.93 (m, 4H), 2.81 (s, 1H), 2.65 (dt, 2H), 2.55-2.47 (m, 1H), 2.30 (dd, 1H), 2.05 (d, 1H), 1.88 (dt, 3H), 1.79-1.64 (m, 7H), 1.40.

Compound 63: (12mg, 30%) as a solid.

LCMS: m/z (ES+), [M+H]⁺ = 451.2


¹H NMR(E05616-153): (400 MHz, Methanol-d₄) δ 7.32-7.22 (m, 4H), 7.18-7.12 (m, 1H), 4.73 (ddd, J = 8.1, 6.0, 4.0 Hz, 2H), 4.61 (t, J = 6.4 Hz, 1H), 4.47 (t, J = 6.4 Hz, 1H), 3.71 (t, J = 8.4 Hz, 1H), 3.60 (dd, J = 16.4, 7.4 Hz, 3H), 3.24 (s, 1H), 3.17-3.03 (m, 1H), 2.99 (s, 3H), 2.96-2.69 (m, 2H), 2.61 (dt, J = 11.3, 5.9 Hz, 2H), 2.40 (s, 1H), 2.06 (s, 1H), 1.87 (d, J = 10.2 Hz, 3H), 1.82-1.60 (m, 7H), 1.31 (d, J = 3.5 Hz, 1H), 0.97 (d, J = 41.5 Hz, 3H).

Compound 64: (13mg, 32.5%) as a solid.

10 LCMS: m/z (ES+), [M+H]⁺ = 451.2

¹H NMR(E05616-154): (400 MHz, Methanol-d₄) δ 7.30-7.22 (m, 4H), 7.15 (ddd, J = 8.6, 6.0, 2.2 Hz, 1H), 4.78-4.69 (m, 2H), 4.61 (t, J = 6.4 Hz, 1H), 4.45 (t, J = 6.3 Hz, 1H), 3.72-3.48 (m, 4H), 3.20 (d, J = 8.0 Hz, 1H), 3.06 (q, J = 7.3 Hz, 1H), 2.99 (s, 3H), 2.84 (d, J = 22.3 Hz, 1H), 2.62 (d, J = 8.6 Hz, 2H), 2.47 (d, J = 39.2 Hz, 2H), 2.08-1.91 (m, 2H), 1.84 (td, J = 11.5, 9.5, 5.2 Hz, 2H), 1.71 (p, J = 7.9 Hz, 7H), 1.39-1.27 (m, 1H), 1.05-0.88 (m, 3H).

Compound 65 and 66



Compound 65 and compound 66 were synthesized using method 3.

20 Compound 65: (27.7 mg, 25.18%) as a solid.

LCMS: m/z (ES+), [M+H]⁺ = 449.6


¹H NMR (400 MHz, Methanol-d₄) δ 7.33 – 7.23 (m, 4H), 7.18 – 7.13 (m, 1H), 4.61 (s, 1H), 3.91 (dt, 1H), 3.77 (dt, 1H), 3.70 – 3.59 (m, 2H), 3.50 – 3.35 (m, 2H), 3.19 – 3.10 (m, 1H), 3.01 (d, 3H), 2.81 (d, 1H), 2.58 (s, 1H), 2.38 – 2.31 (m, 1H), 2.26 – 2.04 (m, 3H), 1.97 – 1.60 (m, 11H).

Compound 66: (32.3 mg, 29%) as a solid.

LCMS: m/z (ES+), $[M+H]^+ = 449.6$

^1H NMR (400 MHz, Methanol- d_4) δ 7.32 – 7.25 (m, 4H), 7.17 – 7.12 (m, 1H), 4.77 (d, 1H), 4.60 (s, 1H), 3.91 (dt, 1H), 3.77 (dt, 1H), 3.70 – 3.59 (m, 2H), 3.47 – 3.35 (m, 2H), 3.18 – 3.10 (m, 1H), 3.01 (d, 3H), 2.81 (d, 1H), 2.58 (s, 1H), 2.41 – 2.14 (m, 3H), 2.08 – 1.62 (m, 13H), 1.31 (s, 1H).

Compound 67 and 68



10 Compound 67 and compound 68 were synthesized using method 3.

Compound 67: (22.9 mg, 22.9%) as a solid.

LCMS: m/z (ES+), $[M+H]^+ = 453.5$


^1H NMR (400 MHz, Methanol- d_4) δ 7.26 (d, 4H), 7.15 (h, 1H), 4.73 (d, 1H), 4.45 (d, 1H), 3.87 (s, 1H), 3.74 – 3.64 (m, 2H), 3.47 (s, 1H), 3.02 (s, 3H), 2.57 (s, 1H), 2.06 (s, 2H), 1.79 (s, 4H), 1.70 – 1.50 (m, 5H), 1.39 – 1.12 (m, 5H).

Compound 68: (22.9 mg, 22.9%) as a solid.

LCMS(E05569-095): m/z (ES+), $[M+H]^+ = 453.59$; NH_4HCO_3 , HPLC $t_R = 1.689$ min.

^1H NMR (400 MHz, Methanol- d_4) δ 7.26 (d, 4H), 7.14 (dt, 1H), 4.73 (d, 1H), 4.32 (s, 1H), 3.87 (s, 1H), 3.69 (d, 2H), 3.02 (s, 4H), 2.97 – 2.51 (m, 2H), 2.06 (s, 4H), 1.73 – 1.51 (m, 7H), 1.25 (d, 5H).

Compound 69 and 70



To a stirred solution of N-[(3R,4S)-1-(1-fluorocyclobutanecarbonyl)-3-[[1-(4-phenylcyclohexyl)methoxy]piperidin-4-yl]methanesulfonamide

5 (95.00 mg, 0.204 mmol, 1.00 eq) and BH_3 . THF

(4.00 mL, 41.796 mmol, 205.29 eq) was stirred for 3 h at room temperature under nitrogen atmosphere. The resulting mixture was concentrated under reduced pressure. To the above mixture was added MeOH (5.00 mL) in portions at room temperature. The resulting

10 mixture was stirred for additional overnight at 50 degrees C. Desired product could be detected by LCMS. The resulting mixture was concentrated under reduced pressure.

The residue was purified by Prep-TLC (PE/EtOAc 1:1) to afford N-[(3R,4S)-1-(1-fluorocyclobutyl)methyl]-3-[[1-(4-phenylcyclohexyl)methoxy]piperidin-4-yl]methanesulfonamide (91 mg, 98.75%) as a solid. The crude product (91

15 mg) was purified by Prep-CHIRAL-HPLC to afford:

Compound 69: (35mg, 39%) as an oil.

LCMS: m/z (ES+), $[M+H]^+ = 452.6$
 ^1H NMR (400 MHz, Methanol- d_4) δ 7.26 (d, 4H), 7.16 (d, 1H), 3.73 (d, 1H), 3.62 – 3.47 (m, 3H), 2.99 (d, 4H), 2.77 (d, 1H), 2.71 (d, 1H), 2.67 – 2.55 (m, 2H), 2.42 (d, 2H), 2.30 – 2.12 (m, 4H), 2.11 – 1.94 (m, 2H), 1.85 (td, 3H), 1.69 (td, 7H), 1.58 – 1.43 (m, 1H), 1.31 (s, 1H).

20

Compound 70: (30mg, 33%) as an oil.

LCMS: m/z (ES+), $[M+H]^+ = 452.6$

^1H NMR (400 MHz, Methanol- d_4) δ 7.30 – 7.21 (m, 4H), 7.15 (dt, 1H), 3.73 (dt, 1H), 3.65 – 3.48 (m, 3H), 2.99 (d, 4H), 2.77 (d, 1H), 2.74 – 2.68 (m, 1H), 2.67 – 2.56 (m, 2H), 2.42 (t,

2H), 2.22 (dt, 4H), 2.05 (d, 1H), 2.02 – 1.93 (m, 1H), 1.93 – 1.76 (m, 3H), 1.76 – 1.60 (m, 7H), 1.52 (d, 1H).


Example 2: Human OX₂R IP1 assay









5 T-Rex CHO cells stably overexpressing the human orexin-2 receptor (OX₂R) are induced overnight with 1 µg/mL of doxycycline in a T225 flask. 24 hours post induction, cells are lifted with accutase and plated into a 384-well proxy plate at 30,000 cells/well. Cells are then treated with different test compounds in 1X stimulation buffer containing 10 mM Hepes, 1 mM CaCl₂, 0.5 mM MgCl₂, 4.2 mM KCl, 146 mM NaCl, 5.5 mM glucose, and 50
10 mM LiCl, pH 7.4, for 1 hr at 37 degrees C. Following incubation, the reaction is terminated by the addition of detection mix, which is composed of IP1-d2 and anti-IP1-cryptate diluted in lysis buffer as well as 1X stimulation buffer. The plates are allowed to incubate for 1 hour at room temperature and are then read in the EnVision® multimode plate reader, measuring inositol phosphate levels.









15 Cisbio IP1 is a cell-based functional assay quantifying the accumulation of inositol monophosphate (IP), a metabolite released as a result of orexin 2 receptor activation through the phospholipase C-Gq signaling pathway. This is a competitive immunoassay in which the IP1 produced by the cells upon receptor activation competes with the IP1 analog coupled to the d2 fluorophore (acceptor) for binding to an anti-IP1 monoclonal antibody labeled with Eu
20 cryptate (donor). The measured HTRF-FRET based signal is inversely proportional to the IP1 concentration produced.








The EC₅₀ values reported in Table 2 were obtained according to the human OX₂R IP1 assay described above. Data are the mean EC₅₀ values ± S.E.M.








25 Table 2.




Structure	Compound No.	EC ₅₀ (nM)
	37	**

	<p>38</p>	<p>**</p>
	<p>39</p>	<p>**</p>
	<p>40</p>	<p>**</p>
	<p>41</p>	<p>*</p>
	<p>42</p>	<p>**</p>
	<p>43</p>	<p>***</p>
	<p>44</p>	<p>*</p>
	<p>45</p>	<p>**</p>

	<p>46</p>	<p>*</p>
	<p>47</p>	<p>**</p>
	<p>48</p>	<p>*</p>
	<p>49</p>	<p>***</p>
	<p>50</p>	<p>**</p>
	<p>51</p>	<p>*</p>
	<p>52</p>	<p>*</p>
	<p>53</p>	<p>*</p>

	<p>54</p>	<p>**</p>
	<p>55</p>	<p>***</p>
	<p>56</p>	<p>**</p>
	<p>57</p>	<p>*</p>
	<p>58</p>	<p>*</p>
	<p>59</p>	<p>**</p>
	<p>60</p>	<p>*</p>

	61	*
	62	*
	63	*
	64	*
	65	***
	66	**
	67	***

	68	**
	69	**
	70	*

*** is $EC_{50} < 100$ nM


** is EC_{50} 100-10,000 nM

* is $EC_{50} > 10,000$ nM

- 5 While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

What is claimed is:

1. A compound of Formula I or a pharmaceutically acceptable salt thereof:



or a pharmaceutically acceptable salt thereof;

5 wherein:

ring A is selected from the group consisting of phenyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, further wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen;

10 X is N or CH;

Y is S(=O)₂, C(=O), or S(=O)(=NR_e);

R_e is selected from the group consisting of H, C₁-C₃ alkyl, or C₃-C₅ cycloalkyl;

E is selected from the group consisting of NR_aR_b, C₁-C₃ alkylene-NR_aR_b, C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-
 15 membered heterocyclyl, C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-membered heteroaryl and C₁-C₃ alkylene-(5- to 7-membered heteroaryl), wherein the C₁-C₃ alkylene-NR_aR_b, C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-membered heterocyclyl, C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-
 20 membered heteroaryl or C₁-C₃ alkylene-(5- to 7-membered heteroaryl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxyl;

R_a and R_b are each, independently, selected from the group consisting of H, C₁-C₃ alkyl, C₃-C₅ cycloalkyl, and 4- to 7-membered heterocyclyl, wherein the C₁-C₃ alkyl, C₃-C₅

cycloalkyl, or 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, C₁-C₃ alkyl, or C₁-C₃ alkoxy;

or, alternatively, R_a and R_b, together with the N atom to which they are attached, form a 4- to 7-membered heterocyclyl or 5- to 7-membered heteroaryl, wherein the 4- to 7-

5 membered heterocyclyl or 5- to 7-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, C₁-C₃ alkyl, C₁-C₃ alkoxy, or C₁-C₃ alkyl substituted with 1-3 halogen;

R₁ is selected from the group consisting of C(=O)-C₁-C₄ alkyl, C(=O)-C₁-C₄ alkoxy, C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), C(=O)-
 10 (CR_cR_d)_n-(C₆-C₁₀ aryl), C(=O)-(CR_cR_d)_n-(5- to 10-membered heteroaryl), C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), (CR_cR_d)_n-(C₆-C₁₀ aryl) and (CR_cR_d)_n-(5- to 10-membered heteroaryl), wherein the C₁-C₄ alkyl, C₁-C₄ alkoxy, C₃-C₈ cycloalkyl, 4- to 7-membered heterocyclyl, C₆-C₁₀ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted
 15 C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

R_c and R_d are each, independently, H, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

n is 0, 1, or 2;

each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H, halogen, unsubstituted C₁-
 20 C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium;

or, alternatively, R₃ and R₆, together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;

or, alternatively, R₄ and R₅, together form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen;

25 m is 0 or 1;

p is 0, 1, 2, 3, or 4; and

each R₉ is, independently, selected from the group consisting of deuterium, halogen, hydroxyl, and cyano.

30 2. The compound according to claim 1, wherein ring A is a phenyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

3. The compound according to claim 1 or 2, wherein ring A is a phenyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
4. The compound according to any of the preceding claims, wherein ring A is a phenyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
5. The compound according to any of the preceding claims, wherein ring A is a phenyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.
6. The compound according to any of the preceding claims, wherein ring A is a phenyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.
7. The compound according to any of the preceding claims, wherein ring A is a phenyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from F or Cl.
8. The compound according to any of the preceding claims, wherein ring A is a phenyl ring, wherein the ring is substituted with 1-2 F substituents.
9. The compound according to any of the preceding claims, wherein ring A is 3-fluorophenyl.
10. The compound of claim 1, wherein ring A is a pyridinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
11. The compound according to claim 1 or claim 10, wherein ring A is a pyridinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
12. The compound according to any one of claims 1, 10, or 11, wherein ring A is a pyridinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
13. The compound according to any one of claims 1 or 10-12, wherein ring A is a pyridinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.

14. The compound according to any one of claims 1 or 10-13, wherein ring A is a pyridinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.
15. The compound according to any one of claims 1 or 10-14, wherein ring A is a pyridinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.
16. The compound according to any one of claims 1 or 10-15, wherein ring A is a pyridinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.
17. The compound according to claim 1, wherein ring A is a pyridazinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
18. The compound according to claim 1 or 17, wherein ring A is a pyridazinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
19. The compound according to any one of claims 1, 17, or 18, wherein ring A is a pyridazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
20. The compound according to any one of claims 1 or 17-19, wherein ring A is a pyridazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.
21. The compound according to any one of claims 1 or 17-20, wherein ring A is a pyridazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.
22. The compound according to any one of claims 1 or 17-19, wherein ring A is a pyridazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.
23. The compound according to any one of claims 1 or 17-19, wherein ring A is a pyridazinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.
24. The compound according to claim 1, wherein ring A is a pyrimidinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.

25. The compound according to claim 1 or 24, wherein ring A is a pyrimidinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
26. The compound according to any one of claims 1, 24, or 25, wherein ring A is a
5 pyrimidinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
27. The compound according to any one of claims 1 or 24-26, wherein ring A is a pyrimidinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.
- 10 28. The compound according to any one of claims 1 or 24-27, wherein ring A is a pyrimidinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.
29. The compound according to any one of claims 1 or 24-28, wherein ring A is a
15 pyrimidinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.
30. The compound according to any one of claims 1 or 24-29, wherein ring A is a pyrimidinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.
31. The compound according to claim 1, wherein ring A is a pyrazinyl ring, wherein the
20 ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
32. The compound according to claim 1 or 31, wherein ring A is a pyrazinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
33. The compound according to any one of claims 1, 31 or 32, wherein ring A is a pyrazinyl
25 ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
34. The compound according to any one of claims 1 or 31-33, wherein ring A is a pyrazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.

35. The compound according to any one of claims 1 or 31-34, wherein ring A is a pyrazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.
36. The compound according to any one of claims 1 or 31-35, wherein ring A is a pyrazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.
37. The compound according to any one of claims 1 or 31-36, wherein ring A is a pyrazinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.
38. The compound according to claim 1, wherein ring A is a triazinyl ring, wherein the ring is unsubstituted or substituted with 1-3 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
39. The compound according to claim 1 or 38, wherein ring A is a triazinyl ring, wherein the ring is unsubstituted or substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
40. The compound according to any one of claims 1, 38, or 39, wherein ring A is a triazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl, C₃-C₅ cycloalkyl, or halogen.
41. The compound according to any one of claims 1 or 38-40, wherein ring A is a triazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl or halogen.
42. The compound according to any one of claims 1 or 38-41, wherein ring A is a triazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from halogen.
43. The compound according to any one of claims 1 or 38-42, wherein ring A is a triazinyl ring, wherein the ring is substituted with 1-2 substituents, each independently selected from C₁-C₃ alkyl.
44. The compound according to any one of claims 1 or 38-43, wherein ring A is a triazinyl ring, wherein the ring is mono-substituted with C₃-C₅ cycloalkyl.
45. The compound according to any one of claims 1-44, wherein p is 0.
46. The compound according to any one of claims 1-44, wherein p is 1.

47. The compound according to any one of claims 1-44, wherein p is 2.
48. The compound according to any one of claims 1-44, wherein p is 3.
49. The compound according to any one of claims 1-44, wherein p is 4.
50. The compound according to any one of claims 1-44, wherein p is 0, 1, or 2.
- 5 51. The compound according to any one of claims 1-44, wherein p is 0 or 1.
52. The compound according to any one of claims 1-51, wherein E is NR_aR_b.
53. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkylene-NR_aR_b.
54. The compound according to any one of claims 1-51, wherein E is unsubstituted C₁-C₃
10 alkyl, unsubstituted C₂-C₄ alkenyl or unsubstituted C₂-C₄ alkynyl.
55. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkyl, C₂-C₄ alkenyl or C₂-C₄ alkynyl substituted with one or more halogen, hydroxyl, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
56. The compound according to any one of claims 1-51, wherein E is unsubstituted C₁-C₃
15 alkyl.
57. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkyl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
58. The compound according to any one of claims 1-51, wherein E is unsubstituted C₃-C₈
20 cycloalkyl.
59. The compound according to any one of claims 1-51, wherein E is C₃-C₈ cycloalkyl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
60. The compound according to any one of claims 1-51, wherein E is unsubstituted C₁-C₃
25 alkylene-(C₃-C₈ cycloalkyl).
61. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkylene-(C₃-C₈ cycloalkyl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.

62. The compound according to any one of claims 1-51, wherein E is unsubstituted 4- to 10-membered heterocyclyl.
63. The compound according to any one of claims 1-51, wherein E is 4- to 10-membered heterocyclyl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
5
64. The compound according to any one of claims 1-51, wherein E is unsubstituted C₁-C₃ alkylene-(4- to 10-membered heterocyclyl).
65. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkylene-(4- to 10-membered heterocyclyl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
10
66. The compound according to any one of claims 1-51, wherein E is unsubstituted C₆-C₁₀ aryl.
67. The compound according to any one of claims 1-51, wherein E is C₆-C₁₀ aryl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
- 15 68. The compound according to any one of claims 1-51, wherein E is unsubstituted C₁-C₃ alkylene-(C₆-C₁₀ aryl).
69. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkylene-(C₆-C₁₀ aryl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
- 20 70. The compound according to any one of claims 1-51, wherein E is unsubstituted 5- to 7-membered heteroaryl.
71. The compound according to any one of claims 1-51, wherein E is 5- to 7-membered heteroaryl substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
- 25 72. The compound according to any one of claims 1-51, wherein E is unsubstituted C₁-C₃ alkylene-(5- to 7-membered heteroaryl).
73. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkylene-(5- to 7-membered heteroaryl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.

74. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-membered heterocyclyl, or C₁-C₃ alkylene-(4- to 10-membered heterocyclyl), wherein the C₁-C₃alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4-
5 to 10-membered heterocyclyl, or C₁-C₃ alkylene-(4- to 10-membered heterocyclyl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
75. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-membered heterocyclyl, or C₁-C₃
10 alkylene-(4- to 10-membered heterocyclyl), wherein the C₁-C₃alkyl, C₃-C₈ cycloalkyl, C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), 4- to 10-membered heterocyclyl, or C₁-C₃ alkylene-(4- to 10-membered heterocyclyl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
76. The compound according to any one of claims 1-51, wherein E is C₁-C₃ alkyl, C₃-C₈
15 cycloalkyl, or C₁-C₃ alkylene-(C₃-C₈ cycloalkyl), wherein the C₁-C₃alkyl, C₃-C₈ cycloalkyl, or C₁-C₃ alkylene-(C₃-C₈ cycloalkyl) is unsubstituted or substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
77. The compound according to any one of claims 1-51, wherein E is methyl.
78. The compound according to any one of claims 1-51, wherein E is methyl substituted
20 with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
79. The compound according to any one of claims 1-51, wherein E is CF₃.
80. The compound according to any one of claims 1-51, wherein E is CHF₂.
81. The compound according to any one of claims 1-51, wherein E is CH₂F.
82. The compound according to any one of claims 1-51, wherein E is NH(CH₃).
- 25 83. The compound according to any one of claims 1-51, wherein E is N(CH₃)₂.
84. The compound according to any one of claims 1-51, wherein E is C₆-C₁₀ aryl, C₁-C₃ alkylene-(C₆-C₁₀ aryl), 5- to 7-membered heteroaryl or C₁-C₃ alkylene-(5- to 7-membered heteroaryl).
85. The compound according to any one of claims 1-51, wherein E is unsubstituted C₆-C₁₀
30 aryl or C₁-C₃ alkylene-(C₆-C₁₀ aryl).

86. The compound according to any one of claims 1-51, wherein E is C₆-C₁₀ aryl or C₁-C₃ alkylene-(C₆-C₁₀ aryl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
87. The compound according to any one of claims 1-51, wherein E is 5- to 7-membered heteroaryl or C₁-C₃ alkylene-(5- to 7-membered heteroaryl).
88. The compound according to any one of claims 1-51, wherein E is unsubstituted 5- to 7-membered heteroaryl or C₁-C₃ alkylene-(5- to 7-membered heteroaryl).
89. The compound according to any one of claims 1-51, wherein E is 5- to 7-membered heteroaryl or C₁-C₃ alkylene-(5- to 7-membered heteroaryl) substituted with one or more halogen, hydroxyl, NR_cR_d, CF₃, CHF₂, CH₂F, C₁-C₃ alkyl, or C₁-C₃ alkoxy.
90. The compound according to any one of claims 1-89, wherein Y is S(=O)₂.
91. The compound according to any one of claims 1-89, wherein Y is C(=O).
92. The compound according to any one of claims 1-89, wherein Y is S(=O)(=NR_e).
93. The compound according to any one of claims 1-92, wherein X is N.
94. The compound according to any one of claims 1-92, wherein X is CH.
95. The compound according to any one of claims 1-94, wherein R₁ is selected from the group consisting of C(=O)-C₁-C₄ alkyl, C(=O)-C₁-C₄ alkoxy, C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), and C(=O)-(CR_cR_d)_n-(5- to 10-membered heteroaryl), wherein the C₁-C₄ alkyl, C₁-C₄ alkoxy, C₃-C₈ cycloalkyl, 4- to 7-membered heterocyclyl, C₆-C₁₀ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.
96. The compound according to any one of claims 1-94, wherein R₁ is selected from the group consisting of C(=O)-C₁-C₄ alkyl, C(=O)-C₁-C₄ alkoxy, and C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, wherein the C₁-C₄ alkyl, C₁-C₄ alkoxy, or C₃-C₈ cycloalkyl, is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.
97. The compound according to any one of claims 1-94, wherein R₁ is selected from the group consisting of C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), and C(=O)-(CR_cR_d)_n-(5- to 10-membered heteroaryl), wherein the 4- to 7-membered

heterocyclyl, C₆-C₁₀ aryl, or 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

5 98. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-C₁-C₄ alkyl, wherein the C₁-C₄ alkyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

10 99. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-C₁-C₄ alkoxy, wherein the C₁-C₄ alkoxy is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

100. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, wherein the C₃-C₈ cycloalkyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

15 101. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), wherein the 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

20 102. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(C₆-C₁₀ aryl), wherein the C₆-C₁₀ aryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

25 103. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(5- to 10-membered heteroaryl), wherein the 5- to 10-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

30 104. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(5- to 7-membered heterocyclyl), wherein the 5- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

105. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5- to 6-membered heterocyclyl), wherein the 5- to 6-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
- 5 106. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5-membered heterocyclyl), wherein the 5-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
107. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (6-
10 membered heterocyclyl), wherein the 6-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
108. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (C_6 aryl), wherein the C_6 aryl is unsubstituted or substituted with one or more halogen, hydroxyl,
15 unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
109. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (C_8 aryl), wherein the C_8 aryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
110. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ -
20 (C_{10} aryl), wherein the C_{10} aryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
111. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5-
25 to 6-membered heteroaryl), wherein the 5- to 6-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
112. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5-
30 membered heteroaryl), wherein the 5-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.

113. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heteroaryl), wherein the 6-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
- 5 114. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (7-membered heteroaryl), wherein the 7-membered heteroaryl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
115. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-O-(CR_cR_d)_n$ -
10 C_3-C_8 cycloalkyl or $C(=O)-O-(CR_cR_d)_n$ - (4- to 7-membered heterocyclyl), wherein the C_3-C_8 cycloalkyl or 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
116. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-O-(CR_cR_d)_n$ -
15 C_3-C_8 cycloalkyl, wherein the C_3-C_8 cycloalkyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
117. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-O-(CR_cR_d)_n$ -
20 (4- to 7-membered heterocyclyl), wherein the 4- to 7-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
118. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-O-(CR_cR_d)_n$ -
25 C_3-C_5 cycloalkyl, wherein the C_3-C_5 cycloalkyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
119. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-O-(CR_cR_d)_n$ -
(5- to 6-membered heterocyclyl), wherein the 5- to 6-membered heterocyclyl is unsubstituted or substituted with one or more halogen, hydroxyl, unsubstituted C_1-C_3 alkyl, or C_1-C_3 alkyl substituted with one or more halogen or deuterium.
- 30 120. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-O-(CR_cR_d)_n$ - (5-membered heterocyclyl), wherein the 5-membered heterocyclyl is unsubstituted or

substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

121. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n- (6-membered heterocyclyl), wherein the 6-membered heterocyclyl is unsubstituted or
5 substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

122. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is unsubstituted.

10 123. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

124. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is unsubstituted and further wherein n is 0.
15

125. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃
20 alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

126. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is unsubstituted and further wherein n is 1.

25 127. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) or (CR_cR_d)_n-(5- to 10-membered heteroaryl) wherein the C₆-C₁₀ aryl or 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.

30 128. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is unsubstituted.

129. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.
130. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is unsubstituted and further wherein n is 0.
131. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.
132. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is unsubstituted and further wherein n is 1.
133. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C₆-C₁₀ aryl) wherein the C₆-C₁₀ aryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.
134. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is unsubstituted.
135. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.
136. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is unsubstituted and further wherein n is 0.
137. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.
138. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is unsubstituted and further wherein n is 1.

139. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5- to 10-membered heteroaryl) wherein the 5- to 10-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium and further wherein n is 1.
- 5 140. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is unsubstituted.
141. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl
10 is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium.
142. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is unsubstituted and further wherein n is 0.
- 15 143. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium and further wherein n is 0.
144. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(phenyl)
20 or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is unsubstituted and further wherein n is 1.
145. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) wherein the phenyl or 5- to 7-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl
25 substituted with one or more halogen or deuterium and further wherein n is 1.
146. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(phenyl) or $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is unsubstituted.
147. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(phenyl)
30 or $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is

substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

148. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(phenyl) or (CR_cR_d)_n-(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is unsubstituted and further wherein n is 0.

149. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(phenyl) or (CR_cR_d)_n-(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

150. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(phenyl) or (CR_cR_d)_n-(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is unsubstituted and further wherein n is 1.

151. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(phenyl) or (CR_cR_d)_n-(6-membered heteroaryl) wherein the phenyl or 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 1.

152. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(6-membered heteroaryl) wherein the 6-membered heteroaryl is unsubstituted.

153. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(6-membered heteroaryl) wherein the 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium.

154. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(6-membered heteroaryl) wherein the 6-membered heteroaryl is unsubstituted and further wherein n is 0.

155. The compound according to any one of claims 1-94, wherein R₁ is (CR_cR_d)_n-(6-membered heteroaryl) wherein the 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C₁-C₃ alkyl, or C₁-C₃ alkyl substituted with one or more halogen or deuterium and further wherein n is 0.

156. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the 6-membered heteroaryl is unsubstituted and further wherein n is 1.
157. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(6-membered heteroaryl) wherein the 6-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium and further wherein n is 1.
158. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5-membered heteroaryl) wherein the 5-membered heteroaryl is unsubstituted.
159. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5-membered heteroaryl) wherein the 5-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium.
160. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5-membered heteroaryl) wherein the 5-membered heteroaryl is unsubstituted and further wherein n is 0.
161. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5-membered heteroaryl) wherein the 5-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium and further wherein n is 0.
162. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5-membered heteroaryl) wherein the 5-membered heteroaryl is unsubstituted and further wherein n is 1.
163. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5-membered heteroaryl) wherein the 5-membered heteroaryl is substituted with one or more halogen, hydroxyl, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium and further wherein n is 1.
164. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)$ - C_1 - C_4 alkyl and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
165. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)$ - C_1 - C_4 alkyl and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H or unsubstituted C_1 - C_3 alkyl.

166. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-C₁-C₄ alkoxy and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
167. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-C₁-C₄ alkoxy and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl.
- 5 168. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
169. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H or unsubstituted C₁-C₃ alkyl.
170. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-C₃-
10 C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
171. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
172. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(4-
to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
- 15 173. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(4-
to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
174. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(4-
to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
175. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(5-
20 membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
176. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(5-
membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
177. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(5-
membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
- 25 178. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(6-
membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
179. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-
(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.

180. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (6-membered heterocyclyl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
181. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (C_6 - C_{10} aryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
- 5 182. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (C_6 - C_{10} aryl), n is 0, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
183. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (C_6 - C_{10} aryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
184. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (C_6 aryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
- 10 185. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (C_6 aryl), n is 0, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
186. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (C_6 aryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
- 15 187. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5- to 7-membered heteroaryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
188. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5- to 7-membered heteroaryl), n is 0, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
189. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5- to 7-membered heteroaryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
- 20 190. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5- membered heteroaryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
191. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5- membered heteroaryl), n is 0, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
- 25 192. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (5- membered heteroaryl), n is 1 or 2, and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.
193. The compound according to any one of claims 1-94, wherein R_1 is $C(=O)-(CR_cR_d)_n$ - (6- membered heteroaryl) and each of $R_2, R_3, R_4, R_5, R_6, R_7,$ and R_8 is H.

194. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(6-membered heteroaryl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
195. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-(CR_cR_d)_n-(6-membered heteroaryl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
- 5 196. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
197. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
198. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-C₃-C₈ cycloalkyl, n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
- 10 199. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
200. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
- 15 201. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(4- to 7-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
202. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
203. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
- 20 204. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(5-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
205. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl) and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
- 25 206. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 0, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.
207. The compound according to any one of claims 1-94, wherein R₁ is C(=O)-O-(CR_cR_d)_n-(6-membered heterocyclyl), n is 1 or 2, and each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.

208. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5- to 10-membered heteroaryl) and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
209. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(5- to 7-membered heteroaryl) and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
- 5 210. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(6-membered heteroaryl) and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
211. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(6-membered heteroaryl), n is 0, and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
212. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(6-
10 membered heteroaryl), n is 1 or 2, and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
213. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C_6 - C_{10} aryl) and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
214. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C_6 aryl) and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
- 15 215. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C_6 aryl), n is 0, and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
216. The compound according to any one of claims 1-94, wherein R_1 is $(CR_cR_d)_n$ -(C_6 aryl), n is 1 or 2, and each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is H.
217. The compound according to any one of claims 1-94, wherein R_1 is 3-pyridazinyl.
- 20 218. The compound according to any one of claims 1-217, wherein each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is, independently, H, halogen, unsubstituted C_1 - C_3 alkyl, or C_1 - C_3 alkyl substituted with one or more halogen or deuterium.
219. The compound according to any one of claims 1-217, wherein each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is, independently, H, halogen, or unsubstituted C_1 - C_3 alkyl.
- 25 220. The compound according to any one of claims 1-217, wherein each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is, independently, H or halogen.
221. The compound according to any one of claims 1-217, wherein each of R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 is, independently, H or fluorine.


222. The compound according to any one of claims 1-217, wherein each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is, independently, H or C₁-C₃ alkyl.


223. The compound according to any one of claims 1-217, wherein each of R₂, R₃, R₄, R₅, R₆, R₇, and R₈ is H.

5 224. The compound according to any one of claims 1-217, wherein R₃ and R₆, together, form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen.


225. The compound according to any one of claims 1-217, wherein R₄ and R₅, together, form an unsubstituted C₁-C₃ alkylene or a C₁-C₃ alkylene substituted with one or more halogen.

226. A compound or a pharmaceutically acceptable salt thereof selected from the group
10 consisting of:







5





5



5







227. A pharmaceutical composition comprising a compound of any one of claims 1-225, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

5 228. A method of treating narcolepsy in a subject in need thereof comprising administering to the subject a compound of any one of claims 1-226 or a pharmaceutically acceptable salt thereof, or a composition according to claim 227.

229. A method of treating cataplexy in a subject in need thereof comprising administering to the subject a compound of any one of claims 1-226 or a pharmaceutically acceptable salt
10 thereof, or a composition according to claim 227.

230. Use of a compound of any one of claims 1-226, or a pharmaceutically acceptable salt thereof, or a composition according to claim 227 for the manufacture of a medicament for narcolepsy.

231. Use of a compound of any one of claims 1-226 or a pharmaceutically acceptable salt
15 thereof, or a composition according to claim 227 for the manufacture of a medicament for cataplexy.

232. A compound of any one of claims 1-226 or a pharmaceutically acceptable salt thereof, or a composition according to claim 227 for use in a method of treating narcolepsy in a subject in need thereof.

20 233. A compound of any one of claims 1-226 or a pharmaceutically acceptable salt thereof, or a composition according to claim 227 for use in a method of treating cataplexy in a subject in need thereof.