



(51) International Patent Classification:

A61P 35/00 (2006.01) A61K 31/519 (2006.01)

C07D 519/00 (2006.01)

(21) International Application Number:

PCT/US2023/062235

(22) International Filing Date:

08 February 2023 (08.02.2023)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/308,424 09 February 2022 (09.02.2022) US

63/368,584 15 July 2022 (15.07.2022) US

63/373,302 23 August 2022 (23.08.2022) US

63/378,843 07 October 2022 (07.10.2022) US

63/384,374 18 November 2022 (18.11.2022) US

TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, CV, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, ME, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))


(71) Applicant: QUANTA THERAPEUTICS, INC. [US/US];
455 Mission Bay Boulevard South, San Francisco, California 94158-2158 (US).

(72) Inventors: LIN, Hong; 455 Mission Bay Boulevard South, San Francisco, California 94158-2158 (US). LUENGO, Juan; 455 Mission Bay Boulevard South, San Francisco, California 94158-2158 (US). JOHNSON, Neil; 455 Mission Bay Boulevard South, San Francisco, California 94158-2158 (US). HOSPITAL, Audrey; 455 Mission Bay Boulevard South, San Francisco, California 94158-2158 (US).

(74) Agent: MARTINEZ, Jose; WILSON SONSINI GOODRICH & ROSATI, P.C., 650 Page Mill Road, Palo Alto, California 94304 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CV, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH,

(54) Title: KRAS MODULATORS AND USES THEREOF



(II)

(57) Abstract: Provided herein are KRAS modulating compounds selected from fused pyrimidines such as compounds of Formula (II) or pharmaceutically acceptable salts, solvates, stereoisomers, atom labelled, or tautomers of any of the foregoing, useful for modulating KRAS G12 and/or other G12 mutants.



KRAS MODULATORS AND USES THEREOF**CROSS-REFERENCE**

[0001] This application claims the benefit of U.S. Provisional Patent Applications Nos. 63/308,424 filed on February 9, 2022; 63/368,584 filed on July 15, 2022; 63/373,302 filed on August 23, 2022; 63/378,843 filed on October 7, 2022; and 63/384,374 filed on November 18, 2022; the entire contents of each of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

[0002] The small GTPase protein Kirsten Rat Sarcoma 2 Viral Oncogene Homolog (KRAS) is a member of the Ras family of cell signaling switches, regulating growth and survival of normal and cancerous cells (e.g., see Cully, M. and J. Downward, SnapShot: Ras Signaling. Cell, 2008. 133(7): p. 1292-1292 e1). KRAS mutations drive approximately 25% of human cancers by aberrant regulation of the mitogen-activated protein kinase (MAPK) signaling cascade and other effector pathways (e.g., see Stephen, A.G., et al., Dragging ras back in the ring. Cancer Cell, 2014. 25(3): p. 272-81). Though Ras has been recognized as a target in cancer for about 40 years, Ras-driven cancers remain among the most difficult to treat due to insensitivity to available targeted therapies. Ras, encoded by the three major genes KRAS, NRAS and HRAS, has the highest frequency of mutation of any oncogene. All oncogenic Ras mutations drive the switch to accumulate in the active GTP-bound state. The most common Ras mutation found across human tumor types is KRAS G12D (e.g., see The AACR Project GENIE Consortium. Cancer Discovery, 2017. 7(8): p. 818-831. Dataset Version 4). Activating mutations in codon 12 impair the small GTPases' ability to perform their role in hydrolyzing GTP. This regulatory impairment is fundamental for initiating and maintaining tumor progression.

[0003] Despite extensive efforts, small molecules have not been identified which block effector binding or restore GTPase activating protein (GAP) sensitivity, though some have been found which block interaction of Ras with the guanine nucleotide exchange factor (GEF), SOS, which activates Ras at the plasma membrane. KRAS G12C mutations, most common in lung adenocarcinoma, have been clinically shown to be susceptible to direct inhibition by covalent modification with small molecule inhibitors trapping the protein in the inactive GDP-bound state. KRAS G12D mutation confers a significantly slower intrinsic rate of GTP hydrolysis than G12C, resulting in more constitutive activation. Thus, pharmacological targeting the of inactive state is unlikely to achieve similar results against G12D, despite the existence of a similar binding pocket in the GDP-state. Additionally, a cysteine present at the site of the activating mutation yields itself

to covalent chemistry, while aspartic acid does not provide typical medicinal chemistry approaches for selective covalent modification.


[0004] In order to potentially exploit the accumulation of KRAS G12D and other mutant variants in the GTP-bound state as a vulnerability to achieve selective inhibition of cancer cells while sparing normal Ras function, it is attractive for small molecule inhibitors to bind selectively to the GTP-state and stabilize a conformation that is incompetent for oncogenic signaling interactions with effector proteins. Furthermore, it has been shown that only constitutive activation of Raf, MEK and ERK kinases in the MAPK cascade downstream of Ras can bypass the requirement for Ras proteins in proliferative signaling (e.g., see Drosten, M., et al., Genetic analysis of Ras signalling pathways in cell proliferation, migration and survival. EMBO J, 2010. 29(6): p. 1091-104). As all evidence has indicated that MAPK signaling is essential for the growth effects of Ras in cancer, KRAS-mutant-selective inhibition in this pathway is considered the critical functional readout for potential clinical benefit of novel therapeutic approaches.

SUMMARY OF THE INVENTION

[0005] There is a need to develop new inhibitors for KRAS-driven cancers that demonstrate inhibition of MAPK signals via a mechanism of action that is selective for binding to the active GTP-bound state over the inactive GDP-bound state.

[0006] The present disclosure relates to Formula (I) or Formula (II) or Formula (III), including stereoisomers, tautomers, solvates, and pharmaceutically acceptable salts thereof, and to uses thereof in, for example, inhibiting KRas G12D and/or other G12 mutants.

[0007] In an aspect, the present disclosure provides a compound represented by the structure of Formula (I):



or a pharmaceutically acceptable salt thereof wherein:

R¹ is selected from C₃-C₁₂ carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl,

C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein the C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*};

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃-C₁₂ carbocycle;

Y is selected from a bond, O, S and NR⁵;

R² is selected from hydrogen, -N(R²¹)₂, -L-N(R²¹)₂, -L-OR²¹, heterocycle, C₁-C₆ alkyl, -L-heterocycle, -L-aryl, -L-heteroaryl, -L-cycloalkyl, -L-NHC(=NH)NH₂, -L-C(O)N(R²¹)₂, -L-C₁-C₆ haloalkyl, -L-OR²¹, -L-NR²¹C(O)-aryl, -L-COOH, -L-NR²¹S(O)₂(R²¹), -L-S(O)₂N(R²¹)₂, -L-N(R²¹)C(O)(OR²¹), -L-OC(O)N(R²¹)₂, and -LC(=O)OC₁-C₆ alkyl, wherein the heterocycle and the aryl portion of -L-NR⁵C(O)-aryl and the heterocycle portion of -L-heterocycle and the cycloalkyl portion of the -L-cycloalkyl are optionally substituted with one or more R⁶, and wherein the aryl or heteroaryl of the -L-aryl and the -L-heteroaryl are optionally substituted with one or more R⁷;

each L is independently selected from a C₁-C₄ alkylene optionally substituted with one or more substituents selected from hydroxy, C₁-C₄ hydroxyalkyl, C₁-C₄ alkyl, C₃-C₆ carbocycle, or 3- to 8-membered heterocycle, wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle or 3- to 8-membered heterocycle wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl;

each R⁴ is independently selected from C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, wherein C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R⁵ is independently selected from hydrogen or C₁-C₆ alkyl;

each R⁶ is independently selected from halogen, hydroxy, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, oxo, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, cyano, =CH₂, =NO-C₁-C₃ alkyl, C₁-C₃ aminoalkyl, -N(R⁵)S(O)₂(R⁵), -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C₁-C₃ alkyl

substituted pyrazolyl, tert-butyldimethylsilyloxyCH₂-, -N(R⁵)₂, (C₁-C₃ alkoxy)C₁-C₃ alkyl-, (C₁-C₃ alkyl)C(=O), oxo, (C₁-C₃ haloalkyl)C(=O)-, -SO₂F, (C₁-C₃ alkoxy)C₁-C₃ alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC₁-C₆ alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O)C₁-C₆ alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂C₁-C₆ alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl), -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl(C₁-C₃ alkyl)N(CH₃)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl, -OC(O)heterocycle, -O-C₁-C₃ alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C₁-C₃ alkyl)(C₁-C₃ alkyl)phenyl are optionally substituted with one or more substituents selected from -C(O)H and OH, and wherein the alkyl of -O-C₁-C₃ alkyl is optionally substituted with substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocycle is optionally substituted with oxo;

each R⁷ is independently selected from halogen, hydroxy, HC(=O)-, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ hydroxyalkyl, or -N(R⁵)₂;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, =NH, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each Q is independently selected from a bond, S, and O;

B is selected from a heterocycle and carbocycle, wherein the heterocycle and carbocycle are optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, =O, -NO₂, C₁-C₄ alkyl, C₁₋₆ aminoalkyl, -S-C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₂-C₄ hydroxyalkynyl, C₁-C₃ cyanoalkyl, triazolyl, C₁-C₃ haloalkyl, -O-C₁-C₃ haloalkyl, -S-C₁-C₃ haloalkyl, C₁-C₃ alkoxy, C₁-C₃ hydroxyalkyl, -CH₂C(=O)N(R⁵)₂, -C₃-C₄ alkynyl(NR⁵)₂, -N(R⁵)₂, (C₁-C₃ alkoxy)haloC₁-C₃ alkyl-, C₁₋₆ alkyl-N(R²⁰)₂, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, -NH₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl, and wherein B forms a spirocycle with Ring A; and


Ring A is selected from a heterocycle and carbocycle, wherein the heterocycle or carbocycle is optionally substituted with one or more substituents selected from R⁴.

[0008] In certain embodiments, the disclosure provides a pharmaceutical composition comprising a compound or salt of Formula (I) and a pharmaceutically acceptable excipient.

[0009] In certain embodiments, the disclosure provides a method of treating a disease or disorder, using a compound or salt of Formula (I). In certain embodiments, the disclosure provides a method of treating a disease or disorder, using a compound or salt of Formula (I) and a pharmaceutically acceptable excipient.

[0010] In certain embodiments, the disclosure provides a method of inhibiting KRas G12D and/or other G12 mutants, using a compound or salt of Formula (I). In certain embodiments, the disclosure provides a method of inhibiting KRas G12D and/or other G12 mutants, using a compound or salt of Formula (I) and a pharmaceutically acceptable excipient.

[0011] In an aspect, the present disclosure provides a compound represented by the structure of Formula (II):



Formula (II)

or a pharmaceutically acceptable salt thereof wherein:

M is selected from O, S, SO, SO₂, and NR³;

R¹ is selected from C₃-C₁₂ carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl-SO₂R²⁰, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein the C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*}; and wherein when M is NR³, Y is O, and R¹ is piperazine, the piperazine is substituted with one or more R⁹;

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃-C₁₂ carbocycle;

Y is selected from a bond, O, S and NR⁵;

R² is selected from -L-N(R²¹)₂, -L-OR²¹, heterocycle, C₁-C₆ alkyl, -L-heterocycle, -L-aryl, -L-heteroaryl, -L-cycloalkyl, -L-N(R²¹)₂, -L-NHC(=NH)NH₂, -L-C(O)N(R²¹)₂, -L-C₁-C₆ haloalkyl, -L-OR²¹, -L-NR²¹C(O)-aryl, -L-COOH, -L-NR²¹S(O)₂(R²¹), -L-S(O)₂N(R²¹)₂, -L-N(R²¹)C(O)(OR²¹), -L-OC(O)N(R²¹)₂, or -LC(=O)OC₁-C₆ alkyl, wherein the heterocycle, the aryl portion of -L-NR²¹C(O)-aryl, the heterocycle portion of -L-heterocycle, the cycloalkyl portion of the -L-cycloalkyl are each optionally substituted with one or more R⁶, and wherein the aryl portion of the -L- aryl and the heteroaryl portion of the -L-heteroaryl are each optionally substituted with one or more R⁷, and wherein when Y is a bond, O, or S, R² is further selected from hydrogen;

each L is independently selected from a C₁-C₄ alkylene optionally substituted with one or more substituents selected from hydroxy, C₁-C₄ hydroxyalkyl, C₁-C₄ alkyl, C₃-C₆ carbocycle, or 3- to 8-membered heterocycle, wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle or 3- to 8-membered heterocycle, wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl;

R³ is selected from hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxyalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, wherein C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

n is selected from 0 to 2;

each R⁴ is independently selected from C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, oxo, hydroxyl, halogen, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, wherein the C₁-C₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from cyano, halogen, —OR⁵, and —N(R⁵)₂;

each R⁵ is independently selected from hydrogen or C₁-C₆ alkyl;

each R⁶ is independently selected from halogen, hydroxy, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, oxo, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, cyano, =CH₂, =NO-C₁-C₃ alkyl, C₁-C₃ aminoalkyl, -N(R⁵)S(O)₂(R⁵), -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C₁-C₃ alkyl substituted pyrazolyl, tert-butyldimethylsilyloxyCH₂-, -N(R⁵)₂, (C₁-C₃ alkoxy)C₁-C₃ alkyl-,

(C₁-C₃ alkyl)C(=O), oxo, (C₁-C₃ haloalkyl)C(=O)-, -SO₂F, (C₁-C₃ alkoxy)C₁-C₃ alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC₁-C₆ alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O)C₁-C₆ alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂C₁-C₆ alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl), -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl(C₁-C₃ alkyl)N(CH₃)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl, -OC(O)heterocycle, -O-C₁-C₃ alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C₁-C₃ alkyl)(C₁-C₃ alkyl)phenyl are each optionally substituted with one or more substituents selected from -C(O)H and OH, and wherein the alkyl of -O-C₁-C₃ alkyl is optionally substituted with substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocycle is optionally substituted with oxo;

each Q is independently selected from a bond, S, and O;

each R⁷ is independently selected from halogen, hydroxy, HC(=O)-, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ hydroxyalkyl, or -N(R⁵)₂;

each R⁹ is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -NR²⁰S(O)₂R²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -NO₂, =O, =NO(R²⁰), -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, =NH, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle; and

B is selected from a heterocycle and carbocycle, wherein the heterocycle and carbocycle are each optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, =O, -NO₂, C₁-C₄ alkyl, C₁₋₆ aminoalkyl, -S-C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₂-C₄ hydroxyalkynyl, C₁-C₃ cyanoalkyl, triazolyl, C₁-C₃ haloalkyl, -O-C₁-C₃ haloalkyl, -S-C₁-C₃ haloalkyl, C₁-C₃ alkoxy, C₁-C₃ hydroxyalkyl, -CH₂C(=O)N(R⁵)₂, -C₃-C₄ alkynyl(NR⁵)₂, -N(R⁵)₂, (C₁-C₃ alkoxy)haloC₁-C₃ alkyl-, C₁₋₆ alkyl-N(R²⁰)₂, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted with one or more substituents selected from halogen,


-OH, -NO₂, -NH₂, =O, =S, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl.

[0012] In certain embodiments, the disclosure provides a pharmaceutical composition comprising a compound or salt of Formula (II) and a pharmaceutically acceptable excipient.

[0013] In certain embodiments, the disclosure provides a method of treating a disease or disorder, using a compound or salt of Formula (II). In certain embodiments, the disclosure provides a method of treating a disease or disorder, using a compound or salt of Formula (II) and a pharmaceutically acceptable excipient.

[0014] In certain embodiments, the disclosure provides a method of inhibiting KRas G12D and/or other G12 mutants, using a compound or salt of Formula (II). In certain embodiments, the disclosure provides a method of inhibiting KRas G12D and/or other G12 mutants, using a compound or salt of Formula (II) and a pharmaceutically acceptable excipient.

[0015] In an aspect, the present disclosure provides a compound represented by the structure of Formula (III):



Formula (III)

or a pharmaceutically acceptable salt thereof wherein:

R¹ is selected from C₃-C₁₂ carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein the C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*};

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃-C₁₂ carbocycle;

R^2 is selected from $-L-NR^{21}S(O)_2(R^{21})$, $-L-S(O)_2N(R^{21})_2$, $-L-N(R^{21})C(O)(OR^{21})$, $-L-OC(O)N(R^{21})_2$, and L-bicyclic heterocycle, wherein the bicyclic heterocycle is optionally substituted with one or more R^6 ;

each L is independently selected from a C_1 - C_4 alkylene optionally substituted with one or more substituents selected from hydroxy, C_1 - C_4 hydroxyalkyl, C_1 - C_4 alkyl, C_3 - C_6 carbocycle, or 3- to 8-membered heterocycle, wherein the C_3 - C_6 carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, $-OH$, $-NO_2$, $=O$, $=S$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle or 3- to 8-membered heterocycle, wherein the C_3 - C_6 carbocycle and 3- to 8-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, $-OH$, $-NO_2$, $=O$, $=S$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl;

n is selected from 0 to 3;

each R^4 is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, oxo, hydroxyl, halogen, C_{3-12} carbocycle, and 3- to 12-membered heterocycle, wherein the C_1 - C_6 alkyl, C_{3-12} carbocycle, and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from cyano, halogen, $-OR^5$, and $-N(R^5)_2$;

B is selected from a heterocycle and carbocycle, wherein the heterocycle or carbocycle is optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, $=O$, $-NO_2$, C_1 - C_4 alkyl, C_{1-6} aminoalkyl, $-S-C_1-C_3$ alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_2 - C_4 hydroxyalkynyl, C_1 - C_3 cyanoalkyl, triazolyl, C_1 - C_3 haloalkyl, $-O-C_1-C_3$ haloalkyl, $-S-C_1-C_3$ haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 hydroxyalkyl, $-CH_2C(=O)N(R^5)_2$, $-C_3$ - C_4 alkynyl(NR^5)₂, $-N(R^5)_2$, $(C_1$ - C_3 alkoxy)halo C_1 - C_3 alkyl-, C_{1-6} alkyl- $N(R^{20})_2$, C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, wherein C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, $-OH$, $-NO_2$, $-NH_2$, $=O$, $=S$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl;

Y is selected from a bond, O, S and NR^5 ;

each R^6 is independently selected from halogen, hydroxy, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, oxo, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, cyano, $=CH_2$, $=NO-C_1-C_3$ alkyl, C_1 - C_3 aminoalkyl, $-N(R^5)S(O)_2(R^5)$, $-Q$ -phenyl, $-Q$ -phenyl SO_2F , $-NHC(O)$ phenyl, $-NHC(O)$ phenyl SO_2F , C_1 - C_3 alkyl substituted pyrazolyl, tert-butyldimethylsilyloxy CH_2- , $-N(R^5)_2$, $(C_1$ - C_3 alkoxy) C_1 - C_3 alkyl-, $(C_1$ - C_3 alkyl) $C(=O)$, oxo, $(C_1$ - C_3 haloalkyl) $C(=O)-$, $-SO_2F$, $(C_1$ - C_3 alkoxy) C_1 - C_3 alkoxy, $-CH_2OC(O)N(R^5)_2$, $-CH_2NHC(O)OC_1$ - C_6 alkyl, $-CH_2NHC(O)N(R^5)_2$, $-CH_2NHC(O)C_1$ - C_6 alkyl, $-CH_2$ (pyrazolyl), $-CH_2NHSO_2C_1$ - C_6 alkyl, $-CH_2OC(O)$ heterocycle, $-OC(O)N(R^5)_2$, $-OC(O)NH(C_1$ - C_3 alkyl) $O(C_1$ - C_3 alkyl), $-OC(O)NH(C_1$ - C_3 alkyl) $O(C_1$ - C_3 alkyl)phenyl(C_1 - C_3

alkyl)N(CH₃)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl, -OC(O)heterocycle, -O-C₁-C₃ alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C₁-C₃ alkyl)(C₁-C₃ alkyl)phenyl are optionally substituted with one or more substituents selected from -C(O)H and OH, and wherein the alkyl of -O-C₁-C₃ alkyl is optionally substituted with substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocycle is optionally substituted with oxo;

each Q is independently selected from a bond, S, and O;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle; and

each R⁵ is independently selected from hydrogen or C₁-C₆ alkyl.

[0016] In certain embodiments, the disclosure provides a pharmaceutical composition comprising a compound or salt of Formula (III) and a pharmaceutically acceptable excipient.

[0017] In certain embodiments, the disclosure provides a method of treating a disease or disorder, using a compound or salt of Formula (III). In certain embodiments, the disclosure provides a method of treating a disease or disorder, using a compound or salt of Formula (III) and a pharmaceutically acceptable excipient.

[0018] In certain embodiments, the disclosure provides a method of inhibiting KRas G12D and/or other G12 mutants, using a compound or salt of Formula (III). In certain embodiments, the disclosure provides a method of inhibiting KRas G12D and/or other G12 mutants, using a compound or salt of Formula (III) and a pharmaceutically acceptable excipient.

INCORPORATION BY REFERENCE

[0019] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

DETAILED DESCRIPTION OF THE INVENTION

[0020] The following description sets forth numerous exemplary configurations, methods, parameters, and the like. It should be recognized, however, that such description is not intended

as a limitation on the scope of the present disclosure, but is instead provided as a description of exemplary embodiments.

[0021] In the following description, certain specific details are set forth in order to provide a thorough understanding of various embodiments of the disclosure. However, one skilled in the art will understand that the disclosure may be practiced without these details.

Definitions

[0022] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. All patents and publications referred to herein are incorporated by reference.

[0023] "Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, and preferably having from one to fifteen carbon atoms (*i.e.*, C₁-C₁₅ alkyl). In certain embodiments, an alkyl comprises one to thirteen carbon atoms (*i.e.*, C₁-C₁₃ alkyl). In certain embodiments, an alkyl comprises one to eight carbon atoms (*i.e.*, C₁-C₈ alkyl). In other embodiments, an alkyl comprises one to five carbon atoms (*i.e.*, C₁-C₅ alkyl). In other embodiments, an alkyl comprises one to four carbon atoms (*i.e.*, C₁-C₄ alkyl). In other embodiments, an alkyl comprises one to three carbon atoms (*i.e.*, C₁-C₃ alkyl). In other embodiments, an alkyl comprises one to two carbon atoms (*i.e.*, C₁-C₂ alkyl). In other embodiments, an alkyl comprises one carbon atom (*i.e.*, C₁ alkyl). In other embodiments, an alkyl comprises five to fifteen carbon atoms (*i.e.*, C₅-C₁₅ alkyl). In other embodiments, an alkyl comprises five to eight carbon atoms (*i.e.*, C₅-C₈ alkyl). In other embodiments, an alkyl comprises two to five carbon atoms (*i.e.*, C₂-C₅ alkyl). In other embodiments, an alkyl comprises three to five carbon atoms (*i.e.*, C₃-C₅ alkyl). In certain embodiments, the alkyl group is selected from methyl, ethyl, 1-propyl (*n*-propyl), 1-methylethyl (*iso*-propyl), 1-butyl (*n*-butyl), 1-methylpropyl (*sec*-butyl), 2-methylpropyl (*iso*-butyl), 1,1-dimethylethyl (*tert*-butyl), 1-pentyl (*n*-pentyl). The alkyl is attached to the rest of the molecule by a single bond.

[0024] The term "C_{x-y}" when used in conjunction with a chemical moiety, such as alkyl, alkenyl, or alkynyl is meant to include groups that contain from x to y carbons in the chain. For example, the term "C₁₋₆alkyl" refers to substituted or unsubstituted saturated hydrocarbon groups, including straight-chain alkyl and branched-chain alkyl groups that contain from 1 to 6 carbons. The term -C_{x-y}alkylene- refers to a substituted or unsubstituted alkylene chain with from x to y carbons in the alkylene chain. For example -C₁₋₆alkylene- may be selected from methylene, ethylene, propylene, butylene, pentylene, and hexylene, any one of which is optionally substituted.

[0025] "Alkoxy" refers to a radical bonded through an oxygen atom of the formula –O-alkyl, where alkyl is an alkyl chain as defined above.

[0026] "Alkenyl" refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one carbon-carbon double bond, and preferably having from two to twelve carbon atoms (*i.e.*, C₂-C₁₂ alkenyl). In certain embodiments, an alkenyl comprises two to eight carbon atoms (*i.e.*, C₂-C₈ alkenyl). In certain embodiments, an alkenyl comprises two to six carbon atoms (*i.e.*, C₂-C₆ alkenyl). In other embodiments, an alkenyl comprises two to four carbon atoms (*i.e.*, C₂-C₄ alkenyl). The alkenyl is attached to the rest of the molecule by a single bond, for example, ethenyl (*i.e.*, vinyl), prop-1-enyl (*i.e.*, allyl), but-1-enyl, pent-1-enyl, penta-1,4-dienyl, and the like.

[0027] "Alkynyl" refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one carbon-carbon triple bond, and preferably having from two to twelve carbon atoms (*i.e.*, C₂-C₁₂ alkynyl). In certain embodiments, an alkynyl comprises two to eight carbon atoms (*i.e.*, C₂-C₈ alkynyl). In other embodiments, an alkynyl comprises two to six carbon atoms (*i.e.*, C₂-C₆ alkynyl). In other embodiments, an alkynyl comprises two to four carbon atoms (*i.e.*, C₂-C₄ alkynyl). The alkynyl is attached to the rest of the molecule by a single bond, for example, ethynyl, propynyl, butynyl, pentynyl, hexynyl, and the like.

[0028] The terms "C_{x-y}alkenyl" and "C_{x-y}alkynyl" refer to substituted or unsubstituted unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond, respectively. The term –C_{x-y}alkenylene- refers to a substituted or unsubstituted alkenylene chain with from x to y carbons in the alkenylene chain. For example, –C₂₋₆alkenylene- may be selected from ethenylene, propenylene, butenylene, pentenylene, and hexenylene, any one of which is optionally substituted. An alkenylene chain may have one double bond or more than one double bond in the alkenylene chain. The term –C_{x-y}alkynylene- refers to a substituted or unsubstituted alkynylene chain with from x to y carbons in the alkenylene chain. For example, –C₂₋₆alkynylene- may be selected from ethynylene, propynylene, butynylene, pentynylene, and hexynylene, any one of which is optionally substituted. An alkynylene chain may have one triple bond or more than one triple bond in the alkynylene chain.

[0029] "Alkylene" or "alkylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing no unsaturation, and preferably having from one to twelve carbon atoms, for example, methylene, ethylene, propylene, *n*-butylene, and the like. The alkylene chain is attached to the rest of the molecule through a single bond and to the radical group through a

single bond. The points of attachment of the alkylene chain to the rest of the molecule and to the radical group may be through any two carbons within the chain. In certain embodiments, an alkylene comprises one to ten carbon atoms (*i.e.*, C₁-C₈ alkylene). In certain embodiments, an alkylene comprises one to eight carbon atoms (*i.e.*, C₁-C₈ alkylene). In other embodiments, an alkylene comprises one to five carbon atoms (*i.e.*, C₁-C₅ alkylene). In other embodiments, an alkylene comprises one to four carbon atoms (*i.e.*, C₁-C₄ alkylene). In other embodiments, an alkylene comprises one to three carbon atoms (*i.e.*, C₁-C₃ alkylene). In other embodiments, an alkylene comprises one to two carbon atoms (*i.e.*, C₁-C₂ alkylene). In other embodiments, an alkylene comprises one carbon atom (*i.e.*, C₁ alkylene). In other embodiments, an alkylene comprises five to eight carbon atoms (*i.e.*, C₅-C₈ alkylene). In other embodiments, an alkylene comprises two to five carbon atoms (*i.e.*, C₂-C₅ alkylene). In other embodiments, an alkylene comprises three to five carbon atoms (*i.e.*, C₃-C₅ alkylene).

[0030] "Alkenylene" or "alkenylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing at least one carbon-carbon double bond, and preferably having from two to twelve carbon atoms. The alkenylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkenylene chain to the rest of the molecule and to the radical group may be through any two carbons within the chain. In certain embodiments, an alkenylene comprises two to ten carbon atoms (*i.e.*, C₂-C₁₀ alkenylene). In certain embodiments, an alkenylene comprises two to eight carbon atoms (*i.e.*, C₂-C₈ alkenylene). In other embodiments, an alkenylene comprises two to five carbon atoms (*i.e.*, C₂-C₅ alkenylene). In other embodiments, an alkenylene comprises two to four carbon atoms (*i.e.*, C₂-C₄ alkenylene). In other embodiments, an alkenylene comprises two to three carbon atoms (*i.e.*, C₂-C₃ alkenylene). In other embodiments, an alkenylene comprises two carbon atom (*i.e.*, C₂ alkenylene). In other embodiments, an alkenylene comprises five to eight carbon atoms (*i.e.*, C₅-C₈ alkenylene). In other embodiments, an alkenylene comprises three to five carbon atoms (*i.e.*, C₃-C₅ alkenylene).

[0031] "Alkynylene" or "alkynylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing at least one carbon-carbon triple bond, and preferably having from two to twelve carbon atoms. The alkynylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkynylene chain to the rest of the molecule and to the radical group may be through any two carbons within the chain. In certain embodiments, an alkynylene comprises two to ten carbon atoms (*i.e.*, C₂-C₁₀ alkynylene). In certain embodiments, an alkynylene comprises two to eight

carbon atoms (*i.e.*, C₂-C₈ alkynylene). In other embodiments, an alkynylene comprises two to five carbon atoms (*i.e.*, C₂-C₅ alkynylene). In other embodiments, an alkynylene comprises two to four carbon atoms (*i.e.*, C₂-C₄ alkynylene). In other embodiments, an alkynylene comprises two to three carbon atoms (*i.e.*, C₂-C₃ alkynylene). In other embodiments, an alkynylene comprises two carbon atom (*i.e.*, C₂ alkynylene). In other embodiments, an alkynylene comprises five to eight carbon atoms (*i.e.*, C₅-C₈ alkynylene). In other embodiments, an alkynylene comprises three to five carbon atoms (*i.e.*, C₃-C₅ alkynylene).

[0032] "Aryl" refers to a radical derived from an aromatic monocyclic or aromatic polycyclic hydrocarbon ring system by removing a hydrogen atom from a ring carbon atom. The aromatic monocyclic or aromatic polycyclic hydrocarbon ring system contains only hydrogen and carbon and from five to eighteen carbon atoms, where at least one of the rings in the ring system is aromatic, *i.e.*, it contains a cyclic, delocalized (4n+2) π -electron system in accordance with the Hückel theory. The ring system from which aryl groups are derived include, but are not limited to, groups such as benzene, fluorene, indane, indene, tetralin and naphthalene.

[0033] "Aralkyl" refers to a radical of the formula -R^c-aryl where R^c is an alkylene chain as defined above, for example, methylene, ethylene, and the like.

[0034] "Aralkenyl" refers to a radical of the formula -R^d-aryl where R^d is an alkenylene chain as defined above. "Aralkynyl" refers to a radical of the formula -R^e-aryl, where R^e is an alkynylene chain as defined above.

[0035] "Carbocycle" refers to a saturated, unsaturated or aromatic rings in which each atom of the ring is carbon. Carbocycle may include 3- to 10-membered monocyclic rings, 6- to 12-membered bicyclic rings, and 6- to 12-membered bridged rings. Each ring of a bicyclic carbocycle may be selected from saturated, unsaturated, and aromatic rings. An aromatic ring, *e.g.*, phenyl, may be fused to a saturated or unsaturated ring, *e.g.*, cyclohexane, cyclopentane, or cyclohexene. Any combination of saturated, unsaturated and aromatic bicyclic rings, as valence permits, are included in the definition of carbocyclic. Exemplary carbocycles include cyclopentyl, cyclohexyl, cyclohexenyl, adamantyl, phenyl, indanyl, and naphthyl. Bicyclic carbocycles may be fused, bridged or spiro-ring systems. In some cases, spiro-ring carbocycles have at least two molecular rings with only one common atom.

[0036] The term "unsaturated carbocycle" refers to carbocycles with at least one degree of unsaturation and excluding aromatic carbocycles. Examples of unsaturated carbocycles include cyclohexadiene, cyclohexene, and cyclopentene.

[0037] "Cycloalkyl" refers to a fully saturated monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which includes fused or bridged ring systems, and preferably having from three to twelve carbon atoms. In certain embodiments, a cycloalkyl

comprises three to ten carbon atoms. In other embodiments, a cycloalkyl comprises five to seven carbon atoms. The cycloalkyl may be attached to the rest of the molecule by a single bond. Examples of monocyclic cycloalkyls include, *e.g.*, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic cycloalkyl radicals include, for example, adamantyl, norbornyl (*i.e.*, bicyclo[2.2.1]heptanyl), norbornenyl, decalanyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like.

[0038] "Cycloalkenyl" refers to an unsaturated non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which includes fused or bridged ring systems, preferably having from three to twelve carbon atoms and comprising at least one double bond. In certain embodiments, a cycloalkenyl comprises three to ten carbon atoms. In other embodiments, a cycloalkenyl comprises five to seven carbon atoms. The cycloalkenyl may be attached to the rest of the molecule by a single bond. Examples of monocyclic cycloalkenyls includes, *e.g.*, cyclopentenyl, cyclohexenyl, cycloheptenyl, and cyclooctenyl.

[0039] "Cycloalkylalkyl" refers to a radical of the formula $-R^c$ -cycloalkyl where R^c is an alkylene chain as described above.

[0040] "Cycloalkylalkoxy" refers to a radical bonded through an oxygen atom of the formula $-O-R^c$ -cycloalkyl where R^c is an alkylene chain as described above.

[0041] "Halo" or "halogen" refers to halogen substituents such as bromo, chloro, fluoro and iodo substituents.

[0042] As used herein, the term "haloalkyl" or "haloalkane" refers to an alkyl radical, as defined above, that is substituted by one or more halogen radicals, for example, trifluoromethyl, dichloromethyl, bromomethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, and the like. In some embodiments, the alkyl part of the fluoroalkyl radical is optionally further substituted. Examples of halogen substituted alkanes ("haloalkanes") include halomethane (*e.g.*, chloromethane, bromomethane, fluoromethane, iodomethane), di- and trihalomethane (*e.g.*, trichloromethane, tribromomethane, trifluoromethane, triiodomethane), 1-haloethane, 2-haloethane, 1,2-dihaloethane, 1-halopropane, 2-halopropane, 3-halopropane, 1,2-dihalopropane, 1,3-dihalopropane, 2,3-dihalopropane, 1,2,3-trihalopropane, and any other suitable combinations of alkanes (or substituted alkanes) and halogens (*e.g.*, Cl, Br, F, I, etc.). When an alkyl group is substituted with more than one halogen radicals, each halogen may be independently selected *e.g.*, 1-chloro,2-fluoroethane.

[0043] "Fluoroalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more fluoro radicals, for example, trifluoromethyl, difluoromethyl, fluoromethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, and the like.

[0044] "Aminoalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more amine radicals, for example, propan-2-amine, butane-1,2-diamine, pentane-1,2,4-triamine and the like.

[0045] "Hydroxyalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more hydroxy radicals, for example, propan-1-ol, butane-1,4-diol, pentane-1,2,4-triol, and the like.

[0046] "Alkoxyalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more alkoxy radicals, for example, methoxymethane, 1,3-dimethoxybutane, 1-methoxypropane, 2-ethoxypentane, and the like.

[0047] "Cyanoalkyl" as used herein refers to an alkyl radical, as defined above, that is substituted by one or more cyano radicals, for example, acetonitrile, 2-ethyl-3-methylsuccinonitrile, butyronitrile, and the like.

[0048] "Heterocycle" as used herein refers to a saturated, unsaturated or aromatic ring comprising one or more heteroatoms. Exemplary heteroatoms include N, O, Si, P, B, and S atoms. The heterocycle may be attached to the rest of the molecule through any atom of the heterocycle, valence permitting, such as a carbon or nitrogen atom of the heterocycle. Heterocycles include 3- to 10-membered monocyclic rings, 6- to 12-membered bicyclic rings, and 6- to 12-membered bridged rings. A bicyclic heterocycle includes any combination of saturated, unsaturated and aromatic bicyclic rings, as valence permits. In an exemplary embodiment, an aromatic ring, e.g., pyridyl, may be fused to a saturated or unsaturated ring, e.g., cyclohexane, cyclopentane, morpholine, piperidine or cyclohexene. A bicyclic heterocycle includes any combination of ring sizes such as 4-5 fused ring systems, 5-5 fused ring systems, 5-6 fused ring systems, 6-6 fused ring systems, 5-7 fused ring systems, 6-7 fused ring systems, 5-8 fused ring systems, and 6-8 fused ring systems. Bicyclic heterocycles may be fused, bridged, or spiro-ring systems. A spiro-ring system may be referred as a "spiroheterocycle", "spiro heterocycle", or "spiro-heterocycle". In some cases, spiro-heterocycles, spiro heterocycles, or spiroheterocycles have at least two molecular rings with only one common atom. The spiro-heterocycle, spiro heterocycle, or spiroheterocycle comprises one or more heteroatoms.

[0049] "Heterocyclene" refers to a divalent heterocycle linking the rest of the molecule to a radical group.

[0050] "Heteroaryl" or "aromatic heterocycle" refers to a radical derived from a heteroaromatic ring radical that comprises one to eleven carbon atoms and at least one heteroatom wherein each heteroatom may be selected from N, O, and S. As used herein, the heteroaryl ring may be selected from monocyclic or bicyclic and fused or bridged ring systems rings wherein at least one of the rings in the ring system is aromatic, *i.e.*, it contains a cyclic, delocalized $(4n+2) \pi$ -

electron system in accordance with the Hückel theory. The heteroatom(s) in the heteroaryl radical may be optionally oxidized. One or more nitrogen atoms, if present, are optionally quaternized. The heteroaryl may be attached to the rest of the molecule through any atom of the heteroaryl, valence permitting, such as a carbon or nitrogen atom of the heteroaryl. Examples of heteroaryls include, but are not limited to, pyridine, pyrimidine, oxazole, furan, pyran, thiophene, isoxazole, benzimidazole, benzthiazole, and imidazopyridine.

[0051] An “X-membered heteroaryl” refers to the number of endocyclic atoms, i.e., X, in the ring. For example, a 5-membered heteroaryl ring or 5-membered aromatic heterocycle has 5 endocyclic atoms, e.g., triazole, oxazole, thiophene, etc.

[0052] The term “unsaturated heterocycle” refers to heterocycles with at least one degree of unsaturation and excluding aromatic heterocycles. Examples of unsaturated heterocycles include dihydropyrrole, dihydrofuran, oxazoline, pyrazoline, and dihydropyridine. Heterocycles may be optionally substituted by one or more substituents such as those substituents described herein.

[0053] The term “substituted” refers to moieties having substituents replacing a hydrogen on one or more carbons or substitutable heteroatoms, e.g., NH, of the structure. It will be understood that “substitution” or “substituted with” includes the implicit proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, i.e., a compound which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc. In certain embodiments, substituted refers to moieties having substituents replacing two hydrogen atoms on the same carbon atom, such as substituting the two hydrogen atoms on a single carbon with an oxo, imino or thioxo group. As used herein, the term “substituted” is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and non-aromatic substituents of organic compounds. The permissible substituents can be one or more and the same or different for appropriate organic compounds. For purposes of this disclosure, the heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valences of the heteroatoms.

[0054] In some embodiments, substituents may include any substituents described herein, for example: halogen, hydroxy, oxo (=O), thioxo (=S), cyano (-CN), nitro (-NO₂), imino (=N-H), oximo (=N-OH), hydrazino (=N-NH₂), -R^b-OR^a, -R^b-OC(O)-R^a, -R^b-OC(O)-OR^a, -R^b-OC(O)-N(R^a)₂, -R^b-N(R^a)₂, -R^b-C(O)R^a, -R^b-C(O)OR^a, -R^b-C(O)N(R^a)₂, -R^b-O-R^c-C(O)N(R^a)₂, -R^b-N(R^a)C(O)OR^a, -R^b-N(R^a)C(O)R^a, -R^b-N(R^a)S(O)_tR^a (where t is 1 or 2), -R^b-S(O)_tR^a (where t is 1 or 2), -R^b-S(O)_tOR^a (where t is 1 or

2), and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2); and alkyl, alkenyl, alkynyl, aryl, aralkyl, aralkenyl, aralkynyl, cycloalkyl, cycloalkylalkyl, and heterocycle, any of which may be optionally substituted by alkyl, alkenyl, alkynyl, halogen, haloalkyl, haloalkenyl, haloalkynyl, oxo ($=O$), thioxo ($=S$), cyano ($-CN$), nitro ($-NO_2$), imino ($=N-H$), oximo ($=N-OH$), hydrazine ($=N-NH_2$), $-R^b-OR^a$, $-R^b-OC(O)-R^a$, $-R^b-OC(O)-OR^a$, $-R^b-OC(O)-N(R^a)_2$, $-R^b-N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2); wherein each R^a is independently selected from hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocycloalkyl, heterocycloalkylalkyl, heteroaryl, or heteroarylalkyl, wherein each R^a , valence permitting, may be optionally substituted with alkyl, alkenyl, alkynyl, halogen, haloalkyl, haloalkenyl, haloalkynyl, oxo ($=O$), thioxo ($=S$), cyano ($-CN$), nitro ($-NO_2$), imino ($=N-H$), oximo ($=N-OH$), hydrazine ($=N-NH_2$), $-R^b-OR^a$, $-R^b-OC(O)-R^a$, $-R^b-OC(O)-OR^a$, $-R^b-OC(O)-N(R^a)_2$, $-R^b-N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2); and wherein each R^b is independently selected from a direct bond or a straight or branched alkylene, alkenylene, or alkynylene chain, and each R^c is a straight or branched alkylene, alkenylene or alkynylene chain.

[0055] As used herein, the term “electrophile” or “electrophilic moiety” is any moiety capable of reacting with a nucleophile (e.g., a moiety having a lone pair of electrons, a negative charge, a partial negative charge and/or an excess of electrons, for example an $-SH$ group). Electrophiles typically are electron poor or comprise atoms which are electron poor. In certain embodiments, an electrophile contains a positive charge or partial positive charge, has a resonance structure which contains a positive charge or partial positive charge, or is a moiety in which delocalization or polarization of electrons results in one or more atoms which contains a positive charge or partial positive charge. In some embodiments, an electrophile comprises a conjugated double bond, for example an α,β -unsaturated carbonyl or α,β -unsaturated thiocarbonyl compound.

[0056] As used herein, the term “optional” or “optionally” means that the subsequently described event or circumstances may or may not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, “optionally substituted aryl” means that the aryl group may or may not be substituted and that the description includes both substituted aryl groups and aryl groups having no substitution.

[0057] As used in the specification and claims, the singular form “a”, “an” and “the” includes plural references unless the context clearly dictates otherwise.

[0058] The term “salt” or “pharmaceutically acceptable salt” refers to salts derived from a variety of organic and inorganic counter ions well known in the art. Pharmaceutically acceptable acid addition salts can be formed with inorganic acids and organic acids. Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. Organic acids from which salts can be derived include, for example, acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid, salicylic acid, and the like. Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases. Inorganic bases from which salts can be derived include, for example, sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum, and the like. Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, basic ion exchange resins, and the like, specifically such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine. In some embodiments, the pharmaceutically acceptable base addition salt is chosen from ammonium, potassium, sodium, calcium, and magnesium salts.

[0059] The phrases “parenteral administration” and “administered parenterally” as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

[0060] The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0061] The phrase “pharmaceutically acceptable excipient” or “pharmaceutically acceptable carrier” as used herein means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium

carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

[0062] In certain embodiments, the term “prevent” or “preventing” as related to a disease or disorder may refer to a compound that, in a statistical sample, reduces the occurrence of the disorder or condition in the treated sample relative to an untreated control sample, or delays the onset or reduces the severity of one or more symptoms of the disorder or condition relative to the untreated control sample.


[0063] The terms “treat,” “treating” or “treatment,” as used herein, may include alleviating, abating or ameliorating a disease or condition symptoms, preventing additional symptoms, ameliorating or preventing the underlying causes of symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition either prophylactically and/or therapeutically.

[0064] The term “G12 mutants”, as used herein, refers to other oncogenic alleles of KRAS at amino acid position 12 (ie. G12X).

Compounds of the disclosure

[0065] The following is a discussion of compounds and salts thereof that may be used in the methods of the disclosure.

[0066] In some aspects, the present disclosure provides a compound represented by the structure of Formula (I):



Formula (I)

or a pharmaceutically acceptable salt thereof wherein:

R¹ is selected from C₃-C₁₂ carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, -

B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein the C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*};

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃-C₁₂ carbocycle;

Y is selected from a bond, O, S and NR⁵;

R² is selected from hydrogen, -N(R²¹)₂, -L-N(R²¹)₂, -L-OR²¹, heterocycle, C₁-C₆ alkyl, -L-heterocycle, -L-aryl, -L-heteroaryl, -L-cycloalkyl, -L-N(R²¹)₂, -L-NHC(=NH)NH₂, -L-C(O)N(R²¹)₂, -L-C₁-C₆ haloalkyl, -L-OR²¹, -L-NR²¹C(O)-aryl, -L-COOH, -L-NR²¹S(O)₂(R²¹), -L-S(O)₂N(R²¹)₂, -L-N(R²¹)C(O)(OR²¹), -L-OC(O)N(R²¹)₂, or -LC(=O)OC₁-C₆ alkyl, wherein the heterocycle and the aryl portion of -L-NR⁵C(O)-aryl and the heterocycle portion of -L-heterocycle and the cycloalkyl portion of the -L-cycloalkyl are optionally substituted with one or more R⁶, and wherein the aryl or heteroaryl of the -L-aryl and the -L-heteroaryl are optionally substituted with one or more R⁷;

each L is independently selected from a C₁-C₄ alkylene optionally substituted with one or more substituents selected from hydroxy, C₁-C₄ hydroxyalkyl, C₁-C₄ alkyl, C₃-C₆ carbocycle, or 3- to 8-membered heterocycle, wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle or 3- to 8-membered heterocycle wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl;

each R⁴ is independently selected from C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, wherein C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from halogen, -

OH, -CN, -NO₂, -NH₂, -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R⁵ is independently selected from hydrogen and C₁₋₆ alkyl;

each R⁶ is independently selected from halogen, hydroxy, C₁₋₃ hydroxyalkyl, C₁₋₃ alkyl, oxo, C₁₋₃ haloalkyl, C₁₋₃ alkoxy, cyano, =CH₂, =NO-C₁₋₃ alkyl, C₁₋₃ aminoalkyl, -N(R⁵)S(O)₂(R⁵), -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C₁₋₃ alkyl substituted pyrazolyl, tert-butyldimethylsilyloxyCH₂-, -N(R⁵)₂, (C₁₋₃ alkoxy)C₁₋₃ alkyl-, (C₁₋₃ alkyl)C(=O), oxo, (C₁₋₃ haloalkyl)C(=O)-, -SO₂F, (C₁₋₃ alkoxy)C₁₋₃ alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC₁₋₆ alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O)C₁₋₆ alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂C₁₋₆ alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C₁₋₃ alkyl)O(C₁₋₃ alkyl), -OC(O)NH(C₁₋₃ alkyl)O(C₁₋₃ alkyl)phenyl(C₁₋₃ alkyl)N(CH₃)₂, -OC(O)NH(C₁₋₃ alkyl)O(C₁₋₃ alkyl)phenyl, -OC(O)heterocycle, -O-C₁₋₃ alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C₁₋₃ alkyl)(C₁₋₃ alkyl)phenyl are optionally substituted with one or more substituents selected from -C(O)H and OH, and wherein the alkyl of -O-C₁₋₃ alkyl is optionally substituted with substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocycle is optionally substituted with oxo;

each R⁷ is independently selected from halogen, hydroxy, HC(=O)-, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ hydroxyalkyl, or -N(R⁵)₂;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, =NH, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each Q is independently selected from a bond, S, and O;

B is selected from a heterocycle and carbocycle, wherein the heterocycle and carbocycle are optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, =O, -NO₂, C₁₋₄ alkyl, C₁₋₆ aminoalkyl, -S-C₁₋₃ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₂₋₄ hydroxyalkynyl, C₁₋₃ cyanoalkyl, triazolyl, C₁₋₃ haloalkyl, -O-C₁₋₃ haloalkyl, -S-C₁₋₃ haloalkyl, C₁₋₃ alkoxy, C₁₋₃ hydroxyalkyl, -CH₂C(=O)N(R⁵)₂, -C₃₋₄ alkynyl(NR⁵)₂, -N(R⁵)₂, (C₁₋₃ alkoxy)haloC₁₋₃ alkyl-, C₁₋₆ alkyl-N(R²⁰)₂, C₃₋₁₂ carbocycle and 5- to 12-membered heterocycle, wherein C₃₋₁₂ carbocycle and 5- to 12-membered heterocycle are each

optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, -NH₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl, and wherein B forms a spirocycle with Ring A; and

Ring A is selected from a heterocycle and carbocycle, wherein the heterocycle or carbocycle is optionally substituted with one or more substituents selected from R⁴.

[0067] In some embodiments, Formula (I) is represented by Formula (II), Formula (II*), or Formula (III).

[0068] In some embodiments, for a compound or salt of Formula (I), Ring A is selected from a heterocycle wherein the heterocycle is optionally substituted with one or more substituents selected from R⁴. In some cases, Ring A includes at least one heteroatom selected from nitrogen, sulfur, and oxygen. In some cases, the heteroatom of Ring A is nitrogen, wherein the nitrogen is optionally substituted with R³, wherein R³ is selected from hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, wherein C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle. In some cases, the heteroatom of Ring A is sulfur, wherein the sulfur is optionally substituted with 1 or 2 oxygen atoms. In some cases, the heteroatom of Ring A is oxygen.

[0069] In some embodiments, for a compound or salt of Formula (I), Ring A is selected from a carbocycle, wherein the carbocycle is optionally substituted with one or more substituents selected from R⁴.

[0070] In some embodiments, for a compound or salt of Formula (I), R² is selected from -L-NR²¹S(O)₂(R²¹) and -L-S(O)₂N(R²¹)₂.

[0071] In some embodiments, for a compound or salt of Formula (I), R² is selected from -L-N(R²¹)C(O)(OR²¹), and -L-OC(O)N(R²¹)₂.

[0072] In some embodiments, for a compound or salt of Formula (I), each R²¹ is independently selected from hydrogen; C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, and oxo. In some cases, each R²¹ is independently selected from hydrogen; C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle. In some cases, each R²¹ is independently selected from hydrogen and C₁₋₆ alkyl.

[0073] In some embodiments, for a compound or salt of Formula (I), R^2 is selected from L-bicyclic heterocycle, wherein the bicyclic heterocycle is optionally substituted with one or more R^6 .

[0074] In some embodiments, for a compound or salt of Formula (I), R^2 is selected from L-pyrrolizine, wherein the pyrrolizine is optionally substituted with one or more R^6 .

[0075] In some embodiments, for a compound or salt of Formula (I), each L is independently selected from an optionally substituted C_1 - C_4 alkylene; and wherein optionally two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle, wherein the C_3 - C_6 carbocycle is optionally substituted with one or more substituents selected from halogen, -OH, - NO_2 , =O, =S, -CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl. In some cases, the optional substituents of L are selected from C_1 - C_4 hydroxyalkyl, C_1 - C_4 alkyl, C_3 - C_6 carbocycle; and wherein optionally two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle or 3- to 8-membered heterocycle wherein the C_3 - C_6 carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen and C_{1-6} haloalkyl.


[0076] In some embodiments, for a compound or salt of Formula (I), each L is independently selected from a substituted C_1 - C_4 alkylene, wherein two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle. In some cases, the C_3 - C_6 carbocycle is optionally substituted with one or more substituents selected from halogen, -OH, - NO_2 , =O, =S, -CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl.

[0077] In some embodiments, for a compound or salt of Formula (I), wherein each L is independently selected from a substituted C_1 - C_4 alkylene, and wherein two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle. In some cases, each L is independently selected from a substituted C_3 alkylene, and wherein two substituents on the same carbon atom of L come together to form a C_3 carbocycle. In some cases, each L is independently


selected from .

[0078] In some embodiments, for a compound or salt of Formula (I), R^2 is selected from -L-heterocycle, wherein the heterocycle portion of -L-heterocycle is optionally substituted with one or more R^6 . In some cases, the heterocycle is a saturated heterocycle. In some cases, the heterocycle has at least one nitrogen atom and at least one sulfur atom. In some cases, the heterocycle has at least one nitrogen atom. In some cases, the heterocycle has at least one sulfur atom.


[0079] In some embodiments, for a compound or salt of Formula (I), R^2 is selected from




[0080] In some embodiments, for a compound or salt of Formula (I), $Y-R^2$ is selected from





[0081] In some embodiments, for a compound or salt of Formula (I), $Y-R^2$ is selected from




[0082] In some embodiments, for a compound or salt of Formula (I), $Y-R^2$ is selected from




[0083] In some embodiments, for a compound or salt of Formula (I), R^2 is selected from -L-saturated heterocycle, wherein the saturated heterocycle portion of the -L-saturated heterocycle is optionally substituted with one or more R^6 , and contains one nitrogen atom and one sulfur atom.



cases, Y-R² is selected from



cases, Y-R² is selected from




[0084] In some embodiments, for a compound or salt of Formula (I), each R⁶ is independently selected from halogen, -OH, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, -CN, C₁-C₃ aminoalkyl, -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C₁-C₃ alkyl substituted pyrazolyl, -N(R⁵)₂, (C₁-C₃ alkoxy)C₁-C₃ alkyl-, (C₁-C₃ alkyl)C(=O), oxo, (C₁-C₃ haloalkyl)C(=O)-, -SO₂F, (C₁-C₃ alkoxy)C₁-C₃ alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC₁-C₆ alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O)C₁-C₆ alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂C₁-C₆ alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl), -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl(C₁-C₃ alkyl)N(CH₃)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl, -OC(O)heterocycle, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C₁-C₃ alkyl)(C₁-C₃ alkyl)phenyl are each optionally substituted with -C(O)H and OH, and wherein the heterocycle of -CH₂heterocycle is optionally substituted with oxo.

[0085] In some embodiments, for a compound or salt of Formula (I), each R⁶ is independently selected from halogen, -OH, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, -CN, and C₁-C₃ aminoalkyl.


[0086] In some embodiments, for a compound or salt of Formula (I), each R⁶ is independently selected from halogen, -OH, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ aminoalkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, -N(R⁵)₂, and oxo. In some cases, each R⁶ is independently selected from -OH, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ aminoalkyl, C₁-C₃ alkoxy, and -N(R⁵)₂. In some cases, each R⁶ is independently selected from C₁-C₃ alkyl, C₁-C₃ alkoxy, and -N(R⁵)₂.

[0087] In some embodiments, for a compound or salt of Formula (I), R⁶ is selected from halogen, -OH, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, -CN, and C₁-C₃ aminoalkyl. In some cases, R⁶ is selected from halogen and C₁-C₃ alkyl. In some cases, R⁶ is halogen. In some cases, R⁶ is C₁-C₃ alkyl. In some cases, R⁶ is selected from halogen and C₁-C₃ alkyl. In some cases, R⁶ is selected from methyl and fluorine.


[0088] In some embodiments, for a compound or salt of Formula (I), R² is selected from




[0089] In some embodiments, for a compound or salt of Formula (I), Y-R² is selected from




[0090] In some embodiments, for a compound or salt of Formula (I), Y-R² is selected from



[0091] In some embodiments, for a compound or salt of Formula (I), Y-R² is



[0092] In some embodiments, for a compound or salt of Formula (I), Y-R² is selected from



[0093] In some embodiments, for a compound or salt of Formula (I), B is an optionally substituted 5- to 15-membered heterocycle or optionally substituted C₃-C₁₅ carbocycle. In some cases, B is an optionally substituted 5- to 15-membered heterocycle. In some cases, B is an optionally substituted C₃-C₁₅ carbocycle.


[0094] In some embodiments, for a compound or salt of Formula (I), B is an optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle. In some cases, B is an optionally substituted 8- to 15-membered fused heterocycle. In some cases, B is an optionally substituted C₈-C₁₅ fused carbocycle.

[0095] In some embodiments, for a compound or salt of Formula (I), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle are each bicyclic or tricyclic. In some cases, for B, the optionally substituted 8- to 15-membered fused heterocycle are each bicyclic or tricyclic. In some cases, for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle are each bicyclic or tricyclic.


[0096] In some embodiments, for a compound or salt of Formula (I), B the heterocycle or carbocycle are each independently bicyclic. In some cases, the heterocycle is bicyclic. In some cases, the carbocycle is bicyclic.

[0097] In some embodiments, for a compound or salt of Formula (I), B the heterocycle or carbocycle are each independently tricyclic. In some cases, the heterocycle is tricyclic. In some cases, the carbocycle is tricyclic.


[0098] In some embodiments, for a compound or salt of Formula (I), B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused



[0099] In some embodiments, for a compound or salt of Formula (I), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused




[00100] In some embodiments, for a compound or salt of Formula (I), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused



[00101] In some embodiments, for a compound or salt of Formula (I), for B, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, C₁-C₃ alkyl, -B(OR²⁰)₂, -OR²⁰, -C(O)N(R²⁰)₂, -N(R²⁰)₂, =O, -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₂₋₆


alkenyl, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, oxo, -NH₂, C₁₋₃ alkyl, -B(OR²⁰)₂, -OH, -C(O)N(R²⁰)₂, =O, -CN, C₁₋₆ alkoxy, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, oxo, -NH₂, C₁₋₃ alkyl, -B(OH)₂, -OH, -C(O)NH₂, -NH₂, =O, -CN, C₁₋₆ alkoxy, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from oxo, -NH₂, -CN, halogen, C₁₋₃ alkyl. In some cases, the one or more optional substituents of the heterocycle or carbocycle are independently selected from oxo, -NH₂, halogen, C₁₋₃ alkyl.

[00102] In some embodiments, for a compound or salt of Formula (I), B is selected from



[00103] In some embodiments, for a compound or salt of Formula (I), for B, the one or more optional substituents of the heterocycle or carbocycle are independently selected from oxo, -NH₂, CN, halogen, C₁₋₃ alkyl.

[00104] In some embodiments, for a compound or salt of Formula (I), B is selected from



[00105] In some embodiments, for a compound or salt of Formula (I), each R^4 is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, oxo, hydroxyl, halogen. In some cases, each R^4 is independently selected from C_{1-6} alkyl, oxo, and halogen.

[00106] In some embodiments, for a compound or salt of Formula (I), n is selected from 1 and 2. In some cases, n is 0.

[00107] In some embodiments, for a compound or salt of Formula (I), Y is O.

[00108] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from optionally substituted 5- to 12-membered heterocycle.

[00109] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, SR^{20} , $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-NR^{20}S(O)_2R^{20}$, $C(O)N(R^{20})_2$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=NO(R^{20})$, CN , C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

[00110] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, $-SR^{20}$, $-S(O)_2(R^{20})$, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}-OR^{20}$, $-S(O)_2N(R^{20})_2$, $-NR^{20}S(O)_2R^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

[00111] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-NR^{20}S(O)_2R^{20}$, $-N(R^{20})C(O)OR^{20}$, and $-OC(O)N(R^{20})_2$.

[00112] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, -OH, $-N(R^{20})_2$, $-NO_2$, $=O$, -CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl.

[00113] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, -OH, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl.


[00114] In some embodiments, for a compound or salt of Formula (I), R^{20} of R^1 is selected from hydrogen and C_{1-3} alkyl.

[00115] In some embodiments, for a compound or salt of Formula (I), the 5- to 12-membered heterocycle of R^1 is an unsaturated heterocycle.

[00116] In some embodiments, for a compound or salt of Formula (I), the 5- to 12-membered heterocycle of R^1 is a saturated heterocycle.


[00117] In some embodiments, for a compound or salt of Formula (I), 5- to 12-membered heterocycle of R^1 is a bridged heterocycle.

[00118] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from




, and , each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, $=O$, -CN, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl.

[00119] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from



, each of which is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, $=O$, $-CN$, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl.

[00120] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from




[00121] In some embodiments, for a compound or salt of Formula (I), L is selected from C_1 - C_4 alkylene.

[00122] In some embodiments, for a compound or salt of Formula (I), L is selected from unsubstituted C_1 - C_4 alkylene.

[00123] In some embodiments, for a compound or salt of Formula (I), R^2 is $-L$ -heterocycle, optionally substituted with one or more R^6 , wherein the heterocycle portion is a bicyclic heterocycle. In some cases, the bicyclic heterocycle contains at least 1 nitrogen atom. In some cases, the bicyclic heterocycle contains at most 1 nitrogen atom.

[00124] In some embodiments, for a compound or salt of Formula (I), $Y-R^2$ is selected from




, wherein the heterocycle portion is optionally substituted with one or more R^6 .

[00125] In some embodiments, for a compound or salt of Formula (I), R^6 of R^2 is independently selected at each occurrence from halogen, hydroxy, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, cyano, and C_1 - C_3 aminoalkyl.


[00126] In some embodiments, for a compound or salt of Formula (I), R^6 of R^2 is independently selected at each occurrence from C_1 - C_3 alkyl and halogen.

[00127] In some embodiments, for a compound or salt of Formula (I), $Y-R^2$ is selected from



[00128] In some aspects, the present disclosure provides a compound represented by the structure of Formula (II):

[00129] A compound of Formula (II):



Formula (II)

or a pharmaceutically acceptable salt thereof wherein:

M is selected from O, S, SO, SO₂, and NR³;

R¹ is selected from C₃-C₁₂ carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl-SO₂R²⁰, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein the C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*}; and wherein when M is NR³, Y is O, and R¹ is piperazine, the piperazine is substituted with one or more R⁹;

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃-C₁₂ carbocycle;

Y is selected from a bond, O, S and NR⁵;

R² is selected from -L-N(R²¹)₂, -L-OR²¹, heterocycle, C₁-C₆ alkyl, -L-heterocycle, -L-aryl, -L-heteroaryl, -L-cycloalkyl, -L-NHC(=NH)NH₂, -L-C(O)N(R²¹)₂, -L-C₁-C₆ haloalkyl, -L-OR²¹, -L-NR²¹C(O)-aryl, -L-COOH, -L-NR²¹S(O)₂(R²¹), -L-S(O)₂N(R²¹)₂, -L-N(R²¹)C(O)(OR²¹), -L-OC(O)N(R²¹)₂, and -LC(=O)OC₁-C₆ alkyl, wherein the heterocycle, the aryl portion of -L-NR²¹C(O)-aryl, the heterocycle portion of -L-heterocycle, the cycloalkyl portion of the -L-cycloalkyl are each optionally substituted with one or more R⁶, and wherein the aryl portion of the -L-aryl and the heteroaryl portion of the -L-heteroaryl are each optionally

substituted with one or more R^7 , and wherein when Y is a bond, O, or S, R^2 is further selected from hydrogen;

each L is independently selected from a C_1 - C_4 alkylene optionally substituted with one or more substituents selected from hydroxy, C_1 - C_4 hydroxyalkyl, C_1 - C_4 alkyl, C_3 - C_6 carbocycle, or 3- to 8-membered heterocycle, wherein the C_3 - C_6 carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle or 3- to 8-membered heterocycle, wherein the C_3 - C_6 carbocycle and 3- to 8-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl;

R^3 is selected from hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkoxyalkyl, C_{3-12} carbocycle, and 3- to 12-membered heterocycle, wherein C_{3-12} carbocycle and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C_{1-6} alkyl)₂, C_{1-10} alkyl, - C_{1-10} haloalkyl, -O- C_{1-10} alkyl, oxo, C_{3-12} carbocycle, and 3- to 12-membered heterocycle;

n is selected from 0 to 2;

each R^4 is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, oxo, hydroxyl, halogen, C_{3-12} carbocycle, and 3- to 12-membered heterocycle, wherein the C_1 - C_6 alkyl, C_{3-12} carbocycle, and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from cyano, halogen, -OR⁵, and -N(R⁵)₂;

each R^5 is independently selected from hydrogen or C_1 - C_6 alkyl;

each R^6 is independently selected from halogen, hydroxy, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, oxo, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, cyano, =CH₂, =NO- C_1 - C_3 alkyl, C_1 - C_3 aminoalkyl, -N(R⁵)S(O)₂(R⁵), -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C_1 - C_3 alkyl substituted pyrazolyl, tert-butyldimethylsilyloxyCH₂-, -N(R⁵)₂, (C_1 - C_3 alkoxy) C_1 - C_3 alkyl-, (C_1 - C_3 alkyl)C(=O), oxo, (C_1 - C_3 haloalkyl)C(=O)-, -SO₂F, (C_1 - C_3 alkoxy) C_1 - C_3 alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC C_1 - C_6 alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O) C_1 - C_6 alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂ C_1 - C_6 alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl), -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl)phenyl(C_1 - C_3 alkyl)N(CH₃)₂, -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl)phenyl, -OC(O)heterocycle, -O- C_1 - C_3 alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C_1 - C_3 alkyl)(C_1 - C_3 alkyl)phenyl are each optionally substituted with one or more substituents selected from -C(O)H and OH, and wherein the alkyl of -O- C_1 - C_3 alkyl is optionally substituted with

substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocyclyl is optionally substituted with oxo;

each Q is independently selected from a bond, S, and O;

each R⁷ is independently selected from halogen, hydroxy, HC(=O)-, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ hydroxyalkyl, or -N(R⁵)₂;

each R⁹ is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -NR²⁰S(O)₂R²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -NO₂, =O, =NO(R²⁰), -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, =NH, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle; and

B is selected from a heterocycle and carbocycle, wherein the heterocycle and carbocycle are each optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, =O, -NO₂, C₁-C₄ alkyl, C₁₋₆ aminoalkyl, -S-C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₂-C₄ hydroxyalkynyl, C₁-C₃ cyanoalkyl, triazolyl, C₁-C₃ haloalkyl, -O-C₁-C₃ haloalkyl, -S-C₁-C₃ haloalkyl, C₁-C₃ alkoxy, C₁-C₃ hydroxyalkyl, -CH₂C(=O)N(R⁵)₂, -C₃-C₄ alkynyl(NR⁵)₂, -N(R⁵)₂, (C₁-C₃ alkoxy)haloC₁-C₃ alkyl-, C₁₋₆ alkyl-N(R²⁰)₂, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, -NH₂, =O, =S, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl.

[00130] In some embodiments, for a compound or salt of Formula (II), R² is selected from -L-NR²¹S(O)₂(R²¹) and -L-S(O)₂N(R²¹)₂.

[00131] In some embodiments, for a compound or salt of Formula (II), R² is selected from -L-N(R²¹)C(O)(OR²¹), and -L-OC(O)N(R²¹)₂.

[00132] In some embodiments, for a compound or salt of Formula (II), each R²¹ is independently selected from hydrogen; C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered

heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, and oxo. In some cases, each R²¹ is independently selected from hydrogen; C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle. In some cases, each R²¹ is independently selected from hydrogen and C₁₋₆ alkyl.

[00133] In some embodiments, for a compound or salt of Formula (II), R³ is selected from hydrogen and C₁₋₆ alkyl.

[00134] In some embodiments, for a compound or salt of Formula (II), M is selected from O, NH, and NMe. In some cases, M is O. In some cases, M is selected from NH and NMe.

[00135] In some embodiments, for a compound or salt of Formula (II), B is an optionally substituted 5- to 15-membered heterocycle or optionally substituted C₃-C₁₅ carbocycle. In some cases, B is an optionally substituted 5- to 15-membered heterocycle. In some cases, B is an optionally substituted C₃-C₁₅ carbocycle. In some cases, B is an optionally substituted 8- to 15-membered heterocycle. In some cases, B is an optionally substituted C₈-C₁₅ carbocycle.

[00136] In some embodiments, for a compound or salt of Formula (II), B is an optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle. In some cases, B is an optionally substituted 8- to 15-membered fused heterocycle. In some cases, B is an optionally substituted C₈-C₁₅ fused carbocycle.



[00137] In some embodiments, for a compound or salt of Formula (II), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle are each bicyclic or tricyclic. In some cases, for B, the optionally substituted 8- to 15-membered fused heterocycle are each bicyclic or tricyclic. In some cases, for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle are each bicyclic or tricyclic.

[00138] In some embodiments, for a compound or salt of Formula (II), for B, the optionally substituted 8- to 15-membered heterocycle contains at least one nitrogen atom. In some cases, the optionally substituted 8- to 15-membered heterocycle contains at least one sulfur atom. In some cases, the optionally substituted 8- to 15-membered heterocycle contains at most one nitrogen atom. In some cases, the optionally substituted 8- to 15-membered heterocycle contains at most one sulfur atom. In some cases, the optionally substituted 8- to 15-membered heterocycle contains at least two heteroatoms.



[00139] In some embodiments, for a compound or salt of Formula (II), B the heterocycle or carbocycle are each independently bicyclic. In some cases, the heterocycle is bicyclic. In some cases, the carbocycle is bicyclic.

[00140] In some embodiments, for a compound or salt of Formula (II), B the heterocycle or carbocycle are each independently tricyclic. In some cases, the heterocycle is tricyclic. In some cases, the carbocycle is tricyclic.



[00141] In some embodiments, for a compound or salt of Formula (II), B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused

carbocycle is selected from , and , each of which is optionally substituted with one or more substituents.

[00142] In some embodiments, for a compound or salt of Formula (II), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused

carbocycle is selected from , and , each of which is optionally substituted with one or more substituents.


[00143] In some embodiments, for a compound or salt of Formula (II), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused

carbocycle is selected from, , and , each of which is optionally substituted with one or more substituents.


[00144] In some embodiments, for a compound or salt of Formula (II), for B, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, C₁-C₃ alkyl, -B(OR²⁰)₂, -OR²⁰, -C(O)N(R²⁰)₂, -N(R²⁰)₂, =O, -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OR²⁰)₂, -OH, -C(O)N(R²⁰)₂, =O, -CN, C₁₋₆ alkoxy, and C₂₋₆ alkynyl. In some cases, the

one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OH)₂, -OH, -C(O)NH₂, -NH₂, =O, -CN, C₁₋₆ alkoxy, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from oxo, -NH₂, -CN, halogen, C₁-C₃ alkyl. In some cases, the one or more optional substituents of the heterocycle or carbocycle are independently selected from oxo, -NH₂, halogen, C₁-C₃ alkyl.

[00145] In some embodiments, for a compound or salt of Formula (II), B is selected from



[00146] In some embodiments, for a compound or salt of Formula (II), B is selected from



[00147] In some embodiments, for a compound or salt of Formula (II), each R^4 is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, oxo, hydroxyl, halogen. In some cases, each R^4 is independently selected from C_{1-6} alkyl, oxo, and halogen.

[00148] In some embodiments, for a compound or salt of Formula (II), n is selected from 1 and 2. In some cases, n is 0.

[00149] In some embodiments, for a compound or salt of Formula (II), Y is O.

[00150] In some embodiments, for a compound or salt of Formula (II), R^1 is selected from optionally substituted 5- to 12-membered heterocycle.

[00151] In some embodiments, for a compound or salt of Formula (II), R^1 is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-N(R^{20})_2$, $-NO_2$, $=O$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl.

[00152] In some embodiments, for a compound or salt of Formula (II), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, SR^{20} , $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=NO(R^{20})$, CN , C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

[00153] In some embodiments, for a compound or salt of Formula (II), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-NR^{20}S(O)_2R^{20}$, $-N(R^{20})C(O)OR^{20}$, and $-OC(O)N(R^{20})_2$.

[00154] In some embodiments, for a compound or salt of Formula (II), R^1 is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally

substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.


[00155] In some embodiments, for a compound or salt of Formula (II), R²⁰ of R¹ is selected from hydrogen and C₁₋₃ alkyl.


[00156] In some embodiments, for a compound or salt of Formula (II), the 5- to 12-membered heterocycle of R¹ is an unsaturated heterocycle.


[00157] In some embodiments, for a compound or salt of Formula (II), the 5- to 12-membered heterocycle of R¹ is a saturated heterocycle.


[00158] In some embodiments, for a compound or salt of Formula (II), 5- to 12-membered heterocycle of R¹ is a bridged heterocycle.


[00159] In some embodiments, for a compound or salt of Formula (II), R¹ is selected from



, and , each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, =O, -CN, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.




In some embodiments, for a compound or salt of Formula (II), R^1 is selected from ,




, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, =O, -CN, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.


[00160] In some embodiments, for a compound or salt of Formula (II), R^1 is selected from





[00161] In some embodiments, for a compound or salt of Formula (II), R^1 is selected from an optionally substituted saturated 6- to 7-membered heterocycle. In some cases, R^1 is selected from an optionally substituted saturated 6-membered heterocycle. In some cases, R^1 is selected from





, which is optionally substituted. In some cases, the optional one or more substituents are independently selected from halogen, -CN, -NHCN, C₁₋₆ cyanoalkyl, and C₁₋₆ alkyl. In some cases, the optional one or more substituents are independently selected from -CN, -NHCN, C₁₋₆ cyanoalkyl, and C₁₋₆ alkyl. In some cases, the optional one or more substituents are independently selected from -CN, -NHCN, C₁₋₆ cyanoalkyl, and C₁₋₆ alkyl. In some cases, the optional one or more substituents are independently selected from -NHCN, and C₁₋₆ alkyl. In




some cases, R^1 is selected from , which is substituted with one or more substituents selected



from -NHCN, and C₁₋₆ alkyl. In some cases, R^1 is selected from , and



[00162] In some embodiments, for a compound or salt of Formula (II), R^1 is selected from a substituted saturated 6-membered heterocycle, wherein the saturated 6-membered heterocycle is substituted with at least one -NHCN, and optionally one or more C_{1-6} alkyl; M is O; n is 0; B is selected from an optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C_8 - C_{15} fused carbocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, oxo, -NH₂, C_1 - C_3 alkyl, -OH, -C(O)NH₂, -NH₂, =O, -CN, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{2-6} alkynyl; Y is O; R^2 is selected from -L-heterocycle, wherein the heterocycle portion is optionally substituted with one or more substituents independently selected from halogen, hydroxy, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 hydroxyalkyl, or -N(R^5)₂; and L is selected from C_1 - C_4 alkylene. In some cases,




[00163] In some embodiments, for a compound or salt of Formula (II), each R^9 is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -NR²⁰S(O)₂R²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -NO₂, =O, =NO(R²⁰), -CN, -NHCN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl. In some cases, each R^9 is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -NR²⁰S(O)₂R²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -NO₂, =O, =NO(R²⁰), -CN, -NHCN. In some cases, each R^9 is independently selected from halogen, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -N(R²⁰)₂, -NO₂, =O, and =NO(R²⁰). In some cases, each R^9 is independently selected from halogen, and -N(R²⁰)₂.

[00164] In some embodiments, for a compound or salt of Formula (II), each L is independently selected from an optionally substituted C₁-C₄ alkylene; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle, wherein the C₃-C₆ carbocycle is optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl. In some cases, the optional substituents of L are selected from C₁-C₄ hydroxyalkyl, C₁-C₄ alkyl, C₃-C₆ carbocycle; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle or 3- to 8-membered heterocycle wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen and C₁₋₆ haloalkyl.



[00165] In some embodiments, for a compound or salt of Formula (II), each L is independently selected from a substituted C₁-C₄ alkylene, wherein two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle. In some cases, the C₃-C₆ carbocycle is optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl.

[00166] In some embodiments, for a compound or salt of Formula (II), wherein each L is independently selected from a substituted C₁-C₄ alkylene, and wherein two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle. In some cases, each L is independently selected from a substituted C₃ alkylene, and wherein two substituents on the same carbon atom of L come together to form a C₃ carbocycle. In some cases, each L is independently


selected from .

[00167] In some embodiments, for a compound or salt of Formula (II), R² is selected from -L-heterocycle, wherein the heterocycle portion of -L-heterocycle is optionally substituted with one or more R⁶. In some cases, the heterocycle is a saturated heterocycle. In some cases, the heterocycle has at least one nitrogen atom and at least one sulfur atom. In some cases, the heterocycle has at least one nitrogen atom. In some cases, the heterocycle has at least one sulfur atom.


[00168] In some embodiments, for a compound or salt of Formula (II), R² is selected from

 and , wherein the heterocycle portion is optionally substituted with one or more R⁶.


[00169] In some embodiments, for a compound or salt of Formula (II), Y-R² is selected from







[00170] In some embodiments, for a compound or salt of Formula (II), Y-R² is selected from



[00171] In some embodiments, for a compound or salt of Formula (II), Y-R² is selected from



[00172] In some embodiments, for a compound or salt of Formula (II), R² is selected from -L-saturated heterocycle, wherein the saturated heterocycle portion of the -L-saturated heterocycle is optionally substituted with one or more R⁶, and contains one nitrogen atom and one sulfur atom.



[00173] In some embodiments, for a compound or salt of Formula (II), each R⁶ is independently selected from halogen, -OH, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, -CN, C₁-C₃ aminoalkyl, -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C₁-C₃


alkyl substituted pyrazolyl, $-N(R^5)_2$, $(C_1-C_3 \text{ alkoxy})C_1-C_3 \text{ alkyl-}$, $(C_1-C_3 \text{ alkyl})C(=O)$, oxo, $(C_1-C_3 \text{ haloalkyl})C(=O)-$, $-SO_2F$, $(C_1-C_3 \text{ alkoxy})C_1-C_3 \text{ alkoxy}$, $-CH_2OC(O)N(R^5)_2$, $-CH_2NHC(O)OC_1-C_6 \text{ alkyl}$, $-CH_2NHC(O)N(R^5)_2$, $-CH_2NHC(O)C_1-C_6 \text{ alkyl}$, $-CH_2(\text{pyrazolyl})$, $-CH_2NHSO_2C_1-C_6 \text{ alkyl}$, $-CH_2OC(O)\text{heterocycle}$, $-OC(O)N(R^5)_2$, $-OC(O)NH(C_1-C_3 \text{ alkyl})O(C_1-C_3 \text{ alkyl})$, $-OC(O)NH(C_1-C_3 \text{ alkyl})O(C_1-C_3 \text{ alkyl})\text{phenyl}(C_1-C_3 \text{ alkyl})N(CH_3)_2$, $-OC(O)NH(C_1-C_3 \text{ alkyl})O(C_1-C_3 \text{ alkyl})\text{phenyl}$, $-OC(O)\text{heterocycle}$, and $-CH_2\text{heterocycle}$, wherein the phenyl of $-NHC(O)\text{phenyl}$ and $-OC(O)NH(C_1-C_3 \text{ alkyl})(C_1-C_3 \text{ alkyl})\text{phenyl}$ are each optionally substituted with $-C(O)H$ and OH , and wherein the heterocycle of $-CH_2\text{heterocycle}$ is optionally substituted with oxo.

[00174] In some embodiments, for a compound or salt of Formula (II), each R^6 is independently selected from halogen, $-OH$, $C_1-C_3 \text{ hydroxyalkyl}$, $C_1-C_3 \text{ alkyl}$, $C_1-C_3 \text{ haloalkyl}$, $C_1-C_3 \text{ alkoxy}$, $-CN$, and $C_1-C_3 \text{ aminoalkyl}$.


[00175] In some embodiments, for a compound or salt of Formula (II), each R^6 is independently selected from halogen, $-OH$, $C_1-C_3 \text{ hydroxyalkyl}$, $C_1-C_3 \text{ alkyl}$, $C_1-C_3 \text{ aminoalkyl}$, $C_1-C_3 \text{ haloalkyl}$, $C_1-C_3 \text{ alkoxy}$, $-N(R^5)_2$, and oxo. In some cases, each R^6 is independently selected from $-OH$, $C_1-C_3 \text{ hydroxyalkyl}$, $C_1-C_3 \text{ alkyl}$, $C_1-C_3 \text{ aminoalkyl}$, $C_1-C_3 \text{ alkoxy}$, and $-N(R^5)_2$. In some cases, each R^6 is independently selected from $C_1-C_3 \text{ alkyl}$, $C_1-C_3 \text{ alkoxy}$, and $-N(R^5)_2$.

[00176] In some embodiments, for a compound or salt of Formula (II), R^6 is selected from halogen, $-OH$, $C_1-C_3 \text{ hydroxyalkyl}$, $C_1-C_3 \text{ alkyl}$, $C_1-C_3 \text{ haloalkyl}$, $C_1-C_3 \text{ alkoxy}$, $-CN$, and $C_1-C_3 \text{ aminoalkyl}$. In some cases, R^6 is selected from halogen and $C_1-C_3 \text{ alkyl}$. In some cases, R^6 is halogen. In some cases, R^6 is $C_1-C_3 \text{ alkyl}$. In some cases, R^6 is selected from halogen and $C_1-C_3 \text{ alkyl}$. In some cases, R^6 is selected from methyl and fluorine.


[00177] In some embodiments, for a compound or salt of Formula (II), R^2 is selected from




[00178] In some embodiments, for a compound or salt of Formula (II), $Y-R^2$ is selected from




[00179] In some embodiments, for a compound or salt of Formula (II), $Y-R^2$ is selected from



[00180] In some embodiments, for a compound or salt of Formula (II), Y-R² is



[00181] In some embodiments, for a compound or salt of Formula (II), Y-R² is selected from



[00182] In some embodiments, for a compound or salt of Formula (II), L is selected from C₁-C₄ alkylene.


[00183] In some embodiments, for a compound or salt of Formula (II), L is selected from unsubstituted C₁-C₄ alkylene.

[00184] In some embodiments, for a compound or salt of Formula (II), R² is -L-heterocycle, optionally substituted with one or more R⁶, wherein the heterocycle portion is a bicyclic heterocycle. In some cases, the bicyclic heterocycle contains at least 1 nitrogen atom. In some cases, the bicyclic heterocycle contains at most 1 nitrogen atom.

[00185] In some embodiments, for a compound or salt of Formula (II), R² is selected from L-bicyclic heterocycle, wherein the bicyclic heterocycle is optionally substituted with one or more R⁶.


[00186] In some embodiments, for a compound or salt of Formula (II), R² is selected from L-pyrrolizine, wherein the pyrrolizine is optionally substituted with one or more R⁶.


[00187] In some embodiments, for a compound or salt of Formula (II), Y-R² is selected from





[00188] In some embodiments, for a compound or salt of Formula (II), R⁶ of R² is independently selected at each occurrence from halogen, hydroxy, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, cyano, and C₁-C₃ aminoalkyl. In some cases, R⁶ of R² is




independently selected at each occurrence from C₁-C₃ alkyl and halogen. In some cases, Y-R² is




[00189] In some embodiments, for a compound or salt of Formula (II), M is selected from NR³. In some cases, M is selected from NH and NMe. In some cases, M is selected from NMe and NCH₂CH₃. In some cases, M is NMe. In some cases, M is selected from C₁₋₆ cyanoalkyl. In some cases, M is selected from C₂ cyanoalkyl. In some cases, M is selected from NH. In some cases, R³ is selected from hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ alkoxyalkyl, and C₁₋₆ haloalkyl. In some cases, R³ is selected from hydrogen, C₁₋₆ alkyl, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ alkoxyalkyl, and C₁₋₆ haloalkyl. In some cases, R³ is selected from C₁₋₆ alkyl, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ alkoxyalkyl, and C₁₋₆ haloalkyl. In some cases, R³ is selected from C₂₋₆ alkyl, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ alkoxyalkyl, and C₁₋₆ haloalkyl. In some cases, R³ is selected from C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ alkoxyalkyl, and C₁₋₆ haloalkyl. In some cases, R³ is selected from C₁₋₆ cyanoalkyl, C₁₋₆ alkoxyalkyl, and C₁₋₆ haloalkyl. In some cases, R³ is selected from hydrogen, C₁₋₆ alkyl, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, and C₁₋₆ haloalkyl. In some cases, R³ is selected from C₁₋₆ alkoxyalkyl, C₁₋₆ cyanoalkyl, and C₁₋₆ alkyl. In some cases, R³ is . In some cases, R³ is selected from C₂₋₆ alkyl.

[00190] In some embodiments, for a compound or salt of Formula (II), M is NR³, B is selected


from  and , each of which is optionally substituted; n is 0; Y is O; R² is selected from L-heterocycle, wherein the heterocycle is optionally substituted with one or more R⁶; R¹ is

selected from ; and , which is optionally substituted. In some cases, R¹ is , which is optionally substituted with one or more substituents independently selected


from halogen, -OH, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxyalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, and C₂₋₆




alkynyl. In some cases, R¹ is



. In some cases, R¹ is



In some cases, R¹ is





. In some cases, R¹ is


. In some cases, the

heterocycle of L-heterocycle is bicyclic. In some cases, the heterocycle of L-heterocycle is monocyclic. In some cases, L is selected from an C₁-C₄ alkylene. In some cases, L is selected from an unsubstituted C₁-C₄ alkylene. In some cases, L is independently selected from a substituted C₁-C₄ alkylene, wherein two substituents on the same carbon atom of L come

together to form a C₃-C₆ carbocycle. In some cases, Y-R² is selected from




and



. In some cases, Y-R² is


selected from




, and





. In some cases, B is selected from



and
















, which is substituted with one or more substituents. In some cases, B is







which is substituted with one or more substituents. In some cases, B is selected from which is substituted with one or more substituents. In some cases, for B, the one or more substituents are independently selected from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OH)₂, -OH, -C(O)NH₂, -NH₂, =O, -CN, -O-C₁-C₃ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₂₋₆


alkynyl. In some cases, B is substituted with at least one halogen. In some cases, B is substituted with at least one chlorine. In some cases, B is substituted with at least one fluorine. In some



cases, B is selected from , , , , , ,






, , , , , , ,





, , and . In some cases, B is selected from 


which is substituted with one or more substituents selected from halogen, -O-C₁-C₃ haloalkyl,

and C₁₋₆ haloalkyl. In some cases, B is , which is substituted with one or more


substituents selected from halogen. In some cases, B is selected from , ,

, , , and . In some cases, B is , which is substituted with one or more substituents selected from fluorine. In some cases, B is selected


from , , and . In some cases, B is , which is substituted with one or more substituents selected from chlorine. In some cases, B is selected from




and . In some cases, R^3 is selected from hydrogen and C_{1-6} alkyl. In some cases, R^3 is selected from C_{1-6} alkyl. In some cases, R^3 is methyl. In some cases, each R^6 is selected from halogen, oxo, and C_{1-6} alkyl. In some cases, each R^6 is selected from halogen, and C_{1-6} alkyl. In some cases, each R^6 is selected from halogen. In some cases, R^1 is selected from




and




. In some cases, R^1 is



. In some cases, B is



[00191] In some aspects, the present disclosure provides a compound represented by the structure of Formula (III):



Formula (III)

or a pharmaceutically acceptable salt thereof wherein:

R^1 is selected from C_3 - C_{12} carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, $-SR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(=NR^{20})N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, wherein the C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*} ;

each R^{1*} is independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, $-SR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6}

aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃₋₁₂ carbocycle;

R² is selected from -L-NR²¹S(O)₂(R²¹), -L-S(O)₂N(R²¹)₂, -L-N(R²¹)C(O)(OR²¹), -L-OC(O)N(R²¹)₂, and L-bicyclic heterocycle, wherein the bicyclic heterocycle is optionally substituted with one or more R⁶;

each L is independently selected from a C₁₋₄ alkylene optionally substituted with one or more substituents selected from hydroxy, C₁₋₄ hydroxyalkyl, C₁₋₄ alkyl, C₃₋₆ carbocycle, or 3- to 8-membered heterocycle, wherein the C₃₋₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃₋₆ carbocycle or 3- to 8-membered heterocycle, wherein the C₃₋₆ carbocycle and 3- to 8-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl;

n is selected from 0 to 3;

each R⁴ is independently selected from C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, oxo, hydroxyl, halogen, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, wherein the C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from cyano, halogen, —OR⁵, and —N(R⁵)₂;

B is selected from a heterocycle and carbocycle, wherein the heterocycle or carbocycle is optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, =O, -NO₂, C₁₋₄ alkyl, C₁₋₆ aminoalkyl, -S-C₁₋₃ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₂₋₄ hydroxyalkynyl, C₁₋₃ cyanoalkyl, triazolyl, C₁₋₃ haloalkyl, -O-C₁₋₃ haloalkyl, -S-C₁₋₃ haloalkyl, C₁₋₃ alkoxy, C₁₋₃ hydroxyalkyl, -CH₂C(=O)N(R⁵)₂, -C₃₋₄ alkynyl(NR⁵)₂, -N(R⁵)₂, (C₁₋₃ alkoxy)haloC₁₋₃ alkyl-, C₁₋₆ alkyl-N(R²⁰)₂, C₃₋₁₂ carbocycle and 5- to 12-membered heterocycle, wherein C₃₋₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, -NH₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl;

Y is selected from a bond, O, S and NR⁵;

each R⁶ is independently selected from halogen, hydroxy, C₁₋₃ hydroxyalkyl, C₁₋₃ alkyl, oxo, C₁₋₃ haloalkyl, C₁₋₃ alkoxy, cyano, =CH₂, =NO-C₁₋₃ alkyl, C₁₋₃ aminoalkyl, -N(R⁵)S(O)₂(R⁵), -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C₁₋₃ alkyl substituted pyrazolyl, tert-butyldimethylsilyloxyCH₂-, -N(R⁵)₂, (C₁₋₃ alkoxy)C₁₋₃ alkyl-, (C₁₋₃ alkyl)C(=O), oxo, (C₁₋₃ haloalkyl)C(=O)-, -SO₂F, (C₁₋₃ alkoxy)C₁₋₃ alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC₁₋₆ alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O)C₁₋₆ alkyl, -

CH₂(pyrazolyl), -CH₂NHSO₂C₁-C₆ alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl), -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl(C₁-C₃ alkyl)N(CH₃)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl, -OC(O)heterocycle, -O-C₁-C₃ alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C₁-C₃ alkyl)(C₁-C₃ alkyl)phenyl are optionally substituted with one or more substituents selected from -C(O)H and OH, and wherein the alkyl of -O-C₁-C₃ alkyl is optionally substituted with substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocycle is optionally substituted with oxo;

each Q is independently selected from a bond, S, and O;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle; and

each R⁵ is independently selected from hydrogen or C₁-C₆ alkyl.

[00192] In some embodiments, for a compound or salt of Formula (III), R¹ is selected from C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -C(O)N(R²⁰)₂, -C(O)NR²⁰-OR²⁰, -S(O)₂N(R²⁰)₂, -NR²⁰S(O)₂R²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =NO(R²⁰), -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

[00193] In some embodiments, for a compound or salt of Formula (III), R² is selected from -L-NR²¹S(O)₂(R²¹) and -L-S(O)₂N(R²¹)₂.

[00194] In some embodiments, for a compound or salt of Formula (III), R² is selected from -L-N(R²¹)C(O)(OR²¹), and -L-OC(O)N(R²¹)₂.

[00195] In some embodiments, for a compound or salt of Formula (III), each R²¹ is independently selected from hydrogen; C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, and oxo. In some cases, each R²¹ is independently selected from hydrogen; C₁₋₆ alkyl,

C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle. In some cases, each R²¹ is independently selected from hydrogen and C₁₋₆ alkyl.

[00196] In some embodiments, for a compound or salt of Formula (III), R² is selected from L-bicyclic heterocycle, wherein the bicyclic heterocycle is optionally substituted with one or more R⁶.

[00197] In some embodiments, for a compound or salt of Formula (III), R² is selected from L-pyrrolizine, wherein the pyrrolizine is optionally substituted with one or more R⁶.

[00198] In some embodiments, for a compound or salt of Formula (III), B is an optionally substituted 5- to 15-membered heterocycle or optionally substituted C₃-C₁₅ carbocycle. In some cases, B is an optionally substituted 5- to 15-membered heterocycle. In some cases, B is an optionally substituted 8- to 15-membered heterocycle. In some cases, B is an optionally substituted C₃-C₁₅ carbocycle. In some cases, B is an optionally substituted C₈-C₁₅ carbocycle.

[00199] In some embodiments, for a compound or salt of Formula (III), B is an optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle. In some cases, B is an optionally substituted 8- to 15-membered fused heterocycle. In some cases, B is an optionally substituted C₈-C₁₅ fused carbocycle.


[00200] In some embodiments, for a compound or salt of Formula (III), for B, the optionally substituted 8- to 15-membered heterocycle contains at least one nitrogen atom. In some cases, the optionally substituted 8- to 15-membered heterocycle contains at least one sulfur atom. In some cases, the optionally substituted 8- to 15-membered heterocycle contains at most one nitrogen atom. In some cases, the optionally substituted 8- to 15-membered heterocycle contains at most one sulfur atom. In some cases, the optionally substituted 8- to 15-membered heterocycle contains at least two heteroatoms.

[00201] In some embodiments, for a compound or salt of Formula (III), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle are each bicyclic or tricyclic. In some cases, for B, the optionally substituted 8- to 15-membered fused heterocycle are each bicyclic or tricyclic. In some cases, for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused carbocycle are each bicyclic or tricyclic.


[00202] In some embodiments, for a compound or salt of Formula (III), B the heterocycle or carbocycle are each independently bicyclic. In some cases, the heterocycle is bicyclic. In some cases, the carbocycle is bicyclic.

[00203] In some embodiments, for a compound or salt of Formula (III), B the heterocycle or carbocycle are each independently tricyclic. In some cases, the heterocycle is tricyclic. In some cases, the carbocycle is tricyclic.


[00204] In some embodiments, for a compound or salt of Formula (III), B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused



[00205] In some embodiments, for a compound or salt of Formula (III), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused




[00206] In some embodiments, for a compound or salt of Formula (III), for B, the optionally substituted 8- to 15-membered fused heterocycle or optionally substituted C₈-C₁₅ fused



[00207] In some embodiments, for a compound or salt of Formula (III), for B, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, C₁-C₃ alkyl, -B(OR²⁰)₂, -OR²⁰, -C(O)N(R²⁰)₂, -N(R²⁰)₂, =O, -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OR²⁰)₂, -OH, -C(O)N(R²⁰)₂, =O, -CN, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OH)₂, -OH, -C(O)NH₂, -NH₂, =O, -CN, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₂₋₆ alkynyl. In some cases,


the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from oxo, -NH₂, -CN, halogen, C₁-C₃ alkyl. In some cases, the one or more optional substituents of the heterocycle or carbocycle are independently selected from oxo, -NH₂, halogen, C₁-C₃ alkyl.

[00208] In some embodiments, for a compound or salt of Formula (III), B is selected from



[00209] In some embodiments, for a compound or salt of Formula (III), for B, the one or more optional substituents of the heterocycle or carbocycle are independently selected from oxo, -NH₂, halogen, C₁-C₃ alkyl.

[00210] In some embodiments, for a compound or salt of Formula (III), B is selected from



[00211] In some embodiments, for a compound or salt of Formula (III), each R^4 is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, oxo, hydroxyl, halogen. In some cases, each R^4 is independently selected from C_{1-6} alkyl, oxo, and halogen.

[00212] In some embodiments, for a compound or salt of Formula (III), n is selected from 1 and 2. In some cases, n is 0.

[00213] In some embodiments, for a compound or salt of Formula (III), Y is O.

[00214] In some embodiments, for a compound or salt of Formula (III), R^1 is selected from optionally substituted 5- to 12-membered heterocycle.

[00215] In some embodiments, for a compound or salt of Formula (III), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, SR^{20} , $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-NR^{20}S(O)_2R^{20}$, $C(O)N(R^{20})_2$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=NO(R^{20})$, CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

[00216] In some embodiments, for a compound or salt of Formula (III), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, SR^{20} , $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-NR^{20}S(O)_2R^{20}$, $C(O)N(R^{20})_2$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=NO(R^{20})$, CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

[00217] In some embodiments, for a compound or salt of Formula (III), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-NR^{20}S(O)_2R^{20}$, $-N(R^{20})C(O)OR^{20}$, and $-OC(O)N(R^{20})_2$.

[00218] In some embodiments, for a compound or salt of Formula (III), R¹ is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, =O, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.

[00219] In some embodiments, for a compound or salt of Formula (III), R¹ is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.


[00220] In some embodiments, for a compound or salt of Formula (III), R²⁰ of R¹ is selected from hydrogen and C₁₋₃ alkyl.

[00221] In some embodiments, for a compound or salt of Formula (III), the 5- to 12-membered heterocycle of R¹ is an unsaturated heterocycle.

[00222] In some embodiments, for a compound or salt of Formula (III), the 5- to 12-membered heterocycle of R¹ is a saturated heterocycle.


[00223] In some embodiments, for a compound or salt of Formula (III), 5- to 12-membered heterocycle of R¹ is a bridged heterocycle.

[00224] In some embodiments, for a compound or salt of Formula (III), R¹ is selected from




independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, =O, -CN, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.

[00225] In some embodiments, for a compound or salt of Formula (III), R^1 is selected from




, each of which is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, $=O$, $-CN$, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl.


[00226] In some embodiments, for a compound or salt of Formula (III), R^1 is selected from




[00227] In some embodiments, for a compound or salt of Formula (III), R^1 is selected from an optionally substituted saturated 6- to 7-membered heterocycle. In some cases, R^1 is selected from an optionally substituted saturated 6-membered heterocycle. In some cases, R^1 is selected from




, which is optionally substituted. In some cases, the optional one or more substituents are independently selected from halogen, $-CN$, $-NHCN$, C_{1-6} cyanoalkyl, and C_{1-6} alkyl. In some cases, the optional one or more substituents are independently selected from $-CN$, $-NHCN$, C_{1-6} cyanoalkyl, and C_{1-6} alkyl. In some cases, the optional one or more substituents are independently selected from $-CN$, $-NHCN$, C_{1-6} cyanoalkyl, and C_{1-6} alkyl. In some cases, the optional one or more substituents are independently selected from $-NHCN$, and C_{1-6} alkyl. In



some cases, R^1 is selected from , which is substituted with one or more substituents selected






from $-NHCN$, and C_{1-6} alkyl. In some cases, R^1 is selected from , and








[00228] In some embodiments, for a compound or salt of Formula (III), R^1 is selected from a substituted saturated 6-membered heterocycle, wherein the saturated 6-membered heterocycle is substituted with at least one $-NHCN$, and optionally one or more C_{1-6} alkyl; M is O ; n is 0; B is

selected from an optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C₈-C₁₅ fused carbocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, oxo, -NH₂, C₁-C₃ alkyl, -OH, -C(O)NH₂, -NH₂, =O, -CN, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₂₋₆ alkynyl; Y is O; R² is selected from -L-bicyclic heterocycle, wherein the bicyclic heterocycle portion is optionally substituted with one or more substituents independently selected from halogen, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ hydroxyalkyl, or -N(R⁵)₂; and L is selected from C₁-C₄ alkylene. In some

cases, R¹ is selected from , and . In some cases, B is selected from



. In some cases, B is selected from , , and . In

some cases, B is .

[00229] In some embodiments, for a compound or salt of Formula (III), L is selected from C₁-C₄ alkylene.

[00230] In some embodiments, for a compound or salt of Formula (III), each L is independently selected from an optionally substituted C₁-C₄ alkylene; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle, wherein the C₃-C₆ carbocycle is optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl. In some cases, the optional substituents of L are selected from C₁-C₄ hydroxyalkyl, C₁-C₄ alkyl, C₃-C₆ carbocycle; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle or 3- to 8-membered heterocycle wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen and C₁₋₆ haloalkyl.



[00231] In some embodiments, for a compound or salt of Formula (III), each L is independently selected from a substituted C₁-C₄ alkylene, wherein two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle. In some cases, the C₃-C₆ carbocycle is optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl.

[00232] In some embodiments, for a compound or salt of Formula (III), wherein each L is independently selected from a substituted C₁-C₄ alkylene, and wherein two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle. In some cases, each L is independently selected from a substituted C₃ alkylene, and wherein two substituents on the same carbon atom of L come together to form a C₃ carbocycle. In some cases, each L is independently



selected from .

[00233] In some embodiments, for a compound or salt of Formula (III), R² is selected from -L-heterocycle, wherein the heterocycle portion of -L-heterocycle is optionally substituted with one or more R⁶. In some cases, the heterocycle is a saturated heterocycle. In some cases, the heterocycle has at least one nitrogen atom and at least one sulfur atom. In some cases, the heterocycle has at least one nitrogen atom. In some cases, the heterocycle has at least one sulfur atom.




[00234] In some embodiments, for a compound or salt of Formula (III), R² is selected from

 and , wherein the heterocycle portion is optionally substituted with one or more R⁶.






[00235] In some embodiments, for a compound or salt of Formula (III), Y-R² is selected from


 and , wherein the heterocycle portion is optionally substituted with one or more R⁶.

[00236] In some embodiments, for a compound or salt of Formula (III), Y-R² is selected from

,  and , wherein the heterocycle portion is optionally substituted with one or more R⁶.




[00237] In some embodiments, for a compound or salt of Formula (III), Y-R² is selected from




, , , ,  and









, wherein the heterocycle portion is optionally substituted with one or more R^6 .

[00238] In some embodiments, for a compound or salt of Formula (III), R^2 is selected from -L-saturated heterocycle, wherein the saturated heterocycle portion of the -L-saturated heterocycle is optionally substituted with one or more R^6 , and contains one nitrogen atom and one sulfur atom.

In some cases, $Y-R^2$ is selected from , , and , wherein the heterocycle portion is optionally substituted with one or more R^6 . In some cases, $Y-R^2$ is

selected from , , and , wherein the heterocycle portion is optionally substituted with one or more substituents selected from C_1 - C_3 alkyl and oxo. In some

cases, $Y-R^2$ is selected from , , and . In some

cases, $Y-R^2$ is selected from , , and .

[00239] In some embodiments, for a compound or salt of Formula (III), each R^6 is independently selected from halogen, -OH, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, -CN, C_1 - C_3 aminoalkyl, -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C_1 - C_3 alkyl substituted pyrazolyl, -N(R^5)₂, (C_1 - C_3 alkoxy) C_1 - C_3 alkyl-, (C_1 - C_3 alkyl)C(=O), oxo, (C_1 - C_3 haloalkyl)C(=O)-, -SO₂F, (C_1 - C_3 alkoxy) C_1 - C_3 alkoxy, -CH₂OC(O)N(R^5)₂, -CH₂NHC(O)OC₁-C₆ alkyl, -CH₂NHC(O)N(R^5)₂, -CH₂NHC(O) C_1 -C₆ alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂ C_1 -C₆ alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R^5)₂, -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl), -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl)phenyl(C_1 - C_3 alkyl)N(CH₃)₂, -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl)phenyl, -OC(O)heterocycle, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C_1 - C_3 alkyl)(C_1 - C_3 alkyl)phenyl are each optionally substituted with -C(O)H and OH, and wherein the heterocycle of -CH₂heterocycle is optionally substituted with oxo.


[00240] In some embodiments, for a compound or salt of Formula (III), each R^6 is independently selected from halogen, -OH, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, -CN, and C_1 - C_3 aminoalkyl.

[00241] In some embodiments, for a compound or salt of Formula (III), each R^6 is independently selected from halogen, -OH, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, C_1 - C_3 aminoalkyl, C_1 - C_3 haloalkyl,


C₁-C₃ alkoxy, -N(R⁵)₂, and oxo. In some cases, each R⁶ is independently selected from -OH, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ aminoalkyl, C₁-C₃ alkoxy, and -N(R⁵)₂. In some cases, each R⁶ is independently selected from C₁-C₃ alkyl, C₁-C₃ alkoxy, and -N(R⁵)₂.

[00242] In some embodiments, for a compound or salt of Formula (III), R⁶ is selected from halogen, -OH, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, -CN, and C₁-C₃ aminoalkyl. In some cases, R⁶ is selected from halogen and C₁-C₃ alkyl. In some cases, R⁶ is halogen. In some cases, R⁶ is C₁-C₃ alkyl. In some cases, R⁶ is selected from halogen and C₁-C₃ alkyl. In some cases, R⁶ is selected from methyl and fluorine.


[00243] In some embodiments, for a compound or salt of Formula (III), R² is selected from




[00244] In some embodiments, for a compound or salt of Formula (III), Y-R² is selected from



[00245] In some embodiments, for a compound or salt of Formula (III), Y-R² is selected from




[00246] In some embodiments, for a compound or salt of Formula (III), Y-R² is



[00247] In some embodiments, for a compound or salt of Formula (III), L is selected from unsubstituted C₁-C₄ alkylene.

[00248] In some embodiments, for a compound or salt of Formula (III), Y-R² is selected from




, wherein the heterocycle portion is optionally substituted with one or more R⁶.

[00249] In some embodiments, for a compound or salt of Formula (III), R⁶ of R² is independently selected at each occurrence from halogen, hydroxy, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, cyano, and C₁-C₃ aminoalkyl.

[00250] In some embodiments, for a compound or salt of Formula (III), R⁶ of R² is independently selected at each occurrence from C₁-C₃ alkyl and halogen.

[00251] In some embodiments, for a compound or salt of Formula (III), Y-R² is selected from



[00252] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the carbocycle of R¹ is selected from C₃-C₁₂ carbocycle, C₃-C₁₀ carbocycle, C₃-C₉ carbocycle, C₃-C₈ carbocycle, or C₃-C₆ carbocycle. In some cases, the carbocycle of R¹ is selected from C₃-C₁₂ carbocycle, C₄-C₁₂ carbocycle, C₅-C₁₂ carbocycle, C₆-C₁₂ carbocycle, C₇-C₁₂ carbocycle, C₈-C₁₂ carbocycle, or C₉-C₁₂ carbocycle.



[00253] In some embodiments, for a compound of Formula (I), the heterocycle of R¹ is a 5- to 12-membered heterocycle, 6- to 12-membered heterocycle, 7- to 12-membered heterocycle, or 8- to 12-membered heterocycle. In some cases, the heterocycle of R¹ is a 5- to 11-membered heterocycle, 5- to 10-membered heterocycle, 5- to 9-membered heterocycle, or 5- to 8-membered heterocycle. In some cases, the heterocycle of R¹ is a 6- to 11-membered heterocycle, 6- to 10-membered heterocycle, 6- to 9-membered heterocycle, or 6- to 8-membered heterocycle. In some cases, the heterocycle of R¹ is a 7- to 11-membered heterocycle, 7- to 10-membered heterocycle, 7- to 9-membered heterocycle, or 7- to 8-membered heterocycle. In some cases, the heterocycle of R¹ is a 5- to 6-membered heterocycle or 5- to 9-membered heterocycle. In some cases, the heterocycle of R¹ is an 8- to 9-membered heterocycle. In some cases, the heterocycle of R¹ is saturated. The heterocycle may be optionally substituted as described elsewhere herein.

[00254] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the heterocycle of R¹ is a 5- to 12-membered monocyclic heterocycle, 6- to 12-membered monocyclic heterocycle, 7- to 12-membered monocyclic heterocycle, or 8- to 12-membered monocyclic heterocycle. In some cases, the heterocycle of R¹ is a 5- to 11-membered monocyclic heterocycle, 5- to 10-membered monocyclic heterocycle, 5- to 9-membered monocyclic heterocycle, or 5- to 8-membered monocyclic heterocycle. In some cases, the heterocycle of R¹ is a 6- to 11-membered monocyclic heterocycle, 6- to 10-membered monocyclic heterocycle, 6- to 9-membered monocyclic heterocycle, or 6- to 8-membered monocyclic heterocycle. In some cases, the heterocycle of R¹ is a monocyclic 7- to 11-membered heterocycle, 7- to 10-membered monocyclic heterocycle, 7- to 9-membered monocyclic heterocycle, or 7- to 8-membered monocyclic heterocycle. In some cases, the heterocycle of R¹ is a 5- to 6-membered monocyclic heterocycle or 5- to 9-membered monocyclic heterocycle. In some cases, the heterocycle of R¹ is an 8- to 9-membered monocyclic heterocycle. In some cases, the heterocycle of R¹ is saturated. The monocyclic heterocycle may be optionally substituted as described elsewhere herein.

[00255] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the heterocycle of R^1 is a 5- to 12-membered bridged heterocycle, 6- to 12-membered bridged heterocycle, 7- to 12-membered bridged heterocycle, or 8- to 12-membered bridged heterocycle. In some cases, the heterocycle of R^1 is a 5- to 11-membered bridged heterocycle, 5- to 10-membered bridged heterocycle, 5- to 9-membered bridged heterocycle, or 5- to 8-membered bridged heterocycle. In some cases, the heterocycle of R^1 is a 6- to 11-membered bridged heterocycle, 6- to 10-membered bridged heterocycle, 6- to 9-membered bridged heterocycle, or 6- to 8-membered bridged heterocycle. In some cases, the heterocycle of R^1 is a bridged 7- to 11-membered heterocycle, 7- to 10-membered bridged heterocycle, 7- to 9-membered bridged heterocycle, or 7- to 8-membered bridged heterocycle. In some cases, the heterocycle of R^1 is a 5- to 6-membered bridged heterocycle or 5- to 9-membered bridged heterocycle. In some cases, the heterocycle of R^1 is an 8- to 9-membered bridged heterocycle. In some embodiments, the heterocycle of R^1 is saturated. The bridged heterocycle may be optionally substituted as described elsewhere herein.

[00256] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is a 5- to 9-membered heterocycle, the 5- to 9-membered heterocycle contains at most 1 nitrogen atom. In some embodiments, R^1 is selected from optionally substituted 5- to 9-membered heterocycle, each of which is optionally substituted.


[00257] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the heterocycle of R^1 contains at most 1 nitrogen atom. In some embodiments, the heterocycle of R^1 contains at most 1 heteroatom atom. In some embodiments, the heterocycle of R^1 contains at most 2 heteroatom atoms. In some cases, the heteroatom is selected from nitrogen, oxygen, and sulfur. In some cases, the heterocycle is a monocyclic heterocycle or a bridged heterocycle. In some cases, the heterocycle is a monocyclic heterocycle. In some cases, the

heterocycle is a bridged heterocycle. In some cases, the heterocycle is selected from  and . In some cases, the heterocycle is selected from , , , , and . In some cases, the bridged heterocycle is selected from , , and .

The heterocycle may be optionally substituted as described elsewhere herein.

[00258] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the spiroheterocycle of R^1 contains at most 1 nitrogen atom. In some embodiments, the spiroheterocycle of R^1 contains at most 2 heteroatom atoms. In some embodiments, the

spiroheterocycle of R^1 contains at most 3 heteroatom atoms. In some embodiments, the spiroheterocycle of R^1 contains at most 1 heteroatom atom. In some cases, the spiroheterocycle of R^1 contains at least 2 heteroatom atoms. In some cases, the spiroheterocycle of R^1 contains at least 3 heteroatom atoms. In some cases, the spiroheterocycle of R^1 contains at least 4 heteroatom atoms. In some cases, the spiroheterocycle of R^1 contains at least 2 nitrogen atoms. In some embodiments, the spiroheterocycle of R^1 contains at most 1 heteroatom atom. In some cases, the spiroheterocycle of R^1 contains at most 1 sulfur atom. In some cases, the heteroatom is selected from nitrogen, oxygen, and sulfur. In some embodiments, the spiroheterocycle of R^1 is



spiroheterocycle may be optionally substituted as described elsewhere herein.


[00259] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from optionally substituted 7- to 8-membered spiroheterocycle. In some cases, R^1 is selected from optionally substituted 7-membered spiroheterocycle. In some cases, R^1 is selected from optionally substituted 8-membered spiroheterocycle.


[00260] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the fused heterocycle of R^1 is a 6- to 12-membered fused heterocycle, 6- to 12-membered fused heterocycle, 7- to 12-membered fused heterocycle, or 8- to 12-membered fused heterocycle. In some cases, the fused heterocycle of R^1 is a 6- to 11-membered fused heterocycle, 6- to 10-membered fused heterocycle, 6- to 9-membered fused heterocycle, or 6- to 8-membered fused heterocycle. In some cases, the fused heterocycle of R^1 is a 7- to 11-membered fused heterocycle, 7- to 10-membered fused heterocycle, 7- to 9-membered fused heterocycle, or 7- to 8-membered fused heterocycle. In some cases, the fused heterocycle of R^1 is an 8- to 11-membered fused heterocycle. In some cases, the fused heterocycle of R^1 is a 6-

membered fused heterocycle. The fused heterocycle may be optionally substituted as described elsewhere herein.

[00261] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the fused heterocycle of R^1 is selected from a 6-, 9-, 10-, 11-, and 12-membered fused heterocycle. In some cases, the fused heterocycle of R^1 is selected from a 9- to 12-membered fused heterocycle. In some cases, the fused heterocycle of R^1 is selected from a 10- to 12-membered fused heterocycle. The fused heterocycle is optionally substituted with one or more substituents independently selected from halogen, $-OR^{20}$, $-N(R^{20})_2$, $-NO_2$, $=O$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. The fused heterocycle is optionally substituted with one or more substituents independently selected from halogen, $-OR^{20}$, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl.

[00262] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the fused heterocycle of R^1 contains at most 1 nitrogen atom. In some embodiments, the fused heterocycle of R^1 contains at most 1 heteroatom atom. In some cases, the heteroatom is



selected from nitrogen, oxygen, and sulfur. In some cases, the fused heterocycle is . The fused heterocycle may be optionally substituted as described elsewhere herein.

[00263] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from C_6 - C_7 carbocycle, 5- to 10-membered heterocycle, 7- to 8-membered spiroheterocycle, and 6-, 9-, 10-, 11-, and 12-membered fused heterocycle, each of which is optionally substituted.

[00264] In some embodiments, for a compound of Formula (I), R^1 is selected from C_6 - C_7 carbocycle, 5- to 10-membered heterocycle, 7- to 8-membered spiroheterocycle, and 6-, 8- to 12-membered fused heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, $-OR^{20}$, $-N(R^{20})_2$, $-NO_2$, $=O$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl.

[00265] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from C_6 - C_7 carbocycle, 5- to 10-membered heterocycle, 7- to 8-membered spiroheterocycle, and 6-, 8- to 12-membered fused heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, $-OR^{20}$, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl.

[00266] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), for R^1 , R^{20} of $-OR^{20}$ and $-N(R^{20})_2$, is selected hydrogen and C_{1-6} alkyl.

[00267] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from C_6 - C_7 carbocycle and 5- to 10-membered heterocycle, each of which is optionally substituted. In some cases, the heterocycle contains at most 1 nitrogen atom. In some cases, R^1 is selected from C_6 - C_7 carbocycle, each of which is optionally substituted.

[00268] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the one or more optional substituents of R^1 are independently selected from halogen, $-OR^{20}$, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, $-N(R^{20})C(O)N(R^{20})_2$, and C_{1-6} alkyl.

[00269] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), one or more optional substituents of R^1 are independently selected from halogen, $-OR^{20}$, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, the one or more optional substituents of R^1 are independently selected from $-OR^{20}$, $-N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{1-6} alkyl. In some cases, the one or more optional substituents of R^1 are independently selected from $-OR^{20}$, $-N(R^{20})_2$, C_{1-6} aminoalkyl, and C_{1-6} hydroxyalkyl. In some cases, the one or more optional substituents of R^1 are independently selected from $-OR^{20}$, $-N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkyl, and C_{1-6} hydroxyalkyl. In some cases, the one or more optional substituents of R^1 are independently selected from $-OR^{20}$, $-N(R^{20})_2$, and C_{1-6} alkyl. In some cases, the one or more optional substituents of R^1 are independently selected from $-N(R^{20})C(O)N(R^{20})_2$.


[00270] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle contains at most 1 nitrogen atom and optionally one or more additional heteroatoms selected from oxygen, boron, and sulfur; or R^1 is further selected from 7-, 8-, 10, 11-membered spiro heterocycle and 6-, 9-, 10-, 11-, and 12-membered fused heterocycle wherein the C_3 - C_{12} carbocycle, 5- to 12-membered heterocycle, 7-, 8-, 10-, 11-membered spiro heterocycle, and 6-, 9-, 10-, 11-, and 12-membered fused heterocycle, are each optionally substituted with one or more substituents independently selected from halogen, $-OR^{20}$, $-SR^{20}$, $-C(O)N(R^{20})_2$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-NO_2$, $=O$, $=N(R^{20})$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

[00271] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle contains at most 1 nitrogen atom and optionally one or more additional heteroatoms selected from oxygen, boron, and sulfur; or R^1 is further selected from 7-, 8-, 10, 11-membered spiro heterocycle and 6-, 9-, 10-, 11-, and 12-membered fused heterocycle


wherein the C₃-C₁₂ carbocycle, 5- to 12-membered heterocycle, 7-, 8-, 10-, 11-membered spiro heterocycle, and 6-, 9-, 10-, 11-, and 12-membered fused heterocycle, are each optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -N(R²⁰)S(O)₂(R²⁰), -C(O)N(R²⁰)₂, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -NO₂, =O, =S, =N(R²⁰), -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.


[00272] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R¹ is selected from 5- to 10-membered heterocycle, 7-, 8-, 10-, 11-membered spiro heterocycle, and 6-, 9-, 10-, 11-, and 12-membered fused heterocycle, and wherein each are optionally substituted with one or more substituents independently selected from halogen, -N(R²⁰)₂, C₁₋₆ alkyl, -OR²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -B(OR²⁰)₂, -N(R²⁰)C(O)N(R²⁰)₂, =O, C₁₋₆ hydroxyalkyl, halogen, -N(R²⁰)C(O)R²⁰, -N(R²⁰)S(O)₂(R²⁰), and C₁₋₆ aminoalkyl.

[00273] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R¹ is selected from 5- to 10-membered heterocycle, 7-, 8-, 10-, 11-membered spiro heterocycle, and 6-, 9-, 10-, 11-, and 12-membered fused heterocycle, and wherein each are optionally substituted with one or more substituents independently selected from halogen, -N(R²⁰)₂, C₁₋₆ alkyl, -OR²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -B(OR²⁰)₂, C₁₋₆ cyanoalkyl, -N(R²⁰)C(O)N(R²⁰)₂, =O, C₁₋₆ hydroxyalkyl, halogen, -N(R²⁰)C(O)R²⁰, -N(R²⁰)S(O)₂(R²⁰), and C₁₋₆ aminoalkyl. In






substituted with one or more substituents independently selected from halogen, -N(R²⁰)₂, C₁₋₆ alkyl, -OR²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -B(OR²⁰)₂, C₁₋₆ cyanoalkyl, -N(R²⁰)C(O)N(R²⁰)₂, =O, C₁₋₆ hydroxyalkyl, halogen, -N(R²⁰)C(O)R²⁰, -N(R²⁰)S(O)₂(R²⁰), and C₁₋₆ aminoalkyl. In some cases,

















[00274] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from 5- to 10-membered heterocycle, wherein the 5- to 10-membered

heterocycle is optionally substituted with one or more substituents independently selected from -OR²⁰, -N(R²⁰)₂, C₁₋₆ alkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ aminoalkyl, -N(R²⁰)C(O)N(R²⁰)₂, -






N(R²⁰)C(O)R²⁰, and -B(OR²⁰)₂. In some cases, R¹ is selected from , , , ,

, and , each of which is optionally substituted with one or more substituents independently selected from -OR²⁰, -N(R²⁰)₂, C₁₋₆ alkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ aminoalkyl, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)R²⁰, and -B(OR²⁰)₂.


[00275] In some embodiments, for a compound of Formula (I), Formula (II), or Formula (III),

R¹ is selected from , , , , , , , , , , , , , and .


[00276] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R¹ is selected from 5- to 10-membered heterocycle, wherein the 5- to 10-membered heterocycle is optionally substituted with one or more substituents independently selected from -N(R²⁰)₂, -OR²⁰, and C₁₋₆ alkyl. In some cases, the 5- to 10-membered heterocycle is optionally substituted with one or more substituents independently selected from -OR²⁰, and C₁₋₆ alkyl. In some embodiments, for a compound of Formula (I), Formula (II), or Formula (III), R¹ is

selected from  and , each of which is optionally substituted with one or more substituents independently selected from -OR²⁰, and C₁₋₆ alkyl. In some cases, R¹ is selected from , , and .


[00277] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula






[00278] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula








[00279] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula


















[00280] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula

(III), R^1 is selected from , , and , each of which are optionally substituted.


[00281] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula

(III), R^1 is selected from , , , , and , each of which are optionally substituted.





[00282] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula


(III), R^1 is selected from , , , , , , , , , , , , , , and , each of which are optionally substituted.

[00283] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula


(III), R^1 is , which is optionally substituted.


[00284] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula

(III), R^1 is selected from , , , and , each of which are optionally substituted.




[00285] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula





[00286] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), wherein the 5- to 12-membered heterocycle of R^1 is unsaturated and a bridged heterocycle. In some cases, R^1 is selected from an optionally substituted 7- to 8-membered unsaturated and



bridged heterocycle. In some cases, R^1 is selected from

[00287] In some embodiments, for a compound of Formula (I) or Formula (II), R^1 is selected from 5- to 10-membered heterocycle, 7-, 8-, 10-, 11-membered spiro heterocycle, and 6-, 9-, 10-, 11-, and 12-membered fused heterocycle, and wherein each are optionally substituted with one or more substituents independently selected from halogen, $-N(R^{20})_2$, C_{1-6} alkyl, $-OR^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-B(OR^{20})_2$, C_{1-6} cyanoalkyl, $-N(R^{20})C(O)N(R^{20})_2$, $=O$, C_{1-6} hydroxyalkyl, halogen, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})S(O)_2(R^{20})$, and C_{1-6} aminoalkyl; R^3 is naphthalene, wherein naphthalene is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-NH_2$, $-NO_2$, $=O$, C_{1-6} alkyl, C_{2-6} alkynyl, C_{1-6} aminoalkyl, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl; R^4 is selected from hydrogen, halogen or C_1-C_3 alkyl; Y is O ; L is independently a C_1-C_4 alkylene; R^2 is selected from $-L$ -heterocycle, wherein the heterocycle portion is optionally substituted with one or more substituents independently selected from halogen, hydroxy, C_1-C_4 alkyl, C_1-C_4 alkoxy, C_1-C_4 haloalkyl, C_1-C_4 hydroxyalkyl, or $-N(R^5)_2$. In some cases, $Y-R^2$ is



selected from




and


, wherein the heterocycle portion is optionally substituted.




[00288] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from an optionally substituted 5- to 12-membered unsaturated heterocycle, wherein the heterocycle has as most one nitrogen atom. In some cases, the 5- to 12-membered unsaturated heterocycle has at least one nitrogen atom. In some cases, the 5- to 12-membered unsaturated heterocycle has at most one nitrogen atom.


[00289] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from 6- to 7-membered heterocycle. In some cases, R^1 is selected from 7-membered heterocycle. In some cases, R^1 is selected from 6-membered heterocycle. In some cases, the 6- to 7-membered heterocycle contains only 1 nitrogen atom and optionally one or more additional heteroatoms selected from oxygen, and sulfur. In some cases, the optionally one or more additional heteroatoms are selected from sulfur. In some cases, the optionally one or more additional heteroatoms are selected from oxygen. In some cases, the 6- to 7-membered heterocycle contains only 1 nitrogen atom and no further additional heteroatoms. In some cases, the 6- to 7-membered heterocycle is a non-aromatic 6- to 7-membered heterocycle. In some cases, the 6- to 7-membered heterocycle of R^1 is bound to Formula (I) via the only 1 nitrogen atom. In some cases, the 6- to 7-membered heterocycle of R^1 is bound to Formula (II) via the only 1 nitrogen atom. In some cases, the 6- to 7-membered heterocycle of R^1 is bound to Formula (III) via the only 1







substituted. In some cases, R¹ is selected from 
 each of which is optionally substituted. In some cases, the one or more optional substituents of R¹ are each independently selected from fluorine, -OH, -C(O)NH₂, -NH-C(O)-(C₁₋₆ alkoxy), -NH-C(O)-(C₁₋₆ hydroxyalkyl), -NH₂, -NH(CN), =O, -CN, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ alkyl, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of R¹ are each independently selected from halogen, -OH, -CN, C₁₋₆ cyanoalkyl, C₁₋₆ alkyl, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of R¹ are each independently selected from halogen, -OH, and -CN. In some cases, the one or more optional substituents of R¹ are each independently selected from fluorine, -OH, -CN, C₁₋₆ cyanoalkyl, C₁₋₆ alkyl, oxo, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of R¹ are each independently selected from fluorine, -OH, -CN, C₁₋₆







R¹ is selected from









some cases, R^1 is selected from an optionally substituted unsaturated 6-membered heterocycle. In some cases, R^1 is selected from an optionally substituted unsaturated 7-membered heterocycle. In some cases, the heterocycle has 1 or 2 double bonds. In some cases, the heterocycle has only 1 double bond. In some cases, the heterocycle has only 2 double bonds. In some cases, R^1 is selected


from , wherein each is optionally substituted with one or more substituents independently selected from halogen, -OH, -NH₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl, and C₁₋₆ alkyl. In some cases, R^1 is




selected from , wherein each is optionally substituted with one or more substituents independently selected from halogen, -OH, -NH₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl, and C₁₋₆ alkyl. In some cases, R^1 is

selected from , wherein each is optionally substituted with one or more substituents independently selected from halogen, -OH, -NH₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl, and C₁₋₆ alkyl. In some cases, R^1 is




selected from ,




, and . In some

cases, R^1 is selected from ,





[00291] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from an unsaturated 6- to 7-membered heterocycle, wherein the unsaturated 6- to 7-membered heterocycle is substituted with one or more substituents selected from halogen. In some cases, the unsaturated 6- to 7-membered heterocycle is substituted with at least one halogen. In some cases, the unsaturated 6- to 7-membered heterocycle is substituted with at only one halogen. In some cases, the unsaturated 7-membered heterocycle is substituted with one fluorine. In some cases, R^1 is selected from an unsaturated 6-membered heterocycle, substituted with at least one halogen. In some cases, R^1 is selected from an unsaturated 7-membered heterocycle,






[00292] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from an optionally substituted unsaturated 6- to 8-membered heterocycle. In some cases, R^1 is selected from an optionally substituted unsaturated 7-membered heterocycle.




[00293] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from an optionally substituted 6-membered heterocycle. In some cases, the 6-membered heterocycle contains only 1 nitrogen atom. In some cases, the 6-membered heterocycle of R^1 is bound to Formula (I) via the only 1 nitrogen atom. In some cases, the 6-membered heterocycle of R^1 is bound to Formula (II) via the only 1 nitrogen atom. In some cases, the 6-membered heterocycle of R^1 is bound to Formula (III) via the only 1 nitrogen atom. In some cases,




R^1 is selected from , and , any of which is optionally substituted. In some cases, the one or more optional substituents of R^1 are each independently selected from halogen, $-OR^{20}$, $-N(R^{20})_2$, $=O$, $-CN$, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} alkyl, and C_{2-6} alkynyl. In some cases, the one or more optional substituents of R^1 are each independently selected from fluorine, $-OH$, $-NH_2$, $-NH(CN)$, $=O$, $-CN$, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} alkyl, and C_{2-6} alkynyl. In some cases, the one or more optional substituents of R^1 are each independently selected from fluorine, $-OH$, $-NH_2$, $-NH(CN)$, $=O$, $-CN$, C_{1-6} hydroxyalkyl, C_{1-6} alkyl, and C_{2-6} alkynyl. In some cases, the 6-membered heterocycle is a partially unsaturated 6-membered heterocycle or a saturated 6-membered heterocycle. In some cases, the 6-membered heterocycle is partially unsaturated. In some cases, the 6-membered heterocycle is a saturated 6-membered heterocycle. In some cases, the 6-membered heterocycle is a monocyclic 6-membered heterocycle. In some cases, the 6-

membered heterocycle is not a bridged heterocycle. In some cases, R^1 is selected from




[00294] In some embodiments, for a compound of Formula (I), Formula (II), or Formula (III), R^1 is selected from an optionally substituted 6-membered unsaturated heterocycle and 6-membered saturated heterocycle.


[00295] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula




(III), R^1 is selected from  and , wherein each is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-NH_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl.




, each of which is optionally substituted with one or more substituents independently selected from halogen, =O, -OH, -CN, -NHCN, -C(O)N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, and C₁₋₆ alkyl. In some cases, each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle. In some cases, R¹ is selected from





[00301] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R¹ is selected from an optionally substituted 10-membered heterocycle. In some cases, the 10-membered heterocycle is a bicyclic heterocycle. In some cases, the 10-membered heterocycle is a spiro heterocycle. In some cases, the 10-membered heterocycle is a fused heterocycle. In some cases, the 10-membered heterocycle is a saturated heterocycle. In some cases, the 10-membered heterocycle is a non-aromatic heterocycle. In some cases, the 10-membered heterocycle contains at least 1 nitrogen atom. In some cases, the 10-membered heterocycle contains at least 2 nitrogen atoms. In some cases, the 10-membered heterocycle contains at least 3 nitrogen atoms. In some cases, the 10-membered heterocycle contains at least




1 sulfur atom. In some cases, R¹ is selected from , and , each of which is optionally substituted with one or more substituents independently selected from halogen, =O, -OH, -CN, -NHCN, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, C₁₋₆ aminoalkyl, C₁₋₆




hydroxyalkyl, C₁₋₆ cyanoalkyl, and C₁₋₆ alkyl. In some cases, R¹ is selected from




. In some cases, R¹ is selected from




, and




. In some cases, R¹ is selected from



and




. In



some cases, R¹ is selected from


. In some cases, R¹ is selected from



, which is optionally substituted with one or more substituents independently

selected from halogen, -OR²⁰, -SR²⁰, -N(R²⁰)₂, -NO₂, =O, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

[00302] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R¹ is selected from an optionally substituted unsaturated 9- to 11-membered heterocycle. In some cases, R¹ is selected from an optionally substituted unsaturated 10-membered heterocycle. In some cases, R¹ is selected from an optionally substituted unsaturated




10-membered fused heterocycle. In some cases, R^1 is , which is optionally substituted.

In some cases, the one or more optional substituents are selected from halogen, -

OH , $-C(O)N(R^{20})_2$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-NO_2$, $=O$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6}

hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, and C_{2-6} alkynyl. In some cases, R^1 is



, optionally substituted with one or more substituents selected from -

$N(R^{20})_2$, $-C(O)R^{20}$, $-NO_2$, $=O$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6}

cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, and C_{2-6} alkynyl. In some cases, each R^{20} is independently

selected from hydrogen; and C_{1-6} alkyl, and C_{3-12} carbocycle, and each of which is optionally

substituted with one or more substituents independently selected from halogen, $-OH$, $-CN$, $-NO_2$,

$-NH_2$, C_{1-10} alkyl, $-C_{1-10}$ haloalkyl, $-O-C_{1-10}$ alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-12} carbocycle,


and 3- to 12-membered heterocycle.

[00303] In some embodiments, for a compound or salt of Formula (I), Formula (II), or

Formula (III), R^1 is selected from a 7- to 11-membered spiro heterocycle. In some cases, R^1 is

selected from a 10-membered spiro heterocycle. In some cases, the spiro heterocycle has at least

3 nitrogen atoms. In some cases, the spiro heterocycle has at least 1 sulfur atom. In some cases,




R^1 is selected from , and , each of which is optionally substituted. In some


cases, the one or more optional substituents are independently selected from halogen, $-OH$, -

$N(R^{20})_2$, $-NO_2$, $=O$, $-CN$, $-NHCN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6}


cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl. In some cases, R^1 is selected



from



, and



. In some cases, R^1 is


. In some cases, R^1 is




. In some cases, M is selected from O , and NR^3 . In some cases, M is selected from

O. In some cases, M is selected from NR^3 . In some cases, R^3 is selected from C_{1-6} alkyl. In some cases, R^3 is selected from C_{1-2} alkyl. In some cases, R^3 is selected from methyl.


[00304] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from an optionally substituted 8- to 10-membered fused heterocycle. In some cases, the 8- to 10-membered fused heterocycle is a bicyclic heterocycle. In some cases, the 8- to 10-membered fused heterocycle is a saturated heterocycle. In some cases, the 8- to 10-membered fused heterocycle is an unsaturated heterocycle. In some cases, the 8- to 10-membered heterocycle is a non-aromatic heterocycle. In some cases, R^1 is selected from an optionally substituted 9-membered fused heterocycle. In some cases, R^1 is selected from an optionally substituted 10-membered fused heterocycle. In some cases, the 10-membered fused heterocycle is a bicyclic heterocycle. In some cases, the 10-membered fused heterocycle is a saturated heterocycle. In some cases, the 9-membered heterocycle is a non-aromatic heterocycle. In some cases, the 10-membered heterocycle is a non-aromatic heterocycle. In some cases, the fused heterocycle has one saturated ring and one aromatic ring. In some cases, the fused heterocycle has one saturated ring and one unsaturated ring. In some cases, the fused heterocycle has two saturated rings. In some cases, the 10-membered heterocycle contains at least 1 nitrogen atom. In some cases, the 10-membered heterocycle contains at least 2 nitrogen atoms. In some cases, the 10-membered heterocycle contains at least 3 nitrogen atoms. In some cases, the 9-membered heterocycle contains at least 1 nitrogen atom. In some cases, the 9-membered heterocycle contains at least 2 nitrogen atoms. In some cases, the 9-membered heterocycle



contains at least 3 nitrogen atoms. In some cases, R^1 is selected from  and , each of which is optionally substituted with one or more substituents. In some cases, R^1 is





, which is optionally substituted with one or more substituents. In some cases, R^1 is







, which is optionally substituted with one or more substituents. In some cases, the one or more optional substituents of R^1 are independently selected from halogen, $-\text{OH}$, $-\text{S}(\text{O})_2(\text{R}^{20})$, -


$\text{S(O)}_2\text{N(R}^{20})_2$, $-\text{S(O)N(R}^{20})_2$, $-\text{S(O)R}^{20}(=\text{NR}^{20})$, $-\text{C}(=\text{NR}^{20})\text{N(R}^{20})_2$, $-\text{C(O)N(R}^{20})_2$, $-\text{C(O)NHOR}^{20}$, $-\text{N(R}^{20})_2$, $-\text{C(O)R}^{20}$, $-\text{NO}_2$, $=\text{O}$, $-\text{CN}$, C_{1-6} alkyl- $\text{N(R}^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkynyl, and 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*} . In some cases, the one or more optional substituents of R^1 are independently selected from halogen, $-\text{OH}$, $-\text{S(O)}_2(\text{R}^{20})$, $-\text{S(O)}_2\text{N(R}^{20})_2$, $-\text{S(O)N(R}^{20})_2$, $-\text{S(O)R}^{20}(=\text{NR}^{20})$, $-\text{C(O)N(R}^{20})_2$, $-\text{C(O)NHOR}^{20}$, $-\text{N(R}^{20})_2$, $-\text{C(O)R}^{20}$, $-\text{NO}_2$, $=\text{O}$, $-\text{CN}$, C_{1-6} alkyl- $\text{N(R}^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, and C_{2-6} alkynyl. In some cases, the optional one or more substituents are independently selected from halogen, $=\text{O}$, $-\text{OH}$, $-\text{CN}$, $-\text{NHCN}$, $-\text{S(O)}_2(\text{R}^{20})$, $-\text{S(O)}_2\text{N(R}^{20})_2$, $-\text{S(O)N(R}^{20})_2$, $-\text{S(O)R}^{20}(=\text{NR}^{20})$, $-\text{C(O)R}^{20}$, $-\text{C(O)N(R}^{20})_2$, $-\text{C(O)NR}^{20}\text{OR}^{20}$, C_{1-6} alkyl- $\text{N(R}^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, and C_{1-6} alkyl. In some cases, the optional one or more substituents are independently selected from halogen, $=\text{O}$, C_{1-6} alkyl- $\text{N(R}^{20})_2$, $-\text{S(O)}_2(\text{R}^{20})$, $-\text{S(O)N(R}^{20})_2$, $-\text{S(O)R}^{20}(=\text{NR}^{20})$, $-\text{C(O)R}^{20}$, $-\text{C(O)N(R}^{20})_2$, and $-\text{C(O)NR}^{20}\text{OR}^{20}$. In some cases, the optional one or more substituents are independently selected from halogen, $=\text{O}$, $-\text{S(O)}_2(\text{R}^{20})$, $-\text{S(O)N(R}^{20})_2$, $-\text{S(O)R}^{20}(=\text{NR}^{20})$, $-\text{C(O)R}^{20}$, $-\text{C(O)N(R}^{20})_2$, and $-\text{C(O)NR}^{20}\text{OR}^{20}$. In some cases, the optional one or more substituents are independently selected from $-\text{C(O)R}^{20}$, $-\text{C(O)N(R}^{20})_2$, and $-\text{C(O)NR}^{20}\text{OR}^{20}$. In some cases, the optional one or more substituents are independently selected from $-\text{S(O)}_2(\text{R}^{20})$, $-\text{S(O)}_2\text{N(R}^{20})_2$, $-\text{S(O)N(R}^{20})_2$, $-\text{S(O)R}^{20}(=\text{NR}^{20})$. In some cases, the optional one or more substituents are independently selected from $-\text{S(O)N(R}^{20})_2$. In some cases, the optional one or more substituents are independently selected from $\text{S(O)}_2(\text{R}^{20})$. In some cases, the optional one or more substituents are independently selected from $\text{S(O)R}^{20}(=\text{NR}^{20})$. In some cases, the optional one or more substituents are independently selected from $-\text{C(O)R}^{20}$. In some cases, the optional one or more substituents are independently selected from $-\text{C(O)N(R}^{20})_2$. In some cases, the optional one or more substituents are independently selected from $-\text{C(O)NR}^{20}\text{OR}^{20}$. In some cases, R^1 is selected from






each of which is further optionally substituted. In some cases, the further one or more optional substituents are selected from halogen, -OH, =O, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, and C₂₋₆ alkynyl. In some cases, the further one or more optional substituents are selected from halogen, -CN, C₂ alkenyl, and C₁₋₆ alkyl. In some cases, the further one or more optional substituents are selected from halogen, and C₁₋₆ alkyl. In some cases, the further one or more optional substituents are selected from halogen. In some cases, each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle. In some cases, each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, and 3- to 12-membered heterocycle. In some cases, each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, and 3- to 12-membered saturated heterocycle. In some cases, each R²⁰ is independently selected from 5- to 6-membered saturated heterocycle. In some cases, the heterocycle of R²⁰ has at least one nitrogen atom. In some cases, the heterocycle of R²⁰ has at least one sulfur atom. In some cases, the heterocycle of R²⁰ has at least one oxygen atom. In some cases, the heterocycle of R²⁰ contains only 1 heteroatom. In some cases, the heterocycle of R²⁰ has at least two heteroatoms. In some cases, the heterocycle of R²⁰ contains only 2 heteroatoms. In some cases, the optional one or more substituents of R¹ are












independently selected from halogen, -CN, C₂ alkenyl, , , , ,



In some cases, the optional one or more substituents of R¹ are independently selected from halogen,



In some cases, the optional one or more substituents of R¹ are


independently selected from , , , , , , , , , , , , , , , and . In some cases, the optional one



or more substituents of R¹ are independently selected from halogen, , ,


Nc1nn[nH]1, Fc1ccc2c(c1)c(c[nH]2)C(=O)N, c1ccc2c(c1)c(c[nH]2)C(=O)N, Cc1ccoc1, Cc1nn[nH]1, Cc1nn[nH]1, Cc1nn[nH]1, and











In some cases, R¹ is selected from





 , and





. In some cases, R^1 is selected from







, and





. In some cases, R^1 is selected from





, and



. In some cases, R^1 is selected from






alkyl, and C_{3-6} carbocycle. In some cases, R^1 is selected







selected from , which is optionally substituted with one more substituents independently selected from halogen, -OH, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)OR²⁰, -C(O)NHOR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxyalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, -OR²⁰, and C₁₋₆ alkyl. In some cases, R¹ is selected




from , which is optionally substituted with one more substituents independently



selected from halogen and C₁₋₆ alkyl. In some cases, R¹ is selected from



[00305] In some embodiments, for a compound or salt of Formula (I), Formula (II), or



Formula (III), R¹ is selected from a , wherein  is selected from a 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted independently with one or more R^{1*}; and R^B is selected from hydrogen, halogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkynyl, and -CN. In some cases, R^B is selected from hydrogen, and halogen. In

some cases, R^B is chloride. In some cases, R^B is hydrogen. In some cases,  has at least 1, 2,


3, or 4 heteroatoms. In some cases,  has at least 1, 2, 3, or 4 nitrogen atoms. In some cases,



 has at least 1 oxygen atom. In some cases,  is a monocyclic heterocycle. In some


cases,  is a bicyclic heterocycle. In some cases,  is selected from an optionally

substituted 5-membered heterocycle. In some cases,  is selected from an optionally

substituted 9-membered heterocycle. In some cases,  is selected from , ,




substituted with one or more R^{1*}. In some cases,  is selected from , ,



optionally substituted with one or more R^{1*} . In some cases, each R^{1*} is independently selected from halogen, $-OR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from

halogen, and C_{1-6} alkyl. In some cases, $\textcircled{B}$ is selected from



[00306] In some embodiments, for a compound or salt of Formula (I), Formula (II), Formula (II*), or Formula (III), when R^1 is substituted with $-C(O)R^{20}$, R^{20} is selected from a 5- to 12-membered heterocycle, which is optionally substituted. In some cases, R^1 is substituted with $-C(O)R^{20}$. In some cases, R^{20} is selected from a 5- to 12-membered unsubstituted heterocycle. In some cases, R^{20} is selected from a 5- to 6-membered heterocycle, which is optionally substituted. In some cases, the heterocycle has at least one nitrogen atom. In some cases, the heterocycle has at least one sulfur atom. In some cases, the heterocycle has at least one oxygen atom. In some cases, the heterocycle has two heteroatoms. In some cases, the heterocycle of R^{20} is selected

from

, each of which is optionally substituted. In some

cases, R^{20} is selected from


. In


some cases, the optional substituents are selected from C_{1-10} alkyl, oxo, and $=NH$.

[00307] In some embodiments, for a compound or salt of Formula (I), Formula (II), Formula (II*), or Formula (III), each R^{20} is independently selected from hydrogen; and C_{1-6} alkyl, C_{3-12} carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one


or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, and =NH. In some cases, each R²⁰ is independently selected from hydrogen; and unsubstituted C₁₋₆ alkyl, and 3- to 12-membered heterocycle which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, and =NH.

[00308] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R¹ is an optionally substituted 12- to 15-membered heterocycle. In some cases, R¹ is an optionally substituted 12-membered heterocycle. In some cases, R¹ is an optionally substituted 13-membered heterocycle. In some cases, R¹ is an optionally substituted 14-membered heterocycle. In some cases, R¹ is an optionally substituted 15-membered heterocycle. In some cases, the heterocycle of R¹ is tricyclic. In some cases, the heterocycle of R¹ contains a fused heterocycle. In some cases, the heterocycle of R¹ contains a spiro-heterocycle. In some cases, the heterocycle of R¹ contains a fused and spiro-heterocycle. In some cases, the heterocycle of R¹ is an unsaturated heterocycle. In some cases, the heterocycle of R¹ is a non-aromatic heterocycle. In some cases, the heterocycle of R¹ has at least one double bond. In some cases, the heterocycle of R¹ has at least two double bonds. In some cases, the heterocycle of R¹ has at least 2 heteroatoms. In some cases, the heterocycle of R¹ has at least 3 heteroatoms. In some cases, the heterocycle of R¹ has at least 4 heteroatoms. In some cases, the heterocycle of R¹ has at least 5 heteroatoms. In some cases, the heterocycle of R¹ has at least 6 heteroatoms. In some cases, the heterocycle of R¹ has at least 7 heteroatoms. In some cases, the heteroatoms are selected from oxygen, nitrogen, and sulfur. In some cases, the heterocycle of R¹ has at least 3, 4, or 5 nitrogen atoms, and at least 1 sulfur atom. In some cases, the heterocycle of R¹ has at least 3, 4, or 5 nitrogen atoms, and at least 1 oxygen atom. In some cases, the heterocycle of R¹ has at least 3, 4, or 5 nitrogen atoms. In some cases, the heterocycle of R¹ has at least 3, 4, or 5 nitrogen atoms and no other heteroatoms. In some cases, the heteroatoms are selected from nitrogen and sulfur. In some cases, the heteroatoms are selected from nitrogen and oxygen. In







each of which is optionally substituted with one or more substituents. In some cases, R^1 is




selected from halogen, -OH, -NHCN, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-C(O)N(R^{20})_2$, $-C(O)NHOR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-NO_2$, $=O$, $=NH$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, and C_{2-6} alkynyl. In some cases, the optional one or more substituents of R^1 are independently selected from halogen, -OH, -NHCN, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-C(O)N(R^{20})_2$, $-C(O)NHOR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-NO_2$, $=O$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, and C_{2-6} alkynyl. In some cases, the optional one or more substituents of R^1 are independently selected from halogen, -OH, C_{1-6} alkyl, and $-C(O)N(R^{20})_2$. In some cases, R^1 is selected from







[00309] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is an optionally substituted 12- to 15-membered heterocycle. In some cases, R^1




is , wherein Ring W is an optionally substituted heterocycle and Ring P is an optionally substituted carbocycle or optionally substituted heterocycle, wherein Ring P forms a spirocycle with Ring W. In some cases, Ring W is an optionally substituted fused heterocycle. In some cases, Ring P and Ring W combine to form a heterocycle having at least 12 atoms and most 15 atoms. In some cases, Ring P and Ring W have in total at least 12 atoms and most 15 atoms. In some cases, Ring W is an optionally substituted 10-membered fused heterocycle. In



some cases, R^1 is , wherein Ring P is an optionally substituted carbocycle or



optionally substituted heterocycle. In some cases, R^1 is . In some cases, Ring P is an optionally substituted carbocycle. In some cases, Ring P is an optionally substituted heterocycle. In some cases, Ring P forms an optionally substituted C₃-C₆ carbocycle or optionally substituted 4-to 6-membered heterocycle. In some cases, Ring P forms an optionally substituted C₃ carbocycle. In some cases, Ring P forms an optionally substituted C₄ carbocycle. In some cases, Ring P forms an optionally substituted C₅ carbocycle. In some cases, Ring P forms an optionally substituted 4-membered heterocycle. In some cases, Ring P forms an optionally substituted 5-membered heterocycle. In some cases, Ring P forms an optionally substituted 5-membered heterocycle. In some cases, Ring P has at least 1, 2, or 3 heteroatoms. In some cases, the heteroatoms are selected from oxygen, nitrogen, and sulfur. In some cases, Ring P has 1 sulfur atom. In some cases, Ring P has 1 nitrogen atom. In some cases, Ring P has 1 oxygen atom. In some cases, the one or more optional substituents of Ring P are independently selected from halogen, -OH, -NHCN, =O, =NR²⁰, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of Ring P are independently selected from halogen, -OH, =O, =NH, -CN, and C₁₋₆ alkyl. In some cases, the one or more optional substituents of Ring W are independently selected from halogen, -OH, -NHCN, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(O)NHOR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, and C₂₋₆ alkynyl. In some cases, the one or more optional substituents of Ring W are independently selected from halogen, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -C(O)NHOR²⁰, -N(R²⁰)₂, -C(O)R²⁰, and C₁₋₆ alkyl. In some cases, the one or more optional substituents of Ring W are independently selected from -C(O)R²⁰. In some cases, Ring P is substituted. In some cases, Ring W is substituted.


[00310] In some embodiments, for a compound or salt of Formula (I), Formula (II), or















[00311] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^2 is selected from optionally substituted -L-heterocycle. In some cases, the heterocycle is


a bicyclic heterocycle. In some cases, the heterocycle is a monocyclic heterocycle. In some cases, the heterocycle has only 1 nitrogen atom. In some cases, the heterocycle has only 1


nitrogen atom and no other heteroatoms. In some cases, Y-R² is selected from




and , wherein the heterocycle portion is optionally substituted. In some cases, R² is


selected from  and , wherein the heterocycle portion is optionally

substituted. In some cases, Y-R² is selected from , wherein the heterocycle

portion is optionally substituted. In some cases, Y-R² is selected from , wherein the heterocycle portion is optionally substituted. In some cases, the heterocycle is optionally substituted with one or more substituent selected from halogen, hydroxy, C₁-C₃ alkyl, -N(R⁵)S(O)₂(R⁵), -OC(O)N(R⁵)₂, oxo, =CH₂, =NO-C₁-C₃ alkyl, -CH₂OC(O)heterocycle, -CH₂heterocycle, -CH₂OC(O)N(R⁵)₂, and -O-C₁-C₃ alkyl, wherein the alkyl of -O-C₁-C₃ alkyl is optionally substituted with substituents selected from heterocycle, oxo, and hydroxy. In some


cases, Y-R² is selected from






of L come together to form a C₃-C₆ carbocycle, wherein the C₃-C₆ carbocycle is optionally


substituted with one or more substituents selected from halogen. In some cases, L is



. In some cases, R²¹ is




. In some cases, Y-R²¹ is




[00313] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), each R²¹ is independently selected from hydrogen. In some cases, each R²¹ is independently selected from hydrogen and C₁₋₆ alkyl. In some cases, each R²¹ is independently selected from C₁₋₆ alkyl. In some cases, each R²¹ is independently selected from hydrogen; and C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo.

[00314] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R¹ is selected from an optionally substituted 8- to 10-membered fused heterocycle. In some cases, the 8- to 10-membered fused heterocycle is a bicyclic heterocycle. In some cases, the 8- to 10-membered fused heterocycle is a saturated heterocycle. In some cases, the 8- to 10-membered heterocycle is a non-aromatic heterocycle. In some cases, R¹ is selected from an optionally substituted 9-membered fused heterocycle. In some cases, R¹ is selected from an optionally substituted 10-membered fused heterocycle. In some cases, the 10-membered fused heterocycle is a bicyclic heterocycle. In some cases, the 10-membered fused heterocycle is a saturated heterocycle. In some cases, the 10-membered heterocycle is a non-aromatic heterocycle. In some cases, the fused heterocycle has one saturated ring and one aromatic ring. In some cases, the fused heterocycle has one saturated ring and one unsaturated ring. In some cases, the fused heterocycle has two saturated rings. In some cases, the 10-membered heterocycle contains at least 1 nitrogen atom. In some cases, the 9-membered heterocycle contains at least 2 nitrogen atoms. In some cases, the 9-membered heterocycle contains at least 3 nitrogen atoms. In some cases, the 10-membered heterocycle contains at least 2 nitrogen atoms. In some cases, the 10-membered heterocycle contains at least 3 nitrogen atoms. In some cases, R¹ is selected from




and


, each of which is optionally substituted with one or more substituents.










In some cases, R^1 is selected from _____, _____, _____, and _____, each of




which is optionally substituted with one or more substituents. In some cases, R^1 is _____, which is optionally substituted with one or more substituents. In some cases, the optional one or more substituents are independently selected from halogen, =O, -OH, -CN, -NHCN, -C(O) R^{20} , -C(O)N(R^{20})₂, -C(O)N R^{20} OR R^{20} , C₁₋₆ aminoalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, and C₁₋₆ alkyl. In some cases, the optional one or more substituents are independently selected from halogen, =O, -C(O) R^{20} , -C(O)N(R^{20})₂, and -C(O)N R^{20} OR R^{20} . In some cases, the optional one or more substituents are independently selected from -C(O) R^{20} , -C(O)N(R^{20})₂, and -C(O)N R^{20} OR R^{20} . In some cases, the optional one or more substituents are independently selected from -C(O) R^{20} . In some cases, the optional one or more substituents are independently selected from -C(O)N(R^{20})₂. In some cases, the optional one or more substituents are independently selected from -C(O)N R^{20} OR R^{20} . In some cases, each R^{20} is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle. In some cases, each R^{20} is independently selected from hydrogen; and C₁₋₆ alkyl, and 3- to 12-membered heterocycle. In some cases, each R^{20} is independently selected from hydrogen; and C₁₋₆ alkyl, and 3- to 12-membered saturated heterocycle. In some cases, the optional one or more substituents of R^1 are







[00315] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from an optionally substituted saturated 6- to 7-membered heterocycle. In some cases, R^1 is selected from an optionally substituted saturated 6-membered heterocycle. In some cases, R^1 is selected from




, which is optionally substituted. In some cases, the optional one or more substituents are independently selected from halogen, -CN, -NHCN, C_{1-6} cyanoalkyl, and C_{1-6} alkyl. In some cases, the optional one or more substituents are independently selected from -CN, -NHCN, C_{1-6} cyanoalkyl, and C_{1-6} alkyl. In some cases, the optional one or more substituents are independently selected from -CN, -NHCN, C_{1-6} cyanoalkyl, and C_{1-6} alkyl. In some cases, the optional one or more substituents are independently selected from -NHCN, and C_{1-6} alkyl. In




some cases, R^1 is selected from , which is substituted with one or more substituents selected




from -NHCN, and C_{1-6} alkyl. In some cases, R^1 is selected from , and






[00316] In some embodiments, for a compound or salt of Formula (I), R^1 is selected from a substituted saturated 6-membered heterocycle, wherein the saturated 6-membered heterocycle is substituted with at least one -NHCN, and optionally one or more C_{1-6} alkyl; B is selected from an optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C_8 - C_{15} fused carbocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, oxo, -NH₂, C_1 - C_3 alkyl, -OH, -C(O)NH₂, -NH₂, =O, -CN, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{2-6} alkynyl; Ring A is selected from an optionally substituted heterocycle; Y is O; R^2 is selected from -L-heterocycle, wherein the heterocycle portion is optionally substituted with one or more substituents independently selected from halogen, hydroxy, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 hydroxyalkyl, or -N(R^5)₂;




and L is selected from C_1 - C_4 alkylene. In some cases, R^1 is selected from , and




. In some cases, B is selected from




[00317] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), B is an optionally substituted 8- to 10-membered fused carbocycle. In some cases, B is a substituted 8- to 10-membered fused carbocycle. In some cases, B is an unsubstituted 8- to 10-membered fused carbocycle. In some cases, B is an optionally substituted 9-membered fused carbocycle. In some cases, B is a substituted 9-membered fused carbocycle. In some cases, B is





, which is optionally substituted with one or more substituents. In some cases, B is










, which is substituted with one or more substituents. In some cases, for B, the one or more substituents are independently selected from halogen, oxo, $-NH_2$, C_1 - C_3 alkyl, $-B(OH)_2$, $-OH$, $-C(O)NH_2$, $-NH_2$, $=O$, $-CN$, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{2-6} alkynyl. In some cases, B is substituted with at least one halogen. In some cases, B is substituted with at least one chlorine. In some cases, B is substituted with at least one fluorine. In some cases, B is selected











selected from halogen and C_{1-6} haloalkyl. In some cases, B is , which is substituted with


one or more substituents selected from halogen. In some cases, B is selected from ,

, , , , and . In some cases, B is ,



which is substituted with one or more substituents selected from fluorine. In some cases, B is

selected from , , and . In some cases, B is , which is substituted with one or more substituents selected from chlorine. In some cases, B is selected

from  and . In some cases, B is a substituted 10-membered fused carbocycle. In some cases, for the 10-membered fused carbocycle of B, the one or more substituents are independently selected from halogen, -NH_2 , $\text{C}_1\text{-C}_3$ alkyl, $\text{-B}(\text{OH})_2$, -OH , $\text{-C}(\text{O})\text{NH}_2$, -NH_2 , $=\text{O}$, -CN , C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{2-6} alkynyl. In some cases, B is




In some cases, for the 10-membered fused carbocycle of B, is substituted with at least one





[00318] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the one or more optional substituents of R^1 are independently selected from halogen, -OH, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NHR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxyalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, and C₁₋₆ alkyl.

[00319] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula





(III), R^1 is , and the one or more optional substituents of R^1 are independently selected from halogen, -OH, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-C(O)N(R^{20})_2$, $-C(=NR^{20})N(R^{20})_2$, $-C(O)NHR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-NO_2$, $=O$, $-CN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} alkoxyalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkynyl, and optionally substituted 5- to 12-membered heterocycle. In some cases, the one or more optional substituents of R^1 are independently selected from halogen, -OH, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-C(O)N(R^{20})_2$, $-C(=NR^{20})N(R^{20})_2$, $-C(O)NHR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-NO_2$, $=O$, $-CN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} alkoxyalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkynyl, and optionally substituted 5- to 12-membered heterocycle. In some cases, the one or more optional substituents of R^1 are independently selected from halogen, -CN, C_{2-6} alkynyl, $-C(=NR^{20})N(R^{20})_2$, and optionally substituted 5- to 12-membered heterocycle. In some cases, the one or more optional substituents of R^1 are independently selected from halogen, $-C(=NR^{20})N(R^{20})_2$, and optionally substituted 5- to 12-membered heterocycle. In some cases, the one or more optional substituents of R^1 are independently selected from optionally substituted 5- to 12-membered heterocycle. In some cases, the one or more optional substituents of R^1 are independently selected from a 5-membered heterocycle and 9-membered heterocycle, each of which is optionally substituted independently with one or more R^{1*} . In some cases, R^1 is substituted with at least one halogen atom and optionally substituted with one or more substituents are independently selected from $-CN$, C_{2-6} alkynyl, $-C(=NR^{20})N(R^{20})_2$, and 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted independently with one or more R^{1*} . In some cases, the heterocycle has at least one nitrogen atom. In some cases, the heterocycle has at least oxygen atom. In some cases, the heterocycle has at least one nitrogen atom and at least one oxygen atom. In some cases, heterocycle has at least two heteroatoms. In some cases, the heterocycle has at least three heteroatoms. In some cases, the heterocycle has at least four heteroatoms. In some cases, the heterocycle of the one or more


optional substituents of R^1 is selected from ,



optionally substituted with one or more R^{1*} . In some cases, the heterocycle of the one or more




optional substituents of R^1 is selected from , which is optionally substituted with one or more R^{1*} . In some cases, each R^{1*} is independently selected from halogen, $-OR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, $-OR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen. In some cases, each R^{1*} is independently selected from C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from $-OR^{20}$. In some cases, each R^{1*} is independently selected from $-OH$. In some cases, each R^{1*} is independently selected from $-OMe$. In some cases, the heterocycle of the one or more optional substituents of R^1 is selected from




[00320] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), the one or more optional substituents of R^1 are independently selected from $-C(=NR^{20})N(R^{20})_2$, and optionally substituted 5- to 12-membered heterocycle. In some cases, the one or more optional substituents of R^1 are independently selected from optionally substituted 5-

to 12-membered heterocycle. In some cases, the heterocycle is selected from



, , , , , , , and , each of which is optionally substituted with one or more R^{1*} . In some cases, the one or more optional






substituents of R^1 is selected from

































[00321] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), each R^{1*} is independently selected from halogen, $-OR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, $-OR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen, and C_{1-6} alkyl. In some cases, each R^{1*} is independently selected from halogen. In some cases, each R^{1*} is independently selected from C_{1-6} alkyl.





[00322] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from 5- to 15-membered heterocycle (preferably 8- to 10-membered heterocycle or preferably 10-membered heterocycle), each of which are optionally substituted with one or more substituents independently selected from halogen, oxo, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-SO_2R^{20}$, $-NHCN$, C_{1-6} cyanoalkyl, C_{1-6} alkyl, C_{1-6} alkyl- $N(R^{20})_2$, C_{2-6} alkynyl, and 5- to 12-membered heterocycle (preferably 5- to 9-

membered heterocycle), wherein the 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*} ; each R^{1*} is independently selected from halogen, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, the 8- to 10-membered heterocycle is bicyclic. In some cases, the 10-membered heterocycle is substituted. In some cases, R^1 is

selected , , , , and , each of which is optionally substituted. In some


cases, R^1 is selected , which is optionally substituted. In some cases, R^1 is selected


, , , , , ,
, , , , , ,
, , , , , ,
, , , , , ,
, , , , and . In some cases, R^1 is selected






, , and . In some cases, R^1 is .






[00323] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from 5- to 15-membered heterocycle (preferably 8- to 10-membered heterocycle or preferably 10-membered heterocycle), each of which are optionally substituted with one or more substituents independently selected from halogen, $-C(O)N(R^{20})_2$, -







$C(O)NR^{20}OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-NHCN$, C_{1-6} cyanoalkyl, C_{1-6} alkyl, C_{2-6} alkynyl, and 5- to 12-membered heterocycle (preferably 5- to 6-membered heterocycle), wherein the 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*} ; each R^{1*} is independently selected from halogen, C_{1-6} haloalkyl, and C_{1-6} alkyl. In some cases, the 8- to 10-membered heterocycle is bicyclic. In some cases, the 10-membered






heterocycle is substituted. In some cases, R^1 is selected , , and , each of which is

optionally substituted. In some cases, R^1 is selected , which is optionally substituted. In

some cases, R^1 is selected , , , , ,



, , , , ,



, , , , , ,

, , and . In some cases, R^1 is selected  and . In


some cases, M is selected from O , and NMe . In some cases, M is O . In some cases, M is NMe .

In some cases, R^2 is selected from $-L-N(R^{21})_2$ and $-L$ -heterocycle, optionally substituted with one

or more R^6 . In some cases, $Y-R^2$ is selected from , ,

, and . In some cases, B is selected from an optionally substituted




carbocycle. In some cases, B is selected from




and

, each of which is optionally


substituted. In some cases, B is selected from



. In some cases, B is



. In some cases, B is




. In some cases, n is 0.

[00324] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^1 is selected from a compound in the Examples section. In some cases, Y is selected from a compound in the Examples section. In some cases, L is selected from a compound in the Examples section. In some cases, R^2 is selected from a compound in the Examples section. In some cases, B is selected from a compound in the Examples section. In some cases, M is selected from a compound in the Examples section. In some cases, the optional substituents of the heterocycle for R^1 is selected from a compound in the Examples section.

[00325] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R^2 is $-L-N(R^{21})_2$. In some cases, R^2 is $-L-OR^{21}$. In some cases, R^2 is heterocycle. In some cases, R^2 is C_1-C_6 alkyl. In some cases, R^2 is $-L$ -heterocycle. In some cases, R^2 is $-L$ -aryl. In some cases, R^2 is $-L$ -heteroaryl. In some cases, R^2 is $-L$ -cycloalkyl. In some cases, R^2 is $-L-N(R^{21})_2$. In some cases, R^2 is $-L-NHC(=NH)NH_2$. In some cases, R^2 is $-L-C(O)N(R^{21})_2$. In some cases, R^2 is $-L-C_1-C_6$ haloalkyl. In some cases, R^2 is $-L-OR^{21}$. In some cases, R^2 is $-L-NR^{21}C(O)$ -aryl. In some cases, R^2 is $-L-COOH$. In some cases, R^2 is $-L-NR^{21}S(O)_2(R^{21})$. In some cases, R^2 is $-L-S(O)_2N(R^{21})_2$. In some cases, R^2 is $-L-N(R^{21})C(O)(OR^{21})$. In some cases, R^2 is $-L-OC(O)N(R^{21})_2$. In some cases, R^2 is or $-LC(=O)OC_1-C_6$ alkyl. In some cases, the heterocycle, the aryl portion of $-L-NR^{21}C(O)$ -aryl, the heterocycle portion of $-L$ -heterocycle, and the cycloalkyl portion of the $-L$ -cycloalkyl are each optionally substituted with one or more R^6 , and wherein the aryl portion of the $-L$ -aryl and the heteroaryl portion of the $-L$ -heteroaryl are each optionally substituted with one or more R^7 . In some cases, when Y is a bond, O, or S, R^2 is further selected from hydrogen

[00326] In some embodiments, Formula (I) or Formula (II) is represented by Formula (II*):



Formula (II*)

or a pharmaceutically acceptable salt thereof wherein:

M is selected from O, and NR³;

R³ is selected from hydrogen, C₁₋₆ alkyl, and C₁₋₆ cyanoalkyl;

R¹ is selected from a 7- to 10-membered heterocycle, wherein the 7- to 10-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted independently with one or more R^{1*};

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃₋₁₂ carbocycle;

B is selected from C₆₋₁₅ carbocycle, wherein the C₆₋₁₅ carbocycle is optionally substituted with one or more substituents independently selected from halogen, C₁₋₃ alkyl, -B(OR²⁰)₂, -OR²⁰, -C(O)N(R²⁰)₂, -N(R²⁰)₂, =O, -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;


R² is selected from -L-heterocycle, wherein the heterocycle of -L-heterocycle is optionally substituted with one or more R⁶;

L is independently selected from a C₁₋₄ alkylene, wherein the C₁₋₄ alkylene is optionally substituted with one or more substituents selected from hydroxy, C₁₋₄ hydroxyalkyl, C₁₋₄ alkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃₋₆ carbocycle;


each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₆ carbocycle, and 3- to 6-membered heterocycle, each of which is optionally substituted with one or more substituents

independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, =NH, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle.


[00327] In some embodiments, for a compound or salt of Formula (II*), R¹ is selected from an optionally substituted 7- to 10-membered spiro heterocycle and optionally substituted 7- to 10-membered fused heterocycle. In some cases, the heterocycle of R¹ has at least one nitrogen atom. In some cases, the at least one nitrogen atom of the heterocycle of R¹ is bound to Formula (II*). In some cases, R¹ is selected from an optionally substituted 10-membered spiro heterocycle and optionally substituted 10-membered fused heterocycle. In some cases, the optional one or more substituents of R¹ are independently selected from halogen, -OH, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)OR²⁰, -C(O)NHR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxyalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, and C₁₋₆ alkyl. In some cases, R¹ is selected from




, which is substituted with one or more substituents independently selected from halogen, -OH, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)OR²⁰, -C(O)NHR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxyalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from




halogen, and C₁₋₆ alkyl. In some cases, R¹ is selected from









halogen, =CH₂, hydroxy, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, cyano, and C₁-C₃ aminoalkyl. In some cases, L of R² is selected from C₁-C₄ alkylene and .

In some cases, R² is selected from




In some cases, B is selected from an optionally substituted


C₉-C₁₀ fused carbocycle. In some cases, B is selected from , and , each of which is optionally substituted. In some cases, B is optionally substituted with one or more substituents independently selected from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OH)₂, -OH, -O-C₁-C₃ haloalkyl, -C(O)NH₂, -NH₂, =O, -CN, C₁-C₆ alkoxy, C₁-C₆ hydroxyalkyl, and C₂-C₆ alkynyl. In some cases, B is optionally substituted with one or more substituents independently selected from halogen. In


some cases, B is . In some cases, B is . In some cases, B is unsubstituted. In some cases, B is substituted.

[00328] In some embodiments, for a compound or salt of Formula (II*), R¹ is selected from an optionally substituted 7- to 10-membered spiro heterocycle and optionally substituted 7- to 10-membered fused heterocycle. In some cases, R¹ is selected from an optionally substituted 10-membered spiro heterocycle and optionally substituted 10-membered fused heterocycle. In some cases, R¹ is selected from an optionally substituted 10-membered spiro heterocycle. In some cases, R¹ is selected from an optionally substituted 10-membered fused heterocycle. In some cases, the heterocycle of R¹ has at least 3 heteroatoms. In some cases, the optional one or more substituents of R¹ are independently selected from halogen, -OH, -S(O)₂(R²⁰), -S(O)N(R²⁰)₂, -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)OR²⁰, -C(O)NHR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁-C₆ alkyl-N(R²⁰)₂, C₁-C₆ aminoalkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxyalkyl, C₁-C₆ hydroxyalkyl, C₁-C₆ cyanoalkyl, C₁-C₆ haloalkyl, C₁-C₆ alkyl, C₂-C₆ alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, and C₁-C₆ alkyl. In




some cases, R^1 is selected from , which is substituted with one or more substituents independently selected from halogen, $-OH$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-C(O)N(R^{20})_2$, $-C(=NR^{20})N(R^{20})_2$, $-C(O)OR^{20}$, $-C(O)NHR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-NO_2$, $=O$, $-CN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} alkoxyalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, and C_{1-6} alkyl. In some cases, R^1 is selected from






and C₁-C₃ aminoalkyl. In some cases, L of R² is selected from C₁-C₄ alkylene and




In some cases, R² is selected from




. In some cases, B is selected from an optionally substituted


C₉-C₁₀ fused carbocycle. In some cases, B is selected from



, and , each of which is optionally substituted. In some cases, B is optionally substituted with one or more substituents independently selected from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OH)₂, -OH, -O-C₁-C₃ haloalkyl, -C(O)NH₂, -NH₂, =O, -CN, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₂₋₆ alkynyl. In some cases, B is optionally substituted with one or more substituents independently selected from halogen. In



some cases, B is



. In some cases, B is . In some cases, B is unsubstituted. In some cases, B is substituted.

[00329] In some embodiments, for a compound or salt of Formula (I), Formula (II), Formula (II*), or Formula (III), the heterocycle or carbocycle of R¹ is not substituted by C₁₋₆ cyanoalkyl. In some embodiments, for a compound or salt of Formula (I), the heterocycle or carbocycle of R¹ is not substituted by C₁₋₆ cyanoalkyl. In some embodiments, for a compound or salt of Formula (II), the heterocycle or carbocycle of R¹ is not substituted by C₁₋₆ cyanoalkyl. In some embodiments, for a compound or salt of Formula (III), the heterocycle or carbocycle of R¹ is not substituted by C₁₋₆ cyanoalkyl.

[00330] In some embodiments, for a compound or salt of Formula (I), Formula (II), Formula (II*), or Formula (III), each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl.

[00331] In some embodiments, for a compound or salt of Formula (I), Formula (II), Formula (II*), or Formula (III), is not substituted by C₁₋₆ cyanoalkyl.

[00332] In some embodiments, for a compound or salt of Formula (I), Formula (II), or Formula (III), R¹ is not a piperazine. In some cases, R¹ is not a substituted piperazine.

[00333] In some embodiments, for a compound of Formula (I), wherein the compound is not a Michael acceptor.

[00334] In some embodiments, for a compound of Formula (I), the compound or salt does not include an electrophilic substituent.

[00335] In some embodiments, for a compound or salt of Formula (I), Formula (II), Formula (II*), or Formula (III), does not contain an electrophile moiety.


[00336] In some embodiments, for a compound or salt of Formula (I), Formula (II), Formula (II*), or Formula (III), does not contain a covalent modifier.

[00337] In some embodiments, for a compound or salt of Formula (I), Formula (II), Formula (II*), or Formula (III), the one or more optional substituents of R¹ are not electrophiles.

[00338] Included in the present disclosure are salts, particularly pharmaceutically acceptable salts, of the compounds described herein. The compounds of the present invention that possess a sufficiently acidic, a sufficiently basic, or both functional groups, can react with any of a number of inorganic bases, and inorganic and organic acids, to form a salt. Alternatively, compounds that are inherently charged, such as those with a quaternary nitrogen, can form a salt with an appropriate counterion, e.g., a halide such as bromide, chloride, or fluoride, particularly bromide.

[00339] Chemical entities having carbon-carbon double bonds or carbon-nitrogen double bonds may exist in *Z*- or *E*- form (or *cis*- or *trans*- form). Furthermore, some chemical entities may exist in various tautomeric forms. Unless otherwise specified, compounds described herein are intended to include all *Z*-, *E*- and tautomeric forms as well.

[00340] A “tautomer” refers to a molecule wherein a proton shift from one atom of a molecule to another atom of the same molecule is possible. The compounds presented herein, in certain embodiments, exist as tautomers. In circumstances where tautomerization is possible, a chemical equilibrium of the tautomers will exist. The exact ratio of the tautomers depends on several factors, including physical state, temperature, solvent, and pH. Some examples of tautomeric equilibrium include:



[00341] The compounds disclosed herein, in some embodiments, are used in different enriched isotopic forms, e.g., enriched in the content of ^2H , ^3H , ^{11}C , ^{13}C and/or ^{14}C . In one particular embodiment, the compound is deuterated in at least one position. Such deuterated forms can be made by the procedure described in U.S. Patent Nos. 5,846,514 and 6,334,997. As described in U.S. Patent Nos. 5,846,514 and 6,334,997, deuteration can improve the metabolic stability and or efficacy, thus increasing the duration of action of drugs.

[00342] Unless otherwise stated, compounds described herein are intended to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by ^{13}C - or ^{14}C -enriched carbon are within the scope of the present disclosure.

[00343] The compounds of the present disclosure optionally contain unnatural proportions of atomic isotopes at one or more atoms that constitute such compounds. For example, the compounds may be labeled with isotopes, such as for example, deuterium (^2H), tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). Isotopic substitution with ^2H , ^{11}C , ^{13}C , ^{14}C , ^{15}C , ^{12}N , ^{13}N , ^{15}N , ^{16}N , ^{16}O , ^{17}O , ^{14}F , ^{15}F , ^{16}F , ^{17}F , ^{18}F , ^{33}S , ^{34}S , ^{35}S , ^{36}S , ^{35}Cl , ^{37}Cl , ^{79}Br , ^{81}Br , and ^{125}I are all contemplated. All isotopic variations of the compounds of the present invention, whether radioactive or not, are encompassed within the scope of the present invention.

[00344] In certain embodiments, the compounds disclosed herein have some or all of the ^1H atoms replaced with ^2H atoms. The methods of synthesis for deuterium-containing compounds are known in the art and include, by way of non-limiting example only, the following synthetic methods.

[00345] Deuterium substituted compounds are synthesized using various methods such as described in: Dean, Dennis C.; Editor. Recent Advances in the Synthesis and Applications of Radiolabeled Compounds for Drug Discovery and Development. [In: Curr., Pharm. Des., 2000; 6(10)] **2000**, 110 pp; George W.; Varma, Rajender S. The Synthesis of Radiolabeled Compounds via Organometallic Intermediates, Tetrahedron, **1989**, 45(21), 6601-21; and Evans, E. Anthony. Synthesis of radiolabeled compounds, J. Radioanal. Chem., **1981**, 64(1-2), 9-32.

[00346] Deuterated starting materials are readily available and are subjected to the synthetic methods described herein to provide for the synthesis of deuterium-containing compounds. Large numbers of deuterium-containing reagents and building blocks are available commercially from chemical vendors, such as Aldrich Chemical Co.

[00347] Compounds of the present invention also include crystalline and amorphous forms of those compounds, pharmaceutically acceptable salts, and active metabolites of these compounds having the same type of activity, including, for example, polymorphs, pseudopolymorphs, solvates, hydrates, unsolvated polymorphs (including anhydrides), conformational polymorphs, and amorphous forms of the compounds, as well as mixtures thereof.

[00348] The compounds described herein may in some cases exist as diastereomers, enantiomers, or other stereoisomeric forms. Where absolute stereochemistry is not specified, the compounds presented herein include all diastereomeric, enantiomeric, and epimeric forms as well as the appropriate mixtures thereof. Separation of stereoisomers may be performed by chromatography or by forming diastereomers and separating by recrystallization, or chromatography, or any combination thereof. (Jean Jacques, Andre Collet, Samuel H. Wilen, "Enantiomers, Racemates and Resolutions", John Wiley And Sons, Inc., 1981, herein incorporated by reference for this disclosure). Stereoisomers may also be obtained by stereoselective synthesis.

[00349] The methods and compositions described herein include the use of amorphous forms as well as crystalline forms (also known as polymorphs). The compounds described herein may be in the form of pharmaceutically acceptable salts. As well, in some embodiments, active metabolites of these compounds having the same type of activity are included in the scope of the present disclosure. In addition, the compounds described herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein.

[00350] In certain embodiments, compounds or salts of the compounds may be prodrugs, e.g., wherein a hydroxyl in the parent compound is presented as an ester or a carbonate, or carboxylic acid present in the parent compound is presented as an ester. The term "prodrug" is intended to

encompass compounds which, under physiologic conditions, are converted into pharmaceutical agents of the present disclosure. One method for making a prodrug is to include one or more selected moieties which are hydrolyzed under physiologic conditions to reveal the desired molecule. In other embodiments, the prodrug is converted by an enzymatic activity of the host animal such as specific target cells in the host animal. For example, esters or carbonates (e.g., esters or carbonates of alcohols or carboxylic acids and esters of phosphonic acids) are preferred prodrugs of the present disclosure.

[00351] Prodrug forms of the herein described compounds, wherein the prodrug is metabolized *in vivo* to produce a compound as set forth herein are included within the scope of the claims. In some cases, some of the herein-described compounds may be a prodrug for another derivative or active compound.

[00352] Prodrugs are often useful because, in some situations, they may be easier to administer than the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent is not. Prodrugs may help enhance the cell permeability of a compound relative to the parent drug. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. Prodrugs may be designed as reversible drug derivatives, for use as modifiers to enhance drug transport to site-specific tissues or to increase drug residence inside of a cell.

[00353] In some embodiments, the design of a prodrug increases the lipophilicity of the pharmaceutical agent. In some embodiments, the design of a prodrug increases the effective water solubility. See, e.g., Fedorak *et al.*, *Am. J. Physiol.*, 269:G210-218 (1995); McLoed *et al.*, *Gastroenterol.*, 106:405-413 (1994); Hochhaus *et al.*, *Biomed. Chrom.*, 6:283-286 (1992); J. Larsen and H. Bundgaard, *Int. J. Pharmaceutics*, 37, 87 (1987); J. Larsen *et al.*, *Int. J. Pharmaceutics*, 47, 103 (1988); Sinkula *et al.*, *J. Pharm. Sci.*, 64:181-210 (1975); T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems*, Vol. 14 of the A.C.S. Symposium Series; and Edward B. Roche, *Bioreversible Carriers in Drug Design*, American Pharmaceutical Association and Pergamon Press, 1987, all incorporated herein for such disclosure). According to another embodiment, the present disclosure provides methods of producing the above-defined compounds. The compounds may be synthesized using conventional techniques.

Advantageously, these compounds are conveniently synthesized from readily available starting materials.

[00354] Synthetic chemistry transformations and methodologies useful in synthesizing the compounds described herein are known in the art and include, for example, those described in R. Larock, *Comprehensive Organic Transformations* (1989); T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 2d. Ed. (1991); L. Fieser and M. Fieser, *Fieser*

and *Fieser's Reagents for Organic Synthesis* (1994); and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis* (1995).

Pharmaceutical Formulations

[00355] Provided herein, in certain embodiments, are compositions comprising a therapeutically effective amount of any compound or salt of any one of Formulas (I), (II), (II*), and (III) (also referred to herein as “a pharmaceutical agent”).

[00356] Pharmaceutical compositions may be formulated using one or more physiologically acceptable carriers including excipients and auxiliaries which facilitate processing of the pharmaceutical agent into preparations which are used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions is found, for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa., Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins, 1999).

[00357] The compositions and methods of the present disclosure may be utilized to treat an individual in need thereof. In certain embodiments, the individual is a mammal such as a human, or a non-human mammal. When administered to an animal, such as a human, the composition or the pharmaceutical agent, is preferably administered as a pharmaceutical composition comprising, for example, a pharmaceutical agent and a pharmaceutically acceptable carrier or excipient. Pharmaceutically acceptable carriers are well known in the art and include, for example, aqueous solutions such as water or physiologically buffered saline or other solvents or vehicles such as glycols, glycerol, oils such as olive oil, or injectable organic esters. In a preferred embodiment, when such pharmaceutical compositions are for human administration, particularly for invasive routes of administration, e.g., routes, such as injection or implantation, that circumvent transport or diffusion through an epithelial barrier, the aqueous solution is pyrogen-free, or substantially pyrogen-free. The excipients can be chosen, for example, to effect delayed release of an agent or to selectively target one or more cells, tissues or organs. The pharmaceutical composition can be in dosage unit form such as tablet, capsule, granule, lyophile for reconstitution, powder, solution, syrup, suppository, injection or the like. The composition can also be present in a transdermal delivery system, e.g., a skin patch. The composition can also be present in a solution suitable for topical administration, such as an eye drop.

[00358] A pharmaceutically acceptable excipient can contain physiologically acceptable agents that act, for example, to stabilize, increase solubility or to increase the absorption of a compound

such as a pharmaceutical agent. Such physiologically acceptable agents include, for example, carbohydrates, such as glucose, sucrose or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins or other stabilizers or excipients. The choice of a pharmaceutically acceptable excipient, including a physiologically acceptable agent, depends, for example, on the route of administration of the composition. The preparation or pharmaceutical composition can be a self emulsifying drug delivery system or a self microemulsifying drug delivery system. The pharmaceutical composition (preparation) also can be a liposome or other polymer matrix, which can have incorporated therein, for example, a compound of the invention. Liposomes, for example, which comprise phospholipids or other lipids, are nontoxic, physiologically acceptable and metabolizable carriers that are relatively simple to make and administer.

[00359] A pharmaceutical composition (preparation) can be administered to a subject by any of a number of routes of administration including, for example, orally, for example, drenches as in aqueous or non-aqueous solutions or suspensions, tablets, capsules, including sprinkle capsules and gelatin capsules, boluses, powders, granules, pastes for application to the tongue; absorption through the oral mucosa, e.g., sublingually; anally, rectally or vaginally, for example, as a pessary, cream or foam; parenterally, including intramuscularly, intravenously, subcutaneously or intrathecally as, for example, a sterile solution or suspension; nasally; intraperitoneally; subcutaneously; transdermally, for example, as a patch applied to the skin; and topically, for example, as a cream, ointment or spray applied to the skin, or as an eye drop. The compound may also be formulated for inhalation. In certain embodiments, a compound may be simply dissolved or suspended in sterile water.

[00360] A pharmaceutical composition may be a sterile aqueous or non-aqueous solution, suspension or emulsion, e.g., a microemulsion. The excipients described herein are examples and are in no way limiting. An effective amount or therapeutically effective amount refers to an amount of the one or more pharmaceutical agents administered to a subject, either as a single dose or as part of a series of doses, which is effective to produce a desired therapeutic effect.

[00361] Subjects may generally be monitored for therapeutic effectiveness using assays and methods suitable for the condition being treated, which assays will be familiar to those having ordinary skill in the art and are described herein. Pharmacokinetics of a pharmaceutical agent, or one or more metabolites thereof, that is administered to a subject may be monitored by determining the level of the pharmaceutical agent or metabolite in a biological fluid, for example, in the blood, blood fraction, e.g., serum, and/or in the urine, and/or other biological sample or biological tissue from the subject. Any method practiced in the art and described

herein to detect the agent may be used to measure the level of the pharmaceutical agent or metabolite during a treatment course.

[00362] The dose of a pharmaceutical agent described herein for treating a disease or disorder may depend upon the subject's condition, that is, stage of the disease, severity of symptoms caused by the disease, general health status, as well as age, gender, and weight, and other factors apparent to a person skilled in the medical art. Pharmaceutical compositions may be administered in a manner appropriate to the disease to be treated as determined by persons skilled in the medical arts. In addition to the factors described herein and above related to use of pharmaceutical agent for treating a disease or disorder, suitable duration and frequency of administration of the pharmaceutical agent may also be determined or adjusted by such factors as the condition of the patient, the type and severity of the patient's disease, the particular form of the active ingredient, and the method of administration. Optimal doses of an agent may generally be determined using experimental models and/or clinical trials. The optimal dose may depend upon the body mass, weight, or blood volume of the subject. The use of the minimum dose that is sufficient to provide effective therapy is usually preferred. Design and execution of pre-clinical and clinical studies for a pharmaceutical agent, including when administered for prophylactic benefit, described herein are well within the skill of a person skilled in the relevant art. When two or more pharmaceutical agents are administered to treat a disease or disorder, the optimal dose of each pharmaceutical agent may be different, such as less than when either agent is administered alone as a single agent therapy. In certain particular embodiments, two pharmaceutical agents in combination may act synergistically or additively, and either agent may be used in a lesser amount than if administered alone. An amount of a pharmaceutical agent that may be administered per day may be, for example, between about 0.01 mg/kg and 100 mg/kg, *e.g.*, between about 0.1 to 1 mg/kg, between about 1 to 10 mg/kg, between about 10-50 mg/kg, between about 50-100 mg/kg body weight. In other embodiments, the amount of a pharmaceutical agent that may be administered per day is between about 0.01 mg/kg and 1000 mg/kg, between about 100-500 mg/kg, or between about 500-1000 mg/kg body weight. The optimal dose, per day or per course of treatment, may be different for the disease or disorder to be treated and may also vary with the administrative route and therapeutic regimen.

[00363] Pharmaceutical compositions comprising a pharmaceutical agent can be formulated in a manner appropriate for the delivery method by using techniques routinely practiced in the art. The composition may be in the form of a solid, *e.g.*, tablet, capsule, semi-solid, *e.g.*, gel, liquid, or gas, *e.g.*, aerosol. In other embodiments, the pharmaceutical composition is administered as a bolus infusion.

[00364] Pharmaceutical acceptable excipients are well known in the pharmaceutical art and described, for example, in Rowe et al., *Handbook of Pharmaceutical Excipients: A Comprehensive Guide to Uses, Properties, and Safety*, 5th Ed., 2006, and in *Remington: The Science and Practice of Pharmacy* (Gennaro, 21st Ed. Mack Pub. Co., Easton, PA (2005)). Exemplary pharmaceutically acceptable excipients include sterile saline and phosphate buffered saline at physiological pH. Preservatives, stabilizers, dyes, buffers, and the like may be provided in the pharmaceutical composition. In addition, antioxidants and suspending agents may also be used. In general, the type of excipient is selected based on the mode of administration, as well as the chemical composition of the active ingredient(s). Alternatively, compositions described herein may be formulated as a lyophilizate. A composition described herein may be lyophilized or otherwise formulated as a lyophilized product using one or more appropriate excipient solutions for solubilizing and/or diluting the pharmaceutical agent(s) of the composition upon administration. In other embodiments, the pharmaceutical agent may be encapsulated within liposomes using technology known and practiced in the art. In certain particular embodiments, a pharmaceutical agent is not formulated within liposomes for application to a stent that is used for treating highly, though not totally, occluded arteries. Pharmaceutical compositions may be formulated for any appropriate manner of administration described herein and in the art.

[00365] A pharmaceutical composition, *e.g.*, for oral administration or for injection, infusion, subcutaneous delivery, intramuscular delivery, intraperitoneal delivery or other method, may be in the form of a liquid. A liquid pharmaceutical composition may include, for example, one or more of the following: a sterile diluent such as water, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils that may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents; antioxidants; chelating agents; buffers and agents for the adjustment of tonicity such as sodium chloride or dextrose. A parenteral composition can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. The use of physiological saline is preferred, and an injectable pharmaceutical composition is preferably sterile. In another embodiment, for treatment of an ophthalmological condition or disease, a liquid pharmaceutical composition may be applied to the eye in the form of eye drops. A liquid pharmaceutical composition may be delivered orally.

[00366] For oral formulations, at least one of the pharmaceutical agents described herein can be used alone or in combination with appropriate additives to make tablets, powders, granules or capsules, and if desired, with diluents, buffering agents, moistening agents, preservatives, coloring agents, and flavoring agents. The pharmaceutical agents may be formulated with a buffering agent to provide for protection of the compound from low pH of the gastric

environment and/or an enteric coating. A pharmaceutical agent included in a pharmaceutical composition may be formulated for oral delivery with a flavoring agent, *e.g.*, in a liquid, solid or semi-solid formulation and/or with an enteric coating.

[00367] A pharmaceutical composition comprising any one of the pharmaceutical agents described herein may be formulated for sustained or slow release, also called timed release or controlled release. Such compositions may generally be prepared using well known technology and administered by, for example, oral, rectal, intradermal, or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain the compound dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane. Excipients for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. The amount of pharmaceutical agent contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release, and the nature of the condition, disease or disorder to be treated or prevented.

[00368] In certain embodiments, the pharmaceutical compositions comprising a pharmaceutical agent are formulated for transdermal, intradermal, or topical administration. The compositions can be administered using a syringe, bandage, transdermal patch, insert, or syringe-like applicator, as a powder/talc or other solid, liquid, spray, aerosol, ointment, foam, cream, gel, paste. This preferably is in the form of a controlled release formulation or sustained release formulation administered topically or injected directly into the skin adjacent to or within the area to be treated, *e.g.*, intradermally or subcutaneously. The active compositions can also be delivered via iontophoresis. Preservatives can be used to prevent the growth of fungi and other microorganisms. Suitable preservatives include, but are not limited to, benzoic acid, butylparaben, ethyl paraben, methyl paraben, propylparaben, sodium benzoate, sodium propionate, benzalkonium chloride, benzethonium chloride, benzyl alcohol, cetylpyridinium chloride, chlorobutanol, phenol, phenylethyl alcohol, thimerosal, and combinations thereof.

[00369] Pharmaceutical compositions comprising a pharmaceutical agent can be formulated as emulsions for topical application. An emulsion contains one liquid distributed in the body of a second liquid. The emulsion may be an oil-in-water emulsion or a water-in-oil emulsion. Either or both of the oil phase and the aqueous phase may contain one or more surfactants, emulsifiers, emulsion stabilizers, buffers, and other excipients. The oil phase may contain other oily pharmaceutically approved excipients. Suitable surfactants include, but are not limited to, anionic surfactants, non-ionic surfactants, cationic surfactants, and amphoteric surfactants. Compositions for topical application may also include at least one suitable suspending agent, antioxidant, chelating agent, emollient, or humectant.

[00370] Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. Liquid sprays may be delivered from pressurized packs, for example, via a specially shaped closure. Oil-in-water emulsions can also be used in the compositions, patches, bandages and articles. These systems are semisolid emulsions, micro-emulsions, or foam emulsion systems.

[00371] In some embodiments, the pharmaceutical agent described herein can be formulated as an inhalant. Inhaled methods can deliver medication directly to the airway. The pharmaceutical agent can be formulated as aerosols, microspheres, liposomes, or nanoparticles. The pharmaceutical agent can be formulated with solvents, gases, nitrates, or any combinations thereof. Compositions described herein are optionally formulated for delivery as a liquid aerosol or inhalable dry powder. Liquid aerosol formulations are optionally nebulized predominantly into particle sizes that can be delivered to the terminal and respiratory bronchioles. Liquid aerosol and inhalable dry powder formulations are preferably delivered throughout the endobronchial tree to the terminal bronchioles and eventually to the parenchymal tissue.

[00372] Aerosolized formulations described herein are optionally delivered using an aerosol forming device, such as a jet, vibrating porous plate or ultrasonic nebulizer, preferably selected to allow the formation of aerosol particles having with a mass medium average diameter predominantly between 1 to 5 μ . Further, the formulation preferably has balanced osmolality ionic strength and chloride concentration, and the smallest aerosolizable volume able to deliver effective dose of the pharmaceutical agent. Additionally, the aerosolized formulation preferably does not impair negatively the functionality of the airways and does not cause undesirable side effects.

[00373] Aerosolization devices suitable for administration of aerosol formulations described herein include, for example, jet, vibrating porous plate, ultrasonic nebulizers and energized dry powder inhalers, that are able to nebulize the formulation into aerosol particle size predominantly in the size range from 1-5 μ . Predominantly in this application means that at least 70% but preferably more than 90% of all generated aerosol particles are within 1-5 μ range. A jet nebulizer works by air pressure to break a liquid solution into aerosol droplets. Vibrating porous plate nebulizers work by using a sonic vacuum produced by a rapidly vibrating porous plate to extrude a solvent droplet through a porous plate. An ultrasonic nebulizer works by a piezoelectric crystal that shears a liquid into small aerosol droplets. A variety of suitable devices are available, including, for example, AeroNebTM and AeroDoseTM vibrating porous plate nebulizers (AeroGen, Inc., Sunnyvale, California), Sidestream[®] nebulizers (Medic-Aid Ltd.,

West Sussex, England), Pari LC® and Pari LC Star® jet nebulizers (Pari Respiratory Equipment, Inc., Richmond, Virginia), and Aerosonic™ (DeVilbiss Medizinische Produkte (Deutschland) GmbH, Heiden, Germany) and UltraAire® (Omron Healthcare, Inc., Vernon Hills, Illinois) ultrasonic nebulizers.

[00374] In some embodiments, the pharmaceutical agent(s) can be formulated with oleaginous bases or ointments to form a semisolid composition with a desired shape. In addition to the pharmaceutical agent, these semisolid compositions can contain dissolved and/or suspended bactericidal agents, preservatives and/or a buffer system. A petrolatum component that may be included may be any paraffin ranging in viscosity from mineral oil that incorporates isobutylene, colloidal silica, or stearate salts to paraffin waxes. Absorption bases can be used with an oleaginous system. Additives may include cholesterol, lanolin (lanolin derivatives, beeswax, fatty alcohols, wool wax alcohols, low HLB (hydrophobellipophobe balance) emulsifiers, and assorted ionic and nonionic surfactants, singularly or in combination.

[00375] Controlled or sustained release transdermal or topical formulations can be achieved by the addition of time-release additives, such as polymeric structures, matrices, that are available in the art. For example, the compositions may be administered through use of hot-melt extrusion articles, such as bioadhesive hot-melt extruded film. The formulation can comprise a cross-linked polycarboxylic acid polymer formulation. A cross-linking agent may be present in an amount that provides adequate adhesion to allow the system to remain attached to target epithelial or endothelial cell surfaces for a sufficient time to allow the desired release of the compound.

[00376] An insert, transdermal patch, bandage or article can comprise a mixture or coating of polymers that provide release of the pharmaceutical agents at a constant rate over a prolonged period of time. In some embodiments, the article, transdermal patch or insert comprises water-soluble pore forming agents, such as polyethylene glycol (PEG) that can be mixed with water insoluble polymers to increase the durability of the insert and to prolong the release of the active ingredients.

[00377] Transdermal devices (inserts, patches, bandages) may also comprise a water insoluble polymer. Rate controlling polymers may be useful for administration to sites where pH change can be used to effect release. These rate controlling polymers can be applied using a continuous coating film during the process of spraying and drying with the active compound. In one embodiment, the coating formulation is used to coat pellets comprising the active ingredients that are compressed to form a solid, biodegradable insert.

[00378] A polymer formulation can also be utilized to provide controlled or sustained release. Bioadhesive polymers described in the art may be used. By way of example, a sustained-release

gel and the compound may be incorporated in a polymeric matrix, such as a hydrophobic polymer matrix. Examples of a polymeric matrix include a microparticle. The microparticles can be microspheres, and the core may be of a different material than the polymeric shell.

Alternatively, the polymer may be cast as a thin slab or film, a powder produced by grinding or other standard techniques, or a gel such as a hydrogel. The polymer can also be in the form of a coating or part of a bandage, stent, catheter, vascular graft, or other device to facilitate delivery of the pharmaceutical agent. The matrices can be formed by solvent evaporation, spray drying, solvent extraction and other methods known to those skilled in the art.

[00379] Kits with unit doses of one or more of the agents described herein, usually in oral or injectable doses, are provided. Such kits may include a container containing the unit dose, an informational package insert describing the use and attendant benefits of the drugs in treating disease, and optionally an appliance or device for delivery of the composition.

Methods of Treatment

[00380] In an aspect, the present disclosure provides compounds that inhibit KRas G12 mutants. In some cases, the method may inhibit KRas G12 mutants activity in a cell. In some cases, inhibiting KRas G12 mutants activity in a cell may include contacting the cell in which inhibition of KRas G12 mutants activity is desired with an effective amount of a compound of Formula (I), Formula (II), Formula (II*), Formula (III), or pharmaceutical compositions containing the compound or pharmaceutically acceptable salt thereof. In some cases, the contacting is in vitro. In some cases, the contacting is in vivo. As used herein, the term "contacting" refers to the bringing together of indicated moieties in an in vitro system or an in vivo system. For example, "contacting" a KRas G12D and/or other G12 mutants with a compound provided herein includes the administration of a compound provided herein to an individual or patient, such as a human, having KRas G12D and/or other G12 mutants, as well as, for example, introducing a compound provided herein into a sample containing a cellular or purified preparation containing the KRas G12D and/or other G12 mutants. In some cases, a cell in which inhibition of KRas G12D and/or other G12 mutants activity is desired is contacted with an effective amount of a compound of Formula (I) or Formula (II) or Formula (II*) or Formula (III) or pharmaceutically acceptable salt thereof to negatively modulate the activity of KRas G12D and/or other G12 mutants. In some cases, by negatively modulating the activity of KRas G12D and/or other G12 mutants, the methods described herein are designed to inhibit undesired cellular proliferation resulting from enhanced KRas G12D and/or other G12 mutants activity within the cell. The cells may be contacted in a single dose or multiple doses in accordance with a particular treatment regimen to effect the desired negative modulation of KRas G12D and/or

other G12 mutants. The ability of compounds to bind KRas G12D and/or other G12 mutants may be monitored in vitro using well known methods.

[00381] In some embodiments, the inhibitory activity of exemplary compounds in cells may be monitored, for example, by measuring the inhibition of KRas G12D and/or other G12 mutants activity of the amount of phosphorylated ERK.

[00382] In another aspect, methods of treating cancer in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a compound of Formula (I), Formula (II), Formula (II*), Formula (III), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising the compound or pharmaceutically acceptable salt thereof are provided. The compositions and methods provided herein may be used for the treatment of a KRas G12D and/or other G12 mutants-associated cancer in a patient in need thereof, comprising administering to the patient a therapeutically effective amount of a compound of Formula (I), Formula (II), Formula (II*), Formula (III), a pharmaceutically acceptable salt any one thereof, or a pharmaceutical composition comprising the compound or pharmaceutically acceptable salt thereof are provided. In some cases, the KRas G12D and/or other G12 mutants associated cancer is lung cancer. The compositions and methods provided herein may be used for the treatment of a wide variety of cancers including tumors such as lung, prostate, breast, brain, skin, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that may be treated by the compositions and methods of the invention include, but are not limited to tumor types such as astrocytic, breast, cervical, colorectal, endometrial, esophageal, gastric, head and neck, hepatocellular, laryngeal, lung, oral, ovarian, prostate and thyroid carcinomas and sarcomas. More specifically, these compounds can be used to treat: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor (nephroblastoma), lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma,

interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Biliary tract: gall bladder carcinoma, ampullary carcinoma, cholangiocarcinoma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma (pinealoma), glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma (serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma), granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia (acute and chronic), acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma (malignant lymphoma); Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. In some cases, the cancer is non-small cell lung cancer, small cell lung cancer, colorectal cancer, rectal cancer or pancreatic cancer. In some cases, the cancer is non-small cell lung cancer. In some cases, the concentration and route of administration to the patient will vary depending on the cancer to be treated. The compounds, pharmaceutically acceptable salts thereof and pharmaceutical compositions comprising such compounds and salts also may be co-administered with other anti-neoplastic compounds, e.g., chemotherapy, or used in combination with other treatments, such as radiation or surgical intervention, either as an adjuvant prior to surgery or post-operatively.

[00383] Also provided herein is a compound of Formula (I), Formula (II), Formula (II*), Formula (III), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof as defined herein for use in therapy.

[00384] Also provided herein is a compound of Formula (I), Formula (II), Formula (II*), Formula (III), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof as defined herein for use in the treatment of cancer.

[00385] Also provided herein is a compound of Formula (I), Formula (II), Formula (II*), Formula (III), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof for use in the inhibition of KRas G12D and/or other G12 mutants.

[00386] Also provided herein is a compound of Formula (I), Formula (II), Formula (III), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof as defined herein, for use in the treatment of a KRas G12D and/or other G12 mutants -associated disease or disorder.

[00387] Also provided herein is the use of a compound of Formula (I), Formula (II), Formula (II*), Formula (III), or a pharmaceutically acceptable salt thereof, as defined herein in the manufacture of a medicament for the treatment of cancer.

[00388] Also provided herein is a use of a compound of Formula (I), Formula (II), Formula (II*), Formula (III), or a pharmaceutically acceptable salt thereof, as defined herein in the manufacture of a medicament for the inhibition of activity of KRas G12D and/or other G12 mutants.

[00389] Also provided herein is the use of a compound of Formula (I), Formula (II), Formula (III), or a pharmaceutically acceptable salt thereof, as defined herein, in the manufacture of a medicament for the treatment of a KRas G12D and/or other G12 mutants-associated disease or disorder.

[00390] In another aspect, the present disclosure provides a method for treating cancer in a patient in need thereof, the method comprising (a) determining that cancer is associated with a KRas G12D mutation and/or other G12 mutants (e.g., a KRas G12D and/or other G12 mutants-associated cancer) (e.g., as determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit); and (b) administering to the patient a therapeutically effective amount of a compound of Formula (I), Formula (II), Formula (II*), Formula (III), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof.

[00391] The compounds described herein can be used in the preparation of medicaments for the prevention or treatment of diseases or conditions. In addition, a method for treating any of the diseases or conditions described herein in a subject in need of such treatment, involves administration of pharmaceutical compositions containing at least one compound described herein, or a pharmaceutically acceptable salt, pharmaceutically acceptable prodrug, or pharmaceutically acceptable solvate thereof, in therapeutically effective amounts to said subject.

[00392] The compositions containing the compound(s) described herein can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, the compositions are administered to a patient already suffering from a disease or condition, in an amount sufficient to cure or at least partially arrest the symptoms of the disease or condition. Amounts effective for this use will depend on the severity and course of the disease or condition, previous therapy, the patient's health status, weight, and response to the drugs, and the judgment of the treating physician.

[00393] In prophylactic applications, compositions containing the compounds described herein are administered to a patient susceptible to or otherwise at risk of a particular disease, disorder or condition. Such an amount is defined to be a "prophylactically effective amount or dose." In this use, the precise amounts also depend on the patient's state of health, weight, and the like. When used in a patient, effective amounts for this use will depend on the severity and course of the disease, disorder or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician.

[00394] In the case wherein the patient's condition does not improve, upon the doctor's discretion the administration of the compounds may be administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disease or condition.

[00395] Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, the dosage or the frequency of administration, or both, can be reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. Patients can, however, require intermittent treatment on a long-term basis upon any recurrence of symptoms.

[00396] The amount of a given agent that will correspond to such an amount will vary depending upon factors such as the particular compound, disease or condition and its severity, the identity (e.g., weight) of the subject or host in need of treatment, but can nevertheless be determined in a manner recognized in the field according to the particular circumstances surrounding the case, including, e.g., the specific agent being administered, the route of administration, the condition being treated, and the subject or host being treated. In general, however, doses employed for adult human treatment will typically be in the range of about 0.02 - about 5000 mg per day, in some embodiments, about 1 - about 1500 mg per day. The desired dose may conveniently be presented in a single dose or as divided doses administered simultaneously (or over a short period of time) or at appropriate intervals, for example as two, three, four or more sub-doses per day.

[00397] The pharmaceutical composition described herein may be in unit dosage forms suitable for single administration of precise dosages. In unit dosage form, the formulation is divided into unit doses containing appropriate quantities of one or more compound. The unit dosage may be in the form of a package containing discrete quantities of the formulation. Non-limiting examples are packaged tablets or capsules, and powders in vials or ampoules. Aqueous suspension compositions can be packaged in single-dose non-reclosable containers.

Alternatively, multiple-dose reclosable containers can be used, in which case it is typical to include a preservative in the composition. By way of example only, formulations for parenteral injection may be presented in unit dosage form, which include, but are not limited to ampoules, or in multi-dose containers, with an added preservative.

[00398] Toxicity and therapeutic efficacy of such therapeutic regimens can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the determination of the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between the toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD₅₀ and ED₅₀. Compounds exhibiting high therapeutic indices are preferred. The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with minimal toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized.

[00399] In certain embodiments, the invention provides a method of treating or preventing a disease, state or condition in a patient in need thereof comprising administering to the patient an effective amount of a compound of any one of embodiments of the invention or a pharmaceutically acceptable salt thereof. The disease, state or condition may be selected from a group as described elsewhere herein.

Bifunctional Compounds

[00400] In some embodiments, compounds herein can adopt to selectively eliminate an over activated KRas signaling which is induced by KRas mutations by directly binding with the mutated KRas protein, either by stabilizing its GDP bound form (the inactive form) or by blocking the interaction between GTP bound form and its downstream target protein. In some embodiments, another way is to hijack the protein degradation mechanism in a cell and leverage E3 ligases' (like VHL, CRBN or IAPs) substrate specificity through a bi-functional molecule called Proteolysis targeting chimera (PROTAC) (Winter GE, Buckley DL, Paulk J, Roberts JM, Souza A, Dhe-Paganon S, Bradner JE. DRUG DEVELOPMENT. Phthalimide conjugation as a strategy for in

vivo target protein degradation. Science. 2015 Jun 19; 348 (6241): 1376-81), which can bind with both mutated KRas protein and E3 ligase, create interactions between those two proteins and induce KRas degradation.

[00401] Disclosed herein is a bifunctional compound composed of a target protein (i.e., KRAS G12D)-binding moiety and an E3 ubiquitin ligase-binding moiety, which may induce proteasome-mediated degradation of selected proteins. In some embodiments, the bifunctional compound comprises a target protein (i.e., KRAS G12D)-binding moiety and an E3 ubiquitin ligase-binding moiety known in the art. In some embodiments, disclosed herein is the use of the compound disclosed herein in the preparation of degrading a target protein compound by using chemical modification of the compound disclosed herein. In some cases, the target protein-binding moiety is derived from a compound of Formula (I), Formula (II), Formula (II*), or Formula (III).

Preparation of Compounds

[00402] The compounds of the present disclosure can generally be prepared in a number of ways well known to those skilled in the art of organic synthesis. By way of example, compounds of the present disclosure can be synthesized using the methods described herein, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereof as appreciated by those skilled in the art. The compounds of the present disclosure may be prepared as described in the schemes and examples described elsewhere herein.

[00403] The following examples further illustrate the invention but, of course, should not be construed as in any way limiting its scope


EXAMPLES

[00404] The following synthetic schemes are provided for purposes of illustration, not limitation. The following examples illustrate the various methods of making compounds described herein. It is understood that one skilled in the art may be able to make these compounds by similar methods or by combining other methods known to one skilled in the art. It is also understood that one skilled in the art would be able to make, in a similar manner as described below by using the appropriate starting materials and modifying the synthetic route as needed. In general, starting materials and reagents can be obtained from commercial vendors or synthesized according to sources known to those skilled in the art or prepared as described herein.

[00405] The present disclosure provides processes for preparing the compounds described herein (described in greater detail below).


General Schemes

Scheme IA




[00406] Treatment of R_1 with methyl 2,4-dichloro-6-methylpyrimidine-5-carboxylate in the presence of a base such as DIEA in DCM can afford **IA-a**, which can be deprotonated with LDA or LiHMDS at $-78\text{ }^\circ\text{C}$ in THF. **IA-b** can be formed by addition of a solution of a ketone in THF at $-78\text{ }^\circ\text{C}$ and the reaction is quenched at $-78\text{ }^\circ\text{C}$. Ester **IA-b** can be reduced by DIBAL or LiAlH_4 to diol **IA-c**, which can be cyclized into **IA-d** via Mitsunobu reaction or in the presence of BuLi and TsCl in THF at $-78\text{ }^\circ\text{C}$. Compound **I** can be prepared via Pd mediated coupling reactions or in the presence of a base such as NaH or LiHMDS.

Scheme IB




[00407] Ethyl 4-chloro-6-methyl-2-(methylthio)pyrimidine-5-carboxylate can be deprotonated with LDA or LiHMDS at $-78\text{ }^\circ\text{C}$ in THF. **IB-a** can be formed by addition of a solution of a ketone in THF at $-78\text{ }^\circ\text{C}$ and the reaction is quenched at $-78\text{ }^\circ\text{C}$. Ester **IB-a** can be reduced by DIBAL or LiAlH_4 to diol **IB-b**, which can be cyclized into **IB-c** via Mitsunobu reaction or in the presence of BuLi and TsCl in THF at $-78\text{ }^\circ\text{C}$. Treatment of R_1 with **IB-c** can afford **IB-d** in the presence of a base such as DIEA in DCM can afford **IB-d**, which can be oxidized to sulfone **IB-e**. Compound **I** can be prepared via Pd mediated coupling reactions or in the presence of a base such as NaH or LiHMDS.

Scheme IIA




[00408] Treatment of t-Butyl-sulfinamide and a ketone in the presence of $\text{Ti}(\text{OEt})_4$ can afford **II A-a**. A solution of **II A-a** in THF can be added to a solution of **IA-a**, deprotonated with LDA or LiHMDS at -78°C in THF, and the reaction can be quenched at -78°C to afford **II A-b**. Ester **II A-b** can be reduced by DIBAL or LiAlH_4 to **II A-c**, which can be cyclized into **II A-d** via a sequence of removal of sulfinamide under HCl, chlorination of OH and cyclization in the presence of a base such as NaOH (ref. Garcia, D.; Moreno, B.; Soler, T.; Foubelo, F.; Yus, M. *Tetrahedron Lett.* 2009, 50, 4710). R_3 of **II A-e** can be introduced via reductive amination or alkylation. Compound **II** can be prepared via Pd mediated coupling reactions or in the presence of a base such as NaH or LiHMDS.

Scheme IIB



[00409] Ethyl 4-chloro-6-methyl-2-(methylthio)pyrimidine-5-carboxylate can be deprotonated with LDA or LiHMDS at -78°C in THF. **II B-a** can be formed by addition of a solution of **II A-a** in THF at -78°C and the reaction is quenched at -78°C . Ester **II B-a** can be reduced by DIBAL or LiAlH_4 to diol **II B-b**, which can be cyclized into **II B-c** as described in **Scheme IIA** above. R_3 of **II B-d** can be introduced via reductive amination or alkylation. Treatment of **R1** with **II B-d** can afford **II B-e** in the presence of a base such as DIEA in DCM. **II B-e** can be oxidized to sulfoxide **II B-f**, which can be converted to **II** in the presence of a base such as NaH or LiHMDS.

Example 1: Exemplary synthesis of 4'-(azepan-1-yl)-4-chloro-2'-(((2*R*,7*aS*)-2-fluorotetrahydro-1*H*-pyrrolizin-7*a*(5*H*)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-*d*]pyrimidine] (**compound 1**)



[00410] Step 1: To a solution of methyl 2,4-dichloro-6-methylpyrimidine-5-carboxylate (1.0 g, 4.52 mmol, 1.0 eq) in ACN (7 mL) was added DIPEA (1.17 g, 9.05 mmol, 2.0 eq) and azepane (449 mg, 4.52 mmol, 1.0 eq) at 20 °C. Then the reaction was stirred at 20 °C for 2 hrs. TLC showed that the reaction was completed. The reaction was concentrated under vacuum. The residue was poured into water (20 mL). The aqueous phase was extracted with EtOAc (5 mL * 2). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under vacuum. The residue was purified by column chromatography on silica gel (PE: EtOAc = 10:1) to give compound **1a** (1.1 g, 3.88 mmol, 85.7% yield) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 3.88 (s, 3H), 3.49-3.52 (m, 4H), 2.36 (s, 3H), 1.75-1.85 (m, 4H), 1.50-1.60 (m, 4H).

[00411] Step 2: To a solution of LDA (2 M, 2.11 mL, 1.2 eq) in THF (30 mL) was added compound **1a** (1.0 g, 3.52 mmol, 1.0 eq) at -60°C. The mixture was stirred at -60 °C for 30 mins. 4-Chloro-2,3-dihydro-1*H*-inden-1-one (587 mg, 3.52 mmol, 1.0 eq) was added at -60 °C. The mixture was stirred at -60 °C for 1 hr. The reaction mixture was quenched by addition HCl (5%, 50 mL) at 0°C, and then extracted with EtOAc (20 mL * 2). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (PE: EtOAc = 5:1) to give compound **1b** (0.5 g, 1.20 mmol, 33.9% yield) as a yellow solid. LCMS calcd for C₂₂H₂₅Cl₂N₃O₃ (M+H)⁺m/z = 450.1; found 450.1. ¹H NMR (400 MHz, CDCl₃) δ 7.22 (d, *J*=8.0 Hz, 1H), 7.13 (t, *J*=7.6 Hz, 1H), 7.03 (d, *J*=7.6 Hz, 1H), 6.06 (s, 1H), 3.69 (s, 3H), 3.47-3.54(m, 4H), 3.08-3.11 (m, 2H), 2.76-2.99 (m, 2H), 2.22-2.26 (m, 2H), 1.70-1.80 (m, 4H), 1.45-1.60 (m, 4H).


[00412] Step 3: To a solution of compound **1b** (490 mg, 1.09 mmol, 1.0 eq) in DCM (5 mL) was added DIBAL-H (1 M, 3.26 mL, 3.0 eq) at 0 °C. Then the reaction was stirred at 0 °C for 1 hr. The reaction mixture was quenched by H₂O (20 mL) at 25 °C. The mixture was extracted with DCM (10 mL * 2). The combined organic layers were washed with brine (15 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (PE/EtOAc=80/1 to 8/1 to give compound **1c** (200 mg, 474 μmol, 43.5% yield) as white solid. LCMS calcd for C₂₁H₂₅Cl₂N₃O₂ (M+H)⁺m/z = 422.1; found 422.1. ¹H NMR (400 MHz, CD₃OD) δ 7.25 (d, *J*=8.0 Hz, 1H), 7.19 (t, *J*=7.6 Hz, 1H), 7.09 (d, *J*=7.6 Hz, 1H), 4.40-4.44 (m, 1H), 4.29-4.32 (m, 1H), 3.78-3.81 (m, 4H), 3.25-3.26 (m, 1H), 2.96-3.03 (m, 2H), 2.84-2.88 (m, 1H), 2.56-2.59 (m, 1H), 2.01-2.04 (m, 1H), 1.75-1.90 (m, 4H), 1.55-1.65 (m, 4H).

[00413] Step 4: To a solution of compound **1c** (200 mg, 474 μmol, 1 eq) in THF (10 mL) was added PPh₃ (149 mg, 568 μmol, 1.2 eq) and DIAD (115 mg, 568 μmol, 1.2 eq) at 0 °C under N₂. Then the reaction was stirred at 0 °C for 2 hrs under N₂. LCMS showed the reaction was completed. The reaction mixture was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (PE: EtOAc = 20:1) to give compound **1d** (100 mg, 247 μmol, 52.2% yield) as a yellow solid. C₂₁H₂₃Cl₂N₃O (M+H)⁺m/z = 404.1; found 404.1. ¹H NMR (400 MHz, CDCl₃) δ 7.31 (d, *J*=8.0 Hz, 1H), 7.17 (t, *J*=7.6 Hz, 1H), 7.12 (d, *J*=7.6 Hz, 1H), 4.65-4.71 (m, 1H), 4.51-4.53 (m, 1H), 3.50-3.65 (m, 4H), 3.07-3.17 (m, 3H), 2.96-3.00 (m, 1H), 2.36-2.38 (m, 1H), 2.15-2.20 (m, 1H), 1.70-1.75 (m, 4H), 1.50-1.65 (m, 4H).

[00414] Step 5: To a solution of ((2*R*,7*aS*)-2-fluorotetrahydro-1*H*-pyrrolizin-7*a*(5*H*)-yl)methanol (37.8 mg, 237 μmol, 1.2 eq) in THF (2 mL) was added NaH (15.8 mg, 396 μmol, 60% purity, 2.0 eq) at 0 °C. The reaction was stirred at 0 °C for 30 min. Then a solution of compound **1d** (80 mg, 198 μmol, 1.0 eq) in THF (0.5 mL) was added to the above solution at 0 °C. The reaction was continued to stir at 60 °C for 8 hrs. LC-MS showed 11% of starting material remained and 52.9% of desired compound was detected. The reaction was quenched with H₂O (10 mL) at 0 °C. The aqueous phase was extracted with EtOAc (5 mL * 2). The combined organic phase was washed with brine (5 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by prep-TLC (SiO₂, DCM: MeOH = 6:1) to give 4'-(azepan-1-yl)-4-chloro-2'-(((2*R*,7*aS*)-2-fluorotetrahydro-1*H*-pyrrolizin-7*a*(5*H*)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-*d*]pyrimidine] (**compound 1**, 11.0 mg, 20.8 μmol, 10.5% yield) as yellow solid. C₂₉H₃₆ClF₂N₄O₂ (M+H)⁺m/z = 526.25; found 527.2. ¹H NMR (400 MHz, CDCl₃) δ 7.10 (d, *J*=8.0 Hz, 1H), 6.90-7.00 (m, 2H), 5.00-5.40 (m, 1H), 4.50-4.53 (m, 1H), 4.32-4.36 (m, 1H), 3.93-3.95 (m, 1H), 3.79-3.81 (m, 1H), 3.38-3.41 (m, 4H), 3.05-3.15 (m, 2H), 2.90-

3.00 (m, 2H), 2.85-2.87(m, 1H), 2.70-2.80 (m, 2H), 2.15-2.25 (m, 1H), 1.95-2.10 (m, 4H), 1.50-1.80 (m, 6H), 1.30-1.50 (m, 6H).

Example 2. Exemplary synthesis of 4'-((1*R*,5*S*)-3,8-diazabicyclo[3.2.1]octan-3-yl)-4-chloro-2'-(((2*R*,7*aS*)-2-fluorotetrahydro-1*H*-pyrrolizin-7*a*(5*H*)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-*d*]pyrimidine] (compound **2**)



[00415] Step 1. Synthesis of tert-butyl 3-(2-chloro-5-methoxycarbonyl-6-methyl-pyrimidin-4-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2a**). To a solution of methyl 2,4-dichloro-6-methyl-pyrimidine-5-carboxylate (200.0 mg, 0.9 mmol) in MeCN (4 mL) was added a mixture of DIEA (233.88 mg, 1.81 mmol) and tert-butyl 3,8-diazabicyclo[3.2.1]octane-3-carboxylate (192.08 mg, 0.9 mmol) in DMF (1 mL) at 0 °C. The reaction mixture was stirred at 0 °C for 30 min under Ar. The solution was diluted with EtOAc (5 ml) and washed with H₂O (5ml x3) and brine (5 ml). The organic phase was dried over Na₂SO₄ and concentrated. The residue was purified by silica gel chromatography (eluting with EtOAc:PE=1:7) to give tert-butyl 3-(2-chloro-5-methoxycarbonyl-6-methyl-pyrimidin-4-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2a**, 280 mg, 0.71 mmol, 78% yield) as a white solid. LCMS (ESI): *m/z* calcd for C₁₈H₂₅ClN₄O₄⁺H: 397.87, found: 397.2.

[00416] Step 2. Synthesis of tert-butyl 3-[2-[[[(2*R*,8*S*)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-5-methoxycarbonyl-6-methyl-pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2b**). To a degassed solution of tert-butyl 3-(2-chloro-5-methoxycarbonyl-6-methyl-pyrimidin-4-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2a**, 400.0 mg, 1.01 mmol) in toluene (15 mL) were added [(2*R*,8*S*)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (332.67 mg, 2.09 mmol), Ruphos (97.5 mg, 0.21 mmol), Pd₂(dba)₃ (95.71 mg, 0.1 mmol) and Cs₂CO₃ (851.06 mg, 2.61mmol). The degassed mixture was stirred at 110 °C for 16 h under Ar. The reaction mixture was concentrated under reduced pressure to give crude product which was purified by silica gel chromatography column (EtOAc:PE = 1:1

to DCM:MeOH=10:1) to obtain tert-butyl 3-[2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-5-methoxycarbonyl-6-methyl-pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2b**, 460mg, 0.88 mmol, 88% yield) as a sticky brown solid. LCMS (ESI): m/z calcd for C₂₆H₃₈FN₅O₅⁺H: 520.2, found: 520.2.

[00417] Step 3. Synthesis of tert-butyl 3-[6-[(4-chloro-1-hydroxy-indan-1-yl)methyl]-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-5-methoxycarbonyl-pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2c**). To a solution of tert-butyl 3-[5-methoxycarbonyl-6-methyl-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2b**, 150.0 mg, 0.29 mmol) in THF (15 mL) at -70 °C was added LDA (0.72 mL, 1.44 mmol, 2 mol/L in THF) dropwise. The reaction mixture was stirred between -70 °C to -20 °C for 1.5 h under Ar. Then a solution of 4-chloroindan-1-one (240.47 mg, 1.44 mmol) in THF (3 mL) was added dropwise to the above resulting solution at -70 °C. The reaction mixture was stirred between -70 °C to 0 °C for 3 h under Ar. The reaction mixture was quenched by aqueous NH₄Cl solution at -50 °C, and extracted with EtOAc, washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated under vacuo to give the crude product which was purified on prep-TLC (DCM:MeOH=10:1) to give tert-butyl 3-[6-[(4-chloro-1-hydroxy-indan-1-yl)methyl]-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-5-methoxycarbonyl-pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2c**, 75 mg, 0.11 mmol, 38% yield) as a light yellow sticky solid. LCMS (ESI): m/z calcd for C₃₅H₄₅ClFN₅O₆⁺H: 686.1, found: 686.2.


[00418] Step 4. Synthesis of tert-butyl 3-[6-[(4-chloro-1-hydroxy-indan-1-yl)methyl]-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-5-(hydroxymethyl)pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2d**). To a degassed solution of tert-butyl 3-[6-[(4-chloro-1-hydroxy-indan-1-yl)methyl]-5-methoxycarbonyl-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2c**, 120.0 mg, 0.17 mmol) in anhydrous THF (8 mL) at -70 °C was added dropwise DIBAL-H (2.91 mL, 4.37 mmol) at -70 °C. The mixture was warmed to 0 °C naturally and stirred for 16 h under Ar. The reaction mixture was quenched by aqueous NH₄Cl solution at 0 °C, and extracted with EtOAc, washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated under *vacuo* to give crude product which was purified on Pre-TLC (DCM:MeOH = 10:1) to give tert-butyl 3-[6-[(4-chloro-1-hydroxy-indan-1-yl)methyl]-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-5-(hydroxymethyl)pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

(**2d**, 60 mg, 0.091 mmol, 52.1% yield) as a colorless sticky solid. LCMS (ESI): m/z calcd for $C_{34}H_{45}ClFN_5O_5^+H$: 658.3, found: 658.3.

[00419] Step 5. Synthesis of tert-butyl 3-[4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl] methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2e**). To a degassed mixture of tert-butyl 3-[6-[(4-chloro-1-hydroxy-indan-1-yl)methyl]-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-5-(hydroxymethyl)pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2d**, 70.0 mg, 0.11 mmol) and DIPEA (0.04 mL, 0.23 mmol) in anhydrous THF (5 mL) at 0 °C was added MsCl (60.92 mg, 0.53 mmol) dropwise. The degassed mixture was warmed to 10 °C and stirred for 16 h under Ar. The mixture was quenched by aqueous $NaHCO_3$ solution and extracted with EtOAc, washed with water and brine. The organics was dried over anhydrous Na_2SO_4 , filtered and concentrated under vacuo. The residue was purified by Pre-TLC (DCM:MeOH = 15:1) to give crude product (**2e**, 65 mg) as a sticky brown solid. LCMS (ESI): m/z calcd for $C_{34}H_{43}ClFN_5O_4+H$: 640.3, found: 640.2.

[00420] Step 6. Synthesis of 4'-((1R,5S)-3,8-diazabicyclo[3.2.1]octan-3-yl)-4-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-d]pyrimidine] (**2**). To a solution of tert-butyl 3-[4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydro-pyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (64.0 mg, 0.1 mmol) in DCM (1.5 mL) was added TFA (2.95 mL, 39.77 mmol). The mixture was stirred at rt. for 20 min. The mixture was concentrated under vacuo to afford the crude product which was purified by Prep-HPLC (eluted with CH_3CN in H_2O (0.1 % TFA) from 5.0% to 95%). Then the preparation solution is treated with 1N hydrochloric acid to replace TFA to give 4'-chloro-4-(3,8-diazabicyclo[3.2.1]octan-3-yl)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane];dihydrochloride (12.32 mg, 0.0141 mmol, 14% yield) as a white solid. LCMS (ESI): m/z calcd for $C_{29}H_{35}ClFN_5O_2+H$: 540.2, found: 540.2. ^1H NMR (400 MHz, CD_3OD) δ 7.37 (m, 3H), 5.59 (dd, $J = 52.0, 13.6$ Hz, 1H), 5.07 (d, $J = 13.9$ Hz, 1H), 4.69 – 4.35 (m, 3H), 4.35 – 4.06 (m, 3H), 3.90 (m, 6H), 3.42 (s, 2H), 3.26 – 3.07 (m, 2H), 3.00 (s, 1H), 2.88 – 1.99 (m, 12H).

Example 3. Exemplary synthesis of (4'-[(1R,5S)-3,8-diazabicyclo[3.2.1]octan-3-yl]-2'-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[2H-indeno[1,2-c]pyrazole-4,7'-5,8-dihydropyrano[4,3-d]pyrimidine] (compound **3**)



[00421] Step 1. Synthesis of 4-hydrazono-2,4-dihydroindeno[1,2-c]pyrazole (**3a**). To a solution of indane-1,3-dione (5.00 g, 34.2 mmol, 1.0 eq) in THF (50 mL) was added DMF-DMA (4.15 g, 34.9 mmol, 1.02 eq) dropwise. The resulting mixture was stirred at room temperature for 1 h. TLC showed SM was consumed. The mixture was concentrated in *vacuo* to afford crude 2-(dimethylaminomethylene)indane-1,3-dione (**3a**, 6.87 g, 34.1 mmol, crude) as a black solid, which was used in next step directly without further purification. LCMS calcd for $C_{12}H_{12}NO_2$ ($M+H$)⁺ m/z = 202.1; found: 202.1.

[00422] Step 2. Synthesis of 4-hydrazono-2,4-dihydroindeno[1,2-c]pyrazole (**3b**). To a solution of 2-(dimethylaminomethylene)indane-1,3-dione (**3a**, 6.87 g, 34.1 mmol, crude) in AcOH (10.0 mL) was added hydrazine hydrate (10.0 mL, 205 mmol, 5.0 eq) drop wise. Then the mixture was stirred at 90 °C for 16 h and the solvent was removed in *vacuo*. The residue was suspended in water (30 mL), and the solid was collected by filtration, washed with water (90 mL) and aq. $NaHCO_3$ (90 mL), and dried in *vacuo* to afford 4-hydrazono-2,4-dihydroindeno[1,2-c]pyrazole (**3b**, 5.23 g, 30.7 mmol, 89.9% yield over two steps) as a yellow solid. LCMS calcd for $C_{10}H_9N_4$ ($M+H$) m/z = 185.0; found: 185.0.

[00423] Step 3. Synthesis of 2H-indeno[1,2-c]pyrazol-4-one (**3c**). A solution of $NaIO_4$ (9.59 g, 44.8 mmol, 2.0 eq) in water (100 mL) and EtOAc (100 mL) was stirred at room temperature for 5 min. Then 2,4-dihydroindeno[1,2-c]pyrazol-4-yl diazene (**3b**, 4.13 g, 22.4 mmol) was added, the resulting mixture was stirred at room temperature for 2h. The mixture was extracted with EtOAc (100 mL*3), the combined organic phase was concentrated and purified by flash chromatography (silica gel, eluted with EtOAc in PE 0 to 30%) to give the 2H-indeno[1,2-c]pyrazol-4-one (**3c**,

1.74 g, 10.2 mmol, 45.6% yield) as a white solid. LCMS calcd for $C_{10}H_7N_2O$ (M+H) m/z = 171.0; found: 171.0.

[00424] Step 4. Synthesis of 2-(2-trimethylsilylethoxymethyl)indeno[1,2-c]pyrazol-4-one (**3d**). To a solution of 2H-indeno[1,2-c]pyrazol-4-one (800 mg, 4.70 mmol, 1.0 eq) in DMF (12 mL) was added NaH (**3c**, 169 mg, 7.05 mmol, 1.5 eq) at 0°C, the mixture was stirred at the same temperature for 30 min, then SEMCl (1.17 g, 7.05 mmol, 1.5 eq) was added. The resulting mixture was allowed to warm to room temperature and stirred for 1h. The reaction was quenched with NH_4Cl (aq) and extracted with EtOAc (50 mL*3), the combined organic phase was concentrated and purified by flash chromatography (silica gel, eluted with EtOAc in PE 0 to 25%) to give a mixture of 2-(2-trimethylsilylethoxymethyl)indeno[1,2-c]pyrazol-4-one and its regioisomer (not shown) (**3d**, 1.38 g, 4.60 mmol, 97% yield) as a yellow solid. LCMS calcd for $C_{16}H_{21}N_2O_2Si$ (M+H) m/z = 301.1; found: 301.0.

[00425] Step 5. Synthesis of tert-butyl (1S,5R)-3-[2-chloro-6-[[4-hydroxy-2-(2-trimethylsilylethoxymethyl)indeno[1,2-c]pyrazol-4-yl]methyl]-5-methoxycarbonyl-pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3e**). To a solution of tert-butyl rac-(1S,5R)-3-(2-chloro-5-methoxycarbonyl-6-methyl-pyrimidin-4-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**2a**, 800 mg, 2.02 mmol, 1.0 eq) in dry THF (6 mL) was added LDA (1.21 mL, 2.42 mmol, 1.2 eq, 2M in THF/n-heptane) dropwise at -70 °C under N_2 , the mixture was stirred at the same temperature for 0.5 h, then 2-(2-trimethylsilylethoxymethyl)indeno[1,2-c]pyrazol-4-one (**3d**, 667 mg, 2.22 mmol, 1.1 eq) in dry THF (6 mL) was added drop wise. The mixture was stirred at the same temperature for 1h and then quenched with NH_4Cl (aq.). The residue was extracted with EtOAc (20 mL*3) and the organic phase was concentrated in vacuum. The residue was purified by flash chromatography (silica gel, Eluant with EtOAc in PE 0 to 50%) to afford tert-butyl (1S,5R)-3-[2-chloro-6-[[4-hydroxy-2-(2-trimethylsilylethoxymethyl)indeno[1,2-c]pyrazol-4-yl]methyl]-5-methoxycarbonyl-pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3e**, 1.20 g, 1.72 mmol, 85.4% yield) as a white solid. LCMS calcd for $C_{34}H_{46}ClN_6O_6Si$ (M+H)⁺ m/z = 697.3; found: 697.5.

[00426] Step 6. Synthesis of 4'-((1R,5S)-3,8-diazabicyclo[3.2.1]octan-3-yl)-2'-((((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-5',8'-dihydro-2H-spiro[indeno[1,2-c]pyrazole-4,7'-pyrano[4,3-d]pyrimidine] (**3f**). To a solution of tert-butyl (1R,5S)-3-[2-chloro-6-[[4-hydroxy-2-(2-trimethylsilylethoxymethyl)indeno[1,2-c]pyrazol-4-yl]methyl]-5-methoxycarbonyl-pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3e**, 670 mg, 0.960 mmol, 1.0 eq) in dry DCM (5 mL) was added DIBAL-H (2.56 mL, 3.84 mmol, 4.0 eq, 1.5 M in toluene) at 0 °C under N_2 , the resulting mixture was stirred at same temperature for 1 h and then quenched with H_2O . The mixture was extracted with EtOAc (50 mL*3) and the organic

phase was concentrated and purified by flash column chromatography (silica gel, eluting EtOAc in PE 0 to 50%). The desired fractions were concentrated to dryness in *vacuo* to afford tert-butyl (1R,5S)-3-[2-chloro-5-(hydroxymethyl)-6-[[4-hydroxy-2-(2-trimethylsilylethoxymethyl)indeno[1,2-c]pyrazol-4-yl]methyl]pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3f**, 320 mg, 0.478 mmol, 49.8% yield). LCMS calcd for $C_{33}H_{46}ClN_6O_5Si$ ($M+H$)⁺ m/z = 669.3; found: 669.2.


[00427] Step 7. Synthesis of tert-butyl (1R,5S)-3-[2-chloro-2'-(2-trimethylsilylethoxymethyl)spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-indeno[1,2-c]pyrazole]-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3g**). To a solution of tert-butyl (1R,5S)-3-[2-chloro-5-(hydroxymethyl)-6-[[4-hydroxy-2-(2-trimethylsilylethoxymethyl)indeno[1,2-c]pyrazol-4-yl]methyl]pyrimidin-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3f**, 67.0 mg, 0.100 mmol, 1.0 eq) in dry THF (1.50 mL) was added n-BuLi (0.042 mL, 0.11 mmol, 1.05 eq, 2.5 M in THF) under N₂ at -78 °C, then stirred for 1 h. A solution of TsCl (21.0 mg, 0.110 mmol, 1.05 eq) in dry THF (1.50 mL) was added slowly, the mixture was allowed warm to 0 °C and stirred for 30 min, then it was cooled to -78 °C and n-BuLi (0.042 mL, 0.11 mmol, 1.05 eq, 2.5 M in THF) was added. After it was stirred at the same temperature for 10 min, the mixture was allowed to warm to room temperature and stirred for 1h. Then it was quenched by the addition of H₂O (2 mL) and extracted with EtOAc (10 mL*3), the organic phase was concentrated and purified by flash column chromatography (silica gel, eluting EtOAc in PE 0 to 25%). The desired fractions were concentrated to dryness in *vacuo* to afford tert-butyl (1R,5S)-3-[2-chloro-2'-(2-trimethylsilylethoxymethyl)spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-indeno[1,2-c]pyrazole]-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3g**, 49.0 mg, 0.0752 mmol, 75.2% yield) as a yellow oil. LCMS calcd for $C_{33}H_{44}ClN_6O_4Si$ ($M+H$)⁺ m/z = 651.3; found: 651.3.

[00428] Step 8. Synthesis of tert-butyl (1R,5S)-3-[2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-2'-(2-trimethylsilylethoxymethyl)spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-indeno[1,2-c]pyrazole]-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3h**). A mixture of [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (35.9 mg, 0.230 mmol, 3.0 eq), Pd₂(dba)₃ (6.89 mg, 0.0100 mmol, 0.1 eq), Ruphos (7.02 mg, 0.0200 mmol, 0.2 eq), Cs₂CO₃ (738 mg, 0.230 mmol, 3.0 eq) and tert-butyl (1R,5S)-3-[2-chloro-2'-(2-trimethylsilylethoxymethyl)spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-indeno[1,2-c]pyrazole]-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3g**, 49.0 mg, 0.0800 mmol, 1.0 eq) in Toluene (3 mL) was charged with N₂ for 3 times. The resulting mixture was heated to 110 °C for 16 h. The resulting solution was cooled to rt and filtered, and then the filtrate was concentrated and

purified by flash column chromatography (silica gel, eluting MeOH in DCM 0 to 10%). The desired fractions were concentrated to dryness in *vacuo* to afford tert-butyl (1R,5S)-3-[2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-2'-(2-trimethylsilylethoxymethyl)spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-indeno[1,2-c]pyrazole]-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**3h**, 51.0 mg, 0.0659 mmol, 87.6% yield) as a yellow solid. LCMS calcd for $C_{41}H_{57}FN_7O_5Si(M+H)^+$ $m/z = 774.4$; found: 774.6.

[00429] Step 9. Synthesis of 4'-((1R,5S)-3,8-diazabicyclo[3.2.1]octan-3-yl)-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-5',8'-dihydro-2H-spiro[indeno[1,2-c]pyrazole-4,7'-pyrano[4,3-d]pyrimidine] (**3**). To a solution of tert-butyl rac-(1R,5S)-3-[2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-2'-(2-trimethylsilylethoxymethyl)spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-indeno[1,2-c]pyrazole]-4-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (51.0 mg, 0.0700 mmol, 1.0 eq) in MeCN (1 mL) was added HCl in dioxane (1.00 mL, 4.00 mmol, 4 M). The resulting mixture was stirred at room temperature for 5h. The solvent was removed by N_2 and the residue was purified by Prep-HPLC on a C18 column (5 μ M, 50 x 150 mm) with mobile phase: H_2O (0.1% NH_4HCO_3) / MeOH at flow rate : 35 mL/min to afford 4'-[(1R,5S)-3,8-diazabicyclo[3.2.1]octan-3-yl]-2'-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[2H-indeno[1,2-c]pyrazole-4,7'-5,8-dihydropyrano[4,3-d]pyrimidine] (7.28 mg, 0.0134 mmol, 20.3% yield) as a yellow solid. LCMS calcd for $C_{30}H_{35}FN_7O_2(M+H)^+$ $m/z = 544.3$; found: 544.5. ^1H NMR (400 MHz, CD_3OD) δ 7.63 (d, $J = 7.2$ Hz, 1H), 7.53 (d, $J = 7.6$ Hz, 1H), 7.40 - 7.47 (m, 2H), 7.36 (t, $J = 7.2$ Hz, 1H), 5.28 (d, $J = 52.0$ Hz, 1H), 4.97 (d, $J = 13.6$ Hz, 1H), 4.79 - 4.82 (m, 1H), 4.03 - 4.21 (m, 3H), 3.73 (d, $J = 12.4$ Hz, 1H), 3.59 (s, 2H), 3.50 (d, $J = 17.6$ Hz, 1H), 3.40 (d, $J = 12.8$ Hz, 1H), 3.11 - 3.26 (m, 4H), 2.95 - 3.05 (m, 1H), 2.63 (d, $J = 17.6$ Hz, 1H), 1.78 - 2.13 (m, 10H).


[00430] **Compound 4.** 1-[7'-bromo-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol



[00431] Compound **4** was prepared similar to that of Ex. 1 as a dihydrochloride salt. LCMS calcd for $C_{30}H_{38}BrFN_4O_3(M+H)^+$ $m/z = 601.2$; found : 601.2. ^1H NMR (400 MHz, CD_3OD) δ 7.65-7.57 (m, 1H), 7.41-7.32 (m, 1H), 7.13-7.04 (m, 1H), 5.69-5.20 (m, 1H), 5.09-


5.00 (m, 1H), 4.85-4.72 (m, 3H), 4.04-3.78 (m, 8H), 3.51-3.39 (m, 1H), 3.19-3.98 (m, 2H), 2.86-2.58 (m, 4H), 2.50-2.41 (m, 1H), 2.41-1.67 (m, 13H).

[00432] Compound 5. 4-(azepan-1-yl)-5'-bromo-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin].




[00433] Compound **5** was prepared similar to that of Ex. 1 as a TFA salt. LCMS (ESI): m/z calcd for $C_{30}H_{38}BrFN_4O_2^+H$: 587.2, found: 587.2. ^1H NMR (400 MHz, CD_3OD) δ 7.55 (d, $J = 7.6$ Hz, 1H), 7.50 (d, $J = 6.9$ Hz, 1H), 7.14 (t, $J = 7.2$ Hz, 1H), 5.59 (d, $J = 51.6$ Hz, 1H), 4.99 (d, $J = 12.0$ Hz, 1H), 4.74 (d, $J = 14.8$ Hz, 3H), 3.89 (dd, $J = 38.2, 24.9$ Hz, 6H), 3.54 – 3.37 (m, 1H), 3.17 (dd, $J = 18.1, 6.4$ Hz, 1H), 3.06 (d, $J = 18.2$ Hz, 1H), 2.92 (d, $J = 17.4$ Hz, 1H), 2.81 – 2.73 (m, 1H), 2.64 (d, $J = 19.6$ Hz, 2H), 2.45 (s, 1H), 2.35 (s, 2H), 2.19 (d, $J = 16.4$ Hz, 3H), 2.06 – 1.81 (m, 6H), 1.65 (d, $J = 28.8$ Hz, 4H).

[00434] Compound **6.** 1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-ol.




[00435] Compound **6** was prepared similar to that of Ex. 1 as a dihydrochloride salt. LCMS (ESI): m/z calcd for $C_{29}H_{37}ClFN_4O_3^+H$: 543.4, found: 543.4. ^1H NMR (400 MHz, CD_3OD) δ 7.37 (d, $J = 8.8$ Hz, 1H), 7.32 – 7.22 (m, 2H), 5.59 (d, $J = 51.4$ Hz, 1H), 4.94 – 4.89 (m, 1H), 4.82 – 4.72 (m, 2H), 4.64 (dd, $J = 14.9, 5.4$ Hz, 1H), 4.09 – 3.73 (m, 8H), 3.52 – 3.41 (m, 1H), 3.20 – 3.07 (m, 3H), 3.05 – 2.94 (m, 1H), 2.77 – 2.58 (m, 3H), 2.49 – 2.29 (m, 5H), 2.27 – 2.07 (m, 3H), 1.97 – 1.74 (m, 4H).

[00436] Compound 7 & compound 8. 4-(azepan-1-yl)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-tetralin]-6'-ol, isomer 1 & 2.




[00437] Compound 7 and compound 8 were prepared similar to that of Ex. 1 separated by reverse phase HPLC. LCMS (ESI): m/z calcd for $C_{30}H_{39}FN_4O_3^+H$: 523.1, found: 523.1. ^1H NMR (400 MHz, CD_3OD , isomer 1, compound 7) δ 6.96 (d, $J = 8.3$ Hz, 1H), 6.79 (d, $J = 2.3$ Hz, 1H), 6.67 (dd, $J = 8.3, 2.4$ Hz, 1H), 5.29 (d, $J = 54.4$ Hz, 1H), 4.81 (d, 1H), 4.58 (d, 1H), 4.17 (d, 1H), 4.08 (d, 1H), 3.65 (t, 4H), 3.30 – 3.13 (m, 3H), 3.01 (dd, 2H), 2.94 – 2.82 (m, 1H), 2.74 (qd, 2H), 2.26 (ddd, 2H), 2.13 – 1.93 (m, 6H), 1.91 – 1.71 (m, 6H), 1.59 (d, $J = 3.3$ Hz, 4H). ^1H NMR (400 MHz, CD_3OD , isomer 2, compound 8) δ 6.94 (d, $J = 8.3$ Hz, 1H), 6.77 (d, $J = 2.3$ Hz, 1H), 6.65 (dd, $J = 8.3, 2.4$ Hz, 1H), 5.27 (d, $J = 54.4$ Hz, 1H), 4.79 (d, $J = 14.2$ Hz, 1H), 4.56 (d, $J = 14.1$ Hz, 1H), 4.15 (d, $J = 10.4$ Hz, 1H), 4.06 (d, $J = 10.4$ Hz, 1H), 3.72 – 3.56 (m, 4H), 3.29 – 3.13 (m, 3H), 3.04 – 2.94 (m, 2H), 2.85 (d, $J = 17.8$ Hz, 1H), 2.80 – 2.63 (m, 2H), 2.32 – 1.77 (m, 14H), 1.65 – 1.51 (m, 4H).

[00438] **Compound 9.** 4-(azepan-1-yl)-7'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,3'-indane]-4'-ol.




[00439] Compound 9 was prepared similar to that of Ex. 1 as a TFA salt. LCMS m/z calcd for $C_{29}H_{36}ClFN_4O_3^+H$: 543.1, found: 543.2. ^1H NMR (400 MHz, CD_3OD) δ 7.13 (d, $J = 8.6$ Hz, 1H), 6.66 (d, $J = 8.6$ Hz, 1H), 5.57 (d, $J = 51.3$ Hz, 1H), 4.75 (d, $J = 14.1$ Hz, 1H), 4.70 – 4.50 (m, 3H), 4.00 – 3.74 (m, 5H), 3.75 – 3.63 (m, 2H), 3.52 – 3.44 (m, 1H), 3.38 – 3.34 (m, 1H), 3.09 – 2.96 (m, 2H), 2.90–2.86 (m, 1H), 2.71 – 2.63 (m, 1H), 2.62 – 2.56 (m, 1H), 2.45 – 2.31 (m, 4H), 2.27 – 2.13 (m, 2H), 2.00 – 1.75 (m, 4H), 1.69 – 1.48 (m, 4H). ^{19}F NMR (377 MHz, CD_3OD) δ -174.17.

[00440] **Compound 10.** 5'-bromo-4-(3,8-diazabicyclo[3.2.1]octan-3-yl)-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]




[00441] Compound **10** was prepared similar to that of Ex. 2 as a dihydrochloride salt. LCMS (ESI): m/z calcd for $C_{30}H_{37}BrFN_5O_2^+H$: 598.2, found: 598.2. ^1H NMR (400 MHz, CD_3OD) δ 7.57 (d, $J = 7.9$ Hz, 1H), 7.52 (d, $J = 7.8$ Hz, 1H), 7.17 (t, $J = 7.8$ Hz, 1H), 5.62 (d, $J = 52.1$ Hz, 1H), 5.01 (dd, 1H), 4.86 (s, 2H), 4.73 (s, 1H), 4.70 (s, 1H), 4.40 (t, 1H), 4.25 (s, 2H), 4.04 – 3.84 (m, 4H), 3.73 (d, 1H), 3.47 (td, 1H), 3.24 (dd, 1H), 3.15 (d, 1H), 2.94 (d, 1H), 2.87 – 2.75 (m, 2H), 2.74 – 2.59 (m, 2H), 2.53 (d, 1H), 2.39 (dd, 2H), 2.27 (s, 2H), 2.15 (s, 5H), 2.03 (t, 1H), 1.88 (s, 1H).

[00442] **Compound 11.** 1-[7'-fluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol; dihydrochloride



[00443] Compound **11** was prepared similar to that of Ex. 1 as a dihydrochloride salt. LCMS m/z calcd for $C_{30}H_{38}F_2N_4O_3^+H$ 541.2, found 541.2. ^1H NMR (400 MHz, CD_3OD) δ 7.22 – 7.12 (m, 2H), 6.98 (dd, $J = 9.7, 7.1$ Hz, 1H), 5.59 (d, $J = 51.7$ Hz, 1H), 5.04 (d, $J = 14.2$ Hz, 1H), 4.77 (d, $J = 9.4$ Hz, 3H), 4.07 – 3.72 (m, 8H), 3.47 (s, 1H), 3.09 (dt, $J = 34.3, 14.2$ Hz, 2H), 2.83 (s, 2H), 2.61 (s, 2H), 2.45 (s, 1H), 2.35 (s, 2H), 2.15 (d, 6H), 1.89 (dd, 6H).


[00444] **Compound 12.** 4-(azepan-1-yl)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]



[00445] Compound **12** was prepared similarly to that of Ex. 1 as a dihydrochloride salt. LCMS (ESI): m/z calcd for $C_{30}H_{39}FN_4O_2^+H$; 507.2, found; 507.2. ^1H NMR (400 MHz, CD_3OD) δ 7.32 (d, $J = 7.7$ Hz, 1H), 7.13 (d, $J = 7.4$ Hz, 1H), 7.07 (t, $J = 7.0$ Hz, 2H), 5.49 (d, $J = 51.3$ Hz, 1H), 4.90 – 4.84 (m, 1H), 4.69 – 4.62 (m, 2H), 4.62 – 4.53 (m, 1H), 3.92 – 3.72

(m, 6H), 3.40 – 3.31 (m, 1H), 3.12 – 3.00 (m, 1H), 2.95-2.89 (m, 1H), 2.82 – 2.69 (m, 2H), 2.63 – 2.55 (m, 1H), 2.54 – 2.48 (m, 1H), 2.41 – 2.31 (m, 1H), 2.30 – 2.19 (m, 2H), 2.14 – 2.06 (m, 2H), 2.00 – 1.91 (m, 2H), 1.84 – 1.45 (m, 10H).

Example 4. Exemplary synthesis of 1-[7'-amino-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol (Compound **13**)




[00446] Step 1. Synthesis of 1-[7'-(benzhydrylideneamino)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol (**13a**). The solution of Pd₂dba₃ (4.57mg, 0.01mmol), diphenylmethanimine (18.08mg, 0.1mmol), t-BuONa (9.58mg, 0.1mmol), BINAP (2.11mg, 0.01mmol) and 1-[7'-bromo-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol (**4**, 30.0mg, 0.05mmol) in Toluene (2mL) was stirred at 100 °C under N₂ for 6h. The mixture was concentrated and purified by prep-HPLC (eluted with CH₃CN in H₂O (0.1% NH₄HCO₃) from 5% to 95%) to obtain 1-[7'-(benzhydrylideneamino)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol (**13a**, 10mg, 0.014mmol, 29% yield) as a brown solid. LCMS calcd for C₄₃H₄₈FN₅O₃ (M+H)⁺ m/z = 702.8; found; 702.4.

[00447] Step 2. Synthesis of 1-[7'-amino-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol (**13**). The solution of NH₂NHOH.HCl (2.97mg, 0.04mmol), AcONa (3.5mg, 0.04mmol) and 1-[7'-(benzhydrylideneamino)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol (**13a**, 10.0mg, 0.01mmol) in Methanol (1mL) was stirred at 25 °C under N₂ for 16h. The reaction was concentrated to get the crude, which was then purified by prep-HPLC (eluted with CH₃CN in H₂O (0.1% TFA) from 5% to 95%), then exchanged with HCl before lyophilization to give 1-[7'-amino-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol; dihydrochloride (**13**, 2.43mg, 0.0037mmol, 26.3% yield) as yellow solid. LCMS calcd for


$C_{30}H_{40}FN_5O_3$ (M+H)⁺ m/z = 538.3; found; 538.3. ¹H NMR (400 MHz, CD₃OD) δ 7.60-7.50 (m, 1H), 7.40-7.21 (m, 2H), 5.69-5.50 (m, 1H), 5.08-4.98 (m, 1H), 4.85-4.65 (m, 4H), 4.05-3.70 (m, 8H), 3.51-3.39 (m, 1H), 3.17-3.06 (m, 2H), 2.95-2.85 (m, 2H), 2.77-2.58 (m, 2H), 2.50-2.00 (m, 9H), 1.96-1.73 (m, 4H).

[00448] Compound 14. 2-[[[(2S,8R)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-4-(4-hydroxyazepan-1-yl)spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-tetralin]-6'-ol




[00449] Compound 14 was prepared similarly to that of Ex. 1 as a formic acid salt. LCMS (ESI): m/z calcd for $C_{30}H_{39}FN_4O_4^+H$; 539.1; found 539.1. ¹H NMR (400 MHz, CD₃OD) δ 8.51 (s, 1H), 6.95 (d, *J* = 8.2 Hz, 1H), 6.79 – 6.72 (m, 1H), 6.66 (dd, *J* = 8.3, 2.4 Hz, 1H), 5.39 (d, *J* = 53.6 Hz, 1H), 4.82 – 4.76 (m, 1H), 4.64 – 4.54 (m, 1H), 4.30 (dd, *J* = 10.9, 2.7 Hz, 1H), 4.25 – 4.16 (m, 1H), 3.91 – 3.51 (m, 6H), 3.49 – 3.39 (m, 2H), 3.23 – 3.10 (m, 1H), 3.00 (d, *J* = 17.6 Hz, 1H), 2.86 (d, *J* = 17.8 Hz, 1H), 2.80 – 2.65 (m, 2H), 2.54 – 2.21 (m, 3H), 2.21 – 1.87 (m, 8H), 1.88 – 1.56 (m, 5H). ¹⁹F NMR (377 MHz, CD₃OD) δ -173.75.

[00450] Compound 15. 1-[5'-chloro-2'-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]azepan-4-ol



[00451] Compound 15 was prepared similarly to that of Ex. 1 as a formic acid salt. LCMS (ESI): m/z calcd for $C_{30}H_{38}ClFN_4O_3^+H$; 557.1; found; 557.1. ¹H NMR (400 MHz, DMSO) δ 7.37 (t, *J* = 6.7 Hz, 2H), 7.21 (t, *J* = 7.9 Hz, 1H), 5.25 (d, *J* = 54.5 Hz, 1H), 4.78 (t, 1H), 4.55 (d, 1H), 4.02 – 3.90 (m, 1H), 3.85 (dd, 1H), 3.66 (s, 3H), 3.13 – 2.66 (m, 10H), 2.11 – 1.41 (m, 16H).


Example 5. Exemplary synthesis of 1-(4-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-d]pyrimidin]-4'-yl)azepane-4-carbonitrile (Compound 16)



[00452] Step 1. Preparation of 1-(4-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-d]pyrimidin]-4'-yl)azepan-4-yl methanesulfonate (**16a**). The mixture of 1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-ol; dihydrochloride (15.0 mg, 0.02 mmol), methanesulfonyl chloride (0.02 mL, 0.29 mmol) and N,N-diethylethanamine (0.05 mL, 0.37 mmol) in DCM (1 mL) was stirred at 25 °C for 30 min. The mixture was concentrated to afford a crude product which was purified on silica gel column eluted with EtOAc in PE from 5% to 50% to afford 1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-yl] methanesulfonate (**16a**, 15 mg, 0.0241 mmol, 99.2% yield) as yellow sticky solid. LCMS calcd for $C_{30}H_{38}ClFN_4O_5S$ ($M+H$)⁺ m/z = 621.3, found; 621.3.


[00453] Step 2. Preparation of 1-(4-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-d]pyrimidin]-4'-yl)azepan-4-carbonitrile (**16**). The mixture of [1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy] spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-yl]methanesulfonate (65.0 mg, 0.1 mmol), 18-crown-6 (27.66 mg, 0.1 mmol) and NaCN (51.29 mg, 1.05 mmol) was stirred at 25 °C for 16 h. The mixture was diluted with EtOAc (2 mL), washed with water (1 mL) and brine (1 mL), dried over Na_2SO_4 , concentrated. The crude product was purified by prep-HPLC (eluted with CH_3CN in H_2O (0.1 % FA)) from 5.0% to 95% to afford 1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-carbonitrile; formic acid (**16**, 11.79 mg, 0.0193 mmol, 18.43 % yield) as white solid. LCMS calcd for $C_{30}H_{36}ClFN_5O_2$ ($M+H$)⁺ m/z = 552.2, found: 552.2. ^1H NMR (400 MHz, CD_3OD) δ 8.57 (s, 1H), 7.31 – 7.10 (m, 3H), 5.34 (d, J = 54.1 Hz, 1H), 4.73 – 4.55 (m, 2H), 4.26 – 4.17 (m, 2H), 3.83 – 3.62 (m, 3H), 3.58 – 3.34 (m, 4H), 3.14 – 2.89 (m, 6H), 2.40 – 1.91 (m, 14H).

[00454] **Compound 17.** 1-(7-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-3,4,5',8'-tetrahydro-2H-spiro[naphthalene-1,7'-pyrano[4,3-d]pyrimidin]-4'-yl)azepan-4-ol




[00455] Compound 17 was prepared similarly to that of Ex. 1 as a formic acid salt. LCMS calcd for $C_{30}H_{38}ClFN_4O_3$ ($M+H$)⁺ m/z = 557.2, found: 557.2. ^1H NMR (400 MHz, CD_3OD) δ 8.52 (s, 1H), 7.31 (d, J = 2.3 Hz, 1H), 7.21 – 7.16 (m, 1H), 7.12 (dd, J = 8.2, 3.4 Hz, 1H), 5.33 (d, J = 53.3 Hz, 1H), 4.65 – 4.60 (m, 1H), 4.24 – 4.14 (m, 2H), 3.74 – 3.55 (m, 6H), 3.15 – 3.03 (m, 1H), 2.94 – 2.87 (m, 2H), 2.94 – 2.89 (m, 2H), 2.88 – 2.79 (m, 2H), 2.29 – 2.15 (m, 1H), 2.12 – 1.92 (m, 8H), 1.91 – 1.63 (m, 6H).

[00456] **Compound 18.** 4'-chloro-2-[[*(2R,8S)*-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-4-(2,3,6,7-tetrahydroazepin-1-yl)spiro[5,8-dihydropyrano[4,3-*d*]pyrimidine-7,1'-indane]



[00457] Compound 18 was prepared similarly to that of Ex. 1 as a TFA salt. LCMS (ESI): m/z calcd for $C_{29}H_{34}ClFN_4O_2^+H$: 525.06, found: 525.2. ^1H NMR (400 MHz, $CDCl_3$) δ 7.27-7.22 (m, 1H), 7.12-7.02 (m, 2H), 7.03 -6.99 (s, 1H), 5.67 (s, 2H), 5.49 (d, J = 53.2 Hz, 1H), 5.04-4.91 (m, 1H), 4.66 – 4.39 (m, 2H), 4.31-4.19 (m, 1H), 4.17-4.03 (m, 2H), 3.81 – 3.63 (m, 4H), 3.47 – 3.36 (m, 2H), 3.31-3.23 (m, 2H), 3.11-3.01 (m, 2H), 2.94-2.78 (m, 2H), 2.38-2.13 (m, 12H).

Example 6. Exemplary synthesis of 1-[4'-chloro-2-[[*rac*-(2*R*,8*S*)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl] methoxy]spiro[5,8-dihydropyrano[4,3-*d*]pyrimidine-7,1'-indane]-4-yl]azepan-4-one (Compound 21) and 2-(1-(4-chloro-2'-(((2*R*,7*aS*)-2-fluorotetrahydro-1*H*-pyrrolizin-7*a*(5*H*)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-*d*]pyrimidin]-4'-yl)azepan-4-yl)acetonitrile (Compound 19)




[00458] Step 1. Synthesis of 1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl] methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-one (**21**). The mixture of 1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl] methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-ol (**6**, 20.0 mg, 0.040 mmol) and DMP (23.43 mg, 0.060 mmol) was stirred at 25 °C for 30 min. The mixture was concentrated and purified by Prep-HPLC eluted with CH₃CN in H₂O (0.1 % FA) from 5.0% to 95% to afford 1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro [5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-one;formic acid (**21**, 1.8 mg, 0.003 mmol, 8.0% yield) as sticky solid. LCMS calcd for C₂₉H₃₅ClFN₄O₃ (M+H)⁺ m/z = 541.1, found: 541.1. ¹H NMR (400 MHz, CD₃OD) δ 8.43 (s, 1H), 7.33 (d, *J* = 7.6 Hz, 1H), 7.20 (t, *J* = 7.6 Hz, 1H), 7.07 (dd, *J* = 7.6, 1.9 Hz, 1H), 5.49 (d, *J* = 51.8 Hz, 1H), 4.73 (d, *J* = 14.6 Hz, 1H), 4.48 (d, *J* = 14.6 Hz, 1H), 4.42 (d, *J* = 11.6 Hz, 1H), 4.35 (d, *J* = 11.6 Hz, 1H), 3.95 (t, *J* = 6.1 Hz, 2H), 3.89 – 3.65 (m, 5H), 3.17 – 3.05 (m, 2H), 3.03 – 2.88 (m, 4H), 2.81 – 2.73 (m, 2H), 2.62 – 2.34 (m, 5H), 2.33 – 2.16 (m, 5H).

[00459] Step 2. Synthesis of 2-(1-(4-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-d]pyrimidin]-4'-yl)azepan-4-ylidene)acetonitrile (**19a**). To a mixture of 1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl] methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-one (**21**, 20.0 mg, 0.04 mmol) and 2-diethoxyphosphorylacetonitrile (6.55 mg, 0.04 mmol) in dried THF (1 mL) was added sodium hydride (1.77 mg, 0.07 mmol) at 25 °C under argon. The mixture was stirred at rt for 1 h. The mixture was diluted with EtOAc (10 mL), washed with water (5 mL) and brine (1 ml), dried over Na₂SO₄, concentrated. The crude product was purified by Prep-TLC to afford 2-[1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-ylidene]acetonitrile (**19a**, 10 mg, 0.018 mmol, 48.0% yield) as yellow sticky solid. LCMS calcd for C₃₁H₃₅ClFN₅O₂ (M+H)⁺ m/z = 564.1, found: 564.1.

[00460] Step 3. Preparation of 2-(1-(4-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-d]pyrimidin]-4'-yl)azepan-4-yl)acetonitrile (**19**). The suspension of 2-[1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl] methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-ylidene]acetonitrile (10.0 mg, 0.02 mmol) and Pd-C (0.19 mg, 0.0018 mmol) in methanol (1 mL) was purged with H₂ three times. The mixture was stirred under H₂ (1 atm) at 25 °C for 16 h. The mixture was filtered and concentrated to afford a crude product which was purified by Prep-HPLC eluted with (eluted with CH₃CN in H₂O (0.1 % FA) from 5.0 % to 95 %) to obtain 2-[1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-


yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-4-yl]acetonitrile;formic acid (**19**, 2.24 mg, 0.0034 mmol, 18.9 % yield) as white sticky solid. LCMS calcd for $C_{31}H_{38}ClFN_5O_2$ (M+H)⁺ m/z = 566.2, found: 566.2. ¹H NMR (400 MHz, CD₃OD) δ 8.47 (s, 1H), 7.41 – 7.08 (m, 3H), 5.45 (d, J = 51.1 Hz, 1H), 4.78 – 4.23 (m, 4H), 3.96 – 3.33 (m, 7H), 3.26 – 3.21 (m, 1H), 3.16 – 2.88 (m, 4H), 2.59 – 2.33 (m, 5H), 2.28 – 1.60 (m, 12H).

Example 7. Exemplary synthesis of 4-(azepan-1-yl)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-tetralin]-6'-carbonitrile (Compound **20**)




[00461] The solution of 4-(azepan-1-yl)-7'-bromo-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin] (**20a**, prepared similarly to that of Ex. 1, 30.mg, 0.05mmol), K₂CO₃ (21.24mg, 0.15mmol), Zn(CN)₂ (18.05mg, 0.15mmol) and Xphos G3 Pd (4.69mg, 0.01mmol) in 1,4-Dioxane (1mL) and Water (0.2mL) was stirred at 100 °C under N₂ for 6h. The reaction mixture was concentrated to get the crude product, which was then purified by prep-HPLC (eluted with CH₃CN in H₂O (0.1% FA) from 5% to 95%) to give 4-(azepan-1-yl)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-tetralin]-6'-carbonitrile;formic acid (**20**, 3.2mg, 0.0042mmol, 8.2% yield) as yellow solid. LCMS calcd for $C_{31}H_{38}FN_5O_2$ (M+H)⁺ m/z = 532.3; found: 532.3. ¹H NMR (400 MHz, CD₃OD) δ 8.51 (s, 1H), 7.72-7.64 (m, 1H), 7.58-7.49 (m, 1H), 7.38-7.26 (m, 1H), 5.47-5.26 (m, 1H), 4.93-4.76 (m, 1H), 4.68-4.52 (m, 1H), 4.33-4.14 (m, 2H), 3.75-3.60 (m, 4H), 3.53-3.33 (m, 3H), 3.19-3.07 (m, 1H), 3.03-2.75 (m, 4H), 2.48-1.90 (m, 9H), 1.87-1.72 (m, 5H), 1.69-1.51 (m, 4H).

[00462] **Example 8.** Exemplary synthesis of syn-1-[4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepane-4,5-diol (Compound **22**)



[00463] To a mixture of 4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl] methoxy]-4-(2,3,6,7-tetrahydroazepin-1-yl)spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane] (**18**, 50.0 mg, 0.1 mmol) in acetone (0.3 mL) and water (0.1 mL) were added $K_2OsO_4 \cdot 2H_2O$ (70.17 mg, 0.19 mmol) and NMO (22.31 mg, 0.19 mmol) at 25 °C. Then the mixture was stirred at 40 °C for 16 h. The solution was quenched with Na_2SO_3 aqueous solution, extracted with EtOAc (3 ml), and concentrated. The residue was purified by prep-HPLC eluting with CH_3CN in H_2O (0.1 % TFA) from 5.0% to 95% to give syn-1-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepane-4,5-diol;2,2,2-trifluoroacetic acid (**22**, 4.71 mg, 0.0065 mmol, 6.8% yield) as a sticky solid. LCMS calcd for $C_{31}H_{37}ClF_4N_4O_6$ ($M+H$)⁺ m/z = 559.24, found: 559.5. ^1H NMR (400 MHz, CD_3OD) δ 7.40-7.38 (dd, J =7.2, 0.8 Hz, 1H), 7.29-7.25 (t, J =7.6 Hz, 1H), 7.29-7.23 (dd, J =7.6 Hz, 1.2 Hz, 1H), 5.67 (s, 2H), 5.60 (d, J =52 Hz, 1H), 4.93 – 4.65 (m, 4H), 4.00 – 3.70 (m, 9H), 3.52 – 3.46 (m, 1H), 3.18 – 3.06 (m, 2H), 3.02 – 2.96 (m, 1H), 2.88 (m, 1 H), 2.75-2.65 (m, 1 H), 2.62 – 2.62 (m, 1 H), 2.50 – 2.12 (m, 8 H), 2.03 – 1.90 (m, 2 H).

Example 9. Exemplary synthesis of (3*R*)-1-(4-chloro-2'-(((2*R*,7*aS*)-2-fluorotetrahydro-1*H*-pyrrolizin-7*a*(5*H*)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-*d*]pyrimidin]-4'-yl)-3-methylpiperidin-3-ol (Compound **23**)



[00464] Step 1. Synthesis of ethyl 4-chloro-6-((4-chloro-1-hydroxy-2,3-dihydro-1*H*-inden-1-yl)methyl)-2-(methylthio)pyrimidine-5-carboxylate (**23a**). To a solution of LiHMDS (1 M, 24.3 mL, 2.0 eq) in THF (10 mL) was added a solution of Ethyl 4-chloro-6-methyl-2-(methylthio)pyrimidine-5-carboxylate (3.0 g, 12.2 mmol, 1.0 eq) in THF (10 mL) at -60 °C. Then the reaction was stirred at -60 °C for 30 minutes. After 30 minutes, 4-chloro-2,3-dihydro-1*H*-inden-1-one (2.03 g, 12.2 mmol, 1.0 eq) in THF (10 mL) was added to the mixture at -60 °C. Then the reaction was stirred at -60 °C for 1 hr. LCMS showed that the reaction was completed.

The residue was poured into water (50 mL). The aqueous phase was extracted with EtOAc (20 mL * 2). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated under vacuum. The residue was purified by column chromatography on silica gel (PE: EtOAc = 80:1 to 30:1) to give ethyl 4-chloro-6-((4-chloro-1-hydroxy-2,3-dihydro-1*H*-inden-1-yl)methyl)-2-(methylthio)pyrimidine-5-carboxylate (**23a**) (3.5 g, 8.47 mmol, 69.64% yield) as a light-yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 7.24-7.25 (m, 1H), 7.17 (t, *J*=7.6 Hz, 1H), 7.06-7.08 (m, 1H), 5.29(s, 1H), 4.31-4.38 (m, 2H), 3.23-3.26 (m, 1H), 3.07-3.11 (m, 1H), 3.06-3.07 (m, 1H), 2.81-2.87 (m, 1H), 2.59 (s, 3H), 2.31-2.37 (m, 3H), 2.12-2.20 (m, 1H), 1.33 (t, *J*=7.2 Hz, 3H).

[00465] Step 2. Synthesis of 4-chloro-1-((6-chloro-5-(hydroxymethyl)-2-(methylthio)pyrimidin-4-yl)methyl)-2,3-dihydro-1*H*-inden-1-ol (**23b**). To a solution of compound **23a** (3.4 g, 8.23 mmol, 1.0 eq) in DCM (35 mL) was added DIBAL-H (1 M, 24.68 mL, 3.0 eq) at 0 °C. The mixture was stirred at 0 °C for 1 hr. LCMS indicated the reaction was completed. The reaction mixture was quenched by water (50 mL) at 0 °C, and then extracted with DCM (20 mL * 2). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (PE: EtOAc = 100:1 to 30:1) to give compound **23b** (2.0 g, 5.39 mmol, 65.48% yield) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 7.26-7.28 (m, 1H), 7.16 (t, *J*=8.0 Hz, 1H), 7.03-7.05 (m, 1H), 4.62-4.73 (m, 2H), 3.34-3.38 (m, 1H), 3.13-3.16 (m, 1H), 3.03-3.06 (m, 1H), 2.89-2.95 (m, 1H), 2.53 (s, 3H), 2.50-2.52 (m, 1H), 2.07-2.12 (m, 1H).

[00466] Step 3. Synthesis of 4,4'-dichloro-2'-(methylthio)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-*d*]pyrimidine] (**23c**). To a solution of compound **23b** (1.7 g, 4.58 mmol, 1.0 eq) in THF (20 mL) was added DIAD (1.39 g, 6.87 mmol, 1.34 mL, 1.5 eq) and PPh₃ (1.8 g, 6.87 mmol, 1.5 eq) at 25 °C. Then the reaction was stirred at 25 °C for 10 minutes. LC-MS showed the starting material was consumed completely. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by column chromatography on silica gel (PE/EtOAc=100/1 to 80/1 to give compound **23c** (1.07 g, 3.03 mmol, 66.15% yield) as off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.30-7.32 (m, 1H), 7.16 (t, *J*=8.0 Hz, 1H), 6.94-6.96 (m, 1H), 4.72-4.76 (m, 1H), 4.56-4.60 (m, 1H), 3.13-3.19 (m, 1H), 3.10 (s, 2H), 2.95-2.96 (m, 1H), 2.58 (s, 3H), 2.36-2.40 (m, 1H), 2.11-2.16 (m, 1H).


[00467] Step 4. Synthesis of (3*R*)-1-(4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-*d*]pyrimidine-7,1'-indane]-4-yl)-3-methyl-piperidin-3-ol (**23d**). A solution of 4,4'-dichloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-*d*]pyrimidine-7,1'-indane] (80.mg, 0.23mmol) in DMF (0.3mL) was added (3*R*)-3-methylpiperidin-3-ol;hydrochloride (37.77mg, 0.25mmol) and DIPEA (87.8mg, 0.68mmol) at room temperature. The mixture was stirred at 100 °C for 1 h. The

solution was diluted with EtOAc (5ml), then washed with water (2 ml x3), dried over Na₂SO₄ and concentrated. The residue was purified with prep-TLC (eluting with EtOAc ether in petroleum from 10 % to 90%) to give (3R)-1-(4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-3-methyl-piperidin-3-ol (**23d**, 80mg, 0.185mmol, 81.8% yield). LCMS calcd for C₂₂H₂₆ClN₃O₂S (M+H)⁺ m/z = 432.98, found: 432.1.

[00468] Step 5. Synthesis of **23e**. A mixture of (3R)-1-(4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-3-methyl-piperidin-3-ol (**23d**, 80. mg, 0.19mmol) in THF (0.4mL) and Water (0.4mL) was added oxone (68.31mg, 0.22mmol) at 25 °C for 1 h. The reaction was quenched with Na₂SO₃ aqueous solution and diluted with EtOAc (3 ml). The organic layer was washed with water (3 ml x3), dried and concentrated to give 90 mg of **23e** as a 5:1 ratio of mixture of sulfoxide and sulfone based on LCMS. LCMS calcd for C₂₂H₂₆ClN₃O₄S (M+H)⁺ m/z = 448.98, 464.98, found: 448.1, 464.1.


[00469] Step 6. Synthesis of (3R)-1-(4-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-2,3,5',8'-tetrahydrospiro[indene-1,7'-pyrano[4,3-d]pyrimidin]-4'-yl)-3-methylpiperidin-3-ol (**23**). A mixture of [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (177.69mg, 1.12mmol) in DMSO (0.3mL) was added NaH (44.64mg, 1.12mmol) at 25 °C for 0.5 h. Then a mixture sulfone and sulfoxide of **23e** in DMSO (0.3mL) was added to the reaction at 25 °C and stirred for 1 h. Acetic acid was added dropwise until pH = 7. The crude mixture was purified by prep-HPLC (eluting with CH₃CN (0.1% TFA)in H₂O from 5 % to 95%) to give (3R)-1-[4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-methyl-piperidin-3-ol; 2,2,2-trifluoroacetic acid (46.03mg, 0.0695mmol, 31.13% yield) as a white solid. LCMS calcd for C₂₉H₃₆ClFN₄O₃ (M+H)⁺ m/z = 543.07, found: 543.1. ¹H NMR (400 MHz, CD₃OD) δ 7.38 (d, J=7.6 Hz, 1H), 7.28-7.20 (m, 2 H), 5.60 (d, J= 52 Hz, 1H), 4.99-4.96 (m, 1H), 4.78-4.50 (m, 4H), 4.35-4.278 (m, 1H), 4.01– 3.75 (m, 4H), 3.52-3.46 (m, 1H), 3.40-3.31 (m, 1 H), 3.25 – 3.07 (m, 3 H), 3.02– 2.96 (m, 1 H), 2.77-2.58 (m, 2 H), 2.49-2.26 (m, 6 H), 2.03-1.94 (m, 1 H), 1.82-1.71 (m, 3 H), 1.24 (s, 3H).

[00470] **Compound 24.** 9-[4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione.




[00471] Compound **24** was prepared similarly to that of Ex. 9. LCMS calcd for $C_{30}H_{34}ClFN_6O_4$ ($M+H$)⁺ m/z = 597.09, found: 597.5. ¹H NMR (400 MHz, CD₃OD) δ 7.34 – 7.27 (m, 1H), 7.20 (t, J = 7.7 Hz, 1H), 7.14 – 7.04 (m, 1H), 5.28 (d, J = 54.2 Hz, 1H), 4.75 – 4.58 (m, 1H), 4.53 – 4.40 (m, 1H), 4.23 – 4.03 (m, 2H), 4.01 – 3.66 (m, 2H), 3.43 – 3.35 (m, 1H), 3.26 – 3.05 (m, 5H), 3.04 – 2.89 (m, 4H), 2.43 – 2.05 (m, 6H), 2.00 – 1.78 (m, 6H).

[00472] **Compound 25.** 4-(azepan-1-yl)-2-[[*(2R,8S)*-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-*d*]pyrimidine-7,4'-isochromane]



[00473] Compound **25** was prepared similarly to that of Ex. 1 as a formic acid salt. LCMS (ESI): m/z calcd for $C_{29}H_{37}FN_4O_3$ ($M+H$)⁺: 509.2, found: 509.2. ¹H NMR (400 MHz, CD₃OD) δ 7.41 – 7.36 (m, 1H), 7.30 – 7.18 (m, 2H), 7.09 (d, J = 7.5 Hz, 1H), 5.39 – 5.22 (m, 1H), 4.80 (m, 3H), 4.70 (d, J = 14.2 Hz, 1H), 4.64 – 4.52 (m, 1H), 4.23 – 4.07 (m, 2H), 3.96 – 3.86 (m, 2H), 3.68 – 3.61 (m, 4H), 3.30 – 3.17 (m, 2H), 3.08 – 2.99 (m, 1H), 2.96 – 2.89 (m, 2H), 2.37 – 1.77 (m, 10H), 1.65 – 1.54 (m, 4H).

[00474] **Example 10.** Exemplary synthesis of 2'-amino-4-(azepan-1-yl)-2-[[*rac*-(*2R,8S*)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-*d*]pyrimidine-7,4'-6,7-dihydro-5H-benzothiophene]-3'-carbonitrile (Compound **26**)



[00475] Step 1. Synthesis of methyl 4-(azepan-1-yl)-2-chloro-6-[(2-ethoxy-1-hydroxy-cyclohex-2-en-1-yl)methyl]pyrimidine-5-carboxylate (**26a**). A mixture of methyl 4-(azepan-1-yl)-2-chloro-6-methyl-pyrimidine-5-carboxylate (**1a**, 1.0 g, 3.52mmol) and LDA (2.11mL, 4.23mmol) in THF (10mL) was stirred at -78 °C for 1 h. Then, a solution of 2-ethoxycyclohex-2-en-1-one (543mg, 3.88mmol) and CeCl₃ (956mg, 3.88mmol), stirred for 4h in THF (5mL), was added at -78 °C. The mixture was stirred at -78 °C for 1 h. LCMS showed 42% of product and 35% of **1a** remained. The mixture was quenched with aq. NH₄Cl (5 mL) at -78 °C, extracted with EtOAc (5 mL x 3), dried, and concentrated. The crude product was purified by flash chromatography (H₂O:CH₃CN = 90:10 to 50:50) to give methyl 4-(azepan-1-yl)-2-chloro-6-[(2-ethoxy-1-hydroxy-cyclohex-2-en-1-yl)methyl]pyrimidine-5-carboxylate (**26a**, 800mg, 1.89mmol, 53.5% yield) was obtained as a yellow sticky oil. LCMS calcd for C₂₁H₃₀ClN₃O₄ (M+H)⁺ m/z = 424.2, found: 424.2.

[00476] Step 2. Synthesis of 1-[[6-(azepan-1-yl)-2-chloro-5-(hydroxymethyl)pyrimidine-4-yl]methyl]-2-ethoxy-cyclohex-2-en-1-ol (**26b**). A suspension of LAH (177mg, 4.67mmol) and lithium chloride (198mg, 4.67mmol) in 2-Methyltetrahydrofuran (6mL) was stirred at room temperature for 20 min. Then methyl 4-(azepan-1-yl)-2-chloro-6-[(2-ethoxy-1-hydroxy-cyclohex-2-en-1-yl)methyl]pyrimidine-5-carboxylate (**26a**, 660.mg, 1.56mmol) was added at 0 °C portion wise. The mixture was stirred at 0 °C for 10 min. The mixture was quenched with NH₄Cl (0.5 mL) at 25 °C, extracted with EtOAc (10 x 3 mL), dried over Na₂SO₄, concentrated. The crude product was purified by silica gel chromatography (eluted with EtOAc in PE from 10% to 30%) to give 1-[[6-(azepan-1-yl)-2-chloro-5-(hydroxymethyl)pyrimidin-4-yl]methyl]-2-ethoxy-cyclohex-2-en-1-ol (**26b**, 230mg, 0.581mmol, 37.3% yield) as a yellow solid. LCMS calcd for C₂₀H₃₀ClN₃O₃ (M+H)⁺ m/z = 396.1, found: 396.1.

[00477] Step 3. Synthesis of 4-(azepan-1-yl)-2-chloro-1'-ethoxy-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,6'-cyclohexene] (**26c**). To a solution of 1-[[6-(azepan-1-yl)-2-chloro-5-(hydroxymethyl)pyrimidin-4-yl]methyl]-2-ethoxy-cyclohex-2-en-1-ol (**26b**, 230mg, 0.58mmol) in THF (2mL) was added n-Butyllithium (0.35mL, 0.87mmol) at -78 °C stirred for 0.5h. then the TSCl (166mg, 0.87mmol) in THF (2mL) was added at -78 °C and the mixture was warmed to -10 °C stirred for 1 h under Ar. Then the temperature was cooled down to -78 °C and the second batch of n-Butyllithium (0.35mL, 0.87mmol) was added, after addition the reaction temperature was allowed to rise to 20 °C and stirred for 1.5 h. The mixture was quenched with NH₄Cl (1 mL) at -78 °C, extracted with EtOAc (10 x 3 mL), dried over Na₂SO₄, concentrated. The crude product was purified by silica gel chromatography (eluted with EtOAc in PE from 1% to 20%) to give 4-(azepan-1-yl)-2-chloro-1'-ethoxy-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,6'-

cyclohexene] (**26c**, 130mg, 0.344mmol, 59.2% yield) as a yellow solid. LCMS calcd for $C_{20}H_{28}ClN_3O_2$ (M+H)⁺ m/z = 378.1, found: 378.1.


[00478] Step 4. Synthesis of 4-(azepan-1-yl)-1'-ethoxy-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,6'-cyclohexene] (**26d**). The mixture of NaH (31.8mg, 0.79mmol) and 2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]methanol (126mg, 0.79mmol) in DMSO (1mL) was stirred at 0 °C for 0.5 h. Then a solution of 4-(azepan-1-yl)-2-chloro-1'-ethoxy-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,6'-cyclohexene] (**26c**, 100.mg, 0.26mmol) in DMSO (0.5mL) was added at 0 °C and the mixture was stirred at 50 °C for 4 h. The mixture was quenched with H₂O (10 mL) at 0 °C, extracted with EtOAc (5 x 3 mL), dried over Na₂SO₄, concentrated. The crude product was purified by silica gel chromatography (eluted with EtOAc in PE from 10% to 30%) to give 4-(azepan-1-yl)-1'-ethoxy-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,6'-cyclohexene] as a yellow solid (**26d**, 80mg, 0.1598mmol, 60.387% yield) was obtained as yellow solid. LCMS calcd for $C_{28}H_{41}FN_4O_3$ (M+H)⁺ m/z = 501.2, found: 501.6.

[00479] Step 5. Synthesis of 4-(azepan-1-yl)-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,2'-cyclohexane]-1'-one (**26e**). The mixture of 4-(azepan-1-yl)-1'-ethoxy-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,6'-cyclohexene] (80.mg, 0.16mmol) and HCl/Dioxane (1.mL, 4.mmol) in Water (0.2mL) was stirred at 25 °C. the mixture was stirred at 25 °C for 10 min. After concentration, the crude product was purified by Prep-TLC (DCM:MeOH = 5:1). LCMS (ESI): m/z calcd for $C_{26}H_{37}FN_4O_3$ (M+H)⁺ m/z = 472.2, found: 473.2.

[00480] Step 6. Synthesis of 2'-amino-4-(azepan-1-yl)-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-6,7-dihydro-5H-benzothiophene]-3'-carbonitrile (Compound **26**). The mixture of 4-(azepan-1-yl)-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,2'-cyclohexane]-1'-one (**26e**, 68.mg, 0.14mmol) and propanedinitrile (9.5mg, 0.14mmol) and octathiocane (36.9mg, 0.14mmol) and morpholine (25mg, 0.29mmol) in Ethanol (0.5mL) was stirred at 65 °C, the mixture was stirred at 65 °C under argon for 7 h. The mixture was filtered and concentrated. The crude product was purified by Prep-HPLC (eluted with CH₃CN in H₂O (0.1 % TFA) from 5.0% to 95%) to give 2'-amino-4-(azepan-1-yl)-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,4'-6,7-dihydro-5H-benzothiophene]-3'-carbonitrile; 2,2,2-trifluoroacetic acid (**26**,


2.9 mg, 0.003 mmol, 2.1 %yield) as a yellow sticky solid. LCMS (ESI): m/z calcd for $C_{29}H_{39}FN_6O_2S$ (M+H)⁺ m/z =553.2, found: 553.2.

[00481] Compound 23A and 23B. (3R)-1-[(7*)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-methyl-piperidin-3-ol (**23A** and **23B**)




[00482] Compound **23** was purified on an DAICELCHIRALCEL®AD (250*25 mm 10 mm) column on a Waters SFC 150 system (Mobile Phase A: Supercritical CO₂, Mobile Phase B: IPA (+0.1% 7.0mol/l Ammonia in MeOH); A:B: 60:40; Flow: 70ml/min) to give faster eluting P1 (3R)-1-[(7*)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-methyl-piperidin-3-ol (**23A**) and slower eluting P2 (**23B**). **Compound 23A:** ¹H NMR (400 MHz, CD₃OD) δ 7.351-7.332 (m, 1H), 7.235-7.916 (m, 1 H), 7.121-7.103 (m, 1H), 5.399-5.256 (m, 1 H), 4.749 (d, J=14 Hz, 1 H), 4.459 (d, J=14 Hz, 1 H), 4.228-4.119 (m, 2 H), 3.665-3.627(m, 1H), 3.422-3.349 (m, 2H), 3.284-3.231 (m, 4 H), 3.165 – 2.927 (m, 5 H), 2.443-1.867 (m, 10 H), 1.768-1.626 (m, 2H), 1.189 (s, 3 H). **Compound 23B:** ¹H NMR (400 MHz, CD₃OD) δ 7.227-7.207 (m, 1H), 7.116-7.077 (m, 1 H), 7.041-7.022 (m, 1H), 5.291-5.155 (m, 1 H), 4.557-4.406 (m, 2 H), 4.150-4.017 (m, 2 H), 3.607-3.544 (m, 1 H), 3.380-3.347 (m, 1H), 3.297-3.255 (m, 1 H), 3.164-2.929 (m, 6 H), 2.884 – 2.802 (m, 3 H), 2.305-2.026 (m, 5H), 1.950-1.73(m, 5 H), 1.646-1.482 (m, 2 H), 1.090 (s, 3 H) .

[00483] Compound 24A. (5*)-9-[(7*)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione




[00484] Compound **24** was purified on an DAICELCHIRALCEL®AD (250*25 mm 10 um) column on a Waters SFC 150 system (Mobile Phase A: Supercritical CO₂, Mobile Phase B: IPA (+0.1% 7.0mol/l Ammonia in MeOH); A:B: 60:40; Flow: 70ml/min) to give 3 peaks and the fastest eluting P1 was assigned to compound **24A**.

Example 11. Exemplary synthesis of 5-[(7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (compound **27A**) and 5-[(7R)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (Compound **27B**)



[00485] Compound **27A** was obtained from **27A-a** similarly to that of compound **27B** as a formic acid salt. The absolute configuration of **27A** was assigned based on a small molecule crystal structure. LCMS calcd for $C_{33}H_{39}ClFN_7O_4$ ($M+H$)⁺ m/z = 636.3, found: 636.3. ¹H NMR (400 MHz, CD₃OD) δ 8.51 (s, 1H), 7.38-7.20 (m, 2H), 7.08-7.00 (m, 1H), 6.53 (s, 1H), 5.46-5.26 (m, 1H), 4.78-4.68 (m, 3H), 4.59-4.46 (m, 3H), 4.26-4.03 (m, 3H), 3.87-3.76 (m, 1H), 3.52-3.31 (m, 7H), 3.18-2.88 (m, 8H), 2.46-1.90 (m, 10H).

[00486] Compound **27B** 5-[(7R)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00487] Compound **23c** was purified on an DAICELCHIRALCEL®AD (25*250 mm, 10 μ m) column on a Waters SFC 150 system (Mobile Phase A: Supercritical CO₂, Mobile Phase B: IPA (+0.2% 7.0mol/l Ammonia in MeOH); A:B: 60:40; Flow: 100ml/min) to give 2 peaks and the fastest eluting P1 was assigned to compound **27A-a** and the second one P2 was assigned as **27B-a**. **27B-a** absolute configuration was assigned based on **27A-a** single crystal structure.

[00488] Step 1. **27B-a** (140.mg, 0.4mmol), N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide;hydrochloride (165.07mg, 0.79mmol) and DIEA (0.21mL, 1.19mmol) in DMF (2mL) was stirred at 100 °C for 2 h. The mixture was concentrated. The


crude product was purified by flash chromatography (eluted with CH₃CN in H₂O from 5.0% to 95%) to get 5-[(7S)-4'-chloro-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide **27b** (173 mg, 0.3295 mmol, 83.14% yield) as yellow solid. LCMS calcd for C₂₆H₂₉ClN₆O₂S (M+H)⁺ m/z = 525.1, found: 525.1. ¹H NMR (400 MHz, CDCl₃) δ 7.37-7.27 (m, 2H), 7.02 (d, J = 7.1 Hz, 1H), 6.49 (s, 1H), 4.69 – 4.39 (m, 7H), 3.78 – 3.65 (m, 1H), 3.35 (s, 3H), 3.23 – 3.11 (m, 3H), 3.09 (s, 3H), 2.99 – 2.92 (m, 1H), 2.52 (s, 3H), 2.41 – 2.13 (m, 4H).

[00489] Step 2. To a solution **27b** (170.mg, 0.32mmol) in THF (2mL) was added a solution of oxone (129.38mg, 0.42mmol) in Water (1mL) at 0 °C, then the mixture was stirred at 25 °C for 1 h. The mixture was added water (10 mL) and extracted with EtOAc (10 mL x 3), washed with a solution of Na₂SO₃ (5 mL), brine (5 ml), dried over Na₂SO₄, concentrated. The crude product was used directly for the next step. The crude product of N,N-dimethyl-5-[(7S)-4'-chloro-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide **27c** (135mg, 0.2495mmol, 77.07% yield) and N,N-dimethyl-5-[(7S)-4'-chloro-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide **27d** (35 mg, 0.0628mmol, 19.41% yield) was obtained as yellow solid. LCMS calcd for **27c** C₂₆H₂₉ClN₆O₃S (M+H)⁺ m/z = 541.1, found: 541.1, 21% of product 3-1. LCMS calcd for **27d** C₂₆H₂₉ClN₆O₄S (M+H)⁺ m/z = 557.1, found: 557.1.

[00490] Step 3. To a solution of [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (88.27mg, 0.55mmol) in DMF (0.5mL) was added NaH (22.18mg, 0.55mmol) at 25 °C. The mixture was stirred at 25 °C for 15 min. A solution of N, N-dimethyl-5-[(7S)-4'-chloro-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide **27c** (100.mg, 0.18mmol) and N,N-dimethyl-5-[(7S)-4'-chloro-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide **27d** (20.mg, 0.04mmol) in DMF (0.5mL) was added the above solution at 25 °C. The mixture was stirred at 25 °C for 15 min. Acetic acid (0.03mL, 0.55mmol) was added the reaction mixture to quench the reaction. The crude product was purified by Prep-HPLC (eluted with CH₃CN in H₂O (0.1% FA) from 5.0% to 95%) to get 5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide;formic acid **27B** (67.2 mg, 0.1052 mmol, 56.92% yield) was obtained as white solid. LCMS calcd for C₃₃H₃₉ClFN₇O₃ (M+H)⁺ m/z = 636.2, found: 636.2. ¹H NMR (400 MHz, CD₃OD) δ 8.52 (s, 1H), 7.33 (d, J = 7.9 Hz, 1H), 7.24 (t, J = 7.7 Hz, 1H), 7.04 (d, J = 7.4 Hz, 1H), 6.53 (s, 1H), 5.42 –

5.24 (m, 1H), 4.77 – 4.68 (m, 3H), 4.57 – 4.48 (m, 3H), 4.23 – 4.10 (m, 2H), 4.09 – 3.76 (m, 2H), 3.45 – 3.34 (m, 2H), 3.31 – 3.29 (m, 4H), 3.16 – 2.89 (m, 8H), 2.43 – 1.90 (m, 10H).

Example 12. Exemplary synthesis of (7S)-4-(azepan-1-yl)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane] (**28A**)



[00491] Step 1. Synthesis of methyl 4-(azepan-1-yl)-6-(((S)-1-(((R)-*tert*-butylsulfinyl)amino)-4-chloro-2,3-dihydro-1*H*-inden-1-yl)methyl)-2-chloropyrimidine-5-carboxylate (**28A-b**). To a solution of LDA (1 M, 4.49 mL) in THF (3 mL) was added compound **1a** (0.85 g, 3.00 mmol) at -65 °C. After the mixture was stirred at -65 °C for 30 minutes, (*R*)-*N*-(4-chloro-2,3-dihydro-1*H*-inden-1-ylidene)-2-methylpropane-2-sulfinamide (**28A-a**, Kobayashi, Jun-ichi; et al. *Bioorganic & Medicinal Chemistry* (**2021**), 30, 115903) (808 mg, 3.00 mmol) was added to the mixture at -65 °C. The mixture was stirred at -65 °C for 1 hr. TLC indicated ~50% of **1a** was remained and one major new spot with larger polarity was detected. The reaction mixture was poured into water (50 mL), the aqueous phase was extracted with EtOAc (20 mL x 2). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (PE: EtOAc = 80:1 to 10:1) to give methyl 4-(azepan-1-yl)-6-(((S)-1-(((R)-*tert*-butylsulfinyl)amino)-4-chloro-2,3-dihydro-1*H*-inden-1-yl)methyl)-2-chloropyrimidine-5-carboxylate as a single diastereomer (**28A-b**, 0.65 g, 1.17 mmol, 39.20% yield) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.23-7.25 (m, 1H), 7.12-7.16 (m, 1H), 6.96-6.98 (m, 1H), 6.88 (s, 1H), 3.72 (s, 3H), 3.51-3.56 (m, 2H), 3.42-3.45 (m, 2H), 3.00-3.03 (m, 1H), 2.85-2.95 (m, 4H), 2.66-2.72 (m, 1H), 2.28-2.33 (m, 1H), 1.80(s, 4H), 1.49-1.56 (m, 4H), 1.29 (s, 9H). Chiral analysis of compound **28A-b** (Chiralpak AD-3, 50×4.6mm I.D., 3μm): 99.66%. The absolute configuration

of **28A-b** was assigned based on the single crystal structure of (*R*)-**28A-d** (HCl salt, not shown, prepared from (*S*)-**28A-a**).

[00492] Step 2. Synthesis of (*R*)-*N*-((*S*)-1-((6-(azepan-1-yl)-2-chloro-5-(hydroxymethyl)pyrimidin-4-yl)methyl)-4-chloro-2,3-dihydro-1*H*-inden-1-yl)-2-methylpropane-2-sulfonamide (**28A-c**). To a solution of compound **28A-b** (0.55 g, 994 μ mol) in DCM (10 mL) was added DIBAL-H (1 M, 2.98 mL) at 0 °C. Then the reaction was stirred at 0 °C for 1 hr. The reaction mixture was quenched by ice water (20 mL) at 25 °C. The aqueous phase was extracted with DCM (10 mL x 2). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by prep-TLC (SiO₂, PE/EtOAc = 3:1) to give compound **28A-c** (0.4 g, 761.13 μ mol, 76.60% yield) as off-white solid. ¹H NMR: (400 MHz, CDCl₃) δ 7.24 (s, 1H), 7.12-7.16 (m, 1H), 7.01-7.03 (m, 1H), 4.23-4.34 (m, 2H), 3.70-3.74 (m, 4H), 3.08-3.19 (m, 3H), 2.93-2.95 (m, 1H), 2.72-2.76 (m, 1H), 2.31-2.37 (m, 1H), 1.75-1.82 (m, 4H), 1.55-1.61 (m, 4H), 1.26 (s, 9H).


[00493] Step 3. Synthesis of (*S*)-4'-(azepan-1-yl)-2',4-dichloro-2,3,5',8'-tetrahydro-6'*H*-spiro[indene-1,7'-pyrido[4,3-*d*]pyrimidine] (**28A-d**). To a solution of compound **28A-c** (0.2 g, 4.58 mmol) in HCl/dioxane (4 M, 2 mL) at 0 °C. The mixture was stirred at 0 °C for 3 hrs. The reaction mixture was adjusted pH to 8 by NaHCO₃ and extracted with EtOAc (5 mL x 2). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give a residue. To a mixture of the residue in CHCl₃ (1.5 mL) was added SOCl₂ (154 mg, 1.29 mmol, 93.86 μ L) at 0 °C, and the mixture was stirred at 50 °C for 2 hrs. The reaction mixture was concentrated under reduced pressure to give a residue. To a mixture of the residue in THF (1.5 mL) was added NaOH (5M, 12 mL) at 25 °C and the mixture was stirred at 25 °C for 1 hr. LC-MS showed the starting material was consumed completely. The mixture was extracted with EtOAc (5 mL x 2). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (PE/EtOAc=1:1) to give compound **28A-d** (0.1 g, 248 μ mol, 65% yield) as a light-yellow solid. ¹H NMR: (400 MHz, CDCl₃) δ 7.25-7.27 (m, 1H), 7.13-7.17 (m, 1H), 7.04-7.06 (m, 1H), 3.99-4.03 (m, 1H), 3.85-3.89 (m, 1H), 3.62-3.65 (m, 4H), 3.07-3.13 (m, 1H), 2.97-3.01 (m, 3H), 2.16-2.24 (m, 2H), 1.80 (s, 4H), 1.68 (m, 8H).

[00494] Step 4. Synthesis of (*S*)-4'-(azepan-1-yl)-2',4-dichloro-6'-methyl-2,3,5',8'-tetrahydro-6'*H*-spiro[indene-1,7'-pyrido[4,3-*d*]pyrimidine] (**28A-e**). To a solution of compound **28A-d** (95 mg, 236 μ mol) in DMF (1 mL) was added NaH (9.42 mg, 235.53 μ mol, 60% purity) and iodomethane (33.4 mg, 236 μ mol, 14.7 μ L) at 25 °C. Then the reaction was stirred at 25 °C for 2 hrs. LC-MS showed the starting material was consumed completely. The reaction mixture

was quenched by water (5 mL), and then extracted with EtOAc (3 mL x 2). The combined organic layers were washed with brine (5 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by prep-TLC (PE/EtOAc=1:1) to give compound **28A-e** (40 mg, 96 μ mol, 41% yield) as light-yellow solid.

[00495] Step 5. Synthesis of (7S)-4-(azepan-1-yl)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane] (**28A**). To a solution of compound **28A-e** (17 mg, 40.73 μ mol) and ((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methanol (12.97 mg, 81.46 μ mol) in toluene (0.2 mL) was added Cs₂CO₃ (39.81 mg, 122.19 μ mol) and RuPhos Pd G3 (3.41 mg, 4.07 μ mol) at 20 °C. Then the reaction was stirred at 100 °C for 5 hrs. LC-MS showed the starting material was consumed completely. The residue was purified by prep-HPLC (column: Waters Xbridge Prep OBD C18 150*40mm*10 μ m; mobile phase: [water(NH₄HCO₃)-ACN]; B%: 60%-80%, 8min) to give **compound 28A** (1.4 mg, 2.59 μ mol, 6.4% yield) as an off-white solid. ¹H NMR (400 MHz, CD₃OD) δ 7.29-7.3 (m, 1H), 7.26-7.28 (m, 1H), 7.17-7.24 (m, 1H), 5.21-5.34 (m, 1H), 4.14-4.17 (m, 1H), 4.03-4.06 (m, 1H), 3.72-3.81 (m, 6H), 3.15-3.21 (m, 3H), 2.98 (m, 3H), 2.80-2.88 (m, 2H), 2.37-2.45 (m, 1H), 2.25-2.30 (m, 1H), 2.17 (m, 3H), 2.15-2.25 (m, 1H), 2.10-2.05 (m, 1H), 1.88 (m, 2H), 1.64-1.65 (m, 6H), 1.28 (m, 4H).


[00496] **Example 13.** Exemplary synthesis of [1-[4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-piperidyl]cyanamide (**29**)



[00497] To the mixture of 1-[4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]piperidin-3-amine (**29a**, prepared similarly to that of **Ex. 11**) (60 mg, 0.11 mmol) and Et₃N (0.08 mL, 0.57 mmol) in DCM (0.4 mL) at 0 °C was added carbononitridic bromide (13 mg, 0.12 mmol) in DCM (0.2 mL) and stirred at 0 °C for 0.5 h. The mixture was taken into water (10 mL), which was extracted with DCM (3 x 5 mL). The organic layers were washed with brine (10 mL), dried over Na₂SO₄, and concentrated. The crude was purified by prep-HPLC (eluted with CH₃CN in H₂O (0.1 % FA) from 5.0% to 95%) to give [1-[4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-


piperidyl]cyanamide; formic acid (**29**, 15.24 mg, 0.0264 mmol, 23.23% yield) as a white solid. LCMS calcd for $C_{29}H_{34}ClFN_6O_2$ ($M+H$)⁺ m/z = 553.1, found: 553.1. ¹H NMR (400 MHz, CD₃OD) δ 8.52 (s, 1H), 7.32 (dd, J = 7.5, 3.1 Hz, 1H), 7.20 (dd, J = 14.7, 7.5 Hz, 1H), 7.13 – 7.09 (m, 1H), 5.44 – 5.23 (m, 1H), 4.78 – 4.69 (m, 1H), 4.61 – 4.53 (m, 2H), 4.52 – 4.39 (m, 1H), 4.28 – 4.15 (m, 2H), 4.02 – 3.95 (m, 1H), 3.57 – 3.46 (m, 1H), 3.42 – 3.34 (m, 2H), 3.25 – 2.93 (m, 7H), 2.45 – 1.83 (m, 10H), 1.69 – 1.54 (m, 2H).

[00498] Compound 30. 9-[6'-hydroxy-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione.




[00499] Compound **30** was prepared similarly to that of **Ex. 9** as a TFA salt. LCMS calcd for $C_{30}H_{35}FN_6O_5$ ($M+H$)⁺ m/z = 579.3, found: 579.1. ¹H NMR (400 MHz, CD₃OD) δ 7.15 – 7.08 (m, 1H), 6.80 – 6.72 (m, 1H), 6.62 – 6.56 (m, 1H), 5.66 – 5.48 (m, 1H), 4.70 – 4.51 (m, 4H), 4.26 – 3.98 (m, 2H), 3.94 – 3.84 (m, 2H), 3.78 – 3.66 (m, 1H), 3.53 – 3.37 (m, 3H), 3.04 – 2.97 (m, 2H), 2.82 – 2.54 (m, 3H), 2.46 – 2.28 (m, 4H), 2.25 – 2.08 (m, 4H), 2.00 – 1.79 (m, 3H).

[00500] Compound 31. N-methoxy-5-[(7S)-4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00501] Compound **31** was prepared from **27B-a** similarly to that of **Ex. 11** as a formic acid salt. LCMS calcd for $C_{32}H_{37}ClFN_7O_4$ ($M+H$)⁺ m/z = 638.2 found: 638.2. ¹H NMR (400 MHz, CD₃OD) δ 8.52 (s, 1H), 7.37 – 7.26 (m, 2H), 7.02 (d, J = 7.3 Hz, 1H), 6.61 (s, 1H), 5.39 – 5.20 (m, 1H), 4.76 – 4.67 (m, 3H), 4.56 – 4.46 (m, 3H), 4.10 (d, J = 17.1 Hz, 2H), 3.78 (s, 3H), 3.26 – 3.19 (m, 2H), 3.16 – 2.80 (m, 6H), 2.49 – 1.78 (m, 12H)

[00502] Compound 32. N-methoxy-5-[4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00503] Compound **32** was prepared similarly to that of **Ex. 9** as a TFA salt. LCMS calcd for $C_{33}H_{40}FN_7O_4$ ($M+H$)⁺ m/z = 618.3, found: 618.2. ¹H NMR (400 MHz, CD₃OD) δ 7.14 (d, J = 8.3 Hz, 1H), 6.77 (dd, J = 8.1, 2.4 Hz, 1H), 6.64 (d, J = 1.6 Hz, 1H), 6.54 (t, J = 2.6 Hz, 1H), 5.69 – 5.51 (m, 1H), 4.96 – 4.90 (m, 2H), 4.86 – 4.74 (m, 2H), 4.68 – 4.57 (m, 1H), 4.55 – 4.46 (m, 3H), 4.07 – 3.83 (m, 5H), 3.55 – 3.42 (m, 1H), 3.38 – 3.35 (m, 3H), 3.15 – 2.90 (m, 6H), 2.82 – 2.57 (m, 3H), 2.52 – 2.31 (m, 4H), 2.28 – 2.04 (m, 4H).

[00504] **Compound 33.** (3aR,6aS)-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3a,4,6,6a-tetrahydropyrrolo[3,4-c]pyrrole-1,3-dione



[00505] Compound **33** was prepared from **27B-a** similarly to that of **Ex. 11** as a TFA salt. LCMS calcd for $C_{29}H_{31}ClFN_5O_4$ ($M+H$)⁺ m/z = 568.2, found: 568.1. ¹H NMR (400 MHz, CD₃OD) δ 7.36 (dd, J = 7.9, 0.7 Hz, 1H), 7.23 (t, J = 7.7 Hz, 1H), 7.10 (d, J = 7.5 Hz, 1H), 5.69 – 5.50 (m, 1H), 4.86 – 4.81 (m, 1H), 4.69 – 4.52 (m, 3H), 4.34 (d, J = 12.9 Hz, 1H), 4.22 (d, J = 12.9 Hz, 1H), 4.03 – 3.87 (m, 3H), 3.83 – 3.70 (m, 2H), 3.58 – 3.45 (m, 3H), 3.19 – 3.09 (m, 1H), 3.05 – 2.91 (m, 3H), 2.75 – 2.57 (m, 2H), 2.48 – 2.31 (m, 4H), 2.27 – 2.17 (m, 2H).


[00506] **Compound 34.** [5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-pyrrolidin-1-yl-methanone



[00507] Compound **34** was prepared from **27B-a** similarly to that of **Ex. 11** as a formic acid salt. LCMS calcd for $C_{30}H_{34}ClFN_6O_4$ ($M+H$)⁺ m/z = 597.09, found: 597.5. ¹H NMR (400 MHz,


CD₃OD) δ 7.34 - 7.27 (m, 1H), (t, J = 7.7 Hz, 1H), 7.14 - 7.04 (m, 1H), 5.28 (d, J = 54.2 Hz, 1H), 4.75 - 4.58 (m, 2H), 4.53 - 4.40 (m, 2H), 4.23 - 4.03 (m, 2H), 4.01 - 3.66 (m, 2H), 3.43 - 3.35 (m, 1H), 3.26 - 3.05 (m, 5H), 3.04 - 2.89 (m, 4H), 2.43 - 2.05 (m, 6H), 2.00 - 1.78 (m, 6H).

[00508] Compound 35. 5-[(7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N-isopropyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00509] Compound **35** was prepared from **27B-a** similarly to that of **Ex. 11** as a formic acid salt. LCMS calcd for C₃₀H₃₄ClFN₆O₄ (M+H)⁺ m/z = 597.09, found: 597.5. ¹H NMR (400 MHz, CD₃OD) δ 7.34 - 7.27 (m, 1H), 7.20 (t, J = 7.7 Hz, 1H), 7.14 - 7.04 (m, 1H), 5.28 (d, J = 54.2 Hz, 1H), 4.75 - 4.58 (m, 1H), 4.53 - 4.40 (m, 1H), 4.23 - 4.03 (m, 3H), 4.01 - 3.66 (m, 3H), 3.43 - 3.35 (m, 1H), 3.26 - 3.05 (m, 5H), 3.04 - 2.89 (m, 4H), 2.43 - 2.05 (m, 6H), 2.00 - 1.78 (m, 6H).

[00510] Compound 36. 3-chloro-5-[(7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00511] Compound **36** was prepared from **27B-a** similarly to that of **Ex. 11** as a formic acid salt. LCMS calcd for C₃₃H₃₈Cl₂FN₇O₃ (M+H)⁺ m/z = 670.5, found: 670.5. ¹H NMR (400 MHz, CD₃OD) δ 8.52 (s, 1H), 7.32 (d, J = 7.1 Hz, 1H), 7.20 (t, J = 7.8 Hz, 1H), 7.11 (d, J = 7.1 Hz, 1H), 5.40 - 5.20 (m, 1H), 4.82 - 4.76 (m, 3H), 4.59 (d, J = 14.3 Hz, 1H), 4.46 - 4.39 (m, 2H), 4.14 - 4.05 (m, 2H), 3.90 - 3.80 (m, 2H), 3.26 - 3.20 (m, 2H), 3.15 - 3.04 (m, 8H), 3.01 - 2.96 (m, 2H), 2.47 - 1.77 (m, 12H).

[00512] Compound 37. 5-[(7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N-hydroxy-N-methyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00513] Compound **37** was prepared from **27B-a** similarly to that of **Ex. 11** as a formic acid salt. LCMS calcd for $C_{32}H_{38}ClFN_7O_4$ ($M+H$)⁺ m/z = 639.1, found: 638.3. ¹H NMR (400 MHz, CD₃OD) δ 7.36 (d, J = 7.9 Hz, 1H), 7.26 (s, 1H), 7.08 (d, J = 7.5 Hz, 1H), 6.82 (s, 1H), 5.30 (d, J = 52.8 Hz, 1H), 4.81 – 4.68 (m, 3H), 4.55 (d, J = 13.8 Hz, 3H), 4.13 (d, J = 10.2 Hz, 1H), 4.03 (s, 2H), 3.87 (s, 1H), 3.44 (s, 2H), 3.29 (s, 2H), 3.22 (s, 1H), 3.17 – 2.91 (m, 5H), 2.40 (dd, J = 10.5, 7.1 Hz, 1H), 2.31 – 2.07 (m, 6H), 1.98 (dd, J = 17.8, 6.1 Hz, 2H), 1.88 (s, 1H).

[00514] **Compound 38.** morpholino-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]methanone;2,2,2-trifluoroacetic acid



[00515] Compound **38** was prepared from **27B-a** similarly to that of **Ex. 11** as a TFA salt. LCMS calcd for $C_{35}H_{41}ClFN_7O_4$ ($M+H$)⁺ m/z = 678.2, found: 678.2. ¹H NMR (400 MHz, CD₃OD) δ 7.37 (d, J = 7.6 Hz, 1H), 7.29-7.25 (m, 1H), 7.11 (d, J = 6.4 Hz, 1H), 6.66 (s, 1H), 5.58 (d, J = 51.6 Hz, 1H), 5.01-4.99 (m, 3H), 4.63-4.52 (m, 5H), 4.12-4.01 (m, 3H), 3.94-3.83 (m, 4H), 3.77-3.68 (m, 6H), 3.47-3.45 (m, 1H), 3.13-3.94 (m, 4H), 2.70 – 2.14 (m, 10H)

[00516] **Example 14.** 5-[7'-hydroxy-6-methyl-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 39**)



[00517] Step 1. To a solution of 7-hydroxytetralin-1-one (30000.mg, 184.97mmol) in DMF (200mL) was added K_2CO_3 (51128.92mg, 369.94mmol) at rt, then BnBr (34798.63mg, 203.47mmol) was added. The mixture was stirred at rt for 16h. Then the mixture was extracted with EtOAc/H₂O. The combined organic layer was washed with brine, dried over Na_2SO_4 and concentrated. The obtained solid is triturated in MeOH and then filtered. The filtrate was purified by flash column chromatography (silica gel, eluting with EtOAc in PE) to afford 7-benzyloxytetralin-1-one (39a) (41760 mg, 166 mmol, 89.48 %yield) as a yellow solid. LCMS calculated for $C_{17}H_{17}O_2$ ($M+H$)⁺ m/z =253.1; found:253.2. ¹H NMR (400 MHz, DMSO) δ 7.18 – 7.49 (m, 8H), 5.14 (s, 2H), 2.87 (t, J = 5.6 Hz, 2H), 2.57 (t, J = 6.4 Hz, 2H), 1.96 – 2.07 (m, 2H).

[00518] Step 2. 7-benzyloxytetralin-1-one (39a) (4320.mg, 17.12mmol), (S)-2-methylpropane-2-sulfinamide (2905.23mg, 23.97mmol) and Ti(IV) ethoxide (10.mL, 51.37mmol) were mixed and stirred under argon at rt. The reaction vessel was placed into the microwave reactor and heated to 100 °C for 120 min. The mixture was quenched with potassium sodium tartrate tetrahydrate (aq, 250 mL), filtered, extract with EtOAc and purified by flash column chromatography (silica gel, eluting with 10% to 30% EtOAc/PE) to afford (NE,S)-N-(7-benzyloxytetralin-1-ylidene)-2-methyl-propane-2-sulfinamide (39b) (3300 mg, 9.28 mmol, 54.22 %yield) as a brown oil. LCMS calculated for $C_{21}H_{26}NO_2S$ ($M+H$)⁺ m/z =356.17; found: 356.3.

[00519] Step 3. To a solution of EtOAc (2478.55mg, 28.13mmol) in THF (16mL) was added LDA (5.60mL, 11.25mmol) portion wise at -70 °C under Ar, the mixture was stirred at -70 °C for 1h, then another 1 equiv LDA was added and the reaction stirred for an additional 1 hr at -70 °C, and (NE,S)-N-(7-benzyloxytetralin-1-ylidene)-2-methyl-propane-2-sulfinamide (39b) (2000.mg, 5.63mmol) in THF (8mL) was added drop wise at -70 °C. The resulting mixture was stirred at -70 °C for 2h, The reaction was heated naturally and overnight. The reaction mixture

was quenched with $\text{NH}_4\text{Cl}(\text{aq.})$ at $-70\text{ }^\circ\text{C}$, and then extracted with EtOAc. The organic layer was washed with water and brine, dried over Na_2SO_4 , concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 6% MeOH/DCM) to afford ethyl 2-[7-benzyloxy-1-[[(S) -tert-butylsulfinyl]amino]tetralin-1-yl]acetate (**39c**) (1200 mg, 2.71 mmol, 48.08 %yield). LCMS calculated for $\text{C}_{25}\text{H}_{33}\text{NNaO}_4\text{S}^+$ ($\text{M}+\text{Na}$) $^+$ m/z = 466.20; found: 466.4.

[00520] Step 4. The mixture of ethyl 2-[7-benzyloxy-1-[[(S) -tert-butylsulfinyl]amino]tetralin-1-yl]acetate (**39c**) (1200.mg, 2.71mmol) and HCl in dioxane (5.mL, 20.mmol) in Ethanol (10mL) was stirred at rt for 1 h. The reaction mixture was concentrated to afford crude ethyl 2-(1-amino-7-benzyloxy-tetralin-1-yl)acetate (**39d**) (918mg, 2.7045mmol, 100% yield). LCMS calculated for $\text{C}_{21}\text{H}_{26}\text{NO}_3$ ($\text{M}+\text{H}$) $^+$ m/z = 340.19; found: 323.4.

[00521] Step 5. A mixture of ethyl 2-(1-amino-7-benzyloxy-tetralin-1-yl)acetate (**39d**) (918.mg, 2.7mmol), ethyl prop-2-enoate (812.33mg, 8.11mmol), TEA (1.13mL, 8.11mmol) and CuO (43.03mg, 0.54mmol) was dissolved in Ethanol (9mL). The vial was sealed, and this mixture was then stirred for 16h at $85\text{ }^\circ\text{C}$. The crude reaction mixture was filtered and purified by Prep-HPLC (FA) to afford ethyl 3-[[7-benzyloxy-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]amino]propanoate (**39e**) (620mg, 1.4106mmol, 52.2% yield) as a colorless oil. LCMS calculated for $\text{C}_{26}\text{H}_{34}\text{NO}_5$ ($\text{M}+\text{H}$) $^+$ m/z = 440.24; found: 440.5.

[00522] Step 6. To a solution of ethyl 3-[[7-benzyloxy-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]amino]propanoate (**39e**) (620.mg, 1.41mmol) in Ethanol (6mL) was added paraformaldehyde (253.9mg, 8.46mmol). Then the mixture was stirred at rt for 1h under N_2 , and NaBH_3CN (265.92mg, 4.23mmol) was added. The resulting mixture was stirred at rt overnight. The reaction mixture was quenched with water, and then extracted with EtOAc. The organic layer was washed with brine, dried over Na_2SO_4 , concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 25% EtOAc/PE) to afford ethyl 3-[[7-benzyloxy-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]-methyl-amino]propanoate (**39f**) (540mg, 1.1906mmol, 84.4% yield) as a colorless oil. LCMS calculated for $\text{C}_{27}\text{H}_{36}\text{NO}_5$ ($\text{M}+\text{H}$) $^+$ m/z = 454.26; found: 454.5.

[00523] Step 7. To a solution of ethyl 3-[[7-benzyloxy-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]-methyl-amino]propanoate (**39f**) (440.mg, 0.97mmol) in THF (6mL) was added LiHMDS (2.91mL, 2.91mmol) portion wise at $-70\text{ }^\circ\text{C}$ under N_2 . Then the mixture was stirred at $-70\text{ }^\circ\text{C}$ for 1h. The reaction mixture was quenched with aqueous NH_4Cl at $-70\text{ }^\circ\text{C}$. The mixture was extracted with EtOAc, washed with brine, dried over Na_2SO_4 , concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 15% EtOAc/PE) to afford ethyl 7'-benzyloxy-1-methyl-4-oxo-spiro[piperidine-6,1'-tetralin]-3-carboxylate (**39g**)

(154mg, 0.3779mmol, 39.0% yield) as a colorless oil. LCMS calculated for $C_{25}H_{30}NO_4$ (M+H)⁺ m/z = 408.22; found: 408.4.

[00524] Step 8. To a solution of ethyl 7'-benzyloxy-1-methyl-4-oxo-spiro[piperidine-6,1'-tetralin]-3-carboxylate (**39g**) (279.mg, 0.68mmol) in ethanol (3mL) was added thiourea (67.75mg, 0.89mmol) and ethoxysodium (139.77mg, 2.05mmol). Then the mixture was stirred at 80 °C for 16h. The reaction mixture was concentrated, diluted with EtOAc. 1M HCl was added to adjust pH=6. The mixture was washed with water and brine, dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 2% to 8% MeOH/DCM) to afford 7'-benzyloxy-6-methyl-2-sulfanyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-ol (**39h**) (250mg, 0.5959mmol, 87.0% yield) as a yellow gel. LCMS calculated for $C_{24}H_{26}N_3O_2S$ (M+H)⁺ m/z = 420.18; found: 420.4.

[00525] Step 9. To a solution of 7'-benzyloxy-6-methyl-2-sulfanyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-ol (**39h**) (250.mg, 0.6mmol) in water (5mL) was added 2-chloroacetic acid (506.81mg, 5.36mmol). Then the mixture was stirred at 100 °C overnight under N₂. The mixture was filtered. The pH of the aqueous layer was adjusted to 8 by adding an aqueous solution of NaHCO₃ aqueous. The resulting precipitate was filtered, and dried in vacuo to afford 7'-benzyloxy-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**39i**) (224mg, 0.5552mmol, 93.2% yield) as a white solid. LCMS calculated for $C_{24}H_{26}N_3O_3$ (M+H)⁺ m/z = 404.20; found: 404.4.

[00526] Step 10. The mixture of 7'-benzyloxy-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**39i**) (224.mg, 0.56mmol) in POCl₃ (3.mL, 32.19mmol) was stirred at 100°C for 16 h. The reaction mixture was concentrated, extracted with EtOAc, washed with brine (10 mL), dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 30% EtOAc/PE) to afford 7'-benzyloxy-2,4-dichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**39j**) (28mg, 0.064mmol, 11% yield) as a white solid. LCMS calculated for $C_{24}H_{24}Cl_2N_3O$ (M+H)⁺ m/z = 440.13, 442.13; found: 440.4, 442.4.


[00527] Step 11. The mixture of 7'-benzyloxy-2,4-dichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**39j**) (28.mg, 0.06mmol), N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide;hydrochloride (18.67mg, 0.08mmol) and DIEA (16.44mg, 0.13mmol) in DMF (1mL) was stirred at rt for 24 h. The reaction mixture was diluted with water, extracted with EtOAc, washed with brine, dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 3% to 7% MeOH/DCM) to afford 5-(7'-benzyloxy-2-chloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl)-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-

2-carboxamide (**39k**) (25mg, 0.041mmol, 64% yield) as a white solid. LCMS calculated for $C_{34}H_{39}ClN_7O_2$ (M+H)⁺ m/z = 612.29; found: 612.4.

[00528] Step 12. The mixture of 5-(7'-benzyloxy-2-chloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl)-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**39k**) (25 mg, 0.04mmol), [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (8.45mg, 0.05mmol), CS_2CO_3 (33.28mg, 0.1mmol), Pd_2dba_3 (3.74mg, 0.01mmol) and RuPhos (3.81mg, 0.01mmol) in toluene (1.5mL) was stirred at 110°C for 10 h under N_2 . The reaction mixture was cooled to room temperature and the solvent was removed under reduced pressure. The mixture was purified by flash column chromatography (silica gel, eluting with 0% to 10% MeOH/DCM) to afford 5-[7'-benzyloxy-6-methyl-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**39l**) (25mg, 0.034mmol, 83% yield) as a white solid. LCMS calculated for $C_{42}H_{52}FN_8O_3$ (M+H)⁺ m/z = 735.42; found: 735.8.


[00529] Step 13. The mixture of 5-[7'-benzyloxy-6-methyl-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**39l**) (25 mg, 0.03mmol) and Pd/C (7.82mg, 0.01mmol) in MeOH (1mL) was stirred at rt for 2 h under H_2 . The reaction mixture was filtered, concentrated and purified by prep-HPLC (0.1% NH_4HCO_3) to afford 5-[7'-hydroxy-6-methyl-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 39**) (7.84mg, 0.0122mmol, 35.7% yield) as a white solid. LCMS calculated for $C_{35}H_{46}FN_8O_3$ (M+H)⁺ m/z = 645.37; found: 645.7. ^1H NMR (400 MHz, CD_3OD) δ 6.92 (dd, J = 7.6, 5.6 Hz, 2H), 6.58 – 6.64 (m, 2H), 5.26 (d, J = 54.4 Hz, 1H), 4.79 – 4.89 (m, 2H), 4.42 – 4.59 (m, 2H), 3.92 – 4.14 (m, 4H), 3.76 (dd, J = 34.0, 14.8 Hz, 2H), 3.32 (s, 3H), 3.14 – 3.27 (m, 3H), 3.06 (s, 3H), 2.87 – 3.03 (m, 3H), 2.65 (dd, J = 9.6, 4.0 Hz, 2H), 2.12 – 2.36 (m, 3H), 2.14 (s, 3H), 1.67 – 2.10 (m, 9H).

[00530] **Compound 39A and 39B.** 5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-7'-hydroxy-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**39A**) and 5-[(7R)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-7'-hydroxy-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**39B**)




[00531] Compound **39** was purified on an DAICELCHIRALCEL®AD (250*25 mm 10 μ m) column on a Waters SFC 150 system (Mobile Phase A: Supercritical CO₂, Mobile Phase B: MeOH (0.2% NH₃ (7M in MeOH))); A:B: 65:35; Flow: 100ml/min) to give faster eluting P1 (**39A**) and slower eluting P2 (**39B**).

[00532] **Compound 40.** 5-[8'-chloro-7'-hydroxy-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide.




[00533] Compound **40** was prepared similarly to that of **Ex. 9** as a white solid. LCMS calculated for C₃₄H₄₂ClFN₇O₄ (M+H)⁺ m/z = 666.3; found: 666.5/668.5. ¹H NMR (400 MHz, CD₃OD) δ 6.94 (d, J = 8.4 Hz, 1H), 6.83 (d, J = 8.4 Hz, 1H), 6.64 (d, J = 2.4 Hz, 1H), 5.32 (d, J = 54.4 Hz, 1H), 5.02 (d, J = 14.0 Hz, 1H), 4.90 (s, 2H), 4.69 (d, J = 14.4 Hz, 1H), 4.44 – 4.53 (m, 2H), 4.00 – 4.20 (m, 3H), 3.84 – 3.92 (m, 1H), 3.58 (d, J = 18.0 Hz, 1H), 3.34 (d, J = 18.0 Hz, 1H), 3.27 (s, 3H), 3.06 (s, 3H), 3.01 – 3.11 (m, 1H), 2.75 – 2.83 (m, 3H), 1.87 – 2.40 (m, 13H), 1.68 – 1.80 (m, 1H).

[00534] **Example 15.** Exemplary synthesis of 5-[6'-(difluoromethoxy)-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**41**)



[00535] To a solution of 5-[6'-hydroxy-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide;2,2,2-trifluoroacetic acid (**32**) (15. mg, 0.02 mmol) in MeCN (0.3 mL) were added 1-[[bromo(difluoro)methyl]-ethoxy-phosphoryl]oxyethane (8.21 mg, 0.03 mmol) and KF (2.38 mg, 0.04 mmol) at 25 °C. The mixture was stirred at 25 °C for 12 h. The crude product was purified by Prep-HPLC (eluted with CH₃CN in H₂O (0.1 % TFA) from 5.0% to 95%) to afford 5-[6'-(difluoromethoxy)-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide;2,2,2-trifluoroacetic acid (**41**) (5.4 mg, 0.0055 mmol, 27 % yield) as white solid. LCMS calcd for C₃₄H₄₀F₃N₇O₄ (M+H)⁺ m/z = 668.3, found: 668.2. ¹H NMR (400 MHz, CD₃OD) δ 7.59-7.26 (m, 1H), 7.12 (d, *J* = 8.1 Hz, 1H), 6.75 (dd, *J* = 8.1, 2.4 Hz, 1H), 6.61 (s, 1H), 6.54 (s, 1H), 5.78 – 5.57 (m, 1H), 4.81 – 4.69 (m, 4H), 4.67 – 4.23 (m, 6H), 4.05 – 3.89 (m, 3H), 3.33 (s, 3H), 3.12 – 2.88 (m, 8H), 2.81 – 2.68 (m, 2H), 2.64 – 2.47 (m, 3H), 2.45 – 2.29 (m, 2H), 2.21 – 2.06 (m, 3H).


[00536] **Compound 42.** 5-[(7S)-4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N,3-trimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00537] Compound **42** was prepared from 27B-a similarly to that of **Ex. 11** as a TFA salt. LCMS calcd for C₃₄H₄₂ClFN₇O₃ (M+H)⁺ m/z = 650.3, found: 650.3. ¹H NMR (400 MHz, CD₃OD) δ 7.36 (dd, *J* = 7.9, 0.8 Hz, 1H), 7.24 (t, *J* = 7.7 Hz, 1H), 7.14 (d, *J* = 7.4 Hz, 1H), 5.60 (d, *J* = 50.7 Hz, 1H), 4.87 (s, 2H), 4.72 (dd, *J* = 86.4, 14.4 Hz, 2H), 4.52 (s, 2H), 4.41 (dd, *J* =


6.6, 3.3 Hz, 2H), 4.00 – 3.83 (m, 5H), 3.47 (td, $J = 11.1, 5.4$ Hz, 1H), 3.21 – 2.91 (m, 10H), 2.85 – 2.53 (m, 2H), 2.49 – 2.09 (m, 11H).

[00538] Compound 43. 5-[5'-chloro-7'-hydroxy-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**43**)




[00539] Compound **43** was prepared similarly to that of **Ex. 9** as a white solid. LCMS calculated for $C_{34}H_{42}ClFN_7O_4$ ($M+H$)⁺ $m/z = 666.3$; found: 666.3/668.3. ¹H NMR (400 MHz, CD₃OD) δ 6.80 (d, $J = 2.4$ Hz, 1H), 6.70 (d, $J = 2.4$ Hz, 1H), 6.55 (d, $J = 2.0$ Hz, 1H), 5.37 (d, $J = 53.6$ Hz, 1H), 4.77 – 4.80 (m, 2H), 4.60 – 4.61 (m, 1H), 4.55 – 4.57 (m, 1H), 4.45 – 4.51 (m, 2H), 4.14 – 4.26 (m, 2H), 3.91 – 3.99 (m, 2H), 3.35 – 3.51 (m, 3H), 3.30 (s, 3H), 3.10 – 3.19 (m, 1H), 3.07 (s, 3H), 2.99 (d, $J = 18.0$ Hz, 1H), 2.91 (d, $J = 18.0$ Hz, 1H), 2.66 – 2.83 (m, 2H), 2.26 – 2.45 (m, 2H), 2.16 – 2.25 (m, 1H), 2.05 – 2.15 (m, 5H), 1.90 – 2.03 (m, 3H), 1.70 – 1.84 (m, 1H).

[00540] Compound 44. [(3R)-1-[(7S)-4'-chloro-2-[[rac-(2R)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]azepan-3-yl]cyanamide




[00541] Compound **44** was prepared similarly to that of **Ex. 11** and **Ex. 13** as a TFA salt. LCMS calcd for $C_{30}H_{36}ClFN_6O_2$ ($M+H$)⁺ $m/z = 567.1$, found 567.2.

[00542] Compound 45. [3-chloro-5-[(7S)-4'-chloro-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-morpholino-methanone




[00543] Compound **45** was prepared from **27B-a** similar to that of **Ex. 11**. LCMS calcd for $C_{35}H_{41}Cl_2FN_7O_4$ ($M+H$)⁺ m/z = 712.3, found: 712.2. ¹H NMR (400 MHz, CD₃OD) δ 7.36 (d, J = 7.4 Hz, 1H), 7.24 (t, J = 7.7 Hz, 1H), 7.13 (d, J = 7.4 Hz, 1H), 5.60 (d, J = 51.9 Hz, 1H), 4.94 – 4.74 (m, 3H), 4.62 (d, J = 14.5 Hz, 1H), 4.45 – 4.44 (m, 2H), 4.42 – 4.31 (m, 2H), 4.03 – 3.81 (m, 6H), 3.77 (s, 3H), 3.69 (s, 3H), 3.46 (td, J = 11.1, 5.9 Hz, 1H), 3.21 – 2.92 (m, 4H), 2.81 – 2.53 (m, 2H), 2.52 – 2.12 (m, 9H).

[00544] **Compound 46A and 46B.** [(3R)-1-[(7*)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-methyl-3-piperidyl]cyanamide;2,2,2-trifluoroacetic acid




[00545] Compound **46A and 46B** were prepared similarly to that of **Ex. 13**. The solution was purified by prep-HPLC (eluting with CH₃CN in H₂O (0.1% TFA) from 5% to 95%) to give faster eluting P1 (**46A**). ¹H NMR (400 MHz, CD₃OD) δ 7.372-7.333 (m, 1 H), 7.281-7.162 (m, 2 H), 5.597 (d, J =50.8 Hz, 1 H), 4.767-4.541 (m, 4 H), 4.146-3.764 (m, 5 H), 3.535-3.405 (m, 1 H), 3.289-2.944 (m, 6 H), 2.770-2.557 (m, 2 H), 2.487-2.203 (m, 6H), 1.971-1.676 (m, 4 H), 1.332 (s, 3 H) and slower eluting P2 (**46B**). ¹H NMR (400 MHz, CD₃OD) δ 7.357-7.322 (m, 1 H), 7.244-7.069 (m, 2 H), 5.574 (d, J =52 Hz, 1 H), 4.635-4.440 (m, 4 H), 4.151-3.969 (m, 2 H), 3.909-3.809 (m, 3 H), 3.496-3.382 (m, 1 H), 3.184-2.912 (m, 6 H), 2.738-2.534 (m, 2 H), 2.455-2.135 (m, 6H), 2.031-1.592 (m, 4 H), 1.293(s, 3 H).

[00546] **Compound 47.** 3-chloro-5-[(7S)-4'-chloro-2-[[1-(morpholinomethyl)cyclopropyl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00547] Compound **47** was prepared from **27B-a** similarly to that of **Ex. 11** as a formic acid salt. LCMS calcd for $C_{33}H_{38}Cl_2FN_7O_3$ ($M+H$)⁺ m/z = 670.4, found: 670.4. ¹H NMR (400 MHz, CD₃OD) δ 8.45 (s, 1H), 7.32 (d, J = 7.1 Hz, 1H), 7.20 (t, J = 7.7 Hz, 1H), 7.12 (d, J = 6.9 Hz, 1H), 4.78 (d, J = 10.8 Hz, 3H), 4.59 (d, J = 14.2 Hz, 1H), 4.43 (d, J = 4.3 Hz, 2H), 4.26 – 4.14 (m, 2H), 3.85 (d, J = 5.7 Hz, 2H), 3.67 (t, J = 4.4 Hz, 4H), 3.10 (t, J = 8.1 Hz, 7H), 3.02 – 2.85 (m, 3H), 2.56 (d, J = 36.2 Hz, 6H), 2.39 (ddd, J = 12.1, 7.8, 4.1 Hz, 1H), 2.23 (dd, J = 12.1, 6.8 Hz, 3H), 0.66 (s, 2H), 0.50 (s, 2H).

[00548] **Compound 48.** 5-[4'-(difluoromethoxy)-2-[[rac-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00549] Compound **48** was prepared similarly to that of **Ex. 9** as a formic acid salt. LCMS calcd for $C_{34}H_{40}F_3N_7O_4$ ($M+H$)⁺ m/z = 668.4, found: 668.4. ¹H NMR (400 MHz, DMSO) δ 7.53 – 7.30 (m, 2H), 7.28 – 7.14 (m, 1H), 7.11 – 6.90 (m, 1H), 6.53 (s, 1H), 5.56 (d, J = 52.8 Hz, 1H), 4.73 (d, J = 14.5 Hz, 3H), 4.45 (dd, J = 30.2, 13.4 Hz, 5H), 4.11 – 3.72 (m, 6H), 3.25 (s, 6H), 3.11 – 2.54 (m, 8H), 2.26 – 1.84 (m, 6H). ¹⁹F NMR (377 MHz, CDCl₃) δ -80.44 (dd, J = 129.9, 23.5 Hz), -171.65 (d, J = 32.3 Hz).

[00550] **Example 16.** Exemplary synthesis of [3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-piperazin-1-yl-methanone (**49**)




[00551] Step 1. A solution of [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (55.23mg, 0.35mmol) in DMSO (1mL) was added sodium hydride (20.82mg, 0.52mmol) at 25 °C for 0.5 h. After 0.5h, a solution of [ethyl 3-chloro-5-[(7S)-4'-chloro-2-methylsulfinyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-

tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**49a**, prepared similarly to that of **27d** of **Ex. 11**) (100.mg, 0.17mmol) in DMSO (1mL) was added to above solution at 25 °C for 0.5 h. The mixture was poured into the mixture of NH₄Cl (10 mL) and iced water (10 mL), extracted with DCM/MeOH(10:1)(10*3 mL), dried over Na₂SO₄, concentrated. Water was added (10 mL) to the residue and the resulting precipitate was filtered, and dried under vacuum to afford 3-chloro-5-[(7S)-4'-chloro-2-[(2R,8R)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methyl]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylic acid (**49b**) (50 mg , 0.0717 mmol, 41.34 % yield) was obtained as a white solid


[00552] Step 2. To a solution of 3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylic acid (**49b**) (40.mg, 0.06mmol) and DIEA (24.1mg, 0.19mmol) in DCM (3mL) was added HATU (30.79uL, 0.12mmol) at 0 °C. The mixture was stirred at 25 °C for 10 min. Then, Piperazine (53.54mg, 0.62mmol) was added at 0 °C. The mixture was stirred at 25 °C for 2 h. The mixture was concentrated to afford a crude product. The crude product was purified by Prep-HPLC. [3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-piperazin-1-yl-methanone (**49**) (11 mg , 0.0135 mmol, 21.68 % yield) was obtained as white solid. LCMS calcd for C₃₅H₄₂Cl₂FN₈O₃ (M+H)⁺ m/z = 711.3, found: 711.3. ¹H NMR (400 MHz, CD₃OD) δ 7.32 (d, *J* = 7.8 Hz, 1H), 7.54 - 7.52 (m, 1H), 7.11 (d, *J* = 7.8 Hz, 1H), 5.36 - 5.19 (m, 1H), 4.82 - 4.74 (m, 2H), 4.58 (d, *J* = 14.2 Hz, 1H), 4.44 - 4.41 (m, 2H), 4.10 - 4.01 (m, 2H), 3.85 (brs, 2H), 3.71 (brs, 2H), 3.56 (brs, 2H), 3.20 - 3.05 (m, 3H), 3.00 - 2.92 (m, 4H), 2.90 - 2.84 (m, 3H), 2.80 (brs, 2H), 2.42 - 2.31 (m, 1H), 2.25 - 2.17 (m, 5H), 2.12 - 1.86 (m, 5H). ¹⁹F NMR (377 MHz, CD₃OD) δ -173.42 (s).

[00553] **Compound 50.** [3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-(4-methylpiperazin-1-yl)methanone




[00554] Compound **50** was prepared similarly to that of **Ex. 16**. LCMS calcd for $C_{36}H_{44}Cl_2FN_8O_3$ ($M+H$)⁺ m/z = 725.3, found: 725.3. ¹H NMR (400 MHz, CD₃OD) δ 7.32 (d, J = 8.0 Hz, 1H), 7.20 (m, 1H), 7.11 (d, J = 8.0 Hz, 1H), 5.35 - 5.22 (m, 1H), 4.88 - 4.74 (m, 3H), 4.58 (d, J = 14.3 Hz, 1H), 4.43 - 4.41 (m, 2H), 4.75 (d, J = 4.0 Hz, 1H), 3.85 (brs, 2H), 3.75 (brs, 2H), 3.63 (brs, 2H), 3.20 - 3.04 (m, 3H), 3.02 - 2.88 (m, 4H), 2.51 (brs, 2H), 2.44 (brs, 2H), 2.40 - 2.35 (m, 1H), 2.32 (s, 3H), 2.29 - 2.17 (m, 5H), 2.12 - 1.82 (m, 6H). ¹⁹F NMR (377 MHz, CD₃OD) δ -173.41 (s).

[00555] **Compound 51A and 51B.** (5R)-9-[(7*)-5'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-7'-hydroxy-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione



[00556] Compound **51A and 51B** were prepared similarly to that of **Ex. 9**. The mixture was purified by Prep-HPLC on a C18 column (5 μ M, 50 x 150 mm) with mobile phase: H₂O (0.1% NH₄HCO₃) / MeCN at flow rate: 35 mL/min to afford to give faster eluting P1 (**51A**): ¹H NMR (400 MHz, CD₃OD) δ 6.78 (d, J = 2.4 Hz, 1H), 6.73 (d, J = 2.4 Hz, 1H), 5.29 (d, J = 54.4 Hz, 1H), 4.78 - 4.82 (m, 2H), 4.12 (s, 2H), 4.07 (d, J = 13.2 Hz, 1H), 3.68 (d, J = 13.2 Hz, 1H), 3.37 (d, J = 13.2 Hz, 1H), 3.15 - 3.28 (m, 4H), 3.01 (s, 1H), 2.95 (d, J = 11.2 Hz, 1H), 2.62 - 2.83 (m, 3H), 2.03 - 2.36 (m, 7H), 1.78 - 1.98 (m, 7H); and slower eluting P2 (**51B**): ¹H NMR (400 MHz, CD₃OD) δ 6.78 (d, J = 2.4 Hz, 1H), 6.76 (d, J = 2.8 Hz, 1H), 5.32 (d, J = 53.6 Hz, 1H), 4.79 - 4.81 (m, 2H), 4.13 - 4.21 (m, 2H), 4.03 (d, J = 13.2 Hz, 1H), 3.76 (d, J = 13.6 Hz, 1H), 3.35 (d, J = 6.0 Hz, 1H), 2.99 - 3.14 (m, 4H), 2.96 (d, J = 8.4 Hz, 1H), 2.62 - 2.83 (m, 4H), 1.83 - 2.21 (m, 14H). Benzyl (*R*)-2,4-dioxo-1,3,7-triazaspiro[4.5]decane-7-carboxylate (P2) was separated on a CHRALCEL OZ-3 column from a racemic mixture. The faster eluting P1 was determined as an *S* enantiomer by a single crystal structure. Removal of Cbz from P2 under hydrogenation conditions afforded (*R*)-1,3,7-triazaspiro[4.5]decane-2,4-dione, which was used to prepare compounds **51A** and **51B**.


[00557] **Example 17.** Exemplary synthesis of [3-chloro-5-[(7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-(1-imino-1-oxo-1,4-thiazinan-4-yl)methanone (**52**)



[00558] Step 1. To a solution of 3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylic acid (**49b**) (50 mg, 0.08 mmol) and DIEA (30.12 mg, 0.23 mmol) in DMF (3 mL) was added HATU (59.09 mg, 0.16 mmol) at 0 °C. The mixture was stirred at 25 °C for 10 min. Then, thiomorpholine (16.03 mg, 0.16 mmol) was added at 0 °C. The mixture was stirred at 25 °C for 2 h. The reaction mixture was diluted with water (20 mL), lots of white solid emerged. The mixture was filtered, and the crude product was recrystallized from EtOAc/PE/MeOH (2 mL/2 mL/0.5 mL) to afford [3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-yl]-thiomorpholino-methanone (**52a**) (35 mg, 0.031 mmol, 40% yield).

[00559] Step 2. The mixture of [3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-yl]-thiomorpholino-methanone (**52a**) (25 mg, 0.03 mmol), Iodobenzene Diacetate (23.21 mg, 0.07 mmol) and Ammonium acetate (4.23 mg, 0.05 mmol) in Methanol (3 mL) was stirred at 25 °C for 2 h. The mixture was quenched with H₂O (10 mL) and Na₂SO₃ (2 mL) at 25 °C, extracted with DCM/MeOH (10/1) (5*3 mL), dried over Na₂SO₄ and concentrated. The crude product was purified by Prep-HPLC to afford [3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-yl]-(1-imino-1-oxo-1,4-thiazinan-4-yl)methanone (**52**) (3 mg, 0.0037 mmol, 11% yield). LCMS calcd for C₃₅H₄₂Cl₂FN₈O₄S (M+H)⁺ m/z = 759.2, found: 759.3. ¹H NMR (400 MHz, CD₃OD) δ 7.32 (d, J = 8.0 Hz, 1H), 7.22 - 7.18 (m, 1H), 7.11 (d, J = 8.0 Hz, 1H), 5.37 - 5.18 (m, 2H), 4.81 - 4.75 (m, 2H), 4.58 (d, J = 14.3 Hz, 2H), 4.48 - 4.41 (m, 2H), 4.38 - 4.18 (m, 2H), 4.12 - 3.92 (m, 4H), 3.89 - 3.81 (m, 2H), 3.25 - 3.05 (m, 6H), 3.05 - 2.88 (m, 4H), 2.35 - 1.88 (m, 11H). ¹⁹F NMR (376 MHz, CD₃OD) δ -173.40 (s).

[00560] **Example 18.** 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**53**)



[00561] Step 1. To a solution of Ethyl Acetate (10.89 mL, 111.2 mmol) in THF (80 mL) was added LDA (55.60 mL, 111.20 mmol) drop wise at -70°C under N_2 . Then the mixture was stirred at -70°C for 3 h, and triisopropoxytitanium (IV) chloride (118.61 mL, 118.61 mmol) was added drop wise at same temperature and stirred for 30 min, then (R,E)-N-(4-chloro-2,3-dihydro-1H-inden-1-ylidene)-2-methylpropane-2-sulfinamide (**28A-a**) (10.00 g, 37.07 mmol) was added drop wise. The resulting mixture was stirred further 3h at -70°C . Then quenched with NH_4Cl solution and extracted with EtOAc. The organic layer was washed with water and brine, dried over Na_2SO_4 , concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 50% EtOAc/PE) to afford the product ethyl 2-[(1S)-1-[(R)-tert-butylsulfinyl]amino]-4-chloro-indan-1-yl]acetate (**53a**) (7.90 g, 22.10 mmol, 59.55 % yield) as yellow solid. LCMS calculated for $\text{C}_{17}\text{H}_{24}\text{ClNO}_3\text{SNa}$ ($\text{M}+\text{Na}$) $^{+}$ m/z = 380.11, 382.10; found: 380.3, 382.3.

[00562] Step 2. To the mixture of ethyl 2-[(1S)-1-[(R)-tert-butylsulfinyl]amino]-4-chloro-indan-1-yl]acetate (**53a**) (7.90 g, 22.07 mmol) in Ethanol (70 mL) was added the solution of HCl/dioxane (22.07 mL, 88.29 mmol) and the mixture was stirred at room temperature for 1 h. The solvent was removed by concentration to afford crude ethyl 2-[(1S)-1-amino-4-chloro-indan-1-yl]acetate hydrochloride (**53b**) (6.40 g, 22.1 mmol, 99.92 % yield), which was used directly in the next step. LCMS calculated for $\text{C}_{13}\text{H}_{18}\text{ClNO}_2$ ($\text{M}+\text{H}$) $^{+}$ m/z = 290.07, 292.07; found: 151.1, 152.1

[00563] Step 3. The mixture of ethyl 2-[(1S)-1-amino-4-chloro-indan-1-yl]acetate;hydrochloride (**53b**) (6.40 g, 22.05 mmol), ethyl prop-2-enoate (15.46 g, 154.38 mmol), Et_3N (9.23 mL, 66.16 mmol) and CuO (0.35 g, 4.41 mmol) in Ethanol (50 mL) was heated to 85°C in a seal tube. The crude reaction mixture was filtered, and filtrate was concentrated in vacuo to give the crude product, which was purified by flash column chromatography (silica gel, eluting with 0% to 50% PE/EtOAc) to afford ethyl 3-[(1S)-4-chloro-1-(2-ethoxy-2-oxo-ethyl)indan-1-yl]amino]propanoate (**53c**) (7.30 g, 20.6 mmol, 93.54 % yield) as a colorless oil. LCMS calculated for $\text{C}_{18}\text{H}_{25}\text{ClNO}_4$ ($\text{M}+\text{H}$) $^{+}$ m/z = 354.15, 356.14; found: 354.3, 356.3.

[00564] Step 4. The mixture of ethyl 3-[[[(1S)-4-chloro-1-(2-ethoxy-2-oxo-ethyl)indan-1-yl]amino]propanoate (**53c**) (2.00 g, 5.65 mmol), Paraformaldehyde (2.04 g, 67.83 mmol) and NaBH₃CN (1.07 g, 16.96 mmol) in Ethanol (20 mL) was stirred at 25 °C for 16 h. The reaction mixture was quenched with water, and then extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 20% EtOAc/PE) to afford ethyl 3-[[[(1S)-4-chloro-1-(2-ethoxy-2-oxo-ethyl)indan-1-yl]-methyl-amino]propanoate (**53d**) (1.90 g, 5.16 mmol, 91.38 % yield) as a yellow oil. LCMS calculated for C₁₉H₂₇ClNO₄ (M+H)⁺ m/z = 368.16, 370.16; found: 368.3, 370.3.

[00565] Step 5. To a solution of ethyl 3-[[[(1S)-4-chloro-1-(2-ethoxy-2-oxo-ethyl)indan-1-yl]-methyl-amino]propanoate (**53d**) (1.10 g, 2.99 mmol) in THF (11 mL) was added [bis(trimethylsilyl)amino]potassium (8.97 mL, 8.97 mmol) dropwise at -70 °C under N₂. The mixture was stirred at same temperature for 2 h. The reaction mixture was quenched with NH₄Cl at -70 °C, and then extracted with EtOAc. The organic layer was washed with water and brine, dried over Na₂SO₄, concentrated, and purified by flash column chromatography (silica gel, eluting with 0% to 20% EtOAc/PE) to afford ethyl (1S)-4-chloro-1'-methyl-4'-oxo-spiro[indane-1,6'-piperidine]-3'-carboxylate (**53e**) (860 mg, 2.67 mmol, 89.37 % yield) as a yellow oil. LCMS calculated for C₁₇H₂₁ClNO₃ (M+H)⁺ m/z = 322.12, 324.12; found: 322.3, 324.3.

[00566] Step 6. To a solution of ethyl (1S)-4-chloro-1'-methyl-4'-oxo-spiro[indane-1,6'-piperidine]-3'-carboxylate (**53e**) (860.00 mg, 2.67 mmol) in Ethanol (10 mL) was added Thiourea (406.86 mg, 5.34 mmol) and C₂H₅ONa (545.58 mg, 8.02 mmol). Then the mixture was stirred at 80 °C under N₂ for 12 h. The reaction mixture was concentrated, then adjust pH to 6 with hydrochloric acid (1N), precipitate the solid, filter and dry the filter cake to afford (7S)-4'-chloro-6-methyl-2-sulfanyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-ol (**53f**) (890.00 mg, 2.67 mmol, 99.76 % yield) as brown solid. LCMS calculated for C₁₆H₁₇ClN₃OS (M+H)⁺ m/z = 334.08, 336.07; found: 334.2, 334.2.

[00567] Step 7. To a solution of (7S)-4'-chloro-6-methyl-2-sulfanyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-ol (**53f**) (890.0 mg, 2.67 mmol) in Water (10 mL) was added Chloroacetic Acid (1.26 g, 13.33 mmol) under N₂. Then the mixture was stirred at 100 °C for 16 h. Upon completion, the reaction was filtered, and the pH of the filtrate was adjusted to 8 with NaHCO₃ saturated aqueous. The solid was collected by filtration and dried to afford (7S)-4'-chloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-2,4-diol (**53g**) (614 mg, 1.93 mmol, 72.48% yield) as a brown solid. LCMS calculated for C₁₆H₁₇ClN₃O₂ (M+H)⁺ m/z = 318.10, 320.10; found: 318.2, 320.3.


[00568] Step 8. The mixture of (7S)-4'-chloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-2,4-diol (**53g**) (590.0 mg, 1.86 mmol) in Phosphorus Oxychloride (4 mL, 42.91 mmol) was stirred at 100 °C for 5 h. LCMS showed SM was consumed, then concentrated, diluted with DCM and purified by flash column chromatography (silica gel, eluting with 0% to 100% EtOAc in PE). Then concentrated and diluted with EtOAc, washed with NaHCO₃ aqueous and brine, dried over Na₂SO₄ and concentrated to afford (7S)-2,4,4'-trichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane] (**53h**) (432.0 mg, 1.22 mmol, 65.60% yield) as a yellow solid. LCMS calculated for C₁₆H₁₅Cl₃N₃ (M+H)⁺ m/z = 354.04, 356.03; found: 354.1, 356.1.

[00569] Step 9. The mixture of (7S)-2,4,4'-trichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane] (**53h**) (150.0 mg, 0.42 mmol), 3-chloro-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide;hydrochloride (118.07 mg, 0.42 mmol) and DIEA (109 mg, 0.85 mmol) in DMF (2 mL) was stirred at room temperature for 30 h. The reaction mixture was diluted with water, extracted with EtOAc, washed with brine, dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 3% to 7% MeOH/DCM) to afford 3-chloro-5-[(7S)-2,4'-dichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**53i**) (120.0 mg, 0.2139 mmol, 50.58 % yield) as a white solid. LCMS calculated for C₂₆H₂₉Cl₃N₇O (M+H)⁺ m/z = 560.15, 562.15; found: 560.3, 562.3.

[00570] Step 10. The mixture of 3-chloro-5-[(7S)-2,4'-dichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**53i**) (20.0 mg, 0.036 mmol), [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (7.38 mg, 0.046 mmol), Cs₂CO₃ (29.06 mg, 0.089 mmol), Pd₂dba₃ (3.27 mg, 0.004 mmol) and Ruphos (3.33 mg, 0.007 mmol) in Toluene (1 mL) was stirred at 110 °C for 10 h under N₂. The reaction mixture was cooled to room temperature and the solvent was removed under reduced pressure. The mixture was purified by flash column chromatography (silica gel, eluting with 0% to 12% MeOH / DCM) followed by prep-HPLC(0.1% NH₄HCO₃) to afford 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**53**) (9.78 mg, 0.015 mmol, 42.25% yield) as a white solid. LCMS calculated for C₃₄H₄₃ClFN₈O₂ (M+H)⁺ m/z = 649.32; found: 649.3. ¹H NMR (400 MHz, CD₃OD) δ 7.13 – 7.34 (m, 4H), 5.26 (d, J = 54.4 Hz, 1H), 4.75 – 4.83 (m, 2H), 4.36 – 4.54 (m, 2H), 3.97 – 4.09 (m,


3H), 3.68 – 3.93 (m, 3H), 3.11 – 3.28 (m, 3H), 3.10 (s, 3H), 3.08 (s, 3H), 2.80 – 3.05 (m, 5H), 2.21 – 2.47 (m, 3H), 2.19 (s, 3H), 1.80 – 2.18 (m, 7H).

[00571] Compound 54. 3-chloro-N,N-dimethyl-5-[rac-(7S)-4'-chloro-2-[(6-methylene-2,3,5,7-tetrahydro-1H-pyrrolizin-8-yl)methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00572] Compound 54 was prepared from **27B-a** similarly to that of **Ex. 11** as a formic acid salt. LCMS calcd for $C_{34}H_{40}Cl_2N_7O_3$ ($M+H$)⁺ m/z = 664.3, found: 664.5/666.5. ¹H NMR (400 MHz, CD₃OD) δ 7.32 (d, J = 7.9 Hz, 1H), 7.20 (t, J = 7.7 Hz, 1H), 7.10 (d, J = 7.4 Hz, 1H), 5.34 – 5.20 (m, 2H), 4.83 – 4.71 (m, 3H), 4.59 (d, J = 14.4 Hz, 1H), 4.48 – 4.32 (m, 4H), 4.28 (d, J = 14.5 Hz, 1H), 3.94 – 3.75 (m, 3H), 3.74 – 3.64 (m, 1H), 3.23 – 3.04 (m, 8H), 3.04 – 2.89 (m, 4H), 2.73 (d, J = 15.9 Hz, 1H), 2.45 – 2.28 (m, 2H), 2.28 – 2.00 (m, 6H).


[00573] Example 19. Exemplary synthesis of 3-chloro-5-((S)-4-chloro-2'-(((2R,7aS)-2-fluorotetrahydro-1H-pyrrolizin-7a(5H)-yl)methoxy)-6'-methyl-2,3,5',8'-tetrahydro-6'H-spiro[indene-1,7'-pyrido[4,3-d]pyrimidin]-4'-yl)-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**55**)



[00574] To a solution of [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (76.63 mg, 0.48 mmol) in THF (1 mL) was added NaH (16.05 mg, 0.4 mmol) at 0°C under N₂. The mixture was stirred at rt for 1 h, and then 3-chloro-5-[(7S)-2,4'-dichloro-6-methyl-spiro[5,8-dihydropyrro[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**53i**) (90.0 mg, 0.16 mmol) was added. The resulting mixture was stirred at 70 °C for 16 h. The reaction mixture was quenched with 1M HCl in MeOH, concentrated, purified by flash column chromatography (silica gel, eluting with 0% to 15% MeOH/DCM), and then purified by prep-HPLC (0.1% NH₄HCO₃) to afford 3-


chloro-5-[(7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**55**) (44.52 mg, 0.0635 mmol, 39.55% yield) as a white solid. LCMS calculated for $C_{34}H_{42}Cl_2FN_8O_2$ ($M+H$)⁺ m/z = 683.28, 685.28; found: 683.4, 685.4. ¹H NMR (400 MHz, CD₃OD) δ 7.23 – 7.34 (m, 2H), 7.17 (dd, J = 7.2, 1.6 Hz, 1H), 5.27 (d, J = 54.0 Hz, 1H), 4.75 – 4.83 (m, 2H), 4.36 – 4.53 (m, 2H), 3.95 – 4.11 (m, 3H), 3.67 – 3.94 (m, 3H), 3.14 – 3.27 (m, 3H), 3.10 (s, 3H), 3.08 (s, 3H), 2.91 – 3.06 (m, 3H), 2.83 – 2.91 (m, 2H), 2.23 – 2.51 (m, 3H), 2.20 (s, 4H), 1.81 – 2.19 (m, 7H).

[00575] Compound 56. [3-chloro-5-[(7R)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-(1,1-dioxo-1,4-thiazinan-4-yl)methanone (**56**)



[00576] Compound 56 was prepared similarly to that of Ex. 16. LCMS calcd for $C_{35}H_{40}Cl_2FN_7O_5S$ ($M+H$)⁺ m/z = 760.2, found: 760.5.

[00577] Example 20. Exemplary synthesis of 1-[3-chloro-5-[(7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-N,N-dimethyl-methanamine (**57**)



[00578] Step 1. To a solution of (7S)-4,4'-dichloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane] (**27B-a P2**) (300.mg, 0.85mmol) in NMP (8mL) were added ethyl 3-chloro-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxylate (517.36mg, 2.12mmol) and DIPEA (329.26mg, 2.55mmol) at 25°C. The mixture was stirred at 100 °C for 16 h. The mixture was diluted with EtOAc (60 mL), washed with water (30*3 mL) and brine (30 ml), dried over Na₂SO₄, concentrated. The crude product was purified by flash chromatography (eluted with EtOAc in PE from 0% to 50%) to afford ethyl 3-chloro-5-[(rac-(7S)-4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-

4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**57a**) (300 mg, 0.535 mmol, 63.03%) yield as a pale white solid. LCMS calcd for $C_{26}H_{27}Cl_2N_5O_3S$ (M+H)⁺ m/z = 560.1, found: 560.2.

[00579] Step 2. To a solution of ethyl 3-chloro-5-[(7S)-4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**57a**) (65 mg, 0.12 mmol) in DCM (1 mL) was added diisobutylaluminum hydride (0.23 mL, 0.23 mmol) at -78 °C. The mixture was stirred at -78 °C for 1 h. IPA (1 mL) and water (1 mL) were added at -78 °C, and the reaction mixture was allowed back to rt and stirred for 1 h. The mixture was diluted with DCM (20 mL) and filtered. The solution was washed with brine (20 mL), dried over Na₂SO₄, concentrated to provide the crude product 3-chloro-5-[(7S)-4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbaldehyde (**57b**) (53 mg, 0.10 mmol, 88.5% yield), which was used directly in the next step. LCMS calcd for $C_{24}H_{23}Cl_2N_5O_2S$ (M+H)⁺ m/z = 516.1 found: 516.2.


[00580] Step 3. To a solution of 3-chloro-5-[(7S)-4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbaldehyde (**57b**) (60 mg, 0.12 mmol) in DCM (2 mL) was added Dimethylamine (0.35 mL, 0.35 mmol) at 25 °C. The mixture was stirred at 25 °C for 1 h. Then Sodium triacetoxyborohydride (73.91 mg, 0.35 mmol) and AcOH (0.7 mg, 0.01 mmol) was added the reaction. The mixture was stirred at 25 °C for 1 h. The mixture was quenched with H₂O (20 mL) at 25 °C, extracted with DCM (20 mL x 3), dried over Na₂SO₄, concentrated. The crude product was purified by Prep-HPLC (eluted with CH₃CN in H₂O (0.1 % NH₄HCO₃) from 5.0% to 95%) to give 1-[3-chloro-5-[(7S)-4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-yl]-N,N-dimethyl-methanamine (**57c**) (34 mg, 0.0623 mmol, 53.6% yield) as a white solid. LCMS calcd for $C_{26}H_{30}Cl_2N_6OS$ (M+H)⁺ m/z = 545.2 found: 545.1.

[00581] Step 4. To a solution of 1-[3-chloro-5-[(7S)-4'-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-yl]-N,N-dimethyl-methanamine (**57c**) (40 mg, 0.07 mmol) in THF (2 mL) was added a solution of oxone (45.08 mg, 0.15 mmol) in Water (1 mL) at 0 °C. Then the mixture was stirred at 25 °C for 3 h. The mixture was added water (30 mL) and extracted with EtOAc (40 mL x 3), washed with a solution of Na₂SO₃ (20 mL), brine (50 mL), dried over Na₂SO₄, concentrated in vacuum to give the crude product. The aqueous layer was adjusted to pH = 9 with the NaHCO₃ solution. The mixture was extracted with EtOAc (40 mL x 3), dried over Na₂SO₄, concentrated in vacuum to give the crude product 1-[3-chloro-5-[(7S)-4'-chloro-2-

methylsulfinyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-N,N-dimethyl-methanamine (**57d**) (37 mg, 0.0659 mmol, 89.86 %yield) as a white solid. LCMS calcd for $C_{26}H_{30}Cl_2N_6O_2S$ (M+H)⁺ m/z = 561.2, found: 561.1.


[00582] Step 5. To a solution of [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (26.22mg, 0.16mmol) in DMF (1mL) was added NaH (5.27mg, 0.13mmol) at 25 °C. The mixture was stirred at 25 °C for 15 min. A solution of 1-[3-chloro-5-[(7S)-4'-chloro-2-methylsulfinyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-N,N-dimethyl-methanamine (**57d**) (37. mg, 0.07mmol) in DMF (0.5mL) was added and the mixture was stirred at 25 °C for 1 h. Acetic acid (0.01mL, 0.13mmol) was added the reaction mixture. The crude product was purified by Prep-HPLC (FA) to afford 1-[3-chloro-5-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-N,N-dimethyl-methanamine;formic acid (**57**) (24.13mg,0.034mmol, 51.60% yield) as a white solid. LCMS calcd for $C_{33}H_{40}Cl_2FN_7O_2$ (M+H)⁺ m/z = 656.26, found: 656.3. ¹H NMR (400 MHz, CD₃OD) δ 8.46 (s, 1H), 7.32 (d, *J* = 7.8 Hz, 1H), 7.20 (t, *J* = 7.7 Hz, 1H), 7.11 (d, *J* = 7.4 Hz, 1H), 5.46 (d, *J* = 52.5 Hz, 1H), 4.79 (d, *J* = 12.4 Hz, 4H), 4.59 (d, *J* = 14.4 Hz, 1H), 4.45 (d, *J* = 5.2 Hz, 2H), 4.34 (s, 2H), 3.93 (s, 2H), 3.87 (s, 2H), 3.79 – 3.48 (m, 3H), 3.28 – 3.20 (m, 1H), 3.09 (dt, *J* = 21.4, 6.8 Hz, 1H), 3.03 – 2.86 (m, 3H), 2.63 (s, 6H), 2.58 – 2.32 (m, 3H), 2.29 – 2.13 (m, 6H), 2.04 (s, 1H).

[00583] **Compound 58.** (5R)-9-[(7S)-4'-chloro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione



[00584] Compound **58** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{31}H_{38}ClFN_7O_3$ (M+H)⁺ m/z = 610.27, 611.27; found: 610.3, 611.3. ¹H NMR (400 MHz, CD₃OD): δ 7.23 – 7.32 (m, 2H), 7.17 (dd, *J* = 6.9, 1.5 Hz, 1H), 5.26 (dd, *J* = 55.4, 3.3 Hz, 1H), 4.10 (q, *J* = 10.4 Hz, 2H), 3.95 (q, *J* = 13.5 Hz, 2H), 3.70 (q, *J* = 14.8 Hz, 2H), 3.39 (d, *J* = 13.2 Hz, 1H), 3.13 – 3.26 (m, 3H), 2.93 – 3.12 (m, 4H), 2.89 (s, 2H), 2.40 – 2.51 (m, 1H), 2.21 – 2.32 (m, 1H), 2.19 (s, 3H), 2.06 – 2.18 (m, 3H), 1.81 – 2.02 (m, 7H). ¹⁹F NMR (376 MHz, CD₃OD): δ -173.76 (s).

Intermediate 1. Synthesis of 3-chloro-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1**)



[00585] Step 1. Synthesis of 2-nitro-N-prop-2-ynyl-benzenesulfonamide (**Int-1a**). A solution of prop-2-yn-1-amine (1.87 mL, 29.15 mmol) and N,N-Diisopropylethylamine (10.15 mL, 58.3 mmol) in DCM (100 mL) was cooled to 0 °C. 2-nitrobenzenesulfonyl chloride (6460 mg, 29.15 mmol) was added portion wise. Upon complete addition, the solution was allowed to warm to room temperature and was further stirred for 1 hr. The mixture was washed with water and brine, dried over Na₂SO₄, concentrated. The crude product was purified by silica gel chromatography (eluted with EtOAc in petroleum ether from 20% to 50%) to afford 2-nitro-N-prop-2-ynyl-benzenesulfonamide (**Int-1a**, 5984 mg, 24.9 mmol, 85.45 % yield) as a yellow solid. LCMS calculated for C₉H₉N₂O₄S (M+H)⁺ m/z = 241.0, found: 241.0.

[00586] Step 2. Synthesis of N-(3-chloropropyl)-2-nitro-N-prop-2-ynyl-benzenesulfonamide (**Int-1b**). To a mixture of 2-nitro-N-prop-2-ynyl-benzenesulfonamide (**Int-1a**, 20 g, 83.25 mmol) and K₂CO₃ (33.37 g, 241.43 mmol) in Acetone (200 mL) was added neat 1-Bromo-3-Chloropropane (41.16 mL, 416.25 mmol) dropwise. Upon complete addition, the reaction mixture was stirred at 25 °C overnight. The mixture was diluted with EtOAc (300 mL), washed with water (100 mL) and brine (20 mL), dried over Na₂SO₄, concentrated. The mixture was concentrated, then washed with water and brine, dried over Na₂SO₄, concentrated. The crude product from 4 batches on the same scale were combined and purified by silica gel chromatography (eluted with EtOAc in petroleum ether from 10% to 90%) to give N-(3-chloropropyl)-2-nitro-N-prop-2-ynyl-benzenesulfonamide (**Int-1b**, 87.00 g, 275 mmol, 82.48 % yield) as a yellow solid. LCMS calcd for C₁₂H₁₄ClN₂O₄S (M+H)⁺ m/z = 317.0, found: 317.0.

[00587] Step 3. Synthesis of ethyl 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-1c**). To solution of N-(3-chloropropyl)-2-nitro-N-prop-2-ynyl-benzenesulfonamide (**Int-1b**, 10 g, 31.57 mmol) and ethyl 2-diazoacetate (5403.14 mg,

47.35 mmol) in Chlorobenzene (80mL) was added N,N-Diisopropylethylamine (5.5 mL, 31.57 mmol) at 140 °C for 1.5 hours. Upon complete addition, cesium carbonate (12.31 g, 37.88 mmol) was added and again heated to 140 °C for 30 minutes. The solvent was concentrated, and the mixture was extracted with EtOAc, washed with brine, dried over Na₂SO₄, concentrated. The crude product was purified by silica gel chromatography (eluted with EtOAc in petroleum ether from 20% to 90%), then triturated in EtOAc to afford ethyl 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-1c**, 5.80 g, 14.7 mmol, 46.58 % yield) as a yellow solid. LCMS calcd for C₁₆H₁₉N₄O₆S (M+H)⁺ m/z = 395.1, found: 395.0.

[00588] Step 4. Synthesis of ethyl 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-1d**). To a solution of ethyl 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-1c**, 12.6 g, 31.95 mmol) in THF (100 mL) and Methanol (25 mL) a was added 1M LiOH in water (128 mL, 127.79 mmol) at 25 °C. The mixture was stirred at 55 °C for 2 h. The mixture was acidified with HCl (1mol/L in H₂O) to pH= 6 and the crude product was triturated in water and filtered. 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylic acid (**Int-1d**, 11.5 g, 31.1 mmol, 97.28 % yield) was obtained as a yellow solid. LCMS calcd for C₁₄H₁₄N₄O₆S (M+H)⁺ m/z = 367.1, found: 367.0.


[00589] Step 5. Synthesis of N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1e**). To a solution of 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylic acid (**Int-1d**, 9.60 g, 26.2 mmol), DIEA (18.26 mL, 104.82 mmol) and HATU (14.946 g, 39.31 mmol) in DMF (90 mL) was added 2M N-methylmethanamine in THF (20 mL) at 30 °C. The mixture was stirred at 30 °C for 2h. The mixture was diluted with DCM (300x2 mL), washed with water (400 mL) and brine (400x2 mL), dried over Na₂SO₄ and concentrated to afford a crude product N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1e**, 10.1g, 24.4 mmol, 93.07 % yield) as a yellow oil. LCMS calcd for C₁₆H₂₀N₄O₃S (M+H)⁺ m/z =394.1, found: 394.2.

[00590] Step 6. Synthesis of 3-chloro-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1f**). To a solution of N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1e**, 10.1 g, 25.67 mmol) in DMF (100 mL) was added N-Chlorosuccinimide (3.428 g, 25.67mmol) at 0 °C under argon. The mixture was stirred at 45 °C for 1h. The mixture was concentrated to afford a crude product. The crude product was triturated in water and filtered to afford 3-chloro-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-

a][1,4]diazepine-2-carboxamide (**Int-1f**, 10.2 g, 23.4 mmol, 91.00 % yield) as a crude yellow solid. LCMS calcd for $C_{16}H_{17}ClN_4O_6S(M+H)^+$ $m/z = 428.1$, found: 428.0.

[00591] Step 7. Synthesis of 3-chloro-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Intermediate 1**). To a solution of 3-chloro-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1f**, 10.2 g, 23.84 mmol), 4-methoxybenzenethiol (8.8 mL, 71.52 mmol) and CS_2CO_3 (31.067 g, 95.36 mmol) in MeCN (100mL). The mixture was stirred at 25 °C for 2 h. The mixture was concentrated and the crude product was purified by silica gel chromatography (eluted with MeOH in DCM from 3% to 10%) to afford 3-chloro-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1**, 3.20 g, 13.2 mmol, 55.30 % yield) as a yellow solid. LCMS calcd for $C_{10}H_{15}ClN_4O(M+H)^+$ $m/z = 243.1$, found: 243.0.

Intermediate 2. Synthesis of 2-methylsulfanyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Int-2**)



[00592] Step 1. Synthesis of 1,1-dimethoxy-4,4-bis(methylsulfanyl)but-3-en-2-one (**Int-2a**). To a solution of sodium hydride (6.77 g, 169.31 mmol) in THF (100 mL) was added Carbon Disulfide (5.1 mL, 84.65 mmol) at 0 °C. The mixture was stirred at 25 °C for 10 min. Then, a solution of 1,1-dimethoxypropan-2-one (10 g, 84.65 mmol) in THF (30 mL) was added at 0 °C. The mixture was stirred at 25 °C for 2 h. Then, a solution of Methyl Iodide (6.35mL, 101.58mmol) in THF (30 mL) was added at 0 °C. The mixture was stirred at 25 °C for 2 h. The mixture was poured into $NaHCO_3$ (70 mL) and iced water (70 mL), extracted with EtOAc (100*3 mL), dried over Na_2SO_4 , concentrated. The crude product was purified by flash chromatography (eluted with EtOAc in petroleum ether from 0% to 20%) to afford 1,1-dimethoxy-4,4-bis(methylsulfanyl)but-3-en-2-one (**Int-2a**, 6.00 g, 27.0 mmol, 31.88%yield) as a pale-yellow oil.

[00593] Step 2. Synthesis of 5-(dimethoxymethyl)-3-methylsulfanyl-1H-pyrazole (**Int-2b**). To a solution of 1,1-dimethoxy-4,4-bis(methylsulfanyl)but-3-en-2-one (**Int-2a**, 11 g, 49.48 mmol) in Ethanol (210 mL) was added hydrazine hydrate (10.11 g, 197.91 mmol) and the reaction mixture was stirred at 25 °C for 16 h. The mixture was poured into iced water (100 mL), extracted with EtOAc (100*3 mL), dried over Na₂SO₄, concentrated. The crude product was purified by flash chromatography (eluted with EtOAc in petroleum ether from 0% to 50%) to afford 5-(dimethoxymethyl)-3-methylsulfanyl-1H-pyrazole (**Int-2b**, 6.00 g, 31.9 mmol, 64.42 %yield) as a pale-yellow oil.

[00594] Step 3. Synthesis of 3-methylsulfanyl-1H-pyrazole-5-carbaldehyde (**Int-2c**). To a solution of 5-(dimethoxymethyl)-3-methylsulfanyl-1H-pyrazole (**Int-2b**, 2.6 g, 13.81 mmol) in Acetone (30 mL) and Water (10 mL) was added p-Toluenesulfonic acid (1189.16 mg, 6.91 mmol) at 25 °C. The mixture was stirred at 50 °C for 2 h. The mixture was poured into the mixture of NH₄Cl (30 mL) and iced water (30 mL). The mixture was filtered, and the filter cake was evaporated. The solid was dissolved with MeOH (20 mL) and filtered. The filter was dried under vacuum to give 3-methylsulfanyl-1H-pyrazole-5-carbaldehyde (**Int-2c**, 1.8 g, 12.3 mmol, 89.19% yield) was obtained as a pale-yellow solid. LCMS calcd for C₅H₆N₂OS (M+H)⁺ m/z = 143.0, found: 143.2.

[00595] Step 4. Synthesis of 3-[(E)-(3-methylsulfanyl-1H-pyrazol-5-yl)methyleneamino]propan-1-ol (**Int-2d**). The solution of 3-methylsulfanyl-1H-pyrazole-5-carbaldehyde (**Int-2c**, 1.1 g, 7.74 mmol) and 3-aminopropan-1-ol (639.21 mg, 8.51 mmol) in Ethanol (10 mL) was stirred at 80 °C for 16 h. The mixture was concentrated to afford a crude product which was used directly in the next step. 3-[(E)-(3-methylsulfanyl-1H-pyrazol-5-yl)methyleneamino]propan-1-ol (**Int-2d**, 1.7 g, 3.23 mmol, 41.80% yield) was obtained as crude yellow oil. LCMS calcd for C₈H₁₃N₃OS (M-H)⁻ m/z = 199.1, found: 199.1.

[00596] Step 5. Synthesis of 3-[(3-methylsulfanyl-1H-pyrazol-5-yl)methylamino]propan-1-ol (**Int-2e**). To a solution of 3-[(E)-(3-methylsulfanyl-1H-pyrazol-5-yl)methyleneamino]propan-1-ol (**Int-2d**, 1.7 g, 8.53 mmol) in Methanol (30 mL) was added Sodium triacetoxyborohydride (3.62 g, 17.06 mmol) at 0 °C. The mixture was stirred at 25 °C for 6 h. The mixture was concentrated, and the crude product was purified by C18 flash chromatography (eluted with ACN in water from 0% to 7%) to give 3-[(3-methylsulfanyl-1H-pyrazol-5-yl)methylamino]propan-1-ol (**Int-2e**, 1.5 g, 7.45 mmol, 87.35% yield). LCMS calcd for C₈H₁₅N₃OS (M+H)⁺ m/z = 202.2, found: 202.2.

[00597] Step 6. Synthesis of tert-butyl N-(3-hydroxypropyl)-N-[(3-methylsulfanyl-1H-pyrazol-5-yl)methyl]carbamate (**Int-2f**). To a solution of 3-[(3-methylsulfanyl-1H-pyrazol-5-yl)methylamino]propan-1-ol (**Int-2e**, 100 mg, 0.5 mmol) and Et₃N (0.14 mL, 0.99 mmol) in


DCM (3 mL) was added Boc₂O (0.14 mL, 0.6 mmol) at 0 °C. The mixture was stirred at 25 °C for 2 h. The mixture was diluted with DCM (10 mL) and filtered. The solution was washed with water (10 mL) and brine (10 mL), dried over Na₂SO₄, concentrated. The crude product was purified by Prep-TLC (eluted with EtOAc 100%) to give tert-butyl N-(3-hydroxypropyl)-N-[(3-methylsulfanyl-1H-pyrazol-5-yl)methyl]carbamate (**Int-2f**, 50 mg, 0.15 mmol, 30% yield) as a colorless viscous semi-solid. LCMS calcd for C₁₃H₂₄N₃O₃S (M+H)⁺ m/z = 302.1, found: 302.1.

[00598] Step 7. Synthesis of 3-[tert-butoxycarbonyl-[(3-methylsulfanyl-1H-pyrazol-5-yl)methyl]amino]propyl methanesulfonate (**Int-2g**). To a solution of tert-butyl N-(3-hydroxypropyl)-N-[(3-methylsulfanyl-1H-pyrazol-5-yl)methyl]carbamate (**Int-2f**, 50 mg, 0.17 mmol) and in DCM (2 mL) was added MsCl (28.5 mg, 0.25 mmol) at 0 °C. The mixture was stirred at 25 °C for 2 h. The mixture was quenched with H₂O (20 mL), extracted with DCM (10*2 mL), dried over Na₂SO₄, concentrated. The crude product was purified by Prep-TLC (eluted with EtOAc in petroleum ether 50%) to give 3-[tert-butoxycarbonyl-[(3-methylsulfanyl-1H-pyrazol-5-yl)methyl]amino]propyl methanesulfonate (**Int-2g**, 27 mg, 0.069 mmol, 42% yield) as a colorless viscous semi-solid. LCMS calcd for C₁₄H₂₅N₃O₅S₂ (M+Na)⁺ m/z = 402.1, found: 402.0.

[00599] Step 8. Synthesis of tert-butyl 2-methylsulfanyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-2h**). To a solution of 3-[tert-butoxycarbonyl-[(3-methylsulfanyl-1H-pyrazol-5-yl)methyl]amino]propyl methanesulfonate (**Int-2g**, 200 mg, 0.53 mmol) in THF (20 mL) was added sodium hydride (63.24 mg, 1.58 mmol) at 0 °C under N₂. The mixture was stirred at 25 °C for 3 h. The mixture was quenched with the mixture of NH₄Cl (20 mL) and iced water (20 mL), extracted with EtOAc (20*3 mL), dried over Na₂SO₄, concentrated. The crude product was purified by flash chromatography (eluted with EtOAc in petroleum ether from 0% to 35%) to give tert-butyl 2-methylsulfanyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-2h**, 50 mg, 0.176 mmol, 33.5% yield) as a yellow oil. LCMS calcd for C₁₃H₂₁N₃O₂ (M+H)⁺ m/z = 284.2, found: 284.2.


[00600] Step 9. Synthesis of 2-methylsulfanyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 2**). To a solution of tert-butyl 2-methylsulfanyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-2h**, 40 mg, 0.14 mmol) in DCM (2 mL) was added TFA (0.5 mL, 6.53 mmol) at 0 °C. The mixture was stirred at 25 °C for 1 h. The mixture was transferred dropwise onto a saturated aqueous solution of NaHCO₃ (20 mL), extracted with DCM (3x10mL), dried over Na₂SO₄, concentrated. the crude product was purified by Prep-TLC (eluted with MeOH in DCM of 10%) to give 2-methylsulfanyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 2**, 8 mg, 0.044 mmol, 31% yield) a yellow oil. LCMS calcd for C₈H₁₃N₃S (M+H)⁺ m/z = 184.1, found: 184.2.

[00601] Compound 59. 1-[3-chloro-5-[(7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-N-methyl-methanamine




[00602] Compound **59** was prepared similarly to that of **Ex. 20**. LCMS calcd for $C_{33}H_{40}Cl_2FN_7O_2$ ($M+H$)⁺ m/z = 642.59, found: 642.3. ¹H NMR (DMSO-d₆, 400Hz): δ 7.39 (d, J = 8.4 Hz, 1H), 7.25 (t, J = 7.6 Hz, 1H), 7.15 (d, J = 7.2 Hz, 1H), 5.32-5.19 (m, 1H), 4.76-4.70 (m, 3H), 4.50-4.47 (m, 1H), 4.32-4.32 (m, 2H), 3.93 (d, J = 10.4 Hz, 1H), 3.84 (d, J = 10.4 Hz, 1H), 3.75-3.74 (m, 3H), 3.51 (s, 3H), 3.07-3.04 (m, 3H), 3.00-2.97 (m, 2H), 2.93-2.90 (m, 3H), 2.84-2.79 (m, 1H), 2.33-2.30 (m, 2H), 2.25 (s, 3H), 2.17-2.10 (m, 1H), 2.09-2.01 (m, 3H), 1.99-1.98 (m, 1H), 1.93-1.91 (m, 1H).

[00603] Compound 60. (5R)-9-[(7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione




[00604] Compound **60** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{31}H_{39}FN_7O_3$ ($M+H$)⁺ m/z = 576.31; found: 576.3. ¹H NMR (400 MHz, CD₃OD) δ 7.17 – 7.30 (m, 4H), 5.32 (d, J = 53.6 Hz, 1H), 4.21, 4.13 (d, J = 10.8 Hz, 2H), 3.99 (d, J = 13.2 Hz, 1H), 3.91 (d, J = 12.8 Hz, 1H), 3.75, 3.66 (d, J = 14.8 Hz, 2H), 3.40 (d, J = 13.2 Hz, 1H), 3.24 – 3.36 (m, 3H), 2.93 – 3.11 (m, 6H), 2.22 – 2.47 (m, 3H), 2.19 (s, 3H), 1.83 – 2.17 (m, 9H).

[00605] Compound 61. 3-chloro-5-[(7S)-4',7'-difluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00606] Compound **61** was prepared similarly to that of **Ex. 18**. LCMS calculated for $C_{34}H_{41}ClF_3N_8O_2$ ($M+H$)⁺ m/z = 685.3; found: 685.3. ¹H NMR (400 MHz, CD₃OD) δ 7.04 (td, J = 8.8, 3.6 Hz, 1H), 6.97 (td, J = 9.2, 4.0 Hz, 1H), 5.27 (d, J = 54.0 Hz, 1H), 4.80 (d, J = 6.8 Hz, 2H), 4.35 – 4.51 (m, 2H), 4.01, 4.07 (d, J = 10.8 Hz, 2H), 3.99–4.10 (m, 1H), 3.83 – 3.92 (m, 1H), 3.65, 3.79 (d, J = 14.8 Hz, 2H), 3.14 – 3.25 (m, 3H), 3.05 – 3.12 (m, 1H), 3.10 (s, 3H), 3.08 (s, 3H), 2.92 – 3.02 (m, 2H), 2.87 (d, J = 18.0 Hz, 1H), 2.30 – 2.58 (m, 3H), 2.28 (s, 3H), 2.03 – 2.23 (m, 4H), 1.82 – 2.00 (m, 4H).

[00607] **Compound 62.** (7S)-4'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-4-(2-methylsulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl)]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]




[00608] Compound **62** was prepared similarly to that of **Ex. 11** using **intermediate 2**. LCMS calcd for $C_{31}H_{36}ClFN_6O_4S$ ($M+H$)⁺ m/z = 643.2, found: 643.3. ¹H NMR (400 MHz, CDCl₃) δ 7.34 (d, J = 7.6 Hz, 1H), 7.26 - 7.23 (m, 1H), 6.97 (d, J = 7.6 Hz, 1H), 6.67 (brs, 1H), 5.31 (d, J = 53.2 Hz, 1H), 4.68 – 4.32 (m, 6H), 4.11 - 4.04 (m, 3H), 3.69 - 3.63 (m, 1H), 3.43 – 3.08 (m, 8H), 3.06 – 2.85 (m, 3H), 2.40 - 2.14 (m, 6H), 2.07 – 1.89 (m, 4H). ¹⁹F NMR (377 MHz, CDCl₃) δ -172.99 (s).

[00609] **Compound 63.** 3-chloro-5-[(7S)-7'-fluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00610] Compound **63** was prepared similarly to that of **Ex. 18**. LCMS calculated for $C_{34}H_{42}ClF_2N_8O_2$ ($M+H$) $^+$ m/z = 667.31, 668.31; found: 667.2, 668.2. ^1H NMR (400 MHz, CD_3OD) δ = 7.31 (td, J =7.9, 5.1, 1H), 7.09 (d, J =7.5, 1H), 6.93 (dd, J =10.1, 8.4, 1H), 5.26 (d, J =54.6, 1H), 4.75-4.81 (m, 2H), 4.35 – 4.52 (m, 2H), 3.96 – 4.09 (m, 3H), 3.83 – 3.92 (m, 1H), 3.73 (d, J = 14.6, 2H), 3.11 – 3.28 (m, 4H), 3.10 (s, 3H), 3.08 (s, 3H), 2.93 – 3.07 (m, 3H), 2.85 (d, J =18.0, 1H), 2.48 (dt, J =13.2, 8.6, 1H), 2.29 – 2.43 (m, 1H), 2.26 (s, 3H), 2.02 – 2.23 (m, 4H), 1.79 – 2.00 (m, 4H). ^{19}F NMR (376 MHz, CD_3OD) δ = -120.88 (s), -173.46 (s).

Example 21. Exemplary synthesis of 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 64**).



[00611] Step 1. Synthesis of (NE,R)-2-methyl-N-tetralin-1-ylidene-propane-2-sulfinamide (**64a**). A mixture of (R)-2-methylpropane-2-sulfinamide (24.87g, 205.21mmol) and tetraethoxytitanium (42.95mL, 205.21mmol) in Toluene (100mL) was stirred and heated to 90 °C. Then the tetralin-1-one (10.g, 68.4mmol) was added to mixture dropwise and stirred for 5h. The mixture was quenched with Potassium sodium tartrate tetrahydrate (aq, 600 mL), filtered and extract with ethyl acetate and purified by flash column chromatography (silica gel, eluting with 10% to 30% EtOAc/PE) to afford (NE,R)-2-methyl-N-tetralin-1-ylidene-propane-2-sulfinamide (**64a**, 13.60 g, 54.5 mmol, 79.73 %yield) as a brown oil. LCMS calculated for $C_{14}H_{20}NOS$ ($M+H$) $^+$ m/z = 250.1; found: 250.1.

[00612] Step 2. Synthesis of ethyl 2-[(1S)-1-[(R)-tert-butylsulfinyl]amino]tetralin-1-yl]acetate (**64b**). To a solution of Diisopropylamine (16.49mL, 117.34mmol) in THF (230mL) was added butyllithium (44.8 mL, 112 mmol) portion wise at 0 °C under N₂. Then the mixture was stirred at 0 °C for 0.5h then cooled to -70 °C to which was added EtOAc (10.44 mL, 106.67 mmol). The solution was stirred for 1 h upon which was added triisopropoxytitanium(IV) chloride (213.34 mL, 213.34 mmol) at -70 °C and stirred for 30 min, then (NE,R)-2-methyl-N-tetralin-1-ylidene-propane-2-sulfinamide (**64a**, 13.3 g, 53.33 mmol) in THF (34 mL) was added drop wise at -70 °C. The resulting mixture was stirred at -70 °C for 2h. Then quenched with NH₄Cl at -70 °C, extracted with EtOAc. The organic layer was washed with water and brine, dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 5% to 34% EtOAc/PE) to afford ethyl 2-[(1S)-1-[(R)-tert-butylsulfinyl]amino]tetralin-1-yl]acetate (**64b**, 15.43 g, 45.7 mmol, 85.74 %yield) as a yellow oil. LCMS calculated for C₁₈H₂₈NO₃S (M+H)⁺ m/z =338.18, found: 338.3.

[00613] Step 3. Synthesis of ethyl 2-[(1S)-1-amino-7-fluoro-indan-1-yl]acetate (**64c**). The mixture of ethyl 2-[(1S)-1-[(S)-tert-butylsulfinyl]amino]tetralin-1-yl]acetate (**64b**, 2.37 g, 7.02 mmol) and HCL/dioxane (7.02 mL, 28.09 mmol) in Ethanol (8 mL) was stirred at rt for 1 h. Aqueous ammonia was added to the reaction mixture until pH adjusted to 9, then the reaction mixture was extracted with EtOAc, The crude reaction mixture was purified by Prep-HPLC to afford crude ethyl 2-[(1S)-1-amino-7-fluoro-indan-1-yl]acetate (**64c**) as a brown oil. LCMS calculated for C₁₄H₁₉NO₂ (M+H)⁺ m/z = 217.2; found: 217.1

[00614] Step 4. Synthesis of ethyl 3-[(1S)-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]amino]propanoate (**64d**). A mixture of ethyl 2-[(1S)-1-aminotetralin-1-yl]acetate (**64c**, 1.55 g, 6.64 mmol), ethyl prop-2-enoate (4.66 g, 46.5 mmol), Et₃N (2.78 mL, 19.93 mmol) and CuO (0.11 g, 1.33 mmol) was dissolved in Ethanol (20 mL). The vial was sealed, and this mixture was then stirred for 8h at 85 °C. The crude reaction mixture was filtered and purified by flash column chromatography (silica gel, eluting with 0% to 50% PE/EA) to afford ethyl 3-[(1S)-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]amino]propanoate (**64d**, 2000 mg, 6.00 mmol, 90.29 %yield) as a light-yellow oil. LCMS calculated for C₁₉H₂₈NO₄ (M+H)⁺ m/z = 334.2; found: 334.2.

[00615] Step 5. Synthesis of ethyl 3-[methyl-[(1S)-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]amino]propanoate (**64e**). To a solution of ethyl 3-[(1S)-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]amino]propanoate (**64d**, 2 g, 6 mmol) in Ethanol (80 mL) was added Paraformaldehyde (1.08 g, 35.99 mmol) and CH₃BNNa (1.13 g, 18 mmol) . Then the mixture was stirred at 25 °C for 16h, the reaction mixture was quenched with water, and then extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 20% EtOAc/PE) to afford ethyl 3-[methyl-[(1S)-1-

(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]amino]propanoate (**64e**, 1680 mg, 4.84 mmol, 80.61 %yield) as a colorless oil. LCMS calculated for $C_{20}H_{30}NO_4$ (M+H)⁺ m/z = 348.2; found: 348.2.

[00616] Step 6. Synthesis of ethyl (6S)-1-methyl-4-oxo-spiro[piperidine-6,1'-tetralin]-3-carboxylate (**64f**). To a solution of [bis(trimethylsilyl)amino]potassium (14.5mL) in THF (8mL) was added ethyl 3-[methyl-[(1S)-1-(2-ethoxy-2-oxo-ethyl)tetralin-1-yl]amino]propanoate (**64e**, 1.68 g, 4.84mmol) portion wise at -70 °C under N₂. Then the mixture was stirred at -70 °C for 2h. The reaction mixture was quenched with NH₄Cl at -70 °C, and then extracted with EtOAc. The organic layer was washed with water and brine, dried over Na₂SO₄, concentrated, and purified by flash column chromatography (silica gel, eluting with 0% to 20% EtOAc/PE) to afford ethyl (6S)-1-methyl-4-oxo-spiro[piperidine-6,1'-tetralin]-3-carboxylate (**64f**, 1300 mg, 4.31 mmol, 89.21 % yield) as a yellow oil. LCMS calculated for $C_{18}H_{23}NO_3$ (M+H)⁺ m/z = 302.2; found: 302.3.

[00617] Step 7. Synthesis of (7S)-6-methyl-2-sulfanyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-ol (**64g**). To a solution of ethyl (6S)-1-methyl-4-oxo-spiro[piperidine-6,1'-tetralin]-3-carboxylate (**64f**, 1300 mg, 4.31 mmol) in Ethanol (10 mL) was added Thiourea (656.69 mg, 8.63 mmol) and C₂H₅ONa (880.6 mg, 12.94 mmol). Then the mixture was stirred at 80 °C under N₂ for 12h. The reaction mixture was concentrated, then pH was adjusted to 6 with hydrochloric acid (1N), the resulting precipitate was filtered to afford (7S)-6-methyl-2-sulfanyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-ol (**64g**, 1000 mg, 3.19 mmol, 73.97 % yield as yellow solid. LCMS calculated for $C_{17}H_{19}N_3OS$ (M+H)⁺ m/z = 314.13; found: 314.0.

[00618] Step 8. Synthesis of (7S)-6-methylspiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**64h**). To a solution of (7S)-6-methyl-2-sulfanyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-ol (**64g**, 1 g, 3.19 mmol) in Water (12 mL) was added Chloroacetic Acid (1.51 g, 15.95 mmol) under N₂. Then the mixture was stirred at 100 °C for 16h. The reaction was filtered, and the pH of the filtrate was adjusted to 8 with NaHCO₃. The solid was collected by filtration and dried to afford (7S)-6-methylspiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**64h**, 913 mg, 3.07 mmol, 96.23 %yield) as a white solid. LCMS calculated for $C_{17}H_{20}N_3O_2$ (M+H)⁺ m/z = 298.15; found: 298.1.


[00619] Step 9. Synthesis of (7S)-2,4-dichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**64i**). The mixture of (7S)-6-methylspiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**64h**, 300 mg, 1.01 mmol) in POCl₃ (3.21 mL, 35.02 mmol) was stirred at 100 °C for 4.5 h. The reaction was concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 100% EtOAc/PE) to afford (7S)-2,4-dichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**64i**, 360 mg, 0.808 mmol, 80.07

%yield) as an orange solid. LCMS calculated for $C_{17}H_{18}Cl_2N_3$ ($M+H$)⁺ m/z = 334.1, 336.1; found: 334.0/336.0

[00620] Step 10. Synthesis of 3-chloro-5-[(7S)-2-chloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**64j**). To a solution of (7S)-2,4-dichloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**64i**, 3 g, 8.98 mmol) in DMSO (30 mL). 3-chloro-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1**, 2724 mg, 11.22 mmol) and DIEA (4.69 mL, 26.93 mmol) were added. The reaction was stirred at 20 °C over weekends. Then the mixture was filtered and purified by Prep-HPLC on a C18 column (5 μ M, 50 x 150 mm) with mobile phase: H₂O (0.1% NH₄HCO₃) / MeCN at flow rate: 65mL/min to afford 3-chloro-5-[(7S)-2-chloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**64j**, 3525 mg, 6.52 mmol, 72.66 % yield) as a white solid. LCMS calculated for $C_{27}H_{31}Cl_2N_7O$ ($M+H$)⁺ m/z = 540.20, 542.20; found: 540.1, 542.1.


[00621] Step 11. Synthesis of 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 64**). To a solution of [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (3.4 g, 21.36 mmol) in DMSO (70 mL) was added sodium hydride (854.27 mg, 21.36 mmol). The reaction was stirred at rt for 40min. Then 3-chloro-5-[(7S)-2-chloro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**64j**, 2600 mg, 4.81 mmol) was added and the reaction was stirred at 40 °C for 1.5h. The reaction mixture was cooled to rt and diluted with H₂O. It was extracted with EtOAc (50 mL*6) and concentrated to get crude product. The reaction was purified with pre-HPLC on a C18 column with a mobile phase (H₂O-NH₄HCO₃/ACN) to afford 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 64**, 2.3 g, 3.451 mmol, 67.33% yield). LCMS calculated for $C_{35}H_{45}ClFN_8O_2$ ($M+H$)⁺ m/z = 663.33 ; found: 663.4 ¹H NMR (400 MHz, CD₃OD) δ 7.49 (d, J = 7.6 Hz, 1H), 7.04 – 7.24 (m, 3H), 5.27 (d, J = 53.6 Hz, 1H), 4.89 – 4.74 (m, 2H), 4.33 – 4.53 (m, 2H), 4.16 – 3.81 (m, 5H), 3.72 – 3.62 (m, 1H), 3.26 – 2.88 (m, 12H), 2.81 – 2.68 (m, 2H), 2.45 – 2.32 (m, 1H), 2.30 – 2.04 (m, 7H), 2.02 – 1.75 (m, 7H).

[00622] Compound 64B. 3-chloro-5-[(7R)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00623] Compound **64B** was isolated as minor isomer from example **21** on Regis (R,R)Whelk-O1 (250*25 mm 10 um) column on a Waters SFC 150 system (Mobile Phase A: Supercritical CO₂, Mobile Phase B: CO₂/EtOH[0.5%NH₃(7M in MeOH)]=70/30; Flow: 100ml/min) to give faster eluting P1 (**64**) and slower eluting P2 (**64B**). LCMS calculated for C₃₅H₄₅ClFN₈O₂ (M+H)⁺ m/z = 663.33; found: 663.3. ¹H NMR (400 MHz, CD₃OD) δ 7.49 (d, *J* = 7.6 Hz, 1H), 7.04–7.25 (m, 3H), 5.25 (d, *J* = 54.8 Hz, 1H), 4.75, 4.90 (d, *J* = 16.8 Hz, 2H), 4.36–4.56 (m, 2H), 3.81–4.15 (m, 5H), 3.67 (d, *J* = 14.8 Hz, 1H), 2.89–3.28 (m, 12H), 2.69–2.80 (m, 2H), 2.32–2.49 (m, 1H), 2.03–2.31 (m, 7H), 1.71–2.03 (m, 7H).

[00624] Compound 65. 3-chloro-5-[(7S)-4'-fluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00625] Compound **65** was prepared similarly to that of **Ex. 18**. LCMS calculated for C₃₄H₄₂ClF₂N₈O₂ (M+H)⁺ m/z = 667.31; found: 667.3. ¹H NMR (400 MHz, CD₃OD) δ 7.29 (td, *J* = 8.0, 5.2 Hz, 1H), 7.05 (d, *J* = 7.6 Hz, 1H), 6.99 (t, *J* = 8.8 Hz, 1H), 5.26 (d, *J* = 54.8 Hz, 1H), 4.73–4.83 (m, 2H), 4.34–4.53 (m, 2H), 3.96–4.10 (m, 3H), 3.66–3.94 (m, 3H), 3.13–3.26 (m, 3H), 3.10 (s, 3H), 3.08 (s, 3H), 2.83–3.06 (m, 5H), 2.24–2.52 (m, 3H), 2.20 (s, 3H), 1.79–2.19 (m, 7H).

[00626] Compound 66. (5R)-9-[(7S)-7'-fluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione




[00627] Compound **66** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{31}H_{38}F_2N_7O_3$ ($M+H$)⁺ m/z = 594.29; found: 594.4. ¹H NMR (400 MHz, CD₃OD) δ 7.25 – 7.40 (m, 1H), 7.10 (d, J = 7.6 Hz, 1H), 6.89 – 6.99 (m, 1H), 5.26 (d, J = 52.4 Hz, 1H), 3.86 – 4.18 (m, 4H), 3.65 (d, J = 14.8 Hz, 2H), 2.93 – 3.37 (m, 9H), 2.82 – 2.91 (m, 1H), 2.42 – 2.55 (m, 1H), 1.80 – 2.32 (m, 14H).

[00628] Compound **67**. (5R)-9-[(7S)-4',7'-difluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione



[00629] Compound **67** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{31}H_{37}F_3N_7O_3$ ($M+H$)⁺ m/z = 612.3; found: 612.3. ¹H NMR (400 MHz, CD₃OD) δ 7.05 (td, J = 8.8, 3.6 Hz, 1H), 6.98 (td, J = 9.2, 4.0 Hz, 1H), 5.23 (d, J = 52.4 Hz, 1H), 4.05, 4.13 (d, J = 10.4 Hz, 2H), 4.03 (d, J = 13.2 Hz, 1H), 3.92 (d, J = 13.2 Hz, 1H), 3.62, 3.68 (d, J = 14.8 Hz, 2H), 3.34 (d, J = 13.63 Hz, 1H), 3.08 – 3.23 (m, 5H), 2.94 – 3.07 (m, 3H), 2.88 (d, J = 18.0 Hz, 1H), 2.48 – 2.58 (m, 1H), 2.26 (s, 3H), 2.19 – 2.31 (m, 1H), 2.03 – 2.18 (m, 3H), 1.82 – 2.00 (m, 7H).


[00630] Compound **68**. (5R)-9-[(7S)-4'-fluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione



[00631] Compound **68** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{31}H_{38}F_2N_7O_3$ ($M+H$)⁺ m/z = 594.29; found: 594.3. ¹H NMR (400 MHz, CD₃OD) δ 7.21 – 7.39 (m, 1H), 6.92 – 7.11 (m, 2H), 5.26 (d, J = 53.6 Hz, 1H), 3.87 – 4.19 (m, 4H), 3.70 (d, J = 14.8


Hz, 2H), 3.39 (d, $J = 13.2$ Hz, 1H), 2.88 – 3.26 (m, 9H), 2.42 – 2.54 (m, 1H), 2.04 – 2.30 (m, 7H), 1.81 – 2.01 (m, 7H).

[00632] Compound 69. 3-chloro-5-[(7S)-7'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00633] Compound **69** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{34}H_{42}Cl_2FN_8O_2$ ($M+H$)⁺ $m/z = 683.28, 685.28$; found: 683.2, 685.2. ¹H NMR (400 MHz, CD_3OD) δ 7.16 – 7.29 (m, 3H), 5.26 (d, $J = 54.7$ Hz, 1H), 4.77, 4.87 (d, $J = 16.6$ Hz, 2H), 4.34 – 4.54 (m, 2H), 4.02 – 4.11 (m, 1H), 3.98, 4.07 (d, $J = 10.5$ Hz, 2H), 3.84 – 3.92 (m, 1H), 3.65, 3.83 (d, $J = 14.4$ Hz, 2H), 3.10 – 3.30 (m, 4H), 3.09 (d, $J = 6.7$ Hz, 6H), 2.88 – 3.05 (m, 3H), 2.67 (d, $J = 17.8$ Hz, 1H), 2.33 – 2.53 (m, 2H), 2.00 – 2.34 (m, 4H), 2.22 (s, 3H), 1.78 – 1.98 (m, 4H).

[00634] Compound 70. (5R)-9-[(7S)-7'-chloro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione




[00635] Compound **70** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{31}H_{38}ClFN_7O_3$ ($M+H$)⁺ $m/z = 610.27, 611.27$; found: 610.3, 611.3. ¹H NMR (400 MHz, DMSO) δ 10.77 (s, 1H), 8.68 (s, 1H), 7.15 – 7.32 (m, 3H), 5.23 (d, $J = 54.0$ Hz, 1H), 3.80 – 3.94 (m, 4H), 3.75 (d, $J = 15.2$ Hz, 1H), 3.46 (d, $J = 14.4$ Hz, 1H), 3.01 – 3.15 (m, 4H), 2.91 – 3.01 (m, 3H), 2.74 – 2.86 (m, 2H), 2.65 (d, $J = 17.6$ Hz, 1H), 2.32 – 2.44 (m, 1H), 2.11 (s, 3H), 1.85 – 2.08 (m, 5H), 1.66 – 1.85 (m, 6H).

[00636] Compound 71. (5R)-9-[(7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-1,3,9-triazaspiro[4.5]decane-2,4-dione.



[00637] Compound **71** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{32}H_{41}FN_7O_3$ ($M+H$)⁺ m/z = 590.32; found: 590.3. ¹H NMR (400 MHz, CD₃OD) δ 7.45 – 7.57 (m, 1H), 7.05 – 7.26 (m, 3H), 5.23 (d, J = 52.0 Hz, 1H), 3.90 – 4.19 (m, 4H), 3.73 (d, J = 15.0 Hz, 2H), 3.38 (d, J = 13.2 Hz, 1H), 2.89 – 3.24 (m, 7H), 2.71 – 2.80 (m, 2H), 1.75 – 2.29 (m, 17H).


[00638] **Compound 72.** 5-[(7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00639] Compound **72** was prepared similarly to that of **Ex. 19**. LCMS calculated for $C_{35}H_{46}FN_8O_2$ ($M+H$)⁺ m/z = 629.37; found: 629.3. ¹H NMR (400 MHz, DMSO) δ 7.45 (d, J = 7.6 Hz, 1H), 7.20 (t, J = 7.6 Hz, 1H), 7.13 (td, J = 7.6, 1.2 Hz, 1H), 7.07 (d, J = 6.4 Hz, 1H), 6.51 (s, 1H), 5.24 (d, J = 54.0 Hz, 1H), 4.79, 4.72 (d, J = 16.4 Hz, 2H), 4.47 (t, J = 5.2 Hz, 2H), 3.77 – 4.00 (m, 4H), 3.61 – 3.73 (m, 2H), 3.24 (s, 3H), 2.96 – 3.10 (m, 3H), 2.93 (s, 3H), 2.66 – 2.90 (m, 5H), 2.03 – 2.22 (m, 2H), 2.00 (s, 3H), 1.59 – 1.98 (m, 10H).

[00640] **Example 22.** Exemplary synthesis of 5-[5-[(7S)-7'-fluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-3-methyl-1,2,4-oxadiazole

Compound 73.




[00641] Step 1. Synthesis of 3-methyl-5-(5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-2-yl)-1,2,4-oxadiazole (**73a**). To a solution of N-hydroxyacetamide (375.66 mg, 5.07 mmol) and 3 Å molecular sieves in THF (5 mL) were added NaH (76.06 mg, 3.17 mmol). The mixture was stirred at R.T. for 15 min. ethyl 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-1c**, 500 mg, 1.27 mmol) was added to the mixture. The mixture was stirred at 25 °C for 3 h. The mixture was filtered over celite and concentrated. The crude product was purified by flash chromatography (eluted with CH₃CN in H₂O from 5.0% to 95%) to afford the desired product 3-methyl-5-(5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-2-yl)-1,2,4-oxadiazole (**73a**, 230 mg, 1.05 mmol, 82.75 %yield) as a yellow solid. LCMS calculated for C₁₀H₁₃N₅O (M+H)⁺ m/z = 220.1, found: 220.1.

[00642] Step 2. Synthesis of 5-[5-[(7S)-2-chloro-7'-fluoro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-3-methyl-1,2,4-oxadiazole (**73c**). The mixture of (7S)-2,4-dichloro-7'-fluoro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane] (**73b**, prepared similarly as **64i**, 50 mg, 0.15 mmol), 3-methyl-5-(5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-2-yl)-1,2,4-oxadiazole (**73a**, 34.03 mg, 0.16 mmol) and DIEA (0.05 mL, 0.3 mmol) in DMSO (1 mL) was stirred at rt for 68 h. The reaction mixture was diluted with water, extracted with EtOAc, washed with brine, dried over Na₂SO₄, concentrated and purified by prep-HPLC (0.1% NH₄HCO₃) to afford to afford 5-[5-[(7S)-2-chloro-7'-fluoro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-3-methyl-1,2,4-oxadiazole (**73c**, 50 mg, 0.0960 mmol, 64.92 %yield) as a white solid. LCMS calculated for C₂₆H₂₇ClFN₈O (M+H)⁺ m/z = 521.20, 523.19; found: 521.1, 523.1

[00643] Step 3. Synthesis of 5-[5-[(7S)-7'-fluoro-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-3-methyl-1,2,4-oxadiazole (**Compound 73**). To a solution of [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (152.79 mg, 0.96 mmol) in DMSO (3 mL) was added NaH (38.39 mg, 0.96 mmol) under N₂ and stirred 20 min, then the mixture was added 5-[5-[(7S)-2-chloro-7'-fluoro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-3-methyl-1,2,4-oxadiazole (**73c**, 50 mg, 0.1 mmol) and stirred at 40 °C for 1 h. The reaction mixture was quenched with water, and extracted with EtOAc, washed with brine, dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 3% to 12% MeOH/DCM) and purified by prep-HPLC (0.1% NH₄HCO₃) to afford 5-[5-[(7S)-7'-fluoro-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-


methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-3-methyl-1,2,4-oxadiazole (**Compound 73**, 32 mg, 0.0487 mmol, 50.76 %yield) as a white solid. LCMS calculated for $C_{34}H_{40}F_2N_9O_2$ ($M+H$)⁺ m/z = 644.33, 645.33; found: 644.3, 645.3. ¹H NMR (400 MHz, CD₃OD) δ = 7.27 – 7.35 (m, 1H), 7.10 (d, J =7.6, 1H), 6.99 (s, 1H), 6.91 – 6.97 (m, 1H), 5.24 (d, J =53.1, 1H), 4.50 – 4.66 (m, 4H), 3.97 – 4.11 (m, 4H), 3.64 – 3.76 (m, 2H), 3.05 – 3.22 (m, 5H), 2.91 – 3.04 (m, 2H), 2.83 (d, J =18.0, 1H), 2.45 – 2.56 (m, 1H), 2.41 (s, 3H), 2.18 – 2.35 (m, 5H), 2.09 – 2.18 (m, 2H), 2.00 – 2.08 (m, 1H), 1.75 – 1.99 (m, 4H).

[00644] Compound 74. 5-[5-[(7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-3-methyl-1,2,4-oxadiazole



[00645] Compound **74** was prepared similarly to that of **Ex. 22**. LCMS calculated for $C_{35}H_{43}FN_9O_2$ ($M+H$)⁺ m/z = 640.25; found: 640.3. ¹H NMR (400 MHz, CD₃OD) δ 7.49 (d, J = 8.0 Hz, 1H), 7.20 (t, J = 7.6 Hz, 1H), 7.14 (td, J = 7.6, 1.2 Hz, 1H), 7.09 (d, J = 7.6 Hz, 1H), 6.99 (s, 1H), 5.25 (d, J = 54.8 Hz, 1H), 4.86 – 4.96 (m, 2H), 4.50 – 4.67 (m, 2H), 3.96 – 4.12 (m, 4H), 3.70 – 3.86 (m, 2H), 3.12 – 3.24 (m, 3H), 2.89 – 3.04 (m, 3H), 2.70 – 2.80 (m, 2H), 2.41 (s, 3H), 2.13 – 2.37 (m, 3H), 2.12 (s, 3H), 1.73 – 2.08 (m, 9H).

[00646] Example 23. Exemplary synthesis of 3-chloro-5-[6'-(difluoromethyl)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 75**)



[00647] Step 1. Synthesis of 5-(6'-bromo-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-3-chloro-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75b**). To the solution of 6'-bromo-4-chloro-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane] (**75a**, 600 mg, 1.51 mmol) DIEA (0.79 mL, 4.53 mmol) and 3-chloro-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (366.15 mg, 1.51 mmol) in DMF (6 mL) and stirred at 100 °C for 2 h. The mixture was extracted with water (20 mL) and EtOAc (3x10 mL), washed with brine (10 mL), dried with Na₂SO₄, concentrated. The crude product was purified by flash chromatography (eluted with CH₃CN in H₂O from 5.0% to 50%) to afford 5-(6'-bromo-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-3-chloro-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75b**, 800 mg, 1.3246 mmol, 87.80% yield) as yellow solid. LCMS calcd for C₂₆H₂₈BrClN₆O₂S (M+H)⁺ m/z = 605.2, found: 605.2.

[00648] Step 2. Synthesis of 3-chloro-N,N-dimethyl-5-(2-methylsulfanyl-6'-vinyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75c**). The mixture of 5-(6'-bromo-2-methylsulfanyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-3-chloro-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75b**, 500 mg, 0.83 mmol), K₂CO₃ (343 mg, 2.48 mmol), Pd(Dppf)Cl₂ (60.52mg, 0.08 mmol) and potassium;trifluoro(vinyl)boranuide (277 mg, 2.07 mmol) was added in DMSO (10mL) and stirred at 100 °C for 16 h under Ar. The crude product was purified by flash chromatography (eluted with CH₃CN in H₂O from 5.0% to 95%) to afford 3-chloro-N,N-dimethyl-5-(2-methylsulfanyl-6'-vinyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-4,6,7,8-tetrahydropyrazolo[1,5-

a][1,4]diazepine-2-carboxamide (**75c**, 360mg, 0.653mmol, 78.9% yield) as a yellow solid.

LCMS calcd for $C_{28}H_{31}ClN_6O_2S$ (M+H)⁺ m/z = 551.2, found: 551.2.

[00649] Step 3. Synthesis of 3-chloro-5-(6'-formyl-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75d**). To a solution of 3-chloro-N,N-dimethyl-5-(2-methylsulfonyl-6'-vinyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75c**, 320 mg, 0.58 mmol), Potassium osmate(VI) dihydrate (9.04 mg, 0.03 mmol) and Sodium Periodate (496.79 mg, 2.32 mmol) was added in THF (4mL) and Water (1mL) at 25 °C. The mixture was stirred at 50 °C for 16 h. The crude product was purified by flash chromatography (eluted with CH₃CN in H₂O from 5.0% to 95%) to afford 3-chloro-5-(6'-formyl-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75d**, 120 mg, 0.205 mmol, 35.3% yield) as a white solid. LCMS calcd for $C_{27}H_{29}ClN_6O_5S$ (M+H)⁺ m/z = 585.3, found: 585.3.


[00650] Step 4. Synthesis of 3-chloro-5-[6'-(difluoromethyl)-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75e**). To a solution of 3-chloro-5-(6'-formyl-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl)-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75d**, 100 mg, 0.17 mmol), and DAST (0.23 mL, 1.71 mmol) was added in DCM (1mL) at 0 °C. The mixture was stirred at 25 °C for 16 h. The mixture was quenched with NH₄Cl (5 mL), then the mixture was extracted with DCM (3x5 mL) and water (10 mL), dried over Na₂SO₄, concentrated to afford the crude of 3-chloro-5-[6'-(difluoromethyl)-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75e**, 100 mg, 0.165 mmol, 96.38% yield) was obtained as a yellow solid. LCMS calcd for $C_{27}H_{29}ClF_2N_6O_4$ (M+H)⁺ m/z = 607.3, found: 607.3.

[00651] Step 5. Synthesis of 3-chloro-5-[6'-(difluoromethyl)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 75**). To a solution of NaH (13.84 mg, 0.35 mmol) and [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (55.07 mg, 0.35 mmol) in DMF (1 mL) and stirred at 25 °C for 10 min, then 3-chloro-5-[6'-(difluoromethyl)-2-methylsulfonyl-spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**75e**, 70 mg, 0.12 mmol) was added at 25 °C. The mixture was stirred at 25 °C for 10 min. The mixture was extracted with water (10 mL) and EA (3x15 mL), then washed with

brine (10 mL), dried over Na_2SO_4 , concentrated. The crude product was purified by Prep-HPLC (eluted with CH_3CN in H_2O (0.1 % FA) from 5.0% to 95%) to afford 3-chloro-5-[6'-(difluoromethyl)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrano[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide;formic acid (**Compound 75**, 25.1 mg, 0.035 mmol, 30.36% yield) as a white solid. LCMS calcd for $\text{C}_{34}\text{H}_{39}\text{ClF}_3\text{N}_7\text{O}_3$ ($\text{M}+\text{H}$) $^+$ m/z = 686.2, found: 686.2. ^1H NMR (400 MHz, CD_3OD) δ 7.45 (dd, J = 18.7, 7.8 Hz, 2H), 7.31 (s, 1H), 6.72 (t, J = 56.3 Hz, 1H), 5.42 – 5.15 (m, 1H), 4.83 – 4.69 (m, 2H), 4.61 – 4.52 (m, 1H), 4.47 – 4.33 (m, 2H), 4.14 – 4.04 (m, 2H), 3.94 – 3.75 (m, 2H), 3.27 – 3.11 (m, 4H), 3.09 (d, J = 8.1 Hz, 6H), 3.04 – 2.88 (m, 4H), 2.42 – 1.82 (m, 11H).

[00652] Example 24. Exemplary synthesis of 3-cyano-5-[(7S)-7'-fluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide

[00653] Compound 76.



[00654] Step 1. Synthesis of 3-iodo-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76a**). To a solution of N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-1e**, 200 mg, 0.51 mmol) in HOAc (2 mL, 34.94 mmol) was added N-Iodosuccinimide (228.74 mg, 1.02 mmol). The mixture was stirred at 80 °C for 2h. The mixture was diluted with water, extracted with DCM, washed with brine, dried over Na_2SO_4 , concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 5% MeOH/DCM) to afford 3-iodo-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-

carboxamide (**76a**, 264 mg, 0.5084 mmol, 100% yield) as a yellow solid. LCMS calcd for $C_{16}H_{19}IN_5O_5S$ (M+H)⁺ m/z = 520.02, found: 520.1.

[00655] Step 2. Synthesis of 3-cyano-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76b**). The mixture of 3-iodo-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76a**, 630 mg, 1.21 mmol), and CuCN (539.85 mg, 6.07 mmol) in DMF (1 mL) was sealed and stirred at 135 °C for 20 h. After completion, the reaction mixture was diluted with water, extracted with EtOAc, washed with brine, dried over Na₂SO₄, concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 5% MeOH/DCM) to afford 3-cyano-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76b**, 340 mg, 0.8126 mmol, 66.98% yield) as a white solid. LCMS calcd for $C_{17}H_{19}N_6O_5S$ (M+H)⁺ m/z = 419.12; found: 419.2.


[00656] Step 3. Synthesis of 3-cyano-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76c**). To a solution of 3-cyano-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76b**, 340 mg, 0.81 mmol), 4-methoxybenzenethiol (341.76 mg, 2.44 mmol) and cesium carbonate (1056.33 mg, 3.25 mmol) in MeCN (4 mL). The mixture was stirred at 20 °C for 2 h. The mixture was filtered, concentrated and the crude product was purified by silica gel chromatography (eluted with MeOH in DCM from 3% to 10%) to afford 3-cyano-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76c**, 164 mg, 0.703 mmol, 86.52% yield) as a white solid. LCMS calcd for $C_{11}H_{16}N_5O$ (M+H)⁺ m/z = 234.14, found: 234.1.

[00657] Step 4. Synthesis of 5-[(7S)-2-chloro-7'-fluoro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-cyano-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76d**). The mixture of (7S)-2,4-dichloro-7'-fluoro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane] (**73b**, 70 mg, 0.21 mmol), 3-cyano-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (53.11 mg, 0.23 mmol) and DIEA (0.14 mL, 0.83 mmol) in DMSO (1.5 mL) was stirred at 30 °C for 20 h. The reaction mixture was filtered and purified by prep-HPLC (0.1% NH₄HCO₃) to afford 5-[(7S)-2-chloro-7'-fluoro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-cyano-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (62 mg, 0.116 mmol, 55.99% yield) as a white solid. LCMS calculated for $C_{27}H_{29}ClFN_8O$ (M+H)⁺ m/z = 535.22; found: 535.3.

[00658] Step 5. The mixture of 5-[(7S)-2-chloro-7'-fluoro-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-3-cyano-N,N-dimethyl-4,6,7,8-


tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (50 mg, 0.09 mmol), [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (22.32 mg, 0.14 mmol), Pd₂dba₃ (12.84 mg, 0.01 mmol), Ruphos (13.08 mg, 0.03 mmol) and cesium carbonate (91.12 mg, 0.28 mmol) in Toluene (2 mL) was charged with N₂ and sealed. The mixture was stirred at 110 °C for 5h. LCMS showed OK. The mixture was diluted with water, extracted with EtOAc, washed with brine, dried over Na₂SO₄, concentrated and purified by prep-HPLC(0.1%NH₄HCO₃) to afford 3-cyano-5-[(7S)-7'-fluoro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (39.09 mg, 0.0587 mmol, 52.99% yield) as a light yellow solid. LCMS calculated for C₃₅H₄₂F₂N₉O₂ (M+H)⁺ m/z = 658.35; found: 658.5. ¹H NMR (400 MHz, CD₃OD) δ 7.31 (td, *J* = 8.0, 5.2 Hz, 1H), 7.09 (d, *J* = 7.6 Hz, 1H), 6.93 (dd, *J* = 10.0, 8.4 Hz, 1H), 5.29 (d, *J* = 53.8 Hz, 1H), 4.87 – 4.97 (m, 2H), 4.37 – 4.57 (m, 2H), 3.89 – 4.12 (m, 4H), 3.66, 3.77 (d, *J* = 14.4 Hz, 2H), 3.32 (s, 3H), 3.12 – 3.28 (m, 4H), 3.10 (s, 3H), 2.94 – 3.08 (m, 3H), 2.86 (d, *J* = 18.0 Hz, 1H), 2.29 – 2.55 (m, 3H), 2.27 (s, 3H), 2.03 – 2.22 (m, 3H), 1.83 – 1.99 (m, 4H).

[00659] Compound 77. 3-cyano-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00660] Compound 77 was prepared similarly to that of **Ex. 23**. LCMS calculated for C₃₆H₄₃FN₉O₂ (M+H)⁺ m/z = 654.37; found: 654.2. ¹H NMR (400 MHz, CD₃OD) δ 7.50 (dd, *J* = 8.0, 1.2 Hz, 1H), 7.22 (td, *J* = 7.4, 1.6 Hz, 1H), 7.14 (td, *J* = 7.4, 1.2 Hz, 1H), 7.08 (dd, *J* = 7.6, 1.2 Hz, 1H), 5.28 (d, *J* = 53.6 Hz, 1H), 4.87 – 4.98 (m, 2H), 4.32 – 4.62 (m, 2H), 4.11 (d, *J* = 10.8 Hz, 1H), 3.93 – 4.06 (m, 2H), 3.83 – 3.92 (m, 2H), 3.70 (d, *J* = 14.8 Hz, 1H), 3.32 (s, 3H), 3.12 – 3.26 (m, 3H), 3.10 (s, 3H), 2.90 – 3.07 (m, 3H), 2.71 – 2.80 (m, 2H), 2.13 – 2.44 (m, 4H), 2.11 (s, 3H), 1.71 – 2.08 (m, 8H).

[00661] Compound 78. 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00662] Compound **78** was prepared similarly to that of **Ex. 21**. LCMS calculated for $C_{33}H_{40}ClFN_7O_3$, $(M+H)^+$ m/z = 636.29, 638.29; found: 636.3, 638.3. ^1H NMR (400 MHz, DMSO) δ = 7.46 (d, J =7.7 Hz, 1H), 7.19 (t, J =7.5 Hz, 1H), 7.12 (td, J =7.4, 1.2 Hz, 1H), 7.07 (d, J =6.8 Hz, 1H), 5.23 (d, J =53.4 Hz, 1H), 4.73 (d, J =16.4 Hz, 2H), 4.34 – 4.53 (m, 2H), 3.87 – 3.98 (m, 1H), 3.83 (s, 2H), 3.68 – 3.80 (m, 2H), 3.60 (d, J =14.9 Hz, 1H), 2.86 – 3.08 (m, 4H), 2.65 – 2.83 (m, 4H), 2.17 – 2.32 (m, 1H), 1.94 – 2.12 (m, 6H), 1.77 – 1.94 (m, 4H), 1.62 – 1.77 (m, 4H).


[00663] **Example 25.** Exemplary synthesis of 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-amine (**Compound 79**).



[00664] To a sealed vial was charged a solution of 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylic acid (**Compound 78**, 30 mg, 0.05 mmol) in 1,4-Dioxane (1 mL), followed by the addition of azido(phenoxy)phosphoryl]oxybenzene (19.47 mg, 0.07 mmol) and *t*-BuOH (59.42 mg, 0.8 mmol) under N_2 . The mixture was heated at 110 °C for 2 hours. The reaction solution was cooled down to rt, and EtOAc was added. The mixture was washed successively with H_2O , $NaHCO_3$ and brine, dried over $MgSO_4$ and concentrated in vacuo. The residue was purified by prep-HPLC to afford 3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-amine (**Compound 79**, 8.0 mg, 0.0132 mmol, 28.05% yield) as a white solid. LCMS calculated for $C_{32}H_{41}ClFN_8O$ $(M+H)^+$ m/z = 607.31, 609.31; found: 607.3, 609.31. ^1H NMR (400 MHz, CD_3OD) δ = 7.49 (d, J =7.1 Hz, 1H), 7.20 (td, J =7.5, 1.8 Hz,


1H), 7.14 (td, $J=7.4$, 1.3 Hz, 1H), 7.09 (d, $J=7.5$ Hz, 1H), 5.26 (d, $J=54.4$ Hz, 1H), 4.68 (d, $J=16.6$ Hz, 2H), 3.78 – 4.23 (m, 7H), 3.67 (d, $J=14.9$ Hz, 1H), 3.11 – 3.28 (m, 3H), 2.88 – 3.07 (m, 3H), 2.70 – 2.80 (m, 2H), 2.23 – 2.37 (m, 1H), 1.70 – 2.21 (m, 14H).

[00665] **Compound 80.** 4-[[1-[(7S)-6-methyl-4-[(1R,5S)-3,8-diazabicyclo[3.2.1]octan-3-yl]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2-yl]oxymethyl]cyclopropyl]methyl]morpholine



[00666] Compound **80** was prepared similarly to that of **Ex. 21**. LCMS calculated for $C_{32}H_{45}N_6O_2$ ($M+H$)⁺ m/z = 545.4; found: 545.3. ¹H NMR (400 MHz, CD_3OD) δ 7.50 (d, J = 7.6 Hz, 1H), 7.19 (t, J = 7.2 Hz, 1H), 7.13 (t, J = 7.2 Hz, 1H), 7.08 (d, J = 6.8 Hz, 1H), 4.14 – 4.33 (m, 3H), 3.56 – 3.77 (m, 9H), 3.43 (d, J = 12.0 Hz, 1H), 2.86 – 3.09 (m, 3H), 2.75 (d, J = 5.2 Hz, 2H), 2.47 (s, 4H), 2.38 (s, 2H), 2.09 (s, 3H), 1.62 – 2.04 (m, 8H), 0.63 (t, J = 5.6 Hz, 2H), 0.45 (t, J = 5.6 Hz, 2H).

[00667] Intermediate 3.




[00668] Step 1. Synthesis of N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-3-(2-trimethylsilylethynyl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide. To a solution of 3-iodo-N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**76a**, 500 mg, 0.96 mmol), Trimethylsilylacetylene (0.71 mL, 5.09 mmol) and Triethylamine (292.28 mg, 2.89 mmol) in DMF (10 mL) was added Bis(Triphenylphosphine)Palladium(II) Chloride (135.16 mg, 0.19 mmol) and Copper (I) Iodide (36.67 mg, 0.19 mmol) under N_2 . The mixture was stirred at 45°C for 20 h. The reaction solution was filtrated, washed by DCM/MeOH=10:1 and concentrated. The residue was purified by flash chromatography (DCM/MeOH=20:1) to give N,N-dimethyl-5-(2-nitrophenyl)sulfonyl-3-(2-trimethylsilylethynyl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-3a**, 345.5 mg, 0.529 mmol, 54.97% yield). LCMS calculated for $C_{21}H_{28}N_5O_5Si$ ($M+H$)⁺ m/z = 490.15; found: 490.0.

[00669] Step 2. Synthesis of N,N-dimethyl-3-(2-trimethylsilylethynyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide. To a solution of N,N-dimethyl-5-(2-


nitrophenyl)sulfonyl-3-(2-trimethylsilylethynyl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-3a**, 350 mg, 0.54 mmol), 4-methoxybenzenethiol (225.5 mg, 1.61 mmol) in CH₃CN (3 mL) was added cesium carbonate (696.97 mg, 2.14 mmol). The mixture was stirred at 20°C for 2 h. The mixture was filtered, concentrated and the crude product was purified by pre-HPLC on a C18 column with a mobile phase (H₂O-NH₄HCO₃/CH₃CN) to afford N,N-dimethyl-3-(2-trimethylsilylethynyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-3b**, 101 mg, 0.331 mmol, 61.88% yield). LCMS calcd for C₁₅H₂₅N₄OSi (M+H)⁺ m/z = 305.17, found: 305.2.

[00670] Step 3. Synthesis of 3-ethynyl-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide. To a solution of N,N-dimethyl-3-(2-trimethylsilylethynyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-3b**, 101 mg, 0.33 mmol) in Methanol (5 mL) was added K₂CO₃ (105.29 mg, 0.76 mmol). The mixture was stirred at 20 °C for 1h. The mixture was filtered, and the filtrate was concentrated and the residue was purified by flash chromatography (DCM/MeOH) to give 3-ethynyl-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Intermediate 3**, 75 mg, 0.323 mmol, 97.33% yield). LCMS calculated for C₁₂H₁₇N₄O (M+H)⁺ m/z = 233.13 ; found: 233.3.

[00671] **Compound 81.** 3-ethynyl-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00672] Compound **81** was prepared similarly to that of **Ex. 21** using **intermediate 3**. LCMS calculated for C₃₇H₄₆FN₈O₂ (M+H)⁺ m/z = 653.36; found: 653.3. ¹H NMR (400 MHz, CD₃OD) δ 7.45 – 7.53 (m, 1H), 7.04 – 7.24 (m, 3H), 5.26 (d, *J* = 54.4 Hz, 1H), 4.71 – 4.90 (m, 2H), 4.44 – 4.54 (m, 1H), 4.30 – 4.40 (m, 1H), 3.80 – 4.16 (m, 6H), 3.66 (d, *J* = 14.8 Hz, 1H), 2.89 – 3.25 (m, 12H), 2.71 – 2.80 (m, 2H), 2.36 – 2.51 (m, 1H), 2.05 – 2.32 (m, 7H), 1.73 – 2.03 (m, 7H).

[00673] Intermediate 4.


[00674] Step 1. Synthesis of [5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]methanol (**Int-4a**). To a solution of ethyl 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-1c** 1500 mg, 3.8 mmol) in THF (15 mL) was added 1M DABAL-H in hexane (11.41 mL, 11.41 mmol) at -68 °C under argon. The mixture was stirred at 25 °C for 2 h. The reaction was quenched with water and isopropyl alcohol (1:1, 12 mL) and extracted with EtOAc (50 mL x 3) and concentrated. The crude product [5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-yl]methanol (**Int-4a**, 850 mg, 2.39 mmol, 62.79 %yield) was carried over without purification. LCMS calcd for $C_{14}H_{17}N_4O_5S$ ($M+H$)⁺ m/z = 353.1, found: 353.1.

[00675] Step 2. Synthesis of 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbaldehyde (**Int-4b**). To a solution of [5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-yl]methanol (**Int-4a**, 850 mg, 2.41 mmol) in DMSO (8 mL) was added IBX (514.6 mg, 3.62 mmol) at 25 °C under argon. The mixture was stirred at 25 °C for 16 h. The mixture was filtered to afford a crude solution. The crude product was purified by flash chromatography (eluted with CH_3CN in H_2O from 50% to 55%). 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbaldehyde (**Int-4b**, 570 mg, 1.59 mmol, 66.10 %yield) was obtained as a white solid. LCMS calcd for $C_{14}H_{15}N_4O_5S$ ($M+H$)⁺ m/z = 351.1, found: 351.1.

[00676] Step 3. Synthesis of 2-(1-methylbenzimidazol-2-yl)-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine (**Int-4c**). The solution of 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbaldehyde (**Int-4b**, 250 mg, 1 mmol) and N2-methylbenzene-1,2-diamine (244.34 mg, 2 mmol) in DMF (2 mL) was stirred at 100 °C for 18 h. The mixture was extracted with EtOAc and water, washed with brine and dried over Na_2SO_4 , concentrated and purified by silica gel chromatography (DCM: MeOH = 50:1) to get the 2-(1-methylbenzimidazol-2-yl)-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine (**Int-4c**, 320 mg, 0.707 mmol, 70.72%yield). LCMS calcd for $C_{21}H_{21}N_6O_4S$ ($M+H$)⁺ m/z = 453.49, found: 453.2.


[00677] Step 4. Synthesis of 2-(1-methylbenzimidazol-2-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 4**). A solution of Cs_2CO_3 (230.42 mg, 0.71 mmol), 2-(1-methylbenzimidazol-2-yl)-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine (**Int-4c**, 160 mg, 0.35 mmol) and 4-Methoxythiophenol (74.36 mg, 0.53 mmol) in CH_3CN (5 mL) was stirred at 25 °C for 3h. The mixture was filtered, and the solvent was removed. The residue was purified by silica gel chromatography (DCM: MeOH=11: 1) to give 2-(1-methylbenzimidazol-2-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 4**, 70 mg, 0.262 mmol, 74.05%yield). LCMS calcd for $\text{C}_{15}\text{H}_{18}\text{N}_5$ ($\text{M}+\text{H}$)⁺ m/z = 268.33, found: 268.3.

[00678] **Compound 82.** (7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-4-[2-(1-methylbenzimidazol-2-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]



[00679] Compound **82** was prepared similarly to that of **Ex. 21** using **intermediate 4**. LCMS calculated for $\text{C}_{40}\text{H}_{47}\text{FN}_9\text{O}$ ($\text{M}+\text{H}$)⁺ m/z = 688.38; found: 688.3. ¹H NMR (400 MHz, CD_3OD) δ 7.61 – 7.68 (m, 1H), 7.45 – 7.52 (m, 2H), 7.23 – 7.33 (m, 2H), 7.03 – 7.21 (m, 3H), 6.97 (s, 1H), 5.14 (d, J = 52.4 Hz, 1H), 4.80 – 4.95 (m, 2H), 4.50 – 4.67 (m, 2H), 3.92 – 4.18 (m, 7H), 3.76, 3.85 (d, J = 14.8 Hz, 2H), 2.85 – 3.15 (m, 6H), 2.67 – 2.78 (m, 2H), 2.29 – 2.42 (m, 1H), 1.92 – 2.22 (m, 9H), 1.69 – 1.90 (m, 5H).


[00680] **Compound 83.** (7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-4-[2-(4-fluoro-1-methyl-benzimidazol-2-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]



[00681] Compound **83** was prepared similarly to **Compound 82**. LCMS calculated for $\text{C}_{40}\text{H}_{46}\text{F}_2\text{N}_9\text{O}$ ($\text{M}+\text{H}$)⁺ m/z = 706.38; found: 706.3. ¹H NMR (400 MHz, CD_3OD) δ 7.49 – 7.51


(m, 1H), 7.35 – 7.37 (m, 1H), 7.25 – 7.30 (m, 1H), 7.17 – 7.21 (m, 1H), 7.11 – 7.15 (m, 1H), 7.07 – 7.09 (m, 1H), 7.04 (s, 1H), 6.98 – 7.03 (m, 1H), 5.20 (d, $J = 53.6$ Hz, 1H), 4.90 – 4.96 (m, 2H), 4.56 – 4.68 (m, 2H), 4.19 (s, 3H), 3.97 – 4.14 (m, 4H), 3.78, 3.86 (d, $J = 14.4$ Hz, 2H), 2.91 – 3.22 (m, 6H), 2.75 – 2.78 (m, 2H), 2.23 – 2.41 (m, 2H), 2.14 (s, 3H), 1.70 – 2.10 (m, 10H).

[00682] Compound 84. (7S)-4-[(1R,5S)-3,8-diazabicyclo[3.2.1]octan-3-yl]-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]




[00683] Compound **84** was prepared similarly to that of **Ex. 21**. LCMS calculated for $C_{31}H_{42}FN_6O$ ($M+H$)⁺ $m/z = 533.3$; found: 533.3. ¹H NMR (400 MHz, CD₃OD) δ 7.49 (d, $J = 7.6$ Hz, 1H), 7.06 – 7.23 (m, 3H), 5.17 – 5.36 (m, 1H), 4.21 (d, $J = 12.0$ Hz, 1H), 4.03, 4.15 (d, $J = 10.4$ Hz, 2H), 3.54 – 3.77 (m, 5H), 3.42 (d, $J = 12.0$ Hz, 1H), 3.13 – 3.28 (m, 3H), 2.87 – 3.09 (m, 4H), 2.71 – 2.80 (m, 2H), 2.11 – 2.33 (m, 2H), 2.09 (s, 3H), 1.67 – 2.07 (m, 12H).

[00684] Compound 85. (1R,5S)-3-[(7S)-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-3-azabicyclo[3.2.1]octan-8-amine



[00685] Compound **85** was prepared similarly to that of **Ex. 21** using **intermediate 3**. LCMS calculated for $C_{32}H_{44}FN_6O$ ($M+H$)⁺ $m/z = 547.4$; found: 547.5. ¹H NMR (400 MHz, DMSO) δ 7.46 (d, $J = 8.0$ Hz, 1H), 7.20 (t, $J = 7.2$ Hz, 1H), 7.13 (t, $J = 7.2$ Hz, 1H), 7.07 (d, $J = 7.6$ Hz, 1H), 5.24 (d, $J = 53.6$ Hz, 1H), 3.97 – 4.09 (m, 1H), 3.79, 3.93 (d, $J = 10.0$ Hz, 2H), 3.46 – 3.66 (m, 4H), 3.15 – 3.23 (m, 1H), 3.03 – 3.11 (m, 3H), 2.61 – 3.02 (m, 8H), 2.04 – 2.12 (m, 3H), 1.98 (s, 3H), 1.98-2.01 (m, 1H), 1.59 – 1.95 (m, 11H), 1.30 – 1.43 (m, 1H).

[00686] Example 26. Exemplary synthesis of [(1R,5S)-3-[(7S)-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-3-azabicyclo[3.2.1]octan-8-yl]cyanamide (**Compound 86**).




[00687] To a solution of (1R,5S)-3-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-3-azabicyclo[3.2.1]octan-8-amine (**Compound 85**, 5 mg, 0.01 mmol) in THF (0.3 mL) were added BrCN (0.93 mg, 0.01 mmol) and Et₃N (1.5 μ L, 0.01 mmol) at -20 °C. The reaction was stirred at

[00688] -20 °C for 2h. Then the mixture was filtered and purified by Prep-HPLC on a C18 column (5 μ M, 50 x 150 mm) with mobile phase: H₂O (0.1% NH₄HCO₃) / CH₃CN at flow rate of 35 mL/min to afford [(1R,5S)-3-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-3-azabicyclo[3.2.1]octan-8-yl]cyanamide (**Compound 86**, 2.22 mg, 0.0035 mmol, 39.58% yield) as a white solid. LCMS calculated for C₃₃H₄₃FN₇O (M+H)⁺ m/z = 572.4; found: 572.5.

[00689] ¹H NMR (400 MHz, DMSO) δ 7.46 (d, *J* = 7.6 Hz, 1H), 7.20 (t, *J* = 6.8 Hz, 1H), 7.13 (t, *J* = 7.2 Hz, 1H), 7.07 (d, *J* = 7.2 Hz, 1H), 6.87 (d, *J* = 2.8 Hz, 1H), 5.25 (d, *J* = 55.6 Hz, 1H), 3.99 – 4.08 (m, 1H), 3.91 – 3.97 (m, 1H), 3.79 – 3.85 (m, 1H), 3.54 – 3.64 (m, 4H), 3.20 – 3.24 (m, 1H), 3.04 – 3.10 (m, 1H), 2.64 – 3.02 (m, 8H), 1.42 – 2.12 (m, 19H).

[00690] **Intermediate 5.** Synthesis of 3-(3-chloro-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-2-yl)-5-methyl-1,2,4-oxadiazole



[00691] Step 1. Synthesis of 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-5a**). To a solution of 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylic acid (**Int-1d**, 870 mg, 2.37 mmol), DIEA (1.65 mL, 9.5 mmol) and HATU (1354.48 mg, 3.56 mmol) in DMF (2 mL) was added NH₄Cl (171.13 mg, 2.85 mmol). The reaction was stirred at 30 °C for 2h. The mixture was diluted with

DCM (30 x 2 mL), washed with water (40 mL) and brine (40 mL), dried over Na₂SO₄. The solvent was removed and the residue was purified by flash chromatography (eluted with CH₃CN in H₂O from 5.0% to 95%) to afford 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-5a**, 760 mg, 2.08 mmol, 87.59 %yield) as a yellowish solid. LCMS calcd for C₁₄H₁₆N₅O₅S (M+H)⁺ m/z = 366.0, found: 366.0.

[00692] Step 2. Synthesis of 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbonitrile (**Int-5b**). To a solution of 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-5a**, 1.5g, 4.11 mmol) and Pyridine (1 mL, 12.32 mmol) in THF (10mL) was added TFAA (2586.98 mg, 12.32 mmol) slowly at 25°C. The mixture was stirred at 25°C for 48 h. The mixture was quenched with H₂O (20 mL), extracted with EtOAc (20 x 3 mL), dried over Na₂SO₄, concentrated. The crude product was purified by silica gel chromatography (eluted with EtOAc in petroleum ether from 10% to 90%) to afford 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbonitrile (**Int-5b**, 0.8 g, 2.303 mmol, 56.10% yield) as a yellow solid. LCMS calcd for C₁₄H₁₄N₅O₄S (M+H)⁺ m/z = 348.07, found: 348.0.


[00693] Step 3. Synthesis of 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbonitrile (**Int-5c**). To a solution of 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbonitrile (500 mg, 1.44 mmol) in CH₃CN (5 mL) were added NCS (192.21 mg, 1.44 mmol). The mixture was stirred at 60 °C for 4h. The mixture was quenched with H₂O (10 mL), extracted with EtOAc (30 mL), dried over Na₂SO₄, concentrated. The crude product was purified by flash chromatography (eluted with EtOAc in petroleum ether from 10% to 90%). The product 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbonitrile (**Int-5c**, 400 mg, 1.05 mmol, 72.78% yield) was obtained as a white solid. LCMS calcd for C₁₄H₁₃ClN₅O₄S (M+H)⁺ m/z = 382.1, found: 382.0.

[00694] Step 4. Synthesis of 3-chloro-N-hydroxy-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-5d**). To a solution of 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbonitrile (**Int-5c**, 400 mg, 1.05mmol) and K₂CO₃ (433.75 mg, 3.14 mmol) in Ethanol (2 mL) was added NH₂OH·HCl (144.58 mg, 2.1 mmol). The reaction was stirred at 80 °C for 12 h. The mixture was cooled to rt, filtered, and washed with water (10ml) to give crude product 3-chloro-N-hydroxy-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-5d**, 183 mg, 0.441 mmol, 42.11% yield) as a white solid. LCMS calcd for C₁₄H₁₆ClN₆O₅S (M+H)⁺ m/z = 415.0, found: 415.2.

[00695] Step 5. Synthesis of 3-[3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-5-methyl-1,2,4-oxadiazole (**Int-5e**). To a solution of 3-chloro-N-hydroxy-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (183 mg, 0.44 mmol) in Acetic acid (2 mL) were added Acetic anhydride (**Int-5d**, 67.56 mg, 0.66 mmol). The reaction was stirred at 100 °C for 16h. The mixture was filtered to give crude product. The crude product was purified by flash chromatography (eluted with CH₃CN in H₂O from 5.0% to 43%). The product 3-[3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-5-methyl-1,2,4-oxadiazole (**Int-5e**, 163 mg, 0.371 mmol, 84.19% yield) was obtained as a white solid. LCMS calcd for C₁₆H₁₆ClN₆O₅S(M+H)⁺ m/z = 439.1, found: 439.2.

[00696] Step 6. Synthesis of 3-(3-chloro-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-2-yl)-5-methyl-1,2,4-oxadiazole (**Intermediate 5**). A solution of 3-[3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-5-methyl-1,2,4-oxadiazole (**Int-5e**, 150 mg, 0.34 mmol), 4-methoxybenzenethiol (0.13 mL, 1.03 mmol) and cesium carbonate (444.34 mg, 1.37 mmol) in CH₃CN (4 mL) was stirred at 20 °C for 1 h. The mixture was filtrated and purified by prep-HPLC on a C18 column with mobile phase: H₂O (0.1% NH₄HCO₃)/CH₃CN to afford 3-(3-chloro-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-2-yl)-5-methyl-1,2,4-oxadiazole (**Intermediate 5**, 80 mg, 0.315 mmol, 92.26% yield) as white solid. LCMS calcd for C₁₀H₁₃ClN₅O (M+H)⁺ m/z = 254.07, found: 254.0.


[00697] **Compound 87**. 3-[3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]-5-methyl-1,2,4-oxadiazole



[00698] Compound **87** was prepared similarly to that of **Ex. 21** using **intermediate 5**. LCMS calculated for C₃₅H₄₂ClFN₉O₂ (M+H)⁺ m/z = 674.31; found: 674.1. ¹H NMR (400 MHz, CD₃OD) δ 7.50 (d, *J* = 7.6 Hz, 1H), 7.00 – 7.27 (m, 3H), 5.21 (d, *J* = 55.2 Hz, 1H), 4.81 – 4.84 (m, 2H), 4.51 – 4.60 (m, 1H), 4.35 – 4.46 (m, 1H), 3.98 – 4.09 (m, 2H), 3.85 – 3.97 (m, 3H), 3.69 (d, *J* = 14.8 Hz, 1H), 2.90 – 3.21 (m, 6H), 2.71 – 2.79 (m, 2H), 2.66 (s, 3H), 2.35 – 2.47 (m, 1H), 1.95 – 2.27 (m, 9H), 1.72 – 1.93 (m, 5H).

[00699]

Intermediate 6.



[00700] Step 1. Synthesis of ethyl 3-(2-pyridyl)-1H-pyrazole-5-carboxylate (**Int-6a**). To a solution of 2-Ethynylpyridine (8000 mg, 77.58 mmol) in Toluene (80 mL) was added ethyl 2-diazoacetate (10622.19 mg, 93.1 mmol) and the mixture was stirred and refluxed at 110 °C for 16h. The reaction was concentrated and purified by flash column chromatography (silica gel, eluting with 27% EtOAc/PE) to afford ethyl 3-(2-pyridyl)-1H-pyrazole-5-carboxylate (9300 mg, 42.8 mmol, 55.19% yield) as a yellow solid. LCMS calculated for $C_{11}H_{12}N_3O_2$ ($M+H$)⁺ m/z =218.1; found:218.1.


[00701] Step 2. Synthesis of ethyl 2-[3-(tert-butoxycarbonylamino)propyl]-5-(2-pyridyl)pyrazole-3-carboxylate (**Int-6b**). To a solution of ethyl 3-(2-pyridyl)-1H-pyrazole-5-carboxylate (6600 mg, 30.38 mmol) in DMF (70 mL) were added tert-butyl N-(3-bromopropyl)carbamate (7958.53 mg, 33.42 mmol), KI (100.88 mg, 0.61 mmol) and cesium carbonate (19749.56 mg, 60.77 mmol). The reaction was stirred at rt overnight. The mixture was diluted with water, extracted with EtOAc, washed with brine, dried over Na_2SO_4 , concentrated. The product ethyl 2-[3-(tert-butoxycarbonylamino)propyl]-5-(2-pyridyl)pyrazole-3-carboxylate (**Int-6b**, 12300 mg) was carried over without further purification. LCMS calculated for $C_{19}H_{27}N_4O_4$ ($M+H$)⁺ m/z =375.5; found:375.1.

[00702] Step 3. Synthesis of 2-(2-pyridyl)-5,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-4-one (**Int-6c**). A mixture of ethyl 2-[3-(tert-butoxycarbonylamino)propyl]-5-(2-pyridyl)pyrazole-3-carboxylate (**Int-6b**, 12.3 g, 32.85 mmol) and HCl in Dioxane (120 mL, 480 mmol) was stirred at rt for 4h. The solvent was removed followed by the addition of Methanol (110 mL) and Et_3N (22.92 mL, 164.25 mmol). The reaction was stirred at 80 °C for 16h. The mixture was concentrated, and the residue was purified by flash column chromatography (silica gel, eluting with 7% MeOH/DCM) to afford 2-(2-pyridyl)-5,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-4-one (12000 mg, 52.6 mmol, 72,54% yield) as a yellow solid. LCMS calculated for $C_{12}H_{13}N_4O$ ($M+H$)⁺ m/z =229.3; found:229.1. ^1H NMR (400 MHz, $CDCl_3$) δ 8.66 (dd, J = 4.8, 0.8 Hz, 1H),

7.85 (d, $J = 8.0$ Hz, 1H), 7.74 (td, $J = 8.0, 2.0$ Hz, 1H), 7.41 (s, 1H), 7.22-7.25 (m, 1H), 7.12 (s, 1H), 4.61 (t, $J = 6.8$ Hz, 2H), 3.41 (s, 2H), 2.27 – 2.38 (m, 2H).


[00703] Step 4. To a solution of 2-(6-methoxy-2-pyridyl)-5,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-4-one (100 mg, 0.39 mmol) in THF (10 mL) was added LiAlH_4 (132.24 mg, 3.48 mmol) at 0 °C. The reaction was stirred at 0 °C to rt overnight. The reaction was quenched with $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$. The mixture was filtered and filtrate was concentrated. The residue was purified by flash column chromatography (silica gel, eluting with 7% MeOH/DCM) to afford 2-(6-methoxy-2-pyridyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (70 mg, 0.287 mmol, 74.01% yield) as a yellow oil. LCMS calculated for $\text{C}_{13}\text{H}_{17}\text{N}_4\text{O}$ ($\text{M}+\text{H}$) $^+$ $m/z = 245.1$; found: 245.1. ^1H NMR (400 MHz, DMSO) δ 7.68 (td, $J = 8.0, 7.6$ Hz, 1H), 7.44 (dd, $J = 7.6, 0.8$ Hz, 1H), 6.67 (dd, $J = 8.4, 0.8$ Hz, 1H), 6.62 (s, 1H), 4.32 – 4.39 (m, 2H), 3.89 (s, 3H), 3.82 (s, 2H), 2.99 – 3.07 (m, 2H), 1.68 – 1.78 (m, 2H), 1.40 – 1.46 (m, 1H).

[00704] **Compound 88.** (7S)-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-4-[2-(2-pyridyl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]




[00705] Compound **88** was prepared similarly to that of **Ex. 21** using **Intermediate 6**. LCMS calculated for $\text{C}_{37}\text{H}_{44}\text{FN}_8\text{O}$ ($\text{M}+\text{H}$) $^+$ $m/z = 635.4$; found: 635.5. ^1H NMR (400 MHz, CD_3OD) δ 8.52 (dd, $J = 4.8, 0.8$ Hz, 1H), 7.93 (d, $J = 8.0$ Hz, 1H), 7.82 (td, $J = 7.6, 1.6$ Hz, 1H), 7.49 (d, $J = 7.6$ Hz, 1H), 7.28 – 7.33 (m, 1H), 7.05 – 7.22 (m, 3H), 6.88 (s, 1H), 5.16 (d, $J = 52.4$ Hz, 1H), 4.47 – 4.60 (m, 2H), 3.94 – 4.15 (m, 4H), 3.73 – 3.89 (m, 2H), 3.07 – 3.22 (m, 3H), 2.87 – 3.03 (m, 3H), 2.71 – 2.80 (m, 2H), 2.19 – 2.40 (m, 3H), 2.12 (s, 3H), 1.69 – 2.11 (m, 11H).

[00706] **Compound 89.** (7S)-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-4-[2-(6-methoxy-2-pyridyl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]



[00707] Compound **89** was prepared similarly to **Compound 88**. LCMS calculated for $C_{38}H_{46}FN_8O_2$ $(M+H)^+$ m/z = 665.4; found: 665.5. ^1H NMR (400 MHz, CD_3OD) δ 7.66 (t, J = 7.6 Hz, 1H), 7.49 (d, J = 7.6 Hz, 1H), 7.45 (dd, J = 7.6, 0.4 Hz, 1H), 7.20 (td, J = 7.2, 1.2 Hz, 1H), 7.14 (td, J = 7.6, 1.2 Hz, 1H), 7.09 (dd, J = 7.6, 1.2 Hz, 1H), 6.87 (s, 1H), 6.68 (d, J = 8.4 Hz, 1H), 5.17 (d, J = 53.2 Hz, 1H), 4.43 – 4.60 (m, 2H), 3.99 – 4.15 (m, 4H), 3.96 (s, 3H), 3.78, 3.87 (d, J = 14.8 Hz, 2H), 3.14 – 3.21 (m, 2H), 3.08 – 3.12 (m, 1H), 2.89 – 3.03 (m, 3H), 2.73 – 2.79 (m, 2H), 2.13 (s, 3H), 1.64 – 2.50 (m, 14H).

[00708] Intermediate 7.



[00709] Step 1. Synthesis of ethyl 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-7a**). A solution of ethyl 5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-1c** 1000 mg, 2.54 mmol) in DMF (10 mL) was added NCS (372.42 mg, 2.79 mmol) at 25 °C. Then the mixture was stirred at 80 °C for 1 h. The solution was extracted with EtOAc (10 ml), the organic phase was concentrated. The residue was purified by silica gel chromatography (eluting with EtOAc in PE from 5% to 85%). The product ethyl 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-7a**, 914 mg, 2.131 mmol, 84.06% yield) was obtained as an oil. LCMS calculated for $C_{16}H_{18}ClN_4O_6S$ $(M+H)^+$ m/z = 429.06, found: 429.2.


[00710] Step 2. Synthesis of 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-7b**). A solution of ethyl 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxylate (**Int-7a**,

500 mg, 1.17 mmol) in Ethanol (5 mL) was added $\text{NH}_2\text{NH}_2\cdot\text{H}_2\text{O}$ (594.85 mg, 11.66 mmol) at 25 °C. The mixture was stirred at 85 °C for 2 h. The solution was extracted with EtOAc (5 ml), the organic phase was concentrated. The residue was purified by silica gel chromatography (eluting with EtOAc in PE from 5% to 95%). The product 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbohydrazide (**Int-7b**, 300 mg, 0.723 mmol, 62.03% yield) was obtained as a white solid. LCMS calculated for $\text{C}_{14}\text{H}_{16}\text{ClN}_6\text{O}_5\text{S}$ ($\text{M}+\text{H}$)⁺ m/z = 415.06, found: 415.0.

[00711] Step 3. Synthesis of 3-chloro-2-(4,5-dimethyl-1,2,4-triazol-3-yl)-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine (**Int-7c**). To a solution of 3-chloro-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carbohydrazide (**Int-7b**, 150 mg, 0.36 mmol) in 1,4-Dioxane (3 mL) was added 1,1,1-trimethoxyethane (65.17 mg, 0.54 mmol), Methylamine in EtOH (0.17 mL, 1.08 mmol) and a drop of AcOH (21.7 mg, 0.36 mmol) at rt. The mixture was stirred at 120 °C for 16 h in a sealed vial. The solution was extracted with EtOAc (2 ml), The water phase was concentrated. The residue was purified by silica gel chromatography (eluting with MeOH in DCM from 1% to 3%). The product 3-chloro-2-(4,5-dimethyl-1,2,4-triazol-3-yl)-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine (95 mg, 0.210 mmol, 58.14% yield) was obtained as a white solid. LCMS calculated for $\text{C}_{17}\text{H}_{19}\text{ClN}_7\text{O}_4\text{S}$ ($\text{M}+\text{H}$)⁺ m/z = 452.09, found: 452.2.


[00712] Step 4. Synthesis of 3-chloro-2-(4,5-dimethyl-1,2,4-triazol-3-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 7**). To a solution of 3-chloro-2-(4,5-dimethyl-1,2,4-triazol-3-yl)-5-(2-nitrophenyl)sulfonyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine (**Int-7c**, 75 mg, 0.17 mmol) in CH_3CN (0.5 mL) was added 4-Methoxythiophenol (44.21 mg, 0.32 mmol) and CS_2CO_3 (137.07 mg, 0.42 mmol) at rt. The mixture was stirred at 25 °C for 2 h. The solution was concentrated. The residue was purified by silica gel on chromatography (eluting with MeOH in DCM from 3 % to 11%). The 3-chloro-2-(4,5-dimethyl-1,2,4-triazol-3-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 7**, 40 mg, 0.15 mmol, 71.33% yield) was obtained as a white solid. LCMS calculated for $\text{C}_{11}\text{H}_{16}\text{ClN}_6$ ($\text{M}+\text{H}$)⁺ m/z = 267.11, found: 267.1.

[00713] **Compound 90.** (7S)-4-[3-chloro-2-(4,5-dimethyl-1,2,4-triazol-3-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl]-2-[[2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]




[00714] Compound **90** was prepared similarly to that of **Ex. 21** using **Intermediate 7**. LCMS calcd for $C_{36}H_{45}ClFN_{10}O$ ($M+H$)⁺ m/z = 687.34; found: 687.4. ¹H NMR (400 MHz, CD₃OD) δ 7.50 (d, J = 8.0 Hz, 1H), 7.04 – 7.24 (m, 3H), 5.23 (d, J = 52.0 Hz, 1H), 4.79 – 4.96 (m, 2H), 4.42 – 4.64 (m, 2H), 3.85 – 4.15 (m, 5H), 3.77 (s, 3H), 3.70 (d, J = 14.8 Hz, 1H), 2.90 – 3.22 (m, 6H), 2.71 – 2.80 (m, 2H), 2.37 – 2.53 (m, 4H), 1.72 – 2.29 (m, 14H).

[00715] **Compound 91.** (7S)-4-(6,8-dihydro-5H-imidazo[1,5-a]pyrazin-7-yl)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]




[00716] Compound **91** was prepared similarly to that of **Ex. 21**. LCMS calculated for $C_{31}H_{39}FN_7O$ ($M+H$)⁺ m/z = 544.31; found: 544.4. ¹H NMR (400 MHz, CD₃OD) δ 7.63 (s, 1H), 7.51 (d, J = 7.6 Hz, 1H), 7.06 – 7.24 (m, 3H), 6.84 (s, 1H), 5.28 (d, J = 53.6 Hz, 1H), 4.77 – 4.83 (m, 2H), 4.30 – 4.40 (m, 1H), 4.05 – 4.25 (m, 4H), 3.68 – 3.78 (m, 3H), 3.15 – 3.27 (m, 3H), 2.91 – 3.09 (m, 3H), 2.71 – 2.81 (m, 2H), 2.15 – 2.31 (m, 2H), 2.05 – 2.14 (m, 4H), 1.91 – 2.04 (m, 4H), 1.74 – 1.90 (m, 3H).

[00717] **Example 27.** Exemplary synthesis of 6-[5-[(7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]pyridin-2-ol (**Compound 92**).



[00718] To a solution of (7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-4-[2-(6-methoxy-2-pyridyl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**Compound 89**, 8 mg, 0.01 mmol) in THF (5 mL) was added HBr (0.5 mL, 0.01 mmol) and the reaction was stirred at 60 °C for 24h. The pH was adjusted to 8 by saturated NaHCO₃ aqueous solution. Then this mixture was concentrated and purified by Prep-HPLC on a C18 column (5 uM, 50 x 150 mm) with mobile phase: H₂O (0.1% NH₄HCO₃) / CH₃CN at flow rate : 35 mL/min to afford 6-[5-[(7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]pyridin-2-ol (**Compound 92**, 3.25 mg, 0.005 mmol, 39.05% yield) as a white solid. LCMS calculated for C₃₇H₄₄FN₈O₂ (M+H)⁺ m/z = 651.4; found: 651.4. ¹H NMR (400 MHz, CD₃OD) δ 7.60 (dd, *J* = 9.2, 7.2 Hz, 1H), 7.49 (dd, *J* = 7.6, 0.8 Hz, 1H), 7.17 – 7.22 (m, 1H), 7.14 (td, *J* = 7.6, 1.6 Hz, 1H), 7.09 (dd, *J* = 7.6, 0.8 Hz, 1H), 6.86 (s, 1H), 6.77 (dd, *J* = 7.2, 1.2 Hz, 1H), 6.46 (dd, *J* = 9.2, 0.8 Hz, 1H), 5.23 (d, *J* = 53.2 Hz, 1H), 4.46 – 4.62 (m, 3H), 3.95 – 4.10 (m, 4H), 3.74, 3.83 (d, *J* = 14.8 Hz, 2H), 3.10 – 3.23 (m, 3H), 2.88 – 3.04 (m, 3H), 2.72 – 2.80 (m, 2H), 2.14 – 2.40 (m, 2H), 2.12 (s, 3H), 1.67 – 2.11 (m, 11H).

[00719] Intermediate 8.




[00720] Step 1. Synthesis of 3-chloro-2-(2-pyridyl)-5,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-4-one (**Int-8a**). To a solution of 2-(2-pyridyl)-5,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-4-one (**Int-6c**, 2500 mg, 10.95 mmol) in DMF (23 mL) was added NCS (1755.05 mg, 13.14 mmol) at 50 °C, then the mixture was stirred at 50 °C for 16h. Then the mixture was concentrate and purified by Prep-HPLC on a C18 column (5 uM, 50 x 150 mm) with mobile phase: H₂O (0.1% HCOOH) / CH₃CN at flow rate : 65mL/min to afford 3-chloro-2-(2-pyridyl)-5,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-4-one (**Int-8a**, 2690 mg, 10.2 mmol, 93.49 %yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 8.79 (d, *J* = 4.4 Hz, 1H), 8.05 (d, *J* = 8.0 Hz, 1H), 7.84 (td, *J* = 8.0, 1.6 Hz, 1H), 7.30 – 7.38 (m, 1H), 6.46 (s, 1H), 4.60 (t, *J* = 7.2 Hz, 2H), 3.34 – 3.44 (m, 2H), 2.24 – 2.37 (m, 2H). LCMS calculated for C₁₂H₁₂ClN₄O (M+H)⁺ m/z = 263.7; found: 263.2/265.2

[00721] Step 2. 3-chloro-2-(2-pyridyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 8**). To a solution of 3-chloro-2-(2-pyridyl)-5,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-4-one (**Int-8a**, 1000 mg, 3.81 mmol) in THF (100 mL)


was added NaBH₄ (845.1 mg, 22.84 mmol) at 0 °C. The mixture was stirred at rt for 1h, followed by the addition of Boron trifluoride diethyl etherate (4322.05 mg, 30.45 mmol). The reaction was allowed to rt and stirred overnight. Methanol was added to quench the reaction at 0 °C and then refluxed at 45 °C for 16h. The mixture was concentrated and purified by flash column chromatography (silica gel, eluting with MeOH/DCM=10/1) to afford 3-chloro-2-(2-pyridyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 8**, 710 mg, 2.85 mmol, 74.99 %yield) as a colorless oil. LCMS calculated for C₁₂H₁₄ClN₄(M+H)⁺ m/z =249.7; found:249.2/251.2.

[00722] Compound 93. (7S)-4-[3-chloro-2-(2-pyridyl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl]-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]



[00723] Compound **93** was prepared similarly to **Compound 88** using **Intermediate 8**. LCMS calculated for C₃₇H₄₃ClFN₈O (M+H)⁺ m/z = 670.3; found:669.4, 671.4. ¹H NMR (400 MHz, CD₃OD) δ 8.61 (d, *J* = 4.8 Hz, 1H), 7.85 – 7.95 (m, 2H), 7.50 (d, *J* = 8.0 Hz, 1H), 7.35 – 7.42 (m, 1H), 7.06 – 7.23 (m, 3H), 5.18 (d, *J* = 55.2 Hz, 1H), 4.77 (d, *J* = 28.4 Hz, 2H), 4.50 – 4.61 (m, 1H), 4.35 – 4.47 (m, 1H), 4.01 – 4.12 (m, 2H), 3.87 – 4.00 (m, 3H), 3.71 (d, *J* = 15.2 Hz, 1H), 2.89 – 3.25 (m, 6H), 2.76 (dd, *J* = 8.0, 2.8 Hz, 2H), 2.37 – 2.49 (m, 1H), 2.14 – 2.28 (m, 2H), 2.11 (s, 3H), 1.70 – 2.08 (m, 9H).

[00724] Intermediate 9.



[00725] Step 1. Synthesis of methyl 3-[(2R)-2-(tert-butoxycarbonylamino)propyl]imidazole-4-carboxylate (**Int-9a**) To a solution of tert-butyl N-[(1R)-2-hydroxy-1-methyl-ethyl]carbamate (1.75 g, 9.99 mmol), methyl 1H-imidazole-4-carboxylate (1.51g, 11.98 mmol) and PPh₃ (3705.38 mg, 16.98 mmol) in THF (100 mL) was added

DIAD (3.34 mL, 16.98 mmol) dropwise at -50°C under N₂. The reaction mixture was allowed to warm to rt and stirred for 16 h. The reaction mixture was concentrated under reduced pressure and then adjusted the pH to 1 by 1M HCl, extracted with EtOAc (3 x 50 mL). The aqueous layer was neutralized to pH 7-8 with sat. aq NaHCO₃, and was extracted with EtOAc (3 x 70 mL). The combined organic phases were dried (Na₂SO₄), concentrated under reduced pressure and the crude residue was purified by silica gel column chromatography (DCM:EtOAc, 1:1) to afford methyl 3-[(2R)-2-(tert-butoxycarbonylamino)propyl]imidazole-4-carboxylate (**Int-9a**, 1.20 g, 4.24 mmol, 42.41 %yield) as a white solid. LCMS calculated for C₁₃H₂₂N₃O₄ (M+H)⁺ m/z = 284.32; found: 284.2.


[00726] Step 2. Synthesis of methyl 3-[(2R)-2-aminopropyl]imidazole-4-carboxylate (**Int-9b**)

[00727] A solution of methyl 3-[(2R)-2-(tert-butoxycarbonylamino)propyl]imidazole-4-carboxylate (**Int-9a**, 1.2g, 4.24mmol) and HCl/Dioxane (6 mL, 24 mmol) in Methanol (4 mL) was stirred at 25 °C for 1 h. The reaction mixture was concentrated under reduced pressure to give the crude methyl 3-[(2R)-2-aminopropyl]imidazole-4-carboxylate (**Int-9b**, 0.80 g) as a white solid.

[00728] Step 3. Synthesis of (6R)-6-methyl-6,7-dihydro-5H-imidazo[1,5-a]pyrazin-8-one (**Int-9c**). A solution of methyl 3-[(2R)-2-aminopropyl]imidazole-4-carboxylate (**Int-9b**, 0.8 g, 4.37 mmol) and TEA (6 mL, 43.25 mmol) in Methanol (30 mL) was stirred at 50 °C overnight. The reaction mixture was concentrated under reduced pressure. The crude residue was purified by column (DCM:MeOH) to afford (6R)-6-methyl-6,7-dihydro-5H-imidazo[1,5-a]pyrazin-8-one (**Int-9c**, 1.20 g, 7.94 mmol, 95.19 %yield) as a white solid.


[00729] Step 4. Synthesis of (**Intermediate 9**). To a solution of (6R)-6-methyl-6,7-dihydro-5H-imidazo[1,5-a]pyrazin-8-one (**Int-9c**, 600 mg, 3.97 mmol) in THF (30mL) was added LiAlH₄ (451.88 mg, 11.91 mmol). The mixture reaction was stirred at 80 °C overnight. The reaction was quenched with Na₂SO₄·10H₂O and dried over MgSO₄. the mixture was filtered, and the filter cake was washed with THF twice. The filtrate was concentrated under reduced pressure to give not the title product(6R)-6-methyl-5,6,7,8-tetrahydroimidazo[1,5-a]pyrazine (**Intermediate 9**, 150 mg, 1.09 mmol, 27.55 %yield) as a white solid. LCMS calculated for C₇H₁₂N₃(M+H)⁺ m/z = 138.10; found: 138.2. ¹H NMR (400 MHz, CD₃OD) δ 7.53 (s, 1H), 6.70 (s, 1H), 4.21 – 4.15 (m, 1H), 4.13 (d, J = 16.0 Hz, 1H), 3.93 (d, J = 15.5 Hz, 1H), 3.58 – 3.50 (m, 1H), 3.18 – 3.08 (m, 1H), 1.25 (d, J = 6.4 Hz, 3H).

[00730] **Compound 94.** (7S)-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-4-[(6R)-6-methyl-6,8-dihydro-5H-imidazo[1,5-a]pyrazin-7-yl]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]




[00731] Compound **94** was prepared similarly to that of **Ex. 21** using **Intermediate 9**. LCMS calculated for $C_{32}H_{41}FN_7O$ ($M+H$)⁺ m/z = 558.3; found: 558.4. ¹H NMR (400 MHz, CD₃OD) δ 7.66 (s, 1H), 7.51 (d, J = 6.8 Hz, 1H), 7.19 – 7.24 (m, 1H), 7.15 (td, J = 7.6, 1.6 Hz, 1H), 7.10 (dd, J = 7.6, 0.8 Hz, 1H), 6.88 (s, 1H), 5.40 (d, J = 52.8 Hz, 1H), 4.76 – 4.82 (m, 3H), 4.42 – 4.49 (m, 1H), 4.15 – 4.36 (m, 3H), 3.89 (d, J = 15.2 Hz, 1H), 3.61 – 3.75 (m, 2H), 3.50 – 3.59 (m, 1H), 3.38 – 3.47 (m, 1H), 3.14 – 3.23 (m, 1H), 2.94 – 3.12 (m, 2H), 2.74 – 2.80 (m, 2H), 2.15 – 2.51 (m, 4H), 2.13 (s, 3H), 1.93 – 2.11 (m, 4H), 1.74 – 1.88 (m, 2H), 1.27 (d, J = 6.8 Hz, 3H).

[00732] **Compound 95.** 3-chloro-N,N-dimethyl-5-[(7S)-6-methyl-2-[[1-(morpholinomethyl)cyclopropyl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00733] Compound **95** was prepared similarly to that of **Ex. 21**. LCMS calculated for $C_{36}H_{48}ClN_8O_3$ ($M+H$)⁺ m/z = 675.3; found: 675.1/677.1. ¹H NMR (400 MHz, CD₃OD) δ 7.50 (d, J = 8.0 Hz, 1H), 7.19 (t, J = 7.2 Hz, 1H), 7.13 (td, J = 7.6, 1.2 Hz, 1H), 7.08 (d, J = 7.6 Hz, 1H), 4.77 (d, J = 16.8 Hz, 1H), 4.35 – 4.61 (m, 2H), 4.22 (d, J = 11.2 Hz, 1H), 4.02 – 4.15 (m, 2H), 3.81 – 3.98 (m, 2H), 3.68 (d, J = 14.4 Hz, 1H), 3.61 (t, J = 4.4 Hz, 4H), 3.09 (d, J = 10.0 Hz, 6H), 2.97 (dd, J = 38.4, 18.4 Hz, 2H), 2.70 – 2.79 (m, 2H), 2.34 – 2.52 (m, 6H), 2.30 (d, J = 12.8 Hz, 1H), 2.13 – 2.24 (m, 1H), 2.11 (s, 3H), 1.92 – 2.06 (m, 3H), 1.71 – 1.86 (m, 2H), 0.54 – 0.66 (m, 2H), 0.37 – 0.48 (m, 2H).

[00734] **Compound 96.** 3-chloro-5-[(7S)-2-[[1-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-7'-hydroxy-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide




[00735] Compound **96** was prepared similarly to that of **Ex. 21**. LCMS calculated for $C_{34}H_{43}ClFN_8O_3$ ($M+H$)⁺ m/z = 665.3; found: 665.8. ¹H NMR (400 MHz, CD₃OD) δ 7.11 (t, J = 8.0 Hz, 1H), 6.73 (d, J = 7.6 Hz, 1H), 6.60 (d, J = 8.0 Hz, 1H), 5.28 (d, J = 54.4 Hz, 1H), 4.80 (d, J = 16.8 Hz, 2H), 4.33 – 4.53 (m, 2H), 3.95 – 4.13 (m, 3H), 3.71 – 3.91 (m, 3H), 3.05 – 3.29 (m, 10H), 2.90 – 3.04 (m, 3H), 2.75 (d, J = 18.4 Hz, 1H), 2.05 – 2.45 (m, 9H), 1.82 – 2.01 (m, 3H), 1.72 – 1.81 (m, 1H).

[00736] **Compound 97.** 3-chloro-5-[(7S)-2-[[1-[(dimethylamino)methyl]-2,2-difluorocyclopropyl]methoxy]-6-methyl-spiro[5,8]-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00737] Compound **97** was prepared similarly to that of **Ex. 21**. LCMS calculated for $C_{34}H_{44}ClF_2N_8O_2$ ($M+H$)⁺ m/z = 669.3; found: 669.0/671.1. ¹H NMR (400 MHz, CD₃OD) δ 7.50 (d, J = 7.6 Hz, 1H), 7.20 (t, J = 7.2 Hz, 1H), 7.13 (td, J = 7.6, 1.2 Hz, 1H), 7.08 (d, J = 7.2 Hz, 1H), 4.87 – 4.93 (m, 1H), 4.75 – 4.82 (m, 1H), 4.47 – 4.54 (m, 1H), 4.37 – 4.46 (m, 2H), 4.23 – 4.29 (m, 1H), 4.03 – 4.13 (m, 1H), 3.84 – 3.98 (m, 2H), 3.68 (d, J = 14.8 Hz, 1H), 3.06 – 3.11 (m, 6H), 2.89 – 3.06 (m, 2H), 2.63 – 2.79 (m, 3H), 2.34 – 2.51 (m, 2H), 2.21 – 2.26 (m, 6H), 2.12 – 2.20 (m, 1H), 2.10 (s, 3H), 1.93 – 2.04 (m, 2H), 1.72 – 1.87 (m, 2H), 1.51 – 1.66 (m, 1H), 1.30 – 1.40 (m, 1H).

[00738] **Intermediate 10.** Synthesis of 2-(3-chloro-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-2-yl)acetonitrile




[00739] Step 1. Synthesis of tert-butyl 3-chloro-2-(methylsulfonyloxymethyl)-4,5,7,8-tetrahydropyrazolo[1,5-d][1,4]diazepine-6-carboxylate (**Int-10a**)

[00740] To a solution of tert-butyl 3-chloro-2-(hydroxymethyl)-4,5,7,8-tetrahydropyrazolo[1,5-d][1,4]diazepine-6-carboxylate (500 mg, 1.66 mmol) in DCM (5 mL) were added Et₃N (0.35 mL, 2.49 mmol) and Methanesulfonyl Chloride (0.15 mL, 1.99 mmol) at 0 °C. The reaction was stirred at 0 °C for 2h. Then the mixture was extracted with EtOAc/H₂O. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by flash column chromatography (silica gel, eluting with 5% MeOH/DCM) to afford tert-butyl 3-chloro-2-(methylsulfonyloxymethyl)-4,5,7,8-tetrahydropyrazolo[1,5-d][1,4]diazepine-6-carboxylate (**Int-10a**, 490 mg, 0.645 mmol, 38.93% yield) as an oil. LCMS calculated for C₁₄H₂₃ClN₃O₅S (M+H)⁺ m/z =380.1; found:380.2/382.2.

[00741] Step 2. Synthesis of tert-butyl 3-chloro-2-(cyanomethyl)-4,5,7,8-tetrahydropyrazolo[1,5-d][1,4]diazepine-6-carboxylate (**Int-10b**). To a solution of tert-butyl 3-chloro-2-(methylsulfonyloxymethyl)-4,5,7,8-tetrahydropyrazolo[1,5-d][1,4]diazepine-6-carboxylate (**Int-10a**, 490 mg, 1.29 mmol) in DMF was added NaCN (316.1 mg, 6.45 mmol) and the reaction was stirred at rt for 2h. Then the mixture was extracted with EtOAc/H₂O. The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by Prep-HPLC on a C18 column (5 uM, 50 x 150 mm) with mobile phase: H₂O (0.1% NH₄HCO₃) / CH₃CN at flow rate: 35 mL/min to afford tert-butyl 3-chloro-2-(cyanomethyl)-4,5,7,8-tetrahydropyrazolo[1,5-d][1,4]diazepine-6-carboxylate (**Int-10b**, 200 mg, 0.644 mmol, 49.89% yield) as a yellow oil. LCMS calculated for C₁₄H₂₀ClN₄O₂ (M+H)⁺ m/z =311.1; found:311.2. ¹H NMR (400 MHz, DMSO) δ 4.49 (s, 2H), 4.35 – 4.42 (m, 2H), 3.96 (s, 2H), 3.61 – 3.70 (m, 2H), 1.72 – 1.85 (m, 2H), 1.36 (s, 9H).


[00742] Step 3. Synthesis of 2-(3-chloro-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-2-yl)acetonitrile (**Intermediate 10**). A mixture of tert-butyl 3-chloro-2-(cyanomethyl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-9b**, 100 mg, 0.32 mmol) and HCl in Dioxane (2 mL, 8 mmol) was stirred at 0 °C to rt for 2h. Then the mixture was concentrated to afford a crude product for the next step. LCMS calculated for C₉H₁₂ClN₄ (M+H)⁺ m/z =211.1; found:211.1/213.1.

[00743] **Compound 98.** 2-[3-chloro-5-[(7S)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-2-yl]acetonitrile



[00744] Compound **98** was prepared similarly to that of **Ex. 21** using **Intermediate 10**. LCMS calculated for $C_{34}H_{41}ClFN_8O$ ($M+H$)⁺ m/z = 631.3; found: 631.1/633.1. ¹H NMR (400 MHz, CD₃OD) δ 7.49 (d, J = 7.2 Hz, 1H), 7.17 – 7.23 (m, 1H), 7.14 (td, J = 7.6, 1.2 Hz, 1H), 7.08 (d, J = 7.6 Hz, 1H), 5.26 (d, J = 54.8 Hz, 1H), 4.74, 4.80 (d, J = 16.4 Hz, 2H), 4.41 – 4.50 (m, 1H), 4.29 – 4.38 (m, 1H), 3.84 – 4.09 (m, 5H), 3.82 (s, 2H), 3.66 (d, J = 14.8 Hz, 1H), 3.09 – 3.28 (m, 3H), 2.89 – 3.06 (m, 3H), 2.71 – 2.81 (m, 2H), 2.12 – 2.42 (m, 4H), 2.10 (s, 3H), 1.93 – 2.08 (m, 5H), 1.74 – 1.92 (m, 3H).

[00745] **Intermediate 11.** Synthesis of 3-chloro-2-(1-methyltetrazol-5-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4] diazepine



[00746] Step 1. Synthesis of tert-butyl 2-(1H-tetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4] diazepine-5-carboxylate (**Int-11a**). To a solution of tert-butyl 2-cyano-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4] diazepine-5-carboxylate (400 mg, 1.52 mmol), sodium azide (495.67 mg, 7.62 mmol) and Triethylamine Hydrochloride (472.28 mg, 3.43 mmol) in Toluene (10 mL). The mixture was stirred at 120 °C for 36 h. The crude product was purified by silica gel chromatography (eluted with MeOH in DCM from 1% to 10%). The crude tert-butyl 2-(1H-tetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4] diazepine-5-carboxylate (**Int-11a**, 275 mg, 0.901 mmol, 59.06% yield) was obtained as yellow solid. LCMS calcd for $C_{13}H_{20}N_7O_2$ ($M+H$)⁺ m/z = 306.1, found: 306.1.


[00747] Step 2. Synthesis of tert-butyl 2-(1-methyltetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-11b**). To a solution of tert-butyl 2-(1H-tetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-11a**, 100 mg, 0.33 mmol) and NaH (15.72 mg, 0.39 mmol) in DMF (0.5 mL) at 0 °C, followed by the addition of MeI (0.02 mL, 0.39 mmol). The reaction was stirred at 25 °C for 16 h. The crude product was purified by prep-HPLC. The product tert-butyl 2-(2-methyltetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (28 mg, 0.0877 mmol, 26.77 % yield) and tert-butyl 2-(1-methyltetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-

carboxylate (**Int-11b**, 60 mg, 0.188 mmol, 57.37 %yield) was obtained as white solid. LCMS calcd for $C_{14}H_{22}N_7O_2$ ($M+H$)⁺ m/z = 320.1, found:320.1.

[00748] Step 3. Synthesis of tert-butyl 3-chloro-2-(1-methyltetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-11c**). To a solution of tert-butyl 2-(1-methyltetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-11b**, 190 mg, 0.44 mmol) in DMF (4 mL) was added NCS (70.24 mg, 0.53 mmol) at 0 °C, then the mixture was stirred at 50 °C for 16h. Then the mixture was concentrated and purified by Prep-HPLC on a C18 column (5 μ M, 50 x 150 mm) with mobile phase: H_2O (0.1% NH_4HCO_3)/MeOH at flow rate : 50 mL/min to afford tert-butyl 3-chloro-2-(1-methyltetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-11c**, 162 mg, 0.458 mmol, 77.09% yield) as a white solid. LCMS calculated for $C_{14}H_{21}ClN_7O_2(M+H)^+$ m/z =354.1; found:354.2/356.1


[00749] Step 4. Synthesis of 3-chloro-2-(1-methyltetrazol-5-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine (**Intermediate 11**). To a solution of tert-butyl 3-chloro-2-(1-methyltetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-5-carboxylate (**Int-11c**, 50 mg, 0.14 mmol) in DCM (1.5 mL) was added HCl in dioxane (1.41 mL, 5.65 mmol) at 0 °C. The reaction was stirred at rt for 1h. Then the mixture was concentrated to afford 70 mg crude of **Intermediate 11**. LCMS calculated for $C_9H_{13}ClN_7$ ($M+H$)⁺ m/z =254.7; found:254.3/256.3.

[00750] **Compound 99**. (7S)-4-[3-chloro-2-(1-methyltetrazol-5-yl)-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepin-5-yl]-2-[[2-(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]-6-methyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]



[00751] **Compound 99** was prepared similarly to that of **Ex. 21** using **Intermediate 11**. LCMS calculated for $C_{34}H_{42}ClFN_{11}O$ ($M+H$)⁺ m/z = 674.3; found: 674.5. ¹H NMR (400 MHz, CD_3OD) δ 7.50 (d, J = 7.6 Hz, 1H), 7.20 (td, J = 7.2, 1.6 Hz, 1H), 7.14 (td, J = 7.6, 1.6 Hz, 1H), 7.09 (d, J = 6.8 Hz, 1H), 5.23 (d, J = 53.2 Hz, 1H), 4.60 – 4.66 (m, 2H), 4.41 – 4.54 (m, 1H), 4.33 (s, 3H), 4.02 – 4.11 (m, 2H), 3.89 – 3.98 (m, 3H), 3.71 (d, J = 14.8 Hz, 1H), 3.10 – 3.20 (m, 2H), 3.00 – 3.10 (m, 2H), 2.90 – 2.99 (m, 2H), 2.76 (d, J = 5.2 Hz, 2H), 2.36 – 2.51 (m, 1H), 2.14 – 2.31 (m, 3H), 2.11 (s, 3H), 1.77 – 2.06 (m, 9H).

[00752] Example 28. Exemplary synthesis of 3-chloro-5-[(7S)-6-ethyl-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 100**).



[00753] Step 1. Synthesis of (7S)-6-ethylspiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**100b**). To a solution of (7S)-spiro[6,8-dihydro-5H-pyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**100a** prepared similarly to that of **64h**, 400 mg, 1.41 mmol) in Ethanol (6.7 mL) was added acetaldehyde (0.8 mL, 14.12 mmol) and NaBH_3CN (266.15 mg, 4.24 mmol). The mixture was stirred at 30 °C for 2 days. NaBH_3CN (266.15 mg, 4.24 mmol) and acetaldehyde (0.8 mL, 14.12 mmol) were added, and the reaction was continued to stir at 30 °C for 6 h. The reaction mixture was quenched with water and filtered, dry over in vacuum to afford (7S)-6-ethylspiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**100b**, 430 mg, 0.884 mmol, 62.60 %yield) as yellow solid. LCMS calculated for $\text{C}_{18}\text{H}_{22}\text{N}_3\text{O}_2$ ($\text{M}+\text{H}$)⁺ m/z = 312.17; found: 312.0.

[00754] Step 2. Synthesis of (7S)-2,4-dichloro-6-ethyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**100c**). A solution of (7S)-6-ethylspiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-2,4-diol (**100b**, 430 mg, 0.88 mmol) in POCl_3 (4065.43 mg, 26.51 mmol) was stirred at 100 °C for 5 h. The reaction mixture was concentrated, diluted with DCM and DIEA, extracted with DCM, dried over Na_2SO_4 , concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 10% EtOAc/PE) to afford (7S)-2,4-dichloro-6-ethyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**100c**, 120 mg, 0.345 mmol, 38.99% yield) as a yellow solid. LCMS calculated for $\text{C}_{18}\text{H}_{20}\text{Cl}_2\text{N}_3$ ($\text{M}+\text{H}$)⁺ m/z = 348.10, 350.10; found: 348.1, 350.1.


[00755] Step 3. Synthesis of 3-chloro-5-[(7S)-2-chloro-6-ethyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**100d**). A solution of (7S)-2,4-dichloro-6-ethyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin] (**100c**, 120mg, 0.34mmol), 3-

chloro-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (83.63 mg, 0.34 mmol) and DIEA (133.59 mg, 1.03 mmol) in DMSO (1.2 mL) was stirred at 30 °C for 16 h. The reaction mixture was filtered, the filtrate was purified by prep-HPLC (0.1% NH₄HCO₃), to afford 3-chloro-5-[(7S)-2-chloro-6-ethyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**100d**, 110 mg, 0.198 mmol, 57.57 %yield) as a white solid. LCMS calculated for C₂₈H₃₄Cl₂N₇O (M+H)⁺ m/z = 554.22, 556.22; found: 554.0, 556.0. ¹H NMR (400 MHz, DMSO) δ = 7.47 (d, J=7.5 Hz, 1H), 7.19 (t, J=6.9 Hz, 1H), 7.13 (t, J=6.8, 1H), 7.07 (d, J=6.9, 1H), 4.85 – 4.71 (m, 2H), 4.51 – 4.38 (m, 2H), 4.02 – 3.86 (m, 2H), 3.74 (dd, J=45.1, 15.6 Hz, 2H), 3.02 – 2.91 (m, 7H), 2.82 (d, J=18.3 Hz, 1H), 2.82 (d, J=18.3 Hz, 1H), 2.76 – 2.62 (m, 2H), 2.37 – 1.99 (m, 4H), 1.94 – 1.64 (m, 4H), 0.84 (t, J=7.0 Hz, 3H).

[00756] Step 4. Synthesis of 3-chloro-5-[(7S)-6-ethyl-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 100**)

[00757] The mixture of 3-chloro-5-[(7S)-2-chloro-6-ethyl-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**100d**, 30 mg, 0.05 mmol), [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (43.07 mg, 0.27 mmol) and Sodium Tert-Butoxide (7.8 mg, 0.08 mmol) in DMSO (1 mL) was stirred at 20°C for 3h until the reaction mixture is clear. The reaction mixture was purified by prep-HPLC (0.1% NH₄HCO₃) to afford 3-chloro-5-[(7S)-6-ethyl-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 100**, 16 mg, 0.0237 mmol, 43.88 %yield) as a white solid. LCMS calculated for C₃₆H₄₆ClFN₈O₂ (M+H)⁺ m/z = 677.35; found: 677.0. ¹H NMR (400 MHz, CD₃OD) δ = 7.52 (d, J=7.7 Hz, 1H), 7.17 (dd, J=10.7, 4.3 Hz, 1H), 7.12 (td, J=7.3, 1.3 Hz, 1H), 7.07 (d, J=6.5 Hz, 1H), 5.26 (d, J=53.8 Hz, 1H), 4.93 – 4.86 (m, 1H), 4.76 (dd, J=16.7, 6.5 Hz, 1H), 4.56 – 4.34 (m, 2H), 4.16 – 3.72 (m, 6H), 3.16 (m, 3H), 3.10 (d, J=2.1 Hz, 3H), 3.08 (s, 3H), 3.06 – 2.86 (m, 3H), 2.75 (d, J=7.0 Hz, 2H), 2.53 – 1.71 (m, 14H), 0.95 (t, J=7.0 Hz, 3H).

[00758] **Example 29.** Exemplary synthesis of 3-chloro-5-[(7S)-6-ethyl-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 101**).



[00759] Step 1. Synthesis of (7S)-2,4-dichloro-7'-fluoro-spiro[6,8-dihydro-5H-pyrido[4,3-d]pyrimidine-7,1'-indane] (**101b**)

[00760] The mixture of (7S)-7'-fluorospiro[6,8-dihydro-5H-pyrido[4,3-d]pyrimidine-7,1'-indane]-2,4-diol (**100a**, 1.2 g, 4.18 mmol) in POCl₃ (12.59 mL, 137.49 mmol) was added DIEA (1.46 mL, 8.35 mmol) and heated at 100 °C for 5 h. The reaction was cooled down to rt, concentrated and dissolved in DCM (20 mL) at 0 °C. The mixture was added DIEA (1.5 mL) and was stirred at 0 °C for 2h, followed by the addition of NH₄HCO₃ (aq, 20mL). The organic layer was dried over Na₂SO₄ and concentrated and purified by flash column chromatography (silica gel, eluting with 0% to 100% EtOAc/PE) to afford (7S)-2,4-dichloro-7'-fluoro-spiro[6,8-dihydro-5H-pyrido[4,3-d]pyrimidine-7,1'-indane] (**101b**, 1.14 g, 3.52 mmol, 84.19 %yield as brown solid. LCMS calculated for C₁₅H₁₃Cl₂FN₃ (M+H)⁺ m/z = 324.1, 326.1; found: 324.1, 326.1.


[00761] Step 2. Synthesis of 3-chloro-5-[(7S)-2-chloro-7'-fluoro-spiro[6,8-dihydro-5H-pyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**101c**). The mixture of (7S)-2,4-dichloro-7'-fluoro-spiro[6,8-dihydro-5H-pyrido[4,3-d]pyrimidine-7,1'-indane] (**101b**, 300 mg, 0.93 mmol), 3-chloro-N,N-dimethyl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Int-**, 224.61 mg, 0.93 mmol) and DIEA (0.64 mL, 3.7 mmol) in DMSO (2 mL) was stirred at 20 °C for 30 h. The reaction mixture was purified by prep-HPLC on a C18 column (20-35 uM, 100 Å, 40 g) with mobile phase: H₂O (0.1% NH₄HCO₃)/MeCN at flowrate: 50 mL/min to give 3-chloro-5-[(7S)-2-chloro-7'-fluoro-spiro[6,8-dihydro-5H-pyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (364 mg, 0.686 mmol, 74.16% yield). LCMS calculated for C₂₅H₂₇Cl₂FN₇O (M+H)⁺ m/z = 530.2/532.2; found: 530.1/532.2

[00762] Step 3. Synthesis of 3-chloro-5-[(7S)-2-chloro-6-(cyanomethyl)-7'-fluoro-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-

tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**101d**). To a solution of 3-chloro-5-[(7S)-2-chloro-7'-fluoro-spiro[6,8-dihydro-5H-pyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**101c**, 364 mg, 0.69 mmol) in DMF (2mL) was added Cesium carbonate (670.78 mg, 2.06 mmol) and bromoacetonitrile (0.1 mL, 1.37 mmol). The reaction was heated at 50 °C overnight. The reaction mixture was purified by prep-HPLC on a C18 column (20-35 uM, 100 Å, 40 g) with mobile phase: H₂O (0.1% NH₄HCO₃)/MeCN to give 3-chloro-5-[(7S)-2-chloro-6-(cyanomethyl)-7'-fluoro-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (370 mg, 0.650 mmol, 94.68% yield). LCMS calculated for C₂₇H₂₈Cl₂FN₈O (M+H)⁺ m/z = 569.17; found: 569.2


[00763] Step 4. Synthesis of 3-chloro-5-[(7S)-6-(cyanomethyl)-7'-fluoro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 101**). The mixture of 3-chloro-5-[(7S)-2-chloro-6-(cyanomethyl)-7'-fluoro-spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**101d**, 30 mg, 0.05 mmol), [(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methanol (41.93 mg, 0.26 mmol) and sodium tert-butoxide (7.59 mg, 0.08 mmol) in DMSO (1 mL) was stirred at 20 °C under Argon for 2 h. The reaction mixture was purified by prep-HPLC H₂O (0.1% NH₄HCO₃)/MeCN to afford 3-chloro-5-[(7S)-6-(cyanomethyl)-7'-fluoro-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide (**Compound 101**, 10 mg, 0.0135 mmol, 25.60 %yield) as a white solid. LCMS calculated for C₃₃H₄₁ClF₂N₉O₂ (M+H)⁺ m/z = 692.3; found: 692.3. ¹H NMR (400 MHz, CD₃OD) δ 7.34 – 7.39 (m, 1H), 7.12 – 7.14 (m, 1H), 6.95 – 7.00 (m, 1H), 5.26 (d, J=53.8, 1H), 4.82, 4.88 (d, J = 16.8 Hz, 2H), 4.39 – 4.53 (m, 2H), 3.87 – 4.16 (m, 6H), 3.56, 3.73 (d, J = 17.6 Hz, 2H), 3.12 – 3.25 (m, 4H), 3.10 (s, 3H), 3.08 (s, 3H), 2.96 – 3.04 (m, 2H), 2.80 (d, J=17.6, 1H), 2.57 – 2.65 (m, 1H), 1.83 – 2.44 (m, 10H).

[00764] **Compound 102.** 3-chloro-5-[(7S)-6-(cyanomethyl)-2-[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-tetralin]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide



[00765] Compound **102** was prepared similarly to that of **Ex. 28**. LCMS calculated for $C_{36}H_{44}ClFN_9O_2$ ($M+H$)⁺ m/z = 688.33, 690.33; found: 689.5, 691.6. ¹H NMR (400 MHz, CD₃OD) δ = 7.50 (d, J =7.8, 1H), 7.26 (dd, J =10.6, 4.4, 1H), 7.20 (td, J =7.4, 1.4, 1H), 7.15 (d, J =7.4, 1H), 5.27 (d, J =54.4, 1H), 4.91 (d, J =16.6, 1H), 4.80 (d, J =16.6, 1H), 4.36 – 4.56 (m, 2H), 4.04 – 4.22 (m, 3H), 3.86 – 4.03 (m, 3H), 3.45 (m, J = 17.2, 2H), 3.12 – 3.26 (m, 3H), 3.10 (s, 3H), 3.07 (s, 3H), 2.92 – 3.06 (m, 3H), 2.73 – 2.83 (m, 2H), 2.33 – 2.47 (m, 1H), 1.73 – 2.33 (m, 11H).

[00766] **Compound 103.** 3-chloro-5-[(7S)-6-ethyl-7'-fluoro-2-[[[(2R,8S)-2-fluoro-1,2,3,5,6,7-hexahydropyrrolizin-8-yl]methoxy]spiro[5,8-dihydropyrido[4,3-d]pyrimidine-7,1'-indane]-4-yl]-N,N-dimethyl-4,6,7,8-tetrahydropyrazolo[1,5-a][1,4]diazepine-2-carboxamide.



[00767] Compound **103** was prepared similarly to that of **Ex. 28**. LCMS calculated for $C_{35}H_{44}ClF_2N_8O_2$ ($M+H$)⁺ m/z = 681.32; found: 681.3, 683.3. ¹H NMR (400 MHz, CD₃OD) δ = 7.26 – 7.31 (m, 1H), 7.04 – 7.08 (m, 1H), 6.89 – 6.93 (m, 1H), 5.26 (d, J =53.8, 1H), 4.73 – 4.82 (m, 2H), 4.36 – 4.52 (m, 2H), 3.66 – 4.14 (m, 6H), 3.14 – 3.25 (m, 3H), 3.10 (s, 3H), 3.08 (s, 3H), 2.91 – 3.04 (m, 3H), 2.80 (d, J =17.6, 1H), 2.55 – 2.65 (m, 1H), 1.81 – 2.46 (m, 12H), 1.07 (t, J =7.2, 3H).

[00768] **Example 30: Nucleotide Exchange Assay:**

[00769] Ras proteins cycle between an active, GTP bound state, and an inactive GDP-bound state. This activity is tightly regulated by GTPase activating proteins (GAPs) and guanine nucleotide exchange factors (GEFs). GEFs, such as SOS1/2, activate Ras proteins by exchanging GDP for GTP, thus returning Ras to its active conformation (Simanshu, Nissley, & McCormick, 2017). Therefore, a small molecule that binds K-Ras in a manner that prevents SOS-mediated nucleotide exchange locks KRas in its inactive state. Homogenous time resolved fluorescence (HTRF) was used to detect SOS-mediated binding of a fluorescent GTP analog, GTP-DY-647P1 (Jena Biosciences NU-820-647P1) to GST-tagged KRAS^{G12D} (2-169, Reaction Biology, MSC-11-539).

[00770] GST-tagged KRAS^{G12D} (2-169) and anti-GST MAb Tb Cryptate Gold (CisBio 61GSTTLB) were diluted into assay buffer (20 mM HEPES, pH 7.3, 150 mM NaCl, 5 mM MgCl₂, 0.05% BSA 0.0025% NP40, 1 mM DTT) to prepare a 2.5X donor solution. 5X

compound was added to the protein mixture and incubated for 1 h at RT. 2.5X acceptor solution containing SOS_{1cat} (564-1049, Reaction Biology MSC-11-502) and GTP-DY-647P1 were then added to the donor KRAS mixture such that the final concentration of the reaction contained 5 nM GST-tagged KRAS^{G12D} (2-169), 20 nM SOS_{cat}, and 150 nM GTP. The reaction was monitored using at RT with the Envision multimode plate reader (Ex/Em 337/665, 620 nM) up to 90 minutes at 5 minute intervals. Data was blanked to reactions without SOS1 and % inhibition was calculated such that DMSO only = 0% and blank = 100%. Curve fitting was done using a 4 parameter fit. NEA KRAS G12D IC₅₀ (uM) values of selected compounds are depicted in Table 2 with compounds having a value <0.01 uM as ++++; > 0.01 uM to 0.1 uM as +++; > 0.1 uM to 1 uM as ++; > 1 uM to 20 uM as +; and >20 uM as NA.

[00771] Example 31: Protein Constructs for Protein-Protein Interaction:

Table 1: Assay, Protein construct, and protein construct sequences

Assay	Protein Construct	Protein Construct Sequence
PPI	Biotinylated Avi-KRAS-G12D (1-169)	SGLNDIFEAQKIEWHEMTEYKLVVVGADGVGKSALTIQLIQ NHFVDEYDPTIEDSYRKQVVIDGETCLLDILDITAGQEEYSA MRDQYMRTGEGFLCVFAINNTKSFEDIHHYREQIKRVKDS EDVPMVLVGKCDLPSRTVDTKQAQDLARSYGIPFIETSAK TRQGVDDAFYTLVREIRKHKEK
PPI	Biotinylated Avi-KRAS-G12V (1-169)	SGLNDIFEAQKIEWHEMTEYKLVVVGAVGVGKSALTIQLIQ NHFVDEYDPTIEDSYRKQVVIDGETCLLDILDITAGQEEYSA MRDQYMRTGEGFLCVFAINNTKSFEDIHHYREQIKRVKDS EDVPMVLVGKCDLPSRTVDTKQAQDLARSYGIPFIETSAK TRQGVDDAFYTLVREIRKHKEK
PPI	Biotinylated Avi-KRAS wt (1-169)	SGLNDIFEAQKIEWHEMTEYKLVVVGAGGVGKSALTIQLI QNHVFDEYDPTIEDSYRKQVVIDGETCLLDILDITAGQEEYSA MRDQYMRTGEGFLCVFAINNTKSFEDIHHYREQIKRVKDS EDVPMVLVGKCDLPSRTVDTKQAQDLARSYGIPFIETSAK TRQGVDDAFYTLVREIRKHKEK
PPI	Biotinylated Avi-NRAS (1-169)	SGLNDIFEAQKIEWHEMTEYKLVVVGAGGVGKSALTIQLI QNHVFDEYDPTIEDSYRKQVVIDGETCLLDILDITAGQEEYSA MRDQYMRTGEGFLCVFAINNSKSFADINLYREQIKRVKDS DDVPMVLVGKCDLPTRTVDTKQAHELAKSYGIPFIETSAK TRQGVDDAFYTLVREIRQYRMK
PPI	His8-RAF1 RBD (52-131)	SHHHHHHHHSSKTSNTIRVFLPNKQRTVVNVRNGMSLHDC LMKALKVRGLQPECCAVFRLLHEHKGKKARLDWNTDAASL IGEELQVDFL

[00772] Example 32. Recombinant Protein Production:

[00773] Biotinylated KRAS wt and KRAS G12D/V proteins were expressed and purified in conditions similar to those previously reported (Tran, et al., 2021) (Zhang, et al., 2020). Briefly, KRAS (1-169) proteins were expressed in E. coli at 18°C with an upstream TEV cleavage site (ENLFYQS) followed an Avi tag sequence (GLNDIFEAQKIEWHE). KRAS expression

constructs contained both a His6 and maltose-binding protein (MBP) tags at the N-terminus for Ni-NTA column purification prior to overnight TEV cleavage and MBP column purification. The avi-tagged NRAS expression construct contained both a His6 tag and SUMO cleavage site at the N-terminus for Ni-NTA column purification followed by His-ULP1 digestion overnight. All avi-tagged RAS proteins were dialyzed into buffer containing ATP, biotin, and BirA followed by purification over a second Ni-NTA column and then run over a size exclusion HiLoad™ 26/600 Superdex™ column in 20 mM HEPES, pH 7.5, 300 mM NaCl, 5 mM MgCl₂, and 1 mM TCEP. Fractions containing the protein of interest were pooled, concentrated, and confirmed by intact mass spectrometry. To prepare 'GTP' loaded KRAS and NRAS, biotinylated KRAS or NRAS was nucleotide exchanged from GDP-bound protein to GppNHp-bound (Jena Biosciences, NU-401-50) protein in the presence of alkaline phosphatase and excess GppNHp as previously described and the resulting nucleotide content was confirmed by HPLC reverse phase analytical chromatography (Donohue, et al., 2019) (Tran, et al., 2021).

[00774] His-tagged RAF1 (52-131) was similarly expressed in *E. coli* at 18°C overnight with an upstream TEV cleavage site. His-tagged RAF1 expression construct contained both a His6 and MBP tags at the N-terminus for Ni-NTA column purification followed by MBP-tagged TEV digestion overnight. RAF1 protein samples were further purified over a MBP column followed by a Ni-NTA column and a second MBP column. The fractions containing the protein of interest were pooled, concentrated, and further purified over a HiLoad™ 16/600 Superdex™ 75 pg size exclusion column into 20mM HEPES, pH8.0, 200mM NaCl, 5mM TCEP.

[00775] **Example 33: Protein-Protein Interaction (PPI) Assay:**

[00776] When RAS proteins are in the active GTP-bound conformation, they bind the effector protein RAF1 at the N-terminus Ras-binding domain (RBD, residues 52-131) (Tran, et al., 2021). Homogenous time resolved fluorescence (HTRF) was used to monitor the interaction between wt or mutant KRAS and RAF1 or wt NRAS and RAF1. Compounds were assayed in the presence of KRAS G12D/V and RAF1 versus wt KRAS to assess activity against mutant and w.t. KRAS. Similarly, compounds were then assayed in the presence of w.t. NRAS and RAF1 to assess RAS isoform selectivity. In all assay formats, His-tagged RAF1 protein was incubated with the HTRF donor, anti-6His Tb Cryptate gold (Cisbio 61DB10RDF), and biotinylated RAS proteins were incubated with the HTRF acceptor, streptavidin-d2 (CisBio 610SADLA). The intensity of the fluorescence signal emitted is proportional to binding between the two proteins. The donor solution was prepared by mixing 16 nM His-tagged RAF1 in protein dilution buffer with 1:100 anti-6His Tb cryptate in PPI-Terbium detection buffer. 16 nM biotinylated RAS protein was diluted into protein dilution buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.1 mM MgCl₂, 1 mM TCEP, 0.005% Tween20) and mixed with 1:2000 Streptavidin-d2 diluted in PPI-Terbium

detection buffer (CisBio 61DB10RDF). 50X compound in DMSO was mixed with 16 nM KRAS-acceptor solution and incubated for 30 minutes at room temperature. After compound pre-incubation with KRAS, the RAF1 donor solution was added to the KRAS-acceptor solution and incubated for 1 hour at room temperature. The fluorescence signal emitted was monitored at 665 nm and 615 nm using an Envision multimode plate reader. The HTRF ratio (665/615) was calculated and normalized to 0% inhibition in the absence of compound and 100% inhibition in the presence of untagged RAF1 protein. PPI KRAS G12D/RAF1, KRAS G12V/RAF1, w.t.KRAS/RAF1 and NRAS/RAF1 IC₅₀ (uM) values of selected compounds are depicted in Table 4 and Table 5 with compounds having a value <0.1 uM as ++++; > 0.1 uM to 1 uM as +++; > 1 uM to 10 uM as ++; > 10 uM to 100 uM as +; and >100 uM as NA.

[00777] Compounds described herein are active against KRAS G12 mutant and other alleles representative by PPI-G12D, PPI-G12V and PPI-w.t.KRAS potency for broad activity against mutant KRAS and wtKRAS amplification driven malignancies.

[00778] **Example 34.** pERK Inhibition cellular HTRF assay in AGS Cell Lines (Method A)

[00779] The Phospho-ERK cellular HTRF assay measures ERK protein phosphorylated at Thr202/Tyr204 as a readout of MAPK pathway activation (Cisbio 64ERKPEH). AGS cells (ATCC CRL-1739) are cultured in the complete medium containing 10% fetal bovine serum and 1x Penicillin/Streptomycin at 37°C in a humid atmosphere of 5% CO₂ in the air (AGS cells: RPMI 1640 medium).

[00780] On day 1, the cells are plated in tissue-culture treated 96-well plates at the specified densities and allowed to attach for overnight (AGS: 30,000 cells/well). On day 2, the cells are treated with the serially diluted compound solutions in a final concentration of 0.5 % DMSO. After the treatment for the specified time (AGS cells: 3 hours), the supernatant is removed, and the cells are lysed by the lysis buffer supplied with the kit. Then, the cell lysates are treated with the detection reagents overnight at 4°C in darkness. On day 3, the fluorescence intensities at the wavelengths 665 and 620 nm are measured by the Envision plate reader (Perkin Elmer). The data are processed and fitted to a 4-parameter logistic model for IC₅₀ calculations (GraphPad Prism 9).

[00781] **Example 35.** pERK In Cell Western (ICW) assay (Method B)

[00782] pERK ICW is a high throughput screening assay to evaluate the cellular potency of mutant KRAS small molecule inhibitors. KRAS mutant cell line AGS (KRAS^{G12D}) were purchased from ATCC and maintained in DMEM and RPMI medium supplemented with 10% fetal bovine serum and Penicillin/Streptomycin.

[00783] Cells grown in exponential phase were trypsinized, resuspended in fresh media, and viable cells were counted using a cell counter with Trypan Blue (BioRad TC20). Cells were seeded into 384-well plate (Greiner 781091) at density of 5,000 cells/well for AGS and allowed to grow

overnight in a 37°C CO₂ incubator. The next day, compounds were dispensed into wells with a ½ log, 10-point serial dilution and top concentration of 10 µM using Tecan D300e dispenser and incubated for 3 hours in a 37 °C CO₂ incubator. Cells were then fixed with paraformaldehyde (Electron Microscopy Sciences, 15710, 4% final concentration) for 30 min, permeabilized with wash buffer (1X PBS + 0.1% Triton X-100) for 30 min and blocked with Odyssey blocking buffer (Li-COR 927-70001) for 1 hour, all at room temperature (RT). Phospho-ERK antibody (CST 4370L) was diluted 1:500 in Odyssey blocking + 0.2% Tween 20 and incubated with cells overnight at 4 °C. The next day, plates were washed 5x with wash buffer, incubated with IRDye 800 CW, Goat anti-Rabbit secondary antibody (Li-COR 926-32211, 1:500) and DRAQ5 (CST 4084L, 1:5,000) diluted in Odyssey blocking + 0.2% Tween 20 for 1 hour, washed 5x, and imaged on an Odyssey CLx imaging system.

[00784] For data analysis, signal intensities from 800 (phospho-ERK) and 700 (DRAQ5) channels were extracted, and phospho-ERK signals were normalized to DRAQ5 signals for each well and percent of DMSO control values were computed. Data were then imported into Graphpad Prism to compute half-maximal inhibitory concentrations (IC₅₀) using a 4-parameter variable slope model. Z-factor for each plate was computed from signals derived from wells treated with either DMSO or 5 µM of Trametinib. AGS pERK ICW (Method B) IC₅₀ (uM) values of selected compounds are depicted in Table 2 with compounds having a value 0.001 uM to 0.01 uM as +++; > 0.01 uM to 0.1 uM as ++; > 0.1 uM to 1 uM as +; 1 uM to 10 uM as + and > 10 uM as NA.

[00785] Table 2 includes NEA KRAS G12D IC₅₀ (uM) values (<0.01 uM as +++; > 0.01 uM to 0.1 uM as ++; > 0.1 uM to 1 uM as +; > 1 uM to 20 uM as +; and >20 uM as NA), PPI KRAS G12D/RAF1 IC₅₀ (uM) values (<0.1 uM as +++; > 0.1 uM to 1 uM as ++; > 1 uM to 10 uM as +; > 10 uM to 100 uM as +; and >100 uM as NA), AGS pERK HTRF (Method A) IC₅₀ (uM) values (<0.01 uM as +++; > 0.01 uM to 0.1 uM as ++; 0.1 uM to 1 uM as +; 1 uM to 20 uM as + and >20 uM as NA), and AGS pERK ICW (Method B) IC₅₀ (uM) values (0.001 uM to 0.01 uM as +++; > 0.01 uM to 0.1 uM as ++; > 0.1 uM to 1 uM as +; 1 uM to 10 uM as + and > 10 uM as NA) of selected compounds. ND indicates not determined.

Table 2: IC₅₀ (uM) values for various assays

cpd#	NEA-G12D (uM)	PPI-G12D (uM)	pERK-AGS (uM) Method B
1	+	NA	NA
2	+++	+	+
3	+	NA	ND
4	+	NA	NA
5	+	+	NA
6	+	NA	NA
7	+	NA	NA

cpd#	NEA-G12D (uM)	PPI-G12D (uM)	pERK-AGS (uM) Method B
8	++	NA	NA
9	+	ND	+
10	+++	NA	NA
11	+	ND	ND
12	+	ND	ND
13	+	+	ND
14	+	NA	NA
15	+	NA	NA
16	+	NA	NA
17	+	+	+
18	NA	+	NA
19	+	NA	ND
20	+	NA	ND
21	+	ND	ND
22	+	ND	ND
23	+	+	ND
24	++	+	ND
21	+	ND	ND
22	+	ND	ND
23	+	+	ND
24	++	+	ND
21	+	ND	ND
22	+	ND	ND
23	+	+	ND
24	++	+	ND
25	+	NA	ND
24A	+++	+	+
26	+	ND	ND
23A	++	NA	ND
23B	+	ND	ND
27A	++	NA	ND
27B	++++	+	++
28A	+	NA	NA
29	++	+	+
30	+	NA	ND
31	+++	NA	ND
32	+++	NA	ND
33	++	+	ND
34	++++	NA	ND
35	+++	NA	ND
36	++++	+	ND
37	++++	NA	ND
38	++++	+	ND
39	++++	NA	+
39A	++++	+	++
39B	+	ND	NA
40	+++	NA	ND
41	+	ND	NA

cpd#	NEA-G12D (uM)	PPI-G12D (uM)	pERK-AGS (uM) Method B
42	++++	+	++
43	+++	+	+
44	+	ND	ND
45	++++	+	++
46A	++	+	NA
46B	+	ND	ND
47	++++	NA	+
48	+	ND	NA
49	++++	+	++
50	++++	+	++
51A	+	ND	ND
51B	+++	+	ND
52	++++	+	ND
53	++++	+	+++
54	++++	+	++
55	++++	+	++
56	+++	ND	ND
58	+++	ND	ND
59	+++	NA	ND
60	++	+	NA
61	++++	+	+++
62	+++	NA	ND
63	++++	+	+++
64	++++	+	+++
64B	++	NA	ND
65	++++	+	+++
66	+++	ND	NA
67	+++	ND	NA
68	+++	ND	NA
69	++++	+	+++
70	+++	ND	+
71	++	NA	ND
72	++++	+	++
73	+++	NA	ND
74	+++	NA	ND
75	++	ND	ND
76	++++	+	ND
77	++++	+	ND
78	++++	+	ND
79	++++	NA	+
80	+++	+	ND
81	++++	+	+++
82	+++	NA	ND
83	+++	NA	ND
84	++++	NA	+
85	+	NA	ND
86	++++	NA	ND
87	++++	+	++

cpd#	NEA-G12D (uM)	PPI-G12D (uM)	pERK-AGS (uM) Method B
88	+++	NA	ND
89	++	NA	ND
90	++++	+	+++
91	+	NA	ND
92	++	NA	ND
93	++++	+	ND
94	+	NA	ND
95	++++	NA	ND
96	++++	+	+++
97	++++	NA	+
98	++++	NA	ND
99	++++	NA	++
100	+++	NA	ND
101	++	ND	ND
102	+++	ND	ND
103	+++	ND	ND

Table 3 includes KRASG12V/RAF1, wtKRAS/RAF1 and wtNRAS/RAF1 PPI IC₅₀ (uM) values of selected compounds; with compounds having a value <0.1 uM as +++++; 0.1 uM to 1 uM as ++++; >1 uM to 10 uM as ++; >10 uM to 100 uM as +; and >100 uM as NA.

Table 3: IC₅₀ (uM) values for KRASG12V/RAF1, wtKRAS/RAF1 and wtNRAS/RAF1 PPI


cpd#	PPI-G12V (uM)	PPI-w.t.KRAS (uM)	PPI-w.t.NRAS (uM)
23	+	NA	NA
24	+	+	NA
53	+	+	NA
64	+	ND	NA
81	+	+	NA
90	+	NA	NA
96	+	NA	NA

[00786] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS

WHAT IS CLAIMED IS:

1. A compound of Formula (II):



Formula (II)

or a pharmaceutically acceptable salt thereof wherein:

M is selected from O, S, SO, SO₂, and NR³;

R¹ is selected from C₃-C₁₂ carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl-SO₂R²⁰, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein the C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*}; and wherein when M is NR³, Y is O, and R¹ is piperazine, the piperazine is substituted with one or more R⁹;

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃-C₁₂ carbocycle;

Y is selected from a bond, O, S and NR⁵;

R² is selected from -L-N(R²¹)₂, -L-OR²¹, heterocycle, C₁-C₆ alkyl, -L-heterocycle, -L-aryl, -L-heteroaryl, -L-cycloalkyl, -L-NHC(=NH)NH₂, -L-C(O)N(R²¹)₂, -L-C₁-C₆ haloalkyl, -L-OR²¹, -L-NR²¹C(O)-aryl, -L-COOH, -L-NR²¹S(O)₂(R²¹), -L-S(O)₂N(R²¹)₂, -L-N(R²¹)C(O)(OR²¹), -L-OC(O)N(R²¹)₂, and -LC(=O)OC₁-C₆ alkyl, wherein the heterocycle, the aryl portion of -L-NR²¹C(O)-aryl, the heterocycle portion of -L-heterocycle, the cycloalkyl portion of the -L-cycloalkyl are each optionally substituted with one or more R⁶, and wherein the aryl portion of the -L-aryl and the heteroaryl portion of the -L-heteroaryl are each optionally

substituted with one or more R^7 , and wherein when Y is a bond, O, or S, R^2 is further selected from hydrogen;

each L is independently selected from a C_1 - C_4 alkylene optionally substituted with one or more substituents selected from hydroxy, C_1 - C_4 hydroxyalkyl, C_1 - C_4 alkyl, C_3 - C_6 carbocycle, or 3- to 8-membered heterocycle, wherein the C_3 - C_6 carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle or 3- to 8-membered heterocycle, wherein the C_3 - C_6 carbocycle and 3- to 8-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl;

R^3 is selected from hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkoxyalkyl, C_{3-12} carbocycle, and 3- to 12-membered heterocycle, wherein C_{3-12} carbocycle and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C_{1-6} alkyl)₂, C_{1-10} alkyl, - C_{1-10} haloalkyl, -O- C_{1-10} alkyl, oxo, C_{3-12} carbocycle, and 3- to 12-membered heterocycle;

n is selected from 0 to 2;

each R^4 is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, oxo, hydroxyl, halogen, C_{3-12} carbocycle, and 3- to 12-membered heterocycle, wherein the C_1 - C_6 alkyl, C_{3-12} carbocycle, and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from cyano, halogen, —OR⁵, and —N(R⁵)₂;

each R^5 is independently selected from hydrogen or C_1 - C_6 alkyl;

each R^6 is independently selected from halogen, hydroxy, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, oxo, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, cyano, =CH₂, =NO- C_1 - C_3 alkyl, C_1 - C_3 aminoalkyl, -N(R⁵)S(O)₂(R⁵), -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C_1 - C_3 alkyl substituted pyrazolyl, tert-butyldimethylsilyloxyCH₂-, -N(R⁵)₂, (C_1 - C_3 alkoxy) C_1 - C_3 alkyl-, (C_1 - C_3 alkyl)C(=O), oxo, (C_1 - C_3 haloalkyl)C(=O)-, -SO₂F, (C_1 - C_3 alkoxy) C_1 - C_3 alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC C_1 - C_6 alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O) C_1 - C_6 alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂ C_1 - C_6 alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl), -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl)phenyl(C_1 - C_3 alkyl)N(CH₃)₂, -OC(O)NH(C_1 - C_3 alkyl)O(C_1 - C_3 alkyl)phenyl, -OC(O)heterocycle, -O- C_1 - C_3 alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C_1 - C_3 alkyl)(C_1 - C_3 alkyl)phenyl are each optionally substituted with one or more substituents selected from -C(O)H and OH, and wherein the alkyl of -O- C_1 - C_3 alkyl is optionally substituted with

substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocyclyl is optionally substituted with oxo;

each Q is independently selected from a bond, S, and O;

each R⁷ is independently selected from halogen, hydroxy, HC(=O)-, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ hydroxyalkyl, or -N(R⁵)₂;

each R⁹ is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -NR²⁰S(O)₂R²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -NO₂, =O, =NO(R²⁰), -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, =NH, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;


each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle; and

B is selected from a heterocycle and carbocycle, wherein the heterocycle and carbocycle are each optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, =O, -NO₂, C₁-C₄ alkyl, C₁₋₆ aminoalkyl, -S-C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₂-C₄ hydroxyalkynyl, C₁-C₃ cyanoalkyl, triazolyl, C₁-C₃ haloalkyl, -O-C₁-C₃ haloalkyl, -S-C₁-C₃ haloalkyl, C₁-C₃ alkoxy, C₁-C₃ hydroxyalkyl, -CH₂C(=O)N(R⁵)₂, -C₃-C₄ alkynyl(NR⁵)₂, -N(R⁵)₂, (C₁-C₃ alkoxy)haloC₁-C₃ alkyl-, C₁₋₆ alkyl-N(R²⁰)₂, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, -NH₂, =O, =S, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl.

2. The compound or salt of claim 1, wherein R³ is selected from hydrogen and C₁₋₆ alkyl.
3. The compound or salt of claims 1 or 2, wherein M is selected from O, NH, and NMe.
4. The compound or salt of claim 3, wherein M is O.
5. The compound or salt of claim 1, wherein M is selected from NR³.
6. The compound or salt of claim 5, wherein M is selected from NH and NMe.
7. The compound or salt of claim 6, wherein M is NMe.


8. The compound or salt of any one of claims 1 to 7, wherein B is selected from an optionally substituted 5- to 15-membered heterocycle and optionally substituted C₃-C₁₅ carbocycle.
9. The compound or salt of claim 8, wherein B is selected from an optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C₈-C₁₅ fused carbocycle.
10. The compound or salt of claim 9, wherein B is an optionally substituted 8- to 15-membered fused heterocycle.
11. The compound or salt of claim 10, wherein B is an optionally substituted unsaturated C₈-C₁₅ fused carbocycle.
12. The compound or salt of any one of claim 9 to 11, wherein for B, the optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C₈-C₁₅ fused carbocycle are each independently bicyclic or tricyclic.
13. The compound or salt of claim 12, wherein for B the heterocycle and carbocycle are each independently bicyclic.
14. The compound or salt of claim 12, wherein for B the heterocycle and carbocycle are each independently tricyclic.

15. The compound or salt of any one of claims 9 to 14, wherein B is selected from






- , , , , , , and , each of which is optionally substituted with one or more substituents.

16. The compound or salt of claim 15, wherein B is selected from




- , , , , and , each of which is optionally substituted with one or more substituents.



17. The compound or salt of claim 16, wherein B is selected from , and , each of which is optionally substituted with one or more substituents.


18. The compound or salt of any of claims 8 to 17, wherein for B, the one or more optional substituents are independently selected from oxo, -NH₂, halogen, C₁-C₃ alkyl.

19. The compound or salt of any one of claims 1 to 18, wherein B is selected from





20. The compound or salt of any of claims 8 to 17, wherein for B, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, C₁-C₃ alkyl, -B(OR²⁰)₂, -OR²⁰, -C(O)N(R²⁰)₂, -N(R²⁰)₂, =O, -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

21. The compound or salt of any of claims 8 to 17, wherein for B, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OH)₂, -OH, -O-C₁-C₃ haloalkyl, -C(O)NH₂, -NH₂, =O, -CN, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₂₋₆ alkynyl.



22. The compound or salt of any of claims 8 to 21, wherein B is selected from





more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.


30. The compound or salt of claims 28 or 29, wherein R²⁰ is selected from hydrogen and C₁₋₃ alkyl.

31. The compound or salt of any one of claims 27 to 29, wherein the 5- to 12-membered heterocycle of R¹ is unsaturated.



32. The compound or salt of any one of claims 27 to 29, wherein the 5- to 12-membered heterocycle of R¹ is saturated.





33. The compound or salt of any one of claims 27 to 29, wherein the 5- to 12-membered heterocycle of R¹ is a bridged heterocycle.



34. The compound or salt of any one of claims 27 to 33, wherein R¹ is selected from




35. The compound or salt of claim 34, wherein the optional one or more substituents are each independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, =O, -CN, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.

36. The compound or salt of claim 34, wherein R^1 is selected from , ,



, , , and , each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, =O, -CN, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.


37. The compound or salt of claim 36, wherein R^1 is selected from , , and



38. The compound or salt of any one of claims 27 to 33, wherein R^1 is selected from 6- to 8-membered heterocycle, which is optionally substituted.




39. The compound or salt of claim 38, wherein R^1 is selected from 7-membered saturated heterocycle, 8-membered bridged heterocycle, and 6- to 7-membered unsaturated heterocycle, each of which is optionally substituted.

40. The compound or salt of claim 39, wherein R^1 is selected from , ,



41. The compound or salt of claim 40, wherein the one or more optional substituents are independently selected from -OH, -CN, oxo, C₁₋₆ cyanoalkyl.

42. The compound or salt of claims 40 or 41, wherein R¹ is selected from



43. The compound or salt of any one of claims 1 to 26, wherein R¹ is selected from an optionally substituted 6- to 7-membered heterocycle.



44. The compound or salt of claim 43, wherein the 6- to 7-membered heterocycle contains only 1 nitrogen atom, and wherein the 6- to 7-membered heterocycle is optionally substituted.

45. The compound or salt of 44, wherein the 6- to 7-membered heterocycle of R¹ is bound to Formula (II) via the only 1 nitrogen atom.



46. The compound or salt of any one of claims 43 to 45, wherein R¹ is selected from an optionally substituted unsaturated 6-membered heterocycle.

47. The compound or salt of any one of claims 43 to 45, wherein R¹ is selected from an optionally substituted unsaturated 7-membered heterocycle.

48. The compound or salt of any one of claims 43 to 47, wherein R¹ is selected from



49. The compound or salt of any one of claims 43 to 48, wherein R¹ is selected from




50. The compound or salt of any one of claims 43 to 49, wherein the one or more optional substituents of R^1 are each independently selected from halogen, $-OR^{20}$, $-N(R^{20})_2$, $=O$, $-CN$, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} alkyl, $-NHCN$, and C_{2-6} alkynyl.


51. The compound or salt of any one of claims 43 to 50, each is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-NH_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl.

52. The compound or salt of any one of claims 43 to 51, wherein the one or more optional substituents of R^1 are each independently selected from halogen.


53. The compound or salt of any one of claims 43 to 52, wherein R^1 is selected from




54. The compound or salt of any one of claims 43 to 51, wherein R^1 is selected from




wherein each is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-NH_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl.




55. The compound or salt of claim 54, wherein R^1 is selected from , and , wherein each is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-NH_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl.





56. The compound or salt of claim 55, wherein R^1 is selected from , and , wherein each is optionally substituted with one or more substituents independently selected from halogen, and C_{1-6} haloalkyl.




57. The compound or salt of claim 56, wherein R^1 is selected from , and ,



58. The compound or salt of any one of claims 43 to 51, wherein R^1 is selected from , which is optionally substituted with one or more substituents independently selected from halogen, -OH, -NH₂, -NO₂, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} haloalkyl, and C_{1-6} alkyl.



59. The compound or salt of claim 58, wherein R^1 is selected from , which is optionally substituted with one or more substituents independently selected from halogen.



60. The compound or salt of claim 59, wherein R^1 is selected from , and .

61. The compound or salt of claim 43, wherein R^1 is selected from 6- to 7-membered heterocycle.


62. The compound or salt of claim 61, wherein the 6- to 7-membered heterocycle contains only 1 nitrogen atom and optionally one or more additional heteroatoms selected from oxygen, and sulfur.

63. The compound or salt of claim 62, wherein the optionally one or more additional heteroatoms are selected from sulfur.


64. The compound or salt of any one of claims 61 to 63, wherein the 6- to 7-membered heterocycle contains only 1 nitrogen atom and no further additional heteroatoms.

65. The compound or salt of any one of claims 61 to 64, wherein the 6- to 7-membered heterocycle is a non-aromatic 6- to 7-membered heterocycle.

66. The compound or salt of any one of claims 62 to 65, wherein the 6- to 7-membered heterocycle of R^1 is bound to Formula (II) via the only 1 nitrogen atom.




67. The compound or salt of any one of claims 61 to 66, wherein R^1 is selected from




68. The compound or salt of claim 67, wherein the one or more optional substituents of R^1 are each independently selected from halogen, -OH, -CN, C_{1-6} cyanoalkyl, -NHCN, C_{1-6} alkyl, oxo, and C_{2-6} alkynyl.

69. The compound or salt of claim 43, wherein R^1 is selected from an optionally substituted 6- to 10-membered heterocycle.




70. The compound or salt of claim 69, wherein R^1 is selected from




optionally substituted.

71. The compound or salt of claim 70, wherein the one or more optional substituents are independently selected from halogen, =O, -OH, -C(O)N(R^{20})₂, C_{2-6} alkynyl, -NHCN, -CN, C_{1-6} aminoalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, and C_{1-6} alkyl.




72. The compound or salt of claim 71, wherein R¹ is selected from




73. The compound or salt of claim 27, wherein R¹ is selected from a 7- to 11-membered spiro heterocycle.

74. The compound or salt of claim 73, wherein R¹ is selected from a 10-membered spiro heterocycle.


75. The compound or salt of claims 73 or 74, wherein the spiro heterocycle has at least 3 nitrogen atoms.




76. The compound or salt of claim 75, wherein R¹ is selected from




, each of which is optionally substituted.




77. The compound or salt of claim 76, wherein R¹ is selected from



, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, =O, -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.




78. The compound or salt of claim 77, wherein R^1 is selected from




79. The compound or salt of claim 27, wherein R^1 is selected from an optionally substituted unsaturated 9- to 11-membered heterocycle.


80. The compound or salt of claim 79, wherein R^1 is selected from an optionally substituted unsaturated 10-membered heterocycle.




81. The compound or salt of claim 80, wherein R^1 is substituted.



82. The compound or salt of claim 81, wherein R^1 is substituted with one or more substituents selected from halogen, -OH, -C(O)N(R^{20})₂, -C(O)OR²⁰, -C(O)NHOR²⁰, -N(R^{20})₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, and C₂₋₆ alkynyl.




83. The compound or salt of claim 82, wherein R^1 is selected from



substituted with one or more substituents selected from halogen, -OH, -N(R^{20})₂, =O, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, and C₂₋₆ alkynyl.

84. The compound or salt of claim 27, wherein R^1 is selected from an optionally substituted unsaturated 9- to 11-membered heterocycle.


85. The compound or salt of claim 84, wherein R^1 is selected from an optionally substituted unsaturated 10-membered heterocycle.



86. The compound or salt of claim 85, wherein R^1 is , which is optionally substituted.


87. The compound or salt of claim 86, wherein the one or more optional substituents of R^1 are independently selected from halogen, -OH, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-C(O)N(R^{20})_2$, $-C(=NR^{20})N(R^{20})_2$, $-C(O)OR^{20}$, $-C(O)NHR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-NO_2$, $=O$, $-CN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} alkoxyalkyl, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, $-OR^{20}$, and C_{1-6} alkyl.


88. The compound or salt of any one of claims 84 to 87, wherein R^1 is selected from



, each of which is optionally substituted.

89. The compound or salt of any one of claims 84 to 88, wherein R¹ is selected from,






90. The compound or salt of any one of claims 84 to 87, wherein the one or more optional substituents of R^1 are independently selected from halogen, $-OH$, $-N(R^{20})_2$, $=O$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, $-C(O)N(R^{20})_2$, C_{1-6} alkyl, and C_{2-6} alkynyl.

91. The compound or salt of claim 90, wherein the one or more optional substituents of R^1 are independently selected from halogen, and $-C(O)N(R^{20})_2$.


92. The compound or salt of claim 91, wherein R^1 is selected from




93. The compound or salt of any one of claims 84 to 87, wherein the one or more optional substituents of R^1 are independently selected from halogen, $-N(R^{20})_2$, $-CN$, C_{1-6} alkyl, C_{1-6} cyanoalkyl, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} alkyl- SO_2 - C_{1-6} alkyl, C_{2-6} alkenyl, $-C(O)NR^{20}OR^{20}$, $-C(O)N(R^{20})_2$, $-C(O)R^{20}$, and 5- to 10-membered heterocycle, wherein the 5- to


10-membered heterocycle is optionally substituted independently with one or more R^{1*} , wherein each R^{1*} is independently selected from halogen, $-OR^{20}$, and C_{1-6} alkyl.


94. The compound or salt of claims 93, wherein the one or more optional substituents of R^1



95. The compound or salt of claim 94, wherein R^1 is selected from








96. The compound or salt of any one of claims 1 to 26, wherein R^1 is an optionally substituted 12- to 15-membered heterocycle.
97. The compound or salt of claim 96, wherein R^1 is an optionally substituted 12-membered heterocycle.
98. The compound or salt of claim 96, wherein R^1 is an optionally substituted 13-membered heterocycle.
99. The compound or salt of claim 96, wherein R^1 is an optionally substituted 14-membered heterocycle.
100. The compound or salt of claim 96, wherein R^1 is an optionally substituted 15-membered heterocycle.
101. The compound or salt of any one of claims 96 to 100, wherein the heterocycle of R^1 is tricyclic.
102. The compound or salt of any one of claims 96 to 101, wherein the heterocycle of R^1 contains a fused heterocycle.
103. The compound or salt of any one of claims 96 to 102, wherein the heterocycle of R^1 contains a spiro heterocycle.
104. The compound or salt of any one of claims 96 to 103, wherein the heterocycle of R^1 contains a fused and spiro-heterocycle.
105. The compound or salt of any one of claims 96 to 104, wherein the heterocycle of R^1 is an unsaturated heterocycle.
106. The compound or salt of any one of claims 96 to 105, wherein the heterocycle of R^1 is a non-aromatic heterocycle.
107. The compound or salt of any one of claims 96 to 106, wherein the heterocycle of R^1 has at least one double bond.

108. The compound or salt of any one of claims 96 to 107, wherein the heterocycle of R¹ has at least two double bonds.

109. The compound or salt of any one of claims 96 to 108, wherein the heterocycle of R¹ has two double bonds.


110. The compound or salt of any one of claims 96 to 109, wherein R¹ is selected from




111. The compound or salt of claim 110, wherein the optional one or more substituents are selected from halogen, -OH, -NHCN, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(O)NHR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, and C₂₋₆ alkynyl.




112. The compound or salt of claim 111, wherein the optional one or more substituents are selected from halogen, -OH, C₁₋₆ alkyl, and -C(O)N(R²⁰)₂.


113. The compound or salt of any one of claims 96 to 112, wherein R¹ is selected from








(preferably 5- to 6-membered heterocycle), wherein the 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*} ; each R^{1*} is independently selected from halogen, C_{1-6} haloalkyl, and C_{1-6} alkyl.

117. The compound or salt of claim 116, wherein R^1 is selected , , and , each of which is optionally substituted.

118. The compound or salt of claim 117, wherein R^1 is selected , which is optionally substituted.

119. The compound or salt of claim 118, wherein R^1 is selected from



120. The compound or salt of claim 117, wherein R^1 is selected  and .

121. The compound or salt of any one of claims 1 to 120, wherein Y is O.

122. The compound or salt of any one of claims 1 to 121, wherein R^2 is -L-heterocycle, optionally substituted with one or more R^6 .


123. The compound or salt of claim 122, wherein L is selected from C_1 - C_4 alkylene.


124. The compound or salt of claim 123, wherein L is selected from unsubstituted C₁-C₄ alkylene.

125. The compound or salt of claims 123, wherein each L is independently selected from a C₁-C₄ alkylene optionally substituted; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle, wherein the C₃-C₆ carbocycle is optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl.

126. The compound or salt of claims 125, wherein the optional substituents of L are selected from C₁-C₄ hydroxyalkyl, C₁-C₄ alkyl, C₃-C₆ carbocycle; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle or 3- to 8-membered heterocycle wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen and C₁₋₆ haloalkyl.

127. The compound or salt of claim 1 to 126, wherein Y-R² is selected from





and , wherein the heterocycle portion is optionally substituted with one or more R⁶.


128. The compound or salt of claim 127, wherein R⁶ of R² is independently selected at each occurrence from halogen, hydroxy, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, cyano, and C₁-C₃ aminoalkyl.

129. The compound or salt of claim 128, wherein R⁶ of R² is independently selected at each occurrence from C₁-C₃ alkyl and halogen.

130. The compound or salt of claim 127 to 129, wherein Y-R² is selected from



131. The compound or salt of any one of claims 121 to 125, wherein Y-R² is selected from




optionally substituted with one or more R⁶.


132. The compound or salt of claim 131, wherein R^6 is selected from halogen, -OH, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, oxo, C_1 - C_3 alkoxy, -CN, and C_1 - C_3 aminoalkyl.

133. The compound or salt of claim 132, wherein R^6 is selected from halogen and C_1 - C_3 alkyl.

134. The compound or salt of claim 133, wherein $Y-R^2$ is selected from



135. The compound or salt of claim 131, $Y-R^2$ selected from



136. The compound or salt of claim 1 to 121, wherein R^2 is selected from -L-heterocycle, wherein the heterocycle portion of -L-heterocycle is optionally substituted with one or more R^6 .

137. The compound or salt of claim 136, wherein each L is independently selected from a substituted C_1 - C_4 alkylene, and wherein two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle, wherein the C_3 - C_6 carbocycle is optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C_1 - C_6 aminoalkyl, C_1 - C_6 alkoxy, C_1 - C_6 hydroxyalkyl, C_1 - C_6 haloalkyl.


138. The compound or salt of claim 137, wherein each L is independently selected from a substituted C_1 - C_4 alkylene, and wherein two substituents on the same carbon atom of L come together to form a C_3 - C_6 carbocycle.


139. The compound or salt of claim 138, wherein each L is independently selected from a substituted C_3 alkylene, and wherein two substituents on the same carbon atom of L come together to form a C_3 carbocycle.

140. The compound or salt of claim 139, wherein each L is independently selected from




141. The compound or salt of claim 140, wherein $Y-R^2$ is






142. The compound or salt of claim 1 to 126, wherein $Y-R^2$ is selected from
wherein the heterocycle is optionally substituted.

143. The compound or salt of claim 142, wherein the heterocycle is optionally substituted with one or more substituent selected from halogen, hydroxy, C_1 - C_3 alkyl, $-N(R^5)S(O)_2(R^5)$, $-OC(O)N(R^5)_2$, $=CH_2$, oxo, $=NO$ - C_1 - C_3 alkyl, $-CH_2OC(O)$ heterocycle, $-CH_2$ heterocycle, $-CH_2OC(O)N(R^5)_2$, and $-O$ - C_1 - C_3 alkyl, wherein the alkyl of $-O$ - C_1 - C_3 alkyl is optionally substituted with substituents selected from heterocycle, oxo, and hydroxy.




144. The compound or salt of claim 143, wherein $Y-R^2$ is selected from




145. The compound or salt of any one of claims 1 to 126 or 136 to 140, wherein R^2 is selected from $-L-N(R^{21})_2$.


146. The compound or salt of any one of claim 145, wherein L is independently selected from a substituted C₁-C₄ alkylene, and wherein two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle, wherein the C₃-C₆ carbocycle is optionally substituted with one or more substituents selected from halogen.



147. The compound or salt of claims 1, 145 or 146, wherein L is





148. The compound or salt of any one of claims 1, or 145 to 147, wherein R² is





149. The compound or salt of any one of claims 1, or 145 to 147, wherein Y-R² is

150. The compound or salt of claim 1, wherein the compound is selected from:










, or a pharmaceutically acceptable salt of any one thereof.

151. A compound of Formula (III):



Formula (III)

or a pharmaceutically acceptable salt thereof wherein:

R¹ is selected from C₃-C₁₂ carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein the C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*};

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃-C₁₂ carbocycle;

R² is selected from -L-NR²¹S(O)₂(R²¹), -L-S(O)₂N(R²¹)₂, -L-N(R²¹)C(O)(OR²¹), -L-OC(O)N(R²¹)₂, and L-bicyclic heterocycle, wherein the bicyclic heterocycle is optionally substituted with one or more R⁶;

each L is independently selected from a C₁-C₄ alkylene optionally substituted with one or more substituents selected from hydroxy, C₁-C₄ hydroxyalkyl, C₁-C₄ alkyl, C₃-C₆ carbocycle, or 3- to 8-membered heterocycle, wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle or 3- to 8-membered heterocycle, wherein the C₃-C₆ carbocycle and 3- to 8-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl;

n is selected from 0 to 3;

each R⁴ is independently selected from C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, oxo, hydroxyl, halogen, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, wherein the C₁-C₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from cyano, halogen, —OR⁵, and —N(R⁵)₂;

B is selected from a heterocycle and carbocycle, wherein the heterocycle or carbocycle is optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, =O, -NO₂, C₁-C₄ alkyl, C₁₋₆ aminoalkyl, -S-C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₂-C₄ hydroxyalkynyl, C₁-C₃ cyanoalkyl, triazolyl, C₁-C₃ haloalkyl, -O-C₁-C₃ haloalkyl, -S-C₁-C₃ haloalkyl, C₁-C₃ alkoxy, C₁-C₃ hydroxyalkyl, -CH₂C(=O)N(R⁵)₂, -C₃-C₄ alkynyl(NR⁵)₂, -N(R⁵)₂, (C₁-C₃ alkoxy)haloC₁-C₃ alkyl-, C₁₋₆ alkyl-N(R²⁰)₂, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, -NH₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl;

Y is selected from a bond, O, S and NR⁵;

each R⁶ is independently selected from halogen, hydroxy, C₁-C₃ hydroxyalkyl, C₁-C₃ alkyl, oxo, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, cyano, =CH₂, =NO-C₁-C₃ alkyl, C₁-C₃ aminoalkyl, -N(R⁵)S(O)₂(R⁵), -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C₁-C₃ alkyl substituted pyrazolyl, tert-butyldimethylsilyloxyCH₂-, -N(R⁵)₂, (C₁-C₃ alkoxy)C₁-C₃ alkyl-, (C₁-C₃ alkyl)C(=O), oxo, (C₁-C₃ haloalkyl)C(=O)-, -SO₂F, (C₁-C₃ alkoxy)C₁-C₃ alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC₁-C₆ alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O)C₁-C₆ alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂C₁-C₆ alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl), -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl(C₁-C₃ alkyl)N(CH₃)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl, -OC(O)heterocycle, -O-C₁-C₃ alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C₁-C₃ alkyl)(C₁-C₃ alkyl)phenyl are optionally substituted with one or more substituents selected from -

C(O)H and OH, and wherein the alkyl of -O-C₁-C₃ alkyl is optionally substituted with substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocyclyl is optionally substituted with oxo;

each Q is independently selected from a bond, S, and O;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle; and

each R⁵ is independently selected from hydrogen or C₁-C₆ alkyl.

152. The compound or salt of claim 151, wherein R² is selected from L-pyrrolizine, wherein the pyrrolizine is optionally substituted with one or more R⁶.

153. The compound or salt of any one of claims 151 to 152, wherein B is selected from an optionally substituted 5- to 15-membered heterocycle and optionally substituted C₃-C₁₅ carbocycle.

154. The compound or salt of claim 153, wherein B is selected from an optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C₈-C₁₅ fused carbocycle.

155. The compound or salt of claim 154, wherein B is selected from an optionally substituted 8- to 15-membered fused heterocycle.


156. The compound or salt of claim 154, wherein B is selected from an optionally substituted unsaturated C₈-C₁₅ fused carbocycle.

157. The compound or salt of any one of claim 153 to 156, wherein for B the optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C₈-C₁₅ fused carbocycle are each bicyclic and tricyclic.


158. The compound or salt of claim 157, wherein for B the heterocycle and carbocycle are each independently bicyclic.

159. The compound or salt of claim 157, wherein for B the heterocycle and carbocycle are each independently tricyclic.


160. The compound or salt of any one of claims 151 to 159, wherein B is selected from



161. The compound or salt of claim 160, wherein B is selected from




162. The compound or salt of claim 161, wherein B is selected from,



163. The compound or salt of any of claims 151 to 162, wherein for B, the one or more optional substituents of the heterocycle and carbocycle are independently selected from oxo, -NH₂, halogen, C₁-C₃ alkyl.

164. The compound or salt of any one of claims 151 to 163, wherein B is selected from




165. The compound or salt of any of claims 151 to 162, wherein for B, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each

occurrence from halogen, C₁-C₃ alkyl, -B(OR²⁰)₂, -OR²⁰, -C(O)N(R²⁰)₂, -N(R²⁰)₂, =O, -CN, -NHCN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

166. The compound or salt of claim 155, wherein for B, the one or more optional substituents of the heterocycle and carbocycle are independently selected at each occurrence from halogen, oxo, -NH₂, C₁-C₃ alkyl, -B(OH)₂, -OH, -C(O)NH₂, -NH₂, =O, -CN, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₂₋₆ alkynyl.

167. The compound or salt of any of claims 165 to 166, wherein B is selected from





168. The compound or salt of any one of claims 151 to 167, wherein each R⁴ is independently selected from C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, oxo, hydroxyl, halogen.

169. The compound or salt of claim 168, wherein each R⁴ is independently selected from C₁₋₆ alkyl, oxo, and halogen.


170. The compound or salt of any one of claims 151 to 169, wherein n is selected from 1 and 2.

171. The compound or salt of any one of claims 151 to 169, wherein n is 0.
172. The compound or salt of any one of claims 151 to 171, wherein Y is O.
173. The compound or salt of any one of claims 151 to 172, wherein R¹ is selected from optionally substituted 5- to 12-membered heterocycle.
174. The compound or salt of claim 173, wherein R¹ is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, =O, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.
175. The compound or salt of claim 174, wherein R¹ is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.
176. The compound or salt of claims 174 or 175, wherein R²⁰ is selected from hydrogen and C₁₋₃ alkyl.
177. The compound or salt of claims 151 to 176, wherein the 5- to 12-membered heterocycle of R¹ is unsaturated.
178. The compound or salt of claims 151 to 176, wherein the 5- to 12-membered heterocycle of R¹ is saturated.
179. The compound or salt of claims 151 to 176, wherein the 5- to 12-membered heterocycle of R¹ is bridged.
180. The compound or salt of claims 151 to 176, wherein the 5- to 12-membered heterocycle of R¹ is a spiro heterocycle.







independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, =O, -CN, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.



, and , each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -N(R²⁰)₂, -NO₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, =O, -CN, C₁₋₆ hydroxyalkyl, and C₁₋₆ haloalkyl.




184. The compound or salt of claims 151 to 183, wherein R² is -L-heterocycle, optionally substituted with one or more R⁶.

185. The compound or salt of 184, wherein L is selected from C₁-C₄ alkylene.

186. The compound or salt of claim 185, wherein L is selected from unsubstituted C₁-C₄ alkylene.

187. The compound or salt of any one of claims 184 to 186, wherein $Y-R^2$ is selected from




, wherein the heterocycle portion is optionally substituted with one or more R^6 .


188. The compound or salt of claim 187, wherein R^6 of R^2 is independently selected at each occurrence from halogen, hydroxy, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, cyano, and C_1 - C_3 aminoalkyl.

189. The compound or salt of claim 188, wherein R^6 of R^2 is independently selected at each occurrence from C_1 - C_3 alkyl and halogen.


190. The compound or salt of claim 189, wherein $Y-R^2$ is selected from



and



191. A compound of Formula (I):



Formula (I)

or a pharmaceutically acceptable salt thereof wherein:

R^1 is selected from C_3 - C_{12} carbocycle and 5- to 15-membered heterocycle, each of which are optionally substituted with one or more substituents independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, $-SR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(=NR^{20})N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, C_{1-6} cyanoalkyl, C_{1-6} haloalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle, wherein the C_3 - C_{12} carbocycle and 5- to 12-membered heterocycle are each optionally substituted independently with one or more R^{1*} ;

each R^{1*} is independently selected from halogen, $-B(OR^{20})_2$, $-OR^{20}$, $-SR^{20}$, $-S(O)_2(R^{20})$, $-S(O)_2N(R^{20})_2$, $-S(O)N(R^{20})_2$, $-S(O)R^{20}(=NR^{20})$, $-NR^{20}S(O)_2R^{20}$, $-C(O)N(R^{20})_2$, $-C(O)NR^{20}OR^{20}$, $-N(R^{20})C(O)R^{20}$, $-N(R^{20})C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-OC(O)R^{20}$, $-OC(O)N(R^{20})_2$, $-NO_2$, $=O$, $=N(R^{20})$, $=NO(R^{20})$, $-CN$, $-NHCN$, C_{1-6} alkyl- $N(R^{20})_2$, C_1 -

₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃₋₁₂ carbocycle;

Y is selected from a bond, O, S and NR⁵;

R² is selected from hydrogen, -N(R²¹)₂, -L-N(R²¹)₂, -L-OR²¹, heterocycle, C₁₋₆ alkyl, -L-heterocycle, -L-aryl, -L-heteroaryl, -L-cycloalkyl, -L-N(R²¹)₂, -L-NHC(=NH)NH₂, -L-C(O)N(R²¹)₂, -L-C₁₋₆ haloalkyl, -L-OR²¹, -L-NR²¹C(O)-aryl, -L-COOH, -L-NR²¹S(O)₂(R²¹), -L-S(O)₂N(R²¹)₂, -L-N(R²¹)C(O)(OR²¹), -L-OC(O)N(R²¹)₂, and -LC(=O)OC₁₋₆ alkyl, wherein the heterocycle and the aryl portion of -L-NR⁵C(O)-aryl and the heterocycle portion of -L-heterocycle and the cycloalkyl portion of the -L-cycloalkyl may be optionally substituted with one or more R⁶, and wherein the aryl or heteroaryl of the -L-aryl and the -L-heteroaryl may be optionally substituted with one or more R⁷;

each L is independently selected from a C₁₋₄ alkylene optionally substituted with one or more substituents selected from hydroxy, C₁₋₄ hydroxyalkyl, C₁₋₄ alkyl, C₃₋₆ carbocycle, or 3- to 8-membered heterocycle, wherein the C₃₋₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃₋₆ carbocycle or 3- to 8-membered heterocycle wherein the C₃₋₆ carbocycle and 3- to 8-membered heterocycle are optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl;

each R⁴ is independently selected from C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, wherein C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle are each optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R⁵ is independently selected from hydrogen or C₁₋₆ alkyl;

each R⁶ is independently selected from halogen, hydroxy, C₁₋₃ hydroxyalkyl, C₁₋₃ alkyl, oxo, C₁₋₃ haloalkyl, C₁₋₃ alkoxy, cyano, =CH₂, =NO-C₁₋₃ alkyl, C₁₋₃ aminoalkyl, -N(R⁵)S(O)₂(R⁵), -Q-phenyl, -Q-phenylSO₂F, -NHC(O)phenyl, -NHC(O)phenylSO₂F, C₁₋₃ alkyl substituted pyrazolyl, tert-butyldimethylsilyloxyCH₂-, -N(R⁵)₂, (C₁₋₃ alkoxy)C₁₋₃ alkyl-, (C₁₋₃ alkyl)C(=O), oxo, (C₁₋₃ haloalkyl)C(=O)-, -SO₂F, (C₁₋₃ alkoxy)C₁₋₃ alkoxy, -CH₂OC(O)N(R⁵)₂, -CH₂NHC(O)OC₁₋₆ alkyl, -CH₂NHC(O)N(R⁵)₂, -CH₂NHC(O)C₁₋₆ alkyl, -CH₂(pyrazolyl), -CH₂NHSO₂C₁₋₆ alkyl, -CH₂OC(O)heterocycle, -OC(O)N(R⁵)₂, -OC(O)NH(C₁₋₃ alkyl)O(C₁₋₃ alkyl), -OC(O)NH(C₁₋₃ alkyl)O(C₁₋₃ alkyl)phenyl(C₁₋₃

alkyl)N(CH₃)₂, -OC(O)NH(C₁-C₃ alkyl)O(C₁-C₃ alkyl)phenyl, -OC(O)heterocycle, -O-C₁-C₃ alkyl, and -CH₂heterocycle, wherein the phenyl of -NHC(O)phenyl and -OC(O)NH(C₁-C₃ alkyl)(C₁-C₃ alkyl)phenyl are optionally substituted with one or more substituents selected from -C(O)H and OH, and wherein the alkyl of -O-C₁-C₃ alkyl is optionally substituted with substituents selected from heterocycle, oxo and hydroxy; and wherein the heterocycle of -CH₂heterocycle is optionally substituted with oxo;

each R⁷ is independently selected from halogen, hydroxy, HC(=O)-, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ hydroxyalkyl, or -N(R⁵)₂;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, =NH, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each R²¹ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

each Q is independently selected from a bond, S, and O;

B is selected from a heterocycle and carbocycle, wherein the heterocycle and carbocycle are optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxy, =O, -NO₂, C₁-C₄ alkyl, C₁₋₆ aminoalkyl, -S-C₁-C₃ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₂-C₄ hydroxyalkynyl, C₁-C₃ cyanoalkyl, triazolyl, C₁-C₃ haloalkyl, -O-C₁-C₃ haloalkyl, -S-C₁-C₃ haloalkyl, C₁-C₃ alkoxy, C₁-C₃ hydroxyalkyl, -CH₂C(=O)N(R⁵)₂, -C₃-C₄ alkynyl(NR⁵)₂, -N(R⁵)₂, (C₁-C₃ alkoxy)haloC₁-C₃ alkyl-, C₁₋₆ alkyl-N(R²⁰)₂, C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle, wherein C₃-C₁₂ carbocycle and 5- to 12-membered heterocycle are each optionally substituted with one or more substituents selected from halogen, -OH, -NO₂, -NH₂, =O, =S, -CN, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ haloalkyl, and wherein B forms a spirocycle with Ring A; and

Ring A is selected from a heterocycle and carbocycle, wherein the heterocycle or carbocycle is optionally substituted with one or more substituents selected from R⁴.

192. The compound or salt of claim 191, wherein B is selected from an optionally substituted 5- to 15-membered heterocycle and optionally substituted C₃-C₁₅ carbocycle.

193. The compound or salt of claim 192, wherein B is selected from an optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C₈-C₁₅ fused carbocycle.

194. The compound or salt of claim 193, wherein B is an optionally substituted 8- to 15-membered fused heterocycle.


195. The compound or salt of claim 193, wherein B is an optionally substituted unsaturated C₈-C₁₅ fused carbocycle.

196. The compound or salt of any one of claim 191 to 195, wherein for B, the optionally substituted 8- to 15-membered fused heterocycle and optionally substituted C₈-C₁₅ fused carbocycle are each independently bicyclic or tricyclic.


197. The compound or salt of claim 196, wherein for B, the heterocycle and carbocycle are each independently bicyclic.

198. The compound or salt of claim 196, wherein for B, the heterocycle and carbocycle are each independently tricyclic.


199. The compound or salt of any one of claims 191 to 198, wherein B is selected from



200. The compound or salt of claim 199, wherein B is selected from




201. The compound or salt of claim 200, wherein B is selected from,



202. The compound or salt of any of claims 191 to 201, wherein for B, the one or more optional substituents of the heterocycle or carbocycle are independently selected from oxo, -NH₂, halogen, C₁-C₃ alkyl.

203. The compound or salt of any one of claims 191 to 202, wherein B is selected from



204. The compound or salt of any one of claims 191 to 203, wherein each R^4 is independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, oxo, hydroxyl, halogen.

205. The compound or salt of claim 204, wherein each R^4 is independently selected from C_{1-6} alkyl, oxo, and halogen.

206. The compound or salt of any one of claims 191 to 205, wherein Y is O.

207. The compound or salt of any one of claims 191 to 206, wherein R^1 is selected from optionally substituted 5- to 5-membered heterocycle.

208. The compound or salt of claim 207, wherein R^1 is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-N(R^{20})_2$, $-NO_2$, $=O$, $-CN$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl.

209. The compound or salt of claim 208, wherein R^1 is selected from 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, $-OH$, $-N(R^{20})_2$, $-NO_2$, C_{1-6} aminoalkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, and C_{1-6} haloalkyl.


210. The compound or salt of claims 208 to 209, wherein R^{20} is selected from hydrogen and C_{1-3} alkyl.

211. The compound or salt of any one of claims 191 to 210, wherein the 5- to 12-membered heterocycle of R^1 is unsaturated.


212. The compound or salt of any one of claims 191 to 210, wherein the 5- to 12-membered heterocycle of R^1 is saturated.

213. The compound or salt of any one of claims 191 to 210, wherein the 5- to 12-membered heterocycle of R^1 is bridged.





216. The compound or salt of claim 215, wherein R^1 is selected from




217. The compound or salt of any one of claims 191 to 216, wherein R^2 is -L-heterocycle, optionally substituted with one or more R^6 .

218. The compound or salt of any one of claims 191 to 217, wherein L is selected from C_1 - C_4 alkylene.

219. The compound or salt of claims 217 or 218, wherein L is selected from unsubstituted C_1 - C_4 alkylene.

220. The compound or salt of any one of claims 181 to 209, wherein $Y-R^2$ is selected from




221. The compound or salt of claim 220, wherein R^6 of R^2 is independently selected at each occurrence from halogen, hydroxy, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, cyano, and C_1 - C_3 aminoalkyl.

222. The compound or salt of claim 221, wherein R^6 of R^2 is independently selected at each occurrence from C_1 - C_3 alkyl and halogen.

223. The compound or salt of claim 222, wherein $Y-R^2$ is selected from



224. A compound of Formula (II*):



Formula (II*)

or a pharmaceutically acceptable salt thereof wherein:

M is selected from O, and NR³;

R³ is selected from hydrogen, C₁₋₆ alkyl, and C₁₋₆ cyanoalkyl;

R¹ is selected from a 7- to 10-membered heterocycle, wherein the 7- to 10-membered heterocycle is optionally substituted with one or more substituents independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted independently with one or more R^{1*};

each R^{1*} is independently selected from halogen, -B(OR²⁰)₂, -OR²⁰, -SR²⁰, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -NR²⁰S(O)₂R²⁰, -C(O)N(R²⁰)₂, -C(O)NR²⁰OR²⁰, -N(R²⁰)C(O)R²⁰, -N(R²⁰)C(O)N(R²⁰)₂, -N(R²⁰)C(O)OR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -C(O)OR²⁰, -OC(O)R²⁰, -OC(O)N(R²⁰)₂, -NO₂, =O, =N(R²⁰), =NO(R²⁰), -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₃-C₁₂ carbocycle;

B is selected from C₆-C₁₅ carbocycle, wherein the C₆-C₁₅ carbocycle is optionally substituted with one or more substituents independently selected from halogen, C₁-C₃ alkyl, -B(OR²⁰)₂, -OR²⁰, -C(O)N(R²⁰)₂, -N(R²⁰)₂, =O, -CN, -NHCN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

R² is selected from -L-heterocycle, wherein the heterocycle of -L-heterocycle is optionally substituted with one or more R⁶;

L is independently selected from a C₁-C₄ alkylene, wherein the C₁-C₄ alkylene is optionally substituted with one or more substituents selected from hydroxy, C₁-C₄ hydroxyalkyl, C₁-C₄ alkyl; and wherein optionally two substituents on the same carbon atom of L come together to form a C₃-C₆ carbocycle;

each R²⁰ is independently selected from hydrogen; and C₁₋₆ alkyl, C₃₋₆ carbocycle, and 3- to 6-membered heterocycle, each of which is optionally substituted with one or more substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -N(C₁₋₆ alkyl)₂, C₁₋₁₀ alkyl, -C₁₋₁₀ haloalkyl, -O-C₁₋₁₀ alkyl, oxo, =NH, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle.

225. The compound or salt of claim 224, wherein R¹ is selected from an optionally substituted 7- to 10-membered spiro heterocycle and optionally substituted 7- to 10-membered fused heterocycle.

226. The compound or salt of claims 224 or 225, wherein the heterocycle of R^1 has at least one nitrogen atom.


227. The compound or salt of claim 226, wherein the at least one nitrogen at of the heterocycle of R^1 is bound to Formula (II*).

228. The compound or salt of anyone of claims 225 to 227, wherein R^1 is selected from an optionally substituted 10-membered spiro heterocycle and optionally substituted 10-membered fused heterocycle.

229. The compound or salt of anyone of claims 225 to 228, wherein the optional one or more substituents of R^1 are independently selected from halogen, -OH, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)OR²⁰, -C(O)NHOR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxyalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, and C₁₋₆ alkyl.


230. The compound or salt of anyone of claims 225 to 229, wherein the optional one or more substituents of R^1 are independently selected from halogen, -OH, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)OR²⁰, -C(O)NHOR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxyalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, and C₁₋₆ alkyl.

231. The compound or salt of anyone of claims 225 to 230, wherein R^1 is selected from



, which is substituted.




232. The compound or salt of anyone of claims 225 to 231, wherein R^1 is selected from




substituted with one or more substituents independently selected from halogen, -OH, -S(O)₂(R²⁰), -S(O)₂N(R²⁰)₂, -S(O)N(R²⁰)₂, -S(O)R²⁰(=NR²⁰), -C(O)N(R²⁰)₂, -C(=NR²⁰)N(R²⁰)₂, -C(O)OR²⁰, -C(O)NHOR²⁰, -N(R²⁰)₂, -C(O)R²⁰, -NO₂, =O, -CN, C₁₋₆ alkyl-N(R²⁰)₂, C₁₋₆ aminoalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxyalkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₂₋₆ alkynyl, 5- to 12-membered heterocycle, wherein the 5- to 12-membered heterocycle is optionally substituted with one or more substituents selected from halogen, and C₁₋₆ alkyl.

[illegible]











235. The compound or salt of claim 228, wherein R^1 is

236. The compound or salt of claim 224, wherein M is selected from O.

237. The compound or salt of claim 224, wherein M is selected from NCH_3 and NCH_2CH_3 .

238. The compound or salt of claim 237, wherein M is NMe.

239. The compound or salt of any one of claims 224 to 238, wherein the heterocycle of R^2 is a saturated heterocycle.


240. The compound or salt of claim 239, wherein the heterocycle of R^2 is a saturated heterocycle.

241. The compound or salt of claim 240, wherein R^6 of R^2 is independently selected at each occurrence from halogen, $=CH_2$, hydroxy, C_1 - C_3 hydroxyalkyl, C_1 - C_3 alkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, cyano, and C_1 - C_3 aminoalkyl.

242. The compound or salt of claim 241, wherein L of R^2 is selected from C_1 - C_4 alkylene and




243. The compound or salt of any one of claims 224 to 242, wherein R^2 is selected from



244. The compound or salt of any one of claims 224 to 243, wherein B is selected from an optionally substituted C_9 - C_{10} fused carbocycle.



245. The compound or salt of any one of claims 224 to 244, wherein B is selected from





246. The compound or salt of any one of claims 224 to 245, wherein B is optionally substituted with one or more substituents independently selected from halogen, oxo, $-NH_2$, C_1 - C_3

alkyl, -B(OH)₂, -OH, -O-C₁-C₃ haloalkyl, -C(O)NH₂, -NH₂, =O, -CN, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, and C₂₋₆ alkynyl.

247. The compound or salt of claim 246, wherein B is optionally substituted with one or more substituents independently selected from halogen.

248. The compound or salt of claim 246, wherein B is selected from  and .

249. The compound or salt of claim 248, wherein B is .

250. The compound or salt of claim 248, wherein B is .

251. The compound or salt of claim 224, wherein B is unsubstituted.

252. The compound or salt of claim 224, wherein B is substituted.

253. A pharmaceutical composition comprising a compound or salt of any one of claims 1 to 252 and a pharmaceutically acceptable excipient.

254. A method of treating a disease or disorder, using a compound or salt of any one of claims 1 to 252 or a pharmaceutical composition of claim 253.

255. A method of inhibiting KRas G12D and/or other G12 mutants, using a compound or salt of any one of claims 1 to 252 or a pharmaceutical composition of claim 253.

256. A method of inhibiting KRas G12D and/or other G12 alleles, using a compound or salt of any one of claims 1 to 252 or a pharmaceutical composition of claim 253.

257. A method of inhibiting KRas G12D and/or other alleles, using a compound or salt of any one of claims 1 to 252 or a pharmaceutical composition of claim 253.

258. Use of a compound of any one of Claims 1-252 thereof in the preparation of a target protein degrading compound by using chemical modification of compound of any one of Claims 1-253.

259. A bifunctional compound composed of a target protein-binding moiety and an E3 ubiquitin ligase-binding moiety, wherein the target protein-binding moiety is derived from the compound of any of claims 1-252.