



(11) (21) (C) **2,065,286**
(86) 1990/08/24
(87) 1991/02/26
(45) 2000/05/16

(72) Stanford, John L., GB

(72) Rook, Graham A. W., GB

(73) STANFORD ROOK LIMITED, GB

(51) Int.Cl.⁵ A61K 39/04

(30) 1989/08/25 (8919321.3) GB

(54) **TRAITEMENT DES ETATS INFLAMMATOIRES CHRONIQUES**

(54) **TREATMENT OF CHRONIC INFLAMMATORY CONDITIONS**

(57) L'invention concerne l'emploi d'une matière antigénique et/ou immunorégulatrice dérivée de *Mycobacterium vaccae*, destinée à être utilisée dans la fabrication d'un agent thérapeutique permettant le traitement d'états pathologiques (différents de la tuberculose, de la lèpre ou de la polyarthrite chronique rhumatismale) chez un patient, dans lesquels l'IgG (immunoglobuline de type G) du patient présente une proportion anormalement élevée d'IgG d'agalactosyle, ou dans le traitement d'un trouble inflammatoire chronique (différent de la polyarthrite chronique rhumatismale) provoqué ou accompagné par la libération anormalement élevée par les macrophages d'interleukine-6 et/ou de facteur de nécrose tumorale.

(57) The invention relates to the use of antigenic and/or immunoregulatory material derived from *Mycobacterium vaccae* for use in the manufacture of a therapeutic agent for the treatment of pathological condition (other than tuberculosis, leprosy or rheumatoid arthritis) in a patient in which the patient's IgG shows an abnormally high proportion of agalactosyl IgG or in the treatment of a chronic inflammatory disorder (other than rheumatoid arthritis) caused or accompanied by an abnormally high release by macrophages of interleukin-6 and/or tumour necrosis factor.



ABSTRACTTREATMENT OF CHRONIC INFLAMMATORY CONDITIONS

The invention relates to the use of antigenic and/or immunoregulatory material derived from Mycobacterium
5 vaccae for use in the manufacture of a therapeutic agent for the treatment of pathological condition (other than tuberculosis, leprosy or rheumatoid arthritis) in a patient in which the patient's IgG shows an abnormally high proportion of agalactosyl IgG or in the treatment of a
10 chronic inflammatory disorder (other than rheumatoid arthritis) caused or accompanied by an abnormally high release by macrophages of interleukin-6 and/or tumour necrosis factor.

2065286

2065286 September 1991

- 1 -

11 09 91

TREATMENT OF CHRONIC INFLAMMATORY CONDITIONS

This invention relates to the treatment of chronic inflammatory conditions, e.g. psoriasis.

British Specification No. 2156673 (International Patent Specification WO85/03639) describes immunotherapeutic agents comprising killed cells of Mycobacterium vaccae. These agents are useful in the immunotherapy of mycobacterial disease, especially tuberculosis and leprosy. It is stated that use of this immunotherapeutic agent facilitates the removal of the persisting bacilli responsible for tuberculosis or leprosy which, as is well known, it is difficult to remove by chemotherapy alone. It is suggested in the specification that the immunotherapeutic agent is believed to act by presenting the "protective" common mycobacterial antigens to advantage and by containing immune suppressor determinants which are active in regulating disadvantageous immune mechanisms. As a consequence, "persister" bacilli are recognized by the immune system by their content of common mycobacterial antigens and effective immune mechanisms are directed against them, in the absence of the tissue necrotic form of immunity usually present in mycobacterial disease.

International Patent Specification PCT/GB85/00183 (WO85/05034) describes compositions for the alleviation of the symptoms of, and for the treatment or diagnosis of, arthritic disease which comprise as active ingredient the whole organism of M. vaccae. It is stated that the

11 September 1991

2065286

- 2 -

11 09 91

preparations of M. vaccae are useful for the treatment of various autoimmune diseases and especially arthritic conditions including rheumatoid arthritis, ankylosing spondylitis or Reiter's syndrome.

5 We have now discovered that compositions comprising antigenic and immunoregulatory material derived from Mycobacterium vaccae are generally useful in the treatment of pathological conditions in which the proportion of agalactosyl IgG (i.e. IgG which lacks terminal galactose
10 from the N-linked oligosaccharides on the heavy chains) is increased. Diseases of this kind include not only the arthritic disease e.g. rheumatoid arthritis, and mycobacterial disease, e.g. tuberculosis and leprosy, mentioned in the specifications referred to above, but also
15 Crohn's disease. Other diseases in which this may play a part but in which an increased level of agalactosyl IgG is not easily detectable by current methods include primary biliary cirrhosis, sarcoidosis, ulcerative colitis, psoriasis, systemic lupus erythematosus (especially when
20 accompanied by Sjogren's syndrome), multiple sclerosis, Guillain-Barré syndrome, primary diabetes mellitus, and perhaps some aspects of graft rejection.

Such diseases may also be described as that class of chronic inflammatory disorder which is caused by, or
25 accompanied by, abnormally high release by macrophages of the cytokines interleukin-6 and/or tumour necrosis factor

2065286

11 September 1991

- 3 -

11 09 91

(cachectin). The specific conditions involved are, of course, the same as those already named.

The present invention accordingly provides a method for the treatment of a pathological condition (other than mycobacterial disease, e.g. tuberculosis or leprosy, and arthritic disease, e.g. rheumatoid arthritis, mentioned in the specifications referred to above) in a patient in which the patient's IgG shows an abnormally high proportion of agalactosyl IgG which comprises administering to the patient suffering from such a condition an effective amount of a therapeutic composition comprising antigenic and immunoregulatory material derived from Mycobacterium vaccae.

The invention also provides a method for the treatment of a chronic inflammatory disorder (other than an arthritic disease, e.g. rheumatoid arthritis) caused or accompanied by an abnormally high release from macrophages of interleukin-6 and/or tumour necrosis factor which comprises administering to a patient suffering from such a disorder an effective amount of the said therapeutic agent.

The invention further provides antigenic and immunoregulatory material derived from M. vaccae for use in the manufacture of a therapeutic agent for the treatment of pathological conditions (other than mycobacterial disease or arthritic disease) in a patient whose IgG shows an abnormally high proportion of agalactosyl IgG. Such antigenic and immunoregulatory material is also provided for use in the manufacture of a therapeutic agent for use in the treatment of a chronic inflammatory disorder (other than arthritic

11 September 09 1991 91

- 4 - 2065286

disease) of the kind mentioned above.

The therapeutic agent of the invention conveniently, and therefore preferably, comprises dead cells of M. vaccae, most preferably cells which have been killed by autoclaving 5 or by irradiation. The therapeutic agent normally comprises more than 10^8 to 10^{11} killed M. vaccae microorganisms per ml of diluent.

The diluent may be pyrogen-free saline for injection alone, or a borate buffer of pH 8.0. The diluent should be 10 sterile. A suitable borate buffer is:

	Na ₂ B ₄ O ₇ .10H ₂ O	3.63 g
	H ₃ BO ₃	5.25 g
	NaCl	6.19 g
	Tween 80	0.0005%
15	Distilled Water	to 1 litre

The preferred strain of M. vaccae is one denoted R877R isolated from mud samples from the Lango district of Central Uganda (J.L. Stanford and R.C. Paul, Ann. Soc. Belge Med, Trop. 1973, 53 141-389). The strain is a stable rough 20 variant and belongs to the aurum sub-species. It can be identified as belonging to M. vaccae by biochemical and antigenic criteria (R. Bonicke, S.E. Juhasz., Zentr abbl. Bakteriол. Parasitenkd. Infection skr. Hyg. Abt. 1, Orig., 1964, 192, 133).

United Kingdom Patent Office
PCT International Application

SUBSTITUTE SHEET

-5-

The strain denoted R877R has been deposited under the Budapest Convention at the National Collection of Type Cultures (NCTC) Central Public Health Laboratory, Colindale Avenue, London NW9 5HT, United Kingdom on February 13th, 1984 under the number NCTC 11659.

For the preparation of the therapeutic agent, the microorganism M. vaccae may be grown on a suitable solid medium. A modified Sauton's liquid medium is preferred (S.V. Boyden and E. Sorkin., J. Immunol, 1955 75, 15) solidified with agar. Preferably the solid medium contains 1.3% agar. The medium inoculated with the microorganisms is incubated aerobically to enable growth of the microorganisms to take place, generally at 32°C for 10 days. The organisms are harvested, then weighed and suspended in a diluent. The diluent may be unbuffered saline but is preferably borate-buffered and contains a surfactant such as Tween 80™ as described above. The suspension is diluted to give 100 mg of microorganism/ml. For further dilution, borate buffered saline is preferably used so that the suspension contains 10 mg wet weight of microorganisms/ml of diluent. The suspension may then be dispensed into 5 ml. multidose vials. Although the microorganisms in the vials may be killed using irradiation e.g. from ⁶⁰Cobalt at a dose of 2.5 megarads, or by any other means, for example chemically, it is preferred to kill the microorganisms by autoclaving, for example at 10 psi (69 kPa) for 10 minutes (115°- 125°C). It has been discovered, unexpectedly, that autoclaving yields a more

effective preparation than irradiation.

The therapeutic agent is in general administered by injection in a volume in the range 0.1-0.2 ml, preferably 0.1 ml, given intradermally. A single dosage will generally
5 contain from 10^7 to 10^{10} killed M. vaccae microorganisms. It is preferred to administer to patients a single dose containing 10^8 to 10^9 killed M. vaccae. However, the dose may be repeated depending on the condition of the patient.

While the present invention does not depend on the
10 truth of this theory it is believed that the active ingredient in the killed M. vaccae may be the 65 kDa mycobacterial heat shock protein (hsp 65) described by Young et al. "Stress proteins are immune targets in leprosy and tuberculosis", Proc. Natl. Acad. Sci. U.S.A. 85 (1988),
15 pp4267-4270 in a form obtained from M. bovis. The preferred autoclaved M. vaccae cells used in the present invention are believed to provide an effective package of the hsp 65 and other substances in a convenient adjuvant.

Although the therapeutic agent will generally be
20 administered by intradermal injection, other routes, e.g. oral administration, can also be used.

It may be advantageous and is within the scope of the invention to use more than one strain of M. vaccae, and/or to include in the immunoprophylactic agent other
25 mycobacterial antigens. Tuberculin may also be included.

The immunoprophylactic agent may also contain BCG (Bacillus Calmette-Guerin) vaccine, in particular the

2065286

- 7 -

freeze-fried form of the vaccine, to promote its effect.

The therapeutic agent can contain further ingredients such as adjuvants, preservatives, stabilisers etc. It may be supplied in sterile injectable liquid form or in sterile freeze-fried form which is reconstituted prior to use.

M. vaccae may be used as such or as an extract or fractioned portion of the organism to manufacture the therapeutic agents according to the invention.

The following Example illustrates the invention.

EXAMPLE

M. vaccae NCTC 11659 is grown on a solid medium comprising modified Sauton's medium solidified with 1.3% agar. The medium is inoculated with the microorganism and incubated for 10 days at 32°C to enable growth of the microorganism to take place. The microorganisms are then harvested by gently scraping the surface of the agar and weighed (without drying) and suspended in M/15 borate buffered saline at pH8 to give 10 mg of microorganisms/ml of saline. The suspension is dispensed into 5 ml vials, and then autoclaved for 10 minutes at 10 psi (69 kPa) to kill the microorganisms. After cooling, 1/10th volume of tuberculin (at the standard concentration of 2 µg/ml) is added. The therapeutic agent thus produced is stored at 44°C before use. A single dose consists of 0.1 ml of the suspension, which should be shaken vigorously immediately

2065286

- 8 -

before use, containing 1 mg wet weight of M. vaccae and 0.02 μ g of tuberculin. The dose is given by intradermal injection normally over the left deltoid muscle.

Only one dose is normally required. The patient
5 should not receive high dose steroids or other immuno-suppressive therapy. Up to six months may elapse before the beneficial effect becomes apparent.

2065286

11 September 1991

- 9 -

11 09 91

CLAIMS

1. Use of antigenic and/or immunoregulatory material derived from Mycobacterium vaccae for use in the manufacture of a therapeutic agent for the treatment of
5 pathological conditions (other than mycobacterial disease or arthritic disease) in a patient in which the patient's IgG shows an abnormally high proportion of agalactosyl IgG or in the treatment of a chronic inflammatory disorder (other than
10 an arthritic disease) caused or accompanied by an abnormally high release by macrophages of interleukin-6 and/or tumour necrosis factor.
2. The use according to claim 1, wherein the antigenic and/or immunoregulatory material derived from
15 M. vaccae comprises dead cells of M. vaccae.
3. The use according to claim 2, wherein the cells of M. vaccae have been killed by autoclaving.
4. The use according to claim 1, wherein the antigenic and/or immunoregulatory material derived for
20 M. vaccae comprises the 65 kDa heat shock protein.
5. The use according to any one of the preceding claims, wherein the material derived from M. vaccae is derived from the strain as deposited at the National
Collection of Type Cultures (NCTC) Central Public Health Laboratory, Colindale Avenue, London NW9 5HT, United Kingdom
25 on February 13th, 1984 under the number NCTC 11659.
6. The use according to any one of the preceding

claims, wherein the therapeutic agent contains, per dose, antigenic and/or immunoregulatory material from 10^7 to 10^{10} M. vaccae microorganisms.

- 5 7. A commercial product comprising, as active ingredient, antigenic and/or immunoregulatory material derived from Mycobacterium vaccae for use in treatment of a pathological condition (other than mycobacterial disease and arthritic disease) in a patient in which the patient's IgG shows an
- 10 abnormally high proportion of agalactosyl IgG or for the treatment of a chronic inflammatory disorder (other than an arthritic disease) caused or accompanied by an abnormally high release from macrophages of interleukin-6 and/or tumour necrosis factor.
- 15
8. A product according to claim 9, wherein the material derived from M.vaccae is as defined in any one of claims 2 to 6.
- 20 9. A pharmaceutical agent for use in the treatment of a pathological condition other than mycobacterial disease and arthritic disease) in a patient in which the patient's IgG shows an abnormally high proportion of agalactosyl IgG or for the treatment of a chronic inflammatory disorder (other than an
- 25 arthritic disease) caused or accompanied by an abnormally high release from macrophages of interleukin-6 and/or tumour

necrosis factor, which agent comprises antigenic and/or immunoregulatory material derived from Mycobacterium vaccae.

10. An agent according to claim 9, wherein the material
5 derived from M.vaccae is as defined in any one of claims 2 to 6.