



(11)

EP 3 909 611 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
17.11.2021 Bulletin 2021/46

(51) Int Cl.:
A61K 47/18^(2017.01) A61K 39/395^(2006.01)
A61K 9/19^(2006.01)

(21) Application number: 21175309.0

(22) Date of filing: 22.10.2015

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR

Designated Extension States:

BA ME

Designated Validation States:

MA

(30) Priority: 23.10.2014 US 201462067637 P

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
15790759.3 / 3 209 332

(71) Applicant: Amgen Inc.
Thousand Oaks, CA 91320-1799 (US)

(72) Inventors:
• SLOEY, Christopher James
Newbury Park, 91320 (US)
• KANAPURAM, Sekhar
Thousand Oaks, 91320 (US)

(74) Representative: Grünecker Patent- und
Rechtsanwälte
PartG mbB
Leopoldstraße 4
80802 München (DE)

Remarks:


This application was filed on 21-05-2021 as a
divisional application to the application mentioned
under INID code 62.

(54) REDUCING VISCOSITY OF PHARMACEUTICAL FORMULATIONS

(57) A method for reducing the viscosity of a pharmaceutical formulation is provided that utilizes a viscosity-reducing concentration of an excipient selected from the group consisting of the n-acetyl arginine, n-acetyl

lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof in combination with a therapeutic protein. A stable pharmaceutical formulation is also provided.

Figure 1



Description**CROSS REFERENCE TO RELATED APPLICATIONS**

5 [0001] This application claims the benefit of U.S. Provisional Application No. 62/067,637 filed October 23, 2014, which is incorporated in its entirety by reference herein.

BACKGROUND

10 [0002] Pharmaceutically active proteins, such as antibodies, are frequently formulated in liquid solutions, particularly for parenteral injection. For products that need to be administered via a subcutaneous route, for example use in self administration; formulations in delivery volumes greater than 1-2 milliliters are not well tolerated. In such cases highly concentrated protein formulations are desirable to meet the limited dose volume. The high dose and small volume requirements such administration means that the protein therapeutic can reach concentrations of upwards of 100 mg/ml
 15 or more. Highly concentrated protein formulations can pose many challenges to the manufacturability and administration of protein therapeutics. One challenge posed by some highly concentrated protein formulations is increased viscosity. High viscosity formulations are difficult to handle during manufacturing, including at the bulk and filling stages. High viscosity formulations are also difficult to draw into a syringe and inject, making administration to the patient difficult and unpleasant. The need to identify compounds that are useful for reducing viscosity of highly concentrated protein formulations,
 20 to develop methods of reducing the viscosity of such formulations, and to provide pharmaceutical formulations with reduced viscosity are well known in the pharmaceutical industry. The present invention provides such compounds, methods and formulations.

SUMMARY

25 [0003] Provided is an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof at selected concentrations for use in reducing the viscosity of protein formulations. Methods for reducing the viscosity of protein formulations by combining a high concentration therapeutic protein with a viscosity-reducing concentration of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof are provided herein. Also provided is lyophilized powder comprising a therapeutic protein and an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof wherein the excipient is present at a weight: weight concentration effective to reduce viscosity upon reconstitution with a diluent. Also provided is a lyophilized powder comprising a therapeutic protein and a diluent for reconstitution that contains an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof.

30 [0004] Provided herein is a method for reducing the viscosity of a liquid pharmaceutical formulation comprising a therapeutic protein at a concentration of at least 70 mg/ml, comprising the step of combining the therapeutic protein with a viscosity-reducing concentration of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof. In one embodiment the viscosity of the formulation is reduced by at least 5%. In another embodiment the viscosity of the formulation is reduced by at least 10%. In another embodiment the viscosity of the formulation is reduced by at least 20%. In another embodiment the viscosity of the formulation is reduced by at least 30%. In another embodiment the viscosity of the formulation is reduced by at least 40%. In another embodiment the viscosity of the formulation is reduced by at least 50%. In another embodiment the viscosity of the formulation is reduced by at least 60%. In another embodiment the viscosity of the formulation is reduced by at least 70%. In another embodiment the viscosity of the formulation is reduced by at least 80%. In a related embodiment, pharmaceutical formulations produced by such methods are provided.

35 [0005] Also provided is a pharmaceutical composition comprising a therapeutic protein at a concentration of at least 70 mg/mL, and an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof. In one embodiment the concentration of the excipient is from about 5 mM to about 700 mM. In a related embodiment the concentration of the excipient is from about 100 mM to about 400 mM. In another related embodiment the concentration of the excipient is from about 200 mM to about 300 mM. In still another related embodiment the concentration of the excipient is from about 140 mM to about 170 mM. Also provided are such pharmaceutical compositions having a pH between about 4.0 to about 8.0. In a related embodiment the pH is about 4.0 to about 6.0. In a further related embodiment the pH is about 4.8 to about 5.4.

40 [0006] Also provided is a method of preparing a lyophilized powder comprising the step of lyophilizing a pharmaceutical formulation as described above.

45 [0007] Provided herein is a lyophilized powder comprising a therapeutic protein and an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof wherein

the excipient is present at a weight:weight concentration effective to reduce viscosity upon reconstitution with a diluent. In one embodiment the excipient is present at a concentration of between about 100 μ g per mg therapeutic protein to about 1 mg per mg therapeutic protein. In a related embodiment the excipient is present at a concentration between about 200 μ g to about 500 μ g per mg therapeutic protein. In a further related embodiment the excipient is present at a concentration between about 150 μ g to about 250 μ g per mg therapeutic protein. Also provided is a method for reconstituting a lyophilized powder as described above comprising the step of adding a sterile aqueous diluent.

[0008] Also provided are therapeutic proteins that are antibodies. Also provided are formulations or compositions as described above wherein the therapeutic protein is an antibody. In addition, also provided herein is a lyophilized powder as described above wherein the therapeutic protein is an antibody.

10 BRIEF DESCRIPTION OF THE DRAWINGS

[0009]

15 Figure 1 is a bar graph showing the effects of various excipients on the viscosity of an antibody solution. Figure 2 is a bar graph showing the effects of various excipients on the viscosity of an antibody solution.

DETAILED DESCRIPTION

20 **[0010]** Reducing the viscosity of high concentration therapeutic protein formulations is of interest to the pharmaceutical industry. The excipients, n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof, were discovered to reduce the viscosity of such formulations. The invention provides the excipient at selected concentrations for use in reducing the viscosity of protein formulations. Methods for reducing the viscosity of protein formulations by combining the therapeutic protein with a viscosity-reducing concentration of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof is provided herein. Also provided is lyophilized powder comprising a therapeutic protein and an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof, wherein the excipient is present at a weight:weight concentration effective to reduce viscosity upon reconstitution with a diluent.

25 **[0011]** Unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001) and Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992), and Harlow and Lane *Antibodies: A Laboratory Manual* Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1990). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The terminology used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques can be used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

30 **[0012]** All patents and other publications identified are expressly incorporated herein by reference in their entirety for the purpose of describing and disclosing, for example, the methodologies described in such publications that might be used in connection with the described.

35 **[0013]** Zwitterions are characterized as having separate positive and negative charges that result in a net zero charge for the compound. Most amino acids are zwitterions at physiological pH. As disclosed herein, pharmaceutical formulations containing zwitterions, in particular, n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof, were found to generally have lower viscosity than polyol containing formulations while having greater or comparable stability.

40 **[0014]** N-acetyl arginine, n-acetyl lysine, n-acetyl histidine, and n-acetyl proline are modified versions of a naturally-occurring amino acids. N-acetyl arginine, n-acetyl lysine, n-acetyl histidine, and n-acetyl proline include both d and l forms of the amino acids, such as n-acetyl-l arginine, n - acetyl-d arginine, n-acetyl-1 lysine, n-acetyl-d lysine, n-acetyl-1 histidine, n-acetyl-d histidine, n-acetyl-1 proline and n-acetyl-d proline.

45 **[0015]** The terms "polypeptide" and "protein" are used interchangeably herein. Exemplary polypeptides contemplated for use in the stable pharmaceutical formulations of the invention include antibodies, peptibodies, immunoglobulin-like proteins, non-antibody proteins and non-immunoglobulin-like proteins. Analogs of naturally occurring proteins are contemplated for inclusion in formulations of the present invention, including polypeptides with modified glycosylation,

polypeptides without glycosylation (unglycosylated). As used herein, "analogs" refers to an amino acid sequence that has insertions, deletions or substitutions relative to the parent sequence, while still substantially maintaining the biological activity of the parent sequence, as determined using biological assays known to one of skill in the art. The formulations of the invention may also include derivatives of naturally occurring or analog polypeptides which have been chemically modified, for example, to attach water soluble polymers (e.g., pegylated), radionuclides, or other diagnostic or targeting or therapeutic moieties.

[0016] Antibodies may be formulated according to the present invention. As used herein, the term "antibody" includes fully assembled antibodies, monoclonal antibodies (including human, humanized or chimeric antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), maxibody, BiTes, DARTs, and antibody fragments that can bind antigen (e.g., Fab', F'(ab)2, Fv, single chain antibodies, diabodies), comprising complementarity determining regions (CDRs) of the foregoing as long as they exhibit the desired biological activity.

[0017] Peptibodies, molecules comprising an antibody Fc domain attached to at least one antigen-binding peptide, are generally described in PCT publication WO 00/24782. Immunoglobulin-like proteins, members of the immunoglobulin superfamily, contain one or more immunoglobulin-like domains which fold in structures similar to portions of the antibody variable region.

[0018] Proteins, including those that bind to one or more of the following, would be useful in the compositions and methods of the present invention. These include CD proteins including, but not limited to, CD3, CD4, CD8, CD19, CD20, CD22, CD30, and CD34; including those that interfere with receptor binding. HER receptor family proteins, including HER2, HER3, HER4, and the EGF receptor. Cell adhesion molecules, for example, LFA-1, Mol, p150, 95, VLA-4, ICAM-1, VCAM, and alpha v/beta 3 integrin. Growth factors, including but not limited to, vascular endothelial growth factor ("VEGF"), growth hormone, thyroid stimulating hormone, follicle stimulating hormone, luteinizing hormone, growth hormone releasing factor, parathyroid hormone, mullerian-inhibiting substance, human macrophage inflammatory protein (MIP-I -alpha), erythropoietin (EPO), nerve growth factor, such as NGF-beta, platelet-derived growth factor (PDGF), fibroblast growth factors, including, for instance, aFGF and bFGF, epidermal growth factor (EGF), transforming growth factors (TGF), including, among others, TGF- α and TGF- β , including TGF- β 1, TGF- β 2, TGF- β 3, TGF- β 4, or TGF- β 5, insulin-like growth factors-I and -II (IGF-I and IGF-II), des(1-3)-IGF-I (brain IGF-I), and osteoinductive factors. Insulins and insulin-related proteins, including but not limited to insulin, insulin A-chain, insulin B-chain, proinsulin, and insulin-like growth factor binding proteins. Coagulation and coagulation-related proteins, such as, among others, factor VIII, tissue factor, von Willebrands factor, protein C, alpha-1-antitrypsin, plasminogen activators, such as urokinase and tissue plasminogen activator ("t-PA"), bombazine, thrombin, and thrombopoietin; (vii) other blood and serum proteins, including but not limited to albumin, IgE, and blood group antigens. Colony stimulating factors and receptors thereof, including the following, among others, M-CSF, GM-CSF, and G-CSF, and receptors thereof, such as CSF-1 receptor (c-fms). Receptors and receptor-associated proteins, including, for example, flk2/flt3 receptor, obesity (OB) receptor, LDL receptor, growth hormone receptors, thrombopoietin receptors ("TPO-R," "c-mpl"), glucagon receptors, interleukin receptors, interferon receptors, T-cell receptors, stem cell factor receptors, such as c-Kit, and other receptors listed herein. Receptor ligands, including, for example, OX40L, the ligand for the OX40 receptor. Neurotrophic factors, including but not limited to, bone-derived neurotrophic factor (BDNF) and neurotrophin-3, -4, -5, or -6 (NT-3, NT-4, NT-5, or NT-6). Relaxin A-chain, relaxin B-chain, and prorelaxin; interferons and interferon receptors, including for example, interferon- α , - β , and - γ , and their receptors. Interleukins and interleukin receptors, including but not limited to IL-1 to IL-33 and IL-1 to IL-33 receptors, such as the IL-8 receptor, among others. Viral antigens, including but not limited to, an AIDS envelope viral antigen. Lipoproteins, calcitonin, glucagon, atrial natriuretic factor, lung surfactant, tumor necrosis factor-alpha and -beta, enkephalinase, RANTES (regulated on activation normally T-cell expressed and secreted), mouse gonadotropin-associated peptide, DNAse, inhibin, and activin. Integrin, protein A or D, rheumatoid factors, immunotoxins, bone morphogenetic protein (BMP), superoxide dismutase, surface membrane proteins, decay accelerating factor (DAF), AIDS envelope, transport proteins, homing receptors, addressins, regulatory proteins, immunoadhesins, antibodies. Myostatins, TALL proteins, including TALL-1, amyloid proteins, including but not limited to amyloid-beta proteins, thymic stromal lymphopietins ("TSLP"), RANK ligand ("OPGL"), c-kit, TNF receptors, including TNF Receptor Type 1, TRAIL-R2, angiopoietins, and biologically active fragments or analogs or variants of any of the foregoing.

[0019] Exemplary proteins and antibodies include Activase® (Alteplase); alirocumab (anti-PCSK9 monoclonal antibody designated as H1H316P, see U.S.P.N. 8,062,640); Aranesp® (Darbepoetin-alfa), EpoGen® (Epoetin alfa, or erythropoietin); Avonex® (Interferon β -1a); Bexxar® (Tositumomab); Betaseron® (Interferon- β); bococizumab (anti-PCSK9 monoclonal antibody designated as L1L3, see U.S.P.N. 8,080,243); Campath® (Alemtuzumab); Dynepo® (Epoetin delta); Velcade® (bortezomib); MLN0002 (anti- α 4 β 7 mAb); MLN1202 (anti-CCR2 chemokine receptor mAb); Enbrel® (etanercept); Eprex® (Epoetin alfa); Erbitux® (Cetuximab); evolocumab (anti-PCSK9 monoclonal antibody designated as 21B12, see U.S.P.N. 8,030,467); Genotropin® (Somatropin); Herceptin® (Trastuzumab); Humatropo® (somatropin [rDNA origin] for injection); Humira® (Adalimumab); Infergen® (Interferon Alfacon-1); Natrecor® (nesiritide); Kineret® (Anakinra); Leukine® (Sargamostim); LymphoCide® (Epratuzumab); Benlysta™ (Belimumab); Metalyse® (Tenecteplase); Mircera® (methoxy polyethylene glycol-epoetin beta); Mylotarg® (Gemtuzumab ozogamicin); Raptiva® (efalizumab); Cimzia®

(certolizumab pegol); Soliris™ (Eculizumab); Pexelizumab (Anti-C5 Complement); MEDI-524 (Numax®); Lucentis® (Ranibizumab); Edrecolomab (Panorex®); Trabio® (lerdelimumab); TheraCim hr3 (Nimotuzumab); Omnitarg (Pertuzumab, 2C4); Osidem® (IDM-1); OvaRex® (B43.13); Nuvion® (visilizumab); Cantuzumab mertansine (huC242-DMI); NeoRecormon® (Epoetin beta); Neumega® (Oprelvekin); Neulasta® (Pegylated filgastrim, pegylated G-CSF, pegylated hu-Met-G-CSF); Neupogen® (Filgrastim); Orthoclone OKT3® (Muromonab-CD3), Procrit® (Epoetin alfa); Remicade® (Infliximab), Reopro® (Abciximab), Actemra® (anti-IL6 Receptor mAb), Avastin® (Bevacizumab), HuMax-CD4 (zanolimumab), Rituxan® (Rituximab); Tarceva® (Erlotinib); Roferon-A®-(Interferon alfa-2a); Simulect® (Basiliximab); Stelara™ (Ustekinumab); Prexige® (lumiracoxib); Synagis® (Palivizumab); 146B7-CHO (anti-IL15 antibody, see U.S.P.N. 7,153,507), Tysabri® (Natalizumab); Valortim® (MDX-1303, anti-B. anthracis Protective Antigen mAb); ABthrax™; Vectibix® (Panitumumab); Xolair® (Omalizumab), ETI211 (anti-MRSA mAb), IL-I Trap (the Fc portion of human IgG1 and the extracellular domains of both IL-I receptor components (the Type I receptor and receptor accessory protein)), VEGF Trap (Ig domains of VEGFR1 fused to IgG1 Fc), Zenapax® (Daclizumab); Zenapax® (Daclizumab), Zevalin® (Ibritumomab tiuxetan), Zetia (ezetimibe), Atacicept (TACI-Ig), anti- α 4 β 7 mAb (vedolizumab); galiximab (anti-CD80 monoclonal antibody), anti-CD23 mAb (lumiliximab); BR2-Fc (huBR3 / huFc fusion protein, soluble BAFF antagonist); Simponi™ (Golimumab); Mapatumumab (human anti-TRAIL Receptor-1 mAb); Ocrelizumab (anti-CD20 human mAb); HuMax-EGFR (zalutumumab); M200 (Volociximab, anti- α 5 β 1 integrin mAb); MDX-010 (Ipilimumab, anti-CTLA-4 mAb and VEGFR-I (IMC-18F1); anti-BR3 mAb; anti-C. difficile Toxin A and Toxin B C mAbs MDX-066 (CDA-I) and MDX-1388); anti-CD22 dsFv-PE38 conjugates (CAT-3888 and CAT-8015); anti-CD25 mAb (HuMax-TAC); anti-TSLP antibodies (see U.S.P.N. 7,982,016, U.S.P.N. 8232372 and U.S. Publication Application 20090186022); anti-TSLP receptor antibody (see U.S.P.N. 8,101,182); anti-TSLP antibody designated as A5 (see U.S.P.N. 7,982,016); (see anti-CD3 mAb (NI-0401); Adecatumumab (MT201, anti-EpCAM-CD326 mAb); MDX-060, SGN-30, SGN-35 (anti-CD30 mAbs); MDX-1333 (anti- IFNAR); HuMax CD38 (anti-CD38 mAb); anti-CD40L mAb; anti-Cripto mAb; anti-CTGF Idiopathic Pulmonary Fibrosis Phase I Fibrogen (FG-3019); anti-CTLA4 mAb; anti-eotaxin1 mAb (CAT-213); anti-FGF8 mAb; anti-ganglioside GD2 mAb; anti-sclerostin antibodies (see, U.S.P.N. 8,715,663 or U.S.P.N. 7,592,429). Anti-sclerostin antibody designated as Ab-5 (see U.S.P.N. 8,715,663 or U.S.P.N. 7,592,429); anti-ganglioside GM2 mAb; anti-GDF-8 human mAb (MYO-029); anti-GM-CSF Receptor mAb (CAM-3001); anti-HepC mAb (HuMax HepC); MEDT-545, MDX-1103 (anti-TFN α mAb); anti-TGFTR mAb; anti-IGF-IR mAb (HuMax-Inflam); anti-IL12/IL23p40 mAb (Briakinumab); anti-IL-23p19 mAb (LY2525623); anti-IL13 mAb (CAT-354); anti-IL-17 mAb (AIN457); anti-IL2Ra mAb (HuMax-TAC); anti-IL5 Receptor mAb; anti-integrin receptors mAb (MDX-O18, CNTO 95); anti-IP10 Ulcerative Colitis mAb (MDX- 1100); anti-LLY antibody; BMS-66513; anti-Mannose Receptor/hCG β mAb (MDX-1307); anti-mesothelin dsFv-PE38 conjugate (CAT-5001); anti-PD1mAb (MDX-1 106 (ONO- 4538)); anti-PDGFR α antibody (IMC-3G3); anti-TGF β mAb (GC-1008); anti-TRAIL Receptor-2 human mAb (HGS-ETR2); anti-TWEAK mAb; anti-VEGFR/Flt-1 mAb; anti-ZP3 mAb (HuMax-ZP3); NVS Antibody #1; NVS Antibody #2; and an amyloid-beta monoclonal antibody comprising sequences, SEQ ID NO:8 and SEQ ID NO:6 (see U.S.P.N. 7,906,625).

[0020] Exemplary protein concentrations in the formulation may range from about 70 mg/ml to about 300 mg/ml, about 120 mg/ml to about 270 mg/ml, from about 140 mg/ml to about 255 mg/ml, from about 140 mg/ml to about 240 mg/ml, or from about 140 mg/ml to about 220 mg/ml, or alternatively from about 190 mg/ml to about 210 mg/ml. The concentration of protein will depend upon the end use of the pharmaceutical formulation and can be easily determined by a person of skill in the art. Particularly contemplated concentrations of protein are at least about 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, 265, 270, 275, 280, 285, 290, 295 and 300 mg/ml and including all values in between.

[0021] As used herein, "pharmaceutical formulation" is a sterile composition of a pharmaceutically active drug, such as a biologically active protein, that is suitable for parenteral administration (including but not limited to intravenous, intramuscular, subcutaneous, aerosolized, intrapulmonary, intranasal or intrathecal) to a patient in need thereof and includes only pharmaceutically acceptable excipients, diluents, and other additives deemed safe by the Federal Drug Administration or other foreign national authorities. Pharmaceutical formulations include liquid, e.g. aqueous, solutions that may be directly administered, and lyophilized powders which may be reconstituted into solutions by adding a diluent before administration. Specifically excluded from the scope of the term "pharmaceutical formulation" are compositions for topical administration to patients, compositions for oral ingestion, and compositions for parenteral feeding.

[0022] "Shelf life", as used herein, means that the storage period during which an active ingredient such as a therapeutic protein in a pharmaceutical formulation has minimal degradation (e.g., not more than about 5% to 10% degradation) when the pharmaceutical formulation is stored under specified storage conditions, for example, 2-8°C. Techniques for assessing degradation vary depending upon the identity of the protein in the pharmaceutical formulation. Exemplary techniques include size-exclusion chromatography (SEC)-HPLC to detect, e.g., aggregation, reverse phase (RP)-HPLC to detect, e.g. protein fragmentation, ion exchange-HPLC to detect, e.g., changes in the charge of the protein, mass spectrometry, fluorescence spectroscopy, circular dichroism (CD) spectroscopy, Fourier transform infrared spectroscopy (FT-IR), and Raman spectroscopy to detect protein conformational changes. All of these techniques can be used singly or in combination to assess the degradation of the protein in the pharmaceutical formulation and determine the shelf life

of that formulation. The pharmaceutical formulations of the present invention preferably exhibit not more than about 5 to 10% increases in degradation (e.g., fragmentation, aggregation or unfolding) over two years when stored at 2-8°C.

[0023] As used herein, "stable" formulations of biologically active proteins are formulations that exhibit reduced aggregation and/or reduced loss of biological activity of at least 20% upon storage at 2-8°C for at least 2 years compared with a control formula sample, or alternatively which exhibit reduced aggregation and/or reduced loss of biological activity under conditions of thermal stress, e.g. 25°C for 1 week to 12 weeks; 40°C for 1 to 12 weeks; 52°C for 7-8 days, etc.

[0024] As used herein, "viscosity" is a fluid's resistance to flow, and may be measured in units of centipoise (cP) or milliPascal-second (mPa-s), where 1 cP=1 mPa-s, at a given shear rate. Viscosity may be measured by using a viscometer, e.g., Brookfield Engineering Dial Reading Viscometer, model LVT, and AR-G2, TA instruments. Viscosity may be measured using any other methods and in any other units known in the art (e.g. absolute, kinematic or dynamic viscosity), understanding that it is the percent reduction in viscosity afforded by use of the excipients described by the invention that is important. Regardless of the method used to determine viscosity, the percent reduction in viscosity in excipient formulations versus control formulations will remain approximately the same at a given shear rate.

[0025] As used herein, a formulation containing an amount of an excipient effective to "reduce viscosity" (or a "viscosity-reducing" amount or concentration of such excipient) means that the viscosity of the formulation in its final form for administration (if a solution, or if a powder, upon reconstitution with the intended amount of diluent) is at least 5% less than the viscosity of an appropriate control formulation, such as water, buffer, other known viscosity-reducing agents such as salt, etc. and those control formulations, for example, exemplified herein. Excipient-free control formulations might also be used even if they may not be implementable as a therapeutic formulation due to hypotonicity, for instance.

[0026] Similarly, a "reduced viscosity" formulation is a formulation that exhibits reduced viscosity compared to a control formulation.

[0027] Protein therapeutics often need to be given at high concentrations which can result in increased viscosity of the solution. It is highly desirable to provide high concentration formulations that do not exhibit the increased viscosity typically seen with such high protein concentrations.

[0028] High viscosity formulations are difficult to handle during manufacturing, including at the bulk and filling stages. High viscosity formulations are also difficult to draw into a syringe and inject, often necessitating use of lower gauge needles which can be unpleasant for the patient. The addition of n-acetyl-1-arginine to solutions of biologically active protein reduced the viscosity of high concentration protein solutions.

[0029] The use of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof permits a higher concentration of therapeutic proteins to be used in the formulation without as great of an increase in viscosity as is typically seen with other excipients. Thus, the invention provides a method for stabilizing or reducing viscosity of protein formulations by adding an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof in an amount effective to reduce viscosity. The invention also provides reduced viscosity formulations of therapeutic proteins, including antibodies, containing effective amounts or concentrations of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof. Also contemplated are methods of screening one or more formulations, each containing different concentrations of the excipient described herein to identify suitable or optimal concentrations that reduce viscosity. Further provided are methods of preparing a lyophilized powder from reduced viscosity solution formulations of the invention, and methods of reconstituting the lyophilized powders of the invention via addition of a sterile diluent.

[0030] Thus, the present invention provides pharmaceutical formulations containing biologically active polypeptides and viscosity-reducing concentrations of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof. The reduction in viscosity is at least about 5-90% versus control formulations. In one embodiment the reduction in viscosity ranges from about 10-80%. In other exemplary embodiments, the reduction in viscosity is at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80% or 85%.

[0031] Formulations of the invention may optionally include pharmaceutically acceptable salts, buffers, surfactants, other excipients, carriers, diluents, and/or other formulation agents.

[0032] Exemplary pharmaceutically acceptable buffers include acetate (e.g. sodium acetate), succinate (such as sodium succinate), glutamic acid, glutamate, gluconate, histidine, citrate or other organic acid buffers. Exemplary buffer concentration can be from about 1 mM to about 200 mM, or from about 10 mM to about 60 mM, depending, for example, on the buffer and the desired tonicity (e.g. isotonic, hypertonic or hypotonic) of the formulation. Exemplary pHs include from about 4.5 to about 8.0, or from about 4.8 to about 5.5, or from about 4 to 6, or about 5 to 5.5, or about 5, greater than about 5, greater than about 5.5, greater than about 6, or greater than about 6.5.

[0033] Suitable diluents, other excipients, or carriers and other agents include, but are not limited to, antioxidants, coloring, flavoring and diluting agents, emulsifying agents, suspending agents, solvents, fillers, bulking agents, buffers, vehicles, diluents and/or pharmaceutical adjuvants. For example, a suitable vehicle may be, physiological saline solution, citrate buffered saline, or artificial CSF, possibly supplemented with other materials common in compositions for parenter-

al administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Those skilled in the art would readily recognize a variety of buffers that could be used in the compositions, and dosage forms used in the invention. Typical buffers include, but are not limited to pharmaceutically acceptable weak acids, weak bases, or mixtures thereof. Exemplary buffer components are water soluble materials such as phosphoric acid, tartaric acids, 5 lactic acid, succinic acid, citric acid, acetic acid, ascorbic acid, aspartic acid, glutamic acid, or salts thereof. Exemplary salts include inorganic and organic acids, or bases such as metals or amines, in exemplary concentrations such as about 50-200 mM, or 100-200 mM, or about 100 mM, or about 150 mM.

[0034] Other excipients or stabilizers may also be included, for example, sugars (e.g., sucrose, glucose, trehalose, 10 fructose, xylose, mannose, fucose), polyols (e.g., glycerol, mannitol, sorbitol, glycol, inositol), amino acids or amino acid derivatives (e.g., arginine, proline, histidine, lysine, glycine, methionine, etc.) or surfactants (e.g., polysorbate, including polysorbate 20, or polysorbate 80, or poloxamer, including poloxamer 188, TPGS (d-alpha tocopheryl poly-ethylene glycol 1000 succinate)). Exemplary concentrations of surfactant may range from about 0.001% to about 1.0%, or from about 0.003% to about 0.5%. Preservatives may also be included, such as benzyl alcohol, phenol, m-cresol, 15 chlorobutanol or benzethonium Cl, e.g. at concentrations ranging from about 0.1% to about 2%, or from about 0.5% to about 1%.

[0035] One or more other pharmaceutically acceptable carriers, excipients or stabilizers such as those described in Remington's Pharmaceutical Sciences 21st edition, Osol, A. Ed. (2005) may be included in the formulation provided that they do not adversely affect the desired characteristics of the formulation.

[0036] The concentration of the therapeutic protein, such as an antibody, in the formulation will depend upon the end 20 use of the pharmaceutical formulation and can be easily determined by a person of skill in the art.

[0037] Therapeutic proteins that are antagonists are frequently administered at higher concentrations than those that are agonists. Particularly contemplated high concentrations of therapeutic proteins (without taking into account the weight 25 of chemical modifications such as pegylation), including antibodies, are at least about 70, 80, 90, 100, 110, 120, 130, 140, 150, 175, 180, 185, 190, 195 200, 250, 300, 350, 400, 450, or 500 mg/ml, and/or less than about 250 300, 350, 400, 450 or 500 mg/ml. Exemplary high concentrations of therapeutic proteins, such as antibodies, in the formulation 30 may range from at least about 100 mg/ml to about 500 mg/ml. Other protein concentrations (without taking into account the weight of chemical modifications such as pegylation), are also contemplated, e.g., at least about 1, 5, 10, 20, 30, 35, 40, 45, 50, 55, 60, 65 or 70 mg/ml. The invention particularly contemplates formulations and methods in which the 35 concentration of therapeutic protein results in a viscosity of at least 6, 8, 10, 12, 14, 16, 18, 20, 25, 30, 35 cP or higher, such as 100, 125, 150, 175 or 200 cP and the inclusion of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof results in the reduction of the viscosity 40 by 5% or greater. For example, a solution with a viscosity of about 30 cP may be difficult to inject with a standard 27 gauge needle. All references to mg/ml concentration of therapeutic protein, weight of therapeutic protein (mg) or molecular weight of therapeutic protein (kD) herein mean the respective weight of the proteinaceous part of the therapeutic protein, excluding any non-proteinaceous modifications.

[0038] The present invention provides a method of reducing the viscosity of and/or improving stability of a liquid pharmaceutical formulation of a therapeutic protein, by combining the therapeutic protein and a viscosity-reducing amount of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof.

[0039] In exemplary embodiments, the therapeutic protein is at a high protein concentration as described above. In some embodiments, the reduction in viscosity is at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80% or 85% compared to control formulations

[0040] In another aspect, the invention provides liquid solutions comprising a therapeutic protein and an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures 45 thereof wherein the formulations exhibit reduced viscosity relative to control formulations. In exemplary embodiments, the therapeutic protein is at a high protein concentration as described above. In some embodiments, the excipient described herein is present at a viscosity-reducing (weight:volume) concentration. Any of these excipients can be used at concentrations up to their solubility limit. Such solutions may further comprise a sugar or other polyol such as sucrose or sorbitol, or an amino acid such as arginine, proline, histidine, lysine, glycine, methionine, etc. in an amount effective 50 to further improve stability, reduce aggregation, and/or make the formulation isotonic, without significantly increasing viscosity.

[0041] In exemplary embodiments, the concentration of the excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof is at least about 50 mM to about 300 mM, or at least about 100 mM to about 250 mM, or at least about 140 mM to about 200 mM. In exemplary embodiments 55 the concentration of the excipient is at least about 50, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 210, 220, 250, or 300 mM or greater. Other exemplary embodiments include concentrations of excipients effective to make the formulation isotonic, without significantly increasing viscosity. Exemplary concentrations include those at least about 180 mM or greater, in further embodiments the amounts are at least

about 200 mM or greater.

[0042] In another aspect, the invention provides lyophilized protein formulations comprising a therapeutic protein and an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof wherein upon reconstitution with the recommended amount of diluent, the formulations exhibit reduced viscosity relative to control formulations. In exemplary embodiments, the therapeutic protein is at a high protein concentration as described above. In some embodiments, the excipient is present at an amount effective to reduce viscosity upon reconstitution with diluent (weight:weight concentration). Such formulations may further comprise a sugar or other polyol such as sucrose or sorbitol, or at least one amino acid such as arginine, proline, histidine, lysine, glycine, methionine, etc., in an amount effective to further improve stability, reduce aggregation, and/or make the formulation isotonic, without significantly increasing viscosity.

[0043] In exemplary embodiments, the concentration of the excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof is at least about 1 μ g per mg therapeutic protein, up to about 1.0 mg per mg therapeutic protein. In some embodiments, the concentration of excipient is at least about 1, 10, 50, 100, 150, 200, 250, 300, 350, 400, 450, 500 or 550 μ g per mg therapeutic protein. In other exemplary embodiments, the concentration of excipient is up to about 600, 650, 700, 750, 800, 850, 900, 950 or 1000 μ g per mg therapeutic protein.

[0044] In yet another embodiment, the present invention provides a method of preventing self-association of proteins in liquid formulations by using n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof as an excipient in any of the amounts or concentrations described herein. Formulations with improved stability (e.g., reduced aggregation) and shelf-life are also provided.

[0045] The invention also provides a kit comprising a liquid protein formulation of the invention, and instructions for its administration, optionally with a container, syringe and/or other administration device. The invention further provides a kit comprising a lyophilized protein formulation of the invention, optionally in a container, and instructions for its reconstitution and administration, optionally with a vial of sterile diluent, and optionally with a syringe or other administration device. Exemplary containers include vials, tubes, bottles, single or multi-chambered pre-filled syringes, or cartridges. Exemplary administration devices include syringes, with or without needles, infusion pumps, jet injectors, pen devices, transdermal injectors, or other needle-free injector, or an aerosolization device for nasal or pulmonary delivery.

[0046] In another aspect, a method is provided for screening for a viscosity-reducing concentration of an excipient comprising the steps of: (1) assessing the viscosity of a first solution comprising a first concentration of an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof and a therapeutic protein, such as an antibody, (2) assessing the viscosity of a second solution comprising a different second concentration of the excipient and the therapeutic protein, and (3) determining that the first concentration of excipient is more viscosity-reducing than the second concentration of excipient if the first solution is less viscous. Viscosity can be determined, e.g., using an Aries ARG2 Rheometer or a Brookfield RV-DVIII Rheometer.

[0047] Similar methods are provided for screening for an aggregation-reducing or stabilizing concentration of an excipient.

[0048] Stability can be assessed in many ways, including monitoring conformational change over a range of temperatures (thermostability) and/or time periods (shelf-life) and/or after exposure to stressful handling situations (e.g. physical shaking). Stability of formulations containing varying concentrations of formulation components can be measured using a variety of methods. For example, the amount of protein aggregation can be measured by visual observation of turbidity, by measuring absorbance at a specific wavelength, by size exclusion chromatography (in which aggregates of a protein will elute in different fractions compared to the protein in its native active state), HPLC, or other chromatographic methods. Other methods of measuring conformational change can be used, including using differential scanning calorimetry (DSC), e.g. to determine the temperature of denaturation, or circular dichroism (CD), which measures the molar ellipticity of the protein. Fluorescence can also be used to analyze the composition. Fluorescence encompasses the release or absorption of energy in the form of light or heat, and changes in the polar properties of light. Fluorescence emission can be intrinsic to a protein or can be due to a fluorescence reporter molecule. For example, ANS is a fluorescent probe that binds to the hydrophobic pockets of partially unfolded proteins. As the concentration of unfolded protein increases, the number of hydrophobic pockets increases and subsequently the concentration of ANS that can bind increases. This increase in ANS binding can be monitored by detection of the fluorescence signal of a protein sample. Other means for measuring stability can be used and are well known to persons of skill in the art.

[0049] The invention will be more fully understood by reference to the following examples which detail exemplary embodiments of the invention. They should not, however, be construed as limiting the scope of the invention. All citations throughout the disclosure are hereby expressly incorporated by reference.

EXAMPLES

EXAMPLE 1

5 [0050] An IgG2 antibody preparation (140 mg/ml)(Antibody A) was dialyzed against 10mM Sodium Glutamate 1% Sucrose pH 5.2 using Snakeskin® pleated dialysis tubing 10,000 MWCO (Thermo Fisher Scientific, Waltham, MA). 120mg/mL Antibody A in 10mM Sodium Glutamate 1% Sucrose pH 5.2 was filled in 3cc vials at 1mL fill each. Vials were lyophilized using a Virtis lyophilizer. Vials were reconstituted with 450uL of various solutions (water - control or 200mM excipient solutions). Following reconstitution, Antibody A concentrations were determined using SOLO-VPE instrument 10 to be 190 mg/mL. Viscosity was measured using Aries ARG2 rheometer at 25 °C and are recorded in Table 1 below.

Table 1: Excipient formulations of Antibody A, 190 mg/mL at 20°C

Buffer Identification	Excipient
A	200mM Proline
B	200mM Arginine Glutamate
C	200mM n-Acetyl-1 Arginine
D	200mM Guanidinopropionate
E	200mM Taurocyamine
F	200mM n-acetyl-1 Proline
G	200mM n-acetyl-1 Glycine
H	200mM n-acetyl-1 Lysine
I	water control

30 [0051] Figure 1 shows the effects of the various excipients on the viscosity of the antibody solution. The data shows that the tested excipients have reduced viscosity relative to the water control.

EXAMPLE 2

35 [0052] An IgG1 antibody preparation (70 mg/ml)(Antibody B) and an IgG2 antibody preparation (70 mg/ml)(Antibody C) were dialyzed against 20mM Sodium Acetate pH 4.7 and 4.9 respectively using 10,000 MWCO dialysis tubing. Antibody samples were concentrated using 30,000 MWCO centrifugal filters and an Allegra X-12R Centrifuge. Antibody sample concentrations were determined using an Agilent UV/Vis Spectrophotometer. Excipient-containing samples were prepared by spiking a bulk concentrated solution with pre-weighed solid excipients. Viscous samples were transferred 40 using positive displacement pipettes. Viscosities were measured using a Brookfield RV-DV III+ Programmable Rheometer at 25°C and are recorded in Tables 2.1 and 2.2 below.

Table 2.1: Antibody B -210mg/mL

Formulation	Viscosity (cP), 225/s, 25°C
Control No Excipient	26.0
200mM n-Acetyl-L Arginine	14.3

Table 2.2: Antibody C - 140 mg/mL

Formulation	Viscosity (cP) 225/s, 25°C
Control No Excipient	22.7
200mM n-Acetyl-L Arginine	10.0

50 [0053] For the excipient, n-acetyl-1 arginine, the decrease in viscosity of Antibody B is approximately 45%, and the decrease in viscosity of Antibody C is approximately 56%.

EXAMPLE 3

[0054] An IgG2 antibody preparation (140 mg/ml)(Antibody D) was dialyzed against 10mM Sodium Glutamate 1% Sucrose pH 5.2 using Snakeskin® pleated dialysis tubing 10,000 MWCO (Thermo Fisher Scientific, Waltham, MA). 100mg/mL Antibody D in 10mM Sodium Glutamate 1% Sucrose pH 5.2 was filled in 3cc vials at 1mL fill each. Vials were lyophilized using Virtis lyophilizer. Vials were reconstituted with 450uL of either water (control - No excipient) or 10mM Glutamate 200mM n-Acetyl-L Arginine pH 5.2. Antibody D concentrations were determined using SOLO-VPE instrument. Viscosity was measured using Aries ARG2 rheometer at 25 °C and are recorded in Table 3.0 below.

10 Table 3.0: Antibody D - 175 mg/mL

Formulation	Viscosity (cP) 1000/s, 25°C
Control No Excipient	38.7
200mM n-Acetyl-1 Arginine	23.4

15 [0055] For the excipient, n-acetyl-1 arginine, the decrease in viscosity of Antibody D is approximately 40%.

EXAMPLE 4

20 [0056] An IgG2 antibody preparation (140 mg/ml)(Antibody E) was concentrated using ultrafiltration and diafiltration to 200 mg/ml with 6 diafiltration volumes of Buffer A (10mM Glutamate 260mM n-acetyl-1 arginine pH 4.6) or Buffer B (10mM Sodium Glutamate pH 4.6). Final pH of UFDF samples was determined to be 5.2. Antibody E concentrations were determined using SOLO-VPE instrument. Viscosity was measured using Aries ARG2 rheometer at 20 °C, and recorded in Table 4.0 below.

25 Table 4.0: Antibody E - 200 mg/mL

Formulation	Viscosity (cP) 1000/s, 20°C
Antibody E in Buffer B (Control)	62.1
Antibody E in Buffer A (260mM n-acetyl - L-arginine)	28.6

30 [0057] For the excipient, n-acetyl-1 arginine, the decrease in viscosity of Antibody E is approximately 54%.

EXAMPLE 5

35 [0058] An IgG2 antibody preparation (140 mg/ml)(Antibody A) was dialyzed against 10mM Sodium Glutamate 1% Sucrose pH 5.2 using Snakeskin® pleated dialysis tubing 10,000 MWCO (Thermo Fisher Scientific, Waltham, MA). 120mg/mL Antibody A in 10mM Sodium Glutamate 1% Sucrose pH 5.2 was filled in 3cc vials at 1mL fill each. Vials were lyophilized using a Virtis lyophilizer. Vials were reconstituted with 450uL of various solutions (water - control or 200mM excipient solutions) (see Table 5 below). Following reconstitution, Antibody A concentrations were determined using SOLO-VPE instrument to be 190 mg/mL. Viscosity was measured using Aries ARG2 rheometer at 25 °C and are recorded in Table 5 below.

40 Table 5: Excipient formulations of Antibody A, 190 mg/mL at 20°C

Buffer Identification	Excipient
A	200mM Histidine
B	200mM n-Acetyl-L-Histidine
C	200mM n-Acetyl-L Arginine
D	control

50 [0059] Figure 2 shows the effects of the various excipients on the viscosity of the antibody solution. The data shows that the tested excipients have reduced viscosity relative to the control, and the excipients, n-acetyl-1 arginine and n-acetyl-1 histidine, have reduced viscosity relative to the excipient histidine alone.

[0060] The invention is further defined by the following items:

1. A method for reducing the viscosity of a liquid pharmaceutical formulation comprising a therapeutic protein at a concentration of at least 70 mg/ml, comprising the step of combining the therapeutic protein with a viscosity-reducing concentration of an excipient selected from the group consisting of the n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof.
2. The method of item 1 wherein the excipient is n-acetyl arginine.
3. The method of item 1 wherein the excipient is n-acetyl lysine.
4. The method of item 1 wherein the excipient is n-acetyl histidine.
5. The method of item 1 wherein the excipient is n-acetyl proline.
6. The method of item 1 wherein viscosity of the formulation is reduced by at least 25%.
7. The method of item 1 wherein viscosity of the formulation is reduced by at least 50%.
8. The method of item 1 wherein viscosity of the formulation is reduced by at least 80%.
9. A pharmaceutical formulation produced by the method of any of items 1-8.
10. A pharmaceutical composition comprising a therapeutic protein at a concentration of at least 70 mg/mL, and an excipient selected from the group consisting of the n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof.
11. The pharmaceutical composition of item 10, wherein the excipient is n-acetyl arginine.
12. The pharmaceutical composition of item 10, wherein the excipient is n-acetyl lysine.
13. The pharmaceutical composition of item 10, wherein the excipient is n-acetyl histidine.
14. The pharmaceutical composition of item 10, wherein the excipient is n-acetyl proline.
15. The pharmaceutical composition of item 10, wherein the concentration of the excipient is less than about 200 mM.
16. The pharmaceutical composition of any of items 10-15 having a pH between about 4.0 to about 8.0.
17. The pharmaceutical composition of item 16 having a pH of about 4.6 to about 5.4.
18. A method of preparing a lyophilized powder comprising the step of lyophilizing a pharmaceutical composition of any of items 10-17.
19. A lyophilized powder comprising a therapeutic protein and an excipient selected from the group consisting of the n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof, wherein the excipient is present at a weight:weight concentration effective to reduce viscosity upon reconstitution with a diluent.
20. A lyophilized powder of item 19 wherein the excipient is present at a concentration of between about 100 µg per mg therapeutic protein to about 1 mg per mg therapeutic protein.
21. A lyophilized powder of item 19 wherein the excipient is present at a concentration between about 200 µg to about 500 µg per mg therapeutic protein to about 1 mg per mg therapeutic protein.
22. A method for reconstituting a lyophilized powder of any of items 19-21, comprising the step of adding a sterile aqueous diluent.
23. The method of any one of items 1-8, 18 or 22 wherein the therapeutic protein is an antibody.
24. A pharmaceutical formulation or pharmaceutical composition of any of items 9-17 wherein the therapeutic protein is an antibody.
25. A lyophilized powder of any of items 19-21 wherein the therapeutic protein is an antibody.

Claims

1. A pharmaceutical composition comprising an antibody at a concentration of at least 70 mg/mL, and an excipient selected from the group consisting of n-acetyl arginine, n-acetyl lysine, n-acetyl histidine, n-acetyl proline and mixtures thereof at a viscosity-reducing concentration, wherein viscosity is reduced by at least 5% versus a control formulation.
2. The pharmaceutical composition of claim 1, wherein the excipient is n-acetyl arginine.
3. The pharmaceutical composition of claim 1, wherein the excipient is n-acetyl lysine.
4. The pharmaceutical composition of claim 1, wherein the excipient is n-acetyl histidine.
5. The pharmaceutical composition of claim 1, wherein the excipient is n-acetyl proline.
6. The pharmaceutical composition of claim 1 wherein viscosity of the formulation is reduced by at least 25%.
7. The pharmaceutical composition of claim 1 wherein viscosity of the formulation is reduced by at least 50%.

8. The pharmaceutical composition of any of claims 2-7 having a pH between about 4.0 to about 8.0, preferably about 4.6 to about 5.4.

5 9. The pharmaceutical composition of any one of claims 1-8, wherein the concentration of the excipient is at least 200 mM.

10. The pharmaceutical composition of any one of claims 1-8, wherein the concentration of the excipient is 200 mM to 300 mM.

10 11. The pharmaceutical composition of any claims 1-10, wherein the antibody is a monoclonal antibody.

12. The pharmaceutical composition of any claims 1-11, wherein the antibody is at a concentration of 70 mg/ml to 300 mg/ml.

15 13. The pharmaceutical composition of any claims 1-11, wherein the antibody is at a concentration of 120 mg/ml to 270 mg/ml.

14. A method of preparing a lyophilized powder comprising the step of lyophilizing a pharmaceutical composition of any of claims 1-13.

20

25

30

35

40

45

50

55

Figure 1





Figure 2





EUROPEAN SEARCH REPORT

Application Number

EP 21 17 5309

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10	A US 6 767 892 B1 (SHIRLEY BRET A [US] ET AL) 27 July 2004 (2004-07-27) * figure 3; example 3 * -----	1-14	INV. A61K47/18 A61K39/395 A61K9/19
15	A EP 1 977 763 A1 (CHUGAI PHARMACEUTICAL CO LTD [JP]) 8 October 2008 (2008-10-08) * examples 7,8; tables 12,13,25 * -----	1-14	
20	A WO 2011/104381 A2 (NOVO NORDISK AS [DK]; PARSHAD HENRIK [DK]; ENGELUND DORTHE KOT [DK]; G) 1 September 2011 (2011-09-01) * example 2; table 2 * -----	1-14	
25			
30			TECHNICAL FIELDS SEARCHED (IPC)
35			A61K
40			
45			
50	1 The present search report has been drawn up for all claims		
55	Place of search Munich	Date of completion of the search 12 August 2021	Examiner Giménez Miralles, J
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 21 17 5309

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

12-08-2021

10	Patent document cited in search report	Publication date		Patent family member(s)	Publication date
15	US 6767892 B1	27-07-2004	US	6767892 B1	27-07-2004
			US	2004209815 A1	21-10-2004
			US	2006074011 A1	06-04-2006
20	EP 1977763 A1	08-10-2008	AR	058888 A1	27-02-2008
			EP	1977763 A1	08-10-2008
			JP	5231810 B2	10-07-2013
			JP	W02007074880 A1	04-06-2009
25			US	2009291076 A1	26-11-2009
			US	2015284466 A1	08-10-2015
			WO	2007074880 A1	05-07-2007
30	WO 2011104381 A2	01-09-2011	AU	2011219715 A1	16-08-2012
			BR	112012021576 A2	25-10-2016
			CA	2789061 A1	01-09-2011
			CN	103037899 A	10-04-2013
			CN	107496917 A	22-12-2017
35			CN	107693791 A	16-02-2018
			EP	2538973 A2	02-01-2013
			EP	3216462 A2	13-09-2017
			EP	3409289 A2	05-12-2018
			EP	3708190 A1	16-09-2020
40			JP	6742290 B2	19-08-2020
			JP	2013520476 A	06-06-2013
			JP	2016047858 A	07-04-2016
			JP	2018100262 A	28-06-2018
			KR	20120130757 A	03-12-2012
			RU	2012139181 A	10-04-2014
			SI	3409289 T1	31-12-2020
			US	2013028907 A1	31-01-2013
			US	2018000935 A1	04-01-2018
45			WO	2011104381 A2	01-09-2011
50					
55					

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 62067637 [0001]
- WO 0024782 A [0017]
- US 8062640 B [0019]
- US 8080243 B [0019]
- US 8030467 B [0019]
- US 7153507 B [0019]
- US 7982016 B [0019]
- US 8232372 B [0019]
- US 20090186022 A [0019]
- US 8101182 B [0019]
- US 8715663 B [0019]
- US 7592429 B [0019]
- US 7906625 B [0019]

Non-patent literature cited in the description

- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 2001 [0011]
- **AUSUBEL et al.** Current Protocols in Molecular Biology. Greene Publishing Associates, 1992 [0011]
- **HARLOW ; LANE.** Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1990 [0011]
- Remington's Pharmaceutical Sciences. 2005 [0035]