

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
23 September 2004 (23.09.2004)

PCT

(10) International Publication Number
WO 2004/080385 A2

(51) International Patent Classification⁷: **A61K**
(21) International Application Number:
PCT/IL2004/000241
(22) International Filing Date: 14 March 2004 (14.03.2004)
(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
60/453,512 12 March 2003 (12.03.2003) US
60/453,514 12 March 2003 (12.03.2003) US

(71) Applicant (for all designated States except US): **RAP-
PAPORT FAMILY INSTITUTE FOR RESEARCH IN
THE MEDICAL SCIENCES** [IL/IL]; Efron Street, P.O.
Box 9697, Haifa 31 096 (IL).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **KARIN, Nathan**
[IL/IL]; 9 Caspari Stree, Haifa 34 677 (IL).

(74) Agent: **G.E. EHRLICH (1995) LTD.**; 11 Menachem Be-
gin Street, Ramat Gan 52 521 (IL).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS AND METHODS FOR DIAGNOSING AND TREATING AN INFLAMMATION

(57) Abstract: A method of reducing an inflammatory response in a subject is provided. The method comprising providing to a subject in need thereof a therapeutically effective amount of an agent capable of reducing activity and/or expression of a scavenger receptor or of an effector thereof, thereby reducing the inflammatory response in the subject.



WO 2004/080385 A2

COMPOSITIONS AND METHODS FOR DIAGNOSING AND TREATING AN INFLAMMATION

FIELD AND BACKGROUND OF THE INVENTION

5 The present invention relates to compositions and methods for diagnosing and treating inflammation. More particularly, the present invention relates to the use of scavenger receptor inhibitors in treatment of an inflammatory response and to methods of diagnosing an inflammatory response via detection of autoantibodies directed at scavenger receptors.

10 Inflammation is a physiological condition characterized in the acute form by the classical signs of pain, heat, redness, swelling and loss of function. Inflammation often accompanies diseases such as Multiple Sclerosis (MS), osteoarthritis, Inflammatory Bowel Disease (IBD) including Crohn's disease and ulcerative colitis, Rheumatoid Arthritis (RA), atherosclerosis, encephalomyelitis, Alzheimer's disease,
15 stroke, traumatic brain injury, Parkinson's disease and others. In most cases, there is no effective cure for inflammation associated with such disease and existing treatments are palliative and largely fail to control the underlying causes of tissue degradation.

 Scavenger receptors (SRs) are cell surface proteins, which are typically found
20 on macrophages and bind various types of chemically modified lipoproteins (1-3), such as low-density lipoprotein (LDL). This family of trans-membrane receptors which are highly varied in structure are involved in receptor-mediated endocytosis, phagocytosis of apoptotic cells and bacteria, as well as in cell adhesion [Peiser L. et al., Curr. Opin. Immun. 14(1):123-128, 2002]. Since the massive receptor-mediated
25 uptake of cholesterol from modified LDL can convert cultured macrophages into cholesteryl ester-filled foam cells, similar to those found in atherosclerotic plaques, it has been postulated that these receptors also function in deposition of LDL cholesterol of macrophages in artery walls during the initial stages of atherosclerotic plaque formation [1].

30 Scavenger receptors (SRs) are termed as such since they mediate the binding of remarkably wide variety of polyanionic ligands [e.g., modified proteins, sulfated polysaccharides and certain polynucleotides [1, 3, 4]. This property led to the hypothesis that these receptors form a part of an innate immune response by

serving as pattern recognition receptors that bind a wide variety of pathogen components [2-5].

SRB-I (also referred to as SR-BI or CLA-I) is a macrophage scavenger molecule and a receptor for high-density lipoprotein (HDL) [2, 3, 6, 7] that mediates cholesterol uptake from cells [Rigotti A. et al., Curr. Opin. Lipidol., 8:181-8, 1997; Rigotti A. et al., Proc. Natl. Acad. Sci., 94:12610-5, 1997]. SRB-I can also serve as a receptor for non-HDL lipoproteins and appears to play an important role in reverse cholesterol transport. *In vivo* experiments showed that this receptor is important for HDL metabolism in mice, and for the metabolism of LDL and HDL cholesterol in humans [Stang H. et al., J. Biol. Chem. 274:32692-8., 1999; Swarnakar S. et al., J. Biol. Chem. 274:29733-9., 1999]. Studies involving the manipulation of SRB-I gene expression in mice, indicate that its expression protects against atherosclerosis [Kozarsky K. F., and Krieger M., Curr. Opin. Lipidol. 10:491-7., 1999; Ueda Y. et al., J. Biol. Chem. 275:20368-73., 2000; Acton S. L. et al., Mol. Med. Today 5:518-24., 1999]. It was also suggested that HDL and particularly its protein fraction Apo-A1 affect the *in vitro* production of pro-inflammatory mediators by macrophages (8). Among mediators derived from macrophages that propagate inflammation are interleukin 12 (IL-12), TNF- α and possibly IL-6 whereas, TGF- β and IL-10 have predominantly anti-inflammatory effects [Kiefer R. et al., Prog. Neurobiol. 64(2):109-27, 2001]. To date, no direct role for scavenger molecules in modulating inflammatory reactions has been shown or suggested.

The present inventor has previously shown that the immune system can selectively generate autoimmunity to chemokines and other proinflammatory mediators when such a response is beneficial for the host [9, 10, 11, 12, 14, 15]. For example, patients suffering from rheumatoid arthritis (RA) but not osteoarthritis (OA) have significant levels of autoantibodies directed to TNF- α , and therapies that neutralize the function of TNF- α suppress RA but not OA. Studies conducted by the present inventor have shown that selective amplification of these beneficial antibodies by targeted DNA vaccines provided protective immunity in experimental models (9, 10, 11, 12, 14, 15).

While reducing the present invention to practice, the present inventor uncovered elevated levels of autoantibodies to scavenger receptor (SR) in serum samples obtained from subjects having an inflammatory disease and showed that

inhibiting SR function can prevent such diseases by altering the cytokine profile produced by macrophages from pro-inflammatory cytokines to anti-inflammatory cytokines. These findings suggest, for the first time, that scavenger receptors can serve as novel targets for diagnostics and treatment of inflammatory diseases.

SUMMARY OF THE INVENTION

According to one aspect of the present invention there is provided a method of reducing an inflammatory response in a subject, the method comprising providing to a subject in need thereof a therapeutically effective amount of an agent capable of reducing activity and/or expression of a scavenger receptor or of an effector thereof, thereby reducing the inflammatory response in the subject.

According to further features in preferred embodiments of the invention described below, the agent is selected from the group consisting of: (i) an oligonucleotide directed to an endogenous nucleic acid sequence expressing the scavenger receptor or the effector thereof; (ii) a chemical inhibitor directed to the scavenger receptor or the effector thereof; (iii) a neutralizing antibody directed at the scavenger receptor or the effector thereof; and (iv) a non-functional derivative of the scavenger receptor or the effector thereof.

According to still further features in the described preferred embodiments the scavenger receptor is a class A scavenger receptor or a class B scavenger receptor.

According to still further features in the described preferred embodiments the class B scavenger receptor is SR-BI.

According to another aspect of the present invention there is provided use of an agent capable of reducing activity and/or expression of scavenger receptor or of an effector thereof for the manufacture of a medicament for the treatment of inflammatory diseases.

According to yet another aspect of the present invention there is provided an article of manufacture comprising packaging material and a pharmaceutical composition identified for treating inflammatory diseases being contained within the packaging material, the pharmaceutical composition including, as an active ingredient, an agent capable of reducing activity and/or expression of scavenger receptor or of an effector thereof and a pharmaceutically acceptable carrier.

According to still another aspect of the present invention there is provided a method of diagnosing predisposition to, or presence of, an inflammatory disease in a subject, the method comprising detecting anti scavenger receptor antibodies in a biological sample obtained from the subject, wherein a level above a predetermined
5 normal threshold of the anti scavenger receptor antibodies in the biological sample is indicative of the inflammatory disease in the subject.

According to still further features in the described preferred embodiments detecting the anti scavenger receptor antibodies in the biological sample is effected by ELISA, RIA and/or dot blot.

10 According to an additional aspect of the present invention there is provided a humanized antibody having an antigen recognition domain capable of specifically binding a scavenger receptor.

The present invention successfully addresses the shortcomings of the presently known configurations by providing novel compositions and methods containing same
15 for diagnosing and treating an inflammatory response.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable
20 methods and materials are described below. In case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

BRIEF DESCRIPTION OF THE DRAWINGS

25 The invention is herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of the preferred embodiments of the present invention only, and are presented in the cause of providing what is believed to be the most useful and
30 readily understood description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show structural details of the invention in more detail than is necessary for a fundamental understanding of the

invention, the description taken with the drawings making apparent to those skilled in the art how the several forms of the invention may be embodied in practice.

In the drawings:

FIGs. 1a-d are graphs depicting the therapeutic effect of recombinant SRB-I injection on EAE-induced Lewis rats. Figure 1a is a plot graph depicting the daily development of clinical manifestations of EAE in Lewis rats previously subjected to SRB-I vaccination. The plots compare rats (6 Lewis rats in each group) subjected to active induction of EAE, two months following last administration of plasmid DNA encoding SRB-I (opened triangles), plasmid DNA encoding β -actin (opened squares), or PBS (closed squares). Data was obtained by an observer blind to the experimental protocol using discrete scoring (0-4) of EAE clinical manifestations as described in Example 1 of the Examples section. Results are presented as mean maximal score $\pm$ SE. Figure 1b is a bar graph depicting antibody titer in Lewis rats sera in response to administration of recombinant SRB-I. Lewis rats were subjected to repeated administration of plasmid DNA encoding SRB-I, plasmid DNA encoding β -actin, or PBS (none). Two months after the last immunization all rats were subjected to active induction of EAE and sera was taken after 13 days and determined for the development of antibody titer to recombinant SRB-I. Data was obtained by ELISA, in which rat anti-sera, detected the recombinant SRB-I (OR soluble β -ACTIN) coated onto 96 well ELISA plates. Antibodies were labeled with Goat anti-rat IgG alkaline phosphatase conjugated antibody and in the presence of p-Nitrophenyl Phosphate (p-NPP) liquid substrate produces absorbance at 405nm. Data was collected using ELISA reader. Results are presented as log₂-Ab titer $\pm$ SE. Figure 1c is a plot graph depicting specific binding of anti SRB-I antibodies to the native form of SRB-I on activated peritoneal macrophages. The graphs represent flow cytometry analysis of rat activated peritoneal macrophages cells (10^6) incubated for 30 minutes with IgG (0.5 μ g/ml) labeled with anti-rat IgG-FITC, purified from rats that were subjected to plasmid DNA encoding SRB-I (Anti SR-BI) or to plasmid DNA encoding β -actin (control IgG) administration. Data was collected using a FACScalibur. Results are presented as relative cell number. Figure 1d is a plot graph depicting the daily development of clinical manifestations of EAE in Lewis rats subjected to IgG administration. The plots compare rats (6 Lewis rats in each group) subjected to active induction of EAE and then repeated (days 5, 7, 9) administration of 100 μ g/rat

of purified IgG from EAE rats treated with plasmid DNA encoding SRB-I (opened triangles), plasmid DNA encoding β -actin (opened squares), or PBS (closed squares). Data was obtained by an observer blind to the experimental protocol using discrete scoring (0-4) of EAE clinical manifestations as described in Example 1 of the Examples section. Results are presented as mean maximal score $\pm$ SE.

FIGs. 2a-c are graphs depicting the effect of anti SRB-I antibodies on cytokine expression by murine peritoneal macrophages. The plots compare levels of IL-12 (Figure 2a), TNF- α (Figure 2b) and IL-10 (Figure 2c) produced by LPS activated murine peritoneal macrophages supplemented with (0, 10, 50 or 100 μ g/ml) purified anti SRB-I polyclonal autoantibodies (closed squares), or control IgG (open squares) from normal rat serum after 48h incubation. Data was obtained by ELISA using commercial kits. Data was collected using ELISA reader. Results are presented in pg/ml as mean triplicates $\pm$ SE.

FIG. 3 is a graph depicting the daily development of clinical manifestations of EAE in C57BL/6 mice treated with SRB-I DNA vaccine. The plots compare treatment of mice (6 in each group) with SRB-I encoding DNA vaccine (closed squares), empty plasmid (closed circles), or PBS (opened squares) 12 days after the onset of disease. Data was obtained by an observer blind to the experimental protocol using discrete scoring (0-4) of EAE clinical manifestations as described in Example 1 of the Examples section. Results are presented as mean maximal score $\pm$ SE.

FIG. 4 is a graph depicting the daily development of clinical manifestations of EAE in Lewis rats treated with anti SRB-I antibodies. The plots compare treatment of rats (6 Lewis rats in each group) with 100 μ g/ml of monoclonal antibody 5D8 (closed triangle), control murine IgG1 (closed circles), anti SRB-I polyclonal antibody (closed squares), control rat IgG2a purified from EAE rats that have been subjected to an empty plasmid DNA vaccination (open circles) or PBS treated control (open squares). Data was obtained by an observer blind to the experimental protocol using discrete scoring (0-4) of EAE clinical manifestations as described in Example 1 of the Examples section. Results are presented as mean maximal score $\pm$ SE.

FIGs. 5a-d are photomicrographs depicting colon histology sections of colitis induced rats. Figure 5a is a representative histology section of colon of an untreated rat (normal colon). Figure 5b is a representative histology section of a rat colon, 15

days post induction of colitis, subjected to repeated administration of PBS (days 6, 8 and 10 post induction of the disease - control sick rat colon). Figure 5c is a representative histology section of a rat colon, 15 days post induction of colitis, subjected to repeated administration of control IgG (days 6, 8 and 10 post induction of the disease - control IgG treated colon). Figure 5d is a representative histology section of a rat colon, 15 days post induction of colitis, subjected to repeated administration of anti SRB-I polyclonal antibodies (days 6, 8 and 10 post induction of the disease - anti SRB-I treated colon). Photos were obtained using a digital camera (Nikon digital camera DXM1200F) and a light microscopy (Nikon TE2000-S).

10

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is of compositions and methods which can be used for the diagnosis and treatment of inflammation. Specifically, the present invention relates to the use of scavenger receptor inhibitors in treating inflammatory response and to methods of diagnosing inflammatory response via detection of autoantibodies to scavenger receptors in subjects.

The principles and operation of the present invention may be better understood with reference to the drawings and accompanying descriptions.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details set forth in the following description or exemplified by the Examples. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

Diseases and disorders which have significant inflammatory components are ubiquitous. Skin disorders, bowel disorders, certain degenerative neurological disorders, arthritis, autoimmune diseases and other illnesses afflict many patients. The factors underlying these disorders are varied and include infectious agents, autoimmune factors, dietary or environmental factors and genetic factors. In the majority of cases, the causative elements have not been defined and many of the key pathophysiological components have not been elucidated. Accordingly, treatment options for the majority of these diseases is suboptimal.

While reducing the present invention to practice, the present inventor uncovered that subjects suffering from inflammatory disease exhibit elevated levels of autoantibodies to scavenger receptor (SR) and showed that inhibiting SR function can prevent such diseases by altering the cytokine profile produced by macrophages from pro-inflammatory cytokines to anti-inflammatory cytokines. These findings suggest, for the first time, that scavenger receptors are involved in inflammatory response and that such receptors are novel targets for diagnostics and treatment of inflammatory diseases.

Thus, according to one aspect of the present invention there is provided a method of reducing an inflammatory response in a subject.

As used herein the term "reducing" refers to preventing, curing, reversing, attenuating, alleviating, minimizing, suppressing or halting the deleterious effects of an inflammatory response.

As used herein the phrase "inflammatory response" refers to an immune response which results in inflammation, typically occurring as a result of injurious stimuli including infection, burns, trauma, neoplasia, autoimmune signals and exposure to chemicals, heat or cold or any other harmful stimulus. An inflammatory response according to the present invention refers to an acute phase response and a chronic inflammation.

As used herein the term "subject" refers to subject who may benefit from the present invention such as a mammal (e.g., canine, feline, ovine, porcine, equine, bovine, human), preferably a human subject.

The method of this aspect of the present invention is effected by providing to a subject in need thereof a therapeutically effective amount of an agent capable of reducing activity and/or expression of a scavenger receptor or of an effector thereof, thereby reducing the inflammatory response in the subject.

As is described in detail hereinbelow, such an agent can directly reduce activity and/or expression of the scavenger receptor, or alternatively activate endogenous components which in turn reduce activity and/or expression of the scavenger receptor (indirect).

As used herein a "scavenger receptor" refers to a gene product (i.e., RNA or protein) of a scavenger receptor, which is known in the Art. Examples of scavenger receptors include but are not limited to class A scavenger receptors, class B scavenger

receptors and class F scavenger receptors. The scavenger receptor is preferably one which is expressed and displayed by macrophages. Preferably, the scavenger receptor of the present invention is SR-BI, a member of the CD36 family, GenBank Accession No. NP_005496, also known as CLA-I or SRB-I.

5 Scavenger receptor activity refers to cell adhesion activity, transporter activity, apoptotic activity, lipid metabolism activity, signal transduction activity and/or preferably cytokine secretion activity.

An effector of a scavenger receptor refers to an endogenous molecule which up-regulates or activates scavenger receptor activity. For example, an effector can be
10 a modified lipid (e.g., oxidized lipid, glycated lipid, alkylated lipid, nitrated lipid, acetylated lipid), which binds to the scavenger receptor and activates signaling therefrom.

A number of agents can be used in accordance with this aspect of the present invention to reduce the activity or expression of a scavenger receptor or an effector
15 thereof. Depending on the type of molecule utilized, an agent can either be directly administered to the subject or expressed in cells thereof as is further described hereinbelow.

Thus, for example the agent can be a neutralizing antibody which inhibits the activity of a scavenger receptor [such as by binding to the extracellular collagenous
20 domain of SR which plays a role in ligand binding. See Acton (1993) J. Biol. Chem. 268(5):3530-7] or an effector thereof. A scavenger receptor neutralizing antibody is described in Example 6 of the Examples section which follows. Other SR-neutralizing antibodies are known in the art, see for example Frolov (2000) J. Biol. Chem. 275(17): 12769-12780.

25 The term "antibody" refers to whole antibody molecules as well as functional fragments thereof, such as Fab, F(ab')₂, and Fv that are capable of binding with antigenic portions of the target polypeptide. These functional antibody fragments constitute preferred embodiments of the present invention, and are defined as follows:
(1) Fab, the fragment which contains a monovalent antigen-binding fragment of an
30 antibody molecule, can be produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain;

- (2) Fab', the fragment of an antibody molecule that can be obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two Fab' fragments are obtained per antibody molecule;
- (3) (Fab')₂, the fragment of the antibody that can be obtained by treating whole antibody with the enzyme pepsin without subsequent reduction; F(ab')₂ is a dimer of two Fab' fragments held together by two disulfide bonds;
- (4) Fv, defined as a genetically engineered fragment containing the variable region of the light chain and the variable region of the heavy chain expressed as two chains; and
- (5) Single chain antibody ("SCA"), a genetically engineered molecule containing the variable region of the light chain and the variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule as described in, for example, U.S. Patent 4,946,778.

Purification of serum immunoglobulin antibodies (polyclonal antisera) or reactive portions thereof can be accomplished by a variety of methods known to those of skill including, precipitation by ammonium sulfate or sodium sulfate followed by dialysis against saline, ion exchange chromatography, affinity or immunoaffinity chromatography as well as gel filtration, zone electrophoresis, etc. (see Goding in, Monoclonal Antibodies: Principles and Practice, 2nd ed., pp. 104-126, 1986, Orlando, Fla., Academic Press). Under normal physiological conditions antibodies are found in plasma and other body fluids and in the membrane of certain cells and are produced by lymphocytes of the type denoted B cells or their functional equivalent. Antibodies of the IgG class are made up of four polypeptide chains linked together by disulfide bonds. The four chains of intact IgG molecules are two identical heavy chains referred to as H-chains and two identical light chains referred to as L-chains. Additional classes include IgD, IgE, IgA, IgM and related proteins.

Methods of generating and isolating monoclonal antibodies are well known in the art, as summarized for example in reviews such as Tramontano and Schloeder, Methods in Enzymology 178, 551-568, 1989. A recombinant scavenger receptor polypeptide may be used to generate antibodies *in vitro* (see Example 6 of the Examples section which follows). In general, a suitable host animal is immunized with the recombinant polypeptide. Advantageously, the animal host used is a mouse of an inbred strain. Animals are typically immunized with a mixture comprising a solution of the recombinant polypeptide in a physiologically acceptable vehicle, and

any suitable adjuvant, which achieves an enhanced immune response to the immunogen. By way of example, the primary immunization conveniently may be accomplished with a mixture of a solution of the recombinant polypeptide and Freund's complete adjuvant, said mixture being prepared in the form of a water in oil emulsion. Typically the immunization will be administered to the animals intramuscularly, intradermally, subcutaneously, intraperitoneally, into the footpads, or by any appropriate route of administration. The immunization schedule of the immunogen may be adapted as required, but customarily involves several subsequent or secondary immunizations using a milder adjuvant such as Freund's incomplete adjuvant. Antibody titers and specificity of binding to the polypeptide can be determined during the immunization schedule by any convenient method including by way of example radioimmunoassay, or enzyme linked immunosorbant assay, which is known as the ELISA assay. When suitable antibody titers are achieved, antibody-producing lymphocytes from the immunized animals are obtained, and these are cultured, selected and cloned, as is known in the art. Typically, lymphocytes may be obtained in large numbers from the spleens of immunized animals, but they may also be retrieved from the circulation, the lymph nodes or other lymphoid organs. Lymphocytes are then fused with any suitable myeloma cell line, to yield hybridomas, as is well known in the art. Alternatively, lymphocytes may also be stimulated to grow in culture, and may be immortalized by methods known in the art including the exposure of these lymphocytes to a virus, a chemical or a nucleic acid such as an oncogene, according to established protocols. After fusion, the hybridomas are cultured under suitable culture conditions, for example in multi-well plates, and the culture supernatants are screened to identify cultures containing antibodies that recognize the hapten of choice. Hybridomas that secrete antibodies that recognize the recombinant polypeptide are cloned by limiting dilution and expanded, under appropriate culture conditions. Monoclonal antibodies are purified and characterized in terms of immunoglobulin type and binding affinity.

Antibody fragments according to the present invention can be prepared by proteolytic hydrolysis of the antibody or by expression in *E. coli* or mammalian cells (e.g. Chinese hamster ovary cell culture or other protein expression systems) of DNA encoding the fragment.

Antibody fragments can be obtained by pepsin or papain digestion of whole antibodies by conventional methods. For example, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted $F(ab')_2$. This fragment can be further cleaved using a thiol reducing agent, and optionally a blocking group for the sulfhydryl groups resulting from cleavage of disulfide linkages, to produce 3.5S Fab' monovalent fragments. Alternatively, an enzymatic cleavage using pepsin produces two monovalent Fab' fragments and an Fc fragment directly. These methods are described, for example, by Goldenberg, in U.S. Pat. Nos. 4,036,945 and 4,331,647, and references contained therein, which patents are hereby incorporated by reference in their entirety (see also Porter, R. R., *Biochem. J.*, 73: 119-126, 1959). Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical, or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

Fv fragments comprise an association of V_H and V_L chains. This association may be noncovalent, as described in Inbar et al. (*Proc. Nat'l Acad. Sci. USA* 69:2659-62, 1972). Alternatively, the variable chains can be linked by an intermolecular disulfide bond or cross-linked by chemicals such as glutaraldehyde. Preferably, the Fv fragments comprise V_H and V_L chains connected by a peptide linker. These single-chain antigen binding proteins (sFv) are prepared by constructing a structural gene comprising DNA sequences encoding the V_H and V_L domains connected by an oligonucleotide. The structural gene is inserted into an expression vector, which is subsequently introduced into a host cell such as *E. coli*. The recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing sFvs are described, for example, by Whitlow and Filpula, *Methods*, 2: 97-105, 1991; Bird et al., *Science* 242:423-426, 1988; Pack et al., *Bio/Technology* 11:1271-77, 1993; and Ladner et al., U.S. Pat. No. 4,946,778, all of which are hereby incorporated, by reference, in entirety.

Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, for example, by using the polymerase chain reaction

to synthesize the variable region from RNA of antibody-producing cells (see, for example, Larrick and Fry Methods, 2: 106-10, 1991).

Humanized forms of non-human (e.g., murine) antibodies are chimeric molecules of immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues form a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues, which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art (see also Example 6 of the Examples section). Generally, a humanized antibody has one or more amino acid residues introduced into it from a source, which is non-human. These non-human amino acid residues are often referred to as import residues, which are typically taken from an import variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature 332:323-327 (1988); Verhoeven et al., Science, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such humanized antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically

human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)]. Similarly, human monoclonal antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., *Bio/Technology* 10, 779-783 (1992); Lonberg et al., *Nature* 368 856-859 (1994); Morrison, *Nature* 368 812-13 (1994); Fishwild et al., *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

An agent for reducing the activity of a scavenger receptor or an effector thereof can also be a non-functional derivative thereof (i.e., dominant negative). For example, artificial dominant negative molecules of scavenger receptors have been previously described by Acton (1993 *Supra*). Such truncation mutants lack the positively charged collagenous extracellular domain and while retain trimerization, post-translational processing, intracellular transport, surface expression, and stability, are unable to bind ligand and have a dominant negative effects over wild-type receptors [see also Dejager et al. *J Clin Invest.* 1993 Aug;92(2):894-902].

It will be appreciated that when available, naturally occurring non-functional derivatives of the pathway can be used. Thus, for example, the present invention can use the natural inhibitor of SR-A isoform which modifies ligand uptake [see Gough *J Lipid Res.* 1998 Mar;39(3):531-43].

Polypeptides of these non-functional derivatives can be synthesized using solid phase peptide synthesis procedures which are well known in the art and further

described by John Morrow Stewart and Janis Dillaha Young, Solid Phase Peptide Syntheses (2nd Ed., Pierce Chemical Company, 1984). Synthetic peptides can be purified by preparative high performance liquid chromatography [Creighton T. (1983) Proteins, structures and molecular principles. WH Freeman and Co. N.Y.] and the composition of which can be confirmed via amino acid sequencing.

In cases where large amounts of the proteins are desired, they can be generated using recombinant techniques such as described by Bitter et al., (1987) Methods in Enzymol. 153:516-544, Studier et al. (1990) Methods in Enzymol. 185:60-89, Brisson et al. (1984) Nature 310:511-514, Takamatsu et al. (1987) EMBO J. 6:307-311, Coruzzi et al. (1984) EMBO J. 3:1671-1680 and Brogli et al., (1984) Science 224:838-843, Gurley et al. (1986) Mol. Cell. Biol. 6:559-565 and Weissbach & Weissbach, 1988, Methods for Plant Molecular Biology, Academic Press, NY, Section VIII, pp 421-463.

Alternatively, these proteins can encoded expressed within the target cell from an exogenous polynucleotides ligated into a nucleic acid expression construct.

It will be appreciated that the nucleic acid construct can be administered to the individual employing any suitable mode of administration, described hereinbelow (i.e., *in vivo* gene therapy). Alternatively, the nucleic acid construct is introduced into a suitable cell via an appropriate gene delivery vehicle/method (transfection, transduction, homologous recombination, etc.) and an expression system as needed and then the modified cells are expanded in culture and returned to the individual (i.e., *ex vivo* gene therapy).

To enable cellular expression of the polynucleotides of the present invention, the nucleic acid construct of the present invention further includes at least one cis acting regulatory element. As used herein, the phrase "cis acting regulatory element" refers to a polynucleotide sequence, preferably a promoter, which binds a trans acting regulator and regulates the transcription of a coding sequence located downstream thereto.

Any suitable promoter sequence can be used by the nucleic acid construct of the present invention.

Preferably, the promoter utilized by the nucleic acid construct of the present invention is active in the specific cell population transformed. Examples of cell type-specific and/or tissue-specific promoters include promoters such as albumin that is

liver specific [Pinkert et al., (1987) Genes Dev. 1:268-277], lymphoid specific promoters [Calame et al., (1988) Adv. Immunol. 43:235-275]; in particular promoters of T-cell receptors [Winoto et al., (1989) EMBO J. 8:729-733] and immunoglobulins; [Banerji et al. (1983) Cell 33:729-740], neuron-specific promoters such as the neurofilament promoter [Byrne et al. (1989) Proc. Natl. Acad. Sci. USA 86:5473-5477], pancreas-specific promoters [Edlun et al. (1985) Science 230:912-916] or mammary gland-specific promoters such as the milk whey promoter (U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). The nucleic acid construct of the present invention can further include an enhancer, which can be adjacent or distant to the promoter sequence and can function in up regulating the transcription therefrom.

The nucleic acid construct of the present invention preferably further includes an appropriate selectable marker and/or an origin of replication. Preferably, the nucleic acid construct utilized is a shuttle vector, which can propagate both in *E. coli* (wherein the construct comprises an appropriate selectable marker and origin of replication) and be compatible for propagation in cells, or integration in a gene and a tissue of choice. The construct according to the present invention can be, for example, a plasmid, a bacmid, a phagemid, a cosmid, a phage, a virus or an artificial chromosome.

Examples of suitable constructs include, but are not limited to, pcDNA3, pcDNA3.1 (+/-), pGL3, PzeoSV2 (+/-), pDisplay, pEF/myc/cyto, pCMV/myc/cyto each of which is commercially available from Invitrogen Co. (www.invitrogen.com). Examples of retroviral vector and packaging systems are those sold by Clontech, San Diego, Calif., including Retro-X vectors pLNCX and pLXSN, which permit cloning into multiple cloning sites and the transgene is transcribed from CMV promoter. Vectors derived from Mo-MuLV are also included such as pBabe, where the transgene will be transcribed from the 5'LTR promoter.

Currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral or non-viral constructs, such as adenovirus, lentivirus, Herpes simplex I virus, or adeno-associated virus (AAV) and lipid-based systems. Useful lipids for lipid-mediated transfer of the gene are, for example, DOTMA, DOPE, and DC-Chol [Tonkinson et al., Cancer Investigation, 14(1): 54-65 (1996)]. The most preferred constructs for use in gene therapy are viruses, most preferably adenoviruses,

AAV, lentiviruses, or retroviruses. A viral construct such as a retroviral construct includes at least one transcriptional promoter/enhancer or locus-defining element(s), or other elements that control gene expression by other means such as alternate splicing, nuclear RNA export, or post-translational modification of messenger. Such vector constructs also include a packaging signal, long terminal repeats (LTRs) or portions thereof, and positive and negative strand primer binding sites appropriate to the virus used, unless it is already present in the viral construct. In addition, such a construct typically includes a signal sequence for secretion of the peptide from a host cell in which it is placed. Preferably, the signal sequence for this purpose is a mammalian signal sequence or the signal sequence of the polypeptide variants of the present invention. Optionally, the construct may also include a signal that directs polyadenylation, as well as one or more restriction sites and a translation termination sequence. By way of example, such constructs will typically include a 5' LTR, a tRNA binding site, a packaging signal, an origin of second-strand DNA synthesis, and a 3' LTR or a portion thereof. Other vectors can be used that are non-viral, such as cationic lipids, polylysine, and dendrimers.

Alternatively, the agent of this aspect of the present invention can be a chemical, which is designed to specifically inhibit the activity or expression of a scavenger receptor or an effector thereof. An example of a scavenger receptor inhibitor is Pitavastatin [NK-104, Circulation. 2004 Feb 17;109(6):790-6], which down-regulates expression of CD36. Further it is well established that TNF- α regulates scavenger receptor expression through MAPK (e.g., ERK, JNK, and p38). Thus it is suggested that short term inhibition of this pathway down-regulates the receptor expression, while long term treatment with TNF- α is expected to have the same effect [Hsu J Biol Chem. 2000 Dec 29;275(52):41035-48]. Signal transduction factors and inhibitors are available from a number of chemical companies including Calbiochem (San Diego, CA, USA) and Sigma-Aldrich Corp. (St Louis, MO, USA). Another agent capable of reducing the expression of an SR or effectors thereof is a small interfering RNA (siRNA) molecule. RNA interference is a two-step process. the first step, which is termed as the initiation step, input dsRNA is digested into 21-23 nucleotide (nt) small interfering RNAs (siRNA), probably by the action of Dicer, a member of the RNase III family of dsRNA-specific ribonucleases, which processes (cleaves) dsRNA (introduced directly or via a transgene or a virus) in an ATP-

dependent manner. Successive cleavage events degrade the RNA to 19-21 bp duplexes (siRNA), each with 2-nucleotide 3' overhangs [Hutvagner and Zamore Curr. Opin. Genetics and Development 12:225-232 (2002); and Bernstein Nature 409:363-366 (2001)].

5 In the effector step, the siRNA duplexes bind to a nuclease complex to form the RNA-induced silencing complex (RISC). An ATP-dependent unwinding of the siRNA duplex is required for activation of the RISC. The active RISC then targets the homologous transcript by base pairing interactions and cleaves the mRNA into 12 nucleotide fragments from the 3' terminus of the siRNA [Hutvagner and Zamore Curr. Opin. Genetics and Development 12:225-232 (2002); Hammond et al. (2001) Nat. Rev. Gen. 2:110-119 (2001); and Sharp Genes. Dev. 15:485-90 (2001)]. Although the mechanism of cleavage is still to be elucidated, research indicates that each RISC contains a single siRNA and an RNase [Hutvagner and Zamore Curr. Opin. Genetics and Development 12:225-232 (2002)].

15 Because of the remarkable potency of RNAi, an amplification step within the RNAi pathway has been suggested. Amplification could occur by copying of the input dsRNAs which would generate more siRNAs, or by replication of the siRNAs formed. Alternatively or additionally, amplification could be effected by multiple turnover events of the RISC [Hammond et al. Nat. Rev. Gen. 2:110-119 (2001), Sharp Genes. Dev. 15:485-90 (2001); Hutvagner and Zamore Curr. Opin. Genetics and Development 12:225-232 (2002)]. For more information on RNAi see the following reviews Tuschl ChemBiochem. 2:239-245 (2001); Cullen Nat. Immunol. 3:597-599 (2002); and Brantl Biochem. Biophys. Act. 1575:15-25 (2002).

25 Synthesis of RNAi molecules suitable for use with the present invention can be effected as follows. First, an SRB-I mRNA sequence (e.g., GenBank Accession No. NP_005496), for example, is scanned downstream of the AUG start codon for AA dinucleotide sequences. Occurrence of each AA and the 3' adjacent 19 nucleotides is recorded as potential siRNA target sites. Preferably, siRNA target sites are selected from the open reading frame, as untranslated regions (UTRs) are richer in regulatory protein binding sites. UTR-binding proteins and/or translation initiation complexes may interfere with binding of the siRNA endonuclease complex [Tuschl ChemBiochem. 2:239-245]. It will be appreciated though, that siRNAs directed at untranslated regions may also be effective, as demonstrated for GAPDH wherein

siRNA directed at the 5' UTR mediated about 90 % decrease in cellular GAPDH mRNA and completely abolished protein level (www.ambion.com/techlib/tn/91/912.html).

Second, potential target sites are compared to an appropriate genomic database (e.g., human, mouse, rat etc.) using any sequence alignment software, such as the BLAST software available from the NCBI server (www.ncbi.nlm.nih.gov/BLAST/). Putative target sites which exhibit significant homology to other coding sequences are filtered out.

Qualifying target sequences are selected as template for siRNA synthesis. Preferred sequences are those including low G/C content as these have proven to be more effective in mediating gene silencing as compared to those with G/C content higher than 55 %. Several target sites are preferably selected along the length of the target gene for evaluation.

For better evaluation of the selected siRNAs, a negative control is preferably used in conjunction. Negative control siRNA preferably include the same nucleotide composition as the siRNAs but lack significant homology to the genome. Thus, a scrambled nucleotide sequence of the siRNA is preferably used, provided it does not display any significant homology to any other gene.

One example of a siRNA molecule directed at a SR (i.e., p120) is described by Nishikawa Eur J Biochem. 2001 Oct;268(20):5295-9.

Another agent capable of downregulating a SR or an effector thereof is a DNAzyme molecule capable of specifically cleaving an mRNA transcript or DNA sequence of interest. DNAzymes are single-stranded polynucleotides which are capable of cleaving both single and double stranded target sequences (Breaker, R.R. and Joyce, G. Chemistry and Biology 1995;2:655; Santoro, S.W. & Joyce, G.F. Proc. Natl, Acad. Sci. USA 1997;943:4262) A general model (the "10-23" model) for the DNAzyme has been proposed. "10-23" DNAzymes have a catalytic domain of 15 deoxyribonucleotides, flanked by two substrate-recognition domains of seven to nine deoxyribonucleotides each. This type of DNAzyme can effectively cleave its substrate RNA at purine:pyrimidine junctions (Santoro, S.W. & Joyce, G.F. Proc. Natl, Acad. Sci. USA 199; for rev of DNAzymes see Khachigian, LM [Curr Opin Mol Ther 4:119-21 (2002)]).

Examples of construction and amplification of synthetic, engineered DNazymes recognizing single and double-stranded target cleavage sites have been disclosed in U.S. Pat. No. 6,326,174 to Joyce et al. DNazymes of similar design directed against the human Urokinase receptor were recently observed to inhibit
5 Urokinase receptor expression, and successfully inhibit colon cancer cell metastasis *in vivo* (Itoh et al., 20002, Abstract 409, Ann Meeting Am Soc Gen Ther www.asgt.org). In another application, DNazymes complementary to bcr-abl oncogenes were successful in inhibiting the oncogenes expression in leukemia cells, and lessening relapse rates in autologous bone marrow transplant in cases of CML
10 and ALL.

Reducing expression of SR or an effector thereof can also be effected by using an antisense polynucleotide capable of specifically hybridizing with an mRNA transcript encoding the proteins of interest.

Design of antisense molecules which can be used to efficiently downregulate a
15 gene product of interest must be effected while considering two aspects important to the antisense approach. The first aspect is delivery of the oligonucleotide into the cytoplasm of the appropriate cells, while the second aspect is design of an oligonucleotide which specifically binds the designated mRNA within cells in a way which inhibits translation thereof.

The prior art teaches of a number of delivery strategies which can be used to
20 efficiently deliver oligonucleotides into a wide variety of cell types [see, for example, Luft J Mol Med 76: 75-6 (1998); Kronenwett et al. Blood 91: 852-62 (1998); Rajur et al. Bioconj Chem 8: 935-40 (1997); Lavigne et al. Biochem Biophys Res Commun 237: 566-71 (1997) and Aoki et al. (1997) Biochem Biophys Res Commun 231: 540-5
25 (1997)].

In addition, algorithms for identifying those sequences with the highest predicted binding affinity for their target mRNA based on a thermodynamic cycle that accounts for the energetics of structural alterations in both the target mRNA and the oligonucleotide are also available [see, for example, Walton et al. Biotechnol Bioeng
30 65: 1-9 (1999)].

Such algorithms have been successfully used to implement an antisense approach in cells. For example, the algorithm developed by Walton et al. enabled scientists to successfully design antisense oligonucleotides for rabbit β -globin (RBG)

and mouse tumor necrosis factor- α (TNF- α) transcripts. The same research group has more recently reported that the antisense activity of rationally selected oligonucleotides against three model target mRNAs (human lactate dehydrogenase A and B and rat gp130) in cell culture as evaluated by a kinetic PCR technique proved effective in almost all cases, including tests against three different targets in two cell types with phosphodiester and phosphorothioate oligonucleotide chemistries.

In addition, several approaches for designing and predicting efficiency of specific oligonucleotides using an *in vitro* system were also published (Matveeva et al., Nature Biotechnology 16: 1374 - 1375 (1998)).

Several clinical trials have demonstrated safety, feasibility and activity of antisense oligonucleotides. For example, antisense oligonucleotides suitable for the treatment of cancer have been successfully used [Holmund et al., Curr Opin Mol Ther 1:372-85 (1999)], while treatment of hematological malignancies via antisense oligonucleotides targeting c-myc gene, p53 and Bcl-2 had entered clinical trials and had been shown to be tolerated by patients [Gerwitz Curr Opin Mol Ther 1:297-306 (1999)].

More recently, antisense-mediated suppression of human heparanase gene expression has been reported to inhibit pleural dissemination of human cancer cells in a mouse model [Uno et al., Cancer Res 61:7855-60 (2001)].

Thus, the current consensus is that recent developments in the field of antisense technology which, as described above, have led to the generation of highly accurate antisense design algorithms and a wide variety of oligonucleotide delivery systems, enable an ordinarily skilled artisan to design and implement antisense approaches suitable for downregulating expression of known sequences without having to resort to undue trial and error experimentation.

SR-specific antisense molecules have been previously described by Rhainds Biochemistry. 2003 Jun 24;42(24):7527-38; Zingg Arterioscler Thromb Vasc Biol. 2002 Mar 1;22(3):412-7; and Imachi Lab Invest. 2000 Feb;80(2):263-70.

Another agent capable of reducing the expression of scavenger receptor or an effector thereof is a ribozyme molecule capable of specifically cleaving an mRNA transcript encoding this gene product. Ribozymes are being increasingly used for the sequence-specific inhibition of gene expression by the cleavage of mRNAs encoding proteins of interest [Welch et al., Curr Opin Biotechnol. 9:486-96 (1998)]. The

possibility of designing ribozymes to cleave any specific target RNA has rendered them valuable tools in both basic research and therapeutic applications. In the therapeutics area, ribozymes have been exploited to target viral RNAs in infectious diseases, dominant oncogenes in cancers and specific somatic mutations in genetic disorders [Welch et al., Clin Diagn Virol. 10:163-71 (1998)]. Most notably, several ribozyme gene therapy protocols for HIV patients are already in Phase 1 trials. More recently, ribozymes have been used for transgenic animal research, gene target validation and pathway elucidation. Several ribozymes are in various stages of clinical trials. ANGIOZYME was the first chemically synthesized ribozyme to be studied in human clinical trials. ANGIOZYME specifically inhibits formation of the VEGF-r (Vascular Endothelial Growth Factor receptor), a key component in the angiogenesis pathway. Ribozyme Pharmaceuticals, Inc., as well as other firms have demonstrated the importance of anti-angiogenesis therapeutics in animal models. HEPTAZYME, a ribozyme designed to selectively destroy Hepatitis C Virus (HCV) RNA, was found effective in decreasing Hepatitis C viral RNA in cell culture assays (Ribozyme Pharmaceuticals, Incorporated - WEB home page).

An additional method of reducing the expression of an SR gene or effectors thereof in cells is via triplex forming oligonucleotides (TFOs). Recent studies have shown that TFOs can be designed which can recognize and bind to polypurine/polypirimidine regions in double-stranded helical DNA in a sequence-specific manner. These recognition rules are outlined by Maher III, L. J., et al., Science, 1989;245:725-730; Moser, H. E., et al., Science, 1987;238:645-630; Beal, P. A., et al, Science, 1992;251:1360-1363; Cooney, M., et al., Science, 1988;241:456-459; and Hogan, M. E., et al., EP Publication 375408. Modification of the oligonucleotides, such as the introduction of intercalators and backbone substitutions, and optimization of binding conditions (pH and cation concentration) have aided in overcoming inherent obstacles to TFO activity such as charge repulsion and instability, and it was recently shown that synthetic oligonucleotides can be targeted to specific sequences (for a recent review see Seidman and Glazer, J Clin Invest 2003;112:487-94).

In general, the triplex-forming oligonucleotide has the sequence correspondence:

		23		
oligo	3'--A	G	G	T
duplex	5'--A	G	C	T
duplex	3'--T	C	G	A

However, it has been shown that the A-AT and G-GC triplets have the greatest
 5 triple helical stability (Reither and Jeltsch, BMC Biochem, 2002, Sept12, Epub). The
 same authors have demonstrated that TFOs designed according to the A-AT and G-
 GC rule do not form non-specific triplexes, indicating that the triplex formation is
 indeed sequence specific.

Thus for any given sequence a triplex forming sequence may be devised.
 10 Triplex-forming oligonucleotides preferably are at least 15, more preferably 25, still
 more preferably 30 or more nucleotides in length, up to 50 or 100 bp.

Transfection of cells (for example, via cationic liposomes) with TFOs, and
 formation of the triple helical structure with the target DNA induces steric and
 functional changes, blocking transcription initiation and elongation, allowing the
 15 introduction of desired sequence changes in the endogenous DNA and resulting in the
 specific downregulation of gene expression. Examples of such suppression of gene
 expression in cells treated with TFOs include knockout of episomal supFG1 and
 endogenous HPRT genes in mammalian cells (Vasquez et al., Nucl Acids Res.
 1999;27:1176-81, and Puri, et al, J Biol Chem, 2001;276:28991-98), and the
 20 sequence- and target specific downregulation of expression of the Ets2 transcription
 factor, important in prostate cancer etiology (Carbone, et al, Nucl Acid Res.
 2003;31:833-43), and the pro-inflammatory ICAM-1 gene (Besch et al, J Biol Chem,
 2002;277:32473-79). In addition, Vuyisich and Beal have recently shown that
 sequence specific TFOs can bind to dsRNA, inhibiting activity of dsRNA-dependent
 25 enzymes such as RNA-dependent kinases (Vuyisich and Beal, Nuc. Acids Res
 2000;28:2369-74).

Additionally, TFOs designed according to the abovementioned principles can
 induce directed mutagenesis capable of effecting DNA repair, thus providing both
 downregulation and upregulation of expression of endogenous genes (Seidman and
 30 Glazer, J Clin Invest 2003;112:487-94). Detailed description of the design, synthesis
 and administration of effective TFOs can be found in U.S. Patent Application Nos.
 2003 017068 and 2003 0096980 to Froehler et al, and 2002 0128218 and 2002
 0123476 to Emanuele et al, and U.S. Pat. No. 5,721,138 to Lawn.

Additional description of oligonucleotide agents is further provided hereinbelow. It will be appreciated that therapeutic oligonucleotides may further include base and/or backbone modifications which may increase bioavailability therapeutic efficacy and reduce cytotoxicity. Such modifications are described in
5 Younes (2002) Current Pharmaceutical Design 8:1451-1466.

For example, the oligonucleotides of the present invention may comprise heterocyclic nucleosides consisting of purines and the pyrimidines bases, bonded in a 3' to 5' phosphodiester linkage.

Preferably used oligonucleotides are those modified in either backbone,
10 internucleoside linkages or bases, as is broadly described hereinunder.

Specific examples of preferred oligonucleotides useful according to this aspect of the present invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. Oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone, as disclosed in U.S. Pat.
15 NOs: 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050.

Preferred modified oligonucleotide backbones include, for example,
20 phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl phosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and
25 boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms can also be used.

Alternatively, modified oligonucleotide backbones that do not include a
30 phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in

part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts, as disclosed in U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623, 070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439.

Other oligonucleotides which can be used according to the present invention, are those modified in both sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for complementation with the appropriate polynucleotide target. An example for such an oligonucleotide mimetic, includes peptide nucleic acid (PNA). A PNA oligonucleotide refers to an oligonucleotide where the sugar-backbone is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The bases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Other backbone modifications, which can be used in the present invention are disclosed in U.S. Pat. No: 6,303,374.

Oligonucleotides of the present invention may also include base modifications or substitutions. As used herein, "unmodified" or "natural" bases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified bases include but are not limited to other synthetic and natural bases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted

adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further bases include those disclosed in U.S. Pat. No: 3,687,808, those disclosed in The Concise Encyclopedia Of Polymer Science And Engineering, pages 858-859, Kroschwitz, J. I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., Angewandte Chemie, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y. S., Chapter 15, Antisense Research and Applications, pages 289-302, Crooke, S. T. and Lebleu, B. , ed., CRC Press, 1993. Such bases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2 °C. [Sanghvi YS et al. (1993) Antisense Research and Applications, CRC Press, Boca Raton 276-278] and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

As is mentioned hereinabove the agent according to this aspect of the present invention can be also a molecule, which indirectly causes reduction of SR activity or expression.

An example of such an agent is a molecule which promotes specific immunogenic response to a scavenger receptor or an effector thereof in the subject. Such a molecule can be an SR-BI protein, a fragment derived therefrom or a nucleic acid sequence encoding same (see Examples 1-5 of the Examples section which follows). Although such a molecule can be provided to the subject per se, the agent is preferably administered with an immunostimulant in an immunogenic composition.

An immunostimulant may be any substance that enhances or potentiates an immune response (antibody and/or cell-mediated) to an exogenous antigen. Examples of immunostimulants include adjuvants, biodegradable microspheres (e.g., polylactic galactide) and liposomes into which the compound is incorporated (see e.g., U.S. Pat. No. 4,235,877).

Vaccine preparation is generally described in, for example, M. F. Powell and M. J. Newman, eds., "Vaccine Design (the subunit and adjuvant approach)," Plenum Press (NY, 1995).

Illustrative immunogenic compositions may contain DNA encoding a scavenger receptor, such that the protein is generated in situ. The DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Gene delivery techniques are well known in the art, such as those described by Rolland, Crit. Rev. Therap. Drug Carrier Systems 15:143-198, 1998, and references cited therein.

Preferably, the DNA is introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, lentivirus or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., Proc. Natl. Acad. Sci. USA 86:317-321, 1989; Flexner et al., Ann. N.Y. Acad. Sci. 569:86-103, 1989; Flexner et al., Vaccine 8:17-21, 1990; U.S. Pat. Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Pat. No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, Biotechniques 6:616-627, 1988; Rosenfeld et al., Science 252:431-434, 1991; Kolls et al., Proc. Natl. Acad. Sci. USA 91:215-219, 1994; Kass-Eisler et al., Proc. Natl. Acad. Sci. USA 90:11498-11502, 1993; Guzman et al., Circulation 88:2838-2848, 1993; and Guzman et al., Cir. Res. 73:1202-1207, 1993.

Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer et al., Science 259:1745-1749, 1993 and reviewed by Cohen, Science 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

It will be appreciated that an immunogenic composition may comprise both a polynucleotide and a polypeptide component. Such immunogenic compositions may provide for an enhanced immune response.

Any of a variety of immunostimulants may be employed in the immunogenic compositions of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism,

such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, Bortadella pertussis or Mycobacterium tuberculosis derived proteins. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, Mich.); Merck Adjuvant 65 (Merck and Company, Inc., Rahway, N.J.); AS-2 (SmithKline Beecham, Philadelphia, Pa.); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF or interleukin-2,-7, or -12, may also be used as adjuvants.

The adjuvant composition may be designed to induce an immune response predominantly of the Th1 type. High levels of Th1-type cytokines (e.g., IFN- γ , TNF- α , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (e.g., IL-4, IL-5, IL-6 and IL-10) tend to favor the induction of humoral immune responses. Following application of an immunogenic composition as provided herein, the subject will support an immune response that includes Th1- and Th2-type responses. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, see Mosmann and Coffinan, Ann. Rev. Immunol. 7:145-173, 1989.

Preferred adjuvants for use in eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL), together with an aluminum salt. MPL adjuvants are available from Corixa Corporation (Seattle, Wash.; see U.S. Pat. Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555, WO 99/33488 and U.S. Pat. Nos. 6,008,200 and 5,856,462. Immunostimulatory DNA sequences are also described, for example, by Sato et al., Science 273:352, 1996. Another preferred adjuvant is a saponin, preferably QS21 (Aquila Biopharmaceuticals Inc., Framingham, Mass.), which may be used alone or in combination with other adjuvants. For example, an enhanced system involves the

combination of a monophosphoryl lipid A and saponin derivative, such as the combination of QS21 and 3D-MPL as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprise an oil-in-water emulsion and tocopherol. A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Other preferred adjuvants include Montanide ISA 720 (Seppic, France), SAF (Chiron, Calif., United States), ISCOMS (CSL), MF-59 (Chiron), the SBAS series of adjuvants (e.g., SBAS-2 or SBAS-4, available from SmithKline Beecham, Rixensart, Belgium), Detox (Corixa, Hamilton, Mont.), RC-529 (Corixa, Hamilton, Mont.) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described in pending U.S. patent application Ser. Nos. 08/853,826 and 09/074,720.

A delivery vehicle can be employed with the immunogenic composition of the present invention in order to facilitate production of an antigen-specific immune response that targets tumor cells. Delivery vehicles include antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or maintenance of the T cell response, to have anti-tumor effects per se and/or to be immunologically compatible with the receiver (i.e., matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, including tumor and peritumoral tissues, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

Dendritic cells are highly potent APCs (Banchereau and Steinman, Nature 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic antitumor immunity (see Timmerman and Levy, Ann. Rev. Med. 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate in situ, with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency and their ability to activate naive T cell responses. Dendritic cells may, of course, be engineered to express specific cell-surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As an alternative to

dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within an immunogenic composition (see Zitvogel et al., Nature Med. 4:594-600, 1998).

Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, tumor-infiltrating cells, peritumoral tissues-infiltrating cells, lymph nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF- α to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells harvested from peripheral blood, umbilical cord blood or bone marrow may be differentiated into dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF- α , CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce differentiation, maturation and proliferation of dendritic cells.

Dendritic cells are categorized as "immature" and "mature" cells, which allows a simple way to discriminate between two well characterized phenotypes. Immature dendritic cells are characterized as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fc γ receptor and mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (e.g., CD54 and CD11) and costimulatory molecules (e.g., CD40, CD80, CD86 and 4-1BB).

APCs may generally be transfected with a polynucleotide encoding a SR, such that SR-BI, or an immunogenic portion thereof, is expressed on the cell surface. Such transfection may take place *ex vivo*, and a composition comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to the subject, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., Immunology and cell Biology 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the SR polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (e.g., vaccinia,

fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (e.g., a carrier molecule) such as described above. Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

It will be appreciated that selection of agents which are capable of reducing the activity or expression of a scavenger receptor or effectors thereof is preferably effected by examining their effect on at least one of the above-described scavenger receptor activities. Preferably modulation of inflammatory cytokine expression in macrophages, such as described in Example 6 of the Examples section.

The above-described agents for reducing expression or activity of a scavenger receptor or of effectors thereof (i.e., active ingredients) can be provided to the subject *per se*, or as part of a pharmaceutical composition where they are mixed with a pharmaceutically acceptable carrier.

As used herein a "pharmaceutical composition" refers to a preparation of one or more of the active ingredients described herein with other chemical components such as physiologically suitable carriers and excipients. The purpose of a pharmaceutical composition is to facilitate administration of a compound to an organism.

Herein the term "active ingredient" refers to the preparation accountable for the biological effect.

Hereinafter, the phrases "physiologically acceptable carrier" and "pharmaceutically acceptable carrier" which may be interchangeably used refer to a carrier or a diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered compound. An adjuvant is included under these phrases. One of the ingredients included in the pharmaceutically acceptable carrier can be for example polyethylene glycol (PEG), a biocompatible polymer with a wide range of solubility in both organic and aqueous media (Mutter et al. (1979).

Herein the term "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate administration of an active ingredient. Examples, without limitation, of excipients include calcium carbonate, calcium

phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

Techniques for formulation and administration of drugs may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition, which is incorporated herein by reference.

Suitable routes of administration may, for example, include oral, rectal, transmucosal, especially transnasal, intestinal or parenteral delivery, including intramuscular, subcutaneous and intramedullary injections as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternately, one may administer a preparation in a local rather than systemic manner, for example, via injection of the preparation directly into a specific region of a patient's body.

Pharmaceutical compositions of the present invention may be manufactured by processes well known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active ingredients into preparations which, can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the active ingredients of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological salt buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for oral ingestion by a patient. Pharmacological preparations for oral use can be made using a solid excipient, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable

auxiliaries if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-
5 cellulose, sodium carbomethylcellulose; and/or physiologically acceptable polymers such as polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose,
10 concentrated sugar solutions may be used which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

15 Pharmaceutical compositions, which can be used orally, include push-fit capsules made of gelatin as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules may contain the active ingredients in admixture with filler such as lactose, binders such as starches, lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the
20 active ingredients may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for the chosen route of administration.

For buccal administration, the compositions may take the form of tablets or
25 lozenges formulated in conventional manner.

For administration by nasal inhalation, the active ingredients for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from a pressurized pack or a nebulizer with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichloro-
30 tetrafluoroethane or carbon dioxide. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in a dispenser may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The preparations described herein may be formulated for parenteral administration, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multidose containers with optionally, an added preservative. The compositions may be suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical compositions for parenteral administration include aqueous solutions of the active preparation in water-soluble form. Additionally, suspensions of the active ingredients may be prepared as appropriate oily or water based injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acids esters such as ethyl oleate, triglycerides or liposomes. Aqueous injection suspensions may contain substances, which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the active ingredients to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water based solution, before use. The preparation of the present invention may also be formulated in rectal compositions such as suppositories or retention enemas, using, e.g., conventional suppository bases such as cocoa butter or other glycerides.

Pharmaceutical compositions suitable for use in context of the present invention include compositions wherein the active ingredients are contained in an amount effective to achieve the intended purpose. More specifically, a therapeutically effective amount means an amount of active ingredients effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. Determination of a therapeutically effective amount is well within the capability of those skilled in the art.

For any preparation used in the methods of the invention, the therapeutically effective amount or dose can be estimated initially from *in vitro* assays. For example, a dose can be formulated in animal models and such information can be used to more accurately determine useful doses in humans.

Toxicity and therapeutic efficacy of the active ingredients described herein can be determined by standard pharmaceutical procedures *in vitro*, in cell cultures or experimental animals. The data obtained from these *in vitro* and cell culture assays and animal studies can be used in formulating a range of dosage for use in human.

5 The dosage may vary depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g., Fingl, et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1).

10 Depending on the severity and responsiveness of the condition to be treated, dosing can be of a single or a plurality of administrations, with course of treatment lasting from several days to several weeks or until cure is effected or diminution of the disease state is achieved.

The amount of a composition to be administered will, of course, be dependent on the subject being treated, the severity of the affliction, the manner of administration, the judgment of the prescribing physician, etc.

15 Compositions including the preparation of the present invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

20 Pharmaceutical compositions of the present invention may, if desired, be presented in a pack or dispenser device, such as an FDA approved kit, which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accommodated by a notice associated with the container in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions or human or veterinary administration. Such notice, for example, may be of labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert.

30 A number of diseases and conditions, which typically cause inflammatory response in individuals can be treated using the methodology described hereinabove. Examples of such diseases and conditions are summarized infra.

Inflammatory diseases - Include, but are not limited to, chronic inflammatory diseases and acute inflammatory diseases.

Inflammatory diseases associated with hypersensitivity

Examples of hypersensitivity include, but are not limited to, Type I hypersensitivity, Type II hypersensitivity, Type III hypersensitivity, Type IV hypersensitivity, immediate hypersensitivity, antibody mediated hypersensitivity, immune complex mediated hypersensitivity, T lymphocyte mediated hypersensitivity and DTH.

Type I or immediate hypersensitivity, such as asthma.

Type II hypersensitivity include, but are not limited to, rheumatoid diseases, rheumatoid autoimmune diseases, rheumatoid arthritis (Krenn V. *et al.*, *Histol Histopathol* 2000 Jul;15 (3):791), spondylitis, ankylosing spondylitis (Jan Voswinkel *et al.*, *Arthritis Res* 2001; 3 (3): 189), systemic diseases, systemic autoimmune diseases, systemic lupus erythematosus (Erikson J. *et al.*, *Immunol Res* 1998;17 (1-2):49), sclerosis, systemic sclerosis (Renaudineau Y. *et al.*, *Clin Diagn Lab Immunol.* 1999 Mar;6 (2):156); Chan OT. *et al.*, *Immunol Rev* 1999 Jun;169:107), glandular diseases, glandular autoimmune diseases, pancreatic autoimmune diseases, diabetes, Type I diabetes (Zimmet P. *Diabetes Res Clin Pract* 1996 Oct;34 Suppl:S125), thyroid diseases, autoimmune thyroid diseases, Graves' disease (Orgiazzi J. *Endocrinol Metab Clin North Am* 2000 Jun;29 (2):339), thyroiditis, spontaneous autoimmune thyroiditis (Braley-Mullen H. and Yu S, *J Immunol* 2000 Dec 15;165 (12):7262), Hashimoto's thyroiditis (Toyoda N. *et al.*, *Nippon Rinsho* 1999 Aug;57 (8):1810), myxedema, idiopathic myxedema (Mitsuma T. *Nippon Rinsho.* 1999 Aug;57 (8):1759); autoimmune reproductive diseases, ovarian diseases, ovarian autoimmunity (Garza KM. *et al.*, *J Reprod Immunol* 1998 Feb;37 (2):87), autoimmune anti-sperm infertility (Diekman AB. *et al.*, *Am J Reprod Immunol.* 2000 Mar;43 (3):134), repeated fetal loss (Tincani A. *et al.*, *Lupus* 1998;7 Suppl 2:S107-9), neurodegenerative diseases, neurological diseases, neurological autoimmune diseases, multiple sclerosis (Cross AH. *et al.*, *J Neuroimmunol* 2001 Jan 1;112 (1-2):1), Alzheimer's disease (Oron L. *et al.*, *J Neural Transm Suppl.* 1997;49:77), myasthenia gravis (Infante AJ. And Kraig E, *Int Rev Immunol* 1999;18 (1-2):83), motor neuropathies (Kornberg AJ. *J Clin Neurosci.* 2000 May;7 (3):191), Guillain-Barre syndrome, neuropathies and autoimmune neuropathies (Kusunoki S. *Am J Med Sci.* 2000 Apr;319 (4):234), myasthenic diseases,

- Lambert-Eaton myasthenic syndrome (Takamori M. *Am J Med Sci.* 2000 Apr;319 (4):204), paraneoplastic neurological diseases, cerebellar atrophy, paraneoplastic cerebellar atrophy, non-paraneoplastic stiff man syndrome, cerebellar atrophies, progressive cerebellar atrophies, encephalitis, Rasmussen's encephalitis, amyotrophic lateral sclerosis, Sydeham chorea, Gilles de la Tourette syndrome, polyendocrinopathies, autoimmune polyendocrinopathies (Antoine JC. and Honnorat J. *Rev Neurol (Paris)* 2000 Jan;156 (1):23); neuropathies, dysimmune neuropathies (Nobile-Orazio E. *et al.*, *Electroencephalogr Clin Neurophysiol Suppl* 1999;50:419); neuromyotonia, acquired neuromyotonia, arthrogryposis multiplex congenita (Vincent A. *et al.*, *Ann N Y Acad Sci.* 1998 May 13;841:482), cardiovascular diseases, cardiovascular autoimmune diseases, atherosclerosis (Matsuura E. *et al.*, *Lupus.* 1998;7 Suppl 2:S135), myocardial infarction (Vaarala O. *Lupus.* 1998;7 Suppl 2:S132), thrombosis (Tincani A. *et al.*, *Lupus* 1998;7 Suppl 2:S107-9), granulomatosis, Wegener's granulomatosis, arteritis, Takayasu's arteritis and Kawasaki syndrome (Praprotnik S. *et al.*, *Wien Klin Wochenschr* 2000 Aug 25;112 (15-16):660); anti-factor VIII autoimmune disease (Lacroix-Desmazes S. *et al.*, *Semin Thromb Hemost.* 2000;26 (2):157); vasculitises, necrotizing small vessel vasculitises, microscopic polyangiitis, Churg and Strauss syndrome, glomerulonephritis, pauci-immune focal necrotizing glomerulonephritis, crescentic glomerulonephritis (Noel LH. *Ann Med Interne (Paris).* 2000 May;151 (3):178); antiphospholipid syndrome (Flamholz R. *et al.*, *J Clin Apheresis* 1999;14 (4):171); heart failure, agonist-like β -adrenoceptor antibodies in heart failure (Wallukat G. *et al.*, *Am J Cardiol.* 1999 Jun 17;83 (12A):75H), thrombocytopenic purpura (Moccia F. *Ann Ital Med Int.* 1999 Apr-Jun;14 (2):114); hemolytic anemia, autoimmune hemolytic anemia (Efremov DG. *et al.*, *Leuk Lymphoma* 1998 Jan;28 (3-4):285), gastrointestinal diseases, autoimmune diseases of the gastrointestinal tract, intestinal diseases, chronic inflammatory intestinal disease (Garcia Herola A. *et al.*, *Gastroenterol Hepatol.* 2000 Jan;23 (1):16), celiac disease (Landau YE. and Shoenfeld Y. *Harefuah* 2000 Jan 16;138 (2):122), autoimmune diseases of the musculature, myositis, autoimmune myositis, Sjogren's syndrome (Feist E. *et al.*, *Int Arch Allergy Immunol* 2000 Sep;123 (1):92); smooth muscle autoimmune disease (Zauli D. *et al.*, *Biomed Pharmacother* 1999 Jun;53 (5-6):234), hepatic diseases, hepatic autoimmune diseases, autoimmune hepatitis (Manns MP. *J Hepatol* 2000

Aug;33 (2):326) and primary biliary cirrhosis (Strassburg CP. *et al.*, Eur J Gastroenterol Hepatol. 1999 Jun;11 (6):595).

Type IV or T cell mediated hypersensitivity, include, but are not limited to, rheumatoid diseases, rheumatoid arthritis (Tisch R, McDevitt HO. Proc Natl Acad Sci U S A 1994 Jan 18;91 (2):437), systemic diseases, systemic autoimmune diseases, systemic lupus erythematosus (Datta SK., Lupus 1998;7 (9):591), glandular diseases, glandular autoimmune diseases, pancreatic diseases, pancreatic autoimmune diseases, Type 1 diabetes (Castano L. and Eisenbarth GS. Ann. Rev. Immunol. 8:647); thyroid diseases, autoimmune thyroid diseases, Graves' disease (Sakata S. *et al.*, Mol Cell Endocrinol 1993 Mar;92 (1):77); ovarian diseases (Garza KM. *et al.*, J Reprod Immunol 1998 Feb;37 (2):87), prostatitis, autoimmune prostatitis (Alexander RB. *et al.*, Urology 1997 Dec;50 (6):893), polyglandular syndrome, autoimmune polyglandular syndrome, Type I autoimmune polyglandular syndrome (Hara T. *et al.*, Blood. 1991 Mar 1;77 (5):1127), neurological diseases, autoimmune neurological diseases, multiple sclerosis, neuritis, optic neuritis (Soderstrom M. *et al.*, J Neurol Neurosurg Psychiatry 1994 May;57 (5):544), myasthenia gravis (Oshima M. *et al.*, Eur J Immunol 1990 Dec;20 (12):2563), stiff-man syndrome (Hiemstra HS. *et al.*, Proc Natl Acad Sci U S A 2001 Mar 27;98 (7):3988), cardiovascular diseases, cardiac autoimmunity in Chagas' disease (Cunha-Neto E. *et al.*, J Clin Invest 1996 Oct 15;98 (8):1709), autoimmune thrombocytopenic purpura (Semple JW. *et al.*, Blood 1996 May 15;87 (10):4245), anti-helper T lymphocyte autoimmunity (Caporossi AP. *et al.*, Viral Immunol 1998;11 (1):9), hemolytic anemia (Sallah S. *et al.*, Ann Hematol 1997 Mar;74 (3):139), hepatic diseases, hepatic autoimmune diseases, hepatitis, chronic active hepatitis (Franco A. *et al.*, Clin Immunol Immunopathol 1990 Mar;54 (3):382), biliary cirrhosis, primary biliary cirrhosis (Jones DE. Clin Sci (Colch) 1996 Nov;91 (5):551), nephric diseases, nephric autoimmune diseases, nephritis, interstitial nephritis (Kelly CJ. J Am Soc Nephrol 1990 Aug;1 (2):140), connective tissue diseases, ear diseases, autoimmune connective tissue diseases, autoimmune ear disease (Yoo TJ. *et al.*, Cell Immunol 1994 Aug;157 (1):249), disease of the inner ear (Gloddek B. *et al.*, Ann N Y Acad Sci 1997 Dec 29;830:266), skin diseases, cutaneous diseases, dermal diseases, bullous skin diseases, pemphigus vulgaris, bullous pemphigoid and pemphigus foliaceus.

Examples of delayed type hypersensitivity include, but are not limited to, contact dermatitis and drug eruption.

Examples of types of T lymphocyte mediating hypersensitivity include, but are not limited to, helper T lymphocytes and cytotoxic T lymphocytes.

Examples of helper T lymphocyte-mediated hypersensitivity include, but are not limited to, T_h1 lymphocyte mediated hypersensitivity and T_h2 lymphocyte mediated hypersensitivity.

Autoimmune diseases

Include, but are not limited to, cardiovascular diseases, rheumatoid diseases, glandular diseases, gastrointestinal diseases, cutaneous diseases, hepatic diseases, neurological diseases, muscular diseases, nephric diseases, diseases related to reproduction, connective tissue diseases and systemic diseases.

Examples of autoimmune cardiovascular diseases include, but are not limited to atherosclerosis (Matsuura E. *et al.*, *Lupus*. 1998;7 Suppl 2:S135), myocardial infarction (Vaarala O. *Lupus*. 1998;7 Suppl 2:S132), thrombosis (Tincani A. *et al.*, *Lupus* 1998;7 Suppl 2:S107-9), Wegener's granulomatosis, Takayasu's arteritis, Kawasaki syndrome (Praprotnik S. *et al.*, *Wien Klin Wochenschr* 2000 Aug 25;112 (15-16):660), anti-factor VIII autoimmune disease (Lacroix-Desmazes S. *et al.*, *Semin Thromb Hemost*.2000;26 (2):157), necrotizing small vessel vasculitis, microscopic polyangiitis, Churg and Strauss syndrome, pauci-immune focal necrotizing and crescentic glomerulonephritis (Noel LH. *Ann Med Interne (Paris)*. 2000 May;151 (3):178), antiphospholipid syndrome (Flamholz R. *et al.*, *J Clin Apheresis* 1999;14 (4):171), antibody-induced heart failure (Wallukat G. *et al.*, *Am J Cardiol*. 1999 Jun 17;83 (12A):75H), thrombocytopenic purpura (Moccia F. *Ann Ital Med Int*. 1999 Apr-Jun;14 (2):114; Semple JW. *et al.*, *Blood* 1996 May 15;87 (10):4245), autoimmune hemolytic anemia (Efremov DG. *et al.*, *Leuk Lymphoma* 1998 Jan;28 (3-4):285; Sallah S. *et al.*, *Ann Hematol* 1997 Mar;74 (3):139), cardiac autoimmunity in Chagas' disease (Cunha-Neto E. *et al.*, *J Clin Invest* 1996 Oct 15;98 (8):1709) and anti-helper T lymphocyte autoimmunity (Caporossi AP. *et al.*, *Viral Immunol* 1998;11 (1):9).

Examples of autoimmune rheumatoid diseases include, but are not limited to rheumatoid arthritis (Krenn V. *et al.*, *Histol Histopathol* 2000 Jul;15 (3):791; Tisch R, McDevitt HO. *Proc Natl Acad Sci units S A* 1994 Jan 18;91 (2):437) and ankylosing spondylitis (Jan Voswinkel *et al.*, *Arthritis Res* 2001; 3 (3): 189).

Examples of autoimmune glandular diseases include, but are not limited to, pancreatic disease, Type I diabetes, thyroid disease, Graves' disease, thyroiditis,

spontaneous autoimmune thyroiditis, Hashimoto's thyroiditis, idiopathic myxedema, ovarian autoimmunity, autoimmune anti-sperm infertility, autoimmune prostatitis and Type I autoimmune polyglandular syndrome. diseases include, but are not limited to autoimmune diseases of the pancreas, Type 1 diabetes (Castano L. and Eisenbarth GS. Ann. Rev. Immunol. 8:647; Zimmet P. Diabetes Res Clin Pract 1996 Oct;34 Suppl:S125), autoimmune thyroid diseases, Graves' disease (Orgiazzi J. Endocrinol Metab Clin North Am 2000 Jun;29 (2):339; Sakata S. *et al.*, Mol Cell Endocrinol 1993 Mar;92 (1):77), spontaneous autoimmune thyroiditis (Braley-Mullen H. and Yu S, J Immunol 2000 Dec 15;165 (12):7262), Hashimoto's thyroiditis (Toyoda N. *et al.*, Nippon Rinsho 1999 Aug;57 (8):1810), idiopathic myxedema (Mitsuma T. Nippon Rinsho. 1999 Aug;57 (8):1759), ovarian autoimmunity (Garza KM. *et al.*, J Reprod Immunol 1998 Feb;37 (2):87), autoimmune anti-sperm infertility (Diekman AB. *et al.*, Am J Reprod Immunol. 2000 Mar;43 (3):134), autoimmune prostatitis (Alexander RB. *et al.*, Urology 1997 Dec;50 (6):893) and Type I autoimmune polyglandular syndrome (Hara T. *et al.*, Blood. 1991 Mar 1;77 (5):1127).

Examples of autoimmune gastrointestinal diseases include, but are not limited to, chronic inflammatory intestinal diseases (Garcia Herola A. *et al.*, Gastroenterol Hepatol. 2000 Jan;23 (1):16), celiac disease (Landau YE. and Shoenfeld Y. Harefuah 2000 Jan 16;138 (2):122), colitis, ileitis and Crohn's disease.

Examples of autoimmune cutaneous diseases include, but are not limited to, autoimmune bullous skin diseases, such as, but are not limited to, pemphigus vulgaris, bullous pemphigoid and pemphigus foliaceus.

Examples of autoimmune hepatic diseases include, but are not limited to, hepatitis, autoimmune chronic active hepatitis (Franco A. *et al.*, Clin Immunol Immunopathol 1990 Mar;54 (3):382), primary biliary cirrhosis (Jones DE. Clin Sci (Colch) 1996 Nov;91 (5):551; Strassburg CP. *et al.*, Eur J Gastroenterol Hepatol. 1999 Jun;11 (6):595) and autoimmune hepatitis (Manns MP. J Hepatol 2000 Aug;33 (2):326).

Examples of autoimmune neurological diseases include, but are not limited to, multiple sclerosis (Cross AH. *et al.*, J Neuroimmunol 2001 Jan 1;112 (1-2):1), Alzheimer's disease (Oron L. *et al.*, J Neural Transm Suppl. 1997;49:77), myasthenia gravis (Infante AJ. And Kraig E, Int Rev Immunol 1999;18 (1-2):83; Oshima M. *et al.*, Eur J Immunol 1990 Dec;20 (12):2563), neuropathies, motor neuropathies (Kornberg AJ. J Clin Neurosci. 2000 May;7 (3):191); Guillain-Barre syndrome and autoimmune

neuropathies (Kusunoki S. *Am J Med Sci.* 2000 Apr;319 (4):234), myasthenia, Lambert-Eaton myasthenic syndrome (Takamori M. *Am J Med Sci.* 2000 Apr;319 (4):204); paraneoplastic neurological diseases, cerebellar atrophy, paraneoplastic cerebellar atrophy and stiff-man syndrome (Hiemstra HS. *et al.*, *Proc Natl Acad Sci* 5 units S A 2001 Mar 27;98 (7):3988); non-paraneoplastic stiff man syndrome, progressive cerebellar atrophies, encephalitis, Rasmussen's encephalitis, amyotrophic lateral sclerosis, Sydeham chorea, Gilles de la Tourette syndrome and autoimmune polyendocrinopathies (Antoine JC. and Honnorat J. *Rev Neurol (Paris)* 2000 Jan;156 (1):23); dysimmune neuropathies (Nobile-Orazio E. *et al.*, *Electroencephalogr Clin* 10 *Neurophysiol Suppl* 1999;50:419); acquired neuromyotonia, arthrogryposis multiplex congenita (Vincent A. *et al.*, *Ann N Y Acad Sci.* 1998 May 13;841:482), neuritis, optic neuritis (Soderstrom M. *et al.*, *J Neurol Neurosurg Psychiatry* 1994 May;57 (5):544) and neurodegenerative diseases.

Examples of autoimmune muscular diseases include, but are not limited to, 15 myositis, autoimmune myositis and primary Sjogren's syndrome (Feist E. *et al.*, *Int Arch Allergy Immunol* 2000 Sep;123 (1):92) and smooth muscle autoimmune disease (Zauli D. *et al.*, *Biomed Pharmacother* 1999 Jun;53 (5-6):234).

Examples of autoimmune nephric diseases include, but are not limited to, 20 nephritis and autoimmune interstitial nephritis (Kelly CJ. *J Am Soc Nephrol* 1990 Aug;1 (2):140).

Examples of autoimmune diseases related to reproduction include, but are not limited to, repeated fetal loss (Tincani A. *et al.*, *Lupus* 1998;7 Suppl 2:S107-9).

Examples of autoimmune connective tissue diseases include, but are not limited 25 to, ear diseases, autoimmune ear diseases (Yoo TJ. *et al.*, *Cell Immunol* 1994 Aug;157 (1):249) and autoimmune diseases of the inner ear (Gloddek B. *et al.*, *Ann N Y Acad Sci* 1997 Dec 29;830:266).

Examples of autoimmune systemic diseases include, but are not limited to, 30 systemic lupus erythematosus (Erikson J. *et al.*, *Immunol Res* 1998;17 (1-2):49) and systemic sclerosis (Renaudineau Y. *et al.*, *Clin Diagn Lab Immunol.* 1999 Mar;6 (2):156); Chan OT. *et al.*, *Immunol Rev* 1999 Jun;169:107).

Infectious diseases

Examples of infectious diseases include, but are not limited to, chronic infectious diseases, subacute infectious diseases, acute infectious diseases, viral

diseases, bacterial diseases, protozoan diseases, parasitic diseases, fungal diseases, mycoplasma diseases and prion diseases.

Graft rejection diseases

Examples of diseases associated with transplantation of a graft include, but are not limited to, graft rejection, chronic graft rejection, subacute graft rejection, hyperacute graft rejection, acute graft rejection and graft versus host disease.

Allergic diseases

Examples of allergic diseases include, but are not limited to, asthma, hives, urticaria, pollen allergy, dust mite allergy, venom allergy, cosmetics allergy, latex allergy, chemical allergy, drug allergy, insect bite allergy, animal dander allergy, stinging plant allergy, poison ivy allergy and food allergy.

Cancerous diseases

Examples of cancer include but are not limited to carcinoma, lymphoma, blastoma, sarcoma, and leukemia. Particular examples of cancerous diseases but are not limited to: Myeloid leukemia such as Chronic myelogenous leukemia. Acute myelogenous leukemia with maturation. Acute promyelocytic leukemia, Acute nonlymphocytic leukemia with increased basophils, Acute monocytic leukemia. Acute myelomonocytic leukemia with eosinophilia; Malignant lymphoma, such as Birkitt's Non-Hodgkin's; Lymphocytic leukemia, such as Acute lymphoblastic leukemia. Chronic lymphocytic leukemia; Myeloproliferative diseases, such as Solid tumors Benign Meningioma, Mixed tumors of salivary gland, Colonic adenomas; Adenocarcinomas, such as Small cell lung cancer, Kidney, Uterus, Prostate, Bladder, Ovary, Colon, Sarcomas, Liposarcoma, myxoid, Synovial sarcoma, Rhabdomyosarcoma (alveolar), Extraskelitel myxoid chondrosarcoma, Ewing's tumor; other include Testicular and ovarian dysgerminoma, Retinoblastoma, Wilms' tumor, Neuroblastoma, Malignant melanoma, Mesothelioma, breast, skin, prostate, and ovarian.

In addition to therapeutic advances pioneered by the present invention, the unprecedented findings that antibodies to scavenger receptor are expressed during an inflammation response may be also employed in diagnostic applications (see Examples 1 of the Examples section which follows).

Thus, according to another aspect of the present invention there is provided a method of diagnosing predisposition to, or presence of, an inflammatory response or diseases related therewith (such as described above) in a subject.

As used herein the term "diagnosing" refers to classifying a disease or a symptom as an inflammatory disease, determining a severity of such a disease,
5 monitoring disease progression, forecasting an outcome of a disease and/or prospects of recovery.

The method is effected by detecting autoantibodies to a scavenger molecule in a biological sample obtained from the subject, wherein a level of the autoantibodies
10 above a predetermined threshold (i.e., the level of the same in a biological sample obtained from a healthy individual) is indicative of the disease in the subject.

As used herein a "biological sample" refers to an antibody-containing sample of cell, tissue or fluid derived from the subject. Antibodies present in the sample are typically found within cytoplasmic membrane-bound compartments (e.g.,
15 endoplasmic reticulum and Golgi apparatus) and on the surface of B lymphocytes (which synthesize antibody molecules) and immune effector cells such as, mononuclear phagocytes, natural killer (NK) cells and mast cells, which express specific receptors for binding antibody molecules. Antibodies are also present in the plasma (i.e., fluid portion) of the blood and in the interstitial fluid of the tissues.
20 Antibodies can also be found in secretory fluids such as mucus, synovial fluid, sperm and milk into which certain types of antibody molecules are specifically transported. Procedures for obtaining biological samples (i.e., biopsying) from individuals are well known in the art. Such procedures include, but are not limited to, blood sampling, joint fluid biopsy, cerebrospinal biopsy and lymph node biopsy. These and other
25 procedures for obtaining tissue or fluid biopsies are described in details in <http://www.healthatoz.com/healthatoz/Atoz/search.asp>.

Regardless of the procedure employed, once the biological sample (i.e., normal and abnormal) is obtained, the titer (number) of antibody molecules for a scavenger receptor in the biological sample is determined.

30 Antibody titer can be determined by techniques which are well known in the art such as ELISA and dot blot using an immobilized antigen (see for Example Abbas, Lichtman and Pober "Cellular and Molecular Immunology". W.B. Saunders International Edition 1994 pages 56-59). Specifically, the antigen is preferably

immobilized on a solid support. To avoid non-specific binding of antibodies, the solid support is preferably coated with a nonantigenic protein as well. A peptide is typically immobilized on a solid matrix by adsorption from an aqueous medium, although other modes of immobilization applicable to proteins and peptides well known to those skilled in the art can be used. Useful solid matrices are also well known in the art. Such materials are water insoluble and include cross-linked dextran (e.g., SEPHADEXTM, Pharmacia Fine Chemicals, Piscataway, N.J.), agarose, polystyrene beads about 1 μ m to about 5 mm in diameter, polyvinyl chloride, polystyrene, cross-linked polyacrylamide, nitrocellulose- or nylon-based webs such as sheets, strips or paddles; or tubes, plates or the wells of a microtiter plate such as those made from polystyrene or polyvinylchloride.

Once the antigens is immobilized antibody-containing samples are added in serial dilutions until binding can no longer be observed. The antibody containing samples can be either a crude sample or immunoglobulin purified samples (e.g., ammonium sulfate precipitated fraction and/or chromatography isolated). Immunocomplexes are allowed to form and the support is washed to remove non-specifically bound antisera. Detection of immunocomplexes can be effected by adding labeled antibody-binding molecules such as staphylococcal protein A. The label can be an enzyme such as horseradish peroxidase (HRP), glucose oxidase, or the like. In cases where the major indicating group is an enzyme such as HRP or glucose oxidase, additional reagents are required to indicate that an immunocomplex has formed. Such additional reagents for HRP include hydrogen peroxide and an oxidation dye precursor such as diaminobenzidine. An additional reagent useful with glucose oxidase is 2,2'-azino-di-(3-ethyl-benzthiazoline-G-sulfonic acid) (ABTS).

Radioactive labels may also be used in accordance with the present invention. An exemplary radiolabeling agent is a radioactive element that produces γ ray emissions, such as ¹²⁵I. Methods of protein labeling are well-known in the art and described in details by Galfre et al., Meth. Enzymol., 73:3-46 (1981). The techniques of protein conjugation or coupling through activated functional groups are also applicable. See, for example, Aurameas et al., Scand. J. Immunol., 8(7):7-23 (1978); Rodwell et al., Biotech., 3:889-894 (1984); and U.S. Pat. No. 4,493,795.

Agents of the present invention (described above) can be included in a diagnostic or therapeutic kit. Thus, for example, antibodies and/or chemicals can be

packaged in a one or more containers with appropriate buffers and preservatives and used for diagnosis or for directing therapeutic treatment.

Preferably, the containers include a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic.

In addition, other additives such as stabilizers, buffers, blockers and the like may also be added.

The kit can also include instructions for determining if the tested subject is suffering from, or is at risk of developing inflammation.

As used herein the term "about" refers to $\pm 10\%$.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, Baltimore, Maryland (1989); Perbal, "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York (1988); Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al. (eds) "Genome Analysis: A Laboratory Manual Series", Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; "Cell Biology: A Laboratory Handbook", Volumes I-III Cellis, J. E., ed. (1994); "Current Protocols in Immunology" Volumes

I-III Coligan J. E., ed. (1994); Stites et al. (eds), "Basic and Clinical Immunology" (8th Edition), Appleton & Lange, Norwalk, CT (1994); Mishell and Shiigi (eds), "Selected Methods in Cellular Immunology", W. H. Freeman and Co., New York (1980); available immunoassays are extensively described in the patent and scientific literature, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 5,011,771 and 5,281,521; "Oligonucleotide Synthesis" Gait, M. J., ed. (1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins S. J., Eds. (1984); "Animal Cell Culture" Freshney, R. I., ed. (1986); "Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., (1984) and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA (1990); Marshak et al., "Strategies for Protein Purification and Characterization - A Laboratory Course Manual" CSHL Press (1996); all of which are incorporated by reference as if fully set forth herein. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

20

EXAMPLE 1

SRB-I vaccine suppresses EAE disease progression

The effect of SRB-I vaccination prior to EAE induction in Lewis rats was evaluated by manifestation of EAE clinical symptoms. The produced SRB-I autoantibodies were used to evaluate the effect of these antibodies on an ongoing disease.

25

MATERIALS AND METHODS

Animals:

Female Lewis rats, approximately six weeks old, were purchased from Harlan (Jerusalem, Israel) and maintained under SPF (specific pathogen-free) conditions in the Technion animal facility (Bruce Rappaport Faculty of Medicine, Technion, Haifa, Israel).

30

Peptide antigens:

Myelin Basic Protein (MBP) p68-86, Myelin Oligodendrocyte Glycoprotein (MOG) p33-55 were all synthesized on a MilliGen 9050 peptide synthesizer by standard 9-fluorenylmethoxycarbonyl chemistry and purified by high performance liquid chromatography. Sequence was confirmed by amino acid analysis and the correct mass was checked by mass spectroscopy. Peptides with over 95 % purity were used.

Immunizations and active disease induction:

Active induction of EAE in each experimental model was done as described before (20, 24, 25). Animals were then monitored for clinical signs daily by an observer blind to the treatment protocol. EAE was scored as follows: 0, clinically normal; 1, flaccid tail; 2, hind limb paralysis; 3, total hind limb paralysis, accompanied by an apparent front limb paralysis; 4, total hind limb and front limb paralysis.

DNA vaccines:

A plasmid DNA vaccine-encoding SRB-I was prepared as described before (10). In brief, RNA extracted from EAE brains of Lewis rats was subjected to RT-PCR using oligonucleotide primers [sense 5'-CCATGGGCGGCAGCTCCAGGGC-3' (SEQ ID NO: 1), anti-sense 5'-CTACAGCTTGGCTTCTTGAC-3' (SEQ ID NO: 2)] complementary to the published sequence of SRB-I (Accession No: AF071495). This RT-PCR reaction mixture was subjected to an amplification program of 1 min at 95 °C, 1 min at 55 °C and 1 min at 72 °C for 25 cycles. The 1.53 kb amplified product (SEQ ID NO:3) comprising nucleotides 10-1539 of SRB1R from *Rattus norvegicus* (Accession No: AF071495), was loaded onto a 5% polyacrylamide gel in TAE buffer, gel purified, sequenced and then ligated into a pcDNA3 plasmid. Prior to its use as a vaccine, the construct was injected to tibia muscle of rats that were sacrificed at different time points and the expression of RNA encoding SRB-I was verified. Under the working condition of the present study, SRB-I was highly transcribed in the leg muscle for not more than 25 days. DNA vaccines were administrated at a dose of 100µg plasmid in 100µl PBS to the tibia muscle.

Production and purification of recombinant SRB-I:

PCR product of SEQ ID NO: 3 was re-cloned into a PQE expression vector (Qaigen, Chatsworth, CA), expressed in *E. coli* and then purified by an NI-NTA-

supper flow affinity purification of 6xHis proteins (Qaigen, Chatsworth, CA). After purification, the purity of recombinant SRB-I was verified by gel electrophoresis followed by N -terminus sequencing (Protein Services Unit of the Technion, Haifa, Israel).

5 ***Evaluation of anti SRB-I antibody titer in sera of vaccinated rats:***

A direct ELISA assay has been utilized to determine the anti SRB-I antibody titer in DNA vaccinated rats. ELISA plates (Nunc, Roskilde, Denmark) were coated with 50 ng/well of the recombinant SRB-I protein produced in the present study as described above, Sera from DNA vaccinated rats were added in serial dilutions from 2⁵ to 2³⁰ to wells that were, or were not, coated previously with recombinant SRB-I. Calculation of each titer was done by comparing the O.D. measured (405nm) in wells coated with SRB-I to those not coated with this recombinant gene product. Goat anti-rat alkaline phosphatase conjugated IgG antibodies (Sigma) were used as a labeled antibody. p-Nitrophenyl Phosphate (p-NPP) (Sigma) was used as a soluble alkaline phosphatase substrate. Results were collected by ELISA reader (TECAN Spectra rainbow thermo absorber mini-plate reader). Results of triplicates were calculated as log₂Ab titer ± SE.

Evaluation of anti β-actin antibody titer in sera of vaccinated rats:

Recombinant soluble β-actin was obtained as described before (11). In brief, cDNA encoding rat β-actin (the natural cytoplasmic soluble form of β-actin, GenBank Accession NO: NM_031144) was PCR amplified using specific oligonucleotide primers [sense 5'-ATGGATGACGATATCGCTGCGCTC-3' (SEQ ID NO:4); anti-sense 5'-CTACCGGCCAGCCAGACG-3' (SEQ ID NO:5)]. Following cloning and sequence verification, the above cDNA was ligated into the pcDNA3 vector to be used as a control DNA vaccine). ELISA test was conducted as described above, but ELISA plates were coated with soluble β-actin rather than recombinant SRB-I.

IgG purification:

IgG was purified as described before (11).

30 ***Anti SRB-I specific antibody purification:***

Recombinant rat SRB-I (5 mg) encoded by SEQ ID NO:3, was bound to a CNBr activated Sepharose Column according to the manufactures instructions (Pharmacia biotech, catalog number 17-0820-01). Anti SRB-I specific antibodies

from sera (IgG fraction) of DNA vaccinated rats were loaded on the column and then eluted by an acidic elution buffer (glycine PH 2.5). Isotype of the purified antibody was determined by an ELISA assay in which anti rat IgG1a, IgG2b and IgG1 (Jackson, USA) were used as detection antibodies. Purified antibody was mostly of the IgG2a Isotype (data not shown).

FACS analysis:

Murine peritoneal macrophages [obtained as described elsewhere (9)], were activated with LPS (1 µg/ml), washed once in FACS buffer (PBS, 0.25 % BSA, 0.05 % sodium azide), and then incubated for 0.5 hours in FACS buffer enriched with 1 % normal rat serum. Cells were then resuspended (4 °C, 0.5 hours) in 96-well U plates (10⁶/well) with 10 µL FACS buffer supplemented with 0.5 µg/ml of IgG purified from rats that were subjected to SRB-I DNA plasmid administration (anti SRB-I), IgG purified from EAE rats treated with β-actin encoding DNA (control IgG) or no antibodies (no Ab). Cells were then washed with FACS buffer three times and incubated (4 °C, 0.5 hours) with 50 µL of FACS buffer supplemented with Goat anti-rat IgG-FITC (#F6258, 1:10,000 dilution; Sigma Chemical Co.). After incubation cells were washed with FACS buffer twice and analyzed in the presence of propidium iodide (PI) using a FACScalibur (Becton Dickinson, Mountain View, California, USA). Data was collected for 10,000 events and analyzed using a Cell Quest program (Becton Dickinson).

Statistical and graphical methods:

Significance of differences was examined using Student's t-test. A value of P<0.05 was considered significant. Mann-Whitney sum of ranks test was used to evaluate significance of differences in mean of maximal clinical score with P<0.05 considered significant.

RESULTS

The effect of SRB-I injections on EAE development:

In the first set of experiments, Lewis rats were subjected to four weekly administration of plasmid DNA encoding SRB-I, plasmid DNA encoding β-actin (as positive control), or PBS. Two months after last immunization EAE was induced. The rats treated with plasmid DNA encoding rat SRB-I (opened triangles) developed a significantly reduced form of disease versus PBS treated rats (closed squares) (Figure 1a, mean maximal score 3±0.28 Vs 1±0.28. p<0.001). DNA vaccine

encoding soluble β -actin had no effect on disease manifestation (Figure 1a, opened squares). At the peak of disease anti SRB-I specific antibody titer was determined in all groups (Figure 1b). Control EAE rats displayed a significant titer against SRB-I (Figure 1b $\log_2$ Ab titer of 10 ± 0.4 Vs 6 ± 0.2 in naïve rats, $p < 0.05$) that continued to
5 persist till 6 days after recovery, and then regressed back to background levels (not shown). DNA plasmid encoding SRB-I dramatically amplified this titer ($\log_2$ Ab titer of 22 ± 0.86 , $p < 0.001$ as compared to each control group) to provide antibody mediated protective immunity (Figure 1d). EAE rats did not develop a notable antibody titer to soluble β -actin and DNA plasmid encoding this gene product could
10 not breakdown tolerance against self (Figure 1b). SRB-I autoantibodies were found capable of specific binding to the recombinant SRB-I (Figure 1b), as well as to the natural form of SRB-I on activated macrophages (Figure 1c) and HEK293 line cells transfected with rat SRB-I (not shown). These autoantibodies could also adoptively transfer EAE resistance to other rats (Figure 1d, mean maximal score of 3.3 ± 0.3 in
15 control rats Vs 1.66 ± 0.18 , $p < 0.01$).

EXAMPLE 2

The anti-inflammatory mechanism of anti SRB-I antibodies.

The mechanism of action of anti SRB-I antibodies was evaluated by testing
20 their ability to modify cytokine production by murine peritoneal macrophages.

MATERIAL AND METHODS

Murine peritoneal macrophages:

Murine peritoneal macrophages were isolated as described before (9). Cells were then activated *in vitro* with $1 \mu\text{g/ml}$ LPS for 48 in the presence of (0, 10, 50 and
25 $100 \mu\text{g/ml}$) anti SRB-I polyclonal autoantibodies (CNBr purified, as described in Example 1 of the Examples section), or control IgG from normal rat serum (purified as described in Example 1 of the Examples section).

Cytokine determination:

Levels of IL-12, TNF- α and IL-10 were determined by ELISA (TECAN
30 Spectra rainbow thermo absorber mini-plate reader) using commercially available kits: mouse IL-12 (R&D system Inc. Minneapolis, MN), TNF- α & IL-10 (Diacclone, Besancon, France).

Statistical and graphical methods:

Statistical methods were as described in Example 1 of the Examples section.

RESULTS

The mechanism of action of anti SRB-I antibodies:

To explore the mechanistic basis by which anti SRB-I antibodies affect the
5 function of the immune system, these antibodies were purified and added to freshly
isolated peritoneal macrophages that were activated *in vitro* with LPS. The presence
of anti SRB-I antibodies in the peritoneal macrophages culture effectively suppressed
IL-12 and TNF- α production (Figure 2a and Figure 2b, closed squares) and at the
same time induced IL-10 production (Figure 2c, closed squares) as compared to
10 control IgG (Figure 2, open squares), all in a dose dependent manner (Figure 2).
Thus anti SRB-I antibodies redirect the polarization of macrophages from a pro-
inflammatory to anti-inflammatory mediators.

EXAMPLE 3

SRB-I vaccine suppresses an ongoing EAE

The effect of SRB-I vaccine on an ongoing disease was evaluated by EAE
C57BL/6 mice vaccination with plasmid DNA encoding SRB-I, 12 days after onset of
the disease.

MATERIALS AND METHODS

Animals:

C57BL/6 mice, approximately six weeks old, were purchased from Harlan
(Jerusalem, Israel) and maintained under SPF conditions in the Technion animal
facility (Bruce Rappaport Faculty of Medicine, Technion, Haifa, Israel).

EAE induction:

25 EAE was induced according to Semi-chronic model of MOG induced EAE in
the C57BL/6 mice as described before (20).

DNA vaccines:

SRB-I plasmid DNA vaccine was prepared as described in Example 1 of the
Examples section.

Mouse vaccination:

30 12 days after the induction of EAE (1-2 days after its onset) sick mice were
separated into 3 group of 6 C57BL/6 mice that were subjected to either a single
administration of plasmid DNA encoding SRB-I (pcDNA3-SRB-I), a single

administration of pcDNA3 (empty vector), or no injection (control). Vaccination was effected by administration of 100µg plasmid in 100µl PBS as described Example 1 of the Examples section.

Statistical and graphical methods:

5 Statistical analysis was effected as described in Example 1 of the Examples section.

RESULTS

The effect of SRB-I injections on the dynamics of an ongoing EAE disease:

Semi-chronic model of MOG induced EAE in the C57BL/6 mice (20) was
10 used to evaluate the effect of a plasmid DNA vaccine encoding SRB-I on the dynamics of an ongoing EAE disease (Figure 3). At this time anti SRB-I specific antibody titer exceeded the level of log₂Ab titer of 11±0.4 (Vs 6±0 in naïve mice) this titer accelerated to log₂Ab titer of 21±0.66 in SRB-I encoding DNA treated mice (p<0.001) (Data not shown). This acceleration was followed by a fast entry to
15 remission in treated mice (Figure 3, p<0.001). These data further suggest that SRB-I encoding DNA vaccine amplifies a pre-existing regulatory response.

EXAMPLE 4

Anti SRB-I monoclonal antibodies suppress ongoing EAE

20 Anti SRB-I monoclonal antibody was produced, and tested for its ability to modulate cytokine production by peritoneal activated macrophages as well as to affect an ongoing EAE disease

MATERIALS AND METHODS

Production of monoclonal anti SRB-I antibody:

25 C57/B6 mice were subsequently immunized (3 weekly immunizations) with the SRB-I encoding DNA plasmid. Two weeks after the last administration, these mice were subjected to active induction of EAE. Spleen cells were obtained for production of monoclonal antibodies two weeks later with SP2 cells (ATCC) as a fusion partner as described before (E. Harlow & D. Lane, Antibodies, Cold Spring
30 Harbor Laboratory Press, 1998). Screening of positive hybridoma was done in two steps of selection. The first one selected positive antibodies producing cells according to the ability to bind SRB-I over expressed by HEK293. Supernatant

isolated from hybridoma clones (1000 wells) was then subjected to FACS analysis for their ability to bind SRB-I.

Antibodies and peritoneal macrophages:

Peritoneal macrophages were obtained from thioglycollate (25 %, 3 ml) injected C57BL/6 mice and co-cultured with 1 µg/ml LPS for 48 in the presence, or
5 absence, of anti SRB-I polyclonal antibody (1:100, Calbichem, San Diego, CA), the monoclonal anti SRB-I antibody generated as described above (Clone 5D8, 10 µg/ml), isotype matched control antibody (IgG1_κ, Sigma), HDL (Chemicon International Temecula, CA) or anti murine CD36 (5 µg/ml, Santa Cruz
10 Biotechnology, Santa Cruz, CA). Cytokine levels were determined by ELISA as described in Example 2 of the Examples section. Results are shown as mean triplicates ± SE.

Antibodies administration to EAE rats:

Lewis rats were subjected to active EAE as described in Example 1 of the
15 Examples section. 1-2 days after the onset of disease these Lewis rats (6 in each group) were treated, every other day, with 100 µg/ml of monoclonal antibody 5D8 (anti SRB-I monoclonal antibody selected from hybridoma, as described above), control murine IgG1 (Sigma), anti SRB-I polyclonal antibody (produced as described in Example 1 of the Examples section), control rat IgG2a purified from EAE rats that
20 have been subjected to an empty plasmid DNA vaccination or PBS. Data was obtained by an observer blind to the experimental protocol as described in Example 1 of the Examples section. Results are presented as mean maximal score ±SE.

RESULTS

***The effect of monoclonal antibody to SRB-I on cytokine production of LPS
25 activated peritoneal macrophages:***

The ability of anti SRB-I specific monoclonal antibody to alter the cytokine profile produced by peritoneal macrophages and by the murine monocytic cell line J774 (ATCC) was evaluated by their cytokines expression. Table 1 compares the competence of one of these antibodies (clone 5D8, IgG1) to decrease TNF-α and
30 increase IL-10 production by LPS activated murine peritoneal macrophages to the effect of commercially available polyclonal antibody to SRB-I, to CD36 and to HDL. Each response was optimized during a preliminary experiment in which different doses were determined for properties to affect these parameters.

Table 1. The effect of anti SRB-I monoclonal antibody on cytokine production of LPS activated peritoneal macrophages

<i>Treatment</i>	<i>TNF-α (pg/ml)</i>	<i>IL-10 (pg/ml)</i>
-	3280 $\pm$ 210	730 $\pm$ 80
Anti SRB-I (polyclonal)	1760 $\pm$ 190	1240 $\pm$ 160
Anti SRB-I (clone 5D8)	1640 $\pm$ 140	1630 $\pm$ 150
Control murine IgG1 κ	3350 $\pm$ 260	710 $\pm$ 100
HDL (100 μ g/ml)	2560 $\pm$ 130	146 $\pm$ 30
Anti CD36 (polyclonal)	4660 $\pm$ 330	330 $\pm$ 90

5 The anti SRB-I monoclonal antibody produced by spleen cells isolated from the mice treated with SRB-I encoding DNA vaccine and the anti SRB-I specific polyclonal antibody could both suppress TNF- α production (1640 $\pm$ 140 and 1760 $\pm$ 190 pg/ml Vs 3280 $\pm$ 210 pg/ml in control samples, $p < 0.001$, Table 1). Control IgG1 κ (Sigma) had no significant effect on the production of these cytokines (Table 1).
 10 Very similar results were obtained using the murine monocytic cell line J774 or peritoneal macrophages from rats (data not shown). HDL could reduce TNF- α production (2560 $\pm$ 130 Vs 3280 $\pm$ 210, $p < 0.05$) but also decreased the production of the anti-inflammatory cytokine IL-10 (Table 1). Anti CD36 antibodies markedly increased TNF- α production (Table 1, $p < 0.01$).

15 ***The effect of anti SRB-I monoclonal antibody on EAE:***

The ability of the isolated anti SRB-I monoclonal antibody to suppress EAE was evaluated in adoptive transfer experiments. The antibodies were transferred to EAE rats just after the onset of disease and led to a significant reduction in disease severity (mean maximal score of 1.66 $\pm$ 0.2 in treated rats Vs 3 $\pm$ 0.3 in control rats,
 20 $p < 0.05$, Figure 4).

EXAMPLE 5

Anti SRB-I antibodies suppress an ongoing Inflammatory Bowel Disease (IBD)

Anti SRB-I antibodies were used to treat an induced IBD in Lewis rats.

25 ***MATERIALS AND METHODS***

Animals:

Female Lewis rats, approximately six weeks old (120-150g), were maintained as described in Example 1 of the Examples section.

Induction of colitis:

Experimental colitis was induced by intrarectal instillation of 250 µl of 125 mg/ml 2,4,6-trinitrobenzene sulfonic acid (TNBS) solution (Fluka, cat# 92822) dissolved in 50 % ethanol, using 8 cm neonate feeding tube as described before
5 [Fiorucci, S. et al., Immunity, 17:769., 2002]. 24 hours post injection all rats developed bloody diarrhea and severe diarrhea in the next day, accompanied with continuous loss of weight.

Anti SRB-I and IgG antibodies:

Specific anti SRB-I polyclonal antibodies and IgG antibodies were purified as
10 described in Example 1 of the Examples section.

Rat vaccination:

5 days after the induction of IBD, when the clinical manifestation of disease was apparent in all rats, they were separated into 3 groups of 6 equally sick rats each. On days 6, 8 and 10 after induction of disease these groups were administered with
15 either rat anti SRB-I polyclonal antibodies (DNA vaccination based antibody, produced as described in Example 1 of the Examples section, 100 µg/rat in PBS), rat IgG from pre-immunized rats (100 µg/rat in PBS), or PBS (control).

Histological sections:

Histological sections were conducted according to Fiorucci et al., [Fiorucci, S., et al., Immunity 17:769., 2002]. Briefly, on day 15 post induction of colitis, Lewis
20 rats were sacrificed and colons (cecum to rectum) were extracted, flushed with PBS and fixed in NBF (neutral buffered formalin). Paraffin sections (6 µm) were made and stained with H&E (hematoxylin and eosin). Each section represents ~250 sections that were screened by an observer blind to the experimental protocol.

RESULTS***Diarrhea symptoms:***

On day 15, post induction of the colitis, all 6 rats treated with anti SRB-I showed no signs of diarrhea and solid feces could be seen in the cage. Control rats (positive control of PBS treated rats and rats treated with control IgG) suffered from
30 severe diarrhea with no solid feces in the cage.

Histological sections of rats colons:

Histological colon sections of untreated rats showed normal mucosa with well defined colonic crypts and glands. Histological colon sections of both control rats

(positive control of PBS treated rats and rats treated with control IgG) taken 15 days post induction of colitis, showed severe signs of inflammation, megacolon, and signs of perforations (Figures 5b-c). Positive control of PBS treated rat sections showed diffuse, mononuclear inflammatory infiltrates in the mucosa and lamina propria, submucosa, and muscularis mucosae (Figure 5b). Sections displayed various degrees of damage to colonic tissue, ranging from heavily infiltrated areas with epithelial exfoliation to lesions with total destruction of mucosal surface, transmural infiltration, necrosis, and loss of tissue architecture. The control IgG colon sections showed massive destruction of colonic tissue with vast necrotic areas, absence of glandular structure and complete loss of tissue architecture (Figure 5c). Congestion and edema and congestion were seen in and around blood vessels. Damage was continuous from rectum and extended proximally up to cecum (Figure 5c). Histological colon sections of rats treated with anti SRB-I antibodies showed very little signs of inflammation, no perforations, and smaller and paler colons (Figure 5d). Sections showed moderate inflammatory infiltration, compared to control IgG treated rats (Figure 5d Vs. Figure 5c). Although infiltrated, colonic mucosa seems intact with visible brush border and glands and no transmural infiltrates. Submucosa was also much less infiltrated with no wall thickening (Figure 5d).

These results indicate that treatment of IBD with anti SRB-I antibodies effectively reduces diarrhea symptoms through reduction of damage to the colon.

EXAMPLE 6

Therapeutic monoclonal human anti SRB-I antibody

A monoclonal human anti SRB-I antibody was produced for therapeutic use.

MATERIALS AND METHODS

SRB-I encoding plasmids:

DNA encoding human SRB-I (CLA-I) was amplified using sense primer: 5' CCATGGGCTGCTCCGCCAAA 3' (SEQ ID NO: 6), and anti-sense primer: 5' CTACAGTTTTGCTTCCTGCAG 3' (SEQ ID NO: 7) The above described reaction mixture was subjected to an amplification program of 1 min at 95 °C, 1 min at 55 °C and 1 min at 72 °C for 25 cycles, generating 1.53 kb DNA fragment of SEQ ID NO:8 (Homo sapiens encoding SRB-I mRNA, nucleotides 70-1599 from accession number :Z22555). After PCR reaction, the mixture was loaded onto a 5% polyacrylamide gel

in TAE buffer. PCR product was gel-purified, cloned into a pUC57/T vector (T-cloning kit K1212; MBI Fermentas, Vilnius, Lithuania) and then used to transform *E. coli* cells. Clones were then sequenced (Sequenase version 2; Upstate Biotechnology, Cleveland, OH) and transferred into a pcDNA3 vector (Invitrogen, San Diego, CA).
5 Large-scale preparation of plasmid DNA was conducted using Mega prep (Qiagen, Chatsworth, CA).

Cells:

HEK293 (ATCC) were transfected with human SRB-I as described before [Scarselli E, et al., EMBO J. 21(19):5017-25, 2002]. Expression was verified by
10 FACS analysis as described before [Scarselli E, et al., EMBO J. 21(19):5017-25, 2002].

Production of monoclonal human anti SRB-I antibody:

Human anti SRB-I monoclonal antibodies were produced according to one of the two following protocols:

Protocol I

C57/B6 mice were subsequently immunized (3 weekly immunizations) with the human SRB-I (SEQ ID NO:8) encoding DNA plasmid. Two weeks after the last administration, these mice were subjected to active induction of EAE. Spleen cells were obtained for production of monoclonal antibodies two weeks later with SP2
20 cells (ATCC) as a fusion partner as described before (E. Harlow & D. Lane, Antibodies, Cold Spring Harbor Laboratory Press, 1998). Screening of positive hybridoma was done in two steps of selection. The first one selected positive antibodies producing cells according to the ability to bind the recombinant SRB-I over expressed by HEK293. Supernatant isolated from hybridoma clones (1000
25 wells) was then subjected to FACS analysis for their ability to bind SRB-I

Protocol II

The cloned human SRB-I (SEQ ID NO:8), obtained as described above, was re-cloned into a pQE expression vector, expressed in *E. coli* (Qiagen) and then purified by an NI-NTA-supper flow affinity purification of 6xHis proteins (Qiagen).
30 After purification, the purity of the recombinant human SRB-I was verified by gel electrophoresis followed by sequencing (N terminus) by the Technion's sequencing services unit (Technion, Haifa, Israel). This recombinant human SRB-I was then injected into BALB/C mice. First immunization was of 50 µg peptide emulsified in

CFA [incomplete Freund's adjuvant (IFA) supplemented with 10 mg/ml heat-killed *Mycobacterium tuberculosis* H37Ra in oil; Difco Laboratories, Detroit, MI] at a total volume of 400 μ l into the peritoneal cavity. Later on, in a 3 weeks interval these mice were administrated with 50 μ g/400 μ l or recombinant human SRB-I emulsified in IFA (Difco Laboratories, Detroit, MI). Three weeks after the third interval mice were injected (intravenous) with 50 μ g of recombinant human SRB-I in 100 μ l PBS. Three days later spleen cells were obtained and preparation of monoclonal antibodies was conducted as described above.

Screening for therapeutic human anti SRB-I monoclonal antibodies:

Best 50 positive clones obtained by each of the above protocols were then subjected to a biological assay in which each antibody was added to freshly isolated monocytes subjected to PMA activation. Levels of IL-12, TNF- α and IL-10 were determined in human THP-1 cell line (ATCC), as described in Example 2 of the Example section. Antibodies capable of down regulating the production TNF- α and at the same time increase the production of IL-10 were selected. Humanized antibodies were produced after re-cloning according to the protocol described by Kostelny, et al., 2001 [Kostelny, et al., Int J Cancer 93:556].

RESULTS

Potentially therapeutic human monoclonal anti SRB-I antibody:

2 pre-clones producing human monoclonal anti SRB-I antibodies (selected following protocol I described above) and 3 pre-clones producing human monoclonal anti SRB-I antibodies (selected following protocol II described above) were found capable of increasing IL-10 and reducing TNF- α production by human THP-1 cells (Data not shown). Clones that are found cross reacting with murine SRB-I can be tested for their ability to suppress EAE (as described in Examples 1-4 of the Examples section) and IBD (as described in Example 5 of the Examples section). These clones are potentially effective in inflammation therapy of human diseases.

REFERENCES CITED

(Additional references are cited in the text)

1. Brown MS, Goldstein JL. Lipoprotein metabolism in the macrophage: implications for cholesterol deposition in atherosclerosis. *Annu Rev Biochem* 1983; 52:223-61.
2. Krieger M. The other side of scavenger receptors: pattern recognition for host defense. *Curr Opin Lipidol* 1997; 8:275-80.
3. Krieger M, Herz J. Structures and functions of multiligand lipoprotein receptors: macrophage scavenger receptors and LDL receptor-related protein (LRP). *Annu Rev Biochem* 1994; 63:601-37.
4. Platt N, Gordon S. Is the class A macrophage scavenger receptor (SR-A) multifunctional? - The mouse's tale. *J Clin Invest* 2001; 108:649-54.
5. Janeway CA. Approaching the asymptote? Evolution and revolution in immunology. *Cold Spring Harb. Symp. Quant. Biol.* 1989; 54:1-13.
6. Acton SL, Scherer PE, Lodish HF, Krieger M. Expression cloning of SR-BI, a CD36-related class B scavenger receptor. *J Biol Chem* 1994; 269:21003-9.
7. Krieger M. Scavenger receptor class B type I is a multiligand HDL receptor that influences diverse physiologic systems. *J Clin Invest* 2001; 108:793-7.
8. Hyka N, Dayer JM, Modoux C, et al. Apolipoprotein A-I inhibits the production of interleukin-1 β and tumor necrosis factor- α by blocking contact-mediated activation of monocytes by T lymphocytes. *Blood* 2001; 97:2381-9.
9. Youssef S, Maor G, Wildbaum G, Grabie N, Gour-Lavie A, Karin N. C-C chemokine-encoding DNA vaccines enhance breakdown of tolerance to their gene products and treat ongoing adjuvant arthritis. *J Clin Invest* 2000; 106:361-71.
10. Youssef S, Wildbaum G, Maor G, et al. Long-lasting protective immunity to experimental autoimmune encephalomyelitis following vaccination with naked DNA encoding C-C chemokines. *J Immunol* 1998; 161:3870-9.
11. Wildbaum G, Netzer N, Karin N. Plasmid DNA encoding IFN- γ -inducible protein 10 redirects antigen-specific T cell polarization and suppresses experimental autoimmune encephalomyelitis. *J Immunol* 2002; 168:5885-92.

12. Wildbaum G, Westermann J, Maor G, Karin N. A targeted DNA vaccine encoding fas ligand defines its dual role in the regulation of experimental autoimmune encephalomyelitis. *J Clin Invest* 2000; 106:671-9.
13. Wildbaum G, Youssef S, Karin N. A targeted DNA vaccine augments the natural immune response to self TNF- α and suppresses ongoing adjuvant arthritis. *J Immunol* 2000; 165:5860-6.
14. Wildbaum G, Karin N. Augmentation of natural immunity to a pro-inflammatory cytokine (TNF- α) by targeted DNA vaccine confers long-lasting resistance to experimental autoimmune encephalomyelitis. *Gene Ther* 1999; 6:1128-38.
15. Salomon I, Netzer N, Wildbaum G, Schiff-Zuck S, Maor G, Karin N. Targeting the Function of IFN- γ -Inducible Protein 10 Suppresses Ongoing Adjuvant Arthritis. *J Immunol* 2002; 169:2685-93.
16. Sato Y, Roman M, Tighe H, et al. Immunostimulatory DNA sequences necessary for effective intradermal gene immunization. *Science* 1996; 273:352-357.
17. Raz E, Tighe H, Sato Y, et al. Preferential induction of a Th1 immune response and inhibition of specific IgE antibody formation by plasmid DNA immunization. *Proc Natl Acad Sci U S A* 1996; 93:5141-5.
18. Hemmi H, Takeuchi O, Kawai T, et al. A Toll-like receptor recognizes bacterial DNA. *Nature* 2000; 408:740-5.
19. Wen-Ming C, Xing Gong K, Zhi-Wei L, et al. DNA-PKcs Is Required for Activation of Innate Immunity by Immunostimulatory DNA. *Cell* 2000; 103:909-918.
20. Mendel I, Kerlero de Rosbo N, Ben-Nun A. A myelin oligodendrocyte glycoprotein peptide induces typical chronic experimental autoimmune encephalomyelitis in H-2b mice: fine specificity and T cell receptor V β expression of encephalitogenic T cells. *European Journal of Immunology* 1995; 25:1951-9.
21. Yednock TA, Cannon C, Fritz LC, Sanchez-Madrid F, Steinman L, Karin N. Prevention of experimental autoimmune encephalomyelitis by antibodies against $\alpha 4 \beta 1$ integrin. *Nature* 1992; 356:63-6.

22. Wildbaum G, Netzer N, Karin N. Tr1 cell-dependent active tolerance blunts the pathogenic effects of determinant spreading. *J Clin Invest* 2002; 110:701-10.
23. Husemann J, Silverstein SC. Expression of scavenger receptor class B, type I, by astrocytes and vascular smooth muscle cells in normal adult mouse and human brain and in Alzheimer's disease brain. *Am J Pathol* 2001; 158:825-32.
24. McRae BL, Kennedy MK, Tan LJ, Dal Canto MC, Picha KS, Miller SD. Induction of active and adoptive relapsing experimental autoimmune encephalomyelitis (EAE) using an encephalitogenic epitope of proteolipid protein. *Journal of Neuroimmunology* 1992; 38:229-40.
25. Karin N, Binah O, Grabie N, et al. Short peptide-based tolerogens without self-antigenic or pathogenic activity reverse autoimmune disease. *J Immunol* 1998; 160:5188-94.

WHAT IS CLAIMED IS:

1. A method of reducing an inflammatory response in a subject, the method comprising providing to a subject in need thereof a therapeutically effective amount of an agent capable of reducing activity and/or expression of a scavenger receptor or of an effector thereof, thereby reducing the inflammatory response in the subject.
2. The method of claim 1, wherein said agent is selected from the group consisting of:
 - (i) an oligonucleotide directed to an endogenous nucleic acid sequence expressing said scavenger receptor or said effector thereof;
 - (ii) a chemical inhibitor directed to said scavenger receptor or said effector thereof;
 - (iii) a neutralizing antibody directed at said scavenger receptor or said effector thereof; and
 - (iv) a non-functional derivative of said scavenger receptor or said effector thereof.
3. The method of claim 1, wherein said scavenger receptor is a class A scavenger receptor or a class B scavenger receptor.
4. The method of claim 3, wherein said class B scavenger receptor is SR-BI.
5. Use of an agent capable of reducing activity and/or expression of scavenger receptor or of an effector thereof for the manufacture of a medicament for the treatment of inflammatory diseases.
6. An article of manufacture comprising packaging material and a pharmaceutical composition identified for treating inflammatory diseases being contained within said packaging material, said pharmaceutical composition including,

as an active ingredient, an agent capable of reducing activity and/or expression of scavenger receptor or of an effector thereof and a pharmaceutically acceptable carrier.

7. A method of diagnosing predisposition to, or presence of, an inflammatory disease in a subject, the method comprising detecting anti scavenger receptor antibodies in a biological sample obtained from the subject, wherein a level above a predetermined normal threshold of said anti scavenger receptor antibodies in said biological sample is indicative of the inflammatory disease in the subject.

8. The method of claim 7, wherein detecting said anti scavenger receptor antibodies in said biological sample is effected by ELISA, RIA and/or dot blot.

9. A humanized antibody having an antigen recognition domain capable of specifically binding a scavenger receptor.

1/6




Fig. 1a




Fig. 1b

2/6




Fig. 1c




Fig. 1d

3/6




Fig. 2a




Fig. 2b




Fig. 2c

4/6




Fig. 3




Fig. 4

5/6

Normal colon




Fig. 5a

Control sick rat (colon)




Fig. 5b

6/6

Control IgG treated (colon)




Fig. 5c

Anti SRB-I therapy (colon)



Fig. 5d

1
SEQUENCE LISTING

<110> Karin, Nathan

<120> COMPOSITIONS AND METHODS FOR DIAGNOSING AND TREATING AN
INFLAMMATION

<130> 27156

<160> 8

<170> PatentIn version 3.2

<210> 1

<211> 22

<212> DNA

<213> Artificial sequence

<220>

<223> Single strand DNA oligonucleotide

<400> 1

ccatgggcgg cagctccagg gc 22

<210> 2

<211> 21

<212> DNA

<213> Artificial sequence

<220>

<223> Single strand DNA oligonucleotide

<400> 2

ctacagcttg gcttcttgca c 21

<210> 3

<211> 1531

<212> DNA

<213> Rattus norvegicus

<400> 3

catgggcggc agctccaggg cacgctgggt ggccttgggg ctgggcgttc tagggctgct 60
gtgtgctgcg ctccggcgta tcgtgattct catggtgcc tcgctcatca aacagcagg 120
gctcaagaat gtccgcatag accccagcag cctgtccttt gggatgtgga aggagatccc 180
tgttcccttc tacttgccg tctactctt cgagggtggtc aacccagcg aggtcctaaa 240
tggccagaag ccagtagtcc gggagcgcg accctatgtc tacagggagt tcagacaaaa 300
ggttaacatc accttcaatg acaatgacac ggtgtcctac atagagaacc gaagccttca 360
tttcagcca gacaggtccc agggctcaga ggtgactac attgtactgc ctaacatcct 420
ggtcctggga ggggcagtga tgatggagga caagcccaca agcctgaagc tgctaataac 480
cttggggttg gtcaccatgg gccagcgggc ctttatgaac cgcacgggtg gtgagatcct 540
gtggggctat gaggatccgt tcgtgaattt cctcagcaaa tatttcccag acatgttccc 600
catcaaaggc aaatttgccc tggtcggttg gatgaaccac tcagagttct ggctcttcac 660
cgtcttacag ggtgtccaga atttcagcaa gatccatctg gtggataagt ggaacggcct 720
cagcgaggtc aaatattggc attcgaaca gtgcaacatg atcaatgta ctgccgggca 780
gatgtgggca ccattcatga caccgaatc ctactggaa ttcttcagcc cagaagcctg 840
cagatctatg aagctcacct accaggaatc aagggtgttc gaaggcatcc ccacttatcg 900
cttcacggcc cccgatactt tgtttgccaa cgggtccgtc taccaccta atgaaggctt 960

2

ctgcccgtgc cgcgagtcgc gcattcagaa tgtcagcacc tgcaggtttg gtgcgcccct 1020
 gtttctctcc cagccccact tctacaatgc tgaccccggtg ctgtcagaag ctgttcttgg 1080
 tctgaaccct gacccaaggg agcattcttt gttccttgac atccaccggtg tcaactgggat 1140
 ccccatgaac tgttccgtga agatgcagct gagtctgtac atcaaatccg tcaagggcgt 1200
 cgggcaaaca gggaagatcg agccagtagt cctgccattg ctgtgggttcg aacagagcgg 1260
 gatgatgggt ggcaagaccc tgaacacgtt ctacacgcag ctggtgctga tgccccaggt 1320
 tcttactac gcgcagtatg tgctgctggg gcttggaggc ctctgctgc tggtgcccat 1380
 catttaccaa ctgcgcagcc aggagaaatg ctttttattt tggagtggta gtaaaaaggg 1440
 ctgcgaggat aaggaggcca tgcaggccta ctctgagtct ctgatgtcac cagctgcca 1500
 gggcacggta gtgcaagaag ccaagctgta g 1531

<210> 4
 <211> 24
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Single strand DNA oligonucleotide

<400> 4
 atggatgacg atacgctgc gctc 24

<210> 5
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Single strand DNA oligonucleotide

<400> 5
 ctaccggcca gccagacg 18

<210> 6
 <211> 20
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Single strand DNA oligonucleotide

<400> 6
 ccatgggctg ctccgcaaa 20

<210> 7
 <211> 21
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Single strand DNA oligonucleotide

<400> 7
 ctacagtttt gcttctgca g 21

<210> 8
 <211> 1530
 <212> DNA
 <213> Homo sapiens

<400> 8

3

atgggctgct ccgccaaagc gcgctgggct gccggggcgc tgggcgtcgc ggggctactg	60
tgcgctgtgc tgggcgctgt catgatcgtg atggtgccgt cgctcatcaa gcagcaggtc	120
cttaagaacg tgcgcatcga cccagtagc ctgtccttca acatgtggaa ggagatccct	180
atcccccttct atctctccgt ctacttcttt gacgtcatga accccagcga gatcctgaag	240
ggcgagaagc cgcagggtgcg ggagcgcggg ccctacgtgt acagggagtc caggcacaaa	300
agcaacatca ccttcaacaa caacgacacc gtgtccttcc tcgagtaccg caccttccag	360
ttccagccct ccaagtccca cggtcggag agcgactaca tcgtcatgcc caacatcctg	420
gtcttgggtg cggcgggtgat gatggagaat aagcccatga ccctgaagct catcatgacc	480
ttggcattca ccacctcgg cgaacgtgcc ttcatgaacc gactgtggg tgagatcatg	540
tggggctaca aggacccct tgtgaatctc atcaacaagt actttccagg catgttcccc	600
ttcaaggaca agttcggatt atttgctgag ctcaacaact ccgactctgg gctcttcacg	660
gtgttcacgg gggtcagaa catcagcagg atccacctcg tggacaagtg gaacgggctg	720
agcaagggtg acttctggca ttccgatcag tgcaacatga tcaatggaac ttctgggcaa	780
atgtggccgc ccttcatgac tctgagtcg tcgctggagt tctacagccc ggaggcctgc	840
cgatccatga agctaata gtaaggagtc ggggtgtttg aaggcatccc cacctatcgc	900
ttcgtggctc caaaaaccct gtttgccaac gggtcctatc acccaccxaa cgaaggcttc	960
tgcccgtgcc tggagtctgg aattcagaac gtcagcacct gcagggttcag tgcccccttg	1020
tttctctccc atcctcactt cctcaacgcc gacccgggtc tggcagaagc ggtgactggc	1080
ctgcacccta accaggaggc acactccttg ttcttggaac tccacccggt caccgggaatc	1140
cccatgaact gctctgtgaa actgcagctg agcctctaca tgaaatctgt cgcaggcatt	1200
ggacaaaactg ggaagattga gcctgtggtc ctgccgctgc tctgggttgc agagagcggg	1260
gccatggagg gggagactct tcacacattc tacactcagc tgggtgtgat gcccaagggtg	1320
atgcactatg cccagtagct cctcctggcg ctgggctgcg tcctgctgct ggtccctgct	1380
atctgccaaa tccggagcca agagaaatgc tatttatattt ggagtagtag taaaaagggc	1440
tcaaaggata aggaggccat tcaggcctat tctgaatccc tgatgacatc agctcccaag	1500
ggctctgtgc tgcaggaagc aaaactgtag	1530