



US 20030119121A1

(19) **United States**

(12) **Patent Application Publication**
Baker et al.

(10) **Pub. No.: US 2003/0119121 A1**

(43) **Pub. Date: Jun. 26, 2003**

(54) **SECRETED AND TRANSMEMBRANE
POLYPEPTIDES AND NUCLEIC ACIDS
ENCODING THE SAME**

Related U.S. Application Data

(75) Inventors: **Kevin P. Baker**, Darnestown, MD
(US); **Dan L. Eaton**, San Rafael, CA
(US); **Ellen Filvaroff**, San Francisco,
CA (US); **Audrey Goddard**, San
Francisco, CA (US); **J. Christopher
Grimaldi**, San Francisco, CA (US);
Austin L. Gurney, Belmont, CA (US);
Victoria Smith, Burlingame, CA (US);
Jean Philippe Stephan, Millbrae, CA
(US); **Colin K. Watanabe**, Moraga, CA
(US); **William I. Wood**, Hillsborough,
CA (US); **Zemin Zhang**, Foster City,
CA (US); **Sherman Fong**, Alameda, CA
(US)

(63) Continuation of application No. 10/197,942, filed on
Jul. 18, 2002, which is a continuation of application
No. PCT/US01/27099, filed on Aug. 29, 2001.

(60) Provisional application No. 60/232,887, filed on Sep.
15, 2000.

Publication Classification

(51) **Int. Cl.⁷** **C12P 21/02**; C12N 5/06;
C07K 14/435; C07H 21/04;
C12N 9/00
(52) **U.S. Cl.** **435/69.1**; 435/183; 435/320.1;
435/325; 536/23.2; 530/350

Correspondence Address:

KNOBBE, MARTENS, OLSON & BEAR, LLP
2040 MAIN STREET
FOURTEENTH FLOOR
IRVINE, CA 92614 (US)

(57) **ABSTRACT**

The present invention is directed to novel polypeptides and
to nucleic acid molecules encoding those polypeptides. Also
provided herein are vectors and host cells comprising those
nucleic acid sequences, chimeric polypeptide molecules
comprising the polypeptides of the present invention fused
to heterologous polypeptide sequences, antibodies which
bind to the polypeptides of the present invention and to
methods for producing the polypeptides of the present
invention.

(73) Assignee: **Genentech, Inc.**

(21) Appl. No.: **10/243,103**

(22) Filed: **Sep. 12, 2002**

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA57694

><subunit 1 of 1, 99 aa, 1 stop

><MW: 11050, pI: 7.47, NX(S/T): 0

MKIPVLPVAVLLSLLVLHSAQGATLGGPEEESTIENYASRPEAFNTPFLNIDKLRSFAKA

DEFLNWHALFESIKRKLPFLNWDAFPCLKGLRSATPDAQ

Important features:

Signal peptide:

Amino acids 1-22

N-myristoylation sites:

Amino acids 22-28;90-96

Homologous region to Peroxidase:

Amino acids 16-48

FIGURE 1

CGGACGCGTGGGTGCGAGGCGAAGGTGACCGGGGACCGAGCATTTCAGATCTGCTCGGTAGA
CCTGGTGCACCACCACC**ATG**TTGGCTGCAAGGCTGGTGTGTCTCCGGACACTACCTTCTAGG
GTTTTCCACCCAGCTTTACCAAGGCCTCCCCTGTTGTGAAGAATTCCATCACGAAGAATCA
ATGGCTGTTAACACCTAGCAGGGAATATGCCACCAAAACAAGAATTGGGATCCGGCGTGGGA
GAACTGGCCAAGAAGCTCAAAGAGGCAGCATTGGAACCATCGATGGAAAAATATTTAAATTT
GATCAGATGGGAAGATGGTTTGTGCTGGAGGGGCTGCTGTTGGTCTTGGAGCATTGTGCTA
CTATGGCTTGGGACTGTCTAATGAGATTGGAGCTATTGAAAAGGCTGTAATTTGGCCTCAGT
ATGTCAAGGATAGAATTCATTCCACCTATATGTACTTAGCAGGGAGTATTGGTTAACAGCT
TTGTCTGCCATAGCAATCAGCAGAACGCCTGTTCTCATGAACTTCATGATGAGAGGCTCTTG
GGTGACAATTGGTGTGACCTTTGCAGCCATGGTTGGAGCTGGAATGCTGGTACGATCAATAC
CATATGACCAGAGCCCAGGCCCAAAGCATCTTGCTTGGTTGCTACATTCTGGTGTGATGGGT
GCAGTGGTGGCTCCTCTGACAATATTAGGGGGTCCCTCTTCTCATCAGAGCTGCATGGGTACAC
AGCTGGCATTTGTGGGAGGCCTCTCCACTGTGGCCATGTGTGCGCCCAGTGAAAAGTTTCTGA
ACATGGGTGCACCCCTGGGAGTGGGCCTGGGTCTCGTCTTTGTGTCTCATTGGGATCTATG
TTTCTTCCACCTACCACCGTGGCTGGTGCCACTCTTTACTCAGTGGCAATGTACGGTGGATT
AGTTCTTTTCAGCATGTTCTTCTGTATGATACCCAGAAAGTAATCAAGCGTGCAGAAGTAT
CACCAATGTATGGAGTTCAAAAATATGATCCCATTAACCTCGATGCTGAGTATCTACATGGAT
ACATTAAATATATTTATGCGAGTTGCAACTATGCTGGCAACTGGAGGCAACAGAAAGAAAT**TG**
AAGTGAATCAGCTTCTGGCTTCTCTGCTACATCAAAATATCTTGTTTAAATGGGGCAGATATGC
ATTAAATAGTTTGTACAAGCAGCTTTCGTTGAAGTTTAGAAGATAAGAAACATGTCATCATA
TTTAAATGTTCCGGTAATGTGATGCCTCAGGTCTGCCTTTTTTTCTGGAGAATAAATGCAGT
AATCCTCTCCCAAAATAAGCACACACATTTTCAATTCTCATGTTTGAGTGATTTTAAATGTT
TTGGTGAATGTGAAAATAAGTTTGTGTGTCATGAGAATGTAAGTCTTTTTTCTACTTTAAAA
TTTAGTAGGTTCACTGAGTAAGTAAATTTAGCAAACCTGTGTTTGCATATTTTTTTGGAGT
GCAGAATATTGTAATTAATGTCATAAGTGATTTGGAGCTTTGGTAAAGGGACCAGAGAGAAG
GAGTCACCTGCAGTCTTTTGTTTTTTTAAATACTTAGAACTTAGCACTTGTGTTATTGATTA
GTGAGGAGCCAGTAAGAAACATCTGGGTATTTGGAAACAAGTGGTCATTGTTACATTCATTT
GCTGAACCTAACAAAACCTGTTTCATCCTGAAACAGGCACAGGTGATGCATTCTCTGCTGTTG
CTTCTCAGTGCTCTCTTCCAATATAGATGTGGTCATGTTTGACTTGTACAGAATGTTAATC
ATACAGAGAATCCTTGATGGAATTATATATGTGTGTTTACTTTTGAATGTTACAAAAGGAA
ATAACTTTAAACTATTTCTCAAGAGAAAATATTCAAAGCATGAAATATGTTGCTTTTTCCAG
AATACAAACAGTATACTCATG

FIGURE 2

MLAARLVCLRTLPSRVFHPAFTKASPVVKNSITKNQWLLTPSREYATKTRIGIRRGRTGQEL
KEAALEPSMEKIFKIDQMGRWFEVAGGAAVGLGALCYGGLSNEIGAIEKAVIWPQYVKDRI
HSTYMYLAGSIGLTALSAIAISRTFVLMNFMMRGSWVTIGVTFAAMVGAGMLVRSIPYDQSP
GPKHLAWLLHSGVMGAVVAPLTILGGPLLIRAAWYTAGIVGGLSTVAMCAPSEKFLNMGAPL
GVGLGLVFVSSLGSMFLPPTTVAGATLYSVAMYGGVLVFSMFLLYDTQKVIKRAEVSPMYGV
QKYDPINSMLSIYMDTLNIFMRVATMLATGGNRKK

FIGURE 3

CCAATCGCCCGGTGCGGTGGTGCAGGGTCTCGGGCTAGTCATGGCGTCCCCGTCTCGGAGACTGCAGACTAAAC
CAGTCATTACTTGTTC AAGAGCGTTCTGCTAATCTACACTTTTATTTTCTGGATCACTGGCGTTATCCTTCTT
GCAGTTGGCATTGTTGGGGCAAGGTGAGCCTGGAGAATTACTTTTCTCTTTTAAATGAGAAGGCCACCAATGTCCC
CTTCGTGCTCATTGCTACTGGTACCGTCATTATTCCTTTGGGCACCTTTGGTTGTTTTGCTACCTGCCGAGCTT
CTGCATGGATGCTAAAACGTATGCAATGTTTCTGACTCTCGTTTTTTTTGGTCGAACCTGGTCGCTGCCATCGTA
GGATTTGTTTTTCAGACATGAGATTAAGAACAGCTTTAAGAATAATTATGAGAAGGCTTTGAAGCAGTATAACTC
TACAGGAGATTATAGAAGCCATGCAGTAGACAAGATCCAAAATACGTTGCATTGTTGTGGTGTCAACCGATTATA
GAGATTGGACAGATACTAATTATTACTCAGAAAAAGGATTTCCTAAGAGTTGCTGTAAACTTGAAGATTGTACT
CCACAGAGAGATGCAGACAAAGTAAACAAAGAAGGTTGTTTTATAAAGGTGATGACCATTATAGAGTCAGAAAT
GGGAGTCGTTGCAGGAATTTCTTTGGAGTTGCTTGCTTCCAACCTGATTGGAATCTTTCTCGCCTACTGCCWCT
CTCGTGCCATAACAAATAACCAGTATGAGATAGTGTATACCCAATGTATCTGTGGGCCTATTCCCTCTCTACCTTT
AAGGACATTTAGGGTCCCCCTGTGAATTAGAAAGTTGCTTGGCTGGAGAACTGACAACACTACTTACTGATAG
ACCAAAAACTACACCAGTAGGTTGATTCAATCAAGATGTATGTAGACCTAAAACTACACCAATAGGCTGATTC
AATCAAGATCCGTGCTCGCAGTGGGCTGATTCAATCAAGATGTATGTTTGCTATGTTCTAAGTCCACCTTCTAT
CCCATTTCATGTTAGATCGTTGAAACCCTGTATCCCTCTGAAACACTGGAAGAGCTAGTAAATTGTAAATGAAGT

FIGURE 4

MASPSRRLQTKPVITCFKSVLLIYTFIFWITGVILLAVGIWGKVSLENYFSLLNEKATNVPF
VLIATGTVIILLGTFGCFATCRASAWMLKLYAMFLTLVFLVELVAAIVGFVFRHEIKNSFKN
NYEKALKQYNSTGDYRSHAVDKIQNTLHCCGVTDYRDWTDTNYYSEKGFPSCKLEDCTPQ
RDADKVNNEGCFIKVMTIIESEMGVVAGISFGVACFQLIGIFLAYCXSRITNNQYEIV

Important features of the protein:

Signal peptide:

amino acids 1-42

Transmembrane domains:

amino acids 19-42, 61-83, 92-114, 209-230,

N-glycosylation site.

amino acids 134-138

Tyrosine kinase phosphorylation site.

amino acids 160-168, 160-169

N-myristoylation site.

amino acids 75-81, 78-84, 210-216, 214-220, 226-232

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 69-80, 211-222

FIGURE 5

GGGGCCGCGGTCTAGGGCGGCTACGTGTGTTGCCATAGCGACCATTTTGCATTAACTGGTTG
GTAGCTTCTATCCTGGGGGCTGAGCGACTGCGGGCCAGCTCTTCCCCTACTCCCTCTCGGCT
CCTTGTGGCCCAAAGGCCTAACCGGGGTCCGGCGGTCTGGCCTAGGGATCTTCCCCGTTGCC
CCTTTGGGGCGGG**ATG**GCTGCGGAAGAAGAAGACGAGGTGGAGTGGGTAGTGGAGAGCATCG
CGGGGTTCTGCGAGGCCAGACTGGTCCATCCCCATCTTGGACTTTGTGGAACAGAAATGT
GAAGTTAACTGCAAAGGAGGGCATGTGATAACTCCAGGAAGCCCAGAGCCGGTGATTTTGGT
GGCCTGTGTTCCCCTTGTTTTTGATGATGAAGAAGAAAGCAAATTGACCTATACAGAGATTC
ATCAGGAATACAAAGAACTAGTTGAAAAGCTGTTAGAAGGTTACCTCAAAGAAATTGGAATT
AATGAAGATCAATTTCAAGAAGCATGCACTTCTCCTCTTGCAAAGACCCATACATCACAGGC
CATTTTGCAACCTGTGTTGGCAGCAGAAGATTTTACTATCTTTAAAGCAATGATGGTCCAGA
AAAACATTGAAATGCAGCTGCAAGCCATTGCAATAATTCAAGAGAGAAATGGTGTATTACCT
GACTGCTTAACCGATGGCTCTGATGTGGTCAGTGACCTTGAACACGAAGAGATGAAAATCCT
GAGGGAAGTTCTTAGAAAAATCAAAAGAGGAATATGACCAGGAAGAAGAAAGGAAGAGGAAAA
AACAGTTATCAGAGGCTAAAACAGAAGAGCCACAGTGCATTCCAGTGAAGCTGCAATAATG
AATAATTTCCAAGGGGATGGTGAACATTTTGCACACCCACCCTCAGAAGTTAAAATGCATTT
TGCTAATCAGTCAATAGAACCTTTGGGAAGAAAAGTGGAAAGGTCAGAACTTCCTCCCTCC
CACAAAAGGCCTGAAGATTCCTGGCTTAGAGCATGCGAGCATTGAAGGACCAATAGCAAAC
TTATCAGTACTTGGAACAGAAGAAGCTTCGGCAACGAGAACACTATCTCAAGCAGAAGAGAGA
TAAGTTGATGTCCATGAGAAAGGATATGAGGACTAAACAGATACAAAATATGGAGCAGAAAG
GAAAACCCACTGGGGAGGTAGAGGAAATGACAGAGAAACCAGAAATGACAGCAGAGGAGAAG
CAAACATTACTAAAGAGGAGATTGCTTGCAGAGAACTCAAAGAAGAAGTTATTAATAAG**TA**
ATAATTAAGAACAATTTAACAAAATGGAAGTTCAAATTGTCTTAAAAATAAATTATTTAGTC
CTTACACTG

FIGURE 6

MAAEEDEVEWVVESIAGFLRGPDWSIPILDFVEQKCEVNCKGGHVITPGSPEPVILVACVP
LVFDDEEESKLTYTEIHQEYKELVEKLLEGYLKEIGINEDQFQEACTSPLAKTHTSQAILQP
VLAADF^TIFKAMMVQKN^IEMQLQAIR^IIQERN^GVLPDCLTDGSDVVS^DLEHEEMKILREVL
RKSKEEYDQEEERKRKKQLSEAKTEEPTVHSSEAAIMNNSQGDGEHFAHPPSEVKMHFANQS
IEPLGRKVERSETSSLPQKGLKIPGLEHASIEGP^IANLSVLGTEELRQREHYLKQKRDKLMS
MRKDMRTKQIQNMEQKGKPTGEVEEMTEKPEMTAEEKQTLLKRLLA^EKLKEEVINK

N-glycosylation sites.

amino acids 224-228, 246-250, 285-289

N-myristoylation site.

amino acids 273-279

Amidation site.

amino acids 252-256

Cytosolic fatty-acid binding proteins.

amino acids 78-108

FIGURE 7

GGGAACGGAAAATGGCGCCTCACGGCCCGGGTAGTCTTACGACCCTGGTGCCCTGGGCTGCCGCCCTGCTCCTC
GCTCTGGGCGTGGAAGGGCTCTGGCGCTACCCGAGATATGCACCCAATGTCCAGGGAGCGTGCAAAATTTGTC
AAAAGTGGCCTTTTATTGTAAAACGACAGGAGAGCTAATGCTGCATGCCCGTTGCTGCCTGAATCAGAAGGGCA
CCATCTTGGGGCTGGATCTCCAGAAGTGTCTCTGGAGGACCCTGGTCCAACTTTTCATCAGGCACATACCACT
GTCATCATAGACCTGCAAGCAAACCCCTCAAAGGTGACTTGGCCAACACCTTCCGTGGCTTTACTCAGCTCCA
GACTCTGATACTGCCACAACATGTCAACTGTCCTGGAGGAATTAATGCCTGGAATACTATCACCTCTTATATAG
ACAACCAAATCTGTCAAGGGCAAAAGAACCTTTGCAATAACACTGGGGACCCAGAAATGTGTCCTGAGAATGGA
TCTTGTTGTACCTGATGGTCCAGGTCTTTTGCACTGTGTTTGTGCTGATGGTTTCCATGGATACAAGTGTATGCC
CCAGGGCTCGTTCTCACTGCTTATGTTCTTCGGGATTCTGGGAGCCACCACTCTATCCGTCTCCATTCTGCTTT
GGCCGACCCAGCGCCGAAAAGCCAAGACTTCATGAACTACATAGGTCTTACCATTGACCTAAGATCAATCTGAA
CTATCTTAGCCCAGTCAGGGAGCTCTGCTTCCTAGAAAGGCATCTTTCGCCAGTGGATTGCGCTCAAGGTTGAG
GCCGCCATTGGAAGATGAAAAATTGCACTCCCTTGGTGTAGACAAATACCAAGTCCCATTGGTGTGTTGCCCTA
TAATAAACACTTTTTCTTTTTTNNAAAAAAAAAAAAAAAAAAAAA

FIGURE 8

Signal Peptide:

Amino acids 1-30

Transmembrane:

Amino acids 198-212

MAPHGPGSLTTLVPWAAALLLALGVERALALPEICTQCPGSVQNLSKVAFYCKTTRELMLHA
RCCLNQKGTILGLDLQNCSEDPGPNFHQAHTTVIIDLQANPLKGDLANTFRGFTQLQTLIL
PQHVNCPGGINAWNTITSYIDNQICQGQKNLCNNTGDPEMCPENGSCVPDGPGLLQCVCADG
FHGYKCMRQGSFSLLMFFGILGATTLVSILLWATQRRKAKTS

FIGURE 9

GGGGGAGAAGGCGGCCGAGCCCCAGCTCTCCGAGCACCGGGTCGGAAGCCGCGACCCGAGCC
 GCGCAGGAAGCTGGGACCGGAACCTCGGCGGACCCGGCCCCACCCAACCTCACCTGCGCAGGT
 CACCAGCACCCCTCGGAACCCAGAGGCCCGCGCTCTGAAGGTGACCCCCCTGGGGAGGAAGGC
GATGCCCCCTGCGAGGACGATGGCCCGCGCCCGCCTCGCCCCGGCCGGCATCCCTGCCGTCTG
 CCTTGTGGCTTCTGTGCACGCTCGGCCTCCAGGGCACCCAGGCCGGGCCACCGCCCCGCGCCC
 CCTGGGCTGCCCCGCGGGAGCCGACTGCCTGAACAGCTTTACCGCCGGGGTGCCTGGCTTCGT
 GCTGGACACCAACGCCTCGGTGAGCAACGGAGCTACCTTCCTGGAGTCCCCACCGTGCGCC
 GGGGCTGGGACTGCGTGCGCGCCTGCTGCACCACCCAGAAGTGAACCTTGGCGCTAGTGGAG
 CTGCAGCCCGACCGCGGGGAGGACGCCATCGCCGCGCTGCTTCCTCATCAACTGCCTCTACGA
 GCAGAACTTCGTGTGCAAGTTCGCGCCAGGGAGGGCTTCATCAACTACCTCACGAGGGAAG
 TGTACCGCTCCTACCGCCAGCTGCGGACCCAGGGCTTTGGAGGGTCTGGGATCCCCAAGGCC
 TGGGCAGGCATAGACTTGAAGGTACAACCCAGGAACCCCTGGTGCTGAAGGATGTGAAAAA
 CACAGATTGGCGCCTACTGCGGGGTGACACGGATGTCAGGGTAGAGAGGAAAGACCCAAACC
 AGGTGGAAGTGTGGGGACTCAAGGAAGGCACCTACCTGTTCCAGCTGACAGTGACTAGTCA
 GACCACCCAGAGGACACGGCCAACGTACAGTCACTGTGCTGTCCACCAAGCAGACAGAAGA
 CTACTGCCTCGCATCCAACAAGGTGGGTGCGTGCCGGGGCTCTTTCCACGCTGGTACTATG
 ACCCCACGGAGCAGATCTGCAAGAGTTTCGTTTATGGAGGGTCTTGGGCAACAAGAACAAC
 TACCTTCGGGAAGAAGAGTGCATTCTAGCCTGTGCGGGTGTGCAAGGTGGGCCTTTGAGAGG
 CAGCTCTGGGGCTCAGGCGACTTTCCCCCAGGGCCCCCTCCATGGAAAGGCGCCATCCAGTGT
 GCTCTGGCACCTGTCAGCCACCCAGTTCGCGTGCAGCAATGGCTGCTGCATCGACAGTTTC
 CTGGAGTGTGACGACACCCCAACTGCCCCGACGCCCTCCGACGAGGCTGCCTGTGAAAAATA
 CACGAGTGGCTTTGACGAGCTCCAGCGCATCCATTTCCCAGTGACAAAGGGCACTGCGTGG
 ACCTGCCAGACACAGGACTCTGCAAGGAGAGCATCCCGCGCTGGTACTACAACCCCTTCAGC
 GAACACTGCGCCCGCTTTACCTATGGTGGTTGTTATGGCAACAAGAACAACCTTTGAGGAAGA
 GCAGCAGTGCCTCGAGTCTTGTGCGGGCATCTCCAAGAAGGATGTGTTTGGCCTGAGGCGGG
 AAATCCCCATTCCCAGCACAGGCTCTGTGGAGATGGCTGTACAGTGTTCCTGGTCATCTGC
 ATTGTGGTGGTGGTAGCCATCTTGGGTACTGCTTCTTCAAGAACCAGAGAAAGGACTTCCA
 CGGACACCACCACCCACCACCCACCCCTGCCAGCTCCACTGTCTCCACTACCGAGGACA
 CGGAGCACCTGGTCTATAACCACACCACCCGGCCCCCT**CTGAG**CCTGGGTCTCACCGGCTCTC
 ACCTGGCCCTGCTTCTGCTTGCCAAGGCAGAGGCCCTGGGCTGGGAAAACTTTGGAACCAG
 ACTCTTGCCTGTTTCCCAGGCCCACTGTGCCTCAGAGACCAGGGCTCCAGCCCTCTTGGAG
 AAGTCTCAGCTAAGCTCACGTCTGAGAAAGCTCAAAGGTTTGGGAAGGAGCAGAAAACCCTT
 GGGCCAGAAGTACCAGACTAGATGGACCTGCCTGCATAGGAGTTTGGAGGAAGTTGGAGTTT
 TGTTCCTCTGTTCAAAGCTGCCTGTCCCTACCCCATGGTGCTAGGAAGAGGAGTGGGGTGG
 TGTCAGACCCTGGAGGGCCCCAACCTGTCTCCCGAGCTCCTCTTCCATGCTGTGCGCCAG
 GGCTGGGAGGAAGGACTTCCCTGTGTAGTTTGTGCTGTAAAGAGTTGCTTTTTGTTTATTTA
 ATGCTGTGGCATGGGTGAAGAGGAGGGGAAGAGGCCGTGTTGGCCTCTCTGTCTCTCTTCC
 TCTTCCCCCAAGATTGAGCTCTCTGCCCTTGATCAGCCCCACCCCTGGCCTAGACCAGCAGAC
 AGAGCCAGGAGAGGCTCAGCTGCATTCGCGAGCCCCCACCCTCAAGGTTCTCCAACATCACA
 GCCCAGCCCACCCACTGGGTAATAAAAGTGGTTTGTGGAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 10

MAPARTMARARLAPAGIPAVALWLLCTLGLQGTQAGPPPAPPGLPAGADCLNSFTAGVPGFV
LDTNASVSNGATFLESPTVRRGWDCVRACCTTQNCNLALVELQPDRGEDAIAACFLINCLYE
QNFVCKFAPREGFINYL TREVYRSYRQLRTQGFGSGIPKAWAGIDLKVQPQEPLVLKDVEN
TDWRLLRGDTDVRVERKDPNQVELWGLKEGTYLFQLTVTSSDHPEDTANVTVTVLSTKQTED
YCLASNKVGRCRGSFPRWYYDPTEQICKSFVYGGCLGNKNNYLREEECILACRGVQGGPLRG
SSGAQATFPQGSPMERRHPVCSGTCQPTQFRCSNGCCIDSFLECDTPNCPDASDEAAACEKY
TSGFDELQRIHFPSDKGHCVDLPDTGLCKESIPRWYYNPFSEHCARFTYGGCYGNKNNFEEE
QQCLESCRGISKKDVFGLRREIPIPISTGSVEMAVTVFLVICIVVVVAILGYCFFKNQRKDFH
GHHHHPPPTPASSTVSTTEDTEHLVYNHTTRPL

signal sequence:

Amino acids 1-35

transmembrane domain:

Amino acids 466-483

N-glycosylation sites:

Amino acids 66-70;235-239;523-527

N-myristoylation sites:

Amino acids 29-35;43-49;161-167;212-218;281-287;282-288;285-291;
310-316;313-319;422-428;423-429;426-432

Cell attachment sequence:

Amino acids 193-199

Pancreatic trypsin inhibitor (Kunitz) family signatures:

Amino acids 278-298;419-438

FIGURE 12

MRAPGCGRLVLPLLLLAAAAALAEGLKEGETPGNFMEDEQWLSSISQYSGKIKHWNFRFRDEVEDDYIKSWE
DNQQGDEALDTTKDPCQKVKCSRHKVCIAGGYQRAMCISRKKLEHRIKQPTVKLHGKNKDSICKPCHMAQLASVC
GSDGHTYSSVCKLEQQACLSSKQLAVRCEGPCPCPTEQAATSTADGKPETCTGQDLADLGDRLRDWFQLLHENS
KQNGSASSVAGPASGLDKSLGASCKDSIGWMFSKLDTSADLFLDQTELAAINLDKYEVCIRPFFNSCDTYKDGR
VSTAECFCFWREKPPCLAELERIQIQEAAKKKPGIFIPSCDEDGYRKMQCDQSSGDCWRVDQLGLELTGTRT
HGSPDCDDIVGFSGDFGSGVGWEDEEEKETEEAGEEAEDEEGEAGEADDGGYIW

FIGURE 13

TGCGGCGACCGTCGTACACC**ATG**GGGCCTCCACCTCCGCCCCTACCGTGTGGGGCTGCTCCCCG
GATGGCCTCCTGTTCCCTCTTGCTGCTGCTAATGCTGCTCGCGGACCCAGCGCTCCCGGCCGG
ACGTCACCCCCCAGTGGTGCTGGTCCCTGGTGATTTGGGTAACCAACTGGAAGCCAAGCTGG
ACAAGCCGACAGTGGTGCACTACCTCTGCTCCAAGAAGACCGAAAGCTACTTCACAATCTGG
CTGAACCTGGAAGTGTGCTGCTGCCCTGTCATCATTGACTGCTGGATTGACAATATCAGGCTGGT
TTACAACAAAACATCCAGGGCCACCCAGTTTCCTGATGGTGTGGATGTACGTGTCCCTGGCT
TTGGGAAGACCTTCTCACTGGAGTTCCTGGACCCAGCAAAAGCAGCGTGGGTTCCTATTTTC
CACACCATGGTGGAGAGCCTTGTGGGCTGGGGCTACACACGGGGTGAGGATGTCCGAGGGGC
TCCCTATGACTGGCGCCGAGCCCCAAATGAAAACGGGGCCTACTTCCTGGCCCTCCGCGAGA
TGATCGAGGAGATGTACCAGCTGTATGGGGGCCCCGTGGTGTGGTTGCCACAGTATGGGC
AACATGTACACGCTCTACTTTCTGCAGCGGCAGCCGAGGCTGGAAGGACAAGTATATCCG
GGCCTTCGTGTCACTGGGTGCGCCCTGGGGGGCGTGGCCAAGACCTGCGCGTCCCTGGCT
CAGGAGACAACAACCGGATCCAGTCACTCGGGCCCCCTGAAGATCCGGGAGCAGCAGCGGCTCA
GCTGTCTCCACCAGCTGGCTGCTGCCCTACAACCTACACATGGTCACTGAGAAGGTGTTCTGT
GCAGACACCCACAATCAACTACACACTGCGGGACTACCGCAAGTTCTTCCAGGACATCGGCT
TTGAAGATGGCTGGCTCATGCGGCAGGACACAGAAGGGCTGGTGGAAGCCACGATGCCACCT
GGCGTGCAGTGCCTCTATGGTACTGGCGTCCCCACACCAGACTCCTTCTACTATGA
GAGCTTCCCTGACCGTGACCCATAAATCTGCTTTGGTGACGGCGATGGTACTGTGAAGTTGA
AGAGTGGCCTGCAGTGCCAGGCTGGCAGAGCCGCCAGGAGCACCAAGTGTGCTGCAGGAG
CTGCCAGGCAGCGAGCACATCGAGATGCTGGCCAACGCCACCACCCTGGCCTATCTGAAACG
TGTGCTCCTTGGGGCC**TGA**CTCCTGTGCCACAGGACTCCTGTGGCTCGGCCGTGGACCTGCT
GTTGGCCTCTGGGGCTGTGATGGCCACGCGTTTTGCAAAGTTTGTGACTCACCATTCAAGG
CCCCGAGTCTTGGACTGTGAAGCATCTGCCATGGGGAAGTGCTGTTTGTATCCTTTCTCTG
TGGCAGTGAAGAAGGAAGAAATGAGAGTCTAGACTCAAGGGACACTGGATGGCAAGAATGCT
GCTGATGGTGGAAGTGTGTGACCTTAGGACTGGCTCCACAGGGTGGACTGGCTGGGCCCTG
GTCCAGTCCCTGCCTGGGGCCATGTGTCCCCCTATTCCTGTGGGCTTTTCATACTTGCCTA
CTGGGCCCTGGCCCCGAGCCTTCCTATGAGGGATGTTACTGGGCTGTGGTCTGTACCCAG
AGGTCCCAGGGATCGGCTCCTGGCCCCCTCGGGTGACCCCTCCCACACACCAGCCACAGATAG
GCCTGCCACTGGTCATGGGTAGCTAGAGCTGCTGGCTTCCCTGTGGCTTAGCTGGTGGCCAG
CCTGACTGGCTTCCCTGGGCGAGCCTAGTAGCTCCTGCAGGCAGGGGAGTTTGTGCGTTCT
TCGTGGTTCCAGGCCCTGGGACATCTCACTCCACTCCTACCTCCCTTACCACCAGGAGCAT
TCAAGCTCTGGATTGGGCAGCAGATGTGCCCCAGTCCCGCAGGCTGTGTTCCAGGGGGCCT
GATTTCCCTCGGATGTGCTATTGGCCCCAGGACTGAAGCTGCCTCCCTTACCCTGGGACTGT
GGTTCCAAGGATGAGAGCAGGGGTGGAGCCATGGCCTTCTGGGAACCTATGGAGAAAGGGA
ATCCAAGGAAGCAGCCAAGGCTGCTCGCAGCTTCCCTGAGCTGCACCTCTTGCTAACCCAC
CATCACACTGCCACCCTGCCCTAGGGTCTCACTAGTACCAAGTGGGTGAGCACAGGGCTGAG
GATGGGGCTCCTATCCACCCTGGCCAGCACCCAGCTTAGTGCTGGGACTAGCCAGAACTT
GAATGGGACCCTGAGAGAGCCAGGGGTCCCCTGAGGCCCCCCTAGGGGCTTTCGTGTGCTGCC
CAGGGTGTCCATGGATCTCCCTGTGGCAGCAGGCATGGAGAGTACAGGGCTGCCTTCATGGC
AGTAGGCTCTAAGTGGGTGACTGGCCACAGGCCGAGAAAAGGGTACAGCCTCTAGGTGGGGT
TCCCAAAGACGCCTTCAGGCTGGACTGAGCTGCTCTCCACAGGGTTTCTGTGCAGCTGGAT
TTTCTCTGTTGCATACATGCCTGGCATCTGTCTCCCCCTGTTTCTGAGTGGCCCCACATGGG
GCTCTGAGCAGGCTGTATCTGGATTCTGGCAATAAAAGTACTCTGGATGCTGTAAAAA
AAAAA

FIGURE 14

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA44189
><subunit 1 of 1, 412 aa, 1 stop
><MW: 46658, pI: 6.65, NX(S/T): 4
MGLHLRPYRVGLLPDGLLFLLLLLMLLADPALPAGRHPPVVLVPGDLGNQLEAKLDKPTV
VHYLCSKKTESYFTIWLNLELLLPVIIDCWIDNIRLVYNKTSRATQFPDGVDRVPGFGK
TFSLEFLDPSKSSVGSYFHTMVESLVGWGYTRGEDVRGAPYDWRRAPNENGPYFLALREM
IEEMYQLYGGPVVLVAHSMGNMYTLYFLQRQPQAWKDKYIRAFVSLGAPWGGVAKTLRVL
ASGDNNRIPVIGPLKIREQQRSVSTSWLLPYNYSPEKVFVQTPTINYTLRDYRKFFQ
DIGFEDGWLMRQDTEGLVEATMPPGVQLHCLYGTGVPTPDSFYYESFPDRDPKICFGDGD
GTVNLKSALQCQAWQSRQEHQVLLQELPGSEHIEMLANATTLAYLKRVLLGP
```

Signal peptide:

Amino acids 1-28

Potential lipid substrate binding site:

Amino acids 147-164

N-glycosylation sites:

Amino acids 99-103;273-277;289-293;398-402

Lipases, serine proteins family:

Amino acids 189-202

Beta-transducin family Trp-Asp repeat:

Amino acids 353-366

Tyrosine kinase phosphorylation site:

Amino acids 165-174;178-186

N-myristoylation sites:

Amino acids 200-206;227-233;232-238;316-322

FIGURE 15

CAGAGCAGATA**ATG**GCAAGCATGGCTGCCGTGCTCACCTGGGCTCTGGCTCTTCTTTTCAGCG
TTTTTCGGCCACCCAGGCACGGAAAGGCTTCTGGGACTACTTCAGCCAGACCAGCGGGGACAA
AGGCAGGGTGGAGCAGATCCATCAGCAGAAGATGGCTCGCGAGCCCGCGACCCTGAAAGACA
GCCTTGAGCAAGACCTCAACAATATGAACAAGTTCCTGGAAAAGCTGAGGCCTCTGAGTGGG
AGCGAGGCTCCTCGGCTCCACAGGACCCGGTGGGCATGCGGCGGCAGCTGCAGGAGGAGTTG
GAGGAGGTGAAGGCTCGCCTCCAGCCCTACATGGCAGAGGCGCACGAGCTGGTGGGCTGGAA
TTTGGAGGGCTTGCGGCAGCAACTGAAGCCCTACACGATGGATCTGATGGAGCAGGTGGCCC
TGCGCGTGCAGGAGCTGCAGGAGCAGTTGCGCGTGGTGGGGGAAGACACCAAGGCCAGTTG
CTGGGGGGCGTGGACGAGGCTTGGGCTTTGCTGCAGGGACTGCAGAGCCGCGTGGTGCACCA
CACCGGCCGCTTCAAAGAGCTCTTCCACCCATACGCCGAGAGCCTGGTGAGCGGCATCGGGC
GCCACGTGCAGGAGCTGCACCGCAGTGTGGCTCCGCACGCCCCCGCCAGCCCCGCGCGCCTC
AGTCGCTGCGTGCAGGTGCTCTCCCGGAAGCTCACGCTCAAGGCCAAGGCCCTGCACGCACG
CATCCAGCAGAACCTGGACCAGCTGCGCGAAGAGCTCAGCAGAGCCTTTGCAGGCACTGGGA
CTGAGGAAGGGGCCGGCCCGGACCC**TAG**ATGCTCTCCGAGGAGGTGCGCCAGCGACTTCAG
GCTTTCCGCCAGGACACCTACCTGCAGATAGCTGCCTTCACTCGCGCCATCGACCAGGAGAC
TGAGGAGGTCCAGCAGCAGCTGGCGCCACCTCCACCAGGCCACAGTGCCTTCGCCCCAGAGT
TTCAACAAACAGACAGTGGCAAGGTTCTGAGCAAGCTGCAGGCCCGTCTGGATGACCTGTGG
GAAGACATCACTCACAGCCTTCATGACCAGGGCCACAGCCATCTGGGGGACCCCTGAGGATC
TACCTGCCCAGGCCCATTCCCAGCTTCTTGTCTGGGGAGCCTTGGCTCTGAGCCTCTAGCAT
GGTTTCAGTCCTTGAAAGTGGCCTGTTGGGTGGAGGGTGGAAAGTCTGTGCAGGACAGGGAG
GCCACCAAAGGGGCTGCTGTCTCCTGCATATCCAGCCTCCTGCGACTCCCCAATCTGGATGC
ATTACATTACCAGGCTTTGCAA
AAAAAA

FIGURE 16

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA48303
><subunit 1 of 1, 274 aa, 1 stop
><MW: 30754, pI: 7.77, NX(S/T): 0
MASMAAVLTWALALLSAFSATQARKGFWDYFSQTS GDKGRVEQIHQQKMAREPATLKDSL
EQDLNNMNKFLEKLRPLSGSEAPRLPQDPVGMRRQLQEELEE VKARLQPYMAEAHELVGW
NLEGLRQQLKPYTMDLMEQVALRVQELQEQLRVVGEDTKAQLLGGVDEAWALLQGLQSRV
VHHTGRFKELFHPYAESLVSGIGRHHVQELHRSVAPHAPASPARLSRCVQVLSRKLTLKAK
ALHARIQQNLDQLREELSRFAFGTGTEEGAGPDP
```

Important features of the protein:

Signal peptide:

Amino acids 1-23

Glycosaminoglycan attachment site:

Amino acids 200-204

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 233-237

N-myristoylation sites:

Amino acids 165-171;265-271

FIGURE 17

CTAAGAGGACAAG**ATG**AGGCCCGGCCTCTCATTCTCCTAGCCCTTCTGTTCTTCCTTGGCC
AAGCTGCAGGGGATTTGGGGGATGTGGGACCTCCAATTTCCAGCCCCGGCTTCAGCTCTTTC
CCAGGTGTTGACTCCAGCTCCAGCTTCAGCTCCAGCTCCAGGTCCGGCTCCAGCTCCAGCCG
CAGCTTAGGCAGCGGAGGTTCTGTGTCCCAGTTGTTTTCCAATTTACCGGCTCCGTGGATG
ACCGTGGGACCTGCCAGTGCTCTGTTTCCCTGCCAGACACCACCTTTCCCGTGGACAGAGTG
GAACGCTTGAATTCACAGCTCATGTTCTTTCTCAGAAGTTTGAGAAAGAACTTTCTAAAGTG
AGGGAATATGTCCAATTAATTAAGTGTGTATGAAAAAGAACTGTTAAACCTAACTGTCCGAAT
TGACATCATGGAGAAGGATACCATTTCTTACACTGAACTGGACTTCGAGCTGATCAAGGTAG
AAGTGAAGGAGATGGAAAACTGGTCATACAGCTGAAGGAGAGTTTTGGTGGAAAGCTCAGAA
ATTGTTGACCAGCTGGAGGTGGAGATAAGAAATATGACTCTCTTGGTAGAGAAGCTTGAGAC
ACTAGACAAAAACAATGTCTTGGCATTGCGCGAGAAATCGTGGCTCTGAAGACCAAGCTGA
AAGAGTGTGAGGCCCTCTAAAGATCAAAACACCCCTGTCTCCACCCCTCCCTCCACTCCAGGG
AGCTGTGGTTCATGGTGGTGTGGTGAACATCAGCAAACCGTCTGTGGTTCAGCTCAACTGGAG
AGGGTTTTCTTATCTATATGGTGTCTGGGGTAGGGATTACTCTCCCCAGCATCCAAACAAAG
GACTGTATTGGGTGGCGCCATTGAATACAGATGGGAGACTGTTGGAGTATTATAGACTGTAC
AACACACTGGATGATTTGCTATTGTATATAAATGCTCGAGAGTTGCGGATCACCTATGGCCA
AGGTAGTGGTACAGCAGTTTACAACAACAACATGTACGTCAACATGTACAACACCGGGAATA
TTGCCAGAGTTAACCTGACCACCAACACGATTGCTGTGACTCAAACCTCTCCCTAATGCTGCC
TATAATAACCGCTTTTCATATGCTAATGTTGCTTGGCAAGATATTGACTTTGCTGTGGATGA
GAATGGATTGTGGGTATTTTATTCAACTGAAGCCAGCACTGGTAACATGGTGAATTAGTAAAC
TCAATGACACCACACTTCAGGTGCTAAACACTTGGTATACCAAGCAGTATAAACCATCTGCT
TCTAACGCCTTCATGGTATGTGGGGTTCTGTATGCCACCCGTAATATGAACACCAGAACAGA
AGAGATTTTTTACTATTATGACACAAACACAGGGAAAGAGGGCAAACCTAGACATTGTAATGC
ATAAGATGCAGGAAAAAGTGCAGAGCATTAACCTATAACCCCTTTTGACCAGAACTTTATGTC
TATAACGATGGTTACCTTCTGAATTATGATCTTTCTGTCTTGAGAAAGCCCCAG**TAA**GCTGT
TTAGGAGTTAGGGTGAAAGAGAAAATGTTTGTGAAAAAATAGTCTTCTCCACTTACTTAGA
TATCTGCAGGGGTGTCTAAAAGTGTGTTTCAATTTGCAGCAATGTTTAGGTGCATAGTTCTAC
CACACTAGAGATCTAGGACATTTGTCTTGATTTGGTGAGTTCTCTTGGAATCATCTGCCTC
TTCAGGCGCATTTTGCATAAAGTCTGTCTAGGGTGGGATTGTGAGGCTTAGGGGCACTG
TGGGCCCTAGTGAAGCCTACTGTGAGGAGGCTTCACTAGAAAGCCTTAAATTAGGAATTAAGGA
ACTTAAACTCAGTATGGCGTCTAGGGATTCTTTGTACAGGAAATATTGCCAATGACTAGT
CCTCATCCATGTAGCACCCTAATTCTTCCATGCCTGGAAGAAACCTGGGGACTTAGTTAGG
TAGATTAATATCTGGAGCTCCTCGAGGGACCAAATCTCCAACCTTTTTTTTCCCCTCACTAGC
ACCTGGAATGATGCTTTGTATGTGGCAGATAAGTAAATTTGGCATGCTTATATATTCTACAT
CTGTAAAGTGCTGAGTTTTATGGAGAGAGGCCTTTTTATGCATTAAATTGTACATGGCAAATAA
ATCCCAGAAGGATCTGTAGATGAGGCACCTGCTTTTTCTTTCTCTCATTTGTCCACCTTACT
AAAAGTCAGTAGAATCTTCTACCTCATAACTTCCTTCCAAAGGCAGCTCAGAAGATTAGAAC
CAGACTTACTAACCAATTCCACCCCCACCAACCCCTTCTACTGCCTACTTTAAAAAAATT
AATAGTTTTCTATGGAACCTGATCTAAGATTAGAAAAATTAATTTTCTTTAATTTTATTATGG
ACTTTTATTTTACATGACTCTAAGACTATAAGAAAATCTGATGGCAGTGACAAAGTGCTAGCA
TTTATTGTTATCTAATAAAGACCTTGGAGCATATGTGCAACTTATGAGTGTATCAGTTGTTG
CATGTAATTTTTGCCTTTGTTTAAAGCCTGGAACCTTGAAGAAAATGAAAATTTAATTTTTTT
TTCTAGGACGAGCTATAGAAAAGCTATTGAGAGTATCTAGTTAATCAGTGCAGTAGTTGGAA
ACCTTGCTGGTGTATGTGATGTGCTTCTGTGCTTTTGAATGACTTTATCATCTAGTCTTTGT
CTATTTTTCTTTGATGTTCAAGTCCTAGTCTATAGGATTGGCAGTTTAAATGCTTTACTCC
CCCTTTTAAATAAATGATTAAATGTGCTTTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 18

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA48320
<subunit 1 of 1, 510 aa, 1 stop
<MW: 57280, pI: 5.61, NX(S/T): 6
MRPGLSFL LALLFFLGQAAGDLGDVGPPIPSPGFSSFPGV DSSSSFS SSSSRSGSSSSRSL
GSGGSVSQ LFSNFTGSVDDRGT CQCSVSLPDTTFPVDRVERLEFTAHVLSQKFEKELSKV
REYVQLISVYEKKLLNLTVRIDIMEKDTISYTELDFELIKVEVKEMEKLVIQLKESFGGS
SEIVDQLEVEIRNMTLLVEKLETLDKNNVLAIRREIVAL KTKLKECEASKDQNTFPVHPP
PTPGSCGHGGVVNISKPSVVQLNWRGFSYLYGAWGRDYSPQH PKNKGLYWVAPLNTDGRLL
EYYRLYNTLDDLLLYINARELRITYGQSGTAVYNNNNMYVNM YNTGNIARVNLTNTIAV
TQTL PNAAYNNRFSYANVAWQDIDFAVDENGLWVIYSTEASTGNMVISK LNDTTLQVLNT
WYTKQYKPSASNAFMVCGVLYATR TMNTRTEEI FYYYDTNTGKEGKL DIVMHKMQEKVQS
INYNPFDQKLYVYNDGYLLNYDLSVLQKPQ
```

Important features:

Signal peptide:

Amino acids 1-20

N-glycosylation sites:

Amino acids 72-76;136-140;193-197;253-257;352-356;
411-415

Tyrosine kinase phosphorylation site:

Amino acids 449-457

N-myristoylation sites:

Amino acids 16-22;39-45;53-59;61-67;63-69;81-87;
249-255;326-332;328-334;438-444

Legume lectins beta-chain proteins:

Amino acids 20-40

HBGF/FGF family proteins:

Amino acids 338-366

FIGURE 19

GCACCGCAGACGGCGCGGATCGCAGGGAGCCGGTCCGCCGCCGGAACGGGAGCCTGGGTGTG
CGTGTGGAGTCCGGA CTCTGGGAGACGATCGCG**ATG**AACACGGTGCTGTCGCGGGCGAACT
CACTGTTTCGCCTTCTCGCTGAGCGTGATGGCGGCGCTCACCTTCGGCTGCTTCATCACCACC
GCCTTCAAAGACAGGAGCGTCCCGGTGCGGCTGCACGTCTCGCGGATCATGCTAAAAAATGT
AGAAGATTTCACTGGACCTAGAGAAAGAAGTGATCTGGGATTTATCACATTTGATATAACTG
CTGATCTAGAGAATATATTTGATTGGAATGTTAAGCAGTTGTTTCTTTATTTATCAGCAGAA
TATTCAACAAAAAATAATGCTCTGAACCAAGTTGTCCTATGGGACAAGATTGTTTTGAGAGG
TGATAATCCGAAGCTGCTGCTGAAAGATATGAAAACAAAATATTTTTCTTTGACGATGGAA
ATGGTCTCAAGGGAAACAGGAATGTCACCTTTGACCCTGTCTTGGAACGTCGTACCAAATGCT
GGAATTCTACCTCTTGTGACAGGATCAGGACACGTATCTGTCCCATTTCCAGATACATATGA
AATAACGAAGAGTTAT**TAA**ATTATTCTGAATTTGAAACAAAA

FIGURE 20

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA56049
><subunit 1 of 1, 180 aa, 1 stop
><MW: 20313, pI: 8.91, NX(S/T): 1
MNTVLSRANSLFAFSLSVMAALTFGCFITTAFKDRSVPVRLHVSRI MLKNVEDFTGPRER
SDLGFITFDITADLENIFDWNVKQLFLYLSAEYSTKNNALNQVVLWDKIVLRGDNPKLLL
KDMKTKYFFFDGNGLKGNRNVTLTLSWNVVPNAGILPLVTGSGHVSVPFPD TYEITKSY
```

Important features of the protein:

Signal peptide:

Amino acids 1-25

Transmembrane domain:

Amino acids 149-164

N-glycosylation site:

Amino acids 141-145

N-myristoylation sites:

Amino acids 25-31;135-141

Cell attachment sequence:

Amino acids 112-115

TonB-dependent receptor proteins signature 1:

Amino acids 1-21

FIGURE 21

AAACTTGACGCC**ATGA**AGATCCCGGTCCTTCCTGCCGTGGTGCTCCTCTCCCTCCTGGTGCT
CCACTCTGCCCAGGGAGCCACCCTGGGTGGTCCTGAGGAAGAAAGCACCATTGAGAATTATG
CGTCACGACCCGAGGCCTTTAACACCCCGTTCCTGAACATCGACAAATTGCGATCTGCGTTT
AAGGCTGATGAGTTCCTGAACTGGCACGCCCTCTTTGAGTCTATCAAAGGAACTTCCTTT
CCTCAACTGGGATGCCTTTCCTAAGCTGAAAGGACTGAGGAGCGCAACTCCTGATGCCCAG**T**
GACCATGACCTCCACTGGAAGAGGGGGCTAGCGTGAGCGCTGATTCTCAACCTACCATAACT
CTTTCCTGCCTCAGGAACTCCAATAAAACATTTCCATCCAAA

FIGURE 22

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA57694
><subunit 1 of 1, 99 aa, 1 stop
><MW: 11050, pI: 7.47, NX(S/T): 0
MKIPVLPVAVLLSLLVLHSAQGATLGGPEEEESTIENYASRPEAFNTPFLNIDKLRSAFKA
DEFLNWHALFESIKRKLPFLNWDAFPKCLKGLRSATPDAQ
```

Important features:

Signal peptide:

Amino acids 1-22

N-myristoylation sites:

Amino acids 22-28;90-96

Homologous region to Peroxidase:

Amino acids 16-48

FIGURE 23

TCTCAGACTCTTGGAAGGGGCTATAC'TAGACACACAAAGACAGCCCCAAGAAGGACGGTGGA
GTAGTGTCTCGCTAAAAGACAGTAGAT**ATG**CAACGCCTCTTGCTCCTGCCCTTTCTCCTGC
TGGGAACAGTTTCTGCTCTTCATCTGGAGAATGATGCCCCCATCTGGAGAGCCTAGAGACA
CAGGCAGACCTAGGCCAGGATCTGGATAGTTCAAAGGAGCAGGAGAGAGACTTGGCTCTGAC
GGAGGAGGTGATTCAGGCAGAGGGAGAGGAGGTCAAGGCTTCTGCCTGTCAAGACAACTTTG
AGGATGAGGAAGCCATGGAGTCGGACCCAGCTGCCTTAGACAAGGACTTCCAGTGCCCCAGG
GAAGAAGACATTGTTGAAGTGCAGGGAAGTCCAAGGTGCAAGACCTGCCGCTACCTATTGGT
GCGGACTCCTAAAAC'TTTTGCAGAAAGCTCAGAATGTCTGCAGCAGATGCTACGGAGGCAACC
TTGTCTCTATCCATGACTTCAACTTCAACTATCGCATTCAAGTGCTGCACTAGCACAGTCAAC
CAAGCCCAGGTCTGGATTGGAGGCAACCTCAGGGGCTGGTTCCTGTGGAAGCGGTTTTGCTGG
ACTGATGGGAGCCACTGGAATTTTGCTTACTGGTCCCCAGGGCAACCTGGGAATGGGCAAGG
CTCCTGTGTGGCCCTATGCACCAAAGGAGGTTATTGGCGACGAGCTCAATGCGACAAGCAAC
TGCCCTTCGTCTGCTCCTTCT**TAAG**CCAGCGGCACGGAGACCCTGCCAGCAGCTCCCTCCCGT
CCCCAACCTCTCCTGCTCATAAATCCAGACTTCCCACAGCAAAAAAAAAAAAAAAAAAAAA

FIGURE 24

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA59208
<subunit 1 of 1, 225 aa, 1 stop
<MW: 25447, pI: 4.79, NX(S/T): 0
MQRLLLLPFLLLCTVSALHLENDAPHLESLETQADLGQDLSSKEQERDLALTEEVIQAE
GEEVKASACQDNFEDEEAMESDPAALDKDFQCPREEDIVEVQGS PRCKTCRYLLV RTPKT
FAEAQNVCSR CYGGNLVSIHDFNFNYRIQCCTSTVNQAQVWIGGNLRGWFLWKRFCWTDG
SHWNFAYWSPGQPGNGQGSCVALCTKGGYWRRACDKQLPFVCSF
```

Important features:

Signal peptide:

Amino acids 1-17

N-myristoylation sites:

Amino acids 13-19;103-109;134-140;164-170;
180-186;191-197;194-200;196-202;
198-204

C-type lectin domain signature:

Amino acids 200-224

FIGURE 25

CAACAGAAGCCAAGAAGGAAGCCGTCTATCTTGTGGCGATC**ATG**TATAAGCTGGCCTCCTGC
TGTTTGCTTTTCACAGGATTCTTAAATCCTCTCTTATCTCTTCCTCTCCTTGACTCCAGGGA
AATATCCTTTCAACTCTCAGCACCTCATGAAGACGCGCGCTTAACTCCGGAGGAGCTAGAAA
GAGCTTCCCTTCTACAGATATTGCCAGAGATGCTGGGTGCAGAAAGAGGGGATATTCTCAGG
AAAGCAGACTCAAGTACCAACATTTTTTAACCCAAGAGGAAATTTGAGAAAGTTTCAGGATTT
CTCTGGACAAGATCCTAACATTTTACTGAGTCATCTTTTGGCCAGAATCTGGAAACCATACA
AGAAACGTGAGACTCCTGATTGCTTCTGGAAATACTGTGTCT**TGA**AGTGAAATAAGCATCTGT
TAGTCAGCTCAGAAACACCCATCTTAGAATATGAAAAATAACACAATGCTTGATTTGAAAAC
AGTGTGGAGAAAACTAGGCAACTACACCCTGTTTATTGTTACCTGGAAAATAAATCCTCT
ATGTTTTGCACAAAAAAAAAAAAAAAAA

FIGURE 26

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA59214
<subunit 1 of 1, 124 aa, 1 stop
<MW: 14284, pI: 8.14, NX(S/T): 0
MYKLASCCLLFTGFLNPLLSLPLLDREISFQLSAPHEDARLTPEELERASLLQILPEML
GAERGDILRKADSSSTNIFNPRGNLRKFQDFSGQDPNILLSHLLARIWKPYKKRETPDCFW
KYCV
```

Important features:

Signal peptide:

Amino acids 1-20

Urotensin II signature:

Amino acids 118-124

Cell attachment sequence:

Amino acids 64-67

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 112-116

N-myristoylation sites:

Amino acids 61-67; 92-98

FIGURE 27

CAAGTAAATGCAGCACTAGTGGGTGGGATTGAGGTATGCCCTGGTGCATAAATAGAGACTCA
GCTGTGCTGGCACAACCTCAGAAGCTTGGACCGCATCCTAGCCGCCGACTCACACAAGGCAGGT
GGGTGAGGAAATCCAGAGTTGCC**ATG**GAGAAAATTCAGTGTGAGCATTTCTTGCTCCTTGTG
GCCCTCTCCTACACTCTGGCCAGAGATACCACAGTCAAACCTGGAGCCAAAAAGGACACAAA
GGACTCTCGACCCAACTGCCCCAGACCCTCTCCAGAGGTTGGGGTGACCAACTCATCTGGA
CTCAGACATATGAAGAAGCTCTATATAAATCCAAGACAAGCAACAAACCCTTGATGATTATT
CATCACTTGGATGAGTGGCCACACAGTCAAGCTTTAAAGAAAGTGTTTGCTGAAAATAAAGA
AATCCAGAAATTGGCAGAGCAGTTTGTCTCCTCAATCTGGTTTATGAAACAACCTGACAAAC
ACCTTTCTCCTGATGGCCAGTATGTCCCCAGGATTATGTTTGTTGACCCATCTCTGACAGTT
AGAGCCGATATCACTGGAAGATATTCAAATCGTCTCTATGCTTACGAACCTGCAGATACAGC
TCTGTTGCTTGACAACATGAAGAAAGCTCTCAAGTTGCTGAAGACTGAATTG**TAAAGAAAAA**
AAATCTCCAAGCCCTTCTGTCTGTCAGGCCTTGAGACTTGAAACCAGAAGAAGTGTGAGAAG
ACTGGCTAGTGTGGAAGCATAGTGAACACACTGATTAGGTTATGGTTTAATGTTACAACAAC
TATTTTTTAAGAAAAACAAGTTTTAGAAATTTGGTTTCAAGTGTACATGTGTGAAAACAATA
TTGTATACTACCATAGTGAGCCATGATTTTCTAAAAAAAAAAAAATAAATGTTA

FIGURE 28

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA59485
><subunit 1 of 1, 175 aa, 1 stop
><MW: 19979, pI: 9.26, NX(S/T): 0
MEKIPVSAFLLLVALSYTLARDTTVKPGAKKDTKDSRPKLPQTLSRGWDQLIWTQTYEE
ALYKSKTSNKPLMIIHHLDECPHSQALKKVFAENKEIQKLAEQFVLLNLVYETTDKHLSP
DGQYVPRIMFVDPSLTVRADITGRYSNRLYAYEPADTALLLDNMKKALKLLKTEL
```

Important features:

Signal peptide:

Amino acids 1-20

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 30-34

FIGURE 29

AAGACCTCTCTTTGCTGTTTGAGAGTCTCTCGGCTCAAGGACCGGGAGGTAAGAGGTT
TGGGACTGCCCCGGCAACTCCAGGGTGTCTGGTCCACGACCTATCCTAGGCGCC**AT**G GGT
GTGATAGGTATACAGCTGGTTGTTACCATGGTGATGGCCAGTGTCATGCAGAAGATTATA
CCTCACTATTCTCTTGCTCGATGGCTACTCTGTAATGGCAGTTTGAGGTGGTATCAACAT
CCTACAGAAGAAGAATTAAGAATTCTTGCAAGGAAACAACAAAAAGGGAAAAACAAAAAA
GATAGGAAATATAATGGTCACATTGAAAGTAAGCCATTAACCATTCCAAAGGATATTGAC
CTTCATCTAGAAACAAAGTCAGTTACAGAAAGTGGATACTTTAGCATTGCATTACTTTCCA
GAATACCAGTGGCTGGTGGATTTCACAGTGGCTGCTACAGTTGTGTATCTAGTAACTGAA
GTCTACTACAATTTTATGAAGCCTACACAGGAAATGAATATCAGCTTAGTCTGGTGCCTA
CTTGTTTTGTCTTTTGCAATCAAAGTTCTATTTTCATTAACACTACACACTATTTTAAAGTA
GAAGATGGTGGTGAAAGATCTGTTTGTGTACCTTTGGATTTTTTTTCTTTGTCAAAGCA
ATGGCAGTGTTGATTGTAACAGAAAATTATCTGGAATTTGGACTTGAAACAGGGTTTACA
AATTTTTTCAGACAGTGCAGTTCAGTTTCTTGAAAAGCAAGGTTTAGAATCTCAGAGTCCT
GTTTCAAACTTACTTTCAAATTTTTCCTGGCTATTTTCTGTTTATTTCATTGGGGCTTTT
TTGACATTTCTGGATTACGACTGGCTCAAATGCATCTGGATGCCCTGAATTTGGCAACA
GAAAAAATTACACAACTTTACTTCATATCAACTTCTTGGCACCTTTATTTATGGTTTTG
CTCTGGGTAAAACCAATCACCAAAGACTACATTATGAACCCACCACTGGGCAAAGAAATT
TCCCCATCTGGAAGAT**TGA**AGATAATAGTATCTAACTCACAAGGTTATCATTGGAATAAAT
GAAAGAACACATGTAATGCAACCAGCTGGAATTAAGTGCTTAATAAATGTTCTTTTCACT
GCTTGCCTCATCAGAATTAATAAGAAATACTTGACTAGT

FIGURE 30

```
</usr/seqdb2/sst/DNA/Dnaseqs.full/ss.DNA64966
<subunit 1 of 1, 307 aa, 1 stop
<MW: 35098, pI: 8.11, NX(S/T): 3
MGVIGIQLVVTMVMASVMQKIIPHYSLARWLLCNGSLRWYQHPTEEELRILAGKQQKGKT
KKDRKYNHIESKPLTIPKDIDLHLETKSVEVDTLALHYFPEYQWLVDFTVAATVVYILV
TEVYYNFMKPTQEMNISLVWCLLVLSFAIKVLFSLTTHYFKVEDGGERSVCVTFGFFFFV
KAMAVLIVTENYLEFGLETGFTNFSDSAMQFLEKQGLSQSPVSKLTFKFFLAIFCSFIG
AFLTFPGLRLAQMHLDALNLATEKITQTLLHINFLAPLFMVLLWVKPITKDYIMNPPLGK
EISPSGR
```

Important features:

Signal peptide:

Amino acids 1-15

Transmembrane domains:

Amino acids 134-157;169-189;230-248;272-285

N-glycosylation sites:

Amino acids 34-38;135-139;203-207

ATP/GTP-binding site motif A (P-loop):

Amino acids 53-61

Tyrosine kinase phosphorylation site:

Amino acids 59-67

N-myristoylation sites:

Amino acids 165-171;196-202;240-246;247-253

FIGURE 31

GTAGCATAGTGTGCAGTTCACCTGGACCAAAGCTTTGGCTGCACCTCTTCTGGAAAGCTGGCC
ATGGGGCTCTTCATGATCATTGCAATTCTGCTGTTCCAGAAACCCACAGTAACCGAACAACT
TAAGAAGTGCTGGAATAACTATGTACAAGGACATTGCAGGAAAATCTGCAGAGTAAATGAAG
TGCCTGAGGCACTATGTGAAAATGGGAGATACTGTTGCCTCAATATCAAGGAAGTGAAGCA
TGTAaaaaaaATTACAAAGCCACCTCGTCCAAAGCCAGCAACACTTGCAGTGAATCTTCAAGA
CTATGTTACAATAATAGAAAATTTCCCAAGCCTGAAGACACAGTCTACAT**TAA**ATCAAATACA
ATTTTCGTTTTCACTTGCTTCTCAACCTAGTCTAATAAACTAAGGTGATGAGATATACATCTT
CTTCCTTCTGGTTTCTTGATCCTTAAAAATGACCTTCGAGCATATTCTAATAAAGTGCATTGC
CAGTTAAAAAAAAAAAA

FIGURE 32

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA82403
><subunit 1 of 1, 99 aa, 1 stop
><MW: 11343, pI: 9.17, NX(S/T): 0
MGLFMIIAILLFQKPTVTEQLKKCWNNYVQGHCRKICRVNEVPEALCENGRYCCLNIKEL
EACKKITKPPRPKPATLALTLDYVTIIENFPSLKTQST
```

cAMP- and cGMP-dependent protein kinase phosphorylation site:
Amino acids 64-68

FIGURE 33

CGGACGCGTGGGCGCTGAGCCCCGGAGGCCAGGGCGTCCGGGGCTGCGCCACTTCCGAGGGC
CGAGCGCTGCCGGTCCCGGCGGTGCGACACGGCCGGGAGGAGGAGAACAACGCAAGGGGCTC
AACCGTCGGTTCGCTGGAGCCCCCCCCGGGGCGTGGCCTCCCGCCCCCTCAGCTGGGGAGGGC
GGGGCTCGCTGCCCCCTGCTGCCGACTGCGACCCTTACAGGGGAGGGAGGGCGCAGGCCGCG
CGGAGATGAGGAGGAGGCTGCGCCTACGCAGGGACGCATTGCTCACGCTGCTCCTTGGCGCC
TCCCTGGGCCTCTTACTCTATGCGCAGCGCGACGGCGCGGCCCGACGGCGAGCGCGCCGCG
AGGGCGAGGGAGGGCGGCACCGAGGGCCACCCCGGACCCGCGCGTTCCAGTTACCCGACG
CGGGTGCAGCCCCGCGGCCTACGAAGGGGACACACCGGCGCCGCCACGCCTACGGGACCC
TTTGACTTCGCCCCGTATTTGCGCGCCAAGGACCAGCGGCGGTTTCCACTGCTCATTAACCA
GCCGCACAAGTGCCGCGGCGACGGCGCACCCGGTGGCCGCCCGGACCTGCTTATTGCTGTCA
AGTCGGTGGCAGAGGACTTCGAGCGGCGCCAAGCCGTGCGCCAGACGTGGGGCGCGGAGGGT
CGCGTGCAGGGGGCGCTGGTGCGCCGCGTGTCTTGCTGGGCGTGCCAGGGGCGCAGGGCTC
GGGCGGGGCGGACGAAGTTGGGGAGGGCGCGCAACCCACTGGCGCGCCCTGCTGCGGGCCG
AGAGCCTTGCGTATGCGGACATCCTGCTCTGGGCCTTCGACGACACCTTTTTTAACCTAACG
CTCAAGGAGATCCACTTTCTAGCCTGGGCCTCAGCTTTCTGCCCCGACGTGCGCTTCGTTTT
TAAGGGCGACGCAGATGTGTTCGTGAACGTGGGAAATCTCCTGGAGTTCTGGCGCCGCGGGAC
CCGGCGCAAGACCTGCTTGCTGGTGACGTAATTGTGCATGCGCGGCCCATCCGCACGCGGGC
TAGCAAGTACTACATCCCCGAGGCCGTGTACGGCCTGCCCGCCTATCCGGCCTACGCGGGCG
GCGGTGGCTTTGTGCTTTCCGGGGCCACGCTGCACCGCCTGGCTGGCGCCTGTGCGCAGGTC
GAGCTCTTCCCCATCGACGACGTCTTTCTGGGCATGTGTCTGCAGCGCCTGCGGCTCACGCC
CGAGCCTCACCTGCCTTCCGCACCTTTGGCATCCCCAGCCTTCAGCCGCGCCGCATTTGA
GCACCTTCGACCCCTGCTTTTTACCGTGAGCTGGTTGTAGTGACGGGCTCTCGGCCGCTGAC
ATCTGGCTTATGTGGCGCCTGCTGCACGGGCCGCATGGGCCAGCCTGTGCGCATCCACAGCC
TGTCGCTGCAGGCCCCCTTCCAATGGGACTCCTAGCTCCCCACTACAGCCCCAAGCTCCTAAC
TCAGACCCAGAATGGAGCCGGTTTCCCAGATTATTGCCGTGTATGTGGTTCTTCCCTGATCA
CCAGGTGCCTGTCTCCACAGGATCCCAGGGGATGGGGGTAAAGCTTGGCTCCTGGCGGTCCA
CCCTGCTGGAACCAAGTTGAAACCCGTGTAATGGTGACCCTTTGAGCGAGCCAAGGCTGGGTG
GTAGATGACCATCTCTTGTCCAACAGGTCCCAGAGCAGTGGATATGTCTGGTCCTCCTAGTA
GCACAGAGGTGTGTTCTGGTGTGGTGGCAGGGACTTAGGGAATCCTACCCTCTGCTGGATT
TGGAACCCCTAGGCTGACGCGGACGTATGCAGAGGCTCTCAAGGCCAGGCCCCACAGGGAG
GTGGAGGGGCTCCGGCCGCCACAGCCTGAATTCATGAACCTGGCAGGCACTTTGCCATAGCT
CATCTGAAAACAGATATTATGCTTCCCACAACCTCTCCTGGGCCAGGTGTGGCTGAGCACC
AGGGATGGAGCCACACATAAGGGACAAATGAGTGCACGGTCTACCTAGTCTTTCCCTCACCT
CCTGAACTCACACAACAATGCCAGTCTCCCACTGGAGGCTGTATCCCCTCAGAGGAGCCAAG
GAATGTCTTCCCCTGAGATGCCACCACTATTAATTTCCCACATATGCTTCAACCACCCCTTG
CTCAAAAAACCAATACCCACACTTACCTTAATACAAACATCCCAGCAACAGCACATGGCAGG
CCATTGCTGAGGGCACAGGTGCTTTATTGGAGAGGGGATGTGGGCAGGGGATAAGGAAGGTTC
CCATTCCAGGAGGATGGGAACAGTCCTGGCTGCCCTGACAGTGGGGATATGCAAGGGGCT
CTGGCCAGGCCACAGTCCAAATGGGAAGACACCAGTCAGTCACAAAAGTCGGGAGCGCCACA
CAAACCTGGCTATAAGGCCCAGGAACCATATAGGAGCCTGAGACAGGTCCCCTGCACATTCA
TCATTAAACTATACAGGATGAGGCTGTACATGAGTTAATTACAAAAGAGTCATATTTACAAA
AATCTGTACACACATTTGAAAAACTCACAAAATTGTCATCTATGTATCACAAGTTGCTAGAC
CCAAAATATTAATAATGGGATAAAATTNNTTTAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAA

FIGURE 34

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA83505
><subunit 1 of 1, 402 aa, 1 stop
><MW: 43751, pI: 9.42, NX(S/T): 1
MRRRLRLRRDALLTLLLGASLGLLLYAQRDGAAPTASAPRGRGRAAPRPTPGPRAFQLPD
AGAAPPAYEGDTPAPPTPTGPFDFARYLRAKDQRRFPLLINQPHKCRGDGAPGGRPDLLI
AVKSV AEDFERRQAVRQTWGAEGRVQ GALVRRVFLLGVPRGAGSGGADEVGEGARTHWRA
LLRAESLAYADILLWAFDDTFFNLT LKEIHFLAWASAFCDVRFVFKGDADVFN VGNLL
EFLAPRDPAQDLLAGDVIVHARPIRTRASKYYIPEAVYGLPAYPAYAGGGGFVLSGATLH
RLAGACAQVELFPIDDVFLGMCLQRLRLTPEPHPAFRTEFGIPQPSAAPHLSTFDPCFYRE
LVVVHGLSAAADIWLMWRLLHGPHGPACAHQPVAAGPFQWDS
```

Important features of the protein:

Signal peptide:

Amino acids 1-27

N-glycosylation site:

Amino acids 203-207

N-myristoylation sites:

Amino acids 18-24;31-37;110-116;157-163;161-167
163-169;366-372

Cell attachment sequence:

Amino acids 107-110

FIGURE 35

AGCAGCCTCTGCCCGACCCGGCTCGTGCGGACCCAGGACCGGGCGCGGGACGCGTGCGTCC
AGCCTCCGGCGCTGCGGAGACCCGCGGCTGGGTCCGGGGAGGCCCAAACCCGCCCCCGCCA
GAACCCCGCCCCAAATTCCCACCTCCTCCAGAAGCCCCGCCCACTCCCGAGCCCCGAGAGCT
CCGCGCACCTGGGCGCCATCCGCCCTGGCTCCGCTGCACGAGCTCCACGCCCGTACCCCGGC
GTCACGCTCAGCCCGCGGTGCTCGCACACCTGAGACTCATCTCGCTTCGACCCCGCCGCCGC
CGCCGCCCCGGCATCCTGAGCACGGAGACAGTCTCCAGCTGCCGTT**CATG**CTTCTCCCCAGC
CTTCCGCAGCCCACCAGGGAAGGGGCGGTAGGAGTGGCCTTTTACCAAAGGGACCGGCGATG
CTCTGCAGGCTGTGCTGGCTGGTCTCGTACAGCTTGGCTGTGCTGTTGCTCGGCTGCCTGCT
CTTCTGAGGAAGGCGGCCAAGCCCGCAGGAGACCCACGGCCCACCAGCCTTCTGGGCTCCC
CCAACACCCCGTCACAGCCGGTGTCCACCCAACCACACAGTGTCTAGCGCCTCTCTGTCCCT
GCCTAGCCGTCACCGTCTCTTCTTGACCTATCGTCACTGCCGAAATTTCTCTATCTTGCTGG
AGCCTTCAGGCTGTTCCAAGGATACCTTCTGCTCCTGGCCATCAAGTCACAGCCTGGTCAC
GTGGAGCGACGTGCGGCTATCCGCAGCACGTGGGGCAGGGTGGGGGGATGGGCTAGGGGGCCG
GCAGCTGAAGCTGGTGTTCCTCCTAGGGGTGGCAGGATCCGCTCCCCAGCCAGCTGCTGG
CCTATGAGAGTAGGGAGTTTGATGACATCCTCCAGTGGGACTTCACTGAGGACTTCTTCAAC
CTGACGCTCAAGGAGCTGCACCTGCAGCGCTGGGTGGTGGCTGCCTGCCCCAGGCCCATTT
CATGCTAAAGGGAGATGACGATGTCTTTGTCCACGTCCCCAACGTGTTAGAGTTCCTGGATG
GCTGGGACCCAGCCAGGACCTCCTGGTGGGAGATGTCATCCGCCAAGCCCTGCCCAACAGG
AACACTAAGGTCAAATACTTCATCCCACCCTCAATGTACAGGGCCACCCACTACCCACCCTA
TGCTGGTGGGGGAGGATATGTCATGTCCAGAGCCACAGTGCGGCGCCTCCAGGCTATCATGG
AAGATGCTGAACCTTTCCCCATTGATGATGTCTTTGTGGGTATGTGCCTGAGGAGGCTGGGG
CTGAGCCCTATGCACCATGCTGGCTTCAAGACATTTGGAATCCGGCGGGCCCTGGACCCCTT
AGACCCCTGCCTGTATAGGGGGCTCCTGCTGGTTACCGCCTCAGCCCCCTCGAGATGTGGA
CCATGTGGGCACTGGTGACAGATGAGGGGCTCAAGTGTGCAGCTGGCCCCATACCCAGCGC
TGAAGGGTGGGTGGGCAACAGCCTGAGAGTGGACTCAGTGTTGATTCTCTATCGTGATGCG
AAATTGATGCCTGCTGCTCTACAGAAAATGCCAACTTGGTTTTTTAACTCCTCTCACCCCTGT
TAGCTCTGATTAAAAACACTGCAACCCAA

FIGURE 36

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA84927
><subunit 1 of 1, 378 aa, 1 stop
><MW: 42310, pI: 9.58, NX(S/T): 3
MLPPQPSAAHQGRGGRSGLLPKGPAMLCRLCWLVSYSIAVLLLGCLLFLRKAAKPAGDPT
AHQPFWAPPTPRHSRCPPNHTVSSASLSLPSRHRLFLTYRHCRNFSILLEPSGCSKDTFL
LLAIKSQPGHVERRAAIRSTWGRVGGWARGRQLKLVFLLGVAGSAPPAQLLAYESREFDD
ILQWDFTEDEFFNLTLKELHLQRWVVAACPQAHFMLKGDDDVFVHVPNVLEFLDGWDPAQD
LLVGDVIRQALPNRNTKVKYFIPPSMYRATHYPPYAGGGGYVMSRATVRRRLQAIMEDAEL
FPIDDVFVGMCLRRLGLSPMHAGFKTFGIRRPLDPLDPCLYRGLLLVHRLSPLEMWTMW
ALVTDEGLKCAAGPIPQR
```

Important features of the protein:

Signal peptide:

Amino acids 1-39

Transmembrane domain:

Amino acids 146-171

N-glycosylation sites:

Amino acids 79-83;104-108;192-196

N-myristoylation sites:

Amino acids 14-20;160-166;367-373

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 35-46

FIGURE 37

ATGAAAGTGATAATCAGGCAGCCCAAATGATTGTTAATAAGGATCAAATGAGATCGTGTATG
TGGGTCCAATCAATTGATTCTACACAAAGGAGCCTGGGGAGGGGGCC**ATG**GTGCCAATGCACT
TACTGGGGAGACTGGAGAAGCCGCTTCTCCTCCTGTGCTGCGCCTCCTTCTACTGGGGCTG
GCTTTGCTGGGCATAAAGACGGACATCACCCCCGTTGCTTATTTCTTTCTCACATTGGGTGG
CTTCTTCTTGTTTGCCTATCTCCTGGTCCGGTTTCTGGAATGGGGGCTTCGGTCCCAGCTCC
AATCAATGCAGACTGAGAGCCCAGGGCCCTCAGGCAATGCACGGGACAATGAAGCCTTTGAA
GTGCCAGTCTATGAAGAGGCCGTGGTGGGACTAGAATCCCAGTGCCGCCCCCAAGAGTTGGA
CCAACCACCCCCCTACAGCACTGTTGTGATACCCCCAGCACCTGAGGAGGAACAACCTAGCC
ATCCAGAGGGGTCCAGGAGAGCCAACTGGAACAGAGGCGAATGGCCTCAGAGGGGTCCATG
GCCCAGGAAGGAAGCCCTGGAAGAGCTCCAATCAACCTTCGGCTTCGGGGACCACGGGCTGT
GTCCACTGCTCCTGATCTGCAGAGCTTGCGGGCAGTCCCCACATTAGAGCCTCTGACTCCAC
CCCCTGCCTATGATGTCTGCTTTGGTCACCCTGATGATGATAGTGTTTTTTATGAGGACAAC
TGGGCACCCCCT**TAA**ATGACTCTCCCAAGATTTCTCTTCTCTCCACACCAGACCTCGTTCAT
TTGACTAACATTTTCCAGCGCCTACTATGTGTGTCAGAAACAAGTGTTTCTGCCTGGACATCAT
AAATGGGGACTTGGACCCTGAGGAGAGTCAGGCCACGGTAAGCCCTTCCCAGCTGAGATATG
GGTGGCATAATTTGAGTCTTCTGGCAACATTTGGTGACCTACCCCATATCCAATATTTCCAG
CGTTAGATTGAGGATGAGGTAGGGAGGTGATCCAGAGAAGGCGGAGAAGGAAGAAGTAACCT
CTGAGTGGCGGCTATTGCTTCTGTTCCAGGTGCTGTTTCGAGCTGTTAGAACCCTTAGGCTTGAC
AGCTTTGTGAGTTATTATTGAAAAATGAGGATTCCAAGAGTCAGAGGAGTTTGATAATGTGC
ACGAGGGCACACTGCTAGTAAATAACATTAAAATAACTGGAATGAA

FIGURE 38

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA92264
><subunit 1 of 1, 216 aa, 1 stop
><MW: 23729, pI: 4.73, NX(S/T): 0
MVPMHLLGRLEKPLLLCCASFLGLALLGIKTDITPVAYFFLTLLGGFFLFAYLLVRFLE
WGLRSQLQSMQTESPGPSGNARDNEAFVVPVYEEAVVGLESQCRPQELDQPPPYSTVVIP
PAPEEEQPSHPEGSSRAKLEQRRMASEGSMAQEGSPGRAPINLRLRGPRAVSTAPDLQSL
AAVPTLEPLTPPPAYDVCFGHPDDDSVFYEDNWAPP
```

Important features of the protein:

Signal peptide:

Amino acids 1-25

Transmembrane domain:

Amino acids 41-59

N-myristoylation site:

Amino acids 133-139

FIGURE 39

CCACACGCTCCGGCGGCTACACACCTAGGTGCGGTGGGCTTCGGGTGGGGGGCCTGCAGTA
GCTGATGGCAAGGGAGGAATAGCAGGGGTGGGGATTGTGGTGTGCGAGAGGTCCC GCGGACG
GGGGGCTCGGGGTCTCTTCAGACGAGATTCCCTTCAGGCTTGGGCCGGGTCCCTTCGCACG
GAGATCCCAATGAACGCGGGCCCCCTGGAGGCCGGTGGTTGGGGCTTCTCCGCGTCGGGGATG
GGGCCGGTACCCTAGCCCGTTTCCAGCGCCTCAGTCGGTTCCCC**ATG**CCCTCAGAGGTGGCC
CGGGGCAAGCGCGCCGCCCTCTTCTTCGCTGCGGTGGCCATCGTGTCTGGGGCTACCGCTCTG
GTGGAAGACCACGGAGACCTACCGGGCCTCGTTGCCTTACTCCCAGATCAGTGGCCTGAATG
CCCTTCAGCTCCGCTCATGGTGCCTGTCACTGTCTGTGTTACGCGGGAGTCAGTGCCCCCTG
GACGACCAGGAGAAGCTGCCCTTCACCGTTGTGCATGAAAGAGAGATTCTCTGAAATACAA
AATGAAAATCAAATGCCGTTTCCAGAAGGCCTATCGGAGGGCTTTGGACCATGAGGAGGAGG
CCCTGTCTATCGGGCAGTGTGCAAGAGGCAGAAAGCCATGTTAGATGAGCCTCAGGAACAAGCG
GAGGGCTCCCTGACTGTGTACGTGATATCTGAACACTCCTCATTCTTCCCCAGGACATGAT
GAGCTACATTGGGCCCCAAGAGGACAGCAGTGGTGCGGGGGATAATGCACCGGGAGGCCCTTTA
ACATCATTGGCCGCCGCATAGTCCAGGTGGCCAGGCCATGTCTTTGACTGAGGATGTGCTT
GTGCTGCTCTGGCTGACCACCTTCCAGAGGACAAGTGGAGCGCTGAGAAGAGGCGGCCCTCT
CAAGTCCAGCTTGGGCTATGAGATCACCTTCAGTTTACTCAACCCAGACCCCAAGTCCCATG
ATGTCTACTGGGACATTGAGGGGGCTGTCCGGCGCTATGTGCAACCTTTCTGAATGCCCTC
GGTGCCGCTGGCAACTTCTCTGTGGACTCTCAGATTCTTTACTATGCAATGTTGGGGGTGAA
TCCCCGCTTTGACTCAGCTTCTCCAGCTACTATTTGGACATGCACAGCCTCCCCCATGTCA
TCAACCCAGTGGAGTCCC GGCTGGGATCCAGTGCTGCCTCCTTGTACCCTGTGCTCAACTTT
CTACTCTACGTGCCTGAGCTTGCACACTCACCGCTGTACATT CAGGACAAGGATGGCGCTCC
AGTGGCCACCAATGCCTTCCATAGTCCCCGCTGGGGTGGCATTATGGTATATAATGTTGACT
CCAAAACCTATAATGCCTCAGTGCTGCCAGTGAGAGTCGAGGTGGACATGGTGCGAGTGATG
GAGGTGTTCTTGGCACAGTTGCGGTTGCTCTTTGGGATTGCTCAGCCCCAGCTGCCTCCAAA
ATGCCTGCTTTCAGGGCCTACGAGTGAAGGGCTAATGACCTGGGAGCTAGACCGGCTGCTCTGG
GCTCGGT CAGTGGAGAACCTTGGCCACAGGCCACCAACCCCTACCTCCCTGGCGCAGCTTCT
GGGCAAGATCAGCAACATTGTCAATTAAGGACGACGTGGCATCTGAGGTGTACAAGGCTGTAG
CTGCCGTC CAGAAGTCGGCAGAAAGTGGCGTCTGGGCACCTGGCATCTGCCTTTGTGCGC
AGCCAGGAAGCTGTGACATCCTCTGAGCTTGCCTTCTTTGACCCGTCACCTCCTCCACCTCCT
TTATTTCCCTGATGACCAGAAGTTTGCCATCTACATCCCACTCTTCTGCTATGGCTGTGC
CCATCCTCCTGTCCCTGGTCAAGATCTTCTGGAGACCCGCAAGTCTGGAGAAAGCCTGAG
AAGACAGACT**TGAG**CAGGGCAGCACCTCCATAGGAAGCCTTCTTTCTGGCCAAGGTGGGCGG
TGTTAGATTGTGAGGCACGTACATGGGGCTGCCGGAATGACTTAAATATTTGTCTCCAGTC
TCCACTGTTGGCTCTCCAGCAACCAAAGTACAACACTCCAAGATGGGTTTCATCTTTTCTTCC
TTTCCCATTCACCTGGCTCAATCCTCCTCCACCACCAGGGGCTCAAAGGCACATCATCCG
GGTCTCCTTATCTTGTGTTGATAAGGCTGCTGCCTGTCTCCCTCTGTGGCAAGGACTGTTTGT
TCTTTTGCCCCATTTCTCAACATAGCACACTTGTGCACTGAGAGGAGGGAGCATTATGGGAA
AGTCCCTGCCTTCCACACCTCTCTCTAGTCCCTGTGGGACAGCCCTAGCCCCTGCTGT CATG
AAGGGGCCAGGCATTGGTCACTGTGGGACCTTCTCCCTCACTCCCCTCCCTCCTAGTTGGC
TTTGTCTGT CAGGTGCAGTCTGGCGGGAGTCAGGAGGCAGCAGCTCAGGACATGGTGTCTGT
GT
GGAATCAAACAGTCTGAATCAAATCCTTGTTTTTTGCACCTATTGTCTGGAGAGCTTTGGA
TAAGGTATTGAATCTCTCTGAGCCTCAGTTTTTTCATTTGTTCAAATGGCACTGATGATGTCT
CCCTTACAAGATGGTTGTGAGGAGTAAATGTGATCAGCATGTAAAGTGTCTGGCGTGTAGTA
GGCTCTTAATAAACACTGGCTGAATATGAATTGGAATGAT

FIGURE 40

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA94713
><subunit 1 of 1, 547 aa, 1 stop
><MW: 61005, pI: 6.34, NX(S/T): 2
MPSEVARGKRAALFFAAVAIVLGLPLWWKTTTETYRASLPYSQISGLNALQLRLMVPVTVV
FTRESVPLDDQEKL PFTVVHEREIPLKYKMKIKCRFQKAYRRALDHEEEALSSGSVQEAE
AMLDEPQEQAEGLTVYVISEHSSLLPQDMMSYIGPKRTAVVRGIMHREAFNIIGRRIVQ
VAQAMSLTEDVLAAALADHLPEDKWSAEKRRPLKSSLGYEITFSLNPDPKSHDVYWDIE
GAVRRYVQPFNLALGAAGNFSVDSQILYYAMLGVNPRFDSASSSYLDMHSLPHVINPVE
SRLGSSAASLYPVLNFLLYVPELAHSPLYIQDKGAPVATNAFHSPRWGGIMVYNVDSKT
YNASVLPVRVEVDMVRVMEVFLAQLRLLFGLIAQPQLPPKCLLSGPTSEGLMTWELDRLLW
ARSVENLATATTTLTSLAQLLGKISNIVIKDDVASEVYKAVAAVQKSAEELASGHLASAF
VASQEAVTSSELAFFDPSLLHLLYFPDDQKFAIYIPLFLPMAVPILLSLVKIFLETRKSW
RKPEKTD
```

Important features of the protein:

Signal peptide:

Amino acids 1-23

Transmembrane domain:

Amino acids 511-530

N-glycosylation sites:

Amino acids 259-263;362-366

N-myristoylation sites:

Amino acids 255-261;304-310;335-341

Amidation sites:

Amino acids 7-11;174-178

FIGURE 41

CCAGCTGCAGAGAGGAGGAGGTGAGCTGCAGAGAAGAGGAGGTTGGTGTGGAGCACAGGCAG
CACCGAGCCTGCCCCGTGAGCTGAGGGCCTGCAGTCTGCGGCTGGAATCAGGATAGACACCA
AGGCAGGACCCCCAGAGATGCTGAAGCCTCTTTGGAAAGCAGCAGTGGCCCCACATGGCCA
TGCTCC**ATG**CCGCCCCCGCCGCCCGTGGGACAGAGAGGCTGGCACGTTGCAGGTCCTGGGAGC
GCTGGCTGTGCTGTGGCTGGGCTCCGTGGCTCTTATCTGCCTCCTGTGGCAAGTGCCCCGTCT
CCCACCTGGGGCCAGGTGCAGCCCAAGGACGTGCCCAGGTCCTGGGAGCATGGCTCCAGCCC
AGCTTGGGAGCCCCTGGAAGCAGAGGCCAGGCAGCAGAGGGACTCCTGCCAGCTTGTCTTG
TGGAAAGCATCCCCAGGACCTGCCATCTGCAGCCGGCAGCCCCCTCTGCCAGCCTCTGGGC
CAGGCCTGGCTGCAGCTGCTGGACACTGCCCAGGAGAGCGTCCACGTGGCTTCATACTACTG
GTCCCTCACAGGGCCTGACATCGGGGTCAACGACTCGTCTTCCCAGCTGGGAGAGGCTCTTC
TGCAGAAGCTGCAGCAGCTGCTGGGCAGGAACATTTCCCTGGCTGTGGCCACCAGCAGCCCG
AACTGGCCAGGACATCCACCGACCTGCAGGTTCTGGCTGCCCGAGGTGCCCATGTACGACA
GGTGCCCATGGGGCGGCTCACAGGGGTGTTTTGCACTCCAAATTCTGGGTTGTGGATGGAC
GGCACATATACATGGGCAGTGCCAACATGGACTGGCGGTCTCTGACGCAGGTGAAGGAGCTT
GGCGCTGTCTATCTATAACTGCAGCCACCTGGCCCAAGACCTGGAGAAGACCTTCCAGACCTA
CTGGGTACTGGGGGTGCCCAAGGCTGTCTCCCCAAAACCTGGCCTCAGAACTTCTCATCTC
ACTTCAACCGTTTCCAGCCCTTCCACGGCCTCTTTGATGGGGTGCCCACTGCTACTTC
TCAGCGTCGCCACCAGCACTCTGTCCCCAGGGCCGCACCCGGGACCTGGAGGCGCTGCTGGC
GGTGATGGGGAGCGCCAGGAGTTCATCTATGCCTCCGTGATGGAGTATTTCCCCACCACGC
GCTTCAGCCACCCCCGAGGTACTGGCCGGTGCTGGACAACGCGCTGCGGGCGGCAGCCTTC
GGCAAGGGCGTGCGCGTGCGCCTGTGGTGGCTGCGGACTCAACACGGACCCACCATGTT
CCCCACCTGCGGTCCCTGCAGGCGCTCAGCAACCCGCGGCCAACGTCTCTGTGGACGTGA
AAGTCTTCATCGTGCCGGTGGGGAACCATTTCCAACATCCCATTACAGCAGGGTGAACACAGC
AAGTTCATGGTCACGGAGAAGGCAGCCTACATAGGCACCTCCAAGTGGTCGGAGGATTACTT
CAGCAGCACGGCGGGGGTGGGCTTGGTGGTCACCCAGAGCCCTGGCGCGCAGCCCGCGGGGG
CCACGGTGCAGGAGCAGCTGCGGCAGCTCTTTGAGCGGGACTGGAGTTCGCGCTACGCCGTC
GGCCTGGACGGACAGGCTCCGGGCCAGGACTGCGTTTGGCAGGGCT**TGA**GGGGGGCCTCTTTT
TCTCTCGGCGACCCCGCCCCGCACGCGCCCTCCCCTCTGACCCCGGCCTGGGCTTCAGCCGC
TTCCTCCCGCAAGCAGCCCGGGTCCGCACTGCGCCAGGAGCCGCCTGCGACCGCCCGGGCGT
CGCAAACCGCCCGCCTGCTCTCTGATTTCCGAGTCCAGCCCCCCTGAGCCCCACCTCCTCC
AGGGAGCCCTCCAGGAAGCCCTTCCCTGACTCCTGGCCACAGGCCAGGCCTAAAAAAAC
TCGTGGCTTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 42

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96869
><subunit 1 of 1, 489 aa, 1 stop
><MW: 53745, pI: 8.36, NX(S/T): 8
MPPRRPWDREAGTLQVLGALAVLWLG SVALICLLWQVPRPPTWGQVQPKDVPRSWEHGSS
PAWEPLEAEARQQORDSCQLVLVESIPQDLPSAAGSPSAQPLGQAWLQLLDTAQESVHVAS
YYWSLTGPDIGVNDSSSQLGEALLQKLQQLLGRNISLAVATSSPTLARTSTD LQVLAARG
AHVRQVPMGRLTRGVLHSEKFWVVDGRHIYMGSANMDWRS LTQVKELGAVIYNCSHLAQDL
EKTFQTYWVLGVPKAVLPKTWPQN FSSHFNRFPFHGLFDGVPTTAYFSASPPALCPQGR
TRDLEALLAVMGSAQEFIIYASVMEYFPTTRFSHP RYWPVLDNALRAAAFGKGVRVRLLV
GCGLNTDPTMFPYLRSLQALS NPAANVSVDVKVFIVPVG NHSNIPFSRVNHSKFMVTEKA
AYIGTSNWSEDIYFSSTAGVGLVVTQSPGAQPAGATVQEQLRQLFERDWSSRYAVGLDGQA
PGQDCVWQG
```

Important features of the protein:

Signal peptide:

Amino acids 1-29

N-glycosylation sites:

Amino acids 133-137;154-158;232-236;264-268;
386-390;400-404;410-414;427-431

N-myristoylation sites:

Amino acids 58-64;94-100;131-137;194-200;251-257;
277-283;281-287;361-367;399-405;
440-446;448-454;478-484

FIGURE 43

GGGCCTGGCGATCCGGATCCCGCAGGCGCGCTGGCTGCGCTGCCCGGCTGTCTGTCTGTC**ATG**
GTGGGGCCCTGGGTGTATCTGGTGGCGGCAGTTTTGCTCATCGGCCTGATCCTCTTCCTGAC
TCGCAGCCGGGGTCGGGCGGCAGCAGCTGACGGAGAACCACTGCACAATGAGGAAGAGAGGG
CAGGAGCAGGCCAGGTAGGCCGCTCTTTGCCCCAGGAGTCTGAAGAACAGAGA**ACT**GGAAGC
AGACCCCGGCGTCGGAGGGACTTGGGCAGCCGTCTACAGGCCCAGCGTCGAGCCCAGCGAGT
GGCCTGGGAAGACGGGGATGAGAATGTGGGTCAA**ACT**GTTATTCCAGCCCAGGAGGAAGAAG
GCATTGAGAAGCCAGCAGAAGTTCACCCAACAGGGAAAATTGGAGCCAAGAA**ACT**ACGGAAG
CTAGAGGAAAAACAGGCTCGAAAGGCTCAGCGAGAGGCAGAGGAGGCTGAACGTGAAGAACG
GAAACGCCTAGAGTCCCAACGTGAGGCCGAATGGAAGAAGGAAGAGGAACGGCTTCGCCTGA
AGGAAGAACAGAAGGAGGAGGAAGAGAGGAAGGCTCAGGAGGAGCAGGCCCGGCGGGATCAC
GAGGAGTACCTGAA**ACT**GAAGGAGGCCTTCGTGGTAGAAGAAGAAGGTGTTAGCGAAACCAT
GACTGAGGAGCAGTCTCACAGCTTCCTGACAGAATTCATCAATTACATCAAGAAGTCCAAGG
TTGTGCTTTTGGAAAGATCTGGCTTTCCAGATGGGCTAAGGACTCAGGACGCCATAA**ACCGC**
ATCCAGGACCTGCTGACGGAGGGGACTCTAACAGGTGTGATTGACGACCGGGGCAAGTTTAT
CTACATAACCC**CAGAGGA**ACTGGCTGCCGTGGCCAATTT**CATCCGACAGCGGGGCCGGGTGT**
CCATCACAGAGCTTGCC**CAGGCCAGCA**ACTCCCTCATCTCCTGGGGCCAGGACCTCCCTGCC
CAGGCTTCAGCC**TGA**CTCCAGTCCTTCCTTGAGTGTATCCTGTGGCCTACATGTGTCTTCAT
CCTTCCCTAATGCCGTCTTGGGGCAGGGATGGAATATGACCAGAAAGTTGTGGATTAAAGGC
CTGTGAATACTGAA

FIGURE 44

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96881
><subunit 1 of 1, 315 aa, 1 stop
><MW: 35963, pI: 5.38, NX(S/T): 0
MVGPPWVYLVAAVLLIGLILFLTRSRGAAAAADGEPLHNEEERAGAGQVGRSLPQEESEEQR
TGSRRPRRRDLGSRLQAQRRRAQRVAVWEDGDENVGQTVIPAQEEEGIEKPAEVHPTGKIGA
KKLRKLEEKQARKAQREAEAAEREERKRLESQREAEWKKEEERLRLKEEQKEEEERKAQE
EQARRDHEEYLLKLEAFVVEEEGVSETMTEEQSHSFLTEFINYIKKSKVVLLLEDLAFQMG
LRTQDAINRIQDLLTEGTLTGVIDDRGKFIYITPEELAAVANFIRQRGRVSITELAQASN
SLISWGQDLPAQASA
```

Important features of the protein:

Signal peptide:

Amino acids 1-26

N-myristoylation sites:

Amino acids 203-209;257-263

FIGURE 45

ACGGGCGCGCAGCGGCAGTGACGTAGGGTTGGCGCACGGATCCGTTGCGGCTGCAGCTCTGCA
 GTCGGGCGCGTTCCTTCGCCGCCGCCAGGGGTAGCGGTGTAGCTGCGCAGCGTCGCGCGCGCT
 ACCGCACCCAGGTTTCGGCCCGTAGGCGTCTGGCAGCCCCGGCGCCATCTTCATCGAGCGCCAT
GGCCGCAGCCTGCGGGCCGGGAGCGGCCGGGTACTGCTTGCTCCTCGGCTTGCAATTTGTTTC
 TGCTGACCGCGGGCCCTGCCCTGGGCTGGAACGACCCCTGACAGAATGTTGCTGCGGGATGTA
 AAAGCTCTTACCCTCCACTATGACCGCTATACCACCTCCCGCAGGCTGGATCCCATCCCACA
 GTTGAAATGTGTTGGAGGCACAGCTGGTTGTGATTCTTATACCCCAAAGTCATACAGTGTC
 AGAACAAAGGCTGGGATGGGTATGATGTACAGTGGGAATGTAAGACGGACTTAGATATTGCA
 TACAAATTTGGAAAACTGTGGTGAGCTGTGAAGGCTATGAGTCCTCTGAAGACCAGTATGT
 ACTAAGAGGTTCTTGTGGCTTGGAGTATAATTTAGATTATACAGAACTTGGCCTGCAGAAAC
 TGAAGGAGTCTGGAAGCAGCACGGCTTTGCCTCTTTCTCTGATTATTATTATAAGTGGTCC
 TCGGCGGATTCCCTGTAACATGAGTGGATTGATTACCATCGTGGTACTCCTTGGGATCGCCTT
 TGTAAGTCTATAAGCTGTTCCCTGAGTGACGGGCAGTATTCTCCTCCACCGTACTCTGAGTATC
 CTCCATTTTCCACCGTTACCAGAGATTACCAACTCAGCAGGACCTCCTCCCCAGGCTTT
 AAGTCTGAGTTACAGGACCACAGAATACTGGCCATGGTGCAACTTCTGGTTTTGGCAGTGC
 TTTTACAGGACAACAAGGATATGAAAATTCAGGACCAGGGTTCTGGACAGGCTTGGGAACTG
 GTGGAATACTAGGATATTTGTTTGGCAGCAATAGAGCGGCAACACCCTTCTCAGACTCGTGG
 TACTACCCGTCCTATCCTCCCTCCTACCCTGGCACGTGGAATAGGGCTTACTCACCCCTTCA
 TGGAGGCTCGGGCAGCTATTCCGTATGTTCAAACCTCAGACACGAAAACCAGAACTGCATCAG
 GATATGGTGGTACCAGGAGACGATTAAAGTAGAAAGTTGGAGTCAAACACTGGATGCAGAAAT
 TTTGGATTTTTCATCACTTTCTCTTTAGAAAAAAAGTACTACCTGTTAACAATTGGGAAAAG
 GGGATATTCAAAGTTCTGTGGTGTTATGTCCAGTGTAGCTTTTTGTATTCTATTATTTGAG
 GCTAAAAGTTGATGTGTGACAAAATACTTATGTGTTGTATGTCAGTGTAAACATGCAGATGTA
 TATTGCAGTTTTTGAAGTGATCATTACTGTGGAATGCTAAAAATACATTAATTTCTAAAAC
 CTGTGATGCCCTAAGAAGCATTAAAGAATGAAGGTGTTGTACTAATAGAACTAAGTACAGAA
 AATTTAGTTTTAGGTGGTTGTAGCTGATGAGTTATTACCTCATAGAGACTATAATATTCTA
 TTTGGTATTATATTATTGATGTTTGCTGTTCTTCAAACATTTAAATCAAGCTTTGGACTAA
 TTATGCTAATTTGTGAGTTCTGATCACTTTTGAGCTCTGAAGCTTTGAATCATTAGTGGTG
 GAGATGGCCTTCTGGTAACTGAATATTACCTTCTGTAGGAAAAGGTGGAAAATAAGCATCTA
 GAAGGTGTTGTGAATGACTCTGTGCTGGCAAAAATGCTTGAAACCTCTATATTTCTTTCGT
 TCATAAGAGGTAAGGTCAAATTTTTCAACAAAAGTCTTTAATAACAAAAGCATGCAGTTCTC
 TGTGAAATCTCAAATATTGTTGTAATAGTCTGTTTCAATCTTAAAAAGAATCA

FIGURE 46

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96889
><subunit 1 of 1, 339 aa, 1 stop
><MW: 36975, pI: 7.85, NX(S/T): 1
MAAACGPGAAGYCLLLGLHLFLLTAGPALGWNDPDRMLLRDVKALTLLHYDRYTTSRRLLDP
IPQLKCVGGTAGCDSYTPKVIQCQNKGWDGYDVQWECKTDLDIAYKFGKTVVSCEGYESS
EDQYVLRGSCGLEYNLDYTELGQLKESGKQHGFAFSFDYKKWSSADSCNMSGITIV
VLLGIAFVVYKFLFLSDGQYSPPPYSEYPPFSHRYQRFTNSAGPPPPGFKSEFTGPQNTGH
GATSGFGSAFTGQQGYENSGPGFWTGLGTGGILGYLFGSNRAATPFSDSWYYPSYPPSY
GTWNRAYSPLHGGSGSYSVCSNSDTKTRTASGYGGTRRR
```

Important features of the protein:

Signal peptide:

Amino acids 1-30

Transmembrane domain:

Amino acids 171-190

N-glycosylation site:

Amino acids 172-176

Glycosaminoglycan attachment sites:

Amino acids 244-248;259-263;331-335

Tyrosine kinase phosphorylation site:

Amino acids 98-106

N-myristoylation sites:

Amino acids 68-74;69-75;131-137;241-247;
247-253;266-272;270-276;278-284;
312-318

FIGURE 47

CCCGGAGCCGGGGAGGGAGGGAGCGAGGTTTCGGACACCGGCGGGCTGCCTGGCCTTTCCA
TAGAGCCCGCGGCGGACCCCTCCGCGCCCCCTCTCGCTCTGCCTCTCCCTCTGCCTCTGCCTC
TGCCTGGCCGCGGCTCTGGGAAGTGCAGTCCGGGTCGTGTAGGGATAAAAAGAACTGTAA
GGTGGTCTTTTCCCAGCAGGAAGTGCAGGAAGCGGCTAACACCCCTGCAGTACCATGTCACCTC
AGGAGAAAGGGACCGAAAGTGCCTTTGAAGGAGAATACACACATCACAAAGATCCTGGAATA
TATAAATGTGTTGTTTGTGGAAGTCCATTGTTTAAAGTCAGAAACCAAATTTGACTCCGGTTC
AGGTTGGCCTTCATTCCACGATGTGATCAATTCTGAGGCAATCACATTCACAGATGACTTTT
CCTATGGGATGCACAGGGTGGAAACAAGCTGCTCTCAGTGTGGTGCTCACCTTGGGCACATT
TTTGATGATGGGCCTCGTCCAAGTGGGAAAAGATACTGCATAAATTCGGCTGCCTTGTCTTT
TACACCTGCGGATAGCAGTGGCACCAGCGGAGGGAGGCAGTGGGGTCGCCAGCCCCGGCCAGG
CAGACAAAGCGGAGCTCT**TA**GAGTAATGGAGAGTATGGAACAAAGTGTACTTAATGCACAG
CTTATTAATAAAATCAAAATGTTATCTTAATAGATATATTTTTCAAAACTATAAGGGCA
GTTTTGTGCTATTGATATTTTTTCTTCTTTTGCTTAAACAGAAGCCCTGGCCATCCATGTAT
TTTGCAATTGACTAGATCAAGAACTGTTTATAGCTTTAGCAAATGGAGACAGCTTTGTGAAA
CTTCTTCACAAGCCACTTATACCCCTTGGCATTCTTTTCTTTGAGCACATGGCTTCTTTTGC
AGTTTTTCCCCCTTTGATTCAGAAAGCAGAGGGTTCATGGTCTTCAAACATGAAAATAGAGAT
CTCCTCTGCAGTGTAGAGACCAGAGCTGGGCAGTGCAGGGCATGGAGACCTGCAAGACACAT
GGCCTTGAGGCCTTTGCACAGACCCACCTAAGATAAGGTTGGAGTGTATGTTTTAATGAGACT
GTCAGCTTTGTGGAAGTTTGAAGCTAAGGTCATTTTTTTTTTCTCACTGAAAGGGTGTGA
AGGTCTAAAGTCTTTCCTTATGTTAAATTGTTGCCAGATCCAAAGGGGCATACTGAGTGTG
TGGCAGAGAAGTAAACATTACCACACTGTTAGGCCTTTATTTTATTTTATTTTCCATCGAAA
GCATTGGAGGCCCAGTGCAATGGCTCACGCCTGTGATCCAGCACTTTGGGAGGCCAAGGCG
GGTGGATCACGAGGTCAGGAGATGGAGACCATCCTGGCTAACATGGTGAAACCCCGTCTCTA
CTAAAAATACGAAAAATTAGCCAGGCGTGGTGGTGGGCACCTGTAGTCCCAGCTACTCAGGAGG
CTGAGGCAGGAGAATGGCGTGAACCCGGAAGGCGGAGCTTGCAGTTAGCCGAGATCATGCCA
CTGCACTCCAGCCTACATGACAATGTGACACTCCATCTCAAAAAATAATAATAATAACAATA
TAAGAACTAGCTGGGCATGGTGGCGCATGCATGTAGTCCCAGCTACTCCTGAGGCTCAGTCA
GGAGAATCGCTTGAAGTTGGGAGGCGGAGGTTGCAGTGAAGTGAAGTGCATACCACTGCACTC
CAGCCTGAACAGAGTGAGATCCTGTCAA

FIGURE 48

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96898
><subunit 1 of 1, 192 aa, 1 stop
><MW: 20702, pI: 7.50, NX(S/T): 0
MSPRRTLPRPLSLCLSLCLCLCLAAALGSAQSGSCRDKKNCKVVFSSQQLRKRLTPLQYH
VTQEKGTESAFEGEYTHHKDPGIYKCVVCGTPLFKSETKFDSGSGWPSFHDVINSEAITF
TDDFSYGMHRVETSCSQCGAHLGHI FDDGPRPTGKRYCINSAALSFTPADSSGTAEGGSG
VASPAQADKAEL
```

Important features of the protein:

Signal peptide:

Amino acids 1-24

Glycosaminoglycan attachment site:

Amino acids 102-106

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 52-56

N-myristoylation sites:

Amino acids 28-34;66-72;82-88;139-145;
173-179;178-184

Amidation site:

Amino acids 153-157

FIGURE 49

CCCAAAGAGGTGAGGAGCCGGCAGCGGGGGCGGCTGTAACGTGTGAGGAAGGCTGCAGAGTGG
CGACGTCTACGCCGTAGGTTGGAGGCTGTGGGGGGTGGCCGGGCGCCAGCTCCCAGGCCGCA
GAAGTGACCTGCGGTGGAGTTCCCTCCTCGCTGCTGGAGAACGGAGGGAGAAGGTTGCTGGC
CGGGTGAAAGTGCCCTCCCTCTGCTTGACGGGGCTGAGGGGGCCGAAGTCTAGGGCGTCCGTA
GTCGCCCCGGCCTCCGTGAAGCCCCAGGTCTAGAGATATGACCCGAGAGTGCCCATCTCCGG
CCCCGGGGCCTGGGGCTCCGCTGAGTGGATCGGTGCTGGCAGAGGCGGCAGTAGTGTGTTGCA
GTGGTGCTGAGCATCCACGCAACCGTATGGGACCGATACTCGTGGTGCGCCGTGGCCCTCGC
AGTGCAGGCCTTCTACGTCCAATACAAGTGGGACCGGCTGCTACAGCAGGGAAGCGCCGTCT
TCCAGTTCCGAATGTCCGCAAACAGTGGCCTATTGCCCGCCTCCATGGTCATGCCTTTGCTT
GGACTAGTCATGAAGGAGCGGTGCCAGACTGCTGGGAACCCGTTCTTTGAGCGTTTTTGGCAT
TGTGGTGGCAGCCACTGGCATGGCAGTGGCCCTCTTCTCATCAGTGTTGGCGCTCGGCATCA
CTCGCCCACTGCCAACCAACACTTGTGTCATCTTGGGCTTGGCTGGAGGTGTTATCATTTAT
ATCATGAAGCACTCGTTGAGCGTGGGGGAGGTGATCGAAGTCCTGGAAGTCCTTCTGATCTT
CGTTTATCTCAACATGATCCTGCTGTACCTGCTGCCCCGCTGCTTACCCCTGGTGAGGCAC
TGCTGGTATTGGGTGGCATTAGCTTTGTCTCAACCAGCTCATCAAGCGCTCTCTGACACTG
GTGGAAAGTCAGGGGGACCCAGTGGACTTCTTCTGCTGGTGGTGGTAGTAGGGATGGTACT
CATGGGCATTTTCTCAGCACTCTGTTTGTCTTCATGGACTCAGGCACCTGGGCCTCCTCCA
TCTTCTTCCACCTCATGACCTGTGTGCTGAGCCTTGGTGTGGTCTTACCCTGGCTGCACCGG
CTCATCCGCAGGAATCCCCTGCTCTGGCTTCTTCAGTTTCTCTTCCAGACAGACACCCGCAT
CTACCTCCTAGCCTATTGGTCTCTGCTGGCCACCTTGGCCTGCCTGGTGGTGTGTACCAGA
ATGCCAAGCGGTATCTTCCGAGTCCAAGAAGCACCAGGCCCCCAACCATCGCCCGAAAGTAT
TTCCACCTCATTGTGGTAGCCACCTACATCCCAGGTATCATCTTTGACCGGCCACTGCTCTAT
GTAGCCGCCACTGTATGCCTGGCGGTCTTCATCTTCTGGAGTATGTGCGCTACTTCCGCAT
CAAGCCTTTGGGTCACTCTACGGAGCTTCCCTGTCCCTTTTTCTGGATGAACGAGACAGTG
GACCACTCATTCTGACACACATCTACCTGCTCCTGGGCATGTCTCTTCCCATCTGGCTGATC
CCCAGACCCTGCACACAGAAGGGTAGCCTGGGAGGAGCCAGGGCCCTCGTCCCCTATGCCGG
TGTCTGGCTGTGGGTGTGGGTGATACTGTGGCCTCCATCTTCGGTAGCACCATGGGGGAGA
TCCGCTGGCCTGGAACCAAAAAGACTTTTGAGGGGACCATGACATCTATATTTGCGCAGATC
ATTTCTGTAGCTCTGATCTTAATCTTTGACAGTGGAGTGGACCTAAACTACAGTTATGCTTG
GATTTTGGGGTCCATCAGCACTGTGTCCCTCCTGGAAGCATACACTACACAGATAGACAATC
TCCTTCTGCCTCTCTACCTCCTGATATTGCTGATGGCCTAGCTGTTACAGTGCAGCAGCACT
GACGGAGGAAACAGACATGGGGAGGGTGAACAGTCCCCACAGCAGACAGCTACTTGGGCATG
AAGAGCCAAGGTGTGAAAAGCAGATTTGATTTTTTCAGTTGATTCAGATTTAAATAAAAAGC
AAAGCTCTCCTAGTTCTA

FIGURE 50

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA97003
><subunit 1 of 1, 538 aa, 1 stop
><MW: 59268, pI: 8.94, NX(S/T): 1
MTRECPSPAPGPGAPLSGSVLAEEAAVVFVAVLSIHATVWDRYSWCAVALAVQAFYVQYKW
DRLLQQGSAVFQFRMSANSGLLPASMVMPLLGLVMKERCQTAGNPFFERFGIVVAATGMA
VALFSSVLALGITRFPVTNTCVILGLAGGVIIYIMKHSLSVGEVIEVLEVLLIFVYLNMI
LLYLLPRCFTPGEALLVLGGISFVLNQLIKRSLTLVESQGDVPDFFLLVVVVGMVLMGIF
FSTLFVFMDSGTWASSIFFHLMTCVLSLGVLPWLHRLIRRNP LLWLLQFLFQTDTRIYL
LAYWSLLATLACLVLVLYQNAKRSSSESKKHQAPTIARKYFHLIVVATYIPGIIFDRPLLY
VAATVCLAVFIFLEYVRYFRIKPLGHTLRSFLSLFLDERDSGPLILTHIYLLLGMSLPIW
LIPRPCTQKGS LGGARALVPYAGVLAVGVGDTVASIFGSTMGEIRWPGTKKTFEGTMTSI
FAQIISVALILIFDSGVDLNYSYAWILGSISTVSLLEAYTTQIDNLLLPLYLLILLMA
```

Important features of the protein:

Signal peptide:

Amino acids 1-36

Transmembrane domains:

Amino acids 77-95;111-133;161-184;225-248;
255-273;299-314;348-373;406-421;
435-456;480-497

N-glycosylation sites:

Amino acids 500-504

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 321-325

N-myristoylation sites:

Amino acids 13-19;18-24;80-86;111-117;
118-124;145-151;238-244;251-257;
430-436;433-439;448-454;458-464;
468-474;475-481;496-502;508-514

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 302-313

FIGURE 51

GCTCTATGCCCGCTACCTTGCTCTCGCCGCTGCTGCCGGAGCCGAAGCAGAGAAGGCAGCGGGTCCCGTGACCG
 TCCCGAGAGCCCCGCGCTCCCGACCAAGGGGGCGGGGGCGGCCCGGGGAGGGCGGGGCAGGGGGCGGGGGAAGA
 AAGGGGGTTTTGTGCTGCGCCGGGAGGGCGGGCGCCCTCTTCCGAATGTCTGCGCCCCAGCCTCTCCTCACG
 CTCGCGCAGTCTCCGCCGAGTCTCAGCTGCAGCTGCAGGACTGAGCCGTGCACCCGGAGGAGACCCCGGAGG
 AGGCGACAACTTCGCACTGCCCGACCCAAACCCAGCCCTGGGTAGCCTGCAGCAATGGGCCAGCTGTTCCCTGC
 CCCTGCTGGCAGCCCTGGTCCTGGCCAGGCTCCTGCAGCTTTAGCAGATGTTCTGGAAGGAGACAGCTCAGAG
 GACCGCGCTTTTCGCGTGCGCATCGCGGGCGACGCGCCACTGCAGGGCGTGCTCGGCGGGCGCCCTCACCATCCC
 TTGCCACGTCCACTACCTGCGGCCACCGCCGAGCCGCGGGCTGTGCTGGGCTCTCCGCGGGTCAAGTGGACTT
 TCCTGTCCCGGGGCGGGAGGCAGAGTGCTGGTGGCGGGGAGTGCGCGTCAAGGTGAACGAGGCCTACCGG
 TTCCGCGTGCGACTGCCTGCGTACCCAGCGTCTGCTCACCAGCTCTCCCTGGCGCTGAGCGAGCTGCGCCCCAA
 CGACTCAGGTATCTATCGCTGTGAGGTCCAGCACGGCATCGATGACAGCAGCGACGCTGTGGAGGTCAAGGTCA
 AAGGGGTCTCTTTCTCTACCGAGAGGGCTCTGCCCGCTATGCTTTCTCCTTTCTGGGGCCAGGAGGCCTGT
 GCCCGCATTTGGAGCCACATCGCCACCCCGAGCAGCTCTATGCCGCTACCTTGGGGGCTATGAGCAATGTGA
 TGCTGGCTGGCTGTGCGATCAGACCGTGAGGTATCCATCCAGACCCACGAGAGGCCTGTTACGGAGACATGG
 ATGGCTTCCCGGGGTCCGGAATATGGTGTGGTGGACCCGGATGACCTCTATGATGTGTACTGTTATGCTGAA
 GACCTAAATGGAGAATGTTCTGGGTGACCTCCAGAGAAGCTGACATTGGAGGAAGCACGGGGCTACTGCCA
 GGAGCGGGTGCAGAGATTGCCACCACGGGCCAACTGTATGCAGCCTGGGATGGTGGCTGGACCACTGCAGCC
 CAGGGTGCTAGCTGATGGCAGTGTCGCTACCCCATCGTCACACCCAGCCAGCGCTGTGGTGGGGGCTTGCCT
 GGTGTCAAGACTCTCTCCTCTTCCCAACCAGACTGGCTTCCCAATAAGCACAGCCGCTTCAACGTCTACTG
 CTTCCGAGACTCGGCCCAGCCTTCTGCCATCCCTGAGGCTTCCAACCCAGCCTCCAACCCAGCCTCTGATGGAC
 TAGAGGCTATCGTCACAGTGACAGAGACCTGGAGGAAGTGCAGCTGCCTCAGGAAGCCACAGAGAGTGAATCC
 CGTGGGGCCATCTATCTCCATCCCCATCATGGAGGACGGAGGAGTGGAAGCTCCACTCCAGAAGACCCAGCAGA
 GGCCCCTAGGACGCTCTAGAAATTTGAAACACAATCCATGGTACCGCCACGGGGTCTCAGAAGAGGAAGGTA
 AGGCATTGGAGGAAGAAGAGAAATATGAAGATGAAGAAGAGAAAGAGGAGGAAGAAGAGGAGGAGGTGGAG
 GATGAGGCTCTGTGGGCATGGCCAGCGAGCTCAGCAGCCGGGGCCCTGAGGCCTCTCTCCCACTGAGCCAGC
 AGCCCAGGAGAAGTCACTCTCCAGGCGCCAGCAAGGGCAGTCTGCAGCCTGGTGCATCACCCTTCCCTGATG
 GAGAGTCAGAAGCTTCCAGGCCCTCCAGGGTCCATGGACCACCTACTGAGACTTGGCCACTCCAGGGGAGAG
 AACCTAGCATCCCCATCACCTTCCACTCTGGTTGAGGCAAGAGAGGTGGGGGAGGCAACTGGTGGTCTGAGCT
 ATCTGGGGTCCCTCGAGGAGAGAGCGAGGAGACAGGAAGCTCCGAGGGTGCCCTTCCCTGCTTCCAGCCACAC
 GGGCCCCCTGAGGGTACCAGGAGCTGGAGGCCCTCTGAAGATAATTCTGGAAGAAGTGGCCAGCAGGGACC
 TCAGTGCAGGCCCAGCCAGTGCTGCCCACTGACAGCGCCAGCCGAGGTGGAGTGGCCGTGGTCCCCGCATCAGG
 TGACTGTGTCCCCAGCCCTGCCACAATGGTGGGACATGCTTGGAGGAGGAGGAAGGGTCCGCTGCCTATGTC
 TGCTGGCTATGGGGGGACCTGTGCGATGTTGGCCTCCGCTTCTGCAACCCCGGCTGGGACGCTTCCAGGGC
 GCCTGCTACAAGCACTTTTCCACACGAAGGAGCTGGGAGGAGGCAGAGACCCAGTGCCGGATGTACGGCGCGCA
 TCTGGCCAGCATCAGCACACCCGAGGAACAGGACTTCATCAACAACCGGTACCGGGAGTACCAGTGGATCGGAC
 TCAACGACAGGACCATCGAAGGCGACTTCTTGTGGTGGATGGCGTCCCCCTGCTCTATGAGAACTGGAACCT
 GGGCAGCCTGACAGTACTTCTGTCTGGAGAGAACTGCGTGGTTCATGGTGTGGCATGATCAGGGACAAATGGAG
 TGACGTGCCCTGCAACTACCACCTGTCTACACCTGCAAGATGGGGCTGGTGTCTGTGGGCCGCCACCGGAGC
 TGCCCTGGCTCAAGTGTTCGGCCGCCACGGCTGCGCTATGAGGTGGACACTGTGCTTTCGCTACCGGTGCCGG
 GAAGGACTGGCCAGCGCAATCTGCCGCTGATCCGATGCCAAGAGAACGGTCGTTGGGAGGCCCCCAGATCTC
 CTGTGTGCCAGAAGACCTGCCGAGCTCTGCACCCAGAGGAGGACCCAGAAGGACGTGAGGGAGGCTACTGG
 GACGCTGGAAGGCGCTGTTGATCCCCCTTCCAGCCCCATGCCAGGTCCCTTAGGGGGCAAGGCCTTGAACACTGCCG
 GCCACAGCACTGCCCTGTACCCAAATTTTCCCTCACACCTTGCGCTCCCGCCACCACAGGAAGTGACAACATG
 ACGAGGGGTGGTGTGGAGTCCAGGTGACAGTTCCTGAAGGGCTTCTGGGAAATACCTAGGAGGCTCCAGCCC
 AGCCCAGGCCCTCTCCCCCTACCCTGGGCACAGATCTTCCATCAGGGCCGGAGTAAATCCCTAAGTGCCTCAA
 CTGCCCTCTCCCTGGCAGCCATCTTGTCCCTCTATTCTCTAGGGAGCACTGTGCCACTCTTCTGGGTTTT
 CCAAGGGAATGGCTTGCAGGATGGAGTGTCTGTAAATCAACAGGAAATAAACTGTGTATGAGCCCA

FIGURE 52

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA98565
><subunit 1 of 1, 911 aa, 1 stop
><MW: 99117, pI: 4.62, NX(S/T): 2
MAQLFLPLLAALVLAQAPAAADVLEGDSSEDRAFRVRIAGDAPLQGVLGALTIPCHVH
YLRPPPSRRRAVLGSPRVKWTFLSRGREAEVLVARGVRVKVNEAYRFRVALPAYPASLTDV
SLALSELRPNDSGIYRCEVQHIGIDDSSDAVEVKVKGVVFLYREGSARYAFSFGAQEACA
RIGAHIAIPEQLYAAAYLGGYEQCDAGWLSQDQTVRYPIQTPREACYGDMGDFPGVRNYGVV
DPDDLYDVYCYAEDLNGELFLGDPPEKLTLEEARAYCQERGAETATTGQLYAAWDGGLDH
CSPGWLADGSVRYPIVTPSQRCGGGLPGVKTLFLFPNQTFGFPNKHRSRNVYCFRDSAQPS
AIPSEASNPASNPASDGLEAIVTVTETLEELQLPQEATESESRGAIYSIPIMEDGGGGSST
PEDPAEAPRTLLEFETQSMVPPTGFSEEEGKALEEEEEKYEDEEEKEEEEEVEDEALW
AWPSELSSPGPEASLPTEPAAQEKSLSQAPARAVLQPGASPLPDGESEASRPPRVHGPPT
ETLPTPRERNLASPSPSTLVEAREVGEATGGPELSGVPRGESEETGSSEGAAPSLPATRA
PEGTRELEAPSEDNSGRTAPAGTSVQAQPVLPDTSASRGGVAVVPASGDCVPSPCNNGGT
CLEEEEGVRCLCLPGYGGDLCDVGLRFCNPGWDAFQGACYKHFSTRRSWEEAETQCRMYG
AHLASISTPEEQDFINNRYREYQWIGLNDRTIEGDFLWSDGVPLLYENWNPGQPDYSYFLS
GENCVVMVWHDQGGQSDVPCNYHLSYCKMGLVSCGPPPELPLAQVFGRPRRLRYEVDTVL
RYRCREGLAQRNILPLIRCQENGRWEAPQISCVPRRPARALHPEEDPEGRQGRLLGRWKAL
LIPPSPPMPGP
```

Important features of the protein:

Signal peptide:

Amino acids 1-15

N-glycosylation sites:

Amino acids 130-134;337-341

Tyrosine kinase phosphorylation sites:

Amino acids 128-136;451-460

N-myristoylation sites:

Amino acids 47-53;50-56;133-139;142-148;
174-180;183-189;281-287;288-294;
297-303;324-330;403-409;414-420;
415-421;576-582;586-592;677-683;
684-690;720-726;772-778;811-817

EGF-like domain cysteine pattern signature:

Amino acids 670-682

C-type lectin domain signature:

Amino acids 784-809

Immunoglobulins and major histocompatibility complex proteins signature:

Amino acids 135-142

Link domain proteins:

Amino acids 166-216;264-314

Calcium-binding EGF-like domain proteins pattern proteins.

Amino acids 655-676

C-type lectin domain proteins:

Amino acids 791-800

FIGURE 53

CTGCCAGGTGACAGCCGCCAAGATGGGGTCTTGGGCCCTGCTGTGGCCTCCCTGCTGTTACCGGGCTGCTCG
TCCGACCCCGGGGACCATGGCCAGGCCAGTACTGCTCTGTGAACAAGGACATCTTTGAAGTAGAGGAGAAC
ACAAATGTACCCGAGCCGCTGGTGGACATCCAGCTCCCGGAGGGCCAGGAGGTGACCTCGGAGCCTTGTCCAC
CCCCTTTGCACTTTCCGATCCAGGGAACCAGCTGTTTCTCAACGTGACTCCTGATTACGAGGAGAAGTCACTGC
TTGAGGCTCAGCTGCTGTGTGAGAGCGGAGGCACATTGGTGACCCAGCTAAGGGTGTCTGTGTCAGTGTGGAC
GTCAATGACAATGCCCCGAATTCCTTTAAGACCAAGGAGATAAGGGTGGAGGAGGACACGAAAGTGAATC
CACCGTCATCCCTGAGACGCAACTGCAGGCTGAGGACCGGACAAGGACGACATTCGTGTTCTACACCTCCAGG
AAATGACAGCAGGTGCCAGTGACTACTTCTCCCTGGTGAGTGTAACCGTCCCGCCCTGAGGCTGGACCGGCC
CTGGACTTCTACGAGCGGCCGAACATGACCTTCTGGCTGCTGGTGCGGGACACTCCAGGGGAGAATGTGGAACC
CAGCCACACTGCCACCGCCACACTAGTGCTGAACGTGGTGCCCGCGGACCTGCGGCCCCCGTGGTTCCCTGCCCT
GCACCTTCTCAGATGGCTACGCTGCAATTCAGCTCAGTACCACGGGGCTGTCCACGGGGCACATACTGCCA
TCTCCCTCGTCTCGCTCCCGGACCCATCTACGCTGAGGACGGAGACCGCGCATCAACCAGCCCATCATCTA
CAGCATCTTTAGGGGAAACGTGAATGGTACATTCATCATCCACCCAGACTCGGSCAACCTCACCCTGGCCAGGA
GTGTCCCCAGCCCCATGACCTTCTTCTGTGTGGTGAAGGGCCAACAGCCGACCTTGCCCCGTACTCAGTGACC
CAGGTCAACGTGGAGGCTGTGGCTGCGGCGGGGAGCCCGCCCTTCTCAGCCTCTGAGGATCCAGGCTCAGGCT
GGCGCGTGGCGCTGGAGCGGGCGTGTGTGTCAGGATGACAGCTGCGCCTTCTCAGCCTCTGAGGATCCAGGCTC
AGGACCCGGAGTTCTCGGACCTCAACTCGGCCATCACATATCGAATTACCAACCACTCACACTTCCGGATGGAG
GGAGAGGTTGTGCTGACCAACCACCACTGGCACAGGCGGGAGCCTTCTACGCAGAGGTTGAGGCCCCAACAC
GCTGACCTCTGGACAGCAACCACAGTCATTGAGATCAAGTTTCCGAACAGGAGCCCCCTCCACAGAGGCTG
GAGGAACAACCTGGGCCCTGGACAGCAACCACAGTCATTGAGATCAAGTTTCCGAACAGGAGCCCCCTCCACAGAGGCTG
AGGACCACTCTGGGGGAGGCACAGGCCCTCATCCACCTCTGGCACAACTCTGAGGCCACCAACCTCGTCCAC
ACCCGGGGGGCCCCCGGTGACAGAAAACAGCACCTCCACCAACCAGCCACTCCCGGTGGGGACACAGCACAGA
CCCCAAGCCAGGAACCTCTCAGCCGATGCCCCCGGTGTGGGAACCAGCACCTCCACCAACCAGCCACACCC
AGTGGGGGACAGCACAGACCCAGAGCCAGGAACCTCTCAGCCGATGCCCCCAGTATGGGAACAGCACCTC
CCACCAACCAGCCACACCCGGTGGGGGACAGCACAGACCCAGAGGCAGGAACCTCTCAGCCGATGCCCCCG
GTATGGGAACAGCACCTCCACCAACCAACCAACCCGGTGGGGGACAGCACAGACCCAGAGCCAGGAAC
TCTCAGCCGATGCCCCCTCAGCAAGAGCACCCATCTTCAGGTGGCGGGCCCTCGGAGGACAAGCGCTTCTCGGT
GGTGGATATGGCGGGCCCTGGGCGGGGTGCTGGGTGCGGTGCTGCTGCTGGCTCTCTTGGCTCGCGCTCCTTG
TCCACAAGCACTATGGCCCCGGCTCAAGTGCTGCTCTGGCAAAGCTCCGGAGCCCCAGCCCCAAGGCTTTGAC
AACCAGGCGTTCTCCTGACCAAGGCCAACTGGGCGCCCGTCCCGAGCCCCACGCACGACCCCAAGCCCGC
GGAGGCACCGATGCCCGCAGAGCCCGCACCCCGGCCCTGCCCTCCCGAGGCGGTGCCCTTGAGCCCCCGCAG
CGGCCCGAGCTGGCGGAAGCCCCACGGCGGTGAGGTCCATCCTGACCAAGCAGCGCGCGCGGAGGGCGGGTAC
AAGGCCGTCTGGTTTGGCGAGGACATCGGGACGGAGGCAGACGTGGTCTCTCAACGCGCCCAACCTGGACGT
GGATGGCGCCAGTGACTCCCGGAGCGGCAGAGGGCGAGGGCGCGGGGAGGGGTGGGGGTCCCTACGATGCAC
CCGTGGTGATGACTCCTACATCTAAGTGGCCCTCCACCTCTCCCCAGCCGCACGGGCACTGGAGGTCTCG
CTCCCCAGCCTCCGACCCGAGGCGAGAATAAAGCAAGGCTCCCGAAACCCAGGCCATGGCGTGGGCGAGCGCG
TGGGTCCCTGGGGGGCCCATTCACCTCAGTCCCCTGTGCTCATTAGCGCTTGAGCCAGGTGTGCAGATGAGGCG
GTGGGTCTGGCCACGCTGTCCCAACCCAAAGCTGCAGCACTTCCCGTAAACCACTGCAGTGCCCGCGCCTT
CCCGAGGCTCTGTGCCAGCTAGTCTGGGAAGTTCCTCTCCCGCTCTAACCACAGCCCGAGGGGGGCTCCCTCC
CCGACCTGCACCAAGAGATCTCAGGCACCGGCTCAACTCAGACCTCCCGCTCCCGACCTACACAGAGATTGC
CTGGGGAGGCTGAGGAGCCGATGCAAAACCCCAAGGCGACGCACTTGGGAGCCGGTGGTCTCAAACACCTGCCG
GGGGTCCCTAGTCCCCTTCTGAAATCTACATGCTTGGGTGGAGCGCAGCAGTAAACACCTTGCCAGTGACCTG
GACTGAGGCGCGCTGGGGTGGGTGCGCGGTGTGGCTGAGCAGGAGCCAGACCAGGAGGCTAGGGGTGAGAG
ACATTTCCCTCGTGTCTCCAAAGCCAGAGCCAGGCTGGGCGCCCATGCCAGAACCATCAAGGGATCCCT
TGCGGCTTGTAGCACTTTCCCTAATGGAAATACACCATTAATTCCTTTCCAAATGTTTT

FIGURE 54

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA102846
><subunit 1 of 1, 839 aa, 1 stop
><MW: 87546, pI: 4.84, NX(S/T): 8
MGSWALLWPPLLFTGLLVRPPGTMAQAQYCSVNKDIFEVEENTNVTEPLVDIHVPEGQEV
TLGALSTPFAFRIQGNQLFLNVTPDYEEKSLLEAQLLCQSGGTLVTQLRVFVSVLDVNDN
APEFPFKTKAIRVEEDTKVNSTVIPETQLQAEDRDKDDILFYTLQEMTAGASDYFSLVSV
NRPALRLDRPLDFYERPNMTFWLLVRDTPGENVEPSHTATATLVNLVVPADLRPPWFLPC
TFSGQYVCIQAQYHGAAPTGHILPSPLVLRPGPIYAEDGDRGINQPIIYSIFRGNVNGTF
IIHPDSGNLTVARSVPSPMTFELLVKGQQADLARYSVTQVTVEAVAAAGSPPRFPQSLYR
GTVARGAGAGVVVKDAAAPSQPLRIQAQDPEFSDLNSAITYRITNHSFRMEGEVVLTTT
TLAQAGAFYAEVEAHNTVTSGTATTVIEIQVSEQEPPSTEAGGTTGPWTSTTSEVPRPPE
PSQGPSTTSSGGGTGPHPPSGTTLRPPTSSTPGGPPGAENSTSHQPATPGGDTAQTTPKPG
TSQPMPPGVGTSTSHQPATPSGGTAQTPEPGTSQPMPPSMGTSTSHQPATPGGGTAQTPE
AGTSQPMPPGMGTSTSHQPTTPGGGTAQTPEPGTSQPMPLSKSTPSSGGGPPSEDKRFESV
DMAALGGVLGALLLLALLGLAVLVHKHYGPRLKCCSGKAPEPQPQGFNDQAFLPDHKANW
APVPSPTHDPKPAEAPMPAEPAPPGPASPGGAPEPAAAARAGGSPTAVRSILTKERRPEG
GYKAVWFGEDIGTEADVVLNAPTLDVDGASDSGSGDEGEGAGRGGGPYDAPGGDDSYI
```

Important features of the protein:

Signal peptide:

Amino acids 1-25

Transmembrane domain:

Amino acids 662-684

N-glycosylation sites:

Amino acids 44-48;140-144;198-202;297-301;
308-312;405-409;520-524

Glycosaminoglycan attachment sites:

Amino acids 490-494;647-651;813-817

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 655-659

Tyrosine kinase phosphorylation sites:

Amino acids 154-163;776-783

N-myristoylation sites:

Amino acids 57-63;102-108;255-261;294-300;
366-372;426-432;441-447;513-519;
517-523;530-536;548-554;550-556;
581-587;592-598;610-616;612-618;
623-629;648-654;666-672;667-673;
762-768;763-769;780-786;809-815;
821-827;833-839

Cadherins extracellular repeated domain signature:

Amino acids 112-123

FIGURE 55

[illegible]

FIGURE 56

MVGFGANRRAGRLPSLVLVLLVVIVVLA FNYSISSRHVLLQEEVAELQCQVQRTTEVAR
GRLEKRNSDLLLLVDTHKKQIDQKEADYGR LSSRLQAREGLGKRCEDDKVKLQNNISYQM
ADIHHLKEQLAELRQEFRLRQEDQLQDYRKNN TYLVKRLEYESFQCGQQMKE LRAQHEENI
KKLADQFLEEQKQETQKIQSNDGKELDINNQVVPKNIPKVAENVADKNEEPSSNHIPHGK
EQIKRGGDAGMPGIEENDLAKVDDLPPALRKPPISVSQHESHQAISHLPTGQPLSPNMPP
DSHINHNGNPGTSKQNPSSPLQRLIPGSNLDSEPRIQTDILKQATKDRVSD FHKLKQNDE
ERELQMDPADYGKQHFNDVL

Important features of the protein:

Signal peptide:

1-29

Transmembrane domain.

None

N-glycosylation site.

115-119

150-154

cAMP- and cGMP-dependent protein kinase phosphorylation site.

65-69

N-myristoylation site.

246-252

253-259

308-314

Amidation site.

101-105

FIGURE 57

GGATGGGCGAGCAGTCTGAATGCCAGAATGGATAACCGTTTTGCTACAGCATTGTGAATTGC
TTGTGTGCTTAGCCTCATTTCACCATCTACATGGCAGCCTCCATTGGCACAGACTTCTGGT
ATGAATATCGAAGTCCAGTTCAAGAAAATTCCAGTGATTTGAATAAAAGCATCTGGGATGAA
TTCATTAGTGATGAGGCAGATGAAAAGACTTATAATGATGCACTTTTTCGATACAATGGCAC
AGTGGGATTGTGGAGACGGTGTATCACCATACCCAAAAACATGCATTGGTATAGCCCACCAG
AAAGGACAGAGTCATTTGATGTGGTCACAAAATGTGTGAGTTTCACACTAACTGAGCAGTTC
ATGGAGAAATTTGTTGATCCCGGAAACCACAATAGCGGGATTGATCTCCTTAGGACCTATCT
TTGGCGTTGCCAGTTCCTTTTACCTTTTGTGAGTTTAGGTTTGATGTGCTTTGGGGCTTTGA
TCGGACTTTGTGCTTGCAATTTGCCGAAGCTTATATCCCACCATTGCCACGGGCATTCTCCAT
CTCCTTGCAAGATACCATGCTGTGAAGTCCAGGCCACATGGAGGTGTCTGTGTAGATGCTCC
AGCTGAAATCCCAAGCTAAGCTCCCAACTGACAGCCAACATCATTTCAGCCATGTGTGGGA
GCCATCCTGGATGTCCAGCCTTAACAAGCCTTGAGAGGACTTCAGCCACAGCTATTATCTTA
CTACATCCTTGTGAGACTCTAATAAAGAACCAACTAGCTGAGCCCAATCAACCTATGGAAGT
ATAGAAATAAAATGAATTGTTGTTTTGTGCCGTT

FIGURE 58

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA102880
><subunit 1 of 1, 184 aa, 1 stop
><MW: 21052, pI: 5.01, NX(S/T): 3
MDNRFATAFVIACVLSLISTIYMAASIGTDFWYEURSPVQENSSDLNKS IWDEFISDEAD
EKTYNDALFRYNGTVGLWRR CITIPKNMHWYSPPERTESFDVVT KCVSFTLTEQFMEKFV
DPGNHNSGIDLLR TYLWRCQFLLPFVSLGLMCFGALIGLCACICRS LYPTIATGILHLLA
DTML
```

Important features of the protein:

Signal peptide:

Amino acids 1-20

Transmembrane domain:

Amino acids 142-163

N-glycosylation sites:

Amino acids 42-46;47-51;72-76;

N-myristoylation sites:

Amino acids 123-129;154-160;158-164

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 152-163

FIGURE 59

[illegible]

FIGURE 60

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA105782
><subunit 1 of 1, 156 aa, 1 stop
><MW: 17472, pI: 10.01, NX(S/T): 1
MAPARAGFCPLLLLLLLGLWVAEIPVSAKPKGMTSSQWFKIQHMQPSPQACNSAMKNINK
HTKRCKDLNTFLHEPFSSVAATCQTPKIACKNGDKNCHQSHGVPVSLTMCKLTSGKYPNCR
YKEKRQNKSYVVACKPPQKKDSQQFHLVPVHLDRLV
```

Important features of the protein:

Signal peptide:

Amino acids 1-22

N-glycosylation site:

Amino acids 127-131

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 139-143

N-myristoylation sites:

Amino acids 18-24;32-38

Pancreatic ribonuclease family signature:

Amino acids 65-72

Pancreatic ribonuclease family proteins:

Amino acids 49-93

FIGURE 61

CGGGTC**ATG**CGCCGCCGCCTGTGGCTGGGCCTGGCCTGGCTGCTGCTGGCGCGGGCGCCGGA
CGCCGCGGGAACCCCGAGCGCGTCGCGGGGACCGCGCAGCTACCCGCACCTGGAGGGCGACGTG
CGCTGGCGGGCGCCTCTTCTCCTCCACTCACTTCTTCCTGCGCGTGGATCCCGGCGGCCGCGT
GCAGGGCACCCGCTGGCGCCACGGCCAGGACAGCATCCTGGAGATCCGCTCTGTACACGTGG
GCGTCGTGGTCATCAAAGCAGTGTCTCAGGCTTCTACGTGGCCATGAACCGCCGGGGCCGC
CTCTACGGGTCGCGACTCTACACCGTGGACTGCAGGTTCCGGGAGCGCATCGAAGAGAACGG
CCACAACACCTACGCCTCACAGCGCTGGCGCCGCCGCGGCCAGCCCATGTTCTTGGCGCTGG
ACAGGAGGGGGGGCCCCGGCCAGGCGGCCGGACGCGGCGGTACCACCTGTCCGCCCACTTC
CTGCCCCTCCTGGTCTCC**TGAG**

FIGURE 62

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA108912
><subunit 1 of 1, 170 aa, 1 stop
><MW: 19663, pI: 11.81, NX(S/T): 0
MRRRLWLGLAWLLLARAPDAAGTPSASRGPRSYPHLEGDVRRRLFSSTHFFLRVDPGGR
VQGTRWRHGGQDSILEIRSVHVGVVVIKAVSSGFYVAMNRRGRLYGSRLYTVDCRFRERIE
ENGHNTYASQRWRRRGQPMFLALDRRGGRPGGRTRRYHLSAHFLPVLVS
```

Important features of the protein:

Signal peptide:

Amino acids 1-17

N-myristoylation site:

Amino acids 22-28

HBGF/FGF family proteins:

Amino acids 74-125;139-166

FIGURE 63

ATCCCTCGACCTCGACCCACGCGTCCGCTGGAAGGTGGCGTGCCCTCCTCTGGCTGGTACCA
TGCAGCTCCCCTGGCCCTGTGTCTCGTCTGCCTGCTGGTACACACAGCCTTCCGTGTAGTG
GAGGGCCAGGGGTGGCAGGCGTTCAAGAATGATGCCACGGAAATCATCCCCGAGCTCGGAGA
GTACCCCGAGCCTCCACCGGAGCTGGAGAACAAAGACCATGAACCGGGCGGAGAACGGAG
GGCGGCCTCCCCACCACCCCTTTGAGACCAAAGACGTGTCCGAGTACAGCTGCCGCGAGCTG
CACTTCACCCGCTACGTGACCGATGGGCGCTGCCGCGAGCGCCAAGCCGGTCACCGAGCTGGT
GTGCTCCGGCCAGTGCGGCCCCGGCGCGCCTGCTGCCAACGCCATCGGCCGCGGCAAGTGGT
GGCGACCTAGTGGGCCCCGACTTCCGCTGCATCCCCGACCGCTACCGCGCGCAGCGCGTGCAG
CTGCTGTGTCCCGGTGGTGAGGCGCGCGCGCGCGCAAGGTGCGCCTGGTGGCCTCGTGCAA
GTGCAAGCGCCTCACCCGCTTCCACAACCAGTCGGAGCTCAAGGACTTCGGGACCGAGGCCG
CTCGGCCGAGAGAAGGGCCGGAAGCCGCGGCCCCGCGCCCCGGAGCGCCAAAGCCAACCAGGCC
GAGCTGGAGAACGCCTACT**TAG**AGCCCCGCCCCGCGCCCCCTCCCCACCGGCGGGCGCCCCGGCCC
TGAACCCGCGCCCCACATTTCTGTCTCTGCGCGTGGTTTGATTGTTTATATTTTATTGTAA
ATGCTGCAACCCAGGGCAGGGGGCTGAGACCTTCCAGGCCCTGAGGAATCCCGGGCGCCGG
CAAGGCCCCCCCTCAGCCCGCCAGCTGAGGGGTCCACGGGGCAGGGGAGGGGAATTGAGAGTC
ACAGACACTGAGCCACGCAGCCCCGCTCTGGGGCGCCTACCTTTGCTGGTCCCCTTTCAG
AGGAGGCAGAAATGGAAGCATTTTCACCGCCCTGGGGTTTTAAGGGAGCGGTGTGGGAGTGG
GAAAGTCCAGGGACTGGTTAAGAAAGTTGGATAAGATTCCCCCTTGACCTCGCTGCCATC
AGAAAGCCTGAGGCGTGCCAGAGCACAAGACTGGGGGCAACTGTAGATGTGGTTTCTAGTCC
TGGCTCTGCCACTAATTTCTGTGTAACCTTGAACCTACACAATTCTCCTTCGGGACCTCAAT
TTCCACTTTGTAAAATGAGGGTGGAGGTGGGAATAGGATCTCGAGGAGACTATTGGCATATG
ATTCCAAGGACTCCAGTGCCTTTTGAATGGGCAGAGGTGAGAGAGAGAGAGAGAAAGAGAGA
GAATGAATGCAGTTGCATTGATTGAGTGCCAAGGTCACCTCCAGAATTCAGAGTTGTGATGC
TCTCTTCTGACAGCCAAAGATGAAAAACAACAGAAAAAAAAAAGTAAAGAGTCTATTTATG
GCTGACATATTTACGGCTGACAACTCCTGGAAGAAGCTATGCTGCTTCCCAGCCTGGCTTC
CCCGGATGTTTGGCTACCTCCACCCCTCCATCTCAAAGAAATAACATCATCCATTGGGGTAG
AAAAGGAGAGGGTCCGAGGGTGGTGGGAGGGGATAGAAATCACATCCGCCCCAACTTCCCAA
GAGCAGCATCCCTCCCCGACCCATAGCCATGTTTTAAAGTCACCTTCCGAAGAGAAGTGAA
AGGTTCAAGGACACTGGCCTTGCAGGCCCCGAGGGAGCAGCCATCACAACTCACAGACCAGC
ACATCCCTTTTGAGACACCGCCTTCTGCCACCACTCACGGACACATTTCTGCCTAGAAAAC
AGCTTCTTACTGCTCTTACATGTGATGGCATATCTTACACTAAAAGAATATTATTGGGGGAA
AACTACAAGTGCTGTACATATGCTGAGAACTGCAGAGCATAATAGCTGCCACCCAAAAAT
CTTTTGAAGATCATTTCCAGACAACCTCTTACTTTCTGTGTAGTTTTTAATTGTTAAAAA
AAAAAGTTTTTAAACAGAAGCACATGACATATGAAAGCCTGCAGGACTGGTCGTTTTTTTGGC
AATTCTTCCACGTGGGACTTGTCCACAAGAATGAAAGTAGTGGTTTTTAAAGAGTTAAGTTA
CATATTTATTTTCTCACTTAAGTTATTTATGCAAAAGTTTTTCTTGTAGAGAATGACAATGT
TAATATTGCTTTATGAATTAACAGTCTGTTCTTCCAGAGTCCAGAGACATTGTTAATAAAGA
CAATGAATCATGAAAAAAAAAAAAAAAAAAAAA

FIGURE 64

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA115253
<subunit 1 of 1, 213 aa, 1 stop
<MW: 24031, pI: 9.59, NX(S/T): 2
MQLPLALCLVCLLVHTAFRVVEGQGWQAFKNDATETIIPELGEYPEPPPELENNKTMNRAE
NGGRPPHHPFETKDVSEYSCRELHFTRYVTDGPCRSAPVTELVCSGQCGPARLLPNAIG
RGKWWRPSGPDFRCIPDRYRAQRVQLLCPGGEAPRARKVRLVASCKCKRLTRFHNQSELK
DFGTEAARPQKGRKPRPRARSAKANQAELENAY
```

Important features of the protein:

Signal peptide:

Amino acids 1-16

N-glycosylation sites:

Amino acids 53-57;175-179

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 168-172

N-myristoylation site:

Amino acids 183-189

Amidation site:

Amino acids 191-195

FIGURE 65

CCCACTCGGCGGTTTGGCGGGAGGGAGGGGCTTTGCGCAGGCCCCGCTCCCGCCCCGCCTCC
ATGCGGCCCCGCCCCGATTGCGCTGTGGCTGCGCCTGGTCTTGGCCCTGGCCCTGTCCGCCC
 CCGGGCTGTGGGGTGGGCCCCGGTCCGAGCCCCCATCTATGTCAGCAGCTGGGCGCTCCAGG
 TGTCCCAGGGTAACCGGGAGGTGAGCGCCTGGCACGCAAATTCGGCTTCGTCAACCTGGGG
 CCGATCTTCTCTGACGGGCAGTACTTTCACCTGCGGCACCGGGGCGTGGTCCAGCAGTCCCT
 GACCCCGCACTGGGGCCACCGCCTGCACCTGAAGAAAAACCCCAAGGTGCAGTGGTTCCAGC
 AGCAGACGCTGCAGCGGCGGGTGAAACGCTCTGTCTGGTGGCCACGGACCCCTGGTTCTCC
 AAGCAGTGGTACATGAACAGCGAGGCCCAACCAGACCTGAGCATCCTGCAGGCCTGGAGTCA
 GGGGCTGTGAGGCCAGGGCATCGTGGTCTCTGTGCTGGACGATGGCATCGAGAAGGACCACC
 CGGACCTCTGGGCCAACTACGACCCCTGGCCAGCTATGACTTCAATGACTACGACCCGGAC
 CCCCAGCCCCGCTACACCCCAAGAGAACCGGCACGGGACCCGCTGTGCTGGGGAGGT
 GGCCGCGATGGCCAACAATGGCTTCTGTGGTGTGGGGTTCGCTTCAACGCCCGAATCGGAG
 GCGTACGGATGGACGGTACCATCACCGATGTGTCATCGAGGCCAGTTCGTGAGCCTGCAG
 CCGCAGCACATCCACATTTACAGCGCCAGTGGGGTCCCGAGGACGAGGCCGCGCAGGTGGA
 CGGCCCCGGCATCCTCACCCGCGAGGCCTTCCGGCGTGGTGTGACCAAGGGCCGCGCGGGC
 TGGGCACGCTCTTCATCTGGGCCTCGGGCAACGGCGGCCTGCACTACGACAACCTGCAACTGC
 GACGGCTACACCAACAGCATCCACACGCTTTCGGTGGGCAGCACCACCCAGCAGGGCCGCGT
 GCCCTGGTACAGCGAAGCCTGCGCCTCCACCCTCACCCACCTACAGCAGCGGCGTGGCCA
 CCGACCCCAAGATCGTCACCACGGACCTGCATCACGGGTGCACAGACCAGCACACGGGCACC
 TCGGCCCTCAGCCCCACTGGCGGCGCGCATGATCGCCCTAGCGCTGGAGGCCAACCCGTTCT
 GACGTGGAGAGACATGCAGCACCTGGTGGTCCGCGCGTCCAAGCCGGCGCACCTGCAGGCCG
 AGGACTGGAGGACCAACGGCGTGGGGCGCCAAGTGAGCCATCACTACGGATACGGGCTGCTG
 GACGCCGGGCTGCTGGTGGACACCGCCCGCACCTGGTGCCCAACCAGCCGAGAGGAAGTG
 CGCCGTCCGGGTCCAGAGCCGCCCCACCCCATCCTGCCGCTGATCTACATCAGGGAAAACG
 TATCGGCCTGCGCCGGCCTCCACAACCTCCATCCGCTCGCTGGAGCACGTGCAGGCGCAGCTG
 ACGCTGTCTACAGCCGGCGCGGAGACC'TGGAGATCTCGCTCACCAGCCCCATGGGCACGCG
 CTCCACACTCGTGGCCATACGACCCTTGGACGTGAGCACTGAAGGCTACAACAACCTGGGTCT
 TCATGTCCACCCACTTCTGGGATGAGAACCACAGGGCGTGTGGACCCTGGGCCTAGAGAAC
 AAGGGCTACTATTTCAACACGGGGACGTTGTACCGCTACACGCTGCTGCTCTATGGGACGGC
 CGAGGACATGACAGCGCGGCCTACAGGCCCCAGGTGACCAGCAGCGCTGTGTGCAGCGGGAC
 ACAGAGGGGCTGTGCCAGGCGTGTGACGGCCCCGCCTACATCCTGGGACAGCTCTGCCTGGC
 CTACTGCCCCCGCGGTTCTTCAACCACACAAGGCTGGTGACCGCTGGGCCTGGGCACACGG
 CGGCGCCCGCGCTGAGGGTCTGCTCCAGCTGCCATGCCCTCCTGCTACACCTGCCGCGGCGGC
 TCCCCGAGGGACTGCACCTCCTGTCCCCCATCCTCCACGCTGGACCAGCAGCAGGGCTCCTG
 CATGGGACCCACCACCCCGACAGCCGCCCCCGGCTTAGAGCTGCCGCC'TGTCCCCACCACCG
 CTGCCCAGCCTCGGCCATGGTGCTGAGCCTCCTGGCCGTGACCCTCGGAGGCCCGCTCCTCT
 GCGGCATGTCCATGGACCTCCCACTATACGCCTGGCTCTCCCGTGCCAGGGCCACCCCCACC
 AAACCCAGGTCTGGCTGCCAGCTGGAACCT**TGA**AGTTGTCAGCTCAGAAAGCGACCTTGCCC
 CCGCCTGGGTCCCTGACAGGCACTGCTGCCATGCTGCCCTCCCCAGGCTGGCCCCAGAGGAGC
 GAGCACCAGCACCCGACGCCTGGCCTGCCAGGGATGGGCCCCGTGGAACCCGAAGCCTGGC
 GGGAGAGAGAGAGAGAGAAAGTCTCCTCTGCATTTTGGGTTTGGGCAGGAGTGGGCTGGGGGG
 AGAGGCTGGAGCACCCCAAAAGCCAGGGGAAAGTGGAGGGAGAGAAACGTGACACTGTCCGT
 CTCGGGCACCGCTCCAACCTCAGAGTTTGCAAATAAAGGTTGCTTAGAAGGTGAA

FIGURE 66

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA119302

><subunit 1 of 1, 755 aa, 1 stop

><MW: 82785, pI: 8.71, NX(S/T): 2

MRPAPIALWLRLLVLAALVRPRAVGWAPVRAPIYVSSWAVQVSQGNREVERLARKFGFVN
 LGPIFSDGQYFHLRHRGVVQQSLTPHWGHRHLKKNPKVQWFQQQTLQRRVKRSVVVPTD
 PWFSKQWYMNSEAQPDLSILQAWSQGLSGQGIVSVLDDGIEKDHPDLWANYDPLASYDF
 NDYDPPDPQPRYTSPKENRHGTRCAGEVAAMANNGFCGVGVAFNARIGGVRMLDGTITDVI
 EAQSLSLQPQHIIHYSASWGPEDDGRTVDGPGILTREAFFRGVTKGRGGLGTLFIWASGN
 GGLHYDNCNCDGYTNSIHTLSVGSTTQQGRVPWYSEACASTLTTTYSSGVATDPQIVTTD
 LHHGCTDQHTGTSASAPLAAGMIALALEANPFLTWRDMQHLVVRASKPAHLQAEDWRTNG
 VGRQVSHHYGYGLLDAGLLVDTARTWLPTQPQRKCAVRVQSRPTPILPLIYIRENVSACA
 GLHNSIRSLEHVQAQLTSLYSRRGDLEISLTSPMGTRSTLVAIRPLDVSTEGYNNWVFMS
 THFWDENPQGVWTLGLENKGYFNTGTLYRYTLLLYGTAEDMTARPTGPQVTSSACVQRD
 TEGLCQACDGPAYILGQLCLAYCPPRFENHTRLVTAGPGHTAAPALRVCSSCHASCYTCTC
 GGSPRDUCTSCPPSSTLDQQQGGSCMGPTTPDSRPLRAAACPHHRCPASAMVLSLLAVTLG
 GPVLCGMSMDLPLYAWLSRARATPTKPQVWLPAGT

Important features of the protein:

Signal peptide:

Amino acids 1-21

Transmembrane domain:

Amino acids 706-730

N-glycosylation sites:

Amino acids 475-479;629-633

Glycosaminoglycan attachment sites:

Amino acids 148-152;298-302

N-myristoylation sites:

Amino acids 151-157;200-206;217-223;219-225;
 282-288;288-294;371-377;432-438;
 481-487;515-521;603-609

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 586-597

Cell attachment sequence:

Amino acids 503-506

Serine proteases, subtilase family, aspartic acid active site:

Amino acids 154-166

Serine proteases, subtilase family, histidine active site:

Amino acids 199-210

Serine proteases, subtilase family, serine active site:

Amino acids 371-382

Cytochrome c family heme-binding site signature:

Amino acids 649-655

FIGURE 67

ATGAGGAAGCTCCAGGGCAGGATGGTTTACCTGCCTGGACAGCAAG**ATG**ATGGCTACACTAG
CCCCATTCTCTGGGCGCCTGGATTTGCCACAGATCTCCTCACCTCTTGCCCTTCACCTC
CTGCTGTACCTACAAGGTCTCCCGATTCTCATCTGCCCATAATCATGGACACAGCCCCAGG
ATGTGCAGGACTCTCAGGGACCATCTGGAGTTCCAGCTGGAATCTGGGCCTGGTGGAGTGGG
AGTGGGGCAGGGGCCTGCATTGGGCTGACTTAGAGAGCACAGTTATTCCATCCATATGGAAA
TAAACATTTTGATTCTGATC

FIGURE 68

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA119536
><subunit 1 of 1, 88 aa, 1 stop
><MW: 9645, pI: 5.45, NX(S/T): 0
MMATLAPILWAPGFAHQISSPLALHLLLYLQGLPDSHLPIIMDTAPGCAGLSGTIWSSSW
NLGLVEWEWGRGLHWADLESTVIPSIIWK
```

Signal sequence:

Amino acids 1-15

N-myristoylation sites:

Amino acids 32-38;50-56;53-59;72-78

FIGURE 69

TTTGACAGTGGGGTCTCCTCTGGCCTCCTGCCCTCCTGCTGCTGCTGCTGCTTCCATTGCT
GGCAGCCCAGGGTGGGGGTGGCCTGCAGGCAGCGCTGCTGGCCCTTGAGGTGGGGCTGGTGG
GTCTGGGGGCTCCTACCTGCTCCTTTGTACAGCCCTGCACCTGCCCTCCAGTCTTTTCCTA
CTCCTGGCCCAGGGTACCGCACTGGGGGCCGTCTGGGCCCTGAGCTGGCGCCGAGGCCT**CAT**
GGGTGTTCCCTTGGGCTTGGAGCTGCCTGGCTCTTAGCTTGGCCAGGCCTAGCTCTACCTC
TGGTGGCTATGGCAGCGGGGGGCAGATGGGTGCGGCAGCAGGGCCCCCGGGTGCGCCGGGGC
ATATCTCGACTCTGGTTGCGGGTCTGCTGCGCCTGTCACCCATGGCCTTCCGGGGCCCTGCA
GGGCTGTGGGGCTGTGGGGGACCGGGGTCTGTTTGCAGTGTACCCCAAACCAACAAGGATG
GCTTCCGCAGCCGCTGCCCGTCCCTGGGGCCCCGGCGGCGTAATCCCCGCACCACCCAACAC
CCATTAGCTCTGTTGGCAAGGGTCTGGGTCTGTGCAAGGGCTGGAAGTGGCGTCTGGCAGC
GGCCAGCCAGGGTTTAGCATCCCACTTGGCCCCGTGGGGCCATCCACACACTGGCCAGCTGGG
GCCTGCTTCGGGGTGAACGGCCCCACCCGAATCCCCCGGCTACTACCACGCAGCCAGCGCCAG
CTAGGGCCCCCTGCCTCCCGCCAGCCACTGCCAGGGACTCTAGCCGGGCGGAGGTACGCAC
CCGCCAGTCCCGGGGCCCTGCCCCCTGGAGGT**TAG**CTGACTCCAGCCCTTCCAGCCCAAATCT
AGAGCATTGAGCACTTTATCTCCACGACTCAGTGAAGTTTCTCCAGTCCCTAGTCCTCTCT
TTTACCCCACTTCTCAGTCTTGCTCACTTACCCAGGGCCAGCCCTTCCGACCTCTAGACA
GGCAGCCTCCTCAGCTGTGGAGTCCAGAGTCACTCTGTGTTCTCCTGGCGCTCCTCCCTA
AGTTATTGCTGTTGCGCCGCTGTGTGTGCTCATCCTCACCCTCATTGACTCAGGCCTGGGGC
CAGGGGTGGTGGAGGGTGGGAAGAGTCATGTTTTTTTTTCTCCTCTTTGATTTTGTTTTCTG
TCTCCCTTCCAACCTGTCCCTTCCCCCACCACAAAAAAXXXXXXXXXXXXXXXXXXXXXXXXXX
XXXXXXXXXXXXXXXXAAAAAAXX

FIGURE 70

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA119542
><subunit 1 of 1, 197 aa, 1 stop
><MW: 21992, pI: 12.18, NX(S/T): 0
MGVPLGLGAAWLLAWPGLALPLVAMAAGGRWVRQQGPRVRRGISRLWLRVLLRLSPMAFR
ALQGCGAVGDRGLEFALYPKTNKDGFRSRLPVPGFRRRNPRTTQHPLALLARVWVLCKGWN
WRLARASQGLASHLPPWAIHTLASWGLLRGERPTRIPRLLPRSQRQLGPPASRQPLPGTL
AGRRSRTRQSRALPPWR
```

Important features of the protein:

Signal peptide:

Amino acids 1-21

N-myristoylation sites:

Amino acids 2-8;6-12;146-152;178-184

Amidation site:

Amino acids 181-185

FIGURE 71

GTTTGGGGGTTGTTTGGGATTAGTGAAGCTACTGCCTTTGCCGCCAGCGCAGCCTCAGAGTT
TGATTATTTGCAATGTCAGGCTTTGAAAACCTAAACACGGATTTCTACCAGACAAGTTACAG
CATCGATGATCAGTCACAGCAGTCCTATGATTATGGAGGAAGTGGAGGACCCTATAGCAAAC
AGTATGCTGGCTATGACTATTTCGCAGCAAGGCAGATTTGTCCCTCCAGACATGATGCAGCCA
CAACAGCCATACACCGGGCAGATTTACCAGCCAACTCAGGCATATACTCCAGCTTCACCTCA
GCCTTTCTATGGAACAACCTTTGAGGATGAGCCACCTTTATTAGAAGAGTTAGGTATCAATTTT
GACCACATCTGGCAAAAAACACTAACAGTATTACATCCGTTAAAAGTAGCAGATGGCAGCAT
CATGAATGAACTGATTTGGCAGGTCCAATGGTTTTTTGCCTTGCTTTTGGAGCCACATTGC
TACTGGCTGGCAAAATCCAGTTTGGCTATGTATACGGGATCAGTGCAATTGGATGTCTAGGA
ATGTTTTGTTTATTAACTTAATGAGTATGACAGGTGTTTCATTTGGTTGTGTGGCAAGTGT
CCTTGGATATTGTCTTCTGCCCATGATCCTACTTTCCAGCTTTGCAGTGATATTTTCTTTGC
AAGGAATGGTAGGAATCATTCTCACTGCTGGGATTATTGGATGGTGTAGTTTTTCTGCTTCC
AAAATATTTATTTCTGCATTAGCCATGGAAGGACAGCAACTTTTAGTAGCATATCCTTGCGC
TTTGTTATATGGAGTCTTTGCCCTGATTTCCGTCTTTTGAAAATTTATCTGGGATGTGGACA
TCAGTGGGCCAGATGTACAAAAGGACCTTGAACCTCTTAAATTGGACCAGCAAACTGCTGCA
GCGCAACTCTCATGCAGATTTACATTTGACTGTTGGAGCAATGAAAGTAAACGTGTATCTCT
TGTTTCATTTTTATAGAACTTTTGCATACTATATTGGATTTACCTGCGGTGTGACTAGCTTTA
AATGTTTGTGTTTATACAGATAAGAAATGCTATTTCTTTCTGGTTCCTGCAGCCATTGAAAA
ACCTTTTTCTTGCAAATTTATAATGTTTTTGATAGATTTTTTATCAACTGTGGGAAACCAAAC
ACAAAGCTGATAACCTTTCTTAAAAACGACCCAGTCACAGTAAAGAAGACACAAGACGGCCG
GGCGTGGTAGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGCGGGCGGATCACAAG
GGCAGGAGATCGAGACCATCCTGGTTAACACGGTGAACCCCGACTCTACTAAAACCTACAAA
AAAAATTAGCTGGGCGTGGTGGCGGGCGCCTGTAGTCCCAGCTACTCAGGAGGCTGAGGCAG
GAGAAGTGTGAACCCAGGAGGCGGAGCTTGCAGTGAGCCGAGATCACACCACTGCACTCCAT
CCAGCCTGGGTGACAGGGTGAGACTCTGTCTCAAAAAAAAAAAAAAAAAAGGAGACACAAGACT
TACTGCAAAAATATTTTCCAAGGATTTAGGAAAGAAAAATTCCTTGTATTCTCAAGTCAG
GTAACCTCAAAGCAAAAAAGTGATCCAAATGTAGAGTATGAGTTTGCAC'TCCAAAAATTTGAC
ATTACTGTAAATTATCTCATGGAAATTTTGTCTAAAATTCAGAGATACGGGAAGTTCACAATC
TACCTCATTTGTAGACATGAAATGCGAACACTTACTTACATATTAATGTTAACTCAACCTTAG
GGACCTGGAATGGTTGCATTAATGCTATAATCGTTGGATCGCCACATTTCCCAAAAAATAATA
AAAAATCACTAACCTTTTTTAAGGAAAATATTTAAAGTTTTACAAAATTCATATTGCAAT
TATCAATGTAAAGTACATTTGAATGCTTATTAAAACCTTTCCCAATTAATTTT

FIGURE 72

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA143498
><subunit 1 of 1, 257 aa, 1 stop
><MW: 27989, pI: 4.16, NX(S/T): 1
MSGFENLNTDFYQTSYSIDDSQQSYDYGSGGPYSKQYAGYDYSQQGRFVPPDMMQPPQQ
PYTGQIYQPTQAYTPASPQPFYGNNEFEDEPPLLEELGINFDHIWQKTLTVLHPLKQVADGS
IMNETDLAGPMVFCLAFGATLLLAGKIQFGYVYGISAIGCLGMFCLLNLMSTGVSFQCV
ASVLGYCLLPMILLSSFAVIFSLQGMVGIILTAGIIGWCSFSASKIFISALAMEGQQLLV
AYPCALLYGVFALISVF
```

Transmembrane domain:

Amino acids 129-145;184-203

N-glycosylation sites:

Amino acids 123-127

N-myristoylation sites:

Amino acids 32-38;119-125;174-180;178-184;208-214

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 150-161;169-180

FIGURE 73

ACACTGGCCAAAACGCGGCTCGCCCTCGGCTGCGCTCGGCTCCCGCGGGCGCTCGGCCCCGA
GCCCCCTCCTCCCCCTACCCGCGGCGGACAGGGAGGAGCCA**ATG**GCTGGGCCTGCCATCCA
CACCGCTCCCATGCTGTTTCCTCGTCCTCCTGCTGCCCCAGCTGAGCCTGGCAGGCGCCCTTG
CACCTGGGACCCCTGCCCCGAACCTCCCTGAGAATCACATTGACCTCCAGGCCAGCGCTG
TGGACGCCTCAGGCCAGCCACCACCGCCGGCGGGGCCCCGGGCAAGAAGGAGTGGGGCCCAGG
CCTGCCCAGCCAGGCCAGGATGGGGCTGTGGTCAACGCCACCAGGCAGGCCTCCAGGCTGC
CAGAGGCTGAGGGGCTGCTGCCTGAGCAGAGTCCTGCAGGCCTGCTGCAGGACAAGGACCTG
CTCCTGGGACTGGCATTGCCCTACCCCGAGAAGGAGAACAGACCTCCAGGTTGGGAGAGGAC
CAGGAAACGCAGCAGGGAGCACAAGAGACGCAGGGACAGGTTGAGGCTGCACCAAGGCCGAG
CCTTGGTCCGAGGTCCCAGCTCCCTGATGAAGAAGGCAGAGCTCTCCGAAGCCCAGGTGCTG
GATGCAGCCATGGAGGAATCCTCCACCAGCCTGGCGCCCACCATGTTCTTTCTCACCACTT
TGAGGCAGCACCTGCCACAGAAGAGTCCCTGATCCTGCCCCGTCACCTCCCTGCGGCCCCAGC
AGGCACAGCCCAGGTCTGACGGGGAGGTGATGCCCACGCTGGACATGGCCTTGTTTCGACTGG
ACCGATTATGAAGACTTAAACCTGATGGTTGGCCCTCTGCAAAGAAGAAAGAGAAACACCG
CGGTAAACTCTCCAGTGATGGTAACGAAACATCACAGCCGAAGGGGAACCATGCGACCATC
ACCAAGACTGCCTGCCAGGGACTTGCTGCGACCTGCGGGAGCATCTCTGCACACCCCAACAAC
CGAGGCCTCAACAACAATGCTTCGATGACTGCATGTGTGTGGAAGGGCTGCGCTGCTATGC
CAAATTCACCCGGAACCGCAGGGTTACACGGAGGAAAGGGCGCTGTGTGGAGCCCGAGACGG
CCAACGGCGACACAGGGATCCTTCATCAACGTC**TAG**CGGCCCCGCGGGACTGGGGACTGAGCC
CAGGAGGTTTGCACAAGCCGGGCGATTTGTTTGTAACTAGCAGTGGGAGATCAAGTTGGGGA
ACAGATGGCTGAGGCTGCAGACTCAGGCCAGGACACTCAACCCC

FIGURE 74

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA145583
><subunit 1 of 1, 348 aa, 1 stop
><MW: 38536, pI: 8.24, NX(S/T): 1
MAGPAIHTAPMLFLVLLLPQLSLAGALAPGTPARNLPENHIDLPGPALWTPQASHHRRRG
PGKKEWGPGLPSQAQDGAVVTATRQASRLPEAEGLLPEQSPAGLLQDKDLLLGLALPYPE
KENRPPGWERTRKRSREHKRRRDRLRLHQGRALVRGPSSLMKKAELSEAQVLDAAMEESS
TSLAPTMMFFLTTFEAAPATEESLILPVTSLRPQQAQPRSDGEVMPITLDMALFDWTDYEDL
KPDGWPSAKKKEKHRGKLSSDGNETSPAEGEPCDHHQDCLPGTCCDLREHLCTPHNRGLN
NKCFDDCMCVEGLRCYAKFHRNRRVTRRKGRCVEPETANGDQGSFINV
```

Important features of the protein:

Signal peptide:

Amino acids 1-24

N-glycosylation site:

Amino acids 263-267

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 132-136;323-327

N-myristoylation sites:

Amino acids 77-83;343-349

Amidation site:

Amino acids 61-65

FIGURE 75

CAGAAGGGCAAAAACATTGACTGCCTCAAGGTCTCAAGCACCAGTCTTCACCGCGGAAAGCA
TGTTGTGGCTGTTCCAATCGCTCCTGTTTGTCTTCTGCTTTGGCCCAGGGAATGTAGTTTCA
CAAAGCAGCTTAACCCCATTTGATGGTGAACGGGATTCTGGGGGAGTCAGTAACCTCTTCCCCT
GGAGTTTCTGCAGGAGAGAAGGTCAACTTCATCACTTGGCTTTTCAATGAAACATCTCTTG
CCTTCATAGTACCCCATGAAACCAAAAGTCCAGAAATCCACGTGACTAATCCGAAACAGGGA
AAGCGACTGAACTTCACCCAGTCCTACTCCCTGCAACTCAGCAACCTGAAGATGGAAGACAC
AGGCTCTTACAGAGCCCAGATATCCACAAAGACCTCTGCAAAGCTGTCCAGTTACACTCTGA
GGATATTAAGACAACTGAGGAACATACAAGTTACCAATCACAGTCAGCTATTTCAGAATATG
ACCTGTGAGCTCCATCTGACTTGCTCTGTGGAGGATGCAGATGACAATGTCTCATTGAGATG
GGAGGCCTTGGGAAACACACTTTCAAGTCAGCCAAACCTCACTGTCTCTGGGACCCCAGGA
TTTCCAGTGAACAGGACTACACCTGCATAGCAGAGAATGCTGTGAGTAATTTATCCTTCTCT
GTCTCTGCCCAGAAGCTTTGCGAAGATGTTAAATTCATATACAGATACCAAAATGATTCT
GTTTATGGTTTCTGGGATATGCATAGTCTTCGGTTTTCATCATACTGCTGTACTTGTTTTGA
GGAAAAGAAGAGATTCCCTATCTTTGTCTACTCAGCGAACACAGGGCCCCGCAGAGTCCGCA
AGGAACCTAGAGTATGTTTCAGTGTCTCCAACGAACAACACTGTGTATGCTTCAGTCACTCA
TTCAAACAGGGAAACAGAAATCTGGACACCTAGAGAAAATGATACTATCACAATTTACTCCA
CAATTAATCATTCCAAAGAGAGTAAACCCACTTTTTCCAGGGCAACTGCCCTTGACAATGTC
GTG**TAA**GTTTGCTGAAAGGCCTCAGAGGAATTCGGGAATGACACGTCTTCTGATCCCATGAGA
CAGAACAAAGAACAGGAAGCTTGGTTCCTGTTGTTTCTGGCAACAGAATTTGAATATCTAGG
ATAGGATGATCACCTCCAGTCCTTCGGACTTAAACCTGCCTACCTGAGTCAAACACCTAAGG
ATAACATCATTTCCAGCATGTGGTTCAAATAATATTTTCCAATCCACTTCAGGCCAAAACAT
GCTAAAGATAACACACCAGCACATTGACTCTCTCTTTGATAACTAAGCAAATGGAATTATGG
TTGACAGAGAGTTTATGATCCAGAAGACAACCACCTTCTCTCCTTTTAGAAAGCAGCAGGATT
GACTTATTGAGAAATAATGCAGTGTGTTGGTTACATGTGTAGTCTCTGGAGTTGGATGGGCC
CATCCTGATACAAGTTGAGCATCCCTTGTCTGAAATGCTTGGGATTAGAAAATGTTTCAGATT
TCAATTTTTTTTTCAGATTTTGGGAATATTTGCATTATATTTAGCGGTTGAGTATCCAAATCCA
AAAATCCAAAATTCAAAATGCTCCAATAAGCATTTCCCTTGAGTTTCATTGATGTGATGCA
GTGCTCAAAATCTCAGATTTTGGAGCAATTTGGATATTGGATTTTTGGATTTGGGATGCTCA
ACTTGTACAATGTTTATTAGACACATCTCCTGGGACATACTGCCTAACCTTTTGGAGCCTTA
GTCTCCAGACTGAAAAAGGAAGAGGATGGTATTACATCAGCTCCATTGTTTGGCCAAGAA
TCTAAGTC

FIGURE 76

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA161000
><subunit 1 of 1, 332 aa, 1 stop
><MW: 37345, pI: 6.72, NX(S/T): 10
MLWLFQSLLEFVFCFGPGNVVSQSSLTPLMVNGILGESVTLPLEFPAGEKVNFFITWLFNET
SLAFIVPHETKSPEIHVTNPKQGKRLNFTQSYSLQLSNLKMEDTGSYRAQISTKTSAKLS
SYTLRILRQLRNIQVTNHSQLFQNMTCELHLTCSVEDADDNVVSFRWEALGNTLSSQPNLT
VSWDPRISSEQDYTCIAENAVSNLSFSVSAQKLCEDVKIQYTDTKMILEFMVSGICIVFGF
IILLLLVLRKRKRDLSLSLSTQRTQGPAESARNLEYVSVSPTNNTVYASVTHSNRETEIWTP
RENDTITIYSTINHSKESKPTFSRATALDNVV
```

Important features of the protein:

Signal peptide:

Amino acids 1-13

Transmembrane domain:

Amino acids 228-247

N-glycosylation sites:

Amino acids 58-62;87-91;137-141;144-148;161-165;
178-182;203-207;281-285;303-307;
313-317

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 251-255

Tyrosine kinase phosphorylation sites:

Amino acids 100-108;186-194

N-myristoylation sites:

Amino acids 17-23;105-111;170-176

Amidation site:

Amino acids 82-86

Immunoglobulin domain:

Amino acids 35-111

FIGURE 77

GATCCCTCGACCTCGACCCACGCGTCCGCTCTTTAATGCTTTCTTTTAAAGAGATCACCTTC
TGACTTCTCACAGAAGAGGTTAACTATTACCTGTGGGAAGTCAGAAGGTGATCTCTTTAATG
CTTTCTTTTAAAGAATTTTTCAAATTGAGACTAATTGCAGAGGTTCCAGTTGACCAGCATT
ATAGGAATGAAGACAAACACAGAGATGGTGTGTCTAAGAACTTCAAAAAGGTGTAGACCTCC
TGACTGAAGCATATTGGATTTATTTAATTTTTTTTCACTGTATTTCTGTCTCCTACAAGGGA
AAGT**CATG**ATTACACTAACTGAGCTAAAAATGCTTAGCAGATGCCAGTCATCTTATCACATC
TTAAAACCATGGTGGGACGTCTTCTGGTATTACATCACACTGATCATGCTGCTGGTGGCCGTG
CTGGCCGGAGCTCTCCAGCTGACGCAGAGCAGGGTTCTGTGCTGTCTTCCATGCAAAGTGGA
ATTTGACAATCACTGTGCCGTGCCTTGGGACATCCTGAAAGCCAGCATGAACACATCCTCTA
ATCCTGGGACACCGCTTCCGCTCCCCCTCCGAATTCAGAATGACCTCCACCGACAGCAGTAC
TCCTATATTGATGCCGTCTGTTACGAGAAACAGCTCCATTGGTTTGCAAAGTTTTTCCCCTA
TCTGGTGCTCTTGACACGCTCATCTTTGCAGCCTGCAGCAACTTTTGCTTCACTACCCCA
GTACCAGTTCAGGCTCGAGCATTTTGTGGCCATCCTTCACAAGTGCTTCGATTCTCCATGG
ACCACCCGCGCCCTTTCAGAAACAGTGGCTGAGCAGTCAGTGAGGCCTCTGAAACTCTCCAA
GTCCAAGATTTTGTCTTCGTCTCAGGGTGTTCAGCTGACATAGATTCCGGCAAACAGTCAT
TGCCCTACCCACAGCCAGGTTTGGAGTCAGCTGGTATAGAAAGCCCAACTTCCAGTGGCCTG
GACAAAGAGGAGGGTGAACAGGCCAAAGCCATCTTTGAAAAAGTGAAAAGATTCCGTCATGCA
TGTGGAGCAGAAGGACATCATTTATAGAGTATATCTGAAACAGATAATAGTCAAAGTCATTT
TGTTTGCTCATCACTTATGTTCCATATTTTTTAACCCACATCACTCTTGAAATCGAC
TGTTCAGTTGATGTGCAGGCTTTTACAGGATATAAGCGCTACCAGTGTGTCTATTCCTTGGC
AGAAATCTTTAAGGCTCTGGCTTCATTTTATGTCATTTTGTTTATACTTTATGGTCTGACCT
CTTCTACAGCCTGTGCTGGATGCTGAGGAGTCCCTGAAGCAATATTCCTTTGAGGCGTTA
AGAGAAAAAAGCAACTACAGTGACATCCCTGATGTCAAGAATGACTTTGCCTTCATCCTTCA
TCTGGCTGATCAGTATGATCCTCTTTATTCCAAACGCTTCTCCATATTCTATCAGAGGTCA
GTGAGAACAACTGAAACAGATCAACCTCAATAATGAATGGACAGTTGAGAACTGAAAAGT
AAGCTTGTGAAAAATGCCAGGACAAGATAGAAGTGCATCTTTTTATGCTCAACGGTCTTCC
AGACAATGTCTTTGAGTTAACTGAAATGGAAGTGCTAAGCCTGGAGCTTATCCAGAGGTGA
AGCTGCCCTCTGCAGTCTCACAGCTGGTCAACCTCAAGGAGCTTCGTGTGTACCATTCATCT
CTGGTCTGAGACCATCTGCAGTGGCCTTTCTAGAGGAGAATTTAAAAATCCTCCGCTGAA
ATTTACTGAAATGGGAAAAATCCACGCTGGGTATTTACCTCAAGAACTCTCAAGGAACCTT
ATCTTTTGGGCTGTGTTCTCCCTGAACAGTTGAGTACTATGCAGTTGGAGGGCTTTCAGGAC
TTAAAAATCTAAGGACCTGTACTTGAAGAGCAGCCTCTCCCGGATCCACAAAGTTGTTACA
GACCTCTGCCCTTCATTGCAGAACTGTCCCTTGATAATGAGGGAAGCAAACCTGGTTGTGTT
GAACAACCTGAAAAAGATGGTCAATCTGAAAAGCCTAGAACTGATCAGCTGTGACCTGGAAC
GCATCCACATTCATTTTTCAGCCTGAATAATTTGCATGAGTTAGACCTAAGGGAAAAATAAC
CTTAAACTGTGGAAGAGATTAGCTTTTCAGCATCTTCAGAATCTTTCCTGCTTAAAGTTGTG
GCACAATAACATTGCTTATATTCTGCACAGATTGGGGCATTATCTAACCTAGAGCAGCTCT
CTTTGGACCATAATAATATTGAGAATCTGCCCTTGCAGCTTTTCTATGCACTAACTACAT
TATTTGGATCTAAGCTATAACCACTTGACCTTCATTCCAGAAGAAATCCAGTATCTGAGTAA
TTTGAGTACTTTGCTGTGACCAACAACAATATTGAGATGCTACCAGATGGGCTGTTTCACT
GCAAAAAGCTGCAGTGTTTACTTTTGGGGAAAAATAGCTTGATGAATTTGTCCCCTCATGTG
GGTGAGCTGTCAAACCTTACTCATCTGGAGCTCATTGGTAATTACCTGGAAACACTTCCTCC
TGAAGTAGAAGGATGTCAGTCCCTAAAACGGAACGTCTGATTGTTGAGGAGAAGCTTGCTCA
ATACTCTTCTCTCCCTGTAACAGAACGTTTACAGACGTGCTTAGACAAATGT**TGA**CTTAAA
GAAAAGAGACCCGTGTTTCAAATCATTTTTTAAAAGTATGCTCGGCCGGGCGTGGTGGCTCA
TGCTATAATCCCAGCACTTTGGGAGGCCAAGATGGGCGGATTGCTTGAGGTCAGGAGTTTCG
AGACCAGTCTGGCCAACCTGGTGAAACCCCATCTCTGCTAAAACCTACAAAAAAATTAGCCAG
GCGTGGTGGCGTGCCTGTAAATCCAGCTACTTGGGAGGCTGACGCAGGGGAATTGCTTGA
ACCAGGGAGGTGGAGGTTGCAGTGAGCCGAGATTGTGCCACTGTACACCAGCCTGGGTGACA
GAGCAAGACTCTTATCTCAAAAAAAAAAAAAA

FIGURE 78

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA161005
><subunit 1 of 1, 802 aa, 1 stop
><MW: 92235, pI: 6.80, NX(S/T): 5
MITLTELKCLADAQSSYHILKPWWDFWYYITLIMLLVAVLAGALQLTQSRVLCCLPCKV
EFDNHCAVPWDILKASMTSSNPGTLPPLPLRIQNDLHRQQYSYIDAVCYEKQLHWFQAFK
FPYLVLLHTLIFAACSNFWLHYPSTSSRLEHFVAILHKCFDSPWTTTRALSETVAEQSVRP
LKLSKSKILLSSSGCSADIDSGKQSLPYPQPGLESAGIESPTSSGLDKKEGEQAKAIFEK
VKRFRMHVEQKDIIRVYLYKQIIVKVILFVLIITYVPYFLTHITLEIDCSVDVQAFTGYK
RYQCVYSLAEIFKVLASFYVILVILYGLTSSYSLWWMRLRSSLKQYSFEALREKSNYS DIP
DVKNDFAFILHLADQYDPLYSKRFSIFLSEVSENKLLKQINLNNEWTVEKLSKLVKNAQD
KIELHLFMLNGLPDNVFELTEMEVLSLELIPEVKLPASVSQVLNKLKELRVYHSSLVVDHP
ALAFLEENLKILRLKFTMGKIPRWVFHLKNLKELYLSGCVLPEQLSTMQLEGFQDLKNL
RTLYLKSSLSRIPQVVTDLLPSLQKLSLDNEGSKLVVLNNLKKMVNLKSLELISCDLERI
PHSIFSLNNLHELDLRENNLKTVEEISFQHLQNLSCLLKWHNNIAYIPAIGALSNEQL
SLDHNNIENLPLQLFLCTKLHYLDLSYNHLTFIPEEIQYLSNLQYFAVTNNNIEMLPDGL
FQCKKLQCLLLGKNSLMNLSPHVGELSNLTHLELIGNYLETLPPPELEGCSLKRNC LIVE
ENLLNTLPLPVTERLQTCLDKC
```

Important features of the protein:

Signal peptide:

Amino acids 1-46

Transmembrane domains:

Amino acids 118-138; 261-281; 311-332

N-glycosylation sites:

Amino acids 78-82; 355-359; 633-637; 748-752

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 382-386

Tyrosine kinase phosphorylation site:

Amino acids 21-30

N-myristoylation sites:

Amino acids 212-218; 327-333; 431-437; 652-658;
719-725

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 125-136

Leucine zipper pattern:

Amino acids 468-490

Leucine Rich Repeat:

Amino acids 609-632; 748-770

FIGURE 79

CGGACGCGTGGGCGCGCTCCCTCACGGCCCCTCGGCGGCGCCCGTCCGGATCCGGCCTCTCT
CTGCGCCCCGGGGCGCGCCACCTCCCCGCCGGAGGTGTCCACGCGTCCGGCCGTCCATCCGT
CCGTCCCTCCTGGGGCCGGCGCTGACCATGCCCAGCGGCTGCCGCTGCCTGCATCTCGTGTG
CCTGTTGTGCATTCTGGGGGCTCCCGTCCAGCCTGTCCGAGCCGATGACTGCAGCTCCCACT
GTGACCTGGCCACGGCTGCTGTGCACCTGACGGCTCCTGCAGGTGTGACCCGGGCTGGGAG
GGGCTGCACTGTGAGCGCTGTGTGAGGATGCCTGGCTGCCAGCACGGTACCTGCCACCAGCC
ATGGCAGTGCATCTGCCACAGTGGCTGGGCAGGCAAGTTCTGTGACAAAGATGAACATATCT
GTACCACGCAGTCCCCCTGCCAGAATGGAGGGCCAGTGCATGTATGACGGGGGCGGTGAGTAC
CATTGTGTGTGCTTACCAGGCTTCCATGGGCGTGACTGCGAGCGCAAGGCTGGACCCTGTGA
ACAGGCAGGCTCCCCATGCCGCAATGGCGGGCAGTGCCAGGACGACCAGGGCTTTGCTCTCA
ACTTCACGTGCCGCTGCTTGGTGGGCTTTGTGGGTGCCCGCTGTGAGGTAAATGTGGATGAC
TGCCTGATGCGGCCTTGTGCTAACGGTGCCACCTGCCTTGACGGCATAAACCGCTTCTCCTG
CCTCTGTCTGAGGGCTTTGCTGGACGCTTCTGCACCATCAACCTGGATGACTGTGCCAGCC
GCCCATGCCAGAGAGGGGGCCCGCTGTCGGGACCGTGTCACGACTTCGACTGCCTCTGCCCC
AGTGGCTATGGTGGCAAGACCTGTGAGCTTGTCTTACCTGTCCCAGACCCCCCAACCACAGTG
GACACCCCTCTAGGGCCCCACCTCAGCTGTAGTGGTACCTGCTACGGGGCCAGCCCCCACAG
CGCAGGGGCTGGTCTGCTGCGGATCTCAGTGAAGGAGGTGGTGC GGAGGCAAGAGGCTGGGC
TAGGTGAGCCTAGCTTGGTGGCCCTGGTGGTGTTTGGGGCCCTCACTGCTGCCCTGGTTCTG
GCTACTGTGTTGCTGACCCTGAGGGCCTGGCGCCGGGGTGTCTGCCCCCTGGACCCTGTTG
CTACCCTGCCCCACACTATGCTCCAGCGTGCCAGGACCAGGAGTGTGAGGTAGCATGCTGC
CAGCAGGGCTCCCCCTGCCACGTGACTTGCCCCCTGAGCCTGGAAAGACCACAGCACTGTGA
TGGAGGTGGGGGCTTTCTGGCCCCCTTCCTCACCTCTTCCACCCCTCAGACTGGAGTGGTCC
GTTCTCACACCCTTCAGCTTGGGTACACACACAGAGGAGACCTCAGCCTCACACCAGAAAT
ATTATTTTTTTAATACACAGAATGTAAGATGGAATTTTATCAAATAAACTATGAAAATGCA
AAAAAAAAAAAAA

FIGURE 80

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA170245
><subunit 1 of 1, 383 aa, 1 stop
><MW: 40548, pI: 6.48, NX(S/T): 1
MPSGCRCLHLVCLLCILGAPGQPVRADDCSSHCDLAHGCCAPDGGSCRCDPGWEGLHCERC
VRMPGCQHGTCHQPWQCICHSGWAGKFCDKDEHICTTQSPCQNGGQCMYDGGGEYHCVCL
PGFHGRDCERKAGPCEQAGSPCRNGGQCQDDQGFALNFTCRCLVGFVGARCEVNVDCLM
RPCANGATCLDGINRFSCLCPEGFAGRECTINLDDCASRPCQRGARCRDRVHDFDCLCPS
GYGGKTCELVLVPDPPTTVDTPLGPTSAVVVPATGPAPHSAGAGLLRISVKEVVRRQEA
GLGEPSSLVALVVFALTAALVLATVLLTLRAWRRGVCPGPGCCYPAPHYAPACQDQECQV
SMLPAGLPLPRDLPPPEPGKTTAL
```

Important features of the protein:

Signal peptide:

Amino acids 1-21

Transmembrane domain:

Amino acids 306-331

N-glycosylation site:

Amino acids 157-160

Glycosaminoglycan attachment site:

Amino acids 240-243

N-myristoylation sites:

Amino acids 44-49;65-70;243-248;314-319

Aspartic acid and asparagine hydroxylation sites:

Amino acids 189-200;227-238

EGF-like domain cysteine pattern signature:

Amino acids 46-57;77-88;117-128;160-171;198-209;
236-247

Zinc finger, C3HC4 type, signature:

Amino acids 7-16

EGF-like domain proteins:

Amino acids 46-58;77-89;117-129;160-172;198-210;
216-228;236-248

FIGURE 81

GTTTGTGCTCAAACCGAGTTCTGGAGAACGCCATCAGCTCGCTGCTTAAAAATTAAACCACA
GGTTCCATT**ATG**GGTTCGACTTGATGGGAAAGTCATCATCCTGACGGCCGCTGCTCAGGGGAT
TGGCCAAGCAGCTGCCTTAGCTTTTGCAAGAGAAGGTGCCAAAGTCATAGCCACAGACATTA
ATGAGTCCAAACTTCAGGAACTGGAAAAGTACCCGGGTATTCAAACCTCGTGTCTTGATGTC
ACAAAGAAGAAACAAATTGATCAGTTTGCCAGTGAAGTTGAGAGACTTGATGTTCTCTTTAAT
GTTGCTGGTTTTGTCCATCATGGAAGTGTCTTGGATTGTGAGGAGAAAGACTGGGACTTCTC
GATGAATCTCAATGTGCGCAGCATGTACCTGATGATCAAGGCATTCTTCTTAAATGCTTG
CTCAGAAATCTGGCAATATTATCAACATGTCTTCTGTGGCTTCCAGCGTCAAAGGAGTTGTG
AACAGATGTGTGTACAGCACAAACCAAGGCAGCCGTGATTGGCCTCACAAAATCTCTGGCTGC
AGATTTTCATCCAGCAGGGCATCAGGTGCAACTGTGTGTGCCAGGAACAGTTGATACGCCAT
CTCTACAAGAAAGAATACAAGCCAGAGGAAATCCTGAAGAGGCACGGAATGATTTCTGAAG
AGACAAAAGACGGGAAGATTCGCAACTGCAGAAGAAATAGCCATGCTCTGCGTGTATTTGGC
TTCTGATGAATCTGCTTATGTAAGTGGTAACCCTGTCATCATTGATGGAGGCTGGAGCTTGT
GATTTTAGGATCTCCATGGTGGGAAGGAAGGCAGGCCCTTCTATCCACAGTGAACCTGGTT
ACGAAGAAAACCTACCAATCATCTCCTTCTGTTAATCACATGTTAATGAAAATAAGCTCTT
TTAATGATGTCACTGTTTGCAAGAGTCTGATTCTTTAAGTATATTAATCTCTTTGTAATCT
CTTCTGAAATCATTGTAAGAAATAAAAAATATTGAACTCAT

FIGURE 82

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA171771
><subunit 1 of 1, 245 aa, 1 stop
><MW: 26711, pI: 8.00, NX(S/T): 2
MGRLDGKVIILTAAQGGIGQAAALAFAREGAKVIATDINESKLQELEKYPGIQTRVLDVT
KKKQIDQFASEVERLDVLEFNVAGFVHHGTVLDCEEKDWDFSMNLNVRSMYLMIKAFLPKM
LAQKSGNIINMSSVASSVKGVNRCVYSTTKAAVIGLTKSLAADFIQQGIRCNCVCPGTV
DTPSLQERIQARGNPPEARNDLKRQKTGRFATAEEIAMLCVYLASDESAYVTGNPVIID
GGWSL
```

Important features of the protein:

Signal peptide:

Amino acids 1-20

N-glycosylation sites:

Amino acids 39-43;130-134

Tyrosine kinase phosphorylation site:

Amino acids 42-50

N-myristoylation sites:

Amino acids 17-23;19-25;126-132;156-162;169-175

Short-chain dehydrogenases/reductases family proteins:

Amino acids 7-19;73-83;127-164; 169-178

Short chain dehydrogenase:

Amino acids 7-183

FIGURE 83

GGGCGGCGGCGGCAGCGGTTGGAGGTTGTAGGACCGGCGAGGAATAGGAATC**ATG**GCGGCTG
CGCTGTTTCGTGCTGCTGGGATTTCGCGCTGCTGGGCACCCACGGAGCCTCCGGGGCTGCCGGC
TTCGTCCAGGCGCCGCTGTCCCAGCAGAGGTGGGTGGGGGCGAGTGTGGAGCTGCACTGCGA
GGCCGTGGGCAGCCCGGTGCCGAGATCCAGTGGTGGTTTGAAGGGCAGGGTCCCAACGACA
CCTGCTCCCAGCTCTGGGACGGCGCCCGGCTGGACCGCGTCCACATCCACGCCACCTACCAC
CAGCACGCGGCCAGCACCATCTCCATCGACACGCTCGTGGAGGAGGACACGGGCACTTACGA
GTGCCGGGCCAGCAACGACCCGGATCGCAACCACCTGACCCGGGCGCCCAGGGTCAAGTGGG
TCCGCGCCAGGCAGTTCGTGCTAGTCCCTGGAACCCGGCACAGTCTTCACTACCGTAGAAGAC
CTTGCTCCAAGATACTCCTCACCTGCTCCTTGAATGACAGCGCCACAGAGGTCACAGGGCA
CCGCTGGCTGAAGGGGGGCGTGGTGCTGAAGGAGGACGCGCTGCCCGGCCAGAAAACGGAGT
TCAAGGTGGACTCCGACGACCACTGGGGAGAGTACTCCTGCGTCTTCCCTCCCGAGCCCATG
GGCACGGCCAACATCCAGCTCCACGGGCTCCAGAGTGAAGGCTGTGAAGTCTGTCAGAACA
CATCAACGAGGGGGAGACGGCCATGCTGGTCTGCAAGTCAGAGTCCGTGCCACCTGTCACTG
ACTGGGCTTGGTACAAGATCACTGACTCTGAGGACAAGGCCCTCATGAACGGCTCCGAGAGC
AGGTTCTTCGTGAGTTCCTCGCAGGGCCGGTCAGAGCTACACATTGAGAACCCTGAACATGGA
GGCCGACCCCGGCCAGTACCGGTGCAACGGCACCAAGCTCCAAGGGCTCCGACCAAGGCCATCA
TCACGCTCCGCGTGCGCAGCCACCTGGCCGCCCTCTGGCCCTTCCCTGGGCATCGTGGCTGAG
GTGCTGGTGCTGGTCACCATCATCTTCATCTACGAGAAGCGCCGGAAGCCCGAGGACGTCTCT
GGATGATGACGACGCCGGCTCTGCACCCCTGAAGAGCAGCGGGCAGCACCAGAATGACAAAG
GCAAGAACGTCCGCCAGAGGAACCTTCCCT**TGA**GGCAGGTGGCCCGAGGACGCTCCCTGCTCC
ACGTCTGCGCCGCCCGCGGAGTCCACTCCAGTGCTTGCAAGATTCCAAGTTCTCACCTCTT
AAAGAAAACCCACCCCGTAGATTCCCATCATACACTTCTTCTTTTTTAAAAAAGTTGGGTT
TTCTCCATTACAGGATTCTGTTCCTTAGGTTTTTTTCTTCTGAAGTGTTCACGAGAGCCCG
GGAGCTGCTGCCCTGCGGCCCGTCTGTGGCTTTCAGCCTCTGGGTCTGAGTCATGGCCGGG
TGGGCGGCACAGCCTTCTCCACTGGCCGGAGTCAGTGCCAGGTCCTTGCCCTTTGTGGAAAGTC
ACAGGTCACACGAGGGGGCCCGTGTCTGCTGTCTGAAGCCAATGCTGTCTGGTTGCGCCA
TTTTTGTGCTTTTATGTTTAATTTTATGAGGGCCACGGGTCTGTGTTGCACTCAGCCTCAGG
GACGACTCTGACCTCTTGCCACAGAGGACTCACTTGCCACACCGAGGGCGACCCCGTCAC
AGCCTCAAGTCACTCCCAAGCCCCCTCCTTGTCTGTGCATCCGGGGGCGAGCTCTGGAGGGGG
TTTGCTGGGGAACCTGGCGCCATCGCCGGGACTCCAGAACCGCAGAAGCCTCCCCAGCTCACC
CCTGGAGGACGGCCGGCTCTCTATAGCACCAGGGCTCACGTGGGAACCCCCCTCCACCCAC
CGCCACAATAAAGATCGCCCCCACCTCCACCCAAAAA

FIGURE 84

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA173157
><subunit 1 of 1, 385 aa, 1 stop
><MW: 42200, pI: 5.57, NX(S/T): 5
MAAALFVLLGFALLGTHGASGAAGFVQAPLSQQRWVGGSVELHCEAVGSPVPEIQWWFEG
QGPNDTCSQLWDGARLDRVHIHATYHQHAASTISIDTLVEEDTGTYECRASNDPDRNHLT
RAPRVKQWVRAQAVVLVLEPGTVFTTVEDLGSKILLTCSLNDSETEVTGHRWLKGGVVLKE
DALPGQKTEFKVDSDDQWGEYSCVFLPEPMGTANIQLHGPPRVKAVKSSEHINEGETAML
VCKSESVPVPTDWAUWKITDSEDKALMNGSESFFVSSSQGRSELHIENLNMEADPGQYR
CNGTSSKGSQDAIITLRVRSHLAALWFLGIVAEVLVLVTIIFIYEKRRKPEDVLDDDDA
GSAPLKSSGQHNDKGNVRQRNSS
```

Important features of the protein:

Signal peptide:

Amino acids 1-18

Transmembrane domain:

Amino acids 320-343

N-glycosylation sites:

Amino acids 64-68;160-164;268-272;302-306

N-myristoylation sites:

Amino acids 15-21;18-24;60-66;104-110;140-146;
297-303;308-314;369-375

Immunoglobulin domain:

Amino acids 37-110;150-205;235-303

FIGURE 85

GGCTCGAGCAAAGACATACGAACAGGGAGGAAGGCCGACTGAAAGAAAGACGGAGAAGAGGA
GAGAGAAGCCAGGGCCGAGCGTGCCAGCAGGCGGATGGAGGGCGGCCTGGTGGAGGAGGAGA
CGTAGTGGCCTGGGCTGAGCTGGGTGGGCCGGGAGAAGCGGGTGCCTCAGAGTGGGGGTGGG
GGC**ATG**GGGAGGGGCAGGCATTCTGCTGCTGCTGCTGGCTGGGGCGGGGTGGTGGTGGCCTGG
AGACCCCAAAGGGAAAGTGTCCCCTGCGCTGCTCCTGCTCTAAAGACAGCGCCCTGTGTGA
GGGCTCCCCGGACCTGCCCCTCAGCTTCTCTCCGACCCTGCTGTCACTCTCACTCGTCAGGA
CGGGAGTCACCCAGCTGAAGGCCGGCAGCTTCCTGAGAATTCCGTCTCTGCACCTGCTCCTC
TTCACCTCCAACCTCCTTCTCCGTGATTGAGGACGATGCATTTGCGGGCCTGTCCCACCTGCA
GTACCTCTTCATCGAGGACAATGAGATTGGCTCCATCTCTAAGAATGCCCTCAGAGGACTTC
GCTCGCTTACACACCTAAGCCTGGCCAATAACCATCTGGAGACCCTCCCCAGATTCCCTGTTT
CGAGGCCTGGACACCCCTTACTCACGTGGACCTCCGCGGGAACCCGTTCCAGTGTGACTGCCG
CGTCCTCTGGCTCCTGCACTGGATGCCACCGTGAATGCCAGCGTGGGGACCGGCGCCTGTG
CGGGCCCCGCCTCCCTGAGCCACATGCAGCTCCACCACCTCGACCCCAAGACTTTCAAGTGC
AGAGCCATAGGTGGGGGGCTTTCCCGATGGGGTGGGAGGCGGGAGATCTGGGGGAAAGGCTG
CCAGGGCCAAGAGGCTCGTCTCACTCCCTGCCCTGCCATTTCCCGAGTGGGAAGACCCTGA
GCAAGCAGCACTGCCTTCCTGAGCCCCAGTTTTCTCATCTG**TAA**AGTGGGGGTAATAAACAG
TGATATAGG

FIGURE 86

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA175734
><subunit 1 of 1, 261 aa, 1 stop
><MW: 28231, pI: 9.28, NX(S/T): 1
MGGAGILLLLLAGAGVVVAWRPPKGKCLRCSCSKDSALCEGSPDLPVSFSPDLLSLSLV
RTGVTQLKAGSFLRIPSLHLLLF'TSNSFSVIEDDAFAGLSHLQYLFIEDNEIGSISKNAL
RGLRSLTHLSLANNHLETLPREFLFRGLDTLTHVDLRGNPFQCDRCVWLWLQWMPTVNASV
GTGACAGPASLSHMLHLLDPKTFKCRAIGGGLSRWGGRRREIWGKGCQGQEARLTPCPAI
SRSGKTLKQHCLEPEPQFSHL
```

Important features of the protein:

Signal peptide:

Amino acids 1-19

N-glycosylation site:

Amino acids 177-181

N-myristoylation sites:

Amino acids 15-21;181-186;210-215

Amidation site:

Amino acids 217-220

Microbodies C-terminal targeting signal:

Amino acids 259-262

ATP/GTP-binding site motif A (P-loop):

Amino acids 239-246

Leucine zipper pattern:

Amino acids 129-150

Leucine Rich Repeat:

Amino acids 53-76; 149-171

Leucine rich repeat C-terminal domain:

Amino acids 158-207

FIGURE 87

CGGACGCGTGGGGCGGCGAGAGCAGCTGCAGTTTCGCATCTCAGGCAGTACCTAGAGGAGCTG
 CCGGTGCC'TCCTCAGAACATCTCCTGATCGCTACCCAGGACCAGGCACCAAGGACAGGGAGT
 CCCAGGCGCACACCCCCCATTTCTGGGTCCCCCAGGCCAGACCCCCACTCTGCCACAGGTTG
 CATCTTGACCTGGTCTCTCCTGCAGAAAGTGGCCCCCTGTGGTCTGCTCTGAGACTCGTCCCTG
 GCGCCCCCTGCAGCCCCCTTTCTATGACTCCATCTGGATTTGGCTGGCTGTGGGGACGCGGTC
 CGAGGGGCGGCTGGCTCTCAGCGTGGTGGCAGCCAGCTCTCTGGCCACCATGGCAAATGCT
 GAGATCTGAGGGGACAAGGCTCTACAGCCTCAGCCAGGGGCCACTCAGCTGTTGTCAGGGTGTG
ATGGAGAACAAGCTATGTACCTACACACCGTCAGCGACTGTGACACCAGCTCCATCTGTGA
 GGATTCCTTTGATGGCAGGAGCCTGTCCAAGCTGAACCTGTGTGAGGATGGTCCATGTCA
 AACGGCGGGCAAGCATCTGCTGTACCCAGCTGGGGTCCCTGTGCGCCCTGAAGCATGCTGTC
 CTGGGGCTCTACCTGCTGGTCTTCTGATTCTTGTGGGCATCTTCATCTTAGCAGGGCCACC
 GGGACCCAAAGGTGATCAGGGGGATGAAGGAAAGGAAGGCAGGCCTGGCATCCCTGGATTGC
 CTGGACTTCGAGGTCTGCCCCGGGGAGAGAGGTACCCAGGATTGCCCGGGCCCAAGGGCGAT
 GATGGGAAGCTGGGGGCCACAGGACCAATGGGCATGCGTGGGTCAAAGGTGACCGAGGCCC
 AAAAGGAGAGAAAGGAGAGAAAGGAGACAGAGCTGGGGATGCCAGTGGCGTGGAGGCCCCGA
 TGATGATCCGCTGGTGAATGGCTCAGGTCCGCACGAGGGCCGCGTGGAAAGTGATCCACGAC
 CGGCGCTGGGGCACCGTGTGTGACGACGGCTGGGACAAGAAGGACGGAGACGTGGTGTGCCG
 CATGCTCGGCTTCCGCGGTGTGGAGGAGGTGTACCGCACAGCTCGATTGGGCAAGGCACTG
 GGAGGATCTGGATGGATGACGTTGCC'TGCAAGGGCACAGAGGAAACCATCTTCCGCTGCAGC
 TTCTCCAAATGGGGGGTGACAACTGTGGACATGCCGAAGATGCCAGCGTGACATGCAACAG
 ACACTGAAAGTGGGCAGAGCCCCAAGTTTCGGGGTCCCTGCACAGAGCACCCCTTGCTGCATCCCT
 GGGTGGGGCACAGCTCGGGGCCACCCTGACCATGCC'TCGACCACACCCCGTCCAGCATTCT
 CAGTCCCTCACACCTGCATCCAGGACCGTGGGGGCCGGTGGTCAATTTCCCTCTTGAACATGT
 GCTCCGAAGTATACTCTGGGACCTACTGCCCCGTCTCTCTCTTCCACAGGTTCCTGCATGA
 GGAGCCCTGATCAACTGGATCACCACTTTGCCACGCCCTCTGAACACCATGCACCAGGCCTCA
 ATATCCCAGTTCCCTTTGGCCCTTTAGTTACAGGTGAATGCTGAGAATGTGTGAGAGACAAG
 TGCAGCAGCAGCGATGGTTGGTAGTATAGATCATTTACTCTTCAGACAATTCCCAAACCTCC
 ATTAGTCCAAGAGTTTCTACATCTTCCCTCCCCAGCAAGAGGCAACGTCAAGTGATGAATTC
 CCCCCTTTACTCTGCTCTGCTTCCCCATTTGCTAGTTTGAGGAAGTGACATAGAGGAGAAGC
 CAGCTGTAGGGGCAAGAGGGAAATGCAAGTCACCTGCAGGAATCCAGCTAGATTTGGAGAAG
 GGAATGAACTAACATTGAATGACTACCATGGCAGCGCTAAATAGTATCTTGGGTGCCAAATTCA
 TGTATCCACTTAGCTGCATTGGTCCAGGGCATGTGAGTCTGGATACAGCC'TTACCTTCAGGT
 AGCACTTAACCTGGTCCATTACCTAGACTGCAAGTAAGAAGACAAAATGACTGAGACCGTGT
 GCCACCTGAACTTATTTGCTTTTACTTGGCCTGAGCTAAAAGCTTGGGTGCAGGACCTGTGT
 AACTAGAAAGTTGCCTACTTCAGAACCTCCAGGGCGTGAGTGCAAGGTCAAACATGACTGGC
 TTCCAGGCCGACCATCAATGTAGGAGGAGAGCTGATGTGGAGGGTGACATGGGGGCTGCCCA
 TGTTAAACCTGAGTCCAGTGCTCTGGCATTTGGGCAGTCACGGTTAAAGCCAAGTCATGTGTG
 TCTCAGCTGTTTGGAGGTGATGATTTTGCATCTTCCAAGCCTCTTCAGGTGTGAATCTGTGG
 TCAGGAAAACACAAGTCCATGGAACCC'TTAGGGGGGAAGGAAATGAAGATTCCCTATAAC
 CTCTGGGGGTGGGGAGTAGGAATAAGGGGCCCTTGGGCCTCCATAAATCTGCAATCTGCACCC
 TCCTCCTAGAGACAGGGAGATCGTGT'TCTGCTTTTACATGAGGAGCAGAAGTGGGCCATAC
 ACGTGT'TCAAGAACTAGGGGAGCTACCTGGTAGCAAGTGAGTGCAGACCCACCTCACCTTGG
 GGAATCTCAAACCTCATAGGCCCTCAGATACACGATCACCTGTATATCAGGTGAGCACTGGC
 CTGCTTGGGGAGAGACCTGGGGCCCTCCAGGTGTAGGAACAGCAACACTCCTGGCTGACAAC
 TAAGCCAATATGGCCCTAGGTCAATCTTGCTTCCAATATGCTTGCCACTCCTTAAATGTCTT
 AATGATGAGAACTCTCTTTCTGACCAATTGCTATGTTTACATAACACGCATGTACTCATGC
 ATCCCTTGCCAGAGCCCATATATGTATGCATATATAAACATAGCACTTTTACTACATAGCT
 CAGCACAT'TGCAAGGTTTGCATTTAAGTT

FIGURE 88

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA176108
><subunit 1 of 1, 270 aa, 1 stop
><MW: 28871, pI: 7.09, NX(S/T): 1
MENKAMYLHTVSDCDTSSICEDSFDGRSLSKLNLCEGDPCHKRRASICCTQLGSLSALKH
AVLGLYLLVFLILVGIFILAGPPGPKGDQGDGKEGRPGIPGLPGLRGLPGERGTPGLPG
PKGDDGKLGATGPMGMRGFKGDRGPKGEKGEKGDAGDASGVEAPMMIRLVNGSGPHEGR
VEVYHRRRWGTVCDDGWDKKGDDVVCRLGFRGVVEVYRTARFGQGTGRIWMDDVACKGT
EETIFRCSFSKWGVTNCGHAEDASVTCNRH
```

Transmembrane domain:

Amino acids 55-80

N-glycosylation site:

Amino acids 172-175

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 43-46

Tyrosine kinase phosphorylation site:

Amino acids 212-218

N-myristoylation sites:

Amino acids 53-58;224-229;239-244;253-258

Speract receptor repeated domain signature:

Amino acids 173-211

Scavenger receptor cysteine-rich domain:

Amino acids 171-268

Collagen Collagen triple helix repeat:

Amino acids 90-149

FIGURE 89

GTCGCCGCGAGGGACGCAGAGAGCACCCCTCCACGCCCAGATGCCTGCGTAGTTTTTGTGACC
 AGTCCGCTCCTGCCTCCCCCTGGGGCAGTAGAGGGGGAGCG**ATG**GAGAACTGGACTGGCAGG
 CCCTGGCTGTATCTGCTGCTGCTTCTGTCCCTCCCTCAGCTCTGCTTGGATCAGGAGGTGTT
 GTCCGGACACTCTCTTCAGACACCTACAGAGGAGGGCCAGGGCCCCGAAGGTGTCTGGGGAC
 CTTGGGTCCAGTGGGCCTCTTGCTCCCAGCCCTGCGGGGTGGGGGTGCAGCGCAGGAGCCGG
 ACATGTCAGCTCCCTACAGTGCAGCTCCACCCGAGTCTGCCCCCTCCCTCCCCGGCCCCCAAG
 ACATCCAGAAGCCCTCCTCCCCCGGGGCCAGGGTCCCAGACCCCAGACTTCTCCAGAAACCC
 TCCCCTTGTAACAGGACACAGTCTCGGGGAAGGGGTGGCCCACTTCGAGGTCCCGCTTCCAC
 CTAGGGAGAGAGGAGACCCAGGAGATTCGAGCGGCCAGGAGGTCCCGGCTTCGAGACCCCAT
 CAAGCCAGGAATGTTTCGGTTATGGGAGAGTGCCCTTTGCATTGCCACTGCACCCGGAACCCGA
 GGCACCCCTCGGAGGCCACCCAGATCTGAGCTGTCCCTGATCTCTTCTAGAGGGGAAGAGGCT
 ATTCGGTCCCTACTCCAAGAGCAGAGCCATTCTCCGCAAACGGCAGCCCCCAAAGTGAAGT
 CCCTCCCACAGAAGTGTCTGTCCACACCCCATCCCCCAAGCAGAACCTCTAAGCCCTGAAA
 CTGCTCAGACAGAGGTGGCCCCCAGAACCAGGCCTGCCCCCTACGGCATCACCCAGAGCC
 CAGGCCTCTGGCACAGAGCCCCCTCACCCACGCACTCCTTAGGAGAAGGTGGCTTCTTCCG
 TGCATCCCCCTCAGCCACGAAGGCCAAGTTCCCAGGGTTGGGCCAGTCCCCAGGTAGCAGGGA
 GACGCCCTGATCCTTTTCCCTTCGGTCCCTCGGGGCCGAGGCCAGCAGGGCCAAGGGCCTTGG
 GGAACGGGGGGGACTCCTCACGGGCCCGCCTGGAGCCTGACCCCTCAGCACCCGGGCGCCTG
 GCTGCCCCCTGCTGAGCAACGGCCCCCATGCCAGCTCCCTCTGGAGCCTCTTTGCTCCAGTA
 GCCCTATTCCAAGATGTTCTGGGGAGAGTGAACAGCTAAGAGCCTGCAGCCAAGCGCCCTGC
 CCCCCTGAGCAGCCAGACCCCCGGGCCCTGCAGTGCGCAGCCTTTAACTCCCAGGAATTTCATG
 GGCCAGCTGTATCAGTGGGAGCCCTTCACTGAAGTCCAGGGCTCCCAGCGCTGTGAAGTAA
 CTGCGCGCCCCGTGGCTTCCGCTTCTATGTCCGTACACTGAAAAGGTCCAGGATGGGACCC
 TGTGTCAGCCTGGAGCCCCCTGACATCTGTGTGGCTGGACGCTGTCTGAGCCCCGGCTGTGAT
 GGGATCCTTGGCTCTGGCAGGCGTCCTGATGGCTGTGGAGTCTGTGGGGGTGATGATTCTAC
 CTGTGCGCCTTGTTCGGGGAACCTCACTGACCGAGGGGGCCCCCTGGGCTATCAGAAGATCT
 TGTGGATTCCAGCGGGAGCCTTGCGGCTCCAGATTGCCAGCTCCGGCCTAGCTCCAAGTAC
 CTGGCACTTCGTGGCCCTGGGGGCCGGTCCATCATCAATGGGAAGTGGGCTGTGGATCCCC
 TGGGTCTTACAGGGCCGGCGGGACCGTCTTTCGATATAACCGTCTTCCCAGGGAGGAGGGCA
 AAGGGGAGAGTCTGTGCGCTGAAGGCCCCACCACCCAGCCTGTGGATGTCTATATGATCTTT
 CAGGAGGAAAACCCAGGCGTTTTTTATCAGTATGTCATCTCTTCACCTCCTCCAATCCTTGA
 GAACCCACCCAGAGCCCCCTGTCCCCAGCTTCAGCCGGAGATTCTGAGGGTGGAGCCCC
 CACTTGCTCCGGCACCCCGCCAGCCCCGACCCAGGCACCTCCAGCGTCAGGTGCGGATC
 CCCCAGATGCCCCCCCCGCCCCATCCCAGGACACCCCTGGGGTCTCCAGCTGCGTACTGGAA
 ACGAGTGGGACACTCTGCATGCTCAGCGTCTTGGGGAAAGGTGTCTGGCGCCCCATTTTCC
 TCTGCATCTCCCGTGAGTCGGGAGAGGAACTGGATGAACGCAGCTGTGCCGCGGGTGGCAGG
 CCCCCAGCCTCCCCTGAACCCCTGCCACGGCACCCCATGCCCCCATACTGGGAGGCTGGCGA
 GTGGACATCCTGCAGCCGCTCCTGTGGCCCCGGCACCCAGCACCGCCAGCTGCAGTGCCGGC
 AGGAATTTGGGGGGGGTGGCTCCTCGGTGCCCCCGGAGCGCTGTGGACATCTCCCCCGGCC
 AACATCACCCAGTCTTGCCAGCTGCGCCTCTGTGGCCATTGGGAAGTTGGCTCTCCTTGGAG
 CCAGTGCTCCGTGCGGTGCGGCCGGGGCCAGAGAAGCCGGCAGGTTGCTGTGTTGGGAACA
 ACGGTGATGAAGTGAGCGAGCAGGAGTGTGCGTCAGGCCCCCACAGCCCCCAGCAGAGAG
 GCCTGTGACATGGGGCCCTGTACTACTGCCTGGTTCCACAGCGACTGGAGCTCCAAGGTGAG
 CCGGAACCCCAAGCATATCCTGCATCCTGGGTAACCATGCCAGGACACCTCAGCCTTTC
 CAGCA**TAG**CTCAATAAAGTGTATTGATC

FIGURE 90

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA190710
><subunit 1 of 1, 877 aa, 1 stop
><MW: 95132, pI: 8.77, NX(S/T): 5
MENWTGRPWLYLLLLLSLPQLCLDQEVLSGHSLSQTPTTEEGQGPEGVWGPWVQWASCSQPC
GVGVQRRSRTCQLPTVQLHPSLPLPPRPPRHPEALLPRGQGPRPQTSPETLPLYRTQSRG
RGGPLRGPPASHLGREETQEIRAARRSRLRDPIKPGMFGYGRVPFALPLHRNRRHPRSPPR
SELSLISSRGEEAIPSTPRAEPPFSANGSPQTELPPTELSVHTPSPQAEPLSPETAQTEV
APRTRPAPLRHHPRAQASGTEPPSPHSLGEGGFFRASQPQRRPSSQGWASPVAGRRPD
PFPSVPRGRGQQGQGPWGTGGTTPHGPRLEPDQHPGAWLPLLSNGPHASSLWSLFAPSSP
IPRCSGESEQLRACSQAPCPPEQPDPRALQCAAFNSQEFMGQLYQWEPFTEVQGSQRCEL
NCRPRGRFRFYVRHTEKVDGTLCPGAPDICVAGRCLSPGCDGILGSGRRPDGCGVCGGD
DSTCRLVSGNLTDRGGPLGYQKILWIPAGALRLQIAQLRPSSNYLALRGPGRRSIINGNW
AVDPPGSYRAGGTVFRYNRPPREEGKGESLSAEGPTTQPDVYIMIFQEENPGVFYQYVIS
SPPPILENPTPEPPVPQLQPEILRVEPPLAPAPRPARTPGTLQRQVRIQMPAPPHPRTF
LGSPAAYWKRVGHSACSASCGKGVWRPIFLCISRESGEELDERSCAAGARPPASPEPCHG
TPCPPYWEAGEWTSCSRSCGPGTQHRQLQCRQEFGGGGSSVPPERCGHLPRPNITQSCQL
RLCGHWEVGSPWSQCSVRCGRGQRSRQVRCVGNNGDEVSEQECASGPPQPPSREACDMGP
CTTAWFHSDWSSKVSPEPPAISCILGNHAQDTSAFPA
```

Important features of the protein:

Signal peptide:

Amino acids 1-24

N-glycosylation sites:

Amino acids 3-6; 490-493; 773-776

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 282-285

N-myristoylation sites:

Amino acids 208-213; 414-419; 463-468; 473-478; 475-480;
478-483; 495-500; 546-551; 662-667; 755-760;
756-761; 789-794

Amidation sites:

Amino acids 295-298; 467-470

Leucine zipper pattern:

Amino acids 504-526

VWFC domain proteins:

Amino acids 53-67; 732-746; 792-806

Thrombospondin type 1 domain:

Amino acids 48-87; 727-783; 787-841

FIGURE 91

CGAGTATTTTCCCACCATTCTCCAGCCGGAAACTGACCAAGAACTCTGAGGCGGATGGC**ATGT**
TCGCGTACGTCTTCCATGATGAGTTCGTGGCCTCGATGATTAAGATCCCTTCGGACACCTTC
ACCATCATCCCTGACTTTGATATCTACTATGTCTATGGTTTTAGCAGTGGCAACTTTGTCTA
CTTTTTTGACCCCTCCAACCTGAGATGGTGTCTCCACCAGGCTCCACCACCAAGGAGCAGGTGT
ATACATCCAAGCTCGTGAGGCTTTGCAAGGAGGACACAGCCTTCAACTCCTATGTAGAGGTG
CCCATTGGCTGTGAGCGCAGTGGGGTGGAGTACCGCCTGCTGCAGGCTGCCTACCTGTCCAA
AGCGGGGGCCGTGCTTGGCAGGACCCTTGGAGTCCATCCAGATGATGACCTGCTCTTCACCG
TCTTCTCCAAGGGCCAGAAGCGGAAAATGAAATCCCTGGATGAGTCGGCCCTGTGCATCTTC
ATCTTGAAGCAGATAAATGACCGCATTAAGGAGCGGCTGCAGTCTTGTTACCGGGGCGAGGG
CACGCTGGACCTGGCCTGGCTCAAGGTGAAGGACATCCCCTGCAGCAGTGCCTCTTAACCA
TTGACGATAA**CTT**CTGTGGCCTGGACATGAATGCTCCCCTGGGAGTGTCCGACATGGTGCCT
GGAATTTCCCGTCTTCACGGAGGACAGGGACCGCATGACGTCTGTATCGCATATGTCTACAA
GAACCACTCTCTGGCCTTTGTGGGCACCAAAAGTGGCAAGCTGAAGAAGGTGCCTGGTACCA
GCCTCTGCCCTACCCTTGAGCTACAGACGGGACCCCGATCCCACAGAGCAACAGTGACTCTG
GAACTCCTGTTCTCCAGCTGTT**CATCAA**CT**TGA**AAAAA**ACTT**CAGAGCTGTGTAGGCTTATT
TAGTGTGTTGT**CAGCCTT**GGATATTGGAAAATGGAAACAGATGAGACACATCTACCTCCCTG
TGACCC**CAGCC**ATACATCATAGCTCATGTCTGCCACCCCAAGTCTTTAGGAAAAAAGACT
TTGGAGAATGTGTCTCTGTCTAGCTTGGCTAGGTAGTGGTCTCTTTCTCTGCCCCAAGCG
TCCCCTGGGTAATTTTGGACAATGGAGTGTAAGGCATGTTTGACTCTTGTGGTGT**TATCA**CTT
GTATATGT**CAGTGA**AACTAACTGATTCTCCCATCGGAATATAGTTATCTCTTGGGCCTGATA
TATGGTAGGATAAACCTTATGCTCATCTGTCCACTTCTGCAGCCAAGTCGCCTGGCCAGTGTG
TGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTATGCTTATCTGTGTTTAAAGGTGTGTG
TGCATACACAGGGCAGAGAGGATGGAGCCCACCGTACTGCAGCATCATGTAATTA**ACTCAGT**
GCTCAGAA**CCATCCC**AGCCTCTGCGGGAAAGAGAAAAAGTAAGCCAACAGTGCCTGATGAGCT
GATCATATGTGCAAAAGCTCTGTTGGCATCTGGTCCAGGAGAGCACCCCAAAAAAAGTTAATT
GGTGT**TGTCCAGTCTCCTTT**CCTTAAGACTATGGTTACAACAAAGCGTGAGCAGTGTCTCCT
GCATGGCCACTATCCAGCACAATTCCATAATTCCCCCATAGAGCCGGTGGGGAGGAGGAGGT
GAGTGGCGAAGGAAGTGAAACACTTGGTGT**CATGTGCTCCTATC**ATTTCTACTAGCTTACT
GGGAAATAAAGTGTA**GTCA**AGAGTGATGAAGGCAAGATGTAA**AA**TTAGCGACTGGTGCTAA
TCTGGTTACTTGA**AA**ACAAGTGAAAGTGCTGTAGATTTGTTCTGTTGCTAAGAAC**ACCACA**
CTAAACCTCGTATAGTT**CCTGG**AGGATATACAACAGTGTAA**TTCTCTTT**AGGGTGTGCCACA
GGTT**CCTGGCCTGTGGG**AGGGAATGAATCAGGAGGGCTCTTGAGAACTT**CATCTGTGTGCT**
TGC**ACTGAA**AGTGAGTCCCAAAGCTGGAGATTTAGTGAGAGCAGGCAACCCCTCTGTGTCTC
ACTGTCCATATTCTGGAGGCAGAGGTTTGTAA**CAGGCCATGTGCACCTGCATAGGGATGGGT**
AAAGCAAGGACTTTGA**AA**AGAGTTGA**AA**AGCATTATAAACAGTTGTT**CAGAAATACGTCCCAG**
GAGTTCCATGTGA**AACTGGCTCTGTGTGCATTGAAGCATGGCTGTTGGGAATCTAACTGGT**
CCAACACTCCTGCA**AA**ACAATGTGTAAATATTTAGGAAGAA**ACTTGA**AAATAGTCAAATCCT
TTGA**ACTGGTGACAATTTT**TAAGAATCAATCTAA**TTTGTTTCAAGGGTAATAATC**ACCA
ATAGACACATTT**CAGCATTTT**ATTTAGTCTATCA**AA**AAATTGGAATTGATATATACACTCATT**T**
ATAGGAGAATGGTTAGGTAGATTTGGTATATTTATGTAGTCATTGAA**AACTTAGT**TTTATA**AA**
GGCCAATCTTGTA**AACTGATTCTTGTGTGATAACATT**CAGTGAA**AA**AGCATGAGACAATTAG**A**
AAGCATGATACAATGAATA**AA**ATA**AAAACTGGAAAGAGA**ACCATCAA**AA**ATGCTAA

FIGURE 92

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA190803
><subunit 1 of 1, 280 aa, 1 stop
><MW: 31222, pI: 7.40, NX(S/T): 1
MFAYVFHDEFVASMIIKIPSDTFTIIPDFDIYYVYGFSSGNFVYFLTTLQPEMVSPPGSTTK
EQVYTSKLVRLCKEDTAFNSYVEVPIGCERSGVEYRLLQAAYLSKAGAVLGRTLGVHPDD
DLLFTVFESKGQKRKMKSLEDESALCIFILKQINDRIKERLQSCYRGEGTLDLAWLKVKDIP
CSSALLTIDDNFCGLDMNAPLGVS DMVRGIPVFTEDRDRMTSVIAYVYKNHSLAFVGTKS
GKLKKVPGTSLCPTLELQTGPRSHRATVTLELLFSSCSSN
```

Important features of the protein:

N-glycosylation site:

Amino acids 230-233

N-myristoylation sites:

Amino acids 87-92;107-112;194-199;237-242

FIGURE 93

CCTTATCAGACAAAGGACGAGATGGAAAATACAAGATAATTTACAGTGGAGAAGAATTAGAA
 TGTAACCTGAAAGATCTTAGACCAGCAACAGATTATCATGTGAGGGTGTATGCC**ATG**TACAA
 TTCCGTAAAGGGATCCTGCTCCGAGCCTGTTAGCTTCACCACCCACAGCTGTGCACCCGAGT
 GTCCTTTCCCCCTAAGCTGGCACATAGGAGCAAAAGTTCAC**T**AACCCTGCAGTGGAAGGCA
 CCAATTGACAACGGTTCAAAAATCACCAACTACCTTTTAGAGTGGGATGAGGGAAAAAGAAA
 TAGTGTTTCAGACAGTGTCTTTCGGGAGCCAGAAGCACTGCAAGTTGACAAAGCTTTGTC
 CGGCAATGGGGTACACATTACGGCTGGCCGCTCGAAACGACATTGGCACCAGTGGTTATAGC
 CAAGAGGTGGTGTGCTACACATTAGGAAATATCCCTCAGATGCCTTCTGCACTAAGGCTGGT
 TCGAGCTGGCATCACATGGGTACGTTGCAGTGGAGTAAGCCAGAAGGCTGTT**C**ACCCGAGG
 AAGTGATCACCTACACCTTGGAAATTCAGGAGGATGAAAATGATAACCTTTCCACCCAAAA
 TACACTGGAGAGGATTTAACCTGTACTGTGAAAAATCTCAAAGAAGCACACAGTATAAATT
 CAGGCTGACTGCTTCTAATACGGAAGGAAAAAGCTGTCCAAGCGAAGTTCTTGTTTGTACGA
 CGAGTCTGACAGGCTGGACCTCCTACCAGACCGCTTGTCAAAGGCCAGTTACATCTCAT
 GGCTTTAGTGTCAAAATGGGATCCCCCTAAGGACAATGGTGGTTCAGAAATCCTCAAGTACTT
 GCTAGAGATTACTGATGGAAATTCTGAAGCGAATCAGTGGGAAGTGGCCTACAGTGGGTTCGG
 CTACCGAATACACCTTCACCCACTTGAAACCAGGCACTTTGTACAAACTCCGAGCATGCTGC
 ATCAGTACCGGCGGACACAGCCAGTGTCTGAAAGTCTCCCTGTTTCGCACACTAAGCATTGC
 ACCAGGTCAATGTCGACCACCGAGGGTTTTGGGTAGACCAAAGCACAAAGAAGTCCACTTAG
 AGTGGGATGTTCTGTCATCGGAAAGTGGCTGTGAGGTCTCAGAGTACAGCGTGGAGATGACG
 GAGCCCGAAGACGTAGCCTCGGAAAGTGTACCATGGCCAGAGCTGGAGTGCACCGTCGGCAA
 CCTGCTTCTGGAACCGTGTATCGCTCCGGGTGAGGGCTCTGAATGATGGAGGGTATGGTC
 CCTATTCTGATGTCTCAGAAATTACCACTGCTGCAGGGCCTCCTGGACAATGCAAAGCACCT
 TGTATTTCTTGACACCTGATGGATGTGTCTTAGTGGGTTGGGAGAGTCTGATAGTTCTGG
 TGCTGACATCTCAGAGTACAGGTTGGAATGGGGAGAAGATGAAGAATCCTTAGAACTCATTT
 ATCATGGGACAGACACCCGTTTTGAAATAAGAGACCTGTTGCCTGCTGCACAGTATTGCTGT
 AGACTACAGGCCTTCAATCAAGCAGGGGCGAGGGCCGTACAGTGAACCTTGCTCTTGCCAGAC
 GCCAGCGTCTGCCCCTGACCCCGTCTCCACTCTCTGTGTCTGGAGGAGGAGCCCTTGATGCC
 TACCCTGATTACCTTCTGCGTGCCTTGTA**T**GAACCTGGGAAGAGCCGTGCAATAACGGATC
 TGAAATCCTTGCTTACACCATTGATCTAGGAGACACTAGCATTACCGTGGGCAACACCACCA
 TGCATGTTATGAAAGATCTCCTTCCAGAAACCACCTACCGGATCAGAATTCAGGCTATAAAT
 GAAATTGGAGCTGGACCATTTAGTCAGTTCATTAAAGCAAAA**A**CTCGGCCATTACCACCCTT
 GCCTCCTAGGCTAGAATGTGCTGCTGCTGGTCCCTCAGAGCCTGAAGCTAAATGGGGAGACA
 GTAACCTCAAGACACATGCTGCTGAGGACATTGTGTACACACTACAGCTGGAGGACAGAAAC
 AAGAGGTTTTATTTCAATCTACAGAGGACCCAGCCACACCTACAAGGTCCAGAGACTGACGGA
 ATTCACATGCTACTCCTTCAGAATCCAGGCAGCAAGCGAGGCTGGAGAAGGGCCCTTCTCAG
 AAACCTATACCTTCAGCACAAACCAAAAGTGTCCCCCACCATCAAAGCACCTCGAGTAACA
 CAGTTAGAAGTAAATTCATGTGAAATTTTATGGGAGACGGTACCATCAATGAAAGGTGACCC
 TGTTAACTACATTCTGCAGGTATTGGTTGGAAGAGAATCTGAGTACAAACAGGTGTACAAGG
 GAGAAGAAGCCACATTCCAAATCTCAGGCCTCCAGACCAACACAGACTACAGGTTCCGCGTA
 TGTGCGTGTGTCGCTGTTTAGACACCTCTCAGGAGCTAAGCGGAGCCTTCAGCCCCCTGTC
 GGCTTTTGATTACAACGAAGTGAGGTACGCTTACAGGGGACATGGGGAGCTTAGATGATC
 CCAAAATGAAGAGCATGATGCCTACTGATGAACAGTTTGACGCCATCATTGTGCTTGGCTTT
 GCAACTTTGTCCATTTTATTTGCCTTTATATTACAGTACTTCTTAATGAAG**TAA**ACCCAACA
 AAAGTAGAGGTATGAATTAATGCTACACATTTTAATACACACATTTATTTCAGATACTCCCT
 TTTTAAAGCCCTTTTGTTTTTTGTATTTATATACTCTGTTTTACAGATTTAGCTAGAAAAAA
 ATGTCAGTGTTTTGGTGCACCTTTTTGAAATGCAAAACTAGGAAAAGGTTAAACTGGATTTT
 TTTTAAAAA

FIGURE 94

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA191064

><subunit 1 of 1, 847 aa, 1 stop

><MW: 93607, pI: 5.33, NX(S/T): 3

MYNSVKGSCSEPVSFTTHSCAPECPFPKLAHRSKSSSLTLQWKAPIDNGSKITNYLLEWD
EGKRNSGFRQCFFGSQKHCKLTKLCPAMGYTFRLAARNDIGTSGYSQEVVVCYTLGNIPQM
PSALRLVRAGITWVTLQWSKPEGCSPEEVITYTLEIQEDENDNLFHPKYTGEDLTCTVKN
LKRSTQYKFRLTASNTEGKSCPSEVLVCTTSPDRPGPPTRPLVKGPVTSHGFSVKWDPPK
DNGGSEILKYLLEITDGNSEANQWEVAYSGSATEYTFTHLKPGTLYKLRACCISTGGHSQ
CSESLPVRTL SIAPGQCRPPRVLGRPKHKEVHLEWDVPASESGCEVSEYSVEMTEPEDVA
SEVYHGPELECTVGNLLPGTVYRFRVRALNDGGYGPYSDVSEITTAAGPPGQCKAPCISC
TPDGCVLVGWESPSSGADISEYRLEWGEDEESLELIYHGTDRFEIRDLLPAAQYCCRL
QAFNQAGAGPYSELVLCQTPASAPDPVSTLCVLEEEPLDAYPDSPSACLVLNWEEPCNNG
SEILAYTIDLGDTSITVGNTTMHVMKDLLPETTYRIRIQAINAIGAGPFSQFIKAKTRPL
PPLPPRLECAAAGPQSLKLKWGDSNSKTHAAEDIVYTLQLEDNRNKRIFISYRGPSHTYKV
QRLTEFTCYSFRIQAASEAGEGPFSEYTFSTTKSVPTIKAPRVLTQLEVNSCEILWETV
PSMKGDPVNYILQVLVGRESEYKQVYKGEEATFQISGLQTNTRYRFRVCACRRCLDTSQE
LSGAFSPSAAFVLQRSEVMLTGDMGSLDDPKMKSMMPPTDEQFAAIIVLGFATLSILFAFI
LQYFLMK

Important features of the protein:

Transmembrane domain:

Amino acids 823-843

N-glycosylation sites:

Amino acids 48-51;539-542;559-562

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 63-66;182-185

Tyrosine kinase phosphorylation sites:

Amino acids 387-394;662-669

N-myristoylation sites:

Amino acids 49-54;257-262;343-348;437-442;757-762

Amidation site:

Amino acids 61-64

ATP/GTP-binding site motif A (P-loop):

Amino acids 193-200

Fibronectin type III domain:

Amino acids 22-106;118-203;215-302;314-398;
410-492;504-590;601-685;697-778

FIGURE 95A

CAATTCGGCCTCGCTCCTTGTGATTGCGCTAAACCTTCCGTCCTCAGCTGAGAACGCTCCACCACCTCCCCGGA
TCGCTCATCTCTTGGCTGCCCCTCCCACTGTTCTGTATGTTATTTTACTCCCCGTATCCCCACTCGTTCCTTAC
AATTCGTAGGTGAGTGGTCCAGCTGGTGCCTGGCCTGTGTCTCTTGGATGCCCTGTGGCTTCAGTCCGTCTC
CTGTTGGCCACCACCTCGTCCCTGGGCGCCTGATACCCAGCCCAACAGCTAAGGTGTGGATGGACAGTAGGG
GGCTGGCTTCTCTCACTGGTCAGGGGTCTTCTCCCCGTGTCTGCCCTCCCGAGCTAGGACTGCAGAGGGGCTAT
CATGGTGTCTGCAGGCCCTCGTGTCTCGCTGTGTCTGCCAGCCTCACACTGCTGGTGTCCACCTCTCCA
GCTCCCAGGATGTCTCCAGTGAGCCAGCAGTGAGCAGCAGCTGTGCCCTTAGCAAGCACCCACCGTGGCC
TTTGAAGACCTGCAGCCGTGGGTCTCTAACTTCACCTACCCTGGAGCCCGGATTTCTCCAGCTGGCTTTGGA
CCCCCTCCGGGAACCACTCATCGTGGGAGCCAGGAACCTACCTCTTCAGACTCAGCCTTGCCAATGTCTCTCTTC
TTCAGGCCACAGAGTGGGCTCCAGTGAGGACACGCGCCGCTCCTGCCAAAGCAAAGGGAAGACTGAGGAGGAG
TGTCAGAACTACGTGCGAGTCTGTATCGTCCGCGCCGGAAGGTGTTATGTGTGGAACCAATGCCTTTTCCCC
CATGTGCACCCAGCAGAGGTGGGAACCTCAGCCGCACTATTGAGAAGATCAATGGTGTGGCCCTGCCCCCT
ATGACCCACGCCACAACCTCCACAGCTGTCATCTCTCTCCAGGGGAGCTCTATGCAGCCACGGTATCGACTTC
TCAGGTCCGGACCCCTGCCATCTACCGCAGCCTGGGCAGTGGGCCACCGCTTCGCACTGCCCAATATAACTCCAAG
TGGCTTAATGAGCCAACTTCGTGGCAGCCTATGATATTGGGCTGTTTGCATACTTCTTCTGCGGGAGAACGC
AGTGGAGCAGACTGTGGACGCACCGTGTACTCTCGCGTGGCCCGCGTGTGCAAGAATGACGTGGGGGGCCGAT
TCCTGCTGGAGGACATGACCACTTTCATGAGGCCCGGCTCAACTGCTCCCGCCGGCGAGGTCCCTCTC
TACTATAACGAGCTGCAGAGTGCCTTCCACTTGGCCGAGCAGGACCTCATCTATGGAGTTTTTACAACCAACGT
AAACAGCATCGCGCTTCTGCTGTCTGCCCTTCAACCTCAGTGTATCTCCAGGCTTTCAATGGCCCATTTTC
GCTACCAGGAGAACCCAGGGCTGCCCTGGCTCCCCATAGCCAACCCCATCCCCAATTTCCAGTGTGGCACCCCTG
CCTGAGACCCGTCCCAACGAGAACTGACGGAGCAGCCTGCAGGACGCGCAGCGCCTCTTCTGTGAGCGA
GGCCGTGCAGCCGGTGACACCCGAGCCCTGTGTACCCAGGACAGCGTGCCTTCTCACACCTCGTGGTGGACC
TGGTGCAGGCTAAAGACACGCTCTACCATGTACTCTACATTGGCACCGAGTCCGGGACCATCTGAAGGCGCTG
TCCACGGCGAGCCGACGCTCCACGGCTGCTACCTGGAGGAGCTGCACGTGCTGCCCGCCGGGCGCCGAGCC
CCTGCGACCCGTGCCATCTCTGCACAGCGCCCGCGCTCTTCTGTGGGCTGAGAGACGGCGTCTGCGGGTCC
CACTGGAGAGGTGCGCGCCTACCGCAGCCAGGGGGCATGCTGGGGGGCCGGGACCCGCTACTGTGGCTGGGAC
GGGAAGCAGCAACGTTGCAGCACACTCGAGGACAGCTCCAACATGAGCCTCTGGACCCAGAACATCACCGCTG
TCCTGTGCGGAATGTGACACGGGATGGGGCTTCGCGCCATGGTCACCATGGCAACCATGTGAGCACTTGGATG
GGGCAACTCAGGCTCTTGCTGTGTGAGCTCGATCTGTGATTCCCCCTCGACCCCGCTGTGGGGGCTTGAC
TGCCTGGGGCCAGCCATCGACATCGCCAACTGCTCCAGGAATGGGGCGTGGACCCCGTATCGTGGCGCT
GTGCAGCAGCTCTGTGGCATCGGCTTCCAGGTCCGCCAGCGAAGTTGCAGCAACCTGCTCCCCGCCACGGGGG
CGCATCTTCTGTGGGCAAGAGCCGGGAGGAACGTTCTGTAATGAGAACACGCTTGGCCGGTGGCCATCTTCTG
GGCTTCTTGGGCTCTTGGAGCAAGTGACAGCAACTGTGGAGGGGATGCAGTCCGGCGTGGGCTGCG
AGAACGGCAACTCCTGCTGGCTGGCTGGCGGAGTTCAAGACGTGCAACCCGAGGGCTGCCCGAAGTGGGCGC
AACACCCCTTGGACGCGCTGGCTGCCGTGAACGTGACGAGGGCGGGGACGGCAGGAGCAGCGTTCCGCTT
CACCTGCCGCGCGCCCTTGCAGACCCGACGGCTGCAGTTCCGAGGAGAAGGACCGAGACGAGGACCTGTC
CCGCGGACGGCTCCGGCTCTGCGACACCGACGCTTGGTGGAGGTCTCCTGCGCAGCGGGAGCACCTCCCCG
CACACGGTGAGCGGGGCTGGGCGCCTGGGCCCCGTGCTCTCTGCTCCCGGAGTGGAGTGGGCTTCCG
CGTCCGCAAGAGAACGTGCACTAACCCGAGCCCGCAACGGGGGCTGCCCTGGCTGGGCGATGCTGCCGAGT
ACCAGGACTGCAACCCCAAGGCTTGCCAGTTCCGGGTGCTTGGTCTGCTGGACCTCATGGTCTCCATGCTCA
GCTTCTGTGGTGGGGTCACTATCAACGCACCCGTTCTGCAACGAGCCCGCACCTCCCCAGGTGAGGACAT
CTGTCTCGGGCTGCACACGGAGGAGCACTATGTGCCACACAGGCTGCCAGGCTGGTCCGCTGGTCTGAGT
GGAGTAAGTGCACTGACGACGGAGCCGAGCCGAAGCCGGCACTGTGAGGAGCTCTCCAGGTTCCAGCGCC
TGTGCTGGAAACAGCAGCCAGAGCCGCCCCCTGCCCTACAGCGAGATTCCCGTCATCTGCCAGCCTCCAGCAT
GGAGGAGGCCACCGACTGTGAGGTAAAAGAAACCGACCTACCTCATGCTCGGCTCTCCAGCCCTCCAGCA
CCCCACTCCAAAGTCTGGACTCTTTCCACATCCTGCTCCAGACAGCCAAGCTTTGTTGGGGTCCCCACTGCTTT
GAGATGGGTTCAATCTCATCCACTTGGTGGCCACGGGCTCTCTGCTTCTTGGGCTCTGGGCTCTTGACCCCTA
GCAGTGTACCTGTCTTGCCAGCACTGCCAGCGTCACTCCAGGAGTCCACACTGGTCCATCTGCCACCCCAACC
ATTTGCACTACAAGGGCGGAGGCACCCGAAGAATGAAAGTACACACCATGGAATTCAGACCCCTGAACAAG
AATAACTTGATCCCTGATGACAGAGCAACTTCTACCCATTGCAGCAGACCAATGTGTACACGACTACTACTA
CCCAAGCCCCCTGAACAAACACAGCTTCCGGCCCCGAGGCTCACCTGGACAACGGTGTCTCCCCAACAGCTGAT
ACCGCCGTCTGGGGACTTGGGCTTCTGCTTTCATAAGGCACAGAGCAGATGGAGATGGGACAGTGGAGCCAG
TTTGGTTTTCTCCCTCTGCACTAGGCCAAGAAGTGTGCTTGCCTTGCCTGTGGGGGTCCCATCCGGCTTCAGAGA
GCTCTGGCTGGCATTGACCATGGGGGAAAGGGCTGGTTTTCAGGCTGACATATGGCCGAGGTCCAGTTAGCCCC
AGGTCTCTCATGGTTATCTTCAACCCACTGTACGCTGACACTATGCTGCCATGCCCTGGGCTGTGGACCTACT
GGGCAATTTGAGGAATGGAGAATGGAGATGGCAAGAGGGCAGGCTTTTAAAGTTTGGGTTGGAGCAACTTCTCT
TGGCCCCCACAAGTGTAGTCTGCTTCTCCAGCTGGCCCCAAAAAAGGCCTTTGTCTACATCTGATTATCTCT
GAAAGTAATCAATCAAGTGGCTCCAGTAGCTCTGGATTTTCTGCCAGGGCTGGGCCATTGTGGTGTGCCCCAG
TATGACATGGGACCAAGGCCAGCGCAGGTTATCCACCTCTGCCTGGAAGTCTATACTCTACCCAGGGCATCCCT
CTGGTCAGAGGCAGTGAAGTGGGAACTGGAGGCTGACCTGTGCTTAGAAGTCTTTAATCTGGGCTGGTACA
GGCCTCAGCCTTGGCCCTCAATGCACGAAAGGTGGCCAGGAGAGGATCAATGCCATAGGAGGCAGAGGCTG
GCCTCTGTGCTCTATGGAGACTATCTTCCAGTTGCTGCTCAACAGAGTTGTTGGCTCAGACCTGCTTGGGAGT

FIGURE 95B

CTCTGCTGGCCCTTCATCTGTTTCAGGAACACACACACACACACTCACACACGCACACACAATCACAATTTGC
TACAGCAACAAAAAGACATTGGGCTGTGGCATTATTAATTAAAGATGATATCCAGTC

FIGURE 96

>>/usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA194909

>>subunit 1 of 1, 1092 aa, 1 stop

><MW: 119324, pI: 8.13, NX(S/T): 14

MPCGFSPSPVAHHLVPGPPDTPAQQLRCGWTVGWLLSLVRGLLPCLPPGARTAEGPIMV
LAGPLAVSLLLPSLTLLVSHLSSSQDVSEPSSEQQLCALSKHPTVAFEDLPWVSNTFY
PGARDFSQLALDPSGNQLIVGARNYLFRLSLANVSLQATEWASSEDTRRSCQSKGKTEE
ECQNYVRVLIVAGRKVFMCGTNAFSPMCTSRQVGNLSRTIEKINGVARCPYDPRHNSTAV
ISSQGEIYAATVIDFSGRDPPIYRSLGSGPPLRTAQYNSKWLNEPNFVAAYDIGLFAYFF
LRENAVEHDCGRTVYSRVARVCKNDVGGRFLEDTWTTFMKARLNCSRPGVEVPFYNELQ
SAFHLPEQDLIYGVTNNVNSIAASAVCAFNLSAISQAFNGPFQYQENPRAAWLPPIANPI
PNFQCGTLPETGPNENLTERSLQDAQRLFLMSEAVQPVTPPCVTQDSVRFSLVVDLVQ
AKDTLYHVLYIGTESGTILKALSTASRLHGCYLEELHVLPPGRREPLRLSLRILHSARAL
FVGLRDGVLRVPLERCAAYRSQGACIGARDPYCGWDGKQRCSTLEDSSNMSLWTQNITA
CPVRNVTRDGGFGPWPQPCHELDGDNSSGCLCRARSCDSPRRCGGDLCLGPAIHIAN
CSRNGAWTPWSSWALCSTSCGIGFQVRQSCSNPAPRHGGRIQVGSREERFCNENTPCP
VPIFWASWGSWSKCSNCGGGMQSRRRACENGNSCLGCGEFKTCNPEGCEVRNTPWTP
WLPVNVTOGGARQEQRFRTCRAPLADPHGLQFGRRTETRTCPADGSGSCDTDALVEVL
LRSGSTSPHTVSGGWAAGWPWSSCSRDELGFRVRKRTCTNPEPRNGGLPCVGDAAEYQD
CNPQACPVRGAWSCWTSWSPCSASC GGHYQRTSCTSPAPSPGEDICLGLHTEALCAT
QACPGWSPWSEWSKCTDDGAQSRSRHCEELLPGSSACAGNSSQSRPCPYSEIPVILPASS
MEEATDCAGKRNRTYIMLRSSQPSSTPLQSLDSFHILLQTAKLCWGPCHCFEMGSISSTWW
PRASPASWALGS

Important features of the protein:

Signal peptide:

Amino acids 1-42

Transmembrane domain:

Amino acids 56-79;373-395

N-glycosylation sites:

Amino acids 117-120;153-156;215-218;236-239;345-348;391-394;
436-439;590-593;597-600;605-608;660-663;785-788;
1000-1003;1032-1035

cAMP- and cGMP-dependent protein kinase phosphorylation sites:

Amino acids 773-776;815-818;875-878

Tyrosine kinase phosphorylation site:

Amino acids 177-185;348-355

N-myristoylation sites:

Amino acids 42-47;50-55;373-378;492-497;543-548;563-568;
630-635;647-652;740-745;810-815;827-832;829-834;
853-858;887-892;910-915;993-998;1073-1078

Amidation sites:

Amino acids 192-195;522-525;813-816;1028-1031

ATP/GTP-binding site motif A (P-loop):

Amino acids 700-707

Cytochrome c oxidase subunit II, copper A binding region signature:

Amino acids 921-929

Growth factor and cytokines receptors family signature 2:

Amino acids 967-973

Sema domain:

Amino acids 126-537

Flexin repeat:

Amino acids 555-602

Thrombospondin type 1 domain:

Amino acids 613-661;668-719;726-769;856-906;913-963;967-1007

FIGURE 97

CAAGCCCTCCCAGCATCCCTCTCCTGTGTTCTCCCGAGTTCTCTACTCAGAGTTGACTGACCAGAGATTTAT
 CAGCTTGGAGGGCTGGAGGTGTGGATCCATGGGGTAGCCTCAACGCATCTGCCCCCACCACCCAGCCAGCTCAT
 GGGCCACGTGGCCTGGCCAGCCTCAGCACCCAGGGCCAGTGAACAGAGCCCTGGCTGGAGTCCAAACATGTGG
 GGCCTGGTGAGGCTCCTGCTGGCCTGGCTGGGTGGCTGGGGCTGCATGGGGCGTCTGGCAGCCCCAGCCCCGGC
 CTGGGCAGGGTCCCAGGAACACCCAGGGCCTGCTCTGCTGCGGACTCGAAGGAGCTGGGTCTGGAACAGTTCT
 TTGTCTATTGAGGAATATGCTGGTCCAGAGCCTGTTCTCATTGGCAAGCTGCACTCGGATGTTGACCGGGGAGAG
 GGCCGCACCAAGTACCTGTTGACCGGGGAGGGGGCAGGCACCGTATTTGTGATTGATGAGGCCACAGGCAATAT
 TCATGTTACCAAGAGCCTTGACCGGGAGGAAAAGCGCAATATGTGCTACTGGCCCAAGCCGTGGACCGAGCCT
 CCAACCGGCCCTGGAGCCCCCATCAGAGTTTCATCATCAAAGTGCAAGACATCAACGACAATCCACCCATTTTT
 CCCCTTGGGCCCTACCATGCCACCGTGCCCGAGATGTCCAATGTGCGGACATCAGTGATCCAGGTGACTGCTCA
 CGATGCTGATGACCCAGCTATGGGAACAGTGCCAAAGCTGGGTGTACACTGTTCTGGATGGACTGCCTTTCTTCT
 CTGTGGACCCCCAGACTGGAGTGGTGCCTACAGCCATCCCCAACATGGACCGGGAGACACAGGAGGAGTTCTTG
 GTGGTGATCCAGGCCAAGGACATGGGCGGCCACATGGGGGGGCTGTGAGGCAGCACTACGGTGACTGTACCGT
 CAGCGATGTCAACGACAACCCCCCAAGTTCACAGAGCCTATACCAGTTCTCCGTGGTGGAGACAGCTGGAC
 CTGGCACACTGGTGGGCGGGCTCCGGGCCAGGACCAGACCTGGGGGACAACGCCCTGATGGCATAACAGCATC
 CTGGATGGGGAGGGGTCTGAGGCCTTCAGCATCAGCACAGACTTGCAAGGGTCGAGACGGGCTCCTCACTGTCCG
 CAAGCCCTAGACTTTGAGAGCCAGCGCTCTACTCTTCCGTGTCGAGGCCACCAACAGCTCATTGACCCAGCC
 TATCTGCGCGAGGGCCCTTCAAGGATGTGGCCTCTGTGCGTGTGGCAGTGCAAGATGCCCCAGAGCCACCTGC
 CTTACCCAGGCTGCCTACCACCTGACAGTGCCGTGAGAACAAGGCCCGGGGACCCTGGTAGGCCAGATCTCCG
 CGGCTGACCTGGACTCCCCCTGCCAGCCCAATCAGATACTCCATCCTCCCCCACTCAGATCCGGAGCGTTGCTTC
 TCTATCCAGCCCCGAGGAAGGCACCATCCATACAGCAGCACCCCTGGATCGCGAGGCTCGCGCCTGGCACAACT
 CACTGTGTGGCTACAGAGCTCGACAGTTCTGCACAGGCCTCGCGCGTGCAAGTGCCCATCCAGACCCTGGATG
 AGAATGACAATGCTCCCCAGCTGGCTGAGCCCTACGATACTTTTGTGTGTGACTCTGCAGCTCCTGGCCAGCTG
 ATTCAGGTCTCCGGGCCCTGGACAGAGATGAAGTTGGCAACAGTAGCCATGTCTCCTTTCAAGGTCTCTGGG
 CCTGATGCCAACTTTTACTGTCCAGGACAACCGAGATGGCTCCGCCAGCCTGTGCTGCCCTCCCGCCCTGCTC
 CACCCCGCCATGCCCCCTACTTGGTTCCCATAGAAGTGTGGGACTGGGGGAGCCGGCGCTGAGCAGCACTGCC
 ACAGTGACTGTTAGTGTGTGCCGCTGCCAGCCTGACGGCTCTGTGGCATCCTGCTGGCCTGAGGCTCACCTCTC
 AGCTGCTGGGCTCAGCACCGGCGCCCTGCTTGCCATCATCACCTGTGTGGGTGCCCTGCTTGCCCTGGTGGTGC
 TCTTCGTGGCCCTGCGGCGGCAGAAAGCAAGAAGCACTGATGGTACTGGAGGAGGAGGACGTCCGAGAGAACATC
 ATCACCTACGACGACGAGGGCGGGCGGAGGAGGACACCGAGGCCTTCGACATCACGGCCTTGCAAGACCCGGA
 CGGGGCGGCCCCCGGCGCGCCGGCCCTCCCGCGCGCGGAGACGTGTGCCCCGGGCGCGGGTGTGCGGCCAGC
 CCAGACCCCCCGGCCCCGCGACGTGGCGCAGCTCCTGGCGCTGCGGCTCCGCGAGGCGGACGAGGACCCGGC
 GTACCCCGTACGACTCGGTGCAGGTGTACGGCTACGAGGGCCGCGGCTCCTCTTGCGGCTCCCTCAGCTCCCT
 GGGCTCCGGCAGCGAAGCGGCGGCGCCCCGCCCCGCGGAGCCGCTGGACGACTGGGTCCGCTCTCCGCACC
 CTGGCCGAGCTGTATGGGGCCAAGGAGCCCCCGGCCCTGAGCGCCCCGGCTGGCCCGGCCACCGCGGGGGG
 GGGGCAGCGGGCACAGGCCTCTGAGTGAGCCCCACGGGTCCAGGCGGGCGGCAGCAGCCAGGGGCCCCAGG
 CCTCCTCCCTGTCTTGTGTCCCTCCTTGCTTCCCCGGGGCACCCCTCGCTCTCACCTCCCTCCTCTGAGTCCG
 TGTGTGTGTCTCTCTCAGGAATCTTGTCTCTATCTGTGACACGCTCCTCTGTCCGGGCTGGGTTCCTGCC
 CTGGCCCTGGCCCTGCGATCTCTCACTGTGATTCTCTCTCTCTCCGTGGCGTTTTGTCTCTGCAGTTCTGAA
 GCTCACACATAGTCTCCCTGCGTCTTCTTGCCCATACACATGCTCTGTGTCTGTCTCTGCCCACATCTCCCT
 TCCTTCTCTCTGGGTCCCTGTGACTGGCTTTTGTGTTTTTCTGTTGTCCATCCCAAAATCAAGAGAACTTCC
 AGCCACTGCTGCCCACCTCCTGCAGGGGATGTTGTGCCCCAGACCTGCCTGCATGGTTCATCCATTACTCAT
 GGCCTCAGCCTCATCTGGCTCCACTGGCCTCCAGCTGAGAGAGGGAACCGCCTGCCTCCAGGGCAAGAGCT
 CCAGCCTCCCGTGTGGCCGCTCCTGGAGCTCTGCCAGCTGCCAGCTTCCCTGGGCATCCAGCCCTGGGC
 ATTGTCTTGTGTGCTTCTGAGGGAGTAGGAAAGGAAAGGGGGAGCGGCTGGGGAAGGGGAAAGAGGGAGGA
 AGGGGAGGGGCTCCATCTCTAATTTCAATAAAACAAACACTTTATTTTGTAAAA

FIGURE 98

MWGLVRLLLLAWLGGWGCMGRLAAPARAWAGSREHFGPALLRTRRSWVWNQFFVIEEYAGP
EPVLIGKLHSDVDRGEGRTKYLLTGEGAGTVFVIDEATGNIHVTKSLDREEKAQYVLLAQ
AVDRASNRPLEPPSEFIIKVQDINDNPPIFPLGPYHATVPEMSNVGTSVIQVTAHDADDP
SYGNSAKLVYTVLDGLPFFSVDPQTGVVRTAIPNMDRETQEEFLVVIQAKDMGGHMGGLS
GSTTVTVTLSDVNDNPPKFPQSLYQFSVVETAGPGTLVGRLRAQDPDLGDNALMAYSILD
GEGSEAFSISTDLQGRDGLLTVRKPLDFESQRSYSFRVEATNTLIDPAYLRRGPFKDVAS
VRVAVQDAPEPPAFTQAAYHLTVPENKAPGTLVGQISAADLDSPASPIRYSILPHSDPER
CFSIQPEEGTIHTAAPLDREARAWNLTVLATELDSSAQASRVQVAIQTLDENDNAPQLA
EPYDTFVCDASAAPQLIQVJRALDRDEVGNSSHVSFQGPLGPDANFTVQDNRDGSASLLL
PSRPAPPRHAPYLVPIELWDWGQPALSSTATVTVSVCRCPDGSVASCWPEAHLAAGLS
TGALLAIITCVGALLALVVLFLVALRRQKQFALMVLEEEDVRENIITYDDEGGGEEDTEAF
DITALQNPDGAAPPAPGPPARRDVLPRARVSRQPRPPGPADVAQLLALRLREADEDPGVP
PYDSVQVYGYEGRGSSCGSLSSLSGSGSEAGGAPGPAEPLDDWGPLFRTLAEELYAKEPPA
P

Signal peptide:

Amino acids 1-16

Transmembrane domain:

Amino acids 597-624

N-glycosylation sites:

Amino acids 446-449;510-513;525-528

N-myristoylation sites:

Amino acids 13-18;206-211;233-238;237-242;238-243;275-280;390-395;
394-399;429-434;583-588;598-603;602-607;612-617;
734-739;738-743;746-751

ATP synthase c subunit signature:

Amino acids 691-712

Cadherins extracellular repeated domain signature:

Amino acids 138-148;247-257

Cadherin domain:

Amino acids 50-141;155-250;264-366;379-470;483-577

Cadherin cytoplasmic region:

Amino acids 625-776

FIGURE 99

[illegible]

FIGURE 100

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA213858
><subunit 1 of 1, 627 aa, 1 stop
><MW: 66189, pI: 7.31, NX(S/T): 5
MAILPLLLCLLPLAPASSPPQSATPSPCPRRCRCQTQSLPLSVLCPGAGLLFVPPSLDRR
AAELRLADNFIASVRRRDLANMTGLLHLSLSRNTIRHVAAGAFADLRALRALHLDGNRLT
SLGEGQLRGLVNLRLHLILSNNQLAALAAGALDDCAETLEDLDLSYNNLEQLPWEALGRLG
NVNTLGLDHNLLASVPGAFSRLHKLARLDMTSNRLTTIPDPLFSRLPLLARPRGSPASA
LVLAFIGGNPLHCNCELVWLRLAREDDLEACASPPALGGRYFWAVGEEEFVCEPPVVTHR
SPPLAVPAGRPAALRCRAVGDPPEPRVRWVSPQGRLLGNSSRARAFPNGTLELLVTEPGDG
GIFTCIAANAAGEATAAVELTVGPPPPPQLANSTSCDPPRDGDPDALTPPSAASASAKVA
DTGPPTDRGVQVTEHGATAALVQWPDQRPIPGIRMYQIQYNSSADDILVYRMIPAESRSE
LLTDLASGRTYDLCVLAVYEDSATGLTATRPVGCARFSTEPALRPGAPHAPFLGGTMI I
ALGGVIVASVLVFI FVLLMRYKVHGGQPPGKAKIPAPVSSVCSQTNGALGPTPTPAPPAP
EPAALRAHTVVQLDCEPWGPGHEPVG
```

Important features of the protein:

Signal peptide:

Amino acids 1-16

Transmembrane domain:

Amino acids 35-55; 536-556

N-glycosylation sites:

Amino acids 81-84; 338-341; 347-350; 392-395; 461-464

N-myristoylation sites:

Amino acids 116-121; 125-130; 180-185; 186-191; 235-240;
360-365; 361-366; 429-434; 436-441; 505-510;
544-549; 566-571

Leucine Rich Repeat:

Amino acids 60-83; 84-107; 108-131; 132-155; 157-180;
181-203; 204-227

Leucine rich repeat C-terminal domain:

Amino acids 248-293

Immunoglobulin domain:

Amino acids 309-367

Fibronectin type III domain:

Amino acids 424-504

FIGURE 101

CGACTCCATAACCGTGGCCTTGGCCCCAGTCCCCCTGACTTCCGGACTTCAGACCAGATACTGCCCCATATCCCC
TTATGAAGTCTTGGCCAGGCCAACCCTAGGGTGTACGTTTCTAAAGATTAAAGAGGCGGTGCTAAGCTGCAGA
CGGACTTGGGACTCAGCCACTGGTGTAAAGTCAGGCGGGAGGTGGCGCCCAATAAGCTCAAGAGAGGAGGCGGGT
5 TCTGGAAAAAGGCCAATAGCCTGTGAAGGCGAGTCTAGCAGCAACCAATAGCTATGAGCGAGAGGCGGGACTCT
GAGGGAAGTCAATCGCTGCCGCAGGTACCGCCAATGGCTTTTGGCGGGGGCGTTCCCCAACCCCTGCCCTCTCTC
ATGACCCCGCTCCGGGATTATGGCCGGGACTGGGCTGCTGGCGCTGCGGACGCTGCCAGGGCCCAGCTGGGTGC
GAGGCTCGGGCCCTTCCGTGCTGAGCCGCTGCAGGACGCGGCCGTGGTGGCGCCTGGCTTCCTGAGCACGGCA
GAGGAGGAGACGCTGAGCCGAGAACTGGAGCCCGAGCTGCCGCCGCCGCGCTACGAATACGATCACTGGGACGC
10 GGCCATCCACGGCTTCCGAGAGACAGAGAAGTCGCGCTGGTCAGAAGCCAGCCGGGCCATCCTGCAGCGCGTGC
AGGCGGCCGCCCTTGGCCCCGGCCAGACCCTGCTCTCCTCCGTGCACGTGCTGGACCTGGAAGCCCGCGGCTAC
ATCAAGCCCCACGTGGACAGCATCAAGTTCTGCGGGGCCACCATCGCCGGCGTGTCTCTCCTGTCTCCAGCGT
TATGCGGCTGGTGCACACCCAGGAGCCGGGGAGTGGCTGGAACCTTTGCTGGAGCCGGGCTCCCTCTACATCC
TTAGGGGCTCAGCCGTTATGACTTCTCCCATGAGATCCTTCGGGATGAAGAGTCTTCTTTGGGGAACGCCGG
15 ATTCCCCGGGGCCGGCGCATCTCCGTGATCTGCCGCTCCCTCCCTGAGGGCATGGGGCCAGGGGAGTCTGGACA
GCGCCCCCAGCCTGCTGACCCCCAGCTTTCTACAGACACCAGATTGTGAATAAAGTTGGGGAATGGACAGCCT

FIGURE 102

MAGTGLLALRTLPGPSWVRGSGPSVLSRLQDAAVVRPGFLSTAEETLSRELEPELRRRRYEYDHWDAAIHGFR
ETEKSRWSEASRAILQRVQAAAFGPGQTLSSVHVLDEARGYIKPHVDSIKFCGATIAGLSLLSPSVMRLVHT
QEPGEWLELLLEPGSLYILRGSARYDFSHEILRDEESFFGERRIPRGRISVICRSLPEGMPGESGQPPAC

Important features of the protein:

Signal peptide:
1-18

Transmembrane domain:
None

cAMP- and cGMP-dependent protein kinase phosphorylation site.
196-199

N-myristoylation site.
20-25
129-134
208-213

Amidation site.
194-197

FIGURE 103

CTCCCCGGCGCCGAGGCAGCGTCCTCCTCCGAAGCAGCTGCACCTGCAACTGGGCAGCCTGGACCCCTCGTGCC
 CTGTTCCCGGACCTCGCGCAGGGGGGCCCCGGGACACCCCTGCGGGCCGGGTGGAGGAGGAAGAGGAGGAG
 GAGGAAGAAGACGTGGACAAGGACCCCCATCCTACCCAGAACACCTGCCTGCGCTGCCGCCACTTCTCTTTAAG
 GGAGAGGAAAAGAGAGCCTAGGAGAACCATGGGGGGCTGCGAAGTCCGGGAATTTCTTTTGCAATTTGGTTTTCT
 TCTTGCCCTCTGCTGACAGCGTGGCCAGGCGACTGCAGTCACGTCTCCAACAACCAAGTTGTGTTGCTTGATACA
 ACAACTGTACTGGGAGAGCTAGGATGGAAAACATATCCATTAAATGGGTGGGATGCCATCACTGAAATGGATGA
 ACATAATAGGCCCATTCACACATACCAGGTATGTAATGTAATGGAACCAACCAAAACAACCTGGCTTCGTACAA
 ACTGGATCTCCCGTGATGCAGCTCAGAAAATTTATGTGGAAATGAAATTCACACTAAGGGATTGTAACAGCATC
 CCATGGGTCTTGGGGACTTGCAAAGAAACATTTAATCTGTTTTATATGGAATCAGATGAGTCCCACGGAATTA
 ATTCAAGCCAAACAGTATACAAAGATCGACACAATTGCTGCTGATGAGAGTTTTACCCAGATGGATTGGGTG
 ATCGCATCCTCAAACCTCAACACTGAAATTCGTGAGGTGGGGCCTATAGAAAGGAAAGGATTTTATCTGGCTTTT
 CAAGACATTGGGGCGTGCAATTGCCCTGGTTTCAGTCCGTGTTTTCTACAAGAAATGCCCTTCACTGTTCTGTA
 CTGGCCATGTTTCTGATACCATTCGAAGGTTGATTCTCTCTCTTTGGTTGAAGTACGGGGTCTTGTGTGA
 AGAGTGCTGAAGAGCGTGACACTCCTAACTGTATTGTGGAGCTGATGGAGATTGGCTGGTCTCTTGGAAAGG
 TGCATCTGCAGTACAGGATATGAAGAAATGAGGGTTCTTGCCATGCTTGCAGACCAGGATTCTATAAGCTTT
 TGCTGGGAACACAAAATGTTCTAAATGTCTCCACACAGTTTAAACATACATGGAAGCAACTTCTGTCTGTGAGT
 GTGAAAAGGGTTATTTCCGAGCTGAAAAAGACCCACCTTCTATGGCATGTACCAGGCCACCTTCACTGCTCCTAGG
 AATGTGGTTTTTAACATCAATGAACAGCCCTTATTTTGAATGGAGCCACCAAGTGACACAGGAGGGAGAAA
 AGATCTCACATACAGTGTATCTGTAAGAAATGTGGCTTAGACACCAGCCAGTGTGAGGACTGTGGTGGAGGAC
 TCCGCTTCATCCCAAGACATACAGGCCCTGATCAACAATTCGTGATAGTACTTGACTTTGTGTCTCACGTGAAT
 TACACCTTTGAAATAGAAGCAATGAATGGAGTTTCTGAGTTGAGTTTTTCTCCAAGCCATTACAGCTATTAC
 AGTGACCACGGATCAAGATGCACCTTCCCTGATAGGTGTGGTAAGGAAGGACTGGGCATCCCAAAATAGCTATGCC
 CTATCATGGCAAGCACCTGCTTTTTTCCAATGGAGCCATTCTGGACTACGAGATCAAGTACTATGAGAAAGAACA
 TGAGCAGCTGACCTACTCTTCCACAAGGTCCAAAGCCCCCAGTGTCTATCATCACAGGTCTTAAGCCAGCCACCA
 AATATGTAATTCACATCCGAGTGAGAACTGCGACAGGATACAGTGGCTACAGTCAGAAATTTGAATTTGAAACA
 GGAGATGAAACTTCTGACATGGCAGCAGAACAAAGGACAGATTCTCGTGATAGCCACCGCCGCTGTTGGCGGATT
 CACTCTCTCTGTCATCTCACTTTATTCTTCTTGATCACTGGGAGATGTGAGTGGTACATAAAAGCCAAGATGA
 AGTCAGAAAGAGAAGAGAAGAAACCACTTACAGAATGGGCATTTGCGCTTCCCGGGAATTAACCTTACATTGAT
 CCAGATACATATGAAGACCCATCCCTAGCAGTCCATGAATTTGCAAAGGAGATTGATCCCTCAAGAATTCGTAT
 TGAGAGAGTCATTGGGGCAGCTGAATTTGGAGAAGTCTGTAGTGGGCGTTTGAAGACACCAGGGAAAAGAGAGA
 TCCAGTTGCCATTAAACTTTGAAAGGTGGCCACATGGATCGGCCAAAGAAGAGATTTTCTAAGAGAAGCTAGT
 ATCATGGGCCAGTTTGACCATCCAACATCATTCGCCCTAGAAGGGGTGTCAACAAAAGATCCTTCCCGGCCAT
 TGGGGTGGAGGCGTTTTTGGCCAGCTTCTGAGGGCAGGGTTTTTAAATAGCATCCAGGCCCGCATCCAGTGC
 CAGGGGGAGGATCTTTGCCCCCAGGATTCCTGCTGGCAGACCAGTAATGATTGTGGTGGAAATATATGGAGAAT
 GGATCCCTAGACTCCTTTTTTGGCGAAGCATGATGGCCACTTCACAGTCATCCAGTTGGTCCGAATGCTCCGAGG
 CATTGCATCAGGCATGAAGTATCTTTCTGATATGGGTTATGTTTCATCGAGACCTAGCGGCTCGGAATATACTGG
 TCAATAGCAACTTAGTATGCAAAGTTTCTGATTTTGGTCTCTCCAGAGTGCTGGAAGATGATCCAGAAGCTGCT
 TATACAACAACTGGTGGAAAAATCCCCATAAGGTGGACAGCCCCAGAAGCCATCGCCTACAGAAAATTTCTCTC
 AGCAAGCGATGCATGGAGCTATGGCATTGTCATGTGGGAGGTGATGCTTATGGAGAGAGACCTTATTGGGAAATG
 TCTAACCAAGATGTCACTCTGTCCATTGAAGAAGGTACAGACTTCCAGCTCCCATGGGCTGTCCAGCATCTCT
 ACACCAGCTGATGCTCCACTGCTGGCAGAAAGGAGAGAAATCACAGACCAAAATTTACTGACATTGTGAGCTTCC
 TTGACAACTGATCCGAAATCCCAGTGCCCTTCACACCCTGGTGGAGGACATCCTTGTAATGCCAGAGTCCCCT
 GGTGAAGTTCCGGAATATCCTTTGTTTGTACAGTTGGTGACTGGCTAGATTCTATAAAGATGGGGCAATACAA
 GAATAACTTCTGTGGCAGCAGGGTTTACAACATTTGACCTGATTTCAAGATGAGCATTGATGACATTAGAAGAA
 TTGGAGTCATACTTATTGGACACCAGAGACGAATAGTCAGCAGCATACAGACTTTACGTTTACACATGATGCAC
 ATACAGGAGAAGGGATTTTCATGTATGAAGTACCACAAGCACCTGTGTTTTGTGCCTCAGCATTTCTAAATGA
 ACGATATCCTCTCTACTACTCTCTCTCTGATTCTCCAAACATCACTTCACAACTGCAGTCTTCTGTTTCAGAC
 TATAGGCACACACCTTATGTTTATGCTTCCAACCAGGATTTAAATCATGCTACATAAATCCGTTCTGAATAA
 CCTGCAACTAAAAAAAAAAAAAAAAA

FIGURE 104

```
>>/usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA222653
><subunit 1 of 1, 1036 aa, 1 stop
><MW: 116379, pI: 6.94, NX(S/T): 5
MGGCEVREFLLQFGFFLLTAWPGDCSHVSNQVLLDTTTVLGEELGWKTYPLNGWDAI
TEMDEHNRPIHTYQVCNVMPEPNQNNWLRTNWISRDAQKIYVEMKFTLRDCNSIPWVLGT
CKETFNLFYMESDESHGKFKPNQYTKIDTIAADESFTQMDLGDRIKLKINTEIREVGPIE
RKGFYLAQFDIGACIALVSVRVFYKKCPFTVRNLAMFPDTIPRVDSSSLVEVRGSCVKSA
EERDTPKLYCGADGDLVPLGRCICSTGYEEIEGSCHACRPGFYKAFAGNTKCSKCPPHS
LTMEATSVCQCEKGYFRAEKDPPSMACRPPSAPRNVVFNINETALILEWSPPSDTGGR
KDLTYSVICKKCGLDTSQCEDCGGGLRFIPRHTGLINNSVIVLDFVSHVNYTTFEIAMNG
VSELSFSPKPFMTAITVTTDQDAPSLIGVVRKDWASQNSIALSWQAPAFSNGAILDYEIKY
YEKEHEQLTYSSTRSKAPSVIITGLKPATKYVFHIRVRTATGYSGYSQKFEFETGDETS
MAAEQGGQILVIATAAVGGFTLLVILTFLITGRCQWYIKAKMKSEEKRRNHLQNGHLRF
PGIKTYIDPDYEDPSLAVHEFAKEIDPSRIRIERVIGAGEFGEVCSGRKTPGKREIPV
AIKTLKGGHMDRQRDFLREASIMGQFDHPNIIIRLEGVVTKRSPFAIGVEAFPCPSFLRAG
FLNSIQAPHVPVPGGSLPPRIIPAGRPVMIVVEYMENGSLDSFLRKHGDHFTVIQLVGMRLR
GIASGMKYLSDMGYVHRDLAARNILVNSNLVCKVSDFGLSRVLEDDPEAAAYTTGCKIP1
RWTAPEAIAIRKFSSASDAWSYGIWMWVMSYGERPYWEMSNQDVILSIEEGYRLPAPMG
CPASLHQLMLHCWKERNHRPKFTDIVSFLDKLIRNPSALHTLVEDILVMPESPGEVPEY
PLFVTVGDWLDSIKMGQYKNNFVAAGFTTFDLISRMSIDDIRRIGVILIGHQRRIVSSIQ
TLRLHMMHIQEKGFHV
```

Important features of the protein:

Signal peptide:

Amino acids 1-22

Transmembrane domain:

Amino acids 551-571

N-glycosylation sites:

Amino acids 343-346;397-400;410-413;756-759

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 851-854

Tyrosine kinase phosphorylation sites:

Amino acids 483-490;604-612;787-794

N-myristoylation sites:

Amino acids 192-197;274-279;289-294;373-378;394-399;504-509;
757-762;777-782;781-786;900-905;976-981

Amidation site:

Amino acids 358-361;653-656

Tyrosine protein kinases specific active-site signature:

Amino acids 794-806

Receptor tyrosine kinase class V signature 1:

Amino acids 192-208

Ephrin receptor ligand binding domain:

Amino acids 34-207

pkinese Protein kinase domain:

Amino acids 631-927

Fibronectin type III domain:

Amino acids 332-425;440-527

SAM domain (Sterile alpha motif):

Amino acids 959-1023

FIGURE 105

GGCGGCGGGCTGCGCGGAGCGGCGTCCCCTGCAGCCGCGGACCGAGGCAGCGGCGGCACCTGCCGCGCGAGCAA
TGCCAACTGAGTACACCTATGTGAACTGAGAAGTGATTGCTCGAGGCCTTCCCCTGCAATGGTACACCCGAGCT
CAAAGCAAGATGAGAAGGCCAGCTTGTTATTAAAGACATCCTCAAATGTACATTGCTTGTTGGAGTGTG
GATCCTTTATATCCTCAAGTTAAATTATACTACTGAAGAATGTGACATGAAAAAATGCATTATGTGGACCCCTG
ACCATGTAAAGAGAGCTCAGAAATATGCTCAGCAAGTCTTGCGAAGGAATGTCGTCCCAAGTTTGCCAGACA
TCAATGGCGCTGTTATTTGAGCACAGGTATAGCGTGGACTTACTCCCTTTTGTGCAGAAGGCCCCCAAAGACAG
TGAAGCTGAGTCCCAAGTACGATCCTCCTTTTGGGTTCGGAAGTTCTCCAGTAAAGTCCAGACCCCTCTTGGAAC
TCTTGCCAGAGCACGACCTCCCTGAACACTTGAAAGCCAAGACCTGTGCGCGCTGTGTGGTTATTGGAAGCGGA
GGAATACTGCACGGATTAGAAGTGGGCCACACCCCTGAACCAGTTCGATGTTGTGATAAGGTTAAACAGTGCACC
AGTTGAGGGATATTGAGAACATGTTGGAAATAAACTACTATAAGGATGACTTATCCAGAGGGCGCACCACTGT
CTGACCTTGAATATTATTCCAATGACTTATTTGTTGCTGTTTTATTTAAGAGTGTTGATTCAACTGGCTTCAA
GCAATGGTAAAAAGGAAACCCTGCCATTCTGGGTACGACTCTTCTTTTGAAGCAGGTGGCAGAAAAAATCCC
ACTGCAGCCAAAACATTTGAGGATTTGAATCCAGTTATCATCAAAGAGACTGCCTTTGACATCCTTCAGTACT
CAGAGCCTCAGTCAAGGTTCTGGGGCCGAGATAAGAACGTCCCCACAATCGGTGTCATTGCCGTTGTCTTAGCC
ACACATCTGTGCGATGAAGTCAGTTTGGCGGGTTTGGATATGACCTCAATCAACCCAGAACACCTTTGCACTA
CTTCGACAGTCAATGCATGGCTGCTATGAACTTTCAGACCATGCATAATGTGACAACGGAAACCAAGTCCCTCT
TAAAGCTGGTCAAAGAGGGAGTGGTGAAAGATCTCAGTGGAGGCATTGATCGTGAATTTGAACACAGAAAACC
TCAGTTGAAAATGCAACTCTAACTCTGAGAGCTGTTTTTGACAGCCTTCTTGATGTATTTCTCCATCCTGCAGA
TACTTTGAAGTGCAGCTCATGTTTTAACTTTTAAATTTAAAAACACAAAAAAATTTTAGCTCTTCCCCTTTT
TTTTTCTATTTATTTGAGGTCAGTGTTTGTGTTTGCACACCATTTTGTAAATGAAACTTAAGAATTGAATTGG
AAAGACTTCTCAAAGAGAATTGTATGTAACGATGTTGTATTGATTTTAAAGAAAGTAATTTAATTTGTAAACT
TCTGCTCGTTTACACTGCACATTGAATACAGGTAACATAATTGGAAGGAGAGGGGAGGTCACCTTTTGTATGGTG
GCCCTGAACCTCATTCTGGTTCCCTGCTGCGCTGCTTGGTGTGACCCACGGAGGATCCACTCCCAGGATGACGT
GCTCCGTAGCTCTGCTGCTGATACTGGGTCTGCGATGCAGCGCGGTGAGGCCTGGGCTGGTTGGAGAAGGTCAC
AACCCTTCTCTGTTGGTCTGCCTTCTGCTGAAAGACTCGAGAACCAACCAGGGAAGCTGTCTGGAGGTCCCTG
GTCGGAGAGGGACATAGAATCTGTGACCTCTGACAACTGTGAAGCCACCCTGGGCTACAGAAACCACAGTCTTC
CCAGCAATTATTACAATTCTTGAATTCCTTGGGGATTTTTTACTGCCCTTTCAAAGCACTTAAGTGTTAGATCT
AACGTGTTCCAGTGTCTGTCTGAGGTGACTTAAAAATCAGAACAAAACCTCTATTATCCAGAGTCATGGGAGA
GTACACCCTTTCCAGGAATAATGTTTTGGGAAACACTGAAATGAAATCTTCCAGTATTATAAATTGTGTATTTAA

FIGURE 106

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA96897
><subunit 1 of 1, 362 aa, 1 stop
><MW: 41736, pI: 8.80, NX(S/T): 3
MRRPSLLLKDILKCTLLVFGVWILYILKLNVTTEECMDMKMHYVDPDHVKRAQKYAQQVLQK
ECRPKFAKTSMAALLFEHRYSDLLPFVQKAPKDSEAESKYDPPFGFRKFSSKVQTLLELLPE
HDLPEHLKAKTCRRCVVIGSGGILHGLELGHTLNQFDVVIRLNSAPVEGYSEHVGNKTTIRM
TYPEGAPLSDLEYYSNDLFVAVLFKSVDFNWLQAMVKKETLFFWVRLFFWKQVAEKIPLQPK
HFRILNPVIIKETAFDILQYSEPQSRFWGRDKNVPTIGVIAVVLATHLCDEVSLAGFGYDLN
QPRTPHLYFDSQCMAAMNFQTMHNVTTETKFLCLKLVKEGVVKDLSGGIDREF
```

Important features of the protein:

Transmembrane domain:

Amino acids 11-27;281-297

N-glycosylation sites:

Amino acids 30-34;180-184;334-338

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 2-6;109-113;223-227

N-myristoylation sites:

Amino acids 146-152;150-156;179-185;191-197

FIGURE 107

TGACGCGGGGCGCCAGCTGCCAACTTCGCGCGCGGAGCTCCCCGGCGGTGCAGTCCCGTCCCGGCGGCGCGG
GCGGCATGAAGACTAGCCGCCGCGGCGGAGCGCTCCTGGCCGTGGCCCTGAACCTGCTGGCGCTGCTGTTTCG
CCACCACCGCTTTCTCTACCACGCACTGGTGCCAGGGCACGCAGCGGGTCCCCAAGCCGGGCTGCGGCCAGG
GCGGGCGCGCCAACTGCCCCAACTCGGGCGCCAACGCCACGGCCAACGGCACCGCCGCCCCCGCCGCCGCCG
CCGCCGCCGCCACCGCCTCGGGGAACGGCCCCCCTGGCGGCGCGCTCTACAGCTGGGAGACCGGCGACGACC
GCTTCCTCTTCAGGAATTTCCACACCGGCATCTGGTACTCGTGCGAGGAGGAGCTCAGCGGGCTTGGTGAAA
AATGTCGCAGCTTCATTGACCTGGCCCCGGCGTCGGAGAAAAGGCCTCCTGGGAATGGTCGCCCACATGATGT
ACACGCAGGTGTTCCAGGTCACCGTGAGCCTCGGTCTTGAGGACTGGAGACCCCATTCTGGGACTACGGGT
GGTCCTTCTGCCTGGCGTGGGGCTCCTTTACCTGCTGCATGGCAGCCTCTGTCAACACGCTCAACTCCTACA
CCAAGACGGTCATTGAGTTCGCGGCACAAGCGCAAGGTCTTTGAGCAGGGCTACCGGGAAGAGCCGACCTTCA
TAGACCTTGAGGCCATCAAGTACTTCCGGGAGAGGATGGAGAAGAGGGACGGGAGCGAGGAGGACTTTCACT
TAGACTGCCGCCACGAGAGATACCCTGCCCGACACCAGCCACACATGGCGGATTCTTGGCCCCGGAGCTCCG
CACAGGAAGCACACAGAGCTGAACCGACAGTGCTGGGTCTTGGGGCACTGGGTGGTGACCAAGACCTCAACCTG
GCCCCGCGGACCTCAGGCCATCGCTGGCACACAGCCCCCTGCTGCAAGACCACAGAGTGGTGCCCCAGAACCC
TGGCCTGTGTGCCGTGAACTCAGTCAGCCTGCGTGGGAGATGCCAGGCCTGTCTGCCCATCGCTGCCTGGG
TCCCATGGCCTTGGAATGGGGCCAGGGCAGGCCCAAGGGAATGCACAGGGCTGCACAGAGTGACTTTGGGA
CAGCAGCCCCGGACTCTTGCCATCATCACATGAGCCCTGCTGGGCACAGCTGCGATGCCAGGAGACACATGG
CCACTGGCCACTGAATGGCTGGCACCCACAAGCCAGTCAGGTGCCCAGAGGGGCAGAGCCCTTTGGGGGGCA
GAGAGTGGCTTCTGAAGGAGGGGGCAGTGGCGCAGGCACTGCAGGGGTGTCACACAGCAGGCACACAGCAG
GGGCTCAATAAATGCTTGTTGAACTTGTTTT

FIGURE 108

MKTSRRGRALLAVALNLLALLFATTAFLTTHWCQGTQRVKPGCGQGGRANCPNSGANATANGTAAPAAAA
AAATASGNGPPGGALYSWETGDDRFLFRNFHTGIWYSCEEELSGLGEKCRSFIDLAPASEKGLLGMVAHMM
YTQVFQVTVSLGPEDWRPHSWDYGWSFCLAWGSFTCCMAASVTTLNSYTKTVIEFRHKRKVFEQGYREEPT
FIDPEAIKYFRERMEKRDGSEEDFHLDCRHERYPARHQPHMADSWPRSSAQEAPELNRQCWWVLGHWV

Important features of the protein:

Signal peptide:

1-26

Transmembrane domain:

169-189

N-glycosylation site.

58-61

62-65

Glycosaminoglycan attachment site.

77-80

114-117

Tyrosine kinase phosphorylation site.

202-208

N-myristoylation site.

43-48

47-52

56-61

84-89

104-109

174-179

FIGURE 109

GATTACCAAGCAAGCAACAGTCTAAATATGAAAGGCCATCATTCATCTTACTCTTCTTGCCTCTCCT
TTCTGTAAACACAGCCACCAACCAAGGCAACTCAGCTGATGCTGTAAACAACACAGAAACTG
CGACTAGTGGTCTTACAGTAGCTGCAGCTGATACCACTGAAACTAATTTCCCTGAAACTGCT
AGCACCACAGCAAATACACCTTCTTTCCCAACAGCTACTTCACCTGCTCCCCCATAAATTAG
TACACATAGTTTCTCCACAATTCCTACACCTGCTCCCCCATAAATTAGTACACATAGTTCTT
CCACAATTCCTATACCTACTGCTGCAGACAGTGAGTCAACCACAAATG'AAATTCATTAGCT
ACCTCTGACATAATCACCGCTTCATCTCCAAATGATGGATTAAATCACAAATGGTTCCTTCTGA
AACACAAAGTAACAATGAAATGTCCCCACCACAGAAGACAATCAATCATCAGGGCCTCCCA
CTGGCACCGCTTTATTGGAGACCAGCACCTTAAACAGCACAGGTCCAGCAATCCTTGCCAA
GATGATCCCTGTGCAGATAATTCGTTATGTGTTAAGCTGCATAATACAAGTTTTTGCCTGTG
TTTAGAAGGGTATTACTACAACCTCTTCTACATGTAAGAAAGGAAAGGTATTCCTGGGAAGATT
TCAGTGACAGTATCAGAAACATTTGACCCAGAAGAGAAACATTCCATGGCCTATCAAGACTT
GCATAGTGAAATTACTAGCTTGTTTTAAAGATGTATTTGGCACATCTGTTTATGGACAGACTG
TAATTCCTTACTGTAAGCACATCTCTGTCCACCAAGATCTGAAATGCGTGCTGATGACAAGTTT
GTTAATGTAACAATAGTAACAATTTTGGCAGAAACCACAAGTGACAATGAGAAGACTGTGAC
TGAGAAAATTAATAAAGCAATTAGAAGTAGCTCAAGCAACTTTCTAAACTATGATTTGACCC
TTCCGTGTGATTATTATGGCTGTAACCAGACTGCGGATGACTGCCTCAATGGTTTAGCATGC
GATTGCAAACTGTACCTGCAAAGGCCTAACCCACAGAGCCCTTTCTGCGTTGCTTCCAGTCT
CAAGTGTCCTGATGGCTGCAACGCACAGCACAAAGCAATGCTTAATAAAGAAGAGTGGTGGGG
CCCCTGAGTGTGCGTGCCTGCCCGGTACCAGGAAGATGCTAATGGGAAGTCCCAAAGTGT
GCATTTGGCTACAGTGGACTCGACTGTAAGGACAAATTCAGCTGATCCTCACTATTGTGGG
CACCATCGCTGGCATTGTCTATTCTCAGCATGATAATTGCATTGATTGTCCAGCAAGATCAA
ATAACAAAACGAAGCATATTGAAGAAGAGAACTTGATTGACGAAGACTTTCAAATCTAAAA
CTGCGGTGACAGGCTTCACCAATCTTGAGAGCAGAAGGGAGCGTCTTTCTAAGGTCAAGAT
AACGGCCTCCAGAGACAGCCAGATGCAAATCCCTATTCAAGCCACAGCAGCATGCCCGCC
CTGACTATAGAAATCATAAGAATGTGGAACCCGCCATGGCCCCCAACCAATGTACAAGCTAT
TATTTAGAGTGTTTAGAAAGACTGATGGAGAAGTGAGCACCAGTAAAGATCTGGCCTCCGGG
GTTTTTCTTCCATCTGACATCTGCCAGCCTCTCTGAATGGAAGTTGTGAATGTTTGCAACGA
ATCCAGCTCACCTTGCTAAATAAGAATCTATGACATTAAATGTAGTAGATGCTATTAGCGCTT
GTCAGAGAGGTGGTTTTCTTCAATCAGTACAAAGTACTGAGACAATGGTTAGGGTTGTTTTCT
TTAATTCTTTTCTGTTAGGGCAACAAGAACCATTTCCAATCTAGAGGAAAGCTCCCCAGCA
TTGCTTGCTCCTGGGCAAACATTGCTCTTGAGTTAAGTGACCTAATTCCTGGGAGACATA
CGCATCAACTGTGGAGGTCCGAGGGGATGAGAAGGGATACCCACCATCTTTCAAGGGTCACA
AGCTCACTCTCTGACAAGTCAGAATAGGGGACACTGCTTCTATCCCTCCAATGGAGAGATTCT
GGCAACCTTTGAACAGCCAGAGCTTGCAACCTAGCCTCACCCAAGAAGACTGGAAAGAGAC
ATATCTCTCAGCTTTTTTTCAGGAGGCGTGCTGGGAATCCAGGAACTTTTTGATGCTAATTAG
AAGGCCTGGACTAAAAATGTCCACTATGGGGTGCACTCTACAGTTTTTGAAATGCTAGGAGG
CAGAAGGGGCAGAGAGTAAAAAACATGACCTGGTAGAAGGAAGAGAGGCAAAGGAAACTGGG
TGGGGAGGATCAATTAGAGAGGAGGCACCTGGGATCCACCTTCTTCCTTAGGTCCCCCTCCTC
CATCAGCAAAGGAGCACTTCTCTAATCATGCCCTCCCGAAGACTGGCTGGGAGAAGGTTTAAAA
ACAAAAAATCCAGGAGTAAGAGCCTTAGGTGAGTTTGAATTTGGAGACAACTGTCTGGCAA
AGGGTGCGAGAGGGAGCTTGTGCTCAGGAGTCCAGCCGCCAGCCTCGGGGTGTAGGTTTCT
GAGGTGTGCCATTGGGGCCTCAGCCTTCTCTGGTGACAGAGGCTCAGCTGTGGCCACCAACA
CACAACCACACACACACACAAACCACACACACAAATGGGGGCAACCACATCCAGTACAAGCTTTT
ACAAATGTTATTAGTGTCTTTTTTTTATTCTAATGCCTTGTCCTCTTAAAGTTATTTTATT
TGTTATTATTATTGTTCTTGACTGTTAATTGTGAATGGTAATGCAATAAAGTGCCTTTGTT
AGATGGTGAA
AA

FIGURE 110

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA142930
><subunit 1 of 1, 512 aa, 1 stop
><MW: 54535, pI: 4.89, NX(S/T): 7
MKAIHLTLLALLSVNTATNQNSADAVTTTETATSGPTVAAADTTETNFPETASTTANT
PSFPTATSPAPPIISTHSSSTIPTPAPPIISTHSSSTIPIPTAADSESTTNVNSLATSDI
ITASSPNDGLITMVPSETQSNNEMSPTTEDNQSSGPPTGTALLETSTLNSTGPSNPCQDD
PCADNSLCVKLHNTSFCLCLEGYYYNSSTCKKGKVFPGKISVTVSETFDPEEKHSMAYQD
LHSEITSLFKDVFGTSVYGQTVILTVSTLSRSEMRADDKFVNVTIVTILAETTS DNEK
TVTEKINKAIRSSSSNFLNYDLTLRCDYYGCNQ TADDCLNGLACDCKSDLQRPNPQSPFC
VASSLKCPDACNAQHKQCLIKKSGGAPECACVPGYQEDANGNCQKCAFGYSGLDCKDKFQ
LILTIVGTIAGIVILSMIILIVTARSNNKTKHIEENLIDEDFQNLKLRSTGFTNLGAE
GSVF PKVRITASRDSQM QNPYSSHSSMPRPDY
```

Important features of the protein:

Signal peptide:

Amino acids 1-17

Transmembrane domain:

Amino acids 421-442

N-glycosylation sites:

Amino acids 151-155;169-173;193-197;206-210;284-288;
332-336;449-453

N-myristoylation sites:

Amino acids 330-336;385-391;427-433;478-484

SEA domain:

Amino acids 212-328

FIGURE 111

CTGGGACTTGGCTTTCTCCGGATAAGCGGCGGCACCGGCGTCAGCGGATGACCGTGCAGAGAC
TCGTGGCCGCGGCCGTGCTGGTGGCCCTGGTCTCACTCATCCTCAACAACGTGGCGGCCTTC
ACCTCCAAC TGGGTGTGCCAGACGCTGGAGGATGGGCGCAGGCGCAGCGTGGGGCTGTGGAG
GTCCTGCTGGCTGGTGGACAGGACCCGGGGAGGGCCGAGCCCTGGGGCCAGAGCCGGCCAGG
TGGACGCACATGACTGTGAGGCGCTGGGCTGGGGCTCCGAGGCAGCCGGCTTCCAGGAGTCC
CGAGGCACCGTCAAAC TGCAGTTCGACATGATGCGCGCCTGCAACCTGGTGGCCACGGCCGC
GCTCACCGCAGGCCAGCTCACCTTCCTCCTGGGGCTGGTGGGCCTGCCCCCTGCTGTCACCCG
ACGCCCCGTGCTGGGAGGAGGCCATGGCCGCTGCATTCCAAC TGGCGAGTTTGTCTCTGGTC
ATCGGGCTCGTGACTTTCTACAGAATTGGCCCATAACCAACCTGTCTCTGGTCCTGTACCT
GAACATTGGCGCCTGCCTTCTGGCCACGCTGGCGGCAGCCATGCTCATCTGGAACATTCTCC
ACAAGAGGGAGGACTGCATGGCCCCCGGGTGATTGTCATCAGCCGCTCCCTGACAGCGCGC
TTTCGCCGTGGGCTGGACAATGACTACGTGGAGTCACCATGCTTGAGTCGCCCTTCTCAGCGC
TCCATCAACGCACACCTGCTATCGTGGAACAGCCTAGAAACCAAGGGACTCCACCACCAAGT
CACTTCCCCTGCTCGTGAGAGGCACGGGATGAGTCTGGGTGACCTCTGCGCCATGCGTGCG
AGACACGTGTGCGTTTACTGTTATGTCGGTCATATGTCTGTACGTGTCTGGGCCAACCTCG
TTCTGCCTCCAGC

FIGURE 112

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA147253
><subunit 1 of 1, 226 aa, 1 stop
><MW: 24540, pI: 8.27, NX(S/T): 1
MTVQRLVAAAVLVSLILNNVAAFTSNWVCQTLEDGRRRSVGLWRSCWLVDRTTRGGPS
PGARAGQVDAHDCEALGWGSEAAGFQESRGTVKLGQFDMMRACNLVATAALTAGQLTFLLG
LVGLPLLSPDAPCWEEAMAAAFQLASFVLVIGLVTFYRIGPYTNLSWSCYLNIGACLLAT
LAAAMLIWNILHKREDCMAPRVIVISRSLTARFRRGLDNDYVESPC
```

Important features of the protein:

Signal peptide:

Amino acids 1-25

Transmembrane domains:

Amino acids 105-125;139-157;169-188

N-glycosylation site:

Amino acids 164-168

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 39-43

Tyrosine kinase phosphorylation site:

Amino acids 214-222

N-myristoylation sites:

Amino acids 44-50;62-68;66-72;79-85

Amidation site:

Amino acids 37-41

FIGURE 113

GACTTTACCACTACTCGCTATAGAGCCCTGGTCAAGTTCTCTCCACCTCTCTATCTATGTCT
CAGTTTCTTCATCTGTAACATCAAATGAATAATAATACCAATCTCCTAGACTTCATAAGAGG
ATTAACAAAGACAAAATATGGGAAAAACATAACATGGCGTCCCATAATTATTAGATCTTATT
ATTGACACTAAAATGGCATTAAAAATTACCAAAAGGAAGACAGCATCTGTTTCCTCTTTGGTC
CTGAGCTGGTTAAAAGGAACACTGGTTGCCTGAACAGTCACACTTGCAACC**ATG**ATGCCTAA
ACATTGCTTTCTAGGCTTCCTCATCAGTTTCTTCCTTACTGGTGTAGCAGGAACTCAGTCAA
CGCATGAGTCTCTGAAGCCTCAGAGGGTACAATTCAGTCCCGAAATTTTCACAACATTTTG
CAATGGCAGCCTGGGAGGGCACTTACTGGCAACAGCAGTGTCTATTTTGTGCAGTACAAAAT
ATATGGACAGAGACAATGGAAAAATAAAGAAGACTGTTGGGGTACTCAAGAACTCTCTTGTG
ACCTTACCAGTGAAACCTCAGACATACAGGAACCTTATTACGGGAGGGTGAGGGCGGCCTCG
GCTGGGAGCTACTCAGAATGGAGCATGACGCCGCGGTTCACTCCCTGGTGGGAAACAAAAAT
AGATCCTCCAGTCATGAATATAACCCAAGTCAATGGCTCTTTGTTGGTAATTCTCCATGCTC
CAAATTTACCATATAGATACCAAAAGGAAAAAAATGTATCTATAGAAGATTACTATGAACTA
CTATACCGAGTTTTTTATAATTAACAATTCACTAGAAAAGGAGCAAAAGGTTTATGAAGGGGC
TCACAGAGCGGTTGAAATTGAAGCTCTAACACCACACTCCAGCTACTGTGTAGTGGCTGAAA
TATATCAGCCCATGTTAGACAGAAGAAGTCAGAGAAGTGAAGAGAGATGTGTGGAAATTCCA
TGACTTGTGGAATTTGGCATTTCAGCAATGTGGAAATTCTAAAGCTCCCTGAGAACAGGATGA
CTCGTGTGTTGAAGGATCTTATTTAAAATTGTTTTTGTATTTTCTTAAAGCAATATTCAGTGT
TACACCTTGGGGACTTCTTTGTTTACCCATTCTTTTATCCTTTATATTTCAATTTGTAACTA
TATTTGAACGACATTCCCCCGAAAAATTGAAATGTAAAGATGAGGCAGAGAATAAAGTGTT
CTATGAAATTCAGAACTTTATTTCTGAATGTAACATCCCTAATAACAACCTTCATTCTTCTA
ATACAGCAAAATAAAAATTTAACAACCAAGGAATAGTATTTAAGAAAATGTTGAAATAATTT
TTTTAAAATAGCATTACAGACTGAG

FIGURE 114

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA149927
><subunit 1 of 1, 231 aa, 1 stop
><MW: 26980, pI: 7.06, NX(S/T): 5
MMPKHCFLGFLISFFLTGVAGTQSTHESLKPQRVQFQSRNFHNLQWQPGRALTGNSSVY
FVQYKIYGQRQWKNKEDCWGTQELSCDLTSETSDIQEPYYGRVRAASAGSYSEWSMTPRF
TPWWETKIDPPVMNITQVNGSLLVILHAPNLPYRYQKEKNVSIEDYYELLYRVFIINNSL
EKEQKVYEGAHRAVEIEALTPHSSYCVVAEIQPMLDRRSQRSEERCVEIP
```

Important features of the protein:

Signal peptide:

Amino acids 1-21

N-glycosylation sites:

Amino acids 56-60;134-138;139-143;160-164;177-181

N-myristoylation sites:

Amino acids 18-24;21-27;189-195

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC ACIDS ENCODING THE SAME

FIELD OF THE INVENTION

[0001] The present invention relates generally to the identification and isolation of novel DNA and to the recombinant production of novel polypeptides.

BACKGROUND OF THE INVENTION

[0002] Extracellular proteins play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of action in the extracellular environment.

[0003] Secreted proteins have various industrial applications, including as pharmaceuticals, diagnostics, biosensors and bioreactors. Most protein drugs available at present, such as thrombolytic agents, interferons, interleukins, erythropoietins, colony stimulating factors, and various other cytokines, are secretory proteins. Their receptors, which are membrane proteins, also have potential as therapeutic or diagnostic agents. Efforts are being undertaken by both industry and academia to identify new, native secreted proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.* 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0004] Membrane-bound proteins and receptors can play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. Such membrane-bound proteins and cell receptors include, but are not limited to, cytokine receptors, receptor kinases, receptor phosphatases, receptors involved in cell-cell interactions, and cellular adhesion molecules like selectins and integrins. For instance, transduction of signals that regulate cell growth and differentiation is regulated in part by phosphorylation of various cellular proteins. Protein tyrosine kinases, enzymes that catalyze that process, can also act as growth factor receptors. Examples include fibroblast growth factor receptor and nerve growth factor receptor.

[0005] Membrane-bound proteins and receptor molecules have various industrial applications, including as pharma-

ceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interactions. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

[0006] Efforts are being undertaken by both industry and academia to identify new, native receptor or membrane-bound proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor or membrane-bound proteins.

SUMMARY OF THE INVENTION

[0007] In one embodiment, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

[0008] In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0009] In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0010] In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0011] Another aspect the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

[0012] Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide

probes. Such nucleic acid fragments are usually at least about 10 nucleotides in length, alternatively at least about 15 nucleotides in length, alternatively at least about 20 nucleotides in length, alternatively at least about 30 nucleotides in length, alternatively at least about 40 nucleotides in length, alternatively at least about 50 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 70 nucleotides in length, alternatively at least about 80 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 100 nucleotides in length, alternatively at least about 110 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 130 nucleotides in length, alternatively at least about 140 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 160 nucleotides in length, alternatively at least about 170 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 190 nucleotides in length, alternatively at least about 200 nucleotides in length, alternatively at least about 250 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 350 nucleotides in length, alternatively at least about 400 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 500 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 700 nucleotides in length, alternatively at least about 800 nucleotides in length, alternatively at least about 900 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

[0013] In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

[0014] In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93%

amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

[0015] In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

[0016] In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

[0017] Another aspect the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

[0018] In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as

defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

[0019] In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

[0020] In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

[0021] Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

[0022] In other embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

[0023] In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

[0024] In another embodiment, the invention provides an antibody which binds, preferably specifically, to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

[0025] In yet other embodiments, the invention provides oligonucleotide probes which may be useful for isolating genomic and cDNA nucleotide sequences, measuring or detecting expression of an associated gene or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences. Preferred probe lengths are described above.

[0026] In yet other embodiments, the present invention is directed to methods of using the PRO polypeptides of the present invention for a variety of uses based upon the functional biological assay data presented in the Examples below.

BRIEF DESCRIPTION OF THE DRAWINGS

[0027] FIG. 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO281 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA16422-1209".

[0028] FIG. 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in FIG. 1.

[0029] FIG. 3 shows a nucleotide sequence (SEQ ID NO:3) of a native sequence PRO1560 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA19902-1669".

[0030] FIG. 4 shows the amino acid sequence (SEQ ID NO:4) derived from the coding sequence of SEQ ID NO:3 shown in FIG. 3.

[0031] FIG. 5 shows a nucleotide sequence (SEQ ID NO:5) of a native sequence PRO189 cDNA, wherein SEQ ID NO:5 is a clone designated herein as "DNA21624-1391".

[0032] FIG. 6 shows the amino acid sequence (SEQ ID NO:6) derived from the coding sequence of SEQ ID NO:5 shown in FIG. 5.

[0033] FIG. 7 shows a nucleotide sequence (SEQ ID NO:7) of a native sequence PRO240 cDNA, wherein SEQ ID NO:7 is a clone designated herein as "DNA34387-1138".

[0034] FIG. 8 shows the amino acid sequence (SEQ ID NO:8) derived from the coding sequence of SEQ ID NO:7 shown in FIG. 7.

[0035] FIG. 9 shows a nucleotide sequence (SEQ ID NO:9) of a native sequence PRO256 cDNA, wherein SEQ ID NO:9 is a clone designated herein as "DNA35880-1160".

[0036] FIG. 10 shows the amino acid sequence (SEQ ID NO:10) derived from the coding sequence of SEQ ID NO:9 shown in FIG. 9.

[0037] FIG. 11 shows a nucleotide sequence (SEQ ID NO:11) of a native sequence PRO306 cDNA, wherein SEQ ID NO:11 is a clone designated herein as "DNA39984-1221".

[0038] FIG. 12 shows the amino acid sequence (SEQ ID NO:12) derived from the coding sequence of SEQ ID NO:11 shown in FIG. 11.

[0039] FIG. 13 shows a nucleotide sequence (SEQ ID NO:13) of a native sequence PRO540 cDNA, wherein SEQ ID NO:13 is a clone designated herein as "DNA44189-1322".

[0040] FIG. 14 shows the amino acid sequence (SEQ ID NO:14) derived from the coding sequence of SEQ ID NO:13 shown in FIG. 13.

[0041] FIG. 15 shows a nucleotide sequence (SEQ ID NO:15) of a native sequence PRO773 cDNA, wherein SEQ ID NO:15 is a clone designated herein as "DNA48303-2829".

[0042] FIG. 16 shows the amino acid sequence (SEQ ID NO:16) derived from the coding sequence of SEQ ID NO:15 shown in FIG. 15.

[0043] FIG. 17 shows a nucleotide sequence (SEQ ID NO:17) of a native sequence PRO698 cDNA, wherein SEQ ID NO:17 is a clone designated herein as "DNA48320-1433".

[0044] FIG. 18 shows the amino acid sequence (SEQ ID NO:18) derived from the coding sequence of SEQ ID NO:17 shown in FIG. 17.

[0045] FIG. 19 shows a nucleotide sequence (SEQ ID NO:19) of a native sequence PRO3567 cDNA, wherein SEQ ID NO:19 is a clone designated herein as "DNA56049-2543".

[0046] FIG. 20 shows the amino acid sequence (SEQ ID NO:20) derived from the coding sequence of SEQ ID NO:19 shown in FIG. 19.

[0047] FIG. 21 shows a nucleotide sequence (SEQ ID NO:21) of a native sequence PRO826 cDNA, wherein SEQ ID NO:21 is a clone designated herein as "DNA57694-1341".

[0048] FIG. 22 shows the amino acid sequence (SEQ ID NO:22) derived from the coding sequence of SEQ ID NO:21 shown in FIG. 21.

[0049] FIG. 23 shows a nucleotide sequence (SEQ ID NO:23) of a native sequence PRO 1002 cDNA, wherein SEQ ID NO:23 is a clone designated herein as "DNA59208-1373".

[0050] FIG. 24 shows the amino acid sequence (SEQ ID NO:24) derived from the coding sequence of SEQ ID NO:23 shown in FIG. 23.

[0051] FIG. 25 shows a nucleotide sequence (SEQ ID NO:25) of a native sequence PRO 1068 cDNA, wherein SEQ ID NO:25 is a clone designated herein as "DNA59214-1449".

[0052] FIG. 26 shows the amino acid sequence (SEQ ID NO:26) derived from the coding sequence of SEQ ID NO:25 shown in FIG. 25.

[0053] FIG. 27 shows a nucleotide sequence (SEQ ID NO:27) of a native sequence PRO 1030 cDNA, wherein SEQ ID NO:27 is a clone designated herein as "DNA59485-1336".

[0054] FIG. 28 shows the amino acid sequence (SEQ ID NO:28) derived from the coding sequence of SEQ ID NO:27 shown in FIG. 27.

[0055] FIG. 29 shows a nucleotide sequence (SEQ ID NO:29) of a native sequence PRO1313 cDNA, wherein SEQ ID NO:29 is a clone designated herein as "DNA64966-1575".

[0056] FIG. 30 shows the amino acid sequence (SEQ ID NO:30) derived from the coding sequence of SEQ ID NO:29 shown in FIG. 29.

[0057] FIG. 31 shows a nucleotide sequence (SEQ ID NO:31) of a native sequence PRO6071 cDNA, wherein SEQ ID NO:31 is a clone designated herein as "DNA82403-2959".

[0058] FIG. 32 shows the amino acid sequence (SEQ ID NO:32) derived from the coding sequence of SEQ ID NO:31 shown in FIG. 31.

[0059] FIG. 33 shows a nucleotide sequence (SEQ ID NO:33) of a native sequence PRO4397 cDNA, wherein SEQ ID NO:33 is a clone designated herein as "DNA83505-2606".

[0060] FIG. 34 shows the amino acid sequence (SEQ ID NO:34) derived from the coding sequence of SEQ ID NO:33 shown in FIG. 33.

[0061] FIG. 35 shows a nucleotide sequence (SEQ ID NO:35) of a native sequence PRO4344 cDNA, wherein SEQ ID NO:35 is a clone designated herein as "DNA84927-2585".

[0062] FIG. 36 shows the amino acid sequence (SEQ ID NO:36) derived from the coding sequence of SEQ ID NO:35 shown in FIG. 35.

[0063] FIG. 37 shows a nucleotide sequence (SEQ ID NO:37) of a native sequence PRO4407 cDNA, wherein SEQ ID NO:37 is a clone designated herein as "DNA92264-2616".

[0064] FIG. 38 shows the amino acid sequence (SEQ ID NO:38) derived from the coding sequence of SEQ ID NO:37 shown in FIG. 37.

[0065] FIG. 39 shows a nucleotide sequence (SEQ ID NO:39) of a native sequence PRO4316 cDNA, wherein SEQ ID NO:39 is a clone designated herein as "DNA94713-2561".

[0066] FIG. 40 shows the amino acid sequence (SEQ ID NO:40) derived from the coding sequence of SEQ ID NO:39 shown in FIG. 39.

[0067] FIG. 41 shows a nucleotide sequence (SEQ ID NO:41) of a native sequence PRO5775 cDNA, wherein SEQ ID NO:41 is a clone designated herein as "DNA96869-2673".

[0068] FIG. 42 shows the amino acid sequence (SEQ ID NO:42) derived from the coding sequence of SEQ ID NO:41 shown in FIG. 41.

[0069] FIG. 43 shows a nucleotide sequence (SEQ ID NO:43) of a native sequence PRO6016 cDNA, wherein SEQ ID NO:43 is a clone designated herein as "DNA96881-2699".

[0070] FIG. 44 shows the amino acid sequence (SEQ ID NO:44) derived from the coding sequence of SEQ ID NO:43 shown in FIG. 43.

[0071] FIG. 45 shows a nucleotide sequence (SEQ ID NO:45) of a native sequence PRO4499 cDNA, wherein SEQ ID NO:45 is a clone designated herein as "DNA96889-2641".

[0072] FIG. 46 shows the amino acid sequence (SEQ ID NO:46) derived from the coding sequence of SEQ ID NO:45 shown in FIG. 45.

[0073] FIG. 47 shows a nucleotide sequence (SEQ ID NO:47) of a native sequence PRO4487 cDNA, wherein SEQ ID NO:47 is a clone designated herein as "DNA96898-2640".

[0074] FIG. 48 shows the amino acid sequence (SEQ ID NO:48) derived from the coding sequence of SEQ ID NO:47 shown in FIG. 47.

[0075] FIG. 49 shows a nucleotide sequence (SEQ ID NO:49) of a native sequence PRO4980 cDNA, wherein SEQ ID NO:49 is a clone designated herein as "DNA97003-2649".

[0076] FIG. 50 shows the amino acid sequence (SEQ ID NO:50) derived from the coding sequence of SEQ ID NO:49 shown in FIG. 49.

[0077] FIG. 51 shows a nucleotide sequence (SEQ ID NO:51) of a native sequence PRO6018 cDNA, wherein SEQ ID NO:51 is a clone designated herein as "DNA98565-2701".

[0078] FIG. 52 shows the amino acid sequence (SEQ ID NO:52) derived from the coding sequence of SEQ ID NO:51 shown in FIG. 51.

[0079] FIG. 53 shows a nucleotide sequence (SEQ ID NO:53) of a native sequence PRO7168 cDNA, wherein SEQ ID NO:53 is a clone designated herein as "DNA102846-2742".

[0080] FIG. 54 shows the amino acid sequence (SEQ ID NO:54) derived from the coding sequence of SEQ ID NO:53 shown in FIG. 53.

[0081] FIG. 55 shows a nucleotide sequence (SEQ ID NO:55) of a native sequence PRO6308 cDNA, wherein SEQ ID NO:55 is a clone designated herein as "DNA102847-2726".

[0082] FIG. 56 shows the amino acid sequence (SEQ ID NO:56) derived from the coding sequence of SEQ ID NO:55 shown in FIG. 55.

[0083] FIG. 57 shows a nucleotide sequence (SEQ ID NO:57) of a native sequence PRO6000 cDNA, wherein SEQ ID NO:57 is a clone designated herein as "DNA102880-2689".

[0084] FIG. 58 shows the amino acid sequence (SEQ ID NO:58) derived from the coding sequence of SEQ ID NO:57 shown in FIG. 57.

[0085] FIG. 59 shows a nucleotide sequence (SEQ ID NO:59) of a native sequence PRO6006 cDNA, wherein SEQ ID NO:59 is a clone designated herein as "DNA 105782-2693".

[0086] FIG. 60 shows the amino acid sequence (SEQ ID NO:60) derived from the coding sequence of SEQ ID NO:59 shown in FIG. 59.

[0087] FIG. 61 shows a nucleotide sequence (SEQ ID NO:61) of a native sequence PRO5800 cDNA, wherein SEQ ID NO:61 is a clone designated herein as "DNA108912-2680".

[0088] FIG. 62 shows the amino acid sequence (SEQ ID NO:62) derived from the coding sequence of SEQ ID NO:61 shown in FIG. 61.

[0089] FIG. 63 shows a nucleotide sequence (SEQ ID NO:63) of a native sequence PRO7476 cDNA, wherein SEQ ID NO:63 is a clone designated herein as "DNA1 15253-2757".

[0090] FIG. 64 shows the amino acid sequence (SEQ ID NO:64) derived from the coding sequence of SEQ ID NO:63 shown in FIG. 63.

[0091] FIG. 65 shows a nucleotide sequence (SEQ ID NO:65) of a native sequence PRO6496 cDNA, wherein SEQ ID NO:65 is a clone designated herein as "DNA1 19302-2737".

[0092] FIG. 66 shows the amino acid sequence (SEQ ID NO:66) derived from the coding sequence of SEQ ID NO:65 shown in FIG. 65.

[0093] FIG. 67 shows a nucleotide sequence (SEQ ID NO:67) of a native sequence PRO7422 cDNA, wherein SEQ ID NO:67 is a clone designated herein as "DNA 119536-2752".

[0094] FIG. 68 shows the amino acid sequence (SEQ ID NO:68) derived from the coding sequence of SEQ ID NO:67 shown in FIG. 67.

[0095] FIG. 69 shows a nucleotide sequence (SEQ ID NO:69) of a native sequence PRO7431cDNA, wherein SEQ ID NO:69 is a clone designated herein as "DNA1 19542-2754".

[0096] FIG. 70 shows the amino acid sequence (SEQ ID NO:70) derived from the coding sequence of SEQ ID NO:69 shown in FIG. 69.

[0097] FIG. 71 shows a nucleotide sequence (SEQ ID NO:71) of a native sequence PRO10275 cDNA, wherein SEQ ID NO:71 is a clone designated herein as "DNA143498-2824".

[0098] FIG. 72 shows the amino acid sequence (SEQ ID NO:72) derived from the coding sequence of SEQ ID NO:71 shown in FIG. 71.

[0099] FIG. 73 shows a nucleotide sequence (SEQ ID NO:73) of a native sequence PRO10268 cDNA, wherein SEQ ID NO:73 is a clone designated herein as "DNA 145583-2820".

[0100] FIG. 74 shows the amino acid sequence (SEQ ID NO:74) derived from the coding sequence of SEQ ID NO:73 shown in FIG. 73.

[0101] FIG. 75 shows a nucleotide sequence (SEQ ID NO:75) of a native sequence PRO20080 cDNA, wherein SEQ ID NO:75 is a clone designated herein as "DNA161000-2896".

[0102] FIG. 76 shows the amino acid sequence (SEQ ID NO:76) derived from the coding sequence of SEQ ID NO:75 shown in FIG. 75.

[0103] FIG. 77 shows a nucleotide sequence (SEQ ID NO:77) of a native sequence PRO21207 cDNA, wherein SEQ ID NO:77 is a clone designated herein as "DNA161005-2943".

[0104] FIG. 78 shows the amino acid sequence (SEQ ID NO:78) derived from the coding sequence of SEQ ID NO:77 shown in FIG. 77.

[0105] FIG. 79 shows a nucleotide sequence (SEQ ID NO:79) of a native sequence PRO28633 cDNA, wherein SEQ ID NO:79 is a clone designated herein as "DNA170245-3053".

[0106] FIG. 80 shows the amino acid sequence (SEQ ID NO:80) derived from the coding sequence of SEQ ID NO:79 shown in FIG. 79.

[0107] FIG. 81 shows a nucleotide sequence (SEQ ID NO:81) of a native sequence PRO20933 cDNA, wherein SEQ ID NO:81 is a clone designated herein as "DNA171771-2919".

[0108] FIG. 82 shows the amino acid sequence (SEQ ID NO:82) derived from the coding sequence of SEQ ID NO:81 shown in FIG. 81.

[0109] FIG. 83 shows a nucleotide sequence (SEQ ID NO:83) of a native sequence PRO21383 cDNA, wherein SEQ ID NO:83 is a clone designated herein as "DNA173157-2981".

[0110] FIG. 84 shows the amino acid sequence (SEQ ID NO:84) derived from the coding sequence of SEQ ID NO:83 shown in FIG. 83.

[0111] FIG. 85 shows a nucleotide sequence (SEQ ID NO:85) of a native sequence PRO21485 cDNA, wherein SEQ ID NO:85 is a clone designated herein as "DNA175734-2985".

[0112] FIG. 86 shows the amino acid sequence (SEQ ID NO:86) derived from the coding sequence of SEQ ID NO:85 shown in FIG. 85.

[0113] FIG. 87 shows a nucleotide sequence (SEQ ID NO:87) of a native sequence PRO28700 cDNA, wherein SEQ ID NO:87 is a clone designated herein as "DNA176108-3040".

[0114] FIG. 88 shows the amino acid sequence (SEQ ID NO:88) derived from the coding sequence of SEQ ID NO:87 shown in FIG. 87.

[0115] FIG. 89 shows a nucleotide sequence (SEQ ID NO:89) of a native sequence PRO34012 cDNA, wherein SEQ ID NO:89 is a clone designated herein as "DNA190710-3028".

[0116] FIG. 90 shows the amino acid sequence (SEQ ID NO:90) derived from the coding sequence of SEQ ID NO:89 shown in FIG. 89.

[0117] FIG. 91 shows a nucleotide sequence (SEQ ID NO:91) of a native sequence PRO34003 cDNA, wherein SEQ ID NO:91 is a clone designated herein as "DNA190803-3019".

[0118] FIG. 92 shows the amino acid sequence (SEQ ID NO:92) derived from the coding sequence of SEQ ID NO:91 shown in FIG. 91.

[0119] FIG. 93 shows a nucleotide sequence (SEQ ID NO:93) of a native sequence PRO34274 cDNA, wherein SEQ ID NO:93 is a clone designated herein as "DNA191064-3069".

[0120] FIG. 94 shows the amino acid sequence (SEQ ID NO:94) derived from the coding sequence of SEQ ID NO:93 shown in FIG. 93.

[0121] FIGS. 95A-95B shows a nucleotide sequence (SEQ ID NO:95) of a native sequence PRO34001 cDNA, wherein SEQ ID NO:95 is a clone designated herein as "DNA194909-3013".

[0122] FIG. 96 shows the amino acid sequence (SEQ ID NO:96) derived from the coding sequence of SEQ ID NO:95 shown in FIGS. 95A-95B.

[0123] FIG. 97 shows a nucleotide sequence (SEQ ID NO:97) of a native sequence PRO34009 cDNA, wherein SEQ ID NO:97 is a clone designated herein as "DNA203532-3029".

[0124] FIG. 98 shows the amino acid sequence (SEQ ID NO:98) derived from the coding sequence of SEQ ID NO:97 shown in FIG. 97.

[0125] FIG. 99 shows a nucleotide sequence (SEQ ID NO:99) of a native sequence PRO34192 cDNA, wherein SEQ ID NO:99 is a clone designated herein as "DNA213858-3060".

[0126] FIG. 100 shows the amino acid sequence (SEQ ID NO:100) derived from the coding sequence of SEQ ID NO:99 shown in FIG. 99.

[0127] FIG. 101 shows a nucleotide sequence (SEQ ID NO:101) of a native sequence PRO34564 cDNA, wherein SEQ ID NO:101 is a clone designated herein as "DNA216676-3083".

[0128] FIG. 102 shows the amino acid sequence (SEQ ID NO:102) derived from the coding sequence of SEQ ID NO:101 shown in FIG. 101.

[0129] FIG. 103 shows a nucleotide sequence (SEQ ID NO:103) of a native sequence PRO35444 cDNA, wherein SEQ ID NO:103 is a clone designated herein as "DNA222653-3104".

[0130] FIG. 104 shows the amino acid sequence (SEQ ID NO:104) derived from the coding sequence of SEQ ID NO:103 shown in FIG. 103.

[0131] FIG. 105 shows a nucleotide sequence (SEQ ID NO:105) of a native sequence PRO5998 cDNA, wherein SEQ ID NO:105 is a clone designated herein as "DNA96897-2688".

[0132] FIG. 106 shows the amino acid sequence (SEQ ID NO:106) derived from the coding sequence of SEQ ID NO:105 shown in FIG. 105.

[0133] FIG. 107 shows a nucleotide sequence (SEQ ID NO:107) of a native sequence PRO 19651 cDNA, wherein SEQ ID NO:107 is a clone designated herein as "DNA 42917-3081".

[0134] FIG. 108 shows the amino acid sequence (SEQ ID NO:108) derived from the coding sequence of SEQ ID NO:107 shown in FIG. 107.

[0135] FIG. 109 shows a nucleotide sequence (SEQ ID NO:109) of a native sequence PRO20221 cDNA, wherein SEQ ID NO:109 is a clone designated herein as "DNA142930-2914".

[0136] FIG. 110 shows the amino acid sequence (SEQ ID NO:110) derived from the coding sequence of SEQ ID NO:109 shown in FIG. 109.

[0137] FIG. 111 shows a nucleotide sequence (SEQ ID NO:111) of a native sequence PRO21434 cDNA, wherein SEQ ID NO:111 is a clone designated herein as "DNA 147253-2983".

[0138] FIG. 112 shows the amino acid sequence (SEQ ID NO:112) derived from the coding sequence of SEQ ID NO:111 shown in FIG. 111.

[0139] FIG. 113 shows a nucleotide sequence (SEQ ID NO:113) of a native sequence PRO 19822 cDNA, wherein SEQ ID NO:113 is a clone designated herein as "DNA 149927-2887".

[0140] FIG. 114 shows the amino acid sequence (SEQ ID NO:114) derived from the coding sequence of SEQ ID NO:113 shown in FIG. 113.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0141] I. Definitions

[0142] The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (i.e., PRO/number) refers to specific

polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. The term "PRO polypeptide" refers to each individual PRO/number polypeptide disclosed herein. All disclosures in this specification which refer to the "PRO polypeptide" refer to each of the polypeptides individually as well as jointly. For example, descriptions of the preparation of, purification of, derivation of, formation of antibodies to or against, administration of, compositions containing, treatment of a disease with, etc., pertain to each polypeptide of the invention individually. The term "PRO polypeptide" also includes variants of the PRO/number polypeptides disclosed herein.

[0143] A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

[0144] The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

[0145] The approximate location of the “signal peptides” of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen et al., *Prot. Eng.* 10:1-6 (1997) and von Heinje et al., *Nucl. Acids. Res.* 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

[0146] “PRO polypeptide variant” means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10 amino acids in length, alternatively at least about 20 amino acids in length, alternatively at least about 30 amino acids in length,

alternatively at least about 40 amino acids in length, alternatively at least about 50 amino acids in length, alternatively at least about 60 amino acids in length, alternatively at least about 70 amino acids in length, alternatively at least about 80 amino acids in length, alternatively at least about 90 amino acids in length, alternatively at least about 100 amino acids in length, alternatively at least about 150 amino acids in length, alternatively at least about 200 amino acids in length, alternatively at least about 300 amino acids in length, or more.

[0147] “Percent (%) amino acid sequence identity” with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific PRO polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

[0148] In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

[0149] where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations using this method, Tables 2 and 3 demonstrate how to calculate the % amino acid sequence identity of the amino

acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO", wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, and "X", "Y" and "Z" each represent different hypothetical amino acid residues.

[0150] Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % amino acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acid residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (i.e., the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an the amino acid sequence A which has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

[0151] Percent amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, Md. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

[0152] In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

[0153] where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid

residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

[0154] "PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleotide acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant polynucleotide will have at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

[0155] Ordinarily, PRO variant polynucleotides are at least about 30 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 210 nucleotides in length, alternatively at least about 240 nucleotides in length, alternatively at least about 270 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 900 nucleotides in length, or more.

[0156] "Percent (%) nucleic acid sequence identity" with respect to PRO-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a

candidate sequence that are identical with the nucleotides in the PRO nucleic acid sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. For purposes herein, however, % nucleic acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V40.D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

[0157] In situations where ALIGN-2 is employed for nucleic acid sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

100 times the fraction W/Z

[0158] where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4 and 5, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA", wherein "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, and "N", "L" and "V" each represent different hypothetical nucleotides.

[0159] Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % nucleic acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold

(T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % nucleic acid sequence identity value is determined by dividing (a) the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (i.e., the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement "an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B", the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

[0160] Percent nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, Md. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

[0161] In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

100 times the fraction W/Z

[0162] where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

[0163] In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding a full-length PRO polypeptide as disclosed herein. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

[0164] "Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components

of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide in situ within recombinant cells, since at least one component of the PRO polypeptide natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

[0165] An “isolated” PRO polypeptide-encoding nucleic acid or other polypeptide-encoding nucleic acid is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide-encoding nucleic acid. An isolated polypeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated polypeptide-encoding nucleic acid molecules therefore are distinguished from the specific polypeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated polypeptide-encoding nucleic acid molecule includes polypeptide-encoding nucleic acid molecules contained in cells that ordinarily express the polypeptide where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

[0166] The term “control sequences” refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

[0167] Nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[0168] The term “antibody” is used in the broadest sense and specifically covers, for example, single anti-PRO monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies), anti-PRO antibody compositions with polypeptidic specificity, single chain anti-PRO antibodies, and fragments of anti-PRO antibodies (see below). The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the

population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

[0169] “Stringency” of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience Publishers, (1995).

[0170] “Stringent conditions” or “high stringency conditions”, as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50° C.; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42° C.; or (3) employ 50% formamide, 5×SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5×Denhardt’s solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C., with washes at 42° C. in 0.2×SSC (sodium chloride/sodium citrate) and 50% formamide at 55° C., followed by a high-stringency wash consisting of 0.1×SSC containing EDTA at 55° C.

[0171] “Moderately stringent conditions” may be identified as described by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37° C. in a solution comprising: 20% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5×Denhardt’s solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1×SSC at about 37–50° C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

[0172] The term “epitope tagged” when used herein refers to a chimeric polypeptide comprising a PRO polypeptide fused to a “tag polypeptide”. The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with

other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

[0173] As used herein, the term “immunoadhesin” designates antibody-like molecules which combine the binding specificity of a heterologous protein (an “adhesin”) with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e. is “heterologous”), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

[0174] “Active” or “activity” for the purposes herein refers to form(s) of a PRO polypeptide which retain a biological and/or an immunological activity of native or naturally-occurring PRO, wherein “biological” activity refers to a biological function (either inhibitory or stimulatory) caused by a native or naturally-occurring PRO other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO and an “immunological” activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO.

[0175] The term “antagonist” is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native PRO polypeptide disclosed herein. In a similar manner, the term “agonist” is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native PRO polypeptides, peptides, antisense oligonucleotides, small organic molecules, etc. Methods for identifying agonists or antagonists of a PRO polypeptide may comprise contacting a PRO polypeptide with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the PRO polypeptide.

[0176] “Treatment” refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

[0177] “Chronic” administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. “Intermittent” administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

[0178] “Mammal” for purposes of treatment refers to any animal classified as a mammal, including humans, domestic

and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

[0179] Administration “in combination with” one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

[0180] “Carriers” as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are non-toxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

[0181] “Antibody fragments” comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (Zapata et al., *Protein Eng.* 8(10): 1057-1062[1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

[0182] Papain digestion of antibodies produces two identical antigen-binding fragments, called “Fab” fragments, each with a single antigen-binding site, and a residual “Fc” fragment, a designation reflecting the ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

[0183] “Fv” is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0184] The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab'), antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0185] The “light chains” of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains.

[0186] Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

[0187] “Single-chain Fv” or “sFv” antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

[0188] The term “diabodies” refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) in the same polypeptide chain (V_H-V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

[0189] An “isolated” antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the

antibody in situ within recombinant cells since at least one component of the antibody’s natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

[0190] An antibody that “specifically binds to” or is “specific for” a particular polypeptide or an epitope on a particular polypeptide is one that binds to that particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

[0191] The word “label” when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a “labeled” antibody. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

[0192] By “solid phase” is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Pat. No. 4,275,149.

[0193] A “liposome” is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as a PRO polypeptide or antibody thereto) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

[0194] A “small molecule” is defined herein to have a molecular weight below about 500 Daltons.

[0195] An “effective amount” of a polypeptide disclosed herein or an agonist or antagonist thereof is an amount sufficient to carry out a specifically stated purpose. An “effective amount” may be determined empirically and in a routine manner, in relation to the stated purpose.

TABLE 1

/*	
*	
* C—C increased from 12 to 15	
* Z is average of EQ	
* B is average of ND	
* match with stop is _M; stop—stop = 0; J (joker) match = 0	
*/	
#define	_M -8 /* value of a match with a stop */
int	_day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */	
/* A */ {2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},	
/* B */ {0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, -2, -5, 0, -3, 1},	
/* C */ {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},	
/* D */ {0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},	
/* E */ {0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},	
/* F */ {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},	

TABLE 1-continued

```

/* G */ {1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ {-1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2}
/* M */ {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ {0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ {_M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, 0, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ {1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ {0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ {1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ {1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ {0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ {0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4},
};
*/
#include <stdio.h>
#include <ctype.h>
#define MAXJMP 16 /* max jumps in a diag */
#define MAXGAP 24 /* don't continue to penalize gaps larger than this */
#define JMPS 1024 /* max jmps in an path */
#define MX 4 /* save if there's at least MX-1 bases since last jmp */
#define DMAT 3 /* value of matching bases */
#define DMIS 0 /* penalty for mismatched bases */
#define DINS0 8 /* penalty for a gap */
#define DINS1 1 /* penalty per base */
#define PINS0 8 /* penalty for a gap */
#define PINS1 4 /* penalty per residue */
struct jmp {
    short n[MAXJMP]; /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
}; /* limits seq to 2 16 -1 */
struct diag {
    int score; /* score at last jmp */
    long offset; /* offset of prev block */
    short ijmp; /* current jmp index */
    struct jmp jp; /* list of jmps */
};
struct path {
    int spc; /* number of leading spaces */
    short n[JMPS]; /* size of jmp (gap) */
    int x[JMPS]; /* loc of jmp (last elem before gap) */
};
char *ofile; /* output file name */
char *names[2]; /* seq names: getseqs() */
char *prog; /* prog name for err msgs */
char *seqs[2]; /* seqs: getseqs() */
int dmax; /* best diag: nw() */
int dmax0; /* final diag */
int dna; /* set if dna: main() */
int endgaps; /* set if penalizing end gaps */
int gapx, gapy; /* total gaps in seqs */
int len0, len1; /* seq lens */
int ngapx, ngapy; /* total size of gaps */
int smax; /* max score: nw() */
int *xbm; /* bitmap for matching */
long offset; /* current offset in jmp file */
struct diag *dx; /* holds diagonals */
struct path pp[2]; /* holds path for seqs */
char *calloc(), *malloc(), *index(), *strcpy();
char *getseq(), *g_calloc();
/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)

```

TABLE 1-continued

```

* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"
static __dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};
static __pbval[26] = {
    1,2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};
main(ac, av)
int    ac;
char   *av[];
{
    prog = av[0];
    if(ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case.\n");
        fprintf(stderr, "Any lines beginning with ';' or '<' are ignored.\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? __dbval : __pbval;
    endgaps = 0; /* 1 to penalize endgaps */
    ofile = "align.out"; /* output file */
    nw(); /* fill in the matrix, get the possible jmps */
    readjmps(); /* get the actual jmps */
    print(); /* print stats, alignment */
    cleanup(0); /* unlink any tmp files */
}
/* do the alignment, return best score: main()
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
* When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/
nw()
{
    char    *px, *py; /* seqs and ptrs */
    int     *ndely, *dely; /* keep track of dely */
    int     ndelx, delx; /* keep track of delx */
    int     *tmp; /* for swapping row0, row1 */
    int     mis; /* score for each type */
    int     ins0, ins1; /* insertion penalties */
    register id; /* diagonal index */
    register ij; /* jmp index */
    register *col0, *col1; /* score for curr, last row */
    register xx, yy; /* index into seqs */
    dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));
    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;
    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0; /* Waterman Bull Math Biol 84 */
    }
}

```

TABLE 1-continued

```

else
    for (yy = 1; yy <= len1; yy++)
        dely[yy] = -ins0;
/* fill in match matrix
*/
for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
    /* initialize first entry in col
    */
    if (endgaps) {
        if (xx == 1)
            col1[0] = delx = -(ins0+ins1);
        else
            col1[0] = delx = col0[0]-ins1;
        ndelx = xx;
    }
    else {
        col1[0] = 0;
        delx = -ins0;
        ndelx = 0;
    }

    for (py = seqy[1], yy = 1; yy <= len1; py++, yy++) {
        mis = col0[yy-1];
        if (dna)
            mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
        else
            mis += _day[*px-'A'][*py-'A'];
        /* update penalty for del in x seq;
        * favor new del over ongoing del
        * ignore MAXGAP if weighting endgaps
        */
        if (endgaps || ndely[yy] < MAXGAP) {
            if (col0[yy] - ins0 >= dely[yy]) {
                dely[yy] = col0[yy] - (ins0+ins1);
                ndely[yy] = 1;
            } else {
                dely[yy] -= ins1;
                ndely[yy]++;
            }
        } else {
            if (col0[yy] - (ins0+ins1) >= dely[yy]) {
                dely[yy] = col0[yy] - (ins0+ins1);
                ndely[yy] = 1;
            } else
                ndely[yy]++;
        }
        /* update penalty for del in y seq;
        * favor new del over ongoing del
        */
        if (endgaps || ndelx < MAXGAP) {
            if (col1[yy-1] - ins0 >= delx) {
                delx = col1[yy-1] - (ins0+ins1);
                ndelx = 1;
            } else {
                delx -= ins1;
                ndelx++;
            }
        } else {
            if (col1[yy-1] - (ins0+ins1) >= delx) {
                delx = col1[yy-1] - (ins0+ins1);
                ndelx = 1;
            } else
                ndelx++;
        }
        /* pick the maximum score; we're favoring
        * mis over any del and delx over dely
        */
        id = xx - yy + len1 - 1;
        if (mis >= delx && mis >= dely[yy])
            col1[yy] = mis;
        else if (delx >= dely[yy]) {
            col1[yy] = delx;
            ij = dx[id].ijmp;
            if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
            && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {

```

...nw

...nw

TABLE 1-continued

```

        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writeimps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
    col1[yy] = dely[yy];
    ij = dx[id].ijmp;
if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writeimps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndely[yy];
    dx[id].jp.x[ij] = xx;
    dx[id].score = dely[yy];
}
if (xx == len0 && yy < len1) {
    /* last col
    */
    if (endgaps)
        col1[yy] -= ins0+ins1*(len1-yy);
    if (col1[yy] > smax) {
        smax = col1[yy];
        dmax = id;
    }
}
}
if (endgaps && xx < len0)
    col1[yy-1] -= ins0+ins1*(len0-xx);
if (col1[yy-1] > smax) {
    smax = col1[yy-1];
    dmax = id;
}
tmp = col0; col0 = col1; col1 = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)col1);
}

/*
*
* print() -- only routine visible outside this module
*
* static:
* getmat() -- trace back best path, count matches: print()
* pr_align() -- print alignment of described in array p[]: print()
* dumpblock() -- dump a block of lines with numbers, stars: pr_align()
* nums() -- put out a number line: dumpblock()
* putline() -- put out a line (name, [num], seq, [num]): dumpblock()
* stars() -- put a line of stars: dumpblock()
* stripname() -- strip any path and prefix from a seqname
*/
#include "nw.h"
#define SPC          3
#define P__LINE      256 /* maximum output line */
#define P__SPC        3 /* space between name or num and seq */
extern      _day[26][26];
int         olen;        /* set output line length */
FILE        *fx;         /* output file */
print()
{
    int         lx, ly, firstgap, lastgap; /* overlap */

```

print

TABLE 1-continued

```

    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) {          /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) {      /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) {          /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) {     /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_alignn();
}
/*
 * trace back the best path, count matches
 */
static
getmat(lx, ly, firstgap, lastgap)                                     getmat
    int      lx, ly;          /* "core" (minus endgaps) */
    int      firstgap, lastgap; /* leading trailing overlap */
{
    int      nm, i0, i1, siz0, siz1;
    char      outx[32];
    double    pct;
    register  n0, n1;
    register char *p0, *p1;
    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;
    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
        else {
            if (xbm[*p0-'A']&xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }
    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */

```

TABLE 1-continued

```

if (endgaps)
    lx = (len0 < len1)? len0 : len1;
else
    lx = (lx < ly)? lx : ly;
pct = 100.*(double)nm/(double)lx;
fprintf(fx, "\n");
fprintf(fx, "<%d match%s in an overlap of %d: %.2f percent similarity\n",
    nm, (nm == 1)? "" : "es", lx, pct);
fprintf(fx, "<gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outx, "(%d %s%s)",
        ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);
}
fprintf(fx, ", gaps in second sequence: %d", gapy);
if (gapy) {
    (void) sprintf(outx, "(%d %s%s)",
        ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
    fprintf(fx, "%s", outx);
}
}
if (dna)
    fprintf(fx,
        "\n<score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
        smax, DMAT, DMIS, DINS0, DINS1);
else
    fprintf(fx,
        "\n<score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
        smax, PINS0, PINS1);
if (endgaps)
    fprintf(fx,
        "<endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
        firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
        lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
else
    fprintf(fx, "<endgaps not penalized\n");
}
static      nm;           /* matches in core -- for checking */
static      lmax;         /* lengths of stripped file names */
static      ij[2];        /* jmp index for a path */
static      nc[2];        /* number at start of current line */
static      ni[2];        /* current elem number -- for gapping */
static      siz[2];
static char  *ps[2];       /* ptr to current element */
static char  *po[2];       /* ptr to next output char slot */
static char  out[2][P_LINE]; /* output line */
static char  star[P_LINE]; /* set by stars() */
/*
 * print alignment of described in struct path pp[]
 */
static
pr_align()
{
    int      nn;           /* char count */
    int      more;
    register i;
    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(namex[i]);
        if (nn > lmax)
            lmax = nn;
        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
    for (nn = nm = 0, more = 1; more;) {
        for (i = more = 0; i < 2; i++) {
            /*
             * do we have more of this sequence?
             */
            if (!*ps[i])
                continue;
            more++;
            if (pp[i].spc) { /* leading space */
                *po[i]++ = ' ';
                pp[i].spc--;
            }
        }
    }
}

```

...getmat

pr_align

...pr_align

TABLE 1-continued

<pre> else if (siz[i]) { /* in a gap */ *po[i]++ = '-'; siz[i]--; } else { /* we're putting a seq element */ *po[i] = *ps[i]; if (islower(*ps[i])) *ps[i] = toupper(*ps[i]); po[i]++; ps[i]++; /* * are we at next gap for this seq? */ if (ni[i] == pp[i].x[ij[i]]) { /* * we need to merge all gaps * at this location */ siz[i] == pp[i].n[ij[i]++]; while (ni[i] == pp[i].x[ij[i]]) siz[i] += pp[i].n[ij[i]++]; } ni[i]++; } } if (++nn == olen !more && nn) { dumpblock(); for (i = 0; i < 2; i++) po[i] = out[i]; nn = 0; } } } /* * dump a block of lines, including numbers, stars: pr_align() */ static dumpblock() { register i; for(i = 0; i < 2; i++) *po[i]-- = '\0'; (void) puts("\n", fx); for (i = 0; i < 2; i++) { if (*out[i] && (*out[i] != ' ' *(po[i]) != ' ')) { if (i == 0) nums(i); if (i == 0 && *out[1]) stars(); putline(i); if (i == 0 && *out[1]) fprintf(fx, star); if (i == 1) nums(i); } } } /* * put out a number line: dumpblock() */ static nums(ix) int ix; /* index in out[] holding seq line */ { char nline[P_LINE]; register i, j; register char *pn, *px, *py; for(pn = nline, i = 0; i < lmax+P_SPC; i++, pn++) *pn = ' '; for (i = nc[ix], py = out[ix]; *py; py++, pn++) { if (*py == ' ' *py == '-') *pn = ' '; else { if (i%10 == 0 (i == 1 && nc[ix] != 1)) {</pre>		
		dumpblock
		...dumpblock
		nums

TABLE 1-continued

<pre>j = (i < 0)? -i : i; for (px = pn; j; j/= 10, px--) *px = j%10 + '0'; if (i < 0) *px = '-'; } else *pn = ' '; i++; } } *pn = '\0'; nc[ix] = i; for (pn = nline; *pn; pn++) (void) putc(*pn, fx); (void) putc('\n', fx); } /* * put out a line (name, [num], seq. [num]): dumpblock() */ static putline(ix) int ix; { int i; register char *px; for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++) (void) putc(*px, fx); for (; i < lmax+P__SPC; i++) (void) putc(' ', fx); /* these count from 1: * ni[] is current element (from 1) * nc[] is number at start of current line */ for (px = out[ix]; *px; px++) (void) putc(*px&0x7F, fx); (void) putc('\n', fx); } /* * put a line of stars (seqs always in out[0], out[1]): dumpblock() */ static stars() { int i; register char *p0, *p1, cx, *px; if (!(*out[0] (*out[0] == ' ' && *(p0[0]) == ' ') !(*out[1] (*out[1] == ' ' && *(p0[1]) == ' '))) return; px = star; for (i = lmax+P__SPC; i; i--) *px++ = ' '; for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) { if (isalpha(*p0) && isalpha(*p1)) { if (xbm[*p0-'A']&xbm[*p1-'A']) { cx = '*'; nm++; } else if (!dna && __day[*p0-'A'][*p1-'A'] > 0) cx = '.'; else cx = ' '; } else cx = ' '; *px++ = cx; } *px++ = '\n'; *px = '\0'; } /* * strip path or prefix from pn, return len: pr__align() */ static stripname(pn) char *pn; /* file name (may be path) */</pre>			putline
			...putline
			stars
			stripname

TABLE 1-continued

<pre>{ register char *px, *py; py = 0; for (px = pn; *px; px++) if (*px == '/') py = px + 1; if (py) (void) strcpy(pn, py); return(strlen(pn)); } /* * cleanup() -- cleanup any tmp file * getseq() -- read in seq, set dna, len, maxlen * g_calloc() -- calloc() with error checkin * readjumps() -- get the good jumps, from tmp file if necessary * writejumps() -- write a filled array of jumps to a tmp file: nw() */ #include "nw.h" #include <sys/file.h> char *jname = "/tmp/homgXXXXXX"; /* tmp file for jumps */ FILE *fj; int cleanup(); /* cleanup tmp file */ long lseek(); /* * remove any tmp file if we blow */ cleanup(i) int i; { if (fj) (void) unlink(jname); exit(i); } /* * read, return ptr to seq, set dna, len, maxlen * skip lines starting with ';', '<', or '>' * seq in upper or lower case */ char * getseq(file, len) char *file; /* file name */ int *len; /* seq len */ { char line[1024], *pseq; register char *px, *py; int natgc, tlen; FILE *fp; if ((fp = fopen(file, "r")) == 0) { fprintf(stderr, "%s: can't read %s\n", prog, file); exit(1); } tlen = natgc = 0; while (fgets(line, 1024, fp)) { if (*line == ';' *line == '<' *line == '>') continue; for (px = line; *px != '\n'; px++) if (isupper(*px) islower(*px)) tlen++; } if ((pseq = malloc((unsigned)(tlen+6))) == 0) { fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file); exit(1); } pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0'; py = pseq + 4; *len = tlen; rewind(fp); while (fgets(line, 1024, fp)) { if (*line == ';' *line == '<' *line == '>') continue; for (px = line; *px != '\n'; px++) { if (isupper(*px)) *py++ = *px; else if (islower(*px)) *py++ = toupper(*px); } } }</pre>		
		cleanup
		getseq
		...getseq

TABLE 1-continued

```

        if (index("ATGCU", *(py-1)))
            natgc++;
    }
}
*py++ = '\0';
*py = '\0';
(void) fclose(fp);
dna = natgc > (tlen/3);
return(pseq+4);
}
char *
g__calloc(msg, nx, sz)                                g__calloc
char      *msg;          /* program, calling routine */
int       nx, sz;        /* number and size of elements */
{
    char      *px, *calloc();
    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g__calloc() failed %s (n= %d, sz= %d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}
/*
* get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
*/
readjmps()                                             readjmps
{
    int      fd = -1;
    int      siz, i0, i1;
    register i, j, xx;
    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;

            if (j < 0 && dx[dmax].offset && fj) {
                (void) lseek(fd, dx[dmax].offset, 0);
                (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
                (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
                dx[dmax].ijmp = MAXJMP-1;
            }
            else
                break;
        }
        if (i >= JMPS) {
            fprintf(stderr, "%s: too many gaps in alignment\n", prog);
            cleanup(1);
        }
        if (j >= 0) {
            siz = dx[dmax].jp.n[j];
            xx = dx[dmax].jp.x[j];
            dmax += siz;
            if (siz < 0) { /* gap in second seq */
                pp[1].n[i1] = -siz;
                xx += siz;
                /* id = xx - yy + len1 - 1
                */
            }
        }
    }
}

```

TABLE 1-continued

<pre> pp[1].x[i1] = xx - dmax + len1 - 1; gapy++; ngapy -= siz; /* ignore MAXGAP when doing endgaps */ siz = (-siz < MAXGAP endgaps)? -siz : MAXGAP; il++; } else if (siz > 0) { /* gap in first seq */ pp[0].n[i0] = siz; pp[0].x[i0] = xx; gapx++; ngapx += siz; /* ignore MAXGAP when doing endgaps */ siz = (siz < MAXGAP endgaps)? siz : MAXGAP; i0++; } } else break; } /* reverse the order of jmps */ for (j = 0, i0--; j < i0; j++, i0--) { i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i; i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i; } for (j = 0, i1--; j < i1; j++, i1--) { i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i; i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i; } if (fd >= 0) (void) close(fd); if (fj) { (void) unlink(jname); fj = 0; offset = 0; } } /* * write a filled jmp struct offset of the prev one (if any): nw() */ writejmps(ix) int ix; { char *mktemp(); if (!fj) { if (mktemp(jname) < 0) { fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname); cleanup(1); } if ((fj = fopen(jname, "w")) == 0) { fprintf(stderr, "%s: can't write %s\n", prog, jname); exit(1); } } (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj); (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj); }</pre>	writejmps
---	-----------

[0196]

TABLE 2

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison	XXXXXXXXYYYYYYY	(Length = 12 amino acids)
Protein		

% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 15 = 33.3%

[0197]

TABLE 3

PRO	XXXXXXXXXXXX	(Length = 10 amino acids)
Comparison	XXXXXXXXYYYYYYZZYZ	(Length = 15 amino acids)
Protein		

% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 10 = 50%

[0198]

TABLE 4

PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLL	(Length = 16 nucleotides)

% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 6 divided by 14 = 42.9%

[0199]

TABLE 5

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLLVV	(Length = 9 nucleotides)

% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 4 divided by 12 = 33.3%

[0200] II. Compositions and Methods of the Invention

[0201] A. Full-Length PRO Polypeptides

[0202] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO polypeptides. In particular, cDNAs encoding various PRO polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by the full length native nucleic acid molecules disclosed herein as well as all further native homologues and variants included in the foregoing definition of PRO, will be referred to as "PRO/number", regardless of their origin or mode of preparation.

[0203] As disclosed in the Examples below, various cDNA clones have been deposited with the ATCC. The actual nucleotide sequences of those clones can readily be determined by the skilled artisan by sequencing of the deposited clone using routine methods in the art. The predicted amino acid sequence can be determined from the nucleotide sequence using routine skill. For the PRO polypeptides and encoding nucleic acids described herein, Applicants have identified what is believed to be the reading frame best identifiable with the sequence information available at the time.

[0204] B. PRO Polypeptide Variants

[0205] In addition to the full-length native sequence PRO polypeptides described herein, it is contemplated that PRO variants can be prepared. PRO variants can be prepared by introducing appropriate nucleotide changes into the PRO DNA, and/or by synthesis of the desired PRO polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

[0206] Variations in the native full-length sequence PRO or in various domains of the PRO described herein, can be

made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Pat. No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO that results in a change in the amino acid sequence of the PRO as compared with the native sequence PRO. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

[0207] PRO polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the PRO polypeptide.

[0208] PRO fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating PRO fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, PRO polypeptide fragments share at least one biological and/or immunological activity with the native PRO polypeptide disclosed herein.

[0209] In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

TABLE 6

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln

TABLE 6-continued

Original Residue	Exemplary Substitutions	Preferred Substitutions
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

[0210] Substantial modifications in function or immunological identity of the PRO polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- [0211] (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- [0212] (2) neutral hydrophilic: cys, ser, thr;
- [0213] (3) acidic: asp, glu;
- [0214] (4) basic: asn, gin, his, lys, arg;
- [0215] (5) residues that influence chain orientation: gly, pro; and
- [0216] (6) aromatic: trp, tyr, phe.

[0217] Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

[0218] The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., *Nucl. Acids Res.*, 13:4331 (1986); Zoller et al., *Nucl. Acids Res.*, 10:6487 (1987)], cassette mutagenesis [Wells et al., *Gene*, 34:315 (1985)], restriction selection mutagenesis [Wells et al., *Philos. Trans. R. Soc. London SerA*, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO variant DNA.

[0219] Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine

is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, *Science*, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, *The Proteins*, (W.H. Freeman & Co., N.Y.); Chothia, *J. Mol. Biol.*, 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

[0220] C. Modifications of PRO

[0221] Covalent modifications of PRO are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues of the PRO. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis-(diazocetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

[0222] Other modifications include deamidation of glutamyl and asparagyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T. E. Creighton, *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

[0223] Another type of covalent modification of the PRO polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

[0224] Addition of glycosylation sites to the PRO polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO (for O-linked glycosylation sites). The PRO amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[0225] Another means of increasing the number of carbohydrate moieties on the PRO polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published Sep. 11, 1987, and in Aplin and Wriston, *CRC Crit. Rev. Biochem.*, pp. 259-306 (1981).

[0226] Removal of carbohydrate moieties present on the PRO polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., *Arch. Biochem. Biophys.*, 259:52 (1987) and by Edge et al., *Anal. Biochem.*, 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., *Meth. Enzymol.*, 138:350 (1987).

[0227] Another type of covalent modification of PRO comprises linking the PRO polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

[0228] The PRO of the present invention may also be modified in a way to form a chimeric molecule comprising PRO fused to another, heterologous polypeptide or amino acid sequence.

[0229] In one embodiment, such a chimeric molecule comprises a fusion of the PRO with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the PRO. The presence of such epitope-tagged forms of the PRO can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., *Mol. Cell. Biol.*, 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., *Molecular and Cellular Biology*, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., *Protein Engineering*, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., *BioTechnology*, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., *Science*, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner et al., *J. Biol. Chem.*, 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., *Proc. Natl. Acad. Sci. USA*, 87:6393-6397 (1990)].

[0230] In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO polypeptide in place of at least one variable region within an Ig molecule. In a particularly

preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG I molecule. For the production of immunoglobulin fusions see also U.S. Pat. No. 5,428,130 issued Jun. 27, 1995.

[0231] D. Preparation of PRO

[0232] The description below relates primarily to production of PRO by culturing cells transformed or transfected with a vector containing PRO nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare PRO. For instance, the PRO sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., *Solid-Phase Peptide Synthesis*, W.H. Freeman Co., San Francisco, Calif. (1969); Merrifield, *J. Am. Chem. Soc.*, 85:2149-2154 (1963)]. In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, Calif.) using manufacturer's instructions. Various portions of the PRO may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO.

[0233] 1. Isolation of DNA Encoding PRO

[0234] DNA encoding PRO may be obtained from a cDNA library prepared from tissue believed to possess the PRO mRNA and to express it at a detectable level. Accordingly, human PRO DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

[0235] Libraries can be screened with probes (such as antibodies to the PRO or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., *PCR Primer: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1995)].

[0236] The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ^{32}P -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

[0237] Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence

identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

[0238] Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

[0239] 2. Selection and Transformation of Host Cells

[0240] Host cells are transfected or transformed with expression or cloning vectors described herein for PRO production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in *Mammalian Cell Biotechnology: a Practical Approach*, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

[0241] Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example, CaCl_2 , CaPO_4 , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., *Gene*, 23:315 (1983) and WO 89/05859 published Jun. 29, 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, *Virology*, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Pat. No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., *J. Bact.*, 130:946 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. (USA)*, 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., *Methods in Enzymology*, 185:527-537 (1990) and Mansour et al., *Nature*, 336:348-352 (1988).

[0242] Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*,

e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published Apr. 12, 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype ton⁺; *E. coli* W3110 strain 9E4, which has the complete genotype tonA ptr3; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan^r; *E. coli* W3110 strain 37D6, which has the complete genotype tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan^r; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant degP deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Pat. No. 4,946,783 issued Aug. 7, 1990. Alternatively, in vitro methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

[0243] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, *Nature*, 290: 140[1981]; EP 139,383 published May 2, 1985); *Kluyveromyces* hosts (U.S. Pat. No. 4,943,529; Fleer et al., *Bio/Technology*, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., *J. Bacteriol.*, 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36,906; Van den Berg et al., *Bio/Technology*, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., *J. Basic Microbiol.*, 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., *Proc. Natl. Acad. Sci. USA*, 76:5259-5263[1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published Oct. 31, 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published Jan. 10, 1991), and *Aspergillus* hosts such as *A. nidulans* (Balance et al., *Biochem. Biophys. Res. Commun.*, 112:284-289 [1983]; Tilburn et al., *Gene*, 26:205-221[1983]; Yelton et al., *Proc. Natl. Acad. Sci. USA*, 81: 1470-1474[1984]) and *A. niger* (Kelly and Hynes, *EMBO J.*, 4:475-479[1985]). Methylotrophic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, *The Biochemistry of Methylotrophs*, 269 (1982).

[0244] Suitable host cells for the expression of glycosylated PRO are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include

Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen. Virol.*, 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

[0245] 3. Selection and Use of a Replicable Vector

[0246] The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

[0247] The PRO may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the PRO-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Pat. No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published Nov. 15, 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

[0248] Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the *2* μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

[0249] Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer

resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

[0250] An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., *Nature*, 282:39 (1979); Kingsman et al., *Gene*, 7:141 (1979); Tschemper et al., *Gene*, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, *Genetics*, 85:12 (1977)].

[0251] Expression and cloning vectors usually contain a promoter operably linked to the PRO-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., *Nature*, 275:615 (1978); Goeddel et al., *Nature*, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, *Nucleic Acids Res.*, 8:4057 (1980); EP 36,776], and hybrid promoters such as the tac promoter [deBoer et al., *Proc. Natl. Acad. Sci. USA*, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO.

[0252] Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., *J. Biol. Chem.*, 255:2073 (1980)] or other glycolytic enzymes [Hess et al., *J. Adv. Enzyme Reg.*, 7:149 (1968); Holland, *Biochemistry*, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

[0253] Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

[0254] PRO transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published Jul. 5, 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

[0255] Transcription of a DNA encoding the PRO by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the PRO coding sequence, but is preferably located at a site 5' from the promoter.

[0256] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO.

[0257] Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO in recombinant vertebrate cell culture are described in Gething et al., *Nature*, 293:620-625 (1981); Mantei et al., *Nature*, 281:40-46 (1979); EP 117,060; and EP 117,058.

[0258] 4. Detecting Gene Amplification/Expression

[0259] Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, *Proc. Natl. Acad. Sci. USA*, 77:5201-5205 (1980)], dot blotting (DNA analysis), or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

[0260] Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope.

[0261] 5. Purification of Polypeptide

[0262] Forms of PRO may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent

solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

[0263] It may be desired to purify PRO from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, *Methods in Enzymology*, 182 (1990); Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO produced.

[0264] E. Uses for PRO

[0265] Nucleotide sequences (or their complement) encoding PRO have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA. PRO nucleic acid will also be useful for the preparation of PRO polypeptides by the recombinant techniques described herein.

[0266] The full-length native sequence PRO gene, or portions thereof, may be used as hybridization probes for a cDNA library to isolate the full-length PRO cDNA or to isolate still other cDNAs (for instance, those encoding naturally-occurring variants of PRO or PRO from other species) which have a desired sequence identity to the native PRO sequence disclosed herein. Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from at least partially novel regions of the full length native nucleotide sequence wherein those regions may be determined without undue experimentation or from genomic sequences including promoters, enhancer elements and introns of native sequence PRO. By way of example, a screening method will comprise isolating the coding region of the PRO gene using the known DNA sequence to synthesize a selected probe of about 40 bases. Hybridization probes may be labeled by a variety of labels, including radionucleotides such as ^{32}P or ^{35}S , or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the PRO gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

[0267] Any EST sequences disclosed in the present application may similarly be employed as probes, using the methods disclosed herein.

[0268] Other useful fragments of the PRO nucleic acids include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA)

capable of binding to target PRO mRNA (sense) or PRO DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of PRO DNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (*Cancer Res.* 48:2659, 1988) and van der Krol et al. (*BioTechniques* 6:958, 1988).

[0269] Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of PRO proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable in vivo (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences.

[0270] Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10048, and other moieties that increases affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

[0271] Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO_4 -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. In a preferred procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either in vivo or ex vivo. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see WO 90/13641).

[0272] Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

[0273] Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target

nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

[0274] Antisense or sense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length, about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length, or more.

[0275] The probes may also be employed in PCR techniques to generate a pool of sequences for identification of closely related PRO coding sequences.

[0276] Nucleotide sequences encoding a PRO can also be used to construct hybridization probes for mapping the gene which encodes that PRO and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as in situ hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

[0277] When the coding sequences for PRO encode a protein which binds to another protein (example, where the PRO is a receptor), the PRO can be used in assays to identify the other proteins or molecules involved in the binding interaction. By such methods, inhibitors of the receptor/ligand binding interaction can be identified. Proteins involved in such binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction. Also, the receptor PRO can be used to isolate correlative ligand(s). Screening assays can be designed to find lead compounds that mimic the biological activity of a native PRO or a receptor for PRO. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

[0278] Nucleic acids which encode PRO or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding PRO. Methods for generating transgenic animals, particu-

larly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for PRO transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding PRO introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding PRO. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

[0279] Alternatively, non-human homologues of PRO can be used to construct a PRO “knock out” animal which has a defective or altered gene encoding PRO as a result of homologous recombination between the endogenous gene encoding PRO and altered genomic DNA encoding PRO introduced into an embryonic stem cell of the animal. For example, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques. A portion of the genomic DNA encoding PRO can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a “knock out” animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knock-out animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the PRO polypeptide.

[0280] Nucleic acid encoding the PRO polypeptides may also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve in vivo synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. “Gene therapy” includes both conventional gene therapy where a lasting effect is achieved by a single treatment, and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes in vivo. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concen-

trations caused by their restricted uptake by the cell membrane. (Zamecnik et al., *Proc. Natl. Acad. Sci. USA* 83:4143-4146[1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups.

[0281] There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells in vitro, or in vivo in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells in vitro include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. The currently preferred in vivo gene transfer techniques include transfection with viral (typically retroviral) vectors and viral coat protein-liposome mediated transfection (Dzau et al., *Trends in Biotechnology* 11, 205-210[1993]). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.* 262, 4429-4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA* 87, 3410-3414 (1990). For review of gene marking and gene therapy protocols see Anderson et al., *Science* 256, 808-813 (1992).

[0282] The PRO polypeptides described herein may also be employed as molecular weight markers for protein electrophoresis purposes and the isolated nucleic acid sequences may be used for recombinantly expressing those markers.

[0283] The nucleic acid molecules encoding the PRO polypeptides or fragments thereof described herein are useful for chromosome identification. In this regard, there exists an ongoing need to identify new chromosome markers, since relatively few chromosome marking reagents, based upon actual sequence data are presently available. Each PRO nucleic acid molecule of the present invention can be used as a chromosome marker.

[0284] The PRO polypeptides and nucleic acid molecules of the present invention may also be used diagnostically for tissue typing, wherein the PRO polypeptides of the present invention may be differentially expressed in one tissue as compared to another, preferably in a diseased tissue as compared to a normal tissue of the same tissue type. PRO nucleic acid molecules will find use for generating probes for PCR, Northern analysis, Southern analysis and Western analysis.

[0285] The PRO polypeptides described herein may also be employed as therapeutic agents. The PRO polypeptides of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby the PRO product hereof is combined in admixture with a pharmaceutically acceptable carrier vehicle. Therapeutic formulations are prepared for storage by mixing the active ingredient having the desired degree of purity with

optional physiologically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone, amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEENTM, PLURONICSTM or PEG.

[0286] The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution.

[0287] Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0288] The route of administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial or intralesional routes, topical administration, or by sustained release systems.

[0289] Dosages and desired drug concentrations of pharmaceutical compositions of the present invention may vary depending on the particular use envisioned. The determination of the appropriate dosage or route of administration is well within the skill of an ordinary physician. Animal experiments provide reliable guidance for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles laid down by Mordenti, J. and Chappell, W. "The use of interspecies scaling in toxicokinetics" In *Toxicokinetics and New Drug Development*, Yacobi et al., Eds., Pergamon Press, New York 1989, pp. 42-96.

[0290] When in vivo administration of a PRO polypeptide or agonist or antagonist thereof is employed, normal dosage amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1 μ g/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting one organ or tissue, for example, may necessitate delivery in a manner different from that to another organ or tissue.

[0291] Where sustained-release administration of a PRO polypeptide is desired in a formulation with release characteristics suitable for the treatment of any disease or disorder requiring administration of the PRO polypeptide, microencapsulation of the PRO polypeptide is contemplated. Microencapsulation of recombinant proteins for sustained

release has been successfully performed with human growth hormone (rhGH), interferon-(rhIFN-), interleukin-2, and MN rgp120. Johnson et al., *Nat. Med.*, 2:795-799 (1996); Yasuda, *Biomed. Ther.*, 27:1221-1223 (1993); Hora et al., *Bio/Technology*, 8:755-758 (1990); Cleland, "Design and Production of Single Immunization Vaccines Using Poly-lactide Polyglycolide Microsphere Systems," in *Vaccine Design: The Subunit and Adjuvant Approach*, Powell and Newman, eds. (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Pat. No. 5,654,010.

[0292] The sustained-release formulations of these proteins were developed using poly-lactic-coglycolic acid (PLGA) polymer due to its biocompatibility and wide range of biodegradable properties. The degradation products of PLGA, lactic and glycolic acids, can be cleared quickly within the human body. Moreover, the degradability of this polymer can be adjusted from months to years depending on its molecular weight and composition. Lewis, "Controlled release of bioactive agents from lactide/glycolide polymer," in: M. Chasin and R. Langer (Eds.), *Biodegradable Polymers as Drug Delivery Systems* (Marcel Dekker: New York, 1990), pp. 1-41.

[0293] This invention encompasses methods of screening compounds to identify those that mimic the PRO polypeptide (agonists) or prevent the effect of the PRO polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO polypeptides encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

[0294] The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

[0295] All assays for antagonists are common in that they call for contacting the drug candidate with a PRO polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

[0296] In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the PRO polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the PRO polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detect-

able label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

[0297] If the candidate compound interacts with but does not bind to a particular PRO polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, *Nature (London)*, 340:245-246 (1989); Chien et al., *Proc. Natl. Acad. Sci. USA*, 88:9578-9582 (1991)) as disclosed by Chevray and Nathans, *Proc. Natl. Acad. Sci. USA*, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-lacZ reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

[0298] Compounds that interfere with the interaction of a gene encoding a PRO polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

[0299] To assay for antagonists, the PRO polypeptide may be added to a cell along with the compound to be screened for a particular activity and the ability of the compound to inhibit the activity of interest in the presence of the PRO polypeptide indicates that the compound is an antagonist to

the PRO polypeptide. Alternatively, antagonists may be detected by combining the PRO polypeptide and a potential antagonist with membrane-bound PRO polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO polypeptide can be labeled, such as by radioactivity, such that the number of PRO polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan et al., *Current Protocols in Immun.*, 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO polypeptide. Transfected cells that are grown on glass slides are exposed to labeled PRO polypeptide. The PRO polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

[0300] As an alternative approach for receptor identification, labeled PRO polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

[0301] In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with labeled PRO polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

[0302] More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with PRO polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO polypeptide.

[0303] Another potential PRO polypeptide antagonist is an antisense RNA or DNA construct prepared using antisense technology, where, e.g., an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a

polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO polypeptides herein, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix—see Lee et al., *Nucl. Acids Res.*, 6:3073 (1979); Cooney et al., *Science*, 241: 456 (1988); Dervan et al., *Science*, 251:1360 (1991)), thereby preventing transcription and the production of the PRO polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into the PRO polypeptide (antisense—Okano, *Neurochem.*, 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression* (CRC Press: Boca Raton, Fla., 1988). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed in vivo to inhibit production of the PRO polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, e.g., between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

[0304] Potential antagonists include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO polypeptide, thereby blocking the normal biological activity of the PRO polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

[0305] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g., Rossi, *Current Biology*, 4:469-471 (1994), and PCT publication No. WO 97/33551 (published Sept. 18, 1997).

[0306] Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, e.g., PCT publication No. WO 97/33551, supra.

[0307] These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

[0308] Diagnostic and therapeutic uses of the herein disclosed molecules may also be based upon the positive functional assay hits disclosed and described below.

[0309] F. Anti-PRO Antibodies

[0310] The present invention further provides anti-PRO antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

[0311] 1. Polyclonal Antibodies

[0312] The anti-PRO antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

[0313] 2. Monoclonal Antibodies

[0314] The anti-PRO antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

[0315] The immunizing agent will typically include the PRO polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

[0316] Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, Calif. and the American Type Culture Collection, Manassas, Va. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies.

ies [Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

[0317] The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against PRO. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, *Anal. Biochem.*, 107:220 (1980).

[0318] After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, *supra*]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown in vivo as ascites in a mammal.

[0319] The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

[0320] The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Pat. No. 4,816,567; Morrison et al., *supra*] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

[0321] The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

[0322] In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce

fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

[0323] 3. Human and Humanized Antibodies

[0324] The anti-PRO antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992)].

[0325] Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeven et al., *Science*, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[0326] Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into

transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., *Bio/Technology* 10, 779-783 (1992); Lonberg et al., *Nature* 368 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild et al., *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

[0327] The antibodies may also be affinity matured using known selection and/or mutagenesis methods as described above. Preferred affinity matured antibodies have an affinity which is five times, more preferably 10 times, even more preferably 20 or 30 times greater than the starting antibody (generally murine, humanized or human) from which the matured antibody is prepared.

[0328] 4. Bispecific Antibodies

[0329] Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

[0330] Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, *Nature*, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published May 13, 1993, and in Traunecker et al., *EMBO J.*, 10:3655-3659 (1991).

[0331] Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 121:210 (1986).

[0332] According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of

the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[0333] Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared can be prepared using chemical linkage. Brennan et al., *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then converted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

[0334] Fab' fragments may be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

[0335] Various technique for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148(5): 1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., *J. Immunol.* 152:5368 (1994). Antibodies with more than two valen-

cies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

[0336] Exemplary bispecific antibodies may bind to two different epitopes on a given PRO polypeptide herein. Alternatively, an anti-PRO polypeptide arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD 16) so as to focus cellular defense mechanisms to the cell expressing the particular PRO polypeptide. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express a particular PRO polypeptide. These antibodies possess a PRO-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the PRO polypeptide and further binds tissue factor (TF).

[0337] 5. Heteroconjugate Antibodies

[0338] Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Pat. No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Pat. No. 4,676,980.

[0339] 6. Effector Function Engineering

[0340] It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp. Med.*, 176: 1191-1195 (1992) and Shopes, *J. Immunol.*, 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff et al. *Cancer Research*, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*, 3: 219-230 (1989).

[0341] 7. Immunoconjugates

[0342] The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

[0343] Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above.

Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ²¹²Bi, ¹³¹I, ¹³¹In, ⁹⁰Y, and ¹⁸⁶Re. Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimide HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis(p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., *Science*, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionuclide to the antibody. See WO94/11026.

[0344] In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is conjugated to a cytotoxic agent (e.g., a radionuclide).

[0345] 8. Immunoliposomes

[0346] The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., *Proc. Natl. Acad. Sci. USA*, 82: 3688 (1985); Hwang et al., *Proc. Natl. Acad. Sci. USA*, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556.

[0347] Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., *J. Biol. Chem.*, 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon et al., *J. National Cancer Inst.*, 81(19): 1484 (1989).

[0348] 9. Pharmaceutical Compositions of Antibodies

[0349] Antibodies specifically binding a PRO polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders in the form of pharmaceutical compositions.

[0350] If the PRO polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco et al., *Proc. Natl. Acad. Sci. USA*, 90: 7889-7893 (1993). The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[0351] The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nano-capsules) or in macroemulsions. Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, supra.

[0352] The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

[0353] Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S—S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

[0354] G. Uses for anti-PRO Antibodies

[0355] The anti-PRO antibodies of the invention have various utilities. For example, anti-PRO antibodies may be used in diagnostic assays for PRO, e.g., detecting its expression (and in some cases, differential expression) in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art may be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases [Zola, *Monoclonal Antibodies: A Manual of Techniques*, CRC Press, Inc. (1987) pp. 147-158]. The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ³H, ¹⁴C, ³²P, ³⁵S, or ¹²⁵I, a fluorescent or chemilumines compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter et al., *Nature*, 144:945 (1962); David et al., *Biochemistry*, 13:1014 (1974); Pain et al., *J. Immunol. Meth.*, 40:219 (1981); and Nygren, *J. Histochem. and Cytochem.*, 30:407 (1982).

[0356] Anti-PRO antibodies also are useful for the affinity purification of PRO from recombinant cell culture or natural sources. In this process, the antibodies against PRO are immobilized on a suitable support, such as a Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the PRO to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the PRO, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the PRO from the antibody.

[0357] The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

[0358] All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

EXAMPLES

[0359] Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, Va.

Example 1

Extracellular Domain Homology Screening to Identify Novel Polypeptides and cDNA Encoding Therefor

[0360] The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (e.g., Dayhoff, GenBank), and proprietary databases (e.g. LIFESEQTM, Incyte

Pharmaceuticals, Palo Alto, Calif.). The search was performed using the computer program BLAST or BLAST-2 (Altschul et al., *Methods in Enzymology*, 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Wash.).

[0361] Using this extracellular domain homology screen, consensus DNA sequences were assembled relative to the other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST or BLAST-2 and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

[0362] Based upon the consensus sequences obtained as described above, oligonucleotides were then synthesized and used to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., *Current Protocols in Molecular Biology*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

[0363] The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, Calif. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or PRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

Example 2

Isolation of cDNA Clones by Amylase Screening

[0364] 1. Preparation of oligo dT Primed cDNA Library

[0365] mRNA was isolated from a human tissue of interest using reagents and protocols from Invitrogen, San Diego, Calif. (Fast Track 2). This RNA was used to generate an oligo dT primed cDNA library in the vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, Md. (Super Script Plasmid System). In this procedure, the double stranded cDNA was sized to greater than 1000 bp and the SalI/NotI Tinkered cDNA was cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

[0366] 2. Preparation of Random Primed cDNA Library

[0367] A secondary cDNA library was generated in order to preferentially represent the 5' ends of the primary cDNA clones. Sp6 RNA was generated from the primary library (described above), and this RNA was used to generate a random primed cDNA library in the vector pSST-AMY.0 using reagents and protocols from Life Technologies (Super Script Plasmid System, referenced above). In this procedure the double stranded cDNA was sized to 500-1000 bp, Tinkered with blunt to NotI adaptors, cleaved with SfiI, and cloned into SfiI/NotI cleaved vector. pSST-AMY.0 is a cloning vector that has a yeast alcohol dehydrogenase promoter preceding the cDNA cloning sites and the mouse amylase sequence (the mature sequence without the secretion signal) followed by the yeast alcohol dehydrogenase terminator, after the cloning sites. Thus, cDNAs cloned into this vector that are fused in frame with amylase sequence will lead to the secretion of amylase from appropriately transfected yeast colonies.

[0368] 3. Transformation and Detection

[0369] DNA from the library described in paragraph 2 above was chilled on ice to which was added electrocompetent DH10B bacteria (Life Technologies, 20 ml). The bacteria and vector mixture was then electroporated as recommended by the manufacturer. Subsequently, SOC media (Life Technologies, 1 ml) was added and the mixture was incubated at 37° C. for 30 minutes. The transformants were then plated onto 20 standard 150 mm LB plates containing ampicillin and incubated for 16 hours (37° C.). Positive colonies were scraped off the plates and the DNA was isolated from the bacterial pellet using standard protocols, e.g. CsCl-gradient. The purified DNA was then carried on to the yeast protocols below.

[0370] The yeast methods were divided into three categories: (1) Transformation of yeast with the plasmid/cDNA combined vector; (2) Detection and isolation of yeast clones secreting amylase; and (3) PCR amplification of the insert directly from the yeast colony and purification of the DNA for sequencing and further analysis.

[0371] The yeast strain used was HD56-5A (ATCC-90785). This strain has the following genotype: MAT alpha, ura3-52, leu2-3, leu2-112, his3-11, his3-15, MAL⁺, SUC⁺, GAL⁺. Pref employed that have deficient post-translational pathways. Such mutants may have translocation deficient alleles in sec71, sec72, sec62, with truncated sec71 being most preferred. Alternatively, antagonists (including antisense nucleotides and/or ligands) which interfere with the normal operation of these genes, other proteins implicated in this post translation pathway (e.g., SEC61p, SEC72p, SEC62p, SEC63p, TDJ1p or SSA1p-4p) or the complex formation of these proteins may also be preferably employed in combination with the amylase-expressing yeast.

[0372] Transformation was performed based on the protocol outlined by Gietz et al., *Nucl. Acid. Res.*, 20:1425 (1992). Transformed cells were then inoculated from agar into YEPD complex media broth (100 ml) and grown overnight at 30° C. The YEPD broth was prepared as described in Kaiser et al., *Methods in Yeast Genetics*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y., p. 207 (1994). The overnight culture was then diluted to about 2×10⁷ cells/ml (approx. OD₆₀₀=1) into fresh YEPD broth (500 ml) and regrown to 1×10⁷ cells OD₆₀₀=0.4-0.5).

[0373] The cells were then harvested and prepared for transformation by transfer into GS3 rotor bottles in a Sorval GS3 rotor at 5,000 rpm for 5 minutes, the supernatant discarded, and then resuspended into sterile water, and centrifuged again in 50 ml falcon tubes at 3,500 rpm in a Beckman GS-6KR centrifuge. The supernatant was discarded and the cells were subsequently washed with LiAc/TE (10 ml, 10 mM Tris-HCl, 1 mM EDTA pH 7.5, 100 mM Li₂OOCCH₃), and resuspended into LiAc/TE (2.5 ml).

[0374] Transformation took place by mixing the prepared cells (100 μ l) with freshly denatured single stranded salmon testes DNA (Lofstrand Labs, Gaithersburg, Md.) and transforming DNA (1 μ g, vol.<10 μ l) in microfuge tubes. The mixture was mixed briefly by vortexing, then 40% PEG/TE (600 μ l, 40% polyethylene glycol-4000, 10 mM Tris-HCl, 1 mM EDTA, 100 mM Li₂OOCCH₃, pH 7.5) was added. This mixture was gently mixed and incubated at 30° C. while agitating for 30 minutes. The cells were then heat shocked at 42° C. for 15 minutes, and the reaction vessel centrifuged in a microfuge at 12,000 rpm for 5-10 seconds, decanted and resuspended into TE (500 μ l, 10 mM Tris-HCl, 1 mM EDTA pH 7.5) followed by recentrifugation. The cells were then diluted into TE (1 ml) and aliquots (200 μ l) were spread onto the selective media previously prepared in 150 mm growth plates (VWR).

[0375] Alternatively, instead of multiple small reactions, the transformation was performed using a single, large scale reaction, wherein reagent amounts were scaled up accordingly.

[0376] The selective media used was a synthetic complete dextrose agar lacking uracil (SCD-Ura) prepared as described in Kaiser et al., *Methods in Yeast Genetics*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y., p. 208-210 (1994). Transformants were grown at 30° C. for 2-3 days.

[0377] The detection of colonies secreting amylase was performed by including red starch in the selective growth media. Starch was coupled to the red dye (Reactive Red-120, Sigma) as per the procedure described by Biely et al., *Anal. Biochem.*, 172:176-179 (1988). The coupled starch was incorporated into the SCD-Ura agar plates at a final concentration of 0.15% (w/v), and was buffered with potassium phosphate to a pH of 7.0 (50-100 mM final concentration).

[0378] The positive colonies were picked and streaked across fresh selective media (onto 150 mm plates) in order to obtain well isolated and identifiable single colonies. Well isolated single colonies positive for amylase secretion were detected by direct incorporation of red starch into buffered SCD-Ura agar. Positive colonies were determined by their ability to break down starch resulting in a clear halo around the positive colony visualized directly.

[0379] 4. Isolation of DNA by PCR Amplification

[0380] When a positive colony was isolated, a portion of it was picked by a toothpick and diluted into sterile water (30 μ l) in a 96 well plate. At this time, the positive colonies were either frozen and stored for subsequent analysis or immediately amplified. An aliquot of cells (5 μ L) was used as a template for the PCR reaction in a 25 μ l volume containing: 0.5 μ l KlenTaq (Clontech, Palo Alto, Calif.); 4.0 μ l 10 mM dNTP's (Perkin Elmer-Cetus); 2.5 μ l Kentaq buffer (Clon-

tech); 0.25 μ l forward oligo 1; 0.25 μ l reverse oligo 2; 12.5 μ l distilled water. The sequence of the forward oligonucleotide 1 was:

[0381] 5'-TGTAACGACGGCCAGTTAAATAGAC-CTGCAATTATTAATCT-3' (SEQ ID NO:115)

[0382] The sequence of reverse oligonucleotide 2 was:

[0383] 5'-CAGGAAACAGCTATGACCACCTGCACACCTGCAAATCCATT-3' (SEQ ID NO:116)

[0384] PCR was then performed as follows:

a.	Denature	92° C., 5 minutes
b. 3 cycles of:	Denature	92° C., 30 seconds
	Anneal	59° C., 30 seconds
	Extend	72° C., 60 seconds
c. 3 cycles of:	Denature	92° C., 30 seconds
	Anneal	57° C., 30 seconds
	Extend	72° C., 60 seconds
d. 25 cycles of:	Denature	92° C., 30 seconds
	Anneal	55° C., 30 seconds
	Extend	72° C., 60 seconds
e.	Hold	4° C.

[0385] The underlined regions of the oligonucleotides annealed to the ADH promoter region and the amylase region, respectively, and amplified a 307 bp region from vector pSST-AMY.0 when no insert was present. Typically, the first 18 nucleotides of the 5' end of these oligonucleotides contained annealing sites for the sequencing primers. Thus, the total product of the PCR reaction from an empty vector was 343 bp. However, signal sequence-fused cDNA resulted in considerably longer nucleotide sequences.

[0386] Following the PCR, an aliquot of the reaction (5 μ l) was examined by agarose gel electrophoresis in a 1% agarose gel using a Tris-Borate-EDTA (TBE) buffering system as described by Sambrook et al., supra. Clones resulting in a single strong PCR product larger than 400 bp were further analyzed by DNA sequencing after purification with a 96 Qiaquick PCR clean-up column (Qiagen Inc., Chatsworth, Calif.).

Example 3

Isolation of cDNA Clones Using Signal Algorithm Analysis

[0387] Various polypeptide-encoding nucleic acid sequences were identified by applying a proprietary signal sequence finding algorithm developed by Genentech, Inc. (South San Francisco, Calif.) upon ESTs as well as clustered and assembled EST fragments from public (e.g., GenBank) and/or private (LIFESEQ®, Incyte Pharmaceuticals, Inc., Palo Alto, Calif.) databases. The signal sequence algorithm computes a secretion signal score based on the character of the DNA nucleotides surrounding the first and optionally the second methionine codon(s) (ATG) at the 5'-end of the sequence or sequence fragment under consideration. The nucleotides following the first ATG must code for at least 35 unambiguous amino acids without any stop codons. If the first ATG has the required amino acids, the second is not examined. If neither meets the requirement, the candidate sequence is not scored. In order to determine whether the EST sequence contains an authentic signal sequence, the

DNA and corresponding amino acid sequences surrounding the ATG codon are scored using a set of seven sensors (evaluation parameters) known to be associated with secretion signals. Use of this algorithm resulted in the identification of numerous polypeptide-encoding nucleic acid sequences.

Example 4

Isolation of cDNA clones Encoding Human PRO Polypeptides

[0388] Using the techniques described in Examples 1 to 3 above, numerous full-length cDNA clones were identified as encoding PRO polypeptides as disclosed herein. These cDNAs were then deposited under the terms of the Budapest Treaty with the American Type Culture Collection, 10801 University Blvd., Manassas, Va. 20110-2209, USA (ATCC) as shown in Table 7 below.

TABLE 7

Material	ATCC Dep. No.	Deposit Date
DNA16422-1209	209929	Jun. 2, 1998
DNA19902-1669	203454	Nov. 3, 1998
DNA21624-1391	209917	Jun. 2, 1998
DNA34387-1138	209260	Sep. 16, 1997
DNA35880-1160	209379	Oct. 16, 1997
DNA39984-1221	209435	Nov. 7, 1997
DNA44189-1322	209699	Mar. 26, 1998
DNA48303-2829	PTA-1342	Feb. 8, 2000
DNA48320-1433	209904	May 27, 1998
DNA56049-2543	203662	Feb. 9, 1999
DNA57694-1341	203017	Jun. 23, 1998
DNA59208-1373	209881	May 20, 1998
DNA59214-1449	203046	Jul. 1, 1998
DNA59485-1336	203015	Jun. 23, 1998
DNA64966-1575	203575	Jan. 12, 1999
DNA82403-2959	PTA-2317	Aug. 1, 2000
DNA83505-2606	PTA-132	May 25, 1999
DNA84927-2585	203865	Mar. 23, 1999
DNA92264-2616	203969	Apr. 27, 1999
DNA94713-2561	203835	Mar. 9, 1999
DNA96869-2673	PTA-255	Jun. 22, 1999
DNA96881-2699	PTA-553	Aug. 17, 1999
DNA96889-2641	PTA-119	May 25, 1999
DNA96898-2640	PTA-122	May 25, 1999
DNA97003-2649	PTA-43	May 11, 1999
DNA98565-2701	PTA-481	Aug. 3, 1999
DNA102846-2742	PTA-545	Aug. 17, 1999
DNA102847-2726	PTA-517	Aug. 10, 1999
DNA102880-2689	PTA-383	Jul. 20, 1999
DNA105782-2683	PTA-387	Jul. 20, 1999
DNA108912-2680	PTA-124	May, 25, 1999
DNA115253-2757	PTA-612	Aug. 31, 1999
DNA119302-2737	PTA-520	Aug. 10, 1999
DNA119536-2752	PTA-551	Aug. 17, 1999
DNA119542-2754	PTA-619	Aug. 31, 1999
DNA143498-2824	PTA-1263	Feb. 2, 2000
DNA145583-2820	PTA-1179	Jan. 11, 2000
DNA161000-2896	PTA-1731	Apr. 18, 2000
DNA161005-2943	PTA-2243	Jun. 27, 2000
DNA170245-3053	PTA-2952	Jan. 23, 2001
DNA171771-2919	PTA-1902	May 23, 2000
DNA173157-2981	PTA-2388	Aug. 8, 2000
DNA175734-2985	PTA-2455	Sep. 12, 2000
DNA176108-3040	PTA-2824	Dec. 19, 2000
DNA190710-3028	PTA-2822	Dec. 19, 2000
DNA190803-3019	PTA-2785	Dec. 12, 2000
DNA191064-3069	PTA-3016	Feb. 6, 2001
DNA194909-3013	PTA-2779	Dec. 12, 2000
DNA203532-3029	PTA-2823	Dec. 19, 2000
DNA213858-3060	PTA-2958	Jan. 23, 2001
DNA216676-3083	PTA-3157	Mar. 6, 2001

TABLE 7-continued

Material	ATCC Dep. No.	Deposit Date
DNA222653-3104	PTA-3330	Apr. 24, 2001
DNA96897-2688	PTA-379	Jul. 20, 1999
DNA142917-3081	PTA-3155	Mar. 6, 2001
DNA142930-2914	PTA-1901	May 23, 2000
DNA147253-2983	PTA-2405	Aug. 22, 2000
DNA149927-2887	PTA-1782	Apr. 25, 2000

[0389] These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Pat. Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

[0390] The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

Example 5

Use of PRO as a Hybridization Probe

[0391] The following method describes use of a nucleotide sequence encoding PRO as a hybridization probe.

[0392] DNA comprising the coding sequence of full-length or mature PRO as disclosed herein is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO) in human tissue cDNA libraries or human tissue genomic libraries.

[0393] Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO-derived probe to the filters is performed in a solution of 50% formamide, 5×SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2×Denhardt's solution, and 10% dextran sulfate at 42° C. for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1×SSC and 0.1% SDS at 42° C.

[0394] DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO can then be identified using standard techniques known in the art.

Example 6

Expression of PRO in *E. coli*

[0395] This example illustrates preparation of an unglycosylated form of PRO by recombinant expression in *E. coli*.

[0396] The DNA sequence encoding PRO is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO coding region, lambda transcriptional terminator, and an argu gene.

[0397] The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., supra. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

[0398] Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

[0399] After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

[0400] PRO may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) Ion galE rpoHts(htpRts) clpP(lacIq). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30° C. with shaking until an O.D.600 of 3-5 is reached. Cultures are then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH₄)₂SO₄, 0.71 g sodium citrate•2H₂O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO₄) and grown for approximately 20-30 hours at 30° C. with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

[0401] *E. coli* paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1 M and 0.02 M, respectively, and the solution is stirred

overnight at 4° C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4° C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

[0402] The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4° C. for 12-36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros RI/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

[0403] Fractions containing the desired folded PRO polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

[0404] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

Example 7

Expression of PRO in Mammalian Cells

[0405] This example illustrates preparation of a potentially glycosylated form of PRO by recombinant expression in mammalian cells.

[0406] The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO DNA using ligation methods such as described in Sambrook et al., supra. The resulting vector is called pRK5-PRO.

[0407] In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 μ g pRK5-PRO DNA is mixed with about 1 μ g DNA encoding the VA RNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500 μ l of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl_2 . To this mixture is added, dropwise, 500 μ l of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO_4 , and a precipitate is allowed to form for 10 minutes at 25° C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37° C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

[0408] Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 μ Ci/ml ^{35}S -cysteine and 200 μ Ci/ml ^{35}S -methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

[0409] In an alternative technique, PRO may be introduced into 293 cells transiently using the dextran sulfate method described by Sompariyac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 μ g pRK5-PRO DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 μ g/ml bovine insulin and 0.1 μ g/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

[0410] In another embodiment, PRO can be expressed in CHO cells. The pRK5-PRO can be transfected into CHO cells using known reagents such as CaPO_4 or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ^{35}S -methionine. After determining the presence of PRO polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO can then be concentrated and purified by any selected method.

[0411] Epitope-tagged PRO may also be expressed in host CHO cells. The PRO may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO insert can then be subcloned into a SV40 driven vector

containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO can then be concentrated and purified by any selected method, such as by Ni^{2+} -chelate affinity chromatography.

[0412] PRO may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

[0413] Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

[0414] Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel et al., *Current Protocols of Molecular Biology*, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., *Nucl. Acids Res.* 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

[0415] Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Qiagen), Dosper® or Eugene® (Boehringer Mannheim). The cells are grown as described in Lucas et al., supra. Approximately 3×10^7 cells are frozen in an ampule for further growth and production as described below.

[0416] The ampules containing the plasmid DNA are thawed by placement into water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 mLs of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 mL of selective media (0.2 μ m filtered PS20 with 5% 0.2 μ m diafiltered fetal bovine serum). The cells are then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells are transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37° C. After another 2-3 days, 250 mL, 500 mL and 2000 mL, spinners are seeded with 3×10^5 cells/mL. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Pat. No. 5,122,469, issued Jun. 16, 1992 may actually be used. A 3L production spinner is seeded at 1.2×10^6 cells/mL. On day 0, the cell number pH is determined. On day 1, the spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33° C., and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at

around 7.2. After 10 days, or until the viability dropped below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22 μ m filter. The filtrate was either stored at 4° C. or immediately loaded onto columns for purification.

[0417] For the poly-His tagged constructs, the proteins are purified using a Ni-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4° C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80° C.

[0418] Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 μ L of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

[0419] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

Example 8

Expression of PRO in Yeast

[0420] The following method describes recombinant expression of PRO in yeast.

[0421] First, yeast expression vectors are constructed for intracellular production or secretion of PRO from the ADH2/GAPDH promoter. DNA encoding PRO and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO. For secretion, DNA encoding PRO can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO.

[0422] Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

[0423] Recombinant PRO can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate

containing PRO may further be purified using selected column chromatography resins.

[0424] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

Example 9

Expression of PRO in Baculovirus-Infected Insect Cells

[0425] The following method describes recombinant expression of PRO in Baculovirus-infected insect cells.

[0426] The sequence coding for PRO is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO or the desired portion of the coding sequence of PRO such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

[0427] Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4-5 days of incubation at 28° C., the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilly et al., *Baculovirus expression vectors: A Laboratory Manual*, Oxford: Oxford University Press (1994).

[0428] Expressed poly-his tagged PRO can then be purified, for example, by Ni²⁺-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., *Nature*, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl₂; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 μ m filter. A Ni²⁺-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A₂₈₀ with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A₂₈₀ baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni²⁺-NTA-conjugated to alkaline phosphatase

(Qiagen). Fractions containing the eluted His₁₀-tagged PRO are pooled and dialyzed against loading buffer.

[0429] Alternatively, purification of the IgG tagged (or Fc tagged) PRO can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

[0430] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

Example 10

Preparation of Antibodies that Bind PRO

[0431] This example illustrates preparation of monoclonal antibodies which can specifically bind PRO.

[0432] Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO, fusion proteins containing PRO, and cells expressing recombinant PRO on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

[0433] Mice, such as Balb/c, are immunized with the PRO immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, Mont.) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO antibodies.

[0434] After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

[0435] The hybridoma cells will be screened in an ELISA for reactivity against PRO. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO is within the skill in the art.

[0436] The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

Example 11

Purification of PRO Polypeptides Using Specific Antibodies

[0437] Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polypeptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

[0438] Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSETM (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

[0439] Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

[0440] A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (e.g., a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

Example 12

Drug Screening

[0441] This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested.

Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

[0442] Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an agent with an PRO polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

[0443] Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on Sept. 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

[0444] This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

Example 13

Rational Drug Design

[0445] The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (i.e., a PRO polypeptide) or of small molecules with which they interact, e.g., agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide in vivo (cf., Hodgson, *Bio/Technology*, 9: 19-21 (1991)).

[0446] In one approach, the three-dimensional structure of the PRO polypeptide, or of an PRO polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant

structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, *Biochemistry*, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda et al., *J. Biochem.*, 113:742-746 (1993).

[0447] It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

[0448] By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

Example 14

Ability of PRO Polypeptides to Stimulate the Release of Proteoglycans from Cartilage (Assay 97)

[0449] The ability of various PRO polypeptides to stimulate the release of proteoglycans from cartilage tissue was tested as follows.

[0450] The metacarpophalangeal joint of 4-6 month old pigs was aseptically dissected, and articular cartilage was removed by free hand slicing being careful to avoid the underlying bone. The cartilage was minced and cultured in bulk for 24 hours in a humidified atmosphere of 95% air, 5% CO₂ in serum free (SF) media (DME/F12 1:1) with 0.1% BSA and 100 U/ml penicillin and 100 µg/ml streptomycin. After washing three times, approximately 100 mg of articular cartilage was aliquoted into micronics tubes and incubated for an additional 24 hours in the above SF media. PRO polypeptides were then added at 1% either alone or in combination with 18 ng/ml interleukin-1 α , a known stimulator of proteoglycan release from cartilage tissue. The supernatant was then harvested and assayed for the amount of proteoglycans using the 1,9-dimethyl-methylene blue (DMB) colorimetric assay (Farndale and Buttle, *Biochem. Biophys. Acta* 883:173-177 (1985)). A positive result in this assay indicates that the test polypeptide will find use, for example, in the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis.

[0451] When various PRO polypeptides were tested in the above assay, the polypeptides demonstrated a marked ability to stimulate release of proteoglycans from cartilage tissue both basally and after stimulation with interleukin-1 α and at 24 and 72 hours after treatment, thereby indicating that these

PRO polypeptides are useful for stimulating proteoglycan release from cartilage tissue. As such, these PRO polypeptides are useful for the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis. PRO6018 polypeptide testing positive in this assay.

Example 15

Human Microvascular Endothelial Cell Proliferation (Assay 146)

[0452] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce proliferation of human microvascular endothelial cells in culture and, therefore, function as useful growth factors.

[0453] On day 0, human microvascular endothelial cells were plated in 96-well plates at 1000 cells/well per 100 microliter and incubated overnight in complete media [EBM-2 growth media, plus supplements: IGF-1; ascorbic acid; VEGF; hEGF; hFGF; hydrocortisone, gentamicin (GA-1000), and fetal bovine serum (FBS, Clonetics)]. On day 1, complete media was replaced by basal media [EBM-2 plus 1% FBS] and addition of PRO polypeptides at 1%, 0.1% and 0.01%. On day 7, an assessment of cell proliferation was performed using the ViaLight HS kit [ATP/luciferase Lumitech]. Results are expressed as % of the cell growth observed with control buffer.

[0454] The following PRO polypeptides stimulated human microvascular endothelial cell proliferation in this assay: PRO1313, PRO20080, and PRO21383.

[0455] The following PRO polypeptides inhibited human microvascular endothelial cell proliferation in this assay: PRO6071, PRO4487, and PRO6006.

Example 16

Microarray Analysis to Detect Overexpression of PRO Polypeptides in Cancerous Tumors

[0456] Nucleic acid microarrays, often containing thousands of gene sequences, are useful for identifying differentially expressed genes in diseased tissues as compared to their normal counterparts. Using nucleic acid microarrays, test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The cDNA probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of each member of the array is known. For example, a selection of genes known to be expressed in certain disease states may be arrayed on a solid support. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. If the hybridization signal of a probe from a test (disease tissue) sample is greater than hybridization signal of a probe from a control (normal tissue) sample, the gene or genes overexpressed in the disease tissue are identified. The implication of this result is that an overexpressed protein in a diseased tissue is useful not only as a diagnostic marker for the presence of the disease condition, but also as a therapeutic target for treatment of the disease condition.

[0457] The methodology of hybridization of nucleic acids and microarray technology is well known in the art. In the present example, the specific preparation of nucleic acids for hybridization and probes, slides, and hybridization conditions are all detailed in U.S. Provisional Patent Application Serial No. 60/193,767, filed on Mar. 31, 2000 and which is herein incorporated by reference.

[0458] In the present example, cancerous tumors derived from various human tissues were studied for PRO polypeptide-encoding gene expression relative to non-cancerous human tissue in an attempt to identify those PRO polypeptides which are overexpressed in cancerous tumors. Cancerous human tumor tissue from any of a variety of different human tumors was obtained and compared to a “universal” epithelial control sample which was prepared by pooling non-cancerous human tissues of epithelial origin, including liver, kidney, and lung. mRNA isolated from the pooled tissues represents a mixture of expressed gene products from these different tissues. Microarray hybridization experiments using the pooled control samples generated a linear plot in a 2-color analysis. The slope of the line generated in a 2-color analysis was then used to normalize the ratios of (test:control detection) within each experiment. The normalized ratios from various experiments were then compared and used to identify clustering of gene expression. Thus, the pooled “universal control” sample not only allowed effective relative gene expression determinations in a simple 2-sample comparison, it also allowed multi-sample comparisons across several experiments.

[0459] In the present experiments, nucleic acid probes derived from the herein described PRO polypeptide-encoding nucleic acid sequences were used in the creation of the microarray and RNA from a panel of nine different tumor tissues (listed below) were used for the hybridization thereto. A value based upon the normalized ratio:experimental ratio was designated as a “cutoff ratio”. Only values that were above this cutoff ratio were determined to be significant. Table 8 below shows the results of these experiments, demonstrating that various PRO polypeptides of the present invention are significantly overexpressed in various human tumor tissues, as compared to a non-cancerous human tissue control or other human tumor tissues. As described above, these data demonstrate that the PRO polypeptides of the present invention are useful not only as diagnostic markers for the presence of one or more cancerous tumors, but also serve as therapeutic targets for the treatment of those tumors.

TABLE 8

Molecule	is overexpressed in:	as compared to normal control:
PRO240	breast tumor	universal normal control
PRO240	lung tumor	universal normal control
PRO256	colon tumor	universal normal control
PRO256	lung tumor	universal normal control
PRO256	breast tumor	universal normal control
PRO306	colon tumor	universal normal control
PRO306	lung tumor	universal normal control
PRO540	lung tumor	universal normal control
PRO540	colon tumor	universal normal control
PRO773	breast tumor	universal normal control
PRO773	colon tumor	universal normal control
PRO698	colon tumor	universal normal control
PRO698	breast tumor	universal normal control
PRO698	lung tumor	universal normal control
PRO698	prostate tumor	universal normal control

TABLE 8-continued

Molecule	is overexpressed in:	as compared to normal control:
PRO698	rectal tumor	universal normal control
PRO3567	colon tumor	universal normal control
PRO3567	breast tumor	universal normal control
PRO3567	lung tumor	universal normal control
PRO826	colon tumor	universal normal control
PRO826	lung tumor	universal normal control
PRO826	breast tumor	universal normal control
PRO826	rectal tumor	universal normal control
PRO826	liver tumor	universal normal control
PRO1002	colon tumor	universal normal control
PRO1002	lung tumor	universal normal control
PRO1068	colon tumor	universal normal control
PRO1068	breast tumor	universal normal control
PRO1030	colon tumor	universal normal control
PRO1030	breast tumor	universal normal control
PRO1030	lung tumor	universal normal control
PRO1030	prostate tumor	universal normal control
PRO1030	rectal tumor	universal normal control
PRO4397	colon tumor	universal normal control
PRO4397	breast tumor	universal normal control
PRO4344	colon tumor	universal normal control
PRO4344	lung tumor	universal normal control
PRO4344	rectal tumor	universal normal control
PRO4407	colon tumor	universal normal control
PRO4407	breast tumor	universal normal control
PRO4407	lung tumor	universal normal control
PRO4407	liver tumor	universal normal control
PRO4407	rectal tumor	universal normal control
PRO4316	colon tumor	universal normal control
PRO5775	colon tumor	universal normal control
PRO6016	colon tumor	universal normal control
PRO4980	breast tumor	universal normal control
PRO4980	colon tumor	universal normal control
PRO4980	lung tumor	universal normal control
PRO6018	colon tumor	universal normal control
PRO7168	colon tumor	universal normal control
PRO6000	colon tumor	universal normal control
PRO6006	colon tumor	universal normal control
PRO5800	colon tumor	universal normal control
PRO5800	breast tumor	universal normal control
PRO5800	lung tumor	universal normal control
PRO5800	rectal tumor	universal normal control
PRO7476	colon tumor	universal normal control
PRO10268	colon tumor	universal normal control
PRO6496	colon tumor	universal normal control
PRO6496	breast tumor	universal normal control
PRO6496	lung tumor	universal normal control
PRO7422	colon tumor	universal normal control
PRO7431	colon tumor	universal normal control
PRO28633	colon tumor	universal normal control
PRO28633	lung tumor	universal normal control
PRO28633	liver tumor	universal normal control
PRO21485	colon tumor	universal normal control
PRO28700	breast tumor	universal normal control
PRO28700	lung tumor	universal normal control
PRO28700	colon tumor	universal normal control
PRO34012	colon tumor	universal normal control
PRO34012	lung tumor	universal normal control
PRO34003	colon tumor	universal normal control
PRO34003	lung tumor	universal normal control
PRO34001	colon tumor	universal normal control
PRO34009	colon tumor	universal normal control
PRO34009	breast tumor	universal normal control
PRO34009	lung tumor	universal normal control
PRO34009	rectal tumor	universal normal control
PRO34192	colon tumor	universal normal control
PRO34564	colon tumor	universal normal control
PRO35444	colon tumor	universal normal control
PRO5998	colon tumor	universal normal control
PRO5998	lung tumor	universal normal control
PRO5998	kidney tumor	universal normal control
PRO19651	colon tumor	universal normal control

TABLE 8-continued

Molecule	is overexpressed in:	as compared to normal control:
PRO20221	liver tumor	universal normal control
PRO21434	liver tumor	universal normal control

Example 17

Fetal Hemoglobin Induction in an Erythroblastic Cell Line (Assay 107)

[0460] This assay is useful for screening PRO polypeptides for the ability to induce the switch from adult hemoglobin to fetal hemoglobin in an erythroblastic cell line. Molecules testing positive in this assay are expected to be useful for therapeutically treating various mammalian hemoglobin-associated disorders such as the various thalassemias. The assay is performed as follows. Erythroblastic cells are plated in standard growth medium at 1000 cells/well in a 96 well format. PRO polypeptides are added to the growth medium at a concentration of 0.2% or 2% and the cells are incubated for 5 days at 37° C. As a positive control, cells are treated with 100 μM hemin and as a negative control, the cells are untreated. After 5 days, cell lysates are prepared and analyzed for the expression of gamma globin (a fetal marker). A positive in the assay is a gamma globin level at least 2-fold above the negative control.

[0461] PRO20080 polypeptide tested positive in this assay.

Example 18

Microarray Analysis to Detect Overexpression of PRO Polypeptides in HUVEC Cells Treated with Growth Factors

[0462] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce angiogenesis by stimulating endothelial cell tube formation in HUVEC cells.

[0463] Nucleic acid microarrays, often containing thousands of gene sequences, are useful for identifying differentially expressed genes in tissues exposed to various stimuli (e.g., growth factors) as compared to their normal, unexposed counterparts. Using nucleic acid microarrays, test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The cDNA probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of each member of the array is known. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. If the hybridization signal of a probe from a test (exposed tissue) sample is greater than hybridization signal of a probe from a control (normal, unexposed tissue) sample, the gene or genes overexpressed in the exposed tissue are identified. The implication of this result is that an overexpressed protein in an exposed tissue may be involved in the functional changes within the tissue following exposure to the stimuli (e.g., tube formation).

[0464] The methodology of hybridization of nucleic acids and microarray technology is well known in the art. In the

present example, the specific preparation of nucleic acids for hybridization and probes, slides, and hybridization conditions are all detailed in U.S. Provisional Patent Application Serial No. 60/193,767, filed on Mar. 31, 2000 and which is herein incorporated by reference.

[0465] In the present example, HUVEC cells grown in either collagen gels or fibrin gels were induced to form tubes by the addition of various growth factors. Specifically, collagen gels were prepared as described previously in Yang et al., *American J. Pathology*, 1999, 155(3):887-895 and Xin et al., *American J. Pathology*, 2001, 158(3): 1111-1120. Following gelation of the HUVEC cells, 1×basal medium containing M199 supplemented with 1% FBS, 1×ITS, 2 mM L-glutamine, 50 µg/ml ascorbic acid, 26.5 mM NaHCO₃, 100 U/ml penicillin and 100 U/ml streptomycin was added. Tube formation was elicited by the inclusion in the culture media of either a mixture of phorbol myristate acetate (50 nM), vascular endothelial cell growth factor (40 ng/ml) and basic fibroblast growth factor (40 ng/ml) ("PMA growth factor mix") or hepatocyte growth factor (40 ng/ml) and vascular endothelial cell growth factor (40 ng/ml) (HGF/VEGF mix) for the indicated period of time. Fibrin Gels were prepared by suspending Huvec (4×10⁵ cells/ml) in M199 containing 1% fetal bovine serum (Hyclone) and human fibrinogen (2.5 mg/ml). Thrombin (50 U/ml) was then added to the fibrinogen suspension at a ratio of 1 part thrombin solution:30 parts fibrinogen suspension. The solution was then layered onto 10 cm tissue culture plates (total volume: 15 ml/plate) and allowed to solidify at 37° C. for 20 min. Tissue culture media (10 ml of BM containing PMA (50 nM), bFGF (40 ng/ml) and VEGF (40 ng/ml)) was then added and the cells incubated at 37° C. in 5% CO₂ in air for the indicated period of time.

[0466] Total RNA was extracted from the HUVEC cells incubated for 0, 4, 8, 24, 40 and 50 hours in the different matrix and media combinations using a TRIzol extraction followed by a second purification using RNeasy Mini Kit (Qiagen). The total RNA was used to prepare cRNA which was then hybridized to the microarrays.

[0467] In the present experiments, nucleic acid probes derived from the herein described PRO polypeptide-encoding nucleic acid sequences were used in the creation of the microarray and RNA from the HUVEC cells described above were used for the hybridization thereto. Pairwise comparisons were made using time 0 chips as a baseline. Three replicate samples were analyzed for each experimental condition and time. Hence there were 3 time 0 samples for each treatment and 3 replicates of each successive time point. Therefore, a 3 by 3 comparison was performed for each time point compared against each time 0 point. This resulted in 9 comparisons per time point. Only those genes that had increased expression in all three non-time-0 replicates in each of the different matrix and media combinations as compared to any of the three time zero replicates were considered positive. Although this stringent method of data analysis does allow for false negatives, it minimizes false positives.

[0468] PRO281, PRO1560, PRO189, PRO4499, PRO6308, PRO6000, PRO10275, PRO21207, PRO20933, and PRO34274 tested positive in this assay.

Example 19

Tumor Versus Normal Differential Tissue Expression Distribution

[0469] Oligonucleotide probes were constructed from some of the PRO polypeptide-encoding nucleotide sequences shown in the accompanying figures for use in quantitative PCR amplification reactions. The oligonucleotide probes were chosen so as to give an approximately 200-600 base pair amplified fragment from the 3' end of its associated template in a standard PCR reaction. The oligonucleotide probes were employed in standard quantitative PCR amplification reactions with cDNA libraries isolated from different human tumor and normal human tissue samples and analyzed by agarose gel electrophoresis so as to obtain a quantitative determination of the level of expression of the PRO polypeptide-encoding nucleic acid in the various tumor and normal tissues tested. β-actin was used as a control to assure that equivalent amounts of nucleic acid was used in each reaction. Identification of the differential expression of the PRO polypeptide-encoding nucleic acid in one or more tumor tissues as compared to one or more normal tissues of the same tissue type renders the molecule useful diagnostically for the determination of the presence or absence of tumor in a subject suspected of possessing a tumor as well as therapeutically as a target for the treatment of a tumor in a subject possessing such a tumor. These assays provided the following results:

[0470] (1) DNA 161005-2943 molecule is very highly expressed in human umbilical vein endothelial cells (HUVEC), substantia nigra, hippocampus and dendrocytes; highly expressed in lymphoblasts; expressed in spleen, prostate, uterus and macrophages; and is weakly expressed in cartilage and heart. Among a panel of normal and tumor tissues examined, it is expressed in esophageal tumor, and is not expressed in normal esophagus, normal stomach, stomach tumor, normal kidney, kidney tumor, normal lung, lung tumor, normal rectum, rectal tumor, normal liver and liver tumor.

[0471] (2) DNA170245-3053 molecule is highly expressed in cartilage, testis, adrenal gland, and uterus, and not expressed in HUVEC, colon tumor, heart, placenta, bone marrow, spleen and aortic endothelial cells. In a panel of tumor and normal tissue samples examined, the DNA170245-3053 molecule was found to be expressed in normal esophagus and esophageal tumor, expressed in normal stomach and in stomach tumor, not expressed in normal kidney, but expressed in kidney tumor, not expressed in normal lung, but expressed in lung tumor, not expressed in normal rectum nor in rectal tumor, and not expressed in normal liver, but is expressed in liver tumor.

[0472] (3) DNA173157-2981 molecule is significantly expressed in the following tissues: cartilage, testis, HUVEC, heart, placenta, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells, and uterus. When these assays were conducted on a tumor tissue panel, it was found that the DNA 173157-2981 molecule is significantly expressed in the following tissues: normal esophagus and esophageal tumor, normal stomach and stomach tumor, normal kidney and kidney tumor, normal lung and lung tumor, normal rectum and rectal tumor, normal liver and liver tumor, and colon tumor.

[0473] (4) DNA175734-2985 molecule is significantly expressed in the adrenal gland and the uterus. The DNA 175734-2985 molecule is not significantly expressed in the following tissues: cartilage, testis, HUVEC, colon tumor, heart, placenta, bone marrow, prostate, spleen and aortic endothelial cells. Screening of a tumor panel revealed that DNA175734-2985 is significantly expressed in normal esophagus but not in esophageal tumor. Similarly, while highly expressed in normal rectum, DNA 175734-2985 is expressed to a lesser extent in rectal tumor. DNA 175734-2985 is expressed equally in normal stomach and stomach tumor as well as normal liver and liver tumor. While not expressed in normal kidney, DNA 175734-2985 is highly expressed in kidney tumor.

[0474] (5) DNA176108-3040 molecule is highly expressed in prostate and uterus, expressed in cartilage, testis, heart, placenta, bone marrow, adrenal gland and spleen, and not significantly expressed in HUVEC, colon tumor, and aortic endothelial cells. In a panel of tumor and normal tissue samples examined, the DNA 176108-3040 molecule was found to be highly expressed in normal esophagus, but expressed at lower levels in esophageal tumor, highly expressed in normal stomach, and expressed at a lower level in stomach tumor, expressed in kidney and in kidney tumor, expressed in normal rectum and at a lower level in rectal tumor, and expressed in normal liver and not expressed in liver tumor.

[0475] (6) DNA191064-3069 molecule is significantly expressed in the following tissues: cartilage, testis, HUVEC, heart, placenta, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells, and uterus and not significantly expressed in colon tumor. In a panel of tumor and normal tissue samples, the DNA191064-3069 molecule was found to be expressed in normal esophagus and in esophageal tumors, expressed in normal stomach and in stomach tumors, expressed in normal kidney and in kidney tumors, expressed in normal lung and in lung tumors, expressed in normal rectum and in rectal tumors, expressed in normal liver and in liver tumors.

[0476] (7) DNA 194909-3013 molecule is highly expressed in placenta, and expressed in cartilage, testis, HUVEC, colon tumor, heart, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells and uterus. In a panel of tumor and normal tissue samples examined, the DNA 194909-3013 molecule was found to be expressed in normal esophagus and expressed at a lower level in esophageal tumor, not expressed in normal stomach nor stomach tumor, expressed in normal kidney and kidney tumor, expressed in normal lung and lung tumor, expressed in normal rectum and rectal tumor, and not expressed in normal liver, but is expressed in liver tumor.

[0477] (8) The PRO34009 encoding genes of the invention (DNA203532-3029) were screened in normal tissues and the following primary tumors and the resulting values are reported below.

[0478] Tumor Panel:

[0479] PRO34009 encoding genes were expressed 39.3 fold higher in lung tumor than normal lung. It is expressed 9.5 fold higher in esophageal tumors than normal esophagus. It is expressed 6.7 fold higher in kidney tumor than normal kidney. It is expressed 4.0 fold higher in colon tumor than

normal colon. It is expressed 2.7 fold higher in stomach tumor than normal stomach. It is expressed at similar levels in normal rectum and rectal tumor, normal liver and liver tumor, normal uterus and uterine tumor.

[0480] Normal Panel:

[0481] For the normal tissue values, the normal tissue with the highest expression, in this case normal thymus, was given a value of 1 and all other normal tissues were given a value of less than 1 and described as expressed, weakly expressed or not expressed, based on their expression relative to thymus. PRO34009 encoding genes were expressed in normal thymus. It is weakly expressed in lymphoblast, spleen, heart, fetal limb, fetal lung, placenta, HUVEC, testis, fetal kidney, uterus, prostate, macrophage, substantia nigra, hippocampus, liver, skin, esophagus, stomach, rectum, kidney, thyroid, skeletal muscle, or fetal articular cartilage. It is not expressed in bone marrow, fetal liver, colon, lung or dendrocytes.

[0482] (9) DNA213858-3060 molecule is not significantly expressed in cartilage, testis, HUVEC, colon tumor, heart, placenta, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells or uterus. In a panel of tumor and normal tissue samples examined, the DNA213858-3060 molecule was found to be expressed in normal esophagus and esophageal tumor, expressed in normal stomach and in stomach tumor, expressed in normal kidney and in kidney tumor, expressed in normal lung and in lung tumor, expressed in normal rectum and in rectal tumor, and expressed in normal liver and in liver tumor.

[0483] (10) DNA216676-3083 molecule is significantly expressed in the following tissues: testis, heart, bone marrow, and uterus, and not significantly expressed in the following tissues: cartilage, HUVEC, colon tumor, placenta, adrenal gland, prostate, spleen, or aortic endothelial cells. In a panel of tumor and normal tissues samples examined, the DNA216676-3083 molecule was found to be expressed in normal esophagus and esophageal tumor, not expressed in normal stomach, but is expressed in stomach tumor, not expressed in normal kidney nor in kidney tumor, not expressed in normal lung, but is expressed in lung tumor, not expressed in normal rectum, but is expressed in rectal tumor, and not expressed in normal liver nor in liver tumor.

[0484] (11) DNA222653-3104 molecule is significantly expressed testis, and not significantly expressed in cartilage, HUVEC, colon tumor, heart, placenta, bone marrow, adrenal gland, prostate, spleen, aortic endothelial cells and uterus. In a panel of tumor and normal tissue samples examined, the DNA222653-3104 molecule was not expressed in normal esophagus, esophageal tumor, normal stomach, stomach tumor, normal kidney, kidney tumor, normal lung, lung tumor, normal rectum, rectal tumor, normal liver and liver tumor.

Example 20

Guinea Pig Vascular Leak (Assay 51)

[0485] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce vascular permeability. Polypeptides testing positive in this assay are expected to be useful for the therapeutic treatment of conditions which would benefit from enhanced

vascular permeability including, for example, conditions which may benefit from enhanced local immune system cell infiltration.

[0486] Hairless guinea pigs weighing 350 grams or more were anesthetized with Ketamine (75-80 mg/kg) and 5 mg/kg Xylazine intramuscularly. Test samples containing the PRO polypeptide or a physiological buffer without the test polypeptide are injected into skin on the back of the test animals with 100 μ l per injection site intradermally. There were approximately 16-24 injection sites per animal. One ml of Evans blue dye (1% in PBS) is then injected intracardially. Skin vascular permeability responses to the compounds (i.e., blemishes at the injection sites of injection) are visually scored by measuring the diameter (in mm) of blue-colored leaks from the site of injection at 1 and 6 hours post administration of the test materials. The mm diameter of blueness at the site of injection is observed and recorded as well as the severity of the vascular leakage. Blemishes of at least 5 mm in diameter are considered positive for the assay when testing purified proteins, being indicative of the ability to induce vascular leakage or permeability. A response greater than 7 mm diameter is considered positive for conditioned media samples. Human VEGF at 0.1 μ g/100 μ l is used as a positive control, inducing a response of 15-23 mm diameter.

[0487] PRO19822 polypeptides tested positive in this assay.

Example 21

Skin Vascular Permeability Assay (Assay 64)

[0488] This assay shows that certain polypeptides of the invention stimulate an immune response and induce inflammation by inducing mononuclear cell, eosinophil and PMN infiltration at the site of injection of the animal. Compounds which stimulate an immune response are useful therapeutically where stimulation of an immune response is beneficial. This skin vascular permeability assay is conducted as follows. Hairless guinea pigs weighing 350 grams or more are anesthetized with ketamine (75-80 mg/Kg) and 5 mg/Kg xylazine intramuscularly (IM). A sample of purified polypeptide of the invention or a conditioned media test sample is injected intradermally onto the backs of the test animals with 100 μ l per injection site. It is possible to have about 10-30, preferably about 16-24, injection sites per animal. One μ l of Evans blue dye (1% in physiologic buffered saline) is injected intracardially. Blemishes at the injection sites are then measured (mm diameter) at 1 hr and 6 hr post injection. Animals were sacrificed at 6 hrs after injection. Each skin injection site is biopsied and fixed in formalin. The skins are then prepared for histopathologic evaluation. Each site is evaluated for inflammatory cell infiltration into the skin. Sites with visible inflammatory cell inflammation are scored as positive. Inflammatory cells may be neutrophilic, eosinophilic, monocytic or lymphocytic. At least a minimal perivascular infiltrate at the injection site is scored as positive, no infiltrate at the site of injection is scored as negative.

[0489] PRO19822 polypeptide tested positive in this assay.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 116

<210> SEQ ID NO 1
<211> LENGTH: 1943
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 1

cggacgcgtg ggtgcgaggc gaaggtgacc ggggaccgag catttcagat	50
ctgctcggtg gacctggtgc accaccacca tgttggtgctgc aaggctggtg	100
tgtctccgga cactaccttc tagggttttc caccagctt tcaccaaggc	150
ctccctgtt gtgaagaatt ccatcacgaa gaatcaatgg ctgttaacac	200
ctagcaggga atatgccacc aaaacaagaa ttgggatccg gcgtgggaga	250
actggccaag aactcaaaga ggcagcattg gaaccatcga tggaaaaaat	300
atttaaaatt gatcagatgg gaagatggtt tgttgctgga ggggtgctg	350
ttggtcttgg agcattgtgc tactatggct tgggactgtc taatgagatt	400
ggagctattg aaaaggctgt aatttggcct cagtatgtca aggatagaat	450
tcattccacc tatatgtact tagcagggag tattggttta acagctttgt	500

-continued

ctgccatagc aatcagcaga acgcctgttc tcatgaactt catgatgaga	550
ggctcttggg tgacaattgg tgtgaccttt gcagccatgg ttggagctgg	600
aatgctggta cgatcaatac catatgacca gagcccaggc ccaaagcatc	650
ttgcttgggt gctacattct ggtgtgatgg gtgcagtggg ggctcctctg	700
acaatattag ggggtcctct tctcatcaga gctgcatggg acacagctgg	750
cattgtggga ggctctcca ctgtggccat gtgtgcgccc agtgaaaagt	800
ttctgaacat ggggtcaccc ctgggagtgg gcctgggtct cgtctttgtg	850
tctctattgg gatctatggt tcttccacct accaccgtgg ctggtgccac	900
tctttactca gtggcaatgt acggtggatt agttcttttc agcatgttcc	950
ttctgtatga taccagaaa gtaatcaagc gtgcagaagt atcaccaatg	1000
tatggagttc aaaaatatga tcccattaac tcgatgctga gtatctacat	1050
ggatacatta aatatattta tgcgagttgc aactatgctg gcaactggag	1100
gcaacagaaa gaaatgaagt gactcagctt ctggcttctc tgctacatca	1150
aatatcttgt ttaatggggc agatatgcat taaatagttt gtacaagcag	1200
ctttcgttga agtttagaag ataagaaaca tgtcatcata tttaaatggt	1250
ccggaatgtg gatgcctcag gtctgccttt tttctggag aataaatgca	1300
gtaatcctct cccaataag cacacacatt ttcaattctc atgtttgagt	1350
gattttaaaa tgttttggtg aatgtgaaaa ctaaagtttg tgtcatgaga	1400
atgtaagtct ttttctact ttaaaattta gtaggttcac tgagtaacta	1450
aaatttagca aacctgtgtt tgcataattt tttggagtgc agaatttgt	1500
aattaatgtc ataagtgatt tggagctttg gtaaaggagc cagagagaag	1550
gagtcacctg cagtcttttg tttttttaa tacttagaac ttagcacttg	1600
tgttattgat tagtgaggag ccagtaagaa acatctgggt atttggaac	1650
aagtggatcat tgttacattc atttgctgaa cttaacaaaa ctgttcatcc	1700
tgaacacaggc acaggtgatg cattctcctg ctgttgcttc tcagtgcctc	1750
ctttccaata tagatgtggt catgtttgac ttgtacagaa tgttaatcat	1800
acagagaatc cttgatggaa ttatatatgt gtgttttact tttgaatgtt	1850
acaaaaggaa ataactttaa aactattctc aagagaaaat attcaaagca	1900
tgaaatatgt tgctttttcc agaatacaaa cagtatactc atg	1943

<210> SEQ ID NO 2

<211> LENGTH: 345

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 2

Met	Leu	Ala	Ala	Arg	Leu	Val	Cys	Leu	Arg	Thr	Leu	Pro	Ser	Arg
1				5					10					15
Val	Phe	His	Pro	Ala	Phe	Thr	Lys	Ala	Ser	Pro	Val	Val	Lys	Asn
			20						25					30
Ser	Ile	Thr	Lys	Asn	Gln	Trp	Leu	Leu	Thr	Pro	Ser	Arg	Glu	Tyr
			35						40					45
Ala	Thr	Lys	Thr	Arg	Ile	Gly	Ile	Arg	Arg	Gly	Arg	Thr	Gly	Gln

-continued

	50		55		60
Glu Leu Lys Glu Ala Ala Leu Glu Pro Ser Met Glu Lys Ile Phe	65		70		75
Lys Ile Asp Gln Met Gly Arg Trp Phe Val Ala Gly Gly Ala Ala	80		85		90
Val Gly Leu Gly Ala Leu Cys Tyr Tyr Gly Leu Gly Leu Ser Asn	95		100		105
Glu Ile Gly Ala Ile Glu Lys Ala Val Ile Trp Pro Gln Tyr Val	110		115		120
Lys Asp Arg Ile His Ser Thr Tyr Met Tyr Leu Ala Gly Ser Ile	125		130		135
Gly Leu Thr Ala Leu Ser Ala Ile Ala Ile Ser Arg Thr Pro Val	140		145		150
Leu Met Asn Phe Met Met Arg Gly Ser Trp Val Thr Ile Gly Val	155		160		165
Thr Phe Ala Ala Met Val Gly Ala Gly Met Leu Val Arg Ser Ile	170		175		180
Pro Tyr Asp Gln Ser Pro Gly Pro Lys His Leu Ala Trp Leu Leu	185		190		195
His Ser Gly Val Met Gly Ala Val Val Ala Pro Leu Thr Ile Leu	200		205		210
Gly Gly Pro Leu Leu Ile Arg Ala Ala Trp Tyr Thr Ala Gly Ile	215		220		225
Val Gly Gly Leu Ser Thr Val Ala Met Cys Ala Pro Ser Glu Lys	230		235		240
Phe Leu Asn Met Gly Ala Pro Leu Gly Val Gly Leu Gly Leu Val	245		250		255
Phe Val Ser Ser Leu Gly Ser Met Phe Leu Pro Pro Thr Thr Val	260		265		270
Ala Gly Ala Thr Leu Tyr Ser Val Ala Met Tyr Gly Gly Leu Val	275		280		285
Leu Phe Ser Met Phe Leu Leu Tyr Asp Thr Gln Lys Val Ile Lys	290		295		300
Arg Ala Glu Val Ser Pro Met Tyr Gly Val Gln Lys Tyr Asp Pro	305		310		315
Ile Asn Ser Met Leu Ser Ile Tyr Met Asp Thr Leu Asn Ile Phe	320		325		330
Met Arg Val Ala Thr Met Leu Ala Thr Gly Gly Asn Arg Lys Lys	335		340		345

<210> SEQ ID NO 3
<211> LENGTH: 1110
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 3

ccaatcgccc ggtgcggtgg tgcagggtct cgggctagtc atggcgctccc	50
cgtctcggag actgcagact aaaccagtca ttacttgttt caagagcgtt	100
ctgctaattct acacttttat tttctggatc actggcggtta tccttcttgc	150
agttggcatt tggggcaagg tgagcctgga gaattacttt tctcttttaa	200
atgagaaggc caccaatgtc cccttcgtgc tcattgctac tggtagcgtc	250

-continued

attattcttt tgggcacctt tggttgtttt gctacctgcc gagcttctgc	300
atgggatgcta aaactgtatg caatgtttct gactctcggt tttttggtcg	350
aactggtcgc tgccatcgta ggatttgttt tcagacatga gattaagaac	400
agctttaaga ataattatga gaaggctttg aagcagtata actctacagg	450
agattataga agccatgcag tagacaagat ccaaaatacg ttgcattgtt	500
gtgggtgtcac cgattataga gattggacag atactaatta ttactcagaa	550
aaaggatttc ctaagagttg ctgtaaactt gaagattgta ctccacagag	600
agatgcagac aaagtaaaca atgaaggttg ttttataaag gtgatgacca	650
ttatagagtc agaaatggga gtcgtgacag gaatttcctt tggagttgct	700
tgcttccaac tgattggaat ctttctcgcc tactgccwct ctcgtgccat	750
aacaaataac cagtatgaga tagtgtaacc caatgtatct gtgggcctat	800
tcctctctac cttaaggac atttagggto cccctgtga attagaaagt	850
tgcttggtcg gagaactgac aacactactt actgatagac caaaaaacta	900
caccagtagg ttgattcaat caagatgtat gtagacctaa aactacacca	950
ataggctgat tcaatcaaga tccgtgctcg cagtgggctg attcaatcaa	1000
gatgtatgtt tgctatgttc taagtcacc ttctatccca ttcatgtag	1050
atcgttgaaa ccctgtatcc ctctgaaaca ctggaagagc tagtaaattg	1100
taaatgaagt	1110

<210> SEQ ID NO 4
<211> LENGTH: 245
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: 233
<223> OTHER INFORMATION: unknown amino acid

<400> SEQUENCE: 4

Met Ala Ser Pro Ser Arg Arg Leu Gln Thr Lys Pro Val Ile Thr	
1 5 10 15	
Cys Phe Lys Ser Val Leu Leu Ile Tyr Thr Phe Ile Phe Trp Ile	
20 25 30	
Thr Gly Val Ile Leu Leu Ala Val Gly Ile Trp Gly Lys Val Ser	
35 40 45	
Leu Glu Asn Tyr Phe Ser Leu Leu Asn Glu Lys Ala Thr Asn Val	
50 55 60	
Pro Phe Val Leu Ile Ala Thr Gly Thr Val Ile Ile Leu Leu Gly	
65 70 75	
Thr Phe Gly Cys Phe Ala Thr Cys Arg Ala Ser Ala Trp Met Leu	
80 85 90	
Lys Leu Tyr Ala Met Phe Leu Thr Leu Val Phe Leu Val Glu Leu	
95 100 105	
Val Ala Ala Ile Val Gly Phe Val Phe Arg His Glu Ile Lys Asn	
110 115 120	
Ser Phe Lys Asn Asn Tyr Glu Lys Ala Leu Lys Gln Tyr Asn Ser	
125 130 135	

-continued

Thr	Gly	Asp	Tyr	Arg	Ser	His	Ala	Val	Asp	Lys	Ile	Gln	Asn	Thr
				140					145				150	
Leu	His	Cys	Cys	Gly	Val	Thr	Asp	Tyr	Arg	Asp	Trp	Thr	Asp	Thr
				155					160				165	
Asn	Tyr	Tyr	Ser	Glu	Lys	Gly	Phe	Pro	Lys	Ser	Cys	Cys	Lys	Leu
				170					175				180	
Glu	Asp	Cys	Thr	Pro	Gln	Arg	Asp	Ala	Asp	Lys	Val	Asn	Asn	Glu
				185					190					195
Gly	Cys	Phe	Ile	Lys	Val	Met	Thr	Ile	Ile	Glu	Ser	Glu	Met	Gly
				200					205					210
Val	Val	Ala	Gly	Ile	Ser	Phe	Gly	Val	Ala	Cys	Phe	Gln	Leu	Ile
				215					220					225
Gly	Ile	Phe	Leu	Ala	Tyr	Cys	Xaa	Ser	Arg	Ala	Ile	Thr	Asn	Asn
				230					235					240
Gln	Tyr	Glu	Ile	Val										
				245										

<210> SEQ ID NO 5
<211> LENGTH: 1373
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 5

ggggccgcgg tctagggcgg ctacgtgtgt tgccatagcg accattttgc	50
attaactggt tggtagcttc tatcctgggg gctgagcgac tgcgggccag	100
ctcttccctc actccctctc ggctccttgt ggcccaaagg cctaaccggg	150
gtccggcggg ctggcctagg gatcttcccc gttgccctt tggggcggga	200
tggtgcgga agaagaagac gaggtggagt gggtagtgga gagcatcgcg	250
gggttcctcg gaggccaga ctggctccatc cccatcttg acttttgga	300
acagaaatgt gaagttaact gcaaaggagg gcatgtgata actccaggaa	350
gcccagagcc ggtgattttg gtggcctgtg ttccccttgt tttgatgat	400
gaagaagaaa gcaaatgac ctatacagag attcatcagg aatacaaaa	450
actagttaa aagctgttag aaggttacct caaagaaatt ggaattaatg	500
aagatcaatt tcaagaagca tgcacttctc ctcttgcaa gaccataca	550
tcacaggcca ttttgaacc tgtgttgga gcagaagatt ttactatctt	600
taaagcaatg atgtccaga aaaacattga aatgcagctg caagccattc	650
gaataattca agagagaaat ggtgtattac ctgactgctt aaccgatggc	700
tctgatgtgg tcagtgcct tgaacacgaa gagatgaaa tcctgaggga	750
agttcttaga aaatcaaaag aggaatatga ccaggaagaa gaaaggaaga	800
ggaaaaaaca gttatcagag gctaaaacag aagagccac agtgcattcc	850
agtgaagctg caataatgaa taattcccaa ggggatggtg aacattttgc	900
acacccacc tcagaagtta aaatgcattt tgctaatacag tcaatagaac	950
ctttgggaag aaaagtggaa aggtctgaaa cttcctccct cccacaaaa	1000
ggcctaaga ttcttgctt agagcatgag agcattgaag gaccaatagc	1050
aaacttatca gtacttggaaga cagaagaact tcggcaacga gaacactatc	1100

-continued

tcaagcagaa gagagataag ttgatgtcca tgagaaagga tatgaggact	1150
aaacagatac aaaatatgga gcagaaagga aaaccctctg gggaggtaga	1200
ggaaatgaca gagaaccag aaatgacagc agaggagaag caaacattac	1250
taaagaggag attgcttgca gagaaactca aagaagaagt tattaataag	1300
taataattaa gaacaattta acaaaatgga agttcaaatt gtcttaaaaa	1350
taaattatatt agtccttaca ctg	1373

<210> SEQ ID NO 6

<211> LENGTH: 367

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 6

Met Ala Ala Glu Glu Glu Asp Glu Val Glu Trp Val Val Glu Ser	
1 5 10 15	
Ile Ala Gly Phe Leu Arg Gly Pro Asp Trp Ser Ile Pro Ile Leu	
20 25 30	
Asp Phe Val Glu Gln Lys Cys Glu Val Asn Cys Lys Gly Gly His	
35 40 45	
Val Ile Thr Pro Gly Ser Pro Glu Pro Val Ile Leu Val Ala Cys	
50 55 60	
Val Pro Leu Val Phe Asp Asp Glu Glu Glu Ser Lys Leu Thr Tyr	
65 70 75	
Thr Glu Ile His Gln Glu Tyr Lys Glu Leu Val Glu Lys Leu Leu	
80 85 90	
Glu Gly Tyr Leu Lys Glu Ile Gly Ile Asn Glu Asp Gln Phe Gln	
95 100 105	
Glu Ala Cys Thr Ser Pro Leu Ala Lys Thr His Thr Ser Gln Ala	
110 115 120	
Ile Leu Gln Pro Val Leu Ala Ala Glu Asp Phe Thr Ile Phe Lys	
125 130 135	
Ala Met Met Val Gln Lys Asn Ile Glu Met Gln Leu Gln Ala Ile	
140 145 150	
Arg Ile Ile Gln Glu Arg Asn Gly Val Leu Pro Asp Cys Leu Thr	
155 160 165	
Asp Gly Ser Asp Val Val Ser Asp Leu Glu His Glu Glu Met Lys	
170 175 180	
Ile Leu Arg Glu Val Leu Arg Lys Ser Lys Glu Glu Tyr Asp Gln	
185 190 195	
Glu Glu Glu Arg Lys Arg Lys Lys Gln Leu Ser Glu Ala Lys Thr	
200 205 210	
Glu Glu Pro Thr Val His Ser Ser Glu Ala Ala Ile Met Asn Asn	
215 220 225	
Ser Gln Gly Asp Gly Glu His Phe Ala His Pro Pro Ser Glu Val	
230 235 240	
Lys Met His Phe Ala Asn Gln Ser Ile Glu Pro Leu Gly Arg Lys	
245 250 255	
Val Glu Arg Ser Glu Thr Ser Ser Leu Pro Gln Lys Gly Leu Lys	
260 265 270	
Ile Pro Gly Leu Glu His Ala Ser Ile Glu Gly Pro Ile Ala Asn	
275 280 285	

-continued

Leu	Ser	Val	Leu	Gly	Thr	Glu	Glu	Leu	Arg	Gln	Arg	Glu	His	Tyr					
									290						295				300
Leu	Lys	Gln	Lys	Arg	Asp	Lys	Leu	Met	Ser	Met	Arg	Lys	Asp	Met					
									305						310				315
Arg	Thr	Lys	Gln	Ile	Gln	Asn	Met	Glu	Gln	Lys	Gly	Lys	Pro	Thr					
									320						325				330
Gly	Glu	Val	Glu	Glu	Met	Thr	Glu	Lys	Pro	Glu	Met	Thr	Ala	Glu					
									335						340				345
Glu	Lys	Gln	Thr	Leu	Leu	Lys	Arg	Arg	Leu	Leu	Ala	Glu	Lys	Leu					
									350						355				360
Lys	Glu	Glu	Val	Ile	Asn	Lys													
														365					

<210> SEQ ID NO 7
<211> LENGTH: 932
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: 911
<223> OTHER INFORMATION: unknown base

<400> SEQUENCE: 7

gggaacggaa aatggcgccct cacggcccgg gtagtcttac gacctggtg	50
ccctgggctg ccgccctgct cctcgctctg ggcgtggaaa gggctctggc	100
gctacccgag atatgcaccc aatgtccagg gagcgtgcaa aatttgtaaa	150
aagtggcctt ttattgtaaa acgacacgag agctaagtct gcatgcccgt	200
tgctgcctga atcagaaggg caccatcttg gggctggatc tccagaactg	250
ttctctggag gacctgggtc caaactttca tcaggcacat accactgtca	300
tcatagacct gcaagcaaac cccctcaaag gtgacttggc caacaccttc	350
cgtaggcttta ctacgtcca gactctgata ctgccacaac atgtcaactg	400
tcctggagga attaatgcct ggaatactat cacctcttat atagacaacc	450
aaatctgtca agggcaaaag aacctttgca ataacactgg ggacccagaa	500
atgtgtcctg agaatggatc ttgtgtacct gatggtccag gtcttttgca	550
gtgtgtttgt gctgatggtt tccatggata caagtgtatg cgccagggct	600
cgttctcact gcttatgttc ttcgggatcc tgggagccac cactctatcc	650
gtctccattc tgctttgggc gaccagcgc cgaaaagcca agacttcatg	700
aactacatag gtcttaccat tgacctaga tcaatctgaa ctatcttagc	750
ccagtcaggg agctctgctt cctagaaagg catctttcgc cagtggatcc	800
gcctcaaggt tgaggccgcc attggaagat gaaaaattgc actcccttgg	850
tgtagacaaa taccagttcc cattggtgtt gttgcctata ataaacactt	900
tttctttttt naaaaaaaaa aaaaaaaaaa aa	932

<210> SEQ ID NO 8
<211> LENGTH: 229
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
<400> SEQUENCE: 8

-continued

Met	Ala	Pro	His	Gly	Pro	Gly	Ser	Leu	Thr	Thr	Leu	Val	Pro	Trp
1				5					10					15
Ala	Ala	Ala	Leu	Leu	Leu	Ala	Leu	Gly	Val	Glu	Arg	Ala	Leu	Ala
			20						25					30
Leu	Pro	Glu	Ile	Cys	Thr	Gln	Cys	Pro	Gly	Ser	Val	Gln	Asn	Leu
			35						40					45
Ser	Lys	Val	Ala	Phe	Tyr	Cys	Lys	Thr	Thr	Arg	Glu	Leu	Met	Leu
			50						55					60
His	Ala	Arg	Cys	Cys	Leu	Asn	Gln	Lys	Gly	Thr	Ile	Leu	Gly	Leu
			65						70					75
Asp	Leu	Gln	Asn	Cys	Ser	Leu	Glu	Asp	Pro	Gly	Pro	Asn	Phe	His
			80						85					90
Gln	Ala	His	Thr	Thr	Val	Ile	Ile	Asp	Leu	Gln	Ala	Asn	Pro	Leu
			95						100					105
Lys	Gly	Asp	Leu	Ala	Asn	Thr	Phe	Arg	Gly	Phe	Thr	Gln	Leu	Gln
			110						115					120
Thr	Leu	Ile	Leu	Pro	Gln	His	Val	Asn	Cys	Pro	Gly	Gly	Ile	Asn
			125						130					135
Ala	Trp	Asn	Thr	Ile	Thr	Ser	Tyr	Ile	Asp	Asn	Gln	Ile	Cys	Gln
			140						145					150
Gly	Gln	Lys	Asn	Leu	Cys	Asn	Asn	Thr	Gly	Asp	Pro	Glu	Met	Cys
			155						160					165
Pro	Glu	Asn	Gly	Ser	Cys	Val	Pro	Asp	Gly	Pro	Gly	Leu	Leu	Gln
			170						175					180
Cys	Val	Cys	Ala	Asp	Gly	Phe	His	Gly	Tyr	Lys	Cys	Met	Arg	Gln
			185						190					195
Gly	Ser	Phe	Ser	Leu	Leu	Met	Phe	Phe	Gly	Ile	Leu	Gly	Ala	Thr
			200						205					210
Thr	Leu	Ser	Val	Ser	Ile	Leu	Leu	Trp	Ala	Thr	Gln	Arg	Arg	Lys
			215						220					225
Ala	Lys	Thr	Ser											

<210> SEQ ID NO 9
<211> LENGTH: 2482
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 9

gggggagaag gcgcccgagc cccagctctc cgagcaccgg gtcggaagcc	50
gcgacccgag ccgcgcagga agctgggacc ggaacctcgg cggaccgggc	100
cccacccaac tcacctgcgc aggtcaccag caccctcgga acccagaggc	150
ccgcgctctg aagtgaccc ccctggggag gaaggcgatg gccctgcga	200
ggacgatggc ccgcgcccgc ctgccccgg ccggcatccc tgccgtcgcc	250
ttgtggcttc tgtgcacgt cggcctccag ggcacccagg ccggggccacc	300
gcccgcgcc cctgggctgc ccgcgggagc cgactgcctg aacagcttta	350
ccgcgggggt gcctggcttc gtgtggaca ccaacgcctc ggtcagcaac	400
ggagctacct tcctggagtc ccccaccgtg cgccggggct gggactgcgt	450
gcgcgcctgc tgcaccaccc agaactgcaa cttggcgcta gtggagctgc	500

-continued

agccccgaccg	cggggaggac	gccatcgccg	cctgcttcct	catcaactgc	550
ctctacgagc	agaacttcgt	gtgcaagttc	gcgcccaggg	agggcttcat	600
caactacctc	acgagggaag	tgtaccgctc	ctaccgccag	ctgcggaccc	650
agggctttgg	agggctctggg	atccccaagg	cctgggcagg	catagacttg	700
aaggtacaac	cccaggaacc	cctggtgctg	aaggatgtgg	aaaacacaga	750
ttggcgcta	ctgcgggggtg	acacggatgt	cagggtagag	aggaaagacc	800
caaaccagg	ggaactgtgg	ggactcaagg	aaggcaccta	cctgttcag	850
ctgacagtga	ctagctcaga	ccaccagag	gacacggcca	acgtcacagt	900
cactgtgctg	tccaccaagc	agacagaaga	ctactgcctc	gcatccaaca	950
aggtgggtcg	ctgcgggggc	tctttccac	gctggtacta	tgacccacg	1000
gagcagatct	gcaagagttt	cgtttatgga	ggctgcttgg	gcaacaagaa	1050
caactacctt	cggaagaag	agtgcattct	agcctgtcgg	ggtgtgcaag	1100
gtgggccttt	gagaggcagc	tctggggctc	aggcgacttt	ccccagggc	1150
ccctccatgg	aaaggcgcca	tccagtgtgc	tctggcacct	gtcagccac	1200
ccagttccgc	tgacgaatg	gctgctgcat	cgacagtttc	ctggagtgtg	1250
acgacacccc	caactgcccc	gacgcctccg	acgaggctgc	ctgtgaaaaa	1300
tacacgagtg	gctttgacga	gctccagcgc	atccatttcc	ccagtgaaca	1350
agggcactgc	gtggacctgc	cagacacagg	actctgcaag	gagagcatcc	1400
cgcgctggta	ctacaacccc	ttagcgaac	actgcgcccg	ctttacctat	1450
ggtggttgtt	atggcaacaa	gaacaacttt	gaggaagagc	agcagtgcc	1500
cgagtcttgt	cgcggcatct	ccaagaagga	tgtgtttggc	ctgaggcggg	1550
aaatccccat	tcccagcaca	ggctctgtgg	agatggctgt	cacagtgttc	1600
ctggtcacat	gcattgtggt	ggtggtagcc	atcttgggtt	actgcttctt	1650
caagaaccag	agaaaggact	tccacggaca	ccaccaccac	ccaccacca	1700
cccctgccag	ctccactgtc	tccactaccg	aggacacgga	gcacctggtc	1750
tataaccaca	ccaccgggcc	cctctgagcc	tgggtctcac	cggctctcac	1800
ctggccctgc	ttcctgcttg	ccaaggcaga	ggcctgggct	gggaaaaact	1850
ttggaaccag	actcttgcct	gtttccacgg	cccactgtgc	ctcagagacc	1900
agggctccag	cccctcttgg	agaagtctca	gctaagctca	cgtcctgaga	1950
aagctcaaag	gtttggaagg	agcagaaaac	ccttgggccca	gaagtaccag	2000
actagatgga	cctgcctgca	taggagtttg	gaggaagttg	gagttttgtt	2050
tcctctgttc	aaagctgcct	gtccctaccc	catggtgcta	ggaagaggag	2100
tggggtggtg	tcagaccctg	gaggcccaaa	ccctgtcctc	ccgagctcct	2150
cttccatgct	gtgcgccag	ggctggggag	aaggacttcc	ctgtgtagtt	2200
tgtgtgttaa	agagttgctt	tttgtttatt	taatgctgtg	gcatgggtga	2250
agaggagggg	aagaggcctg	tttggcctct	ctgtcctctc	ttcctcttcc	2300
cccaagattg	agctctctgc	ccttgatcag	ccccaccctg	gcctagacca	2350
gcagacagag	ccaggagagg	ctcagctgca	ttccgcagcc	cccaccccca	2400

-continued

aggttctcca acatcacagc ccagcccacc cactgggtaa taaaagtgtg 2450

ttgtggaaaa aaaaaaaaaa aaaaaaaaaa aa 2482

<210> SEQ ID NO 10

<211> LENGTH: 529

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 10

Met Ala Pro Ala Arg Thr Met Ala Arg Ala Arg Leu Ala Pro Ala
1 5 10 15

Gly Ile Pro Ala Val Ala Leu Trp Leu Leu Cys Thr Leu Gly Leu
20 25 30

Gln Gly Thr Gln Ala Gly Pro Pro Pro Ala Pro Pro Gly Leu Pro
35 40 45

Ala Gly Ala Asp Cys Leu Asn Ser Phe Thr Ala Gly Val Pro Gly
50 55 60

Phe Val Leu Asp Thr Asn Ala Ser Val Ser Asn Gly Ala Thr Phe
65 70 75

Leu Glu Ser Pro Thr Val Arg Arg Gly Trp Asp Cys Val Arg Ala
80 85 90

Cys Cys Thr Thr Gln Asn Cys Asn Leu Ala Leu Val Glu Leu Gln
95 100 105

Pro Asp Arg Gly Glu Asp Ala Ile Ala Ala Cys Phe Leu Ile Asn
110 115 120

Cys Leu Tyr Glu Gln Asn Phe Val Cys Lys Phe Ala Pro Arg Glu
125 130 135

Gly Phe Ile Asn Tyr Leu Thr Arg Glu Val Tyr Arg Ser Tyr Arg
140 145 150

Gln Leu Arg Thr Gln Gly Phe Gly Gly Ser Gly Ile Pro Lys Ala
155 160 165

Trp Ala Gly Ile Asp Leu Lys Val Gln Pro Gln Glu Pro Leu Val
170 175 180

Leu Lys Asp Val Glu Asn Thr Asp Trp Arg Leu Leu Arg Gly Asp
185 190 195

Thr Asp Val Arg Val Glu Arg Lys Asp Pro Asn Gln Val Glu Leu
200 205 210

Trp Gly Leu Lys Glu Gly Thr Tyr Leu Phe Gln Leu Thr Val Thr
215 220 225

Ser Ser Asp His Pro Glu Asp Thr Ala Asn Val Thr Val Thr Val
230 235 240

Leu Ser Thr Lys Gln Thr Glu Asp Tyr Cys Leu Ala Ser Asn Lys
245 250 255

Val Gly Arg Cys Arg Gly Ser Phe Pro Arg Trp Tyr Tyr Asp Pro
260 265 270

Thr Glu Gln Ile Cys Lys Ser Phe Val Tyr Gly Gly Cys Leu Gly
275 280 285

Asn Lys Asn Asn Tyr Leu Arg Glu Glu Glu Cys Ile Leu Ala Cys
290 295 300

Arg Gly Val Gln Gly Gly Pro Leu Arg Gly Ser Ser Gly Ala Gln
305 310 315

Ala Thr Phe Pro Gln Gly Pro Ser Met Glu Arg Arg His Pro Val

-continued

320					325					330				
Cys	Ser	Gly	Thr	Cys	Gln	Pro	Thr	Gln	Phe	Arg	Cys	Ser	Asn	Gly
335					340					345				
Cys	Cys	Ile	Asp	Ser	Phe	Leu	Glu	Cys	Asp	Asp	Thr	Pro	Asn	Cys
350					355					360				
Pro	Asp	Ala	Ser	Asp	Glu	Ala	Ala	Cys	Glu	Lys	Tyr	Thr	Ser	Gly
365					370					375				
Phe	Asp	Glu	Leu	Gln	Arg	Ile	His	Phe	Pro	Ser	Asp	Lys	Gly	His
380					385					390				
Cys	Val	Asp	Leu	Pro	Asp	Thr	Gly	Leu	Cys	Lys	Glu	Ser	Ile	Pro
395					400					405				
Arg	Trp	Tyr	Tyr	Asn	Pro	Phe	Ser	Glu	His	Cys	Ala	Arg	Phe	Thr
410					415					420				
Tyr	Gly	Gly	Cys	Tyr	Gly	Asn	Lys	Asn	Asn	Phe	Glu	Glu	Glu	Gln
425					430					435				
Gln	Cys	Leu	Glu	Ser	Cys	Arg	Gly	Ile	Ser	Lys	Lys	Asp	Val	Phe
440					445					450				
Gly	Leu	Arg	Arg	Glu	Ile	Pro	Ile	Pro	Ser	Thr	Gly	Ser	Val	Glu
455					460					465				
Met	Ala	Val	Thr	Val	Phe	Leu	Val	Ile	Cys	Ile	Val	Val	Val	Val
470					475					480				
Ala	Ile	Leu	Gly	Tyr	Cys	Phe	Phe	Lys	Asn	Gln	Arg	Lys	Asp	Phe
485					490					495				
His	Gly	His	His	His	His	Pro	Pro	Pro	Thr	Pro	Ala	Ser	Ser	Thr
500					505					510				
Val	Ser	Thr	Thr	Glu	Asp	Thr	Glu	His	Leu	Val	Tyr	Asn	His	Thr
515					520					525				
Thr Arg Pro Leu														

<210> SEQ ID NO 11
<211> LENGTH: 1899
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 11

gtgctgggct ttttcagaca agtgcattct ctaaccaggt cacatttcag	50
ccgcgaccca ctctccgccca gtcaccggag gcagaccgcg ggaggagagc	100
tgaggacagc cgcgtgcgct tcgccagcag cgggggtggga ggaaggacat	150
taaaatactg cagaagtcaa gaccccccca ggtcgaacct agaccacgat	200
gcgcgccccg ggctgcgggc ggctggtgct gccgctgctg ctctgggccg	250
cggcagccct ggccgaaggc gacgccaagg ggctcaagga ggcgagagcc	300
cccggcaatt tcatggagga cgagcaatgg ctgtcgtcca tctcgagta	350
cagcggcaag atcaagcact ggaaccgctt ccgagacgaa gtggaggatg	400
actatatcaa gagctgggag gacaatcagc aaggagatga agccctggat	450
accaccaagg acccctgccca gaagtggaag tgcagccgcc acaaggtgtg	500
cattgccagc ggctaccagc gggccatgtg catcagtcgc aagaagctgg	550
agcacaggat caagcagccg accgtgaaac tccatggaaa caaagactcc	600
atctgcaagc cctgccacat ggcccagctt gcctctgtct gcggctcaga	650

-continued

tggccacact tacagctctg tgtgtaagct ggagcaacag gcgtgcctga 700
gcagcaagca gctggcggtg cgatgcgagg gccctgccc ctgccccacg 750
gagcaggctg ccacctccac cgccgatggc aaaccagaga cttgcaccgg 800
tcaggacctg gctgacctgg gagatcggct gcgggactgg ttccagctcc 850
ttcatgagaa ctccaagcag aatggctcag ccagcagtgt agccggcccg 900
gccagcgggc tggacaagag cctggggggc agctgcaagg actccattgg 950
ctggatgttc tccaagctgg acaccagtgc tgacctcttc ctggaccaga 1000
cggagctggc cgccatcaac ctggacaagt acgaggtctg catccgtccc 1050
ttcttcaact cctgtgacac ctacaaggat ggccgggtct ctactgctga 1100
gtggtgcttc tgcttctgga gggagaagcc cccctgcctg gcagagctgg 1150
agcgcattcca gatccaggag gccgccaaga agaagccagg catcttcac 1200
ccgagctgcg acgaggatgg ctactaccgg aagatgcagt gtgaccagag 1250
cagcgggtgac tgctggcgtg tggaccagct gggcctggag ctgactggca 1300
cgcgcacgca tgggagcccc gactgcgatg acatcgtggg cttctcgggg 1350
gactttggaa gcggtgtcgg ctgggaggat gaggaggaga aggagacgga 1400
ggaagcaggc gaggaggccg aggaggagga gggcgaggca ggcgaggctg 1450
acgacggggg ctacatctgg tagacgccct caggagccgg ctgccggggg 1500
ggactcaaca gcagagctct gagcagcagc aggcaacttc gagaacggat 1550
ccagaaatgc agtcagaagg accctgctcc acctgggggg actgggagtg 1600
tgagtgtgca tggcatgtgt gtggcacaga tggctgggac ggtgacagt 1650
gtgagtgcac gtgtgcatgc atgtgtgtat gtgtgtgtgt gtgtggcatg 1700
cgctgacaaa tgtgtccttg atccacactg ctctggcag agtgagtcac 1750
ccaaaggccc ctcggcctc ctgtagctg ttttctttcc tttgttgtt 1800
ggttttaaaa tacattcaca cacaaataca aaaaaaaaaa aaaaaaaaaa 1850
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1899

<210> SEQ ID NO 12
<211> LENGTH: 424
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 12

Met Arg Ala Pro Gly Cys Gly Arg Leu Val Leu Pro Leu Leu Leu
1 5 10 15
Leu Ala Ala Ala Ala Leu Ala Glu Gly Asp Ala Lys Gly Leu Lys
20 25 30
Glu Gly Glu Thr Pro Gly Asn Phe Met Glu Asp Glu Gln Trp Leu
35 40 45
Ser Ser Ile Ser Gln Tyr Ser Gly Lys Ile Lys His Trp Asn Arg
50 55 60
Phe Arg Asp Glu Val Glu Asp Asp Tyr Ile Lys Ser Trp Glu Asp
65 70 75
Asn Gln Gln Gly Asp Glu Ala Leu Asp Thr Thr Lys Asp Pro Cys
80 85 90

-continued

Gln	Lys	Val	Lys	Cys	Ser	Arg	His	Lys	Val	Cys	Ile	Ala	Gln	Gly	95	100	105
Tyr	Gln	Arg	Ala	Met	Cys	Ile	Ser	Arg	Lys	Lys	Leu	Glu	His	Arg	110	115	120
Ile	Lys	Gln	Pro	Thr	Val	Lys	Leu	His	Gly	Asn	Lys	Asp	Ser	Ile	125	130	135
Cys	Lys	Pro	Cys	His	Met	Ala	Gln	Leu	Ala	Ser	Val	Cys	Gly	Ser	140	145	150
Asp	Gly	His	Thr	Tyr	Ser	Ser	Val	Cys	Lys	Leu	Glu	Gln	Gln	Ala	155	160	165
Cys	Leu	Ser	Ser	Lys	Gln	Leu	Ala	Val	Arg	Cys	Glu	Gly	Pro	Cys	170	175	180
Pro	Cys	Pro	Thr	Glu	Gln	Ala	Ala	Thr	Ser	Thr	Ala	Asp	Gly	Lys	185	190	195
Pro	Glu	Thr	Cys	Thr	Gly	Gln	Asp	Leu	Ala	Asp	Leu	Gly	Asp	Arg	200	205	210
Leu	Arg	Asp	Trp	Phe	Gln	Leu	Leu	His	Glu	Asn	Ser	Lys	Gln	Asn	215	220	225
Gly	Ser	Ala	Ser	Ser	Val	Ala	Gly	Pro	Ala	Ser	Gly	Leu	Asp	Lys	230	235	240
Ser	Leu	Gly	Ala	Ser	Cys	Lys	Asp	Ser	Ile	Gly	Trp	Met	Phe	Ser	245	250	255
Lys	Leu	Asp	Thr	Ser	Ala	Asp	Leu	Phe	Leu	Asp	Gln	Thr	Glu	Leu	260	265	270
Ala	Ala	Ile	Asn	Leu	Asp	Lys	Tyr	Glu	Val	Cys	Ile	Arg	Pro	Phe	275	280	285
Phe	Asn	Ser	Cys	Asp	Thr	Tyr	Lys	Asp	Gly	Arg	Val	Ser	Thr	Ala	290	295	300
Glu	Trp	Cys	Phe	Cys	Phe	Trp	Arg	Glu	Lys	Pro	Pro	Cys	Leu	Ala	305	310	315
Glu	Leu	Glu	Arg	Ile	Gln	Ile	Gln	Glu	Ala	Ala	Lys	Lys	Lys	Pro	320	325	330
Gly	Ile	Phe	Ile	Pro	Ser	Cys	Asp	Glu	Asp	Gly	Tyr	Tyr	Arg	Lys	335	340	345
Met	Gln	Cys	Asp	Gln	Ser	Ser	Gly	Asp	Cys	Trp	Arg	Val	Asp	Gln	350	355	360
Leu	Gly	Leu	Glu	Leu	Thr	Gly	Thr	Arg	Thr	His	Gly	Ser	Pro	Asp	365	370	375
Cys	Asp	Asp	Ile	Val	Gly	Phe	Ser	Gly	Asp	Phe	Gly	Ser	Gly	Val	380	385	390
Gly	Trp	Glu	Asp	Glu	Glu	Glu	Lys	Glu	Thr	Glu	Glu	Ala	Gly	Glu	395	400	405
Glu	Ala	Glu	Glu	Glu	Glu	Gly	Glu	Ala	Gly	Glu	Ala	Asp	Asp	Gly	410	415	420
Gly	Tyr	Ile	Trp														

<210> SEQ ID NO 13

<211> LENGTH: 2680

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 13

-continued

tgccgacgacc gtcgtacacc atgggcctcc acctccgccc ctaccgtgtg	50
gggctgctcc cggatggcct cctgttcctc ttgtctgtgc taatgtgtct	100
cgcggaccca gcgctcccg cggagcgtca cccccagtg gtgtgggtcc	150
ctggtgattt gggtaaccaa ctggaagcca agctggacaa gccgacagt	200
gtgcactacc tctgtccaa gaagaccgaa agctacttca caatctggct	250
gaacctggaa ctgtctgtgc ctgtcatcat tgactgtgtg attgacaata	300
tcaggctggt ttacaacaaa acatccaggg cccccagtt tcctgatggt	350
gtggatgtac gtgtccctgg ctttgggaag accttctcac tggagtccct	400
ggaccccagc aaaagcagcg tgggttccta tttccacacc atgggtggaga	450
gccttggtgg ctggggctac acacgggtg aggatgtccg aggggtccc	500
tatgactggc gccgagcccc aaatgaaac gggccctact tcctggccct	550
ccgcgagatg atcgaggaga tgtaccagt gtatgggggc cccgtggtgc	600
tggttgccca cagtatgggc aacatgtaca cgctctactt tctgcagcg	650
cagccgcagg cctggaagga caagtatac cgggccttcg tgtcaactgg	700
tgcgccctgg gggggcgtgg ccaagaccct gcgcgtcctg gcttcaggag	750
acaacaaccg gatcccagtc atcgggcccc tgaagatccg ggagcagcag	800
cggtcagctg tctccaccag ctggctgtgc ccctacaact acacatggtc	850
acctgagaag gtgttcgtgc agacaccac aatcaactac acactgcggg	900
actaccgcaa gttcttccag gacatcggct ttgaagatgg ctggctcatg	950
cggcaggaca cagaagggct ggtggaagcc acgatgccac ctggcgtgca	1000
gctgcactgc ctctatggta ctggcgtccc cacaccagac tccttctact	1050
atgagagctt ccctgaccgt gaccctaaaa tctgctttgg tgacggcgat	1100
ggtactgtga acttgaagag tgccctgcag tgccaggcct ggcagagccg	1150
ccaggagcac caagtgttgc tgcaggagct gccaggcagc gagcacatcg	1200
agatgctggc caacgccacc accctggcct atctgaaacg tgtgtccctt	1250
gggcctgac tcctgtgcca caggactcct gtggtcggc cgtggacctg	1300
ctgttggcct ctggggctgt catggcccac gcgttttgca aagtttgtga	1350
ctcaccattc aaggccccga gtcttggaact gtgaagcatc tgccatgggg	1400
aagtgtgtgt tgttatcctt tctctgtggc agtgaagaag gaagaaatga	1450
gagtctagac tcaagggaca ctggatggca agaagctgc tgatggtgga	1500
actgtgtgta ccttaggact ggctccacag ggtggactgg ctgggacctg	1550
gtcccagtc ctgcctgggg ccatgtgtcc ccctattcct gtgggctttt	1600
catacttgcc tactggggcc tggccccga gccttcctat gagggatgtt	1650
actgggctgt ggtcctgtac ccagaggtcc cagggatcgg ctccctggccc	1700
ctcgggtgac ccttcccaca caccagccac agataggcct gccactggtc	1750
atgggtagct agagctgctg gcttccctgt ggcttagctg gtggccagcc	1800
tgactggctt cctgggcgag cctagtagct cctgcaggca ggggcagttt	1850
gttgcggtct tcgtggttcc caggccctgg gacatctcac tccactccta	1900

-continued

cctcccttac caccaggagc attcaagctc tggattgggc agcagatgtg	1950
cccccagtc cgcaggctgt gttccagggg ccctgatttc ctcggatgtg	2000
ctattggccc caggactgaa gctgcctccc ttcaccctgg gactgtgggt	2050
ccaaggatga gagcaggggt tggagccatg gccttctggg aacctatgga	2100
gaaagggaa ccaaggaagc agccaaggct gctcgagct tccctgagct	2150
gcacctcttg ctaacccac catcacactg ccaccctgcc ctagggtctc	2200
actagtacca agtgggtcag cacagggctg aggatggggc tcctatccac	2250
cctggccagc acccagctta gtgctgggac tagcccagaa acttgaatgg	2300
gacctgaga gagccagggg tcccctgagg ccccctagg ggctttctgt	2350
ctgccccagg gtgctccatg gatctccctg tggcagcagg catggagagt	2400
cagggtgccc ttcattggcag taggtcttaa gtgggtgact ggccacaggc	2450
cgagaaaagg gtacagcctc taggtggggg tcccaaagac gccttcaggc	2500
tggactgagc tgctctccca cagggtttct gtgcagctgg attttctctg	2550
ttgcatacat gcctggcatc tgtctcccct tgttctgag tggccccaca	2600
tggggctctg agcaggctgt atctggatc tggcaataaa agtactctgg	2650
atgctgtaaa aaaaaaaaaa aaaaaaaaaa	2680

<210> SEQ ID NO 14
<211> LENGTH: 412
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 14

Met Gly Leu His Leu Arg Pro Tyr Arg Val Gly Leu Leu Pro Asp	
1 5 10 15	
Gly Leu Leu Phe Leu Leu Leu Leu Met Leu Leu Ala Asp Pro	
20 25 30	
Ala Leu Pro Ala Gly Arg His Pro Pro Val Val Leu Val Pro Gly	
35 40 45	
Asp Leu Gly Asn Gln Leu Glu Ala Lys Leu Asp Lys Pro Thr Val	
50 55 60	
Val His Tyr Leu Cys Ser Lys Lys Thr Glu Ser Tyr Phe Thr Ile	
65 70 75	
Trp Leu Asn Leu Glu Leu Leu Leu Pro Val Ile Ile Asp Cys Trp	
80 85 90	
Ile Asp Asn Ile Arg Leu Val Tyr Asn Lys Thr Ser Arg Ala Thr	
95 100 105	
Gln Phe Pro Asp Gly Val Asp Val Arg Val Pro Gly Phe Gly Lys	
110 115 120	
Thr Phe Ser Leu Glu Phe Leu Asp Pro Ser Lys Ser Ser Val Gly	
125 130 135	
Ser Tyr Phe His Thr Met Val Glu Ser Leu Val Gly Trp Gly Tyr	
140 145 150	
Thr Arg Gly Glu Asp Val Arg Gly Ala Pro Tyr Asp Trp Arg Arg	
155 160 165	
Ala Pro Asn Glu Asn Gly Pro Tyr Phe Leu Ala Leu Arg Glu Met	
170 175 180	

-continued

Ile	Glu	Glu	Met	Tyr	Gln	Leu	Tyr	Gly	Gly	Pro	Val	Val	Leu	Val
				185					190					195
Ala	His	Ser	Met	Gly	Asn	Met	Tyr	Thr	Leu	Tyr	Phe	Leu	Gln	Arg
				200					205					210
Gln	Pro	Gln	Ala	Trp	Lys	Asp	Lys	Tyr	Ile	Arg	Ala	Phe	Val	Ser
				215					220					225
Leu	Gly	Ala	Pro	Trp	Gly	Gly	Val	Ala	Lys	Thr	Leu	Arg	Val	Leu
				230					235					240
Ala	Ser	Gly	Asp	Asn	Asn	Arg	Ile	Pro	Val	Ile	Gly	Pro	Leu	Lys
				245					250					255
Ile	Arg	Glu	Gln	Gln	Arg	Ser	Ala	Val	Ser	Thr	Ser	Trp	Leu	Leu
				260					265					270
Pro	Tyr	Asn	Tyr	Thr	Trp	Ser	Pro	Glu	Lys	Val	Phe	Val	Gln	Thr
				275					280					285
Pro	Thr	Ile	Asn	Tyr	Thr	Leu	Arg	Asp	Tyr	Arg	Lys	Phe	Phe	Gln
				290					295					300
Asp	Ile	Gly	Phe	Glu	Asp	Gly	Trp	Leu	Met	Arg	Gln	Asp	Thr	Glu
				305					310					315
Gly	Leu	Val	Glu	Ala	Thr	Met	Pro	Pro	Gly	Val	Gln	Leu	His	Cys
				320					325					330
Leu	Tyr	Gly	Thr	Gly	Val	Pro	Thr	Pro	Asp	Ser	Phe	Tyr	Tyr	Glu
				335					340					345
Ser	Phe	Pro	Asp	Arg	Asp	Pro	Lys	Ile	Cys	Phe	Gly	Asp	Gly	Asp
				350					355					360
Gly	Thr	Val	Asn	Leu	Lys	Ser	Ala	Leu	Gln	Cys	Gln	Ala	Trp	Gln
				365					370					375
Ser	Arg	Gln	Glu	His	Gln	Val	Leu	Leu	Gln	Glu	Leu	Pro	Gly	Ser
				380					385					390
Glu	His	Ile	Glu	Met	Leu	Ala	Asn	Ala	Thr	Thr	Leu	Ala	Tyr	Leu
				395					400					405
Lys	Arg	Val	Leu	Leu	Gly	Pro								
				410										

<210> SEQ ID NO 15
<211> LENGTH: 1371
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 15

cagagcagat aatggcaagc atggctgccg tgctcacctg ggctctggct	50
cttctttcag cgttttcggc caccaggca cggaaaggct tctgggacta	100
cttcagccag accagcgggg acaaaggcag ggtggagcag atccatcagc	150
agaagatggc tcgcgagccc gcgacctga aagacagcct tgagcaagac	200
ctcaacaata tgaacaagtt cctggaaaag ctgaggcctc tgagtgggag	250
cgaggctcct cggctcccac aggaccgggt gggcatgcgg cggcagctgc	300
aggaggaggt ggaggaggtg aaggctcgcc tocagcccta catggcagag	350
gcgcacgagc tggtagggctg gaatttgag ggcttgcggc agcaactgaa	400
gccctacacg atggatctga tggagcaggt ggcctgcgc gtgcaggagc	450
tgacaggagca gttgcgcgtg gtgggggaag acaccaaggc ccagttgctg	500

-continued

```
gggggctgg acgaggcttg ggctttgctg cagggactgc agagccgcgt      550
gggtgaccac accggccgct tcaaagagct cttccacca tacgccgaga      600
gcctggtgag cggcatcggg cgccacgtgc aggagctgca ccgcagtgtg      650
gctccgcacg ccccgccag ccccgcgcg ctcagtcgct gcgtgcaggt      700
gctctccggg aagctcacgc tcaaggccaa ggccctgcac gcacgcatcc      750
agcagaacct ggaccagctg cgcaagagc tcagcagagc ctttgcaggc      800
actgggactg aggaaggggc cgcccgagc ccctagatgc tctccgagga      850
gggtgcgccg cgacttcagg ctttccgcca ggacacctac ctgcagatag      900
ctgccttcac tcgcgccatc gaccaggaga ctgaggaggt ccagcagcag      950
ctggcgccac ctccaccagg ccacagtgc ttcgcccag agtttcaaca     1000
aacagacagt ggcaaggttc tgagcaagct gcaggcccg ctggatgacc     1050
tgttggaaga catcactcac agccttcag accagggcca cagccatctg     1100
ggggaccctt gaggatctac ctgcccaggc ccattcccag cttcttgtct     1150
ggggagcctt ggctctgagc ctctagcatg gttcagtcct tgaagtggc     1200
ctgttgggtg gagggtgga ggctctgtgc aggacaggga ggccaccaa     1250
ggggctgctg tctcctgcat atccagcctc ctgcgactcc ccaatctgga     1300
tgcattacat tcaccaggct ttgcaaaaaa aaaaaaaaaa aaaaaaaaaa     1350
aaaaaaaaa aaaaaaaaaa a                                     1371
```

<210> SEQ ID NO 16

<211> LENGTH: 274

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 16

```
Met Ala Ser Met Ala Ala Val Leu Thr Trp Ala Leu Ala Leu Leu
 1             5             10             15
Ser Ala Phe Ser Ala Thr Gln Ala Arg Lys Gly Phe Trp Asp Tyr
          20             25             30
Phe Ser Gln Thr Ser Gly Asp Lys Gly Arg Val Glu Gln Ile His
          35             40             45
Gln Gln Lys Met Ala Arg Glu Pro Ala Thr Leu Lys Asp Ser Leu
          50             55             60
Glu Gln Asp Leu Asn Asn Met Asn Lys Phe Leu Glu Lys Leu Arg
          65             70             75
Pro Leu Ser Gly Ser Glu Ala Pro Arg Leu Pro Gln Asp Pro Val
          80             85             90
Gly Met Arg Arg Gln Leu Gln Glu Glu Leu Glu Glu Val Lys Ala
          95             100            105
Arg Leu Gln Pro Tyr Met Ala Glu Ala His Glu Leu Val Gly Trp
          110            115            120
Asn Leu Glu Gly Leu Arg Gln Gln Leu Lys Pro Tyr Thr Met Asp
          125            130            135
Leu Met Glu Gln Val Ala Leu Arg Val Gln Glu Leu Gln Glu Gln
          140            145            150
Leu Arg Val Val Gly Glu Asp Thr Lys Ala Gln Leu Leu Gly Gly
```

-continued

	155		160		165
Val Asp Glu Ala Trp Ala Leu Leu Gln Gly Leu Gln Ser Arg Val					
	170		175		180
Val His His Thr Gly Arg Phe Lys Glu Leu Phe His Pro Tyr Ala					
	185		190		195
Glu Ser Leu Val Ser Gly Ile Gly Arg His Val Gln Glu Leu His					
	200		205		210
Arg Ser Val Ala Pro His Ala Pro Ala Ser Pro Ala Arg Leu Ser					
	215		220		225
Arg Cys Val Gln Val Leu Ser Arg Lys Leu Thr Leu Lys Ala Lys					
	230		235		240
Ala Leu His Ala Arg Ile Gln Gln Asn Leu Asp Gln Leu Arg Glu					
	245		250		255
Glu Leu Ser Arg Ala Phe Ala Gly Thr Gly Thr Glu Glu Gly Ala					
	260		265		270
Gly Pro Asp Pro					
<210> SEQ ID NO 17					
<211> LENGTH: 2854					
<212> TYPE: DNA					
<213> ORGANISM: Homo Sapien					
<400> SEQUENCE: 17					
ctaagaggac aagatgaggc ccggcctctc atttctccta gcccttctgt					50
tcttccttgg ccaagctgca ggggatttgg gggatgtggg acctccaatt					100
cccagccccc gcttcagctc tttccaggt gttgactcca gctccagctt					150
cagctccagc tccaggtcgg gctccagctc cagccgcagc ttaggcagcg					200
gaggttctgt gtcccagttg ttttccaatt tcaccggctc cgtggatgac					250
cgtggggacct gccagtgtc tgtttccctg ccagacacca cctttcccgt					300
ggacagagtg gaacgcttgg aattcacagc tcattgttctt tctcagaagt					350
ttgagaaaga actttctaaa gtgagggaat atgtccaatt aattagtgtg					400
tatgaaaaga aactgttaaa cctaactgtc cgaattgaca tcatggagaa					450
ggataccatt tcttacactg aactggactt cgagctgata aaggtagaag					500
tgaaggagat ggaaaaactg gtcatacagc tgaaggagag ttttggtgga					550
agctcagaaa ttgttgacca gctggagggt gagataagaa atatgactct					600
cttggtagag aagcttgaga cactagacaa aaacaatgtc cttgccattc					650
gccgagaaat cgtggctctg aagaccaagc tgaagagtg tgaggcctct					700
aaagatcaaa acaccctgt cgtccaccct cctcccactc cagggagctg					750
tggtcatggt ggtgtggtga acatcagcaa accgtctgtg gttcagctca					800
actggagagg gttttcttat ctatatggtg cttggggtag ggattactct					850
ccccagcatc caaacaaggt actgtatttg gtggcgccat tgaatacaga					900
tgggagactg ttggagtatt atagactgta caacacactg gatgatttgc					950
tattgtatat aaatgctcga gagttgcgga tcacctatgg ccaaggtagt					1000
ggtacagcag tttaacaaca caacatgtac gtcaacatgt acaacaccgg					1050
gaatatggcc agagttaacc tgaccaccaa cagattgct gtgactcaaa					1100

-continued

ctctccctaa tgctgcctat aataaccgct tttcatatgc taatgttgct	1150
tggaagata ttgactttgc tgtggatgag aatggattgt gggttattta	1200
ttcaactgaa gccagcactg gtaacatggt gattagtaaa ctcaatgaca	1250
ccacacttca ggtgctaaac acttggtata ccaagcagta taaaccatct	1300
gcttctaacg ccttcatggt atgtgggggt ctgtatgcc cccgtactat	1350
gaacaccaga acagaagaga ttttttacta ttatgacaca aacacaggga	1400
aagagggcaa actagacatt gtaatgcata agatgcagga aaaagtgcag	1450
agcattaact ataacccttt tgaccagaaa ctttatgtct ataacgatgg	1500
ttaccttctg aattatgac tttctgtctt gcagaagccc cagtaagctg	1550
tttaggagtt aggtgaaa agaaaatggt tgttgaaaa atagtcttct	1600
ccacttactt agatatctgc aggggtgtct aaaagtgtgt tcattttgca	1650
gcaatgttta ggtgcatagt tctaccacac tagagatcta ggacatttgt	1700
cttgatttgg tgagttctct tgggaatcat ctgcctcttc aggcgcattt	1750
tgcaataaag tctgtctagg gtgggattgt cagaggctca ggggcactgt	1800
gggcctagtg aagcctactg tgaggaggct tcactagaag ccttaaatta	1850
ggaattaagg aacttaaac tcagtatggc gtctagggat tctttgtaca	1900
ggaaatattg cccaatgact agtcctcatc catgtagcac cactaattct	1950
tccatgcctg gaagaaacct ggggacttag ttaggtagat taatatctgg	2000
agctcctcga gggaccaaat ctccaacttt tttttcccct cactagcacc	2050
tggaatgatg ctttgtatgt ggcagataag taaatttggc atgottatat	2100
attctacatc tgtaaaagtc tgagttttat ggagagaggc ctttttatgc	2150
attaaattgt acatggcaaa taaatcccag aaggatctgt agatgaggca	2200
cctgcttttt cttttctctc attgtccacc ttactaaaag tcagtagaat	2250
cttctacctc ataacttct tccaaaggca gctcagaaga ttagaaccag	2300
acttactaac caattccacc cccaccaaac ccccttctac tgcctacttt	2350
aaaaaaatta atagttttct atggaactga tctaagatta gaaaaattaa	2400
ttttctttaa tttcattatg gacttttatt tacatgactc taagactata	2450
agaaaaatctg atggcagtga caaagtgcta gcattttattg ttatctaata	2500
aagaccttgg agcatatgtg caacttatga gtgtatcagt tgttgcattg	2550
aatttttgcc tttgtttaag cctggaaact gtaagaaaat gaaaatttaa	2600
tttttttttc taggacgagc tatagaaaag ctattgagag tatctagtta	2650
atcagtgcag tagttggaaa ccttgctggt gtatgtgatg tgcttctgtg	2700
cttttgaatg actttatcat ctagtctttg tctatttttc ctttgatgtt	2750
caagtcctag tctataggat tggcagttta aatgctttac tccccctttt	2800
aaaataaatg attaaaatgt gctttgaaaa aaaaaaaaaa aaaaaaaaaa	2850
aaaa	2854

<210> SEQ ID NO 18

<211> LENGTH: 510

-continued

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 18

```

Met Arg Pro Gly Leu Ser Phe Leu Leu Ala Leu Leu Phe Phe Leu
 1           5           10           15

Gly Gln Ala Ala Gly Asp Leu Gly Asp Val Gly Pro Pro Ile Pro
          20           25           30

Ser Pro Gly Phe Ser Ser Phe Pro Gly Val Asp Ser Ser Ser Ser
          35           40           45

Phe Ser Ser Ser Ser Arg Ser Gly Ser Ser Ser Ser Arg Ser Leu
          50           55           60

Gly Ser Gly Gly Ser Val Ser Gln Leu Phe Ser Asn Phe Thr Gly
          65           70           75

Ser Val Asp Asp Arg Gly Thr Cys Gln Cys Ser Val Ser Leu Pro
          80           85           90

Asp Thr Thr Phe Pro Val Asp Arg Val Glu Arg Leu Glu Phe Thr
          95          100          105

Ala His Val Leu Ser Gln Lys Phe Glu Lys Glu Leu Ser Lys Val
          110          115          120

Arg Glu Tyr Val Gln Leu Ile Ser Val Tyr Glu Lys Lys Leu Leu
          125          130          135

Asn Leu Thr Val Arg Ile Asp Ile Met Glu Lys Asp Thr Ile Ser
          140          145          150

Tyr Thr Glu Leu Asp Phe Glu Leu Ile Lys Val Glu Val Lys Glu
          155          160          165

Met Glu Lys Leu Val Ile Gln Leu Lys Glu Ser Phe Gly Gly Ser
          170          175          180

Ser Glu Ile Val Asp Gln Leu Glu Val Glu Ile Arg Asn Met Thr
          185          190          195

Leu Leu Val Glu Lys Leu Glu Thr Leu Asp Lys Asn Asn Val Leu
          200          205          210

Ala Ile Arg Arg Glu Ile Val Ala Leu Lys Thr Lys Leu Lys Glu
          215          220          225

Cys Glu Ala Ser Lys Asp Gln Asn Thr Pro Val Val His Pro Pro
          230          235          240

Pro Thr Pro Gly Ser Cys Gly His Gly Gly Val Val Asn Ile Ser
          245          250          255

Lys Pro Ser Val Val Gln Leu Asn Trp Arg Gly Phe Ser Tyr Leu
          260          265          270

Tyr Gly Ala Trp Gly Arg Asp Tyr Ser Pro Gln His Pro Asn Lys
          275          280          285

Gly Leu Tyr Trp Val Ala Pro Leu Asn Thr Asp Gly Arg Leu Leu
          290          295          300

Glu Tyr Tyr Arg Leu Tyr Asn Thr Leu Asp Asp Leu Leu Leu Tyr
          305          310          315

Ile Asn Ala Arg Glu Leu Arg Ile Thr Tyr Gly Gln Gly Ser Gly
          320          325          330

Thr Ala Val Tyr Asn Asn Asn Met Tyr Val Asn Met Tyr Asn Thr
          335          340          345

Gly Asn Ile Ala Arg Val Asn Leu Thr Thr Asn Thr Ile Ala Val
          350          355          360

```

-continued

Thr	Gln	Thr	Leu	Pro	Asn	Ala	Ala	Tyr	Asn	Asn	Arg	Phe	Ser	Tyr
			365						370					375
Ala	Asn	Val	Ala	Trp	Gln	Asp	Ile	Asp	Phe	Ala	Val	Asp	Glu	Asn
			380						385					390
Gly	Leu	Trp	Val	Ile	Tyr	Ser	Thr	Glu	Ala	Ser	Thr	Gly	Asn	Met
			395						400					405
Val	Ile	Ser	Lys	Leu	Asn	Asp	Thr	Thr	Leu	Gln	Val	Leu	Asn	Thr
			410						415					420
Trp	Tyr	Thr	Lys	Gln	Tyr	Lys	Pro	Ser	Ala	Ser	Asn	Ala	Phe	Met
			425						430					435
Val	Cys	Gly	Val	Leu	Tyr	Ala	Thr	Arg	Thr	Met	Asn	Thr	Arg	Thr
			440						445					450
Glu	Glu	Ile	Phe	Tyr	Tyr	Tyr	Asp	Thr	Asn	Thr	Gly	Lys	Glu	Gly
			455						460					465
Lys	Leu	Asp	Ile	Val	Met	His	Lys	Met	Gln	Glu	Lys	Val	Gln	Ser
			470						475					480
Ile	Asn	Tyr	Asn	Pro	Phe	Asp	Gln	Lys	Leu	Tyr	Val	Tyr	Asn	Asp
			485						490					495
Gly	Tyr	Leu	Leu	Asn	Tyr	Asp	Leu	Ser	Val	Leu	Gln	Lys	Pro	Gln
			500						505					510

<210> SEQ ID NO 19
<211> LENGTH: 663
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 19

gcaccgcaga cggcgcggat cgcagggagc cggtcgcgcg ccggaacggg	50
agcctgggtg tgcgtgtgga gtccggactc gtgggagacg atcgcgatga	100
acacggtgct gtcgcggggc aactcactgt tcgccttctc gctgagcgtg	150
atggcgggcg tcaccttcgg ctgcttcac accaccgcct tcaaagacag	200
gagcgctccc gtgcggctgc acgtctcgcg gatcatgcta aaaaatgtag	250
aagatttcac tggacctaga gaaagaagtg atctgggatt tatcacattt	300
gataatactg ctgatctaga gaatatattt gattggaatg ttaagcagtt	350
gtttctttat ttatcagcag aatattcaac aaaaaataat gctctgaacc	400
aagttgtcct atgggacaag attgttttga gaggtgataa tccgaagctg	450
ctgctgaaag atatgaaaac aaaatatatt ttctttgacg atggaaatgg	500
tctcaaggga aacaggaatg tcactttgac cctgtcttgg aacgtcgtac	550
caaatgctgg aattctacct ctgtgtgacg gatcaggaca cgtatctgtc	600
ccatttccag atacatatga aatacgaag agttattaaa ttattctgaa	650
tttgaacaaa aaa	663

<210> SEQ ID NO 20
<211> LENGTH: 180
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 20

Met	Asn	Thr	Val	Leu	Ser	Arg	Ala	Asn	Ser	Leu	Phe	Ala	Phe	Ser
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

-continued

1	5	10	15
Leu Ser Val Met	Ala Ala Leu Thr Phe	Gly Cys Phe Ile Thr	Thr
	20	25	30
Ala Phe Lys Asp	Arg Ser Val Pro Val Arg	Leu His Val Ser	Arg
	35	40	45
Ile Met Leu Lys	Asn Val Glu Asp Phe Thr	Gly Pro Arg Glu	Arg
	50	55	60
Ser Asp Leu Gly	Phe Ile Thr Phe Asp	Ile Thr Ala Asp	Leu Glu
	65	70	75
Asn Ile Phe Asp	Trp Asn Val Lys Gln	Leu Phe Leu Tyr	Leu Ser
	80	85	90
Ala Glu Tyr Ser	Thr Lys Asn Asn Ala	Leu Asn Gln Val	Val Leu
	95	100	105
Trp Asp Lys Ile	Val Leu Arg Gly Asp	Asn Pro Lys Leu	Leu Leu
	110	115	120
Lys Asp Met Lys	Thr Lys Tyr Phe Phe	Phe Asp Asp Gly	Asn Gly
	125	130	135
Leu Lys Gly Asn	Arg Asn Val Thr Leu	Thr Leu Ser Trp	Asn Val
	140	145	150
Val Pro Asn Ala	Gly Ile Leu Pro Leu	Val Thr Gly Ser	Gly His
	155	160	165
Val Ser Val Pro	Phe Pro Asp Thr Tyr	Glu Ile Thr Lys	Ser Tyr
	170	175	180

<210> SEQ ID NO 21
<211> LENGTH: 415
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 21

aaacttgacg ccatgaagat cccggtcctt cctgccgtgg tgetcctctc	50
cctcctgggt ctccactctg cccagggagc caccctgggt ggtcctgagg	100
aagaaagcac cattgagaat tatgcgtcac gacccgaggc ctttaacacc	150
ccgttcctga acatcgacaa attgcgatct gcgtttaagg ctgatgagtt	200
cctgaactgg cagccctctt ttgagtctat caaaaggaaa cttcctttcc	250
tcaactggga tgcctttcct aagctgaaag gactgaggag cgcaactcct	300
gatgccagtg gaccatgacc tccactggaa gagggggcta gcgtgagcgc	350
tgattctcaa cctaccataa ctctttcctg cctcaggaac tccaataaaa	400
cattttccat ccaaa	415

<210> SEQ ID NO 22
<211> LENGTH: 99
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 22

Met Lys Ile Pro Val Leu Pro Ala Val Val Leu Leu Ser Leu Leu	
1	5 10 15
Val Leu His Ser Ala Gln Gly Ala Thr Leu Gly Gly Pro Glu Glu	
	20 25 30
Glu Ser Thr Ile Glu Asn Tyr Ala Ser Arg Pro Glu Ala Phe Asn	

-continued

				35				40						45
Thr	Pro	Phe	Leu	Asn	Ile	Asp	Lys	Leu	Arg	Ser	Ala	Phe	Lys	Ala
				50					55					60
Asp	Glu	Phe	Leu	Asn	Trp	His	Ala	Leu	Phe	Glu	Ser	Ile	Lys	Arg
				65					70					75
Lys	Leu	Pro	Phe	Leu	Asn	Trp	Asp	Ala	Phe	Pro	Lys	Leu	Lys	Gly
				80					85					90
Leu	Arg	Ser	Ala	Thr	Pro	Asp	Ala	Gln						
				95										

<210> SEQ ID NO 23
<211> LENGTH: 866
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 23

tctcagactc	ttggaagggg	ctatactaga	cacacaaaga	cagccccaag	50
aaggacggtg	gagtagtgtc	ctcgctaaaa	gacagtagat	atgcaacgcc	100
tcttgctcct	gccctttctc	ctgctgggaa	cagtttctgc	tcttcatctg	150
gagaatgatg	ccccccatct	ggagagccta	gagacacagg	cagacctagg	200
ccaggatctg	gatagttcaa	aggagcagga	gagagacttg	gctctgacgg	250
aggaggtgat	tcaggcagag	ggagaggagg	tcaaggcttc	tgctgtctaa	300
gacaactttg	aggatgagga	agccatggag	tcggacccag	ctgccttaga	350
caaggacttc	cagtgcacca	gggaagaaga	cattgttgaa	gtgcagggaa	400
gtccaaggtg	caagacctgc	cgctacctat	tgggtcggac	tcctaaaact	450
tttgcagaag	ctcagaatgt	ctgcagcaga	tgctacggag	gcaaccttgt	500
ctctatccat	gacttcaact	tcaactatcg	cattcagtgc	tgactagca	550
cagtcaacca	agcccaggtc	tggattggag	gcaacctcag	gggctggttc	600
ctgtggaagc	ggttttgctg	gactgatggg	agccactgga	attttgctta	650
ctgggtcccca	gggcaacctg	ggaatgggca	aggctcctgt	gtggccctat	700
gcaccaaagg	aggttattgg	cgacgagctc	aatgcgacaa	gcaactgccc	750
ttcgtctgct	ccttctaagc	cagcggcacg	gagaccctgc	cagcagctcc	800
ctcccgtccc	ccaacctctc	ctgctcataa	atccagactt	cccacagcaa	850
aaaaaaaaaa	aaaaaa				866

<210> SEQ ID NO 24
<211> LENGTH: 225
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 24

Met	Gln	Arg	Leu	Leu	Leu	Leu	Pro	Phe	Leu	Leu	Leu	Gly	Thr	Val
1				5					10					15
Ser	Ala	Leu	His	Leu	Glu	Asn	Asp	Ala	Pro	His	Leu	Glu	Ser	Leu
				20					25					30
Glu	Thr	Gln	Ala	Asp	Leu	Gly	Gln	Asp	Leu	Asp	Ser	Ser	Lys	Glu
				35					40					45
Gln	Glu	Arg	Asp	Leu	Ala	Leu	Thr	Glu	Glu	Val	Ile	Gln	Ala	Glu

-continued

	50		55		60
Gly Glu Glu Val Lys Ala Ser Ala Cys Gln Asp Asn Phe Glu Asp	65		70		75
Glu Glu Ala Met Glu Ser Asp Pro Ala Ala Leu Asp Lys Asp Phe	80		85		90
Gln Cys Pro Arg Glu Glu Asp Ile Val Glu Val Gln Gly Ser Pro	95		100		105
Arg Cys Lys Thr Cys Arg Tyr Leu Leu Val Arg Thr Pro Lys Thr	110		115		120
Phe Ala Glu Ala Gln Asn Val Cys Ser Arg Cys Tyr Gly Gly Asn	125		130		135
Leu Val Ser Ile His Asp Phe Asn Phe Asn Tyr Arg Ile Gln Cys	140		145		150
Cys Thr Ser Thr Val Asn Gln Ala Gln Val Trp Ile Gly Gly Asn	155		160		165
Leu Arg Gly Trp Phe Leu Trp Lys Arg Phe Cys Trp Thr Asp Gly	170		175		180
Ser His Trp Asn Phe Ala Tyr Trp Ser Pro Gly Gln Pro Gly Asn	185		190		195
Gly Gln Gly Ser Cys Val Ala Leu Cys Thr Lys Gly Gly Tyr Trp	200		205		210
Arg Arg Ala Gln Cys Asp Lys Gln Leu Pro Phe Val Cys Ser Phe	215		220		225

<210> SEQ ID NO 25
<211> LENGTH: 584
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 25

caacagaagc caagaaggaa gccgtctatc ttgtggcgat catgtataag	50
ctggcctcct gctgtttgct tttcacagga ttcttaaadc ctctcttadc	100
tcttctcttc cttgactcca gggaaatadc ctttcaactc tcagcacctc	150
atgaagacgc gcgcttaact ccggaggagc tagaaagagc ttcccttcta	200
cagatattgc cagagatgct ggggtgcagaa agaggggata ttctcaggaa	250
agcagactca agtaccaaca tttttaaccc aagaggaaat ttgagaaagt	300
ttcaggattt ctctggacaa gatcctaaca ttttactgag tcatcttttg	350
gccagaatct ggaaccata caagaaacgt gagactcctg attgcttctg	400
gaaatactgt gtctgaagtg aaataagcat ctgttagtca gctcagaaac	450
acccatctta gaatatgaaa aataacacaa tgcttgattt gaaaacagt	500
tggaagaaaa ctaggcaaac tacacctgt tcattgttac ctggaaaata	550
aatcctctat gttttgcaca aaaaaaaaaa aaaa	584

<210> SEQ ID NO 26
<211> LENGTH: 124
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 26

Met Tyr Lys Leu Ala Ser Cys Cys Leu Leu Phe Thr Gly Phe Leu

-continued

[illegible]

<210> SEQ ID NO 27

<211> LENGTH: 920

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 27

caagtaaatg	cagcactagt	gggtgggatt	gaggtatgcc	ctggtgcata	50
aatagagact	cagctgtgct	ggcacactca	gaagcttgga	ccgcataccta	100
gccgccgact	cacacaagcg	aggtgggtga	ggaaatccag	agttgccatg	150
gagaaaaattc	cagtgtcagc	attcttgtct	cttgtggccc	tctctacac	200
tctggccaga	gataccacag	tcaaacctgg	agccaaaaag	gacacaaagg	250
actctcgacc	caaactgccc	cagaccctct	ccagaggttg	gggtgaccaa	300
ctcatctgga	ctcagacata	tgaagaagct	ctatataaat	ccaagacaag	350
caacaaaccc	ttgatgatta	ttcatcactt	ggatgagtgc	ccacacagtc	400
aagcttttaa	gaaagtgttt	gctgaaaata	aagaaatcca	gaaattggca	450
gagcagtttg	tctctctcaa	tctgttttat	gaaacaactg	acaaacacct	500
ttctcctgat	ggccagtatg	tcccaggat	tatgtttgtt	gacccatctc	550
tgacagttag	agccgatatc	actggaagat	attcaaatcg	tctctatgct	600
tacgaacctg	cagatacagc	tctgttgctt	gacaacatga	agaaagctct	650
caagttgctg	aagactgaat	tgtaaagaaa	aaaaatctcc	aagcccttct	700
gtctgtcagg	ccttgagact	tgaaaccaga	agaagtgtga	gaagactggc	750
tagtgtggaa	gcatagtgaa	cacactgatt	aggttatggt	ttaatgttac	800
aacaactatt	ttttaagaaa	aacaagtttt	agaaatttgg	tttcaagtgt	850
acatgtgtga	aaacaatatt	gtatactacc	atagtgagcc	atgattttct	900
aaaaaaaaaa	ataaatgtta				920

<210> SEQ ID NO 28

```
<210> SEQ ID NO: 2
<211> LENGTH: 175
```

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

-continued

<400> SEQUENCE: 28

Met	Glu	Lys	Ile	Pro	Val	Ser	Ala	Phe	Leu	Leu	Leu	Val	Ala	Leu
1				5					10					15
Ser	Tyr	Thr	Leu	Ala	Arg	Asp	Thr	Thr	Val	Lys	Pro	Gly	Ala	Lys
			20						25					30
Lys	Asp	Thr	Lys	Asp	Ser	Arg	Pro	Lys	Leu	Pro	Gln	Thr	Leu	Ser
			35						40					45
Arg	Gly	Trp	Gly	Asp	Gln	Leu	Ile	Trp	Thr	Gln	Thr	Tyr	Glu	Glu
			50						55					60
Ala	Leu	Tyr	Lys	Ser	Lys	Thr	Ser	Asn	Lys	Pro	Leu	Met	Ile	Ile
			65						70					75
His	His	Leu	Asp	Glu	Cys	Pro	His	Ser	Gln	Ala	Leu	Lys	Lys	Val
			80						85					90
Phe	Ala	Glu	Asn	Lys	Glu	Ile	Gln	Lys	Leu	Ala	Glu	Gln	Phe	Val
			95						100					105
Leu	Leu	Asn	Leu	Val	Tyr	Glu	Thr	Thr	Asp	Lys	His	Leu	Ser	Pro
			110						115					120
Asp	Gly	Gln	Tyr	Val	Pro	Arg	Ile	Met	Phe	Val	Asp	Pro	Ser	Leu
			125						130					135
Thr	Val	Arg	Ala	Asp	Ile	Thr	Gly	Arg	Tyr	Ser	Asn	Arg	Leu	Tyr
			140						145					150
Ala	Tyr	Glu	Pro	Ala	Asp	Thr	Ala	Leu	Leu	Leu	Asp	Asn	Met	Lys
			155						160					165
Lys	Ala	Leu	Lys	Leu	Leu	Lys	Thr	Glu	Leu					
			170						175					

<210> SEQ ID NO 29

<211> LENGTH: 1181

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 29

aagaccctct ctttcgctgt ttgagagtct ctcggctcaa ggaccgggag	50
gtaagaggtt tgggactgcc ccggcaactc caggggtgtct ggtccacgac	100
ctatcctagg cgccatgggt gtgataggta tacagctggt tgttaccatg	150
gtgatggcca gtgtcatgca gaagattata cctcactatt ctcttgctcg	200
atggctactc tgtaatggca gtttgagggt gtatcaacat cctacagaag	250
aagaattaag aattcttgca gggaaacaac aaaagggaa aaccaaaaaa	300
gataggaaat ataatgttca cattgaaagt aagccattaa ccattccaaa	350
ggatattgac cttcatctag aaacaaagtc agttacagaa gtggatactt	400
tagcattgca ttactttcca gaataccagt ggctgggtgga ttccacagt	450
gctgctacag ttgtgtatct agtaactgaa gtctactaca attttatgaa	500
gcctacacag gaaatgaata tcagcttagt ctggtgccta cttgttttgt	550
cttttgcaat caaagttcta ttttcattaa ctacacacta ttttaaagta	600
gaagatggtg gtgaaagatc tgtttgtgtc acctttggat ttttttctt	650
tgtcaaagca atggcagtgt tgattgtaac agaaaattat ctggaatttg	700
gacttgaaac agggtttaca aatttttcag acagtgcgat gcagtttctt	750

-continued

```

gaaaagcaag gtttagaatc tcagagtcct gtttcaaac ttactttcaa      800
atttttcctg gctattttct gttcattcat tggggctttt ttgacatttc      850
ctggattacg actggctcaa atgcatctgg atgccctgaa tttggcaaca      900
gaaaaaatta cacaaacttt acttcatatc aacttcttgg cacctttatt      950
tatggttttg ctctgggtaa aaccaatcac caaagactac attatgaacc     1000
caccactggg caaagaaatt tcccatctg gaagatgaag ataatagtat     1050
ctaactcaca aggttatcat tggaataaat gaaagaacac atgtaatgca     1100
accagctgga attaagtgtc taataaatgt tcttttctact gctttgcctc     1150
atcagaatta aaatagaaat acttgactag t                          1181

```

<210> SEQ ID NO 30

<211> LENGTH: 307

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 30

```

Met Gly Val Ile Gly Ile Gln Leu Val Val Thr Met Val Met Ala
  1           5           10           15
Ser Val Met Gln Lys Ile Ile Pro His Tyr Ser Leu Ala Arg Trp
          20           25           30
Leu Leu Cys Asn Gly Ser Leu Arg Trp Tyr Gln His Pro Thr Glu
          35           40           45
Glu Glu Leu Arg Ile Leu Ala Gly Lys Gln Gln Lys Gly Lys Thr
          50           55           60
Lys Lys Asp Arg Lys Tyr Asn Gly His Ile Glu Ser Lys Pro Leu
          65           70           75
Thr Ile Pro Lys Asp Ile Asp Leu His Leu Glu Thr Lys Ser Val
          80           85           90
Thr Glu Val Asp Thr Leu Ala Leu His Tyr Phe Pro Glu Tyr Gln
          95          100          105
Trp Leu Val Asp Phe Thr Val Ala Ala Thr Val Val Tyr Leu Val
          110          115          120
Thr Glu Val Tyr Tyr Asn Phe Met Lys Pro Thr Gln Glu Met Asn
          125          130          135
Ile Ser Leu Val Trp Cys Leu Leu Val Leu Ser Phe Ala Ile Lys
          140          145          150
Val Leu Phe Ser Leu Thr Thr His Tyr Phe Lys Val Glu Asp Gly
          155          160          165
Gly Glu Arg Ser Val Cys Val Thr Phe Gly Phe Phe Phe Phe Val
          170          175          180
Lys Ala Met Ala Val Leu Ile Val Thr Glu Asn Tyr Leu Glu Phe
          185          190          195
Gly Leu Glu Thr Gly Phe Thr Asn Phe Ser Asp Ser Ala Met Gln
          200          205          210
Phe Leu Glu Lys Gln Gly Leu Glu Ser Gln Ser Pro Val Ser Lys
          215          220          225
Leu Thr Phe Lys Phe Phe Leu Ala Ile Phe Cys Ser Phe Ile Gly
          230          235          240
Ala Phe Leu Thr Phe Pro Gly Leu Arg Leu Ala Gln Met His Leu

```

-continued

245									250					255															
Asp	Ala	Leu	Asn	Leu	Ala	Thr	Glu	Lys	Ile	Thr	Gln	Thr	Leu	Leu															
260									265					270															
His	Ile	Asn	Phe	Leu	Ala	Pro	Leu	Phe	Met	Val	Leu	Leu	Trp	Val															
275									280					285															
Lys	Pro	Ile	Thr	Lys	Asp	Tyr	Ile	Met	Asn	Pro	Pro	Leu	Gly	Lys															
290									295					300															
Glu	Ile	Ser	Pro	Ser	Gly	Arg																							
305																													

<210> SEQ ID NO 31
<211> LENGTH: 513
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 31

gtagcatagt gtgcagttca ctggacacaa agctttggct gcacctottc	50
tggaagctg gccatggggc ttttcgatg cattgcaatt ctgctgttcc	100
agaaaccac agtaaccgaa caacttaaga agtgctggaa taactatgta	150
caaggacatt gcaggaaaat ctgcagagta aatgaagtgc ctgaggcact	200
atgtgaaaaa gggagatact gttgcctcaa tatcaaggaa ctggaagcat	250
gtaaaaaaat tacaagcca cctcgtccaa agccagcaac acttgactg	300
actcttcaag actatgttac aataatagaa aatttcccaa gcctgaagac	350
acagtctaca taaatcaaat acaatttcgt tttcacttgc ttctcaacct	400
agtctaataa actaaggtga tgagatatac atcttcttcc ttctggtttc	450
ttgatcctta aaatgacctt cgagcatatt ctaataaagt gcattgccag	500
ttaaaaaaaa aaa	513

<210> SEQ ID NO 32
<211> LENGTH: 99
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 32

Met Gly Leu Phe Met Ile Ile Ala Ile Leu Leu Phe Gln Lys Pro	1	5	10	15
Thr Val Thr Glu Gln Leu Lys Lys Cys Trp Asn Asn Tyr Val Gln	20	25	30	
Gly His Cys Arg Lys Ile Cys Arg Val Asn Glu Val Pro Glu Ala	35	40	45	
Leu Cys Glu Asn Gly Arg Tyr Cys Cys Leu Asn Ile Lys Glu Leu	50	55	60	
Glu Ala Cys Lys Lys Ile Thr Lys Pro Pro Arg Pro Lys Pro Ala	65	70	75	
Thr Leu Ala Leu Thr Leu Gln Asp Tyr Val Thr Ile Ile Glu Asn	80	85	90	
Phe Pro Ser Leu Lys Thr Gln Ser Thr	95			

<210> SEQ ID NO 33
<211> LENGTH: 2684

-continued

<212> TYPE: DNA
<213> ORGANISM: Homo Sapien
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: 2636-2637
<223> OTHER INFORMATION: unknown base

<400> SEQUENCE: 33

cggacgcgtg ggcgctgagc cccggaggcc agggcgctccg gggctgcgcc	50
acttccgagg gccgagcgt gctcaaccgt cggctgcgtg agcccccccc	100
ggaggagaac aacgcaagg gctcaaccgt cggctgcgtg agcccccccc	150
ggggcgctgg ctcgcgcccc ctcagctggg gagggcgggg ctcgctgcgc	200
cctgctgcgc actgcgaccc ttacagggga ggagggcgcg aggccgcgcg	250
gagatgagga ggaggctgcg cctacgcagg gacgcattgc tcacgctgct	300
ccttggcgcc tccctgggcc tcttactcta tgcgcagcgc gacggcgcg	350
ccccgacggc gagcgcgccg cgaggcgag ggagggcggc accgaggccc	400
acccccggac cccgcgcgtt ccagttaccc gacgcgggtg cagccccgcc	450
ggcctacgaa ggggacacac cggcgccgcc cagcctacg ggaccctttg	500
acttcgcccc ctatttgcgc gccaaaggacc agcggcggtt tccactgctc	550
attaaccagc cgcacaagt cgcggcgac ggcgcaccgc gtggcgcccc	600
ggacctgtt attgctgtca agtcggtggc agaggacttc gagcgcgccc	650
aagccgtgcg ccagacgtg ggcgcggagg gtcgcgtgca gggggcgctg	700
gtgcgcccg tggtcttctt gggcgctgcc agggcgcgag gtcggggcg	750
ggccgacgaa gttggggagg gcgcgcgaac cactggcggc gccctgctgc	800
gggcgagag ccttgcgtat gcggacatcc tgctctgggc cttcgacgac	850
acctttttta acctaacgt caaggagatc cactttctag cctgggcctc	900
agctttctgc cccgacgtgc gcttcgtttt taaggcgac gcagatgtgt	950
tcgtgaacct gggaaatctc ctggagttcc tggcgccgcg ggaccggcg	1000
caagacctgc ttgctggtga cgtaattgtg catgcgcggc ccatccgcac	1050
gcgggctagc aagtactaca tccccaggc cgtgtacggc ctgccgcct	1100
atccggccta cgcggggcg ggtggctttg tgctttccgg gccacgctg	1150
caccgcctgg ctggcgccctg tgcgcaggtc gagctcttcc ccatcgacga	1200
cgtctttctg ggcattgtgc tgcagcgct cgcgctcacg cccgagcctc	1250
accctgcctt ccgcacctt ggcaccccc agccttcagc cgcgcgcgat	1300
ttgagcacct tcgaccctg cttttaccgt gagctggttg tagtgacgg	1350
gctctcgccc gctgacatct ggcttatgtg gcgcctgctg caggggccgc	1400
atgggccagc ctgtgcgat ccacagcctg tcgctgcagg ccccttccaa	1450
tgggactcct agctccccac tacagcccca agctcctaac tcagaccag	1500
aatggagcgc gtttcccaga ttattgccgt gtatgtggtt cttocctgat	1550
caccaggctc ctgtctccac aggatccag gggatggggg ttaagcttg	1600
ctcctggcgg tccaccctgc tggaaccagt tgaaaccctg gtaatggtga	1650
ccctttgagc gagccaaggc tgggtggtag atgaccatct cttgtccaac	1700

-continued

agggtcccaga gcagtggata tgtctgggtcc tcctagtagc acagagggtgt	1750
gttctgtgtgt ggtggcaggg acttagggaa tcctaccact ctgctggatt	1800
tggaaccccc taggctgacg cggacgtatg cagaggctct caaggccagg	1850
ccccacaggg aggtggaggg gctccggcgg ccacagcctg aattcatgaa	1900
cctggcaggg actttgccat agctcatctg aaaacagata ttatgcttcc	1950
cacaacctct cctgggcca ggtgtggctg agcaccaggg atggagccac	2000
acataagga caaatgagt caccgtccta cctagtcttt cctcacctcc	2050
tgaactcaca caacaatgcc agtctccac tggaggctgt atcccctcag	2100
aggagccaag gaatgtcttc ccctgagatg ccaccactat taatttcccc	2150
atatgcttca accacccct tgctcaaaaa accaataccc acacttacct	2200
taatacaaac atcccagcaa cagcacatgg caggccattg ctgagggcac	2250
agggtgcttta ttggagaggg gatgtgggca ggggataagg aaggttcccc	2300
cattccagga ggatgggaac agtcctggct gccctgaca gtggggatat	2350
gcaaggggct ctggccaggc cacagtccaa atgggaagac accagtcagt	2400
cacaaaagtc gggagcgcca cacaaacctg gctataaggc ccaggaacca	2450
tataggagcc tgagacaggt cccctgcaca ttcattatta aactatacag	2500
gatgaggctg tacatgagtt aattacaaaa gagtcattat tacaaaaatc	2550
tgtacacaca ttgaaaaaac tcacaaaatt gtcattatg tatcacaagt	2600
tgctagaccc aaaaatattaa aaatgggata aaatnnnttt aaaaaaaaaa	2650
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa	2684

<210> SEQ ID NO 34
<211> LENGTH: 402
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 34

Met Arg Arg Arg Leu Arg Leu Arg Arg Asp Ala Leu Leu Thr Leu	1	5	10	15
Leu Leu Gly Ala Ser Leu Gly Leu Leu Leu Tyr Ala Gln Arg Asp	20	25	30	
Gly Ala Ala Pro Thr Ala Ser Ala Pro Arg Gly Arg Gly Arg Ala	35	40	45	
Ala Pro Arg Pro Thr Pro Gly Pro Arg Ala Phe Gln Leu Pro Asp	50	55	60	
Ala Gly Ala Ala Pro Pro Ala Tyr Glu Gly Asp Thr Pro Ala Pro	65	70	75	
Pro Thr Pro Thr Gly Pro Phe Asp Phe Ala Arg Tyr Leu Arg Ala	80	85	90	
Lys Asp Gln Arg Arg Phe Pro Leu Leu Ile Asn Gln Pro His Lys	95	100	105	
Cys Arg Gly Asp Gly Ala Pro Gly Gly Arg Pro Asp Leu Leu Ile	110	115	120	
Ala Val Lys Ser Val Ala Glu Asp Phe Glu Arg Arg Gln Ala Val	125	130	135	

-continued

Arg	Gln	Thr	Trp	Gly	Ala	Glu	Gly	Arg	Val	Gln	Gly	Ala	Leu	Val
				140					145					150
Arg	Arg	Val	Phe	Leu	Leu	Gly	Val	Pro	Arg	Gly	Ala	Gly	Ser	Gly
				155					160					165
Gly	Ala	Asp	Glu	Val	Gly	Glu	Gly	Ala	Arg	Thr	His	Trp	Arg	Ala
				170					175					180
Leu	Leu	Arg	Ala	Glu	Ser	Leu	Ala	Tyr	Ala	Asp	Ile	Leu	Leu	Trp
				185					190					195
Ala	Phe	Asp	Asp	Thr	Phe	Phe	Asn	Leu	Thr	Leu	Lys	Glu	Ile	His
				200					205					210
Phe	Leu	Ala	Trp	Ala	Ser	Ala	Phe	Cys	Pro	Asp	Val	Arg	Phe	Val
				215					220					225
Phe	Lys	Gly	Asp	Ala	Asp	Val	Phe	Val	Asn	Val	Gly	Asn	Leu	Leu
				230					235					240
Glu	Phe	Leu	Ala	Pro	Arg	Asp	Pro	Ala	Gln	Asp	Leu	Leu	Ala	Gly
				245					250					255
Asp	Val	Ile	Val	His	Ala	Arg	Pro	Ile	Arg	Thr	Arg	Ala	Ser	Lys
				260					265					270
Tyr	Tyr	Ile	Pro	Glu	Ala	Val	Tyr	Gly	Leu	Pro	Ala	Tyr	Pro	Ala
				275					280					285
Tyr	Ala	Gly	Gly	Gly	Gly	Phe	Val	Leu	Ser	Gly	Ala	Thr	Leu	His
				290					295					300
Arg	Leu	Ala	Gly	Ala	Cys	Ala	Gln	Val	Glu	Leu	Phe	Pro	Ile	Asp
				305					310					315
Asp	Val	Phe	Leu	Gly	Met	Cys	Leu	Gln	Arg	Leu	Arg	Leu	Thr	Pro
				320					325					330
Glu	Pro	His	Pro	Ala	Phe	Arg	Thr	Phe	Gly	Ile	Pro	Gln	Pro	Ser
				335					340					345
Ala	Ala	Pro	His	Leu	Ser	Thr	Phe	Asp	Pro	Cys	Phe	Tyr	Arg	Glu
				350					355					360
Leu	Val	Val	Val	His	Gly	Leu	Ser	Ala	Ala	Asp	Ile	Trp	Leu	Met
				365					370					375
Trp	Arg	Leu	Leu	His	Gly	Pro	His	Gly	Pro	Ala	Cys	Ala	His	Pro
				380					385					390
Gln	Pro	Val	Ala	Ala	Gly	Pro	Phe	Gln	Trp	Asp	Ser			
				395					400					

<210> SEQ ID NO 35
<211> LENGTH: 1643
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 35

agcagcctct gccgaccgc gctcgtgcgg accccaggac cgggcgcggg	50
acgcgtgcgt ccagcctccg gcgctgcgga gaccgcggc tgggtccggg	100
gaggcccaaa acccgcccc gccagaacc cgcccaaat tccacctcc	150
tccagaagcc ccgcccactc ccgagcccg agagctccgc gcacctggg	200
gccatccgcc ctggtccgc tgcacgagct ccacgccgt accccggcgt	250
cacgctcagc ccgcggtgct cgcacacctg agactcatct cgcttcgacc	300
ccgcgcgcgc cgcgcgccg catcctgagc acggagacag tctccagctg	350

-continued

ccgttcacgc	ttcctcccca	gccttccgca	gcccaccagg	gaaggggcgg	400
taggagtggc	cttttaccaa	agggaccggc	gatgctctgc	aggetgtgct	450
ggctggtctc	gtacagcttg	gctgtgctgt	tgctcggctg	cctgctcttc	500
ctgaggaagg	cggccaagcc	cgcaggagac	cccacggccc	accagccttt	550
ctgggctccc	ccaacacccc	gtcacagccg	gtgtccaccc	aaccacacag	600
tgtctagcgc	ctctctgtcc	ctgcctagcc	gtcaccgtct	cttcttgacc	650
tatcgtcact	gccgaaatth	ctctatcttg	ctggagcctt	caggctgttc	700
caaggatacc	ttcttgctcc	tggccatcaa	gtcacagcct	ggtcacgtgg	750
agcgacgtgc	ggctatccgc	agcacgtggg	gcaggggtggg	gggatgggct	800
aggggccggc	agctgaagct	ggtgttcctc	ctaggggtgg	caggatccgc	850
tccccagcc	cagctgctgg	cctatgagag	tagggagtth	gatgacatcc	900
tccagtggga	cttcactgag	gacttcttca	acctgacgct	caaggagctg	950
cacctgcagc	gctgggtggt	ggctgcctgc	ccccaggccc	atttcatgct	1000
aaaggagat	gacgatgtct	ttgtccacgt	ccccaacgtg	ttagagtthc	1050
tggatggctg	ggaccacgcc	caggacctcc	tgggtgggaga	tgtcatccgc	1100
caagccctgc	ccaacaggaa	cactaaggtc	aaatacttca	tcccaccctc	1150
aatgtacagg	gccaccact	accacccta	tgctgggtggg	ggaggatatg	1200
tcatgtccag	agccacagtg	cggcgccctc	aggctatcat	ggaagatgct	1250
gaactcttcc	ccattgatga	tgtctttgtg	ggtatgtgcc	tgaggaggct	1300
ggggctgagc	cctatgcacc	atgctggctt	caagacatth	ggaatccggc	1350
ggcccctgga	ccccttagac	ccctgcctgt	atagggggct	cctgctggtt	1400
caccgcctca	gccccctcga	gatgtggacc	atgtgggcac	tggtgacaga	1450
tgaggggctc	aagtgtgcag	ctggcccat	acccagcgc	tgaagggtgg	1500
gttgggcaac	agcctgagag	tggactcagt	gttgattctc	tatcgtgatg	1550
cgaaattgat	gcctgctgct	ctacagaaaa	tgccaacttg	gttttttaac	1600
tcctctcacc	ctgttagctc	tgattaataa	caactgcaacc	caa	1643

<210> SEQ ID NO 36
<211> LENGTH: 378
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 36

Met	Leu	Pro	Pro	Gln	Pro	Ser	Ala	Ala	His	Gln	Gly	Arg	Gly	Gly
1				5					10					15
Arg	Ser	Gly	Leu	Leu	Pro	Lys	Gly	Pro	Ala	Met	Leu	Cys	Arg	Leu
			20						25					30
Cys	Trp	Leu	Val	Ser	Tyr	Ser	Leu	Ala	Val	Leu	Leu	Leu	Gly	Cys
			35						40					45
Leu	Leu	Phe	Leu	Arg	Lys	Ala	Ala	Lys	Pro	Ala	Gly	Asp	Pro	Thr
			50						55					60
Ala	His	Gln	Pro	Phe	Trp	Ala	Pro	Pro	Thr	Pro	Arg	His	Ser	Arg
			65						70					75
Cys	Pro	Pro	Asn	His	Thr	Val	Ser	Ser	Ala	Ser	Leu	Ser	Leu	Pro

-continued

80										85					90				
Ser	Arg	His	Arg	Leu	Phe	Leu	Thr	Tyr	Arg	His	Cys	Arg	Asn	Phe					
				95					100					105					
Ser	Ile	Leu	Leu	Glu	Pro	Ser	Gly	Cys	Ser	Lys	Asp	Thr	Phe	Leu					
				110					115					120					
Leu	Leu	Ala	Ile	Lys	Ser	Gln	Pro	Gly	His	Val	Glu	Arg	Arg	Ala					
				125					130					135					
Ala	Ile	Arg	Ser	Thr	Trp	Gly	Arg	Val	Gly	Gly	Trp	Ala	Arg	Gly					
				140					145					150					
Arg	Gln	Leu	Lys	Leu	Val	Phe	Leu	Leu	Gly	Val	Ala	Gly	Ser	Ala					
				155					160					165					
Pro	Pro	Ala	Gln	Leu	Leu	Ala	Tyr	Glu	Ser	Arg	Glu	Phe	Asp	Asp					
				170					175					180					
Ile	Leu	Gln	Trp	Asp	Phe	Thr	Glu	Asp	Phe	Phe	Asn	Leu	Thr	Leu					
				185					190					195					
Lys	Glu	Leu	His	Leu	Gln	Arg	Trp	Val	Val	Ala	Ala	Cys	Pro	Gln					
				200					205					210					
Ala	His	Phe	Met	Leu	Lys	Gly	Asp	Asp	Asp	Val	Phe	Val	His	Val					
				215					220					225					
Pro	Asn	Val	Leu	Glu	Phe	Leu	Asp	Gly	Trp	Asp	Pro	Ala	Gln	Asp					
				230					235					240					
Leu	Leu	Val	Gly	Asp	Val	Ile	Arg	Gln	Ala	Leu	Pro	Asn	Arg	Asn					
				245					250					255					
Thr	Lys	Val	Lys	Tyr	Phe	Ile	Pro	Pro	Ser	Met	Tyr	Arg	Ala	Thr					
				260					265					270					
His	Tyr	Pro	Pro	Tyr	Ala	Gly	Gly	Gly	Gly	Tyr	Val	Met	Ser	Arg					
				275					280					285					
Ala	Thr	Val	Arg	Arg	Leu	Gln	Ala	Ile	Met	Glu	Asp	Ala	Glu	Leu					
				290					295					300					
Phe	Pro	Ile	Asp	Asp	Val	Phe	Val	Gly	Met	Cys	Leu	Arg	Arg	Leu					
				305					310					315					
Gly	Leu	Ser	Pro	Met	His	His	Ala	Gly	Phe	Lys	Thr	Phe	Gly	Ile					
				320					325					330					
Arg	Arg	Pro	Leu	Asp	Pro	Leu	Asp	Pro	Cys	Leu	Tyr	Arg	Gly	Leu					
				335					340					345					
Leu	Leu	Val	His	Arg	Leu	Ser	Pro	Leu	Glu	Met	Trp	Thr	Met	Trp					
				350					355					360					
Ala	Leu	Val	Thr	Asp	Glu	Gly	Leu	Lys	Cys	Ala	Ala	Gly	Pro	Ile					
				365					370					375					

Pro Gln Arg

<210> SEQ ID NO 37
<211> LENGTH: 1226
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 37

atgaaagtga taatcaggca gcccaaatga ttgttaataa ggatcaaatg	50
agatcgtgta tgtgggtcca atcaattgat tctacacaaa ggagcctggg	100
gagggggccat ggtgccaatg cacttactgg ggagactgga gaagccgctt	150
ctcctcctgt gctgcgcctc cttcctactg gggctggcctt tgctgggcatt	200

-continued

aaagacggac atcacccccc ttgcttattt ctttctcaca ttgggtggct 250
tcttcttggt tgctatctc ctggtccggt ttctggaatg ggggttcgg 300
tcccagctcc aatcaatgca gactgagagc ccagggccct caggcaatgc 350
acgggacaat gaagcctttg aagtgccagt ctatgaagag gccgtggtg 400
gactagaatc ccagtgcgcg ccccaagagt tggaccaacc acccccctac 450
agcactgttg tgataccccc agcacctgag gaggaacaac ctagccatcc 500
agaggggtcc aggagagcca aactggaaca gaggcgaatg gcctcagag 550
ggtccatggc ccaggaagga agccctggaa gagctccaat caaccttcgg 600
cttcggggac cacgggctgt gtccactgct cctgatctgc agagcttggc 650
ggcagtcgcc acattagagc ctctgactcc accccctgcc tatgatgtct 700
gctttggtca ccctgatgat gatagtgttt tttatgagga caactgggca 750
ccccctaaa tgactctccc aagatttctc ttctctccac accagacctc 800
gttcatttga ctaacatttt ccagcgccta ctatgtgtca gaaacaagt 850
ttctgcctg gacatcataa atggggactt ggaccctgag gagagtcagg 900
ccacggttag cccttccag ctgagatatg ggtggcataa tttgagtctt 950
ctggcaacat ttggtgacct accccatctc caatatttcc agcgtagat 1000
tgaggatgag gtagggaggt gatccagaga aggcggagaa ggaagaagta 1050
acctctgagt ggcggctatt gcttctgttc caggtgctgt tcgagctgtt 1100
agaaccctta ggcttgacag ctttgtgagt tattattgaa aaatgaggat 1150
tccaagatgc agaggagtgt gataatgtgc acgagggcac actgctagta 1200
aataacatta aaataactgg aatgaa 1226

<210> SEQ ID NO 38
<211> LENGTH: 216
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 38

Met Val Pro Met His Leu Leu Gly Arg Leu Glu Lys Pro Leu Leu
1 5 10 15
Leu Leu Cys Cys Ala Ser Phe Leu Leu Gly Leu Ala Leu Leu Gly
20 25 30
Ile Lys Thr Asp Ile Thr Pro Val Ala Tyr Phe Phe Leu Thr Leu
35 40 45
Gly Gly Phe Phe Leu Phe Ala Tyr Leu Leu Val Arg Phe Leu Glu
50 55 60
Trp Gly Leu Arg Ser Gln Leu Gln Ser Met Gln Thr Glu Ser Pro
65 70 75
Gly Pro Ser Gly Asn Ala Arg Asp Asn Glu Ala Phe Glu Val Pro
80 85 90
Val Tyr Glu Glu Ala Val Val Gly Leu Glu Ser Gln Cys Arg Pro
95 100 105
Gln Glu Leu Asp Gln Pro Pro Pro Tyr Ser Thr Val Val Ile Pro
110 115 120
Pro Ala Pro Glu Glu Glu Gln Pro Ser His Pro Glu Gly Ser Arg

-continued

125	130	135
Arg Ala Lys Leu Glu Gln Arg Arg Met	Ala Ser Glu Gly Ser Met	
140	145	150
Ala Gln Glu Gly Ser Pro Gly Arg Ala	Pro Ile Asn Leu Arg Leu	
155	160	165
Arg Gly Pro Arg Ala Val Ser Thr Ala	Pro Asp Leu Gln Ser Leu	
170	175	180
Ala Ala Val Pro Thr Leu Glu Pro Leu Thr	Pro Pro Pro Ala Tyr	
185	190	195
Asp Val Cys Phe Gly His Pro Asp Asp	Asp Ser Val Phe Tyr Glu	
200	205	210
Asp Asn Trp Ala Pro Pro		
215		

<210> SEQ ID NO 39
<211> LENGTH: 2770
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 39

cccacgcgtc cggcggctac acacctaggt gcggtgggct tcgggtgggg	50
ggcctgcagc tagctgatgg caagggagga atagcagggg tggggattgt	100
ggtgtgcgag aggtcccgcg gacggggggc tcgggggtct cttcagacga	150
gattcccttc aggccttgggc cgggtccctt cgcacggaga tcccaatgaa	200
cgcggggccc tggaggccgg tggttggggc ttctccgcgt cggggatggg	250
gccgtaccc tagccgttt ccagcgcctc agtcgggtcc ccatgccctc	300
agaggtggcc cggggcaagc gcgccccct cttcttcgct gcggtggcca	350
tcgtgctggg gctaccgctc tggtggaaga ccacggagac ctaccgggcc	400
tcgttgccct actcccagat cagtggcctg aatgcccttc agctccgcct	450
catggtgcct gtcactgtcg tgtttacgcg ggagtcagtg cccttggaag	500
accaggagaa gctgcccttc accgttgtgc atgaaagaga gattcctctg	550
aaatacaaaa tgaaaatcaa atgccgtttc cagaaggcct atcggagggc	600
tttgaccat gaggaggagg ccctgtcatc gggcagtgtg caagaggcag	650
aagccatggt agatgagcct caggaacaag cggagggctc cctgactgtg	700
tacgtgatat ctgaacactc ctcaacttct cccaggaca tgatgagcta	750
cattggggccc aagaggacag cagtgggtcg ggggataatg caccgggagg	800
cctttaacat cattggccgc cgcatagtcc aggtggccca ggccatgtct	850
ttgactgagg atgtgcttgc tgctgctctg gctgaccacc ttccagagga	900
caagtggagc gctgagaaga ggccgcctct caagtccagc ttgggctatg	950
agatcacctt cagtttactc aaccagacc ccaagtccca tgatgtctac	1000
tgggacattg agggggctgt ccggcgctat gtgcaacctt tcctgaatgc	1050
cctcggtgcc gctggcaact tctctgtgga ctctcagatt ctttactatg	1100
caatgttggg ggtgaatccc cgctttgact cagcttcctc cagctactat	1150
ttggacatgc acagcctccc ccatgtcatc aaccagtggt agtccgggct	1200

-continued

gggatccagt gctgcctcct tgtaccctgt gctcaacttt ctactctacg	1250
tgacctgagct tgcacactca ccgctgtaca ttcaggacaa ggatggcgct	1300
ccagtggcca ccaatgcctt ccatagtccc cgctgggggtg gcattatggt	1350
atataatggt gactccaaaa cctataatgc ctcagtgtcg ccagtgagag	1400
tcgaggtgga catggtgcga gtgatggagg tgttcctggc acagttgcgg	1450
ttgtcttttg ggattgctca gcccagctg cctccaaaat gcctgctttc	1500
agggcctacg agtgaagggc taatgacctg ggagctagac cgctgctct	1550
gggctcggtc agtggagaac ctggccacag ccaccaccac ccttacctcc	1600
ctggcgcgagc ttctgggcaa gatcagcaac attgtcatta aggacgacgt	1650
ggcatctgag gtgtacaagg ctgtagctgc cgtccagaag tcggcagaag	1700
agttggcgtc tgggcacctg gcatctgcct ttgtcgccag ccaggaagct	1750
gtgacatcct ctgagcttgc cttctttgac ccgtcactcc tccacctcct	1800
ttatttccct gatgaccaga agtttgccat ctacatccca ctcttcctgc	1850
ctatggctgt gccatcctc ctgtccctgg tcaagatctt cctggagacc	1900
cgcaagtccct ggagaaagcc tgagaagaca gactgagcag ggcagcacct	1950
ccataggaag ccttcctttc tggccaaggt gggcggtgtt agattgtgag	2000
gcacgtacat ggggcctgcc ggaatgactt aaatatttgt ctccagtctc	2050
cactgtttgc tctccagcaa ccaaagtaca acactccaag atgggttcat	2100
cttttcttcc tttccattc acctggctca atcctcctcc accaccaggg	2150
gcctcaaaag gcacatcatc cgggtctcct tatcttgttt gataaggctg	2200
ctgcctgtct ccctctgtgg caaggactgt ttgttctttt gccccatttc	2250
tcaacatagc acacttgtgc actgagagga gggagcatta tgggaaagtc	2300
cctgccttcc acacctctct ctagtccctg tgggacagcc ctagcccctg	2350
ctgtcatgaa ggggccaggc attggtcacc tgtgggacct tctccctcac	2400
tcccctccct cctagttggc tttgtctgtc aggtgcagtc tggcgggagt	2450
ccaggaggca gcagctcagg acatggtgct gtgtgtgtgt gtgtgtgtgt	2500
gtgtgtgtgt gtgtgtgtca gaggttcag aaagttccag atttgaatc	2550
aaacagtccct gaattcaaat ccttgttttt gcacttattg tctggagagc	2600
tttgataag gtattgaatc tctctgagcc tcagtttttc atttgttcaa	2650
atggcactga tgatgtctcc cttaacaagat ggttgtagag agtaaatgtg	2700
atcagcatgt aaagtgtctg gcgtgtagta ggctcttaat aaacactggc	2750
tgaatatgaa ttggaatgat	2770

<210> SEQ ID NO 40
<211> LENGTH: 547
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 40

Met	Pro	Ser	Glu	Val	Ala	Arg	Gly	Lys	Arg	Ala	Ala	Leu	Phe	Phe
1				5				10					15	
Ala	Ala	Val	Ala	Ile	Val	Leu	Gly	Leu	Pro	Leu	Trp	Trp	Lys	Thr

-continued

20										25					30				
Thr	Glu	Thr	Tyr	Arg	Ala	Ser	Leu	Pro	Tyr	Ser	Gln	Ile	Ser	Gly					
				35					40					45					
Leu	Asn	Ala	Leu	Gln	Leu	Arg	Leu	Met	Val	Pro	Val	Thr	Val	Val					
				50					55					60					
Phe	Thr	Arg	Glu	Ser	Val	Pro	Leu	Asp	Asp	Gln	Glu	Lys	Leu	Pro					
				65					70					75					
Phe	Thr	Val	Val	His	Glu	Arg	Glu	Ile	Pro	Leu	Lys	Tyr	Lys	Met					
				80					85					90					
Lys	Ile	Lys	Cys	Arg	Phe	Gln	Lys	Ala	Tyr	Arg	Arg	Ala	Leu	Asp					
				95					100					105					
His	Glu	Glu	Glu	Ala	Leu	Ser	Ser	Gly	Ser	Val	Gln	Glu	Ala	Glu					
				110					115					120					
Ala	Met	Leu	Asp	Glu	Pro	Gln	Glu	Gln	Ala	Glu	Gly	Ser	Leu	Thr					
				125					130					135					
Val	Tyr	Val	Ile	Ser	Glu	His	Ser	Ser	Leu	Leu	Pro	Gln	Asp	Met					
				140					145					150					
Met	Ser	Tyr	Ile	Gly	Pro	Lys	Arg	Thr	Ala	Val	Val	Arg	Gly	Ile					
				155					160					165					
Met	His	Arg	Glu	Ala	Phe	Asn	Ile	Ile	Gly	Arg	Arg	Ile	Val	Gln					
				170					175					180					
Val	Ala	Gln	Ala	Met	Ser	Leu	Thr	Glu	Asp	Val	Leu	Ala	Ala	Ala					
				185					190					195					
Leu	Ala	Asp	His	Leu	Pro	Glu	Asp	Lys	Trp	Ser	Ala	Glu	Lys	Arg					
				200					205					210					
Arg	Pro	Leu	Lys	Ser	Ser	Leu	Gly	Tyr	Glu	Ile	Thr	Phe	Ser	Leu					
				215					220					225					
Leu	Asn	Pro	Asp	Pro	Lys	Ser	His	Asp	Val	Tyr	Trp	Asp	Ile	Glu					
				230					235					240					
Gly	Ala	Val	Arg	Arg	Tyr	Val	Gln	Pro	Phe	Leu	Asn	Ala	Leu	Gly					
				245					250					255					
Ala	Ala	Gly	Asn	Phe	Ser	Val	Asp	Ser	Gln	Ile	Leu	Tyr	Tyr	Ala					
				260					265					270					
Met	Leu	Gly	Val	Asn	Pro	Arg	Phe	Asp	Ser	Ala	Ser	Ser	Ser	Tyr					
				275					280					285					
Tyr	Leu	Asp	Met	His	Ser	Leu	Pro	His	Val	Ile	Asn	Pro	Val	Glu					
				290					295					300					
Ser	Arg	Leu	Gly	Ser	Ser	Ala	Ala	Ser	Leu	Tyr	Pro	Val	Leu	Asn					
				305					310					315					
Phe	Leu	Leu	Tyr	Val	Pro	Glu	Leu	Ala	His	Ser	Pro	Leu	Tyr	Ile					
				320					325					330					
Gln	Asp	Lys	Asp	Gly	Ala	Pro	Val	Ala	Thr	Asn	Ala	Phe	His	Ser					
				335					340					345					
Pro	Arg	Trp	Gly	Gly	Ile	Met	Val	Tyr	Asn	Val	Asp	Ser	Lys	Thr					
				350					355					360					
Tyr	Asn	Ala	Ser	Val	Leu	Pro	Val	Arg	Val	Glu	Val	Asp	Met	Val					
				365					370					375					
Arg	Val	Met	Glu	Val	Phe	Leu	Ala	Gln	Leu	Arg	Leu	Leu	Phe	Gly					
				380					385					390					
Ile	Ala	Gln	Pro	Gln	Leu	Pro	Pro	Lys	Cys	Leu	Leu	Ser	Gly	Pro					
				395					400					405					

-continued

Thr Ser Glu Gly Leu Met Thr Trp Glu Leu Asp Arg Leu Leu Trp
410 415 420

Ala Arg Ser Val Glu Asn Leu Ala Thr Ala Thr Thr Thr Leu Thr
425 430 435

Ser Leu Ala Gln Leu Leu Gly Lys Ile Ser Asn Ile Val Ile Lys
440 445 450

Asp Asp Val Ala Ser Glu Val Tyr Lys Ala Val Ala Ala Val Gln
455 460 465

Lys Ser Ala Glu Glu Leu Ala Ser Gly His Leu Ala Ser Ala Phe
470 475 480

Val Ala Ser Gln Glu Ala Val Thr Ser Ser Glu Leu Ala Phe Phe
485 490 495

Asp Pro Ser Leu Leu His Leu Leu Tyr Phe Pro Asp Asp Gln Lys
500 505 510

Phe Ala Ile Tyr Ile Pro Leu Phe Leu Pro Met Ala Val Pro Ile
515 520 525

Leu Leu Ser Leu Val Lys Ile Phe Leu Glu Thr Arg Lys Ser Trp
530 535 540

Arg Lys Pro Glu Lys Thr Asp
545

<210> SEQ ID NO 41
<211> LENGTH: 1964
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 41

ccagctgcag agaggaggag gtgagctgca gagaagagga ggttggtgtg 50

gagcacaggc agcaccgagc ctgccccgtg agctgagggc ctgcagtctg 100

cggttggaat caggatagac accaaggcag gacccccaga gatgctgaag 150

cctcttttga aagcagcagt ggccccaca tggccatgct ccatgccgcc 200

ccgccgcccg tgggacagag aggtggcac gttgcaggtc ctgggagcgc 250

tggtgtgtgt gtggctgggc tccgtggctc ttatctgcct cctgtggcaa 300

gtgccccctc ctccacctg gggccagggt cagcccaagg acgtgccag 350

gtcctgggag catggctcca gcccgcttg ggagccccctg gaagcagagg 400

ccaggcagca gagggactcc tgccagcttg tccttgtgga aagcatcccc 450

caggacctgc catctgcagc cggcagcccc tctgcccagc ctctgggcca 500

ggcctgggtg cagctgctgg aactgcccc ggagagcgtc cacgtggctt 550

catactactg gtccctcaca gggcctgaca tcgggggtcaa cgactcgtct 600

tcccagctgg gagaggctct tctgcagaag ctgcagcagc tgctgggcag 650

gaacatttcc ctggctgtgg ccaccagcag cccgacactg gccaggacat 700

ccaccgacct gcaggttctg gctgcccag gtgcccattgt acgacagggtg 750

ccctgggggc ggctcaccag ggggtgttttg cactccaaat tctgggttgt 800

ggatggacgg cacatataca tgggcagtgc caacatggac tggcgggtctc 850

tgacgcaggt gaaggagctt ggcgctgtca tctataactg cagccacctg 900

gcccaagacc tggagaagac cttccagacc tactgggtac tgggggtgcc 950

-continued

caaggctgtc ctccccaaaa cctggcctca gaacttctca tctcacttca	1000
accgttttcca gcccttccac ggctcttttg atggggtgcc caccactgcc	1050
tactttctcag cgtcgccacc agcactctgt ccccagggcc gcacccggga	1100
cctggaggcg ctgctggcgg tgatggggag cgcccaggag ttcatctatg	1150
cctccgtgat ggagtatttc cccaccacgc gcttcagcca cccccgagg	1200
tactggccgg tgctggacaa cgcgtgcgg gcggcagcct tcggcaaggg	1250
cgtagcgcgtg cgctgctgg tcggctgcgg actcaacacg gacccaccaa	1300
tgttccccta cctgcggtcc ctgcaggcgc tcagcaacct cgcgccaac	1350
gtctctgtgg acgtgaaagt cttcatcgtg ccggtgggga accattccaa	1400
catcccattc agcagggtga accacagcaa gttcatggtc acggagaagg	1450
cagcctacat aggcacctcc aactggtcgg aggattactt cagcagcacg	1500
gcgggggtgg gcttggtggt caccagagc cctggcgcgc agcccgcggg	1550
ggccacggtg caggagcagc tgccgcagct ctttgagcgg gactggagtt	1600
cgcgctacgc cgtcggcctg gacggacagg ctccgggcca ggactgcgtt	1650
tggcagggct gaggggggcc tctttttctc tcggcgacct cgccccgcac	1700
gcgcctccc ctctgacccc ggcttgggt tcagccgctt cctcccgcaa	1750
gcagcccggg tccgcactgc gccaggagcc gctgcgacc gcccgggcgt	1800
cgcaaacgc ccgcctgctc tctgatttcc gagtccagcc cccctgagc	1850
cccacctct ccagggagcc ctccaggaag ccccttccct gactcctggc	1900
ccacaggcca ggcctaaaaa aaactcgtgg cttcaaaaaa aaaaaaaaaa	1950
aaaaaaaaaa aaaa	1964

<210> SEQ ID NO 42
<211> LENGTH: 489
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 42

Met Pro Pro Arg Arg Pro Trp Asp Arg Glu Ala Gly Thr Leu Gln	
1 5 10 15	
Val Leu Gly Ala Leu Ala Val Leu Trp Leu Gly Ser Val Ala Leu	
20 25 30	
Ile Cys Leu Leu Trp Gln Val Pro Arg Pro Pro Thr Trp Gly Gln	
35 40 45	
Val Gln Pro Lys Asp Val Pro Arg Ser Trp Glu His Gly Ser Ser	
50 55 60	
Pro Ala Trp Glu Pro Leu Glu Ala Glu Ala Arg Gln Gln Arg Asp	
65 70 75	
Ser Cys Gln Leu Val Leu Val Glu Ser Ile Pro Gln Asp Leu Pro	
80 85 90	
Ser Ala Ala Gly Ser Pro Ser Ala Gln Pro Leu Gly Gln Ala Trp	
95 100 105	
Leu Gln Leu Leu Asp Thr Ala Gln Glu Ser Val His Val Ala Ser	
110 115 120	
Tyr Tyr Trp Ser Leu Thr Gly Pro Asp Ile Gly Val Asn Asp Ser	

-continued

125					130					135				
Ser	Ser	Gln	Leu	Gly	Glu	Ala	Leu	Leu	Gln	Lys	Leu	Gln	Gln	Leu
				140					145					150
Leu	Gly	Arg	Asn	Ile	Ser	Leu	Ala	Val	Ala	Thr	Ser	Ser	Pro	Thr
				155					160					165
Leu	Ala	Arg	Thr	Ser	Thr	Asp	Leu	Gln	Val	Leu	Ala	Ala	Arg	Gly
				170					175					180
Ala	His	Val	Arg	Gln	Val	Pro	Met	Gly	Arg	Leu	Thr	Arg	Gly	Val
				185					190					195
Leu	His	Ser	Lys	Phe	Trp	Val	Val	Asp	Gly	Arg	His	Ile	Tyr	Met
				200					205					210
Gly	Ser	Ala	Asn	Met	Asp	Trp	Arg	Ser	Leu	Thr	Gln	Val	Lys	Glu
				215					220					225
Leu	Gly	Ala	Val	Ile	Tyr	Asn	Cys	Ser	His	Leu	Ala	Gln	Asp	Leu
				230					235					240
Glu	Lys	Thr	Phe	Gln	Thr	Tyr	Trp	Val	Leu	Gly	Val	Pro	Lys	Ala
				245					250					255
Val	Leu	Pro	Lys	Thr	Trp	Pro	Gln	Asn	Phe	Ser	Ser	His	Phe	Asn
				260					265					270
Arg	Phe	Gln	Pro	Phe	His	Gly	Leu	Phe	Asp	Gly	Val	Pro	Thr	Thr
				275					280					285
Ala	Tyr	Phe	Ser	Ala	Ser	Pro	Pro	Ala	Leu	Cys	Pro	Gln	Gly	Arg
				290					295					300
Thr	Arg	Asp	Leu	Glu	Ala	Leu	Leu	Ala	Val	Met	Gly	Ser	Ala	Gln
				305					310					315
Glu	Phe	Ile	Tyr	Ala	Ser	Val	Met	Glu	Tyr	Phe	Pro	Thr	Thr	Arg
				320					325					330
Phe	Ser	His	Pro	Pro	Arg	Tyr	Trp	Pro	Val	Leu	Asp	Asn	Ala	Leu
				335					340					345
Arg	Ala	Ala	Ala	Phe	Gly	Lys	Gly	Val	Arg	Val	Arg	Leu	Leu	Val
				350					355					360
Gly	Cys	Gly	Leu	Asn	Thr	Asp	Pro	Thr	Met	Phe	Pro	Tyr	Leu	Arg
				365					370					375
Ser	Leu	Gln	Ala	Leu	Ser	Asn	Pro	Ala	Ala	Asn	Val	Ser	Val	Asp
				380					385					390
Val	Lys	Val	Phe	Ile	Val	Pro	Val	Gly	Asn	His	Ser	Asn	Ile	Pro
				395					400					405
Phe	Ser	Arg	Val	Asn	His	Ser	Lys	Phe	Met	Val	Thr	Glu	Lys	Ala
				410					415					420
Ala	Tyr	Ile	Gly	Thr	Ser	Asn	Trp	Ser	Glu	Asp	Tyr	Phe	Ser	Ser
				425					430					435
Thr	Ala	Gly	Val	Gly	Leu	Val	Val	Thr	Gln	Ser	Pro	Gly	Ala	Gln
				440					445					450
Pro	Ala	Gly	Ala	Thr	Val	Gln	Glu	Gln	Leu	Arg	Gln	Leu	Phe	Glu
				455					460					465
Arg	Asp	Trp	Ser	Ser	Arg	Tyr	Ala	Val	Gly	Leu	Asp	Gly	Gln	Ala
				470					475					480
Pro	Gly	Gln	Asp	Cys	Val	Trp	Gln	Gly						
				485										

-continued

<211> LENGTH: 1130
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 43
gggcctggcg atccggatcc cgcaggcgcg ctggctgcgc tgcccggctg 50
tctgtcgta tgggtggggc ctgggtgtat ctgggtggcg cagttttgct 100
catcggcctg atcctcttcc tgactcgag cgggggtcgg gcggcagcag 150
ctgacggaga accactgcac aatgaggaag agagggcagg agcaggccag 200
gtaggccgct ctttgcccca ggagtctgaa gaacagagaa ctggaagcag 250
accccggcgt cggagggact tgggcagccg tctacaggcc cagcgtcgag 300
cccacgagat ggcctgggaa gacggggatg agaatgtggg tcaaaactgtt 350
attccagccc aggaggaaga aggcattgag aagccagcag aagttcaccc 400
aacagggaaa attggagcca agaaactacg gaagctagag gaaaaacagg 450
ctcgaaggcg tcagcgagag gcagaggagg ctgaacgtga agaacggaaa 500
cgcctagagt cccaacgtga ggccgaatgg aagaaggaa aggaacggct 550
tcgcctgaag gaagaacaga aggaggagga agagaggaa gctcaggagg 600
agcaggcccg gcgggatcac gaggagtacc tgaaactgaa ggaggccttc 650
gtggtagaag aagaaggtgt tagcgaaacc atgactgagg agcagtctca 700
cagcttcctg acagaattca tcaattacat caagaagtcc aaggttgtgc 750
ttttggaaga tctggctttc cagatgggcc taaggactca ggacgccata 800
aaccgcatcc aggacctgct gacggagggg actctaacag gtgtgattga 850
cgaccggggc aagtttatct acataacccc agaggaactg gctgccgtgg 900
ccaatttcat ccgacagcgg gcccggtgt ccatcacaga gcttgcccag 950
gccagcaact ccctcatctc ctggggccag gacctcctg cccaggcttc 1000
agcctgactc cagtccttcc ttgagtgtat cctgtggcct acatgtgtct 1050
tcacctctcc ctaatgccgt cttggggcag ggatggaata tgaccagaaa 1100
gttgtggatt aaaggcctgt gaatactgaa 1130

<210> SEQ ID NO 44
<211> LENGTH: 315
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 44
Met Val Gly Pro Trp Val Tyr Leu Val Ala Ala Val Leu Leu Ile
1 5 10 15
Gly Leu Ile Leu Phe Leu Thr Arg Ser Arg Gly Arg Ala Ala Ala
20 25 30
Ala Asp Gly Glu Pro Leu His Asn Glu Glu Glu Arg Ala Gly Ala
35 40 45
Gly Gln Val Gly Arg Ser Leu Pro Gln Glu Ser Glu Glu Gln Arg
50 55 60
Thr Gly Ser Arg Pro Arg Arg Arg Arg Asp Leu Gly Ser Arg Leu
65 70 75
Gln Ala Gln Arg Arg Ala Gln Arg Val Ala Trp Glu Asp Gly Asp

-continued

80										85										90									
Glu	Asn	Val	Gly	Gln	Thr	Val	Ile	Pro	Ala	Gln	Glu	Glu	Glu	Gly															
				95						100																			
Ile	Glu	Lys	Pro	Ala	Glu	Val	His	Pro	Thr	Gly	Lys	Ile	Gly	Ala															
				110						115																			
Lys	Lys	Leu	Arg	Lys	Leu	Glu	Glu	Lys	Gln	Ala	Arg	Lys	Ala	Gln															
				125						130																			
Arg	Glu	Ala	Glu	Glu	Ala	Glu	Arg	Glu	Glu	Arg	Lys	Arg	Leu	Glu															
				140						145																			
Ser	Gln	Arg	Glu	Ala	Glu	Trp	Lys	Lys	Glu	Glu	Glu	Arg	Leu	Arg															
				155						160																			
Leu	Lys	Glu	Glu	Gln	Lys	Glu	Glu	Glu	Glu	Arg	Lys	Ala	Gln	Glu															
				170						175																			
Glu	Gln	Ala	Arg	Arg	Asp	His	Glu	Glu	Tyr	Leu	Lys	Leu	Lys	Glu															
				185						190																			
Ala	Phe	Val	Val	Glu	Glu	Glu	Gly	Val	Ser	Glu	Thr	Met	Thr	Glu															
				200						205																			
Glu	Gln	Ser	His	Ser	Phe	Leu	Thr	Glu	Phe	Ile	Asn	Tyr	Ile	Lys															
				215						220																			
Lys	Ser	Lys	Val	Val	Leu	Leu	Glu	Asp	Leu	Ala	Phe	Gln	Met	Gly															
				230						235																			
Leu	Arg	Thr	Gln	Asp	Ala	Ile	Asn	Arg	Ile	Gln	Asp	Leu	Leu	Thr															
				245						250																			
Glu	Gly	Thr	Leu	Thr	Gly	Val	Ile	Asp	Asp	Arg	Gly	Lys	Phe	Ile															
				260						265																			
Tyr	Ile	Thr	Pro	Glu	Glu	Leu	Ala	Ala	Val	Ala	Asn	Phe	Ile	Arg															
				275						280																			
Gln	Arg	Gly	Arg	Val	Ser	Ile	Thr	Glu	Leu	Ala	Gln	Ala	Ser	Asn															
				290						295																			
Ser	Leu	Ile	Ser	Trp	Gly	Gln	Asp	Leu	Pro	Ala	Gln	Ala	Ser	Ala															
				305						310																			

<210> SEQ ID NO 45
<211> LENGTH: 1977
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 45

acggggccgca gcggcagtga cgtagggttg gcgcacggat ccgttgccggc	50
tgcagctcttg cagtcggggcc gttccttcgc cgccgccagg ggtagcggtg	100
tagctgcgca gcgtcgcgcg cgctaccgca cccaggttcg gcccgtaggc	150
gtctggcagc ccggcgccat cttcatcgag cgccatggcc gcagcctgcg	200
ggccgggagc ggccgggtac tgcttgctcc tcggttgca tttgtttctg	250
ctgaccgcgg gccctgccct gggctggaac gacctgaca gaatgttgct	300
gcgggatgta aaagctctta ccctccacta tgaccgctat accacctccc	350
gcaggtgga tcccatccca cagttgaaat gtgttgagg cacagctggt	400
tgtgattctt ataccctaaa agtcatacag tgtcagaaca aaggctggga	450
tgggtatgat gtacagtggg aatgtaagac ggacttagat attgcataca	500
aatttgga aaactgtggtg agctgtgaag gctatgagtc ctctgaagac	550

-continued

cagtatgtac taagaggttc ttgtggcttg gagtataatt tagattatac 600
agaacttggc ctgcagaaac tgaaggagtc tggaaagcag cacggctttg 650
cctctttctc tgattattat tataagtggc cctcggcgga ttctctgtaac 700
atgagtggat tgattacccat cgtggctactc cttgggatcg cctttgtagt 750
ctataagctg ttctctgagt acgggcagta ttctcctcca cgtactctg 800
agtatcctcc attttccac cgttaccaga gattcaccaa ctcagcagga 850
cctcctcccc caggctttaa gtctgagttc acaggaccac agaatactgg 900
ccatggtgca acttctgggt ttggcagtc ttttacagga caacaaggat 950
atgaaaattc aggaccaggg ttctggacag gcttgggaac tggtggaata 1000
ctagatatt tgtttggcag caatagagcg gcaacaccct tctcagactc 1050
gtggtactac ccgtcctatc ctccctccta cctgggcacg tggaataggg 1100
cttactcacc ccttcatgga ggctcgggca gctattcggg atgttcaaac 1150
tcagacacga aaaccagaac tgcacagga tatggtggtg ccaggagacg 1200
ataaagtaga aagttggagt caaacactgg atgcagaaat tttggatttt 1250
tcatcacttt ctcttttaga aaaaagtact acctgttaac aattgggaaa 1300
aggggatatt caaaagttct gtggtgttat gtccagtgtg gctttttgta 1350
ttctattatt tgaggctaaa agttgatgtg tgacaaaata cttatgtgtt 1400
gtatgtcagt gtaacatgca gatgtatatt gcagtttttg aaagtgatca 1450
ttactgtgga atgctaaaaa tacattaatt tctaaaacct gtgatgccct 1500
aagaagcatt aagaatgaag gtgtgtgtact aatagaaact aagtacagaa 1550
aatttcagtt ttaggtgggt gtagctgatg agttattacc tcatagagac 1600
tataatattc tatttgggat tatattattt gatgtttgct gttcttcaaa 1650
catttaaatac aagctttgga ctaattatgc taatttgtga gttctgatca 1700
cttttgagct ctgaagcttt gaatcattca gtggtggaga tggccttctg 1750
gtaactgaat attaccttct gtaggaaaag gtggaaaata agcatctaga 1800
aggttgttgt gaatgactct gtgctggcaa aaatgcttga aacctctata 1850
tttctttcgt tcataagagg taaaggtcaa atttttcaac aaaagtcttt 1900
taataacaaa agcatgcagt tctctgtgaa atctcaaata ttgttgtaat 1950
agtctgtttc aatcttaaaa agaataca 1977

<210> SEQ ID NO 46
<211> LENGTH: 339
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 46

Met Ala Ala Ala Cys Gly Pro Gly Ala Ala Gly Tyr Cys Leu Leu
1 5 10 15
Leu Gly Leu His Leu Phe Leu Leu Thr Ala Gly Pro Ala Leu Gly
20 25 30
Trp Asn Asp Pro Asp Arg Met Leu Leu Arg Asp Val Lys Ala Leu
35 40 45

-continued

Thr	Leu	His	Tyr	Asp	Arg	Tyr	Thr	Thr	Ser	Arg	Arg	Leu	Asp	Pro	
				50					55					60	
Ile	Pro	Gln	Leu	Lys	Cys	Val	Gly	Gly	Thr	Ala	Gly	Cys	Asp	Ser	
				65					70					75	
Tyr	Thr	Pro	Lys	Val	Ile	Gln	Cys	Gln	Asn	Lys	Gly	Trp	Asp	Gly	
				80					85					90	
Tyr	Asp	Val	Gln	Trp	Glu	Cys	Lys	Thr	Asp	Leu	Asp	Ile	Ala	Tyr	
				95					100					105	
Lys	Phe	Gly	Lys	Thr	Val	Val	Ser	Cys	Glu	Gly	Tyr	Glu	Ser	Ser	
				110					115					120	
Glu	Asp	Gln	Tyr	Val	Leu	Arg	Gly	Ser	Cys	Gly	Leu	Glu	Tyr	Asn	
				125					130					135	
Leu	Asp	Tyr	Thr	Glu	Leu	Gly	Leu	Gln	Lys	Leu	Lys	Glu	Ser	Gly	
				140					145					150	
Lys	Gln	His	Gly	Phe	Ala	Ser	Phe	Ser	Asp	Tyr	Tyr	Tyr	Lys	Trp	
				155					160					165	
Ser	Ser	Ala	Asp	Ser	Cys	Asn	Met	Ser	Gly	Leu	Ile	Thr	Ile	Val	
				170					175					180	
Val	Leu	Leu	Gly	Ile	Ala	Phe	Val	Val	Tyr	Lys	Leu	Phe	Leu	Ser	
				185					190					195	
Asp	Gly	Gln	Tyr	Ser	Pro	Pro	Pro	Tyr	Ser	Glu	Tyr	Pro	Pro	Phe	
				200					205					210	
Ser	His	Arg	Tyr	Gln	Arg	Phe	Thr	Asn	Ser	Ala	Gly	Pro	Pro	Pro	
				215					220					225	
Pro	Gly	Phe	Lys	Ser	Glu	Phe	Thr	Gly	Pro	Gln	Asn	Thr	Gly	His	
				230					235					240	
Gly	Ala	Thr	Ser	Gly	Phe	Gly	Ser	Ala	Phe	Thr	Gly	Gln	Gln	Gly	
				245					250					255	
Tyr	Glu	Asn	Ser	Gly	Pro	Gly	Phe	Trp	Thr	Gly	Leu	Gly	Thr	Gly	
				260					265					270	
Gly	Ile	Leu	Gly	Tyr	Leu	Phe	Gly	Ser	Asn	Arg	Ala	Ala	Thr	Pro	
				275					280					285	
Phe	Ser	Asp	Ser	Trp	Tyr	Tyr	Pro	Ser	Tyr	Pro	Pro	Ser	Tyr	Pro	
				290					295					300	
Gly	Thr	Trp	Asn	Arg	Ala	Tyr	Ser	Pro	Leu	His	Gly	Gly	Ser	Gly	
				305					310					315	
Ser	Tyr	Ser	Val	Cys	Ser	Asn	Ser	Asp	Thr	Lys	Thr	Arg	Thr	Ala	
				320					325					330	
Ser	Gly	Tyr	Gly	Gly	Thr	Arg	Arg	Arg							
				335											

<210> SEQ ID NO 47
<211> LENGTH: 1766
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 47

cccggagccg	gggaggagg	gagcgaggtt	cggacaccgg	cggcggtgc	50
ctggcctttc	catgagccc	cggggagccc	tcccgcgccc	cctctcgtc	100
tgccctctcc	tctgcctctg	cctctgcctg	gccgcggctc	tggaagtgc	150
gcagtcggg	tcgtgtagg	ataaaaagaa	ctgtaagtg	gtcttttccc	200

-continued	
agcaggaact gaggaagcgg ctaacacccc tgcagtacca tgtcactcag	250
gagaaaggga ccgaaagtgc ctttgaagga gaatacacac atcacaaaga	300
tccttggaata tataaatgtg ttgtttgtgg aactccattg tttaaagtcag	350
aaaccaaatt tgactccggt tcaggttggc cttcattcca cgatgtgatc	400
aattctgagg caatcacatt cacagatgac ttttcctatg ggatgcacag	450
ggtggaagaa agctgctctc agtgtggtgc tcaccttggg cacatttttg	500
atgatgggccc tcgtccaact gggaaaagat actgcataaa ttcggctgcc	550
ttgtctttta cacctgcgga tagcagtggc accgccgagg gaggcagtgg	600
ggtcgccagc ccgcccagg cagacaaagc ggagctctag agtaatggag	650
agtgatggaa acaaagtgtg cttaatgcac agcttattaa aaaaatcaaa	700
attgttatct taatagatat attttttcaa aaactataag ggcagttttg	750
tgctattgat attttttctt cttttgctta aacagaagcc ctggccatcc	800
atgtattttg caattgacta gatcaagaac tgtttatagc tttagcaaat	850
ggagacagct ttgtgaaact tcttcacaag ccacttatac cctttggcat	900
tcttttcttt gagcacatgg cttcttttgc agtttttccc cctttgattc	950
agaagcagag ggttcatggt cttcaaacat gaaaatagag atctcctctg	1000
cagtgtagag accagagctg ggcagtgcag ggcattggaga cctgcaagac	1050
acatggcctt gaggcctttg cacagaccca cctaagataa ggttggagtg	1100
atgttttaat gagactgttc agctttgtgg aaagtttgag ctaaggtcac	1150
tttttttttt ctactgaaa ggggtgtgaag gtctaaagtc tttccttatg	1200
ttaaattgtt gccagatcca aaggggcata ctgagtgttg tggcagagaa	1250
gtaaacatta ccacactgtt aggcctttat tttattttat tttccatcga	1300
aagcattgga ggcccagtgc aatggctcac gcctgtgatc ccagcacttt	1350
gggaggccaa ggcgggtgga tcacgaggtc aggagatgga gaccatcctg	1400
gctaacatgg tgaacccccg tctctactaa aaatacgaag aattagccag	1450
gcgtgggtgtt gggcacctgt agtcccagct actcaggagg ctgaggcagg	1500
agaatggcgt gaacccggaa ggcggagctt gcagttagcc gagatcatgc	1550
cactgcactc cagcctacat gacaatgtga cactccatct caaaaaataa	1600
taataataac aatataagaa ctagctgggc atggtggcgc atgcatgtag	1650
tcccagctac tcctgaggct cagtcaggag aatcgcttga acttgggagg	1700
cggagggttg agtgagctga gctcatacca ctgcactcca gcctgaacag	1750
agtgagatcc tgtcaa	1766
<210> SEQ ID NO 48	
<211> LENGTH: 192	
<212> TYPE: PRT	
<213> ORGANISM: Homo Sapien	
<400> SEQUENCE: 48	
Met Ser Pro Arg Arg Thr Leu Pro Arg Pro Leu Ser Leu Cys Leu	
1 5 10 15	
Ser Leu Cys Leu Cys Leu Cys Leu Ala Ala Ala Leu Gly Ser Ala	

-continued

20										25										30									
Gln	Ser	Gly	Ser	Cys	Arg	Asp	Lys	Lys	Asn	Cys	Lys	Val	Val	Phe															
				35					40					45															
Ser	Gln	Gln	Glu	Leu	Arg	Lys	Arg	Leu	Thr	Pro	Leu	Gln	Tyr	His															
				50					55					60															
Val	Thr	Gln	Glu	Lys	Gly	Thr	Glu	Ser	Ala	Phe	Glu	Gly	Glu	Tyr															
				65					70					75															
Thr	His	His	Lys	Asp	Pro	Gly	Ile	Tyr	Lys	Cys	Val	Val	Cys	Gly															
				80					85					90															
Thr	Pro	Leu	Phe	Lys	Ser	Glu	Thr	Lys	Phe	Asp	Ser	Gly	Ser	Gly															
				95					100					105															
Trp	Pro	Ser	Phe	His	Asp	Val	Ile	Asn	Ser	Glu	Ala	Ile	Thr	Phe															
				110					115					120															
Thr	Asp	Asp	Phe	Ser	Tyr	Gly	Met	His	Arg	Val	Glu	Thr	Ser	Cys															
				125					130					135															
Ser	Gln	Cys	Gly	Ala	His	Leu	Gly	His	Ile	Phe	Asp	Asp	Gly	Pro															
				140					145					150															
Arg	Pro	Thr	Gly	Lys	Arg	Tyr	Cys	Ile	Asn	Ser	Ala	Ala	Leu	Ser															
				155					160					165															
Phe	Thr	Pro	Ala	Asp	Ser	Ser	Gly	Thr	Ala	Glu	Gly	Gly	Ser	Gly															
				170					175					180															
Val	Ala	Ser	Pro	Ala	Gln	Ala	Asp	Lys	Ala	Glu	Leu																		
				185					190																				

<210> SEQ ID NO 49
<211> LENGTH: 2065
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 49

cccaaagagg tgaggagccg gcagcggggg cggctgtaac tgtgaggaag	50
gctgcagagt ggcgacgtct acgccgtagg ttggaggctg tgggggggtg	100
ccgggcgcga gctcccaggc cgcagaagtg acctgcggtg gagttccctc	150
ctcgtctgtg gagaacggag ggagaaggtt gctggccggg tgaaagtgcc	200
tccctctgct tgacggggct gaggggcccg aagtctaggg cgtccgtagt	250
cgccccggcc tccgtgaagc ccaggtcta gagatatgac ccgagagtgc	300
ccatctccgg ccccggggcc tggggctccg ctgagtggat cggtgctggc	350
agaggcgcga gtagtgtttg cagtgggtgt gagcatccac gcaaccgtat	400
gggaccgata ctctgtgtgc gccgtggccc tcgcagtga ggccttctac	450
gtccaataca agtgggaccg gctgctacag cagggaagcg cgtcttcca	500
gttcgaatg tccgcaaaca gtggcctatt gccgcctcc atggtcatgc	550
ctttgcttgg actagtcatg aaggagcggg gccagactgc tgggaacccg	600
ttctttgagc gttttggcat tgtgtgggca gccactggca tggcagtggc	650
cctcttctca tcagtgttgg cgctcggcat cactcgccca gtgccaacca	700
acacttgtgt catcttgggc ttggctggag gtgttatcat ttatatcatg	750
aagcactcgt tgagcgtggg ggaggtgatc gaagtcctgg aagtccttct	800
gatcttcgtt tatctcaaca tgatcctgct gtacctgtg ccccgctgct	850

-continued

tcacccctgg tgaggcactg ctggtattgg gtggcattag ctttgcctc 900
aaccagctca tcaagcgctc tctgacactg gtggaaagtc agggggaccc 950
agtggacttc ttctgtctgg tgggtggtagt agggatggta ctcatgggca 1000
ttttcttcag cactctgttt gtcttcatgg actcaggcac ctgggcctcc 1050
tccatcttct tccacctcat gacctgtgtg ctgagccttg gtgtggtcct 1100
accctggctg caccggctca tccgcaggaa tcccctgctc tggcttcttc 1150
agtttctctt ccagacagac acccgcatct acctcctagc ctattggtct 1200
ctgtgggcca ccttggcctg cctggtggtg ctgtaccaga atgccaagcg 1250
gtcatcttcc gagtccaaga agcaccaggc cccaccatc gcccgaaagt 1300
atttccacct cattgtggta gccacctaca tcccaggat catctttgac 1350
cggcactcgc tctatgtagc cgccactgta tgcctggcgg tcttcatctt 1400
cctggagtat gtgcgctact tccgcatcaa gcctttgggt cacactctac 1450
ggagcttctt gtcccttttt ctggatgaac gagacagtgg accactcatt 1500
ctgacacaca tctacctgct cctgggcatg tctcttccca tctggtgat 1550
ccccagacct tgcacacaga agggtagcct gggaggagcc agggccctcg 1600
tcccctatgc cgggtgcctg gctgtgggtg tgggtgatac tgtggcctcc 1650
atcttcggta gcaccatggg ggagatccgc tggcctggaa ccaaaaagac 1700
ttttgagggg accatgacat ctatatattg gcagatcatt tctgtagctc 1750
tgatcttaat ctttgacagt ggagtggacc taaactacag ttatgcttgg 1800
attttggggt ccatcagcac tgtgtccctc ctggaagcat aactacaca 1850
gatagacaa ctcttctgct ctctctacct cctgatattg ctgatggcct 1900
agctgttaca gtgcagcagc agtgacggag gaaacagaca tggggagggg 1950
gaacagtccc cacagcagac agctacttgg gcataagag ccaaggtgtg 2000
aaaagcagat ttgatttttc agttgattca gatttaaaat aaaaagcaaa 2050
gctctcctag ttcta 2065

<210> SEQ ID NO 50
<211> LENGTH: 538
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 50

Met Thr Arg Glu Cys Pro Ser Pro Ala Pro Gly Pro Gly Ala Pro
1 5 10 15
Leu Ser Gly Ser Val Leu Ala Glu Ala Ala Val Val Phe Ala Val
20 25 30
Val Leu Ser Ile His Ala Thr Val Trp Asp Arg Tyr Ser Trp Cys
35 40 45
Ala Val Ala Leu Ala Val Gln Ala Phe Tyr Val Gln Tyr Lys Trp
50 55 60
Asp Arg Leu Leu Gln Gln Gly Ser Ala Val Phe Gln Phe Arg Met
65 70 75
Ser Ala Asn Ser Gly Leu Leu Pro Ala Ser Met Val Met Pro Leu
80 85 90

-continued

Leu Gly Leu Val Met Lys Glu Arg Cys Gln Thr Ala Gly Asn Pro	95	100	105
Phe Phe Glu Arg Phe Gly Ile Val Val Ala Ala Thr Gly Met Ala	110	115	120
Val Ala Leu Phe Ser Ser Val Leu Ala Leu Gly Ile Thr Arg Pro	125	130	135
Val Pro Thr Asn Thr Cys Val Ile Leu Gly Leu Ala Gly Gly Val	140	145	150
Ile Ile Tyr Ile Met Lys His Ser Leu Ser Val Gly Glu Val Ile	155	160	165
Glu Val Leu Glu Val Leu Leu Ile Phe Val Tyr Leu Asn Met Ile	170	175	180
Leu Leu Tyr Leu Leu Pro Arg Cys Phe Thr Pro Gly Glu Ala Leu	185	190	195
Leu Val Leu Gly Gly Ile Ser Phe Val Leu Asn Gln Leu Ile Lys	200	205	210
Arg Ser Leu Thr Leu Val Glu Ser Gln Gly Asp Pro Val Asp Phe	215	220	225
Phe Leu Leu Val Val Val Gly Met Val Leu Met Gly Ile Phe	230	235	240
Phe Ser Thr Leu Phe Val Phe Met Asp Ser Gly Thr Trp Ala Ser	245	250	255
Ser Ile Phe Phe His Leu Met Thr Cys Val Leu Ser Leu Gly Val	260	265	270
Val Leu Pro Trp Leu His Arg Leu Ile Arg Arg Asn Pro Leu Leu	275	280	285
Trp Leu Leu Gln Phe Leu Phe Gln Thr Asp Thr Arg Ile Tyr Leu	290	295	300
Leu Ala Tyr Trp Ser Leu Leu Ala Thr Leu Ala Cys Leu Val Val	305	310	315
Leu Tyr Gln Asn Ala Lys Arg Ser Ser Ser Glu Ser Lys Lys His	320	325	330
Gln Ala Pro Thr Ile Ala Arg Lys Tyr Phe His Leu Ile Val Val	335	340	345
Ala Thr Tyr Ile Pro Gly Ile Ile Phe Asp Arg Pro Leu Leu Tyr	350	355	360
Val Ala Ala Thr Val Cys Leu Ala Val Phe Ile Phe Leu Glu Tyr	365	370	375
Val Arg Tyr Phe Arg Ile Lys Pro Leu Gly His Thr Leu Arg Ser	380	385	390
Phe Leu Ser Leu Phe Leu Asp Glu Arg Asp Ser Gly Pro Leu Ile	395	400	405
Leu Thr His Ile Tyr Leu Leu Leu Gly Met Ser Leu Pro Ile Trp	410	415	420
Leu Ile Pro Arg Pro Cys Thr Gln Lys Gly Ser Leu Gly Gly Ala	425	430	435
Arg Ala Leu Val Pro Tyr Ala Gly Val Leu Ala Val Gly Val Gly	440	445	450
Asp Thr Val Ala Ser Ile Phe Gly Ser Thr Met Gly Glu Ile Arg	455	460	465

-continued

Trp	Pro	Gly	Thr	Lys	Lys	Thr	Phe	Glu	Gly	Thr	Met	Thr	Ser	Ile
				470					475					480
Phe	Ala	Gln	Ile	Ile	Ser	Val	Ala	Leu	Ile	Leu	Ile	Phe	Asp	Ser
				485					490					495
Gly	Val	Asp	Leu	Asn	Tyr	Ser	Tyr	Ala	Trp	Ile	Leu	Gly	Ser	Ile
				500					505					510
Ser	Thr	Val	Ser	Leu	Leu	Glu	Ala	Tyr	Thr	Thr	Gln	Ile	Asp	Asn
				515					520					525
Leu	Leu	Leu	Pro	Leu	Tyr	Leu	Leu	Ile	Leu	Leu	Met	Ala		
				530					535					

<210> SEQ ID NO 51
<211> LENGTH: 3476
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 51

gctctatgcc gcctaccttg ctctcgccgc tgctgccgga gccgaagcag	50
agaaggcagc ggggcccttg accgtcccga gagccccgcg ctcccgacca	100
ggggggcggg gcggccccgg ggaggggcgg gcaggggcgg ggggaagaaa	150
gggggttttg tgctgcgcgc ggagggccgg cgccctcttc cgaatgtcct	200
gcggccccag cctctcctca cgctcgcgca gtctccgcgc cagtctcagc	250
tgacagtgoa ggactgagcc gtgcacccgg aggagacccc cggaggaggc	300
gacaaacttc gcagtgcgcg gacccaaccc cagccctggg tagcctgcag	350
catggccccg ctgttcctgc ccctgctggc agccctggtc ctggccccag	400
ctctcgcagc tttagcagat gttctggaag gagacagctc agaggaccgc	450
gcttttcgcg tgcgcatcgc gggcgacgcg ccaactgcagg gcgtgctcgg	500
cgggcgccctc accatccctt gccacgtcca ctacctgcgg ccaccgccga	550
gccgcggggc tgtgctgggc tctccgcggg tcaagtggac ttctctgtcc	600
cggggccggg aggcagaggt gctggtggcg cggggagtgc gcgtcaaggt	650
gaacgaggcc taccggttcc gcgtggcact gcctgcgtac ccagcgtcgc	700
tcaccgacgt ctccctggcg ctgagcgagc tgcgccccaa cgactcaggt	750
atctatcgct gtgaggtcca gcacggcatc gatgacagca gcgacgctgt	800
ggaggtcaag gtcaaagggg tcgtctttct ctaccgagag ggtctgccc	850
gctatgcttt ctctttttct ggggcccgag aggcctgtgc ccgcattgga	900
gcccacatcg ccaccccgga gcagctctat gccgcctacc ttggggggcta	950
tgagcaatgt gatgctggct ggctgtcgga tcagaccgtg aggtatccca	1000
tccagacccc acgagaggcc tgttacggag acatggatgg cttccccggg	1050
gtccggaact atggtgtggt ggaccggat gacctctatg atgtgtactg	1100
ttatgctgaa gacctaaatg gagaactgtt cctgggtgac cctccagaga	1150
agctgacatt ggaggaagca cgggcgtact gccaggagcg gggcgcagag	1200
attgccacca cgggccaaact gtatgcagcc tgggatggtg gcctggacca	1250
ctgcagccca ggggtggctag ctgatggcag tgtgcgctac cccatcgtca	1300
caccagccca gcgctgtggt gggggcttgc ctggtgtcaa gactctcttc	1350

-continued

ctcttcccca accagactgg cttccccaat aagcacagcc gcttcaacgt	1400
ctactgcttc cgagactcgg cccagccttc tgccatccct gaggcctcca	1450
accagcctc caaccagcc tctgatggac tagaggctat cgtcacagtg	1500
acagagacc tggaggaact gcagctgcct caggaagcca cagagagtga	1550
atcccgtggg gccatctact ccattcccat catggaggac ggaggaggtg	1600
gaagctccac tccagaagac ccagcagagg cccctaggac gctcctagaa	1650
tttgaacac aatccatggt accgcccacg gggttctcag aagaggaagg	1700
taaggcattg gaggaagaag agaaatatga agatgaagaa gaaaaagag	1750
aggaagaaga agaggaggag gtggaggatg aggctctgtg ggcatggccc	1800
agcgagctca gcagcccggg ccttgaggcc tctctcccca ctgagccagc	1850
agcccaggag aagtcaactt cccaggcgcc agcaagggca gtcctgcagc	1900
ctggtgcac accacttcct gatggagagt cagaagcttc caggcctcca	1950
aggggccatg gaccacctac tgagactctg cccactccca gggagaggaa	2000
cctagcatcc ccatcacctt ccactctggt tgaggcaaga gaggtggggg	2050
aggcaactgg tggctcctgag ctatctgggg tccctcgagg agagagcgag	2100
gagacagaa gctccgaggg tgccccttcc ctgcttcag ccacacgggc	2150
ccctgagggg accagggagc tggaggcccc ctctgaagat aattctggaa	2200
gaactgcccc agcagggacc tcagtgcagg cccagccagt gctgcccact	2250
gacacggcca gccgaggtgg agtggccgtg gtccccgcat cagggtgactg	2300
tgccccagc ccctgccaca atggtgggac atgcttgagg gaggaggaag	2350
gggtccgctg cctatgtctg cctggctatg ggggggacct gtgcgatgtt	2400
ggcctccgct tctgcaaccc cggctgggac gccttcagg gcgcctgcta	2450
caagcacttt tccacacgaa gagctggga ggaggcagag acccagtgcc	2500
ggatgtacgg cgcgcactct gccagcatca gcacaccga ggaacaggac	2550
ttcatcaaca accggtaccg ggagtaccag tggatcggac tcaacgacag	2600
gaccatcgaa ggcgacttct tgtggtcgga tggcgcccc ctgctctatg	2650
agaactggaa ccctgggcag cctgacagct acttcctgtc tggagagaa	2700
tgcggtgcca tgggtgtggca tgatcaggga caatggagtg acgtgccctg	2750
caactaccac ctgtcctaca cctgcaagat ggggctggtg tcctgtgggc	2800
cgccaccgga gctgcccctg gctcaagtgt tcggccgccc acggctgcgc	2850
tatgaggtgg aactgtgct tcgtaccgg tgccgggaag gactggccca	2900
gcgcaatctg ccgctgatcc gatgccaa gaacgggtcgt tgggagggcc	2950
cccagatctc ctgtgtgccc agaagacctg cccgagctct gccaccagag	3000
gaggaccag aaggacgtca ggggaggcta ctgggacgct ggaaggcgct	3050
gttgatcccc cctccagcc catgccagg tccctagggg gcaaggcctt	3100
gaacactgcc ggccacagca ctgcctgtc acccaaattt tccctcacac	3150
cttgcgctcc cgccaccaca ggaagtgaca acatgacgag ggtgtgtgct	3200
ggagtccagg tgacagttcc tgaaggggct tctgggaaat acctaggagg	3250

-continued

ctccagccca gccagggccc tctcccccta ccctgggcac cagatcttcc 3300
atcagggccg gagtaaatcc ctaagtcct caactgccct ctccctggca 3350
gccatcttgt cccctctatt cctctaggga gcaactgtgcc cactctttct 3400
gggtttttcca agggaatggg cttgcaggat ggagtgtctg taaatcaac 3450
aggaaataaa actgtgtatg agccca 3476

<210> SEQ ID NO 52
<211> LENGTH: 911
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 52

Met Ala Gln Leu Phe Leu Pro Leu Leu Ala Ala Leu Val Leu Ala
1 5 10 15
Gln Ala Pro Ala Ala Leu Ala Asp Val Leu Glu Gly Asp Ser Ser
20 25 30
Glu Asp Arg Ala Phe Arg Val Arg Ile Ala Gly Asp Ala Pro Leu
35 40 45
Gln Gly Val Leu Gly Gly Ala Leu Thr Ile Pro Cys His Val His
50 55 60
Tyr Leu Arg Pro Pro Pro Ser Arg Arg Ala Val Leu Gly Ser Pro
65 70 75
Arg Val Lys Trp Thr Phe Leu Ser Arg Gly Arg Glu Ala Glu Val
80 85 90
Leu Val Ala Arg Gly Val Arg Val Lys Val Asn Glu Ala Tyr Arg
95 100 105
Phe Arg Val Ala Leu Pro Ala Tyr Pro Ala Ser Leu Thr Asp Val
110 115 120
Ser Leu Ala Leu Ser Glu Leu Arg Pro Asn Asp Ser Gly Ile Tyr
125 130 135
Arg Cys Glu Val Gln His Gly Ile Asp Asp Ser Ser Asp Ala Val
140 145 150
Glu Val Lys Val Lys Gly Val Val Phe Leu Tyr Arg Glu Gly Ser
155 160 165
Ala Arg Tyr Ala Phe Ser Phe Ser Gly Ala Gln Glu Ala Cys Ala
170 175 180
Arg Ile Gly Ala His Ile Ala Thr Pro Glu Gln Leu Tyr Ala Ala
185 190 195
Tyr Leu Gly Gly Tyr Glu Gln Cys Asp Ala Gly Trp Leu Ser Asp
200 205 210
Gln Thr Val Arg Tyr Pro Ile Gln Thr Pro Arg Glu Ala Cys Tyr
215 220 225
Gly Asp Met Asp Gly Phe Pro Gly Val Arg Asn Tyr Gly Val Val
230 235 240
Asp Pro Asp Asp Leu Tyr Asp Val Tyr Cys Tyr Ala Glu Asp Leu
245 250 255
Asn Gly Glu Leu Phe Leu Gly Asp Pro Pro Glu Lys Leu Thr Leu
260 265 270
Glu Glu Ala Arg Ala Tyr Cys Gln Glu Arg Gly Ala Glu Ile Ala
275 280 285

-continued

Thr	Thr	Gly	Gln	Leu	Tyr	Ala	Ala	Trp	Asp	Gly	Gly	Leu	Asp	His
				290					295					300
Cys	Ser	Pro	Gly	Trp	Leu	Ala	Asp	Gly	Ser	Val	Arg	Tyr	Pro	Ile
				305					310					315
Val	Thr	Pro	Ser	Gln	Arg	Cys	Gly	Gly	Gly	Leu	Pro	Gly	Val	Lys
				320					325					330
Thr	Leu	Phe	Leu	Phe	Pro	Asn	Gln	Thr	Gly	Phe	Pro	Asn	Lys	His
				335					340					345
Ser	Arg	Phe	Asn	Val	Tyr	Cys	Phe	Arg	Asp	Ser	Ala	Gln	Pro	Ser
				350					355					360
Ala	Ile	Pro	Glu	Ala	Ser	Asn	Pro	Ala	Ser	Asn	Pro	Ala	Ser	Asp
				365					370					375
Gly	Leu	Glu	Ala	Ile	Val	Thr	Val	Thr	Glu	Thr	Leu	Glu	Glu	Leu
				380					385					390
Gln	Leu	Pro	Gln	Glu	Ala	Thr	Glu	Ser	Glu	Ser	Arg	Gly	Ala	Ile
				395					400					405
Tyr	Ser	Ile	Pro	Ile	Met	Glu	Asp	Gly	Gly	Gly	Gly	Ser	Ser	Thr
				410					415					420
Pro	Glu	Asp	Pro	Ala	Glu	Ala	Pro	Arg	Thr	Leu	Leu	Glu	Phe	Glu
				425					430					435
Thr	Gln	Ser	Met	Val	Pro	Pro	Thr	Gly	Phe	Ser	Glu	Glu	Glu	Gly
				440					445					450
Lys	Ala	Leu	Glu	Glu	Glu	Glu	Lys	Tyr	Glu	Asp	Glu	Glu	Glu	Lys
				455					460					465
Glu	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Val	Glu	Asp	Glu	Ala	Leu	Trp
				470					475					480
Ala	Trp	Pro	Ser	Glu	Leu	Ser	Ser	Pro	Gly	Pro	Glu	Ala	Ser	Leu
				485					490					495
Pro	Thr	Glu	Pro	Ala	Ala	Gln	Glu	Lys	Ser	Leu	Ser	Gln	Ala	Pro
				500					505					510
Ala	Arg	Ala	Val	Leu	Gln	Pro	Gly	Ala	Ser	Pro	Leu	Pro	Asp	Gly
				515					520					525
Glu	Ser	Glu	Ala	Ser	Arg	Pro	Pro	Arg	Val	His	Gly	Pro	Pro	Thr
				530					535					540
Glu	Thr	Leu	Pro	Thr	Pro	Arg	Glu	Arg	Asn	Leu	Ala	Ser	Pro	Ser
				545					550					555
Pro	Ser	Thr	Leu	Val	Glu	Ala	Arg	Glu	Val	Gly	Glu	Ala	Thr	Gly
				560					565					570
Gly	Pro	Glu	Leu	Ser	Gly	Val	Pro	Arg	Gly	Glu	Ser	Glu	Glu	Thr
				575					580					585
Gly	Ser	Ser	Glu	Gly	Ala	Pro	Ser	Leu	Leu	Pro	Ala	Thr	Arg	Ala
				590					595					600
Pro	Glu	Gly	Thr	Arg	Glu	Leu	Glu	Ala	Pro	Ser	Glu	Asp	Asn	Ser
				605					610					615
Gly	Arg	Thr	Ala	Pro	Ala	Gly	Thr	Ser	Val	Gln	Ala	Gln	Pro	Val
				620					625					630
Leu	Pro	Thr	Asp	Ser	Ala	Ser	Arg	Gly	Gly	Val	Ala	Val	Val	Pro
				635					640					645
Ala	Ser	Gly	Asp	Cys	Val	Pro	Ser	Pro	Cys	His	Asn	Gly	Gly	Thr
				650					655					660
Cys	Leu	Glu	Glu	Glu	Glu	Gly	Val	Arg	Cys	Leu	Cys	Leu	Pro	Gly

-continued

665									670					675				
Tyr	Gly	Gly	Asp	Leu	Cys	Asp	Val	Gly	Leu	Arg	Phe	Cys	Asn	Pro				
680									685					690				
Gly	Trp	Asp	Ala	Phe	Gln	Gly	Ala	Cys	Tyr	Lys	His	Phe	Ser	Thr				
695									700					705				
Arg	Arg	Ser	Trp	Glu	Glu	Ala	Glu	Thr	Gln	Cys	Arg	Met	Tyr	Gly				
710									715					720				
Ala	His	Leu	Ala	Ser	Ile	Ser	Thr	Pro	Glu	Glu	Gln	Asp	Phe	Ile				
725									730					735				
Asn	Asn	Arg	Tyr	Arg	Glu	Tyr	Gln	Trp	Ile	Gly	Leu	Asn	Asp	Arg				
740									745					750				
Thr	Ile	Glu	Gly	Asp	Phe	Leu	Trp	Ser	Asp	Gly	Val	Pro	Leu	Leu				
755									760					765				
Tyr	Glu	Asn	Trp	Asn	Pro	Gly	Gln	Pro	Asp	Ser	Tyr	Phe	Leu	Ser				
770									775					780				
Gly	Glu	Asn	Cys	Val	Val	Met	Val	Trp	His	Asp	Gln	Gly	Gln	Trp				
785									790					795				
Ser	Asp	Val	Pro	Cys	Asn	Tyr	His	Leu	Ser	Tyr	Thr	Cys	Lys	Met				
800									805					810				
Gly	Leu	Val	Ser	Cys	Gly	Pro	Pro	Pro	Glu	Leu	Pro	Leu	Ala	Gln				
815									820					825				
Val	Phe	Gly	Arg	Pro	Arg	Leu	Arg	Tyr	Glu	Val	Asp	Thr	Val	Leu				
830									835					840				
Arg	Tyr	Arg	Cys	Arg	Glu	Gly	Leu	Ala	Gln	Arg	Asn	Leu	Pro	Leu				
845									850					855				
Ile	Arg	Cys	Gln	Glu	Asn	Gly	Arg	Trp	Glu	Ala	Pro	Gln	Ile	Ser				
860									865					870				
Cys	Val	Pro	Arg	Arg	Pro	Ala	Arg	Ala	Leu	His	Pro	Glu	Glu	Asp				
875									880					885				
Pro	Glu	Gly	Arg	Gln	Gly	Arg	Leu	Leu	Gly	Arg	Trp	Lys	Ala	Leu				
890									895					900				
Leu	Ile	Pro	Pro	Ser	Ser	Pro	Met	Pro	Gly	Pro								
905									910									

<210> SEQ ID NO 53
<211> LENGTH: 3316
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 53

ctgccaggtg acagccgccca agatgggggtc ttggggccctg ctgtggcctc	50
ccctgctgtt caccgggctg ctcgccgcac ccccggggac catggcccag	100
gcccagtact gctctgtgaa caaggacatc tttgaagtag aggagaacac	150
aaatgtcacc gagccgctgg tggacatcca cgtcccggag ggccaggagg	200
tgacctctcg agccttgtcc accccctttg catttcggat ccagggaaac	250
cagctgtttc tcaacgtgac tcctgattac gaggagaagt cactgcttga	300
ggctcagctg ctgtgtcaga gcggaggcac attggtgacc cagctaaggg	350
tgttcgtgtc agtgctggac gtcaatgaca atgccccga attccccttt	400
aagaccaagg agataagggt ggaggaggac acgaaagtga actccaccgt	450

-continued

catccctgag acgcaactgc aggetgagga ccgcgacaag gacgacattc	500
tgttctacac cctccaggaa atgacagcag gtgccagtga ctactttctc	550
ctggtgagtg taaacctgcc cgccctgagg ctggaccggc ccctggactt	600
ctacgagcgg ccgaacatga cttctggct gctggtgcgg gacactccag	650
gggagaatgt ggaacccagc cacactgcca ccgccacact agtgctgaac	700
gtggtgcccg ccgacctgcg gccccctgg ttctgcctt gcaccttctc	750
agatggctac gtctgcattc aagctcagta ccacggggct gtccccacgg	800
ggcacatact gccatctccc ctgcctctgc gtcccgacc catctacgt	850
gaggacggag accgcggcat caaccagccc atcatctaca gcattcttag	900
gggaaactgt aatggtacat tcatcatcca ccagactcg ggcaacctca	950
ccgtggccag gagtgtcccc agccccatga cttctctct gctggtgaag	1000
ggccaacagg ccgacctgc ccgctactca gtgaccagg tcacctgga	1050
ggctgtggct gcggccggga gcccgcctc cttccccag agcctgtatc	1100
gtggaccgt ggcgcgtggc gctggagcgg gcgttggtt caaggatgca	1150
gtgccccct ctacgctct gaggatccag gctcaggacc cggagtctc	1200
ggacctcaac tcggccatca catatcgaat taccaaccac tcacacttcc	1250
ggatggagg agaggttgt ctgaccacca ccacactggc acaggcggga	1300
gccttctacg cagaggttga ggcccacaac acggtgacct ctggcaccgc	1350
aaccacagtc attgagatac aagtttccga acaggagccc ccctccacag	1400
aggctggagg aacaactggg ccctggacca gcaccacttc cgagggtccc	1450
agacccccct agccctccca gggaccctcc acgaccagct ctgggggagg	1500
cacaggccct catccacct ctggcacaac totgaggcca ccaacctcgt	1550
ccacaccggg ggggcccccg ggtgcagaaa acagcacctc ccaccaacca	1600
gccactcccg gtggggacac agcacagacc ccaaagccag gaacctctca	1650
gccgatgcc cccggtgtgg gaaccagcac ctcccaccaa ccagccacac	1700
ccagtggggg cacagcacag accccagagc caggaaacct tcagccgatg	1750
ccccccagta tgggaaccag cacctcccac caaccagcca cccccgttg	1800
gggcacagca cagaccccag aggcaggaac ctctcagccg atgcccccg	1850
gtatgggaac cagcacctcc caccaaccaa ccacaccgg tgggggcaca	1900
gcacagacc cagagccagg aacctctcag ccgatgcccc tcagcaagag	1950
cacccatct tcaggtggcg gcccctcggg ggacaagcgc ttctcgttg	2000
tggtatggc ggccctgggc ggggtgctgg gtgcgctgct gctgctggct	2050
ctccttgcc tcgccgtcct tgtccacaag cactatggcc cccggctcaa	2100
gtgctgctct ggcaaagctc cggagcccca gcccgaaggc ttgacaacc	2150
aggcgttct ccctgaccac aaggccaact gggcgccgt cccagcccc	2200
acgcacgacc ccaagccgc ggaggcacc atgcccgcag agccgcacc	2250
ccccggcct gcctccccag gcggtgcccc tgagcccccc gcagcgccc	2300
gagctggcg aagccccac gcggtgaggt ccactctgac caaggagcgg	2350

-continued

cgggcggagg gcggtacaa ggcgtcttg tttggcgagg acatcgggac	2400
ggaggcagac gtggtcgttc tcaacgcgc caccctggac gtggatggcg	2450
ccagtactc cggcagcggc gacgagggcg agggcgcggg gaggggtggg	2500
ggtccctacg atgcaccgg tggtgatgac tcctacatct aagtggcccc	2550
tccaccctct cccccagccg caccggcact ggaggtctcg ctccccagc	2600
ctccgaccgg aggcagaata aagcaaggct cccgaaacc aggccatggc	2650
gtggggcagg cgcgtgggtc cctgggggccc ccattcactc agtcccctgt	2700
cgtcattagc gcttgagccc aggtgtgcag atgaggcggg gggctctggc	2750
acgtgtctcc caccccaagg ctgcagcact tcccgtaaac cacctgcagt	2800
gcccgcgcc ttcccagggc tctgtgccag ctagtctggg aagttcctct	2850
cccgtcttaa ccacagccc aggggggctc cctcccccgc acctgcacca	2900
gagatctcag gcacccggct caactcagac ctcccgtctc cgaccctaca	2950
cagagattgc ctggggaggc tgaggagccg atgcaaacc ccaaggcgac	3000
gcacttgga gccggtggc tcaaacacct gccgggggtc ctagtcccct	3050
tctgaaatct acatgcttg gttggagcgc agcagtaaac accctgcccc	3100
gtgacctgga ctgaggcgcg ctgggggtgg gtgcgccgtg tggcctgagc	3150
aggagccaga ccaggaggcc taggggtgag agacacattc ccctcgctgc	3200
tcccaaagcc agagcccagg ctgggcgccc atgccagaa ccatcaaggg	3250
atcccttgog gcttgtcagc actttcccta atggaaatac accattaatt	3300
cctttccaaa tgtttt	3316

<210> SEQ ID NO 54

<211> LENGTH: 839

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 54

Met Gly Ser Trp Ala Leu Leu Trp Pro Pro Leu Leu Phe Thr Gly	
1 5 10 15	
Leu Leu Val Arg Pro Pro Gly Thr Met Ala Gln Ala Gln Tyr Cys	
20 25 30	
Ser Val Asn Lys Asp Ile Phe Glu Val Glu Glu Asn Thr Asn Val	
35 40 45	
Thr Glu Pro Leu Val Asp Ile His Val Pro Glu Gly Gln Glu Val	
50 55 60	
Thr Leu Gly Ala Leu Ser Thr Pro Phe Ala Phe Arg Ile Gln Gly	
65 70 75	
Asn Gln Leu Phe Leu Asn Val Thr Pro Asp Tyr Glu Glu Lys Ser	
80 85 90	
Leu Leu Glu Ala Gln Leu Leu Cys Gln Ser Gly Gly Thr Leu Val	
95 100 105	
Thr Gln Leu Arg Val Phe Val Ser Val Leu Asp Val Asn Asp Asn	
110 115 120	
Ala Pro Glu Phe Pro Phe Lys Thr Lys Glu Ile Arg Val Glu Glu	
125 130 135	
Asp Thr Lys Val Asn Ser Thr Val Ile Pro Glu Thr Gln Leu Gln	

-continued

140					145					150				
Ala	Glu	Asp	Arg	Asp	Lys	Asp	Asp	Ile	Leu	Phe	Tyr	Thr	Leu	Gln
				155					160					165
Glu	Met	Thr	Ala	Gly	Ala	Ser	Asp	Tyr	Phe	Ser	Leu	Val	Ser	Val
				170					175					180
Asn	Arg	Pro	Ala	Leu	Arg	Leu	Asp	Arg	Pro	Leu	Asp	Phe	Tyr	Glu
				185					190					195
Arg	Pro	Asn	Met	Thr	Phe	Trp	Leu	Leu	Val	Arg	Asp	Thr	Pro	Gly
				200					205					210
Glu	Asn	Val	Glu	Pro	Ser	His	Thr	Ala	Thr	Ala	Thr	Leu	Val	Leu
				215					220					225
Asn	Val	Val	Pro	Ala	Asp	Leu	Arg	Pro	Pro	Trp	Phe	Leu	Pro	Cys
				230					235					240
Thr	Phe	Ser	Asp	Gly	Tyr	Val	Cys	Ile	Gln	Ala	Gln	Tyr	His	Gly
				245					250					255
Ala	Val	Pro	Thr	Gly	His	Ile	Leu	Pro	Ser	Pro	Leu	Val	Leu	Arg
				260					265					270
Pro	Gly	Pro	Ile	Tyr	Ala	Glu	Asp	Gly	Asp	Arg	Gly	Ile	Asn	Gln
				275					280					285
Pro	Ile	Ile	Tyr	Ser	Ile	Phe	Arg	Gly	Asn	Val	Asn	Gly	Thr	Phe
				290					295					300
Ile	Ile	His	Pro	Asp	Ser	Gly	Asn	Leu	Thr	Val	Ala	Arg	Ser	Val
				305					310					315
Pro	Ser	Pro	Met	Thr	Phe	Leu	Leu	Leu	Val	Lys	Gly	Gln	Gln	Ala
				320					325					330
Asp	Leu	Ala	Arg	Tyr	Ser	Val	Thr	Gln	Val	Thr	Val	Glu	Ala	Val
				335					340					345
Ala	Ala	Ala	Gly	Ser	Pro	Pro	Arg	Phe	Pro	Gln	Ser	Leu	Tyr	Arg
				350					355					360
Gly	Thr	Val	Ala	Arg	Gly	Ala	Gly	Ala	Gly	Val	Val	Val	Lys	Asp
				365					370					375
Ala	Ala	Ala	Pro	Ser	Gln	Pro	Leu	Arg	Ile	Gln	Ala	Gln	Asp	Pro
				380					385					390
Glu	Phe	Ser	Asp	Leu	Asn	Ser	Ala	Ile	Thr	Tyr	Arg	Ile	Thr	Asn
				395					400					405
His	Ser	His	Phe	Arg	Met	Glu	Gly	Glu	Val	Val	Leu	Thr	Thr	Thr
				410					415					420
Thr	Leu	Ala	Gln	Ala	Gly	Ala	Phe	Tyr	Ala	Glu	Val	Glu	Ala	His
				425					430					435
Asn	Thr	Val	Thr	Ser	Gly	Thr	Ala	Thr	Thr	Val	Ile	Glu	Ile	Gln
				440					445					450
Val	Ser	Glu	Gln	Glu	Pro	Pro	Ser	Thr	Glu	Ala	Gly	Gly	Thr	Thr
				455					460					465
Gly	Pro	Trp	Thr	Ser	Thr	Thr	Ser	Glu	Val	Pro	Arg	Pro	Pro	Glu
				470					475					480
Pro	Ser	Gln	Gly	Pro	Ser	Thr	Thr	Ser	Ser	Gly	Gly	Gly	Thr	Gly
				485					490					495
Pro	His	Pro	Pro	Ser	Gly	Thr	Thr	Leu	Arg	Pro	Pro	Thr	Ser	Ser
				500					505					510
Thr	Pro	Gly	Gly	Pro	Pro	Gly	Ala	Glu	Asn	Ser	Thr	Ser	His	Gln
				515					520					525

-continued

Pro Ala Thr Pro Gly Gly Asp Thr Ala Gln Thr Pro Lys Pro Gly
530 535 540

Thr Ser Gln Pro Met Pro Pro Gly Val Gly Thr Ser Thr Ser His
545 550 555

Gln Pro Ala Thr Pro Ser Gly Gly Thr Ala Gln Thr Pro Glu Pro
560 565 570

Gly Thr Ser Gln Pro Met Pro Pro Ser Met Gly Thr Ser Thr Ser
575 580 585

His Gln Pro Ala Thr Pro Gly Gly Gly Thr Ala Gln Thr Pro Glu
590 595 600

Ala Gly Thr Ser Gln Pro Met Pro Pro Gly Met Gly Thr Ser Thr
605 610 615

Ser His Gln Pro Thr Thr Pro Gly Gly Gly Thr Ala Gln Thr Pro
620 625 630

Glu Pro Gly Thr Ser Gln Pro Met Pro Leu Ser Lys Ser Thr Pro
635 640 645

Ser Ser Gly Gly Gly Pro Ser Glu Asp Lys Arg Phe Ser Val Val
650 655 660

Asp Met Ala Ala Leu Gly Gly Val Leu Gly Ala Leu Leu Leu Leu
665 670 675

Ala Leu Leu Gly Leu Ala Val Leu Val His Lys His Tyr Gly Pro
680 685 690

Arg Leu Lys Cys Cys Ser Gly Lys Ala Pro Glu Pro Gln Pro Gln
695 700 705

Gly Phe Asp Asn Gln Ala Phe Leu Pro Asp His Lys Ala Asn Trp
710 715 720

Ala Pro Val Pro Ser Pro Thr His Asp Pro Lys Pro Ala Glu Ala
725 730 735

Pro Met Pro Ala Glu Pro Ala Pro Pro Gly Pro Ala Ser Pro Gly
740 745 750

Gly Ala Pro Glu Pro Pro Ala Ala Ala Arg Ala Gly Gly Ser Pro
755 760 765

Thr Ala Val Arg Ser Ile Leu Thr Lys Glu Arg Arg Pro Glu Gly
770 775 780

Gly Tyr Lys Ala Val Trp Phe Gly Glu Asp Ile Gly Thr Glu Ala
785 790 795

Asp Val Val Val Leu Asn Ala Pro Thr Leu Asp Val Asp Gly Ala
800 805 810

Ser Asp Ser Gly Ser Gly Asp Glu Gly Glu Gly Ala Gly Arg Gly
815 820 825

Gly Gly Pro Tyr Asp Ala Pro Gly Gly Asp Asp Ser Tyr Ile
830 835

<210> SEQ ID NO 55
<211> LENGTH: 3846
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 55

gcagctgggt tctcccggtt cccttgggca ggtgcagggt cgggttcaaa 50

gcctccggaa cgcgttttgg cctgatttga ggaggggggc ggggaggac 100

-continued

ctgcggcttg	cgcccccgcc	cccttctccg	gctcgcagcc	gaccggtaag	150
cccgcctcct	ccctcggccg	gccttggggc	cgtgtccgcc	gggcaactcc	200
agccgagggc	tgggcttctg	cctgcagggtg	tctgcggcga	ggcccctag	250
gtacagcccg	atttggtccc	atggtgggtt	tcggggccaa	ccggcgggct	300
ggccgcctgc	cctctctcgt	gctggtgggtg	ctgctgggtg	tgatcgtcgt	350
cctcgccttc	aactactgga	gcattctctc	ccgccacgtc	ctgcttcagg	400
aggaggtggc	cgagctgcag	ggccagggtcc	agcgcaccga	agtggcccgc	450
ggcgggctgg	aaaagcgcaa	ttcggacctc	ttgctgttgg	tggacacgca	500
caagaaacag	atcgaccaga	aggagggcga	ctacggccgc	ctcagcagcc	550
ggctgcaggc	cagagagggc	ctcgggaaga	gatgcgagga	tgacaagggt	600
aaactacaga	acaacatata	gtatcagatg	gcagacatac	atcatttaaa	650
ggagcaactt	gctgagcttc	gtcaggaatt	tcttcgacaa	gaagaccagc	700
ttcaggacta	taggaagaac	aatacttacc	ttgtgaagag	gttagaatat	750
gaaagttttc	agtgtggaca	gcagatgaag	gaattgagag	cacagcatga	800
agaaaaatatt	aaaaagttag	cagaccagtt	tttagaggaa	caaaagcaag	850
agacccaaaa	gattcaatca	aatgatggaa	aggaattgga	tataaacaat	900
caagtagtac	ctaaaaatat	tccaaaagta	gctgagaatg	ttgcagataa	950
gaatgaagaa	ccctcaagca	atcatattcc	acatgggaaa	gaacaaatca	1000
aaagagggtg	tgatgcaggg	atgcctggaa	tagaagagaa	tgacctagca	1050
aaagttagat	atcttcccc	tgctttaagg	aagcctccta	tttcagtttc	1100
tcaacatgaa	agtcatacaag	caatctccca	tcttccaact	ggacaacctc	1150
tctcccaaaa	tatgcctcca	gattcacaca	taaaccacaa	tggaaccccc	1200
ggtacttcaa	aacagaatcc	ttccagtcct	cttcagcgtt	taattccagg	1250
ctcaaaacttg	gacagtgaac	ccagaattca	aacagatata	ctaaagcagg	1300
ctaccaagga	cagagtcagt	gatttccata	aattgaagca	aaatgatgaa	1350
gaacgagagc	ttcaaatgga	tcctgcagac	tatggaaagc	aacatttcaa	1400
tgatgtcctt	taagtcctaa	aggaatgctt	cagaaaacct	aaagtgtctgt	1450
aaaatgaaat	cattctactt	tgtcctttct	gacttttggt	gtaaagacga	1500
attgtatcag	ttgtaaagat	acattgagat	agaattaagg	aaaaacttta	1550
atgaaggaat	gtacccatgt	acatatgtga	actttttcat	attgtattat	1600
caaggtatag	acttttttgg	ttatgataca	gttaagccaa	aaacagctaa	1650
tctttgcata	taaagcaaac	taatgtatat	ttcacatttt	attgagccga	1700
cttattttcca	caaatagata	aacaggacaa	aatagttgta	caggttatat	1750
gtggcatagc	ataaccacag	taagaacaga	acagatattc	agcagaaaac	1800
tttttatact	ctaattcttt	tttttttttt	tttgagacag	agtttttagtc	1850
ttgtttccca	ggctggagtg	caatggcaca	atcttggctc	actgcaacct	1900
ccgcctcctg	ggttcaggca	attttcctgc	ctcagcctcc	caagtagctg	1950
ggattacagg	caccaccacc	catgccagc	taatttttgt	atttttaata	2000

-continued

gagagctaataattgtatat ttaataaaga cgggtttcac catgttggcc	2050
aggctggtct tgaactcctg acctcaggtg atcctcctgc attggcctcc	2100
caaagtgtcg gaattccagg catgagccac tgcgcccagt ctacacacta	2150
attcttgtta gcccaacagc tgttctgttc tatctacccc tcatttcacg	2200
ctcaaggagt catacctaga atagttacac acaagaggga aactggaagc	2250
caaacactgt acagtattgt gtagaaagtc acctccctac tccttttatt	2300
ttacatgagt gctgatgtgt ttggcagat gagctttcag ctgaggcctg	2350
atggaaattg agataacctg caaagacata acagtattta tgagttatat	2400
cttagttctt gaaattgtgg aatgcatgat tgacaatata tttttaattt	2450
ttattttttc aagtaatacc agtactgttt aactatagcc agaactggct	2500
aaaattttta tattttcaga gttgaagttg gtgaagacat tcatgattta	2550
aacaccagat cctgaaaggg gttaaatcta ctttgaaatg aatctgcaat	2600
cagtatttca aagcttttct ggtaatttta gtgatcttat ttgattagac	2650
tttttcagaa gtactaaata aggaatttta acaggttttt attaatgcac	2700
agataaatag aagtacagtg aggtctatag ccatttttatt aaaatagctt	2750
aaaagtttgt aaaaaaatga atctttgtaa ttacttaata tgttagttaa	2800
gaacccgtca agcttatatt tgctagactt acaaattatt ttaaatgcat	2850
ttatcttttt tgacactatt cagtggaaatg tgtaagctag ctaattcttg	2900
ttttctgatt taaagcactt ttaaacttta tcctgcccc taaaaacaaa	2950
agggtttgat cacaagggga aatttaagat tgtaaccctt gtttttcaga	3000
agggtacttg ttaattgcac ataaacatga aatgtgtttt ccctgtgta	3050
ctaacacatt ctaggcaaaa ttcaacttta tagtggtaaa gaaacagggt	3100
gttctctgc tgagggtgcaa aaattcttaa gacttctggt tgaaattgct	3150
caatgactag gaaaagatgt agtagtttac taaaattggt tttctaccat	3200
atcaaattaa acaattcatg cttttatagg gtcaggccta caatgaatag	3250
gtatgggtgt ttcacagaat tttaaaatag agttaagggt aagtgatgta	3300
catttcgggg gcattagggt agggagatga atcaaaaaat acccctagta	3350
atgctttata ttttaatact gcaaaagctt tacaatgga aacctgcaa	3400
ttacctgcct tagttctttt gtcataaaaa caatcacttg gttggttgta	3450
ttgtagctat tacttataca gcaacatttc ttcaattagc agtctagaca	3500
ttttataaac agaaatcttg gaccaattga taatatttct gactgtatta	3550
atattttagt gctataaaat actatgtgaa tctcttaaaa atctgacatt	3600
ttacagtctg tatttagacat actgttttta taatgtttta cttctgcctt	3650
aagattttag ttttttaaat gtatttttgc cctgaattaa gtgttaattt	3700
gatggaaact ctgcttttaa aatcatcatt tactgggttc taataaatta	3750
aaaattaaac ttgaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	3800
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa	3846

-continued

<211> LENGTH: 380

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 56

```

Met Val Gly Phe Gly Ala Asn Arg Arg Ala Gly Arg Leu Pro Ser
 1          5          10          15

Leu Val Leu Val Val Leu Leu Val Val Ile Val Val Leu Ala Phe
 20          25          30

Asn Tyr Trp Ser Ile Ser Ser Arg His Val Leu Leu Gln Glu Glu
 35          40          45

Val Ala Glu Leu Gln Gly Gln Val Gln Arg Thr Glu Val Ala Arg
 50          55          60

Gly Arg Leu Glu Lys Arg Asn Ser Asp Leu Leu Leu Leu Val Asp
 65          70          75

Thr His Lys Lys Gln Ile Asp Gln Lys Glu Ala Asp Tyr Gly Arg
 80          85          90

Leu Ser Ser Arg Leu Gln Ala Arg Glu Gly Leu Gly Lys Arg Cys
 95          100          105

Glu Asp Asp Lys Val Lys Leu Gln Asn Asn Ile Ser Tyr Gln Met
 110          115          120

Ala Asp Ile His His Leu Lys Glu Gln Leu Ala Glu Leu Arg Gln
 125          130          135

Glu Phe Leu Arg Gln Glu Asp Gln Leu Gln Asp Tyr Arg Lys Asn
 140          145          150

Asn Thr Tyr Leu Val Lys Arg Leu Glu Tyr Glu Ser Phe Gln Cys
 155          160          165

Gly Gln Gln Met Lys Glu Leu Arg Ala Gln His Glu Glu Asn Ile
 170          175          180

Lys Lys Leu Ala Asp Gln Phe Leu Glu Glu Gln Lys Gln Glu Thr
 185          190          195

Gln Lys Ile Gln Ser Asn Asp Gly Lys Glu Leu Asp Ile Asn Asn
 200          205          210

Gln Val Val Pro Lys Asn Ile Pro Lys Val Ala Glu Asn Val Ala
 215          220          225

Asp Lys Asn Glu Glu Pro Ser Ser Asn His Ile Pro His Gly Lys
 230          235          240

Glu Gln Ile Lys Arg Gly Gly Asp Ala Gly Met Pro Gly Ile Glu
 245          250          255

Glu Asn Asp Leu Ala Lys Val Asp Asp Leu Pro Pro Ala Leu Arg
 260          265          270

Lys Pro Pro Ile Ser Val Ser Gln His Glu Ser His Gln Ala Ile
 275          280          285

Ser His Leu Pro Thr Gly Gln Pro Leu Ser Pro Asn Met Pro Pro
 290          295          300

Asp Ser His Ile Asn His Asn Gly Asn Pro Gly Thr Ser Lys Gln
 305          310          315

Asn Pro Ser Ser Pro Leu Gln Arg Leu Ile Pro Gly Ser Asn Leu
 320          325          330

Asp Ser Glu Pro Arg Ile Gln Thr Asp Ile Leu Lys Gln Ala Thr
 335          340          345

Lys Asp Arg Val Ser Asp Phe His Lys Leu Lys Gln Asn Asp Glu

```

-continued

350										355										360									
Glu	Arg	Glu	Leu	Gln	Met	Asp	Pro	Ala	Asp	Tyr	Gly	Lys	Gln	His															
				365						370				375															
Phe	Asn	Asp	Val	Leu																									
				380																									

<210> SEQ ID NO 57
<211> LENGTH: 841
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 57

ggatgggcga gcagtctgaa tgccagaatg gataaccgtt ttgctacagc	50
atttgtaatt gcttgtgtgc ttagcctcat ttccaccatc tacatggcag	100
cctccattgg cacagacttc tggatgaat atcgaagtcc agttcaagaa	150
aattccagtg atttgaataa aagcatctgg gatgaattca ttagtgatga	200
ggcagatgaa aagacttata atgatgcact ttttcgatac aatggcacag	250
tgggattgtg gagacggtgt atcacctac ccaaaaacat gcattggtat	300
agccccaccg aaaggacaga gtcatttgat gtggtcacia aatgtgtgag	350
tttcacacta actgagcagt tcatggagaa atttgttgat cccggaacc	400
acaatagcgg gattgatctc cttagacct atctttggcg ttgccagttc	450
cttttacctt ttgtgagttt aggtttgatg tgctttgggg ctttgatcgg	500
actttgtgot tgcatttgcc gaagcttata tcccaccatt gccacgggca	550
ttctccatct ccttcagat accatgctgt gaagtccagg ccacatggag	600
gtgtcctgtg tagatgctcc agctgaaatc ccaagctaag ctcccaactg	650
acagccaaca tcatttccag ccatgtgtgg gagccatcct ggatgtccag	700
ccttaacaag ccttcagagg acttcagcca cagctattat cttactacat	750
ccttgatgaga ctctaataaa gaaccaacta gctgagccca atcaacctat	800
ggaactgata gaaataaaat gaattgttgt tttgtgccgt t	841

<210> SEQ ID NO 58
<211> LENGTH: 184
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 58

Met	Asp	Asn	Arg	Phe	Ala	Thr	Ala	Phe	Val	Ile	Ala	Cys	Val	Leu	
1				5					10					15	
Ser	Leu	Ile	Ser	Thr	Ile	Tyr	Met	Ala	Ala	Ser	Ile	Gly	Thr	Asp	
			20					25						30	
Phe	Trp	Tyr	Glu	Tyr	Arg	Ser	Pro	Val	Gln	Glu	Asn	Ser	Ser	Asp	
			35					40						45	
Leu	Asn	Lys	Ser	Ile	Trp	Asp	Glu	Phe	Ile	Ser	Asp	Glu	Ala	Asp	
			50					55						60	
Glu	Lys	Thr	Tyr	Asn	Asp	Ala	Leu	Phe	Arg	Tyr	Asn	Gly	Thr	Val	
			65					70						75	
Gly	Leu	Trp	Arg	Arg	Cys	Ile	Thr	Ile	Pro	Lys	Asn	Met	His	Trp	
			80					85						90	

-continued

Tyr	Ser	Pro	Pro	Glu	Arg	Thr	Glu	Ser	Phe	Asp	Val	Val	Thr	Lys
				95					100					105
Cys	Val	Ser	Phe	Thr	Leu	Thr	Glu	Gln	Phe	Met	Glu	Lys	Phe	Val
				110					115					120
Asp	Pro	Gly	Asn	His	Asn	Ser	Gly	Ile	Asp	Leu	Leu	Arg	Thr	Tyr
				125					130					135
Leu	Trp	Arg	Cys	Gln	Phe	Leu	Leu	Pro	Phe	Val	Ser	Leu	Gly	Leu
				140					145					150
Met	Cys	Phe	Gly	Ala	Leu	Ile	Gly	Leu	Cys	Ala	Cys	Ile	Cys	Arg
				155					160					165
Ser	Leu	Tyr	Pro	Thr	Ile	Ala	Thr	Gly	Ile	Leu	His	Leu	Leu	Ala
				170					175					180
Asp	Thr	Met	Leu											

<210> SEQ ID NO 59
<211> LENGTH: 997
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 59

gcgtggacac cacctcagcc cactgagcag gagtcacagc acgaagacca	50
agcgcaaagc gacccttgcc ctccatcctg actgctcctc ctaagagaga	100
tggcaccggc cagagcagga ttctgcccc ttctgctgct tctgctgctg	150
gggctgtggg tggcagagat cccagtcagt gccaaagcca agggcatgac	200
ctcatcacag tggtttaaaa ttcagcacat gcagccagc cctcaagcat	250
gcaactcagc catgaaaaac attaacaagc acacaaaacg gtgcaaagac	300
ctcaacacct tcctgcacga gcctttctcc agtgtggccg ccacctgcca	350
gaccccaaaa atagcctgca agaatggcga taaaaactgc caccagagcc	400
acgggcccgt gtccctgacc atgtgtaagc tcacctcagg gaagtatccg	450
aactgcaggc acaaagagaa gcgacagaac aagtcttacg tagtggcctg	500
taagcctccc cagaaaaagg actctcagca attccacctg gttcctgtac	550
acttgacagc agtcctttag gtttcagac tggcttgctc tttggctgac	600
cttcaattcc ctctccagga ctccgcacca ctcccctaca cccagagcat	650
tctcttcccc tcatctcttg gggctgttcc tgggtcagcc tctgctggga	700
ggctgaagct gacactctgg tgagctgagc tctagaggga tggcttttca	750
tctttttgtt gctgttttcc cagatgctta tccccaaaga acagcaagct	800
caggtctgtg ggttccctgg tctatgccat tgcacatgtc tcccctgccc	850
cctggcatta gggcagcatg acaaggagag gaaataaatg gaaagggggc	900
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	950
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa	997

<210> SEQ ID NO 60
<211> LENGTH: 156
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 60

-continued

Met	Ala	Pro	Ala	Arg	Ala	Gly	Phe	Cys	Pro	Leu	Leu	Leu	Leu	Leu
1				5					10					15
Leu	Leu	Gly	Leu	Trp	Val	Ala	Glu	Ile	Pro	Val	Ser	Ala	Lys	Pro
			20						25					30
Lys	Gly	Met	Thr	Ser	Ser	Gln	Trp	Phe	Lys	Ile	Gln	His	Met	Gln
			35						40					45
Pro	Ser	Pro	Gln	Ala	Cys	Asn	Ser	Ala	Met	Lys	Asn	Ile	Asn	Lys
			50						55					60
His	Thr	Lys	Arg	Cys	Lys	Asp	Leu	Asn	Thr	Phe	Leu	His	Glu	Pro
			65						70					75
Phe	Ser	Ser	Val	Ala	Ala	Thr	Cys	Gln	Thr	Pro	Lys	Ile	Ala	Cys
			80						85					90
Lys	Asn	Gly	Asp	Lys	Asn	Cys	His	Gln	Ser	His	Gly	Pro	Val	Ser
			95						100					105
Leu	Thr	Met	Cys	Lys	Leu	Thr	Ser	Gly	Lys	Tyr	Pro	Asn	Cys	Arg
			110						115					120
Tyr	Lys	Glu	Lys	Arg	Gln	Asn	Lys	Ser	Tyr	Val	Val	Ala	Cys	Lys
			125						130					135
Pro	Pro	Gln	Lys	Lys	Asp	Ser	Gln	Gln	Phe	His	Leu	Val	Pro	Val
			140						145					150
His	Leu	Asp	Arg	Val	Leu									
					155									

<210> SEQ ID NO 61
<211> LENGTH: 520
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 61	
cggggtcatgc gccgccgcct gtggctgggc ctggcctggc tgctgctggc	50
gcggggcgccg gacgccgcgg gaaccccgag cgcgtcgcgg ggaccgcgca	100
gctaccgcga cctggagggc gacgtgcgct ggcggcgccct cttctcctcc	150
actcacttct tcctgcgcgt ggatcccggc ggccgcgtgc agggcaccgc	200
ctggcgccac ggccaggaca gcatcctgga gatccgctct gtacacgtgg	250
gcgtcgtggt catcaaagca gtgtcctcag gcttctacgt ggccatgaac	300
cgccggggcc gcctctacgg gtgcgcgactc tacaccgtgg actgcaggtt	350
ccgggagcgc atcgaagaga acggccacaa cacctacgcc tcacagcgct	400
ggcgccgcgc cggccagccc atgttcctgg cgctggacag gagggggggg	450
ccccggccag gcggccggac gcggcggtac cacctgtccg cccacttcct	500
gcccgtcctg gtctcctgag	520

<210> SEQ ID NO 62
<211> LENGTH: 170
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 62														
Met	Arg	Arg	Arg	Leu	Trp	Leu	Gly	Leu	Ala	Trp	Leu	Leu	Leu	Ala
1				5					10					15
Arg	Ala	Pro	Asp	Ala	Ala	Gly	Thr	Pro	Ser	Ala	Ser	Arg	Gly	Pro
				20					25					30

-continued

Arg Ser Tyr Pro His Leu Glu Gly Asp Val Arg Trp Arg Arg Leu
35 40 45
Phe Ser Ser Thr His Phe Phe Leu Arg Val Asp Pro Gly Gly Arg
50 55 60
Val Gln Gly Thr Arg Trp Arg His Gly Gln Asp Ser Ile Leu Glu
65 70 75
Ile Arg Ser Val His Val Gly Val Val Val Ile Lys Ala Val Ser
80 85 90
Ser Gly Phe Tyr Val Ala Met Asn Arg Arg Gly Arg Leu Tyr Gly
95 100 105
Ser Arg Leu Tyr Thr Val Asp Cys Arg Phe Arg Glu Arg Ile Glu
110 115 120
Glu Asn Gly His Asn Thr Tyr Ala Ser Gln Arg Trp Arg Arg Arg
125 130 135
Gly Gln Pro Met Phe Leu Ala Leu Asp Arg Arg Gly Gly Pro Arg
140 145 150
Pro Gly Gly Arg Thr Arg Arg Tyr His Leu Ser Ala His Phe Leu
155 160 165
Pro Val Leu Val Ser
170

<210> SEQ ID NO 63
<211> LENGTH: 2329
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 63

atccctcgac ctcgaccac gcgtccgctg gaaggtggcg tgcctctctc 50
tggtgtgttac catgcagctc ccactggccc tgtgtctcgt ctgcctgctg 100
gtacacacag ccttccgtgt agtggagggc caggggtggc aggcgttcaa 150
gaatgatgcc acggaatatc tccccgagct cggagagtac cccgagcctc 200
caccggagct ggagaacaac aagaccatga accgggcgga gaacggaggg 250
cggcctcccc accaccctt tgagacaaa gacgtgtccg agtacagctg 300
ccgcgagctg cacttcaccc gctacgtgac cgatgggccc tgccgcagcg 350
ccaagccggt caccgagctg gtgtgctccg gccagtgcgg cccggcgcg 400
ctgctgcccc acgccatcgg ccgcggcaag tggtggcgac ctagtgggcc 450
cgacttcgcg tgcaccccc accgtaccg cgcgagcgcg gtgcagctgc 500
tgtgtccccg tggtagaggc ccgcgcgcgc gcaaggtgcg cctgggtggc 550
tcgtgcaagt gcaagcgct caccgcctt cacaaccagt cggagctcaa 600
ggacttcggg accgaggccg ctgcggcgca gaagggccg aagcccgggc 650
cccgcgccc gagcgccaaa gccaacagg ccgagctgga gaacgcctac 700
tagagcccg ccgcgcccct cccaccggc gggcgcccc gccctgaacc 750
cgcgccccac atttctgtcc tctgcgctg gtttgattgt ttatatattca 800
ttgtaaatgc ctgcaacca gggcagggg ctgagacctt ccaggccctg 850
aggaatcccg ggcgcgggca agggccccct cagcccgcca gctgaggggt 900
cccacggggc aggggaggga attgagagtc acagacactg agccacgcag 950

-continued

```
ccccgcctct ggggccgcct acctttgctg gtcccacttc agaggaggca      1000
gaaatggaag cattttcacc gccctggggt ttttaaggag cgggtgtggga      1050
gtgggaaagt ccagggaactg gttaagaaag ttggataaga tcccccttg      1100
cacctcgctg cccatcagaa agcctgaggc gtgccagag cacaagactg      1150
ggggcaactg tagatgtggt ttctagtctt ggctctgcc ctaacttcct      1200
gtgtaacctt gaactacaca attctccttc gggacctcaa tttccacttt      1250
gtaaaatgag ggtggagggt ggaataggat ctcgaggaga ctattggcat      1300
atgattccaa ggactccagt gccttttgaa tgggcagagg tgagagagag      1350
agagagaaag agagagaatg aatgcagttg cattgattca gtgccaaggt      1400
cacttcagaa attcagagtt gtgatgctct cttctgacag ccaaagatga      1450
aaaaacaaca gaaaaaaaaa agtaaagagt ctatttatgg ctgacatatt      1500
tacggctgac aaactcctgg aagaagctat gctgcttccc agcctggcct      1550
ccccgtagt ttggctacct ccaccctcc atctcaaaga aataacatca      1600
tccattgggg tagaaaagga gagggtccga ggggtgtggg agggatagaa      1650
atcacatccg cccaacttc ccaaagagca gcacccctcc cccgacctat      1700
agccatgttt taaagtcacc ttccgaagag aagtgaaggt ttcaaggaca      1750
ctggccttgc agggccgagg gagcagccat caaaaactca cagaccagca      1800
catccctttt gagacaccgc cttctgccca ccactcacgg acacatttct      1850
gcctagaaaa cagcttctta ctgctcttac atgtgatggc atatcttaca      1900
ctaaaagaat attattgggg gaaaaactac aagtgtgtga catatgtgta      1950
gaaactgcag agcataatag ctgccacca aaaatctttt tgaaaatcat      2000
ttccagacaa cctcttactt tctgtgtagt ttttaattgt taaaaaaaaa      2050
aagttttaa cagaagcaca tgacatatga aagcctgcag gactggtcgt      2100
ttttttgcca attcttccac gtgggacttg tccacaagaa tgaaagtagt      2150
ggttttttaa gagttaagtt acatatttat tttctcactt aagttattta      2200
tgcaaaagtt tttctgtag agaatgacaa tgtaaatatt gctttatgaa      2250
ttaacagtct gttcttccag agtccagaga cattgttaat aaagacaatg      2300
aatcatgaaa aaaaaaaaaa aaaaaaaaaa      2329
```

<210> SEQ ID NO 64

<211> LENGTH: 213

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 64

```
Met Gln Leu Pro Leu Ala Leu Cys Leu Val Cys Leu Leu Val His
 1             5             10             15
Thr Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys
                20             25             30
Asn Asp Ala Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu
                35             40             45
Pro Pro Pro Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu
                50             55             60
```

-continued

Asn	Gly	Gly	Arg	Pro	Pro	His	His	Pro	Phe	Glu	Thr	Lys	Asp	Val
				65					70					75
Ser	Glu	Tyr	Ser	Cys	Arg	Glu	Leu	His	Phe	Thr	Arg	Tyr	Val	Thr
				80					85					90
Asp	Gly	Pro	Cys	Arg	Ser	Ala	Lys	Pro	Val	Thr	Glu	Leu	Val	Cys
				95					100					105
Ser	Gly	Gln	Cys	Gly	Pro	Ala	Arg	Leu	Leu	Pro	Asn	Ala	Ile	Gly
				110					115					120
Arg	Gly	Lys	Trp	Arg	Pro	Ser	Gly	Pro	Asp	Phe	Arg	Cys	Ile	
				125					130					135
Pro	Asp	Arg	Tyr	Arg	Ala	Gln	Arg	Val	Gln	Leu	Leu	Cys	Pro	Gly
				140					145					150
Gly	Glu	Ala	Pro	Arg	Ala	Arg	Lys	Val	Arg	Leu	Val	Ala	Ser	Cys
				155					160					165
Lys	Cys	Lys	Arg	Leu	Thr	Arg	Phe	His	Asn	Gln	Ser	Glu	Leu	Lys
				170					175					180
Asp	Phe	Gly	Thr	Glu	Ala	Ala	Arg	Pro	Gln	Lys	Gly	Arg	Lys	Pro
				185					190					195
Arg	Pro	Arg	Ala	Arg	Ser	Ala	Lys	Ala	Asn	Gln	Ala	Glu	Leu	Glu
				200					205					210
Asn	Ala	Tyr												

<210> SEQ ID NO 65
<211> LENGTH: 2663
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 65

cccactcggc	ggtttgccgg	gagggagggg	ctttgcgcag	gccccgctcc	50
cgccccgcct	ccatgcggcc	cgccccgatt	gcgctgtggc	tgcgccctggt	100
cttggccctg	gcccttgtcc	gccccggggc	tgtgggggtg	gccccggtcc	150
gagcccccat	ctatgtcagc	agctggggcg	tccagggtgc	ccagggtaac	200
cgggaggtcg	agcgccctgg	acgcaaattc	ggcttcgtca	acctggggcc	250
gatcttctct	gacgggcagt	actttcacct	gcggcaccgg	ggcgtggctc	300
agcagtcctc	gaccccgcac	tggggccacc	gcctgcacct	gaagaaaaac	350
cccaaggctc	agtggttcca	gcagcagacg	ctgcagcggc	gggtgaaacg	400
ctctgtcgtg	gtgcccacgg	acccctggtt	ctccaagcag	tggtacatga	450
acagcgaggc	ccaaccagac	ctgagcatcc	tgcaggcctg	gagtcagggg	500
ctgtcaggcc	agggcatcgt	ggtctctgtg	ctggacgatg	gcatcgagaa	550
ggaccaccgc	gacctctggg	ccaactacga	cccctgggcc	agctatgact	600
tcaatgacta	cgaccgggac	ccccagcccc	gctacacccc	cagcaaagag	650
aaccggcacg	ggaccgcgtg	tgctggggag	gtggccgcga	tggccaacaa	700
tggcttctgt	ggtgtggggg	tcgctttcaa	cgcccgaaatc	ggaggcgtag	750
ggatgctgga	cggtaccatc	accgatgtca	tcgaggccca	gtcgctgagc	800
ctgcagccgc	agcacatcca	catttacagc	gccagctggg	gtcccagagga	850
cgacggccgc	acggtggacg	gccccggcat	cctcaccgcg	gaggccttcc	900

-continued

ggcgtggtgt gaccaagggc cgcggcgggc tgggcacgct cttcatctgg	950
gcctcgggca acggcggcct gcactacgac aactgcaact gcgacggcta	1000
caccaacagc atccacacgc ttccgtggg cagcaccacc cagcagggcc	1050
gcgtgccctg gtacagcgaa gcctgcgcct ccaccctcac caccacctac	1100
agcagcggcg tggccaccga cccccagatc gtcaccacgg acctgcatca	1150
cgggtgcaca gaccagcaca cgggcaccto ggcctcagcc ccactggcgg	1200
ccggcatgat cgccctagcg ctggaggcca acccgttcct gacgtggaga	1250
gacatgcagc acctggtggt ccgcgcgtcc aagccggcgc acctgcaggc	1300
cgaggactgg aggaccaacg gcgtggggcg ccaagtgagc catcactacg	1350
gatacgggct gctggacgcc gggctgctgg tggacaccgc ccgcacctgg	1400
ctgcccaccc agccgcagag gaagtgcgcc gtccgggtcc agagccgcc	1450
cacccccatc ctgccgtga tctacatcag ggaaaacgta tcggcctgcg	1500
ccggcctcca caactccatc cgctcgctgg agcacgtgca ggcgcagctg	1550
acgtgtcct acagccggcg cggagacctg gagatctcgc tcaccagccc	1600
catgggcacg cgctccacac tcgtggccat acgacccttg gacgtcagca	1650
ctgaaggcta caacaactgg gtcttcatgt cccccactt ctgggatgag	1700
aaccacaggg gcgtgtggac cctgggccta gagaacaagg gctactattt	1750
caacacgggg acgttgtacc gctacacgct gctgctctat gggacggccg	1800
aggacatgac agcgcggcct acaggccccc aggtgaccag cagcgcgtgt	1850
gtgcagcggg acacagaggg gctgtgccag gcgtgtgacg gccccgcta	1900
catcctggga cagctctgcc tggcctactg cccccgcgg ttcttcaacc	1950
acacaaggct ggtgaccgct gggcctgggc acacggcggc gcccgcgctg	2000
agggctctgt ccagctgcca tgctcctgc tacacctgcc gcggcggctc	2050
cccaggagac tgcacctcct gtccccatc ctccacgctg gaccagcagc	2100
agggctcctg catgggaccc accacccccg acagccgccc ccggcttaga	2150
gctgccgcct gtccccacca ccgctgccca gcctcgcca tgggtgtgag	2200
cctcctggcc gtgacctcgc gagggcccg cctctgcggc atgtccatgg	2250
acctcccaact atacgcctgg ctctcccgtg ccagggccac cccaccaaa	2300
ccccaggctc ggttgccagc tggaaacctga agttgtcagc tcagaaagcg	2350
accttgcccc cgctgggct cctgacagc actgctgcca tgctgcctcc	2400
ccaggctggc ccagaggag cgagcaccag caccgacgc ctggcctgcc	2450
agggatgggc cccgtggaac cccgaagcct ggcgggagag agagagagag	2500
aagtctctc tgcatcttgg gtttgggcag gagtgggctg gggggagagg	2550
ctggagcacc ccaaaagcca ggggaaagt gagggagaga aacgtgacac	2600
tgtccgtctc gggcacgcgc tccaacctca gagtttgcaa ataaaggttg	2650
cttagaaggt gaa	2663

<210> SEQ ID NO 66

<211> LENGTH: 755

-continued

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 66

```

Met Arg Pro Ala Pro Ile Ala Leu Trp Leu Arg Leu Val Leu Ala
 1           5           10           15

Leu Ala Leu Val Arg Pro Arg Ala Val Gly Trp Ala Pro Val Arg
 20           25           30

Ala Pro Ile Tyr Val Ser Ser Trp Ala Val Gln Val Ser Gln Gly
 35           40           45

Asn Arg Glu Val Glu Arg Leu Ala Arg Lys Phe Gly Phe Val Asn
 50           55           60

Leu Gly Pro Ile Phe Ser Asp Gly Gln Tyr Phe His Leu Arg His
 65           70           75

Arg Gly Val Val Gln Gln Ser Leu Thr Pro His Trp Gly His Arg
 80           85           90

Leu His Leu Lys Lys Asn Pro Lys Val Gln Trp Phe Gln Gln Gln
 95           100          105

Thr Leu Gln Arg Arg Val Lys Arg Ser Val Val Val Pro Thr Asp
110           115          120

Pro Trp Phe Ser Lys Gln Trp Tyr Met Asn Ser Glu Ala Gln Pro
125           130          135

Asp Leu Ser Ile Leu Gln Ala Trp Ser Gln Gly Leu Ser Gly Gln
140           145          150

Gly Ile Val Val Ser Val Leu Asp Asp Gly Ile Glu Lys Asp His
155           160          165

Pro Asp Leu Trp Ala Asn Tyr Asp Pro Leu Ala Ser Tyr Asp Phe
170           175          180

Asn Asp Tyr Asp Pro Asp Pro Gln Pro Arg Tyr Thr Pro Ser Lys
185           190          195

Glu Asn Arg His Gly Thr Arg Cys Ala Gly Glu Val Ala Ala Met
200           205          210

Ala Asn Asn Gly Phe Cys Gly Val Gly Val Ala Phe Asn Ala Arg
215           220          225

Ile Gly Gly Val Arg Met Leu Asp Gly Thr Ile Thr Asp Val Ile
230           235          240

Glu Ala Gln Ser Leu Ser Leu Gln Pro Gln His Ile His Ile Tyr
245           250          255

Ser Ala Ser Trp Gly Pro Glu Asp Asp Gly Arg Thr Val Asp Gly
260           265          270

Pro Gly Ile Leu Thr Arg Glu Ala Phe Arg Arg Gly Val Thr Lys
275           280          285

Gly Arg Gly Gly Leu Gly Thr Leu Phe Ile Trp Ala Ser Gly Asn
290           295          300

Gly Gly Leu His Tyr Asp Asn Cys Asn Cys Asp Gly Tyr Thr Asn
305           310          315

Ser Ile His Thr Leu Ser Val Gly Ser Thr Thr Gln Gln Gly Arg
320           325          330

Val Pro Trp Tyr Ser Glu Ala Cys Ala Ser Thr Leu Thr Thr Thr
335           340          345

Tyr Ser Ser Gly Val Ala Thr Asp Pro Gln Ile Val Thr Thr Asp
350           355          360

```

-continued

Leu	His	His	Gly	Cys	Thr	Asp	Gln	His	Thr	Gly	Thr	Ser	Ala	Ser	365	370	375
Ala	Pro	Leu	Ala	Ala	Gly	Met	Ile	Ala	Leu	Ala	Leu	Glu	Ala	Asn	380	385	390
Pro	Phe	Leu	Thr	Trp	Arg	Asp	Met	Gln	His	Leu	Val	Val	Arg	Ala	395	400	405
Ser	Lys	Pro	Ala	His	Leu	Gln	Ala	Glu	Asp	Trp	Arg	Thr	Asn	Gly	410	415	420
Val	Gly	Arg	Gln	Val	Ser	His	His	Tyr	Gly	Tyr	Gly	Leu	Leu	Asp	425	430	435
Ala	Gly	Leu	Leu	Val	Asp	Thr	Ala	Arg	Thr	Trp	Leu	Pro	Thr	Gln	440	445	450
Pro	Gln	Arg	Lys	Cys	Ala	Val	Arg	Val	Gln	Ser	Arg	Pro	Thr	Pro	455	460	465
Ile	Leu	Pro	Leu	Ile	Tyr	Ile	Arg	Glu	Asn	Val	Ser	Ala	Cys	Ala	470	475	480
Gly	Leu	His	Asn	Ser	Ile	Arg	Ser	Leu	Glu	His	Val	Gln	Ala	Gln	485	490	495
Leu	Thr	Leu	Ser	Tyr	Ser	Arg	Arg	Gly	Asp	Leu	Glu	Ile	Ser	Leu	500	505	510
Thr	Ser	Pro	Met	Gly	Thr	Arg	Ser	Thr	Leu	Val	Ala	Ile	Arg	Pro	515	520	525
Leu	Asp	Val	Ser	Thr	Glu	Gly	Tyr	Asn	Asn	Trp	Val	Phe	Met	Ser	530	535	540
Thr	His	Phe	Trp	Asp	Glu	Asn	Pro	Gln	Gly	Val	Trp	Thr	Leu	Gly	545	550	555
Leu	Glu	Asn	Lys	Gly	Tyr	Tyr	Phe	Asn	Thr	Gly	Thr	Leu	Tyr	Arg	560	565	570
Tyr	Thr	Leu	Leu	Leu	Tyr	Gly	Thr	Ala	Glu	Asp	Met	Thr	Ala	Arg	575	580	585
Pro	Thr	Gly	Pro	Gln	Val	Thr	Ser	Ser	Ala	Cys	Val	Gln	Arg	Asp	590	595	600
Thr	Glu	Gly	Leu	Cys	Gln	Ala	Cys	Asp	Gly	Pro	Ala	Tyr	Ile	Leu	605	610	615
Gly	Gln	Leu	Cys	Leu	Ala	Tyr	Cys	Pro	Pro	Arg	Phe	Phe	Asn	His	620	625	630
Thr	Arg	Leu	Val	Thr	Ala	Gly	Pro	Gly	His	Thr	Ala	Ala	Pro	Ala	635	640	645
Leu	Arg	Val	Cys	Ser	Ser	Cys	His	Ala	Ser	Cys	Tyr	Thr	Cys	Arg	650	655	660
Gly	Gly	Ser	Pro	Arg	Asp	Cys	Thr	Ser	Cys	Pro	Pro	Ser	Ser	Thr	665	670	675
Leu	Asp	Gln	Gln	Gln	Gly	Ser	Cys	Met	Gly	Pro	Thr	Thr	Pro	Asp	680	685	690
Ser	Arg	Pro	Arg	Leu	Arg	Ala	Ala	Ala	Cys	Pro	His	His	Arg	Cys	695	700	705
Pro	Ala	Ser	Ala	Met	Val	Leu	Ser	Leu	Leu	Ala	Val	Thr	Leu	Gly	710	715	720
Gly	Pro	Val	Leu	Cys	Gly	Met	Ser	Met	Asp	Leu	Pro	Leu	Tyr	Ala	725	730	735

-continued

Trp Leu Ser Arg Ala Arg Ala Thr Pro Thr Lys Pro Gln Val Trp
740 745 750

Leu Pro Ala Gly Thr
755

<210> SEQ ID NO 67
<211> LENGTH: 332
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 67

atgaggaagc tccagggcag gatggtttac ctgcctggac agcaagatga 50
tggtacact agcccccatt ctctgggcgc ctggatttgc ccaccagatc 100
tcctcacctc ttgcccttca cctcctgctg tacctacaag gtctccccga 150
ttctcatctg ccataatca tggacacagc cccaggatgt gcaggactct 200
cagggaccat ctggagtcc agctggaatc tgggcctggt ggagtgggag 250
tggggcaggg gcctgcattg ggctgactta gagagcacag ttattccatc 300
catatggaaa taaacatttt ggattcctga tc 332

<210> SEQ ID NO 68
<211> LENGTH: 88
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 68

Met Met Ala Thr Leu Ala Pro Ile Leu Trp Ala Pro Gly Phe Ala
1 5 10 15
His Gln Ile Ser Ser Pro Leu Ala Leu His Leu Leu Leu Tyr Leu
20 25 30
Gln Gly Leu Pro Asp Ser His Leu Pro Ile Ile Met Asp Thr Ala
35 40 45
Pro Gly Cys Ala Gly Leu Ser Gly Thr Ile Trp Ser Ser Ser Trp
50 55 60
Asn Leu Gly Leu Val Glu Trp Glu Trp Gly Arg Gly Leu His Trp
65 70 75
Ala Asp Leu Glu Ser Thr Val Ile Pro Ser Ile Trp Lys
80 85

<210> SEQ ID NO 69
<211> LENGTH: 1302
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: 1218-1253
<223> OTHER INFORMATION: unknown base

<400> SEQUENCE: 69

tttgcatggtt ggtcctcctc tggcctcctg cccctcctgc tgetgtgtgt 50
gcttcatttg ctggcagccc aggggtgggg tggcctgcag gcagcgctgc 100
tggcccttga ggtggggctg gtgggtctg gggcctccta cctgtcctt 150
tgtacagccc tgcacctgcc ctccagtctt ttcctactcc tggcccaggg 200
taccgcactg ggggccgtcc tgggcctgag ctgggcgcga ggcctcatgg 250

-continued	
gtgttccccct gggccttggg gctgcctggc tcttagcttg gccaggccta	300
gctctacctc tggtagctat ggcagcgggg ggcagatggg tgcggcagca	350
gggccccccg gtgcgcgggg gcatactctg actctggttg cgggttctgc	400
tgcgcctgtc acccatggcc ttccggggccc tgcagggttg tggggctgtg	450
ggggaccggg gtctgtttgc actgtacccc aaaaccaaca aggatggctt	500
ccgcagccgc ctgcccgtcc ctgggccccg gcgcgtaat ccccgcacca	550
cccaacaccc attagctctg ttggcaaggg tctgggtcct gtgcaagggc	600
tggaaactggc gtctggcacg ggccagccag ggtttagcat ccacttgcc	650
cccgtgggcc atccacacac tggccagctg gggcctgctt cggggtgaac	700
ggcccaccgc aatcccccg ctactaccac gcagccagcg ccagctaggg	750
ccccctgcct ccgcccagcc actgccaggg actctagccg ggcggaggtc	800
acgcaccgc cagtcccggg ccctgcccc ctggaggtag ctgactccag	850
cccttcacgc ccaaacttag agcattgagc actttatctc ccacgactca	900
gtgaagtttc tccagtccct agtcctctct tttcacccac cttoctcagt	950
ttgtcactt accccaggcc cagcccttcg gacctctaga caggcagcct	1000
cctcagctgt ggagtccagc agtcactctg tgttctcctg gcgtcctcc	1050
cctaagtatt tgctgttcgc ccgctgtgtg tgctcctcct caccctcatt	1100
gactcaggcc tggggccagg ggtggtggag ggtgggaaga gtcatgttt	1150
ttttctctc tttgattttg tttttctgtc tcccttccaa cctgtcccct	1200
tccccccacc aaaaaaannnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	1250
nnnaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	1300
aa	1302

<210> SEQ ID NO 70
<211> LENGTH: 197
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 70

Met Gly Val Pro Leu Gly Leu Gly Ala Ala Trp Leu Leu Ala Trp
1 5 10 15

Pro Gly Leu Ala Leu Pro Leu Val Ala Met Ala Ala Gly Gly Arg
20 25 30

Trp Val Arg Gln Gln Gly Pro Arg Val Arg Arg Gly Ile Ser Arg
35 40 45

Leu Trp Leu Arg Val Leu Leu Arg Leu Ser Pro Met Ala Phe Arg
50 55 60

Ala Leu Gln Gly Cys Gly Ala Val Gly Asp Arg Gly Leu Phe Ala
65 70 75

Leu Tyr Pro Lys Thr Asn Lys Asp Gly Phe Arg Ser Arg Leu Pro
80 85 90

Val Pro Gly Pro Arg Arg Arg Asn Pro Arg Thr Thr Gln His Pro
95 100 105

Leu Ala Leu Leu Ala Arg Val Trp Val Leu Cys Lys Gly Trp Asn
110 115 120

-continued

Trp	Arg	Leu	Ala	Arg	Ala	Ser	Gln	Gly	Leu	Ala	Ser	His	Leu	Pro
			125						130					135
Pro	Trp	Ala	Ile	His	Thr	Leu	Ala	Ser	Trp	Gly	Leu	Leu	Arg	Gly
			140						145					150
Glu	Arg	Pro	Thr	Arg	Ile	Pro	Arg	Leu	Leu	Pro	Arg	Ser	Gln	Arg
			155						160					165
Gln	Leu	Gly	Pro	Pro	Ala	Ser	Arg	Gln	Pro	Leu	Pro	Gly	Thr	Leu
			170						175					180
Ala	Gly	Arg	Arg	Ser	Arg	Thr	Arg	Gln	Ser	Arg	Ala	Leu	Pro	Pro
			185						190					195
Trp	Arg													

<210> SEQ ID NO 71
<211> LENGTH: 1976
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 71

gtttgggggt	tgtttgggat	tagtgaagct	actgcctttg	ccgccagcgc	50
agcctcagag	tttgattatt	tgcaatgtca	ggctttgaaa	acttaaacac	100
ggattttctac	cagacaagtt	acagcatcga	tgatcagtca	cagcagtcct	150
atgattatgg	aggaagtgga	ggaccctata	gcaaacagta	tgctggctat	200
gactatttcg	agcaaggcag	atttgtccct	ccagacatga	tgcaagccaca	250
acagccatac	accgggcaga	tttaccagcc	aactcaggca	tatactccag	300
cttcacctca	gcctttctat	ggaaacaact	ttgaggatga	gccaccttta	350
ttagaagagt	taggtatcaa	ttttgaccac	atctggcaaa	aaacactaac	400
agtattacat	ccgttaaaa	tagcagatgg	cagcatcatg	aatgaaactg	450
atttggcagg	tccaatgggt	ttttgccttg	cttttgagc	cacattgcta	500
ctggctggca	aaatccagtt	tggctatgta	tacgggatca	gtgcaattgg	550
atgtctagga	atgttttgtt	tattaaactt	aatgagtatg	acaggtgttt	600
catttggttg	tgtggcaagt	gtccttggat	attgtcttct	gcccattgatc	650
ctactttcca	gctttgcagt	gatattttct	ttgcaaggaa	tggttaggaat	700
cattctcact	gctgggatta	ttggatgttg	tagtttttct	gcttccaaaa	750
tatttatattc	tgcattagcc	atggaaggac	agcaactttt	agtagcatat	800
ccttgcgctt	tggtatatgg	agtctttgcc	ctgatttccg	tcttttgaaa	850
atztatctgg	gatgtggaca	tcagtgggcc	agatgtacaa	aaaggacctt	900
gaactcttaa	attggaccag	caaactgctg	cagcgcaact	ctcatgcaga	950
tttacatttg	actgttgag	caatgaaagt	aaacgtgtat	ctcttgttca	1000
tttttataga	acttttgcatt	actatattgg	atttacctgc	ggtgtgacta	1050
gctttaaatg	tttgtgttta	tacagataag	aaatgctatt	tctttctggt	1100
tcctgcagcc	attgaaaaac	ctttttcctt	gcaaattata	atgtttttga	1150
tagattttta	tcaactgtgg	gaaaccaaac	acaaagctga	taacctttct	1200
taaaaacgac	ccagtcacag	taaagaagac	acaagacggc	cgggcgtggg	1250
agctcacgcc	tgtaaatccca	gcactttggg	aggccgaggg	gggcggatca	1300

-continued

```
caagggcagg agatcgagac catcctggtt aacacggtga aaccccgact      1350
ctactaaaaa tacaaaaaaa attagctggg cgtggtggcg ggcgcctgta      1400
gtcccagcta ctcaggaggc tgaggcagga gaagtgtgaa cccaggaggc      1450
ggagcttgca gtgagccgag atcacaccac tgcactccat ccagcctggg      1500
tgacaggggtg agactctgtc tcaaaaaaaa aaaaaaaagg agacacaaga      1550
cttactgcaa aaatattttt ccaaggattt aggaaagaaa aattgccttg      1600
tattctcaag tcaggttaact caaagcaaaa aagtgatcca aatgtagagt      1650
atgagtttgc actccaaaaa tttagacatta ctgtaaatta tctcatggaa      1700
tttttgctaa aattcagaga tacgggaagt tcacaatcta cctcattgta      1750
gacatgaaat gcgaacactt acttacatat taatgttaac tcaaccttag      1800
ggacctggaa tggttgcatt aatgctataa tcgttggatc gccacatttc      1850
ccaaaaataa taaaaaaatc actaaccttt ttttaagaaa atatttaaag      1900
ttttacaaaa ttcaatattg caattatcaa tgtaaagtac atttgaatgc      1950
ttattaaaac ttccaatt aattttt                                1976
```

<210> SEQ ID NO 72
<211> LENGTH: 257
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 72

```
Met Ser Gly Phe Glu Asn Leu Asn Thr Asp Phe Tyr Gln Thr Ser
 1          5          10          15
Tyr Ser Ile Asp Asp Gln Ser Gln Gln Ser Tyr Asp Tyr Gly Gly
          20          25          30
Ser Gly Gly Pro Tyr Ser Lys Gln Tyr Ala Gly Tyr Asp Tyr Ser
          35          40          45
Gln Gln Gly Arg Phe Val Pro Pro Asp Met Met Gln Pro Gln Gln
          50          55          60
Pro Tyr Thr Gly Gln Ile Tyr Gln Pro Thr Gln Ala Tyr Thr Pro
          65          70          75
Ala Ser Pro Gln Pro Phe Tyr Gly Asn Asn Phe Glu Asp Glu Pro
          80          85          90
Pro Leu Leu Glu Glu Leu Gly Ile Asn Phe Asp His Ile Trp Gln
          95          100         105
Lys Thr Leu Thr Val Leu His Pro Leu Lys Val Ala Asp Gly Ser
          110         115         120
Ile Met Asn Glu Thr Asp Leu Ala Gly Pro Met Val Phe Cys Leu
          125         130         135
Ala Phe Gly Ala Thr Leu Leu Leu Ala Gly Lys Ile Gln Phe Gly
          140         145         150
Tyr Val Tyr Gly Ile Ser Ala Ile Gly Cys Leu Gly Met Phe Cys
          155         160         165
Leu Leu Asn Leu Met Ser Met Thr Gly Val Ser Phe Gly Cys Val
          170         175         180
Ala Ser Val Leu Gly Tyr Cys Leu Leu Pro Met Ile Leu Leu Ser
          185         190         195
```

-continued

Ser	Phe	Ala	Val	Ile	Phe	Ser	Leu	Gln	Gly	Met	Val	Gly	Ile	Ile
			200						205				210	
Leu	Thr	Ala	Gly	Ile	Ile	Gly	Trp	Cys	Ser	Phe	Ser	Ala	Ser	Lys
			215						220				225	
Ile	Phe	Ile	Ser	Ala	Leu	Ala	Met	Glu	Gly	Gln	Gln	Leu	Leu	Val
			230						235				240	
Ala	Tyr	Pro	Cys	Ala	Leu	Leu	Tyr	Gly	Val	Phe	Ala	Leu	Ile	Ser
			245						250				255	
Val	Phe													

<210> SEQ ID NO 73
<211> LENGTH: 1285
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 73

acactggcca aaacgcggct cgccctcggc tgcgctcggc tcccgcgggc	50
gctcgggccc gagcccctcc tccccctacc cgccggcccg acagggagga	100
gccaatggct gggcctgcca tccacaccgc tcccatgctg ttcctcgtcc	150
tcctgctgcc ccagctgagc ctggcaggcg cccttgacc tgggacccct	200
gcccgaacc tccctgagaa tcacattgac ctcccaggcc cagcgctgtg	250
gacgcctcag gccagccacc accgcgcggc gggcccgggc aagaaggagt	300
ggggcccagg cctgccagc caggcccagg atggggctgt ggtcaccgcc	350
accaggcagg cctccaggct gccagaggct gaggggctgc tgcctgagca	400
gagtctgca ggcctgctgc aggacaagga cctgctcctg ggactggcat	450
tgccctaccc cgagaaggag aacagacctc caggttgga gaggaccagg	500
aaacgcagca gggagcaca gagacgcagg gacaggttga ggctgcacca	550
aggccgagcc ttggtccgag gtcccagctc cctgatgaag aaggcagagc	600
tctccgaagc ccagggtctg gatgcagcca tggaggaatc ctccaccagc	650
ctggcgccca ccatgttctt tctcaccacc tttgaggcag cacctgccac	700
agaagagtcc ctgatcctgc ccgtcacctc cctgcggccc cagcaggcac	750
agcccaggtc tgacggggag gtgatgcca cgctggacat ggccttgttc	800
gactggaccg attatgaaga cttaaacct gatggttggc cctctgcaaa	850
gaagaaagag aaacaccgcg gtaaaacttc cagtgatggt aacgaacat	900
caccagccga aggggaacca tgcgaccatc accaagactg cctgccaggg	950
acttgctgcg acctgcggga gcatctctgc acaccccaca accgaggcct	1000
caacaacaaa tgcttcgatg actgcatgtg tgtggaaggg ctgcgctgct	1050
atgccaaatt ccaccggaac cgcagggtta cacggaggaa agggcgctgt	1100
gtggagcccg agacggccaa cggcgaccag ggatccttca tcaacgtcta	1150
gcggccccgc gggactgggg actgagccca ggaggtttgc acaagccggg	1200
cgatttgttt gtaactagca gtgggagatc aagttgggga acagatggct	1250
gaggctgcag actcaggccc aggacactca accccc	1285

<210> SEQ ID NO 74

-continued

<211> LENGTH: 348

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 74

```

Met Ala Gly Pro Ala Ile His Thr Ala Pro Met Leu Phe Leu Val
 1          5          10          15
Leu Leu Leu Pro Gln Leu Ser Leu Ala Gly Ala Leu Ala Pro Gly
 20          25          30
Thr Pro Ala Arg Asn Leu Pro Glu Asn His Ile Asp Leu Pro Gly
 35          40          45
Pro Ala Leu Trp Thr Pro Gln Ala Ser His His Arg Arg Arg Gly
 50          55          60
Pro Gly Lys Lys Glu Trp Gly Pro Gly Leu Pro Ser Gln Ala Gln
 65          70          75
Asp Gly Ala Val Val Thr Ala Thr Arg Gln Ala Ser Arg Leu Pro
 80          85          90
Glu Ala Glu Gly Leu Leu Pro Glu Gln Ser Pro Ala Gly Leu Leu
 95          100         105
Gln Asp Lys Asp Leu Leu Leu Gly Leu Ala Leu Pro Tyr Pro Glu
110          115         120
Lys Glu Asn Arg Pro Pro Gly Trp Glu Arg Thr Arg Lys Arg Ser
125          130         135
Arg Glu His Lys Arg Arg Arg Asp Arg Leu Arg Leu His Gln Gly
140          145         150
Arg Ala Leu Val Arg Gly Pro Ser Ser Leu Met Lys Lys Ala Glu
155          160         165
Leu Ser Glu Ala Gln Val Leu Asp Ala Ala Met Glu Glu Ser Ser
170          175         180
Thr Ser Leu Ala Pro Thr Met Phe Phe Leu Thr Thr Phe Glu Ala
185          190         195
Ala Pro Ala Thr Glu Glu Ser Leu Ile Leu Pro Val Thr Ser Leu
200          205         210
Arg Pro Gln Gln Ala Gln Pro Arg Ser Asp Gly Glu Val Met Pro
215          220         225
Thr Leu Asp Met Ala Leu Phe Asp Trp Thr Asp Tyr Glu Asp Leu
230          235         240
Lys Pro Asp Gly Trp Pro Ser Ala Lys Lys Lys Glu Lys His Arg
245          250         255
Gly Lys Leu Ser Ser Asp Gly Asn Glu Thr Ser Pro Ala Glu Gly
260          265         270
Glu Pro Cys Asp His His Gln Asp Cys Leu Pro Gly Thr Cys Cys
275          280         285
Asp Leu Arg Glu His Leu Cys Thr Pro His Asn Arg Gly Leu Asn
290          295         300
Asn Lys Cys Phe Asp Asp Cys Met Cys Val Glu Gly Leu Arg Cys
305          310         315
Tyr Ala Lys Phe His Arg Asn Arg Arg Val Thr Arg Arg Lys Gly
320          325         330
Arg Cys Val Glu Pro Glu Thr Ala Asn Gly Asp Gln Gly Ser Phe
335          340         345
Ile Asn Val

```

-continued

<210> SEQ ID NO 75

<211> LENGTH: 1868

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 75

cagaagggca aaaacattga ctgcctcaag gtctcaagca ccagtcttca	50
ccgcggaaag catgttgtagg ctgttccaat cgctcctggt tgtcttctgc	100
tttgccccag ggaatgtagt ttcacaaagc agcttaaccc cattgatggt	150
gaacgggatt ctgggggagt cagtaactct tcccctggag tttcctgcag	200
gagagaaggt caacttcata acttggcttt tcaatgaaac atctcttgcc	250
ttcatagtag cccatgaaac caaaagtcca gaaatccacg tgactaatcc	300
gaaacaggga aagcgactga acttcaccca gtctactccc ctgcaactca	350
gcaacctgaa gatggaagac acaggctctt acagagccca gatatccaca	400
aagacctctg caaagctgtc cagttacact ctgaggatat taagacaact	450
gaggaacata caagttacca atcacagtca gctatttcag aatatgacct	500
gtgagctcca tctgacttgc tctgtggagg atgcagatga caatgtctca	550
ttcagatggg aggccttggg aaacacactt tcaagtcagc caaacctcac	600
tgtctcctgg gaccccagga tttccagtga acaggactac acctgcatag	650
cagagaatgc tgtcagtaat ttatccttct ctgtctctgc ccagaagctt	700
tgcgaaagat ttaaaattca atatacagat accaaaatga ttctgtttat	750
ggtttctggg atatgcatag tcttcgggtt catcatactg ctgttacttg	800
ttttgaggaa aagaagagat tccctatctt tgtctactca gcgaacacag	850
ggccccgcag agtccgcaag gaacctagag tatgtttcag tgtctccaac	900
gaacaacact gtgtatgctt cagtcactca ttcaaacagg gaaacagaaa	950
tctggacacc tagagaaaat gatactatca caatttactc cacaattaat	1000
cattccaaag agagtaaacc cactttttcc agggcaactg cccttgacaa	1050
tgctgtgtaa gttgctgaaa ggcctcagag gaattcggga atgacacgtc	1100
ttctgatccc atgagacaga acaaagaaca ggaagcttgg ttctgttgt	1150
tcctggcaac agaatttgaa tatctaggat aggatgatca cctccagtcc	1200
ttcggactta aacctgccta cctgagtcaa acacctaagg ataacatcat	1250
ttccagcatg tggttcaaat aatattttcc aatccacttc aggccaaaac	1300
atgctaaaga taacacacca gcacattgac tctctctttg ataactaagc	1350
aaatggaatt atggttgaca gagagtttat gatccagaag acaaccactt	1400
ctctcctttt agaaagcagc aggattgact tattgagaaa taatgcagtg	1450
tggttggttac atgtgtagtc tctggagttg gatgggcca tcctgataca	1500
agttgagcat cccttgtctg aaatgcttgg gattagaaat gtttcagatt	1550
tcaatttttt ttcatgtttt ggaatatttg cattatatat agcggttgag	1600
tatccaaatc caaaaatcca aaattcaaaa tgctccaata agcatttccc	1650
ttgagtttca ttgatgtoga tgcagtgtct aaaatctcag attttgagac	1700

-continued

aatttgata ttgattttt ggatttgga tgctcaactt gtacaatgtt 1750
tattagacac atctcctggg acatactgcc taaccttttg gagccttagt 1800
ctcccagact gaaaaaggaa gaggatggta ttacatcagc tccattgttt 1850
gagccaagaa tctaagtc 1868

<210> SEQ ID NO 76
<211> LENGTH: 332
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 76

Met Leu Trp Leu Phe Gln Ser Leu Leu Phe Val Phe Cys Phe Gly
1 5 10 15
Pro Gly Asn Val Val Ser Gln Ser Ser Leu Thr Pro Leu Met Val
20 25 30
Asn Gly Ile Leu Gly Glu Ser Val Thr Leu Pro Leu Glu Phe Pro
35 40 45
Ala Gly Glu Lys Val Asn Phe Ile Thr Trp Leu Phe Asn Glu Thr
50 55 60
Ser Leu Ala Phe Ile Val Pro His Glu Thr Lys Ser Pro Glu Ile
65 70 75
His Val Thr Asn Pro Lys Gln Gly Lys Arg Leu Asn Phe Thr Gln
80 85 90
Ser Tyr Ser Leu Gln Leu Ser Asn Leu Lys Met Glu Asp Thr Gly
95 100 105
Ser Tyr Arg Ala Gln Ile Ser Thr Lys Thr Ser Ala Lys Leu Ser
110 115 120
Ser Tyr Thr Leu Arg Ile Leu Arg Gln Leu Arg Asn Ile Gln Val
125 130 135
Thr Asn His Ser Gln Leu Phe Gln Asn Met Thr Cys Glu Leu His
140 145 150
Leu Thr Cys Ser Val Glu Asp Ala Asp Asp Asn Val Ser Phe Arg
155 160 165
Trp Glu Ala Leu Gly Asn Thr Leu Ser Ser Gln Pro Asn Leu Thr
170 175 180
Val Ser Trp Asp Pro Arg Ile Ser Ser Glu Gln Asp Tyr Thr Cys
185 190 195
Ile Ala Glu Asn Ala Val Ser Asn Leu Ser Phe Ser Val Ser Ala
200 205 210
Gln Lys Leu Cys Glu Asp Val Lys Ile Gln Tyr Thr Asp Thr Lys
215 220 225
Met Ile Leu Phe Met Val Ser Gly Ile Cys Ile Val Phe Gly Phe
230 235 240
Ile Ile Leu Leu Leu Leu Val Leu Arg Lys Arg Arg Asp Ser Leu
245 250 255
Ser Leu Ser Thr Gln Arg Thr Gln Gly Pro Ala Glu Ser Ala Arg
260 265 270
Asn Leu Glu Tyr Val Ser Val Ser Pro Thr Asn Asn Thr Val Tyr
275 280 285
Ala Ser Val Thr His Ser Asn Arg Glu Thr Glu Ile Trp Thr Pro
290 295 300

-continued

Arg Glu Asn Asp Thr Ile Thr Ile Tyr Ser Thr Ile Asn His Ser
305 310 315
Lys Glu Ser Lys Pro Thr Phe Ser Arg Ala Thr Ala Leu Asp Asn
320 325 330
Val Val

<210> SEQ ID NO 77
<211> LENGTH: 3073
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 77

gatccctcga cctcgacca cgcgtccgct ctttaatgct ttcttttttaa	50
gagatcacct tctgacttct cacagaagag gttaactatt acctgtggga	100
agtcagaagg tgatctcttt aatgctttct ttttaagaat ttttcaaatt	150
gagactaatt gcagaggttc cagttgacca gcattcatag gaatgaagac	200
aaacacagag atggtgtgtc taagaaactt caaaagggtg agacctcctg	250
actgaagcat attggattta tttaattttt ttcactgtat ttctgtctctc	300
ctacaaggga aagtcatgat tacactaact gagctaaaat gcttagcaga	350
tgcccagtc tcttatcaca tcttaaaacc atggtgggac gtcttctggt	400
attacatcac actgatcatg ctgctgggtg cagtgtctgc cggagctctc	450
cagctgacgc agagcagggt tctgtgctgt cttccatgca aagtggaatt	500
tgacaatcac tgtgccgtgc cttgggacat cctgaaagcc agcatgaaca	550
catcctctaa tcctgggaca ccgcttccgc tccccctccg aattcagaat	600
gacctccacc gacagcagta ctctatatatt gatccgtct gttacgagaa	650
acagctccat tggtttgcaa agtttttccc ctatctggtg ctcttgca	700
cgtctcatctt tgcagcctgc agcaactttt ggcttcaacta cccagttacc	750
agttccaggc tcgagcattt tgtggccatc cttcacaagt gcttcgattc	800
tccatggacc acccgcgccc tttcagaaac agtggctgag cagtcagtga	850
ggcctctgaa actctccaag tccaagattt tgctttcgtc ctcagggtgt	900
tcagctgaca tagattccgg caaacagtca ttgccctacc cacagccagg	950
tttgagtgca gctggtatag aaagcccaac ttccagtggc ctggacaaga	1000
aggagggtga acaggccaaa gccatctttg aaaaagtga aagattccgc	1050
atgcatgtgg agcagaagga catcatttat agagtatatc tgaacagat	1100
aatagtcaaa gtcattttgt ttgtgctcat cataacttat gttccatatt	1150
ttttaaccca catcactctt gaaatcgact gttcagttga tgtgcaggct	1200
tttacaggat ataagcgcta ccagtgtgtc tattccttgg cagaaatctt	1250
taaggtcctg gcttcatttt atgtcatttt gggtatactt tatggctctga	1300
cctcttccta cagcctgtgg tggatgctga ggagttccct gaagcaatat	1350
tcctttgagg cgttaagaga aaaaagcaac tacagtgaca tcctgatgt	1400
caagaatgac ttgaccttca tccttcatct ggctgatcag tatgatcctc	1450
tttattccaa acgcttctcc atattcctat cagaggtcag tgagaacaaa	1500

-continued

ctgaacaga tcaacctcaa taatgaatgg acagttgaga aactgaaaag	1550
taagcttgtg aaaaatgccc aggacaagat agaactgcat ctttttatgc	1600
tcaacggctct tccagacaat gtctttgagt taactgaaat ggaagtgcta	1650
agcctggagc ttatcccaga ggtgaagctg ccctctgcag tctcacagct	1700
ggtcaacctc aaggagcttc gtgtgtacca ttcattctctg gtcgtagacc	1750
atcctgcact ggcctttcta gaggagaatt taaaaatcct ccgcctgaaa	1800
tttactgaaa tgggaaaaat ccacagctgg gtatttcacc tcaagaatct	1850
caaggaactt tatctttcgg gctgtgttct ccctgaacag ttgagtacta	1900
tgcaagtggg gggctttcag gacttaaaaa atctaaggac cctgtacttg	1950
aagagcagcc tctcccgat ccacaaagt gttacagacc tctgccttc	2000
attgcagaaa ctgtcccttg ataagaggg aagcaactg gttgtgtga	2050
acaacttgaa aaagatggtc aatctgaaaa gcctagaact gatcagctgt	2100
gacctggaac gcatcccaca ttccattttc agcctgaata attgcatga	2150
gttagacctt agggaaaata accttaaaac tgtggaagag attagctttc	2200
agcatcttca gaatctttcc tgcttaaaat tgtggcacia taacattgct	2250
tatattcctg cacagattgg ggcattatct aacctagagc agctctcttt	2300
ggaccataat aatattgaga atctgccctt gcagcttttc ctatgacta	2350
aactacatta tttggatcta agctataacc acttgacctt cattccagaa	2400
gaaatccagt atctgagtaa ttgagctac ttgtgtgtga ccaacaacaa	2450
tattgagatg ctaccagatg ggctgtttca gtgcaaaaag ctgcagtggt	2500
tacttttggg gaaaaatagc ttgatgaatt tgtccctca tgtgggtgag	2550
ctgtcaaaac ttactcatct ggagctcatt ggtaattacc tggaaacact	2600
tctcctgaa ctagaaggat gtcagtcctt aaaacggaac tgtctgattg	2650
ttgaggagaa cttgctcaat actcttcctc tccctgtaac agaacgttta	2700
cagacgtgct tagacaaatg ttgacttaaa gaaaagagac ccgtgtttca	2750
aaatcatttt taaaagtatg ctgcggccgg cgtgggtggct catgcctata	2800
atcccagcac tttgggaggc caagatgggc ggattgcttg aggtcaggag	2850
ttcgagacca gtctggccaa cctggtgaaa ccccatctct gctaaaaacta	2900
caaaaaaatt agccaggcgt ggtggcgtgc gcctgtaac ccagctactt	2950
gggaggctga cgcaggggaa ttgcttgaac caggaggtg gaggttgag	3000
tgagccgaga ttgtgccact gtacaccagc ctgggtgaca gagcaagact	3050
cttatctcaa aaaaaaaaaa aaa	3073

<210> SEQ ID NO 78

<211> LENGTH: 802

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 78

Met	Ile	Thr	Leu	Thr	Glu	Leu	Lys	Cys	Leu	Ala	Asp	Ala	Gln	Ser
1				5					10					15

-continued

Ser	Tyr	His	Ile	Leu	Lys	Pro	Trp	Trp	Asp	Val	Phe	Trp	Tyr	Tyr	20	25	30
Ile	Thr	Leu	Ile	Met	Leu	Leu	Val	Ala	Val	Leu	Ala	Gly	Ala	Leu	35	40	45
Gln	Leu	Thr	Gln	Ser	Arg	Val	Leu	Cys	Cys	Leu	Pro	Cys	Lys	Val	50	55	60
Glu	Phe	Asp	Asn	His	Cys	Ala	Val	Pro	Trp	Asp	Ile	Leu	Lys	Ala	65	70	75
Ser	Met	Asn	Thr	Ser	Ser	Asn	Pro	Gly	Thr	Pro	Leu	Pro	Leu	Pro	80	85	90
Leu	Arg	Ile	Gln	Asn	Asp	Leu	His	Arg	Gln	Gln	Tyr	Ser	Tyr	Ile	95	100	105
Asp	Ala	Val	Cys	Tyr	Glu	Lys	Gln	Leu	His	Trp	Phe	Ala	Lys	Phe	110	115	120
Phe	Pro	Tyr	Leu	Val	Leu	Leu	His	Thr	Leu	Ile	Phe	Ala	Ala	Cys	125	130	135
Ser	Asn	Phe	Trp	Leu	His	Tyr	Pro	Ser	Thr	Ser	Ser	Arg	Leu	Glu	140	145	150
His	Phe	Val	Ala	Ile	Leu	His	Lys	Cys	Phe	Asp	Ser	Pro	Trp	Thr	155	160	165
Thr	Arg	Ala	Leu	Ser	Glu	Thr	Val	Ala	Glu	Gln	Ser	Val	Arg	Pro	170	175	180
Leu	Lys	Leu	Ser	Lys	Ser	Lys	Ile	Leu	Leu	Ser	Ser	Ser	Gly	Cys	185	190	195
Ser	Ala	Asp	Ile	Asp	Ser	Gly	Lys	Gln	Ser	Leu	Pro	Tyr	Pro	Gln	200	205	210
Pro	Gly	Leu	Glu	Ser	Ala	Gly	Ile	Glu	Ser	Pro	Thr	Ser	Ser	Gly	215	220	225
Leu	Asp	Lys	Lys	Glu	Gly	Glu	Gln	Ala	Lys	Ala	Ile	Phe	Glu	Lys	230	235	240
Val	Lys	Arg	Phe	Arg	Met	His	Val	Glu	Gln	Lys	Asp	Ile	Ile	Tyr	245	250	255
Arg	Val	Tyr	Leu	Lys	Gln	Ile	Ile	Val	Lys	Val	Ile	Leu	Phe	Val	260	265	270
Leu	Ile	Ile	Thr	Tyr	Val	Pro	Tyr	Phe	Leu	Thr	His	Ile	Thr	Leu	275	280	285
Glu	Ile	Asp	Cys	Ser	Val	Asp	Val	Gln	Ala	Phe	Thr	Gly	Tyr	Lys	290	295	300
Arg	Tyr	Gln	Cys	Val	Tyr	Ser	Leu	Ala	Glu	Ile	Phe	Lys	Val	Leu	305	310	315
Ala	Ser	Phe	Tyr	Val	Ile	Leu	Val	Ile	Leu	Tyr	Gly	Leu	Thr	Ser	320	325	330
Ser	Tyr	Ser	Leu	Trp	Trp	Met	Leu	Arg	Ser	Ser	Leu	Lys	Gln	Tyr	335	340	345
Ser	Phe	Glu	Ala	Leu	Arg	Glu	Lys	Ser	Asn	Tyr	Ser	Asp	Ile	Pro	350	355	360
Asp	Val	Lys	Asn	Asp	Phe	Ala	Phe	Ile	Leu	His	Leu	Ala	Asp	Gln	365	370	375
Tyr	Asp	Pro	Leu	Tyr	Ser	Lys	Arg	Phe	Ser	Ile	Phe	Leu	Ser	Glu	380	385	390
Val	Ser	Glu	Asn	Lys	Leu	Lys	Gln	Ile	Asn	Leu	Asn	Asn	Glu	Trp			

-continued

395					400					405				
Thr	Val	Glu	Lys	Leu	Lys	Ser	Lys	Leu	Val	Lys	Asn	Ala	Gln	Asp
				410					415					420
Lys	Ile	Glu	Leu	His	Leu	Phe	Met	Leu	Asn	Gly	Leu	Pro	Asp	Asn
				425					430					435
Val	Phe	Glu	Leu	Thr	Glu	Met	Glu	Val	Leu	Ser	Leu	Glu	Leu	Ile
				440					445					450
Pro	Glu	Val	Lys	Leu	Pro	Ser	Ala	Val	Ser	Gln	Leu	Val	Asn	Leu
				455					460					465
Lys	Glu	Leu	Arg	Val	Tyr	His	Ser	Ser	Leu	Val	Val	Asp	His	Pro
				470					475					480
Ala	Leu	Ala	Phe	Leu	Glu	Glu	Asn	Leu	Lys	Ile	Leu	Arg	Leu	Lys
				485					490					495
Phe	Thr	Glu	Met	Gly	Lys	Ile	Pro	Arg	Trp	Val	Phe	His	Leu	Lys
				500					505					510
Asn	Leu	Lys	Glu	Leu	Tyr	Leu	Ser	Gly	Cys	Val	Leu	Pro	Glu	Gln
				515					520					525
Leu	Ser	Thr	Met	Gln	Leu	Glu	Gly	Phe	Gln	Asp	Leu	Lys	Asn	Leu
				530					535					540
Arg	Thr	Leu	Tyr	Leu	Lys	Ser	Ser	Leu	Ser	Arg	Ile	Pro	Gln	Val
				545					550					555
Val	Thr	Asp	Leu	Leu	Pro	Ser	Leu	Gln	Lys	Leu	Ser	Leu	Asp	Asn
				560					565					570
Glu	Gly	Ser	Lys	Leu	Val	Val	Leu	Asn	Asn	Leu	Lys	Lys	Met	Val
				575					580					585
Asn	Leu	Lys	Ser	Leu	Glu	Leu	Ile	Ser	Cys	Asp	Leu	Glu	Arg	Ile
				590					595					600
Pro	His	Ser	Ile	Phe	Ser	Leu	Asn	Asn	Leu	His	Glu	Leu	Asp	Leu
				605					610					615
Arg	Glu	Asn	Asn	Leu	Lys	Thr	Val	Glu	Glu	Ile	Ser	Phe	Gln	His
				620					625					630
Leu	Gln	Asn	Leu	Ser	Cys	Leu	Lys	Leu	Trp	His	Asn	Asn	Ile	Ala
				635					640					645
Tyr	Ile	Pro	Ala	Gln	Ile	Gly	Ala	Leu	Ser	Asn	Leu	Glu	Gln	Leu
				650					655					660
Ser	Leu	Asp	His	Asn	Asn	Ile	Glu	Asn	Leu	Pro	Leu	Gln	Leu	Phe
				665					670					675
Leu	Cys	Thr	Lys	Leu	His	Tyr	Leu	Asp	Leu	Ser	Tyr	Asn	His	Leu
				680					685					690
Thr	Phe	Ile	Pro	Glu	Glu	Ile	Gln	Tyr	Leu	Ser	Asn	Leu	Gln	Tyr
				695					700					705
Phe	Ala	Val	Thr	Asn	Asn	Asn	Ile	Glu	Met	Leu	Pro	Asp	Gly	Leu
				710					715					720
Phe	Gln	Cys	Lys	Lys	Leu	Gln	Cys	Leu	Leu	Leu	Gly	Lys	Asn	Ser
				725					730					735
Leu	Met	Asn	Leu	Ser	Pro	His	Val	Gly	Glu	Leu	Ser	Asn	Leu	Thr
				740					745					750
His	Leu	Glu	Leu	Ile	Gly	Asn	Tyr	Leu	Glu	Thr	Leu	Pro	Pro	Glu
				755					760					765
Leu	Glu	Gly	Cys	Gln	Ser	Leu	Lys	Arg	Asn	Cys	Leu	Ile	Val	Glu
				770					775					780

-continued

Glu Asn Leu Leu Asn Thr Leu Pro Leu Pro Val Thr Glu Arg Leu
785 790 795
Gln Thr Cys Leu Asp Lys Cys
800

<210> SEQ ID NO 79
<211> LENGTH: 1504
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 79

cggaacgcgtg	ggccgcgcctc	cctcacggcc	cctcggcggc	gcccgtcgga	50
tccggcctct	ctctgcgccc	cggggcgcgc	cacctccccg	ccggaggtgt	100
ccacgcgtcc	ggcgtcccat	ccgtccgtcc	ctcctggggc	cgccgctgac	150
catgcccagc	ggctgcgcgt	gcctgcattc	cgtgtgcctg	ttgtgcattc	200
tgggggctcc	cggtcagcct	gtccgagccg	atgactgcag	ctcccactgt	250
gacctggccc	acggctgctg	tgcacctgac	ggctcctgca	ggtgtgaccc	300
gggctgggag	gggctgcact	gtgagcgctg	tgtgaggatg	cctggctgcc	350
agcacgggtac	ctgccaccag	ccatggcagt	gcattctgca	cagtggctgg	400
gcaggaagt	tctgtgacaa	agatgaacat	atctgtacca	cgagtcctcc	450
ctgcagaaat	ggaggccagt	gcattgtatg	cgggggcggt	gagtagcatt	500
gtgtgtgctt	accaggcttc	catggcgctg	actgcgagcg	caaggctgga	550
ccctgtgaac	aggcaggctc	cccatgccgc	aatggcgggc	agtgcaggga	600
cgaccagggc	tttgctctca	acttcacgtg	ccgctgcttg	gtgggctttg	650
tgggtgcccc	ctgtgaggta	aatgtggatg	actgcctgat	gcggccttgt	700
gctaacgggtg	ccacctgcct	tgacggcata	aaccgcttct	cctgcctctg	750
tcctgagggc	tttgctggac	gctttcgcac	catcaacctg	gatgactgtg	800
ccagccgccc	atgccagaga	ggggcccgcct	gtcgggaccg	tgtccacgac	850
ttcgactgcc	tctgccccag	tggctatggt	ggcaagacct	gtgagcttgt	900
cttacctgtc	ccagaccccc	caaccacagt	ggacaccctc	ctagggccca	950
cctcagctgt	agtggtacct	gtacggggc	cagcccccca	cagcgagggg	1000
gctggtctgc	tgcggatctc	agtgaaggag	gtggtgcgga	ggcaagaggc	1050
tgggctaggt	gagcctagct	tgggtggcct	ggtggtgttt	ggggccctca	1100
ctgctgccct	ggttctggct	actgtgttgc	tgaccctgag	ggcctggcgc	1150
cggggtgtct	gccccctgg	accctgttgc	taccctgccc	cacactatgc	1200
tccacgctgc	caggaccagg	agtgtcaggt	tagcatgctg	ccagcagggc	1250
ttccctctgc	acgtgacttg	ccccctgagc	ctggaaagac	cacagcactg	1300
tgatggaggt	gggggctttc	tgccccctt	cctcacctct	tccacccctc	1350
agactggagt	ggtccgttct	caccaccctt	cagcttgggt	acacacacag	1400
aggagacctc	agcctcacac	cagaaatatt	atTTTTTTaa	tacacagaat	1450
gtaagatgga	atTTTatcaa	ataaaactat	gaaaatgcaa	aaaaaaaaaa	1500
aaaa					1504

-continued

<210> SEQ ID NO 80

<211> LENGTH: 383

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 80

Met	Pro	Ser	Gly	Cys	Arg	Cys	Leu	His	Leu	Val	Cys	Leu	Leu	Cys
1				5					10					15
Ile	Leu	Gly	Ala	Pro	Gly	Gln	Pro	Val	Arg	Ala	Asp	Asp	Cys	Ser
				20					25					30
Ser	His	Cys	Asp	Leu	Ala	His	Gly	Cys	Cys	Ala	Pro	Asp	Gly	Ser
				35					40					45
Cys	Arg	Cys	Asp	Pro	Gly	Trp	Glu	Gly	Leu	His	Cys	Glu	Arg	Cys
				50					55					60
Val	Arg	Met	Pro	Gly	Cys	Gln	His	Gly	Thr	Cys	His	Gln	Pro	Trp
				65					70					75
Gln	Cys	Ile	Cys	His	Ser	Gly	Trp	Ala	Gly	Lys	Phe	Cys	Asp	Lys
				80					85					90
Asp	Glu	His	Ile	Cys	Thr	Thr	Gln	Ser	Pro	Cys	Gln	Asn	Gly	Gly
				95					100					105
Gln	Cys	Met	Tyr	Asp	Gly	Gly	Gly	Glu	Tyr	His	Cys	Val	Cys	Leu
				110					115					120
Pro	Gly	Phe	His	Gly	Arg	Asp	Cys	Glu	Arg	Lys	Ala	Gly	Pro	Cys
				125					130					135
Glu	Gln	Ala	Gly	Ser	Pro	Cys	Arg	Asn	Gly	Gly	Gln	Cys	Gln	Asp
				140					145					150
Asp	Gln	Gly	Phe	Ala	Leu	Asn	Phe	Thr	Cys	Arg	Cys	Leu	Val	Gly
				155					160					165
Phe	Val	Gly	Ala	Arg	Cys	Glu	Val	Asn	Val	Asp	Asp	Cys	Leu	Met
				170					175					180
Arg	Pro	Cys	Ala	Asn	Gly	Ala	Thr	Cys	Leu	Asp	Gly	Ile	Asn	Arg
				185					190					195
Phe	Ser	Cys	Leu	Cys	Pro	Glu	Gly	Phe	Ala	Gly	Arg	Phe	Cys	Thr
				200					205					210
Ile	Asn	Leu	Asp	Asp	Cys	Ala	Ser	Arg	Pro	Cys	Gln	Arg	Gly	Ala
				215					220					225
Arg	Cys	Arg	Asp	Arg	Val	His	Asp	Phe	Asp	Cys	Leu	Cys	Pro	Ser
				230					235					240
Gly	Tyr	Gly	Gly	Lys	Thr	Cys	Glu	Leu	Val	Leu	Pro	Val	Pro	Asp
				245					250					255
Pro	Pro	Thr	Thr	Val	Asp	Thr	Pro	Leu	Gly	Pro	Thr	Ser	Ala	Val
				260					265					270
Val	Val	Pro	Ala	Thr	Gly	Pro	Ala	Pro	His	Ser	Ala	Gly	Ala	Gly
				275					280					285
Leu	Leu	Arg	Ile	Ser	Val	Lys	Glu	Val	Val	Arg	Arg	Gln	Glu	Ala
				290					295					300
Gly	Leu	Gly	Glu	Pro	Ser	Leu	Val	Ala	Leu	Val	Val	Phe	Gly	Ala
				305					310					315
Leu	Thr	Ala	Ala	Leu	Val	Leu	Ala	Thr	Val	Leu	Leu	Thr	Leu	Arg
				320					325					330
Ala	Trp	Arg	Arg	Gly	Val	Cys	Pro	Pro	Gly	Pro	Cys	Cys	Tyr	Pro

-continued

														335	340	345
Ala	Pro	His	Tyr	Ala	Pro	Ala	Cys	Gln	Asp	Gln	Glu	Cys	Gln	Val		
				350					355					360		
Ser	Met	Leu	Pro	Ala	Gly	Leu	Pro	Leu	Pro	Arg	Asp	Leu	Pro	Pro		
				365					370					375		
Glu	Pro	Gly	Lys	Thr	Thr	Ala	Leu									
				380												
<210> SEQ ID NO 81																
<211> LENGTH: 1034																
<212> TYPE: DNA																
<213> ORGANISM: Homo Sapien																
<400> SEQUENCE: 81																
gtttgttgct caaaccgagt tctggagaac gccatcagct cgctgcttaa															50	
aattaaacca caggttccat tatgggtcga cttgatggga aagtcacat															100	
cctgacggcc gctgctcagg ggattggcca agcagctgcc ttagcttttg															150	
caagagaagg tgccaaagtc atagccacag acattaatga gtccaaactt															200	
caggaactgg aaaagtaccc gggatttcaa actcgtgtcc ttgatgtcac															250	
aaagaagaaa caaattgatc agtttgccag tgaagttgag agacttgatg															300	
ttctctttaa tgttgctggg ttgtccatc atggaactgt cctggattgt															350	
gaggagaaag actgggactt ctogatgaat ctcaatgtgc gcagcatgta															400	
cctgatgac aaggcattcc ttctaaaaat gcttgctcag aaatctggca															450	
atattatcaa catgtcttct gtggcttcca gcgtcaaagg agttgtgaac															500	
agatgtgtgt acagcacaac caaggcagcc gtgattggcc tcacaaaatc															550	
tctggctgca gatttcatcc agcagggcat caggtgcaac tgtgtgtgcc															600	
caggaacagt tgatacgcca tctctacaag aaagaataca agccagagga															650	
aatcctgaag aggcacggaa tgatttcctg aagagacaaa agacgggaag															700	
attcgcaact gcagaagaaa tagccatgct ctgcgtgtat ttggcttctg															750	
atgaatctgc ttatgtaact ggtaaccctg tcatcattga tggaggctgg															800	
agcttgtgat tttaggatct ccatgggtgg aaggaaggca ggccttcct															850	
atccacagtg aacctgggta cgaagaaaac tcaccaatca tctccttcct															900	
gttaatcaca tgtaaatgaa aataagctct ttttaatgat gtcactgttt															950	
gcaagagtct gattctttaa gtatattaat ctctttgtaa tctcttctga															1000	
aatcattgta aagaaataaa aatattgaac tcat															1034	
<210> SEQ ID NO 82																
<211> LENGTH: 245																
<212> TYPE: PRT																
<213> ORGANISM: Homo Sapien																
<400> SEQUENCE: 82																
Met	Gly	Arg	Leu	Asp	Gly	Lys	Val	Ile	Ile	Leu	Thr	Ala	Ala	Ala		
1				5					10					15		
Gln	Gly	Ile	Gly	Gln	Ala	Ala	Ala	Leu	Ala	Phe	Ala	Arg	Glu	Gly		
				20					25					30		
Ala	Lys	Val	Ile	Ala	Thr	Asp	Ile	Asn	Glu	Ser	Lys	Leu	Gln	Glu		

-continued

	35		40		45
Leu Glu Lys Tyr Pro Gly Ile Gln Thr Arg Val Leu Asp Val Thr	50		55		60
Lys Lys Lys Gln Ile Asp Gln Phe Ala Ser Glu Val Glu Arg Leu	65		70		75
Asp Val Leu Phe Asn Val Ala Gly Phe Val His His Gly Thr Val	80		85		90
Leu Asp Cys Glu Glu Lys Asp Trp Asp Phe Ser Met Asn Leu Asn	95		100		105
Val Arg Ser Met Tyr Leu Met Ile Lys Ala Phe Leu Pro Lys Met	110		115		120
Leu Ala Gln Lys Ser Gly Asn Ile Ile Asn Met Ser Ser Val Ala	125		130		135
Ser Ser Val Lys Gly Val Val Asn Arg Cys Val Tyr Ser Thr Thr	140		145		150
Lys Ala Ala Val Ile Gly Leu Thr Lys Ser Leu Ala Ala Asp Phe	155		160		165
Ile Gln Gln Gly Ile Arg Cys Asn Cys Val Cys Pro Gly Thr Val	170		175		180
Asp Thr Pro Ser Leu Gln Glu Arg Ile Gln Ala Arg Gly Asn Pro	185		190		195
Glu Glu Ala Arg Asn Asp Phe Leu Lys Arg Gln Lys Thr Gly Arg	200		205		210
Phe Ala Thr Ala Glu Glu Ile Ala Met Leu Cys Val Tyr Leu Ala	215		220		225
Ser Asp Glu Ser Ala Tyr Val Thr Gly Asn Pro Val Ile Ile Asp	230		235		240
Gly Gly Trp Ser Leu	245				

<210> SEQ ID NO 83
<211> LENGTH: 1961
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 83

gggcggcggc ggcagcgggt ggaggttgta ggaccggcga ggaataggaa	50
tcatggcggc tgcgctgttc gtgctgctgg gattcgcgct gctgggcacc	100
cacggagcct ccggggctgc cggcttcgtc caggcgccgc tgtcccagca	150
gaggtgggtg gggggcagtg tggagctgca ctgcgaggcc gtgggcagcc	200
cggtgcccga gatccagtgg tggtttgaag ggcagggtcc caacgacacc	250
tgctcccagc tctgggacgg cgcccggctg gaccgcgtcc acatccacgc	300
cacctaccac cagcacgcgg ccagcaccat ctccatcgac acgctcgtgg	350
aggaggacac gggcacttac gagtgccggg ccagcaacga cccggatcgc	400
aaccacctga cccgggcgcc cagggtcaag tgggtccgcg cccaggcagt	450
cgtgctagtc ctggaaccgc gcacagtctt cactaccgta gaagaccttg	500
gtccaagat actcctcacc tgctccttga atgacagcgc cacagaggtc	550
acagggcacc gctggctgaa ggggggcgtg gtgctgaagg aggacgcgct	600

-continued

gccccggccag aaaacggagt tcaaggtgga ctccgacgac cagtggggag	650
agtactcctg cgtcttcctc cccgagccca tgggcacggc caacatccag	700
ctccacgggc ctcccagagt gaaggctgtg aagtcgtcag aacacatcaa	750
cgagggggag acggccatgc tggctgcaa gtcagagtcc gtgccacctg	800
tcactgactg ggcctggtac aagatcactg actctgagga caaggccctc	850
atgaacggct ccgagagcag gttcttcgtg agttcctcgc agggccggtc	900
agagctacac attgagaacc tgaacatgga ggccgacccc ggccagtacc	950
ggtgcaacgg caccagctcc aagggtccg accaggccat catcacgctc	1000
cgcgtgcgca gccacctggc cgcctctctg cccttccttg gcatcgtggc	1050
tgaggtgctg gtgctggtca ccatcatctt catctacgag aagcgccgga	1100
agcccagagga cgtcctggat gatgacgacg ccggtctctg acccctgaag	1150
agcagcgggc agcaccagaa tgacaaaggc aagaacgtcc gccagaggaa	1200
ctcttcctga ggcaggtggc ccgaggacgc tcctgtctcc acgtctgcgc	1250
cgcgcgggga gtccactccc agtgcttgca agattccaag ttctcacctc	1300
ttaaagaaaa cccaccccggt agattcccat catacacttc cttctttttt	1350
aaaaagttg ggttttctcc attcaggatt ctgttcctta ggtttttttc	1400
cttctgaagt gtttcacgag agcccgggag ctgctgcctt gcggccccgt	1450
ctgtggcttt cagcctctgg gtctgagtca tggccgggtg ggcggcacag	1500
cttctccac tggccggagt cagtgccagg tccttgccct ttgtggaag	1550
tcacaggcca cagaggggc ccggtgtcct gcctgtctga agccaatgct	1600
gtctggttgc gccatttttg tgcttttatg ttttaattta tgagggccac	1650
gggtctgtgt tcgactcagc ctcagggacg actctgacct cttggccaca	1700
gaggactcac ttgcccacac cgaggcgac cccgtcacag cctcaagtca	1750
ctcccaagcc ccctccttgt ctgtgcatcc gggggcagct ctggaggggg	1800
tttgctgggg aactggcgcc atcgccggga ctccagaacc gcagaagcct	1850
ccccagctca ccctggagg acggccggct ctctatagca ccagggctca	1900
cgtgggaacc cccctccac ccaccgccac aataaagatc gccccacct	1950
ccacccaaaa a	1961

<210> SEQ ID NO 84

<211> LENGTH: 385

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 84

Met	Ala	Ala	Ala	Leu	Phe	Val	Leu	Leu	Gly	Phe	Ala	Leu	Leu	Gly
1				5					10					15
Thr	His	Gly	Ala	Ser	Gly	Ala	Ala	Gly	Phe	Val	Gln	Ala	Pro	Leu
				20					25					30
Ser	Gln	Gln	Arg	Trp	Val	Gly	Gly	Ser	Val	Glu	Leu	His	Cys	Glu
				35					40					45
Ala	Val	Gly	Ser	Pro	Val	Pro	Glu	Ile	Gln	Trp	Trp	Phe	Glu	Gly
				50					55					60

-continued

Gln	Gly	Pro	Asn	Asp	Thr	Cys	Ser	Gln	Leu	Trp	Asp	Gly	Ala	Arg	65	70	75
Leu	Asp	Arg	Val	His	Ile	His	Ala	Thr	Tyr	His	Gln	His	Ala	Ala	80	85	90
Ser	Thr	Ile	Ser	Ile	Asp	Thr	Leu	Val	Glu	Glu	Asp	Thr	Gly	Thr	95	100	105
Tyr	Glu	Cys	Arg	Ala	Ser	Asn	Asp	Pro	Asp	Arg	Asn	His	Leu	Thr	110	115	120
Arg	Ala	Pro	Arg	Val	Lys	Trp	Val	Arg	Ala	Gln	Ala	Val	Val	Leu	125	130	135
Val	Leu	Glu	Pro	Gly	Thr	Val	Phe	Thr	Thr	Val	Glu	Asp	Leu	Gly	140	145	150
Ser	Lys	Ile	Leu	Leu	Thr	Cys	Ser	Leu	Asn	Asp	Ser	Ala	Thr	Glu	155	160	165
Val	Thr	Gly	His	Arg	Trp	Leu	Lys	Gly	Gly	Val	Val	Leu	Lys	Glu	170	175	180
Asp	Ala	Leu	Pro	Gly	Gln	Lys	Thr	Glu	Phe	Lys	Val	Asp	Ser	Asp	185	190	195
Asp	Gln	Trp	Gly	Glu	Tyr	Ser	Cys	Val	Phe	Leu	Pro	Glu	Pro	Met	200	205	210
Gly	Thr	Ala	Asn	Ile	Gln	Leu	His	Gly	Pro	Pro	Arg	Val	Lys	Ala	215	220	225
Val	Lys	Ser	Ser	Glu	His	Ile	Asn	Glu	Gly	Glu	Thr	Ala	Met	Leu	230	235	240
Val	Cys	Lys	Ser	Glu	Ser	Val	Pro	Pro	Val	Thr	Asp	Trp	Ala	Trp	245	250	255
Tyr	Lys	Ile	Thr	Asp	Ser	Glu	Asp	Lys	Ala	Leu	Met	Asn	Gly	Ser	260	265	270
Glu	Ser	Arg	Phe	Phe	Val	Ser	Ser	Ser	Gln	Gly	Arg	Ser	Glu	Leu	275	280	285
His	Ile	Glu	Asn	Leu	Asn	Met	Glu	Ala	Asp	Pro	Gly	Gln	Tyr	Arg	290	295	300
Cys	Asn	Gly	Thr	Ser	Ser	Lys	Gly	Ser	Asp	Gln	Ala	Ile	Ile	Thr	305	310	315
Leu	Arg	Val	Arg	Ser	His	Leu	Ala	Ala	Leu	Trp	Pro	Phe	Leu	Gly	320	325	330
Ile	Val	Ala	Glu	Val	Leu	Val	Leu	Val	Thr	Ile	Ile	Phe	Ile	Tyr	335	340	345
Glu	Lys	Arg	Arg	Lys	Pro	Glu	Asp	Val	Leu	Asp	Asp	Asp	Asp	Ala	350	355	360
Gly	Ser	Ala	Pro	Leu	Lys	Ser	Ser	Gly	Gln	His	Gln	Asn	Asp	Lys	365	370	375
Gly	Lys	Asn	Val	Arg	Gln	Arg	Asn	Ser	Ser						380	385	

<210> SEQ ID NO 85

<211> LENGTH: 1002

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 85

ggctcgagca aagacatacg aacaggaggagg aaggccgact gaaagaaaga

50

-continued

cggagaagag gagagagaag ccagggccga gcgtgccagc aggcggatgg	100
agggcggcct ggtggaggag gagacgtagt ggcctgggct gagctgggtg	150
ggccgggaga agcgggtgcc tcagagtggg ggtgggggca tgggaggggc	200
aggcattctg ctgctgctgc tggctggggc ggggggtgtg gtggcctgga	250
gacccccaaa gggaaagtgt cccctgcgct gctcctgctc taaagacagc	300
gccctgtgtg agggctcccc ggacctgccc gtcagcttct ctccgaccct	350
gctgtcactc tcactcgtca ggacgggagt caccagctg aaggccggca	400
gttctctgag aattccgtct ctgcacctgc tcctcttcac ctccaactcc	450
ttctccgtga ttgaggacga tgcatttgcg ggcctgtccc acctgcagta	500
cctcttcacg gaggacaatg agattggctc catctctaag aatgccctca	550
gaggacttcg ctcgcttaca cacctaagcc tggccaataa ccatctggag	600
accctcccca gattcctgtt ccgaggcctg gacaccctta ctcacgtgga	650
cctccgcggg aacccgttcc agtgtgactg ccgcgtcctc tggtcctgc	700
agtgatgcc caccgtgaat gccagcgtgg ggaccggcgc ctgtgcgggc	750
cccgcctccc tgagccacat gcagctccac cacctcgacc ccaagacttt	800
caatgcaga gccataggtg gggggccttc ccgatggggt gggaggcggg	850
agatctgggg gaaaggctgc cagggccaaag aggctcgtct cactccctgc	900
cctgccattt cccggagtgg gaagaccctg agcaagcagc actgccttcc	950
tgagccccag ttttctcatc tgtaaagtgg gggaataaaa cagtgatata	1000
gg	1002

<210> SEQ ID NO 86
<211> LENGTH: 261
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 86

Met Gly Gly Ala Gly Ile Leu Leu Leu Leu Ala Gly Ala Gly	
1 5 10 15	
Val Val Val Ala Trp Arg Pro Pro Lys Gly Lys Cys Pro Leu Arg	
20 25 30	
Cys Ser Cys Ser Lys Asp Ser Ala Leu Cys Glu Gly Ser Pro Asp	
35 40 45	
Leu Pro Val Ser Phe Ser Pro Thr Leu Leu Ser Leu Ser Leu Val	
50 55 60	
Arg Thr Gly Val Thr Gln Leu Lys Ala Gly Ser Phe Leu Arg Ile	
65 70 75	
Pro Ser Leu His Leu Leu Leu Phe Thr Ser Asn Ser Phe Ser Val	
80 85 90	
Ile Glu Asp Asp Ala Phe Ala Gly Leu Ser His Leu Gln Tyr Leu	
95 100 105	
Phe Ile Glu Asp Asn Glu Ile Gly Ser Ile Ser Lys Asn Ala Leu	
110 115 120	
Arg Gly Leu Arg Ser Leu Thr His Leu Ser Leu Ala Asn Asn His	
125 130 135	
Leu Glu Thr Leu Pro Arg Phe Leu Phe Arg Gly Leu Asp Thr Leu	

-continued

140									145					150				
Thr	His	Val	Asp	Leu	Arg	Gly	Asn	Pro	Phe	Gln	Cys	Asp	Cys	Arg				
155									160					165				
Val	Leu	Trp	Leu	Leu	Gln	Trp	Met	Pro	Thr	Val	Asn	Ala	Ser	Val				
170									175					180				
Gly	Thr	Gly	Ala	Cys	Ala	Gly	Pro	Ala	Ser	Leu	Ser	His	Met	Gln				
185									190					195				
Leu	His	His	Leu	Asp	Pro	Lys	Thr	Phe	Lys	Cys	Arg	Ala	Ile	Gly				
200									205					210				
Gly	Gly	Leu	Ser	Arg	Trp	Gly	Gly	Arg	Arg	Glu	Ile	Trp	Gly	Lys				
215									220					225				
Gly	Cys	Gln	Gly	Gln	Glu	Ala	Arg	Leu	Thr	Pro	Cys	Pro	Ala	Ile				
230									235					240				
Ser	Arg	Ser	Gly	Lys	Thr	Leu	Ser	Lys	Gln	His	Cys	Leu	Pro	Glu				
245									250					255				
Pro	Gln	Phe	Ser	His	Leu													
260																		

<210> SEQ ID NO 87
<211> LENGTH: 2945
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 87

cggaacgcgtg gggcggcgag agcagctgca gttcgcatct caggcagtac	50
ctagaggagc tgccgggtgcc tcctcagaac atctcctgat cgctacccag	100
gaccaggcac caaggacagg gagtcccagg cgcacacccc ccattctggg	150
tccccaggc ccagaccccc actctgccac aggttgcatc ttgacctggt	200
cctcctgcag aagtggcccc tgtggtcctg ctctgagact cgtccctggg	250
cgccctctca gcccttttct atgactccat ctggatttgg ctggctgtgg	300
ggacgcggtc cgagggcgcg cctggctctc agcgtggtgg cagccagctc	350
tctggccacc atggcaaag ctgagatctg aggggacaag gctctacagc	400
ctcagccagg ggactcagc tgttcagagg tgtgatggag aacaaagcta	450
tgtacctaca caccgtcagc gactgtgaca ccagctccat ctgtgaggat	500
tcctttgatg gcaggagcct gtccaagctg aacctgtgtg aggatgggcc	550
atgtcacaaa cggcgggcaa gcatctgctg taccagctg gggccctgt	600
cgccctgaa gcatgctgtc ctggggctct acctgctggt cttcctgatt	650
cttgtgggca tcttcatctt agcaggggca cgggaccca aaggtgatca	700
gggggatgaa ggaaggaag gcaggcctgg catccctgga ttgcctggac	750
ttcgaggctc gcccggggag agaggtaccc caggattgcc cgggccaag	800
ggcgtatgat ggaagctggg ggccacagga ccaatgggca tgcgtgggtt	850
caaaggtgac cgaggcccaa aaggagagaa aggagagaaa ggagacagag	900
ctggggatgc cagtggcgtg gaggccccga tgatgatccg cctggtgaat	950
ggctcaggtc cgcacgaggg ccgctgggaa gtgtaccacg accggcgtg	1000
gggcaccgtg tgtgacgacg gctgggacaa gaaggacgga gacgtggtgt	1050

-continued

gccgcatgct cggcttccgc ggtgtggagg aggtgtaccg cacagctcga	1100
ttcgggcaag gcactgggag gatctggatg gatgacgttg cctgcaaggg	1150
cacagaggaa accatcttcc gctgcagctt ctccaaatgg ggggtgacaa	1200
actgtggaca tgccgaagat gccagcgtga catgcaacag aactgaaag	1250
tgggcagagc ccaagttcgg ggtcctgcac agagcacctt tgctgcatcc	1300
ctgggggtggg gcacagctcg gggccacctt gaccatgcct cgaccacacc	1350
ccgtccagca ttctcagtcc tcacacctgc atcccaggac cgtggggggc	1400
ggtcgtcatt tccctcttga acatgtgctc cgaagtataa ctctgggacc	1450
tactgcccggt ctctctcttc caccaggttc ctgcatgagg agccctgatc	1500
aactggatca ccactttgcc cagcctctga acaccatgca ccaggcctca	1550
atatcccagt tccctttggc cttttagtta cagggtaatg ctgagaatgt	1600
gtcagagaca agtgcagcag cagcgatggt tggtagtata gatcatttac	1650
tcttcagaca attcccaaac ctccattagt ccaagagttt ctacatcttc	1700
ctccccagca agaggcaacg tcaagtgatg aatttcccc ctttactctg	1750
cctctgtctc ccatttgcta gtttgaggaa gtgacataga ggagaagcca	1800
gctgtagggg caagagggaa atgcaagtca cctgcaggaa tccagctaga	1850
tttgagaag ggaatgaaac taacattgaa tgactacat ggacgctaa	1900
atagtatctt ggggtccaaa ttcattgata cacttagctg cattggtcca	1950
gggcatgtca gtctggatac agccttacct tcaggtagca cttaactggt	2000
ccattcacct agactgcaag taagaagaca aaatgactga gaccgtgtgc	2050
ccacctgaac ttattgtctt tacttggcct gagctaaaag cttgggtgca	2100
ggacctgtgt aactagaaag ttgcctactt cagaacctcc agggcgtgag	2150
tgcaagggtca aacatgactg gcttccaggc cgaccatcaa tgtaggagga	2200
gagctgatgt ggagggtgac atgggggctg cccatgttaa acctgagtcc	2250
agtgtctctg cattgggcag tcacggttaa agccaagtca tgtgtgtctc	2300
agctgtttgg aggtgatgat ttgcatctt ccaagcctct tcagggtgta	2350
atctgtggtc aggaaaacac aagtccaat ggaaccctta ggggggaagg	2400
aaatgaagat tccctataac ctctgggggt ggggagtagg aataaggggc	2450
cttgggcctc cataaatctg caatctgcac cctcctccta gagacaggga	2500
gatcgtgttc tgctttttac atgaggagca gaactgggcc atacacgtgt	2550
tcaagaacta ggggagctac ctggtagcaa gtgagtgcag acccacctca	2600
ccttggggga atctcaaaact catagccctc agatacacga tcacctgtca	2650
tatcagggtga gcactggcct gcttggggag agacctgggc ccctccaggt	2700
gtaggaaacg caacactcct ggctgacaac taagccaata tggccctagg	2750
tcatctctgc ttccaatatg cttgccactc cttaaagtgc ctaatgatga	2800
gaaactctct ttctgaccaa ttgctatgtt tacataacac gcatgtactc	2850
atgcatccct tgccagagcc catatatgta tgcataata aacatagcac	2900
tttttactac atagctcagc acattgcaag gtttgcattt aagtt	2945

-continued

<210> SEQ ID NO 88
<211> LENGTH: 270
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 88

Met	Glu	Asn	Lys	Ala	Met	Tyr	Leu	His	Thr	Val	Ser	Asp	Cys	Asp	
1				5					10					15	
Thr	Ser	Ser	Ile	Cys	Glu	Asp	Ser	Phe	Asp	Gly	Arg	Ser	Leu	Ser	
			20						25					30	
Lys	Leu	Asn	Leu	Cys	Glu	Asp	Gly	Pro	Cys	His	Lys	Arg	Arg	Ala	
			35						40					45	
Ser	Ile	Cys	Cys	Thr	Gln	Leu	Gly	Ser	Leu	Ser	Ala	Leu	Lys	His	
			50						55					60	
Ala	Val	Leu	Gly	Leu	Tyr	Leu	Leu	Val	Phe	Leu	Ile	Leu	Val	Gly	
			65						70					75	
Ile	Phe	Ile	Leu	Ala	Gly	Pro	Pro	Gly	Pro	Lys	Gly	Asp	Gln	Gly	
			80						85					90	
Asp	Glu	Gly	Lys	Glu	Gly	Arg	Pro	Gly	Ile	Pro	Gly	Leu	Pro	Gly	
			95						100					105	
Leu	Arg	Gly	Leu	Pro	Gly	Glu	Arg	Gly	Thr	Pro	Gly	Leu	Pro	Gly	
			110						115					120	
Pro	Lys	Gly	Asp	Asp	Gly	Lys	Leu	Gly	Ala	Thr	Gly	Pro	Met	Gly	
			125						130					135	
Met	Arg	Gly	Phe	Lys	Gly	Asp	Arg	Gly	Pro	Lys	Gly	Glu	Lys	Gly	
			140						145					150	
Glu	Lys	Gly	Asp	Arg	Ala	Gly	Asp	Ala	Ser	Gly	Val	Glu	Ala	Pro	
			155						160					165	
Met	Met	Ile	Arg	Leu	Val	Asn	Gly	Ser	Gly	Pro	His	Glu	Gly	Arg	
			170						175					180	
Val	Glu	Val	Tyr	His	Asp	Arg	Arg	Trp	Gly	Thr	Val	Cys	Asp	Asp	
			185						190					195	
Gly	Trp	Asp	Lys	Lys	Asp	Gly	Asp	Val	Val	Cys	Arg	Met	Leu	Gly	
			200						205					210	
Phe	Arg	Gly	Val	Glu	Glu	Val	Tyr	Arg	Thr	Ala	Arg	Phe	Gly	Gln	
			215						220					225	
Gly	Thr	Gly	Arg	Ile	Trp	Met	Asp	Asp	Val	Ala	Cys	Lys	Gly	Thr	
			230						235					240	
Glu	Glu	Thr	Ile	Phe	Arg	Cys	Ser	Phe	Ser	Lys	Trp	Gly	Val	Thr	
			245						250					255	
Asn	Cys	Gly	His	Ala	Glu	Asp	Ala	Ser	Val	Thr	Cys	Asn	Arg	His	
			260						265					270	

<210> SEQ ID NO 89
<211> LENGTH: 2758
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 89

gtcgccgcga gggacgcaga gagcaccctc cacgcccaga tgcctgcgta	50
gttttttgtga ccagtccgct cctgcctccc cctggggcag tagaggggga	100
gcgatggaga actggactgg caggccctgg ctgtatctgc tgctgcttct	150

-continued

gtccctccct cagctctgct tggatcagga ggtgtgtgcc ggacactctc	200
ttcagacacc tacagaggag ggccagggcc ccgaaggtgt ctggggacct	250
tgggtccagt gggcctcttg ctcccagccc tgcggggtgg ggtgcagcg	300
caggagccgg acatgtcagc tccctacagt gcagctccac ccgagtctgc	350
cctccctcc ccggccccc agacatccag aagccctcct cccccgggc	400
cagggtccca gaccccagac ttctccagaa accctcccct tgtacaggac	450
acagtctcgg ggaaggggtg gcccacttcg aggtcccgt tccacctag	500
ggagagagga gaccagagag attcagcgg ccaggaggtc ccggcttcga	550
gacccatca agccaggaat gttcgttat gggagagtgc cttttgcatt	600
gccactgcac cggaaccgca ggcacctcg gagccaccc agatctgagc	650
tgtccctgat ctcttctaga ggggaagagg ctattccgtc ccctactcca	700
agagcagagc cattctccgc aaacggcagc ccccaaactg agctccctcc	750
cacagaactg tctgtccaca ccccatcccc ccaagcagaa cctetaagcc	800
ctgaaactgc tcagacagag gtggccccc gaaccaggcc tgcccccta	850
cggcatcacc ccagagccca ggcctctggc acagagcccc cctcaccac	900
gcactcctta ggagaagggt gcttcttccg tgcattccct cagccaagaa	950
ggccaagtcc ccagggttg gccagtcccc aggtagcagg gagacgcct	1000
gatccttttc cttcgggtccc tcggggccga ggccagcagg gccaagggcc	1050
ttggggaacg ggggggactc ctacggggc ccgcctggag cctgacctc	1100
agcaccgcgg cgcctggctg cccctgctga gcaacggccc ccatgccagc	1150
tccctctgga gcctctttgc tcccagtagc cctattccaa gatgttctgg	1200
ggagagtga cagctaagag cctgcagcca agcgcctgc cccctgagc	1250
agccagaccc ccgggcccctg cagtgcgcag cctttaactc ccaggaattc	1300
atgggccagc tgtatcagtg ggagcccttc actgaagtcc agggctccca	1350
gcgtgtgaa ctgaactgcc ggcccgttg cttccgttc tatgtccgtc	1400
acactgaaaa ggtccaggat gggaccctgt gtcagcctgg agcccctgac	1450
atctgtgtgg ctggacgctg tctgagcccc ggctgtgatg ggatccttg	1500
ctctggcagg cgtcctgatg gctgtggagt ctgtgggggt gatgattcta	1550
cctgtcgctt tgtttcgggg aacctcactg accgaggggg cccctgggc	1600
tatcagaaga tcttgtggat tccagcggga gccttgccgc tccagattgc	1650
ccagctccgg cctagctcca actacctggc acttcgtggc cctggggggc	1700
ggtccatcat caatgggaac tgggctgtgg atccccctgg gtcctacag	1750
gccggcggga ccgtctttcg atataaccgt cctcccaggg aggagggcaa	1800
aggggagagt ctgtcggctg aaggccccc caccagcct gtggatgtct	1850
atatgatctt tcaggaggaa aaccagcg ttttttatca gtatgtcatc	1900
tcttcacctc ctccaatcct tgagaacccc accccagagc cccctgtccc	1950
ccagcttcag ccggagattc tgagggtgga gcccactt gctccggcac	2000
cccggcagc ccggacccc ggcacctcc agcgtcaggt gcggatcccc	2050

-continued

cagatgcccg ccccgcccca tcccaggaca cccctggggt ctccagctgc	2100
gtactggaaa cgagtgggac actctgcatg ctccagctcc tgcgggaaaag	2150
gtgtctggcg ccccatcttc ctctgcatct cccgtgagtc gggagaggaa	2200
ctggatgaac gcagctgtgc cgcgggtgcc agggccccag cctcccctga	2250
accctgccac ggcaccccat gcccccata ctgggaggct ggcgagtga	2300
catctgcag ccgctcctgt ggccccgga cccagcaccg ccagctgcag	2350
tgcggcagg aatttggggg ggggtgctcc tcggtgcccc cggagcgctg	2400
tggacatctc ccccgccca acatcaccca gtcttgccag ctgcgcctct	2450
gtggccattg ggaagttggc tctccttga gccagtgtc cgtgcggtgc	2500
ggccggggcc agagaagccg gcaggttcgc tgtgttgga acaacggtga	2550
tgaagtgagc gagcaggagt gtgcgtcagg cccccacag cccccagca	2600
gagaggcctg tgacatgggg ccctgtacta ctgcctggtt ccacagcgac	2650
tggagctcca aggtgagccc ggaaccccca gccatcctt gcacctggg	2700
taaccatgcc caggacacct cagcctttcc agcatagctc aataaacttg	2750
tattgatc	2758

<210> SEQ ID NO 90

<211> LENGTH: 877

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 90

Met	Glu	Asn	Trp	Thr	Gly	Arg	Pro	Trp	Leu	Tyr	Leu	Leu	Leu	Leu	1	5	10	15
Leu	Ser	Leu	Pro	Gln	Leu	Cys	Leu	Asp	Gln	Glu	Val	Leu	Ser	Gly	20	25	30	
His	Ser	Leu	Gln	Thr	Pro	Thr	Glu	Glu	Gly	Gln	Gly	Pro	Glu	Gly	35	40	45	
Val	Trp	Gly	Pro	Trp	Val	Gln	Trp	Ala	Ser	Cys	Ser	Gln	Pro	Cys	50	55	60	
Gly	Val	Gly	Val	Gln	Arg	Arg	Ser	Arg	Thr	Cys	Gln	Leu	Pro	Thr	65	70	75	
Val	Gln	Leu	His	Pro	Ser	Leu	Pro	Leu	Pro	Pro	Arg	Pro	Pro	Arg	80	85	90	
His	Pro	Glu	Ala	Leu	Leu	Pro	Arg	Gly	Gln	Gly	Pro	Arg	Pro	Gln	95	100	105	
Thr	Ser	Pro	Glu	Thr	Leu	Pro	Leu	Tyr	Arg	Thr	Gln	Ser	Arg	Gly	110	115	120	
Arg	Gly	Gly	Pro	Leu	Arg	Gly	Pro	Ala	Ser	His	Leu	Gly	Arg	Glu	125	130	135	
Glu	Thr	Gln	Glu	Ile	Arg	Ala	Ala	Arg	Arg	Ser	Arg	Leu	Arg	Asp	140	145	150	
Pro	Ile	Lys	Pro	Gly	Met	Phe	Gly	Tyr	Gly	Arg	Val	Pro	Phe	Ala	155	160	165	
Leu	Pro	Leu	His	Arg	Asn	Arg	Arg	His	Pro	Arg	Ser	Pro	Pro	Arg	170	175	180	
Ser	Glu	Leu	Ser	Leu	Ile	Ser	Ser	Arg	Gly	Glu	Glu	Ala	Ile	Pro	185	190	195	

-continued

Ser	Pro	Thr	Pro	Arg	Ala	Glu	Pro	Phe	Ser	Ala	Asn	Gly	Ser	Pro	200	205	210
Gln	Thr	Glu	Leu	Pro	Pro	Thr	Glu	Leu	Ser	Val	His	Thr	Pro	Ser	215	220	225
Pro	Gln	Ala	Glu	Pro	Leu	Ser	Pro	Glu	Thr	Ala	Gln	Thr	Glu	Val	230	235	240
Ala	Pro	Arg	Thr	Arg	Pro	Ala	Pro	Leu	Arg	His	His	Pro	Arg	Ala	245	250	255
Gln	Ala	Ser	Gly	Thr	Glu	Pro	Pro	Ser	Pro	Thr	His	Ser	Leu	Gly	260	265	270
Glu	Gly	Gly	Phe	Phe	Arg	Ala	Ser	Pro	Gln	Pro	Arg	Arg	Pro	Ser	275	280	285
Ser	Gln	Gly	Trp	Ala	Ser	Pro	Gln	Val	Ala	Gly	Arg	Arg	Pro	Asp	290	295	300
Pro	Phe	Pro	Ser	Val	Pro	Arg	Gly	Arg	Gly	Gln	Gln	Gly	Gln	Gly	305	310	315
Pro	Trp	Gly	Thr	Gly	Gly	Thr	Pro	His	Gly	Pro	Arg	Leu	Glu	Pro	320	325	330
Asp	Pro	Gln	His	Pro	Gly	Ala	Trp	Leu	Pro	Leu	Leu	Ser	Asn	Gly	335	340	345
Pro	His	Ala	Ser	Ser	Leu	Trp	Ser	Leu	Phe	Ala	Pro	Ser	Ser	Pro	350	355	360
Ile	Pro	Arg	Cys	Ser	Gly	Glu	Ser	Glu	Gln	Leu	Arg	Ala	Cys	Ser	365	370	375
Gln	Ala	Pro	Cys	Pro	Pro	Glu	Gln	Pro	Asp	Pro	Arg	Ala	Leu	Gln	380	385	390
Cys	Ala	Ala	Phe	Asn	Ser	Gln	Glu	Phe	Met	Gly	Gln	Leu	Tyr	Gln	395	400	405
Trp	Glu	Pro	Phe	Thr	Glu	Val	Gln	Gly	Ser	Gln	Arg	Cys	Glu	Leu	410	415	420
Asn	Cys	Arg	Pro	Arg	Gly	Phe	Arg	Phe	Tyr	Val	Arg	His	Thr	Glu	425	430	435
Lys	Val	Gln	Asp	Gly	Thr	Leu	Cys	Gln	Pro	Gly	Ala	Pro	Asp	Ile	440	445	450
Cys	Val	Ala	Gly	Arg	Cys	Leu	Ser	Pro	Gly	Cys	Asp	Gly	Ile	Leu	455	460	465
Gly	Ser	Gly	Arg	Arg	Pro	Asp	Gly	Cys	Gly	Val	Cys	Gly	Gly	Asp	470	475	480
Asp	Ser	Thr	Cys	Arg	Leu	Val	Ser	Gly	Asn	Leu	Thr	Asp	Arg	Gly	485	490	495
Gly	Pro	Leu	Gly	Tyr	Gln	Lys	Ile	Leu	Trp	Ile	Pro	Ala	Gly	Ala	500	505	510
Leu	Arg	Leu	Gln	Ile	Ala	Gln	Leu	Arg	Pro	Ser	Ser	Asn	Tyr	Leu	515	520	525
Ala	Leu	Arg	Gly	Pro	Gly	Gly	Arg	Ser	Ile	Ile	Asn	Gly	Asn	Trp	530	535	540
Ala	Val	Asp	Pro	Pro	Gly	Ser	Tyr	Arg	Ala	Gly	Gly	Thr	Val	Phe	545	550	555
Arg	Tyr	Asn	Arg	Pro	Pro	Arg	Glu	Glu	Gly	Lys	Gly	Glu	Ser	Leu	560	565	570

-continued

Ser	Ala	Glu	Gly	Pro	Thr	Thr	Gln	Pro	Val	Asp	Val	Tyr	Met	Ile
				575					580					585
Phe	Gln	Glu	Glu	Asn	Pro	Gly	Val	Phe	Tyr	Gln	Tyr	Val	Ile	Ser
				590					595					600
Ser	Pro	Pro	Pro	Ile	Leu	Glu	Asn	Pro	Thr	Pro	Glu	Pro	Pro	Val
				605					610					615
Pro	Gln	Leu	Gln	Pro	Glu	Ile	Leu	Arg	Val	Glu	Pro	Pro	Leu	Ala
				620					625					630
Pro	Ala	Pro	Arg	Pro	Ala	Arg	Thr	Pro	Gly	Thr	Leu	Gln	Arg	Gln
				635					640					645
Val	Arg	Ile	Pro	Gln	Met	Pro	Ala	Pro	Pro	His	Pro	Arg	Thr	Pro
				650					655					660
Leu	Gly	Ser	Pro	Ala	Ala	Tyr	Trp	Lys	Arg	Val	Gly	His	Ser	Ala
				665					670					675
Cys	Ser	Ala	Ser	Cys	Gly	Lys	Gly	Val	Trp	Arg	Pro	Ile	Phe	Leu
				680					685					690
Cys	Ile	Ser	Arg	Glu	Ser	Gly	Glu	Glu	Leu	Asp	Glu	Arg	Ser	Cys
				695					700					705
Ala	Ala	Gly	Ala	Arg	Pro	Pro	Ala	Ser	Pro	Glu	Pro	Cys	His	Gly
				710					715					720
Thr	Pro	Cys	Pro	Pro	Tyr	Trp	Glu	Ala	Gly	Glu	Trp	Thr	Ser	Cys
				725					730					735
Ser	Arg	Ser	Cys	Gly	Pro	Gly	Thr	Gln	His	Arg	Gln	Leu	Gln	Cys
				740					745					750
Arg	Gln	Glu	Phe	Gly	Gly	Gly	Gly	Ser	Ser	Val	Pro	Pro	Glu	Arg
				755					760					765
Cys	Gly	His	Leu	Pro	Arg	Pro	Asn	Ile	Thr	Gln	Ser	Cys	Gln	Leu
				770					775					780
Arg	Leu	Cys	Gly	His	Trp	Glu	Val	Gly	Ser	Pro	Trp	Ser	Gln	Cys
				785					790					795
Ser	Val	Arg	Cys	Gly	Arg	Gly	Gln	Arg	Ser	Arg	Gln	Val	Arg	Cys
				800					805					810
Val	Gly	Asn	Asn	Gly	Asp	Glu	Val	Ser	Glu	Gln	Glu	Cys	Ala	Ser
				815					820					825
Gly	Pro	Pro	Gln	Pro	Pro	Ser	Arg	Glu	Ala	Cys	Asp	Met	Gly	Pro
				830					835					840
Cys	Thr	Thr	Ala	Trp	Phe	His	Ser	Asp	Trp	Ser	Ser	Lys	Val	Ser
				845					850					855
Pro	Glu	Pro	Pro	Ala	Ile	Ser	Cys	Ile	Leu	Gly	Asn	His	Ala	Gln
				860					865					870
Asp	Thr	Ser	Ala	Phe	Pro	Ala								
				875										

<210> SEQ ID NO 91
<211> LENGTH: 2597
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 91

cgagtatatttt cccaccatct ccagccggaa actgaccaag aactctgag	50
cggatggcat gttcgcgtac gtcttccatg atgagttcgt ggcctcgatg	100
attaagatcc cttcggacac cttcaccatc atccctgact ttgatatacta	150

-continued

ctatgtctat ggttttagca gtggcaactt tgtctacttt ttgacctcc	200
aacctgagat ggtgtctcca ccaggctcca ccaccaagga gcagggtgat	250
acatccaagc tcgtgaggct ttgcaaggag gacacagcct tcaactccta	300
tgtagagggtg ccattgggt gtgagcgag tggggtggag taccgcctgc	350
tgcaggctgc ctacctgtcc aaagcggggg ccgtgcttgg caggaccctt	400
ggagtccatc cagatgatga cctgtctctc accgtcttct ccaagggccca	450
gaagcggaat atgaaatccc tggatgagtc ggccctgtgc atcttcatct	500
tgaagcagat aaatgaccgc attaaggagc ggctgcagtc ttgttacgg	550
ggcaggggca cgctggacct ggctggctc aaggtgaagg acatcccctg	600
cagcagtgcg ctcttaacca ttgacgataa cttctgtggc ctggacatga	650
atgtccccc gggagtgctc gacatggctc gtggaattcc cgtcttcacg	700
gaggacagg accgcatgac gtctgtcatc gcatatgtct acaagaacca	750
ctctctggcc tttgtgggca ccaaaagtgg caagctgaag aaggtgctg	800
gtaccagcct ctgccctacc cttgagctac agacgggacc ccgatccac	850
agagcaacag tgactctgga actcctgttc tccagctgtt catcaaactg	900
agaaaaactt cagagctgtg taggcttatt tagtgtgttg tcagccttgg	950
atattggaaa atggaaacag atgagacaca tctacctccc tgtgaccca	1000
gccatacatc atagctcatg tcctgccacc ccaagtcctt agggaaaaaa	1050
gactttggag aatgtgtctc tgcttagctt ggctaggtag ttggtctctt	1100
ttctctgccc caagcgtccc ctgggtaatt ttggacaatg gagtgtaggc	1150
atgtttgact cttgtggtgt tatcacttgt atatgtcagt gaaactaact	1200
gatttcccca tcggaatata gttatctctt gggcctgata tatggtagga	1250
taaccttatg ctcatctgtc cactcttgca gccaaagtcg ctggccagtg	1300
tgtgtgtgtg tgtgtgtgtg tgtgtgtgtg tgtgtgtatg cttatctgtg	1350
tttaaagggtg tgtgtgcata cacaggcgag agaggatgga gcccaccgta	1400
ctgcagcatc atgtaattaa ctcagtgtc agaaccatcc cagcctctgc	1450
gggaaagaga aaagtaagcc aacagtgcct gatgagctga tcatatgtgc	1500
aaaagctctg ttggcatctg gtccaggaga gcacccaaaa aaagttaatt	1550
ggtgttgtcc agtctccttt ccttaagact atggttaca caaagcgtga	1600
gcagtgtctc ctgcatggcc actatccagc acaattccat aattccccca	1650
tagagccggt ggggaggag aggtgagtgg cgaagggaagt ggaacactt	1700
ggtgtcatgt gctcctatca tttctactag cttactggga aataaagtgt	1750
agtcaagagt gtatgaaggc aagatgtaaa attagcgact ggtgctaatac	1800
tggttacttg aaaacaagtg aaagtgtgt agatttgttc tgttgctaag	1850
aaccaccaca ctaaacctcg tatagttcct ggaggatata caacagtga	1900
attctcttta ggggtgtcca caggttcctg gcctgtggga gggaatgaat	1950
caggagggtc cttgagaacc ttcactctgt tgcttgcaact gaaagtgagt	2000
cccaaagctg gagatttagt gagagcaggc aaccctctg tgtctcactg	2050

-continued

tccatattct ggaggcagag gtttgtaaca ggccatgtgc acctgcatag	2100
ggatgggtaa agcaaggact ttgaaagagt tgaaaagcat tataaacagt	2150
tgttcagaaa tacgtcccag gaggttccatg tgaaactggc tctgtgtgca	2200
ttgaagcatg gctgttggga attctaactg gtccaacact cctgcaaaac	2250
aatgtgtaaa tatttaggaa gaaacttgaa aatagtcaaa tcctttgaac	2300
tggtgacaat tttttaaaga atcaattcta atttgtttca agggtaataa	2350
tcaccaagat acacatttca gcatttattt agtctatcaa aaattggaat	2400
tgatatatac actcatttat aggagaatgg ttaggtagat ttggtatatt	2450
tatgtagtca ttgaaaactt agtttataaa ggccaatcct gtaactgatt	2500
cttgtgtgat aacattcagt gaaaagcat gagacaatta gaaagcatga	2550
tacaatgaat aaaataaaaa ctggaaagag aaccatcaaa atgctaa	2597

<210> SEQ ID NO 92

<211> LENGTH: 280

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 92

Met Phe Ala Tyr Val Phe His Asp Glu Phe Val Ala Ser Met Ile	1 5 10 15
Lys Ile Pro Ser Asp Thr Phe Thr Ile Ile Pro Asp Phe Asp Ile	20 25 30
Tyr Tyr Val Tyr Gly Phe Ser Ser Gly Asn Phe Val Tyr Phe Leu	35 40 45
Thr Leu Gln Pro Glu Met Val Ser Pro Pro Gly Ser Thr Thr Lys	50 55 60
Glu Gln Val Tyr Thr Ser Lys Leu Val Arg Leu Cys Lys Glu Asp	65 70 75
Thr Ala Phe Asn Ser Tyr Val Glu Val Pro Ile Gly Cys Glu Arg	80 85 90
Ser Gly Val Glu Tyr Arg Leu Leu Gln Ala Ala Tyr Leu Ser Lys	95 100 105
Ala Gly Ala Val Leu Gly Arg Thr Leu Gly Val His Pro Asp Asp	110 115 120
Asp Leu Leu Phe Thr Val Phe Ser Lys Gly Gln Lys Arg Lys Met	125 130 135
Lys Ser Leu Asp Glu Ser Ala Leu Cys Ile Phe Ile Leu Lys Gln	140 145 150
Ile Asn Asp Arg Ile Lys Glu Arg Leu Gln Ser Cys Tyr Arg Gly	155 160 165
Glu Gly Thr Leu Asp Leu Ala Trp Leu Lys Val Lys Asp Ile Pro	170 175 180
Cys Ser Ser Ala Leu Leu Thr Ile Asp Asp Asn Phe Cys Gly Leu	185 190 195
Asp Met Asn Ala Pro Leu Gly Val Ser Asp Met Val Arg Gly Ile	200 205 210
Pro Val Phe Thr Glu Asp Arg Asp Arg Met Thr Ser Val Ile Ala	215 220 225

-continued

Tyr	Val	Tyr	Lys	Asn	His	Ser	Leu	Ala	Phe	Val	Gly	Thr	Lys	Ser
				230					235					240
Gly	Lys	Leu	Lys	Lys	Val	Pro	Gly	Thr	Ser	Leu	Cys	Pro	Thr	Leu
				245					250					255
Glu	Leu	Gln	Thr	Gly	Pro	Arg	Ser	His	Arg	Ala	Thr	Val	Thr	Leu
				260					265					270
Glu	Leu	Leu	Phe	Ser	Ser	Cys	Ser	Ser	Asn					
				275					280					

<210> SEQ ID NO 93
<211> LENGTH: 2883
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 93

ccttatcaga caaaggacga gatggaaaat acaagataat ttacagtgga	50
gaagaattag aatgtaacct gaaagatctt agaccagcaa cagattatca	100
tgtaggggtg tatgccatgt acaattccgt aaagggatcc tgctccgagc	150
ctgttagctt caccacccac agctgtgcac cagagtgtcc ttccccccct	200
aagctggcac ataggagcaa aagttcacta accctgcagt ggaaggcacc	250
aattgacaac ggttcaaaaa tcaccaacta ccttttagag tgggatgagg	300
gaaaagaaa tagtggtttc agacagtgtc tcttcgggag ccagaagcac	350
tgcaagtga caaagctttg tccggcaatg gggtagacat tcaggctggc	400
cgctcgaaac gacattggca ccagtgggta tagccaagag gtgggtgtgt	450
acacattagg aaatatccct cagatgcctt ctgcactaag gctgggtcga	500
gctggcatca catgggtcac gttgcagtgg agtaagccag aaggctgttc	550
acccgaggaa gtgatcacct acaccttgga aattcaggag gatgaaaatg	600
ataacctttt ccacccaaaa tacactggag aggatttaac ctgtactgtg	650
aaaaatctca aaagaagcac acagtataaa ttcaggctga ctgcttctaa	700
tacggaagga aaaagctgtc caagcgaagt tcttgtttgt acgacgagtc	750
ctgacaggcc tggacctcct accagaccgc ttgtcaaagg cccagttaca	800
tctcatggct ttagtgtcaa atgggatccc cctaaggaca atgggtggtc	850
agaaatcctc aagtacttgc tagagattac tgatggaaat tctgaagcga	900
atcagtggga agtggcctac agtgggtcgg ctaccgaata caccttcacc	950
cacttgaaac caggcacttt gtacaaactc cgagcatgct gcatcagtac	1000
cggcggacac agccagtgtt ctgaaagtct ccctgttcgc acactaagca	1050
ttgcaccagg tcaatgtcga ccaccgaggg ttttgggtag accaaagcac	1100
aaagaagtcc acttagagtg ggatgttcct gcatcggaag gtggctgtga	1150
ggtctcagag tacagcgtgg agatgacgga gcccgagac gtagcctcgg	1200
aagtgtacca tggcccagag ctggagtgca ccgtcgga cctgcttcct	1250
ggaaccgtgt atcgcttcgg ggtgagggct ctgaatgatg gagggtatgg	1300
tccctattct gatgtctcag aaattaccac tgctgcaggg cctcctggac	1350
aatgcaaaag accttgtatt tcttgtacac ctgatggatg tgtcttagtg	1400

-continued

ggttgggaga gtcctgatag ttctgggtgct gacatctcag agtacaggtt	1450
ggaatgggga gaagatgaag aatccttaga actcatttat catgggacag	1500
acacccgttt tgaataaga gacctgttgc ctgctgcaca gtattgctgt	1550
agactacagg ccttcaatca agcaggggca gggccgtaca gtgaacttgt	1600
cctttgccag acgccagcgt ctgccctga ccccgctctcc actctctgtg	1650
tcctggagga ggagcccctt gatgcctacc ctgattcacc ttctgcgtgc	1700
cttgactact actgggaaga gccgtgcaat aacggatctg aaatccttgc	1750
ttacaccatt gatctaggag acactagcat taccgtgggc aacaccacca	1800
tgcatgttat gaaagatctc cttccagaaa ccacctaccg gatcagaatt	1850
caggctataa atgaaattgg agctggacca tttagttagt tcattaaagc	1900
aaaaactcgg ccattaccac ctttgccctc taggctagaa tgtgctgctg	1950
ctggctccta gagcctgaag ctaaaatggg gagacagtaa ctccaagaca	2000
catgctgctg aggacattgt gtacacacta cagctggagg acagaaacaa	2050
gaggtttatt tcaatctaca gaggaccag ccacacctac aaggtccaga	2100
gactgacgga attcacatgc tactccttca gaatccaggc agcaagcag	2150
gctggagaag ggcccttctc agaaacctat accttcagca caaccaaag	2200
tgtccccccc accatcaaag cacctcgagt aacacagtta gaagtaaatt	2250
catgtgaaat tttatgggag acggtacat caatgaaagg tgaccctgtt	2300
aactacattc tgcagggtatt ggttggaaga gaatctgagt acaaacagg	2350
gtacaaggga gaagaagcca cattccaaat ctgaggcctc cagaccaaca	2400
cagactacag gttccgcgta tgtgcgtgtc gtcgctgttt agacacctct	2450
caggagctaa gcggagcctt cagccctctc gcggtttttg tattacaacg	2500
aagtgaggtc atgcttacag gggacatggg gagcttagat gatcccaaaa	2550
tgaagagcat gatgcctact gatgaacagt ttgcagccat cattgtgctt	2600
ggctttgcaa ctttgtccat tttatttgcc tttatattac agtacttctt	2650
aatgaagtaa acccaacaaa actagaggta tgaattaatg ctacacattt	2700
taatacacac atttattcag atactcccct ttttaaagcc cttttgtttt	2750
ttgatttata tactctgttt tacagattta gctagaaaaa aaatgtcagt	2800
gttttggtgc acctttttga aatgcaaaac taggaaaagg ttaaactgga	2850
ttttttttta aaaaaaaaaa aaaaaaaaaa aaa	2883

<210> SEQ ID NO 94

<211> LENGTH: 847

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 94

Met	Tyr	Asn	Ser	Val	Lys	Gly	Ser	Cys	Ser	Glu	Pro	Val	Ser	Phe
1				5					10					15

Thr	Thr	His	Ser	Cys	Ala	Pro	Glu	Cys	Pro	Phe	Pro	Pro	Lys	Leu
				20					25					30

Ala	His	Arg	Ser	Lys	Ser	Ser	Leu	Thr	Leu	Gln	Trp	Lys	Ala	Pro
				35				40						45

-continued

Ile Asp Asn Gly Ser Lys Ile Thr Asn Tyr Leu Leu Glu Trp Asp	50	55	60
Glu Gly Lys Arg Asn Ser Gly Phe Arg Gln Cys Phe Phe Gly Ser	65	70	75
Gln Lys His Cys Lys Leu Thr Lys Leu Cys Pro Ala Met Gly Tyr	80	85	90
Thr Phe Arg Leu Ala Ala Arg Asn Asp Ile Gly Thr Ser Gly Tyr	95	100	105
Ser Gln Glu Val Cys Tyr Thr Leu Gly Asn Ile Pro Gln Met	110	115	120
Pro Ser Ala Leu Arg Leu Val Arg Ala Gly Ile Thr Trp Val Thr	125	130	135
Leu Gln Trp Ser Lys Pro Glu Gly Cys Ser Pro Glu Glu Val Ile	140	145	150
Thr Tyr Thr Leu Glu Ile Gln Glu Asp Glu Asn Asp Asn Leu Phe	155	160	165
His Pro Lys Tyr Thr Gly Glu Asp Leu Thr Cys Thr Val Lys Asn	170	175	180
Leu Lys Arg Ser Thr Gln Tyr Lys Phe Arg Leu Thr Ala Ser Asn	185	190	195
Thr Glu Gly Lys Ser Cys Pro Ser Glu Val Leu Val Cys Thr Thr	200	205	210
Ser Pro Asp Arg Pro Gly Pro Pro Thr Arg Pro Leu Val Lys Gly	215	220	225
Pro Val Thr Ser His Gly Phe Ser Val Lys Trp Asp Pro Pro Lys	230	235	240
Asp Asn Gly Gly Ser Glu Ile Leu Lys Tyr Leu Leu Glu Ile Thr	245	250	255
Asp Gly Asn Ser Glu Ala Asn Gln Trp Glu Val Ala Tyr Ser Gly	260	265	270
Ser Ala Thr Glu Tyr Thr Phe Thr His Leu Lys Pro Gly Thr Leu	275	280	285
Tyr Lys Leu Arg Ala Cys Cys Ile Ser Thr Gly Gly His Ser Gln	290	295	300
Cys Ser Glu Ser Leu Pro Val Arg Thr Leu Ser Ile Ala Pro Gly	305	310	315
Gln Cys Arg Pro Pro Arg Val Leu Gly Arg Pro Lys His Lys Glu	320	325	330
Val His Leu Glu Trp Asp Val Pro Ala Ser Glu Ser Gly Cys Glu	335	340	345
Val Ser Glu Tyr Ser Val Glu Met Thr Glu Pro Glu Asp Val Ala	350	355	360
Ser Glu Val Tyr His Gly Pro Glu Leu Glu Cys Thr Val Gly Asn	365	370	375
Leu Leu Pro Gly Thr Val Tyr Arg Phe Arg Val Arg Ala Leu Asn	380	385	390
Asp Gly Gly Tyr Gly Pro Tyr Ser Asp Val Ser Glu Ile Thr Thr	395	400	405
Ala Ala Gly Pro Pro Gly Gln Cys Lys Ala Pro Cys Ile Ser Cys	410	415	420

-continued

Thr	Pro	Asp	Gly	Cys	Val	Leu	Val	Gly	Trp	Glu	Ser	Pro	Asp	Ser	
				425					430					435	
Ser	Gly	Ala	Asp	Ile	Ser	Glu	Tyr	Arg	Leu	Glu	Trp	Gly	Glu	Asp	
				440					445					450	
Glu	Glu	Ser	Leu	Glu	Leu	Ile	Tyr	His	Gly	Thr	Asp	Thr	Arg	Phe	
				455					460					465	
Glu	Ile	Arg	Asp	Leu	Leu	Pro	Ala	Ala	Gln	Tyr	Cys	Cys	Arg	Leu	
				470					475					480	
Gln	Ala	Phe	Asn	Gln	Ala	Gly	Ala	Gly	Pro	Tyr	Ser	Glu	Leu	Val	
				485					490					495	
Leu	Cys	Gln	Thr	Pro	Ala	Ser	Ala	Pro	Asp	Pro	Val	Ser	Thr	Leu	
				500					505					510	
Cys	Val	Leu	Glu	Glu	Glu	Pro	Leu	Asp	Ala	Tyr	Pro	Asp	Ser	Pro	
				515					520					525	
Ser	Ala	Cys	Leu	Val	Leu	Asn	Trp	Glu	Glu	Pro	Cys	Asn	Asn	Gly	
				530					535					540	
Ser	Glu	Ile	Leu	Ala	Tyr	Thr	Ile	Asp	Leu	Gly	Asp	Thr	Ser	Ile	
				545					550					555	
Thr	Val	Gly	Asn	Thr	Thr	Met	His	Val	Met	Lys	Asp	Leu	Leu	Pro	
				560					565					570	
Glu	Thr	Thr	Tyr	Arg	Ile	Arg	Ile	Gln	Ala	Ile	Asn	Glu	Ile	Gly	
				575					580					585	
Ala	Gly	Pro	Phe	Ser	Gln	Phe	Ile	Lys	Ala	Lys	Thr	Arg	Pro	Leu	
				590					595					600	
Pro	Pro	Leu	Pro	Pro	Arg	Leu	Glu	Cys	Ala	Ala	Ala	Gly	Pro	Gln	
				605					610					615	
Ser	Leu	Lys	Leu	Lys	Trp	Gly	Asp	Ser	Asn	Ser	Lys	Thr	His	Ala	
				620					625					630	
Ala	Glu	Asp	Ile	Val	Tyr	Thr	Leu	Gln	Leu	Glu	Asp	Arg	Asn	Lys	
				635					640					645	
Arg	Phe	Ile	Ser	Ile	Tyr	Arg	Gly	Pro	Ser	His	Thr	Tyr	Lys	Val	
				650					655					660	
Gln	Arg	Leu	Thr	Glu	Phe	Thr	Cys	Tyr	Ser	Phe	Arg	Ile	Gln	Ala	
				665					670					675	
Ala	Ser	Glu	Ala	Gly	Glu	Gly	Pro	Phe	Ser	Glu	Thr	Tyr	Thr	Phe	
				680					685					690	
Ser	Thr	Thr	Lys	Ser	Val	Pro	Pro	Thr	Ile	Lys	Ala	Pro	Arg	Val	
				695					700					705	
Thr	Gln	Leu	Glu	Val	Asn	Ser	Cys	Glu	Ile	Leu	Trp	Glu	Thr	Val	
				710					715					720	
Pro	Ser	Met	Lys	Gly	Asp	Pro	Val	Asn	Tyr	Ile	Leu	Gln	Val	Leu	
				725					730					735	
Val	Gly	Arg	Glu	Ser	Glu	Tyr	Lys	Gln	Val	Tyr	Lys	Gly	Glu	Glu	
				740					745					750	
Ala	Thr	Phe	Gln	Ile	Ser	Gly	Leu	Gln	Thr	Asn	Thr	Asp	Tyr	Arg	
				755					760					765	
Phe	Arg	Val	Cys	Ala	Cys	Arg	Arg	Cys	Leu	Asp	Thr	Ser	Gln	Glu	
				770					775					780	
Leu	Ser	Gly	Ala	Phe	Ser	Pro	Ser	Ala	Ala	Phe	Val	Leu	Gln	Arg	
				785					790					795	
Ser	Glu	Val	Met	Leu	Thr	Gly	Asp	Met	Gly	Ser	Leu	Asp	Asp	Pro	

-continued

[illegible]

-continued

ccccatcccc aatttccagt gtggcaccct gcctgagacc ggtcccaacg	1500
agaacctgac ggagcgagc ctgcaggacg cgcagcgctt cttcctgatg	1550
agcgaggccg tgcagccggt gacacccgag ccctgtgtca ccagggacag	1600
cgtgcgcttc tcacacctcg tggtagacct ggtgcaggct aaagacacgc	1650
tctaccatgt actctacatt ggcaccgagt cgggcacccat cctgaaggcg	1700
ctgtccacgg cgagccgag cctccacggc tgctacctgg aggagctgca	1750
cgtgctgccc cccgggcgcc gcgagccctt gcgcagcctg cgcacccctg	1800
acagcgcccc cgcgctcttc gtggggctga gagacggcgt cctgcgggtc	1850
ccactggaga ggtgcgccc ctaccgcagc cagggggcat gcctgggggc	1900
ccgggacccg tactgtggct gggacgggaa gcagcaacgt tgcagcacac	1950
tcgaggacag ctccaacatg agcctctgga ccagaacat caccgcctgt	2000
cctgtgcgga atgtgacacg ggatgggggc ttcggcccat ggtcaccatg	2050
gcaacctatg gagcacttgg atggggacaa ctacggctct tgccctgtgtc	2100
gagctcgatc ctgtgattcc cctcgacccc gctgtggggg ccttgactgc	2150
ctggggccag ccatccacat cgccaactgc tccaggaatg ggcgctggac	2200
cccgtgtgca tcgtgggggc tgtgcagcac gtccctgtggc atcggtcttc	2250
aggtccgcga gcgaagtgc agcaaccctg ctccccgcca cgggggcccgc	2300
atcttcgtgg gcaagagccg ggaggaacgg ttctgtaatg agaacacgcc	2350
ttgcccgttg cccatcttct gggcttcctg gggctcctgg agcaagtgca	2400
gcagcaactg tggagggggc atgcagtcgc ggcgtcgggc ctgcgagaac	2450
ggcaactcct gcctgggctg cggcgagtgc aagacgtgca accccgaggg	2500
ctgccccgaa gtgcggcgca acaccccctg gacgccgtgg ctgcccgtga	2550
acgtgacgca gggcggggca cggcaggagc agcggttccg cttcacctgc	2600
cgcgcgcccc ttgcagacct gcacggcctg cagttcggca ggagaaggac	2650
cgagacgagg acctgtcccg cggacggctc cggctcctgc gacaccgacg	2700
ccctggtgga ggtcctcctg cgcagcggga gcacctcccc gcacacggtg	2750
agcgggggct gggccgcctg gggcccgctg tcgtcctgct cccgggactg	2800
cgagctgggc ttccgcgtcc gcaagagaac gtgcactaac ccggagcccc	2850
gcaacggggg cctgccctgc gtgggcgatg ctgccagta ccaggactgc	2900
aacccccagg cttgccctgt tcggggtgct tggctcctgct ggacctcatg	2950
gtctccatgc tcagcttcct gtggtggggg tcaactatcaa cgcacccgtt	3000
cctgcaccag ccccgacccc tcccagggtg aggacatctg tctcgggctg	3050
cacacggagg aggcactatg tgccacacag gcctgccag gctggtcgcc	3100
ctggtctgag tggagtaagt gcaactgacg cggagcccag agccgaagcc	3150
ggcactgtga ggagctcctc ccagggtcca gcgcctgtgc tggaaacagc	3200
agccagagcc gccctgccc ctacagcgag attcccgta tctgcccagc	3250
ctccagcatg gaggaggcca ccgactgtgc aggtaaaaga aaccggacct	3300
acctcatgct gcggtcctcc cagccctcca gcaccccact ccaaagtctg	3350

-continued

gactctttcc acatcctgct ccagacagcc aagctttgtt ggggtcccca 3400
ctgctttgag atgggttcaa tctcatccac ttggtggcca cgggcatctc 3450
ctgcttcttg ggctctgggc tcctgaccct agcagtgtac ctgtcttgcc 3500
agcactgcca gcgtcagtcc caggagtcca cactgggtcca tcctgccacc 3550
cccaaccatt tgcactacaa gggcggaggc accccgaaga atgaaaagta 3600
cacacccatg gaattcaaga ccctgaacaa gaataacttg atccctgatg 3650
acagagccaa cttctaccca ttgcagcaga ccaatgtgta cagcactact 3700
tactacccaa gccccctgaa caaacacagc ttccggcccg aggctcacc 3750
tggaacaacgg tgcttcccca acagctgata ccgccgtcct ggggacttgg 3800
gcttcttgcc ttcataaggc acagagcaga tggagatggg acagtggagc 3850
cagtttggtt ttctccctct gcactaggcc aagaacttg cgccttgctt 3900
gtgggggggtc ccatccggct tcagagagct ctggctggca ttgaccatgg 3950
gggaagggc tggtttcagg ctgacatatg gccgcaggtc cagttcagcc 4000
caggctctctc atggttatct tccaaccac tgtcacgctg acactatgct 4050
gccatgcctg ggctgtggac ctactgggca tttgaggaat tggagaatgg 4100
agatggcaag agggcaggct tttaagtttg ggttgagac aacttcctgt 4150
ggccccaca agctgagtct ggcttctcc agctggcccc aaaaaggcc 4200
tttgctacat cctgattatc tctgaaagta atcaatcaag tggctccagt 4250
agctctggat tttctgccag ggctgggcca ttgtggtgct gccccagtat 4300
gacatgggac caaggccagc gcaggttatc cacctctgcc tggaagtcta 4350
tactctaccc agggcatccc tctggtcaga ggcagtgagt actgggaact 4400
ggaggctgac ctgtgcttag aagtccttta atctgggctg gtacaggcct 4450
cagccttgcc ctcaatgcac gaaagggtgc ccaggagaga ggatcaatgc 4500
cataggaggc agaagtctgg cctctgtgcc tctatggaga ctatcttcca 4550
gttgctgctc aacagagttg ttggctgaga cctgcttggg agtctctgct 4600
ggcccttcat ctgttcagga acacacacac acacacactc acacacgcac 4650
acacaatcac aatttgctac agcaacaaaa aagacattgg gctgtggcat 4700
tattaattaa agatgatatc cagtc 4725

<210> SEQ ID NO 96

<211> LENGTH: 1092

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 96

Met Pro Cys Gly Phe Ser Pro Ser Pro Val Ala His His Leu Val
1 5 10 15
Pro Gly Pro Pro Asp Thr Pro Ala Gln Gln Leu Arg Cys Gly Trp
20 25 30
Thr Val Gly Gly Trp Leu Leu Ser Leu Val Arg Gly Leu Leu Pro
35 40 45
Cys Leu Pro Pro Gly Ala Arg Thr Ala Glu Gly Pro Ile Met Val
50 55 60

-continued

Leu Ala Gly Pro	Leu Ala Val Ser	Leu	Leu	Leu	Pro Ser	Leu Thr	65	70	75
Leu Leu Val Ser	His Leu Ser Ser	Ser	Gln Asp	Val Ser	Ser Glu	80	85	90	
Pro Ser Ser Glu	Gln Gln Leu Cys	Ala	Leu Ser	Lys His	Pro Thr	95	100	105	
Val Ala Phe Glu	Asp Leu Gln Pro	Trp	Val Ser	Asn Phe	Thr Tyr	110	115	120	
Pro Gly Ala Arg	Asp Phe Ser Gln	Leu	Ala Leu	Asp Pro	Ser Gly	125	130	135	
Asn Gln Leu Ile	Val Gly Ala Arg	Asn	Tyr Leu	Phe Arg	Leu Ser	140	145	150	
Leu Ala Asn Val	Ser Leu Leu Gln	Ala	Thr Glu	Trp Ala	Ser Ser	155	160	165	
Glu Asp Thr Arg	Arg Ser Cys Gln	Ser	Lys Gly	Lys Thr	Glu Glu	170	175	180	
Glu Cys Gln Asn	Tyr Val Arg Val	Leu	Ile Val	Ala Gly	Arg Lys	185	190	195	
Val Phe Met Cys	Gly Thr Asn Ala	Phe	Ser Pro	Met Cys	Thr Ser	200	205	210	
Arg Gln Val Gly	Asn Leu Ser Arg	Thr	Ile Glu	Lys Ile	Asn Gly	215	220	225	
Val Ala Arg Cys	Pro Tyr Asp Pro	Arg	His Asn	Ser Thr	Ala Val	230	235	240	
Ile Ser Ser Gln	Gly Glu Leu Tyr	Ala	Ala Thr	Val Ile	Asp Phe	245	250	255	
Ser Gly Arg Asp	Pro Ala Ile Tyr	Arg	Ser Leu	Gly Ser	Gly Pro	260	265	270	
Pro Leu Arg Thr	Ala Gln Tyr Asn	Ser	Lys Trp	Leu Asn	Glu Pro	275	280	285	
Asn Phe Val Ala	Ala Tyr Asp Ile	Gly	Leu Phe	Ala Tyr	Phe Phe	290	295	300	
Leu Arg Glu Asn	Ala Val Glu His	Asp	Cys Gly	Arg Thr	Val Tyr	305	310	315	
Ser Arg Val Ala	Arg Val Cys Lys	Asn	Asp Val	Gly Gly	Arg Phe	320	325	330	
Leu Leu Glu Asp	Thr Trp Thr Thr	Phe	Met Lys	Ala Arg	Leu Asn	335	340	345	
Cys Ser Arg Pro	Gly Glu Val Pro	Phe	Tyr Tyr	Asn Glu	Leu Gln	350	355	360	
Ser Ala Phe His	Leu Pro Glu Gln	Asp	Leu Ile	Tyr Gly	Val Phe	365	370	375	
Thr Thr Asn Val	Asn Ser Ile Ala	Ala	Ser Ala	Val Cys	Ala Phe	380	385	390	
Asn Leu Ser Ala	Ile Ser Gln Ala	Phe	Asn Gly	Pro Phe	Arg Tyr	395	400	405	
Gln Glu Asn Pro	Arg Ala Ala Trp	Leu	Pro Ile	Ala Asn	Pro Ile	410	415	420	
Pro Asn Phe Gln	Cys Gly Thr Leu	Pro	Glu Thr	Gly Pro	Asn Glu	425	430	435	

-continued

Asn	Leu	Thr	Glu	Arg	Ser	Leu	Gln	Asp	Ala	Gln	Arg	Leu	Phe	Leu	440	445	450
Met	Ser	Glu	Ala	Val	Gln	Pro	Val	Thr	Pro	Glu	Pro	Cys	Val	Thr	455	460	465
Gln	Asp	Ser	Val	Arg	Phe	Ser	His	Leu	Val	Val	Asp	Leu	Val	Gln	470	475	480
Ala	Lys	Asp	Thr	Leu	Tyr	His	Val	Leu	Tyr	Ile	Gly	Thr	Glu	Ser	485	490	495
Gly	Thr	Ile	Leu	Lys	Ala	Leu	Ser	Thr	Ala	Ser	Arg	Ser	Leu	His	500	505	510
Gly	Cys	Tyr	Leu	Glu	Glu	Leu	His	Val	Leu	Pro	Pro	Gly	Arg	Arg	515	520	525
Glu	Pro	Leu	Arg	Ser	Leu	Arg	Ile	Leu	His	Ser	Ala	Arg	Ala	Leu	530	535	540
Phe	Val	Gly	Leu	Arg	Asp	Gly	Val	Leu	Arg	Val	Pro	Leu	Glu	Arg	545	550	555
Cys	Ala	Ala	Tyr	Arg	Ser	Gln	Gly	Ala	Cys	Leu	Gly	Ala	Arg	Asp	560	565	570
Pro	Tyr	Cys	Gly	Trp	Asp	Gly	Lys	Gln	Gln	Arg	Cys	Ser	Thr	Leu	575	580	585
Glu	Asp	Ser	Ser	Asn	Met	Ser	Leu	Trp	Thr	Gln	Asn	Ile	Thr	Ala	590	595	600
Cys	Pro	Val	Arg	Asn	Val	Thr	Arg	Asp	Gly	Gly	Phe	Gly	Pro	Trp	605	610	615
Ser	Pro	Trp	Gln	Pro	Cys	Glu	His	Leu	Asp	Gly	Asp	Asn	Ser	Gly	620	625	630
Ser	Cys	Leu	Cys	Arg	Ala	Arg	Ser	Cys	Asp	Ser	Pro	Arg	Pro	Arg	635	640	645
Cys	Gly	Gly	Leu	Asp	Cys	Leu	Gly	Pro	Ala	Ile	His	Ile	Ala	Asn	650	655	660
Cys	Ser	Arg	Asn	Gly	Ala	Trp	Thr	Pro	Trp	Ser	Ser	Trp	Ala	Leu	665	670	675
Cys	Ser	Thr	Ser	Cys	Gly	Ile	Gly	Phe	Gln	Val	Arg	Gln	Arg	Ser	680	685	690
Cys	Ser	Asn	Pro	Ala	Pro	Arg	His	Gly	Gly	Arg	Ile	Phe	Val	Gly	695	700	705
Lys	Ser	Arg	Glu	Glu	Arg	Phe	Cys	Asn	Glu	Asn	Thr	Pro	Cys	Pro	710	715	720
Val	Pro	Ile	Phe	Trp	Ala	Ser	Trp	Gly	Ser	Trp	Ser	Lys	Cys	Ser	725	730	735
Ser	Asn	Cys	Gly	Gly	Gly	Met	Gln	Ser	Arg	Arg	Arg	Ala	Cys	Glu	740	745	750
Asn	Gly	Asn	Ser	Cys	Leu	Gly	Cys	Gly	Glu	Phe	Lys	Thr	Cys	Asn	755	760	765
Pro	Glu	Gly	Cys	Pro	Glu	Val	Arg	Arg	Asn	Thr	Pro	Trp	Thr	Pro	770	775	780
Trp	Leu	Pro	Val	Asn	Val	Thr	Gln	Gly	Gly	Ala	Arg	Gln	Glu	Gln	785	790	795
Arg	Phe	Arg	Phe	Thr	Cys	Arg	Ala	Pro	Leu	Ala	Asp	Pro	His	Gly	800	805	810
Leu	Gln	Phe	Gly	Arg	Arg	Arg	Thr	Glu	Thr	Arg	Thr	Cys	Pro	Ala			

-continued

815								820					825				
Asp	Gly	Ser	Gly	Ser	Cys	Asp	Thr	Asp	Ala	Leu	Val	Glu	Val	Leu			
830								835					840				
Leu	Arg	Ser	Gly	Ser	Thr	Ser	Pro	His	Thr	Val	Ser	Gly	Gly	Trp			
845								850					855				
Ala	Ala	Trp	Gly	Pro	Trp	Ser	Ser	Cys	Ser	Arg	Asp	Cys	Glu	Leu			
860								865					870				
Gly	Phe	Arg	Val	Arg	Lys	Arg	Thr	Cys	Thr	Asn	Pro	Glu	Pro	Arg			
875								880					885				
Asn	Gly	Gly	Leu	Pro	Cys	Val	Gly	Asp	Ala	Ala	Glu	Tyr	Gln	Asp			
890								895					900				
Cys	Asn	Pro	Gln	Ala	Cys	Pro	Val	Arg	Gly	Ala	Trp	Ser	Cys	Trp			
905								910					915				
Thr	Ser	Trp	Ser	Pro	Cys	Ser	Ala	Ser	Cys	Gly	Gly	Gly	His	Tyr			
920								925					930				
Gln	Arg	Thr	Arg	Ser	Cys	Thr	Ser	Pro	Ala	Pro	Ser	Pro	Gly	Glu			
935								940					945				
Asp	Ile	Cys	Leu	Gly	Leu	His	Thr	Glu	Glu	Ala	Leu	Cys	Ala	Thr			
950								955					960				
Gln	Ala	Cys	Pro	Gly	Trp	Ser	Pro	Trp	Ser	Glu	Trp	Ser	Lys	Cys			
965								970					975				
Thr	Asp	Asp	Gly	Ala	Gln	Ser	Arg	Ser	Arg	His	Cys	Glu	Glu	Leu			
980								985					990				
Leu	Pro	Gly	Ser	Ser	Ala	Cys	Ala	Gly	Asn	Ser	Ser	Gln	Ser	Arg			
995								1000					1005				
Pro	Cys	Pro	Tyr	Ser	Glu	Ile	Pro	Val	Ile	Leu	Pro	Ala	Ser	Ser			
1010								1015					1020				
Met	Glu	Glu	Ala	Thr	Asp	Cys	Ala	Gly	Lys	Arg	Asn	Arg	Thr	Tyr			
1025								1030					1035				
Leu	Met	Leu	Arg	Ser	Ser	Gln	Pro	Ser	Ser	Thr	Pro	Leu	Gln	Ser			
1040								1045					1050				
Leu	Asp	Ser	Phe	His	Ile	Leu	Leu	Gln	Thr	Ala	Lys	Leu	Cys	Trp			
1055								1060					1065				
Gly	Pro	His	Cys	Phe	Glu	Met	Gly	Ser	Ile	Ser	Ser	Thr	Trp	Trp			
1070								1075					1080				
Pro	Arg	Ala	Ser	Pro	Ala	Ser	Trp	Ala	Leu	Gly	Ser						
1085								1090									

<210> SEQ ID NO 97
<211> LENGTH: 3391
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 97

caagccctcc cagcatcccc tctcctgtgt tcctccccag ttctctactc	50
agagttgact gaccagagat ttatcagctt ggagggtctg aggtgtggat	100
ccatggggta gcctcaacgc atctgccctt ccaccccagc cagctcatgg	150
gccacgtggc ctggcccagc ctcagcaccc agggccagtg aacagagccc	200
tggtctggagt ccaaacatgt ggggccttgt gaggtcctg ctggcctggc	250
tggtgtggctg gggctgcatg gggcgtctgg cagccccagc ccgggcctgg	300

-continued

gcaggggtccc	gggaacaccc	agggcctgct	ctgctgcgga	ctcgaaggag	350
ctgggtctgg	aaccagttct	ttgtcattga	ggaatatgct	ggtccagagc	400
ctgttctcat	tggcaagctg	cactcggatg	ttgaccgggg	agagggccgc	450
accaagtacc	tgttgaccgg	ggagggggca	ggcaccgtat	ttgtgattga	500
tgaggccaca	ggcaatatc	atgttaccaa	gagccttgac	cgggaggaaa	550
agggcgaata	tgtgctactg	gcccagccg	tggaccgagc	ctccaaccgg	600
ccccctggagc	ccccatcaga	gttcacatc	aaagtgaag	acatcaacga	650
caatccaccc	atttttcccc	ttgggcccct	ccatgccacc	gtgcccagaga	700
tgtccaatgt	cgggacatca	gtgatccagg	tgactgctca	cgatgctgat	750
gacccagct	atgggaacag	tgccaagctg	gtgtacactg	ttctggatgg	800
actgcctttc	ttctctgtgg	acccccagac	tggagtgggtg	cgtacagcca	850
tccccaacat	ggaccgggag	acacaggagg	agttcttggt	ggtgatccag	900
gccaaagaca	tgggcgccca	catggggggg	ctgtcaggca	gcaactacgg	950
gactgtcacg	ctcagcgatg	tcaacgacaa	ccccccaag	ttcccacaga	1000
gcctatacca	gttctccgtg	gtggagacag	ctggaccctg	cacactgggtg	1050
ggccggctcc	gggcccagga	cccagacctg	ggggacaacg	ccctgatggc	1100
atacagcatc	ctggatgggg	aggggtctga	ggccttcagc	atcagcacag	1150
acttgacagg	tcgagacggg	ctcctcactg	tccgcaagcc	cctagacttt	1200
gagagccagc	gctcctactc	cttccgtgtc	gaggccacca	acacgctcat	1250
tgacccagcc	tatctgcggc	gagggcccct	caaggatgtg	gcctctgtgc	1300
gtgtggcagt	gcaagatgcc	ccagagccac	ctgccttcac	ccaggctgcc	1350
taccacctga	cagtgcctga	gaacaaggcc	ccggggaccc	tggtaggcca	1400
gatctccgcg	gctgacctgg	actcccctgc	cagcccaatc	agatactcca	1450
tcctccccca	ctcagatccg	gagcgttgct	tctctatcca	gcccagaggaa	1500
ggcaccatcc	atacagcagc	acccctggat	cgcgaggctc	gcgcctggca	1550
caacctcact	gtgctggcta	cagagctcga	cagttctgca	caggcctcgc	1600
gcgtgcaagt	ggccatccag	accctggatg	agaatgacaa	tgctccccag	1650
ctggctgagc	cctacgatac	ttttgtgtgt	gactctgcag	ctcctggcca	1700
gctgattcag	gtcatccggg	ccctggacag	agatgaagtt	ggcaacagta	1750
gccatgtctc	ctttcaaggt	cctctggggc	ctgatgcaa	ctttactgtc	1800
caggacaacc	gagatggctc	cgccagcctg	ctgctgcctc	cccgccctgc	1850
tccaccccgc	catgccccct	acttggttcc	catagaactg	tgggactggg	1900
ggcagccggc	gctgagcagc	actgccacag	tgactgttag	tgtgtgccgc	1950
tgccagcctg	acggctctgt	ggcatcctgc	tggcctgagg	ctcacctctc	2000
agctgctggg	ctcagcaccg	gcgcctctgt	tgccatcatc	acctgtgtgg	2050
gtgccctgct	tgccctgggt	gtgctcttcg	tggccctgcg	gcggcagaag	2100
caagaagcac	tgatggtaact	ggaggaggag	gacgtccgag	agaacatcat	2150
cacctacgac	gacgagggcg	gcggcgagga	ggacaccgag	gccttcgaca	2200

-continued

tcacggcctt gcagaaccgc gacggggcgg cccccccgc gcccgccct	2250
cccgcgcc gagacgtgtt gccccggcc cgggtgtcgc gccagccag	2300
ccccccgc cccgccgcg tggcgagct cctggcgtg cggctccgcg	2350
aggcggacga ggaccccgcc gtaccccgct acgactcgt gcagggtgtac	2400
ggctacgagg gccgcggctc ctcttgccgc tccctcagct ccctgggctc	2450
cggcagcgaa gccggcggcg cccccggccc cgcggagccg ctggacgact	2500
ggggtcctgt cttccgcacc ctggccgagc tgtatggggc caaggagccc	2550
ccggccccct gagcgcccg gctggcccg cccaccgcgg ggggggggca	2600
gcgggcacag gccctctgag tgagccccc ggggtccagg cgggcggcag	2650
cagcccagg gccccaggcc tcctccctgt ccttgtgtcc ctcttgctt	2700
ccccggggca ccctcgtct caccctccctc ctctgagtc ggtgtgtgtg	2750
tctctctcca ggaatctttg tctctatctg tgacacgctc ctctgtccg	2800
gcctgggttt cctgccctgg ccctggccct gogatctctc actgtgattc	2850
ctctccttcc tccgtggcgt tttgtctctg cagttctgaa gctcacacat	2900
agtctccctg cgtcttcctt gccatacac atgctctgtg tctgtctcct	2950
gccacatct cccttccttc tctctgggtc cctgtgactg gctttttgtt	3000
ttttctgtt gtccatccca aaatcaagag aaacttcag ccactgctgc	3050
ccaccctcct gcaggggatg ttgtgcccc gacctgcctg catggttcca	3100
tccattactc atggcctcag cctcatcctg gctccactgg cctccagctg	3150
agagagggaa ccagcctgcc tcccagggca agagctccag cctcccggtg	3200
ggccgcctcc ctggagctct gccagctgc cagcttcccc tgggcacccc	3250
agccctgggc attgtcttgt gtgcttcctg agggagtagg gaaaggaaag	3300
ggggaggcgg ctggggaagg ggaaagagg aggaaggga ggggcctcca	3350
tctctaattt cataataaac aaacacttta ttttgtaaaa c	3391

<210> SEQ ID NO 98

<211> LENGTH: 781

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 98

Met Trp Gly Leu Val Arg Leu Leu Leu Ala Trp Leu Gly Gly Trp	
1 5 10 15	
Gly Cys Met Gly Arg Leu Ala Ala Pro Ala Arg Ala Trp Ala Gly	
20 25 30	
Ser Arg Glu His Pro Gly Pro Ala Leu Leu Arg Thr Arg Arg Ser	
35 40 45	
Trp Val Trp Asn Gln Phe Phe Val Ile Glu Glu Tyr Ala Gly Pro	
50 55 60	
Glu Pro Val Leu Ile Gly Lys Leu His Ser Asp Val Asp Arg Gly	
65 70 75	
Glu Gly Arg Thr Lys Tyr Leu Leu Thr Gly Glu Gly Ala Gly Thr	
80 85 90	
Val Phe Val Ile Asp Glu Ala Thr Gly Asn Ile His Val Thr Lys	
95 100 105	

-continued

Ser	Leu	Asp	Arg	Glu	Glu	Lys	Ala	Gln	Tyr	Val	Leu	Leu	Ala	Gln
				110					115					120
Ala	Val	Asp	Arg	Ala	Ser	Asn	Arg	Pro	Leu	Glu	Pro	Pro	Ser	Glu
				125					130					135
Phe	Ile	Ile	Lys	Val	Gln	Asp	Ile	Asn	Asp	Asn	Pro	Pro	Ile	Phe
				140					145					150
Pro	Leu	Gly	Pro	Tyr	His	Ala	Thr	Val	Pro	Glu	Met	Ser	Asn	Val
				155					160					165
Gly	Thr	Ser	Val	Ile	Gln	Val	Thr	Ala	His	Asp	Ala	Asp	Asp	Pro
				170					175					180
Ser	Tyr	Gly	Asn	Ser	Ala	Lys	Leu	Val	Tyr	Thr	Val	Leu	Asp	Gly
				185					190					195
Leu	Pro	Phe	Phe	Ser	Val	Asp	Pro	Gln	Thr	Gly	Val	Val	Arg	Thr
				200					205					210
Ala	Ile	Pro	Asn	Met	Asp	Arg	Glu	Thr	Gln	Glu	Glu	Phe	Leu	Val
				215					220					225
Val	Ile	Gln	Ala	Lys	Asp	Met	Gly	Gly	His	Met	Gly	Gly	Leu	Ser
				230					235					240
Gly	Ser	Thr	Thr	Val	Thr	Val	Thr	Leu	Ser	Asp	Val	Asn	Asp	Asn
				245					250					255
Pro	Pro	Lys	Phe	Pro	Gln	Ser	Leu	Tyr	Gln	Phe	Ser	Val	Val	Glu
				260					265					270
Thr	Ala	Gly	Pro	Gly	Thr	Leu	Val	Gly	Arg	Leu	Arg	Ala	Gln	Asp
				275					280					285
Pro	Asp	Leu	Gly	Asp	Asn	Ala	Leu	Met	Ala	Tyr	Ser	Ile	Leu	Asp
				290					295					300
Gly	Glu	Gly	Ser	Glu	Ala	Phe	Ser	Ile	Ser	Thr	Asp	Leu	Gln	Gly
				305					310					315
Arg	Asp	Gly	Leu	Leu	Thr	Val	Arg	Lys	Pro	Leu	Asp	Phe	Glu	Ser
				320					325					330
Gln	Arg	Ser	Tyr	Ser	Phe	Arg	Val	Glu	Ala	Thr	Asn	Thr	Leu	Ile
				335					340					345
Asp	Pro	Ala	Tyr	Leu	Arg	Arg	Gly	Pro	Phe	Lys	Asp	Val	Ala	Ser
				350					355					360
Val	Arg	Val	Ala	Val	Gln	Asp	Ala	Pro	Glu	Pro	Pro	Ala	Phe	Thr
				365					370					375
Gln	Ala	Ala	Tyr	His	Leu	Thr	Val	Pro	Glu	Asn	Lys	Ala	Pro	Gly
				380					385					390
Thr	Leu	Val	Gly	Gln	Ile	Ser	Ala	Ala	Asp	Leu	Asp	Ser	Pro	Ala
				395					400					405
Ser	Pro	Ile	Arg	Tyr	Ser	Ile	Leu	Pro	His	Ser	Asp	Pro	Glu	Arg
				410					415					420
Cys	Phe	Ser	Ile	Gln	Pro	Glu	Glu	Gly	Thr	Ile	His	Thr	Ala	Ala
				425					430					435
Pro	Leu	Asp	Arg	Glu	Ala	Arg	Ala	Trp	His	Asn	Leu	Thr	Val	Leu
				440					445					450
Ala	Thr	Glu	Leu	Asp	Ser	Ser	Ala	Gln	Ala	Ser	Arg	Val	Gln	Val
				455					460					465
Ala	Ile	Gln	Thr	Leu	Asp	Glu	Asn	Asp	Asn	Ala	Pro	Gln	Leu	Ala
				470					475					480

-continued

Glu	Pro	Tyr	Asp	Thr	Phe	Val	Cys	Asp	Ser	Ala	Ala	Pro	Gly	Gln	
			485						490					495	
Leu	Ile	Gln	Val	Ile	Arg	Ala	Leu	Asp	Arg	Asp	Glu	Val	Gly	Asn	
			500						505					510	
Ser	Ser	His	Val	Ser	Phe	Gln	Gly	Pro	Leu	Gly	Pro	Asp	Ala	Asn	
			515						520					525	
Phe	Thr	Val	Gln	Asp	Asn	Arg	Asp	Gly	Ser	Ala	Ser	Leu	Leu	Leu	
			530						535					540	
Pro	Ser	Arg	Pro	Ala	Pro	Pro	Arg	His	Ala	Pro	Tyr	Leu	Val	Pro	
			545						550					555	
Ile	Glu	Leu	Trp	Asp	Trp	Gly	Gln	Pro	Ala	Leu	Ser	Ser	Thr	Ala	
			560						565					570	
Thr	Val	Thr	Val	Ser	Val	Cys	Arg	Cys	Gln	Pro	Asp	Gly	Ser	Val	
			575						580					585	
Ala	Ser	Cys	Trp	Pro	Glu	Ala	His	Leu	Ser	Ala	Ala	Gly	Leu	Ser	
			590						595					600	
Thr	Gly	Ala	Leu	Leu	Ala	Ile	Ile	Thr	Cys	Val	Gly	Ala	Leu	Leu	
			605						610					615	
Ala	Leu	Val	Val	Leu	Phe	Val	Ala	Leu	Arg	Arg	Gln	Lys	Gln	Glu	
			620						625					630	
Ala	Leu	Met	Val	Leu	Glu	Glu	Glu	Asp	Val	Arg	Glu	Asn	Ile	Ile	
			635						640					645	
Thr	Tyr	Asp	Asp	Glu	Gly	Gly	Gly	Glu	Glu	Asp	Thr	Glu	Ala	Phe	
			650						655					660	
Asp	Ile	Thr	Ala	Leu	Gln	Asn	Pro	Asp	Gly	Ala	Ala	Pro	Pro	Ala	
			665						670					675	
Pro	Gly	Pro	Pro	Ala	Arg	Arg	Asp	Val	Leu	Pro	Arg	Ala	Arg	Val	
			680						685					690	
Ser	Arg	Gln	Pro	Arg	Pro	Pro	Gly	Pro	Ala	Asp	Val	Ala	Gln	Leu	
			695						700					705	
Leu	Ala	Leu	Arg	Leu	Arg	Glu	Ala	Asp	Glu	Asp	Pro	Gly	Val	Pro	
			710						715					720	
Pro	Tyr	Asp	Ser	Val	Gln	Val	Tyr	Gly	Tyr	Glu	Gly	Arg	Gly	Ser	
			725						730					735	
Ser	Cys	Gly	Ser	Leu	Ser	Ser	Leu	Gly	Ser	Gly	Ser	Glu	Ala	Gly	
			740						745					750	
Gly	Ala	Pro	Gly	Pro	Ala	Glu	Pro	Leu	Asp	Asp	Trp	Gly	Pro	Leu	
			755						760					765	
Phe	Arg	Thr	Leu	Ala	Glu	Leu	Tyr	Gly	Ala	Lys	Glu	Pro	Pro	Ala	
			770						775					780	
Pro															

<210> SEQ ID NO 99
<211> LENGTH: 2855
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 99

gccaaactg gccaaacata tggggctgga atctcaacat cggtcactgg	50
gacctcaata tttggagccg gaacccccaca atttgggaaca cagaccccaa	100
tatttggagc agaaccctaa gatttgacat ctaaaacctc aagcctggag	150

-continued

ctgaactctg aattctgggc ctgggacctt gaaatctggg actggatttc	200
cagtactgta ccctggaacc cactcttggg gacctgaacc ctgggattca	250
ggcctcaaat tccaagatct ggactgtggg attccaaggg gcctgaaccc	300
gagtttgggc ctgaagtcct tgctgcagac ctgagtgtt aaatctggg	350
cttgagacct cccaatcttg actcagcacc ccaatatctg aatgcagaac	400
cccgggatcg gatctcagac tctaaacccc accgtttggc tgcttagcat	450
cccaagactg gacctgggag accctgaccc tgaacaaccc aaactggacc	500
cgtaaaactg gaccctagag gcccaatatt taggggtctg gaaccccgag	550
tattaaggtc tggagactcc gttgccacag atttgagccg agtcaggaca	600
cagtccctct acagaagcct tggggacagg aaaagcatga ccagatgtc	650
cctccagagc cctgacctct gactcccctg gagctaggac tctgtccct	700
ggggtgtctt ctgctcagg acaccctgc ccgcatggc catcctcccg	750
ttgtctctgt gcctgtgtcc gctggcccct gctcatccc caccctcgtc	800
agccacaccc agcccatgtc ccgcgcgtg ccgctgccag acacagtgc	850
tgcccctaag cgtgtgtgtc ccaggggcag gcctcctgtt cgtgccaccc	900
tcgttggaac gccgggcagc cgagctgcgg ctggcagaca acttcacgc	950
ctccgtgtgc cgccgcgacc tggccaacat gacaggcctg ctgcatctga	1000
gcctgtcgcg gaacaccatc cgccacgtgg ctgccggcgc cttcgccgac	1050
ctgccccgcc tgcgtgccct gcacctggat ggcaaccggc tgacctact	1100
gggcgagggc cagctgcgcg gcctgtgtcaa cttgcgccac ctcatcctca	1150
gcaacaacca gctggcagcg ctggcgcccg gcgccctgga tgattgtgcc	1200
gagacactgg aggacctcga cctctctac aacaacctcg agcagctgcc	1250
ctgggaggcc ctgggcgcgc tgggcaacgt caacacgttg ggcctcgacc	1300
acaacctgct ggcttctgtg ccggcgctt tttcccgcct gcacaagctg	1350
gcccggctgg acatgacctc caaccgcctg accacaatcc caccgaccc	1400
actcttctcc cgctgcccc tgctcgcag gcccggggc tcgccgcct	1450
ctgccctggt gctggccttt ggcggaacc ccctgactg caactgcgag	1500
ctggtgtggc tgcgtgcctt ggcgcgggag gacgacctc aggcctgcgc	1550
gtccccacct gctctgggcg gccgctactt ctggcggtg ggcgaggag	1600
agtttgtctg cgagccgcc gtggtgactc accgctcacc acctctggt	1650
gtgcccgag gtccggcggc tgccctgcgc tgccgggcag tgggggaccc	1700
agagccccgt gtgcgttggg tgtacccca gggccggtg ctaggcaact	1750
caagccgtgc ccgcgccttc cccaatggga cgctggagct gctggtcacc	1800
gagccgggtg atggtggcat cttcacctgc attgcggcca atgcagctgg	1850
cgagccaca gctgtgtgg agctgactgt gggccccca ccacctctc	1900
agctagccaa cagcaccagc tgtgaccccc cgcgggacgg ggcctctgat	1950
gtctctaccc caccctcgc tgctctgtct totgccaagg tggccgacac	2000
tgggccccct accgacctg gcgtccaggt gactgagcac ggggccacag	2050

-continued

ctgctcttgt ccagtggccg gatcagcggc ctatcccggg catccgcatg	2100
taccagatcc agtacaacag ctcggtgat gacatcctcg tctacaggat	2150
gatcccggcg gagagccgct cgttctctgt gacggacctg gcgtcaggcc	2200
ggacctacga tctgtgcgtg ctgcgcgtgt atgaggacag cgccacgggg	2250
ctcacggcca cgcggcctgt gggctgcgcc cgcttctcca ccgaacctgc	2300
gctgcggcca tgcggggcgc cgcacgctcc cttcctgggc ggcacgatga	2350
tcatcgcgct gggcggcgct atcgtagcct cggtactggt cttcatcttc	2400
gtgctgctaa tgcgctacaa ggtgcacggc ggccagcccc ccggcaaggc	2450
caagattccc gcgcctgtta gcagcgtttg ctcccagacc aacggcgccc	2500
tgggccccac gccacgccc gccccgccg ccccgagacc cgcggcgctc	2550
agggcccaca ccgtggtcca gctggactgc gagccctggg ggcccggcca	2600
cgaacctgtg ggaccctagc caggcgcccc cccctctaag ggtcctctg	2650
ccccacggag agcaggaccc ggacaccctg tgggacctgg cctcaactc	2700
accaaatacg tcatggtttt taaaactctg atggggaggg tgtcggggac	2750
accggggcaa aacaagaaag tcctattttt ccaaaaaaaaa aaaaaaaaaa	2800
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	2850
aaaaaa	2855

<210> SEQ ID NO 100
<211> LENGTH: 627
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 100

Met Ala Ile Leu Pro Leu Leu Leu Cys Leu Leu Pro Leu Ala Pro	1 5 10 15
Ala Ser Ser Pro Pro Gln Ser Ala Thr Pro Ser Pro Cys Pro Arg	20 25 30
Arg Cys Arg Cys Gln Thr Gln Ser Leu Pro Leu Ser Val Leu Cys	35 40 45
Pro Gly Ala Gly Leu Leu Phe Val Pro Pro Ser Leu Asp Arg Arg	50 55 60
Ala Ala Glu Leu Arg Leu Ala Asp Asn Phe Ile Ala Ser Val Arg	65 70 75
Arg Arg Asp Leu Ala Asn Met Thr Gly Leu Leu His Leu Ser Leu	80 85 90
Ser Arg Asn Thr Ile Arg His Val Ala Ala Gly Ala Phe Ala Asp	95 100 105
Leu Arg Ala Leu Arg Ala Leu His Leu Asp Gly Asn Arg Leu Thr	110 115 120
Ser Leu Gly Glu Gly Gln Leu Arg Gly Leu Val Asn Leu Arg His	125 130 135
Leu Ile Leu Ser Asn Asn Gln Leu Ala Ala Leu Ala Ala Gly Ala	140 145 150
Leu Asp Asp Cys Ala Glu Thr Leu Glu Asp Leu Asp Leu Ser Tyr	155 160 165
Asn Asn Leu Glu Gln Leu Pro Trp Glu Ala Leu Gly Arg Leu Gly	

-continued

170					175					180				
Asn	Val	Asn	Thr	Leu	Gly	Leu	Asp	His	Asn	Leu	Leu	Ala	Ser	Val
				185					190					195
Pro	Gly	Ala	Phe	Ser	Arg	Leu	His	Lys	Leu	Ala	Arg	Leu	Asp	Met
				200					205					210
Thr	Ser	Asn	Arg	Leu	Thr	Thr	Ile	Pro	Pro	Asp	Pro	Leu	Phe	Ser
				215					220					225
Arg	Leu	Pro	Leu	Leu	Ala	Arg	Pro	Arg	Gly	Ser	Pro	Ala	Ser	Ala
				230					235					240
Leu	Val	Leu	Ala	Phe	Gly	Gly	Asn	Pro	Leu	His	Cys	Asn	Cys	Glu
				245					250					255
Leu	Val	Trp	Leu	Arg	Arg	Leu	Ala	Arg	Glu	Asp	Asp	Leu	Glu	Ala
				260					265					270
Cys	Ala	Ser	Pro	Pro	Ala	Leu	Gly	Gly	Arg	Tyr	Phe	Trp	Ala	Val
				275					280					285
Gly	Glu	Glu	Glu	Phe	Val	Cys	Glu	Pro	Pro	Val	Val	Thr	His	Arg
				290					295					300
Ser	Pro	Pro	Leu	Ala	Val	Pro	Ala	Gly	Arg	Pro	Ala	Ala	Leu	Arg
				305					310					315
Cys	Arg	Ala	Val	Gly	Asp	Pro	Glu	Pro	Arg	Val	Arg	Trp	Val	Ser
				320					325					330
Pro	Gln	Gly	Arg	Leu	Leu	Gly	Asn	Ser	Ser	Arg	Ala	Arg	Ala	Phe
				335					340					345
Pro	Asn	Gly	Thr	Leu	Glu	Leu	Leu	Val	Thr	Glu	Pro	Gly	Asp	Gly
				350					355					360
Gly	Ile	Phe	Thr	Cys	Ile	Ala	Ala	Asn	Ala	Ala	Gly	Glu	Ala	Thr
				365					370					375
Ala	Ala	Val	Glu	Leu	Thr	Val	Gly	Pro	Pro	Pro	Pro	Pro	Gln	Leu
				380					385					390
Ala	Asn	Ser	Thr	Ser	Cys	Asp	Pro	Pro	Arg	Asp	Gly	Asp	Pro	Asp
				395					400					405
Ala	Leu	Thr	Pro	Pro	Ser	Ala	Ala	Ser	Ala	Ser	Ala	Lys	Val	Ala
				410					415					420
Asp	Thr	Gly	Pro	Pro	Thr	Asp	Arg	Gly	Val	Gln	Val	Thr	Glu	His
				425					430					435
Gly	Ala	Thr	Ala	Ala	Leu	Val	Gln	Trp	Pro	Asp	Gln	Arg	Pro	Ile
				440					445					450
Pro	Gly	Ile	Arg	Met	Tyr	Gln	Ile	Gln	Tyr	Asn	Ser	Ser	Ala	Asp
				455					460					465
Asp	Ile	Leu	Val	Tyr	Arg	Met	Ile	Pro	Ala	Glu	Ser	Arg	Ser	Phe
				470					475					480
Leu	Leu	Thr	Asp	Leu	Ala	Ser	Gly	Arg	Thr	Tyr	Asp	Leu	Cys	Val
				485					490					495
Leu	Ala	Val	Tyr	Glu	Asp	Ser	Ala	Thr	Gly	Leu	Thr	Ala	Thr	Arg
				500					505					510
Pro	Val	Gly	Cys	Ala	Arg	Phe	Ser	Thr	Glu	Pro	Ala	Leu	Arg	Pro
				515					520					525
Cys	Gly	Ala	Pro	His	Ala	Pro	Phe	Leu	Gly	Gly	Thr	Met	Ile	Ile
				530					535					540
Ala	Leu	Gly	Gly	Val	Ile	Val	Ala	Ser	Val	Leu	Val	Phe	Ile	Phe
				545					550					555

-continued

Val	Leu	Leu	Met	Arg	Tyr	Lys	Val	His	Gly	Gly	Gln	Pro	Pro	Gly
			560						565					570
Lys	Ala	Lys	Ile	Pro	Ala	Pro	Val	Ser	Ser	Val	Cys	Ser	Gln	Thr
			575						580					585
Asn	Gly	Ala	Leu	Gly	Pro	Thr	Pro	Thr	Pro	Ala	Pro	Pro	Ala	Pro
			590						595					600
Glu	Pro	Ala	Ala	Leu	Arg	Ala	His	Thr	Val	Val	Gln	Leu	Asp	Cys
			605						610					615
Glu	Pro	Trp	Gly	Pro	Gly	His	Glu	Pro	Val	Gly	Pro			
			620						625					

<210> SEQ ID NO 101
<211> LENGTH: 1111
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 101

cgactccata accgtggcct tggccccagt ccccttgact tccggacttc	50
agaccagata ctgcccatac ccccttatga agtcttgcc aggcaacccc	100
taggggtgtac gttttctaaa gattaaagag gcggtgctaa gctgcagacg	150
gacttgcgac tcagccactg gtgtaagtca ggcgggaggt ggcgccaat	200
aagctcaaga gaggaggcgg gttctggaaa aaggccaata gcctgtgaag	250
gcgagtctag cagcaaccaa tagctatgag cgagaggcgg gactctgagg	300
gaagtcaatc gctgccgcag gtaccgcaa tggcttttgg cgggggcgtt	350
ccccaacctt gccctctctc atgacccgc tccgggatta tggccgggac	400
tgggctgctg gcgctgcgga cgctgccagg gccagctgg gtgcgaggct	450
cgggcccctt cgtgctgagc cgctgcagg acgcggccgt ggtgcggcct	500
ggcttcctga gcacggcaga ggaggagacg ctgagccgag aactggagcc	550
cgagctgcgc cgccgcgcgt acgaatacga tcaactgggac gcggccatcc	600
acggcttcgc agagacagag aagtcgcgct ggtcagaagc cagccgggcc	650
atcctgcagc gcgtgcaggc ggccgccttt ggccccggcc agaccctgct	700
ctcctccgtg cacgtgctgg acctggaagc ccgcggctac atcaagcccc	750
acgtggacag catcaagttc tgcggggcca ccacgcgcg cctgtctctc	800
ctgtctccca gcgttatgcg gctggtgcac acccaggagc cgggggagtg	850
gctggaactc ttgctggagc cgggctccct ctacatcctt aggggctcag	900
cccgttatga cttctcccat gagatccttc gggatgaaga gtccttcttt	950
ggggaacgcc ggattccccg gggccggcgc atctccgtga tctgccgctc	1000
cctccctgag ggcattggggc caggggagtc tggacagccg cccccagcct	1050
gctgaccccc agctttctac agacaccaga tttgtgaata aagttgggga	1100
atggacagcc t	1111

<210> SEQ ID NO 102
<211> LENGTH: 221
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

-continued

<400> SEQUENCE: 102

Met	Ala	Gly	Thr	Gly	Leu	Leu	Ala	Leu	Arg	Thr	Leu	Pro	Gly	Pro
1				5					10					15
Ser	Trp	Val	Arg	Gly	Ser	Gly	Pro	Ser	Val	Leu	Ser	Arg	Leu	Gln
				20					25					30
Asp	Ala	Ala	Val	Val	Arg	Pro	Gly	Phe	Leu	Ser	Thr	Ala	Glu	Glu
				35					40					45
Glu	Thr	Leu	Ser	Arg	Glu	Leu	Glu	Pro	Glu	Leu	Arg	Arg	Arg	Arg
				50					55					60
Tyr	Glu	Tyr	Asp	His	Trp	Asp	Ala	Ala	Ile	His	Gly	Phe	Arg	Glu
				65					70					75
Thr	Glu	Lys	Ser	Arg	Trp	Ser	Glu	Ala	Ser	Arg	Ala	Ile	Leu	Gln
				80					85					90
Arg	Val	Gln	Ala	Ala	Ala	Phe	Gly	Pro	Gly	Gln	Thr	Leu	Leu	Ser
				95					100					105
Ser	Val	His	Val	Leu	Asp	Leu	Glu	Ala	Arg	Gly	Tyr	Ile	Lys	Pro
				110					115					120
His	Val	Asp	Ser	Ile	Lys	Phe	Cys	Gly	Ala	Thr	Ile	Ala	Gly	Leu
				125					130					135
Ser	Leu	Leu	Ser	Pro	Ser	Val	Met	Arg	Leu	Val	His	Thr	Gln	Glu
				140					145					150
Pro	Gly	Glu	Trp	Leu	Glu	Leu	Leu	Leu	Glu	Pro	Gly	Ser	Leu	Tyr
				155					160					165
Ile	Leu	Arg	Gly	Ser	Ala	Arg	Tyr	Asp	Phe	Ser	His	Glu	Ile	Leu
				170					175					180
Arg	Asp	Glu	Glu	Ser	Phe	Phe	Gly	Glu	Arg	Arg	Ile	Pro	Arg	Gly
				185					190					195
Arg	Arg	Ile	Ser	Val	Ile	Cys	Arg	Ser	Leu	Pro	Glu	Gly	Met	Gly
				200					205					210
Pro	Gly	Glu	Ser	Gly	Gln	Pro	Pro	Pro	Ala	Cys				
				215					220					

<210> SEQ ID NO 103
<211> LENGTH: 3583
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 103

ctccccggcg	ccgcaggcag	cgctctcctc	cgaagcagct	gcacctgcaa	50
ctggggcagcc	tggaccctcg	tgccctgttc	ccgggacctc	gcgcaggggg	100
cgccccggga	cacccctgc	gggccgggtg	gaggaggaag	aggaggagga	150
ggaagaagac	gtggacaagg	acccccatcc	taccagaac	acctgcctgc	200
gtgcccgcc	cttctcttta	agggagagga	aaagagagcc	taggagaacc	250
atggggggct	gcgaagtccg	ggaatttctt	ttgcaatttg	gtttcttctt	300
gcctctgctg	acagcgtggc	caggcgactg	cagtcacgtc	tccaacaacc	350
aagttgtgtt	gcttgataca	acaactgtac	tgggagagct	aggatggaaa	400
acatatccat	taaatgggtg	ggatgccatc	actgaaatgg	atgaacataa	450
taggcccatt	cacacatacc	aggtatgtaa	tgtaatggaa	ccaaaccaa	500
acaactggct	tcgtacaaac	tggatctccc	gtgatgcagc	tcagaaaatt	550

-continued

tatgtggaaa tgaaattcac actaagggat tgtaacagca tcccatgggt	600
cttggggact tgcaaagaaa catttaatct gttttatatg gaatcagatg	650
agtcccacgg aattaaattc aagccaaacc agtatacaaa gatcgacaca	700
attgctgctg atgagagttt taccagatg gatttgggtg atcgcatcct	750
caaaactcaac actgaaattc gtgagggtgg gcctatagaa aggaaaggat	800
tttatctggc ttttcaagac attggggcgt gcattgccct ggtttcagtc	850
cgtgttttct acaagaaatg ccccttcaact gttcgtaact tggccatgtt	900
tcctgatacc attccaaggg ttgattcctc ctctttgggt gaagtacggg	950
gttcttgtgt gaagagtgtt gaagagcgtg acactcctaa actgtattgt	1000
ggagctgatg gagattggct ggttcctctt ggaaggtgca tctgcagtac	1050
aggatatgaa gaaattgagg gttcttgcca tgcttgacga ccaggattct	1100
ataaagcttt tgctgggaac aaaaatgtt ctaaatgtcc tccacacagt	1150
ttaacataca tggaagcaac ttctgtctgt cagtgtgaaa agggttattt	1200
ccgagctgaa aaagaccac cttctatggc atgtaccagg ccaccttcag	1250
ctcctaggaa tgtgggtttt aacatcaatg aaacagccct tattttggaa	1300
tggagcccac caagtgcac agggaggaga aaagatctca catacagtgt	1350
aatctgtaag aatgtggct tagacaccag ccagtgtgag gactgtggtg	1400
gaggactcgg cttcatccca agacatacag gcctgatcaa caattccgtg	1450
atagtacttg actttgtgtc tcacgtgaat tacacctttg aaatagaagc	1500
aatgaatgga gtttctgagt tgagtttttc tccaagcca ttcacagcta	1550
ttacagtgcac cacggatcaa gatgcacctt ccctgatagg tgtggaagg	1600
aaggactggg catcccaaaa tagcattgcc ctatcatggc aagcacctgc	1650
tttttccaat ggagccattc tggactacga gatcaagtac tatgagaaa	1700
aacatgagca gctgacctac tcttcacaaa ggtccaaagc cccagtgctc	1750
atcatcacag gtcttaagcc agccacaaaa tatgtatttc acatccgagt	1800
gagaactcgg acaggataga gtggctacag tcagaaattt gaatttgaaa	1850
caggagatga aacttctgac atggcagcag aacaaggaca gattctcgtg	1900
atagccaccg ccgctgttgg cggattcact ctctctgtca tcctcacttt	1950
attcttctgt atcactggga gatgtcagtg gtacataaaa gccaaagtga	2000
agtcagaaga gaagagaaga aacctttac agaattggga tttgcgcttc	2050
ccgggaatta aaacttacat tgatccagat acatatgaag acccatccct	2100
agcagtcocat gaatttgcaa aggagattga tccctcaaga attcgtattg	2150
agagagtcat tggggcaggt gaatttggag aagtctgtag tgggcgtttg	2200
aagacaccag ggaaaagaga gatccagtt gccattaaaa ctttgaaagg	2250
tggccacatg gatcggaaca gaagagattt totaagagaa gctagtatca	2300
tgggccagtt tgacctcca aacatcattc gcctagaagg ggttgcacc	2350
aaaagatcct tcccggccat tgggggtggag gcgttttgcc ccagcttcct	2400
gagggcaggg tttttaata gcatccaggc cccgcattca gtgccagggg	2450

-continued

gaggatcttt gccccccagg attcctgctg gcagaccagt aatgattgtg	2500
gtggaatata tggagaatgg atccctagac tcctttttgc ggaagcatga	2550
tggccacttc acagtcatcc agttggtcgg aatgctccga ggcattgcat	2600
caggcatgaa gtatctttct gatatgggtt atgttcatcg agacctagcg	2650
gctcggaaata tactgggtcaa tagcaactta gtatgcaaag tttctgattt	2700
tggctctctcc agagtgtctg aagatgatcc agaagctgct tatacaacaa	2750
ctggtggaaa aatccccata aggtggacag cccagaagc catcgcttac	2800
agaaaattct cctcagcaag cgatgcattg agctatggca ttgtcatgtg	2850
ggaggtcatg tcctatggag agagacctta ttgggaaatg tctaaccaag	2900
atgtcattct gtccattgaa gaagggtaca gacttcacgc tcccatgggc	2950
tgtccagcat ctctacacca gctgatgctc cactgctggc agaaggagag	3000
aaatcacaga ccaaaattta ctgacattgt cagcttcctt gacaaactga	3050
tccgaaatcc cagtgcctt cacaccctgg tggaggacat ccttgtaatg	3100
ccagagtccc ctggtgaagt tccggaatat cctttgtttg tcacagtgg	3150
tgactggcta gattctataa agatggggca atacaagaat aacttcgtgg	3200
cagcaggggt tacaacattt gacctgattt caagaatgag cattgatgac	3250
attagaagaa ttggagtcac acttatttga caccagagac gaatagtcag	3300
cagcatacag actttacgtt tacacatgat gcacatacag gagaagggat	3350
ttcatgtatg aaagtaccac aagcacctgt gttttgtgcc tcagcatttc	3400
taaaatgaac gatatcctct ctactactct ctcttctgat tctccaaaca	3450
tcacttcaca aactgcagtc ttctgttcag actataggca cacaccttat	3500
gtttatgctt ccaaccagga ttttaaaatc atgctacata aatccgttct	3550
gaataacctg caactaaaaa aaaaaaaaaa aaa	3583

<210> SEQ ID NO 104

<211> LENGTH: 1036

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 104

Met Gly Gly Cys Glu Val Arg Glu Phe Leu Leu Gln Phe Gly Phe	
1 5 10 15	
Phe Leu Pro Leu Leu Thr Ala Trp Pro Gly Asp Cys Ser His Val	
20 25 30	
Ser Asn Asn Gln Val Val Leu Leu Asp Thr Thr Thr Val Leu Gly	
35 40 45	
Glu Leu Gly Trp Lys Thr Tyr Pro Leu Asn Gly Trp Asp Ala Ile	
50 55 60	
Thr Glu Met Asp Glu His Asn Arg Pro Ile His Thr Tyr Gln Val	
65 70 75	
Cys Asn Val Met Glu Pro Asn Gln Asn Asn Trp Leu Arg Thr Asn	
80 85 90	
Trp Ile Ser Arg Asp Ala Ala Gln Lys Ile Tyr Val Glu Met Lys	
95 100 105	

-continued

Phe	Thr	Leu	Arg	Asp	Cys	Asn	Ser	Ile	Pro	Trp	Val	Leu	Gly	Thr
				110					115					120
Cys	Lys	Glu	Thr	Phe	Asn	Leu	Phe	Tyr	Met	Glu	Ser	Asp	Glu	Ser
				125					130					135
His	Gly	Ile	Lys	Phe	Lys	Pro	Asn	Gln	Tyr	Thr	Lys	Ile	Asp	Thr
				140					145					150
Ile	Ala	Ala	Asp	Glu	Ser	Phe	Thr	Gln	Met	Asp	Leu	Gly	Asp	Arg
				155					160					165
Ile	Leu	Lys	Leu	Asn	Thr	Glu	Ile	Arg	Glu	Val	Gly	Pro	Ile	Glu
				170					175					180
Arg	Lys	Gly	Phe	Tyr	Leu	Ala	Phe	Gln	Asp	Ile	Gly	Ala	Cys	Ile
				185					190					195
Ala	Leu	Val	Ser	Val	Arg	Val	Phe	Tyr	Lys	Lys	Cys	Pro	Phe	Thr
				200					205					210
Val	Arg	Asn	Leu	Ala	Met	Phe	Pro	Asp	Thr	Ile	Pro	Arg	Val	Asp
				215					220					225
Ser	Ser	Ser	Leu	Val	Glu	Val	Arg	Gly	Ser	Cys	Val	Lys	Ser	Ala
				230					235					240
Glu	Glu	Arg	Asp	Thr	Pro	Lys	Leu	Tyr	Cys	Gly	Ala	Asp	Gly	Asp
				245					250					255
Trp	Leu	Val	Pro	Leu	Gly	Arg	Cys	Ile	Cys	Ser	Thr	Gly	Tyr	Glu
				260					265					270
Glu	Ile	Glu	Gly	Ser	Cys	His	Ala	Cys	Arg	Pro	Gly	Phe	Tyr	Lys
				275					280					285
Ala	Phe	Ala	Gly	Asn	Thr	Lys	Cys	Ser	Lys	Cys	Pro	Pro	His	Ser
				290					295					300
Leu	Thr	Tyr	Met	Glu	Ala	Thr	Ser	Val	Cys	Gln	Cys	Glu	Lys	Gly
				305					310					315
Tyr	Phe	Arg	Ala	Glu	Lys	Asp	Pro	Pro	Ser	Met	Ala	Cys	Thr	Arg
				320					325					330
Pro	Pro	Ser	Ala	Pro	Arg	Asn	Val	Val	Phe	Asn	Ile	Asn	Glu	Thr
				335					340					345
Ala	Leu	Ile	Leu	Glu	Trp	Ser	Pro	Pro	Ser	Asp	Thr	Gly	Gly	Arg
				350					355					360
Lys	Asp	Leu	Thr	Tyr	Ser	Val	Ile	Cys	Lys	Lys	Cys	Gly	Leu	Asp
				365					370					375
Thr	Ser	Gln	Cys	Glu	Asp	Cys	Gly	Gly	Gly	Leu	Arg	Phe	Ile	Pro
				380					385					390
Arg	His	Thr	Gly	Leu	Ile	Asn	Asn	Ser	Val	Ile	Val	Leu	Asp	Phe
				395					400					405
Val	Ser	His	Val	Asn	Tyr	Thr	Phe	Glu	Ile	Glu	Ala	Met	Asn	Gly
				410					415					420
Val	Ser	Glu	Leu	Ser	Phe	Ser	Pro	Lys	Pro	Phe	Thr	Ala	Ile	Thr
				425					430					435
Val	Thr	Thr	Asp	Gln	Asp	Ala	Pro	Ser	Leu	Ile	Gly	Val	Val	Arg
				440					445					450
Lys	Asp	Trp	Ala	Ser	Gln	Asn	Ser	Ile	Ala	Leu	Ser	Trp	Gln	Ala
				455					460					465
Pro	Ala	Phe	Ser	Asn	Gly	Ala	Ile	Leu	Asp	Tyr	Glu	Ile	Lys	Tyr
				470					475					480
Tyr	Glu	Lys	Glu	His	Glu	Gln	Leu	Thr	Tyr	Ser	Ser	Thr	Arg	Ser

-continued

485					490					495				
Lys	Ala	Pro	Ser	Val	Ile	Ile	Thr	Gly	Leu	Lys	Pro	Ala	Thr	Lys
				500					505					510
Tyr	Val	Phe	His	Ile	Arg	Val	Arg	Thr	Ala	Thr	Gly	Tyr	Ser	Gly
				515					520					525
Tyr	Ser	Gln	Lys	Phe	Glu	Phe	Glu	Thr	Gly	Asp	Glu	Thr	Ser	Asp
				530					535					540
Met	Ala	Ala	Glu	Gln	Gly	Gln	Ile	Leu	Val	Ile	Ala	Thr	Ala	Ala
				545					550					555
Val	Gly	Gly	Phe	Thr	Leu	Leu	Val	Ile	Leu	Thr	Leu	Phe	Phe	Leu
				560					565					570
Ile	Thr	Gly	Arg	Cys	Gln	Trp	Tyr	Ile	Lys	Ala	Lys	Met	Lys	Ser
				575					580					585
Glu	Glu	Lys	Arg	Arg	Asn	His	Leu	Gln	Asn	Gly	His	Leu	Arg	Phe
				590					595					600
Pro	Gly	Ile	Lys	Thr	Tyr	Ile	Asp	Pro	Asp	Thr	Tyr	Glu	Asp	Pro
				605					610					615
Ser	Leu	Ala	Val	His	Glu	Phe	Ala	Lys	Glu	Ile	Asp	Pro	Ser	Arg
				620					625					630
Ile	Arg	Ile	Glu	Arg	Val	Ile	Gly	Ala	Gly	Glu	Phe	Gly	Glu	Val
				635					640					645
Cys	Ser	Gly	Arg	Leu	Lys	Thr	Pro	Gly	Lys	Arg	Glu	Ile	Pro	Val
				650					655					660
Ala	Ile	Lys	Thr	Leu	Lys	Gly	Gly	His	Met	Asp	Arg	Gln	Arg	Arg
				665					670					675
Asp	Phe	Leu	Arg	Glu	Ala	Ser	Ile	Met	Gly	Gln	Phe	Asp	His	Pro
				680					685					690
Asn	Ile	Ile	Arg	Leu	Glu	Gly	Val	Val	Thr	Lys	Arg	Ser	Phe	Pro
				695					700					705
Ala	Ile	Gly	Val	Glu	Ala	Phe	Cys	Pro	Ser	Phe	Leu	Arg	Ala	Gly
				710					715					720
Phe	Leu	Asn	Ser	Ile	Gln	Ala	Pro	His	Pro	Val	Pro	Gly	Gly	Gly
				725					730					735
Ser	Leu	Pro	Pro	Arg	Ile	Pro	Ala	Gly	Arg	Pro	Val	Met	Ile	Val
				740					745					750
Val	Glu	Tyr	Met	Glu	Asn	Gly	Ser	Leu	Asp	Ser	Phe	Leu	Arg	Lys
				755					760					765
His	Asp	Gly	His	Phe	Thr	Val	Ile	Gln	Leu	Val	Gly	Met	Leu	Arg
				770					775					780
Gly	Ile	Ala	Ser	Gly	Met	Lys	Tyr	Leu	Ser	Asp	Met	Gly	Tyr	Val
				785					790					795
His	Arg	Asp	Leu	Ala	Ala	Arg	Asn	Ile	Leu	Val	Asn	Ser	Asn	Leu
				800					805					810
Val	Cys	Lys	Val	Ser	Asp	Phe	Gly	Leu	Ser	Arg	Val	Leu	Glu	Asp
				815					820					825
Asp	Pro	Glu	Ala	Ala	Tyr	Thr	Thr	Thr	Gly	Gly	Lys	Ile	Pro	Ile
				830					835					840
Arg	Trp	Thr	Ala	Pro	Glu	Ala	Ile	Ala	Tyr	Arg	Lys	Phe	Ser	Ser
				845					850					855
Ala	Ser	Asp	Ala	Trp	Ser	Tyr	Gly	Ile	Val	Met	Trp	Glu	Val	Met
				860					865					870

-continued

Ser	Tyr	Gly	Glu	Arg	Pro	Tyr	Trp	Glu	Met	Ser	Asn	Gln	Asp	Val
				875					880					885
Ile	Leu	Ser	Ile	Glu	Glu	Gly	Tyr	Arg	Leu	Pro	Ala	Pro	Met	Gly
				890					895					900
Cys	Pro	Ala	Ser	Leu	His	Gln	Leu	Met	Leu	His	Cys	Trp	Gln	Lys
				905					910					915
Glu	Arg	Asn	His	Arg	Pro	Lys	Phe	Thr	Asp	Ile	Val	Ser	Phe	Leu
				920					925					930
Asp	Lys	Leu	Ile	Arg	Asn	Pro	Ser	Ala	Leu	His	Thr	Leu	Val	Glu
				935					940					945
Asp	Ile	Leu	Val	Met	Pro	Glu	Ser	Pro	Gly	Glu	Val	Pro	Glu	Tyr
				950					955					960
Pro	Leu	Phe	Val	Thr	Val	Gly	Asp	Trp	Leu	Asp	Ser	Ile	Lys	Met
				965					970					975
Gly	Gln	Tyr	Lys	Asn	Asn	Phe	Val	Ala	Ala	Gly	Phe	Thr	Thr	Phe
				980					985					990
Asp	Leu	Ile	Ser	Arg	Met	Ser	Ile	Asp	Asp	Ile	Arg	Arg	Ile	Gly
				995					1000					1005
Val	Ile	Leu	Ile	Gly	His	Gln	Arg	Arg	Ile	Val	Ser	Ser	Ile	Gln
				1010					1015					1020
Thr	Leu	Arg	Leu	His	Met	Met	His	Ile	Gln	Glu	Lys	Gly	Phe	His
				1025					1030					1035
Val														

<210> SEQ ID NO 105
<211> LENGTH: 2148
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 105

ggcggcgggc	tgcgcgagc	ggcgccccct	gcagccgcgg	accgaggcag	50
cgcgggcacc	tgccggccga	gcaatgccaa	gtgagtacac	ctatgtgaaa	100
ctgagaagtg	attgctcgag	gccttccctg	caatgggtaca	cccgaagctca	150
aagcaagatg	agaaggccca	gcttggttatt	aaaagacatc	ctcaaatgta	200
cattgcttgt	gtttggagtg	tggtaccttt	atatcctcaa	gttaaattat	250
actactgaag	aatgtgacat	gaaaaaaatg	cattatgtgg	accctgacca	300
tgtaaagaga	gctcagaaat	atgctcagca	agtcttgag	aaggaatgtc	350
gtccaagtt	tgccaagaca	tcaatggcgc	tgttatttga	gcacaggtat	400
agcgtggact	tactcccttt	tgtgcagaag	gcccccaaag	acagtgaagc	450
tgagtccaag	tacgacctc	cttttggtt	ccggaagttc	tccagtaaag	500
tccagaccct	cttgaactc	ttgccagagc	acgacctccc	tgaacacttg	550
aaagccaaga	cctgtcggcg	ctgtgtggtt	attggaagcg	gaggaatact	600
gcacggatta	gaactgggcc	acacctgaa	ccagttcgat	gttgtgataa	650
ggttaaacag	tgaccagtt	gagggatatt	cagaacatgt	tggaataaaa	700
actactataa	ggatgactta	tccagagggc	gcaccactgt	ctgaccttga	750
atattattcc	aatgacttat	ttgttgctgt	tttatttaag	agtgttgatt	800

-continued

tcaactggct tcaagcaatg gtaaaaaagg aaaccctgcc attctgggta	850
cgactcttct tttggaagca ggtggcagaa aaaatcccac tgcagccaaa	900
acatttcagg attttgaatc cagttatcat caaagagact gcctttgaca	950
tccttcagta ctgagagcct cagtcaaggt tctggggccg agataagaac	1000
gtccccacaa tcggtgtcat tgccgttgto ttagccacac atctgtgcga	1050
tgaagtcagt ttggcgggtt ttggatatga cctcaatcaa ccagaaacac	1100
ctttgcacta cttcgacagt caatgcattg ctgctatgaa ctttcagacc	1150
atgcataatg tgacaacgga aaccaagtto ctcttaaagc tggtaaaga	1200
gggagtggtg aaagatctca gtggaggcat tgatcgtgaa ttttgaacac	1250
agaaaacctc agttgaaaat gcaactctaa ctctgagagc tgtttttgac	1300
agccttcttg atgtatttct ccatcctgca gatactttga agtgcagctc	1350
atgtttttaa cttttaattt aaaaacacaa aaaaaatttt agctcttccc	1400
actttttttt tcctatttat ttgaggtcag tgtttgtttt tgcacaccat	1450
tttgtaaatg aaacttaaga attgaattgg aaagacttct caaagagaat	1500
tgatgttaac gatgttgat tgatttttaa gaaagtaatt taatttgtaa	1550
aacttctgct cgtttacct gcacattgaa tacaggtaac taattggaag	1600
gagaggggag gtcactcttt tgatggtggc cctgaacctc attctggttc	1650
cctgctgcgc tgcttggtgt gaccacagga ggatccactc ccaggatgac	1700
gtgtcccgta gctctgctgc tgatactggg totgcgatgc agcggcgtga	1750
ggcctgggct ggttgagaa ggtcacacc cttctctggt ggtctgcctt	1800
ctgtgaaaag actcgagaac caaccaggga agctgtcctg gaggtccctg	1850
gtcggagagg gacatagaat ctgtgacctc tgacaactgt gaagccaccc	1900
tgggctacag aaaccacagt cttccagca attattacaa ttcttgaatt	1950
ccttggggat tttttactgc cttttcaaag cacttaagt ttagatctaa	2000
cgtgttccag tgtctgtctg aggtgactta aaaaatcaga acaaaacttc	2050
tattatccag agtcatggga gagtacacc tttccaggaa taatgttttg	2100
ggaaacactg aaatgaaatc ttcccagtat tataaattgt gtatttaa	2148

<210> SEQ ID NO 106

<211> LENGTH: 362

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 106

Met Arg Arg Pro Ser Leu Leu Leu Lys Asp Ile Leu Lys Cys Thr	
1 5 10 15	
Leu Leu Val Phe Gly Val Trp Ile Leu Tyr Ile Leu Lys Leu Asn	
20 25 30	
Tyr Thr Thr Glu Glu Cys Asp Met Lys Lys Met His Tyr Val Asp	
35 40 45	
Pro Asp His Val Lys Arg Ala Gln Lys Tyr Ala Gln Gln Val Leu	
50 55 60	
Gln Lys Glu Cys Arg Pro Lys Phe Ala Lys Thr Ser Met Ala Leu	
65 70 75	

-continued

Leu	Phe	Glu	His	Arg	Tyr	Ser	Val	Asp	Leu	Leu	Pro	Phe	Val	Gln
			80						85					90
Lys	Ala	Pro	Lys	Asp	Ser	Glu	Ala	Glu	Ser	Lys	Tyr	Asp	Pro	Pro
			95						100					105
Phe	Gly	Phe	Arg	Lys	Phe	Ser	Ser	Lys	Val	Gln	Thr	Leu	Leu	Glu
			110						115					120
Leu	Leu	Pro	Glu	His	Asp	Leu	Pro	Glu	His	Leu	Lys	Ala	Lys	Thr
			125						130					135
Cys	Arg	Arg	Cys	Val	Val	Ile	Gly	Ser	Gly	Gly	Ile	Leu	His	Gly
			140						145					150
Leu	Glu	Leu	Gly	His	Thr	Leu	Asn	Gln	Phe	Asp	Val	Val	Ile	Arg
			155						160					165
Leu	Asn	Ser	Ala	Pro	Val	Glu	Gly	Tyr	Ser	Glu	His	Val	Gly	Asn
			170						175					180
Lys	Thr	Thr	Ile	Arg	Met	Thr	Tyr	Pro	Glu	Gly	Ala	Pro	Leu	Ser
			185						190					195
Asp	Leu	Glu	Tyr	Tyr	Ser	Asn	Asp	Leu	Phe	Val	Ala	Val	Leu	Phe
			200						205					210
Lys	Ser	Val	Asp	Phe	Asn	Trp	Leu	Gln	Ala	Met	Val	Lys	Lys	Glu
			215						220					225
Thr	Leu	Pro	Phe	Trp	Val	Arg	Leu	Phe	Phe	Trp	Lys	Gln	Val	Ala
			230						235					240
Glu	Lys	Ile	Pro	Leu	Gln	Pro	Lys	His	Phe	Arg	Ile	Leu	Asn	Pro
			245						250					255
Val	Ile	Ile	Lys	Glu	Thr	Ala	Phe	Asp	Ile	Leu	Gln	Tyr	Ser	Glu
			260						265					270
Pro	Gln	Ser	Arg	Phe	Trp	Gly	Arg	Asp	Lys	Asn	Val	Pro	Thr	Ile
			275						280					285
Gly	Val	Ile	Ala	Val	Val	Leu	Ala	Thr	His	Leu	Cys	Asp	Glu	Val
			290						295					300
Ser	Leu	Ala	Gly	Phe	Gly	Tyr	Asp	Leu	Asn	Gln	Pro	Arg	Thr	Pro
			305						310					315
Leu	His	Tyr	Phe	Asp	Ser	Gln	Cys	Met	Ala	Ala	Met	Asn	Phe	Gln
			320						325					330
Thr	Met	His	Asn	Val	Thr	Thr	Glu	Thr	Lys	Phe	Leu	Leu	Lys	Leu
			335						340					345
Val	Lys	Glu	Gly	Val	Val	Lys	Asp	Leu	Ser	Gly	Gly	Ile	Asp	Arg
			350						355					360

Glu Phe

<210> SEQ ID NO 107
<211> LENGTH: 1399
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 107

tgacgcgggg cgccagctgc caacttcgcg cgcggagctc cccggcggtg	50
cagtccccgtc ccggcgggcgc gggcggcgatg aagactagcc gccgcggccg	100
agcgctcctg gccgtggccc tgaacctgct ggcgctgctg ttcgccacca	150
ccgctttcct caccacgcac tggtgccagg gcacgcagcg ggtccccaag	200

-continued

```
ccgggctgcg gccagggcgg gcgcgccaac tgccccaact cgggcgcca 250
cgccacggcc aacggcaccc cgccccccgc cgccgccgcc gccgcgcca 300
cgccctcggg gaacggcccc cctggcgggc cgctctacag ctgggagacc 350
ggcgacgacc gcttctctct caggaatttc cacaccggca tctggtactc 400
gtgcgaggag gagctcagcg ggcttggtga aaaatgtcgc agcttcattg 450
acctggcccc ggctcgagg aaaggcctcc tgggaatggc cgccacatg 500
atgtacacgc aggtgttcca ggtcaccgtg agcctcggtc ctgaggactg 550
gagaccccat tcctgggact acgggtggtc cttctgcctg gcgtggggct 600
cctttacctg ctgcatggca gcctctgtca ccacgctcaa ctctacacc 650
aagacggtea ttgagttccg gcacaagcgc aaggtctttg agcagggcta 700
ccgggaagag ccgaccttca tagaccctga ggccatcaag tacttccggg 750
agaggatgga gaagagggac gggagcgagg aggactttca cttagactgc 800
cgccacgaga gataccctgc ccgacaccag ccacacatgg cggattcctg 850
gccccggagc tccgcacagg aagcaccaga gctgaaccga cagtgtggg 900
tcttggggca ctgggtgtga ccaagacctc aacctggccc gcggacctca 950
ggccatcgct ggcaccagcc cctgctgcaa gaccaccaga gtggtgcccc 1000
cagaacctg gcctgtgtgc cgtgaactca gtcagcctgc gtgggagatg 1050
ccaggcctgt cctgcccata gctgcctggg tcccatggcc ttggaaatgg 1100
ggccagggca ggcccaaggg aatgcacagg gctgcacaga gtgactttgg 1150
gacagcagcc ccggactctt gccatcatca catgagccct gctgggcaca 1200
gctgcgatgc caggagacac atggccactg gccactgaat ggctggcacc 1250
cacaagccag tcaggtgccc agaggggcag agccctttgg ggggcagaga 1300
gtggcttcct gaaggagggg gcagtggcgc aggcactgca ggggtgtcac 1350
acagcaggca cacagcaggg gctcaataaa tgcttgttga acttgtttt 1399
```

<210> SEQ ID NO 108

<211> LENGTH: 280

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 108

```
Met Lys Thr Ser Arg Arg Gly Arg Ala Leu Leu Ala Val Ala Leu
 1             5             10             15
Asn Leu Leu Ala Leu Leu Phe Ala Thr Thr Ala Phe Leu Thr Thr
      20             25             30
His Trp Cys Gln Gly Thr Gln Arg Val Pro Lys Pro Gly Cys Gly
      35             40             45
Gln Gly Gly Arg Ala Asn Cys Pro Asn Ser Gly Ala Asn Ala Thr
      50             55             60
Ala Asn Gly Thr Ala Ala Pro Ala Ala Ala Ala Ala Ala Thr
      65             70             75
Ala Ser Gly Asn Gly Pro Pro Gly Gly Ala Leu Tyr Ser Trp Glu
      80             85             90
Thr Gly Asp Asp Arg Phe Leu Phe Arg Asn Phe His Thr Gly Ile
      95             100            105
```

-continued

Trp	Tyr	Ser	Cys	Glu	Glu	Glu	Leu	Ser	Gly	Leu	Gly	Glu	Lys	Cys
				110					115					120
Arg	Ser	Phe	Ile	Asp	Leu	Ala	Pro	Ala	Ser	Glu	Lys	Gly	Leu	Leu
				125					130					135
Gly	Met	Val	Ala	His	Met	Met	Tyr	Thr	Gln	Val	Phe	Gln	Val	Thr
				140					145					150
Val	Ser	Leu	Gly	Pro	Glu	Asp	Trp	Arg	Pro	His	Ser	Trp	Asp	Tyr
				155					160					165
Gly	Trp	Ser	Phe	Cys	Leu	Ala	Trp	Gly	Ser	Phe	Thr	Cys	Cys	Met
				170					175					180
Ala	Ala	Ser	Val	Thr	Thr	Leu	Asn	Ser	Tyr	Thr	Lys	Thr	Val	Ile
				185					190					195
Glu	Phe	Arg	His	Lys	Arg	Lys	Val	Phe	Glu	Gln	Gly	Tyr	Arg	Glu
				200					205					210
Glu	Pro	Thr	Phe	Ile	Asp	Pro	Glu	Ala	Ile	Lys	Tyr	Phe	Arg	Glu
				215					220					225
Arg	Met	Glu	Lys	Arg	Asp	Gly	Ser	Glu	Glu	Asp	Phe	His	Leu	Asp
				230					235					240
Cys	Arg	His	Glu	Arg	Tyr	Pro	Ala	Arg	His	Gln	Pro	His	Met	Ala
				245					250					255
Asp	Ser	Trp	Pro	Arg	Ser	Ser	Ala	Gln	Glu	Ala	Pro	Glu	Leu	Asn
				260					265					270
Arg	Gln	Cys	Trp	Val	Leu	Gly	His	Trp	Val					
				275					280					

<210> SEQ ID NO 109

<211> LENGTH: 2964

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 109

gattaccaag caagaacagc taaaatgaaa gccatcattc atcttactct	50
tcttgctctc ctttctgtaa acacagccac caaccaaggc aactcagctg	100
atgctgtaac aaccacagaa actgcgacta gtggtcctac agtagctgca	150
gctgatacca ctgaaactaa tttccctgaa actgctagca ccacagcaaa	200
tacaccttct ttcccaacag ctacttcacc tgctcccccc ataattagta	250
cacatagttc ctccacaatt cctacacctg ctccccccat aattagtaca	300
catagttcct ccacaattcc tatactact gctgcagaca gtgagtcaac	350
cacaaatgta aattcattag ctacctctga cataatcacc gcttcattctc	400
caaatgatgg attaatacaca atggttcctt ctgaaacaca aagtaacaat	450
gaaatgtccc ccaccacaga agacaatcaa tcatcagggc ctcccactgg	500
caccgcctta ttggagacca gcaccctaaa cagcacaggt cccagcaatc	550
cttgccaaga tgatccctgt gcagataatt cgttatgtgt taagctgcat	600
aatacaagtt tttgcctgtg tttagaaggg tattactaca actottctac	650
atgtaagaaa ggaaaggtat tcctctggaa gatttcagtg acagtatcag	700
aaacatttga ccagaagag aaacattcca tggcctatca agacttgcac	750
agtgaatata ctgacttggt taaagatgta tttggcacat ctgtttatgg	800

-continued

acagactgta attcttactg taagcacatc tctgtcacca agatctgaaa	850
tgcggtgctga tgacaagttt gttaatgtaa caatagtaac aattttggca	900
gaaaccacaa gtgacaatga gaagactgtg actgagaaaa ttaataaagc	950
aattagaagt agctcaagca actttctaaa ctatgatttg acccttcggt	1000
gtgattatta tggctgtaac cagactgcgg atgactgcct caatggttta	1050
gcatgcgatt gcaaactctga cctgcaaagg cctaaccac agagcccttt	1100
ctgcgttgct tccagtctca agtgtcctga tgcctgcaac gcacagcaca	1150
agcaatgctt aataaagaag agtggtaggg cccctgagtg tgcgtgcgtg	1200
cccggctacc aggaagatgc taatgggaac tgccaaaagt gtgcatttg	1250
ctacagtgga ctgcactgta aggacaaatt tcagctgatc ctactattg	1300
tgggcaccat cgctggcatt gtcattctca gcatgataat tgcattgatt	1350
gtcacagcaa gatcaaataa caaacgaag catattgaag aagagaactt	1400
gattgacgaa gactttcaaa atctaaaact gcggtcgaca ggcttcacca	1450
atcttgagc agaaggagc gtccttccta aggtcaggat aacggcctcc	1500
agagacagcc agatgcaaaa tccctattca agccacagca gcatgccccg	1550
ccctgactat tagaatcata agaattgtga acccgccatg gcccccaacc	1600
aatgtacaag ctattattta gagtgtttag aaagactgat ggagaagtga	1650
gcaccagtaa agatctggcc tccggggttt ttcttccatc tgacatctgc	1700
cagcctctct gaattggaagt tgtgaatgtt tgcaacgaat ccagctcact	1750
tgctaaataa gaatctatga cattaaatgt agtagatgct attagcgctt	1800
gtcagagagg tggttttctt caatcagtac aaagtactga gacaatgggt	1850
agggttgttt tcttaattct tttcctggta gggcaacaag aaccatttcc	1900
aatctagagg aaagctcccc agcattgctt gtcctgaggc aaacattgct	1950
cttgagttaa gtgacctaat tcccctggga gacatacgca tcaactgtgg	2000
aggtccgagg ggatgagaag ggataccac catctttcaa gggtcacaag	2050
ctcactctct gacaagtcag aatagggaca ctgcttctat ccctccaatg	2100
gagagattct ggcaaccttt gaacagccca gagcttgcaa cctagcctca	2150
cccaagaaga ctggaaagag acatatctct cagctttttc aggaggcgtg	2200
cctgggaatc caggaaactt ttgatgctaa ttagaaggcc tggactaaaa	2250
atgtccacta tggggtgcac tctacagttt ttgaaatgct aggaggcaga	2300
aggggcagag agtaaaaaac atgacctggt agaagggaaga gaggcaaagg	2350
aaactgggtg gggaggatca attagagagg aggcacctgg gatccacctt	2400
cttccttagg tcccctctc catcagcaaa ggagcacttc tctaactatg	2450
ccctcccgaa gactggctgg gagaaggttt aaaaacaaa aatccaggag	2500
taagagcctt aggtcagttt gaaattggag acaactgtc tggcaaaggg	2550
tgcgagaggg agcttgctct caggagtcca gccgccagc ctcggggtgt	2600
aggtttctga ggtgtgcca tggggcctca gccttctctg gtgacagagg	2650
ctcagctgtg gccaccaaca cacaaccaca cacacacaac cacacacaca	2700

-continued

aatgggggca accacatcca gtacaagctt ttacaaatgt tattagtgtc	2750
cttttttatt tctaatagcct tgcctcttta aaagttatit tatttgttat	2800
tattatttgt tcttgactgt taattgtgaa tggtaatgca ataaagtgcc	2850
tttgttagat ggtgaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	2900
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	2950
aaaaaaaaaa aaaa	2964

<210> SEQ ID NO 110

<211> LENGTH: 512

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 110

Met	Lys	Ala	Ile	Ile	His	Leu	Thr	Leu	Leu	Ala	Leu	Leu	Ser	Val
1				5					10					15
Asn	Thr	Ala	Thr	Asn	Gln	Gly	Asn	Ser	Ala	Asp	Ala	Val	Thr	Thr
				20					25					30
Thr	Glu	Thr	Ala	Thr	Ser	Gly	Pro	Thr	Val	Ala	Ala	Ala	Asp	Thr
				35					40					45
Thr	Glu	Thr	Asn	Phe	Pro	Glu	Thr	Ala	Ser	Thr	Thr	Ala	Asn	Thr
				50					55					60
Pro	Ser	Phe	Pro	Thr	Ala	Thr	Ser	Pro	Ala	Pro	Pro	Ile	Ile	Ser
				65					70					75
Thr	His	Ser	Ser	Ser	Thr	Ile	Pro	Thr	Pro	Ala	Pro	Pro	Ile	Ile
				80					85					90
Ser	Thr	His	Ser	Ser	Ser	Thr	Ile	Pro	Ile	Pro	Thr	Ala	Ala	Asp
				95					100					105
Ser	Glu	Ser	Thr	Thr	Asn	Val	Asn	Ser	Leu	Ala	Thr	Ser	Asp	Ile
				110					115					120
Ile	Thr	Ala	Ser	Ser	Pro	Asn	Asp	Gly	Leu	Ile	Thr	Met	Val	Pro
				125					130					135
Ser	Glu	Thr	Gln	Ser	Asn	Asn	Glu	Met	Ser	Pro	Thr	Thr	Glu	Asp
				140					145					150
Asn	Gln	Ser	Ser	Gly	Pro	Pro	Thr	Gly	Thr	Ala	Leu	Leu	Glu	Thr
				155					160					165
Ser	Thr	Leu	Asn	Ser	Thr	Gly	Pro	Ser	Asn	Pro	Cys	Gln	Asp	Asp
				170					175					180
Pro	Cys	Ala	Asp	Asn	Ser	Leu	Cys	Val	Lys	Leu	His	Asn	Thr	Ser
				185					190					195
Phe	Cys	Leu	Cys	Leu	Glu	Gly	Tyr	Tyr	Tyr	Asn	Ser	Ser	Thr	Cys
				200					205					210
Lys	Lys	Gly	Lys	Val	Phe	Pro	Gly	Lys	Ile	Ser	Val	Thr	Val	Ser
				215					220					225
Glu	Thr	Phe	Asp	Pro	Glu	Glu	Lys	His	Ser	Met	Ala	Tyr	Gln	Asp
				230					235					240
Leu	His	Ser	Glu	Ile	Thr	Ser	Leu	Phe	Lys	Asp	Val	Phe	Gly	Thr
				245					250					255
Ser	Val	Tyr	Gly	Gln	Thr	Val	Ile	Leu	Thr	Val	Ser	Thr	Ser	Leu
				260					265					270
Ser	Pro	Arg	Ser	Glu	Met	Arg	Ala	Asp	Asp	Lys	Phe	Val	Asn	Val

-continued

275										280										285									
Thr	Ile	Val	Thr	Ile	Leu	Ala	Glu	Thr	Thr	Ser	Asp	Asn	Glu	Lys															
				290					295					300															
Thr	Val	Thr	Glu	Lys	Ile	Asn	Lys	Ala	Ile	Arg	Ser	Ser	Ser	Ser															
				305					310					315															
Asn	Phe	Leu	Asn	Tyr	Asp	Leu	Thr	Leu	Arg	Cys	Asp	Tyr	Tyr	Gly															
				320					325					330															
Cys	Asn	Gln	Thr	Ala	Asp	Asp	Cys	Leu	Asn	Gly	Leu	Ala	Cys	Asp															
				335					340					345															
Cys	Lys	Ser	Asp	Leu	Gln	Arg	Pro	Asn	Pro	Gln	Ser	Pro	Phe	Cys															
				350					355					360															
Val	Ala	Ser	Ser	Leu	Lys	Cys	Pro	Asp	Ala	Cys	Asn	Ala	Gln	His															
				365					370					375															
Lys	Gln	Cys	Leu	Ile	Lys	Lys	Ser	Gly	Gly	Ala	Pro	Glu	Cys	Ala															
				380					385					390															
Cys	Val	Pro	Gly	Tyr	Gln	Glu	Asp	Ala	Asn	Gly	Asn	Cys	Gln	Lys															
				395					400					405															
Cys	Ala	Phe	Gly	Tyr	Ser	Gly	Leu	Asp	Cys	Lys	Asp	Lys	Phe	Gln															
				410					415					420															
Leu	Ile	Leu	Thr	Ile	Val	Gly	Thr	Ile	Ala	Gly	Ile	Val	Ile	Leu															
				425					430					435															
Ser	Met	Ile	Ile	Ala	Leu	Ile	Val	Thr	Ala	Arg	Ser	Asn	Asn	Lys															
				440					445					450															
Thr	Lys	His	Ile	Glu	Glu	Glu	Asn	Leu	Ile	Asp	Glu	Asp	Phe	Gln															
				455					460					465															
Asn	Leu	Lys	Leu	Arg	Ser	Thr	Gly	Phe	Thr	Asn	Leu	Gly	Ala	Glu															
				470					475					480															
Gly	Ser	Val	Phe	Pro	Lys	Val	Arg	Ile	Thr	Ala	Ser	Arg	Asp	Ser															
				485					490					495															
Gln	Met	Gln	Asn	Pro	Tyr	Ser	Ser	His	Ser	Ser	Met	Pro	Arg	Pro															
				500					505					510															

Asp Tyr

<210> SEQ ID NO 111
<211> LENGTH: 943
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 111

ctgggacttg gctttctccg gataagcggc ggcaccggcg tcagcgatga	50
ccgtgcagag actcgtggcc gcggccgtgc tggtagccct ggtctcactc	100
atcctcaaca acgtggcggc cttcacctcc aactgggtgt gccagacgct	150
ggaggatggg cgcaggcgca gcgtggggct gtggaggtcc tgctggctgg	200
tggacagagc ccggggaggg ccgagccctg gggccagagc cggccagggtg	250
gacgcacatg actgtgaggc gctgggctgg ggctccgagg cagccggctt	300
ccaggagtcc cgaggcaccg tcaaactgca gtctgacatg atgcgcgcct	350
gcaacctggt ggccacggcc gcgctcaccg caggccagct caccttcttc	400
ctggggctgg tgggcctgcc cctgctgtca cccgacgcc cgtgctggga	450
ggaggccatg gccgctgcat tccaactggc gagttttgtc ctggtcatcg	500

-continued

ggctcgtgac tttctacaga attggcccat acaccaacct gtccctgggtcc 550
tgctaccta acattggcgc ctgccttctg gccacgctgg cggcagccat 600
gtcatctgg aacattctcc acaagagga ggactgcatg gccccccggg 650
tgattgtcat cagccgctcc ctgacagcgc gctttcgccg tgggctggac 700
aatgactacg tggagtcacc atgctgagtc gcccttctca gcgtccatc 750
aacgcacacc tgctatcgtg gaacagccta gaaaccaagg gactccacca 800
ccaagtcaact tcccctgctc gtgcagaggc acgggatgag tctgggtgac 850
ctctgcgccca tgcgtgcgag acacgtgtgc gtttactgtt atgtcggta 900
tatgtctgta cgtgtcgtgg gccaacctcg ttctgcctcc agc 943

<210> SEQ ID NO 112

<211> LENGTH: 226

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 112

Met Thr Val Gln Arg Leu Val Ala Ala Ala Val Leu Val Ala Leu
1 5 10 15
Val Ser Leu Ile Leu Asn Asn Val Ala Ala Phe Thr Ser Asn Trp
20 25 30
Val Cys Gln Thr Leu Glu Asp Gly Arg Arg Arg Ser Val Gly Leu
35 40 45
Trp Arg Ser Cys Trp Leu Val Asp Arg Thr Arg Gly Gly Pro Ser
50 55 60
Pro Gly Ala Arg Ala Gly Gln Val Asp Ala His Asp Cys Glu Ala
65 70 75
Leu Gly Trp Gly Ser Glu Ala Ala Gly Phe Gln Glu Ser Arg Gly
80 85 90
Thr Val Lys Leu Gln Phe Asp Met Met Arg Ala Cys Asn Leu Val
95 100 105
Ala Thr Ala Ala Leu Thr Ala Gly Gln Leu Thr Phe Leu Leu Gly
110 115 120
Leu Val Gly Leu Pro Leu Leu Ser Pro Asp Ala Pro Cys Trp Glu
125 130 135
Glu Ala Met Ala Ala Ala Phe Gln Leu Ala Ser Phe Val Leu Val
140 145 150
Ile Gly Leu Val Thr Phe Tyr Arg Ile Gly Pro Tyr Thr Asn Leu
155 160 165
Ser Trp Ser Cys Tyr Leu Asn Ile Gly Ala Cys Leu Leu Ala Thr
170 175 180
Leu Ala Ala Ala Met Leu Ile Trp Asn Ile Leu His Lys Arg Glu
185 190 195
Asp Cys Met Ala Pro Arg Val Ile Val Ile Ser Arg Ser Leu Thr
200 205 210
Ala Arg Phe Arg Arg Gly Leu Asp Asn Asp Tyr Val Glu Ser Pro
215 220 225

Cys

<210> SEQ ID NO 113

-continued

<211> LENGTH: 1389
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 113

gactttacca ctactcgcta tagagccctg gtcaagttct ctccacctct 50
ctatctatgt ctcagtttct tcatctgtaa catcaaatga ataataatac 100
caatctccta gacttcataa gaggattaac aaagacaaaa tatgggaaaa 150
acataacatg gcgtcccata attattagat cttattattg aactaaaaat 200
ggcattaaaa ttaccaaaag gaagacagca tctgtttcct ctttggtcct 250
gagctggtta aaaggaacac tggttgcctg aacagtcaca cttgcaacca 300
tgatgcctaa acattgcttt ctaggcttcc tcatcagttt ctctcttact 350
ggtgtagcag gaactcagtc aacgcatgag tctctgaagc ctgagagggt 400
acaattttcag tcccgaaatt ttcacaacat tttgcaatgg cagcctggga 450
gggcacttac tggcaacagc agtgtctatt ttgtgcagta caaaatatat 500
ggacagagac aatggaaaaa taaagaagac tgttggggta ctcaagaact 550
ctcttgtgac cttaccagtg aaacctcaga catacaggaa ccttattacg 600
ggaggggtgag ggcggcctcg gctgggagct actcagaatg gagcatgacg 650
ccgcggttca ctccctgggtg ggaaacaaaa atagatcctc cagtcatgaa 700
tataacccaa gtcaatggct ctttgttggg aattctccat gtcctcaatt 750
taccatatag ataccaaaag gaaaaaatg tatctataga agattactat 800
gaactactat accgagtttt tataattaac aattcactag aaaaggagca 850
aaaggtttat gaaggggctc acagagcggg tgaaattgaa gctctaacac 900
cacactccag ctactgtgta gtggctgaaa tatacagcc catggttagac 950
agaagaagtc agagaagtga agagagatgt gtggaaattc catgacttgt 1000
ggaattttggc attcagcaat gtggaaattc taaagctccc tgagaacagg 1050
atgactcgtg tttgaaggat cttattttaa attgtttttg tattttctta 1100
aagcaatatt cactgttaca ccttggggac ttctttgttt acccattctt 1150
ttatccttta tatttcattt gtaaaactata tttgaacgac attccccccg 1200
aaaaattgaa atgtaaagat gaggcagaga ataaagtgtt ctatgaaatt 1250
cagaacttta tttctgaatg taacatccct aataacaacc ttcattcttc 1300
taatacagca aaataaaaat ttaacaacca aggaatagta tttaagaaaa 1350
tgttgaaata atttttttaa aatagcatta cagactgag 1389

<210> SEQ ID NO 114
<211> LENGTH: 231
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 114

Met Met Pro Lys His Cys Phe Leu Gly Phe Leu Ile Ser Phe Phe
1 5 10 15

Leu Thr Gly Val Ala Gly Thr Gln Ser Thr His Glu Ser Leu Lys
20 25 30

-continued

Pro	Gln	Arg	Val	Gln	Phe	Gln	Ser	Arg	Asn	Phe	His	Asn	Ile	Leu
			35						40					45
Gln	Trp	Gln	Pro	Gly	Arg	Ala	Leu	Thr	Gly	Asn	Ser	Ser	Val	Tyr
			50						55					60
Phe	Val	Gln	Tyr	Lys	Ile	Tyr	Gly	Gln	Arg	Gln	Trp	Lys	Asn	Lys
			65						70					75
Glu	Asp	Cys	Trp	Gly	Thr	Gln	Glu	Leu	Ser	Cys	Asp	Leu	Thr	Ser
			80						85					90
Glu	Thr	Ser	Asp	Ile	Gln	Glu	Pro	Tyr	Tyr	Gly	Arg	Val	Arg	Ala
			95						100					105
Ala	Ser	Ala	Gly	Ser	Tyr	Ser	Glu	Trp	Ser	Met	Thr	Pro	Arg	Phe
			110						115					120
Thr	Pro	Trp	Trp	Glu	Thr	Lys	Ile	Asp	Pro	Pro	Val	Met	Asn	Ile
			125						130					135
Thr	Gln	Val	Asn	Gly	Ser	Leu	Leu	Val	Ile	Leu	His	Ala	Pro	Asn
			140						145					150
Leu	Pro	Tyr	Arg	Tyr	Gln	Lys	Glu	Lys	Asn	Val	Ser	Ile	Glu	Asp
			155						160					165
Tyr	Tyr	Glu	Leu	Leu	Tyr	Arg	Val	Phe	Ile	Ile	Asn	Asn	Ser	Leu
			170						175					180
Glu	Lys	Glu	Gln	Lys	Val	Tyr	Glu	Gly	Ala	His	Arg	Ala	Val	Glu
			185						190					195
Ile	Glu	Ala	Leu	Thr	Pro	His	Ser	Ser	Tyr	Cys	Val	Val	Ala	Glu
			200						205					210
Ile	Tyr	Gln	Pro	Met	Leu	Asp	Arg	Arg	Ser	Gln	Arg	Ser	Glu	Glu
			215						220					225
Arg	Cys	Val	Glu	Ile	Pro									
			230											

<210> SEQ ID NO 115
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 115

tgtaaaacga cggccagtta aatagacctg caattattaa tct

43

<210> SEQ ID NO 116
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 116

caggaaacag ctatgaccac ctgcacacct gcaaatccat t

41

What is claimed is:

1. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of the amino acid sequence shown in FIG. 2 (SEQ ID NO:2), FIG. 4 (SEQ ID NO:4), FIG. 6 (SEQ ID NO:6), FIG. 8 (SEQ ID NO:8), FIG. 10 (SEQ ID NO:10), FIG. 12 (SEQ

ID NO:12), FIG. 14 (SEQ ID NO:14), FIG. 16 (SEQ ID NO:16), FIG. 18 (SEQ ID NO:18), FIG. 20 (SEQ ID NO:20), FIG. 22 (SEQ ID NO:22), FIG. 24 (SEQ ID NO:24), FIG. 26 (SEQ ID NO:26), FIG. 28 (SEQ ID NO:28), FIG. 30 (SEQ ID NO:30), FIG. 32 (SEQ ID NO:32), FIG. 34 (SEQ ID NO:34), FIG. 36 (SEQ ID NO:36), FIG. 38 (SEQ ID NO:38), FIG. 40 (SEQ ID

NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) and **FIG. 114** (SEQ ID NO:114).

2. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the nucleotide sequence shown in **FIG. 1** (SEQ ID NO:1), **FIG. 3** (SEQ ID NO:3), **FIG. 5** (SEQ ID NO:5), **FIG. 7** (SEQ ID NO:7), **FIG. 9** (SEQ ID NO:9), **FIG. 11** (SEQ ID NO:11), **FIG. 13** (SEQ ID NO:13), **FIG. 15** (SEQ ID NO:15), **FIG. 17** (SEQ ID NO:17), **FIG. 19** (SEQ ID NO:19), **FIG. 21** (SEQ ID NO:21), **FIG. 23** (SEQ ID NO:23), **FIG. 25** (SEQ ID NO:25), **FIG. 27** (SEQ ID NO:27), **FIG. 29** (SEQ ID NO:29), **FIG. 31** (SEQ ID NO:31), **FIG. 33** (SEQ ID NO:33), **FIG. 35** (SEQ ID NO:35), **FIG. 37** (SEQ ID NO:37), **FIG. 39** (SEQ ID NO:39), **FIG. 41** (SEQ ID NO:41), **FIG. 43** (SEQ ID NO:43), **FIG. 45** (SEQ ID NO:45), **FIG. 47** (SEQ ID NO:47), **FIG. 49** (SEQ ID NO:49), **FIG. 51** (SEQ ID NO:51), **FIG. 53** (SEQ ID NO:53), **FIG. 55** (SEQ ID NO:55), **FIG. 57** (SEQ ID NO:57), **FIG. 59** (SEQ ID NO:59), **FIG. 61** (SEQ ID NO:61), **FIG. 63** (SEQ ID NO:63), **FIG. 65** (SEQ ID NO:65), **FIG. 67** (SEQ ID NO:67), **FIG. 69** (SEQ ID NO:69), **FIG. 71** (SEQ ID NO:71), **FIG. 73** (SEQ ID NO:73), **FIG. 75** (SEQ ID NO:75), **FIG. 77** (SEQ ID NO:77), **FIG. 79** (SEQ ID NO:79), **FIG. 81** (SEQ ID NO:81), **FIG. 83** (SEQ ID NO:83), **FIG. 85** (SEQ ID NO:85), **FIG. 87** (SEQ ID NO:87), **FIG. 89** (SEQ ID NO:89), **FIG. 91** (SEQ ID NO:91), **FIG. 93** (SEQ ID NO:93), **FIGS. 95A-95B** (SEQ ID NO:95), **FIG. 97** (SEQ ID NO:97), **FIG. 99** (SEQ ID NO:99), **FIG. 101** (SEQ ID NO:101), **FIG. 103** (SEQ ID NO:103), **FIG. 105** (SEQ ID NO:105), **FIG. 107** (SEQ ID NO:107), **FIG. 109** (SEQ ID NO:109), **FIG. 111** (SEQ ID NO:111) and **FIG. 113** (SEQ ID NO:113).

3. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the full-length coding sequence of the nucleotide sequence shown in **FIG. 1** (SEQ ID NO:1), **FIG. 3** (SEQ ID NO:3), **FIG. 5** (SEQ ID NO:5), **FIG. 7** (SEQ ID NO:7), **FIG. 9** (SEQ ID NO:9), **FIG. 11** (SEQ ID NO:11), **FIG. 13** (SEQ ID NO:13), **FIG. 15** (SEQ ID NO:15), **FIG. 17** (SEQ ID NO:17), **FIG. 19** (SEQ ID NO:19), **FIG. 21** (SEQ ID NO:21), **FIG. 23** (SEQ ID NO:23), **FIG. 25** (SEQ ID NO:25), **FIG. 27** (SEQ ID NO:27), **FIG. 29** (SEQ ID NO:29), **FIG. 31** (SEQ ID NO:31), **FIG. 33** (SEQ ID NO:33), **FIG. 35** (SEQ ID NO:35), **FIG. 37** (SEQ ID NO:37), **FIG. 39** (SEQ ID NO:39), **FIG. 41** (SEQ ID NO:41), **FIG. 43** (SEQ ID NO:43), **FIG. 45** (SEQ ID

NO:45), **FIG. 47** (SEQ ID NO:47), **FIG. 49** (SEQ ID NO:49), **FIG. 51** (SEQ ID NO:51), **FIG. 53** (SEQ ID NO:53), **FIG. 55** (SEQ ID NO:55), **FIG. 57** (SEQ ID NO:57), **FIG. 59** (SEQ ID NO:59), **FIG. 61** (SEQ ID NO:61), **FIG. 63** (SEQ ID NO:63), **FIG. 65** (SEQ ID NO:65), **FIG. 67** (SEQ ID NO:67), **FIG. 69** (SEQ ID NO:69), **FIG. 71** (SEQ ID NO:71), **FIG. 73** (SEQ ID NO:73), **FIG. 75** (SEQ ID NO:75), **FIG. 77** (SEQ ID NO:77), **FIG. 79** (SEQ ID NO:79), **FIG. 81** (SEQ ID NO:81), **FIG. 83** (SEQ ID NO:83), **FIG. 85** (SEQ ID NO:85), **FIG. 87** (SEQ ID NO:87), **FIG. 89** (SEQ ID NO:89), **FIG. 91** (SEQ ID NO:91), **FIG. 93** (SEQ ID NO:93), **FIGS. 95A-95B** (SEQ ID NO:95), **FIG. 97** (SEQ ID NO:97), **FIG. 99** (SEQ ID NO:99), **FIG. 101** (SEQ ID NO:101), **FIG. 103** (SEQ ID NO:103), **FIG. 105** (SEQ ID NO:105), **FIG. 107** (SEQ ID NO:107), **FIG. 109** (SEQ ID NO:109), **FIG. 111** (SEQ ID NO:111) and **FIG. 113** (SEQ ID NO:113).

4. Isolated nucleic acid having at least 80% nucleic acid sequence identity to the full-length coding sequence of the DNA deposited under any ATCC accession number shown in Table 7.

5. A vector comprising the nucleic acid of claim 1.

6. A host cell comprising the vector of claim 5.

7. The host cell of claim 6, wherein said cell is a CHO cell.

8. The host cell of claim 6, wherein said cell is an *E. coli*.

9. The host cell of claim 6, wherein said cell is a yeast cell.

10. A process for producing a PRO polypeptide comprising culturing the host cell of claim 6 under conditions suitable for expression of said PRO polypeptide and recovering said PRO polypeptide from the cell culture.

11. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence selected from the group consisting of the amino acid sequence shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) and **FIG. 114** (SEQ ID NO:114).

12. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence encoded by the full-length coding sequence of the DNA deposited under any ATCC accession number shown in Table 7.

13. A chimeric molecule comprising a polypeptide according to claim 11 fused to a heterologous amino acid sequence.

14. The chimeric molecule of claim 13, wherein said heterologous amino acid sequence is an epitope tag sequence.

15. The chimeric molecule of claim 13, wherein said heterologous amino acid sequence is a Fc region of an immunoglobulin.

16. An antibody which specifically binds to a polypeptide according to claim 11.

17. The antibody of claim 16, wherein said antibody is a monoclonal antibody, a humanized antibody or a single-chain antibody.

18. Isolated nucleic acid having at least 80% nucleic acid sequence identity to:

(a) a nucleotide sequence encoding the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), lacking its associated signal peptide;

(b) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID

NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), with its associated signal peptide; or

(c) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), lacking its associated signal peptide.

19. An isolated polypeptide having at least 80% amino acid sequence identity to:

(a) an amino acid sequence of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34**

(SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), lacking its associated signal peptide;

(b) an amino acid sequence of an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24** (SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), with its associated signal peptide; or

(c) an amino acid sequence of an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:6), **FIG. 8** (SEQ ID NO:8), **FIG. 10** (SEQ ID NO:10), **FIG. 12** (SEQ ID NO:12), **FIG. 14** (SEQ ID NO:14), **FIG. 16** (SEQ ID NO:16), **FIG. 18** (SEQ ID NO:18), **FIG. 20** (SEQ ID NO:20), **FIG. 22** (SEQ ID NO:22), **FIG. 24**

(SEQ ID NO:24), **FIG. 26** (SEQ ID NO:26), **FIG. 28** (SEQ ID NO:28), **FIG. 30** (SEQ ID NO:30), **FIG. 32** (SEQ ID NO:32), **FIG. 34** (SEQ ID NO:34), **FIG. 36** (SEQ ID NO:36), **FIG. 38** (SEQ ID NO:38), **FIG. 40** (SEQ ID NO:40), **FIG. 42** (SEQ ID NO:42), **FIG. 44** (SEQ ID NO:44), **FIG. 46** (SEQ ID NO:46), **FIG. 48** (SEQ ID NO:48), **FIG. 50** (SEQ ID NO:50), **FIG. 52** (SEQ ID NO:52), **FIG. 54** (SEQ ID NO:54), **FIG. 56** (SEQ ID NO:56), **FIG. 58** (SEQ ID NO:58), **FIG. 60** (SEQ ID NO:60), **FIG. 62** (SEQ ID NO:62), **FIG. 64** (SEQ ID NO:64), **FIG. 66** (SEQ ID NO:66), **FIG. 68** (SEQ ID NO:68), **FIG. 70** (SEQ ID NO:70), **FIG. 72** (SEQ ID NO:72), **FIG. 74** (SEQ ID NO:74), **FIG. 76** (SEQ ID NO:76), **FIG. 78** (SEQ ID NO:78), **FIG. 80** (SEQ ID NO:80), **FIG. 82** (SEQ ID NO:82), **FIG. 84** (SEQ ID NO:84), **FIG. 86** (SEQ ID NO:86), **FIG. 88** (SEQ ID NO:88), **FIG. 90** (SEQ ID NO:90), **FIG. 92** (SEQ ID NO:92), **FIG. 94** (SEQ ID NO:94), **FIG. 96** (SEQ ID NO:96), **FIG. 98** (SEQ ID NO:98), **FIG. 100** (SEQ ID NO:100), **FIG. 102** (SEQ ID NO:102), **FIG. 104** (SEQ ID NO:104), **FIG. 106** (SEQ ID NO:106), **FIG. 108** (SEQ ID NO:108), **FIG. 110** (SEQ ID NO:110), **FIG. 112** (SEQ ID NO:112) or **FIG. 114** (SEQ ID NO:114), lacking its associated signal peptide.

20. A method for stimulating the proliferation or differentiation of chondrocyte cells, said method comprising contacting said cells with a PRO6018 polypeptide, wherein the proliferation or differentiation of said cells is stimulated.

21. A method for stimulating the proliferation of human microvascular endothelial cells, said method comprising contacting said cells with a PRO1313, PRO20080 or PRO21383 polypeptide, wherein the proliferation of said cells is stimulated.

24. A method for inhibiting the proliferation of human microvascular endothelial cells, said method comprising contacting said cells with a PRO6071, PRO4487 or PRO6006 polypeptide, wherein the proliferation of said cells is inhibited.

25. A method for detecting the presence of tumor in a mammal, said method comprising comparing the level of expression of any PRO polypeptide shown in Table 8 in (a) a test sample of cells taken from said mammal and (b) a control sample of normal cells of the same cell type, wherein a higher level of expression of said PRO polypeptide in the test sample as compared to the control sample is indicative of the presence of tumor in said mammal.

26. The method of claim 25, wherein said tumor is lung tumor, colon tumor, breast tumor, prostate tumor, rectal tumor, kidney tumor or liver tumor.

27. A method for inducing endothelial cell tube formation comprising administering to the endothelial cell a PRO281, PRO1560, PRO189, PRO4499, PRO6308, PRO6000, PRO10275, PRO21207, PRO20933 or PRO34274 polypeptide, or agonist thereof, wherein tube formation in said endothelial cell is induced.

28. An oligonucleotide probe derived from any of the nucleotide sequences shown in the accompanying figures.

* * * * *