

(19) **DANMARK**

(10) **DK/EP 3406258 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 38/14 (2006.01)** **A 61 K 31/7012 (2006.01)** **A 61 K 31/715 (2006.01)**
A 61 K 36/05 (2006.01) **A 61 K 36/21 (2006.01)** **A 61 K 36/315 (2006.01)**
A 61 K 36/48 (2006.01) **A 61 K 36/898 (2006.01)** **A 61 K 38/10 (2006.01)**
A 61 K 38/16 (2006.01) **A 61 P 19/06 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2022-05-30**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2022-03-02**
- (86) Europæisk ansøgning nr.: **17743673.0**
- (86) Europæisk indleveringsdag: **2017-01-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2018-11-28**
- (86) International ansøgning nr.: **CN2017071864**
- (87) Internationalt publikationsnr.: **WO2017129058**
- (30) Prioritet: **2016-01-29 CN 201610063691**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Shandong Zhonghai Pharmaceutical CO., Ltd., 10298 East Jiankang Street, Weifang, Shandong Province, Kina**
- (72) Opfinder: **CHENG, Qian, No.10298 East Jiankang Street, High Tech Zone, Weifang, Shandong 261041, Kina**
CHENG, Long, No.10298 East Jiankang Street, High Tech Zone, Weifang, Shandong 261041, Kina
XU, Baozhen, No.10298 East Jiankang Street, High Tech Zone, Weifang, Shandong 261041, Kina
- (74) Fuldmægtig i Danmark: **Novagraaf Brevets, Bâtiment O2, 2 rue Sarah Bernhardt CS90017, F-92665 Asnières-sur-Seine cedex, Frankrig**
- (54) Benævnelse: **MEDIKAMENT TIL ANVENDELSE TIL BEHANDLING AF URINSYREGIGT**
- (56) Fremdragne publikationer:
EP-A1- 1 506 785
CN-A- 1 045 586
CN-A- 101 002 873
CN-A- 102 077 933
CN-A- 105 709 207
JP-A- 2010 235 517
US-A1- 2010 048 507
CARDOSO SUSANA M ET AL: "Bioproducts from Seaweeds: A Review with Special Focus on the Iberian Peninsula", CURRENT ORGANIC CHEM., vol. 18, no. 7, 1 April 2014 (2014-04-01), pages 896-917, XP009190894,
GO, H. et al.: "A Glycoprotein from Laminaria Japonica Induces Apoptosis in HT-29 Colon Cancer Cells", TOXICOLOGY IN VITRO, vol. 24, no. 6, 30 September 2010 (2010-09-30), pages 1546-1553, XP055532530,

Fortsættes ...

DESCRIPTION

TECHNICAL FIELD

[0001] The present invention relates to a medicament for use in treating gout, and belongs to the technical field of medical technology.

BACKGROUND TECHNOLOGY

[0002] Gout is a disease caused by a disorder of purine metabolism. Its clinical manifestations are characteristic acute arthritis, tophi formation, and gouty chronic arthritis caused by hyperuricemia and urate crystal deposition, patients may experience urate nephropathy and uric acid-related urinary calculus; in severe cases, patients may experience joint disability and renal insufficiency. The prerequisite for the onset of gout is hyperuricemia. Therefore, hyperuricemia is the most important biochemical basis for gout, followed by inflammatory response caused by urate deposition. With the improvement of people's living standards, the incidence of hyperuricemia and primary gout has been increasing on a year-to-year basis due to the changes in dietary structure, especially the increase in alcohol, especially beer, as well as the decrease in physical activity; the age of onset is increasingly younger. Since the uric acid is supersaturated in the blood, in addition to direct damage to bones and joints, it will also deposit in the kidneys, causing kidney disease, which poses a greater risk to human health and has drawn widespread attention in the medical community.

[0003] At present, western medicine treats hyperuricemia and gout by stages. Western medicine usually uses non-steroidal anti-inflammatory drugs (such as etoricoxib, celecoxib), uric acid lowering drugs (such as allopurinol, febuxostat, probenecid and benzbromarone), glucocorticoids and colchicine etc. during the acute phase; as for the treatments during the intermittent and chronic stages, patients are often treated with diet control and uric-acid-lowering drugs; although nonsteroidal anti-inflammatory drugs can quickly control the polar attack of gout, the drug only exerts an analgesic role during the onset stage of gout, but a relieve in pain does not cure the disease, the treatment only focuses on the relieving of symptoms and ignores the elimination of cause root, and these drugs have serious toxicity.

[0004] Patent: ASTER GLEHNI EXTRACTS, FRACTIONS OR COMPOUNDS ISOLATED THEREFROM FOR THE TREATMENT OR PREVENTION OF HYPERURICEMIA OR GOUT Patent Number is US2015337001 (A1), it is disclosed that: the invention relates to an aster glehni extract capable of preventing and improving hyperuricemia or gout, with the effect of inhibiting uric acid production and reducing its concentration.

[0005] Patent: THE USE OF BUCILLAMINE IN THE TREATMENT OF GOUT, Patent No. US2015306054 (A1), it is disclosed that: disclosed are pharmaceutical compositions

comprising, bucillamine, including bucillamine and allopurinol or colchicine, or pharmaceutically acceptable salts or solvates thereof, together with one or more pharmaceutically acceptable carriers, diluents and excipients. It is disclosed that methods for use of the said compositions in the treatment of gout and metabolic syndrome are also disclosed.

[0006] Patent: a traditional Chinese medicine composition for treating gout, Patent No. CN201310601617.4, it is disclosed that: A traditional Chinese medicine composition for treating gout relates to a traditional Chinese medicine composition. The invention aims to solve the problem that the western medicine has many side effects on the human body in gout treatment and the treatment only focuses on the relieving of symptoms and ignores the elimination of cause root. The traditional Chinese medicine composition for treating gout hereof comprises, by weight, 5-10 portions of sophora flavescens, 10-15 portions of epimedium, 5-10 portions of prepared aconite root, 4-8 portions of ligusticum wallichii, 8-12 portions of pyrrosia lingua, 6-9 parts of papaya and 3-6 portions of fructus amomi. The traditional Chinese medicine composition for treating gout has small side effects on the human body, and is capable of relieving the symptoms and eliminating cause root, with an overall clinical response rate of 96.7%. The invention is used for the treatment of gout.

[0007] JP 2010 235517 A discloses a material for prophylaxis of lifestyle-related diseases containing the protease-treated material of chlorella as an active ingredient. The material for the prophylaxis of lifestyle-related diseases ameliorates any of adiposis, hyperlipemia (a neutral fat or a free fatty acid), or hyperglycemia, and has any of an insulin sensitivity-ameliorating action, a hepatic function (GPT)-ameliorating action, or a renal function (uric acid)-ameliorating action. Besides, as lifestyle-related diseases can be prevented by small amount ingestion, the material is suitable for daily ingestion for long periods, particularly as a food.

[0008] EP 1 506 785 A1 discloses a composition for treating or preventing hyperuricemia which contains a chondroitin sulfate protein complex as an effective ingredient. Because of having an effect of lowering serum uric acid level, this composition makes it possible to provide a novel method of the treatment, the prevention and the relief of gout.

[0009] CN 101 002 873 A discloses special dietary seafood special for hyperuricemia patients. The seafood is characterized by being prepared by taking food-derived marine bioactive polysaccharides as main raw materials, adding algae powder and mixing the food-derived marine bioactive polysaccharides and the algae powder with the auxiliary materials including nutritional amino acids, oligosaccharides, vitamins and minerals. The seafood can be powdery or granular and is eaten after being dissolved in warm water. The seafood is easy to dissolve, has maximum biological absorbance and maximum bioavailability in human bodies, provides marine active factors and nutritional components for human bodies, supplements various vitamins and minerals essential to human bodies, is beneficial to improving the nutritional status of the hyperuricemia patients and can assist in reducing the uric acid level and controlling the hyperuricemia level, improving the immune of the human bodies and preventing gout through synergy of such factors as bioactive components in the food-derived marine

bioactive polysaccharides, the algae powder, the nutritional amino acids and the like, thus being suitable for the hyperuricemia patients.

[0010] US 2010/048507 A1 discloses a food/drink or pharmaceutical composition for oral administration for improving an acidic urine, comprising fucoidan or a fucoidan-containing material as an active ingredient, characterized in that a urinal pH is persistently increased; use of fucoidan or a fucoidan-containing material for production of the food/drink or the pharmaceutical composition for oral administration; and a method for improving an acidic urine by persistently increasing a urinal pH of a patient, which comprises orally administering fucoidan or a fucoidan-containing material to a patient.

[0011] The mechanism of action of traditional Chinese medicine for treating gout is complicated, and the clinical efficacy remains unclear. At present, there is no clinically recommended Chinese patent medicine for treating gout, and domestic gout patients western practically rely on western medicine for treatment.

CONTENT OF INVENTION

[0012] The present invention provides a medicament for use in treating gout in order to solve the deficiencies in the prior art for the purpose to achieve the following objectives hereof:

1. (1) The medicament hereof has the characteristics of high efficiency, safety and no side effects, and has good therapeutic effect on human gout, with a total effective rate of 98% and a low recurrence rate.
2. (2) The medicament hereof reduces the content of urine protein in gout rat model;
3. (3) The medicament hereof reduces the content of creatinine in gout rat model;
4. (4) The medicament hereof reduces the content of urea nitrogen in gout rat model;
5. (5) The medicament hereof reduces the content of blood uric acid in hyperuricemia mice model.

[0013] The present invention is defined in the appended claims.

[0014] Compared with the prior art, the advantages of the present invention are:

1. (1) The medicament hereof has significant therapeutic effect on human gout, with a total effective rate of 98%, of which the rate of well-healed is 85%, the rate of marked efficacy is 13% and the rate of inefficacy is only 2%. As for the healed patients, the recurrence rate in 5 years is only 1%;
2. (2) The medicament hereof reduces the content of urine protein in gout rat model; after four weeks administration of the gout rat model, the urine protein value of the rat is 50.89-168.78 mg/L; the urine protein value of the rat in the model control group is 364.83 mg/L;

3. (3) The medicament hereof reduces the content of creatinine in gout rat model; after four weeks administration of gout rat model, the creatinine value of the rat is 152.45-172.82 mg/L; the creatinine value of the rat in the model control group is 288.01 mmol/L;
4. (4) The medicament hereof reduces the content of urea nitrogen in gout rat model; after four weeks administration of gout rat model, the urea nitrogen value of the rat is 11.58-15.64 mmol/L; the urea nitrogen value in the model control group is 56.07 mmol/L;
5. (5) The medicament hereof reduces the content of blood uric acid in hyperuricemia mice model; the mice administered with the medicament in the invention has a blood uric acid value of 147.89-160.5 μ mol/L; the mice in the model control group have a blood uric acid value of 250.4 μ mol/L.

SPECIFIC EMBODIMENTS

[0015] Herein disclosed is:

Embodiment 1 A medicament for use in treating gout (not covered by the claims)

[0016] Wherein the medicament is a marine algal glycoprotein;

The marine algal glycoprotein comprises, by weight content, 1% sugar and 99% protein;

The molecular weight is 0.2 kDa;

The said sugar is a polysaccharide;

The said marine algae is blue-green algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 2 A medicament for use in treating gout (not covered by the claims)

[0017]

Wherein the medicament is a marine algal glycoprotein;

The marine algal glycoprotein comprises, by weight content, 9% sugar and 88% protein;

The molecular weight is 3 kDa;

The said sugar is a polysaccharide;

The said marine algae is blue-green algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 3 A medicament for use in treating gout (not covered by the claims)

[0018]

Wherein the medicament is a marine algal glycoprotein;

The marine algal glycoprotein comprises, by weight content, 16% sugar and 80% protein;

The molecular weight is 25 kDa;

The said sugar is a polysaccharide;

The said marine algae is chlorella;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 4 A medicament for use in treating gout (not covered by the claims)

[0019]

Wherein the medicament is a marine algal glycoprotein;

The marine algal glycoprotein comprises, by weight content, 30% sugar and 70% protein;

The molecular weight is 40 kDa;

The said sugar is a polysaccharide;

The said marine algae is red algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 5 A medicament for use in treating gout (not covered by the claims)

[0020]

Wherein the medicament is a marine algal glycoprotein;

The marine algal glycoprotein comprises, by weight content, 50% sugar and 50% protein;

The molecular weight is 100 kDa;

The said sugar is a polysaccharide;

The said marine algae is brown algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 6 A medicament for use in treating gout (not covered by the claims)

[0021]

Wherein the medicament is a marine algal glycoprotein;

The marine algal glycoprotein comprises, by weight content, 99% sugar and 1% protein;

The molecular weight is 3000 kDa;

The said sugar is a polysaccharide;

The said marine algae is gold algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

The glycoprotein said in these above embodiments 1-6 further includes a pigment; the said pigment is a natural pigment contained in algal substances;

These above embodiments 1-6 could be summarized as:

A medicament for use in treating gout**[0022]**

The said medicament is a glycoprotein;

The glycoprotein comprises, by weight content, 1-99% sugar and 1-99% protein;

The molecular weight is 0.2-30000 kDa;

The said sugar is a polysaccharide;

The said medicament includes synthetic glycoproteins and synthetic polysaccharides and proteins.

[0023] The said protein comprises 20 kinds of amino acids and 8 kinds of synthetic amino acids; The preparation method of the said medicament: the glycoprotein is prepared into capsules and tablets etc. according to a conventional process; the mixture of the polysaccharide and the protein is prepared into capsules and tablets etc. according to a conventional process.

Embodiment 7 Application of the said medicament in treating gout (not covered by the claims)**(1) Therapeutic effect of the medicament in the present invention on gout rats model induced by streptozotocin**

[0024] In model control groups and the invention groups, the streptozotocin was first used to induce gout in rats, and then the rats in the invention groups were administered the medicament prepared by the present invention and administered intragastrically three times a day; the rats in control group were administered the distilled water, and the value of urine protein, and the creatinine and urea nitrogen content were measured four weeks later, with the results shown in Table 1 and Table 2.

Table 1 Effect of the medicament on gout rats model induced by streptozotocin

Group	Dose (g/day)	Urine protein(mg/L)	Creatinine(mmol/L)	Urea nitrogen(mmol/L)
compared group			133.39±13.91	6.37±2.48
compared		364.83±20.62	288.01±12.35	56.07±3.00

Group	Dose (g/day)	Urine protein(mg/L)	Creatinine(mmol/L)	Urea nitrogen(mmol/L)
group				
Embodiment 1	3	110.43±21.32	169.65±10.05	14.67±2.30
Embodiment 2	3	50.89±22.14	152.45±10.89	11.58±2.60
Embodiment 3	3	62.13±23.96	158.48±13.59	12.12±3.29
Embodiment 4	3	107.45±22.76	162.23±10.67	13.75±2.50
Embodiment 5	3	130.12±21.23	166.18±11.25	14.41±3.30
Embodiment 6	3	168.78±22.70	172.82±11.10	15.64±2.93

Table 2 Effect of the medicament at different doses on gout rats model induced by streptozotocin

Group	Dose (g/day)	Urine protein(mg/L)	Creatinine(mmol/L)	Urea nitrogen(mmol/L)
Normal compared group	-	-	133.39±13.91	6.37±2.48
Model compared group	-	364.83±20.62	288.01±12.35	56.07±3.00
Embodiment 2	0.4	80.12±20.23	178.22±11.15	15.24±2.63
Embodiment 2	0.8	65.45±22.26	162.38±13.19	12.28±2.29
Embodiment 2	3	50.89±22.14	152.45±10.89	11.58±2.60
Embodiment 2	4	70.22±21.25	155.12±11.20	12.64±2.33
Embodiment 2	6	92.32±22.34	162.48±12.19	13.12±2.59
Embodiment 2	8	108.25±20.89	179.82±10.32	14.64±2.41

(2) Therapeutic effect of the medicament on hyperuricemia mice model induced by xanthine

A. Experimental materials

[0025]

Animals: 50 clean-grade Kunming mice, male, 20g ± 2g

Medicament: the medicament described in Embodiments 1-6

Xanthine: Sinopharm Chemical Reagent Co., Ltd., batch number of WL20101109

Blood uric acid test kit: Shanghai Rongsheng Biotech Co., Ltd., batch number of 20100406

B. Experimental method

[0026] Mice were randomly divided into 8 groups after entering the laboratory, namely normal control group, model control group, and present invention groups 1-6, 10 rats in each group, weighed and numbered;

[0027] After one-week adaptive feeding, the mice in the normal control group and the model control group were administered distilled water by means of intragastric administration, and mice 1-6 in the invention group were administered the medicament in the present invention at a dose of 3 g/day, administered intragastrically, three times a day for three consecutive days.

[0028] At one hour after the last intragastric administration, the mice in normal control group were intraperitoneally injected with normal saline 0.1mL/10g, and the mice in the other 7 groups were intraperitoneally injected with 10% xanthine suspension 0.1mL/10g for modeling. After 0.5h, the blood was taken by eyeball removal method, the uric acid value was measured and the measurement results were statistically analyzed. The experimental results are shown in Table 3.

Table 3 Effect of the medicament on blood uric acid in hyperuricemia mice model

Group	Dose (g/day)	Blood Uric acid(μ mol/L)
Normal compared group	-	126.4 $\pm$ 15.48
Model compared group	-	250.4 $\pm$ 17.62
Embodiment 1	3	160.13 $\pm$ 16.96
Embodiment 2	3	147.89 $\pm$ 18.14
Embodiment 3	3	150.43 $\pm$ 18.32
Embodiment 4	3	153.45 $\pm$ 17.76
Embodiment 5	3	158.12 $\pm$ 18.23
Embodiment 6	3	160.5 $\pm$ 16.70

Table 4 Effect of the medicament at different doses on blood uric acid in hyperuricemia mice model

Group	Dose (g/day)	Blood Uric acid (μ mol/L)
Normal compared group	-	126.4 $\pm$ 15.48
Model compared group	-	250.4 $\pm$ 17.62

Group	Dose (g/day)	Blood Uric acid (μ mol/L)
Embodiment 2	0.4	175.5 $\pm$ 15.80
Embodiment 2	0.8	150.13 $\pm$ 16.36
Embodiment 2	3	147.89 $\pm$ 17.84
Embodiment 2	4	152.12 $\pm$ 18.03
Embodiment 2	6	157.45 $\pm$ 17.46
Embodiment 2	8	172.89 $\pm$ 17.25

(3) Human clinical trials

[0029] The human clinical trials are as follows:

A total of 100 cases were included, 52 males and 48 females, aged from 32 years old to 61 years old, with a course of disease ranging from 6 months to 8 years, mostly 3 to 5 years; as the first metatarsophalangeal joint swelling and pain, 50 cases had the unilateral involvement and 50 cases had the bilateral involvement; wherein 60 cases of unilateral arthritis, 40 cases of pain and inflammation involving bilateral joints, and wherein 10 cases of the presence of arthritic calculus. All cases passed the examination and all the 100 cases had an increase in blood uric acid. For all of them, the possibility of being caused by medications or other diseases was ruled out.

[0030] Treatment methods:

All the gout-treating medicament in Embodiment 3 were administered 3 g per day, in 3 times, 3 times per day, to be taken after being mixed with warm boiled water. 21 days remained a course of treatment, usually a course of treatment was needed, and the efficacy was observed at the end of the treatment.

[0031] Efficacy assessment criteria:

Healing: all clinical symptoms disappeared, joints could move freely, the blood uric acid decreased to a value within the normal range; marked efficacy: clinical symptoms improved, joint activity was flexible, and the blood uric acid decreased by more than 10%; no response: no change in symptoms and laboratory test indicators.

[0032] Results:

Healed: 85 cases, accounting for 85%;

Marked efficacy: 13 cases, accounting for 13%;

No response: 2 cases, accounting for 2%;

Total effective rate: 98%;

The recurrence rate is low, and as for the healed patients, the recurrence rate in 5 years is only 1%.

Embodiment 8 A medicament for use in treating gout (not covered by the claims)

[0033] It comprises, on the basis of weight, the following components:

1 portion of marine algal glycoprotein and 1 portion of glucuronic acid;

The marine algal glycoprotein comprises, by weight content, 9% sugar and 88% protein;

The molecular weight is 8 kDa;

The said sugar is a polysaccharide;

The said marine algae is spirulina;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 9 A medicament for use in treating gout (not covered by the claims)

[0034] It comprises, on the basis of weight, the following components:

15 portions of marine algal glycoprotein and 6 portions of glucuronic acid;

The marine algal glycoprotein comprises, by weight content, 9% sugar and 88% protein;

The molecular weight is 12 kDa;

The said sugar is a polysaccharide;

The said marine algae is chlorella;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 10 A medicament for use in treating gout (not covered by the claims)

[0035] It comprises, on the basis of weight, the following components:

40 portions of marine algal glycoprotein and 13 portions of glucuronic acid;

The marine algal glycoprotein comprises, by weight content, 30% sugar and 70% protein;

The molecular weight is 20 kDa;

The said sugar is a polysaccharide;

The said marine algae is *Bangia atropurpurea* (Roth) C. Agardh;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 11 A medicament for use in treating gout (not covered by the claims)

[0036] It comprises, on the basis of weight, the following components:

70 portions of marine algal glycoprotein and 17 portions of glucuronic acid;

The marine algal glycoprotein comprises, by weight content, 50% sugar and 50% protein;

The molecular weight is 8 kDa;

The said sugar is a polysaccharide;

The said marine algae is gulfweed;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 12 A medicament for use in treating gout (not covered by the claims)

[0037] It comprises, on the basis of weight, the following components:

99 portions of marine algal glycoprotein and 25 portions of glucuronic acid;

The marine algal glycoprotein comprises, by weight content, 99% sugar and 1% protein;

The molecular weight is 20 kDa;

The said sugar is a polysaccharide;

The said marine algae is synuraceae urelin;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 13 A medicament for use in treating gout (not covered by the claims)

[0038] It comprises, on the basis of weight, the following components:

1 portion of marine algal glycoprotein, 1 portion of glucuronic acid and 2 portions of indigo naturalis;

The marine algal glycoprotein comprises, by weight content, 9% sugar and 88% protein;

The molecular weight is 6 kDa;

The said sugar is a polysaccharide;

The said marine algae is blue-green algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 14 A medicament for use in treating gout (not covered by the claims)

[0039] It comprises, on the basis of weight, the following components:

27 portions of marine algal glycoprotein, 6 portions of glucuronic acid and 4 portions of indigo naturalis;

The marine algal glycoprotein comprises, by weight content, 9% sugar and 88% protein;

The molecular weight is 20 kDa;

The said sugar is a polysaccharide;

The said marine algae is blue-green algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 15 A medicament for use in treating gout (not covered by the claims)

[0040] It comprises, on the basis of weight, the following components:

52 portions of marine algal glycoprotein, 14 portions of glucuronic acid and 5 portions of indigo naturalis;

The marine algal glycoprotein comprises, by weight content, 30% sugar and 70% protein;

The molecular weight is 200 kDa;

The said sugar is a polysaccharide;

The said marine algae is blue-green algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 16 A medicament for use in treating gout (not covered by the claims)

[0041] It comprises, on the basis of weight, the following components:

99 portions of marine algal glycoprotein, 25 portions of glucuronic acid and 9 portions of indigo naturalis;

The marine algal glycoprotein comprises, by weight content, 50% sugar and 50% protein;

The molecular weight is 3000 kDa;

The said sugar is a polysaccharide;

The said marine algae is blue-green algae;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Application of the said medicament in Embodiment 8 - Embodiment 16 in treating gout

Using the test method said in Embodiment 7, the medicament said in Embodiment 8-Embodiment

[0042] 16 in this invention groups have the following application effects:

(1) Effects of the medicament on gout rats model induced by streptozotocin

[0043]

Table 5 Effects of the medicament on gout rats model induced by streptozotocin

Group	Dose (g/day)	Urine protein(mg/L)	Creatinine(mmol/L)	Urea nitrogen(mmol/L)
Normal compared group	-	-	130.39±13.91	6.57±2.48
Model compared group	-	364.83±20.62	288.01±12.35	56.07±3.00
Embodiment 8	3	48.89±20.14	150.15±10.29	10.20±2.50
Embodiment 9	3	48.60±20.02	149.80±10.14	10.00±2.23
Embodiment 10	3	30.5±19.54	135.56±10.28	7.89±1.95
Embodiment 11	3	48.50±19.82	149.45±10.21	9.94±2.20
Embodiment 12	3	48.60±19.24	149.25±10.18	9.92±2.14
Embodiment 13	3	47.65±19.75	148.15±10.00	9.90±1.98
Embodiment 14	3	47.29±19.85	150.15±9.90	9.95±2.00
Embodiment 15	3	32.24±19.56	136.15±10.16	7.70±1.96
Embodiment 16	3	47.59±19.47	150.15±10.27	9.89±1.95

(2) Therapeutic effect of the medicament on hyperuricemia mice model induced by xanthine

[0044]

Table 6 Effect of the medicament on hyperuricemia mice model

Group	Dose (g/day)	Blood Uric acid (μ mol/L)
Normal compared group	-	127.4±15.48
Model compared group	-	250.4±17.62

Group	Dose (g/day)	Blood Uric acid (μ mol/L)
Embodiment 8	3	146.21 $\pm$ 17.54
Embodiment 9	3	145.01 $\pm$ 17.37
Embodiment 10	3	135.25 $\pm$ 17.08
Embodiment 11	3	145.21 $\pm$ 17.44
Embodiment 12	3	146.04 $\pm$ 17.21
Embodiment 13	3	146.11 $\pm$ 16.95
Embodiment 14	3	146.00 $\pm$ 17.08
Embodiment 15	3	134.34 $\pm$ 17.12
Embodiment 16	3	146.24 $\pm$ 17.04

(3) Human clinical trials

[0045] Results:

Healed: 88 cases, accounting for 88%;

Marked efficacy: 10 cases, accounting for 10%;

No response: 2 cases, accounting for 2%;

Total effective rate: 98%;

The recurrence rate is low, and as for the healed patients, the recurrence rate in 6 years is only 1%.

Embodiment 17 A medicament for use in treating gout

[0046] It comprises, on the basis of weight, the following components:

1 portion of marine algal glycoprotein, 8 portions of indigo naturalis, and 7 portions of twotooth achyranthes root;

The marine algal glycoprotein comprises, by weight content, 10% sugar and 85% protein;

The molecular weight is 16 kDa;

The said sugar is a polysaccharide;

The said marine algae is spirulina;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 18 A medicament for use in treating gout

[0047] It comprises, on the basis of weight, the following components:

40 portions of marine algal glycoprotein, 11 portions of indigo naturalis, and 9 portions of twotooth achyranthes root;

[0050] The marine algal glycoprotein comprises, by weight content, 8% sugar and 88% protein;

The molecular weight is 14 kDa;

The said sugar is a polysaccharide;

The said marine algae is chlorella;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 19 A medicament for use in treating gout

[0048] It comprises, on the basis of weight, the following components:

99 portions of marine algal glycoprotein, 14 portions of indigo naturalis, and 13 portions of twotooth achyranthes root;

The marine algal glycoprotein comprises, by weight content, 30% sugar and 70% protein;

The molecular weight is 38 kDa;

The said sugar is a polysaccharide;

The said marine algae is Gelidium amansii Lamx;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 20 A medicament for use in treating gout

[0049] It comprises, on the basis of weight, the following components:

1 portion of marine algal glycoprotein, 8 portions of indigo naturalis, 7 portions of twotooth achyranthes root and 1 portion of glucuronic acid;

The marine algal glycoprotein comprises, by weight content, 10% sugar and 85% protein;

The molecular weight is 16 kDa;

The said sugar is a polysaccharide;

The said marine algae is spirulina;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 21 A medicament for use in treating gout

[0050] It comprises, on the basis of weight, the following components:

40 portions of marine algal glycoprotein, 11 portions of indigo naturalis, 9 portions of twotooth achyranthes root and 8 glucuronic acid;

[0056] The marine algal glycoprotein comprises, by weight content, 8% sugar and 88% protein;

The molecular weight is 14 kDa;

The said sugar is a polysaccharide;

The said marine algae is chlorella;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Embodiment 22 A medicament for use in treating gout

[0051] It comprises, on the basis of weight, the following components:

99 portions of marine algal glycoprotein, 14 portions of indigo naturalis, 13 portions of twotooth achyranthes root and 15 portions of glucuronic acid;

The marine algal glycoprotein comprises, by weight content, 30% sugar and 70% protein;

The molecular weight is 200 kDa;

The said sugar is a polysaccharide;

The said marine algae is *Gelidium amansii* Lamx;

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

Application of the said medicament in Embodiment 17 - Embodiment 22 in treating gout

Using the test method said in Embodiment 7, the medicament said in Embodiment 17- Embodiment

[0052] 22 in this invention groups have the following application effects:

(1) Effects of the medicament in the present invention on gout rats model induced by streptozotocin

[0053]

Table 7 Effects of the medicament in the present invention on gout rats model induced by streptozotocin

Group	Dose (g/day)	Urine protein(mg/L)	Creatinine(mmol/L)	Urea nitrogen(mmol/L)
Normal compared group	-	-	130.39±13.91	6.57±2.48
Model compared group	-	364.83±20.62	288.01±12.35	56.07±3.00
Embodiment 17	3	46.60±19.68	147.80±10.00	9.80±2.14
Embodiment 18	3	28.7±19.27	132.56±10.02	6.69±1.92
Embodiment 19	3	46.24±19.75	147.65±9.97	9.84±1.97
Embodiment 20	3	45.45±19.68	145.25±10.05	9.70±2.14

Group	Dose (g/day)	Urine protein(mg/L)	Creatinine(mmol/L)	Urea nitrogen(mmol/L)
Embodiment 21	3	27.5±19.54	130.36±10.28	6.64±1.86
Embodiment 22	3	45.30±19.68	145.67±10.19	9.74±2.04

(2) Therapeutic effect of the medicament in the present invention on hyperuricemia mice model induced by xanthine

[0054]

Table 8 Effect of the medicament in the present invention on hyperuricemia mice model

Group	Dose (g/day)	Blood Uric acid (μ mol/L)
Normal compared group	-	126.4±15.48
Model compared group	-	250.4±17.62
Embodiment 17	3	143.31±17.57
Embodiment 18	3	128.20±17.42
Embodiment 19	3	143.27±17.35
Embodiment 20	3	142.89±17.05
Embodiment 21	3	126.57±17.18
Embodiment 22	3	142.57±17.17

(3) Human clinical trials

[0055] Results:

Healed: 90 cases, accounting for 90%;

Marked efficacy: 9 cases, accounting for 9%;

No response: 1 case, accounting for 1%;

Total effective rate: 99%;

The recurrence rate is low, and as for the healed patients, no recurrence was observed in 5 years.

Embodiment 23 A medicament for use in treating gout (not covered by the claims)

[0056] It comprises, on the basis of weight, the following components:

80 portions of marine algal glycoprotein, 10 portions of radix sophorae flavescentis, 5 portions of radices saussureae, 5 portions of rhododendron mariae, 7 portions of unispice kyllinga herb, 8 portions of fructus evodiae, 10 portions of pyrola herb;

The marine algal glycoprotein comprises, by weight content, 10% sugar and 85% protein;

The molecular weight is 15 kDa;

The said marine algae is green algae;

The said sugar, by weight, comprises the following components: 8 portions of glucose, 5 portions of galactose, and 11 portions of carubiose;

The protein mentioned, by weight, includes the following components: 7 portions of serine, 9 portions of threonine, and 15 portions of hydroxylysine.

Embodiment 24 A medicament for use in treating gout (not covered by the claims)

[0057] It comprises, on the basis of weight, the following components:

75 portions of marine algal glycoprotein, 12 portions of epimedium, 8 portions of nutmeg, 7 portions of gallnut, 9 portions of radix aconiti lateralis preparata, 6 portions of gastrodia elata, and 5 portions of caulis sinomenii;

The marine algal glycoprotein comprises, by weight content, 12% sugar and 75% protein;

The molecular weight is 8 kDa;

The said marine algae is blue-green algae;

The said sugar, by weight, comprises the following components: 20 portions of glucose, 8 portions of galactose, and 15 portions of carubiose;

The protein mentioned, by weight, includes the following components: 10 portions of serine, 15 portions of threonine, and 17 portions of hydroxylysine.

Embodiment 25 A medicament for use in treating gout (not covered by the claims)

[0058] It comprises, on the basis of weight, the following components:

70 portions of marine algal glycoprotein, 8 portions of monkshood, 6 portions of semen dolichoris, 5 portions of bombyx batryticatus, 3 portions of pinellia ternata, 7 portions of atractylodes rhizome, and 8 portions of pubescent angelica root;

The marine algal glycoprotein comprises, by weight content, 14% sugar and 72% protein;

The molecular weight is 20 kDa;

The said marine algae is blue-green algae;

The said sugar, by weight, comprises the following components: 20 portions of glucose, 8 portions of galactose, and 15 portions of carubinoase;

The protein mentioned, by weight, includes the following components: 10 portions of serine, 15 portions of threonine, and 17 portions of hydroxylysine.

Embodiment 26 A medicament for use in treating gout (not covered by the claims)

[0059] It comprises, on the basis of weight, the following components:

78 portions of marine algal glycoprotein, 10 portions of ligusticum wallichii, 8 portions of alismatis rhizoma, 6 portions of cimicifuga foetida, 7 portions of glabrousleaf pittosporum leaf, 7 portions of notoperygium root, and 5 portions of Chinese clematis root;

The marine algal glycoprotein comprises, by weight content, 22% sugar and 69% protein;

The molecular weight is 100 kDa;

The said marine algae is blue-green algae;

The said sugar, by weight, comprises the following components: 20 portions of glucose, 8 portions of galactose, and 15 portions of carubinoase;

The protein mentioned, by weight, includes the following components: 10 portions of serine, 15 portions of threonine, and 17 portions of hydroxylysine.

Embodiment 27 A medicament for use in treating gout (not covered by the claims)

[0060]

Step 1: Weighing

Weigh the marine algal glycoprotein and all Chinese medicine components according to the formula;

Step 2: Extraction of Chinese medicine**1. (1) Washing**

Wash all Chinese medicine components with clear water, and remove the impurities;

2. (2) Crash and microwave extraction

The Chinese medicine is pulverized into 100-mesh medicinal material powder, 8 times of 50% ethanol is added, the temperature is controlled at 60 °C, microwave radiation is performed at the microwave power of 260 W, microwave wavelength of 130 mm, a frequency of 1200 MHz for 5 min, then filtration is carried out, and finally the filtrate is collected;

The medicine dregs are separated, 6 times of clear water is added, the temperature is controlled at 50 °C, microwave radiation is performed at the microwave power of 200W, microwave wavelength of 140 mm, a frequency of 1250 MHz for 5 min, then filtration is carried out, and finally the filtrate is collected;

Pool the filtrate collected from the two procedures; atomize and dry to prepare them into Chinese medicine powder;

Step 3 Add marine algal glycoprotein.

[0061] Mixing the powder of marine algal glycoprotein with the above prepared Chinese medicine powder to produce different dosage forms such as capsules, tablets and so on.

Application of the said medicament in Embodiment 23 - Embodiment 26 in treating gout

[0062] Applying the test method specified in Embodiment 7, this invention group is the medicament mentioned in Embodiment 23-Embodiment 26, the applicable effects are as follows:

(1) Effects of the medicament on gout rats model induced by streptozotocin**[0063]**

Table 9 Effects of the medicament on gout rats model induced by streptozotocin

Group	Dose(g/day)	Urine protein(mg/L)	Creatinine(mmol/L)	Urea nitrogen(mmol/L)
Normal compared group	-	-	130.39±13.91	6.57±2.48
Model compared group	-	364.83±20.62	288.01±12.35	56.07±3.00
Embodiment 23	3	40.86±20.02	130.25±10.00	6.68±2.25
Embodiment 24	3	40.26±20.12	130.15±10.07	6.65±2.15
Embodiment 25	3	40.56±20.25	131.23±10.00	6.45±2.34
Embodiment 26	3	40.16±20.35	130.05±10.37	6.28±2.25

(2) Therapeutic effect of the medicament on hyperuricemia mice model induced by xanthine

[0064]

Table 10 Effect of the medicament on hyperuricemia mice model

Group	Dose (g/day)	Blood Uric acid (μ mol/L)
Normal compared- group	-	126.4±15.48
Model compared group	-	250.4±17.62
Embodiment 23	3	128.3±12.25
Embodiment 24	3	127.56±12.35
Embodiment 25	3	129.25±12.24
Embodiment 26	3	130.89±12.19

(3) Human clinical trials

[0065] Results:

Healed: 90 cases, accounting for 90%;

Marked efficacy: 9 cases, accounting for 9%;

No response: 1 case, accounting for 1%;

Total effective rate: 99%;

The recurrence rate is low, and as for the healed patients, no recurrence was observed in 5 years.

[0066] For the medicament mentioned in this invention, pH is between 5.3-9.8, and 6.5-7.5 is preferred.

[0067] The invention has been subjected to a large number of experiments, and we have carried out multiple tests using a mixture of marine shells, bones of livestock and poultry, a mixture of glycoprotein, polysaccharides and proteins extracted from the skeleton of marine animals, and the objectives of the invention have also been achieved.

Embodiment 28 A medicament for use in treating gout (not covered by the claims)

[0068] The said medicament is a mixture of polysaccharides and proteins;

The said medicament comprises, by weight content, 1-99% polysaccharide and 1-99% protein.

The said polysaccharide comprises: glucose, galactose, mannose and rhamnose;

The said protein comprises: arginine, lysine, serine, and threonine.

[0069] As for the said mixture of polysaccharide and the protein, the polysaccharide has a molecular weight of 0.2-3000 kDa and the protein has a molecular weight of 0.2-3000 kDa.

[0070] The mixture of polysaccharides and proteins, further a mixture of algal polysaccharides and algal proteins;

The said mixture of the algal polysaccharide and the algal protein also comprises a pigment;

The said pigment is a natural pigment contained in the algal substance;

The said algal protein may be phycocyanin, phycoerythrin or algae xanthoprotein. The said glycoprotein includes synthetic glycoprotein, synthetic polysaccharide and protein.

[0071] The medicament mentioned in this invention has a No Observed Adverse Effect Level (NOAEL) of 1.6 g/kg for 12-week oral administration for dogs, which is equivalent to 50 times

the equivalent dose for humans, so it is concluded that the safety of the clinical trial can be guaranteed.

[0072] The medicament described in this invention can also be a health care product or a food.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US2015337001A1 **[0004]**
- US2015306054A1 **[0005]**
- CN201310601617 **[0006]**
- JP2010235517A **[0007]**
- EP1506785A1 **[0008]**
- CN101002873A **[0009]**
- US2010048507A1 **[0010]**

PATENTKRAV

1. Medikament til anvendelse til behandling af urinsyregigt,
kendetegnet ved, at medikamentet i forhold til vægtindhold omfatter 1-99 andele af et havalge-glycoprotein, 8-14 andele indigo naturalis og 7-5 13 andele Achyranthes bidentata-rod.
2. Medikament til anvendelse til behandling af urinsyregigt ifølge krav 1, der endvidere omfatter 1-15 andele glucuronsyre.
3. Medikament til anvendelse til behandling af urinsyregigt ifølge krav 1, hvor havalge-glycoproteinet i forhold til vægtindhold omfatter 1-99 % sukkerart
10 og 1-99 % protein.
4. Medikament til anvendelse til behandling af urinsyregigt ifølge krav 1, hvor en molekylvægt af havalge-glycoproteinet er 0,2 til 3000 kDa.
5. Medikament til anvendelse til behandling af urinsyregigt ifølge krav 1, hvor algerne omfatter en eller flere slags blågrønalger, grønalger, rødalger,
15 gulalger og brunalger.