




(86) Date de dépôt PCT/PCT Filing Date: 2015/11/03
(87) Date publication PCT/PCT Publication Date: 2016/05/12
(45) Date de délivrance/Issue Date: 2024/04/16
(85) Entrée phase nationale/National Entry: 2017/04/28
(86) N° demande PCT/PCT Application No.: US 2015/058878
(87) N° publication PCT/PCT Publication No.: 2016/073514
(30) Priorité/Priority: 2014/11/03 (US62/074,374)

(51) Cl.Int./Int.Cl. A61K 38/49 (2006.01),
A61P 9/00 (2006.01), C12Q 1/60 (2006.01)
(72) Inventeur/Inventor:
GUREWICH, VICTOR, US
(73) Propriétaire/Owner:
THROMBOLYTIC SCIENCE, LLC, US
(74) Agent: SMART & BIGGAR LP

(54) Titre : METHODES ET COMPOSITIONS POUR THROMBOLYSE SURE ET EFFICACE
(54) Title: METHODS AND COMPOSITIONS FOR SAFE AND EFFECTIVE THROMBOLYSIS



(57) Abrégé/Abstract:

Provided herein are methods, compositions, and kits for safe and effective thrombolysis, e.g., in therapy for a potential stroke or acute myocardial infarction ("AMI").

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2016/073514 A3

(43) International Publication Date

12 May 2016 (12.05.2016)

(51) International Patent Classification:
A61P 9/00 (2006.01) *C12Q 1/60* (2006.01)

(21) International Application Number:
 PCT/US2015/058878

(22) International Filing Date:
 3 November 2015 (03.11.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
 62/074,374 3 November 2014 (03.11.2014) US

(71) Applicant: THROMBOLYTIC SCIENCE, LLC [US/US]; 763D Concord Avenue, Cambridge, Massachusetts 02138 (US).

(72) Inventor: GUREWICH, Victor; 11 Reservoir Street, Cambridge, Massachusetts 02138 (US).

(74) Agent: FASSE, J. Peter; Fish & Richardson P.C., P.O. Box 1022, Minneapolis, Minnesota 55440-1022 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:

— with international search report (Art. 21(3))

(88) Date of publication of the international search report: 11 August 2016

(54) Title: METHODS AND COMPOSITIONS FOR SAFE AND EFFECTIVE THROMBOLYSIS

FIG. 5A



(57) Abstract: Provided herein are methods, compositions, and kits for safe and effective thrombolysis, e.g., in therapy for a potential stroke or acute myocardial infarction ("AMI").

WO 2016/073514 A3

Methods and Compositions For Safe and Effective Thrombolysis

TECHNICAL FIELD

This invention relates to methods and compositions for safe and effective
5 thrombolysis.

BACKGROUND

Thrombosis occurs when a blood clot (thrombus) forms inside a blood vessel and obstructs the flow of blood through the circulatory system. When a blood vessel is injured, the body uses platelets and fibrin to form a blood clot to seal injured vessels and prevent 10 blood loss. This process is called hemostasis. Even when a blood vessel is not injured, blood clots may form in the body under certain circumstances. Blood clots consist largely of aggregated platelets and a mesh of cross-linked fibrin, which is a natural polymer of fibrinogen in the blood. When a thrombus is large enough to reduce the blood flow to a tissue, hypoxia or anoxia can occur, leading to tissue damage or even tissue death. 15 Depending on the location in the arterial system, i.e., heart, brain, or leg, a thrombus can trigger heart attack, stroke, or peripheral gangrene. In the venous circulation the same process can cause thrombophlebitis (deep vein thrombosis) or a pulmonary embolism. Together, these cardiovascular diseases constitute the leading causes of death and disability in industrialized countries.

20 Thrombolysis mainly involves the use of thrombolytic drugs to dissolve the disease-causing blood clot and restore blood flow. Thrombolytic drugs such as tPA and its derivatives in current use function by activating the proenzyme plasminogen to the protease plasmin, which degrades the fibrin mesh in the blood clot and makes the clot soluble, thus restoring blood flow through occluded blood vessels. However, the present thrombolytic 25 drugs also induce degradation of hemostatic fibrin that seals wounds, and generate plasmin in plasma which degrades three clotting factors, fibrinogen, factor V, and factor VII (hemophilic factor). Thus, current thrombolytic therapy carries the risk of causing bleeding from hemophilia-like side effects or from the degradation of hemostatic fibrin. This imposes serious limitations on the number of patients eligible for treatment and limits the dose of

thrombolytic that can be used. As a result, only about 5% of patients with stroke receive treatment and the efficacy of this treatment is limited.

A stroke can be caused either by a blood clot or a bleeding vessel in the brain. However, properly diagnosing the specific cause of a stroke requires a CT scan or MRI, 5 which can delay treatment if not immediately available. Absent such a proper diagnosis, it can be very dangerous to administer a clot dissolving agent, such as tissue plasminogen activator (“tPA”), if the cause happens to be a bleed rather than a blood clot, as the results of administering tPA or other thrombolytic agent to a patient who has had a brain bleed can be fatal.

10

SUMMARY

The present disclosure is based, at least in part, on the discovery that the distinct and different specificities of tPA and prourokinase mutant (“proUK mutant or “mproUK”) means that small doses of each in combination induces thrombolysis that is faster and safer, i.e., more specific and with lower incidence of bleeding complications, e.g., for treating stroke or 15 acute myocardial infarction (“AMI”), than is possible using either one alone (monotherapy) with high doses that achieve a maximum blood clot lysis rate, but with a high risk of bleeding complications.

The mproUK can comprise a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase, referred to herein as “M5.” mproUK, like proUK, is a 20 proenzyme, i.e., the inactive precursor of the active enzyme. It differs from proUK in that its enzymatic form mutant urokinase or mUK, unlike urokinase (UK), is inhibited by a plasma inhibitor, C1-inhibitor, which additionally to the lower doses necessary, helps to reduce the bleeding side effects seen with pro-UK, which are due to UK for which there is insufficient inhibitor (plasminogen activator inhibitor-1, PAI-1) in the plasma.

25 For patients with stroke or AMI, the time needed for thrombolysis and reperfusion is critical to clinical outcome and survival rate. This means that treatment must be safe enough to be given outside the hospital without preliminary diagnostic testing. However, this is possible only if the hemorrhagic risk is greatly reduced or eliminated, as is the case with the new methods described herein including the administration of a mini-dose of tPA (e.g., a bolus of less than 5.0 mg, e.g., less than or equal to 4.5 mg, 4.0 mg, 3.5 mg, 3.0 mg, 2.5 mg, 30 or 2.0 mg) and low dose of mproUK (e.g., an infusion over 60 to 90 minutes (e.g., 60, 70, 80,

or 90 minutes) at a rate of 60 to 120 mg/hour, e.g., 60 to 90 mg/hour, e.g., 60, 65, 70, 75, 80, 85, or 90 mg/hour) in combination.

If the endogenous concentration of C1-inhibitor is insufficient in a given subject, it can be supplemented in the new methods by administering an effective amount of a C1-inhibitor, e.g., a commercially available C1-inhibitor such as CINRYZE®, CETOR®, BERINERT®, and RUCONEST®. The C1-inhibitor is an additional precaution in case some conversion of mproUK to mutant UK occurs, but may not be necessary due to the low dose of mproUK required and the fact that C1-inhibitor is an acute phase reactant, which is automatically elevated by the body immediately after a heart attack or stroke.

Provided herein in a first aspect are methods of treating a subject, e.g., a human patient, with symptoms of a stroke or an acute myocardial infarction (AMI) at a maximum rate of blood clot lysis and with minimal associated hemorrhagic side effects, the methods include (a) identifying a subject who potentially had a stroke or AMI by observing one or more symptoms of a stroke or AMI without determining the cause of the stroke; and (b) administering to the subject a bolus of a first composition including a mini-dose, i.e., less than 5 mg, of tissue plasminogen activator (tPA), e.g., 2.0, 2.5, 3.0, 3.5, 4.0, or 4.5 mg of tPA, followed by an intravenous infusion of a second composition including a low dose of pro-urokinase mutant (mproUK) infused over 60 to 90 minutes, e.g., 60, 65, 70, 75, 80, 85, or 90 minutes, at a low dosage rate of 60 to 120 mg, e.g., 60, 70, 75, 80, 85, 90, 100, or 120 mg/hour; wherein a maximum clot lysis rate is achieved with minimal associated hemorrhagic side effects.

In some embodiments of these methods the minimal associated hemorrhagic side effects can be determined as a level of fibrinogen degradation in the subject's blood of less than about 30 percent, e.g., less than 27.5%, 25%, 22.5%, or 20% fibrinogen degradation (whereas fibrinogen degradation is about 50 to 80% with monotherapy by either composition alone). In some embodiments, the maximum clot lysis rate is indicated as having been achieved by a Thrombolysis In Myocardial Infarction (TIMI) score of 2 or higher. In other embodiments, the maximum clot lysis rate is indicated as having been achieved by a lysis of about 50% of the mass of at least one clot in the subject achieved within 75 minutes, e.g., within 70, 60, 50, 40, 30 minutes.

In these methods, the pro-urokinase mutant can include a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase. The methods can include

starting administration of the second composition within five, 10, or 15 minutes after the administration of the first composition. In some embodiments, the first composition and the second composition together lyse 50% of a mass of at least one blood clot in the subject in less than one hour.

5 In certain embodiments, the methods can further include administering to the subject a third composition including a bolus of C1-inhibitor. The third composition can be administered to the subject before or at about the same time, e.g., within 5 minutes, of the administration of the second composition. In certain embodiments, the third composition is administered in an amount sufficient to establish a concentration of C1-inhibitor that is about 10 500-750 μ g/ml in the subject's blood. In some embodiments, the third composition is a bolus of 500-1500 mg of C1-inhibitor.

In some of these methods the first composition and the second composition together lyse the blood clots in the presence of the C1-inhibitor with less than 30% fibrinogen degradation when compared to monotherapy by either tPA or pro-urokinase alone.

15 In another aspect, the disclosure provides kits that include a first composition in a first container including 2-5 mg of tissue plasminogen activator (tPA); and a second composition in a second container including 60-120 mg of a pro-urokinase mutant (mproUK) having a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase. In these kits, the first composition can be formulated suitable for administration as 20 a bolus and/or the second composition can be formulated suitable for intravenous infusion. In certain embodiments, the kits further include a third composition including 500-1500 mg of C1-inhibitor, e.g., formulated suitable for administration as a bolus.

In another aspect, the disclosure provides compositions for use in any of the methods described herein. In certain embodiments, the compositions are for use in treating a subject 25 with symptoms of a stroke or acute myocardial infarction (AMI) at a maximum rate of clot lysis and with minimal associated hemorrhagic side effects. These compositions include a first composition including less than 5 mg of tissue plasminogen activator (tPA), wherein the first composition is or is prepared to be administered to a subject in a dosage regime of a bolus; and a second composition including a pro-urokinase mutant (mproUK), wherein the 30 second composition is or is prepared to be administered to a subject by an intravenous infusion in a dosage regime of 60 to 120 mg/hour for 60 to 90 minutes following the administration of the first composition; wherein the subject is identified as potentially having

had a stroke or AMI by observing one or more symptoms of a stroke or AMI without determining the cause of the stroke prior to administering the first composition; and wherein a maximum clot lysis rate is achieved with minimal associated hemorrhagic side effects. These compositions can include all of the features and elements described herein.

5 The term "treatment" or "therapeutic treatment" means the administration of one or more pharmaceutical agents to a subject or the performance of a medical procedure on the body of a subject. The term therapeutic treatment also includes an adjustment (e.g., increase or decrease) in the dose or frequency of one or more pharmaceutical agents that a subject can be taking, the administration of one or more new pharmaceutical agents to the subject, or the
10 removal of one or more pharmaceutical agents from the subject's treatment plan.

As used herein, a "subject" or "patient" is a human.

An "effective amount" used herein is an amount sufficient to achieve thrombolysis without causing significant hemorrhagic or hemophilia-like side effects. An effective amount can be administered in one or more administrations, applications, or dosages. A
15 therapeutically effective amount of a pharmaceutical composition (i.e., an effective dosage) depends on the pharmaceutical composition selected.

The "maximum lysis rate" of blood clots is defined herein as a rate of lysis of blood clots at least as fast as what can be achieved by monotherapy with tPA, proUK or mproUK at a maximally effective dosage (i.e., a dose of the drug at which no faster clot lysis can be
20 achieved by adding additional drug, e.g., a plateau in the clot lysis rate is shown at the maximally effective dosage) without any associated hemorrhagic side effects. It is important to note that the maximum lysis rate achieved using monotherapy, e.g., with tPA or M5, would cause significant bleeding in a patient, as indicated, for example, by a level of fibrinogen
25 degradation of greater than 30%. Surprisingly, the present therapeutic methods achieve the maximum lysis rate without hemorrhagic side effects, which can be indicated by a fibrinogen degradation level of less than 30%, which would not be associated with excess bleeding and would be clinically acceptable.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention
30 belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below.

In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

5 The invention as claimed relates to:

- use of tissue plasminogen activator (tPA) comprised in a first composition and a pro-urokinase mutant (mproUK) comprised in a second composition for treating a subject with symptoms of a stroke or an acute myocardial infarction (AMI) at a maximum rate of blood clot lysis and with minimal associated hemorrhagic side effects, wherein: the subject is identified as potentially having had a stroke or AMI by

10 observing one or more symptoms of a stroke or AMI without determining the cause of the stroke prior to the administration of the first composition; the first composition comprises 2 mg to less than 5 mg of tissue plasminogen activator (tPA) and is for administration in a dosage regime of a bolus, the second composition is for administration by an intravenous infusion in a dosage regime of 60 to 120 mg/hour for 60 to 90 minutes following the administration of the first composition, the pro-urokinase mutant (mproUK) comprises a
15 substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase; and a maximum clot lysis rate is achieved with minimal associated hemorrhagic side effects;

- a kit comprising: a first composition in a first container comprising 2-5 mg of tissue plasminogen activator (tPA), formulated for administration as a bolus; and a second composition in a second container comprising 60-120 mg of a pro-urokinase mutant (mproUK) comprising a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase; and

20 - a kit for use in treating a subject with symptoms of a stroke or acute myocardial infarction (AMI) at a maximum rate of clot lysis and with minimal associated hemorrhagic side effects, the kit comprising: a first composition comprising 2 mg to less than 5 mg of tissue plasminogen activator (tPA), wherein the first composition is for administration to a subject in a dosage regime of a bolus; and a second composition comprising a pro-urokinase mutant (mproUK) comprising a substitution of histidine for lysine at
25 amino acid position 300 (Lys300→His) of pro-urokinase, wherein the second composition is for administration to a subject by an intravenous infusion in a dosage regime of 60 to 120 mg/hour for 60 to 90 minutes following the administration of the first composition; wherein the subject is identified as potentially having had a stroke or AMI by observing one or more symptoms of a stroke or AMI without determining the
30 cause of the stroke prior to the administration of the first composition; and wherein a maximum clot lysis rate is achieved with minimal associated hemorrhagic side effects.

DESCRIPTION OF THE DRAWINGS

FIG. 1A is a line graph showing the lysis of fluorescein-labeled plasma clots by tPA at three doses: 1, 2, or 3 $\mu\text{g}/\text{ml}$. The time used by tPA to lyse 50% of the plasma clots at a saturated dose is 60 minutes. FIG. 1B is a bar graph showing remaining fibrinogen at the end of clot lysis, expressed as percentage of the baseline (BL).

FIG. 2A is a line graph showing the lysis of fluorescein-labeled plasma clots by mproUK (M5) at three doses: 10, 12.5, or 15 $\mu\text{g}/\text{ml}$. The time used by M5 to lyse 50% of the plasma clots at a saturated dose is 50 minutes. FIG. 2B is a bar graph showing remaining fibrinogen at the end of clot lysis, expressed as percentage of the baseline (BL). FIG. 2C is a line graph showing the time used by mproUK (M5) to lyse fluorescein-labeled plasma clots was unaffected by the C1-inhibitor (750 $\mu\text{g}/\text{ml}$). FIG. 2D is a bar graph showing fibrinogenolysis by M5 was prevented by the C1-inhibitor (750 $\mu\text{g}/\text{ml}$).

FIG. 3 is a line graph showing maximum clot lysis rate by a combination of tPA (0.2 $\mu\text{g}/\text{ml}$) and M5 (6 $\mu\text{g}/\text{ml}$) (circle). The combination induced clot lysis much faster than 0.2 $\mu\text{g}/\text{ml}$ tPA (square) or 6 $\mu\text{g}/\text{ml}$ M5 (triangle) alone. Results are representative of a single experiment performed in triplicate.

FIG. 4A is a line graph showing clot lysis by a combination of tPA (0.2 $\mu\text{g}/\text{ml}$) and M5 (6 $\mu\text{g}/\text{ml}$) (circle). The combination induced clot lysis much faster than 0.2 $\mu\text{g}/\text{ml}$ tPA (square) or 6 $\mu\text{g}/\text{ml}$ M5 (triangle) alone. FIG. 4B is a bar graph showing the remaining plasma fibrinogen at the end of clot lysis, expressed as percentage of baseline (BL) fibrinogen. Results are representative of 10 experiments performed in triplicate.

FIG. 5A is a line graph showing maximum clot lysis rate by a combination of tPA (0.2 $\mu\text{g}/\text{ml}$) and M6 (6 $\mu\text{g}/\text{ml}$) in addition to C1-inhibitor (750 $\mu\text{g}/\text{ml}$). The addition of C1-inhibitor to the combination did not inhibit lysis, however it did inhibit lysis by tPA alone. FIG. 5B is a bar graph showing the remaining plasma fibrinogen at the end of clot lysis,

expressed as percentage of baseline (BL) fibrinogen. The C1-inhibitor attenuated fibrinogenolysis.

FIG. 6 is a line graph showing tPA doses higher than 0.2 $\mu\text{g}/\text{ml}$ do not further enhance clot lysis rate when combined with 6 $\mu\text{g}/\text{ml}$ M5.

5 FIG. 7 is a bar graph showing the average time required to lyse 50% of the plasma clots by the combination of 0.2 $\mu\text{g}/\text{ml}$ tPA and 6 $\mu\text{g}/\text{ml}$ M5 (“tPA+M5”), 0.2 $\mu\text{g}/\text{ml}$ tPA (“tPA 0.2”), 6 $\mu\text{g}/\text{ml}$ M5 (“M5 6”), 15 $\mu\text{g}/\text{ml}$ M5 (“M5 15”), 3 $\mu\text{g}/\text{ml}$ tPA (“tPA 3”). Data are pooled from multiple experiments (the number of experiments is shown above each bar).

10 FIG. 8 is a bar graph showing the remaining fibrinogen (% BL) at the end of clot lysis by a combination of 0.2 $\mu\text{g}/\text{ml}$ tPA and 6 $\mu\text{g}/\text{ml}$ M5 under the different conditions tested, namely, (1) in 2 ml plasma; (2) in 2 ml plasma plus C1-inhibitor (500 $\mu\text{g}/\text{ml}$); and (3) in 5 ml plasma. Increases in plasma volume naturally decrease fibrinogen degradation (plasma volumes in human subjects are about 3000 ml on average) and thus the use of a C1-inhibitor is not likely required *in vivo*, but is useful in *in vitro* studies to better mimic *in vivo* conditions.

15 FIG. 9 is a line graph showing that in the absence of a blood clot, no fibrinogenolysis by the combination of 0.2 $\mu\text{g}/\text{ml}$ tPA and 6 $\mu\text{g}/\text{ml}$ M5 occurred for at least 3 hours, which suggests that in the absence of a clot *in vivo* the combination would also be inactive.

20 FIG. 10A is a bar graph showing the hematoma volume following intracerebral hemorrhage in a rat model. Hematoma volume was measured by hemoglobin content at 2 hours post dosing. No significant difference ($p = 0.5194$) was determined between the animals dosed with C1 inhibitor/M5 and those dosed with saline/saline. The C1-inhibitor was included since the rat is missing this inhibitor of M5/UK.

25 FIG. 10B is a bar graph showing the histological determination of hemorrhage volume following intracerebral hemorrhage in a rat model. No significant difference ($p = 0.6899$) in hemorrhage volume was determined between the animals dosed with C1 inhibitor/M5 and those dosed with saline/saline.

DETAILED DESCRIPTION

30 The present disclosure is based, at least in part, on the discovery that thrombolysis can be achieved at a maximum clot lysis rate to achieve increased blood flow, e.g., at a Thrombolysis In Myocardial Infarction (TIMI) score of 2 or better within about 75 minutes and without the expected hemorrhagic risk by administering a combination of a mini-dose of

tissue plasminogen activator (tPA) and a low dose of pro-urokinase mutant (“pro-UK mutant” or “mproUK”), for example, a mproUK that comprises a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase, referred to herein as “M5.” For patients with stroke or acute myocardial infarction, the time needed for thrombolysis and 5 reperfusion is critical to clinical outcome and survival rate. To make this possible, treatment must also be safe enough so that patients can be treated before hospitalization. Provided herein are methods of treating a subject who potentially had a stroke or acute myocardial infarction.

These methods can be used to treat patients with symptoms of stroke or acute 10 myocardial infarction without the delay caused by time-consuming diagnostic procedures, thereby enhancing the chances of better clinical outcome and survival rate for those patients. Since fibrinolysis can be achieved with mini-doses of tPA and low doses of mproUK in the combination, non-specific activation of plasminogen and the hemophiliac-like side effects 15 associated with high dose plasminogen activator regimens can be reduced to minimal levels while still achieving a maximum clot lysis rate. Also provided herein are kits that include a first composition comprising tPA and a second composition comprising mproUK, for use in the methods described herein.

Blood clots are lysed in three steps by the protease plasmin, which is the activated 20 form of plasminogen. Step 1: plasminogen present in the blood plasma binds to the intact fibrin clot at a specific binding site on the D-domain that is adjacent to the tPA binding site. tPA activates plasminogen in this so-called “ternary complex”, which consists of fibrin, 25 plasminogen, and tPA, and initiates fibrinolysis. Step 2: plasmin cleaves fibrin preferentially after a lysine residue, creating carboxy terminal lysines which represent new plasminogen binding sites. One of these newly created binding sites is a high affinity binding site consisting of three C-terminal lysines in the fibrin E-domain. When plasminogen binds to 30 this site in the fibrin E-domain, it undergoes a special conformational change which enables it to be activated by the intrinsic catalytic activity of pro-UK and mproUK. Step 3: activation of the plasminogen by pro-UK/mproUK is accompanied by reciprocal activation of pro-UK/mproUK by plasmin to the enzyme urokinase (UK)/mUK. UK/mUK then activates the remaining fibrin-bound plasminogen, thereby completing fibrinolysis (see US Patent Nos. 5,055,295; 5,472,692; 5,626,841; 5,759,542; 7,074,401; 7,837,992; 8,187,592; Pannell, J.

Clin. Invest. 81: 853-859, 1988; Zarich, J Am Coll Cardiol 1995; 26: 374-379; Lee, AJNR Am J Neuroradiol 25: 1470-1475, 2004).

The present inventor has discovered how the new combination therapy methods described herein are supported by the steps of fibrinolysis described above. Only tPA can 5 efficiently perform step 1, since only tPA binds to fibrin, forms a ternary, complex with plasminogen and is specifically promoted by intact fibrin or by fibrin fragment D. Pro-UK/mproUK does not bind to fibrin *in vivo* under normal conditions (i.e., it can bind to fibrin only at very high, non-specific doses). Thus, step 2 can be performed efficiently only by pro-UK/mproUK, because it has a high substrate affinity for the plasminogen confirmation that 10 forms when it binds to the fibrin E-domain of degraded fibrin. Fibrin fragment E has no effect on tPA, which does not bind to this domain, and promotes plasminogen activation only by proUK/mproUK. Thus, step 3 is also promoted only by pro-UK/mproUK, because, when 15 proUK activates plasminogen bound to the fibrin E-domain, there is reciprocal activation of proUK to UK by plasmin and UK completes lysis by activating the remaining fibrin-bound plasminogen (by contrast, tPA undergoes no activation, because its one and two-chain forms have identical activities). The only way tPA can perform steps 2 and 3 is at high non-specific 20 doses that are associated with a high bleeding risk.

The present inventor has discovered that fibrinolysis can be attained with a combination of a mini-dose of tPA (only 2-5% of the standard 100 mg dose) plus a low dose 20 of a mproUK, e.g., M5 (40-50% of the mono-therapy dose) at a maximum rate of clot lysis with a minimal level of fibrinogen degradation of less than about 30% (e.g., less than about 25 or 20%). This tPA-mproUK combination achieves fibrinolysis at a maximum lysis rate 25 that is at least the same as the maximum rate of lysis that can be achieved by monotherapy with either tPA or mproUK at a maximum dose, but with far lower levels of fibrinogen degradation that are safe and clinically acceptable, as no bleeding risks arise at these lower levels, which is not possible with known monotherapies. Thus, the present methods described herein provide thrombolysis that is clearly superior to any alleged synergistic combination, which would not include the newly discovered dual benefits of a maximum 30 blood clot lysis rate with maximum safety for the patient, i.e., minimal associated hemorrhagic side effects.

The “maximum lysis rate” of blood clots is defined herein as a rate of lysis of blood clots at least as fast as what can be achieved by monotherapy with tPA, proUK or mproUK at

a maximally effective dosage (i.e., a dose of the drug at which no faster clot lysis can be achieved by adding additional drug, e.g., a plateau in the clot lysis rate is shown at the maximally effective dosage). However, it is important to note that the maximum lysis rate achieved using monotherapy, e.g., with tPA or M5, would cause significant bleeding in a patient, as indicated, for example, by a level of fibrinogen degradation of greater than 30%. Surprisingly, the “maximum safety” achieved by the new methods described herein is defined as a fibrinogen degradation of less than 30%, which would not be associated with excess bleeding and would be clinically acceptable.

For example, the tPA-mproUK combination can lyse 50% of the mass of at least one blood clot, e.g., multiple blood clots, in less than an hour, e.g., 48, 50, 55, 60, 65, 70, 75, or 80 minutes. The combination also allows significantly lower doses to be used, which are far more fibrin-specific, safer, and more economical. mproUK, e.g., M5, is used instead of native pro-UK because mproUK is more stable in plasma at therapeutic doses and remains in its fibrin-specific pro-enzyme form, while native pro-UK tends to spontaneously convert into urokinase, which is a non-specific plasminogen activator that can cause bleeding complications. This is why pro-UK was denied market approval by the EMEA in Europe, following which its development also ceased in this country.

As shown in the examples below, when incubated in plasma in the absence of a blood clot, the very low doses used in the tPA-mproUK combination do not induce any degradation of fibrinogen, and it is thus expected that the same combination will have little or no effect on coagulation or wound healing. In addition, since M5 is a proenzyme and requires fibrin for its activation, it is expected to be safe even in the presence of bleeding, e.g., during a hemorrhagic stroke. Furthermore, C1-inhibitor, which is available as a pharmacological agent, can be used to quench any non-specific activity of any mUK that may be formed during fibrinolysis (Pannell, J Thromb Haemost. 5(5):1047-54, 2007), e.g., in tests done *in vitro* as well as in therapy *in vivo*, if required. This C1-inhibitor effect is not shared by UK, and is unique to the proUK mutant.

Clinically, the tPA-mproUK combination therapy can be used to treat patients with stroke or heart attack symptoms without the delay caused by time-consuming diagnostic procedures, which are currently mandatory due to the significant bleeding risk associated with tPA in treating stroke. Treatment can be administered on suspicion or in the ambulance. Since time is of the essence for survival and clinical outcome of those patients, the

combination therapy offers better efficacy and outcome than tPA as a monotherapy, which is the only therapy currently approved and available. The therapeutic benefit of tPA monotherapy in treating stroke remains controversial, and its authorization has been recently questioned (Sandercock P, Lancet. 2014 Aug 23; 384(9944):660-1). The mproUK 5 preferentially activates plasminogens in the “bad” occlusive clots while sparing plasminogens in the “good” wound-sealing clots. Therefore, there is little risk of opening up a hemostatic site. By contrast, tPA targets such site which are made up of intact fibrin, as described above in Step 1 of fibrinolysis. Free plasminogen in the plasma is also protected by the action of C1-inhibitor, so there is less risk of inducing a hemophilia-like state.

10 The maximum clot lysis rate of the present methods can be determined *in vivo* can be determined, for example, by standard techniques for assessing blood vessel recanalization. For example, the grade of blood vessel occlusion can be assessed in analogy to the Thrombolysis In Myocardial Infarction (TIMI) score, wherein a TIMI score of 0 is complete occlusion, TIMI of 1 is minimal perfusion, TIMI of 2 is partial flow (recanalization), and a 15 TIMI score of 3 is complete flow. The TIMI study group developed this grading scale for coronary blood flow based on visual assessment of the rate of contrast opacification of the infarct artery (see, e.g., The TIMI Study Group. The Thrombolysis in Myocardial Infarction (TIMI) trial: phase I findings. N Engl J Med. 1984;309:523-530; and The TIMI Study Group. Comparison of invasive and conservative strategies after treatment with intravenous tissue 20 plasminogen activator in acute myocardial infarction: results of the Thrombolysis in Myocardial Infarction (TIMI) phase II trial. N Engl J Med. 1989;320:618-627).

25 The TIMI flow grade has become the standard for semi-quantitative evaluation of myocardial perfusion before and after coronary reperfusion therapies as well as for determining therapies for stroke. Both TIMI flow grades 2 and 3 have been considered indicative of successful reperfusion. Thus, as used herein, a maximum clot lysis rate is expected to be associated in human subjects or patients with achieving a TIMI score of 2 or better within about 75 minutes or less, e.g., within 70, 65, 60, 55, 50, 45, 40, 35, or 30 minutes.

30

Plasminogen Conformations and Distinction between “Good” and “Bad” Blood

Clots

Plasminogen can take on at least three different conformations before converting to active plasmin. The first conformation is the native “closed” conformation, i.e., unbound to any fibrin, and plasminogen exists in the blood in this first conformation. Urokinase can activate plasminogen in this first conformation and may cause non-specific hemorrhagic diathesis, i.e., a hemophilia-like state. Since pro-UK at therapeutic doses is unstable and readily converts to urokinase at therapeutic concentration, pro-UK can also cause hemophilia-like side-effects.

When bound to fibrin, plasminogen can adopt two or three different “open” conformations, providing a basis to distinguish the good wound-healing clots from bad occlusive clots. The first of these is the plasminogen conformation that takes place when plasminogen binds to an internal lysine in the D-domain of intact fibrin. The second of these is the plasminogen conformation that occurs when plasminogen binds to the three C-terminal lysines on fibrin fragment E, which is exposed only after some fibrin degradation as described above in Step 2 of fibrinolysis.

When hemostatic fibrin forms to seal an injury, it acts like a bandage and causes no interference with blood flow. The intact fibrin in such a hemostatic clot has only an internal plasminogen-binding site located in the D-domain, which induces the first of these fibrin-bound conformations. In this conformation, plasminogen is susceptible to activation by tPA, whose fibrin binding site is adjacent, but not by pro-UK, which does not bind to fibrin and has substrate binding due to the second conformation induced by the fibrin E-domain.

When a thrombus forms in a blood vessel, it impedes or arrests blood flow and triggers the local release of tPA from the occluded blood vessel wall and leads to fibrin degradation. Fibrin degradation exposes a new plasminogen-binding site on fibrin fragment E. Plasminogen binds to this new binding site on fibrin fragment E. Pro-UK preferentially activates plasminogen bound to fibrin fragment E.

Utilization of this important distinction between the “good” and the “bad” fibrin clots can effectively achieve thrombolysis while protecting hemostatic plugs that seal injuries. Pro-UK mutant has the same mode of action as pro-UK.

A third fibrin-bound plasminogen conformation exist when plasminogen binds to a single C-terminal site, believed to be on the fibrin gamma chain.

Tissue Plasminogen Activator

Tissue plasminogen activator is a serine protease stored in endothelial cells lining the blood vessel wall. When a thrombus occludes a blood vessel, tPA is released from the blood vessel wall and lyses fibrin clots.

5 Currently most therapeutic thrombolysis is performed using tissue plasminogen activator (tPA) and its derivatives, however, tPA can cause hemorrhagic side effects. For example, tissue plasminogen activator (tPA) at a dose of 150 mg has been shown to induce superior coronary thrombolysis, but has been accompanied by an unacceptable incidence of intracranial hemorrhage, obliging the adoption of a less effective dose of 100 mg (Braunwald, 10 J Amer Coll Cardiol. 9: 467, 1987; Grossbard, J Amer Coll Cardiol. 9:467, 1987). In comparative clinical trials in acute myocardial infarction (AMI) patients, results with percutaneous coronary intervention (PCI) were significantly better than intravenous administration of tPA, although PCI is more costly, technically demanding, and time-consuming. This clinical outcome was surprising, but can be explained by the following tPA 15 properties: (1) the therapeutic dose of tPA is limited by the intracranial hemorrhage complications; and (2) tPA's efficacy is undermined by a relatively high coronary rethrombosis rate, which is associated with hematological evidence of thrombin generation (Verheugt, J Am Coll Cardiol 1996, 27: 766-773; Gurewich, Circulation 1993, 87: 1759-1761; Rapold, Blood 1991, 78: 1490-1495; Gulba, Lancet 1988, 2: 97; Gulba, Circulation 20 1991, 83: 937-944).

In ischemic stroke, a further dose reduction was required due to a 20% incidence of intracranial hemorrhage complications when administering tPA at doses equivalent to those used in AMI (Hacke, JAMA 1995, 274: 1017-1025). The use of heparin, which is used with tPA in AMI, is precluded in stroke, as reocclusion rates of 14-31% have been reported 25 (Alexandrov, Neurology 2002, 59: 862-867; Rubiera, Stroke 2005, 36: 1452-1456; Saqqur, Stroke 2007; 38: 69-74). The net result has been that only about 2-5% of patients with ischemic stroke are treated with tPA in the United States (Kleindorfer, Stroke 2008; 39: 924-928).

tPA-induced bleeding is believed to be primarily related to lysis of hemostatic fibrin 30 needed to repair injury sites in the vessel wall, which are usually occult and unpredictable, but tPA dose-dependent. proUK/M5 spares these sites due its different mode of action, as described above.

Pro-Urokinase and Pro-Urokinase Mutants

Pro-urokinase (pro-UK) is less well-known as a thrombolytic drug, but Phase 3 clinical studies in acute myocardial infarction have been completed (Michels R, J Thromb Thrombolysis 1995, 2: 117-124; PRIMI Trial Study Group. Lancet 1989, 1: 863-867; Tcbbc U, J Am Coll Cardiol 1998, 31: 487-493). Pro-UK induced little (5%) or no coronary rethrombosis and no hematological evidence of thrombin generation in these studies (PRIMI Trial Study Group. Lancet 1989, 1: 863-867; Weaver, J Am Coll Cardiol 1994, 241: 242-1248). Unfortunately, at therapeutic doses, pro-UK became vulnerable to spontaneous activation into the enzyme form, two-chain urokinase (tcUK), in plasma. When this occurred, it incurs a bleeding risk, and for this reason, marketing approval was denied and pro-UK development was abandoned in the West.

The instability of pro-UK was related to its relatively high intrinsic catalytic activity. Structure-function studies revealed the charged residues in a flexible loop consisting of amino acid residues 297-313 in the catalytic domain are responsible for this activity. Mutagenesis in the flexible loop region resulted in modulation of the intrinsic activity of pro-UK. Exemplary pro-UK “flexible loop” mutants with reduced intrinsic catalytic activity are described in U.S. Pat. No. 5,472,692, such as Gly299→Ala mutant, Lys300→His mutant (known as “M5” or “M5 mutant”), Lys300→Ala mutant, and Glu301→His mutant.

One of these pro-UK “flexible loop” mutants, M5 (Lys300→His), has been tested both *in vitro* and *in vivo*, and was shown to dissolve blood clots much faster than native pro-UK (Liu et al., Circulation Research, 90:757-763, 2002). The intrinsic activity of the single-chain M5 mutant is five-fold lower than pro-UK, so M5 is more stable in blood than native pro-UK and less likely to spontaneously convert into active enzyme form and cause hemophilia-like side effects (Liu, Biochemistry 1996, 35: 14070-14076). The activity of the two-chain enzymatic form of mproUKs, e.g., M5, and the mode of action of mproUKs, e.g., M5, remain the same as native pro-UK (Sun Z, J Biol Chem 1997, 272: 23818-23823; Liu, Circu. Res. 2002, 90: 757-763). mproUKs like M5 possess another superior property; they can be inhibited by endogenous plasma C1-inhibitor, providing protection against non-specific side effects without interfering with fibrinolysis by mproUK (Gurewich, J Thrombos Haemost, 2006, 4: 1559-1565; Pannell, J Thromb Haemost, 2007, 5: 1047-1054; Gurewich, Thromb Haemost, 2009, 102: 279-286). Importantly, mproUKs such as M5 do not show any

hemorrhagic side effects normally associated with thrombolytic agents as described in U.S. Pat. No. 7,074,401. In addition, mproUKs such as M5 can be synthesized according to the methods described in U.S. Pat. No. 7,070,958.

5 M5 and other mproUKs are expected to be safe for human administration, because (1) they are essentially a natural human protein (99.8% similarity to native pro-UK), (2) they are free of antigenic (immunologic) reactions, and (3) naturally occurring human pro-UK and recombinant human pro-UK from *E. coli* have already been safely administered to about 5,000 human patients in Phase III clinical studies.

10

Combination Therapy of Low Dose tPA and mproUK

By utilizing the complementary mechanism of action of tPA and pro-UK on plasminogen action, the present inventor has demonstrated that fibrinolysis at a maximum clot lysis rate can be attained with a combination of a mini-dose of tPA (at 2-5% of the standard 100 mg dose, e.g., 1.0, 2.0, 2.5, 3.0, 3.5, 4.0, or 4.5 mg bolus) plus an infusion of mproUK, e.g., M5, (at 40-50% of monotherapy dose, e.g., 60 to 120 ug/ml infused over 60-90 minutes) giving a maximum clot lysis rate, which is at least as fast as the maximum clot lysis rate that can be achieved by monotherapy with either tPA or mproUK alone at their maximal effective dose (FIGs. 1A to 5B). For example, the tPA-mproUK combination can lysis 50% of the mass of blood clots in less than an hour, e.g., 48 minutes, on average (see FIG. 7, tPa + MF). Although tPA and mproUK (M5) alone can achieve similar lysis time at very high doses, e.g., tPA at 3 μ g/ml or mproUK (M5) at 15 μ g/ml (FIG. 7), non-specific activation of plasminogen and resultant fibrinogen degradation of about 77% for M5 (see FIG. 2B) and about 80% for tPA (see FIG. 1B) can occur at those doses and cause significant and clinically unacceptable hemophiliac-like side effects.

20 mproUK such as M5 is used instead of native pro-UK, because mproUK is more stable in plasma and remains in proenzyme form while native pro-UK tends to spontaneously convert into urokinase and cause hemophiliac-like side effects. Thus, the maximum clot lysis rate can be achieved by the tPA-mproUK combination with only a fraction of the monotherapy doses of tPA and mproUK.

25 When incubated in plasma in the absence of a blood clot, the tPA-mproUK combination does not induce any degradation of fibrinogen (FIG. 9), and thus is expected to

have no effect on coagulation and wound healing *in vivo*. Furthermore, C1-inhibitor can be used to quench any non-specific activity of mUK (the enzymatic form of mproUK) in plasma (FIGs. 2D and 6), adding further protection against bleeding complications.

Clinically, the tPA-mproUK combination therapy can be used to treat patients with
5 stroke or heart attack symptoms without the delay caused by time-consuming diagnostic
procedures. Provided herein are methods of treating a subject with symptoms of stroke or
acute myocardial infarction by (a) identifying a subject who potentially had a stroke or acute
myocardial infarction by observing one or more symptoms without determining the
underlying cause of the stroke or acute myocardial infarction; and (b) administering to the
10 subject a bolus of a first composition comprising 5 mg or less of tPA followed by an infusion
of a second composition of mproUK at a rate of 60 to 120 mg/hour infused over 60 to 90
minutes. The second composition can be administered about 5, 10, or 15 minutes after the
administration of the first composition. The first composition and the second composition are
administered in an amount to lyse any blood clot that causes the symptoms of stroke or acute
15 myocardial infarction at a maximum lysis rate. The first composition can include 2 to 5 mg
of tPA, e.g., about 2 mg, about 2.5mg, about 3 mg, about 3.5mg, about 4 mg, about 4.5mg,
about 5 mg. The second composition that includes an mproUK like M5 can be administered
by intravenous infusion. The intravenous infusion of mproUK like M5 can be performed at a
dose of about 60-120 mg/hour (e.g., 60-100, 70-90, or 75-85 mg/hour) for 60, 70, 80, or 90
20 minutes.

The methods described herein can be used to treat stroke. A stroke can be ischemic or
hemorrhagic. Ischemic stroke is caused by a thrombus obstructing blood flow, while
hemorrhagic stroke is caused by a broken blood vessel. About 85% of the time, a stroke is
ischemic, e.g., caused by a blood clot, and is therefore amenable to treatment by a
25 thrombolytic agent. The timing of reperfusion after an ischemic stroke is critical, because the
longer the brain cells are without oxygenated blood, the more brain cells are lost. However, it
is difficult and time consuming to fully diagnose the cause of a stroke, yet an accurate
diagnosis is critical for treatment with currently available thrombolytic agents due to the high
risk of hemorrhagic side effects. Administering a thrombolytic agent to an ischemic stroke
30 patient may be a proper therapy, but administering the same thrombolytic agent to a
hemorrhagic stroke patient will exacerbate the problem and can kill the patient. Although it
takes time to confirm a diagnosis, the basic symptoms of stroke exhibited by a person (such

as sudden onset of one-sided paralysis) can be readily determined by one skilled in the medical field, such as an EMT, a nurse, or a doctor, or even a layperson with minimal training.

A subject with symptoms of stroke can be treated using the methods described herein 5 by administration of a bolus of a low dose of tPA followed by an infusion of mproUK. The very low dose of tPA in the combination reduces potential hemorrhagic risk. Since the tPA in the combination is a mini-dose and the mproUK can lyse a thrombus but spare hemostatic fibrin, the combination can be used to treat patients with a possible ischemic stroke safely with little risk of aggravating bleeding in the brain. Thus, it is possible to initiate treatment in 10 the ambulance on the basis of a clinical suspicion of the diagnosis.

The methods described herein can also be used to treat heart attack, e.g., acute myocardial infarction. A heart attack occurs when one of the coronary arteries is blocked, e.g., by a blood clot. The timing of reperfusion after a heart attack is critical, because the longer the heart muscle is without oxygenated blood, the more muscle cells are damaged or 15 lost. The present treatment of choice is by catheterization and angioplasty, which requires hospitalization, an available catheterization room and staff at the ready. This delays treatment and carries a high cost. This first one hour after a coronary occlusion has been called the “Golden Hour,” because it is the time during which the maximum salvage of heart muscle and the maximum reduction in mortality is possible. Pretreatment with tPA to gain time 20 before catheterization has been generally abandoned since multiple studies have shown that tPA significantly increase the complication rate with catheterization.

A subject with symptoms of a heart attack can be treated by administration of a bolus of a low dose of tPA followed by an infusion of an mproUK such as M5 as described herein. Experience has shown that pretreatment with pro-UK is not associated with post 25 catheterization complication, so that pretreatment with the combination should be well tolerated and may also reduce the need for the subsequent catheterization.

C1-inhibitor

C1-inhibitor is a 104 KD serine protease inhibitor with a normal plasma concentration 30 of about 250 µg/ml and a half- life of about 28 hours. Deficiency of this protein has been associated with a disease called hereditary angioedema. C1-inhibitor has long been

administered clinically for the treatment of hereditary angioedema. Commercially available C1-inhibitors include CINRYZE®, CETOR®, BERINERT®, and RUCONEST®.

The active enzymes two-chain UK or mUK generated during fibrinolysis are eventually released into the plasma where plasminogen activation becomes a liability.

5 Quenching this activity requires plasma inhibitors. As described in U.S. Pat. No. 7,837,992, an inhibitor complex of C1-inhibitor appeared within minutes of the incubation of enzyme mUK with human or dog plasma. C1-inhibitor is present endogenously in human body and can be supplemented pharmacologically. The endogenous C1-inhibitor was shown to be very effective in quenching activity of mUK, e.g.,
10 tcM5, but not UK activity (Gurewich V, J Thrombos Haemost, 2006; 4: 1559-1565; Pannell R, J Thromb Haemost, 2007; 5: 1047-1054; Gurewich, Thromb Haemost, 2009; 102: 279-286). As shown in U.S. Pat. No. 7,837,992, the C1-inhibitor by inhibiting mUK effectively stabilized the mproUK like M5 in plasma and allowed a higher concentration of mproUK like M5 to be tolerated without compromising fibrin-specificity.

15 The methods of treating a subject with symptoms of stroke or acute myocardial infarction disclosed herein can further include administration of a third composition comprising a C1-inhibitor to the subject. The C1-inhibitor can be administered as a bolus in an amount sufficient to establish a concentration of C1-inhibitor in the subject's plasma that is within the range of two to three times above normal physiological level of C1-inhibitor
20 (about 250 µg/ml), i.e., about 500-750 µg/ml. For example, the third composition can include 500-1500 mg of C1-inhibitor. In some embodiments, the third composition can be administered to the subject before the administration of the second composition. In some embodiments, the third composition can be administered to the subject simultaneously with the second composition.

25

Pharmaceutical Compositions, Dosage Regimens, and Methods of Administration

Pharmaceutical compositions provided herein can include specific dosages of tPA and a pro-UK mutant such as M5 as active ingredients. The active ingredient of a pharmaceutical composition, e.g., tPA or mproUK, can be formulated for delivery by intravenous injections.
30

Methods of formulating suitable pharmaceutical compositions are known in the art, see, e.g., Remington: The Science and Practice of Pharmacy, 21st ed., 2005; and the books in

the series Drugs and the Pharmaceutical Sciences: a Series of Textbooks and Monographs (Dekker, NY). For example, solutions or suspensions used for parenteral administration can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerin, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injection can include sterile aqueous solutions (where water soluble), dispersions, and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ), or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants.

Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, and sodium chloride in the composition. Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle, which contains a basic dispersion medium and the required other ingredients from those enumerated

above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying, which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

5 The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

Dosage regimens can be adjusted to provide the optimum therapeutic response. See e.g., Physicians' Desk Reference, 63rd edition, Thomson Reuters, November 30, 2008. For example, Dosage, toxicity and therapeutic efficacy of the therapeutic compounds can be 10 determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit high therapeutic indices are preferred. While compounds that 15 exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies 20 preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma 25 concentration range that includes the IC50 (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

The tPA bolus can include 2-5 mg of tPA, e.g., about 2 mg, about 2.5 mg, about 3 mg, 30 about 3.5 mg, about 4 mg, about 4.5 mg, or about 5 mg. The intravenous dose of mproUK, e.g., M5, can be 60-120 mg/hour (e.g., 60-100, 70-90, or 75-85 mg/hour) for 60, 65, 70, 75, 80, 85, or 90 minutes. The C1-inhibitor can be administered as a bolus in an amount

sufficient to establish a concentration of C1-inhibitor in the subject's plasma that is about 500-750 μ g/ml. For example, the C1-inhibitor bolus can include 500-1500 mg of C1-inhibitor.

5 For patients with stroke or acute myocardial infarction, the time needed to reperfusion is critical to survival and clinical outcome. These findings suggest that a combination of mini-dose tPA and M5 can achieve therapeutic thrombolysis in a safer and more effective way. Thus, patients with stroke or acute myocardial infarction can be treated with a combination of tPA and M5 with little or no delay.

10 Efficacy of thrombolysis is defined by the rate of lysis. However, the clinical utility requires that efficacy be divided by the incidence of bleeding complications from that rate. With the combination therapy described herein, the evidence is that there is no maximum clinical utility, since the optimum combination induces a maximum rate of lysis without significant fibrinogen degradation (see Pannell et al., PLOS ONE, DOI:10.1371/journal.pone.0122018 March 26, 2015).

15 This makes for a far superior clinical utility index, in fact it gives the maximum utility index possible, i.e., the maximum clot lysis rate possible for plasminogen activators without side effects.

Kits

20 Also provided herein are kits that include at least a composition comprising tissue plasminogen activator (tPA) in one container and another composition comprising an mproUK, e.g., M5, in a separate container. The kits are used to carry out the therapeutic methods described herein. The first composition can be formulated suitable for administration as a bolus, and can include 2-5 mg of tPA. The second composition can be 25 formulated suitable for intravenous infusion. The second composition can include 60-120 mg (e.g., 60, 65, 70, 75, 80, 85, or 90 mg) of the pro-UK mutant, for infusion over a time period of 60-90 minutes. The kit can also include a third composition comprising C1-inhibitor. The third composition can be formulated suitable for administration as a bolus, and can include about 500-1500 mg of C1-inhibitor.

30 Kits generally include the following major elements: packaging, reagents comprising binding compositions as described above, optionally a control, and instructions. Packaging can be a box-like structure for holding a vial (or number of vials) containing the

compositions, and instructions for use in a method described herein. Individuals skilled in the art can readily modify the packaging to suit individual needs.

Compositions and kits provided herein can be used in accordance with any of the methods (e.g., treatment methods) described above. For example, compositions and kits containing a composition comprising tPA and another composition comprising a pro-UK mutant, e.g., the M5 mutant, can be used to treat stroke, heart attack, or other cardiovascular diseases caused by a thrombus such as peripheral gangrene. Those skilled in the art will be aware of other suitable uses for compositions and kits provided herein, and will be able to employ the compositions and kits for such uses.

10

EXAMPLES

The invention is further described in the following examples, which do not limit the scope of the invention described in the claims.

Example 1. Maximum Clot Lysis Rate Thrombolysis *in Vitro* by tPA / M5

15 The fibrinolytic and fibrinogenolytic effects of specific combinations of tPA and M5 were studied *in vitro* in a human plasma milieu.

Materials

20 The mproUK that comprises a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase (M5) was prepared from *E. coli* by PxTherapeutics (Grenoble, France). Tissue plasminogen activator (tPA) was obtained from Genentech (South San Francisco, CA). Human Fibrinogen, Kabi Grade L, was obtained from Chromogenix, Milan, Italy. Aprotinin and fluorescein isothiocyanate were from Sigma Chemicals, St. Louis, MO. Thrombin (ThromboMax, 100 NIH U per ml) was obtained from Sigma (St Louis, MO). Clinical grade C1-inhibitor was obtained from CSL Behring, 25 Marburg, Germany. Human outdated blood bank plasma, pooled from four donors, was used in the experiments.

Note that various animal models have different sensitivities to human pro-UK (see, e.g., Table 1 in Gurewich et al., *J. Clin. Invest.*, 73:1731-1739 (1984)).

30 Thus, animal dosages cannot be used to determine human dosages, and the dosages described here are based on experience using the individual components in human tests. *In vitro* studies in human plasma show that 2x as much M5 is needed than proUK for the same lysis rate. The monotherapy dose of proUK/M5 is known

from Phase-3 studies and the newly discovered mini-dose tPA plus the low dose of M5 in the combination therapy methods described herein are based on this 2x factor.

Blood Clot Lysis Experiments

5 Fibrinolysis by M5 was studied in an inhibitor-containing plasma milieu. Blood clots were made by recalcification of 0.2 ml pooled blood bank plasma with 35 mM of calcium in the presence of a trace amount of thromboplastin and a fluoresceinated fibrinogen (see Dmitry V, J Biol Chem, 1996; 271: 2133-2138). The blood clots were then incubated for an hour at 37°C, followed by overnight incubation at room temperature. The following day, the 10 blood clots were placed into 2 ml of blood bank plasma, followed by addition of tPA, M5, or the combination of tPA and M5. Clot lysis was monitored by taking 50 µl samples of the plasma at certain time points and measuring fluorescence emission. The fluorescence emission in the sample represents the amount of fibrin degradation products released from the blood clots. In some experiments, the volume of plasma was increased to 5 ml, and 15 unlabeled fibrinogen was added to make up for the dilution of fluoresceinated fibrinogen. In those experiments, the end of blood clot lysis was determined visually.

Each blood clot lysis experiment was performed in triplicate. Graph Pad Prism was used to prepare graphs and conduct statistical analysis.

20 The lysis curves were plotted as a percentage lysis of blood clots over time. The 100% point was obtained from the mean of the highest readings. The base line reading was obtained at onset by subtracting the amount of fluorescence shown by the plasma (~15% of the full signal) to obtain the zero point. The time to 50% lysis for each experimental condition was determined from the lysis graph and was used as the principal endpoint.

Determination of Remaining Fibrinogen after Blood Clot Lysis

After blood clot lysis was completed, a final plasma sample (1.0 ml) was obtained for determining the remaining fibrinogen. Aprotinin (200 KIU/ml) was added to the sample to prevent further proteolysis. The remaining fibrinogen was recorded as the percentage of the baseline (BL) fibrinogen.

30 Fibrinogen was measured as a thrombin-clottable protein. After dilution of the plasma sample with an equal amount of phosphate-buffered saline, 200 µl of thrombin (1,000 NIH units/ml solution) was added. The solution was mixed gently, and incubated for an hour

at 37°C, followed by overnight incubation at room temperature. The following morning, each blood clot was wound onto a thin, long stemmed plastic transfer pipette tip, to which the gel adhered, and the serum content expressed by pressure against the test tube wall and then against a paper towel. The white fibrin on the pipette stem was then placed in at least 5 ml of 5
saline for at least an hour to allow diffusion of any remaining serum proteins. The fibrin was then peeled off the tip and placed into 1 ml of 5% NaOH, boiled for one minute, and then kept at room temperature until all fibrin had gone into solution. The protein in the solution was measured spectrophotometrically at 280 nm.

10 Determination of the Shortest Time to Lysis by tPA or M5 Alone

The blood clot lysis experiments were performed as described above in the presence of 1, 2, or 3 μ g/ml of tPA. The lysis curves were plotted as a percentage lysis of blood clots over time, and the time used to lyse 50% blood clots for each experimental condition was determined from the lysis curve. The shortest time to lysis (from which the maximum clot lysis rate can be determined) was defined as the time at which there is no further dose-
15 dependent shortening of the lysis time. The results of a representative experiment are shown in FIG. 1A and the shortest time used to lyse 50% blood clots by tPA, was found to be 60 minutes when 3 μ g/ml of tPA was used. Thus, the maximum clot lysis rate is 50% lysis at one hour.

20 FIG. 1B shows the percentage of the remaining fibrinogen of the baseline level at the end of lysis for each tPA dose tested, ranging from 19% to 45%. Thus, tPA alone caused 55%-81% fibrinogen degradation.

25 For M5, the blood clot lysis experiments were performed as described above in the presence of 10, 12.5, 15 μ g/ml of M5. The shortest time used to lyse 50% blood clots by M5, was determined to be 50 minutes, when 15 μ g/ml of mproUK was used (FIG. 2A). Thus, the maximum clot lysis rate is 50% clot lysis at 50 minutes.

The percentage of the remaining fibrinogen of the baseline level at the end of lysis for each M5 dose, ranging from 25% to 55%, is shown in FIG. 2B. Thus, M5 alone caused 45%-75% fibrinogen degradation.

30

Effect of C1-Inhibitor on Fibrinogenolysis by M5

To prevent fibrinogenolysis, 750 μ g/ml of C1-inhibitor was added to the plasma prior to the addition of 10, 12.5, 15 μ g/ml of M5 in the clot lysis experiments. The presence of 750 μ g/ml of C1-inhibitor does not affect the time used to lyse 50% blood clots by M5, which 5 was still 50 minutes (FIG. 2C). However, as shown in FIG. 2D, little fibrinogen degradation occurred in the presence of C1-inhibitor.

The C1-inhibitor experiment was not repeated for tPA, since C1-inhibitor has already been shown to inhibit blood blot lysis by tPA (Tomasi S, PLoS One., 2011; 6: e21999).

10 **Maximum Clot Lysis Rate by a Combination of Mini-Dose tPA and Low Dose M5**

To determine the lowest dose of tPA and M5 that is needed to achieve the shortest lysis time (and maximum clot lysis rate) when used in combination, the blood clot lysis experiments were performed using various combinations and ratios of tPA and M5. The lowest doses of tPA and mproUK when used in combination to consistently achieve the 15 maximum clot lysis rate were determined to be 0.2 μ g/ml of tPA and 6 μ g/ml of M5. These doses corresponded to 6% of the tPA dose required to achieve the shortest lysis time plus 40% of the M5 dose needed to achieve that when they were used alone in monotherapy. FIG. 3 shows the results of a representative clot lysis experiment, where the time used to lyse 50% blood clots by a combination of 0.2 μ g/ml tPA and 6 μ g/ml M5 was 53 minutes, whereas the 20 time used to lyse 50% blood clots by 6 μ g/ml M5 alone was 135 minutes and the time used to lyse 50% blood clots by 0.2 μ g/ml tPA alone was 225 minutes. These results demonstrate that a small dose of tPA, e.g., 0.2 μ g/ml, greatly shortened the clot lysis time by M5. This 25 observation is consistent with tPA's function in initiating fibrinolysis.

The results of ten representative clot lysis experiments are shown in FIG. 4A, using 25 the combination (0.2 μ g/ml tPA + 6 μ g/ml M5) (circle), which induced an average 50% lysis time of approximately 75 minutes. At this dose of M5 alone (triangle), the lysis time was 135 minutes, and tPA alone (square) induced a lysis time of 225 minutes. The findings show that a mini-dose tPA caused about a 45% shortening of the onset of lysis by M5. FIG. 4B shows the plasma fibrinogens at the end of lysis, expressed as percentage baseline (BL) fibrinogen. 30 All experiments were performed in triplicate.

To test if tPA's function in the combination is limited to the initiation of fibrinolysis, blood clot lysis experiments were performed using four tPA-M5 combinations. Each tPA-M5

combination includes a fixed dose of 6 $\mu\text{g}/\text{ml}$ of M5 and a different tPA dose selected from 0.2, 0.6, 1.0, and 3.0 $\mu\text{g}/\text{ml}$. The time used to lyse 50% blood clots is about 48-60 minutes for all four combinations tested and a tPA dose higher than 0.2 $\mu\text{g}/\text{ml}$ does not further shorten the time needed to lyse 50% blood clots (FIG. 6). These findings are consistent with tPA's function in the combination being essential for the initiation of fibrinolysis but not contribute to fibrinolysis beyond the initiation.

The average time used to lyse 50% blood clots *in vitro*, from multiple blood clot experiments were tabulated and compared. FIG. 7 shows that the mean time used to lyse 50% blood clots by the combination of 0.2 $\mu\text{g}/\text{ml}$ tPA plus 6 $\mu\text{g}/\text{ml}$ M5 was 48 (± 2.5) minutes ($n = 10$); the time used to lyse 50% blood clots by 0.2 $\mu\text{g}/\text{ml}$ tPA alone was 156 (± 5.3) minutes ($n = 5$); for 6 $\mu\text{g}/\text{ml}$ of M5 alone, it was 118 (± 7.2) minutes ($n = 7$); for 15 $\mu\text{g}/\text{ml}$ of M5 alone, it was 48 (± 1.6) minutes ($n = 9$); and for 3 $\mu\text{g}/\text{ml}$ of tPA alone, it was 55 (± 5) minutes ($n = 6$). The time used to lyse 50% blood clots by a combination of 0.2 $\mu\text{g}/\text{ml}$ tPA plus 6 $\mu\text{g}/\text{ml}$ M5 is significantly less than monotherapy of 0.2 $\mu\text{g}/\text{ml}$ tPA alone or 6 $\mu\text{g}/\text{ml}$ M5 alone (FIG. 7). Although tPA and mproUK alone can achieve similar clot lysis time at very high doses, e.g., tPA at 3 $\mu\text{g}/\text{ml}$ or M5 at 15 $\mu\text{g}/\text{ml}$ (FIG. 7), non-specific activation of plasminogen can occur at those doses and cause hemophiliac-like side effects.

Effect of Volume and C1-inhibitor on Fibrinogenolysis by the tPA and M5 Combination

During fibrinolysis, mproUK is activated by plasmin and converted to mUK, which can then diffuse into the plasma. Confining proteolysis to the clot becomes a function of plasma inhibitors, e.g., C1-inhibitor. In a test tube, the limited plasma volume *in vitro* relative to the volume *in vivo* may be influential. Therefore, clot lysis experiments were performed under three conditions: (1) a control plasma volume of 2 ml, (2) an increased plasma volume of 5 ml, and (3) 2 ml plasma with 500 $\mu\text{g}/\text{ml}$ C1-inhibitor. Unlabeled blood clots were used for those experiments and the time used to lyse 100% blood clots was determined to be 75-80 minutes in all three conditions tested. In the standard 2 ml plasma volume, the tPA-M5 combination degraded 70% of fibrinogen, reflecting the rapid tcM5 generation rate by plasmin (FIG. 8). When C1-inhibitor was added, fibrinogenolysis was reduced to 45%, with 55% remaining fibrinogen (FIG. 8). A similar effect was observed in 5 ml plasma volume, probably reflecting dilution of tcM5 in the immediate clot environment

(FIG. 8). This volume effect suggests that fibrinogenolysis by the tPA-M5 combination may be further attenuated *in vivo* where the plasma volume to clot ratio is considerably greater.

The more modest effect of C1-inhibitor when compared to FIG. 2D, are related to the more rapid fibrin-dependent plasmin generation induced by the tPA-M5 combination compared with M5 monotherapy. It has been verified by additional studies that the C1-inhibitor's ability to inhibit fibrinogenolysis by the tPA-M5 combination is reduced when compared to its ability to inhibit fibrinogenolysis by M5 alone. C1-inhibitor is a relatively slow inhibitor and thus is less able to quench the more rapidly generated tcM5 from the more rapid fibrinolysis achieved by the tPA-M5 combination. Higher concentration of C1-inhibitor might be needed to sufficiently quench tcM5 generated from the tPA-M5 combination.

The results of representative clot lysis experiments are shown in FIG. 5A, using the combination (0.2 μ g/ml tPA + 6 μ g/ml M5) (circle) in addition to C1-inhibitor. C1-inhibitor (750 μ g/ml) was added 30 minutes after the addition of the activators. As shown in FIG. 5A, this did not inhibit lysis (circles), which on average was shortened to approximately 30 minute, but did inhibit lysis by tPA alone (squares). FIG. 5B shows the plasma fibrinogens at the end of lysis, expressed as percentage baseline (BL) fibrinogen. As shown in FIG. 5B, the C1-inhibitor attenuated fibrinogenolysis. All experiments were performed in triplicate.

Fibrinogenolysis by the tPA-M5 Combination in the Absence of a Clot

To test the effect of the tPA-M5 combination on fibrinogenolysis in the absence of a blood clot, the combination of 0.2 μ g/ml of tPA plus 6 μ g/ml of M5 was incubated with plasma at 37°C and samples were taken for fibrinogen determination after 2-5 hours. As shown in FIG. 9, there was no fibrinogenolysis for at least 3 hours, well beyond the duration of therapeutic thrombolysis. These findings indicate that the plasmin generation induced by the tPA-M5 combination was fibrin-dependent and that in the absence of a blood clot, this fibrinolytic combination does not induce plasminogen activation.

Example 2. Thrombolysis *In Vivo* by a Combination of tPA and M5

Male, mongrel dogs weighing 10-15 kg are anesthetized with pentobarbital sodium and maintained breathing room air. Blood clots are formed from 1 ml of native whole dog blood as described in U.S. Pat. No. 7,074,401, to which radiolabeled fibrinogen (1.9 μ Ci, 0.75mCi/mg protein) and thrombin (10 units) are added. After 20 minutes, the clots are washed with saline

three times and then cut into small (about 1 mm³) pieces and injected through a 16-gauge needle into the femoral vein. After 15 minutes, a blood sample is obtained from a cannula in the contralateral femoral vein for the measurement of baseline radioactivity.

5 The dogs are divided into four groups and injected with (1) saline, (2) a bolus of 2-5 mg of tPA, (3) intravenous infusion of M5 (20 µg/kg/min) for 60 minutes, or (4) a bolus of 2-5 mg of tPA followed by intravenous infusion of M5 (20 µg/kg/min) for 60 minutes. At intervals during the infusions, blood samples were obtained and measured for radioactivity and fibrinogen. Time used to lyse blood clots is determined and compared among the four groups.

10 **Example 3. Characterization of C1 Inhibitor and M5 in a Rat Model of Intracerebral Hemorrhage (ICH)**

The purpose of this study was to investigate the effect of M5 on intra-cerebral hemorrhage (ICH) volume.

15 Twenty adult, male Sprague-Dawley rats were used for the study. Rats were randomly selected for use on surgical days. Rats were given a unique identification number by tail marking. Immediately prior to initiation of surgery Cefazolin Sodium intraperitoneal injection (40 mg/kg; Hospira 101C049) and Buprenorphine subcutaneous (1 mg/kg; Reckitt Benckiser, 219202) were administered to the animals. While the rats were under isoflurane anesthesia (1.5% to 2%) with spontaneous respiration in a nitrous oxide/oxygen mixture (2:1), a small burr 20 hole was drilled, and a 30-gauge 10 microliter microinjection needle (Hamilton, 700 series) was slowly lowered into the right striatum at the following coordinates from the bregma: 0.0 mm anterior, 3 mm lateral, and 6 mm depth. During a period of 3 minutes, 3 microliters of saline containing 0.45 U collagenase VII-S (Sigma, St. Louis, MO) was injected. The needle was left 25 in place for 2 minutes and then slowly removed over 5 minutes. Afterward, the scalp was stapled closed, and the rats were allowed to recover. The whole surgical procedure lasted about 20 minutes for each rat. A heating pad (37 ± 1°C) is used to maintain animal body temperatures.

The C1 inhibitor and the test article (M5) were formulated prior to dosing. Test solutions were kept on ice during the daily usage. The remaining unused solutions were kept at -20°C. Immediately after a bolus at 4 mL/kg (IV), animals were dosed by intravenous infusion (over 30 minutes), starting at 15 minutes following ICH at 4 mL/kg. At 2 hours after the beginning of infusion, animals were humanely killed under isoflurane in 100% N₂O inhalation. Brains were removed and cut in slices according to a 2-mm rat brain matrix. Under standardized

conditions, images of the brain slices were taken with a digital camera. Hematoma size was calculated by ImageJ software (available online at rsb.info.nih.gov/ij).

Two hours after hemorrhage induction, rats were sacrificed under deep (5%) isoflurane anesthesia (100% N₂O). Brains were removed and placed in phosphate-buffered saline (PBS) on ice. Brains were sliced into 7 sections of 2mm each and photographed for a visual assessment of the hematoma. A histological assessment was not conducted on the individual sections. Hemoglobin content was determined as a quantitative measurement of hemorrhagic volume. The hemorrhaged side (inclusive of all 7 sections) was isolated from the normal side and placed into 1.5 mL of cold PBS. After 30 seconds of homogenization (manually with a 5 Polytron PT2100), ultrasound was applied for 1 minute to lyse erythrocytic membranes. After centrifugation for 30 minutes (13000 rpm, 4°C), 200 µL of supernatant was added to 800 µL of Drabkins reagent (Sigma, St. Louis, MO) and allowed to sit for 10 minutes at room temperature. With use of a photometer, absorption rates were determined at 540 nm, and hemorrhagic blood 10 volumes were calculated for the injured half of the brain on the basis of a standard curve. The 15 standard curve was generated by using eight naïve brain hemispheres from age and weight matched (to the average study subject). These naïve brain hemispheres were spiked with pooled blood from the same individuals at increasing volumes from 0 to 192µL. Statistical significance was assessed by analysis of variance (ANOVA).

The timing at which the size of the hemorrhage was measured (2 hours post hemorrhage 20 induction) was determined from previous work optimized at Biotrofix.

Rats were dosed as described with a bolus (4ml/kg) of vehicle (saline) (4 ml/kg) followed immediately with a 30 minute infusion of vehicle, or a bolus of C1 inhibitor (200U/4ml/kg) followed immediately with a 30 minute infusion of M5 (10 mg/4 ml/kg) at fifteen minutes post-ICH.

25 Clinical Observations and Survival

All animals survived the study period for this study. Animal #24 was not included for the analysis as there was insufficient hemorrhage induced in this animal to be included.

Histological Determination of Hemorrhage Volume

30 M5 (immediately) following a C1inhibitor bolus injection was studied in a rat model of ICH. In this model, an intracerebral hemorrhage was induced by injection of collagenase into the right striatum of the brain. Fifteen minutes post-injury M5 (at a dose of 10mg/4ml/kg) or vehicle was administered by a 30 minute intravenous infusion immediately following bolus injections as

described above. No significant difference ($p=0.6899$) was determined between the animals dosed with C1inh/M5 and those dosed with saline/saline (Figure 1).

Direct Hematoma Volume

As with the histological determination for hemorrhage volume, no significant difference (5 $p = 0.5194$) was determined between the animals dosed with C1inh/M5 and those dosed with saline/saline (FIGs. 10A and 10B). FIG. 10A is a bar graph showing the hematoma volume following intracerebral hemorrhage in a rat model. Intracerebral hemorrhage was induced by stereotactic injection of collagenase. Rats were dosed at 15 minutes post injury with a C1inhibitor bolus injection followed immediately with a 30 minute infusion of M5 (at a dose of 10 10 mg/4ml/kg). Hematoma volume was measured by hemoglobin content at 2 hours post dosing. No significant difference ($p = 0.5194$) was determined between the animals dosed with C1inh/M5 and those dosed with saline/saline.

FIG. 10B is a bar graph showing the histological determination of hemorrhage volume following intracerebral hemorrhage in a rat model. Intracerebral hemorrhage was induced by stereotactic injection of collagenase. Rats were dosed at 15 minutes post injury with a C1inhibitor bolus injection followed immediately with a 30 minute infusion of M5 (at a dose of 15 10 mg/4 ml/kg). No significant difference ($p = 0.6899$) in hemorrhage volume was determined between the animals dosed with C1inh/M5 and those dosed with saline/saline.

In this rat model of collagenase induced intracerebral hemorrhage, the administration of 20 C1 inhibitor / M5 administered 15 minutes post ICH induction did not differ in hematoma volume or hemorrhage size relative to saline/saline treated animals. Thus, M5 did not induce or augment bleeding in this stroke model and thus should be safe to use in bleeding events such as hemorrhagic stroke.

25 **Example 4. Clinical Trial of a Combination of tPA and M5**

Healthy male subjects, aged between 18 and 35 years inclusive, and with a body weight of at least 60 kg and a body mass index (BMI) between 18.5 and 25 kg/m² inclusive are enrolled in the clinical trial. The subjects have normal endogenous C1-inhibitor, α 2-antiplasmin, and fibrinogen levels, a negative serology for HIV, HBsAg, and HCV, and a 30 negative test for alcohol and drugs of abuse at screening and on study day 1. The subjects cannot have clinically significant abnormalities.

A subject is not included if he fulfils one or more of the following criteria: (1) the subject has a known or suspected inherited, congenital, or acquired disease or condition that affects the haemostatic or coagulation pathways or that is associated with an increased bleeding tendency; (2) the subject has a reasonable chance of developing a clinically significant bleeding event or a bleeding event that may go undetected for a considerable amount of time during the study, for example, the subject (a) has undergone major (internal) surgery or trauma within the last three months of the anticipated dosing day, (b) has an intestinal or cerebral vascular malformation, or (c) has participated in high impact contact sports, such as kick-boxing, within two weeks of the anticipated dosing day; (3) the subject has received any systemically absorbed drug or substance (including prescription, over-the-counter, or alternative remedies) that is not permitted by this protocol prior to dosing without undergoing a wash-out period of at least seven times the elimination half-life of the product; (4) the subject has smoked tobacco in any form within three months of dosing, or has ever smoked more than five cigarettes per day (or equivalent) on average; (5) the subject has received blood or plasma derivatives in the year preceding the administration day; (6) the subject has lost blood or plasma outside the limits of the local blood donation service (i.e. Sanquin) three months prior to dosing; (7) the subject has a known hypersensitivity to any of the investigational material or related compounds; (8) the subject has a history of severe hypersensitivity or of an allergy with severe reactions; (9) the subject has a history of substance abuse, including caffeine, tobacco, and alcohol; (10) the subject has a condition or demonstrates an attitude that in the opinion of the investigator might jeopardize the subject's health or well-being, or the scientific integrity of the study results; (11) the subject is mentally or legally incapacitated to provide informed consent.

The enrolled subjects are randomly allocated to one of the following treatment arms:

(1) Injection of a bolus of 2.5 mg of tPA followed by intravenous infusion of an mproUK, e.g., M5, for 60-90 minutes at approximately 80 mg/hour (50% of the monotherapy dose);

(2) Injection of a bolus of 2.5 mg of tPA followed by intravenous infusion of a placebo for 60-90 minutes;

(3) Injection of a placebo bolus followed by intravenous infusion of an mproUK, e.g., M5, for 60-90 minutes at approximately 160 mg/hour; or

(4) Injection of a bolus of 2.5 mg of tPA followed by a C1-inhibitor (Berinert[®]) bolus (e.g. 1000 EU (2 vials)), and intravenous infusion of an mproUK, e.g., M5, for 60-90 minutes at approximately 80 mg/hour (50% of the monotherapy dose).

5 The dose for an mproUK ranges from 60-120 mg/hour. The C1-inhibitor bolus consists of 25-100 U/kg of Berinert[®] (based on body weight of the subject). This study ends either when plasminaemia occurs or when sufficient data has been gathered.

10 Based on data gathered from this study, the overall safety and tolerability of the combination of tPA and mproUK such as M5 are evaluated. The effect of a mini-dose of tPA on mproUK-induced coagulation changes are assessed. The effect of a single dose of C1-inhibitor on the overall safety and tolerability of tPA-mproUK combination and its effect on tPA-mproUK-induced coagulation changes are assessed.

OTHER EMBODIMENTS

15 It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

CLAIMS:

1. Use of tissue plasminogen activator (tPA) comprised in a first composition and a pro-urokinase mutant (mproUK) comprised in a second composition for treating a subject with symptoms of a stroke or an acute myocardial infarction (AMI) at a maximum rate of blood clot lysis
5 and with minimal associated hemorrhagic side effects, wherein:

the subject is identified as potentially having had a stroke or AMI by observing one or more symptoms of a stroke or AMI without determining the cause of the stroke prior to the administration of the first composition;

10 the first composition comprises 2 mg to less than 5 mg of tissue plasminogen activator (tPA) and is for administration in a dosage regime of a bolus,

the second composition is for administration by an intravenous infusion in a dosage regime of 60 to 120 mg/hour for 60 to 90 minutes following the administration of the first composition,

the pro-urokinase mutant (mproUK) comprises a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase; and

15 a maximum clot lysis rate is achieved with minimal associated hemorrhagic side effects.

2. The use of claim 1, wherein the subject has symptoms of a stroke.

3. The use of claim 1, wherein the minimal associated hemorrhagic side effects are determined as a level of fibrinogen degradation in the subject's blood of less than about 30 percent.

4. The use of any one of claims 1 to 3, wherein the maximum clot lysis rate is indicated as
20 having been achieved by a Thrombolysis In Myocardial Infarction (TIMI) score of 2 or higher.

5. The use of any one of claims 1 to 4, wherein the maximum clot lysis rate is indicated as having been achieved by a lysis of about 50% of the mass of at least one clot in the subject achieved within 75 minutes.

6. The use of any one of claims 1 to 5, wherein the bolus comprises 2 to 4.5 mg of tPA.

25 7. The use of claim 6, wherein the bolus comprises 2 to 4.0 mg of tPA.

8. The use of claim 6, wherein the bolus comprises 2 mg of tPA.

9. The use of any one of claims 1 to 8, wherein the second composition is for administration as an intravenous infusion at a rate of 60-90 mg/hour of the mproUK for 60-90 minutes.
10. The use of claim 9, wherein the second composition is for administration as an intravenous infusion at a rate of 60-80 mg/hour of the mproUK for 60 minutes.
- 5 11. The use of any one of claims 1 to 10, wherein the administration of the second composition begins within five minutes after the administration of the first composition.
12. The use of any one of claims 1 to 11, wherein the first composition and the second composition together lyse 50% of a mass of at least one blood clot in the subject in less than one hour.
- 10 13. The use of any one of claims 1 to 12, further comprising use of a third composition comprising a bolus of C1-inhibitor.
14. The use of claim 13, wherein the third composition is for administration to the subject before the administration of the second composition.
15. The use of claim 13, wherein the third composition is for administration to the subject at about the same time as the second composition.
16. The use of claim 13, wherein the third composition comprises a bolus of 500-1500 mg of C1-inhibitor.
17. The use of claim 13, wherein the first composition and the second composition together lyse the blood clots in the presence of the C1-inhibitor with less than 30% fibrinogen degradation
- 20 when compared to monotherapy by either tPA or pro-urokinase alone.
18. A kit comprising:
 - a first composition in a first container comprising 2-5 mg of tissue plasminogen activator (tPA), formulated for administration as a bolus; and
 - a second composition in a second container comprising 60-120 mg of a pro-urokinase
- 25 mutant (mproUK) comprising a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase.
19. The kit of claim 18, wherein the second composition is formulated for intravenous infusion.

20. The kit of claim 18, further comprising a third composition comprising 500-1500 mg of C1-inhibitor.

21. The kit of claim 20, wherein the third composition is formulated for administration as a bolus.

5 22. A kit for use in treating a subject with symptoms of a stroke or acute myocardial infarction (AMI) at a maximum rate of clot lysis and with minimal associated hemorrhagic side effects, the kit comprising:

a first composition comprising 2 mg to less than 5 mg of tissue plasminogen activator (tPA), wherein the first composition is for administration to a subject in a dosage regime of a bolus; 10 and

a second composition comprising a pro-urokinase mutant (mproUK) comprising a substitution of histidine for lysine at amino acid position 300 (Lys300→His) of pro-urokinase, wherein the second composition is for administration to a subject by an intravenous infusion in a dosage regime of 60 to 120 mg/hour for 60 to 90 minutes following the administration of the 15 first composition;

wherein the subject is identified as potentially having had a stroke or AMI by observing one or more symptoms of a stroke or AMI without determining the cause of the stroke prior to the administration of the first composition; and

wherein a maximum clot lysis rate is achieved with minimal associated hemorrhagic 20 side effects.

23. The kit of claim 22, for use in treating a subject with symptoms of a stroke.

24. The kit of claim 22 or claim 23, wherein the minimal associated hemorrhagic side effects are determined as a level of fibrinogen degradation in the subject's blood of less than about 30 percent.

25. 25. The kit of claim 22 or claim 23, wherein the maximum clot lysis rate is indicated as having been achieved by a Thrombolysis In Myocardial Infarction (TIMI) score of 2 or higher.

26. The kit of claim 22 or claim 23, wherein the maximum clot lysis rate is indicated as having been achieved by a lysis of about 50% of the mass of at least one clot in the subject achieved within 75 minutes.
27. The kit of any one of claims 22 to 26, wherein the bolus comprises 2 to 4.5 mg of tPA.
- 5 28. The kit of any one of claims 22 to 26, wherein the bolus comprises 2 to 4.0 mg of tPA.
29. The kit of claim 28, wherein the bolus comprises 2 mg of tPA.
30. The kit of any one of claims 22 to 29, wherein the second composition is prepared for administration as an intravenous infusion at a rate of 60-90 mg/hour of the mproUK for 60-90 minutes.
- 10 31. The kit of claim 30, wherein the second composition is prepared for administration as an intravenous infusion at a rate of 60-80 mg/hour of the mproUK for 60 minutes.
32. The kit of any one of claims 22 to 31, further comprising a third composition comprising a bolus of C1-inhibitor.
33. The kit of claim 32, wherein the third composition comprises a bolus of 500-1500 mg of C1-inhibitor.
- 15



FIG. 1A**FIG. 1B**



FIG. 2A**FIG. 2B**



FIG. 2C**FIG. 2D**


FIG. 3



FIG. 4A**FIG. 4B**



FIG. 5A**FIG. 5B**



FIG. 6**FIG. 7**




FIG. 8**FIG. 9**

FIG. 10A**FIG. 10B**