



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>6</sup> : <b>A61K 38/18, 31/185</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 99/17797</b></p> <p>(43) International Publication Date: 15 April 1999 (15.04.99)</p>
<p>(21) International Application Number: PCT/US98/21199</p> <p>(22) International Filing Date: 6 October 1998 (06.10.98)</p> <p>(30) Priority Data: 08/946,092 6 October 1997 (06.10.97) US</p> <p>(71) Applicant: THE JOHNS HOPKINS UNIVERSITY SCHOOL OF MEDICINE [US/US]; 720 Rutland Avenue, Baltimore, MD 21205 (US).</p> <p>(72) Inventor: LEE, Se-Jin; 6711 Chokeberry Road, Baltimore, MD 21209 (US).</p> <p>(74) Agent: HAILE, Lisa, A.; Fish &amp; Richardson P.C., Suite 1400, 4225 Executive Square, La Jolla, CA 92037 (US).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report.</i></p>	
<p>(54) Title: USE OF GROWTH DIFFERENCIATION FACTOR-9 (GDF-9) AS A CONTRACEPTIVE</p>		
<p>(57) Abstract</p> <p>Growth differentiation factor-9 (GDF-9) is disclosed along with its polynucleotide sequence and amino acid sequence. Also disclosed is a method for inhibiting oocyte maturation using GDF-9 inhibitors or by inducing an immune response to GDF-9 polypeptide or fragments thereof.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AL</b>	Albania	<b>ES</b>	Spain	<b>LS</b>	Lesotho	<b>SI</b>	Slovenia
<b>AM</b>	Armenia	<b>FI</b>	Finland	<b>LT</b>	Lithuania	<b>SK</b>	Slovakia
<b>AT</b>	Austria	<b>FR</b>	France	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>AU</b>	Australia	<b>GA</b>	Gabon	<b>LV</b>	Latvia	<b>SZ</b>	Swaziland
<b>AZ</b>	Azerbaijan	<b>GB</b>	United Kingdom	<b>MC</b>	Monaco	<b>TD</b>	Chad
<b>BA</b>	Bosnia and Herzegovina	<b>GE</b>	Georgia	<b>MD</b>	Republic of Moldova	<b>TG</b>	Togo
<b>BB</b>	Barbados	<b>GH</b>	Ghana	<b>MG</b>	Madagascar	<b>TJ</b>	Tajikistan
<b>BE</b>	Belgium	<b>GN</b>	Guinea	<b>MK</b>	The former Yugoslav Republic of Macedonia	<b>TM</b>	Turkmenistan
<b>BF</b>	Burkina Faso	<b>GR</b>	Greece	<b>ML</b>	Mali	<b>TR</b>	Turkey
<b>BG</b>	Bulgaria	<b>HU</b>	Hungary	<b>MN</b>	Mongolia	<b>TT</b>	Trinidad and Tobago
<b>BJ</b>	Benin	<b>IE</b>	Ireland	<b>MR</b>	Mauritania	<b>UA</b>	Ukraine
<b>BR</b>	Brazil	<b>IL</b>	Israel	<b>MW</b>	Malawi	<b>UG</b>	Uganda
<b>BY</b>	Belarus	<b>IS</b>	Iceland	<b>MX</b>	Mexico	<b>US</b>	United States of America
<b>CA</b>	Canada	<b>IT</b>	Italy	<b>NE</b>	Niger	<b>UZ</b>	Uzbekistan
<b>CF</b>	Central African Republic	<b>JP</b>	Japan	<b>NL</b>	Netherlands	<b>VN</b>	Viet Nam
<b>CG</b>	Congo	<b>KE</b>	Kenya	<b>NO</b>	Norway	<b>YU</b>	Yugoslavia
<b>CH</b>	Switzerland	<b>KG</b>	Kyrgyzstan	<b>NZ</b>	New Zealand	<b>ZW</b>	Zimbabwe
<b>CI</b>	Côte d'Ivoire	<b>KP</b>	Democratic People's Republic of Korea	<b>PL</b>	Poland		
<b>CM</b>	Cameroon	<b>KR</b>	Republic of Korea	<b>PT</b>	Portugal		
<b>CN</b>	China	<b>KZ</b>	Kazakstan	<b>RO</b>	Romania		
<b>CU</b>	Cuba	<b>LC</b>	Saint Lucia	<b>RU</b>	Russian Federation		
<b>CZ</b>	Czech Republic	<b>LI</b>	Liechtenstein	<b>SD</b>	Sudan		
<b>DE</b>	Germany	<b>LK</b>	Sri Lanka	<b>SE</b>	Sweden		
<b>DK</b>	Denmark	<b>LR</b>	Liberia	<b>SG</b>	Singapore		
<b>EE</b>	Estonia						

**USE OF GROWTH DIFFERENTIATION FACTOR-9 (GDF-9)**  
**AS A CONTRACEPTIVE**

**BACKGROUND OF THE INVENTION**

5 1. ***Field of the Invention***

The invention relates generally to growth factors and specifically to a new member of the transforming growth factor beta (TGF- $\beta$ ) superfamily, which is denoted, growth differentiation factor-9 (GDF-9) and methods of use as a contraceptive.

2. ***Description of Related Art***

10 The transforming growth factor  $\beta$  (TGF- $\beta$ ) superfamily encompasses a group of structurally-related proteins which affect a wide range of differentiation processes during embryonic development. The family includes, Mullerian inhibiting substance (MIS), which is required for normal male sex development (Behringer, *et al.*, *Nature*, 345:167, 1990), *Drosophila* decapentaplegic (DPP) gene product, which is required for dorsal-  
15 ventral axis formation and morphogenesis of the imaginal disks (Padgett, *et al.*, *Nature*, 325:81-84, 1987), the *Xenopus* Vg-1 gene product, which localizes to the vegetal pole of eggs ((Weeks, *et al.*, *Cell*, 51:861-867, 1987), the activins (Mason, *et al.*, *Biochem, Biophys. Res. Commun.*, 135:957-964, 1986), which can induce the formation of mesoderm and anterior structures in *Xenopus* embryos (Thomsen, *et al.*, *Cell*, 63:485,  
20 1990), and the bone morphogenetic proteins (BMPs, osteogenin, OP-1) which can induce *de novo* cartilage and bone formation (Sampath, *et al.*, *J. Biol. Chem.*, 265:13198, 1990). The TGF- $\beta$ s can influence a variety of differentiation processes, including adipogenesis, myogenesis, chondrogenesis, hematopoiesis, and epithelial cell differentiation (for review, see Massague, *Cell* 49:437, 1987).

- 2 -

The proteins of the TGF- $\beta$  family are initially synthesized as a large precursor protein which subsequently undergoes proteolytic cleavage at a cluster of basic residues approximately 110-140 amino acids from the C-terminus. The C-terminal regions of the proteins are all structurally related and the different family members can be classified  
5 into distinct subgroups based on the extent of their homology. Although the homologies within particular subgroups range from 70% to 90% amino acid sequence identity, the homologies between subgroups are significantly lower, generally ranging from only 20% to 50%. In each case, the active species appears to be a disulfide-linked dimer of C-terminal fragments. For most of the family members that have been studied, the  
10 homodimeric species has been found to be biologically active, but for other family members, like the inhibins (Ling, *et al.*, *Nature*, 321:779, 1986) and the TGF- $\beta$ s (Cheifetz, *et al.*, *Cell*, 48:409, 1987), heterodimers have also been detected, and these appear to have different biological properties than the respective homodimers.

The inhibins and activins were originally purified from follicular fluid and shown to have  
15 counteracting effects on the release of follicle-stimulating hormone by the pituitary gland. Although the mRNAs for all three inhibin/activin subunits ( $\alpha$ ,  $\beta$ A and  $\beta$ B) have been detected in the ovary, none of these appear to be ovary-specific (Meunier, *et al.*, *Proc.Natl.Acad.Sci. USA*, 85:247, 1988). MIS has also been shown to be expressed by granulosa cells and the effects of MIS on ovarian development have been documented  
20 both *in vivo* in transgenic mice expressing MIS ectopically (Behringer, *supra*) and *in vitro* in organ culture (Vigier, *et al.*, *Development*, 100:43, 1987).

Identification of new factors that are tissue-specific in their expression pattern will provide a greater understanding of that tissue's development and function.

- 3 -

**SUMMARY OF THE INVENTION**

The present invention provides a cell growth and differentiation factor, GDF-9, a polynucleotide sequence which encodes the factor and antibodies which are immunoreactive with the factor. This factor appears to relate to various cell proliferative disorders, especially those involving ovarian tumors, such as granulosa cell tumors. GDF-9 is exclusively expressed in oocytes and plays a role in maturation of oocytes.

Thus, in one embodiment, the invention provides a method for detecting a cell proliferative disorder of ovarian origin and which is associated with GDF-9. In another embodiment, the invention provides a method of treating a cell proliferative disorder associated with abnormal levels of expression of GDF-9, by suppressing or enhancing GDF-9 activity.

In another embodiment, the invention provides a method for inhibiting oocyte maturation. The method includes contacting oocytes with an inhibiting effective amount of an inhibitor of GDF-9, thereby inhibiting oocyte maturation as compared to untreated oocytes. Inhibition of oocyte maturation provides contraception for fertile female subjects.

The invention also provides a method for inducing an immune response in a subject to GDF-9 whereby oocyte maturation is inhibited, comprising administering to a subject GDF-9 polypeptide or immunogenic fragment thereof and inducing an anti-GDF-9 immune response in the subject.

- 4 -

**BRIEF DESCRIPTION OF THE DRAWINGS**

FIGURE 1 shows expression of GDF-9 mRNA in adult tissues.

FIGURES 2a-2c shows nucleotide and predicted amino acid sequence of GDF-9. Consensus N-glycosylation signals are denoted by plain boxes. The putative tetrabasic  
5 processing sites are denoted by stippled boxes. The in-frame termination codons upstream of the putative initiating ATG and the consensus polyadenylation signals are underlined. The poly A tails are not shown. Numbers indicate nucleotide position relative to the 5' end.

FIGURES 3a-3b show the alignment of the C-terminal sequences of GDF-9 with other  
10 members of the TGF- $\beta$  family. The conserved cysteine residues are shaded. Dashes denote gaps introduced in order to maximize alignment.

FIGURES 4a-4b show amino acid homologies among the different members of the TGF- $\beta$  superfamily. Numbers represent percent amino acid identities between each pair calculated from the first conserved cysteine to the C-terminus. Boxes represent homolo-  
15 gies among highly-related members within particular subgroups.

FIGURES 5a-5c show the immunohistochemical localization of GDF-9 protein. Adjacent sections of an adult ovary were either stained with hematoxylin and eosin (FIGURE 5a) or incubated with immune (FIGURE 5b) or pre-immune (FIGURE 5c) serum at a dilution of 1:500. Anti-GDF-9 antiserum was prepared by expressing the C-  
20 terminal portion of murine GDF-9 (residues 308-441) in bacteria, excising GDF-9 protein from preparative SDS gels, and immunizing rabbits. Sites of antibody binding were visualized using the Vectastain ABC kit (Vector Labs).

- 5 -

FIGURE 6 shows a comparison of the predicted amino acid sequences of murine (top lines) and human (bottom lines) GDF-9. Numbers represent amino acid positions relative to the N-termini. Vertical lines represent sequence identities. Dots represent gaps introduced in order to maximize the alignment. The clear box shows the predicted proteolytic processing sites. The shaded boxes show the cysteine residues in the mature region of the proteins. The bars at the bottom show a schematic of the pre-(clear) and mature (shaded) regions of GDF-9 with the percent sequence identities between the murine and human sequences shown below.

FIGURES 7a-7d show *in situ* hybridization to adult ovary sections using a GDF-9 RNA probe. [<sup>35</sup>S]-labeled anti-sense (FIGURE 7a and 7c) or sense (FIGURE 7b and 7d) GDF-9 RNA probes were hybridized to adjacent paraffin-embedded sections of ovaries fixed in 4% paraformaldehyde. Sections were dipped in photographic emulsion, exposed, developed, and then stained with hematoxylin and eosin. Two representative fields are shown.

FIGURES 8a-8b show *in situ* hybridization to a postnatal day 4 ovary section using an antisense GDF-9 RNA probe. Sections were prepared as described for FIGURE 7. Following autoradiography and staining, the section was photographed under bright-field (FIGURE 8a) or dark-field (FIGURE 8b) illumination.

FIGURES 9a-9b show *in situ* hybridization to postnatal day 8 ovary sections using an antisense (FIGURE 9a) or sense (FIGURE 9b) GDF-9 RNA probe. Sections were prepared as described for FIGURE 7.

FIGURES 10a-10b show *in situ* hybridization to adult oviduct sections using an antisense (FIGURE 10a) or sense (FIGURE 10b) GDF-9 RNA probe. Sections were prepared as described for FIGURE 7.

- 6 -

FIGURES 11a-11b show *in situ* hybridization to an adult oviduct (0.5 days following fertilization) section using an antisense GDF-9 RNA probe. Sections were prepared as described for FIGURE 7. Following autoradiography and staining, the section was photographed under bright-field (FIGURE 11a) or dark-field (FIGURE 11b) illumination.

### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a growth and differentiation factor, GDF-9 and a polynucleotide sequence encoding GDF-9. Unlike other members of the TGF- $\beta$  superfamily, GDF-9 expression is highly tissue specific, being expressed in cells primarily in ovarian tissue. In one embodiment, the invention provides a method for detection of a cell proliferative disorder of the ovary, which is associated with GDF-9 expression. In another embodiment, the invention provides a method for treating a cell proliferative disorder associated with abnormal expression of GDF-9 by using an agent which suppresses or enhances GDF-9 activity. In another embodiment, the invention provides a method for inhibiting oocyte maturation.

Due to the expression of GDF-9 in the reproductive tract, there are a variety of applications using the polypeptide, polynucleotide and antibodies of the invention, related to contraception, fertility and pregnancy. GDF-9 could play a role in regulation of the menstrual cycle and, therefore, could be useful in various contraceptive regimens.

The TGF- $\beta$  superfamily consists of multifunctional polypeptides that control proliferation, differentiation, and other functions in many cell types. Many of the peptides have regulatory, both positive and negative, effects on other peptide growth factors. The structural homology between the GDF-9 protein of this invention and the members of the TGF- $\beta$  family, indicates that GDF-9 is a new member of the family of growth and differentiation factors. Based on the known activities of many of the other

- 7 -

members, it can be expected that GDF-9 will also possess biological activities that will make it useful as a diagnostic and therapeutic reagent.

For example, another regulatory protein that has been found to have structural homology with TGF- $\beta$  is inhibin, a specific and potent polypeptide inhibitor of the pituitary secretion of FSH. Inhibin has been isolated from ovarian follicular fluid. Because of its suppression of FSH, inhibin has potential to be used as a contraceptive in both males and females. GDF-9 may possess similar biological activity since it is also an ovarian specific peptide. Inhibin has also been shown to be useful as a marker for certain ovarian tumors (Lappohn, *et al.*, *N. Engl. J. Med.*, 321:790, 1989). GDF-9 may also be useful as a marker for identifying primary and metastatic neoplasms of ovarian origin. Similarly, GDF-9 may be useful as an indicator of developmental anomalies in prenatal screening procedures.

Another peptide of the TGF- $\beta$  family is MIS, produced by the testis and responsible for the regression of the Mullerian ducts in the male embryo. MIS has been shown to inhibit the growth of human ovarian cancer in nude mice (Donahoe, *et al.*, *Ann. Surg.*, 194:472, 1981). GDF-9 may function similarly and may, therefore, be useful as an anti-cancer agent, such as for the treatment of ovarian cancer.

GDF-9 may also function as a growth stimulatory factor and, therefore, be useful for the survival of various cell populations *in vitro*. In particular, if GDF-9 plays a role in oocyte maturation, it may be useful in *in vitro* fertilization procedures, e.g., in enhancing the success rate. Many of the members of the TGF- $\beta$  family are also important mediators of tissue repair. TGF- $\beta$  has been shown to have marked effects on the formation of collagen and causes a striking angiogenic response in the newborn mouse (Roberts, *et al.*, *Proc. Natl. Acad. Sci. USA*, 83:4167, 1986). GDF-9 may also have similar activities and may be useful in repair of tissue injury caused by trauma or burns for example.

- 8 -

The term "substantially pure" as used herein refers to GDF-9 which is substantially free of other proteins, lipids, carbohydrates or other materials with which it is naturally associated. One skilled in the art can purify GDF-9 using standard techniques for protein purification. The substantially pure polypeptide will yield a single major band on a non-reducing polyacrylamide gel. The purity of the GDF-9 polypeptide can also be  
5 determined by amino-terminal amino acid sequence analysis. GDF-9 polypeptide includes functional fragments of the polypeptide, as long as the activity of GDF-9 remains. Smaller peptides containing the biological activity of GDF-9 are included in the invention.

10 The invention provides polynucleotides encoding the GDF-9 protein. These polynucleotides include DNA, cDNA and RNA sequences which encode GDF-9. It is understood that all polynucleotides encoding all or a portion of GDF-9 are also included herein, as long as they encode a polypeptide with GDF-9 activity. Such polynucleotides include naturally occurring, synthetic, and intentionally manipulated polynucleotides.  
15 For example, GDF-9 polynucleotide may be subjected to site-directed mutagenesis. The polynucleotide sequence for GDF-9 also includes antisense sequences. The polynucleotides of the invention include sequences that are degenerate as a result of the genetic code. There are 20 natural amino acids, most of which are specified by more than one codon. Therefore, all degenerate nucleotide sequences are included in the  
20 invention as long as the amino acid sequence of GDF-9 polypeptide encoded by the nucleotide sequence is functionally unchanged.

Specifically disclosed herein is a cDNA sequence for GDF-9 which is 1712 base pairs in length and contains an open reading frame beginning with a methionine codon at nucleotide 29. The encoded polypeptide is 441 amino acids in length with a molecular  
25 weight of about 49.6 kD, as determined by nucleotide sequence analysis. The GDF-9 sequence contains a core of hydrophobic amino acids near the N-terminus, suggestive of a signal sequence for secretion. GDF-9 contains four potential N-glycosylation sites at asparagine residues 163, 229, 258, and 325 and a putative tetrabasic proteolytic

- 9 -

processing site (RRRR) at amino acids 303-306. The mature C-terminal fragment of GDF-9 is predicted to be 135 amino acids in length and have an unglycosylated molecular weight of about 15.6 kD, as determined by nucleotide sequence analysis. One skilled in the art can modify, or partially or completely remove the glycosyl groups from  
5 the GDF-9 protein using standard techniques. Therefore, the functional protein or fragments thereof of the invention includes glycosylated, partially glycosylated and unglycosylated species of GDF-9.

The degree of sequence identity of GDF-9 with known TGF- $\beta$  family members ranges from a minimum of 21% with Mullerian inhibiting substance (MIS) to a maximum of  
10 34% with bone morphogenetic protein-4 (BMP-4). GDF-9 specifically disclosed herein differs from the known family members in its pattern of cysteine residues in the C-terminal region. GDF-9 lacks the fourth cysteine of the seven cysteines present in other family members; in place of cysteine at this position, the GDF-9 sequence contains a serine residue. This GDF-9 does not contain a seventh cysteine residue elsewhere in the  
15 C-terminal region.

Minor modifications of the recombinant GDF-9 primary amino acid sequence may result in proteins which have substantially equivalent activity as compared to the GDF-9 polypeptide described herein. Such modifications may be deliberate, as by site-directed mutagenesis, or may be spontaneous. All of the polypeptides produced by these  
20 modifications are included herein as long as the biological activity of GDF-9 still exists. Further, deletion of one or more amino acids can also result in a modification of the structure of the resultant molecule without significantly altering its biological activity. This can lead to the development of a smaller active molecule which would have broader utility. For example, one can remove amino or carboxy terminal amino acids which are  
25 not required for GDF-9 biological activity.

- 10 -

The nucleotide sequence encoding the GDF-9 polypeptide of the invention includes the disclosed sequence and conservative variations thereof. The term "conservative variation" as used herein denotes the replacement of an amino acid residue by another, biologically similar residue. Examples of conservative variations include the substitution  
5 of one hydrophobic residue such as isoleucine, valine, leucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acids, or glutamine for asparagine, and the like. The term "conservative variation" also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid provided that antibodies raised to the substituted  
10 polypeptide also immunoreact with the unsubstituted polypeptide.

DNA sequences of the invention can be obtained by several methods. For example, the DNA can be isolated using hybridization techniques which are well known in the art. These include, but are not limited to: 1) hybridization of genomic or cDNA libraries with probes to detect homologous nucleotide sequences and 2) antibody screening of  
15 expression libraries to detect cloned DNA fragments with shared structural features.

Preferably the GDF-9 polynucleotide of the invention is derived from a mammalian organism, and most preferably from a mouse, rat, or human. Screening procedures which rely on nucleic acid hybridization make it possible to isolate any gene sequence from any organism, provided the appropriate probe is available. Oligonucleotide probes, which  
20 correspond to a part of the sequence encoding the protein in question, can be synthesized chemically. This requires that short, oligopeptide stretches of amino acid sequence must be known. The DNA sequence encoding the protein can be deduced from the genetic code, however, the degeneracy of the code must be taken into account. It is possible to perform a mixed addition reaction when the sequence is degenerate. This includes a  
25 heterogeneous mixture of denatured double-stranded DNA. For such screening, hybridization is preferably performed on either single-stranded DNA or denatured double-stranded DNA. Hybridization is particularly useful in the detection of cDNA clones derived from sources where an extremely low amount of mRNA sequences

- 11 -

relating to the polypeptide of interest are present. In other words, by using stringent hybridization conditions directed to avoid non-specific binding, it is possible, for example, to allow the autoradiographic visualization of a specific cDNA clone by the hybridization of the target DNA to that single probe in the mixture which is its complete  
5 complement (Wallace, *et al.*, *Nucl. Acid Res.*, 9:879, 1981).

The development of specific DNA sequences encoding GDF-9 can also be obtained by:  
1) isolation of double-stranded DNA sequences from the genomic DNA; 2) chemical  
manufacture of a DNA sequence to provide the necessary codons for the polypeptide of  
interest; and 3) *in vitro* synthesis of a double-stranded DNA sequence by reverse  
10 transcription of mRNA isolated from a eukaryotic donor cell. In the latter case, a double-  
stranded DNA complement of mRNA is eventually formed which is generally referred  
to as cDNA.

Of the three above-noted methods for developing specific DNA sequences for use in  
recombinant procedures, the isolation of genomic DNA isolates is the least common.  
15 This is especially true when it is desirable to obtain the microbial expression of  
mammalian polypeptides due to the presence of introns.

The synthesis of DNA sequences is frequently the method of choice when the entire  
sequence of amino acid residues of the desired polypeptide product is known. When the  
entire sequence of amino acid residues of the desired polypeptide is not known, the direct  
20 synthesis of DNA sequences is not possible and the method of choice is the synthesis of  
cDNA sequences. Among the standard procedures for isolating cDNA sequences of  
interest is the formation of plasmid- or phage-carrying cDNA libraries which are derived  
from reverse transcription of mRNA which is abundant in donor cells that have a high  
level of genetic expression. When used in combination with polymerase chain reaction  
25 technology, even rare expression products can be cloned. In those cases where  
significant portions of the amino acid sequence of the polypeptide are known, the  
production of labeled single or double-stranded DNA or RNA probe sequences

- 12 -

duplicating a sequence putatively present in the target cDNA may be employed in DNA/DNA hybridization procedures which are carried out on cloned copies of the cDNA which have been denatured into a single-stranded form (Jay, *et al.*, *Nucl. Acid Res.*, 11:2325, 1983).

- 5 A cDNA expression library, such as lambda gt11, can be screened indirectly for GDF-9 peptides having at least one epitope, using antibodies specific for GDF-9. Such antibodies can be either polyclonally or monoclonally derived and used to detect expression product indicative of the presence of GDF-9 cDNA.

DNA sequences encoding GDF-9 can be expressed *in vitro* by DNA transfer into a  
10 suitable host cell. "Host cells" are cells in which a vector can be propagated and its DNA expressed. The term also includes any progeny of the subject host cell. It is understood that all progeny may not be identical to the parental cell since there may be mutations that occur during replication. However, such progeny are included when the term "host cell" is used. Methods of stable transfer, meaning that the foreign DNA is continuously  
15 maintained in the host, are known in the art.

In the present invention, the GDF-9 polynucleotide sequences may be inserted into a recombinant expression vector. The term "recombinant expression vector" refers to a plasmid, virus or other vehicle known in the art that has been manipulated by insertion or incorporation of the GDF-9 genetic sequences. Such expression vectors contain a  
20 promoter sequence which facilitates the efficient transcription of the inserted genetic sequence of the host. The expression vector typically contains an origin of replication, a promoter, as well as specific genes which allow phenotypic selection of the transformed cells. Vectors suitable for use in the present invention include, but are not limited to the T7-based expression vector for expression in bacteria (Rosenberg, *et al.*,  
25 *Gene* 56:125, 1987), the pMSXND expression vector for expression in mammalian cells (Lee and Nathans, *J. Biol. Chem.*, 263:3521, 1988) and baculovirus-derived vectors for expression in insect cells. The DNA segment can be present in the vector operably

- 13 -

linked to regulatory elements, for example, a promoter (e.g., T7, metallothionein I, or polyhedrin promoters).

Polynucleotide sequences encoding GDF-9 can be expressed in either prokaryotes or eukaryotes. Hosts can include microbial, yeast, insect and mammalian organisms.

5 Methods of expressing DNA sequences having eukaryotic or viral sequences in prokaryotes are well known in the art. Biologically functional viral and plasmid DNA vectors capable of expression and replication in a host are known in the art. Such vectors are used to incorporate DNA sequences of the invention.

Transformation of a host cell with recombinant DNA may be carried out by conventional  
10 techniques as are well known to those skilled in the art. Where the host is prokaryotic, such as *E. coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently treated by the  $\text{CaCl}_2$  method using procedures well known in the art. Alternatively,  $\text{MgCl}_2$  or  $\text{RbCl}$  can be used. Transformation can also be performed after forming a protoplast of the host cell  
15 if desired.

When the host is a eukaryote, such methods of transfection of DNA as calcium phosphate co-precipitates, conventional mechanical procedures such as microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used. Eukaryotic cells can also be cotransformed with DNA sequences encoding the  
20 GDF-9 of the invention, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral vector, such as simian virus 40 (SV40) or bovine papilloma virus, to transiently infect or transform eukaryotic cells and express the protein. (see for example, *Eukaryotic Viral Vectors*, Cold Spring Harbor Laboratory, Gluzman ed., 1982).

- 14 -

Isolation and purification of microbial expressed polypeptide, or fragments thereof, provided by the invention, may be carried out by conventional means including preparative chromatography and immunological separations involving monoclonal or polyclonal antibodies.

- 5 The invention includes antibodies immunoreactive with GDF-9 polypeptide or functional fragments thereof. Antibody which consists essentially of pooled monoclonal antibodies with different epitopic specificities, as well as distinct monoclonal antibody preparations are provided. Monoclonal antibodies are made from antigen containing fragments of the protein by methods well known to those skilled in the art (Kohler, *et al.*, *Nature*,  
10 256:495, 1975). The term antibody as used in this invention is meant to include intact molecules as well as fragments thereof, such as Fab and F(ab')<sub>2</sub>, which are capable of binding an epitopic determinant on GDF-9.

The preparation of polyclonal antibodies is well-known to those skilled in the art. See, for example, Green *et al.*, *Production of Polyclonal Antisera*, in IMMUNOCHEMICAL  
15 PROTOCOLS (Manson, ed.), pages 1-5 (Humana Press 1992); Coligan *et al.*, *Production of Polyclonal Antisera in Rabbits, Rats, Mice and Hamsters*, in CURRENT PROTOCOLS IN IMMUNOLOGY, section 2.4.1 (1992), which are hereby incorporated by reference.

The preparation of monoclonal antibodies likewise is conventional. See, for example,  
20 Kohler & Milstein, *Nature* 256:495 (1975); Coligan *et al.*, sections 2.5.1-2.6.7; and Harlow *et al.*, ANTIBODIES: A LABORATORY MANUAL, page 726 (Cold Spring Harbor Pub. 1988), which are hereby incorporated by reference. Briefly, monoclonal antibodies can be obtained by injecting mice with a composition comprising an antigen, verifying the presence of antibody production by removing a serum sample, removing  
25 the spleen to obtain B lymphocytes, fusing the B lymphocytes with myeloma cells to produce hybridomas, cloning the hybridomas, selecting positive clones that produce antibodies to the antigen, and isolating the antibodies from the hybridoma cultures.

- 15 -

Monoclonal antibodies can be isolated and purified from hybridoma cultures by a variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A Sepharose, size-exclusion chromatography, and ion-exchange chromatography. See, e.g., Coligan *et al.*, sections 2.7.1-2.7.12 and sections 5 2.9.1-2.9.3; Barnes *et al.*, *Purification of Immunoglobulin G (IgG)*, in METHODS IN MOLECULAR BIOLOGY, VOL. 10, pages 79-104 (Humana Press 1992).

Methods of *in vitro* and *in vivo* multiplication of monoclonal antibodies is well-known to those skilled in the art. Multiplication *in vitro* may be carried out in suitable culture media such as Dulbecco's Modified Eagle Medium or RPMI 1640 medium, optionally 10 replenished by a mammalian serum such as fetal calf serum or trace elements and growth-sustaining supplements such as normal mouse peritoneal exudate cells, spleen cells, bone marrow macrophages. Production *in vitro* provides relatively pure antibody preparations and allows scale-up to yield large amounts of the desired antibodies. Large scale hybridoma cultivation can be carried out by homogenous suspension culture in an 15 airlift reactor, in a continuous stirrer reactor, or in immobilized or entrapped cell culture. Multiplication *in vivo* may be carried out by injecting cell clones into mammals histocompatible with the parent cells, e.g., osyngeneic mice, to cause growth of antibody-producing tumors. Optionally, the animals are primed with a hydrocarbon, especially oils such as pristane (tetramethylpentadecane) prior to injection. After one to three 20 weeks, the desired monoclonal antibody is recovered from the body fluid of the animal.

Therapeutic applications for antibodies disclosed herein are also part of the present invention. For example, antibodies of the present invention may also be derived from subhuman primate antibody. General techniques for raising therapeutically useful antibodies in baboons can be found, for example, in Goldenberg *et al.*, International 25 Patent Publication WO 91/11465 (1991) and Losman *et al.*, *Int. J. Cancer* 46:310 (1990), which are hereby incorporated by reference.

- 16 -

A therapeutically useful anti-GDF-9 antibody may be derived from a "humanized" monoclonal antibody. Humanized monoclonal antibodies are produced by transferring mouse complementarity determining regions from heavy and light variable chains of the mouse immunoglobulin into a human variable domain, and then substituting human residues in the framework regions of the murine counterparts. The use of antibody components derived from humanized monoclonal antibodies obviates potential problems associated with the immunogenicity of murine constant regions. General techniques for cloning murine immunoglobulin variable domains are described, for example, by Orlandi *et al.*, *Proc. Nat'l Acad. Sci. USA* 86:3833 (1989), which is hereby incorporated in its entirety by reference. Techniques for producing humanized monoclonal antibodies are described, for example, by Jones *et al.*, *Nature* 321: 522 (1986); Riechmann *et al.*, *Nature* 332: 323 (1988); Verhoeven *et al.*, *Science* 239: 1534 (1988); Carter *et al.*, *Proc. Nat'l Acad. Sci. USA* 89: 4285 (1992); Sandhu, *Crit. Rev. Biotech.* 12: 437 (1992); and Singer *et al.*, *J. Immunol.* 150: 2844 (1993), which are hereby incorporated by reference.

Antibodies of the invention also may be derived from human antibody fragments isolated from a combinatorial immunoglobulin library. See, for example, Barbas *et al.*, *METHODS: A COMPANION TO METHODS IN ENZYMOLOGY*, VOL. 2, page 119 (1991); Winter *et al.*, *Ann. Rev. Immunol.* 12: 433 (1994), which are hereby incorporated by reference. Cloning and expression vectors that are useful for producing a human immunoglobulin phage library can be obtained, for example, from STRATAGENE Cloning Systems (La Jolla, CA).

In addition, antibodies of the present invention may be derived from a human monoclonal antibody. Such antibodies are obtained from transgenic mice that have been "engineered" to produce specific human antibodies in response to antigenic challenge.

In this technique, elements of the human heavy and light chain loci are introduced into strains of mice derived from embryonic stem cell lines that contain targeted disruptions of the endogenous heavy and light chain loci. The transgenic mice can synthesize human antibodies specific for human antigens, and the mice can be used to produce human

- 17 -

antibody-secreting hybridomas. Methods for obtaining human antibodies from transgenic mice are described by Green *et al.*, *Nature Genet.* 7:13 (1994); Lonberg *et al.*, *Nature* 368:856 (1994); and Taylor *et al.*, *Int. Immunol.* 6:579 (1994), which are hereby incorporated by reference.

- 5 Antibody fragments of the present invention can be prepared by proteolytic hydrolysis of the antibody or by expression in *E. coli* of DNA encoding the fragment. Antibody fragments can be obtained by pepsin or papain digestion of whole antibodies by conventional methods. For example, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted F(ab')<sub>2</sub>. This  
10 fragment can be further cleaved using a thiol reducing agent, and optionally a blocking group for the sulfhydryl groups resulting from cleavage of disulfide linkages, to produce 3.5S Fab' monovalent fragments. Alternatively, an enzymatic cleavage using pepsin produces two monovalent Fab' fragments and an Fc fragment directly. These methods are described, for example, by Goldenberg, U.S. patents No. 4,036,945 and No.  
15 4,331,647, and references contained therein. These patents are hereby incorporated in their entireties by reference. *See also* Nisonhoff *et al.*, *Arch. Biochem. Biophys.* 89:230 (1960); Porter, *Biochem. J.* 73:119 (1959); Edelman *et al.*, *METHODS IN ENZYMOLOGY*, VOL. 1, page 422 (Academic Press 1967); and Coligan *et al.* at sections 2.8.1-2.8.10 and 2.10.1-2.10.4.
- 20 Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical, or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

For example, Fv fragments comprise an association of V<sub>H</sub> and V<sub>L</sub> chains. This  
25 association may be noncovalent, as described in Inbar *et al.*, *Proc. Nat'l Acad. Sci. USA* 69:2659 (1972). Alternatively, the variable chains can be linked by an intermolecular disulfide bond or cross-linked by chemicals such as glutaraldehyde. *See, e.g.*, Sandhu,

- 18 -

*supra*. Preferably, the Fv fragments comprise V<sub>H</sub> and V<sub>L</sub> chains connected by a peptide linker. These single-chain antigen binding proteins (sFv) are prepared by constructing a structural gene comprising DNA sequences encoding the V<sub>H</sub> and V<sub>L</sub> domains connected by an oligonucleotide. The structural gene is inserted into an expression vector, which  
5 is subsequently introduced into a host cell such as *E. coli*. The recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing sFvs are described, for example, by Whitlow *et al.*, METHODS: A COMPANION TO METHODS IN ENZYMOLOGY, VOL. 2, page 97 (1991); Bird *et al.*, *Science* 242:423-426 (1988); Ladner *et al.*, U.S. patent No. 4,946,778; Pack *et al.*,  
10 *Bio/Technology* 11: 1271-77 (1993); and Sandhu, *supra*.

Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, for example, by using the polymerase chain reaction to synthesize the variable  
15 region from RNA of antibody-producing cells. See, for example, Larrick *et al.*, METHODS: A COMPANION TO METHODS IN ENZYMOLOGY, VOL. 2, page 106 (1991).

The term "cell-proliferative disorder" denotes malignant as well as non-malignant cell populations which often appear to differ from the surrounding tissue both  
20 morphologically and genotypically. The GDF-9 polynucleotide that is an antisense molecule is useful in treating malignancies of the various organ systems, particularly, for example, the ovaries. Essentially, any disorder which is etiologically linked to altered expression of GDF-9 could be considered susceptible to treatment with a GDF-9 suppressing reagent.

25 The invention provides a method for detecting a cell proliferative disorder of the ovary which comprises contacting an anti-GDF-9 antibody with a cell suspected of having a GDF-9 associated disorder and detecting binding to the antibody. The antibody reactive

- 19 -

with GDF-9 is labeled with a compound which allows detection of binding to GDF-9. For purposes of the invention, an antibody specific for GDF-9 polypeptide may be used to detect the level of GDF-9 in biological fluids and tissues. Any specimen containing a detectable amount of antigen can be used. A preferred sample of this invention is tissue  
5 of ovarian origin, specifically tissue containing granulosa cells or ovarian follicular fluid. The level of GDF-9 in the suspect cell can be compared with the level in a normal cell to determine whether the subject has a GDF-9-associated cell proliferative disorder. Preferably the subject is human.

In another embodiment, the invention provides a contraceptive method which includes  
10 inhibiting oocyte maturation. The method comprises contacting oocytes with an inhibiting effective amount of an inhibitor of GDF-9, thereby inhibiting oocyte maturation as compared to untreated oocytes. The term "inhibiting effective amount" as used herein means that amount of an inhibitor of GDF-9 that reduces maturation of an oocyte by about 50%, more preferably by about 75%, and most preferably by about 95%.  
15 One of skill in the art could use any of several methods for determining whether the amount of inhibitor of GDF-9 is effective. For example, maturation of an oocyte results in ovulation, therefore, prevention of ovulation is indicative of an effective amount. For a period just prior to ovulation, a female's body temperature increases, indicating a fertile period. An absence of a change in body temperature is indicative of prevention of oocyte  
20 maturation. Alternatively, one could measure the level of serum progesterone (see Example 6) in the subject prior to and following treatment with the inhibitor of GDF-9, where a low level of progesterone maintained throughout the period is indicative of inhibition of maturation. Levels of luteinizing hormone (LH) generally increase at the time of ovulation as well, and continuous low levels of LH can serve as an indicator of  
25 inhibition of maturation of oocytes as well. Other events associated with ovulation can be monitored in a similar manner and will be known to those of skill in the art.

An "inhibitor of GDF-9" as used herein, refers to GDF-antibodies or fragments thereof as described above, antisense molecules, or peptide or peptidomimetics of GDF-9

- 20 -

polypeptide, for example. Other small molecules or chemical compounds may also be GDF-9 inhibitors. Also included are compounds identified using combinatorial chemistry methods for identifying chemical compounds that bind to GDF-9 or GDF-9 receptor. Preferably, when the inhibitor is a GDF-9 antibody or fragment thereof, the  
5 antibody is administered in a dose range of 0.1 ug to 100 mg. Preferably, the inhibitor is administered *in vivo* and preferably, the subject is a mammal, such as a human. The term "contacting" refers to administering the GDF-9 inhibitor in such a way that the inhibitor inhibits GDF-9. Methods of administration are described in detail below.

The method of the invention is also useful for inducing an immune response in a subject  
10 to GDF-9 whereby oocyte maturation is inhibited, comprising administering to a subject GDF-9 polypeptide or immunogenic fragment thereof and inducing an anti-GDF-9 immune response in the subject. Methods of producing immunogenic fragments of a polypeptide are well known in the art.

Polypeptides or peptides, including GDF-9 variants useful for the present invention  
15 comprise analogs, homologs, muteins and mimetics of GDF-9 that retain the ability to induce a specific immune response to GDF-9 or to inhibit oocyte maturation. Peptides of GDF-9 refer to portions of the amino acid sequence of GDF-9 that also retain this ability. The variants can be generated directly from GDF-9 itself by chemical modification, by proteolytic enzyme digestion, or by combinations thereof. Additionally,  
20 genetic engineering techniques, as well as methods of synthesizing polypeptides directly from amino acid residues, can be employed.

Peptides of the invention can be synthesized by such commonly used methods as t-BOC or Fmoc protection of alpha-amino groups. Both methods involve stepwise syntheses whereby a single amino acid is added at each step starting from the C terminus of the  
25 peptide (See, Coligan, *et al.*, *Current Protocols in Immunology*, Wiley Interscience, 1991, Unit 9). Peptides of the invention can also be synthesized by the well known solid phase peptide synthesis methods described Merrifield, *J. Am. Chem. Soc.*, 85:2149,

- 21 -

1962), and Stewart and Young, *Solid Phase Peptides Synthesis*, (Freeman, San Francisco, 1969, pp.27-62), using a copoly(styrene-divinylbenzene) containing 0.1-1.0 mMol amines/g polymer. On completion of chemical synthesis, the peptides can be deprotected and cleaved from the polymer by treatment with liquid HF-10% anisole for about 1/4-1  
5 hours at 0°C. After evaporation of the reagents, the peptides are extracted from the polymer with 1% acetic acid solution which is then lyophilized to yield the crude material. This can normally be purified by such techniques as gel filtration on Sephadex G-15 using 5% acetic acid as a solvent. Lyophilization of appropriate fractions of the column will yield the homogeneous peptide or peptide derivatives, which can then be  
10 characterized by such standard techniques as amino acid analysis, thin layer chromatography, high performance liquid chromatography, ultraviolet absorption spectroscopy, molar rotation, solubility, and quantitated by the solid phase Edman degradation. Alternatively, peptides can be produced by recombinant methods.

The term "substantially purified" as used herein refers to a molecule, such as a peptide  
15 that is substantially free of other proteins, lipids, carbohydrates, nucleic acids, and other biological materials with which it is naturally associated. For example, a substantially pure molecule, such as a polypeptide, can be at least 60%, by dry weight, the molecule of interest. One skilled in the art can purify GDF-9 peptides using standard protein purification methods and the purity of the polypeptides can be determined using standard  
20 methods including, *e.g.*, polyacrylamide gel electrophoresis (*e.g.*, SDS-PAGE), column chromatography (*e.g.*, high performance liquid chromatography (HPLC)), and amino-terminal amino acid sequence analysis.

Non-peptide compounds that mimic the binding and function of GDF-9 ("mimetics") can be produced by the approach outlined in Saragovi *et al.*, *Science* 253: 792-95 (1991).  
25 Mimetics are molecules which mimic elements of protein secondary structure. See, for example, Johnson *et al.*, "Peptide Turn Mimetics," in BIOTECHNOLOGY AND PHARMACY, Pezzuto *et al.*, Eds., (Chapman and Hall, New York 1993). The underlying rationale behind the use of peptide mimetics is that the peptide backbone of

- 22 -

proteins exists chiefly to orient amino acid side chains in such a way as to facilitate molecular interactions. For the purposes of the present invention, appropriate mimetics can be considered to be the equivalent of GDF-9 itself.

Longer peptides can be produced by the "native chemical" ligation technique which links  
5 together peptides (Dawson, *et al.*, *Science*, 266:776, 1994). Variants can be created by recombinant techniques employing genomic or cDNA cloning methods. Site-specific and region-directed mutagenesis techniques can be employed. See CURRENT PROTOCOLS IN MOLECULAR BIOLOGY vol. 1, ch. 8 (Ausubel *et al.* eds., J. Wiley & Sons 1989 & Supp. 1990-93); PROTEIN ENGINEERING (Oxender & Fox eds., A.  
10 Liss, Inc. 1987). In addition, linker-scanning and PCR-mediated techniques can be employed for mutagenesis. See PCR TECHNOLOGY (Erlich ed., Stockton Press 1989); CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, vols. 1 & 2, *supra*. Protein sequencing, structure and modeling approaches for use with any of the above techniques are disclosed in PROTEIN ENGINEERING, *loc. cit.*, and CURRENT PROTOCOLS IN  
15 MOLECULAR BIOLOGY, vols. 1 & 2, *supra*.

The invention also includes various pharmaceutical compositions that prevent maturation of an oocyte. The pharmaceutical compositions according to the invention are prepared by bringing an antibody against GDF-9, a peptide or peptide derivative of GDF-9, a GDF-9 mimetic, or a GDF-9-binding agent according to the present invention into a form  
20 suitable for administration to a subject using carriers, excipients and additives or auxiliaries. Frequently used carriers or auxiliaries include magnesium carbonate, titanium dioxide, lactose, mannitol and other sugars, talc, milk protein, gelatin, starch, vitamins, cellulose and its derivatives, animal and vegetable oils, polyethylene glycols and solvents, such as sterile water, alcohols, glycerol and polyhydric alcohols.  
25 Intravenous vehicles include fluid and nutrient replenishers. Preservatives include antimicrobial, anti-oxidants, chelating agents and inert gases. Other pharmaceutically acceptable carriers include aqueous solutions, non-toxic excipients, including salts, preservatives, buffers and the like, as described, for instance, in *Remington's*

- 23 -

*Pharmaceutical Sciences*, 15th ed. Easton: Mack Publishing Co., 1405-1412, 1461-1487 (1975) and *The National Formulary XIV.*, 14th ed. Washington: American Pharmaceutical Association (1975), the contents of which are hereby incorporated by reference. The pH and exact concentration of the various components of the  
5 pharmaceutical composition are adjusted according to routine skills in the art. See *Goodman and Gilman's The Pharmacological Basis for Therapeutics* (7th ed.).

The methods involves administering to a subject a therapeutically or inhibiting effective dose of a pharmaceutical composition containing the compounds of the present invention and a pharmaceutically acceptable carrier. "Administering" the pharmaceutical  
10 composition of the present invention may be accomplished by any means known to the skilled artisan. By "subject" is meant any mammal, preferably a human.

The pharmaceutical compositions are preferably prepared and administered in dose units. Solid dose units are tablets, capsules and suppositories. For treatment of a patient, depending on activity of the compound, manner of administration, nature and severity  
15 of the disorder, age and body weight of the patient, different daily doses are necessary. Under certain circumstances, however, higher or lower daily doses may be appropriate. The administration of the daily dose can be carried out both by single administration in the form of an individual dose unit or else several smaller dose units and also by multiple administration of subdivided doses at specific intervals.

20 The pharmaceutical compositions according to the invention are in general administered topically, intravenously, orally or parenterally or as implants, but even rectal use is possible in principle. Suitable solid or liquid pharmaceutical preparation forms are, for example, granules, powders, tablets, coated tablets, (micro)capsules, suppositories, syrups, emulsions, suspensions, creams, aerosols, drops or injectable solution in ampule  
25\* form and also preparations with protracted release of active compounds, in whose preparation excipients and additives and/or auxiliaries such as disintegrants, binders, coating agents, swelling agents, lubricants, flavorings, sweeteners or solubilizers are

- 24 -

customarily used as described above. The pharmaceutical compositions are suitable for use in a variety of drug delivery systems. For a brief review of present methods for drug delivery, see Langer, *Science*, 249: 1527-1533 (1990), which is incorporated herein by reference.

5 The pharmaceutical compositions according to the invention may be administered locally or systemically. By "therapeutically effective dose" is meant the quantity of a compound according to the invention necessary to prevent, to cure or at least partially arrest the symptoms of the disease and its complications. An "inhibiting effective amount" of GDF-9 inhibitor has been defined above for use as in contraception regimes. Amounts  
10 effective for this use will, of course, depend on the severity of the disease and the weight and general state of the patient. Typically, dosages used *in vitro* may provide useful guidance in the amounts useful for *in situ* administration of the pharmaceutical composition, and animal models may be used to determine effective dosages for treatment of particular disorders. Various considerations are described, *e.g.*, in Gilman  
15 *et al.* (eds.) (1990) GOODMAN AND GILMAN'S: THE PHARMACOLOGICAL BASES OF THERAPEUTICS, 8th ed., Pergamon Press; and REMINGTON'S PHARMACEUTICAL SCIENCES, 17th ed. (1990), Mack Publishing Co., Easton, Pa., each of which is herein incorporated by reference.

The pharmaceutical compositions of the invention may also be delivered by a slow  
20 release vehicle. Compounds of the invention can be further diluted by addition of suspending medium or other biologically acceptable carrier to obtain injectable or implantable slow release formulations of any therapeutically effective total dosage. One approach which has been used to provide controlled release compositions for drug delivery is liposome encapsulation. Among the main types of liposomes, multivesicular  
25 liposomes (Kim, *et al.*, *Biochim. Biophys. Acta*; 728:339-348, 1983), are uniquely different from unilamellar liposomes (Huang, *Biochemistry*; 8:334-352, 1969; Kim, *et al.*, *Biochim. Biophys. Acta*; 646:1-10, 1981), multilamellar liposomes (Bangham, *et al.*, *J. Mol. Bio.*, 13:238-252, 1965), and stable plurilamellar liposomes (U.S. Patent No.

- 25 -

4,522,803). U.S. Patent No. 5,077,056 discloses studies that show the rate of release of the encapsulated biological agent from liposomes into an aqueous environment can be modulated by introducing protonophores or ionophores into liposomes to create a membrane potential. In addition, a method is known (U.S. Patent No. 5,186,941) for  
5 controlling the release rate of drugs from vesicle compositions wherein the liposomes containing a therapeutic agent encapsulated are suspended in a solution containing sufficient solute to provide an osmolarity substantially isotonic with respect to that of the solution within the vesicles, and hypertonic with respect to physiological saline. In multivesicular liposomes, it is also known (WO 96/08253) to control the rate of release  
10 of active agents by introducing an osmotic spacer into the aqueous solution in which the active agent is dissolved prior to formation of the multivesicular liposomes.

The antibodies of the invention can be used in any subject in which it is desirable to administer *in vitro* or *in vivo* immunodiagnosis or immunotherapy. The antibodies of the invention are suited for use, for example, in immunoassays in which they can be utilized  
15 in liquid phase or bound to a solid phase carrier. In addition, the antibodies in these immunoassays can be detectably labeled in various ways. Examples of types of immunoassays which can utilize antibodies of the invention are competitive and non-competitive immunoassays in either a direct or indirect format. Examples of such immunoassays are the radioimmunoassay (RIA) and the sandwich (immunometric) assay.  
20 Detection of the antigens using the antibodies of the invention can be done utilizing immunoassays which are run in either the forward, reverse, or simultaneous modes, including immunohistochemical assays on physiological samples. Those of skill in the art will know, or can readily discern, other immunoassay formats without undue experimentation.

25 The antibodies of the invention can be bound to many different carriers and used to detect the presence of an antigen comprising the polypeptide of the invention. Examples of well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, agaroses and

- 26 -

magnetite. The nature of the carrier can be either soluble or insoluble for purposes of the invention. Those skilled in the art will know of other suitable carriers for binding antibodies, or will be able to ascertain such, using routine experimentation.

There are many different labels and methods of labeling known to those of ordinary skill  
5 in the art. Examples of the types of labels which can be used in the present invention include enzymes, radioisotopes, fluorescent compounds, colloidal metals, chemiluminescent compounds, phosphorescent compounds, and bioluminescent compounds. Those of ordinary skill in the art will know of other suitable labels for binding to the antibody, or will be able to ascertain such, using routine experimentation.

10 Another technique which may also result in greater sensitivity consists of coupling the antibodies to low molecular weight haptens. These haptens can then be specifically detected by means of a second reaction. For example, it is common to use such haptens as biotin, which reacts with avidin, or dinitrophenyl, puridoxal, and fluorescein, which can react with specific antihapten antibodies.

15 In using the monoclonal antibodies of the invention for the *in vivo* detection of antigen, the detectably labeled antibody is given a dose which is diagnostically effective. The term "diagnostically effective" means that the amount of detectably labeled monoclonal antibody is administered in sufficient quantity to enable detection of the site having the antigen comprising a polypeptide of the invention for which the monoclonal antibodies  
20 are specific.

The concentration of detectably labeled monoclonal antibody which is administered should be sufficient such that the binding to those cells having the polypeptide is detectable compared to the background. Further, it is desirable that the detectably labeled monoclonal antibody be rapidly cleared from the circulatory system in order to  
25 give the best target-to-background signal ratio.

- 27 -

As a rule, the dosage of detectably labeled monoclonal antibody for *in vivo* diagnosis will vary depending on such factors as age, sex, and extent of disease of the individual. Such dosages may vary, for example, depending on whether multiple injections are given, antigenic burden, and other factors known to those of skill in the art.

- 5 For *in vivo* diagnostic imaging, the type of detection instrument available is a major factor in selecting a given radioisotope. The radioisotope chosen must have a type of decay which is detectable for a given type of instrument. Still another important factor in selecting a radioisotope for *in vivo* diagnosis is that deleterious radiation with respect to the host is minimized. Ideally, a radioisotope used for *in vivo* imaging will lack a  
10 particle emission, but produce a large number of photons in the 140-250 keV range, which may readily be detected by conventional gamma cameras.

For *in vivo* diagnosis radioisotopes may be bound to immunoglobulin either directly or indirectly by using an intermediate functional group. Intermediate functional groups which often are used to bind radioisotopes which exist as metallic ions to  
15 immunoglobulins are the bifunctional chelating agents such as diethylenetriaminepentaacetic acid (DTPA) and ethylenediaminetetraacetic acid (EDTA) and similar molecules. Typical examples of metallic ions which can be bound to the monoclonal antibodies of the invention are  $^{111}\text{In}$ ,  $^{97}\text{Ru}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{89}\text{Zr}$ , and  $^{201}\text{Tl}$ .

The monoclonal antibodies of the invention can also be labeled with a paramagnetic  
20 isotope for purposes of *in vivo* diagnosis, as in magnetic resonance imaging (MRI) or electron spin resonance (ESR). In general, any conventional method for visualizing diagnostic imaging can be utilized. Usually gamma and positron emitting radioisotopes are used for camera imaging and paramagnetic isotopes for MRI. Elements which are particularly useful in such techniques include  $^{157}\text{Gd}$ ,  $^{55}\text{Mn}$ ,  $^{162}\text{Dy}$ ,  $^{52}\text{Cr}$ , and  $^{56}\text{Fe}$ .

- 28 -

The monoclonal antibodies of the invention can be used *in vitro* and *in vivo* to monitor the course of amelioration of a GDF-9-associated disease in a subject. Thus, for example, by measuring the increase or decrease in the number of cells expressing antigen comprising a polypeptide of the invention or changes in the concentration of such antigen present in various body fluids, it would be possible to determine whether a particular therapeutic regimen aimed at ameliorating the GDF-9-associated disease is effective. The term "ameliorate" denotes a lessening of the detrimental effect of the GDF-9-associated disease in the subject receiving therapy. Similarly, such techniques can be used to follow a contraceptive regime in a subject.

- 10 The present invention identifies a nucleotide sequence that can be expressed in an altered manner as compared to expression in a normal cell, therefore, it is possible to design appropriate therapeutic or diagnostic techniques directed to this sequence. Thus, where a cell-proliferative disorder is associated with the expression of GDF-9, nucleic acid sequences that interfere with GDF-9 expression at the translational level can be used.
- 15 This approach utilizes, for example, antisense nucleic acid and ribozymes to block translation of a specific GDF-9 mRNA, either by masking that mRNA with an antisense nucleic acid or by cleaving it with a ribozyme. In addition, antisense molecules or ribozymes are useful in the contraceptive methods of the invention where it is desirable to inhibit GDF-9 expression. As a further mechanism of action, in "transcription arrest"
- 20 it appears that some polynucleotides can form "triplex," or triple-helical structures with double stranded genomic DNA containing the gene of interest, thus interfering with transcription by RNA polymerase. Giovannangeli *et al.*, *Proc. Natl. Acad. Sci.* **90**:10013 (1993); Ebbinghaus *et al. J. Clin. Invest.* **92**:2433 (1993).

Antisense nucleic acids are DNA or RNA molecules that are complementary to at least a portion of a specific mRNA molecule (Weintraub, *Scientific American*, 262:40, 1990). In the cell, the antisense nucleic acids hybridize to the corresponding mRNA, forming a double-stranded molecule. The antisense nucleic acids interfere with the translation of the mRNA, since the cell will not translate a mRNA that is double-stranded.

- 29 -

Antisense oligomers of about 15 nucleotides are preferred, since they are easily synthesized and are less likely to cause problems than larger molecules when introduced into the target GDF-9-producing cell. The use of antisense methods to inhibit the *in vitro* translation of genes is well known in the art (Marcus-Sakura, *Anal. Biochem.*, 172:289, 5 1988).

Ribozymes are RNA molecules possessing the ability to specifically cleave other single-stranded RNA in a manner analogous to DNA restriction endonucleases. Through the modification of nucleotide sequences which encode these RNAs, it is possible to engineer molecules that recognize specific nucleotide sequences in an RNA molecule and 10 cleave it (Cech, *J. Amer. Med. Assn.*, 260:3030, 1988). A major advantage of this approach is that, because they are sequence-specific, only mRNAs with particular sequences are inactivated.

There are two basic types of ribozymes namely, *tetrahymena*-type (Hasselhoff, *Nature*, 334:585, 1988) and "hammerhead"-type. *Tetrahymena*-type ribozymes recognize 15 sequences which are four bases in length, while "hammerhead"-type ribozymes recognize base sequences 11-18 bases in length. The longer the recognition sequence, the greater the likelihood that the sequence will occur exclusively in the target mRNA species. Consequently, hammerhead-type ribozymes are preferable to *tetrahymena*-type ribozymes for inactivating a specific mRNA species and 18-based recognition sequences 20 are preferable to shorter recognition sequences.

The present invention also provides gene therapy for the treatment of cell proliferative disorders which are mediated by GDF-9 protein or where it is desirable to inhibit GDF-9 expression for purposes of contraception. Such therapy would achieve its therapeutic effect by introduction of the GDF-9 antisense polynucleotide into cells having the 25 proliferative disorder or introduction of antisense into oocytes. Delivery of antisense or sense GDF-9 polynucleotide, GDF-9 polypeptide or peptides can be achieved using a recombinant expression vector such as a chimeric virus or a colloidal dispersion system.

- 30 -

Especially preferred for therapeutic delivery of antisense sequences is the use of targeted liposomes.

Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, herpes virus, vaccinia, or, preferably, an RNA virus such as a retrovirus.

5 Preferably, the retroviral vector is a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these  
10 vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a GDF-9 sequence of interest into the viral vector, along with another gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target specific. Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a sugar, a glycolipid,  
15 or a protein. Preferred targeting is accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector containing the GDF-9 antisense polynucleotide.

20 Since recombinant retroviruses are defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA  
25 transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include, but are not limited to  $\Psi$ 2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are

- 31 -

replaced by other genes of interest, the vector can be packaged and vector virion produced.

Alternatively, NIH 3T3 or other tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes *gag*, *pol* and *env*, by conventional  
5 calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral vector into the culture medium.

Another targeted delivery system for GDF-9 polynucleotides or polypeptides is a colloidal dispersion system. Colloidal dispersion systems include macromolecule  
10 complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles *in vitro* and *in vivo*. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0  $\mu\text{m}$  can encapsulate a  
15 substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, *et al.*, *Trends Biochem. Sci.*, 6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an  
20 efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic  
25 information (Mannino, *et al.*, *Biotechniques*, 6:682, 1988).

- 32 -

The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of  
5 divalent cations.

Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebrosides, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms,  
10 particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine.

The targeting of liposomes can be classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished  
15 based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries. Active targeting, on the other hand, involves alteration of the liposome by coupling the liposome to a specific ligand such as a  
20 monoclonal antibody, sugar, glycolipid, or protein, or by changing the composition or size of the liposome in order to achieve targeting to organs and cell types other than the naturally occurring sites of localization.

The surface of the targeted delivery system may be modified in a variety of ways. In the case of a liposomal targeted delivery system, lipid groups can be incorporated into the  
25 lipid bilayer of the liposome in order to maintain the targeting ligand in stable association with the liposomal bilayer. Various linking groups can be used for joining the lipid chains to the targeting ligand.



- 34 -

replicate filters were hybridized to probes representing known members of the family, and DNA was prepared from non-hybridizing colonies for sequence analysis.

The primer combination of SJL160 and SJL153, yielded three known sequences (inhibin  $\beta$ B, BMP-2, and BMP-4) and one novel sequence (designated GDF-9) among 145  
5 subclones analyzed.

RNA isolation and Northern analysis were carried out as described previously (Lee, S.J., *Mol. Endocrinol.* 4:1034, 1990). An oligo dT-primed cDNA library was prepared from 2.5-3  $\mu$ g of ovary poly A-selected RNA in the lambda ZAP II vector according to the instructions provided by Stratagene. The ovary library was not amplified prior to  
10 screening. Filters were hybridized as described previously (Lee, S.-J., *Proc. Natl. Acad. Sci. USA*, 88:4250-4254, 1991). DNA sequencing of both strands was carried out using the dideoxy chain termination method (Sanger, *et al.*, *Proc. Natl. Acad. Sci., USA*, 74:5463-5467, 1977) and a combination of the S1 nuclease/exonuclease III strategy (Henikoff, S., *Gene*, 28:351-359, 1984) and synthetic oligonucleotide primers.

15

## EXAMPLE 2

### EXPRESSION PATTERN AND SEQUENCE OF GDF-9

To determine the expression pattern of GDF-9, RNA samples prepared from a variety of adult tissues were screened by Northern analysis. Five micrograms of twice polyA-selected RNA prepared from each tissue were electrophoresed on formaldehyde gels,  
20 blotted and probed with GDF-9. As shown in Figure 1, the GDF-9 probe detected a 1.7 kb mRNA expressed exclusively in the ovary.

A mouse ovary cDNA library of  $1.5 \times 10^6$  recombinant phage was constructed in lambda ZAP II and screened with a probe derived from the GDF-9 PCR product. The nucleotide sequence of the longest of nineteen hybridizing clones is shown in Figure 2. Consensus  
25 N-glycosylation signals are denoted by plain boxes. The putative tetrabasic processing

- 35 -

sites are denoted by stippled boxes. The in-frame termination codons upstream of the putative initiating ATG and the consensus polyadenylation signals are underlined. The poly A tails are not shown. Numbers indicate nucleotide position relative to the 5' end. The 1712 bp sequence contains a long open reading frame beginning with a methionine codon at nucleotide 29 and potentially encoding a protein 441 amino acids in length with a molecular weight of 49.6 kD. Like other TGF- $\beta$  family members, the GDF-9 sequence contains a core of hydrophobic amino acids near the N-terminus suggestive of a signal sequence for secretion. GDF-9 contains four potential N-glycosylation sites at asparagine residues 163, 229, 258, and 325 and a putative tetrabasic proteolytic processing site (RRRR) at amino acids 303-306. The mature C-terminal fragment of GDF-9 is predicted to be 135 amino acids in length and have an unglycosylated molecular weight of 15.6 kD. Although the C-terminal portion of GDF-9 clearly shows homology with the other family members, the sequence of GDF-9 is significantly diverged from those of the other family members (Figures 3 and 4). Figure 3 shows the alignment of the C-terminal sequences of GDF-9 with the corresponding regions of human GDF-1 (Lee, *Proc. Natl. Acad. Sci. USA*, 88:4250-4254, 1991), *Xenopus* Vg-1 (Weeks, *et al.*, *Cell*, 51:861-867, 1987), human Vgr-1 (Celeste, *et al.*, *Proc. Natl. Acad. Sci. USA*, 87:9843-9847, 1990), human OP-1 (Ozkaynak, *et al.*, *EMBO J.*, 9:2085-2093, 1990), human BMP-5 (Celeste, *et al.*, *Proc. Natl. Acad. Sci. USA*, 87:9843-9847, 1990), *Drosophila* 60A (Wharton, *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:9214-9218, 1991), human BMP-2 and 4 (Wozney, *et al.*, *Science*, 242:1528-1534, 1988), *Drosophila* DPP (Padgett, *et al.*, *Nature*, 325:81-84, 1987), human BMP-3 (Wozney, *et al.*, *Science*, 242:1528-1534, 1988), human MIS (Cate, *et al.*, *Cell*, 45:685-698, 1986), human inhibin ,  $\beta$ A, and  $\beta$ B (Mason, *et al.*, *Biochem. Biophys. Res. Commun.*, 135:957-964, 1986), human TGF- $\beta$ 1 (Derynck, *et al.*, *Nature*, 316:701-705, 1985), human TGF- $\beta$ 2 (deMartin, *et al.*, *EMBO J.*, 6:3673-3677, 1987), human TGF- $\beta$ 3 (ten Dijke, *et al.*, *Proc. Natl. Acad. Sci. USA*, 85:4715-4719, 1988), chicken TGF- $\beta$ 4 (Jakowlew, *et al.*, *Mol. Endocrinol.*, 2:1186-1195, 1988), and *Xenopus* TGF- $\beta$ 5 (Kondaiah, *et al.*, *J. Biol. Chem.*, 265:1089-1093, 1990). The conserved cysteine residues are shaded. Dashes denote gaps introduced in order to maximize the alignment.

- 36 -

Figure 4 shows the amino acid homologies among the different members of the TGF- $\beta$  superfamily. Numbers represent percent amino acid identities between each pair calculated from the first conserved cysteine to the C-terminus. Boxes represent homologies among highly-related members within particular subgroups.

5 The degree of sequence identify with known family members ranges from a minimum of 21% with MIS to a maximum of 34% with BMP-4. Hence, GDF-9 is comparable to MIS in its degree of sequence divergence from the other members of this superfamily. Moreover, GDF-9 shows no significant sequence homology to other family members in the pro-region of the molecule. GDF-9 also differs from the known family members in  
10 its pattern of cysteine residues in the C-terminal region. GDF-9 lacks the fourth cysteine of the seven cysteines that are present in all other family members; in place of cysteine at this position, the GDF-9 sequence contains a serine residue. In addition, GDF-9 does not contain a seventh cysteine residue elsewhere in the C-terminal region.

### EXAMPLE 3

#### 15 IMMUNOCHEMICAL LOCALIZATION OF GDF-9 IN THE ZONA PELLUCIDA

To determine whether GDF-9 mRNA was translated, sections of adult ovaries were incubated with antibodies directed against recombinant GDF-9 protein. In order to raise antibodies against GDF-9, portions of GDF-9 cDNA spanning amino acids 30 to 295  
20 (pro-region) or 308 to 441 (mature region) were cloned into the T7-based pET3 expression vector (provided by F.W. Studier, Brookhaven National Laboratory), and the resulting plasmids were transformed into the BL21 (DE3) bacterial strain. Total cell extracts from isopropyl  $\beta$ -D-thiogalactoside-induced cells were electrophoresed on SDS/polyacrylamide gels, and the GDF-9 protein fragments were excised, mixed with  
25 Freund's adjuvant, and used to immunize rabbits by standard methods known to those of skill in the art. All immunizations were carried out by Spring Valley Lab (Sykesville, MD). The presence of GDF-9 reactive antibodies in the sera of these rabbits were assessed by Western analysis of bacterially-expressed protein by Western analysis.

- 37 -

For immunohistochemical studies, ovaries were removed from adult mice, fixed in 4% paraformaldehyde, embedded in paraffin, and sectioned. Sites of antibody binding were detected by using the Vectastain ABC kit, according to the instructions provided by Vector Laboratories. FIGURE 5 shows the immunohistochemical localization of GDF-9 protein. Adjacent sections of an adult ovary were either stained with hematoxylin and eosin (FIGURE 5a) or incubated with immune (FIGURE 5b) or pre-immune (FIGURE 5c) serum at a dilution of 1:500. As show in FIGURE 5b, the antiserum detected protein solely in oocytes. No staining was detected using pre-immune serum (FIGURE 5c). Hence, GDF-9 protein appears to translated *in vivo* by oocytes.

10

**EXAMPLE 4****ISOLATION OF HUMAN GDF-9**

In order to isolate a cDNA clone encoding human GDF-9, a cDNA library was constructed in lambda ZAP II usign poly A-selected RNA prepared from an adult human ovary. From this library, a cDNA clone containing the entire human GDF-9 coding sequence was identified using standard screening techniques as in Example 1 and using the murine GDF-9 clone as a probe. A comparison of the predicted amino acid sequences of murine (top lines) and human (bottom lines) GDF-9 is shown in FIGURE 6. Numbers represent amino acid positions relative to the N-termini. Vertical lines represent sequence identities. Dots represent gaps introduced in order to maximize the alignment. The clear box shows the predicted proteolytic processing sites. The shaded boxes show the cysteine residues in the mature region of the proteins. The bars at the bottom show a schematic of the pre-(clear) and mature (shaded) regions of GDF-9 with the percent sequence identities between the murine and human sequences shown below.

Like murine GDF-9, human GDF-9 contains a hydrophobic leader sequence, a putative RXXR proteolytic cleavage site, and a C-terminal region containing the hallmarks of other TGF- $\beta$  family members. Murine and human GDF-9 are 64% identical in the pro-region and 90% identical in the predicted mature region of the molecule. The high degree

25

- 38 -

of homology between the two sequences suggest that human GDF-9 plays an important role during embryonic development and/or in the adult ovary.

### **EXAMPLE 5**

#### **NUCLEIC ACID DETECTION OF EXPRESSION OF GDF-9 IN OOCYTES**

5 In order to localize the expression of GDF-9 in the ovary, *in situ* hybridization to mouse ovary sections was carried out using an antisense GDF-9 RNA probe. FIGURE 7 shows *in situ* hybridization to adult ovary sections using a GDF-9 RNA probe. [<sup>35</sup>S]-labeled anti-sense (FIGURE 7a and 7c) or sense (FIGURE 7b and 7d) GDF-9 RNA probes were hybridized to adjacent paraffin-embedded sections of ovarians fixed in 4%  
10 paraformaldehyde. Sections were dipped in photographic emulsion, exposed, developed, and then stained with hematoxylin and eosin. Two representative fields are shown.

As shown in FIGURES 7a and 7c, GDF-9 mRNA was detected primarily in oocytes in adult ovaries. Every oocyte (regardless of the stage of follicular development) examined showed GDF-9 expression, and no expression was detected in any other cell types. No  
15 hybridization was seen using a control GDF-9 sense RNA probe (FIGURE 7b and 7d). Hence, GDF-9 expression appears to be oocyte-specific in adult ovaries.

To determine the pattern of expression of GDF-9 mRNA during ovarian development, sections of neonatal ovaries were probed with a GDF-9 RNA probe. FIGURE 8 shows *in situ* hybridization to a postnatal day 4 ovary section using an antisense GDF-9 RNA  
20 probe. Sections were prepared as described for FIGURE 7. Following autoradiography and staining, the section was photographed under bright-field (FIGURE 8a) or dark-field (FIGURE 8b) illumination.

FIGURE 9 shows *in situ* hybridization to postnatal day 8 ovary sections using an antisense (FIGURE 9a) or sense (FIGURE 9b) GDF-9 RNA probe. Sections were  
25 prepared as described for FIGURE 7.

- 39 -

GDF-9 mRNA expression was first detected at the onset of follicular development. This was most clearly evident at postnatal day 4, where only oocytes that were present in follicles showed GDF-9 expression (FIGURE 8): no expression was seen in oocytes that were not surrounded by granulosa cells. By postnatal day 8, every oocyte appeared to  
5 have undergone follicular development, and every oocyte showed GDF-9 expression (FIGURE 9).

To determine whether GDF-9 was also expressed following ovulation, sections of mouse oviducts were examined by *in situ* hybridization. FIGURE 10 shows *in situ* hybridization to adult oviduct sections using an antisense (FIGURE 10a) or sense  
10 (FIGURE 10b) GDF-9 RNA probe. Sections were prepared as described for FIGURE 7.

FIGURE 11 shows *in situ* hybridization to adult oviduct (0.5 days following fertilization) section using an antisense GDF-9 RNA probe. Sections were prepared as described for FIGURE 7. Following autoradiography and staining, the section was photographed  
15 under bright-field (FIGURE 11a) or dark-field (FIGURE 11b) illumination.

As shown in FIGURE 10, GDF-9 was expressed by oocytes that had been released into the oviduct. However, the expression of GDF-9 mRNA turned off rapidly following fertilization of the oocytes; by day 0.5 following fertilization, only some embryos (such as the one shown in FIGURE 11) expressed GDF-9 mRNA, and by day 1.5, all embryos  
20 were negative for GDF-9 expression.

- 40 -

**EXAMPLE 6**  
**CONTRACEPTIVE TARGET**

GDF-9 is a novel protein growth factor which regulates the early development and the maturation of the egg (oocyte). Experimental results demonstrate that this factor is essential to the development of the egg. Several lines of evidence support this including:  
5 1) GDF-9 mRNA is expressed exclusively by developing and maturing oocytes; 2) GDF-9 protein is secreted by maturing oocytes and is essential for the normal development of the follicle; and 3) GDF-9 deficient (knockout) female mice do not develop mature eggs or follicles and are completely sterile (Dong *et al.*, *Nature*, 383,531, 1996).

10 GDF-9 is a highly specific and novel target for the development of a female contraceptive or as a nonsurgical method of sterilization. There are several different approaches one may use to develop a contraceptive using GDF-9 as a target: 1) the use of the GDF-9 protein or a peptide fragment as a “vaccine”; 2) use of an anti-GDF-9 antibody which binds to and effectively inhibit or neutralize the biological actions of the  
15 GDF-9 molecule; and 3) the use of a GDF-9 receptor antagonist. The term “vaccine” as used herein refers to a composition or compound that induces a protective immune response. For example, a GDF-9 “vaccine” protects against conception or pregnancy and acts as a contraceptive.

The following examples include different ways that oocyte development and  
20 maturation may be regulated in order to prevent ovulation or to cause permanent, nonsurgical sterilization in humans and in other species. These examples are intended to illustrate but not limit the invention. While they are typical of those that might be used, other procedures known to those skilled in the art may alternatively be used.

- 41 -

1. Active Immunization: GDF-9 "vaccines"

The intact GDF-9 protein, or a fragment thereof, is an effective source of the composition used to induce an immune response directed toward oocytes (*e.g.*, vaccine). The intact GDF-9 protein or a monomeric subunit can be economically produced in mammalian cell  
5 culture, yeast or microbial culture (such as *Bacillus* or *E. coli*). Alternatively, a peptide fragment of GDF-9 is also a suitable source for producing a vaccine. Typically, such fragments are 10 to 30 amino acids long, are exposed on the exterior portions of the GDF-9 molecule, and may be located within or near the receptor binding portion of the GDF-9 molecule.

10 Several methods can be used to enhance the antigenicity of the GDF-9, or a fragment thereof. Such methods are well known and include forming a chemical complex between a hapten (GDF-9 or a fragment thereof) and a larger carrier molecule (hapten-carrier); mixing the antigen with an adjuvant; or denaturing the protein with heat, detergent, or chaotrophic agent (urea or guanidine-HCl).

15 The following experiment can be performed to demonstrate that a GDF-9 composition is an effective contraceptive. Using New Zealand white rabbits as a model, the rabbits are immunized with either the intact GDF-9, a GDF-9 monomer, or one of the GDF-9 fragments listed in the table below. The route of administration is subcutaneous when an adjuvant is used. In the case when an adjuvant is not used, the routes of  
20 administration may include subcutaneous, intramuscular, intravenous or intraperitoneal. The preferred route is subcutaneous.

Typically, single or multiple immunization may be required to produce an immune response to GDF-9. The humoral immune response may be monitored by routine serologic methods. Briefly, one to two weeks following the first immunization, a serum  
25 sample is obtained and the antibody titer to the GDF-9 protein is determined by either

- 42 -

ELISA or RIA methods. A second or third immunization may be required every 10-14 days in order to produce an effective immune response.

Table 1

GDF-9 vaccines, proteins, concentrations and formulations

	<b>Vaccine</b>	<b>Protein concentration (mg)</b>	<b>Formulation</b>
5	Intact GDF-9 homodimer	0.1, 1.0, 10	neutral buffered saline solution with or without adjuvant <sup>1</sup>
	Heat denatured or chemically denatured GDF-9	0.1, 1.0, 10	neutral buffered saline solution with or without adjuvant
	GDF-9 monomer	0.1, 1.0, 10	neutral buffered saline solution with or without adjuvant
10	GDF-9 monomer conjugated to carrier protein <sup>2</sup>	0.1, 1.0, 10	neutral buffered saline solution with or without adjuvant
	GDF-9 fragments 1, 2 or 3 (predicted exposed site within putative receptor binding domain)	0.1, 1.0, 10	neutral buffered saline solution with adjuvant
15	GDF-9 fragments 1, 2 or 3 conjugated to a carrier protein	0.1, 1.0, 10	neutral buffered saline solution with or without adjuvant

<sup>1</sup>Adjuvants suitable for use in animals include Freund's complete or incomplete adjuvant. Adjuvants suitable for use in humans will be known to those of skill in the art.

<sup>2</sup>Monomer is covalently coupled to a suitable carrier protein such as heat denatured immunoglobulin, denatured albumin or KLH.

- 44 -

Rabbits are a good model to demonstrate the effectiveness of a GDF-9 vaccine. Rabbits are immunized with GDF-9 or a GDF-9 fragment in an adjuvant. Antibody titers to GDF-9 are determined in order to assure that a humoral response has been achieved. Gonadotrophin is administered in order to induce ovulation in the rabbit. Following  
5 gonadotrophin administration serum progesterone levels are measured. Low serum progesterone levels are an indication that ovulation is inhibited by hyperimmunizing the rabbits with GDF-9 vaccines.

Histologic evaluation of the ovaries from GDF-9 vaccinated rabbits will show that the ovaries are intact. A mild mixed focal inflammatory infiltrate within the medulla of each  
10 ovary is observed. The ovaries will be devoid of mature oocytes and follicles. The cortex will appear normal. Histologic examination of the fallopian tubes and uterus will reveal no abnormalities or inflammation.

## 2. Passive Immunization

Passive immunization is a second approach to developing a GDF-9 specific  
15 contraceptive. Using this approach, antibodies are produced which have high affinity of GDF-9. The antibodies are purified by conventional affinity chromatography methods (*e.g.*, protein-A). In this instance, either a immunoglobulin enriched serum fraction (gamma globulin) or a monoclonal antibody generated against GDF-9 is suitable. In these instances, the anti-GDF-9 gamma globulin or the monoclonal antibody will have  
20 the property of specifically binding to GDF-9. The antibody may either block the binding of the GDF-9 to its natural receptor or function to increase the rate of clearance of the GDF-9 from the ovary. In either case, the objective is to allow the antibody to contact the GDF-9 produced by the oocytes and prevent it from contacting its natural receptor.

- 45 -

For use in humans, the preferred antibody is a humanized monoclonal antibody. Production of murine monoclonal antibodies is well established and familiar to those skilled in the art. Briefly, mice are immunized with GDF-9 homodimer, monomer or a fragment (Table 1). Once a humoral response is produced, the spleen or lymph node  
5 cells are removed and fused with a mouse myeloma cell, typically SP2/0 or P-2 cells. The fused cells (hybridoma) are cultured in a selection media, cloned and subcloned to generate an immortal cell line producing an antibody to GDF-9.

Due to its immunogenicity, the murine monoclonal antibody is not most desirable to use as a therapeutic for human use. The antigenicity of a murine antibody can be reduced by  
10 replacing the murine immunoglobulin heavy and light chain constant domains with the equivalent human antibody domains (chimeric antibody). The chimeric antibody can be further "humanized" by replacing intervening structural murine sequences within the antibodies hypervariable regions. This is referred to as a "fully humanized" antibody. The fully humanized antibody is preferred since this antibody can be used repeatedly in  
15 a human subject without causing an immune response to the antibody.

### 3. GDF-9 Receptor Antagonist

A third approach to suppressing GDF-9 function is to develop a receptor antagonist. The approach is to first identify cell surface receptor(s) which are responsible for signal transduction in the ovary. These receptors are likely to be expressed on the egg or  
20 associated accessory cells (*e.g.*, granulosa or thecal). Once a target cell population is identified, a receptor binding assay is used to identify peptides or small organic compounds which will bind to the GDF-9 receptor and prevent the GDF-9 from binding to its receptor. The use of the receptor antagonist has the advantage of being a specific inhibitor of GDF-9 induction of oocyte maturation and the potential to be a long-term,  
25 reversible therapeutic.

- 46 -

Although the invention has been described with reference to the presently preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

**SUMMARY OF SEQUENCES**

Sequence ID No. 1 is the nucleotide sequence for the primer, SJL160, for GDF-9 (page 24, lines 15 and 16);

Sequence ID No. 2 is the nucleotide sequence for the primer, SJL153, for GDF-9 (page 5 24, lines 17 and 18);

Sequence ID No. 3 is the nucleotide and deduced amino acid sequence for GDF-9 (Figure 2);

Sequence ID No. 4 is the deduced amino acid sequence for GDF-9 (Figure 2);

Sequence ID No. 5 is the amino acid sequence of the C-terminus of GDF-3 (Figure 3);

10 Sequence ID No. 6 is the amino acid sequence of the C-terminus of GDF-9 (Figure 3);

Sequence ID No. 7 is the amino acid sequence of the C-terminus of GDF-1 (Figure 3);

Sequence ID No. 8 is the amino acid sequence of the C-terminus of Vg-1 (Figure 3);

Sequence ID No. 9 is the amino acid sequence of the C-terminus of Vgr-1 (Figure 3);

Sequence ID No. 10 is the amino acid sequence of the C-terminus of OP-1 (Figure 3);

15 Sequence ID No. 11 is the amino acid sequence of the C-terminus of BMP-5 (Figure 3);

Sequence ID No. 12 is the amino acid sequence of the C-terminus of 60A (Figure 3);

Sequence ID No. 13 is the amino acid sequence of the C-terminus of BMP-2 (Figure 3);

- 48 -

Sequence ID No. 14 is the amino acid sequence of the C-terminus of BMP-4 (Figure 3);

Sequence ID No. 15 is the amino acid sequence of the C-terminus of DPP (Figure 3);

Sequence ID No. 16 is the amino acid sequence of the C-terminus of BMP-3 (Figure 3);

Sequence ID No. 17 is the amino acid sequence of the C-terminus of MIS (Figure 3);

5 Sequence ID No. 18 is the amino acid sequence of the C-terminus of inhibin (Figure 3);

Sequence ID No. 19 is the amino acid sequence of the C-terminus of  $\beta$ A (Figure 3);

Sequence ID No. 20 is the amino acid sequence of the C-terminus of  $\beta$ B (Figure 3);

Sequence ID No. 21 is the amino acid sequence of the C-terminus of TGF- $\beta$ 1 (Figure 3);

Sequence ID No. 22 is the amino acid sequence of the C-terminus of TGF- $\beta$ 2 (Figure 3);

10 Sequence ID No. 23 is the amino acid sequence of the C-terminus of TGF- $\beta$ 3 (Figure 3);

Sequence ID No. 24 is the amino acid sequence of the C-terminus of TGF- $\beta$ 4 (Figure 3);

Sequence ID No. 25 is the amino acid sequence of the C-terminus of TGF- $\beta$ 5 (Figure 3);

and

Sequence ID No. 26 is the amino acid sequence of the C-terminus of GDF-9 (Figure 6).

**CLAIMS**

1. A method for inhibiting oocyte maturation comprising contacting oocytes with an inhibiting effective amount of an inhibitor of GDF-9, thereby inhibiting oocyte maturation as compared to untreated oocytes.
2. The method of claim 1, wherein the inhibitor is a GDF-9 antibody or fragment thereof.
3. The method of claim 2, wherein the antibody is a monoclonal antibody.
4. The method of claim 3, wherein the antibody is a humanized antibody.
5. The method of claim 2, wherein the antibody is administered to a subject *in vivo*.
6. The method of claim 2, wherein the antibody is administered in a dose range of about 0.1 ug to 100 mg.
7. The method of claim 1, wherein the inhibitor is a GDF-9 antisense molecule.
8. A method for inducing an immune response in a subject to GDF-9 whereby oocyte maturation is inhibited, comprising administering to a subject GDF-9 polypeptide or immunogenic fragment thereof and inducing an anti-GDF-9 immune response in the subject.
9. The method of claim 8, wherein the GDF-9 polypeptide or fragment thereof is administered with an adjuvant.
10. The method of claim 8, wherein the GDF-9 polypeptide or fragment thereof is administered by injection or by implantation in a slow release delivery vehicle.

11. The method of claim 8, wherein the GDF-9 polypeptide or fragment thereof is administered in a dose range of about 0.1 ug to 100 mg.




FIG. 1



601 AAGAGCACCGTACTCATTCACCCCTGAAGAACAACACAGATGGATTGAGATTGATGTGACCTC 660  
 R A P Y S F T L K K H R W I E I D V T S  
 661 CCTCCTTCAGCCCCCTAGTGACCTCCAGCGAGAGGAGCATTTCACCTGTCTGTCAATTTTAC 720  
 L L Q P L V T S S E R S I H L S V N F T  
 721 ATGCACAAGAAGACCAGGTGCCAGAGGACGGAGTGTTTAGCATGCCCTCTCTCAGTGCCTCC 780  
 C T K D Q V P E D G V F S M P L S V P P  
 781 TTCCCTCATCTGTATCTCAACGACACAAGCACCCAGGCCCTACCACCTCTTGGCAGTCTCT 840  
 S L I L Y L N D T S T Q A Y H S W Q S L  
 841 TCAGTCCACCTGGAGGCCCTTTACAGCATCCCCGGCCAGGCCGGTGTGGCTGCCCGTCCCCGT 900  
 Q S T W R P L Q H P G Q A G V A A R P V 3/15  
 901 GAAAGAGGAGTACTGAGGTGGAAGATCTCCCGGCCGTCGAGGGCAGAAAGCCAT 960  
 K E E A T E V E R S P R R R R G Q K A I  
 961 CCGCTCCGAAGCGAAGGGCCACTTCTTACAGCATCCCTTCAACCTCAGCGAATACTTCAA 1020  
 R S E A K G P L L T A S F N L S E Y F K  
 1021 ACAGTTTCTTTCCCCCAAAACGAGTGTGAATCCACTCCACTTCAGACTGAGTTTGTAGTCA 1080  
 Q F L F P Q N E C E L H D F R L S F S Q  
 1081 GCTCAAATGGGACAACCTGGATCGTGGCCCCCGCACAGGTACAACCCCTAGGTACTGTAAAGG 1140  
 L K W D N W I V A P H R Y N P R Y C K G

FIG.2b

1141 GGACTGTCCCTAGGGCGGTCAGGCATCGGTATGGCTCCTCTGTCACACCATGGTCCAGAA 1200  
       D C P R A V R H R Y G S . P V H T M V Q N  
 1201 TATAATCTATGAGAAGCTGGACCCCTTCAGTGCCAAGGCCCTTCGTGTGTCGCCGGCAAGTA 1260  
       I I Y E K L D P S V P R P S C V P G K Y  
 1261 CAGCCCCCTGAGTGTGTGACCCATTGAACCCGACGGCTCCATCGCTTACAAAGAGTACGA 1320  
       S P L S V L T I E P D G S I A Y K E Y E  
 1321 AGACATGATAGCTACGAGGTGCCACCTGTCTCGTTAGCATTGGGGCCCACTTCAACAAGCCCTGC 1380  
       D M I A T R C T C R \*  
 1381 CTGGCAGAGCAATGCTGTGGGCCCTTAGAGTGCCTGGCAGAGAGCTTCCCTGTGACCCAGTC 1440  
 1441 TCTCCGTCCTGCTCAGTGCACACTGTGTGAGCCGGGGAAGTGTGTGTGTGATGAGCA 1500  
 1501 CATCGAGTGCAGTGTCCGTAGGTGTAAGGGCACACTCAGTGGTCCATAAACCAA 1560  
 1561 GTGAAATGTAACCTCATTTGGAGAGCTCTTTCTCCCCACGAGTGTAGTTTTTCAGTGGACAG 1620  
 1621 ATTTGTTAGCATAAGTCTCGAGTAGAATGTAGCTGTGAACATGTCAGAGTGTCTGTGGTTT 1680  
 1681 TATGTGACCGGAAGAATAAACTGTTGATGGCAT 1712

4/15

FIG.2c

GDF-3 KRRAAISVPKGFCC--RNFCHRHQLFINF--QDLGWHKWVIAPKGFMANVCHGECFFSMTTYLNS--  
GDF-9 FNLSEYFKQFLFP--QNECELHDFRLSF--SQLKWDNWIVAPHRYNPRYCKGDCPRAVRHRYS--  
GDF-1 PRDAEPVLGGP--GGAORARRLYVSF--REVGWHRWVIAPRGFLANYOQCCALPVALSGSGGP  
Vg-1 RRKRSYKLPFTA--SNICKRRHLYVEF--KDVGWQNWVIAPQGYMANYCYGECYPLTEILNG--  
Vgr-1 RVSSASDYNSEL--KTACKRKHELIVSF--QDLGWQDWIIAPKGYAANYDGECSFPLNAHMNA--  
OP-1 RMANVAENSSDQ--RQACKKHELIVSF--RDLGWQDWIIAPEGYAAYYCEGCAFPPLNSYMNA--  
BMP-5 RMSSVGDYNTSEQ--KQACKKHELIVSF--RDLGWQDWIIAPEGYAAFYDGECSFPLNAHMNA--  
60A SPNNVPLLEPMES--TRSQMQTLYIDF--KDLGWHDWIIAPEGYGAFYCSGECNFPPLNAHMNA--  
BMP-2 EKRQAKHKQRKRL--KSSCKRHPLYVDF--SDVGWNDWIVAPPGYHAFYCHGECFFPLADHLNS--  
BMP-4 RSPKHSQRARKK--NKNORRHSLYVDF--SDVGWNDWIVAPPGYQAFYCHGDCFPPLADHLNS--  
DPP KRHARRPTRRKNH--DDTORRHSLYVDF--SDVGWDDWIVAPLGYDAYYCHGKCFPLADHFNS--  
BMP-3 QTLKARRKQWIE--PRNCARRYLKVDF--ADIGWSEWIIISPKSFDAYYCSGACQFPMPKSLKP--  
MIS PGRAQRSAGATAA--DGFCALRELSVDL---RAERSVLIPETYQANNQGVCGWPQSDRNPRY--  
Inhibin  $\alpha$  LRLLRPPEEPAA--HANCHRVALNISF--QELGWERWIVYPPSFIHYCHGCGLHIPPNLSLPV  
Inhibin  $\beta$ A RRRRRGLECDGKV---NICCKKQFFVSF--KDIGWNDWIIAPSGYHANYCEGECPSHIAGTSGSSL  
Inhibin  $\beta$ B RIRKRGLECDGRT---NLCCRQFFIDF--RLIGWNDWIIAPTGYGNYCEGCPAYLAGVPGSAS  
TGF- $\beta$ 1 RRALDTNYCFSST--EKNCVQRQLYIDFRKDLGWK--WIHEPKGYHANFCLGFCPIWSLD-----  
TGF- $\beta$ 2 KRALDAAYCFRNV--QDNCLRPLYIDFKRDLGWK--WIHEPKGYANFCAGACPYLWSSD-----  
TGF- $\beta$ 3 KRALDTNYCFRNL--EENCVVRPLYIDFRQDLGWK--WVHEPKGYANFCSGRCPYLRSAD-----  
TGF- $\beta$ 4 RRDLDTDYCFGPGTDEKNCVVRPLYIDFRKDLQWK--WIHEPKGYMANFOMGFCPIWSAD-----  
TGF- $\beta$ 5 KRGVGQEYCFGNN--GPNCCVKPLYINFRKDLGWK--WIHEPKGYEANYCLGNCPYIWSMD-----

FIG.3a

GDF-3 --SNYAFMQALMHH--ADPKVPKAVCV--PTKLSPI SMLYQ-DSDKNVILRHYEDMVVDECCCG  
 GDF-9 --PVHTWQNI IYE--KLDPSVPRFSCV--PGKYSPLSVLTI-EPDGSIA YKEYEDMIATRCIOR

GDF-1 PALNHAVLRALMHA--AAPGAADLEFCV--PARLSPISVLF--DNSDNVVL RQYEDMVVDECCOR  
 Vg-1 --SNHAILQTLVHS--IEPEDIPLFCV--PTKMSPISMLFY-DNNDNVVLRHYENMAVDECCOR  
 Vgr-1 --TNHAI VQTLVHL--MNPEYVVPKFCV--PTKLN AISVLYF-DDNSNVILK KYRNMVVRACCH  
 OP-1 --TNHAI VQTLVHF--INPETVVPKFCV--PTQLNAISVLYF-DDSSNVILK KYRNMVVRACCH  
 BMP-5 --TNHAI VQTLVHL--MFPDHVVPKFCV--PTKLN AISVLYF-DDSSNVILK KYRNMVVRACCH  
 60A --TNHAI VQTLVHL--LEPKKVPKFCV--PTRLGALPVLYH-LNDENVNLK KYRNMVVRACCH  
 BMP-2 --TNHAI VQTLVNS---VNSKI PKACV--PTELSAISMLYL-DENEKVV LKNYQDMVVECCOR  
 BMP-4 --TNHAI VQTLVNS---VNSSI PKACV--PTELSAISMLYL-DEYDKVV LKNYQEMVVECCOR  
 DPP --TNHAVVQTLVNN--MNP GKVPKACV--PTQLDSVAMLYL-NDQSTVVLK NYQEMTVVCCOR  
 BMP-3 --SNHATI QSI VRA-VGVVPGIPEFCV--PEKMSSLSILFF-DENKNVVLK VY PNMVTVESCCOR  
 MIS --GNHVVLL LKMQA--RGAALARPFCV--PTAYAGKLLI SLSEER--ISAHHV PNMVATECCOR  
 Inhibin  $\alpha$  --PGAPPTPAQ PYS---LLPGAQFCV--AALPGTMRPLHVRTTSDGGY SFKYETVPNLLTQHACI  
 Inhibin  $\beta$ A --SFHSTVINHYRMRGHS PFANLKS CV--PTKLRPMSMLY-DDGQNI IKKDIQNMIVECCOS  
 Inhibin  $\beta$ B --SFHTAVVNQYRMRGLNPGT-VNSCCI--PTKLS TMSMLYF-DDEYNIVKRDV PNMIVECCCA  
 TGF- $\beta$ 1 --TQYSKVLALYNQ--HNP GASAFPCCV--PQALEPLPIVY-VGRKPKV-EQLSNMIVRSKOS  
 TGF- $\beta$ 2 --TQHSRVL SLYNT--INPEASAFPCCV--SQDLEPLTILY-IGKTPKI-EQLSNMIVRSKOS  
 TGF- $\beta$ 3 --TTHSTVGLYNT--LNPEASAFPCCV--PQDLEPLTILY-VGRTPKV-EQLSNMIVRSKOS  
 TGF- $\beta$ 4 --TQYTKVLALYNQ--HNP GASAFPCCV--PQTLDPLPIIY-VGRNVRV-EQLSNMIVRSKOS  
 TGF- $\beta$ 5 --TQYSKVL SLYNQ--NNPGASISFCV--PDVLEPLPIIY-VGRTAKV-EQLSNMIVRSKOS

FIG.3b








FIG. 5c




FIG. 5b




FIG. 5a

10/15

1 MALPSNFLGVCFFAWLCFLSSLSSQASTEESQSGASENVESEADPWSLL 50  
 || | ||| ||||| || |||| | | ||| | |||||

1 MARPNKFLWFCFFAWLCFPISLGSQASGGEAQIAASAELESGAMPWSLL 50

51 LPVDGTDRSGLLPPLFKVLSDRRGETPKLQPDSTRALYMKKLYKTYATKE 100  
 | || |||| ||||| || | ||||| |||||

51 QHIDERDRAGLLPALFKVLSVGRGGSPRLQPDSTRALHYMKKLYKTYATKE 100

101 GVPKPSRSHLYNTVRLFSPCAQQEQAPSNQVTGPLPMVDLLFNLDRTAM 150  
 | || ||||| ||| ||| ||||| || | ||||| |

101 GIPKSNRSHLYNTVRLFTPCTRHKQAPGDQVTGILPSVELLNFNLDRTTV 150

151 EHLKSVLLYTLNNSASSSTVTTCMCDLVVKEAMSSGRAPPRAPYSFTL. 199  
 ||||| ||| | | | | | | | | | | | | |

151 EHLKSVLLYNINNSVSFSSAVKVCNLMIKEPKSSRTLGRAPYSFTFN 200

200 .....KKHRWIEIDVTSLLQPLVTSSERSIHLSVNFTCTKDQV....PE 239  
 ||| || ||||| || | |||| | |||| |||

201 SQFEFGKHKHWIQIDVTSLLQPLVASNKRSIHMSINFTCMKDQLEHPSAQ 250

240 DGVFSMPLSVPPSLILYLNDTSTQAYHSWQSLQSTWRPLQHPGQA.GVAA 283  
 | | | | | ||||| ||||| || | | | | | | |

251 NGLFNMTL.VSPSLILYLNDTSAQAYHSWYSLHYKRRPSQGPDQERSLSA 299

289 RPVKEEATEVERSP..RRRRCQKAIRSEAKGPLLTASFNLSEYFKQFLFP 336  
 || ||| | || |||| | | | | ||||| ||| |

300 YPVGEEAAEDGRSSHRRRRCQETVSSELKKPLGPASFNLSEYFRQFLLP 349

337 QNECELHDFRLSFSQLKWDNWIVAPHRYNPRYCKGDPRAVRHRYGSPVH 386  
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

350 QNECELHDFRLSFSQLKWDNWIVAPHRYNPRYCKGDPRAVGHRYGSPVH 399

387 TMVQNIIEKLDPSVPRPSVPGKYSPLSVLTIEPDGSIAYKEYEDMIAT 436  
 ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

400 TMVQNIIEKLDSSVPRPSVPAKYSPLSVLTIEPDGSIAYKEYEDMIAT 449

437 RCTCR 441  
 ||||

450 KCTCR 454  
 ||||



64%

FIG. 6

90%

FIG. 7a




FIG. 7b



FIG. 7c



FIG. 7d



FIG. 8a




FIG. 8b




FIG. 9b




FIG. 9a



FIG. 10a



FIG. 10b



FIG. 11a



FIG. 11b

## --SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (I) APPLICANT: The Johns Hopkins University School of Medicine
- (ii) TITLE OF INVENTION: GROWTH DIFFERENTIATION FACTOR-9
- (iii) NUMBER OF SEQUENCES: 26
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Fish & Richardson
  - (B) STREET: 4225 Executive Square, Suite 1400
  - (C) CITY: San Diego
  - (D) STATE: California
  - (E) COUNTRY: USA
  - (F) ZIP: 92037
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 08/946,092
  - (B) FILING DATE: 06-OCT-1997
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 08/491,835
  - (B) FILING DATE: 23-OCT-1995
  - (C) CLASSIFICATION:
- (vi) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: PCT/US94/00685
  - (B) FILING DATE: 12-JAN-1994
  - (C) CLASSIFICATION:
- (vi) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 08/003,303
  - (B) FILING DATE: 12-JAN-1993
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Lisa A. Haile Ph.D., J.D.
  - (B) REGISTRATION NUMBER: 38,347
  - (C) REFERENCE/DOCKET NUMBER: 07265/138W01
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (619) 678-5070
  - (B) TELEFAX: (619) 678-5099

## (2) INFORMATION FOR SEQ ID NO:1:

- (I) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 35 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: SJL160
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 1..35
  - (D) OTHER INFORMATION: /note= "Where "B" occurs, B = inosine"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCGGAATTCG GBTGGVANVA NTGGRTBRTB KCBCC 35

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 33 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: SJL153

- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 1..33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CCGGAATTCR CADSCRCADC YNBTDGYDRY CAT 33

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1712 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: GDF-9

- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 29..1351

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATGCGTTCCT TCTTAGTTCT TCCAAGTC ATG GCA CTT CCC AGC AAC TTC CTG 52  
 Met Ala Leu Pro Ser Asn Phe Leu  
 1 5

TTG	GGG	GTT	TGC	TGC	TTT	GCC	TGG	CTG	TGT	TTT	CTT	AGT	AGC	CTT	AGC	100
Leu	Gly	Val	Cys	Cys	Phe	Ala	Trp	Leu	Cys	Phe	Leu	Ser	Ser	Leu	Ser	
	10					15					20					
TCT	CAG	GCT	TCT	ACT	GAA	GAA	TCC	CAG	AGT	GGA	GCC	AGT	GAA	AAT	GTG	148
Ser	Gln	Ala	Ser	Thr	Glu	Glu	Ser	Gln	Ser	Gly	Ala	Ser	Glu	Asn	Val	
25					30					35					40	
GAG	TCT	GAG	GCA	GAC	CCC	TGG	TCC	TTG	CTG	CTG	CCT	GTA	GAT	GGG	ACT	196
Glu	Ser	Glu	Ala	Asp	Pro	Trp	Ser	Leu	Leu	Leu	Pro	Val	Asp	Gly	Thr	
				45					50					55		
GAC	AGG	TCT	GGC	CTC	TTG	CCC	CCC	CTC	TTT	AAG	GTT	CTA	TCT	GAT	AGG	244
Asp	Arg	Ser	Gly	Leu	Leu	Pro	Pro	Leu	Phe	Lys	Val	Leu	Ser	Asp	Arg	
			60					65					70			
CGA	GGT	GAG	ACC	CCT	AAG	CTG	CAG	CCT	GAC	TCC	AGA	GCA	CTC	TAC	TAC	292
Arg	Gly	Glu	Thr	Pro	Lys	Leu	Gln	Pro	Asp	Ser	Arg	Ala	Leu	Tyr	Tyr	
		75					80					85				
ATG	AAA	AAG	CTC	TAT	AAG	ACG	TAT	GCT	ACC	AAA	GAG	GGG	GTT	CCC	AAA	340
Met	Lys	Lys	Leu	Tyr	Lys	Thr	Tyr	Ala	Thr	Lys	Glu	Gly	Val	Pro	Lys	
	90					95					100					
CCC	AGC	AGA	AGT	CAC	CTC	TAC	AAT	ACC	GTC	CGG	CTC	TTC	AGT	CCC	TGT	388
Pro	Ser	Arg	Ser	His	Leu	Tyr	Asn	Thr	Val	Arg	Leu	Phe	Ser	Pro	Cys	
105					110					115					120	
GCC	CAG	CAA	GAG	CAG	GCA	CCC	AGC	AAC	CAG	GTG	ACA	GGA	CCG	CTG	CCG	436
Ala	Gln	Gln	Glu	Gln	Ala	Pro	Ser	Asn	Gln	Val	Thr	Gly	Pro	Leu	Pro	
				125					130					135		
ATG	GTG	GAC	CTG	CTG	TTT	AAC	CTG	GAC	CGG	GTG	ACT	GCC	ATG	GAA	CAC	484
Met	Val	Asp	Leu	Leu	Phe	Asn	Leu	Asp	Arg	Val	Thr	Ala	Met	Glu	His	
			140					145					150			
TTG	CTC	AAA	TCG	GTC	TTG	CTA	TAC	ACT	CTG	AAC	AAC	TCT	GCC	TCT	TCC	532
Leu	Leu	Lys	Ser	Val	Leu	Leu	Tyr	Thr	Leu	Asn	Asn	Ser	Ala	Ser	Ser	
		155					160					165				
TCC	TCC	ACT	GTG	ACC	TGT	ATG	TGT	GAC	CTT	GTG	GTA	AAG	GAG	GCC	ATG	580
Ser	Ser	Thr	Val	Thr	Cys	Met	Cys	Asp	Leu	Val	Val	Lys	Glu	Ala	Met	
	170					175					180					
TCT	TCT	GGC	AGG	GCA	CCC	CCA	AGA	GCA	CCG	TAC	TCA	TTC	ACC	CTG	AAG	628
Ser	Ser	Gly	Arg	Ala	Pro	Pro	Arg	Ala	Pro	Tyr	Ser	Phe	Thr	Leu	Lys	
185					190					195					200	
AAA	CAC	AGA	TGG	ATT	GAG	ATT	GAT	GTG	ACC	TCC	CTC	CTT	CAG	CCC	CTA	676
Lys	His	Arg	Trp	Ile	Glu	Ile	Asp	Val	Thr	Ser	Leu	Leu	Gln	Pro	Leu	
				205					210					215		
GTG	ACC	TCC	AGC	GAG	AGG	AGC	ATT	CAC	CTG	TCT	GTC	AAT	TTT	ACA	TGC	724
Val	Thr	Ser	Ser	Glu	Arg	Ser	Ile	His	Leu	Ser	Val	Asn	Phe	Thr	Cys	
			220					225					230			
ACA	AAA	GAC	CAG	GTG	CCA	GAG	GAC	GGA	GTG	TTT	AGC	ATG	CCT	CTC	TCA	772
Thr	Lys	Asp	Gln	Val	Pro	Glu	Asp	Gly	Val	Phe	Ser	Met	Pro	Leu	Ser	
		235					240					245				

GTG CCT CCT TCC CTC ATC TTG TAT CTC AAC GAC ACA AGC ACC CAG GCC	820
Val Pro Pro Ser Leu Ile Leu Tyr Leu Asn Asp Thr Ser Thr Gln Ala	
250 255 260	
TAC CAC TCT TGG CAG TCT CTT CAG TCC ACC TGG AGG CCT TTA CAG CAT	868
Tyr His Ser Trp Gln Ser Leu Gln Ser Thr Trp Arg Pro Leu Gln His	
265 270 275 280	
CCC GGC CAG GCC GGT GTG GCT GCC CGT CCC GTG AAA GAG GAA GCT ACT	916
Pro Gly Gln Ala Gly Val Ala Ala Arg Pro Val Lys Glu Glu Ala Thr	
285 290 295	
GAG GTG GAA AGA TCT CCC CGG CGC CGT CGA GGG CAG AAA GCC ATC CGC	964
Glu Val Glu Arg Ser Pro Arg Arg Arg Arg Gly Gln Lys Ala Ile Arg	
300 305 310	
TCC GAA GCG AAG GGG CCA CTT CTT ACA GCA TCC TTC AAC CTC AGC GAA	1012
Ser Glu Ala Lys Gly Pro Leu Leu Thr Ala Ser Phe Asn Leu Ser Glu	
315 320 325	
TAC TTC AAA CAG TTT CTT TTC CCC CAA AAC GAG TGT GAA CTC CAT GAC	1060
Tyr Phe Lys Gln Phe Leu Phe Pro Gln Asn Glu Cys Glu Leu His Asp	
330 335 340	
TTC AGA CTG AGT TTT AGT CAG CTC AAA TGG GAC AAC TGG ATC GTG GCC	1108
Phe Arg Leu Ser Phe Ser Gln Leu Lys Trp Asp Asn Trp Ile Val Ala	
345 350 355 360	
CCG CAC AGG TAC AAC CCT AGG TAC TGT AAA GGG GAC TGT CCT AGG GCG	1156
Pro His Arg Tyr Asn Pro Arg Tyr Cys Lys Gly Asp Cys Pro Arg Ala	
365 370 375	
GTC AGG CAT CGG TAT GGC TCT CCT GTG CAC ACC ATG GTC CAG AAT ATA	1204
Val Arg His Arg Tyr Gly Ser Pro Val His Thr Met Val Gln Asn Ile	
380 385 390	
ATC TAT GAG AAG CTG GAC CCT TCA GTG CCA AGG CCT TCG TGT GTG CCG	1252
Ile Tyr Glu Lys Leu Asp Pro Ser Val Pro Arg Pro Ser Cys Val Pro	
395 400 405	
GGC AAG TAC AGC CCC CTG AGT GTG TTG ACC ATT GAA CCC GAC GGC TCC	1300
Gly Lys Tyr Ser Pro Leu Ser Val Leu Thr Ile Glu Pro Asp Gly Ser	
410 415 420	
ATC GCT TAC AAA GAG TAC GAA GAC ATG ATA GCT ACG AGG TGC ACC TGT	1348
Ile Ala Tyr Lys Glu Tyr Glu Asp Met Ile Ala Thr Arg Cys Thr Cys	
425 430 435 440	
CGT TAGCATGGGG GCCACTTCAA CAAGCCTGCC TGGCAGAGCA ATGCTGTGGG	1401
Arg	
CCTTAGAGTG CCTGGGCAGA GAGCTTCCTG TGACCAGTCT CTCCGTGCTG CTCAGTGCAC	1461
ACTGTGTGAG CGGGGGAAGT GTGTGTGTGT GGATGAGCAC ATCGAGTGCA GTGTCCGTAG	1521
GTGTAAAGGG CACACTCACT GGTTCGTTGCC ATAAACCAAG TGAAATGTAA CTCATTTGGA	1581
GAGCTCTTTC TCCCCACGAG TGTAGTTTTT AGTGGACAGA TTTGTTAGCA TAAGTCTCGA	1641
GTAGAATGTA GCTGTGAACA TGTCAGAGTG CTGTGGTTTT ATGTGACGGA AGAATAAACT	1701
GTTGATGGCA T	1712

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 441 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Ala Leu Pro Ser Asn Phe Leu Leu Gly Val Cys Cys Phe Ala Trp
1           5           10           15
Leu Cys Phe Leu Ser Ser Leu Ser Ser Gln Ala Ser Thr Glu Glu Ser
20           25           30
Gln Ser Gly Ala Ser Glu Asn Val Glu Ser Glu Ala Asp Pro Trp Ser
35           40           45
Leu Leu Leu Pro Val Asp Gly Thr Asp Arg Ser Gly Leu Leu Pro Pro
50           55           60
Leu Phe Lys Val Leu Ser Asp Arg Arg Gly Glu Thr Pro Lys Leu Gln
65           70           75           80
Pro Asp Ser Arg Ala Leu Tyr Tyr Met Lys Lys Leu Tyr Lys Thr Tyr
85           90           95
Ala Thr Lys Glu Gly Val Pro Lys Pro Ser Arg Ser His Leu Tyr Asn
100          105          110
Thr Val Arg Leu Phe Ser Pro Cys Ala Gln Gln Glu Gln Ala Pro Ser
115          120          125
Asn Gln Val Thr Gly Pro Leu Pro Met Val Asp Leu Leu Phe Asn Leu
130          135          140
Asp Arg Val Thr Ala Met Glu His Leu Leu Lys Ser Val Leu Leu Tyr
145          150          155          160
Thr Leu Asn Asn Ser Ala Ser Ser Ser Ser Thr Val Thr Cys Met Cys
165          170          175
Asp Leu Val Val Lys Glu Ala Met Ser Ser Gly Arg Ala Pro Pro Arg
180          185          190
Ala Pro Tyr Ser Phe Thr Leu Lys Lys His Arg Trp Ile Glu Ile Asp
195          200          205
Val Thr Ser Leu Leu Gln Pro Leu Val Thr Ser Ser Glu Arg Ser Ile
210          215          220
His Leu Ser Val Asn Phe Thr Cys Thr Lys Asp Gln Val Pro Glu Asp
225          230          235          240
Gly Val Phe Ser Met Pro Leu Ser Val Pro Pro Ser Leu Ile Leu Tyr
245          250          255
Leu Asn Asp Thr Ser Thr Gln Ala Tyr His Ser Trp Gln Ser Leu Gln
260          265          270

```

Ser Thr Trp Arg Pro Leu Gln His Pro Gly Gln Ala Gly Val Ala Ala  
 275 280 285

Arg Pro Val Lys Glu Glu Ala Thr Glu Val Glu Arg Ser Pro Arg Arg  
 290 295 300

Arg Arg Gly Gln Lys Ala Ile Arg Ser Glu Ala Lys Gly Pro Leu Leu  
 305 310 315 320

Thr Ala Ser Phe Asn Leu Ser Glu Tyr Phe Lys Gln Phe Leu Phe Pro  
 325 330 335

Gln Asn Glu Cys Glu Leu His Asp Phe Arg Leu Ser Phe Ser Gln Leu  
 340 345 350

Lys Trp Asp Asn Trp Ile Val Ala Pro His Arg Tyr Asn Pro Arg Tyr  
 355 360 365

Cys Lys Gly Asp Cys Pro Arg Ala Val Arg His Arg Tyr Gly Ser Pro  
 370 375 380

Val His Thr Met Val Gln Asn Ile Ile Tyr Glu Lys Leu Asp Pro Ser  
 385 390 395 400

Val Pro Arg Pro Ser Cys Val Pro Gly Lys Tyr Ser Pro Leu Ser Val  
 405 410 415

Leu Thr Ile Glu Pro Asp Gly Ser Ile Ala Tyr Lys Glu Tyr Glu Asp  
 420 425 430

Met Ile Ala Thr Arg Cys Thr Cys Arg  
 435 440

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 117 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: GDF-3

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..117

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Lys Arg Arg Ala Ala Ile Ser Val Pro Lys Gly Phe Cys Arg Asn Phe  
 1 5 10 15

Cys His Arg His Gln Leu Phe Ile Asn Phe Gln Asp Leu Gly Trp His  
 20 25 30

Lys Trp Val Ile Ala Pro Lys Gly Phe Met Ala Asn Tyr Cys His Gly  
 35 40 45

Glu Cys Pro Phe Ser Met Thr Thr Tyr Leu Asn Ser Ser Asn Tyr Ala  
 50 55 60  
 Phe Met Gln Ala Leu Met His Met Ala Asp Pro Lys Val Pro Lys Ala  
 65 70 75 80  
 Val Cys Val Pro Thr Lys Leu Ser Pro Ile Ser Met Leu Tyr Gln Asp  
 85 90 95  
 Ser Asp Lys Asn Val Ile Leu Arg His Tyr Glu Asp Met Val Val Asp  
 100 105 110  
 Glu Cys Gly Cys Gly  
 115

## (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 118 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vii) IMMEDIATE SOURCE:
- (B) CLONE: GDF-9
- (ix) FEATURE:
- (A) NAME/KEY: Protein
  - (B) LOCATION: 1..118
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
- Phe Asn Leu Ser Glu Tyr Phe Lys Gln Phe Leu Phe Pro Gln Asn Glu  
 1 5 10 15  
 Cys Glu Leu His Asp Phe Arg Leu Ser Phe Ser Gln Leu Lys Trp Asp  
 20 25 30  
 Asn Trp Ile Val Ala Pro His Arg Tyr Asn Pro Arg Tyr Cys Lys Gly  
 35 40 45  
 Asp Cys Pro Arg Ala Val Arg His Arg Tyr Gly Ser Pro Val His Thr  
 50 55 60  
 Met Val Gln Asn Ile Ile Tyr Glu Lys Leu Asp Pro Ser Val Pro Arg  
 65 70 75 80  
 Pro Ser Cys Val Pro Gly Lys Tyr Ser Pro Leu Ser Val Leu Thr Ile  
 85 90 95  
 Glu Pro Asp Gly Ser Ile Ala Tyr Lys Glu Tyr Glu Asp Met Ile Ala  
 100 105 110  
 Thr Arg Cys Thr Cys Arg  
 115

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 122 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: GDF-1

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..122

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

```

Pro Arg Arg Asp Ala Glu Pro Val Leu Gly Gly Gly Pro Gly Gly Ala
1           5           10
Cys Arg Ala Arg Arg Leu Tyr Val Ser Phe Arg Glu Val Gly Trp His
          20           25           30
Arg Trp Val Ile Ala Pro Arg Gly Phe Leu Ala Asn Tyr Cys Gln Gly
          35           40           45
Gln Cys Ala Leu Pro Val Ala Leu Ser Gly Ser Gly Gly Pro Pro Ala
          50           55           60
Leu Asn His Ala Val Leu Arg Ala Leu Met His Ala Ala Ala Pro Gly
65           70           75           80
Ala Ala Asp Leu Pro Cys Cys Val Pro Ala Arg Leu Ser Pro Ile Ser
          85           90           95
Val Leu Phe Phe Asp Asn Ser Asp Asn Val Val Leu Arg Gln Tyr Glu
          100          105          110
Asp Met Val Val Asp Glu Cys Gly Cys Arg
          115          120
  
```

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 118 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: Vg-1

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..118

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

Arg Arg Lys Arg Ser Tyr Ser Lys Leu Pro Phe Thr Ala Ser Asn Ile
1      5      10      15
Cys Lys Lys Arg His Leu Tyr Val Glu Phe Lys Asp Val Gly Trp Gln
20      25      30
Asn Trp Val Ile Ala Pro Gln Gly Tyr Met Ala Asn Tyr Cys Tyr Gly
35      40      45
Glu Cys Pro Tyr Pro Leu Thr Glu Ile Leu Asn Gly Ser Asn His Ala
50      55      60
Ile Leu Gln Thr Leu Val His Ser Ile Glu Pro Glu Asp Ile Pro Leu
65      70      75      80
Pro Cys Cys Val Pro Thr Lys Met Ser Pro Ile Ser Met Leu Phe Tyr
85      90      95
Asp Asn Asn Asp Asn Val Val Leu Arg His Tyr Glu Asn Met Ala Val
100     105     110
Asp Glu Cys Gly Cys Arg
115

```

## (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 118 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:  
 (B) CLONE: Vgr-1

(ix) FEATURE:  
 (A) NAME/KEY: Protein  
 (B) LOCATION: 1..118

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Arg Val Ser Ser Ala Ser Asp Tyr Asn Ser Ser Glu Leu Lys Thr Ala
1      5      10      15
Cys Arg Lys His Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Gln
20      25      30
Asp Trp Ile Ile Ala Pro Lys Gly Tyr Ala Ala Asn Tyr Cys Asp Gly
35      40      45
Glu Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His Ala
50      55      60
Ile Val Gln Thr Leu Val His Leu Met Asn Pro Glu Tyr Val Pro Lys
65      70      75      80

```



(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:  
(B) CLONE: BMP-5

(ix) FEATURE:  
(A) NAME/KEY: Protein  
(B) LOCATION: 1..118

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```

Arg Met Ser Ser Val Gly Asp Tyr Asn Thr Ser Glu Gln Lys Gln Ala
1           5           10           15
Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg Asp Leu Gly Trp Gln
20           25           30
Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala Phe Tyr Cys Asp Gly
35           40           45
Glu Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His Ala
50           55           60
Ile Val Gln Thr Leu Val His Leu Met Phe Pro Asp His Val Pro Lys
65           70           75           80
Pro Cys Cys Ala Pro Thr Lys Leu Asn Ala Ile Ser Val Leu Tyr Phe
85           90           95
Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr Arg Asn Met Val Val
100          105          110
Arg Ser Cys Gly Cys His
115

```

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 118 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:  
(B) CLONE: 60A

(ix) FEATURE:  
(A) NAME/KEY: Protein  
(B) LOCATION: 1..118

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

Ser Pro Asn Asn Val Pro Leu Leu Glu Pro Met Glu Ser Thr Arg Ser
1           5           10           15
Cys Gln Met Gln Thr Leu Tyr Ile Asp Phe Lys Asp Leu Gly Trp His
20           25           30

```



(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 117 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: BMP-4

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..117

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```

Arg Ser Pro Lys His His Ser Gln Arg Ala Arg Lys Lys Asn Lys Asn
 1           5           10
Cys Arg Arg His Ser Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asn
 20          25
Asp Trp Ile Val Ala Pro Pro Gly Tyr Gln Ala Phe Tyr Cys His Gly
 35          40          45
Asp Cys Pro Phe Pro Leu Ala Asp His Leu Asn Ser Thr Asn His Ala
 50          55          60
Ile Val Gln Thr Leu Val Asn Ser Val Asn Ser Ser Ile Pro Lys Ala
 65          70          75
Cys Cys Val Pro Thr Glu Leu Ser Ala Ile Ser Met Leu Tyr Leu Asp
 85          90          95
Glu Tyr Asp Lys Val Val Leu Lys Asn Tyr Gln Glu Met Val Val Glu
100         105         110
Gly Cys Gly Cys Arg
115
    
```

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 118 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: DPP

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..118

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

```

Lys Arg His Ala Arg Arg Pro Thr Arg Arg Lys Asn His Asp Asp Thr
1           5           10
Cys Arg Arg His Ser Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asp
20           25
Asp Trp Ile Val Ala Pro Leu Gly Tyr Asp Ala Tyr Tyr Cys His Gly
35           40           45
Lys Cys Pro Phe Pro Leu Ala Asp His Phe Asn Ser Thr Asn His Ala
50           55           60
Val Val Gln Thr Leu Val Asn Asn Met Asn Pro Gly Lys Val Pro Lys
65           70           75
Ala Cys Cys Val Pro Thr Gln Leu Asp Ser Val Ala Met Leu Tyr Leu
85           90           95
Asn Asp Gln Ser Thr Val Val Leu Lys Asn Tyr Gln Glu Met Thr Val
100          105          110
Val Gly Cys Gly Cys Arg
115
    
```

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 119 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: BMP-3

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..119

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

```

Gln Thr Leu Lys Lys Ala Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn
1           5           10
Cys Ala Arg Arg Tyr Leu Lys Val Asp Phe Ala Asp Ile Gly Trp Ser
20           25           30
Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp Ala Tyr Tyr Cys Ser Gly
35           40           45
Ala Cys Gln Phe Pro Met Pro Lys Ser Leu Lys Pro Ser Asn His Ala
50           55           60
Thr Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro
65           70           75
    
```



(vii) IMMEDIATE SOURCE:  
 (B) CLONE: Inhibin alpha

(ix) FEATURE:  
 (A) NAME/KEY: Protein  
 (B) LOCATION: 1..121

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

```

Leu Arg Leu Leu Gln Arg Pro Pro Glu Glu Pro Ala Ala His Ala Asn
1           5           10
Cys His Arg Val Ala Leu Asn Ile Ser Phe Gln Glu Leu Gly Trp Glu
20          25          30
Arg Trp Ile Val Tyr Pro Pro Ser Phe Ile Phe His Tyr Cys His Gly
35          40          45
Gly Cys Gly Leu His Ile Pro Pro Asn Leu Ser Leu Pro Val Pro Gly
50          55          60
Ala Pro Pro Thr Pro Ala Gln Pro Tyr Ser Leu Leu Pro Gly Ala Gln
65          70          75          80
Pro Cys Cys Ala Ala Leu Pro Gly Thr Met Arg Pro Leu His Val Arg
85          90          95
Thr Thr Ser Asp Gly Gly Tyr Ser Phe Lys Tyr Glu Thr Val Pro Asn
100         105         110
Leu Leu Thr Gln His Cys Ala Cys Ile
115         120
    
```

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 121 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:  
 (B) CLONE: Inhibin betaA

(ix) FEATURE:  
 (A) NAME/KEY: Protein  
 (B) LOCATION: 1..121

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

```

Arg Arg Arg Arg Arg Gly Leu Glu Cys Asp Gly Lys Val Asn Ile Cys
1           5           10          15
Cys Lys Lys Gln Phe Phe Val Ser Phe Lys Asp Ile Gly Trp Asn Asp
20          25          30
    
```

Trp Ile Ile Ala Pro Ser Gly Tyr His Ala Asn Tyr Cys Glu Gly Glu  
 35 40 45  
 Cys Pro Ser His Ile Ala Gly Thr Ser Gly Ser Ser Leu Ser Phe His  
 50 55 60  
 Ser Thr Val Ile Asn His Tyr Arg Met Arg Gly His Ser Pro Phe Ala  
 65 70 75 80  
 Asn Leu Lys Ser Cys Cys Val Pro Thr Lys Leu Arg Pro Met Ser Met  
 85 90 95  
 Leu Tyr Tyr Asp Asp Gly Gln Asn Ile Ile Lys Lys Asp Ile Gln Asn  
 100 105 110  
 Met Ile Val Glu Glu Cys Gly Cys Ser  
 115 120

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 120 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: Inhibin betaB

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..120

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Arg Ile Arg Lys Arg Gly Leu Glu Cys Asp Gly Arg Thr Asn Leu Cys  
 1 5 10 15  
 Cys Arg Gln Gln Phe Phe Ile Asp Phe Arg Leu Ile Gly Trp Asn Asp  
 20 25 30  
 Trp Ile Ile Ala Pro Thr Gly Tyr Tyr Gly Asn Tyr Cys Glu Gly Ser  
 35 40 45  
 Cys Pro Ala Tyr Leu Ala Gly Val Pro Gly Ser Ala Ser Ser Phe His  
 50 55 60  
 Thr Ala Val Val Asn Gln Tyr Arg Met Arg Gly Leu Asn Pro Gly Thr  
 65 70 75 80  
 Val Asn Ser Cys Cys Ile Pro Thr Lys Leu Ser Thr Met Ser Met Leu  
 85 90 95  
 Tyr Phe Asp Asp Glu Tyr Asn Ile Val Lys Arg Asp Val Pro Asn Met  
 100 105 110

Ile Val Glu Glu Cys Gly Cys Ala  
 115 120

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 114 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: TGF-beta1

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..114

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Arg	Arg	Ala	Leu	Asp	Thr	Asn	Tyr	Cys	Phe	Ser	Ser	Thr	Glu	Lys	Asn
1				5					10					15	
Cys	Cys	Val	Arg	Gln	Leu	Tyr	Ile	Asp	Phe	Arg	Lys	Asp	Leu	Gly	Trp
			20					25					30		
Lys	Trp	Ile	His	Glu	Pro	Lys	Gly	Tyr	His	Ala	Asn	Phe	Cys	Leu	Gly
		35					40					45			
Pro	Cys	Pro	Tyr	Ile	Trp	Ser	Leu	Asp	Thr	Gln	Tyr	Ser	Lys	Val	Leu
	50					55					60				
Ala	Leu	Tyr	Asn	Gln	His	Asn	Pro	Gly	Ala	Ser	Ala	Ala	Pro	Cys	Cys
65					70					75					80
Val	Pro	Gln	Ala	Leu	Glu	Pro	Leu	Pro	Ile	Val	Tyr	Tyr	Val	Gly	Arg
				85					90					95	
Lys	Pro	Lys	Val	Glu	Gln	Leu	Ser	Asn	Met	Ile	Val	Arg	Ser	Cys	Lys
			100					105					110		
Cys	Ser														

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 114 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:

(B) CLONE: TGF-beta2

(ix) FEATURE:

(A) NAME/KEY: Protein  
(B) LOCATION: 1..114

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```

Lys Arg Ala Leu Asp Ala Ala Tyr Cys Phe Arg Asn Val Gln Asp Asn
1          5          10          15
Cys Cys Leu Arg Pro Leu Tyr Ile Asp Phe Lys Arg Asp Leu Gly Trp
20          25          30
Lys Trp Ile His Glu Pro Lys Gly Tyr Asn Ala Asn Phe Cys Ala Gly
35          40          45
Ala Cys Pro Tyr Leu Trp Ser Ser Asp Thr Gln His Ser Arg Val Leu
50          55          60
Ser Leu Tyr Asn Thr Ile Asn Pro Glu Ala Ser Ala Ser Pro Cys Cys
65          70          75          80
Val Ser Gln Asp Leu Glu Pro Leu Thr Ile Leu Tyr Tyr Ile Gly Lys
85          90          95
Thr Pro Lys Ile Glu Gln Leu Ser Asn Met Ile Val Lys Ser Cys Lys
100         105         110
Cys Ser

```

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 114 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:

(B) CLONE: TGF-beta3

(ix) FEATURE:

(A) NAME/KEY: Protein  
(B) LOCATION: 1..114

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

```

Lys Arg Ala Leu Asp Thr Asn Tyr Cys Phe Arg Asn Leu Glu Glu Asn
1          5          10          15
Cys Cys Val Arg Pro Leu Tyr Ile Asp Phe Arg Gln Asp Leu Gly Trp
20          25          30
Lys Trp Val His Glu Pro Lys Gly Tyr Tyr Ala Asn Phe Cys Ser Gly
35          40          45

```

Pro Cys Pro Tyr Leu Arg Ser Ala Asp Thr Thr His Ser Thr Val Leu  
 50 55 60  
 Gly Leu Tyr Asn Thr Leu Asn Pro Glu Ala Ser Ala Ser Pro Cys Cys  
 65 70 75 80  
 Val Pro Gln Asp Leu Glu Pro Leu Thr Ile Leu Tyr Tyr Val Gly Arg  
 85 90 95  
 Thr Pro Lys Val Glu Gln Leu Ser Asn Met Val Val Lys Ser Cys Lys  
 100 105 110  
 Cys Ser

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 116 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vii) IMMEDIATE SOURCE:
 

- (B) CLONE: TGF-beta4

(ix) FEATURE:
 

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..116

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Arg Arg Asp Leu Asp Thr Asp Tyr Cys Phe Gly Pro Gly Thr Asp Glu  
 1 5 10 15  
 Lys Asn Cys Cys Val Arg Pro Leu Tyr Ile Asp Phe Arg Lys Asp Leu  
 20 25 30  
 Gln Trp Lys Trp Ile His Glu Pro Lys Gly Tyr Met Ala Asn Phe Cys  
 35 40 45  
 Met Gly Pro Cys Pro Tyr Ile Trp Ser Ala Asp Thr Gln Tyr Thr Lys  
 50 55 60  
 Val Leu Ala Leu Tyr Asn Gln His Asn Pro Gly Ala Ser Ala Ala Pro  
 65 70 75 80  
 Cys Cys Val Pro Gln Thr Leu Asp Pro Leu Pro Ile Ile Tyr Tyr Val  
 85 90 95  
 Gly Arg Asn Val Arg Val Glu Gln Leu Ser Asn Met Val Val Arg Ala  
 100 105 110  
 Cys Lys Cys Ser  
 115

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:




Met Ala Arg Pro Asn Lys Phe Leu Leu Trp Phe Cys Cys Phe Ala Trp  
 1 5 10 15  
 Leu Cys Phe Pro Ile Ser Leu Gly Ser Gln Ala Ser Gly Gly Glu Ala  
 20 25 30  
 Gln Ile Ala Ala Ser Ala Glu Leu Glu Ser Gly Ala Met Pro Trp Ser  
 35 40 45  
 Leu Leu Gln His Ile Asp Glu Arg Asp Arg Ala Gly Leu Leu Pro Ala  
 50 55 60  
 Leu Phe Lys Val Leu Ser Val Gly Arg Gly Gly Ser Pro Arg Leu Gln  
 65 70 75 80  
 Pro Asp Ser Arg Ala Leu His Tyr Met Lys Lys Leu Tyr Lys Thr Tyr  
 85 90 95  
 Ala Thr Lys Glu Gly Ile Pro Lys Ser Asn Arg Ser His Leu Tyr Asn  
 100 105 110  
 Thr Val Arg Leu Phe Thr Pro Cys Thr Arg His Lys Gln Ala Pro Gly  
 115 120 125  
 Asp Gln Val Thr Gly Ile Leu Pro Ser Val Glu Leu Leu Phe Asn Leu  
 130 135 140  
 Asp Arg Ile Thr Thr Val Glu His Leu Leu Lys Ser Val Leu Leu Tyr  
 145 150 155 160  
 Asn Ile Asn Asn Ser Val Ser Phe Ser Ser Ala Val Lys Cys Val Cys  
 165 170 175  
 Asn Leu Met Ile Lys Glu Pro Lys Ser Ser Ser Arg Thr Leu Gly Arg  
 180 185 190  
 Ala Pro Tyr Ser Phe Thr Phe Asn Ser Gln Phe Glu Phe Gly Lys Lys  
 195 200 205  
 His Lys Trp Ile Gln Ile Asp Val Thr Ser Leu Leu Gln Pro Leu Val  
 210 215 220  
 Ala Ser Asn Lys Arg Ser Ile His Met Ser Ile Asn Phe Thr Cys Met  
 225 230 235 240  
 Lys Asp Gln Leu Glu His Pro Ser Ala Gln Asn Gly Leu Phe Asn Met  
 245 250 255  
 Thr Leu Val Ser Pro Ser Leu Ile Leu Tyr Leu Asn Asp Thr Ser Ala  
 260 265 270  
 Gln Ala Tyr His Ser Trp Tyr Ser Leu His Tyr Lys Arg Arg Pro Ser  
 275 280 285  
 Gln Gly Pro Asp Gln Glu Arg Ser Leu Ser Ala Tyr Pro Val Gly Glu  
 290 295 300  
 Glu Ala Ala Glu Asp Gly Arg Ser Ser His His Arg His Arg Arg Gly  
 305 310 315 320  
 Gln Glu Thr Val Ser Ser Glu Leu Lys Lys Pro Leu Gly Pro Ala Ser  
 325 330 335



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/21199

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) :A61K 38/18, 31/185 US CL :424/85.1, 130.1, 139.1, 141.1, 142.1; 514/44 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/85.1, 130.1, 139.1, 141.1, 142.1; 514/44  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, GenEMBL sequence databases		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DONG, J. et al. Growth differentiation factor-9 is required during early ovarian folliculogenesis. Nature. 10 October 1996, Vol. 383, pages 531-535, see entire document.	1-11
A	MASSAGUE, J. The TGF- $\beta$ family of growth and differentiation factors. Cell. March 1987, Vol. 49, pages 437-438, see entire document.	1-11
A,P	FITZPATRICK, S. L. et al. Expression of growth differentiation factor-9 messenger ribonucleic acid in ovarian and nonovarian rodent and human tissues. Endocrinology. 1998, Vol. 139, No. 5, pages 2571-2578, see entire document.	1-11
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *p* document published prior to the international filing date but later than the priority date claimed		*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *&* document member of the same patent family
Date of the actual completion of the international search 31 DECEMBER 1998		Date of mailing of the international search report 26 JAN 1999
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer ELIZABETH C. KEMMERER  Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/21199

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2.  Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3.  Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/21199

**BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING**

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-6, drawn to a method for inhibiting oocyte maturation comprising administering a GDF-9 antibody.

Group II, claim(s) 1 and 7, drawn to a method for inhibiting oocyte maturation comprising administering GDF-9 antisense molecule.

Group III, claim(s) 8-11, drawn to a method for inducing an immune response in a subject to GDF-9 comprising administering GDF-9 or a fragment thereof.

The inventions listed as Groups I-III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the special technical feature of Group I is the active method step of administering a GDF-9 antibody. The other Groups do not require administration of an antibody, and thus do not share a special technical feature with Group I. Furthermore, the agents administered in the methods of Groups II and III have completely different physical and chemical properties.