



(86) Date de dépôt PCT/PCT Filing Date: 2000/03/29
(87) Date publication PCT/PCT Publication Date: 2000/10/12
(45) Date de délivrance/Issue Date: 2007/05/01
(85) Entrée phase nationale/National Entry: 2001/08/28
(86) N° demande PCT/PCT Application No.: EP 2000/002769
(87) N° publication PCT/PCT Publication No.: 2000/060061
(30) Priorités/Priorities: 1999/03/31 (US09/282,995);
2000/02/11 (US09/502,852)

(51) Cl.Int./Int.Cl. *C12N 5/10* (2006.01),
A01H 1/00 (2006.01), *A01H 5/00* (2006.01),
A01N 63/00 (2006.01), *C12N 15/29* (2006.01),
C12N 15/54 (2006.01), *C12N 15/63* (2006.01),
C12N 15/82 (2006.01), *C12N 5/04* (2006.01),
C12N 9/10 (2006.01)

(72) Inventeurs/Inventors:
HOHN, THOMAS M., US;
PETERS, CHERYL, US;
SALMERON, JOHN MANUEL, US;
REED, JANET N., US;
DAWSON, JOHN LUTHER, US

(73) Propriétaire/Owner:
SYNGENTA PARTICIPATIONS AG, CH

(54) Titre : PLANTES TRANSGENIQUES ET PROCEDES

(54) Title: TRANSGENIC PLANT RESISTANT TO MYCOTOXINS AND METHODS

(57) Abrégé/Abstract:

The invention relates to a plant cell comprising a heterologous polynucleotide encoding a gene product that is expressed in the plant cell wherein the gene product comprises trichothecene resistance activity



(74) **Agent:** FETHERSTONHAUGH & CO.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 October 2000 (12.10.2000)

PCT

(10) International Publication Number
WO 00/60061 A3

(51) International Patent Classification⁷: **C12N 9/10**,
15/54, 15/29, 15/82, 15/63, 5/04, A01H 5/00

Luther [US/US]; 214 East Avondale Drive, Greensboro,
NC 27403 (US).

(21) International Application Number: PCT/EP00/02769

(74) Agent: BECKER, Konrad; Syngenta Participations AG,
Intellectual Property, P.O. Box, CH-4002 Basel (CH).

(22) International Filing Date: 29 March 2000 (29.03.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/282,995 31 March 1999 (31.03.1999) US
09/502,852 11 February 2000 (11.02.2000) US

(71) Applicant (*for AT, US only*): NOVARTIS AG [CH/CH];
Schwarzwaldallee 215, CH-4058 Basle (CH).

(71) Applicant (*for AT only*): NOVARTIS-ERFINDUN-
GEN VERWALTUNGSEGESELLSCHAFT M.B.H.
[AT/AT]; Brunner Strasse 59,, 1230 Vienna (AT).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): HOHN, Thomas,
M. [US/US]; 102 Blakely Drive, Chapel Hill, NC 27514
(US). PETERS, Cheryl [US/US]; 6308 Valley Estates
Drive, Raleigh, NC 27612 (US). SALMERON, John,
Manuel [US/US]; 1308 Blackberry Lane, Hillsborough,
NC 27278 (US). REED, Janet, N. [US/US]; 5415 Lake
Vista Drive, Durham, NC 27712 (US). DAWSON, John,

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— *With international search report.*

(88) Date of publication of the international search report:
19 April 2001

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: TRANSGENIC PLANT RESISTANT TO MYCOTOXINS AND METHODS

(57) Abstract: The invention relates to a plant cell comprising a heterologous polynucleotide encoding a gene product that is ex-
pressed in the plant cell wherein the gene product comprises trichothecene resistance activity

WO 00/60061 A3

WO 00/60061

PCT/EP00/02769

Transgenic Plant and Methods

The present invention relates to transgenic hosts particularly transgenic plants, plant tissues, seeds and cells that are trichothecene resistant and methods of making and using the same. The present invention further relates to methods of preventing and/or reducing fungal growth on a plant, plant tissue, seed or plant cell. The present invention further relates to preventing and/or reducing mycotoxin contamination of a plant, plant tissue or seed. The present invention further relates to using trichothecenes as selective agents in transformation protocols.

Numerous fungi are serious pests of economically important agricultural crops. Further, crop contamination by fungal toxins is a major problem for agriculture throughout the world. Mycotoxins are toxic fungal metabolites, often found in agricultural products that are characterized by their ability to cause health problems for vertebrates. Trichothecenes are sesquiterpene epoxide mycotoxins produced by species of *Fusarium*, *Trichothecium*, and *Myrothecium* that act as potent inhibitors of eukaryotic protein synthesis. *Fusarium* species that produce such trichothecenes include *F. acuminatum*, *F. crookwellense*, *F. culmorum*, *F. equiseti*, *F. graminearum* (*Gibberella zeae*), *F. lateritium*, *F. poae*, *F. sambucinum* (*G. pulicaris*), and *F. sporotrichioides* (Marasas, W.F.O., Nelson, P.E., and Toussoun, T.A. 1984).

As previously described (A. E.Desjardins and T. M Hohn, Mycotoxins in plant pathogenesis.Mol.Plant-Microbe Interact. 10 (2):147-152, 1997), both acute and chronic mycotoxicoses in farm animals and in humans have been associated with consumption of wheat, rye, barley, oats, rice and maize contaminated with *Fusarium* species that produce trichothecene mycotoxins. Experiments with chemically pure trichothecenes at low dosage levels have reproduced many of the features observed in moldy-grain toxicoses in animals, including anemia and immunosuppression, hemorrhage, emesis and feed refusal. Historical and epidemiological data from human populations indicate an association between certain disease epidemics and consumption of grain infected with *Fusarium* species that produce trichothecenes. In particular, outbreaks of a fatal disease known as alimentary toxic aleukia, which has occurred in Russia since the nineteenth century, have been associated with consumption of over-wintered grains contaminated with *Fusarium* species that produce the trichothecene T-2 toxin. In Japan, outbreaks of a similar disease called akakabi-byo or red

mold disease have been associated with grain infected with *Fusarium* species that produce the trichothecene, deoxynivalenol (hereinafter "DON"). Trichothecenes were detected in the toxic grain samples responsible for recent human disease outbreaks in India and Japan. There exists, therefore, a need for agricultural methods for preventing and, crops having reduced levels of, mycotoxin contamination.

Further, trichothecene-producing *Fusarium* species are destructive pathogens and attack a wide range of plant species. The acute phytotoxicity of trichothecenes and their occurrence in plant tissues also suggest that these mycotoxins play a role in the pathogenesis of *Fusarium* on plants. This implies that mycotoxins play a role in disease and, therefore, reducing their toxicity to the plant may also prevent or reduce disease in the plant. Further, reduction in disease levels may have the additional benefit of reducing mycotoxin contamination on the plant and particularly in grain where the plant is a cereal plant.

Various methods of controlling diseases in plants, such as corn ear rot, stock rot or wheat head blight, have been used with varying degrees of success. One method of controlling plant disease has been to apply an antimicrobial chemical to crops. This method has numerous, art-recognized problems. Alternatively, a more recent method involves the use of biological control organisms ("biocontrol") which are natural competitors or inhibitors of the pest organism. However, it is difficult to apply biocontrol to large areas, and even more difficult to cause those living organisms to remain in the treated area for an extended period of time. More recently, techniques in recombinant DNA have provided the opportunity to insert into plant cells cloned genes, which express antimicrobial compounds. However, this technology has given rise to concerns about eventual microbial resistance to well-known, naturally occurring antimicrobials. Thus, a continuing need exists to identify naturally occurring antimicrobial agents, such as proteins, which can be formed by plant cells directly by translation of a single gene.

A trichothecene 3-O-acetyltransferase that catalyzes the acetylation of a number of different *Fusarium* trichothecenes including DON at the C3 hydroxyl group has been identified in *Fusarium sporotrichioides*. (S. P. McCormick, N. J. Alexander, S. C. Trapp, and T. M. Hohn. Disruption of TRI101, the gene encoding trichothecene 3-O-acetyltransferase, from *Fusarium sporotrichioides*. *Applied Environ. Microbiol.* 65 (12):5252-5256, 1999.) Acetylation of trichothecenes at the C3-OH significantly reduces their toxicity in vertebrates and plants and results in the reaction product 3-acetyldeoxyvalenol (hereinafter "3ADON") See, Kimura et al. below.

WO 00/60061

PCT/EP00/02769

The sequence of structural genes encoding trichothecene 3-O-acetyl transferases from *Fusarium graminearum*, *Fusarium sporotrichioides* as well as sequences of other orthologs has been published. See, e.g. Kimura et al., Biosci. Biotechnol. Biochem., 62 (5) 1033-1036 (1998), and Kimura et al., FEBS Letters, 435, 163-168 (1998). Further, it has been speculated that the gene from *Fusarium sporotrichioides* encoding a trichothecene 3-O-acetyl transferase may be useful in developing plant varieties with increased resistance to *Fusarium*. See., e.g. Hohn, T.M. et al. Molecular Genetics of Host-Specific Toxins in Plant Disease, 17-24 (1998), and Kimura et al. J.Biological Chemistry, 273(3) 1654-1661 (1998).

Prior to the present invention, however, many uncertainties rendered it far from obvious whether expressing trichothecene 3-O-acetyl transferases in a plant would actually lead to trichothecene resistant plants. For example, the reaction catalyzed by the *Fusarium sporotrichioides* trichothecene 3-O-acetyl transferase is reversible and might, therefore have failed to protect plant cells from trichothecenes such as DON. It was also uncertain whether there might be esterases in plant cells that would compete with the 3-O-acetyl transferase activities to generate toxic DON from 3ADON. It was also uncertain how the metabolism of the reaction product 3ADON might affect the plant, e.g. whether introduction of the trichothecene 3-O-acetyltransferase would alter plant growth and development in ways that would negate any positive contribution of the acetyltransferase by for example, interfering with the plant's natural disease resistance mechanisms. It was also uncertain whether 3ADON could be metabolized by the plant to form a novel secondary metabolite with toxic effects. It was also uncertain, even if DON produced by an invading fungus was efficiently converted to 3ADON, whether this conversion would impart enhanced pathogen resistance upon the plant. The above are but a few of the uncertainties in the art before the time of the present invention.

It is an object of the invention to provide a plant cell or cells comprising a heterologous polynucleotide encoding a gene product that is expressed in the plant cell wherein the gene product comprises trichothecene resistance activity.

Another object of the invention is to provide a plant comprising the above described plant cell wherein the plant is resistant to a trichothecene.

Another object of the invention is to provide a plant of the invention that is resistant to a trichothecene where the trichothecene comprises a C-3 hydroxyl group.

Another object of the invention is to provide a plant of the invention that is resistant to a fungus that produces a trichothecene, preferably a trichothecene comprising a C-3 hydroxyl

WO 00/60061

PCT/EP00/02769

group.

Another object of the invention is to provide a plant of the invention that is resistant to *Fusarium*, *Trichothecium* or *Myrothecium*.

Another object of the invention is to provide a plant of the invention that is resistant to *Fusarium*, in particular but not limited to *Fusarium graminearum*, *Fusarium culmorum*, *Fusarium sporotrichioides*, *Fusarium poae*, *Fusarium sambucinum*, *Fusarium equiseti*, *Fusarium acuminatum*, *Fusarium lateritium*, and *Fusarium pseudograminearum*.

Another object of the invention is to provide a plant of the invention, wherein the plant is resistant to *Fusarium graminearum*.

Another object of the invention is to provide a plant of the invention comprising a heterologous polynucleotide encoding a gene product that is expressed in the plant cell wherein the gene product comprises trichothecene resistance activity, wherein the heterologous polynucleotide is a microbial polynucleotide, preferably a yeast or fungal polynucleotide.

Another object of the invention is to provide a plant of the invention, wherein the fungal polynucleotide is a *Fusarium* polynucleotide, preferably is a *Fusarium graminearum* or *Fusarium sporotrichioides* polynucleotide.

Another object of the invention is to provide a plant of the invention wherein the heterologous polynucleotide comprises a sequence substantially similar to the nucleic acid sequence of SEQ ID NOs:1, 5 or 7.

Another object of the invention is to provide a plant of the invention wherein the heterologous polynucleotide comprises the nucleic acid sequence of SEQ ID NO:1, 5 or 7 or homologs thereof.

Another object of the invention is to provide a plant of the invention comprising a heterologous polynucleotide, which comprises a consecutive at least 80 base pair portion identical in sequence to a consecutive 80 base pair portion set forth in SEQ ID NO:1, 5 or 7, preferably a consecutive at least 50 base pair portion identical in sequence to a consecutive 50 base pair portion set forth in SEQ ID NO:1, 5 or 7, and more preferably a consecutive at least 21 base pair portion identical in sequence to a consecutive 21 base pair portion set forth in SEQ ID NO:1, 5 or 7, and most preferably a consecutive 18 base pair portion identical in sequence to a consecutive 18 base pair portion set forth in SEQ ID NO:1, 5 or 7.

WO 00/60061

PCT/EP00/02769

Another object of the invention is to provide a plant of the invention resistant to a trichothecene selected from the group consisting T-2 toxin, HT-2 toxin, isotrichodermol, 4,15-diacetoxyscripenol (hereinafter "DAS"), 3-deacetylcalonectrin, 3,15-dideacetylcalonectrin, scirpentriol, neosolaniol; type B: 15-acetyldeoxynivalenol, nivalenol, 4-acetylnivalenol (fusarenone-X), 4,15-diacetylnivalenol, 4,7,15-acetylnivalenol, and DON.

Another object of the invention is to provide a plant of the invention resistant to DAS or DON.

Another object of the invention is to provide a plant wherein the gene product is encoded by a polynucleotide according to the invention.

Another object of the invention is to provide a plant of the invention wherein the gene product is a 3-O-acetyltransferase.

Another object of the invention is to provide a plant of the invention wherein the gene product is a polypeptide comprising a sequence substantially similar to SEQ ID NO:2, 6 or 8.

Another object of the invention is to provide a seed of any of the plants of the invention.

Another object of the invention is to provide anyone of the above-described plants wherein the plant is a wheat, maize, barley or rice plant.

Another object of the invention is to provide a method for producing a trichothecene resistant plant comprising the steps of:

- (a) transforming a plant cell with a heterologous gene encoding a gene product, wherein the gene product increases resistance to a trichothecene; and
- (b) expressing the gene product at a biologically significant level.
- (c) regenerating the plant cell into a plant; and
- (d) selecting a plant having increased resistance to a trichothecene.

Another object of the invention is to provide a method as described above further comprising the step of selecting a plant on which there is reduced growth of a fungus where the fungus produces a trichothecene.

Another object of the invention is to provide a method as described above wherein the fungus is of the genus *Fusarium*.

Another object of the invention is to provide a trichothecene resistant plant obtained according to the above-described methods.

WO 00/60061

PCT/EP00/02769

Another object of the invention is to provide a seed produced by selfing or outcrossing a plant of the invention as described above, wherein a plant grown from the seed has an increased resistance to trichothecene.

Another object of the invention is to provide a method of preventing mycotoxin contamination of a plant, including of a plant's seed, comprising growing a plant of the invention as described above, preferably in an area with moderate to severe fungal infestation, wherein the plant is preferably a crop plant.

Another object of the invention is to provide a method of preventing fungal growth on a plant, preferably growth of a fungus of the genus *Fusarium* comprising growing a plant of the invention that produces a trichothecene, preferably a trichothecene comprising a C-3 hydroxyl group as described above, preferably in an area with moderate to severe fungal infestation, wherein the plant is preferably a crop plant.

Another object of the invention is to provide a method of producing a fungal resistant plant comprising

- (a) transforming a plant cell with a heterologous gene encoding a gene product, wherein the gene product increases resistance to a trichothecene;
- (b) expressing the gene product at a biologically significant level;
- (c) regenerating the plant cell into a plant; and
- (d) selecting a plant with increased resistance to a trichothecene; and
- (e) optionally, selfing or outcrossing the plant obtained in step (d)

Another object of the invention is to provide the method, wherein the fungus is of the genus *Fusarium*.

Another object of the invention is to provide a method of producing seed wherein the plant grown from the seed is fungal resistant, comprising selfing or outcrossing the plant of the invention.

Another object of the invention is to provide the above method, wherein the seed is resistant to fungi to the genus *Fusarium*.

Another object of the invention is to provide a method of selecting transformed host cells, the method comprising:

transforming a host cell with a nucleic acid construct encoding a trichothecene 3-O-acetyltransferase, and

WO 00/60061

PCT/EP00/02769

growing the transformed host cell in the presence of a trichothecene selective agent.

Another object of the invention is to provide a method of selecting transformed host cells wherein the host cells are plant cells, or microbial cells, particularly where the microbial cells are fungal cells.

Another object of the invention is to provide a method of selecting transformed host cells as described above where the host cell is further transformed with a second polynucleotide of interest.

Another object of the invention is to provide a method of selecting transformed host cells wherein in the trichothecene is selected from the group the group consisting T-2 toxin, HT-2 toxin, isotrichodermol, DAS, 3-deacetylcalonecetrin, 3,15-dideacetylcalonecetrin, scirpentriol, neosolaniol; type B: 15-acetyldeoxynivalenol, nivalenol, 4-acetylnivalenol (fusarenone-X), 4,15-diacetylnivalenol, 4,7,15-acetylnivalenol, and DON.

Another object of the invention is to provide a trichothecene resistant plant obtainable by the method of the invention.

Another object of the invention is to provide a fungal resistant plant obtainable by the method of the invention.

Another object of the invention is to provide a plant seed obtainable by the method of the invention.

For clarity, certain terms used in the specification are defined and presented as follows:

"Expression" refers to the transcription and/or translation of an endogenous gene or a transgene in plants. In the case of antisense constructs, for example, expression may refer to the transcription of the antisense DNA only.

"Operably linked/associated" when referring to a regulatory DNA sequence being "operably linked to" or "associated with" a DNA sequence that codes for an RNA or a protein refers to the two sequences being situated such that the regulatory DNA sequence affects expression of the coding DNA sequence.

The term "heterologous polynucleotide" or "heterologous DNA" as used herein each refers to a nucleic acid molecule not naturally associated with a host cell into which it is introduced, including genetic constructs, non-naturally occurring multiple copies of a naturally occurring nucleic acid molecule; and an otherwise homologous nucleic acid molecule operatively linked to a non-native nucleic acid molecule. Thus, a heterologous gene in a host cell includes a gene that is endogenous to the particular host cell but has

WO 00/60061

PCT/EP00/02769

been modified through, for example, the use of DNA shuffling. Thus, the term encompasses a DNA segment that is foreign or heterologous to the cell, or homologous to the cell but in a position within the host cell nucleic acid in which the element is not ordinarily found.

The terms "nucleic acid" or "polynucleotide" refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides which have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.* degenerate codon substitutions) and complementary sequences and as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer *et al.*, *Nucleic Acid Res.* 19: 5081 (1991); Ohtsuka *et al.*, *J. Biol. Chem.* 260: 2605-2608 (1985); Rossolini *et al.*, *Mol. Cell. Probes* 8: 91-98 (1994)). The terms "nucleic acid" or "nucleic acid sequence" or "polynucleotide" may also be used interchangeably with gene, cDNA, and mRNA encoded by a gene.

In its broadest sense, the term "substantially similar", when used herein with respect to a nucleic acid molecule, means a nucleic acid molecule corresponding to a reference nucleotide sequence, wherein the corresponding nucleic acid molecule encodes a polypeptide having substantially the same structure and function as the polypeptide encoded by the reference nucleotide sequence, *e.g.* where only changes in amino acids not affecting the polypeptide function occur. Desirably the substantially similar nucleic acid molecule encodes the polypeptide encoded by the reference nucleotide sequence. The term "substantially similar" is specifically intended to include nucleic acid molecules wherein the sequence has been modified to optimize expression in particular cells, *e.g.* in plant cells. The percentage of identity between the substantially similar nucleic acid molecule and the reference nucleotide sequence desirably is at least 45%, more desirably at least 65%, more desirably at least 75%, preferably at least 85%, more preferably at least 90%, still more preferably at least 95%, yet still more preferably at least 99%. Preferably, the percentage of identity exists over a region of the sequences that is at least about 50 residues in length, more preferably over a region of at least about 100 residues, and most preferably the sequences are substantially similar over at least about 150 residues. In a most preferred

WO 00/60061

PCT/EP00/02769

embodiment, the sequences are substantially similar over the entire length of the coding regions. Sequence comparisons may be carried out using a Smith-Waterman sequence alignment algorithm and as described in more detail below (see e.g. Waterman, M.S. Introduction to Computational Biology: Maps, sequences and genomes. Chapman & Hall. London: 1995. ISBN 0-412-99391-0, or at <http://www-hto.usc.edu/software/seqaln/index.html>). The local S program, version 1.16, is used with following parameters: match: 1, mismatch penalty: 0.33, open-gap penalty: 2, extended-gap penalty: 2.

Another indication that a nucleic acid sequences is a substantially similar nucleic acid of the invention is that it hybridizes to a nucleic acid molecule of the invention under stringent conditions. The phrase "hybridizing specifically to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent conditions when that sequence is present in a complex mixture (e.g., total cellular) DNA or RNA. "Bind(s) substantially" refers to complementary hybridization between a probe nucleic acid and a target nucleic acid and embraces minor mismatches that can be accommodated by reducing the stringency of the hybridization media to achieve the desired detection of the target nucleic acid sequence.

"Stringent hybridization conditions" and "stringent hybridization wash conditions" in the context of nucleic acid hybridization experiments such as Southern and Northern hybridizations are sequence dependent, and are different under different environmental parameters. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes* part I chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays" Elsevier, New York. Generally, highly stringent hybridization and wash conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. Typically, under "stringent conditions" a probe will hybridize to its target subsequence, but to no other sequences.

The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent conditions are selected to be equal to the T_m for a particular probe. An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is 50% formamide with 1 mg of heparin at 42°C, with the hybridization being carried out overnight. An example of highly stringent wash conditions is 0.1 5M NaCl at 72°C for about 15 minutes. An example of

WO 00/60061

PCT/EP00/02769

stringent wash conditions is a 0.2x SSC wash at 65°C for 15 minutes (see, Sambrook, *infra*, for a description of SSC buffer). Often, a high stringency wash is preceded by a low stringency wash to remove background probe signal. An example medium stringency wash for a duplex of, e.g., more than 100 nucleotides, is 1x SSC at 45°C for 15 minutes. An example low stringency wash for a duplex of, e.g., more than 100 nucleotides, is 4-6x SSC at 40°C for 15 minutes. For short probes (e.g., about 10 to 50 nucleotides), stringent conditions typically involve salt concentrations of less than about 1.0M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3, and the temperature is typically at least about 30°C. Stringent conditions can also be achieved with the addition of destabilizing agents such as formamide. In general, a signal to noise ratio of 2x (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization. Nucleic acids that do not hybridize to each other under stringent conditions are still substantially similar if the proteins that they encode are substantially similar. This occurs, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

The following are examples of sets of hybridization/wash conditions that may be used to identify homologous nucleotide sequences that are substantially similar to reference nucleotide sequences of the present invention: a test sequence that hybridizes to the reference nucleotide sequence in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 2X SSC, 0.1% SDS at 50°C, more desirably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 1X SSC, 0.1% SDS at 50°C, more desirably still in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.5X SSC, 0.1% SDS at 50°C, preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 50°C, more preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 65°C. The polynucleotide of the invention that hybridize under the above conditions preferably comprises at least 80 base pairs, more preferably at least 50 base pairs and particularly at least 21, and more particularly 18 base pairs. Preferred homologs of use in the invention include nucleic acid molecules that encode an amino acid sequence that is at least 45% identical to SEQ ID NO:2, 6 or 8 as measured, using the parameters described below, wherein the amino acid sequence encoded by the homolog has trichothecene resistance activity, e.g. 3-O-acetyl transferase activity.

The term "substantially similar", when used herein with respect to a protein, means a protein corresponding to a reference protein, wherein the protein has substantially the same

WO 00/60061

PCT/EP00/02769

structure and function as the reference protein, e.g. where only changes in amino acids sequence not affecting the polypeptide function occur. When used for a protein or an amino acid sequence the percentage of identity between the substantially similar and the reference protein or amino acid sequence desirably is at least 45% identity, more desirably at least 65%, more desirably at least 75%, preferably at least 85%, more preferably at least 90%, still more preferably at least 95%, yet still more preferably at least 99%, using default BLAST analysis parameters and as described in more detail below.

Preferred homologs of the polypeptide of use in the invention comprise those having amino acid sequences that are at least 45% identical to SEQ ID NO:2, 6 or 8, wherein the amino acid sequence encoded by the homolog has trichothecene resistance activity, e.g. 3-O-acetyl transferase activity.

Optimal alignment of nucleic acid or protein sequences for comparison can be conducted as described above and, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2: 482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48: 443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (*see generally*, Ausubel *et al.*, *infra*).

One example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul *et al.*, *J. Mol. Biol.* 215: 403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, 1990). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when the cumulative alignment score falls off by the quantity X from its maximum achieved value, the cumulative score goes to

WO 00/60061

PCT/EP00/02769

zero or below due to the accumulation of one or more negative-scoring residue alignments, or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89: 10915 (1989)).

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90: 5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a test nucleic acid sequence is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid sequence to the reference nucleic acid sequence is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

A further indication that two nucleic acid sequences or proteins are substantially similar is that the protein encoded by the first nucleic acid is immunologically cross reactive with, or specifically binds to, the protein encoded by the second nucleic acid. Thus, a protein is typically substantially similar to a second protein, for example, where the two proteins differ only by conservative substitutions.

The phrase "specifically (or selectively) binds to an antibody," or "specifically (or selectively) immunoreactive with," when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the protein in the presence of a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein and do not bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, antibodies raised to the protein with the amino acid sequence encoded by any of the nucleic acid sequences of the invention can be selected to obtain antibodies specifically immunoreactive with that protein and not with other proteins except for polymorphic variants. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays, Western blots, or immunohistochemistry are routinely used to select

WO 00/60061

PCT/EP00/02769

monoclonal antibodies specifically immunoreactive with a protein. See Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York "Harlow and Lane"), for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity. Typically a specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background.

"Conservatively modified variations" of a particular nucleic acid sequence refers to those nucleic acid sequences that encode identical or essentially identical amino acid sequences, or where the nucleic acid sequence does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given polypeptide. For instance the codons CGT, CGC, CGA, CGG, AGA, and AGG all encode the amino acid arginine. Thus, at every position where an arginine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded protein. Such nucleic acid variations are "silent variations" which are one species of "conservatively modified variations." Every nucleic acid sequence described herein which encodes a protein also describes every possible silent variation, except where otherwise noted. One of skill will recognize that each codon in a nucleic acid (except ATG, which is ordinarily the only codon for methionine) can be modified to yield a functionally identical molecule by standard techniques. Accordingly, each "silent variation" of a nucleic acid which encodes a protein is implicit in each described sequence.

Furthermore, one of skill will recognize that individual substitutions deletions or additions that alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 1%) in an encoded sequence are "conservatively modified variations," where the alterations result in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. The following five groups each contain amino acids that are conservative substitutions for one another: Aliphatic: Glycine (G), Alanine (A), Valine (V), Leucine (L), Isoleucine (I); Aromatic: Phenylalanine (F), Tyrosine (Y), Tryptophan (W); Sulfur-containing: Methionine (M), Cysteine (C); Basic: Arginine (R), Lysine (K), Histidine (H); Acidic: Aspartic acid (D), Glutamic acid (E), Asparagine (N), Glutamine (Q). *See also*, Creighton (1984) *Proteins*, W.H. Freeman and Company. In addition, individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids in an encoded sequence are also "conservatively modified variations."

A "subsequence" refers to a sequence of nucleic acids or amino acids that comprise a part of a longer sequence of nucleic acids or amino acids (e.g., protein) respectively.

Nucleic acids are "elongated" when additional nucleotides (or other analogous molecules) are incorporated into the nucleic acid. Most commonly, this is performed with a polymerase (e.g., a DNA polymerase), e.g., a polymerase which adds sequences at the 3' terminus of the nucleic acid.

Two nucleic acids are "recombined" when sequences from each of the two nucleic acids are combined in a progeny nucleic acid. Two sequences are "directly" recombined when both of the nucleic acids are substrates for recombination. Two sequences are "indirectly recombined" when the sequences are recombined using an intermediate such as a cross-over oligonucleotide. For indirect recombination, no more than one of the sequences is an actual substrate for recombination, and in some cases, neither sequence is a substrate for recombination.

A "specific binding affinity" between two molecules, for example, a ligand and a receptor, means a preferential binding of one molecule for another in a mixture of molecules. The binding of the molecules can be considered specific if the binding affinity is about $1 \times 10^4 \text{ M}^{-1}$ to about $1 \times 10^6 \text{ M}^{-1}$ or greater.

Substrate: a substrate is the molecule that an enzyme naturally recognizes and converts to a product in the biochemical pathway in which the enzyme naturally carries out its function, or is a modified version of the molecule, which is also recognized by the enzyme and is converted by the enzyme to a product in an enzymatic reaction similar to the naturally-occurring reaction.

Transformation: a process for introducing heterologous DNA into a cell, tissue, or insect. Transformed cells, tissues, or insects are understood to encompass not only the end product of a transformation process, but also transgenic progeny thereof.

"Transformed," "transgenic," and "recombinant" refer to a host organism such as a bacterium or a plant into which a heterologous nucleic acid molecule has been introduced. The nucleic acid molecule can be stably integrated into the genome of the host or the nucleic acid molecule can also be present as an extrachromosomal molecule. Such an extrachromosomal molecule can be auto-replicating. Transformed cells, tissues, or plants are understood to encompass not only the end product of a transformation process, but also transgenic progeny thereof. A "non-transformed," "non-transgenic," or "non-recombinant" host refers to a wild-type organism, e.g., a bacterium or plant, which does not contain the heterologous nucleic acid molecule.

The present invention relates to transgenic hosts particularly, transgenic plants, plant tissues, plant seeds, and plant cells comprising a heterologous polynucleotide encoding a gene product where the gene product comprises trichothecene resistance activity and methods of making and using the same. Trichothecene resistance activity as used herein refers to an activity that reduces or inhibits the phytotoxicity of a trichothecene, particularly to a fungus and/or plant, in a particular embodiment of the invention trichothecene resistance activity refers to an activity that transfers an acetate to the C-3 position of a trichothecene.

The present invention further relates to transgenic hosts, particularly, transgenic plants, plant tissues, plant seeds, and plant cells expressing a heterologous polynucleotide encoding a gene product, the gene product having trichothecene resistance activity, particularly an acetyl transferase gene product, more particularly a 3-O-acetyl transferase gene product, more particularly trichothecene 3-O- acetyl transferase gene product and methods of making and using the same. Expression of the heterologous polynucleotide of the invention comprises the synthesis of RNA and may be detected by northern blot analysis. Particularly, expression of the heterologous polynucleotide of the invention may be detected where a labeled probe derived from a heterologous nucleotide of the invention, in particular embodiments, from SEQ ID NOs. 1, 5 or 7, hybridizes with RNA isolated from a transgenic plant of the invention in 7% sodium dodecyl sulfate (SDS), 0.5 M Sodium phosphate pH 7.0, 1 mM EDTA, 10 mg/ml BSA at 65°C with washing in .5% BSA (fraction V), 5% SDS, 40 mM Sodium phosphate pH 7.0, 1 mM EDTA, .25 M sodium chloride at 65 ° C, preferably in 1% SDS, 40 mM Sodium phosphate pH 7.0, 1 mM EDTA, .125 M sodium chloride at 65 ° C, and preferably in 1% SDS, 40 mM Sodium phosphate pH 7.0, 1 mM EDTA at 65 ° C.

The present invention further relates to transgenic plants plant tissues, plant seeds, and plant cells, expressing a heterologous polynucleotide of the invention where the plant, plant cell, plant tissue or plant seed is trichothecene resistant. Trichothecene resistant plants, plant cells, plant tissues and plant seeds as used herein are those which are capable of metabolism in the presence of a trichothecene which may be determined as described in Example 7 below. In a particular embodiment, trichothecene resistant plants, plant tissues, plant cells and plant seeds which have a specific enzyme activity of at least 10 nmol triacetoxyscirpenol (hereinafter "TAS ")/microgram protein/15 min incubation at saturating substrate levels, more particularly at least 5 nmol TAS/microgram protein/15 min, more particularly at least 1 nmol TAS/microgram protein/15 min, more particularly at least 0.8 nmol TAS/microgram protein/15 min more particularly at least 0.5 nmol TAS/microgram protein/15 min, more particularly a specific activity of 0.25 nmol TAS/microgram protein/ 15 minute,

WO 00/60061

PCT/EP00/02769

more particularly a specific activity of 0.1 nmol TAS/microgram protein/15 min., more particularly a specific activity of 0.05 nmol TAS/microgram protein/15 min and even more particularly a specific activity of 0.01 nmol TAS/microgram protein/15 min above background levels of activity that occur naturally in a wild type control, particularly as determined in an assay as described in Example 6 below.

Trichothecene resistant plants of the invention comprise those of which a greater percentage of the seed germinate and form roots in the presence of a trichothecene than the seed from a wild type control where the trichothecene is present at a concentration of at least 5 microgram/ml, more preferably at least 10 microgram/ml, more preferably at least 15 microgram/ml, more preferably at least 20 microgram/ml and more preferably at least 25 microgram/ml. In a particularly preferred embodiment, trichothecene resistant plants of the invention comprise those of which at least 10% more seed, more preferably at least 20% more seed, more preferably at least 30% more seed, more preferably at least 40% more seed, more preferably at least 50% more seed, more preferably at least 60% more seed, more preferably at least 70% more seed, more preferably at least 80% more seed and more preferably at least 90% more seed germinate and form roots in the presence of a trichothecene than the seed of a wild type control.

Trichothecenes are frequently divided into several different structural groups. A particular embodiment of the present invention is drawn to resistance to group A and B trichothecenes. Groups A and B comprise the *Fusarium* trichothecenes and are differentiated primarily by the absence (group A) or presence (group B) of a carbonyl functional group at position C-8. The group B trichothecene DON, accordingly, comprises a carbonyl group at the C-8 position.

The present invention is more particularly drawn to resistance to trichothecenes, which contain a C-3 hydroxyl. Such trichothecenes include T-2 toxin, HT-2 toxin, isotrichodermol, DAS, 3-deacetylcalonecetrin, 3,15-dideacetylcalonecetrin, scirpentriol, neosolaniol, 15-acetyldeoxynivalenol, nivalenol, 4-acetylnivalenol (fusarenone-X), 4,15-diacetylnivalenol, 4,7,15-acetylnivalenol, and DON and their various acetylated derivatives.

In a particular embodiment, the trichothecene resistant plant, cell, tissue or seed thereof is resistant to a trichothecene producing fungus, particularly a fungus of the genus *Fusarium*. Fungus resistance as used herein refers to no initiation of infection after fungal inoculation or reduced spread of the infection after fungal inoculation compared to a wild type control.

In a preferred embodiment, a fungal resistant transgenic plant of the present invention is a cereal plant and under fungal challenge comprises less infected kernels or

WO 00/60061

PCT/EP00/02769

seeds compared to a wild type control, preferably at least a 10% decrease of infected kernels or seeds compared to the same number of kernels or seeds evaluated in a wild type control, more preferably at least a 20% decrease, more preferably at least a 40% decrease and more preferably at least a 50% decrease of infected kernels compared to the same number of kernels or seeds in a wild type control. The fungal resistant transgenic cereal plants of the invention comprise but are not limited to maize, wheat, barley, rice, and oats.

In wheat, fungal spread in the head may be evaluated as described in Example 9 below, by counting the number of symptomatic and asymptomatic spikelets on each inoculated head and calculating the percentage of spikelets on each head that are symptomatic. In a preferred embodiment, fungal resistant wheat of the present invention comprises, under fungal challenge, less infected spikelets than the wild type control, preferably at least a 10% decrease of infected spikelets compared to the same number of spikelets evaluated in a wild type control, more preferably at least a 20% decrease, more preferably at least a 40% decrease and more preferably at least a 50% decrease of infected spikelets compared to the same number of spikelets in a wild type control.

In maize, fungal spread in the ear may be evaluated by visual estimation of the percentage of infected kernels as described further in Example 9 below. In a preferred embodiment, fungal resistant maize of the invention, under fungal challenge, comprise less infected kernels than the wild type control, preferably at least a 10% decrease in infected kernels compared to the number of infected kernels in the same number of ears visibly estimated in a wild type control, more preferably at least a 20% decrease, more preferably at least 30% decrease, more preferably at least a 40 % decrease and more preferably at least a 50% decrease in infected kernels compared to the same number of ears visibly estimated in a wild type control. In maize, internal fungal spread in the stalk may be visually evaluated by splitting open the stalk and assessing the amount of discoloration. In a preferred embodiment of the invention, the transgenic maize of the invention comprises less internal and/or external discoloration of the stalk compared to a wild type control.

In another, preferred embodiment fungal resistant plants of the invention comprise those of which a greater percentage of seed germinate in the presence of fungal challenge than germinate in the wild type control. In a particularly preferred embodiment, fungal resistant plants of the invention comprise those of which at least 10% more seed, more preferably at least 20% more seed, more preferably at least 30% more seed, more preferably at least 40% more seed, more preferably at least 50% more seed, more preferably at least 60% more seed, more preferably at least 70% more seed, more preferably at least 80% more seed and more preferably at least 90% more seed, more

WO 00/60061

PCT/EP00/02769

preferably at least 100% more seed, more preferably at least 150% more seed germinates in the presence of *Fusarium* than does seed from the wild type control.

In another preferred embodiment, fungal resistant transgenic plants producing seed or kernels having less mycotoxin, e.g. trichothecene contamination, than the seed of a wild type control are provided. In a particularly preferred embodiment crop plants and more particularly cereal plants producing seed having at least 10% less trichothecene, more preferable at least 20% less trichothecene, more preferably at least 30% less trichothecene, more preferably at least 40% less trichothecene, more preferably at least 50% less trichothecene, more preferably at least 60% less trichothecene, more preferably at least 70% less trichothecene and more preferably at least 80% less trichothecene contamination than a wild type control are provided. Trichothecene contamination may be determined as described in Example 10 below.

The polynucleotides of use in the invention include heterologous polynucleotides encoding acetyl transferases, particularly those encoding acetyl transferases capable of conferring trichothecene resistance, more particularly those encoding trichothecene 3-O-acetyltransferases. In a particular embodiment, the heterologous polynucleotide of the invention may be derived from but is not limited to fungal origin, more particularly from *Fusarium*, *Trichothecium*, and *Myrothecium* origin, more particularly from a *Fusarium* species such as *F. acuminatum*, *F. crookwellense*, *F. culmorum*, *F. equiseti*, *F. graminearum* (*Gibberella zeae*), *F. lateritium*, *F. poae*, *F. sambucinum* (*G. pulicaris*), and *F. sporotrichioides*. Heterologous polynucleotides of use in the invention include SEQ ID NO:1, 5 and/or 7 and sequences substantially similar to SEQ ID NO:1, 5 and/or 7.

A polynucleotide of use in the invention can be incorporated into host cells, such as plant, fungal or bacterial cells, using conventional recombinant DNA technology. Generally, this involves inserting the polynucleotide into an expression system to which the polynucleotide is heterologous using standard cloning procedures known in the art. The vector contains the necessary elements for the transcription and translation of the polynucleotide of use in the invention in a host cell containing the vector. A large number of vector systems known in the art can be used, such as plasmids, bacteriophage viruses and other modified viruses. The components of the expression system may also be modified to increase expression. For example, truncated sequences, nucleotide substitutions, nucleotide optimization or other modifications may be employed. Expression systems known in the art can be used to transform virtually any crop plant cell under suitable conditions. A heterologous polynucleotide of the inventions is preferably stably transformed and integrated into the genome of the host cells. In another preferred embodiment, the heterologous polynucleotide

WO 00/60061

PCT/EP00/02769

of the inventions is located on a self-replicating vector. Examples of self-replicating vectors are viruses, in particular Gemini viruses. Transformed cells can be regenerated into whole plants such that the chosen form of the polynucleotide of the invention confers trichothecene resistance in the transgenic plants.

A polynucleotide of the invention intended for expression in transgenic plants is first assembled in an expression cassette behind a suitable promoter expressible in plants. The expression cassettes may also comprise any further sequences required or selected for the expression of the heterologous polynucleotide of the invention. Such sequences include, but are not restricted to, transcription terminators, extraneous sequences to enhance expression such as introns, vital sequences, and sequences intended for the targeting of the gene product to specific organelles and cell compartments. These expression cassettes can then be easily transferred to the plant transformation vectors described *infra*. The following is a description of various components of typical expression cassettes.

The selection of the promoter used in expression cassettes will determine the spatial and temporal expression pattern of the heterologous polynucleotide of the invention in the transformed plant. Selected promoters will express heterologous polynucleotides of the invention in specific cell types (such as leaf epidermal cells, mesophyll cells, root cortex cells) or in specific tissues or organs (roots, leaves or flowers, for example) and the selection will reflect the desired location of accumulation of the gene product. Alternatively, the selected promoter may drive expression of the gene under various inducing conditions. Promoters vary in their strength, i.e., ability to promote transcription. Depending upon the host cell system utilized, any one of a number of suitable promoters known in the art can be used. For example, for constitutive expression, the CaMV 35S promoter, the rice actin promoter, or the ubiquitin promoter may be used. For regulatable expression, the chemically inducible PR-1 promoter from tobacco or *Arabidopsis* may be used (see, e.g., U.S. Patent No. 5,689,044).

A variety of transcriptional terminators are available for use in expression cassettes. These are responsible for the termination of transcription beyond the heterologous polynucleotide of the invention and its correct polyadenylation. Appropriate transcriptional terminators are those that are known to function in plants and include the CaMV 35S terminator, the *tm1* terminator, the nopaline synthase terminator and the pea *rbcS* E9 terminator. These can be used in both monocotyledonous and dicotyledonous plants.

Numerous sequences have been found to enhance gene expression from within the transcriptional unit and these sequences can be used in conjunction with the polynucleotides of this invention to increase their expression in transgenic plants. For example, various intron sequences such as introns of the maize *Adhl* gene have been shown to enhance expression, particularly in monocotyledonous cells. In addition, a number of non-translated leader sequences derived from viruses are also known to enhance expression, and these are particularly effective in dicotyledonous cells.

The coding sequence of the selected gene optionally is genetically engineered by altering the coding sequence for optimal expression in the crop species of interest. Methods for modifying coding sequences to achieve optimal expression in a particular crop species are well known (see, e.g. Perlak *et al.*, *Proc. Natl. Acad. Sci. USA* 88: 3324 (1991); and Koziel *et al.*, *Bio/technol.* 11: 194 (1993); Fennoy and Bailey-Serres. *Nucl. Acids Res.* 21: 5294-5300 (1993). Methods for modifying coding sequences by taking into account codon usage in plant genes and in higher plants, green algae, and cyanobacteria are well known (see table 4 in: Murray *et al.* *Nucl. Acids Res.* 17: 477-498 (1989); Campbell and Gowri *Plant Physiol.* 92: 1-11(1990).

Various mechanisms for targeting gene products are known to exist in plants and the sequences controlling the functioning of these mechanisms have been characterized in some detail. For example, the targeting of gene products to the chloroplast is controlled by a signal sequence found at the amino terminal end of various proteins which is cleaved during chloroplast import to yield the mature protein (e.g. Comai *et al.* *J. Biol. Chem.* 263: 15104-15109 (1988)). Other gene products are localized to other organelles such as the mitochondrion and the peroxisome (e.g. Unger *et al.* *Plant Molec. Biol.* 13: 411-418 (1989)). The cDNAs encoding these products can also be manipulated to effect the targeting of heterologous products encoded by DNA sequences to these organelles. In addition, sequences have been characterized which cause the targeting of products encoded by DNA sequences to other cell compartments. Amino terminal sequences are responsible for targeting to the ER, the apoplast, and extracellular secretion from aleurone cells (Koehler & Ho, *Plant Cell* 2: 769-783 (1990)). Additionally, amino terminal sequences in conjunction with carboxy terminal sequences are responsible for vacuolar targeting of gene products (Shinshi *et al.* *Plant Molec. Biol.* 14: 357-368 (1990)). By the fusion of the appropriate

30041-231

targeting sequences described above to a heterologous polynucleotide of the invention, it is possible to direct a resulting product to any organelle or cell compartment.

Numerous transformation vectors available for plant transformation are known to those of ordinary skill in the plant transformation arts, and the polynucleotides pertinent to this invention can be used in conjunction with any such vectors. The selection of vector will depend upon the preferred transformation technique and the target species for transformation. For certain target species, different selection markers may be preferred. Selection markers used routinely in transformation include the *nptII* gene, which confers resistance to kanamycin and related antibiotics (Messing & Vierra. *Gene* 19: 259-268 (1982); Bevan et al., *Nature* 304:184-187 (1983)), the *bar* gene, which confers resistance to the herbicide phosphinothricin (White et al., *Nucl. Acids Res* 18: 1062 (1990), Spencer et al. *Theor. Appl. Genet* 79: 625-631 (1990)), the *hph* gene, which confers resistance to the antibiotic hygromycin (Blochinger & Diggelmann, *Mol Cell Biol* 4: 2929-2931), and the *dhfr* gene, which confers resistance to methotrexate (Bourouis et al., *EMBO J.* 2(7): 1099-1104 (1983)), and the EPSPS gene, which confers resistance to glyphosate (U.S. Patent Nos. 4,940,935 and 5,188,642), phosphomannose isomerase gene, *manA*, which confers a selective metabolic advantage in the presence of mannose (U.S. Pat. Ser. No. 5,767,378 and Miles & Guest, *GENE*, 32:41-48 (1984)). PAT selectable marker that confers resistance to BASTA (Sung H. Park et al., *In Vitro Cell.Dev.Biol.-Plant*, 34: 117-121 (1998)).

Many vectors are available for transformation using *Agrobacterium tumefaciens*. These typically carry at least one T-DNA border sequence and include vectors such as pBIN19 (Bevan, *Nucl. Acids Res.* (1984)). Typical vectors suitable for *Agrobacterium* transformation include the binary vectors pCIB200 and pCIB2001, as well as the binary vector pCIB10 and hygromycin selection derivatives thereof. (See, for example, U.S. Patent No. 5,639,949).

Transformation without the use of *Agrobacterium tumefaciens* circumvents the requirement for T-DNA sequences in the chosen transformation vector and consequently vectors lacking these sequences can be utilized in addition to vectors such as the ones described above which contain T-DNA sequences. Transformation techniques that do not rely on *Agrobacterium* include transformation via particle bombardment, protoplast uptake (e.g. PEG and electroporation) and microinjection. The choice of vector depends largely on

30041-231

the preferred selection for the species being transformed. Typical vectors suitable for non-*Agrobacterium* transformation include pClB3064, pSOG19, and pSOG35. (See, for example, U.S. Patent No. 5,639,949).

Once the polynucleotide of interest has been cloned into an expression system, it is transformed into a plant cell. Methods for transformation and regeneration of plants are well known in the art. For example, Ti plasmid vectors have been utilized for the delivery of foreign DNA, as well as direct DNA uptake, liposomes, electroporation, micro-injection, and microprojectiles. In addition, bacteria from the genus *Agrobacterium* can be utilized to transform plant cells.

Transformation techniques for dicotyledons are well known in the art and include *Agrobacterium*-based techniques and techniques that do not require *Agrobacterium*. Non-*Agrobacterium* techniques involve the uptake of exogenous genetic material directly by protoplasts or cells. This can be accomplished by PEG or electroporation mediated uptake, particle bombardment-mediated delivery, or microinjection. In each case the transformed cells are regenerated to whole plants using standard techniques known in the art.

Transformation of most monocotyledon species has now also become routine. Preferred techniques include direct gene transfer into protoplasts using PEG or electroporation techniques, particle bombardment into callus tissue, as well as *Agrobacterium*-mediated transformation. Target tissue may be derived from such sources as wheat cultivar UC703 or maize genotype CG000526. For example, *Agrobacterium* mediated transformation of maize may be carried out as described in U.S. patent application 09/089,111 which correspondingly published as WO 98/54961, and of barley may be carried out as described by: M. Cho, J. Wong, C. Marx, W. Jiang, P. Lemaux and B. Buchanan (1999). Overexpression of thioredoxin h leads to enhanced activity of starch debranching enzyme (pullulanase) in barley grain. PNAS 96: 14641-14646; S. Zhang, M. Cho, T. Koprek, R. Yun, P. Bregitzer and P. Lemaux (1999). Genetic transformation of commercial cultivars of oat (*Avena sativa* L.) and barley (*Hordeum vulgare* L.) using in vitro shoot meristematic cultures derived from germinated seedlings. Plant Cell Rep. 18: 959-966; P. Bregitzer, S. Harlbert and P. Lemaux (1998). Somaclonal variation in the progeny of transgenic barley. TAG 96: 421-425; M. Cho, W. Jiang and P. Lemaux (1998). Transformation of recalcitrant barley cultivars through improvement of regenerability and decreased albinism. Plant sci. 138: 229-244; P. Lemaux, M. Cho, S. Zhang, and P. Bregitzer (1998). Transgenic cereals: *Hordeum vulgare* L. – current status and future prospects. In: Vasil I, Phillips R (eds) Molecular Improvement of

WO 00/60061

PCT/EP00/02769

Cereal Crops, Kluwer Academic Publ, Dordrecht, The Netherlands, pp 255-316; S. Zhang, R. Williams-Carrier, D. Jackson, and P. Lemaux (1998). Expression of CDC2Zm and KNOTTED1 during in vitro axillary shoot meristem proliferation and adventitious shoot meristem formation in maize (*Zea mays* L.) and barley (*Hordeum vulgare* L.). *Planta* 204: 542-549; D. McElroy, J. Louwerse, S. McElroy and P. Lemaux (1997). Development of a simple transient assay for Ac/Ds activity in cells of intact barley tissue. *Plant J.* 11: 157-165; S. Tingay, D. McElroy, R. Kalla, S. Fieg, M. Wang, S. Thornton and R. Brettell (1997). *Agrobacterium tumefaciens*-mediated barley transformation. *The Plant J.* 11: 1369-1376; J. Qureshi, Z. Basri, R. Singh, R. Burton, M. Dalton, J. Kollmorgen and G. Fincher. 1988. *Agrobacterium*-mediated transformation of two varieties of barley (*Hordeum vulgare* L.) *Proc. 42nd. Conference of Australian Society for Biochemistry and Molecular Biology*, September 28-October 1, 1998, Adelaide, Australia; J. Qureshi, R. Singh, Z. Basri, R. Stewart, R. Burton, J. Kollmorgen and G. Fincher (1997). Strategies for genetic transformation of elite Australian barley varieties. *Proc. 8th. Aust. Barley Technical symp.* Gold Coast, Queensland, 7-12 September 1997. 2:8.9-11; P. Lemaux, M. Cho, J. Louwerse, R. Williams and Y. Wan (1996). Bombardment-mediated transformation methods for barley. *Bio-Rad US/EG Bull* 2007: 1-6; T. Koprek, R. Hansch, A. Nerlich, R. Mendel and J. Schulze (1996). Fertile transgenic barley of different cultivars obtained by adjustment of bombardment conditions to tissue response. *Plant Sci.* 119: 79-91; T. Hagio, T. Hirabayashi, H. Machii and H. Tomutsune (1995). Production of fertile transgenic barley (*Hordeum vulgare* L.) plants using the hygromycin-resistance marker. *Plant Cell Rep.* 14: 329-334; H. Funatsuki, H. Kuroda, M. Kihara, P. Lazzeri, E. Muller, H. Lorz and I. Kishinami (1995). Fertile transgenic barley regenerated by direct DNA transfer to protoplasts. *TAG* 91: 707-712; A. Jahne, D. Becker, R. Brettschneider and H. Lorz (1994). Regeneration of transgenic, microspore-derived, fertile barley. *TAG* 89: 525-533; Y. Wan and P. Lemaux (1994). Generation of large numbers of independently transformed fertile barley plants. *Plant Physiol.* 104: 37-48;

The polynucleotides of the invention can be utilized to confer trichothecene resistance to a wide variety of plant cells, including those of gymnosperms, monocots, and dicots. Although the heterologous polynucleotide of the invention can be inserted, e.g. transformed into any plant cell falling within these broad classes, it is particularly useful in crop plant cells, such as rice, wheat, barley, rye, corn, oats, potato, sweet potato, turnip, squash, pumpkin, zucchini, melon, soybean, and sorghum. The polynucleotides of use in the invention rendering a plant trichothecene resistant may be used in combination with other characteristics important for production and quality. The polynucleotides of the

WO 00/60061

PCT/EP00/02769

invention can be incorporated into plant lines through breeding approaches and techniques known in the art.

Where a trichothecene resistant gene allele is obtained by transformation into a crop plant or plant cell culture from which a crop plant can be regenerated, it is moved into commercial varieties using traditional breeding techniques to develop a trichothecene resistant crop without the need for genetically engineering the allele and transforming it into the plant.

The various breeding steps are characterized by well-defined human intervention such as selecting the lines to be crossed, directing pollination of the parental lines, or selecting appropriate descendant plants. Depending on the desired properties different breeding measures are taken. The relevant techniques are well known in the art and include but are not limited to hybridization, inbreeding, backcross breeding, multiline breeding, variety blend, interspecific hybridization, aneuploid techniques, etc. . Hybridization techniques also include the sterilization of plants to yield male or female sterile plants by mechanical, chemical or biochemical means. Cross pollination of a male sterile plant with pollen of a different line assures that the genome of the male sterile but female fertile plant will uniformly obtain properties of both parental lines. Thus, the transgenic seeds and plants according to the invention can be used for the breeding of improved plant lines exhibiting no or reduced fungal growth on the plant, plant tissue, or seed thereby preventing and/or reducing mycotoxin contamination of said plant, plant tissue or seed.

The trichothecene resistance engineered into the transgenic seeds and plants mentioned above are passed on by sexual reproduction or vegetative growth and can thus be maintained and propagated in descendant plants. Generally said maintenance and propagation make use of known agricultural methods developed to fit specific purposes such as tilling, sowing or harvesting. As the growing crop is vulnerable to attack and damages caused by insects or infections, measures are undertaken to control plant diseases, insects, nematodes, and other adverse conditions to improve yield. These include mechanical measures such a tillage of the soil or removal of infected plants, as well as the application of agrochemicals such as fungicides, gametocides, nematicides, growth regulants, ripening agents and insecticides.

. In seeds production germination quality and uniformity of seeds are essential product characteristics, whereas germination quality and uniformity of seeds harvested and sold by the farmer is not important. As it is difficult to keep a crop free from other crop and weed seeds, to control seedborne diseases, and to produce seed with good germination, fairly

extensive and well-defined seed production practices have been developed by seed producers, who are experienced in the art of growing, conditioning and marketing of pure seed. Thus, it is common practice for the farmer to buy certified seed meeting specific quality standards instead of using seed harvested from his own crop. Propagation material to be used as seeds is customarily treated with a protectant coating comprising herbicides, insecticides, fungicides, bactericides, nematocides, molluscicides or mixtures thereof. Customarily used protectant coatings comprise compounds such as captan, carboxin, thiram (TMTD[®]), methalaxyl (Apron[®]), and pirimiphos-methyl (Actellic[®]). If desired these compounds are formulated together with further carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation to provide protection against damage caused by bacterial, fungal or animal pests. The protectant coatings may be applied by impregnating propagation material with a liquid formulation or by coating with a combined wet or dry formulation. Other methods of application are also possible such as treatment directed at the buds or the fruit.

It is a further aspect of the present invention to provide new agricultural methods such as the methods exemplified above which are characterized by the use of transgenic plants, transgenic plant material, or transgenic seed according to the present invention.

In another embodiment, the present invention relates to a transgenic plant cell, tissue, organ, seed or plant parts obtained from the transgenic plant. Also included within the invention are transgenic descendants of the plant as well as transgenic plant cells, tissues, organs, seeds and plant parts obtained from the descendants.

In another embodiment, the heterologous polynucleotide of use in the invention, can also be used as a selectable marker in transformation procedures. In this aspect the host cell is transformed with a second heterologous polynucleotide of interest as well as a heterologous polynucleotide of the invention which encodes a gene product comprising trichothecene resistance activity, using expressions cassettes and transformation techniques exemplified above and known in the art. After transformation, the transformed cells are selected for their ability to survive when exposed to a trichothecene, particularly DAS or DON or T-2 toxin. The host cell may be a eukaryotic or prokaryotic host cell using transformation and expression systems known in the art. The host cell may be a plant cell, a fungal cell, a bacterial cell, a yeast cell, an animal cell, or an insect cell.

30041-231

In a particularly preferred embodiment of the invention, a polynucleotide which encodes a gene product comprising trichothecene resistance activity is used as a selectable marker in plant cell transformation methods. For example, plants, plant tissue, plant seeds, or plant cells expressing at least a second heterologous DNA sequence of interest can also be transformed to express a sequence encoding a polypeptide comprising a sequence substantially similar to that of SEQ ID NO:2, 6 or 8. The transformed cells are transferred to medium containing a phytotoxic trichothecene, particularly DAS and/or DON and/or T-2 toxin, in an amount sufficient to inhibit the growth or survivability of plant cells not expressing the polypeptide substantially similar to that having the amino acid sequence of SEQ ID NO:2, 6 or 8, wherein only the transformed cells will grow or will be unstunted. Concentrations of trichothecenes useful for selection of plants expressing the polypeptide substantially similar to that having the amino acid sequence of SEQ ID NO:2, 6 or 8 range from 1 µg/ml to 90 µg/ml. The method is applicable to any plant cell capable of expressing a polynucleotide comprising a nucleotide sequence substantially similar to that of SEQ ID NO: 1, 5 or 7, and can be used with any heterologous DNA sequence of interest. Expression of the second heterologous DNA sequence and the heterologous polynucleotide of the invention can be driven by the same promoter functional in plant cells, or by separate promoters.

Description of the Sequences:

- | | |
|---------------|--|
| SEQ ID NO:1 | is a cDNA sequence from <i>Fusarium sporotrichioides</i> encoding a polypeptide of the invention having trichothecene resistance activity. |
| SEQ ID NO:2 | is the polypeptide having trichothecene resistance activity encoded by SEQ ID NO:1. |
| SEQ ID NO: 3 | is a DNA primer. |
| SEQ ID NO 4: | is a DNA primer. |
| SEQ ID NO: 5 | is a DNA sequence from <i>Fusarium graminearum</i> encoding a polypeptide of the invention having trichothecene resistance activity. |
| SEQ ID NO: 6 | is the polypeptide having trichothecene resistance activity encoded by SEQ ID NO. 5. |
| SEQ ID NO. 7 | is a DNA sequence from <i>Saccharomyces cerevisiae</i> encoding a polypeptide of the invention having trichothecene resistance activity. |
| SEQ ID NO. 8 | is the polypeptide having trichothecene resistance activity encoded by SEQ ID NO. 7. |
| SEQ ID NO. 9 | is the DNA sequence of pCIB9818. |
| SEQ ID NO. 10 | is the DNA sequence of pAgroTRlr. |

WO 00/60061

PCT/EP00/02769

SEQ ID NO. 11 is the DNA sequence of pNOV1704.

Examples

The following examples further describe the materials and methods used in carrying out the invention and the subsequent results. They are offered by way of illustration, and their recitation should not be considered as a limitation of the claimed invention.

Example 1: Composition of pNOV1700, pCIB9818, pAgroTRlr and pNov 1704

1. pNOV1700:

pNOV1700 was deposited under the terms of the Budapest Treaty on March 19, 1999, with the Agricultural Research Service, Patent Culture Collection (NRRL), Northern Regional Research Center, 1815 Northern University Street, Peoria, Illinois 61604, USA and assigned accession number NRRL B-30117.

Accordingly, pNOV1700 comprises SEQ ID NO. 1 operably linked to the *Z.mays* ubiquitin promoter, including a portion of the exon and intron, and to the nopaline synthase polyadenylation signal.

2. pCIB9818

Plasmid pCIB9818 is a 6111 base pair circular plasmid having a DNA sequence according to SEQ ID NO. 9. The *Z. mays* ubiquitin promoter, base 12 to 1993 of SEQ ID NO.9, including a portion of the exon, base 896 to 1011, and the intron, base 1047 to 1993, is operably linked to the phosphate mannose isomerase selectable marker, base pair 2090 to 3193, the inverted PEPC intron #9 from base 3248 to 3355 and the termination sequence of the CaMV 35S gene, base 3357 to 3434.

3. pAgroTRlr

Plasmid pAgroTRlr is a 13,737 base pair circular binary vector having a DNA sequence according to SEQ ID NO. 10. Accordingly, pAgroTRlr comprises a selectable marker operable linked to a promoter and termination sequence and the polynucleotide

WO 00/60061

PCT/EP00/02769

region of SEQ ID NO: 1 behind and in frame with the *Arabidopsis thaliana* UBI 3 promoter (S. Norris, S. Meyer, and J. Callus, Plant Molecular Biology 22:895-906, (1993)) and in front of and in frame with the nos polyadenylation signal.

4. pNov1704

Plasmid pNOV1704 is a 12949 base pair circular binary vector having a DNA sequence according to SEQ ID NO. 11. The *Z. mays* ubiquitin promoter, base 11 to 1992 of SEQ ID NO.11(including exon 1: 895 to 1010 and intron 1: 1046 to 1992), is operably linked to the phosphate mannose isomerase selectable marker sequence, base 2089 to 3192, and the nopaline synthase termination sequence, base 3557 to 3688. pNOV1704, further comprises the *Z. mays* ubiquitin promoter, base 9218 to 11218 (including exon 1: 10110 to 10224 and intron 1: 10225 to 11218) operably linked to the trichothecene 3-O-acetyl transferase sequence of SEQ ID NO.1, at base 11234 to 12662 and the nos termination sequence 12667 to 12935.

Example 2: Wheat transformation , selection and regeneration

Transformation

Immature zygotic embryos (0.75-1.25mm) are dissected from surface sterilized wheat caryopses (10% Chlorox X 10 minutes) then plated scutellum up onto an MS based medium (Murashige and Skoog, (1962) Physiol. Plant 15:473-439) supplemented with 3% sucrose, 3mg/liter 2,4-D (dichlorophenoxyacetic acid), 150mg/l glutamine, 75mg/l asparagine and solidified with 0.7% phytagar (3MS3S medium). The embryos are incubated in the dark at 28°C for 5-10 days prior to bombardment. The optimal time for bombardment is 6-7 days post-plating. Four hours prior to bombardment, the embryos are placed on a plasmolysis medium (same medium described above but with 15% maltose added in place of the sucrose) and arranged in a 2.5cm diameter circle with scutellum facing up.

pNOV1700, described in Example 1 above, is digested with PvuII and XmnI and a 4117bp fragment comprising a polynucleotide region having a sequence according to SEQ ID NO:1 as well as the ubiquitin promoter and NOS polyadenylation signal is isolated. pCIB9818, also described in Example 1 above, is digested with AscI and the 4246 bp fragment comprising the UBI maize promoter, selectable marker and CaMV 35S termination sequence is isolated.

The isolated DNA fragments are precipitated onto 0.3 micro meter gold particles using the standard Sanford method. While continuously vortexing , 5 microgram of the

WO 00/60061

PCT/EP00/02769

Isolated fragment DNA per construct , 50ul of 2.5 M CaCl_2 and 20ul of 0.1M spermidine are added to an eppendorf tube that contains 50ul of 50% glycerol and 3mg gold. The DNA/gold mixture is given two ethanol washes. After discarding the supernatant from the last wash, ethanol is added to make a final volume of 70ul. This provides six shots per tube of gold/DNA. The target plates are shot twice so that gives a delivery of approximately 3microgram DNA (if 5microgram each of 2 constructs is used which is usually the case) and 1.0 mg gold per target plate. The rupture pressure used is 1100 psi. After bombardment, the target plates are returned to the dark overnight. After approximately 24 hours of plasmolysis, the embryos are removed to 3MS3S and returned to the dark for 3 weeks of callus initiation. No subculturing is done during this time.

Selection/Regeneration

The embryogenic tissue that develops during the 3-week initiation period is dissected away from non-embryogenic tissue and placed on a regeneration/selection medium. The basic regeneration medium is 3MS3S without the 2,4-D but with 1mg/l GA3 (Gibberellin A₃) NAA (1-Naphtheleneacetic Acid) and NAA added instead (called NG medium). 10g/l mannose and 5g/l sucrose is added (NG1M.5S). The tissue is subjected to this initial phase of regeneration and selection for 2 weeks. For most of the 2 week period the tissue is in the light room. Shoot and root development begins during this phase and after 2 weeks all tissue is taken to the next stage.

For the second phase of regeneration and selection with mannose selection, mannose is decreased to 5g/l and the sucrose increased to 20g/l (MS2S.5M). The tissue normally stays on these media for approximately 4 weeks time during which further shoot and root development occurs.

Vigorously growing plantlets with good color, and root and shoot development are removed from plates and placed in larger containers called GA7's. This is the final stage of selection and regeneration. The medium contains only 1/2MS salts and 15g/l mannose. The best indicator that a plant may be transformed is the observance of active root growth into the medium. Leaf tissue from actively growing plantlets is collected and PCR is done for either the gene of interest or selectable marker before transferring to the green house.

EXAMPLE 3: Arabidopsis transformation

The binary vector pAgroTRlr constructs described in Example 1 above is transformed into *Agrobacterium tumefaciens* strain GV3101 (Bechtold, N. *et al.*, CR Acad. Sci. Paris, Sciences de la vie, 316:1194-1199 (1993)) by electroporation (Dower, W.J., Mol. Biol. Rep 1:5 (1987)). A 25 ml culture from single colonies of GV3101 agrobacterium containing pAgroTRlr plasmids in YEB + Rifampicin 100 and Kanomycin 100 is incubated at 30 degrees overnight. Large cultures are started by inoculating 500 ml of the same media with 5 mls of the small culture and are incubated overnight at 30 degrees. The OD at 600 nm of cultures is determined and the cultures are then spun down at 5 K in the GSA rotor for 15 minutes. Cells are resuspended in "IM Modified infiltration media" to achieve a final O. D. at 600 nm of .08. 200 microliters of Silwet per liter of suspended cells is added. Three pots of bolting Arabidopsis var Columbia about 4 plants per pot, are inverted in about 500 ml of cell suspension. The flowers are shaken in the cell suspension to dislodge the air bubbles and the plants are incubated in the cell suspension for 15 minutes. A dome is placed on the tray to keep the plants humid overnight.

Plants are allowed to grow about 3-4 weeks after which the plants are not watered for up to 1 week. Seed pods are collected and dried in drying room for about a week and a half. The seeds are planted and allowed to grow for about 2 weeks. The plants are sprayed with the selection agent and then sprayed again 2 days later and 4 days later. After about three days surviving plants can be transplanted to new pots.

EXAMPLE 4: Maize biolistic transformation.

Type I embryogenic callus cultures (Green et al 1983, Somatic cell genetic systems in corn. A. Fazelahmad, K. Downey, J. Schultz, RW Voellmy, eds. Advances in Gene Technology: Molecular Genetics of Plants and Animals. Miami Winter Symposium Series, Vol. 20. Academic Press, NY.) are initiated from immature maize embryos, that are 1.5 – 2.0 mm in length, from greenhouse grown material. Embryos are aseptically excised from surface-sterilized ears approximately 14 days after pollination. The embryos are placed on D callus initiation media (Duncan et al, (1985) Planta 165:pp322-332) with 2% sucrose and 5mg/L chloramben. Embryos and embryogenic cultures are subsequently cultured in the dark. Embryogenic responses are excised from the explants after about 14 days. Responses are placed onto D callus maintenance media with 2% sucrose and 0.5mg/L 2,4-D. After about 6 weeks of weekly selective subculture to fresh maintenance media, high quality compact embryogenic cultures are established. Actively growing embryogenic callus pieces are selected as target tissue for gene delivery. The callus pieces are plated onto

WO 00/60061

PCT/EP00/02769

target plates containing maintenance medium with 12% sucrose approximately 4 hours prior to gene delivery.

The callus pieces are arranged in circles, with radii of 8 and 10mm from the center of the target plate.

pNOV1700, described in Example 1 above, is digested with PvuII and XmnI and a 4117bp fragment comprising a polynucleotide region having a sequence according to SEQ ID NO:1 isolated as well as promoter and polyadenylation signal. pCIB9818, also described in Example 1 above, is digested with AscI and the 4246 bp fragment comprising the marker gene, promoter and termination signal is isolated. The isolated DNA fragments are precipitated onto gold microcarriers as described in the DuPont Biolistics manual. Two to three μ g for each plasmid construct is used in each 6 shot microcarrier preparation. Polynucleotides of the invention are delivered to the target tissue cells using the PDS-1000He Biolistics device. The settings on the Biolistics device are as follows: 8 mm between the rupture disk and the macrocarrier, 10 mm between the macrocarrier and the stopping screen and 7 cm between the stopping screen and the target. Each target plate is shot twice using 650psi rupture disks. A 200 X 200 stainless steel mesh (McMaster-Carr, New Brunswick, NJ) is placed between the stopping screen and the target tissue. Seven days after gene delivery, target tissue pieces are transferred from the high osmotic medium to selection medium.

The target tissue is placed onto maintenance medium containing no sucrose and 1% mannose. After 3 to 5 weeks, growing callus pieces are subcultured to the maintenance medium containing no sucrose and 1.5% mannose. Embryogenic callus growing on selection media is subcultured every 2 weeks for 6 to 10 weeks until enough callus is produced to generate 10-20 plants. Tissue surviving selection from an original target tissue piece is subcultured as a single colony and designated as an independent transformation event. Colonies are transferred to a modified MS medium (Murashige and Skoog, 1962(1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant* 15: 473-497.) containing 2% sucrose and 1% mannose (MS2S+1M) with 0.25mg/L ancymidol and 0.5mg/L kinetin. After 2 weeks, regenerating colonies are then transferred to MS2S+1M without hormones. Regenerating shoots with or without roots from all colonies are transferred to Magenta boxes containing MS3S medium and small plants with roots are recovered and transferred to soil in the greenhouse.

WO 00/60061

PCT/EP00/02769

EXAMPLE 5: Analyses of transgenic plant expression

Tissue from transformed plants is analyzed for the presence of a polynucleotide comprising the sequence of SEQ ID NO:1. DNA is extracted from transformed plant and PCR analyses are performed according to standard protocols. The primers used for amplification of the gene constructs are (5'-acgaatcattcaccgaggag-3') (SEQ ID No. 3) and (5'-ctcacactctcaggcttacc-3') (SEQ ID NO. 4). A 650 nt fragment within the sequence of SEQ ID NO:1 in wheat obtained according to Example 2 above is detected.

b. Northern analysis

Transformed plants are analyzed for the presence of RNA by northern blot hybridization. For northern blot analysis, RNA extracted from plant tissue is size separated and blotted onto a nylon membrane. This membrane is subsequently hybridized with a radioactive probe, derived from the 429 nt Styl fragment of the polynucleotide according to SEQ ID NO:1 is used as the probe. RNA is detected in wheat and arabadopsis plants transformed according to examples 2 and 3 above.

EXAMPLE 6: Enzymatic assay for trichothecene 3-O-acetyltransferase activity.

a.) Extraction of plant tissue for enzyme assays: Three 1 x 1/8 in pieces of leaf (about 50 mg) from transgenic plants of the invention including those transformed and regenerated according to Examples 2-4 above are selected.

(b) Glass Bead Mill: Tissue is placed in 2 ml round bottomed tube and the cap closed. The tube is immersed in liquid nitrogen and is incubated overnight at -80 °C. Tube is shaken on saws-all 24 seconds and 0.4 ml sodium phosphate buffer is added. The tube is vortexed about 10 seconds and is placed on ice. The tube is vortexed another 5 minutes and then is spun at 14,000 rpm in Eppendorf centrifuge 5 min. The supernatant is removed and is placed in a clean tube.

The following components are mixed

trichothecene substrate, 2 microliters of DAS (20% acetone in 50 mM Sodium phosphate buffer pH 7.0). DON may also be used.

Acetyl CoA substrate, 2 micro liters of [¹⁴C]-acetyl CoA NEN cat. # NEC313 (60 mCi/millimole and 0.02 mCi/ml)

WO 00/60061

PCT/EP00/02769

Buffer, to a final volume of 50 μ l with sodium phosphate buffer pH 7.0

The assay is initiated by adding the following enzyme preparation and is incubated at 30 °C for 15 minutes.

Enzyme preparation, 10 microliter plant extract in sodium phosphate buffer pH 7.0

After 15 minutes, 100 microliters ethyl acetate is added and the tube is vortexed twice for several seconds. The tube is spun for 2 minutes at 14,000 rpm in an Eppendorf centrifuge. 50 microliters of the ethyl acetate phase is removed and is added to a vial containing scintillation cocktail. The tube is counted for 2 min. using a scintillation counter.

Twenty separate wheat plants obtained according to Example 2 above and having specific activities of 0.60 to 13.4 nmol acetylated product/ μ g protein/15 min are identified. The specific activities of the transformed wheat plants are significantly greater than the negative control. The negative control is a non-transformed wheat cultivar, which has a specific activity of 0.1 to 0.2 nmol acetylated product/ μ g protein/15 min.

Five separate Arabidopsis plants obtained according to Example 3 above and having specific activities ranging from 3.8 to 28 nmol acetylated product/ μ g protein/15 min are identified. The specific activities of the transformed plants are significantly greater than that of the negative control. The negative control is an *Arabidopsis thaliana* var columbia transformed with a nucleic acid construct for expressing the selectable marker which has a specific activity of less than 0.1 nmol acetylated product/ μ g protein/15 min.

Maize plants from at least two different transformants obtained according to Example 4 above and having specific activities ranging from 11.1 to 17.9 nmol acetylated product/ μ g protein/15 min are identified. The specific activities of the transformed plants are significantly greater than that of the negative control. The negative control is a non-transformed *maize* genotype that has a specific activity of less than 0.2 nmol acetylated product/ μ g protein/15 min.

WO 00/60061

PCT/EP00/02769

Maize plants from at least 16 different transformants obtained using *Agrobacterium* mediated transformation of pNOV 1704, having specific activities ranging from 17 to 183 nmol/microgram/15 min are identified.

EXAMPLE 7: Bioassay for trichothecene resistance in transgenic plants.

250 ml of CPR media having the following components is prepared and the pH adjusted to 6.5 with KOH.

½ MS salts 0.54 g
 ½ MS vitamins 1.25 ml
 Sucrose 1% (optional) 2.50 g

Agarose is added to the above media to a concentration of 1% (2.50 g) and the media is autoclaved. 25 ml of 50 mg/ml chlorophenol red is added to the autoclaved media.

While media is maintained at 55 °C, DAS or DON is added in acetone at various concentrations. (i. e. DON at 4, 8, or 16 microliters 10 mg/ml or DAS at 2, 4, or 6 microliters 50 mg/ml DAS per 1.7 ml). About 0.5 ml of media is aliquoted to each well in a 48 well microtiter plate.

1/3 x 1/8 inch pieces of transformed plant tissue are added to microtiter plate wells as well as control tissue from untransformed wild type controls. The leaf pieces are allowed to fall into a petri dish and are pushed into the microtiter plate well media with tweezers. The microtiter plates are incubated 2 to 4 days at 20 °C under lights. Leaf piece metabolism results in color change (drop in pH) from red to yellow. Trichothecene resistance activity or reduced sensitivity to trichothecenes by transformants, results in yellow colored wells in the presence of DAS or DON.

A color change from red to yellow compared to the control that remains red is observed in wheat and maize plants of the invention. Furthermore, the individual leaf pieces have significantly less chlorosis than the corresponding control.

EXAMPLE 8: Germination Assay

A. Trichothecene Resistance Germination Assay

30041-231

Seed from transgenic plants of the invention is grown under selective pressure from the selection agent and the resulting plants are selfed. The resulting seed is plated on MS3S medium (MS salts 4.3 g/L, MS vitamins 100X, Sucrose 30 g/L, and phytagar 8 g/L) and supplemented with either DAS or DON (at 20 mg/ml) at a density of 1000 to 1200 seeds/petri dish (100 mm diameter). After incubation in the light for four days the plates are examined for seedling growth.

Arabidopsis seed from plants obtained according to Example 3 above and grown in media comprising DAS, has numerous plants with both root and shoot development. While control seed (parental Arabidopsis line, var. Columbia) germinates poorly and no roots form when grown in DAS supplemented media. No differences are observed between transformed and control seeds grown in the same media without DAS.

B. Fungal Resistance Germination Assay for Detecting Resistance to Seedling Blight

1. Wheat Fungal Resistance Germination Assay:

Fungal resistance germination assays in wheat are carried out substantially as described by R. H. Proctor, T. M. Hohn, and S. P. McCormick. Reduced virulence of *Gibberella zeae* caused by disruption of a trichothecene toxin biosynthetic gene. *Mol. Plant-Microbe Interact.* 8 (4):593-601, 1995.).

Inoculum consists of macroconidia of *F. graminearum* diluted in water to 1×10^6 conidia per ml. Inoculum is prepared by washing the macroconidia from V-8 juice agar cultures grown under white and near UV fluorescent lights for 7-10 days. In seedling assays, seeds of two different transgenic wheat events from Example 2 above and the wild type control are surface sterilized by washing in a 10% bleach and .05% Tween solution for approximately 15 min. and rinsed five times with sterile distilled water. The seeds are soaked in a suspension of macroconidia for approximately 10 min and then sown in vermiculite contained in 10 cm plastic pots (20 seeds per pot). Prior to sowing, the pots are filled approximately $\frac{3}{4}$ full with vermiculite and set in 2-4 cm of water until the top of the vermiculite was wet. After sowing, seeds are covered with an additional 1-2 cm of vermiculite and pots are placed individually into plastic bags and incubated in a growth chamber at 22 °C with 16 h light and 8 h dark for week. After approximately one week the pots are removed from the bags, and after two weeks, disease is evaluated by counting the number of seedlings that emerge in each pot. Controls are treated as described above except that the seeds are soaked in sterile water and 40 seeds are used.

WO 00/60061

PCT/EP00/02769

50% and 43% of the seed from the two different transgenic plant events germinate as compared to the same transgenic seed treated with water, whereas, 17% of the wild type control germinate compared to the same seed treated with water.

2. Maize Fungal Resistance Germination Assay:

Inoculum is produced from *F. graminearum* cultures grown on mung bean agar (made with liquor from boiled mung beans) under 12h alternating light and dark cycles at 25°C. Spores are harvested by first flooding the plate with sterile water and then scraping the plate using a glass rod. The solution is collected and the spore concentration adjusted to 1×10^6 spores/ml with double distilled, sterile water on the day of inoculation.

Soil consisting of a sterilized mix of soil, peat, and vermiculite is inoculated with 1 ml of spore solution/liter of soil in 5 liter flats and the inoculated soil is mixed in a cement mixer for 2 minutes per load. Control treatments consist of non-infested soil.. Flats are planted with 30 kernels each of the transgenic seed of the invention or wild type control and incubated in growth chambers kept at 55°F under semi-saturated soil conditions for up to 4 weeks. Light levels are under 24-hour darkness for 14 days post planting, and 12-hour light 15-24 days post planting. Labeled stakes are added to flats, flats are moved to the growth chamber and randomized.

Plant counts are initiated as soon as emergence begins, and are performed daily or every other day until there is no longer a change in plant emergence.

Symptoms of visual discoloration and damping-off of seedling emergence are determined and used to characterize the degree of plant resistance as compared to wild type controls. Plants having less visual discoloration and/or damping off of seedling emergence than the wild type control are selected.

EXAMPLE 9: Fungal Resistance Assay

A. Testing of transgenic wheat plants

Wheat head blight or head scab is caused by *Fusarium graminearum* (teleomorph: *Gibberella zeae*). *F. graminearum* cultures are grown on V-8 agar medium (made with V-8 juice) or on mung bean agar (made with liquor from boiled mung beans) under 12h alternating light and dark cycles at 25°C. Spores are harvested by first flooding the plate with sterile water and then scraping the plate using a glass rod. The solution is collected and the spore concentration adjusted to 5×10^4 spores/ml with double distilled, sterile water on the day of inoculation.

WO 00/60061

PCT/EP00/02769

Transgenic plants are obtained as described in Example 2 above, and control plants may be grown in the greenhouse until anthesis or heading. Heads are inoculated by injecting approximately 20 ul (about 1000 spores) of inoculum between the lemma and palea of one floret near the middle of each head. Some heads are left uninoculated or are inoculated with sterile water as controls. Plants are then moved to a growth chamber and incubated under high humidity for up to 21 days or plants are first incubated in a mist chamber for 72 h at 65 to 70° F and then incubated in the greenhouse for an additional 18 days.

Transgenic plants of the invention and control plant may also be grown in the field, where heads are inoculated by spraying.

Disease is evaluated by counting the number of symptomatic and asymptomatic spikelets on each inoculated head on a representative number of transgenic heads and wild type control heads and from this calculating the percentage of spikelets on each head that are symptomatic. Symptoms consist of premature whitening as compared to the control plants and in some cases necrosis of spikelets. Plants having fewer symptomatic spikelets than the wild type control are selected.

Six different transgenic plants having different copy numbers of SEQ ID NO.1 and different numbers of insertion sites according to southern data have average percent of symptomatic spikelets per head ranging from 10.40% to 31.20% compared to 44.75% for the wild type control, where the transgenic plants and controls are incubated in the greenhouse as described above. These same transgenic plants have enzymatic activities, measured as described in example 6 above, ranging from 0.874 to 29.1 nmol/microgram/15 min.

B. Testing of transgenic corn plants

Corn ear rot assay.

Corn ear rot is caused by *Fusarium graminearum* (teleomorph: *Gibberella zeae*). *F. graminearum* cultures are grown on V-8 agar medium (made with V-8 juice) under 12h alternating light and dark cycles at 25°C. Spores are harvested by first flooding the plate with sterile water and then scraping the plate using a glass rod. The solution is collected and the spore concentration adjusted to 5×10^5 spores/ml with double distilled, sterile water) on the day of inoculation. Transgenic plants and control plants are grown in the greenhouse or field. Where grown in the greenhouse, the transgenic and control plants are maintained in the greenhouse until four to seven days post silk emergence when a 2 ml spore suspension is introduced into the silk channel (inside the husk cavity and above the cob). This is accomplished using an 18-gauge stainless steel hypodermic needle attached to a large

WO 00/60061

PCT/EP00/02769

syringe. In addition to silk channel inoculations, a kernel inoculation method is also used to assay disease resistance. Kernel inoculation involves the introduction of the spore suspension (approx. 0.4 ml) into a group of four kernels through multiple injections with an 18-gauge needle attached to a syringe. Disease is evaluated by visual inspection of ears harvested 5 to 7 weeks post-inoculation for visibly infected kernels. The disease rating scale for husked ears is based on a visual estimation of the percentage of visibly infected kernels on an ear as follows: 1 is 0%; 2 is 1 to 3%; 3 is 4 to 10%; 4 is 11 to 25%; 5 is 26 to 50%; 6 is 51 to 75%; 7 is 76 to 100%. Maize plants are selected that have a lower percentage of visibly infected kernels compared to the wild type control.

EXAMPLE 10: Mycotoxin Contamination Assay

Samples are prepared for mycotoxin concentration analysis as follows. Seed is collected from transgenic plants of the invention weighed and bulked together. Where wheat seed is being assayed, wheat seed is collected from the heads of the same transgenic plants of the invention and weighed and bulked together. Where transgenic maize is being assayed, corn ears are dried to low moisture levels, ears are hand-shelled and kernels from ears of the same transgenic plant are weighed and bulked together. Each seed or kernel sample is mixed thoroughly to promote a random distribution of seed. A 50 g seed or kernel sample is ground to a fine powder in a mill (e.g. Retsch ultra centrifugal mill type ZM1, Brinkman Instruments, Inc., Rexdale, Ontario, Romer Series II Mill, Union, Missouri, USA). The concentration of the mycotoxin of interest such as, DON is then determined using the commercially available tests such as DONtest TAG™ mycotoxin testing system (VICAM, LP, 313 Pleasant Street, Watertown, MA 02472) or analyzed by a commercial analysis company (e.g. Romer Labs, Inc, Union, Missouri, USA or Trilogy Analytical Laboratory, Inc., Beaufort, Missouri, USA). The manufacturer's instructions are followed for all aspects of the analysis. For DONtest TAG™ mycotoxin testing system, a final fluorometric measurement for DON is conducted. Plants producing seed or kernels having less mycotoxin, such as DON, than the wild type control are selected.

Example 11: Use of polynucleotide according to SEQ ID NO:1 as a selectable marker.

A. Selectable Marker in Fungal Cells.

Ashbya gossypi is transformed using standard fungal transformation techniques with a DNA construct comprising a polynucleotide having the sequence of SEQ ID NO:1 operably linked to the galactosidase promoter. Transformed cells grow in media comprising DAS at a

WO 00/60061

PCT/EP00/02769

concentration ranging from 1.56ng/ml to 196 pg/ml whereas as the untransformed wild type fungal cells do not.

B. Selectable Marker in Plant Cells.

Seed from Arabidopsis plants transformed according to Example 3 above but not yet subjected to selection is plated out in .1% agarose medium containing 0, 5, or 10 µg /ml DAS. After incubation in a growth room at 22 C with 16 hours of light and 8 hours of darkness for 2 weeks, the larger unstunted plants are transplanted from a DAS plate, and a corresponding number are transplanted from the control plate.

Leaves of Arabidopsis plants transplanted from the 5 microgram/ml plate, are assayed for enzymatic activity after a 2 week growth period, and showed 11 out of 11 unstunted plants were enzymatically active as measured by Example 6 while 9 out of 10 plants not selected by DAS were negative in the same assay. The one non-selected plant that was enzymatically active was much less active than any of the DAS selected plants assayed.

The above-disclosed embodiments are illustrative. This disclosure of the invention will place one skilled in the art in possession of many variations of the invention. All such obvious and foreseeable variations are intended to be encompassed by the appended claims.

WO 00/60061

PCT/EP00/02769

SEQUENCE LISTING

<110> Hohn, T.
 Salmeron, J.
 Peters, C.
 Kendra, D.
 Reinders, J.
 Kuznia, R.
 Dill-Mackey, R.

<120> Transgenic Plant and Methods

<130> sequencelist

<140>

<141>

<160> 11

<170> PatentIn Ver. 2.0

<210> 1

<211> 1403

<212> DNA

<213> Fusarium sporotrichioides

<400> 1

```

atcaaaatgg ccgcaacaag cagcacaagc agccagtctt ttgacataga gctcgacatc 60
atcggccagc aaccgcctct tctttcaatc tacacccaga tcagtctcgt ttaccccgtc 120
tctgatccct cccagtatcc caccatcgtc agcacccttg aggaaggcct aaaacgcctc 180
tctcaaacct tcccatgggt cgcgggccag gtcaagaccg agggcatcag cgaaggaaac 240
acaggaactt ccaagatcat tccatatgag gagacacccc gtcttgtggt gaaagacctc 300
cgtgatgatt cctcagcgcc aacgatcgag gggttgagaa aggcgggttt ccccttagag 360
atgtttgacg agaacgctcg cgctccgagg aagacattag ctatcggacc tggcaatggc 420
cccaacgacc cgaagcctgt gttgctattg cagctcaact tcattaaggg cggactcatt 480
ctcacggtca acggacaaca tgggtgctatg gacatgacag gacaagatgc aattattcgt 540
cttctctcca aggcgtgccg caacgaatca ttcaccgagg aggaaatctc ggccatgaac 600
ctcgatcgca agacggtagt ccctctcctt gaaaactaca aagttgggtc tgagctagac 660
caccagatcg ccaaacctgc gcctgctggc gacgctccac ccgcaccggc caaggcaagc 720
tgggcgttct tttcattcac tcccaaggcc ctctcggagc tgaaagacgc agccacaaag 780
actcttgacg cgtcgtccaa gtttgtgtca actgatgatg ctctttcggc gtttatctgg 840
caatcaacct cgcgcgtagc tctcgcaaga ttggatgctt ccacacctac tgaattctgc 900
cgcgctgtcg acatgcgggg cccaatgggc gtatcaagca catacccagg ctttcttcaa 960
aacatgacct accatgactc gaccgtcgcc gaaatcgcca acgaaccact tggcgcaaca 1020
gcatcacgcc tgcgctcgga actcaacagt gatcgtttgc gcagacgaac acaagctttg 1080
gcgacgtaca tgcatggcct gcctgacaag tcgagcgtct ccctgaccgc cgatgcgaat 1140
ccgtcaagca gcatcatgct gagttcctgg gccaaagggtg gatgctggga gtatgacttt 1200
gggtttggac tgggtaagcc tgagagtgtg agaagacctc gctttgaacc ttttgagagt 1260
ttgatgtact ttatgcccac gaagcctgat ggggagttta cggcgtccat ttctctgagg 1320
gatgaggata tggagagact aaaggcggat gaggagtgga caaagtacgc aaagtatatt 1380
gggtagatag ttactagac tac                                     1403

```

<210> 2

<211> 459

<212> PRT

<213> Fusarium sporotrichioides

<400> 2

```

Met Ala Ala Thr Ser Ser Thr Ser Ser Gln Ser Phe Asp Ile Glu Leu
  1                      5                      10                      15

```

WO 00/60061

PCT/EP00/02769

Asp Ile Ile Gly Gln Gln Pro Pro Leu Leu Ser Ile Tyr Thr Gln Ile
 20 25 30
 Ser Leu Val Tyr Pro Val Ser Asp Pro Ser Gln Tyr Pro Thr Ile Val
 35 40 45
 Ser Thr Leu Glu Glu Gly Leu Lys Arg Leu Ser Gln Thr Phe Pro Trp
 50 55 60
 Val Ala Gly Gln Val Lys Thr Glu Gly Ile Ser Glu Gly Asn Thr Gly
 65 70 75 80
 Thr Ser Lys Ile Ile Pro Tyr Glu Glu Thr Pro Arg Leu Val Val Lys
 85 90 95
 Asp Leu Arg Asp Asp Ser Ser Ala Pro Thr Ile Glu Gly Leu Arg Lys
 100 105 110
 Ala Gly Phe Pro Leu Glu Met Phe Asp Glu Asn Val Val Ala Pro Arg
 115 120 125
 Lys Thr Leu Ala Ile Gly Pro Gly Asn Gly Pro Asn Asp Pro Lys Pro
 130 135 140
 Val Leu Leu Leu Gln Leu Asn Phe Ile Lys Gly Gly Leu Ile Leu Thr
 145 150 155 160
 Val Asn Gly Gln His Gly Ala Met Asp Met Thr Gly Gln Asp Ala Ile
 165 170 175
 Ile Arg Leu Leu Ser Lys Ala Cys Arg Asn Glu Ser Phe Thr Glu Glu
 180 185 190
 Glu Ile Ser Ala Met Asn Leu Asp Arg Lys Thr Val Val Pro Leu Leu
 195 200 205
 Glu Asn Tyr Lys Val Gly Pro Glu Leu Asp His Gln Ile Ala Lys Pro
 210 215 220
 Ala Pro Ala Gly Asp Ala Pro Pro Ala Pro Ala Lys Ala Ser Trp Ala
 225 230 235 240
 Phe Phe Ser Phe Thr Pro Lys Ala Leu Ser Glu Leu Lys Asp Ala Ala
 245 250 255
 Thr Lys Thr Leu Asp Ala Ser Ser Lys Phe Val Ser Thr Asp Asp Ala
 260 265 270
 Leu Ser Ala Phe Ile Trp Gln Ser Thr Ser Arg Val Arg Leu Ala Arg
 275 280 285
 Leu Asp Ala Ser Thr Pro Thr Glu Phe Cys Arg Ala Val Asp Met Arg
 290 295 300
 Gly Pro Met Gly Val Ser Ser Thr Tyr Pro Gly Leu Leu Gln Asn Met
 305 310 315 320
 Thr Tyr His Asp Ser Thr Val Ala Glu Ile Ala Asn Glu Pro Leu Gly
 325 330 335
 Ala Thr Ala Ser Arg Leu Arg Ser Glu Leu Asn Ser Asp Arg Leu Arg

WO 00/60061

PCT/EP00/02769

340	345	350
Arg Arg Thr Gln Ala Leu Ala Thr Tyr Met His Gly Leu Pro Asp Lys		
355	360	365
Ser Ser Val Ser Leu Thr Ala Asp Ala Asn Pro Ser Ser Ser Ile Met		
370	375	380
Leu Ser Ser Trp Ala Lys Val Gly Cys Trp Glu Tyr Asp Phe Gly Phe		
385	390	395
Gly Leu Gly Lys Pro Glu Ser Val Arg Arg Pro Arg Phe Glu Pro Phe		
405	410	415
Glu Ser Leu Met Tyr Phe Met Pro Lys Lys Pro Asp Gly Glu Phe Thr		
420	425	430
Ala Ser Ile Ser Leu Arg Asp Glu Asp Met Glu Arg Leu Lys Ala Asp		
435	440	445
Glu Glu Trp Thr Lys Tyr Ala Lys Tyr Ile Gly		
450	455	

<210> 3
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: DNA Primer

<400> 3
 acgaatcatt caccgaggag 20

<210> 4
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: DNA Primer

<400> 4
 ctcacactct caggcttacc 20

<210> 5
 <211> 1356
 <212> DNA
 <213> Fusarium graminearum

<400> 5
 atggctttca agatacagct cgacaccctc ggccagctac caggcctcct ttcgatctac 60
 acccaaata gttcctctta ccccgctctc gattcctctc aatatccac tattgtcagc 120
 accttcgagc aaggtcttaa gcgcttctcc gaagccgtcc catgggtcgc aggccaggtc 180
 aaagccgagg gcattagcga gggaaacaca ggaacttctt ttatcgtccc ttttgaggac 240
 gttcctcgtg ttgtagtga agacctccgc gatgacctt cagcgccac gatcgagggt 300
 atgagaaagg cgggataccc tatggcgatg tttgacgaga acatcatcgc gccaaggaag 360
 acgttaccta ttggacctgg tactgggtccc gacgacccaa agcctgtaat tctattgcag 420
 ctcaacttca tcaagggcgg actcatcctc actgtcaacg gacagcacgg tgctatggat 480
 atggtaggcc aagatgcggt gatccgtcta ctctccaagg cgtgccgtaa cgaccattc 540

WO 00/60061

PCT/EP00/02769

```

accgaagagg aaatgacggc catgaacctc gatcgcaaga cgatagttcc ttaccttgaa 600
aactatacga ttggccccga ggtagatcat cagattgtca aagctgatgt agctgggtgg 660
gacgctgttc tcacgccggt cagtgcgaagc tgggcgttct tcacattcag cccaaggcc 720
atgtcagagc tcaaggatgc tgctaccaag actcttgacg catcaacaaa gttcgtgtcg 780
actgacgatg ctctttcggc gttcatctgg aaatcggcct ctgcgtgcg tctcgaaaga 840
atcgatggct ctgcacctac cgagttctgc cgtgctgttg atgctcgacc ggcaatgggt 900
gtctcgaaca actaccagc ctttcttcaa aacatgacct accacaactc gaccatcggc 960
gaaatcgcca acgagtcact cggcgcaaca gcatcacgcc ttcgttcaga actcgacccc 1020
gcgagcatgc gccagcgaac aagaggtctc gcgacgtacc tgcacaacaa ccccgacaag 1080
tccaacgtat ccctgacggc tgatgcggac ccatctacca gcgtcatgct gagttcttgg 1140
gccaaggtgg gactctggga ttacgacttt gggctcggac tgggtaagcc cgagactgtg 1200
agacggccaa tctttgagcc tggtgagagc ttgatgtact ttatgcccaa gaagcctgat 1260
ggcgagttct gtgcggcgct ttctctgagg gatgaggata tggaccgatt gaaggcggat 1320
aaggagtgga ccaagtatgc gcagtacgtt ggtag 1356

```

<210> 6

<211> 451

<212> PRT

<213> *Fusarium graminearum*

<400> 6

```

Met Ala Phe Lys Ile Gln Leu Asp Thr Leu Gly Gln Leu Pro Gly Leu
  1                      5                      10                      15

Leu Ser Ile Tyr Thr Gln Ile Ser Leu Leu Tyr Pro Val Ser Asp Ser
                20                      25                      30

Ser Gln Tyr Pro Thr Ile Val Ser Thr Phe Glu Gln Gly Leu Lys Arg
                35                      40                      45

Phe Ser Glu Ala Val Pro Trp Val Ala Gly Gln Val Lys Ala Glu Gly
                50                      55                      60

Ile Ser Glu Gly Asn Thr Gly Thr Ser Phe Ile Val Pro Phe Glu Asp
  65                      70                      75                      80

Val Pro Arg Val Val Val Lys Asp Leu Arg Asp Asp Pro Ser Ala Pro
                85                      90                      95

Thr Ile Glu Gly Met Arg Lys Ala Gly Tyr Pro Met Ala Met Phe Asp
                100                      105                      110

Glu Asn Ile Ile Ala Pro Arg Lys Thr Leu Pro Ile Gly Pro Gly Thr
                115                      120                      125

Gly Pro Asp Asp Pro Lys Pro Val Ile Leu Leu Gln Leu Asn Phe Ile
                130                      135                      140

Lys Gly Gly Leu Ile Leu Thr Val Asn Gly Gln His Gly Ala Met Asp
  145                      150                      155                      160

Met Val Gly Gln Asp Ala Val Ile Arg Leu Leu Ser Lys Ala Cys Arg
                165                      170                      175

Asn Asp Pro Phe Thr Glu Glu Glu Met Thr Ala Met Asn Leu Asp Arg
                180                      185                      190

Lys Thr Ile Val Pro Tyr Leu Glu Asn Tyr Thr Ile Gly Pro Glu Val
                195                      200                      205

Asp His Gln Ile Val Lys Ala Asp Val Ala Gly Gly Asp Ala Val Leu

```

WO 00/60061

PCT/EP00/02769

210	215	220
Thr Pro Val Ser Ala Ser Trp Ala Phe Phe Thr Phe Ser Pro Lys Ala 225 230 235 240		
Met Ser Glu Leu Lys Asp Ala Ala Thr Lys Thr Leu Asp Ala Ser Thr 245 250 255		
Lys Phe Val Ser Thr Asp Asp Ala Leu Ser Ala Phe Ile Trp Lys Ser 260 265 270		
Ala Ser Arg Val Arg Leu Glu Arg Ile Asp Gly Ser Ala Pro Thr Glu 275 280 285		
Phe Cys Arg Ala Val Asp Ala Arg Pro Ala Met Gly Val Ser Asn Asn 290 295 300		
Tyr Pro Gly Leu Leu Gln Asn Met Thr Tyr His Asn Ser Thr Ile Gly 305 310 315 320		
Glu Ile Ala Asn Glu Ser Leu Gly Ala Thr Ala Ser Arg Leu Arg Ser 325 330 335		
Glu Leu Asp Pro Ala Ser Met Arg Gln Arg Thr Arg Gly Leu Ala Thr 340 345 350		
Tyr Leu His Asn Asn Pro Asp Lys Ser Asn Val Ser Leu Thr Ala Asp 355 360 365		
Ala Asp Pro Ser Thr Ser Val Met Leu Ser Ser Trp Ala Lys Val Gly 370 375 380		
Leu Trp Asp Tyr Asp Phe Gly Leu Gly Leu Gly Lys Pro Glu Thr Val 385 390 395 400		
Arg Arg Pro Ile Phe Glu Pro Val Glu Ser Leu Met Tyr Phe Met Pro 405 410 415		
Lys Lys Pro Asp Gly Glu Phe Cys Ala Ala Leu Ser Leu Arg Asp Glu 420 425 430		
Asp Met Asp Arg Leu Lys Ala Asp Lys Glu Trp Thr Lys Tyr Ala Gln 435 440 445		
Tyr Val Gly 450		

<210> 7

<211> 1425

<212> DNA

<213> *Saccharomyces cerevisiae*

<400> 7

```

atgttttagag tcaagatcat ctctcagaaa cgtacaaaaa gtgtacagat gctagaaaac 60
gatcaacttg atattttggg acaacaacct tcgctataca aactatacac tcaaatatgc 120
tctatctacc gtgtaccaga tccttctgct catgaccata tcgtaaatac cttaacaaga 180
ggacttgaaa cattggctaa aaatttccag tggctagcag gaaatgtcgt aaatgaaggt 240
gctgacgaag gtaacactgg tacctacaga attgtcccg tccacttatc 300
gtccaagatc ttcgagaaga tctgtctgcc ccaacaatgg attcgcttga aaaagctgac 360
tttcctatct acatgtaga cgaaaagact tttgcgcctt gcatgactat caatccacct 420

```

WO 00/60061

PCT/EP00/02769

```

ggaaacacta taggtatggc cgccaagagt gggcctgtat ttgcagttca agcaaacttt 480
atctccggcg gcctcgtctt aactattgtc gggcagcaca atattatgga tataacagga 540
caggaaagta tcatcaactt gctcaataaa tcttgccacc aaaaaccttt ctctgatgaa 600
gaactgctca ttggaaatat agataaaaagc aaatctattc ctttgtttga tgaaacttgg 660
gaacccgaca ccacgctagt tcatgaaata gtggaaacct ctagaaatac aagtggagag 720
gaaaaggaac agtcttggtc ttcgaactct acttgggctt atgttgaatt ttctgctatc 780
tcattgcaga atctgaggat tttggcaatg cagacatgta cttctggcac aaaatttgtc 840
tccactgatg atatcgtcac tgctttcatc tggaaatcag tttctcgagc ccgtttatct 900
cgacttaaac cagaaacgaa atcaaattta gggcgtgctg tggatgtag aaaacggcta 960
ggactccccg aaacgtatcc agggttatta gtcaacatga cttttaatac aggttccctg 1020
aaaagcttgg atcataaaag tttgggcggt cttgcatcac agattcgcag gaagctagac 1080
cctaaagtct tcgatttggc ctataataca tgcgcacttg ctacgctcct tagccgatgc 1140
ccggacaaga ctaagggttc tatacctcaa ccaattgata ctttatctgg aattatggtc 1200
agttcgtggg caaaagtcag cctgtatgac gttgatttca atctagggct tgggaagccc 1260
aagagtgtac gacggccgcg cttcatttcc cttgagagcc taatatattt tatgcctaga 1320
tcctccagag gtgaaatggg ggttgctctt tgccttagag ataaagattg ggagtgcctg 1380
aatgcggata aagaatggac aaattatgct acacatatag gatga 1425

```

<210> 8

<211> 474

<212> PRT

<213> *Saccharomyces cerevisiae*

<400> 8

```

Met Phe Arg Val Lys Ile Ile Ser Gln Lys Arg Thr Lys Ser Val Gln
  1                      5                      10                      15

Met Leu Glu Asn Asp Gln Leu Asp Ile Leu Gly Gln Gln Pro Ser Leu
          20                      25                      30

Tyr Lys Leu Tyr Thr Gln Ile Cys Ser Ile Tyr Arg Val Pro Asp Pro
          35                      40                      45

Ser Ala His Asp His Ile Val Asn Thr Leu Thr Arg Gly Leu Glu Thr
          50                      55                      60

Leu Ala Lys Asn Phe Gln Trp Leu Ala Gly Asn Val Val Asn Glu Gly
          65                      70                      75                      80

Ala Asp Glu Gly Asn Thr Gly Thr Tyr Arg Ile Val Pro Ser Asp Lys
          85                      90                      95

Ile Pro Leu Ile Val Gln Asp Leu Arg Glu Asp Leu Ser Ala Pro Thr
          100                     105                     110

Met Asp Ser Leu Glu Lys Ala Asp Phe Pro Ile Tyr Met Leu Asp Glu
          115                     120                     125

Lys Thr Phe Ala Pro Cys Met Thr Ile Asn Pro Pro Gly Asn Thr Ile
          130                     135                     140

Gly Met Ala Ala Lys Ser Gly Pro Val Phe Ala Val Gln Ala Asn Phe
          145                     150                     155                     160

Ile Ser Gly Gly Leu Val Leu Thr Ile Val Gly Gln His Asn Ile Met
          165                     170                     175

Asp Ile Thr Gly Gln Glu Ser Ile Ile Asn Leu Leu Asn Lys Ser Cys
          180                     185                     190

His Gln Lys Pro Phe Ser Asp Glu Glu Leu Leu Ile Gly Asn Ile Asp

```

WO 00/60061

PCT/EP00/02769

195					200					205					
Lys	Ser	Lys	Ser	Ile	Pro	Leu	Phe	Asp	Glu	Thr	Trp	Glu	Pro	Asp	Thr
	210					215					220				
Thr	Leu	Val	His	Glu	Ile	Val	Glu	Thr	Ser	Arg	Asn	Thr	Ser	Gly	Glu
225					230					235					240
Glu	Lys	Glu	Gln	Ser	Cys	Ser	Ser	Asn	Ser	Thr	Trp	Ala	Tyr	Val	Glu
				245					250					255	
Phe	Ser	Ala	Ile	Ser	Leu	Gln	Asn	Leu	Arg	Ile	Leu	Ala	Met	Gln	Thr
			260					265					270		
Cys	Thr	Ser	Gly	Thr	Lys	Phe	Val	Ser	Thr	Asp	Asp	Ile	Val	Thr	Ala
		275					280					285			
Phe	Ile	Trp	Lys	Ser	Val	Ser	Arg	Ala	Arg	Leu	Ser	Arg	Leu	Lys	Pro
	290					295					300				
Glu	Thr	Lys	Ser	Asn	Leu	Gly	Arg	Ala	Val	Asp	Val	Arg	Lys	Arg	Leu
305					310					315					320
Gly	Leu	Pro	Glu	Thr	Tyr	Pro	Gly	Leu	Leu	Val	Asn	Met	Thr	Phe	Asn
				325					330					335	
Thr	Gly	Ser	Leu	Lys	Ser	Leu	Asp	His	Lys	Ser	Leu	Gly	Val	Leu	Ala
			340					345					350		
Ser	Gln	Ile	Arg	Arg	Lys	Leu	Asp	Pro	Lys	Val	Phe	Asp	Leu	Ala	Tyr
		355					360					365			
Asn	Thr	Cys	Ala	Leu	Ala	Thr	Leu	Leu	Ser	Arg	Cys	Pro	Asp	Lys	Thr
	370					375					380				
Lys	Val	Ser	Ile	Pro	Gln	Pro	Ile	Asp	Thr	Leu	Ser	Gly	Ile	Met	Val
385					390					395					400
Ser	Ser	Trp	Ala	Lys	Val	Ser	Leu	Tyr	Asp	Val	Asp	Phe	Asn	Leu	Gly
				405					410					415	
Leu	Gly	Lys	Pro	Lys	Ser	Val	Arg	Arg	Pro	Arg	Phe	Ile	Ser	Leu	Glu
			420					425					430		
Ser	Leu	Ile	Tyr	Phe	Met	Pro	Arg	Ser	Ser	Arg	Gly	Glu	Met	Val	Val
		435					440					445			
Ala	Leu	Cys	Leu	Arg	Asp	Lys	Asp	Trp	Glu	Cys	Leu	Asn	Ala	Asp	Lys
	450					455					460				
Glu	Trp	Thr	Asn	Tyr	Ala	Thr	His	Ile	Gly						
465					470										

<210> 9

<211> 6111

<212> DNA

<213> Plasmid

<400> 9

aagcttgcat gcctgcagtg cagcgtgacc cggtcgtgcc cctctctaga gataatgagc 60

```

attgcatgtc taagttataa aaaattacca catatttttt ttgtcacact tgtttgaagt 120
gcagtttata tatctttata catatattta aactttactc tacgaataat ataacttata 180
gtactacaat aatatcagtg ttttagagaa tcatataaat gaacagttag acatggtcta 240
aaggacaatt gagtatTTTtg acaacaggac tctacagttt tatcttttta gtgtgcatgt 300
gttctccttt ttttttgcaa atagcttcac ctatataata cttcatccat tttattagta 360
catccattta gggtttaggg ttaatggttt ttatagacta attttttttag tacatctatt 420
ttattctatt ttagcctcta aattaagaaa actaaaactc tatttttagtt tttttattta 480
ataatttaga tataaaatag aataaaataa agtgactaaa aattaaacaa atacccttta 540
agaaattaaa aaaactaagg aaacattttt cttgtttcga gtagataatg ccagcctgtt 600
aaacgccgtc gacgagtcta acggacacca accagcgaac cagcagcgtc gcgtcggggc 660
aagcgaagca gacggcacgg catctctgtc gctgcctctg gacccctctc gagagtccg 720
ctccaccgtt ggacttgctc cgctgtcggc atccagaaat tgcgtggcgg agcggcagac 780
gtgagccggc acggcaggcg gcctcctcct cctctcacgg caccggcagc tacgggggat 840
tcctttccca ccgctccttc gctttccctt cctcgcccg cgtataaat agacaccccc 900
tccacaccct ctttcccaa cctcgtgttg ttcggagcgc acacacacac aaccagatct 960
cccccaaatc caccgtcgg cactccgct tcaaggtag ccgctcgtcc tcccccccc 1020
cccctctcta ccttctctag atcggcgttc cgggtccatgg ttagggcccg gtagttctac 1080
ttctgttcat gtttgtgta gatccgtgtt tgtgttagat ccgtgctgct agcgttcgta 1140
cacggatgcg acctgtacgt cagacacgtt ctgattgcta acttgccagt gtttctcttt 1200
ggggaatcct gggatggctc tagccgttcc gcagacggga tcgatttcat gatttttttt 1260
gtttcgttgc atagggtttg gtttgccctt ttcctttatt tcaatatatg ccgtgcactt 1320
gtttgtcggg tcatcttttc atgctttttt ttgtcttggg tgtgatgatg tggctctggt 1380
gggcggtcgt tctagatcgg agtagaatc tgtttcaaac tacctgggtg atttattaat 1440
tttggtatctg tatgtgtgtg ccatacatat tcatagttag gaattgaaga tgatggatgg 1500
aaatatcgat ctaggatagg tatacatgtt gatgcgggtt ttactgatgc atatacagag 1560
atgctttttg ttcgcttggg tgtgatgatg tgggtgtggt gggcggtcgt tcattcgttc 1620
tagatcggag tagaatactg tttcaaaacta cctggtgtat ttattaattt tggaaactgta 1680
tgtgtgtgtc atacatcttc atagttacga gtttaagatg gatggaaata tcgatctagg 1740
ataggtatac atgttgatgt gggttttact gatgcatata catgatggca tatgcagcat 1800
ctattcatat gctctaacct tgagtaccta tctattataa taaacaagta tgttttataa 1860
ttattttgat cttgatatac ttggatgatg gcatatgcag cagctatatg tggatttttt 1920
tagccctgcc ttcatacgtt atttatttgc ttggtactgt ttcttttgtc gatgctcacc 1980
ctgttggttg gtgttacttc tgcagggatc cccgatcatg caaaaactca ttaactcagt 2040
gcaaaactat gcctggggca gcaaaacggc gttgactgaa ctttatggta tggaaaatcc 2100
gtccagccag ccgatggccg agctgtggat gggcgcacat ccgaaaagca gttcacgagt 2160
gcagaatgcc gccggagata tcgtttcact gcgtgatgtg attgagagtg ataaatcgac 2220
tctgctcggg gaggccgttg ccaaacgctt tggcgaactg cctttcctgt tcaaagtatt 2280
atgcgcagca cagccactct ccattcaggt tcatccaaac aaacacaatt ctgaaatcgg 2340
ttttgccaaa gaaaatgccg caggtatccc gatggatgcc gccgagcgta actataaaga 2400
tcctaaccac aagccggagc tggtttttgc gctgacgcct ttccttgcca tgaacgcgtt 2460
tcgtgaattt tccgagattg tctcctact ccagccggtc gcaggtgcac atccggcgat 2520
tgctcacttt ttacaacagc ctgatgccga acgtttaagc gaactgttcg ccagcctgtt 2580
gaatatgcag ggtgaagaaa aatcccgcgc gctggcgatt ttaaaatcgg ccctcgatag 2640
ccagcagggg gaaccgtggc aaacgattcg ttttaatttct gaattttacc cggaagacag 2700
cggctctgtt tccccgctat tgctgaatgt ggtgaaattg aaccctggcg aagcgatgtt 2760
cctgttcgct gaaacaccgc acgcttacct gcaaggcgtg gcgctggaag tgatggcaaa 2820
ctccgataac gtgctgcgtg cgggtctgac gcctaaatac attgatattc cggaactggg 2880
tgccaatgtg aaattcgaag ccaaaccggc taaccagttg ttgaccagc cggtgaaaca 2940
aggtgcagaa ctggacttcc cgattccagt ggatgatttt gccttctcgc tgcattgacct 3000
tagtgataaa gaaaccacca ttagccagca gagtgccgcc attttgttct gcgtcgaagg 3060
cgatgcaacg ttgtggaaag gttctcagca gttacagctt aaaccgggtg aatcagcgtt 3120
tattgccgcc aacgaatcac cggtgactgt caaaggccac ggccgtttag cgctgtttta 3180
caacaagctg taagagctta ctgaaaaaat taacatctct tgctaagctg ggagctctag 3240
atctgttctg caciaagtgg agtagtcagt catcgatcag gaaccagaca ccagactttt 3300
attcatacag tgaagtgaag tgaagtgcag tgcagtgagt tgctggtttt tgtacaactt 3360
agtatgtatt tgtatttgta aaatacttct atcaataaaa tttctaattc ctaaaaccaa 3420
aatccagtgg gtaccgaatt cactggccgt cgtttttaca cgctcgtgact gggaaaaccc 3480
tggcgttacc caacttaatc gccttgacgc acatccccct ttcgccagct ggcgtaatag 3540
cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg gcgaatggcg 3600
cctgatgcgg tattttctcc ttacgcatct gtgcgggtatt tcacaccgca tatggtgcac 3660
tctcagtaca atctgctctg atgccgcata gttaagccag ccccgacacc cgccaacacc 3720

```

WO 00/60061

PCT/EP00/02769

```

cgctgacgcg ccctgacggg cttgtctgct cccggcatcc gcttacagac aagctgtgac 3780
cgtctccggg agctgcatgt gtcagagggt ttcaccgtca tcaccgaaac gcgcgagacg 3840
aaagggcctc gtgatacgcc tatttttata ggttaatgtc atgataataa tggtttctta 3900
gacgtcaggt ggcacttttc ggggaaatgt gcgcggaacc cctatttgtt tatttttcta 3960
aatacattca aatatgtatc cgctcatgag acaataaccc tgataaatgc ttcaatggcg 4020
cgccgcggcc gcttaagaat attgaaaaag gaagagtatg agtattcaac atttccgtgt 4080
cgcccttatt cccttttttg cggcattttg ccttcctgtt tttgctcacc cagaaacgct 4140
ggtgaaagta aaagatgctg aagatcagtt ggggtgcacga gtgggttaca tcgaactgga 4200
tctcaacagc ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc caatgatgag 4260
cactttttaa gttctgctat gtggcgcggt attatcccggt attgacgccg ggcaagagca 4320
actcggtcgc cgcatacact attctcagaa tgacttggtt gagtactcac cagtcacaga 4380
aaagcatctt acggatggca tgacagtaag agaattatgc agtgctgcca taaccatgag 4440
tgataacact gcggccaact tacttctgac aacgatcggg ggaccgaagg agctaaccgc 4500
ttttttgcac aacatggggg atcatgtaac tcgccttgat cgttggggaa cggagctgaa 4560
tgaagccata ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg caacaacggt 4620
gcgcaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat taatagactg 4680
gatggaggcg gataaagttg caggaccact tctgcgctcg gcccttcagg ctggctgggt 4740
tattgctgat aaatctggag ccggtgagcg tgggtctcgc ggtatcattg cagcactggg 4800
gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc aggcaactat 4860
ggatgaacga aatagacaga tcgctgagat aggtgcctca ctgattaagc attggtaact 4920
gtcagaccaa gtttactcat atatacttta gattgattta aaacttcatt tttaatttaa 4980
aaggatctag gtgaagatcc tttttgataa tctcatgacc aaaatccctt aacgtgagtt 5040
ttcgttccac tgagcgtcag accccgtaga aaagatcaaa ggatcttctt gagatccttt 5100
ttttctgcgc gtaatctgct gcttgcaaac aaaaaaacca ccgctaccag cggtggtttg 5160
tttgccggat caagagctac caactctttt tccgaaggta actggcttca gcagagcgca 5220
gataccaaat actgtccttc tagtgtagcc gtagttaggc caccacttca agaactctgt 5280
agcaccgcct acatacctcg ctctgcta at cctgttacca gtggctgctg ccagtggcga 5340
taagtcgtgt cttaccgggt tggactcaag acgatagtta ccggataagg cgcagcggtc 5400
gggctgaacg ggggggttcgt gcacacagcc cagcttgagc cgaacgacct acaccgaact 5460
gagataccta cagcgtgagc tatgagaaag cgccacgctt cccgaaggga gaaaggcgga 5520
caggtatccg gtaagcggca gggtcggaac aggagagcgc acgagggagc ttccaggggg 5580
aaacgcctgg tatctttata gtctgtcgg gtttcgccac ctctgacttg agcgtcgatt 5640
tttgtgatgc tcgtcagggg ggcggagcct atggaaaaac gccagcaacg cggccttttt 5700
acggttcctg gcctttttgt ggcctttttg tcacatgttc tttcctgctg tatcccctga 5760
ttctgtggat aaccgtatta ccgcctttga gtgagctgat accgctcgcc gcagccgaac 5820
gaccgagcgc agcgagtcag tgagcgagga agcggaagag cttaaagcggc cgcggcgcgc 5880
cgcccaatac gcaaaccgcc tctccccgcg cgttgggcga ttcattaatg cagctggcac 5940
gacaggtttc ccgactggaa agcgggcagt gagcgcaacg caattaatgt gagttagctc 6000
actcattagg caccaccaggc tttacacttt atgcttccgg ctctgtatgt gtgtggaatt 6060
gtgagcggat aacaatttca cacaggaaac agctatgacc atgattacgc c 6111

```

<210> 10

<211> 13737

<212> DNA

<213> Plasmid

<220>

<223> Description of Artificial Sequence:Plasmid

<400> 10

```

gatccagaat tcgtgatcaa atggccgcaa caagcagcac aagcagccag tcttttgaca 60
tagagctcga catcatcggc cagcaaccgc ctcttctttc aatctacacc cagatcagtc 120
tcgtttaccc cgtctctgat ccctcccagt atcccaccat cgtcagcacc cttgagggaag 180
gcctaaaacg cctctctcaa accttcccat gggtcgcggg ccagggtcaag accgagggca 240
tcagcgaagg aaacacagga acttccaaga tcattccata tgaggagaca ccccgctcttg 300
tggtgaaaga cctccgtgat gattcctcag cgccaacgat cgagggggtt agaaaggcgg 360
gtttccccct agagatgttt gacgagaacg tcgtcgctcc gaggaagaca ttagctatcg 420
gacctggcaa tggccccaac gaccogaagc ctgtgttgct attgcagctc aacttcatta 480
agggcggact cattctcacc gtcaacggac aacatggtgc tatggacatg acaggacaag 540
atgcaattat tcgtcttctc tccaaggcgt gccgcaacga atcattcacc gaggaggaaa 600
tctcggccat gaacctcgat cgcaagacgg tagtccctct ccttgaaaac taaaagttg 660

```

WO 00/60061

PCT/EP00/02769

gtcctgagct	agaccaccag	atcgccaaac	ctgcgcctgc	tggcgacgct	ccaccgcgac	720
cggccaaggc	aagctgggcg	ttctttttcat	tcactcccaa	ggccctctcg	gagctgaaag	780
acgcagccac	aaagactctt	gacgcgtcgt	ccaagtttgt	gtcaactgat	gatgctcttt	840
cggcgtttat	ctggcaatca	acctcgcgcg	tacgtctcgc	aagattggat	gcttccacac	900
ctactgaatt	ctgccgcgct	gtcgacatgc	ggggcccaat	gggcgtatca	agcacatacc	960
caggccttct	tcaaaacatg	acctaccatg	actcgaccgt	cgccgaaatc	gccaacgaac	1020
cacttggcgc	aacagcatca	cgcctgcgct	cggaaactcaa	cagtgatcgt	ttgcgcagac	1080
gaacacaagc	tttggcgacg	tacatgcatg	gcctgcctga	caagtcgagc	gtctccctga	1140
ccgccgatgc	gaatccgtca	agcagcatca	tgctgagttc	ctgggccaag	gtgggatgct	1200
gggagtatga	ctttgggttt	ggactgggta	agcctgagag	tgtgagaaga	cctcgctttg	1260
aaccttttga	gagtttgatg	tacttttatgc	ccaagaagcc	tgatggggag	tttacggcgt	1320
ccattttctct	gagggatgag	gatatggaga	gactaaaggc	ggatgaggag	tggacaaagt	1380
acgcaaagta	tattgggtag	atagtttact	agactactgc	agggatatcg	tggatcccc	1440
gaatttcccc	gatcgttcaa	acatttggca	ataaagtttc	ttaagattga	atcctgttgc	1500
cggctcttgcg	atgattatca	tctaatttct	gttgaattac	gttaagcatg	taataattaa	1560
catgtaatgc	atgacgttat	ttatgagatg	ggtttttatg	attagagtc	cgcaattata	1620
catttaatac	gcgatagaaa	acaaaatata	gcgcgcaaac	taggataaat	tatcgcgcg	1680
ggtgtcatct	atgttactag	atccgggaat	tggcgcgcc	caattgat	aaatggccgc	1740
tgcggccaat	tcctgcagcg	ttgcggttct	gtcagttcca	aacgtaaaac	ggcttgtccc	1800
gcgtcatcgg	cgggggtcat	aacgtgactc	ccttaattct	ccgctcatga	tcagattgtc	1860
gtttcccgc	ttcagtttaa	actatcagtg	tttgacagga	tatattggcg	ggtaaacct	1920
agagaaaaga	gcgtttatta	gaataatcgg	atatttaaaa	gggcgtgaaa	aggtttatcc	1980
gttcgtccat	ttgtatgtgc	atgccaacca	cagggttccc	cagatctggc	gccggccagc	2040
gagacgagca	agattggccg	ccgcccgaaa	cgatccgaca	gcgcgccag	cacagggtgc	2100
caggcaaatt	gcaccaacgc	atacagcgcc	agcagaatgc	catagtgggc	ggtgacgtcg	2160
ttcgagtga	ccagatcgcg	caggaggccc	ggcagcaccg	gcataatcag	gccgatgccg	2220
acagcgtcga	gcgcgacagt	gctcagaatt	acgatcaggg	gtatgttggg	tttcacgtct	2280
ggcctccgga	ccagcctccg	ctgggtccgat	tgaacgcgcg	gattctttat	cactgataag	2340
ttggtggaca	tattatgttt	atcagtgata	aagtgtcaag	catgacaaag	ttgcagccga	2400
atacagtgat	ccgtgccgcc	ctggacctgt	tgaacgaggt	cggcgtagac	ggtctgacga	2460
cacgcaaact	ggcggaaacg	ttgggggttc	agcagccggc	gctttactgg	cacttcagga	2520
acaagcgggc	gctgctcgac	gcactggccg	aagccatgct	ggcggagaat	catacgcat	2580
cgggtgccgag	agccgacgac	gactggcgct	catttctgat	cgggaatgcc	cgcagcttca	2640
ggcaggcgct	gctcgcttac	cgcgatggcg	cgcgcatcca	tgccggcacg	cgaccgggcg	2700
caccgcagat	ggaaacggcc	gacgcgcagc	ttcgcttcc	ctgcgaggcg	ggtttttcgg	2760
ccggggacgc	cgtcaatgcg	ctgatgacaa	tcagctactt	cactgttggg	gccgtgcttg	2820
aggagcaggc	cggcgacagc	gatgccggcg	agcgcggcgg	caccgttgaa	caggctccgc	2880
tctcgccgct	gttgcggggc	gcgatagacg	ccttcgacga	agccgggtccg	gacgcagcgt	2940
tcgagcaggg	actcgcggtg	attgtcgatg	gattggcgaa	aaggaggctc	gttgtcagga	3000
acgttgaagg	accgagaaag	ggtgacgatt	gatcaggacc	gctgccggag	cgcaaccac	3060
tcactacagc	agagccatgt	agacaacatc	ccctccccct	ttccaccgcg	tcagacgccc	3120
gtagcagccc	gctacgggct	ttttcatgcc	ctgccctagc	gtccaagcct	cacggccgcg	3180
ctcggcctct	ctggcggcct	tctggcgctc	ttccgcttcc	tcgctcactg	actcgctgcg	3240
ctcggtcggt	cggctgcggc	gagcgggtatc	agctcactca	aaggcggtaa	tacggttatc	3300
cacagaatca	ggggataacg	caggaaagaa	catgtgagca	aaaggccagc	aaaaggccag	3360
gaaccgtaaa	aaggccgcgt	tgctggcggt	tttccatagg	ctccgcccc	ctgacgagca	3420
tcacaaaaat	cgacgctcaa	gtcagagggtg	gcgaaacccg	acaggactat	aaagatacca	3480
ggcgtttccc	cctggaagct	ccctcgtgcg	ctctcctggt	ccgaccctgc	cgcttaccgg	3540
atacctgtcc	gcctttctcc	cttcgggaag	cgtggcgctt	ttccgctgca	taaccctgct	3600
tcgggggtcat	tatagcgatt	ttttcggtat	atccatcctt	tttcgcacga	tatacaggat	3660
tttgccaaag	ggttcgtgta	gactttcctt	ggtgtatcca	acggcgctcag	ccgggcagga	3720
taggtgaagt	aggcccaccc	gcgagcgggt	gttccttctt	cactgtccct	tattcgacc	3780
tggcggtgct	caacgggaat	cctgctctgc	gaggctggcc	ggctaccgcc	ggcgtaacag	3840
atgagggcaa	gcggatggct	gatgaaacca	agccaaccag	gaagggcagc	ccacctatca	3900
aggtgtactg	ccttccagac	gaacgaagag	cgattgagga	aaaggcggcg	gcggccggca	3960
tgagcctgtc	ggcctacctg	ctggccgctcg	gccagggcta	caaaatcacg	ggcgctcgtg	4020
actatgagca	cgtccgcgag	ctggcccgcga	tcaatggcga	cctgggcccgc	ctgggcccgc	4080
tgctgaaact	ctggctcacc	gacgacccgc	gcacggcgcg	gttcgggtgat	gccacgatcc	4140
tcgccctgct	ggcgaagatc	gaagagaagc	aggacgagct	tggcaaggct	atgatgggcg	4200
tggctccgcc	gagggcagag	ccatgacttt	tttagccgct	aaaacggccg	gggggtgcgc	4260
gtgattgcca	agcacgtccc	catgcgctcc	atcaagaaga	gcgacttcgc	ggagctgggtg	4320

WO 00/60061

PCT/EP00/02769

aagtacatca	ccgacgagca	aggcaagacc	gagcgccttt	gcgacgctca	ccgggctggt	4380
tgccctcgcc	gctgggctgg	cggccgtcta	tggccctgca	aacgcgccag	aaacgccgtc	4440
gaagccgtgt	gcgagacacc	gcggccggcc	gccggcgttg	tggatacctc	gcggaaaact	4500
tggccctcac	tgacagatga	ggggcgagcg	ttgacacttg	agggggccgac	tcacccggcg	4560
cggcgttgac	agatgagggg	caggctcgat	ttcggccggc	gacgtggagc	tggccagcct	4620
cgaaatcgg	cgaaaacgcc	tgattttacg	cgagtttccc	acagatgatg	tggacaagcc	4680
tggggataag	tgccctgcgg	tattgacact	tgagggggcg	gactactgac	agatgagggg	4740
cgcgacccct	gacacttgag	gggcagagtg	ctgacagatg	aggggcgcac	ctattgacat	4800
ttgaggggct	gtccacaggc	agaaaatcca	gcatttgcaa	gggtttccgc	ccgtttttcg	4860
gccaccgcta	acctgtcttt	taacctgctt	ttaaaccaat	atttataaac	cttggttttta	4920
accagggtg	cgccctgtgc	gcgtgaccgc	gcacgccgaa	gggggggtgcc	cccccttctc	4980
gaaccctccc	ggcccgcctaa	cgcggggcctc	ccatcccccc	aggggctgcg	cccctcggcc	5040
gcgaacggcc	tcaccccaaa	aatggcagcg	ctggcagtc	ttgccattgc	cgggatcggg	5100
gcagtaacgg	gatgggagat	cagcccgagc	gcgacgccc	gaagcattga	cgtgccgcag	5160
gtgctggcat	cgacattcag	cgaccaggtg	ccgggcagtg	agggcgggcg	cctgggtggc	5220
ggcctgccct	tcacttcggc	cgtcggggca	ttcacggact	tcattggcggg	gccggcaatt	5280
tttaccttgg	gcattcttgg	catagtggtc	gcgggtgccg	tgctcgtgtt	cgggggtgcg	5340
ataaaccag	cgaaccattt	gaggtgatag	gtaagattat	accgaggtat	gaaaacgaga	5400
attggacctt	tacagaatta	ctctatgaag	cgccatattt	aaaaagctac	caagacgaag	5460
aggatgaaga	ggatgaggag	gcagattgcc	ttgaatatat	tgacaatact	gataagataa	5520
tatatctttt	atatagaaga	tatcgccgta	tgtaaggatt	tcagggggca	aggcataggc	5580
agcgcgctta	tcaatatatc	tatagaatgg	gcaaagcata	aaaacttgca	tggactaatg	5640
cttgaaaccc	aggacaataa	ccttatagct	tgtaaattct	atcataattg	ggtaatgact	5700
ccaacttatt	gatagtgttt	tatgttcaga	taatgcccg	tgactttgtc	atgcagctcc	5760
accgattttg	agaacgacag	cgacttcctg	cccagccgtg	ccaggtgctg	cctcagattc	5820
aggttatgcc	gctcaattcg	ctgcgtatat	cgcttgctga	ttacgtgcag	ctttcccttc	5880
aggcgggatt	catacagcgg	ccagccatcc	gtcatccata	tcaccacgtc	aaagggtgac	5940
agcaggctca	taagacgccc	cagcgtcgcc	atagtgcgtt	caccgaatac	gtgcgcaaca	6000
accgtcttcc	ggagactgtc	atacgcgtaa	aacagccagc	gctggcgcg	tttagccccg	6060
acatagcccc	actgttcgtc	catttccgcg	cagacgatga	cgtcactgcc	cggctgtatg	6120
cgcgaggtta	ccgactgcgg	cctgagtttt	ttaagtgcg	taaaatcgtg	ttgaggccaa	6180
cgcccataat	gcgggctgtt	gcccggcctc	caacgccatt	catggccata	tcaatgattt	6240
tctggtgcgt	accgggttga	gaagcgggtg	aagtgaactg	cagttgccat	gtttttacggc	6300
agtgcgagca	gagatagcgc	tgatgtccgg	cggtgctttt	gccgttacgc	accaccccg	6360
cagtagctga	acaggaggga	cagctgatag	acacagaagc	cactggagca	cctcaaaaac	6420
accatcatac	actaaatcag	taagttggca	gcatacccca	taattgtggt	ttcaaaatcg	6480
gctccgtcga	tactatgtta	tacgcccaact	ttgaaaacaa	ctttgaaaaa	gctgttttct	6540
ggtattttaag	gtttttagaat	gcaaggaaca	gtgaattgga	gttcgtcttg	ttataattag	6600
cttcttgggg	tatcttttaa	tactgtagaa	aagaggaagg	aaataataaa	tggctaaaat	6660
gagaatatca	ccggaattga	aaaaactgat	cgaaaaatac	cgctgcgtaa	aagatacgg	6720
aggaatgtct	cctgctaagg	tatataagct	ggtgggagaa	aatgaaaacc	tatatttaaa	6780
aatgacggac	agccggtata	aagggaccac	ctatgatgtg	gaacgggaaa	aggacatgat	6840
gctatggctg	gaaggaaagc	tgccctgttc	aaaggctcct	cactttgaac	ggcatgatgg	6900
ctggagcaat	ctgctcatga	gtgaggccga	tggcgtcctt	tgctcggaag	agtatgaaga	6960
tgaacaaagc	cctgaaaaga	ttatcgagct	gtatgcggag	tgcatcaggc	tctttcactc	7020
catcgacata	tcggattgtc	cctatacgaa	tagcttagac	agccgcttag	ccgaattgga	7080
ttacttactg	aataacgatc	tggccgatgt	ggattgcgaa	aactgggaag	aagacactcc	7140
atttaaagat	ccgcgcgagc	tgtatgattt	tttaaagacg	gaaaagcccg	aagaggaact	7200
tgtcttttcc	cacggcgacc	tgggagacag	caacatcttt	gtgaaagatg	gcaaagtaag	7260
tggctttatt	gatcttggga	gaagcggcag	ggcggaacaag	tgggtatgaca	ttgccttctg	7320
cgtccggctg	atcagggagg	atatcgggga	agaacagtat	gtcgagctat	tttttgactt	7380
actggggatc	aagcctgatt	gggagaaaat	aaaatattat	attttactgg	atgaattgtt	7440
ttagtaccta	gatgtggcgc	aacgatgccg	gcgacaagca	ggagcgcacc	gacttcttcc	7500
gcatacaagt	ttttggctct	caggccgagg	cccacggcaa	gtatttgggc	aagggggtcgc	7560
tgggtattcgt	gcagggcaag	attcggaata	ccaagtacga	gaaggacggc	cagacgggtc	7620
acgggaccga	cttcattgcc	gataaggtgg	attatctgga	caccaaggca	ccaggcgggt	7680
caaatcagga	ataagggcac	attgccccgg	cgtgagtcgg	ggcaatccc	caaggagggt	7740
gaatgaatcg	gacgtttgac	cggaaggcat	acaggcaaga	actgatcgac	gcgggggtttt	7800
ccgccgagga	tgccgaaacc	atcgcaagcc	gcaccgtcat	gcgtgcgccc	cgcgaaacct	7860
tccagtcctg	cggctcgatg	gtccagcaag	ctacggccaa	gatcgagcgc	gacagcgtgc	7920
aactggctcc	ccctgccctg	cccgcgccat	cggccgcctg	ggagcgttcg	cgtcgtctcg	7980

WO 00/60061

PCT/EP00/02769

aacaggaggc	ggcagggtttg	gcgaagtcga	tgaccatcga	cacgcgagga	actatgacga	8040
ccaagaagcg	aaaaaccgcc	ggcgaggacc	tggcaaaaca	ggtcagcgag	gccaagcagg	8100
ccgcgttgct	gaaacacacg	aagcagcaga	tcaaggaaat	gcagctttcc	ttgttcgata	8160
ttgcgcctg	gccggacacg	atgcgagcga	tgccaaacga	cacggcccgc	tctgccctgt	8220
tcaccacgcg	caacaagaaa	atccccgcgcg	aggcgctgca	aaacaaggtc	attttccacg	8280
tcaacaagga	cgtgaagatc	acctacaccg	gcgtcgagct	gcgggcccgc	gatgacgaac	8340
tgggtgtggca	gcaggtgttg	gagtacgcga	agcgcacccc	tatcggcgag	ccgatcacct	8400
tcacgttcta	cgagctttgc	caggacctgg	gctggctgat	caatggcccg	tattacacga	8460
aggccgagga	atgcctgtcg	cgcctacagg	cgacggcgat	gggcttcacg	tccgaccgcg	8520
ttgggcacct	ggaatcggtg	tcgctgctgc	accgcttcgc	cgtcctggac	cgtggcaaga	8580
aaacgtcccg	ttgccagggtc	ctgatcgacg	aggaaatcgt	cgtgctgttt	gctggcgacc	8640
actacacgaa	attcatatgg	gagaagtacc	gcaagctgtc	gccgacggcc	cgacggatgt	8700
tcgactattt	cagctcgcac	cgggagccgt	acccgctcaa	gctggaaacc	ttccgcctca	8760
tgtgcggatc	ggattccacc	cgctgaaga	agtggcgcg	gcaggtcggc	gaagcctgcg	8820
aagagttgcg	aggcagcggc	ctggtggaac	acgcctgggt	caatgatgac	ctggtgcatt	8880
gcaaacgcta	gggccttgtg	gggtcagttc	cggctggggg	ttcagcagcc	agcgttttac	8940
tggcatttca	ggaacaagcg	ggcactgctc	gacgcacttg	cttcgctcag	tatcgctcgg	9000
gacgcacggc	gcgctctacg	aactgccgat	aaacagagga	ttaaaattga	caattgtgat	9060
taaggctcag	attcgacggc	ttggagcggc	cgacgtgcag	gatttccgcg	agatccgatt	9120
gtcggccctg	aagaaagctc	cagagatggt	cgggtccgtt	tacgagcacg	aggagaaaaa	9180
gcccattggag	gcgttcgctg	aacggttgcg	agatgccgtg	gcattcggcg	cctacatcga	9240
cggcgagatc	attgggctgt	cggctctcaa	acaggaggac	ggccccaagg	acgctcacia	9300
ggcgcatctg	tccggcgttt	tcgtggagcc	cgaacagcga	ggccgagggg	tcgcccgtat	9360
gctgctgcgg	gcgttgccgg	cgggtttatt	gctcgtgatg	atcgtccgac	agattccaac	9420
gggaatctgg	tggatgcgca	tcttcatcct	cggcgcactt	aatatttcgc	tattctggag	9480
cttgttgttt	atttcgggtc	accgcctgcc	gggcgggggtc	gcggcgacgg	taggcgctgt	9540
gcagccgctg	atggtcgtgt	tcattctctgc	cgctctgcta	ggtagcccga	tacgattgat	9600
ggcggctcctg	ggggctattt	gcggaactgc	gggcgtggcg	ctggtgggtg	tgacaccaaa	9660
cgcagcgcta	gatcctgtcg	gcgtcgcagc	gggcctggcg	ggggcggttt	ccatggcggt	9720
cggaaaccgtg	ctgaccgcga	agtggcaacc	tcccgtgcct	ctgctcacct	ttaccgcctg	9780
gcaactggcg	gccggaggac	ttctgctcgt	tccagtagct	ttagtggttg	atccgccaat	9840
cccgatgcct	acaggaacca	atgttctcgg	cctggcggtg	ctcggcctga	tcggagcggg	9900
tttaacctac	ttccttttgt	tccgggggat	ctcgcgactc	gaacctacag	ttgtttcctt	9960
actgggcttt	ctcagcccca	gatctgggggt	cgatcagccg	gggatgcac	aggccgacag	10020
tcggaacttc	gggtccccga	cctgtaccat	tcggtgagca	atggataggg	gagttgatat	10080
cgtcaacggt	cacttctaaa	gaaatagcgc	cactcagctt	cctcagcggc	tttatccagc	10140
gatttcctat	tatgtcggca	tagttctcaa	gatcgacagc	ctgtcacggg	taagcgagaa	10200
atgaataaga	aggctgataa	ttcggatctc	tgcgagggag	atgatatttg	atcacaggca	10260
gcaacgctct	gtcatcggtt	caatcaacat	gctaccctcc	gcgagatcat	ccgtgtttca	10320
aaccgcggcag	cttagttgcc	gttcttccga	atagcatcgg	taacatgagc	aaagtctgcc	10380
gccttacaac	ggctctcccg	ctgacgcctg	cccggactga	tgggctgcct	gtatcgagtg	10440
gtgattttgt	gccgagctgc	cggtcgggga	gctgttggct	ggctgggtggc	aggatatatt	10500
gtggtgtaaa	caaattgacg	cttagacaac	ttaataacac	attgcggacg	tttttaaatg	10560
actgcggtac	ggccatgctg	gccgcccggg	caccggtaaa	tttcctgcag	ggctagcgaa	10620
ttcgagctcg	gtaccctctg	attttggttt	taggaattag	attattgata	gaagtatttt	10680
acaaatacaa	atacatacta	agggtttctt	atatgctcaa	cacatgagcg	aaaccctata	10740
agaaccctaa	ttcccttatc	tgggaactac	tcacacatta	ttatagagag	agatagattt	10800
gtagagagag	actggtgatt	tcagcgggca	tgcttgcagg	tcgactcaga	tctgggtaac	10860
tggcctaact	ggccttggag	gagctggcaa	ctcaaaatcc	ctttgccaaa	aaccaacatc	10920
atgccatcca	ccatgcttgt	atccagctgc	gcgcaatgta	ccccgggctg	tgtatcccaa	10980
agcctcatgc	aacctaacag	atggatcggt	tgggaaggcct	ataacagcaa	ccacagactt	11040
aaaaccttgc	gcctccatag	acttaagcaa	atgtgtgtac	aatgtggatc	ctaggcccaa	11100
cctttgatgc	ctatgtgaca	cgtaaacagt	actctcaact	gtccaatcgt	aagcgttcct	11160
agccttccag	ggcccagcgt	aagcaatacc	agccacaaca	ccctcaacct	cagcaaccaa	11220
ccaagggtat	ctatcttgca	acctctctag	atcatcaatc	cactcttgtg	gtgtttgtgg	11280
ctctgtccta	aagttcactg	tagacgtctc	aatgtaatgg	ttaacgatat	cacaaaccgc	11340
ggccatatca	gctgctgtag	ctggcctaata	ctcaactggg	ctcctctccg	gagacatgtc	11400
gactctagag	gatccccggg	taccctgtcc	tctccaaatg	aatgaactt	ccttatatag	11460
aggaagggtc	ttgcgaagga	tagtgggatt	gtgcgtcatc	ccttacgtca	gtggagatat	11520
cacatcaatc	cacttgcttt	gaagacgtgg	ttggaacgtc	ttctttttcc	acgatgctcc	11580
tcgtgggtgg	gggtccatct	ttgggaccac	tgctcggcaga	ggcatcttca	acgatggcct	11640

WO 00/60061

PCT/EP00/02769

```

ttcctttatc gcaatgatgg catttgtagg agccaccttc cttttccact atcttcacaa 11700
taaagtgaca gatagctggg caatggaatc cgaggagggt tccggatatt accctttggt 11760
gaaaagtctc aattgccctt tggctctctg agactgtatc tttgatattt ttggagtaga 11820
caagcgtgtc gtgctccacc atgttgacga agatattctt cttgtcattg agtcgtaaga 11880
gactctgtat gaactgttcg ccagtcttta cggcgagttc tgttggtcct ctatttgaat 11940
ctttgactcc atgggaattg agatctctcg aggtttaaac gggccacgcc tgcggccgcc 12000
tcgagggtacc ggatttggag ccaagtctca taaacgccat tgtggaagaa agtcttgagt 12060
tggtggtaat gtaacagagt agtaagaaca gagaagagag agagtgtgag atacatgaat 12120
tgtcgggcaa caaaaatcct gaacatctta ttttagcaaa gagaaagagt tccgagtctg 12180
tagcagaaga gtgaggagaa atttaagctc ttggacttgt gaattgttcc gcctcttgaa 12240
tacttcttca atcctcatat attcttcttc tatgttacct gaaaaccggc atttaatctc 12300
gcggttttat tccggttcaa catttttttt gttttgagtt attatctggg ctttaataacg 12360
caggcctgaa ataaattcaa ggcccaactg tttttttttt taagaagttg ctgttaaaaa 12420
aaaaaaaaagg gaattaacaa caacaacaaa aaaagataaa gaaaataata acaattactt 12480
taattgtaga ctaaaaaaac atagatttta tcatgaaaaa aagagaaaag aaataaaaaac 12540
ttggatcaaa aaaaaaaca tacagatctt ctaattatta acttttctta aaaattaggt 12600
cctttttccc aacaattagg ttttagagtt tgggaattaaa ccaaaaagat tgttctaaaa 12660
aatactcaaa tttggttagat aagtttcctt attttaatta gtcaatggta gatacttttt 12720
tttcttttct ttattagagt agattagaat cttttatgcc aagttttgat aaattaaatc 12780
aagaagataa actatcataa tcaacatgaa attaaaagaa aaatctcata tatagtatta 12840
gtattctcta tatatattat gattgcttat tcttaatggg ttgggttaac caagacatag 12900
tcttaatgga aagaatcttt tttgaacttt ttccttattg attaaattct tctatagaaa 12960
agaaagaaat tatttgagga aaagtatata caaaaagaaa aatagaaaaa tgtcagtga 13020
gcagatgtaa tggatgacct aatccaacca ccaccatagg atgtttctac ttgagtcggt 13080
cttttaaaaa cgcacggtgg aaaatatgac acgtatcata tgattccttc ctttagtttc 13140
gtgataataa tcctcaactg atatcttctt ttttttgttt tggctaaaga tattttattc 13200
tcattaatag aaaagacggt tttgggcttt tggtttgcca tataaagaag accttcgtgt 13260
ggaagataat aattcatcct ttcgtctttt tctgactctt caatctctcc caaagcctaa 13320
agcgatctct gcaaatctct cgcgactctc tctttcaagg tatattttct gattcttttt 13380
gtttttgatt cgtatctgat ctccaatttt tgttatgtgg attattgaat cttttgtata 13440
aattgctttt gacaatattg ttcgtttcgt caatccagct tctaaatttt gtcctgatta 13500
ctaagatata gattcgtagt gtttacatct gtgtaatttc ttgcttgatt gtgaaattag 13560
gattttcaag gacgatctat tcaatttttg tgttttcttt gttcgattct ctctgtttta 13620
ggtttcttat gtttagatcc gtttctcttt ggtgttgttt tgatttctct tacggctttt 13680
gatttggtat atgttcgctg attggtttct acttgttcta ttgttttatt tcagggtg 13737

```

<210> 11

<211> 12949

<212> DNA

<213> Plasmid

<400> 11

```

agcttgcagt cctgcagtgc agcgtgaccc ggtcgtgccc ctctctagag ataatgagca 60
ttgcatgtct aagttataaa aaattaccac atattttttt tgtcacactt gtttgaagtg 120
cagtttatct atctttatac atatatttaa actttactct acgaataata taatctatag 180
tactacaata atatcagtgt tttagagaat catataaatg aacagttaga catggtctaa 240
aggacaattg agtattttga caacaggact ctacagtttt atcttttttag tgtgcatgtg 300
ttctcctttt tttttgcaaa tagcttcacc tatataatac ttcattccatt ttattagtag 360
atccatttag ggttttaggg taatggtttt tatagactaa ttttttttagt acatctattt 420
tattctattt tagcctctaa attaagaaaa ctaaaactct atttttagttt ttttatttaa 480
taatttagat ataaaataga ataaaataaa gtgactaaaa attaaacaaa taccctttaa 540
gaaattaaaa aaactaagga aacatttttc ttgtttcgag tagataatgc cagcctgtta 600
aacgccgtcg acgagtctaa cggacaccaa ccagcgaacc agcagcgtcg cgtcgggcca 660
agcgaagcag acggcacggc atctctgtcg ctgcctctgg accctctctg agagttccgc 720
tccaccgttg gacttgctcc gctgtcggca tccagaaatt gcgtggcgga gcggcagacg 780
tgagccggca cggcaggcgg cctcctcttc ctctcacggc accggcagct acgggggatt 840
cctttcccac cgctccttcg ctttcccttc ctgcgccgcc gtaataaata gacacccctt 900
ccacaccctc tttccccaac ctctgtgtgt tcggagcgca cacacacaca accagatctc 960
cccaaatacc acccgtcggc acctccgctt caaggtacgc cgctcgtcct ccccccccc 1020
ccctctctac ctctcttaga tcggcggttc ggtccatggg tagggcccg tagttctact 1080
tctgttcatg tttgtgttag atccgtgttt gtgttagatc cgtgctgcta gcgttcgtac 1140

```

WO 00/60061

PCT/EP00/02769

acggatgcga	cctgtacgtc	agacacgttc	tgattgctaa	cttgccagt	tttctctttg	1200
gggaatcctg	ggatggctct	agccgttccg	cagacgggat	cgatttcatg	atTTTTTTTg	1260
tttcgttgca	tagggtttgg	tttgcccttt	tcctttatTT	caatatatgc	cgtgcacttg	1320
tttgtcgggt	catcttttca	tgcttttttt	tgtcttggtt	gtgatgatgt	ggtctgggtg	1380
ggcggtcgtt	ctagatcgga	gtagaattct	gtttcaaact	acctgggtga	tttattaatt	1440
ttggatctgt	atgtgtgtgc	catacatatt	catagttacg	aattgaagat	gatggatgga	1500
aatatcgatc	taggataggt	atacatgttg	atgcgggttt	tactgatgca	tatacagaga	1560
tgctttttgt	tcgcttggtt	gtgatgatgt	ggtgtgggtg	ggcggtcgtt	cattcgttct	1620
agatcggagt	agaatactgt	ttcaaactac	ctgggtgtatt	tattaatttt	ggaactgtat	1680
gtgtgtgtca	tacatcttca	tagttacgag	tttaagatgg	atggaaatat	cgatctagga	1740
taggtataca	tgttgatgtg	ggttttactg	atgcatatac	atgatggcat	atgcagcatc	1800
tattcatatg	ctctaacctt	gagtacctat	ctattataat	aaacaagtat	gtttttataat	1860
tattttgatc	ttgatatact	tggatgatgg	catatgcagc	agctatatgt	ggattttttt	1920
agccctgcct	tcatacgcta	tttattttgt	tggtactgtt	tcttttgctg	atgctcacc	1980
tggtgtttgg	tgttacttct	gcagggatcc	ccgatcatgc	aaaaactcat	taactcagtg	2040
caaaactatg	cctggggcag	caaaacggcg	ttgactgaac	tttatgggat	ggaaaatccg	2100
tccagccagc	cgatggccga	gctgtggatg	ggcgcacatc	cgaaaagcag	ttcacgagtg	2160
cagaatgccg	ccggagatat	cgtttctactg	cgtgatgtga	ttgagagtga	taaatcgact	2220
ctgctcggag	aggccgttgc	caaacgcttt	ggcgaactgc	ctttcctgtt	caaagtatta	2280
tgcgacgac	agccactctc	cattcagggtt	catccaaaca	aacacaattc	tgaaatcggg	2340
tttgccaaag	aaaatgccgc	aggtatcccg	atggatgccg	ccgagcgtaa	ctataaagat	2400
cctaaccaca	agccggagct	ggttttttg	ctgacgcctt	tccttgcgat	gaacgcgttt	2460
cgtgaatttt	ccgagattgt	ctccctactc	cagccggctg	caggtgcaca	tccggcgatt	2520
gctcactttt	tacaacagcc	tgatgccgaa	cgtttaagcg	aactgttcgc	cagcctgttg	2580
aatatgcagg	gtgaagaaaa	atcccgcgcg	ctggcgattt	taaaatcggc	cctcgatagc	2640
cagcaggggt	aaccgtggca	aacgattcgt	ttaattttctg	aatttttacc	ggaagacagc	2700
ggtctgttct	ccccgctatt	gctgaatgtg	gtgaaattga	accctggcga	agcgatgttc	2760
ctgttcgctg	aaacaccgca	cgcttacctg	caaggcggtg	cgctggaagt	gatggcaaac	2820
tccgataacg	tgctgcgtgc	gggtctgacg	cctaaataca	ttgatattcc	ggaactgggt	2880
gccaatgtga	aattcgaagc	caaaccggct	aaccagttgt	tgaccagacc	ggtgaaacaa	2940
ggtgcagaac	tggacttccc	gattccagtg	gatgattttg	ccttctcgct	gcatgacctt	3000
agtataaag	aaaccaccat	tagccagcag	agtgccgcca	ttttgttctg	cgtcgaaggc	3060
gatgcaacgt	tgtggaaagg	ttctcagcag	ttacagctta	aaccgggtga	atcagcggtt	3120
attgccgcca	acgaatcacc	ggtgactgtc	aaaggccacg	gccgttttag	gcgtgtttac	3180
aacaagctgt	aagagcttac	tgaaaaaatt	aacatctctt	gctaagctgg	gagctcgatc	3240
cgtcgacctg	cagatcgttc	aaacattttg	caataaagtt	tcttaagatt	gaatcctgtt	3300
gccggtcttg	cgatgattat	catataattt	ctgttgaatt	acgttaagca	tgtaataatt	3360
aacatgtaat	gcatgacgtt	atttatgaga	tgggttttta	tgattagagt	cccgcatta	3420
tacatttaat	acgcgataga	aaacaaaata	tagcgcgcaa	actaggataa	attatcgcg	3480
gcggtgtcat	ctatgttact	agatctgcta	gccctgcagg	aaatttaccg	gtgcccgggc	3540
ggccagcatg	gccgtatccg	caatgtgtta	ttaagtgtgc	taagcgtcaa	tttgtttaca	3600
ccacaatata	tcctgccacc	agccagccaa	cagctccccg	accggcagct	cggcacaaaa	3660
tcaccactcg	atacaggcag	cccatcagaa	ttaattctca	tgtttgacag	cttatcatcg	3720
actgcacggg	gcaccaatgc	ttctggcgct	aggcagccat	cggaagctgt	ggtatggctg	3780
tgcaggtcgt	aaatcactgc	ataattcgtg	tcgctcaagg	cgcactcccg	ttctggataa	3840
tgtttttttg	gccgacatca	taacgggttct	ggcaaatatt	ctgaaatgag	ctgttgacaa	3900
ttaatcatcc	ggctcgtata	atgtgtggaa	ttgtgagcgg	ataacaattt	cacacaggaa	3960
acagaccatg	aggggaagcg	tgatcgccga	agtatcgact	caactatcag	aggtagttag	4020
cgtcatcgag	cgccatctcg	aaccgacgtt	gctggccgta	catttgtagc	gctccgcagt	4080
ggatggcggc	ctgaagccac	acagtgatat	tgatttgctg	gttacgggtg	ccgtaaggct	4140
tgatgaaaca	acgcggcgag	ctttgatcaa	cgaccttttg	gaaacttcgg	cttccccctg	4200
agagagcgag	attctccgcg	ctgtagaagt	caccattgtt	gtgcacgacg	acatcattcc	4260
gtggcggttat	ccagctaagc	gcgaactgca	atttgagaaa	tggcagcgca	atgacattct	4320
tgcaggtatc	ttcgagccag	ccacgatcga	cattgatctg	gctatcttgc	tgacaaaagc	4380
aagagaacat	agcgttgcc	tggtaggtcc	agcggcggag	gaactctttg	atccgggttc	4440
tgaacaggat	ctatttgagg	cgctaaatga	aaccttaacg	ctatggaaact	cgcgcgccga	4500
ctgggctggc	gatgagcgaa	atgtagtgct	tacgttgctc	cgcatttggt	acagcgcagt	4560
aaccggcaaa	atcgcgccga	aggatgtcgc	tgccgactgg	gcaatggagc	gcctgccggc	4620
ccagtatcag	cccgtcatac	ttgaagctag	gcaggcttat	cttggaacaag	aagatcgctt	4680
ggcctcgcg	gcagatcagt	tggaagaatt	tgttcactac	gtgaaaggcg	agatcaccaa	4740
agtagtcggc	aaataaagct	ctagtggatc	tccgtacccc	cgggggatct	ggctcgcggc	4800

ggacgcacga	cgccggggcg	agaccatag	cgatctccta	aatcaatagt	agctgtaacc	4860
tcgaagcgtt	tcacttgtaa	caacgattga	gaatttttgt	cataaaattg	aaatacttgg	4920
ttcgcathtt	tgtcatccgc	ggtcagccgc	aattctgacg	aactgcccac	ttagctggag	4980
atgattgtac	atccttcacg	tgaaaatttc	tcaagcgctg	tgaacaaggg	ttcagathtt	5040
agattgaaag	gtgagccggt	gaaacacggt	cttcttgctg	atgacgacgt	cgctatgcgg	5100
catcttatta	ttgaatacct	tacgatccac	gccttcaaag	tgaccgcggt	agccgacagc	5160
accagttca	caagagtact	ctcttccgcg	acggtcgatg	tcgtggttgt	tgatctaaat	5220
ttaggtcgtg	aagatgggct	cgagatcggt	cgtaatctgg	cggcaaagtc	tgatattcca	5280
atcataatta	tcagtggcga	ccgccttgag	gagacggata	aagttggtgc	actcgagcta	5340
ggagcaagtg	attttatcgc	taagccgttc	agtatcagag	agtttctagc	acgcattcgg	5400
gttgccctgc	gcgtgcgccc	caacgttggt	cgctccaaag	accgacgggt	tttttgtttt	5460
actgactgga	cacttaattc	caggcaacgt	cgcttgatgt	ccgaagctgg	cggtgaggtg	5520
aaacttacgg	caggtgagtt	caatcttctc	ctcgcgtttt	tagagaaacc	ccgcgacggt	5580
ctatcgcgcg	agcaacttct	cattgccagt	cgagtacgcg	acgaggaggt	ttatgacagg	5640
agtatagatg	ttctcatttt	gaggctgcgc	cgcaaacttg	aggcagatcc	gtcaagccct	5700
caactgataa	aaacagcaag	aggtgccggt	tatttctttg	acgcggacgt	gcaggtttcg	5760
cacgggggga	cgatggcagc	ctgagccaat	tcccagatcc	ccgaggaatc	ggcgtgagcg	5820
gtcgcaaacc	atccggcccg	gtacaaatcg	gcgcggcgct	gggtgatgac	ctggtggaga	5880
agttgaaggc	cgcgccaggc	gccagcgggc	aacgcacatc	ggcagaagca	cgccccggtg	5940
aatcgtggca	agcggccgct	gatcgaatcc	gcaaagaatc	ccggcaaccg	ccggcagccg	6000
gtgcgcccgc	gattaggaag	ccgcccgaag	gcgacgagca	accagatttt	ttcgttccga	6060
tgctctatga	cgtgggcacc	cgcgatagtc	gcagcatcat	ggacgtggcc	gttttccgct	6120
tgtcgaagcg	tgaccgacga	gctggcgagg	tgatccgcta	cgagcttcca	gacgggcacg	6180
tagaggtttc	cgccgggccc	gccggcatgg	ccagtgtgtg	ggattacgac	ctggtactga	6240
tggcggtttc	ccatctaacc	gaatccatga	accgataccg	ggaagggaag	ggagacaagc	6300
ccggccgctg	gttccgtcca	cacgttgccg	acgtactcaa	gttctgccgg	cgagccgatg	6360
gcggaaagca	gaaagacgac	ctggtagaaa	cctgcattcg	gttaaaccac	acgcacgttg	6420
ccatgcagcg	tacgaagaag	gccaaagaac	gccgcctggt	gacggatatcc	gaggggtgaag	6480
ccttgattag	ccgctacaag	atcgtaaaga	gcgaaaccgg	gcggccggag	tacatcgaga	6540
tcgagctagc	tgattggatg	taccgcgaga	tcacagaagg	caagaaccgg	gacgtgctga	6600
cggttcacc	cgattacttt	ttgatcgatc	ccggcatcgg	ccgttttctc	taccgcctgg	6660
cacgccgcgc	cgccaggcaag	gcagaagcca	gatggtttgt	caagacgatc	tacgaacgca	6720
gtggcagcgc	cggagagttc	aagaagtctc	gtttcaccgt	gcgcaagctg	atcgggtcaa	6780
atgacctgcc	ggagtacgat	ttgaaggagg	aggcggggca	ggctggcccg	atcctagtca	6840
tgcgctaccg	caacctgatc	gagggcgaa	catccgcggg	ttcctaattg	acggagcaga	6900
tgctagggca	aattgcccta	gcaggggaaa	aaggtcgaaa	aggtctcttt	cctgtggata	6960
gcacgtacat	tggaaccaca	aagccgtaca	ttgggaaccg	gaaccggtac	attgggaacc	7020
caaagccgta	cattgggaac	cggtcacaca	tgtaagtga	tgatataaaa	gagaaaaaag	7080
gcgatttttc	cgcctaaaac	tctttaaaa	ttattaaaac	tcttaaaaac	cgcctggcct	7140
gtgcataact	gtctggccag	cgccagcccg	aagagctgca	aaaagcgcc	acccttcggt	7200
cgctgcgctc	cctacgcccc	gccgcttcgc	gtcggcctat	cgccggccgct	ggccgctcaa	7260
aatggctgg	cctacggcca	ggcaatctac	cagggcgcg	acaagccg	ccgtcgccac	7320
tcgaccgccc	gcgctgaggt	ctgcctcggt	aagaagggtg	tgctgactca	taccaggcct	7380
gaatcgcccc	atcatccagc	cagaaagtga	gggagccacg	gttgatgaga	gctttgttgt	7440
aggtggacca	gttggtgatt	ttgaactttt	gctttgccac	ggaacggtct	gcgttgctcg	7500
gaagatgcgt	gatctgatcc	ttcaactcag	caaaagtctg	atttattcaa	caaagccgcc	7560
gtcccgtcaa	gtcagcgtaa	tgctctgcca	gtgttacaac	caattaacca	attctgatta	7620
gaaaaactca	tcgagcatca	aatgaaactg	caattttatt	atatcaggat	tatcaatacc	7680
atatttttga	aaaagccggt	tctgtaatga	aggagaaaac	tcaccgaggc	agttccatag	7740
gatggcaaga	tcctgggtatc	ggctctgcgat	tccgactcgt	ccaacatcaa	tacaacctat	7800
taattttccc	tcgtcaaaaa	taagggtatc	aagtgagaaa	tcaccatgag	tgacgactga	7860
atccggtgag	aatggcaaaa	gctctgcatt	aatgaatcgg	ccaacgcg	gggagaggcg	7920
gtttgcgtat	tgggcgctct	tccgcttcct	cgctcactga	ctcgctgcgc	tcggtcgctc	7980
ggctgcggcg	agcggtatca	gctcactcaa	aggcggtaat	acggttatcc	acagaatcag	8040
gggataacgc	aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	8100
aggccgcgtt	gctggcggtt	ttccataggg	tccgcccccc	tgacgagcat	cacaaaaatc	8160
gacgctcaag	tcagaggtgg	cgaaaccgga	caggactata	aagataccag	gcgtttcccc	8220
ctggaagctc	cctcgtgcgc	tctcctgttc	cgaccctgcc	gcttaccgga	tacctgtccg	8280
cctttctccc	ttcgggaagc	gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	8340
cgggtgtaggt	cgttcgctcc	aagctggggt	gtgtgcacga	accccccggt	cagccccgac	8400
gctgcgcctt	atccggtaac	tatcgtcttg	agccaacccc	ggtaagacac	gacttatcgc	8460

cactggcagc	agccactggt	aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	8520
agttcttgaa	gtggtggcct	aactacggct	acactagaag	aacagtatct	ggtatctgcg	8580
ctctgctgaa	gccagttacc	ttcggaaaaa	gagttggtag	ctcttgatcc	ggcaaacaaa	8640
ccaccgctgg	tagcgggtgt	ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	8700
gatctcaaga	agatcctttg	atcttttcta	cggggtctga	cgctcagtgg	aacgaaaact	8760
cacgttaagg	gatttttggtc	atgagattat	caaaaaggat	cttcacctag	atccttttga	8820
tccggaatta	attcctgtgg	ttggcatgca	catacaaagt	gacgaacgga	taaacccttt	8880
cacgcccttt	taaatatccg	attatttcta	taaacgctct	tttctcttag	gtttaccctg	8940
caatatatcc	tgtcaaacac	tgatagttta	aactgaaggc	gggaaacgac	aatctgatca	9000
tgagcggaga	attaagggag	tcacgttatg	acccccgccc	atgacgcggg	acaagccggt	9060
ttacgttttg	aactgacaga	accgcaacgc	tgcaggaatt	ggccgcagcg	gccatttaaa	9120
tggtacctta	attaacgtac	gaagcttgca	tgcacgcggg	ctagagcggc	cgcctcgagg	9180
taccggggccc	cccctcgagg	tcgacgggat	cgataagctt	gcatgcctgc	agtgcagcgt	9240
gaccgggtcg	tgccccctct	tagagataat	gagcattgca	tgtctaagtt	ataaaaaaatt	9300
accacatatt	ttttttgtca	cacttggttg	aagtgcagtt	tatctatctt	tatacatata	9360
tttaaacttt	actctacgaa	taatataatc	tatagtacta	caataatatc	agtgttttag	9420
agaatcatat	aaatgaacag	ttagacatgg	tctaaaggac	aattgagtat	tttgacaaca	9480
ggactctaca	gtttttatct	tttagtgtgc	atgtgttctc	cttttttttt	gcaaataagct	9540
tcacctatat	aatacttcat	ccatttttatt	agtacatcca	tttagggttt	agggttaatg	9600
gttttttatag	actaattttt	ttagtacatc	tatttttatc	tatttttagcc	tctaaattaa	9660
gaaaactaaa	actctatttt	agttttttta	tttaataatt	tagatataaa	atagaataaa	9720
ataaagtgac	taaaaattaa	acaaataccc	tttaagaaat	taaaaaaact	aaggaaacat	9780
ttttcttggt	tcgagtagat	aatgccagcc	tgttaaacgc	cgtcgacgag	tctaacggac	9840
accaaccagc	gaaccagcag	cgctcgctcg	ggccaagcga	agcagacggc	acggcatctc	9900
tgctcgctgc	tctggacccc	tctcgagagt	tccgctccac	cgttggactt	gctccgctgt	9960
cggcatccag	aaattgcgtg	gcggagcggc	agacgtgagc	cggcacggca	ggcggcctcc	10020
tcctcctctc	acggcacggc	agctacgggg	gattcctttc	ccaccgctcc	ttcgctttcc	10080
cttcctcgcc	cgccgtaata	aatagacacc	ccctccacac	cctctttccc	caacctcggt	10140
ttgttcggag	cgcacacaca	cacaaccaga	tctcccccaa	atccaccctg	cggcacctcc	10200
gcttcaaggt	acgccgctcg	tcctcccccc	ccccccctct	ctacctctc	tagatcggcg	10260
ttccgggtcca	tggttagggc	ccggtagttc	tacttctggt	catgtttgtg	ttagatccgt	10320
gtttgtgtta	gatccgtgct	gctagcgttc	gtacacggat	gcgacctgta	cgtcagacac	10380
gttctgattg	ctaacttgcc	agtgtttctc	tttggggaat	cctgggatgg	ctctagccgt	10440
tccgcagacg	ggatcgatct	catgattttt	tttggttctg	tgcataagggt	ttgggtttg	10500
cttttccttt	atctcaatat	atgccgtgca	cttggttgct	gggtcatctt	ttcatgcttt	10560
tttttgctct	ggttgtgatg	atgtggtctg	gttggtgctg	cgttctagat	cggagtagaa	10620
ttctgtttca	aactacctgg	tggattttat	aattttggat	ctgtatgtgt	gtgccataca	10680
tattcatagt	tacgaattga	agatgatgga	tggaaatatc	gatctaggat	aggtatacat	10740
gttgatgcgg	gttttactga	tgcatataca	gagatgcttt	ttgttcgctt	ggttgtgatg	10800
atgtggtgtg	gttgggcggg	cgttcattcg	ttctagatcg	gagtagaata	ctgtttcaaa	10860
ctacctgggtg	tattttattaa	ttttggaact	gtatgtgtgt	gtcatacatc	ttcatagtta	10920
cgagtttaag	atggatggaa	atatcgatct	aggataggta	tacatgttga	tgtgggtttt	10980
actgatgcat	atacatgatg	gcataatgcag	catctattca	tatgctctaa	ccttgagtac	11040
ctatctatta	taataaacia	gtatgtttta	taattatttt	gatcttgata	tacttggtatg	11100
atggcatatg	cagcagctat	atgtggattt	tttttagccct	gccttcatac	gctattttatt	11160
tgcttggtac	tgttttcttt	gtcgatgctc	accctgttgt	ttgggtgttac	ttctgcagggt	11220
cgactctaga	ggatccagaa	ttcgtgatca	aatggccgca	acaagcagca	caagcagcca	11280
gtcttttgac	atagagctcg	acatcatcgg	ccagcaaccg	cctcttcttt	caatctacac	11340
ccagatcagt	ctcgtttacc	ccgtctctga	tccctcccag	tatcccacca	tcgtcagcac	11400
ccttgaggaa	ggcctaaaac	gcctctctca	aaccttccca	tgggtcgcgg	gccaggtcaa	11460
gaccgagggc	atcagcgaag	gaaacacagg	aacttccaag	atcattccat	atgaggagac	11520
accccgctct	gtggtgaaag	acctccgtga	tgattcctca	gcgccaacga	tcgaggggtt	11580
gagaaaggcg	ggtttcccct	tagagatgtt	tgacgagaac	gtcgtcgtct	cgaggaagac	11640
attagctatc	ggacctggca	atggccccaa	cgacccgaag	cctgtgttgc	tattgcagct	11700
caacttcatt	aagggcggac	tcattctcac	cgtcaacgga	caacatgggt	ctatggacat	11760
gacaggacaa	gatgcaatta	ttcgtcttct	ctccaaggcg	tgccgcaacg	aatcattcac	11820
cgaggaggaa	atctcggcca	tgaacctcga	tcgcaagacg	gtagtccctc	tccttgaaaa	11880
ctacaaagtt	ggtcctgagc	tagaccacca	gacgcgcaaa	cctgcgctct	ctggcgacgc	11940
tccaccgcga	ccggccaagg	caagctgggc	gttcttttca	ttcactccca	aggccctctc	12000
ggagctgaaa	gacgcagcca	caaagactct	tgacgcgtcg	tccaagtgtg	tgtcaactga	12060
tgatgctctt	tcggcggtta	tctggcaatc	aacctcgcgc	gtacgtctcg	caagattgga	12120

WO 00/60061

PCT/EP00/02769

tgcttccaca	cctactgaat	tctgccgcgc	tgtegcacatg	cggggcccaa	tgggcgtatc	12180
aagcacatac	ccaggccttc	ttcaaaacat	gacctaccat	gactcgaccg	tcgccgaaat	12240
cgccaacgaa	ccacttggcg	caacagcatc	acgcctgcgc	tcggaactca	acagtgatcg	12300
tttgcgcaga	cgaacacaag	ctttggcgac	gtacatgcat	ggcctgcctg	acaagtcgag	12360
cgtctccctg	accgccgatg	cgaatccgtc	aagcagcatc	atgctgagtt	cctgggccaa	12420
ggtgggatgc	tgggagtatg	actttgggtt	tggactgggt	aagcctgaga	gtgtgagaag	12480
acctcgcttt	gaaccttttg	agagtttgat	gtactttatg	ccaagaagc	ctgatgggga	12540
gtttacggcg	tccatttctc	tgagggatga	ggatatggag	agactaaagg	cggatgagga	12600
gtggacaaag	tacgcaaagt	atattgggta	gatagtttac	tagactactg	caggatatcg	12660
tggatccccg	aatttccccg	atcggtcaaa	catttggcaa	taaagtttct	taagattgaa	12720
tcctgttgcc	ggtcttgcca	tgattatcat	ataatttctg	ttgaattacg	ttaagcatgt	12780
aataattaac	atgtaatgca	tgacgttatt	tatgagatgg	gtttttatga	ttagagtccc	12840
gcaattatac	atttaatacg	cgatagaaaa	caaatatag	cgcgcaaact	aggataaatt	12900
atcgcgcgcg	gtgtcatcta	tgttactaga	tcgggaattc	ggcgcgcca		12949

30041-231

CLAIMS:

1. A plant cell comprising a heterologous polynucleotide encoding a trichothecene 3-O-acetyltransferase that is expressed in the plant cell, said polynucleotide comprising:
 - a) the nucleotide sequence of SEQ ID NO: 1 or a sequence having at least 65% sequence identity to SEQ ID NO: 1; or
 - b) a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2 or an amino acid sequence having at least 65% identity to said amino acid sequence;said plant cell being resistant to a trichothecene comprising a C-3 hydroxyl group.
2. The plant cell of claim 1, wherein the plant cell is resistant to a fungus that produces a trichothecene comprising a C-3 hydroxyl group.
3. The plant cell of claim 2, wherein the plant cell is resistant to a fungus of the genus *Fusarium*.
4. The plant cell of claim 3, wherein the plant cell is resistant to *Fusarium graminearum*.
5. The plant cell of claim 1, wherein the heterologous polynucleotide is a microbial polynucleotide.
6. The plant cell of claim 5, wherein the heterologous polynucleotide is a yeast or fungal polynucleotide.

30041-231

7. The plant cell of claim 6, wherein the fungal polynucleotide is a *Fusarium* polynucleotide.
8. The plant cell of claim 7, wherein the fungal polynucleotide is a *Fusarium graminearum* or *Fusarium sporotrichioides* polynucleotide.
9. The plant cell of any one of claims 1 to 8, wherein the trichothecene is selected from the group consisting of T-2 toxin, HT-2 toxin, isotrichodermol, 4,15-diacetoxyscripenol (DAS), 3-deacetylcalonectrin, 3,15-dideacetylcalonectrin, scirpentriol, neosolaniol; type B: 15-acetyldeoxynivalenol, nivalenol, 4-acetylnivalenol (fusarenone-X), 4,15-diacetylnivalenol, 4,7,15-acetylnivalenol, and deoxynivalenol (DON).
10. The plant cell of claim 9, wherein the trichothecene is DON or DAS.
11. The plant cell according to any one of claims 1 to 10, wherein the plant cell is a wheat, maize, barley or rice plant cell.
12. A method for producing a trichothecene resistant plant comprising the steps of:
 - a) transforming a plant cell with a heterologous polynucleotide encoding a trichothecene 3-O-acetyltransferase that increases resistance to a trichothecene comprising a C-3 hydroxyl group, when compared to a wild type control plant, said heterologous polynucleotide comprising:
 - i) the nucleotide sequence of SEQ ID NO: 1 or a sequence having at least 65% sequence identity to SEQ ID NO: 1; or
 - ii) a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2 or an amino acid sequence having at least 65% identity to said amino acid

30041-231

sequence;

- b) expressing the trichothecene 3-O-acetyltransferase at a biologically significant level;
 - c) regenerating the plant cell into a plant; and
 - d) selecting a plant with increased resistance to a trichothecene when compared to a wild type control plant.
13. The method of claim 12 further comprising the step of selecting a plant on which growth of fungus, wherein the fungus produces a trichothecene virulence factor comprising a C-3 hydroxyl group, is reduced when compared to a wild type control plant.
14. The method of claim 13, wherein the fungus is of the genus *Fusarium*.
15. A method of preventing contamination of a plant, including of a plant's seed, with a trichothecene comprising a C-3 hydroxyl group, comprising growing a plant derived from a plant cell of any one of claims 1 to 11 or a plant produced by a method according to claims 12 to 14.
16. A method for reducing growth of a fungus when compared to a wild type control plant and/or preventing the growth of a fungus that produces a trichothecene comprising a C-3 hydroxyl group, on a plant, comprising growing a plant derived from a plant cell of any one of claims 1 to 11 or a plant produced by a method according to claims 12 to 14.
17. The method of claim 15 or claim 16, wherein the plant is grown in an area with moderate to severe fungal infestation.

30041-231

18. The method of any one of claims 12 to 17, wherein the plant is a crop plant.
19. A method of producing a fungal resistant plant comprising
- a) transforming a plant cell with a heterologous polynucleotide encoding a trichothecene 3-O-acetyltransferase that increases resistance to a trichothecene comprising a C-3 hydroxyl group compared to a wild type control plant, said heterologous polynucleotide comprising:
 - i) the nucleotide sequence of SEQ ID NO: 1 or a sequence having at least 65% sequence identity to SEQ ID NO: 1; or
 - ii) a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2 or an amino acid sequence having at least 65% identity to said amino acid sequence;
 - b) expressing the trichothecene 3-O-acetyltransferase at a biologically significant level;
 - c) regenerating the plant cell into a plant; and
 - d) selecting a plant with increased resistance to a trichothecene when compared to a wild type control plant.
20. The method of claim 19, which further comprises selfing or outcrossing the plant obtained in step (d).
21. The method of claim 19 or claim 20, wherein the fungus is of the genus *Fusarium*.
22. A method of producing seed wherein the plant grown from the seed is resistant to a fungus that produces a trichothecene comprising a C-3 hydroxyl group, comprising selfing or outcrossing a plant derived from a plant cell of any one of claims 1 to 11

30041-231

or a plant produced by a method according to claims 19 to 21 and collecting the seed, wherein said seed comprises a plant cell of any one of claims 1 to 11.

23. The method of claim 22, wherein the seed is
5 resistant to fungi to the genus *Fusarium*.

FETHERSTONHAUGH & CO.

OTTAWA, CANADA

PATENT AGENTS