



(51) International Patent Classification: Not classified

(21) International Application Number:
PCT/US2010/054846(22) International Filing Date:
29 October 2010 (29.10.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/255,980 29 October 2009 (29.10.2009) US(71) Applicant (for all designated States except US):
TRUSTEES OF DARTMOUTH COLLEGE [US/US];
11 Rope Ferry Road, #6210, Hanover, NH 03755-1404
(US).

(72) Inventor; and


(75) Inventor/Applicant (for US only): **SENTMAN, Charles,**
L. [US/US]; 11 Pearl St., West Lebanon, NH 03784 (US).(74) Agents: **TESKIN, Robin, L.** et al.; Hunton & Williams,
Intellectual Property Department, 1900 K Street, N.W.,
Washington, DC 20006 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,
ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report (Rule 48.2(g))

(54) Title: T CELL RECEPTOR-DEFICIENT T CELL COMPOSITIONS

Figure 1



(57) Abstract: The invention is directed to modified T cells, methods of making and using isolated, modified T cells, and methods of using these isolated, modified T cells to address diseases and disorders. In one embodiment, this invention broadly relates to TCR-deficient T cells, isolated populations thereof, and compositions comprising the same. In another embodiment of the invention, these TCR-deficient T cells are designed to express a functional non-TCR receptor. The invention also pertains to methods of making said TCR-deficient T cells, and methods of reducing or ameliorating, or preventing or treating, diseases and disorders using said TCR-deficient T cells, populations thereof, or compositions comprising the same.



T CELL RECEPTOR-DEFICIENT T CELL COMPOSITIONS

[0001] This invention was made with government support under contract number CA 130911 awarded by the National Institutes of Health. The government has certain rights in the invention.

CROSS-REFERENCE TO RELATED APPLICATIONS

[0002] This application claims the benefit of priority to U.S. provisional patent application no. 61/255,980, filed October 29, 2009, the disclosure of which is herein incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

[0003] The invention is directed to TCR-deficient T cells, methods of making and using TCR-deficient T cells, and methods of using these TCR-deficient T cells to address diseases and disorders. In one embodiment, the invention broadly relates to TCR-deficient T cells, isolated populations thereof, and compositions comprising the same. In another embodiment of the invention, said TCR-deficient T cells are further designed to express a functional non-TCR receptor. The invention also pertains to methods of making said TCR-deficient T cells, and methods of reducing or ameliorating, or preventing or treating, diseases and disorders using said TCR-deficient T cells, populations thereof, or compositions comprising the same.

Description of Related Art

[0004] The global burden of cancer doubled between 1975 and 2000, and cancer is expected to become the leading cause of death worldwide by 2010. According to the American Cancer Society, it is projected to double again by 2020 and to triple by 2030. Thus, there is a need for more effective therapies to treat various forms of cancer. Ideally, any cancer therapy should be effective (at killing cancerous cells), targeted (i.e. selective, to

avoid killing healthy cells), permanent (to avoid relapse and metastasis), and affordable. Today's standards of care for most cancers fall short in some or all of these criteria.

[0005] Cellular immunotherapy has been shown to result in specific tumor elimination and has the potential to provide specific and effective cancer therapy (Ho, W.Y. et al. 2003. *Cancer Cell* 3:1318-1328; Morris, E.C. et al. 2003. *Clin. Exp. Immunol.* 131:1-7; Rosenberg, S.A. 2001. *Nature* 411:380-384; Boon, T. and P. van der Bruggen. 1996. *J. Exp. Med.* 183:725-729). T cells have often been the effector cells of choice for cancer immunotherapy due to their selective recognition and powerful effector mechanisms. T cells recognize specific peptides derived from internal cellular proteins in the context of self-major histocompatibility complex (MHC) using their T cell receptors (TCR).

[0006] It is recognized in the art that the TCR complex associates in precise fashion by the formation of dimers and association of these dimers (TCR-alpha/beta, CD3-gamma/epsilon, CD3-delta/epsilon, and CD3-zeta dimer) into one TCR complex that can be exported to the cell surface. The inability of any of these complexes to form properly will inhibit TCR assembly and expression (Call, M.E. et al., (2007) *Nature Rev. Immunol.*, 7:841-850; Call, M.E. et al., (2005) *Annu. Rev. Immunol.*, 23:101-125).

[0007] Particular amino acid residues in the respective TCR chains have been identified as important for proper dimer formation and TCR assembly. In particular, for TCR-alpha, these key amino acids in the transmembrane portion are arginine (for association with CD3-zeta) and lysine (for association with the CD3-epsilon/delta dimer). For TCR-beta, the key amino acid in the transmembrane portion is a lysine (for association with CD3-epsilon/gamma dimer). For CD3-gamma, the key amino acid in the transmembrane portion is a glutamic acid. For CD3-delta, the key amino acid in the transmembrane portion is an aspartic acid. For CD3-epsilon, the key amino acid in the transmembrane portion is an aspartic acid. For CD3-zeta, the key amino acid in the transmembrane portion is an aspartic acid (Call, M.E. et al., (2007) *Nature Rev. Immunol.*, 7:841-850; Call, M.E. et al., (2005) *Annu. Rev. Immunol.*, 23:101-125).

[0008] Peptides derived from altered or mutated proteins in tumors can be recognized by specific TCRs. Several key studies have led to the identification of antigens associated with specific tumors that have been able to induce effective cytotoxic T lymphocyte (CTL)

responses in patients (Ribas, A. et al. 2003. *J. Clin. Oncol.* 21:2415-2432). T cell effector mechanisms include the ability to kill tumor cells directly and the production of cytokines that activate other host immune cells and change the local tumor microenvironment. Theoretically, T cells could identify and destroy a tumor cell expressing a single mutated peptide. Adoptive immunotherapy with CTL clones specific for MART1 or gp100 with low dose IL-2 has been effective in reduction or stabilization of tumor burden in some patients (Yee, C. et al. 2002. *Proc. Natl. Acad. Sci. USA* 99:16168-16173). Other approaches use T cells with a defined anti-tumor receptor. These approaches include genetically modifying CTLs with new antigen-specific T cell receptors that recognize tumor peptides and MHC, chimeric antigen receptors (CARs) derived from single chain antibody fragments (scFv) coupled to an appropriate signaling element, or the use of chimeric NK cell receptors (Ho, W.Y. et al. 2003. *Cancer Cell* 3:431-437; Eshhar, Z. et al. 1993. *Proc. Natl. Acad. Sci. USA* 90:720-724; Maher, J. and E.T. Davies. 2004. *Br. J. Cancer* 91:817-821; Zhang, T. et al. 2005. *Blood* 106:1544-1551).

[0009] Cell-based therapies are used in patients who have failed conventional chemotherapy or radiation treatments, or have relapsed, often having attempted more than one type of therapy. The immune cells from patients with advanced cancer, who may have gone through rounds of chemotherapy, do not respond as robustly as healthy individuals. Moreover, cancer patients are often elderly and may suffer from other diseases that may limit the potential of their immune cells to become primed effector cells, even after *in vitro* activation and expansion. In addition, each cancer patient must provide a sufficient number of their own immune cells in order for them to be engineered to express a new immune receptor. Because each therapy must be custom made for the patient, this process requires weeks from the time the decision to undertake such therapy is made; meanwhile, the cancer continues to grow. U.S. patent application publication no. US 2002/0039576 discloses a method for modulating T cell activity, where the T cells used have a phenotype of CD3⁺-αβ-TcR⁺CD4⁻CD8⁻CD28⁻NK1.1⁻. U.S. patent application publication no. US 2006/0166314 discloses use of mutated T cells to treat cancer where the T cells are ones with a T cell response-mediating MDM2 protein-specific αβ-T cell receptor.

[0010] Cancer is not the only disease wherein T cell manipulation could be effective therapy. It is known that active T cell receptors on T cells are critical to the response of the body to stimulate immune system activity. For example, it has been shown that T cell receptor diversity plays a role in graft-versus-host-disease (GVHD), in particular chronic GVHD (Anderson *et al.* (2004) *Blood* 104:1565-1573). In fact, administration of T cell receptor antibodies has been shown to reduce the symptoms of acute GVHD (Maeda *et al.* (2005) *Blood* 106:749-755).

[0011] There remains a need for more effective T cell-based therapies for the treatment of certain diseases and disorders, and methods of treatment based on the design of new types of T cells.

BRIEF SUMMARY OF THE INVENTION

[0012] In one embodiment, this invention broadly relates to isolated, modified T cells that do not express a functional T cell receptor (TCR). In this embodiment, the T cells are TCR-deficient in the expression of a functional TCR. In another embodiment of the invention, TCR-deficient T cells are engineered to express a functional non-TCR receptor, such as for example a chimeric receptor. These cells also function as a platform to allow the expression of other targeting receptors, receptors that may be useful in specific diseases, while retaining the effector functions of T cells, albeit without a functioning TCR.

[0013] The invention contemplates populations of TCR-deficient T cells, and compositions comprising the same. The invention also contemplates methods of making said TCR-deficient T cells, and methods of reducing or ameliorating, or preventing or treating, diseases and disorders using said TCR-deficient T cells, populations thereof, or therapeutic compositions comprising the same. In one embodiment, this composition can be used to treat cancer, infection, one or more autoimmune disorders, radiation sickness, or to prevent or treat graft versus host disease (GVHD) or transplantation rejection in a subject undergoing transplant surgery.

BRIEF DESCRIPTION OF THE DRAWING

[0014] Figure 1 illustrates chimeric NK receptors described herein. Extracellular (EC), transmembrane (TM), and cytoplasmic (Cyp) portions are indicated. Wild-type (WT) and chimeric (CH) forms of the receptors are indicated, wherein NH₂ denotes the N-terminus and COOH denotes the C-terminus.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Definitions

[0015] It is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, animal species or genera, and reagents described, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

[0016] As used herein the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a plurality of such cells and reference to "the protein" includes reference to one or more proteins and equivalents thereof known to those skilled in the art, and so forth. All technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs unless clearly indicated otherwise.

[0017] In the context of the present invention, by a "TCR-deficient T cell", or a similar phrase is intended an isolated T cell(s) that lacks expression of a functional TCR, is internally capable of inhibiting its own TCR production, and further wherein progeny of said T cell(s) may also be reasonably expected to be internally capable of inhibiting their own TCR production. Internal capability is important in the context of therapy where TCR turnover timescales (~ hours) are much faster than demonstrable efficacy timescales (days-months), i.e., internal capability is required to maintain the desired phenotype during the therapeutic period. This may e.g., be accomplished by different means as described *infra*, e.g., by

engineering a T cell such that it does not express any functional TCR on its cell surface, or by engineering the T cell such that it does not express one or more of the subunits that comprise a functional TCR and therefore does not produce a functional TCR or by engineering a T cell such that it produces very little functional TCR on its surface, or which expresses a substantially impaired TCR, e.g by engineering the T cell to express mutated or truncated forms of one or more of the subunits that comprise the TCR, thereby rendering the T cell incapable of expressing a functional TCR or resulting in a cell that expresses a substantially impaired TCR. The different subunits that comprise a functional TCR are described *infra*. Whether a cell expresses a functional TCR may be determined using known assay methods such as are known in the art described herein. By a “substantially impaired TCR” applicants mean that this TCR will not substantially elicit an adverse immune reaction in a host, e.g., a GVHD reaction.

[0018] As described in detail *infra*, optionally these TCR-deficient cells may be engineered to comprise other mutations or transgenes that e.g., mutations or transgenes that affect T cell growth or proliferation, result in expression or absence of expression of a desired gene or gene construct, e.g., another receptor or a cytokine or other immunomodulatory or therapeutic polypeptide or a selectable marker such as a dominant selectable marker gene, e.g., DHFR or neomycin transferase.

[0019] “Allogeneic T cell” refers to a T cell from a donor having a tissue HLA type that matches the recipient. Typically, matching is performed on the basis of variability at three or more loci of the HLA gene, and a perfect match at these loci is preferred. In some instances allogeneic transplant donors may be related (usually a closely HLA matched sibling), *syngeneic* (a monozygotic 'identical' twin of the patient) or unrelated (donor who is not related and found to have very close degree of HLA matching). The HLA genes fall in two categories (Type I and Type II). In general, mismatches of the Type-I genes (i.e. HLA-A, HLA-B, or HLA-C) increase the risk of graft rejection. A mismatch of an HLA Type II gene (i.e. HLA-DR, or HLA-DQB1) increases the risk of graft-versus-host disease.

[0020] In the context of the present invention, a “bank of tissue matched TCR-deficient T cells” refers to different compositions each containing T cells of a specific HLA allotype which are rendered TCR-deficient according to the invention. Ideally this bank will

comprise compositions containing T cells of a wide range of different HLA types that are representative of the human population. Such a bank of engineered TCR-deficient T cells will be useful as it will facilitate the availability of T cells suitable for use in different recipients such as cancer patients.

[0021] In the context of the present invention, a “therapeutically effective amount” is identified by one of skill in the art as being an amount of TCR-deficient T cells that, when administered to a patient, alleviates the signs and or symptoms of the disease (e.g., cancer, infection or GVHD). The actual amount to be administered can be determined based on studies done either *in vitro* or *in vivo* where the functional TCR-deficient T cells exhibit pharmacological activity against disease. For example, the functional TCR-deficient T cells may inhibit tumor cell growth either *in vitro* or *in vivo* and the amount of functional TCR-deficient T cells that inhibits such growth is identified as a therapeutically effective amount.

[0022] A “pharmaceutical composition” refers to a chemical or biological composition suitable for administration to a mammal. Such compositions may be specifically formulated for administration *via* one or more of a number of routes, including but not limited to buccal, intraarterial, intracardial, intracerebroventricular, intradermal, intramuscular, intraocular, intraperitoneal, intraspinal, intrathecal, intravenous, oral, parenteral, rectally *via* an enema or suppository, subcutaneous, subdermal, sublingual, transdermal, and transmucosal. In addition, administration can occur by means of injection, liquid, gel, drops, or other means of administration.

[0023] As used herein, a nucleic acid construct or nucleic acid sequence is intended to mean a DNA molecule which can be transformed or introduced into a T cell and be transcribed and translated to produce a product (e.g., a chimeric receptor or a suicide protein).

[0024] Nucleic acids are “operably linked” when placed into a functional relationship with another nucleic acid sequence. For example, DNA for a signal sequence is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and

in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites or alternatively via a PCR/recombination method familiar to those skilled in the art (Gateway[®] Technology; Invitrogen, Carlsbad California). If such sites do not exist, the synthetic oligonucleotide adapters or linkers are used in accordance with conventional practice.

[0025] The invention contemplates compositions and methods for reducing or ameliorating, or preventing or treating, diseases or conditions such as cancer, infectious disease, GVHD, transplantation rejection, one or more autoimmune disorders, or radiation sickness. In a non-limiting embodiment, the compositions are based on the concept of providing an allogeneic source of isolated human T cells, namely TCR-deficient T cells, that can be manufactured in advance of patient need and inexpensively. The ability to create a single therapeutic product at a single site using processes that are well controlled is attractive in terms of both cost and quality considerations. The change from an autologous to an allogeneic source for T cells offers significant advantages. For example, it has been estimated that a single healthy donor could supply T cells sufficient to treat dozens of patients after transduction and expansion.

[0026] According to the present invention, modified allogeneic T cells are produced that do not express functional T cell receptors (TCRs). It is to be understood that some, or even all, of the TCR subunits/dimers may be expressed on the cell surface, but that the T cell does not express enough functional TCR to induce an undesirable reaction in the host. Without functional TCRs on their surface, the allogeneic T cells fail to mount an undesired immune response to host cells. As a result, these TCR-deficient T cells fail to cause GVHD, for example, as they cannot recognize the host MHC molecules. Additionally, these TCR-deficient T cells can be engineered to simultaneously express functional, non-TCR, disease-specific receptors.

[0027] As is well known to one of skill in the art, various methods are readily available for isolating allogeneic T cells from a subject. For example, using cell surface marker expression or using commercially available kits (e.g., ISOCELL[™] from Pierce, Rockford, Ill.).

[0028] For cancer therapy, the approach encompasses producing an isolated pool of TCR-deficient T effector cells, e.g., of a desired tissue allotype that do not express a functional form of their endogenous TCR or which express substantially reduced levels of endogenous TCR compared to wild type T cells such that they do not elicit an immune response upon administration (such as GVHD), but instead express a functional, non-TCR receptor that recognizes tumor cells, or express another polypeptide that does not appreciably, or at all, attack non-disease associated cells, e.g., normal (non-tumorigenic) cells that do not express the antigen or ligand recognized by the tumor specific receptor or which express said antigen or ligand at reduced levels relative to tumor cells. It is understood by those skilled in the art that certain tumor-associated antigens are expressed in non-cancerous tissues, but they are viable therapeutic targets in a tumor-bearing host. With respect thereto it is generally understood by those skilled in the art that certain non-TCR, tumor-specific receptors are expressed in non-cancerous tissues, but are viable therapeutic targets in a tumor-bearing host as they may be expressed at significantly reduced levels in normal than tumor cells.

[0029] While not necessary for most therapeutic usages of the subject TCR-deficient T cells, in some instances it may be desirable to remove some or all of the donor T cells from the host shortly after they have mediated their anti-tumor effect. This may be facilitated by engineering the T cells to express additional receptors or markers that facilitate their removal and/or identification in the host such as GFP and the like. While the present invention should substantially eliminate any possibility of GVHD or other adverse immune reaction in the recipient this may be desired in some individuals. This should not compromise efficacy as it has already been shown that donor T cells do not need to remain long in the host for a long-term anti-tumor effect to be initiated (Zhang, T., *et al.* 2007. *Cancer Res.* 67:11029-11036; Barber, A. *et al.* 2008. *J. Immunol.* 180:72-78).

[0030] In one embodiment of the invention, nucleic acid constructs introduced into engineered T cells further contains a suicide gene such as thymidine kinase (TK) of the HSV virus (herpesvirus) type I (Bonini, *et al.* (1997) *Science* 276:1719-1724), a Fas-based "artificial suicide gene" (Thomis, *et al.* (2001) *Blood* 97:1249-1257), or E. coli cytosine deaminase gene which are activated by gancyclovir, AP1903, or 5-fluorocytosine, respectively. The suicide gene is advantageously included in the nucleic acid construct of the

present invention to provide for the opportunity to ablate the transduced T cells in case of toxicity and to destroy the chimeric construct once a tumor has been reduced or eliminated. The use of suicide genes for eliminating transformed or transduced cells is well-known in the art. For example, Bonini, *et al.* ((1997) Science 276:1719-1724) teach that donor lymphocytes transduced with the HSV-TK suicide gene provide antitumor activity in patients for up to one year and elimination of the transduced cells is achieved using ganciclovir. Further, Gonzalez, *et al.* ((2004) J. Gene Med. 6:704-711) describe the targeting of neuroblastoma with cytotoxic T lymphocyte clones genetically modified to express a chimeric scFvFc:zeta immunoreceptor specific for an epitope on L1-CAM, wherein the construct further expresses the hygromycin thymidine kinase (HyTK) suicide gene to eliminate the transgenic clones.

[0031] It is contemplated that the suicide gene can be expressed from the same promoter as the shRNA, minigene, or non-TCR receptor, or from a different promoter. Generally, however, nucleic acid sequences encoding the suicide protein and shRNA, minigene, or non-TCR receptor reside on the same construct or vector. Expression of the suicide gene from the same promoter as the shRNA, minigene, or non-TCR receptor can be accomplished using any well-known internal ribosome entry site (IRES). Suitable IRES sequences which can be used in the nucleic acid construct of the present invention include, but are not limited to, IRES from EMCV, c-myc, FGF-2, poliovirus and HTLV-1. By way of illustration only, a nucleic acid construct for expressing a chimeric receptor can have the following structure: promoter->chimeric receptor->IRES->suicidal gene. Alternatively, the suicide gene can be expressed from a different promoter than that of the chimeric receptor (e.g., promoter 1->chimeric receptor->promoter 2->suicidal gene).

[0032] Because of the broad application of T cells for cell therapies, and the improved nature of the T cells of the invention, the present invention encompasses any method or composition wherein T cells are therapeutically desirable. Such compositions and methods include those for reducing or ameliorating, or preventing or treating cancer, GVHD, transplantation rejection, infection, one or more autoimmune disorders, radiation sickness, or other diseases or conditions that are based on the use of T cells derived from an allogeneic source that lack expression of functional TCR.

[0033] As indicated, further embodiments of the invention embrace recombinant expression of receptors in said TCR-deficient T cells, such as chimeric NKG2D, chimeric Fv domains, NKG2D, or any other receptor to initiate signals to T cells, thereby creating potent, specific effector T cells. One of skill in the art can select the appropriate receptor to be expressed by the TCR-deficient T cell based on the disease to be treated. For example, receptors that can be expressed by the TCR-deficient T cell for treatment of cancer would include any receptor to a ligand that has been identified on cancer cells. Such receptors include, but are not limited to, NKG2D, NKG2A, NKG2C, NKG2F, LLT1, AICL, CD26, NKRP1, NKp30, NKp44, NKp46, CD244 (2B4), DNAM-1, and NKp80.

[0034] In another embodiment of the invention, such receptors include, but not limited to, chimeric receptors comprising a ligand binding domain obtained from NKG2D, NKG2A, NKG2C, NKG2F, LLT1, AICL, CD26, NKRP1, NKp30, NKp44, NKp46, CD244 (2B4), DNAM-1, and NKp80, or an anti-tumor antibody such as anti-Her2neu or anti-EGFR, and a signaling domain obtained from CD3-zeta, Dap10, CD28, 41BB, and CD40L. In one embodiment of the invention, the chimeric receptor binds MIC-A, MIC-B, Her2neu, EGFR, mesothelin, CD38, CD20, CD19, PSA, MUC1, MUC2, MUC3A, MUC3B, MUC4, MUC5AC, MUC5B, MUC6, MUC7, MUC8, MUC12, MUC13, MUC15, MUC16, MUC17, MUC19, MUC20, estrogen receptor, progesterone receptor, RON, or one or more members of the ULBP/RAET1 family including ULBP1, ULBP2, ULBP3, ULBP4, ULBP5, and ULBP6.

[0035] In the methods of the present invention a patient suffering from cancer, GVHD, transplantation rejection, infection, one or more autoimmune disorders, or radiation sickness is administered a therapeutically effective amount of a composition comprising said TCR-deficient T cells. In another embodiment of the invention, a therapeutically effective amount of a composition comprising said TCR-deficient T cells is administered to prevent, treat, or reduce GVHD, transplantation rejection, or cancer.

Methods of Producing TCR-deficient T-cells

[0036] T cells stably lacking expression of a functional TCR according to the invention may be produced using a variety of approaches. T cells internalize, sort, and degrade the

entire T cell receptor as a complex, with a half-life of about 10 hours in resting T cells and 3 hours in stimulated T cells (von Essen, M. et al. 2004. *J. Immunol.* 173:384-393). Proper functioning of the TCR complex requires the proper stoichiometric ratio of the proteins that compose the TCR complex. TCR function also requires two functioning TCR zeta proteins with ITAM motifs. The activation of the TCR upon engagement of its MHC-peptide ligand requires the engagement of several TCRs on the same T cell, which all must signal properly. Thus, if a TCR complex is destabilized with proteins that do not associate properly or cannot signal optimally, the T cell will not become activated sufficiently to begin a cellular response.

[0037] In one embodiment of the invention, TCR expression is eliminated using small-hairpin RNAs (shRNAs) that target nucleic acids encoding specific TCRs (*e.g.*, TCR- α and TCR- β) and/or CD3 chains in primary T cells. By blocking expression of one or more of these proteins, the T cell will no longer produce one or more of the key components of the TCR complex, thereby destabilizing the TCR complex and preventing cell surface expression of a functional TCR. Even though some TCR complexes can be recycled to the cell surface, the shRNA will prevent new production of TCR proteins resulting in degradation and removal of the entire TCR complex, resulting in the production of a T cell having a stable deficiency in functional TCR expression.

[0038] Expression of shRNAs in primary T cells can be achieved using any conventional expression system, *e.g.*, a lentiviral expression system. Although lentiviruses are useful for targeting resting primary T cells, not all T cells will express the shRNAs. Some of these T cells may not express sufficient amounts of the shRNAs to allow enough inhibition of TCR expression to alter the functional activity of the T cell. Thus, T cells that retain moderate to high TCR expression after viral transduction can be removed, *e.g.*, by cell sorting or separation techniques, so that the remaining T cells are deficient in cell surface TCR or CD3, enabling the expansion of an isolated population of T cells deficient in expression of functional TCR or CD3.

[0039] In a non-limiting embodiment of the invention, exemplary targeting shRNAs have been designed for key components of the TCR complex as set forth below (Table 1).

Table 1

Target	Target base	shRNA Sequence	GC%	SEQ ID NO:
TCR- β	18 ^a	AGTGCGAGGAGATTTCGGCAGCTTAT	52	1
	21 ^a	GCGAGGAGATTTCGGCAGCTTATTTC	52	2
	48 ^a	CCACCATCCTCTATGAGATCTTGCT	48	3
	54 ^a	TCCTCTATGAGATCTTGCTAGGGAA	44	4
TCR- α	3 ^b	TCTATGGCTTCAACTGGCTAGGGTG	52	5
	76 ^b	CAGGTAGAGGCCTTGTCACCTAAT	52	6
	01 ^b	GCAGCAGACACTGCTTCTTACTTCT	48	7
	07 ^b	GACACTGCTTCTTACTTCTGTGCTA	44	8
CD3- ϵ	89 ^c	CCTCTGCCTCTTATCAGTTGGCGTT	52	9
	27 ^c	GAGCAAAGTGGTTATTATGTCTGCT	40	10
	62 ^c	AAGCAAACCAGAAGATGCGAACTTT	40	11
	45	GACCTGTATTCTGGCCTGAATCAGA	48	12
		GGCCTCTGCCTCTTATCAGTT	52	13
		GCCTCTGCCTCTTATCAGTTG	52	14
		GCCTCTTATCAGTTGGCGTTT	48	15
		AGGATCACCTGTCACTGAAGG	52	16
		GGATCACCTGTCACTGAAGGA	52	17
		GAATTGGAGCAAAGTGGTTAT	38	18
		GGAGCAAAGTGGTTATTATGT	38	19
		GCAAACCAGAAGATGCGAACT	48	20
		ACCTGTATTCTGGCCTGAATC	48	21
		GCCTGAATCAGAGACGCATCT	52	22
		CTGAAATACTATGGCAACACAATGATAAA	31	23
		AAACATAGGCAGTGATGAGGATCACCTGT	45	24
		ATTGTCATAGTGGACATCTGCATCACTGG	45	25
		CTGTATTCTGGCCTGAATCAGAGACGCAT	48	26
CD3- δ ^d		GATACCTATAGAGGAACTTGA	38	27
		GACAGAGTGTTTGTGAATTGC	43	28
		GAACACTGCTCTCAGACATTA	43	29
		GGACCCACGAGGAATATATAG	48	30
		GGTGTAATGGGACAGATATAT	38	31
		GCAAGTTCATTATCGAATGTG	38	32
		GGCTGGCATCATTGTCACTGA	52	33
		GCTGGCATCATTGTCACTGAT	48	34
		GCATCATTGTCACTGATGTCA	43	35

		GCTTTGGGAGTCTTCTGCTTT	48	36
		TGGAACATAGCACGTTTCTCTCTGGCCTG	52	37
		CTGCTCTCAGACATTACAAGACTGGACCT	48	38
		ACCGTGGCTGGCATCATTGTCACTGATGT	52	39
		TGATGCTCAGTACAGCCACCTTGGAGGAA	52	40
CD3- γ^c		GGCTATCATTCTTCTTCAAGG	43	41
		GCCCAGTCAATCAAAGGAAAC	48	42
		GGTTAAGGTGTATGACTATCA	38	43
		GGTTCGGTACTTCTGACTTGT	48	44
		GAATGTGTCAGAACTGCATTG	43	45
		GCAGCCACCATATCTGGCTTT	52	46
		GGCTTTCTCTTTGCTGAAATC	43	47
		GCTTTCTCTTTGCTGAAATCG	43	48
		GCCACCTTCAAGGAAACCAGT	52	49
		GAAACCAGTTGAGGAGGAATT	43	50
		GGCTTTCTCTTTGCTGAAATCGTCAGCAT	45	51
		AGGATGGAGTTCGCCAGTCGAGAGCTTCA	55	52
		CCTCAAGGATCGAGAAGATGACCAGTACA	48	53
		TACAGCCACCTTCAAGGAAACCAGTTGAG	48	54

^a With reference to Accession No. EU030678.

^b With reference to Accession No. AY247834.

^c With reference to Accession No. NM_000733.

^d With reference to Accession No. NM_000732.

^e With reference to Accession No. NM_000073.

[0040] TCR-alpha, TCR-beta, TCR-gamma, TCR-delta, CD3-gamma, CD3-delta, CD3-epsilon, or CD3-zeta mRNAs can be targeted separately or together using a variety of targeting shRNAs. The TCR- β and TCR- α chains are composed of variable and constant portions. Several targeting shRNAs have been designed for the constant portions of these TCR/CD3 sequences. One or a combination of shRNAs can be used for each molecular target to identify the most efficient inhibitor of TCR expression. Using established protocols, each shRNA construct is cloned into, *e.g.*, a pLko.1 plasmid, with expression controlled by a promoter routinely used in the art, *e.g.*, the U6p promoter. The resulting construct can be screened and confirmed for accuracy by sequencing. The shRNA expression plasmid can then be transfected into any suitable host cell (*e.g.*, 293T), together with a packaging plasmid and an envelope plasmid for packaging. Primary human peripheral blood mononuclear cells (PBMCs) are isolated from healthy donors and activated with low dose soluble anti-CD3 and

25U/ml rhIL-2 for 48 hours. Although it is not required to activate T cells for lentiviral transduction, transduction works more efficiently and allows the cells to continue to expand in IL-2. The activated cells are washed and transduced, *e.g.*, using a 1 hour spin-infection at 30°C, followed by a 7 hour resting period.

[0041] In another embodiment of the invention, over-expression of a dominant-negative inhibitor protein is capable of interrupting TCR expression or function. In this embodiment of the invention, a minigene that incorporates part, or all, of a polynucleotide encoding for one of the TCR components (*e.g.*, TCR-alpha, TCR-beta, CD3-gamma, CD3-delta, CD3-epsilon, or CD3-zeta) is prepared, but is modified so that: (1) it lacks key signaling motifs (*e.g.* an ITAM) required for protein function; (2) is modified so it does not associate properly with its other natural TCR components; or (3) can associate properly but does not bind ligands (*e.g.* a truncated TCR beta minigene).

[0042] These minigenes may also encode a portion of a protein that serves as a means to identify the over-expressed minigene. For example, polynucleotides encoding a truncated CD19 protein, which contains the binding site for anti-CD19 mAbs, can be operably linked to the minigene so that the resulting cell that expresses the minigene will express the encoded protein and can be identified with anti-CD19 mAbs. This identification enables one to determine the extent of minigene expression and isolate cells expressing this protein (and thus lack a functional TCR).

[0043] In one embodiment of the invention, over-expression of a minigene lacking a signaling motif(s) lead to a TCR complex that cannot signal properly when the TCR is engaged by its MHC-peptide ligand on an opposing cell. In a non-limiting embodiment of the invention, high expression of this minigene (and the encoded polypeptide) outcompetes the natural complete protein when the TCR components associate, resulting in a TCR complex that cannot signal. In another embodiment of the invention, the minigene comprises, or alternatively consists of, a polynucleotide encoding full or partial CD3-zeta, CD3-gamma, CD3-delta, or CD3-epsilon polypeptides lacking the ITAM motifs required for signaling. The CD3-zeta protein contains three ITAM motifs in the cytoplasmic portion, and in one embodiment of the invention, removal of all of these through truncation inhibits proper TCR signaling in any complexes where this modified protein is incorporated. The

construct may incorporate ITIM or other signaling motifs, which are known to alter cell signaling and promote inhibitory signals through the recruitment of phosphatases such as SHP1 and SHP2.

[0044] In another embodiment of the invention, over-expression of a minigene is modified so that the encoded polypeptide can associate with some, but not all, of its natural partners, creating competition with the normal protein for those associating proteins. In another non-limiting hypothesis of the invention, high level expression of the minigene (and the encoded polypeptide) outcompetes the natural partner proteins and prevents assembly of a functional TCR complex, which requires all components to associate in the proper ratios and protein-protein interactions. In another embodiment of the invention, minigenes comprise, or alternatively consist of, all or part of the polynucleotides encoding full-length proteins (e.g., TCR-alpha, TCR-beta, CD3-gamma, CD3-delta, CD3-epsilon, or CD3-zeta), but containing selected deletions in the sequence coding for amino acids in the transmembrane portion of the protein that are known to be required for assembly with other TCR/CD3 proteins.

[0045] In a preferred embodiment of the invention, selected deletions in the sequence coding for amino acids in the transmembrane portion of the protein known to be required for assembly with other TCR/CD3 proteins include, but are not limited to: the arginine residue at position 5 in the TCR-alpha transmembrane region; the lysine residue at position 10 in the TCR-alpha transmembrane region; the lysine residue at position 9 in the TCR-beta transmembrane region; the glutamic acid residue in the transmembrane region of CD3-gamma; the aspartic acid residue in the transmembrane region of CD3-delta-epsilon; the aspartic acid residue in the transmembrane region of CD3-epsilon; and the aspartic acid residue in the transmembrane region of CD3-zeta.

[0046] Over-expression of a truncated TCR-alpha, TCR-beta, TCR-gamma, or TCR-delta protein results in a TCR complex that cannot bind to MHC-peptide ligands, and thus will not function to activate the T cell. In another embodiment of the invention, minigenes comprise, or alternatively consist of, polynucleotides encoding the entire cytoplasmic and transmembrane portions of these proteins and portions of the extracellular region, but lacks polynucleotides encoding all or part of the first extracellular domain (i.e., the most outer

domain containing the ligand binding site). In a preferred embodiment, said minigene polynucleotides do not encode Valpha and Vbeta polypeptides of the TCR-alpha and TCR-beta chains. In one embodiment, the minigene polynucleotides may be operably linked to polynucleotides encoding a protein epitope tag (e.g. CD19), thereby allowing mAb identification of cells expressing these genes.

[0047] In another embodiment, these minigenes can be expressed using a strong viral promoter, such as the 5'LTR of a retrovirus, or a CMV or SV40 promoter. Typically, this promoter is immediately upstream of the minigene and leads to a high expression of the minigene mRNA. In another embodiment, the construct encodes a second polynucleotide sequence under the same promoter (using for example an IRES DNA sequence between) or another promoter. This second polynucleotide sequence may encode for a functional non-TCR receptor providing specificity for the T cell. Examples of this polynucleotide include, but are not limited to, chimeric NKG2D, chimeric NKp30, chimeric NKp46, or chimeric anti-Her2neu. In a further embodiment, promoter-minigenes are constructed into a retroviral or other suitable expression plasmid and transfected or transduced directly into T cells using standard methods (Zhang, T. *et al.*, (2006) *Cancer Res.*, 66(11) 5927-5933; Barber, A. *et al.*, (2007) *Cancer Res.*, 67(10):5003-5008).

[0048] After viral transduction and expansion using any of the methods discussed previously, any T cells that still express TCR/CD3 are removed using anti-CD3 mAbs and magnetic beads using Miltenyi selection columns as previously described (Barber, A. *et al.*, (2007) *Cancer Res.*, 67(10):5003-5008). The T cells are subsequently washed and cultured in IL-2 (25U/ml) for 3 to 7 days to allow expansion of the effector cells in a similar manner as for use of the cells *in vivo*.

[0049] The expression of TCR $\alpha\beta$ and CD3 can be evaluated by flow cytometry and quantitative real-time PCR (QRT-PCR). Expression of TCR- α , TCR- β , CD3 ϵ , CD3- ζ , and GAPDH (as a control) mRNA can be analyzed by QRT-PCR using an ABI7300 real-time PCR instrument and gene-specific TAQMAN® primers using methods similar to those used in Sentman, C.L. *et al.* ((2004) *J. Immunol.* 173:6760-6766). Changes in cell surface expression can be determined using antibodies specific for TCR- α , TCR- β , CD3 ϵ , CD8, CD4, CD5, and CD45.

[0050] It is possible that a single shRNA species may not sufficiently inhibit TCR expression on the cell surface. In this case, multiple TCR shRNAs may be used simultaneously to target multiple components of the TCR complex. Each component is required for TCR complex assembly at the cell surface, so a loss of one of these proteins can result in loss of TCR expression at the cell surface. While some or even all TCR expression may remain, it is the receptor function which determines whether the receptor induces an immune response. The functional deficiency, rather than complete cell surface absence, is the critical measure. In general, the lower the TCR expression, the less likely sufficient TCR cross-linking can occur to lead to T cell activation *via* the TCR complex. While particular embodiments embrace the targeting of TCR-alpha, TCR-beta, and CD3-epsilon, other components of the TCR complex, such as CD3-gamma, CD3-delta, or CD3-zeta, can also be targeted.

[0051] The primary aim of removing the TCR from the cell surface is to prevent the activation of the T cell to incompatible MHC alleles. To determine whether the reduction in TCR expression with each shRNA or minigene construct is sufficient to alter T cell function, the T cells can be tested for: (1) cell survival *in vitro*; (2) proliferation in the presence of mitomycin C-treated allogeneic PBMCs; and (3) cytokine production in response to allogeneic PBMCs, anti-CD3 mAbs, or anti-TCR mAbs.

[0052] To test cell survival, transduced T cells are propagated in complete RPMI medium with rhuIL-2 (25U/ml). Cells are plated at similar densities at the start of culture and a sample may be removed for cell counting and viability daily for 7 or more days. To determine whether the T cells express sufficient TCR to induce a response against allogeneic cells, transduced or control T cells are cultured with mitomycin C-treated allogeneic or syngeneic PBMCs. The T cells are preloaded with CFSE, which is a cell permeable dye that divides equally between daughter cells after division. The extent of cell division can be readily determined by flow cytometry. Another hallmark of T cell activation is production of cytokines. To determine whether each shRNA construct inhibits T cell function, transduced T cells are cultured with anti-CD3 mAbs (25 ng/ml). After 24 hours, cell-free supernatants are collected and the amount of IL-2 and IFN- γ produced is quantified by ELISA.

PMA/ionomycin are used as a positive control to stimulate the T cells and T cells alone are used as a negative control.

[0053] It is possible that removal of TCR-alpha or TCR-beta components may allow the preferential expansion of TCR-gamma/delta T cells. These T cells are quite rare in the blood, however the presence of these cells can be determined with anti-TCR-gamma/delta antibodies. If there is an outgrowth of these cells, the targeting of CD3-epsilon, which is required for cell surface expression of both TCR-alpha/beta and TCR-gamma/delta at the cell surface, can be used. Both IL-2 and IFN- γ are key effector cytokines that drive T cell expansion and macrophage activation. Therefore, lack of production of these cytokines is a sign of functional inactivation. It is also possible to measure changes in other cytokines, such as TNF- α . Any reduction in T cell survival upon elimination of TCR expression can be determined by culturing the TCR-deficient T cells with PBMCs, which better reflects the *in vivo* environment and provides support for T cell survival.

[0054] In another embodiment of the invention, The T cells stably deficient in functional TCR expression express a functional, non-TCR receptor. In this embodiment, the removal of TCR function (as described previously) is further combined with expression of one or more exogenous non-TCR targeting receptors (such as for example chimeric NKG2D (chNKG2D) or Fv molecules). This embodiment provides "universal" cell products, which can be stored for future therapy of any patient with any type of cancer, provided a suitable targeting receptor is employed.

[0055] Further embodiments of the invention embrace recombinant expression of receptors in said TCR-deficient T cells, such as chNKG2D, chimeric Fv domains, NKG2D, or any other receptor to initiate signals to T cells, thereby creating potent, specific effector T cells. One of skill in the art can select the appropriate receptor to be expressed by the TCR-deficient T cell based on the disease to be treated. For example, receptors that can be expressed by the TCR-deficient T cell for treatment of cancer would include any receptor to a ligand that has been identified on cancer cells. Such receptors include, but are not limited to, NKG2D (GENBANK accession number BC039836), NKG2A (GENBANK accession number AF461812), NKG2C (GENBANK accession number AJ001684), NKG2F, LLT1, AICL, CD26, NKRP1, NKp30 (e.g., GENBANK accession number AB055881), NKp44

(e.g., GENBANK accession number AJ225109), NKp46 (e.g., GENBANK accession number AJ001383), CD244 (2B4), DNAM-1, and NKp80.

[0056] In another embodiment of the invention, such receptors include, but not limited to, chimeric receptors comprising a ligand binding domain obtained from NKG2D, NKG2A, NKG2C, NKG2F, LIT1, AICL, CD26, NKRP1, NKp30, NKp44, NKp46, CD244 (2B4), DNAM-1, and NKp80, or an anti-tumor antibody, such as anti-Her2neu and anti-EGFR, and a signaling domain obtained from CD3-zeta (CD3 ζ) (e.g., GENBANK accession number human NM__198053), Dap10 (e.g., GENBANK accession number AF072845), CD28, 41BB, and/or CD40L.

[0057] In a further embodiment of the invention, the chimeric receptor binds MIC-A, MIC-B, Her2neu, EGFR, mesothelin, CD38, CD20, CD19, PSA, MUC1, MUC2, MUC3A, MUC3B, MUC4, MUC5AC, MUC5B, MUC6, MUC7, MUC8, MUC12, MUC13, MUC15, MUC16, MUC17, MUC19, MUC20, estrogen receptor, progesterone receptor, RON, or one or more members of the ULBP/RAET1 family including ULBP1, ULBP2, ULBP3, ULBP4, ULBP5, and ULBP6.

[0058] By way of illustration only, shRNAs or minigenes shown to eliminate cell surface expression of the TCR complex are co-expressed with the chNKG2D receptor *via* one or more viral vectors. To achieve co-expression in one vector, the shRNA can be driven by a U6 promoter and the chNKG2D receptor by a PGK promoter. In another embodiment, if an IRES sequence is used to separate the genetic elements then only one promoter is used.

[0059] A C-type lectin-like NK cell receptor protein particularly suitable for use in the chimeric receptor includes a receptor expressed on the surface of natural killer cells, wherein upon binding to its cognate ligand(s) it alters NK cell activation. The receptor can work alone or in concert with other molecules. Ligands for these receptors are generally expressed on the surface of one or more tumor cell types, e.g., tumors associated with cancers of the colon, lung, breast, kidney, ovary, cervix, and prostate; melanomas; myelomas; leukemias; and lymphomas (Wu, *et al.* (2004) *J. Clin. Invest.* 114:60-568; Groh, *et al.* (1999) *Proc. Natl. Acad. Sci. USA* 96:6879-6884; Pende, *et al.* (2001) *Eur. J. Immunol.* 31:1076-1086) and are not widely expressed on the surface of cells of normal tissues.

[0060] Examples of such ligands include, but are not limited to, MIC-A, MIC-B, heat shock proteins, ULBP binding proteins (e.g., ULPBs 1-4), and non-classical HLA molecules such as HLA-E and HLA-G, whereas classical MHC molecules such as HLA-A, HLA-B, or HLA-C and alleles thereof are not generally considered strong ligands of the C-type lectin-like NK cell receptor protein of the present invention. C-type lectin-like NK cell receptors which bind these ligands generally have a type II protein structure, wherein the N-terminal end of the protein is intracellular. In addition to any NK cell receptors previously listed above, further exemplary NK cell receptors of this type include, but are not limited to, Dectin-1 (GENBANK accession number AJ312373 or AJ312372), Mast cell function-associated antigen (GENBANK accession number AF097358), HNKRP-1A (GENBANK accession number U11276), LLT1 (GENBANK accession number AF133299), CD69 (GENBANK accession number NM__001781), CD69 homolog, CD72 (GENBANK accession number NM__001782), CD94 (GENBANK accession number NM__002262 or NM__007334), KLRF1 (GENBANK accession number NM__016523), Oxidised LDL receptor (GENBANK accession number NM__002543), CLEC-1, CLEC-2 (GENBANK accession number NM__016509), NKG2D (GENBANK accession number BC039836), NKG2C (GENBANK accession number AJ001684), NKG2A (GENBANK accession number AF461812), NKG2E (GENBANK accession number AF461157), WUGSC:H_DJ0701016.2, or Myeloid DAP12-associating lectin (MDL-1; GENBANK accession number AJ271684). In a preferred embodiment of the invention, the NK cell receptor is human NKG2D (SEQ ID NO:58) or human NKG2C (SEQ ID NO:59).

[0061] Similar type I receptors which would be useful in the chimeric receptor include NKp46 (GENBANK accession number AJ001383), NKp30 (GENBANK accession number AB055881), or NKp44 (GENBANK accession number AJ225109).

[0062] As an alternative to the C-type lectin-like NK cell receptor protein, a protein associated with a C-type lectin-like NK cell receptor protein can be used in the chimeric receptor protein. In general, proteins associated with C-type lectin-like NK cell receptor are defined as proteins that interact with the receptor and transduce signals therefrom. Suitable human proteins which function in this manner further include, but are not limited to, DAP10

(e.g., GENBANK accession number AF072845)(SEQ ID NO:60), DAP12 (e.g., GENBANK accession number AF019562)(SEQ ID NO:61) and FcR gamma.

[0063] To the N-terminus of the C-type lectin-like NK cell receptor is fused an immune signaling receptor having an immunoreceptor tyrosine-based activation motif (ITAM), (Asp/Glu)-Xaa-Xaa-Tyr*-Xaa-Xaa-(Ile/Leu)-Xaa₆₋₈-Tyr*-Xaa-Xaa-(Ile/Leu) (SEQ ID NOS: 55-57) which is involved in the activation of cellular responses *via* immune receptors. Similarly, when employing a protein associated with a C-type lectin-like NK cell receptor, an immune signaling receptor can be fused to the C-terminus of said protein (FIG. 1). Suitable immune signaling receptors for use in the chimeric receptor of the present invention include, but are not limited to, the zeta chain of the T-cell receptor, the eta chain which differs from the zeta chain only in its most C-terminal exon as a result of alternative splicing of the zeta mRNA, the delta, gamma and epsilon chains of the T-cell receptor (CD3 chains) and the gamma subunit of the FcR1 receptor. In particular embodiments, in addition to immune signaling receptors identified previously, the immune signaling receptor is CD3-zeta (CD3ζ) (e.g., GENBANK accession number human NM__198053)(SEQ ID NO:62), or human Fc epsilon receptor-gamma chain (e.g., GENBANK accession number M33195)(SEQ ID NO:63) or the cytoplasmic domain or a splicing variant thereof.

[0064] In particular embodiments, a chimeric receptor of the present invention is a fusion between NKG2D and CD3-zeta, or Dap10 and CD3-zeta.

[0065] In the nucleic acid construct of the present invention, the promoter is operably linked to the nucleic acid sequence encoding the chimeric receptor of the present invention, i.e., they are positioned so as to promote transcription of the messenger RNA from the DNA encoding the chimeric receptor. The promoter can be of genomic origin or synthetically generated. A variety of promoters for use in T cells are well-known in the art (e.g., the CD4 promoter disclosed by Marodon, *et al.* (2003) Blood 101(9):3416-23). The promoter can be constitutive or inducible, where induction is associated with the specific cell type or a specific level of maturation. Alternatively, a number of well-known viral promoters are also suitable. Promoters of interest include the β-actin promoter, SV40 early and late promoters, immunoglobulin promoter, human cytomegalovirus promoter, retrovirus promoter, and the Friend spleen focus-forming virus promoter. The promoters may or may not be associated

with enhancers, wherein the enhancers may be naturally associated with the particular promoter or associated with a different promoter.

[0066] The sequence of the open reading frame encoding the chimeric receptor can be obtained from a genomic DNA source, a cDNA source, or can be synthesized (e.g., *via* PCR), or combinations thereof. Depending upon the size of the genomic DNA and the number of introns, it may be desirable to use cDNA or a combination thereof as it is found that introns stabilize the mRNA or provide T cell-specific expression (Barthel and Goldfeld (2003) *J. Immunol.* 171(7):3612-9). Also, it may be further advantageous to use endogenous or exogenous non-coding regions to stabilize the mRNA.

[0067] For expression of a chimeric receptor of the present invention, the naturally occurring or endogenous transcriptional initiation region of the nucleic acid sequence encoding N-terminal component of the chimeric receptor can be used to generate the chimeric receptor in the target host. Alternatively, an exogenous transcriptional initiation region can be used which allows for constitutive or inducible expression, wherein expression can be controlled depending upon the target host, the level of expression desired, the nature of the target host, and the like.

[0068] Likewise, the signal sequence directing the chimeric receptor to the surface membrane can be the endogenous signal sequence of N-terminal component of the chimeric receptor. Optionally, in some instances, it may be desirable to exchange this sequence for a different signal sequence. However, the signal sequence selected should be compatible with the secretory pathway of T cells so that the chimeric receptor is presented on the surface of the T cell.

[0069] Similarly, a termination region can be provided by the naturally occurring or endogenous transcriptional termination region of the nucleic acid sequence encoding the C-terminal component of the chimeric receptor. Alternatively, the termination region can be derived from a different source. For the most part, the source of the termination region is generally not considered to be critical to the expression of a recombinant protein and a wide variety of termination regions can be employed without adversely affecting expression.

[0070] As will be appreciated by one of skill in the art, in some instances, a few amino acids at the ends of the C-type lectin-like natural killer cell receptor (or protein associated

therewith) or immune signaling receptor can be deleted, usually not more than 10, more usually not more than 5 residues. Also, it may be desirable to introduce a small number of amino acids at the borders, usually not more than 10, more usually not more than 5 residues. The deletion or insertion of amino acids will usually be as a result of the needs of the construction, providing for convenient restriction sites, ease of manipulation, improvement in levels of expression, or the like. In addition, the substitute of one or more amino acids with a different amino acid can occur for similar reasons, usually not substituting more than about five amino acids in any one domain.

[0071] The chimeric construct, which encodes the chimeric receptor can be prepared in conventional ways. Since, for the most part, natural sequences are employed, the natural genes are isolated and manipulated, as appropriate (e.g., when employing a Type II receptor, the immune signaling receptor component may have to be inverted), so as to allow for the proper joining of the various components. Thus, the nucleic acid sequences encoding for the N-terminal and C-terminal proteins of the chimeric receptor can be isolated by employing the polymerase chain reaction (PCR), using appropriate primers which result in deletion of the undesired portions of the gene. Alternatively, restriction digests of cloned genes can be used to generate the chimeric construct. In either case, the sequences can be selected to provide for restriction sites which are blunt-ended, or have complementary overlaps.

[0072] The various manipulations for preparing the chimeric construct can be carried out *in vitro* and in particular embodiments the chimeric construct is introduced into vectors for cloning and expression in an appropriate host using standard transformation or transfection methods. Thus, after each manipulation, the resulting construct from joining of the DNA sequences is cloned, the vector isolated, and the sequence screened to insure that the sequence encodes the desired chimeric receptor. The sequence can be screened by restriction analysis, sequencing, or the like.

[0073] It is contemplated that the chimeric construct can be introduced into T cells as naked DNA or in a suitable vector. Methods of stably transfecting T cells by electroporation using naked DNA are known in the art. *See*, e.g., U.S. Pat. No. 6,410,319. Naked DNA generally refers to the DNA encoding a chimeric receptor of the present invention contained in a plasmid expression vector in proper orientation for expression. Advantageously, the use

of naked DNA reduces the time required to produce T cells expressing the chimeric receptor of the present invention.

[0074] Alternatively, a viral vector (e.g., a retroviral vector, adenoviral vector, adeno-associated viral vector, or lentiviral vector) can be used to introduce the chimeric construct into T cells. Suitable vectors for use in accordance with the method of the present invention are non-replicating in the subject's T cells. A large number of vectors are known which are based on viruses, where the copy number of the virus maintained in the cell is low enough to maintain the viability of the cell. Illustrative vectors include the pFB-neo vectors (STRATAGENETM) as well as vectors based on HIV, SV40, EBV, HSV or BPV. Once it is established that the transfected or transduced T cell is capable of expressing the chimeric receptor as a surface membrane protein with the desired regulation and at a desired level, it can be determined whether the chimeric receptor is functional in the host cell to provide for the desired signal induction (e.g., production of Rantes, Mip1-alpha, GM-CSF upon stimulation with the appropriate ligand).

[0075] As described above, primary human PBMCs are isolated from healthy donors and activated with low-dose soluble anti-CD3 and rhIL-2, anti-CD3/anti-CD28 beads and rhIL-2, or irradiated antigen presenting cells and rhIL-2. Although it is not required to activate T cells for lentiviral transduction, transduction is more efficient and the cells continue to expand in IL-2. The activated cells are washed and transduced as described herein, followed by a resting period. The cells are washed and cultured in IL-2 for 3 to 7 days to allow expansion of the effector cells in a similar manner as for use of the cells *in vivo*.

[0076] The expression of TCR $\alpha\beta$, CD3, and NKG2D can be evaluated by flow cytometry and quantitative QRT-PCR as discussed herein. The number of CD4⁺ and CD8⁺ T cells can also be determined. Overall cell numbers and the percentage of TCR complex-deficient, TCR-competent, and chNKG2D-expressing T cells can be determined by flow cytometry. These numbers can be compared to PBMCs that have been transduced with the shRNA or chNKG2D genes alone (as controls). Vector-only transduced cells can also be included as controls.

[0077] After viral transduction and expansion, the TCR⁺ and TCR⁻ cells can be separated by mAbs with magnetic beads over Miltenyi columns and TCR-deficient T cells expressing

the chNKG2D receptor are identified and isolated. For example, chNKG2D expression can be verified by QRT-PCR using specific primers for the chNKG2D receptor (Zhang, T. *et al.* (2007) *Cancer Res.* 67:11029-11036; Barber, A. *et al.* (2008) *J. Immunol.* 180:72-78). Function of these TCR-deficient chNKG2D⁺ cells can be determined by culturing the cells with allogeneic PBMCs or tumor cells that express NKG2D ligands. T cell proliferation and cytokine production (IFN- γ and IL-2) can be determined by flow cytometry and ELISA, respectively. To determine whether the T cells that have lost TCR function and retained chNKG2D function, transduced or control T cells will be cultured with anti-CD3 (25 ng/ml) or mitomycin C-treated allogeneic PBMCs, or syngeneic PBMCs. The extent of cytokine production (IFN- γ and IL-2) can be determined by ELISA. The T cells can be preloaded with CFSE, which is a cell permeable dye that divides equally between daughter cells after division. The extent of cell division can be readily determined by flow cytometry.

[0078] Another hallmark of T cell activation is production of cytokines. To determine whether TCR-deficient chNKG2D⁺ cells induce T cell activation, the T cells are cocultured with mitomycin C-treated allogeneic PBMCs, syngeneic PBMCs, or tumor cells: P815-MICA (a murine tumor expressing human MICA, a ligand for NKG2D), P815, A2008 (a human ovarian tumor cell, NKG2D ligand⁺), and U266 (a human myeloma cell line, NKG2D ligand⁺). After 24 hours, cell-free supernatants are collected and the amount of IL-2 and IFN- γ produced is quantified by ELISA. T cells alone and culture with syngeneic PBMCs are used as a negative control.

[0079] Subsequently, the transduced T cells are reintroduced or administered to the subject to activate anti-tumor responses in said subject. To facilitate administration, the transduced T cells according to the invention can be made into a pharmaceutical composition or made implant appropriate for administration in vivo, with appropriate carriers or diluents, which further can be pharmaceutically acceptable. The means of making such a composition or an implant have been described in the art (see, for instance, Remington's Pharmaceutical Sciences, 16th Ed., Mack, ed. (1980)). Where appropriate, the transduced T cells can be formulated into a preparation in semisolid or liquid form, such as a capsule, solution, injection, inhalant, or aerosol, in the usual ways for their respective route of administration. Means known in the art can be utilized to prevent or minimize release and absorption of the

composition until it reaches the target tissue or organ, or to ensure timed-release of the composition. Desirably, however, a pharmaceutically acceptable form is employed which does not ineffectuate the cells expressing the chimeric receptor. Thus, desirably the transduced T cells can be made into a pharmaceutical composition containing a balanced salt solution, preferably Hanks' balanced salt solution, or normal saline.

Methods of Ameliorating or Reducing Symptoms of, or Treating, or Preventing, Diseases and Disorders Using TCR-deficient T-cells

[0080] The invention is also directed to methods of reducing or ameliorating, or preventing or treating, diseases and disorders using the TCR-deficient T cells described herein, isolated populations thereof, or therapeutic compositions comprising the same. In one embodiment, the TCR-deficient T cells described herein, isolated populations thereof, or therapeutic compositions comprising the same are used to reduce or ameliorate, or prevent or treat, cancer, infection, one or more autoimmune disorders, radiation sickness, or to prevent or treat graft versus host disease (GVHD) or transplantation rejection in a subject undergoing transplant surgery.

[0081] The TCR-deficient T cells described herein, isolated populations thereof, or therapeutic compositions comprising the same are useful in altering autoimmune or transplant rejection because these effector cells can be grown in TGF- β during development and will differentiate to become induced T regulatory cells. In one embodiment, the functional non-TCR is used to give these induced T regulatory cells the functional specificity that is required for them to perform their inhibitory function at the tissue site of disease. Thus, a large number of antigen-specific regulatory T cells are grown for use in patients. The expression of FoxP3, which is essential for T regulatory cell differentiation, can be analyzed by flow cytometry, and functional inhibition of T cell proliferation by these T regulatory cells can be analyzed by examining decreases in T cell proliferation after anti-CD3 stimulation upon co-culture.

[0082] Another embodiment of the invention is directed to the use of TCR-deficient T cells described herein, isolated populations thereof, or therapeutic compositions comprising the same for the prevention or treatment of radiation sickness. One challenge after radiation

treatment or exposure (e.g. dirty bomb exposure, radiation leak) or other condition that ablates bone marrow cells (certain drug therapies) is to reconstitute the hematopoietic system. In patients undergoing a bone marrow transplant, the absolute lymphocyte count on day 15 post-transplant is correlated with successful outcome. Those patients with a high lymphocyte count reconstitute well, so it is important to have a good lymphocyte reconstitution. The reason for this effect is unclear, but it may be due to lymphocyte protection from infection and/or production of growth factors that favors hematopoietic reconstitution.

[0083] In this embodiment, TCR-deficient T cells described herein, isolated populations thereof, or therapeutic compositions comprising the same result in the production of a large number of T cells that are unable to respond to allogeneic MHC antigens. Hence, these T cells may be used to reconstitute people and offer protection from infection, leading to faster self-reconstitution of people suffering from full or partial bone marrow ablation due to radiation exposure. In the event of a catastrophic or unexpected exposure to high doses of radiation, TCR-deficient T cells described herein having another functional receptor, isolated populations thereof, or therapeutic compositions comprising the same can be infused rapidly into patients to offer some reconstitution of their immune response and growth factor production for days to weeks until their own hematopoietic cells have reconstituted themselves, or until the person has been treated with an additional source of hematopoietic stem cells (e.g. a bone marrow transplant).

[0084] One of skill would understand how to treat cancer, infection, transplantation rejection, one or more autoimmune disorders, radiation sickness, or GVHD based on their experience with use of other types of T cells.

[0085] In addition to the illustrative TCR-deficient chNKG2D⁺ T cells described herein, it is contemplated that TCR-deficient T cells can be modified or developed to express other functional receptors useful in treatment of diseases such as cancer or infection as described previously. Briefly, the treatment methods of the invention contemplate the use of TCR-deficient T cells expressing functional non-TCR receptors, such as chNKG2D, chimeric Fv domains, NKG2D, or any other receptor to initiate signals to T cells, thereby creating potent, specific effector T cells. One of skill in the art can select the appropriate receptor to be

expressed by the TCR-deficient T cell based on the disease to be treated. For example, receptors that can be expressed by the TCR-deficient T cell for treatment of cancer would include any receptor that binds to a ligand that has been identified on cancer cells. Such receptors include, but are not limited to, NKG2D, NKG2A, NKG2C, NKG2F, LIT1, AICL, CD26, NKRP1, NKp30, NKp44, NKp46, CD244 (2B4), DNAM-1, and NKp80.

[0086] In another embodiment of the invention, such receptors include, but not limited to, chimeric receptors comprising a ligand binding domain obtained from NKG2D, NKG2A, NKG2C, NKG2F, LIT1, AICL, CD26, NKRP1, NKp30, NKp44, NKp46, CD244 (2B4), DNAM-1, and NKp80, or an anti-tumor antibody such as anti-Her2neu and anti-EGFR, and a signaling domain obtained from CD3zeta, Dap10, CD28, 41BB, and CD40L.

[0087] In a further embodiment of the invention, the chimeric receptor binds MIC-A, MIC-B, Her2neu, EGFR, mesothelin, CD38, CD20, CD19, PSA, MUC1, MUC2, MUC3A, MUC3B, MUC4, MUC5AC, MUC5B, MUC6, MUC7, MUC8, MUC12, MUC13, MUC15, MUC16, MUC17, MUC19, MUC20, estrogen receptor, progesterone receptor, RON, or one or more members of the ULBP/RAET1 family including ULBP1, ULBP2, ULBP3, ULBP4, ULBP5, and ULBP6.

[0088] Also embraced by the present invention are TCR-deficient T cells that express a non-TCR pathogen-associated receptor and the use of the TCR-deficient T cells expressing the pathogen receptor to treat or prevent infectious disease. In this embodiment, the non-TCR receptor binds to virus antigen or viral-induced antigen found on the surface of an infected cell. The infection to be prevented or treated, for example may be caused by a virus, bacteria, protozoa, or parasite. Viruses which can be treated include, but are not limited to, HCMV, EBV, hepatitis type A, hepatitis type B (HBV), hepatitis type C (HCV), ebola virus, VSV, influenza, varicella, adenovirus, herpes simplex type I (HSV-1), herpes simplex type 2 (HSV-2), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, cytomegalovirus (CMV), echinovirus, arbovirus, hantavirus, coxsackie virus, mumps virus, measles virus, rubella virus, polio virus, and/or human immunodeficiency virus type 1 or type 2 (HIV-1, HIV-2). Non-viral infections which can be treated using the TCR-deficient T cells include, but are not limited to, infectious *Staphylococcus* sp., *Enterococcus* sp., *Bacillus anthracis*, *Lactobacillus* sp., *Listeria* sp., *Corynebacterium diphtheriae*, *Nocardia*

sp., *Streptococcus* sp., *Pseudomonas* sp., *Gardnerella* sp., *Streptomyces* sp., *Thermoactinomyces vulgaris*, *Treponema* sp., *Campylobacter* sp., *Raeruginosa* sp., *Legionella* sp., *N. gonorrhoeae*, *N. meningitides*, *F. meningosepticum*, *F. odoratum*, *Brucella* sp., *B. pertussis*, *B. bronchiseptica*, *E. coli*, *Klebsiella*, *Enterobacter*, *S. marcescens*, *S. liquefaciens*, *Edwardsiella*, *P. mirabilis*, *P. vulgaris*, *Streptobacillus*, *R. fickettsii*, *C. psittaci*, *C. trachomatis*, *M. tuberculosis*, *M. intracellulare*, *M. fortuitum*, *M. laprae*, *M. avium*, *M. bovis*, *M. africanum*, *M. kansasii*, *M. lepraemurium*, trypanosomes, Chlamydia, or rickettsia.

[0089] Efficacy of the compositions of the present invention can be demonstrated in the most appropriate *in vivo* model system depending on the type of drug product being developed. The medical literature provides detailed disclosure on the advantages and uses of a wide variety of such models. For example, there are many different types of cancer models that are used routinely to examine the pharmacological activity of drugs against cancer such as xenograft mouse models (*e.g.*, Mattern, J. et al. 1988. *Cancer Metastasis Rev.* 7:263-284; Macor, P. et al. 2008. *Curr. Pharm. Des.* 14:2023-2039) or even the inhibition of tumor cell growth *in vitro*. In the case of GVHD, there are models in mice of both acute GVHD (*e.g.*, He, S. et al. 2008. *J. Immunol.* 181:7581-7592) and chronic GVHD (*e.g.*, Xiao, Z.Y. et al. 2007. *Life Sci.* 81:1403-1410).

[0090] Once the compositions of the present invention have been shown to be effective *in vivo* in animals, clinical studies may be designed based on the doses shown to be safe and effective in animals. One of skill in the art can design such clinical studies using standard protocols as described in textbooks such as Spilker (2000. *Guide to Clinical Trials*. Lippincott Williams & Wilkins: Philadelphia).

Administration

[0091] In one embodiment of the invention, the TCR-deficient T cells are administered to a recipient subject at an amount of between about 10^6 to 10^{11} cells. In a preferred embodiment of the invention, the TCR-deficient T cells are administered to a recipient subject at an amount of between 10^8 to 10^9 cells. In a preferred embodiment of the invention, the TCR-deficient T cells are administered to a recipient subject with a frequency of once

every twenty-six weeks or less, such as once every sixteen weeks or less, once every eight weeks or less, or once every four weeks or less.

[0092] These values provide general guidance of the range of transduced T cells to be utilized by the practitioner upon optimizing the method of the present invention for practice of the invention. The recitation herein of such ranges by no means precludes the use of a higher or lower amount of a component, as might be warranted in a particular application. For example, the actual dose and schedule can vary depending on whether the compositions are administered in combination with other pharmaceutical compositions, or depending on interindividual differences in pharmacokinetics, drug disposition, and metabolism. One skilled in the art readily can make any necessary adjustments in accordance with the exigencies of the particular situation.

[0093] A person of skill in the art would be able to determine an effective dosage and frequency of administration based on teachings in the art or through routine experimentation, for example guided by the disclosure herein and the teachings in Goodman, L. S., Gilman, A., Brunton, L. L., Lazo, J. S., & Parker, K. L. (2006). Goodman & Gilman's the pharmacological basis of therapeutics. New York: McGraw-Hill; Howland, R. D., Mycek, M. J., Harvey, R. A., Champe, P. C., & Mycek, M. J. (2006). Pharmacology. Lippincott's illustrated reviews. Philadelphia: Lippincott Williams & Wilkins; and Golan, D. E. (2008). Principles of pharmacology: the pathophysiologic basis of drug therapy. Philadelphia, Pa., [etc.]: Lippincott Williams & Wilkins. The dosing schedule can be based on well-established cell-based therapies (see, e.g., Topalian and Rosenberg (1987) *Acta Haematol.* 78 Suppl 1:75-6; U.S. Pat. No. 4,690,915) or an alternate continuous infusion strategy can be employed.

[0094] In another embodiment of the invention, the TCR-deficient T cells are administered to a subject in a pharmaceutical formulation.

[0095] In one embodiment of the invention, the TCR-deficient T cells may be optionally administered in combination with one or more active agents. Such active agents include analgesic, antipyretic, anti-inflammatory, antibiotic, antiviral, and anti-cytokine agents. Active agents include agonists, antagonists, and modulators of TNF- α , IL-2, IL-4, IL-6, IL-10, IL-12, IL-13, IL-18, IFN- α , IFN- γ , BAFF, CXCL13, IP-10, VEGF, EPO, EGF, HRG,

Hepatocyte Growth Factor (HGF), Hecidin, including antibodies reactive against any of the foregoing, and antibodies reactive against any of their receptors. Active agents also include 2-Arylpropionic acids, Aceclofenac, Acemetacin, Acetylsalicylic acid (Aspirin), Alclofenac, Alminoprofen, Amoxiprin, Ampyrone, Arylalkanoic acids, Azapropazone, Benorylate/Benorilate, Benoxaprofen, Bromfenac, Carprofen, Celecoxib, Choline magnesium salicylate, Clofezone, COX-2 inhibitors, Dexibuprofen, Dexketoprofen, Diclofenac, Diflunisal, Droxicam, Ethenzamide, Etodolac, Etoricoxib, Faislamine, fenamic acids, Fenbufen, Fenoprofen, Flufenamic acid, Flunoxaprofen, Flurbiprofen, Ibuprofen, Ibuprofen, Indometacin, Indoprofen, Kebuzone, Ketoprofen, Ketorolac, Lornoxicam, Loxoprofen, Lumiracoxib, Magnesium salicylate, Meclofenamic acid, Mefenamic acid, Meloxicam, Metamizole, Methyl salicylate, Mofebutazone, Nabumetone, Naproxen, N-Arylanthranilic acids, Nerve Growth Factor (NGF), Oxametacin, Oxaprozin, Oxicams, Oxyphenbutazone, Parecoxib, Phenazone, Phenylbutazone, Phenylbutazone, Piroxicam, Pirprofen, profens, Proglumetacin, Pyrazolidine derivatives, Rofecoxib, Salicyl salicylate, Salicylamide, Salicylates, Sulfinpyrazone, Sulindac, Suprofen, Tenoxicam, Tiaprofenic acid, Tolfenamic acid, Tolmetin, and Valdecoxib.

[0096] Antibiotics include Amikacin, Aminoglycosides, Amoxicillin, Ampicillin, Ansamycins, Arsphenamine, Azithromycin, Azlocillin, Aztreonam, Bacitracin, Carbacephem, Carbapenems, Carbenicillin, Cefaclor, Cefadroxil, Cefalexin, Cefalothin, Cefalotin, Cefamandole, Cefazolin, Cefdinir, Cefditoren, Cefepime, Cefixime, Cefoperazone, Cefotaxime, Cefoxitin, Cefpodoxime, Cefprozil, Ceftazidime, Ceftibuten, Ceftizoxime, Ceftobiprole, Ceftriaxone, Cefuroxime, Cephalosporins, Chloramphenicol, Cilastatin, Ciprofloxacin, Clarithromycin, Clindamycin, Cloxacillin, Colistin, Co-trimoxazole, Dalfopristin, Demeclocycline, Dicloxacillin, Dirithromycin, Doripenem, Doxycycline, Enoxacin, Ertapenem, Erythromycin, Ethambutol, Flucloxacillin, Fosfomycin, Furazolidone, Fusidic acid, Gatifloxacin, Geldanamycin, Gentamicin, Glycopeptides, Herbimycin, Imipenem, Isoniazid, Kanamycin, Levofloxacin, Lincomycin, Linezolid, Lomefloxacin, Loracarbef, Macrolides, Mafenide, Meropenem, Meticillin, Metronidazole, Mezlocillin, Minocycline, Monobactams, Moxifloxacin, Mupirocin, Nafcillin, Neomycin, Netilmicin, Nitrofurantoin, Norfloxacin, Ofloxacin, Oxacillin, Oxytetracycline, Paromomycin, Penicillin,

Penicillins, Piperacillin, Platensimycin, Polymyxin B, Polypeptides, Prontosil, Pyrazinamide, Quinolones, Quinupristin, Rifampicin, Rifampin, Roxithromycin, Spectinomycin, Streptomycin, Sulfacetamide, Sulfamethizole, Sulfanilimide, Sulfasalazine, Sulfisoxazole, Sulfonamides, Teicoplanin, Telithromycin, Tetracycline, Tetracyclines, Ticarcillin, Tinidazole, Tobramycin, Trimethoprim, Trimethoprim-Sulfamethoxazole, Troleandomycin, Trovafloxacin, and Vancomycin.

[0097] Active agents also include Aldosterone, Beclometasone, Betamethasone, Corticosteroids, Cortisol, Cortisone acetate, Deoxycorticosterone acetate, Dexamethasone, Fludrocortisone acetate, Glucocorticoids, Hydrocortisone, Methylprednisolone, Prednisolone, Prednisone, Steroids, and Triamcinolone. Any suitable combination of these active agents is also contemplated.

[0098] A “pharmaceutical excipient” or a “pharmaceutically acceptable excipient” is a carrier, usually a liquid, in which an active therapeutic agent is formulated. In one embodiment of the invention, the active therapeutic agent is a population of TCR-deficient T cells. In one embodiment of the invention, the active therapeutic agent is a population of TCR-deficient T cells expressing a functional, non-TCR receptor. The excipient generally does not provide any pharmacological activity to the formulation, though it may provide chemical and/or biological stability. Exemplary formulations can be found, for example, in Remington’s Pharmaceutical Sciences, 19th Ed., Grennaro, A., Ed., 1995 which is incorporated by reference.

[0099] As used herein “pharmaceutically acceptable carrier” or “excipient” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents that are physiologically compatible. In one embodiment, the carrier is suitable for parenteral administration. Alternatively, the carrier can be suitable for intravenous, intraperitoneal, intramuscular, or sublingual administration. Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions for the extemporaneous preparation of sterile injectable solutions or dispersions. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the

pharmaceutical compositions of the invention is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[0100] In a particularly preferred embodiment of the invention, appropriate carriers include, but are not limited to, Hank's Balanced Salt Solution (HBSS), Phosphate Buffered Saline (PBS), or any freezing medium having for example 10% DMSO and 90% human serum.

[0101] Pharmaceutical compositions typically must be sterile and stable under the conditions of manufacture and storage. The invention contemplates that the pharmaceutical composition is present in lyophilized form. The composition can be formulated as a solution. The carrier can be a dispersion medium containing, for example, water.

[0102] For each of the recited embodiments, the compounds can be administered by a variety of dosage forms. Any biologically-acceptable dosage form known to persons of ordinary skill in the art, and combinations thereof, are contemplated. Examples of such dosage forms include, without limitation, liquids, solutions, suspensions, emulsions, injectables (including subcutaneous, intramuscular, intravenous, and intradermal), infusions, and combinations thereof.

[0103] The above description of various illustrated embodiments of the invention is not intended to be exhaustive or to limit the invention to the precise form disclosed. While specific embodiments of, and examples for, the invention are described herein for illustrative purposes, various equivalent modifications are possible within the scope of the invention, as those skilled in the relevant art will recognize. The teachings provided herein of the invention can be applied to other purposes, other than the examples described above.

[0104] These and other changes can be made to the invention in light of the above detailed description. In general, in the following claims, the terms used should not be construed to limit the invention to the specific embodiments disclosed in the specification and the claims. Accordingly, the invention is not limited by the disclosure, but instead the scope of the invention is to be determined entirely by the following claims.

[0105] The invention may be practiced in ways other than those particularly described in the foregoing description and examples. Numerous modifications and variations of the

invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

[0106] Certain teachings related to T-cell receptor deficient T-cell compositions and methods of use thereof were disclosed in U.S. Provisional patent application no. 61/255,980, filed October 29, 2009, the disclosure of which is herein incorporated by reference in its entirety.

[0107] Certain teachings related to the production of T cells expressing chimeric receptors and methods of use thereof were disclosed in U.S. patent application publication no. US 2010/0029749, published February 4, 2010, the disclosure of which is herein incorporated by reference in its entirety.

[0108] Certain polynucleotide sequences useful in the production of T-cell receptor deficient T-cells of the invention are disclosed in the sequence listing accompanying this patent application filing, and the disclosure of said sequence listing is herein incorporated by reference in its entirety.

[0109] The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is herein incorporated by reference in their entireties.

[0110] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the subject invention, and are not intended to limit the scope of what is regarded as the invention. Efforts have been made to ensure accuracy with respect to the numbers used (*e.g.* amounts, temperature, concentrations, *etc.*) but some experimental errors and deviations should be allowed for. Unless otherwise indicated, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees centigrade; and pressure is at or near atmospheric.

EXAMPLES

Example 1: Production of T cell receptor (TCR)-deficient T cells

[0111] Minigenes are encoded on a retrovirus expression plasmid (e.g. pFB-neo or pSFG) containing 5' and 3' LTR sequences. The plasmids are packaged in a retroviral packaging cell line, such as PT67 or PG13, and viral particles are collected once the packaging cells have grown to confluence. T cells are then activated by PHA, anti-CD3, or anti-CD3/28 mAbs for 1 to 3 days in complete medium (or serum free medium) plus rIL-2 (25 U/ml), and T cells are transduced by spinoculation at 32°C in the presence of retronectin or polybrene. After resting for some 5 to 7 hours, the cells are washed and placed in fresh medium plus IL-2 for 2 to 7 days. Cells are counted periodically to avoid excessive cell concentration (i.e., $> 2 \times 10^6$ cells/ml) and re-plated at 7×10^5 cells/ml. Selection medium to remove non-transduced T cells is optionally used after 2 days for a period of 3 to 5 days. Live cells are harvested by Lymphoprep™ (Sentinel, Milan, Italy) gradient and further expanded for 1 to 3 days.

[0112] Following incubation, cells are analyzed for expression and function of the TCR. Functional non-TCR receptor expression may also be analyzed at this time, if appropriate. Flow cytometry is used to test for TCR/CD3 expression using fluorochrome-labeled antibodies. Live cells are stained with antibodies against CD5, CD8, and CD4, in combination with an antibody against CD3ε, TCRα, TCRβ, TCRγ, or TCRδ. If the expression of either the CD3 or TCR genes is used, the expression of both TCR proteins and CD3 proteins should be severely reduced compared to control vector treated T cells. Isotype control antibodies are used to control for background fluorescence. To identify T cells, cells are gated on CD5, then expression of CD4, CD8, CD3, and TCR is determined. Multiple samples are used for each treatment and appropriate compensation of fluorochrome emission spectra is used. The expression of another receptor (e.g. chNKG2D) is determined using specific antibodies and flow cytometry, as previously described in the art (Zhang, T. et al., (2006) Cancer Res., 66(11) 5927-5933; Barber, A. et al., (2007) Cancer Res., 67(10):5003-5008).

[0113] To test for functional deficiency of the TCR, anti-CD3 stimulation of effector cells is used at the end of culture to measure interferon (IFN)-gamma production after 24 hours. T cells (2×10^5) are cultured with soluble anti-CD3 (OKT3) mAbs in complete medium. After 24 hours, cell-free conditioned medium is collected and assayed by ELISA for IFN-gamma. Changes in TCR expression or function should be reflected in reduced IFN-gamma production.

[0114] To test for the function of the functional non-TCR, specific cytokine production by T cells incubated with tumor cells that do, or do not, express their specific ligand is used. For example, to test the function of chNKG2D, 10^5 T cells are incubated with 10^5 P815-MICA tumor cells (ligand+), 10^5 P815 (ligand-) cells, 10^5 RPMI8226 cells (ligand+) or T cells alone. After 24 hours, cell-free conditioned medium is collected and IFN-g measured by ELISA. Chimeric NKG2D T cells produce IFN-g after culture with ligand-expressing tumor cells (Zhang, T. et al., (2006) Cancer Res., 66(11) 5927-5933; Barber, A. et al., (2007) Cancer Res., 67(10):5003-5008). It is also possible to test cellular cytotoxicity against ligand+ tumor cells, as previously described in the art (Zhang, T. et al., (2006) Cancer Res., 66(11) 5927-5933). Specificity is shown using ligand-tumor cells or specific receptor blocking mAbs.

Example 2: Production of T cell receptor (TCR)-deficient T cells expressing chNKG2D

[0115] In this example, simultaneous expression of a chNKG2D receptor and inhibition of endogenous TCR expression is performed. In this example, a murine chNKG2D receptor is used, composed of NKG2D in combination with a N-terminally attached CD3-zeta. The chNKG2D receptor is generated and expressed in murine T-cells. NKG2D is a type II protein, in which the N-terminus is located intracellularly (Raulet (2003) Nat. Rev. Immunol. 3:781-790), whereas the CD3-zeta chain is type I protein with the C-terminus in the cytoplasm (Weissman, et al. (1988) Proc. Natl. Acad. Sci. USA 85:9709-9713). To generate a chimeric NKG2D-CD3-zeta fusion protein, an initiation codon ATG is placed ahead of the coding sequence for the cytoplasmic region of the CD3-zeta chain (without a stop codon TAA) followed by a wild-type NKG2D gene. Upon expression, the orientation of the CD3-

zeta portion is reversed inside the cells. The extracellular and transmembrane domains are derived from NKG2D. A second chimeric gene encoding the Dap10 gene followed by a fragment coding for the CD3-zeta cytoplasmic domain is also constructed. Figure 1 presents the structures of the chimeric and wild-type receptors.

[0116] An shRNA is operably linked in a lentiviral vector with the chNKG2D receptor. To achieve expression of both genes, the shRNA is driven by a U6 promoter and the chNKG2D receptor by a PGK promoter. Primary human PBMCs are isolated from healthy donors and activated with low-dose soluble anti-CD3 and 25U/ml rhuIL-2 for 48 hours. Although it is not required to activate T cells for lentiviral transduction, the transduction will work more efficiently and allow the cells to continue to expand in IL-2. The activated cells are washed and transduced using a 1 h spin-infection at 30°C, followed by a resting period for 7 h. The cells are washed and cultured in 25U/ml IL-2 for 3 to 7 d to allow expansion of the effector cells in a similar manner as we do for use of the cells *in vivo*. The expression of TCR $\alpha\beta$, CD3, and NKG2D is evaluated by flow cytometry and quantitative realtime-PCR (QRT-PCR). The number of CD4⁺ and CD8⁺ T cells are determined by flow cytometry. Overall cell numbers and the percentage of TCR complex deficient and expressing T cells are determined by flow cytometry. These are compared to PBMCs that are transduced with the shRNA or chNKG2D genes alone (as controls). Vector-only transduced cells are also included as controls.

[0117] It is anticipated that those cells with no or little TCR expression at the cell surface will express higher amounts of cell surface NKG2D because of co-expression of the chNKG2D receptor.

[0118] As an alternative, transduction may occur with two viruses at the same time, one with the shRNA construct and one with the chNKG2D receptor. A larger amount of the chNKG2D virus is used to ensure high expression of chNKG2D in those T cells that lack TCR expression. TCR⁺ T cells that may remain are removed to obtain TCR⁻, chNKG2D⁺ T cells.

[0119] After viral transduction and expansion, the TCR⁺ and TCR⁻ cells are separated by mAbs with magnetic beads over Miltenyi columns. Verification of chNKG2D expression is performed by QRT-PCR using specific primers for the chNKG2D receptor.

[0120] To determine whether the T cells have lost TCR function and retained chNKG2D function, transduced or control T cells are cultured with mitomycin C-treated allogeneic PBMCs or syngeneic PBMCs. The T cells are preloaded with CFSE, which is a cell permeable dye that divides equally between daughter cells after division. The extent of cell division can be easily determined by flow cytometry.

[0121] To determine whether the shRNA construct can inhibit TCR function and allow chNKG2D receptor function, transduced T cells are cultured with mitomycin C-treated allogeneic PBMCs, syngeneic PBMCs, or tumor cells: P815-MICA (a murine tumor expressing human MICA, a ligand for NKG2D), P815, A2008 (a human ovarian tumor cell, NKG2D ligand+), and U266 (a human myeloma cell line, NKG2D ligand+). After 48 hours, cell-free supernatants are collected and the amount of IL-2 and IFN- γ produced will be quantitated by ELISA. T cells alone are used as a negative control.

Example 3: *In vivo administration of T cell receptor (TCR)-deficient T cells expressing chNKG2D*

[0122] In this example, the TCR-deficient T cells expressing a murine chNKG2D receptor as produced in Example 2 are administered to mice to evaluate the *in vivo* therapeutic potential of said T cells on certain cancers. The chimeric NKG2D-bearing T cells (10^6) are co-injected with RMA/Rae-1 β tumor cells (10^5) subcutaneously to C57BL/6 mice. Chimeric NKG2D-bearing, TCR-deficient T cell-treated mice that are tumor-free or have tumor-inhibited growth of RMA/Rae-1 β tumors after 30 days reflects therapeutic anti-cancer activity in these mice.

[0123] In a second and more stringent model, transduced T cells (10^7) are adoptively transferred i.v. into B6 mice one day before RMA/Rae-1 β s.c. tumor inoculation in the right flank. Suppression of the growth of the RMA/Rae-1 β tumors (s.c.) compared with control vector-modified T cells reflects therapeutic anti-cancer activity in these mice. As for toxicity of treatment with chimeric NKG2D-modified T cells, it is anticipated that the animals will not show any overt evidence of inflammatory damage (i.e., ruffled hair, hunchback or diarrhea, etc.) when treated with chimeric NKG2D-bearing T cells, which would be reflective of a lack of overt toxicity.

[0124] In a more stringent model of established ovarian tumors (ID8), transduced chNKG2D T cells (5×10^6 T cells, i.p.) are injected into mice bearing tumors for 5 weeks. Mice are further injected with T cells at 7 and 9 weeks following tumor challenge. Under these conditions, mice treated with chNKG2D T cells will remain tumor-free for more than 250 days, whereas mice treated on a similar schedule with control T cells will die from tumor growth within 100 days. As for toxicity of treatment with chimeric NKG2D-modified T cells, it is anticipated that the animals will not show any overt evidence of inflammatory damage (i.e., ruffled hair, hunchback or diarrhea, etc.) when treated with chimeric NKG2D-bearing T cells, which would be reflective of a lack of overt toxicity.

[0125] In a model of multiple myeloma, mice bearing 5T33MM tumor cells are treated on day 12 post tumor cell infusion with chNKG2D T cells (5×10^6 cells, i.v.). This treatment will result in an increased life-span of all mice and about half of these mice will be long-term, tumor-free survivors. Mice treated with control T cells will succumb to their tumors within 30 days. No overt evidence of toxicity will be observed due to treatment with the chNKG2D T cells.

[0126] Because the immune system can select for tumor variants, the most effective immunotherapies for cancer are likely going to be those that induce immunity against multiple tumor antigens. In a third experiment, it is tested whether treatment with chimeric NKG2D-bearing T cells will induce host immunity against wild-type tumor cells. Mice that are treated with chimeric NKG2D-bearing T cells and 5T33MM tumor cells, and are tumor-free after 80 days, are challenged with 5T33MM tumor cells. Tumor-free surviving mice are resistant to a subsequent challenge of 5T33MM cells (3×10^5), compared to control naive mice which succumb to the tumor within an average of 27days. However, tumor-free surviving mice are not resistant to a subsequent challenge of RMA-Rae1 tumor cells (3×10^5), and succumb to the tumor in a similar time-span as naïve mice (20days). This indicates that adoptive transfer of chimeric NKG2D-bearing T cells will allow hosts to generate tumor-specific T cell memory.

CLAIMS

What is claimed is:

1. An isolated TCR-deficient T cell.
2. The isolated TCR-deficient T cell of claim 1, wherein said T cell lacks cell-surface expression of at least one component of the T cell receptor complex.
3. The isolated TCR-deficient T cell of claim 1, wherein said T cell lacks cell surface expression of all components of the T cell receptor complex.
4. An isolated TCR-deficient T cell, wherein said T cell expresses at least one functional non-TCR receptor and does not express a functional TCR.
5. An isolated T cell of any one of claims 1-3, wherein said T cell expresses at least one functional non-TCR receptor.
6. The isolated T cell of claims 4 or 5, wherein the at least one functional non-TCR receptor is a chimeric receptor comprising a ligand binding domain attached to a signaling domain.
7. The isolated T cell of claim 6, wherein the ligand binding domain is obtained from NKG2D, NKG2A, NKG2C, NKG2F, LLT1, AICL, CD26, NKRP1, NKp30, NKp44, NKp46, CD244 (2B4), DNAM-1, or NKp80.
8. The isolated T cell of claim 6, wherein the ligand binding domain is obtained from an anti-tumor chimeric antigen receptor or anti-tumor antibody.
9. The isolated T cell of claim 8, wherein the anti-tumor antibody is obtained from an anti-Her2neu antibody or an anti-EGFR antibody.

10. The isolated T cell of claim 6, wherein the signaling domain is obtained from CD3zeta, Dap10, CD28, 41BB, or CD40L.
11. The isolated T cell of claims 4 or 5, wherein the at least one functional non-TCR receptor binds MIC-A, MIC-B, Her2neu, EGFR, mesothelin, CD38, CD20, CD19, PSA, MUC1, MUC2, MUC3A, MUC3B, MUC4, MUC5AC, MUC5B, MUC6, MUC7, MUC8, MUC12, MUC13, MUC15, MUC16, MUC17, MUC19, MUC20, estrogen receptor, progesterone receptor, RON, or one or more members of the ULBP/RAET1 family.
12. The isolated T cell of claim 11, wherein the one or more members of the ULBP/RAET1 family include ULBP1, ULBP2, ULBP3, ULBP4, ULBP5, and ULBP6.
13. The isolated T cell of claims 4 or 5, wherein the at least one functional non-TCR receptor binds to virus antigen or viral-induced antigen found on the surface of an infected cell.
14. The isolated T cell of claim 13, wherein the virus is selected from HCMV, HIV, EBV, HBV, HCV, Ebola virus, a hantavirus, or VSV.
15. The isolated T cell of claim 1 which is human.
16. The isolated T cell of claim 1 which comprises a gene that encodes a dominant selectable marker.
17. A composition comprising an isolated T cell of any one of claims 1-16.
18. The composition of claim 17, comprising a therapeutically-effective amount of said isolated TCR-deficient T cells.

19. The composition of claim 17, comprising a pharmaceutical carrier.
20. A method of preventing or treating cancer in an individual in need thereof, comprising administering the composition of claim 18 to said individual.
21. A method of preventing or treating graft versus host disease in an individual having transplant surgery, comprising administering the composition of claim 18 to said individual prior to the transplant surgery.
22. A method of treating infection in an individual in need thereof, comprising administering the composition of claim 18 to said individual.
23. A method of treating one or more autoimmune disorders in an individual in need thereof, comprising administering the composition of claim 18 to said individual.
24. A method of preventing or treating transplantation rejection in an individual having transplant surgery, comprising administering the composition of claim 18 to said individual prior to the transplant surgery.
25. A method of treating radiation sickness in an individual in need thereof, comprising administering the composition of claim 18 to said individual.
26. A method of preventing or treating cancer in an individual in need thereof, comprising administering a therapeutically-effective amount of the isolated T cells of claim 4 or 5 to said individual.
27. A method of preventing or treating graft versus host disease in an individual having transplant surgery, comprising administering a therapeutically-effective amount of the isolated T cells of claim 4 or 5 to said individual prior to the transplant surgery.

28. A method of treating infection in an individual in need thereof, comprising administering a therapeutically-effective amount of the isolated T cells of claim 4 or 5 to said individual.
29. A method of treating one or more autoimmune disorders in an individual in need thereof, comprising administering a therapeutically-effective amount of the isolated T cells of claim 4 or 5 to said individual.
30. A method of preventing or treating transplantation rejection in an individual having transplant surgery, comprising administering a therapeutically-effective amount of the isolated T cells of claim 4 or 5 to said individual prior to the transplant surgery.
31. A method of treating radiation sickness in an individual in need thereof, comprising administering a therapeutically-effective amount of the isolated T cells of claim 4 or 5 to said individual.
32. A T cell bank comprising TCR-deficient T cells of different tissue HLA types, wherein said T cell bank comprises different T cell containing compositions each composition containing T cells of a specific HLA type, which are TCR-deficient.
33. The T cell bank of claim 32 which comprises T cells of at least 10 different HLA tissue types.
34. The T cell bank of claim 32 which comprises T cells of from about 50-100 different HLA tissue types.
35. The T cell bank of claim 32 which contains T cells that are engineered to stably lack cell-surface expression of at least one component of the T cell receptor complex.

Figure 1

