

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 October 2007 (04.10.2007)

PCT

(10) International Publication Number
WO 2007/110629 A1

(51) International Patent Classification:

A61K 31/00 (2006.01) A61P 25/16 (2006.01)
A61K 31/5415 (2006.01) A61K 49/00 (2006.01)
A61P 25/28 (2006.01)

(74) Agents: KREMER, Simon et al.; Mewburn Ellis LLP,
York House, 23 Kingsway, London Greater London WC2B
6HP (GB).

(21) International Application Number:

PCT/GB2007/001105

(22) International Filing Date: 28 March 2007 (28.03.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/786,700 29 March 2006 (29.03.2006) US

(71) Applicant (for all designated States except US): WISTA
LABORATORIES LTD [SG/SG]; 51 Ayer Rajah Cres-
cent, #07-01/02 Singapore, Singapore 139948 (SG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): WISCHIK, Claude,
Michel [FR/GB]; 20 The Chanonry, Old Aberdeen, Ab-
erdeen Aberdeenshire AB24 1RQ (GB). RICKARD,
Janet, Elizabeth [GB/GB]; 1 Donmouth Gardens, Bridge
of Don, Aberdeen Aberdeenshire AB23 8DS (GB). HAR-
RINGTON, Charles, Robert [GB/GB]; 59 Argyll Place,
Aberdeen Aberdeenshire AB25 2HU (GB). HORSLEY,
David [GB/GB]; 4 Denend Cottages, Udney, Ellon Ab-
erdeenshire AB41 6RU (GB).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES,
FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN,
IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR,
LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY,
MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS,
RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL,
PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF PROTEIN AGGREGATION

(57) Abstract: The invention relates generally to the use of diaminophenothiazine compounds to inhibit or reverse the aggregation of synuclein, and for their use in the manufacture of medicaments for this purpose (e.g. for the treatment of Parkinson's Disease). Also provided are related methods of detecting or labelling of aggregated synuclein.

WO 2007/110629 A1

- 1 -

INHIBITORS OF PROTEIN AGGREGATION

TECHNICAL FIELD

- 5 The present invention generally concerns the aggregation of α -synuclein proteins associated with neurodegenerative disease such as Parkinson's disease (PD) and diaminophenothiazine compounds capable of modulating such aggregation.

BACKGROUND TO INVENTION

- 10 Parkinson's disease is a common human neurodegenerative movement disorder and affects 1% of the elderly population (see discussion by Kapurniotu (2004) Chemistry & Biology 11, pp 1476-1478). Primary clinical symptoms of PD are bradykinesia, resting tremor, muscular rigidity, and difficulty with balance. PD is neuropathologically
15 characterized by a marked and progressive degeneration of dopaminergic neurons and by the presence of fibrillar cytoplasmic inclusions (Lewy bodies [LBs]) and dystrophic neurites (Lewy neurites [LNs]) in the *substantia nigra* and other regions of the brain (Recchia *et al.* (2004) FASEB J 18: 617-626).

- 20 Although the loss of dopamine neurons is certainly related to the major clinical symptoms of PD, the causes and the pathogenesis of this multifactorial disease as well as that of related "synucleinopathies" are still largely unknown.

- The major components of both LBs and LNs are fibrillar aggregates of α -synuclein.
25 α -Synuclein is a widely expressed, neuronal presynaptic protein that appears to play a role in membrane-associated processes and synaptic plasticity and has been linked to learning and development processes. While the mechanism(s) of formation of LBs and LNs and their association with PD are yet not understood, several lines of evidence suggest that α -synuclein fibrillization is associated with PD and that α -synuclein fibrillization causes
30 toxicity (see e.g. Masliah *et al.*, Science, 287:1265-1269 (2000); Feany *et al.*, Nature 404:394-8 (2000)).

- 2 -

In addition to α -synuclein, β -synuclein has also been implicated in neurodegenerative synucleinopathies. Human β -synuclein is a 134-residue neuronal protein that is 78% homologous to α -synuclein. The α - and β -synucleins share a conserved C-terminus with three identically placed tyrosine residues. In addition to α -synuclein-containing LBs and LNs, the development of PD and dementia with LBs is accompanied by the appearance of novel α - and β -synuclein-positive lesions in hippocampus (Galvin et al. 1999) implicating β -synuclein, in addition to α -synuclein, in the onset and progression of these diseases. It has been indicated that β -synuclein may regulate α -synuclein fibrillation, perhaps acting as a chaperone to minimize the aggregation of α -synuclein (Hashimoto et al. 2001; Uversky et al. 2002; Park and Lansbury, 2002). Thus a decrease in the levels of β -synuclein has been considered as a possible factor in the PD etiology (Uversky et al. 2002).

Thus the inhibition or reversal of synuclein aggregation is believed to be of therapeutic benefit.

Li *et al.* (2004) Chemistry & Biology 11: pp 1513-1521 discuss the inhibition of α -synuclein fibrillization, and the disaggregation of fibrils, by the antibiotic rifampicin.

Zhu *et al.* (2004) Journal of Biological Chemistry 279, 26: pp 26846-26857 discuss the inhibition of α -synuclein fibrillization, and the disaggregation of fibrils, by the flavanoid baicalein.

There are a number of other publications in the art said to be concerned with the inhibitors of such aggregation. These include "Compositions for inhibiting the aggregation pathway of alpha-synuclein" (US6780971 - 2004-08-24); "Polyhydroxylated aromatic compounds for the treatment of amyloidosis and alpha-synuclein fibril diseases" (US2004152760 - 2004-08-05); Peptide and peptide derivatives for the treatment of alpha-synuclein related diseases (WO2004009625 - 2004-01-29); Proanthocyanidins for the treatment of amyloid and alpha-synuclein diseases (EP1377287 - 2004-01-07); Methods for preventing neural tissue damage and for the treatment of alpha-synuclein diseases (CN1440420T - 2003-09-03).

- 3 -

However, it will be understood that the provision of compounds not previously known to be capable of inhibiting synuclein aggregation would provide a contribution to the art.

DISCLOSURE OF THE INVENTION

5

The present inventors have demonstrated for first the time that diaminophenothiazine compounds can be used to inhibit the aggregation of synuclein proteins.

10

Briefly, the inventors expressed and purified two forms of α -synuclein and used them in assays for self-assembly and fibril formation. A truncated form of α -synuclein (tsyn) was found to be particularly effective in the fibril-formation assay, and such assembled tsyn was shown to enhance the fluorescence of thioflavine T. The inventors assayed the fibril-disrupting activity of diaminophenothiazines, as well as other compounds. The diaminophenothiazines were found to disrupt assembled α -synuclein at less than 1 μ M.

15

A solid phase assay for synuclein binding was also devised, and was used to show that diaminophenothiazines such as thioninium chlorides, and flavones, inhibited the binding.

20

As will be appreciated by those skilled in the art, in the light of the present disclosure, these results demonstrate utility for such compounds *inter alia* in the treatment of the underlying cause of diseases (such as PD and others discussed herein) associated with synuclein aggregation.

25

Piotrowski, G. (1936) "The treatment of parkinsonian tremor. *Medical Record*, 144:322-323" reported the symptomatic relief of Parkinsonian tremor in a study of 4 individuals using Methylene blue (methyl thioninium chloride – MTC). The MTC was administered intravenously at 1 or 2 mg/kg doses. An oral administration of 8 grains (=518 mg)/day was discontinued due to side effects. The reported effects on tremor were not strong and lasted for a limited time only, while a different symptom (rigidity) was not greatly affected.

30

A separate test with thionin gave no result. The essence of the disclosure is thus that MTC specifically, which was known to have a parasympathetic action, had a limited effect on one symptom of Parkinson's disease i.e. the "parkinsonian tremor".

- 4 -

By contrast the present invention concerns a treatment directed as the underlying disease process itself rather than a symptomatic manifestation of the disease.

5 Diaminophenothiazines have previously been shown to inhibit tau protein aggregation and to disrupt the structure of PHFs, and reverse the proteolytic stability of the PHF core (see WO 96/30766, F Hoffman-La Roche). Such compounds were disclosed for use in the treatment of various diseases, including Alzheimer's disease and Lewy Body Disease.

10 Additionally WO 02/055720 (The University Court of the University of Aberdeen) discusses the use of reduced forms of diaminophenothiazines specifically for the treatment of a variety of protein aggregating diseases, although the disclosure is primarily concerned with tauopathies.

15 WO 2005/030676 (The University Court of the University of Aberdeen) discusses radiolabelled phenothiazines, and their use in diagnosis and therapy e.g. of tauopathies.





However none of these publications specifically disclose the use of diaminophenothiazines, particularly non-reduced forms, for the inhibition or reversal of α -synuclein aggregation in particular.

20

Diaminophenothiazine Compounds

25 The invention pertains to certain diaminophenothiazine compounds and analogs thereof, having one of the following formulae, and pharmaceutically acceptable salts, hydrates, and solvates thereof (collectively referred to herein as "diaminophenothiazines" or "diaminophenothiazine compounds"):

- 5 -



5

Formula (1) depicts compounds in a reduced form, whereas each of Formulae (2), (3), and (4) depicts compounds in an oxidized form.

10

In one embodiment, the compounds are selected from compounds of formula (1), and pharmaceutically acceptable salts, hydrates, and solvates thereof.




In one embodiment, the compounds are selected from compounds of formula (2) or (3), and pharmaceutically acceptable salts, hydrates, and solvates thereof.

15 In one embodiment, the compounds are selected from compounds of formula (4), and pharmaceutically acceptable salts, hydrates, and solvates thereof.

- 6 -

Each one of the above structures is only one of many equivalent resonance structures, and all of which are intended to be encompassed by that representative structure. For example, structure (4) is only one of many equivalent resonance structures, some of which are shown below, and all of which are intended to be encompassed by structure (4):

5



10 Carbon Ring Atom Substituents

In each one of the above formulae, each one of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is independently selected from:

- H;
- F; -Cl; -Br; -I;
- OH; -OR;
- SH; -SR;
- NO₂;
- C(=O)R;
- C(=O)OH; -C(=O)OR;

- 7 -

-C(=O)NH₂; -C(=O)NHR; -C(=O)NR₂; -C(=O)NR^{N1}R^{N2};
 -NH₂; -NHR; -NR₂; -NR^{N1}R^{N2};
 -NHC(=O)H; -NRC(=O)H; -NHC(=O)R; -NRC(=O)R;
 -R;

5 wherein each R is independently selected from:

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;
 unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;
 unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;
 unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;
 10 unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;
 unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

wherein, in each group -NR^{N1}R^{N2}, independently, R^{N1} and R^{N2} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

15 Examples of groups -NR^{N1}R^{N2}, wherein R^{N1} and R^{N2} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms, include: pyrrolidino, piperidino, piperazino, morpholino, pyrrolyl, and substituted forms, such as N-substituted forms, such as N-methyl piperazino.

20 In one embodiment, each one of R¹, R², R⁴, R⁶, R⁸, and R⁹ is independently selected from:

-H;
 -F; -Cl; -Br; -I;
 -OH; -OR;
 -C(=O)OH; -C(=O)OR;
 25 -R.

In one embodiment, each one of R¹, R², R⁴, R⁶, R⁸, and R⁹ is independently selected from:

-H;
 -R.

30

In one embodiment, each R is independently selected from:

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;
 unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;

- 8 -

unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl.

In one embodiment, each R is independently selected from:

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl.

5

In one embodiment, each R is independently selected from: -Me, -Et, -nPr, and -iPr.

In one embodiment, each R is independently selected from: -Me and -Et.

In one embodiment, the C₁₋₆alkyl group is a C₁₋₄alkyl group.

10 In one embodiment, the C₂₋₆alkenyl group is a C₂₋₄alkenyl group.

In one embodiment, the C₃₋₆cycloalkyl group is a C₃₋₄cycloalkyl group.

Examples of unsubstituted aliphatic C₁₋₆alkyl groups include: methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, iso-pentyl, tert-pentyl,

15 neo-pentyl, hexyl, iso-hexyl, etc.

Examples of unsubstituted aliphatic C₂₋₆alkenyl groups include: propen-1-yl, propen-2-yl, buten-1-yl, buten-2-yl, buten-3-yl, etc.

20 Examples of unsubstituted C₃₋₆cycloalkyl groups include: cyclopropyl, cyclopropyl-methyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.

In one embodiment, the C₆₋₁₀carboaryl group is a C₆carboaryl group.

In one embodiment, the C₅₋₁₀heteroaryl group is a C₅₋₆heteroaryl group.

25 In one embodiment, the C₆₋₁₀carboaryl-C₁₋₄alkyl group is a C₆carboaryl-C₁₋₂alkyl group.

Examples of unsubstituted C₆₋₁₀carboaryl groups include: phenyl, naphthyl.

Examples of unsubstituted C₅₋₁₀heteroaryl groups include: pyrrolyl, thienyl, furyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl.

30

Examples of unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl groups include: benzyl, phenylethyl.

- 9 -

In one embodiment, optional substituents (e.g., on aliphatic C₁₋₆alkyl, aliphatic C₁₋₆alkenyl, C₃₋₆cycloalkyl, C₆₋₁₀carboaryl, C₅₋₁₀heteroaryl, C₆₋₁₀carboaryl-C₁₋₄alkyl) are independently selected from:

- 5 -F; -Cl; -Br; -I;
 -OH; -OR';
 -SH; -SR';
 -NO₂;
 -C(=O)R';
 10 -C(=O)OH; -C(=O)OR';
 -C(=O)NH₂; -C(=O)NHR'; -C(=O)NR'₂; -C(=O)NR'^{N1}R'^{N2};
 -NH₂; -NHR'; -NR'₂; -NR'^{N1}R'^{N2};
 -NHC(=O)H; -N'RC(=O)H; -NHC(=O)'R; -N'RC(=O)'R;
 -R';

- 15 wherein each R' is independently selected from:

- unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;
 unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;
 unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;
 unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;
 20 unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;
 unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

wherein, in each group -NR'^{N1}R'^{N2}, independently, R'^{N1} and R'^{N2} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

- 25 In one embodiment, optional substituents (e.g., on aliphatic C₁₋₆alkyl, aliphatic C₁₋₆alkenyl, C₃₋₆cycloalkyl, C₆₋₁₀carboaryl, C₅₋₁₀heteroaryl, C₆₋₁₀carboaryl-C₁₋₄alkyl) are independently selected from:

- F; -Cl; -Br; -I;
 -OH; -OR;
 30 -C(=O)OH; -C(=O)OR';
 -R'.

- 10 -

In one embodiment, optional substituents (e.g., on aliphatic C₁₋₆alkyl, aliphatic C₁₋₆alkenyl, C₃₋₆cycloalkyl, C₆₋₁₀carboaryl, C₅₋₁₀heteroaryl, C₆₋₁₀carboaryl-C₁₋₄alkyl) are as defined above, except that each R' is independently selected from:

unsubstituted aliphatic C₁₋₆alkyl;
unsubstituted aliphatic C₂₋₆alkenyl;
unsubstituted C₃₋₆cycloalkyl;
unsubstituted C₆₋₁₀carboaryl;
unsubstituted C₅₋₁₀heteroaryl;
unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl.

In one embodiment, optional substituents (e.g., on aliphatic C₁₋₆alkyl, aliphatic C₁₋₆alkenyl, C₃₋₆cycloalkyl, C₆₋₁₀carboaryl, C₅₋₁₀heteroaryl, C₆₋₁₀carboaryl-C₁₋₄alkyl) are as defined above, except that each R' is independently selected from:

unsubstituted aliphatic C₁₋₆alkyl;
unsubstituted aliphatic C₂₋₆alkenyl;
unsubstituted C₃₋₆cycloalkyl.

In one embodiment, optional substituents (e.g., on aliphatic C₁₋₆alkyl, aliphatic C₁₋₆alkenyl, C₃₋₆cycloalkyl, C₆₋₁₀carboaryl, C₅₋₁₀heteroaryl, C₆₋₁₀carboaryl-C₁₋₄alkyl) are as defined above, except that each R' is independently selected from:

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl.

In one embodiment, optional substituents (e.g., on aliphatic C₁₋₆alkyl, aliphatic C₁₋₆alkenyl, C₃₋₆cycloalkyl, C₆₋₁₀carboaryl, C₅₋₁₀heteroaryl, C₆₋₁₀carboaryl-C₁₋₄alkyl) are as defined above, except that each R' is independently selected from: -Me, -Et, -nPr, and -iPr.

In one embodiment, optional substituents (e.g., on aliphatic C₁₋₆alkyl, aliphatic C₁₋₆alkenyl, C₃₋₆cycloalkyl, C₆₋₁₀carboaryl, C₅₋₁₀heteroaryl, C₆₋₁₀carboaryl-C₁₋₄alkyl) are as defined above, except that each R' is independently selected from: -Me and -Et.

In one embodiment, each one of R¹, R², R⁴, R⁶, R⁸, and R⁹ is independently selected from: -H, -Me, -Et, -nPr, and -iPr.

- 11 -

In one embodiment, each one of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is independently selected from: -H, -Me, and -Et.

5 In one embodiment, each one of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is independently selected from: -H and -Me.

In one embodiment, all except four of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is -H.

In one embodiment, all except two of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is -H.

In one embodiment, all except one of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is -H.

10 In one embodiment, each of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is -H.

Amino Groups

15 In each one of the above formulae, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently -H or as defined above for R; or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

20 For example, in one embodiment, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently as defined above for R; or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

For example, in one embodiment, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently selected from:

25 -H;
 unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;
 unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl;
 unsubstituted C_{6-10} carboaryl; substituted C_{6-10} carboaryl;
 unsubstituted C_{5-10} heteroaryl; substituted C_{5-10} heteroaryl;
 30 unsubstituted C_{6-10} carboaryl- C_{1-4} alkyl; substituted C_{6-10} carboaryl- C_{1-4} alkyl;
 or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

- 12 -

For example, in one embodiment, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently selected from:

unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;
 5 unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl;
 unsubstituted C_{6-10} carboaryl; substituted C_{6-10} carboaryl;
 unsubstituted C_{5-10} heteroaryl; substituted C_{5-10} heteroaryl;
 unsubstituted C_{6-10} carboaryl- C_{1-4} alkyl; substituted C_{6-10} carboaryl- C_{1-4} alkyl;

or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a
 10 ring having from 3 to 7 ring atoms.

In another example, in one embodiment, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently selected from:

-H;

15 unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;
 unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl;

or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a
 ring having from 3 to 7 ring atoms.

20 In another example, in one embodiment, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently selected from:

unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;
 25 unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl;

or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a
 ring having from 3 to 7 ring atoms.

30 In another example, in one embodiment, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently selected from:

-H;

unsubstituted aliphatic C_{1-6} alkyl;
 unsubstituted aliphatic C_{2-6} alkenyl;

- 13 -

unsubstituted C₃₋₆cycloalkyl;
or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

- 5 In another example, in one embodiment, in each group -NR^{3NA}R^{3NB}, if present, each one of R^{3NA} and R^{3NB} is independently selected from:

unsubstituted aliphatic C₁₋₆alkyl;
unsubstituted aliphatic C₂₋₆alkenyl;
unsubstituted C₃₋₆cycloalkyl;

- 10 or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

In another example, in one embodiment, in each group -NR^{3NA}R^{3NB}, if present, each one of R^{3NA} and R^{3NB} is independently selected from: -H, -Me, -Et, -nPr, and -iPr.

- 15 In another example, in one embodiment, in each group -NR^{3NA}R^{3NB}, if present, each one of R^{3NA} and R^{3NB} is independently selected from: -H, -Me, and -Et (e.g., -NR^{3NA}R^{3NA} is -NH₂, -NHMe, -NMe₂, -NH₂Et, -NEt₂, or -NMeEt).

- 20 In another example, in one embodiment, in each group -NR^{3NA}R^{3NB}, if present, each one of R^{3NA} and R^{3NB} is independently selected from: -H and -Me (e.g., -NR^{3NA}R^{3NA} is -NH₂, -NHMe, or -NMe₂).

- In precise analogy, in each one of the above formulae, in each group -NR^{7NA}R^{7NB}, if present, each one of R^{7NA} and R^{7NB} is independently -H or as defined above for R; or R^{7NA} and R^{7NB} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

- 25 For example, in one embodiment, in each group -NR^{7NA}R^{7NB}, if present, each one of R^{7NA} and R^{7NB} is independently as defined above for R; or R^{7NA} and R^{7NB} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

30 In one embodiment, -NR^{3NA}R^{3NB} and -NR^{7NA}R^{7NB}, if both present, are the same.

- 14 -

In one embodiment, $-NR^{3NA}R^{3NB}$ and $-NR^{7NA}R^{7NB}$, if both present, are different.

In each one of the above formulae, in each group $=NR^{3NC}$, if present, R^{3NC} is independently -H or as defined above for R.

5

For example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently as defined above for R.

10

For example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from:

-H;
 unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;
 unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl;
 15 unsubstituted C_{6-10} carboaryl; substituted C_{6-10} carboaryl;
 unsubstituted C_{5-10} heteroaryl; substituted C_{5-10} heteroaryl;
 unsubstituted C_{6-10} carboaryl- C_{1-4} alkyl; substituted C_{6-10} carboaryl- C_{1-4} alkyl.

20

For example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from:

unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;
 unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl;
 unsubstituted C_{6-10} carboaryl; substituted C_{6-10} carboaryl;
 25 unsubstituted C_{5-10} heteroaryl; substituted C_{5-10} heteroaryl;
 unsubstituted C_{6-10} carboaryl- C_{1-4} alkyl; substituted C_{6-10} carboaryl- C_{1-4} alkyl.

30

In another example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from:

-H;
 unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;
 unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl.

- 15 -

In another example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from:

- unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
- 5 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;
- unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl.

In another example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from:

- 10 -H;
- unsubstituted aliphatic C_{1-6} alkyl;
- unsubstituted aliphatic C_{2-6} alkenyl;
- unsubstituted C_{3-6} cycloalkyl.

15 In another example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from:

- unsubstituted aliphatic C_{1-6} alkyl;
- unsubstituted aliphatic C_{2-6} alkenyl;
- unsubstituted C_{3-6} cycloalkyl.

20

In another example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from: -H, -Me, -Et, -nPr, and -iPr.

25 In another example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from: -H, -Me, and -Et (e.g., $=NR^{3NC}$ is =NH, =NMe, or =NEt).

In another example, in one embodiment, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from: -H and -Me (e.g., $=NR^{3NC}$ is =NH or =NMe).

30 In precise analogy, in each one of the above formulae, in each group $=NR^{7NC}$, if present, R^{7NC} is independently as defined above for R^{3NC} .

- 16 -

Nitrogen Ring Atom Substituent

Also, in precise analogy, in each one of the above formulae, R^{N10} , if present, is independently as defined above for R^{3NC} (or R^{7NC}).

5

For example, in one embodiment, R^{N10} , if present, is independently selected from: -H and unsubstituted aliphatic C_{1-6} alkyl.

10 For example, in one embodiment, R^{N10} , if present, is independently selected from: -H, -Me, and -Et.

For example, in one embodiment, R^{N10} , if present, is independently selected from: -H and -Me.

15 For example, in one embodiment, R^{N10} , if present, is independently -H.

Counter Ion

X^- , if present, is one or more anionic counter ions to achieve electrical neutrality.

20

Examples of suitable anionic counter ions are discussed below under the heading "Salts".

In one embodiment, X^- is independently a halogen anion (i.e., a halide).

In one embodiment, X^- is independently Cl^- , Br^- , or I^- .

25 In one embodiment, X^- is independently Cl^- .

In one embodiment, X^- is independently NO_3^- .

Combinations

30

All plausible combinations of the embodiments described above are disclosed herein as if each combination was individually and explicitly recited.


- 17 -

Isomers

Certain compounds may exist in one or more particular geometric, optical, enantiomeric, diastereomeric, epimeric, atropic, stereoisomeric, tautomeric, conformational, or anomeric forms, including but not limited to, cis- and trans-forms; E- and Z-forms; c-, t-, and r- forms; endo- and exo-forms; R-, S-, and meso-forms; D- and L-forms; d- and l-forms; (+) and (-) forms; keto-, enol-, and enolate-forms; syn- and anti-forms; synclinal- and anticlinal-forms; α - and β -forms; axial and equatorial forms; boat-, chair-, twist-, envelope-, and halfchair-forms; and combinations thereof, hereinafter collectively referred to as "isomers" (or "isomeric forms").

Note that, except as discussed below for tautomeric forms, specifically excluded from the term "isomers," as used herein, are structural (or constitutional) isomers (i.e., isomers which differ in the connections between atoms rather than merely by the position of atoms in space). For example, a reference to a methoxy group, $-\text{OCH}_3$, is not to be construed as a reference to its structural isomer, a hydroxymethyl group, $-\text{CH}_2\text{OH}$. Similarly, a reference to ortho-chlorophenyl is not to be construed as a reference to its structural isomer, meta-chlorophenyl. However, a reference to a class of structures may well include structurally isomeric forms falling within that class (e.g., C_{1-7} alkyl includes n-propyl and iso-propyl; butyl includes n-, iso-, sec-, and tert-butyl; methoxyphenyl includes ortho-, meta-, and para-methoxyphenyl).

The above exclusion does not pertain to tautomeric forms, for example, keto-, enol-, and enolate-forms, as in, for example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/amidine, nitroso/oxime, thioketone/enethiol, N-nitroso/hydroxyazo, and nitro/aci-nitro.



Note that specifically included in the term "isomer" are compounds with one or more isotopic substitutions. For example, H may be in any isotopic form, including ^1H , ^2H (D),

- 18 -

and ^3H (T); C may be in any isotopic form, including ^{11}C , ^{12}C , ^{13}C , and ^{14}C ; O may be in any isotopic form, including ^{16}O and ^{18}O ; and the like.

Unless otherwise specified, a reference to a particular compound includes all such isomeric forms, including (wholly or partially) racemic and other mixtures thereof. Methods for the preparation (e.g., asymmetric synthesis) and separation (e.g., fractional crystallisation and chromatographic means) of such isomeric forms are either known in the art or are readily obtained by adapting the methods taught herein, or known methods, in a known manner.

10 Salts

It may be convenient or desirable to prepare, purify, and/or handle a corresponding salt of the compound, for example, a pharmaceutically-acceptable salt. Examples of pharmaceutically acceptable salts are discussed in Berge *et al.*, 1977, "Pharmaceutically Acceptable Salts," J. Pharm. Sci., Vol. 66, pp. 1-19.

For example, if the compound is anionic, or has a functional group which may be anionic (e.g., $-\text{COOH}$ may be $-\text{COO}^-$), then a salt may be formed with a suitable cation. Examples of suitable inorganic cations include, but are not limited to, alkali metal ions such as Na^+ and K^+ , alkaline earth cations such as Ca^{2+} and Mg^{2+} , and other cations such as Al^{+3} .

Examples of suitable organic cations include, but are not limited to, ammonium ion (i.e., NH_4^+) and substituted ammonium ions (e.g., NH_3R^+ , NH_2R_2^+ , NHR_3^+ , NR_4^+). Examples of some suitable substituted ammonium ions are those derived from: ethylamine, diethylamine, dicyclohexylamine, triethylamine, butylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, benzylamine, phenylbenzylamine, choline, meglumine, and tromethamine, as well as amino acids, such as lysine and arginine. An example of a common quaternary ammonium ion is $\text{N}(\text{CH}_3)_4^+$.

If the compound is cationic, or has a functional group which may be cationic (e.g., $-\text{NH}_2$ may be $-\text{NH}_3^+$), then a salt may be formed with a suitable anion. Examples of suitable inorganic anions include, but are not limited to, those derived from the following inorganic acids: hydrochloric, hydrobromic, hydroiodic, sulfuric, sulfurous, nitric, nitrous, phosphoric, and phosphorous.

- 19 -

Examples of suitable organic anions include, but are not limited to, those derived from the following organic acids: 2-acetyoxybenzoic, acetic, ascorbic, aspartic, benzoic, camphorsulfonic, cinnamic, citric, edetic, ethanedisulfonic, ethanesulfonic, fumaric, 5 glucheptonic, gluconic, glutamic, glycolic, hydroxymaleic, hydroxynaphthalene carboxylic, isethionic, lactic, lactobionic, lauric, maleic, malic, methanesulfonic, mucic, oleic, oxalic, palmitic, pamoic, pantothenic, phenylacetic, phenylsulfonic, propionic, pyruvic, salicylic, stearic, succinic, sulfanilic, tartaric, toluenesulfonic, and valeric. Examples of suitable polymeric organic anions include, but are not limited to, those derived from the following 10 polymeric acids: tannic acid, carboxymethyl cellulose.

The compound may also be provided in the form of a mixed salt (i.e., the compound in combination with a salt, or another salt). For example, methyl-thioninium chloride zinc chloride mixed salt (MTZ) is a mixed salt of methyl-thioninium chloride (MTC), a chloride 15 salt, and another salt, zinc chloride. Such mixed salts are intended to be encompassed by the term "and pharmaceutically acceptable salts thereof".

Unless otherwise specified, a reference to a particular compound also includes salt forms thereof.

Hydrates and Solvates

It may be convenient or desirable to prepare, purify, and/or handle a corresponding solvate of the active compound. The term "solvate" is used herein in the conventional sense to 25 refer to a complex of solute (e.g., compound, salt of compound) and solvent. If the solvent is water, the solvate may be conveniently referred to as a hydrate, for example, a mono-hydrate, a di-hydrate, a tri-hydrate, etc.





Unless otherwise specified, a reference to a particular compound also includes solvate 30 forms thereof.

- 20 -






Some Preferred Examples

Some preferred diaminophenothiazines include the following, and pharmaceutically acceptable salts, hydrates, and solvates thereof:






5

A	MTC	 <p>methyl-thioninium chloride</p>
B	ETC	 <p>ethyl-thioninium chloride</p>
C	DMMTC	 <p>1,9-dimethyl-methyl-thioninium chloride</p>
D	DEMTC	 <p>1,9-diethyl-methyl-thioninium chloride</p>





- 21 -

E	DMETC	 <p>1,9-dimethyl-ethyl-thioninium chloride</p>
F	DEETC	 <p>1,9-diethyl-ethyl-thioninium chloride</p>
G	MTZ	 <p>methyl-thioninium chloride zinc chloride mixed salt</p>
H	ETZ	 <p>ethyl-thioninium chloride zinc chloride mixed salt</p>
I	MTI	 <p>methyl-thioninium iodide</p>

- 22 -

J	MTI.HI	 <p>methyl-thioninium iodide hydrogen iodide mixed salt</p>
K	ETI	 <p>ethyl-thioninium iodide</p>
L	ETI.HI	 <p>ethyl-thioninium iodide hydrogen iodide mixed salt</p>
M	MTN	 <p>methyl-thioninium nitrate</p>
N	ETN	 <p>ethyl-thioninium nitrate</p>

- 23 -

O		 <p>Tolonium chloride</p>
P		 <p>Thionine</p>
Q		 <p>Azure A</p>
R		 <p>Azure B</p>

In one embodiment, the diaminophenothiazine is selected from: MTC, ETC, DEMTC, DEETC, Thionine, and Tolonium Chloride (also known as Toluidine Blue O).

- 5 Preferred compounds of the present invention are those which show high activity in the assays described herein.

- 24 -

Inhibition of aggregation

In all therapeutic and other aspects of the invention, it is preferred that the diaminophenothiazine is in substantially oxidised form e.g. at least 50, 60, 70, 80, 90, 95,
5 99, or 100% oxidised form.

Thus in a first aspect of the present invention there is disclosed use of a diaminophenothiazine to inhibit the aggregation of synuclein, in particular, α -synuclein, for example in a cell.

10 The aggregation may be in the context of a disease state manifested as neurodegeneration and/or clinical dementia.

In another aspect, the invention provides a diaminophenothiazine for use in a method of
15 treatment or therapy of the human or animal body, e.g., in a method of treatment or prophylaxis of a neurodegenerative disease and/or clinical dementia associated with synuclein, particularly α -synuclein, aggregation.

In another aspect, the invention provides for use of a diaminophenothiazine in the
20 manufacture of a medicament to inhibit the aggregation of synuclein, particularly α -synuclein, which aggregation is associated with a disease state manifested as neurodegeneration and/or clinical dementia, e.g., a medicament for the treatment or prophylaxis of a neurodegenerative disease and/or clinical dementia associated with synuclein aggregation.

25 In another aspect, the invention provides a method of treatment or prophylaxis of a neurodegenerative disease and/or clinical dementia associated with synuclein aggregation, particularly α -synuclein, which method comprises administering to a subject a prophylactically or therapeutically effective amount of a diaminophenothiazine, or
30 therapeutic composition comprising the same, so as to inhibit the aggregation of the synuclein.

- 25 -

In another aspect, the invention provides a method of regulating the aggregation of synuclein, particularly α -synuclein, in the brain of a mammal, which aggregation is associated with a disease state as described below, the treatment comprising the step of administering to said mammal in need of said treatment, a prophylactically or
5 therapeutically effective amount of a diaminophenothiazine.

In another aspect, the invention provides a method of inhibiting production of synuclein, particularly α -synuclein, aggregates in the brain of a mammal, the treatment being as described above.

10 In another aspect, the invention provides a drug product for the treatment of a disease associated with synuclein, particularly α -synuclein, aggregation in a mammal suffering therefrom, comprising a container labeled or accompanied by a label indicating that the drug product is for the treatment of said disease, the container containing one or more
15 dosage units each comprising at least one pharmaceutically acceptable excipient and, as an active ingredient, an isolated pure diaminophenothiazine compound selected from those described above.

20 Diaminophenothiazines may be administered alone, or in combination with other treatments, either simultaneously or sequentially, dependent upon the condition or disease to be treated. In particular it may be desired to use or formulate diaminophenothiazines with other inhibitors of the relevant protein aggregation reaction.

25 Preferred combinations are any one or more of the diaminophenothiazine compounds discussed above plus a compound that modulates dopamine levels in the mammal to be treated. Such additional compounds may include levo-DOPA and dopaminergic agonists such as ropinirole (see e.g. Olanow, C.W. 2004, The scientific basis for the current treatment of Parkinson's disease, Ann. Rev. Med. 55:41-60)

30 In each case, preferably the mammal is a human.

- 26 -

Ligands

Diaminophenothiazine compounds discussed herein that are capable of inhibiting the aggregation of α -synuclein will also be capable of acting as ligands or labels of α -synuclein (or aggregated α -synuclein). Thus, in one embodiment, the diaminophenothiazine compound is a ligand, e.g., a ligand of synuclein (or aggregated synuclein), particularly α -synuclein.

Such diaminophenothiazine compounds (ligands) may incorporate, be conjugated to, be chelated with, or otherwise be associated with, other chemical groups, such as detectable labels, such as stable and unstable detectable isotopes, radioisotopes, positron-emitting atoms, magnetic resonance labels, dyes, fluorescent markers, antigenic groups, therapeutic moieties, or any other moiety that may aid in a prognostic, diagnostic or therapeutic application.

For example, in one embodiment, the diaminophenothiazine compound is as defined above, but with the additional limitation that the compound incorporates, is conjugated to, is chelated with, or is otherwise associated with one or more (e.g., 1, 2, 3, 4, etc.) isotopes, radioisotopes, positron-emitting atoms, magnetic resonance labels, dyes, fluorescent markers, antigenic groups, or therapeutic moieties.

In one embodiment, the diaminophenothiazine compound is a ligand as well as a label, e.g., a label for α -synuclein (or aggregated α -synuclein), and incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels.

For example, in one embodiment, the diaminophenothiazine compound is as defined above, but with the additional limitation that the compound incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels.

In one embodiment, the detectable label is, or incorporates, a stable detectable isotope, an unstable detectable isotope, a radioisotope (e.g., ^{99}Tc), a positron-emitting atom (e.g., ^{11}C ,

- 27 -

¹⁸F), a magnetic resonance label (e.g., ¹⁹F), a dye, a fluorescent group, or an antigenic group.

5 Labelled diaminophenothiazine compounds (e.g., when ligated to α -synuclein or aggregated α -synuclein) may be visualised or detected by any suitable means, and the skilled person will appreciate that any suitable detection means as is known in the art may be used.

10 For example, the diaminophenothiazine compound (ligand-label) may be suitably detected by incorporating a positron-emitting atom (e.g., ¹¹C) (e.g., as a carbon atom of one or more alkyl group substituents, e.g., methyl group substituents) and detecting the compound using positron emission tomography (PET) as is known in the art.





15 For example, in one embodiment, the diaminophenothiazine compound is as defined above, but with the additional limitation that at least one (e.g., 1, 2, 3, 4, etc.) of the ring carbon atoms of the diaminophenothiazine compound is a positron-emitting carbon atom, e.g., ¹¹C; and/or at least one (e.g., 1, 2, 3, 4, etc.) of the carbon atoms of at least one (e.g., 1, 2, 3, 4, etc.) of the substituents R¹, R², R⁴, R⁶, R⁸, R⁹, R^{3NA}, R^{3NB}, R^{3NC}, R^{7NA}, R^{7NB}, R^{7NC}, and R^{N10} is a positron-emitting carbon atom, e.g., ¹¹C.

20 In one embodiment, at least one (e.g., 1, 2, 3, 4, etc.) of the carbon atoms of at least one (e.g., 1, 2, 3, 4) of the substituents R^{3NA}, R^{3NB}, R^{3NC}, R^{7NA}, R^{7NB}, and R^{7NC} is a positron-emitting carbon atom, e.g., ¹¹C.

25 In one embodiment, at least one (e.g., 1, 2, 3, 4, etc.) of the substituents R^{3NA}, R^{3NB}, R^{3NC}, R^{7NA}, R^{7NB}, and R^{7NC} is -¹¹CH₃.

- 28 -

Examples of such diaminophenothiazine compounds (i.e., which incorporate a positron-emitting atom detectable by PET) include the following:



Suitable methods for preparing these and similar ^{11}C labelled diaminophenothiazines are shown, for example, in WO 02/075318 (see Figures 11a, 11b, 12) and WO 2005/030676.

Alternatively, or in addition, the diaminophenothiazine compound may be conjugated to a chelating group (i.e., a moiety suitable for conjugation to another molecule or atom or ion by complex or chelate formation) (e.g., a radioisotope-chelating group, e.g., a technetium-chelating group, e.g., a diethylenetriaminepentaacetic acid group) that is chelated to a detectable label (e.g., a radioisotope, e.g., ^{99}Tc).

- 29 -

For example, in one embodiment, the diaminophenothiazine compound is as defined above, but with the additional limitation that at least one (e.g., 1, 2, 3, 4, etc.) of the substituents R^1 , R^2 , R^4 , R^6 , R^8 , R^9 , R^{3NA} , R^{3NB} , R^{3NC} , R^{7NA} , R^{7NB} , R^{7NC} , and R^{N10} is, or
5 incorporates, a chelating group (e.g., a technetium-chelating group, e.g., a diethylenetriaminepentaacetic acid group) that is able to chelate a detectable label (e.g., a radioisotope, e.g., ^{99}Tc).

Alternatively, or in addition, the diaminophenothiazine compound may incorporate a
10 magnetic resonance label (e.g., ^{19}F), and so be suitable for MRI imaging (see e.g. Higuchi et al. Nat Neurosci. 2005 Apr; 8(4):527-33).

For example, in one embodiment, the diaminophenothiazine compound is as defined above, but with the additional limitation that at least one (e.g., 1, 2, 3, 4, etc.) of the
15 substituents R^1 , R^2 , R^4 , R^6 , R^8 , R^9 , R^{3NA} , R^{3NB} , R^{3NC} , R^{7NA} , R^{7NB} , R^{7NC} , and R^{N10} is, or incorporates, a magnetic resonance label (e.g., ^{19}F , for example, as ^{-19}F , $-\text{C}(^{19}\text{F})_3$, etc.)

Thus, in one aspect, the present invention provides a method of labelling synuclein (or aggregated synuclein), particularly α -synuclein, comprising the steps of: contacting the
20 synuclein (or aggregated synuclein) with a diaminophenothiazine compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, a detectable label.

In another aspect, the present invention provides a method of detecting synuclein (or aggregated synuclein), particularly α -synuclein, comprising the steps of: contacting the
25 synuclein (or aggregated synuclein) with a diaminophenothiazine compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, a detectable label, and detecting the presence and/or amount of said compound bound to synuclein (or aggregated synuclein).

30 In another aspect, the present invention provides a method of diagnosis or prognosis of a synucleinopathy in a subject believed to suffer from the disease, comprising the steps of:

- 30 -

(i) introducing into the subject a diaminophenothiazine compound capable of labelling synuclein or aggregated synuclein, particularly α -synuclein (e.g., a diaminophenothiazine compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, a detectable label),

5 (ii) determining the presence and/or amount of said compound bound to synuclein or aggregated synuclein in the brain of the subject,

(iii) correlating the result of the determination made in (ii) with the disease state of the subject.

10 In another aspect, the present invention provides a diaminophenothiazine compound capable of labelling synuclein or aggregated synuclein, particularly α -synuclein, (e.g., a diaminophenothiazine compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, a detectable label), for use in a method of diagnosis or prognosis of a synucleinopathy.

15 In another aspect, the present invention provides use of a diaminophenothiazine compound capable of labelling synuclein or aggregated synuclein, particularly α -synuclein (e.g., a diaminophenothiazine compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, a detectable label), in a method of manufacture of a
20 diagnostic or prognostic reagent for use in the diagnosis or prognosis of a synucleinopathy.

Those skilled in the art will appreciate that instead of administering diaminophenothiazine ligands/labels directly, they could be administered in a precursor form, for conversion to the active form (e.g., ligating form, labelling form) by an activating agent present in, or
25 administered to, the same subject.

Diseases

The disease states with which the present invention is concerned are synucleinopathies.

30 As those skilled in the art will be aware, the term synucleinopathies is used to name a group of neurodegenerative disorders characterized by fibrillary aggregates of synuclein protein, particularly α -synuclein, in the cytoplasm of selective populations of neurons and

- 31 -

glia, and in particular in which the presence of synuclein-containing inclusions are pathognomic for the disease.

This should be distinguished from non-synucleinopathy disorders in which synuclein-
5 containing inclusions may or may not be present in addition to other pathologies.

The synucleinopathies currently consist of the following disorders: Parkinson's disease (PD), dementia with Lewy bodies (DLB), multiple system atrophy (MSA), drug-induced parkinsonism (e.g. produced by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine [MPTP] or
10 pesticides such as rotenone), and pure autonomic failure (PAF).

Non-synucleinopathy disorders in which Lewy bodies may be found include the following: Alzheimer's disease, Pick's/frontotemporal dementia, Creutzfeldt-Jakob disease, ataxia telangectasia, corticobasal degeneration, dystonia, progressive supranuclear palsy,
15 neuraxonal dystrophy, subacute sclerosing panencephalitis, amyotrophic lateral sclerosis, ALS-dementia Guam complex, Meige's syndrome and Hallervorden-Spatz disease (HSD) (neurodegeneration with brain iron). LBs occur commonly in a variety of neurodegenerative diseases. Studies have indicated that the morphology of α -synuclein fibrils in neuronal LBs show basic similarities regardless of the underlying disease.

20 Parkinson's disease has a high prevalence (ca. 100 per 100,000) compared with MSA (4 per 100,000).

DLB was adopted as the consensus name to cover several other ones that had existed
25 earlier (McKeith, I. G. et al. (1996). "Consensus guidelines for the clinical and pathologic diagnosis of dementia with Lewy bodies (DLB): report of the consortium on DLB international workshop. *Neurology*, 47, 1113-1124." *Neurology* 47: 1113-1124). These include senile dementia of the Lewy body type, Lewy body variant of Alzheimer's disease, cortical Lewy body dementia and Lewy body dementia. MSA encompasses Shy Drager syndrome, olivopontocerebellar atrophy and striatonigral degeneration. DLB has been
30 reported to be the second most common form of dementia in the elderly after Alzheimer's disease.

- 32 -

Parkinson's disease is characterised by LBs in substantia nigra, but they may also be found in cortex. DLB is characterised by more frequent occurrence of cortical LBs. In MSA, filamentous inclusions, termed glial cytoplasmic inclusions (GCIs) are found mainly in oligodendrocytes. The nature of the component filaments in LBs was unknown until 1997, when two findings established the major component: (i) a missense mutation in α -synuclein was found to cause a rare form of familial PD (Polymeropoulos, M. H. et al. (1997).

"Mutation in the α -synuclein gene identified in families with Parkinson's disease." Science 276: 2045-2047) and (ii) LBs and LNs in idiopathic PD and in DLB were found to be immunoreactive for α -synuclein (Spillantini, M. G. et al. (1997). " α -Synuclein in Lewy bodies." Nature 388: 839-840). Recombinant α -synuclein can form filaments in vitro. The protein is a natively unfolded protein. In diseases where it aggregates, it forms fibrils with β -sheet structure.

Preferably the compounds of the present invention are used in respect of a synucleinopathy selected from PD, PAF, MSA and HSD.

Choice of subject

The ligands disclosed herein may be used as part of a method of diagnosis or prognosis. It may be used to select a patient for treatment, or to assess the effectiveness of a treatment or a therapeutic e.g. an inhibitor of α -synuclein association administered to the subject.

Suitable subjects for the method may be selected on the basis of conventional factors.

Thus the initial selection of a patient may involve any one or more of: rigorous evaluation by experienced clinician; exclusion of non-AD diagnosis as far as possible by

supplementary laboratory and other investigations; objective evaluation of level of cognitive function using neuropathologically validated battery.

Dosage units, and formulation and administration of compounds

Administration of compounds, compositions or medicaments as described herein is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual.

- 33 -

For ligands the amount will be a diagnostically effective amount which will give rise to detectable binding in the patient suffering from a synucleinopathy.

- 5 For medicaments the actual amount administered, and rate and time-course of administration, will depend on the nature and severity of the disease being treated. Prescription of treatment, e.g. decisions on dosage etc., is within the responsibility of general practitioners and other medical doctors, and typically takes account of the disorder to be treated, the condition of the individual patient, the site of delivery, the method of
10 administration and other factors known to practitioners.

Typically the mammal will be human, although use in animals (e.g. for test purposes, or veterinary therapeutic purposes) is also embraced by the invention.

- 15 Example phenothiazines of the present invention are known in the art and may be manufactured by the processes referred to in standard texts (e.g. Merck Manual, Houben-Weyl, Beilstein E III/IV 27, 1214 ff, J. Heterocycl. Chem 21, 613 (1984), etc.). The compounds of the above formulae, their pharmaceutically acceptable salts, or other compounds found to have the properties defined in the assays provided, could be used as
20 medicaments after further testing for toxicity (e.g. in the form of pharmaceutical preparations).

- The prior pharmaceutical use of methylene blue in a wide range of medical indications has been described, including treatment of methaemoglobineamia and the prophylaxis of manic
25 depressive psychosis (Naylor (1986) Biol. Psychiatry 21, 915-920), and CNS penetration following systemic administration has been described (Muller (1992) Acta Anat., 144, 39-44). The production of Azure A and B occur as normal metabolic degradation products of methylene blue (Disanto and Wagner (1972a) J. Pharm. Sci. 61, 598-602; Disanto and Wagner (1972b) J. Pharm. Sci. 61 1086-1094). The administration of pharmaceuticals can
30 be effected parentally such as orally, in the form of tablets, coated tablets, dragees, hard and soft gelatine capsules, solutions, emulsions or suspensions), nasally (e.g. in the form of nasal sprays) or rectally (e.g. in the form of suppositories). However, the administration

- 34 -

can also be effected parentally such as intramuscularly or intravenously (e.g. in the form of injection solutions).

5 The compositions may include, in addition to the above constituents, pharmaceutically-acceptable excipients, preserving agents, solubilizers, viscosity-increasing substances, stabilising agents, wetting agents, emulsifying agents, sweetening agents, colouring agents, flavouring agents, salts for varying the osmotic pressure, buffers, or coating agents. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material may depend on the route of administration. Examples of techniques and protocols can be found in 10 "*Remington's Pharmaceutical Sciences*", 16th edition, Osol, A. (ed.), 1980.

Where the composition is formulated into a pharmaceutical composition, the administration thereof can be effected parentally such as orally, in the form of powders, tablets, coated 15 tablets, dragees, hard and soft gelatine capsules, solutions, emulsions or suspensions, nasally (e.g. in the form of nasal sprays) or rectally (e.g. in the form of suppositories). However, the administration can also be effected parentally such as intramuscularly, intravenously, cutaneously, subcutaneously, or intraperitoneally (e.g. in the form of injection solutions).

20 Thus, for example, where the pharmaceutical composition is in the form of a tablet, it may include a solid carrier such as gelatine or an adjuvant. For the manufacture of tablets, coated tablets, dragees and hard gelatine capsules, the active compounds and their pharmaceutically-acceptable acid addition salts can be processed with pharmaceutically inert, inorganic or organic excipients. Lactose, maize, starch or derivatives thereof, talc, 25 stearic acid or its salts etc. can be used, for example, as such excipients for tablets, dragees and hard gelatine capsules. Suitable excipients for soft gelatine capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols etc. Where the composition is in the form of a liquid pharmaceutical formulation, it will generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. 30 Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may also be included. Other suitable excipients for the manufacture of solutions and syrups are, for example, water,

- 35 -

polyols, saccharose, invert sugar, glucose, trihalose, etc. Suitable excipients for injection solutions are, for example, water, alcohols, polyols, glycerol, vegetable oils, etc. For intravenous, cutaneous or subcutaneous injection, or intracatheter infusion into the brain, the active ingredient will be in the form of a parenterally-acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers and/or other additives may be included, as required.

Uses of the compounds herein as ligands may utilise similar carriers or compositions.

Thus in aspects of the invention wherein a diaminophenothiazine (for example MTC) is used in a method of treatment or therapy of the human or animal body, that method will preferably involve administration of the effective amount of diaminophenothiazine orally.

Preferably the medicament is adapted for oral administration, and preferably is in solid dosage unit form.

Preferably the dosage will be administered orally. Preferably it will be less than or equal to 400, 300, 200, or 100 mg daily total dose. For example it may consist of dosage units of 10, 20, 30, 40, 50, 60, 60, 80, 90, 100, 110, 120, or 130 mg t.i.d. (three times a day)

Alternatively it may consist of dosage units of 10, 20, 30, 40, 50, 60, 60, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or 200 mg b.i.d. (twice a day).

Preferably the treatment is continued for equal to or at least 2, 3, or 4 weeks.

Instructions in respect of these dosages may be included in written form on or within the container of a drug product of the invention.

Where administration is in intravenous, it is preferred that the diaminophenothiazine is not MTC.

- 36 -

The disclosure of any cross-reference made herein, inasmuch as it may be required by one skilled in the art to supplement the present disclosure, is hereby specifically incorporated herein.

- 5 The invention will now be further described with reference to the following non-limiting Figures and Examples. Other embodiments of the invention will occur to those skilled in the art in the light of these.

Figures

10

Figure 1: *Samples from the purification of tsyn by ammonium sulphate fractionation and Ni-affinity chromatography.* Samples were analysed by 15% SDS-PAGE and staining with Coomassie Blue.

- 15 Figure 2. *Samples from the purification of fsyn by ammonium sulphate fractionation and DEAE-Sepharose anion exchange chromatography.* Samples were analysed by 15% SDS-PAGE and staining with Coomassie Blue.

- Figure 3. *Samples from the purification of fsyn by DEAE-Sepharose anion exchange and*
20 *CM-Sepharose cation exchange chromatography.* Samples were analysed by 15% SDS-PAGE and staining with Coomassie Blue.

- Figure 4. *Time course of fibril formation by tsyn and fsyn monitored by fluorescence of thioflavine T and primulin.* tsyn-8 at 1 mg/ml (A) and fsyn-14 (B) at 2 mg/ml in 20 mM
25 Tris.HCl, pH 7.5, plus 50 µg/ml heparin where indicated, were incubated at 37 °C with mixing. Fibril formation was assayed by adding 10 µl of the incubation to 100 µl total of 1 µM thioflavine T or primulin and the excitation spectra were measured, with the emission wavelength at 480 nm. The signal for the fluorophore alone was subtracted from the spectra before measurement of the signal at the excitation peak (420 nm for primulin, 450
30 nm for thioflavine T).

Figure 5. *The effect of MTC and ETC on the fluorescence of thioflavine T or primulin induced by assembled tsyn.* Tsyn-13 at 1 mg/ml (~95 µM) plus 50 µg/ml heparin was

- 37 -

incubated at 37 °C for 20 h, then aliquots were mixed with MTC (A, C) or ETC (B, D) to give the concentrations shown (μM) and incubated a further 1 h. 10 μl of the protein was added to 100 μl total volume of 1 μM thioflavine T (A, B) or primulin (C, D) and the excitation spectra were measured, with the emission wavelength at 480 nm. The traces shown are the result of subtracting the signal for the fluorophore plus compound from the signal for protein plus fluorophore and compound.

Figure 6. *The effect of MTC and ETC on the fluorescence of primulin induced by assembled fsyn-14.* Protein at 2 mg/ml ($\sim 140 \mu\text{M}$) plus 50 $\mu\text{g/ml}$ heparin was incubated at 37 °C for 47 h, then aliquots were mixed with MTC (A, C) or ETC (B, D) to give the concentrations shown (μM) and incubated a further 1 h. 10 μl of the protein was added to 100 μl total volume of primulin and the excitation spectra were measured, with the emission wavelength at 480 nm. The traces shown are the result of subtracting the signal for the fluorophore plus compound from the signal for protein plus fluorophore and compound.

Figure 7. *The effect of MTC and ETC on the fluorescence of thioflavine T or primulin induced by assembled tsyn or fsyn.* Fluorescence values were measured from the traces shown in Figure 5 for tsyn-8 (A) or Figure 6 for fsyn-14 (B) and normalised to the value measured without compound.

Figure 8. *The effect of MTC and ETC on the fluorescence of thioflavine T or primulin induced by assembled tsyn, assayed at different concentrations of thioflavine T and primulin.* Tsyn-16 was assembled and assayed for the effect of MTC or ETC as described for Figure 5, except that thioflavine T and primulin were at 0.2, 1 or 5 μM . Peak fluorescence values after correction for background were normalized to the value measured without compound and are plotted as a function of concentration of compound. The effect of MTC (A, B) or ETC (C, D) was monitored with thioflavine T (A, C) or primulin (B, D).

Figure 9. *Binding of aqueous phase fsyn to solid phase tsyn.* fsyn-20 at 0-10 μM was incubated with tsyn-13 bound on the solid phase at 0-2 μM and bound fsyn was detected using antibody 211.

- 38 -

Figure 10. *The effect of compounds on synuclein-synuclein binding.* Aqueous phase fsyn-10 at 5 μ M was incubated with tsyn-13 at 1 μ M in the solid phase in the presence of the compounds shown.

5

Figure 13. *Inhibitory action of DEETC on SSFsyn expression in NIE cells.* Each drug concentration (0-100 nM) was carried out in triplicate. DEETC added with dbcAMP and cells analysed after 2 days by immunoblot with mAb 42.

10 Figure 14. *Presence of additional protein bands that are detected with mAb 42 but not by mAb 211.* Lane 1, untreated; lanes 2, 3, 4, three independent plates which were differentiated using dbcAMP.

15 Figure 15. *Aggregated α -synuclein in NIE cells expressing SSFsyn.* Left panel, SSFsyn cells; right panel, non-transfected, control NIE-115 cells after dbcAMP treatment.

Figure 16. *Aggregated α -synuclein in NIE cells expressing SSFsyn.* SSFsyn cells stained with Texas red-labelled anti- α -synuclein (left); right panel, primulin labelling; middle panel, merged image showing co-localisation of antibody and primulin labelling.

20

Figure 17. *The effect of MTC on the polymerisation of tsyn.* tsyn-16 at 1 mg/ml (95 μ M) in 20 mM Tris.HCl pH 7.5, 50 μ g/ml heparin was incubated at 37 °C with mixing, in the presence of MTC at the concentrations shown. Samples (10 μ l) were taken at various times and assayed for their effect on the fluorescence of either thioflavine T (upper panel) or primulin (lower panel) at 1 μ M.

25

Figure 18. *Binding curves for different syn preparations.* Tsyn-13 at 1 μ M bound to ELISA plates was incubated with dilution series of three different syn preparations as shown. The aqueous phase buffer was 20 mM Tris.HCl, 50 mM NaCl, pH 7.5, 0.05% Tween-20, 1% fish skin gelatine.

30

- 39 -

Figure 19A and 19B: *Binding curves for syn-10 in different buffers.* Tsyn-13 at 1 μ M bound to ELISA plates was incubated with dilution series of syn-10. The aqueous phase buffers were all 20 mM, and included 50 mM NaCl, 0.05% Tween-20, 1% fish skin gelatine.

5 Figure 20: *Binding curves for different syn preparations.* Tsyn-13 at 1 μ M bound to ELISA plates was incubated with dilution series of syn preparations as shown. The aqueous phase buffer was 20 mM Tris.HCl, pH 7.0, 50 mM NaCl, 0.05% Tween-20, 1% fish skin gelatine.

10 Figure 21: *Binding curves for different syn preparations.* Tsyn-13 at 1 μ M bound to ELISA plates was incubated with dilution series of syn preparations as shown. The aqueous phase buffer was 20 mM Tris.HCl, pH 7.0, 50 mM NaCl, 0.05% Tween-20, 1% fish skin gelatine.

15 Figure 22. *Binding curves for fsyn and 20 fsyn-22 in different buffers.* A. Tsyn-13 at 1 μ M bound to ELISA plates was incubated with dilution series of fsyn-20 in 20 mM Tris pH 7.0 or 50 mM Na phosphate pH 6.0 or 5.5. B. Tsyn-13 at 1 μ M bound to ELISA plates was incubated with dilution series of fsyn-20 or fsyn-22 in Na phosphate pH 6.0. In both cases, buffers also contained 50 mM NaCl, 0.05% Tween-20 and 1% fish skin gelatine.

20

Examples


Chemical Synthesis

25 The following syntheses are provided solely for illustrative purposes and are not intended to limit the scope of the invention, as described herein.

- 40 -

Synthesis 1

Ethyl-thioninium chloride (ETC)

*N,N*-diethyl-*p*-phenylenediamine dihydrochloride

5

N,N-diethyl-*p*-phenylenediamine (5 g, 30.4 mmol) was dissolved in diethyl ether (25 cm³) and hydrochloric acid (6 cm³, 10 M) was added and the mixture was concentrated to give the title compound (7.22 g, 100%) as a red/brown solid. δ_{H} (250 MHz; D₂O): 7.68 (4H, m, ArH), 3.69 (4H, q, 7.32, NCH₂), 1.11 (6H, t, 7.32, CH₃); δ_{C} (62.9 MHz; D₂O): 12.1 (CH₃), 56.4 (NCH₂), 126.8 (ArC), 127.6 (ArC), 135.5 (ArC), 139.1 (ArC).

10

Ethyl-thioninium chloride

N,N-diethyl-*p*-phenylenediamine dihydrochloride (7.22 g, 30.4 mmol) was dissolved in water (250 cm³) and the pH adjusted to 1.6 with HCl, to which sodium sulphide (>60%) (3.95 g, 30.4 mmol) was added portionwise. The suspension was stirred until all the sodium sulphide had dissolved. A solution of iron (III) chloride (27.15 g, 100 mmol) in water (200 cm³) was prepared and half the solution was added to the mixture. An immediate colour change from light yellow to blue occurred. The solution was then aerated for 1 hour before the remaining iron (III) chloride solution was added. The mixture was cooled to 5°C and filtered to remove a light green sludge. Aqueous HCl (15 cm³, 6 M) was added to the filtrate, followed by sodium chloride (60 g), and the suspension stirred for 5 minutes before filtering to give a solid product, which was dissolved in DCM, dried over magnesium sulphate, filtered, and concentrated to give a purple/green solid (1.28 g, 22%). This purple/green solid was loaded onto a prepared C18 reverse phase column and washed with water (1 L) or until the yellow colour ceased. The product was washed off the

15

20

25


- 41 -

column with MeOH/HCl (pH 2) and concentrated to give the title compound (0.64 g, 11%) as a sticky purple solid. δ_{H} (250 MHz; D_2O): 1.26(12H, t, 6.5, CH_3), 3.56 (8H, q, 6.5, NCH_2), 7.01 (2H, s, ArH), 7.20 (2H, d, 9.25, ArH), 7.54 (2H, d, 9.25, ArH); m/z (ESI) 340.2 (100%, $[\text{M}-\text{Cl}]^+$).

5

Synthesis 2

1,9-Dimethyl-methyl-thioninium chloride (DMMTC)

10 *3-Methyl-N,N-dimethylphenylenediamine dihydrochloride*

To a 250cm^3 round bottom flask was added water (100cm^3) and the temperature was reduced to 5°C with an ice bath. To this cooled solution was carefully added sulphuric acid (98%, 22.5 g). To this solution was added 3-methyl-N,N-dimethylaniline (10 g, 74 mmol) and then sodium nitrite (5.6 g, 81.4 mmol), and the solution was stirred at room temperature for 1 hour. Iron (Fe) filings (12.8 g, 229 mmol) were added and the mixture stirred for a further 2 hours. The solution was filtered and then neutralized with saturated sodium hydrogen carbonate solution and the organics were extracted into ethyl acetate ($3 \times 100\text{cm}^3$). The extracts were dried over magnesium sulphate, filtered, and concentrated to give a brown oil. The oil was dissolved in diethyl ether (100cm^3) and concentrated hydrochloric acid (50cm^3) was added. The solution was evaporated to dryness to give the title compound (10 g, 60%) as a light tan solid. ν_{max} (KBr)/ cm^{-1} : 2849 (CH), 2821 (CH), 2543 (CH), 2444 (CH), 1586 (C=N), 1487 (CH), 1445 (CH), 1415 (CH), 1138 (CH); δ_{H} (250 MHz; D_2O): 7.59 (1H, s, ArH), 7.50 (2H, s, ArH), 3.24 (6H, s, CH_3), 2.39 (3H, s, CH_3);

- 42 -

δ_c (62.9 MHz; D₂O) 18.9 (CH₃), 48.8 (CH₃), 122.1 (ArC), 126.2 (ArC), 127.6 (ArC), 133.7 (ArC), 137.4 (ArC), 144.4 (ArC).

Dimethylmethythioninium Chloride

5

To a 500 cm³ round bottom flask was added 3-methyl-*N,N*-dimethyl-phenylene-diamine dihydrochloride (0.9 g, 4.03 mmol) which was dissolved in aqueous hydrochloric acid (50 cm³, 3 M) before sodium sulphide (>60%) (0.52 g, 4.03 mmol) was added. Iron (III) chloride hexahydrate (7.26 g, 27 mmol) was dissolved in water (50 cm³) and half of this solution was poured into the reaction mixture, giving an immediate blue colour. The solution was then aerated for 2 hours before the remaining aqueous iron (III) chloride solution was added. The mixture was cooled to 5°C and filtered; the precipitate was dissolved in boiling water (60 cm³), filtered, and cooled. Hydrochloric acid (10 cm³, 6 M) was added to the cooled solution, which was then filtered to yield the title compound

10

15


(0.22 g, 16%) as a purple/blue solid. ν_{\max} (KBr)/cm⁻¹: 2926 (CH), 1604 (C=N), 1535, 1496, 1444 (CH), 1404 (CH), 1315 (CH), 1185 (CH); δ_H (250 MHz; DMSO): 7.29 (2H, s, ArH), 7.23 (2H, s, ArH), 3.29 (12H, s, CH₃), 2.55 (6H, s, CH₃); δ_c (62.9 MHz; DMSO): 18.9 (CH₃), 41.5 (CH₃), 105.7 (ArC), 118.7 (ArC), 133.6 (ArC), 134.5 (ArC), 147.2 (ArC), 154.2 (ArC); Anal. Calcd. for C₁₈H₂₂N₃S.3H₂O: C, 51.98; H, 6.74; N, 10.11; S, 7.70. Found: C, 52.03; H, 6.59; N, 10.05; S, 7.66.

20

- 43 -

Synthesis 3

1,9-Diethyl-methyl-thioninium chloride (DEMTTC)

*N,N*-Dimethyl-*m*-ethylaniline

5

To a 100 cm³ round bottom flask was added 3-ethylaniline (10 g, 82.5 mmol), ethanol (15 cm³), sodium carbonate (11.81 g, 111.4 mmol). Methyl iodide (31.63 g, 222 mmol) was added dropwise. The mixture was then heated at 45°C for 10 hours before cooling to room temperature and adding water (100 cm³). The mixture was extracted into diethyl ether (3 x 100 cm³) and the extracts were dried over magnesium sulphate, filtered, and concentrated to give the title compound (4.68 g, 38%) as a light yellow oil. ν_{max} (neat)/cm⁻¹: 3045 (CH), 2960 (CH), 2920 (CH), 2891 (CH), 2797 (CH), 1597 (C=N), 1494 (CH), 1438 (CH), 1352 (CH), 1225 (CH); δ_{H} (250 MHz; CDCl₃): 7.22 (1H, t, 7.75, ArH), 6.63 (3H, m, ArH), 2.97 (6H, s, NCH₃), 2.63 (2H, q, 7.5, CH₂), 1.27 (3H, t, 7.5, CH₃); δ_{C} (62.9 MHz; CDCl₃): 15.8 (CH₃), 29.5 (NCH₂), 40.8 (NCH₃), 110.3 (ArC), 112.4 (ArC), 116.5 (ArC), 129.1 (ArC), 145.3 (ArC), 150.9 (ArC).

N,N-Dimethyl-*m*-ethyl-*p*-phenylenediamine dihydrochloride

To a 250 cm³ round bottom flask was added *N,N*-dimethyl-*m*-ethylaniline (4.68 g, 31.3 mmol), water (100 cm³) and hydrochloric acid (8.5 cm³, 37%) and the solution was

- 44 -

cooled to 5°C. An aqueous (80 cm³) solution of sodium nitrite (2.46 g, 3.57 mmol) was then added dropwise to the aniline mixture and stirred for 3 hours at room temperature. Iron (Fe) fillings (5.24 g, 94 mmol) and hydrochloric acid (8.5 cm³, 37%) were added and the mixture was stirred at room temperature for 3 hours. The suspension was filtered and the filtrate adjusted to pH 7 with sodium bicarbonate solution before extraction into ethyl acetate (3 x 50 cm³). The combined extracts were dried over magnesium sulphate, filtered, and concentrated to yield a brown oil. The oil was dissolved in ethanol (100 cm³) and diethyl ether (80 cm³) and hydrochloric acid (7 cm³, 37%) was added carefully to give the title compound (7.42 g, 72%) as a light tan solid. ν_{\max} (KBr)/cm⁻¹: 2976 (CH), 2894 (CH), 2859 (CH), 2753 (CH), 1583 (C=N), 1508 (CH), 1486 (CH), 1459 (CH), 1183 (CH); δ_{H} (250 MHz; D₂O): 7.66 (1H, s, ArH), 7.56 (2H, s, ArH), 3.29 (6H, s, NCH₃), 2.74 (2H, q, 7.5, CH₂), 1.25 (3H, t, 7.5, CH₃); δ_{C} (62.9 MHz; CDCl₃): 15.5 (CH₃), 25.6 (NCH₂), 48.9 (NCH₃), 122.1 (ArC), 124.6 (ArC), 128.1 (ArC), 132.6 (ArC), 143.3 (ArC), 144.9 (ArC).


15 1,9-Diethyl Methylthioninium chloride

N,N-Dimethyl-*m*-ethyl-*p*-phenylenediamine dihydrochloride (1.3 g, 5.5 mmol) was dissolved in water (50 cm³) and the solution adjusted to pH 1.6. Sodium sulphide >60% (0.71 g, 5.5 mmol) was then added portionwise to the pink solution. To the suspension was added an aqueous solution of iron (III) chloride (2.23 g, 8.2 mmol in 50 cm³ of water) and there was an immediate colour change to purple. The solution was then aerated for 1 hour before a second portion of iron (III) chloride solution (2.23 g, 8.2 mmol in 50 cm³ of water) was added. The solution was cooled to 5°C before filtering and washing the precipitate with water. To the filtrate was added sodium chloride (50 g) and the solution was stirred for 10 minutes, and the colour changed to red/purple as the product was salted out. The suspension was filtered and the solid dissolved in dichloromethane (100 cm³) and methanol (10 cm³) before drying over magnesium sulphate. Filtration and concentration gave the title compound (0.15 g, 15%) as a green solid. ν_{\max} (KBr)/cm⁻¹: 3408 (CH), 2613 (CH), 1606 (C=N), 1399 (CH), 1316 (CH); δ_{H} (250 MHz; D₂O): 6.55 (2H, s, ArH), 6.23 (2H, s, ArH), 2.92 (12H, s, NCH₃), 2.56 (4H, q, 7.5, CH₂), 0.99 (6H, t, 7.5, CH₃); (ESI), 340.4 (100%, [M - Cl]⁺). Optionally, flash column chromatography was performed to remove iron chloride residues, with 10% methanol: 90% dichloromethane as eluent and using silica 40-63 μ 60 Å.

- 45 -

Synthesis 4

1,9-Dimethyl-ethyl-thioninium chloride (DMETC)




- 5 *N,N*-Diethyl-3-methyl-4-phenylenediamine dihydrochloride (10.74 g, 50 mmol) was dissolved in water (400 cm³) and the pH adjusted to 1.6, which then had sodium sulphide (>60%) (3.90 g, 50 mmol) added. Iron (III) chloride (20.28 g, 75 mmol) was added as an aqueous solution (175 cm³) giving an immediate colour change from yellow to deep blue. The mixture was aerated for 1 hour before a second aliquot of aqueous iron (III) chloride
- 10 (20.28 g, 75 mmol in 175 cm³) was added. The solution was cooled to 5°C and held at that temperature for 1 hour before filtering. The filtrate had sodium chloride (200 g) added and was filtered to yield the crude product as a blue/purple solid. The crude solid was purified by column chromatography (eluent being 10% MeOH, 90% DCM using silica 40-63μ 60Å) to give the title compound (0.80 g, 4%) as a green/purple solid. ν_{\max} (KBr)/cm⁻¹: 2971 (CH), 2921 (CH), 2865 (CH), 1600 (C=N), 1412 (CH), 1326 (CH); δ_{H} (250 MHz; D₂O): 6.62 (2H, s, ArH), 6.39 (2H, s, ArH), 3.30 (8H, q, NCH₂), 1.89 (6H, s, ArCH₃), 1.09 (12H, t, CH₃);
- 15 δ_{C} (62.9 MHz; D₂O) 12.6 (CH₃), 18.0 (CH₃), 46.2 (NCH₂), 103.6 (ArC), 117.1 (ArC), 132.3 (ArC), 133.9 (ArC), 147.3 (ArC), 151.9 (ArC); m/z (ESI) 368.1 (100%, [M-Cl]⁺).

- 46 -

Synthesis 5

1,9-Diethyl-ethyl-thioninium chloride (DEETC)

5 *N,N*-Diethyl-*m*-ethylaniline

To a 100 cm³ round bottom flask was added 3-ethylaniline (5.0 g, 41.3 mmol), ethanol (7.5 cm³), sodium carbonate (5.9 g, 55.7 mmol). Ethyl iodide (17.38 g, 111.4 mmol) was added dropwise. The mixture was then heated at 45°C for 12 hours before cooling to room temperature and adding water (50 cm³). The mixture was extracted into diethyl ether (3 x 50 cm³) the extracts were dried over magnesium sulphate, filtered, and concentrated to give the title compound (7.03 g, 96%) as a light yellow oil. δ_{H} (250 MHz; CDCl_3): 7.20 (1H, dd, 9, 7.25, ArH), 6.60 (3H, m, ArH), 3.43 (4H, q, 7, NCH_2), 2.69 (2H, q, 7.25, CH_2), 1.32 (3H, t, 7.5, CH_3), 1.23 (6H, t, 7, CH_3); δ_{C} (62.9 MHz; CDCl_3): 12.7 (CH_3), 15.8 (CH_3), 29.5 (CH_2), 44.4 (NCH_3), 109.4 (ArC), 111.4 (ArC), 115.1 (ArC), 129.2 (ArC), 145.4 (ArC), 147.9 (ArC).

N,N-Diethyl-*m*-ethyl-*p*-phenylenediamine dihydrochloride

To a 250 cm³ round bottom flask was added *N,N*-diethyl-*m*-ethylaniline (5 g, 28.2 mmol), water (50 cm³) and hydrochloric acid (9 cm³, 37%) and the solution was cooled to 5°C. An

- 47 -

aqueous (20 cm³) solution of sodium nitrite (2.14 g, 31.0 mmol) was then added dropwise to the aniline mixture and stirred for 1 hour at low temperature. Iron (Fe) fillings (4.72 g, 84.6 mmol) and hydrochloric acid (9 cm³, 37%) were added and the mixture stirred below 30°C for 2 hours. The suspension was filtered and the filtrate adjusted to pH 7 with sodium bicarbonate solution before extraction into ethyl acetate (3 x 50 cm³). The combined extracts were dried over magnesium sulphate, filtered, and concentrated to yield a brown oil. The crude oil was purified by column chromatography (eluent being ethyl acetate using silica 40-63µ 60Å) giving the phenylenediamine as a brown oil (2.2 g, 41%). The oil was dissolved in diethyl ether (50 cm³) and hydrochloric acid added (2.5 cm³, 37%) and the solution was concentrated to give the title compound (2.76 g, 41%) as a light brown solid. δ_H (250 MHz; D₂O): 7.50 (3H, m, ArH), 3.59 (4H, q, 7.25, NCH₂), 2.69 (2H, q, 7.5, CH₂), 1.20 (3H, t, 7.5, CH₃), 1.03 (6H, t, 7.25, CH₃); δ_C (62.9 MHz; D₂O): 12.1 (CH₃), 15.5 (CH₃), 25.5 (CH₂), 56.3 (NCH₂), 123.9 (ArC), 126.0 (ArC), 127.9 (ArC), 133.1 (ArC), 139.4 (ArC), 143.3 (ArC).


1,9-Diethyl Ethylthioninium chloride

N,N-Diethyl-*m*-ethyl-*p*-phenylenediamine dihydrochloride (2 g, 7.5 mmol) was dissolved in water (75 cm³) and the solution adjusted to pH 1.6. The pink solution then had sodium sulphide (>60%) (1.35g, 10.4mmol) added portion-wise. To the suspension was added an aqueous solution of iron (III) chloride (4.22 g, 15.6 mmol in 35 cm³ of water) where there was an immediate colour change to purple. The solution was then aerated for 1 hour before a second portion of iron (III) chloride (4.22 g, 15.6 mmol in 35 cm³ of water) solution was added. The solution was cooled to 5°C before filtering and washing the precipitate with water. The precipitate was also washed with ethanol and the ethanol concentrated to give a sticky purple solid. To the aqueous filtrate was added sodium chloride (50 g) and the solution was stirred for 10 minutes whereby the colour changed to red/purple as the product was salted out. The suspension was filtered and the solid dissolved in dichloromethane (100 cm³) and methanol (10 cm³) before drying over magnesium sulphate. Filtering and concentration with the ethanol soluble product gave the title compound (0.06 g, 3%) as a purple solid. δ_H (250 MHz; D₂O): 6.73 (2H, s, ArH), 6.48 (2H, s, ArH), 3.45 (8H, brdq, NCH₂), 2.46 (4H, q, 7.5, CH₂), 1.17 (12H, brdt, CH₃), 0.93 (6H, t, 7.5, CH₃); *m/z* (ESI) 396.2 (100%, [M-Cl]⁺). Optionally, flash column chromatography was

- 48 -

performed to remove iron-chloride residues, with 10% methanol: 90% dichloromethane as eluent and using silica 40-63 μ 60Å.

Synthesis 6




- A stirred mixture of *N,N*-diethyl-*p*-phenylenediamine (5.0 g, 30.4 mmol) in H₂O (100 cm³) and H₂SO₄ (conc., '98 %', 1 cm³) was treated with non-reducing ZnCl₂ solution (ZnCl₂, 7.60 g, 55 mmol in 15 cm³ of H₂O with Na₂Cr₂O₇·2H₂O, 100 mg) to produce a reddish reaction mixture.
- 10 Additions of Al₂(SO₄)₃·16H₂O solution (5.80 g, 9.2 mmol in 10 cm³ of H₂O), Na₂S₂O₃·5H₂O solution (8.0 g, 32.2 mmol in 10 cm³ H₂O) and one-third of a solution of Na₂Cr₂O₇·2H₂O (8.7 g, 29.2 mmol in 15 cm³ of H₂O) were followed by a rapid rise in temperature to 40°C. A solution of *N,N*-diethylaniline (3.0 g, 20.1 mmol in conc. HCl, 4 cm³) was added, and followed by an addition of the remaining Na₂Cr₂O₇·2H₂O solution.
- 15 A dark green precipitate was observed. The temperature was rapidly raised to 75°C, after which a slurry of activated MnO₂ (3.80 g, 44.7 mmol in 5 cm³ of H₂O) was added. The temperature was raised to 85°C, and left to stir at that temperature for 30 minutes. A blue solution with precipitate was observed. The reaction mixture was cooled to 50°C and H₂SO₄ (conc., 11cm³) was slowly added. The reaction was further cooled to 20°C, and
- 20 vacuum filtered to recover the precipitate, which was then washed with brine (saturated salt water). This black solid was re-dissolved in H₂O (250 cm³) at 100°C, and cooled, followed by vacuum filtration to remove insolubles. The filtrate was treated with ZnCl₂ (4 g) and

- 49 -

NaCl (23 g) and left in the refrigerator for 16 hours, after which the resulting precipitate was recovered by vacuum filtration, washed with brine (30 cm³), and dried in a vacuum oven for 3 hours, to give the title compound (5.7 g, 71 %) as a rusty red powder. δ_H (250 MHz, D₂O): 1.20 (12H, br t, CH₃), 3.50 (8H, br q, CH₂), 6.80 (2H, s, ArH), 7.05 (2H, br d, ArH) and 7.30 (2H, br d, ArH). See, for example, Fierz-David and Blangley, 1949, "F. Oxazine and Thiazine Dyes," in: Fundamental Processes of Dye Chemistry, published by Interscience (London, UK), pp. 308-314.

Synthesis 7

Methyl-thioninium Iodide (MTI)




Methyl-thioninium chloride (2.00 g, 6.25 mmol) was dissolved in water (50 cm³) and potassium iodide (1.56 g, 9.4 mmol) was added with stirring. A precipitate formed, which was filtered and the solid was recrystallised from boiling water (50 cm³) to yield the title compound (1.98 g, 77%) as fine green needles. δ_H (250 MHz; DMSO): 7.88 (2H, br d, ArH), 7.49 (4H, br s, ArH), 3.37 (12H, s, CH₃). Analysis for C₁₆H₁₈N₃SI: C, 46.72; H, 4.41; N, 10.22; S, 7.80; I, 30.85; Found: C, 46.30; H, 4.21; N, 10.14; S, 7.86; I, 29.34.

- 50 -

Synthesis 8

Methyl-thioninium Iodide Hydrogen Iodide Mixed Salt (MTI.HI)




- Methyl-thioninium iodide (0.50 g, 1.22 mmol) was dissolved in methanol (20 cm³) and
- 5 methyl iodide (1.90 g, 13.37 mmol) was added while stirring. The mixture was heated at reflux for 18 hours before additional methyl iodide (0.42 g, 6.69 mmol) was added and the mixture was once again heated to reflux and stirred for 8 hours. The mixture was cooled to room temperature, giving a solid that was filtered and washed with methanol to yield the title compound (0.30 g, 46%) as bronze green solid. δ_H (250 MHz; DMSO): 7.82 (2H, d, J =
- 10 8.5, ArH), 7.42 (4H, s, ArH), 3.34 (12H, s, CH₃). δ_C (62.9 MHz; DMSO): 153.8 (ArC), 137.9 (ArC), 134.9 (ArC), 133.5 (ArC), 119.1 (ArC), 118.8 (ArC), 106.9 (ArC), 106.6 (ArC), 41.1 (NCH₃).

- 51 -

Synthesis 9

Ethyl-thioninium iodide (ETI)




- A stirred mixture of *N,N*-diethyl-*p*-phenylenediamine (10.0 g, 61 mmol) in aqueous
- 5 hydrochloric acid (0.5 M, 200 cm³) was adjusted to pH 2 with aqueous sodium hydroxide (10%). The diamine solution was cooled to 5°C before the addition of aqueous $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$ (16.65 g, 67 mmol in 20 cm³ H₂O). An aqueous solution of $\text{Na}_2\text{Cr}_2\text{O}_7 \cdot 2\text{H}_2\text{O}$ (7.27 g, 24 mmol in 35 cm³ of H₂O) was added dropwise to the mixture over a 15 minute period giving a black suspension. The suspension was stirred at 5°C for 1 hour (pH = 8.07, T = 3.7°C).
- 10 A solution of *N,N*-diethylaniline (8.25 g, 61 mmol), H_2SO_4 (6 g) and water (10 cm³) was cooled to 5°C before addition to the suspension. An aqueous solution of $\text{Na}_2\text{Cr}_2\text{O}_7 \cdot 2\text{H}_2\text{O}$ (19.09 g, 64 mmol in 50 cm³ of H₂O) was then added dropwise to the mixture over a 20 minute period giving a thick dark green suspension. The mixture was stirred at 5°C for 2 hours (pH = 6.75, T = 6°C) before filtering. The green purple solid
- 15 obtained was washed with water (2 x 50 cm³). The solid was slurried in aqueous hydrochloric acid (300 cm³, pH 2) giving a suspension with a pH = 6.37 at 22°C. To the suspension was added CuSO_4 (1.52 g, 6.1 mmol) and the mixture heated to 90°C where a deep blue solution formed. After stirring at this temperature for 1 hour the mixture was cooled to 25°C and filtered. The solid was washed with water (2 x 50 cm³), the filtrate was
- 20 adjusted from pH 6.33 to pH 2.00, T = 25°C with hydrochloric acid (5 M). The deep blue solution was heated to 80°C and potassium iodide (14 g) was added and upon cooling an orange purple precipitate was deposited. Filtration gave a purple powder (8.8 g, 31%),

- 52 -

which was recrystallised from hot ethanol (400 cm³) to give the title compound as fine purple needles. Mp 211°C; ν_{\max} (KBr)/cm⁻¹: 3574 (CH), 3484 (CH), 3028 (CH), 2965 (CH), 1662 (C=C), 1539 (CH), 1474 (CH), 1346 (CH); δ_c (62.9 MHz, CDCl₃): 1.33 (12H, t, 7, CH₃), 3.72 (8H, q, 7, NCH₂), 7.23 (2H, d, 9.75, ArH), 7.41 (2H, s, ArH), 7.83 (2H, d, 9.75, ArH); δ_H (62.9 MHz, CDCl₃): 152.4, 138.8, 135.7, 135.2, 118.3, 106.4, 46.8, 13.2.

Synthesis 10

Ethyl-thioninium iodide Hydrogen Iodide Mixed Salt (ETI.HI)




- 10 Ethyl-thioninium iodide (2.00 g, 4.28 mmol) was dissolved in ethanol (100 cm³) and ethyl iodide (27.35 g, 175 mmol) was added while stirring. The mixture was heated at reflux for 18 hours, then cooled to room temperature, giving a precipitate that was filtered and washed with ethanol to yield the title compound (1.02 g, 40%) as a bronze solid. δ_H (250 MHz; D₂O): 7.90 (2H, br d, ArH), 7.42 (4H, s, ArH), 2.45 (8H, br q, NCH₂), 1.23 (12H, br t, CH₃).

- 53 -

Synthesis 11

Ethyl-thioninium nitrate (ETN)



- A stirred mixture of *N,N*-diethyl-*p*-phenylenediamine (10.0 g, 61 mmol) in aqueous
- 5 hydrochloric acid (0.5 M, 200 cm³) was adjusted to pH 2 with aqueous sodium hydroxide (10%). The diamine solution was cooled to 5°C before the addition of aqueous $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$ (16.65 g, 67 mmol in 20 cm³ H₂O). An aqueous solution of $\text{Na}_2\text{Cr}_2\text{O}_7 \cdot 2\text{H}_2\text{O}$ (7.27 g, 24 mmol in 35 cm³ of H₂O) was added dropwise to the mixture over a 15 minute period giving a black suspension. The suspension was stirred at 5°C for 1 hour (pH = 8.07,
 - 10 T = 3.7°C). A solution of *N,N*-diethylaniline (8.25 g, 61 mmol), H_2SO_4 (6 g) and water (10 cm³) was cooled to 5°C before addition to the suspension. An aqueous solution of $\text{Na}_2\text{Cr}_2\text{O}_7 \cdot 2\text{H}_2\text{O}$ (19.09 g, 64 mmol in 50 cm³ of H₂O) was then added dropwise to the mixture over a 20 minute period giving a thick dark green suspension. The mixture was stirred at 5°C for 2 hours (pH = 6.75, T = 6°C) before filtering. The green purple solid
 - 15 obtained was washed with water (2 x 50 cm³). The solid was slurried in aqueous hydrochloric acid (300 cm³, pH 2) giving a suspension with a pH = 6.37 at 22°C. To the suspension was added CuSO_4 (1.52 g, 6.1 mmol) and the mixture heated to 90°C wherein a deep blue solution formed. After stirring at this temperature for 1 hour, the mixture was cooled to 25°C and filtered. The solid was washed with water (2 x 50 cm³), the and the
 - 20 filtrate was adjusted from pH 6.33 to pH 2.00, T = 25°C with hydrochloric acid (5 M). The deep blue solution was heated to 80°C and had sodium nitrate (50 g) added and was allowed to cool to 25°C slowly while stirring gently. The product was filtered as green

- 54 -

needles (6.80 g, 28%). δ_H (250 MHz, $CDCl_3$): 1.36 (12H, t, 7, CH_3), 3.72 (8H, q, 7, NCH_2), 7.23 (2H, d, 9.5, ArH), 7.39 (2H, s, ArH), 7.89 (2H, d, 9.5, ArH); δ_H (62.9 MHz, $CDCl_3$): 152.5, 138.8, 135.7, 135.6, 118.1, 106.4, 46.6, 12.9.

5 Biological Studies

Methods

Purification of α -synuclein proteins

10

Two plasmids for expression of α -synuclein in *E. coli* were constructed. The core aggregation domain of α -synuclein (amino acids 31-109) was expressed with an N-terminal polyhistidine tag (tsyn), which allows its purification on a Ni-chelating column. Full-length α -synuclein (syn) was expressed without a tag, and purified by ion exchange

15 chromatography on DEAE Sepharose, and in some cases followed by purification on CM-Sepharose. For both proteins, the bacterial extract was first enriched by taking a 30-50% ammonium sulphate cut. The proteins eluted from the column were dialysed against 20 mM CAPS, pH 9.5 or 20 mM Tris.HCl, pH 7.5, 50 mM NaCl (see Table 1 for details), and stored at -70 °C.

20

Fluorescence assay for filament assembly

α -Synuclein proteins (tsyn or fsyn) were incubated at 37°C for the times indicated in Figure legends with mixing to induce fibril formation. In some cases, 50 μ g/ml heparin was

25 included to enhance fibril formation.

Samples of 10 μ l were then diluted to 100 μ l with water, plus thioflavine T or primulin at 1 μ M, or 0.2 or 5 μ M in some cases. Fluorescence excitation spectra were measured in 96 well plates in a Varian Carey Eclipse Fluorescence Spectrophotometer, with the

30 emission wavelength at 480 nm. Excitation spectra were corrected for the signal measured without tsyn and the peak signal measured from the spectra. The data were normalised to the value measured without compound and P50 values measured from plots of normalised fluorescence against concentration of compound.

- 55 -

ELISA assay for synuclein-synuclein binding

A solid phase assay was used to measure self-association of α -synuclein. tsyn diluted in carbonate buffer (pH 8.5) was bound to the assay plate, and full-length α -synuclein (fsyn) was added in the aqueous phase. The aqueous phase binding buffer was 50 mM Na-phosphate, pH 6.0, 20 mM NaCl, 0.05% Tween-20, 1% fish skin gelatine. Bound fsyn was detected using a commercial antibody (211) that does not recognise tsyn.

Example 1. Purification of α -synuclein

Figure 1 shows samples from the purification of tsyn analysed by SDS-PAGE and staining with Coomassie Blue. The Ni-affinity column provides a very efficient purification; the final purified protein (tsyn-8 in Fig. 1) is greater than 95% pure, with a yield of 44 mg protein from a 750 ml culture of bacteria.

Figure 2 shows the purification of fsyn on DEAE-Sepharose. The final protein using this method is not so pure (fsyn-9 in Fig. 2).

Figure 3 shows the purification of fsyn on DEAE Sepharose followed by CM-Sepharose. This method produces protein of >95% purity, but the yield is lower (12 mg protein compared to 85 mg with DEAE Sepharose alone).

Several different preparations of α -synuclein have been used in the assays described below, and brief details of their purification are summarised in Table 1.

Table 1. *Purification details of synuclein preparations*

Prep	Purification details
tsyn-8	AS cut, Ni column, dialysed 20 mM CAPS, pH 9.5
fsyn-9	AS cut, DEAE column, dialysed 20 mM Tris, pH 7.5, 50 mM NaCl
tsyn-16	AS cut, Ni column, dialysed 20 mM Tris, pH 7.5
fsyn-10,17,18,19	AS cut, DEAE column, CM column, dialysed 20 mM Tris.HCl, 50 mM NaCl, pH 7.5
fsyn-14	AS cut, DEAE column, dialysed 20 mM Tris.HCl, pH 7.5
fsyn-15	AS cut, DEAE column, SP column, dialysed 20 mM Tris.HCl, pH 7.5
fsyn-20,22	AS cut, DEAE column, CM column, dialysed 20 mM CAPS, pH 9.5

Protein prepared by heat treatment was inactive in the assay. Protein prepared by DEAE ion-exchange chromatography was active. A further purification step on CM- or SP-Sephacrose was therefore carried out. It was found that CM- Sepharose gave the cleanest preparation, but with lower yield than SP-Sephacrose. These preparations were compared for binding activity (see Example 9). Table 1 summarises the synuclein preparations used for these experiments.

Example 2. Assay of synuclein assembly by fluorescence

It has been reported that assembly and fibril formation of α -synuclein enhances the fluorescence of thioflavine T. We tested the effect of α -synuclein on the fluorescence of this and also primulin. The proteins were induced to assemble by incubation at 37°C, with and without 50 μ g/ml heparin and samples were assayed with 1 μ M thioflavine T or primulin at various time points.

Figure 4 shows the time course of assembly of tsyn and fsyn preparations. In the absence of heparin, there is very little appearance of thioflavine T fluorescence with either protein (Fig. 4A, 4B). There is appearance of a primulin signal over 20-30 h with tsyn protein in the

- 57 -

absence of heparin, in the presence of heparin there is no lag phase for appearance of the primulin signal but the final extent of the fluorescence is similar with and without heparin (Fig. 4A). Heparin stimulates appearance of a thioflavine T signal for tsyn to a similar extent as primulin (Fig. 4A). There is a very slow appearance of both thioflavine T and primulin signals for fsyn in the absence of heparin. In the presence of heparin, there is appearance of a signal with both fluorophores, but with a longer lag phase than seen with tsyn (Fig. 4B). The lag phase for appearance of the primulin signal is shorter than that for thioflavine T, but the final extent of the thioflavine T signal is greater than that for primulin (Fig. 4B). The difference in signal between primulin and thioflavine T indicates that these two fluorophores are detecting different assembly states of synuclein, and is consistent with the idea that primulin detects an early precursor state of assembly prior to fibril formation, which is detected by thioflavine T.

Example 3. Assay of fibril disruption by MTC and ETC

The fluorescence effects have been used to assay the effect of compounds MTC and ETC on assembled α -synuclein.

Figure 5 shows the effect of MTC and ETC on the fluorescence signal of either thioflavine T or primulin induced by assembled tsyn, and the peak fluorescence values from the traces are shown as a function of concentration of compound in Figure 7A. Figure 6 shows the effect of MTC and ETC on the fluorescence signal of either thioflavine T or primulin induced by assembled fsyn, and the peak fluorescence values from the traces are shown as a function of concentration of compound in Figure 7B. The P50 values measured from the graphs in Figure 7 are summarised in Table 2.

- 58 -

Table 2. *P50 values for the inhibition of α -synuclein-dependent fluorescence of thioflavine T or primulin by MTC and ETC.*

protein	fluorophore	compound	P50 (μ M)
tsyn-13	thioflavine T	MTC	18.5
tsyn-13	thioflavine T	ETC	30.3
tsyn-13	primulin	MTC	12.9
tsyn-13	primulin	ETC	24.1
fsyn-14	thioflavine T	MTC	3.8
fsyn-14	thioflavine T	ETC	25.3
fsyn-14	primulin	MTC	23.5
fsyn-14	primulin	ETC	30.6

- 5 The effect of the compounds on thioflavine T and primulin fluorescence could be due to competition for the fluorescence ligand rather than disruption of the fibrils. To test this, the experiment was done at three different concentrations of fluorophore, since the P50 will be dependent on fluorophore concentration only if the effect is due to competition. Data from one experiment is shown in Figure 8, and the mean P50s from all experiments is shown in
- 10 Table 3. There is not a significant difference in P50 values measured over a 25-fold difference in fluorophore concentrations, indicating that the effect of the compounds is due to fibril disruption.

Table 3. *P50 values for the inhibition of α -synuclein-dependent fluorescence of thioflavine T or primulin by MTC and ETC. Effect of fluorophore concentration.*

compound	fluorophore	[fluorophore] (μ M)	P50 (μ M)	SEM	N
MTC	thioflavine T	0.2	11.6	6.7	3
MTC	thioflavine T	1.0	12.7	3.9	3
MTC	thioflavine T	5.0	20.2	0.0	2
MTC	primulin	0.2	41.7	8.2	2
MTC	primulin	1.0	35.1	2.2	3
MTC	primulin	5.0	31.2	0.7	2
ETC	thioflavine T	0.2	3.2	2.5	3
ETC	thioflavine T	1.0	9.0	6.6	3
ETC	thioflavine T	5.0	17.6	2.3	2
ETC	primulin	0.2	41.7	8.2	2
ETC	primulin	1.0	34.0	3.3	3
ETC	primulin	5.0	32.3	-	1

5 Example 4. The effect of compounds on the assembly of α -synuclein aggregates in vitro.

As well as affecting assembled α -synuclein aggregates, MTC also inhibits the assembly of α -synuclein into aggregates, as determined by competition of the binding of primulin.

Optimal conditions for the assembly of aggregates from tsyn and fsyn has been determined and the inhibitory effect of MTC is shown in the Figure (New Figure A). tsyn (1mg/ml in 20 mM Tris.HCl, pH 7.5 + 50 μ g/ml heparin) was assembled at 37° C for 24 hr. MTC inhibits tsyn assembly at concentrations greater than 5 μ M (\circ , open circles). fsyn was assembled under the same conditions, except that the concentration of fsyn was 2 mg/ml and incubation was for 120 hr. MTC shows a greater inhibitory effect with fsyn than with tsyn, with inhibition occurring at 0.05 μ M with the former (\bullet , closed circles).

Inhibition of α -synuclein aggregation (fsyn; using the primulin assay as described above) by MTC and ETC was comparable and greater than that observed with DEMTC and DEETC. All these compounds completely inhibited assembly at a concentration of 50 μ M.

- 60 -

Table 4. *Effect of diaminophenothiazines on α -synuclein aggregation using fsyn.*

		Fluorescence with drug concentration (μ M):			
		0	0.5	5.0	50
5	Compound				
	MTC	311.3	145.9	24.2	1.2
10	ETC	311.3	190.6	44.6	1.5
	DEMTC	311.3	295.7	23.4	0.0
15	DEETC	311.3	369.2	18.2	0.0

Thioflavine T has also been used to monitor fsyn assembly. The evolution of the thioflavine T signal is slower than the primulin signal, but reaches a higher level and appears to be reporting elongation of fibrils rather than formation of aggregates. At later stages of assembly (160 h) when the thioflavine T signal has reached a plateau, the thioflavine T signal is more sensitive to inhibition by MTC than the primulin signal, with a significant effect being observed at 0.05 μ M. This is shown in Figure 12.

Example 5. Assay of α -synuclein binding by a solid phase ELISA assay

The two α -synuclein proteins were also used in a binding assay. The tsyn is bound in the solid phase, and full-length fsyn is added in the aqueous phase. An antibody against a C-terminal epitope in α -synuclein that does not recognise tsyn is used to quantify bound fsyn.

Figure 9 shows the aqueous and solid phase binding curves for fsyn-20 binding to tsyn-13. Binding of fsyn-20 plateaus at $\sim 5 \mu$ M and tsyn-13 binding plateaus at $\sim 2 \mu$ M. These

- 61 -

concentrations were used to test the effect of various thioninium chlorides and flavones on synuclein-synuclein binding.

The inhibition curves are shown in Figure 10, and the B50 values calculated from the curves are summarised in Table 4.

All the flavones tested in the binding assay have good inhibitory activity, whereas although most of the thioninium chlorides are active, namely MTC, ETC, DMMTC, DEMTC, DMETC, DEETC, thionine and tolonium chloride, others such as azure A and azure B are inactive.

Table 5. P50 and B50 values for the inhibition of synuclein-synuclein binding. P50 measured with 1 µg/ml tsyn-16 assembled in 20 µM Tris.HCl (pH 7.5) with heparin (50 µg/ml) and assayed with either of two fluorophores (1 µM), thioflavine T or primulin. B50 measured with 1 µM tsyn-16 (solid phase) and 5 µM fsyn-20 (aqueous phase), using 50 mM sodium phosphate buffer (pH 6.0) containing 20 mM NaCl.

Compound	P50 (µM) using thioflavone T	P50 (µM) using primulin	B50 (µM) in α-synuclein binding assay
MTC	15.4 ± 3.7 (6)	30.4 ± 3.0 (6)	130.8 ± 15.0 (5)
ETC	7.3 ± 3.8 (5)	26.0 ± 6.0 (5)	3.8 ± 0.3 (2)
DMMTC	0.2 (1)	20.7 ± 7.8 (2)	4.9 (1)
DEMTC			0.8 ± 0.4 (2)
DMETC	0.9 ± 0.2 (2)	5.0 ± 0.9 (3)	0.5 ± 0.1 (2)
DEETC			0.4 (1)
Azure A	3.8 ± 0.9 (2)	17.9 ± 2.0 (2)	> 500
Azure B			> 500
Thionine	3.8 ± 0.3 (5)	23.0 ± 2.2 (5)	32.2 ± 11.7 (2)
Tolonium chloride	1.5 ± 0.2 (5)	8.8 ± 3.3 (5)	3.3 ± 0.3 (2)
Baicalein	47.6	28.2	2.5 (1)
7-Hydroxy-flavone	> 50	> 50	35.3 (1)
7-hydroxy-3-methyl-flavone			141.2 (1)
Fisetin			88.2 (1)
Geraldol	>50	> 50	60.8 (1)

- 62 -

Compound	P50 (μ M) using thioflavone T	P50 (μ M) using primulin	B50 (μ M) in α -synuclein binding assay
Rhamnetin	39.4	34.7	11.4 \pm 9.7 (2)
Robinetin	21.1	27.6	39.2 (1)
3,3',4'-Trihydroxy-flavone			43.1 (1)
Primulin		(1.0)	
Thioflavine T	(1.0)		

Example 6. Cell based assay for α -synuclein aggregation

- The cell based assay employed a mouse neuroblastoma cell line NIE-115 that had been engineered to express full-length α -synuclein incorporating an N-terminal signal sequence (SSfsyn) to target incorporation of the protein into the membrane (see WO02/059150). When the cells were differentiated with dibutyryl cyclicAMP (dbcAMP) (1mM) expression of the α -synuclein protein was increased.
- α -synuclein protein was detected by immunoblot using various anti- α -synuclein antibodies. These included: mAb 42 (BD Biosciences Cat No. 610787) that recognises an epitope within tsyn (residues 31-109 of α -synuclein). In addition to α -synuclein, mAb 42 also reacts non-specifically with a protein of higher molecular mass (the protein is not recognised by other anti- α -synuclein antibodies). This protein was used as an estimate of cell numbers as the level of this protein in cells correlates with cell density. Drugs were tested with these cells and an inhibitory activity (EC50) was calculated by determining the drug concentration in which the ratio of α -synuclein to the non-specific band fell to 50% of the value for cells treated with dbcAMP alone.
- The timing of the addition of the dbcAMP and drug was varied and the length of time that the cells are left in the presence of drug + dbcAMP before collecting cells was also be varied. A typical result is shown in Fig 13 for DEETC.

MTC was inhibitory when the cells had been left for more than 2 days in the presence of MTC and dbcAMP. The most effective compound was DEETC that inhibited in the nM

- 63 -

range; DEMTC, DMETC and ETC also show inhibitory activity (Table 6). The flavone, rhamnetin, was also inhibitory.

Table 6. *Inhibition of FSyn expression in NIE cells differentiated with dbcAMP by phenothiazine compounds.*

Compound	EC50 (μ M)	LD50 (μ M)
MTC	1.17 ± 0.3 (7)	62
ETC	0.08 ± 0.02 (13)	38
DMMTC	0.10 ± 0.02 (4)	1.2
DMETC	0.05 ± 0.04 (2)	6
DEMTC	0.012 ± 0.005 (3)	3
DEETC	0.004 ± 0.001 (3)	1.0
Tolonium chloride	0.26 ± 0.05 (3)	1.2
Azure A	0.32	8.8
Azure B	0.25	0.8
Rhamnetin	2.5	31

Example 7. Truncation and aggregation of α -synuclein in a cell-based assay.

When DH60.21 NIE cells were differentiated using dbcAMP, there was increased expression of SSFsyn, as detected using mAb 42 (recognising the core of α -synuclein) or mAb 211 (recognising a C-terminal epitope of α -synuclein). In addition, two lower molecular mass bands of approximately 15 and 16 kDa were produced. The latter may correspond to Fsyn lacking the signal sequence. While the larger of these two proteins was detected by both of these antibodies, the 15 kDa band was, at best, only weakly detected using mAb 211. A typical example is shown in Figure 14. This suggests that this is a protein that has been C-terminally truncated. A further 22 kDa band with an apparent mobility greater than FSyn was also observed, but only using mAb 42 and not mAb 211 (Figure 14). The ratio of the 22-kDa to SSFsyn bands differed significantly depending upon the antibody used ($p < 0.001$; Table 7). This suggests the presence of aggregated

- 64 -

synuclein that has been truncated at both N- and C-termini. This 22-kDa band was also observed in SH-SY5Y neuroblastoma cells transfected with SSFsyn.

Table 7. *Presence of aggregated and truncated α -synuclein in cells as demonstrated by the absence of reactivity with the C-terminal mAb 211.*

Detection antibody	Mean ratio of 22 kDa:SSFsyn (\pm SEM; n=30)
mAb 42	0.14 \pm 0.02
mAb 211	0.01 \pm 0.003

When cells expressing SSFsyn were examined by fluorescence microscopy, abundant expression was observed, including material of a granular nature, suggestive of aggregated protein (Figure 15). Furthermore the aggregates observed in cells were simultaneously recognised by primulin, a fluorophore that binds to aggregated proteins (Figure 16). This confirms other studies that demonstrate that α -synuclein aggregation arises after differentiation of SH-SY5Y neuroblastoma cells (Hasegawa et al. 2004; Brain Res. 1013:51-59)

Example 8. Effect of MTC on α -synuclein oligomer assembly and on α -synuclein binding of in vitro.

Referring to the the results of P50 and B50 measurements in synuclein assays in Table 5, it can be seen that thioninium chlorides generally have lower P50s than flavones in the synuclein assays, although some of the flavones have low B50s, comparable to the thioninium chlorides. This suggests that although both classes of compound are effective in inhibiting the synuclein self aggregation reaction, only thioninium chlorides have the ability to disrupt preformed aggregates.

As a further assay of the activity of MTC, its effect was measured over the time course of assembly of tsyn (Figure 17). As shown in Figure 17, in the absence of MTC, there is a lag phase before assembly of \sim 2 hours. The fluorescence signal peaks at 4-5 hours, then

- 65 -

gradually decreases over 20 hours. The time course of appearance of fluorescence is similar for the two fluorophores. In the presence of a low concentration of MTC (0.05 μ M), the lag phase before start of assembly is reduced to 1 hour and the final fluorescence signal after 24 hours is higher. In particular, with primulin, the maximum fluorescence is higher with primulin compared to thioflavine T. In the presence of 0.5 μ M MTC, assembly is slower than control, but the final fluorescence level is higher than control. At 5 μ M MTC, the time course for assembly over the first 4-5 hours is similar to the control, but then there is a more rapid fall off in the fluorescence, although the final fluorescence level is similar to control. At 50 μ M MTC, there is no assembly over the time course of the experiment. The data in Figure 17 show that MTC has a complex effect on the assembly of synuclein; at low concentrations (molar ratio synuclein:MTC 2000:1) MTC apparently stimulates assembly to some degree, whereas at higher concentrations (maximum molar ratio of synuclein:MTC 2:1), it completely inhibits assembly. Without wishing to be bound by theory, it is surmised that MTC provides a ligand cross-linking effect at a low molar ratio that is insufficient to allow inhibition of aggregation but allows binding of the compound to more than one synuclein molecule to promote aggregation.

Example 9. Optimisation of the α -synuclein solid phase binding assay

Figure 18 shows the binding curves for three different synuclein preparations, as described in Example 1. The best binding was shown using syn-10, which was the most pure preparation, and the worst binding using syn-14, which was the least pure. This suggests that the second purification step removes a contaminant that inhibits binding, and that the protein should be purified with a second CM-Sepharose step, even though this reduces the yield.

Syn-10 gave the best binding of the three preparations tested, but the maximum extent of binding is still quite low. Two different solid phase preparations of tsyn were tested and showed no difference, and the concentration of tsyn used was shown to be optimal (data not shown). Different buffer conditions for the aqueous phase step were therefore tested.

Figure 19A shows that the buffers, HEPES and MES, completely abolished binding of syn. Figure 19B shows that Tris buffer at pH 7.0 allowed better binding than the pH 7.5 used

- 66 -

previously. Tris at pH 8.0 gave worse binding, so a lower pH appeared to be better for syn-
binding. Since Tris cannot be used to buffer at lower pHs, phosphate buffer was also tried,
to allow testing at pH 6.5. Phosphate at pH 6.5 gave better binding than phosphate at pH
7.0. Notwithstanding the variation in binding between the experiments, these results show
5 that the chemical nature of the buffer, as well as the pH, influences binding of syn, and that
the best buffer was Tris, pH 7.0.

The data in Figure 18 show that the best purification method for syn for solid phase binding
assays is DEAE- followed by CM- chromatography. Three more preparations of syn,
10 purified in this way, were also tested in Tris (pH 7.0) buffer (Figure 20). The results
demonstrate that there is some variability in the binding characteristics between
preparations. In a further method, the protein was dialysed against a high pH buffer (syn-
20). The binding of this protein is compared with syn-19 in Figure 21. The fall off in binding
after 2 μ M is due to a high non-specific binding, the value of which is subtracted from the
15 binding. Syn-20 showed significantly better binding than syn-19, validating the change of
dialysis buffer.

In order to try and reduce variability, the use of phosphate buffer at lower pH values was
also investigated. Figure 22A shows that fsyn-20 assayed in phosphate buffer at pH 6.0
20 gives better binding than Tris at pH 7.0 or phosphate at pH 5.5. Importantly, there is no fall
off of the curve at higher fsyn concentration due to increased background. A second fsyn
preparation (fsyn-22) also dialysed against high pH CAPS was tested in phosphate buffer
at pH 6.0 and gave similar binding to fsyn-20 (Fig. 22B). Fsyn prepared with a final high
pH dialysis and assayed in phosphate buffer at pH 6.0 was used for solid phase inhibition
25 assays (Example 5).

References

- Galvin, J. E., Uryu, K., Lee, V. M., and Trojanowski, J. Q. (1999) Axon pathology in
30 Parkinson's disease and Lewy body dementia hippocampus contains α -, β -, and γ -
synuclein, Proc. Natl. Acad. Sci. U.S.A. 96, 13450-13455.
- Uversky, V. N., Li, J., Souillac, P. O., Millett, I. S., Doniach, S., Jakes, R., Goedert, M., and
Fink, A. L. (2002) Biophysical properties of the synucleins and their propensities to

- 67 -

fibrillate: Inhibition of α -synuclein assembly by β - and γ -synucleins, J. Biol. Chem. 277, 11970-11978.





Park, J. Y., and Lansbury, P. T., Jr. (2003) β -Synuclein inhibits formation of α -synuclein protofibrils: A possible therapeutic strategy against Parkinson's disease, 5 Biochemistry 42, 3696-3700.

Hashimoto, M., Rockenstein, E., Mante, M., Mallory, M., and Masliah, E. (2001) β -Synuclein inhibits α -synuclein aggregation: A possible role as an anti-parkinsonian factor, Neuron 32, 213-223.

- 68 -

Claims

1 Use of a diaminophenothiazine compound in the manufacture of a medicament to inhibit or reverse the aggregation of synuclein, wherein said compound is selected from
 5 compounds of the following formulae:



10 wherein each one of R¹, R², R⁴, R⁶, R⁸, and R⁹ is independently selected from:

- H;
- F; -Cl; -Br; -I;
- OH; -OR;
- 15 -SH; -SR;
- NO₂;
- C(=O)R;
- C(=O)OH; -C(=O)OR;

- 69 -

-C(=O)NH₂; -C(=O)NHR; -C(=O)NR₂; -C(=O)NR^{N1}R^{N2};
 -NH₂; -NHR; -NR₂; -NR^{N1}R^{N2};
 -NHC(=O)H; -NRC(=O)H; -NHC(=O)R; -NRC(=O)R;
 -R;

5 wherein each R is independently selected from:

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;
 unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;
 unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;
 unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;
 10 unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;
 unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

wherein, in each group -NR^{N1}R^{N2}, independently, R^{N1} and R^{N2} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms;

15 and wherein, in each group -NR^{3NA}R^{3NB}, if present:

each one of R^{3NA} and R^{3NB} is independently selected from:

-H;
 unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;
 unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;
 20 unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;
 unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;
 unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;
 unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

or: R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a
 25 ring having from 3 to 7 ring atoms;

and wherein, in each group =NR^{3NC}, if present, R^{3NC} is independently selected from:

-H;
 unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;
 30 unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;
 unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;
 unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;
 unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;

- 70 -

unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

and wherein, in each group -NR^{7NA}R^{7NA}, if present:

each one of R^{7NA} and R^{7NB} is independently selected from:

5 -H;

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;

unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;

unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;

unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;

10 unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;

unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

or: R^{7NA} and R^{7NB} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms;

15 and wherein, in each group =NR^{7NC}, if present, R^{7NC} is independently selected from:

-H;

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;

unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;

unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;

20 unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;

unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;

unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

and wherein R^{N10}, if present, is independently selected from:

25 -H;

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;

unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;

unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;

unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;

30 unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;

unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

and wherein X⁻, if present, is one or more anionic counter ions to achieve electrical

- 71 -

neutrality;

and pharmaceutically acceptable salts, mixed salts, hydrates, and solvates thereof.

5 2. The use according to claim 1, wherein said compound is selected from compounds of formula (2), and pharmaceutically acceptable salts, mixed salts, hydrates, and solvates thereof.

10 3. The use according to claim 1, wherein said compound is selected from compounds of formula (3), and pharmaceutically acceptable salts, mixed salts, hydrates, and solvates thereof.

15 4. The use according to claim 1, wherein said compound is selected from compounds of formula (4), and pharmaceutically acceptable salts, mixed salts, hydrates, and solvates thereof.

5. The use according to any one of claims 1 to 4, wherein each one of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is independently selected from:

20 -H;
-F; -Cl; -Br; -I;
-OH; -OR;
-G(=O)OH; -C(=O)OR;
-R.

25 6. The use according to any one of claims 1 to 4, wherein each one of R^1 , R^2 , R^4 , R^6 , R^8 , and R^9 is independently selected from:

-H;
-R.

30 7. The use according to any one of claims 1 to 6, wherein each R is independently selected from:

unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;
unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;

- 72 -

unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl.

8. The use according to any one of claims 1 to 6, wherein each R is independently selected from:

5 unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl.

9. The use according to any one of claims 1 to 8, wherein substituents on R, if present, are independently selected from:

10 -F; -Cl; -Br; -I;
-OH; -OR';
-SH; -SR';
-NO₂;
-C(=O)R';
-C(=O)OH; -C(=O)OR';
15 -C(=O)NH₂; -C(=O)NHR'; -C(=O)NR'₂; -C(=O)NR'^{N1}R'^{N2};
-NH₂; -NHR'; -NR'₂; -NR'^{N1}R'^{N2};
-NHC(=O)H; -N'RC(=O)H; -NHC(=O)R'; -N'RC(=O)R';
-R';

wherein each R' is independently selected from:

20 unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;
unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;
unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;
unsubstituted C₆₋₁₀carboaryl; substituted C₆₋₁₀carboaryl;
unsubstituted C₅₋₁₀heteroaryl; substituted C₅₋₁₀heteroaryl;
25 unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl; substituted C₆₋₁₀carboaryl-C₁₋₄alkyl;

wherein, in each group -NR'^{N1}R'^{N2}, independently, R'^{N1} and R'^{N2} taken together with the nitrogen atom to which they are attached form a ring having from 3 to 7 ring atoms.

10. The use according to any one of claims 1 to 8, wherein substituents on R, if present, are independently selected from:

30 -F; -Cl; -Br; -I;
-OH; -OR;
-C(=O)OH; -C(=O)OR';

- 73 -

-R'.

11. The use according to claim 9 or 10, wherein each R' is independently selected from:

- 5 unsubstituted aliphatic C₁₋₆alkyl;
 unsubstituted aliphatic C₂₋₆alkenyl;
 unsubstituted C₃₋₆cycloalkyl;
 unsubstituted C₆₋₁₀carboaryl;
 unsubstituted C₅₋₁₀heteroaryl;
 10 unsubstituted C₆₋₁₀carboaryl-C₁₋₄alkyl.

12. The use according to any one of claims 1 to 6, wherein each R is independently selected from: -Me, -Et, -nPr, and -iPr.

15 13. The use according to any one of claims 1 to 4, wherein each one of R¹, R², R⁴, R⁶, R⁸, and R⁹ is independently selected from: -H, -Me, -Et, -nPr, and -iPr.

14. The use according to any one of claims 1 to 4, wherein each one of R¹, R², R⁴, R⁶, R⁸, and R⁹ is independently selected from: -H, -Me, and -Et.

20

15. The use according to any one of claims 1 to 4, wherein wherein each one of R¹, R², R⁴, R⁶, R⁸, and R⁹ is independently selected from: -H and -Me.

16. The use according to any one of claims 1 to 4, wherein wherein each of R¹, R², R⁴, R⁶, R⁸, and R⁹ is -H.

25

17. The use according to any one of claims 1 to 16, wherein, in each group -NR^{3NA}R^{3NB}, if present:

-H;

30

unsubstituted aliphatic C₁₋₆alkyl; substituted aliphatic C₁₋₆alkyl;
 unsubstituted aliphatic C₂₋₆alkenyl; substituted aliphatic C₂₋₆alkenyl;
 unsubstituted C₃₋₆cycloalkyl; substituted C₃₋₆cycloalkyl;

or R^{3NA} and R^{3NB} taken together with the nitrogen atom to which they are attached form a

- 74 -

ring having from 3 to 7 ring atoms.

18. The use according to any one of claims 1 to 16, wherein, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently selected from: -H, -Me, -Et, -nPr,
5 and -iPr.

19. The use according to any one of claims 1 to 16, wherein, in each group $-NR^{3NA}R^{3NB}$, if present, each one of R^{3NA} and R^{3NB} is independently selected from: -H and -Me.

10 20. The use according to any one of claims 1 to 16, wherein, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from:

-H;

unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;

unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;

15 unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl.

21. The use according to any one of claims 1 to 16, wherein, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from: -H, -Me, -Et, -nPr, and -iPr.

20 22. The use according to any one of claims 1 to 16, wherein, in each group $=NR^{3NC}$, if present, R^{3NC} is independently selected from: -H and -Me.

23. The use according to any one of claims 1 to 22, wherein, in each group $-NR^{7NA}R^{7NB}$, if present:

25 -H;

unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;

unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;

unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl;

or R^{7NA} and R^{7NB} taken together with the nitrogen atom to which they are attached form a
30 ring having from 3 to 7 ring atoms.

24. The use according to any one of claims 1 to 22, wherein, in each group $-NR^{7NA}R^{7NB}$, if present, each one of R^{7NA} and R^{7NB} is independently selected from: -H, -Me, -Et, -nPr,

- 75 -

and -iPr.

25. The use according to any one of claims 1 to 22, wherein, in each group $-NR^{7NA}R^{7NB}$, if present, each one of R^{7NA} and R^{7NB} is independently selected from: -H and -Me.

5

26. The use according to any one of claims 1 to 22, wherein, in each group $=NR^{7NC}$, if present, R^{7NC} is independently selected from:

-H;

unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;

10 unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;

unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl.

27. The use according to any one of claims 1 to 22, wherein, in each group $=NR^{7NC}$, if present, R^{7NC} is independently selected from: -H, -Me, -Et, -nPr, and -iPr.

15

28. The use according to any one of claims 1 to 22, wherein, in each group $=NR^{7NC}$, if present, R^{7NC} is independently selected from: -H and -Me.

29. The use according to any one of claims 1 to 28, wherein R^{N10} , if present, is independently selected from:

20

-H;

unsubstituted aliphatic C_{1-6} alkyl; substituted aliphatic C_{1-6} alkyl;

unsubstituted aliphatic C_{2-6} alkenyl; substituted aliphatic C_{2-6} alkenyl;

unsubstituted C_{3-6} cycloalkyl; substituted C_{3-6} cycloalkyl.

25

30. The use according to any one of claims 1 to 28, wherein R^{N10} , if present, is independently selected from: -H and unsubstituted aliphatic C_{1-6} alkyl.

31. The use according to any one of claims 1 to 28, wherein R^{N10} , if present, is independently selected from: -H, -Me, and -Et.





30

32. The use according to any one of claims 1 to 28, wherein R^{N10} , if present, is independently -H.






33. The use according to any one of claims 1 to 32, wherein X^- , if present, is one or more anionic counter ions to achieve electrical neutrality, optionally selected from Cl^- , Br^- , or I^- .

5






34. The use according to claim 1, wherein the compound is selected from the following compounds, and pharmaceutically acceptable salts, mixed salts, hydrates, and solvates thereof:

A	MTC	 <p>methyl-thioninium chloride</p>
B	ETC	 <p>ethyl-thioninium chloride</p>
C	DMMTC	 <p>1,9-dimethyl-methyl-thioninium chloride</p>
D	DEMTTC	 <p>1,9-diethyl-methyl-thioninium chloride</p>

- 77 -

E	DMETC	 <p>1,9-dimethyl-ethyl-thioninium chloride</p>
F	DEETC	 <p>1,9-diethyl-ethyl-thioninium chloride</p>
G	MTZ	 <p>methyl-thioninium chloride zinc chloride mixed salt</p>
H	ETZ	 <p>ethyl-thioninium chloride zinc chloride mixed salt</p>
I	MTI	 <p>methyl-thioninium iodide</p>

- 78 -

J	MTI.HI	 I^- HI <p>methyl-thioninium iodide hydrogen iodide mixed salt</p>
K	ETI	 I^- <p>ethyl-thioninium iodide</p>
L	ETI.HI	 I^- HI <p>ethyl-thioninium iodide hydrogen iodide mixed salt</p>
M	MTN	 NO_3^- <p>methyl-thioninium nitrate</p>
N	ETN	 NO_3^- <p>ethyl-thioninium nitrate</p>

- 79 -

O		<div data-bbox="635 365 1214 600" data-label="Chemical-Block"> </div> <p data-bbox="798 633 1038 667">Tolonium Chloride</p>
P		<div data-bbox="635 701 1214 936" data-label="Chemical-Block"> </div> <p data-bbox="861 969 973 1003">Azure A</p>
Q		<div data-bbox="635 1037 1214 1272" data-label="Chemical-Block"> </div> <p data-bbox="858 1305 976 1339">Azure B</p>

35 Use of a diaminophenothiazine compound as described in any of claims 1 to 34 for the inhibition or reversal of synuclein aggregation.

5 36 The use according to any one of claims 1 to 35 wherein the aggregation is associated with a disease state manifested as neurodegeneration and/or clinical dementia.

37 Use of a diaminophenothiazine compound as described in any of claims 1 to 34 in the manufacture of a medicament for the treatment or prophylaxis of a neurodegenerative
10 disease state and/or clinical dementia associated with synuclein aggregation in a subject.

38 A method of treatment or prophylaxis of a neurodegenerative disease state and/or clinical dementia associated with synuclein aggregation, which method comprises

- 80 -

administering to a subject a prophylactically or therapeutically effective amount of a diaminophenothiazine compound as described in any of claims 1 to 34, or a therapeutic composition comprising the same, so as to inhibit the aggregation of the synuclein.

5 39 A method of regulating or inhibiting the aggregation of synuclein in the brain of a mammal, which aggregation is associated with a disease state manifested as neurodegeneration and/or clinical dementia, the treatment comprising the step of administering to said mammal in need of said treatment, a prophylactically or therapeutically effective amount of a diaminophenothiazine compound as described in any
10 of claims 1 to 34.

40 A drug product for the treatment of a disease state associated with synuclein aggregation in a mammal suffering therefrom, comprising a container labeled or accompanied by a label indicating that the drug product is for the treatment of said disease,
15 the container containing one or more dosage units each comprising at least one pharmaceutically acceptable excipient and, as an active ingredient, an isolated pure diaminophenothiazine compound as described in any of claims 1 to 34.

41 The use, method, or product according to any one of claims 37 to 40 wherein said
20 treatment comprises administering said diaminophenothiazine compound in combination with a compound that modulates dopamine levels in the mammal to be treated.

42 The use, method, or product according to any one of claims 37 to 41 wherein in
25 said treatment the diaminophenothiazine compound is administered orally.

43 The use, method, or product according to any one of claims 37 to 42 wherein more than or equal to 400, 300, 200, or 100 mg daily total dose is administered.

44 The use, method, or product according to claim 43 wherein dosage units of about
30 10, 20, 30, 40, 50, 60, 60, 80, 90, 100, 110, 120, or 130 mg are administered three times a day.

45 The use, method, or product according to claim 43 wherein dosage units of about

- 81 -

10, 20, 30, 40, 50, 60, 60, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or 200 mg or administered twice a day

46 The use, method, or product of any one of claims 1 to 45 wherein less than 50% of
5 the total amount of diaminophenothiazine compound in the medicament is a compound of formula (1), or is a pharmaceutically acceptable salts, mixed salts, hydrates, and solvates thereof.

47 Use of a diaminophenothiazine compound capable of labelling aggregated
10 synuclein, in a method of manufacture of a diagnostic or prognostic reagent for use in the diagnosis or prognosis of a synucleinopathy disease state,

wherein said diaminophenothiazine compound is a compound as described in any one of claims 1 to 34 and incorporates, is conjugated to, is chelated with, or is otherwise associated with one or more isotopes, radioisotopes, positron-emitting atoms, magnetic
15 resonance labels, dyes, fluorescent markers, or antigenic groups.





48 The use according to claim 47 wherein at least one ring carbon atoms of the diaminophenothiazine compound is a positron-emitting carbon atom and/or at least one of the carbon atoms of at least one of the substituents R^1 , R^2 , R^4 , R^6 , R^8 , R^9 , R^{3NA} , R^{3NB} , R^{3NC} ,
20 R^{7NA} , R^{7NB} , R^{7NC} , and R^{N10} is a positron-emitting carbon atom.

49. The use according to claim 47 or claim 48, wherein at least one ring carbon atoms of the diaminophenothiazine compound is ^{11}C and/or at least one of the carbon atoms of at least one of the substituents R^1 , R^2 , R^4 , R^6 , R^8 , R^9 , R^{3NA} , R^{3NB} , R^{3NC} , R^{7NA} , R^{7NB} , R^{7NC} , and
25 R^{N10} is ^{11}C .

50. The use according to any one of claims 47 to 49, wherein at least one of the carbon atoms of at least one of the substituents R^{3NA} , R^{3NB} , R^{3NC} , R^{7NA} , R^{7NB} , and R^{7NC} is ^{11}C .

- 82 -

51. The use according to claim 48, selected from the following compounds, and pharmaceutically acceptable salts, mixed salts, hydrates, and solvates thereof:



52 A method of labelling aggregated synuclein comprising the steps of: contacting the aggregated synuclein with a diaminophenothiazine compound as described in any one of claims 47 to 51.

53 A method of detecting aggregated synuclein comprising the steps of: contacting the aggregated synuclein with a diaminophenothiazine compound as described in any one of claims 47 to 51 and detecting the presence and/or amount of said compound bound to aggregated synuclein.

- 83 -

54 A method of diagnosis or prognosis of a synucleinopathy in a subject believed to suffer from the disease, comprising the steps of:

(i) introducing into the subject a diaminophenothiazine compound as described in any one

5 of claims 47 to 51,

(ii) determining the presence and/or amount of said compound bound to synuclein or aggregated synuclein in the brain of the subject,

(iii) correlating the result of the determination made in (ii) with the disease state of the subject.

10

55 The use, method, or product according to any one of claims 37 to 45, or 54, wherein the subject or mammal is a human.

56 The use, method, or product according to any one of claims 36 to 45, or 47 to 51, or
15 54, wherein the disease is selected from: Parkinson's disease (PD), dementia with Lewy bodies (DLB), multiple system atrophy (MSA), drug-induced parkinsonism; pure autonomic failure (PAF).

57 The use, method, or product according to claim 56 wherein the disease is selected
20 from PD, PAF, MSA and HSD.

58 The use, method, or product according to any one of the preceding claims wherein the synuclein is α -synuclein.

25 59 The use, method, or product according to any one of the preceding claims wherein the synuclein is β -synuclein.




Figure 1




Figure 2




Figure 3

2/11




Figure 4




Figure 5

3/11




Figure 6




Figure 7




Figure 8




Figure 9

5/11




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17

9/11




Figure 18




Figure 19

10/11




Figure 20




Figure 21

11/11




Figure 22

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2007/001105

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/00 A61K31/5415 A61P25/28 A61P25/16 A61K49/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, EMBASE, BIOSIS, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ZHANG XIAN ET AL: "Methylene blue prevents neurodegeneration caused by rotenone in the retina." NEUROTOXICITY RESEARCH JAN 2006, vol. 9, no. 1, January 2006 (2006-01), pages 47-57, XP008079753 ISSN: 1029-8428 page 47, column 2, paragraph 2 - page 48, column 1, paragraph 2 page 48, column 1, last paragraph - column 2, paragraph 1 page 52, column 2, paragraph 2 - page 54, column 1, paragraph 1 page 54, column 2, last paragraph - page 55, column 1, paragraph 2 page 56, paragraph 2 - paragraph 3 ----- -/--</p>	<p>1,4-19, 23-25, 33-39,56</p>



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

3 July 2007

Date of mailing of the international search report

13/07/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kerkmann, Miren

INTERNATIONAL SEARCH REPORT

International application No

PCT/GB2007/001105

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/03972 A (HUNTER FLEMING LTD [GB]; WUELFERT ERNEST [BE]; AKTINSON ANTHONY [GB];) 17 January 2002 (2002-01-17) claims 1,7-9 page 7, lines 2-4 page 10, lines 1-7,16-27 -----	1,5-19, 23-25, 29-32, 35-39, 42,56,57
X	WO 02/04025 A (HUNTER FLEMING LTD [GB]; WUELFERT ERNEST [BE]; ATKINSON ANTHONY [GB];) 17 January 2002 (2002-01-17) page 10, lines 1-5 -----	1,5-19, 23-25, 29-32, 35-39, 42-57
X	WO 02/055720 A2 (UNIV ABERDEEN [GB]; WISCHIK CLAUDE MICHEL [GB]; HORSLEY DAVID [GB]; RI) 18 July 2002 (2002-07-18) claims 2,39,46-49,51,52 page 9, line 26 - page 10, line 3 page 11, line 6 - line 20 page 38, lines 7-27 -----	1,4-19, 23-25, 33-39, 42,56,57
X	WO 2005/095958 A (ANGESMG INC [JP]; SATO NAOYUKI [JP]; OKOUCHI MASAYASU [JP]; TANIYAMA Y) 13 October 2005 (2005-10-13) abstract -----	1,4-19, 23-25, 33-39, 42,56,57
P,X	EP 1 739 425 A (ANGES MG INC [JP]) 3 January 2007 (2007-01-03) claim 17 paragraphs [0046], [0063], [0068] -----	1,4-19, 23-25, 33-39, 42,56,57
P,X	MASUDA MASAMI ET AL: "Small molecule inhibitors of alpha-synuclein filament assembly." BIOCHEMISTRY 16 MAY 2006, vol. 45, no. 19, 16 May 2006 (2006-05-16), pages 6085-6094, XP002436919 ISSN: 0006-2960 page 6085, column 1 - column 2, paragraph 1 page 6086, column 1, paragraph 4 page 6087; table 1 -----	1,4-19, 23-25, 33-39, 56-58

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2007/001105

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.1

Although claims 38, 39, 41-46, 55-59 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition. Although claims 52-59 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/GB2007/001105

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0203972	A	17-01-2002	AU 6931401 A	21-01-2002
			AU 7077801 A	21-01-2002
			EP 1301181 A2	16-04-2003
			EP 1299125 A1	09-04-2003
			WO 0204025 A1	17-01-2002
			JP 2004502728 T	29-01-2004
			JP 2004502743 T	29-01-2004
			US 2004033936 A1	19-02-2004
			US 2003181389 A1	25-09-2003
WO 0204025	A	17-01-2002	AU 6931401 A	21-01-2002
			AU 7077801 A	21-01-2002
			EP 1301181 A2	16-04-2003
			EP 1299125 A1	09-04-2003
			WO 0203972 A2	17-01-2002
			JP 2004502728 T	29-01-2004
			JP 2004502743 T	29-01-2004
			US 2004033936 A1	19-02-2004
			US 2003181389 A1	25-09-2003
WO 02055720	A2	18-07-2002	CA 2434067 A1	18-07-2002
			CN 1496404 A	12-05-2004
			EP 1352077 A2	15-10-2003
			JP 2004524831 T	19-08-2004
			US 2004110250 A1	10-06-2004
WO 2005095958	A	13-10-2005	EP 1739425 A1	03-01-2007
EP 1739425	A	03-01-2007	WO 2005095958 A1	13-10-2005