



(51) International Patent Classification:

A61K 31/167 (2006.01) A61K 31/337 (2006.01)
A61K 31/18 (2006.01)

(21) International Application Number:

PCT/US2022/028429

(22) International Filing Date:

10 May 2022 (10.05.2022)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/201,749 11 May 2021 (11.05.2021) US

(71) Applicant: **ABBVIE INC.** [US/US]; 1 North Waukegan Road, North Chicago, IL 60064 (US).

(72) Inventors: **HAYSLIP, John**; 490 Lanterback Island Drive, Satellite Beach, FL 32937 (US). **KYE, Steve, H.**; 538 Council Circle, Vernon Hills, IL 60061 (US). **AINSWORTH, William, B.**; 1667 Church Street, Evanston, IL 60201 (US).

(74) Agent: **RICH, Roger, C.** et al.; Jones Day, 250 Vesey Street, New York, NY 10281-1047 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH,

KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: VENETOCLAX DOSING REGIMENS FOR USE IN TREATING MYELODYSPLASTIC SYNDROMES IN COMBINATION WITH AZACITIDINE




FIG. 1

(57) Abstract: The invention described herein relates to therapeutic dosing regimens comprising administering venetoclax in combination with azacitidine for treating myelodysplastic syndromes (MDS).

WO 2022/240786 A1

**VENETOCLAX DOSING REGIMENS FOR USE IN TREATING MYELODYSPLASTIC
SYNDROMES IN COMBINATION WITH AZACITIDINE**

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 63/201,749, filed May 11, 2021, the disclosure of which is incorporated by reference herein in its entirety.

FIELD OF THE INVENTION

[0002] This invention relates to methods for treating myelodysplastic syndromes (MDS) in a human subject comprising administering to the subject venetoclax in combination with azacitidine.

BACKGROUND OF THE INVENTION

[0003] Myelodysplastic Syndromes (MDS) represent a heterogeneous group of clonal hematopoietic stem cell disorders with significant morbidity and high mortality. These syndromes are characterized by ineffective hematopoiesis that manifest clinically as cytopenias and a variable rate of transformation to acute myeloid leukemia (secondary or sAML). Although about one-third of all MDS patients later develop AML, MDS are not considered to be an early form of AML. The primary reason for death in MDS patients is not because of AML transformation, but due to consequences of bone marrow failure, and in particular neutropenia leading to infections, including septic shock, or thrombocytopenia leading to bleeding.

[0004] Approximately half (45%) of MDS patients present with higher-risk MDS risk (International Prognostic Scoring System (IPSS) overall score > 1.5) and have a median survival less than one year with best supportive care. The only curative treatment for higher-risk MDS is an allogeneic stem cell or bone marrow transplantation. However, not all patients are eligible for this intensive treatment approach. If bone marrow transplantation is not possible, patients are typically treated with hypomethylating agents such as azacitidine. Currently, azacitidine is the only drug shown to prolong survival in treatment-naïve higher-risk MDS, however, overall outcomes need to be improved.

[0005] Venetoclax is an oral small molecule inhibitor of B-cell lymphoma 2 (BCL-2) that rapidly induces multiple hallmarks of apoptotic cell death. Venetoclax is being investigated in

clinical oncology studies as a monotherapy and in combination with a variety of compounds for the treatment of a number of hematologic malignancies, including chronic lymphocytic leukemia (CLL) and acute myeloid leukemia (AML). However, the dosing regimen used in the first clinical trial of venetoclax in MDS produced deleterious side effects in certain patients. For example, two subjects developed fatal sepsis in the setting of severe neutropenia. Thus, there exists in the art a need for dosing regimen for MDS patients experiencing certain side effects.

BRIEF SUMMARY OF THE INVENTION

[0006] The present disclosure relates to methods for treating myelodysplastic syndromes in a human subject, and in some aspects, more specifically treatment-naïve higher-risk myelodysplastic syndromes.

[0007] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle. In some aspects, the daily dose of azacitidine is reduced according to the bone marrow cellularity, the absolute neutrophil count nadir decrease, the white blood cell count nadir decrease, and the platelet count nadir count decrease.

[0008] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 50\%$ but no less than 33% during a next 28 day dosing cycle. The human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$. The human subject has no improvement in cell line differentiation and the human subject has at least one condition selected

from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction. In some aspects, the daily dose of azacitidine is reduced according to the bone marrow cellularity.

[0009] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$, and a platelet count nadir of $\leq 50 \times 10^9/L$; the daily dose of azacitidine is reduced from 75 mg/m² to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle.

[0010] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject, comprising administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50% during a next 28 day dosing cycle. The human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$. The human subject also has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.0 \times 10^9/L$, and a platelet count nadir of $\leq 50 \times 10^9/L$.

[0011] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to ≤ 50 mg/m² but no less than 36 mg/m² during a next 28 day dosing cycle. The human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and a previous response of complete remission, partial remission, or marrow complete remission at the beginning of a prior dosing cycle. The human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 0.500 \times 10^9/L$, a platelet count nadir of $\leq 50 \times 10^9/L$; wherein the baseline platelet was $>$

$100 \times 10^9/L$; and a platelet count nadir of $< 50 \%$; wherein the baseline platelet was $\leq 100 \times 10^9/L$.

BRIEF DESCRIPTION OF THE FIGURES

[0012] FIG. 1 is a plot of the mean value of the absolute neutrophil count versus study day cycle. The number of observations is shown in Table 11.

[0013] FIG. 2 is a plot of the mean value of the platelet count versus study day cycle. The number of observations is shown in Table 11.

[0014] FIGs. 3A-3F are bar graphs of hematologic toxicity showing the number of patients with worsening common terminology criteria grade over baseline per cycle. FIG. 3A is anemia. FIG. 3B is febrile neutropenia. FIG. 3C is leukopenia. FIG. 3D is neutropenia. FIG. 3E is thrombocytopenia. FIG. 3F is infections.

[0015] FIGs. 4A-4C are bar graphs of gastrointestinal toxicity showing the number of patients with worsening common terminology criteria grade over baseline per cycle. FIG. 4A is diarrhea. FIG. 4B is vomiting. FIG. 4C is nausea.

DETAILED DESCRIPTION OF THE INVENTION

[0016] This present disclosure relates to methods for treating treatment-naïve higher-risk myelodysplastic syndromes (MDS) in a human subject comprising administering to the subject venetoclax in combination with azacitidine.

[0017] Although venetoclax has been administered to patients with AML who had a prior history of MDS (sAML), herein is the first disclosure evaluating venetoclax in combination with azacitidine in subjects with MDS, more specifically, in those subjects with treatment-naïve higher-risk MDS, in which the dosing regimen is modified to account for subjects who have developed hematological toxicities after beginning treatment. Apart from the toxicity-specific dosing modifications disclosed herein, other significant differences between dosing venetoclax in combination with azacitidine for MDS versus AML include reducing the duration of venetoclax dosing from 28 to 14 days in the 28 day cycle for MDS, as well the lack of any dosing ramp up for subjects with MDS.

[0018] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily

dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle.

[0019] "Venetoclax" is 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-*N*-({3nitro-4-[(tetrahydro-2*H*-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1*H*-pyrrolo[2,3-*b*]pyridin5-yloxy)benzamide. Venetoclax is a selective Bcl-2 inhibitor approved for adult patients with CLL and adult patients with newly diagnosed AML who are 75 years or older, or who are ineligible for intensive induction chemotherapy.

[0020] Azacitidine is 4-amino-1-β-D-ribofuranosyl-s-triazin-2(1*H*)-one. Azacitidine is supplied in a sterile form for reconstitution as a suspension for subcutaneous injection or reconstitution as a solution with further dilution for intravenous infusion.

[0021] The term "AE" as used herein refers to adverse event.

[0022] The term "AML" as used herein refers to acute myeloid leukemia.

[0023] The term "ANC" as used herein refers to absolute neutrophil count.

[0024] The term "CLL" as used herein refers to chronic lymphocytic leukemia.

[0025] The term "CML" as used herein refers to chronic myeloid leukemia.

[0026] The term "CMML" as used herein refers to chronic myelomonocytic leukemia.

[0027] The term "CR" as used herein refers to complete remission.

[0028] The term "CTC" as used herein refers to common terminology criteria.

[0029] The term "ECOG" as used herein refers to Eastern Cooperative Oncology Group.

[0030] The term "G-CSF" as used herein refers to granulocyte colony-stimulating factor.

[0031] The term "HRQoL" as used herein refers to health-related quality of life.

[0032] The term "HMAs" as used herein refers to hypomethylating agents.

[0033] The term "HR-MDS" as used herein refers to higher risk myelodysplastic syndromes.

[0034] The term "IPSS" as used herein refers to International Prognostic Scoring System.

[0035] The term "IPSS-R" as used herein refers to Revised International Prognostic Scoring System.

[0036] The term "JMML" as used herein refers to juvenile myelomonocytic leukemia.

[0037] The term "mCR" as used herein refers to marrow complete remission.

[0038] The term "MDS" as used herein refers to myelodysplastic syndromes.

[0039] The term "MPN" as used herein refers to myeloproliferative neoplasm.

[0040] The term "OS" as used herein refers to overall survival.

[0041] The term "PR" as used herein refers to partial remission.

[0042] The term "RAEB" as used herein refers to refractory anemia with excess blasts.

[0043] The term "sAML" as used herein refers to secondary acute myeloid leukemia.

[0044] The term "SE1" as used herein refers to Safety Expansion Cohort 1.

[0045] The term "SE2" as used herein refers to Safety Expansion Cohort 2.

[0046] The term "TEAE" as used herein refers to treatment-emergent adverse events.

[0047] The term "tMDS" as used herein refers to treatment-related or therapy-related myelodysplastic syndromes.

[0048] No improvement in cell line differentiation refers to the lack of clear improvement in cell differentiation at the time of the next cycle. For example, the percentage of mature granulocytes is lower and the absolute neutrophil count is lower than at onset of the dosing cycle.

[0049] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle.

[0050] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-

risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 75% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 30-60%; and wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 75\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 75\%$ from the baseline white blood cell count; and a platelet count nadir decrease of $> 75\%$ from the baseline platelet count. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0051] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 75% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 30-60%; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 75\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 75\%$

from the baseline white blood cell count; and a platelet count nadir decrease of $> 75\%$ from the baseline platelet count; and wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day.

[0052] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 75% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 30-60%; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 75\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 75\%$ from the baseline white blood cell count; and a platelet count nadir decrease of $> 75\%$ from the baseline platelet count; wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood

cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir; wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; and wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a subsequent absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0053] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 50% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 15 to $< 30\%$; and wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of \geq

50% from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0054] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 50% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 15 to $< 30\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; and wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day.

[0055] In certain embodiments, a method for treating myelodysplastic syndromes in a

human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 50% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 15 to $< 30\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir; wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; and wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a subsequent absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a subsequent white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a subsequent platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the

azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0056] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 50% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 15 to $< 30\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; and wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $> 75\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $> 75\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $> 75\%$ from the baseline platelet count. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0057] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75

mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 50% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 15 to $< 30\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; and wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir decrease of 50-75% from the baseline absolute neutrophil count, a white blood cell count nadir decrease of 50-75% from the baseline white blood cell, and a platelet count nadir decrease of 50-75% from the baseline platelet count. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0058] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ during but no less than 33% a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 33% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity $< 15\%$; and wherein the human subject has at least one of a condition

selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0059] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 33% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity $< 15\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; and wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day.

[0060] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ during but no less than 33% a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 33% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity $< 15\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir; wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; and wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a next subsequent neutrophil count of $> 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a subsequent white blood cell count of $> 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a subsequent platelet count of $> 25\%$ above the platelet count nadir and relative to a

difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0061] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 33% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity $< 15\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; and wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; and wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a subsequent neutrophil count of $> 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a subsequent white blood cell count of $> 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a subsequent platelet count of $> 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the

azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0062] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ during but no less than 33% a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 33% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity $< 15\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; and wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $> 75\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $> 75\%$ from the baseline white blood cell count, and a platelet count decrease of $> 75\%$ from the baseline platelet count. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0063] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75

mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; no improvement in cell line differentiation; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 33% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity $< 15\%$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count, a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count, and a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count; and wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir decrease of 50-75% from the baseline absolute neutrophil count, a white blood cell count nadir decrease of 50-75% from the baseline white blood cell count relative to the white blood cell count nadir, and a platelet count nadir decrease of 50-75% from the baseline platelet count. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0064] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 50\%$ during but no less than 33% a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; wherein the human subject has no improvement in cell line differentiation; and wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction.

[0065] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-

risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 50\%$ but no less than 33% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; wherein the human subject has no improvement in cell line differentiation; and wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction.

[0066] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 50\%$ but no less than 33% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; wherein the human subject has no improvement in cell line differentiation; wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 50% during the next 28 day dosing cycle; and wherein the human subject has a bone marrow cellularity of 15-50%. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0067] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 50\%$ but no less than 33% during a next 28 day dosing cycle; wherein the

human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; wherein the human subject has no improvement in cell line differentiation; wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; wherein the daily dose of azacitidine of 75 mg/m^2 is reduced to 50% but no less than 33% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 15-50%; and wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count increase of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count increase of $< 50\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count increase of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day.

[0068] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m^2 is reduced to $\leq 50\%$ but no less than 33% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; wherein the human subject has no improvement in cell line differentiation; wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; wherein the daily dose of azacitidine of 75 mg/m^2 is reduced to 50% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of 15-50%;

wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count increase of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count increase of $< 50\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count increase of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir; and wherein the daily dose of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a subsequent absolute neutrophil count of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a subsequent white blood cell count of $< 50\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a subsequent platelet count of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0069] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m^2 is reduced to $\leq 50\%$ but no less than 33% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/\text{L}$, a baseline white blood cell count of $< 3 \times 10^9/\text{L}$, or a baseline platelet count of $< 75 \times 10^9/\text{L}$; wherein the human subject has no improvement in cell line differentiation; wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$

reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; wherein the daily dose of azacitidine of 75 mg/m^2 is reduced to 33% during the next 28 day dosing cycle; and wherein the human subject has a bone marrow cellularity of $< 15\%$. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0070] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m^2 is reduced to $\leq 50\%$ but no less than 33% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/\text{L}$, a baseline white blood cell count of $< 3 \times 10^9/\text{L}$, or a baseline platelet count of $< 75 \times 10^9/\text{L}$; wherein the human subject has no improvement in cell line differentiation; wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; wherein the daily dose of azacitidine of 75 mg/m^2 is reduced to 33% during the next 28 day dosing cycle; and wherein the human subject has a bone marrow cellularity of $< 15\%$; wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count increase of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count increase of $< 50\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir; and a next platelet count increase of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day.

[0071] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a

daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 50\%$ but no less than 33% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$; wherein the human subject has no improvement in cell line differentiation; wherein the human subject has at least one condition selected from the group consisting of: an absolute neutrophil count nadir of $\geq 50\%$ reduction, a white blood cell count nadir of $\geq 50\%$ reduction, and a platelet count nadir of $\geq 50\%$ reduction; wherein the daily dose of azacitidine of 75 mg/m² is reduced to 33% during the next 28 day dosing cycle; wherein the human subject has a bone marrow cellularity of $< 15\%$; wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count increase of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count increase of $< 50\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir; and a next platelet count increase of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir; and wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a subsequent absolute neutrophil count of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a subsequent white blood cell count of $< 50\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a subsequent platelet count of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0072] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has: a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; the daily dose of azacitidine is reduced from 75 mg/m² to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle.

[0073] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has: a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; the daily dose of azacitidine is reduced from 75 mg/m² to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle.

[0074] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; ; the daily dose of azacitidine is reduced from 75 mg/m² to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 67%; and wherein the human subject has at least one condition selected from the group consisting of: the absolute neutrophil count nadir is from $0.5 \times 10^9/L$ to $1.5 \times 10^9/L$;

and the platelet count nadir is from $25 \times 10^9/L$ to $50 \times 10^9/L$. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0075] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; ; the daily dose of azacitidine is reduced from 75 mg/m² to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle;; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 67%; wherein the human subject has at least one condition selected from the group consisting of: the absolute neutrophil count nadir is from $0.5 \times 10^9/L$ to $1.5 \times 10^9/L$; and the platelet count nadir is from $25 \times 10^9/L$ to $50 \times 10^9/L$; and wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and a next platelet count of $\leq 50 \times 10^9/L$. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day.

[0076] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; ; the daily dose of azacitidine is reduced from 75 mg/m² to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle;; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 67%; wherein the human subject has at least one condition selected

from the group consisting of: the absolute neutrophil count nadir is from $0.5 \times 10^9/L$ to $1.5 \times 10^9/L$; and the platelet count nadir is from $25 \times 10^9/L$ to $50 \times 10^9/L$; wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and a next platelet count of $\leq 50 \times 10^9/L$; wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; and wherein the human subject has prior to a start of the subsequent 28 day dosing cycle at least one condition selected from the group consisting of: a subsequent absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and a subsequent platelet count of $\leq 50 \times 10^9/L$. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0077] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m^2 of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; ; the daily dose of azacitidine is reduced from 75 mg/m^2 to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle;; wherein the daily dose of azacitidine is reduced from 75 mg/m^2 to 50%; and wherein the human subject has at least one condition selected from the group consisting of: the absolute neutrophil count nadir is $< 0.5 \times 10^9/L$; and the platelet count nadir is $< 25 \times 10^9/L$. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0078] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75

mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; ; the daily dose of azacitidine is reduced from 75 mg/m² to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50%; wherein the human subject has at least one condition selected from the group consisting of: the absolute neutrophil count nadir is $< 0.5 \times 10^9/L$; and the platelet count nadir is $< 25 \times 10^9/L$; and wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and a next platelet count of $\leq 50 \times 10^9/L$. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day.

[0079] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein if the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; ; the daily dose of azacitidine is reduced from 75 mg/m² to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50%; wherein the human subject has at least one condition selected from the group consisting of: the absolute neutrophil count nadir is $< 0.5 \times 10^9/L$; and the platelet count nadir is $< 25 \times 10^9/L$; wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and a next platelet count of $\leq 50 \times 10^9/L$; wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; and wherein the human subject has prior to a start of the subsequent 28 day dosing cycle at least one condition

selected from the group consisting of: a subsequent absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and a subsequent platelet count of $\leq 50 \times 10^9/L$. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0080] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.0 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$.

[0081] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.0 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously.

[0082] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75

mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.0 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; and wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a subsequent absolute neutrophil count increase of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir; and a subsequent platelet count increase of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day.

[0083] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50% during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.0 \times 10^9/L$; and a platelet count nadir of $\leq 50 \times 10^9/L$; wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of: a subsequent absolute neutrophil count increase of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir; and a subsequent platelet count increase of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir; wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle; and wherein the human subject has prior to the start of the subsequent 28 day

dosing cycle, at least one condition selected from the group consisting of: a subsequent absolute neutrophil count increase of $< 50\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir; and a subsequent platelet count increase of $< 50\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In some aspects, the azacitidine is administered intravenously. In other aspects, the azacitidine is administered subcutaneously. In yet other aspects, the method further comprises a delay prior to the subsequent 28 day dosing cycle of > 1 day. In still other aspects, the method further comprises a delay prior to the next 28 day dosing cycle of ≥ 1 day; and further comprises a delay prior to the subsequent 28 day dosing cycle of ≥ 1 day.

[0084] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to ≤ 50 mg/m² but no less than 36 mg/m² during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; wherein the human subject has a previous response of complete remission, partial remission, or marrow complete remission at the beginning of a prior dosing cycle; and wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 0.500 \times 10^9/L$, a platelet count nadir of $\leq 50 \times 10^9/L$; wherein the baseline platelet was $> 100 \times 10^9/L$, and a platelet count nadir of $< 50 \%$; wherein the baseline platelet was $\leq 100 \times 10^9/L$.

[0085] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to ≤ 50 mg/m² but no less than 36 mg/m² during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; wherein the human subject has a previous response of complete remission, partial remission, or marrow complete remission at the beginning of a prior

dosing cycle; and wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 0.500 \times 10^9/L$, and a platelet count nadir of $\leq 50 \times 10^9/L$; wherein the baseline platelet was $> 100 \times 10^9/L$, a platelet count nadir of $< 50 \%$; wherein the baseline platelet was $\leq 100 \times 10^9/L$.

[0086] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to ≤ 50 mg/m² but no less than 36 mg/m² during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; wherein the human subject has a previous response of complete remission, partial remission, or marrow complete remission at the beginning of a prior dosing cycle; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 0.500 \times 10^9/L$, and a platelet count nadir of $\leq 50 \times 10^9/L$; wherein the baseline platelet was $> 100 \times 10^9/L$, a platelet count nadir of $< 50 \%$; wherein the baseline platelet was $\leq 100 \times 10^9/L$; and wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50 mg/m².

[0087] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to ≤ 50 mg/m² but no less than 36 mg/m² during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; wherein the human subject has a previous response of complete remission, partial remission, or marrow complete remission at the beginning of a prior dosing cycle; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 0.500 \times 10^9/L$, and a platelet count nadir of $\leq 50 \times 10^9/L$; wherein the baseline platelet was $> 100 \times 10^9/L$, a platelet count nadir of $< 50 \%$; wherein the baseline platelet was $\leq 100 \times 10^9/L$; wherein the daily dose of azacitidine is reduced

from 75 mg/m² to 50 mg/m²; and wherein the daily dose of azacitidine is reduced from 50 mg/m² to 36 mg/m² in a successive 28 day dosing cycle.

[0088] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. The method comprises administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle; wherein the daily dose of azacitidine is reduced from 75 mg/m² to ≤ 50 mg/m² but no less than 36 mg/m² during a next 28 day dosing cycle; wherein the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; wherein the human subject has a previous response of complete remission, partial remission, or marrow complete remission at the beginning of a prior dosing cycle; wherein the human subject has at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 0.500 \times 10^9/L$, and a platelet count nadir of $\leq 50 \times 10^9/L$; wherein the baseline platelet was $> 100 \times 10^9/L$, a platelet count nadir of < 50 %; wherein the baseline platelet was $\leq 100 \times 10^9/L$; wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50 mg/m²; wherein the daily dose of azacitidine is reduced from 50 mg/m² to 36 mg/m² in a successive 28 day dosing cycle; and wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle.

[0089] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises (a) obtaining a baseline absolute neutrophil count, a baseline white blood cell count, and a baseline platelet count from the human subject, (b) treating the human subject with an initial dose of venetoclax and an initial dose of azacitidine during a first treatment cycle, (c) comparing the baseline absolute neutrophil count, baseline white blood cell count, and baseline platelet count with a subsequent absolute neutrophil count, subsequent white blood cell count, and subsequent platelet count obtained after the first treatment cycle, and (d) if the subsequent absolute neutrophil count nadir is $\geq 50\%$ of the baseline neutrophil count, the subsequent white blood cell count nadir is $\geq 50\%$ of the baseline neutrophil count, or the subsequent platelet count nadir is $\geq 50\%$ of the baseline platelet count, treating the human subject with a dose of azacitidine that is $\leq 75\%$ but no less than 33% of the initial dose of azacitidine in a subsequent treatment cycle. In some aspects, the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. In other aspects, the

initial dose of venetoclax is 400 mg and the initial dose of azacitidine is 75 mg/m². In yet other aspects, the dosing cycle is 28 days. In yet other aspects, the human subject is administered a daily dose of venetoclax for 14 days in a 28 day dosing cycle and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle. In yet other aspects, the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a further subsequent 28 day dosing cycle if the human subject has prior to a start of the further subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In yet other aspects, the azacitidine is administered intravenously. In yet other aspects, the azacitidine is administered subcutaneously.

[0090] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises (a) obtaining a baseline absolute neutrophil count, a baseline white blood cell count, and a baseline platelet count from the human subject, (b) treating the human subject with an initial dose of venetoclax and an initial dose of azacitidine during a first treatment cycle, (c) comparing the baseline absolute neutrophil count, baseline white blood cell count, and baseline platelet count with a subsequent absolute neutrophil count, subsequent white blood cell count, and subsequent platelet count obtained after the first treatment cycle, and (d) if the subsequent absolute neutrophil count nadir is $\geq 50\%$ of the baseline neutrophil count, the subsequent white blood cell count nadir is $\geq 50\%$ of the baseline neutrophil count, or the subsequent platelet count nadir is $\geq 50\%$ of the baseline platelet count, treating the human subject with a dose of azacitidine that is $\leq 50\%$ but no less than 33% of the initial dose of azacitidine in a subsequent treatment cycle. In some aspects, the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. In other aspects, the initial dose of venetoclax is 400 mg and the initial dose of azacitidine is 75 mg/m². In yet other aspects, the dosing cycle is 28 days. In yet other aspects, the human subject is administered a daily dose of venetoclax for 14 days in a 28 day dosing cycle and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle. In yet other aspects, the daily dose of 400 mg

of venetoclax is reduced from 14 to 7 days in a further subsequent 28 day dosing cycle if the human subject has prior to a start of the further subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In yet other aspects, the azacitidine is administered intravenously. In yet other aspects, the azacitidine is administered subcutaneously.

[0091] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises (a) obtaining a baseline absolute neutrophil count, a baseline white blood cell count, and a baseline platelet count from the human subject, (b) treating the human subject with an initial dose of venetoclax and an initial dose of azacitidine during a first treatment cycle, (c) comparing the baseline absolute neutrophil count, baseline white blood cell count, and baseline platelet count with a subsequent absolute neutrophil count, subsequent white blood cell count, and subsequent platelet count obtained after the first treatment cycle, and (d) if the human subject has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/L$, and a platelet count nadir of $\leq 50 \times 10^9/L$; the dose of azacitidine is reduced from 75 mg/m^2 to $\leq 67\%$ but no less than 50% during a next 28 day dosing cycle. In some aspects, the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. In other aspects, the initial dose of venetoclax is 400 mg and the initial dose of azacitidine is 75 mg/m^2 . In yet other aspects, the dosing cycle is 28 days. In yet other aspects, the human subject is administered a daily dose of venetoclax for 14 days in a 28 day dosing cycle and a daily dose of 75 mg/m^2 of azacitidine for 7 days in the 28 day dosing cycle. In yet other aspects, the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a further subsequent 28 day dosing cycle if the human subject has prior to a start of the further subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to

a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir. In yet other aspects, the azacitidine is administered intravenously. In yet other aspects, the azacitidine is administered subcutaneously.

[0092] In certain embodiments, a method for treating myelodysplastic syndromes in a human subject is provided. The method comprises (a) obtaining a baseline absolute neutrophil count, a baseline white blood cell count, and a baseline platelet count from the human subject, (b) treating the human subject with an initial dose of venetoclax and an initial dose of azacitidine during a first treatment cycle, (c) comparing the baseline absolute neutrophil count, baseline white blood cell count, and baseline platelet count with a subsequent absolute neutrophil count, subsequent white blood cell count, and subsequent platelet count obtained after the first treatment cycle, and (d) has a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/L$, a baseline white blood cell count of $\geq 3 \times 10^9/L$, or a baseline platelet count of $\geq 75 \times 10^9/L$; and at least one of a condition selected from the group consisting of: an absolute neutrophil count nadir of $\leq 1.0 \times 10^9/L$, and a platelet count nadir of $\leq 50 \times 10^9/L$, the dose of azacitidine is reduced from 75 mg/m² to 50% during a next 28 day dosing cycle. In some aspects, the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes. In other aspects, the initial dose of venetoclax is 400 mg and the initial dose of azacitidine is 75 mg/m². In yet other aspects, the dosing cycle is 28 days. In yet other aspects, the human subject is administered a daily dose of venetoclax for 14 days in a 28 day dosing cycle and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle. In yet other aspects, the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a further subsequent 28 day dosing cycle if the human subject has prior to a start of the further subsequent 28 day dosing cycle, at least one condition selected from the group consisting of: a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir, a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir, and a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet

count nadir. In yet other aspects, the azacitidine is administered intravenously. In yet other aspects, the azacitidine is administered subcutaneously.

EXAMPLES

[0093] In order that the invention described herein may be more fully understood, the following examples are set forth.

[0094] A multicenter, non-randomized Phase 1b study in adults with previously untreated higher-risk MDS, defined as IPSS risk categories of Int-2 or High (IPSS overall score ≥ 1.5) was initiated. The original protocol randomized patients into 1 of 3 treatment groups: venetoclax 800 mg + azacitidine, venetoclax 400 mg + azacitidine, and azacitidine monotherapy. Under the original protocol, venetoclax was administered on Days 1 through 28 of each 28-day-cycle and dosing was initiated according to a ramp-up dosing schedule in Cycle 1. With this dosing schedule, 2 patients developed fatal sepsis in the setting of severe neutropenia, after which the study was placed on partial clinical hold and enrollment was suspended. The partial clinical hold was lifted based on a revised protocol, which ultimately resulted in a lower incidence of infections and leukopenia events.

Study Design

Subject Inclusion Criteria

[0095] Subjects 18 years or older diagnosed with treatment-naïve IPSS intermediate-2 or high-risk myelodysplastic syndromes with ECOG ≤ 2 were enrolled. For this study, inclusion criteria included the following:

1. Subject must be ≥ 18 years of age.
2. Subject must have documented diagnosis of previously untreated *de novo* MDS with:
 - a. International Prognostic Scoring System (IPSS) risk categories Intermediate-2 or High (i.e., minimum IPSS score of 1.5) OR Revised IPSS (IPSS-R) categories Intermediate, High or Very High (score of > 3); and
 - b. Presence of $< 20\%$ bone marrow blasts per bone marrow biopsy/aspirate.
3. Subject must have an Eastern Cooperative Oncology Group (ECOG) performance

score of ≤ 2 .

[0096] According to the International Prognostic Scoring System, myelodysplastic syndromes patients are grouped into two major risk groups, low risk and higher-risk. Higher-risk myelodysplastic syndromes as used herein are defined as an International Prognostic Scoring System (IPSS) risk categories Int-2 or High (i.e., minimum IPSS score of 1.5) or Revised IPSS (IPSS-R) categories Intermediate, High or Very High (overall score of > 3).

Table 1: International Prognostic Scoring System (IPSS)

Category	Score				
	0	0.5	1.0	1.5	2.0
Bone Marrow Blasts (%)	< 5	5 to 10	-	11 to 20	21 to 30
Karyotype	Normal, Y-, 5q-, 20q-	All other cytogenetic abnormalities	Abnormal chromosome 7 or 3 or more abnormalities		
Cytopenias defined as: <ul style="list-style-type: none"> • Hemoglobin < 10 g/dL • absolute neutrophil count < 1800/μL • platelet count < 100,000 μL 	No cytopenia or cytopenia of 1 cell type	Cytopenia of 2 or 3 cell types			

[0097] Subjects with higher-risk MDS are classified into the Revised International Prognostic Scoring System (IPSS-R) categories of Intermediate, High and Very High. This patient population largely corresponds to IPSS Intermediate risk-2 and High groups and World Health Organization (WHO) histologic subtypes of refractory anemia with excess blasts (RAEB)-1 and RAEB-2. The IPSS-R is now also considered a well-validated assessment tool to identify patients who are commonly considered clinically appropriate to receive active treatment. The Revised International Prognostic Scoring System (IPSS-R) is shown in Table 2 and includes a refined classification of cytogenetic abnormalities, more specific cut-offs for bone marrow blast counts and cytopenias, and is weighted for their severity. The Revised International Prognostic Scoring System risk groups for myelodysplastic syndromes are defined based an overall score. The overall score is calculated as the sum of the blast score, cytogenetics score,

hemoglobin score, platelets score, and absolute neutrophil count score.

Table 2: Revised International Prognostic Scoring System (IPSS-R) Criteria and Scoring for MDS

Parameter	Criteria	Score	Risk Groups (Overall Score ¹ and Median Survival Time)
Blasts in bone marrow (%)	≤ 2	0	Very Low: Overall Risk Score ≤ 1.5 Median survival time = 8.8 years
	> 2 to < 5	1.0	
	5 to 10	2.0	
	> 10	3.0	
Cytogenetics	Very good	0	Low: Overall Risk Score > 1.5 to 3.0 Median survival time = 5.3 years
	Good	1.0	
	Intermediate	2.0	
	Poor	3.0	
	Very Poor	4.0	
Hemoglobin (g/dL)	≥ 10	0	Intermediate: Overall Risk Score > 3.0 to 4.5 Median survival time = 3 years
	8 to < 10	1.0	
	< 8	1.5	
Platelets (10 ⁹ / L)	≥ 100	0	High: Overall Risk Score > 4.5 to 6.0 Median survival time = 1.6 years
	50 to < 100	0.5	
	< 50	1.0	
ANC (10 ⁹ / L)	≥ 0.8	0	Very High: Overall Risk Score > 6.0 Median survival time = 0.8 years
	< 8	0.5	

¹ Overall score is calculated as the blast score + cytogenetics score + hemoglobin score + platelets score + absolute neutrophil count score

[0098] ECOG performance status was assessed using the criteria in Table 3.

Table 3: ECOG performance status

Grade	Description
0	Fully active, able to carry on all pre-disease performance without restriction.
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature, e.g., light housework, office work.
2	Ambulatory and capable of all self-care but unable to carry out any work activities. Up and about more than 50% of waking hours.
3	Capable of only limited self-care, confined to bed or chair more than 50% of waking hours.

4	Completely disabled. Cannot carry on any self-care. Totally confined to bed or chair.
---	---

Key Exclusion Criteria

1. Subject has received prior therapy for MDS.
2. Subject has received prior therapy with a BH3 mimetic
3. Subject has a diagnosis other than previously untreated de novo MDS, including:
 - a. MDS with IPSS risk categories Low or Int-1 (overall IPSS score < 1.5)
 - b. Therapy-related MDS (t-MDS)
 - c. MDS evolving from a pre-existing myeloproliferative neoplasm (MPN)
 - d. MDS/MPN including chronic myelomonocytic leukemia (CMML), atypical chronic myeloid leukemia (CML), juvenile myelomonocytic leukemia (JMML) and unclassifiable MDS/MPN.
4. Subject has received strong or moderate CYP3A inducers within 7 days prior to the first dose of study drug.
5. Subject enrolled in a Dose-Escalation cohort has received strong or moderate CYP3A inhibitors within 3 days prior to the first dose of study drug with the exception of Safety Expansion cohorts.

[0099] Azacitidine 75 mg/m² (intravenous or subcutaneous daily) was administered for 7 days and venetoclax was administered at 400 mg for 14 days in each 28 day cycle. In both cohorts, dose modification during Cycle 1 was not recommended. Dose modifications in subsequent cycles were prescribed for adverse events. In Safety Expansion Cohort 1 (SE1), venetoclax was initially reduced for significant neutrophil or platelet toxicity. Dose reductions per protocol were 33% for azacitidine or 50% for venetoclax for 14 days in each cycle. In subsequent cycles, venetoclax duration could be shortened to 9 days of each cycle. In Safety Expansion Cohort 2 (SE2), dose modification guidelines recommended stepwise reductions, first in azacitidine dose (first to 50 mg/m², then 36 mg/m²) and subsequently in venetoclax duration to 7 days of each cycle (venetoclax 400 mg) as shown in Table 4. The impact of each dose modification strategy on safety and efficacy in SE1 versus SE2 was compared. Worsening of treatment-emergent adverse events grades from baseline was analyzed by cycle. Responses were evaluated using IWG 2006 criteria. Analyses included all subject who received ≥1 dose of study

drug.

Table 4: SE2 Dose Modification Guidelines

	Azacitidine	Venetoclax
Planned Dose	75 mg/m ² x 7 days	400 mg x 14 days
1 st Dose Reduction	50 mg/m ² x 7 days	400 mg x 14 days
2 nd Dose Reduction	36 mg/m ² x 7 days	400 mg x 14 days
3 rd Dose Reduction	36 mg/m ² x 7 days	400 mg x 7 days

[0100] After a previous interruption of treatment, delay starting the next cycle, onset of adverse events associated with hematological toxicity, significant reduction of neutrophils, or significant reduction of platelets, a treatment dose reduction may be indicated. Stepwise azacitidine dose modification occurred followed by adjustment of the venetoclax treatment from 14 to 7 days at the last step after all azacitidine dose modification steps had occurred. Venetoclax and azacitidine were resumed on the same day after any delays or interruptions in treatment.

[0101] Subjects who initiate a cycle of treatment with neutropenia, absolute neutrophil count $< 1.5 \times 10^9 / L$, or thrombocytopenia, platelets $< 75 \times 10^9 / L$, may be especially sensitive to cytopenias due to abnormal hematopoiesis resulting from the underlying absolute neutrophil count. Accordingly, such subjects do not typically require dose reductions in response to uncomplicated nadir cytopenias. However, subjects who initiate a cycle of treatment with absolute neutrophil count $\geq 1.5 \times 10^9 / L$ and platelets $\geq 75 \times 10^9 / L$ and have a previous response of complete remission, partial remission, or marrow complete remission may require dose reduction, if subsequent absolute neutrophil count nadir $< 0.500 \times 10^9 / L$, or platelets nadir $< 50 \times 10^9 / L$, if baseline was $> 100 \times 10^9 / L$, or platelets $< 50\%$ if baseline was $\leq 100 \times 10^9 / L$.

Results

[0102] Baseline characteristics for SE1 and SE2 are shown in Table 5. 22 Subjects in SE1 and 21 subjects in SE2 were compared with median (range) follow-up of 7.5 (1.0–8.9) and 7.9 (1.8–10.1) months, respectively, as shown in Table 6.

Table 5. SE1 and SE2 Baseline Characteristics

n (%)	SE1 n=22	SE2 n=21
Male	15 (68)	13 (62)
Median age, years [range]	70 [55–79]	68 [59–87]
ECOG performance score		
0	7 (32)	11 (52)
1	13 (59)	9 (43)
2	2 (9)	1 (5)
Bone marrow blasts		
≤ 5%	1 (4.5)	2 (9.5)
> 5% to ≤ 10%	7 (32)	6 (29)
> 10% to ≤ 20%	14 (64)	13 (62)
Baseline transfusion dependence		
RBC or platelet	13 (59)	10 (48)
RBC	13 (59)	9 (43)
Platelet	3 (14)	2 (10)
IPSS risk classification		
Intermediate-2	17 (77)	15 (71)
High	5 (23)	6 (29)
IPSS-R risk classification		
Intermediate	5 (23)	3 (14)
High	5 (23)	7 (33)
Very high	12 (54)	11 (52)
Baseline cytopenias (Grade ≥3)		
Neutropenia	14 (64)	14 (67)
Thrombocytopenia	6 (27)	7 (33)
Leukopenia	10 (46)	10 (48)
Anemia	3 (14)	3 (14)

Table 6: Follow-up Time and Dose Administration

	SE1 n=22	SE2 n=21
Median follow-up, months (min, max)	7.5 (1.0, 8.9)	7.9 (1.8, 10.1)
Azacitidine		
Median duration, days	156	184
Cycles, median (range)	4 (1–17)	5 (1–11)
Patients with dose reductions in Azacitidine, n (%):		
0 reductions	16 (73)	14 (67)
1 reduction	5 (23)	5 (24)
2 reductions	0	2 (10)

>2 reductions	1 (4.5)	0
Median days to dose modification	125.5	115
Venetoclax		
Median duration, days	156	184
Median days dosed	56	70
# of reductions in Venetoclax duration*	3 (14)	0
Patients with ≥ 1 Ven dose amount reduction, n (%)	10 (45)	10 (48)
Median days to duration reduction	52	n/a

*Reductions due to hematologic toxicity

[0103] The summary of adverse events in >20% of subjects is shown in Table 7. A similar frequency of \geq Grade 3 hematologic treatment-emergent adverse events (approximate %) were reported in SE1 and SE2, respectively, including anemia (14% and 33%), febrile neutropenia (46% and 48%), leukopenia (36% and 19%), neutropenia (55% and 48%) and thrombocytopenia (32% and 38%). Infections (59% and 38%) and leukopenia (36% and 19%) were more frequent in SE1 than SE2.

Table 7: Summary of Adverse Events

n (%)	SE1 n=22	SE2 n=21
Any Adverse Event		
Infections	13 (59)	8 (38)
Neutropenia	12 (55)	10 (48)
Febrile neutropenia	10 (46)	10 (48)
Thrombocytopenia	8 (36)	10 (48)
Leukopenia	8 (36)	4 (19)
Anemia	3 (14)	10 (48)
Nausea	7 (32)	13 (62)
Constipation	8 (36)	9 (43)
Vomiting	8 (36)	11 (52)
Diarrhea	6 (27)	12 (57)
Grade ≥ 3 Adverse Events		
Neutropenia	12 (55)	10 (48)
Febrile neutropenia	10 (46)	10 (48)
Infections	9 (41)	4 (19)
Leukopenia	8 (36)	4 (19)
Thrombocytopenia	7 (32)	8 (38)
Anemia	3 (14)	7 (33)
Serious Adverse Events^a		
Febrile neutropenia	8 (36)	9 (43)

Infections	8 (36)	3 (14)
GI disorders	3 (13)	3 (14)

^a Includes death, life-threatening, requiring hospitalization or surgical intervention, persistent/significant disability.

^b SE1: abdominal pain, diverticular perforation, and gastroesophageal reflux disease; SE2: nausea, pancreatitis, vomiting, and gastrointestinal hemorrhage

[0104] Worsening of treatment-emergent adverse events grades from baseline was analyzed by cycle. As shown in **FIGs. 3A-3F** and **FIGs. 4A-4C**, adverse event progression remains low after the first few cycles, such as cycles 1 and 2. **FIGs. 3A-3F** are hematologic toxicity showing the number of patients with worsening common terminology criteria grade over baseline per cycle. **FIGs. 4A-4C** are gastrointestinal toxicity showing the number of patients with worsening common terminology criteria grade over baseline per cycle.

[0105] Response rates were identical for SE1 and SE2 as shown in Table 8: 86% of subjects in both SE1 and SE2 had complete remission (CR) or marrow complete remission (mCR). For subjects with mCR, hematologic improvement occurred in 50% of SE1 and 46% of SE2 subjects.

Table 8: Response Rates

n (%)	SE1 n=22	SE2 n=21
Overall Response Rate (CR+mCR+PR)	86.30%	85.7%
Best Response		
Complete remission	7 (32)	5 (24)
Marrow complete remission	12 (55)	13 (62)
+ hematologic improvement	6 (50)	6 (46)
- hematologic improvement	6 (50)	7 (54)
Partial remission	0	0
Stable disease	2 (9)	2 (10)
Progressive disease	0	0
Not evaluable	1 (5)	1 (5)
Time on study, median months [min, max]	7.5 [1.0, 8.9]	7.9 [1.8, 10.1]

[0106] The summary of cycle delays is shown in Table 9. Cycle delays were comparable between SE1 and SE2, with a slightly longer duration for SE1 in early cycles.

Table 9: Summary of Cycle Delays

Venetoclax dose delay by end of each cycle	SE1 n=22		SE2 n=21	
	n (%)	Median # days (range)	n (%)	Median # days (range)
Cycle 2	14 (63)	7.5 (1-21)	13 (62)	8 (5-21)
Cycle 3	12 (55)	14 (3-28)	10 (48)	8.5 (1-48)
Cycle 4	8 (36)	17 (1-54)	6 (28)	9 (6-43)
Cycle 5	7 (32)	14 (3-28)	9 (43)	14 (1-97)
Cycle 6	7 (32)	7 (1-21)	5 (24)	14 (6-91)

[0107] The mean value of absolute neutrophil count and platelet count is shown in **FIGs. 1** and **2**, respectively. The number of count observations per study cycle day for both absolute neutrophil count and platelet count is shown in Table 10.

Table 10: Number of Count Observations per Study Cycle Day

Study Cycle Day	Absolute Neutrophil Count Observations		Platelet Count Observations	
	SE1	SE2	SE1	SE2
Baseline	22	21	19	19
C1D4	16	19	15	18
C1D7	11	13	9	12
C1D9	9	8	8	7
C1D15	20	20	20	17
C1D22	20	18	19	16
C2D1	8	12	8	12
C2D4	5	6	5	6
C2D7	9	12	9	11
C2D9	7	6	7	6
C2D15	17	17	17	16
C2D22	19	16	20	17
C3D1	15	14	15	13
C3D4	3	3	3	3
C3D7	9	10	8	10
C3D9	4	4	4	4
C3D15	13	16	12	16
C3D22	11	13	11	12
C4D1	15	11	15	12
C4D7	11	5	10	5
C4D9	3	5	3	5

C4D22	13	11	13	12
C5D1	8	5	8	6
C5D7	6	6	6	6
C5D9	2	2	2	2
C5D22	13	12	13	12
C6D1	4	8	4	7
C6D7	7	5	7	5
C6D9	0	2	0	2
C6D22	10	9	10	9

[0108] 86% of patients in both SE1 and SE2 had complete remission or marrow complete remission.

[0109] It is understood that the foregoing detailed description and accompanying examples are merely illustrative and are not to be taken as limitations upon the scope of the invention, which is defined solely by the appended claims and their equivalents. Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art. Such changes and modifications, including those relating to the methods of use of the invention, may be made without departing from the spirit and scope thereof. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

WE CLAIM:

1. A method for treating myelodysplastic syndromes in a human subject, comprising administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m² of azacitidine for 7 days in the 28 day dosing cycle;
wherein, if the human subject has:
a baseline absolute neutrophil count of $< 1.5 \times 10^9/L$, a baseline white blood cell count of $< 3 \times 10^9/L$, or a baseline platelet count of $< 75 \times 10^9/L$;
no improvement in cell line differentiation; and
at least one of a condition selected from the group consisting of:
 - a. an absolute neutrophil count nadir of $\geq 50\%$ reduction;
 - b. a white blood cell count nadir of $\geq 50\%$ reduction; and
 - c. a platelet count nadir of $\geq 50\%$ reduction;the daily dose of azacitidine of 75 mg/m² is reduced to $\leq 75\%$ but no less than 33% during a next 28 day dosing cycle.
2. The method of claim 1, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes.
3. The method of claim 1 or 2, wherein the daily dose of azacitidine of 75 mg/m² is reduced to 75% during the next 28 day dosing cycle;
wherein the human subject has a bone marrow cellularity of 30-60%; and
wherein the human subject has at least one of a condition selected from the group consisting of:
 - a. an absolute neutrophil count nadir decrease of $\geq 75\%$ from the baseline absolute neutrophil count;
 - b. a white blood cell count nadir decrease of $\geq 75\%$ from the baseline white blood cell count; and
 - c. a platelet count nadir decrease of $> 75\%$ from the baseline platelet count.
4. The method of claim 1-3, wherein the azacitidine is administered intravenously.
5. The method of claim 1-3, wherein the azacitidine is administered subcutaneously.
6. The method of claim 3-5, wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of:

- a. a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir;
 - b. a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir; and
 - c. a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir.
7. The method of claim 6, further comprising a delay prior to the next 28 day dosing cycle of ≥ 1 day.
8. The method of claim 6 or 7, wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle;
wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of:
- a. a subsequent absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir;
 - b. a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir; and
 - c. a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir.
9. The method of claim 1 or 2, wherein the daily dose of azacitidine of 75 mg/m² is reduced to 50% during the next 28 day dosing cycle;
wherein the human subject has a bone marrow cellularity of 15 to $< 30\%$; and
wherein the human subject has at least one of a condition selected from the group consisting of:
- a. an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count;

- b. a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count; and
 - c. a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count.
10. The method of claim 9, wherein the azacitidine is administered intravenously.
11. The method of claim 9, wherein the azacitidine is administered subcutaneously.
12. The method of claim 9-11, wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of:
- a. a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir;
 - b. a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir; and
 - c. a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir.
13. The method of claim 12, further comprising a delay prior to the next 28 day dosing cycle of ≥ 1 day.
14. The method of claim 12 or 13, wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle;
- wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of:
- a. a subsequent absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir;
 - b. a subsequent white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir; and
 - c. a subsequent platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir.
15. The method of claim 1 or 2, wherein the daily dose of azacitidine of 75 mg/m² is reduced to 33% during the next 28 day dosing cycle;

wherein the human subject has a bone marrow cellularity $< 15\%$; and

wherein the human subject has at least one of a condition selected from the group consisting of:

- a. an absolute neutrophil count nadir decrease of $\geq 50\%$ from the baseline absolute neutrophil count;
- b. a white blood cell count nadir decrease of $\geq 50\%$ from the baseline white blood cell count; and
- c. a platelet count nadir decrease of $\geq 50\%$ from the baseline platelet count.

16. The method of claim 15, wherein the azacitidine is administered intravenously.

17. The method of claim 15, wherein the azacitidine is administered subcutaneously.

18. The method of claim 15-17, wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of:

- a. a next absolute neutrophil count of $\leq 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir;
- b. a next white blood cell count of $\leq 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir; and
- c. a next platelet count of $\leq 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir.

19. The method of claim 18, further comprising a delay prior to the next 28 day dosing cycle of ≥ 1 day.

20. The method of claim 18 or 19, wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle;

wherein the human subject has prior to a start of the subsequent 28 day dosing cycle, at least one condition selected from the group consisting of:

- a. a subsequent neutrophil count of $> 25\%$ above the absolute neutrophil count nadir and relative to a difference of the baseline absolute neutrophil count and the absolute neutrophil count nadir;

- b. a subsequent white blood cell count of $> 25\%$ above the white blood cell count nadir and relative to a difference of the baseline white blood cell count and the white blood cell count nadir; and
 - c. a subsequent platelet count of $> 25\%$ above the platelet count nadir and relative to a difference of the baseline platelet count and the platelet count nadir.
21. A method for treating myelodysplastic syndromes in a human subject, comprising administering to the human subject a daily dose of 400 mg of venetoclax for 14 days in a 28 day dosing cycle, and a daily dose of 75 mg/m^2 of azacitidine for 7 days in the 28 day dosing cycle;
- wherein, if the human subject has:
- a baseline absolute neutrophil count of $\geq 1.5 \times 10^9/\text{L}$,
 - a baseline white blood cell count of $\geq 3 \times 10^9/\text{L}$, or
 - a baseline platelet count of $\geq 75 \times 10^9/\text{L}$; and
- at least one of a condition selected from the group consisting of:
- a. an absolute neutrophil count nadir of $\leq 1.5 \times 10^9/\text{L}$; and
 - b. a platelet count nadir of $\leq 50 \times 10^9/\text{L}$;
- the daily dose of azacitidine is reduced from 75 mg/m^2 to $\leq 67\%$ but no less than 50% during the next 28 day dosing cycle.
22. The method of claim 21, wherein the myelodysplastic syndromes are treatment-naïve higher-risk myelodysplastic syndromes.
23. The method of claim 21 or 22, wherein the daily dose of azacitidine is reduced from 75 mg/m^2 to 67%; and
- wherein the human subject has at least one condition selected from the group consisting of:
- a. the absolute neutrophil count nadir is from $0.5 \times 10^9/\text{L}$ to $1.5 \times 10^9/\text{L}$; and
 - b. the platelet count nadir is from $25 \times 10^9/\text{L}$ to $50 \times 10^9/\text{L}$.
24. The method of claim 21-23, wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of:
- a. a next absolute neutrophil count of $\leq 1.5 \times 10^9/\text{L}$; and
 - b. a next platelet count of $\leq 50 \times 10^9/\text{L}$.

25. The method of claim 24, further comprising a delay prior to the next 28 day dosing cycle of \geq 1 day.
26. The method of claim 21-24, wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle;
wherein the human subject has prior to a start of the subsequent 28 day dosing cycle at least one condition selected from the group consisting of:
- a subsequent absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and
 - a subsequent platelet count of $\leq 50 \times 10^9/L$.
27. The method of claim 21 or 22, wherein the daily dose of azacitidine is reduced from 75 mg/m² to 50%; and
wherein the human subject has at least one condition selected from the group consisting of:
- the absolute neutrophil count nadir is $< 0.5 \times 10^9/L$; and
 - the platelet count nadir is $< 25 \times 10^9/L$.
28. The method of claim 27, wherein the human subject has prior to a start of the next 28 day dosing cycle, at least one condition selected from the group consisting of:
- a next absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and
 - a next platelet count of $\leq 50 \times 10^9/L$.
29. The method of claim 28, further comprising a delay prior to the next 28 day dosing cycle of \geq 1 day.
30. The method of claim 28, wherein the daily dose of 400 mg of venetoclax is reduced from 14 to 7 days in a subsequent 28 day dosing cycle;
wherein the human subject has prior to a start of the subsequent 28 day dosing cycle at least one condition selected from the group consisting of:
- a subsequent absolute neutrophil count of $\leq 1.5 \times 10^9/L$; and
 - a subsequent platelet count of $\leq 50 \times 10^9/L$.




FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E




FIG. 3F




FIG. 4A




FIG. 4B




FIG. 4C

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 22/28429

A. CLASSIFICATION OF SUBJECT MATTER

IPC - A61K 31/167; A61K 31/18; A61K 31/337 (2022.01)

CPC - A61K 31/167; A61K 31/18; A61K 31/337; A61K 31/4025

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Samra et al., "Venetoclax-Based Combinations in Acute Myeloid Leukemia: Current Evidence and Future Directions" 05 November 2020 (05.11.2020) entire document especially Page 7, Para 3; Page 10, Para 1	1-3, 9-11, 15-17, 21-23, and 27-30
A	DiNardo et al., "Azacitidine and Venetoclax in Previously Untreated Acute Myeloid Leukemia" 13 August 2020 (13.08.2020) entire document especially Page 618, Para 1; Page 618, Para 5; Page 618, Para 6	1-3, 9-11, 15-17, 21-23, and 27-30
A	WO 2020/257671 A1 (CELGENE CORPORATION) 24 December 2020 (24.12.2020) entire document especially Claim 1; Claim 14; Claim 15	1-3, 9-11, 15-17, 21-23, and 27-30
A	US 2016/0235779 A1 (Cantex Pharmaceuticals, Inc.) 18 August 2016 (18.08.2016) entire document	1-3, 9-11, 15-17, 21-23, and 27-30
A	US 2021/0023237 A1 (ImmunoGen, Inc.) 28 January 2021 (28.01.2021) entire document	1-3, 9-11, 15-17, 21-23, and 27-30

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"D" document cited by the applicant in the international application

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

30 June 2022

Date of mailing of the international search report

AUG 01 2022

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Kari Rodriguez

Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 22/28429

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 1-3, 9-11, 15-17, 21-23, and 27-30
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.