

(19) **DANMARK**

(10) **DK/EP 2590949 T5**



(12) **Rettet oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 D 215/56 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-05-23**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2015-11-25**
- (86) Europæisk ansøgning nr.: **11732421.0**
- (86) Europæisk indleveringsdag: **2011-07-07**
- (87) Den europæiske ansøgnings publiceringsdag: **2013-05-15**
- (86) International ansøgning nr.: **EP2011061490**
- (87) Internationalt publikationsnr.: **WO2012004338**
- (30) Prioritet: **2010-07-09 EP 10169162** **2010-07-09 US 362849 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Active Biotech AB, Box 724, 220 07 Lund, Sverige**
- (72) Opfinder: **HOLMBERG, Pär Henning, Gustavsgatan 20A, 691 34 Karlskoga, Sverige**
BOCK, Lillemor Maria, HögÅysen 812, S-691 93 Karlskoga, Sverige
JANSSON, Karl-Erik, SpannmÅylsvägen 11, S-247 55 Dalby, Sverige
- (74) Fuldmægtig i Danmark: **Chas. Hude A/S, H.C. Andersens Boulevard 33, 1780 København V, Danmark**
- (54) Benævnelse: **FREMGANGSMÅDE TIL FREMSTILLING AF QUINOLIN-3-CARBOXAMIDER**
- (56) Fremdragne publikationer:
US-A- 3 868 379
US-B2- 6 875 869
TEN HAKEN: "Use of molecular sieves for affecting the equilibrium of reactions", CHEMISTRY & INDUSTRY, SOCIETY OF CHEMICAL INDUSTRY. LONDON, GB, no. 7, 1973, page 325, XP008130938, ISSN: 0009-3068

Description**Field of the invention**

5

[0001] The present invention relates to the field of organic synthesis. In particular, it relates to the field of chemical synthesis of certain derivatives of N-alkyl-N-phenyl-quinoline-3-carboxamides.

Background of the invention




10

[0002] N-alkyl-N-phenyl-quinoline-3-carboxamides are a promising class of compounds being evaluated in clinical trials against different classes of diseases, e.g. autoimmune diseases and cancer.

15

[0003] N-alkyl-N-phenyl-quinoline-3-carboxamides such as paquinimod (herein below also referred to as **A**), laquinimod (herein below also referred to as **B**), and tasquinimod (herein below also referred to as **C**), have been prepared by a method involving distillation of the volatiles from a mixture comprising an ester,


20 aniline and an aliphatic solvent such as n-heptane or n-octane.

Paquinimod (**A**)Laquinimod (**B**)Tasquinimod (**C**)

[0004] This method is described in US patent No. 6,875,869.

25 **[0005]** The prior art synthetic protocol (Org. Process. Res&Dev. 2007, 11, 674-680) for N-alkyl-N-phenyl-quinoline-3-carboxamides such as paquinimod (**A**), laquinimod (**B**), and tasquinimod (**C**) is exemplified with synthesis of paquinimod in **Scheme 1**. The route starts with an an-thranilic acid **1** which is

transformed into an isatoic anhydride **2**. The isatoic anhydride is methylated to give **3**, which is condensed with dimethylmalonate to give the corresponding methyl ester **4**. The methyl ester is subsequently condensed with the appropriate aniline, to give the desired final compound.




Scheme 1. The synthetic route to N-alkyl-N-phenyl-quinoline-3-carboxamides, exemplified with the synthesis of paquinimod (**A**) via the corresponding methyl ester **4**.

[0006] This manufacturing method is short and avoids the use of expensive reagents. All intermediates are stable and easy to isolate in high purity by precipitation and filtration. The main impurity in the final condensation step is remaining alkyl ester. Alternative methods, such as carbodiimide mediated coupling between a carboxylic acid and aniline, or the condensation of N-alkyl-N-phenyl-malonamic acid methyl ester with an isatoic anhydride (US patent No. 5,912,349) are either longer or yield product of lower purity.

[0007] The final condensation step is an equilibrium (**Scheme 2**) that favors the alkyl ester and a reaction mechanism involving a ketene intermediate **5** has been strongly indicated (J. Org. Chem. 2006, 71, 1658-1667 and J Phys. Chem.

3



Scheme 2. An intermediate ketene **5** is involved in the equilibrium between **4** and **A**.

[0008] Usually formation of amides from esters and anilines requires very high reaction temperatures that also cause extensive byproduct formation. The above-mentioned reaction is enabled at moderate temperatures by the ability of **4** to unimolecularly form the ketene intermediate **5** instead of a tetrahedral intermediate. The method, which is described e.g. in US patent No. 6,875,869, comprises charging the reactor with an appropriate ester and an aniline derivative in an aliphatic solvent such as n-heptane or n-octane. The equilibrium is driven towards the desired product by distilling off the solvent and any formed alcohol. After complete reaction the mixture is cooled and the precipitated raw product is isolated by filtration.


[0009] Unlike most other reactions where esters are transformed into thermodynamically more stable amides this particular reaction needs a very efficient removal of formed alcohol in order to give a high yield.

[0010] The method described above has been used for GMP (Good Manufacturing Practice) manufacturing of paquinimod, laquinimod and tasquinimod.

Summary of the invention

[0011] As noted herein above, in the reaction described in Org. Process. Res&Dev. 2007, 11, 674-680 for preparing an N-alkyl-N-phenyl-quinoline-3-carboxamide, the main impurity in the isolated raw product is the alkyl ester,

used as the starting material in the final condensation step. This condensation may be represented by to the following general reaction scheme:



5 **[0012]** In the reaction as represented herein above, the remaining ester (II) may be easily removed from the product by recrystallisation, if small amounts of the ester are present. On the other hand, the present inventors have discovered that if larger amounts of remaining ester are present in the raw product, only one recrystallisation may not be enough for achieving a pure product. It is
 10 therefore important to drive the equilibrium as far as possible to the product side, by removal of the alcohol (IV) that is formed as a byproduct in the reaction.


[0013] The present inventors also have discovered, as a further problem, that the removal of the alcohol (IV) is scale-dependent. Thus, when the
 15 condensation reaction is performed on a small scale (e.g. less than 20 g), most of the ester (II) is transformed into product (I) within a few hours of distillation. In this case, the precipitated product contains only low, and acceptable, amounts of the ester as an impurity. However, on a larger scale, e.g. kilogram scale, the process becomes less predictable and in some cases a substantial amount of
 20 the ester (II) still persists in the mixture even after 20 hours of distillation. In such cases more solvent has to be added and the distillation continued until most of the ester (II) is transformed into the desired product (I). A process that consumes large and unpredictable amounts of the aliphatic solvent in order to get full conversion is far from optimal for drug manufacturing.

25

[0014] The present inventors therefore set out to solve the problem of providing an improved method for preparing a compound of formula (I), as defined herein, in a high yield, advantageously without the above-mentioned drawbacks. In particular, this method should allow preparing the compound according to

formula (I) on a large scale in high yield and in high purity, advantageously without the need for prolonged reaction times and excessive solvent and energy consumption.

- 5 **[0015]** Accordingly, a method is provided for preparing a compound of formula (I)



wherein


R1 is ethyl, R2 is ethyl and R3 is H; or

- 10 R1 is chloro, R2 is ethyl and R3 is H; or

R1 is methoxy, R2 is methyl and R3 is trifluoromethyl;

by

1. (i) reacting a compound of formula (II)




- 15 wherein

R1 is ethyl, chloro or methoxy; and

R4 is C1-C4 alkyl;

with a compound of formula (III)



- 20 wherein

R2 is ethyl and R3 is H; or

R2 is methyl and R3 is trifluoromethyl;

in a refluxing mixture containing an aliphatic solvent or a mixture of aliphatic solvents, said aliphatic solvent(s) having a boiling point in the range of 68-191

- 25 °C;

2. (ii) condensing vapors of the refluxing mixture;
3. (iii) treating the condensed vapors with an alcohol scavenging agent selected from molecular sieves or with an alcohol scavenging agent which is a mixture of molecular sieves; and
- 5 4. (iv) returning the condensed vapors back to the reaction mixture.

[0016] The present inventors have found that by the method as defined herein, a higher reaction yield may be obtained in a shorter reaction time, compared to
10 the prior art methods. This in turn results in reduced energy consumption which, in addition to a reduced solvent consumption during the reaction, is very advantageous from an economic and environmental point of view. Even more importantly, by the method of the present invention, a reaction product of a higher purity is obtainable. Very advantageously, a reaction product at a higher
15 yield and improved purity is obtained in a shorter reaction time and at a lower solvent and energy consumption than according to the prior art methods, in particular when performing the reaction on a large scale.

[0017] Other benefits of the present invention are that the use of an alcohol
20 scavenging agent enables a higher distillation rate without reducing the reaction volume, thus a shorter reaction time is obtained; the solvent consumption is decreased to a minimum, by recycling the solvent; there is no need for careful monitoring of the distillation rate and there is an almost complete conversion of the ester, resulting in higher purity of the isolated product.

25 **Brief description of the drawings**

[0018]

Figure 1 represents the modified Soxhlet-apparatus used in the experiments for the evaluation of scavenging agents; and

30 **Figure 2** represents an equipment set-up in pilot plant for production of Paquinimod.

Detailed Description of the Invention

The alcohol scavenging agent

5

[0019] In the method of the present invention, the distillate is treated with an alcohol scavenging agent before being recycled back to the reaction mixture. The alcohol scavenging agent is selected from molecular sieves, or a mixture of any of these.

10

[0020] The present inventors have tested numerous alcohol scavenging agents in combination with different solvents and have found that molecular sieves are particularly useful for the manufacturing of compounds of formula (I) as defined herein.

15

[0021] Molecular sieves are crystalline metal alumina silicates having a three dimensional interconnecting network of silica and alumina tetrahedra. Molecular sieves are capable of adsorbing various polar compounds into the cavities formed by the network. Different qualities of molecular sieves exist, such as the powder forms of the 3A, 4A, 5A and 13X sieves, which may be purchased e.g. from Sigma-Aldrich Co.

20

[0022] According to the present invention, the alcohol scavenging agent is selected from molecular sieves or is a mixture of molecular sieves. For example, the alcohol scavenging agent may be selected from 3A, 4A, 5A, 13X molecular sieves or may be a mixture of any of said molecular sieves.

25

[0023] The composition of the 3A molecular sieve is approximately $0.6 \text{ K}_2\text{O} : 0.40 \text{ Na}_2\text{O} : 1 \text{ Al}_2\text{O}_3 : 2.0 \pm 0.1 \text{ SiO}_2 : x \text{ H}_2\text{O}$. This form is obtained by substituting potassium cations for the inherent sodium ions of the 4A structure, giving a pore size of approximately $3\text{\AA}$. The 3A molecular sieve is known mainly for use in removing moisture.

30

[0024] The 4A molecular sieve has a composition that is approximately 1 Na₂O : 1 Al₂O₃ : 2.0 ± 0.1 SiO₂ : x H₂O. 4A molecular sieve has an effective pore opening of about 4 angstroms (0.4nm). This sieve is generally considered as a universal drying agent in polar and nonpolar media.

[0025] The 5A molecular sieve is of the approximate composition 0.80 CaO : 0.20 Na₂O : 1 Al₂O₃ : 2.0 ± 0.1 SiO₂ : x H₂O and the pore size is about 5Å.

10 [0026] The 13X molecular sieve is of the approximate composition 1 Na₂O : 1 Al₂O₃ : 2.8 ± 0.2 SiO₂ : xH₂O. The pore size of the 13X molecular sieve is about 10 Å.

[0027] Other molecular sieves may equally well be used and it is contemplated that the skilled person will be able, by simple routine experimentation, to identify molecular sieves or mixture of molecular sieves that are suitable for use in the present invention. A further description of molecular sieves may be found in the book Molecular Sieves, R. Szostak, ed. Blackie Academic & Professional (ISBN 0 75140480 2).

20

[0028] The amount of alcohol scavenging agent used in the method of the invention suitably is such that the weight ratio of alcohol scavenging agent to ester (II) is at least 1:1. Preferably, the alcohol scavenging agent is used in excess over the ester (II), e.g. at a weight ratio of alcohol scavenging agent to ester (II) of 2:1, or 3:1, or 5:1, or even higher.

25

The aliphatic solvent

[0029] The aliphatic solvent or mixture of aliphatic solvents for use in the method of the invention should have a boiling point in the range of 68-191 °C, preferably 80-150 °C, more preferably 95-130 °C. Such solvents are, for example selected from C6-C10 branched or linear alkanes and cycloalkanes, e.g. C7 or C8 branched or linear alkanes and cycloalkanes.

30

[0030] For the purpose of the present invention, a C6-C10 branched or linear alkane or cycloalkane refers to an alkane or cycloalkane having in total from 6 to 10 atoms.

5

[0031] A C7 or C8 branched or linear alkane refers to an alkane having in total 7 or 8 carbon atoms, e.g. n-heptane, n-octane, 2,2,4-trimethylpentane etc.

[0032] The cycloalkane may be mono- or polycyclic, e.g. bicyclic. Furthermore, the cycloalkane may be either unsubstituted or substituted (i.e. branched) with one or several, branched or linear, alkyl groups, it being understood that the total number of carbon atoms in the ring and any alkyl substituent, taken together, is 6 to 10, in particular 7 or 8. An example of a C10 cycloalkane is decalin (i.e. decahydronaphthalene). Examples of C7-C8 cycloalkanes are methylcyclohexane and cyclooctane.

15


[0033] Other isomers of alkanes and cycloalkanes, as may easily come to the mind of the skilled reader, are equally possible as long as the above-mentioned boiling point requirement is fulfilled.

20

[0034] In one embodiment, the aliphatic solvent is selected from n-heptane, n-octane, methylcyclohexane, 2,2,4-trimethylpentane and cyclooctane, or a mixture of any of these. In another embodiment, the solvent is selected from n-heptane and n-octane or a mixture of any of these.

25 ***The compound of formula (II)***

[0035] In the compound of formula (II)




[0036] R4 is selected from C1-C4 alkyl, and may be branched or linear.

[0037] In one embodiment, R4 is selected from methyl, ethyl, n-propyl, iso-propyl and n-butyl. In another embodiment, R4 is C1-C3 alkyl, i.e. methyl, ethyl, n-propyl, and iso-propyl. In still another embodiment, R4 is methyl or ethyl, e.g. R4 is methyl.

[0038] In the condensation reaction between compounds (II) and (III), the aliphatic alcohol (IV) is formed as a byproduct and this aliphatic alcohol is withdrawn from the condensed vapors containing the aliphatic solvent or mixture of aliphatic solvents by treatment of the condensed vapors with an alcohol scavenging agent or a mixture of alcohol scavenging agents.

[0039] Some non-limiting examples of esters according to formula (II) are compounds 6-11, represented herein below.




[0040] When using e.g. 4 as starting compound (II), the formed alcohol (IV) will be methanol, when using e.g. 10 as starting compound (II), the formed alcohol (IV) will be ethanol etc.

The compound of formula (I)

[0041] The product obtained in the method of the invention is a compound of formula (I),

11



wherein

R1 is ethyl, R2 is ethyl and R3 is H; or

R1 is chloro, R2 is ethyl and R3 is H; or

5 R1 is methoxy, R2 is methyl and R3 is trifluoromethyl.

[0042] In one embodiment of the invention, R1 is ethyl, R2 is ethyl and R3 is H, i.e. the compound prepared is paquinimod.

10

[0043] In one other embodiment of the invention, R1 is chloro, R2 is ethyl and R3 is H, i.e. the compound prepared is laquinimod.

[0044] In still another embodiment of the invention, R1 is methoxy, R2 is methyl and R3 is trifluoromethyl, i.e. the compound prepared is tasquinimod.

15

Some preferred embodiments

[0045] In one suitable embodiment of the invention, R1 is ethyl, R2 is ethyl, R3 is H, R4 is methyl or ethyl; the alcohol scavenging agent is a 3A, 4A, 5A or 13X molecular sieve, or is a mixture thereof; and the aliphatic solvent(s) is/are selected from n-heptane and n-octane.

20

[0046] In one particular embodiment of the invention, R1 is ethyl, R2 is ethyl, R3 is H, R4 is methyl, the alcohol scavenging agent is a 4A molecular sieve, and the aliphatic solvent(s) is/are selected from n-heptane and n-octane.

25

[0047] In another particular embodiment, R1 is ethyl, R2 is ethyl, R3 is H, R4 is

ethyl, the alcohol scavenging agent is a 5A molecular sieve, and the aliphatic solvent is n-octane.

5 [0048] In still another embodiment, R1 is ethyl, R2 is ethyl, R3 is H, R4 is ethyl, the alcohol scavenging agent is a 3A molecular sieve, and the aliphatic solvent is n-octane.

10 [0049] In another embodiment, R1 is chloro, R2 is ethyl, R3 is H, R4 is methyl, the alcohol scavenging agent is a 3A, 4A, 5A, or 13X molecular sieve, and the aliphatic solvent(s) is/are selected from n-heptane and n-octane, and in particular is n-heptane.

15 [0050] In one other embodiment, R1 is methoxy, R2 is methyl, R3 is trifluoromethyl, R4 is methyl or ethyl; the alcohol scavenging agent is a 3A, 4A, 5A or 13X molecular sieve, or is a mixture thereof; and the aliphatic solvent is n-octane.

20 [0051] In one particular embodiment, R1 is methoxy, R2 is methyl, R3 is trifluoromethyl, R4 is methyl, the alcohol scavenging agent is a 4A molecular sieve, and the aliphatic solvent is n-octane.

25 [0052] Herein below, the invention will be illustrated by some non-limiting examples and comparative examples, whereby the alkyl esters of formula (II), used as starting material, will be referred to by the numbering indicated herein above (i.e. compounds Nos. 4 and 6-11).

Examples

30 [0053] Initial experiments were conducted on a 5-gram scale using the equipment shown in **Figure 1**. A special Soxhlet apparatus (not part of the invention), was designed for the evaluation. This Soxhlet apparatus cools the distillate before passing through the alcohol-scavenging material. In the classical Soxhlet apparatus the adsorption chamber has a temperature a few

degrees below the boiling point of the solvent. As the experiments confirm, molecular sieves are more effective scavenging agents at lower temperatures.

- 5 [0054] In using the modified Soxhlet apparatus, the round-bottom flask is charged with ester, the appropriate aniline and a solvent. The mixture is heated to reflux and a condensate is formed in the two liquid-cooled condensers. The condensate is passed by gravity through a pad of molecular sieves and then back into the reaction mixture.
- 10 [0055] The results are summarized in **Table 1**. Molecular sieves are better at scavenging alcohol, e.g. methanol, at lower temperatures (**Entry 17** and **20**). The scavenging of alcohol from aliphatic solvents proceeded more efficiently, than if the solvent was aromatic e.g. toluene. This can be concluded from **Entry 2** and **9** or **Entry 17** and **21**. The process was faster with molecular
- 15 sieves, consumed less solvent, was easier to control and yielded a product of higher purity than the prior art method (**Entry 17** and **22** or **Entry 25** and **26**, example 1 and 4 in experimental part, and US patent No. 6,875,869). Methyl esters gave an excellent result with molecular sieves of the types 3A, 4A, 5A, and 13X (**Entry 1, 2, 4, 17, 18, 25, 27, and 28**). Molecular sieves of the type 3A
- 20 and 5A were also effective when using an ethyl ester (**Entry 8 and 19**). The excellent result with the condensation of a methyl or an ethyl ester (**Entry 1** and **19**) in the presence of 3A molecular sieves is despite the discrepancy between the molecular size of methanol and ethanol and the 3 Angstrom cavity in the sieves. According to theory the critical diameter of methanol and ethanol
- 25 respectively is too large for 3A molecular sieves. However, 3A molecular sieves were not effective when an n-propyl ester was used as the starting material (**Entry 5**). The 13X molecular sieves were effective when using an n-butyl ester in the condensation reaction (**Entry 6**). The molecular sieves turned out to be very effective as alcohol-scavenging material in comparison with other tested
- 30 materials.

Table 1. Evaluation of scavenging agents.

Entry	Alkyl ester used as starting material	Conditions ^a	Product	Remaining ^b alkyl ester in product (mol%)	Yield ^c (weight%)
1	6	3A MS/n-heptane/4h	B	<1	95
2	6	Hot 4A MS/n-heptane/4h (classical Soxhlet)	B	<1	98
4	6	13X MS/n-heptane/4h	B	<1	94
5	7	3A MS/n-heptane/2.5h	B	23	
6	8	13X MS/n-octane/2.5h	B	<1	92
7	9	4A MS/n-heptane/2h	B	1.5	94
8	10	5A MS/n-octane/4h	A	<1	91
9	6	4A MS/toluene/4h	B	12	
11	6	4 h reflux in n-heptane without scavenging methanol	B	52	
17	4	4A MS/n-heptane/2h	A	2	82
18	4	4A MS/n-octane/3.5h	A	<1	89
19	10	3A MS/n-	A	<1	92

Entry	Alkyl ester used as starting material	Conditions ^a	Product	Remaining ^b alkyl ester in product (mol%)	Yield ^c (weight%)
		octane/2.5h			
20	4	Hot 4A MS/n-heptane/6h (classical Soxhlet)	A	5	
21	4	4A MS/toluene/ 2h	A	10	
22	4	2 hours distillation from n-heptane according to prior art	A	25	
24	11	4A MS/n-heptane/2h	C	87 ^d	
25	11	4A MS/n-octane/2h	C	1	99
26	11	2 hours distillation from n-octane according to prior art	C	4	94
27	6	5A MS/n-heptane/4h	B	<1	97
28	4	5A MS/n-octane/3.25h	A	<1	93

[0056] ^aThe starting material (5.0 g) and 2.0 equiv. of the appropriate aniline were suspended in the indicated solvent (150 mL) and the mixture was refluxed in the described Soxhlet-type apparatus, illustrated in **Figure 1**, containing 22 gram of the alcohol-scavenging material. The mixture was then cooled and the

precipitated product was isolated by filtration. In all entries the only detected impurity besides the desired product was the ester used as starting material.

^bRemaining alkyl ester (mol%) in isolated product according to ¹H-NMR.

^cIsolated yields are only given when the isolated product contained less than

5 5% remaining starting material. ^dLow conversion of ester is partly due to limited solubility of the starting material.

[0057] Compounds **A** (paquinimod), **B** (laquinimod), and **C** (tasquinimod) were described in J. Med. Chem. 2004, 47, 2075-2088. Large-scale synthesis of
10 compound **B** was described in Org. Process. Res&Dev. 2007, 11, 674-680.

[0058] Compound **6** (1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxylic acid methyl ester) was prepared from 2-amino-6-chlorobenzoic acid according to **Scheme 1** and further as described in Org. Process.
15 Res&Dev. 2007,11,674-680.

¹H NMR (CDCl₃) 14.9 (s, 1H), 7.52 (t, 1H), 7.27 (m, 2H), 4.04 (s, 3H), 3.66 (s, 3H).

[0059] Compound **7** (1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxylic acid n-propyl ester) was prepared from compound **B** by refluxing in
20 n-propanol overnight and adding dilute aqueous hydrochloric acid in order to precipitate compound **7**. ¹H NMR (CDCl₃) 15.06 (s, 1H), 7.50 (t, 1H), 7.27 (d, 1H), 7.23 (d, 1H), 4.39 (t, 2H), 3.64 (s, 3H), 1.87 (pentet, 2H), 1.07 (t, 3H).

[0060] Compound **8** (1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxylic acid n-butyl ester) was prepared from compound **B** by heating in n-butanol at 100 °C overnight and adding dilute aqueous hydrochloric acid and methanol in order to precipitate compound **8**. ¹H NMR (CDCl₃) 15.08 (s, 1H),
25 7.50 (t, 1H), 7.28 (d, 1H), 7.24 (d, 1H), 4.43 (t, 2H), 3.64 (s, 3H), 1.82 (pentet, 2H), 1.52 (sextet, 2H), 0.98 (t, 3H).
30

[0061] Compound **4** (1,2-dihydro-4-hydroxy-5-ethyl-1-methyl-2-oxo-quinoline-3-carboxylic acid methyl ester) was prepared from 2-amino-6-ethylbenzoic acid

according to **Scheme 1** and further as described for compound **6** in Org. Process. Res&Dev. 2007, 11, 674-680. ¹H NMR (CDCl₃) 14.9 (s, 1H), 7.57 (t, 1H), 7.22 (d, 1H), 7.08 (d, 1H), 4.06 (s, 3H), 3.68 (s, 3H), 3.27 (q, 2H), 1.31 (t, 3H).

5

[0062] Compound **10** (1,2-dihydro-4-hydroxy-5-ethyl-1-methyl-2-oxo-quinoline-3-carboxylic acid ethyl ester) was prepared from compound **A** by refluxing in ethanol overnight and adding cold dilute aqueous hydrochloric acid in order to precipitate compound **10**. Compound **10** is also described in J. Med. Chem. 2004, 47, 2075-2088. ¹H NMR (CDCl₃) 15.02 (s, 1H), 7.53 (t, 1H), 7.18 (d, 1H), 7.04 (d, 1H), 4.50 (q, 2H), 3.64 (s, 3H), 3.25 (q, 2H), 1.48 (t, 3H), 1.28 (t, 3H).

[0063] Compound **11** (1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxylic acid methyl ester) was prepared from 2-methylamino-6-methoxybenzoic acid (US patent No. 6,133,285) according to **Scheme 1** and as described for compound **6** in Org. Process. Res&Dev. 2007, 11, 674-680. ¹H NMR (CDCl₃) 13.8 (s, 1H), 7.57 (t, 1H), 6.93 (d, 1H), 6.73 (d, 1H), 4.02 (s, 3H), 4.01 (s, 3H), 3.63 (s, 3H).

[0064] Compound **9** (1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxylic acid isopropyl ester) was prepared from compound **B** by refluxing in 2-propanol overnight and adding dilute aqueous hydrochloric acid in order to precipitate compound **9**. ¹H NMR (CDCl₃) 15.2 (s, 1H), 7.51 (t, 1H), 7.28 (d, 1H), 7.24 (d, 1H), 5.37 (septet, 1H), 3.67 (s, 3H), 1.47 (d, 6H).

25 **Example 1**

Preparation of compound A (Entry 17)

[0065] A mixture of **4** (5.00 g) and N-ethylaniline (4.7 g) in n-heptane (200 mL) was refluxed in a Soxhlet extraction apparatus (**Figure 1**) containing 4A molecular sieves (22.9 g). The reflux was stopped after 2.0 hours and the mixture was cooled to room temperature. The crystalline suspension was

30

filtered. The precipitate was washed with n-heptane and dried in vacuum to furnish 5.53 g (82 %) of N-ethyl-N-phenyl-5-ethyl-1,2-dihydro-4-hydroxy-1-methyl-2-oxo-quinoline-3-carboxamide (**A**). ¹H-NMR analysis on the isolated product revealed 2% remaining ester. ¹H-NMR (d-DMSO), all broad peaks; 11.1 (1H), 7.41 (1H), 7.29 (2H), 7.21 (3H), 7.14 (1H), 6.96 (1H), 3.80 (2H), 3.42 (3H), 3.08 (2H), 1.07 (3H), 1.06 (3H).

[0066] When the same reaction was performed by the traditional distillation from n-heptane (**Entry 22**) during 2 hours according to prior art US patent No. 6,875,869 the product was isolated in 85 % yield (based on compound **A**) and found by ¹H-NMR to be a mixture of compound **A** (75 mol%) and the ester used as starting material (25 mol%).

Example 2

15 Preparation of compound B (Entry 2)

[0067] A mixture of **6** (5.00 g) and N-ethylaniline (5.0 g) in n-heptane (150 mL) was refluxed for 4.0 hours in a Soxhlet apparatus containing 4A molecular sieves (22.9 g). After cooling the mixture the product was isolated as above to furnish 6.54 g (98 %) of N-ethyl-N-phenyl-5-chloro-1,2-dihydro-4-hydroxy-1-methyl-2-oxo-quinoline-3-carboxamide (**B**). ¹H-NMR analysis on the isolated product revealed no impurities. ¹H-NMR (CDCl₃) 12.6 (s, 1H), 7.41 (t, 1H), 7.08-7.26 (m, 7H), 3.98 (q, 2H), 3.30 (s, 3H), 1.22 (t, 3H).

25 Example 3

Preparation of compound C (Entry 25)

[0068] A mixture of **11** (5.00 g, 18.9 mmol), N-methyl-p-trifluoromethylaniline (5.13 g, 28.4 mmol) and n-octane (200 mL) were refluxed in a Soxhlet extraction apparatus containing 4A molecular sieves (22.9 g) for 2 hours. After cooling the mixture the product was isolated as above to furnish 7.6 g (99 %) of

4-hydroxy-5-methoxy-N,1-dimethyl-2-oxo-N-[(4-trifluoromethyl)phenyl]-1,2-dihydroquinoline-3-carboxamide (**C**). ¹H-NMR analysis on the isolated product revealed no impurities other than 1 mol% remaining ester **11**. ¹H-NMR (CDCl₃) 9.9 (s, 1H), 7.50 (bs, 4H), 7.46 (t, 1H), 6.94 (d, 1H), 6.70 (d, 1H), 4.06 (s, 3H),
5 3.54 (s, 3H), 3.48 (s, 3H).

[0069] When the same reaction was performed by the traditional distillation from n-octane (**Entry 26**) during 2 hours according to prior art US patent No. 6,875,869 the product was isolated in 94 % yield and determined by ¹H-NMR
10 analysis to consist of a mixture of compound **C** (96 mol%) and the starting material **11** (4 mol%).

Example 4

Large-Scale Manufacturing of Compound A

15

[0070] The equipment set-up illustrated in **Figure 2** was used in a large-scale manufacturing of Paquinimod. A glass-lined reactor was charged with **4** (11.6 kg) through the manhole, followed by the addition of n-octane (212 kg) and n-ethyl aniline (13.4 kg). The mixture was heated to reflux temperature, i.e. 125°C. n-Octane was distilled off to a glass-sphere (156) at atmospheric pressure. The distillate in pos 156 was pumped through a bag filter (194) filled with 4A molecular sieves (7.5 kg), when the volume had become 70 L. The distillation was continuous during the operation. The distillate was re-circulated
20 back to the reactor as well as to the glass sphere to allow a continuous flow through the filter. After 7 hours the mixture was cooled to ≤ 60°C. An in process control (**IPC**) was withdrawn to check the conversion according to actual specification. Immediately after the sampling the mixture was heated to reflux temperature. Distillation and re-circulation through a second bag filter (195)
25 filled with molecular sieve (7.5 kg) were continued. Since the first sample was not approved a second sample was withdrawn after additional 4.5 hours of re-circulation. Sample no. 2 was approved according to the specification. The results are presented in the **Table 2**.
30

Table 2. IPC results.

Sample	IPC analysis	Criteria (%)	Result (%)
1 (7 hours)	Starting material 4	≤2	8.6
2 (11.5 hours)			1.3

- [0071]** Additional distillation and re-circulation through a third bag filter (196) filled with molecular sieve (7 kg) were started before the result of the IPC analysis was received. Since the result was approved the re-circulation was interrupted. Distillation was continued until a set volume (110 ± 10 L) in the reactor.
- 10 **[0072]** In another campaign of manufacturing **A** using the prior art method a reactor was charged with **4** (6.3 kg), n-octane (124 kg) and n-ethyl aniline (6.9 kg). After distilling off 88 kg solvent during 16 hours IPC showed that 2.7% of **4** remained.

References Cited

15

[0073]

US patent No. 6,875,869

US patent No. 5,912,349

20 US patent No. 6,133,285

Jansson et al., J. Org. Chem. 2006, 71, 1658-1667

Lu Jin et al., J. Phys. Chem. A 2008, 112, 4501-4510


Wennerberg et al., Org. Process. Res&Dev. 2007, 11, 674-680

Jönsson et al., J. Med. Chem. 2004, 47, 2075-2088

25

Patentkrav

1. Fremgangsmåde til fremstilling af en forbindelse med formen (I)



- 5 hvor


R1 er ethyl, R2 er ethyl, og R3 er H; eller

R1 er chlor, R2 er ethyl, og R3 er H; eller

R1 er methoxy, R2 er methyl, og R3 er trifluormethyl;

- 10 ved


(i) omsætning af en forbindelse med formen (II)



hvor

R1 er ethyl, chlor eller methoxy; og R4 er (C₁-C₄)-alkyl;

- 15 med en forbindelse med formen (III)



hvor

R2 er ethyl, og R3 er H; eller

R2 er methyl, og R3 er trifluormethyl;

i en tilbagesvalende blanding, som indeholder et alifatisk opløsningsmiddel eller en blanding af alifatiske opløsningsmidler, hvor det eller de alifatiske opløsningsmidler har et kogepunkt i intervallet 68-191°C;

5 (ii) kondensering af dampe fra den tilbagesvalende blanding;

(iii) blanding af de kondenserede dampe med et alkoholbindende middel valgt blandt molekylsigter eller med et alkoholbindende middel, som er en blanding af molekylsigter; og

10

(iv) returnering af de kondenserede dampe til reaktionsblandingen.

2. Fremgangsmåde ifølge krav 1, hvor det eller de alifatiske opløsningsmidler har et kogepunkt i intervallet 80-150°C.

15

3. Fremgangsmåde ifølge krav 1, hvor det eller de alifatiske opløsningsmidler har et kogepunkt i intervallet 95-130°C.

4. Fremgangsmåde ifølge krav 1, hvor det alkoholbindende middel er valgt
20 blandt 3A-, 4A-, 5A-, 13X-molekylsigter eller er en blanding af hvilke som helst af de nævnte molekylsigter.

5. Fremgangsmåde ifølge et hvilket som helst af kravene 1-4, hvor det eller de alifatiske opløsningsmidler er valgt blandt lineære eller forgrenede (C₆-C₁₀)-
25 alkaner og -cycloalkaner.

6. Fremgangsmåde ifølge et hvilket som helst af kravene 1-5, hvor det eller de alifatiske opløsningsmidler er valgt blandt lineære eller forgrenede (C₇-C₈)-
alkaner og -cycloalkaner.

30

7. Fremgangsmåde ifølge krav 6, hvor det eller de alifatiske opløsningsmidler er valgt blandt n-heptan, n-octan, methylcyclohexan, 2,2,4-trimethylpentan og cyclooctan.

8. Fremgangsmåde ifølge krav 7, hvor det eller de alifatiske opløsningsmidler er valgt blandt n-heptan og n-octan.
- 5 9. Fremgangsmåde ifølge et hvilket som helst af kravene 1-8, hvor R4 er valgt blandt methyl, ethyl, n-propyl, iso-propyl og n-butyl.
10. Fremgangsmåde ifølge krav 9, hvor R4 er valgt blandt methyl og ethyl.
- 10 11. Fremgangsmåde ifølge krav 1, hvor R1 er ethyl, R2 er ethyl, R3 er H, R4 er methyl eller ethyl; det alkoholbindende middel er en 3A-, 4A-, 5A- eller 13X-molekylsigte eller en blanding deraf; og det eller de alifatiske opløsningsmidler er valgt blandt n-heptan og n-octan.
- 15 12. Fremgangsmåde ifølge krav 1, hvor R1 er chlor, R2 er ethyl, R3 er H, R4 er methyl, det alkoholbindende middel er en 3A-, 4A-, 5A- eller 13X-molekylsigte, og det eller de alifatiske opløsningsmidler er valgt blandt n-heptan og n-octan.
- 20 13. Fremgangsmåde ifølge krav 1, hvor R1 er methoxy, R2 er methyl, R3 er trifluormethyl, R4 er methyl eller ethyl; det alkoholbindende middel er en 3A-, 4A, 5A- eller 13X-molekylsigte eller en blanding deraf; og det alifatiske opløsningsmiddel er n-octan.

1




Fig. 1




Fig. 2