




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 498/18, A61K 31/435 // (C07D 498/18, 311:00, 273:00, 221:00)	A1	(11) International Publication Number: WO 95/28406 (43) International Publication Date: 26 October 1995 (26.10.95)
(21) International Application Number: PCT/US95/04603 (22) International Filing Date: 14 April 1995 (14.04.95) (30) Priority Data: 08/229,261 18 April 1994 (18.04.94) US (71) Applicant: AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US). (72) Inventors: SKOTNICKI, Jerauld, Stanley; 4 Tyler Court, Allentown, NJ 08501 (US). LEONE, Christina, Louise; 99 Bayard Lane, Princeton, NJ 08540 (US). SCHIEHSER, Guy, Alan; 658 Bayberry Lane, Yardley, PA 19067 (US). (74) Agents: ALICE, Ronald, W.; American Home Products Corporation, Five Giralda Farms, Madison, NJ 07940-0874 (US) et al.	(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>	

(54) Title: RAPAMYCIN HYDROXYESTERS, PROCESS FOR THEIR PREPARATION AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

(57) Abstract

A compound of structure (I) wherein R^1 and R^2 are each, independently, hydrogen or $-\text{CO}(\text{CR}^3\text{R}^4)_b(\text{CR}^5\text{R}^6)_d\text{CR}^7\text{R}^8\text{R}^9$; R^3 and R^4 are each, independently, hydrogen, alkyl, alkenyle, alkynyl, trifluoromethyl, or $-\text{F}$; R^5 and R^6 are each, independently, hydrogen, alkyl, alkenyl, alkynyl, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, $-\text{F}$, or $-\text{CO}_2\text{R}^{11}$, or R^5 and R^6 may be taken together to form X or a cycloalkyl ring that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$; R^7 is hydrogen, alkyl, alkenyl, alkynyl, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, $-\text{F}$, or $-\text{CO}_2\text{R}^{11}$; R^8 and R^9 are each, independently, hydrogen, alkyl, alkenyl, alkynyl, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, $-\text{F}$, or $-\text{CO}_2\text{R}^{11}$, or R^8 and R^9 may be taken together to form X or a cycloalkyl ring that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$; R^{10} is hydrogen, alkyl, alkenyl, alkynyl, tri-(alkyl)silyl, tri-(alkyl)silylethyl, triphenylmethyl, benzyl, alkoxymethyl, tri-(alkyl)silylethoxymethyl, chloroethyl, or tetrahydropyranyl; R^{11} is hydrogen, alkyl, alkenyl, alkynyl, or phenylalkyl; X is 5-(2,2-dialkyl)[1,3]dioxanyl, 5-(2-spiro-cycloalkyl)[1,3]dioxanyl, 4-(2,2-dialkyl)[1,3]dioxanyl, 4-(2-spiro-cycloalkyl)[1,3]dioxanyl, 4-(2,2-dialkyl)[1,3]dioxalanyl, or 4-(2-spiro-cycloalkyl)[1,3]dioxalanyl; $b = 0-6$; $d = 0-6$; and $f = 0-6$ with the proviso that R^1 and R^2 are both not hydrogen and further provided that either R^1 or R^2 contains at least one $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, X, or $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$ substituted cycloalkyl group, or a pharmaceutically acceptable salt thereof which is useful as an immunosuppressive, antiinflammatory, antifungal, antiproliferative, and antitumor agent.



(I)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

RAPAMYCIN HYDROXYESTERS, PROCESS FOR THEIR PREPARATION AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

BACKGROUND OF THE INVENTION

5 This invention relates to hydroxyesters of rapamycin and a method for using them for inducing immunosuppression, and in the treatment of transplantation rejection, graft vs. host disease, autoimmune diseases, diseases of inflammation, adult T-cell leukemia/lymphoma, solid tumors, fungal infections, and hyperproliferative vascular disorders.

10

 Rapamycin is a macrocyclic triene antibiotic produced by Streptomyces hygroscopicus, which was found to have antifungal activity, particularly against Candida albicans, both in vitro and in vivo [C. Vezina et al., J. Antibiot. 28, 721 (1975); S.N. Sehgal et al., J. Antibiot. 28, 727 (1975); H. A. Baker et al., J. Antibiot. 15 31, 539 (1978); U.S. Patent 3,929,992; and U.S. Patent 3,993,749].

 Rapamycin alone (U.S. Patent 4,885,171) or in combination with picibanil (U.S. Patent 4,401,653) has been shown to have antitumor activity. R. Martel et al. [Can. J. Physiol. Pharmacol. 55, 48 (1977)] disclosed that rapamycin is effective in the experimental allergic encephalomyelitis model, a model for multiple sclerosis; in the 20 adjuvant arthritis model, a model for rheumatoid arthritis; and effectively inhibited the formation of IgE-like antibodies.

 The immunosuppressive effects of rapamycin have been disclosed in FASEB 3, 3411 (1989). Cyclosporin A and FK-506, other macrocyclic molecules, also have been shown to be effective as immunosuppressive agents, therefore useful in 25 preventing transplant rejection [FASEB 3, 3411 (1989); FASEB 3, 5256 (1989); R. Y. Calne et al., Lancet 1183 (1978); and U.S. Patent 5,100,899].

 Rapamycin has also been shown to be useful in preventing or treating systemic lupus erythematosus [U.S. Patent 5,078,999], pulmonary inflammation [U.S. Patent 5,080,899], insulin dependent diabetes mellitus [Fifth Int. Conf. Inflamm. Res. Assoc. 30 121 (Abstract), (1990)], smooth muscle cell proliferation and intimal thickening following vascular injury [Morris, R. J. Heart Lung Transplant 11 (pt. 2): 197 (1992)], adult T-cell leukemia/lymphoma [European Patent Application 525,960 A1], and ocular inflammation [European Patent Application 532,862 A1].


 Mono- and diacylated derivatives of rapamycin (esterified at the 28 and 43 35 positions) have been shown to be useful as antifungal agents (U.S. Patent 4,316,885) and used to make water soluble aminoacyl prodrugs of rapamycin (U.S. Patent

- 2 -

4,650,803). Recently, the numbering convention for rapamycin has been changed; therefore according to Chemical Abstracts nomenclature, the esters described above would be at the 31- and 42- positions.

5 DESCRIPTION OF THE INVENTION

This invention provides derivatives of rapamycin which are useful as immunosuppressive, antiinflammatory, antifungal, antiproliferative, and antitumor agents having the structure



10

wherein R^1 and R^2 are each, independently, hydrogen or $-\text{CO}(\text{CR}^3\text{R}^4)_b(\text{CR}^5\text{R}^6)_d\text{CR}^7\text{R}^8\text{R}^9$;

R^3 and R^4 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, trifluoromethyl, or -F;

R^5 and R^6 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^5 and R^6 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;

R^7 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$;

R^8 and R^9 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^8 and R^9 may be taken together to form X or a cycloalkyl ring of

25

- 3 -

3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(CR^3R^4)_fOR^{10}$;

5 R^{10} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxymethyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, or tetrahydropyranyl;

R^{11} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenylalkyl of 7-10 carbon atoms;

10 X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 5-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]-dioxanyl, 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxalanyl, or 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxalanyl;

15 b = 0-6;

d = 0-6; and

f = 0-6


with the proviso that R^1 and R^2 are both not hydrogen and further provided that either R^1 or R^2 contains at least one $-(CR^3R^4)_fOR^{10}$, X, or $-(CR^3R^4)_fOR^{10}$ substituted
20 cycloalkyl of 3-8 carbon atoms group, or a pharmaceutically acceptable salt thereof.

The pharmaceutically acceptable salts are those derived from such inorganic cations such as sodium, potassium, and the like; and organic bases such as: mono-, di-, and trialkyl amines of 1-6 carbon atoms, per alkyl group and mono-, di-, and
25 trihydroxyalkyl amines of 1-6 carbon atoms per alkyl group, and the like.

The terms alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, and alkynyl of 2-7 carbon atoms, include both straight chain as well as branched carbon chains. As the compounds of this invention can contain more than one $-(CR^3R^4)_fOR^{10}$ group,
30 R^3 , R^4 , f, and R^{10} can be the same or different. Similarly, when other generic substituent descriptions are repeated in the same structure, they can be the same or different.


For a compound in which R^1 contains R^8 and R^9 taken together to form X, where X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, the alkyl group of X
35 contains 1 carbon atom, and d = 0, R^1 would have the following structure.

- 4 -



Similarly, for a compound in which R^1 contains R^8 and R^9 taken together to form X, where X is 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, the cycloalkyl group of X contains 6 carbon atoms, and $d = 0$, R^1 would have the following structure.

5



For compounds containing X, preferred compounds include those in which the alkyl group of X, if present, is methyl and the cycloalkyl group of X, if present, is cyclohexyl.

10

When R^{10} is not hydrogen, alkyl, alkenyl, or alkynyl, it is intended that R^{10} is a group that can serve as an alcohol protecting group. Thus, these groups are intermediates of free hydroxylated compounds, as well as being biologically active in their own right. R^{10} covers tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxyethyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, and tetrahydropyranyl groups. Other alcohol protecting groups are known by one skilled in the art and are also considered part of this invention.

20

Of the compounds of this invention preferred members are those in which R^2 is hydrogen; those in which R^2 is hydrogen, $b = 0$, and $d = 0$; those in which R^2 is hydrogen, $b = 0$, $d = 0$, and R^8 and R^9 are each, independently hydrogen, alkyl, or $-(CR^3R^4)_fOR^{10}$, or are taken together to form X.

25

Compounds of this invention having the ester group $-CO(CR^3R^4)_b(CR^5R^6)_dCR^7R^8R^9$ at the 42- or 31,42-positions can be prepared by acylation of rapamycin using protected hydroxy and polyhydroxy acids, alkoxy or polyalkoxy carboxylic acids that have been activated, followed by removal of the


- 5 -

alcohol protecting groups, if so desired. Several procedures for carboxylate activation are known in the art, but the preferred methods utilize carbodiimides, mixed anhydrides, or acid chlorides. For example, an appropriately substituted carboxylic acid can be activated as a mixed anhydride, with an acylating group such as 2,4,6-trichlorobenzoyl chloride. Treatment of rapamycin with the mixed anhydride under mildly basic condition provides the desired compounds. Alternatively, the acylation reaction can be accomplished with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and dimethylaminopyridine. Mixtures of 42- and 31,42-esters can be separated by chromatography.

10

Accordingly, the invention also provides a process for preparing the rapamycin compounds of this invention. In particular this invention provides a process for preparing hydroxy esters of rapamycin including those of formula I as defined above which comprises:

- 15 a) acylating rapamycin or a functional derivative or analogue thereof with an acylating agent;
or
b) sequentially acylating rapamycin or a functional derivative or analogue thereof with two acylating agents;
20 said acylating agent(s) being selected from acids of formula



or a reactive derivative thereof wherein $\text{R}^3\text{-R}^9$, b and d are as defined above providing that free hydroxy groups are protected, if desired protecting the 42-position of rapamycin or functional derivative with an appropriate protecting group and after the reaction removing any protecting groups present as required.

25

The reaction may be carried out in the presence of a coupling reagent, such as a suitably substituted carbodiimide coupling reagent. The above-mentioned compounds of this invention can also be prepared by acylation using reactive derivatives of the acid of formula II such as an anhydride, a mixed anhydride, or an acid halide such as the chloride.

30

The 31-ester-42-hydroxy compounds of this invention can be prepared by protecting the 42-alcohol of rapamycin with a protecting group, such as with a tert-butyl dimethylsilyl group, followed by esterification of the 31-position by the

- 6 -

procedures described above. The preparation of rapamycin 42-silyl ethers is described in U.S. Patent B1 5,120,842, which is hereby incorporated by reference. Removal of the protecting group provides the 31-esterified compounds. In the case of the tert-butyl dimethylsilyl protecting group, deprotection can be accomplished under mildly acidic
5 conditions, such as acetic acid / water / THF. The deprotection procedure is described in Example 15 of U.S. Patent 5,118,678, which is hereby incorporated by reference.

Having the 31-position esterified and the 42-position deprotected, the 42-position can be esterified using a different acylating agent than was reacted with the 31-alcohol, to give compounds having different esters at the 31- and 42- positions.
10 Alternatively, the 42-esterified compounds, prepared as described above, can be reacted with a different acylating agent to provide compounds having different esters at the 31- and 42-positions.

This invention also covers analogous hydroxy esters of other rapamycins such
15 as, but not limited to, 29-demethoxyrapamycin, [U.S. Patent 4,375,464, 32-demethoxyrapamycin under C.A. nomenclature]; rapamycin derivatives in which the double bonds in the 1-, 3-, and/or 5-positions have been reduced [U.S. Patent 5,023,262]; 29-desmethylrapamycin [U.S. Patent 5,093,339, 32-desmethylrapamycin under C.A. nomenclature]; 7,29-bisdesmethylrapamycin [U.S. Patent 5,093,338,
20 7,32-desmethylrapamycin under C.A. nomenclature]; and 15-hydroxyrapamycin [U.S. Patent 5,102,876]. The disclosures in the above cited U.S. Patents are hereby incorporated by reference.

Immunosuppressive activity for representative compounds of this invention was
25 evaluated in an in vitro standard pharmacological test procedure to measure the inhibition of lymphocyte proliferation (LAF) and in two in vivo standard pharmacological test procedures. The pinch skin graft test procedure measures the immunosuppressive activity of the compound tested as well as the ability of the compound tested to inhibit or treat transplant rejection. The adjuvant arthritis standard
30 pharmacological test procedure, which measures the ability of the compound tested to inhibit immune mediated inflammation. The adjuvant arthritis test procedure is a standard pharmacological test procedure for rheumatoid arthritis. The procedures for these standard pharmacological test procedures are provided below.

35 The comitogen-induced thymocyte proliferation procedure (LAF) was used as an in vitro measure of the immunosuppressive effects of representative compounds.

- 7 -

Briefly, cells from the thymus of normal BALB/c mice are cultured for 72 hours with PHA and IL-1 and pulsed with tritiated thymidine during the last six hours. Cells are cultured with and without various concentrations of rapamycin, cyclosporin A, or test compound. Cells are harvested and incorporated radioactivity is determined. Inhibition of lymphoproliferation is assessed as percent change in counts per minute from non-drug treated controls. For each compound evaluated, rapamycin was also evaluated for the purpose of comparison. An IC₅₀ was obtained for each test compound as well as for rapamycin. When evaluated as a comparator for the representative compounds of this invention, rapamycin had an IC₅₀ ranging from 0.6 -1.5 nM. The results obtained are provided as an IC₅₀ and as the percent inhibition of T-cell proliferation at 0.1 μM. The results obtained for the representative compounds of this invention were also expressed as a ratio compared with rapamycin. A positive ratio indicates immunosuppressive activity. A ratio of greater than 1 indicates that the test compound inhibited thymocyte proliferation to a greater extent than rapamycin. Calculation of the ratio is shown below.

$$\frac{\text{IC}_{50} \text{ of Rapamycin}}{\text{IC}_{50} \text{ of Test Compound}}$$

20

Representative compounds of this invention were also evaluated in an *in vivo* test procedure designed to determine the survival time of pinch skin graft from male BALB/c donors transplanted to male C₃H(H-2K) recipients. The method is adapted from Billingham R.E. and Medawar P.B., J. Exp. Biol. 28:385-402, (1951). Briefly, a pinch skin graft from the donor was grafted on the dorsum of the recipient as a allograft, and an isograft was used as control in the same region. The recipients were treated with either varying concentrations of test compounds intraperitoneally or orally. Rapamycin was used as a test control. Untreated recipients serve as rejection control. The graft was monitored daily and observations were recorded until the graft became dry and formed a blackened scab. This was considered as the rejection day. The mean graft survival time (number of days ± S.D.) of the drug treatment group was compared with the control group. The following table shows the results that were obtained. Results are expressed as the mean survival time in days. Untreated (control) pinch skin grafts are usually rejected within 6-7 days. Compounds were tested using a dose of 4 mg/kg.

35

- 8 -

The adjuvant arthritis standard pharmacological test procedure measures the ability of test compounds to prevent immune mediated inflammation and inhibit or treat rheumatoid arthritis. The following briefly describes the test procedure used. A group of rats (male inbred Wistar Lewis rats) are pre-treated with the compound to be tested (1 h prior to antigen) and then injected with Freud's Complete Adjuvant (FCA) in the right hind paw to induce arthritis. The rats are then orally dosed on a Monday, Wednesday, Friday schedule from day 0-14 for a total of 7 doses. Both hind paws are measured on days 16, 23, and 30. The difference in paw volume (mL) from day 16 to day 0 is determined and a percent change from control is obtained. The left hind paw (uninjected paw) inflammation is caused by T-cell mediated inflammation and is recorded in the above table (% change from control). The right hind paw inflammation, on the other hand, is caused by nonspecific inflammation. Compounds were tested at a dose of 5 mg/kg. The results are expressed as the percent change in the uninjected paw at day 16 versus control; the more negative the percent change, the more potent the compound. Rapamycin provided between -70% and -90% change versus control, indicating that rapamycin treated rats had between 70-90% less immune induced inflammation than control rats.

The results obtained in these standard pharmacological test procedures are provided following the procedure for making the specific compounds that were tested.

The results of these standard pharmacological test procedures demonstrate immunosuppressive activity both *in vitro* and *in vivo* for the compounds of this invention. The results obtained in the LAF test procedure indicates suppression of T-cell proliferation, thereby demonstrating the immunosuppressive activity of the compounds of this invention. Further demonstration of the utility of the compounds of this invention as immunosuppressive agents was shown by the results obtained in the skin graft and adjuvant arthritis standard pharmacological test procedures. Additionally, the results obtained in the skin graft test procedure further demonstrates the ability of the compounds of this invention to treat or inhibit transplantation rejection. The results obtained in the adjuvant arthritis standard pharmacological test procedure further demonstrate the ability of the compounds of this invention to treat or inhibit rheumatoid arthritis.

Based on the results of these standard pharmacological test procedures, the compounds are useful in the treatment or inhibition of transplantation rejection such as kidney, heart, liver, lung, bone marrow, pancreas (islet cells), cornea, small bowel,

and skin allografts, and heart valve xenografts; in the treatment or inhibition of autoimmune diseases such as lupus, rheumatoid arthritis, diabetes mellitus, myasthenia gravis, and multiple sclerosis; and diseases of inflammation such as psoriasis, dermatitis, eczema, seborrhea, inflammatory bowel disease, pulmonary inflammation
5 (including asthma, chronic obstructive pulmonary disease, emphysema, acute respiratory distress syndrome, bronchitis, and the like), and eye uveitis.

Because of the activity profile obtained, the compounds of this invention also are considered to have antitumor, antifungal activities, and antiproliferative activities. The compounds of this invention therefore also useful in treating solid tumors, adult T-
10 cell leukemia/lymphoma, fungal infections, and hyperproliferative vascular diseases such as restenosis and atherosclerosis. When used for restenosis, it is preferred that the compounds of this invention are used to treat restenosis that occurs following an angioplasty procedure. When used for this purpose, the compounds of this invention can be administered prior to the procedure, during the procedure, subsequent to the
15 procedure, or any combination of the above.

When administered for the treatment or inhibition of the above disease states, the compounds of this invention can be administered to a mammal orally, parenterally, intranasally, intrabronchially, transdermally, topically, intravaginally, or rectally.

20 It is contemplated that when the compounds of this invention are used as an immunosuppressive or antiinflammatory agent, they can be administered in conjunction with one or more other immunoregulatory agents. Such other immunoregulatory agents include, but are not limited to azathioprine, corticosteroids, such as prednisone and methylprednisolone, cyclophosphamide, rapamycin, cyclosporin A, FK-506,
25 OKT-3, and ATG. By combining the compounds of this invention with such other drugs or agents for inducing immunosuppression or treating inflammatory conditions, the lesser amounts of each of the agents are required to achieve the desired effect. The basis for such combination therapy was established by Stepkowski whose results showed that the use of a combination of rapamycin and cyclosporin A at subtherapeutic
30 doses significantly prolonged heart allograft survival time. [Transplantation Proc. 23: 507 (1991)].

The compounds of this invention can be formulated neat or with a pharmaceutical carrier to a mammal in need thereof. The pharmaceutical carrier may be
35 solid or liquid. When formulated orally, it has been found that 0.01% Tween 80 in

- 10 -

PHOSAL PG-50 (phospholipid concentrate with 1,2-propylene glycol, A. Nattermann & Cie. GmbH) provides an acceptable oral formulation.

5 A solid carrier can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents; it can also be an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of
10 the active ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

Liquid carriers are used in preparing solutions, suspensions, emulsions,
15 syrups, elixirs and pressurized compositions. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fats. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening
20 agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (partially containing additives as above, e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g. glycols) and their derivatives, and oils (e.g. fractionated coconut oil and arachis oil). For parenteral administration, the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are useful
25 in sterile liquid form compositions for parenteral administration. The liquid carrier for pressurized compositions can be halogenated hydrocarbon or other pharmaceutically acceptable propellant.

30 Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. The compound can also be administered orally either in liquid or solid composition form.

35 The compounds of this invention may be administered rectally in the form of a conventional suppository. For administration by intranasal or intrabronchial inhalation or insufflation, the compounds of this invention may be formulated into an aqueous or

- 11 -

partially aqueous solution, which can then be utilized in the form of an aerosol. The compounds of this invention may also be administered transdermally through the use of a transdermal patch containing the active compound and a carrier that is inert to the active compound, is non toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier may take any number of forms such as creams and ointments, pastes, gels, and occlusive devices. The creams and ointments may be viscous liquid or semisolid emulsions of either the oil-in-water or water-in-oil type. Pastes comprised of absorptive powders dispersed in petroleum or hydrophilic petroleum containing the active ingredient may also be suitable. A variety of occlusive devices may be used to release the active ingredient into the blood stream such as a semipermeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix containing the active ingredient. Other occlusive devices are known in the literature.

In addition, the compounds of this invention may be employed as a solution, cream, or lotion by formulation with pharmaceutically acceptable vehicles containing 0.1 - 5 percent, preferably 2%, of active compound which may be administered to a fungally affected area.

The dosage requirements vary with the particular compositions employed, the route of administration, the severity of the symptoms presented and the particular subject being treated. Based on the results obtained in the standard pharmacological test procedures, projected daily dosages of active compound would be 0.1 $\mu\text{g}/\text{kg}$ - 100 mg/kg, preferably between 0.001 - 25 mg/kg, and more preferably between 0.01 - 5 mg/kg. Treatment will generally be initiated with small dosages less than the optimum dose of the compound. Thereafter the dosage is increased until the optimum effect under the circumstances is reached; precise dosages for oral, parenteral, nasal, or intrabronchial administration will be determined by the administering physician based on experience with the individual subject treated. Preferably, the pharmaceutical composition is in unit dosage form, e.g. as tablets or capsules. In such form, the composition is sub-divided in unit dose containing appropriate quantities of the active ingredient; the unit dosage forms can be packaged compositions, for example, packeted powders, vials, ampoules, prefilled syringes or sachets containing liquids. The unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.

The following examples illustrate the preparation and biological activities of representative compounds of this invention.

- 12 -

Example 1Rapamycin 42-ester with (tetrahydropyran-2-yloxy)acetic acid

2,4,6-Trichlorobenzoyl chloride (0.55 mL, 3.51 mmol) was added via syringe to a solution of the glycolic acid THP-ether (0.562 g, 3.51 mmol) and triethylamine (0.49 mL, 3.51 mmol) in 10 mL THF at 0 °C under nitrogen. The mixture was stirred for 4 h at room temperature, and a white precipitate formed. The white precipitate was removed by vacuum filtration and the filtrate was concentrated with a stream of nitrogen and warm water bath. The residue was dissolved in 10 mL benzene, then rapamycin (2.92 g, 3.19 mmol) and DMAP (0.429 g, 3.51 mmol) were added and the mixture was stirred overnight at room temperature. The mixture was diluted with EtOAc, washed with cold 1N HCl (aq), saturated NaHCO₃ (aq) and brine, dried over MgSO₄, filtered and concentrated to an oily yellow solid. Flash chromatography (2X with 65% EtOAc-hexane) afforded the title compound (1.114 g, 33%) as a white solid.

(-)FAB-MS *m/z* 1055.5 (M⁻), 590.3 (southern fragment), 463.2 (northern fragment).
¹H NMR (400 MHz, d-6 DMSO) δ 4.60 (m, 1 H, C(42)H), 4.66 (m, 1 H), 4.14 (s, 2 H), 3.73 (m, 1 H), 3.42 (m, 1 H).
¹³C NMR (100.6 MHz, d-6 DMSO) δ 169.2, 97.4, 63.5, 61.2, 29.7, 24.8, 18.8.

Example 2Rapamycin 42-ester with hydroxyacetic acid

p-Toluenesulfonic acid (10 mg) was added to a solution of the product of Example 1 (306 mg, 0.29 mmol) in 10 mL CH₃OH at 0 °C. The solution was stirred 2 h at room temperature, then quenched with saturated NaHCO₃ solution. The aqueous phase was extracted 3X with EtOAc and the combined organic phases were washed with brine, dried over MgSO₄, filtered and concentrated to a white solid. Purification by flash chromatography (2X with EtOAc) afforded the title compound (145 mg, 51%) as a white solid.

(-) FAB-MS *m/z* 971.3 (M⁻), 590 (southern fragment), 379.1 (northern fragment).
¹H NMR (400 MHz, d-6 DMSO) δ 4.60 (m, 1 H, C(42)H), 3.98 (s, 2 H).
¹³C NMR (100.6 MHz, d-6 DMSO) δ 172.1, 59.7.

Results obtained in standard pharmacological test procedures:

LAF IC₅₀: 1.80 nM

LAF ratio: 0.83

Percent change in adjuvant arthritis versus control: -88%

- 13 -

Example 3**Rapamycin 42-ester with 2,2-dimethyl-3-(tetrahydropyran-2-yloxy)propionic acid**

To a solution of the 2,2-dimethyl-3-hydroxypropionic acid THP-ether (0.319 g, 5 1.58 mmol) and triethylamine (0.22 mL, 1.58 mmol) in 5 mL dry THF at 0 °C under nitrogen was added 2, 4, 6-trichlorobenzoyl chloride (0.25 mL, 1.58 mmol) dropwise via syringe. The mixture was stirred 4.5 h at room temperature. The white precipitate was removed by vacuum filtration and the filtrate was concentrated with a stream of nitrogen and a warm water bath. The residue was dissolved in 5 mL benzene, then 10 rapamycin (1.31 g, 1.43 mmol) and DMAP (0.193 g, 1.58 mmol) were added. The mixture was stirred overnight at room temperature, diluted with EtOAc, washed with 1N HCl (aq), saturated NaHCO₃ (aq), H₂O and brine, dried over MgSO₄, filtered and concentrated to a yellow oily solid. Flash chromatography (1X with 60% EtOAc-hexane, 1X with 55% EtOAc-hexane) afforded the title compound (0.356 g, 23%), as a 15 white solid.

(-)-FAB-MS *m/z* 1097.7 (M⁻), 590.4 (southern fragment), 505.3 (northern fragment).

¹H NMR (400 MHz, d-6 DMSO) δ 4.55 (m, 1 H, C(42)H), 4.55 (m, 1 H), 3.69 (m, 1 H), 3.60 (m, 2 H), 3.42 (m, 1 H), 1.13 (s, 3 H), 1.11 (s, 3 H).

20 ¹³C NMR (100.6 MHz, d-6 DMSO) δ 175.0, 98.0, 73.8, 60.7, 42.6, 30.0, 24.9, 22.0, 21.6, 18.7.

Results obtained in standard pharmacological test procedures:

LAF IC₅₀: 7.10 nM

25 LAF ratio: 0.34

Example 4**Rapamycin 42-ester with 3-hydroxy-2,2-dimethylpropionic acid**

p-Toluenesulfonic acid (10 mg) was added to a solution of the product of 30 Example 3 (250 mg, 0.23 mmol) in 10 mL CH₃OH at 0 °C. The solution was stirred 2 hours at room temperature, then quenched with saturated NaHCO₃ solution. The aqueous phase was extracted 3X with EtOAc and the combined organic phases were washed with brine, dried over MgSO₄, filtered and concentrated to a white solid. Purification by flash chromatography (2X with 75% EtOAc-hexane) afforded the title 35 compound (103 mg, 45%) as a white solid.

- 14 -

(-) FAB-MS m/z 1013.3 (M^-), 590.2 (southern fragment), 421.1 (northern fragment).
 ^1H NMR (400 MHz, d_6 DMSO) δ 4.48 (m, 1 H, C(42)H), 3.39 (d, 2 H), 1.06 (s, 6 H).
 ^{13}C NMR (100.6 MHz, d_6 DMSO) δ 175.5, 68.0, 44.1, 21.7.

5

Results obtained in standard pharmacological test procedures:

LAF IC_{50} : 0.80 nM

LAF ratio: 1.25

Skin graft survival time: 10.7 ± 0.5 days

10

Examples 5 and 6

Rapamycin 42-ester with 2,2-dimethyl[1,3]dioxalane-4-carboxylic acid (Ex. 5)

Rapamycin 31,42-diester with 2,2-dimethyl[1,3]dioxalane-4-carboxylic acid (Ex. 6)

2,4,6-Trichlorobenzoyl chloride (0.56 mL, 3.61 mmol) was added via syringe
15 to a solution of the 2,3-dihydroxypropionic acid isopropylidene ketal (0.527 g, 3.61 mmol) and triethylamine (0.50 mL, 3.61 mmol) in 10 mL THF at 0 °C under nitrogen. The mixture was stirred 4 h at room temperature. The white precipitate was removed by vacuum filtration and the filtrate was concentrated with a stream of nitrogen and warm water bath. The residue was dissolved in 15 mL benzene and rapamycin (3.00
20 g, 3.28 mmol), then DMAP (0.441 g, 3.61 mmol) were added and the mixture was stirred overnight at room temperature. The mixture was diluted with EtOAc, washed with cold 1N HCl (aq), saturated NaHCO_3 (aq) and brine, dried over MgSO_4 , filtered and concentrated to a yellow foam. Flash chromatography on silica gel (gradient elution: 50-60-75-100% EtOAc-hexane, 4X with 65% EtOAc-hexane) afforded the title
25 compounds. The less polar 31,42-diester (0.415 g) eluted first and the more polar 42-monoester (0.601 g, 16%) eluted second, and were isolated as white solids.

Example 5

(-)FAB-MS m/z 1041.4 (M^-), 590.3 (southern fragment), 449.2 (northern fragment).
30 ^1H NMR (400 MHz, d_6 DMSO) δ 4.6 (m, 1 H, C(42)H), 4.6 (m, 1 H), 4.20 (dd, 1 H), 3.96 (m, 1 H), 1.36 (s, 3 H), 1.30 (s, 3 H).
 ^{13}C NMR (100.6 MHz, d_6 DMSO) δ 170.5, 110.2, 73.4, 66.6, 25.7, 25.4.

- 15 -

Example 6

(-)FAB-MS m/z 1169.6 (M^-).

1H NMR (400 MHz, d-6 DMSO) δ 5.3 (m, 1 H, C(31)H), 4.6 (m, 1 H, C(42)H), 4.6 (m, 2 H), 4.19 (t, 1 H), 4.13 (t, 1 H), 3.9 (m, 2 H), 1.36 (s, 3 H), 1.33 (s, 3 H),
5 1.30 (s, 3 H), 1.28 (s, 3 H).
 ^{13}C NMR (100.6 MHz, d-6 DMSO) δ 170.5, 169.2, 110.3, 110.2, 73.4, 66.6, 66.5,
25.8, 25.7, 25.4, 25.1.

Results obtained in standard pharmacological test procedures:

10 Example 5

LAF IC_{50} : 1.20 nM

LAF ratio: 0.74

Example 6

15 LAF IC_{50} : 1.30 nM

LAF ratio: 0.5

Example 7

Rapamycin 42-ester with 2,3-dihydroxypropionic acid

20 A solution of the product of Example 5 (351 mg, 0.34 mmol) in 10 mL THF and 10 mL 1N HCl was stirred at room temperature for 6 h. The mixture was diluted with EtOAc, washed with saturated $NaHCO_3$ solution and brine, dried over $MgSO_4$, filtered and concentrated to an oil. Flash chromatography (1X with EtOAc, 1X with 10% MeOH- CH_2Cl_2 , 1X with 5% MeOH-EtOAc) afforded the title compound (78 mg,
25 23%) as a white solid.

(-)FAB-MS m/z 1001.2 (M^-), 590.2 (southern fragment), 409.1 (northern fragment).

1H NMR (400 MHz, d-6 DMSO) δ 4.5 (m, 1 H, C(42)H), 3.60 (m, 1 H), 3.45 (m,
30 2 H).

Results obtained in standard pharmacological test procedures:

LAF IC_{50} : 1.4 nM

LAF ratio: 0.40

- 16 -

Example 8Rapamycin 42-ester with 2,2-dimethyl[1,3]dioxane-5-carboxylic acid

2,4,6-Trichlorobenzoyl chloride (0.98 mL, 6.27 mmol) was added via syringe to a solution of the 2-(hydroxymethyl)-3-hydroxypropionic acid isopropylidene ketal (1.000 g, 6.24 mmol) and triethylamine (0.90 mL, 6.46 mmol) in 20 mL THF at 0 °C under nitrogen. The mixture was stirred for 4 h at room temperature, and a white precipitate formed. The white precipitate was removed by vacuum filtration and the filtrate was concentrated with a stream of nitrogen and warm water bath. The residue was dissolved in 20 mL benzene, then rapamycin (5.70 g, 6.24 mmol) and DMAP (0.762 g, 6.24 mmol) were added and the mixture was stirred overnight at room temperature. The mixture was diluted with EtOAc, washed with H₂O and brine, dried over MgSO₄, filtered and concentrated to a yellow solid. Flash chromatography (75% EtOAc-hexane) afforded the title compound (4.17 g, 63%) as a white solid.

(-)-FAB-MS *m/z* 1055.8 (*M*⁻), 590.5 (southern fragment), 463.4 (northern fragment).
¹H NMR (400 MHz, d-6 DMSO) δ 4.55 (m, 1 H, C(42)H), 3.95 (m, 4 H), 1.30 (s, 6 H).
¹³C NMR (100.6 MHz, d-6 DMSO) δ 170.1, 97.4, 59.5, 24.8, 22.5.

Results obtained in standard pharmacological test procedures:
LAF IC₅₀: 0.76 nM
LAF ratio: 0.45

Example 9Rapamycin 42-ester with 3-hydroxy-2-hydroxymethylpropionic acid

A solution of the product of Example 8 (3.30 g, 3.12 mmol) in 50 mL THF and 25 mL 1N HCl was stirred 2 h at room temperature. The solution was diluted with saturated NaHCO₃ solution and extracted with EtOAc (3X). The combined organic phases were washed with saturated NaCl (aq), dried over MgSO₄, filtered and concentrated to a yellow foam. Purification by flash chromatography (1X with EtOAc; 2X with 5% EtOH-EtOAc) afforded the title compound (1.68 g, 53 %) as a white solid.

(-)-FAB-MS *m/z* 1015.5 (*M*⁻), 590.3 (southern fragment), 423.3 (northern fragment).
¹H NMR (400 MHz, d-6 DMSO) δ 4.6 (br s, 2 H), 4.55 (m, 1 H, C(42)H), 3.55 (m, 4 H), 2.57-2.53 (m, 1 H).
¹³C NMR (100.6 MHz, d-6 DMSO) δ 172.2, 59.3, 51.5.

- 17 -

Results obtained in standard pharmacological test procedures:

LAF IC₅₀: 0.84 nM

LAF ratio: 0.57

5

Example 10

Rapamycin 42-ester with 2,2,5-trimethyl[1,3]dioxane-5-carboxylic acid

To a solution of the 2,2-bis(hydroxymethyl)propionic acid isopropylidene ketal (1.041 g, 5.98 mmol) (prepared according to the procedure of Bruice, J. Am. Chem. Soc. 89: 3568 (1967)) and triethylamine (0.83 mL, 5.98 mmol) in 20 mL anhydrous THF at 0 °C under nitrogen was added 2, 4, 6-trichlorobenzoyl chloride (0.93 mL, 5.98 mmol) and the resultant white suspension was stirred 5 h at room temperature. The precipitate was removed by vacuum filtration, rinsing the flask and filter cake with an additional 10 mL dry THF. The filtrate was concentrated by rotary evaporation to a white solid. The residue was dissolved in 20 mL dry benzene, then rapamycin (5.47 g, 5.98 mmol) and DMAP (0.731 g, 5.98 mmol) were added. After stirring overnight at room temperature, the mixture was diluted with EtOAc, washed with H₂O and saturated NaCl (aq), dried over MgSO₄, filtered and evaporated to a yellow oil. Flash chromatography (5X with 60% EtOAc-hexane) afforded the title compound (2.2 g, 34 %) as a white solid.

15

20

(-)FAB-MS *m/z* 1069.5 (M⁻), 590.3 (southern fragment), 477.2 (northern fragment).

¹H NMR (400 MHz, d-6 DMSO) δ 4.57 (m, 1 H, C(42)H), 4.02 (d, 2 H), 3.60 (d, 2 H), 1.34 (s, 3 H), 1.24 (s, 3 H), 1.06 (s, 3 H).

¹³C NMR (100.6 MHz, d-6 DMSO) δ 173.2, 99.0, 65.0, 22.2, 18.1.

25 Results obtained in standard pharmacological test procedures:

LAF IC₅₀: 4.90 nM

LAF ratio: 0.41

Skin graft survival time: 11.0 ± 1.3 days

30

Example 11

Rapamycin 42-ester with 2,2-bis-(hydroxymethyl)propionic acid

A solution of the product of Example 10 (2.8 g, 2.65 mmol) in 50 mL THF and 25 mL 1N HCl was stirred at room temperature for 4 h. The mixture was diluted with water and extracted three times with EtOAc. The combined organic phases were washed with saturated NaHCO₃ solution, saturated NaCl solution, dried over MgSO₄,

35

- 18 -

filtered and evaporated to a yellow oily solid. Purification by flash chromatography (3X with EtOAc) afforded the title compound (1.6 g, 59 %).

(-)FAB-MS m/z 1029.6 (M^-), 590.4 (southern fragment), 437.3 (northern fragment).
 1H NMR (400 MHz, d-6 DMSO) δ 4.5 (m, 1 H, C(42)H), 3.45 (s, 4 H), 1.04 (s,
5 3 H).
 ^{13}C NMR (100.6 MHz, d-6 DMSO) δ 174.2, 63.7, 63.6, 49.9, 16.8.

Results obtained in standard pharmacological test procedures:

LAF IC_{50} : 0.80 and 1.80 nM

LAF ratio: 1.00 and 0.44

10 Skin graft survival time: 11.4 ± 1.5 and 12.0 ± 1.1 days
Percent change in adjuvant arthritis versus control: -88%

Example 12

15 Rapamycin 42-ester with 2,2-dimethyl-5-(2-trimethylsilylethoxymethyl)[1.3]-
dioxane-5-carboxylic acid

2,4,6-Trichlorobenzoyl chloride (0.14 mL, 0.86 mmol) was added via syringe to a solution of the 2,2-bis(hydroxymethyl)-2-(2-trimethylsilylethoxy)propionic acid isopropylidene ketal (0.250 g, 0.86 mmol) and triethylamine (0.12 mL, 0.86 mmol) in 2 mL THF at 0 °C under nitrogen. The mixture was stirred for 4 h at room
20 temperature, and a white precipitate formed. The white precipitate was removed by vacuum filtration and the filtrate was concentrated with a stream of nitrogen and warm water bath. The residue was dissolved in 2 mL benzene, then rapamycin (0.786 g, 0.86 mmol) and DMAP (0.105 g, 0.86 mmol) were added and the mixture was stirred overnight at room temperature. The mixture was diluted with EtOAc, washed with
25 H_2O and brine, dried over $MgSO_4$, filtered and concentrated to a yellow solid. Flash chromatography (gradient elution: 40-60-80-100% EtOAc-hexane) afforded the title compound (0.559 g, 54%) as a white solid.

(-)FAB-MS m/z 1185.2 (M^-), 590.1 (southern fragment), 593 (northern fragment).
 1H NMR (400 MHz, d-6 DMSO) δ 4.55 (m, 1 H, C(42)H), 3.73 (m, 4 H), 3.57 (s,
30 2 H), 3.43 (t, 2 H), 1.29 (s, 6 H), 0.79 (t, 2 H), -0.04 (s, 9 H).
 ^{13}C NMR (100.6 MHz, d-6 DMSO) δ 171.1, 97.7, 70.2, 68.1, 61.3, 46.0, 24.6, 22.1, 14.6, -1.3.

Results obtained in standard pharmacological test procedures:

LAF IC_{50} : 7.20 nM

35 LAF ratio: 0.05

Examples 13 and 14

Rapamycin 42-ester with 3-methyl-1,5-dioxaspiro[5.5]undecane 3-carboxylic acid (Ex. 13)

5 Rapamycin 31,42-diester with 3-methyl-1,5-dioxaspiro[5.5]undecane 3-carboxylic acid (Ex. 14)

2,4,6-Trichlorobenzoyl chloride (0.16 mL, 1.0 mmol) was added via syringe to a solution of the 2,3-dihydroxypropionic acid cyclohexylidene ketal (0.214 g, 1.0 mmol) and triethylamine (0.14 mL, 1.0 mmol) in 2.5 mL THF at 0 °C under nitrogen. The mixture was stirred 4 h at room temperature. The white precipitate was removed
10 by vacuum filtration and the filtrate was concentrated with a stream of nitrogen and warm water bath. The residue was dissolved in 3 mL benzene and rapamycin (0.457 g, 0.5 mmol), then DMAP (0.061 g, 0.5 mmol) were added and the mixture was stirred overnight at room temperature. The mixture was diluted with EtOAc, washed with cold 1N HCl (aq), saturated NaHCO₃ (aq) and brine, dried over MgSO₄, filtered
15 and concentrated to a yellow foam. Flash chromatography on silica gel (45-50% EtOAc-hexane) afforded the title compounds. The 31,42-diester (0.168 g, 26%) eluted first and the more polar 42-monoester (0.301 g, 52%) eluted second, and the products were isolated as white solids.

Example 13

20 (-)FAB-MS *m/z* 1109.5 (M⁻), 590.3 (southern fragment), 517.3 (northern fragment).
¹H NMR (400 MHz, d-6 DMSO) δ 4.55 (m, 1 H, C(42)H), 3.61 (t, 4 H), 1.04 (s, 3 H).
¹³C NMR (100.6 MHz, d-6 DMSO) δ 173.3, 97.2, 64.2.

Example 14

25 (-)FAB-MS *m/z* 1305.6 (M⁻).
¹H NMR (400 MHz, d-6 DMSO) δ 5.25 (m, 1 H, C(31)H), 4.55 (m, 1 H, C(42)H), 3.64-3.54 (m, 8 H), 1.05 (s, 3 H), 0.97 (s, 3 H).
¹³C NMR (100.6 MHz, d-6 DMSO) δ 173.2, 172.1, 97.3, 97.2, 64.3, 64.2, 63.9.

Results obtained in standard pharmacological test procedures:

30 Example 13

LAF IC₅₀: 0.6 nM

LAF ratio: 2.00

Example 14


LAF: inhibited T-cell proliferation by 43% at 0.1 μM

- 20 -

CLAIMS

What is claimed is:

1. A compound of the structure



- 5 wherein R^1 and R^2 are each, independently, hydrogen or $-\text{CO}(\text{CR}^3\text{R}^4)_b(\text{CR}^5\text{R}^6)_d\text{CR}^7\text{R}^8\text{R}^9$;
 R^3 and R^4 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, trifluoromethyl, or -F;
 R^5 and R^6 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of
 10 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^5 and R^6 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;
 R^7 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-
 15 7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$;
 R^8 and R^9 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^8 and R^9 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with
 20 $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;
 R^{10} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxyethyl of 2-7 carbon atoms,

- 21 -

tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, or tetrahydropyranyl;

R¹¹ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenylalkyl of 7-10 carbon atoms;

5 X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 5-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]-dioxanyl, 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxalanyl, or 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxalanyl;

10 b = 0-6;

d = 0-6; and

f = 0-6

with the proviso that R¹ and R² are both not hydrogen and further provided that either R¹ or R² contains at least one -(CR³R⁴)_fOR¹⁰, X, or -(CR³R⁴)_fOR¹⁰ substituted
15 cycloalkyl of 3-8 carbon atoms group, or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, wherein R² is hydrogen or a pharmaceutically acceptable salt thereof.

20 3. The compound of claim 2, wherein b = 0 and d = 0 or a pharmaceutically acceptable salt thereof.

4. The compound of claim 3, wherein R⁸ and R⁹ are each, independently hydrogen, alkyl, or -(CR³R⁴)_fOR¹⁰, or are taken together to form X or a
25 pharmaceutically acceptable salt thereof.

5. The compound of claim 1 which is rapamycin 42-ester with (tetrahydropyran-2-yloxy)acetic acid or a pharmaceutically acceptable salt thereof.

30 6. The compound of claim 1 which is rapamycin 42-ester with hydroxyacetic acid or a pharmaceutically acceptable salt thereof.

7. The compound of claim 1 which is rapamycin 42-ester with 2,2-dimethyl-3-(tetrahydropyran-2-yloxy)propionic acid or a pharmaceutically acceptable salt thereof.


35

- 22 -

8. The compound of claim 1 which is rapamycin 42-ester with 3-hydroxy-2,2-dimethylpropionic acid or a pharmaceutically acceptable salt thereof.
9. The compound of claim 1 which is rapamycin 42-ester with 2,2-dimethyl[1,3]dioxalane-4-carboxylic acid or a pharmaceutically acceptable salt thereof.
10. The compound of claim 1 which is rapamycin 31,42-diester with 2,2-dimethyl[1,3]dioxalane-4-carboxylic acid or a pharmaceutically acceptable salt thereof.
- 10 11. The compound of claim 1 which is rapamycin 42-ester with 2,3-dihydroxypropionic acid or a pharmaceutically acceptable salt thereof.
12. The compound of claim 1 which is rapamycin 42-ester with 2,2-dimethyl[1,3]dioxane-5-carboxylic acid or a pharmaceutically acceptable salt thereof.
- 15 13. The compound of claim 1 which is rapamycin 42-ester with 3-hydroxy-2-hydroxymethylpropionic acid or a pharmaceutically acceptable salt thereof.
14. The compound of claim 1 which is rapamycin 42-ester with 2,2,5-trimethyl[1,3]dioxane-5-carboxylic acid or a pharmaceutically acceptable salt thereof.
- 20 15. The compound of claim 1 which is rapamycin 42-ester with 2,2-bis-(hydroxymethyl)propionic acid or a pharmaceutically acceptable salt thereof.
- 25 16. The compound of claim 1 which is rapamycin 42-ester with 2,2-dimethyl-5-(2-trimethylsilanyloxyethyl)[1,3]-dioxane-5-carboxylic acid or a pharmaceutically acceptable salt thereof.
17. The compound of claim 1 which is rapamycin 42-ester with 3-methyl-1,5-dioxaspiro[5.5]undecane 3-carboxylic acid or a pharmaceutically acceptable salt thereof.
- 30 18. The compound of claim 1 which is rapamycin 31,42-diester with 3-methyl-1,5-dioxaspiro[5.5]undecane 3-carboxylic acid or a pharmaceutically acceptable salt thereof.
- 35

- 23 -

19. A method of treating transplantation rejection or graft vs. host disease in a mammal in need thereof, which comprises administering to said mammal an antirejection effective amount of a compound of the structure



- 5 wherein R^1 and R^2 are each, independently, hydrogen or $-\text{CO}(\text{CR}^3\text{R}^4)_b(\text{CR}^5\text{R}^6)_d\text{CR}^7\text{R}^8\text{R}^9$;
- R^3 and R^4 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, trifluoromethyl, or -F;
- 10 R^5 and R^6 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^5 and R^6 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;
- 15 R^7 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$;
- R^8 and R^9 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^8 and R^9 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;
- 20 R^{10} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxymethyl of 2-7 carbon atoms,

- 24 -

tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, or tetrahydropyranyl;

R¹¹ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenylalkyl of 7-10 carbon atoms;

5 X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 5-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]-dioxanyl, 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxalanyl, or 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxalanyl;

10 b = 0-6;


d = 0-6; and

f = 0-6

with the proviso that R¹ and R² are both not hydrogen and further provided that either R¹ or R² contains at least one -(CR³R⁴)_fOR¹⁰, X, or -(CR³R⁴)_fOR¹⁰ substituted

15 cycloalkyl of 3-8 carbon atoms group, or a pharmaceutically acceptable salt thereof.

20. A method of treating a fungal infection in a mammal in need thereof, which comprises administering to said mammal an antifungal effective amount of a compound of the structure



20

wherein R¹ and R² are each, independently, hydrogen or -CO(CR³R⁴)_b(CR⁵R⁶)_dCR⁷R⁸R⁹;


R³ and R⁴ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, trifluoromethyl, or -F;

- 25 -

- 5 R⁵ and R⁶ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(CR^3R^4)_fOR^{10}$, $-CF_3$, $-F$, or $-CO_2R^{11}$, or R⁵ and R⁶ may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(CR^3R^4)_fOR^{10}$;
- 10 R⁷ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(CR^3R^4)_fOR^{10}$, $-CF_3$, $-F$, or $-CO_2R^{11}$;
- R⁸ and R⁹ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(CR^3R^4)_fOR^{10}$, $-CF_3$, $-F$, or $-CO_2R^{11}$, or R⁸ and R⁹ may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(CR^3R^4)_fOR^{10}$;
- 15 R¹⁰ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxymethyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, or tetrahydropyranyl;
- R¹¹ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenylalkyl of 7-10 carbon atoms;
- 20 X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 5-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxalanyl, or 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxalanyl;
- 25 b = 0-6;
d = 0-6; and
f = 0-6
- with the proviso that R¹ and R² are both not hydrogen and further provided that either R¹ or R² contains at least one $-(CR^3R^4)_fOR^{10}$, X, or $-(CR^3R^4)_fOR^{10}$ substituted
- 30 cycloalkyl of 3-8 carbon atoms group, or a pharmaceutically acceptable salt thereof.

- 26 -

21. A method of treating rheumatoid arthritis in a mammal in need thereof, which comprises administering to said mammal an antiarthritis effective amount of a compound of the structure



- 5 wherein R^1 and R^2 are each, independently, hydrogen or $-\text{CO}(\text{CR}^3\text{R}^4)_b(\text{CR}^5\text{R}^6)_d\text{CR}^7\text{R}^8\text{R}^9$;
- R^3 and R^4 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, trifluoromethyl, or -F;
- R^5 and R^6 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^5 and R^6 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;
- R^7 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$;
- R^8 and R^9 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^8 and R^9 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;
- R^{10} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxyethyl of 2-7 carbon atoms,

- 27 -

tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, or tetrahydropyranyl;

R^{11} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenylalkyl of 7-10 carbon atoms;

5 X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 5-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]-dioxanyl, 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxalanyl, or 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxalanyl;

10 b = 0-6;


d = 0-6; and

f = 0-6

with the proviso that R^1 and R^2 are both not hydrogen and further provided that either R^1 or R^2 contains at least one $-(CR^3R^4)_fOR^{10}$, X, or $-(CR^3R^4)_fOR^{10}$ substituted

15 cycloalkyl of 3-8 carbon atoms group, or a pharmaceutically acceptable salt thereof.

22. A method of treating restenosis in a mammal in need thereof, which comprises administering to said mammal an antiproliferative effective amount of a compound of the structure



20

wherein R^1 and R^2 are each, independently, hydrogen or $-(CO(CR^3R^4)_b(CR^5R^6)_dCR^7R^8R^9)$;

R^3 and R^4 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, trifluoromethyl, or -F;

- 28 -

5 R⁵ and R⁶ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, -(CR³R⁴)_fOR¹⁰, -CF₃, -F, or -CO₂R¹¹, or R⁵ and R⁶ may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with -(CR³R⁴)_fOR¹⁰;

R⁷ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, -(CR³R⁴)_fOR¹⁰, -CF₃, -F, or -CO₂R¹¹;

10 R⁸ and R⁹ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, -(CR³R⁴)_fOR¹⁰, -CF₃, -F, or -CO₂R¹¹, or R⁸ and R⁹ may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with -(CR³R⁴)_fOR¹⁰;

15 R¹⁰ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxymethyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, or tetrahydropyranyl;

R¹¹ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenylalkyl of 7-10 carbon atoms;

20 X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 5-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxalanyl, or 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxalanyl;

25 b = 0-6;


d = 0-6; and

f = 0-6

30 with the proviso that R¹ and R² are both not hydrogen and further provided that either R¹ or R² contains at least one -(CR³R⁴)_fOR¹⁰, X, or -(CR³R⁴)_fOR¹⁰ substituted cycloalkyl of 3-8 carbon atoms group, or a pharmaceutically acceptable salt thereof.

- 29 -

23. A method of treating pulmonary inflammation in a mammal in need thereof, which comprises administering to said mammal an antiinflammatory effective amount of a compound of the structure



- 5 wherein R^1 and R^2 are each, independently, hydrogen or $-\text{CO}(\text{CR}^3\text{R}^4)_b(\text{CR}^5\text{R}^6)_d\text{CR}^7\text{R}^8\text{R}^9$;
- R^3 and R^4 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, trifluoromethyl, or -F;
- 10 R^5 and R^6 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^5 and R^6 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;
- 15 R^7 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$;
- R^8 and R^9 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, -F, or $-\text{CO}_2\text{R}^{11}$, or R^8 and R^9 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;
- 20 R^{10} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxyethyl of 2-7 carbon atoms,

- 30 -

tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, or tetrahydropyranyl;

R¹¹ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenylalkyl of 7-10 carbon atoms;

5 X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 5-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]-dioxanyl, 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxalanyl, or 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxalanyl;

10 b = 0-6;


d = 0-6; and

f = 0-6

with the proviso that R¹ and R² are both not hydrogen and further provided that either R¹ or R² contains at least one -(CR³R⁴)_fOR¹⁰, X, or -(CR³R⁴)_fOR¹⁰ substituted

15 cycloalkyl of 3-8 carbon atoms group, or a pharmaceutically acceptable salt thereof.

24. A pharmaceutical composition which comprises a compound of the structure



20 wherein R¹ and R² are each, independently, hydrogen or -CO(CR³R⁴)_b(CR⁵R⁶)_dCR⁷R⁸R⁹;

R³ and R⁴ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, trifluoromethyl, or -F;

R⁵ and R⁶ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, -(CR³R⁴)_fOR¹⁰, -CF₃, -F, or

- 31 -

$-\text{CO}_2\text{R}^{11}$, or R^5 and R^6 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;

5 R^7 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, $-\text{F}$, or $-\text{CO}_2\text{R}^{11}$;

R^8 and R^9 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, $-\text{CF}_3$, $-\text{F}$, or $-\text{CO}_2\text{R}^{11}$, or R^8 and R^9 may be taken together to form X or a cycloalkyl ring of 3-8 carbon atoms that is optionally mono-, di-, or tri-substituted with $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$;

10 R^{10} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silyl, tri-(alkyl of 1-6 carbon atoms)silylethyl, triphenylmethyl, benzyl, alkoxymethyl of 2-7 carbon atoms, tri-(alkyl of 1-6 carbon atoms)silylethoxymethyl, chloroethyl, or tetrahydropyranyl;

15 R^{11} is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-7 carbon atoms, alkynyl of 2-7 carbon atoms, or phenylalkyl of 7-10 carbon atoms;

20 X is 5-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 5-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxanyl, 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxanyl, 4-(2,2-di-(alkyl of 1-6 carbon atoms))[1,3]dioxalanyl, or 4-(2-spiro(cycloalkyl of 3-8 carbon atoms))[1,3]dioxalanyl;

b = 0-6;

d = 0-6; and

25 f = 0-6

with the proviso that R^1 and R^2 are both not hydrogen and further provided that either R^1 or R^2 contains at least one $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$, X, or $-(\text{CR}^3\text{R}^4)_f\text{OR}^{10}$ substituted cycloalkyl of 3-8 carbon atoms group, or a pharmaceutically acceptable salt thereof, and a pharmaceutical carrier.

30


25. A process for preparing hydroxy esters of rapamycin including those of formula I as defined in Claim 1 which comprises:

a) acylating rapamycin or a functional derivative or analogue thereof with an acylating agent;

35 or

- 32 -

b) sequentially acylating rapamycin or a functional derivative or analogue thereof with two acylating agents; said acylating agent(s) being selected from acids of formula



- 5 or a reactive derivative thereof wherein R³-R⁹, b and d are as defined in Claim 1 providing that free hydroxy groups are protected, if desired protecting the 42-position of rapamycin or functional derivative with an appropriate protecting group and after the reaction removing any protecting groups present as required.

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/US 95/04603

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D498/18 A61K31/435 //(C07D498/18,311:00,273:00,221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,4 650 803 (V. J. STELLA ET AL) 17 March 1987 cited in the application see claims 1,5 ---	1,24
A	WO,A,92 05179 (AMERICAN HOME PRODUCTS CORPORATION) 2 April 1992 see claims 1,35,36 ---	1,24
A	US,A,5 233 036 (P. F. HUGHES ET AL) 3 August 1993 see column 1 - column 2; claim 1 --- -/--	1,24

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

26 June 1995

Date of mailing of the international search report

29 . 06 . 95

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Voyiazoglou, D

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 95/04603

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TETRAHEDRON LETTERS., vol.35, no.7, 14 February 1994, OXFORD GB pages 1019 - 1022 M. ADAMCZYK ET AL 'Lipase mediated hydrolysis of rapamycin 42-hemisuccinate benzyl and methyl esters' see page 1020</p> <p style="text-align: center;">-----</p>	1

1

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: 1 Application No PCT/US 95/04603
--

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4650803	17-03-87	AU-B- 583439	27-04-89
		AU-A- 6608086	11-06-87
		CA-A- 1273920	11-09-90
		CA-A- 1312076	29-12-92
		DE-A- 3684574	30-04-92
		DK-B- 169409	24-10-94
		EP-A, B 0227355	01-07-87
		EP-A- 0429436	29-05-91
		GB-A, B 2183647	10-06-87
		JP-B- 6070066	07-09-94
		JP-A- 62215592	22-09-87
		JP-A- 6263765	20-09-94
		JP-B- 7047593	24-05-95
		KR-B- 9404072	11-05-94
		KR-B- 9404073	11-05-94
WO-A-9205179	02-04-92	AU-B- 653175	22-09-94
		AU-A- 8659991	15-04-92
		CA-A- 2051781	20-03-92
		EP-A- 0549727	07-07-93
		HU-A- 65763	28-07-94
		JP-T- 6501012	27-01-94
		US-A- 5358944	25-10-94
		US-A- 5378696	03-01-95
		US-A- 5221670	22-06-93
		CA-A- 2051782	29-03-92
		US-A- 5130307	14-07-92
US-A-5233036	03-08-93	NONE	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 95/04603

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 19-23 are directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.