

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2005295774 B2**

- (54) Title
Isomeric mixtures of dinitro-octylphenyl esters and synergistic fungicidal mixtures therefrom
- (51) International Patent Classification(s)
A01N 37/06 (2006.01) **A01N 47/06** (2006.01)
- (21) Application No: **2005295774** (22) Date of Filing: **2005.10.14**
- (87) WIPO No: **WO06/044460**
- (30) Priority Data
- | (31) Number | (32) Date | (33) Country |
|-------------------|-------------------|--------------|
| 60/695,232 | 2005.06.29 | US |
| 60/624,704 | 2004.11.03 | US |
| 60/618,919 | 2004.10.14 | US |
- (43) Publication Date: **2006.04.27**
- (44) Accepted Journal Date: **2012.03.08**
- (71) Applicant(s)
Dow AgroSciences LLC
- (72) Inventor(s)
Green, Elizabeth Ann;Sivieri, Franco;Davies, John;Ehr, Robert Joseph;Mathieson, John Todd;Distler, Bernhard;Billington, Richard
- (74) Agent / Attorney
Freehills Patent & Trade Mark Attorneys, Level 43 101 Collins Street, Melbourne, VIC, 3000
- (56) Related Art
US 3715384
FAO AND WHO: "1969 EVALUATIONS OF SOME PESTICIDE RESIDUES IN FOOD" IPCS INCHEM WO 2003/103393

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 April 2006 (27.04.2006)

PCT

(10) International Publication Number
WO 2006/044460 A1

(51) International Patent Classification:
A01N 37/06 (2006.01) *A01N 47/06* (2006.01)

(74) Agents: **ZETTLER, Lynn, M.** et al.; Dow AgroSciences LLC, 9330 Zionsville Road, Indianapolis, IN 46268 (US).

(21) International Application Number:
PCT/US2005/036679

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 14 October 2005 (14.10.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/618,919 14 October 2004 (14.10.2004) US
60/624,704 3 November 2004 (03.11.2004) US
60/695,232 29 June 2005 (29.06.2005) US

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **DOW AGROSCIENCES LLC** [US/US]; 9330 Zionsville Road, Indianapolis, IN 46268-1054 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BILLINGTON, Richard** [GB/GB]; Mokundange, East Challow, Wanrage, OX 129RW (GB). **GREEN, Elizabeth, Ann** [GB/GB]; Cander, 43A New Road, East Hagbourne, OX11 9JX (GB). **MATHIESON, John, Todd** [US/US]; 6527 Donnelly Drive, Brownsburg, IN 46112 (US). **SIVIERI, Franco** [IT/IT]; Via Matteotti 12, I-24047 Treviglio (IT). **DAVIES, John** [GB/US]; 2462 Quail Run, Midland, MI 48642 (US). **DISTLER, Bernhard** [DE/DE]; Stuerzer Str. 8A, 83043 Bad Aibling (DE). **EHR, Robert, Joseph** [US/US]; 8693 Highwoods Lane, Indianapolis, IN 46278 (US).

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ISOMERIC MIXTURES OF DINITRO-OCTYLPHENYL ESTERS AND SYNERGISTIC FUNGICIDAL MIXTURES THEREFROM

(57) Abstract: The present invention relates to an isomeric composition comprising isomers of dinitro-octylphenyl esters, wherein the 2,6-dinitro-4-(1-propylpentyl)phenyl ester isomer is present in an amount of less than 0.1 weight percent, based on the total weight of the isomeric composition and to synergistic fungicidal mixtures therefrom.



WO 2006/044460 A1

ISOMERIC MIXTURES OF DINITRO-OCTYLPHENYL ESTERS AND SYNERGISTIC FUNGICIDAL MIXTURES THEREFROM

The present invention relates to isomeric compositions of dinitro-octylphenyl
5 esters and synergistic fungicidal compositions.

Compositions comprising 2,4-dinitro-6-octylphenyl crotonates and 2,6-dinitro-4-octylphenyl crotonates have been used as fungicides for control of powdery mildew, or certain species of plant parasitic mites. Such compositions, also known as dinocap, are the active ingredients used in Karathane™ Fungicide/Miticide. Dinocap is a mixture of
10 six isomers, each having a cis and trans component, and contains as its active ingredients a mixture of 2,4- and 2,6-dinitro-octylphenyl crotonates in an approximate 2:1 ratio, wherein 'octyl' refers to a mixture of 1-methylheptyl, 1-ethylhexyl, and 1-propylpentyl isomers. Various studies have been published relating to 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate, including a report on the dermal absorption and
15 mammalian developmental effects of dinocap. "Dinocap Dermal Absorption in Female Rabbits and Rhesus Monkeys", Biological Monitoring for Pesticide Exposure, Chapter 11, pages 137-151, December 4, 1987, discloses a study related to absorption data for 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate isomer at 97 percent purity. Another study, "Developmental Toxicity of Dinocap in the Mouse is Not Due to Two Isomers of
20 the Major Active Ingredients", Teratogenesis, Carcinogenesis, and Mutagenesis 7:341-346(1987) reports the use of Dinocap as a fungicide and reports toxicity data for 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate isomer at 95 percent purity.

Dinocap is known to form synergistic mixtures with other fungicides, including mixtures with benzophenones as taught in WO 02/067679 and U.S. 6,346,535; with
25 phenylbenzylethers and/or carbamates as taught in U.S. 6,528,536; with amides as taught in U.S. 6,515,000; with benzamidoximes as taught in WO 2004091298; with 1-((N-(2,3-dichloro-4-hydroxyphenyl)amino)-carbonyl)-1-methyl cyclohexanes as taught in U.S. 6,207,691; with chlorothalonil and ditalimfos as taught in FR 2445696; with N-(p-

fluorophenyl)-2,3-dichloromaleimide as taught in GB 2003032; with
tetrachloroisophthalonitrile as taught in U.S. 3,456,055; with phenylbenzylethers or
carbamates as taught in U.S. 6,489,360; with aminomethyl heterocycles as taught in U.S.
5,569,656; with fused pyrimidinones as taught in WO2003103393; and the like, wherein
5 all of the U.S. patents recited above are incorporated herein by reference.

Dinocap is a mixture of isomers with varying levels of toxicological effects, e.g.
teratogenicity in mice and retinopathy in dogs. Thus, there continues to be a need to
produce similar fungicides having more favorable toxicological properties, while
maintaining or increasing fungicidal efficacy and in developing synergistic fungicidal
10 mixtures therefrom.

The present invention relates to an isomeric composition comprising isomers of
dinitro-octylphenyl esters, wherein the 2,6-dinitro-4-(1-propylpentyl)phenyl ester isomer
is present in an amount of less than 0.1 weight percent, based on the total weight of the
isomeric composition; and to synergistic fungicidal mixtures therefrom.

15 It has been surprisingly discovered that the isomeric composition of the present
invention containing less than 0.1 weight percent 2,6-dinitro-4-(1-propylpentyl)phenyl
ester isomer has very favorable toxicological properties while maintaining fungicidal
efficacy.

Typically, a dinitro-octylphenyl ester exists as a mixture of isomers which include
20 forms of the following isomers:

2,4-dinitro-6-(1-methylheptyl)phenyl ester,
2,4-dinitro-6-(1-ethylhexyl)phenyl ester,
2,4-dinitro-6-(1-propylpentyl)phenyl ester,
2,6-dinitro-4-(1-methylheptyl)phenyl ester,
25 2,6-dinitro-4-(1-ethylhexyl)phenyl ester, and
2,6-dinitro-4-(1-propylpentyl)phenyl ester;

wherein the ester is defined as any ester functional group which is capable of hydrolyzing
under environmental conditions to form the corresponding phenol. Environmental

conditions include those conditions which occur naturally in the environment of agricultural crops or within the crops themselves, including the presence of moisture.

It has been surprisingly discovered that an isomeric composition comprising less than 0.1 weight percent of 2,6-dinitro-4-(1-propylpentyl)phenyl ester, based on the total weight of the isomeric composition, has no teratogenic or retinopathic effects, while
5 maintaining fungicidal efficacy.

For example, dinocap is an isomeric composition which refers to a mixture of isomers of dinitro-octylphenyl crotonates which include both cis and trans forms of the following isomers:

- 10 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate,
- 2,4-dinitro-6-(1-ethylhexyl)phenyl crotonate,
- 2,4-dinitro-6-(1-propylpentyl)phenyl crotonate,
- 2,6-dinitro-4-(1-methylheptyl)phenyl crotonate,
- 2,6-dinitro-4-(1-ethylhexyl)phenyl crotonate, and
- 15 2,6-dinitro-4-(1-propylpentyl)phenyl crotonate.

It is well known in the art that the crotonate ester, as in dinocap, hydrolyzes with exposure to the plant or environment to the corresponding dinitro-octylphenol and that the phenol is the active ingredient having fungicidal effects. Therefore, any ester or analog of these compounds in which the ester is derivatized to form a related substituent
20 that can be transformed within plants or the environment to the phenol, possess essentially the same fungicidal effect and are within the scope of the present invention. Preferably, the isomeric composition of the present invention is a mixture of dinitro-octylphenyl crotonates.

In general, the isomeric composition of the present invention can comprise a
25 dinitro-octylphenyl ester in any combination and amounts of isomers, as long as the amount of 2,6-dinitro-4-(1-propylpentyl)phenyl ester is below 0.1 weight percent based on the total weight of the isomeric composition and as long as the improved toxicology and fungicidal efficacy is maintained. In other words, the isomeric composition can

comprise any number of dinitro-octylphenyl ester isomers, each isomer in any amount, so long as the amount of 2,6-dinitro-4-(1-propylpentyl)phenyl ester is below 0.1 weight percent based on the total weight of the isomeric composition. Generally, the isomeric composition will comprise:

- 5 from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-methylheptyl)phenyl ester,
- from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-ethylhexyl)phenyl ester,
- from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-propylpentyl)phenyl ester,
- from 0.1 to 99.9 weight percent 2,6-dinitro-4-(1-methylheptyl)phenyl ester,
- from 0.1 to 99.9 weight percent 2,6-dinitro-4-(1-ethylhexyl)phenyl ester, and
- 10 less than 0.1 weight percent 2,6-dinitro-4-(1-propylpentyl)phenyl ester.

In one embodiment, the isomeric composition of the present invention comprises at least 20, generally at least 30, more generally at least 40, typically at least 50, more typically at least 60, preferably at least 70, more preferably at least 80, even more
15 preferably at least 90 and most preferably at least 95 weight percent 2,4-dinitro-6-(1-methylheptyl)phenyl ester, based on the total weight of all of the isomers which are present in the composition.

In another embodiment, the isomeric composition of the present invention comprises from 1, generally from 1.5, typically from 2, preferably from 2.5, more
20 preferably from 3 and most preferably from 5 to 45, generally to 40, typically to 35, preferably to 30, more preferably to 27 and most preferably to 25 weight percent 2,4-dinitro-6-(1-ethylhexyl)phenyl ester, based on the total weight of all of the isomers which are present in the composition.

In another embodiment, the isomeric composition of the present invention
25 comprises from 0.1, generally from 0.5, typically from 1, preferably from 1.5, more preferably from 2.0 and most preferably from 2.5 to 45, generally to 40, typically to 35, preferably to 30, more preferably to 27 and most preferably to 25 weight percent 2,4-

dinitro-6-(1-propylpentyl)phenyl ester, based on the total weight of all of the isomers which are present in the composition.

In yet another embodiment, the isomeric composition of the present invention comprises from 0.1, generally from 0.5, typically from 0.7, preferably from 1.0, more preferably from 1.5 and most preferably from 2.0 to 45, generally to 40, typically to 35, preferably to 30, more preferably to 27 and most preferably to 25 weight percent 2,6-dinitro-4-(1-methylheptyl)phenyl ester, based on the total weight of all of the isomers which are present in the composition.

In another embodiment, the isomeric composition of the present invention comprises 0.1, generally from 0.5, typically from 1.0, preferably from 1.5, more preferably from 2.0 and most preferably from 2.5 to 45, generally to 40, typically to 35, preferably to 30, more preferably to 27 and most preferably to 25 weight percent or less 2,4-dinitro-6-(1-ethylhexyl)phenyl ester.

The isomeric composition of the present invention comprises less than 0.1, generally, less than 0.08, more generally, less than 0.06, preferably less than 0.04, more preferably less than 0.02 and most preferably less than 0.001 weight percent of 2,6-dinitro-4-(1-propylpentyl)phenyl ester. In a most preferred embodiment, the isomeric composition of the present invention is a mixture of isomers in the absence of 2,6-dinitro-4-(1-propylpentyl)phenyl ester. 'The absence of' refers to an undetectable amount as measured using High Pressure Liquid Chromatography.

In another embodiment of the present invention, the isomeric compositions comprise:

from 80 to 98 weight percent 2,4-dinitro-6-(1-methylheptyl)phenyl ester,
less than 1.5 weight percent 2,4-dinitro-6-(1-ethylhexyl)phenyl ester,
less than 0.1 weight percent 2,4-dinitro-6-(1-propylpentyl)phenyl ester,
less than 0.1 weight percent 2,6-dinitro-4-(1-methylheptyl)phenyl ester,
less than 0.1 weight percent 2,6-dinitro-4-(1-ethylhexyl)phenyl ester, and
less than 0.1 weight percent 2,6-dinitro-4-(1-propylpentyl)phenyl ester.

The isomeric composition of the present invention can be made by a variety of methods as long as the final isomeric composition contains less than 0.1 wt. percent of 2,6-dinitro-4-(1-propylpentyl)phenyl ester. Dinocap is typically obtained by conversion of a mixture of dinitro-octylphenols to dinitro-octylphenyl crotonates by reaction with crotonyl chloride, as is well known in the art and described in "E.Y. Guide to the
5 Chemicals Used in Crop Protection." 7th ed. Publication 1093, Research Institute, Agriculture Canada, Ottawa, Canada: Information Canada, 1982. 229. Other esters can also be made by substituting the appropriate material for the crotonyl chloride in order to place a different ester on the phenyl ring. The isomeric composition can be obtained
10 using the above process by utilizing a mixture of dinitro-octylphenols, containing less than 0.1 weight percent 2,6-dinitro-4-(1-propylpentyl)phenol in the reaction, which can in turn be produced by the dinitration of octylphenols containing less than 0.1 weight percent 4-(1-propylpentyl)phenol. Highly pure starting materials can be obtained by any known process including purification and separation processes necessary to achieve the
15 desired purity. For example, methods contemplated include any known method for producing 2,4-dinitro-6-(1-methylheptyl)phenol or any isomer of 2,4-dinitro-6-octylphenol, 2,6-dinitro-4-octylphenol, or mixture thereof, followed by appropriate purification and/or separation methods in order to obtain the desired highly pure starting material having less than 0.1 weight percent 2,6-dinitro-4-(1-propylpentyl)phenol.
20 Alternatively, highly pure methylheptyl phenol can be nitrated to produce a highly pure dinitro-octylphenol. Highly pure methylheptyl phenol can be obtained by any known process for alkylating phenol, followed by purification such as distillation, chromatography and the like.

Still other methods contemplated herein include any known method for producing
25 isomeric mixtures of 2,4-dinitro-6-octylphenyl esters and 2,6-dinitro-4-octylphenyl esters, followed by appropriate purification and/or separation methods in order to obtain the isomeric composition of the present invention. Purification and separation techniques can be any known technique to one skilled in the art, including distillation,

chromatography, and the like. Other related methods of producing such isomeric mixtures also include those disclosed in U.S. Patent 2,810,767 and U.S. Patent 2,526,660, which are both incorporated herein by reference.

In one embodiment, 2,4-dinitro-6-(1-methylheptyl)phenol having a purity of at least 98.5 percent and having less than 0.1 weight percent 2,6-dinitro-4-(1-propylpentyl)phenol is utilized in preparing the isomeric composition of the present invention.

Additionally, another embodiment of the present invention is a fungicidal composition, useful for protecting a plant against attack by a phytopathogenic organism and/or treatment of a plant infested by a phytopathogenic organism, comprising

I) an isomeric composition of dinitro-octylphenyl esters containing less than 0.1 weight percent of 2,6-dinitro-4-(1-propylpentyl)phenyl esters, based on the total weight of the isomeric composition, and

II) a phytologically acceptable carrier material.

The isomeric composition of the present invention contemplates all vehicles by which the isomeric composition can be formulated for delivery and used as a fungicide. Typically, formulations are applied as aqueous suspensions or emulsions. Such suspensions or emulsions are produced from water-soluble, water suspendable, or emulsifiable formulations which are (1) solids, usually known as wettable powders or water dispersible granules or (2) liquids, usually known as emulsifiable concentrates, aqueous emulsions, suspension concentrates and water suspended capsules containing the isomeric composition. As will be readily appreciated, any material to which the isomeric composition can be added may be used, provided they yield the desired utility without significant interference with the activity of the isomeric composition as antifungal agents.

Wettable powders, which may be compacted, extruded or processed through a dispersion in water followed by spray drying or fluid bed agglomeration to form water dispersible granules, comprise an intimate mixture of the isomeric composition, an inert carrier and surfactants. The concentration of the isomeric composition in the wettable

powder is usually from 10 percent to 90 percent by weight based on the total weight of the wettable powder, more preferably 25 wt. percent to 75 wt. percent. In the preparation of wettable powder formulations, the isomeric composition can be compounded with any finely divided solid, such as prophyllite, talc, chalk, gypsum, Fuller's earth, bentonite, attapulgite, starch, casein, gluten, montmorillonite clays, diatomaceous earths, purified silicates or the like. In such operations, the finely divided carrier and surfactants are typically blended with the isomeric composition and milled.

Emulsifiable concentrates of the isomeric composition comprise a convenient concentration, such as from 5 wt. percent to 75 wt. percent of the isomeric composition, in a suitable liquid, based on the total weight of the concentrate. The isomeric composition is dissolved in an inert carrier, which is either water, a water miscible solvent, a water immiscible solvent, or a mixture thereof and emulsifiers. The concentrates may be diluted with water and oil to form spray mixtures in the form of oil-in-water emulsions. Useful organic solvents include aromatics, especially the high-boiling naphthalenic and olefinic portions of petroleum such as heavy aromatic naphtha. Other organic solvents may also be used, such as, for example, terpenic solvents, including rosin derivatives, aliphatic ketones, such as cyclohexanone, and complex alcohols, such as 2-ethoxyethanol.

Emulsifiers which can be advantageously employed herein can be readily determined by those skilled in the art and include various nonionic, anionic, cationic and amphoteric emulsifiers, or a blend of two or more emulsifiers. Examples of nonionic emulsifiers useful in preparing the emulsifiable concentrates include the polyalkylene glycol ethers and condensation products of alkyl and aryl phenols, aliphatic alcohols, aliphatic amines or fatty acids with ethylene oxide, propylene oxides such as the ethoxylated alkyl phenols and carboxylic esters solubilized with the polyol or polyoxyalkylene. Cationic emulsifiers include quaternary ammonium compounds and fatty amine salts. Anionic emulsifiers include the oil-soluble salts (e.g., calcium) of

alkylaryl sulphonic acids, oil soluble salts or sulfated polyglycol ethers and appropriate salts of phosphated polyglycol ether.

Representative organic liquids which can be employed in preparing the emulsifiable concentrates of the isomeric composition are the aromatic liquids such as xylene, propyl benzene fractions; or mixed naphthalene fractions, mineral oils, substituted aromatic organic liquids such as dioctyl phthalate; kerosene; dialkyl amides of various fatty acids, particularly the dimethyl amides of fatty glycols and glycol derivatives such as the n-butyl ether, ethyl ether or methyl ether of diethylene glycol, and the methyl ether of triethylene glycol and the like. Mixtures of two or more organic liquids may also be employed in the preparation of the emulsifiable concentrate. Preferred organic liquids include xylene, and propyl benzene fractions, with propylbenzene fractions being most preferred. Surface-active emulsifying agents are typically employed in liquid formulations and in an amount of from 0.1 to 20 percent by weight based on the combined weight of the emulsifying agent with the isomeric composition. The formulations comprising the isomeric composition of the present invention can also contain other compatible additives, for example, miticides, insecticides, plant growth regulators, other fungicides, and other biologically active compounds used in agriculture.

Aqueous suspensions comprise suspensions of the isomeric composition, dispersed in an aqueous vehicle at a concentration in the range from 5 to 50 weight percent, based on the total weight of the aqueous suspension. Aqueous suspensions are prepared by vigorously mixing the isomeric composition of the present invention, or its solution, into a vehicle comprised of water and surfactants chosen from the same types discussed above. Other components, such as inorganic salts and synthetic or natural gums, may also be added to increase the density and viscosity of the aqueous vehicle. Examples of aqueous suspensions include suspensions of oil droplets (EW's), solids (SC's), and capsules (CS's).

The isomeric composition can also be applied as granular formulations, which are particularly useful for applications to the soil. Granular formulations usually contain from 0.5 to 10 wt. percent, based on the total weight of the granular formulation of the isomeric composition, dispersed in an inert carrier which consists entirely or in large part of coarsely divided inert material such as attapulgite, bentonite, diatomite, clay or a similar inexpensive substance. Such formulations are usually prepared by diluting the isomeric composition in a suitable solvent and applying it to a granular carrier which has been preformed to the appropriate particle size, in the range of from 0.5 to 3 mm. A suitable solvent is a solvent in which the compound is substantially or completely soluble. Such formulations may also be prepared by making a dough or paste of the carrier and the isomeric composition and solvent, and crushing and drying to obtain the desired granular particle.

The isomeric composition of the present invention can also be applied as a water dispersible granule, or dry flowable formulation. Water dispersible granules typically contain from 10 to 70 percent of the isomeric composition, based on the total weight of the formulation. Such formulations are typically obtained through mixing and/or spraying the isomeric mixture onto a carrier with the addition of a dispersing and/or wetting agent, and combining with water to form a mixture suitable for further processing using well known granulation technologies, such as pan granulation, extrusion, spray-drying, fluid bed agglomeration, and the like.

Dusts containing the isomeric composition can be prepared by intimately mixing the isomeric composition with a suitable dusty agricultural carrier, such as, for example, kaolin clay, ground volcanic rock, and the like. Dusts can suitably contain from 1 to 10 wt. percent of the isomeric composition or other fungicide, based on the total weight of the dust. Dusts may also be prepared by impregnating the isomeric composition onto a carrier in a similar manner to that described for granules above.

The formulations of the present invention may additionally contain adjuvant surfactants to enhance deposition, wetting and penetration of the isomeric composition

onto the target crop and organism. These adjuvant surfactants may optionally be employed as a component of the formulation or as a tank mix. The amount of adjuvant surfactant will typically vary from 0.01 to 1.0 percent by volume, based on a spray-volume of water, preferably 0.05 to 0.5 volume percent. Suitable adjuvant surfactants
5 include, but are not limited to ethoxylated nonyl phenols, ethoxylated synthetic or natural alcohols, salts of the esters or sulphosuccinic acids, ethoxylated organosilicones, ethoxylated fatty amines and blends of surfactants with mineral or vegetable oils.

One embodiment of the present invention relates to a synergistic mixture of fungicides comprising the isomeric composition of the present invention and at least one
10 other fungicide, and its' use for protection of a plant against attack by a phytopathogenic organism or the treatment of a plant already infested by a phytopathogenic organism, comprising applying the synergistic mixture, to soil, a plant, a part of a plant, foliage, flowers, and/or fruit.

Fungicidal compounds are often applied in conjunction with one or more other
15 fungicides to control a wider variety of undesirable diseases. When used in conjunction with other fungicide(s), the presently claimed isomeric composition can be formulated with the other fungicide(s), tank mixed with the other fungicide(s) or applied sequentially with the other fungicide(s). Such other fungicides include amisulbrom 2-(thiocyanatomethylthio)-benzothiazole, 2-phenylphenol, 8-hydroxyquinoline sulfate,
20 antimycin, Ampelomyces, quisqualis, azaconazole, azoxystrobin, Bacillus subtilis, benalaxyl, benomyl, benthiavalicarb-isopropyl, benzylaminobenzene-sulfonate (BABS) salt, bicarbonates, biphenyl, bismethiazol, bitertanol, blastidicid-S, borax, Bordeaux mixture, boscalid, bromuconazole, bupirimate, calcium polysulfide, captafol, captan, carbendazim, carboxin, carpropamid, carvone, chloroneb, chlorothalonil, chlozolinate,
25 Coniothyrium minitans, copper hydroxide, copper octanoate, copper oxychloride, copper sulfate, copper sulfate (tribasic), coumarins, cuprous oxide, cyazofamid, cyflufenamid, cymoxanil, cyproconazole, cyprodinil, dazomet, debacarb, diammonium ethylenebis-(dithiocarbamate), dichlofluanid, dichlorophen, diclocymet, diclomezine, dichloran,

diethofencarb, difenoconazole, difenzoquat ion, diflumetorim, dimethomorph,
dimoxystrobin, diniconazole, diniconazole-M, dinobuton, diphenylamine, dithianon,
dodemorph, dodemorph acetate, dodine, dodine free base, edifenphos, enestrobin,
epoxiconazole, ethaboxam, ethoxyquin, etridiazole, famoxadone, fenamidone, fenarimol,
5 fenbuconazole, fenfuram, fenhexamid, fenoxanil, fenciclonil, fenpropidin,
fenpropimorph, fentin, fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam,
fluidioxonil, flumorph, fluopicolide, fluoroimide, fluoxastrobin, fluquinconazole,
flusilazole, flusulfamide, flutolanil, flutriafol, folpet, formaldehyde, fosetyl, fosetyl-
aluminium, fuberidazole, furalaxyl, furametpyr, guazatine, guazatine acetates, GY-81,
10 hexachlorobenzene, hexaconazole, hymexazol, imazalil, imazalil sulfate,
imibenconazole, iminoctadine, iminoctadine triacetate, iminoctadine tris(albesilate),
ipconazole, iprobenfos, iprodione, iprovalicarb, isoprothiolane, kasugamycin,
kasugamycin hydrochloride hydrate, kresoxim-methyl, mancozeb, mancozeb,
mandipropamid, maneb, mepanipyrim, mepronil, mercuric chloride, mercuric oxide,
15 mercurous chloride, metalaxyl, mefenoxam, metalaxyl-M, metam, metam-ammonium,
metam-potassium, metam-sodium, metconazole, methasulfocarb, methyl iodide, methyl
isothiocyanate, metiram, metominostrobin, metrafenone, mildiomyacin, myclobutanil,
nabam, nitrothal-isopropyl, nuarimol, octhilinone, ofurace, oleic acid (fatty acids),
orysastrobin, oxadixyl, oxine-copper, oxpoconazole fumarate, oxycarboxin, pefurazoate,
20 penconazole, pencycuron, pentachlorophenol, pentachlorophenyl laurate, penthiopyrad,
phenylmercury acetate, phosphonic acid, phthalide, picoxystrobin, polyoxin B, polyoxins,
polyoxorim, potassium bicarbonate, potassium hydroxyquinoline sulfate, probenazole,
prochloraz, procymidone, propamocarb, propamocarb hydrochloride, propiconazole,
propineb, proquinazid, prothioconazole, pyraclostrobin, pyrazophos, pyributicarb,
25 pyrifenox, pyrimethanil, pyroquilon, quinochloramine, quinoxifen, quintozone, Reynoutria
sachalinensis extract, silthiofam, simeconazole, sodium 2-phenylphenoxide, sodium
bicarbonate, sodium pentachlorophenoxide, spiroxamine, sulfur, SYP-Z071, SYP-048,
tar oils, tebuconazole, tecnazene, tetraconazole, thiabendazole, thifluzamide, thiophanate-

methyl, thiram, tiadinil, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol,
 triazolopyrimidine, triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole,
 triforine, triticonazole, validamycin, vinclozolin, zineb, ziram, zoxamide, *Candida*
oleophila, *Fusarium oxysporum*, *Gliocladium* spp., *Phlebiopsis gigantea*, *Streptomyces*
 5 *griseoviridis*, *Trichoderma* spp., (RS)-N-(3,5-dichlorophenyl)-2-(methoxymethyl)-
 succinimide, 1,2-dichloropropane, 1,3-dichloro-1,1,3,3-tetrafluoroacetone hydrate, 1-
 chloro-2,4-dinitronaphthalene, 1-chloro-2-nitropropane, 2-(2-heptadecyl-2-imidazolin-1-
 yl)ethanol, 2,3-dihydro-5-phenyl-1,4-dithiane 1,1,4,4-tetraoxide, 2-methoxyethylmercury
 acetate, 2-methoxyethylmercury chloride, 2-methoxyethylmercury silicate, 3-(4-
 10 chlorophenyl)-5-methylrhodanine, 4-(2-nitroprop-1-enyl)phenyl thiocyanate:
 ampropylfos, anilazine, azithiram, barium polysulfide, Bayer 32394, benodanil,
 benquinox, bentazon, benzamocil; benzamocil-isobutyl, benzamorf, binapacryl,
 bis(methylmercury) sulfate, bis(tributyltin) oxide, buthiobate, cadmium calcium copper
 zinc chromate sulfate, carbamorph, CECA, chlorobenzothiazole, chloranilmethane,
 15 chlorfenazole, chlorquinox, clomazone, copper bis(3-phenylsalicylate), copper zinc
 chromate, cufraneb, cupric hydrazinium sulfate, cuprobum, cyclafuramid, cypendazole,
 cyproflumuron, decafenit, dichlorone, dichlorzoline, diclobutrazol, dimethirimol, dinocron,
 dinosulfon, dinoterbon, dipyrithione, ditalimfos, dodicin, drazoxolon, EBP, ESBP,
 etaconazole, etem, ethirim, fenaminosulf, fenapanil, fenitropan,
 20 fluotrimazole, furcarbanil, furconazole, furconazole-cis, furmecycloz, furophanate,
 glyodine, griseofulvin, halacrinat, Hercules 3944, hexylthiofos, ICIA0858, isopamphos,
 isovaldione, mebenil, mecarbinid, metazoxolon, methfuroxam, methylmercury
 dicyandiamide, metsulfovax, milneb, mucochloric anhydride, myclozolin, N-3,5-
 dichlorophenyl-succinimide, N-3-nitrophenylitaconimide, natamycin, N-ethylmercurio-4-
 25 toluenesulfonanilide, nickel bis(dimethyldithiocarbamate), OCH, phenylmercury
 dimethyldithiocarbamate, phenylmercury nitrate, phosdiphen, prothiocarb; prothiocarb
 hydrochloride, pyracarbolid, pyridinitril, pyroxychlor, pyroxyfur, quinacetol; quinacetol
 sulfate, quinazamid, quinconazole, rabenzazole, salicylanilide, SSF-109, sultropen,

tecoram, thiadifluor, thicyofen, thiochlorfenphim, thiophanate, thioquinox, tioxymid, triamiphos, triarimol, triazbutil, trichlamide, urbacid, XRD-563, and zarilamid, IK-1140, NC-224, and any combinations thereof.

Synergy has been found previously with dinocap as disclosed in references such as, but not limited to WO 02/067679; WO2003103393; WO 2004091298; FR 2445696; GB 2003032; U.S. 6,346,535; U.S. 6,528,536; U.S. 6,515,000; U.S. 6,207,691; U.S. 3,456,055; and U.S. 6,489,360. However, synergies with the composition of the present invention have not been previously reported.

Another aspect of the present invention is related to synergistic mixtures of the isomeric composition described herein with at least one other fungicide, which surprisingly have improved toxicity properties. The synergistic composition of the present invention can comprise 1) any fungicide which is synergistic with dinocap and 2) the isomeric composition as detailed herein. Exemplary fungicides known to show synergistic effects include, but are not limited to, QoI fungicides such as, methoxyacrylates including azoxystrobin, and oximino acetates including kresoxim methyl; SBI fungicides: class I such as triazoles including myclobutanil; class II such as spiroketal-amines including spiroxamine; and class III such as hydroxyanilides including fenhexamid; benzophenones such as metrafenone; quinolines such as quinoxifen; dithiocarbamates and derivatives such as mancozeb; chloronitriles such as chlorothalonil; respiration inhibition at complex II succinate dehydrogenase carboxamides including boscalid, and the like.

In one specific embodiment, the present invention is a synergistic mixture comprising:

- i) an isomeric composition comprising isomers of dinitro-(octyl)phenyl esters, wherein the 2,6-dinitro-4-(1-propylpentyl)phenyl ester isomer is present in an amount of less than 0.1 weight percent, based on the total weight of the isomeric composition, and
- ii) QoI fungicide (commonly referred to as strobilurins and related chemistry).

QoI fungicides include compounds such as azoxystrobin, dimoxystrobin, fluoxastrobin, orysastrobin, picoxystrobin, pyraclostrobin, trifloxystrobin, kresoxim-methyl, famoxadone, fenamidone and the like.

In another embodiment, the present invention is a synergistic mixture comprising:

- 5 i) an isomeric composition comprising isomers of dinitro-(octyl)phenyl esters, wherein the 2,6-dinitro-4-(1-propylpentyl)phenyl ester isomer is present in an amount of less than 0.1 weight percent, based on the total weight of the isomeric composition, and
- ii) a quinoline or cinnoline.

Quinolines and cinnolines, such as Quinoxifen are reported in U.S. 5,145,843, which is
10 incorporated herein by reference.

In another embodiment, the present invention is a synergistic mixture comprising:

- i) an isomeric composition comprising isomers of dinitro-(octyl)phenyl esters, wherein the 2,6-dinitro-4-(1-propylpentyl)phenyl ester isomer is present in an amount of less than 0.1 weight percent, based on the total weight of the isomeric composition, and
- 15 ii) a dithiocarbamate.

Dithiocarbamates include compounds such as mancozeb (a coordination product of zinc and manganese ethylene bisdithiocarbamate), maneb (manganese ethylenebisdithiocarbamate) and zineb (zinc ethylenebisdithiocarbamate), ziram (zinc dimethyldithiocarbamate), propineb ([[1-methyl-1,2-ethanediy]bis[carbamatothioato]](2-))zinc
20 homopolymer), metiram)(tris[amine-[ethylene bis(dithiocarbamate)]-zinc(II)-[tetrahydro-1,2,4,7-dithiadiazocine-3,8-dithione]polymer), ferbam (ferric dimethyldithiocarbamate), metham (sodium N-methyldithiocarbamate), and thiram (bis(dimethylthiocarbamoyl) disulfide) and the like.

In another embodiment, the present invention is a synergistic mixture comprising:

- 25 i) an isomeric composition comprising isomers of dinitro-(octyl)phenyl esters, wherein the 2,6-dinitro-4-(1-propylpentyl)phenyl ester isomer is present in an amount of less than 0.1 weight percent, based on the total weight of the isomeric composition, and
- ii) a triazole.

Triazoles include compounds such as azaconazole, bitertanol, bromuconazole, chlorfenazole, climbazole, cypendazole, cyproconazole, diclobutrazol, difenoconazole, diniconazole, diniconazole-M, epoxiconazole, etaconazole, etridiazole, fenbuconazole, fluotrimazole, fluquinconazole, flusilazole, flutriafol, fuberidazole, furconazole, and
5 furconazole-cis. hexaconazole, imazalil, imibenconazole, ipconazole, metconazole, myclobutanil, pefurazoate, penconazole, probenazole, prochloraz, propiconazole, prothioconazole, quinconazole, rabenzazole, simeconazole, tebuconazole, tetraconazole, triadimefon, triadimenol, tricyclazole, triflumazole, triticonazole and the like.

In another embodiment, the present invention is a synergistic mixture comprising:

- 10 i) an isomeric composition comprising isomers of dinitro-(octyl)phenyl esters, wherein the 2,6-dinitro-4-(1-propylpentyl)phenyl ester isomer is present in an amount of less than 0.1 weight percent, based on the total weight of the isomeric composition, and
ii) spiroxamine.

The amount of the isomeric mixture and other fungicide within the synergistic
15 mixture can vary depending on the application desired and diseases for desired control. Generally, the isomeric composition is present in amounts of from 0.1, more generally from 1, even more generally from 5, most generally from 10, typically from 15, more typically from 20, even more typically from 30, most typically from 40, and routinely from 45 weight percent to generally 99.9, more generally to 99, even more generally to
20 95, most generally to 90, typically to 85, more typically to 80, even more typically to 70, most typically to 60, and routinely to 55 weight percent based on the total weight of the isomeric composition and other fungicide(s).

Generally, the at least one other fungicide is present in amounts of from 99.9, more generally 99, even more generally 95, most generally 90, typically 85, more
25 typically 80, even more typically 70, most typically 60, and routinely 55 to 0.1, more generally 1, even more generally 5, most generally 10, typically 15, more typically 20, even more typically 30, most typically 40, and routinely 45 weight percent based on the total weight of the isomeric composition and other fungicide(s).

Additionally, the isomeric compositions and synergistic mixtures can be combined with other pesticides including insecticides, nematocides, miticides, arthropodicides, bactericides or combinations thereof that are compatible with the isomeric composition in the medium selected for application, and not antagonistic to the activity of the isomeric composition. Accordingly, in such embodiments, the other pesticidal compound is employed as a supplemental toxicant for a different pesticidal use. When used in conjunction with other pesticides, the isomeric composition or synergistic mixture can be formulated with the other pesticide(s), tank mixed with the other pesticide(s) or applied sequentially with the other pesticide(s). Typical insecticides include, but are not limited to: antibiotic insecticides such as allosamidin and thuringiensin; macrocyclic lactone insecticides such as spinosad; avermectin insecticides such as abamectin, doramectin, emamectin, eprinomectin, ivermectin and selamectin; milbemycin insecticides such as lepipmectin, milbemectin, milbemycin oxime and moxidectin; arsenical insecticides such as calcium arsenate, copper acetoarsenite, copper arsenate, lead arsenate, potassium arsenite and sodium arsenite; botanical insecticides such as anabasine, azadirachtin, d-limonene, nicotine, pyrethrins, cinerins, cinerin I, cinerin II, jasmolin I, jasmolin II, pyrethrin I, pyrethrin II, quassia, rotenone, ryania and sabadilla; carbamate insecticides such as bendiocarb and carbaryl; benzofuranyl methylcarbamate insecticides such as benfuracarb, carbofuran, carbosulfan, decarbofuran and furathiocarb; dimethylcarbamate insecticides dimitan, dimetilan, hyquincarb and pirimicarb; oxime carbamate insecticides such as alanycarb, aldicarb, aldoxycarb, butocarboxim, butoxycarboxim, methomyl, nitrilacarb, oxamyl, tazimcarb, thiocarboxime, thiodicarb and thiofanox; phenyl methylcarbamate insecticides such as allyxycarb, aminocarb, bufencarb, butacarb, carbanolate, cloethocarb, dicresyl, dioxacarb, EMPC, ethiofencarb, fenethacarb, fenobucarb, isoprocarb, methiocarb, metolcarb, mexacarbate, promacyl, promecarb, propoxur, trimethacarb, XMC and xylylcarb; dinitrophenol insecticides such as dinex, dinoprop, dinosam and DNOC; fluorine insecticides such as barium hexafluorosilicate, cryolite, sodium fluoride, sodium

hexafluorosilicate and sulfluramid; formamidine insecticides such as amitraz, chlordimeform, formetanate and formparanate; fumigant insecticides such as acrylonitrile, carbon disulfide, carbon tetrachloride, chloroform, chloropicrin, para-dichlorobenzene, 1,2-dichloropropane, ethyl formate, ethylene dibromide, ethylene
5 dichloride, ethylene oxide, hydrogen cyanide, iodomethane, methyl bromide, methylchloroform, methylene chloride, naphthalene, phosphine, sulfuryl fluoride and tetrachloroethane; inorganic insecticides such as borax, calcium polysulfide, copper oleate, mercurous chloride, potassium thiocyanate and sodium thiocyanate; chitin synthesis inhibitors such as bistrifluron, buprofezin, chlorfluazuron, cyromazine,
10 diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, noviflumuron, penfluron, teflubenzuron and triflumuron; juvenile hormone mimics such as epofenonane, fenoxycarb, hydroprene, kinoprene, methoprene, pyriproxyfen and triprene; juvenile hormones such as juvenile hormone I, juvenile hormone II and juvenile hormone III; moulting hormone agonists such as chromafenozide, halofenozide,
15 methoxyfenozide and tebufenozide; moulting hormones such as α -ecdysone and ecdysterone; moulting inhibitors such as diofenolan; precocenes such as precocene I, precocene II and precocene III; unclassified insect growth regulators such as dicyclanil; nereistoxin analogue insecticides such as bensultap, cartap, thiocyclam and thiosultap; nicotinoid insecticides such as flonicamid; nitroguanidine insecticides such as
20 clothianidin, dinotefuran, imidacloprid and thiamethoxam; nitromethylene insecticides such as nitenpyram and nithiazine; pyridylmethylaniline insecticides such as acetamiprid, imidacloprid, nitenpyram and thiacloprid; organochlorine insecticides such as bromo-DDT, camphechlor, DDT, pp'-DDT, ethyl-DDD, HCH, gamma-HCH, lindane, methoxychlor, pentachlorophenol and TDE; cyclodiene insecticides such as aldrin,
25 bromocyclen, chlorbicyclen, chlordane, chlordecone, dieldrin, dilor, endosulfan, endrin, HEOD, heptachlor, HHDN, isobenzan, isodrin, kelevan and mirex; organophosphate insecticides such as bromfenvinfos, chlorfenvinphos, crotoxyphos, dichlorvos, dicrotophos, dimethylvinphos, fospirate, heptenophos, methocrotophos, mevinphos,

monocrotophos, naled, naftalofos, phosphamidon, propaphos, TEPP and tetrachlorvinphos; organothiophosphate insecticides such as dioxabenzofos, fosmethilan and phenthoate; aliphatic organothiophosphate insecticides such as acethion, amiton, cadusafos, chlorethoxyfos, chlormephos, demephion, demephion-O, demephion-S, 5 demeton, demeton-O, demeton-S, demeton-methyl, demeton-O-methyl, demeton-S-methyl, demeton-S-methylsulphon, disulfoton, ethion, ethoprophos, IPSP, isothioate, malathion, methacrifos, oxydemeton-methyl, oxydeprofos, oxydisulfoton, phorate, sulfotep, terbufos and thiometon; aliphatic amide organothiophosphate insecticides such as amidithion, cyanthoate, dimethoate, ethoate-methyl, formothion, mecarbam, 10 omethoate, prothoate, sophamide and vamidothion; oxime organothiophosphate insecticides such as chlorphoxim, phoxim and phoxim-methyl; heterocyclic organothiophosphate insecticides such as azamethiphos, coumaphos, coumithoate, dioxathion, endothion, menazon, morphothion, phosalone, pyraclofos, pyridaphenthion and quinothion; benzothiopyran organothiophosphate insecticides such as dithicrofos and 15 thicrofos; benzotriazine organothiophosphate insecticides such as azinphos-ethyl and azinphos-methyl; isoindole organothiophosphate insecticides such as dialifos and phosmet; isoxazole organothiophosphate insecticides such as isoxathion and zolaprofos; pyrazolopyrimidine organothiophosphate insecticides such as chlorprazophos and pyrazophos; pyridine organothiophosphate insecticides such as chlorpyrifos and 20 chlorpyrifos-methyl; pyrimidine organothiophosphate insecticides such as butathiofos, diazinon, etrimfos, lirimfos, pirimiphos-ethyl, pirimiphos-methyl, primidophos, pyrimitate and tebupirimfos; quinoxaline organothiophosphate insecticides such as quinalphos and quinalphos-methyl; thiadiazole organothiophosphate insecticides such as athidathion, lythidathion, methidathion and prothidathion; triazole organothiophosphate 25 insecticides such as isazofos and triazophos; phenyl organothiophosphate insecticides such as azothoate, bromophos, bromophos-ethyl, carbophenothion, chlorthiophos, cyanophos, cythioate, dicapthon, dichlofenthion, etaphos, famphur, fenchlorphos, fenitrothion fensulfothion, fenthion, fenthion-ethyl, heterophos, jodfenphos, mesulfenfos,

parathion, parathion-methyl, phenkapton, phosnichlor, profenofos, prothiofos, sulprofos, temephos, trichlormetaphos-3 and trifenofos; phosphonate insecticides such as butonate and trichlorfon; phosphonothioate insecticides such as mecarphon; phenyl ethylphosphonothioate insecticides such as fonofos and trichloronat; phenyl

5 phenylphosphonothioate insecticides such as cyanofenphos, EPN and leptophos; phosphoramidate insecticides such as crufomate, fenamiphos, fosthietan, mephosfolan, phosfolan and pirimetaphos; phosphoramidothioate insecticides such as acephate, isocarbophos, isofenphos, methamidophos and propetamphos; phosphorodiamide insecticides such as dimefox, mazidox, mipafox and schradan; oxadiazine insecticides

10 such as indoxacarb; phthalimide insecticides such as dialifos, phosmet and tetramethrin; pyrazole insecticides such as acetoprole, ethiprole, fipronil, pyrafluprole, pyriprole, tebufenpyrad, tolfenpyrad and vanilprole; pyrethroid ester insecticides such as acrinathrin, allethrin, bioallethrin, barthrin, bifenthrin, bioethanomethrin, cyclethrin, cycloprothrin, cyfluthrin, beta-cyfluthrin, cyhalothrin, gamma-cyhalothrin, lambda-

15 cyhalothrin, cypermethrin, alpha-cypermethrin, beta-cypermethrin, theta-cypermethrin, zeta-cypermethrin, cyphenothrin, deltamethrin, dimefluthrin, dimethrin, empenthrin, fenfluthrin, fenpirithrin, fenpropathrin, fenvalerate, esfenvalerate, flucythrinate, fluvalinate, tau-fluvalinate, furethrin, imiprothrin, metofluthrin, permethrin, biopermethrin, transpermethrin, phenothrin, prallethrin, profluthrin, pyresmethrin,

20 resmethrin, bioresmethrin, cismethrin, tefluthrin, terallethrin, tetramethrin, tralomethrin and transfluthrin; pyrethroid ether insecticides such as etofenprox, flufenprox, halfenprox, protrifenbute and silafluofen; pyrimidinamine insecticides such as flufenerim and pyrimidifen; pyrrole insecticides such as chlorfenapyr; tetrone acid insecticides such as spiromesifen, spirodiclofen and spirotetramet; thiourea insecticides such as

25 diafenthiuron; urea insecticides such as flucofuron and sulcofuron; and unclassified insecticides such as closantel, crotamiton, EXD, fenazaflor, fenoxacrim, flubendiamide, hydramethylnon, isoprothiolane, malonoben, metaflumizone, metoxadiazone, nifluridide,

pyridaben, pyridalyl, rafoxanide, triarathene and triazamate, and any combinations thereof.

The amount of the isomeric composition or synergistic mixture thereof, and other pesticide within a pesticidal mixture can vary depending on the application desired and diseases for desired control. Generally, the isomeric composition or synergistic mixture of the present invention is present in amounts of from 0.1, more generally from 1, even more generally from 5, most generally from 10, typically from 15, more typically from 20, even more typically from 30, most typically from 40, and routinely from 45 weight percent to generally 99.9, more generally to 99, even more generally to 95, most generally to 90, typically to 85, more typically to 80, even more typically to 70, most typically to 60, and routinely to 55 weight percent, based on the total weight of the isomeric composition or synergistic mixture and other pesticide(s).

Generally, when employed, the other pesticide is present in amounts of from 99.9, more generally 99, even more generally 95, most generally 90, typically 85, more typically 80, even more typically 70, most typically 60, and routinely 55 to 0.1, more generally 1, even more generally 5, most generally 10, typically 15, more typically 20, even more typically 30, most typically 40, and routinely 45 weight percent, based on the total weight of the isomeric composition or synergistic mixture and other fungicide(s).

Another embodiment of the present invention is a method for the control or prevention of fungal attack. This method comprises applying to the plant, foliage, locus of the fungus, or to a locus in which the infestation is to be prevented (for example applying to cereal or grape plants), a fungicidal effective amount of the isomeric composition or synergistic mixture therefrom. The isomeric composition or synergistic mixture of the present invention is suitable for treatment of various plants at fungicidal levels, while exhibiting no teratogenicity or retinopathy. The isomeric composition or synergistic mixture is useful both in a protectant, curative and/or an eradicant fashion.

The isomeric composition and synergistic mixture therefrom have been found to have significant fungicidal effect particularly for agricultural use and is particularly

effective for use with agricultural crops and horticultural plants. In particular, the isomeric composition and synergistic mixture therefrom effectively controls a variety of undesirable fungi that infect useful plant crops, such as powdery mildews in pome fruit, stone fruit, citrus fruit, soft fruit, vines, cucurbits, ornamentals, tobacco, hops, and some
5 vegetables, and American gooseberry mildew in gooseberries and currants.

The exact amount of isomeric composition or synergistic mixture therefrom to be applied is dependent on the particular action desired, the fungal species to be controlled, the stage of growth thereof, as well as the part of the plant or other product to be contacted with the isomeric composition.

10 The isomeric composition and synergistic mixture therefrom are effective in use with plants in a disease-inhibiting and phytologically acceptable amount. The term "disease inhibiting and phytologically acceptable amount" refers to an amount of an active compound that kills or inhibits the plant disease for which control is desired, but is not significantly toxic to the plant. This amount will generally be from 1 to 1000
15 ppm(parts per million), with 10 to 500 ppm being preferred. The exact concentration of isomeric composition or synergistic mixture required varies with the fungal disease to be controlled, the type of formulations employed, the method of application, the particular plant species, climate conditions, and the like. A suitable application rate is typically in the range from 0.10 to 4 pounds/acre (0.01 to 0.45 grams per square meter g/m^2).

20 It has also been surprisingly discovered that the isomeric composition and synergistic mixtures therefrom of the present invention can have a decreased acaracidal effect, which is advantageous in maintaining beneficial mites.

EXAMPLES

These examples are provided to further illustrate the invention and are not meant to be
25 construed as limiting. As disclosed herein, all temperatures are given in degrees Celsius and all percentages are weight percentages, except for percent yields which are mole percentages, unless otherwise stated.

Preparation of 2,4-Dinitro-6-(1-methylheptyl)phenol

2-(1-methylheptyl)phenol (206.3 gm, 98.5 percent assay) is stirred and heated to 60°C. Concentrated sulfuric acid (315.2 gm) is added dropwise over 1 hour, maintaining the temperature in the range 65 – 72°C.

5

In a separate vessel, a 35 percent aqueous solution of sodium nitrate (553.2 gm) is stirred and heated to 70°C. The sulfonated phenol reaction mixture is then added dropwise, over 3 hours to the sodium nitrate solution, maintaining the temperature at 78°C. The mixture is stirred at 78°C for one hour, and then cooled to 35°C. The mixture is allowed to settle, and the aqueous layer separated. The organic layer is washed twice with water containing a little methanol, and then dried under vacuum at 75°C to leave 252 gm of 2,4-dinitro-6-(1-methylheptyl)phenol product as an orange oil.

10

EXAMPLE 1

15

Preparation of 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate

2,4-Dinitro-6-(1-methylheptyl)phenol (98.5 percent pure) (252 gm) is dissolved in heptane (225 gm). Benzyldimethylamine (140.8 gm) is added dropwise over 40 minutes, keeping the temperature of the mixture below 40°C, and the mixture is stirred for 15 minutes. Crotonyl chloride (139 gm) is then added dropwise over 30 minutes, keeping the temperature below 60°C, and the mixture is stirred at 60°C for 30 minutes and cooled to less than 50°C. The mixture is washed twice with water and then cooled to 20°C. The mixture is washed with a mixture of methanol and aqueous sodium hydroxide, with water and methanol, and finally with dilute hydrochloric acid. The organic layer is stripped at 70°C under vacuum to remove heptane and leave 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate (282.6 gm) as an orange/brown oil.

20

25

For analysis, normal phase liquid chromatography is utilized using the following method:

HPLC Conditions

Column temp 40°C

Flow volume 1ml/min

5 Wavelength 235 nm

Injection volume 10 microliters

HPLC Milton Roy Column 250x4.6mm, packed with 7 µ LICHROSORB SI 60 (Merck)

Detection: Detector Milton Roy SM 4000

10 Integration system Shimadzu Mega2.

Sample solution:

A 0.45-0.50 g sample of 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate and 0.09-0.11 g of acetophenone is placed in a vial. Hexane (20 ml), is added, the vial is capped and
15 sonicated for 2 minutes. The solution (1ml) is transferred into a vial and 20 ml of hexane is added, the vial capped and shaken to produce a sample solution.

Standard solution:

0.48-0.52 g of Karathane™ Working Standard and 0.09-0.11 g of acetophenone (I.S.) are
20 placed into a vial. Hexane (20 ml), is added, the vial is capped and sonicated for 2 minutes. The solution (1ml) is transferred into a vial and 20 ml of hexane is added, the vial capped and shaken to produce a standard solution.

10 microliters of standard solution is injected into the HPLC to obtain a chromatogram. Identically, 10 microliters of sample solution is injected into the HPLC to obtain a
25 chromatogram. The chromatograms are compared.

For calculating response factors and ingredient content, the peaks are grouped as:

Internal Standard (I.S.) =Acetophenone

Working Standard=Karathane™

Test Sample= sample being evaluated for isomeric content

Group 1: Sum of the area for 2,4-dinitro-6-octylphenol

5 Group 2: Sum of the area for 2,4-dinitro-6-octylphenyl crotonates

Group 3: Sum of the area for 2,6-dinitro-4-octylphenyl crotonates

$$RF = \frac{\text{Area of I.S. peak} \times \text{wt. of Working Standard}}{\text{Area of the I.S. peak} \times \text{weight of Test Sample}} \times (\text{percent of Group in Working Standard})$$

10
$$\text{Area Group} \times \text{weight I.S.} \times 100$$

An RF factor is calculated for each group.

15
$$\text{Content (percent)} = \frac{\text{Area of the Group peak} \times \text{weight of I.S.} \times RF \times 100}{\text{Area of the I.S. peak} \times \text{weight of Test Sample}}$$

percent Content is calculated for each group using the appropriate RF factor.

For Group 3, the content would be 0 if there were no peaks associated with Group 3.

20 The isomeric percent of Group 2 is then determined by comparing the relative peaks within the group, i.e. 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate 98.5 percent and 2,4-dinitro-6-(1-ethylhexyl)phenyl crotonate 1.2 percent.

The analysis of the product using this normal phase, liquid chromatographic method, is as follows:

25	2,4-dinitro-6-(1-methylheptyl)phenyl crotonate:	98.5 percent
	2,4-dinitro-6-(1-ethylhexyl)phenyl crotonate:	1.2 percent
	2,4-dinitro-6-(1-methylheptyl)phenol:	0.3 percent

EXAMPLES 2 AND 3**Additional Isomer Analysis of Samples prepared according to Example I**

In a separate analysis, two additional samples prepared according to Example I were analyzed for the active ingredient and impurities using normal phase HPLC with internal
5 standard calibration as outlined below.

Normal Phase High Performance Liquid Chromatography (HPLC)**Preparation of Acetophenone Internal Standard Solution:**

10 Approximately 1 g of Acetophenone was weighed into a 500-mL volumetric flask, the weight was recorded to the nearest 0.1 mg and the flask was filled to volume with hexane.

Preparation of calibration solutions:

15 Calibration Solution #1 was prepared in duplicate by weighing approximately 50 mg of 2,4-dinitro-6-(1-methylheptyl)phenyl crotonate (DNOP) standard, into a vial and 20 mL of the acetophenone internal standard solution was added using a volumetric pipette. A 1 mL aliquot was added to 5 mL of hexane.

20 Calibration Solution #2 was prepared by weighing approximately 15 mg each of 2,6-dinitro-4-(1-propylpentyl)phenyl crotonate, 2,4-dinitro-6-(1-propylpentyl)phenyl crotonate (propylpentyl isomer), 2,4-dinitro-6-(1-ethylhexyl)phenyl crotonate (Ethylhexyl isomer) and 2,4-dinitro-6-(1-methylheptyl)phenol into a vial and adding 10 mL of hexane. A 1 mL aliquot of the resulting solution was added to a vial containing 20 mL of
25 the acetophenone internal standard solution. A 1 mL aliquot of this solution was added to 5 mL of hexane.

Calibration Solution #3 was prepared by weighing approximately 10 mg of Dinocap, (mixture of 12 isomers, purity 95.3 percent) into a vial and adding 20 mL of the acetophenone internal standard solution using a volumetric pipette. A 1 mL aliquot of the resulting solution was added to 5 mL of hexane.

5

Calibration Procedure:

The calibration solutions were injected before, during and after the samples, using the following conditions:

10 Column: Lichrosorb SI-60 5 μ m; 250 mm x 4.6 mm
Autosampler: Alcott 728
Pump: Varian 9012
Injection: 10 μ L
Detector: Applied Biosystems 757

15 Column Temperature: 30°C
Column Oven: Phenomenex
Data System: PE/Nelson Access*Chrom
Mobile Phase: 720 mL hexane/280 mL dichloromethane/1 mL polar mix (85 mL dichloromethane/10 mL acetic acid/5 mL ethanol)

20

Approximate Retention Times:

2,4-dinitro-6-(1-propylpentyl) phenol	5.79 minutes
2,4-dinitro-6-(1-ethylhexyl) phenol	5.90 minutes
2,4-dinitro-6-(1-methylhexyl) phenol	6.13 minutes
25 unknown peak	6.42 minutes
2,6-dinitro-4-(1-propylpentyl) phenyl crotonate cis	7.49 minutes
2,6-dinitro-4-(1-ethylhexyl) phenyl crotonate cis	7.92 minutes
2,6-dinitro-4-(1-methylhexyl) phenyl crotonate cis	8.15 minutes

	2,6-dinitro-4-(1-propylpentyl) phenyl crotonate trans	9.70 minutes
	2,6-dinitro-4-(1-ethylhexyl) phenyl crotonate trans	10.40 minutes
	2,6-dinitro-4-(1-methylhexyl) phenyl crotonate trans	10.89 minutes
	2,4-dinitro-6-(1-propylpentyl) phenyl crotonate cis	11.30 minutes
5	2,4-dinitro-6-(1-ethylhexyl) phenyl crotonate cis	11.53 minutes
	2,4-dinitro-6-(1-methylhexyl) phenyl crotonate cis	12.04 minutes
	2,4-dinitro-6-(1-propylpentyl) phenyl crotonate trans	15.50 minutes
	2,4-dinitro-6-(1-ethylhexyl) phenyl crotonate trans	16.00 minutes
	2,4-dinitro-6-(1-methylhexyl) phenyl crotonate trans	16.98 minutes
10	Acetophenone (ISTD)	31.82 minutes

Response factors were calculated for each as described previously. The average of the response factors was used for calibration.

15 Preparation of samples for analysis:

Duplicate weighings of approximately 50 mg (± 10 mg), recorded to the nearest 0.1 mg, of EXAMPLES 2 and 3, prepared according to Example 1 were placed into separate vials. To each vial, 20 mL of the acetophenone internal standard solution was added using a volumetric pipette. A 1 mL aliquot of the resulting solution was added to 5 mL
20 of hexane. The samples were analyzed using the same conditions as the standard.
Weight percent analysis of EXAMPLES 2 and # are listed in TABLE I.

TABLE I

Component	EXAMPLE 2 (wt percent)	EXAMPLE 3 (wt percent)
2,4-dinitro-6-(1-methylheptyl)phenyl crotonate	97.9	98.0
2,4-dinitro-6-(1-propylpentyl) phenol	ND	ND
2,4-dinitro-6-(1-ethylhexyl) phenol	ND	ND
2,4-dinitro-6-(1-methylheptyl) phenol	0.23	0.35
unknown at 6.4 min	0.02	0.03
2,6-dinitro-4-(1-propylpentyl) phenyl crotonate cis	ND	ND
2,6-dinitro-4-(1-ethylhexyl) phenyl crotonate cis	ND	ND
2,6-dinitro-4-(1-methylheptyl) phenyl crotonate cis	ND	ND
2,6-dinitro-4-(1-propylpentyl) phenyl crotonate trans	0.07	0.03
2,6-dinitro-4-(1-ethylhexyl) phenyl crotonate trans	0.03	0.007
2,6-dinitro-4-(1-methylheptyl) phenyl crotonate trans	ND	ND
2,4-dinitro-6-(1-propylpentyl) phenyl crotonate cis	ND	ND
2,4-dinitro-6-(1-ethylhexyl) phenyl crotonate cis	ND	ND
2,4-dinitro-6-(1-propylpentyl) phenyl crotonate trans	ND	ND
2,4-dinitro-6-(1-ethylhexyl) phenyl crotonate trans (LC)	1.4	1.3
Other	1.45	1.283
Mass Balance (Wt. percent)	101.1	101.0

Not Detected (ND)

Limit of Detection (LOD) = 0.02 percent

5

Testing on plants:

The composition of Example 1 was evaluated for fungicidal efficacy in a greenhouse on 9 diseases caused by plant pathogenic fungi (TABLE II). The plant cultivars used in this

study were Ugni Blanc (grape), Bovowinka (crab apple), HH88 (sugar beet), M-9 (rice) and Yuma (wheat). Broad leaf plants and rice were grown in a soil-less peat-based potting mixture (Metromix), and wheat was grown in a 50/50 mix of mineral soil and Metromix. The wheat was grown at a temperature of 20°C and all the other plants were grown at 25°C. Apple seeds were soaked in water, incubated for a short time in 200 ppm solution of benlate and washed again with water. The moist seeds were stored in a closed container at 7°C (45 °F) and washed once a week with water until the radicle had emerged 3 mm (approximately 30 days). The seeds were then planted and seedlings with 4-6 leaves were typically ready for inoculation 3 weeks later.

High volume application studies.

The composition of Example 1 was applied to seedlings at 100, 50, 25, and 12.5 ppm using a high volume spray application. The composition was formulated in water and serial dilutions made and then brought up to volume by adding water solution. Plants were sprayed to run off with 15 mL of solution on a turn-table sprayer with two opposing 4JAUPM air atomization nozzles (Spraying Systems) at a pressure of 138 kPa. Volume of spray solution on a per hectare basis is approximately 4000 L/ha. Plants were inoculated with the pathogens 1 day after application of the composition of Example 1 (1 day protectant test), incubated in a dew chamber for 24 hours with 100 percent relative humidity and then moved to an appropriate environment for disease expression for the duration of the test. Three replicate pots were sprayed for each pathogen/composition combination. Plants were evaluated for percent disease incidence 7-14 days after inoculation and the average percent disease control determined. Results are listed in TABLES III and IV.

Pathogen propagation and host inoculation.

Plants were inoculated with various pathogens 1 day after application of the composition of EXAMPLES 1, 4-5 (protectant tests) and 2 days before application for EXAMPLES 6-

7 (curative tests). For all wheat trials, compositions were applied at growth stage 1.2, when the second leaf was expanded to ½ of its final size (12 days after seeds were first watered). Information on the growth stages of other plant species at the time composition application and on the propagation and inoculation procedures associated with each pathogen is given below.

TABLE II. Plant pathogens and host plants used in greenhouse evaluation.			
Host Plant	Disease Name	Causal Organism	Bayer Code
Wheat	Powdery Mildew	<i>Erysiphe graminis</i> f.sp <i>tritici</i>	ERYSGT
Wheat	Brown Rust	<i>Puccinia recondita</i>	PUCCRT
Apple	Scab	<i>Venturia inaequalis</i>	VENTIN
Grapes	Powdery mildew	<i>Uncinula necator</i>	UNCINE
Cucumbers	Powdery mildew	<i>Erysiphe cichoracearum</i>	ERYSCI

ERYSGT: Wheat seedlings were infected with fresh spores from the obligate pathogen ERYSGT by shaking heavily infected wheat plants over them. Plants that had been dusted with ERYSGT spores were incubated in the greenhouse at 22°C until disease symptoms had fully developed (usually 7 days).

ERYSCI: Cucumber seedlings were infected with fresh spores from the obligate pathogen ERYSCI by spraying a spore suspension on them. Plants were sprayed to run-off. The seedlings were incubated in the greenhouse at 22°C until disease symptoms had fully developed (usually 10 days).

PUCCRT: Spores of the obligate pathogen PUCCRT were collected from infected plants with a vacuum apparatus and stored at 4°C. Approximately 0.1 g of fresh spores (stored at 4°C for less than 30 days) was mixed with several drops of Tween 20. The thick spore

paste was diluted to 100 ml with water and sprayed to run-off on wheat seed seedlings. Plants inoculated with Puccinia were kept in a 20°C dew chamber overnight and then transferred to a 20°C growth chamber where symptoms developed in 8-9 days.

- 5 PYRRIOR: Inoculum was produced on PDA grown under white light at 24°C for 10-12 days. The mycelium was scraped off in water, minced in a blender and expressed through several layers of cheesecloth. Spore concentration was adjusted to 50,000/ml and three large drops of Tween 20 were then added for each 100 ml of volume. Twelve day-old rice plants (second leaf fully expanded) were sprayed to run-off with the aqueous spore
10 suspension, placed in a 24°C dew chamber for 24 hr and then moved to a 24°C growth chamber until disease symptoms had fully developed (7 days).

- VENTIN: Conidia that were previously harvested (up to 6 months earlier) by washing infected leaves in tap water and then freezing the solution at a concentration of 400,000
15 spores/ml, were used for inoculations. Apple seedlings were sprayed to run-off with the VENTIN inoculum, incubated 24 hr in a 20°C dew chamber and then moved to an 18°C growth chamber until disease symptoms developed (8 days). Disease severity was evaluated only on the two youngest leaves at the time of inoculation because older leaves become highly resistant to infection with this pathogen.

20

UNCINE: Six week-old grape plants, pruned to maintain two fully expanded leaves, were infected with fresh spores from this obligate pathogen by shaking infected plants over them. Plants that had been dusted with UNCINE spores were incubated in the greenhouse at 22°C until disease symptoms developed (usually 10 days).

25

TABLE III. High volume test with three plant pathogenic fungi in a high volume screen.

Pathogen	Rate (ppm)	Percent disease control
ERYSCI	200	**
PUCCRT	200	***
PYRIOR	200	***

* is 0-49 percent control of disease

** is 50-79 percent control of disease

*** is 80-100 percent control of disease

TABLE IV. Percent Disease Control

Material	Rate ppm	ERYSGT	PUCCRT	UNCINE	VENTIN ₁₀
Ex. 1	100	***	***	***	***
Ex. 1	50	***	**	***	**
Ex. 1	25	NT	*	***	**
Ex. 1	12.5	NT	*	***	** 15
Azoxystrobin	25	NT	NT	***	NT
Fenarimol	25	NT	NT	NT	***
untreated	*	*	*	*	* 20

* is 0-49 percent control of disease

** is 50-79 percent control of disease

*** is 80-100 percent control of disease

NT is not tested

SYNERGY EXAMPLES

Combinations of Example 1 and named fungicides show synergistic activity on phytopathogenic disease of crop plants. A synergistic effect exists whenever the action of an active ingredient combination is greater than the sum of the action of the individual components.

The action to be expected E for a given active ingredient combination obeys the so-called COLBY formula and can be calculated as follows (COLBY, S.R. "Calculating synergistic and antagonistic responses of herbicide combination". Weeds, Vol. 15, pages 20-22; 1967)

5

X= percent action by active ingredient I using p ppm of active ingredient

Y = percent action by active ingredient II using q ppm of active ingredient

According to Colby the expected additive action of active ingredients I+II using p+q ppm of active ingredients is

10
$$E = X+Y - (X*Y)/100$$

If the action actual observed (O) is greater than the expected action (E), then the action of the combination is super additive, that is, there is a synergistic effect.

Alternatively the synergistic action may also be determined from the dose response curves using the so called WADLEY method. (EPPO – Bulletin 16, 1986, 651-657)

15

EXAMPLE 4.

Synergistic combination of Example 1 and Mancozeb.

Protectant activity of combinations against wheat brown rust (PUCCRT).

1 day protectant glasshouse studies.

TABLE V.

	Mixing partner Rate (ppm)	Ex. 1 (ppm)	Observed Percent CONTROL	Predicted Percent CONTROL	SF Colby
Ex. 1	none	75.00	68.63		
Ex. 1	none	50.00	33.33		
Ex. 1	none	25.00	11.76		
Ex. 1	none	12.50	0.00		
Mancozeb	200.00	none	88.24		
Mancozeb	100.00	none	58.82		
Mancozeb	50.00	none	5.88		
Mancozeb	25.00	none	0.00		
Mancozeb	100.00	12.50	82.35	58.82	1.40
Mancozeb	50.00	12.50	52.94	5.88	9.00
Mancozeb	25.00	12.50	25.49	0.00	25.49
Mancozeb	100.00	25.00	86.27	63.67	1.36
Mancozeb	50.00	25.00	80.39	16.96	4.74
Mancozeb	25.00	25.00	33.33	11.76	2.83
Mancozeb	100.00	50.00	97.65	72.55	1.35
Mancozeb	50.00	50.00	86.67	37.25	2.33
Mancozeb	25.00	50.00	76.47	33.33	2.29
Mancozeb	100.00	75.00	97.25	87.08	1.12
Mancozeb	50.00	75.00	94.12	70.47	1.34
Mancozeb	25.00	75.00	76.47	68.63	1.11

Efficacy of the combinations of mancozeb and Example 1 with ratios ranging from 8:1 to 3:1 showed synergistic activity.

5

EXAMPLE 5. Synergistic combination of Example 1 and Myclobutanil.

Protectant activity of combinations against wheat brown rust (Puccinia).

1 day protectant glasshouse studies.

TABLE VI

	Mixing partner rate (ppm)	Ex. 1 (ppm)	Observed Percent CONTROL	Predicted Percent CONTROL	SF Colby
Ex. 1	none	75.00	68.63		
Ex. 1	none	50.00	33.33		
Ex. 1	none	25.00	11.76		
Ex. 1	none	12.50	0.00		
Myclobutanil	18.75	none	84.31		
Myclobutanil	12.50	none	74.51		
Myclobutanil	6.25	none	27.45		
Myclobutanil	3.13	none	0.00		
Myclobutanil	12.50	12.50	72.55	74.51	0.97
Myclobutanil	6.25	12.50	56.86	27.45	2.07
Myclobutanil	3.13	12.50	11.76	0.00	11.76
Myclobutanil	12.50	25.00	84.31	77.51	1.09
Myclobutanil	6.25	25.00	70.59	35.99	1.96
Myclobutanil	3.13	25.00	45.10	11.76	3.83
Myclobutanil	12.50	50.00	98.04	83.01	1.18
Myclobutanil	6.25	50.00	96.47	51.63	1.87
Myclobutanil	3.13	50.00	92.94	33.33	2.79
Myclobutanil	6.25	75.00	92.94	77.24	1.20
Myclobutanil	3.13	75.00	84.31	68.63	1.23

Synergistic activity for Sterol biosynthesis inhibition; C14-demethylase inhibition (SBI class I), illustrative example myclobutanil. Efficacy of the combinations of myclobutanil and Example 1 with ratios ranging from 1:24 to 1:4 showed synergistic activity.

EXAMPLE 6. Synergistic combination of Example 1 and Spiroxamine.

Curative activity of combinations against wheat powdery mildew (ERYSGT).

2 day curative glasshouse studies.

TABLE VII

	Mixing partner rate (ppm)	Ex. 1 (ppm)	Observed Percent CONTROL	Predicted Percent CONTROL	SF Colby
Ex. 1	none	75.00	23.00		
Spiroxamine	12.50	none	15.00		
Spiroxamine	12.50	75.00	65.00	34.75	1.88

Confirmation of synergistic activity for Sterol biosynthesis inhibition; D14 reductase and D8 isomerase in sterol biosynthesis (SBI class II), illustrative example spiroxamine. Efficacy of the combinations of spiroxamine and Example 1 with a 1:4 ratio showed synergistic activity.

5

EXAMPLE 7. Synergistic combination of Example 1 and Azoxystrobin.

Curative activity of combinations against wheat powdery mildew (ERYSGT).

2 day curative glasshouse studies.

10

TABLE VIII.

	Mixing partner rate (ppm)	Ex. 1 (ppm)	Observed Percent CONTROL	Predicted Percent CONTROL	SF Colby
Ex. 1	none	75.00	23.00		
Azoxystrobin	6.25	none	13.00		
Azoxystrobin	12.50	none	15.00		
Azoxystrobin	25	none	15		
Azoxystrobin	75	none	52		
Azoxystrobin	100	none	79		
Azoxystrobin	6.25	75	42	33.41	1.27
Azoxystrobin	12.50	75	54	34.91	1.54
Azoxystrobin	25	75	73	34.91	2.09
Azoxystrobin	75	75	97	63.02	1.54
Azoxystrobin	100	75	98	83.77	1.17

Synergistic activity for fungicides acting as respiration inhibitors at complex III;

15 cytochrome bc1 at Qo site. (QoI fungicides); illustrative example azoxystrobin. Efficacy of the combinations of azoxystrobin and Example 1 with ratios ranging from 1:12 to 4:3 showed synergistic activity.

800828881

2005295774 17 Feb 2012

As used herein, except where the context requires otherwise, the term "comprise" and variations of the term, such as "comprising", "comprises" and "comprised", are not intended to exclude other additives, components, integers or steps.

5 Reference to any prior art in the specification is not, and should not be taken as, an acknowledgment, or any form of suggestion, that this prior art forms part of the common general knowledge in Australia or any other jurisdiction or that this prior art could reasonably be expected to be ascertained, understood and regarded as relevant by a person skilled in the art.

800828801

2005295774 17 Feb 2012

The claims defining the invention are as follows:

1. An isomeric composition comprising dinitro-octylphenyl esters, wherein 2,6-dinitro-4-(1-propylpentyl)phenyl ester is present in an amount of less than 0.1 weight percent, based on the total weight of the isomeric composition.
- 5 2. The isomeric composition of Claim 1 comprising:
 from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-methylheptyl)phenyl ester,
 from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-ethylhexyl)phenyl ester,
 from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-propylpentyl)phenyl ester,
 from 0.1 to 99.9 weight percent 2,6-dinitro-4-(1-methylheptyl)phenyl ester,
 10 from 0.1 to 99.9 weight percent 2,6-dinitro-4-(1-ethylhexyl)phenyl ester, and
 less than 0.1 weight percent 2,6-dinitro-4-(1-propylpentyl)phenyl ester,
 based on the total weight of the isomeric composition.
3. The isomeric composition of Claim 1 comprising:
 -at least 95 weight percent of 2,4-dinitro-6-(1-methylheptyl)phenyl ester,
 15 -less than 3 weight percent of 2,4-dinitro-6-(1-ethylhexyl)phenyl ester, and
 -less than 0.1 weight percent of 2,6-dinitro-4-(1-propylpentyl)phenyl ester,
 based on the total weight of the isomeric composition.
4. The isomeric composition of any one of the preceding claims wherein each ester of the isomeric composition is a crotonate.
- 20 5. A fungicidal composition, useful for protecting a plant against attack by a phytopathogenic organism and/or treatment of a plant infested by a phytopathogenic organism, comprising
 - I) the isomeric composition of Claim 1, and
 - II) a phytologically acceptable carrier material.
- 25 6. The fungicidal composition of Claim 5 wherein the isomeric composition comprises:
 from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-methylheptyl)phenyl ester,
 from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-ethylhexyl)phenyl ester,
 from 0.1 to 99.9 weight percent 2,4-dinitro-6-(1-propylpentyl)phenyl ester,
 from 0.1 to 99.9 weight percent 2,6-dinitro-4-(1-methylheptyl)phenyl ester,
 30 from 0.1 to 99.9 weight percent 2,6-dinitro-4-(1-ethylhexyl)phenyl ester, and

000628881

2005295774 17 Feb 2012

less than 0.1 weight percent 2,6-dinitro-4-(1-propylpentyl)phenyl ester,
based on the total weight of the isomeric composition.

7. The fungicidal composition of Claim 5 wherein, the isomeric composition is
 - at least 95 weight percent of 2,4-dinitro-6-(1-methylheptyl)phenyl ester,
 - less than 3 weight percent of 2,4-dinitro-6-(1-ethylhexyl)phenyl ester, and
 - less than 0.1 weight percent of 2,6-dinitro-4-(1-propylpentyl)phenyl ester,
 based on the total weight of the isomeric composition.
8. The fungicidal composition of any one of Claims 5 to 7 wherein each ester of the isomeric composition is a crotonate.
9. A synergistic fungicidal composition comprising the isomeric composition of any one of the preceding claims and at least one other fungicide.
10. The synergistic fungicidal composition of Claim 9 wherein the at least one other fungicide is selected from the group consisting of: QoI fungicides, strobilurins, sterol biosynthesis inhibitors, benzophenones, quinoxifen or dithiocarbamates.
11. The synergistic fungicidal composition of Claim 10 wherein the at least one other fungicide is selected from QoI fungicides.
12. The synergistic fungicidal composition of Claim 10 wherein the at least one other fungicide is selected from strobilurins.
13. The synergistic fungicidal composition of Claim 10 wherein the at least one other fungicide is selected from sterol biosynthesis inhibitors.
14. The synergistic fungicidal composition of Claim 10 wherein the at least one other fungicide is selected from benzophenones.
15. The synergistic fungicidal composition of Claim 10 wherein the at least one other fungicide is quinoxifen.
16. The synergistic fungicidal composition of Claim 10 wherein the at least one other fungicide is selected from dithiocarbamates.

800828561

2005295774 17 Feb 2012

17. A method for the control or prevention of fungal attack comprising applying to the plant, foliage, locus of the fungus, or to a locus in which the infestation is to be prevented, a fungicidal effective amount of the isomeric composition of any one of Claims 1 to 4 or the synergistic fungicidal composition of any one of Claims 9 to 16.
- 5 18. The method according to Claim 17, wherein the isomeric composition comprises:
-at least 95 weight percent of 2,4-dinitro-6-(1-methylheptyl)phenyl ester,
-less than 3 weight percent of 2,4-dinitro-6-(1-ethylhexyl)phenyl ester, and
-less than 0.1 weight percent of 2,6-dinitro-4-(1-propylpentyl)phenyl ester,
based on the total weight of the isomeric composition.
- 10 19. The method according to Claim 17 wherein each ester of the isomeric composition is a crotonate.
20. An isomeric composition according to Claim 1, substantially as hereinbefore described with reference to any one of the examples.
- 15 21. A synergistic fungicidal composition according to Claim 9, substantially as hereinbefore described with reference to any one of the examples.