



US00RE44971E

(19) **United States**
 (12) **Reissued Patent**
Wan et al.

(10) **Patent Number:** **US RE44,971 E**
 (45) **Date of Reissued Patent:** **Jun. 24, 2014**

(54) **METHOD FOR SELECTION OF TRANSFORMED CELLS**

(71) Applicant: **Monsanto Technology LLC**, St. Louis, MO (US)

(72) Inventors: **Yuenchun Wan**, Madison, WI (US);
Ronald J. Brinker, Ellisville, MO (US);
Paul C. C. Feng, Wildwood, MO (US)

(73) Assignee: **Monsanto Technology LLC**, St. Louis, MO (US)

(21) Appl. No.: **13/855,631**

(22) Filed: **Apr. 2, 2013**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **7,851,670**
 Issued: **Dec. 14, 2010**
 Appl. No.: **11/758,656**
 Filed: **Jun. 5, 2007**

U.S. Applications:

(60) Provisional application No. 60/811,190, filed on Jun. 6, 2006.

(51) **Int. Cl.**
C12N 15/82 (2006.01)
A01H 5/00 (2006.01)
C07H 21/04 (2006.01)
C12N 5/04 (2006.01)

(52) **U.S. Cl.**
 USPC **800/278; 800/298; 800/300; 800/312;**
536/23.2; 536/23.7; 435/415

(58) **Field of Classification Search**
 None
 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,816,403 A 3/1989 Roy
 5,094,945 A 3/1992 Comai
 5,188,642 A 2/1993 Shah et al.
 5,254,799 A 10/1993 De Greve et al.
 5,362,865 A 11/1994 Austin
 5,445,962 A 8/1995 Atallah et al.
 5,463,175 A 10/1995 Barry et al.
 5,491,084 A 2/1996 Chalfie et al.
 5,561,236 A 10/1996 Leemans et al.
 5,627,061 A 5/1997 Barry et al.
 5,633,435 A 5/1997 Barry et al.
 5,633,437 A 5/1997 Bernasconi et al.
 5,656,422 A 8/1997 Crawford et al.
 5,659,122 A 8/1997 Austin
 5,670,454 A 9/1997 Grossmann et al.
 5,728,925 A 3/1998 Herrera-Estrella et al.
 5,850,019 A 12/1998 Maiti et al.
 5,939,602 A 8/1999 Volrath et al.
 6,040,497 A 3/2000 Spencer et al.
 6,146,826 A 11/2000 Chalfie et al.
 6,268,549 B1 7/2001 Sailland et al.
 6,628,549 B2 7/2001 Sailland et al.
 6,376,754 B1 4/2002 Schillinger et al.
 6,414,222 B1 7/2002 Gengenbach et al.
 6,586,367 B2 7/2003 Lee et al.

6,613,963 B1 9/2003 Gingera et al.
 7,022,896 B1 4/2006 Weeks et al.
 RE39,247 E 8/2006 Barry et al.
 7,105,724 B2 9/2006 Weeks et al.
 7,230,163 B2 6/2007 Becton et al.
 7,385,106 B2 6/2008 Stein et al.
 7,405,074 B2 7/2008 Castle et al.
 7,405,347 B2 7/2008 Hammer et al.
 7,407,913 B2 8/2008 Lee et al.
 7,429,691 B2 9/2008 Zhang et al.
 7,462,481 B2 12/2008 Castle et al.
 7,622,641 B2 11/2009 McCutchen et al.
 7,714,188 B2 5/2010 Castle et al.
 7,786,041 B2 8/2010 Rosinger et al.
 7,812,224 B2 10/2010 Weeks et al.
 7,838,729 B2 11/2010 Feng et al.
 7,838,733 B2 11/2010 Wright et al.
 7,851,670 B2 12/2010 Wan et al.
 7,855,326 B2 12/2010 Feng et al.
 7,884,262 B2 2/2011 Clemente et al.
 7,939,721 B2 5/2011 Arnevik et al.
 8,022,272 B2 9/2011 Heim et al.
 8,084,666 B2 12/2011 Feng et al.
 8,119,380 B2 2/2012 Weeks et al.
 8,207,092 B2 6/2012 Bhatti et al.
 8,420,888 B2 4/2013 Feng et al.
 2001/0046945 A1 11/2001 Lee et al.
 2002/0168680 A1 11/2002 Barry et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CA 2165036 6/1996
 CA 2165036 A1 6/1996

(Continued)

OTHER PUBLICATIONS

U.S. Appl. No. 60/811,190, filed Jun. 6, 2006, Feng et al.
 U.S. Appl. No. 60/811,276, filed Jun. 6, 2006, Feng et al.
 U.S. Appl. No. 60/884,854, filed Jan. 12, 2007, D'Ordine et al.
 U.S. Appl. No. 12/440,173, filed Mar. 5, 2009, Bhatti et al.
 U.S. Appl. No. 12/875,747, filed Sep. 3, 2010, Weeks et al.
 U.S. Appl. No. 12/914,901, filed May 26, 2011, Feng et al.
 U.S. Appl. No. 12/942,905, filed Jun. 23, 2011, Feng et al.
 Al-Khatib et al., "Foliar absorption and translocation of dicamba from aqueous solution and dicamba-treated soil deposits," *Weed Technology*, 6:57-61, 1992.
 "Banvel Herbicide," In: Crop Protection Reference, 11th Edition, pp. 1803-1821, 1995.
 Baker, "Response of cotton (*Gossypium hirsutum*) to preplant-applied hormone-type herbicides," *Weed Technology*, 7:150-153, 1993.

(Continued)

Primary Examiner — Eileen B O Hara

(74) *Attorney, Agent, or Firm* — Dentons US LLP; Pamela Sisson

(57) **ABSTRACT**

The invention provides methods for the selection of transgenic cells. The invention relates to the unexpected finding that cells expressing a gene conferring tolerance to auxin-like herbicides such as dicamba may be directly selected from non-transgenic cells using auxin-like herbicides as a selective agent. In this manner, plants exhibiting tolerance to auxin-like herbicides can be directly produced without the need for separate selectable markers.

10 Claims, 9 Drawing Sheets

(56)

References Cited

U.S. PATENT DOCUMENTS

2003/0041357	A1	2/2003	Jepson et al.
2003/0083480	A1	5/2003	Castle et al.
2003/0115626	A1	6/2003	Weeks et al.
2003/0135879	A1	7/2003	Weeks et al.
2004/0082770	A1	4/2004	Castle et al.
2004/0097373	A1	5/2004	Lee et al.
2004/0177399	A1	9/2004	Hammer et al.
2005/0235379	A1	10/2005	Luo et al.
2006/0019828	A1	1/2006	Becher et al.
2006/0111240	A1	5/2006	Hacker et al.
2006/0235215	A1	10/2006	Cooper
2007/0010398	A1	1/2007	Rosinger et al.
2007/0079393	A1	4/2007	McCutchen et al.
2008/0015110	A1	1/2008	Clemente et al.
2008/0119361	A1	5/2008	Feng et al.
2008/0120739	A1	5/2008	Wan et al.
2008/0305952	A1	12/2008	Arnevik et al.
2009/0029861	A1	1/2009	Feng et al.
2009/0081760	A1	3/2009	D'Ordine et al.
2009/0093366	A1	4/2009	Wright et al.
2009/0105077	A1	4/2009	Bhatti et al.
2010/0279866	A1	11/2010	Bhatti et al.
2011/0061137	A1	3/2011	Weeks et al.
2011/0126307	A1	5/2011	Feng et al.
2011/0152096	A1	6/2011	Feng et al.
2011/0203017	A1	8/2011	Wright et al.
2011/0245080	A1	10/2011	Arnevik et al.
2012/0151620	A1	6/2012	Feng et al.
2013/0198886	A1	8/2013	Feng et al.

FOREIGN PATENT DOCUMENTS

CN	1236394	A	11/1999
CN	1332800	A	1/2002
CN	1541270	A	10/2004
DE	199 19 993	A1	11/2000
WO	WO 97/41228		11/1997
WO	WO 98/20144		5/1998
WO	WO 98/39419		9/1998
WO	WO 98/45424		10/1998
WO	WO 98/45424	A1	10/1998
WO	WO 00/08936		2/2000
WO	WO 00/08939		2/2000
WO	WO 00/26371		5/2000
WO	WO 00/029596		5/2000
WO	WO 02/26995		4/2002
WO	WO 02/068607		9/2002
WO	WO 2003/006660		1/2003
WO	WO 03/024221		3/2003
WO	WO 03/034813		5/2003
WO	WO 03/034813	A2	5/2003
WO	WO 03/096811		11/2003
WO	WO 2004/009761		1/2004
WO	WO 2004/074443		9/2004
WO	WO/2004/101797		11/2004
WO	WO 2005/003362		1/2005
WO	WO 2005/107437		11/2005
WO	WO 2005/107437	A2	11/2005
WO	WO 2006/065815		6/2006
WO	WO 2007/143690		12/2007
WO	WO 2007/143690	A2	12/2007
WO	WO 2007/146706		12/2007
WO	WO 2007/146706	A2	12/2007
WO	WO 2008/048964		4/2008
WO	WO 2008/048964	A2	4/2008
WO	WO 2008/051633		5/2008
WO	WO 2008/051633	A2	5/2008
WO	WO 2008/105890		9/2008
WO	WO 2008/105890	A2	9/2008

OTHER PUBLICATIONS

Batie et al., "Purification and characterization of phthalate oxygenase and phthalate oxygenase reductase from *Pseudomonas cepacia*," *J. of Bio. Chem.*, 262(4):1510-1518, 1987.

Batie et al., "Phthalate dioxygenase reductase and related flavin-iron-sulfur containing electron transferases," In: Chemistry and Biochemistry of Flavoproteins, Muller (Ed.), CRC Press, Boca Raton, FL, pp. 543-556, 1992.

Behrens et al., "Dicamba resistance: enlarging and preserving biotechnology-based weed management strategies," *Science*, 316:1185-1188, 2007.

Bernhardt et al., "A 4-methoxybenzoate O-demethylase from *Pseudomonas putida*. A new type of monooxygenase system," *Eur. J. Biochem.*, 57(1):241-256, 1975.

Bevan, "Structure and transcription of the nopaline synthase gene region of T-DNA," *Nucleic Acids Res.*, 11(2):369-385, 1983.

Buchanan-Wollaston, et al., Detoxification of the herbicide Dalapon by transformed plants, *J. Cell. Biochem.*, 13D, Abstract No. M503, p. 330, 1989.

Butler et al., "Structure-function analysis of the bacterial aromatic ring-hydroxylating dioxygenases," *Advances in Microbial Physiology*, 38:47-85, 1997.

Comai et al., "Expression in plants of a mutant *aroA* gene from *Salmonella typhimurium* confers tolerance to glyphosate," *Nature*, 317:741-744, 1985.

Cork et al., "Microbial transformations of herbicides and pesticides," *Adv. Appl. Microbiology*, 36:1-67, 1991.

Cork et al., "Detection, isolation, and stability of megaplasmid-encoded chloroaromatic herbicide-degrading genes within *Pseudomonas* species," *Adv. Appl. Microbiol.*, 40:289-321, 1995.

Coruzzi, et al., "Nucleotide Sequences of Two Pea cDNA Clones Encoding the Small Subunit of Ribulose 1.5-Bisphosphate Carboxylase and the Major Chlorophyll a/b-binding Thylakoid Polypeptide," *J. of Biological Chem.* 258(3):1399-1402, 1983.

Coruzzi, et al., "Tissue-specific and light-regulated expression of pea nuclear gene encoding the small subunit of ribulose-1,5-bisphosphate carboxylase," *EMBO J.*, (8):1671-1679, 1984.

Creissen et al., "Molecular characterization of glutathione reductase cDNAs from pea (*Pisum sativum* L.)," *Plant J.*, 2(1):129-131, 1992.

Creissen et al., "Simultaneous targeting of pea glutathione reductase and of a bacterial fusion protein to chloroplasts and mitochondria in transgenic tobacco," *Plant J.*, 8:167-175, 1995.

De Block et al., "Engineering herbicide resistance in plants by expression of a detoxifying enzyme," *EMBO J.*, 6(9):2513-2518, 1987.

Dehmel et al., "Cloning, nucleotide sequence and expression of the gene encoding a novel dioxygenase involved in metabolism of carboxydiphenyl ethers in *Pseudomonas pseudoalcaligenes* POB310," *Arch. Microbiol.*, 163:35-41, 1995.

Depicker et al., "Nopaline synthase: transcript mapping and DNA sequence," *J. Mol. Appl. Genet.*, 1:561-573, 1982.

Desvaux et al., "Genomic analysis of the protein secretion systems in *Clostridium acetobutylicum* ATCC 824," *Biochimica et Biophysica Acta*, 1745:223-253, 2005.

Eckes et al., "Isolation and characterization of a light-inducible, organ-specific gene from potato and analysis of its expression after tagging and transfer into tobacco and potato shoots," *Mol. Gen. Genet.*, 205:14-22, 1986.

Fogarty et al., "Microbiological degradation of the herbicide dicamba," *J. of Industrial Microbiology*, 14:365-370, 1995.

Fukumori et al., "Purification and characterization of 2,-dichlorophenoxyacetate/ α -ketoglutarate dioxygenase," *J. Biol. Chem.*, 268:24311-24317, 1993.

Gasser et al., "Structure, expression, and evolution of the 5-enolpyruvylshikimate-3-phosphate synthase genes of petunia and tomato," *J. Biol. Chem.*, 263:4280-4287, 1988.

GenBank Accession No. E01312, Nov. 4, 2005.

GenBank Accession No. V00087, Mar. 18, 1996.

GenBank Accession No. AY786443, dated Jun. 29, 2005.

GenBank Accession No. AAT48249, Jun. 25, 2004.

Gibson et al., "Aromatic hydrocarbon dioxygenases in environmental biotechnology," *Current Opinion in Biotechnology*, 11:236-243, 2000.

Gurbel et al., Active site-structure of Rieske-type proteins: electron nuclear double resonance studies of isotopically labeled phthalate

(56)

References Cited

OTHER PUBLICATIONS

- dioxygenase from *Pseudomonas cepacia* and Rieske protein from rhodobacter capsulatus and molecular modeling studies of a, Biochemistry 1996 Rieske center.
- Hajdukiewicz et al., "The small, versatile pPZP family of agrobacterium binary vectors for plant transformation," *Plant Mol. Biol.*, 25:989-994, 1994.
- Hamill et al., "Weed control in glufosinate-resistant corn (*Zea mays*)," *Weed Technology* 14(3):578-585, 2000.
- Herman et al., "A three-component dicamba O-demethylase from *Pseudomonas maltophilia*, Strain Di-6," *J. of Biological Chemistry*, 280(26):24759-24767, 2005.
- Khalil et al., "Plasmid-mediated catabolism of dicamba by *Pseudomonas* species strain PXM," *Microbios*, 102:183-191, 2000.
- Klee et al., "Cloning of an *Arabidopsis thaliana* gene encoding 5-enolpyruvylshikimate-3-phosphate synthase: sequence analysis and manipulation to obtain glyphosate-tolerant plants," *Mol. Gen. Genet.*, 210:437-442, 1987.
- Krueger et al., "Isolation and identification of microorganisms for the degradation of dicamba," *J. Agric. Food Chem.*, 37:534-538, 1989.
- Krueger et al., "Use of dicamba-degrading microorganisms to protect dicamba susceptible plant species," *J. of Agri. and Food Chem.*, 39(5):1000-10003, 1991.
- Magnusson et al., "Tolerance of soybean (*Glycine max*) and sunflower (*Helianthus annuus*) to fall-applied dicamba," *Weed Sci.*, 35:846-852, 1987.
- Markus et al., "Purification and some properties of component A of the 4-chlorophenylacetate 3,4-dioxygenase from *Pseudomonas* species strain CBS," *J. of Biol. Chem.*, 261(27):12883-12888, 1986.
- Mason et al., "The electron-transport proteins of hydroxylating bacterial dioxygenases," *Ann. Rev. of Microbiology*, 46:277-305, 1992.
- Mazur et al., "Sequence of a genomic DNA clone for the small subunit of ribulose bis-phosphate carboxylase-oxygenase from tobacco," *Nucleic Acids Res.*, 13(7):2373-2386, 1985.
- Misawa et al., "Functional expression of the *Erwinia uredovora* carotenoid biosynthesis gene *crtI* in transgenic plants showing an increase of beta-carotene biosynthesis activity and resistance to the bleaching herbicide norflurazon," *Plant J.*, 4:833-840, 1993.
- Misawa et al., "Expression of an *Erwinia phytoene* desaturase gene not only confers multiple resistance to herbicides interfering with carotenoid biosynthesis but also alters xanthophyll metabolism in transgenic plants," *Plant J.*, 6:481-489, 1994.
- Mueller et al., *Weed Technology* 19:924-933, 2005.
- Padgett et al., "Development, identification and characterization of a glyphosate-tolerant soybean line," *Crop Sci.*, 35:1451-1461, 1995.
- Peniuk et al., "Physiological investigations into the resistance of a wild mustard (*Sinapis arvensis* L.) biotype to auxinic herbicides," *Weed Research*, 33:431-440, 1993.
- Sarpe et al., "Researches on resistance of maize hybrids and inbred lines to the herbicides based on 2,4-D and dicamba," *Fragmenta Herbologica Jugoslavica*, 16(1-2):299-305, 1987.
- Sathasivan et al., "Nucleotide sequence of a mutant acetolactate synthase gene from an imidazolinone-resistant *Arabidopsis thaliana* var. Columbia," *Nucl. Acids Res.*, 18(8):2188, 1990.
- Schroeder et al., "Soft red winter wheat (*Triticum aestivum*) response to dicamba and dicamba plus 2,4-D," *Weed Technology*, 3:67-71, 1989.
- Spencer et al., "Segregation of transgenes in maize," *Plant Molecular Biology*, 18:201-210, 1992.
- Sprague, "Avoid herbicide spray tank contamination," *IPM News*, ipmnews.msu.edu/fieldcrop/tabid/56, Mar. 24, 2010.
- Stalker et al., "Herbicide resistance in transgenic plants expressing a bacterial detoxification gene," *Science*, 242:419-423, 1988.
- Streber et al., "Transgenic tobacco plants expressing a bacterial detoxifying enzyme are resistant to 2,4-D," *Bio/Technology*, 7:811, 1989.
- Svab et al., "Aminoglycoside-3'-adenyltransferase confers resistance to spectinomycin and streptomycin in *Nicotiana tabacum*," *Plant Mol. Biol.*, 14:197, 1990.
- Thompson et al., "Soybean tolerance to early preplant applications of 2,4-D ester, 2,4-D amine, and dicamba," *Weed Technology*, 21:882-885, 2007.
- Treadaway-Ducar et al., "Site-specific weed management in corn (*Zea mays*)," *Weed Technology* 17(4):711-717, 2003.
- Wang et al., "A three-component enzyme system catalyzes the O demethylation of the herbicide dicamba in *Pseudomonas maltophilia* DI-6," *Applied and Environmental Microbiology*, 63(4):1623-1626, 1997.
- Wang, "Characterization of cellular and enzymatic degradation of dicamba by *Pseudomonas maltophilia*, Strain DI-6," Thesis, University of Nebraska, Aug. 1996.
- Weeks et al., "Characterization of a bacterial system capable of degrading dicamba and evaluation of its potential in the development of herbicide-tolerant crops," *J. of Cellular Biochemistry*, Supplement 18A:91, 1994.
- Zuver et al., "Evaluation of postemergence weed control strategies in herbicide-resistant isolines of corn (*Zea mays*)," *Weed Technology* 29(1):172-178, 2006.
- Office Action regarding U.S. Appl. No. 10/330,662 dated Apr. 18, 2006.
- Interview Summary regarding U.S. Appl. No. 10/330,662, dated Sep. 13, 2006.
- Declaration of Donald P. Weeks regarding U.S. Appl. No. 10/330,662, dated Oct. 18, 2006.
- Terminal filed in U.S. Appl. No. 10/330,662, dated Oct. 18, 2006.
- Amendment and Response to Office Action regarding U.S. Appl. No. 10/330,662, dated Oct. 18, 2006.
- Final Office Action regarding U.S. Appl. No. 10/330,662, dated Jan. 10, 2007.
- Amendment and Response to Office Action regarding U.S. Appl. No. 10/330,662, dated Jul. 9, 2007.
- Amendment regarding U.S. Appl. No. 10/330,662, dated Jul. 20, 2007.
- Office Action regarding U.S. Appl. No. 10/330,662, dated Sep. 21, 2007.
- Declaration of Donald P. Weeks regarding U.S. Appl. No. 10/330,662, dated Feb. 20, 2008.
- Amendment and Response to Office Action regarding U.S. Appl. No. 10/330,662, dated Mar. 20, 2008.
- Office Action regarding U.S. Appl. No. 10/330,662, dated Jul. 9, 2008.
- Amendment and Remarks regarding U.S. Appl. No. 10/330,662, dated Jan. 9, 2009.
- Final Office Action regarding U.S. Appl. No. 10/330,662, dated Apr. 24, 2009.
- Office Action regarding U.S. Appl. No. 11/758,657 dated Sep. 2, 2009.
- Office Action regarding U.S. Appl. No. 11/758,656 dated Sep. 15, 2009.
- Amendment and Remarks regarding U.S. Appl. No. 10/330,662, dated Sep. 24, 2009.
- Office Action regarding U.S. Appl. No. 11/758,659 dated Nov. 24, 2009.
- Response to Office Action regarding U.S. Appl. No. 11/758,656 dated Dec. 17, 2009.
- Office Action regarding U.S. Appl. No. 11/758,653 dated Dec. 29, 2009.
- Response to Office Action regarding U.S. Appl. No. 11/758,657 dated Jan. 4, 2010.
- Office Action regarding U.S. Appl. No. 10/330,662, dated Jan. 11, 2010.
- Interview Summary regarding U.S. Appl. No. 10/330,662, dated Mar. 19, 2010.
- Amendment and Response to Office Action regarding U.S. Appl. No. 11/758,653, dated Mar. 29, 2010.
- Final Office Action regarding U.S. Appl. No. 11/758,656, dated Apr. 14, 2010.
- Final Office Action regarding U.S. Appl. No. 11/758,657, dated Apr. 14, 2010.
- Office Action regarding U.S. Appl. No. 11/758,660, dated Apr. 28, 2010.

(56)

References Cited

OTHER PUBLICATIONS

- Amendment and Remarks regarding U.S. Appl. No. 10/330,662, dated May 4, 2010.
- Amendment and Response to Office Action regarding U.S. Appl. No. 11/758,659, dated May 24, 2010.
- Final Office Action regarding U.S. Appl. No. 11/758,653, dated Jun. 24, 2010.
- Notice of Allowance regarding U.S. Appl. No. 10/330,662, dated Jul. 12, 2010.
- Response to Final Office Action regarding U.S. Appl. No. 11/758,657, dated Jul. 14, 2010.
- Notice of Allowance regarding U.S. Appl. No. 11/758,659, dated Aug. 3, 2010.
- Declaration of Yuechun Wan Under 37 C.F.R. §1.132, dated Aug. 11, 2010.
- Amendment and Response to Final Office Action regarding U.S. Appl. No. 11/758,653, dated Aug. 16, 2010.
- Notice of Allowance regarding U.S. Appl. No. 11/758,657, dated Sep. 10, 2010.
- Response to Office Action regarding U.S. Appl. No. 11/758,660, dated Sep. 27, 2010.
- Notice of Allowance regarding U.S. Appl. No. 11/758,656, dated Oct. 4, 2010.
- Notice of Allowance regarding U.S. Appl. No. 11/758,653, dated Oct. 5, 2010.
- Office Action dated Dec. 22, 2010, in Honduran Patent Application No. 2009-00795.
- English translation of Office Action dated Dec. 22, 2010, in Honduran Patent Application No. 2009-00795.
- Office Action regarding U.S. Appl. No. 12/914,901 dated Apr. 26, 2011.
- Response to Office Action regarding U.S. Appl. No. 12/914,901 dated Jun. 8, 2011.
- Office Action regarding U.S. Appl. No. 12/942,905 dated Jul. 7, 2011.
- Notice of Allowance regarding U.S. Appl. No. 12/914,901 dated Aug. 25, 2011.
- Response to Office Action regarding U.S. Appl. No. 12/942,905 dated Aug. 30, 2011.
- Supplemental Information Disclosure Statement regarding U.S. Appl. No. 13/751,021, dated Mar. 22, 2013.
- Supplemental Information Disclosure Statement regarding U.S. Appl. No. 12/942,905, dated Mar. 22, 2013.
- Supplemental Information Disclosure Statement regarding U.S. Appl. No. 13/035,902, dated Mar. 22, 2013.
- Supplemental Information Disclosure Statement regarding U.S. Appl. No. 13/774,794, dated Mar. 22, 2013.
- USPTO; Non-final Office Action in U.S. Appl. No. 13/774,794, dated Oct. 4, 2013.
- U.S. Appl. No. 13/751,021, Feng et al.
- U.S. Appl. No. 13/774,794, Feng et al.
- Dernoeden et al., "Fenoxaprop Activity Influenced by Auxin-like Herbicide Application Timing," Department of Agronomy, University of Maryland, *HortScience* 29(12):1518-1519, 1994.
- Jordan et al., "Italian Ryegrass Control with Preplant Herbicides," *The J of Cotton Sciences*; 5:268-274; 2001.
- Morgan et al., "Chemical cotton stalk destruction: what will be the herbicide options in the future," presentation given at the 57th annual Beltwide Cotton Production Conference, Wednesday, Jan. 4, 2012.
- Petition for Determination of Nonregulated Status for Dicamba-tolerant soybean MON87708, pp. 1, 2, and 77, Oct. 10, 2011.
- Terminal Disclaimer filed in U.S. Appl. No. 12/942,905 dated Aug. 30, 2011.
- USPTO; Notice of Non-compliant amendment in U.S. Appl. No. 12/942,905 dated Sep. 14, 2011.
- Response to Notice of Non-compliant amendment in U.S. Appl. No. 12/942,905 dated Sep. 28, 2011.
- USPTO; Notice of Allowance in U.S. Appl. No. 12/942,905 dated Oct. 24, 2011.
- Request for Continued Examination and Amendment in U.S. Appl. No. 12/942,905 dated Jan. 24, 2012.
- USPTO; Notice of Allowance in U.S. Appl. No. 12/942,905 dated Sep. 17, 2013.
- USPTO; Office Action in U.S. Appl. No. 11/758,660 dated Dec. 2, 2010.
- Response to Office Action in U.S. Appl. No. 11/758,660 dated Jan. 13, 2011.
- Terminal Disclaimer in U.S. Appl. No. 11/758,660 dated Jan. 13, 2011.
- USPTO; Notice of Allowance in U.S. Appl. No. 11/758,660 dated Jan. 28, 2011.
- USPTO; Office Action in U.S. Appl. No. 13/035,902 dated Dec. 15, 2011.
- Response to Office Action in U.S. Appl. No. 13/035,902 dated Feb. 7, 2012.
- USPTO; Final Office Action in U.S. Appl. No. 13/035,902 dated Apr. 2, 2012.
- Request for Continued Examination and Amendment in U.S. Appl. No. 13/035,902 dated Jul. 2, 2012.
- USPTO; Office Action in U.S. Appl. No. 13/326,204 dated Sep. 17, 2012.
- Response to Office Action in U.S. Appl. No. 13/326,204 dated Nov. 6, 2012.
- USPTO; Notice of Allowance in U.S. Appl. No. 13/326,204 dated Dec. 10, 2012.
- Third Party Submission in a Published Application Under 37 C.F.R. § 1.99 dated Dec. 2, 2011.
- European Search Report issued in Application No. 12181146.7 dated Nov. 14, 2012.
- Response to Office Action regarding U.S. Appl. No. 13/751,021, filed Nov. 22, 2013.
- Response to Office Action regarding U.S. Appl. No. 13/774,794, filed Dec. 19, 2013.
- USPTO; Notice of Allowance regarding U.S. Appl. No. 13/774,794, issued on Mar. 21, 2014.
- USPTO; Notice of Allowance regarding U.S. Appl. No. 13/751,021, issued on Mar. 12, 2014.
- Carrington et al., "Cap-independent enhancement of translation by a plant potyvirus 5' nontranslated region," *J. of Virology*, 4:1590-1597, 1990.
- Cork and Krueger, "Microbial Transformations of Herbicides and Pesticides," *Adv. Appl. Microbiology*, 36:1-67, 1991.
- De Block et al., "Engineering herbicide resistance in plants by expression of a detoxifying enzyme," *EMBO J.*, 6:2513-2518, 1987.
- Krueger et al., "Isolation and identification of microorganisms for the degradation of dicamba," *J. Agric. Food Chem.*, 37:534, 1989.
- Gurbel et al., "Active site structure of Rieske-type proteins: electron nuclear double resonance of studies of isotopically labeled phthalate dioxygenase from *Pseudomonas cepacia* and Rieske protein from *rhodobacter capsulatus* and molecular modeling studies of a Rieske center," *Biochemistry*, 35(24):7834-7845, 1996 (Abstract).
- Krueger et al., "Use of dicamba-degrading microorganisms to protect dicamba susceptible plant species," *J. of Agri. and Food Chem.*, 39(5):1000-1003, 1991.



FIG. 1




FIG. 2




FIG. 3



FIG. 4




FIG. 5




FIG. 6





FIG. 7

**FIG. 8**




FIG. 9

METHOD FOR SELECTION OF TRANSFORMED CELLS

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.

This application claims the priority of U.S. Provisional Patent Application 60/811,190, filed Jun. 6, 2006, the disclosure of which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The invention relates generally to the field of plant biotechnology. More specifically, the invention relates to methods for selecting transformed plant cells using auxin-like herbicides as a selective agent.

2. Description of the Related Art

Transgenic crops are currently grown on more than 80.0 million hectares world-wide. Improved traits provided by transgenes have significantly increased productivity and in many instances decreased reliance on herbicides and insecticides that can potentially contaminate the environment. However, for transgenic crops to continue to be competitive in the market place, new value-added traits will be required.

In the production of transgenic plants, a particularly important step is the selection of transgenic cells. This is because only a small percentage of cells are typically transformed in any given transformation protocol. The use of a selectable marker gene allows those cells containing a marker gene to be selected away from those that do not. In attempts to stack multiple transgenes in a single plant, this can become particularly difficult, as multiple selectable marker genes are required. Additionally, while a number of selectable markers have previously been described, many do not confer a trait of any practical agronomic value and thus needlessly complicate regulatory approval. Alternatively, labor intensive steps must be taken to attempt to breed selectable markers out of a given transgenic plant. A selectable marker gene with dual functions of a selectable marker and a trait would thus be especially valuable.

Commonly used selectable marker genes for plant transformation are neomycin phosphotransferase II, isolated from Tn5 and conferring resistance to kanamycin (Fraley et al., 1983) and hygromycin phosphotransferase, which confers resistance to the antibiotic hygromycin (Vanden Elzen et al., 1985). Additional selectable marker genes of bacterial origin that confer resistance to antibiotics include gentamycin acetyl transferase, streptomycin phosphotransferase, aminoglycoside-3'-adenyl transferase, and the bleomycin resistance determinant (Hayford et al., 1988; Jones et al., 1987; Svab et al., 1990; Hille et al., 1986).

Other selectable marker genes for plant transformation not of bacterial origin are available. These genes include, for example, mouse dihydrofolate reductase, plant 5-enolpyruvylshikimate-3-phosphate synthase and plant acetolactate synthase (Eichholtz et al., 1987; Shah et al., 1986; Charest et al., 1990). Among some herbicides that selectable marker genes confer resistance to are glyphosate, glufosinate, or bromoxynil (Comai et al., 1985; Gordon-Kamm et al., 1990; Stalker et al., 1988).

Genes encoding enzymes which inactivate herbicides and other xenophobic compounds have previously been isolated from a variety of prokaryotic and eukaryotic organisms. In some cases, these genes have been genetically engineered for successful expression in plants. Through this approach, plants have been developed which are tolerant to the herbicides 2,4-dichlorophenoxyacetic acid (Streber and Willmitzer, 1989), bromoxynil (Stalker et al., 1988), glyphosate (Comai et al., 1985) and phosphinothricin (De Block et al., 1987). While these plants have proven valuable in a commercial setting, plants tolerant to other herbicides are needed to avoid over reliance on any single herbicide and to increase options for managing difficult to control weed species.

In addition to the foregoing herbicides, there are auxin-like herbicides that mimic or act like natural plant growth regulators called auxins. Auxin-like herbicides appear to affect cell wall plasticity and nucleic acid metabolism, which can lead to uncontrolled cell division and growth. The injury symptoms caused by auxin-like herbicides includes epinastic bending and twisting of stems and petioles, leaf cupping and curling, and abnormal leaf shape and venation.

Dicamba is one example of an auxin-like herbicide and is used as a pre-emergent and post-emergent herbicide for the control of annual and perennial broadleaf weeds and several grassy weeds in corn, sorghum, small grains, pasture, hay, rangeland, sugarcane, asparagus, turf, and grass seed crops (Crop Protection Reference, 1995). Unfortunately, dicamba can injure many commercial crops and dicot plants such as soybeans, cotton, peas, potatoes, sunflowers, and canola are particularly sensitive to the herbicide. Despite this, auxin-like herbicides are very effective in controlling weed growth and thus are an important tool in agriculture. This is underscored by the development of weeds tolerant to other herbicides.

Recently, a gene for dicamba monooxygenase (DMO) was isolated from *Pseudomonas maltophilia* that confers tolerance to dicamba (US Patent Appln. 20030135879). DMO is involved in conversion of herbicidal dicamba (3,6-dichloro-o-anisic acid) to a non-toxic 3,6-dichlorosalicylic acid. This gene provides tolerance to dicamba in plants expressing the DMO gene. However, transformants containing the gene had to date only been selected using a separate selectable marker gene and techniques enabling use of a DMO gene as a direct selectable marker were not described. The need to use a separate selectable marker complicates the production of plants tolerant to auxin-like herbicides by requiring an additional gene on transformation vectors used and also presents regulatory hurdles.

Thus, there is a need in the art for new selectable marker genes and new herbicide tolerance genes. Particularly needed is a method for the selection of cells expressing a gene conferring tolerance to dicamba and other auxin-like herbicides that can be directly selected. A selectable marker gene with the dual function of a marker and a trait would eliminate the costs associated with preparing and tracking of two expression units during the development of a product and would facilitate the production of plants having valuable new traits.

SUMMARY OF THE INVENTION

In one aspect, the invention provides a method for selecting a transformed plant cell comprising the steps of: a) contacting a population of plant cells comprising a transgenic plant cell transformed with a polynucleotide encoding dicamba monooxygenase with medium comprising auxin-like herbicide in an amount that inhibits the growth of cells from the population lacking the polynucleotide, wherein the polynucleotide comprises a nucleic acid sequence selected from:

(1) a nucleic acid sequence encoding the polypeptide of SEQ ID NO:8, (2) a nucleic acid sequence comprising the sequence of SEQ ID NO:7, (3) a nucleic acid sequence that hybridizes to a complement of the nucleic acid sequence of SEQ ID NO:7 under conditions of 5×SSC, 50% formamide and 42° C., (4) a nucleic acid sequence having at least 70% sequence identity to the nucleic acid sequence of SEQ ID NO:7, and (5) a nucleic acid sequence encoding a polypeptide having at least 70% sequence identity to the polypeptide sequence of SEQ ID NO:8; and b) selecting the transformed plant cell from the population of plant cells based on tolerance exhibited by the transformed cell to the auxin-like herbicide. The population of cells may be contacted with medium comprising auxin-like herbicide any amount of time that allows selection of the transgenic cell. In certain embodiments, this may comprise at least 1-3 hours or may be carried out indefinitely, for example, for tens or even hundreds of days. In one embodiment, the method may comprise culturing the population of plant cells on a medium lacking the auxin-like herbicide prior to step a) and/or between step a) and step b). The medium lacking the auxin-like herbicide may contain a cytokinin such as 6-benzyl amino purine (BAP). In particular embodiments, 6-benzyl amino purine may be in a concentration of about 10 mg/l of medium or less, including about 8, 6, 5, 4.5, 4, 3.5, 3, 2.5, 2, 1.5, 1, and about 0.5 mg/l or less.

In certain embodiments of the invention, a polynucleotide encoding dicamba monooxygenase is not genetically linked to a selectable or screenable marker gene other than dicamba monooxygenase. The polynucleotide encoding dicamba monooxygenase may be operatively linked to a chloroplast transit peptide. A method of the invention may also further comprise the step of: regenerating a transgenic plant from the transformed plant cell. In certain aspects of the invention, the transformed plant cell is from a dicot or monocot plant. Examples of dicot plants include alfalfa, beans, broccoli, cabbage, carrot, cauliflower, cotton, pea, rapeseed, and soybean and monocots include corn, onion, rice, sorghum, and wheat. In specific embodiments, the plant is a cotton, soybean or canola plant.

In certain aspects, an auxin-like herbicide is selected from the group consisting of a phenoxy carboxylic acid compound, a benzoic acid compound, a pyridine carboxylic acid compound, a quinoline carboxylic acid compound, and a benzoxazinyl compound. In one embodiment, a phenoxy carboxylic acid compound is selected from the group consisting of 2,4-dichlorophenoxyacetic acid, 4-(2,4-dichlorophenoxy) butyric acid, and (4-chloro-2-methylphenoxy)acetic acid. In specific embodiments, a 2,4-dichlorophenoxyacetic compound, 4-(2,4-dichlorophenoxy) butyric acid, and/or (4-chloro-2-methylphenoxy)acetic acid is contained in the medium at a concentration of from about 0.001 mg/l to about 10 mg/l, including, for example, from about 0.01 mg/l to about 10 mg/l, from about 0.01 mg/l to about 5 mg/l, from about 0.1 mg/l to about 5 mg/l, from about 1 mg/l to about 5 mg/l, from about 1 mg/l to about 10 mg/l, from about 5 mg/l to about 10 mg/l, and from about 0.1 mg/l to about 3 mg/l. In other embodiments the benzoic acid is dicamba (3,6-dichloro-o-anisic acid) and is contained in the medium at a concentration of from about 0.001 mg/l to about 10 mg/l, including, for example, from about 0.01 mg/l to about 10 mg/l, from about 0.01 mg/l to about 3 mg/l, from about 0.001 mg/l to about 0.1 mg/l, from about 1 mg/l to about 10 mg/l, from about 2 mg/l to about 10 mg/l, and from about 0.001 mg/l to about 1 mg/l. In particular embodiments, the medium contains at least two auxin-like herbicides, for example, dicamba and 2,4-dichlorophenoxyacetic acid. In a method of the invention the population of cells may comprise a cotyle-

don explant and the transformed plant cell may be prepared by Agrobacterium-mediated transformation.

In another aspect, the invention provides a transgenic plant cell comprising a polynucleotide encoding dicamba monooxygenase and capable of growing in medium comprising 0.01 mg/l dicamba, wherein the dicamba monooxygenase is not genetically linked to a selectable or screenable marker gene and wherein the polynucleotide encoding dicamba monooxygenase comprises a nucleic acid sequence selected from the group consisting of (1) a nucleic acid sequence encoding the polypeptide of SEQ ID NO:8, (2) a nucleic acid sequence comprising the sequence of SEQ ID NO:7, (3) a nucleic acid sequence that hybridizes to a complement of the nucleic acid sequence of SEQ ID NO:7 under conditions of 5×SSC, 50% formamide and 42° C., (4) a nucleic acid sequence having at least 70% sequence identity to the nucleic acid sequence of SEQ ID NO:7, and (5) a nucleic acid sequence encoding a polypeptide having at least 70% sequence identity to the polypeptide sequence of SEQ ID NO:8. The cell may be defined in particular embodiments as prepared by a selection method disclosed herein. The invention also provides a tissue culture comprising such a cell. The tissue culture may comprise the cell in a media comprising auxin-like herbicide in an amount that inhibits the growth of a plant cell of the same genotype as the transgenic plant cell that lacks the polynucleotide. The invention still further provides a transgenic plant regenerated from the transgenic plant cell.

BRIEF DESCRIPTION OF THE FIGURES

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1. Response of soybean explants to dicamba with or without addition of BAP. (A) On medium without dicamba (left) or with 0.1 mg/l dicamba (right), 13DAT. (B) Explants were inoculated and co-cultivated with Agrobacterium for 3 days, and then cultured on medium with 0 (top left), 0.1 (top center), 0.5 (top right), 1.0 (bottom left), 5.0 (bottom center) and 10 (bottom right) mg/l dicamba, 11DAT (14DAI). (C) Explants were also inoculated and co-cultivated with Agrobacterium for 3 d, and then cultured on medium with different levels of dicamba combined with BAP. From left to right: 0, 0.1, 1.0, and 5.0 mg/l dicamba. From top to bottom: 0, 1.0, 3.0, 5.0 mg/l BAP.

FIG. 2. Examples of explants with GFP+small bud (top) or sectors (bottom) in experiment (Exp508) with dicamba selection. The pictures were taken at 45 DAI under regular bright field (left) or UV light for detecting GFP expression (right).

FIG. 3. GFP-positive event from dicamba selection. (A) A small shoot observed 29 DAI under regular dissecting microscope. (B) The same bud showed GFP-expression as observed under fluorescent light for detecting GFP. (C) The small shoot in A&B developed into a resistant elongated shoot (arrow), 48DAI.

FIG. 4. Response of explants cultured on medium containing 0.01 (left), 0.02 (center) and 0.05 mg/l dicamba, 23DAT (29 DAI).

FIG. 5. Detached resistant shoots were cultured on the liquid rooting medium with small glass beads (A) as support material and almost all of the shoots could produce roots. (B) Semi-solid medium can also be used for root induction.

5

FIG. 6. (A) Young soybean flowers from a transgenic plant with CP4 and GUS gene (top) and a plant transformed with pMON73691 through dicamba selection and also carrying a GFP gene (bottom). (B) The same two flowers observed under a dissecting microscope equipped with fluorescent light to detect GFP expression. GFP expression was observed on the flower transformed with pMON73691, which contains DMO and GFP genes. (C, D) The same flower transformed with pMON73691 was opened to show GFP expression in various floral structures.

FIG. 7. Soybean explants cultured on selection medium containing 0.01 mg/l dicamba after being inoculated with *Agrobacterium* harboring different constructs containing different versions of DMO gene driven with different CTP. (A) pMON73696, DMO-w, with CTP1. (B and D) pMON73698, DMO-c with CTP1. (C) pMON73691, DMO-c, with CTP2. The pictures were taken 39 (A&B) and 54 DAI (C&D), respectively. Resistant shoots are shown by arrow in panels A and C.

FIG. 8. Shows susceptibility of wild type *Arabidopsis* to various concentration of dicamba in culture medium.

FIG. 9. Shows recovery of dicamba tolerant *Arabidopsis* plants transformed with a DMO-encoding polynucleotide.

DETAILED DESCRIPTION OF THE INVENTION

The invention provides in one aspect methods for the selection of transformed cells with auxin-like herbicides such as dicamba. The invention overcomes deficiencies in the prior art that previously required coupling of a gene conferring tolerance to auxin-like herbicides to a separate selectable marker gene in order to recover transformants. Direct selection eliminates the need for extraneous selectable marker genes, which can complicate transformation procedures and subsequent regulatory approval of transgenic plants. Efficient selection of transgenic cells is crucial because typically only a small number of cells are transformed in a transformation protocol. Cells that survive exposure to the selective agent may then be cultured in media that supports regeneration of plants to produce transgenic plants. By use of a nucleic acid encoding dicamba monooxygenase (DMO) in particular, the invention allows the selection and creation of transgenic plants exhibiting tolerance to auxin-like herbicides, which can be applied to fields containing herbicide tolerant plants for effective weed control.

Selection of transformed cells in accordance with the invention may be carried out, for example, by first introducing a DMO-encoding polynucleotide molecule into a selected target plant tissue; contacting cells containing the transformed plant cell with a medium containing an auxin-like herbicide in an amount that inhibits the growth of plant cells of the same genotype as the transformed plant cell not containing the DMO-encoding polynucleotide; and selecting a plant cell capable of growing in the medium. In this manner, a transgenic cell can be selected from a large population of non-transgenic cells. In an exemplary embodiment, selective media may be modified by including further substances such as growth regulators. Tissue may be maintained on a basic media with growth regulators until sufficient tissue is available to begin plant regeneration efforts, or following repeated rounds of manual selection, until the morphology of the tissue is suitable for regeneration, typically at least 2 weeks, then transferred to media conducive to maturation into plants. Cultures may be transferred every 2 weeks on this medium. Shoot development will signal the time to transfer to medium lacking growth regulators.

6

Numerous plant tissues are amenable to transformation. The plant cell may in certain embodiments come from a plant explant, which refers to a part excised from a plant that is capable of being transformed and subsequently regenerated into a transgenic plant. Typical explants include cell suspensions, meristems, mature or immature embryos, dry embryos, wet embryos, dried embryos, seeds, callus, cotyledons, cotyledonary nodes, leaves, or stems.

Once a transgenic cell has been selected and tissues grown therefrom, the presence of the exogenous DNA or "transgene(s)" in the regenerating tissue or plants can be confirmed using a variety of assays. Such assays include, for example, "molecular biological" assays, such as Southern and northern blotting and PCRTM; "biochemical" assays, such as detecting the presence of a protein product, e.g., by immunological means (ELISAs and western blots) or by enzymatic function; plant part assays, such as leaf or root assays; and also, by analyzing the phenotype of the whole regenerated plant.

A. Nucleic Acids and Recombinant Constructs

1. Dicamba Monooxygenase (DMO)

In one embodiment of the present invention, a DNA construct expressing a dicamba monooxygenase (DMO) polypeptide is used as a selectable marker gene in plant cells. Exemplary DMO polypeptides are provided herein as SEQ ID Nos: 2, 4, 6, 8, 10 or 12. Exemplary nucleic acids encoding these sequences are provided as SEQ ID Nos: 1, 3, 5, 7, 9, or 11. Thus, in one embodiment of the invention, these sequences are used for the selection of transformed cells. As is well known in the art, homologous sequences and derivatives of these sequences may readily be prepared and used. For example, a nucleic acid may be used that encodes a DMO polypeptide having at least 70% sequence identity to a polypeptides provided as SEQ ID No: 2, 4, 6, 8, 10 or 12, including at least about 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or greater identity to such sequences. A nucleic acid may be also be used that exhibits at least 70% sequence identity to a nucleic acid sequence provided as SEQ ID No: 1, 3, 5, 7, 9, or 11, including at least about 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99% or greater identity to such sequences. In one embodiment, the identity is determined using the Sequence Analysis software package of the GCG Wisconsin Package (Accelrys, San Diego, Calif.), MEGAlign (DNASTar, Inc., 1228 S. Park St., Madison, Wis. 53715) with default parameters. Such software matches similar sequences by assigning degrees of similarity or identity.

A polynucleotide molecule that expresses a DMO polypeptide can be obtained by techniques well known in the art. Variants of DMOs having a capability to degrade auxin-like herbicides can readily be prepared and assayed for activity according to standard methods. Such sequences can also be identified by techniques known in the art, for example, from suitable organisms including bacteria that degrade auxin-like herbicides such as dicamba (U.S. Pat. No. 5,445,962; Krueger et al., 1989; Cork and Krueger, 1991; Cork and Khalil, 1995). One means of isolating a DMO sequence is by nucleic acid hybridization, for example, to a library constructed from the source organism, or by RT-PCR using mRNA from the source organism and primers based on the disclosed DMO. The invention therefore encompasses use of nucleic acids hybridizing under stringent conditions to a DMO encoding sequence described herein. One of skill in the art understands that conditions may be rendered less stringent by increasing salt concentration and decreasing temperature. Thus, hybridization conditions can be readily manipulated, and thus will

generally be a method of choice depending on the desired results. An example of high stringency conditions is 5×SSC, 50% formamide and 42° C. By conducting a wash under such conditions, for example, for 10 minutes, those sequences not hybridizing to a particular target sequence under these conditions can be removed. One embodiment of the invention thus comprises use of a DMO-encoding nucleic acid that is defined as hybridizing under wash conditions of 5×SSC, 50% formamide and 42° C. for 10 minutes to a nucleic acid selected from SEQ ID NOS: 1, 3, 5, 7, 9, or 11.

SEQ ID NO: 1 shows DMO from *Pseudomonas maltophilia* optimized for expression in dicots using *Arabidopsis thaliana* codon usage. The polypeptide, predicted to have an Ala, Thr, Cys at positions 2, 3, 112, respectively, is given in SEQ ID NO:2. SEQ ID NO:3 shows another *Pseudomonas maltophilia* DMO optimized for expression in dicots and encoding the polypeptide of SEQ ID NO:4, predicted to have an Leu, Thr, Cys at positions 2, 3, 112, respectively. SEQ ID NO:5 shows the coding sequence and SEQ ID NO:6 the polypeptide for dicot optimized DMO predicted to have a Leu, Thr, Trp at positions 2, 3, 112, respectively. SEQ ID NOS:7 and 8 show the coding and polypeptide sequences for DMO predicted to have an Ala, Thr, Cys at position 2, 3, 112, respectively. SEQ ID NOS:9 and 10 show the dicot-optimized coding sequence and polypeptide sequences for DMO predicted to have an Ala, Thr, Trp at positions 2, 3, 112, respectively. SEQ ID NOS: 11 and 12 show coding sequence and polypeptide sequences for DMO from *Pseudomonas maltophilia* (US Patent Application No: 20030135879).

Variants can also be chemically synthesized using the known DMO polynucleotide sequences according to techniques well known in the art. For instance, DNA sequences may be synthesized by phosphoramidite chemistry in an automated DNA synthesizer. Chemical synthesis has a number of advantages. In particular, chemical synthesis is desirable because codons preferred by the host in which the DNA sequence will be expressed may be used to optimize expression. Not all of the codons need to be altered to obtain improved expression, but preferably at least the codons rarely used in the host are changed to host-preferred codons. High levels of expression can be obtained by changing greater than about 50%, most preferably at least about 80%, of the codons to host-preferred codons. The codon preferences of many host cells are known (PCT WO 97/31115; PCT WO 97/11086; EP 646643; EP 553494; and U.S. Pat. Nos. 5,689,052; 5,567,862; 5,567,600; 5,552,299 and 5,017,692). The codon preferences of other host cells can be deduced by methods known in the art. Also, using chemical synthesis, the sequence of the DNA molecule or its encoded protein can be readily changed to, for example, optimize expression (for example, eliminate mRNA secondary structures that interfere with transcription or translation), add unique restriction sites at convenient points, and delete protease cleavage sites.

Modification and changes may be made to the polypeptide sequence of a protein such as the DMO sequences provided herein while retaining enzymatic activity. The following is a discussion based upon changing the amino acids of a protein to create an equivalent, or even an improved, modified polypeptide and corresponding coding sequences. In particular embodiments of the invention, DMO sequences may be altered in this manner and used in the methods of the invention. The amino acid changes may be achieved by changing the codons of the DNA sequence.

It is known, for example, that certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as binding sites on substrate molecules. Since

it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, the underlying DNA coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated that various changes may be made in the DMO peptide sequences described herein and corresponding DNA coding sequences without appreciable loss of their biological utility or activity.

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte et al., 1982). It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics (Kyte et al., 1982), these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8); tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

It is known in the art that amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, i.e., still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Pat. No. 4,554,101 states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein. As detailed in U.S. Pat. No. 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 $\pm$ 1); glutamate (+3.0 $\pm$ 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (−0.4); proline (−0.5 $\pm$ 1); alanine (−0.5); histidine (−0.5); cysteine (−1.0); methionine (−1.3); valine (−1.5); leucine (−1.8); isoleucine (−1.8); tyrosine (−2.3); phenylalanine (−2.5); tryptophan (−3.4). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. Exemplary substitutions which take these and various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

2. Transformation Constructs

A DMO-encoding polynucleotide used in accordance with the invention as a selectable marker will typically be introduced into a cell as a construct comprising expression control elements necessary for the efficient expression of DMO. Methods of operatively linking expression control elements to coding sequences are well known in the art (Maniatis et al.,

1982; Sambrook et al., 1989). Expression control sequences are DNA sequences involved in any way in the control of transcription. Suitable expression control sequences and methods of using them are well known in the art. A promoter in particular may be used, with or without enhancer elements, 5' untranslated region, transit or signal peptides for targeting of a protein or RNA product to a plant organelle, particularly to a chloroplast and 3' untranslated regions such as polyadenylation sites. One skilled in the art will know that various enhancers, promoters, introns, transit peptides, targeting signal sequences, and 5' and 3' untranslated regions (UTRs) are useful in the design of effective plant expression vectors, such as those disclosed, for example, in U.S. Patent Application Publication 2003/01403641.

Promoters suitable for the current and other uses are well known in the art. Examples describing such promoters include U.S. Pat. No. 6,437,217 (maize RS81 promoter), U.S. Pat. No. 5,641,876 (rice actin promoter), U.S. Pat. No. 6,426,446 (maize RS324 promoter), U.S. Pat. No. 6,429,362 (maize PR-1 promoter), U.S. Pat. No. 6,232,526 (maize A3 promoter), U.S. Pat. No. 6,177,611 (constitutive maize promoters), U.S. Pat. Nos. 5,322,938, 5,352,605, 5,359,142 and 5,530,196 (35S promoter), U.S. Pat. No. 6,433,252 (maize L3 oleosin promoter), U.S. Pat. No. 6,429,357 (rice actin 2 promoter as well as a rice actin 2 intron), U.S. Pat. No. 5,837,848 (root specific promoter), U.S. Pat. No. 6,294,714 (light inducible promoters), U.S. Pat. No. 6,140,078 (salt inducible promoters), U.S. Pat. No. 6,252,138 (pathogen inducible promoters), U.S. Pat. No. 6,175,060 (phosphorus deficiency inducible promoters), U.S. Pat. No. 6,635,806 (gamma-coixin promoter), and U.S. patent application Ser. No. 09/757,089 (maize chloroplast aldolase promoter). Additional promoters that may find use are a nopaline synthase (NOS) promoter (Ebert et al., 1987), the octopine synthase (OCS) promoter (which is carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*), the caulimovirus promoters such as the cauliflower mosaic virus (CaMV) 19S promoter (Lawton et al., 1987), the CaMV 35S promoter (Odell et al., 1985), the figwort mosaic virus 35S-promoter (Walker et al., 1987), the sucrose synthase promoter (Yang et al., 1990), the R gene complex promoter (Chandler et al., 1989), and the chlorophyll a/b binding protein gene promoter, etc. Particularly beneficial promoters for use with the present invention are CaMV35S, FMV35S, PCISV, AtAnt1 and P-AGRTu.nos promoters (also see Table 1).

Benefit may be obtained for the expression of heterologous genes by use of a sequence coding for a transit peptide. Transit peptides generally refer to peptide molecules that when linked to a protein of interest directs the protein to a particular tissue, cell, subcellular location, or cell organelle. Examples include, but are not limited to, chloroplast transit peptides, nuclear targeting signals, and vacuolar signals. A chloroplast transit peptide is of particular utility in the present invention for directing expression of a DMO enzyme to the chloroplasts. It is anticipated that DMO function will be facilitated by endogenous reductases and ferredoxins found in plant cells to degrade dicamba. Plant chloroplasts are particularly rich in reductases and ferredoxins. Accordingly, in a preferred embodiment for the production of transgenic dicamba-tolerant plants a sequence coding for a peptide may be used that will direct dicamba-degrading oxygenase into chloroplasts. Alternatively or in addition, heterologous reductase and/or ferredoxin can also be expressed in a cell.

DNA coding for a chloroplast targeting sequence may preferably be placed upstream (5') of a sequence coding for DMO, but may also be placed downstream (3') of the coding sequence, or both upstream and downstream of the coding

sequence. A chloroplast transit peptide (CTP) in particular can be engineered to be fused to the N-terminus of proteins that are to be targeted into the plant chloroplast. Many chloroplast-localized proteins are expressed from nuclear genes as precursors and are targeted to the chloroplast by a CTP that is removed during the import steps. Useful CTPs can be identified from the primary amino acid sequence of such polypeptides. Examples of chloroplast proteins include the small subunit (RbcS2) of ribulose-1,5-bisphosphate carboxylase, ferredoxin, ferredoxin oxidoreductase, the light-harvesting complex protein I and protein II, and thioredoxin F. It has been demonstrated in vivo and in vitro that non-chloroplast proteins may be targeted to the chloroplast by use of protein fusions with a CTP and that a CTP is sufficient to target a protein to the chloroplast. For example, incorporation of a suitable chloroplast transit peptide, such as, the *Arabidopsis thaliana* EPSPS CTP (Klee et al., 1987), and the *Petunia hybrida* EPSPS CTP (della-Cioppa et al., 1986) has been shown to target heterologous EPSPS protein sequences to chloroplasts in transgenic plants. Other exemplary chloroplast targeting sequences include the maize cab-m7 signal sequence (Becker et al., 1992; PCT WO 97/41228) and the pea glutathione reductase signal sequence (Creissen et al., 1991; PCT WO 97/41228). In the present invention, AtRbcS4 (CTP1), AtShkG (CTP2), AtShkGZm (CTP2synthetic), and PsRbcS, as well as others, disclosed in U.S. Provisional Appln. Ser. No. 60/891,675, in particular may be of benefit, for instance with regard to expression of a DMO polypeptide (e.g. SEQ ID NOs:17-28 for peptide sequences of CTPs and nucleic acid sequences that encode them).

A 5' UTR that functions as a translation leader sequence is a DNA genetic element located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences include maize and petunia heat shock protein leaders (U.S. Pat. No. 5,362,865), plant virus coat protein leaders, plant rubisco leaders, among others (Turner and Foster, 1995). In the present invention, 5' UTRs that may in particular find benefit are GmHsp, PhdNaK, AtAnt1, TEV, and L-Atnos (also see Table 1).

The 3' non-translated sequence, 3' transcription termination region, or poly adenylation region means a DNA molecule linked to and located downstream of a structural polynucleotide molecule and includes polynucleotides that provide polyadenylation signal and other regulatory signals capable of affecting transcription, mRNA processing or gene expression. The polyadenylation signal functions in plants to cause the addition of polyadenylate nucleotides to the 3' end of the mRNA precursor. The polyadenylation sequence can be derived from the natural gene, from a variety of plant genes, or from T-DNA genes. An example of a 3' transcription termination region is the nopaline synthase 3' region (nos 3'; Fraley et al., 1983). The use of different 3' nontranslated regions is exemplified (Ingelbrecht et al., 1989). Polyadenylation molecules from a *Pisum sativum* RbcS2 gene (Ps.RbcS2-E9; Coruzzi et al., 1984) and T-AGRTu.nos (Rojiyaa et al., 1987, Genbank Accession E01312) in particular may be of benefit for use with the invention.

A DMO-encoding polynucleotide molecule expression unit can be linked to a second polynucleotide molecule in an expression unit containing genetic elements for a screenable/scorable marker or for a gene conferring a desired trait. Commonly used genes for screening presumptively transformed cells include β -glucuronidase (GUS), β -galactosidase,

luciferase, and chloramphenicol acetyltransferase (Jefferson, 1987; Teeri et al., 1989; Koncz et al., 1987; De Block et al., 1984), green fluorescent protein (GFP) (Chalfie et al., 1994; Haseloff et al., 1995; and PCT application WO 97/41228). An AvGFP interrupted by StLS1 was used in the working examples for obtaining expression only in plant cells (also see Table 1).

The second polynucleotide molecule includes, but is not limited to, a gene that provides a desirable characteristic associated with plant morphology, physiology, growth and development, yield, nutritional enhancement, disease or pest resistance, or environmental or chemical tolerance and may include genetic elements comprising herbicide resistance (U.S. Pat. Nos. 6,803,501; 6,448,476; 6,248,876; 6,225,114; 6,107,549; 5,866,775; 5,804,425; 5,633,435; 5,463,175), increased yield (U.S. Pat. RE38,446; U.S. Pat. No. 6,716,474; U.S. Pat. No. 6,663,906; U.S. Pat. No. 6,476,295; U.S. Pat. No. 6,441,277; U.S. Pat. No. 6,423,828; U.S. Pat. No. 6,399,330; U.S. Pat. No. 6,372,211; U.S. Pat. No. 6,235,971; U.S. Pat. No. 6,222,098; U.S. Pat. No. 5,716,837), insect control (U.S. Pat. Nos. 6,809,078; 6,713,063; 6,686,452; 6,657,046; 6,645,497; 6,642,030; 6,639,054; 6,620,988; 6,593,293; 6,555,655; 6,538,109; 6,537,756; 6,521,442; 6,501,009; 6,468,523; 6,326,351; 6,313,378; 6,284,949; 6,281,016; 6,248,536; 6,242,241; 6,221,649; 6,177,615; 6,156,573; 6,153,814; 6,110,464; 6,093,695; 6,063,756; 6,063,597; 6,023,013; 5,959,091; 5,942,664; 5,942,658; 5,880,275; 5,763,245; 5,763,241), fungal disease resistance (U.S. Pat. Nos. 6,653,280; 6,573,361; 6,506,962; 6,316,407; 6,215,048; 5,516,671; 5,773,696; 6,121,436; 6,316,407; 6,506,962), virus resistance (U.S. Pat. Nos. 6,617,496; 6,608,241; 6,015,940; 6,013,864; 5,850,023; 5,304,730), nematode resistance (U.S. Pat. No. 6,228,992), bacterial disease resistance (U.S. Pat. No. 5,516,671), plant growth and development (U.S. Pat. Nos. 6,723,897; 6,518,488), starch production (U.S. Pat. Nos. 6,538,181; 6,538,179; 6,538,178; 5,750,876; 6,476,295), modified oils production (U.S. Pat. Nos. 6,444,876; 6,426,447; 6,380,462), high oil production (U.S. Pat. Nos. 6,495,739; 5,608,149; 6,483,008; 6,476,295), modified fatty acid content (U.S. Pat. Nos. 6,828,475; 6,822,141; 6,770,465; 6,706,950; 6,660,849; 6,596,538; 6,589,767; 6,537,750; 6,489,461; 6,459,018), high protein production (U.S. Pat. No. 6,380,466), fruit ripening (U.S. Pat. No. 5,512,466), enhanced animal and human nutrition (U.S. Pat. Nos. 6,723,837; 6,653,530; 6,5412,59; 5,985,605; 6,171,640), biopolymers (U.S. Pat. RE37,543; U.S. Pat. No. 6,228,623; U.S. Pat. No. 5,958,745 and U.S. Patent Publication No. US20030028917), environmental stress resistance (U.S. Pat. No. 6,072,103), pharmaceutical peptides and secretable peptides (U.S. Pat. Nos. 6,812,379; 6,774,283; 6,140,075; 6,080,560), improved processing traits (U.S. Pat. No. 6,476,295), improved digestibility (U.S. Pat. No. 6,531,648) low raffinose (U.S. Pat. No. 6,166,292), industrial enzyme production (U.S. Pat. No. 5,543,576), improved flavor (U.S. Pat. No. 6,011,199), nitrogen fixation (U.S. Pat. No. 5,229,114), hybrid seed production (U.S. Pat. No. 5,689,041), fiber production (U.S. Pat. Nos. 6,576,818; 6,271,443; 5,981,834; 5,869,720) and biofuel production (U.S. Pat. No. 5,998,700). Any of these or other genetic elements, methods, and transgenes may be used with the invention as will be appreciated by those of skill in the art in view of the instant disclosure.

Alternatively, the second polynucleotide molecule can affect the above mentioned plant characteristic or phenotype by encoding a RNA molecule that causes the targeted inhibition of expression of an endogenous gene, for example, via antisense, inhibitory RNA (RNAi), or cosuppression-mediated mechanisms. The RNA could also be a catalytic RNA

molecule (i.e., a ribozyme) engineered to cleave a desired endogenous mRNA product. Thus, any polynucleotide molecule that encodes a transcribed RNA molecule that affects a phenotype or morphology change of interest may be useful for the practice of the present invention.

Expression units may be provided on T-DNAs between right border (RB) and left border (LB) regions of a first plasmid together with a second plasmid carrying T-DNA transfer and integration functions in *Agrobacterium*. The constructs may also contain plasmid backbone DNA segments that provide replication function and antibiotic selection in bacterial cells, for example, an *Escherichia coli* origin of replication such as ori322, a broad host range origin of replication such as oriV or oriRi, and a coding region for a selectable marker such as Spec/Strp that encodes for Tn7 aminoglycoside adenylyltransferase (*aadA*) conferring resistance to spectinomycin or streptomycin, or a gentamicin (Gm, Gent) selectable marker gene. For plant transformation, the host bacterial strain is often *Agrobacterium tumefaciens* ABI, C58, or LBA4404. However, other strains known to those skilled in the art of plant transformation can function in the present invention.

3. Preparation of Transgenic Cells

Transforming plant cells can be achieved by any of the techniques known in the art for introduction of transgenes into cells (see, for example, Miki et al., 1993). Examples of such methods are believed to include virtually any method by which DNA can be introduced into a cell. Methods that have been described include electroporation as illustrated in U.S. Pat. No. 5,384,253; microprojectile bombardment as illustrated in U.S. Pat. Nos. 5,015,580; 5,550,318; 5,538,880; 6,160,208; 6,399,861; and 6,403,865; *Agrobacterium*-mediated transformation as illustrated in U.S. Pat. Nos. 5,635,055; 5,824,877; 5,591,616; 5,981,840; and 6,384,301; and protoplast transformation as illustrated in U.S. Pat. No. 5,508,184. Through the application of techniques such as these, the cells of virtually any plant species may be stably transformed and selected according to the invention and these cells developed into transgenic plants.

The most widely utilized method for introducing an expression vector into plants is based on the natural transformation system of *Agrobacterium* (for example, Horsch et al., 1985). *A. tumefaciens* and *A. rhizogenes* are plant pathogenic soil bacteria which genetically transform plant cells. The Ti and Ri plasmids of *A. tumefaciens* and *A. rhizogenes*, respectively, carry genes responsible for genetic transformation of the plant (for example, Kado, 1991). Descriptions of *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer are provided by numerous references, including Gruber et al., supra, Miki et al., supra, Moloney et al., 1989, and U.S. Pat. Nos. 4,940,838 and 5,464,763. Other bacteria such as *Sinorhizobium*, *Rhizobium*, and *Mesorhizobium* that interact with plants naturally can be modified to mediate gene transfer to a number of diverse plants. These plant-associated symbiotic bacteria can be made competent for gene transfer by acquisition of both a disarmed Ti plasmid and a suitable binary vector (e.g. Broothaerts et al, 2005; U.S. patent application Ser. No. 11/749,583).

B. Tissue Cultures and Media

In accordance with the invention transgenic cells may be selected by using media containing an amount of an auxin-like herbicide that inhibits the growth of a cell lacking expression of a DMO polypeptide. "Media" refers to the numerous nutrient mixtures that are used to grow cells in vitro, that is, outside of the intact living organism. The medium is usually

a suspension of various categories of ingredients (salts, amino acids, growth regulators, sugars, buffers) that are required for growth of most cell types. However, each specific cell type requires a specific range of ingredient proportions for growth, and an even more specific range of formulas for optimum growth. Rate of cell growth will also vary among cultures initiated with the array of media that permit growth of that cell type.

Regenerating a transformed plant cell can be achieved by first culturing the explant on a shooting medium and subsequently on a rooting medium. In accordance with the invention these media generally include an auxin-like herbicide such as dicamba as the selection agent besides nutrients and growth regulators. A variety of media and transfer requirements can be implemented and optimized for each plant system for plant transformation and recovery of transgenic plants. Consequently, such media and culture conditions disclosed in the present invention can be modified or substituted with nutritionally equivalent components, or similar processes for selection and recovery of transgenic events, and still fall within the scope of the present invention.

Nutrient media is prepared as a liquid, but this may be solidified by adding the liquid to materials capable of providing a solid support. Agar is most commonly used for this purpose. Bactoagar, Hazelton agar, Gelrite, and Gelgro are specific types of solid support that are suitable for growth of plant cells in tissue culture. Some cell types will grow and divide either in liquid suspension or on solid media.

Recipient cell targets include, but are not limited to, meristem cells, callus, immature embryos and gametic cells such as microspores pollen, sperm and egg cells. Any cell from which a transgenic plant, including a fertile transgenic plant, may be regenerated may be used in certain embodiments. For example, immature embryos may be transformed followed by selection and initiation of callus and subsequent regeneration of transgenic plants. Direct transformation of immature embryos obviates the need for long term development of recipient cell cultures. Meristematic cells (i.e., plant cells capable of continual cell division and characterized by an undifferentiated cytological appearance, normally found at growing points or tissues in plants such as root tips, stem apices, lateral buds, etc.) may also be used as a recipient plant cell. Because of their undifferentiated growth and capacity for organ differentiation and totipotency, a single transformed meristematic cell could be recovered as a whole transformed plant.

Somatic cells are of various types. Embryogenic cells are one example of somatic cells which may be induced to regenerate a plant through embryo formation. Non-embryogenic cells are those which typically will not respond in such a fashion.

Certain techniques may be used that enrich recipient cells within a cell population. For example, Type II callus development, followed by manual selection and culture of friable, embryogenic tissue, generally results in an enrichment of recipient cells for use in, for example, micro-projectile transformation.

Selection in culture may be carried out following plant cell transformation using a variety of transformation methods. Agrobacterium transformation followed by selection is described in the working examples below. In addition, exemplary procedures for selection of transformed cells prepared by microprojectile bombardment are provided as follows:

1. Tissue (suspension) is plated on filters, microprojectile bombarded and then filters transferred to culture medium. After 2-7 days, filters are transferred to selective medium.

Approximately 3 weeks after bombardment, tissue is picked from filters as separate callus clumps onto fresh selective medium.

2. As in 1 above, except after bombardment the suspension is put back into liquid—subjected to liquid selection for 7-14 days and then pipetted at a low density onto fresh selection plates.

3. Callus is bombarded while sitting directly on medium or on filters. Cells are transferred to selective medium 1-14 days after particle bombardment. Tissue is transferred on filters 1-3 times at 2 weeks intervals to fresh selective medium. Callus is then briefly put into liquid to disperse the tissue onto selective plates at a low density.

4. Callus tissue is transferred onto selective plates one to seven days after DNA introduction. Tissue is subcultured as small units of callus on selective plates until transformants are identified.

In certain embodiments, recipient cells are selected following growth in culture. Cultured cells may be grown either on solid supports or in the form of liquid suspensions. In either instance, nutrients may be provided to the cells in the form of media, and environmental conditions controlled. There are many types of tissue culture media comprised of amino acids, salts, sugars, growth regulators and vitamins. Most of the media employed in the practice of the invention will have some similar components, while the media can differ in composition and proportions of ingredients according to known tissue culture practices. For example, various cell types usually grow in more than one type of media, but will exhibit different growth rates and different morphologies, depending on the growth media. In some media, cells survive but do not divide. Media composition is also frequently optimized based on the species or cell type selected.

Various types of media suitable for culture of plant cells have been previously described. Examples of such media are defined below. In some embodiments, it may be preferable to use a media with a somewhat lower ammonia/nitrate ratio such as N6 to promote generation of recipient cells by maintaining cells in a proembryonic state capable of sustained divisions. In certain embodiments of the present invention, Woody Plant Medium (WPM) was used (Lloyd and McCown, 1981).

The method of maintenance of cell cultures may contribute to their utility as sources of recipient cells for transformation. Manual selection of cells for transfer to fresh culture medium, frequency of transfer to fresh culture medium, composition of culture medium, and environment factors including, but not limited to, light quality and quantity and temperature are all factors in maintaining callus and/or suspension cultures that are useful as sources of recipient cells. Alternating callus between different culture conditions may be beneficial in enriching for recipient cells within a culture. For example, cells may be cultured in suspension culture, but transferred to solid medium at regular intervals. After a period of growth on solid medium, cells can be manually selected for return to liquid culture medium. Repeating this sequence of transfers to fresh culture medium may be used to enrich for recipient cells. Passing cell cultures through a 1.9 mm sieve may also be useful to maintain the friability of a callus or suspension culture and enriching for transformable cells when such cell types are used.

C. Transgenic Plants

Once a transgenic cell has been selected, the cell can be regenerated into a transgenic plant using techniques well known in the art. The transformed plants can be subsequently

analyzed to determine the presence or absence of a particular nucleic acid of interest contained on a DNA construct. Molecular analyses can include, but are not limited to, Southern blots (Southern, 1975), northern blot analysis, western blot analysis, or PCR analyses, immunodiagnostic approaches, and field evaluations. These and other well known methods can be performed to confirm the stability of the transformed plants produced by the methods disclosed. These methods are well known to those of skill in the art (Sambrook et al., 1989).

Transgenic plants tolerant to auxin-like herbicides can be produced. In particular, economically important plants, including crops, fruit trees, and ornamental plants and trees that are currently known to be injured by auxin-like herbicides can be transformed with DNA constructs of the present invention so that they become tolerant to the herbicide. Plants that are currently considered tolerant to auxin-like herbicides can be transformed to increase their tolerance to the herbicide. Examples of plants that may in particular find use with the current invention include, but are not limited to, alfalfa, beans, broccoli, cabbage, carrot, cauliflower, celery, cotton, cucumber, eggplant, lettuce, melon, pea, pepper, pumpkin, radish, rapeseed, spinach, soybean, squash, tomato, watermelon, corn, onion, rice, sorghum, wheat, rye, millet, sugarcane, oat, triticale, switchgrass, and turfgrass.

Once a transgenic plant containing a transgene has been prepared, that transgene can be introduced into any plant sexually compatible with the first plant by crossing, without the need for ever directly transforming the second plant. Therefore, as used herein the term "progeny" denotes the offspring of any generation of a parent plant prepared in accordance with the instant invention, wherein the progeny comprises a selected DNA construct prepared in accordance with the invention. A "transgenic plant" may thus be of any generation. "Crossing" a plant to provide a plant line having one or more added transgenes or alleles relative to a starting plant line, as disclosed herein, is defined as the techniques that result in a particular sequence being introduced into a plant line by crossing a starting line with a donor plant line that comprises a transgene or allele of the invention. To achieve this one could, for example, perform the following steps: (a) plant seeds of the first (starting line) and second (donor plant line that comprises a desired transgene or allele) parent plants; (b) grow the seeds of the first and second parent plants into plants that bear flowers; (c) pollinate a flower from the first parent plant with pollen from the second parent plant; and (d) harvest seeds produced on the parent plant bearing the fertilized flower.

D. Definitions

As used herein, the term "transformed" refers to a cell, tissue, organ, or organism into which has been introduced a foreign polynucleotide molecule, such as a construct. The introduced polynucleotide molecule may be integrated into the genomic DNA of the recipient cell, tissue, organ, or organism such that the introduced polynucleotide molecule is inherited by subsequent progeny. A "transgenic" or "transformed" cell or organism also includes progeny of the cell or organism and progeny produced from a breeding program employing such a transgenic plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of a foreign polynucleotide molecule.

"Contacting" the transformed plant cell with a tissue culture medium containing an auxin-like herbicide can be achieved by culturing the plant cell in a plant tissue culture medium containing an auxin-like herbicide.

"Tissue culture medium" refers to liquid, semi-solid, or solid medium used to support plant growth and development in a non-soil environment. Suitable plant tissue culture media are known to one of skill in the art may include MS-based media (Murashige and Skoog, 1962) or N6-based media (Chu et al., 1975) supplemented with additional plant growth regulators such as auxins, cytokinins, kinetin, ABA, and gibberellins. Other media additives can include but are not limited to amino acids, macroelements, iron, microelements, inositol, vitamins and organics, carbohydrates, undefined media components such as casein hydrolysates, with or without an appropriate gelling agent such as a form of agar, such as a low melting point agarose or Gelrite® if desired for preparing semi-solid or solid medium. Those of skill in the art are familiar with the variety of tissue culture media, which when supplemented appropriately, support plant tissue growth and development and are suitable for plant transformation and regeneration. These tissue culture media can either be purchased as a commercial preparation or custom prepared and modified. Examples of such media would include but are not limited to Murashige and Skoog (1962), N6 (Chu et al., 1975), Linsmaier and Skoog (1965), Uchimiya and Murashige (1962), Gamborg's media (Gamborg et al., 1968), D medium (Duncan et al., 1985), McCown's Woody plant media (McCown and Lloyd, 1981), Nitsch and Nitsch (1969), and Schenk and Hildebrandt (1972) or derivations of these media supplemented accordingly. Those of skill in the art are aware that media and media supplements such as nutrients and growth regulators for use in transformation and regeneration and other culture conditions such as light intensity during incubation, pH, and incubation temperatures can be optimized for a plant of interest.

"Auxin-like herbicides" are also called auxinic or growth regulator herbicides or Group 4 herbicides (based on their mode of action). These herbicides mimic or act like the natural plant growth regulators called auxins. Auxins include natural hormones such as indole acetic acid and naphthalene acetic acid, both of which are responsible for cell elongation in plants. The mode of action of the auxinic herbicides is that they appear to affect cell wall plasticity and nucleic acid metabolism, which can lead to uncontrolled cell division and growth. The group of auxin-like herbicides includes four chemical families: phenoxy, carboxylic acid (or pyridine), benzoic acid, and the newest family quinaline carboxylic acid. Phenoxy carboxylic acids: the phenoxy herbicides are most common and have been used as herbicides since the 1940s when (2,4-dichlorophenoxy)acetic acid (2,4-D) was discovered. Other examples include 4-(2,4-dichlorophenoxy) butyric acid (2,4-DB), 2-(2,4-dichlorophenoxy) propanoic acid (2,4-DP), (2,4,5-trichlorophenoxy)acetic acid (2,4,5-T), 2-(2,4,5-Trichlorophenoxy) Propionic Acid (2,4,5-TP), 2-(2,4-dichloro-3-methylphenoxy)-N-phenylpropanamide (clomeprop), (4-chloro-2-methylphenoxy)acetic acid (MCPA), 4-(4-chloro-o-tolyloxy) butyric acid (MCPB), and 2-(4-chloro-2-methylphenoxy) propanoic acid (MCPBP).

Pyridine carboxylic acids: the next largest chemical family is the carboxylic acid herbicides, also called pyridine herbicides. Examples include 3,6-dichloro-2-pyridinecarboxylic acid (Clopyralid), 4-amino-3,5,6-trichloro-2-pyridinecarboxylic acid (picloram), (2,4,5-trichlorophenoxy)acetic acid (triclopyr), and 4-amino-3,5-dichloro-6-fluoro-2-pyridylloxyacetic acid (fluoroxypyr). Benzoic acids: Examples include 3,6-dichloro-o-anisic acid (dicamba) and 3-amino-2,5-dichlorobenzoic acid (choramben). Quinaline carboxylic acids: the fourth and newest chemical family of the auxinic herbicides is the quinaline carboxylic acid family. Example includes 3,7-dichloro-8-quinolinecarboxylic acid (quinclor-

rac). This herbicide is unique in that it also will control some grass weeds, unlike the other auxin-like herbicides which essentially control only broadleaf or dicotyledonous plants. The other herbicide in this category is 7-chloro-3-methyl-8-quinolinecarboxylic acid (quinmerac).

"Auxin-like herbicide effect" means injury symptoms caused by auxin-like herbicides. These include epinastic bending and twisting of stems and petioles, leaf cupping and curling, and abnormal leaf shape and venation. All of these herbicides translocate, with some translocating more than others. Some of these herbicides have soil activity and some can persist in soil for fairly long time periods. Due to their effect, they are used widely on many crops including small grain cereals, corn, rice, and other grass crops, turf, rangeland, non-crop, and industrial sites.

"Selecting" the transformed plant cell that is tolerant to an auxin-like herbicide can be achieved by methods described in the present invention. Briefly, at least some of the plant cells in a population of starting cells are transformed with a DNA construct containing a DMO-encoding polynucleotide molecule. The resulting population of plant cells is placed in a culture medium containing an auxin-like herbicide at a concentration selected so that transformed plant cells will grow, whereas untransformed plant cells will not. Suitable concentrations of an auxin-like herbicide can be determined empirically. Before selecting, explants may be cultured on a medium without auxin-like herbicide. Such medium is called delay medium. Explants may be placed on delay medium to allow for some time to grow before being placed on the selection medium. Selection regimes could be optimized depending upon a particular auxin-like herbicide and the explant system. Often multiple steps of selection are used and varying amounts of selection agent can be used in each step.

"Tolerant" means that transformed plant cells are able to survive and regenerate into plants when placed in a culture medium containing a level of an auxin-like herbicide that prevents untransformed cells from doing so. "Tolerant" also means that transformed plants are able to grow after application of an amount of an auxin-like herbicide that inhibits the growth of untransformed plants.

EXAMPLES

The following examples are included to illustrate embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent techniques discovered by the inventor to function well in the practice of the invention. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

Example 1

Preparation of DMO-Encoding Polynucleotide Constructs

Several binary vectors were prepared for testing the ability of DMO-encoding polynucleotide molecules to allow selec-

tion of transformed soybean cells. Genetic elements used for preparing the binary vectors are given in Table 1 and include a CaMV 35S promoter and enhancer (U.S. Pat. Nos. 5,322,938; 5,352,605; 5,359,142; and 5,530,196); GmHsp untranslated leader from the Hsp17.9 gene of *Glycine max* (U.S. Pat. No. 5,659,122); AvGFPI coding region for the first 126.3 amino acids of the GFP protein from *Aequorea victoria* (U.S. Pat. Nos. 5,491,084; 6,146,826) with a serine to threonine change at amino acid 65 and optimized for plant expression; a StLS1 second intron from the LS1 gene of *Solanum tuberosum* (Eckes et al., 1986); an AvGFPII coding region for the last 112.6 amino acids of the GFP protein from *Aequorea victoria* (U.S. Pat. Nos. 5,491,084; 6,146,826) optimized for plant expression; a T-Atnos 3' untranslated region of the nopaline synthetase gene from *Agrobacterium tumefaciens* (Rojiyaa et al., 1987, GenBank Accession E01312); a FMV Figwort Mosaic Virus 35S promoter (U.S. Pat. Nos. 6,051,753; 5,378,619); a PhDnaK untranslated leader from Hsp70 gene of *Petunia hybrida* (U.S. Pat. No. 5,362,865); and an AtRbcS4 (CTP1) coding region for *Arabidopsis* SSU1A transit peptide. The latter element includes the transit peptide, 24 amino acids of the mature protein, and a repeat of the last 6 amino acids of the transit peptide (U.S. Pat. No. 5,728,925). Also used were an AtShkG (CTP2) coding region for *Arabidopsis thaliana* 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) transit peptide. This element varies from the wild-type sequence (Klee et al., 1987) in that the last codon was changed from GAG (glutamic acid) to TGC (cysteine). An AtShkGzm codon (CTP2syn) element was used which is the same as AtShkG (CTP2) but optimized for plant expression using *Zea mays* codons (see SEQ ID NO:14 of WO04009761). A PmDMOCys112Atcodon region for dicamba monooxygenase from *Pseudomonas maltophilia* was used (US Patent Application 20030115626) having a cysteine at 112 position and optimized for dicot expression using *Arabidopsis thaliana* codons (SEQ ID NOs: 1, 3, 7). Also used for construct design were a PmDMOTrp112Atcodon coding region for dicamba monooxygenase from *Pseudomonas maltophilia* (US patent application 20030115626) having a tryptophan (Trp) at 112 position and optimized for dicot expression using *Arabidopsis thaliana* codons (SEQ ID NOs: 5, 9); a PsRbcS2: 3' polyadenylation region from the RbcS2-E9 gene of *Pisum sativum* (Coruzzi et al., 1984); an AtAnt1 promoter/intron and leader of adenine nucleotide translocator 1 gene from *Arabidopsis thaliana*; an AtaroA-CP4 coding region for non-naturally occurring aroA-CP4 (U.S. Pat. No. 5,633,435) engineered for expression in plants; a TEV 5' untranslated leader from the Tobacco Etch RNA virus (Carrington and Freed, 1990); a PsRbcS chloroplast transit peptide from ribulose 1,5-bisphosphate carboxylase small subunit of pea and first 24 amino acids of the mature rubisco protein (Coruzzi et al., 1984); a P-Atnos promoter for nopaline synthetase of *Agrobacterium tumefaciens* pTiT37 (GenBank Accession V00087; Depicker et al, 1982; Bevan et al., 1983); a L-At.nos 5' untranslated region from the nopaline synthetase gene of *Agrobacterium tumefaciens* pTiT37 (GenBank Accession V00087; Bevan et al., 1983), and a PCISV promoter for the full length transcript of peanut chlorotic streak virus. The latter element has a duplication of 179 nt in direct repeats with 6 nt between the repeat followed by the 70 bp region containing the TATA box (U.S. Pat. No. 5,850,019). Different CTPs and DMO-encoding polynucleotide molecule variants are summarized in Table 2.

TABLE 1

Genetic elements used for constructing T-DNAs											
Expression Unit 1						Expression Unit 2					
Construct	Pro-moter	5'UTR	CR	Intron	CR	PolyA	Pro-moter	5' UL	TS	CR	PolyA
pMON73690	CaMV	GmHsp	AvGFPI	StLS1	AvGFPII	T-Atnos	FMV	PhDnaK	None	Pm-DMOCys ₁₁₂ -Atcodon	PsRbcS2
pMON73691	CaMV	GmHsp	AvGFPI	StLS1	AvGFPII	T-Atnos	FMV	PhDnaK	AtShkG	Pm-DMOCys ₁₁₂ -Atcodon	PsRbcS2
pMON73696	CaMV	GmHsp	AvGFPI	StLS1	AvGFPII	T-Atnos	FMV	PhDnaK	AtRbcS4	Pm-DMOTrp ₁₁₂ -Atcodon	PsRbcS2
pMON73698	CaMV	GmHsp	AvGFPI	StLS1	AvGFPII	T-Atnos	FMV	PhDnaK	AtRbcS4	Pm-DMOCys ₁₁₂ -Atcodon	PsRbcS2
	Pro-moter	5'UR	TS	CR			Pro-moter	5' UL	TS	CR	PolyA
pMON73724	AtAnt1	AtAnt1	At.ShkG	At.aroA-CP4	PsRbcS2		PC1SV	TEV	AtShkGZ-meodon	PmDMOTrp ₁₁₂ Atcodon	Atnos
First T-DNA						Second T-DNA					
	Promoter	5'UTR	TS	CR	PolyA	Promoter	5' UL	TS	CR	PolyA	
pMON58498	PC1SV	TEV	PsRbcS	PmDMO Cys ₁₁₂	PsRbcS2	FMV	PhDnaK	AtShkG	AtaroA-CP4	PsRbcS2	
pMON84254	PC1SV	TEV	PsRbcS	PmDMO Cys ₁₁₂	PsRbcS2	P-Atnos	L-Atnos	None	Sh.bar	T-Atnos	

Key:

5'UTR: 5' untranslated region;

CR: coding region;

Poly A: polyadenylation region; and

TS: transit sequence.

TABLE 2

Chloroplast transit peptides and DMO-encoding polynucleotides used in binary vectors.						
Construct	CTP variant	DMO variant	SEQ ID	Length	Predicted aa at position 2	Predicted aa at position 112
pMON73690	None	DMO-Cys ₁₁₂ and codon optimized for dicots	1	1023	Ala	Cys
pMON73691	CTP2	DMO-Cys ₁₁₂ and codon optimized for dicots	3	1023	Leu	Cys
pMON73696	CTP1	DMO-Trp ₁₁₂ and codon optimized for dicots	5	1023	Leu	Trp
pMON73698	CTP1	DMO-Cys ₁₁₂ and codon optimized for dicots	3	1023	Leu	Cys
pMON58498	PsRbcS	DMO-Cys ₁₁₂	7	1023	Ala	Cys
pMON84254	PsRbcS	DMO-Cys ₁₁₂	7	1023	Ala	Cys
pMON73724	CTP2Zm	DMO-Trp ₁₁₂ and codon optimized for dicots	9	1023	Ala	Trp

In the case of pMON73690, pMON73691, pMON73696, and pMON73698, the DMO-encoding polynucleotide molecule was linked to the screenable marker GFP and provided on the same T-DNA to show that the DMO-encoding polynucleotide molecule can be used with another gene. In case of pMON58498, pMON84254, and pMON73724, the DMO-encoding polynucleotide molecule was unlinked from the other transgene (selectable marker or agronomic trait gene) by separating them on two T-DNAs.

Example 2

Development of Selection Method

Mature seeds of soybean [*Glycine max* (L.) Merrill] cv. A3525 were imbibed, sterilized, and germinated at room

temperature as set forth below. Other examples of soybean genotypes that can readily be used include, but are not limited to, Jack, Williams, Bert, Thorne, Granite, Lambert, Chapman, and Kunitz. Briefly, dry seeds (about 770 g) were soaked for 3 min in 2 L of 200 ppm sodium hypochlorite solution made from commercially available Clorox. The solution was drained and the seeds were set side for about 2 h. About 2 L of bean sterilization/germination medium was then added to the seeds. After about 9-10 h, seeds were ready for hand excision of explants. The bean germination medium contained the following in mg/L—NH₄NO₃: 240, KNO₃: 505, CaCl₂·2H₂O: 176, MgSO₄·7H₂O: 493, KH₂PO₄: 27, H₃BO₃: 1.86, Na₂MoO₄·2H₂O: 0.216, MnSO₄·H₂O: 5.07, ZnSO₄·7H₂O: 2.58, FeSO₄·7H₂O: 2.502, KI: 0.249, Na₂EDTA·2H₂O: 3.348, CuSO₄·5H₂O: 0.0008, CoCl₂·6H₂O: 0.0008, B1: 1.34, B3: 0.5, B6: 0.82, Bravo (75% WP; Diamond Sham-

rock Company, Cleveland, Ohio): 30, Captan (50% WP; Micro Flo Company, Lakeland, Fla.): 30, Cefotaxime: 125, and Sucrose: 25000, pH 5.8).

For machine excision of the explants, seeds were treated with 2 L of 200 ppm sodium hypochlorite solution for 15 min. After draining the solution the seeds were rinsed with 2 L of sterile distilled water for 1 min. The machine and method for mechanical excision are described in the US Patent Appln. Pub. 20050005321. Briefly, imbibed seeds were run through three sets of rollers in the machine, with sterile distilled water running over them, and crushed. A mixture of cotyledons, seed coats and the explants (embryo axis) is collected and sieved by either hand or by using an auto-sieving device to recover the explants. The explants were rinsed with 0.05% ethanol for 1 min, followed by two rinses with sterile distilled water for removing more debris.

The binary vectors described above were mobilized into disarmed *Agrobacterium tumefaciens* strain C58 (ABI). *Agrobacterium* inoculum for infection was prepared as follows: 250 ml of LB medium (Luria-Bertani; Difco, Detroit, Mich.) containing 50 mg/l kanamycin (Sigma, St. Louis, Mo.) and 75 mg/l spectinomycin (Sigma, St. Louis, Mo.) was inoculated with 0.5 ml of *Agrobacterium* stock in glycerol (Acros Organics, Geel, Belgium) and was shaken at 200 rpm at 28° C. for approximately 20-22 h until the OD₆₆₀ reached 0.8 to 1.0. The *Agrobacterium* broth was then centrifuged for 25 min at 3500 rpm (about 3565 g) at 2-4° C. After removing the supernatant, the *Agrobacterium* pellet was re-suspended in inoculation medium containing $\frac{2}{5} \times$ of the macro nutrients, $\frac{1}{10} \times$ of the micro nutrients and vitamins of Gamborg's B5 medium, supplemented with 3.9 g/l MES (Sigma, St. Louis, Mo.), and 30 g/l glucose (PhytoTechnology Laboratories, Shawnee Mission, Kans.), pH 5.4. Lipoic acid (Sigma, St. Louis, Mo.) was added to the *Agrobacterium* suspension to a final concentration of 0.25 mM after the density of the *Agrobacterium* cell suspension was adjusted to an OD₆₆₀ of 0.30 to 0.35.

Agrobacterium-infection and co-cultivation of the explants were conducted as follows: about 100 excised explants were dispensed into the lid of a sterile plastic culture vessel PLANTCON (MP Biomedicals, LLC, Irvine, Calif.). Five ml of *Agrobacterium* inoculum was added to the explants in each PLANTCON lid. The explants were then sonicated for 20 sec in a sonicator (Ultrasonic Multi Cleaner, Model W-113, Honda, Japan). One piece of Whatmann #1 filter paper (Whatman Inc., Clifton, N.J.) cut to the size of the PLANTCON bottom was placed in the bottom part of the PLANTCON. The explants were transferred from the lid onto the filter paper with the *Agrobacterium* inoculum. The PLANTCONs were then incubated in a Percival incubator at 16 h light (at about 85-90 μ E) and 8 h dark photoperiod and at 23° C. for 2 to 4 d for co-cultivation.

After a co-cultivation period of 2-4 days, explants were first cultured on a medium without dicamba (delay medium) for 3-5 d before being transferred to the selection medium with dicamba. Until transfer from delay medium to selection medium, the explants were kept in the same PLANTCON used for co-cultivation, but 10 or 12 ml of the delay medium was added to each PLANTCON. Alternatively, the explants were transferred to new PLANTCONs, each containing one piece of autoclaved felt (Jo-Ann Fabrics & Crafts, Madison, Wis.) and 30 ml of the delay medium. The delay medium contained modified wood plants medium (Lloyd and McCown, 1981) supplemented with 1 or 5 mg/l BAP (6-benzyl Amino Purine), 200 mg/l carbenicillin (Phyto Technology Laboratories, Shawnee Mission, Kans.), 200 mg/l cefotaxime (Hospira, Lake Forest, Ill.) and 100 mg/l ticarcillin (Duchefa,

The Netherlands). BAP may help maintain the auxin-cytokinin ratio as dicamba is an auxin herbicide, and promote production of multiple shoots from the apical meristem. Other suitable cytokinins that can be useful in practicing the present invention include: Adenine cytokinins (e.g., kinetin, zeatin, benzyl adenine (i.e. 6-Benzyl aminopurine), adenine and Phenylurea cytokinins (e.g., N,N'-diphenylurea), and Thidiazuron (TDZ).

Selection was conducted in a liquid or on a semi-solid medium. The selection medium was the delay medium absent BAP and contained different concentrations of dicamba. For selection in liquid medium, 50 or 60 ml of the selection medium and one piece of foam sponge (Wisconsin Foam Products, Madison, Wis.) having 5 parallel slits were placed in each Plantcon. Twenty-five explants were implanted into the slits in an upward position such that apical meristem faced upward. Every two to three weeks, old medium was replaced with the fresh medium.

Semi-solid medium was prepared by adding 4 g/l AgarGel (Sigma, St. Louis, Mo.) to the liquid medium. For selection on semi-solid selection medium, the explants were individually implanted into the medium in PLANTCONs. At the late stage of the selection and shoot development (approximately 4 weeks on the selection medium), 20 ml of the liquid selection medium was optionally overlaid on the semi-solid medium. Elongated shoots with expanded trifoliate foliage leaves started to develop after the explants had been cultured on the selection medium for about 4 to 5 weeks. These tolerant shoots were detached from the original explants when they were about and over 2 cm long and transferred to the liquid or semi-solid root induction medium.

The medium for root induction was the same as for shoot development and was also supplemented with dicamba to reduce the frequency of escapes. Alternatively, Bean Rooting Medium (BRM) supplemented with 0.01 mg/l dicamba was used for root induction. This medium contained $\frac{1}{2}$ strength of MS salts, MS vitamins, 100 mg/l inositol, 100 mg/l cysteine, 30 mg/l sucrose and 100 mg/l ticarcillin and was solidified with 8 g/l washed agar. For root induction in the liquid medium, enough small glass beads (Inotech Biosystems International Inc., Dottikon, Switzerland) and 60 ml of the rooting medium were placed in each PLANTCON such that the medium and beads were at the same level. Up to nine detached shoots were inserted into the beads for liquid root induction or in semi-solid medium in each PLANTCON. Almost all shoots could produce roots on the rooting medium in 1-2 weeks (FIG. 5). However, only those shoots in which the existing and newly developed leaves remained expanded and grew vigorously were transferred to soil for growing to maturity. All cultures were kept under fluorescent light with a photoperiod of 16 h with light intensity of about 20-70 μ E at 27-28° C. until R₀ plants were transferred to the soil.

In one study, soybean cells transformed with pMON73691 were selected on 0.01 to 0.1 mg/l of dicamba in selection medium (FIGS. 1 A & B; FIG. 4). Shoots coming out of explants grown on selection medium with 0.05 or 0.1 mg/l dicamba did not have much growth and eventually bleached out and no tolerant shoots were obtained. However, in selection medium containing 0.01 mg/l dicamba, 30 dicamba-tolerant shoots were harvested from 800 explants. Twelve of these formed roots on rooting medium and were transferred to the soil. Ten of these were tested for DMO-encoding polynucleotides and seven were found to be positive. At a dicamba level of 0.02 mg/l, few tolerant shoots were harvested. These results suggested that a dicamba concentration of 0.01 mg/l or lower was most efficacious for selecting dicamba-tolerant shoots. This level could readily be altered by one skill in the

art for particular studies, however, depending upon the nature of the explant, construct, and other variables.

In order to demonstrate the selection of tolerant shoots containing a linked gene, a plant expressible DMO-encoding nucleic acid coupled to a plant expressible GFP-encoding nucleic acid and introduced into cells following selection. The cultures were examined for GFP expression 45 days after inoculation (DAI). GFP-positive small buds were observed on several explants, suggesting that these buds originated from cells transformed with the linked DMO-encoding polynucleotide molecule (FIG. 2). Several of these buds developed into GFP positive shoots and were positive for GFP gene (FIGS. 3, 6). These results demonstrated that a DMO-encoding polynucleotide molecule can be used as a selectable marker and be used for recovery of transformants containing and expressing a linked gene. Confirmation that the GFP transgene was inherited in the progeny was found by self-pollinating R₀ plants transformed with pMON73691. Immature R₁ seeds (about 4 mm in length) were collected and cut into two halves to expose the cotyledon tissue. GFP expression was detected in the cotyledon tissue of seeds under a dissecting microscope equipped with fluorescent light.

Rooting was also accomplished in Oasis Growing Medium i.e. plugs (Smithers-Oasis North America, Kent, Ohio, USA). A total of 102 Dicamba-selected shoots were inserted into Oasis plugs for inducing roots. The plugs were surrounded by a liquid medium containing 0.01 mg/l dicamba. The shoots in the plugs were kept in culture room at 28° C. and 16-h light. Thirty shoots developed roots and appeared to be resistant to dicamba showing relatively expanded new leaves. The plants with roots were tested by invader assay and 19 plants were found to contain both DMO and GUS genes. The escape rate on the liquid medium was about 33%, which was much lower than the 53% escape rate when the roots were induced in the semi-solid medium. The negative phenotypes of the shoots i.e., cupping leaves could be seen sooner in the liquid selection medium than in the semi-solid medium. This suggested that rooting in the liquid selection medium could be a more efficient method to eliminate escapes.

Example 3

Molecular Analysis of Transformed Soybean Plants

In order to confirm that the dicamba tolerant plants obtained were the result of transfer of DMO-encoding polynucleotides, leaf tissue was collected from each R₀ or R₁ plant, DNA was extracted, and the presence of the DMO-encoding polynucleotide was confirmed by Invader™ technology (Third Wave Technologies, Madison, Wis.) and Southern blot analysis using non-radioactive probe kit from Roche (Indianapolis, Ind.).

For the Invader assay, the primers used were: primary probe 5'-acggacgcggag ATGCTCAACTTCATCGC-3' (SEQ

ID NO: 13) and Invader oligo 5'-TCCGCTGGAACAAGGT-GAGCGCGT-3' (SEQ ID NO: 14). The sequence in lower case letters in the primary probe is the 5' flap sequence which is cleaved and is not complimentary to the target sequence.

For Southern blot analysis a DNA fragment of 897 bp was used to prepare the probe. The forward primer 5'-GTCGCT-GCCCTGCTTGATATT-3' (SEQ ID NO: 15) and the reverse primer 5'-CGCCGCTTCTAGTTGTTTC-3' (SEQ ID NO: 16) were used to amplify the 897 bp DNA fragment. A total of 12 rooted shoots selected on 0.01 mg/l dicamba shoots were transferred to soil. Ten plants were assayed by Invader and/or Southern analysis. Seven of these showed the presence of the DMO-encoding nucleic acid (Table 3). Several of these were also positive for GFP-encoding polynucleotides confirming the ability to use the DMO-encoding nucleic acid as a selectable marker for recovery of transformants containing a linked gene.

TABLE 3

Testing of R0 plants for DMO-encoding nucleic acid by Invader and/or Southern Analysis.			
Plant Name	Origin (Exp-Trt)	Invader	Southern
GM_A4755D	533-1	+	+
GM_A4756D	533-1	-	-
GM_A4757D	533-1	-	-
GM_A4758D	533-1	-	-
GM_A4763D	533-1	+	+
GM_A4759D	534-1	+	+
GM_A4760D	534-1	-	+
GM_A4761D	534-1	+	+
GM_A4764D	534-1	N/T	+
GM_A5087D	534-1	N/T	+

Example 4

Selection of Dicamba-Tolerant Plants Transformed with DMO-Encoding Polynucleotide Molecule Variants

Two DMO-encoding polynucleotide molecule variants were used to obtain dicamba tolerant plants. The first variant had cysteine at amino acid position 112 (DMOCys₁₁₂; pMON73698) and the second had tryptophan at amino acid position 112 (DMOTrp₁₁₂; pMON73696). Selection using both variants resulted into dicamba tolerant shoots (FIG. 7). Following selection and shoot and root induction, the rooted plants were moved to soil for growing to maturity and assayed by Invader™ and/or Southern analysis for the presence of DMO-encoding nucleic acid. Several plants transformed with pMON73696 were found to be positive for the DMO gene in R₀ and R₁ generation indicating germline transformation using the method of the present invention.

TABLE 4

Selection of dicamba-tolerant shoots transformed with DMO-encoding nucleic acid variants.					
Medium	Construct	# Explants	# Tolerant shoots harvested	# Rooted shoots moved to soil	# plants with DMO gene (# plants assayed)
Liquid	73696 (DMOTrp ₁₁₂)	1022	6	1	0
	73698 (DMOCys ₁₁₂)	869	0	0	0
	73696 (DMOTrp ₁₁₂)	1200	25	9	3
	73698 (DMOCys ₁₁₂)	1200	0	0	0

TABLE 4-continued

Selection of dicamba-tolerant shoots transformed with DMO-encoding nucleic acid variants.					
Medium	Construct	# Explants	# Tolerant shoots harvested	# Rooted shoots moved to soil	# plants with DMO gene (# plants assayed)
Semisolid	73696 (DMOTrp ₁₁₂)	1536	94	50	28 (47)
	73698 (DMOCys ₁₁₂)	1845	3	0	0 (0)
	73696 (DMOTrp ₁₁₂)	450	27	18	10 (16)
	73698 (DMOCys ₁₁₂)	475	0	0	0 (0)

Example 5

Selection of Dicamba-Tolerant Plants Transformed
with DMO-Encoding Polynucleotide Molecules
Combined with Different Chloroplast Transit
Peptides

It is known that different chloroplast transit peptides (CTPs) target a foreign polypeptide to chloroplasts with different efficiencies. The effect of different types of CTPs was therefore tested by transforming soybean explants with DMO-encoding polynucleotide molecules targeted either by CTP2 (pMON73691) or CTP1 (pMON73698) or not targeted to chloroplasts (pMON73690). As shown in Table 5, in general shoots were harvested with constructs containing either CTP2 or CTP1 (also see FIG. 7). The rooted plants were moved to soil for growing to maturity and assayed by Invader™ and/or Southern analysis for the presence of DMO-encoding nucleic acid. Several plants transformed with pMON73691 were found to be positive for the DMO gene in R₀ and R₁ generation indicating germline transformation using the method of the present invention.

Several transgenic plants carrying either a PCISV/RbcS/DMO-Wdc/Nos or PCISV/CTP2nat/DMO-Cnative/Nos expression unit were also found to be tolerant to a dicamba treatment at the rate of 1 lb/A (Clarity, BASF) at V3-4 when analysed 18 DAT in a greenhouse study.

TABLE 5

Selection of dicamba tolerant plants transformed with DMO-encoding nucleic acid combined with different chloroplast transit peptides.					
Exp- Trt	Construct	# Explants	# tolerant shoots harvested	# rooted shoots moved to soil	# plant with DMO gene (# plants assayed)
576-1	73691 (CTP2/DMOCys ₁₁₂)	1350	74	14	10 (14)
625-3	73691 (CTP2/DMOCys ₁₁₂)	500	17	6	4 (5)
576-2	73698 (CTP1/DMOCys ₁₁₂)	1050	22	1	1 (1)
625-1	73690 (DMOCys ₁₁₂)	531	1	0	0
625-2	73690 (DMOCys ₁₁₂)	531	2	0	0

Example 6

20 Use of DMO-Encoding Polynucleotide Molecule as
a Selectable Marker in Combination with an
Agronomic Trait Gene

25 One beneficial use of a DMO-encoding polynucleotide molecule as a selectable marker is the recovery of transformants containing a genetically linked gene, for example, conferring an improved agronomic trait. This ability was demonstrated by transforming soybean explants with pMON58498 having 2-DNAs: a first T-DNA having a DMO-encoding polynucleotide molecule and a second T-DNA having a CP4 gene for glyphosate tolerance. Transgenic plants selected on semi-solid medium were transferred to soil and assayed by Invader™ and/or Southern analysis to show the presence of DMO and CP4 nucleic acids.

While both the DMO and CP4-encoding polynucleotide molecules could be used as a selectable marker, it was shown that transformants comprising a CP4 transgene could be selected using dicamba selection alone. As shown in Table 6, all but one regenerated plant from each of the two treatments had both DMO and CP4 genes. This study therefore demonstrates the ability to use DMO as a selectable marker for the recovery of agronomic genes. It is understood that any genethat is genetically linked to a selectable DMO marker as introduced into a genome, e.g., present within 50 cM, can be selected in this manner and that such genes need not necessarily be introduced on the same vector.

27

TABLE 6

Use of DMO as a selectable marker in combination with an agronomic trait gene.						
Exp-Trt	BAP in delay medium (mg/l)	# Explants	# tolerant shoots harvested	# shoots rooted in soil	# plants with DMO gene (# plants assayed)	# plants with CP4 gene (# plants assayed)
566-1 & 567-1	1	1654	51	37	19 (37)	18 (37)
566-2 & 567-2	5	1800	35	11	3 (11)	2 (11)

Example 7

Tolerance of Plants Containing DMO-Encoding Polynucleotide Molecule to Other Auxin-Like Herbicides

An analysis was carried out to determine whether soybean plants having DMO-encoding polynucleotide could deactivate other auxin-like herbicides in addition to dicamba. This was carried out by applying various concentrations of commercially available formulations of other auxin-like herbicides such as 2,4-D (Helena, Collierville, Tenn.), MCPA (Agrilience, St. Paul, Minn.), triclopyr (GARLON 3A; Dow Elanco, Indianapolis, Ind.), clopyralid (STINGER; Dow Elanco, Indianapolis, Ind.), picloram (TORDON 22K; Dow Elanco, Indianapolis, Ind.), or Banvel or Clarity (BASF, Raleigh, N.C.) to DMO containing plant tissues or plants.

Transgenic soybean plants were obtained by Agrobacterium-mediated transformation of soybean explants with a DMO-encoding polynucleotide as described above for the events designated Events 1-4. A non-transgenic line was used as a control. Non-transgenic and transgenic soybean seeds were planted into 3,5-inch square plastic pots containing Redi-earth™ (Scotts-Sierra Horticultural Products Co., Marysville, Ohio). The pots were placed on capillary matting in 35 inch×60 inch fiberglass watering trays for overhead and/or sub-irrigation for the duration of the test period so as to maintain optimum soil moisture for plant growth. The pots were fertilized with Osmocote (14-14-14 slow release; Scotts-Sierra Horticultural Products Co., Marysville, Ohio) at the rate of 100 gm/cu.ft. to sustain plant growth for the duration of greenhouse trials. The plants were grown in greenhouses at 27°/21° C. day/night temperature, with relative humidity between 25%-75% to simulate warm season growing conditions of late spring. A 14 h minimum photoperiod was provided with supplemental light at about 600 µE as needed.

All herbicide applications were made with the track sprayer using a Teejet 9501E flat fan nozzle (Spraying Systems Co, Wheaton, Ill.) with air pressure set at a minimum of 24 psi (165 kpa). The spray nozzle was kept at a height of about 16 inches above the top of plant material for spraying. The spray volume was 10 gallons per acre or 93 liters per hectare. Applications were made when plants had reached V-3 stage. All trials were established in a randomized block design (randomized by rate) with 4 to 6 replications of each treatment depending on plant quality, availability and to account for any environmental variability that may have occurred within the confines of each greenhouse.

All treated plants in greenhouse trials were visually assessed at about 4, 14, 18, and 21 days after treatment (DAT) for injury on a scale of 0 to 100 percent relative to untreated

28

control plants, with zero representing “no” injury and 100% representing “complete” injury or death. Data were collected using a palm top computer and analyzed using standard statistical methods. The results shown in Table 9 clearly indicate tolerance of transgenic soybean to other auxin-like herbicides such as 2,4-D and MCPA relative to the non-transgenic line.

TABLE 7

Percentage injury relative to un-treated controls at 25 DAT post-V3 applications of different auxin-like herbicides to non-transgenic or transgenic soybean plants.*				
Herbicide	Plant/trial	280	561	1120
% injury at shown rates (g ae/ha**) at 21 DAT				
Dicamba (Clarity)	Non-transgenic		100	100
	Event 1		0.0	1.2
	Event 2		0.0	1.7
	Event 3		0.0	0.7
	Event 4		0.0	1.5
Dicamba (Banvel)	Non-transgenic		100.0	100.0
	Event 1		0.0	1.5
	Event 2		0.0	0.7
	Event 3		0.0	0.5
	Event 4		0.0	1.3
2,4-D	Non-transgenic	86.8	100.0	100.0
	Event 1	58.3	75.0	100.0
	Event 2	64.2	94.7	100.0
	Event 3	40.0	85.0	100.0
	Event 4	45.8	84.2	100.0
MCPA	Non-transgenic	93.0	98.3	100.0
	Event 1	72.5	99.3	100.0
	Event 2	55.0	95.0	99.7
	Event 3	55.0	95.8	100.0
	Event 4	88.3	98.8	100.0
LSD		16.3	10.6	3.7
% injury shown rates (g ae/ha**) at 14 DAT				
Triclopyr	Non-transgenic	86.7	97.3	98.7
	Event 1	88.3	95.7	99.3
	Event 2	86.7	98.7	99.3
	Event 3	86.7	94.0	96.3
	Event 4	90.8	98.0	99.2
Clopyralid	Non-transgenic	99.3	100.0	100.0
	Event 1	99.2	100.0	100.0
	Event 2	98.2	99.7	100.0
	Event 3	99.3	100.0	100.0
	Event 4	99.7	100.0	100.0
Picloram	Non-transgenic	99.3	100.0	100.0
	Event 1	99.7	100.0	100.0
	Event 2	99.3	100.0	100.0
	Event 3	99.3	99.7	100.0
	Event 4	99.3	100.0	100.0
LSD		2.9	1.8	1.4

*The % injury was represented as ANOVA mean comparisons.

**grams of active acid equivalent/hectare

This example shows that transgenic soybean plants exhibit tolerance to other auxin-like herbicides, indicating a likely common deactivation mechanism for dicamba and other auxin-like herbicides such as 2,4-D and MCPA. In case of triclopyr, clopyralid, and picloram, the application rate of 280 g ae/ha appeared too stringent in this study and thus lower concentrations may be desired in a most settings to reduce plant damage. The results indicate that auxin-like herbicides may be used for selecting plant cells transformed with DMO-encoding polynucleotide molecules, especially in case of plants that are very sensitive to dicamba, for example, cotton. The appropriate concentration of the auxin-like herbicide for selection under a given set of conditions may be optimized using a test grid of treatments followed by observation of treated plant tissues. An example of such a grid analyzes the

29

effect of concentrations of from about 0.001 mg/l to about 10 mg/l, including 0.01, 0.1, 1.0, 2.0, and 5.0 mg/L.

Another auxin-like herbicide Butyrac 200 (2,4-DB; Albaugh) was also tested on transgenic soybean plants carrying a DMO gene for testing the plants tolerance to it. The herbicide was applied as a post-emergence treatment at three application rates on two transgenic soybean events and compared with a non-transgenic line for total crop injury across all three application rates: 280 g/ha (0.25 lb/a), 561 g/ha (0.5 lb/a) and 841 g/ha (0.75 lb/a) (see Table 8). Both transgenic soybean lines showed low level of tolerance to 2,4-DB. This example demonstrates that dicamba tolerant soybean is also tolerant to low levels of 2,4-DB and should be useful in managing damage from spray drift from the same or neighboring fields to prevent crop losses, and would exhibit tolerance to residual levels of 2,4-DB following incomplete washing of herbicide delivery equipment.

TABLE 8

Percentage injury relative to the untreated control at 16 DAT by the application of 2,4-DB to non-transgenic or transgenic soybean plants.					
Herbicide	Plant	% injury at shown rates (g ae/ha) at 16 DAT			
		280	561	1120	
2,4-DB (Butyrac 200)	Non-transgenic NE3001	59.2	70.0	79.2	
	462-1-21	25.0	43.3	75.8	
	469-13-19	18.3	37.5	70.0	

Example 8

Use of DMO Gene as a Selectable Marker Against Other Auxin-Like Herbicides

Freshly isolated soybean explants (mature embryo axes without cotyledons) were inoculated with Agrobacterium strain ABI harboring pMON73691 (containing DMO and GFP genes). After 3-d co-culture with Agrobacterium at 23° C. and a photoperiod of 16-h light and 8-h dark, the explants were cultured in liquid delay medium which contained modified woody plant medium supplemented with 5 mg/l BAP, 200 mg/l carbenicillin, 200 mg/l cefotaxime and 100 mg/l

30

sterile microscope equipped for detecting GFP expressing tissues. At 48 days after inoculation (DAI), GFP-expressing (GFP+) buds or young shoots were observed on number of explants in the treatments with 0.01 mg/L dicamba, 0.01, 0.1 or 1.0 mg/L 2,4-D, but not on the explants treated with 2 mg/L 2,4-D. Extensive callus development was observed on the explants in treatments with 1 or 2 mg/L 2,4-D. In the treatment with 0.01 or 0.1 mg/L 2,4-D, the explants had extensive shoot growth, and a few had elongated GFP+ shoots.

TABLE 9

Summary of experiment using DMO as a selectable marker and 2,4-D as the selective agent.			
Treatment #	Selective agent and concentration	# Explants inoculated	# Explants w/GFP + buds/young shoots at 48DAI
710-1	0.01 mg/L dicamba (control)	375	28
710-2	0.01 mg/L 2,4-D	375	33
710-3	0.1 mg/L 2,4-D	375	19
710-4	1 mg/L 2,4-D	375	4
710-5	2 mg/L 2,4-D	375	0

Example 9

Selection of Dicamba-Tolerant Plants Transformed with DMO-Encoding Polynucleotide without a Delay Step

Transgenic plants with a DMO gene without a delay-to-selection step were produced in three studies. As an example, explants were infected and co-cultivated with Agrobacterium harboring pMON73696. After the co-culture period, the explants were cultured on liquid medium containing 5 mg/l BAP and 0.01 mg/l dicamba for 4 day, and then transferred onto to the liquid or semi-solid selection medium with 0.01 mg/l dicamba. As shown in Table 10, dicamba tolerant shoots could be obtained from the treatments (717-2 and 757-2) that utilized selection immediately after co-culture with Agrobacterium.

TABLE 10

Selection of dicamba-tolerant plants transformed with DMO-encoding polynucleotide without a delay step.						
Experimental Treatment	Number of days delayed to selection	Construct (pMON)	# Explants	# Plants moved to soil	# Plants assayed w/Invader	# Plants w/the gene
717-1	4	73696	608	15	15	4
717-2	0	73696	665	14	12	6
757-1	4	73696	542	13	11	6
757-2	0	73696	542	7	7	7

ticarcillin. The explants were in the delay medium for 4 days. They were then transferred to liquid selection medium in PLANTCONs. The selection medium was the same as the delay medium except of addition of various levels of 2,4-D (0.01, 0.1, 1.0 or 2.0 mg/L) or 0.01 mg/L dicamba as shown in Table 9 below. Each PLANTCON contained 60 ml of the selection medium and one piece of foam sponge with 5 slits. Twenty-five explants were evenly inserted into the slits. The cultures were maintained at 28° C. and a photoperiod of 16-h light and 8-h dark, and were examined periodically under a

Example 10

Use of DMO as a Selectable Marker for Arabidopsis

The susceptibility of Arabidopsis to different levels of dicamba was first tested. Wild type Arabidopsis var. Columbia seeds were plated on plant tissue culture medium containing various levels of dicamba. FIG. 8 shows that wild type Arabidopsis was quite susceptible to 1.0 mg/L in the culture medium. Arabidopsis plants were then transformed with the

constructs containing DMO polynucleotides using the floral dip method (Clough and Bent, 1998). R₁ seeds were plated on the culture medium containing up to 4 mg/L of dicamba. FIG. 9, for example, shows recovery of dicamba tolerant plants (shown by arrows) after transformation with pMON 73696. These dicamba tolerant plants were found to contain one or more copies of DMO nucleotide as ascertained by the Invader™ test. The example demonstrated the utility of DMO gene in producing dicamba tolerant plants of other plant species.

All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents that are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The references listed below are incorporated herein by reference to the extent that they supplement, explain, provide a background for, or teach methodology, techniques, and/or compositions employed herein.

U.S. Pat. No. 4,554,101; U.S. Pat. No. 4,940,838; U.S. Pat. No. 5,017,692; U.S. Pat. No. 5,229,114; U.S. Pat. No. 5,302,523; U.S. Pat. No. 5,304,730; U.S. Pat. No. 5,322,938; U.S. Pat. No. 5,352,605; U.S. Pat. No. 5,359,142; U.S. Pat. No. 5,362,865; U.S. Pat. No. 5,384,253; U.S. Pat. No. 5,445,962; U.S. Pat. No. 5,463,175; U.S. Pat. No. 5,464,763; U.S. Pat. No. 5,464,765; U.S. Pat. No. 5,491,084; U.S. Pat. No. 5,512,466; U.S. Pat. No. 5,516,671; U.S. Pat. No. 5,530,196; U.S. Pat. No. 5,538,877; U.S. Pat. No. 5,538,880; U.S. Pat. No. 5,543,576; U.S. Pat. No. 5,545,818; U.S. Pat. No. 5,550,318; U.S. Pat. No. 5,552,299; U.S. Pat. No. 5,563,055; U.S. Pat. No. 5,567,600; U.S. Pat. No. 5,567,862; U.S. Pat. No. 5,591,616; U.S. Pat. No. 5,608,149; U.S. Pat. No. 5,633,435; U.S. Pat. No. 5,641,876; U.S. Pat. No. 5,659,122; U.S. Pat. No. 5,689,041; U.S. Pat. No. 5,689,052; U.S. Pat. No. 5,716,837; U.S. Pat. No. 5,750,876; U.S. Pat. No. 5,763,241; U.S. Pat. No. 5,763,245; U.S. Pat. No. 5,773,696; U.S. Pat. No. 5,804,425; U.S. Pat. No. 5,837,848; U.S. Pat. No. 5,850,019; U.S. Pat. No. 5,850,023; U.S. Pat. No. 5,866,775; U.S. Pat. No. 5,869,720; U.S. Pat. No. 5,880,275; U.S. Pat. No. 5,942,658; U.S. Pat. No. 5,942,664; U.S. Pat. No. 5,958,745; U.S. Pat. No. 5,959,091; U.S. Pat. No. 5,981,834; U.S. Pat. No. 5,985,605; U.S. Pat. No. 5,998,700; U.S. Pat. No. 6,011,199; U.S. Pat. No. 6,013,864; U.S. Pat. No. 6,015,940; U.S. Pat. No. 6,023,013; U.S. Pat. No. 6,051,753; U.S. Pat. No. 6,063,597; U.S. Pat. No. 6,063,756; U.S. Pat. No. 6,072,103; U.S. Pat. No. 6,080,560; U.S. Pat. No. 6,093,695; U.S. Pat. No. 6,107,549; U.S. Pat. No. 6,110,464; U.S. Pat. No. 6,121,436; U.S. Pat. No. 6,140,075; U.S. Pat. No. 6,140,078; U.S. Pat. No. 6,146,826; U.S. Pat. No. 6,153,814; U.S. Pat. No. 6,156,573; U.S. Pat. No. 6,166,292; U.S. Pat. No. 6,171,640; U.S. Pat. No. 6,175,060; U.S. Pat. No. 6,177,611; U.S. Pat. No. 6,177,615; U.S. Pat. No. 6,215,048; U.S. Pat. No. 6,221,649; U.S.

Pat. No. 6,222,098; U.S. Pat. No. 6,225,114; U.S. Pat. No. 6,228,623; U.S. Pat. No. 6,228,992; U.S. Pat. No. 6,232,526; U.S. Pat. No. 6,235,971; U.S. Pat. No. 6,242,241; U.S. Pat. No. 6,248,536; U.S. Pat. No. 6,248,876; U.S. Pat. No. 6,252,138; U.S. Pat. No. 6,271,443; U.S. Pat. No. 6,281,016; U.S. Pat. No. 6,284,949; U.S. Pat. No. 6,294,714; U.S. Pat. No. 6,313,378; U.S. Pat. No. 6,316,407; U.S. Pat. No. 6,326,351; U.S. Pat. No. 6,372,211; U.S. Pat. No. 6,380,462; U.S. Pat. No. 6,380,466; U.S. Pat. No. 6,399,330; U.S. Pat. No. 6,423,828; U.S. Pat. No. 6,426,446; U.S. Pat. No. 6,426,447; U.S. Pat. No. 6,429,357; U.S. Pat. No. 6,429,362; U.S. Pat. No. 6,433,252; U.S. Pat. No. 6,437,217; U.S. Pat. No. 6,441,277; U.S. Pat. No. 6,444,876; U.S. Pat. No. 6,448,476; U.S. Pat. No. 6,459,018; U.S. Pat. No. 6,468,523; U.S. Pat. No. 6,476,295; U.S. Pat. No. 6,483,008; U.S. Pat. No. 6,489,461; U.S. Pat. No. 6,495,739; U.S. Pat. No. 6,501,009; U.S. Pat. No. 6,506,962; U.S. Pat. No. 6,518,488; U.S. Pat. No. 6,521,442; U.S. Pat. No. 6,531,648; U.S. Pat. No. 6,537,750; U.S. Pat. No. 6,537,756; U.S. Pat. No. 6,538,109; U.S. Pat. No. 6,538,178; U.S. Pat. No. 6,538,179; U.S. Pat. No. 6,538,181; U.S. Pat. No. 6,541,259; U.S. Pat. No. 6,555,655; U.S. Pat. No. 6,573,361; U.S. Pat. No. 6,576,818; U.S. Pat. No. 6,589,767; U.S. Pat. No. 6,593,293; U.S. Pat. No. 6,596,538; U.S. Pat. No. 6,608,241; U.S. Pat. No. 6,617,496; U.S. Pat. No. 6,620,988; U.S. Pat. No. 6,635,806; U.S. Pat. No. 6,639,054; U.S. Pat. No. 6,642,030; U.S. Pat. No. 6,645,497; U.S. Pat. No. 6,653,280; U.S. Pat. No. 6,653,530; U.S. Pat. No. 6,657,046; U.S. Pat. No. 6,660,849; U.S. Pat. No. 6,663,906; U.S. Pat. No. 6,686,452; U.S. Pat. No. 6,706,950; U.S. Pat. No. 6,713,063; U.S. Pat. No. 6,716,474; U.S. Pat. No. 6,723,837; U.S. Pat. No. 6,723,897; U.S. Pat. No. 6,770,465; U.S. Pat. No. 6,774,283; U.S. Pat. No. 6,803,501; U.S. Pat. No. 6,809,078; U.S. Pat. No. 6,812,379; U.S. Pat. No. 6,822,141; U.S. Pat. No. 6,828,475

U.S. Provisional Appln. Ser. No. 60/891,675

U.S. Patent Pub. 2005/0005321; U.S. Patent Pub. 2003/0115626; U.S. Patent Pub. 2003/01403641; U.S. Patent Pub. 2003/0135879; U.S. Patent Pub. 2003/0028917

U.S. patent application Ser. No. 09/757,089; U.S. patent application Ser. No. 11/749,583

U.S. Pat. RE37,543

U.S. Pat. RE38,446

Becker et al., *Plant Mol. Biol.*, 20:49, 1992.

Bevan et al., *NAR*, 11: 369, 1983.

Broothaerts et al., *Nature*, 433: 629-633, 2005.

Carrington and Freed, *J. Virology*, 64:1590, 1990.

Chalfie et al., *Science*, 263:802, 1994.

Chandler et al., *Plant Cell*, 1:1175-1183, 1989.

Charest et al., *Plant Cell Rep.*, 8:643, 1990.

Chu et al., *Scientia Sinica*, 18:659, 1975.

Clough and Bent, *Plant J.*, 16:735, 1998.

Comai et al., *Nature*, 317:741, 1985.

Cork and Khalil, *Adv. Appl. Microbiol.*, 40: 289, 1995.

Cork and Krueger, *Adv. Appl. Microbiol.*, 38:1, 1991.

Coruzzi et al., *EMBO J.*, 3: 1671, 1984.

Creissen et al., *Plant J.*, 2:129, 1991.

Crop Protection Reference, *Chemical & Pharmaceutical Press, Inc.*, NY, 11th Ed., 1803-1821,

De Block et al., *EMBO J.*, 3:1681, 1984.

De Block et al., *EMBO J.*, 6:2513, 1987.

della-Cioppa et al., *Proc. Natl. Acad. Sci. USA*, 83:6873-6877, 1986.

Depicker et al., *J. Mol. Appl. Genet.*, 1: 561, 1982.

Duncan et al., *Planta*, 165:322, 1985.

Ebert et al., *Proc. Natl. Acad. Sci. USA*, 84:5745-5749, 1987.

Eckes et al., *Mol. Gen. Genet.*, 205:14, 1986.

33

Eichholtz et al., Somatic Cell Mol. Genet., 13:67, 1987.
 EP Appln. 553494
 EP Appln. 646643
 Fraley et al., Proc. Natl. Acad. Sci. USA, 80:4803, 1983.
 Gamborg et al., Exp. Cell Res., 50:151, 1968.
 Gordon-Kamm et al., Plant Cell, 2:603, 1990.
 Haseloff et al., TIG, 11:328-329, 1995.
 Hayford et al., Plant Physiol., 86:1216, 1988.
 Hille et al., Plant Mol. Biol., 7:171, 1986.
 Horsch et al., Science, 227:1229, 1985.
 Ingelbrecht et al., Plant Cell, 1:671, 1989.
 Jefferson, Plant Mol. Biol. Rep., 5:387, 1987.
 Jones et al., Mol. Gen. Genet., 210:86, 1987.
 Kado, Crit. Rev. Plant. Sci., 10:1, 1991.
 Klee et al., Mol. Gen. Genet., 210:437-442, 1987.
 Koncz et al., Proc. Natl. Acad. Sci., USA, 84:131, 1987.
 Krueger et al., J. Agric. Food Chem., 37:534, 1989.
 Kyte and Doolittle, J. Mol. Biol., 157(1):105-132, 1982.
 Lawton et al., Plant Mol. Biol., 9:315-324, 1987.
 Linsmaier and Skoog, Physio. Plant, 18:100, 1965.
 Lloyd and McCown, Proc. Int. Plant Prop. Soc., 30:421, 1981.
 Maniatis, et al., Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, N.Y., 1982.
 McCown and Lloyd, Hort. Science, 16:453, 1981.
 Miki et al., In: Methods in Plant Molecular Biology and Biotechnology, Glick and Thompson (Eds.), CRC Press, 67-88, 1993.

34

Moloney et al., Plant Cell Reports, 8:238, 1989.
 Murashige and Skoog, Physiol. Plant, 15:473-497, 1962.
 Nitsch and Nitsch, Science, 163:85, 1969.
 Odell et al., Nature, 313:810-812, 1985.
 5 PCT Appln. WO 97/11086
 PCT Appln. WO 95/24492
 PCT Appln. WO 97/31115
 PCT Appln. WO 97/41228
 10 Rojiyaa et al., 1987 (JP 1987201527-A), GenBank Accession E01312.
 Sambrook et al., In: Molecular cloning: a laboratory manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.
 15 Schenk and Hildebrandt, Can. J. Bot., 50:199, 1972.
 Shah et al., Science, 233:478, 1986.
 Southern, Mol. Biol., 98:503, 1975.
 Stalker et al., Science, 242:419, 1988.
 Streber and Willmitzer, Bio/Technology, 7:811, 1989.
 20 Svab et al., Plant Mol. Biol., 14:197, 1990.
 Teeri et al., EMBO J., 8:343, 1989.
 Turner and Foster, Molec. Biotechn., 3:225, 1995.
 Uchimiya and Murashige, Plant Physiol., 15:473, 1962.
 Vanden Elzen et al., Plant Mol. Biol., 5:299, 1985.
 25 Walker et al., Proc. Natl. Acad. Sci. USA, 84:6624-6628, 1987.
 Yang et al., Proc. Natl. Acad. Sci. USA, 87:4144-4148, 1990.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 29

<210> SEQ ID NO 1

<211> LENGTH: 1023

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Based on dicamba monooxygenase gene from *Pseudomonas maltophilia*

<400> SEQUENCE: 1

```

atggccactt tcgtagaaaa cgcttggtac gttgctgcac ttctgagga gttgagcgag      60
aagcctctag gaagaactat cctcgatact ccactagetc tctatcgta acctgacgga      120
gttgctcgctg ccctgcttga tatttgctcg catcgcttcg ctccgttgag tgacgggtatt      180
ctagtcaacg gacatctcca gtgtccatat cacggctctg aatttgacgg aggtggccag      240
tgtgtccaca acccgacagg caacggagcc cgccctgctt ctctgaacgt gcgateatcc      300
cctgtcgctg aaagagacgc attgatctgg atctgccctg gagatccagc actcgacagat      360
cccgtgctta tcctgactt tgggtgtcgt gttgatccag cttaccgtac tgcgcgaggt      420
tacggtcacg tggactgcaa ctacaagetc cttgtggata acctcatgga tcttgacac      480
gtcagtagcg tgcacgcgcg taacgccccaa acagacgcct tcgatagact tgagcgtgag      540
gtgatcgctg gcgacggcga gatccaggcg ctcatgaaga tccttggtgg cacaccctca      600
gttctcatgg ctaagttctt gcgtggtgct aacacaccag ttgacgcctg gaacgacatc      660
cgggtgaata aggtgtcggc tatgctgaac ttcatcgcgg tcgcgcggga agggacgcgg      720
aaggagcagt caatccactc ccgaggaacc catatcctta ctccctgagac cgaggcaagc      780
tgccattact tcttcggtag ttcccgaac ttcggtatag acgatccaga gatggacggt      840
gttctcagga gctggcaagc tcaagccctg gtgaaggagg acaaagtggc cggtgaagct      900
atcgaaagcg ggagggttta cgtcgaagcg aacgggatca gaaccgccat gttgtcctgc      960

```

-continued

gacgaggcag ccgtcagggt atccaggag attgagaagc tcgaacaact agaagcggcg 1020

tga 1023

<210> SEQ ID NO 2

<211> LENGTH: 340

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
Pseudomonas maltophilia

<400> SEQUENCE: 2

Met Ala Thr Phe Val Arg Asn Ala Trp Tyr Val Ala Ala Leu Pro Glu
1 5 10 15Glu Leu Ser Glu Lys Pro Leu Gly Arg Thr Ile Leu Asp Thr Pro Leu
20 25 30Ala Leu Tyr Arg Gln Pro Asp Gly Val Val Ala Ala Leu Leu Asp Ile
35 40 45Cys Pro His Arg Phe Ala Pro Leu Ser Asp Gly Ile Leu Val Asn Gly
50 55 60His Leu Gln Cys Pro Tyr His Gly Leu Glu Phe Asp Gly Gly Gly Gln
65 70 75 80Cys Val His Asn Pro His Gly Asn Gly Ala Arg Pro Ala Ser Leu Asn
85 90 95Val Arg Ser Phe Pro Val Val Glu Arg Asp Ala Leu Ile Trp Ile Cys
100 105 110Pro Gly Asp Pro Ala Leu Ala Asp Pro Gly Ala Ile Pro Asp Phe Gly
115 120 125Cys Arg Val Asp Pro Ala Tyr Arg Thr Val Gly Gly Tyr Gly His Val
130 135 140Asp Cys Asn Tyr Lys Leu Leu Val Asp Asn Leu Met Asp Leu Gly His
145 150 155 160Ala Gln Tyr Val His Arg Ala Asn Ala Gln Thr Asp Ala Phe Asp Arg
165 170 175Leu Glu Arg Glu Val Ile Val Gly Asp Gly Glu Ile Gln Ala Leu Met
180 185 190Lys Ile Pro Gly Gly Thr Pro Ser Val Leu Met Ala Lys Phe Leu Arg
195 200 205Gly Ala Asn Thr Pro Val Asp Ala Trp Asn Asp Ile Arg Trp Asn Lys
210 215 220Val Ser Ala Met Leu Asn Phe Ile Ala Val Ala Pro Glu Gly Thr Pro
225 230 235 240Lys Glu Gln Ser Ile His Ser Arg Gly Thr His Ile Leu Thr Pro Glu
245 250 255Thr Glu Ala Ser Cys His Tyr Phe Phe Gly Ser Ser Arg Asn Phe Gly
260 265 270Ile Asp Asp Pro Glu Met Asp Gly Val Leu Arg Ser Trp Gln Ala Gln
275 280 285Ala Leu Val Lys Glu Asp Lys Val Val Val Glu Ala Ile Glu Arg Arg
290 295 300Arg Ala Tyr Val Glu Ala Asn Gly Ile Arg Pro Ala Met Leu Ser Cys
305 310 315 320Asp Glu Ala Ala Val Arg Val Ser Arg Glu Ile Glu Lys Leu Glu Gln
325 330 335

-continued

Leu Glu Ala Ala
340

<210> SEQ ID NO 3
<211> LENGTH: 1023
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
Pseudomonas maltophilia

<400> SEQUENCE: 3

```

atgctcactt tcgtagaaa cgcttggtac gttgctgcac ttcctgagga gttgagcgag    60
aagcctctag gaagaactat cctcgatact ccactagctc tctatcgtea acctgacgga    120
gttgctcgctg ccctgcttga tatttgctcg catcgcttcg ctccgttgag tgacgggtatt    180
ctagtcaacg gacatctcca gtgtccatat cacggtctgg aatttgacgg aggtggccag    240
tgtgtccaca acccgcaagg caacggagcc cgccctgctt ctctgaacgt gcgatcattc    300
cctgtcgtgg aaagagacgc attgatctgg atctgccctg gagatccagc actcgagat    360
cccggtgcta tccctgactt tgggtgtcgt gttgatccag cttaccgtac tgcggaggt    420
tacggtcacg tggactgcaa ctacaagctc cttgtggata acctcatgga tcttgacac    480
gctcagtagc tgcaccgcgc taacgcccaa acagacgcct tcgatagact tgagcgtgag    540
gtgatcggtg gcgacggcga gatccaggcg ctcatgaaga tccctggtgg cacaccctca    600
gttctcatgg ctaagttctt gcgtggtgct aacacaccag ttgacgcctg gaacgacatc    660
cggtggaata aggtgtcggc tatgtgaac ttcatcgcgg tcgcgccgga agggacgccg    720
aaggagcagt caatccactc ccgaggaacc catatcctta ctctgagac cgaggcaagc    780
tgccattact tcttcggtag tccccgaac ttcggtatag acgatccaga gatggacggt    840
gttctcagga gctggcaagc tcaagccctg gtgaaggagg acaaagtggc cgttgaagct    900
atcgaaaggc ggagggctta cgtcgaagcg aacgggatca gacccgccat gttgtcctgc    960
gacgaggcag ccgtcagggt atccagggag attgagaagc tcgaacaact agaagcggcg    1020
tga                                                                    1023

```

<210> SEQ ID NO 4
<211> LENGTH: 340
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
Pseudomonas maltophilia

<400> SEQUENCE: 4

```

Met Leu Thr Phe Val Arg Asn Ala Trp Tyr Val Ala Ala Leu Pro Glu
1           5           10          15

Glu Leu Ser Glu Lys Pro Leu Gly Arg Thr Ile Leu Asp Thr Pro Leu
20          25          30

Ala Leu Tyr Arg Gln Pro Asp Gly Val Val Ala Ala Leu Leu Asp Ile
35          40          45

Cys Pro His Arg Phe Ala Pro Leu Ser Asp Gly Ile Leu Val Asn Gly
50          55          60

His Leu Gln Cys Pro Tyr His Gly Leu Glu Phe Asp Gly Gly Gly Gln
65          70          75          80

Cys Val His Asn Pro His Gly Asn Gly Ala Arg Pro Ala Ser Leu Asn
85          90          95

```

-continued

Val Arg Ser Phe Pro Val Val Glu Arg Asp Ala Leu Ile Trp Ile Cys
 100 105 110
 Pro Gly Asp Pro Ala Leu Ala Asp Pro Gly Ala Ile Pro Asp Phe Gly
 115 120 125
 Cys Arg Val Asp Pro Ala Tyr Arg Thr Val Gly Gly Tyr Gly His Val
 130 135 140
 Asp Cys Asn Tyr Lys Leu Leu Val Asp Asn Leu Met Asp Leu Gly His
 145 150 155 160
 Ala Gln Tyr Val His Arg Ala Asn Ala Gln Thr Asp Ala Phe Asp Arg
 165 170 175
 Leu Glu Arg Glu Val Ile Val Gly Asp Gly Glu Ile Gln Ala Leu Met
 180 185 190
 Lys Ile Pro Gly Gly Thr Pro Ser Val Leu Met Ala Lys Phe Leu Arg
 195 200 205
 Gly Ala Asn Thr Pro Val Asp Ala Trp Asn Asp Ile Arg Trp Asn Lys
 210 215 220
 Val Ser Ala Met Leu Asn Phe Ile Ala Val Ala Pro Glu Gly Thr Pro
 225 230 235 240
 Lys Glu Gln Ser Ile His Ser Arg Gly Thr His Ile Leu Thr Pro Glu
 245 250 255
 Thr Glu Ala Ser Cys His Tyr Phe Phe Gly Ser Ser Arg Asn Phe Gly
 260 265 270
 Ile Asp Asp Pro Glu Met Asp Gly Val Leu Arg Ser Trp Gln Ala Gln
 275 280 285
 Ala Leu Val Lys Glu Asp Lys Val Val Val Glu Ala Ile Glu Arg Arg
 290 295 300
 Arg Ala Tyr Val Glu Ala Asn Gly Ile Arg Pro Ala Met Leu Ser Cys
 305 310 315 320
 Asp Glu Ala Ala Val Arg Val Ser Arg Glu Ile Glu Lys Leu Glu Gln
 325 330 335
 Leu Glu Ala Ala
 340

<210> SEQ ID NO 5
 <211> LENGTH: 1023
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
 Pseudomonas maltophilia

<400> SEQUENCE: 5

atgctcactt tcgtagaaa cgcttggtac gttgctgcac ttctgagga gttgagcgag	60
aagcctctag gaagaactat cctcgatact ccactagetc tctatcgtea acctgacgga	120
gttgtecgctg ccctgcttga tatttgctcg catcgcttcg ctccgttgag tgacgggtatt	180
ctagtcaacg gacatctcca gtgtccatat cacggtctgg aatttgacgg aggtggccag	240
tggtgccaca acccgacagg caacggagcc cgccctgctt ctctgaacgt gcgatcattc	300
cctgtcgtgg aaagagacgc attgatctgg atctggcctg gagatccagc actcgagat	360
cccgggtgcta tccctgactt tgggtgtcgt gttgatccag cttaccgtac tgctggaggt	420
tacggtcacg tggactgcaa ctacaagctc cttgtggata acctcatgga tcttgacac	480
gctcagtagc tgcaccgcgc taacgcccaa acagacgcct tcgatagact tgagcgtgag	540
gtgatcggtg ggcacggcga gatccaggcg ctcatagaaga tccctgggtg cacacctca	600

-continued

```

gtttctcatgg ctaagttctt gcgtgggtgct aacacaccag ttgacgcctg gaacgacatc 660
cggtggaata aggtgtcggc tatgtgaac ttcacgcgg tcgcgccgga agggacgccg 720
aaggagcagt caatccactc ccgaggaacc catatcctta ctctgagac cgaggcaagc 780
tgccattact tcttcggtag tccccgaac ttcggtatag acgatccaga gatggacggt 840
gtttctcagga gctggcaagc tcaagccctg gtgaaggagg acaaagtggc cgttgaagct 900
atcgaaagcg ggaggcctta cgtcgaagcg aacgggatca gaccgcccat gttgtcctgc 960
gacgaggcag ccgtcagggt atccagggag attgagaagc tcgaacaact agaagcggcg 1020
tga 1023

```

```

<210> SEQ ID NO 6
<211> LENGTH: 340
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
Pseudomonas maltophilia

```

```

<400> SEQUENCE: 6

```

```

Met Leu Thr Phe Val Arg Asn Ala Trp Tyr Val Ala Ala Leu Pro Glu
1          5          10          15
Glu Leu Ser Glu Lys Pro Leu Gly Arg Thr Ile Leu Asp Thr Pro Leu
20         25         30
Ala Leu Tyr Arg Gln Pro Asp Gly Val Val Ala Ala Leu Leu Asp Ile
35         40         45
Cys Pro His Arg Phe Ala Pro Leu Ser Asp Gly Ile Leu Val Asn Gly
50         55         60
His Leu Gln Cys Pro Tyr His Gly Leu Glu Phe Asp Gly Gly Gly Gln
65         70         75         80
Cys Val His Asn Pro His Gly Asn Gly Ala Arg Pro Ala Ser Leu Asn
85         90         95
Val Arg Ser Phe Pro Val Val Glu Arg Asp Ala Leu Ile Trp Ile Trp
100        105        110
Pro Gly Asp Pro Ala Leu Ala Asp Pro Gly Ala Ile Pro Asp Phe Gly
115        120        125
Cys Arg Val Asp Pro Ala Tyr Arg Thr Val Gly Gly Tyr Gly His Val
130        135        140
Asp Cys Asn Tyr Lys Leu Leu Val Asp Asn Leu Met Asp Leu Gly His
145        150        155        160
Ala Gln Tyr Val His Arg Ala Asn Ala Gln Thr Asp Ala Phe Asp Arg
165        170        175
Leu Glu Arg Glu Val Ile Val Gly Asp Gly Glu Ile Gln Ala Leu Met
180        185        190
Lys Ile Pro Gly Gly Thr Pro Ser Val Leu Met Ala Lys Phe Leu Arg
195        200        205
Gly Ala Asn Thr Pro Val Asp Ala Trp Asn Asp Ile Arg Trp Asn Lys
210        215        220
Val Ser Ala Met Leu Asn Phe Ile Ala Val Ala Pro Glu Gly Thr Pro
225        230        235        240
Lys Glu Gln Ser Ile His Ser Arg Gly Thr His Ile Leu Thr Pro Glu
245        250        255
Thr Glu Ala Ser Cys His Tyr Phe Phe Gly Ser Ser Arg Asn Phe Gly
260        265        270
Ile Asp Asp Pro Glu Met Asp Gly Val Leu Arg Ser Trp Gln Ala Gln

```

-continued

275	280	285	
Ala Leu Val Lys Glu Asp Lys Val Val Val Glu Ala Ile Glu Arg Arg			
290	295	300	
Arg Ala Tyr Val Glu Ala Asn Gly Ile Arg Pro Ala Met Leu Ser Cys			
305	310	315	320
Asp Glu Ala Ala Val Arg Val Ser Arg Glu Ile Glu Lys Leu Glu Gln			
	325	330	335
Leu Glu Ala Ala			
340			

<210> SEQ ID NO 7
 <211> LENGTH: 1023
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
 Pseudomonas maltophilia

<400> SEQUENCE: 7

atggccacct tcgtccgcaa tgccctggtat gtggcggcgc tgcccagga actgtccgaa	60
aagccgctcg gccggacgat tctcgacaca ccgctcgcgc tctaccgcca gcccgacggt	120
gtggtcgcgg cgctgctcga catctgtccg caccgcttcg cgccgctgag cgacggcatc	180
ctcgtcaacg gccatctcca atgccctat cacgggctgg aattcgatgg cgcgggcgag	240
tgctccata acccgcacgg caatggcgcc cgcccggtt cgctcaacgt ccgtcccttc	300
ccggtggtgg agcgcgacgc gctgatctgg atctgtccc gcgatccggc gctggccgat	360
cctggggcga tccccgaatt cggtgcgcgc gtgatcccg cctatcggac cgtcggcggc	420
tatgggcatg tcgactgcaa ctacaagctg ctggtcgaca acctgatgga cctcggccac	480
gcccaatatg tccatcgcgc caacgcccag accgacgcct tcgaccggt ggagcgcgag	540
gtgatcgtcg gcgacggtga gatacaggcg ctgatgaaga ttcccggcgg cagcccgagc	600
gtgctgatgg ccaagttcct gcgcggcgcc aatacccccg tcgacgcttg gaacgacatc	660
cgctggaaca aggtgagcgc gatgtcaac ttcatcgcgg tggcgccgga aggcaccccg	720
aaggagcaga gcatccactc gcgcggtacc catatcctga ccccgagac ggaggcgagc	780
tgccattatt tcttcggctc ctgcgcgaat ttcggcatcg acgatccgga gatggacggc	840
gtgctgcgca gctggcaggc tcaggcgctg gtcaaggagg acaaggctgt cgtcgaggcg	900
atcgagcgcc gccgcgccta tgctgaggcg aatggcatcc gccggcgat gctgtcgtgc	960
gacgaagccg cagtcctgtg cagccgcgag atcgagaagc ttgagcagct cgaagccgcc	1020
tga	1023

<210> SEQ ID NO 8
 <211> LENGTH: 340
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
 Pseudomonas maltophilia

<400> SEQUENCE: 8

Met Ala Thr Phe Val Arg Asn Ala Trp Tyr Val Ala Ala Leu Pro Glu	
1	15
Glu Leu Ser Glu Lys Pro Leu Gly Arg Thr Ile Leu Asp Thr Pro Leu	
20	30
Ala Leu Tyr Arg Gln Pro Asp Gly Val Val Ala Ala Leu Leu Asp Ile	

-continued

35	40	45
Cys Pro His Arg Phe Ala Pro Leu Ser Asp Gly Ile Leu Val Asn Gly		
50	55	60
His Leu Gln Cys Pro Tyr His Gly Leu Glu Phe Asp Gly Gly Gly Gln		
65	70	75 80
Cys Val His Asn Pro His Gly Asn Gly Ala Arg Pro Ala Ser Leu Asn		
	85	90 95
Val Arg Ser Phe Pro Val Val Glu Arg Asp Ala Leu Ile Trp Ile Cys		
	100	105 110
Pro Gly Asp Pro Ala Leu Ala Asp Pro Gly Ala Ile Pro Asp Phe Gly		
	115	120 125
Cys Arg Val Asp Pro Ala Tyr Arg Thr Val Gly Gly Tyr Gly His Val		
	130	135 140
Asp Cys Asn Tyr Lys Leu Leu Val Asp Asn Leu Met Asp Leu Gly His		
	145	150 155 160
Ala Gln Tyr Val His Arg Ala Asn Ala Gln Thr Asp Ala Phe Asp Arg		
	165	170 175
Leu Glu Arg Glu Val Ile Val Gly Asp Gly Glu Ile Gln Ala Leu Met		
	180	185 190
Lys Ile Pro Gly Gly Thr Pro Ser Val Leu Met Ala Lys Phe Leu Arg		
	195	200 205
Gly Ala Asn Thr Pro Val Asp Ala Trp Asn Asp Ile Arg Trp Asn Lys		
	210	215 220
Val Ser Ala Met Leu Asn Phe Ile Ala Val Ala Pro Glu Gly Thr Pro		
	225	230 235 240
Lys Glu Gln Ser Ile His Ser Arg Gly Thr His Ile Leu Thr Pro Glu		
	245	250 255
Thr Glu Ala Ser Cys His Tyr Phe Phe Gly Ser Ser Arg Asn Phe Gly		
	260	265 270
Ile Asp Asp Pro Glu Met Asp Gly Val Leu Arg Ser Trp Gln Ala Gln		
	275	280 285
Ala Leu Val Lys Glu Asp Lys Val Val Val Glu Ala Ile Glu Arg Arg		
	290	295 300
Arg Ala Tyr Val Glu Ala Asn Gly Ile Arg Pro Ala Met Leu Ser Cys		
	305	310 315 320
Asp Glu Ala Ala Val Arg Val Ser Arg Glu Ile Glu Lys Leu Glu Gln		
	325	330 335
Leu Glu Ala Ala		
	340	

<210> SEQ ID NO 9

<211> LENGTH: 1023

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
Pseudomonas maltophilia

<400> SEQUENCE: 9

atggccactt tcgtagaaa cgcttggtac gttgctgcac ttcctgagga gttgagcgag	60
aagcctctag gaagaactat cctcgatact ccactagctc tctatcgtea acctgacgga	120
gtgtgcgctg ccctgcttga tatttgctcg catcgcttcg ctccggttag tgacggattt	180
ctagtcaacg gacatctcca gtgtccatat cacggtcttg aatttgacgg aggtggccag	240
tgtgtccaca acccgacgag caacggagcc cgcctgctt ctctgaacgt gcgatcattc	300

-continued

```

cctgtcgtgg aaagagacgc attgatctgg atctggcctg gagatccagc actcgcagat 360
cccggtgcta tccctgactt tgggtgtcgt gttgatccag cttaccgtac tgcgggaggt 420
tacggtcacg tggactgcaa ctacaagctc cttgtggata acctcatgga tcttggaacac 480
gtcagtaacg tgcaccgcgc taacgcccaa acagacgcct tcgatagact tgagcgtgag 540
gtgatcgttg gcgacggcga gatccaggcg ctcatgaaga tccctggtgg cacaccctca 600
gttctcatgg ctaagttctt gcgtggtgct aacacaccag ttgacgcctg gaacgacatc 660
cggtggaata aggtgtcggc tatgtgaac ttcacgcggg tcgcgccgga agggacgcgg 720
aaggagcagt caatccactc ccgaggaacc catatcctta ctctgagac cgaggcaagc 780
tgccattact tcttcggtag tccccgaac ttcggtatag acgatccaga gatggacggt 840
gttctcagga gctggcaagc tcaagccctg gtgaaggagg acaaagtggc cgttgaagct 900
atcgaaaggc ggagggtcta cgtcgaagcg aacgggatca gacccgccat gttgtcctgc 960
gacgaggcag ccgtcagggt atccagggag attgagaagc tcgaacaact agaagcggcg 1020
tga 1023

```

```

<210> SEQ ID NO 10
<211> LENGTH: 340
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Based on dicamba monooxygenase gene from
Pseudomonas maltophilia

```

```

<400> SEQUENCE: 10

```

```

Met Ala Thr Phe Val Arg Asn Ala Trp Tyr Val Ala Ala Leu Pro Glu
1      5      10      15
Glu Leu Ser Glu Lys Pro Leu Gly Arg Thr Ile Leu Asp Thr Pro Leu
20     25     30
Ala Leu Tyr Arg Gln Pro Asp Gly Val Val Ala Ala Leu Leu Asp Ile
35     40     45
Cys Pro His Arg Phe Ala Pro Leu Ser Asp Gly Ile Leu Val Asn Gly
50     55     60
His Leu Gln Cys Pro Tyr His Gly Leu Glu Phe Asp Gly Gly Gly Gln
65     70     75     80
Cys Val His Asn Pro His Gly Asn Gly Ala Arg Pro Ala Ser Leu Asn
85     90     95
Val Arg Ser Phe Pro Val Val Glu Arg Asp Ala Leu Ile Trp Ile Trp
100    105    110
Pro Gly Asp Pro Ala Leu Ala Asp Pro Gly Ala Ile Pro Asp Phe Gly
115    120    125
Cys Arg Val Asp Pro Ala Tyr Arg Thr Val Gly Gly Tyr Gly His Val
130    135    140
Asp Cys Asn Tyr Lys Leu Leu Val Asp Asn Leu Met Asp Leu Gly His
145    150    155    160
Ala Gln Tyr Val His Arg Ala Asn Ala Gln Thr Asp Ala Phe Asp Arg
165    170    175
Leu Glu Arg Glu Val Ile Val Gly Asp Gly Glu Ile Gln Ala Leu Met
180    185    190
Lys Ile Pro Gly Gly Thr Pro Ser Val Leu Met Ala Lys Phe Leu Arg
195    200    205
Gly Ala Asn Thr Pro Val Asp Ala Trp Asn Asp Ile Arg Trp Asn Lys
210    215    220

```

-continued

Val Ser Ala Met Leu Asn Phe Ile Ala Val Ala Pro Glu Gly Thr Pro
 225 230 235 240

Lys Glu Gln Ser Ile His Ser Arg Gly Thr His Ile Leu Thr Pro Glu
 245 250 255

Thr Glu Ala Ser Cys His Tyr Phe Phe Gly Ser Ser Arg Asn Phe Gly
 260 265 270

Ile Asp Asp Pro Glu Met Asp Gly Val Leu Arg Ser Trp Gln Ala Gln
 275 280 285

Ala Leu Val Lys Glu Asp Lys Val Val Val Glu Ala Ile Glu Arg Arg
 290 295 300

Arg Ala Tyr Val Glu Ala Asn Gly Ile Arg Pro Ala Met Leu Ser Cys
 305 310 315 320

Asp Glu Ala Ala Val Arg Val Ser Arg Glu Ile Glu Lys Leu Glu Gln
 325 330 335

Leu Glu Ala Ala
 340

<210> SEQ ID NO 11
 <211> LENGTH: 1020
 <212> TYPE: DNA
 <213> ORGANISM: Pseudomonas maltophilia

<400> SEQUENCE: 11

```

atgaccttcg tccgcaatgc ctggtatgtg gcggcgctgc ccgaggaact gtccgaaaag    60
ccgctcggcc ggacgattct cgacacaccg ctgcgcgtct accgccagcc cgacggtgtg    120
gtcgcggcgc tgctcgacat ctgtccgcac cgcttcgcgc cgctgagcga cggcaccctc    180
gtcaacggcc atctccaatg cccctatcac gggctggaat tcgatggcgg cgggcagtgc    240
gtccataacc cgcacggcaa tggcgcccgc cgggcttcgc tcaacgtccg ctccctcccg    300
gtggtggagc gcgacgcgct gatctggatc tggcccggcg atccggcgct ggccgaccc    360
ggggcgatcc ccgacttcgg ctgccgcgtc gatccgcct atccggaccgt cggcggttat    420
gggcgatgtc actgcaacta caagctgctg gtcgacaacc tgatggacct cggccacgcc    480
caatatgtcc atcgcgccaa cgcccagacc gacgccttcg accggctgga gcgcgaggtg    540
atcgtcggcg acggtgagat acaggcgctg atgaagattc ccggcggcac gccgagcgtg    600
ctgatggcca agttcctcgc cggcgccaat acccccgtcg acgcttgga cgcacatccgc    660
tggaacaagg tgagcgcgat gctcaacttc atcgcggtgg cgccggaagg caccgccgaag    720
gagcagagca tccactcgcg cggtaccat atcctgaccc ccgagacgga ggcgagctgc    780
cattatttct tcggctcctc gcgcaatttc ggcatcgacg atccggagat ggacggcgtg    840
ctgcgcagct ggcaggetca ggcgctggtc aaggaggaca aggtcgtcgt cgaggcgatc    900
gagcgccgcc ggcctatgt cgaggcgaat ggcacccgcc cggcgatgct gtcgtgcgac    960
gaagcgcgag tccgtgtcag ccgcgagatc gagaagcttg agcagctcga agccgcctga   1020

```

<210> SEQ ID NO 12
 <211> LENGTH: 339
 <212> TYPE: PRT
 <213> ORGANISM: Pseudomonas maltophilia

<400> SEQUENCE: 12

Met Thr Phe Val Arg Asn Ala Trp Tyr Val Ala Ala Leu Pro Glu Glu
 1 5 10 15

Leu Ser Glu Lys Pro Leu Gly Arg Thr Ile Leu Asp Thr Pro Leu Ala

-continued

20					25					30					
Leu	Tyr	Arg	Gln	Pro	Asp	Gly	Val	Val	Ala	Ala	Leu	Leu	Asp	Ile	Cys
		35					40					45			
Pro	His	Arg	Phe	Ala	Pro	Leu	Ser	Asp	Gly	Ile	Leu	Val	Asn	Gly	His
	50					55					60				
Leu	Gln	Cys	Pro	Tyr	His	Gly	Leu	Glu	Phe	Asp	Gly	Gly	Gly	Gln	Cys
65						70					75				80
Val	His	Asn	Pro	His	Gly	Asn	Gly	Ala	Arg	Pro	Ala	Ser	Leu	Asn	Val
				85					90					95	
Arg	Ser	Phe	Pro	Val	Val	Glu	Arg	Asp	Ala	Leu	Ile	Trp	Ile	Trp	Pro
			100					105					110		
Gly	Asp	Pro	Ala	Leu	Ala	Asp	Pro	Gly	Ala	Ile	Pro	Asp	Phe	Gly	Cys
		115					120					125			
Arg	Val	Asp	Pro	Ala	Tyr	Arg	Thr	Val	Gly	Gly	Tyr	Gly	His	Val	Asp
		130					135					140			
Cys	Asn	Tyr	Lys	Leu	Leu	Val	Asp	Asn	Leu	Met	Asp	Leu	Gly	His	Ala
145						150					155				160
Gln	Tyr	Val	His	Arg	Ala	Asn	Ala	Gln	Thr	Asp	Ala	Phe	Asp	Arg	Leu
				165					170					175	
Glu	Arg	Glu	Val	Ile	Val	Gly	Asp	Gly	Glu	Ile	Gln	Ala	Leu	Met	Lys
			180					185					190		
Ile	Pro	Gly	Gly	Thr	Pro	Ser	Val	Leu	Met	Ala	Lys	Phe	Leu	Arg	Gly
		195					200					205			
Ala	Asn	Thr	Pro	Val	Asp	Ala	Trp	Asn	Asp	Ile	Arg	Trp	Asn	Lys	Val
		210					215					220			
Ser	Ala	Met	Leu	Asn	Phe	Ile	Ala	Val	Ala	Pro	Glu	Gly	Thr	Pro	Lys
225						230					235				240
Glu	Gln	Ser	Ile	His	Ser	Arg	Gly	Thr	His	Ile	Leu	Thr	Pro	Glu	Thr
				245					250					255	
Glu	Ala	Ser	Cys	His	Tyr	Phe	Phe	Gly	Ser	Ser	Arg	Asn	Phe	Gly	Ile
			260					265					270		
Asp	Asp	Pro	Glu	Met	Asp	Gly	Val	Leu	Arg	Ser	Trp	Gln	Ala	Gln	Ala
		275					280					285			
Leu	Val	Lys	Glu	Asp	Lys	Val	Val	Val	Glu	Ala	Ile	Glu	Arg	Arg	Arg
		290					295					300			
Ala	Tyr	Val	Glu	Ala	Asn	Gly	Ile	Arg	Pro	Ala	Met	Leu	Ser	Cys	Asp
305						310					315				320
Glu	Ala	Ala	Val	Arg	Val	Ser	Arg	Glu	Ile	Glu	Lys	Leu	Glu	Gln	Leu
				325					330					335	
Glu	Ala	Ala													

<210> SEQ ID NO 13
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Primer

<400> SEQUENCE: 13

acggacgcgg agatgctcaa cttcatcgc

29

<210> SEQ ID NO 14
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Primer

<400> SEQUENCE: 14

tccgctggaa caaggtgagc gcgt 24

<210> SEQ ID NO 15

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Primer

<400> SEQUENCE: 15

gtcgtgccc tgcttgatat t 21

<210> SEQ ID NO 16

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Primer

<400> SEQUENCE: 16

cgccgcttct agttgttc 18

<210> SEQ ID NO 17

<211> LENGTH: 57

<212> TYPE: PRT

<213> ORGANISM: Pisum sativum

<400> SEQUENCE: 17

Met Ala Ser Met Ile Ser Ser Ser Ala Val Thr Thr Val Ser Arg Ala
 1 5 10 15

Ser Arg Gly Gln Ser Ala Ala Met Ala Pro Phe Gly Gly Leu Lys Ser
 20 25 30

Met Thr Gly Phe Pro Val Arg Lys Val Asn Thr Asp Ile Thr Ser Ile
 35 40 45

Thr Ser Asn Gly Gly Arg Val Lys Cys
 50 55

<210> SEQ ID NO 18

<211> LENGTH: 85

<212> TYPE: PRT

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 18

Met Ala Ser Ser Met Leu Ser Ser Ala Thr Met Val Ala Ser Pro Ala
 1 5 10 15

Gln Ala Thr Met Val Ala Pro Phe Asn Gly Leu Lys Ser Ser Ala Ala
 20 25 30

Phe Pro Ala Thr Arg Lys Ala Asn Asn Asp Ile Thr Ser Ile Thr Ser
 35 40 45

Asn Gly Gly Arg Val Asn Cys Met Gln Val Trp Pro Pro Ile Glu Lys
 50 55 60

Lys Lys Phe Glu Thr Leu Ser Tyr Leu Pro Asp Leu Thr Asp Ser Gly
 65 70 75 80

Gly Arg Val Asn Cys
 85

-continued

<210> SEQ ID NO 19
 <211> LENGTH: 76
 <212> TYPE: PRT
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 19

```

Met Ala Gln Val Ser Arg Ile Cys Asn Gly Val Gln Asn Pro Ser Leu
1           5           10           15

Ile Ser Asn Leu Ser Lys Ser Ser Gln Arg Lys Ser Pro Leu Ser Val
          20           25           30

Ser Leu Lys Thr Gln Gln His Pro Arg Ala Tyr Pro Ile Ser Ser Ser
          35           40           45

Trp Gly Leu Lys Lys Ser Gly Met Thr Leu Ile Gly Ser Glu Leu Arg
          50           55           60

Pro Leu Lys Val Met Ser Ser Val Ser Thr Ala Cys
65           70           75

```

<210> SEQ ID NO 20
 <211> LENGTH: 76
 <212> TYPE: PRT
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 20

```

Met Ala Gln Val Ser Arg Ile Cys Asn Gly Val Gln Asn Pro Ser Leu
1           5           10           15

Ile Ser Asn Leu Ser Lys Ser Ser Gln Arg Lys Ser Pro Leu Ser Val
          20           25           30

Ser Leu Lys Thr Gln Gln His Pro Arg Ala Tyr Pro Ile Ser Ser Ser
          35           40           45

Trp Gly Leu Lys Lys Ser Gly Met Thr Leu Ile Gly Ser Glu Leu Arg
          50           55           60

Pro Leu Lys Val Met Ser Ser Val Ser Thr Ala Cys
65           70           75

```

<210> SEQ ID NO 21
 <211> LENGTH: 72
 <212> TYPE: PRT
 <213> ORGANISM: Petunia hybrida

<400> SEQUENCE: 21

```

Met Ala Gln Ile Asn Asn Met Ala Gln Gly Ile Gln Thr Leu Asn Pro
1           5           10           15

Asn Ser Asn Phe His Lys Pro Gln Val Pro Lys Ser Ser Ser Phe Leu
          20           25           30

Val Phe Gly Ser Lys Lys Leu Lys Asn Ser Ala Asn Ser Met Leu Val
          35           40           45

Leu Lys Lys Asp Ser Ile Phe Met Gln Lys Phe Cys Ser Phe Arg Ile
          50           55           60

Ser Ala Ser Val Ala Thr Ala Cys
65           70

```

<210> SEQ ID NO 22
 <211> LENGTH: 69
 <212> TYPE: PRT
 <213> ORGANISM: Triticum aestivum

<400> SEQUENCE: 22

```

Met Ala Ala Leu Val Thr Ser Gln Leu Ala Thr Ser Gly Thr Val Leu

```

-continued

1	5	10	15	
Ser Val Thr	Asp Arg Phe Arg Arg	Pro Gly Phe Gln Gly	Leu Arg Pro	
	20	25	30	
Arg Asn Pro	Ala Asp Ala Ala Leu	Gly Met Arg Thr Val	Gly Ala Ser	
	35	40	45	
Ala Ala Pro	Lys Gln Ser Arg Lys	Pro His Arg Phe Asp	Arg Arg Cys	
	50	55	60	
Leu Ser Met	Val Val			
65				

<210> SEQ ID NO 23
 <211> LENGTH: 171
 <212> TYPE: DNA
 <213> ORGANISM: Pisum sativum

<400> SEQUENCE: 23

atggcttcta tgatattctc ttccgctgtg acaacagtca gccgtgcctc tagggggcaa	60
tccgcgcgcaa tggctccatt cggcggcctc aaatccatga ctggattccc agtgaggaag	120
gtcaacactg acattacttc cattacaagc aatggtggaa gagtaaagtg c	171

<210> SEQ ID NO 24
 <211> LENGTH: 255
 <212> TYPE: DNA
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 24

atggcttctc ctatgtcttc ttccgtact atggttgccct ctccggctca ggccactatg	60
gtcgtctcct tcaacggact taagtccctc gctgccttcc cagccaccgc caaggctaac	120
aacgacatta cttccatcac aagcaacggc ggaagagtta actgtatgca ggtgtggcct	180
ccgattgaaa agaagaagt tgagactctc tcttaccttc ctgaccttac cgattccggt	240
ggtcgcgtca actgc	255

<210> SEQ ID NO 25
 <211> LENGTH: 228
 <212> TYPE: DNA
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 25

atggcgcaag ttagcagaat ctgcaatggt gtgcagaacc catctcttat ctccaatctc	60
tcgaaatcca gtcaacgcaa atctccctta tcggtttctc tgaagacgca gcagcatcca	120
cgagcttata cgatttcgtc gtcgtgggga ttgaagaaga gtgggatgac gttaattggc	180
tctgagcttc gtctctttaa ggcatgtct tctgtttcca cggcgtgc	228

<210> SEQ ID NO 26
 <211> LENGTH: 228
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Artificial primer

<400> SEQUENCE: 26

atggcgcaag ttagcagaat ctgcaatggt gtgcagaacc catctcttat ctccaatctc	60
tcgaaatcca gtcaacgcaa atctccctta tcggtttctc tgaagacgca gcagcatcca	120
cgagcttata cgatttcgtc gtcgtgggga ttgaagaaga gtgggatgac gttaattggc	180

-continued

tctgagcttc gtctctcttaa ggcatgtct tctgtttcca cggcgtgc 228

<210> SEQ ID NO 27
 <211> LENGTH: 216
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Artificial
 primer

<400> SEQUENCE: 27

atggcccaga tcaacaacat ggcccagggc atccagaccc tgaaccctaa ctctaacttc 60
 cacaagccgc aagtgcccaa gtctagctcc ttctctgtgt tcggctccaa gaagctcaag 120
 aatagcgcca attccatgct ggtcctgaag aaagactcga tcttcattgca gaagttctgc 180
 tccttttcga tcagtgtctc ggttgcgact gcttgc 216

<210> SEQ ID NO 28
 <211> LENGTH: 207
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Artificial
 primer

<400> SEQUENCE: 28

atggcggcac tggtagacct ccagctcgcg acaagcggca ccgtcctgtc ggtgacggac 60
 cgcttcgggc gtcccggttc ccagggaact agggccacgga acccagccga tgcgctctc 120
 gggatgagga cgggtggcgc gtccgcggct cccaagcaga gcaggaagcc acaccgtttc 180
 gaccgcgggt gcttgagcat ggtcgtc 207

<210> SEQ ID NO 29
 <211> LENGTH: 433
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Artificial
 primer

<400> SEQUENCE: 29

agatcttgag ccaatcaaag aggagtgatg tagacctaaa gcaataatgg agccatgacg 60
 taagggtcta cggccatacg aaataattaa aggctgatgt gacctgtcgg tctctcagaa 120
 cctttacttt ttatgttttg cgtgtatatt taaatttcca cggcaatgac gatgtgacct 180
 aacgagatct tgagccaatc aaagaggagt gatgtgacct taaagcaata atggagccat 240
 gacgtaaggg cttacgcccc tacgaaataa ttaaaggctg atgtgacctg tcggtctctc 300
 agaaccttta ctttttatat ttggcgtgta tttttaaatt tccacggcaa tgacgatgtg 360
 acctgtgcat ccgctttgcc tataaataag ttttagtttg tattgatoga cagggtcgag 420
 aagacacggc cat 433

What is claimed is:

1. A method for selecting a transformed soybean plant cell 60
 comprising the steps of:

- a) contacting a population of plant cells comprising a trans-
 genic soybean plant cell transformed with a polynucle-
 otide encoding dicamba monooxygenase with medium 65
 comprising [auxin-like herbicide] *dicamba* in an amount
 that inhibits the growth of cells from the population
 lacking the polynucleotide, wherein the polynucleotide

comprises a nucleic acid sequence selected from: (1) a
 nucleic acid sequence encoding the polypeptide of SEQ
 ID NO: 6, (2) a nucleic acid sequence comprising the
 sequence of SEQ ID NO: 5, and (3) a nucleic acid
 sequence encoding a polypeptide with at least 90%
 sequence identity to the polypeptide of SEQ ID NO:6,
 wherein the polypeptide has dicamba monooxygenase
 activity; and

61

b) selecting the transformed soybean plant cell from the population of plant cells based on tolerance exhibited by the transformed cell to the [auxin-like] *dicamba* herbicide

further wherein:

said polynucleotide encoding dicamba monooxygenase is operatively linked to a chloroplast transit peptide coding sequence.

2. The method of claim 1, comprising culturing the population of plant cells on a medium lacking the [auxin-like herbicide] *dicamba* prior to step a) and/or between step a) and step b).

3. The method of claim 2, wherein the medium lacking the [auxin-like herbicide] *dicamba* contains a cytokinin.

4. The method of claim 2, wherein the medium lacking the [auxin-like herbicide] *dicamba* contains 6-benzyl amino purine (BAP).

5. The method of claim 4, wherein the 6-benzyl amino purine is in a concentration of about 10 mg/l of medium or less.

62

6. The method of claim 1, wherein the polynucleotide encoding dicamba monooxygenase is not genetically linked to a selectable or screenable marker gene other than dicamba monooxygenase.

5 7. The method of claim 1, further comprising the step of:
c) regenerating a fertile transgenic soybean plant from the transformed soybean plant cell.

[8. The method of claim 1, wherein the medium contains at least two auxin-like herbicides.]

10 [9. The method of claim 8, wherein the medium contains dicamba and 2,4-dichlorophenoxyacetic acid.]

10. The method of claim 1, wherein the population of cells comprises a cotyledon explant.

11. The method of claim 10, wherein the transformed plant cell is prepared by Agrobacterium-mediated transformation.

15 [12. The method of claim 1, wherein said auxin-like herbicide comprises 2,4-D, MCPA or 2,4-DB.]

13. The method of claim 1, wherein said [auxin-like herbicide] *dicamba* comprises dicamba in a concentration of from about 0.001 mg/L to about 0.02 mg/L.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION


PATENT NO. : RE44,971 E
APPLICATION NO. : 13/855631
DATED : June 24, 2014
INVENTOR(S) : Yuechun Wan et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title Page, Item (72), Please delete “**Yuenchun**”, and insert --**Yuechun**--

Signed and Sealed this
Thirtieth Day of December, 2014

A handwritten signature in black ink, reading "Michelle K. Lee". The signature is fluid and cursive, with the first letters of each name being capitalized and prominent.

Michelle K. Lee
Deputy Director of the United States Patent and Trademark Office