



(86) Date de dépôt PCT/PCT Filing Date: 2015/04/22
(87) Date publication PCT/PCT Publication Date: 2015/10/29
(45) Date de délivrance/Issue Date: 2023/03/07
(85) Entrée phase nationale/National Entry: 2016/10/19
(86) N° demande PCT/PCT Application No.: EP 2015/058633
(87) N° publication PCT/PCT Publication No.: 2015/162143
(30) Priorités/Priorities: 2014/04/23 (US61/982,893);
2014/04/23 (US61/982,895); 2014/04/23 (US61/982,894);
2014/04/23 (US61/982,897); 2014/04/23 (US61/982,896);
2014/04/23 (US61/982,903); 2014/04/23 (US61/982,898);
2014/04/23 (US61/982,899); 2014/04/23 (US61/982,901);
2014/04/23 (US61/982,904);

(51) Cl.Int./Int.Cl. *C12N 5/10* (2006.01),
A23D 9/00 (2006.01), *A23K 10/30* (2016.01),
C12N 15/54 (2006.01), *C12N 15/82* (2006.01),
C12N 5/04 (2006.01), *C12N 9/10* (2006.01)
(72) Inventeurs/Inventors:
TRESCH, STEFAN, DE;
SCHACHTSCHABEL, DOREEN, DE;
SISAY, MIHIRET TEKESTE, DE;
LERCHL, JENS, DE;
MAJOR, JULIA, DE;
...
(73) Propriétaire/Owner:
BASF SE, DE
(74) Agent: ROBIC

(54) Titre : PLANTES PRESENTANT UNE TOLERANCE ACCRUE AUX HERBICIDES
(54) Title: PLANTS HAVING INCREASED TOLERANCE TO HERBICIDES

(57) Abrégé/Abstract:

The present invention refers to a plant or plant part comprising a polynucleotide encoding a wildtype or mutated cellulose synthase (CESA) polypeptide, the expression of said polynucleotide confers to the plant or plant part tolerance to CESA-inhibiting herbicides, such as azines.

(30) Priorités(suite)/Priorities(continued): 2014/04/23 (US61/982,900)

(72) Inventeurs(suite)/Inventors(continued): VOGT, FLORIAN, DE; CALO, FREDERICK, DE; PAULIK, JILL MARIE, US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2015/162143 A1

(43) International Publication Date
29 October 2015 (29.10.2015)

(51) International Patent Classification:

C12N 15/82 (2006.01) *A01H 5/10* (2006.01)
A01H 5/00 (2006.01) *C12N 5/04* (2006.01)

(21) International Application Number:

PCT/EP2015/058633

(22) International Filing Date:

22 April 2015 (22.04.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/982,893	23 April 2014 (23.04.2014)	US
61/982,895	23 April 2014 (23.04.2014)	US
61/982,894	23 April 2014 (23.04.2014)	US
61/982,897	23 April 2014 (23.04.2014)	US
61/982,896	23 April 2014 (23.04.2014)	US
61/982,903	23 April 2014 (23.04.2014)	US
61/982,898	23 April 2014 (23.04.2014)	US
61/982,899	23 April 2014 (23.04.2014)	US
61/982,901	23 April 2014 (23.04.2014)	US
61/982,904	23 April 2014 (23.04.2014)	US
61/982,900	23 April 2014 (23.04.2014)	US

(71) Applicant: **BASF SE** [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors: **TRESCH, Stefan**; Rieslingweg 18, 67281 Kirchheim (DE). **SCHACHTSCHABEL, Doreen**; Nietzsche Straße 16, 68165 Mannheim (DE). **SISAY, Mihiret Tekeste**; Gontardstr. 6, 68163 Mannheim (DE). **LERCHL, Jens**; Golmer Fichten 5, 14476 Potsdam OT Golm (DE).

MAJOR, Julia; Hinter den Rüstern 15, 67251 Freinsheim (DE). **VOGT, Florian**; Lange Rötterstr. 64, 68167 Mannheim (DE). **CALO, Frederick**; Kopernikusstraße 36, 40223 Düsseldorf (DE). **PAULIK, Jill Marie**; 215 King George Loop, Cary, NC 27511 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: PLANTS HAVING INCREASED TOLERANCE TO HERBICIDES

(57) Abstract: The present invention refers to a plant or plant part comprising a polynucleotide encoding a wildtype or mutated cellulose synthase (CESA) polypeptide, the expression of said polynucleotide confers to the plant or plant part tolerance to CESA-inhibiting herbicides, such as azines.

WO 2015/162143 A1

PLANTS HAVING INCREASED TOLERANCE TO HERBICIDES

This application claims priority to US provisional applications numbers US 61/982893, US 61/982894, US 61/982895, US 61/982896, US 61/982897, US 61/982898, US

- 5 61/982899, US 61/982900, US 61/982901, US 61/982903, US 61/982904, which were all filed on April 23, 2014.

FIELD OF THE INVENTION

- 10 The present invention relates in general to methods for conferring on plants agricultural level tolerance to herbicides. Particularly, the invention refers to plants having an increased tolerance to herbicides, more specifically to herbicides which inhibit the enzyme cellulose synthase (CESA), also known cellulose biosynthesis inhibitors (CBI). More specifically, the present invention relates to methods and plants obtained by
- 15 mutagenesis and cross-breeding and transformation that have an increased tolerance to herbicides, particularly CESA-inhibiting herbicides.

BACKGROUND OF THE INVENTION

- 20 Plant cell walls are complex structures composed of high-molecular-weight polysaccharides, proteins, and lignins. Among the wall polysaccharides, cellulose, a hydrogen-bonded β -1,4-linked glucan microfibril, is the main load-bearing wall component and a key precursor for industrial applications. Cellulose is synthesized by large multimeric cellulose synthase (CESA) complexes (E.C.2.4.1.12), tracking along
- 25 cortical microtubules at the plasma membrane. The only known components of these complexes are the cellulose synthase proteins. Recent studies have identified tentative interaction partners for the CESAs and shown that the migratory patterns of the CESA complexes depend on phosphorylation status (for review see Endler and Persson, Molecular Plant, 2011, Volume 4 , Number 2, Pages 199–211, and references contained
- 30 therein). For example, cotton cellulose synthase genes, termed *CESA1* and *CESA2*, were identified in a collection of expressed sequence tag (EST) sequences on the basis of weak sequence similarity to genes for cellulose synthase from bacteria (Richmond and Somerville. Plant Physiology, 2000, Vol. 124, 495-498; and references contained
- 35 therein) In addition, the genes were expressed at high levels in cotton fibers at the onset of secondary wall synthesis and a purified fragment of one of the corresponding proteins as shown to bind UDP-Glc, the proposed substrate for cellulose biosynthesis. The conclusion that the cotton *CESA* genes are cellulose synthases is supported by results obtained with two cellulose-deficient Arabidopsis mutants, *rsw1* and *irx3* (Richmond and Somerville. Plant Physiology, Vol. 124, 2000, 495-498; and references
- 40 contained therein). The genes corresponding to the *RSW1* and *IRX3* loci exhibit a high degree of sequence similarity to the cotton *CESA* genes and are considered orthologs. Ten full-length *CESA* genes have been sequenced from Arabidopsis, and there is a genome survey sequence that may indicate one additional

family member. Reiterative database searches using the Arabidopsis Rsw1 (AtCESA1) and the cotton CESA polypeptide sequences as the initial query sequences revealed a large superfamily of at least 41 *CESA*-like genes in Arabidopsis. Based on predicted protein sequences, these genes were grouped into seven clearly distinguishable families

5 (Richmond and Somerville. Plant Physiology, Vol. 124, 2000, 495-498; and references contained therein): the *CESA* family, which includes *RSW1* and *IRX3* (*AtCESA7*), and six families of structurally related genes of unknown function designated as the "cellulose synthase-like" genes (*CsIA*, *CsIB*, *CsIC*, *CsID*, *CsIE*, and *CsIG*).

10 WO 2013/142968 describes plant cellulose synthase (CESA) alleles identified by mutagenizing plants and screening said plants with a cellulose biosynthetic inhibitor (CBI). CBIs employed in WO 2013/142968 include dichlobenil, chlorthiamid, isoxaben, flupoxam, and quinclorac, particularly isoxaben or flupoxam (named fpx1-1 to fpx1-3 [*CESA3*], fpx2-1 to fpx2-3 [*CESA1*] and ixr1-1 to ixr1-7 [*CESA3*], ixr2-1 to ixr2-2 [*CESA6*] mutants of
15 Arabidopsis CESA wildtype enzymes)

The inventors of the present invention have now surprisingly found that over-expression of the mutant cellulose synthase forms disclosed in WO 2013/142968 confers in plants tolerance/resistance to particular classes of CESA-inhibiting herbicides (cellulose
20 biosynthesis inhibitors; CBIs) as compared to the non-transformed and/or non-mutagenized plants or plant cells, respectively. More specifically, the inventors of the present invention have found that CESA expression confers tolerance/resistance to azines. More specifically, the inventors of the present invention have found that modifications of the C-terminal part of CESA proteins confer tolerance/resistance to azines.

25 The problem of the present invention can be seen as to the provision of novel traits by identifying target polypeptides, the manipulation of which makes plants tolerant to herbicides.

30 Three main strategies are available for making plants tolerant to herbicides, i.e. (1) detoxifying the herbicide with an enzyme which transforms the herbicide, or its active metabolite, into non-toxic products, such as, for example, the enzymes for tolerance to bromoxynil or to basta (EP242236, EP337899); (2) mutating the target enzyme into a functional enzyme which is less sensitive to the herbicide, or to its active metabolite, such
35 as, for example, the enzymes for tolerance to glyphosate (EP293356, Padgett S. R. et al., J.Biol. Chem., 266, 33, 1991); or (3) overexpressing the sensitive enzyme so as to produce quantities of the target enzyme in the plant which are sufficient in relation to the herbicide, in view of the kinetic constants of this enzyme, so as to have enough of the functional enzyme available despite the presence of its inhibitor.

40 The problem is solved by the subject-matter of the present invention.

SUMMARY OF THE INVENTION

Accordingly, in one aspect, the present invention provides a plant or plant part comprising a polynucleotide encoding a wildtype or mutated CESA polypeptide, the expression of said
5 polynucleotide confers to the plant or plant part tolerance to CESA-inhibiting herbicides.

In some aspects, the present invention provides a seed capable of germination into a plant comprising in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA
10 polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

In one aspect, the present invention provides a plant cell of or capable of regenerating a plant comprising in at least some of its cells a polynucleotide operably linked to a promoter
15 operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides, wherein the plant cell comprises the polynucleotide operably linked to a promoter.

20 In another aspect, the present invention provides a plant cell comprising a polynucleotide operably linked to a promoter operable in a cell, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

25 In other aspects, the present invention provides a plant product prepared from a plant or plant part comprising in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or
30 mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

In some aspects, the present invention provides a progeny or descendant plant derived from a plant comprising in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated
35 CESA polypeptide encoded by the polynucleotide, wherein the progeny or descendant plant comprises in at least some of its cells the recombinant polynucleotide operably linked to the promoter, the expression of the wildtype or mutated CESA polypeptide conferring to the progeny or descendant plant tolerance to the CESA-inhibiting herbicides.

40 In other aspects, the present invention provides a method for controlling weeds at a locus for growth of a plant, the method comprising: (a) applying an herbicide composition comprising CESA-inhibiting herbicides to the locus; and (b) planting a seed at the locus,

wherein the seed is capable of producing a plant that comprises in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

In some aspects, the present invention provides a method for controlling weeds at a locus for growth of a plant, the method comprising: applying an herbicidal composition comprising CESA-inhibiting herbicides to the locus; wherein said locus is: (a) a locus that contains: a plant or a seed capable of producing said plant; or (b) a locus that is to be after said applying is made to contain the plant or the seed; wherein the plant or the seed comprises in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

In one aspect, step (a) occurs before, after, or concurrently with step (b).

In other aspects, the present invention provides a method of producing a plant having tolerance to CESA-inhibiting herbicides, the method comprising regenerating a plant from a plant cell transformed with a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

In one aspect, the present invention provides a method of producing a progeny plant having tolerance to CESA-inhibiting herbicides, the method comprising: crossing a first CESA-inhibiting herbicides- tolerant plant with a second plant to produce a CESA-inhibiting herbicides- tolerant progeny plant, wherein the first plant and the progeny plant comprise in at least some of their cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

In addition, the present invention refers to a method for identifying a CESA-inhibiting herbicide by using a wildtype or mutated CESA of the present invention encoded by a nucleic acid which comprises the nucleotide sequence of SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or a variant, homologue, paralogue or orthologue thereof.

Said method comprises the steps of:

- a) generating a transgenic cell or plant comprising a nucleic acid encoding a wildtype or mutated CESA of the present invention, wherein the wildtype or mutated CESA of the

present invention is expressed;

- b) applying a CESA-inhibiting herbicide to the transgenic cell or plant of a) and to a control cell or plant of the same variety;
- c) determining the growth or the viability of the transgenic cell or plant and the control cell or plant after application of said test compound, and
- d) selecting test compounds which confer reduced growth to the control cell or plant as compared to the growth of the transgenic cell or plant.

Another object refers to a method of identifying a nucleotide sequence encoding a wildtype or mutated CESA which is resistant or tolerant to a CESA-inhibiting herbicide, the method comprising:

- a) generating a library of wildtype or mutated CESA-encoding nucleic acids,
- b) screening a population of the resulting wildtype or mutated CESA-encoding nucleic acids by expressing each of said nucleic acids in a cell or plant and treating said cell or plant with a CESA-inhibiting herbicide,
- c) comparing the CESA-inhibiting herbicide-tolerance levels provided by said population of wildtype or mutated CESA encoding nucleic acids with the CESA-inhibiting herbicide-tolerance level provided by a control CESA-encoding nucleic acid,
- d) selecting at least one wildtype or mutated CESA-encoding nucleic acid that provides a significantly increased level of tolerance to a CESA-inhibiting herbicide as compared to that provided by the control CESA-encoding nucleic acid.

In a preferred embodiment, the wildtype or mutated CESA-encoding nucleic acid selected in step d) provides at least 2-fold as much tolerance to a CESA-inhibiting herbicide as compared to that provided by the control CESA-encoding nucleic acid.

The resistance or tolerance can be determined by generating a transgenic plant comprising a nucleic acid sequence of the library of step a) and comparing said transgenic plant with a control plant.

Another object refers to a method of identifying a plant or algae containing a nucleic acid encoding a wildtype or mutated CESA which is resistant or tolerant to a CESA-inhibiting herbicide, the method comprising:

- a) identifying an effective amount of a CESA-inhibiting herbicide in a culture of plant cells or green algae.
- b) treating said plant cells or green algae with a mutagenizing agent,
- c) contacting said mutagenized cells population with an effective amount of CESA-inhibiting herbicide, identified in a),
- d) selecting at least one cell surviving these test conditions,
- e) PCR-amplification and sequencing of CESA genes from cells selected in d) and comparing such sequences to wild-type CESA gene sequences, respectively.

In a preferred embodiment, the mutagenizing agent is ethylmethanesulfonate.

Another object refers to an isolated recombinantly produced, and/or chemically synthesized nucleic acid encoding a wildtype or mutated CESA, the nucleic acid comprising the
5 sequence of SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or a variant thereof, as defined hereinafter.

10 In a preferred embodiment, the nucleic acid being identifiable by a method as defined above.

Another object refers to an isolated, recombinantly produced, and/or chemically synthesized wildtype or mutated CESA polypeptide, the polypeptide comprising the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23,
24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47,
15 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, a variant, derivative, orthologue, paralogue or homologue thereof, as defined hereinafter.

In still further aspects, the present invention provides a plant or plant part comprising in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells,
20 the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides, wherein the plant or plant part further exhibits a second or third herbicide-tolerant trait.

25 In another embodiment, the invention refers to a plant cell transformed by and expressing a a wildtype or mutated CESA nucleic acid according to the present invention or a plant which has been mutated to obtain a plant expressing, preferably over-expressing a a wildtype or mutated CESA nucleic acid according to the present invention, wherein expression of said nucleic acid in the plant cell results in increased resistance or tolerance to a CESA -
30 inhibiting herbicide as compared to a wild type variety of the plant cell

In another embodiment, the invention refers to a plant comprising a plant cell according to the present invention, wherein expression of the nucleic acid in the plant results in the plant's increased resistance to CESA-inhibiting herbicide as compared to a wild type variety
35 of the plant.

The plants of the present invention can be transgenic or non-transgenic.

Preferably, the expression of the nucleic acid of the invention in the plant results in the
40 plant's increased resistance to CESA-inhibiting herbicides as compared to a wild type variety of the plant. In another embodiment, the invention refers to a seed produced by a transgenic plant comprising a plant cell of the present invention, wherein the seed is true

breeding for an increased resistance to a CESA-inhibiting herbicide as compared to a wild type variety of the seed.

In another embodiment, the invention refers to a method of producing a transgenic plant cell
5 with an increased resistance to a CESA-inhibiting herbicide as compared to a wild type
variety of the plant cell comprising, transforming the plant cell with an expression cassette
comprising a a polynucleotide operably linked to a promoter operable in plant cells, the
promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the
10 polynucleotide.

In another embodiment, the invention refers to a method of producing a transgenic plant
comprising, (a) transforming a plant cell with an expression cassette comprising a a
polynucleotide operably linked to a promoter operable in plant cells, the promoter capable
of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, and
15 (b) generating a plant with an increased resistance to CESA-inhibiting herbicide from the
plant cell.

Preferably, the expression cassette further comprises a transcription initiation regulatory
region and a translation initiation regulatory region that are functional in the plant.

20

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 A, B, C

25 Figure 1 shows phylogenetic trees of cellulose synthase homologues in corn (A), soy (B)
and rice (C).

Figure 2 A, B, C

30 Figure 2 shows alignment of all cellulose synthase homologues (A), arabidopsis CESA1,
CESA3 and CESA 6 homologues (B) and Arabidopsis CESA1 and CESA3 homologues (C).

KEY TO SEQUENCE LISTING

Table 1

SEQ ID NO	
Amino acid	Sequence / Origin
1	At_CESA1
2	At_CESA2
3	At_CESA3
4	At_CESA4
5	At_CESA5

6	At_CESA6
7	At_CESA7
8	At_CESA8
9	At_CESA9
10	At_CESA10
11	GRMZM2G027723 GRMZM2G027723_T01
12	GRMZM2G055795 GRMZM2G055795_T01
13	GRMZM2G039454 GRMZM2G039454_T01
14	GRMZM2G025231 GRMZM2G025231_T02
15	GRMZM2G142898 GRMZM2G142898_T01
16	GRMZM2G424832 GRMZM2G424832_T01
17	GRMZM2G177631 GRMZM2G177631_T01
18	GRMZM2G150404 GRMZM2G150404_T01
19	GRMZM2G028353 GRMZM2G028353_T01
20	GRMZM2G018241 GRMZM2G018241_T01
21	GRMZM2G037413 GRMZM2G037413_T01
22	GRMZM2G082580 GRMZM2G082580_T01
23	GRMZM2G112336 GRMZM2G112336_T01
24	GRMZM2G111642 GRMZM2G111642_T01
25	GRMZM2G113137 GRMZM2G113137_T01
26	Glyma02g36720 Glyma02g36720.1
27	Glyma02g08920 Glyma02g08920.1
28	Glyma12g36570 Glyma12g36570.1
29	Glyma12g17730 Glyma12g17730.1
30	Glyma06g06870 Glyma06g06870.1
31	Glyma06g30860 Glyma06g30860.1
32	Glyma06g47420 Glyma06g47420.1
33	Glyma06g30850 Glyma06g30850.1
34	Glyma06g07320 Glyma06g07320.1
35	Glyma04g07220 Glyma04g07220.1
36	Glyma04g06780 Glyma04g06780.1
37	Glyma04g23530 Glyma04g23530.2
38	Glyma08g15380 Glyma08g15380.1
39	Glyma08g12400 Glyma08g12400.1
40	Glyma17g08000 Glyma17g08000.1
41	Glyma13g27250 Glyma13g27250.3
42	Glyma05g29240 Glyma05g29240.2
43	Glyma05g32100 Glyma05g32100.1
44	Glyma15g43040 Glyma15g43040.1
45	Glyma09g15620 Glyma09g15620.2

46	Glyma10g36790 Glyma10g36790.1
47	Glyma16g28080 Glyma16g28080.2
48	Os01g54620.1
49	Os03g59340.1
50	Os03g62090.1
51	Os05g08370.1
52	Os06g02180.1
53	Os06g22980.1
54	Os06g39970.1
55	Os07g10770.1
56	Os07g14850.1
57	Os07g24190.1
58	Os07g24190.2
59	Os07g24190.3
60	Os08g25710.1
61	Os09g25490.1
62	Os10g32980.1
63	Os10g42750.1
64	Os12g36890.1

DETAILED DESCRIPTION

5 The articles "a" and "an" are used herein to refer to one or more than one (i.e., to at least one) of the grammatical object of the article. By way of example, "an element" means one or more elements.

10 As used herein, the word "comprising," or variations such as "comprises" or "comprising," will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

The term "control of undesired vegetation or weeds" is to be understood as meaning the killing of weeds and/or otherwise retarding or inhibiting the normal growth of the weeds.

15 Weeds, in the broadest sense, are understood as meaning all those plants which grow in locations where they are undesired. The weeds of the present invention include, for example, dicotyledonous and monocotyledonous weeds. Dicotyledonous weeds include, but are not limited to, weeds of the genera: Sinapis, Lepidium, Galium, Stellaria, Matricaria, Anthemis, Galinsoga, Chenopodium, Urtica, Senecio, Amaranthus, Portulaca, Xanthium,

20 Convolvulus, Ipomoea, Polygonum, Sesbania, Ambrosia, Cirsium, Carduus, Sonchus, Solanum, Rorippa, Rotala, Lindernia, Lamium, Veronica, Abutilon, Emex, Datura, Viola, Galeopsis, Papaver, Centaurea, Trifolium, Ranunculus, and Taraxacum. Monocotyledonous

- weeds include, but are not limited to, weeds of the genera: *Echinochloa*, *Setaria*, *Panicum*, *Digitaria*, *Phleum*, *Poa*, *Festuca*, *Eleusine*, *Brachiaria*, *Lolium*, *Bromus*, *Avena*, *Cyperus*, *Sorghum*, *Agropyron*, *Cynodon*, *Monochoria*, *Fimbristylis*, *Sagittaria*, *Eleocharis*, *Scirpus*, *Paspalum*, *Ischaemum*, *Sphenoclea*, *Dactyloctenium*, *Agrostis*, *Alopecurus*, and
- 5 *Apera*. In addition, the weeds of the present invention can include, for example, crop plants that are growing in an undesired location. For example, a volunteer maize plant that is in a field that predominantly comprises soybean plants can be considered a weed, if the maize plant is undesired in the field of soybean plants.
- 10 The term “plant” is used in its broadest sense as it pertains to organic material and is intended to encompass eukaryotic organisms that are members of the Kingdom Plantae, examples of which include but are not limited to vascular plants, vegetables, grains, flowers, trees, herbs, bushes, grasses, vines, ferns, mosses, fungi and algae, etc, as well as clones, offsets, and parts of plants used for asexual propagation (e.g. cuttings, pipings, shoots,
- 15 rhizomes, underground stems, clumps, crowns, bulbs, corms, tubers, rhizomes, plants/tissues produced in tissue culture, etc.). The term “plant” further encompasses whole plants, ancestors and progeny of the plants and plant parts, including seeds, shoots, stems, leaves, roots (including tubers), flowers, florets, fruits, pedicles, peduncles, stamen, anther, stigma, style, ovary, petal, sepal, carpel, root tip, root cap, root hair, leaf hair, seed hair,
- 20 pollen grain, microspore, cotyledon, hypocotyl, epicotyl, xylem, phloem, parenchyma, endosperm, a companion cell, a guard cell, and any other known organs, tissues, and cells of a plant, and tissues and organs, wherein each of the aforementioned comprise the gene/nucleic acid of interest. The term “plant” also encompasses plant cells, suspension cultures, callus tissue, embryos, meristematic regions, gametophytes, sporophytes, pollen
- 25 and microspores, again wherein each of the aforementioned comprises the gene/nucleic acid of interest.

- Plants that are particularly useful in the methods of the invention include all plants which belong to the superfamily Viridiplantae, in particular monocotyledonous and dicotyledonous
- 30 plants including fodder or forage legumes, ornamental plants, food crops, trees or shrubs selected from the list comprising *Acer* spp., *Actinidia* spp., *Abelmoschus* spp., *Agave sisalana*, *Agropyron* spp., *Agrostis stolonifera*, *Allium* spp., *Amaranthus* spp., *Ammophila arenaria*, *Ananas comosus*, *Annona* spp., *Apium graveolens*, *Arachis* spp., *Artocarpus* spp., *Asparagus officinalis*, *Avena* spp. (e.g. *Avena sativa*, *Avena fatua*, *Avena byzantina*, *Avena fatua* var. *sativa*, *Avena hybrida*), *Averrhoa carambola*, *Bambusa* sp., *Benincasa hispida*, *Bertholletia excelsa*, *Beta vulgaris*, *Brassica* spp. (e.g. *Brassica napus*, *Brassica rapa* ssp. [canola, oilseed rape, turnip rape]), *Cadaba farinosa*, *Camellia sinensis*, *Canna indica*, *Cannabis sativa*, *Capsicum* spp., *Carex elata*, *Carica papaya*, *Carissa macrocarpa*, *Carya* spp., *Carthamus tinctorius*, *Castanea* spp., *Ceiba pentandra*, *Cichorium endivia*,
- 35 *Cinnamomum* spp., *Citrullus lanatus*, *Citrus* spp., *Cocos* spp., *Coffea* spp., *Colocasia esculenta*, *Cola* spp., *Corchorus* sp., *Coriandrum sativum*, *Corylus* spp., *Crataegus* spp., *Crocus sativus*, *Cucurbita* spp., *Cucumis* spp., *Cynara* spp., *Daucus carota*, *Desmodium*
- 40

spp., *Dimocarpus longan*, *Dioscorea* spp., *Diospyros* spp., *Echinochloa* spp., *Elaeis* (e.g. *Elaeis guineensis*, *Elaeis oleifera*), *Eleusine coracana*, *Eragrostis tef*, *Erianthus* sp., *Eriobotrya japonica*, *Eucalyptus* sp., *Eugenia uniflora*, *Fagopyrum* spp., *Fagus* spp., *Festuca arundinacea*, *Ficus carica*, *Fortunella* spp., *Fragaria* spp., *Ginkgo biloba*, *Glycine* spp. (e.g. *Glycine max*, *Soja hispida* or *Soja max*), *Gossypium hirsutum*, *Helianthus* spp. (e.g. *Helianthus annuus*), *Hemerocallis fulva*, *Hibiscus* spp., *Hordeum* spp. (e.g. *Hordeum vulgare*), *Ipomoea batatas*, *Juglans* spp., *Lactuca sativa*, *Lathyrus* spp., *Lens culinaris*, *Linum usitatissimum*, *Litchi chinensis*, *Lotus* spp., *Luffa acutangula*, *Lupinus* spp., *Luzula sylvatica*, *Lycopersicon* spp. (e.g. *Lycopersicon esculentum*, *Lycopersicon lycopersicum*, *Lycopersicon pyriforme*), *Macrotyloma* spp., *Malus* spp., *Malpighia emarginata*, *Mammea americana*, *Mangifera indica*, *Manihot* spp., *Manilkara zapota*, *Medicago sativa*, *Melilotus* spp., *Mentha* spp., *Miscanthus sinensis*, *Momordica* spp., *Morus nigra*, *Musa* spp., *Nicotiana* spp., *Olea* spp., *Opuntia* spp., *Ornithopus* spp., *Oryza* spp. (e.g. *Oryza sativa*, *Oryza latifolia*), *Panicum miliaceum*, *Panicum virgatum*, *Passiflora edulis*, *Pastinaca sativa*, *Pennisetum* sp., *Persea* spp., *Petroselinum crispum*, *Phalaris arundinacea*, *Phaseolus* spp., *Phleum pratense*, *Phoenix* spp., *Phragmites australis*, *Physalis* spp., *Pinus* spp., *Pistacia vera*, *Pisum* spp., *Poa* spp., *Populus* spp., *Prosopis* spp., *Prunus* spp., *Psidium* spp., *Punica granatum*, *Pyrus communis*, *Quercus* spp., *Raphanus sativus*, *Rheum rhabarbarum*, *Ribes* spp., *Ricinus communis*, *Rubus* spp., *Saccharum* spp., *Salix* sp., *Sambucus* spp., *Secale cereale*, *Sesamum* spp., *Sinapis* sp., *Solanum* spp. (e.g. *Solanum tuberosum*, *Solanum integrifolium* or *Solanum lycopersicum*), *Sorghum bicolor*, *Spinacia* spp., *Syzygium* spp., *Tagetes* spp., *Tamarindus indica*, *Theobroma cacao*, *Trifolium* spp., *Tripsacum dactyloides*, *Triticosecale rimpau*, *Triticum* spp. (e.g. *Triticum aestivum*, *Triticum durum*, *Triticum turgidum*, *Triticum hybernum*, *Triticum macha*, *Triticum sativum*, *Triticum monococcum* or *Triticum vulgare*), *Tropaeolum minus*, *Tropaeolum majus*, *Vaccinium* spp., *Vicia* spp., *Vigna* spp., *Viola odorata*, *Vitis* spp., *Zea mays*, *Zizania palustris*, *Ziziphus* spp., amaranth, artichoke, asparagus, broccoli, Brussels sprouts, cabbage, canola, carrot, cauliflower, celery, collard greens, flax, kale, lentil, oilseed rape, okra, onion, potato, rice, soybean, strawberry, sugar beet, sugar cane, sunflower, tomato, squash, tea and algae, amongst others. According to a preferred embodiment of the present invention, the plant is a crop plant. Examples of crop plants include inter alia soybean, sunflower, canola, alfalfa, rapeseed, cotton, tomato, potato or tobacco. Further preferably, the plant is a monocotyledonous plant, such as sugarcane. Further preferably, the plant is a cereal, such as rice, maize, wheat, barley, millet, rye, sorghum or oats.

35

Generally, the term "herbicide" is used herein to mean an active ingredient that kills, controls or otherwise adversely modifies the growth of plants. The preferred amount or concentration of the herbicide is an "effective amount" or "effective concentration." By "effective amount" and "effective concentration" is intended an amount and concentration, respectively, that is sufficient to kill or inhibit the growth of a similar, wild-type, plant, plant tissue, plant cell, or host cell, but that said amount does not kill or inhibit as severely the growth of the herbicide-resistant plants, plant tissues, plant cells, and host cells of the

40

present invention. Typically, the effective amount of a herbicide is an amount that is routinely used in agricultural production systems to kill weeds of interest. Such an amount is known to those of ordinary skill in the art. Herbicidal activity is exhibited by herbicides useful for the the present invention when they are applied directly to the plant or to the locus of the
5 plant at any stage of growth or before planting or emergence. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of
10 chemical applied. These and other factors can be adjusted as is known in the art to promote non-selective or selective herbicidal action. Generally, it is preferred to apply the herbicide postemergence to relatively immature undesirable vegetation to achieve the maximum control of weeds.

15 By a "herbicide-tolerant" or "herbicide-resistant" plant, it is intended that a plant that is tolerant or resistant to at least one herbicide at a level that would normally kill, or inhibit the growth of, a normal or wild-type plant. Levels of herbicide that normally inhibit growth of a non-tolerant plant are known and readily determined by those skilled in the art. Examples include the amounts recommended by manufacturers for application. The maximum rate is
20 an example of an amount of herbicide that would normally inhibit growth of a non-tolerant plant. For the present invention, the terms "herbicide-tolerant" and "herbicide-resistant" are used interchangeably and are intended to have an equivalent meaning and an equivalent scope. Similarly, the terms "herbicide-tolerance" and "herbicide-resistance" are used interchangeably and are intended to have an equivalent meaning and an equivalent scope.
25 Similarly, the terms "tolerant" and "resistant" are used interchangeably and are intended to have an equivalent meaning and an equivalent scope. As used herein, in regard to an herbicidal composition useful in various embodiments hereof, terms such as CESA-inhibiting herbicides, and the like, refer to those agronomically acceptable herbicide active ingredients (A.I.) recognized in the art. Similarly, terms such as fungicide, nematicide,
30 pesticide, and the like, refer to other agronomically acceptable active ingredients recognized in the art.

When used in reference to a particular mutant enzyme or polypeptide, terms such as herbicide-tolerant and herbicide-tolerance refer to the ability of such enzyme or polypeptide
35 to perform its physiological activity in the presence of an amount of an herbicide A.I. that would normally inactivate or inhibit the activity of the wild-type (non-mutant) version of said enzyme or polypeptide. For example, when used specifically in regard to a CESA enzyme, it refers specifically to the ability to tolerate a CESA-inhibitor. By "herbicide-tolerant wildtype or mutated CESA protein" or "herbicide -resistant wildtype or mutated CESA protein", it is
40 intended that such a CESA protein displays higher CESA activity, relative to the CESA activity of a wild-type CESA protein, when in the presence of at least one herbicide that is known to interfere with CESA activity and at a concentration or level of the herbicide that is

known to inhibit the CESA activity of the wild-type CESA protein. Furthermore, the CESA activity of such a herbicide-tolerant or herbicide-resistant wildtype or mutated CESA protein may be referred to herein as "herbicide-tolerant" or "herbicide-resistant" CESA activity.

- 5 As used herein, "recombinant," when referring to nucleic acid or polypeptide, indicates that such material has been altered as a result of human application of a recombinant technique, such as by polynucleotide restriction and ligation, by polynucleotide overlap-extension, or by genomic insertion or transformation. A gene sequence open reading frame is recombinant if that nucleotide sequence has been removed from its natural text and
10 cloned into any type of artificial nucleic acid vector. The term recombinant also can refer to an organism having a recombinant material, e.g., a plant that comprises a recombinant nucleic acid can be considered a recombinant plant.

- The term "transgenic plant" refers to a plant that comprises a heterologous polynucleotide.
15 Preferably, the heterologous polynucleotide is stably integrated within the genome such that the polynucleotide is passed on to successive generations. The heterologous polynucleotide may be integrated into the genome alone or as part of a recombinant expression cassette. "Transgenic" is used herein to refer to any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been so altered by the presence of
20 heterologous nucleic acid including those transgenic organisms or cells initially so altered, as well as those created by crosses or asexual propagation from the initial transgenic organism or cell. In some embodiments, a "recombinant" organism is a "transgenic" organism. The term "transgenic" as used herein is not intended to encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding
25 methods (e.g., crosses) or by naturally occurring events such as, e.g., self-fertilization, random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

- As used herein, "mutagenized" refers to an organism or DNA thereof having alteration(s) in
30 the biomolecular sequence of its native genetic material as compared to the sequence of the genetic material of a corresponding wild-type organism or DNA, wherein the alteration(s) in genetic material were induced and/or selected by human action. Examples of human action that can be used to produce a mutagenized organism or DNA include, but are not limited to, as illustrated in regard to herbicide tolerance: tissue culture of plant cells
35 (e.g., calli) and selection thereof with herbicides (e.g., CESA-inhibiting herbicides), treatment of plant cells with a chemical mutagen such as EMS and subsequent selection with herbicide(s); or by treatment of plant cells with x-rays and subsequent selection with herbicide(s). Any method known in the art can be used to induce mutations. Methods of inducing mutations can induce mutations in random positions in the genetic material or can
40 induce mutations in specific locations in the genetic material (i.e., can be directed mutagenesis techniques), such as by use of a genoplasty technique.

As used herein, a "genetically modified organism" (GMO) is an organism whose genetic characteristics contain alteration(s) that were produced by human effort causing transfection that results in transformation of a target organism with genetic material from another or "source" organism, or with synthetic or modified-native genetic material, or an organism that is a descendant thereof that retains the inserted genetic material. The source organism can be of a different type of organism (e.g., a GMO plant can contain bacterial genetic material) or from the same type of organism (e.g., a GMO plant can contain genetic material from another plant). As used herein in regard to plants and other organisms, "recombinant," "transgenic," and "GMO" are considered synonyms and indicate the presence of genetic material from a different source; in contrast, "mutagenized" is used to refer to a plant or other organism, or the DNA thereof, in which no such transgenic material is present, but in which the native genetic material has become mutated so as to differ from a corresponding wild-type organism or DNA.

As used herein, "wild-type" or "corresponding wild-type plant" means the typical form of an organism or its genetic material, as it normally occurs, as distinguished from, e.g., mutagenized and/or recombinant forms. Similarly, by "control cell" or "similar, wild-type, plant, plant tissue, plant cell or host cell" is intended a plant, plant tissue, plant cell, or host cell, respectively, that lacks the herbicide-resistance characteristics and/or particular polynucleotide of the invention that are disclosed herein. The use of the term "wild-type" is not, therefore, intended to imply that a plant, plant tissue, plant cell, or other host cell lacks recombinant DNA in its genome, and/or does not possess herbicide-resistant characteristics that are different from those disclosed herein.

As used herein, "descendant" refers to any generation plant. In some embodiments, a descendant is a first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, or tenth generation plant.

As used herein, "progeny" refers to a first generation plant.

The term "seed" comprises seeds of all types, such as, for example, true seeds, caryopses, achenes, fruits, tubers, seedlings and similar forms. In the context of Brassica and Sinapis species, "seed" refers to true seed(s) unless otherwise specified. For example, the seed can be seed of transgenic plants or plants obtained by traditional breeding methods.

Examples of traditional breeding methods can include cross-breeding, selfing, back-crossing, embryo rescue, in-crossing, out-crossing, inbreeding, selection, asexual propagation, and other traditional techniques as are known in the art.

Although exemplified with reference to specific plants or plant varieties and their hybrids, in various embodiments, the presently described methods using CESA-inhibiting herbicides can be employed with a variety of commercially valuable plants. CESA-inhibiting herbicides-tolerant plant lines described as useful herein can be employed in weed control methods

either directly or indirectly, i. e. either as crops for herbicide treatment or as CESA-inhibiting herbicides-tolerance trait donor lines for development, as by traditional plant breeding, to produce other varietal and/or hybrid crops containing such trait or traits. All such resulting variety or hybrids crops, containing the ancestral CESA-inhibiting herbicides-tolerance trait or traits can be referred to herein as progeny or descendant of the ancestral, CESA-inhibiting herbicides-tolerant line(s). Such resulting plants can be said to retain the "herbicide tolerance characteristic(s)" of the ancestral plant, i.e. meaning that they possess and express the ancestral genetic molecular components responsible for the trait.

10 In one aspect, the present invention provides a plant or plant part comprising a polynucleotide encoding a wildtype or mutated CESA polypeptide, the expression of said polynucleotide confers to the plant or plant part tolerance to CESA-inhibiting herbicides.

15 In a preferred embodiment, the plant has been previously produced by a process comprising recombinantly preparing a plant by introducing and over-expressing a wildtype or mutated CESA transgene according to the present invention, as described in greater detail hereinafter.

20 In another preferred embodiment, the plant has been previously produced by a process comprising in situ mutagenizing plant cells or seeds, to obtain plant cells or plants which express a wildtype or mutated CESA.

25 In another embodiment, the polynucleotide encoding the wildtype or mutated CESA polypeptide comprises the nucleic acid sequence set forth in SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or a variant or derivative thereof.

30 In other embodiments, the wildtype or mutated CESA polypeptide for use according to the present invention is a functional variant having, over the full-length of the variant, at least about 80%, illustratively, at least about 80%, 90%, 95%, 98%, 99% or more amino acid sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64.

35 In another embodiment, the wildtype or mutated CESA polypeptide for use according to the present invention is a functional fragment of a polypeptide having the amino acid sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64.

40 It is recognized that the CESA polynucleotide molecules and CESA polypeptides of the invention encompass polynucleotide molecules and polypeptides comprising a nucleotide or an amino acid sequence that is sufficiently identical to nucleotide sequences set forth in

SEQ ID Nos: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or to the amino acid sequences set forth in SEQ ID Nos: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64.

5 The term "sufficiently identical" is used herein to refer to a first amino acid or nucleotide sequence that contains a sufficient or minimum number of identical or equivalent (e.g., with a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences have a common structural domain and/or common functional activity.

10

Generally, "sequence identity" refers to the extent to which two optimally aligned DNA or amino acid sequences are invariant throughout a window of alignment of components, e.g., nucleotides or amino acids. An "identity fraction" for aligned segments of a test sequence and a reference sequence is the number of identical components that are shared by the two aligned sequences divided by the total number of components in reference sequence segment, i.e., the entire reference sequence or a smaller defined part of the reference sequence. "Percent identity" is the identity fraction times 100. Optimal alignment of sequences for aligning a comparison window are well known to those skilled in the art and may be conducted by tools such as the local homology algorithm of Smith and Waterman, 15 the homology alignment algorithm of Needleman and Wunsch, the search for similarity method of Pearson and Lipman, and preferably by computerized implementations of these algorithms such as GAP, BESTFIT, FASTA, and TFASTA available as part of the GCG. Wisconsin Package. (Accelrys Inc. Burlington, Mass.)

25 Polynucleotides and Oligonucleotides

By an "isolated polynucleotide", including DNA, RNA, or a combination of these, single or double stranded, in the sense or antisense orientation or a combination of both, dsRNA or otherwise, we mean a polynucleotide which is at least partially separated from the polynucleotide sequences with which it is associated or linked in its native state. Preferably, 30 the isolated polynucleotide is at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated. As the skilled addressee would be aware, an isolated polynucleotide can be an exogenous polynucleotide present in, for example, a transgenic organism which does not naturally comprise the polynucleotide. Furthermore, the terms "polynucleotide(s)", "nucleic acid sequence(s)", "nucleotide sequence(s)", "nucleic acid(s)", "nucleic acid molecule" are used interchangeably herein and refer to nucleotides, either ribonucleotides or deoxyribonucleotides or a combination of both, in a polymeric unbranched form of any length.

40

The term "mutated CESA nucleic acid" refers to a CESA nucleic acid having a sequence that is mutated from a wild-type CESA nucleic acid and that confers increased CESA-

inhibiting herbicide tolerance to a plant in which it is expressed. Furthermore, the term "mutated cellulose synthase (mutated CESA)" refers to the replacement of an amino acid of the wild-type primary sequences of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, or a variant, a derivative, a homologue, an orthologue, or paralogue thereof, with another amino acid. The expression "mutated amino acid" will be used below to designate the amino acid which is replaced by another amino acid, thereby designating the site of the mutation in the primary sequence of the protein.

10

In a preferred embodiment, the CESA nucleotide sequence encoding a mutated CESA comprises the sequence of SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or a variant or derivative thereof

15 Furthermore, it will be understood by the person skilled in the art that the CESA nucleotide sequences encompass homologues, paralogues and orthologues of SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, as defined hereinafter.

The term "variant" with respect to a sequence (e.g., a polypeptide or nucleic acid sequence such as – for example – a transcription regulating nucleotide sequence of the invention) is intended to mean substantially similar sequences. For nucleotide sequences comprising an open reading frame, variants include those sequences that, because of the degeneracy of the genetic code, encode the identical amino acid sequence of the native protein. Naturally occurring allelic variants such as these can be identified with the use of well-known molecular biology techniques, as, for example, with polymerase chain reaction (PCR) and hybridization techniques. Variant nucleotide sequences also include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis and for open reading frames, encode the native protein comprising the sequence of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, as well as those that encode a polypeptide having amino acid substitutions relative to the native protein, e.g. the mutated CESA according to the present invention as disclosed herein. Generally, nucleotide sequence variants of the invention will have at least 30, 40, 50, 60, to 70%, e.g., preferably 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, to 79%, generally at least 80%, e.g., 81%-84%, at least 85%, e.g., 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, to 98% and 99% nucleotide "sequence identity" to the nucleotide sequence of SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77. The % identity of a polynucleotide is determined by GAP (Needleman and Wunsch, 1970) analysis (GCG program) with a gap creation penalty=5, and a gap extension penalty=0.3. Unless stated otherwise, the query sequence is at least 45 nucleotides in length, and the GAP analysis aligns the two sequences over a region of at least 45 nucleotides. Preferably, the query

40

sequence is at least 150 nucleotides in length, and the GAP analysis aligns the two sequences over a region of at least 150 nucleotides. More preferably, the query sequence is at least 300 nucleotides in length and the GAP analysis aligns the two sequences over a region of at least 300 nucleotides. Even more preferably, the GAP analysis aligns the two sequences over their entire length.

Polypeptides

By "substantially purified polypeptide" or "purified" a polypeptide is meant that has been separated from one or more lipids, nucleic acids, other polypeptides, or other contaminating molecules with which it is associated in its native state. It is preferred that the substantially purified polypeptide is at least 60% free, more preferably at least 75% free, and more preferably at least 90% free from other components with which it is naturally associated. As the skilled addressee will appreciate, the purified polypeptide can be a recombinantly produced polypeptide. The terms "polypeptide" and "protein" are generally used interchangeably and refer to a single polypeptide chain which may or may not be modified by addition of non-amino acid groups. It would be understood that such polypeptide chains may associate with other polypeptides or proteins or other molecules such as co-factors. The terms "proteins" and "polypeptides" as used herein also include variants, mutants, modifications, analogous and/or derivatives of the polypeptides of the invention as described herein.

The % identity of a polypeptide is determined by GAP (Needleman and Wunsch, 1970) analysis (GCG program) with a gap creation penalty=5, and a gap extension penalty=0.3.

The query sequence is at least 25 amino acids in length, and the GAP analysis aligns the two sequences over a region of at least 25 amino acids. More preferably, the query sequence is at least 50 amino acids in length, and the GAP analysis aligns the two sequences over a region of at least 50 amino acids. More preferably, the query sequence is at least 100 amino acids in length and the GAP analysis aligns the two sequences over a region of at least 100 amino acids. Even more preferably, the query sequence is at least 250 amino acids in length and the GAP analysis aligns the two sequences over a region of at least 250 amino acids. Even more preferably, the GAP analysis aligns the two sequences over their entire length.

With regard to a defined polypeptide, it will be appreciated that % identity figures higher than those provided above will encompass preferred embodiments. Thus, where applicable, in light of the minimum % identity figures, it is preferred that the CESA polypeptide of the invention comprises an amino acid sequence which is at least 40%, more preferably at least 45%, more preferably at least 50%, more preferably at least 55%, more preferably at least 60%, more preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 91%, more preferably at least 92%, more preferably at least

93%, more preferably at least 94%, more preferably at least 95%, more preferably at least 96%, more preferably at least 97%, more preferably at least 98%, more preferably at least 99%, more preferably at least 99.1%, more preferably at least 99.2%, more preferably at least 99.3%, more preferably at least 99.4%, more preferably at least 99.5%, more preferably at least 99.6%, more preferably at least 99.7%, more preferably at least 99.8%, and even more preferably at least 99.9% identical to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64.

By "variant" polypeptide is intended a polypeptide derived from the protein of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, by deletion (so-called truncation) or addition of one or more amino acids to the N-terminal and/or C-terminal end of the native protein; deletion or addition of one or more amino acids at one or more sites in the native protein; or substitution of one or more amino acids at one or more sites in the native protein. Such variants may result from, for example, genetic polymorphism or from human manipulation. Methods for such manipulations are generally known in the art.

"Derivatives" of a protein encompass peptides, oligopeptides, polypeptides, proteins and enzymes having amino acid substitutions, deletions and/or insertions relative to the unmodified protein in question and having similar biological and functional activity as the unmodified protein from which they are derived. Thus, functional variants and fragments of the CESA polypeptides, and nucleic acid molecules encoding them, also are within the scope of the present invention, and unless specifically described otherwise, irrespective of the origin of said polypeptide and irrespective of whether it occurs naturally. Various assays for functionality of a CESA polypeptide can be employed. For example, a functional variant or fragment of the CESA polypeptide can be assayed to determine its ability to confer CESA-inhibiting herbicides tolerance. By way of illustration, a CESA-inhibiting herbicides tolerance can be defined as insensitivity to CESA inhibiting herbicides sufficient to provide a determinable increase in tolerance to CESA-inhibiting herbicides in a plant or plant part comprising a recombinant polynucleotide encoding the variant or fragment of the CESA polypeptide, wherein the plant or plant part expresses the variant or fragment at up to about 0.5%, illustratively, about 0.05 to about 0.5%, about 0.1 to about 0.4%, and about 0.2 to about 0.3%, of the total cellular protein relative to a similarly treated control plant that does not express the variant or fragment.

In a preferred embodiment, the mutated CESA polypeptide is a functional variant or fragment of a cellulose synthase having the amino acid sequence set forth in SEQ ID NO: 1 or 3, wherein the functional variant or fragment has at least about 80% amino acid sequence identity to SEQ ID NO:1 or 3.

In other embodiments, the functional variant or fragment further has a CESA-inhibiting herbicides tolerance defined as insensitivity to CESA inhibiting herbicides sufficient to provide a determinable increase in tolerance to CESA-inhibiting herbicides in a plant or
5 plant part comprising a recombinant polynucleotide encoding the variant or fragment, wherein the plant or plant part expresses the variant or fragment at up to about 0.5% of the total cellular protein to a similarly treated control plant that does not express the variant or fragment.

10 "Homologues" of a protein encompass peptides, oligopeptides, polypeptides, proteins and enzymes having amino acid substitutions, deletions and/or insertions relative to the unmodified protein in question and having similar biological and functional activity as the unmodified protein from which they are derived.

15 In addition, one of ordinary skill in the art will further appreciate that changes can be introduced by mutation into the nucleotide sequences of the invention thereby leading to changes in the amino acid sequence of the encoded proteins without altering the biological activity of the proteins. Thus, for example, an isolated polynucleotide molecule encoding a mutated CESA polypeptide having an amino acid sequence that differs from that of SEQ ID
20 NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, can be created by introducing one or more nucleotide substitutions, additions, or deletions into the corresponding nucleotide sequence, such that one or more amino acid substitutions, additions or deletions are
25 introduced into the encoded protein. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR- mediated mutagenesis. Such variant nucleotide sequences are also encompassed by the present invention. For example, preferably, conservative amino acid substitutions may be made at one or more predicted preferably nonessential amino acid residues. A "nonessential" amino acid residue is a
30 residue that can be altered from the wild-type sequence of a protein without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity.

A deletion refers to removal of one or more amino acids from a protein.

35

An insertion refers to one or more amino acid residues being introduced into a predetermined site in a protein. Insertions may comprise N-terminal and/or C-terminal fusions as well as intra-sequence insertions of single or multiple amino acids. Generally, insertions within the amino acid sequence will be smaller than N- or C-terminal fusions, of
40 the order of about 1 to 10 residues. Examples of N- or C-terminal fusion proteins or peptides include the binding domain or activation domain of a transcriptional activator as used in the yeast two-hybrid system, phage coat proteins, (histidine)-6-tag, glutathione S-

transferase-tag, protein A, maltose-binding protein, dihydrofolate reductase, Tag•100 epitope, c-myc epitope, FLAG®-epitope, lacZ, CMP (calmodulin-binding peptide), HA epitope, protein C epitope and VSV epitope.

- 5 A substitution refers to replacement of amino acids of the protein with other amino acids having similar properties (such as similar hydrophobicity, hydrophilicity, antigenicity, propensity to form or break α -helical structures or β -sheet structures). Amino acid substitutions are typically of single residues, but may be clustered depending upon functional constraints placed upon the polypeptide and may range from 1 to 10 amino acids;
- 10 insertions will usually be of the order of about 1 to 10 amino acid residues. A conservative amino acid substitution is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine,
- 15 tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Such substitutions would not be made for conserved amino acid residues, or for amino acid
- 20 residues residing within a conserved motif. Conservative substitution tables are well known in the art (see for example Creighton (1984) Proteins. W.H. Freeman and Company (Eds).

Amino acid substitutions, deletions and/or insertions may readily be made using peptide synthetic techniques well known in the art, such as solid phase peptide synthesis and the

25 like, or by recombinant DNA manipulation. Methods for the manipulation of DNA sequences to produce substitution, insertion or deletion variants of a protein are well known in the art. For example, techniques for making substitution mutations at predetermined sites in DNA are well known to those skilled in the art and include M13 mutagenesis, T7-Gen in vitro mutagenesis (USB, Cleveland, OH), QuickChange Site Directed mutagenesis (Stratagene,

30 San Diego, CA), PCR-mediated site-directed mutagenesis or other site-directed mutagenesis protocols.

“Derivatives” further include peptides, oligopeptides, polypeptides which may, compared to the amino acid sequence of the naturally-occurring form of the protein, such as the protein

35 of interest, comprise substitutions of amino acids with non-naturally occurring amino acid residues, or additions of non-naturally occurring amino acid residues. “Derivatives” of a protein also encompass peptides, oligopeptides, polypeptides which comprise naturally occurring altered (glycosylated, acylated, prenylated, phosphorylated, myristoylated, sulphated etc.) or non-naturally altered amino acid residues compared to the amino acid

40 sequence of a naturally-occurring form of the polypeptide. A derivative may also comprise one or more non-amino acid substituents or additions compared to the amino acid sequence from which it is derived, for example a reporter molecule or other ligand,

covalently or non-covalently bound to the amino acid sequence, such as a reporter molecule which is bound to facilitate its detection, and non-naturally occurring amino acid residues relative to the amino acid sequence of a naturally-occurring protein. Furthermore, “derivatives” also include fusions of the naturally-occurring form of the protein with tagging peptides such as FLAG, HIS6 or thioredoxin (for a review of tagging peptides, see Terpe, Appl. Microbiol. Biotechnol. 60, 523-533, 2003).

“Orthologues” and “paralogues” encompass evolutionary concepts used to describe the ancestral relationships of genes. Paralogues are genes within the same species that have originated through duplication of an ancestral gene; orthologues are genes from different organisms that have originated through speciation, and are also derived from a common ancestral gene. A non-limiting list of examples of such orthologues is shown in Table 1. It will be understood by the person skilled in the art that the sequences of SEQ ID NOs: 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, as listed in Table 1 represent orthologues and paralogues to SEQ ID NO:1.

It is well-known in the art that paralogues and orthologues may share distinct domains harboring suitable amino acid residues at given sites, such as binding pockets for particular substrates or binding motifs for interaction with other proteins.

The term “domain” refers to a set of amino acids conserved at specific positions along an alignment of sequences of evolutionarily related proteins. While amino acids at other positions can vary between homologues, amino acids that are highly conserved at specific positions indicate amino acids that are likely essential in the structure, stability or function of a protein. Identified by their high degree of conservation in aligned sequences of a family of protein homologues, they can be used as identifiers to determine if any polypeptide in question belongs to a previously identified polypeptide family.

The term “motif” or “consensus sequence” refers to a short conserved region in the sequence of evolutionarily related proteins. Motifs are frequently highly conserved parts of domains, but may also include only part of the domain, or be located outside of conserved domain (if all of the amino acids of the motif fall outside of a defined domain).

In a preferred embodiment, the CESA polypeptide useful for the present invention, comprises one or more of the following motifs:

i) Motif 1a:

[V/I][A/V]G[V/I/F][S/T][Y/D/N/A]A[V/I/L][N/S/G][S/N]G[Y/F/E][Q/D/G/E/H][S/A]WG[P/A]L[F/M/L]G[K/R][L/V][F/L]F (SEQ ID NO: 78).

Preferably said motif is

[V/I]AG[V/I]S[Y/D/N]A[V/I][N/S][S/N]G[Y/F][Q/D]SWGPL[F/M/L]G[K/R]L[F/L]F (motif 1b; SEQ ID NO: 79).

More preferably said motif is

VAG[V/I]SYA[V/I]NSGYQSWGPL[F/M]GKL[F/L]F (motif 1c; SEQ ID NO:80)

5

ii) Motif 2a:

[V/L/I]W[S/A][V/A/I]LL[A/S]S[I/F/V][F/L][S/T][L/V][L/M/V/I]WV[R/K][I/V][N/D]PF (SEQ ID NO: 81)

Preferably, said motif is

10 VW[S/A][V/A/I]LL[A/S]S[I/F][F/L][S/T][L/V][L/M]WV[R/K][I/V][N/D]PF (motif 2b; SEQ ID NO: 82);

More preferably said motif is

VW[S/A][V/A/I]LLASIFSL[L/M]WV[R/K][I/V][N/D]PF (motif 2c; SEQ ID NO: 83)

15 Motifs 1a-c, 2a-c, given above were derived using the ClustalW algorithm to generate the alignments of cellulose synthase sequences (Figure 2 A – C) (Larkin et al., Bioinformatics 23:21 (2007) 2947 - 2948 pp. 28-36.). The motifs were essentially derived based on sequence alignment; highly conserved regions were identified that contain the site of mutations conferring azine-herbicide tolerance. Residues within square brackets represent
20 alternatives.

In a preferred embodiment, a CESA polypeptide as applied herein comprises, at least 1, at least 2, selected from the group comprising motifs 1a, 2a, as given above. Alternatively or in addition, in another preferred embodiment, a CESA polypeptide as applied herein

25 comprises at least 1, at least 2, motifs selected from the group comprising motifs 1b, 2b, as given above. Alternatively or in addition, in another preferred embodiment, a CESA polypeptide as applied herein comprises at least 1, at least 2, motifs selected from the group comprising motifs 1c, 2c, as given above.

30 Additionally or alternatively, the homologue of a CESA protein has in increasing order of preference at least 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%,
35 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% overall sequence identity to the amino acid represented by SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, provided that the homologous
40 protein comprises any one or more of the conserved motifs 1 and/or 2 as outlined above.

The overall sequence identity is determined using a global alignment algorithm, such as the Needleman Wunsch algorithm in the program GAP (GCG Wisconsin Package, Accelrys),

- preferably with default parameters and preferably with sequences of mature proteins (i.e. without taking into account secretion signals or transit peptides). Compared to overall sequence identity, the sequence identity will generally be higher when only conserved domains or motifs are considered. Preferably the motifs in a CESA polypeptide have, in
- 5 increasing order of preference, at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to any one or more of the motifs represented by SEQ ID NO: 78, 79, 80, 81, 82, and 83 (Motifs 1a, 1b, 1c, 2a, 2b, 2c).
- 10 Specialist databases exist for the identification of domains, for example, SMART (Schultz et al. (1998) *Proc. Natl. Acad. Sci. USA* 95, 5857-5864; Letunic et al. (2002) *Nucleic Acids Res* 30, 242-244), InterPro (Mulder et al., (2003) *Nucl. Acids. Res.* 31, 315-318), Prosite (Bucher and Bairoch (1994), A generalized profile syntax for biomolecular sequences motifs and its function in automatic sequence interpretation. (In) *ISMB-94; Proceedings 2nd*
- 15 *International Conference on Intelligent Systems for Molecular Biology*. Altman R., Brutlag D., Karp P., Lathrop R., Searls D., Eds., pp53-61, AAAI Press, Menlo Park; Hulo et al., *Nucl. Acids. Res.* 32:D134-D137, (2004)), or Pfam (Bateman et al., *Nucleic Acids Research* 30(1): 276-280 (2002)). A set of tools for in silico analysis of protein sequences is available on the ExPASy proteomics server (Swiss Institute of Bioinformatics (Gasteiger et al.,
- 20 ExPASy: the proteomics server for in-depth protein knowledge and analysis, *Nucleic Acids Res.* 31:3784-3788(2003)). Domains or motifs may also be identified using routine techniques, such as by sequence alignment.

Methods for the alignment of sequences for comparison are well known in the art, such

25 methods include GAP, BESTFIT, BLAST, FASTA and TFASTA. GAP uses the algorithm of Needleman and Wunsch ((1970) *J Mol Biol* 48: 443-453) to find the global (i.e. spanning the complete sequences) alignment of two sequences that maximizes the number of matches and minimizes the number of gaps. The BLAST algorithm (Altschul et al. (1990) *J Mol Biol* 215: 403-10) calculates percent sequence identity and performs a statistical analysis of the

30 similarity between the two sequences. The software for performing BLAST analysis is publicly available through the National Centre for Biotechnology Information (NCBI). Homologues may readily be identified using, for example, the ClustalW multiple sequence alignment algorithm (version 1.83), with the default pairwise alignment parameters, and a scoring method in percentage (See Figure 1). Global percentages of similarity and identity

35 may also be determined using one of the methods available in the MatGAT software package (Campanella et al., *BMC Bioinformatics*. 2003 Jul 10;4:29. MatGAT: an application that generates similarity/identity matrices using protein or DNA sequences.). Minor manual editing may be performed to optimise alignment between conserved motifs, as would be apparent to a person skilled in the art. Furthermore, instead of using full-length sequences

40 for the identification of homologues, specific domains may also be used. The sequence identity values may be determined over the entire nucleic acid or amino acid sequence or over selected domains or conserved motif(s), using the programs mentioned above using

the default parameters. For local alignments, the Smith-Waterman algorithm is particularly useful (Smith TF, Waterman MS (1981) J. Mol. Biol 147(1);195-7).

The proteins of the invention may be altered in various ways including amino acid
 5 substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985) PNAS, 82:488-492; Kunkel et al. (1987) Methods in Enzymol. 154:367-382; U.S. Patent No. 4,873,192; Walker and Gaastra,
 10 eds. (1983) Techniques in Molecular Biology (MacMillan Publishing Company, New York) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff et al. (1978) Atlas of Protein Sequence and Structure (Natl. Biomed. Res. Found., Washington, D. C). Conservative substitutions, such as exchanging one amino acid with
 15 another having similar properties, may be preferable.

Alternatively, variant nucleotide sequences can be made by introducing mutations randomly along all or part of a coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened to identify mutants that encode proteins that retain activity. For
 20 example, following mutagenesis, the encoded protein can be expressed recombinantly, and the activity of the protein can be determined using standard assay techniques.

The inventors of the present invention have found that by substituting one or more of the key amino acid residues of the CESA enzyme of SEQ ID NO: 1 or 3, e.g. by
 25 employing one of the above described methods to mutate the CESA encoding nucleic acids, the tolerance or resistance to particular CESA-inhibiting herbicides, collectively named azines, and described in greater detail herein below, could be remarkably increased Preferred substitutions of mutated CESA are those that increase the herbicide tolerance of the plant, but leave the biological activity of the cellulose
 30 synthase activity substantially unaffected.

Accordingly, in another object of the present invention refers to a CESA polypeptide, comprising the sequence of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40,
 35 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, a variant, derivative, orthologue, paralogue or homologue thereof, the key amino acid residues of which is substituted by any other amino acid.

It will be understood by the person skilled in the art that amino acids located in a close
 40 proximity to the positions of amino acids mentioned below may also be substituted. Thus, in another embodiment the variant of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62,

63, or 64, a variant, derivative, orthologue, paralogue or homologue thereof comprises a mutated CESA, wherein an amino acid ± 3 , ± 2 or ± 1 amino acid positions from a key amino acid is substituted by any other amino acid.

- 5 Based on techniques well-known in the art, a highly characteristic sequence pattern can be developed, by means of which further of mutated CESA candidates with the desired activity may be searched.

10 Searching for further mutated CESA candidates by applying a suitable sequence pattern would also be encompassed by the present invention. It will be understood by a skilled reader that the present sequence pattern is not limited by the exact distances between two adjacent amino acid residues of said pattern. Each of the distances between two neighbours in the above patterns may, for example, vary independently of each other by up to ± 10 , ± 5 , ± 3 , ± 2 or ± 1 amino acid positions without substantially affecting the desired
15 activity.

Furthermore, by applying the method of site directed mutagenesis, in particular saturation mutagenesis (see e.g. Schenk et al., Biospektrum 03/2006, pages 277-279) PCR based site-directed mutagenesis (e.g. directed mutagenesis kit, Stratagene, California, USA or
20 GeneArt Mutagenesis Service, ThermoFisher Scientific Inc., Massachusetts, USA) or systematic mutagenesis (GeneArt Systematic Mutagenesis Service, ThermoFisher Scientific Inc., Massachusetts, USA), the inventors of the present invention have identified and generated specific amino acid substitutions and combinations thereof, which - when introduced into a plant by transforming and expressing the respective mutated CESA
25 encoding nucleic acid - confer increased herbicide resistance or tolerance to a CESA inhibiting herbicide to said plant.

Thus, in preferred embodiment, the variant or derivative of the CESA polypeptide refers to a mutated CESA polypeptide which comprises one or more of the following motifs:

30

- i) Motif 1a:
[V/I] [A/V] G [V/I/F] [S/T] [Y/D/N/A] A [V/I/L] [N/S/G] [S/N] G [Y/F/E] [Q/D/G/E/H]
[S/A] WG [P/A] L [F/M/L] G [K/R] [L/V] [F/L] F (SEQ ID NO: 78).

Preferably said motif is

35

[V/I] A G [V/I] S [Y/D/N] A [V/I] [N/S] [S/N] G [Y/F] [Q/D] SWGPL [F/M/L] G [K/R] L
[F/L] F (motif 1b; SEQ ID NO: 79).

More preferably said motif is

VAG [V/I] SYA [V/I] NSGYQSWGPL [F/M] GKL [F/L] F (motif 1c; SEQ ID NO:80);

40

Wherein the amino acid at position 5, 16, 17, and/or 20 within said motif is substituted by any other amino acid.

- ii) Motif 2a:
 [V/L/I]W [S/A] [V/A/I]LL [A/S]S [I/F/V] [F/L][S/T] [L/V] [L/M/V/I]WV [R/K] [I/V]
 [N/D]PF (SEQ ID NO: 81)
 Preferably, said motif is
 5 VW [S/A] [V/A/I]LL [A/S]S [I/F] [F/L][S/T][L/V] [L/M]WV [R/K] [I/V] [N/D]PF (motif
 2b; SEQ ID NO: 82);
 More preferably said motif is
 VW [S/A] [V/A/I]LLASIFSL [L/M]WV [R/K]I [N/D]PF (motif 2c; SEQ ID NO: 83)
 10 Wherein the amino acid at position 8, and/or 11 within said motif is substituted by
 any other amino acid

In a more preferred embodiment, the amino acid corresponding to position 5 of motif 1a, 1b,
 or 1c is:

Arg, His, Lys, Asp, Glu, Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp;

15 And/or

the amino acid corresponding to position 16 of motif 1a, 1b, or 1c is

Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

And/or

the amino acid corresponding to position 17 of motif 1a, 1b, or 1c is

20 Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Gly, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

And/or

the amino acid corresponding to position 20 of motif 1a, 1b, or 1c is

Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

25 In another more preferred embodiment, the amino acid corresponding to position 8 of motif
 2a, 2b, or 2c is:

Arg, His, Lys, Asp, Glu, Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp;

And/or

the amino acid corresponding to position 11 of motif 2a, 2b, or 2c is

30 Arg, His, Lys, Asp, Glu, Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

In a particularly preferred embodiment,

the amino acid corresponding to position 5 of motif 1a, 1b, or 1c is Phe,

And/or

the amino acid corresponding to position 16 of motif 1a, 1b, or 1c is Asp,

35 And/or

the amino acid corresponding to position 17 of motif 1a, 1b, or 1c is Leu,

And/or

the amino acid corresponding to position 20 of motif 1a, 1b, or 1c is

Arg.

40

In another more preferred embodiment,

the amino acid corresponding to position 8 of motif 2a, 2b, or 2c is Phe,

And/or

the amino acid corresponding to position 11 of motif 2a, 2b, or 2c is Leu,

In another preferred embodiment, the variant or derivative of the CESA polypeptide refers to a CESA polypeptide comprising SEQ ID NO: 1, an orthologue, paralogue, or homologue thereof, wherein the amino acid sequence differs from the wildtype amino acid sequence of a CESA polypeptide at one or more positions corresponding to the following positions of SEQ ID NO:1:

998, 1009, 1010, 1013, 1052, 1055.

Examples of differences at these amino acid positions include, but are not limited to, one or more of the following:

the amino acid at or corresponding to position 998 is other than serine;

the amino acid at or corresponding to position 1009 is other than glycine;

the amino acid at or corresponding to position 1010 is other than proline;

the amino acid at or corresponding to position 1013 is other than glycine,

the amino acid at or corresponding to position 1052 is other than serine,

the amino acid at or corresponding to position 1055 is other than serine,

In some embodiments, the mutated CESA enzyme comprising SEQ ID NO: 1, an orthologue, paralogue, or homologue thereof, comprises one or more of the following:

the amino acid corresponding to position 998 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp;

the amino acid corresponding to position 1009 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

the amino acid corresponding to position 1010 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Gly, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

the amino acid corresponding to position 1013 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

the amino acid corresponding to position 1052 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

the amino acid corresponding to position 1055 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp.

In a preferred embodiment, the amino acid corresponding to position 1009 of SEQ ID NO: 1 is Asp.

In another preferred embodiment, the amino acid corresponding to position 1010 of SEQ ID NO: 1 is Leu.

5 In another preferred embodiment, the amino acid corresponding to position 1013 of SEQ ID NO: 1 is Arg.

In another preferred embodiment, the amino acid corresponding to position 983 of SEQ ID NO: 3 is Phe.

10 In another preferred embodiment, the amino acid corresponding to position 1037 of SEQ ID NO: 3 is Phe.

In another preferred embodiment, the amino acid corresponding to position 1040 of SEQ ID NO: 3 is Leu.

15

It will be within the knowledge of the skilled artisan to identify conserved regions and motifs shared between the homologues, orthologues and paralogues encoded by SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, such as those depicted in Table 1. Having identified such conserved regions that may represent suitable binding motifs, amino acids
20 corresponding to the amino acids listed below in Table 2, can be chosen to be substituted by any other amino acid, for example by conserved amino acids, preferably by the amino acid substitutions described SUPRA using SEQ ID NO:1 as reference.

25 **Table 2** provides an overview of positions in the orthologues and homologues to SEQ ID NO:1, i.e. the corresponding positions in SEQ ID NOs: 1 to 64.

Table 2

ID	Pos 1	Pos 2	Pos 3	Pos 4	Pos 5	Pos 6
1	S998	G1009	P1010	G1013	S1052	S1055
2	S1001	G1012	P1013	G1016	S1055	T1058
3	S983	G994	P995	G998	S1037	S1040
4	S967	G978	P979	G982	S1021	S1024
5	S987	G998	P999	G1002	S1041	T1044
6	S1002	G1013	P1014	G1017	S1056	T1059
7	S944	G955	P956	G959	S998	S1001
8	S901	G912	P913	G916	S955	S958
9	S1005	G1016	P1017	G1020	S1059	T1062
10	S985	G996	P997	G1000	S1039	S1042
11	S991	G1002	P1003	G1006	S1045	S1048
12	S900	G911	P912	G915	S954	S957
13	S991	G1002	P1003	G1006	S1045	S1048

14	S1003	G1014	P1015	G1018	S1057	S1060
15	S974	G985	P986	G989	S1028	S1031
16	S995	G1006	P1007	G1010	S1049	S1052
17	S1011	G1022	P1023	G1026	S1065	S1068
18	S514	G525	P526	G529	S568	S571
19	S1000	G1011	P1012	G1015	S1054	S1057
20	S997	G1008	P1009	G1012	S1051	S1054
21	S900	G911	P912	G915	S954	S957
22	-	-	-	-	-	-
23	S992	G1003	P1004	G1007	S1046	S1049
24	S994	G1005	P1006	G1009	S1048	S1051
25	S1006	G1017	P1018	G1021	S1060	S1063
26	S951	G962	P963	G966	S1005	S1008
27	S996	G1007	P1008	G1011	S1050	T1053
28	S997	G1008	P1009	G1012	S1051	S1054
29	T891	G902	A903	G906	S945	S948
30	S891	G902	P903	G906	S945	S948
31	S957	G968	P969	G972	S1011	S1014
32	S884	G895	P896	G899	S938	S941
33	T908	G919	A920	G923	S962	S965
34	S1002	G1013	P1014	G1017	S1056	S1059
35	S1002	G1013	P1014	G1017	S1056	S1059
36	S892	G903	P904	G907	S946	S949
37	S957	G968	P969	G972	S1011	S1014
38	S1013	G1024	P1025	G1028	S1067	T1070
39	S905	G916	P917	G920	S959	S962
40	S951	G962	P963	G966	S1005	S1008
41	S998	G1009	P1010	G1013	S1052	S1055
42	S874	G885	P886	G889	S928	S931
43	S1013	G1024	P1025	G1028	S1067	T1070
44	S991	G1002	P1003	G1006	S1045	S1048
45	S992	G1003	P1004	G1007	S1046	S1049
46	S1013	G1024	P1025	G1028	S1067	S1070
47	S996	G1007	P1008	G1011	S1050	T1053
48	S906	G917	P918	G921	S960	S963
49	S991	G1002	P1003	G1006	S1045	S1048
50	S1009	G1020	P1021	G1024	S1063	S1066
51	S993	G1004	P1005	G1008	S1047	S1050
52	S1091	S1102	K1103	G1106	I1145	S1148
53	S935	S946	K947	G950	I989	S992

54	A778	G790	A791	A794	S834	S837
55	S999	G1010	P1011	G1014	S1053	S1056
56	S1009	G1020	P1021	G1024	S1063	S1066
57	S1010	G1021	P1022	G1025	S1064	S1067
58	S754	G765	P766	G769	S808	S811
59	S883	G894	P895	G898	S937	S940
60	A1033	G1044	K1045	G1048	I1087	S1090
61	S973	G984	P985	G988	S1027	S1030
62	S981	G992	P993	G996	S1035	S1038
63	S1046	S1057	K1058	G1061	I1100	S1103
64	A1134	S1145	K1146	G1149	M1188	S1191

Another object refers to a method of identifying a nucleotide sequence encoding a mutated CESA which is resistant or tolerant to a CESA-inhibiting herbicide, the method comprising:

- a) generating a library of mutated CESA-encoding nucleic acids,
- 5 b) screening a population of the resulting mutated CESA-encoding nucleic acids by expressing each of said nucleic acids in a cell or plant and treating said cell or plant with a CESA-inhibiting herbicide,
- c) comparing the CESA-inhibiting herbicide-tolerance levels provided by said population of mutated CESA encoding nucleic acids with the CESA-inhibiting herbicide-tolerance
- 10 level provided by a control CESA-encoding nucleic acid,
- d) selecting at least one mutated CESA-encoding nucleic acid that provides a significantly increased level of tolerance to a CESA-inhibiting herbicide as compared to that provided by the control CESA-encoding nucleic acid.

- 15 In a preferred embodiment, the mutated CESA-encoding nucleic acid selected in step d) provides at least 2-fold as much resistance or tolerance of a cell or plant to a CESA-inhibiting herbicide as compared to that provided by the control CESA-encoding nucleic acid.

- 20 In a further preferred embodiment, the mutated CESA-encoding nucleic acid selected in step d) provides at least 2-fold, at least 5-fold, at least 10-fold, at least 20-fold, at least 50-fold, at least 100-fold, at least 500-fold, as much resistance or tolerance of a cell or plant to a CESA-inhibiting herbicide as compared to that provided by the control CESA-encoding nucleic acid.

25

The resistance or tolerance can be determined by generating a transgenic plant or host cell, preferably a plant cell, comprising a nucleic acid sequence of the library of step a) and comparing said transgenic plant with a control plant or host cell, preferably a plant cell.

- 30 Another object refers to a method of identifying a plant or algae containing a nucleic acid

comprising a nucleotide sequence encoding a wildtype or mutated CESA which is resistant or tolerant to a CESA-inhibiting herbicide, the method comprising:

- a) identifying an effective amount of a CESA-inhibiting herbicide in a culture of plant cells or green algae that leads to death of said cells.
- 5 b) treating said plant cells or green algae with a mutagenizing agent,
- c) contacting said mutagenized cells population with an effective amount of CESA-inhibiting herbicide, identified in a),
- d) selecting at least one cell surviving these test conditions,
- e) PCR-amplification and sequencing of CESA genes from cells selected in d) and
- 10 comparing such sequences to wild-type CESA gene sequences, respectively.

In a preferred embodiment, said mutagenizing agent is ethylmethanesulfonate (EMS).

Many methods well known to the skilled artisan are available for obtaining suitable candidate nucleic acids for identifying a nucleotide sequence encoding a wildtype or mutated CESA from a variety of different potential source organisms including microbes, plants, fungi, algae, mixed cultures etc. as well as environmental sources of DNA such as soil. These methods include inter alia the preparation of cDNA or genomic DNA libraries, the use of suitably degenerate oligonucleotide primers, the use of probes based upon

20 known sequences or complementation assays (for example, for growth upon tyrosine) as well as the use of mutagenesis and shuffling in order to provide recombined or shuffled wildtype or mutated CESA-encoding sequences.

Nucleic acids comprising candidate and control CESA encoding sequences can be expressed in yeast, in a bacterial host strain, in an alga or in a higher plant such as tobacco or Arabidopsis and the relative levels of inherent tolerance of the CESA encoding sequences screened according to a visible indicator phenotype of the transformed strain or plant in the presence of different concentrations of the selected CESA-inhibiting herbicide. Dose responses and relative shifts in dose responses associated with these indicator

30 phenotypes (formation of necrosis, growth inhibition, herbicidal effect etc) are conveniently expressed in terms, for example, of GR50 (concentration for 50% reduction of growth) or MIC (minimum inhibitory concentration) values where increases in values correspond to increases in inherent tolerance of the expressed CESA. For example, in a relatively rapid assay system based upon transformation of Arabidopsis as described in the Example section (Example 7), each wildtype or mutated CESA encoding sequence may be expressed, for example, as a DNA sequence under expression control of a suitable promoter and T1 plants can be selected for differential tolerance to selected CESA-inhibiting herbicides, measured by growth.

In another embodiment, candidate nucleic acids are transformed into plant material to generate a transgenic plant, regenerated into morphologically normal fertile plants which

40 are then measured for differential tolerance to selected CESA-inhibiting herbicides as described in the Example section hereinafter. Many suitable methods for transformation

using suitable selection markers such as kanamycin, binary vectors such as from Agrobacterium and plant regeneration as, for example, from tobacco leaf discs are well known in the art. Optionally, a control population of plants is likewise transformed with a nucleic acid expressing the control CESA. Alternatively, an untransformed dicot plant such as Arabidopsis or Tobacco can be used as a control since this, in any case, expresses its own endogenous CESA. The average, and distribution, of herbicide tolerance levels of a range of primary plant transformation events or their progeny to CESA-inhibiting herbicides described supra are evaluated in the normal manner based upon plant damage, meristematic bleaching symptoms etc. at a range of different concentrations of herbicides. These data can be expressed in terms of, for example, GR50 values derived from dose/response curves having "dose" plotted on the x-axis and "percentage kill", "herbicidal effect", "numbers of emerging green plants" etc. plotted on the y-axis where increased GR50 values correspond to increased levels of inherent tolerance of the expressed CESA. Herbicides can suitably be applied pre-emergence or post-emergence.

Another object of the present invention refers to an isolated and or recombinantly produced and/or synthetic nucleic acid encoding a mutated CESA as disclosed SUPRA, wherein the nucleic acid comprises the nucleotide sequence of SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or a variant or derivative thereof.

In one embodiment, the nucleic acid is identifiable by a method as defined above.

For the purposes of the invention "recombinant" means with regard for example to a nucleic acid sequence, an expression cassette (= gene construct, nucleic acid construct) or a vector containing the nucleic acid sequence according to the invention or an organism transformed by said nucleic acid sequences, expression cassette or vector according to the invention all those constructions produced by genetic engineering methods in which either (a) the nucleic acid sequence comprising the sequence of SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or a homolog thereof, or its derivatives or parts thereof; or

(b) a genetic control sequence functionally linked to the nucleic acid sequence described under (a), for example a 3'- and/or 5'- genetic control sequence such as a promoter or terminator, or

(c) (a) and (b);

are not found in their natural, genetic environment or have been modified by genetic engineering methods, wherein the modification may by way of example be a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues.

"Natural genetic environment" means the natural genomic or chromosomal locus in the organism of origin or inside the host organism or presence in a genomic library. In the case of a genomic library the natural genetic environment of the nucleic acid sequence is preferably retained at least in part. The environment borders the nucleic acid sequence at

least on one side and has a sequence length of at least 50 bp, preferably at least 500 bp, particularly preferably at least 1,000 bp, most particularly preferably at least 5,000 bp. A naturally occurring expression cassette – for example the naturally occurring combination of the natural promoter of the nucleic acid sequence according to the invention with the
 5 corresponding gene – turns into a transgenic expression cassette when the latter is modified by unnatural, synthetic (“artificial”) methods such as by way of example a mutagenation. Appropriate methods are described by way of example in US 5,565,350 or WO 00/15815

10 In a preferred embodiment, the encoded mutated CESA is a variant of SEQ ID NO: 1, which includes one or more of the following:

the amino acid corresponding to position 998 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp;

15 the amino acid corresponding to position 1009 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

the amino acid corresponding to position 1010 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Gly, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

20

the amino acid corresponding to position 1013 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, Ser, Thr, Asn, Gln, Cys, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

the amino acid corresponding to position 1052 of SEQ ID NO:1 is

25 Arg, His, Lys, Asp, Glu, Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp,

the amino acid corresponding to position 1055 of SEQ ID NO:1 is

Arg, His, Lys, Asp, Glu, , Thr, Asn, Gln, Cys, Gly, Pro, Ala, Val, Leu, Ile, Met, Phe, Tyr, or Trp.

30 In a preferred embodiment, the amino acid corresponding to position 1009 of SEQ ID NO: 1 is Asp.

In another preferred embodiment, the amino acid corresponding to position 1010 of SEQ ID NO: 1 is Leu.

35

In another preferred embodiment, the amino acid corresponding to position 1013 of SEQ ID NO: 1 is Arg.

In another preferred embodiment, the amino acid corresponding to position 983 of SEQ ID
 40 NO: 3 is Phe.

In another preferred embodiment, the amino acid corresponding to position 1037 of SEQ ID

NO: 3 is Phe.

In another preferred embodiment, the amino acid corresponding to position 1040 of SEQ ID NO: 3 is Leu.

5

In other aspects, the present invention encompasses a progeny or a descendant of a CESA-inhibiting herbicides-tolerant plant of the present invention as well as seeds derived from the CESA-inhibiting herbicides-tolerant plants of the invention and cells derived from the CESA-inhibiting herbicides-tolerant plants of the invention.

10

In some embodiments, the present invention provides a progeny or descendant plant derived from a plant comprising in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, wherein the progeny or descendant plant comprises in at least some of its cells the recombinant polynucleotide operably linked to the promoter, the expression of the wildtype or mutated CESA polypeptide conferring to the progeny or descendant plant tolerance to the CESA-inhibiting herbicides.

15

20

In one embodiment, seeds of the present invention preferably comprise the CESA-inhibiting herbicides-tolerance characteristics of the CESA-inhibiting herbicides-tolerant plant. In other embodiments, a seed is capable of germination into a plant comprising in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the progeny or descendant plant tolerance to the CESA-inhibiting herbicides.

25

In some embodiments, plant cells of the present invention are capable of regenerating a plant or plant part. In other embodiments, plant cells are not capable of regenerating a plant or plant part. Examples of cells not capable of regenerating a plant include, but are not limited to, endosperm, seed coat (testa & pericarp), and root cap.

30

In another embodiment, the present invention provides a plant cell of or capable of regenerating a plant comprising in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to the CESA-inhibiting herbicides, wherein the plant cell comprises the recombinant polynucleotide operably linked to a promoter.

35

40

In other embodiments, the present invention provides a plant cell comprising a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable

of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the cell tolerance to the CESA-inhibiting herbicides.

5 In another embodiment, the invention refers to a plant cell transformed by a nucleic acid encoding a wildtype or mutated CESA polypeptide according to the present invention or to a plant cell which has been mutated to obtain a plant expressing a nucleic acid encoding a wildtype or mutated CESA polypeptide according to the present invention, wherein expression of the nucleic acid in the plant cell results in increased resistance or tolerance to a CESA-inhibiting herbicide as compared to a wild type variety of the plant cell. Preferably, the wildtype or mutated CESA polypeptide encoding nucleic acid comprises a polynucleotide sequence selected from the group consisting of: a) a polynucleotide as shown in SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or a variant or derivative thereof; b) a polynucleotide encoding a polypeptide as shown in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, or a variant or derivative thereof; c) a polynucleotide comprising at least 60 consecutive nucleotides of any of a) or b); and d) a polynucleotide complementary to the polynucleotide of any of a) through c).

20

In some aspects, the present invention provides a plant product prepared from the CESA-inhibiting herbicides-tolerant plants hereof. In some embodiments, examples of plant products include, without limitation, grain, oil, and meal. In one embodiment, a plant product is plant grain (e.g., grain suitable for use as feed or for processing), plant oil (e.g., oil suitable for use as food or biodiesel), or plant meal (e.g., meal suitable for use as feed).

25

In one embodiment, a plant product prepared from a plant or plant part is provided, wherein the plant or plant part comprises in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the a plant or plant part tolerance to the CESA-inhibiting herbicides.

30

In another embodiment, the invention refers to a method of producing a transgenic plant cell with an increased resistance to a CESA-inhibiting herbicide as compared to a wild type variety of the plant cell comprising, transforming the plant cell with an expression cassette comprising a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide.

40

In another embodiment, the invention refers to a method of producing a transgenic plant comprising, (a) transforming a plant cell with an expression cassette comprising a

polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, and (b) generating a plant with an increased resistance to CESA-inhibiting herbicide from the plant cell.

5

In some aspects, the present invention provides a method for producing a CESA-inhibiting herbicides-tolerant plant. In one embodiment, the method comprises: regenerating a plant from a plant cell transformed with a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to the CESA-inhibiting herbicides.

10

The term “expression/expressing” or “gene expression” means the transcription of a specific gene or specific genes or specific genetic construct. The term “expression” or “gene expression” in particular means the transcription of a gene or genes or genetic construct into structural RNA (rRNA, tRNA) or mRNA with or without subsequent translation of the latter into a protein. The process includes transcription of DNA and processing of the resulting mRNA product.

15

To obtain the desired effect, i.e. plants that are tolerant or resistant to the CESA-inhibiting herbicide derivative herbicide of the present invention, it will be understood that the at least one nucleic acid is “over-expressed” by methods and means known to the person skilled in the art.

20

The term “increased expression” or “overexpression” as used herein means any form of expression that is additional to the original wild-type expression level. Methods for increasing expression of genes or gene products are well documented in the art and include, for example, overexpression driven by appropriate promoters, the use of transcription enhancers or translation enhancers. Isolated nucleic acids which serve as promoter or enhancer elements may be introduced in an appropriate position (typically upstream) of a non-heterologous form of a polynucleotide so as to upregulate expression of a nucleic acid encoding the polypeptide of interest. For example, endogenous promoters may be altered in vivo by mutation, deletion, and/or substitution (see, Kmiec, US 5,565,350; Zarling et al., WO9322443), or isolated promoters may be introduced into a plant cell in the proper orientation and distance from a gene of the present invention so as to control the expression of the gene. If polypeptide expression is desired, it is generally desirable to include a polyadenylation region at the 3'-end of a polynucleotide coding region. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA. The 3' end sequence to be added may be derived from, for example, the nopaline synthase or octopine synthase genes, or alternatively from another plant gene, or less preferably from any other eukaryotic gene. An intron sequence may also be added to the 5' untranslated region (UTR) or the coding sequence of the partial coding sequence

25

30

35

40

to increase the amount of the mature message that accumulates in the cytosol. Inclusion of a spliceable intron in the transcription unit in both plant and animal expression constructs has been shown to increase gene expression at both the mRNA and protein levels up to 1000-fold (Buchman and Berg (1988) *Mol. Cell Biol.* 8: 4395-4405; Callis et al. (1987) *Genes Dev* 1:1183-1200). Such intron enhancement of gene expression is typically greatest when placed near the 5' end of the transcription unit. Use of the maize introns Adh1-S intron 1, 2, and 6, the Bronze-1 intron are known in the art. For general information see: *The Maize Handbook*, Chapter 116, Freeling and Walbot, Eds., Springer, N.Y. (1994).

10

Where appropriate, nucleic acid sequences may be optimized for increased expression in a transformed plant. For example, coding sequences that comprise plant-preferred codons for improved expression in a plant can be provided. See, for example, Campbell and Gowri (1990) *Plant Physiol.*, 92: 1-11 for a discussion of host-preferred codon usage. Methods also are known in the art for preparing plant-preferred genes. See, for example, U.S. Patent Nos. 5,380,831, and 5,436,391, and Murray et al. (1989) *Nucleic Acids Res.* 17:477-498.

15

Consequently, wildtype or mutated CESA nucleic acids of the invention are provided in expression cassettes for expression in the plant of interest. The cassette will include regulatory sequences operably linked to a wildtype or mutated CESA nucleic acid sequence of the invention. The term "regulatory element" as used herein refers to a polynucleotide that is capable of regulating the transcription of an operably linked polynucleotide. It includes, but not limited to, promoters, enhancers, introns, 5' UTRs, and 3' UTRs. By "operably linked" is intended a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame. The cassette may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes.

25

30

Such an expression cassette is provided with a plurality of restriction sites for insertion of the wildtype or mutated CESA nucleic acid sequence to be under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes. The expression cassette of the present invention will include in the 5'-3' direction of transcription, a transcriptional and translational initiation region (i.e., a promoter), a wildtype or mutated CESA encoding nucleic acid sequence of the invention, and a transcriptional and translational termination region (i.e., termination region) functional in plants. The promoter may be native or analogous, or foreign or heterologous, to the plant host and/or to the wildtype or mutated CESA nucleic acid sequence of the invention. Additionally, the promoter may be the natural sequence or

40

alternatively a synthetic sequence. Where the promoter is "foreign" or "heterologous" to the plant host, it is intended that the promoter is not found in the native plant into which the promoter is introduced. Where the promoter is "foreign" or "heterologous" to the wildtype or mutated CESA nucleic acid sequence of the invention, it is intended that the promoter is not the native or naturally occurring promoter for the operably linked wildtype or mutated CESA nucleic acid sequence of the invention. As used herein, a chimeric gene comprises a coding sequence operably linked to a transcription initiation region that is heterologous to the coding sequence. While it may be preferable to express the wildtype or mutated CESA nucleic acids of the invention using heterologous promoters, the native promoter sequences may be used. Such constructs would change expression levels of the wildtype or mutated CESA protein in the plant or plant cell. Thus, the phenotype of the plant or plant cell is altered.

The termination region may be native with the transcriptional initiation region, may be native with the operably linked wildtype or mutated CESA sequence of interest, may be native with the plant host, or may be derived from another source (i.e., foreign or heterologous to the promoter, the wildtype or mutated CESA nucleic acid sequence of interest, the plant host, or any combination thereof). Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also Guerineau et al. (1991) *Mol. Gen. Genet.* 262: 141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon et al. (1991) *Genes Dev.* 5: 141-149; Mogen et al. (1990) *Plant Cell* 2: 1261-1272; Munroe et al. (1990) *Gene* 91: 151-158; Ballas et al. (1989) *Nucleic Acids Res.* 17:7891-7903; and Joshi et al. (1987) *Nucleic Acid Res.* 15:9627-9639. Where appropriate, the gene(s) may be optimized for increased expression in the transformed plant. That is, the genes can be synthesized using plant-preferred codons for improved expression. See, for example, Campbell and Gowri (1990) *Plant Physiol.* 92: 1-11 for a discussion of host-preferred codon usage. Methods are available in the art for synthesizing plant-preferred genes. See, for example, U.S. Patent Nos. 5,380,831, and 5,436,391, and Murray et al. (1989) *Nucleic Acids Res.* 17:477-498.

While the polynucleotides of the invention may find use as selectable marker genes for plant transformation, the expression cassettes of the invention can include another selectable marker gene for the selection of transformed cells. Selectable marker genes, including those of the present invention, are utilized for the selection of transformed cells or tissues. Marker genes include, but are not limited to, genes encoding antibiotic resistance, such as those encoding neomycin phosphotransferase II (NEO) and hygromycin phosphotransferase (HPT), as well as genes conferring resistance to herbicidal compounds, such as glufosinate ammonium, bromoxynil, imidazolinones, and 2,4-dichlorophenoxyacetate (2,4-D). See generally, Yarranton (1992) *Curr. Opin. Biotech.* 3 :506-511 ; Christopherson et al (1992) *Proc. Natl. Acad. Sci. USA* 89:6314-6318; Yao et al. (1992) *Cell* 71:63-72; Reznikoff (1992) *Mol Microbiol* 6:2419-2422; Barkley et al (1980) in *The Operon*, pp. 177-220; Hu et al (1987) *Cell* 48:555-566; Brown et al (1987) *Cell* 49:603-612; Figge et al (1988) *Cell* 52:713-722; Deuschle et al (1989)

- Proc. Natl Acad. Sci USA 86:5400-5404; Fuerst et al (1989) Proc. Natl Acad. Sci USA 86:2549-2553; Deuschle et al (1990) Science 248:480-483; Gossen (1993) Ph.D. Thesis, University of Heidelberg; Reines et al (1993) Proc. Natl Acad. Sci USA 90: 1917-1921; Labow et al (1990) Mol Cell Biol 10:3343-3356; Zambretti et al (1992) Proc. Natl Acad. Sci USA 89:3952-3956; Bairn et al (1991) Proc. Natl Acad. Sci USA 88:5072-5076; Wyborski et al (1991) Nucleic Acids Res. 19:4647-4653; Hillenand-Wissman (1989) Topics Mol Struct. Biol 10: 143- 162; Degenkolb et al (1991) Antimicrob. Agents Chemother. 35: 1591-1595; Kleinschmidt et al (1988) Biochemistry 27: 1094-1104; Bonin (1993) Ph.D. Thesis, University of Heidelberg; Gossen et al (1992) Proc. Natl Acad. Sci USA 89:5547-5551; Oliva et al (1992) Antimicrob. Agents Chemother. 36:913-919; Hlavka et al (1985) Handbook of Experimental Pharmacology, Vol. 78 (Springer-Verlag, Berlin); Gill et al (1988) Nature 334:721-724. The above list of selectable marker genes is not meant to be limiting. Any selectable marker gene can be used in the present invention.
- Further, additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats, and other such well -characterized sequences that may be deleterious to gene expression. The G-C content of the sequence may be adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. Also, if desired, sequences can be readily modified to avoid predicted hairpin secondary mRNA structures. Nucleotide sequences for enhancing gene expression can also be used in the plant expression vectors. These include, for example, introns of the maize Adh gene Adh1-S intron 1, 2, and 6 (Callis et al. Genes and Development 1 : 1183-1200, 1987), and leader sequences, (W-sequence) from the Tobacco Mosaic virus (TMV), Maize Chlorotic Mottle Virus and Alfalfa Mosaic Virus (Gallie et al. Nucleic Acid Res. 15:8693-8711 , 1987 and Skuzeski et al. Plant Mol. Biol. 15:65-79, 1990). The first intron from the shrunken-1 locus of maize has been shown to increase expression of genes in chimeric gene constructs. U.S. Pat. Nos. 5,424,412 and 5,593,874 disclose the use of specific introns in gene expression constructs, and Gallie et al. (Plant Physiol. 106:929-939, 1994) also have shown that introns are useful for regulating gene expression on a tissue specific basis. To further enhance or to optimize gene expression, the plant expression vectors of the invention also may contain DNA sequences containing matrix attachment regions (MARs). Plant cells transformed with such modified expression systems, then, may exhibit overexpression or constitutive expression of a nucleotide sequence of the invention.

The invention further provides an isolated recombinant expression vector comprising the expression cassette containing a wildtype or mutated CESA nucleic acid nucleic acid as described above, wherein expression of the vector in a host cell results in increased tolerance to a CESA-inhibiting herbicide as compared to a wild type variety of the host cell. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid," which

refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors." In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses, and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operably linked to the nucleic acid sequence to be expressed. Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cells and those that direct expression of the nucleotide sequence only in certain host cells or under certain conditions. It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of polypeptide desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce polypeptides or peptides, including fusion polypeptides or peptides, encoded by nucleic acids as described herein (e.g., wildtype or mutated CESA polypeptides, fusion polypeptides, etc.)

Expression vectors may additionally contain 5' leader sequences in the expression construct. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyo carditis 5' noncoding region) (Elroy-Stein et al. (1989) PNAS, 86:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Gallie et al. (1995) Gene 165(2):233-238), MDMV leader (Maize Dwarf Mosaic Virus) (Virology 154:9-20), and human immunoglobulin heavy-chain binding protein (BiP) (Macejak et al. (1991) Nature 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4) (Jobling et al. (1987) Nature 325:622-625); tobacco mosaic virus leader (TMV) (Gallie et al. (1989) in Molecular Biology of RNA, ed. Cech (Liss, New York), pp. 237-256); and maize chlorotic mottle virus leader (MCMV) (Lommel et al. (1991) Virology 81 :382-385). See also, Della-Cioppa et al. (1987) Plant Physiol. 84:965-968.

Other methods known to enhance translation also can be utilized, for example, introns, and the like. In preparing an expression vector, the various nucleic acid fragments may be manipulated, so as to provide for the nucleic acid sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be employed to join the nucleic acid fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous nucleic acid, removal of restriction sites, or the like. For this purpose, in vitro mutagenesis, primer repair, restriction, annealing, resubstitutions, e.g., transitions and transversions, may be involved.

- 10 A number of promoters can be used in the practice of the invention. The promoters can be selected based on the desired outcome. The nucleic acids can be combined with constitutive, tissue-preferred, or other promoters for expression in plants.

- 15 Constitutive promoters include, for example, the core promoter of the Rsyn7 promoter and other constitutive promoters disclosed in WO 99/43838 and U.S. Patent No. 6,072,050; the core CaMV 35S promoter (Odell et al. (1985) *Nature* 313:810-812); rice actin (McElroy et al. (1990) *Plant Cell* 2: 163-171); ubiquitin (Christensen et al. (1989) *Plant Mol. Biol.* 12:619-632 and Christensen et al. (1992) *Plant Mol. Biol.* 18:675-689); pEMU (Last et al. (1991) *Theor. Appl. Genet.* 81:581- 588); MAS (Velten et al. (1984) *EMBO J.* 3:2723- 2730); ALS promoter (U.S. Patent No. 5,659,026), and the like. Other constitutive promoters include, for example, U.S. Patent Nos. 5,608,149; 5,608, 144; 5,604,121 ; 5,569,597; 5,466,785; 5,399,680; 5,268,463; 5,608,142; and 6,177,611.

- 25 Tissue-preferred promoters can be utilized to target enhanced expression within a particular plant tissue. Such tissue-preferred promoters include, but are not limited to, leaf- preferred promoters, root-preferred promoters, seed- preferred promoters, and stem-preferred promoters. Some examples of tissue-preferred promoters are described by, e.g., Yamamoto et al. (1997) *Plant J.* 12(2):255-265; Kawamata et al. (1997) *Plant Cell Physiol.* 38(7):792-803; Hansen et al. (1997) *Mol. Gen Genet.* 254(3):337-343; Russell et al. (1997) *Transgenic Res.* 6(2): 157-168; Rinehart et al. (1996) *Plant Physiol.* 1 12(3): 1331-1341; Van Camp et al. (1996) *Plant Physiol.* 112(2):525-535; Canevascini et al. (1996) *Plant Physiol.* 1 12(2):513- 524; Yamamoto et al. (1994) *Plant Cell Physiol.* 35(5):773-778; Lam (1994) *Results Probl. Cell Differ.* 20:181- 196; Orozco ef al. (1993) *Plant Mol Biol.* 23(6): 1 129-1138; Matsuoka et al. (1993) *Voc Natl. Acad. ScL USA* 90(20):9586-9590; and 35 Guevara-Garcia et al. (1993) *Plant J* 4(3):495-505. Promoters can be modified, if necessary, for weak expression.

- 40 In some embodiments, the nucleic acids of interest can be targeted to the chloroplast for expression. In this manner, where the nucleic acid of interest is not directly inserted into the chloroplast, the expression vector will additionally contain a chloroplast- targeting sequence comprising a nucleotide sequence that encodes a chloroplast transit peptide to direct the gene product of interest to the chloroplasts. Such transit peptides are known in the art. With

respect to chloroplast-targeting sequences, "operably linked" means that the nucleic acid sequence encoding a transit peptide (i.e., the chloroplast-targeting sequence) is linked to the desired coding sequence of the invention such that the two sequences are contiguous and in the same reading frame. See, for example, Von Heijne et al. (1991)

5 Plant Mol. Biol. Rep. 9: 104-126; Clark et al. (1989) J Biol. Chem. 264:17544-17550; Della-Cioppa et al. (1987) Plant Physiol. 84:965-968; Romer et al. (1993) Biochem. Biophys. Res. Commun. 196: 1414-1421 ; and Shah et al. (1986) Science 233:478-481. For example, a chloroplast transit peptide known in the art can be fused to the amino acid sequence of a CESA polypeptide of the invention by operably linking a chloroplast-targeting sequence to the 5'-end of a nucleotide sequence encoding the CESA
10 polypeptide.

Chloroplast targeting sequences are known in the art and include the chloroplast small subunit of ribulose-1,5-bisphosphate carboxylase (Rubisco) (de Castro Silva Filho et al.
15 (1996) Plant Mol. Biol. 30:769-780; Schnell et al. (1991) J Biol. Chem. 266(5):3335-3342); EPSPS (Archer et al. (1990) J Bioenerg. Biomemb. 22(6):789-810); tryptophan synthase (Zhao et al. (1995) J Biol. Chem. 270(11):6081-6087); plastocyanin (Lawrence et al. (1997) J Biol. Chem. 272(33):20357-20363); chorismate synthase (Schmidt et al. (1993) J Biol. Chem. 268(36):27447-27457); and the light harvesting chlorophyll a/b
20 binding protein (LHBP) (Lamppa et al. (1988) J Biol. Chem. 263: 14996-14999). See also Von Heijne et al. (1991) Plant Mol. Biol. Rep. 9: 104- 126; Clark et al. (1989) J Biol. Chem. 264: 17544-17550; Della-Cioppa et al. (1987) Plant Physiol. 84:965-968; Romer et al. (1993) Biochem Biophys. Res. Commun. 196: 1414-1421; and Shah et al. (1986) Science 233:478-481.

25 Methods for transformation of chloroplasts are known in the art. See, for example, Svab et al. (1990) Proc. Natl. Acad. Sci. USA 87:8526-8530; Svab and Maliga (1993) Proc. Natl. Acad. Sci. USA 90:913-917; Svab and Maliga (1993) EMBO J. 12:601-606. The method relies on particle gun delivery of DNA containing a selectable marker and
30 targeting of the DNA to the plastid genome through homologous recombination. Additionally, plastid transformation can be accomplished by transactivation of a silent plastid-borne transgene by tissue-preferred expression of a nuclear-encoded and plastid-directed RNA polymerase. Such a system has been reported in McBride et al. (1994) Proc. Natl. Acad. Sci. USA 91:7301- 7305.

35 The nucleic acids of interest to be targeted to the chloroplast may be optimized for expression in the chloroplast to account for differences in codon usage between the plant nucleus and this organelle. In this manner, the nucleic acids of interest may be synthesized using chloroplast-preferred codons. See, for example, U.S. Patent No. 5,380,831.

40 Numerous plant transformation vectors and methods for transforming plants are available. See, for example, An, G. et al. (1986) Plant Physiol., 81 :301-305; Fry, J., et al. (1987) Plant Cell Rep. 6:321-325; Block, M. (1988) Theor. Appl. Genet. 16: 161 -1 1 A; Hinchey, et al.

(1990) Stadler. Genet. Symp. 203:203-212; Cousins, et al. (1991) Aust. J. Plant Physiol. 18:481-494; Chee, P. P. and Slightom, J. L. (1992) Gene. 118:255-260; Christou, et al. (1992) Trends. Biotechnol. 10:239-246; Halluin, et al. (1992) Bio/Technol. 10:309-314; Dhir, et al. (1992) Plant Physiol. 99:81-88; Casas et al. (1993) Proc. Nat. Acad. Sci. USA 90: 1212-1216; Christou, P. (1993) In Vitro Cell. Dev. Biol.-Plant; 29P:119-124; Davies, et al. (1993) Plant Cell Rep. 12: 180-183; Dong, J. A. and McHughen, A. (1993) Plant Sci. 91: 139-148; Franklin, C. I. and Trieu, T. N. (1993) Plant. Physiol. 102: 167; Golovkin, et al. (1993) Plant Sci. 90:41-52; Guo Chin Sci Bull. 38:2072-2078; Asano, et al. (1994) Plant Cell Rep. 13; Ayeres N. M. and Park, W. D. (1994) Crit. Rev. Plant. Sci. 13:219-239; Barcelo, et al. (1994) Plant. J. 5:583-592; Becker, et al. (1994) Plant. J. 5:299-307; Borkowska et al. (1994) Acta. Physiol. Plant. 16:225-230; Christou, P. (1994) Agro. Food. Ind. Hi Tech. 5: 17-27; Eapen et al. (1994) Plant Cell Rep. 13:582-586; Hartman, et al. (1994) Bio-Technology 12: 919-923; Ritala, et al. (1994) Plant. Mol. Biol. 24:317-325; and Wan, Y. C. and Lemaux, P. G. (1994) Plant Physiol. 104:3748.

15

In some embodiments, the methods of the invention involve introducing a polynucleotide construct into a plant. By "introducing" is intended presenting to the plant the polynucleotide construct in such a manner that the construct gains access to the interior of a cell of the plant. The methods of the invention do not depend on a particular method for introducing a polynucleotide construct to a plant, only that the polynucleotide construct gains access to the interior of at least one cell of the plant. Methods for introducing polynucleotide constructs into plants are known in the art including, but not limited to, stable transformation methods, transient transformation methods, and virus-mediated methods. The term "introduction" or "transformation" as referred to herein further means the transfer of an exogenous polynucleotide into a host cell, irrespective of the method used for transfer. Plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a genetic construct of the present invention and a whole plant regenerated there from. The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (e.g., apical meristem, axillary buds, and root meristems), and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem). The polynucleotide may be transiently or stably introduced into a host cell and may be maintained non-integrated, for example, as a plasmid. Alternatively, it may be integrated into the host genome. The resulting transformed plant cell may then be used to regenerate a transformed plant in a manner known to persons skilled in the art.

By "stable transformation" is intended that the polynucleotide construct introduced into a plant integrates into the genome of the plant and is capable of being inherited by descendent thereof. By "transient transformation" is intended that a polynucleotide construct introduced into a plant does not integrate into the genome of the plant.

For the transformation of plants and plant cells, the nucleotide sequences of the invention are inserted using standard techniques into any vector known in the art that is suitable for expression of the nucleotide sequences in a plant or plant cell. The selection of the vector depends on the preferred transformation technique and the target plant species to be transformed. In an embodiment of the invention, the encoding nucleotide sequence is operably linked to a plant promoter, e.g. a promoter known in the art for high-level expression in a plant cell, and this construct is then introduced into a plant cell that is susceptible to CESA-inhibiting herbicides; and a transformed plant is regenerated. In some embodiments, the transformed plant is tolerant to exposure to a level of CESA-inhibiting herbicides that would kill or significantly injure a plant regenerated from an untransformed cell. This method can be applied to any plant species or crops.

Methodologies for constructing plant expression vectors and introducing foreign nucleic acids into plants are generally known in the art. For example, foreign DNA can be introduced into plants, using tumor-inducing (Ti) plasmid vectors. Other methods utilized for foreign DNA delivery involve the use of PEG mediated protoplast transformation, electroporation, microinjection whiskers, and biolistics or microprojectile bombardment for direct DNA uptake. Such methods are known in the art. (U.S. Pat. No. 5,405,765 to Vasil et al.; Bilang et al. (1991) *Gene* 100: 247-250; Scheid et al., (1991) *Mol. Gen. Genet.*, 228: 104-112; Guerche et al., (1987) *Plant Science* 52: 1-16; Neuhaus et al., (1987) *Theor. Appl. Genet.* 75: 30-36; Klein et al., (1987) *Nature* 327: 70-73; Howell et al., (1980) *Science* 208: 1265; Horsch et al., (1985) *Science* 227: 1229-1231; DeBlock et al., (1989) *Plant Physiology* 91: 694-701; *Methods for Plant Molecular Biology* (Weissbach and Weissbach, eds.) Academic Press, Inc. (1988) and *Methods in Plant Molecular Biology* (Schuler and Zielinski, eds.) Academic Press, Inc. (1989).

Other suitable methods of introducing nucleotide sequences into plant cells include microinjection as described by e.g., Crossway et al. (1986) *Biotechniques* 4:320-334, electroporation as described by e.g., Riggs et al. (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606, *Agrobacterium*-mediated transformation as described by e.g., Townsend et al., U.S. Patent No. 5,563,055, Zhao et al., U.S. Patent No. 5,981,840, direct gene transfer as described by, e.g., Paszkowski et al. (1984) *EMBO J.* 3:2717-2722, and ballistic particle acceleration as described by, e.g., U.S. Patent Nos. 4,945,050; 5,879,918; 5,886,244; and 5,932,782; Tomes et al. (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); McCabe et al. (1988) *Biotechnology* 6:923-926; and *Led transformation* (WO 00/28058). Also see, Weissinger et al., (1988) *Ann. Rev. Genet.* 22:421-477; Sanford et al., (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou et al., (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe et al., (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.* 27P: 175-182 (soybean); Singh et al., (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta et al., (1990) *Biotechnology* 8:736-740 (rice); Klein et al.,

- (1988) PNAS, 85:4305-4309 (maize); Klein et al., (1988) Biotechnology 6:559-563 (maize); U.S. Patent Nos. 5,240,855; 5,322,783; and 5,324,646; Tomes et al., (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in Plant Cell, Tissue, and Organ Culture: Fundamental Methods, ed. Gamborg (Springer- Verlag, Berlin) (maize); Klein et al., (1988) Plant Physiol. 91 :440-444 (maize); Fromm et al., (1990) Biotechnology 8:833-839 (maize); Hooykaas-Van Slogteren et al., (1984) Nature (London) 311 :763-764; Bowen et al, U.S. Patent No. 5,736,369 (cereals); Bytebier et al, (1987) PNAS 84:5345- 5349 (Liliaceae); De Wet et al., (1985) in The Experimental Manipulation of Ovule Tissues, ed. Chapman et al, (Longman, New York), pp. 197-209 (pollen); Kaeppler et al., (1990) Plant Cell Reports 9:415-418 and Kaeppler et al., (1992) Theor. Apph Genet. 84:560-566 (whisker-mediated transformation); D'Halluin et al., (1992) Plant Cell 4: 1495-1505 (electroporation); Li et al., (1993) Plant Cell Reports 12:250- 255 and Christou and Ford (1995) Annals of Botany 75:407-413 (rice); Osjoda et al, (1996) Nature Biotechnology 14:745-750 (maize via *Agrobacterium tumefaciens*).
- Transgenic plants, including transgenic crop plants, are preferably produced via *Agrobacterium*-mediated transformation. An advantageous transformation method is the transformation in planta. To this end, it is possible, for example, to allow the *agrobacteria* to act on plant seeds or to inoculate the plant meristem with *agrobacteria*. It has proved particularly expedient in accordance with the invention to allow a suspension of transformed *agrobacteria* to act on the intact plant or at least on the flower primordia. The plant is subsequently grown on until the seeds of the treated plant are obtained (Clough and Bent, Plant J. (1998) 16, 735-743). Methods for *Agrobacterium*-mediated transformation of rice include well known methods for rice transformation, such as those described in any of the following: European patent application EP 1198985 A1, Aldemita and Hodges (Planta 199: 612-617, 1996); Chan et al. (Plant Mol Biol 22 (3): 491-506, 1993), Hiei et al. (Plant J 6 (2): 271-282, 1994). In the case of corn transformation, the preferred method is as described in either Ishida et al. (Nat. Biotechnol 14(6): 745-50, 1996) or Frame et al. (Plant Physiol 129(1): 13-22, 2002). Said methods are further described by way of example in B. Jenes et al., Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, eds. S.D. Kung and R. Wu, Academic Press (1993) 128-143 and in Potrykus Annu. Rev. Plant Physiol. Plant Molec. Biol. 42 (1991) 205-225). The nucleic acids or the construct to be expressed is preferably cloned into a vector, which is suitable for transforming *Agrobacterium tumefaciens*, for example pBin19 (Bevan et al., Nucl. Acids Res. 12 (1984) 8711). *Agrobacteria* transformed by such a vector can then be used in known manner for the transformation of plants, such as plants used as a model, like *Arabidopsis* (*Arabidopsis thaliana* is within the scope of the present invention not considered as a crop plant), or crop plants such as, by way of example, tobacco plants, for example by immersing bruised leaves or chopped leaves in an *agrobacterial* solution and then culturing them in suitable

media. The transformation of plants by means of *Agrobacterium tumefaciens* is described, for example, by Höfgen and Willmitzer in Nucl. Acid Res. (1988) 16, 9877 or is known inter alia from F.F. White, Vectors for Gene Transfer in Higher Plants; in Transgenic Plants, Vol. 1, Engineering and Utilization, eds. S.D. Kung and R. Wu, Academic Press, 1993, pp. 15-38.

One transformation method known to those of skill in the art is the dipping of a flowering plant into an *Agrobacteria* solution, wherein the *Agrobacteria* contains the CESA nucleic acid, followed by breeding of the transformed gametes. *Agrobacterium* mediated plant transformation can be performed using for example the GV3101(pMP90) (Koncz and Schell, 1986, Mol. Gen. Genet. 204:383-396) or LBA4404 (Clontech) *Agrobacterium tumefaciens* strain. Transformation can be performed by standard transformation and regeneration techniques (Deblaere et al., 1994, Nucl. Acids. Res. 13:4777-4788; Gelvin, Stanton B. and Schilperoort, Robert A, Plant Molecular Biology Manual, 2nd Ed. - Dordrecht: Kluwer Academic Publ., 1995. - in Sect., Ringbuc Zentrale Signatur: BT11-P ISBN 0-7923-2731-4; Glick, Bernard R. and Thompson, John E., Methods in Plant Molecular Biology and Biotechnology, Boca Raton : CRC Press, 1993 360 S., ISBN 0-8493-5164-2). For example, rapeseed can be transformed via cotyledon or hypocotyl transformation (Moloney et al., 1989, Plant Cell Report 8:238-242; De Block et al., 1989, Plant Physiol. 91:694-701). Use of antibiotics for *Agrobacterium* and plant selection depends on the binary vector and the *Agrobacterium* strain used for transformation. Rapeseed selection is normally performed using kanamycin as selectable plant marker. *Agrobacterium* mediated gene transfer to flax can be performed using, for example, a technique described by Mlynarova et al., 1994, Plant Cell Report 13:282-285. Additionally, transformation of soybean can be performed using for example a technique described in European Patent No. 0424 047, U.S. Patent No. 5,322,783, European Patent No. 0397 687, U.S. Patent No. 5,376,543, or U.S. Patent No. 5,169,770. Transformation of maize can be achieved by particle bombardment, polyethylene glycol mediated DNA uptake, or via the silicon carbide fiber technique. (See, for example, Freeling and Walbot "The maize handbook" Springer Verlag: New York (1993) ISBN 3-540-97826-7). A specific example of maize transformation is found in U.S. Patent No. 5,990,387, and a specific example of wheat transformation can be found in PCT Application No. WO 93/07256.

In some embodiments, polynucleotides of the present invention may be introduced into plants by contacting plants with a virus or viral nucleic acids. Generally, such methods involve incorporating a polynucleotide construct of the invention within a viral DNA or RNA molecule. It is recognized that the polypeptides of the invention may be initially synthesized as part of a viral polyprotein, which later may be processed by proteolysis in vivo or in vitro to produce the desired recombinant polypeptide. Further, it is recognized that promoters of the invention also encompass promoters utilized for transcription by viral RNA polymerases. Methods for introducing polynucleotide constructs into plants and expressing a protein encoded therein, involving viral DNA or RNA molecules, are known in the art. See, for

example, U.S. Patent Nos. 5,889,191, 5,889,190, 5,866,785, 5,589,367 and 5,316,931. The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick et al. (1986) Plant Cell Reports 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure expression of the desired phenotypic characteristic has been achieved.

10

The present invention may be used for transformation of any plant species, including, but not limited to, monocots and dicots. Examples of plant species of interest include, but are not limited to, corn or maize (*Zea mays*), *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa

15

(*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet e.g., pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria italica*), finger millet (*Eleusine coracana*), sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), wheat (*Triticum aestivum*, *T.*

20

Turgidum ssp. *durum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Coffea* spp.), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees (*Citrus* spp.), cocoa (*Theobroma cacao*), tea (*Camellia sinensis*), banana (*Musa* spp.), avocado (*Persea americana*), fig (*Ficus casica*), guava (*Psidium guajava*), mango

25

(*Mangifera indica*), olive (*Olea europaea*), papaya (*Carica papaya*), cashew (*Anacardium occidentale*), macadamia (*Macadamia integrifolia*), almond (*Prunus amygdalus*), sugar beets (*Beta vulgaris*), sugarcane (*Saccharum* spp.), oats, barley, vegetables, ornamentals, and conifers. Preferably, plants of the present invention are crop plants (for example, sunflower, *Brassica* sp., cotton, sugar, beet, soybean, peanut, alfalfa, safflower, tobacco, corn, rice, wheat, rye, barley triticales, sorghum, millet, etc.).

30

In addition to the transformation of somatic cells, which then have to be regenerated into intact plants, it is also possible to transform the cells of plant meristems and in particular those cells which develop into gametes. In this case, the transformed gametes follow the natural plant development, giving rise to transgenic plants. Thus, for example, seeds of *Arabidopsis* are treated with agrobacteria and seeds are obtained from the developing plants of which a certain proportion is transformed and thus transgenic [Feldman, KA and Marks MD (1987). Mol Gen Genet 208:274-289; Feldmann K (1992). In: C Koncz, N-H Chua and J Shell, Eds, Methods in *Arabidopsis* Research. Word Scientific, Singapore, pp. 274-289]. Alternative methods are based on the repeated removal of the inflorescences and incubation of the excision site in the center of the rosette with transformed agrobacteria, whereby transformed seeds can likewise be obtained at a later point in time (Chang (1994).

40

Plant J. 5: 551-558; Katavic (1994). Mol Gen Genet, 245: 363-370). However, an especially effective method is the vacuum infiltration method with its modifications such as the "floral dip" method. In the case of vacuum infiltration of Arabidopsis, intact plants under reduced pressure are treated with an agrobacterial suspension [Bechthold, N (1993). C R Acad Sci Paris Life Sci, 316: 1194-1199], while in the case of the "floral dip" method the developing floral tissue is incubated briefly with a surfactant-treated agrobacterial suspension [Clough, SJ and Bent AF (1998) The Plant J. 16, 735-743]. A certain proportion of transgenic seeds are harvested in both cases, and these seeds can be distinguished from non-transgenic seeds by growing under the above-described selective conditions. In addition the stable transformation of plastids is of advantages because plastids are inherited maternally in most crops reducing or eliminating the risk of transgene flow through pollen. The transformation of the chloroplast genome is generally achieved by a process which has been schematically displayed in Klaus et al., 2004 [Nature Biotechnology 22 (2), 225-229]. Briefly the sequences to be transformed are cloned together with a selectable marker gene between flanking sequences homologous to the chloroplast genome. These homologous flanking sequences direct site specific integration into the plastome. Plastid transformation has been described for many different plant species and an overview is given in Bock (2001) Transgenic plastids in basic research and plant biotechnology. J Mol Biol. 2001 Sep 21; 312 (3):425-38 or Maliga, P (2003) Progress towards commercialization of plastid transformation technology. Trends Biotechnol. 21, 20-28. Further biotechnological progress has recently been reported in form of marker free plastid transformants, which can be produced by a transient co-integrated marker gene (Klaus et al., 2004, Nature Biotechnology 22(2), 225-229). The genetically modified plant cells can be regenerated via all methods with which the skilled worker is familiar. Suitable methods can be found in the abovementioned publications by S.D. Kung and R. Wu, Potrykus or Höfgen and Willmitzer.

Generally after transformation, plant cells or cell groupings are selected for the presence of one or more markers which are encoded by plant-expressible genes co-transferred with the gene of interest, following which the transformed material is regenerated into a whole plant. To select transformed plants, the plant material obtained in the transformation is, as a rule, subjected to selective conditions so that transformed plants can be distinguished from untransformed plants. For example, the seeds obtained in the above-described manner can be planted and, after an initial growing period, subjected to a suitable selection by spraying. A further possibility consists in growing the seeds, if appropriate after sterilization, on agar plates using a suitable selection agent so that only the transformed seeds can grow into plants. Alternatively, the transformed plants are screened for the presence of a selectable marker such as the ones described above.

Following DNA transfer and regeneration, putatively transformed plants may also be evaluated, for instance using Southern analysis, for the presence of the gene of interest, copy number and/or genomic organisation. Alternatively or additionally, expression levels of the newly introduced DNA may be monitored using Northern and/or Western analysis, both

techniques being well known to persons having ordinary skill in the art.

The generated transformed plants may be propagated by a variety of means, such as by clonal propagation or classical breeding techniques. For example, a first generation (or T1) transformed plant may be selfed and homozygous second-generation (or T2) transformants selected, and the T2 plants may then further be propagated through classical breeding techniques. The generated transformed organisms may take a variety of forms. For example, they may be chimeras of transformed cells and non-transformed cells; clonal transformants (e.g., all cells transformed to contain the expression cassette); grafts of transformed and untransformed tissues (e.g., in plants, a transformed rootstock grafted to an untransformed scion).

Preferably, the expression of the nucleic acid in the plant results in the plant's increased tolerance to CESA-inhibiting herbicide as compared to a wild type variety of the plant.

In another embodiment, the invention refers to a plant, comprising a plant cell according to the present invention, wherein expression of the nucleic acid in the plant results in the plant's increased resistance to CESA-inhibiting herbicide as compared to a wild type variety of the plant.

The plants described herein can be either transgenic crop plants or non-transgenic plants.

In addition to the general definition, give SUPRA, "transgenic", "transgene" or "recombinant" means with regard to, for example, a nucleic acid sequence, an expression cassette, gene construct or a vector comprising the nucleic acid sequence or an organism transformed with the nucleic acid sequences, expression cassettes or vectors according to the invention, all those constructions brought about by recombinant methods in which either

- (a) the nucleic acid sequences encoding proteins useful in the methods of the invention, or
- (b) genetic control sequence(s) which is operably linked with the nucleic acid sequence according to the invention, for example a promoter, or
- (c) a) and b)

are not located in their natural genetic environment or have been modified by recombinant methods, it being possible for the modification to take the form of, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues in order to allow for the expression of the wildtype or mutated CESA of the present invention. The natural genetic environment is understood as meaning the natural genomic or chromosomal locus in the original plant or the presence in a genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least in part. The environment flanks the nucleic acid sequence at least on one side and has a sequence length of at least 50 bp, preferably at least 500 bp, especially preferably at least 1000 bp, most preferably at least 5000 bp. A naturally

occurring expression cassette – for example the naturally occurring combination of the natural promoter of the nucleic acid sequences with the corresponding nucleic acid sequence encoding a polypeptide useful in the methods of the present invention, as defined above – becomes a transgenic expression cassette when this expression cassette is
5 modified by non-natural, synthetic ("artificial") methods such as, for example, mutagenic treatment. Suitable methods are described, for example, in US 5,565,350 or WO 00/15815.

A transgenic plant for the purposes of the invention is thus understood as meaning, as above, that the nucleic acids of the invention are not at their natural locus in the genome of
10 said plant, it being possible for the nucleic acids to be expressed homologously or heterologously. However, as mentioned, transgenic also means that, while the nucleic acids according to the invention or used in the inventive method are at their natural position in the genome of a plant, the sequence has been modified with regard to the natural sequence, and/or that the regulatory sequences of the natural sequences have been modified.

15 Transgenic is preferably understood as meaning the expression of the nucleic acids according to the invention at an unnatural locus in the genome, i.e. homologous or, preferably, heterologous expression of the nucleic acids takes place. Preferred transgenic plants are mentioned herein. Furthermore, the term "transgenic" refers to any plant, plant cell, callus, plant tissue, or plant part, that contains all or part of at least one recombinant
20 polynucleotide. In many cases, all or part of the recombinant polynucleotide is stably integrated into a chromosome or stable extra-chromosomal element, so that it is passed on to successive generations. For the purposes of the invention, the term "recombinant polynucleotide" refers to a polynucleotide that has been altered, rearranged, or modified by genetic engineering. Examples include any cloned polynucleotide, or polynucleotides, that
25 are linked or joined to heterologous sequences. The term "recombinant" does not refer to alterations of polynucleotides that result from naturally occurring events, such as spontaneous mutations, or from non-spontaneous mutagenesis followed by selective breeding.

30 Plants containing mutations arising due to non-spontaneous mutagenesis and selective breeding are referred to herein as non-transgenic plants and are included in the present invention. In embodiments wherein the plant is transgenic and comprises multiple wildtype or mutated CESA nucleic acids, the nucleic acids can be derived from different genomes or from the same genome. Alternatively, in embodiments wherein the plant is non-transgenic
35 and comprises multiple wildtype or mutated CESA nucleic acids, the nucleic acids are located on different genomes or on the same genome.

In certain embodiments, the present invention involves herbicide-resistant plants that are produced by mutation breeding. Such plants comprise a polynucleotide encoding a wildtype
40 or mutated CESA and are tolerant to one or more CESA-inhibiting herbicides. Such methods can involve, for example, exposing the plants or seeds to a mutagen, particularly a chemical mutagen such as, for example, ethyl methanesulfonate (EMS) and selecting for

plants that have enhanced tolerance to at least one or more CESA-inhibiting herbicide [see Example 1].

However, the present invention is not limited to herbicide-tolerant plants that are
 5 produced by a mutagenesis method involving the chemical mutagen EMS. Any
 mutagenesis method known in the art may be used to produce the herbicide-resistant
 plants of the present invention. Such mutagenesis methods can involve, for example,
 the use of any one or more of the following mutagens: radiation, such as X-rays,
 Gamma rays (e.g., cobalt 60 or cesium 137), neutrons, (e.g., product of nuclear fission
 10 by uranium 235 in an atomic reactor), Beta radiation (e.g., emitted from radioisotopes
 such as phosphorus 32 or carbon 14), and ultraviolet radiation (preferably from 250 to
 290 nm), and chemical mutagens such as base analogues (e.g., 5-bromo-uracil), related
 compounds (e.g., 8-ethoxy caffeine), antibiotics (e.g., streptonigrin), alkylating agents
 (e.g., sulfur mustards, nitrogen mustards, epoxides, ethylenamines, sulfates, sulfonates,
 15 sulfones, lactones), azide, hydroxylamine, nitrous acid, or acridines. Herbicide-resistant
 plants can also be produced by using tissue culture methods to select for plant cells
 comprising herbicide-resistance mutations and then regenerating herbicide-resistant
 plants therefrom. See, for example, U.S. Patent Nos. 5,773,702 and 5,859,348. Further
 details of mutation breeding can be found in "Principals of Cultivar Development" Fehr,
 20 1993 Macmillan Publishing Company.

The plant of the present invention comprises at least one wildtype or mutated CESA
 nucleic acid or over-expressed wild-type CESA nucleic acid, and has increased
 tolerance to a CESA-inhibiting herbicide as compared to a wild-type variety of the plant.
 25 It is possible for the plants of the present invention to have multiple wildtype or
 mutated CESA nucleic acids from different genomes since these plants can contain
 more than one genome. For example, a plant contains two genomes, usually referred to
 as the A and B genomes. Because CESA is a required metabolic enzyme, it is assumed
 that each genome has at least one gene coding for the CESA enzyme (i.e. at least one
 30 CESA gene). As used herein, the term "CESA gene locus" refers to the position of a
 CESA gene on a genome, and the terms "CESA gene" and "CESA nucleic acid" refer to a
 nucleic acid encoding the CESA enzyme. The CESA nucleic acid on each genome differs
 in its nucleotide sequence from a CESA nucleic acid on another genome. One of skill in
 the art can determine the genome of origin of each CESA nucleic acid through genetic
 35 crossing and/or either sequencing methods or exonuclease digestion methods known to
 those of skill in the art.

The present invention includes plants comprising one, two, three, or more wildtype or
 mutated CESA alleles, wherein the plant has increased tolerance to a CESA-inhibiting
 40 herbicide as compared to a wild-type variety of the plant. The wildtype or mutated
 CESA alleles can comprise a nucleotide sequence selected from the group consisting of
 a polynucleotide as defined in SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or

77, or a variant or derivative thereof, a polynucleotide encoding a polypeptide as defined in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, or 64, or a variant or derivative,
 5 homologue, orthologue, paralogue thereof, a polynucleotide comprising at least 60 consecutive nucleotides of any of the aforementioned polynucleotides; and a polynucleotide complementary to any of the aforementioned polynucleotides.

10 “Alleles” or “allelic variants” are alternative forms of a given gene, located at the same chromosomal position. Allelic variants encompass Single Nucleotide Polymorphisms (SNPs), as well as Small Insertion/Deletion Polymorphisms (INDELs). The size of INDELs is usually less than 100 bp. SNPs and INDELs form the largest set of sequence variants in naturally occurring polymorphic strains of most organisms

15 The term “variety” refers to a group of plants within a species defined by the sharing of a common set of characteristics or traits accepted by those skilled in the art as sufficient to distinguish one cultivar or variety from another cultivar or variety. There is no implication in either term that all plants of any given cultivar or variety will be genetically identical at either the whole gene or molecular level or that any given plant will be homozygous at all loci. A
 20 cultivar or variety is considered “true breeding” for a particular trait if, when the true-breeding cultivar or variety is self-pollinated, all of the progeny contain the trait. The terms “breeding line” or “line” refer to a group of plants within a cultivar defined by the sharing of a common set of characteristics or traits accepted by those skilled in the art as sufficient to distinguish one breeding line or line from another breeding line or line. There is no
 25 implication in either term that all plants of any given breeding line or line will be genetically identical at either the whole gene or molecular level or that any given plant will be homozygous at all loci. A breeding line or line is considered “true breeding” for a particular trait if, when the true-breeding line or breeding line is self-pollinated, all of the progeny contain the trait. In the present invention, the trait arises from a mutation in a CESA gene of
 30 the plant or seed.

The herbicide-resistant plants of the invention that comprise polynucleotides encoding wildtype or mutated CESA polypeptides also find use in methods for increasing the herbicide-resistance of a plant through conventional plant breeding involving sexual
 35 reproduction. The methods comprise crossing a first plant that is a herbicide-resistant plant of the invention to a second plant that may or may not be resistant to the same herbicide or herbicides as the first plant or may be resistant to different herbicide or herbicides than the first plant. The second plant can be any plant that is capable of producing viable progeny plants (i.e., seeds) when crossed with the first plant. Typically, but not necessarily, the first
 40 and second plants are of the same species. The methods can optionally involve selecting for progeny plants that comprise the wildtype or mutated CESA polypeptides of the first plant and the herbicide resistance characteristics of the second plant. The progeny plants

produced by this method of the present invention have increased resistance to a herbicide when compared to either the first or second plant or both. When the first and second plants are resistant to different herbicides, the progeny plants will have the combined herbicide tolerance characteristics of the first and second plants. The methods of the invention can

5 further involve one or more generations of backcrossing the progeny plants of the first cross to a plant of the same line or genotype as either the first or second plant. Alternatively, the progeny of the first cross or any subsequent cross can be crossed to a third plant that is of a different line or genotype than either the first or second plant. The present invention also provides plants, plant organs, plant tissues, plant cells, seeds, and non-human host cells

10 that are transformed with the at least one polynucleotide molecule, expression cassette, or transformation vector of the invention. Such transformed plants, plant organs, plant tissues, plant cells, seeds, and non-human host cells have enhanced tolerance or resistance to at least one herbicide, at levels of the herbicide that kill or inhibit the growth of an untransformed plant, plant tissue, plant cell, or non-human host cell, respectively.

15 Preferably, the transformed plants, plant tissues, plant cells, and seeds of the invention are *Arabidopsis thaliana* and crop plants.

It is to be understood that the plant of the present invention can comprise a wild type CESA nucleic acid in addition to a mutated CESA nucleic acid. It is contemplated that the CESA-

20 inhibiting herbicide tolerant lines may contain a mutation in only one of multiple CESA isoenzymes. Therefore, the present invention includes a plant comprising one or more mutated CESA nucleic acids in addition to one or more wild type CESA nucleic acids.

In another embodiment, the invention refers to a seed produced by a transgenic plant

25 comprising a plant cell of the present invention, wherein the seed is true breeding for an increased resistance to a CESA-inhibiting herbicide as compared to a wild type variety of the seed.

In other aspects, CESA-inhibiting herbicides-tolerant plants of the present invention can be

30 employed as CESA-inhibiting herbicides-tolerance trait donor lines for development, as by traditional plant breeding, to produce other varietal and/or hybrid crops containing such trait or traits. All such resulting variety or hybrids crops, containing the ancestral CESA-inhibiting herbicides-tolerance trait or traits can be referred to herein as progeny or descendant of the ancestral, CESA-inhibiting herbicides-tolerant line(s).

35 In other embodiments, the present invention provides a method for producing a CESA-inhibiting herbicides-tolerant plant. The method comprises: crossing a first CESA-inhibiting herbicides-tolerant plant with a second plant to produce a CESA-inhibiting herbicides-tolerant progeny plant, wherein the first plant and the progeny plant comprise in at least

40 some of their cells a polynucleotide operably linked to a promoter operable in plant cells, the recombinant polynucleotide being effective in the cells of the first plant to express a wildtype or mutated CESA polypeptide encoded by the polynucleotide, the expression of

the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

5 In some embodiments, traditional plant breeding is employed whereby the CESA-inhibiting herbicides-tolerant trait is introduced in the progeny plant resulting therefrom. In one embodiment, the present invention provides a method for producing a CESA-inhibiting herbicides-tolerant progeny plant, the method comprising: crossing a parent plant with a CESA-inhibiting herbicides-tolerant plant to introduce the CESA-inhibiting herbicides-tolerance characteristics of the CESA-inhibiting herbicides-tolerant plant into the germplasm
10 of the progeny plant, wherein the progeny plant has increased tolerance to the CESA-inhibiting herbicides relative to the parent plant. In other embodiments, the method further comprises the step of introgressing the CESA-inhibiting herbicides-tolerance characteristics through traditional plant breeding techniques to obtain a descendent plant having the CESA-inhibiting herbicides-tolerance characteristics.

15 In other aspects, plants of the invention include those plants which, in addition to being CESA-inhibiting herbicides-tolerant, have been subjected to further genetic modifications by breeding, mutagenesis or genetic engineering, e.g. have been rendered tolerant to applications of specific other classes of herbicides, such as AHAS inhibitors; auxinic
20 herbicides; bleaching herbicides such as hydroxyphenylpyruvate dioxygenase (HPPD) inhibitors or phytoene desaturase (PDS) inhibitors; EPSPS inhibitors such as glyphosate; glutamine synthetase (GS) inhibitors such as glufosinate; lipid biosynthesis inhibitors such as acetyl CoA carboxylase (ACCase) inhibitors; or oxynil {i.e. bromoxynil or ioxynil} herbicides as a result of conventional methods of breeding or genetic engineering. Thus,
25 CESA-inhibiting herbicides-tolerant plants of the invention can be made resistant to multiple classes of herbicides through multiple genetic modifications, such as resistance to both glyphosate and glufosinate or to both glyphosate and a herbicide from another class such as HPPD inhibitors, AHAS inhibitors, or ACCase inhibitors. These herbicide resistance technologies are, for example, described in Pest Management Science (at volume, year,
30 page): 61, 2005, 246; 61, 2005, 258; 61, 2005, 277; 61, 2005, 269; 61, 2005, 286; 64, 2008, 326; 64, 2008, 332; Weed Science 57, 2009, 108; Australian Journal of Agricultural Research 58, 2007, 708; Science 316, 2007, 1185; and references quoted therein. For example, CESA-inhibiting herbicides-tolerant plants of the invention, in some embodiments, may be tolerant to ACCase inhibitors, such as "dims" {e.g., cycloxydim, sethoxydim,
35 clethodim, or tepraloxymid), "fops" {e.g., clodinafop, diclofop, fluazifop, haloxyfop, or quizalofop), and "dens" (such as pinoxaden); to auxinic herbicides, such as dicamba; to EPSPS inhibitors, such as glyphosate; to other CESA inhibitors; and to GS inhibitors, such as glufosinate.

40 In addition to these classes of inhibitors, CESA-inhibiting herbicides-tolerant plants of the invention may also be tolerant to herbicides having other modes of action, for example, chlorophyll/carotenoid pigment inhibitors, cell membrane disrupters, photosynthesis

inhibitors, cell division inhibitors, root inhibitors, shoot inhibitors, and combinations thereof.

Such tolerance traits may be expressed, e.g. : as mutant or wildtype HPPD proteins, as mutant AHASL proteins, mutant ACCase proteins, mutant EPSPS proteins, or mutant
 5 glutamine synthetase proteins; or as mutant native, inbred, or transgenic aryloxyalkanoate dioxygenase (AAD or DHT), haloarylnitrilase (BXN), 2,2-dichloropropionic acid dehalogenase (DEH), glyphosate-N- acetyltransferase (GAT), glyphosate decarboxylase (GDC), glyphosate oxidoreductase (GOX), glutathione-S-transferase (GST),
 10 phosphinothricin acetyltransferase (PAT or bar), or CYP450s proteins having an herbicide-degrading activity. CESA-inhibiting herbicides- tolerant plants hereof can also be stacked with other traits including, but not limited to, pesticidal traits such as Bt Cry and other proteins having pesticidal activity toward coleopteran, lepidopteran, nematode, or other pests; nutrition or nutraceutical traits such as modified oil content or oil profile traits, high protein or high amino acid concentration traits, and other trait types known in the art.

15 Furthermore, in other embodiments, CESA-inhibiting herbicides-tolerant plants are also covered which are, by the use of recombinant DNA techniques and/or by breeding and/or otherwise selected for such characteristics, rendered able to synthesize one or more insecticidal proteins, especially those known from the bacterial genus *Bacillus*, particularly
 20 from *Bacillus thuringiensis*, such as [delta]-endotoxins, e.g. CryIA(b), CryIA(c), CryIF, CryIF(a2), CryIIA(b), CryIIIA, CryIIIB(b) or Cry9c; vegetative insecticidal proteins (VIP), e.g. VIP1, VIP2, VIP3 or VIP3A; insecticidal proteins of bacteria colonizing nematodes, e.g. *Photorhabdus* spp. or *Xenorhabdus* spp.; toxins produced by animals, such as scorpion toxins, arachnid toxins, wasp toxins, or other insect-specific neurotoxins; toxins produced by
 25 fungi, such as streptomycete toxins; plant lectins, such as pea or barley lectins; agglutinins; proteinase inhibitors, such as trypsin inhibitors, serine protease inhibitors, patatin, cystatin or papain inhibitors; ribosome-inactivating proteins (RIP), such as ricin, maize-RIP, abrin, luffin, saporin or bryodin; steroid metabolism enzymes, such as 3-hydroxy-steroid oxidase, ecdysteroid-IDP-glycosyl-transferase, cholesterol oxidases, ecdysone inhibitors or HMG-
 30 CoA-reductase; ion channel blockers, such as blockers of sodium or calcium channels; juvenile hormone esterase; diuretic hormone receptors (helicokinin receptors); stilben synthase, bibenzyl synthase, chitinases or glucanases. In the context of the present invention these insecticidal proteins or toxins are to be understood expressly also as pre-toxins, hybrid proteins, truncated or otherwise modified proteins. Hybrid proteins are
 35 characterized by a new combination of protein domains, (see, e.g. WO 02/015701). Further examples of such toxins or genetically modified plants capable of synthesizing such toxins are disclosed, e.g., in EP-A 374 753, WO 93/007278, WO 95/34656, EP-A 427 529, EP-A 451 878, WO 03/18810 und WO 03/52073. The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e.g.
 40 in the publications mentioned above. These insecticidal proteins contained in the genetically modified plants impart to the plants producing these proteins tolerance to harmful pests from all taxonomic groups of arthropods, especially to beetles (Coleoptera), two-winged

insects (Diptera), and moths (Lepidoptera) and to nematodes (Nematoda).

In some embodiments, expression of one or more protein toxins (e.g., insecticidal proteins) in the CESA-inhibiting herbicides-tolerant plants is effective for controlling organisms that include, for example, members of the classes and orders: Coleoptera such as the American bean weevil *Acanthoscelides obtectus*; the leaf beetle *Agelastica alni*; click beetles (*Agriotes lineatus*, *Agriotes obscurus*, *Agriotes bicolor*); the grain beetle *Ahasverus advena*; the summer schaffer *Amphimallon solstitialis*; the furniture beetle *Anobium punctatum*; *Anthonomus* spp. (weevils); the Pygmy mangold beetle *Atomaria linearis*; carpet beetles (*Anthrenus* spp., *Attagenus* spp.); the cowpea weevil *Callosobruchus maculatus*; the fried fruit beetle *Carpophilus hemipterus*; the cabbage seedpod weevil *Ceutorhynchus assimilis*; the rape winter stem weevil *Ceutorhynchus picipitarsis*; the wireworms *Conoderus vespertinus* and *Conoderus falli*; the banana weevil *Cosmopolites sordidus*; the New Zealand grass grub *Costelytra zealandica*; the June beetle *Cotinis nitida*; the sunflower stem weevil *Cylindrocopturus adspersus*; the larder beetle *Dermestes lardarius*; the corn rootworms *Diabrotica virgifera*, *Diabrotica virgifera virgifera*, and *Diabrotica barberi*; the Mexican bean beetle *Epilachna varivestis*; the old house borer *Hylotropes bajulus*; the lucerne weevil *Hypera postica*; the shiny spider beetle *Gibbium psyllodes*; the cigarette beetle *Lasioderma serricorne*; the Colorado potato beetle *Leptinotarsa decemlineata*; *Lyctus* beetles {*Lyctus* spp. , the pollen beetle *Meligethes aeneus*; the common cockshafer *Melolontha melolontha*; the American spider beetle *Mezium americanum*; the golden spider beetle *Niptus hololeucus*; the grain beetles *Oryzaephilus surinamensis* and *Oryzaephilus Mercator*; the black vine weevil *Otiorhynchus sulcatus*; the mustard beetle *Phaedon cochleariae*, the crucifer flea beetle *Phyllotreta cruciferae*; the striped flea beetle *Phyllotreta striolata*; the cabbage steam flea beetle *Psylliodes chrysocephala*; *Ptinus* spp. (spider beetles); the lesser grain borer *Rhizopertha dominica*; the pea and bean weevil *Sitona lineatus*; the rice and granary beetles *Sitophilus oryzae* and *Sitophilus granaries*; the red sunflower seed weevil *Smicronyx fulvus*; the drugstore beetle *Stegobium paniceum*; the yellow mealworm beetle *Tenebrio molitor*, the flour beetles *Tribolium castaneum* and *Tribolium confusum*; warehouse and cabinet beetles {*Trogoderma* spp.}; the sunflower beetle *Zygogramma exclamationis*; Dermaptera (earwigs) such as the European earwig *Forficula auricularia* and the striped earwig *Labidura riparia*; Dictyoptera such as the oriental cockroach *Blatta orientalis*; the greenhouse millipede *Oxidus gracilis*; the beet fly *Pegomyia betae*; the frit fly *Oscinella frit*; fruitflies (*Dacus* spp., *Drosophila* spp.); Isoptera (termites) including species from the families Hodotermitidae, Kalotermitidae, Mastotermitidae, Rhinotermitidae, Serritermitidae, Termitidae, Termopsidae; the tarnished plant bug *Lygus lineolaris*; the black bean aphid *Aphis fabae*; the cotton or melon aphid *Aphis gossypii*; the green apple aphid *Aphis pomi*; the citrus spiny whitefly *Aleurocanthus spiniferus*; the sweet potato whitefly *Bemisia tabaci*; the cabbage aphid *Brevicoryne brassicae*; the pear psylla *Cacopsylla pyricola*; the currant aphid *Cryptomyzus ribis*; the grape phylloxera *Daktulosphaira vitifoliae*; the citrus psylla *Diaphorina citri*; the potato leafhopper *Empoasca fabae*; the bean

leafhopper *Empoasca Solana*; the vine leafhopper *Empoasca vitis*; the woolly aphid *Eriosoma lanigerum*; the European fruit scale *Eulecanium corni*; the mealy plum aphid *Hyalopterus arundinis*; the small brown planthopper *Laodelphax striatellus*; the potato aphid *Macrosiphum euphorbiae*; the green peach aphid *Myzus persicae*; the green rice

5 leafhopper *Nephotettix cincticeps*; the brown planthopper *Nilaparvata lugens*; the hop aphid *Phorodon humuli*; the bird-cherry aphid *Rhopalosiphum padi*; the grain aphid *Sitobion avenae*; Lepidoptera such as *Adoxophyes orana* (summer fruit tortrix moth); *Archips podana* (fruit tree tortrix moth); *Bucculatrix pyrivorella* (pear leafminer); *Bucculatrix thurberiella* (cotton leaf perforator); *Bupalus piniarius* (pine looper); *Carpocapsa pomonella*

10 (codling moth); *Chilo suppressalis* (striped rice borer); *Choristoneura fumiferana* (eastern spruce budworm); *Cochylis hospes* (banded sunflower moth); *Diatraea grandiosella* (southwestern corn borer); *Eupoecilia ambiguella* (European grape berry moth); *Helicoverpa armigera* (cotton bollworm); *Helicoverpa zea* (cotton bollworm); *Heliothis virescens* (tobacco budworm), *Homeosoma electellum* (sunflower moth); *Homona magnanima*

15 (oriental tea tree tortrix moth); *Lithocolletis blancardella* (spotted tentiform leafminer); *Lymantria dispar* (gypsy moth); *Malacosoma neustria* (tent caterpillar); *Mamestra brassicae* (cabbage armyworm); *Mamestra configurata* (Bertha armyworm); *Operophtera brumata* (winter moth); *Ostrinia nubilalis* (European corn borer), *Panolis flammea* (pine beauty moth), *Phyllocnistis citrella* (citrus leafminer); *Pieris brassicae* (cabbage white butterfly);

20 *Rachiplusia ni* (soybean looper); *Spodoptera exigua* (beet armyworm); *Spodoptera littoralis* (cotton leafworm); *Sylepta derogata* (cotton leaf roller); *Trichoplusia ni* (cabbage looper); Orthoptera such as the common cricket *Acheta domesticus*, tree locusts (*Anacridium* spp.), the migratory locust *Locusta migratoria*, the two-striped grasshopper *Melanoplus bivittatus*, the differential grasshopper *Melanoplus differentialis*, the redlegged grasshopper

25 *Melanoplus femurrubrum*, the migratory grasshopper *Melanoplus sanguinipes*, the northern mole cricket *Neocurtilla hexadactyla*, the red locust *Nomadacris septemfasciata*, the shortwinged mole cricket *Scapteriscus abbreviatus*, the southern mole cricket *Scapteriscus borellii*, the tawny mole cricket *Scapteriscus vicinus*, and the desert locust *Schistocerca gregaria*; Symphyla such as the garden symphylan *Scutigera immaculata*; Thysanoptera

30 such as the tobacco thrips *Frankliniella fusca*, the flower thrips *Frankliniella intonsa*, the western flower thrips *Frankliniella occidentalis*, the cotton bud thrips *Frankliniella schultzei*, the banded greenhouse thrips *Hercinothrips femoralis*, the soybean thrips *Neohydatothrips variabilis*, Kelly's citrus thrips *Pezothrips kellyanus*, the avocado thrips *Scirtothrips perseae*, the melon thrips *Thrips palmi*, and the onion thrips *Thrips tabaci*; and the like, and

35 combinations comprising one or more of the foregoing organisms.

In some embodiments, expression of one or more protein toxins (e.g., insecticidal proteins) in the CESA-inhibiting herbicides-tolerant plants is effective for controlling flea beetles, i.e. members of the flea beetle tribe of family Chrysomelidae, preferably against *Phyllotreta*

40 spp., such as *Phyllotreta cruciferae* and/or *Phyllotreta triolata*. In other embodiments, expression of one or more protein toxins (e.g., insecticidal proteins) in the CESA-inhibiting herbicides- tolerant plants is effective for controlling cabbage seedpod weevil, the Bertha

armyworm, Lygus bugs, or the diamondback moth.

Furthermore, in one embodiment, CESA-inhibiting herbicides-tolerant plants are also covered which are, e.g. by the use of recombinant DNA techniques and/or by breeding
 5 and/or otherwise selected for such traits, rendered able to synthesize one or more proteins to increase the resistance or tolerance of those plants to bacterial, viral or fungal pathogens. The methods for producing such genetically modified plants are generally known to the person skilled in the art.

10 Furthermore, in another embodiment, CESA-inhibiting herbicides-tolerant plants are also covered which are, e.g. by the use of recombinant DNA techniques and/or by breeding and/or otherwise selected for such traits, rendered able to synthesize one or more proteins to increase the productivity (e.g. oil content), tolerance to drought, salinity or other growth-limiting environmental factors or tolerance to pests and fungal, bacterial or viral pathogens
 15 of those plants.

Furthermore, in other embodiments, CESA-inhibiting herbicides-tolerant plants are also covered which are, e.g. by the use of recombinant DNA techniques and/or by breeding and/or otherwise selected for such traits, altered to contain a modified amount of one or
 20 more substances or new substances, for example, to improve human or animal nutrition, e.g. oil crops that produce health-promoting long-chain omega-3 fatty acids or unsaturated omega-9 fatty acids (e.g. Nexera(R) rape, Dow Agro Sciences, Canada).

Furthermore, in some embodiments, CESA-inhibiting herbicides-tolerant plants are also covered which are, e.g. by the use of recombinant DNA techniques and/or by breeding and/or otherwise selected for such traits, altered to contain increased amounts of vitamins and/or minerals, and/or improved profiles of nutraceutical compounds.

In one embodiment, CESA-inhibiting herbicides-tolerant plants of the present invention, relative to a wild-type plant, comprise an increased amount of, or an improved profile of, a compound selected from the group consisting of: glucosinolates (e.g., glucoraphanin (4-methylsulfinylbutyl-glucosinolate), sulforaphane, 3-indolylmethyl-glucosinolate (glucobrassicin), 1-methoxy-3-indolylmethyl-glucosinolate (neoglucobrassicin)); phenolics (e.g., flavonoids (e.g., quercetin, kaempferol),
 30 hydroxycinnamoyl derivatives (e.g., 1,2,2'-trisinapoylgentiobiose, 1,2-diferuloylgentiobiose, 1,2'-disinapoyl-2-feruloylgentiobiose, 3-O-caffeoyl-quinic (neochlorogenic acid)); and vitamins and minerals (e.g., vitamin C, vitamin E, carotene, folic acid, niacin, riboflavin, thiamine, calcium, iron, magnesium, potassium, selenium, and zinc).

40 In another embodiment, CESA-inhibiting herbicides-tolerant plants of the present invention, relative to a wild-type plant, comprise an increased amount of, or an improved profile of, a compound selected from the group consisting of: progoitrin; isothiocyanates; indoles

(products of glucosinolate hydrolysis); glutathione; carotenoids such as beta-carotene, lycopene, and the xanthophyll carotenoids such as lutein and zeaxanthin; phenolics comprising the flavonoids such as the flavonols (e.g. quercetin, rutin), the flavans/tannins (such as the procyanidins comprising coumarin, proanthocyanidins, catechins, and anthocyanins); flavones; phytoestrogens such as coumestans, lignans, resveratrol, isoflavones e.g. genistein, daidzein, and glycitein; resorcylic acid lactones; organosulphur compounds; phytosterols; terpenoids such as carnosol, rosmarinic acid, glycyrrhizin and saponins; chlorophyll; chlorophyllin, sugars, anthocyanins, and vanilla.

10 In other embodiments, CESA-inhibiting herbicides-tolerant plants of the present invention, relative to a wild-type plant, comprise an increased amount of, or an improved profile of, a compound selected from the group consisting of: vincristine, vinblastine, taxanes (e.g., taxol (paclitaxel), baccatin III, 10-desacetylbaccatin III, 10-desacetyl taxol, xylosyl taxol, 7-epitaxol, 7-epibaccatin III, 10-desacetylcephalomannine, 7-epicephalomannine, taxotere, cephalomannine, xylosyl cephalomannine, taxagifine, 8-benxoyloxy taxagifine, 9-acetyloxy taxusin, 9-hydroxy taxusin, taiwanxam, taxane Ia, taxane Ib, taxane Ic, taxane Id, GMP paclitaxel, 9-dihydro 13-acetylbaccatin III, 10-desacetyl-7-epitaxol, tetrahydrocannabinol (THC), cannabidiol (CBD), genistein, diadzein, codeine, morphine, quinine, shikonin, ajmalacine, serpentine, and the like.

20

In other aspects, a method for treating a plant of the present invention is provided.

In some embodiments, the method comprises contacting the plant with an agronomically acceptable composition. In one embodiment, the agronomically acceptable composition comprises a CESA inhibiting herbicide A. I, such as an azine as described herein.

25

In another aspect, the present invention provides a method for preparing a descendent seed. The method comprises planting a seed of or capable of producing a plant of the present invention. In one embodiment, the method further comprises growing a descendent plant from the seed; and harvesting a descendant seed from the descendent plant. In other embodiments, the method further comprises applying a CESA-inhibiting herbicides herbicidal composition to the descendent plant.

30

In another embodiment, the invention refers to harvestable parts of the plant according to the present invention. Preferably, the harvestable parts comprise the CESA nucleic acid or CESA protein of the present invention. The harvestable parts may be seeds, roots, leaves and/or flowers comprising the CESA nucleic acid or CESA protein or parts thereof. Preferred parts of soy plants are soy beans comprising the CESA nucleic acid or CESA protein.

40

In another embodiment, the invention refers to products derived from a plant according to the present invention, parts thereof or harvestable parts thereof. A preferred plant product is

fodder, seed meal, oil, or seed-treatment-coated seeds. Preferably, the meal and/or oil comprise the CESA nucleic acids or CESA proteins.

- 5 In another embodiment, the invention refers to a method for the production of a product, which method comprises
- a) growing the plants of the invention or obtainable by the methods of invention and
 - b) producing said product from or by the plants of the invention and/or parts, e.g. seeds, of these plants.
- 10 In a further embodiment the method comprises the steps
- a) growing the plants of the invention,
 - b) removing the harvestable parts as defined above from the plants and
 - c) producing said product from or by the harvestable parts of the invention.
- 15 The product may be produced at the site where the plant has been grown, the plants and/or parts thereof may be removed from the site where the plants have been grown to produce the product. Typically, the plant is grown, the desired harvestable parts are removed from the plant, if feasible in repeated cycles, and the product made from the harvestable parts of the plant. The step of growing the plant may be performed only once each time the methods
- 20 of the invention is performed, while allowing repeated times the steps of product production e.g. by repeated removal of harvestable parts of the plants of the invention and if necessary further processing of these parts to arrive at the product. It is also possible that the step of growing the plants of the invention is repeated and plants or harvestable parts are stored until the production of the product is then performed once for the accumulated plants or
- 25 plant parts. Also, the steps of growing the plants and producing the product may be performed with an overlap in time, even simultaneously to a large extent or sequentially. Generally the plants are grown for some time before the product is produced.

30 In one embodiment the products produced by said methods of the invention are plant products such as, but not limited to, a foodstuff, feedstuff, a food supplement, feed supplement, fiber, cosmetic and/or pharmaceutical. Foodstuffs are regarded as compositions used for nutrition and/or for supplementing nutrition. Animal feedstuffs and animal feed supplements, in particular, are regarded as foodstuffs.

35 In another embodiment the inventive methods for the production are used to make agricultural products such as, but not limited to, plant extracts, proteins, amino acids, carbohydrates, fats, oils, polymers, vitamins, and the like.




40 It is possible that a plant product consists of one or more agricultural products to a large extent.






Herbicides

As described above, the present invention provides nucleic acids, polypeptides, conferring tolerance of plants to compounds / herbicides interfering or inhibiting cell wall (cellulose) biosynthesis by interfering with the activity of cellulose synthase (“CESA-inhibiting herbicides”), also known to the person skilled in the art as Cellulose Biosynthesis Inhibitors (CBI).

Examples of herbicides which can be used according to the present invention, i.e. to which the plants according to the present invention are tolerant / resistant to, are compounds known to the skilled artisan as azines. Examples of Azines are described in detail in the following patent applications depicted in the following Table 1.


Table 1

No.:	Structural Formula	Publication or Application number/Internal reference
1	 I,	WO 2014/064094 PF74283
2	 (I),	WO 2015/007711 PF75365
3	 (II)	PCT/EP2015/056711 PF76068

4	 (I)	EP 14163356.0 PF76069
5	 (I)	EP 14163742.1 PF76635
6	 (I)	EP 14163743.9 PF76636
7		EP 14165565.4 PF76857
8		EP 14165624.9 PF76888

9		EP 14164431.0 PF76890
10		EP 14164434.4 PF76930
11	(I)	EP 14164433.6 PF77027
12		Indaziflam
13		Triazofenamid

Examples of preferred CESA inhibiting herbicides from the group of so-called azines which can be used according to the present invention are compounds having the Formula (I), known to the skilled artisan as azines.



wherein

A is phenyl, which is substituted by two to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₁-C₆-haloalkynyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl;

R¹ H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl or phenylsulfonyl,

wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl and C₁-C₆-alkoxy;

R² H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, OH, C₁-C₆-alkoxy or C₁-C₆-alkoxy-C₁-C₆-alkyl;

R³ H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxy;

R⁴ H, halogen, CN, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or

R³ and R⁴ together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl and three- to six-membered heterocyclyl,

wherein the C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, or three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy; and

R⁵ H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl or phenylsulfonyl,

wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl and C₁-C₆-alkoxy;

including their agriculturally acceptable salts or N-oxides.

Preferably the present invention provides azines of formula (I), wherein

A is 2-fluoro-phenyl, which is substituted by one to four substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl and (C₁-C₆-alkoxy)-carbonyl;

R¹ H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-

- C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl or phenylsulfonyl, wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl and C₁-C₆-alkoxy;
- 5 R² H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, OH, C₁-C₆-alkoxy or C₁-C₆-alkoxy-C₁-C₆-alkyl;
- R³ H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxy;
- R⁴ H, halogen, CN, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or
- R³ and R⁴ together with the carbon atom to which they are attached form a moiety
- 10 selected from the group consisting of carbonyl, C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl and three- to six-membered heterocyclyl, wherein the C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl or and three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy; and
- 15 R⁵ H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl or phenylsulfonyl, wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl and C₁-C₆-alkoxy;
- 20 including their agriculturally acceptable salts or N-oxides.

Useful for the present invention are also agrochemical compositions comprising at least one azines of formula (I) and auxiliaries customary for formulating crop protection agents.

- 25 The present invention also provides the use of azines of formula (I) as herbicides, i.e. for controlling harmful plants.

If the azines of formula (I) as described herein are capable of forming geometrical isomers, for example E/Z isomers, it is possible to use both, the pure isomers and mixtures thereof,

30 in the compositions according to the invention.

If the azines of formula (I) as described herein have one or more centres of chirality and, as a consequence, are present as enantiomers or diastereomers, it is possible to use both, the pure enantiomers and diastereomers and their mixtures, in the compositions according to

35 the invention.

If the azines of formula (I) as described herein have ionizable functional groups, they can also be employed in the form of their agriculturally acceptable salts. Suitable are, in general, the salts of those cations and the acid addition salts of those acids whose cations and

40 anions, respectively, have no adverse effect on the activity of the active compounds.

Preferred cations are the ions of the alkali metals, preferably of lithium, sodium and

potassium, of the alkaline earth metals, preferably of calcium and magnesium, and of the transition metals, preferably of manganese, copper, zinc and iron, further ammonium and substituted ammonium in which one to four hydrogen atoms are replaced by C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, hydroxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl or benzyl, preferably ammonium, methylammonium, isopropylammonium, dimethylammonium, diisopropylammonium, trimethylammonium, heptylammonium, dodecylammonium, tetradecylammonium, tetramethylammonium, tetraethylammonium, tetrabutylammonium, 2-hydroxyethylammonium (olamine salt), 2-(2-hydroxyeth-1-oxy)eth-1-ylammonium (diglycolamine salt), di(2-hydroxyeth-1-yl)ammonium (diolamine salt), tris(2-hydroxyethyl)ammonium (trolamine salt), tris(2-hydroxypropyl)ammonium, benzyltrimethylammonium, benzyltriethylammonium, N,N,N-trimethylethanolammonium (choline salt), furthermore phosphonium ions, sulfonium ions, preferably tri(C₁-C₄-alkyl)sulfonium, such as trimethylsulfonium, and sulfoxonium ions, preferably tri(C₁-C₄-alkyl)sulfoxonium, and finally the salts of polybasic amines such as N,N-bis-(3-aminopropyl)methylamine and diethylenetriamine.

Anions of useful acid addition salts are primarily chloride, bromide, fluoride, iodide, hydrogensulfate, methylsulfate, sulfate, dihydrogenphosphate, hydrogenphosphate, nitrate, bicarbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate and also the anions of C₁-C₄-alkanoic acids, preferably formate, acetate, propionate and butyrate.

The organic moieties mentioned in the definition of the variables, e.g. R¹ to R⁵, are - like the term halogen - collective terms for individual enumerations of the individual group members. The term halogen denotes in each case fluorine, chlorine, bromine or iodine. All hydrocarbon chains, i.e. all alkyl, haloalkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, (alkyl)amino, di(alkyl)amino chains can be straight-chain or branched, the prefix C_n-C_m denoting in each case the possible number of carbon atoms in the group.

Examples of such meanings are:

- C₁-C₄-alkyl: for example CH₃, C₂H₅, n-propyl, CH(CH₃)₂, n-butyl, CH(CH₃)-C₂H₅, CH₂-CH(CH₃)₂ and C(CH₃)₃;
- C₁-C₆-alkyl and also the C₁-C₆-alkyl moieties of (C₁-C₆-alkyl)carbonyl, C₁-C₆-alkoxy-C₁-C₆-alkyl: C₁-C₄-alkyl as mentioned above, and also, for example, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl or 1-ethyl-2-methylpropyl, preferably methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1,1-dimethylethyl, n-pentyl or n-hexyl;
- C₁-C₄-haloalkyl: a C₁-C₄-alkyl radical as mentioned above which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, for example, chloromethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl,

- chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, bromomethyl, iodomethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-iodoethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl, pentafluoroethyl, 2-fluoropropyl, 3-fluoropropyl, 2,2-difluoropropyl, 2,3-difluoropropyl, 2-chloropropyl, 3-chloropropyl, 2,3-dichloropropyl, 2-bromopropyl, 3-bromopropyl, 3,3,3-trifluoropropyl, 3,3,3-trichloropropyl, 2,2,3,3,3-pentafluoropropyl, heptafluoropropyl, 1-(fluoromethyl)-2-fluoroethyl, 1-(chloromethyl)-2-chloroethyl, 1-(bromomethyl)-2-bromoethyl, 4-fluorobutyl, 4-chlorobutyl, 4-bromobutyl, nonafluorobutyl, 1,1,2,2,-tetrafluoroethyl and 1-trifluoromethyl-1,2,2,2-tetrafluoroethyl;
- 10 - C₁-C₆-haloalkyl: C₁-C₄-haloalkyl as mentioned above, and also, for example, 5-fluoropentyl, 5-chloropentyl, 5-bromopentyl, 5-iodopentyl, undecafluoropentyl, 6-fluorohexyl, 6-chlorohexyl, 6-bromohexyl, 6-iodohexyl and dodecafluorohexyl;
- C₃-C₆-cycloalkyl: monocyclic saturated hydrocarbons having 3 to 6 ring members, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl;
- 15 - C₂-C₆-alkenyl: for example ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-
- 20 2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-
- 25 methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-
- 30 butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl and 1-ethyl-2-methyl-2-propenyl;
- C₃-C₆-cycloalkenyl: 1-cyclopropenyl, 2-cyclopropenyl, 1-cyclobutenyl, 2-cyclobutenyl, 1-cyclopentenyl, 2-cyclopentenyl, 1,3-cyclopentadienyl, 1,4-cyclopentadienyl, 2,4-cyclopentadienyl, 1-cyclohexenyl, 2-cyclohexenyl, 3-cyclohexenyl, 1,3-cyclohexadienyl, 1,4-
- 35 cyclohexadienyl, 2,5-cyclohexadienyl;
- C₃-C₆-alkynyl: for example 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-methyl-2-propynyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-methyl-2-butylnyl, 1-methyl-3-butylnyl, 2-methyl-3-butylnyl, 3-methyl-1-butylnyl, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentylnyl, 1-
- 40 methyl-3-pentylnyl, 1-methyl-4-pentylnyl, 2-methyl-3-pentylnyl, 2-methyl-4-pentylnyl, 3-methyl-1-pentylnyl, 3-methyl-4-pentylnyl, 4-methyl-1-pentylnyl, 4-methyl-2-pentylnyl, 1,1-dimethyl-

2-butynyl, 1,1-dimethyl-3-butynyl, 1,2-dimethyl-3-butynyl, 2,2-dimethyl-3-butynyl, 3,3-dimethyl-1-butynyl, 1-ethyl-2-butynyl, 1-ethyl-3-butynyl, 2-ethyl-3-butynyl and 1-ethyl-1-methyl-2-propynyl;

- C₁-C₄-alkoxy: for example methoxy, ethoxy, propoxy, 1-methylethoxy butoxy, 5 1-methylpropoxy, 2-methylpropoxy and 1,1-dimethylethoxy;
- C₁-C₆-alkoxy and also the C₁-C₆-alkoxy moieties of (C₁-C₆-alkoxy)carbonyl, C₁-C₆-alkoxy-C₁-C₆-alkyl: C₁-C₄-alkoxy as mentioned above, and also, for example, pentoxy, 1-methylbutoxy, 2-methylbutoxy, 3-methoxybutoxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hexoxy, 1-methylpentoxy, 2-methylpentoxy, 3-methylpentoxy, 4-methylpentoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy and 1-ethyl-2-methylpropoxy.
- C₁-C₄-alkylthio: for example methylthio, ethylthio, propylthio, 1-methylethylthio, 15 butylthio, 1-methylpropylthio, 2-methylpropylthio and 1,1-dimethylethylthio;
- C₁-C₆-alkylthio: C₁-C₄-alkylthio as mentioned above, and also, for example, pentylthio, 1-methylbutylthio, 2-methylbutylthio, 3-methylbutylthio, 2,2-dimethylpropylthio, 1-ethylpropylthio, hexylthio, 1,1-dimethylpropylthio, 1,2-dimethylpropylthio, 1-methylpentylthio, 2-methylpentylthio, 3-methylpentylthio, 4-methylpentylthio, 1,1-dimethylbutylthio, 1,2-dimethylbutylthio, 1,3-dimethylbutylthio, 2,2-dimethylbutylthio, 2,3-dimethylbutylthio, 3,3-dimethylbutylthio, 1-ethylbutylthio, 2-ethylbutylthio, 1,1,2-trimethylpropylthio, 1,2,2-trimethylpropylthio, 1-ethyl-1-methylpropylthio and 1-ethyl-2-methylpropylthio;
- C₁-C₆-alkylsulfinyl (C₁-C₆-alkyl-S(=O)-): z.B. methylsulfinyl, ethylsulfinyl, propylsulfinyl, 1-methylethylsulfinyl, butylsulfinyl, 1-methylpropylsulfinyl, 2-methylpropylsulfinyl, 1,1-dimethylethylsulfinyl, pentylsulfinyl, 1-methylbutylsulfinyl, 2-methylbutylsulfinyl, 3-methylbutylsulfinyl, 2,2-dimethylpropylsulfinyl, 1-ethylpropylsulfinyl, 1,1-dimethylpropylsulfinyl, 1,2-dimethylpropylsulfinyl, hexylsulfinyl, 1-methylpentylsulfinyl, 2-methylpentylsulfinyl, 3-methylpentylsulfinyl, 4-methylpentylsulfinyl, 1,1-dimethylbutylsulfinyl, 1,2-dimethylbutylsulfinyl, 1,3-dimethylbutylsulfinyl, 2,2-dimethylbutylsulfinyl, 2,3-dimethylbutylsulfinyl, 3,3-dimethylbutylsulfinyl, 1-ethylbutylsulfinyl, 2-ethylbutylsulfinyl, 1,1,2-trimethylpropylsulfinyl, 1,2,2-trimethylpropylsulfinyl, 1-ethyl-1-methylpropylsulfinyl and 1-ethyl-2-methylpropylsulfinyl;
- C₁-C₆-alkylsulfonyl (C₁-C₆-alkyl-S(O)₂-): for example methylsulfonyl, ethylsulfonyl, propylsulfonyl, 1-methylethylsulfonyl, butylsulfonyl, 1-methylpropylsulfonyl, 2-methylpropylsulfonyl, 1,1-dimethylethylsulfonyl, pentylsulfonyl, 1-methylbutylsulfonyl, 2-methylbutylsulfonyl, 3-methylbutylsulfonyl, 1,1-dimethylpropylsulfonyl, 1,2-dimethylpropylsulfonyl, 2,2-dimethylpropylsulfonyl, 1-ethylpropylsulfonyl, hexylsulfonyl, 1-methylpentylsulfonyl, 2-methylpentylsulfonyl, 3-methylpentylsulfonyl, 4-methylpentylsulfonyl, 1,1-dimethylbutylsulfonyl, 1,2-dimethylbutylsulfonyl, 1,3-dimethylbutylsulfonyl, 2,2-dimethylbutylsulfonyl, 2,3-dimethylbutylsulfonyl, 3,3-dimethylbutylsulfonyl, 1-ethylbutylsulfonyl, 2-ethylbutylsulfonyl, 1,1,2-trimethylpropylsulfonyl, 1,2,2-trimethylpropylsulfonyl, 1-ethyl-1-methylpropylsulfonyl and 1-ethyl-2-methylpropylsulfonyl;
- C₁-C₆-alkylsulfonyl (C₁-C₆-alkyl-S(O)₂-): for example methylsulfonyl, ethylsulfonyl, propylsulfonyl, 1-methylethylsulfonyl, butylsulfonyl, 1-methylpropylsulfonyl, 2-methylpropylsulfonyl, 1,1-dimethylethylsulfonyl, pentylsulfonyl, 1-methylbutylsulfonyl, 2-methylbutylsulfonyl, 3-methylbutylsulfonyl, 1,1-dimethylpropylsulfonyl, 1,2-dimethylpropylsulfonyl, 2,2-dimethylpropylsulfonyl, 1-ethylpropylsulfonyl, hexylsulfonyl, 1-methylpentylsulfonyl, 2-methylpentylsulfonyl, 3-methylpentylsulfonyl, 4-methylpentylsulfonyl, 1,1-dimethylbutylsulfonyl, 1,2-dimethylbutylsulfonyl, 1,3-dimethylbutylsulfonyl, 2,2-dimethylbutylsulfonyl, 2,3-dimethylbutylsulfonyl, 3,3-dimethylbutylsulfonyl, 1-ethylbutylsulfonyl, 2-ethylbutylsulfonyl, 1,1,2-trimethylpropylsulfonyl, 1,2,2-trimethylpropylsulfonyl, 1-ethyl-1-methylpropylsulfonyl and 1-ethyl-2-methylpropylsulfonyl;

- (C₁-C₄-alkyl)amino: for example methylamino, ethylamino, propylamino, 1-methylethylamino, butylamino, 1-methylpropylamino, 2-methylpropylamino or 1,1-dimethylethylamino;
- (C₁-C₆-alkyl)amino: (C₁-C₄-alkylamino) as mentioned above, and also, for example, pentylamino, 1-methylbutylamino, 2-methylbutylamino, 3-methylbutylamino, 2,2-dimethylpropylamino, 1-ethylpropylamino, hexylamino, 1,1-dimethylpropylamino, 1,2-dimethylpropylamino, 1-methylpentylamino, 2-methylpentylamino, 3-methylpentylamino, 4-methylpentylamino, 1,1-dimethylbutylamino, 1,2-dimethylbutylamino, 1,3-dimethylbutylamino, 2,2-dimethylbutylamino, 2,3-dimethylbutylamino, 3,3-dimethylbutylamino, 1-ethylbutylamino, 2-ethylbutylamino, 1,1,2-trimethylpropylamino, 1,2,2-trimethylpropylamino, 1-ethyl-1-methylpropylamino or 1-ethyl-2-methylpropylamino;
- di(C₁-C₄-alkyl)amino: for example N,N-dimethylamino, N,N-diethylamino, N,N-di(1-methylethyl)amino, N,N-dipropylamino, N,N-dibutylamino, N,N-di(1-methylpropyl)amino, N,N-di(2-methylpropyl)amino, N,N-di(1,1-dimethylethyl)amino, N-ethyl-N-methylamino, N-methyl-N-propylamino, N-methyl-N-(1-methylethyl)amino, N-butyl-N-methylamino, N-methyl-N-(1-methylpropyl)amino, N-methyl-N-(2-methylpropyl)amino, N-(1,1-dimethylethyl)-N-methylamino, N-ethyl-N-propylamino, N-ethyl-N-(1-methylethyl)amino, N-butyl-N-ethylamino, N-ethyl-N-(1-methylpropyl)amino, N-ethyl-N-(2-methylpropyl)amino, N-ethyl-N-(1,1-dimethylethyl)amino, N-(1-methylethyl)-N-propylamino, N-butyl-N-propylamino, N-(1-methylpropyl)-N-propylamino, N-(2-methylpropyl)-N-propylamino, N-(1,1-dimethylethyl)-N-propylamino, N-butyl-N-(1-methylethyl)amino, N-(1-methylethyl)-N-(1-methylpropyl)amino, N-(1-methylethyl)-N-(2-methylpropyl)amino, N-(1,1-dimethylethyl)-N-(1-methylethyl)amino, N-butyl-N-(1-methylpropyl)amino, N-butyl-N-(2-methylpropyl)amino, N-butyl-N-(1,1-dimethylethyl)amino, N-(1-methylpropyl)-N-(2-methylpropyl)amino, N-(1,1-dimethylethyl)-N-(1-methylpropyl)amino or N-(1,1-dimethylethyl)-N-(2-methylpropyl)amino;
- di(C₁-C₆-alkyl)amino: di(C₁-C₄-alkyl)amino as mentioned above, and also, for example, N-methyl-N-pentylamino, N-methyl-N-(1-methylbutyl)amino, N-methyl-N-(2-methylbutyl)amino, N-methyl-N-(3-methylbutyl)amino, N-methyl-N-(2,2-dimethylpropyl)amino, N-methyl-N-(1-ethylpropyl)amino, N-methyl-N-hexylamino, N-methyl-N-(1,1-dimethylpropyl)amino, N-methyl-N-(1,2-dimethylpropyl)amino, N-methyl-N-(1-methylpentyl)amino, N-methyl-N-(2-methylpentyl)amino, N-methyl-N-(3-methylpentyl)amino, N-methyl-N-(4-methylpentyl)amino, N-methyl-N-(1,1-dimethylbutyl)amino, N-methyl-N-(1,2-dimethylbutyl)amino, N-methyl-N-(1,3-dimethylbutyl)amino, N-methyl-N-(2,2-dimethylbutyl)amino, N-methyl-N-(2,3-dimethylbutyl)amino, N-methyl-N-(3,3-dimethylbutyl)amino, N-methyl-N-(1-ethylbutyl)amino, N-methyl-N-(2-ethylbutyl)amino, N-methyl-N-(1,1,2-trimethylpropyl)amino, N-methyl-N-(1,2,2-trimethylpropyl)amino, N-methyl-N-(1-ethyl-1-methylpropyl)amino, N-methyl-N-(1-ethyl-2-methylpropyl)amino, N-ethyl-N-pentylamino, N-ethyl-N-(1-methylbutyl)amino, N-ethyl-N-(2-methylbutyl)amino, N-ethyl-N-(3-methylbutyl)amino, N-ethyl-N-(2,2-dimethylpropyl)amino, N-ethyl-N-(1-ethylpropyl)amino, N-ethyl-N-hexylamino, N-ethyl-N-(1,1-dimethylpropyl)amino, N-ethyl-N-(1,2-dimethylpropyl)amino, N-ethyl-N-(1-methylpentyl)amino, N-ethyl-N-(2-methylpentyl)amino, N-ethyl-N-(3-methylpentyl)amino, N-ethyl-N-(4-methylpentyl)amino, N-ethyl-N-(1,1-dimethylbutyl)amino, N-ethyl-N-(1,2-dimethylbutyl)amino, N-ethyl-N-(1,3-dimethylbutyl)amino, N-ethyl-N-(2,2-

- dimethylbutyl)amino, N-ethyl-N-(2,3-dimethylbutyl)amino, N-ethyl-N-(3,3-dimethylbutyl)amino, N-ethyl-N-(1-ethylbutyl)amino, N-ethyl-N-(2-ethylbutyl)amino, N-ethyl-N-(1,1,2-trimethylpropyl)amino, N-ethyl-N-(1,2,2-trimethylpropyl)amino, N-ethyl-N-(1-ethyl-1-methylpropyl)amino, N-ethyl-N-(1-ethyl-2-methylpropyl)amino, N-propyl-N-pentylamino,
- 5 N-butyl-N-pentylamino, N,N-dipentylamino, N-propyl-N-hexylamino, N-butyl-N-hexylamino, N-pentyl-N-hexylamino or N,N-dihexylamino;
- three- to six-membered heterocyclyl: monocyclic saturated or partially unsaturated hydrocarbon having three to six ring members as mentioned above which, in addition to carbon atoms, contains one or two heteroatoms selected from O, S and N;
- 10 for example 2-oxiranyl, 2-oxetanyl, 3-oxetanyl, 2-aziridinyl, 3-thietanyl, 1-azetidiny, 2-azetidiny,
- for example 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydrothienyl, 3-tetra-
- 15 hydrothienyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 3-isoxazolidinyl, 4-isoxazolidinyl, 5-isoxazolidinyl, 3-isothiazolidinyl, 4-isothiazolidinyl, 5-isothiazolidinyl, 3-pyrazolidinyl, 4-pyrazolidinyl, 5-pyrazolidinyl, 2-oxazolidinyl, 4-oxazolidinyl, 5-oxazolidinyl, 2-thiazolidinyl, 4-thiazolidinyl, 5-thiazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl;
- 20 for example 2,3-dihydrofur-2-yl, 2,3-dihydrofur-3-yl, 2,4-dihydrofur-2-yl, 2,4-dihydrofur-3-yl, 2,3-dihydrothien-2-yl, 2,3-dihydrothien-3-yl, 2,4-dihydrothien-2-yl, 2,4-dihydrothien-3-yl, 4,5-dihydropyrrol-2-yl, 4,5-dihydropyrrol-3-yl, 2,5-dihydropyrrol-2-yl, 2,5-dihydropyrrol-3-yl, 4,5-dihydroisoxazol-3-yl, 2,5-dihydroisoxazol-3-yl, 2,3-dihydroisoxazol-3-yl, 4,5-dihydroisoxazol-4-yl, 2,5-dihydroisoxazol-4-yl, 2,3-dihydroisoxazol-4-yl, 4,5-dihydroisoxazol-5-yl, 2,5-
- 25 dihydroisoxazol-5-yl, 2,3-dihydroisoxazol-5-yl, 4,5-dihydroisothiazol-3-yl, 2,5-dihydroisothiazol-3-yl, 2,3-dihydroisothiazol-3-yl, 4,5-dihydroisothiazol-4-yl, 2,5-dihydroisothiazol-4-yl, 2,3-dihydroisothiazol-4-yl, 4,5-dihydroisothiazol-5-yl, 2,5-dihydroisothiazol-5-yl, 2,3-dihydroisothiazol-5-yl, 2,3-dihydropyrazol-2-yl, 2,3-dihydropyrazol-3-yl, 2,3-dihydropyrazol-4-yl, 2,3-dihydropyrazol-5-yl, 3,4-dihydropyrazol-3-
- 30 yl, 3,4-dihydropyrazol-4-yl, 3,4-dihydropyrazol-5-yl, 4,5-dihydropyrazol-3-yl, 4,5-dihydropyrazol-4-yl, 4,5-dihydropyrazol-5-yl, 2,3-dihydroimidazol-2-yl, 2,3-dihydroimidazol-3-yl, 2,3-dihydroimidazol-4-yl, 2,3-dihydroimidazol-5-yl, 4,5-dihydroimidazol-2-yl, 4,5-dihydroimidazol-4-yl, 4,5-dihydroimidazol-5-yl, 2,5-dihydroimidazol-2-yl, 2,5-dihydroimidazol-4-yl, 2,5-dihydroimidazol-5-yl, 2,3-dihydrooxazol-3-yl, 2,3-dihydrooxazol-4-yl, 2,3-
- 35 dihydrooxazol-5-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 3,4-dihydrooxazol-5-yl, 2,3-dihydrothiazol-3-yl, 2,3-dihydrothiazol-4-yl, 2,3-dihydrothiazol-5-yl, 3,4-dihydrothiazol-3-yl, 3,4-dihydrothiazol-4-yl, 3,4-dihydrothiazol-5-yl, 3,4-dihydrothiazol-2-yl, 3,4-dihydrothiazol-3-yl, 3,4-dihydrothiazol-4-yl;
- 40 for example 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1,3-dioxan-2-yl, 1,3-dioxan-4-yl, 1,3-dioxan-5-yl, 1,4-dioxan-2-yl, 1,3-dithian-2-yl, 1,3-dithian-4-yl, 1,4-dithian-2-yl, 1,3-dithian-5-

yl, 2-tetrahydropyranyl, 3-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothiopyranyl, 3-tetrahydrothiopyranyl, 4-tetrahydrothiopyranyl, 3-hexahydropyridazinyl, 4-hexahydropyridazinyl, 2-hexahydropyrimidinyl, 4-hexahydropyrimidinyl, 5-hexahydropyrimidinyl, 2-piperazinyl, tetrahydro-1,3-oxazin-2-yl, tetrahydro-1,3-oxazin-6-yl, 2-morpholinyl, 3-morpholinyl;

for example 2H-pyran-2-yl, 2H-pyran-3-yl, 2H-pyran-4-yl, 2H-pyran-5-yl, 2H-pyran-6-yl, 3,6-dihydro-2H-pyran-2-yl, 3,6-dihydro-2H-pyran-3-yl, 3,6-dihydro-2H-pyran-4-yl, 3,6-dihydro-2H-pyran-5-yl, 3,6-dihydro-2H-pyran-6-yl, 3,4-dihydro-2H-pyran-3-yl, 3,4-dihydro-2H-pyran-4-yl, 3,4-dihydro-2H-pyran-6-yl, 2H-thiopyran-2-yl, 2H-thiopyran-3-yl, 2H-thiopyran-4-yl, 2H-thiopyran-5-yl, 2H-thiopyran-6-yl, 5,6-dihydro-4H-1,3-oxazin-2-yl;

The preferred embodiments of the invention mentioned herein below have to be understood as being preferred either independently from each other or in combination with one another.

15

According to a preferred embodiment of the invention preference is also given to those azines of formula (I), wherein the variables, either independently of one another or in combination with one another, have the following meanings:

20 Preferred are the azines of formula (I), wherein

A is phenyl, which is substituted by two to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl;

25

particularly preferred phenyl, which is substituted by two to five substituents selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;

particularly preferred selected from halogen and CN;

30

also particularly preferred selected from the group consisting of F, Cl, CN and CH₃;

especially preferred selected from the group consisting of F, Cl and CN;

especially preferred phenyl, which is substituted by two to four substituents

35 selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl;

particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;

40

especially preferred selected from halogen and CN;

also especially preferred selected from the group consisting of F, Cl, CN and

CH₃;

more preferred selected from the group consisting of F, Cl and CN;

more preferred phenyl, which is substituted by two substituents

5 selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl;
 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
 10 especially preferred selected from halogen and CN;
 also especially preferred selected from the group consisting of F, Cl, CN and CH₃;
 more preferred selected from the group consisting of F, Cl and CN;


15 also more preferred phenyl, which is substituted by three substituents

selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl;
 20 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
 especially preferred selected from halogen and CN;
 also especially preferred selected from the group consisting of F, Cl, CN and CH₃;
 25 more preferred selected from the group consisting of F, Cl and CN;

also more preferred phenyl, which is substituted by four substituents

selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl;
 30 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
 especially preferred selected from halogen and CN;
 also especially preferred selected from the group consisting of F, Cl, CN and CH₃;
 35 more preferred selected from the group consisting of F, Cl and CN.

40 Also preferred are the azines of formula (I), wherein
 A is



wherein

R^a and R^e independently of one another are halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl; and

R^b , R^c and R^d independently of one another are hydrogen, halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl;

particularly preferred R^a and R^e independently of one another are halogen, CN, C₁-C₆-alkyl or C₁-C₆-alkoxy; and

R^b , R^c and R^d independently of one another are hydrogen, halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxy;

especially preferred R^a and R^e independently of one another are halogen or CN; and

R^b , R^c and R^d independently of one another are hydrogen, halogen, CN, C₁-C₆-alkyl or C₁-C₆-alkoxy;

more preferred R^a and R^e are halogen; and

R^b , R^c and R^d independently of one another are hydrogen, halogen or CN;

most preferred R^a and R^e are halogen; and

R^b , R^c and R^d are hydrogen;


also most preferred R^a , R^b , R^d and R^e are halogen; and

R^c hydrogen;

also most preferred R^a , R^b , R^c , R^d and R^e are halogen.

Also preferred are the azines of formula (I), wherein

A is



wherein R^a is halogen or CN;
 R^b and R^d are H, halogen or CN;
 R^c is H or halogen;
 R^e is halogen, CN or C₁-C₆-alkyl;

5

particularly preferred R^a is halogen;
 R^b , R^c and R^d are H or halogen; and
 R^e is halogen or CN;


10

especially preferred R^a , R^b , R^d and R^e are halogen; and
 R^c is H or halogen;


more preferred R^a , R^b , R^d and R^e are F; and
 R^c is H or F.

15


Especially preferred are the azines of formula (I), wherein A is selected from the group consisting of (A.1.1), (A.1.2) and (A.1.3);
more preferred selected from the group consisting of (A.1.2) and (A.1.3);



(A.1.1)



(A.1.2)



(A.1.3)

20

wherein

R^a and R^e independently of one another are halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl; and

25

R^b and R^d independently of one another are halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl;

30

particularly preferred R^a and R^e independently of one another are halogen, CN, C₁-C₆-alkyl or C₁-C₆-alkoxy; and
 R^b and R^d independently of one another are halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxy;

35

especially preferred R^a and R^e independently of one another halogen or CN; and
 R^b and R^d independently of one another are halogen, CN, C₁-C₆-alkyl or C₁-C₆-

alkoxy;

more preferred R^a and R^e are halogen; and


R^b and R^d independently of one another are halogen or CN;

5

most preferred R^a , R^b , R^d and R^e are halogen.

Also especially preferred are the azines of formula (I), wherein

A is




10

wherein R^a , R^b , R^d and R^e have the meanings, in particular the preferred meanings, as defined above.

Also especially preferred are the azines of formula (I), wherein


15 A is



wherein R^a , R^b and R^e have the meanings, in particular the preferred meanings, as defined above.

20 Also especially preferred are the azines of formula (I), wherein

A is



wherein R^a and R^e have the meanings, in particular the preferred meanings, as defined above.

25

Also preferred are the azines of formula (I), wherein

A is 2-fluoro-phenyl, which is substituted by one to four substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl and (C₁-C₆-alkoxy)carbonyl;

30

- particularly preferred 2-fluoro-phenyl, which is substituted by one to four substituents selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
- 5 particularly preferred selected from halogen and CN;
also particularly preferred selected from the group consisting of F, Cl, CN and CH₃;
especially preferred selected from the group consisting of F, Cl and CN;
- 10 especially preferred 2-fluoro-phenyl, which is substituted by one to three substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl and (C₁-C₆-alkoxy)carbonyl;
- 15 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
especially preferred selected from halogen and CN;
also especially preferred selected from the group consisting of F, Cl, CN and CH₃;
- 20 more preferred selected from the group consisting of F, Cl and CN;
- more preferred 2-fluoro-phenyl, which is substituted by one substituent selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl and (C₁-C₆-alkoxy)carbonyl;
- 25 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
especially preferred selected from halogen and CN;
- 30 also especially preferred selected from the group consisting of F, Cl, CN and CH₃;
more preferred selected from the group consisting of F, Cl and CN;
- also more preferred 2-fluoro-phenyl, which is substituted by two substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl and (C₁-C₆-alkoxy)carbonyl;
- 35 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
- 40 especially preferred selected from halogen and CN;
also especially preferred selected from the group consisting of F, Cl, CN and

CH₃;

more preferred selected from the group consisting of F, Cl and CN;

also more preferred 2-fluoro-phenyl, which is substituted by three substituents

5 selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl and (C₁-C₆-alkoxy)carbonyl;

10 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;

especially preferred selected from halogen and CN;


also especially preferred selected from the group consisting of F, Cl, CN and CH₃;

more preferred selected from the group consisting of F, Cl and CN.

15

Also preferred are the azines of formula (I), wherein

A is



wherein

20 R^a is halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl; and

25 R^b, R^c and R^d independently of one another are hydrogen, halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl;

particularly preferred R^a is halogen, CN, C₁-C₆-alkyl or C₁-C₆-alkoxy; and

30 R^b, R^c and R^d independently of one another are hydrogen, halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxy;

especially preferred R^a is halogen or CN; and

35 R^b, R^c and R^d independently of one another are hydrogen, halogen, CN, C₁-C₆-alkyl or C₁-C₆-alkoxy;

more preferred R^a is halogen; and

R^b, R^c and R^d independently of one another are hydrogen, halogen or CN;

most preferred R^a is halogen; and

R^b , R^c and R^d are hydrogen;

also most preferred R^a , R^b and R^d are halogen; and


R^c is hydrogen;

5

also most preferred R^a , R^b , R^c and R^d are halogen.

Also preferred are the azines of formula (I), wherein

A is



10

wherein R^a is halogen, CN or C_1 - C_6 -alkyl;

R^b and R^d are H, halogen or CN; and

R^c is H or halogen;

15

particularly preferred R^a is halogen or CN; and

R^b , R^c and R^d are H or halogen;

especially preferred R^a , R^b and R^d are halogen; and

R^c is H or halogen;

20

Also especially preferred R^a , R^b and R^d are halogen; and

R^c is H, F, Br or I;

more preferred R^a , R^b and R^d are F; and

25


R^c is F, Br or I;

also more preferred R^a , R^b and R^d are F; and


R^c is H or F.

30 Especially preferred are the azines of formula (I), wherein A is selected from the group consisting of (A.1a.1), (A.1a.2) and (A.1a.3);


more preferred selected from the group consisting of (A.1.2) and (A.1.3);



(A.1a.1)



(A.1a.2)



(A.1a.3)

wherein

R^a is halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl; and

R^b and R^d independently of one another are halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl;

particularly preferred R^a is halogen, CN, C₁-C₆-alkyl or C₁-C₆-alkoxy; and

R^b and R^d independently of one another are halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxy;

especially preferred R^a is halogen or CN; and

R^b and R^d independently of one another are halogen, CN, C₁-C₆-alkyl or C₁-C₆-alkoxy;


more preferred R^a is halogen; and

R^b and R^d independently of one another are halogen or CN;

most preferred R^a , R^b and R^d are halogen.

Also especially preferred are the azines of formula (I), wherein

A is




(A.1a.1)

wherein R^a , R^b and R^d have the meanings, in particular the preferred meanings, as defined above.


Also especially preferred are the azines of formula (I), wherein

A is



wherein R^a and R^b have the meanings, in particular the preferred meanings, as defined above.

- 5 Also especially preferred are the azines of formula (I), wherein A is



wherein R^a has the meanings, in particular the preferred meanings, as defined above.

- 10 Also preferred are the azines of formula (I), wherein R^1 is H, CN, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, (C_1 - C_6 -alkyl)carbonyl or (C_1 - C_6 -alkyl)sulfonyl;
 particularly preferred H, CN, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, (C_1 - C_6 -alkyl)carbonyl or (C_1 - C_6 -alkyl)sulfonyl;
 15 especially preferred H, CN, CH_3 , CH_2OCH_3 , OCH_3 , $COCH_3$ or SO_2CH_3 ;
 more preferred hydrogen.

Also preferred are the azines of formula (I), wherein

R^2 is H, halogen, C_1 - C_6 -alkyl or C_1 - C_6 -haloalkyl;

- 20 particularly preferred halogen, C_1 - C_6 -alkyl or C_1 - C_6 -haloalkyl;
 also particularly preferred H, F, Cl, CH_3 or CF_3 .

Also preferred are the azines of formula (I), wherein

R^3 and R^4 are

- 25 independently of one another H, halogen, C_1 - C_6 -alkyl or C_1 - C_6 -haloalkyl; or
 together with the carbon atom to which they are attached form a moiety selected from
 the group consisting of C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl and three- to six-
 membered heterocyclyl,
 wherein the C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl or the three- to six-membered
 30 heterocyclyl is unsubstituted or substituted by one to three substituents selected
 from halogen, CN, C_1 - C_6 -alkyl and C_1 - C_6 -alkoxy;

independently of one another particularly preferred H, halogen, C_1 - C_6 -alkyl or C_1 - C_6 -

haloalkyl; or

together with the carbon atom to which they are attached form a moiety selected from the group consisting of C₃-C₆-cycloalkyl and C₃-C₆-cycloalkenyl,

wherein the C₃-C₆-cycloalkyl or C₃-C₆-cycloalkenyl is unsubstituted or
 5 substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;

independently of one another especially preferred H, halogen, C₁-C₆-alkyl or C₁-C₆-haloalkyl;

10

independently of one another more preferred H, halogen or C₁-C₆-alkyl.

Also preferred are the azines of formula (I), wherein

R² is H, halogen, C₁-C₆-alkyl; and

15 R³ and R⁴ are independently of one another H, halogen, C₁-C₆-alkyl, or together with the carbon atom to which they are attached form a C₃-C₆-cycloalkyl;

particularly preferred R² is H, halogen or C₁-C₆-alkyl;

R³ is C₁-C₆-alkyl;

20

R⁴ is H, halogen or C₁-C₆-alkyl;

R³ and R⁴ together with the carbon atom to which they are attached form a C₃-C₆-cycloalkyl;

especially preferred R² is halogen or C₁-C₆-alkyl;

25

R³ is C₁-C₆-alkyl;

R⁴ is H or C₁-C₆-alkyl;

more preferred R² is halogen; and

R³ and R⁴ are C₁-C₆-alkyl.

30

Also preferred are the azines of formula (I), wherein

R⁵ is H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl or (C₁-C₆-alkyl)sulfonyl;

particularly preferred H, CN, C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl or (C₁-C₆-alkyl)sulfonyl;

35

especially preferred H, CN, CH₃, CH₂OCH₃, OCH₃, COCH₃ or SO₂CH₃;

more preferred hydrogen.

Also preferred are the azines of formula (I), wherein

40 A is phenyl, which is substituted by two to five substituents

selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;

particularly preferred selected from halogen and CN;
 also particularly preferred selected from the group consisting of F, Cl, CN and CH₃;
 especially preferred selected from the group consisting of F, Cl and CN;

5

particularly preferred phenyl, which is substituted by two to four substituents
 selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl;
 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
 especially preferred selected from halogen and CN;
 also especially preferred selected from the group consisting of F, Cl, CN and CH₃;
 more preferred selected from the group consisting of F, Cl and CN;

10

15

especially preferred phenyl, which is substituted by two substituents
 selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl;
 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
 especially preferred selected from halogen and CN;
 also especially preferred selected from the group consisting of F, Cl, CN and CH₃;
 more preferred selected from the group consisting of F, Cl and CN;

20

25

30

also especially preferred phenyl, which is substituted by three substituents
 selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl;
 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;
 especially preferred selected from halogen and CN;
 also especially preferred selected from the group consisting of F, Cl, CN and CH₃;
 more preferred selected from the group consisting of F, Cl and CN;

35

40

also specially preferred phenyl, which is substituted by four substituents

selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl;

5 particularly preferred selected from the group consisting of halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;

especially preferred selected from halogen and CN;

also especially preferred selected from the group consisting of F, Cl, CN and CH₃;

10 more preferred selected from the group consisting of F, Cl and CN;

R¹ is H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl or (C₁-C₆-alkyl)sulfonyl;

15 particularly preferred H, CN, C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl or (C₁-C₆-alkyl)sulfonyl;

especially preferred H, CN, CH₃, CH₂OCH₃, OCH₃, COCH₃ or SO₂CH₃;

more preferred hydrogen.

R² is H, halogen, C₁-C₆-alkyl or C₁-C₆-haloalkyl;

20 particularly preferred halogen, C₁-C₆-alkyl or C₁-C₆-haloalkyl;

also particularly preferred H, F, CH₃ or CF₃;

R³ and R⁴ are independently of one another H, halogen, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or together with the carbon atom to which they are attached form a moiety selected from the group consisting of C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl and three- to six-

25 membered heterocyclyl,

wherein the C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl or the three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;

30 independently of one another particularly preferred H, halogen, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or

together with the carbon atom to which they are attached form a moiety selected from the group consisting of C₃-C₆-cycloalkyl and C₃-C₆-cycloalkenyl,

35 wherein the C₃-C₆-cycloalkyl or C₃-C₆-cycloalkenyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy;


40 independently of one another especially preferred H, halogen, C₁-C₆-alkyl or C₁-C₆-haloalkyl;

independently of one another more preferred H, halogen or C₁-C₆-alkyl;

and

R^5 is H, CN, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, (C_1 - C_6 -alkyl)carbonyl or (C_1 - C_6 -alkyl)sulfonyl;
 particularly preferred H, CN, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, (C_1 - C_6 -alkyl)carbonyl or (C_1 - C_6 -alkyl)sulfonyl;
 especially preferred H, CN, CH_3 , CH_2OCH_3 , OCH_3 , $COCH_3$ or SO_2CH_3 ;
 more preferred hydrogen.

- 10 Particular preference is given to azines of formula (I.a), which correspond to azines of formula (I) wherein A is (A.1) and R^1 and R^5 are H:



- 15 wherein the variables R^a , R^b , R^c , R^d , R^e , R^2 , R^3 and R^4 have the meanings, in particular the preferred meanings, as defined above;

- special preference is given to the azines of the formulae (I.a.1) to (I.a.1406) of Table A, where the definitions of the variables R^a , R^b , R^c , R^d , R^e , R^2 , R^3 and R^4 are of particular importance for the compounds according to the invention not only in combination with one
 20 another but in each case also on their own:

Table A

No.	R^a	R^b	R^c	R^d	R^e	R^2	R^3	R^4
I.a.1	F	H	H	H	F	CH_3	H	H
I.a.2	Cl	H	H	H	F	CH_3	H	H
I.a.3	Br	H	H	H	F	CH_3	H	H
I.a.4	CN	H	H	H	F	CH_3	H	H
I.a.5	CH_3	H	H	H	F	CH_3	H	H
I.a.6	F	H	H	F	F	CH_3	H	H
I.a.7	Cl	H	H	F	F	CH_3	H	H
I.a.8	F	H	H	Cl	F	CH_3	H	H
I.a.9	Cl	H	H	F	F	CH_3	H	H
I.a.10	CN	H	H	F	F	CH_3	H	H
I.a.11	F	H	H	CN	F	CH_3	H	H
I.a.12	CN	H	H	F	F	CH_3	H	H
I.a.13	F	H	F	H	F	CH_3	H	H

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.14	Cl	H	F	H	F	CH ₃	H	H
I.a.15	CN	H	F	H	F	CH ₃	H	H
I.a.16	F	F	F	H	F	CH ₃	H	H
I.a.17	Cl	F	F	H	F	CH ₃	H	H
I.a.18	F	Cl	F	H	F	CH ₃	H	H
I.a.19	Cl	F	F	H	F	CH ₃	H	H
I.a.20	CN	F	F	H	F	CH ₃	H	H
I.a.21	F	CN	F	H	F	CH ₃	H	H
I.a.22	CN	F	F	H	F	CH ₃	H	H
I.a.23	F	F	H	F	F	CH ₃	H	H
I.a.24	Cl	F	H	F	F	CH ₃	H	H
I.a.25	F	Cl	H	F	F	CH ₃	H	H
I.a.26	CN	F	H	F	F	CH ₃	H	H
I.a.27	F	CN	H	F	F	CH ₃	H	H
I.a.28	F	F	F	F	F	CH ₃	H	H
I.a.29	Cl	F	F	F	F	CH ₃	H	H
I.a.30	F	Cl	F	F	F	CH ₃	H	H
I.a.31	CN	F	F	F	F	CH ₃	H	H
I.a.32	F	CN	F	F	F	CH ₃	H	H
I.a.33	H	F	F	F	F	CH ₃	H	H
I.a.34	F	F	Br	F	F	CH ₃	H	H
I.a.35	F	F	C≡CH	F	F	CH ₃	H	H
I.a.36	CF ₃	Cl	H	H	F	CH ₃	H	H
I.a.37	F	F	I	F	F	CH ₃	H	H
I.a.38	F	H	H	H	F	CH ₃	CH ₃	H
I.a.39	Cl	H	H	H	F	CH ₃	CH ₃	H
I.a.40	Br	H	H	H	F	CH ₃	CH ₃	H
I.a.41	CN	H	H	H	F	CH ₃	CH ₃	H
I.a.42	CH ₃	H	H	H	F	CH ₃	CH ₃	H
I.a.43	F	H	H	F	F	CH ₃	CH ₃	H
I.a.44	Cl	H	H	F	F	CH ₃	CH ₃	H
I.a.45	F	H	H	Cl	F	CH ₃	CH ₃	H
I.a.46	Cl	H	H	F	F	CH ₃	CH ₃	H
I.a.47	CN	H	H	F	F	CH ₃	CH ₃	H
I.a.48	F	H	H	CN	F	CH ₃	CH ₃	H
I.a.49	CN	H	H	F	F	CH ₃	CH ₃	H
I.a.50	F	H	F	H	F	CH ₃	CH ₃	H
I.a.51	Cl	H	F	H	F	CH ₃	CH ₃	H
I.a.52	CN	H	F	H	F	CH ₃	CH ₃	H

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.53	F	F	F	H	F	CH ₃	CH ₃	H
I.a.54	Cl	F	F	H	F	CH ₃	CH ₃	H
I.a.55	F	Cl	F	H	F	CH ₃	CH ₃	H
I.a.56	Cl	F	F	H	F	CH ₃	CH ₃	H
I.a.57	CN	F	F	H	F	CH ₃	CH ₃	H
I.a.58	F	CN	F	H	F	CH ₃	CH ₃	H
I.a.59	CN	F	F	H	F	CH ₃	CH ₃	H
I.a.60	F	F	H	F	F	CH ₃	CH ₃	H
I.a.61	Cl	F	H	F	F	CH ₃	CH ₃	H
I.a.62	F	Cl	H	F	F	CH ₃	CH ₃	H
I.a.63	CN	F	H	F	F	CH ₃	CH ₃	H
I.a.64	F	CN	H	F	F	CH ₃	CH ₃	H
I.a.65	F	F	F	F	F	CH ₃	CH ₃	H
I.a.66	Cl	F	F	F	F	CH ₃	CH ₃	H
I.a.67	F	Cl	F	F	F	CH ₃	CH ₃	H
I.a.68	CN	F	F	F	F	CH ₃	CH ₃	H
I.a.69	F	CN	F	F	F	CH ₃	CH ₃	H
I.a.70	H	F	F	F	F	CH ₃	CH ₃	H
I.a.71	F	F	Br	F	F	CH ₃	CH ₃	H
I.a.72	F	F	C≡CH	F	F	CH ₃	CH ₃	H
I.a.73	CF ₃	Cl	H	H	F	CH ₃	CH ₃	H
I.a.74	F	F	I	F	F	CH ₃	CH ₃	H
I.a.75	F	H	H	H	F	CH ₃	CH ₃	CH ₃
I.a.76	Cl	H	H	H	F	CH ₃	CH ₃	CH ₃
I.a.77	Br	H	H	H	F	CH ₃	CH ₃	CH ₃
I.a.78	CN	H	H	H	F	CH ₃	CH ₃	CH ₃
I.a.79	CH ₃	H	H	H	F	CH ₃	CH ₃	CH ₃
I.a.80	F	H	H	F	F	CH ₃	CH ₃	CH ₃
I.a.81	Cl	H	H	F	F	CH ₃	CH ₃	CH ₃
I.a.82	F	H	H	Cl	F	CH ₃	CH ₃	CH ₃
I.a.83	Cl	H	H	F	F	CH ₃	CH ₃	CH ₃
I.a.84	CN	H	H	F	F	CH ₃	CH ₃	CH ₃
I.a.85	F	H	H	CN	F	CH ₃	CH ₃	CH ₃
I.a.86	CN	H	H	F	F	CH ₃	CH ₃	CH ₃
I.a.87	F	H	F	H	F	CH ₃	CH ₃	CH ₃
I.a.88	Cl	H	F	H	F	CH ₃	CH ₃	CH ₃
I.a.89	CN	H	F	H	F	CH ₃	CH ₃	CH ₃
I.a.90	F	F	F	H	F	CH ₃	CH ₃	CH ₃
I.a.91	Cl	F	F	H	F	CH ₃	CH ₃	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.92	F	Cl	F	H	F	CH ₃	CH ₃	CH ₃
I.a.93	Cl	F	F	H	F	CH ₃	CH ₃	CH ₃
I.a.94	CN	F	F	H	F	CH ₃	CH ₃	CH ₃
I.a.95	F	CN	F	H	F	CH ₃	CH ₃	CH ₃
I.a.96	CN	F	F	H	F	CH ₃	CH ₃	CH ₃
I.a.97	F	F	H	F	F	CH ₃	CH ₃	CH ₃
I.a.98	Cl	F	H	F	F	CH ₃	CH ₃	CH ₃
I.a.99	F	Cl	H	F	F	CH ₃	CH ₃	CH ₃
I.a.100	CN	F	H	F	F	CH ₃	CH ₃	CH ₃
I.a.101	F	CN	H	F	F	CH ₃	CH ₃	CH ₃
I.a.102	F	F	F	F	F	CH ₃	CH ₃	CH ₃
I.a.103	Cl	F	F	F	F	CH ₃	CH ₃	CH ₃
I.a.104	F	Cl	F	F	F	CH ₃	CH ₃	CH ₃
I.a.105	CN	F	F	F	F	CH ₃	CH ₃	CH ₃
I.a.106	F	CN	F	F	F	CH ₃	CH ₃	CH ₃
I.a.107	H	F	F	F	F	CH ₃	CH ₃	CH ₃
I.a.108	F	F	Br	F	F	CH ₃	CH ₃	CH ₃
I.a.109	F	F	C≡CH	F	F	CH ₃	CH ₃	CH ₃
I.a.110	CF ₃	Cl	H	H	F	CH ₃	CH ₃	CH ₃
I.a.111	F	F	I	F	F	CH ₃	CH ₃	CH ₃
I.a.112	F	H	H	H	F	F	F	F
I.a.113	Cl	H	H	H	F	F	F	F
I.a.114	Br	H	H	H	F	F	F	F
I.a.115	CN	H	H	H	F	F	F	F
I.a.116	CH ₃	H	H	H	F	F	F	F
I.a.117	F	H	H	F	F	F	F	F
I.a.118	Cl	H	H	F	F	F	F	F
I.a.119	F	H	H	Cl	F	F	F	F
I.a.120	Cl	H	H	F	F	F	F	F
I.a.121	CN	H	H	F	F	F	F	F
I.a.122	F	H	H	CN	F	F	F	F
I.a.123	CN	H	H	F	F	F	F	F
I.a.124	F	H	F	H	F	F	F	F
I.a.125	Cl	H	F	H	F	F	F	F
I.a.126	CN	H	F	H	F	F	F	F
I.a.127	F	F	F	H	F	F	F	F
I.a.128	Cl	F	F	H	F	F	F	F
I.a.129	F	Cl	F	H	F	F	F	F
I.a.130	Cl	F	F	H	F	F	F	F

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.131	CN	F	F	H	F	F	F	F
I.a.132	F	CN	F	H	F	F	F	F
I.a.133	CN	F	F	H	F	F	F	F
I.a.134	F	F	H	F	F	F	F	F
I.a.135	Cl	F	H	F	F	F	F	F
I.a.136	F	Cl	H	F	F	F	F	F
I.a.137	CN	F	H	F	F	F	F	F
I.a.138	F	CN	H	F	F	F	F	F
I.a.139	F	F	F	F	F	F	F	F
I.a.140	Cl	F	F	F	F	F	F	F
I.a.141	F	Cl	F	F	F	F	F	F
I.a.142	CN	F	F	F	F	F	F	F
I.a.143	F	CN	F	F	F	F	F	F
I.a.144	H	F	F	F	F	F	F	F
I.a.145	F	F	Br	F	F	F	F	F
I.a.146	F	F	C≡CH	F	F	F	F	F
I.a.147	CF ₃	Cl	H	H	F	F	F	F
I.a.148	F	F	I	F	F	F	F	F
I.a.149	F	H	H	H	F	F	CF ₃	F
I.a.150	Cl	H	H	H	F	F	CF ₃	F
I.a.151	Br	H	H	H	F	F	CF ₃	F
I.a.152	CN	H	H	H	F	F	CF ₃	F
I.a.153	CH ₃	H	H	H	F	F	CF ₃	F
I.a.154	F	H	H	F	F	F	CF ₃	F
I.a.155	Cl	H	H	F	F	F	CF ₃	F
I.a.156	F	H	H	Cl	F	F	CF ₃	F
I.a.157	Cl	H	H	F	F	F	CF ₃	F
I.a.158	CN	H	H	F	F	F	CF ₃	F
I.a.159	F	H	H	CN	F	F	CF ₃	F
I.a.160	CN	H	H	F	F	F	CF ₃	F
I.a.161	F	H	F	H	F	F	CF ₃	F
I.a.162	Cl	H	F	H	F	F	CF ₃	F
I.a.163	CN	H	F	H	F	F	CF ₃	F
I.a.164	F	F	F	H	F	F	CF ₃	F
I.a.165	Cl	F	F	H	F	F	CF ₃	F
I.a.166	F	Cl	F	H	F	F	CF ₃	F
I.a.167	Cl	F	F	H	F	F	CF ₃	F
I.a.168	CN	F	F	H	F	F	CF ₃	F
I.a.169	F	CN	F	H	F	F	CF ₃	F

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.170	CN	F	F	H	F	F	CF ₃	F
I.a.171	F	F	H	F	F	F	CF ₃	F
I.a.172	Cl	F	H	F	F	F	CF ₃	F
I.a.173	F	Cl	H	F	F	F	CF ₃	F
I.a.174	CN	F	H	F	F	F	CF ₃	F
I.a.175	F	CN	H	F	F	F	CF ₃	F
I.a.176	F	F	F	F	F	F	CF ₃	F
I.a.177	Cl	F	F	F	F	F	CF ₃	F
I.a.178	F	Cl	F	F	F	F	CF ₃	F
I.a.179	CN	F	F	F	F	F	CF ₃	F
I.a.180	F	CN	F	F	F	F	CF ₃	F
I.a.181	H	F	F	F	F	F	CF ₃	F
I.a.182	F	F	Br	F	F	F	CF ₃	F
I.a.183	F	F	C≡CH	F	F	F	CF ₃	F
I.a.184	CF ₃	Cl	H	H	F	F	CF ₃	F
I.a.185	F	F	I	F	F	F	CF ₃	F
I.a.186	F	H	H	H	F	F	CH ₃	F
I.a.187	Cl	H	H	H	F	F	CH ₃	F
I.a.188	Br	H	H	H	F	F	CH ₃	F
I.a.189	CN	H	H	H	F	F	CH ₃	F
I.a.190	CH ₃	H	H	H	F	F	CH ₃	F
I.a.191	F	H	H	F	F	F	CH ₃	F
I.a.192	Cl	H	H	F	F	F	CH ₃	F
I.a.193	F	H	H	Cl	F	F	CH ₃	F
I.a.194	Cl	H	H	F	F	F	CH ₃	F
I.a.195	CN	H	H	F	F	F	CH ₃	F
I.a.196	F	H	H	CN	F	F	CH ₃	F
I.a.197	CN	H	H	F	F	F	CH ₃	F
I.a.198	F	H	F	H	F	F	CH ₃	F
I.a.199	Cl	H	F	H	F	F	CH ₃	F
I.a.200	CN	H	F	H	F	F	CH ₃	F
I.a.201	F	F	F	H	F	F	CH ₃	F
I.a.202	Cl	F	F	H	F	F	CH ₃	F
I.a.203	F	Cl	F	H	F	F	CH ₃	F
I.a.204	Cl	F	F	H	F	F	CH ₃	F
I.a.205	CN	F	F	H	F	F	CH ₃	F
I.a.206	F	CN	F	H	F	F	CH ₃	F
I.a.207	CN	F	F	H	F	F	CH ₃	F
I.a.208	F	F	H	F	F	F	CH ₃	F

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.209	Cl	F	H	F	F	F	CH ₃	F
I.a.210	F	Cl	H	F	F	F	CH ₃	F
I.a.211	CN	F	H	F	F	F	CH ₃	F
I.a.212	F	CN	H	F	F	F	CH ₃	F
I.a.213	F	F	F	F	F	F	CH ₃	F
I.a.214	Cl	F	F	F	F	F	CH ₃	F
I.a.215	F	Cl	F	F	F	F	CH ₃	F
I.a.216	CN	F	F	F	F	F	CH ₃	F
I.a.217	F	CN	F	F	F	F	CH ₃	F
I.a.218	H	F	F	F	F	F	CH ₃	F
I.a.219	F	F	Br	F	F	F	CH ₃	F
I.a.220	F	F	C≡CH	F	F	F	CH ₃	F
I.a.221	CF ₃	Cl	H	H	F	F	CH ₃	F
I.a.222	F	F	I	F	F	F	CH ₃	F
I.a.223	F	H	H	H	F	F	CH ₃	H
I.a.224	Cl	H	H	H	F	F	CH ₃	H
I.a.225	Br	H	H	H	F	F	CH ₃	H
I.a.226	CN	H	H	H	F	F	CH ₃	H
I.a.227	CH ₃	H	H	H	F	F	CH ₃	H
I.a.228	F	H	H	F	F	F	CH ₃	H
I.a.229	Cl	H	H	F	F	F	CH ₃	H
I.a.230	F	H	H	Cl	F	F	CH ₃	H
I.a.231	Cl	H	H	F	F	F	CH ₃	H
I.a.232	CN	H	H	F	F	F	CH ₃	H
I.a.233	F	H	H	CN	F	F	CH ₃	H
I.a.234	CN	H	H	F	F	F	CH ₃	H
I.a.235	F	H	F	H	F	F	CH ₃	H
I.a.236	Cl	H	F	H	F	F	CH ₃	H
I.a.237	CN	H	F	H	F	F	CH ₃	H
I.a.238	F	F	F	H	F	F	CH ₃	H
I.a.239	Cl	F	F	H	F	F	CH ₃	H
I.a.240	F	Cl	F	H	F	F	CH ₃	H
I.a.241	Cl	F	F	H	F	F	CH ₃	H
I.a.242	CN	F	F	H	F	F	CH ₃	H
I.a.243	F	CN	F	H	F	F	CH ₃	H
I.a.244	CN	F	F	H	F	F	CH ₃	H
I.a.245	F	F	H	F	F	F	CH ₃	H
I.a.246	Cl	F	H	F	F	F	CH ₃	H
I.a.247	F	Cl	H	F	F	F	CH ₃	H

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.248	CN	F	H	F	F	F	CH ₃	H
I.a.249	F	CN	H	F	F	F	CH ₃	H
I.a.250	F	F	F	F	F	F	CH ₃	H
I.a.251	Cl	F	F	F	F	F	CH ₃	H
I.a.252	F	Cl	F	F	F	F	CH ₃	H
I.a.253	CN	F	F	F	F	F	CH ₃	H
I.a.254	F	CN	F	F	F	F	CH ₃	H
I.a.255	H	F	F	F	F	F	CH ₃	H
I.a.256	F	F	Br	F	F	F	CH ₃	H
I.a.257	F	F	C≡CH	F	F	F	CH ₃	H
I.a.258	CF ₃	Cl	H	H	F	F	CH ₃	H
I.a.259	F	F	I	F	F	F	CH ₃	H
I.a.260	F	H	H	H	F	F	CH ₃	CH ₃
I.a.261	Cl	H	H	H	F	F	CH ₃	CH ₃
I.a.262	Br	H	H	H	F	F	CH ₃	CH ₃
I.a.263	CN	H	H	H	F	F	CH ₃	CH ₃
I.a.264	CH ₃	H	H	H	F	F	CH ₃	CH ₃
I.a.265	F	H	H	F	F	F	CH ₃	CH ₃
I.a.266	Cl	H	H	F	F	F	CH ₃	CH ₃
I.a.267	F	H	H	Cl	F	F	CH ₃	CH ₃
I.a.268	Cl	H	H	F	F	F	CH ₃	CH ₃
I.a.269	CN	H	H	F	F	F	CH ₃	CH ₃
I.a.270	F	H	H	CN	F	F	CH ₃	CH ₃
I.a.271	CN	H	H	F	F	F	CH ₃	CH ₃
I.a.272	F	H	F	H	F	F	CH ₃	CH ₃
I.a.273	Cl	H	F	H	F	F	CH ₃	CH ₃
I.a.274	CN	H	F	H	F	F	CH ₃	CH ₃
I.a.275	F	F	F	H	F	F	CH ₃	CH ₃
I.a.276	Cl	F	F	H	F	F	CH ₃	CH ₃
I.a.277	F	Cl	F	H	F	F	CH ₃	CH ₃
I.a.278	Cl	F	F	H	F	F	CH ₃	CH ₃
I.a.279	CN	F	F	H	F	F	CH ₃	CH ₃
I.a.280	F	CN	F	H	F	F	CH ₃	CH ₃
I.a.281	CN	F	F	H	F	F	CH ₃	CH ₃
I.a.282	F	F	H	F	F	F	CH ₃	CH ₃
I.a.283	Cl	F	H	F	F	F	CH ₃	CH ₃
I.a.284	F	Cl	H	F	F	F	CH ₃	CH ₃
I.a.285	CN	F	H	F	F	F	CH ₃	CH ₃
I.a.286	F	CN	H	F	F	F	CH ₃	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.287	F	F	F	F	F	F	CH ₃	CH ₃
I.a.288	Cl	F	F	F	F	F	CH ₃	CH ₃
I.a.289	F	Cl	F	F	F	F	CH ₃	CH ₃
I.a.290	CN	F	F	F	F	F	CH ₃	CH ₃
I.a.291	F	CN	F	F	F	F	CH ₃	CH ₃
I.a.292	H	F	F	F	F	F	CH ₃	CH ₃
I.a.293	F	F	Br	F	F	F	CH ₃	CH ₃
I.a.294	F	F	C≡CH	F	F	F	CH ₃	CH ₃
I.a.295	CF ₃	Cl	H	H	F	F	CH ₃	CH ₃
I.a.296	F	F	I	F	F	F	CH ₃	CH ₃
I.a.297	F	H	H	H	F	Cl	CH ₃	CH ₃
I.a.298	Cl	H	H	H	F	Cl	CH ₃	CH ₃
I.a.299	Br	H	H	H	F	Cl	CH ₃	CH ₃
I.a.300	CN	H	H	H	F	Cl	CH ₃	CH ₃
I.a.301	CH ₃	H	H	H	F	Cl	CH ₃	CH ₃
I.a.302	F	H	H	F	F	Cl	CH ₃	CH ₃
I.a.303	Cl	H	H	F	F	Cl	CH ₃	CH ₃
I.a.304	F	H	H	Cl	F	Cl	CH ₃	CH ₃
I.a.305	Cl	H	H	F	F	Cl	CH ₃	CH ₃
I.a.306	CN	H	H	F	F	Cl	CH ₃	CH ₃
I.a.307	F	H	H	CN	F	Cl	CH ₃	CH ₃
I.a.308	CN	H	H	F	F	Cl	CH ₃	CH ₃
I.a.309	F	H	F	H	F	Cl	CH ₃	CH ₃
I.a.310	Cl	H	F	H	F	Cl	CH ₃	CH ₃
I.a.311	CN	H	F	H	F	Cl	CH ₃	CH ₃
I.a.312	F	F	F	H	F	Cl	CH ₃	CH ₃
I.a.313	Cl	F	F	H	F	Cl	CH ₃	CH ₃
I.a.314	F	Cl	F	H	F	Cl	CH ₃	CH ₃
I.a.315	Cl	F	F	H	F	Cl	CH ₃	CH ₃
I.a.316	CN	F	F	H	F	Cl	CH ₃	CH ₃
I.a.317	F	CN	F	H	F	Cl	CH ₃	CH ₃
I.a.318	CN	F	F	H	F	Cl	CH ₃	CH ₃
I.a.319	F	F	H	F	F	Cl	CH ₃	CH ₃
I.a.320	Cl	F	H	F	F	Cl	CH ₃	CH ₃
I.a.321	F	Cl	H	F	F	Cl	CH ₃	CH ₃
I.a.322	CN	F	H	F	F	Cl	CH ₃	CH ₃
I.a.323	F	CN	H	F	F	Cl	CH ₃	CH ₃
I.a.324	F	F	F	F	F	Cl	CH ₃	CH ₃
I.a.325	Cl	F	F	F	F	Cl	CH ₃	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.326	F	Cl	F	F	F	Cl	CH ₃	CH ₃
I.a.327	CN	F	F	F	F	Cl	CH ₃	CH ₃
I.a.328	F	CN	F	F	F	Cl	CH ₃	CH ₃
I.a.329	H	F	F	F	F	Cl	CH ₃	CH ₃
I.a.330	F	F	Br	F	F	Cl	CH ₃	CH ₃
I.a.331	F	F	C≡CH	F	F	Cl	CH ₃	CH ₃
I.a.332	CF ₃	Cl	H	H	F	Cl	CH ₃	CH ₃
I.a.333	F	F	I	F	F	Cl	CH ₃	CH ₃
I.a.334	F	H	H	H	F	F	C ₂ H ₅	CH ₃
I.a.335	Cl	H	H	H	F	F	C ₂ H ₅	CH ₃
I.a.336	Br	H	H	H	F	F	C ₂ H ₅	CH ₃
I.a.337	CN	H	H	H	F	F	C ₂ H ₅	CH ₃
I.a.338	CH ₃	H	H	H	F	F	C ₂ H ₅	CH ₃
I.a.339	F	H	H	F	F	F	C ₂ H ₅	CH ₃
I.a.340	Cl	H	H	F	F	F	C ₂ H ₅	CH ₃
I.a.341	F	H	H	Cl	F	F	C ₂ H ₅	CH ₃
I.a.342	Cl	H	H	F	F	F	C ₂ H ₅	CH ₃
I.a.343	CN	H	H	F	F	F	C ₂ H ₅	CH ₃
I.a.344	F	H	H	CN	F	F	C ₂ H ₅	CH ₃
I.a.345	CN	H	H	F	F	F	C ₂ H ₅	CH ₃
I.a.346	F	H	F	H	F	F	C ₂ H ₅	CH ₃
I.a.347	Cl	H	F	H	F	F	C ₂ H ₅	CH ₃
I.a.348	CN	H	F	H	F	F	C ₂ H ₅	CH ₃
I.a.349	F	F	F	H	F	F	C ₂ H ₅	CH ₃
I.a.350	Cl	F	F	H	F	F	C ₂ H ₅	CH ₃
I.a.351	F	Cl	F	H	F	F	C ₂ H ₅	CH ₃
I.a.352	Cl	F	F	H	F	F	C ₂ H ₅	CH ₃
I.a.353	CN	F	F	H	F	F	C ₂ H ₅	CH ₃
I.a.354	F	CN	F	H	F	F	C ₂ H ₅	CH ₃
I.a.355	CN	F	F	H	F	F	C ₂ H ₅	CH ₃
I.a.356	F	F	H	F	F	F	C ₂ H ₅	CH ₃
I.a.357	Cl	F	H	F	F	F	C ₂ H ₅	CH ₃
I.a.358	F	Cl	H	F	F	F	C ₂ H ₅	CH ₃
I.a.359	CN	F	H	F	F	F	C ₂ H ₅	CH ₃
I.a.360	F	CN	H	F	F	F	C ₂ H ₅	CH ₃
I.a.361	F	F	F	F	F	F	C ₂ H ₅	CH ₃
I.a.362	Cl	F	F	F	F	F	C ₂ H ₅	CH ₃
I.a.363	F	Cl	F	F	F	F	C ₂ H ₅	CH ₃
I.a.364	CN	F	F	F	F	F	C ₂ H ₅	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.365	F	CN	F	F	F	F	C ₂ H ₅	CH ₃
I.a.366	H	F	F	F	F	F	C ₂ H ₅	CH ₃
I.a.367	F	F	Br	F	F	F	C ₂ H ₅	CH ₃
I.a.368	F	F	C≡CH	F	F	F	C ₂ H ₅	CH ₃
I.a.369	CF ₃	Cl	H	H	F	F	C ₂ H ₅	CH ₃
I.a.370	F	F	I	F	F	F	C ₂ H ₅	CH ₃
I.a.371	F	H	H	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.372	Cl	H	H	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.373	Br	H	H	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.374	CN	H	H	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.375	CH ₃	H	H	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.376	F	H	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.377	Cl	H	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.378	F	H	H	Cl	F	F	C ₂ H ₅	C ₂ H ₅
I.a.379	Cl	H	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.380	CN	H	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.381	F	H	H	CN	F	F	C ₂ H ₅	C ₂ H ₅
I.a.382	CN	H	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.383	F	H	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.384	Cl	H	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.385	CN	H	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.386	F	F	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.387	Cl	F	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.388	F	Cl	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.389	Cl	F	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.390	CN	F	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.391	F	CN	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.392	CN	F	F	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.393	F	F	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.394	Cl	F	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.395	F	Cl	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.396	CN	F	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.397	F	CN	H	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.398	F	F	F	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.399	Cl	F	F	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.400	F	Cl	F	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.401	CN	F	F	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.402	F	CN	F	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.403	H	F	F	F	F	F	C ₂ H ₅	C ₂ H ₅

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.404	F	F	Br	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.405	F	F	C≡CH	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.406	CF ₃	Cl	H	H	F	F	C ₂ H ₅	C ₂ H ₅
I.a.407	F	F	I	F	F	F	C ₂ H ₅	C ₂ H ₅
I.a.408	F	H	H	H	F	H	-(CH ₂) ₂ -	
I.a.409	Cl	H	H	H	F	H	-(CH ₂) ₂ -	
I.a.410	Br	H	H	H	F	H	-(CH ₂) ₂ -	
I.a.411	CN	H	H	H	F	H	-(CH ₂) ₂ -	
I.a.412	CH ₃	H	H	H	F	H	-(CH ₂) ₂ -	
I.a.413	F	H	H	F	F	H	-(CH ₂) ₂ -	
I.a.414	Cl	H	H	F	F	H	-(CH ₂) ₂ -	
I.a.415	F	H	H	Cl	F	H	-(CH ₂) ₂ -	
I.a.416	Cl	H	H	F	F	H	-(CH ₂) ₂ -	
I.a.417	CN	H	H	F	F	H	-(CH ₂) ₂ -	
I.a.418	F	H	H	CN	F	H	-(CH ₂) ₂ -	
I.a.419	CN	H	H	F	F	H	-(CH ₂) ₂ -	
I.a.420	F	H	F	H	F	H	-(CH ₂) ₂ -	
I.a.421	Cl	H	F	H	F	H	-(CH ₂) ₂ -	
I.a.422	CN	H	F	H	F	H	-(CH ₂) ₂ -	
I.a.423	F	F	F	H	F	H	-(CH ₂) ₂ -	
I.a.424	Cl	F	F	H	F	H	-(CH ₂) ₂ -	
I.a.425	F	Cl	F	H	F	H	-(CH ₂) ₂ -	
I.a.426	Cl	F	F	H	F	H	-(CH ₂) ₂ -	
I.a.427	CN	F	F	H	F	H	-(CH ₂) ₂ -	
I.a.428	F	CN	F	H	F	H	-(CH ₂) ₂ -	
I.a.429	CN	F	F	H	F	H	-(CH ₂) ₂ -	
I.a.430	F	F	H	F	F	H	-(CH ₂) ₂ -	
I.a.431	Cl	F	H	F	F	H	-(CH ₂) ₂ -	
I.a.432	F	Cl	H	F	F	H	-(CH ₂) ₂ -	
I.a.433	CN	F	H	F	F	H	-(CH ₂) ₂ -	
I.a.434	F	CN	H	F	F	H	-(CH ₂) ₂ -	
I.a.435	F	F	F	F	F	H	-(CH ₂) ₂ -	
I.a.436	Cl	F	F	F	F	H	-(CH ₂) ₂ -	
I.a.437	F	Cl	F	F	F	H	-(CH ₂) ₂ -	
I.a.438	CN	F	F	F	F	H	-(CH ₂) ₂ -	
I.a.439	F	CN	F	F	F	H	-(CH ₂) ₂ -	
I.a.440	H	F	F	F	F	H	-(CH ₂) ₂ -	
I.a.441	F	F	Br	F	F	H	-(CH ₂) ₂ -	
I.a.442	F	F	C≡CH	F	F	H	-(CH ₂) ₂ -	

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.443	CF ₃	Cl	H	H	F	H	-(CH ₂) ₂ -	
I.a.444	F	F	I	F	F	H	-(CH ₂) ₂ -	
I.a.445	F	H	H	H	F	H	-(CH ₂) ₃ -	
I.a.446	Cl	H	H	H	F	H	-(CH ₂) ₃ -	
I.a.447	Br	H	H	H	F	H	-(CH ₂) ₃ -	
I.a.448	CN	H	H	H	F	H	-(CH ₂) ₃ -	
I.a.449	CH ₃	H	H	H	F	H	-(CH ₂) ₃ -	
I.a.450	F	H	H	F	F	H	-(CH ₂) ₃ -	
I.a.451	Cl	H	H	F	F	H	-(CH ₂) ₃ -	
I.a.452	F	H	H	Cl	F	H	-(CH ₂) ₃ -	
I.a.453	Cl	H	H	F	F	H	-(CH ₂) ₃ -	
I.a.454	CN	H	H	F	F	H	-(CH ₂) ₃ -	
I.a.455	F	H	H	CN	F	H	-(CH ₂) ₃ -	
I.a.456	CN	H	H	F	F	H	-(CH ₂) ₃ -	
I.a.457	F	H	F	H	F	H	-(CH ₂) ₃ -	
I.a.458	Cl	H	F	H	F	H	-(CH ₂) ₃ -	
I.a.459	CN	H	F	H	F	H	-(CH ₂) ₃ -	
I.a.460	F	F	F	H	F	H	-(CH ₂) ₃ -	
I.a.461	Cl	F	F	H	F	H	-(CH ₂) ₃ -	
I.a.462	F	Cl	F	H	F	H	-(CH ₂) ₃ -	
I.a.463	Cl	F	F	H	F	H	-(CH ₂) ₃ -	
I.a.464	CN	F	F	H	F	H	-(CH ₂) ₃ -	
I.a.465	F	CN	F	H	F	H	-(CH ₂) ₃ -	
I.a.466	CN	F	F	H	F	H	-(CH ₂) ₃ -	
I.a.467	F	F	H	F	F	H	-(CH ₂) ₃ -	
I.a.468	Cl	F	H	F	F	H	-(CH ₂) ₃ -	
I.a.469	F	Cl	H	F	F	H	-(CH ₂) ₃ -	
I.a.470	CN	F	H	F	F	H	-(CH ₂) ₃ -	
I.a.471	F	CN	H	F	F	H	-(CH ₂) ₃ -	
I.a.472	F	F	F	F	F	H	-(CH ₂) ₃ -	
I.a.473	Cl	F	F	F	F	H	-(CH ₂) ₃ -	
I.a.474	F	Cl	F	F	F	H	-(CH ₂) ₃ -	
I.a.475	CN	F	F	F	F	H	-(CH ₂) ₃ -	
I.a.476	F	CN	F	F	F	H	-(CH ₂) ₃ -	
I.a.477	H	F	F	F	F	H	-(CH ₂) ₃ -	
I.a.478	F	F	Br	F	F	H	-(CH ₂) ₃ -	
I.a.479	F	F	C≡CH	F	F	H	-(CH ₂) ₃ -	
I.a.480	CF ₃	Cl	H	H	F	H	-(CH ₂) ₃ -	
I.a.481	F	F	I	F	F	H	-(CH ₂) ₃ -	

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.482	F	H	H	H	F	H	-(CH ₂) ₄ -	
I.a.483	Cl	H	H	H	F	H	-(CH ₂) ₄ -	
I.a.484	Br	H	H	H	F	H	-(CH ₂) ₄ -	
I.a.485	CN	H	H	H	F	H	-(CH ₂) ₄ -	
I.a.486	CH ₃	H	H	H	F	H	-(CH ₂) ₄ -	
I.a.487	F	H	H	F	F	H	-(CH ₂) ₄ -	
I.a.488	Cl	H	H	F	F	H	-(CH ₂) ₄ -	
I.a.489	F	H	H	Cl	F	H	-(CH ₂) ₄ -	
I.a.490	Cl	H	H	F	F	H	-(CH ₂) ₄ -	
I.a.491	CN	H	H	F	F	H	-(CH ₂) ₄ -	
I.a.492	F	H	H	CN	F	H	-(CH ₂) ₄ -	
I.a.493	CN	H	H	F	F	H	-(CH ₂) ₄ -	
I.a.494	F	H	F	H	F	H	-(CH ₂) ₄ -	
I.a.495	Cl	H	F	H	F	H	-(CH ₂) ₄ -	
I.a.496	CN	H	F	H	F	H	-(CH ₂) ₄ -	
I.a.497	F	F	F	H	F	H	-(CH ₂) ₄ -	
I.a.498	Cl	F	F	H	F	H	-(CH ₂) ₄ -	
I.a.499	F	Cl	F	H	F	H	-(CH ₂) ₄ -	
I.a.500	Cl	F	F	H	F	H	-(CH ₂) ₄ -	
I.a.501	CN	F	F	H	F	H	-(CH ₂) ₄ -	
I.a.502	F	CN	F	H	F	H	-(CH ₂) ₄ -	
I.a.503	CN	F	F	H	F	H	-(CH ₂) ₄ -	
I.a.504	F	F	H	F	F	H	-(CH ₂) ₄ -	
I.a.505	Cl	F	H	F	F	H	-(CH ₂) ₄ -	
I.a.506	F	Cl	H	F	F	H	-(CH ₂) ₄ -	
I.a.507	CN	F	H	F	F	H	-(CH ₂) ₄ -	
I.a.508	F	CN	H	F	F	H	-(CH ₂) ₄ -	
I.a.509	F	F	F	F	F	H	-(CH ₂) ₄ -	
I.a.510	Cl	F	F	F	F	H	-(CH ₂) ₄ -	
I.a.511	F	Cl	F	F	F	H	-(CH ₂) ₄ -	
I.a.512	CN	F	F	F	F	H	-(CH ₂) ₄ -	
I.a.513	F	CN	F	F	F	H	-(CH ₂) ₄ -	
I.a.514	H	F	F	F	F	H	-(CH ₂) ₄ -	
I.a.515	F	F	Br	F	F	H	-(CH ₂) ₄ -	
I.a.516	F	F	C≡CH	F	F	H	-(CH ₂) ₄ -	
I.a.517	CF ₃	Cl	H	H	F	H	-(CH ₂) ₄ -	
I.a.518	F	F	I	F	F	H	-(CH ₂) ₄ -	
I.a.519	F	H	H	H	F	H	-(CH ₂) ₅ -	
I.a.520	Cl	H	H	H	F	H	-(CH ₂) ₅ -	

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.521	Br	H	H	H	F	H	-(CH ₂) ₅ -	
I.a.522	CN	H	H	H	F	H	-(CH ₂) ₅ -	
I.a.523	CH ₃	H	H	H	F	H	-(CH ₂) ₅ -	
I.a.524	F	H	H	F	F	H	-(CH ₂) ₅ -	
I.a.525	Cl	H	H	F	F	H	-(CH ₂) ₅ -	
I.a.526	F	H	H	Cl	F	H	-(CH ₂) ₅ -	
I.a.527	Cl	H	H	F	F	H	-(CH ₂) ₅ -	
I.a.528	CN	H	H	F	F	H	-(CH ₂) ₅ -	
I.a.529	F	H	H	CN	F	H	-(CH ₂) ₅ -	
I.a.530	CN	H	H	F	F	H	-(CH ₂) ₅ -	
I.a.531	F	H	F	H	F	H	-(CH ₂) ₅ -	
I.a.532	Cl	H	F	H	F	H	-(CH ₂) ₅ -	
I.a.533	CN	H	F	H	F	H	-(CH ₂) ₅ -	
I.a.534	F	F	F	H	F	H	-(CH ₂) ₅ -	
I.a.535	Cl	F	F	H	F	H	-(CH ₂) ₅ -	
I.a.536	F	Cl	F	H	F	H	-(CH ₂) ₅ -	
I.a.537	Cl	F	F	H	F	H	-(CH ₂) ₅ -	
I.a.538	CN	F	F	H	F	H	-(CH ₂) ₅ -	
I.a.539	F	CN	F	H	F	H	-(CH ₂) ₅ -	
I.a.540	CN	F	F	H	F	H	-(CH ₂) ₅ -	
I.a.541	F	F	H	F	F	H	-(CH ₂) ₅ -	
I.a.542	Cl	F	H	F	F	H	-(CH ₂) ₅ -	
I.a.543	F	Cl	H	F	F	H	-(CH ₂) ₅ -	
I.a.544	CN	F	H	F	F	H	-(CH ₂) ₅ -	
I.a.545	F	CN	H	F	F	H	-(CH ₂) ₅ -	
I.a.546	F	F	F	F	F	H	-(CH ₂) ₅ -	
I.a.547	Cl	F	F	F	F	H	-(CH ₂) ₅ -	
I.a.548	F	Cl	F	F	F	H	-(CH ₂) ₅ -	
I.a.549	CN	F	F	F	F	H	-(CH ₂) ₅ -	
I.a.550	F	CN	F	F	F	H	-(CH ₂) ₅ -	
I.a.551	H	F	F	F	F	H	-(CH ₂) ₅ -	
I.a.552	F	F	Br	F	F	H	-(CH ₂) ₅ -	
I.a.553	F	F	C≡CH	F	F	H	-(CH ₂) ₅ -	
I.a.554	CF ₃	Cl	H	H	F	H	-(CH ₂) ₅ -	
I.a.555	F	F	I	F	F	H	-(CH ₂) ₅ -	
I.a.556	F	H	H	H	F	CH ₃	-(CH ₂) ₂ -	
I.a.557	Cl	H	H	H	F	CH ₃	-(CH ₂) ₂ -	
I.a.558	Br	H	H	H	F	CH ₃	-(CH ₂) ₂ -	
I.a.559	CN	H	H	H	F	CH ₃	-(CH ₂) ₂ -	

I.a.560	CH ₃	H	H	H	F	CH ₃	-(CH ₂) ₂ -
I.a.561	F	H	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.562	Cl	H	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.563	F	H	H	Cl	F	CH ₃	-(CH ₂) ₂ -
I.a.564	Cl	H	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.565	CN	H	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.566	F	H	H	CN	F	CH ₃	-(CH ₂) ₂ -
I.a.567	CN	H	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.568	F	H	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.569	Cl	H	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.570	CN	H	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.571	F	F	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.572	Cl	F	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.573	F	Cl	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.574	Cl	F	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.575	CN	F	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.576	F	CN	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.577	CN	F	F	H	F	CH ₃	-(CH ₂) ₂ -
I.a.578	F	F	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.579	Cl	F	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.580	F	Cl	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.581	CN	F	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.582	F	CN	H	F	F	CH ₃	-(CH ₂) ₂ -
I.a.583	F	F	F	F	F	CH ₃	-(CH ₂) ₂ -
I.a.584	Cl	F	F	F	F	CH ₃	-(CH ₂) ₂ -
I.a.585	F	Cl	F	F	F	CH ₃	-(CH ₂) ₂ -
I.a.586	CN	F	F	F	F	CH ₃	-(CH ₂) ₂ -
I.a.587	F	CN	F	F	F	CH ₃	-(CH ₂) ₂ -
I.a.588	H	F	F	F	F	CH ₃	-(CH ₂) ₂ -
I.a.589	F	F	Br	F	F	CH ₃	-(CH ₂) ₂ -
I.a.590	F	F	C≡CH	F	F	CH ₃	-(CH ₂) ₂ -
I.a.591	CF ₃	Cl	H	H	F	CH ₃	-(CH ₂) ₂ -
I.a.592	F	F	I	F	F	CH ₃	-(CH ₂) ₂ -
I.a.593	F	H	H	H	F	CH ₃	-(CH ₂) ₃ -
I.a.594	Cl	H	H	H	F	CH ₃	-(CH ₂) ₃ -
I.a.595	Br	H	H	H	F	CH ₃	-(CH ₂) ₃ -
I.a.596	CN	H	H	H	F	CH ₃	-(CH ₂) ₃ -
I.a.597	CH ₃	H	H	H	F	CH ₃	-(CH ₂) ₃ -
I.a.598	F	H	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.599	Cl	H	H	F	F	CH ₃	-(CH ₂) ₃ -

I.a.600	F	H	H	Cl	F	CH ₃	-(CH ₂) ₃ -
I.a.601	Cl	H	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.602	CN	H	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.603	F	H	H	CN	F	CH ₃	-(CH ₂) ₃ -
I.a.604	CN	H	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.605	F	H	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.606	Cl	H	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.607	CN	H	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.608	F	F	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.609	Cl	F	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.610	F	Cl	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.611	Cl	F	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.612	CN	F	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.613	F	CN	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.614	CN	F	F	H	F	CH ₃	-(CH ₂) ₃ -
I.a.615	F	F	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.616	Cl	F	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.617	F	Cl	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.618	CN	F	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.619	F	CN	H	F	F	CH ₃	-(CH ₂) ₃ -
I.a.620	F	F	F	F	F	CH ₃	-(CH ₂) ₃ -
I.a.621	Cl	F	F	F	F	CH ₃	-(CH ₂) ₃ -
I.a.622	F	Cl	F	F	F	CH ₃	-(CH ₂) ₃ -
I.a.623	CN	F	F	F	F	CH ₃	-(CH ₂) ₃ -
I.a.624	F	CN	F	F	F	CH ₃	-(CH ₂) ₃ -
I.a.625	H	F	F	F	F	CH ₃	-(CH ₂) ₃ -
I.a.626	F	F	Br	F	F	CH ₃	-(CH ₂) ₃ -
I.a.627	F	F	C≡CH	F	F	CH ₃	-(CH ₂) ₃ -
I.a.628	CF ₃	Cl	H	H	F	CH ₃	-(CH ₂) ₃ -
I.a.629	F	F	I	F	F	CH ₃	-(CH ₂) ₃ -
I.a.630	F	H	H	H	F	CH ₃	-(CH ₂) ₄ -
I.a.631	Cl	H	H	H	F	CH ₃	-(CH ₂) ₄ -
I.a.632	Br	H	H	H	F	CH ₃	-(CH ₂) ₄ -
I.a.633	CN	H	H	H	F	CH ₃	-(CH ₂) ₄ -
I.a.634	CH ₃	H	H	H	F	CH ₃	-(CH ₂) ₄ -
I.a.635	F	H	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.636	Cl	H	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.637	F	H	H	Cl	F	CH ₃	-(CH ₂) ₄ -
I.a.638	Cl	H	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.639	CN	H	H	F	F	CH ₃	-(CH ₂) ₄ -

I.a.640	F	H	H	CN	F	CH ₃	-(CH ₂) ₄ -
I.a.641	CN	H	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.642	F	H	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.643	Cl	H	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.644	CN	H	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.645	F	F	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.646	Cl	F	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.647	F	Cl	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.648	Cl	F	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.649	CN	F	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.650	F	CN	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.651	CN	F	F	H	F	CH ₃	-(CH ₂) ₄ -
I.a.652	F	F	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.653	Cl	F	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.654	F	Cl	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.655	CN	F	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.656	F	CN	H	F	F	CH ₃	-(CH ₂) ₄ -
I.a.657	F	F	F	F	F	CH ₃	-(CH ₂) ₄ -
I.a.658	Cl	F	F	F	F	CH ₃	-(CH ₂) ₄ -
I.a.659	F	Cl	F	F	F	CH ₃	-(CH ₂) ₄ -
I.a.660	CN	F	F	F	F	CH ₃	-(CH ₂) ₄ -
I.a.661	F	CN	F	F	F	CH ₃	-(CH ₂) ₄ -
I.a.662	H	F	F	F	F	CH ₃	-(CH ₂) ₄ -
I.a.663	F	F	Br	F	F	CH ₃	-(CH ₂) ₄ -
I.a.664	F	F	C≡CH	F	F	CH ₃	-(CH ₂) ₄ -
I.a.665	CF ₃	Cl	H	H	F	CH ₃	-(CH ₂) ₄ -
I.a.666	F	F	I	F	F	CH ₃	-(CH ₂) ₄ -
I.a.667	F	H	H	H	F	CH ₃	-(CH ₂) ₅ -
I.a.668	Cl	H	H	H	F	CH ₃	-(CH ₂) ₅ -
I.a.669	Br	H	H	H	F	CH ₃	-(CH ₂) ₅ -
I.a.670	CN	H	H	H	F	CH ₃	-(CH ₂) ₅ -
I.a.671	CH ₃	H	H	H	F	CH ₃	-(CH ₂) ₅ -
I.a.672	F	H	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.673	Cl	H	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.674	F	H	H	Cl	F	CH ₃	-(CH ₂) ₅ -
I.a.675	Cl	H	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.676	CN	H	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.677	F	H	H	CN	F	CH ₃	-(CH ₂) ₅ -
I.a.678	CN	H	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.679	F	H	F	H	F	CH ₃	-(CH ₂) ₅ -

I.a.680	Cl	H	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.681	CN	H	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.682	F	F	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.683	Cl	F	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.684	F	Cl	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.685	Cl	F	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.686	CN	F	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.687	F	CN	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.688	CN	F	F	H	F	CH ₃	-(CH ₂) ₅ -
I.a.689	F	F	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.690	Cl	F	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.691	F	Cl	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.692	CN	F	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.693	F	CN	H	F	F	CH ₃	-(CH ₂) ₅ -
I.a.694	F	F	F	F	F	CH ₃	-(CH ₂) ₅ -
I.a.695	Cl	F	F	F	F	CH ₃	-(CH ₂) ₅ -
I.a.696	F	Cl	F	F	F	CH ₃	-(CH ₂) ₅ -
I.a.697	CN	F	F	F	F	CH ₃	-(CH ₂) ₅ -
I.a.698	F	CN	F	F	F	CH ₃	-(CH ₂) ₅ -
I.a.699	H	F	F	F	F	CH ₃	-(CH ₂) ₅ -
I.a.700	F	F	Br	F	F	CH ₃	-(CH ₂) ₅ -
I.a.701	F	F	C≡CH	F	F	CH ₃	-(CH ₂) ₅ -
I.a.702	CF ₃	Cl	H	H	F	CH ₃	-(CH ₂) ₅ -
I.a.703	F	F	I	F	F	CH ₃	-(CH ₂) ₅ -
I.a.704	F	H	H	H	F	F	-(CH ₂) ₂ -
I.a.705	Cl	H	H	H	F	F	-(CH ₂) ₂ -
I.a.706	Br	H	H	H	F	F	-(CH ₂) ₂ -
I.a.707	CN	H	H	H	F	F	-(CH ₂) ₂ -
I.a.708	CH ₃	H	H	H	F	F	-(CH ₂) ₂ -
I.a.709	F	H	H	F	F	F	-(CH ₂) ₂ -
I.a.710	Cl	H	H	F	F	F	-(CH ₂) ₂ -
I.a.711	F	H	H	Cl	F	F	-(CH ₂) ₂ -
I.a.712	Cl	H	H	F	F	F	-(CH ₂) ₂ -
I.a.713	CN	H	H	F	F	F	-(CH ₂) ₂ -
I.a.714	F	H	H	CN	F	F	-(CH ₂) ₂ -
I.a.715	CN	H	H	F	F	F	-(CH ₂) ₂ -
I.a.716	F	H	F	H	F	F	-(CH ₂) ₂ -
I.a.717	Cl	H	F	H	F	F	-(CH ₂) ₂ -
I.a.718	CN	H	F	H	F	F	-(CH ₂) ₂ -
I.a.719	F	F	F	H	F	F	-(CH ₂) ₂ -

I.a.720	Cl	F	F	H	F	F	-(CH ₂) ₂ -
I.a.721	F	Cl	F	H	F	F	-(CH ₂) ₂ -
I.a.722	Cl	F	F	H	F	F	-(CH ₂) ₂ -
I.a.723	CN	F	F	H	F	F	-(CH ₂) ₂ -
I.a.724	F	CN	F	H	F	F	-(CH ₂) ₂ -
I.a.725	CN	F	F	H	F	F	-(CH ₂) ₂ -
I.a.726	F	F	H	F	F	F	-(CH ₂) ₂ -
I.a.727	Cl	F	H	F	F	F	-(CH ₂) ₂ -
I.a.728	F	Cl	H	F	F	F	-(CH ₂) ₂ -
I.a.729	CN	F	H	F	F	F	-(CH ₂) ₂ -
I.a.730	F	CN	H	F	F	F	-(CH ₂) ₂ -
I.a.731	F	F	F	F	F	F	-(CH ₂) ₂ -
I.a.732	Cl	F	F	F	F	F	-(CH ₂) ₂ -
I.a.733	F	Cl	F	F	F	F	-(CH ₂) ₂ -
I.a.734	CN	F	F	F	F	F	-(CH ₂) ₂ -
I.a.735	F	CN	F	F	F	F	-(CH ₂) ₂ -
I.a.736	H	F	F	F	F	F	-(CH ₂) ₂ -
I.a.737	F	F	Br	F	F	F	-(CH ₂) ₂ -
I.a.738	F	F	C≡CH	F	F	F	-(CH ₂) ₂ -
I.a.739	CF ₃	Cl	H	H	F	F	-(CH ₂) ₂ -
I.a.740	F	F	I	F	F	F	-(CH ₂) ₂ -
I.a.741	F	H	H	H	F	F	-(CH ₂) ₃ -
I.a.742	Cl	H	H	H	F	F	-(CH ₂) ₃ -
I.a.743	Br	H	H	H	F	F	-(CH ₂) ₃ -
I.a.744	CN	H	H	H	F	F	-(CH ₂) ₃ -
I.a.745	CH ₃	H	H	H	F	F	-(CH ₂) ₃ -
I.a.746	F	H	H	F	F	F	-(CH ₂) ₃ -
I.a.747	Cl	H	H	F	F	F	-(CH ₂) ₃ -
I.a.748	F	H	H	Cl	F	F	-(CH ₂) ₃ -
I.a.749	Cl	H	H	F	F	F	-(CH ₂) ₃ -
I.a.750	CN	H	H	F	F	F	-(CH ₂) ₃ -
I.a.751	F	H	H	CN	F	F	-(CH ₂) ₃ -
I.a.752	CN	H	H	F	F	F	-(CH ₂) ₃ -
I.a.753	F	H	F	H	F	F	-(CH ₂) ₃ -
I.a.754	Cl	H	F	H	F	F	-(CH ₂) ₃ -
I.a.755	CN	H	F	H	F	F	-(CH ₂) ₃ -
I.a.756	F	F	F	H	F	F	-(CH ₂) ₃ -
I.a.757	Cl	F	F	H	F	F	-(CH ₂) ₃ -
I.a.758	F	Cl	F	H	F	F	-(CH ₂) ₃ -
I.a.759	Cl	F	F	H	F	F	-(CH ₂) ₃ -

I.a.760	CN	F	F	H	F	F	-(CH ₂) ₃ -
I.a.761	F	CN	F	H	F	F	-(CH ₂) ₃ -
I.a.762	CN	F	F	H	F	F	-(CH ₂) ₃ -
I.a.763	F	F	H	F	F	F	-(CH ₂) ₃ -
I.a.764	Cl	F	H	F	F	F	-(CH ₂) ₃ -
I.a.765	F	Cl	H	F	F	F	-(CH ₂) ₃ -
I.a.766	CN	F	H	F	F	F	-(CH ₂) ₃ -
I.a.767	F	CN	H	F	F	F	-(CH ₂) ₃ -
I.a.768	F	F	F	F	F	F	-(CH ₂) ₃ -
I.a.769	Cl	F	F	F	F	F	-(CH ₂) ₃ -
I.a.770	F	Cl	F	F	F	F	-(CH ₂) ₃ -
I.a.771	CN	F	F	F	F	F	-(CH ₂) ₃ -
I.a.772	F	CN	F	F	F	F	-(CH ₂) ₃ -
I.a.773	H	F	F	F	F	F	-(CH ₂) ₃ -
I.a.774	F	F	Br	F	F	F	-(CH ₂) ₃ -
I.a.775	F	F	C≡CH	F	F	F	-(CH ₂) ₃ -
I.a.776	CF ₃	Cl	H	H	F	F	-(CH ₂) ₃ -
I.a.777	F	F	I	F	F	F	-(CH ₂) ₃ -
I.a.778	F	H	H	H	F	F	-(CH ₂) ₄ -
I.a.779	Cl	H	H	H	F	F	-(CH ₂) ₄ -
I.a.780	Br	H	H	H	F	F	-(CH ₂) ₄ -
I.a.781	CN	H	H	H	F	F	-(CH ₂) ₄ -
I.a.782	CH ₃	H	H	H	F	F	-(CH ₂) ₄ -
I.a.783	F	H	H	F	F	F	-(CH ₂) ₄ -
I.a.784	Cl	H	H	F	F	F	-(CH ₂) ₄ -
I.a.785	F	H	H	Cl	F	F	-(CH ₂) ₄ -
I.a.786	Cl	H	H	F	F	F	-(CH ₂) ₄ -
I.a.787	CN	H	H	F	F	F	-(CH ₂) ₄ -
I.a.788	F	H	H	CN	F	F	-(CH ₂) ₄ -
I.a.789	CN	H	H	F	F	F	-(CH ₂) ₄ -
I.a.790	F	H	F	H	F	F	-(CH ₂) ₄ -
I.a.791	Cl	H	F	H	F	F	-(CH ₂) ₄ -
I.a.792	CN	H	F	H	F	F	-(CH ₂) ₄ -
I.a.793	F	F	F	H	F	F	-(CH ₂) ₄ -
I.a.794	Cl	F	F	H	F	F	-(CH ₂) ₄ -
I.a.795	F	Cl	F	H	F	F	-(CH ₂) ₄ -
I.a.796	Cl	F	F	H	F	F	-(CH ₂) ₄ -
I.a.797	CN	F	F	H	F	F	-(CH ₂) ₄ -
I.a.798	F	CN	F	H	F	F	-(CH ₂) ₄ -
I.a.799	CN	F	F	H	F	F	-(CH ₂) ₄ -

I.a.800	F	F	H	F	F	F	-(CH ₂) ₄ -
I.a.801	Cl	F	H	F	F	F	-(CH ₂) ₄ -
I.a.802	F	Cl	H	F	F	F	-(CH ₂) ₄ -
I.a.803	CN	F	H	F	F	F	-(CH ₂) ₄ -
I.a.804	F	CN	H	F	F	F	-(CH ₂) ₄ -
I.a.805	F	F	F	F	F	F	-(CH ₂) ₄ -
I.a.806	Cl	F	F	F	F	F	-(CH ₂) ₄ -
I.a.807	F	Cl	F	F	F	F	-(CH ₂) ₄ -
I.a.808	CN	F	F	F	F	F	-(CH ₂) ₄ -
I.a.809	F	CN	F	F	F	F	-(CH ₂) ₄ -
I.a.810	H	F	F	F	F	F	-(CH ₂) ₄ -
I.a.811	F	F	Br	F	F	F	-(CH ₂) ₄ -
I.a.812	F	F	C≡CH	F	F	F	-(CH ₂) ₄ -
I.a.813	CF ₃	Cl	H	H	F	F	-(CH ₂) ₄ -
I.a.814	F	F	I	F	F	F	-(CH ₂) ₄ -
I.a.815	F	H	H	H	F	F	-(CH ₂) ₅ -
I.a.816	Cl	H	H	H	F	F	-(CH ₂) ₅ -
I.a.817	Br	H	H	H	F	F	-(CH ₂) ₅ -
I.a.818	CN	H	H	H	F	F	-(CH ₂) ₅ -
I.a.819	CH ₃	H	H	H	F	F	-(CH ₂) ₅ -
I.a.820	F	H	H	F	F	F	-(CH ₂) ₅ -
I.a.821	Cl	H	H	F	F	F	-(CH ₂) ₅ -
I.a.822	F	H	H	Cl	F	F	-(CH ₂) ₅ -
I.a.823	Cl	H	H	F	F	F	-(CH ₂) ₅ -
I.a.824	CN	H	H	F	F	F	-(CH ₂) ₅ -
I.a.825	F	H	H	CN	F	F	-(CH ₂) ₅ -
I.a.826	CN	H	H	F	F	F	-(CH ₂) ₅ -
I.a.827	F	H	F	H	F	F	-(CH ₂) ₅ -
I.a.828	Cl	H	F	H	F	F	-(CH ₂) ₅ -
I.a.829	CN	H	F	H	F	F	-(CH ₂) ₅ -
I.a.830	F	F	F	H	F	F	-(CH ₂) ₅ -
I.a.831	Cl	F	F	H	F	F	-(CH ₂) ₅ -
I.a.832	F	Cl	F	H	F	F	-(CH ₂) ₅ -
I.a.833	Cl	F	F	H	F	F	-(CH ₂) ₅ -
I.a.834	CN	F	F	H	F	F	-(CH ₂) ₅ -
I.a.835	F	CN	F	H	F	F	-(CH ₂) ₅ -
I.a.836	CN	F	F	H	F	F	-(CH ₂) ₅ -
I.a.837	F	F	H	F	F	F	-(CH ₂) ₅ -
I.a.838	Cl	F	H	F	F	F	-(CH ₂) ₅ -
I.a.839	F	Cl	H	F	F	F	-(CH ₂) ₅ -

I.a.840	CN	F	H	F	F	F	-(CH ₂) ₅ -
I.a.841	F	CN	H	F	F	F	-(CH ₂) ₅ -
I.a.842	F	F	F	F	F	F	-(CH ₂) ₅ -
I.a.843	Cl	F	F	F	F	F	-(CH ₂) ₅ -
I.a.844	F	Cl	F	F	F	F	-(CH ₂) ₅ -
I.a.845	CN	F	F	F	F	F	-(CH ₂) ₅ -
I.a.846	F	CN	F	F	F	F	-(CH ₂) ₅ -
I.a.847	H	F	F	F	F	F	-(CH ₂) ₅ -
I.a.848	F	F	Br	F	F	F	-(CH ₂) ₅ -
I.a.849	F	F	C≡CH	F	F	F	-(CH ₂) ₅ -
I.a.850	CF ₃	Cl	H	H	F	F	-(CH ₂) ₅ -
I.a.851	F	F	I	F	F	F	-(CH ₂) ₅ -
I.a.852	F	H	H	H	F	Cl	-(CH ₂) ₂ -
I.a.853	Cl	H	H	H	F	Cl	-(CH ₂) ₂ -
I.a.854	Br	H	H	H	F	Cl	-(CH ₂) ₂ -
I.a.855	CN	H	H	H	F	Cl	-(CH ₂) ₂ -
I.a.856	CH ₃	H	H	H	F	Cl	-(CH ₂) ₂ -
I.a.857	F	H	H	F	F	Cl	-(CH ₂) ₂ -
I.a.858	Cl	H	H	F	F	Cl	-(CH ₂) ₂ -
I.a.859	F	H	H	Cl	F	Cl	-(CH ₂) ₂ -
I.a.860	Cl	H	H	F	F	Cl	-(CH ₂) ₂ -
I.a.861	CN	H	H	F	F	Cl	-(CH ₂) ₂ -
I.a.862	F	H	H	CN	F	Cl	-(CH ₂) ₂ -
I.a.863	CN	H	H	F	F	Cl	-(CH ₂) ₂ -
I.a.864	F	H	F	H	F	Cl	-(CH ₂) ₂ -
I.a.865	Cl	H	F	H	F	Cl	-(CH ₂) ₂ -
I.a.866	CN	H	F	H	F	Cl	-(CH ₂) ₂ -
I.a.867	F	F	F	H	F	Cl	-(CH ₂) ₂ -
I.a.868	Cl	F	F	H	F	Cl	-(CH ₂) ₂ -
I.a.869	F	Cl	F	H	F	Cl	-(CH ₂) ₂ -
I.a.870	Cl	F	F	H	F	Cl	-(CH ₂) ₂ -
I.a.871	CN	F	F	H	F	Cl	-(CH ₂) ₂ -
I.a.872	F	CN	F	H	F	Cl	-(CH ₂) ₂ -
I.a.873	CN	F	F	H	F	Cl	-(CH ₂) ₂ -
I.a.874	F	F	H	F	F	Cl	-(CH ₂) ₂ -
I.a.875	Cl	F	H	F	F	Cl	-(CH ₂) ₂ -
I.a.876	F	Cl	H	F	F	Cl	-(CH ₂) ₂ -
I.a.877	CN	F	H	F	F	Cl	-(CH ₂) ₂ -
I.a.878	F	CN	H	F	F	Cl	-(CH ₂) ₂ -
I.a.879	F	F	F	F	F	Cl	-(CH ₂) ₂ -

I.a.880	Cl	F	F	F	F	Cl	-(CH ₂) ₂ -
I.a.881	F	Cl	F	F	F	Cl	-(CH ₂) ₂ -
I.a.882	CN	F	F	F	F	Cl	-(CH ₂) ₂ -
I.a.883	F	CN	F	F	F	Cl	-(CH ₂) ₂ -
I.a.884	H	F	F	F	F	Cl	-(CH ₂) ₂ -
I.a.885	F	F	Br	F	F	Cl	-(CH ₂) ₂ -
I.a.886	F	F	C≡CH	F	F	Cl	-(CH ₂) ₂ -
I.a.887	CF ₃	Cl	H	H	F	Cl	-(CH ₂) ₂ -
I.a.888	F	F	I	F	F	Cl	-(CH ₂) ₂ -
I.a.889	F	H	H	H	F	Cl	-(CH ₂) ₃ -
I.a.890	Cl	H	H	H	F	Cl	-(CH ₂) ₃ -
I.a.891	Br	H	H	H	F	Cl	-(CH ₂) ₃ -
I.a.892	CN	H	H	H	F	Cl	-(CH ₂) ₃ -
I.a.893	CH ₃	H	H	H	F	Cl	-(CH ₂) ₃ -
I.a.894	F	H	H	F	F	Cl	-(CH ₂) ₃ -
I.a.895	Cl	H	H	F	F	Cl	-(CH ₂) ₃ -
I.a.896	F	H	H	Cl	F	Cl	-(CH ₂) ₃ -
I.a.897	Cl	H	H	F	F	Cl	-(CH ₂) ₃ -
I.a.898	CN	H	H	F	F	Cl	-(CH ₂) ₃ -
I.a.899	F	H	H	CN	F	Cl	-(CH ₂) ₃ -
I.a.900	CN	H	H	F	F	Cl	-(CH ₂) ₃ -
I.a.901	F	H	F	H	F	Cl	-(CH ₂) ₃ -
I.a.902	Cl	H	F	H	F	Cl	-(CH ₂) ₃ -
I.a.903	CN	H	F	H	F	Cl	-(CH ₂) ₃ -
I.a.904	F	F	F	H	F	Cl	-(CH ₂) ₃ -
I.a.905	Cl	F	F	H	F	Cl	-(CH ₂) ₃ -
I.a.906	F	Cl	F	H	F	Cl	-(CH ₂) ₃ -
I.a.907	Cl	F	F	H	F	Cl	-(CH ₂) ₃ -
I.a.908	CN	F	F	H	F	Cl	-(CH ₂) ₃ -
I.a.909	F	CN	F	H	F	Cl	-(CH ₂) ₃ -
I.a.910	CN	F	F	H	F	Cl	-(CH ₂) ₃ -
I.a.911	F	F	H	F	F	Cl	-(CH ₂) ₃ -
I.a.912	Cl	F	H	F	F	Cl	-(CH ₂) ₃ -
I.a.913	F	Cl	H	F	F	Cl	-(CH ₂) ₃ -
I.a.914	CN	F	H	F	F	Cl	-(CH ₂) ₃ -
I.a.915	F	CN	H	F	F	Cl	-(CH ₂) ₃ -
I.a.916	F	F	F	F	F	Cl	-(CH ₂) ₃ -
I.a.917	Cl	F	F	F	F	Cl	-(CH ₂) ₃ -
I.a.918	F	Cl	F	F	F	Cl	-(CH ₂) ₃ -
I.a.919	CN	F	F	F	F	Cl	-(CH ₂) ₃ -

I.a.920	F	CN	F	F	F	Cl	-(CH ₂) ₃ -
I.a.921	H	F	F	F	F	Cl	-(CH ₂) ₃ -
I.a.922	F	F	Br	F	F	Cl	-(CH ₂) ₃ -
I.a.923	F	F	C≡CH	F	F	Cl	-(CH ₂) ₃ -
I.a.924	CF ₃	Cl	H	H	F	Cl	-(CH ₂) ₃ -
I.a.925	F	F	I	F	F	Cl	-(CH ₂) ₃ -
I.a.926	F	H	H	H	F	Cl	-(CH ₂) ₄ -
I.a.927	Cl	H	H	H	F	Cl	-(CH ₂) ₄ -
I.a.928	Br	H	H	H	F	Cl	-(CH ₂) ₄ -
I.a.929	CN	H	H	H	F	Cl	-(CH ₂) ₄ -
I.a.930	CH ₃	H	H	H	F	Cl	-(CH ₂) ₄ -
I.a.931	F	H	H	F	F	Cl	-(CH ₂) ₄ -
I.a.932	Cl	H	H	F	F	Cl	-(CH ₂) ₄ -
I.a.933	F	H	H	Cl	F	Cl	-(CH ₂) ₄ -
I.a.934	Cl	H	H	F	F	Cl	-(CH ₂) ₄ -
I.a.935	CN	H	H	F	F	Cl	-(CH ₂) ₄ -
I.a.936	F	H	H	CN	F	Cl	-(CH ₂) ₄ -
I.a.937	CN	H	H	F	F	Cl	-(CH ₂) ₄ -
I.a.938	F	H	F	H	F	Cl	-(CH ₂) ₄ -
I.a.939	Cl	H	F	H	F	Cl	-(CH ₂) ₄ -
I.a.940	CN	H	F	H	F	Cl	-(CH ₂) ₄ -
I.a.941	F	F	F	H	F	Cl	-(CH ₂) ₄ -
I.a.942	Cl	F	F	H	F	Cl	-(CH ₂) ₄ -
I.a.943	F	Cl	F	H	F	Cl	-(CH ₂) ₄ -
I.a.944	Cl	F	F	H	F	Cl	-(CH ₂) ₄ -
I.a.945	CN	F	F	H	F	Cl	-(CH ₂) ₄ -
I.a.946	F	CN	F	H	F	Cl	-(CH ₂) ₄ -
I.a.947	CN	F	F	H	F	Cl	-(CH ₂) ₄ -
I.a.948	F	F	H	F	F	Cl	-(CH ₂) ₄ -
I.a.949	Cl	F	H	F	F	Cl	-(CH ₂) ₄ -
I.a.950	F	Cl	H	F	F	Cl	-(CH ₂) ₄ -
I.a.951	CN	F	H	F	F	Cl	-(CH ₂) ₄ -
I.a.952	F	CN	H	F	F	Cl	-(CH ₂) ₄ -
I.a.953	F	F	F	F	F	Cl	-(CH ₂) ₄ -
I.a.954	Cl	F	F	F	F	Cl	-(CH ₂) ₄ -
I.a.955	F	Cl	F	F	F	Cl	-(CH ₂) ₄ -
I.a.956	CN	F	F	F	F	Cl	-(CH ₂) ₄ -
I.a.957	F	CN	F	F	F	Cl	-(CH ₂) ₄ -
I.a.958	H	F	F	F	F	Cl	-(CH ₂) ₄ -
I.a.959	F	F	Br	F	F	Cl	-(CH ₂) ₄ -

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.960	F	F	C≡CH	F	F	Cl	-(CH ₂) ₄ -	
I.a.961	CF ₃	Cl	H	H	F	Cl	-(CH ₂) ₄ -	
I.a.962	F	F	I	F	F	Cl	-(CH ₂) ₄ -	
I.a.963	F	H	H	H	F	Cl	-(CH ₂) ₅ -	
I.a.964	Cl	H	H	H	F	Cl	-(CH ₂) ₅ -	
I.a.965	Br	H	H	H	F	Cl	-(CH ₂) ₅ -	
I.a.966	CN	H	H	H	F	Cl	-(CH ₂) ₅ -	
I.a.967	CH ₃	H	H	H	F	Cl	-(CH ₂) ₅ -	
I.a.968	F	H	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.969	Cl	H	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.970	F	H	H	Cl	F	Cl	-(CH ₂) ₅ -	
I.a.971	Cl	H	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.972	CN	H	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.973	F	H	H	CN	F	Cl	-(CH ₂) ₅ -	
I.a.974	CN	H	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.975	F	H	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.976	Cl	H	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.977	CN	H	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.978	F	F	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.979	Cl	F	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.980	F	Cl	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.981	Cl	F	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.982	CN	F	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.983	F	CN	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.984	CN	F	F	H	F	Cl	-(CH ₂) ₅ -	
I.a.985	F	F	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.986	Cl	F	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.987	F	Cl	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.988	CN	F	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.989	F	CN	H	F	F	Cl	-(CH ₂) ₅ -	
I.a.990	F	F	F	F	F	Cl	-(CH ₂) ₅ -	
I.a.991	Cl	F	F	F	F	Cl	-(CH ₂) ₅ -	
I.a.992	F	Cl	F	F	F	Cl	-(CH ₂) ₅ -	
I.a.993	CN	F	F	F	F	Cl	-(CH ₂) ₅ -	
I.a.994	F	CN	F	F	F	Cl	-(CH ₂) ₅ -	
I.a.995	H	F	F	F	F	Cl	-(CH ₂) ₅ -	
I.a.996	F	F	Br	F	F	Cl	-(CH ₂) ₅ -	
I.a.997	F	F	C≡CH	F	F	Cl	-(CH ₂) ₅ -	
I.a.998	CF ₃	Cl	H	H	F	Cl	-(CH ₂) ₅ -	

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.999	F	F	I	F	F	Cl	-(CH ₂) ₅ -	
I.a.1000	F	H	H	H	F	C ₂ H ₅	CH ₃	H
I.a.1001	Cl	H	H	H	F	C ₂ H ₅	CH ₃	H
I.a.1002	Br	H	H	H	F	C ₂ H ₅	CH ₃	H
I.a.1003	CN	H	H	H	F	C ₂ H ₅	CH ₃	H
I.a.1004	CH ₃	H	H	H	F	C ₂ H ₅	CH ₃	H
I.a.1005	F	H	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1006	Cl	H	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1007	F	H	H	Cl	F	C ₂ H ₅	CH ₃	H
I.a.1008	Cl	H	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1009	CN	H	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1010	F	H	H	CN	F	C ₂ H ₅	CH ₃	H
I.a.1011	CN	H	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1012	F	H	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1013	Cl	H	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1014	CN	H	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1015	F	F	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1016	Cl	F	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1017	F	Cl	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1018	Cl	F	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1019	CN	F	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1020	F	CN	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1021	CN	F	F	H	F	C ₂ H ₅	CH ₃	H
I.a.1022	F	F	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1023	Cl	F	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1024	F	Cl	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1025	CN	F	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1026	F	CN	H	F	F	C ₂ H ₅	CH ₃	H
I.a.1027	F	F	F	F	F	C ₂ H ₅	CH ₃	H
I.a.1028	Cl	F	F	F	F	C ₂ H ₅	CH ₃	H
I.a.1029	F	Cl	F	F	F	C ₂ H ₅	CH ₃	H
I.a.1030	CN	F	F	F	F	C ₂ H ₅	CH ₃	H
I.a.1031	F	CN	F	F	F	C ₂ H ₅	CH ₃	H
I.a.1032	H	F	F	F	F	C ₂ H ₅	CH ₃	H
I.a.1033	F	F	Br	F	F	C ₂ H ₅	CH ₃	H
I.a.1034	F	F	C≡CH	F	F	C ₂ H ₅	CH ₃	H
I.a.1035	CF ₃	Cl	H	H	F	C ₂ H ₅	CH ₃	H
I.a.1036	F	F	I	F	F	C ₂ H ₅	CH ₃	H
I.a.1037	F	H	H	H	F	C ₂ H ₅	C ₂ H ₅	H

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1038	Cl	H	H	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1039	Br	H	H	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1040	CN	H	H	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1041	CH ₃	H	H	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1042	F	H	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1043	Cl	H	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1044	F	H	H	Cl	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1045	Cl	H	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1046	CN	H	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1047	F	H	H	CN	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1048	CN	H	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1049	F	H	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1050	Cl	H	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1051	CN	H	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1052	F	F	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1053	Cl	F	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1054	F	Cl	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1055	Cl	F	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1056	CN	F	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1057	F	CN	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1058	CN	F	F	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1059	F	F	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1060	Cl	F	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1061	F	Cl	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1062	CN	F	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1063	F	CN	H	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1064	F	F	F	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1065	Cl	F	F	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1066	F	Cl	F	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1067	CN	F	F	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1068	F	CN	F	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1069	H	F	F	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1070	F	F	Br	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1071	F	F	C≡CH	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1072	CF ₃	Cl	H	H	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1073	F	F	I	F	F	C ₂ H ₅	C ₂ H ₅	H
I.a.1074	F	H	H	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1075	Cl	H	H	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1076	Br	H	H	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1077	CN	H	H	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1078	CH ₃	H	H	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1079	F	H	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1080	Cl	H	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1081	F	H	H	Cl	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1082	Cl	H	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1083	CN	H	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1084	F	H	H	CN	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1085	CN	H	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1086	F	H	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1087	Cl	H	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1088	CN	H	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1089	F	F	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1090	Cl	F	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1091	F	Cl	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1092	Cl	F	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1093	CN	F	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1094	F	CN	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1095	CN	F	F	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1096	F	F	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1097	Cl	F	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1098	F	Cl	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1099	CN	F	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1100	F	CN	H	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1101	F	F	F	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1102	Cl	F	F	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1103	F	Cl	F	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1104	CN	F	F	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1105	F	CN	F	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1106	H	F	F	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1107	F	F	Br	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1108	F	F	C≡CH	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1109	CF ₃	Cl	H	H	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1110	F	F	I	F	F	C ₂ H ₅	C ₂ H ₅	CH ₃
I.a.1111	F	H	H	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1112	Cl	H	H	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1113	Br	H	H	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1114	CN	H	H	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1115	CH ₃	H	H	H	F	C ₂ H ₅	CH ₃	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1116	F	H	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1117	Cl	H	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1118	F	H	H	Cl	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1119	Cl	H	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1120	CN	H	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1121	F	H	H	CN	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1122	CN	H	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1123	F	H	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1124	Cl	H	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1125	CN	H	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1126	F	F	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1127	Cl	F	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1128	F	Cl	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1129	Cl	F	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1130	CN	F	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1131	F	CN	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1132	CN	F	F	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1133	F	F	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1134	Cl	F	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1135	F	Cl	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1136	CN	F	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1137	F	CN	H	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1138	F	F	F	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1139	Cl	F	F	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1140	F	Cl	F	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1141	CN	F	F	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1142	F	CN	F	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1143	H	F	F	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1144	F	F	Br	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1145	F	F	C≡CH	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1146	CF ₃	Cl	H	H	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1147	F	F	I	F	F	C ₂ H ₅	CH ₃	CH ₃
I.a.1148	F	H	H	H	F	Cl	CH ₃	H
I.a.1149	Cl	H	H	H	F	Cl	CH ₃	H
I.a.1150	Br	H	H	H	F	Cl	CH ₃	H
I.a.1151	CN	H	H	H	F	Cl	CH ₃	H
I.a.1152	CH ₃	H	H	H	F	Cl	CH ₃	H
I.a.1153	F	H	H	F	F	Cl	CH ₃	H
I.a.1154	Cl	H	H	F	F	Cl	CH ₃	H

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1155	F	H	H	Cl	F	Cl	CH ₃	H
I.a.1156	Cl	H	H	F	F	Cl	CH ₃	H
I.a.1157	CN	H	H	F	F	Cl	CH ₃	H
I.a.1158	F	H	H	CN	F	Cl	CH ₃	H
I.a.1159	CN	H	H	F	F	Cl	CH ₃	H
I.a.1160	F	H	F	H	F	Cl	CH ₃	H
I.a.1161	Cl	H	F	H	F	Cl	CH ₃	H
I.a.1162	CN	H	F	H	F	Cl	CH ₃	H
I.a.1163	F	F	F	H	F	Cl	CH ₃	H
I.a.1164	Cl	F	F	H	F	Cl	CH ₃	H
I.a.1165	F	Cl	F	H	F	Cl	CH ₃	H
I.a.1166	Cl	F	F	H	F	Cl	CH ₃	H
I.a.1167	CN	F	F	H	F	Cl	CH ₃	H
I.a.1168	F	CN	F	H	F	Cl	CH ₃	H
I.a.1169	CN	F	F	H	F	Cl	CH ₃	H
I.a.1170	F	F	H	F	F	Cl	CH ₃	H
I.a.1171	Cl	F	H	F	F	Cl	CH ₃	H
I.a.1172	F	Cl	H	F	F	Cl	CH ₃	H
I.a.1173	CN	F	H	F	F	Cl	CH ₃	H
I.a.1174	F	CN	H	F	F	Cl	CH ₃	H
I.a.1175	F	F	F	F	F	Cl	CH ₃	H
I.a.1176	Cl	F	F	F	F	Cl	CH ₃	H
I.a.1177	F	Cl	F	F	F	Cl	CH ₃	H
I.a.1178	CN	F	F	F	F	Cl	CH ₃	H
I.a.1179	F	CN	F	F	F	Cl	CH ₃	H
I.a.1180	H	F	F	F	F	Cl	CH ₃	H
I.a.1181	F	F	Br	F	F	Cl	CH ₃	H
I.a.1182	F	F	C≡CH	F	F	Cl	CH ₃	H
I.a.1183	CF ₃	Cl	H	H	F	Cl	CH ₃	H
I.a.1184	F	F	I	F	F	Cl	CH ₃	H
I.a.1185	F	H	H	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1186	Cl	H	H	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1187	Br	H	H	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1188	CN	H	H	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1189	CH ₃	H	H	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1190	F	H	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1191	Cl	H	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1192	F	H	H	Cl	F	CH ₂ Cl	Cl	CH ₃
I.a.1193	Cl	H	H	F	F	CH ₂ Cl	Cl	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1194	CN	H	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1195	F	H	H	CN	F	CH ₂ Cl	Cl	CH ₃
I.a.1196	CN	H	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1197	F	H	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1198	Cl	H	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1199	CN	H	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1200	F	F	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1201	Cl	F	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1202	F	Cl	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1203	Cl	F	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1204	CN	F	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1205	F	CN	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1206	CN	F	F	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1207	F	F	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1208	Cl	F	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1209	F	Cl	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1210	CN	F	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1211	F	CN	H	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1212	F	F	F	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1213	Cl	F	F	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1214	F	Cl	F	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1215	CN	F	F	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1216	F	CN	F	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1217	H	F	F	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1218	F	F	Br	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1219	F	F	C≡CH	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1220	CF ₃	Cl	H	H	F	CH ₂ Cl	Cl	CH ₃
I.a.1221	F	F	I	F	F	CH ₂ Cl	Cl	CH ₃
I.a.1222	F	H	H	H	F	CN	CH ₃	CH ₃
I.a.1223	Cl	H	H	H	F	CN	CH ₃	CH ₃
I.a.1224	Br	H	H	H	F	CN	CH ₃	CH ₃
I.a.1225	CN	H	H	H	F	CN	CH ₃	CH ₃
I.a.1226	CH ₃	H	H	H	F	CN	CH ₃	CH ₃
I.a.1227	F	H	H	F	F	CN	CH ₃	CH ₃
I.a.1228	Cl	H	H	F	F	CN	CH ₃	CH ₃
I.a.1229	F	H	H	Cl	F	CN	CH ₃	CH ₃
I.a.1230	Cl	H	H	F	F	CN	CH ₃	CH ₃
I.a.1231	CN	H	H	F	F	CN	CH ₃	CH ₃
I.a.1232	F	H	H	CN	F	CN	CH ₃	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1233	CN	H	H	F	F	CN	CH ₃	CH ₃
I.a.1234	F	H	F	H	F	CN	CH ₃	CH ₃
I.a.1235	Cl	H	F	H	F	CN	CH ₃	CH ₃
I.a.1236	CN	H	F	H	F	CN	CH ₃	CH ₃
I.a.1237	F	F	F	H	F	CN	CH ₃	CH ₃
I.a.1238	Cl	F	F	H	F	CN	CH ₃	CH ₃
I.a.1239	F	Cl	F	H	F	CN	CH ₃	CH ₃
I.a.1240	Cl	F	F	H	F	CN	CH ₃	CH ₃
I.a.1241	CN	F	F	H	F	CN	CH ₃	CH ₃
I.a.1242	F	CN	F	H	F	CN	CH ₃	CH ₃
I.a.1243	CN	F	F	H	F	CN	CH ₃	CH ₃
I.a.1244	F	F	H	F	F	CN	CH ₃	CH ₃
I.a.1245	Cl	F	H	F	F	CN	CH ₃	CH ₃
I.a.1246	F	Cl	H	F	F	CN	CH ₃	CH ₃
I.a.1247	CN	F	H	F	F	CN	CH ₃	CH ₃
I.a.1248	F	CN	H	F	F	CN	CH ₃	CH ₃
I.a.1249	F	F	F	F	F	CN	CH ₃	CH ₃
I.a.1250	Cl	F	F	F	F	CN	CH ₃	CH ₃
I.a.1251	F	Cl	F	F	F	CN	CH ₃	CH ₃
I.a.1252	CN	F	F	F	F	CN	CH ₃	CH ₃
I.a.1253	F	CN	F	F	F	CN	CH ₃	CH ₃
I.a.1254	H	F	F	F	F	CN	CH ₃	CH ₃
I.a.1255	F	F	Br	F	F	CN	CH ₃	CH ₃
I.a.1256	F	F	C≡CH	F	F	CN	CH ₃	CH ₃
I.a.1257	CF ₃	Cl	H	H	F	CN	CH ₃	CH ₃
I.a.1258	F	F	I	F	F	CN	CH ₃	CH ₃
I.a.1259	F	H	H	H	F	OCH ₃	H	H
I.a.1260	Cl	H	H	H	F	OCH ₃	H	H
I.a.1261	Br	H	H	H	F	OCH ₃	H	H
I.a.1262	CN	H	H	H	F	OCH ₃	H	H
I.a.1263	CH ₃	H	H	H	F	OCH ₃	H	H
I.a.1264	F	H	H	F	F	OCH ₃	H	H
I.a.1265	Cl	H	H	F	F	OCH ₃	H	H
I.a.1266	F	H	H	Cl	F	OCH ₃	H	H
I.a.1267	Cl	H	H	F	F	OCH ₃	H	H
I.a.1268	CN	H	H	F	F	OCH ₃	H	H
I.a.1269	F	H	H	CN	F	OCH ₃	H	H
I.a.1270	CN	H	H	F	F	OCH ₃	H	H
I.a.1271	F	H	F	H	F	OCH ₃	H	H


No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1272	Cl	H	F	H	F	OCH ₃	H	H
I.a.1273	CN	H	F	H	F	OCH ₃	H	H
I.a.1274	F	F	F	H	F	OCH ₃	H	H
I.a.1275	Cl	F	F	H	F	OCH ₃	H	H
I.a.1276	F	Cl	F	H	F	OCH ₃	H	H
I.a.1277	Cl	F	F	H	F	OCH ₃	H	H
I.a.1278	CN	F	F	H	F	OCH ₃	H	H
I.a.1279	F	CN	F	H	F	OCH ₃	H	H
I.a.1280	CN	F	F	H	F	OCH ₃	H	H
I.a.1281	F	F	H	F	F	OCH ₃	H	H
I.a.1282	Cl	F	H	F	F	OCH ₃	H	H
I.a.1283	F	Cl	H	F	F	OCH ₃	H	H
I.a.1284	CN	F	H	F	F	OCH ₃	H	H
I.a.1285	F	CN	H	F	F	OCH ₃	H	H
I.a.1286	F	F	F	F	F	OCH ₃	H	H
I.a.1287	Cl	F	F	F	F	OCH ₃	H	H
I.a.1288	F	Cl	F	F	F	OCH ₃	H	H
I.a.1289	CN	F	F	F	F	OCH ₃	H	H
I.a.1290	F	CN	F	F	F	OCH ₃	H	H
I.a.1291	H	F	F	F	F	OCH ₃	H	H
I.a.1292	F	F	Br	F	F	OCH ₃	H	H
I.a.1293	F	F	C≡CH	F	F	OCH ₃	H	H
I.a.1294	CF ₃	Cl	H	H	F	OCH ₃	H	H
I.a.1295	F	F	I	F	F	OCH ₃	H	H
I.a.1296	F	H	H	H	F	OCH ₃	CH ₃	H
I.a.1297	Cl	H	H	H	F	OCH ₃	CH ₃	H
I.a.1298	Br	H	H	H	F	OCH ₃	CH ₃	H
I.a.1299	CN	H	H	H	F	OCH ₃	CH ₃	H
I.a.1300	CH ₃	H	H	H	F	OCH ₃	CH ₃	H
I.a.1301	F	H	H	F	F	OCH ₃	CH ₃	H
I.a.1302	Cl	H	H	F	F	OCH ₃	CH ₃	H
I.a.1303	F	H	H	Cl	F	OCH ₃	CH ₃	H
I.a.1304	Cl	H	H	F	F	OCH ₃	CH ₃	H
I.a.1305	CN	H	H	F	F	OCH ₃	CH ₃	H
I.a.1306	F	H	H	CN	F	OCH ₃	CH ₃	H
I.a.1307	CN	H	H	F	F	OCH ₃	CH ₃	H
I.a.1308	F	H	F	H	F	OCH ₃	CH ₃	H
I.a.1309	Cl	H	F	H	F	OCH ₃	CH ₃	H
I.a.1310	CN	H	F	H	F	OCH ₃	CH ₃	H

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1311	F	F	F	H	F	OCH ₃	CH ₃	H
I.a.1312	Cl	F	F	H	F	OCH ₃	CH ₃	H
I.a.1313	F	Cl	F	H	F	OCH ₃	CH ₃	H
I.a.1314	Cl	F	F	H	F	OCH ₃	CH ₃	H
I.a.1315	CN	F	F	H	F	OCH ₃	CH ₃	H
I.a.1316	F	CN	F	H	F	OCH ₃	CH ₃	H
I.a.1317	CN	F	F	H	F	OCH ₃	CH ₃	H
I.a.1318	F	F	H	F	F	OCH ₃	CH ₃	H
I.a.1319	Cl	F	H	F	F	OCH ₃	CH ₃	H
I.a.1320	F	Cl	H	F	F	OCH ₃	CH ₃	H
I.a.1321	CN	F	H	F	F	OCH ₃	CH ₃	H
I.a.1322	F	CN	H	F	F	OCH ₃	CH ₃	H
I.a.1323	F	F	F	F	F	OCH ₃	CH ₃	H
I.a.1324	Cl	F	F	F	F	OCH ₃	CH ₃	H
I.a.1325	F	Cl	F	F	F	OCH ₃	CH ₃	H
I.a.1326	CN	F	F	F	F	OCH ₃	CH ₃	H
I.a.1327	F	CN	F	F	F	OCH ₃	CH ₃	H
I.a.1328	H	F	F	F	F	OCH ₃	CH ₃	H
I.a.1329	F	F	Br	F	F	OCH ₃	CH ₃	H
I.a.1330	F	F	C≡CH	F	F	OCH ₃	CH ₃	H
I.a.1331	CF ₃	Cl	H	H	F	OCH ₃	CH ₃	H
I.a.1332	F	F	I	F	F	OCH ₃	CH ₃	H
I.a.1333	F	H	H	H	F	OCH ₃	CH ₃	CH ₃
I.a.1334	Cl	H	H	H	F	OCH ₃	CH ₃	CH ₃
I.a.1335	Br	H	H	H	F	OCH ₃	CH ₃	CH ₃
I.a.1336	CN	H	H	H	F	OCH ₃	CH ₃	CH ₃
I.a.1337	CH ₃	H	H	H	F	OCH ₃	CH ₃	CH ₃
I.a.1338	F	H	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1339	Cl	H	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1340	F	H	H	Cl	F	OCH ₃	CH ₃	CH ₃
I.a.1341	Cl	H	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1342	CN	H	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1343	F	H	H	CN	F	OCH ₃	CH ₃	CH ₃
I.a.1344	CN	H	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1345	F	H	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1346	Cl	H	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1347	CN	H	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1348	F	F	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1349	Cl	F	F	H	F	OCH ₃	CH ₃	CH ₃

No.	R ^a	R ^b	R ^c	R ^d	R ^e	R ²	R ³	R ⁴
I.a.1350	F	Cl	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1351	Cl	F	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1352	CN	F	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1353	F	CN	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1354	CN	F	F	H	F	OCH ₃	CH ₃	CH ₃
I.a.1355	F	F	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1356	Cl	F	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1357	F	Cl	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1358	CN	F	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1359	F	CN	H	F	F	OCH ₃	CH ₃	CH ₃
I.a.1360	F	F	F	F	F	OCH ₃	CH ₃	CH ₃
I.a.1361	Cl	F	F	F	F	OCH ₃	CH ₃	CH ₃
I.a.1362	F	Cl	F	F	F	OCH ₃	CH ₃	CH ₃
I.a.1363	CN	F	F	F	F	OCH ₃	CH ₃	CH ₃
I.a.1364	F	CN	F	F	F	OCH ₃	CH ₃	CH ₃
I.a.1365	H	F	F	F	F	OCH ₃	CH ₃	CH ₃
I.a.1366	F	F	Br	F	F	OCH ₃	CH ₃	CH ₃
I.a.1367	F	F	C≡CH	F	F	OCH ₃	CH ₃	CH ₃
I.a.1368	CF ₃	Cl	H	H	F	OCH ₃	CH ₃	CH ₃
I.a.1369	F	F	I	F	F	OCH ₃	CH ₃	CH ₃
I.a.1370	F	H	H	H	F	H	-O(CH ₂) ₃ -	
I.a.1371	Cl	H	H	H	F	H	-O(CH ₂) ₃ -	
I.a.1372	Br	H	H	H	F	H	-O(CH ₂) ₃ -	
I.a.1373	CN	H	H	H	F	H	-O(CH ₂) ₃ -	
I.a.1374	CH ₃	H	H	H	F	H	-O(CH ₂) ₃ -	
I.a.1375	F	H	H	F	F	H	-O(CH ₂) ₃ -	
I.a.1376	Cl	H	H	F	F	H	-O(CH ₂) ₃ -	
I.a.1377	F	H	H	Cl	F	H	-O(CH ₂) ₃ -	
I.a.1378	Cl	H	H	F	F	H	-O(CH ₂) ₃ -	
I.a.1379	CN	H	H	F	F	H	-O(CH ₂) ₃ -	
I.a.1380	F	H	H	CN	F	H	-O(CH ₂) ₃ -	
I.a.1381	CN	H	H	F	F	H	-O(CH ₂) ₃ -	
I.a.1382	F	H	F	H	F	H	-O(CH ₂) ₃ -	
I.a.1383	Cl	H	F	H	F	H	-O(CH ₂) ₃ -	
I.a.1384	CN	H	F	H	F	H	-O(CH ₂) ₃ -	
I.a.1385	F	F	F	H	F	H	-O(CH ₂) ₃ -	
I.a.1386	Cl	F	F	H	F	H	-O(CH ₂) ₃ -	
I.a.1387	F	Cl	F	H	F	H	-O(CH ₂) ₃ -	
I.a.1388	Cl	F	F	H	F	H	-O(CH ₂) ₃ -	

I.a.1389	CN	F	F	H	F	H	-O(CH ₂) ₃ -
I.a.1390	F	CN	F	H	F	H	-O(CH ₂) ₃ -
I.a.1391	CN	F	F	H	F	H	-O(CH ₂) ₃ -
I.a.1392	F	F	H	F	F	H	-O(CH ₂) ₃ -
I.a.1393	Cl	F	H	F	F	H	-O(CH ₂) ₃ -
I.a.1394	F	Cl	H	F	F	H	-O(CH ₂) ₃ -
I.a.1395	CN	F	H	F	F	H	-O(CH ₂) ₃ -
I.a.1396	F	CN	H	F	F	H	-O(CH ₂) ₃ -
I.a.1397	F	F	F	F	F	H	-O(CH ₂) ₃ -
I.a.1398	Cl	F	F	F	F	H	-O(CH ₂) ₃ -
I.a.1399	F	Cl	F	F	F	H	-O(CH ₂) ₃ -
I.a.1400	CN	F	F	F	F	H	-O(CH ₂) ₃ -
I.a.1401	F	CN	F	F	F	H	-O(CH ₂) ₃ -
I.a.1402	H	F	F	F	F	H	-O(CH ₂) ₃ -
I.a.1403	F	F	Br	F	F	H	-O(CH ₂) ₃ -
I.a.1404	F	F	C≡CH	F	F	H	-O(CH ₂) ₃ -
I.a.1405	CF ₃	Cl	H	H	F	H	-O(CH ₂) ₃ -
I.a.1406	F	F	I	F	F	H	-O(CH ₂) ₃ -

In another preferred embodiment, the azines useful for the present invention are azines of formula (I)



5 wherein

- A is heteroaryl, which is substituted by 1, 2, 3, 4, 5 or 6 substituents R^A selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22 aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups;
- 15 R¹ is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-

alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or completely halogenated,

5 phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylcarbonyl and phenoxy carbonyl,

wherein phenyl in the last 6 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

10 R² is H, halogen, OH, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, C₃-C₆-cycloalkoxy or (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 9 aforementioned radicals are unsubstituted, partly or completely halogenated,

15 phenyl, phenyl-C₁-C₆-alkyl,

wherein phenyl in the last 2 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

R³ is selected from the group consisting of H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

20 R⁴ is selected from the group consisting of H, halogen, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₃-C₆-cycloalkenyl and C₁-C₆-alkoxy-C₁-C₆-alkyl, where the aliphatic and cycloaliphatic parts of the 7 aforementioned radicals are unsubstituted, partly or completely halogenated; or

25 R³ and R⁴ together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl and three- to six-membered heterocyclyl,

wherein the C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, or three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy; and

30 R⁵ is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or completely halogenated,

35 phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylcarbonyl and phenoxy carbonyl,

40 wherein phenyl in the last 6 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

and wherein the variables A, R¹, R², R³, R⁴ and R⁵ are in particular:

- A is heteroaryl, which is substituted by 1, 2, 3, 4, 5 or 6 substituents R^A selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22 aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups;
- R¹ is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or completely halogenated, phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylcarbonyl and phenoxycarbonyl, wherein phenyl in the last 6 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;
- R² is H, halogen, OH, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, C₃-C₆-cycloalkoxy or (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 9 aforementioned radicals are unsubstituted, partly or completely halogenated or phenyl, wherein phenyl is unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;
- R³ is selected from the group consisting of H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;
- R⁴ is selected from the group consisting of H, halogen, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₃-C₆-cycloalkenyl and C₁-C₆-alkoxy-C₁-C₆-alkyl, where the aliphatic and cycloaliphatic parts of the 7 aforementioned radicals are unsubstituted, partly or completely halogenated; or
- R³ and R⁴ together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl and three- to six-membered heterocyclyl,

wherein the C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, or three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy; and

5 R⁵ is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or
10 completely halogenated,
phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylcarbonyl and phenoxycarbonyl,
wherein phenyl in the last 6 mentioned radicals are unsubstituted or substituted by 1, 2, 3,
4 or 5 identical or different substituents selected from the group consisting of
15 halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

and wherein the variables A, R¹, R², R³, R⁴ and R⁵ are in more particular:

A is heteroaryl, which is substituted by 1, 2, 3, 4, 5 or 6 substituents R^A selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl,
20 C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-
25 C₄-alkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22 aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups;


R¹ is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or
30 completely halogenated,
phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylcarbonyl and phenoxycarbonyl,

wherein phenyl in the last 6 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or
5 identical or different substituents selected from the group consisting of halogen,
40 CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

- 5 R^2 is H, halogen, OH, CN, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_3 - C_6 -alkynyl, (C_1 - C_6 -alkoxy)- C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, (C_3 - C_6 -cycloalkyl)- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxy, C_2 - C_6 -alkenyloxy, C_2 - C_6 -alkynyloxy, C_3 - C_6 -cycloalkoxy or (C_3 - C_6 -cycloalkyl)- C_1 - C_4 -alkoxy, where the aliphatic and cycloaliphatic parts of the 9 aforementioned radicals are unsubstituted, partly or completely halogenated,
- 10 R^3 is selected from the group consisting of H, halogen, CN, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy and C_1 - C_6 -haloalkoxy;
- R^4 is selected from the group consisting of H, halogen, CN, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_3 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, (C_3 - C_6 -cycloalkyl)- C_1 - C_4 -alkyl, C_3 - C_6 -cycloalkenyl and C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, where the aliphatic and cycloaliphatic parts of the 7 aforementioned radicals are unsubstituted, partly or completely halogenated; or
- R^3 and R^4 together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, C_2 - C_6 -alkenyl, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl and three- to six-membered heterocyclyl,
- 15 wherein the C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, or three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C_1 - C_6 -alkyl and C_1 - C_6 -alkoxy; and
- R^5 is selected from the group consisting of H, OH, $S(O)_2NH_2$, CN, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, (C_3 - C_6 -cycloalkyl)- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxy, (C_1 - C_6 -alkoxy)- C_1 - C_6 -alkyl, (C_1 - C_6 -alkyl)-carbonyl, (C_1 - C_6 -alkoxy)carbonyl, (C_1 - C_6 -alkyl)sulfonyl, (C_1 - C_6 -alkylamino)carbonyl, di(C_1 - C_6 -alkyl)aminocarbonyl, (C_1 - C_6 -alkylamino)sulfonyl, di(C_1 - C_6 -alkyl)aminosulfonyl and (C_1 - C_6 -alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or completely halogenated,
- 25 phenyl, phenyl- C_1 - C_6 -alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylcarbonyl and phenoxycarbonyl,
- wherein phenyl in the last 6 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO_2 , C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy and C_1 - C_6 -haloalkoxy;
- 30 and wherein the variables A, R^1 , R^2 , R^3 , R^4 and R^5 are preferred:
- A is heteroaryl, which is substituted by one to six substituents selected from the group consisting of halogen, CN, NO_2 , C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, OH, C_1 - C_6 -alkoxy, C_1 - C_6 -alkylthio, (C_1 - C_6 -alkyl)sulfinyl, (C_1 - C_6 -alkyl)sulfonyl, amino, (C_1 - C_6 -alkyl)amino,
- 35 di(C_1 - C_6 -alkyl)amino, (C_1 - C_6 -alkyl)carbonyl, (C_1 - C_6 -alkoxy)carbonyl;
- R^1 is H, CN, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, (C_1 - C_6 -alkyl)carbonyl, (C_1 - C_6 -alkoxy)carbonyl, (C_1 - C_6 -alkyl)sulfonyl or phenylsulfonyl,
- wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO_2 , C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl and C_1 - C_6 -alkoxy;
- 40

- R^2 is H, halogen, OH, CN, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_3 - C_6 -alkynyl, (C_1 - C_6 -alkoxy)- C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, (C_3 - C_6 -cycloalkyl)- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxy, C_2 - C_6 -alkenyloxy, C_2 - C_6 -alkynyloxy, C_3 - C_6 -cycloalkoxy or (C_3 - C_6 -cycloalkyl)- C_1 - C_4 -alkoxy, where the aliphatic and cycloaliphatic parts of the 9
- 5 R^3 is H, halogen, CN, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl or C_1 - C_6 -alkoxy;
- R^4 is H, halogen, CN, C_1 - C_6 -alkyl or C_1 - C_6 -haloalkyl; or
- R^3 and R^4 together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, C_2 - C_6 -alkenyl, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl
- 10 and three- to six-membered heterocyclyl,
- wherein the C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, or three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C_1 - C_6 -alkyl and C_1 - C_6 -alkoxy; and
- R^5 is H, CN, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, (C_1 - C_6 -alkyl)carbonyl,
- 15 (C_1 - C_6 -alkoxy)carbonyl, (C_1 - C_6 -alkyl)sulfonyl or phenylsulfonyl,
- wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO_2 , C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl and C_1 - C_6 -alkoxy;
- 20 including their agriculturally acceptable salts or N-oxides.

In another preferred embodiment, the azines useful for the present invention comprise a diaminotriazine compound of formula (I)



wherein

p is 1 or 2;

q is 0, 1, 2 or 3 provided that p+q is 1, 2, 3 or 4;

30 Q is a chemical bond, O, $S(O)_m$, $CR^{q1}R^{q2}$, NR^{q3} , C(O), C(O)O, $CR^{q1}R^{q2}$ -O, $S(O)_mNR^{q3}$ or C(O) NR^{q3} ,

wherein

m is 0, 1 or 2;

R^{q1} , R^{q2} are hydrogen, halogen or C_1 - C_4 -alkyl;

35 R^{q3} is H, CN, C_1 - C_6 -alkyl, (C_1 - C_6 -alkoxy)- C_1 - C_6 -alkyl, (C_1 - C_6 -alkyl)-carbonyl, (C_1 - C_6 -alkoxy)carbonyl, (C_1 - C_6 -alkyl)sulfonyl, where the aliphatic parts of the 6 aforementioned radicals are unsubstituted, partly or completely halogenated;

- Ar is phenyl, which is unsubstituted or carries 1, 2, 3, 4 or 5 radicals R^A which are selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22
aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups,
phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylaminocarbonyl, phenyl(C₁-C₆-alkyl)aminocarbonyl, phenylcarbonyl and phenoxy carbonyl,
wherein phenyl in the last 8 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy,
it being possible that R^A are identical or different;
- R^a is selected from the group consisting of hydrogen, halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or completely halogenated;
- R^b is selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22
aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups,
for q = 2 or 3 it being possible that R^b are identical or different;
- R¹ is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₃-C₆-cycloalkyl)-carbonyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-

alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 15 aforementioned radicals are unsubstituted, partly or completely halogenated,

5 phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylaminocarbonyl, phenyl(C₁-C₆-alkyl)aminocarbonyl, phenylcarbonyl and phenoxy carbonyl,

wherein phenyl in the last 8 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

10 R² is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₃-C₆-cycloalkyl)-carbonyl C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 15 aforementioned radicals are unsubstituted, partly or completely halogenated,

20 phenyl, phenylsulfonyl, phenylaminosulfonyl, phenylaminocarbonyl, phenyl(C₁-C₆-alkyl)aminocarbonyl, phenyl-C₁-C₆ alkyl, phenoxy, phenylcarbonyl and phenoxy carbonyl,

wherein phenyl in the last 9 mentioned radicals is unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

25 X is a radical selected from the group consisting of CR³R⁴R⁵,

phenyl, which is unsubstituted or carries 1, 2, 3, 4 or 5 radicals R^{Ar} which are identical or different;

30 NR^{3a}R^{3b},

OR^{3c} and

S(O)_kR^{3d} with k being 0,1 or 2

wherein

35 R³ is selected from the group consisting of H, halogen, OH, CN, C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 9 aforementioned radicals are unsubstituted, partly or completely halogenated;

40 R⁴ is selected from the group consisting of H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

R⁵ is selected from the group consisting of halogen, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₃-C₆-

cycloalkenyl and C₁-C₆-alkoxy-C₁-C₆-alkyl, where the aliphatic and cycloaliphatic parts of the 7 aforementioned radicals are unsubstituted, partly or completely halogenated;

R⁴ and R⁵ together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, three- to six-membered saturated or partially unsaturated heterocyclyl, and the moiety >C=CR^xR^y, where R^x and R^y are hydrogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R^{Ar} selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22 aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups,


R^{3a}, R^{3b}, R^{3c} or R^{3d} are independently of one another are selected from the group consisting of H, CN, S(O)₂NH₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₃-C₆-cycloalkyl)-carbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 15 aforementioned radicals are unsubstituted, partly or completely halogenated, phenyl, phenylsulfonyl, phenyl-C₁-C₆ alkyl, phenylaminosulfonyl, phenylcarbonyl and phenoxycarbonyl, wherein phenyl in the last 6 mentioned radicals is unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy, or

R^{3a}, R^{3b} together with the nitrogen atom, to which they are bound, form an N-bound, mono – or bicyclic heterocyclic radical, which may have 1, 2, 3 or 4 further heteroatoms which are selected from N, O and S, which is unsubstituted or substituted by one or more identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy, one of R^{3a}, R^{3b} may also be OH, C₁-C₆-alkoxy, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, where the aliphatic and cycloaliphatic parts of the 7 aforementioned radicals are unsubstituted, partly or completely halogenated, or phenoxy, which is unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or

different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

including their agriculturally acceptable salts

- 5 In another preferred embodiment, the azines useful for the present invention comprises a diaminotriazine compound of the formula:



10 wherein

A is a fused saturated or unsaturated, 5- or 6-membered carbocycle or a fused saturated or unsaturated, 5- or 6-membered heterocycle having 1, 2 or 3 heteroatoms or heteroatom moieties, selected from O, S, S(O)_p, N or NR^c as ring members,

15 where the carbocycle and the heterocycle are unsubstituted or carry 1, 2, 3 or 4 radicals R^A;

p is 0, 1 or 2

q is 0, 1, 2 or 3;

R^A is selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22 aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups,

30 it being possible that R^A are identical or different, it being possible that two radicals R^A which are bound at the same carbon atom may together be =O or =NR^d;

R^b is selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₃-C₆-

35

cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22 aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups,

5

for q = 2 or 3 it being possible that R^b are identical or different;

R^c

is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 16 aforementioned radicals are unsubstituted, partly or completely halogenated,

10

R^d

is selected from the group consisting of H, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, where the aliphatic and cycloaliphatic parts of the 8 aforementioned radicals are unsubstituted, partly or completely halogenated;

15

R¹

is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₃-C₆-cycloalkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 17 aforementioned radicals are unsubstituted, partly or completely halogenated, phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylaminocarbonyl, phenyl(C₁-C₆-alkyl)aminocarbonyl, phenylcarbonyl and phenoxy carbonyl,

20

25

30

wherein phenyl in the last 8 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

R²

is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)carbonyl, (C₃-C₆-cycloalkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl, di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 17 aforementioned radicals are unsubstituted, partly or completely halogenated,

35

40

phenyl, phenylsulfonyl, phenylaminosulfonyl, phenyl-C₁-C₆ alkyl, phenoxy, phenylaminocarbonyl, phenyl(C₁-C₆-alkyl)aminocarbonyl, phenylcarbonyl and phenoxy carbonyl,

wherein phenyl in the last 8 mentioned radicals is unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

X is a radical selected from the group consisting of CR³R⁴R⁵

phenyl, which is unsubstituted or carries 1, 2, 3, 4 or 5 radicals R^{Ar} which are identical or different;

NR^{3a}R^{3b},

OR^{3c} and

S(O)_kR^{3d} with k being 0, 1 or 2 ,

wherein

R³ is selected from the group consisting of H, halogen, OH, CN, C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 9 aforementioned radicals are unsubstituted, partly or completely halogenated;

R⁴ is selected from the group consisting of H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;

R⁵ is selected from the group consisting of halogen, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₃-C₆-cycloalkenyl and C₁-C₆-alkoxy-C₁-C₆-alkyl, where the aliphatic and cycloaliphatic parts of the 7 aforementioned radicals are unsubstituted, partly or completely halogenated;

R⁴ and R⁵ together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, thiocarbonyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, three- to six-membered saturated or partially unsaturated heterocyclyl, and the moiety >C=CR^xR^y, where R^x and R^y are hydrogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl;

R^{Ar} selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkoxy, C₂-C₆-alkenyloxy, C₂-C₆-alkynyloxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, (C₁-C₆-alkyl)-carbonyloxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkoxy, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkoxy, where the aliphatic and cycloaliphatic parts of the 22 aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups,

R^{3a} , R^{3b} , R^{3c} or R^{3d} are independently of one another are selected from the group consisting of H, CN, $S(O)_2NH_2$, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, (C_3 - C_6 -cycloalkyl)- C_1 - C_6 -alkyl, (C_1 - C_6 -alkoxy)- C_1 - C_6 -alkyl, (C_1 - C_6 -alkyl)-carbonyl, (C_3 - C_6 -cycloalkyl)-carbonyl, (C_1 - C_6 -alkoxy)carbonyl, (C_1 - C_6 -alkyl)sulfonyl, (C_1 - C_6 -alkylamino)carbonyl, di(C_1 - C_6 -alkyl)aminocarbonyl, (C_1 - C_6 -alkylamino)sulfonyl, di(C_1 - C_6 -alkyl)aminosulfonyl and (C_1 - C_6 -alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 16 aforementioned radicals are unsubstituted, partly or completely halogenated,


phenyl, phenylsulfonyl, phenyl- C_1 - C_6 alkyl, phenylaminosulfonyl, phenylcarbonyl and phenoxy carbonyl, wherein phenyl in the last 6 mentioned radicals is unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO_2 , C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy and C_1 - C_6 -haloalkoxy, or

R^{3a} , R^{3b} together with the nitrogen atom, to which they are bound, form an N-bound, mono – or bicyclic heterocyclic radical, which may have 1, 2, 3 or 4 further heteroatoms which are selected from N, O and S, which is unsubstituted or substituted by one or more identical or different substituents selected from the group consisting of halogen, CN, NO_2 , C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy and C_1 - C_6 -haloalkoxy,

one of R^{3a} , R^{3b} may also be OH, C_1 - C_6 -alkoxy, C_3 - C_6 -cycloalkoxy, (C_3 - C_6 -cycloalkyl)- C_1 - C_4 -alkoxy, C_2 - C_6 -alkenyloxy, C_2 - C_6 -alkynyloxy, (C_1 - C_6 -alkoxy)- C_1 - C_6 -alkoxy, where the aliphatic and cycloaliphatic parts of the 6 aforementioned radicals are unsubstituted, partly or completely halogenated, or phenoxy, which is unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO_2 , C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy and C_1 - C_6 -haloalkoxy;

including their agriculturally acceptable salts.

In another preferred embodiment, the azines useful for the present invention comprise a diaminotriazine compound of formula (I)



wherein

- 5 A is phenyl, which is substituted by fluorine in the ortho-position and which may additionally carry 1, 2, 3 or 4 identical or different substituents R^A selected from the group consisting of halogen, OH, CN, amino, NO₂, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₁-C₆-alkoxy)-C₂-C₆-alkyl, (C₁-C₆-alkoxy)-C₂-C₆-alkenyl, (C₁-C₆-alkoxy)-C₂-C₆-alkynyl, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)-carbonyl, C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, where the aliphatic and cycloaliphatic parts of the 11 aforementioned radicals are unsubstituted, partly or completely halogenated and where the cycloaliphatic parts of the last 4
- 10 mentioned radicals may carry 1, 2, 3, 4, 5 or 6 methyl groups;
- R¹ is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)-carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl,
- 15 di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or completely halogenated,
- phenyl, phenyl-C₁-C₆-alkyl, phenylsulfonyl, phenylaminosulfonyl, phenylcarbonyl and phenoxy carbonyl,
- 20 wherein phenyl in the last 6 mentioned radicals are unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;
- R² is selected from the group consisting of H, OH, S(O)₂NH₂, CN, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl, (C₁-C₆-alkylamino)carbonyl, di(C₁-C₆-alkyl)aminocarbonyl, (C₁-C₆-alkylamino)sulfonyl,
- 25 di(C₁-C₆-alkyl)aminosulfonyl and (C₁-C₆-alkoxy)sulfonyl, where the aliphatic and cycloaliphatic parts of the 14 aforementioned radicals are unsubstituted, partly or completely halogenated,
- phenyl, phenylsulfonyl, phenylaminosulfonyl, phenyl-C₁-C₆ alkyl, phenoxy, phenylcarbonyl and phenoxy carbonyl,
- 30 wherein phenyl in the last 6 mentioned radicals is unsubstituted or substituted by 1, 2, 3, 4 or 5 identical or different substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy;
- 35 R³ is selected from the group consisting of C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, (C₃-C₆-cycloalkyl)-C₁-C₄-alkyl, (C₁-C₆-alkoxy)-C₁-C₆-alkyl, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkylamino)carbonyl and di(C₁-C₆-alkyl)aminocarbonyl, where the aliphatic and cycloaliphatic parts of the 9 aforementioned radicals are unsubstituted, partly or completely halogenated,
- 40 R⁴ and R⁵, together with the carbon atom, to which they are bound form a saturated 3-, 4-, 5- or 6-membered carbocyclic radical or a saturated 3-, 4-, 5- or 6-membered heterocyclic radical, where the carbocyclic radical and the heterocyclic radical are

unsubstituted, partly or completely halogenated or carry from 1 to 6 C₁-C₆-alkyl groups;

including their agriculturally acceptable salts.

5

The herbicidal compounds (component A) useful for the present invention as disclosed SUPRA may further be used in conjunction with additional herbicides to which the crop plant is naturally tolerant or to which has been made tolerant by mutagenesis as described SUPRA, or to which it is resistant via expression of one or more additional transgenes as mentioned supra. The CESA-inhibiting herbicides useful for the present invention are often best applied in conjunction with one or more other herbicides to obtain control of a wider variety of undesirable vegetation. When used in conjunction with other herbicides (hereinafter referred to a compound B), the presently claimed compounds can be formulated with the other herbicide or herbicides, tank mixed with the other herbicide or herbicides, or applied sequentially with the other herbicide or herbicides.

The further herbicidal compound B (component B) is in particular selected from the herbicides of class b1) to b15):

- b1) lipid biosynthesis inhibitors;
- 20 b2) acetolactate synthase inhibitors (ALS inhibitors);
- b3) photosynthesis inhibitors;
- b4) protoporphyrinogen-IX oxidase inhibitors,
- b5) bleacher herbicides;
- b6) enolpyruvyl shikimate 3-phosphate synthase inhibitors (EPSP inhibitors);
- 25 b7) glutamine synthetase inhibitors;
- b8) 7,8-dihydropteroate synthase inhibitors (DHP inhibitors);
- b9) mitosis inhibitors;
- b10) inhibitors of the synthesis of very long chain fatty acids (VLCFA inhibitors);
- b11) cellulose biosynthesis inhibitors;
- 30 b12) decoupler herbicides;
- b13) auxinic herbicides;
- b14) auxin transport inhibitors; and
- b15) other herbicides selected from the group consisting of bromobutide, chlorflurenol, chlorflurenol-methyl, cinmethylin, cumyluron, dalapon, dazomet, difenzoquat, difenzoquat-metilsulfate, dimethipin, DSMA, dymron, endothal and its salts,
- 35 etobenzanid, flamprop, flamprop-isopropyl, flamprop-methyl, flamprop-M-isopropyl, flamprop-M-methyl, flurenol, flurenol-butyl, flurprimidol, fosamine, fosamine-ammonium, indanofan, indaziflam, maleic hydrazide, mefluidide, metam, methiozolin (CAS 403640-27-7), methyl azide, methyl bromide, methyl-dymron, methyl iodide,
- 40 MSMA, oleic acid, oxaziclomefone, pelargonic acid, pyributicarb, quinoclamine, triaziflam, tridiphane and 6-chloro-3-(2-cyclopropyl-6-methylphenoxy)-4-pyridazinol (CAS 499223-49-3) and its salts and esters;

including their agriculturally acceptable salts or derivatives such as ethers, esters or amides.

Preference is given to those compositions according to the present invention comprising at least one herbicide B selected from herbicides of class b1, b6, b9, b10 and b11.

Examples of herbicides B which can be used in combination with the compounds of formula (I) according to the present invention are:

b1) from the group of the lipid biosynthesis inhibitors:

ACC-herbicides such as alloxydim, alloxydim-sodium, butroxydim, clethodim, clodinafop, clodinafop-propargyl, cycloxydim, cyhalofop, cyhalofop-butyl, diclofop, diclofop-methyl, fenoxaprop, fenoxaprop-ethyl, fenoxaprop-P, fenoxaprop-P-ethyl, fluazifop, fluazifop-butyl, fluazifop-P, fluazifop-P-butyl, haloxyfop, haloxyfop-methyl, haloxyfop-P, haloxyfop-P-methyl, metamifop, pinoxaden, profoxydim, propaquizafop, quizalofop, quizalofop-ethyl, quizalofop-tefuryl, quizalofop-P, quizalofop-P-ethyl, quizalofop-P-tefuryl, sethoxydim, tepraloxydim, tralkoxydim, 4-(4'-Chloro-4-cyclopropyl-2'-fluoro[1,1'-biphenyl]-3-yl)-5-hydroxy-2,2,6,6-tetramethyl-2H-pyran-3(6H)-one (CAS 1312337-72-6); 4-(2',4'-Dichloro-4-cyclopropyl[1,1'-biphenyl]-3-yl)-5-hydroxy-2,2,6,6-tetramethyl-2H-pyran-3(6H)-one (CAS 1312337-45-3); 4-(4'-Chloro-4-ethyl-2'-fluoro[1,1'-biphenyl]-3-yl)-5-hydroxy-2,2,6,6-tetramethyl-2H-pyran-3(6H)-one (CAS 1033757-93-5); 4-(2',4'-Dichloro-4-ethyl[1,1'-biphenyl]-3-yl)-2,2,6,6-tetramethyl-2H-pyran-3,5(4H,6H)-dione (CAS 1312340-84-3); 5-(Acetyloxy)-4-(4'-chloro-4-cyclopropyl-2'-fluoro[1,1'-biphenyl]-3-yl)-3,6-dihydro-2,2,6,6-tetramethyl-2H-pyran-3-one (CAS 1312337-48-6); 5-(Acetyloxy)-4-(2',4'-dichloro-4-cyclopropyl- [1,1'-biphenyl]-3-yl)-3,6-dihydro-2,2,6,6-tetramethyl-2H-pyran-3-one; 5-(Acetyloxy)-4-(4'-chloro-4-ethyl-2'-fluoro[1,1'-biphenyl]-3-yl)-3,6-dihydro-2,2,6,6-tetramethyl-2H-pyran-3-one (CAS 1312340-82-1); 5-(Acetyloxy)-4-(2',4'-dichloro-4-ethyl[1,1'-biphenyl]-3-yl)-3,6-dihydro-2,2,6,6-tetramethyl-2H-pyran-3-one (CAS 1033760-55-2); 4-(4'-Chloro-4-cyclopropyl-2'-fluoro[1,1'-biphenyl]-3-yl)-5,6-dihydro-2,2,6,6-tetramethyl-5-oxo-2H-pyran-3-yl carbonic acid methyl ester (CAS 1312337-51-1); 4-(2',4'-Dichloro -4-cyclopropyl- [1,1'-biphenyl]-3-yl)-5,6-dihydro-2,2,6,6-tetramethyl-5-oxo-2H-pyran-3-yl carbonic acid methyl ester; 4-(4'-Chloro-4-ethyl-2'-fluoro[1,1'-biphenyl]-3-yl)-5,6-dihydro-2,2,6,6-tetramethyl-5-oxo-2H-pyran-3-yl carbonic acid methyl ester (CAS 1312340-83-2); 4-(2',4'-Dichloro-4-ethyl[1,1'-biphenyl]-3-yl)-5,6-dihydro-2,2,6,6-tetramethyl-5-oxo-2H-pyran-3-yl carbonic acid methyl ester (CAS 1033760-58-5); and non ACC herbicides such as benfuresate, butylate, cycloate, dalapon, dimepiperate, EPTC, esprocarb, ethofumesate, flupropanate, molinate, orbencarb, pebulate, prosulfocarb, TCA, thiobencarb, tiocarbazil, triallate and vernolate;

b2) from the group of the ALS inhibitors:

sulfonylureas such as amidosulfuron, azimsulfuron, bensulfuron, bensulfuron-methyl, chlorimuron, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethametsulfuron, ethametsulfuron-methyl, ethoxysulfuron, flazasulfuron, flucetosulfuron, flupyrsulfuron, flupyrsulfuron-methyl-sodium, foramsulfuron, halosulfuron, halosulfuron-methyl, imazosulfuron, iodosulfuron, iodosulfuron-methyl-sodium, iofensulfuron,


- iofensulfuron-sodium, mesosulfuron, metazosulfuron, metsulfuron, metsulfuron-methyl, nicosulfuron, orthosulfamuron, oxasulfuron, primisulfuron, primisulfuron-methyl, propyrisulfuron, prosulfuron, pyrazosulfuron, pyrazosulfuron-ethyl, rimsulfuron, sulfometuron, sulfometuron-methyl, sulfosulfuron, thifensulfuron, thifensulfuron-methyl, 5 triasulfuron, tribenuron, tribenuron-methyl, trifloxysulfuron, triflusulfuron, triflusulfuron-methyl and tritosulfuron,
- imidazolinones such as imazamethabenz, imazamethabenz-methyl, imazamox, imazapic, imazapyr, imazaquin and imazethapyr, triazolopyrimidine herbicides and sulfonanilides such as cloransulam, cloransulam-methyl, diclosulam, flumetsulam, florasulam, metosulam, 10 penoxsulam, pyrimisulfan and pyroxsulam,
- pyrimidinylbenzoates such as bispyribac, bispyribac-sodium, pyribenzoxim, pyriftalid, pyriminobac, pyriminobac-methyl, pyriithiobac, pyriithiobac-sodium, 4-[[[2-[(4,6-dimethoxy-2-pyrimidinyl)oxy]phenyl]methyl]amino]-benzoic acid-1-methylethyl ester (CAS 420138-41-6), 4-[[[2-[(4,6-dimethoxy-2-pyrimidinyl)oxy]phenyl]methyl]amino]-benzoic acid propyl ester 15 (CAS 420138-40-5), N-(4-bromophenyl)-2-[(4,6-dimethoxy-2-pyrimidinyl)oxy]benzenemethanamine (CAS 420138-01-8),
- sulfonylaminocarbonyl-triazolinone herbicides such as flucarbazone, flucarbazone-sodium, propoxycarbazon, propoxycarbazon-sodium, thienicarbazone and thienicarbazone-methyl; and triafamone;
- 20 among these, a preferred embodiment of the invention relates to those compositions comprising at least one imidazolinone herbicide;
- b3) from the group of the photosynthesis inhibitors:
- amicarbazone, inhibitors of the photosystem II, e.g. triazine herbicides, including of chlorotriazine, triazinones, triazindiones, methylthiotriazines and pyridazinones such as 25 ametryn, atrazine, chloridazone, cyanazine, desmetryn, dimethametryn, hexazinone, metribuzin, prometon, prometryn, propazine, simazine, simetryn, terbutometon, terbuthylazin, terbutryn and trietazin, aryl urea such as chlorobromuron, chlorotoluron, chloroxuron, dimefuron, diuron, fluometuron, isoproturon, isouron, linuron, metamitron,
- methabenzthiazuron, metobenzuron, metoxuron, monolinuron, neburon, siduron, 30 tebuthiuron and thiadiazuron, phenyl carbamates such as desmedipham, karbutilat, phenmedipham, phenmedipham-ethyl, nitrile herbicides such as bromofenoxim, bromoxynil and its salts and esters, ioxynil and its salts and esters, uraciles such as bromacil, lenacil and terbacil, and bentazon and bentazon-sodium, pyridate, pyridafol, pentanochlor and propanil and inhibitors of the photosystem I such as diquat, diquat-dibromide, paraquat, 35 paraquat-dichloride and paraquat-dimethylsulfate. Among these, a preferred embodiment of the invention relates to those compositions comprising at least one aryl urea herbicide.
- Among these, likewise a preferred embodiment of the invention relates to those compositions comprising at least one triazine herbicide. Among these, likewise a preferred embodiment of the invention relates to those compositions comprising at least one nitrile 40 herbicide;
- b4) from the group of the protoporphyrinogen-IX oxidase inhibitors:
- acifluorfen, acifluorfen-sodium, azafenidin, bencarbazone, benzfendizone, bifenox,

- butafenacil, carfentrazone, carfentrazone-ethyl, chlormethoxyfen, cinidon-ethyl, fluazolate, flufenpyr, flufenpyr-ethyl, flumiclorac, flumiclorac-pentyl, flumioxazin, fluoroglycofen, fluoroglycofen-ethyl, fluthiacet, fluthiacet-methyl, fomesafen, halosafen, lactofen, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, profluzol, pyraclonil, pyraflufen, pyraflufen-ethyl,
- 5 saflufenacil, sulfentrazone, thidiazimin, tiafenacil, ethyl [3-[2-chloro-4-fluoro-5-(1-methyl-6-trifluoromethyl-2,4-dioxo-1,2,3,4-tetrahydropyrimidin-3-yl)phenoxy]-2-pyridyloxy]acetate (CAS 353292-31-6; S-3100), N-ethyl-3-(2,6-dichloro-4-trifluoromethylphenoxy)-5-methyl-1*H*-pyrazole-1-carboxamide (CAS 452098-92-9), N-tetrahydrofurfuryl-3-(2,6-dichloro-4-trifluoromethylphenoxy)-5-methyl-1*H*-pyrazole-1-carboxamide (CAS 915396-43-9), N-ethyl-
- 10 3-(2-chloro-6-fluoro-4-trifluoromethylphenoxy)-5-methyl-1*H*-pyrazole-1-carboxamide (CAS 452099-05-7), N-tetrahydrofurfuryl-3-(2-chloro-6-fluoro-4-trifluoromethylphenoxy)-5-methyl-1*H*-pyrazole-1-carboxamide (CAS 452100-03-7), 3-[7-fluoro-3-oxo-4-(prop-2-ynyl)-3,4-dihydro-2*H*-benzo[1,4]oxazin-6-yl]-1,5-dimethyl-6-thioxo-[1,3,5]triazinan-2,4-dione, 1,5-dimethyl-6-thioxo-3-(2,2,7-trifluoro-3-oxo-4-(prop-2-ynyl)-3,4-dihydro-2*H*-
- 15 benzo[b][1,4]oxazin-6-yl)-1,3,5-triazinane-2,4-dione (CAS 1258836-72-4), 2-(2,2,7-Trifluoro-3-oxo-4-prop-2-ynyl-3,4-dihydro-2*H*-benzo[1,4]oxazin-6-yl)-4,5,6,7-tetrahydro-isoindole-1,3-dione, 1-Methyl-6-trifluoromethyl-3-(2,2,7-trifluoro-3-oxo-4-prop-2-ynyl-3,4-dihydro-2*H*-benzo[1,4]oxazin-6-yl)-1*H*-pyrimidine-2,4-dione, methyl (*E*)-4-[2-chloro-5-[4-chloro-5-(difluoromethoxy)-1*H*-methyl-pyrazol-3-yl]-4-fluoro-phenoxy]-3-methoxy-but-2-enoate [CAS
- 20 948893-00-3], and 3-[7-Chloro-5-fluoro-2-(trifluoromethyl)-1*H*-benzimidazol-4-yl]-1-methyl-6-(trifluoromethyl)-1*H*-pyrimidine-2,4-dione (CAS 212754-02-4);
- b5) from the group of the bleacher herbicides:
- PDS inhibitors: beflubutamid, diflufenican, fluridone, flurochloridone, flurtamone, norflurazon, picolinafen, and 4-(3-trifluoromethylphenoxy)-2-(4-trifluoromethylphenyl)-
- 25 pyrimidine (CAS 180608-33-7), HPPD inhibitors: benzobicyclon, benzofenap, clomazone, fenquintrione, isoxaflutole, mesotrione, pyrasulfotole, pyrazolynate, pyrazoxyfen, sulcotrione, tefuryltrione, tembotrione, topramezone and bicyclopyrone, bleacher, unknown target: aclonifen, amitrole and flumeturon;
- b6) from the group of the EPSP synthase inhibitors:
- 30 glyphosate, glyphosate-isopropylammonium, glyposate-potassium and glyphosate-trimesium (sulfosate);
- b7) from the group of the glutamine synthase inhibitors:
- bilanaphos (bialaphos), bilanaphos-sodium, glufosinate, glufosinate-P and glufosinate-ammonium;
- 35 b8) from the group of the DHP synthase inhibitors:
- asulam;
- b9) from the group of the mitosis inhibitors:
- compounds of group K1: dinitroanilines such as benfluralin, butralin, dinitramine, ethalfluralin, fluchloralin, oryzalin, pendimethalin, prodiamine and trifluralin,
- 40 phosphoramidates such as amiprofos, amiprofos-methyl, and butamiphos, benzoic acid herbicides such as chlorthal, chlorthal-dimethyl, pyridines such as dithiopyr and thiazopyr, benzamides such as propyzamide and tebutam; compounds of group K2: chlorpropham,


propham and carbetamide, among these, compounds of group K1, in particular dinitroanilines are preferred;

b10) from the group of the VLCFA inhibitors:


- chloroacetamides such as acetochlor, alachlor, butachlor, dimethachlor, dimethenamid,
 5 dimethenamid-P, metazachlor, metolachlor, metolachlor-S, pethoxamid, pretilachlor,
 propachlor, propisochlor and thenylchlor, oxyacetanilides such as flufenacet and
 mefenacet, acetanilides such as diphenamid, naproanilide, napropamide and napropamide-M,
 tetrazolinones such fentrazamide, and other herbicides such as anilofos, cafenstrole,
 fenoxasulfone, ipfencarbazone, piperophos, pyroxasulfone and isoxazoline compounds of
 10 the formulae II.1, II.2, II.3, II.4, II.5, II.6, II.7, II.8 and II.9




II.1




II.2




II.3




II.4




II.5




II.6



II.7



II.8



II.9

- the isoxazoline compounds of the formula (I) are known in the art, e.g. from WO
 20 2006/024820, WO 2006/037945, WO 2007/071900 and WO 2007/096576;
 among the VLCFA inhibitors, preference is given to chloroacetamides and oxyacetamides;
 b11) from the group of the cellulose biosynthesis inhibitors:
 chlorthiamid, dichlobenil, flupoxam, isoxaben and 1-Cyclohexyl-5-pentafluorophenyl-14-
 [1,2,4,6]thiatiazin-3-ylamine;

b12) from the group of the decoupler herbicides:

dinoseb, dinoterb and DNOC and its salts;

b13) from the group of the auxinic herbicides:

2,4-D and its salts and esters such as clacyfos, 2,4-DB and its salts and esters,

- 5 aminocyclopyrachlor and its salts and esters, aminopyralid and its salts such as aminopyralid-dimethylammonium, aminopyralid-tris(2-hydroxypropyl)ammonium and its esters, benazolin, benazolin-ethyl, chloramben and its salts and esters, clomeprop, clopyralid and its salts and esters, dicamba and its salts and esters, dichlorprop and its salts and esters, dichlorprop-P and its salts and esters, fluroxypyr, fluroxypyr-butometyl, fluroxypyr-meptyl, halauxifen and its salts and esters (CAS 943832-60-8); MCPA and its salts and esters, MCPA-thioethyl, MCPB and its salts and esters, mecoprop and its salts and esters, mecoprop-P and its salts and esters, picloram and its salts and esters, quinclorac, quinmerac, TBA (2,3,6) and its salts and esters and triclopyr and its salts and esters;

- 15 b14) from the group of the auxin transport inhibitors: diflufenzopyr, diflufenzopyr-sodium, naptalam and naptalam-sodium;

- b15) from the group of the other herbicides: bromobutide, chlorflurenol, chlorflurenol-methyl, cinmethylin, cumyluron, cyclopyrimorate (CAS 499223-49-3) and its salts and esters, dalapon, dazomet, difenzoquat, difenzoquat-metilsulfate, dimethipin, DSMA, dymron, endothal and its salts, etobenzanid, flamprop, flamprop-isopropyl, flamprop-methyl, flamprop-M-isopropyl, flamprop-M-methyl, flurenol, flurenol-butyl, flurprimidol, fosamine, fosamine-ammonium, indanofan, indaziflam, maleic hydrazide, mefluidide, metam, methiozolin (CAS 403640-27-7), methyl azide, methyl bromide, methyl-dymron, methyl iodide, MSMA, oleic acid, oxaziclomefone, pelargonic acid, pyributicarb, quinoclamine, triaziflam and tridiphan..

Active compounds B and C having a carboxyl group can be employed in the form of the acid, in the form of an agriculturally suitable salt as mentioned above or else in the form of an agriculturally acceptable derivative in the compositions according to the invention.

- 30 In the case of dicamba, suitable salts include those, where the counterion is an agriculturally acceptable cation. For example, suitable salts of dicamba are dicamba-sodium, dicamba-potassium, dicamba-methylammonium, dicamba-dimethylammonium, dicamba-isopropylammonium, dicamba-diglycolamine, dicamba-olamine, dicamba-diolamine, dicamba-trolamine, dicamba-N,N-bis-(3-aminopropyl)methylamine and dicamba-diethylenetriamine. Examples of a suitable ester are dicamba-methyl and dicamba-butotyl.

- 35 Suitable salts of 2,4-D are 2,4-D-ammonium, 2,4-D-dimethylammonium, 2,4-D-diethylammonium, 2,4-D-diethanolammonium (2,4-D-diolamine), 2,4-D-triethanolammonium, 2,4-D-isopropylammonium, 2,4-D-triisopropanolammonium, 2,4-D-heptylammonium, 2,4-D-dodecylammonium, 2,4-D-tetradecylammonium, 2,4-D-triethylammonium, 2,4-D-tris(2-hydroxypropyl)ammonium, 2,4-D-tris(isopropyl)ammonium, 2,4-D-trolamine, 2,4-D-lithium, 2,4-D-sodium. Examples of suitable esters of 2,4-D are 2,4-D-butotyl, 2,4-D-2-butoxypropyl, 2,4-D-3-butoxypropyl, 2,4-D-butyl, 2,4-D-ethyl, 2,4-D-

ethylhexyl, 2,4-D-isobutyl, 2,4-D-isooctyl, 2,4-D-isopropyl, 2,4-D-meptyl, 2,4-D-methyl, 2,4-D-octyl, 2,4-D-pentyl, 2,4-D-propyl, 2,4-D-tefuryl and clacyfos.

Suitable salts of 2,4-DB are for example 2,4-DB-sodium, 2,4-DB-potassium and 2,4-DB-dimethylammonium. Suitable esters of 2,4-DB are for example 2,4-DB-butyl and 2,4-DB-isooctyl.

Suitable salts of dichlorprop are for example dichlorprop-sodium, dichlorprop-potassium and dichlorprop-dimethylammonium. Examples of suitable esters of dichlorprop are dichlorprop-butotyl and dichlorprop-isooctyl.

Suitable salts and esters of MCPA include MCPA-butotyl, MCPA-butyl, MCPA-dimethylammonium, MCPA-diolamine, MCPA-ethyl, MCPA-thioethyl, MCPA-2-ethylhexyl, MCPA-isobutyl, MCPA-isooctyl, MCPA-isopropyl, MCPA-isopropylammonium, MCPA-methyl, MCPA-olamine, MCPA-potassium, MCPA-sodium and MCPA-trolamine.

A suitable salt of MCPB is MCPB sodium. A suitable ester of MCPB is MCPB-ethyl.

Suitable salts of clopyralid are clopyralid-potassium, clopyralid-olamine and clopyralid-tris-(2-hydroxypropyl)ammonium. Example of suitable esters of clopyralid is clopyralid-methyl.

Examples of a suitable ester of fluroxypyr are fluroxypyr-meptyl and fluroxypyr-2-butoxy-1-methylethyl, wherein fluroxypyr-meptyl is preferred.

Suitable salts of picloram are picloram-dimethylammonium, picloram-potassium, picloram-triisopropanolammonium, picloram-triisopropylammonium and picloram-trolamine. A suitable ester of picloram is picloram-isooctyl.

A suitable salt of triclopyr is triclopyr-triethylammonium. Suitable esters of triclopyr are for example triclopyr-ethyl and triclopyr-butotyl.

Suitable salts and esters of chloramben include chloramben-ammonium, chloramben-diolamine, chloramben-methyl, chloramben-methylammonium and chloramben-sodium. Suitable salts and esters of 2,3,6-TBA include 2,3,6-TBA-dimethylammonium, 2,3,6-TBA-lithium, 2,3,6-TBA-potassium and 2,3,6-TBA-sodium.

Suitable salts and esters of aminopyralid include aminopyralid-potassium, aminopyralid-dimethylammonium, and aminopyralid-tris(2-hydroxypropyl)ammonium.

Suitable salts of glyphosate are for example glyphosate-ammonium, glyphosate-diammonium, glyphosate-dimethylammonium, glyphosate-isopropylammonium, glyphosate-potassium, glyphosate-sodium, glyphosate-trimesium as well as the ethanolamine and diethanolamine salts, preferably glyphosate-diammonium, glyphosate-isopropylammonium and glyphosate-trimesium (sulfosate).

A suitable salt of glufosinate is for example glufosinate-ammonium.

A suitable salt of glufosinate-P is for example glufosinate-P-ammonium.

Suitable salts and esters of bromoxynil are for example bromoxynil-butyrate, bromoxynil-heptanoate, bromoxynil-octanoate, bromoxynil-potassium and bromoxynil-sodium.

Suitable salts and esters of ioxonil are for example ioxonil-octanoate, ioxonil-potassium and ioxonil-sodium.

Suitable salts and esters of mecoprop include mecoprop-butotyl, mecoprop-

dimethylammonium, mecoprop-diolamine, mecoprop-ethadyl, mecoprop-2-ethylhexyl, mecoprop-isooctyl, mecoprop-methyl, mecoprop-potassium, mecoprop-sodium and mecoprop-trolamine.

- Suitable salts of mecoprop-P are for example mecoprop-P-butotyl, mecoprop-P-dimethylammonium, mecoprop-P-2-ethylhexyl, mecoprop-P-isobutyl, mecoprop-P-potassium and mecoprop-P-sodium.

A suitable salt of diflufenzopyr is for example diflufenzopyr-sodium.

A suitable salt of naptalam is for example naptalam-sodium.

- Suitable salts and esters of aminocyclopyrachlor are for example aminocyclopyrachlor-dimethylammonium, aminocyclopyrachlor-methyl, aminocyclopyrachlor-triisopropanolammonium, aminocyclopyrachlor-sodium and aminocyclopyrachlor-potassium.

A suitable salt of quinclorac is for example quinclorac-dimethylammonium.

A suitable salt of quinmerac is for example quinclorac-dimethylammonium.

- A suitable salt of imazamox is for example imazamox-ammonium.

Suitable salts of imazapic are for example imazapic-ammonium and imazapic-isopropylammonium.

Suitable salts of imazapyr are for example imazapyr-ammonium and imazapyr-isopropylammonium.

- A suitable salt of imazaquin is for example imazaquin-ammonium.

Suitable salts of imazethapyr are for example imazethapyr-ammonium and imazethapyr-isopropylammonium.

A suitable salt of topramezone is for example topramezone-sodium.

- Particularly preferred herbicidal compounds B are the herbicides B as defined above; in particular the herbicides B.1 - B.189 listed below in table B:

Table B:

	Herbicide B
B.1	clethodim
B.2	clodinafop-propargyl
B.3	cycloxydim
B.4	cyhalofop-butyl
B.5	fenoxaprop-ethyl
B.6	fenoxaprop-P-ethyl
B.7	metamifop
B.8	pinoxaden
B.9	profoxydim
B.10	sethoxydim
B.11	tepraloxym
B.12	tralkoxydim

	Herbicide B
B.13	esprocarb
B.14	ethofumesate
B.15	molinate
B.16	prosulfocarb
B.17	thiobencarb
B.18	triallate
B.19	bensulfuron-methyl
B.20	bispyribac-sodium
B.21	cloransulam-methyl
B.22	chlorsulfuron
B.23	clorimuron
B.24	cyclosulfamuron
B.25	diclosulam

	Herbicide B
B.26	florasulam
B.27	flumetsulam
B.28	flupyrsulfuron-methyl-sodium
B.29	foramsulfuron
B.30	imazamox
B.31	imazamox-ammonium
B.32	imazapic
B.33	imazapic-ammonium
B.34	imazapic-isopropylammonium
B.35	imazapyr
B.36	imazapyr-ammonium
B.37	imazapyr-isopropylammonium
B.38	imazaquin
B.39	imazaquin-ammonium
B.40	imazethapyr
B.41	imazethapyr-ammonium
B.42	imazethapyr- isopropylammonium
B.43	imazosulfuron
B.44	iodosulfuron-methyl-sodium
B.45	iofensulfuron
B.46	iofensulfuron-sodium
B.47	mesosulfuron-methyl
B.48	metazosulfuron
B.49	metsulfuron-methyl
B.50	metosulam
B.51	nicosulfuron
B.52	penoxsulam
B.53	propoxycarbazon-sodium
B.54	pyrazosulfuron-ethyl
B.55	pyribenzoxim
B.56	pyriftalid
B.57	pyroxsulam
B.58	propyrisulfuron
B.59	rimsulfuron
B.60	sulfosulfuron
B.61	thiencarbazone-methyl
B.62	thifensulfuron-methyl

	Herbicide B
B.63	tribenuron-methyl
B.64	tritosulfuron
B.65	triafamone
B.66	ametryne
B.67	atrazine
B.68	bentazon
B.69	bromoxynil
B.70	bromoxynil-octanoate
B.71	bromoxynil-heptanoate
B.72	bromoxynil-potassium
B.73	diuron
B.74	fluometuron
B.75	hexazinone
B.76	isoproturon
B.77	linuron
B.78	metamitron
B.79	metribuzin
B.80	propanil
B.81	simazin
B.82	terbuthylazine
B.83	terbutryn
B.84	paraquat-dichloride
B.85	acifluorfen
B.86	butafenacil
B.87	carfentrazone-ethyl
B.88	flumioxazin
B.89	fomesafen
B.90	oxadiargyl
B.91	oxyfluorfen
B.92	saflufenacil
B.93	sulfentrazone
B.94	ethyl [3-[2-chloro-4-fluoro-5-(1-methyl-6-trifluoromethyl-2,4-dioxo-1,2,3,4-tetrahydro-pyrimidin-3-yl)phenoxy]-2-pyridyloxy]acetate (CAS 353292-31-6)

	Herbicide B
B.95	1,5-dimethyl-6-thioxo-3-(2,2,7-trifluoro-3-oxo-4-(prop-2-ynyl)-3,4-dihydro-2H-benzo[b][1,4]-oxazin-6-yl)-1,3,5-triazinane-2,4-dione (CAS 1258836-72-4)
B.96	benzobicyclon
B.97	clomazone
B.98	diflufenican
B.99	flurochloridone
B.100	isoxaflutole
B.101	mesotrione
B.102	norflurazone
B.103	picolinafen
B.104	sulcotrione
B.105	tefuryltrione
B.106	tembotrione
B.107	topramezone
B.108	topramezone-sodium
B.109	bicyclopypyrone
B.110	amitrole
B.111	fluometuron
B.112	fenquintrione
B.113	glyphosate
B.114	glyphosate-ammonium
B.115	glyphosate-dimethylammonium
B.116	glyphosate-isopropylammonium
B.117	glyphosate-trimesium (sulfosate)
B.118	glyphosate-potassium
B.119	glufosinate
B.120	glufosinate-ammonium
B.121	glufosinate-P
B.122	glufosinate-P-ammonium
B.123	pendimethalin
B.124	trifluralin

	Herbicide B
B.125	acetochlor
B.126	butachlor
B.127	cafenstrole
B.128	dimethenamid-P
B.129	fentrazamide
B.130	flufenacet
B.131	mefenacet
B.132	metazachlor
B.133	metolachlor
B.134	S-metolachlor
B.135	pretilachlor
B.136	fenoxasulfone
B.137	isoxaben
B.138	ipfencarbazone
B.139	pyroxasulfone
B.140	2,4-D
B.141	2,4-D-isobutyl
B.142	2,4-D-dimethylammonium
B.143	2,4-D-N,N,N-trimethylethanolammonium
B.144	aminopyralid
B.145	aminopyralid-methyl
B.146	aminopyralid-dimethyl-ammonium
B.147	aminopyralid-tris(2-hydroxypropyl)ammonium
B.148	clopyralid
B.149	clopyralid-methyl
B.150	clopyralid-olamine
B.151	dicamba
B.152	dicamba-butotyl
B.153	dicamba-diglycolamine
B.154	dicamba-dimethylammonium
B.155	dicamba-diolamine
B.156	dicamba-isopropylammonium
B.157	dicamba-potassium
B.158	dicamba-sodium
B.159	dicamba-trolamine

	Herbicide B
B.160	dicamba-N,N-bis-(3-aminopropyl)methylamine
B.161	dicamba-diethylenetriamine
B.162	fluroxypyr
B.163	fluroxypyr-meptyl
B.164	MCPA
B.165	MCPA-2-ethylhexyl
B.166	MCPA-dimethylammonium
B.167	quinclorac
B.168	quinclorac-dimethylammonium
B.169	quinmerac
B.170	quinmerac-dimethylammonium
B.171	aminocyclopyrachlor
B.172	aminocyclopyrachlor-potassium
B.173	aminocyclopyrachlor-methyl
B.174	diflufenzopyr
B.175	diflufenzopyr-sodium
B.176	dymron
B.177	indanofan
B.178	indaziflam
B.179	oxaziclomefone
B.180	triaziflam
B.181	II.1
B.182	II.2
B.183	II.3
B.184	II.4
B.185	II.5
B.186	II.6
B.187	II.7
B.188	II.8
B.189	II.9

Moreover, it may be useful to apply the compounds of formula (I) in combination with safeners and optionally with one or more further herbicides. Safeners are chemical compounds which prevent or reduce damage on useful plants without having a major impact on the herbicidal action of the compounds of the formula (I) towards unwanted plants. They can be applied either before sowings (e.g. on seed treatments, shoots or seedlings) or in the pre-emergence application or post-emergence application of the useful plant. The safeners and the compounds of formula (I) and optionally the herbicides B can be applied simultaneously or in succession.

Suitable safeners are e.g. (quinolin-8-oxy)acetic acids, 1-phenyl-5-haloalkyl-1H-1,2,4-triazol-3-carboxylic acids, 1-phenyl-4,5-dihydro-5-alkyl-1H-pyrazol-3,5-dicarboxylic acids, 4,5-dihydro-5,5-diaryl-3-isoxazol carboxylic acids, dichloroacetamides, alpha-oximinophenylacetoneitriles, acetophenonoximes, 4,6-dihalo-2-phenylpyrimidines, N-[[4-(aminocarbonyl)phenyl]sulfonyl]-2-benzoic amides, 1,8-naphthalic anhydride, 2-halo-4-(haloalkyl)-5-thiazol carboxylic acids, phosphorothiolates and N-alkyl-O-phenylcarbamates and their agriculturally acceptable salts and their agriculturally acceptable derivatives such as amides, esters, and thioesters, provided they have an acid group.

Examples of preferred safeners C are benoxacor, cloquintocet, cyometrinil, cyprosulfamide, dichlormid, dicyclonon, dietholate, fenchlorazole, fenclorim, flurazole, fluxofenim, furilazole, isoxadifen, mefenpyr, mephenate, naphthalic anhydride, oxabetrinil, 4-(dichloroacetyl)-1-oxa-4-azaspiro[4.5]decane (MON4660, CAS 71526-07-3), 2,2,5-trimethyl-3-(dichloroacetyl)-1,3-oxazolidine (R-29148, CAS 52836-31-4) and N-(2-Methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide (CAS 129531-12-0).

Particularly preferred safeners C are the following compounds C.1 to C.17

C.1	benoxacor	C.2	cloquintocet
C.3	cloquintocet-mexyl	C.4	cyprosulfamide
C.5	dichlormid	C.6	fenchlorazole
C.7	fenchlorazole-ethyl	C.8	fenclorim
C.9	furilazole	C.10	isoxadifen
C.11	isoxadifen-ethyl	C.12	mefenpyr
C.13	mefenpyr-diethyl	C.14	naphthalic acid anhydride
C.15	4-(dichloroacetyl)-1-oxa-4-azaspiro[4.5]decane	C.16	2,2,5-trimethyl-3-(dichloroacetyl)-1,3-oxazolidine
C.17	N-(2-Methoxybenzoyl)-4-[(methylaminocarbonyl)amino]benzenesulfonamide		

The active compounds B of groups b1) to b15) and the safener compounds C are known herbicides and safeners, see, for example, The Compendium of Pesticide Common Names (<http://www.alanwood.net/pesticides/>); Farm Chemicals Handbook 2000 volume 86, Meister Publishing Company, 2000; B. Hock, C. Fedtke, R. R. Schmidt, Herbicide [Herbicides], Georg Thieme Verlag, Stuttgart 1995; W. H. Ahrens, Herbicide Handbook, 7th

edition, Weed Science Society of America, 1994; and K. K. Hatzios, Herbicide Handbook, Supplement for the 7th edition, Weed Science Society of America, 1998. 2,2,5-Trimethyl-3-(dichloroacetyl)-1,3-oxazolidine [CAS No. 52836-31-4] is also referred to as R-29148. 4-(Dichloroacetyl)-1-oxa-4-azaspiro[4.5]decane [CAS No. 71526-07-3] is also referred to as AD-67 and MON 4660.

The assignment of the active compounds to the respective mechanisms of action is based on current knowledge. If several mechanisms of action apply to one active compound, this substance was only assigned to one mechanism of action.

It is generally preferred to use the compounds of the invention in combination with herbicides that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate employed. It is further generally preferred to apply the compounds of the invention and other complementary herbicides at the same time, either as a combination formulation or as a tank mix.

In another embodiment, the present invention refers to a method for identifying a CESA-inhibiting herbicide by using a wildtype or mutated CESA encoded by a nucleic acid which comprises the nucleotide sequence of SEQ ID NO: 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, or 77, or a variant or derivative thereof.

Said method comprises the steps of:

- a) generating a transgenic cell or plant comprising a nucleic acid encoding a wildtype or mutated CESA, wherein the wildtype or mutated CESA is expressed;
- b) applying a CESA-inhibiting herbicide to the transgenic cell or plant of a) and to a control cell or plant of the same variety;
- c) determining the growth or the viability of the transgenic cell or plant and the control cell or plant after application of said CESA-inhibiting herbicide, and
- d) selecting "CESA-inhibiting herbicides" which confer reduced growth to the control cell or plant as compared to the growth of the transgenic cell or plant.

As described above, the present invention teaches compositions and methods for increasing the CESA-inhibiting tolerance of a crop plant or seed as compared to a wild-type variety of the plant or seed. In a preferred embodiment, the CESA-inhibiting tolerance of a crop plant or seed is increased such that the plant or seed can withstand a CESA-inhibiting herbicide application of preferably approximately 1-1000 g ai ha⁻¹, more preferably 1-200 g ai ha⁻¹, even more preferably 5-150 g ai ha⁻¹, and most preferably 10-100 g ai ha⁻¹. As used herein, to "withstand" a CESA-inhibiting herbicide application means that the plant is either not killed or only moderately injured by such application. It will be understood by the person skilled in the art that the application rates may vary, depending on the environmental conditions such as temperature or humidity, and depending on the chosen kind of herbicide (active ingredient ai).

Pre- and/or Post-emergent weed control methods useful in various embodiments hereof utilize about $>0.3x$ application rates of CESA-inhibiting herbicides; in some embodiments, this can be about, for example, $>0.3x$, $>0.4x$, $>0.5x$, $>0.6x$, $>0.7x$, $>0.8x$, $>0.9x$, or $>1x$ of CESA-inhibiting herbicides. In one embodiment, CESA-inhibiting herbicides-tolerant plants of the present invention have tolerance to a pre- and / or post-emergent application of a CESA-inhibiting herbicides at an amount of about 25 to about 500 g ai/ha. In some embodiments, wherein the CESA-inhibiting herbicides-tolerant plant is a dicot (e.g., soy, cotton), the pre- and/or post-emergent application of the CESA-inhibiting herbicides is at an amount of about 25 - 250 g ai/ha. In another embodiment, wherein the CESA-inhibiting herbicides-tolerant plant is a monocot (e.g., maize, rice, sorghum), the pre- and/or post-emergent application of the CESA-inhibiting herbicides is at an amount of about 50 - 500 g ai/ha. In other embodiments, wherein the CESA-inhibiting herbicides-tolerant plant is a Brassica (e.g., canola), the pre- and/or post-emergent application of the CESA-inhibiting herbicides is at an amount of about 25-200 g ai/ha. In pre- and/or post-emergent weed control methods hereof, in some embodiments, the method can utilize CESA-inhibiting herbicides application rates at pre-emergent and /or about 7 to 10 days post-emergent. In another embodiment, the application rate can exceed $8x$ CESA-inhibiting herbicides; in some embodiments, the rate can be up to $4x$ CESA-inhibiting herbicides, though more typically it will be about $2.5x$ or less, or about $2x$ or less, or about $1x$ or less.

Furthermore, the present invention provides methods that involve the use of at least one CESA-inhibiting herbicide, optionally in combination with one or more herbicidal compounds B, and, optionally, a safener C, as described in detail supra.

In these methods, the CESA-inhibiting herbicide can be applied by any method known in the art including, but not limited to, seed treatment, soil treatment, and foliar treatment. Prior to application, the CESA-inhibiting herbicide can be converted into the customary formulations, for example solutions, emulsions, suspensions, dusts, powders, pastes and granules. The use form depends on the particular intended purpose; in each case, it should ensure a fine and even distribution of the compound according to the invention.

By providing plants having increased tolerance to CESA-inhibiting herbicide, a wide variety of formulations can be employed for protecting plants from weeds, so as to enhance plant growth and reduce competition for nutrients. A CESA-inhibiting herbicide can be used by itself for pre-emergence, post-emergence, pre-planting, and at-planting control of weeds in areas surrounding the crop plants described herein, or a CESA-inhibiting herbicide formulation can be used that contains other additives. The CESA-inhibiting herbicide can also be used as a seed treatment. Additives found in a CESA-inhibiting herbicide formulation include other herbicides, detergents, adjuvants, spreading agents, sticking agents, stabilizing agents, or the like. The CESA-inhibiting herbicide formulation can be a wet or dry preparation and can include, but is not limited to, flowable powders, emulsifiable concentrates, and liquid concentrates. The CESA-inhibiting herbicide and herbicide

formulations can be applied in accordance with conventional methods, for example, by spraying, irrigation, dusting, or the like.

Suitable formulations are described in detail in PCT/EP2009/063387 and
 5 PCT/EP2009/063386.

As disclosed herein, the CESA nucleic acids of the invention find use in enhancing the CBI herbicide tolerance of plants that comprise in their genomes a gene encoding a herbicide-tolerant wildtype or mutated CESA protein. Such a gene may be an
 10 endogenous gene or a transgene, as described above. Additionally, in certain embodiments, the nucleic acids of the present invention can be stacked with any combination of polynucleotide sequences of interest in order to create plants with a desired phenotype. For example, the nucleic acids of the present invention may be stacked with any other polynucleotides encoding polypeptides having pesticidal and/or
 15 insecticidal activity, such as, for example, the *Bacillus thuringiensis* toxin proteins (described in U.S. Patent Nos. 5,366,892; 5,747,450; 5,737,514; 5,723,756; 5,593,881; and Geiser et al (1986) Gene 48: 109), 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), Glyphosate acetyl transferase (GAT), cytochrome P450 monooxygenase, phosphinothricin acetyltransferase (PAT), Acetohydroxyacid synthase (AHAS; EC
 20 4.1.3.18, also known as acetolactate synthase or ALS), hydroxyphenyl pyruvate dioxygenase (HPPD), Phytoene desaturase (PD), Protoporphyrinogen oxidase (PPO) and dicamba degrading enzymes as disclosed in WO 02/068607, or phenoxyaceticacid- and phenoxypropionicacid-derivative degrading enzymes as disclosed in WO 2008141154 or WO 2005107437. The combinations generated can also include multiple
 25 copies of any one of the polynucleotides of interest.

Consequently, Herbicide-tolerant plants of the invention can be used in conjunction with an herbicide to which they are tolerant. Herbicides can be applied to the plants of the invention using any techniques known to those skilled in the art. Herbicides can be
 30 applied at any point in the plant cultivation process. For example, herbicides can be applied pre-planting, at planting, pre-emergence, post-emergence or combinations thereof. Herbicides may be applied to seeds and dried to form a layer on the seeds.

In some embodiments, seeds are treated with a safener, followed by a post- emergent application of a CESA-inhibiting herbicides. In one embodiment, the post-emergent application of the CESA-inhibiting herbicides is about 7 to 10 days following planting of safener-treated seeds. In some embodiments, the safener is cloquintocet, dichlormid, fluxofenim, or combinations thereof.

40 Methods of controlling weeds or undesired vegetation

In other aspects, the present invention provides a method for controlling weeds at a locus for growth of a plant or plant part thereof, the method comprising: applying a composition

comprising a CESA-inhibiting herbicides to the locus.

In some aspects, the present invention provides a method for controlling weeds at a locus for growth of a plant, the method comprising: applying an herbicide composition comprising
5 CESA-inhibiting herbicides to the locus; wherein said locus is: (a) a locus that contains: a plant or a seed capable of producing said plant; or (b) a locus that is to be after said applying is made to contain the plant or the seed; wherein the plant or the seed comprises in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a wildtype or mutated CESA polypeptide encoded
10 by the polynucleotide, the expression of the wildtype or mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides.

Herbicide compositions hereof can be applied, e.g., as foliar treatments, soil treatments, seed treatments, or soil drenches. Application can be made, e.g., by spraying, dusting,
15 broadcasting, or any other mode known useful in the art.

In one embodiment, herbicides can be used to control the growth of weeds that may be found growing in the vicinity of the herbicide-tolerant plants invention. In embodiments of this type, an herbicide can be applied to a plot in which herbicide-tolerant plants of the
20 invention are growing in vicinity to weeds. An herbicide to which the herbicide-tolerant plant of the invention is tolerant can then be applied to the plot at a concentration sufficient to kill or inhibit the growth of the weed. Concentrations of herbicide sufficient to kill or inhibit the growth of weeds are known in the art and are disclosed above.

In other embodiments, the present invention provides a method for controlling weeds in the vicinity of a CESA-inhibiting herbicides-tolerant plant of the invention. The method comprises applying an effective amount of a CESA-inhibiting herbicides to the weeds and to the herbicide- tolerant plant, wherein the plant has increased tolerance to CESA-inhibiting herbicide when compared to a wild-type plant. In some embodiments, the CESA-inhibiting
25 herbicides-tolerant plants of the invention are preferably crop plants, including, but not limited to, sunflower, alfalfa, Brassica sp., soybean, cotton, safflower, peanut, tobacco, tomato, potato, wheat, rice, maize, sorghum, barley, rye, millet, and sorghum.
30

In other aspects, herbicide(s) (e.g., CESA-inhibiting herbicides) can also be used as a seed
35 treatment. In some embodiments, an effective concentration or an effective amount of herbicide(s), or a composition comprising an effective concentration or an effective amount of herbicide(s) can be applied directly to the seeds prior to or during the sowing of the seeds. Seed Treatment formulations may additionally comprise binders and optionally colorants.

40

Binders can be added to improve the adhesion of the active materials on the seeds after treatment. In one embodiments, suitable binders are block copolymers EO/PO surfactants

but also polyvinylalcohols, polyvinylpyrrolidones, polyacrylates, polymethacrylates, polybutenes, polyisobutylenes, polystyrene, polyethyleneamines, polyethyleneamides, polyethyleneimines (Lupasol(R), Polymin(R)), polyethers, polyurethans, polyvinylacetate, tylose and copolymers derived from these polymers. Optionally, also colorants can be included in the formulation. Suitable colorants or dyes for seed treatment formulations are Rhodamin B, C.I. Pigment Red 112, C.I. Solvent Red 1, pigment blue 15:4, pigment blue 15:3, pigment blue 15:2, pigment blue 15: 1, pigment blue 80, pigment yellow 1, pigment yellow 13, pigment red 1 12, pigment red 48:2, pigment red 48: 1, pigment red 57: 1, pigment red 53:1, pigment orange 43, pigment orange 34, pigment orange 5, pigment green 36, pigment green 7, pigment white 6, pigment brown 25, basic violet 10, basic violet 49, acid red 51, acid red 52, acid red 14, acid blue 9, acid yellow 23, basic red 10, basic red 108.

The term seed treatment comprises all suitable seed treatment techniques known in the art, such as seed dressing, seed coating, seed dusting, seed soaking, and seed pelleting. In one embodiment, the present invention provides a method of treating soil by the application, in particular into the seed drill: either of a granular formulation containing the CESA-inhibiting herbicides as a composition/formulation (e.g., a granular formulation), with optionally one or more solid or liquid, agriculturally acceptable carriers and/or optionally with one or more agriculturally acceptable surfactants. This method is advantageously employed, for example, in seedbeds of cereals, maize, cotton, and sunflower.

The present invention also comprises seeds coated with or containing with a seed treatment formulation comprising CESA-inhibiting herbicides and at least one other herbicide such as, e.g., an AHAS-inhibitor selected from the group consisting of amidosulfuron, azimsulfuron, bensulfuron, chlorimuron, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethametsulfuron, ethoxysulfuron, flazasulfuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriminobac, propoxycarbazone, flucarbazone, pyribenzoxim, pyriftalid and pyriithiobac.

The term "coated with and/or containing" generally signifies that the active ingredient is for the most part on the surface of the propagation product at the time of application, although a greater or lesser part of the ingredient may penetrate into the propagation product, depending on the method of application. When the said propagation product is (re)planted, it may absorb the active ingredient.

In some embodiments, the seed treatment application with CESA-inhibiting herbicides or

with a formulation comprising the CESA-inhibiting herbicides is carried out by spraying or dusting the seeds before sowing of the plants and before emergence of the plants.

5 In other embodiments, in the treatment of seeds, the corresponding formulations are applied by treating the seeds with an effective amount of CESA-inhibiting herbicides or a formulation comprising the CESA-inhibiting herbicides.

10 In other aspects, the present invention provides a method for combating undesired vegetation or controlling weeds comprising contacting the seeds of the CESA-inhibiting herbicides-tolerant plants of the present invention before sowing and/or after pregermination with CESA-inhibiting herbicides. The method can further comprise sowing the seeds, for example, in soil in a field or in a potting medium in greenhouse. The method finds particular use in combating undesired vegetation or controlling weeds in the immediate vicinity of the seed. The control of undesired vegetation is understood as the
15 killing of weeds and/or otherwise retarding or inhibiting the normal growth of the weeds. Weeds, in the broadest sense, are understood as meaning all those plants which grow in locations where they are undesired.

The weeds of the present invention include, for example, dicotyledonous and
20 monocotyledonous weeds. Dicotyledonous weeds include, but are not limited to, weeds of the genera: Sinapis, Lepidium, Galium, Stellaria, Matricaria, Anthemis, Galinsoga, Chenopodium, Urtica, Senecio, Amaranthus, Portulaca, Xanthium, Convolvulus, Ipomoea, Polygonum, Sesbania, Ambrosia, Cirsium, Carduus, Sonchus, Solarium, Rorippa, Rotala, Lindernia, Lamium, Veronica, Abutilon, Emex, Datura, Viola, Galeopsis, Papaver,
25 Centaurea, Trifolium, Ranunculus, and Taraxacum. Monocotyledonous weeds include, but are not limited to, weeds of the genera: Echinochloa, Setaria, Panicum, Digitaria, Phleum, Poa, Festuca, Eleusine, Brachiaria, Lolium, Bromus, Avena, Cyperus, Sorghum, Agropyron, Cynodon, Monochoria, Fimbristylis, Sagittaria, Eleocharis, Scirpus, Paspalum, Ischaemum, Sphenoclea, Dactyloctenium, Agrostis, Alopecurus, and Apera.

30 In addition, the weeds of the present invention can include, for example, crop plants that are growing in an undesired location. For example, a volunteer maize plant that is in a field that predominantly comprises soybean plants can be considered a weed, if the maize plant is undesired in the field of soybean plants.

35 In other embodiments, in the treatment of seeds, the corresponding formulations are applied by treating the seeds with an effective amount of CESA-inhibiting herbicides or a formulation comprising the CESA-inhibiting herbicides.

In still further aspects, treatment of loci, plants, plant parts, or seeds of the present invention
40 comprises application of an agronomically acceptable composition that does not contain an A.I. In one embodiment, the treatment comprises application of an agronomically acceptable composition that does not contain a CESA-inhibiting herbicides A.I. In some

embodiments, the treatment comprises application of an agronomically acceptable composition that does not contain a CESA-inhibiting herbicides A.L, wherein the composition comprises one or more of agronomically-acceptable carriers, diluents, excipients, plant growth regulators, and the like. In other embodiments, the treatment

5 comprises application of an agronomically acceptable composition that does not contain a CESA-inhibiting herbicides A.I., wherein the composition comprises an adjuvant. In one embodiment, the adjuvant is a surfactant, a spreader, a sticker, a penetrant, a drift-control agent, a crop oil, an emulsifier, a compatibility agent, or combinations thereof.

10 It should also be understood that the foregoing relates to preferred embodiments of the present invention and that numerous changes may be made therein without departing from the scope of the invention. The invention is further illustrated by the following examples, which are not to be construed in any way as imposing limitations upon the scope thereof. On the contrary, it is to be clearly understood that resort may be had to various other

15 embodiments, modifications, and equivalents thereof, which, after reading the description herein, may suggest themselves to those skilled in the art without departing from the spirit of the present invention and/or the scope of the appended claims.

EXAMPLES

20

EXAMPLE 1: Identification of cellulose biosynthesis inhibitor (CBI; CESA inhibitor) resistant plants

For selection of cellulose biosynthesis inhibitor resistant plants EMS mutagenized seed populations of *Arabidopsis thaliana* are used. EMS mutagenized seed populations are

25 either bought from Lehle Seeds (1102 South Industrial Blvd. Suite D, Round Rock, Texas USA) or are generated as described elsewhere (Heim DR, *et. al.* (1989) Plant Physiol. 90: 146-150). Causative mutations in CESA wildtype sequences (e.g. SEQ ID NO: 1 or 3) are identified as described by Scheible WR *et. al.* (2001, Proc. Natl. Acad. of Sci. 98: 10079-10084), McCourt *et. al.* (WO2013/142968) or with the use of next generation sequencing

30 methods as described by Austin RS, *et. al.* (2011) Plant Journal 67: 715-725.

Selected *Arabidopsis thaliana* lines were assayed for improved resistance to azines like 6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine; 6-(1-fluoro-1-methyl-ethyl)-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine; 6-cyclohexyl-N2-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine; 6-(2,6-difluorophenyl)-N2-(2,3,5,6-

35 tetrafluorophenyl)-1,3,5-triazine-2,4-diamine; N2-(4-chloro-3,5,6-trifluoro-2-pyridyl)-6-(1-fluoro-1-methyl-ethyl)-1,3,5-triazine-2,4-diamine in 48-well plates. Therefore, M2 or M3 seeds are surface sterilized by stirring for 5 min in ethanol + water (70+30 by volume), rinsing one time with ethanol + water (70+30 by volume) and two times with sterile, deionized water. The seeds are resuspended in 0.1% agar dissolved in water (w/v) Four to

40 five seeds per well are plated on solid nutrient medium consisting of half-strength murashige skoog nutrient solution, pH 5.8 (Murashige and Skoog (1962) *Physiologia Plantarum* 15: 473-497). Compounds are dissolved in dimethylsulfoxid (DMSO) and added

to the medium prior solidification (final DMSO concentration 0.1%). Multi well plates are incubated in a growth chamber at 22°C, 75% relative humidity and 110 $\mu\text{mol Phot} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ with 14 : 10 h light : dark photoperiod. Growth inhibition is evaluated seven to ten days after seeding in comparison to wild type plants. Tolerance factors are calculated based on IC50 values of growth inhibition of transformed versus non-transformed Arabidopsis plants (Table 3).

Table 3: Tolerance of cellulose synthase mutant lines compared to wildtype plants. Relative tolerance rates of mutant Arabidopsis lines compared to a wildtype Arabidopsis plants (**wildtype** = 1.0), treated with various cellulose biosynthesis inhibitors. Growth inhibition is evaluated seven to ten days after seeding in comparison to wild type plants.

Mutant	6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine	6-(1-fluoro-1-methyl-ethyl)-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine	6-cyclohexyl-N2-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine	6-(2,6-difluorophenyl)-N2-(2,3,5,6-tetrafluorophenyl)-1,3,5-triazine-2,4-diamine	N2-(4-chloro-3,5,6-trifluoro-2-pyridyl)-6-(1-fluoro-1-methyl-ethyl)-1,3,5-triazine-2,4-diamine
wildtype	1	1	1	1	1
Cesa3_S1040L	56	2	10	13	4
Cesa3_S1037F	75	4	1	1	10
Cesa3_S983F	75	1	1	1	4
Cesa1_G1013R	25	4	4	13	10
Cesa1_P1010L	3	1	1	2	1
Cesa1_G1009D	3	2	1	1	1

Additionally, M2 or M3 *Arabidopsis* plants are tested for improved tolerance to cellulose biosynthesis-inhibiting herbicides in greenhouse studies (e. g. with the following cellulose biosynthesis-inhibiting herbicides: 6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine; 6-(1-fluoro-1-methyl-ethyl)-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine; N4-(4-chloro-3,5,6-trifluoro-2-pyridyl)-6-(1-fluoro-1-methyl-ethyl)-1,3,5-triazine-2,4-diamine; 6-(2,6-difluorophenyl)-N4-(2,3,5,6-tetrafluorophenyl)-1,3,5-triazine-2,4-diamine; N4-(4-bromo-2,3,5,6-tetrafluoro-phenyl)-6-(1-fluoro-1-methyl-ethyl)-1,3,5-triazine-2,4-diamine.

Table 4: Phytotox values of cellulose synthase mutant lines to various cellulose biosynthesis inhibitors when treated pre-emergent. Shown are phytotox values on a scale from 0 – 100, were 100 is 100% damage.

compound	g ai/ha	A. thaliana WT L. erecta	Cesa3_S1040L, fpx1-1	Cesa3_S1037F, fpx1-2	Cesa3_S983F, fpx1-3	Cesa1_G1013R, fpx2-1	Cesa1_P1010L, fpx2-2	Cesa1_G1009D, fpx2-3
6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine	15.63	100	40	5	40	15	98	88
	3.91	99	25	15	20	25	63	58
	0.98	80	20	23	5	15	33	25
	0.25	38	5	0	0	5	0	5
1-(m-tolyl)-5-phenyl-1,2,4-triazole-3-carboxamide	250	100	25	68	30	15	35	45
	62.5	100	0	23	5	20	13	30
	15.63	63	30	30	8	15	20	18
	3.91	0	38	0	0	5	0	0
6-(1-fluoro-1-methyl-ethyl)-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine	3.91	100	98	80	98	97	100	100
	0.98	100	75	30	75	63	99	97
	0.25	78	65	20	45	65	50	78
	0.0625	38	20	10	78	13	43	65
N4-(4-chloro-3,5,6-trifluoro-2-pyridyl)-6-(1-fluoro-1-methyl-ethyl)-1,3,5-triazine-2,4-diamine	3.91	99	53	25	13	80	98	99
	0.98	83	78	25	48	70	78	93
	0.25	60	15	15	55	50	25	43
	0.0625	45	13	5	23	40	50	30
6-(2,6-difluorophenyl)-N4-(2,3,5,6-tetrafluorophenyl)-1,3,5-triazine-2,4-diamine	62.5	100	28	99	33	40	98	95
	15.63	90	25	73	43	28	65	18
	3.91	70	8	65	35	35	35	38
	0.98	30	13	28	18	15	48	65
N4-(4-bromo-2,3,5,6-tetrafluoro-phenyl)-6-(1-fluoro-1-methyl-ethyl)-1,3,5-triazine-2,4-diamine	15.63	100	93	30	45	73	99	99
	3.91	97	63	33	25	35	88	68
	0.98	35	5	18	23	10	25	35
	0.25	8	13	28	35	10	25	50

Table 5: Phytotox values of cellulose synthase mutant lines to various cellulose biosynthesis inhibitors when treated post-emergent, 6 weeks after sowing. Shown are phytotox values on a scale from 0 – 100, where 100 is 100% damage.

5

compound	g ai/ha	A. thaliana WT L. erecta	Cesa3_S1040L, fpx1-1	Cesa3_S1037F, fpx1-2	Cesa3_S983F, fpx1-3	Cesa1_G1013R, fpx2-1	Cesa1_P1010L, fpx2-2	Cesa1_G1009D, fpx2-3
6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-	63	98	68	78	73	80	93	75
	31.5	78	35	70	55	65	78	73

diamine	15.75	80	5	58	65	55	70	63
	7,88	68	0	55	25	38	63	45
1-(m-tolyl)-5-phenyl-1,2,4-triazole-3-carboxamide								
	250	95	55	68	53	40	48	45
	125	83	23	50	23	33	15	33
	62.5	78	0	30	28	23	15	23

EXAMPLE 2: Identification of homologue cellulose synthase isoforms in crop plants

To identify homologue cellulose synthase genes from soy, corn and rice, BLAST searches using the protein sequences of Arabidopsis thaliana cellulose synthase isoforms were performed (Altschul et al. (1990) J Mol Biol 215: 403-10). Cellulose synthase protein encoding genes from corn, soy and rice were analyzed regarding their phylogenetic relationship by the R software library phangorn (Schliep KP. (2011) Bioinformatics 27: 592-593). Bootstrap analyses to statistically confirm monophyletic groups were calculated. In addition genes were classified by their expression level and expression pattern (Hruz T, et. al. (2008) Adv. in Bioinformatics 2008: 1-5) (Figure 1 - 3) and selected for plant transformation.

EXAMPLE 3: Tissue Culture Conditions.

An *in vitro* tissue culture mutagenesis assay is developed to isolate and characterize plant tissue (e.g., maize, rice) that is tolerant to cellulose synthase inhibiting herbicides, (e. g. 6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine, and photosynthesis inhibitor diuron as negative control). The assay utilizes the somaclonal variation that is found in *in vitro* tissue culture. Spontaneous mutations derived from somaclonal variation can be enhanced by chemical mutagenesis (e. g. ethylmethane sulfonate; ethylethane sulfonate, N-nitroso-N-ethyl urea, ethylnitrosourea, nitrose acid, bromouracil, 2.-aminopurine, 5-fluorodeoxyuridine, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine) and subsequent selection in a stepwise manner, on increasing concentrations of herbicide.

The present invention provides tissue culture conditions for encouraging growth of friable, embryogenic maize or rice callus that is regenerable. Calli are initiated from 4 different maize or rice cultivars encompassing *Zea mays* and Japonica (Taipei 309, Nipponbare, Koshihikari) and Indica (Indica 1) varieties, respectively. Seeds are surface sterilized in 70% ethanol for approximately 1 min followed by 20% commercial Clorox bleach for 20 minutes. Seeds are rinsed with sterile water and plated on callus induction media. Various callus induction media are tested. The ingredient lists for the media tested are presented in Table 6.

Table 6

Ingredient	Supplier	R001M	R025M	R026M	R327M	R008M	MS711R
B5 Vitamins	Sigma					1.0 X	
MS salts	Sigma			1.0 X	1.0 X	1.0 X	1.0 X
MS Vitamins	Sigma			1.0 X	1.0 X		

Ingredient	Supplier	R001M	R025M	R026M	R327M	R008M	MS711R
N6 salts	Phytotech	4.0 g/L	4.0g/L				
N6 vitamins	Phytotech	1.0 X	1.0 X				
L-Proline	Sigma	2.9 g/L	0.5 g/L				1.2 g/L
Casamino Acids	BD	0.3 g/L	0.3 g/L	2 g/L			
Casein Hydrolysate	Sigma						1.0 g/L
L-Asp Monohydrate	Phytotech						150 mg/L
Nicotinic Acid	Sigma						0.5 mg/L
Pyridoxine HCl	Sigma						0.5 mg/L
Thiamine HCl	Sigma						1.0 mg/L
Myo-inositol	Sigma						100 mg/L
MES	Sigma	500 mg/L	500 mg/L	500 mg/L	500 mg/L	500 mg/L	500 mg/L
Maltose	VWR	30 g/L	30 g/L	30 g/L	30 g/L		
Sorbitol	Duchefa			30 g/L			
Sucrose	VWR					10 g/L	30 g/L
NAA	Duchefa					50 µg/L	
2,4-D	Sigma	2.0 mg/L					1.0 mg/L
MgCl ₂ ·6H ₂ O	VWR					750 mg/L	
→pH		5.8	5.8	5.8	5.8	5.8	5.7
Gelrite	Duchefa	4.0 g/L				2.5 g/L	
Agarose Type1	Sigma		7.0 g/L	10 g/L	10 g/L		
→Autoclave		15 min	15 min	15 min	15 min	15 min	20 min
Kinetin	Sigma		2.0 mg/L	2.0 mg/L			
NAA	Duchefa		1.0 mg/L	1.0 mg/L			
ABA	Sigma		5.0 mg/L				
Cefotaxime	Duchefa		0.1 g/L	0.1 g/L	0.1 g/L		
Vancomycin	Duchefa		0.1 g/L	0.1 g/L	0.1 g/L		
G418 Disulfate	Sigma		20 mg/L	20 mg/L	20 mg/L		

R001M callus induction media was selected after testing numerous variations. Cultures were kept in the dark at 30°C. Embryogenic callus was subcultured to fresh media after 10-14 days.

5

EXAMPLE 4: Selection of Herbicide-tolerant Calli.

Once tissue culture conditions were determined, further establishment of selection conditions are established through the analysis of tissue survival in kill curves with azine

herbicides e. g. like 6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine, and photosynthesis inhibitor diuron as negative control. Careful consideration of accumulation of the herbicide in the tissue, as well as its persistence and stability in the cells and the culture media are performed. Through these experiments, a sub-lethal dose is established for the initial selection of mutated material. After the establishment of the starting dose of azine herbicides like e. g. 6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine, and photosynthesis inhibitor diuron as negative control in selection media, the tissues were selected in a step-wise fashion by increasing the concentration of the cellulose synthase inhibitor with each transfer until cells are recovered that grew vigorously in the presence of toxic doses. The resulting calli are further subcultured every 3-4 weeks to R001M with selective agent. Over 26,000 calli are subjected to selection for 4-5 subcultures until the selective pressure is above toxic levels as determined by kill curves and observations of continued culture. Alternatively, liquid cultures initiated from calli in MS711R with slow shaking and weekly subcultures. Once liquid cultures are established, selection agent is added directly to the flask at each subculture. Following 2-4 rounds of liquid selection, cultures are transferred to filters on solid R001M media for further growth.

EXAMPLE 5: Regeneration of Plants.

Tolerant tissue is regenerated and characterized molecularly for cellulose synthase gene sequences mutations. In addition, genes involved directly and/or indirectly in cell wall biosynthesis and/or metabolism pathways are also sequenced to characterize mutations. Finally, enzymes that change the fate (e.g. metabolism, translocation, transportation) are also sequenced to characterize mutations. Following herbicide selection, calli are regenerated using a media regime of R025M for 10 – 14 days, R026M for ca. 2 weeks, R327M until well formed shoots were developed, and R008S until shoots are well rooted for transfer to the greenhouse. Regeneration was carried out in the light. No selection agent is included during regeneration. Once strong roots are established, M0 regenerants are transplant to the greenhouse in square or round pots. Transplants are maintained under a clear plastic cup until they were adapted to greenhouse conditions. The greenhouse was set to a day/night cycle of 27°C/21°C (80°F/70°F) with 600W high pressure sodium lights supplementing light to maintain a 14 hour day length. Plants are watered according to need, depending in the weather and fertilized daily.

EXAMPLE 6: Sequence Analysis.

Leaf tissue is collected from clonal plants separated for transplanting and analyzed as individuals. Genomic DNA is extracted using a Wizard® 96 Magnetic DNA Plant System kit (Promega, US Patent Nos. 6,027,945 & 6,368,800) as directed by the manufacturer. Isolated DNA is PCR amplified using the appropriate forward and reverse primer.

PCR amplification is performed using Hotstar Taq DNA Polymerase (Qiagen) using touchdown thermocycling program as follows: 96°C for 15 min, followed by 35 cycles (96°C,

30 sec; 58°C – 0.2 °C per cycle, 30 sec; 72°C, 3 min and 30 sec), 10 min at 72°C. PCR products were verified for concentration and fragment size via agarose gel electrophoresis. Dephosphorylated PCR products are analyzed by direct sequence using the PCR primers (DNA Landmarks). Chromatogram trace files (.scf) are analyzed for mutation relative to the wild-type gene using Vector NTI Advance 10™ (Invitrogen). Based on sequence information, mutations are identified in several individuals. Sequence analysis is performed on the representative chromatograms and corresponding AlignX alignment with default settings and edited to call secondary peaks.

EXAMPLE 7: Engineering azine-tolerant *Arabidopsis* plants having wildtype or mutated cellulose synthase sequences.

For transformation of *Arabidopsis thaliana*, wildtype or mutated cellulose synthase sequences based on one of the following sequences SEQ ID NO: 72, 73, 74, 75, 76, or 77, are cloned with standard cloning techniques as described in Sambrook *et al.* (Molecular cloning (2001) Cold Spring Harbor Laboratory Press) in a binary vector containing resistance marker gene cassette (AHAS) and mutated cellulose synthase sequence (marked as GOI) in between ubiquitin promoter (PcUbi) and nopaline synthase terminator (NOS) sequence. Binary plasmids are introduced to *Agrobacterium tumefaciens* for plant transformation. *Arabidopsis thaliana* are transformed with wildtype or mutated cellulose synthase sequences by floral dip method as described by McElver and Singh (WO 2008/124495). Transgenic *Arabidopsis* plants are subjected to TaqMan analysis for analysis of the number of integration loci.

Transgenic *Arabidopsis thaliana* plants are assayed for improved tolerance to azine herbicides like e. g. 6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine in 48-well plates. Therefore, T2 seeds are surface sterilized by stirring for 5 min in ethanol + water (70+30 by volume), rinsing one time with ethanol + water (70+30 by volume) and two times with sterile, deionized water. The seeds are resuspended in 0.1% agar dissolved in water (w/v) Four to five seeds per well are plated on solid nutrient medium consisting of half-strength murashige skoog nutrient solution, pH 5.8 (Murashige and Skoog (1962) *Physiologia Plantarum* 15: 473-497). Compounds are dissolved in dimethylsulfoxid (DMSO) and added to the medium prior solidification (final DMSO concentration 0.1%). Multi well plates are incubated in a growth chamber at 22°C, 75% relative humidity and 110 $\mu\text{mol Phot} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ with 14 : 10 h light : dark photoperiod. Growth inhibition is evaluated seven to ten days after seeding in comparison to wild type plants. Tolerance factors are calculated based on IC50 values of growth inhibition of transformed versus non-transformed *Arabidopsis* plants.

Table 7: Relative tolerance rates of transgenic *Arabidopsis* plants as compared to a non-transgenic *Arabidopsis* plant (**non-transgenic = 1.0**), treated with various cellulose biosynthesis inhibitors. Growth inhibition is evaluated seven to ten days after seeding in comparison to wild type plants.

construct	AtCESA1 wt	AtCESA1_G1013R	AtCESA3_wt	AtCESA3_S1040L	AtCESA3_S1037F
SEQ ID NO	1	1	3	3	3
substitution	-	G1013R	-	S1040L	S1037F
1-(m-tolyl)-5-phenyl-1,2,4-triazole-3-carboxamide	3	33	4	2	1333
6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine	1	3	1	1	17

Additionally, transgenic T2 *Arabidopsis* plants are tested for improved tolerance to CESA-inhibiting herbicides in greenhouse studies with azine compounds like e. g. 6-cyclopentyl-N4-(2,3,4,5,6-pentafluorophenyl)-1,3,5-triazine-2,4-diamine.

5

EXAMPLE 8: Soybean transformation and cellulose biosynthesis Inhibitor tolerance testing.

Binary vectors are generated as described in EXAMPLE 7. Soybean cv Jake are transformed as previously described by Siminszky et al., *Phytochem Rev.* 5:445-458 (2006). After regeneration, transformants are transplanted to soil in small pots, placed in growth chambers (16 hr day/ 8 hr night; 25°C day/ 23°C night; 65% relative humidity; 130-150 microE m⁻² s⁻¹) and subsequently tested for the presence of the T-DNA via Taqman analysis. After a few weeks, healthy, transgenic positive, single copy events are transplanted to larger pots and allowed to grow in the growth chamber. An optimal shoot for cutting is about 3-4 inches tall, with at least two nodes present. Each cutting is taken from the original transformant (mother plant) and dipped into rooting hormone powder (indole-3-butyric acid, IBA). The cutting is then placed in oasis wedges inside a bio-dome. The mother plant is taken to maturity in the greenhouse and harvested for seed. Wild type cuttings are also taken simultaneously to serve as negative controls. The cuttings are kept in the bio-dome for 5-7 days. 3-4 days after transfer to oasis wedges, the shoots are treated via nutrient solution with the herbicide. Typical phytotox symptoms, like club shaped root, are evaluated 3-4 days after treatment. Less or no injury of transgenic plants compared to wildtype plants are interpreted as herbicide tolerance.

EXAMPLE 9: Engineering cellulose biosynthesis inhibitor tolerant corn or rice plants having mutated cellulose synthase sequences.

Immature embryos can be transformed according to the procedure outlined in Peng et al. (WO2006/136596). Plants are tested for the presence of the T-DNA by Taqman analysis with the target being the nos terminator which is present in all constructs. Healthy looking plants are sent to the greenhouse for hardening and subsequent spray testing. The plants are individually transplanted into MetroMix 360 soil in 4" pots. Once in the greenhouse (day/night cycle of 27°C /21°C with 14 hour day length supported by 600W high pressure

sodium lights), they are allowed to grow for 14 days. Transformation of *Oryza sativa* (rice) are done by protoplast transformation as described by Peng et al. (US 6653529).

Transgenic corn and rice plants are cultivated to T1 seeds for herbicide tolerance testing.

5 **EXAMPLE 10: Demonstration of herbicide tolerance**

T0 or T1 transgenic plant of soybean, corn and rice containing cellulose synthase sequences or mutated gene variants thereof are tested for improved tolerance to herbicides in greenhouse studies and mini-plot studies with azine herbicides. For the pre-emergence treatment, the herbicides are applied directly after sowing by means of finely distributing nozzles. The containers are irrigated gently to promote germination and growth and subsequently covered with transparent plastic hoods until the plants have rooted. This cover causes uniform germination of the test plants, unless this has been impaired by the herbicides. For post emergence treatment, the test plants are first grown to a height of 3 to 15 cm, depending on the plant habit, and only then treated with the herbicides. For this purpose, the test plants are either sown directly, and grown in the same containers or they are first grown separately and transplanted into the test containers a few days prior to treatment.

Herbicide injury evaluations are taken at 2 and 3 weeks after treatment. Plant injury is rated on a scale of 0% to 100%, 0% being no injury and 100% being complete death.

The following gives a definition of the injury scores measured above:

Score	Description of injury
25 0	No Injury
1	Minimal injury, only a few patches of leaf injury or chlorosis.
2	Minimal injury with slightly stronger chlorosis. Overall growth points remain undamaged.
30 3	Slightly stronger injury on secondary leaf tissue, but primary leaf and growth points are still undamaged.
4	Overall plant morphology is slightly different, some chlorosis and necrosis in secondary growth points and leaf tissue. Stems are intact. Regrowth is highly probable within 1 week.
35 5	Overall plant morphology is clearly different, some chlorosis and necrosis on a few leaves and growth points, but primary growth point is intact. Stem tissue is still green. Regrowth is highly probably within 1 week.
6	Strong injury can be seen on the new leaflet growth. Plant has a high probability to survive only through regrowth at different growth points. Most of the leaves are chlorotic/ necrotic but stem tissue is still green. May have regrowth but with noticeable injured appearance.
40 7	Most of the active growth points are necrotic. There may be a single growth point that could survive and may be partially chlorotic or green and partially

- necrotic. Two leaves may still be chlorotic with some green; the rest of the plant including stem is necrotic.
- 8 Plant will likely die, and all growth points are necrotic. One leaf may still be chlorotic with some green. The remainder of the plant is necrotic.
- 5 9 Plant is dead.
- * Not tested

10 In some aspects, embodiments of the present invention as described herein include the following items:

1. A method for controlling weeds at a locus for growth of a plant, the method comprising: (a) applying an herbicide composition comprising CESA-inhibiting
- 15 herbicides to the locus; and (b) planting a seed at the locus, wherein the seed is capable of producing a plant that comprises in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a mutated CESA polypeptide encoded by the polynucleotide, the expression of the mutated CESA polypeptide conferring to the plant tolerance to CESA-
- 20 inhibiting herbicides, wherein the mutated CESA polypeptide comprises one or more of the following sequences:


i) [V/I] [A/V]G [V/I/F] [S/T] [Y/D/N/A]A [V/I/L] [N/S/G] [S/N]G
[Y/F/E] [Q/D/G/E/H] [S/A]WG [P/A]L [F/M/L]G [K/R] [L/V] [F/L]F (SEQ
25 ID NO: 78),

wherein the amino acid at position 5, 16, 17, and/or 20 within said motif is substituted by any other amino acid; and

ii) [V/L/I]W [S/A] [V/A/I]LL [A/S]S [I/F/V] [F/L][S/T] [L/V] [L/M/V/I]
30]WV [R/K] [I/V] [N/D]PF (SEQ ID NO: 81),

wherein the amino acid at position 8, and/or 11 within said motif is substituted by any other amino acid,

wherein the CESA-inhibiting herbicide comprises a compound having the Formula
35 (I):



wherein

A is phenyl, which is substituted by two to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₁-C₆-haloalkynyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, and (C₁-C₆-alkoxy)carbonyl;

R¹ is H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl or phenylsulfonyl,

wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl and C₁-C₆-alkoxy;

R² is H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, OH, C₁-C₆-alkoxy or C₁-C₆-alkoxy-C₁-C₆-alkyl;

R³ is H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxy;

R⁴ is H, halogen, CN, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or

R³ and R⁴ together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl and three- to six-membered heterocyclyl,

wherein the C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, or three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy; and

R⁵ is H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl or phenylsulfonyl,

wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl and C₁-C₆-alkoxy;

including their agriculturally acceptable salts or N-oxides.

2. The method of item 1, wherein the mutated CESA polypeptide comprises the following sequence [V/I]AG [V/I]S [Y/D/N]A [V/I] [N/S] [S/N]G [Y/F] [Q/D] [SWGPL] [F/M/L]G [K/R]L [F/L]F (SEQ ID NO: 79).


3. The method of item 1, wherein the mutated CESA polypeptide comprises the following sequence VAG [V/I]SYA [V/I]NSGYQSWGPL [F/M]GKL [F/L]F (motif 1c;

SEQ ID NO:80).

4. The method of any one of items 1 to 3, wherein the mutated CESA polypeptide comprises the following sequence VW [S/A] [V/A/I]LL [A/S]S [I/F] [F/L][S/T][L/V]
5 [L/M]WV [R/K] [I/V] [N/D]PF (SEQ ID NO: 82).
5. The method of any one of items 1 to 3, wherein the mutated CESA polypeptide comprises the following sequence VW [S/A] [V/A/I]LLASIFSL [L/M]WV [R/K]I [N/D]PF (SEQ ID NO: 83).
10
6. The method of any one of items 1 to 3, wherein herbicide composition is applied to the weeds and to the plant produced by the seed.

Claims:

1. A method for controlling weeds at a locus for growth of a plant, the method comprising: (a) applying an herbicide composition comprising CESA-inhibiting herbicides to the locus; and (b) planting a seed at the locus, wherein the seed is capable of producing a plant that comprises in at least some of its cells a polynucleotide operably linked to a promoter operable in plant cells, the promoter capable of expressing a mutated CESA polypeptide encoded by the polynucleotide, the expression of the mutated CESA polypeptide conferring to the plant tolerance to CESA-inhibiting herbicides, wherein the mutated CESA polypeptide comprises one or more of the following sequences:
- i) [V/I] [A/V] G [V/I/F] [S/T] [Y/D/N/A] A [V/I/L] [N/S/G] [S/N] G [Y/F/E] [Q/D/G/E/H] [S/A] W G [P/A] L [F/M/L] G [K/R] [L/V] [F/L] F (SEQ ID NO: 78),
wherein the amino acid at position 5, 16, 17, and/or 20 within said motif is substituted by any other amino acid; and
- ii) [V/L/I] W [S/A] [V/A/I] L L [A/S] S [I/F/V] [F/L] [S/T] [L/V] [L/M/V/I] W V [R/K] [I/V] [N/D] P F (SEQ ID NO: 81),
wherein the amino acid at position 8, and/or 11 within said motif is substituted by any other amino acid,
- wherein the CESA-inhibiting herbicide comprises a compound having the Formula (I):



wherein

- A is phenyl, which is substituted by two to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-haloalkenyl, C₂-C₆-alkynyl, C₁-C₆-haloalkynyl, OH, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (C₁-C₆-alkyl)sulfinyl, (C₁-C₆-alkyl)sulfonyl, amino, (C₁-C₆-alkyl)amino, di(C₁-C₆-alkyl)amino, (C₁-C₆-alkyl)carbonyl, and (C₁-C₆-alkoxy)carbonyl;
- R¹ is H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl or phenylsulfonyl,

wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl and C₁-C₆-alkoxy;

- 5 R² is H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, OH, C₁-C₆-alkoxy or C₁-C₆-alkoxy-C₁-C₆-alkyl;
- 10 R³ is H, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxy;
- 10 R⁴ is H, halogen, CN, C₁-C₆-alkyl or C₁-C₆-haloalkyl; or
- 15 R³ and R⁴ together with the carbon atom to which they are attached form a moiety selected from the group consisting of carbonyl, C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl and three- to six-membered heterocyclyl, wherein the C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, or three- to six-membered heterocyclyl is unsubstituted or substituted by one to three substituents selected from halogen, CN, C₁-C₆-alkyl and C₁-C₆-alkoxy; and
- 20 R⁵ is H, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, (C₁-C₆-alkyl)carbonyl, (C₁-C₆-alkoxy)carbonyl, (C₁-C₆-alkyl)sulfonyl or phenylsulfonyl, wherein the phenyl is unsubstituted or substituted by one to five substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₆-alkyl, C₁-C₆-haloalkyl and C₁-C₆-alkoxy;
- 25


including their agriculturally acceptable salts or N-oxides.

- 30 2. The method of claim 1, wherein the mutated CESA polypeptide comprises the following sequence [V/I]AG [V/I]S [Y/D/N]A [V/I] [N/S] [S/N]G [Y/F] [Q/D] [SWGPL [F/M/L]G [K/R]L [F/L]F (SEQ ID NO: 79).
- 35 3. The method of claim 1, wherein the mutated CESA polypeptide comprises the following sequence VAG [V/I]SYA [V/I]NSGYQSWGPL [F/M]GKL [F/L]F (motif 1c; SEQ ID NO:80).
- 40 4. The method of any one of claims 1 to 3, wherein the mutated CESA polypeptide comprises the following sequence VW [S/A] [V/A/I]LL [A/S]S [I/F] [F/L][S/T][L/V] [L/M]WV [R/K] [I/V] [N/D]PF (SEQ ID NO: 82).
5. The method of any one of claims 1 to 3, wherein the mutated CESA polypeptide comprises the following sequence VW [S/A] [V/A/I]LLASIFSL [L/M]WV [R/K]I [N/D]PF (SEQ ID NO: 83).

6. The method of any one of claims 1 to 3, wherein herbicide composition is applied to the weeds and to the plant produced by the seed.


1/13

Figure 1A




2/13

Figure 1B



3/13

Figure 1C



4/13

Figure 2A

		*	20	*	40	*	60	
At_CesA1	:	VYPITS	LIAYCILPAFLCLITDRFI	PEISNYASIWFI	LLFISIAV	TGILELRWSGV	SL	: 60
At_CesA10	:		L...M.....	NT.....	L..LC..M...	A...YASA...	K...D...AL	: 60
Glyma04g07220	:	...F.....	T.....	L.NK.....	F...M.....	V...FT.S.....		: 60
Glyma06g07320	:	...F.....	T.....	L.NK.....	F...M.....	V...FT.S.....		: 60
GRMZM2G027723	:	V.....	V.....	I..L.NK.....	GMF.....	A...FA.....	G.....	: 60
GRMZM2G112336	:	V.....	V.....	I..L.NK.....	GMF.....	A...FA.....	G.....	: 60
Os05g08370	:	V.....	V.....	I..L.NK.....	GMF.....	A...FA.....	G.....	: 60
GRMZM2G039454	:	V.....	V.....	I..L.NK.....	GAF.....	A...FA.....	G.....	: 60
At_CesA3	:	I.....	LM..T.....	V..F.NQ.....	Q...I.....	LS...L...FA.....	M.....	: 60
Glyma09g15620	:	I..V.....	LM..T.....	V..L.NK.....	Q...I.....	S...L...FA.....	M.....	: 60
Glyma15g43040	:	I..V.....	LM..T.....	V..L.NK.....	Q...I.....	S...L...FA.....	M.....	: 60
Glyma12g36570	:	I..V.A.....	LI.....	V..L.NK.....	Q...L.....	S...L...FA.....	M.....	: 60
Glyma13g27250	:	I..V.A.....	LI.....	V..L.NK.....	Q...L.....	S...L...FA.....	M.....	: 60
GRMZM2G018241	:	I..L..L.....	LI.....	I..L.GK.....	F.....	S...FA.....	M.....	: 60
GRMZM2G424832	:	I..L..L.....	LI.....	I..L.GK.....	F.....	S...FA.....	M.....	: 60
Os07g10770	:	I..L..L.....	LI..V.....	I..L.GK.....	F.....	S...FA.....	M.....	: 60
GRMZM2G150404	:	I..L..L.....	LI.....	I..L.GK.....	F.....	S...FA.....	M.....	: 60
GRMZM2G111642	:	I..L..L.....	LI.....	V..L.GK.....	K...LE.V.....	S...FA.....	M.....	: 60
Os03g59340	:	I..L..L.....	LI.....	I..L.GK.....	F.....	S...L...FA.....	M.....	: 60
At_CesA4	:	...F.....	L...TI..V.....	L.GK.....	T.N.F.....	LA...L...IA.A.....		: 60
Os10g32980	:	...F.....	L...TI..V.....	L.GK.....	TLN.L.....	A...L...IA.V.....		: 60
At_CesA8	:	...F.L.V.....	T.....	I..L.GK.....	TL..L..ML.LG.....	L...I.L.SV.....		: 60
Glyma04g06780	:	...F.L.V.....	T.....	I..L.GK.....	TL..L..AL.LG.....	L...I.L.SV.....		: 60
Glyma06g06870	:	...F.L.V.....	T.....	I..L.GK.....	TL..L..AL.LG.....	L...I.L.SV.....		: 60
Glyma05g29240	:	...F.L.V.....	S.....	I..L.GK.....	TL..V..VL.LG.....	L...II.SV.....		: 60
Glyma08g12400	:	...F.L.V.....	S.....	I..L.GK.....	TL..V..VL.LG.....	L...II.SV.....		: 60
GRMZM2G037413	:	...F.L.V.....	C.....	I..L.GK.....	TL..A.T..LG.....	M...I.L.SV.....	IG.....	: 60
GRMZM2G055795	:	...F.L.V.....	C.....	I..L.GK.....	TL..A.T..LG.....	M...I.L.SV.....	IG.....	: 60
Os01g54620	:	...F.L.V.....	C.....	I..L.GK.....	TL..A.T..LG.....	M...I.L.SV.....	IG.....	: 60
Glyma06g30850	:	...F.....	LI..TI..I..L..K..T.SVGTF	LI..A.....	FASA.....	RL..L.....		: 60
Glyma12g17730	:	...FS.....	LI..I..I..L..K..T.SVGTF	LI..A.....	FASA.....	L.....		: 60
At_CesA7	:	I..F.....	L.....	I..L..K..M.P..TF..LF..S..M..I.....				: 60
Glyma02g36720	:	...F.....	V...V...V...L..K..M.P..TF..GLY.VA..S..IA..L..K.....					: 60
Glyma17g08000	:	...F.....	V...V...V...L..K..M.P..TF..GLY.VA..S..IA..L..K.....					: 60
Glyma04g23530	:	...F.....	V...V...V...L..K..M.P..TF..GLY.VA..S..IA..L..K.....					: 60
Glyma06g30860	:	...F.....	V...V...V...L..K..M.P..TF..GLY.VA..S..IA..L..K.....					: 60
GRMZM2G142898	:	I..F..L..L...T.....	V..L.GK.....	M.S..TF..LF..A..M..FA.....	M.....			: 60
Os09g25490	:	I..F..L..L...T.....	V..L.GK.....	M.P..TF..LF..A.....	FA.....	M.....		: 60
At_CesA2	:	...W..L..V..S.....	V..L.GK.....	V.....	G.L.M.M.....	MQ.G.....	G.....	: 60
At_CesA9	:	...W..L..LV..S.....	I..L.GK.....	V.....	G.L.L.M.M.....	MQ.GKIG.....		: 60
At_CesA5	:	...W..L..LV..S.....	I..L.GK.....	V.....	L.MA.....	G.....	MQ.GKIG.....	: 60
At_CesA6	:	...W..L..V..S.....	I..L.GK.....	V.....	L.MA.....	S...I.....	MQ.GKIG.....	: 60
Glyma05g32100	:	...W..L..LV..T.....	I..L.GK.....	V.....	V.MA.....	A.....	MQ.G.....	: 60
Glyma08g15380	:	...W..L..LV..T.....	I..L.GK.....	V.....	LV.MA.....	A.....	MQ.G.....	: 60
Glyma02g08920	:	...L.....	A...V..L.GK.....	V.....	I.MA.....	A.....	MQ.G.....	: 60
Glyma16g28080	:	...L.....	A...V..L.GK.....	V.....	I.MA.....	A.....	MQ.G.....	: 60
Glyma10g36790	:	I..L..L.....	T.....	V..L.GK.....	V.....	I.MA.....	S...MQ.G.....	: 60
Glyma06g47420	:	...W.....	VV..T.....	I..L.GK.....	L..A.GM..VS..FC.SV..M.....	TV.....		: 60
GRMZM2G177631	:	...W.....	L...T.....	I..L.GK.....	T..LT.V.....	MA...C.S.....	M.....	: 60
Os07g14850	:	...W.....	L...T.....	I..L.GK.....	T..LT.V..L..MS...	C.F.....	M.....	: 60
Os07g24190	:	...F.....	L...T.....	I..L.GK.....	T..LT.V..L..MS...	C.FA.....	M.....	: 60
Os07g24190	:	...F.....	L...T.....	I..L.GK.....	T..LT.V..L..MS...	C.FA.....	M.....	: 60
Os07g24190	:	...F.....	L...T.....	I..L.GK.....	T..LT.V..L..MS...	C.FA.....	M.....	: 60
GRMZM2G025231	:	...W.....	L...T.....	I..L.GK.....	T..LN.V..L..MS...	C.FA.S.....	M.....	: 60
GRMZM2G028353	:	...W.....	L...T.....	I..L.GK.....	T..LN.V..L..MS...	C.FA.S.....	M.....	: 60
GRMZM2G113137	:	...W.....	L...T.....	I..L.GK.....	T..LN.V..L..MS...	C.FA.S.....	M.....	: 60
Os03g62090	:	...W.....	L...T.....	I..L.GK.....	T..LT.I..L..MS...	C.FA.....	M.....	: 60

5/13

Figure 2A continued

		*	80	*	100	*	120	
At_CesA1	:	EDWWRNEQFWVIGG	ISAHLF	AVFQGLLKVLAGID	---	INFIVTSKAT-DE	GDFAELLYIF	: 116
At_CesA10	:			F	---	S	V	: 116
Glyma04g07220	:				---	S	V	: 116
Glyma06g07320	:				---	S	V	: 116
GRMZM2G027723	:				---	S	V	: 116
GRMZM2G112336	:				---	S	V	: 116
Os05g08370	:				---	S	V	: 116
GRMZM2G039454	:				---	D	V	: 116
At_CesA3	:	DE	V	I	---	S	L	: 116
Glyma09g15620	:	DE	V		---	S	M	: 116
Glyma15g43040	:	DE	V		---	S	L	: 116
Glyma12g36570	:	DE	V		---	S	M	: 116
Glyma13g27250	:	DE	V		---	S	M	: 116
GRMZM2G018241	:	DE	I		---	S	M	: 116
GRMZM2G424832	:	DE	I		---	S	M	: 116
Os07g10770	:	DE	I		---	S	M	: 116
GRMZM2G150404	:	DE	I		---	S	M	: 116
GRMZM2G111642	:	DE	I		---	S	E	: 116
Os03g59340	:	DE	I		---	S	E	: 116
At_CesA4	:	N	L	V	F	V	GAS	: 117
Os10g32980	:		V		G	V	AA	: 117
At_CesA8	:	L	V	F	M	L	TAD	: 115
Glyma04g06780	:	AL	V	F	M	V	AD	: 115
Glyma06g06870	:	AL	V	F	M	V	AD	: 115
Glyma05g29240	:	L	V		M	V		: 98
Glyma08g12400	:	L	V		M	V	AE	: 115
GRMZM2G037413	:		V	I	MI	L	AD	: 115
GRMZM2G055795	:		V	I	MI	L	AD	: 115
Os01g54620	:		V	I	MI	L	AD	: 115
Glyma06g30850	:	E	SQ	SV	N	L	IMGA	: 118
Glyma12g17730	:	E	SQ	SV	N	LL	IMRA	: 117
At_CesA7	:	E		I	V	I		: 115
Glyma02g36720	:	E		V	I		D	: 115
Glyma17g08000	:	E		V	I		D	: 115
Glyma04g23530	:	E		V	I		D	: 115
Glyma06g30860	:	E		V	I		D	: 115
GRMZM2G142898	:	E		V			D	: 117
Os09g25490	:	E		V			D	: 117
At_CesA2	:	D		A	S	L	VN	: 115
At_CesA9	:	D		V	S	L	VS	: 115
At_CesA5	:	D		V		L	VE	: 115
At_CesA6	:	D		V		L	VN	: 115
Glyma05g32100	:	D		V	S	L	VN	: 115
Glyma08g15380	:	D		V	S	L	VN	: 115
Glyma02g08920	:	H		A	S	L	VN	: 115
Glyma16g28080	:	H		A	S	L	VN	: 115
Glyma10g36790	:	H		A	S	L	VN	: 115
Glyma06g47420	:	DE		V	FL	L	MF	: 114
GRMZM2G177631	:	D		V			F	: 115
Os07g14850	:	D		V	S		V	: 115
Os07g24190	:	D		V	S		L	: 115
Os07g24190	:	D		V	S		L	: 115
Os07g24190	:	D		V	S		L	: 115
GRMZM2G025231	:	D		V	S		I	: 115
GRMZM2G028353	:	D		V	S		I	: 115
GRMZM2G113137	:	D		V	S		I	: 115
Os03g62090	:	D		V	S		I	: 115

Figure 2A continued

	*	140	*	160	*	180	
At_CesA1	:	KWTALLIPPTTVLLVNLIGIVAGVSYAVNSGVQSWGPLEGKLEFFALWVLAHLYPFLKGLL	:	176			
At_CesA10	:	S.....I.....V.....I.....M.....L.....F.....V.....	:	176			
Glyma04g07220	:	S.....I.....V.....I.....I.....I.....I.....I.....	:	176			
Glyma06g07320	:	S.....I.....V.....I.....I.....I.....I.....I.....	:	176			
GRMZM2G027723	:	S.....VI.....VM.....I.....I.....I.....SI.....L.....M.....	:	176			
GRMZM2G112336	:	S.....VI.....VM.....I.....I.....I.....SI.....L.....M.....	:	176			
Os05g08370	:	S.....VI.....VM.....I.....I.....I.....SI.....L.....M.....	:	176			
GRMZM2G039454	:	T.....VI.....V.....I.....I.....I.....I.....L.....M.....	:	176			
At_CesA3	:	T.....LII.....VV.....I.....I.....I.....F.....V.....M.....	:	176			
Glyma09g15620	:	T.....LII.....VV.....I.....I.....I.....F.....I.....M.....	:	176			
Glyma15g43040	:	T.....LII.....VV.....I.....I.....I.....F.....I.....M.....	:	176			
Glyma12g36570	:	T.....LII.....VV.....I.....I.....I.....F.....I.....M.....	:	176			
Glyma13g27250	:	T.....LII.....M.....V.....I.....I.....I.....F.....I.....M.....	:	176			
GRMZM2G018241	:	T.....I.....II.....VV.....I.....I.....I.....F.....V.....M.....	:	176			
GRMZM2G424832	:	T.....I.....II.....VV.....I.....I.....I.....F.....V.....M.....	:	176			
Os07g10770	:	T.....I.....II.....VV.....I.....I.....I.....F.....V.....M.....	:	176			
GRMZM2G150404	:	T.....I.....II.....VV.....I.....I.....I.....F.....V.....M.....	:	176			
GRMZM2G111642	:	T.....I.....II.....VV.....I.....I.....I.....F.....V.....M.....	:	176			
Os03g59340	:	T.....I.....II.....VV.....I.....I.....I.....F.....V.....M.....	:	176			
At_CesA4	:	T.....LIIILMV.....V.....D.....I.....N.....G.....F.....V.....M.....	:	177			
Os10g32980	:	T.....V.....LIIII MV.....D.....N.....G.....SF.....L.....M.....	:	177			
At_CesA8	:	T.....SLII.....V.....V.....FD.....L.....K.....EA.....V.....F.....L.....M.....	:	175			
Glyma04g06780	:	T.....LIIII IV.....V.....FD.....L.....G.....E.....V.....F.....F.....M.....	:	175			
Glyma06g06870	:	T.....LIIII MV.....V.....FD.....L.....G.....E.....V.....F.....F.....M.....	:	175			
Glyma05g29240	:	T.....LII.....MV.....V.....FD.....L.....G.....E.....V.....F.....F.....M.....	:	157			
Glyma08g12400	:	T.....LIV.....MV.....V.....FD.....L.....G.....E.....V.....F.....F.....M.....	:	175			
GRMZM2G037413	:	TV.....SI.....VL.....VV.....FA.....L.....E.....V.....M.....M.....M.....	:	175			
GRMZM2G055795	:	TV.....SI.....VL.....VV.....FA.....L.....E.....V.....M.....M.....M.....	:	175			
Os01g54620	:	TV.....SI.....VL.....VV.....FD.....L.....E.....V.....M.....M.....M.....	:	175			
Glyma06g30850	:	R.....IIII.....I.....FTD.....I.....EH.....A.....L.....S.....V.....M.....	:	178			
Glyma12g17730	:	R.....IIII.....I.....FTD.....I.....EH.....A.....L.....S.....VI.....M.....	:	177			
At_CesA7	:	T.....II.....IV.....V.....ID.....I.....N.....SF.....V.....M.....	:	175			
Glyma02g36720	:	T.....I.....II.....IV.....V.....ID.....I.....N.....SF.....V.....M.....	:	175			
Glyma17g08000	:	T.....I.....II.....IV.....V.....ID.....I.....N.....SF.....V.....M.....	:	175			
Glyma04g23530	:	T.....I.....II.....IV.....V.....ID.....I.....N.....SF.....V.....M.....	:	175			
Glyma06g30860	:	T.....I.....II.....IV.....V.....ID.....I.....N.....SF.....V.....M.....	:	175			
GRMZM2G142898	:	T.....LII.....V.....V.....ID.....I.....N.....F.....V.....M.....	:	177			
Os09g25490	:	T.....LIL.....I.....V.....D.....I.....N.....SEA.....F.....V.....M.....	:	177			
At_CesA2	:	T.....LII.....I.....VIV.....D.....ISN.....D.....R.....V.....M.....	:	175			
At_CesA9	:	S.....LII.....IV.....VIV.....D.....I.....N.....D.....R.....V.....	:	175			
At_CesA5	:	S.....LII.....V.....VIV.....I.....D.....ISN.....D.....R.....F.....L.....	:	175			
At_CesA6	:	S.....M.....LII.....V.....VIV.....D.....ISN.....D.....R.....I.....	:	175			
Glyma05g32100	:	S.....M.....LIM.....IV.....V.....V.....I.....D.....I.....N.....D.....R.....L.....	:	175			
Glyma08g15380	:	S.....M.....LIM.....IV.....V.....V.....I.....D.....I.....N.....D.....R.....L.....	:	175			
Glyma02g08920	:	S.....L.....LII.....I.....VIV.....D.....I.....N.....D.....R.....V.....VM.....	:	175			
Glyma16g28080	:	S.....L.....LIL.....I.....VIV.....D.....I.....N.....D.....R.....V.....VM.....	:	175			
Glyma10g36790	:	S.....L.....LII.....I.....VIV.....D.....I.....N.....D.....R.....V.....VM.....	:	175			
Glyma06g47420	:	T.....I.....L.....VL.....I.....AV.....I.....N.....FE.....L.....LS.....L.....MI.....	:	174			
GRMZM2G177631	:	T.....L.....L.....LF.....V.....I.....N.....I.....N.....E.....F.....V.....V.....	:	175			
Os07g14850	:	T.....L.....L.....LF.....V.....N.....I.....N.....E.....F.....V.....V.....	:	175			
Os07g24190	:	T.....L.....L.....LF.....V.....N.....I.....N.....E.....F.....V.....V.....	:	175			
Os07g24190	:	T.....L.....L.....LF.....V.....N.....I.....N.....E.....F.....V.....V.....	:	175			
Os07g24190	:	T.....L.....L.....LF.....V.....N.....I.....N.....E.....F.....V.....V.....	:	175			
GRMZM2G025231	:	T.....L.....L.....LF.....V.....N.....I.....N.....E.....F.....V.....V.....	:	175			
GRMZM2G028353	:	T.....L.....L.....LF.....V.....N.....I.....N.....E.....F.....V.....V.....	:	175			
GRMZM2G113137	:	T.....L.....L.....LF.....V.....I.....N.....I.....N.....E.....F.....V.....V.....	:	175			
Os03g62090	:	T.....L.....L.....LF.....V.....N.....I.....N.....E.....F.....V.....V.....	:	175			

7/13

Figure 2 A continued

		*	200	*	
At_CesA1	:	GRQNRTPTIVIVWSVLLASIFSLWVRINPF	:	207	
At_CesA10	:	A.....	:	207	
Glyma04g07220	:	D.....	:	207	
Glyma06g07320	:	D.....	:	207	
GRMZM2G027723	:	I.....K.D.	:	207	
GRMZM2G112336	:	I.....K.D.	:	207	
Os05g08370	:	I.....K.D.	:	207	
GRMZM2G039454	:	K.....K.D.	:	207	
At_CesA3	:	V.....D.	:	207	
Glyma09g15620	:	V.....D.	:	207	
Glyma15g43040	:	V.....D.	:	207	
Glyma12g36570	:	V.....D.	:	207	
Glyma13g27250	:	V.....D.	:	207	
GRMZM2G018241	:	V.AI.....D.	:	207	
GRMZM2G424832	:	V.AI.....D.	:	207	
Os07g10770	:	V.AI.....D.	:	207	
GRMZM2G150404	:	V.AI.....D.	:	207	
GRMZM2G111642	:	K.....V.AI.....M.....D.	:	207	
Os03g59340	:	V.AI.....D.	:	207	
At_CesA4	:	VL.I.....V.....D.	:	208	
Os10g32980	:	VL.I.....V.....D.	:	208	
At_CesA8	:	L.I.....V.....V.....D.	:	206	
Glyma04g06780	:	L.....V.....V.....K.....D.	:	206	
Glyma06g06870	:	L.....V.....V.....K.....D.	:	206	
Glyma05g29240	:	VL.....V.....V.....K.....D.	:	188	
Glyma08g12400	:	L.....V.....I.....K.....D.	:	206	
GRMZM2G037413	:	VL.....V.....K.....D.	:	206	
GRMZM2G055795	:	VL.....V.....K.....D.	:	206	
Os01g54620	:	VL.....V.....K.....D.	:	206	
Glyma06g30850	:	LIVI.....V.....VD.....D.	:	209	
Glyma12g17730	:	LIVI.....V.....VD.....D.	:	208	
At_CesA7	:	VI.....D.	:	206	
Glyma02g36720	:	VI.....D.	:	206	
Glyma17g08000	:	VI.....D.	:	206	
Glyma04g23530	:	VI.....D.	:	206	
Glyma06g30860	:	VI.....D.	:	206	
GRMZM2G142898	:	V.VI.I.....D.	:	208	
Os09g25490	:	VI.....D.	:	208	
At_CesA2	:	K.DKM.....IV.....LT.....V.....D.	:	206	
At_CesA9	:	K.D.V.....IL.....LT.....V.....D.	:	206	
At_CesA5	:	K.D.M.....IL.....LT.....V.....D.	:	206	
At_CesA6	:	K.D.M.....IV.....LT.....V.....D.	:	206	
Glyma05g32100	:	K.D.M.....IL.....LT.M.....D.	:	206	
Glyma08g15380	:	K.D.M.....IL.....LT.M.....D.	:	206	
Glyma02g08920	:	K.EGV.....IL.AI.....S.....LT.....D.	:	206	
Glyma16g28080	:	K.EGV.....IL.AI.....LT.....D.	:	206	
Glyma10g36790	:	K.EGV.....IL.AI.....D.	:	206	
Glyma06g47420	:	H.....L.AI.....F.V.....K.D.	:	205	
GRMZM2G177631	:	I.....VD.....D.	:	206	
Os07g14850	:	I.....D.	:	206	
Os07g24190	:	I.....D.	:	206	
Os07g24190	:	I.....D.	:	206	
Os07g24190	:	I.....D.	:	206	
GRMZM2G025231	:	I.....D.	:	206	
GRMZM2G028353	:	I.....D.	:	206	
GRMZM2G113137	:	I.....D.	:	206	
Os03g62090	:	I.....D.	:	206	

	*	20	*	40	*	60	
At Cesa1	:	VYPITSIPLIAYCILPAFLCIITDRFIIP	EISNYAS	IWFILLFISIAVTGILELRWSGV	SI	:	60
At Cesa10	:	 L . M NT	L . LC . M . . A . YASA . . . K . D . AL	:	:	:	60
Glyma04g07220	:	. . F T LNK	F . M V . FT . S	:	:	:	60
Glyma06g07320	:	. . F T LNK	F . M V . FT . S	:	:	:	60
GRMZM2G027723	:	 V I . LNK	GMF A . FA G .	:	:	:	60
GRMZM2G112336	:	 V I . LNK	GMF A . FA G .	:	:	:	60
Oso5g08370	:	 V I . LNK	GMF A . FA G .	:	:	:	60
GRMZM2G039454	:	 V . V I . LNK	A . GAF A . FA G .	:	:	:	60
At Cesa3	:	I LM . T . V . F . NC . Q . I . .	LS . L . FA . . M . . G .	:	:	:	60
Glyma09g15620	:	I . V . . LM . T . V . LNK . Q . I . .	S . L . FA . . M . . G .	:	:	:	60
Glyma15g43040	:	I . V . . LM . T . V . LNK . Q . I . .	S . L . FA . . M . . G .	:	:	:	60
Glyma12g36570	:	I . V . A . LI . . . V . LNK . Q . L . .	S . L . FA . . M . . G .	:	:	:	60
Glyma13g27250	:	I . V . A . LI . . . V . LNK . Q . L . .	S . L . FA . . M . . G .	:	:	:	60
GRMZM2G018241	:	I . L . L . LI . . . I . LGK . F . . S . .	FA . . M . . G .	:	:	:	60
GRMZM2G424832	:	I . L . . LI . . . I . LGK . F . . S . .	FA . . M . . G .	:	:	:	60
Oso7g10770	:	I . L . . LI . V . . I . LGK . F . . S . .	FA . . M . . G .	:	:	:	60
GRMZM2G150404	:	I . L . L . LI . . . I . LGK . F . . S . .	FA . . M . . G .	:	:	:	60
GRMZM2G111642	:	I . L . . LL . . . V . LGK . K . LE . V . S . .	FA . . M . . G .	:	:	:	60
Oso3g59340	:	I . L . . LL . . . I . LGK . F . . S . L .	FA . . M . . G .	:	:	:	60
At Cesa2	:	. . W . L . V . S . . V . LGK . V	G . L . M . M MQ . G . G .	:	:	:	60
At Cesa9	:	. . W . L . LV . S . . I . LGK . V	G . L . L . M . M MQ . GKIG .	:	:	:	60
At Cesa5	:	. . W . . LV . S . . I . LGK . V	L . MA . G MQ . GK . G .	:	:	:	60
At Cesa6	:	. . W . L . V . S . . I . LGK . V	L . MA . S . . I . . MQ . GK . G .	:	:	:	60
Glyma05g32100	:	. . W . L . LV . T . . I . LGK . V	V . MA A . . MQ . G . G .	:	:	:	60
Glyma08g15380	:	. . W . L . LV . T . . I . LGK . V	LV . MA A . . MQ . G . G .	:	:	:	60
Glyma02g08920	:	. . L A . V . LGK . V	I . MA A . . MQ . G . G .	:	:	:	60
Glyma16g28080	:	. . L A . V . LGK . V	I . MA A . . MQ . G . G .	:	:	:	60
Glyma10g36790	:	I . L . L . . . T . V . LGK . V	I . MA S . . MQ . G . G .	:	:	:	60
Glyma06g47420	:	. . W . . VV . T . . I . LGK L . A . GM . VS . FC . FT . SV . M . . TV	:	:	:	:	60
GRMZM2G177631	:	. . W . . L . T . . I . LGK . T . LT . V . .	MA . . C . S M . . A .	:	:	:	60
Oso7g14850	:	. . W . . L . T . . I . LGK . T . LT . V . L . MS . . C . F . .	M . . A . : 60				
Oso7g24190	:	. . F . . L . T . . I . LGK . T . LT . V . L . MS . . C . FA . .	M . . G . : 60				
GRMZM2G025231	:	. . W . . L . T . . I . LGK . T . LN . V . L . MS . . C . FA . S .	M . . G . : 60				
GRMZM2G028353	:	. . W . . L . T . . I . LGK . T . LN . V . L . MS . . C . FA . S .	M . . G . : 60				
GRMZM2G113137	:	. . W . . L . T . . I . LGK . T . LN . V . L . MS . . C . FA . S .	M . . G . : 60				
Oso3g62090	:	. . W . . L . T . . I . LGK . T . LT . I . L . MS . . C . FA . .	M . . G . : 60				

9/13

Figure 2B continued

		*	80	*	100	*	120				
At_CesA1	:	EDWWRNEQFWVIGGT	SAHLFAVFOGLLKVL	AGIDINFTVTS	KAIDEDGDFAEL	YIFKWTIA	:	120			
At_CesA10	:			F.....	S.....	V.....	S	: 120			
Glyma04g07220	:				S.....	V.....	S	: 120			
Glyma06g07320	:				S.....	V.....	S	: 120			
GRMZM2G027723	:				S.....	V.....	S	: 120			
GRMZM2G112336	:				S.....	V.....	S	: 120			
Os05g08370	:				S.....	V.....	S	: 120			
GRMZM2G039454	:				..D.....	V.....	T	: 120			
At_CesA3	:	DE.....	V.....	I.....	S.....	L.....	T	: 120			
Glyma09g15620	:	DE.....	V.....		S.....	M.....	T	: 120			
Glyma15g43040	:	DE.....	V.....		S.....	L.....	T	: 120			
Glyma12g36570	:	DE.....	V.....		S.....	M.....	T	: 120			
Glyma13g27250	:	DE.....	V.....		S.....	M.....	T	: 120			
GRMZM2G018241	:	DE.....	I.....		S.....	M.....	T	: 120			
GRMZM2G424832	:	DE.....	I.....		S.....	M.....	T	: 120			
Os07g10770	:	DE.....	I.....		S.....	M.....	T	: 120			
GRMZM2G150404	:	DE.....	I.....		S.....	M.....	T	: 120			
GRMZM2G111642	:	DE.....	I.....		S.....	E.....	M.....	T	: 120		
Os03g59340	:	DE.....	I.....		S.....	E.....	M.....	T	: 120		
At_CesA2	:	D.....	A.S.....	L.....	VN.....	-AD.....	A.S.....	T	: 119		
At_CesA9	:	D.....	V.S.....	L.....	VS.....	-AD.....	E.S.....	S	: 119		
At_CesA5	:	D.....	V.....	L.....	VE.....	-AD.....	E.S.....	S	: 119		
At_CesA6	:	D.....	V.....	L.....	V.....	-AD.....	E.SD.....	L.....	S	: 119	
Glyma05g32100	:	D.....	V.S.....	L.....	VN.....	-AD.....	E.S.....	S	: 119		
Glyma08g15380	:	D.....	V.S.....	L.....	VN.....	-AD.....	E.S.....	S	: 119		
Glyma02g08920	:	H.....	A.S.....	L.....	VN.....	-AD.....	E.....D.....	S	: 119		
Glyma16g28080	:	H.....	A.S.....	L.....	VN.....	-AD.....	E.....	S	: 119		
Glyma10g36790	:	H.....	A.S.....	L.....	VN.....	-ADG.....		L.....	S	: 119	
Glyma06g47420	:	DE.....	V.....	FL.....	L.MF.....	VK.....	I.A.....	--VD.....	KEHSNMFAL.....	T	: 118
GRMZM2G177631	:	D.....	V.....		F.....	S.....	-GD.....	EE.....	S.....	T	: 119
Os07g14850	:	D.....	V.S.....		V.....	S.....	-GD.....	EE.....	S.....	T	: 119
Os07g24190	:	D.....	V.S.....	L.....	I.....	S.....	G-GD.....	EE.....	S.....	T	: 119
GRMZM2G025231	:	D.....	V.S.....		I.V.....	S.....	G-GD.....	EE.....	S.....	T	: 119
GRMZM2G028353	:	D.....	V.S.....		I.V.....	S.....	G-GD.....	DE.....	S.....	T	: 119
GRMZM2G113137	:	D.....	V.S.....		I.V.....	S.....	G-GD.....	EE.....	S.....	T	: 119
Os03g62090	:	D.....	V.S.....		I.....	S.....	G-GD.....	EE.....	S.....	T	: 119

10/13

Figure 2B continued

	*	140	*	160	*	180	
At_CesA1	:	LLIPPTTVLLVNLIGIVAGVSYAVNSGYQSWGPLEFGKLEFFALWVIAHLYPFLKGLLGRON	:	180			
At_CesA10	:	I.....V.....I.....M.....L.....F.....V.....	:	180			
Glyma04g07220	:	I.....V.....I.....I.....I.....I.....I.....I.....	:	180			
Glyma06g07320	:	I.....V.....I.....I.....I.....I.....I.....I.....	:	180			
GRMZM2G027723	:	VI.....V.....M.....I.....I.....I.....I.....SI.....L.....M.....	:	180			
GRMZM2G112336	:	VI.....V.....M.....I.....I.....I.....I.....SI.....L.....M.....	:	180			
Os05g08370	:	VI.....V.....M.....I.....I.....I.....I.....SI.....L.....M.....	:	180			
GRMZM2G039454	:	VI.....V.....I.....I.....I.....I.....I.....I.....L.....M.....K.....	:	180			
At_CesA3	:	L.....I.....V.....V.....I.....I.....I.....I.....F.....V.....M.....	:	180			
Glyma09g15620	:	L.....II.....V.....V.....I.....I.....I.....I.....F.....I.....M.....	:	180			
Glyma15g43040	:	L.....II.....V.....V.....I.....I.....I.....I.....F.....I.....M.....	:	180			
Glyma12g36570	:	L.....II.....V.....V.....I.....I.....I.....I.....F.....I.....M.....	:	180			
Glyma13g27250	:	L.....II.....M.....V.....I.....I.....I.....I.....F.....I.....M.....	:	180			
GRMZM2G018241	:	I.....II.....V.....V.....I.....I.....I.....I.....F.....V.....M.....	:	180			
GRMZM2G424832	:	I.....II.....V.....V.....I.....I.....I.....I.....F.....V.....M.....	:	180			
Os07g10770	:	I.....II.....V.....V.....I.....I.....I.....I.....F.....V.....M.....	:	180			
GRMZM2G150404	:	I.....II.....V.....V.....I.....I.....I.....I.....F.....V.....M.....	:	180			
GRMZM2G111642	:	I.....II.....V.....V.....I.....I.....I.....I.....F.....V.....M.....K.....	:	180			
Os03g59340	:	I.....II.....V.....V.....I.....I.....I.....I.....F.....V.....M.....	:	180			
At_CesA2	:	L.....II.....I.....VIV.....D.....ISN.....D.....R.....V.....M.....K.....D.....	:	179			
At_CesA9	:	L.....II.....IV.....VIV.....D.....I.....N.....D.....R.....V.....K.....D.....	:	179			
At_CesA5	:	L.....II.....V.....VIV.....I.....D.....ISN.....D.....R.....F.....L.....K.....D.....	:	179			
At_CesA6	:	M.....L.....II.....V.....VIV.....D.....ISN.....D.....R.....I.....K.....D.....	:	179			
Glyma05g32100	:	M.....L.....IM.....IV.....V.....V.....D.....I.....N.....D.....R.....L.....K.....D.....	:	179			
Glyma08g15380	:	M.....L.....IM.....IV.....V.....V.....I.....D.....I.....N.....D.....R.....L.....K.....D.....	:	179			
Glyma02g08920	:	L.....L.....II.....I.....VIV.....D.....I.....N.....D.....R.....V.....VM.....K.....E.....	:	179			
Glyma16g28080	:	L.....L.....IL.....I.....VIV.....D.....I.....N.....D.....R.....V.....VM.....K.....E.....	:	179			
Glyma10g36790	:	L.....L.....II.....I.....VIV.....D.....I.....N.....D.....R.....V.....VM.....K.....E.....	:	179			
Glyma06g47420	:	I.....L.....VL.....I.....AV.....I.....N.....FE.....L.....L.....S.....L.....MI.....H.....	:	178			
GRMZM2G177631	:	L.....L.....L.....F.....V.....I.....N.....I.....N.....E.....F.....V.....V.....	:	179			
Os07g14850	:	L.....L.....L.....F.....V.....N.....I.....N.....E.....F.....V.....V.....	:	179			
Os07g24190	:	L.....L.....L.....F.....V.....N.....I.....N.....E.....F.....V.....V.....	:	179			
GRMZM2G025231	:	L.....L.....L.....F.....V.....N.....I.....N.....E.....F.....V.....V.....	:	179			
GRMZM2G028353	:	L.....L.....L.....F.....V.....N.....I.....N.....E.....F.....V.....V.....	:	179			
GRMZM2G113137	:	L.....L.....L.....F.....V.....I.....N.....I.....N.....E.....F.....V.....V.....	:	179			
Os03g62090	:	L.....L.....L.....F.....V.....N.....I.....N.....E.....F.....V.....V.....	:	179			

11/13

Figure 2B continued

		*	200		
At_CesA1	:	RTPTIVIVWSVLLASIFSELLWVRINPF	:	207	
At_CesA10	:	A.....	:	207	
Glyma04g07220	:	D..	:	207	
Glyma06g07320	:	D..	:	207	
GRMZM2G027723	:	K.D..	:	207	
GRMZM2G112336	:	K.D..	:	207	
Os05g08370	:	K.D..	:	207	
GRMZM2G039454	:	K.D..	:	207	
At_CesA3	:	V.....	:	207	
Glyma09g15620	:	V.....	:	207	
Glyma15g43040	:	V.....	:	207	
Glyma12g36570	:	V.....	:	207	
Glyma13g27250	:	V.....	:	207	
GRMZM2G018241	:	V..AI.....	:	207	
GRMZM2G424832	:	V..AI.....	:	207	
Os07g10770	:	V..AI.....	:	207	
GRMZM2G150404	:	V..AI.....	:	207	
GRMZM2G111642	:	V..AI.....M.....	:	207	
Os03g59340	:	V..AI.....	:	207	
At_CesA2	:	KM...IV...I...LT...V...	:	206	
At_CesA9	:	.V...IL...I...LT...V...	:	206	
At_CesA5	:	.M...IL...I...LT...V...	:	206	
At_CesA6	:	.M...IV...I...LT...V...	:	206	
Glyma05g32100	:	.M...IL...I...LT.M.....	:	206	
Glyma08g15380	:	.M...IL...I...LT.M.....	:	206	
Glyma02g08920	:	GV...IL..AI...S...LT.....	:	206	
Glyma16g28080	:	GV...IL..AI...LT.....	:	206	
Glyma10g36790	:	GV...IL..AI.....	:	206	
Glyma06g47420	:	L..AI.....F..V...K.D..	:	205	
GRMZM2G177631	:	I.....V.D..	:	206	
Os07g14850	:	I.....	:	206	
Os07g24190	:	I.....	:	206	
GRMZM2G025231	:	I.....	:	206	
GRMZM2G028353	:	I.....	:	206	
GRMZM2G113137	:	I.....	:	206	
Os03g62090	:	I.....	:	206	

12/13

Figure 2C

		*	20	*	40	*	60				
At_CesA1	:	VYPITS	ILPIAYC	ILPAFCL	ITDRFI	PEISNY	ASINWF	ILLFISIAV	TGILELRWSGVST	:	60
At_CesA10	:		L...M...		NT.....		L..LC.M...	A..YNSA...	...K..D.AL	:	60
Glyma04g07220	:	...F.....	...T.....	...L.NK.....	...F.M.....	...V..FT.S.....				:	60
Glyma06g07320	:	...F.....	...T.....	...L.NK.....	...F.M.....	...V..FT.S.....				:	60
GRMZM2G027723	:		V.....	...I..L.NK.....		GMF...A..FA.....				:	60
GRMZM2G112336	:		...V.....	...I..L.NK.....		GMF...A..FA.....				:	60
Os05g08370	:		...V.....	...I..L.NK.....		GMF...A..FA.....				:	60
GRMZM2G039454	:		V...V...	...I..L.NK.....	A...	GAF...A..FA.....				:	60
At_CesA3	:	I.....	LM...T...	V...F.NQ...	Q...	I.....	LS...L..FA.....	M.....	G.....	:	60
Glyma09g15620	:	I..V.....	LM...T...	V...L.NK.....	Q...	I.....	S...L..FA.....	M.....	G.....	:	60
Glyma15g43040	:	I..V.....	LM...T...	V...L.NK.....	Q...	I.....	S...L..FA.....	M.....	G.....	:	60
Glyma12g36570	:	I..V.A...	LI.....	V...L.NK.....	Q...	L.....	S...L..FA.....	M.....	G.....	:	60
Glyma13g27250	:	I..V.A...	LI.....	V...L.NK.....	Q...	L.....	S...L..FA.....	M.....	G.....	:	60
GRMZM2G018241	:	I..L...L..	LI.....	I...L.GK.....	F.....	S.....	FA.....	M.....	G.....	:	60
GRMZM2G424832	:	I..L...LI..		I...L.GK.....	F.....	S.....	FA.....	M.....	G.....	:	60
Os07g10770	:	I..L...LI..	V.....	I...L.GK.....	F.....	S.....	FA.....	M.....	G.....	:	60
GRMZM2G150404	:	I..L...L..	LI.....	I...L.GK.....	F.....	S.....	FA.....	M.....	G.....	:	60
GRMZM2G111642	:	I..L...LL..		V...L.GK.....	K...	LE.V...	S.....	FA.....	M.....	:	60
Os03g59340	:	I..L...LL..		I...L.GK.....	F.....	S.....	L..FA.....	M.....	G.....	:	60

		*	80	*	100	*	120			
At_CesA1	:	EDWWRNEQFWVIGG	TS	SAHLFAV	FQGLLKVLAGID	TNFTVTSKAT	DEDGDFAELYIFK	WTA	:	120
At_CesA10	:				F.....			V.....S	:	120
Glyma04g07220	:							V.....S	:	120
Glyma06g07320	:							V.....S	:	120
GRMZM2G027723	:							V.....S	:	120
GRMZM2G112336	:							V.....S	:	120
Os05g08370	:							V.....S	:	120
GRMZM2G039454	:							V.....T	:	120
At_CesA3	:	DE.....	V.....	I.....				L.....T	:	120
Glyma09g15620	:	DE.....	V.....	I.....				M.....T	:	120
Glyma15g43040	:	DE.....	V.....	I.....				L.....T	:	120
Glyma12g36570	:	DE.....	V.....	I.....				M.....T	:	120
Glyma13g27250	:	DE.....	V.....	I.....				M.....T	:	120
GRMZM2G018241	:	DE.....	I.....					M.....T	:	120
GRMZM2G424832	:	DE.....	I.....					M.....T	:	120
Os07g10770	:	DE.....	I.....					M.....T	:	120
GRMZM2G150404	:	DE.....	I.....					M.....T	:	120
GRMZM2G111642	:	DE.....	I.....		S.....		E.....	M.....T	:	120
Os03g59340	:	DE.....	I.....		S.....		E.....	M.....T	:	120

13/13

Figure 2C continued

		*		140	*		160	*		180		
At_CesA1	:	LLIPPTTV	LLVNI	IGIVAGV	SYAVNS	GYQSWG	PLE	FGKLE	FALWV	LAHLYP	FLKGLLGRON	: 180
At_CesA10	:	I	V	I	I	M	L	F	V			: 180
Glyma04g07220	:	I	V	I	I			I				: 180
Glyma06g07320	:	I	V	I	I			I				: 180
GRMZM2G027723	:	VI	V	M	I			SI	L		M	: 180
GRMZM2G112336	:	VI	V	M	I			SI	L		M	: 180
Os05g08370	:	VI	V	M	I			SI	L		M	: 180
GRMZM2G039454	:	VI	V					I	L		M	: 180
At_CesA3	:	L	I	V	V			F	V		M	: 180
Glyma09g15620	:	L	II	V	V	I		F	I		M	: 180
Glyma15g43040	:	L	II	V	V	I		F	I		M	: 180
Glyma12g36570	:	L	II	V	V	I		F	I		M	: 180
Glyma13g27250	:	L	II	M	V	I		F	I		M	: 180
GRMZM2G018241	:	I	II	V	V	I		F	V		M	: 180
GRMZM2G424832	:	I	II	V	V	I		F	V		M	: 180
Os07g10770	:	I	II	V	V	I		F	V		M	: 180
GRMZM2G150404	:	I	II	V	V	I		F	V		M	: 180
GRMZM2G111642	:	I	II	V	V	I		F	V		M	: 180
Os03g59340	:	I	II	V	V	I		F	V		M	: 180

		*		200							
At_CesA1	:	RTPTIIV	VNSVLL	ASIFSLL	NVRINPF	:	207				
At_CesA10	:	A				:	207				
Glyma04g07220	:	D				:	207				
Glyma06g07320	:	D				:	207				
GRMZM2G027723	:	I		K	D	:	207				
GRMZM2G112336	:	I		K	D	:	207				
Os05g08370	:	I		K	D	:	207				
GRMZM2G039454	:	K		D		:	207				
At_CesA3	:	V		D		:	207				
Glyma09g15620	:	V	I	D		:	207				
Glyma15g43040	:	V	I	D		:	207				
Glyma12g36570	:	V		D		:	207				
Glyma13g27250	:	V	I	D		:	207				
GRMZM2G018241	:	V	AI	D		:	207				
GRMZM2G424832	:	V	AI	D		:	207				
Os07g10770	:	V	AI	D		:	207				
GRMZM2G150404	:	V	AI	D		:	207				
GRMZM2G111642	:	V	AI	M	D	:	207				
Os03g59340	:	V	AI	D		:	207				