



(86) **Date de dépôt PCT/PCT Filing Date:** 2007/03/30
(87) **Date publication PCT/PCT Publication Date:** 2007/11/01
(45) **Date de délivrance/Issue Date:** 2016/09/20
(85) **Entrée phase nationale/National Entry:** 2008/10/20
(86) **N° demande PCT/PCT Application No.:** JP 2007/057109
(87) **N° publication PCT/PCT Publication No.:** 2007/122975
(30) **Priorité/Priority:** 2006/04/20 (JP2006-116498)

(51) **Cl.Int./Int.Cl. C07K 14/475** (2006.01),
A61K 38/00 (2006.01), **A61P 1/00** (2006.01),
A61P 1/16 (2006.01), **A61P 11/00** (2006.01),
A61P 13/12 (2006.01), **A61P 17/00** (2006.01),
A61P 17/16 (2006.01), **A61P 25/00** (2006.01),
A61P 35/00 (2006.01), **A61P 37/06** (2006.01),
A61P 7/00 (2006.01), ...

(72) **Inventeurs/Inventors:**
NAKAMURA, TOSHIKAZU, JP;
MATSUMOTO, KUNIO, JP;
FUKUTA, KAZUHIRO, JP;
ADACHI, KIICHI, JP;
HAYATA, DAICHIKA, JP

(73) **Propriétaires/Owners:**

(54) **Titre : VARIANTE GENETIQUE DE LA PROTEINE PRECURSEUR DU HGF ET PROTEINE ACTIVE CONNEXE**
(54) **Title: HGF PRECURSOR PROTEIN VARIANT AND ACTIVE PROTEIN THEREOF**

(57) **Abrégé/Abstract:**

An HGF precursor protein variant, in which a peptide structure comprises a sequence including a peptide chain X inserted between an α chain of HGF or a polypeptide where 1 to 20 amino-acid residues from the C-terminus of the α chain are deleted, and a β chain of HGF or a polypeptide where 1 to 20 amino-acid residues from the N-terminus of the β chain are deleted; wherein (i) the peptide chain X has an amino-acid sequence of at least two residues, (ii) the peptide chain X can be cleaved by a protease reaction or a chemical reaction, and (iii) a protein obtained by cleaving at least one site of the peptide chain X has HGF action.



(51) **Cl.Int./Int.Cl. (suite/continued)** *C07K 19/00* (2006.01), *C12P 21/02* (2006.01), *C12P 21/06* (2006.01),
C12N 15/09 (2006.01)

(73) **Propriétaires(suite)/Owners(continued)**:KRINGLE PHARMA INC., JP; OSAKA UNIVERSITY, JP

(74) **Agent**: TORYS LLP

ABSTRACT

An HGF precursor protein variant, in which a peptide structure comprises a sequence including a peptide chain X
5 inserted between an α chain of HGF or a polypeptide where 1 to 20 amino-acid residues from the C-terminus of the α chain are deleted, and a β chain of HGF or a polypeptide where 1 to 20 amino-acid residues from the N-terminus of the β chain are deleted; wherein (i) the peptide chain X has an amino-acid
10 sequence of at least two residues, (ii) the peptide chain X can be cleaved by a protease reaction or a chemical reaction, and (iii) a protein obtained by cleaving at least one site of the peptide chain X has HGF action.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

DESCRIPTION

HGF PRECURSOR PROTEIN VARIANT AND ACTIVE PROTEIN THEREOF

5 Technical Field

The present invention relates to an HGF precursor protein variant that can be activated without serum. More specifically, the present invention relates to a single-stranded HGF precursor protein variant comprising a sequence including a peptide chain
10 X, which has an amino-acid sequence composed of at least two residues that can be cleaved by a protease reaction or a chemical reaction, inserted between the α chain and the β chain of the HGF. The present invention also relates to an active HGF protein variant obtained from the variant by cleaving one site in the
15 amino-acid sequence of the inserted peptide chain X. Further, the present invention relates to a method for producing an active HGF protein variant.

Background Art

20 Hepatocyte growth factor (HGF) was found as a protein having hepatocyte proliferation action, and subsequent studies have shown that the HGF is a physiologically active protein having various kinds of pharmacological activities other than the proliferation action of hepatocytes. The pharmacological
25 activities are described, for example, in Jikken Igaku zoukan 10(3) pp.330-339 (1992).

Based on the diverse activities of HGF, other names of HGF including SF (scatter factor) and TCF (tumor cytotoxic factor) are used. However, in the present invention, these known

proteins having proliferation action of hepatocytes are collectively referred to as HGF. Because of its pharmacological activities, expected is the development of HGF as a cirrhosis drug, a renal disease drug, an epithelial cell proliferation accelerant, an anticancer drug, an anti-side-effect drug for cancer therapy, a lung disorder drug, a gastroduodenal damage drug, a cranial nerve disorder drug, an immunosuppressive anti-side-effect drug, a collagen-degrading promoter, a cartilage disorder drug, an arterial disease drug, a pulmonary fibrosis drug, a liver disorder drug, a blood coagulation disorder drug, a hypoproteinemia drug, a wound healing drug, a neuropathy drug, a hematopoietic stem cell augments and hair growth promotion drug, etc. (see, for example, JP-A No. 4-18028, JP-A No. 4-49246, JP-A No. 7-179356, JP-A No. 6-25010, JP-A No. 6-340546, JP-A No. 6-172207, JP-A No. 7-89869, JP-A No. 6-40934, JP-A No. 6-503949, JP-A No. 6-40935, JP-A No. 6-56692, JP-A No. 7-41429, JP3395181 and JP-A No. 5-213721).

HGF is secreted from organs such as the liver, kidneys, lungs, brain, bone marrow, spleen, and placenta, or from blood cells such as platelets and leukocytes. However, as the in vivo content of HGF is infinitesimal, in order to use HGF as a medicinal preparation, a large amount of HGF should be produced by a genetic engineering technique using cells. It is conventionally known that HGF can be produced using animal cells such as Chinese hamster ovary (CHO) cells (see, for example, JP-A No. 11-4696 and JP-A No. 10-191991).

In cultivating animal cells, fetal bovine serum has conventionally been added. However, recently, serum-free

cultivation has been advanced. Therefore, to produce a protein for use as a medicinal preparation with the use of animal cells such as CHO cells, cultivation is generally performed under a serum-free condition. This is because by not using fetal bovine serum, production costs can be reduced and the risk of contamination of viruses and abnormal prions derived from fetal bovine serum can be avoided. Even when HGF is produced using animal cells such as CHO cells, serum-free cultivation can be employed. In this case, however, there is a problem that HGF can be produced only as an inactive HGF precursor protein.

In the biosynthesis of HGF, single-stranded HGF precursor protein is first synthesized and secreted from cells. This HGF precursor protein is an inactive precursor. It is not until the HGF precursor protein is cleaved by the action of a protease called HGF activator (HGFA) and converted into a double stranded structure that the HGF becomes active. This activated HGF is a heterodimer comprising an α chain and a β chain. HGFA itself is also originally biosynthesized as an inactive single-stranded precursor (hereinafter also referred to as pro-HGFA), and is normally circulating in the form of pro-HGFA in the plasma. In the event of tissue injury, working with blood-clotting system and the like, the pro-HGFA is cleaved by the action of thrombin and becomes an active double-stranded HGFA to activate the HGF precursor protein. As serum is in a state where blood-clotting system has already worked, HGFA exists in its active form in serum. Therefore, when CHO cells into which DNA encoding HGF is introduced are cultivated in the presence of serum, HGF produced in culture medium is in its active form by the action of active HGFA in serum. Meanwhile,

when the CHO cells are cultivated under a serum-free condition, because of the absence of HGFA, HGF is produced only as an inactive HGF precursor protein. Although it might be a possible option that HGFA, instead of serum, is added to the cultivation system of CHO cells, there is difficulty in obtaining an active HGFA in the absence of serum because of a cascade that, as described above, HGFA is also secreted as an inactive single-stranded pro-HGFA and then converted into an active HGFA in conjunction with blood-clotting system. Therefore, with conventional technique, an active HGF cannot be efficiently produced unless serum is added.

Consequently, development of a method to activate HGF precursor protein without adding serum has been desired. If such a method becomes available, an active HGF can be safely produced even when CHO cells are cultivated under a serum-free condition, and the risk of contamination of viruses and abnormal prions can be avoided. In addition, serum-free host systems in which yeast and an individual insect, etc. are used as hosts for recombinant production of HGF become available, and the method can be applied to a production system in which HGF expression at a higher level than in CHO cells can be expected.

However, such a method to activate HGF precursor protein without adding serum was previously unknown.

25 Disclosure of the invention

Problems to be solved by the invention

An object of the present invention is to provide an HGF precursor protein variant that can be converted without serum into an active HGF protein variant (an active heterodimer), an

active HGF protein thereof and a preparation method thereof.

Means for solving the problems

To achieve the above-mentioned object, the present
5 inventors have intensively carried out investigations on the
activation of HGF precursor protein, and finally found that an
HGF precursor protein variant comprising a peptide chain X,
which has an amino-acid sequence of at least two residues that
can be cleavage by a protease reaction or a chemical reaction,
10 inserted between the α chain and the β chain thereof can be
converted without serum into an active double-stranded HGF
protein variant. Cleaving the sequence of the peptide chain
X inserted between the α chain and the β chain using a protease
or a chemical treatment agent capable of cleaving the cleavable
15 sequence converts the HGF precursor protein variant into an
active HGF protein variant, which is a disulfide (S-S)-bound
heterodimer, showing the bioaction of HGF. Based on the above
findings, the present inventors have carried out further
investigations and completed the present invention.

20 That is, the present invention provides an HGF precursor
protein variant that can be converted without serum into an
active heterodimer, an active HGF protein thereof and a
preparation method thereof. Further, the present invention
provides a medicinal preparation comprising an active HGF
25 protein variant as an active ingredient.

Namely, the present invention relates to:

(1) an HGF precursor protein variant, in which
a peptide structure comprises a sequence including a peptide

chain X inserted between an α chain of HGF or a polypeptide where 1 to 20 amino-acid residues from the C-terminus of the α chain are deleted, and a β chain of HGF or a polypeptide where 1 to 20 amino-acid residues from the N-terminus of the β chain are deleted; wherein (i) the peptide chain X has an amino-acid sequence of at least two residues, (ii) the peptide chain X can be cleaved by a protease reaction or a chemical reaction, and (iii) a protein obtained by cleaving at least one site of the peptide chain X has HGF action;

10 (2) the HGF precursor protein variant according to the above-mentioned (1), wherein the peptide chain X has a protease recognition sequence;

(3) the HGF precursor protein variant according to the above-mentioned (2), wherein the protease recognition sequence is at least one protease recognition sequence selected from the group consisting of Genenase I recognition sequence, Enterokinase recognition sequence, blood coagulation factor Xa recognition sequence, thrombin recognition sequence, TEV protease recognition sequence, Rhinovirus 3C protease recognition sequence and Furin recognition sequence;

(4) the HGF precursor protein variant according to the above-mentioned (2), wherein the protease recognition sequence is His-Tyr or Tyr-His;

(5) the HGF precursor protein variant according to any of the above-mentioned (1) to (4), wherein the HGF is of human, canine or feline origin;

(6) the HGF precursor protein variant according to any of the above-mentioned (1) to (4), wherein the HGF is of human origin;

(7) the HGF precursor protein variant according to the above-mentioned (6), wherein the HGF is

(a) a protein comprising an amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2;

5 (b) a protein comprising an amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2 wherein one or several amino-acid residues are deleted, substituted or added, and having substantially the same action as that of HGF; or

(c) a protein comprising an amino-acid sequence having a
10 80% homology or more with the amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2, and having substantially the same action as that of HGF;

(8) the HGF precursor protein variant according to the above-mentioned (6), wherein the α -chain is

15 (a) an amino-acid sequence from the 32nd to the 494th of the sequence represented by SEQ ID NO: 1, and the β -chain is

(b) an amino-acid sequence from the 495th to the 728th of the sequence represented by SEQ ID NO: 1;

20 or alternatively the α -chain is

(c) an amino-acid sequence from the 32nd to the 489th of the sequence represented by SEQ ID NO: 2, and the β -chain is

(d) an amino-acid sequence from the 490th to the 723th of
25 the sequence represented by SEQ ID NO: 2;

(9) an active HGF protein variant obtained by cleaving at least one site of the peptide chain X in the HGF precursor protein variant according to any of the above-mentioned (1) to (8);

(10) the active HGF protein variant according to the

above-mentioned (9), wherein the cleavage is caused by a protease treatment or a chemical treatment;

(11) the active HGF protein variant according to the above-mentioned (10), wherein the protease is at least one
5 protease selected from the group consisting of Genenase I, Enterokinase, blood coagulation factor Xa, thrombin, TEV protease, Rhinovirus 3C protease and Furin;

(12) the active HGF protein variant according to the above-mentioned (10), wherein the cleavage occurs at the C
10 terminus of His-Tyr or Tyr-His;

(13) the active HGF protein variant according to the above-mentioned (10) or (12), wherein the cleavage is caused by Genenase I treatment;

(14) the active HGF protein variant according to the above-mentioned (9), wherein the cleavage is caused by a
15 chemical cleavage method;

(15) a method for producing the active HGF protein variant, comprising producing the HGF precursor protein variant according to any of the above-mentioned (1) to (8), and
20 simultaneously or subsequently converting the HGF precursor protein variant into an active HGF protein variant by cleaving at least one site of the peptide chain X;

(16) the method for producing the active HGF protein variant according to the above-mentioned (15), wherein the cleavage is
25 caused by a protease treatment or a chemical treatment;

(17) the method for producing the active HGF protein variant according to the above-mentioned (16), wherein the protease is at least one protease selected from the group consisting of Genenase I, Enterokinase, blood coagulation factor Xa, thrombin,

TEV protease, Rhinovirus 3C protease and Furin;

(18) the method for producing the active HGF protein variant according to the above-mentioned (15), comprising producing a single-stranded HGF precursor protein variant having a sequence
5 into which a peptide chain X having a sequence of His-Tyr or Tyr-His and 2 to 20 amino acid residues, is inserted, and simultaneously or subsequently treating the HGF precursor protein variant with Genenase I;

(19) the method for producing the active HGF protein variant
10 according to any of the above-mentioned (16) to (18), wherein a DNA coding for the HGF precursor protein variant according to any of the above-mentioned (1) to (8) and a DNA coding for a protease to cleave the peptide chain X are simultaneously introduced into a host, the HGF precursor protein variant and
15 the protease to cleave the peptide chain X are simultaneously expressed, and the peptide chain X is cleaved by the protease; and

(20) a medicament comprising the active HGF protein variant according to any of the above-mentioned (9) to (14) as an active
20 ingredient.

Effect of the invention

The HGF precursor protein variant according to the present invention can be converted into an active heterodimer bound by
25 a disulfide (S-S) bond (an active HGF protein variant) under a serum-free condition, and therefore, the risk of contamination of abnormal prions derived from fetal bovine serum can be avoided. Also, according to the invention, an active HGF protein variant can be produced in a serum-free expression system, resulting

in low-cost production of an active HGF protein variant, which is an economical advantage. The active HGF protein variant produced according to the invention can be used as an alternative medicament of HGF because the variant has substantially the same
5 action as that of HGF.

Brief description of the drawings

Fig.1 shows the results of western blot analysis using an anti-HGF polyclonal antibody performed after subjecting the
10 samples of an HGF precursor protein variant treated with or without Genenase I to SDS-PAGE under reducing conditions. In this figure, Pro-HGF represents an HGF-NG precursor protein or an HGF precursor protein variant, α represents the α chain of HGF-NG or the chain of an active HGF protein variant
15 corresponding to the α chain, and β represents the β chain of HGF-NG or the chain of an active HGF protein variant corresponding to the β chain.

Fig.2 shows the results of western blot analysis using an anti-HGF polyclonal antibody performed after subjecting the
20 samples of an HGF-NG precursor protein and an HGF precursor protein variant treated with (+) or without (-) Genenase I to SDS-PAGE under nonreducing conditions.

Fig. 3 shows the evaluation results of the scattering action of the sample on the MDCK cells. The samples were obtained by
25 subjecting HGF-G1 precursor protein, HGF-G2 precursor protein and HGF-G3 precursor protein to Genenase I treatment (+).

Best mode for carrying out the invention

The present invention will, hereinafter, be described in

detail. The peptide chain X inserted between the α chain and the β chain of HGF is not particularly limited as long as the peptide chain has an amino-acid sequence composed of at least two residues, the amino-acid sequence being capable of
5 undergoing peptide cleavage by a protease reaction or a chemical reaction. The number of amino-acid residues which compose peptide chain X is preferably about 20 or less, more preferably about 2 to 10, and most preferably about 2 to 6.

Preferable examples of the amino-acid sequence capable of
10 undergoing peptide cleavage by a protease reaction include a protease recognition sequence (a sequence capable of undergoing peptide cleavage by protease), and the like. The protease recognition sequence is not particularly limited as long as it is recognized by a protease having high substrate specificity.
15 However, it is further preferred that the protease recognition sequence is not included in the amino-acid sequence of HGF. Preferable examples of the protease recognition sequence include His-Tyr or Tyr-His, which is a recognition sequence for Genenase I (Carter, P. et al., Proteins, 6, 240-248(1989));
20 Asp-Asp-Asp-Lys (SEQ ID NO: 3), which is a recognition sequence for Enterokinase (Kunitz, M. ,J. Gen. Physiol. 22, 429-446 (1939), LaVallie, E. R. et al. Journal of Biological Chemistry, 268, 23311-23317 (1993), Voza, L. A. et al. Biotechnology (NY). 14, 77-81 (1996)); Ile-Glu-Gly-Arg (SEQ ID NO: 4) or
25 Ile-Asp-Gly-Arg (SEQ ID NO: 5), which is a recognition sequence for blood coagulation factor Xa; Leu-Val-Pro-Arg-Gly-Ser (SEQ ID NO: 6), which is a recognition sequence for thrombin; Glu-Xaa-Xaa-Tyr-Phe-Gln-Ser (Xaa represents any amino-acid residue; SEQ ID NO: 7) or Glu-Xaa-Xaa-Tyr-Phe-Gln-Gly (Xaa

represents any amino-acid residue; SEQ ID NO: 8), which is a recognition sequence for TEV (tobacco etch virus) protease (Dougherty WG et al., Microbiological Reviews, 57, 781-822 (1992)); Leu-Glu-Val-Leu-Phe-Gln-Gly-Pro (SEQ ID NO: 9), which is a recognition sequence for Rhinovirus 3C protease (Walker PA, et al., Biotechnology (NY), 12(6), 601-605 (1994)); Arg-Xaa-Lys-Arg (Xaa represents any amino-acid residue; SEQ ID NO: 10) or Arg-Xaa-Arg-Arg (Xaa represents any amino-acid residue; SEQ ID NO: 11), which is a recognition sequence for Furin (Hosaka M, et al., Journal of Biological Chemistry, 266, 12127-12130 (1991)), and the like. His-Tyr or Tyr-His, which is recognition sequence for Genenase I, is particularly preferable because of its shortness. The recognition by Genenase I can be improved by extending the N-terminus of His-Tyr or Tyr-His and thereby converting the His-Tyr or Tyr-His into Pro-Gly-Ala-Ala-His-Tyr (SEQ ID NO: 12) or Pro-Gly-Ala-Ala-Tyr-His (SEQ ID NO: 13), etc. The any amino-acid residue Xaa may be selected from 20 kinds of natural amino acids and unnatural amino acids. The unnatural amino acid may be any compound as long as it has an amino group and a carboxyl group, and for example, it may be a γ -amino butyric acid, etc.

Examples of the amino-acid sequence being capable of undergoing peptide cleavage by a chemical reaction include Asn-Gly cleaved by hydroxylamine reaction, and Asp-Pro cleaved by acetic acid containing guanidine hydrochloride.

The HGF precursor protein variant relating to the present invention can be designed based on the amino-acid sequences of the HGF of human origin or mammal (for example, feline, canine, rat, mouse, bovine, chimpanzee, equine, porcine, sheep, or the

like) origin. Examples of the HGF include human-derived HGF (for example, Accession No. NP_001010932, P14210, BAA14348, AAC71655, and the like), mouse-derived HGF (for example, Accession No. AAB31855, NP_034557, BAA01065, BAA01064, and the like), rat-derived HGF (for example, Accession No. NP_58713, and the like), bovine-derived HGF (for example, Accession No. NP_001026921, XP874086, BAD02475, and the like), feline-derived HGF (for example, Accession No. NP_001009830, BAC10545, BAB21499, and the like), canine-derived HGF (for example, Accession No. NP_001002964, BAC57560, and the like), and chimpanzee-derived HGF (for example, Accession No. XP519174, and the like) registered in, for example, the NCBI database or the like, but are not limited thereto. In addition, as long as the HGF has substantially the same action as HGF, one or several (for example about 2 to 30, preferably about 2 to 20, more preferably about 2 to 10, further more preferably 2 to 5; hereinafter the same) amino-acid residues may be substituted, deleted or added in the amino-acid sequence of the HGF. Similarly, a sugar chain may be substituted, deleted or added. Examples of the HGF include HGF (Accession No. NP_001010932), in which 5 amino-acid residues of the HGF registered as Accession No. P14210 are deleted, and the like. Also, when the sequence existing in the inserted peptide chain X and being capable of undergoing peptide cleavage by a protease reaction or a chemical reaction is included in the amino-acid sequence of HGF, as long as the HGF has substantially the same action as HGF, one or several amino-acid residues of the peptide cleavage sequence may be substituted, deleted or added by a known method. Examples of the known method include site-directed mutagenesis described

below. With regard to the amino-acid sequence here, "one or several amino-acid residues of the peptide-cleavage sequence may be substituted, deleted, or added" means that a certain number (one or several, the definition of several is same as above) of the amino-acid residues is substituted, deleted or added, with the proviso that the number can be given by a known method such as genetic engineering technique site-directed mutagenesis or naturally. Examples of the HGF where a sugar chain is substituted, deleted or added include an HGF obtained by depriving a natural HGF of a sugar chain by the treatment of an enzyme etc., an HGF having a mutated amino-acid sequence at a glycosylation site to prevent glycosylation, an HGF having a mutated amino-acid sequence to induce glycosylation at a different site from the natural glycosylation site, and the like. Specific examples include an HGF obtained by respectively substituting the 289th Asn with Gln, 397th Asn with Gln, 471th Thr with Gly, 561th Asn with Gln, and 648th Asn with Gln in the glycosylation site of HGF (Accession No.NP_001010932 (SEQ ID NO: 2)) to prevent glycosylation (SEQ ID NO: 14; Fukuta K et al., Biochemical Journal, 388, 555-562 (2005)), and the like. In addition, a protein having at least about 80% homology or more with the amino-acid sequence of HGF, preferably about 90% homology or more, more preferably about 95% homology or more, and having substantially the same action as HGF, is also included in the above-mentioned HGF.

With regard to the above amino-acid sequence, "homology" means, in comparison of primary structures of proteins, the extent of correspondence of amino acids composing each sequence (hereinafter the same).

Specifically, the above HGF may be an amino-acid sequence represented by SEQ ID NO: 1 (Accession No. P14210) or SEQ ID NO: 2 (Accession No. NP_001010932), or the like. Further, examples of the HGF amino-acid sequence of the present invention include an amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2 where one or several amino acids are inserted or deleted, where one or more amino acid residues are substituted with another (or more) amino acid residue(s), or where one or more amino acid residues are modified, with the proviso that the amino-acid sequence constitutes a protein which has substantially the same action as HGF. The HGF represented by SEQ ID NO: 2 lacks 5 amino-acid residues from the 162th phenylalanine residue to the 166th serine residue of the amino-acid sequence represented by SEQ ID NO: 1. Therefore, the HGF represented by SEQ ID NO: 2 may be called 5-amino-acid-deleted-type human HGF. The inserted or substituted amino-acid residue may be unnatural amino acids other than 20 kinds of natural amino acids encoded by DNA. The unnatural amino acid may be any compound as long as it has an amino group and a carboxyl group, and for example, it may be a γ -amino butyric acid, etc. Also, examples of the HGF amino-acid sequence of the present invention include an amino-acid sequence having at least about 80% or more, preferably about 90% or more, more preferably about 95% or more homology with the amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2, and having substantially the same action as HGF. The amino-acid sequence having at least about 80% homology or more with the amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2 may be the HGF amino-acid sequence registered

as Accession No. NP_001010934, BAA14348, AAC71655, AAB31855,
NP_034557, BAA01065, BAA01064, NP_58713, NP_001026921,
XP874086, BAD02475, NP_001009830, BAC10545, BAB21499,
NP_001002964, BAC57560, XP519174 or the like in the NCBI
5 database or the like, but not limited thereto.

The HGF precursor protein variant according to the present
invention can be produced by a method comprising the following
steps 1 to 4, but steps are not limited thereto as long as the
HGF precursor protein variant can be produced.

10 Step 1:

In Step 1, DNA encoding HGF precursor protein variant is
prepared. The step includes a step to prepare a recombinant
expression vector including DNA encoding HGF precursor protein
variant by inserting the peptide chain X between the α chain
15 and the β chain of HGF.

The insertion of the peptide chain X between the α chain
and the β chain of HGF may be only inserting the peptide chain
X into the boundary between the α chain and the β chain of a
natural HGF. In this case, the total number of amino-acid
20 residues of the HGF is increased by the number of the amino-acid
residues included in the inserted sequence. Also, the
insertion may be inserting the peptide chain X into the region
at either side of the boundary between the α chain and the β
chain of HGF where about 1 to 20 amino-acid residues from the
25 C-terminus of the α chain and/or about 1 to 20 amino-acid
residues from the N-terminus of the β chain are deleted.

The α chain and the β chain of HGF may be, for example,
the α chain consisting of from the 32nd to the 494th of the
amino-acid sequence represented by SEQ ID NO: 1 and the β chain

consisting of from the 495th to the 728th of the amino-acid
sequence represented by SEQ ID NO: 1; or the α chain consisting
of from the 32nd to the 489th of the amino-acid sequence
represented by SEQ ID NO: 2 and the β chain consisting of from
5 the 490th to the 723rd of the amino-acid sequence represented
by SEQ ID NO: 2. The amino-acid residues constituting the
boundary between the α chain and the β chain are, for example,
in the case of the HGF represented by SEQ ID NO: 1, the arginine
at the 494th and the valine at the 495th. The amino-acid
10 residues constituting the boundary between the α chain and the
 β chain are, for example, in the case of the
5-amino-acid-deleted-type human HGF represented by SEQ ID NO:
2, the arginine at the 489th and the valine at the 490th.

The insertion of the peptide chain X between the α chain
15 and the β chain of HGF can be performed by site-directed
mutagenesis of protein or the like. The mutagenesis method may
be, for example, inserting a base sequence encoding the aimed
peptide chain X into the base-sequence region at the boundary
between the DNA encoding the α chain and the DNA encoding the
20 β chain of HGF, or introducing a mutation to induce substitution
for an aimed sequence. The method to mutagenize a base sequence
may be, for example, synthesizing a mutant primer corresponding
to the site to mutate, and then employing a known method, for
example, the Kunkel method (Kunkel, T. A. Proc. Natl. Acad. Sci.
25 U.S.A. 82, 488-492 (1985)), etc. By using a
commercially-available mutagenesis kit or the like, mutation
can be easily introduced. Examples of the mutagenesis kit
include, for example, GeneMorph Random Mutagenesis Kit (made
by Stratagene) containing Mutazyme DNA polymerase, GeneTailor

(Trademark) Site-Directed Mutagenesis System (made by Invitrogen), Mutan (Trademark) -Super Express Km (made by Takara Bio), QuikChange (Trademark) XL Site-Directed Mutagenesis Kit (made by Toyobo), GeneEditor in vitro Site-Directed Mutagenesis System (made by Promega), and the like, but are not limited thereto. The DNA mutated by the above-mentioned Kunkel method or with the mutagenesis kit, etc. (DNA having a base sequence encoding HGF precursor protein variant; hereinafter referred to as DNA encoding HGF precursor protein variant) can be excised by a restriction enzyme from a plasmid or phage that has produced the DNA through mutation and/or amplification. The excised DNA may be purified by a known method, or directly inserted into an appropriate expression vector described below. The purification may be performed using a commercially-available kit, for example, QIAquick Gel extraction Kit (made by Qiagene), S.N.A.P. UV-Free Gel Purification Kit (made by Invitrogen), or the like, but the kit is not limited thereto. The DNA encoding HGF precursor protein variant can also be produced by a known chemical synthetic method. Examples of the chemical synthetic method include chemical synthesis using a DNA synthesizer such as DNA synthesizer adopting the phosphoramidite method.

The expression vector having DNA encoding HGF precursor protein variant can be produced by ligating a DNA encoding HGF precursor protein variant to downstream of a promoter in an appropriate vector for HGF expression using a restriction enzyme and DNA ligase. The recombinant expression vector contains a promoter, a ribosome binding site, an initiation codon, a termination codon, a terminator, and the like, as needed. It is preferred that the recombinant expression vector is

constructed to contain (1) a promoter, (2) a ribosome binding site, (3) an initiation codon, (4) a DNA encoding HGF precursor protein variant of the present invention, (5) a termination codon and (6) a terminator in order toward downstream of transcription. The expression vector used in the present invention may be, when the host is *Escherichia coli*, a plasmid of pBR322, pUC18, pUC19 (made by Toyobo), or the like; when the host is *Bacillus subtilis*, a plasmid of pUB110 (made by Sigma), or the like; and when the host is yeast, a plasmid of pYES2 (made by Invitrogen), pRB15 (ATCC37062), or the like. A vector for an animal cell may be pCAGGS and pCXN2 (Niwa, H., Yamamura, K. and Miyazaki, J., *Gene*, Vol. 108, pp. 193-200 (1991) and JP-A No.3-168087) or pcDL-SR α (Takebe, Y. et al., *Mol. Cell. Biol.*, Vol. 8, pp. 466-472 (1988)) or the like. Besides, the expression vector may be bacteriophage λ gt10 or λ gt11 (made by Stratagene), virus SV40 (made by BRL), BPV (ATCC VR-703), or a vector derived from retroviral gene, but not limited thereto. Any vector may be used as long as it can be replicated and/or amplified in a host.

The promoter and the terminator are not also particularly limited as long as they are corresponding to the host used for expression of DNA encoding the aimed HGF precursor protein variant. Examples of the promoter include, when the host is *Escherichia coli*, trp promoter, lac promoter, recA promoter, λ PL promoter, lpp promoter, and the like; and when the host is yeast, PH05 promoter, PGK promoter, GAP promoter, ADH promoter, and the like. When the host is an animal cell, in addition to SR α promoter; CAG promoter; a promoter obtained from viral genome of Rous sarcoma virus (RSV), MPSV, polyoma virus, fowlpox

virus, adenovirus, bovine papilloma virus, fowl sarcoma virus, cytomegalovirus (CMV), hepatitis B virus, simian virus 40 (SV40), and vaccinia virus; metallo-thioneine promoter; heat shock promoter; or the like is included. In the case of using a higher mammal host, an enhancer is preferably introduced into a vector. Introduction of an enhancer may increase transcription. The enhancer may be SV40 enhancer, initial promoter/enhancer of cytomegalovirus, polyoma enhancer, adenovirus enhancer, or the like. The terminator may be, when the host is *Escherichia coli*, trp terminator, lpp terminator, or the like; when the host is *Bacillus subtilis*, amyF terminator or the like; when the host is yeast, CYC1 terminator or the like; or when the host is an animal cell, SV40 terminator, HSV1TK terminator, or the like. These promoters and terminators are appropriately combined depending on the host used.

Step 2:

In Step 2, HGF precursor protein variant is synthesized. The step includes a step to produce a transformant by introducing the recombinant expression vector prepared in the Step 1 into a host, and then synthesize HGF precursor protein variant in the transformant.

The recombinant expression vector having DNA encoding HGF precursor protein variant, the vector which has been prepared in the Step 1, is introduced into a host, by a competent cell method (J. Mol. Biol., vol. 53, p. 154, (1970)), protoplast method (Proc. Natl. Acad. Sci. USA, vol. 75, p. 1929 (1978)), calcium phosphate method (Science, vol. 221, p. 551 (1983)), DEAE dextran method (Science, vol. 215, p. 166 (1982)), electric pulse method (Proc. Natl. Acad. Sci. USA, vol. 81, p. 7161

(1984)), in vitro packaging method (Proc. Nat. Acad. Sci. USA, vol. 72, p. 581 (1975)), viral vector method (Cell, vol. 37, p. 1053 (1984)), micro injection method (Exp. Cell. Res., vol. 153, p. 347 (1984)), or the like, to produce a transformant.

5 The cell which can be used as a host is not particularly limited, and eukaryotic cells such as an animal, a plant, an insect, a eukaryotic microorganism, or the like; or prokaryotic cells such as a prokaryotic microorganism or the like, may be used. These cells may form an individual, and a host may be
10 an animal individual, a plant individual, or an insect individual. The eukaryotic cell may be an adherent cell or a floating cell, for example, may be a eukaryotic cell producing HGF precursor protein variant and accumulating it in the cell, or a eukaryotic cell producing HGF precursor protein variant
15 and secreting it out of the cell. The animal cell may be, for example, a CHO cell (Chinese hamster ovary cell), a COS cell, a BHK cell, a mouse C127 cell, a Hela cell, or the like. The plant cell may be, for example, cells of rice, tobacco, *Arabidopsis thaliana*, or the like, and the insect cell may be,
20 for example, Sf9 cell, Sf21 cell, or the like. The insect individual may be, for example, silk worm (*Bombyx mori*). The eukaryotic microorganism may be, yeast such as *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Candida boidinii*, or *Pichia pastoris*; or a filamentous fungus such as *Aspergillus*,
25 *Trichoderma* or *Mucor*. The prokaryotic microorganism may be, *Escherichia coli*, *Bacillus subtilis*, or the like.

The resultant transformant is preferably cultured in a medium appropriate for the host for the purpose of producing an aimed HGF precursor protein variant. The medium contains

a carbon source, a nitrogen source, inorganic substances, vitamins, serum, agents, and the like, necessary for growth of the transformant. When the host of the transformant is Escherichia coli, the medium may be LB medium (Nissui
5 Pharmaceutical Co., Ltd.), M9 medium (J. Exp. Mol. Genet., Cold Spring Laboratory, New York, p. 431(1972)), or the like. When the host is yeast, the medium may be YEPD medium (Genetic Engineering, vol. 1, Plenum Press, New York, p. 117(1979)) or the like. When the host is an animal cell, the medium may be
10 MEM medium (Minimum Essential Medium), DMEM medium (Dulbecco's Modified Eagle's Medium), RPMI 1640 medium (Nissui Pharmaceutical Co., Ltd.) containing 20% or less by volume of fetal bovine serum, or the like. A transformant is usually cultured at a temperature of 20 to 45°C and a pH of 5 to 8, with
15 ventilation or stirring as required. When the host is an animal adherent cell or the like, carriers such as glass beads, collagen beads or acetyl cellulose hollow fiber can be preferably used. A transformant can be cultured even with any other medium composition or under other culturing conditions as long as the
20 transformant can grow, and therefore, the composition and culturing conditions are not limited to the above-mentioned examples.

The obtained transformant can express DNA encoding HGF precursor protein variant and synthesize HGF precursor protein
25 variant.

Further, the HGF precursor protein variant of the present invention can be obtained also by a cell-free protein synthesis system. The cell-free protein synthesis system includes a method of synthesizing protein not using a live cell but using

DNA or mRNA encoding the aimed protein as a template, by using cell extract prepared from *Escherichia coli*, rabbit reticulocyte, wheat germ, or the like; or by using protein synthesis factors contained in the cell extract solution.

5 Since a cell extract solution contains molecules necessary for protein synthesis such as ribosome, tRNA, and translation factors, adding an energy source such as ATP and GTP; and substrate amino acids thereto synthesizes a protein. Instead of the cell extract solution, a mixture of protein synthesis

10 factors contained in the cell extract solution may be used.

Step 3:

In Step 3, HGF precursor protein variant (hereinafter also referred to just as precursor) is converted into an active HGF protein variant.

15 Since the HGF precursor protein variant synthesized in the Step 2 is inactive, it is preferable to convert the precursor into an active HGF protein variant by cleaving the precursor to convert it into a double-stranded chain by cleaving at least one site of the peptide chain X inserted between the α chain

20 and the β chain.

The cleaving method is not particularly limited as long as a specific amino-acid sequence in the peptide chain X on the precursor is cleaved. The method includes, for example, a protease treatment or a chemical treatment.

25 The protease capable of cleaving a specific amino-acid sequence is not particularly limited, and preferably used is, for example, Genenase I, Enterokinase, blood coagulation factor Xa, thrombin, TEV protease, Rhinovirus 3C protease, Furin, or the like, which has a high substrate specificity to recognize

a specific amino-acid sequence. For example, when a precursor has an inserted Genenase I recognition sequence, the single-stranded precursor can be activated by the action of Genenase I. In this case, the cleavage occurs between His-Tyr or Tyr-His, which is a Genenase I recognition sequence. When a precursor has an inserted Enterokinase recognition sequence, the single-stranded precursor can be activated by the action of Enterokinase. In this case, the cleavage occurs at the C-terminus of the lysine residue of Asp-Asp-Asp-Lys (SEQ ID NO: 3), which is an Enterokinase recognition sequence. Similarly, when a precursor has an inserted recognition sequence for blood coagulation factor Xa, the single-stranded precursor can be activated by the action of blood coagulation factor Xa. In this case, the cleavage occurs between Gly-Arg in the Ile-Glu-Gly-Arg (SEQ ID NO: 4) or Ile-Asp-Gly-Arg (SEQ ID NO: 5), which is a recognition sequence for blood coagulation factor Xa. When a precursor has an inserted thrombin recognition sequence, the single-stranded precursor can be activated by thrombin. In this case, the cleavage occurs between Arg-Gly of the Leu-Val-Pro-Arg-Gly-Ser (SEQ ID NO: 6), which is a thrombin recognition sequence. When a precursor has an inserted TEV protease recognition sequence, the single-stranded precursor can be activated by the action of TEV protease. In this case, the cleavage occurs at the C-terminus of the serine residue of Glu-Xaa-Xaa-Tyr-Phe-Gln-Ser (SEQ ID NO: 7), which is a TEV protease recognition sequence, or at the C-terminus of the glycine residue of Glu-Xaa-Xaa-Tyr-Phe-Gln-Gly (SEQ ID NO: 8), which is also a TEV protease recognition sequence. When a precursor has an inserted Rhinovirus 3C protease recognition

sequence, the single-stranded precursor can be activated by Rhinovirus 3C protease. In this case, the cleavage occurs between Gln-Gly of the Leu-Gln-Val-Leu-Phe-Gln-Gly-Pro (SEQ ID NO: 9), which is a Rhinovirus 3C protease recognition sequence.

5 When a precursor has an inserted Furin recognition sequence, the single-stranded precursor can be activated by Furin. In this case, the cleavage occurs at the C-terminus of the C-terminal arginine residue of Arg-Xaa-Lys-Arg (SEQ ID NO: 10) or Arg-Xaa-Arg-Arg (SEQ ID NO: 11), which is a Furin recognition
10 sequence.

The protease includes a protease isolated from a living body, cells or fungus and then purified, a protease purified as a recombinant protein, or a commercially-available protease. The protease may be added to the above-mentioned medium for a
15 transformant, or reacted with HGF precursor protein variant extracted from a transformant or the like and then purified.

Examples of other methods to react a protease with HGF precursor protein variant include a method where DNA encoding a protease is also implanted into the transformant producing
20 HGF precursor protein variant in order to allow the transformant to express the protease and HGF precursor protein variant at the same time. In this case, it is preferred that the DNA to be implanted is capable of encoding a protease recognizing the protease recognition sequence included in the peptide chain X.
25 Since the transformant produces the protease and HGF precursor protein variant at the same time, the HGF precursor protein variant produced by the transformant can be automatically cleaved at the site of protease recognition sequence in the peptide chain X and activated.

Examples of the method to cleave the peptide chain X site in HGF precursor protein variant by a chemical treatment include a method to react a chemical treatment agent such as hydroxylamine or acetic acid containing guanidine hydrochloride (preferably, for example, about 10% by volume acetic acid containing about 7 M guanidine hydrochloride (approximately pH 2.5)) with HGF precursor protein variant. Protein cleavage can occur when, for example, Asn-Gly binding is cleaved by the reaction of hydroxylamine, or when Asp-Pro binding is cleaved by acetic acid containing guanidine hydrochloride. The above-mentioned chemical treatment agent may be added to the above-mentioned medium for a transformant, as long as the transformant can grow.

The protease treatment or the chemical treatment may be performed in the presence of, for example, a reducing agent (for example, dithiothreitol (DTT), β -mercaptoethanol, or the like) or a denaturation agent (for example, SDS, urea, guanidine hydrochloride, or the like). In this case, it is preferred that a renaturation reaction is performed after the peptide chain X is cleaved. The renaturation reaction may be performed according to a known method, for example, a method described in *Molecular Cloning: A Laboratory Manual*, 3rd Edition. Sambrook, J. and Russell, D. W., eds. (2001) Cold Spring Harbor Press, pp. A4. 39.

Step 4:

In Step 4, an active HGF precursor protein variant is isolated and purified. The active HGF precursor protein variant produced in the culture supernatant of a transformant or in a transformant can be isolated and purified by a known

salting out method, solvent precipitation method, dialysis method, ultrafiltration method, gel electrophoresis method or gel filtration chromatography, ion exchange chromatography, reverse phase chromatography, affinity chromatography, or the like, or by a combination thereof. Particularly, a combination of a salting out method using ammonium sulfate, S-sepharose ion chromatography, heparin sepharose affinity chromatography and phenylsepharose hydrophobic chromatography; and a combination of a salting out method using ammonium sulfate, S-sepharose ion chromatography and anti-HGF antibody sepharose affinity chromatography; and the like, are preferable and effective purification methods.

The active HGF precursor protein variant of the present invention has substantially the same action as HGF, and therefore, like HGF, can be used as a protein medicament, namely as a therapeutic drug or a preventive drug for various diseases of human and other mammals (canine, feline, rat, mouse, rabbit, horse, bovine, sheep, guinea pig, and the like). Examples of the uses thereof include a liver disease drug, a renal disease drug, a wound healing drug, a cutaneous ulcer drug, a hair root cell proliferating drug, an anticancer drug, a lung disease drug, and an anti-side-effect drug for cancer therapy. More specifically, the active HGF precursor protein variant is useful for preventing and/or treating diseases to which HGF is applicable, including liver diseases (for example, hepatitis, cirrhosis, liver failure, liver regeneration after surgery, and the like), renal disease (for example, glomerular nephritis, kidney failure, nephrogenic anemia, diabetic nephropathy, kidney disorder after drug administration, and the like),

cutaneous diseases (for example, white spot lesion, burn injury, bed sore, cutaneous ulcer, calvities, and the like), blood diseases (for example, thrombocytopenia, bone marrow transplantation, and the like), eye diseases (for example, 5 corneal ulcer, and the like), lung diseases (for example, pneumonia, emphysema, pulmonary tuberculosis, chronic obstructive lung disease, pneumoconiosis, fibroid lung, and the like), gastroduodenal diseases (for example, gastric inflammation, gastric ulcer, duodenal ulcer, and the like), 10 cancers and related diseases (for example, various cancers; side effects of cancer therapy such as liver toxicity, kidney toxicity, nausea, vomit, thrombocytopenia, hair loss; and the like), bone diseases (for example, osteoporosis, osteodysplasty, osteoarthritis, and the like), and central 15 nervous system diseases (for example, abnormal neural differentiation, and the like).

The medicinal preparation containing the active HGF precursor protein variant of the present invention is used in the form of a general medicinal preparation. As the form of 20 the medicinal preparation, various dosage forms (for example, liquid, solid, capsule, and the like) may be adopted. In general, the active HGF precursor protein variant as an active ingredient and binding substances are solely used or used in combination with a customary carrier to give an injection, inhalant, 25 suppository or oral agent, and an injection is preferable. The injection may be an aqueous or oily injection. The injection can be prepared by a known method. For example, an aqueous injection can be prepared as follows: to an aqueous solvent such as water for injection and purified water, optionally added is

a pharmaceutically acceptable excipient, for example a tonicity agent (such as sodium chloride, potassium chloride, glycerin, mannitol, sorbitol, boric acid, borax, glucose, propylene glycol), a buffering agent (such as phosphate buffer solution, acetate buffer solution, borate buffer solution, carbonate buffer solution, citrate buffer solution, Tris-buffer solution, glutamic acid buffer solution, epsilon-aminocaproic acid buffer solution), a preservative (such as methyl parahydroxybenzoate, ethyl parahydroxybenzoate, propyl parahydroxybenzoate, butyl parahydroxybenzoate, chlorobutanol, benzyl alcohol, benzalkonium chloride, sodium dehydroacetate, sodium edetate, boric acid, borax), a thickener (such as hydroxyethyl cellulose, hydroxypropyl cellulose, polyvinyl alcohol, polyethylene glycol), a stabilizer (such as albumin, globulin, gelatine, alanine, glycine, mannitol, glucose, dextran, sorbitol, ethylene glycol, sodium bisulfite, sodium thiosulfate, sodium edetate, sodium citrate, ascorbic acid, dibutyl hydroxytoluene), a pH adjuster (such as hydrochloric acid, sodium hydroxide, phosphoric acid, acetic acid) or the like. Next, after the active HGF precursor protein variant is dissolved in the resulting solution, the solution is sterile-filtered through a filter or the like. Finally, the filtered solution is filled into a sterile container. Additionally, an appropriate solubilizing agent, for example an alcohol (such as ethanol), a polyalcohol (such as propylene glycol and polyethylene glycol), a nonionic surfactant (such as polysorbate 80 and polyoxyethylene (50) hydrogenated castor oil) or the like may be also incorporated. To prepare an oily injection, sesame oil, soy bean oil or the like may be used as

an oily solvent and benzyl benzoate, benzyl alcohol or the like may be incorporated as a solubilizing agent. The prepared injection is usually filled into an appropriate ampule or vial, etc. The injection is prepared in an amount of the active HGF precursor protein variant ranging usually from about 0.0002 to 3% by mass, preferably about 0.001 to 2% by mass. It is preferred that a liquid preparation such as an aqueous injection is frozen for preservation or stored after removing moisture by lyophilization or the like. The lyophilized preparation can be used by adding distilled water for injection or the like as needed and redissolving the preparation.

The oral drug is formulated into for example, a tablet (including sugarcoated tablet, filmcoated tablet, and enteric tablet), granule, fine granule, powder, soft or hard capsule (including enteric capsule), liquid, emulsion, suspension, syrup or the like. These preparations can be prepared by an ordinary method for preparation.

The active HGF precursor protein variant to be used in the present invention together with a biodegradable polymer can be prepared in the form of a sustained-release preparation, for example, a depot preparation. Especially, a depot preparation of the active HGF precursor protein variant can be expected to reduce the dose frequency, prolong the effect and reduce the side effect, etc. The sustained-release preparation can be prepared by known methods. The biodegradable polymer to be used in the sustained-release preparation can be appropriately selected from known biodegradable polymers, for example, a polysaccharide such as starch, dextran or chitosan; a protein such as collagen or gelatin; a polyamino acid such as

polyglutamic acid, polylysine, polyleucine, polyalanine or polymethionine ; a polyester such as polylactic acid, polyglycolic acid, lactic acid-glycolic acid copolymer, polycaprolactone, poly- β -hydroxybutyric acid, polymalic acid, 5 polyanhydride or fumaric acid-polyethylene glycol-vinylpyrrolidone copolymer; a polyalkyl cyanoacrylate such as a polyortho ester or polymethyl- α -cyanoacrylate; or a polycarbonate such as polyethylene carbonate or polypropylene carbonate. Preferred is a polyester and more preferred is 10 polylactic acid or lactic acid-glycolic acid copolymer. When lactic acid-glycolic acid copolymer is used as a biodegradable polymer, the proportion based on the mole percentage (lactic acid/glycolic acid) depends on the duration of sustained release. For example, when the duration of sustained release is from about 15 2 weeks to 3 months, preferably from about 2 weeks to 1 month, the preferable proportion is from about 100/0 to 50/50. In general, the weight-average molecular weight of the polylactic acid or lactic acid-glycolic acid copolymer is preferably from about 5,000 to 20,000. The polylactic acid or lactic 20 acid-glycolic acid copolymer can be prepared by known synthesis methods, for example the method disclosed by JP-A No. 61-28521. The proportion of the active HGF precursor protein variant to the biodegradable polymer is not particularly limited, but a preferable example of the proportion is from about 0.01 to 30% 25 by mass of the active HGF precursor protein variant relative to the biodegradable polymer.

The inhalant can also be prepared according to normal means for preparation. The amount of the active HGF precursor protein variant in a preparation can be appropriately adjusted depending

on dosage form, disease to be treated and the like.

A spray can also be prepared according to normal means for preparation. To prepare a spray, any excipient may be incorporated into the spray as long as the excipient is usually used for an inhaled preparation. For example, in addition to a propellant, the above-mentioned solvent, preservative, stabilizer, tonicity agent or pH adjuster, etc. can be incorporated. Examples of the propellant include a liquefied gas propellant and a compressed gas. Examples of the liquefied gas propellant include a fluorohydrocarbon such as a substitute for chlorofluorocarbons (HCFC22, HCFC-123, HCFC-134a, HCFC142 or the like), liquefied petroleum, dimethylether or the like. Examples of the compressed gas include a soluble gas such as carbon dioxide gas and nitrous oxide gas or an insoluble gas such as nitrogen gas.

The suppository can also be prepared by an ordinary method for preparation using a conventional base (for example, cacao butter, lauric butter, glycerogelatine, Macrogol, Witepsol and the like). In formulation, a stabilizer is preferably added. In addition, a preparation of the present invention may contain other essential excipients, for example, fillers, solubilizers, antioxidants, soothing agents, isotonic agents, or the like.

The preparation of the present invention can be administered via a suitable administration route corresponding to its dosage form. For example, it can be formed into an injection and administered intravenously, intraarterially, subcutaneously, intramuscularly, or the like. The dose thereof is appropriately adjusted depending on disease, symptom, age, body weight or the like of a patient, and for example, it is

from 0.01 mg to 500 mg of HGF for an adult, preferably from 0.05 mg to 100 mg, more preferably from 0.05 mg to 50 mg, and most preferably from 0.05 mg to 20 mg. This dose is preferably administered once or in several divided portions daily.

5 The present invention will hereinafter be described in more detail by the following examples, but the scope of the invention is not limited thereto.

The meanings of abbreviations used in the examples are shown below.

10 HGF: hepatocyte growth factor
LB medium: Luria-Bertani medium
DMEM medium: Dulbecco's Modified Eagle's Medium
Amp: ampicillin
FCS: fetal calf serum
15 Tris: tris (hydroxymethyl) aminomethane
Tween 80: polyoxyethylene (20) sorbitan monooleate
SDS: sodium dodecylsulfate
PAGE: polyacrylamide gel electrophoresis
PVDF: polyvinylidene fluoride
20 A: Adenine
C: Cytosine
G: Guanine
T: Thymine
Ala: Alanine
25 Arg: Arginine
Gly: Glycine
Gln: Glutamine
Lys: Lysine
Thr: Threonine

Leu: Leucine

His: Histamine

Tyr: Tyrosine

Pro: Proline

5 Also, "%" means % by mass unless otherwise stated.

Example 1

To the both ends of the base sequence encoding the HGF represented by SEQ ID NO: 14 (5-amino-acid-deleted-type
10 /glycosylation-deficient human HGF; HGF having a natural α chain- β chain boundary sequence Arg (489th)-Val (490th) and having a mutation at a glycosylation site; hereinafter referred to as HGF-NG) (SEQ ID NO: 15), a base sequence including BamHI recognition sequence (GGATCC) and a base sequence including XbaI
15 recognition sequence (TCTAGA) were added, and this sequence was incorporated between the BamHI site and the XbaI site of pCDNA3.1 (+) vector (made by Invitrogen). The obtained vector is called pCDNA-dHGF-NG.

As variants having an introduced Genenase I recognition
20 sequence (His-Tyr) at the α chain- β chain boundary of HGF-NG, a variant where the two C-terminal residues (Leu-Arg) of the HGF α chain were converted into His-Tyr (referred to as HGF-G1), a variant where the C-terminal residue (Arg) of the HGF α chain was converted into Tyr and the N-terminal residue (Val) of the
25 HGF β chain was converted into His (referred to as HGF-G2), and a variant where the six C-terminal residues (the 484th to 489th of the SEQ ID NO: 14; Lys-Thr-Lys-Gln-Leu-Arg) of the HGF α chain were converted into Pro-Gly-Ala-Ala-His-Tyr (SEQ ID NO: 12) (referred to as HGF-G3), were prepared as follows.

First, vectors for expressing HGF-G1, HGF-G2 and HGF-G3 were prepared. For this purpose, using the above-mentioned pCDNA-dHGF-NG as a template, a base sequence region encoding the HGF α chain- β chain boundary was mutated by the Kunkel method, and the mutated chain was amplified. Specifically, to prepare vectors to express HGF-G1, HGF-G2, and HGF-G3, the primer of the SEQ ID NO: 16 (5'-phosphorylated), the primer of the SEQ ID NO: 17 (5'-phosphorylated), and the primer of the SEQ ID NO: 18 (5'-phosphorylated), were used respectively. The mutated chain was elongated and amplified using pCDNA-dHGF-NG as a template and KOD Plus (made by Toyobo) as DNA polymerase. Then, after the template DNA was digested by DpnI treatment, the remaining mutated chain was used for transformation of Escherichia DH5 α competent cells (made by Nippon Gene) by the calcium chloride method to prepare the desired mutated vector.

Table 1

Primer	Sequence listing
5' -CCAAAACGAAACAACACTATGTTGTAAATGGGATTCCAACACG-3'	SEQ ID NO: 16
5' -CGAAACAATTGTATCACGTAAATGGGATTCCAACACG-3'	SEQ ID NO: 17
5' -GTAATATCTTGTGCCCCAGGGCCGACACTATGTTGTAAATGG-3'	SEQ ID NO: 18

On LB/Amp plates, Amp-resistant colonies were picked, and each mutated vector was extracted from each obtained clone using QIAprep Spin Miniprep Kit (made by Qiagen). By analyzing the base sequence encoding HGF-NG in each mutated vector, the target clone was chosen. Specifically, the base sequence region encoding the HGF represented by the SEQ ID NO: 14 in a mutated

vector was subjected to sequence reaction using Big Dye Terminator v3.1 Cycle Sequence Kit (made by Applied Biosystems), and then analyzed with 3100 Genetic Analyzer (made by Applied Biosystems). A vector of which the HGF-NG α chain- β chain boundary was confirmed to be successfully mutated was chosen and used in later experiments.

The mutated vector where the Leu-Arg at the 488th to 489th of the SEQ ID NO: 14 is substituted with His-Tyr by using the mutated primer of the SEQ ID NO: 16 is referred to as pCDNA-dHGF-NG-G1. The mutated vector where the Arg-Val at the 489th to 490th of the SEQ ID NO: 14 is substituted with Tyr-His by using the mutated primer of the SEQ ID NO: 17 is referred to as pCDNA-dHGF-NG-G2. The mutated vector where the Lys-Thr-Lys-Gln-Leu-Arg at the 484th to 489th of the SEQ ID NO: 14 is substituted with Pro-Gly-Ala-Ala-His-Tyr by using the mutated primer of the SEQ ID NO: 18 is referred to as pCDNA-dHGF-NG-G3.

Next, pCDNA-dHGF-NG and each mutated vector (pCDNA-dHGF-NG-G1, pCDNA-dHGF-NG-G2 and pCDNA-dHGF-NG-G3), were respectively transfected into Human embryonic kidney 293T cells (DuBridge RB, et al., Molecular Cellular Biology, 7, 379-387 (1987)). For the transfection, the 293T cells were precultured in the Dulbecco's Modified Eagle's Medium (DMEM) to which 10% by volume of fetal calf serum (FCS) is added. This DMEM was replaced with the serum-free DMEM just before transfection, and transfection was performed by the lipofection method using LIPOFECTAMINE 2000 (made by Invitrogen). After transfection was complete, the culture in the serum-free DMEM was continued, and 6 hours after the transfection, heparin was

added to a concentration of 1 µg/mL. The culture was further continued for three days to allow HGF-NG precursor protein or HGF precursor protein variant (the HGF-G1 precursor protein, HGF-G2 precursor protein, or HGF-G3 precursor protein) produced
5 by each vector to accumulate in the serum-free DMEM, respectively. Three days after, the DMEM was collected from three petri dishes respectively, mixed, filtered through a 0.22 µm filter, and kept at -80°C until being purified. The concentration of the HGF-NG precursor protein, HGF-G1 precursor
10 protein, HGF-G2 precursor protein, or HGF-G3 precursor protein was analyzed by ELISA. The ELISA method was performed by using the Immunis kit (made by Institute of Immunology) in accordance with the protocol described in the kit.

The above-mentioned medium was thawed and filtered again
15 through a 0.22 µm filter. To this, Heparin Sepharose resin (made by Amersham Biosciences) equilibrated with 50 mM Tris-HCL (pH 7.5), 0.01% Tween 80 and 0.3 M NaCl was added in order to allow the HGF-NG precursor protein, HGF-G1 precursor protein, HGF-G2 precursor protein, or HGF-G3 precursor protein to bind to the
20 resin at room temperature. The resin was then washed with 50 mM Tris-HCL (pH 7.5), 0.01% Tween 80 and 0.3 M NaCl and the HGF-NG precursor protein, HGF-G1 precursor protein, HGF-G2 precursor protein, or HGF-G3 precursor protein was eluted with 50 mM Tris-HCL (pH 7.5), 0.01% Tween 80 and 2 M NaCl for partial
25 purification.

To the HGF-NG precursor protein, HGF-G1 precursor protein, HGF-G2 precursor protein, or HGF-G3 precursor protein (200 ng each) partially purified with the heparin resin, half the amount (100 ng) of Genenase I (made by New England Laboratory) was added.

The mixture was treated in 50 mM Tris-HCL (pH 7.5), 0.01% Tween 80 and 2 M NaCl at 25°C for 12 hours. Meanwhile, HGF-NG precursor protein having a natural α chain- β chain boundary sequence, to which any Genenase I recognition sequence was not inserted, was
5 incubated in 50 mM Tris-HCL (pH 7.5), 0.01% Tween 80 and 2 M NaCl with 1% by volume of FCS added thereto at 37°C for 12 hours.

Each active HGF precursor protein variant (active HGF-G1, active HGF-G2, or active HGF-G3) obtained after Genenase I treatment was subjected to the Western blot under reducing or
10 nonreducing condition as follows.

Western blot: Active HGF-G1, active HGF-G2, or active HGF-G3 after Genenase I treatment was subjected to SDS-PAGE under reducing condition (in the presence of 100 mM DTT) or nonreducing condition, and transferred onto PVDF membrane. The
15 protein transferred onto the PVDF membrane was detected using, as a probe, human HGF polyclonal antibody prepared by immunizing a rabbit with human HGF (Matsumoto K, et al., Proceedings for National Academy of Science of the United States of America, 89, 3800-3804 (1992)).

20 Fig. 1 shows the results of western blot analysis in the SDS-PAGE under reducing condition. With regard to HGF-NG precursor protein having a natural α chain- β chain boundary sequence, to which any Genenase I recognition sequence was not inserted, Genenase I treatment did not affect band positions.
25 In either case of treatment with or without Genenase I, a band was found at the position of single-stranded HGF-NG precursor protein (Lanes 1 and 2 in Fig. 1). From this, it was confirmed that HGF-NG precursor protein having a natural α chain- β chain boundary sequence was not cleaved by Genenase I. Meanwhile,

after FCS was added to this HGF-NG precursor protein and the protein was treated at 37°C for 12 hours, two bands of α chain and β chain were detected, which confirmed that the protein was activated (Lane 3). In each case of HGF-G1 precursor protein, HGF-G2 precursor protein, and HGF-G3 precursor protein to which a Genenase I recognition sequence was inserted, without Genenase I treatment, a band was found at the same position as the single-stranded HGF-NG precursor protein (Lanes 4, 6 and 8 in Fig. 1). However, after Genenase I treatment, bands were detected at the same positions as those of the α chain and β chain of HGF-NG (Lanes 5, 7 and 9 in Fig. 1), which confirmed that each protein was converted into a double-stranded structure.

Fig. 2 shows the results of western blot analysis in the case of SDS-PAGE under nonreducing condition. HGF-NG precursor protein having a natural α chain- β chain boundary sequence was detected as a single band in the cases not only before but also after activation treatment with FCS (Lanes 1 and 3 in Fig. 2). This means that the active HGF-NG structure was held with S-S bond. The HGF-NG was not activated by Genenase I and did not undergo any other cleavage. Therefore, it remained as a single band (Lane 2 in Fig. 2). Also, in each case of HGF-G1 precursor protein, HGF-G2 precursor protein, and HGF-G3 precursor protein to which a Genenase I recognition sequence was inserted, not only before (Lanes 4, 6 and 8 in Fig. 2) but also after (Lanes 5, 7 and 9 in Fig. 2) Genenase I treatment, only a single band was detected. These facts showed that the HGF precursor protein variants were held with S-S bond even after being cleaved by Genenase I.

Example 2

The active HGF-G1 or active HGF-G3 prepared in Example 1 was subjected to SDS-PAGE under reducing condition, and transferred onto PVDF membrane. The protein transferred onto the PVDF membrane was stained with Coomassie Brilliant Blue, and the 30 kDa band corresponding to the HGF β chain was cut out. The sequence of the N-terminal amino acid was analyzed using an amino-acid sequencer (Procise 491 cLC made by Applied Biosystems). The N-terminal amino acid sequences of the bands thought to be the β chains of HGF-G1 and HGF-G3 were both VVNGI (Val-Val-Asn-Gly-Ile). This confirmed that the HGF precursor protein variant to which a Genenase I recognition sequence was inserted was cleaved at the site of the Genenase I recognition sequence as designed.

Table 2

	Amino-acid sequence	Sequence listing
HGF	...CAKTKQLR↓VVNGI...	SEQ ID NO: 14;482..494
HGF-G1	...CAKTKQHY↓VVNGI...	SEQ ID NO: 19;482..494
HGF-G3	...CAPGAAHY↓VVNGI...	SEQ ID NO: 20;482..494
HGF β chain	VVNGI...	SEQ ID NO: 14;490..494
HGF-G1 β chain	VVNGI...	SEQ ID NO: 19;490..494
HGF-G3 β chain	VVNGI...	SEQ ID NO: 20;490..494

Example 3

Canine kidney epithelial cells (MDCK) (Montesano R, et al., Cell, 66, 697-711 (1991)) were suspended in DMEM (containing 10% by volume of FCS) and plated into each well of a 24-well

plate at 1×10^4 cells/well (480 μL /well). A test sample in an amount of 20 μL containing active 5-amino-acid-deleted-type human HGF, the active HGF-G1, active HGF-G2 or active HGF-G3 prepared in Example 1, was added thereto. After incubation at 5 37°C for 20 hours, the presence or absence of scattering was observed with a microscope (Fig.3). The test sample was prepared so as that the concentration of 5-amino-acid-deleted-type human HGF, active HGF-G1, active HGF-G2 or active HGF-G3 was 2, 5 or 10 ng/mL after 20 μL of the 10 test sample was added to the medium.

None of HGF-G1 precursor protein, HGF-G2 precursor protein and HGF-G3 precursor protein to which a Genenase I recognition sequence was inserted showed cell migration action before Genenase I treatment (Fig.3; -). After Genenase I treatment 15 (Fig.3; +), all of active HGF-G1, active HGF-G2 precursor protein and HGF-G3 showed cell migration action equivalent to active 5-amino-acid-deleted-type human HGF. This confirmed that the HGF precursor protein variant to which a Genenase I recognition sequence was inserted was converted into a 20 double-stranded chain structure by Genenase I, and therefore, became an active HGF precursor protein variant having HGF action.

Industrial applicability

25 The active HGF protein variant of the present invention can be used as an alternative medicament of HGF because it has substantially the same action as HGF.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

Claims

1. A Hepatocyte growth factor (HGF) precursor protein variant, in which a peptide structure comprises a sequence including a peptide chain X inserted between an α chain of HGF where 1 to 20 amino-acid residues from the C-terminus of the α chain are deleted, and a β chain of HGF where 1 to 20 amino-acid residues from the N-terminus of the β chain are deleted; wherein

- (i) the peptide chain X has an amino-acid sequence of 2 to 20 residues,
- (ii) the peptide chain X can be cleaved by a protease reaction,
- (iii) a protein obtained by cleaving at one site of the peptide chain X has HGF action, and
- (iv) the peptide chain X has a protease recognition sequence,

wherein the protease recognition sequence is at least one protease recognition sequence that is Genenase I recognition sequence, Enterokinase recognition sequence, blood coagulation factor Xa recognition sequence, thrombin recognition sequence, TEV protease recognition sequence, Rhinovirus 3C protease recognition sequence or Furin recognition sequence.

2. The HGF precursor protein variant according to Claim 1, wherein the protease recognition sequence is His-Tyr or Tyr-His.

3. The HGF precursor protein variant according to Claim 1 or 2, wherein the HGF is of human, canine or feline origin.

4. The HGF precursor protein variant according to Claim 1 or 2, wherein the HGF is of human origin.

5. The HGF precursor protein variant according to Claim 4, wherein the HGF is

- (a) a protein comprising an amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2;
- (b) a protein comprising an amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2 wherein one or several amino-acid residues are deleted, substituted or added, and having hepatocyte proliferation action of HGF; or

- (c) a protein comprising an amino-acid sequence having a 80% homology or more with the amino-acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2, and having hepatocyte proliferation action of HGF.
6. The HGF precursor protein variant according to Claim 4, wherein the α -chain is
- (a) an amino-acid sequence from the 32nd to the 494th of the sequence represented by SEQ ID NO: 1,
and the β -chain is
- (b) an amino-acid sequence from the 495th to the 728th of the sequence represented by SEQ ID NO: 1;
or alternatively the α -chain is
- (c) an amino-acid sequence from the 32nd to the 489th of the sequence represented by SEQ ID NO: 2,
and the β -chain is
- (d) an amino-acid sequence from the 490th to the 723th of the sequence represented by SEQ ID NO: 2.
7. An active HGF protein variant obtained by cleaving one site of the peptide chain X in the HGF precursor protein variant according to any one of Claims 1 to 6.
8. The active HGF protein variant according to Claim 7, wherein the cleavage is caused by Genenase I treatment.
9. An *in vitro* method for producing the active HGF protein variant as defined in claim 7, comprising producing the HGF precursor protein variant according to Claim 1, and simultaneously or subsequently converting the HGF precursor protein variant into an active HGF protein variant by cleaving at least one site of the peptide chain X.
10. The method for producing the active HGF protein variant according to Claim 9, wherein the cleavage is caused by a protease treatment or a chemical treatment.
11. The method for producing the active HGF protein variant according to Claim 10, wherein the protease is at least one protease that is Genenase I, Enterokinase, blood coagulation factor Xa, thrombin, TEV protease, Rhinovirus 3C protease or Furin.

12. The method for producing the active HGF protein variant according to Claim 9, comprising producing a single-stranded HGF precursor protein variant having a sequence into which a peptide chain X having a sequence of His-Tyr or Tyr-His and 2 to 20 amino acid residues, is inserted, and simultaneously or subsequently treating the HGF precursor protein variant with Genenase I.
13. The method for producing the active HGF protein variant according to Claim 9, wherein a DNA encoding the HGF precursor protein variant according to Claim 1 and a DNA encoding a protease to cleave the peptide chain X are simultaneously introduced into a host, the HGF precursor protein variant and the protease to cleave the peptide chain X are simultaneously expressed, and the peptide chain X is cleaved by the protease.
14. A medicament comprising the active HGF protein variant according to Claim 7 or 8.

Fig.1




Fig.2




Fig.3

