



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : C07D 205/04, 207/16, 211/34	A1	(11) International Publication Number: WO 90/02732 (43) International Publication Date: 22 March 1990 (22.03.90)
(21) International Application Number: PCT/US89/03960 (22) International Filing Date: 12 September 1989 (12.09.89) (30) Priority data: 244,203 13 September 1988 (13.09.88) US (71) Applicant: SEPRACOR, INC. [US/US]; 33 Locke Drive, Marlborough, MA 01752 (US). (72) Inventor: ZEPP, Charles, M. ; 19 Highland Street, Berlin, MA 01502 (US). (74) Agent: MISROCK, S., Leslie; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US). (81) Designated States: AT (European patent), AU, BB, BE (Eu- ropean patent), BF (OAPI patent), BG, BJ (OAPI pa- tent), BR, CF (OAPI patent), CG (OAPI patent), CH (European patent), CM (OAPI patent), DE (European patent), DK,		FI, FR (European patent), GA (OAPI patent), GB (European patent), HU, IT (European patent), JP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL (European patent), NO, RO, SD, SE (European pa- tent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent). Published <i>With international search report.</i>
(54) Title: METHODS FOR PREPARING CAPTOPRIL AND ITS ANALOGUES		
(57) Abstract This invention relates to novel methods for converting a diastereomeric mixture of S-protected derivatives of an orally ac- tive inhibitor of an angiotensin-converting enzyme (ACE) and its analogues into its separate optically resolved diastereomeric components. Specifically the invention relates to methods for the preparation of optically purified captopril and its analogs from racemic precursors. This resolution process is achieved through the fractional crystallization of S-protected derivatives of capto- pril and its precursors, which derivatives are useful for the reason that they are (1) easily prepared from novel precursors, (2) re- solvable to their optically purified stereoisomeric species and (3) convertible to non-derivatized stereoisomeric species which correspond to the pharmacologically active inhibitor and its analogues. Novel methods for preparing the derivatives and their precursors are also noted herein. In addition, the novel derivatives and their precursors are also described herein.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

METHODS FOR PREPARING
CAPTOPRIL AND ITS ANALOGUES

1.0 INTRODUCTION

This invention relates to novel methods for converting a diastereomeric mixture of S-protected derivatives of an orally active inhibitor of an angiotensin-converting enzyme (ACE) and its analogues into its separate optically resolved diastereomeric components. Specifically the invention relates to methods for the preparation of optically purified captopril for the preparation of optically purified captopril and its analogs from racemic precursors. This resolution process is achieved through the fractional crystallization of S-protected derivatives of captopril and its precursors, which derivatives are useful for the reason that they are (1) easily prepared from novel precursors, (2) resolvable to their optically purified stereoisomeric species and (3) convertible to non-derivatized stereoisomeric species which correspond to the pharmacologically active inhibitor and its analogues.

Novel methods for preparing the derivatives and their precursors are also noted herein. In addition, the novel derivatives and their precursors are also described herein.

2.0 BACKGROUND OF THE INVENTION

Enzymatic resolution and fractional crystallization procedures have long been known and exploited for resolving racemic mixtures. Unfortunately, many chiral compounds of commercial significance are not adequately prepared using these procedures.

2.1 SIGNIFICANCE OF OPTICAL PURITY

Many organic compounds exist in optically active forms, *i.e.*, they have the ability to rotate the plane of plane-polarized light. The property of optical activity is due to molecular asymmetry about carbon atoms that are linked to four different atoms or chemical groups. In describing an optically active compound, the prefixes D and L or R and S are used to denote the configuration of the molecule about its chiral center(s). The prefixes (+) and (-) or d and l are employed to designate the sign of rotation of plane-polarized light by the compound, with (-) or l meaning that the compound is levorotatory. A compound prefixed with (+) or d is dextrorotatory. Compounds of a given chemical structure, which differ from one another only in the configuration of chemical groups about one asymmetric carbon atom, or chiral center as it is sometimes called, are called stereoisomers. Where there are *n* asymmetric carbons or chiral centers, the number of potential stereoisomers increases to 2^n . Thus, a molecule with three chiral centers would have eight possible stereoisomers.

Where these stereoisomers are identical except that they are non-superimposable mirror images of one another, the molecules are referred to as enantiomers. Stereoisomers which are not non-superimposable mirror images of other stereoisomers of the same compound are described as diastereomers (e.g. 2R,3R-tartaric acid and 2R,3S-tartaric acid are diastereomers whereas 2R,3R-tartaric acid and 2S,3S-tartaric acid are enantiomers). A mixture of enantiomers is called an enantiomeric or racemic mixture, and as used herein this term is applied to mixtures of enantiomers in any proportion. A mixture of

diastereomers is referred to as a diastereomeric mixture, where as the term is used herein the diastereomers may be present in any proportion.

While the structural differences between stereoisomers are subtle and of little consequence in ordinary chemical reactions, they may be profound where biological systems are concerned, *i.e.*, if the compounds are utilized in enzyme-catalyzed reactions. Thus, the L-amino acids are metabolized in humans but the corresponding D-analogs are not, and only D-glucose can be phosphorylated and processed into glycogen or degraded by the glycolytic and oxidative pathways of intermediary metabolism. Similarly, beta blockers, pheromones, prostaglandins, steroids, flavoring and fragrance agents, pharmaceuticals, pesticides, herbicides and many other compounds exhibit critical stereospecificity. In the field of pesticides, Tessier [Chemistry and Industry, March 19, 1984, p. 199] has shown that only two of the eight stereoisomers of deltamethrin, a pyrethroid insecticide, have any biological activity. The same statement concerning the concentration of bioactivity in a single isomer can be made about many other pesticides, including the phenoxypropanoates and halopropanoate derivatives, each containing one chiral center and existing in the form of two optical isomers.

Stereochemical purity is of equal importance in the field of pharmaceuticals, where 12 of the 20 most prescribed drugs exhibit chirality. A case in point is provided by naproxen, or (+)-S-2-(6-methoxy-2-naphthyl)propanoic acid, which is one of the two most important members of a class of 2-arylpropanoic acids with non-steroidal anti-inflammatory activity used, for instance, in the management of arthritis. In this case, the S(+) enantiomer of the drug is known to be 28 times more therapeutically potent than its R(-) counterpart.

Still another example of chiral pharmaceuticals is provided by the family of beta-blockers; the L-form of propranolol is known to be 100 times more potent than the D-enantiomer.

In particular, interest in captopril exists since it belongs to a new class of antihypertensive agents. Its specific action resides in its acting as a specific competitive inhibitor of angiotensin I-converting enzyme (ACE) which converts angiotensin I to angiotensin II. Although the actual mechanism of its action is not fully elucidated, it has been shown that captopril reduces blood pressure and results in beneficial hemodynamic effects in patients with congestive heart failure. More specifically, captopril (commercially available from Squibb as CAPOTEN[™]), contains a 2-methyl-3-thiolpropanoyl segment linked via an amide bond to L-proline. When the chiral carbon atom in the 2-methyl-3-thiopropionyl segment has the S-configuration, this diastereomer is known to be 100 times more active than the corresponding diastereomer where the segment has the R-configuration. Consequently, a variety of methods to produce captopril have been undertaken, including preparation by enzymatic resolution and by fractional crystallization of its diastereomers.

Synthesis of compounds with asymmetric centers by standard organic synthetic techniques generally leads to a racemic or diastereomeric mixture which, in the aggregate, may have a relatively low specific bioactivity since certain of the stereoisomers in the mixture are likely to be biologically inactive. As a result, larger quantities of the material must be used to obtain an effective dose, and manufacturing costs are increased due to the co-production of stereochemically "incorrect" and hence, inactive ingredients.

In some instances, certain isomers may actually be deleterious rather than simply inert. For example, the D-enantiomer of thalidomide was a safe and effective sedative when prescribed for the control of morning sickness during pregnancy. However, its L-thalidomide counterpart was discovered to be a potent teratogen.

2.2 CONVENTIONAL MEANS OF OBTAINING OPTICALLY PURE COMPOUNDS

Methods are available for stereoselective synthesis. For example, a synthetic pathway to optically pure deltamethrin [(1R,3R)-3-(2,2-dibromovinyl)-2,2-dimethyl cyclopropane carboxylic acid] has been developed, but the process is lengthy, complex and costly [Tessier, J., Chem. and Ind., March 9, 1984, p.199]. Moreover, a synthetic scheme capable of producing one specific stereoisomer cannot be applied in a general way to obtain other optically active compounds. What is needed is a generalized approach to the separation of racemic and diastereomeric mixtures produced by nonstereoselective chemical reactions, and a number of approaches have been used.

The terms "racemic mixture" and "diastereomeric mixture" as used herein, refer to mixtures of a first and a second stereoisomer in any proportions, such that the first and second stereoisomers are enantiomers or diastereomers, respectively. Further, it is intended that the term "resolution" as used herein will refer to the transformation of a racemic or diastereomeric mixture, as defined above, into two product mixtures, in each of which the proportions of the two above defined stereoisomers may be different from both the starting racemic or diastereomeric mixture and from each other, the proportion being greater in one and necessarily smaller in the other.

The term "resolved" is intended to refer to a quantity of any compound, capable of resolution, which has undergone the process of resolution defined above to yield an optically active product material. Finally, the terms "stereospecific" and "stereoselective" as used herein are synonymous.

A widely used approach has been the selective precipitation of desired stereoisomeric compounds from diastereomeric mixtures. Such diastereomeric mixtures are created by addition of a chiral and optically purified resolving agent to a racemic mixture, which then induces fractional crystallization. Such chiral and optically purified resolving agents form complexes, salts or covalent compounds with the racemic species to be resolved. Said diastereomeric complexes, salts and covalent compounds now possess two or more chiral centers wherein one chiral center in one diastereomer has the same configuration as the corresponding chiral center in another diastereomer, while the diastereomers' other corresponding chiral centers have the opposite configuration. The individual diastereomers have different physical properties (e.g., solubility) and are hence readily separable from one another. For example, Yoshioka et al., [U.S. Patent No. 3,879,451] treated a mixture of (+)-cis- and (+)-trans-chrysanthemic acids with an optically active aromatic amine and recovered the resulting diastereomeric amine salts of (+)-cis- and (+)-trans-chrysanthemic acids by crystallization. Paven et al., [U.S. Patent No. 4,257,976] resolved D,L-cis-chrysanthemic acid and D,L-trans-chrysanthemic acid by treating the mixtures with L or D N-methyl-ephedrine to form the corresponding salts, which were then hydrolyzed after isolation to produce the resolved acids. Halmos [U.S. Patent No. 4,151,198] treated a mixture of N-acyl-D,L(+)-phenylalanine isomers with D(-

) -2-(2,5-dimethylbenzylamino)-1-butanol to obtain a crystalline salt, from which N-acyl-L(+)-phenylalanine could be recovered. Kameswaran [U.S. Patent No. 4,454,344] isolated (+) 2-(p-difluoromethoxyphenyl)-3-methylbutanoic acid by treating the racemic acid with an optically active amine. The S(+)-enantiomer of naproxen can be obtained by stereoselective crystallization of a diastereomeric salt formed with an amine resolving agent such as cinchonidine, glucamine, or N-methylglucamine [Harrison, I.T. et al., J. Med. Chem., 13:203 (1970); Felder, E. et al., U.K. Patent Appl. GB2025968A (1980)].

In some cases, a two-step method has been used, as when Dannenberg et al. [U.S. Patent No. 4,285,884] resolved racemic D,L-alpha-aminocarboxylic acids by first treating the mixture with an aromatic o-hydroxyaldehyde to obtain an azomethine derivative. This derivative was then immediately treated with an optically active amine base to produce a salt which was isolated. By subjecting the salt to acid hydrolysis, the desired alpha-aminocarboxylic acid isomer was obtained.

Regarding captopril, fractional crystallization methods have been noted in resolving captopril, captopril analogues and their precursors, generally based on the use of optically active amines capable of forming diastereomers with the acid precursors of captopril (e.g., 2-methyl-3-thiolpropanoic acid) and its analogues. M.A. Ondetti in U.S. Patent Nos. 4,046,889 (Col. 4, lines 52-55); 4,316,906 (Col. 3, lines 7-11); 4,105,776 (Col. 8, lines 59-63) and 4,154,840 (Col. 8, lines 55-59) discloses that fractional crystallization techniques are available to resolve the racemic precursors of captopril. In U.S. Patent No. 4,346,045, N. De Heij utilized a fractional crystallization process to isolate D-(-)-S-benzoyl- β -mercaptoisobutyric acid for subsequent conversion to captopril. D-(+)-N-

benzyl- α -phenethylamine is used as one of the fractional crystallization reagents (i.e., resolving agents). J. Houbiers in U.S. Patent No. 4,585,595, also prepared the same captopril precursor. Cinchonidine, D-(-)-2-aminobutanol or a derivative thereof were used as fractional crystallization reagents. Interestingly enough, De Heij also discloses in European Patent Application No. 0,008,831 A1 that cinchonidine is a suitable resolving agent as described in his Dutch Patent Application No. 7,809,121. Also, 1,2-diphenylethylamine and 2-amino-1-butanol have been employed in resolutions (J. Iwao, et al., Jpn. Kokai Tokkyo Koho 79: 151912 (1979); H.-J.-P. Marie. Jpn. Kokai Tokkyo Koho 80: 38386 (1980); N. Ohashi, et al., Jpn. Kokai Tokkyo Koho 81: 7756 (1981)). However, N. Ohashi, et al., in U.S. Patent No. 4,297,282 noted that racemic- α -methyl- β -acylthiopropionic acids are not given to fractional crystallization using standard optically active amines such as quinidine, ephedrine, α -methylbenzylamine and brucine (Col. 1, lines 44 to Col. 2, line 2). In addition, even though the use by N. Ohashi, et al. of optically active amines such as 1-(α -naphthyl)-ethylamine and α,β -diphenylethylamine proved to be more successful, it was noted that the procedures were very expensive, cumbersome to effect, and capable of producing only relatively low yields (C.J. Sih, PCT No. WO87/05328 p. 2).

In addition, non-optically active amines have been employed in fractional crystallization resolution steps to prepare captopril. D.H. Naur, et al., J. Pharm. Sci. 73(12): 1843-4 (1984) used dicyclohexylamine to resolve the diastereomers of racemic N-(L-proline)-3-halo-2-methylpropanamide precursor used in a captopril synthesis. Following the isolation of that resolved N-[3-halo-2-methylpropanoyl]proline, the amide can be converted to the corresponding diastereomeric S-protected derivatives

of captopril by treating the amide with either a sodium dialkyldithiocarbamate or potassium xanthogenate, as demonstrated by D.K. Kim in U.K. patent No. 2,170,806A. Dicyclohexylamine was also used by I. Castellet-Linan, Spain Patent No. 548154, to resolve the benzoylthio-protected captopril. However, the use of non-optically active amines also incurs an added expense and additional synthetic steps in using the amine.


Consequently, although some of the above procedures successfully resolved racemic or diastereomeric mixtures, not all could be successfully applied to captopril. Moreover, some of these fractional crystallization procedures using optically active reagents did not produce diastereomers with sufficiently different physical properties when applied to captopril and its precursors. Thus, fractional crystallization using such reagents is not always possible.

2.3 STEREOSPECIFIC RESOLUTIONS BY ENZYME-CATALYZED BIOCONVERSIONS OF ORGANIC COMPOUNDS


Various enzymes have been used for bioconversions of assorted organic compounds. Many different classes of enzymes have been used for the resolution of stereoisomers, including hydrolases (especially the lipases, proteases and esterases such as chymotrypsin), lyases and oxidoreductases (e.g., amino acid oxidases and alcohol reductases).

Regarding captopril, enzymatic resolution procedures have been preferably undertaken since there have been problems with fractional crystallization methods. M. Shimazaki, et al., Chem. Pharm. Bull. 30(9): 3139-45 (1982) used a microbiological step developed by Cohen et al., J. Org. Chem. 41, 3505 (1976) to convert isobutyric acid by a bacterial hydroxylation to D-3-hydroxy-2-

methylpropanoic acid which was used as a precursor to synthesize captopril. A. Sakimae *et al.*, U.S. Patent No. 4,629,701 also prepared captopril and captopril analogue precursors represented by the formula,



wherein Y_2 is an alkyl group, preferably C_{1-6} alkyl group, an aralkyl group, preferably C_{7-18} aralkyl group or an aryl group, preferably C_{6-26} aryl group; Y_1 is an alkyl group, preferably C_{1-6} alkyl group; and Y is 1 or 2, by allowing a source containing an enzyme capable of asymmetrically hydrolyzing an ester to act on the ester represented by the formula,



wherein Y_2 , Y_1 , and Y have the same meanings as those mentioned above; and R_3 is an alkyl group, preferably C_{1-6} alkyl group. In PCT No. WO 87/05328 by C.J. Sih, a microbiological deesterification similar to the one employed by Sakimae was used except that Y_2 was either an acyl radical in straight chain, branched chain or cyclic configuration having 1 to about 12 carbon atoms; cycloalkane radicals having 5 to 7 carbon atoms; or benzoyl, naphthoyl, biphenoyl and carbobenzoxy radicals containing substituents such as nitro, halogen, methyl or alkoxy groups on the aromatic ring.

However, the efficiency of these methods is problematic since the enzyme has to be replenished, a cosolvent or emulsifier may have to be employed to facilitate the substrate susceptibility to the enzymatic processing, and some of the enzymes used in these processes are expensive.

Consequently, neither enzymatic resolution nor the previously noted fractional crystallization steps appear to be without certain disadvantages. Therefore, another approach to resolving captopril and its analogues is needed.

3.0 SUMMARY OF THE INVENTION

This invention relates to novel methods for converting a diastereomeric mixture of S-protected derivatives of an orally active inhibitor of an angiotensin-converting enzyme (ACE) and its analogues into its separate optically resolved diastereomeric components. Specifically the invention relates to methods for the preparation of optically purified captopril and its analogs from racemic precursors. This resolution process is achieved through the fractional crystallization of S-protected derivatives of captopril and its precursors, which derivatives are useful for the reason that they are (1) easily prepared from novel precursors, (2) resolvable to their optically purified stereoisomeric species and (3) convertible to non-derivatized stereoisomeric species which correspond to the pharmacologically active inhibitor and its analogues.

Novel methods for preparing the derivatives and their precursors are also noted herein. In addition, the novel derivatives and their precursors are also described herein.

4.0 BRIEF DESCRIPTION OF THE FIGURES

This invention may be more readily understood by reference to the following detailed description of the invention and figures in which

FIG. 1. is a schematic representation of (1) the methods for preparing S-protected captopril and its analogues, (2) the use of the precursors of the S-protected captopril and its analogues and (3) the methods for preparing captopril and its analogues from their S-protected derivatives. In addition, it is noted that the undesired protected diastereomeric amide can be hydrolyzed to isolate the amide's amine for recycling to the coupling (amidization) step.

5.0 DETAILED DESCRIPTION OF THE INVENTION

It is an object of the present invention to cover novel methods for resolving a diastereomeric mixture of S-protected derivatives of captopril and its analogs and for subsequently de-protecting one of the separated diastereomeric species to yield the optically purified ACE inhibitor.

Compositions of matter regarding the S-protected derivatives and their precursors, and methods of producing these compositions of matter are also noted herein.

The amide compounds which are the primary subject matter of this invention contain two chiral centers wherein one of the chiral centers is located alpha to the carboxyl carbon in the thiol alkanoyl segment of the amide and the other chiral center is alpha to the nitrogen which forms the amide. Consequently, the compounds may be produced as mixtures of diastereomers. The compounds are useful in that they directly, without further chemical modification, undergo fractional crystallization to allow separation of their component diastereomers. This property of the subject compounds is described herein as "self-fractional crystallization".

Self-fractional crystallization is to be distinguished from the usual practice of fractional crystallization to resolve stereoisomers whereby the desired stereoisomer is separated as a diastereomeric derivative (for example, resolution of a racemic mixture of a carboxylic acid by fractional crystallization of a diastereomeric mixture of salts formed between the former and an optically active amine). This usual practice requires regeneration of the underivatized compound and recovery of the amine "resolving agent". In the present invention it is an unexpected property of the subject compounds that they undergo self-fractional crystallization with sufficiently high selectivity and efficiency that the S-protected drug precursor requires no further stereochemical purification. This self-fractional crystallization behavior is attributed to the physical properties resulting from the novel thiol protecting group.

Specifically, racemic S-functionalized carboxylic acid compounds (precursors) are described herein, as are the S-functionalized amide compounds which result from their reactions with alpha-amino acids or alpha-amino acid derivatives. Said S-functionalized amide compounds may undergo self-fractional crystallization and be further converted by removal of the thiol protecting group to the desired single optical isomer of the desired compound. As a specific example of the utility of the claimed compounds, reaction of the claimed racemic S-functionalized carboxylic acid compound (after conversion to a suitable acylating agent) with L-proline results in formation of a diastereomeric mixture of amide compounds which undergoes self-fractional crystallization and is further converted by de-protection to captopril.


Embodiments of this invention, therefore, pertain to novel methods for preparing optically purified captopril and its analogues from a diastereomeric mixture of the S-protected derivatives, which resolved derivatives are de-protected to form captopril and its analogues.

5.1 NOVEL COMPOSITIONS OF MATTER PERTAINING TO DERIVATIVES AND PRECURSORS OF CAPTOPRIL AND ITS ANALOGUES


5.1.1 THIOL-PROTECTED DERIVATIVES OF CAPTOPRIL AND ITS ANALOGUES

Regarding the thiol-protected derivatives of captopril and its analogues, which are described herein, the importance of these compositions of matter resides in the protecting group's facilitation of the compositions' self-fractional crystallization characteristics. Consequently, a diastereomeric mixture of the thiol-protected derivatives is separable to the individual diastereomeric species, one of which is then converted to the corresponding diastereomeric de-protected captopril species and its analogues by a deprotection step.

Regarding heterocyclic amine compounds or amine segments in cyclic amides herein represented by the formulae




wherein R₅ is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, halide, phenyl and substituted phenyl, and p is 2, 3, or 4; and



wherein R is selected from the group consisting of hydroxy, amino and lower alkoxy, R_3 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, arylthio, halide, phenyl and substituted phenyl, and m is 2, 3, or 4, both of the formulae represent rings ranging in sizes from four to six. More specifically, where m is 3 regarding formula II it corresponds to the substituent group R_7 described later herein as being selected from the group consisting of L-proline (where R is hydroxyl and R_3 is hydrogen) and substituted L-proline selected from the group consisting of 4S-(phenylthio)-L-proline, 4S-hydroxy L-proline, lower alkyl ester of L-proline and amide of L-proline. In addition, the R_5 and R_3 groups are each represented as such to note that they can substitute for any methylene ($-\text{CH}_2-$) hydrogen(s) of the ring. The substitutions pertain to those replacements which either provide a heterocyclic amine which is non-optically active or which replacement involves a single stereoconfiguration at that chiral carbon which is the site of substitution. Examples of the former are geminal substitutions or those which provide meso compounds (*i.e.*, having a plane of symmetry). An example of the latter is provided by substituted L-proline wherein the substituent group R_3 is 4S-(phenylthio).

One of the novel compositions of matter prepared herein is a compound of the formula III



wherein R is selected from the group consisting of hydroxy, amino and lower alkoxy; R_1 is selected from the group consisting of lower alkyl, lower alkoxy, hydroxy, halide, phenyl and substituted phenyl; R_2 is selected from the group consisting of O, S and imino; R_3 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, arylthio, halide, phenyl and substituted phenyl; R_4 is a cyclic secondary amino; m is 2, 3 or 4; C_a has an S or R configuration; C_b has a D or L configuration; n is 1, 2 or 3; and basic salts thereof.

The compound of formula III also encompasses the compound wherein said cyclic secondary amino has the formula I as described above; said R_2 is an imino which has the formula NR_6 wherein R_6 is selected from the group consisting of hydrogen, hydroxy, lower alkoxy, lower alkyl, phenyl and substituted phenyl; said phenyl substituent is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy; and said lower alkoxy and lower alkyl groups have up to seven carbon atoms.


Three further descriptions of the compound include the compound of the formula III:

- 1) wherein R is hydroxy, R_1 is lower alkyl wherein said lower alkyl is methyl, R_2 is S, R_3 is hydrogen, 4S-hydroxy or 4S-phenylthio, n is 1, m is 3, R_4 is a cyclic secondary amino of the formula I above wherein R_5 is hydrogen and p is 3, and C_b has an L configuration;

2) wherein R is lower alkoxy wherein said lower alkoxy is methoxy or t-butoxy, R_1 is lower alkyl wherein said lower alkyl is methyl, R_2 is S, R_3 is hydrogen, 4S-hydroxy or 4S-phenylthio, n is 1, m is 3, R_4 is a cyclic secondary amino of the formula of I above wherein R_5 is hydrogen and p is 3, and C_b has an L configuration; and

3) wherein R is lower alkoxy wherein said lower alkoxy is methoxy, R_1 is lower alkyl wherein said lower alkyl is methyl, R_2 is O, R_3 is hydrogen, 4S-hydroxy or 4S-phenylthio, n is 1, m is 3, R_4 is a cyclic secondary amino of the formula of I above, wherein R_5 is hydrogen and p is 3, and C_b has an L configuration.

The novel compositions of matter also cover a compound of the formula IV



and basic salts thereof wherein the substituents are defined as above.

Further descriptions of the compound include the compound of the formula IV:


1) wherein R_2 is S, R_4 is a cyclic secondary amino of formula I above wherein R_5 is hydrogen and p is 3, and R_7 is a) L-proline or b) substituted L-(proline) wherein the substituted L-proline is

- (i) 4S-(phenylthio)proline,
- (ii) 4S-hydroxyproline or

(iii) lower alkyl ester of L-proline wherein said lower alkyl ester of L-proline is methyl ester of L-proline or t-butyl ester of L-proline; and

2) wherein R_2 is O, R_4 is a cyclic secondary amino of formula I above wherein R_5 is hydrogen and p is 3, and R_7 is substituted L-proline wherein said substituted L-proline is lower alkyl ester of L-proline wherein said lower alkyl ester of L-proline is methyl ester of L-proline.

More particularly, the novel compositions of matter pertain to a compound of the formula V



wherein R_8 is selected from the group consisting of O and S; R_9 is pyrrolidino; and basic salts thereof wherein all the other substituents are defined as above.

Further descriptions of the compound include a compound:

- 1) wherein R_8 is S, and R_7 is
 - a) L-proline or
 - b) substituted L-proline wherein the substituted L-proline is
 - (i) 4S-(phenylthio)-L-proline
 - (ii) 4S-hydroxy-L-proline or
 - (iii) lower alkyl ester of L-proline wherein said lower alkyl ester of L-proline is

methyl ester of L-proline or
t-butyl ester of L-proline;
and


2) wherein R_2 is O, and R_7 is substituted L-proline wherein said substituted L-proline is lower alkyl ester of L-proline wherein said lower alkyl ester of L-proline is methyl ester of L-proline.

5.1.2 PRECURSORS OF THE THIOL-PROTECTED DERIVATIVES OF CAPTOPRIL AND ITS ANALOGUES

Regarding the significance of the precursors of the S-protected derivatives of captopril and its analogues, the S-protected precursor provides a novel and simple substrate for forming S-protected derivatives (thiol-protected captopril and its analogues) which undergo self-fractional crystallization upon their formation from that precursor.

In addition, it is noted that the amidization of the precursor with optically active amines other than proline or its analogues is useful for forming amides which also undergo self-fractional crystallization to separate diastereomeric amide species. More specifically, these resulting individual diastereomeric amides are separately hydrolyzable to the corresponding optically resolved amine species. Consequently, the precursor has utility for preparing individual diastereomeric amide species of which it is part.


Novel compositions of matter, therefore, pertain to a compound of the formula VI



wherein R_{10} is selected from the group consisting of lower alkoxy or hydroxy, and basic salts thereof wherein the other substituents are defined as above.

Further descriptions of this compound include the compound of the formula VI wherein: R_1 is lower alkyl wherein said lower alkyl is methyl, R_2 is S, n is 1, R_4 is a cyclic secondary amino of the formula I as defined above wherein R_5 is hydrogen and p is 3, and R_{10} is hydroxy.


Novel compositions of matter also pertain to a compound of the formula VII



and basic salts thereof wherein the substituent groups are defined above.


Further descriptions of this compound include the compound of the formula VII wherein: R_2 is S, R_4 is a cyclic secondary amino of formula I defined above wherein R_5 is hydrogen and p is 3, and R_{10} is hydroxy.

More specifically, novel compositions of matter pertinent to this invention comprise a compound of the formula VIII




and basic salts thereof wherein the substituent groups are defined above.

a) reacting the compound of the formula X



with a compound of the formula $\text{R}_8=\text{C}=\text{S}$ (XI) and a cyclic secondary amine to form the compound of the formula XII




wherein the substituents in the compounds are defined above;

b) converting said formed compound of the formula XII of step a) to an acylating agent by reacting it, for example, with the acid chloride of an inorganic acid;

c) reacting said formed acylating agent of step b) with a compound of the formula II to form said compound of the formula IX; and


d) isolating said compound of the formula IX.

A method for making a compound of the following formula XIII




and basic salts thereof wherein all of the substituents are defined above, comprises:

a) reacting the compound of the formula XIV



wherein L is a leaving group, with a compound of the formula XI and a cyclic secondary amine to form the compound of the formula XV




wherein the other substituents in the compounds are defined above;

b) converting said compound of the formula XV of step a) to an acylating agent by reacting it, for example, with the acid chloride of an inorganic acid;

c) reacting said formed acylating agent of step b) with a compound of the formula II to form said compound of the formula XIII; and


d) isolating said compound of the formula XIII.

A method for making a compound of the following formula XVI



and basic salts thereof, wherein R_{11} is imino, and the other substituents are defined above, comprises:

a) reacting the compound of the formula XIV and a compound of the formula $R_4C(S)NHR_6$ (XVII) to form the compound of the formula XVIII




wherein the substituents in the compounds are defined above;

b) converting said compound of the formula XVIII of step a) to an acylating agent by reacting it, for example, with the acid chloride of an inorganic acid;

c) reacting said formed acylating agent of step b) with a compound of the formula II to form said compound of the formula XVI; and


d) isolating said compound of the formula XVI.

A further method for making a compound of the following formula IX




and basic salts thereof wherein all of the substituent groups are defined above, comprises:

a) reacting the compound of the formula XIX



wherein R_{12} is lower alkoxy, with a compound of the formula XI and a cyclic secondary amine to form the compound of the formula XX



wherein the other substituents in the compounds are defined above;


b) hydrolyzing said compound of the formula XX whereby said R_{12} group is replaced by a hydroxyl group to form a compound of the formula XII;

c) converting said formed compound of the formula XII of step b) to an acylating agent by reacting it with, for example, the acid chloride of an inorganic acid;

d) reacting said formed acylating agent of step c) with a compound of the formula II to form said compound of the formula IX; and


e) isolating said compound of the formula IX.

Finally, an additional method for making a compound of the following formula XVI




and basic salts thereof wherein all of the substituents are defined above, comprises:

a) reacting the compound of the formula XXI



and a compound of the formula XVII to form the compound of the formula XXII



wherein the substituents in the compounds are defined above;

b) hydrolyzing said compound of the formula XXII whereby said R_{12} group is replaced by a hydroxyl group to form a compound of the formula XVIII;

c) converting said compound of the formula XVIII of step b) to an acylating agent by reacting it, for example, with the acid chloride of an inorganic acid;

d) reacting said formed acylating agent of step c) with a compound of the formula II to form said compound of the formula XVI; and

e) isolating said compound of the formula XVI.

With regard to the prior noted methods for preparing the compounds which comprise compound III, the methods also pertain to the preparation of compounds wherein the protecting group portion of the compound is variable as noted regarding the compositions of matter, whereas the alkanoyl and amino portions of the amide compound are limited respectively to 2-methyl-3-

thiopropionyl and to L-proline and substituted L-proline, including but not limited to the case wherein the substituted L-proline is 4S-(phenylthio)-L-proline. In addition, the methods are useful in the situation wherein the protecting group is limited to pyrrolidinocarbonyl or pyrrolidinothioxomethyl, and the thioalkylcarboxamide portion of the compound is limited as just noted.

Compound III may also be prepared by alternating the preparative sequences noted in prior methods. Here the initial step again necessitates condensing the carboxylic acid compounds of step (a) of the prior noted methods with an amine using known methods to produce an amide. For example, one such method involves reacting the carboxylic acid compounds of step (a) with the acid halide of an inorganic acid to convert the carboxylic acid to an acylating agent. However, the order of subsequent reaction steps can be changed. In these alternative methods, the formed acylating agent is then subjected to an amidization reaction as noted in the prior noted preparative methods. Once the amide is formed, the carboxylic acid portion of the amide undergoes reaction steps as noted in the prior discussed preparative methods whereby the S-functionalized thiol group is incorporated in the carboxylic acid portion of the amide to form compound III.

More specifically, as an example, a derivative and precursor of captopril were prepared as follows. The S-protected 2-methyl-3-thiopropionic acid corresponding to formula VII is formed where pyrrolidine, carbon disulfide and methacrylic acid are allowed to react -- preferably in equimolar amounts -- either with, or without a solvent present. Typically, the reactants are dissolved in a solvent, for example, 2-propanol or ethyl acetate, in the amounts of about 5 to about 40 weight percent. This solution is then either refluxed for about 1 to about 10

hours or kept at room temperature for from about 1 to about 5 days. The desired product either crystallizes spontaneously from solution or is induced to crystallize by the addition of a non-solvent. After chilling to about -5°C to about 15°C for about 1 to about 12 hours the S-protected 2-methyl-3-thiopropionic acid precursor is isolated by filtration or centrifugation, washed with a small amount of crystallization solvent and air dried or dried in an oven or vacuum oven. This material is pure enough to be used in the coupling reaction with L-proline or another nucleophile to form the S-protected 2-methyl-3-thiopropionic acid. For such a reaction the compound is dissolved in methylene chloride or another suitable solvent at a concentration of about 2 to about 15 weight percent and is treated with about 0.9 to about 1.1 equivalent of an inorganic acid chloride, typically thionyl chloride at about -5°C to about 25°C . After about 10 minutes to about 1 hour L-proline or another nucleophile is added followed by the addition of about 2 to about 5 equivalents of an organic base, for example, pyridine at a temperature of from about -5°C to about 25°C . After about 0.5 to about 2 hours, the solution is washed with aqueous acid to remove the base. The product is (i) crystallized directly from solution, (ii) induced to crystallize by addition of a non-solvent, (iii) collected by evaporating the reaction solvent followed by dissolving the product in a solvent from which it can be recrystallized, or (iv) crystallized by a combination of the above.


5.2.2 PREPARATIONS OF PRECURSORS OF THIOL-PROTECTED DERIVATIVES OF CAPTOPRIL AND ITS ANALOGUES

Described herein are also methods that relate to the production of a useful precursor for combining with an amine whereby the resulting amide undergoes self-fractional

crystallization. The reaction of the precursor with carboxyl-substituted heterocyclic amine compounds of the formula II and more specifically proline or substituted proline, whereby the protected thiol derivatives of captopril and its analogues are easily prepared, is of particular interest. In particular, the precursor is useful since it is incorporated with the S-functionalized substituent which facilitates the previously noted derivative's susceptibility to self-fractional crystallization upon its formation from that precursor.

In the following methods for preparation of the captopril precursors and their analogues, the designation of the substituents is the same as noted in the discussion of Section 5.1.1 and the discussion regarding genus formula except where noted. More limited descriptions of the compounds are also described therein.

A method for preparing a precursor which is useful for preparing derivatives of captopril and its analogues pertains to a method for making a compound of the formula XXIII



wherein the substituent groups are defined above. The method comprises:


- a) reacting the compound of the formula XXIV



with a compound of the formula XI and a cyclic secondary amine to form the compound of the formula XXIII wherein the substituent groups in the compounds are defined above; and

b) isolating said compound of the formula XXIII.

Another method for preparing a precursor which is useful for preparing derivatives of captopril and its analogues pertains to a method for making a compound of the formula XXV



wherein the substituents are defined above. The method comprises:


a) reacting the compound of the formula XXVI



with a compound of the formula XI and a cyclic secondary amine to form the compound of the formula XXV wherein the substituent groups in the compounds are defined above; and

b) isolating said compound of the formula XXV.

Another method for preparing a precursor which is useful for preparing derivatives of captopril and its analogues pertains to a method for making a compound of the formula XXVII



wherein the substituent groups are defined above. The method comprises:

a) reacting the compound of the formula XXVI and a compound of the formula XVII to form the compound of the formula XXVII wherein the substituent groups in the compound are defined above; and

b) isolating said compound of the formula XXVII.

With regard to the methods for preparing these precursor compounds, the methods also relate to the preparation of compounds wherein the protecting group portion of the compound is variable as noted whereas the thioalkanoyl portion of the compound is limited to 2-methyl-3-thiopropionyl. In addition, the methods are useful in the situation wherein the protecting group is limited to pyrrolidinocarbonyl or pyrrolidinethoxymethyl, and the thioalkanoyl portion of the compound is limited as noted.

5.2.3 PREPARATIONS OF CAPTOPRIL AND ITS ANALOGUES FROM THEIR THIOL-PROTECTED DERIVATIVES

The preparation of captopril and captopril analogues as the racemic mixture or chiral species is also described herein. The desired racemic species is obtained by affecting the removal of the thiol protecting group. Of particular significance regarding the described method is that the precursor can be separated as a pure diastereomer by its self-fractional crystallization, and subsequently treated to remove the thiol protecting group to obtain captopril or a captopril analogue. Specific details regarding the preparation of captopril and its analogues are exemplified by the following procedures used in preparing captopril.

The preparation of a mixture of diastereomers is accomplished starting from non-optically active raw materials that condense to form a racemic mixture of *S*-protected 2-methyl-3-thiopropionic acid. Such syntheses of *S*-protected captopril and its analogues are noted previously herein. This material is then coupled with *L*-proline to form *S*-protected captopril as a mixture of diastereomers. This can be conducted by two general processes. One involves isolation of the racemic acid precursor from one set of solvents (e.g. isopropanol/water) and subsequent coupling to *L*-proline in another solvent (e.g. methylene chloride). The other general approach conducts both reactions in one solvent (e.g., ethyl acetate) without precursor isolation.

When *L*-proline is coupled with the *S*-protected 2-methyl-3-thiopropionic acid, diastereomers of *S*-protected captopril are formed. A solution comprised of these diastereomers having (*S,S*) and (*R,S*) configurations is prepared in the desired crystallization solvent (e.g. ethyl acetate), at a concentration of about 6 to about 35%, by solvent exchange from methylene chloride or directly from the coupling reaction after base (pyridine) removal. Crystallization of the desired diastereomer is accomplished by reducing the temperature from about -20°C to about 10°C, and either adding non-solvent, removing solvent, or some combination thereof.


The (*S,S*) diastereomer will crystallize from the solution while the remaining mother liquor contains predominately the other diastereomer. The solid can be collected by filtration or centrifugation. After washing with said crystallizing solvent, the solid is further converted by de-protection to captopril while the filtrate is sent for proline recovery and recycling. The solvent-wet crystals, isolated from the above fractional

crystallization, represent 40 to 49% recovery of the (S,S) diastereomer of the protected captopril (or captopril analog).

Captopril and captopril analogs can be produced from their S-protected derivatives by direct hydrolysis of the S-protected derivative. The S-protected material is first dissolved in a concentrated aqueous solution of an alkali metal hydroxide (e.g., potassium or sodium hydroxide) -- preferably with the hydroxide at a concentration from about 5 wt % to about 25 wt % -- to yield a solution containing about 5 wt % to about 25 wt % of the S-protected captopril derivative. A preferred hydroxide solution is aqueous potassium hydroxide. This solution is then refluxed for from about 1 hour to about 24 hours, preferably under an inert gas (e.g., nitrogen) atmosphere. After cooling, the solution is acidified to a pH of about 0 to about 3, and then it is extracted with an organic solvent. Suitable organic solvents include methylene chloride, ethyl acetate, and chloroform. Finally, the solvent is removed or concentrated, and the desired (S, S) diastereomer of captopril or captopril analog is isolated by crystallization, chromatography, or yet another standard purification method.

The filtrate from the asymmetric crystallization step contains predominately the (R,S) diastereomer of the S-protected captopril. The crystallizing solvent is evaporated and replaced with a solution of a strong acid (e.g., hydrochloric acid). This mixture is heated to reflux for about 1 to about 10 hours to hydrolyze the amide bond. The proline is then recovered by conventional means and recycled as the free amino acid or its hydrochloride salt.

The preferred embodiment of this invention which is useful in preparing captopril and its analogues pertains to a method for the production of a compound of the formula XXVIII



and basic salts thereof, wherein the substituent groups are defined above, comprising:

- a) preparing a mixture of diastereomers of a compound of formula III wherein the diastereomeric mixture contains two stereoisomers which have the same configuration at C_b and opposite configurations at C_a ;
- b) preparing a solution of said mixture of said diastereomers in a solvent;
- c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution, said solution becoming enriched in the other diastereomer;
- d) separating said crystallized and resolved diastereomer from said resulting solution containing said other resolved diastereomer;
- e) removing the thiol protecting substituent of the formula XXIX



wherein the substituent groups are defined above, from either of said separated and resolved diastereomers, to form either resolved diastereomer of formula XXVIII; and


f) isolating either resolved diastereomer of formula XXVIII;

whereby either resolved diastereomer of the compound of formula XXVIII is obtained.

A further description of the compounds prepared by this method includes the compounds of the formula XXVII:

1. wherein R is hydroxy, R_1 is lower alkyl wherein said lower alkyl is methyl, R_3 is hydrogen, 4 \underline{S} -hydroxy or 4 \underline{S} -arylthio, n is 1, m is 3, C_a has an \underline{S} configuration and C_b has an \underline{L} configuration.

Another embodiment of this invention which is useful in preparing captopril and its analogues pertains to a method for the production of a compound of the formula XXX



and basic salts thereof, wherein the substituent groups are defined above, comprising:

a) preparing a mixture of diastereomers of a compound of formula IV wherein the diastereomeric mixture contains two stereoisomers which have opposite configurations at C_a ; and basic salts thereof;

b) preparing a solution of said mixture of said diastereomers in a solvent;

c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution, said solution becoming enriched in the other diastereomer;

d) separating said crystallized and resolved diastereomer from said resulting solution containing said other resolved diastereomer;

e) removing the thiol protecting substituent of the formula XXIX from either of said separated and resolved diastereomers, to form either resolved diastereomer of formula XXX, and

f) isolating either resolved diastereomer of formula XXX,

whereby either resolved diastereomer of the compound of formula XXX is obtained.


A further description of the compounds prepared by this method includes the compounds of the formula XXVII:

wherein R_7 is selected from the group consisting of L-proline and substituted L-proline wherein the substituted L-proline is selected from the group consisting of 4S-(phenylthio)-L-proline and 4S-hydroxy L-proline.

A further embodiment of this invention which is useful in preparing captopril and its analogues pertains to a method for the production of a compound of the formula XXX and basic salts thereof, wherein all of the substituent groups are defined above, comprising:

a) preparing a mixture of diastereomers of a compound of formula V wherein the diastereomeric mixture contains two stereoisomers which have opposite configurations at C_a ; and basic salts thereof;

- b) preparing a solution of said mixture of said diastereomers in a solvent;
- c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution, said solution becoming enriched in the other diastereomer;
- d) separating said crystallized and resolved diastereomer from said resulting solution containing said other resolved diastereomer;
- e) removing the thiol protecting substituent of the formula XXXI



from either of said separated and resolved diastereomers, to form either resolved diastereomer of formula XXX; and

- f) isolating either resolved diastereomer of formula XXX;

whereby either resolved diastereomer of the compound of formula XXX is obtained.

A further description of the compounds prepared by this method includes the compounds of the formula XXVII:

wherein R_7 is selected from the group consisting of L-proline and substituted L-proline wherein the substituted L-proline is selected from the group consisting of 4S-(phenylthio)-L-proline and 4S-hydroxy L-proline.

6.0 EXAMPLES

6.1 Preparation of 2-Methyl-3-(S-pyrrolidinothioxomethyl)thiopropionic Acid (XXXII)

8.6 g (0.1 m) of methacrylic acid was added to 50 ml of isopropyl alcohol, cooled on ice, and treated with stirring, with 7.1 g (8.3 ml, 0.1 m) of pyrrolidine. This mixture was, in turn, treated under the same conditions with 7.6 g (6.00 ml, 0.1 m) of carbon disulfide. The solution was then placed in an oil bath at 110°C and refluxed for 2.5 hours. The heat was removed and 50 ml of water was added. Upon cooling, the product crystallized as a white solid. This was collected by filtration, washed two times with cold 50/50 H₂O/isopropyl alcohol and dried to yield 17 g of material at 73% yield and with a melting point of 135-136°C.

6.2 Preparation of N-[2-Methyl-3-(S-pyrrolidinothioxomethyl)thiopropionyl] L-Proline (XXXIII) Using Two Reaction Solvents

A solution of the compound of formula XXXII (750 g, 3.21 m) was prepared in 4.0 l of dichloromethane and was chilled in an ice bath to 15°C. Thionyl chloride (382 g, 234.5 ml, 3.21 m) was run into this stirred solution over the course of 15 minutes. The slurry thus formed was stirred on ice for 1/2 hour, after which L-proline (370 g, 3.21 m) was added. The pyridine (763 g) was dripped in over a 45 minute period, the temperature being kept at less than 25°C by use of an ice bath. Upon complete addition of the pyridine, the solution was stirred at room temperature for 1 hour. The resulting solution was washed 2 times with dilute HCl to remove pyridine. The dichloromethane

solution was diluted with 4 liters of ethyl acetate and solvent was distilled from the flask. Water which was present was removed as the azeotrope with dichloromethane. Distillation was continued until the vapor temperature reached 65°C, during which time product began to crystallize. After cooling to room temperature overnight, the solid was isolated by filtration to give 420 gm of a white product with a rotation of -1.375° ($c=1/\text{ethanol}$). This material was recrystallized from 1.6 liters of 2-methoxyethanol to provide 350 g of pure product with a rotation of -1.443° ($c=1/\text{ethanol}$); the melting point of the material was 190-191°C.

6.3 Preparation of N-[2-Methyl-3-(S-pyrrolidino-thioxomethyl)thiopropionyl] L-Proline (XXXIII)
Using A Single Reaction Solvent

17.2 g (16.94 ml, 0.2 m) of methacrylic acid was added to 0.3 l of ethyl acetate and was stirred as pyrrolidine (16.6 ml, 0.2 m) was slowly added. Then to the stirred solution was slowly added carbon disulfide (12.0 ml, 0.2 m). The mix became warm as a solid began to precipitate. The mix was refluxed for 2.5 hours to give a pale yellow solution. The solution was cooled to 25° and then treated with thionyl chloride (23.8 g, 14.6 ml, 0.2 m) with stirring. After 15 minutes, the slurry was cooled in ice. With stirring, L-proline (23 g, 0.2 m) was added followed by the dropwise addition of 20 ml of pyridine. After 1/2 hour 0.1 l H₂O and 20 ml concentrated HCl were added, and the mixture was stirred 5 minutes and cooled on ice, this causing the product to precipitate as a white powder. The yield was 8.5 g (26%) and the rotation was -1.003° ($C=1/\text{ethanol}$). Further cooling of the filtrates gave an additional 8.7 g of the product which had a rotation of -0.982° ($C=1/\text{ethanol}$).

6.4 Preparation of Methyl Ester of N-[2-Methyl-3-(S-pyrrolidinothioxomethyl)thiopropionyl] L-Proline (XXXIV) Using Esterified Reactant

7.0 gm (0.03 m) of the protected mercapto-isobutyric acid from above (XXIX) was dissolved in 100 ml of methylene chloride, and was treated with thionyl chloride (3.6 g, 2.2 ml, 0.03 m). After 15 minutes the slurry was cooled to 0°C, and L-proline methyl ester hydrochloride (5.0 g, 0.03 m) was added with stirring. To this stirred mixture, pyridine (10 ml) was slowly dripped in with stirring. After 1 hour the pale yellow solution was diluted with 200 ml of ethyl acetate and this solution was extracted with water (2 x 100 ml). After drying over magnesium sulfate, the solvents were stripped to give an oil which solidified upon cooling. This solid was recrystallized from isopropyl alcohol to yield 3.0 g (29.0%) of white crystals. The melting point of the product was 124-125°C and its rotation was -1.526° (C=1/ethanol).

6.5 Preparation of Methyl Ester of N-[2-methyl-3-(S-pyrrolidinothioxomethyl)thiopropionyl] L-Proline (XXXIV) By Esterification Step

The resolved, protected captopril derivative (11.0 g, 0.033 m) from above (XXXIII) was dissolved in 250 ml of methanol. This solution was treated with gaseous hydrogen chloride and left overnight. The excess methanol was stripped and the residue was poured into water. The white solid thus formed was filtered, and washed well with water. The damp filter cake was recrystallized from isopropyl alcohol to yield 9.3 g (81.0%) of white crystals. The product's melting point was 124-125°C and its rotation was -1.493° (C=1/ethanol).

6.6 Preparation of the t-Butyl Ester of N-[2-Methyl-3-(S-pyrrolidinothioxomethyl)thiopropionyl] L-Proline (XXXV)

15.3 gm (0.46 m) of the protected captopril derivative from above (XXXIII) was mixed with 150 ml of methylene chloride and 1.0 ml of methanesulfonic acid. This was cooled in dry ice/acetone and 50 ml of liquid isobutylene was added. The bottle was stoppered and then shaken on a Parr apparatus for 48 hours. The resulting clear solution was cooled on dry ice/acetone before opening. The contents were poured into 1.0 N sodium hydroxide and the organic material which separated was taken into diethyl ether. After drying over magnesium sulfate, the solvents were stripped to give a white solid. This was dissolved in a small amount of methylene chloride. Hexane was added to the cloud point. Upon cooling and scratching, 13.5 g (75.9%) of the ester crystallized as a white powder. The rotation was -1.430° (C=1/ethanol) and its melting point was 128-129°C.

6.7 Preparation of Methyl Ester of N-[2-Methyl-3-(S-pyrrolidinocarbonyl)thiopropionyl] L-Proline (XXXVI)

3.44 g (0.01 m) of (XXXIV) was treated with dimethyl sulfate (1.26 g, 0.01 m, 0.95 ml) and was heated in an oil bath at 90°C for 1 hour. The clear product was dissolved in 25 ml of 4:1 H₂O/HCl and was stirred at room temperature for 15 minutes. This solution was saturated with NaCl to precipitate some starting material which was extracted into ethyl acetate. The aqueous layer was treated with concentrated NH₄OH to a pH of 11-12. After 5 minutes the precipitated oil was taken into ethyl acetate, dried and stripped to give 2.5 g of a clear oil. This was dissolved in 10 ml ethyl acetate and treated with hexane to

the cloud point. Upon cooling, a yield of 1.5 g of a white solid crystallized from the solution. The melting point of the product is 90-91°C and its rotation -1.323° (C=1/ethanol).

6.8 Preparation of N-[2-Methyl-3-(thiopropionyl)]
L-Proline (Captopril) (XXXVII)


A solution of potassium hydroxide (70.0 g, 1.25 m) in 250 ml of water was prepared, and de-oxygenated with nitrogen. This solution was stirred as 50 gm (0.15 m) of the resolved protected captopril compound (XXXVIII) ($\alpha = -1.428^\circ$ c=1/ethanol) was added. The resulting solution was kept under nitrogen as it was brought to reflux. After 2 hours, the solution was cooled to room temperature and then placed in an ice bath. Concentrated HCl (about 80 to about 85 ml) was dripped in with stirring causing the evolution of CO₂ and H₂S. When the solution became acidic (pH about 1 to about 2) a white precipitate formed. This precipitate was removed by filtration, and the filtrates were extracted four times with about 150 ml portions of methylene chloride. The extracts were combined, dried over magnesium sulfate and stripped to provide a clear oil in a yield of 31 gm. HPLC analysis showed the oil to be about 75% captopril wherein the overall chemical yield was 23.3 gm (71%).

A portion of the crude product was purified by chromatography on silica gel using ethyl acetate (94)/methanol (5)/acetic acid (1) as the eluent. The fractions containing captopril were pooled and evaporated to provide captopril as an oil. This oil was solidified by trituration in diethyl ether to provide purified captopril. Its rotation was -1.281° (c=1/ethanol).

The present invention is not intended to be limited in scope by the above examples concerning the preparation of captopril and captopril analogues, described herein, since each is intended merely as an illustration of the invention. In addition, any method which is functionally equivalent to those set forth herein is intended to be within the scope of the invention. Indeed various modifications of the invention, in addition to those shown and described herein, will become apparent to those skilled in the art from the foregoing description and accompanying specification. Such modifications are intended to fall within the scope of the appended claims.


I claim:

1. The method for preparing a compound of formula



wherein R is selected from the group consisting of hydroxy, amino and lower alkoxy; R_1 is selected from the group consisting of lower alkyl, lower alkoxy, hydroxy, halide, phenyl and substituted phenyl; R_3 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, arylthio, halide, phenyl and substituted phenyl; C_a has a R or S configuration; C_b has a D or L configuration; m is 2, 3 or 4; n is 1, 2 or 3; and basic salts thereof; comprising

- a) preparing a mixture of two diastereomers of a compound of formula



wherein the diastereomers have the same configuration at C_b and opposite configurations at C_a ; R is selected from the group consisting of hydroxy, amino and lower alkoxy; R_1 is selected from the group consisting of lower alkyl, lower alkoxy, hydroxy, halide, phenyl and substituted phenyl; R_2 is selected from the group consisting of O, S and imino; R_3 is


selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, arylthio, halide, phenyl and substituted phenyl; R_4 is a cyclic secondary amino; m is 2, 3 or 4; n is 1, 2 or 3; and basic salts thereof;

b) preparing a solution of said mixture of said diastereomers in a solvent;


c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution, said solution becoming enriched in the other diastereomer;

d) separating said crystallized and resolved diastereomer from said resulting solution containing said other resolved diastereomer;

e) removing the thiol protecting substituent of the formula




wherein R_2 is selected from the group consisting of O, S and imino, and R_4 is a cyclic secondary amino, from said separated crystallized and resolved diastereomer of step d) to form the resolved diastereomer of formula




and

f) isolating the resolved diastereomer of formula




whereby the resolved diastereomer of formula




is obtained.

2. A method for preparing a compound of formula,




wherein R is selected from the group consisting of hydroxy, amino and lower alkoxy; R_1 is selected from the group consisting of lower alkyl, lower alkoxy, hydroxy, halide, phenyl and substituted phenyl; R_3 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, arylthio, halide, phenyl and substituted phenyl; C_a has a R or S configuration; C_b has a D or L configuration; m is 2, 3 or 4; n is 1, 2 or 3; and basic salts thereof; comprising

a) preparing a mixture of two diastereomers of a compound of formula




wherein the diastereomers have the same configuration at C_b and opposite configurations at C_a ; R is selected from the group consisting of hydroxy, amino and lower alkoxy; R_1 is selected from the group consisting of lower alkyl, lower alkoxy, hydroxy, halide, phenyl and substituted phenyl; R_2 is selected from the group consisting of O, S and imino; R_3 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, arylthio, halide, phenyl and substituted phenyl; R_4 is a cyclic secondary amino; m is 2, 3 or 4; n is 1, 2 or 3; and basic salts thereof;

- b) preparing a solution of said mixture of said diastereomers in a solvent;
- c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution, said solution becoming enriched in the other diastereomer;
- d) separating said other resolved diastereomer from said resulting solution and said crystallized and resolved diastereomer;
- e) removing the thiol protecting substituent of the formula




wherein R_2 is selected from the group consisting of O, S and imino, and R_4 is a cyclic secondary amino, from said other separated and resolved diastereomer of step d) to form the resolved diastereomer of formula;




and

f) isolating the resolved diastereomer of formula




whereby the resolved diastereomer of formula




is obtained.

3. The method of claim 1 or 2, wherein said cyclic secondary amino of the compound of formula




has the formula




wherein R_5 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, halide, arylthio and substituted phenyl and p is 2, 3 or 4.

4. The method of claim 1 or 2, wherein said imino group of compound of formula




has the formula NR_6 wherein R_6 is selected from the group consisting of hydrogen, hydroxy, lower alkoxy, lower alkyl, phenyl and substituted phenyl.

5. The method of claims 1 or 2, wherein said phenyl substituent of compound of formula




is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy.

6. The method of claims 3, wherein said phenyl substituent of compound of formula




is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy.

7. The method of claims 4, wherein said phenyl substituent of compound of formula



is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy.

8. The method of claims 1 or 2, wherein said phenyl substituent of compound of formula



is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy.

9. The method of claim 1 or 2, wherein said solvent of step b) is ethyl acetate.


10. The method of claim 1 or 2, wherein said crystallization of step c) further comprises facilitating said crystallization by a method selected from the group consisting of reducing the temperature, adding non-solvent, removing solvent and some combination thereof.


11. The method of claim 10, wherein said reduction of the temperature is to about 10°C to about -20°C.

12. The method of claim 1 or 2, wherein said solution of step b) contains a concentration of said mixture of said diastereomers from about 6% to about 35%.

13. The method of claim 1 or 2, wherein said separation of step d) is effected by a method selected from the group consisting of filtration and centrifugation.


14. The method of claim 1, wherein said crystallized and resolved diastereomer of formula





bb) acidifying said alkali metal hydroxide solution of step aa) by the addition of an inorganic acid to said alkali metal hydroxide solution to form an acidified solution; and


cc) isolating said compound of formula




from the acidified solution of step bb).

17. The method of claim 2, wherein said thiol protecting substituent in step e) is removed by the process comprising:


aa) hydrolyzing said diastereomer of formula



with an alkali metal hydroxide solution to form the compound of formula



- bb) acidifying said alkali metal hydroxide solution of step aa) by the addition of an inorganic acid to said alkali metal hydroxide solution to form an acidified solution; and
- cc) isolating said compound of formula



from the acidified solution of step bb).

18. The method of claim 16 or 17, wherein said alkali metal hydroxide solution of step aa) consists of water and an alkali metal hydroxide which is selected from the group consisting of sodium hydroxide and potassium hydroxide.
19. The method of claim 16 or 17, wherein solution of step aa) has an alkali metal concentration from about 5 wt % to about 25 wt %.
20. The method of claim 16 or 17, wherein said hydrolyzing of step aa) occurs by refluxing the solution from about 1 hour to about 24 hours.
21. The method of claim 20, wherein said hydrolyzing of step aa) occurs under an inert gas.

22. The method of claim 16 or 17, wherein said inorganic acid of step bb) is selected from the group consisting of hydrochloric acid and sulfuric acid.


23. The method of claim 16 or 17, wherein said addition of said inorganic acid of step bb) takes place until said acidified solution has pH from about 0 to about 3.

24. The method of claim 16 or 17, wherein said isolation of step cc) proceeds by extraction by an organic solvent.

25. The method of claim 16 or 17, wherein said organic solvent of step cc) is selected from the group consisting of methylene chloride, ethyl acetate and chloroform.


15

26. The method of claim 16, wherein said diastereomer of formula



of step aa) has an S configuration at C_a and L configuration at C_b and said compound of formula

25




30


of step cc) has an S configuration at C_a and L configuration at C_b .

35

27. The method of claim 1 or 2, wherein in said compound of formula




R is hydroxy, R_1 is lower alkyl wherein said lower alkyl is methyl, R_2 is S, R_3 is hydrogen, 4S-hydroxy or 4S-arylthio, n is 1, m is 3, R_4 is a cyclic secondary amino of the formula




wherein R_5 is hydrogen and p is 3, and C_b has an L configuration.

28. The method of claim 1, wherein in said compound of formula



R is hydroxy, R_1 is lower alkyl wherein said lower alkyl is methyl, R_3 is hydrogen, 4S-hydroxy or 4S-arylthio, n is 1, m is 3, C_a has an S configuration, and C_b has an L configuration.

29. The method of claim 2, wherein in said compound of formula



R is hydroxy, R_1 is lower alkyl wherein said lower alkyl is methyl, R_3 is hydrogen, 4S-hydroxy or 4S-arylthio, n is 1, m is 3, C_a has an R configuration, and C_b has an L configuration.

30. The method of claims 1 or 2, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.

31. The method of claim 3, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.

32. The method of claim 4, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.


33. The method of claim 5, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.

34. The method of claim 6, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.

35. The method of claim 7, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.


36. The method of claim 8, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.

37. The method for preparing a compound of formula



wherein C_a has a R or S configuration; R₇ is selected from the group consisting of L-(proline) and substituted L-(proline) and basic salts thereof, comprising:

a) preparing a mixture of diastereomers of a compound of formula




wherein the diastereomers have opposite configurations at C_a; R₂ is selected from the group consisting of O, S and imino; R₄ is a cyclic secondary amino; R₇ is selected from the group consisting of L-(proline) and substituted L-(proline) and basic salt thereof;

b) preparing a solution of said mixture of said diastereomers in a solvent;


c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution, said solution becoming enriched in the other diastereomer;

d) separating said crystallized and resolved diastereomer from said resulting solution containing said other resolved diastereomer;

e) removing the thiol protecting substituent of the formula




wherein R_2 is selected from the group consisting of O, S and imino, and R_4 is a cyclic secondary amino, from said separated crystallized and resolved diastereomer of step d) to form the resolved diastereomer of formula




and

f) isolating the resolved diastereomer of formula




whereby the resolved diastereomer of formula




is obtained.

38. A method for preparing a compound of formula



wherein C_a has a R or S configuration; R_7 is selected from the group consisting of L-(proline) and substituted L-(proline) and basic salts thereof, comprising:

a) preparing a mixture of diastereomers of a compound of formula




wherein the diastereomers have opposite configurations at C_a ; R_2 is selected from the group consisting of O, S and imino; R_4 is a secondary amino; R_7 is selected from the group consisting of L-(proline) and substituted L-(proline) and basic salt thereof;

b) preparing a solution of said mixture of said diastereomers in a solvent;


c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution, said solution becoming enriched in the other diastereomer;

d) separating said other resolved diastereomer from said resulting solution and said crystallized and resolved diastereomer;

e) removing the thiol protecting substituent of the formula




wherein R_2 is selected from the group consisting of O, S and imino, and R_4 is a cyclic secondary amino from said other separated and resolved diastereomer of step d) to form the resolved diastereomer of formula




and

f) isolating the resolved diastereomer of formula




whereby the resolved diastereomer of formula




is obtained.

39. The method of claim 37 or 38, wherein said cyclic secondary amino of the compound of formula




has the formula




wherein R_5 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, phenoxy, hydroxy, thiol, alkylthio, halide, arylphenyl and substituted phenyl and p is 2, 3 or 4.

40. The method of claim 37 or 38, wherein said imino group of compound of formula




has the formula NR_6 wherein R_6 is selected from the group consisting of hydrogen, hydroxy, lower alkoxy, lower alkyl, phenyl and substituted phenyl.

41. The method of claims 37 or 38, wherein said phenyl substituent of compound of formula




is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy.

42. The method of claims 39, wherein said phenyl substituent of compound of formula




is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy.

43. The method of claims 40, wherein said phenyl substituent of compound of formula



is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy.

44. The method of claims 37 or 38, wherein said phenyl substituent of compound of formula



is selected from the group consisting of halide, lower alkyl, hydroxy and lower alkoxy.

45. The method of claim 37 or 38, wherein said solvent of step b) is ethyl acetate.


46. The method of claim 37 or 38, wherein said crystallization of step c) further comprises facilitating said crystallization by a method selected from the group consisting of reducing the temperature, adding non-solvent, removing solvent and some combination thereof.

47. The method of claim 46, wherein said reduction of the temperature is to about 10°C to about -20°C.

48. The method of claim 37 or 38, wherein said solution of step b) contains a concentration of said mixture of said diastereomers from about 6% to about 35%.


49. The method of claim 37 or 38, wherein said separation of step d) is effected by a method selected from the group consisting of filtration and centrifugation.

50. The method of claim 37, wherein said crystallized and resolved diastereomer of formula



of step d) has an S configuration at C_a.


51. The method of claim 38, wherein said other resolved diastereomer of formula



of step d) has an R configuration at C_a.


52. The method of claim 37, wherein said thiol protecting substituent in step e) is removed by the process comprising:

aa) hydrolyzing said diastereomer with an alkali metal hydroxide solution to form the compound of formula



bb) acidifying said alkali metal hydroxide solution of step aa) by the addition of an inorganic acid to said alkali metal hydroxide solution to form an acidified solution; and


cc) isolating said compound of formula



from the acidified solution of step bb).


53. The method of claim 38, wherein said thiol protecting substituent in step e) is removed by the process comprising:

aa) hydrolyzing said diastereomer with an alkali metal hydroxide solution to form the compound of formula



bb) acidifying said alkali metal hydroxide solution of step aa) by the addition of an inorganic acid to said alkali metal hydroxide solution to form an acidified solution; and

cc) isolating said compound of formula



from the acidified solution of step bb).

54. The method of claim 52 or 53, wherein said alkali metal hydroxide solution of step aa) consists of water and an alkali metal hydroxide which is selected from the group consisting of sodium hydroxide and potassium hydroxide.

55. The method of claim 52 or 53, wherein solution of step aa) has an alkali metal concentration from about 5 wt % to about 25 wt %.

56. The method of claim 52 or 53, wherein said hydrolyzing of step aa) occurs by refluxing the solution from about 1 hour to about 24 hours.

57. The method of claim 56, wherein said hydrolyzing of step aa) occurs under an inert gas.


58. The method of claim 52 or 53, wherein said inorganic acid of step bb) is selected from the group consisting of hydrochloric acid and sulfuric acid.

59. The method of claim 52 or 53, wherein said addition of said inorganic acid of step bb) takes place until said acidified solution has pH from about 0 to about 3.


60. The method of claim 52 or 53, wherein said isolation of step cc) proceeds by extraction by an organic solvent.

61. The method of claim 52 or 53, wherein said organic solvent of step cc) is selected from the group consisting of methylene chloride, ethyl acetate and chloroform.

62. The method of claim 52, wherein said diastereomer of formula




of step aa) has an S configuration at C_a and said compound of formula




of step c) has an S configuration at C_a.

63. The method of claim 37 or 38, wherein in said compound of formula




R₂ is S, and R₄ is a cyclic secondary amino of the formula




wherein R₅ is hydrogen and p is 3.

64. The method of claim 37, wherein said diastereomer of formula




has an S configuration at C_a.

65. The method of claim 38, wherein said diastereomer of formula




has an R configuration at C_a.

66. The method of claim 37 or 38, wherein said substituted L-proline is 4S-(phenylthio)-L-proline or 4S-hydroxy-L-proline.
67. The method of claim 63, wherein said substituted L-proline is 4S-(phenylthio)-L-proline or 4S-hydroxy-L-proline.
68. The method of claims 37 or 38, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.
69. The method of claim 39, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.
70. The method of claim 40, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.
71. The method of claim 41, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.
72. The method of claim 42, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.
73. The method of claim 43, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.
74. The method of claim 44, wherein said lower alkoxy and lower alkyl groups have up to seven carbon atoms.
75. The method for preparing a compound of formula



wherein C_a has an S or R configuration; R_7 is selected from the group consisting of L-(proline) and substituted L-(proline) and basic salt thereof, comprising:

a) preparing a mixture of diastereomers of a compound of formula




wherein the diastereomers have opposite configurations at C_a ; R_7 is selected from the group consisting of L-(proline) and substituted L-(proline); R_8 is selected from the group consisting of O and S; R_9 is pyrrolidino; and basic salts thereof;

b) preparing a solution of said mixture of said diastereomers in a solvent;


c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution, said solution becoming enriched in the other diastereomer;

d) separating said crystallized and resolved diastereomer from said resulting solution containing said other resolved diastereomer;

e) removing the thiol protecting substituent of the formula




wherein R_8 is selected from the group consisting of O and S, and R_9 is pyrrolidino, from said separated crystallized and resolved diastereomer of step d) to form the resolved diastereomer of formula




and

f) isolating the resolved diastereomer of formula




whereby the resolved diastereomer of formula




is obtained.

76. A method for preparing a compound of formula,



wherein C_a has an S or R configuration; R_7 is selected from the group consisting of L-(proline) and substituted L-(proline) and basic salt thereof, comprising:

a) preparing a mixture of diastereomers of a compound of formula




wherein the diastereomers have opposite configurations at C_a ; R_7 is selected from the group consisting of L-(proline) and substituted L-(proline); R_8 is selected from the group consisting of O and S; R_9 is pyrrolidino; and basic salts thereof;

b) preparing a solution of said mixture of said diastereomers in a solvent;


c) allowing predominantly one diastereomer of said mixture to crystallize from the resulting solution containing the other diastereomer;

d) separating said other resolved diastereomer from said resulting solution and said crystallized and resolved diastereomer;

e) removing the thiol protecting substituent of the formula




wherein R_8 is selected from the group consisting of O and S, and R_9 is pyrrolidino, from said other separated and resolved diastereomer of step d) to form the resolved diastereomer of formula




and

f) isolating the resolved diastereomer of formula



whereby the resolved diastereomer of formula



is obtained.

77. The method of claim 75 or 76, wherein said solvent of step b) is ethyl acetate.


78. The method of claim 75 or 76, wherein said crystallization of step c) further comprises facilitating said crystallization by a method selected from the group consisting of reducing the temperature, adding non-solvent, removing solvent and some combination thereof.

79. The method of claim 78, wherein said reduction of the temperature is to about 10°C to about -20°C.

80. The method of claim 75 or 76, wherein said solution of step b) contains a concentration of said mixture of said diastereomers from about 6% to about 35%.


81. The method of claim 75 or 76, wherein said separation of step d) is effected by a method selected from the group consisting of filtration and centrifugation.

82. The method of claim 75, wherein said crystallized and resolved diastereomer of formula



of step d) has an S configuration at C_a.


83. The method of claim 76, wherein said crystallized and resolved diastereomer of formula



of step d) has an R configuration at C_a.


84. The method of claim 75, wherein said thiol protecting substituent of step e) is removed by the process comprising:

aa) hydrolyzing said diastereomer with an alkali metal hydroxide solution to form the compound of formula



bb) acidifying said alkali solution of step aa) by the addition of an inorganic acid to said alkali metal hydroxide solution to form an acidified solution; and


cc) isolating said compound of formula



from the acidified solution of step bb).


85. The method of claim 76, wherein said thiol protecting substituent of step e) is removed by the process comprising:

aa) hydrolyzing said diastereomer with an alkali metal hydroxide solution to form the compound of formula



bb) acidifying said alkali solution of step aa) by the addition of an inorganic acid to said alkali metal hydroxide solution to form an acidified solution; and

cc) isolating said compound of formula



from the acidified solution of step bb).

86. The method of claim 84 or 85, wherein said alkali metal hydroxide solution of step aa) consists of water and an alkali metal hydroxide which is selected from the group consisting of sodium hydroxide and potassium hydroxide.

87. The method of claim 84 or 85, wherein solution of step aa) has an alkali metal concentration from about 5 wt % to about 25 wt %.

88. The method of claim 84 or 85, wherein said hydrolyzing of step aa) occurs by refluxing the solution from about 1 hour to about 24 hours.

89. The method of claim 88, wherein said hydrolyzing of step aa) occurs under an inert gas.


90. The method of claim 84 or 85, wherein said inorganic acid of step bb) is selected from the group consisting of hydrochloric acid and sulfuric acid.

91. The method of claim 84 or 85, wherein said addition of said inorganic acid of step bb) takes place until said acidified solution has pH from about 0 to about 3.


92. The method of claim 84 or 85, wherein said isolation of step cc) proceeds by extraction by an organic solvent.

93. The method of claim 84 or 85, wherein said organic solvent of step cc) is selected from the group consisting of methylene chloride, ethyl acetate and chloroform.

94. The method of claim 84, wherein said diastereomer of formula




of step aa) has an S configuration at C_a and said compound of formula




of step cc) has an S configuration at C_a.

95. The method of claim 75, wherein said diastereomer of formula




has an S configuration at C_a.

96. The method of claim 76, wherein said diastereomer of formula



has an R configuration at C_a.

97. The method of claim 75 or 76, wherein in said compound of formula



R_8 is S.

98. The method of claim 75 or 76, wherein said substituted L-proline is 4S-(phenylthio)-L-proline or 4S-hydroxy-L-proline.

99. The method of claim 97, wherein said substituted L-proline is 4S-(phenylthio)-L-proline or 4S-hydroxy-L-proline.





FIG. 1

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US89/03960

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC(4): C07D 205/04; C07D 207/16; C07D 211/34 U.S.C.I.: 546/226; 548/533; 548/537; 548/953		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S.	546/226 548/533	548/537 548/953
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X,Y	US, A, 4,105,776 (ONDETTI ET AL) 8 August 1978; Note: column 8, lines 52 to 67.	1 to 99
X,Y	US A 4,332,725 (FISCHER ET AL) 1 June 1982; Note: column 1, line 52 to column 2 line 11.	1 to 99
X,Y	US, A 4,506,082 (CROSSLEY) 19 March 1985; Note: examples.	1 to 99
A	US A 4,297,282 (OHASHI ET AL) 27 October 1981. See entire document.	1 to 99
A	US A 4 261,895 (WISKOTT) 14 April 1981. See entire document.	1 to 99
A	US A 4 154,934 (BERNSTEIN ET AL) 15 May 1979. See entire document.	1 to 99
A	US, A 4,046,889 (ONDETTI ET AL) 6 September 1977. See entire document.	1 to 99
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
21 NOVEMBER 1989	<div style="font-size: 2em; font-weight: bold; margin: 0;">15 DEC 1989</div>	
International Searching Authority	Signature of Authorized Officer	
ISA/US	 JOSEPH P. BRUST	