

(43) International Publication Date
21 January 2016 (21.01.2016)(51) International Patent Classification:
A61K 39/395 (2006.01) *C07K 16/28* (2006.01)(74) Agents: **ZHOU, Jie** et al.; Morrison & Foerster LLP, 425 Market Street, San Francisco, CA 94105-2482 (US).(21) International Application Number:
PCT/US2015/040582(22) International Filing Date:
15 July 2015 (15.07.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/024,988 15 July 2014 (15.07.2014) US(71) Applicant (for all designated States except AL, AT, BA, BE, BG, CH, CN, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IN, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR): **GENENTECH, INC.** [US/US]; 1 Dna Way, South San Francisco, CA 94080-4990 (US).(71) Applicant (for AL, AT, BA, BE, BG, CH, CN, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IN, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR only): **F.HOFFMAN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).(72) Inventor: **JUNTTILA, Melissa**; C/o Genentech, Inc., 1 Dna Way, South San Francisco, CA 94080 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: COMPOSITIONS FOR TREATING CANCER USING PD-1 AXIS BINDING ANTAGONISTS AND MEK INHIBITORS




FIG. 1

(57) Abstract: The present invention describes combination treatment comprising a PD-1 axis binding antagonist and a MEK inhibitor and methods for use thereof, including methods of treating conditions where enhanced immunogenicity is desired such as increasing tumor immunogenicity for the treatment of cancer.

COMPOSITIONS FOR TREATING CANCER USING PD-1 AXIS BINDING ANTAGONISTS AND MEK INHIBITORS

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority benefit of U.S. provisional application Serial No. 62/024,988, filed July 15, 2014, the contents of which are incorporated herein by reference in its entirety.

SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

[0002] The content of the following submission on ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing (file name: 146392027540SeqList.txt, date recorded: July 8, 2015, size: 22 KB).

BACKGROUND

[0003] The provision of two distinct signals to T-cells is a widely accepted model for lymphocyte activation of resting T lymphocytes by antigen-presenting cells (APCs). Lafferty et al, Aust. J. Exp. Biol. Med. ScL 53: 27-42 (1975). This model further provides for the discrimination of self from non-self and immune tolerance. Bretscher et al, Science 169: 1042-1049 (1970); Bretscher, P.A., P.N.A.S. USA 96: 185-190 (1999); Jenkins et al, J. Exp. Med. 165: 302-319 (1987). The primary signal, or antigen specific signal, is transduced through the T- cell receptor (TCR) following recognition of foreign antigen peptide presented in the context of the major histocompatibility-complex (MHC). The second or co-stimulatory signal is delivered to T-cells by co-stimulatory molecules expressed on antigen-presenting cells (APCs), and induces T-cells to promote clonal expansion, cytokine secretion and effector function. Lenschow et al., Ann. Rev. Immunol. 14:233 (1996). In the absence of co-stimulation, T-cells can become refractory to antigen stimulation, do not mount an effective immune response, and further may result in exhaustion or tolerance to foreign antigens.

[0004] In the two-signal model T-cells receive both positive and negative secondary co-stimulatory signals. The regulation of such positive and negative signals is critical to maximize the host's protective immune responses, while maintaining immune tolerance and preventing autoimmunity. Negative secondary signals seem necessary for induction of T-cell tolerance, while positive signals promote T-cell activation. While the simple two-signal model still

provides a valid explanation for naive lymphocytes, a host's immune response is a dynamic process, and co-stimulatory signals can also be provided to antigen-exposed T-cells. The mechanism of co-stimulation is of therapeutic interest because the manipulation of co-stimulatory signals has shown to provide a means to either enhance or terminate cell-based immune response. Recently, it has been discovered that T cell dysfunction or anergy occurs concurrently with an induced and sustained expression of the inhibitory receptor, programmed death 1 polypeptide (PD-1). As a result, therapeutic targeting of PD-1 and other molecules which signal through interactions with PD-1, such as programmed death ligand 1 (PD-L1) and programmed death ligand 2 (PD-L2) are an area of intense interest.

[0005] PD-L1 is overexpressed in many cancers and is often associated with poor prognosis (Okazaki T et al., *Intern. Immun.* 2007 19(7):813) (Thompson RH et al., *Cancer Res* 2006, 66(7):3381). Interestingly, the majority of tumor infiltrating T lymphocytes predominantly express PD-1, in contrast to T lymphocytes in normal tissues and peripheral blood T lymphocytes indicating that up-regulation of PD-1 on tumor-reactive T cells can contribute to impaired antitumor immune responses (*Blood* 2009 114(8):1537). This may be due to exploitation of PD-L1 signaling mediated by PD-L1 expressing tumor cells interacting with PD-1 expressing T cells to result in attenuation of T cell activation and evasion of immune surveillance (Sharpe et al., *Nat Rev* 2002) (Keir ME et al., 2008 *Annu. Rev. Immunol.* 26:677). Therefore, inhibition of the PD-L1/PD-1 interaction may enhance CD8+ T cell-mediated killing of tumors.

[0006] The inhibition of PD-1 axis signaling through its direct ligands (e.g., PD-L1, PD-L2) has been proposed as a means to enhance T cell immunity for the treatment of cancer (e.g., tumor immunity). Moreover, similar enhancements to T cell immunity have been observed by inhibiting the binding of PD-L1 to the binding partner B7-1. Furthermore, combining inhibition of PD-1 signaling with other signaling pathways (e.g. MAPK pathway, “MEK”) that are deregulated in tumor cells may further enhance treatment efficacy. However, an optimal therapeutic treatment would combine blockade of PD-1 receptor/ligand interaction with an agent that directly inhibited tumor growth, optionally further including unique immune enhancing properties not provided by PD-1 blockade alone. There remains a need for such an optimal therapy for treating, stabilizing, preventing, and/or delaying development of various cancers.

[0007] All references, publications, and patent applications disclosed herein are hereby incorporated by reference in their entirety.

BRIEF SUMMARY

[0008] Provided herein are methods for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor, wherein the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist. In some embodiments, the method further comprises diagnosing the individual as having a cancer that is resistant to a B-raf antagonist, wherein the diagnosing occurs prior to administering the effective amount of the PD-1 axis binding antagonist and the MEK inhibitor. In some embodiments, the method further comprises selecting an individual for treatment based on the individual having cancer that is resistant to a B-raf antagonist or assessing that the individual is at risk of developing cancer that is resistant to a B-raf antagonist, wherein the selecting occurs prior to administering the effective amount of the PD-1 axis binding antagonist and the MEK inhibitor. In some embodiments, the individual has not been previously treated with a B-raf antagonist. In some embodiments, the individual has been previously treated with a B-raf antagonist.

[0009] In another aspect, provided herein are methods for treating or delaying progression of cancer in an individual comprising (a) diagnosing the individual as having a cancer that is resistant to a B-raf antagonist; and (b) administering to the individual an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor, wherein the administering occurs after diagnosing the individual. In some embodiments, the individual has not been previously treated with a B-raf antagonist. In some embodiments, the individual has been previously treated with a B-raf antagonist.

[0010] In another aspect, provided herein are methods for treating or delaying progression of cancer in an individual comprising (a) selecting an individual for treatment based on the individual having cancer that is resistant to a B-raf antagonist or assessing that the individual is at risk of developing cancer that is resistant to a B-raf antagonist; and (b) administering to the individual an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor, wherein the administering occurs after selecting the individual. In some embodiments, the individual has not been previously treated with a B-raf antagonist. In some embodiments, the individual has been previously treated with a B-raf antagonist.

[0011] In another aspect, provided herein are methods for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a PD-1

axis binding antagonist and a MEK inhibitor, wherein the individual has been previously treated with a B-raf antagonist for cancer.

[0012] In some embodiments, the cancer in the individual has progressed within 1 month, 6 months, 1 year, or 5 years after completing a B-raf antagonist-based therapy regimen. In some embodiments, the B-raf antagonist is a small molecule inhibitor, an antibody, a peptide, a peptidomimetic, an aptamer or a polynucleotide. In some embodiments, the B-raf antagonist is dabrafenib, vemurafenib, GSK 2118436, RAF265, XL281, ARQ736, BAY73-4506, sorafenib, PLX4720, PLX-3603, GSK2118436, GDC-0879, or N-(3-(5-(4-chlorophenyl)-1H-pyrrolo[2,3-b]pyridine-3-carbonyl)-2,4-difluorophenyl)propane-1-sulfonamide. In some embodiments, the B-raf antagonist is a selective B-raf antagonist of B-raf V600. In some embodiments, the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600E. In some embodiments, the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600E, B-raf V600K, and/or V600D. In some embodiments, the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600R.

[0013] In some embodiments, the cancer contains a BRAF V600E mutation, a BRAF wildtype, a KRAS wildtype, or an activating KRAS mutation. In some embodiments, the treatment results in a sustained response in the individual after cessation of the treatment. In some embodiments, the individual has colorectal cancer, melanoma, lung cancer, ovarian cancer, breast cancer, pancreatic cancer, hematological malignancy, bladder cancer, and/or renal cell carcinoma. In some embodiments, the cancer is metastatic.

[0014] In some embodiments, the PD-1 axis binding antagonist is selected from the group consisting of a PD-1 binding antagonist, a PD-L1 binding antagonist and a PD-L2 binding antagonist. In some embodiments, the PD-1 axis binding antagonist is a PD-1 binding antagonist. In some embodiments, the PD-1 binding antagonist inhibits the binding of PD-1 to its ligand binding partners. In some embodiments, the PD-1 binding antagonist inhibits the binding of PD-1 to PD-L1, PD-1 to PD-L2, or PD-1 to both PD-L1 and PD-L2. In some embodiments, the PD-1 binding antagonist is an antibody. In some embodiments, the PD-1 binding antagonist is MDX-1106, Merck 3745, CT-011, MEDI-0680, PDR001, REGN2810, BGB-108, BGB-A317, or AMP-224. In some embodiments, the PD-1 binding antagonist is nivolumab, pembrolizumab, pidilizumab, MEDI-0680, PDR001, REGN2810, BGB-108, BGB-A317, or AMP-224. In some embodiments, the PD-1 axis binding antagonist is a PD-L1 binding antagonist. In some embodiments, the PD-L1 binding antagonist inhibits the binding of

PD-L1 to PD-1, PD-L1 to B7-1, or PD-L1 to both PD-1 and B7-1. In some embodiments, the PD-L1 binding antagonist is an anti-PD-L1 antibody. In some embodiments, the anti-PD-L1 antibody is a monoclonal antibody. In some embodiments, the anti-PD-L1 antibody is an antibody fragment selected from the group consisting of Fab, Fab'-SH, Fv, scFv, and (Fab')₂ fragments. In some embodiments, the anti-PD-L1 antibody is a humanized antibody or a human antibody. In some embodiments, the PD-L1 binding antagonist is selected from the group consisting of: YW243.55.S70, MPDL3280A, MEDI4736, MDX-1105, and MSB0010718C. In some embodiments, the PD-L1 binding antagonist is selected from the group consisting of: YW243.55.S70, atezolizumab, durvalumab, MDX-1105, and avelumab. In some embodiments, the antibody comprises a heavy chain comprising HVR-H1 sequence of SEQ ID NO:15, HVR-H2 sequence of SEQ ID NO:16, and HVR-H3 sequence of SEQ ID NO:3; and a light chain comprising HVR-L1 sequence of SEQ ID NO:17, HVR-L2 sequence of SEQ ID NO:18, and HVR-L3 sequence of SEQ ID NO:19. In some embodiments, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:24 or 28 and a light chain variable region comprising the amino acid sequence of SEQ ID NO:21. In some embodiments, the PD-1 axis binding antagonist is a PD-L2 binding antagonist. In some embodiments, the PD-L2 binding antagonist is an immunoadhesin. In some embodiments, the MEK inhibitor is a competitive inhibitor of MEK. In some embodiments, the MEK inhibitor is more selective against an activating KRAS mutation. In some embodiments, the MEK inhibitor is an allosteric inhibitor of MEK. In some embodiments, the MEK inhibitor is more selective against an activating BRAF mutation. In some embodiments, the MEK inhibitor is a compound of the formula (I), (II), (III), (IV), (V), (VI) or (VII), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the MEK inhibitor is selected from the group consisting of G02442104, G-38963, G02443714, G00039805 and GDC-0973, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the MEK inhibitor is G02443714, G02442104 or G00039805.

[0015] In some embodiments, the MEK inhibitor is administered continuously. In some embodiments, the MEK inhibitor is administered intermittently. In some embodiments, the MEK inhibitor is administered before the PD-1 axis binding antagonist. In some embodiments, the MEK inhibitor is administered simultaneous with the PD-1 axis binding antagonist. In some embodiments, the MEK inhibitor is administered after the PD-1 axis binding antagonist. In

some embodiments, the PD-1 axis binding antagonist and/or the MEK inhibitor is administered intravenously, intramuscularly, subcutaneously, topically, orally, transdermally, intraperitoneally, intraorbitally, by implantation, by inhalation, intrathecally, intraventricularly, or intranasally.

[0016] In another aspect, provided herein are kits comprising a PD-1 axis binding antagonist and a package insert comprising instructions for using the PD-1 axis binding antagonist in combination with a MEK inhibitor to treat or delay progression of cancer in an individual, wherein the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist. In another aspect, provided herein are kits comprising a PD-1 axis binding antagonist and a MEK inhibitor, and a package insert comprising instructions for using the PD-1 axis binding antagonist and the MEK inhibitor to treat or delay progression of cancer in an individual, wherein the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist. In another aspect, provided herein are kits comprising a MEK inhibitor and a package insert comprising instructions for using the MEK inhibitor in combination with a PD-1 axis binding antagonist to treat or delay progression of cancer in an individual, wherein the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist. In another aspect, provided herein are kits comprising a PD-1 axis binding antagonist and a package insert comprising instructions for using the PD-1 axis binding antagonist in combination with a MEK inhibitor to treat or delay progression of cancer in an individual, wherein the individual has been previously treated with a B-raf antagonist for cancer. In another aspect, provided herein are kits comprising a PD-1 axis binding antagonist and a MEK inhibitor, and a package insert comprising instructions for using the PD-1 axis binding antagonist and the MEK inhibitor to treat or delay progression of cancer in an individual, wherein the individual has been previously treated with a B-raf antagonist for cancer. In another aspect, provided herein are kits comprising a MEK inhibitor and a package insert comprising instructions for using the MEK inhibitor in combination with a PD-1 axis binding antagonist to treat or delay progression of cancer in an individual, wherein the individual has been previously treated with a B-raf antagonist for cancer.

[0017] In some embodiments, the individual is a human.

[0018] It is to be understood that one, some, or all of the properties of the various embodiments described herein may be combined to form other embodiments of the present invention. These and other aspects of the invention will become apparent to one of skill in the

art. These and other embodiments of the invention are further described by the detailed description that follows.

BRIEF DESCRIPTION OF THE DRAWINGS

[0019] **FIG. 1** shows tumor re-growth upon treatment with anti-PDL1, a MEK inhibitor, or both. The graph shows the percent tumor volume change over time during treatment with the indicated agent(s).

[0020] **FIG. 2** shows individual animal responses to treatment with Vemurafenib followed by anti-PDL1, a MEK inhibitor, or both. Each bar depicts the percent change in tumor growth upon crossover from Vemurafenib to the indicated treatment in an individual animal.

DETAILED DESCRIPTION

I. General techniques

[0021] The techniques and procedures described or referenced herein are generally well understood and commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized methodologies described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* 3d edition (2001) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; *Current Protocols in Molecular Biology* (F.M. Ausubel, et al. eds., (2003)); the series *Methods in Enzymology* (Academic Press, Inc.): *PCR 2: A Practical Approach* (M.J. MacPherson, B.D. Hames and G.R. Taylor eds. (1995)), Harlow and Lane, eds. (1988) *Antibodies, A Laboratory Manual*, and *Animal Cell Culture* (R.I. Freshney, ed. (1987)); *Oligonucleotide Synthesis* (M.J. Gait, ed., 1984); *Methods in Molecular Biology*, Humana Press; *Cell Biology: A Laboratory Notebook* (J.E. Cellis, ed., 1998) Academic Press; *Animal Cell Culture* (R.I. Freshney), ed., 1987); *Introduction to Cell and Tissue Culture* (J.P. Mather and P.E. Roberts, 1998) Plenum Press; *Cell and Tissue Culture: Laboratory Procedures* (A. Doyle, J.B. Griffiths, and D.G. Newell, eds., 1993-8) J. Wiley and Sons; *Handbook of Experimental Immunology* (D.M. Weir and C.C. Blackwell, eds.); *Gene Transfer Vectors for Mammalian Cells* (J.M. Miller and M.P. Calos, eds., 1987); *PCR: The Polymerase Chain Reaction*, (Mullis et al., eds., 1994); *Current Protocols in Immunology* (J.E. Coligan et al., eds., 1991); *Short Protocols in Molecular Biology* (Wiley and Sons, 1999); *Immunobiology* (C.A. Janeway and P. Travers, 1997); *Antibodies* (P. Finch, 1997); *Antibodies: A Practical Approach* (D. Catty., ed., IRL Press, 1988-1989); *Monoclonal Antibodies: A Practical Approach* (P. Shepherd and C.

Dean, eds., Oxford University Press, 2000); *Using Antibodies: A Laboratory Manual* (E. Harlow and D. Lane (Cold Spring Harbor Laboratory Press, 1999); *The Antibodies* (M. Zanetti and J. D. Capra, eds., Harwood Academic Publishers, 1995); and *Cancer: Principles and Practice of Oncology* (V.T. DeVita et al., eds., J.B. Lippincott Company, 1993).

II. Definitions

[0022] The term "*antagonist*" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native polypeptide disclosed herein. In a similar manner, the term "*agonist*" is used in the broadest sense and includes any molecule that mimics a biological activity of a native polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native polypeptides, peptides, antisense oligonucleotides, small organic molecules, etc. Methods for identifying agonists or antagonists of a polypeptide may comprise contacting a polypeptide with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the polypeptide.

[0023] The term "*aptamer*" refers to a nucleic acid molecule that is capable of binding to a target molecule, such as a polypeptide. For example, an aptamer of the invention can specifically bind to a B-raf polypeptide, or to a molecule in a signaling pathway that modulates the expression or activity of B-raf. The generation and therapeutic use of aptamers are well established in the art. See, e.g., U.S. Pat. No. 5,475,096, and the therapeutic efficacy of Macugen® (Eyetech, New York) for treating age-related macular degeneration.

[0024] The term "*PD-1 axis binding antagonist*" is a molecule that inhibits the interaction of a PD-1 axis binding partner with either one or more of its binding partner, so as to remove T-cell dysfunction resulting from signaling on the PD-1 signaling axis – with a result being to restore or enhance T-cell function (e.g., proliferation, cytokine production, target cell killing). As used herein, a PD-1 axis binding antagonist includes a PD-1 binding antagonist, a PD-L1 binding antagonist and a PD-L2 binding antagonist.

[0025] The term "*PD-1 binding antagonists*" is a molecule that decreases, blocks, inhibits, abrogates or interferes with signal transduction resulting from the interaction of PD-1 with one or more of its binding partners, such as PD-L1, PD-L2. In some embodiments, the PD-1 binding antagonist is a molecule that inhibits the binding of PD-1 to its binding partners. In a specific

aspect, the PD-1 binding antagonist inhibits the binding of PD-1 to PD-L1 and/or PD-L2. For example, PD-1 binding antagonists include anti-PD-1 antibodies, antigen binding fragments thereof, immunoadhesins, fusion proteins, oligopeptides and other molecules that decrease, block, inhibit, abrogate or interfere with signal transduction resulting from the interaction of PD-1 with PD-L1 and/or PD-L2. In one embodiment, a PD-1 binding antagonist reduces the negative co-stimulatory signal mediated by or through cell surface proteins expressed on T lymphocytes mediated signaling through PD-1 so as render a dysfunctional T-cell less dysfunctional (e.g., enhancing effector responses to antigen recognition). In some embodiments, the PD-1 binding antagonist is an anti-PD-1 antibody. In a specific aspect, a PD-1 binding antagonist is MDX-1106 described herein. In another specific aspect, a PD-1 binding antagonist is Merck 3745 described herein. In another specific aspect, a PD-1 binding antagonist is CT-011 described herein.

[0026] The term “*PD-L1 binding antagonists*” is a molecule that decreases, blocks, inhibits, abrogates or interferes with signal transduction resulting from the interaction of PD-L1 with either one or more of its binding partners, such as PD-1, B7-1. In some embodiments, a PD-L1 binding antagonist is a molecule that inhibits the binding of PD-L1 to its binding partners. In a specific aspect, the PD-L1 binding antagonist inhibits binding of PD-L1 to PD-1 and/or B7-1. In some embodiments, the PD-L1 binding antagonists include anti-PD-L1 antibodies, antigen binding fragments thereof, immunoadhesins, fusion proteins, oligopeptides and other molecules that decrease, block, inhibit, abrogate or interfere with signal transduction resulting from the interaction of PD-L1 with one or more of its binding partners, such as PD-1, B7-1. In one embodiment, a PD-L1 binding antagonist reduces the negative co-stimulatory signal mediated by or through cell surface proteins expressed on T lymphocytes mediated signaling through PD-L1 so as to render a dysfunctional T-cell less dysfunctional (e.g., enhancing effector responses to antigen recognition). In some embodiments, a PD-L1 binding antagonist is an anti-PD-L1 antibody. In a specific aspect, an anti-PD-L1 antibody is YW243.55.S70 described herein. In another specific aspect, an anti-PD-L1 antibody is MDX-1105 described herein. In still another specific aspect, an anti-PD-L1 antibody is MPDL3280A (atezolizumab) described herein.

[0027] The term “*PD-L2 binding antagonists*” is a molecule that decreases, blocks, inhibits, abrogates or interferes with signal transduction resulting from the interaction of PD-L2 with either one or more of its binding partners, such as PD-1. In some embodiments, a PD-L2 binding antagonist is a molecule that inhibits the binding of PD-L2 to its binding partners. In a

specific aspect, the PD-L2 binding antagonist inhibits binding of PD-L2 to PD-1. In some embodiments, the PD-L2 antagonists include anti-PD-L2 antibodies, antigen binding fragments thereof, immunoadhesins, fusion proteins, oligopeptides and other molecules that decrease, block, inhibit, abrogate or interfere with signal transduction resulting from the interaction of PD-L2 with either one or more of its binding partners, such as PD-1. In one embodiment, a PD-L2 binding antagonist reduces the negative co-stimulatory signal mediated by or through cell surface proteins expressed on T lymphocytes mediated signaling through PD-L2 so as render a dysfunctional T-cell less dysfunctional (e.g., enhancing effector responses to antigen recognition). In some embodiments, a PD-L2 binding antagonist is an immunoadhesin.

[0028] The term “*dysfunction*” in the context of immune dysfunction, refers to a state of reduced immune responsiveness to antigenic stimulation. The term includes the common elements of both *exhaustion* and/or *anergy* in which antigen recognition may occur, but the ensuing immune response is ineffective to control infection or tumor growth.

[0029] The term “*dysfunctional*”, as used herein, also includes refractory or unresponsive to antigen recognition, specifically, impaired capacity to translate antigen recognition into downstream T-cell effector functions, such as proliferation, cytokine production (e.g., IL-2) and/or target cell killing.

[0030] The term “*anergy*” refers to the state of unresponsiveness to antigen stimulation resulting from incomplete or insufficient signals delivered through the T-cell receptor (e.g. increase in intracellular Ca^{+2} in the absence of ras-activation). T cell anergy can also result upon stimulation with antigen in the absence of co-stimulation, resulting in the cell becoming refractory to subsequent activation by the antigen even in the context of costimulation. The unresponsive state can often be overridden by the presence of Interleukin-2. Anergic T-cells do not undergo clonal expansion and/or acquire effector functions.

[0031] The term “*exhaustion*” refers to T cell exhaustion as a state of T cell dysfunction that arises from sustained TCR signaling that occurs during many chronic infections and cancer. It is distinguished from anergy in that it arises not through incomplete or deficient signaling, but from sustained signaling. It is defined by poor effector function, sustained expression of inhibitory receptors and a transcriptional state distinct from that of functional effector or memory T cells. Exhaustion prevents optimal control of infection and tumors. Exhaustion can result from both extrinsic negative regulatory pathways (e.g., immunoregulatory cytokines) as well as cell intrinsic negative regulatory (costimulatory) pathways (PD-1, B7-H3, B7-H4, etc.).

[0032] “*Enhancing T-cell function*” means to induce, cause or stimulate a T-cell to have a sustained or amplified biological function, or renew or reactivate exhausted or inactive T-cells. Examples of enhancing T-cell function include: increased secretion of γ -interferon from CD8⁺ T-cells, increased proliferation, increased antigen responsiveness (e.g., viral, pathogen, or tumor clearance) relative to such levels before the intervention. In one embodiment, the level of enhancement is at least 50%, alternatively 60%, 70%, 80%, 90%, 100%, 120%, 150%, or 200%. The manner of measuring this enhancement is known to one of ordinary skill in the art.

[0033] A “*T cell dysfunctional disorder*” is a disorder or condition of T-cells characterized by decreased responsiveness to antigenic stimulation. In a particular embodiment, a T-cell dysfunctional disorder is a disorder that is specifically associated with inappropriate increased signaling through PD-1. In another embodiment, a T-cell dysfunctional disorder is one in which T-cells are anergic or have decreased ability to secrete cytokines, proliferate, or execute cytolytic activity. In a specific aspect, the decreased responsiveness results in ineffective control of a pathogen or tumor expressing an immunogen. Examples of T cell dysfunctional disorders characterized by T-cell dysfunction include unresolved acute infection, chronic infection and tumor immunity.

[0034] “*Tumor immunity*” refers to the process in which tumors evade immune recognition and clearance. Thus, as a therapeutic concept, tumor immunity is “treated” when such evasion is attenuated, and the tumors are recognized and attacked by the immune system. Examples of tumor recognition include tumor binding, tumor shrinkage and tumor clearance.

[0035] “*Immunogenicity*” refers to the ability of a particular substance to provoke an immune response. Tumors are immunogenic and enhancing tumor immunogenicity aids in the clearance of the tumor cells by the immune response. Examples of enhancing tumor immunogenicity include treatment with anti-PDL antibodies and a MEK inhibitor.

[0036] “*Sustained response*” refers to the sustained effect on reducing tumor growth after cessation of a treatment. For example, the tumor size may remain to be the same or smaller as compared to the size at the *beginning* of the administration phase. In some embodiments, the sustained response has a duration at least the same as the treatment duration, at least 1.5X, 2.0X, 2.5X, or 3.0X length of the treatment duration.

[0037] As used herein, “*cancer*” and “*cancerous*” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Included in this definition are benign and malignant cancers as well as dormant tumors or micrometastases.

Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include but are not limited to squamous cell cancer, lung cancer (including small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, and squamous carcinoma of the lung), cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer (including gastrointestinal cancer), pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer, as well as B-cell lymphoma (including low grade/follicular non-Hodgkin's lymphoma (NHL); small lymphocytic (SL) NHL; intermediate grade/follicular NHL; intermediate grade diffuse NHL; high grade immunoblastic NHL; high grade lymphoblastic NHL; high grade small non-cleaved cell NHL; bulky disease NHL; mantle cell lymphoma; AIDS-related lymphoma; and Waldenstrom's Macroglobulinemia); chronic lymphocytic leukemia (CLL); acute lymphoblastic leukemia (ALL); Hairy cell leukemia; chronic myeloblastic leukemia; and post-transplant lymphoproliferative disorder (PTLD), as well as abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain tumors), and Meigs' syndrome. Examples of cancer may include primary tumors of any of the above types of cancer or metastatic tumors at a second site derived from any of the above types of cancer.

[0038] The term “*antibody*” includes monoclonal antibodies (including full length antibodies which have an immunoglobulin Fc region), antibody compositions with polyepitopic specificity, multispecific antibodies (e.g., bispecific antibodies, diabodies, and single-chain molecules, as well as antibody fragments (e.g., Fab, F(ab')₂, and Fv). The term “*immunoglobulin*” (Ig) is used interchangeably with “*antibody*” herein.

[0039] The basic 4-chain antibody unit is a heterotetrameric glycoprotein composed of two identical light (L) chains and two identical heavy (H) chains. An IgM antibody consists of 5 of the basic heterotetramer units along with an additional polypeptide called a J chain, and contains 10 antigen binding sites, while IgA antibodies comprise from 2-5 of the basic 4-chain units which can polymerize to form polyvalent assemblages in combination with the J chain. In the case of IgGs, the 4-chain unit is generally about 150,000 daltons. Each L chain is linked to an H chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. Each H and L chain also has regularly

spaced intrachain disulfide bridges. Each H chain has at the N-terminus, a variable domain (V_H) followed by three constant domains (C_H) for each of the α and γ chains and four C_H domains for μ and ϵ isotypes. Each L chain has at the N-terminus, a variable domain (V_L) followed by a constant domain at its other end. The V_L is aligned with the V_H and the C_L is aligned with the first constant domain of the heavy chain (C_H1). Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains. The pairing of a V_H and V_L together forms a single antigen-binding site. For the structure and properties of the different classes of antibodies, see *e.g.*, *Basic and Clinical Immunology*, 8th Edition, Daniel P. Sties, Abba I. Terr and Tristram G. Parsolw (eds), Appleton & Lange, Norwalk, CT, 1994, page 71 and Chapter 6. The L chain from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains. Depending on the amino acid sequence of the constant domain of their heavy chains (C_H), immunoglobulins can be assigned to different classes or isotypes. There are five classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, having heavy chains designated α , δ , ϵ , γ and μ , respectively. The γ and α classes are further divided into subclasses on the basis of relatively minor differences in the C_H sequence and function, *e.g.*, humans express the following subclasses: IgG1, IgG2A, IgG2B, IgG3, IgG4, IgA1 and IgA2.

[0040] The “*variable region*” or “*variable domain*” of an antibody refers to the amino-terminal domains of the heavy or light chain of the antibody. The variable domains of the heavy chain and light chain may be referred to as “ VH ” and “ VL ”, respectively. These domains are generally the most variable parts of the antibody (relative to other antibodies of the same class) and contain the antigen binding sites.

[0041] The term “*variable*” refers to the fact that certain segments of the variable domains differ extensively in sequence among antibodies. The V domain mediates antigen binding and defines the specificity of a particular antibody for its particular antigen. However, the variability is not evenly distributed across the entire span of the variable domains. Instead, it is concentrated in three segments called hypervariable regions (HVRs) both in the light-chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a beta-sheet configuration, connected by three HVRs, which form loops connecting, and in some cases forming part of, the beta-sheet structure. The HVRs in each chain are held together in close proximity by the FR regions and, with the

HVRs from the other chain, contribute to the formation of the antigen binding site of antibodies (see Kabat *et al.*, *Sequences of Immunological Interest*, Fifth Edition, National Institute of Health, Bethesda, MD (1991)). The constant domains are not involved directly in the binding of antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

[0042] The term “*monoclonal antibody*” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translation modifications (*e.g.*, isomerizations, amidations) that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. In contrast to polyclonal antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier “*monoclonal*” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including, for example, the hybridoma method (*e.g.*, Kohler and Milstein, *Nature*, 256:495-97 (1975); Hongo *et al.*, *Hybridoma*, 14 (3): 253-260 (1995), Harlow *et al.*, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling *et al.*, in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981)), recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567), phage-display technologies (see, *e.g.*, Clackson *et al.*, *Nature*, 352: 624-628 (1991); Marks *et al.*, *J. Mol. Biol.* 222: 581-597 (1992); Sidhu *et al.*, *J. Mol. Biol.* 338(2): 299-310 (2004); Lee *et al.*, *J. Mol. Biol.* 340(5): 1073-1093 (2004); Fellouse, *Proc. Natl. Acad. Sci. USA* 101(34): 12467-12472 (2004); and Lee *et al.*, *J. Immunol. Methods* 284(1-2): 119-132 (2004), and technologies for producing human or human-like antibodies in animals that have parts or all of the human immunoglobulin loci or genes encoding human immunoglobulin sequences (see, *e.g.*, WO 1998/24893; WO 1996/34096; WO 1996/33735; WO 1991/10741; Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 2551 (1993); Jakobovits *et al.*, *Nature* 362: 255-258 (1993); Bruggemann *et al.*, *Year in Immunol.* 7:33 (1993); U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425;

and 5,661,016; Marks *et al.*, *Bio/Technology* 10: 779-783 (1992); Lonberg *et al.*, *Nature* 368: 856-859 (1994); Morrison, *Nature* 368: 812-813 (1994); Fishwild *et al.*, *Nature Biotechnol.* 14: 845-851 (1996); Neuberger, *Nature Biotechnol.* 14: 826 (1996); and Lonberg and Huszar, *Intern. Rev. Immunol.* 13: 65-93 (1995).

[0043] The term “*naked antibody*” refers to an antibody that is not conjugated to a cytotoxic moiety or radiolabel.

[0044] The terms “*full-length antibody*,” “*intact antibody*” or “*whole antibody*” are used interchangeably to refer to an antibody in its substantially intact form, as opposed to an antibody fragment. Specifically whole antibodies include those with heavy and light chains including an Fc region. The constant domains may be native sequence constant domains (e.g., human native sequence constant domains) or amino acid sequence variants thereof. In some cases, the intact antibody may have one or more effector functions.

[0045] An “*antibody fragment*” comprises a portion of an intact antibody, preferably the antigen binding and/or the variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂ and Fv fragments; diabodies; linear antibodies (see U.S. Patent 5,641,870, Example 2; Zapata *et al.*, *Protein Eng.* 8(10): 1057-1062 [1995]); single-chain antibody molecules and multispecific antibodies formed from antibody fragments. Papain digestion of antibodies produced two identical antigen-binding fragments, called "Fab" fragments, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. The Fab fragment consists of an entire L chain along with the variable region domain of the H chain (V_H), and the first constant domain of one heavy chain (C_{H1}). Each Fab fragment is monovalent with respect to antigen binding, *i.e.*, it has a single antigen-binding site. Pepsin treatment of an antibody yields a single large F(ab')₂ fragment which roughly corresponds to two disulfide linked Fab fragments having different antigen-binding activity and is still capable of cross-linking antigen. Fab' fragments differ from Fab fragments by having a few additional residues at the carboxy terminus of the C_{H1} domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0046] The Fc fragment comprises the carboxy-terminal portions of both H chains held together by disulfides. The effector functions of antibodies are determined by sequences in the

Fc region, the region which is also recognized by Fc receptors (FcR) found on certain types of cells.

[0047] “Fv” is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This fragment consists of a dimer of one heavy- and one light-chain variable region domain in tight, non-covalent association. From the folding of these two domains emanate six hypervariable loops (3 loops each from the H and L chain) that contribute the amino acid residues for antigen binding and confer antigen binding specificity to the antibody.

However, even a single variable domain (or half of an Fv comprising only three HVRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0048] “*Single-chain Fv*” also abbreviated as “*sFv*” or “*scFv*” are antibody fragments that comprise the V_H and V_L antibody domains connected into a single polypeptide chain. Preferably, the sFv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of the sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenburg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

[0049] “*Functional fragments*” of the antibodies of the invention comprise a portion of an intact antibody, generally including the antigen binding or variable region of the intact antibody or the Fc region of an antibody which retains or has modified FcR binding capability. Examples of antibody fragments include linear antibody, single-chain antibody molecules and multispecific antibodies formed from antibody fragments.

[0050] The term “*diabodies*” refers to small antibody fragments prepared by constructing sFv fragments (see preceding paragraph) with short linkers (about 5-10) residues) between the V_H and V_L domains such that inter-chain but not intra-chain pairing of the V domains is achieved, thereby resulting in a bivalent fragment, *i.e.*, a fragment having two antigen-binding sites.

Bispecific diabodies are heterodimers of two “crossover” sFv fragments in which the V_H and V_L domains of the two antibodies are present on different polypeptide chains. Diabodies are described in greater detail in, for example, EP 404,097; WO 93/11161; Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 6444-6448 (1993).

[0051] The monoclonal antibodies herein specifically include “*chimeric*” antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or

belonging to a particular antibody class or subclass, while the remainder of the chain(s) is(are) identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)). Chimeric antibodies of interest herein include PRIMATIZED® antibodies wherein the antigen-binding region of the antibody is derived from an antibody produced by, *e.g.*, immunizing macaque monkeys with an antigen of interest. As used herein, “humanized antibody” is used a subset of “chimeric antibodies.”

[0052] “Humanized” forms of non-human (*e.g.*, murine) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. In one embodiment, a humanized antibody is a human immunoglobulin (recipient antibody) in which residues from an HVR (hereinafter defined) of the recipient are replaced by residues from an HVR of a non-human species (donor antibody) such as mouse, rat, rabbit or non-human primate having the desired specificity, affinity, and/or capacity. In some instances, framework (“FR”) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications may be made to further refine antibody performance, such as binding affinity. In general, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin sequence, and all or substantially all of the FR regions are those of a human immunoglobulin sequence, although the FR regions may include one or more individual FR residue substitutions that improve antibody performance, such as binding affinity, isomerization, immunogenicity, *etc.* The number of these amino acid substitutions in the FR are typically no more than 6 in the H chain, and in the L chain, no more than 3. The humanized antibody optionally will also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see, *e.g.*, Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992). See also, for example, Vaswani and Hamilton, *Ann. Allergy, Asthma & Immunol.* 1:105-115 (1998); Harris, *Biochem. Soc. Transactions* 23:1035-1038 (1995); Hurle and Gross, *Curr. Op. Biotech.* 5:428-433 (1994); and U.S. Pat. Nos. 6,982,321 and 7,087,409.

[0053] A “*human antibody*” is an antibody that possesses an amino-acid sequence corresponding to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. Human antibodies can be produced using various techniques known in the art, including phage-display libraries. Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks *et al.*, *J. Mol. Biol.*, 222:581 (1991). Also available for the preparation of human monoclonal antibodies are methods described in Cole *et al.*, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985); Boerner *et al.*, *J. Immunol.*, 147(1):86-95 (1991). See also van Dijk and van de Winkel, *Curr. Opin. Pharmacol.*, 5: 368-74 (2001). Human antibodies can be prepared by administering the antigen to a transgenic animal that has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled, *e.g.*, immunized xenomice (see, *e.g.*, U.S. Pat. Nos. 6,075,181 and 6,150,584 regarding XENOMOUSETM technology). See also, for example, Li *et al.*, *Proc. Natl. Acad. Sci. USA*, 103:3557-3562 (2006) regarding human antibodies generated via a human B-cell hybridoma technology.

[0054] The term “*hypervariable region*,” “*HVR*,” or “*HV*,” when used herein refers to the regions of an antibody variable domain which are hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six HVRs; three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). In native antibodies, H3 and L3 display the most diversity of the six HVRs, and H3 in particular is believed to play a unique role in conferring fine specificity to antibodies. See, *e.g.*, Xu *et al.*, *Immunity* 13:37-45 (2000); Johnson and Wu, in *Methods in Molecular Biology* 248:1-25 (Lo, ed., Human Press, Totowa, NJ, 2003). Indeed, naturally occurring camelid antibodies consisting of a heavy chain only are functional and stable in the absence of light chain. See, *e.g.*, Hamers-Casterman *et al.*, *Nature* 363:446-448 (1993); Sheriff *et al.*, *Nature Struct. Biol.* 3:733-736 (1996).

[0055] A number of HVR delineations are in use and are encompassed herein. The Kabat Complementarity Determining Regions (CDRs) are based on sequence variability and are the most commonly used (Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). Chothia refers instead to the location of the structural loops (Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987)). The AbM HVRs represent a compromise between the Kabat HVRs and Chothia

structural loops, and are used by Oxford Molecular's AbM antibody modeling software. The "contact" HVRs are based on an analysis of the available complex crystal structures. The residues from each of these HVRs are noted below.

Loop	Kabat	AbM	Chothia	Contact
L1	L24-L34	L24-L34	L26-L32	L30-L36
L2	L50-L56	L50-L56	L50-L52	L46-L55
L3	L89-L97	L89-L97	L91-L96	L89-L96
H1	H31-H35B	H26-H35B	H26-H32	H30-H35B (Kabat numbering)
H1	H31-H35	H26-H35	H26-H32	H30-H35 (Chothia numbering)
H2	H50-H65	H50-H58	H53-H55	H47-H58
H3	H95-H102	H95-H102	H96-H101	H93-H101

[0056] HVRs may comprise "extended HVRs" as follows: 24-36 or 24-34 (L1), 46-56 or 50-56 (L2) and 89-97 or 89-96 (L3) in the VL and 26-35 (H1), 50-65 or 49-65 (H2) and 93-102, 94-102, or 95-102 (H3) in the VH. The variable domain residues are numbered according to Kabat *et al.*, *supra*, for each of these definitions.

[0057] The expression "*variable-domain residue-numbering as in Kabat*" or "*amino-acid-position numbering as in Kabat*," and variations thereof, refers to the numbering system used for heavy-chain variable domains or light-chain variable domains of the compilation of antibodies in Kabat *et al.*, *supra*. Using this numbering system, the actual linear amino acid sequence may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or HVR of the variable domain. For example, a heavy-chain variable domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 of H2 and inserted residues (*e.g.* residues 82a, 82b, and 82c, *etc.* according to Kabat) after heavy-chain FR residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a "standard" Kabat numbered sequence.

[0058] "*Framework*" or "FR" residues are those variable-domain residues other than the HVR residues as herein defined.

[0059] A "*human consensus framework*" or "*acceptor human framework*" is a framework that represents the most commonly occurring amino acid residues in a selection of human immunoglobulin VL or VH framework sequences. Generally, the selection of human immunoglobulin VL or VH sequences is from a subgroup of variable domain sequences. Generally, the subgroup of sequences is a subgroup as in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda,

MD (1991). Examples include for the VL, the subgroup may be subgroup kappa I, kappa II, kappa III or kappa IV as in Kabat *et al.*, *supra*. Additionally, for the VH, the subgroup may be subgroup I, subgroup II, or subgroup III as in Kabat *et al.*, *supra*. Alternatively, a human consensus framework can be derived from the above in which particular residues, such as when a human framework residue is selected based on its homology to the donor framework by aligning the donor framework sequence with a collection of various human framework sequences. An acceptor human framework “derived from” a human immunoglobulin framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may contain pre-existing amino acid sequence changes. In some embodiments, the number of pre-existing amino acid changes are 10 or less, 9 or less, 8 or less, 7 or less, 6 or less, 5 or less, 4 or less, 3 or less, or 2 or less.

[0060] A “VH subgroup III consensus framework” comprises the consensus sequence obtained from the amino acid sequences in variable heavy subgroup III of Kabat *et al.*, *supra*. In one embodiment, the VH subgroup III consensus framework amino acid sequence comprises at least a portion or all of each of the following sequences: EVQLVESGGGLVQPGGSLRLSCAAS (HC-FR1)(SEQ ID NO:4), WVRQAPGKGLEWV (HC-FR2), (SEQ ID NO:5), RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR (HC-FR3, SEQ ID NO:6), WGQGTLVTVSA (HC-FR4), (SEQ ID NO:7).

[0061] A “VL kappa I consensus framework” comprises the consensus sequence obtained from the amino acid sequences in variable light kappa subgroup I of Kabat *et al.*, *supra*. In one embodiment, the VH subgroup I consensus framework amino acid sequence comprises at least a portion or all of each of the following sequences: DIQMTQSPSSLSASVGDRVITIC (LC-FR1) (SEQ ID NO:11), WYQQKPGKAPKLLIY (LC-FR2) (SEQ ID NO:12), GVPSRFSGSGSGTDFLTISLQPEDFATYYC (LC-FR3)(SEQ ID NO:13), FGQGTKVEIKR (LC-FR4)(SEQ ID NO:14).

[0062] An “*amino-acid modification*” at a specified position, *e.g.* of the Fc region, refers to the substitution or deletion of the specified residue, or the insertion of at least one amino acid residue adjacent the specified residue. Insertion “adjacent” to a specified residue means insertion within one to two residues thereof. The insertion may be N-terminal or C-terminal to the specified residue. The preferred amino acid modification herein is a substitution.

[0063] An “*affinity-matured*” antibody is one with one or more alterations in one or more HVRs thereof that result in an improvement in the affinity of the antibody for antigen, compared

to a parent antibody that does not possess those alteration(s). In one embodiment, an affinity-matured antibody has nanomolar or even picomolar affinities for the target antigen. Affinity-matured antibodies are produced by procedures known in the art. For example, Marks *et al.*, *Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH- and VL-domain shuffling. Random mutagenesis of HVR and/or framework residues is described by, for example: Barbas *et al.* *Proc Nat. Acad. Sci. USA* 91:3809-3813 (1994); Schier *et al.* *Gene* 169:147-155 (1995); Yelton *et al.* *J. Immunol.* 155:1994-2004 (1995); Jackson *et al.*, *J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al.* *J. Mol. Biol.* 226:889-896 (1992).

[0064] As use herein, the term "*specifically binds to*" or is "*specific for*" refers to measurable and reproducible interactions such as binding between a target and an antibody, which is determinative of the presence of the target in the presence of a heterogeneous population of molecules including biological molecules. For example, an antibody that specifically binds to a target (which can be an epitope) is an antibody that binds this target with greater affinity, avidity, more readily, and/or with greater duration than it binds to other targets. In one embodiment, the extent of binding of an antibody to an unrelated target is less than about 10% of the binding of the antibody to the target as measured, *e.g.*, by a radioimmunoassay (RIA). In certain embodiments, an antibody that specifically binds to a target has a dissociation constant (Kd) of $\leq 1\mu\text{M}$, $\leq 100\text{ nM}$, $\leq 10\text{ nM}$, $\leq 1\text{ nM}$, or $\leq 0.1\text{ nM}$. In certain embodiments, an antibody specifically binds to an epitope on a protein that is conserved among the protein from different species. In another embodiment, specific binding can include, but does not require exclusive binding.

[0065] As used herein, the term "*immunoadhesin*" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (*i.e.*, is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2 (including IgG2A and IgG2B), IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM. The Ig fusions preferably include the substitution of a domain of a polypeptide or antibody described herein in

the place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also US Patent No. 5,428,130 issued June 27, 1995. For example, useful immunoadhesins as second medicaments useful for combination therapy herein include polypeptides that comprise the extracellular or PD-1 binding portions of PD-L1 or PD-L2 or the extracellular or PD-L1 or PD-L2 binding portions of PD-1, fused to a constant domain of an immunoglobulin sequence, such as a PD-L1 ECD – Fc, a PD-L2 ECD – Fc, and a PD-1 ECD - Fc, respectively. Immunoadhesin combinations of Ig Fc and ECD of cell surface receptors are sometimes termed soluble receptors.

[0066] A “*fusion protein*” and a “*fusion polypeptide*” refer to a polypeptide having two portions covalently linked together, where each of the portions is a polypeptide having a different property. The property may be a biological property, such as activity *in vitro* or *in vivo*. The property may also be simple chemical or physical property, such as binding to a target molecule, catalysis of a reaction, *etc.* The two portions may be linked directly by a single peptide bond or through a peptide linker but are in reading frame with each other.

[0067] A “*PD-1 oligopeptide*,” “*PD-L1 oligopeptide*,” or “*PD-L2 oligopeptide*” is an oligopeptide that binds, preferably specifically, to a PD-1, PD-L1 or PD-L2 negative costimulatory polypeptide, respectively, including a receptor, ligand or signaling component, respectively, as described herein. Such oligopeptides may be chemically synthesized using known oligopeptide synthesis methodology or may be prepared and purified using recombinant technology. Such oligopeptides are usually at least about 5 amino acids in length, alternatively at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 amino acids in length or more. Such oligopeptides may be identified using well known techniques. In this regard, it is noted that techniques for screening oligopeptide libraries for oligopeptides that are capable of specifically binding to a polypeptide target are well known in the art (see, *e.g.*, U.S. Patent Nos. 5,556,762, 5,750,373, 4,708,871, 4,833,092, 5,223,409, 5,403,484, 5,571,689, 5,663,143; PCT Publication Nos. WO 84/03506 and WO84/03564; Geysen *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 81:3998-4002 (1984); Geysen *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 82:178-182

(1985); Geysen *et al.*, in *Synthetic Peptides as Antigens*, 130-149 (1986); Geysen *et al.*, *J. Immunol. Meth.*, 102:259-274 (1987); Schoofs *et al.*, *J. Immunol.*, 140:611-616 (1988), Cwirla, S. E. *et al.* *Proc. Natl. Acad. Sci. USA*, 87:6378 (1990); Lowman, H.B. *et al.* *Biochemistry*, 30:10832 (1991); Clackson, T. *et al.* *Nature*, 352: 624 (1991); Marks, J. D. *et al.*, *J. Mol. Biol.*, 222:581 (1991); Kang, A.S. *et al.* *Proc. Natl. Acad. Sci. USA*, 88:8363 (1991), and Smith, G. P., *Current Opin. Biotechnol.*, 2:668 (1991).

[0068] A “blocking” antibody or an “antagonist” antibody is one that inhibits or reduces a biological activity of the antigen it binds. In some embodiments, blocking antibodies or antagonist antibodies substantially or completely inhibit the biological activity of the antigen. The anti-PD-L1 antibodies of the invention block the signaling through PD-1 so as to restore a functional response by T-cells (*e.g.*, proliferation, cytokine production, target cell killing) from a dysfunctional state to antigen stimulation.

[0069] An “agonist” or activating antibody is one that enhances or initiates signaling by the antigen to which it binds. In some embodiments, agonist antibodies cause or activate signaling without the presence of the natural ligand.

[0070] The term “Fc region” herein is used to define a C-terminal region of an immunoglobulin heavy chain, including native-sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy-chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue. Suitable native-sequence Fc regions for use in the antibodies of the invention include human IgG1, IgG2 (IgG2A, IgG2B), IgG3 and IgG4.

[0071] “Fc receptor” or “FcR” describes a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the Fc γ RI, Fc γ RII, and Fc γ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors, Fc γ RII receptors include Fc γ RIIA (an “activating receptor”) and Fc γ RIIB (an “inhibiting

receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor Fc γ RIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor Fc γ RIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (see M. Daëron, *Annu. Rev. Immunol.* 15:203-234 (1997). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9: 457-92 (1991); Capel *et al.*, *Immunomethods* 4: 25-34 (1994); and de Haas *et al.*, *J. Lab. Clin. Med.* 126: 330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein.

[0072] The term "*Fc receptor*" or "*FcR*" also includes the neonatal receptor, *FcRn*, which is responsible for the transfer of maternal IgGs to the fetus. Guyer *et al.*, *J. Immunol.* 117: 587 (1976) and Kim *et al.*, *J. Immunol.* 24: 249 (1994). Methods of measuring binding to FcRn are known (see, *e.g.*, Ghetie and Ward, *Immunol. Today* 18: (12): 592-8 (1997); Ghetie *et al.*, *Nature Biotechnology* 15 (7): 637-40 (1997); Hinton *et al.*, *J. Biol. Chem.* 279 (8): 6213-6 (2004); WO 2004/92219 (Hinton *et al.*). Binding to FcRn in vivo and serum half-life of human FcRn high-affinity binding polypeptides can be assayed, *e.g.*, in transgenic mice or transfected human cell lines expressing human FcRn, or in primates to which the polypeptides having a variant Fc region are administered. WO 2004/42072 (Presta) describes antibody variants which improved or diminished binding to FcRs. See also, *e.g.*, Shields *et al.*, *J. Biol. Chem.* 9(2): 6591-6604 (2001).

[0073] The phrase "*substantially reduced*," or "*substantially different*," as used herein, denotes a sufficiently high degree of difference between two numeric values (generally one associated with a molecule and the other associated with a reference/comparator molecule) such that one of skill in the art would consider the difference between the two values to be of statistical significance within the context of the biological characteristic measured by said values (*e.g.*, Kd values). The difference between said two values is, for example, greater than about 10%, greater than about 20%, greater than about 30%, greater than about 40%, and/or greater than about 50% as a function of the value for the reference/comparator molecule.

[0074] The term "*substantially similar*" or "*substantially the same*," as used herein, denotes a sufficiently high degree of similarity between two numeric values (for example, one associated with an antibody of the invention and the other associated with a reference/comparator antibody), such that one of skill in the art would consider the difference between the two values to be of little or no biological and/or statistical significance within the context of the biological

characteristic measured by said values (e.g., Kd values). The difference between said two values is, for example, less than about 50%, less than about 40%, less than about 30%, less than about 20%, and/or less than about 10% as a function of the reference/comparator value.

[0075] "Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers that are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

[0076] A "package insert" refers to instructions customarily included in commercial packages of medicaments that contain information about the indications customarily included in commercial packages of medicaments that contain information about the indications, usage, dosage, administration, contraindications, other medicaments to be combined with the packaged product, and/or warnings concerning the use of such medicaments, *etc.*

[0077] As used herein, the term "*treatment*" refers to clinical intervention designed to alter the natural course of the individual or cell being treated during the course of clinical pathology. Desirable effects of treatment include decreasing the rate of disease progression, ameliorating or palliating the disease state, and remission or improved prognosis. For example, an individual is successfully "treated" if one or more symptoms associated with cancer are mitigated or eliminated, including, but are not limited to, reducing the proliferation of (or destroying) cancerous cells, decreasing symptoms resulting from the disease, increasing the quality of life of those suffering from the disease, decreasing the dose of other medications required to treat the disease, delaying the progression of the disease, and/or prolonging survival of individuals.

[0078] As used herein, "*delaying progression of a disease*" means to defer, hinder, slow, retard, stabilize, and/or postpone development of the disease (such as cancer). This delay can be of varying lengths of time, depending on the history of the disease and/or individual being treated. As is evident to one skilled in the art, a sufficient or significant delay can, in effect,

encompass prevention, in that the individual does not develop the disease. For example, a late stage cancer, such as development of metastasis, may be delayed.

[0079] As used herein, “*reducing or inhibiting cancer relapse*” means to reduce or inhibit tumor or cancer relapse or tumor or cancer progression. As disclosed herein, cancer relapse and/or cancer progression include, without limitation, cancer metastasis.

[0080] An “*effective amount*” is at least the minimum concentration required to effect a measurable improvement or prevention of a particular disorder. An effective amount herein may vary according to factors such as the disease state, age, sex, and weight of the patient, and the ability of the antibody to elicit a desired response in the individual. An effective amount is also one in which any toxic or detrimental effects of the treatment are outweighed by the therapeutically beneficial effects. For prophylactic use, beneficial or desired results include results such as eliminating or reducing the risk, lessening the severity, or delaying the onset of the disease, including biochemical, histological and/or behavioral symptoms of the disease, its complications and intermediate pathological phenotypes presenting during development of the disease. For therapeutic use, beneficial or desired results include clinical results such as decreasing one or more symptoms resulting from the disease, increasing the quality of life of those suffering from the disease, decreasing the dose of other medications required to treat the disease, enhancing effect of another medication such as via targeting, delaying the progression of the disease, and/or prolonging survival. In the case of cancer or tumor, an effective amount of the drug may have the effect in reducing the number of cancer cells; reducing the tumor size; inhibiting (*i.e.*, slow to some extent or desirably stop) cancer cell infiltration into peripheral organs; inhibit (*i.e.*, slow to some extent and desirably stop) tumor metastasis; inhibiting to some extent tumor growth; and/or relieving to some extent one or more of the symptoms associated with the disorder. An effective amount can be administered in one or more administrations. For purposes of this invention, an effective amount of drug, compound, or pharmaceutical composition is an amount sufficient to accomplish prophylactic or therapeutic treatment either directly or indirectly. As is understood in the clinical context, an effective amount of a drug, compound, or pharmaceutical composition may or may not be achieved in conjunction with another drug, compound, or pharmaceutical composition. Thus, an “*effective amount*” may be considered in the context of administering one or more therapeutic agents, and a single agent may be considered to be given in an effective amount if, in conjunction with one or more other agents, a desirable result may be or is achieved.

[0081] As used herein, “in conjunction with” refers to administration of one treatment modality in addition to another treatment modality. As such, “in conjunction with” refers to administration of one treatment modality before, during, or after administration of the other treatment modality to the individual.

[0082] As used herein, “complete response” or “CR” refers to disappearance of all target lesions; “partial response” or “PR” refers to at least a 30% decrease in the sum of the longest diameters (SLD) of target lesions, taking as reference the baseline SLD; and “stable disease” or “SD” refers to neither sufficient shrinkage of target lesions to qualify for PR, nor sufficient increase to qualify for PD, taking as reference the smallest SLD since the treatment started.

[0083] As used herein, “progressive disease” or “PD” refers to at least a 20% increase in the SLD of target lesions, taking as reference the smallest SLD recorded since the treatment started or the presence of one or more new lesions.

[0084] As used herein, “progression free survival” (PFS) refers to the length of time during and after treatment during which the disease being treated (e.g., cancer) does not get worse. Progression-free survival may include the amount of time patients have experienced a complete response or a partial response, as well as the amount of time patients have experienced stable disease.

[0085] As used herein, "overall response rate" (ORR) refers to the sum of complete response (CR) rate and partial response (PR) rate.

[0086] As used herein, "overall survival" refers to the percentage of individuals in a group who are likely to be alive after a particular duration of time.

[0087] A “*chemotherapeutic agent*” is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXAN®); alkyl sulfonates such as busulfan, improsulfan, and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylololomelamine; acetogenins (especially bullatacin and bullatacinone); delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN®), CPT-11 (irinotecan, CAMPTOSAR®), acetylcamptothecin, scopolectin, and 9-aminocamptothecin); bryostatin; pemetrexed; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide;

cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; TLK-286; CDP323, an oral alpha-4 integrin inhibitor; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, chlophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (*e. g.*, calicheamicin, especially calicheamicin gammaI and calicheamicin omegaI (see, *e.g.*, Nicolaou *et al.*, *Angew. Chem Int. Ed. Engl.*, 33: 183-186 (1994)); dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabacin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (including ADRIAMYCIN®, morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin, doxorubicin HCl liposome injection (DOXIL®) and doxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate, gemcitabine (GEMZAR®), tegafur (UFTORAL®), capecitabine (XELODA®), an epothilone, and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprime, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, and imatinib (a 2-phenylaminopyrimidine derivative), as well as other c-Kit inhibitors; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as folinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; el fornithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; moperanomol; niraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine;

trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine (ELDISINE®, FILDESIN®); dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); thiotepa; taxoids, *e.g.*, paclitaxel (TAXOL®), albumin-engineered nanoparticle formulation of paclitaxel (ABRAXANE™), and doxetaxel (TAXOTERE®); chlorambucil; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine (VELBAN®); platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN®); oxaliplatin; leucovorin; vinorelbine (NAVELBINE®); novantrone; edatrexate; daunomycin; aminopterin; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone, and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATIN™) combined with 5-FU and leucovorin.

[0088] Additional examples of chemotherapeutic agents include anti-hormonal agents that act to regulate, reduce, block, or inhibit the effects of hormones that can promote the growth of cancer, and are often in the form of systemic, or whole-body treatment. They may be hormones themselves. Examples include anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX® tamoxifen), raloxifene (EVISTA®), droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (FARESTON®); anti-progesterones; estrogen receptor down-regulators (ERDs); estrogen receptor antagonists such as fulvestrant (FASLODEX®); agents that function to suppress or shut down the ovaries, for example, leutinizing hormone-releasing hormone (LHRH) agonists such as leuprolide acetate (LUPRON® and ELIGARD®), goserelin acetate, buserelin acetate and triptorelin; anti-androgens such as flutamide, nilutamide and bicalutamide; and aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, megestrol acetate (MEGASE®), exemestane (AROMASIN®), formestan, fadrozole, vorozole (RIVISOR®), letrozole (FEMARA®), and anastrozole (ARIMIDEX®). In addition, such definition of chemotherapeutic agents includes bisphosphonates such as clodronate (for example, BONEFOS® or OSTAC®), etidronate (DIDROCAL®), NE-58095, zoledronic acid/zoledronate (ZOMETA®), alendronate (FOSAMAX®), pamidronate (AREDIA®), tiludronate (SKELID®),

or risedronate (ACTONEL®); as well as troxacicabine (a 1,3-dioxolane nucleoside cytosine analog); anti-sense oligonucleotides, particularly those that inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Raf, H-Ras, and epidermal growth factor receptor (EGF-R); vaccines such as THERATOPE® vaccine and gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEUVECTIN® vaccine, and VAXID® vaccine; topoisomerase 1 inhibitor (e.g., LURTOTECAN®); an anti-estrogen such as fulvestrant; a Kit inhibitor such as imatinib or EXEL-0862 (a tyrosine kinase inhibitor); EGFR inhibitor such as erlotinib or cetuximab; an anti-VEGF inhibitor such as bevacizumab; arinotecan; rmRH (e.g., ABARELIX®); lapatinib and lapatinib ditosylate (an ErbB-2 and EGFR dual tyrosine kinase small-molecule inhibitor also known as GW572016); 17AAG (geldanamycin derivative that is a heat shock protein (Hsp) 90 poison), and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0089] As used herein, the term “*cytokine*” refers generically to proteins released by one cell population that act on another cell as intercellular mediators or have an autocrine effect on the cells producing the proteins. Examples of such cytokines include lymphokines, monokines; interleukins (“ILs”) such as IL-1, IL-1 α , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL10, IL-11, IL-12, IL-13, IL-15, IL-17A-F, IL-18 to IL-29 (such as IL-23), IL-31, including PROLEUKIN® rIL-2; a tumor-necrosis factor such as TNF- α or TNF- β , TGF- β 1-3; and other polypeptide factors including leukemia inhibitory factor (“LIF”), ciliary neurotrophic factor (“CNTF”), CNTF-like cytokine (“CLC”), cardiotrophin (“CT”), and kit ligand (“KL”).

[0090] As used herein, the term “*chemokine*” refers to soluble factors (e.g., cytokines) that have the ability to selectively induce chemotaxis and activation of leukocytes. They also trigger processes of angiogenesis, inflammation, wound healing, and tumorigenesis. Example chemokines include IL-8, a human homolog of murine keratinocyte chemoattractant (KC).

[0091] As used herein and in the appended claims, the singular forms “a,” “or,” and “the” include plural referents unless the context clearly dictates otherwise.

[0092] Reference to “about” a value or parameter herein includes (and describes) variations that are directed to that value or parameter *per se*. For example, description referring to “about X” includes description of “X”.

[0093] The term “*alkyl*” as used herein refers to a saturated linear or branched-chain monovalent hydrocarbon radical of one to twelve carbon atoms. Examples of alkyl groups include, but are not limited to, methyl (Me, -CH₃), ethyl (Et, -CH₂CH₃), 1-propyl (n-Pr, n-propyl,

-CH₂CH₂CH₃), 2-propyl (i-Pr, i-propyl, -CH(CH₃)₂), 1-butyl (n-Bu, n-butyl, -CH₂CH₂CH₂CH₃), 2-methyl-1-propyl (i-Bu, i-butyl, -CH₂CH(CH₃)₂), 2-butyl (s-Bu, s-butyl, -CH(CH₃)CH₂CH₃), 2-methyl-2-propyl (t-Bu, t-butyl, -C(CH₃)₃), 1-pentyl (n-pentyl, -CH₂CH₂CH₂CH₂CH₃), 2-pentyl (-CH(CH₃)CH₂CH₂CH₃), 3-pentyl (-CH(CH₂CH₃)₂), 2-methyl-2-butyl (-C(CH₃)₂CH₂CH₃), 3-methyl-2-butyl (-CH(CH₃)CH(CH₃)₂), 3-methyl-1-butyl (-CH₂CH₂CH(CH₃)₂), 2-methyl-1-butyl (-CH₂CH(CH₃)CH₂CH₃), 1-hexyl (-CH₂CH₂CH₂CH₂CH₂CH₃), 2-hexyl (-CH(CH₃)CH₂CH₂CH₂CH₃), 3-hexyl (-CH(CH₂CH₃)(CH₂CH₂CH₃)), 2-methyl-2-pentyl (-C(CH₃)₂CH₂CH₂CH₃), 3-methyl-2-pentyl (-CH(CH₃)CH(CH₃)CH₂CH₃), 4-methyl-2-pentyl (-CH(CH₃)CH₂CH(CH₃)₂), 3-methyl-3-pentyl (-C(CH₃)(CH₂CH₃)₂), 2-methyl-3-pentyl (-CH(CH₂CH₃)CH(CH₃)₂), 2,3-dimethyl-2-butyl (-C(CH₃)₂CH(CH₃)₂), 3,3-dimethyl-2-butyl (-CH(CH₃)C(CH₃)₃, 1-heptyl, 1-octyl, and the like.

[0094] The term "alkenyl" refers to linear or branched-chain monovalent hydrocarbon radical of two to twelve carbon atoms with at least one site of unsaturation, i.e., a carbon-carbon, sp² double bond, wherein the alkenyl radical includes radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations. Examples include, but are not limited to, ethylenyl or vinyl (-CH=CH₂), allyl (-CH₂CH=CH₂), and the like.

[0095] The term "alkynyl" refers to a linear or branched monovalent hydrocarbon radical of two to twelve carbon atoms with at least one site of unsaturation, i.e., a carbon-carbon, sp triple bond. Examples include, but are not limited to, ethynyl (-C≡CH), propynyl (propargyl, -CH₂C≡CH), and the like.

[0096] The terms "carbocycle", "carbocyclyl", "carbocyclic ring" and "cycloalkyl" refer to a monovalent non-aromatic, saturated or partially unsaturated ring having 3 to 12 carbon atoms as a monocyclic ring or 7 to 12 carbon atoms as a bicyclic ring. Bicyclic carbocycles having 7 to 12 atoms can be arranged, for example, as a bicyclo [4,5], [5,5], [5,6] or [6,6] system, and bicyclic carbocycles having 9 or 10 ring atoms can be arranged as a bicyclo [5,6] or [6,6] system, or as bridged systems such as bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane and bicyclo[3.2.2]nonane. Examples of monocyclic carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, cyclohexadienyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, cycloundecyl, cyclododecyl, and the like.

[0097] "Aryl" means a monovalent aromatic hydrocarbon radical of 6-18 carbon atoms derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic

ring system. Some aryl groups are represented in the exemplary structures as "Ar". Aryl includes bicyclic radicals comprising an aromatic ring fused to a saturated, partially unsaturated ring, or aromatic carbocyclic or heterocyclic ring. Typical aryl groups include, but are not limited to, radicals derived from benzene (phenyl), substituted benzenes, naphthalene, anthracene, indenyl, indanyl, 1,2-dihydronaphthalene, 1,2,3,4-tetrahydronaphthyl, and the like.

[0098] The terms "heterocycle," "heterocyclyl" and "heterocyclic ring" are used interchangeably herein and refer to a saturated or a partially unsaturated (i.e., having one or more double and/or triple bonds within the ring) carbocyclic radical of 3 to 18 ring atoms in which at least one ring atom is a heteroatom selected from nitrogen, oxygen and sulfur, the remaining ring atoms being C, where one or more ring atoms is optionally substituted independently with one or more substituents described below. A heterocycle may be a monocycle having 3 to 7 ring members (2 to 6 carbon atoms and 1 to 4 heteroatoms selected from N, O, P, and S) or a bicycle having 7 to 10 ring members (4 to 9 carbon atoms and 1 to 6 heteroatoms selected from N, O, P, and S), for example: a bicyclo [4,5], [5,5], [5,6], or [6,6] system. Heterocycles are described in Paquette, Leo A.; "Principles of Modern Heterocyclic Chemistry" (W.A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; "The Chemistry of Heterocyclic Compounds, A series of Monographs" (John Wiley & Sons, New York, 1950 to present), in particular Volumes 13, 14, 16, 19, and 28; and J. Am. Chem. Soc. (1960) 82:5566. "Heterocyclyl" also includes radicals where heterocycle radicals are fused with a saturated, partially unsaturated ring, or aromatic carbocyclic or heterocyclic ring. Examples of heterocyclic rings include, but are not limited to, pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, thioxanyl, piperazinyl, homopiperazinyl, azetidinyl, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepinyl, diazepinyl, thiazepinyl, 2-pyrrolinyl, 3-pyrrolinyl, indolinyl, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazolinyl, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidinylimidazolinyl, imidazolidinyl, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, and azabicyclo[2.2.2]hexanyl. Spiro moieties are also included within the scope of this definition. Examples of a heterocyclic group wherein ring atoms are substituted with oxo (=O) moieties are pyrimidinonyl and 1,1-dioxo-thiomorpholinyl.

[0099] The term "heteroaryl" refers to a monovalent aromatic radical of 5- or 6-membered rings, and includes fused ring systems (at least one of which is aromatic) of 5-18 atoms,

containing one or more heteroatoms independently selected from nitrogen, oxygen, and sulfur. Examples of heteroaryl groups are pyridinyl (including, for example, 2-hydroxypyridinyl), imidazolyl, imidazopyridinyl, pyrimidinyl (including, for example, 4-hydroxypyrimidinyl), pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indolizinyl, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, triazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, and furopyridinyl.

[0100] The heterocycle or heteroaryl groups may be carbon (carbon-linked) or nitrogen (nitrogen-linked) attached where such is possible. By way of example and not limitation, carbon bonded heterocycles or heteroaryls are bonded at position 2, 3, 4, 5, or 6 of a pyridine, position 3, 4, 5, or 6 of a pyridazine, position 2, 4, 5, or 6 of a pyrimidine, position 2, 3, 5, or 6 of a pyrazine, position 2, 3, 4, or 5 of a furan, tetrahydrofuran, thiofuran, thiophene, pyrrole or tetrahydropyrrole, position 2, 4, or 5 of an oxazole, imidazole or thiazole, position 3, 4, or 5 of an isoxazole, pyrazole, or isothiazole, position 2 or 3 of an aziridine, position 2, 3, or 4 of an azetidine, position 2, 3, 4, 5, 6, 7, or 8 of a quinoline or position 1, 3, 4, 5, 6, 7, or 8 of an isoquinoline.

[0101] By way of example and not limitation, nitrogen bonded heterocycles or heteroaryls are bonded at position 1 of an aziridine, azetidine, pyrrole, pyrrolidine, 2-pyrroline, 3-pyrroline, imidazole, imidazolidine, 2-imidazoline, 3-imidazoline, pyrazole, pyrazoline, 2-pyrazoline, 3-pyrazoline, piperidine, piperazine, indole, indoline, 1H-indazole, position 2 of a isoindole, or isoindoline, position 4 of a morpholine, and position 9 of a carbazole, or β -carboline.

[0102] The heteroatoms present in heteroaryl or heterocyclyl include the oxidized forms such as $\text{N}^+\rightarrow\text{O}^-$, $\text{S}(\text{O})$ and $\text{S}(\text{O})_2$.

[0103] The term "halo" refers to F, Cl, Br or I.

[0104] The phrase "pharmaceutically acceptable salt" as used herein, refers to pharmaceutically acceptable organic or inorganic salts of a compound of the invention. Exemplary salts include, but are not limited, to sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate, isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate "mesylate", ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, pamoate

(i.e., 1,1'-methylene-bis -(2-hydroxy-3-naphthoate)) salts, alkali metal (e.g., sodium and potassium) salts, alkaline earth metal (e.g., magnesium) salts, and ammonium salts. A pharmaceutically acceptable salt may involve the inclusion of another molecule such as an acetate ion, a succinate ion or other counter ion. The counter ion may be any organic or inorganic moiety that stabilizes the charge on the parent compound. Furthermore, a pharmaceutically acceptable salt may have more than one charged atom in its structure. Instances where multiple charged atoms are part of the pharmaceutically acceptable salt can have multiple counter ions. Hence, a pharmaceutically acceptable salt can have one or more charged atoms and/or one or more counter ion.

[0105] If the compound of the invention is a base, the desired pharmaceutically acceptable salt may be prepared by any suitable method available in the art, for example, treatment of the free base with an inorganic acid, such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, methanesulfonic acid, phosphoric acid and the like, or with an organic acid, such as acetic acid, maleic acid, succinic acid, mandelic acid, fumaric acid, malonic acid, pyruvic acid, oxalic acid, glycolic acid, salicylic acid, a pyranosidyl acid, such as glucuronic acid or galacturonic acid, an alpha hydroxy acid, such as citric acid or tartaric acid, an amino acid, such as aspartic acid or glutamic acid, an aromatic acid, such as benzoic acid or cinnamic acid, a sulfonic acid, such as p-toluenesulfonic acid or ethanesulfonic acid, or the like.

[0106] If the compound of the invention is an acid, the desired pharmaceutically acceptable salt may be prepared by any suitable method, for example, treatment of the free acid with an inorganic or organic base, such as an amine (primary, secondary or tertiary), an alkali metal hydroxide or alkaline earth metal hydroxide, or the like. Illustrative examples of suitable salts include, but are not limited to, organic salts derived from amino acids, such as glycine and arginine, ammonia, primary, secondary, and tertiary amines, and cyclic amines, such as piperidine, morpholine and piperazine, and inorganic salts derived from sodium, calcium, potassium, magnesium, manganese, iron, copper, zinc, aluminum and lithium.

[0107] The phrase "pharmaceutically acceptable" indicates that the substance or composition must be compatible chemically and/or toxicologically, with the other ingredients comprising a formulation, and/or the mammal being treated therewith.

[0108] A "solvate" refers to an association or complex of one or more solvent molecules and a compound of the invention. Examples of solvents that form solvates include, but are not limited

to, water, isopropanol, ethanol, methanol, DMSO, ethyl acetate, acetic acid, and ethanolamine. The term "hydrate" refers to the complex where the solvent molecule is water.

[0109] It is understood that aspects and variations of the invention described herein include "consisting of" and/or "consisting essentially of" aspects and variations.

III. Methods

[0110] In one aspect, provided herein is a method for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor.

[0111] The methods of this invention may find use in treating conditions where enhanced immunogenicity is desired such as increasing tumor immunogenicity for the treatment of cancer. A variety of cancers may be treated, or their progression may be delayed, including but are not limited to a cancer that may contain a B-raf V600E mutation, a cancer that may contain a B-raf wildtype, a cancer that may contain a KRAS wildtype, or a cancer that may contain an activating KRAS mutation.

[0112] In some embodiments, the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist. In some embodiments, the individual has been previously treated with a B-raf antagonist for cancer. In some embodiments, the individual has not been previously treated with a B-raf antagonist. B-raf is a serine-threonine kinase known to be frequently mutated in cancer, *e.g.*, malignant melanoma, colorectal, ovarian, and thyroid cancer. Typically, B-raf mutations observed in cancer include activating mutations, such as the V600E mutation. Without wishing to be bound to theory, it is thought that activating mutations in B-raf promote deregulated MAPK/ERK signaling, leading to tumor cell proliferation and survival.

[0113] While B-raf antagonists (*e.g.*, B-raf inhibitors) are known to produce effective short-term increases in patient survival and tumor regression, resistance to B-raf inhibition is frequently observed (see, *e.g.*, Tentori, L., *et al.* Trends Pharmacol. Sci. 34(12):656-66 (2013)). Resistance to B-raf inhibition may be characterized by numerous phenomena. In some embodiments, resistance to B-raf inhibition may be characterized by one or more of MAPK pathway reactivation, PI3K activation, CRAF upregulation, NRAS mutation, PDGFR overexpression, COT1 overexpression, IGFR-1 overexpression, MEK1 mutation, HGF expression, PD-L1 overexpression, MEK2 mutation, MITF focal amplification, AKT mutation (*e.g.*, AKT1 or AKT3), B-raf amplification, and the formation of RAF dimers (*e.g.*,

CRAF/CRAF dimers, CRAF/B-raf dimers, and mutated dimerizing B-raf such as a B-raf variant lacking exons 4-8).

[0114] In some embodiments, resistance to B-raf inhibition may refer to a cancer cell or tumor that is refractory to B-raf inhibition. Resistance to B-raf inhibition is used herein in the broadest sense and may include any cancer cell that was previously or may be expected to be sensitive to B-raf inhibition through any particular mechanism or at any particular dose of a B-raf antagonist of the present disclosure. Resistance to B-raf inhibition may refer to B-raf activity in the presence of a B-raf antagonist. Resistance to B-raf inhibition may refer to a cell that grows in the presence of a B-raf antagonist when it is not expected to grow under such a condition, regardless of the enzymatic activity of B-raf that may be present.

[0115] In some embodiments, the B-raf antagonist is a small molecule inhibitor, an antibody, a peptide, a peptidomimetic, an aptamer or a polynucleotide.

[0116] In some embodiments, the individual has previously been treated with a B-raf antagonist. In some embodiments, a B-raf antagonist may include vemurafenib (also known as ZELBORAF®), dabrafenib (also known as TAFINLAR®), LGX818, GSK 2118436, RAF265, XL281, ARQ736, BAY73-4506, sorafenib, PLX4720, PLX-3603, GSK2118436, GDC-0879, or N-(3-(5-(4-chlorophenyl)-1H-pyrrolo[2,3-b]pyridine-3-carbonyl)-2,4-difluorophenyl)propane-1-sulfonamide. In some embodiments, a B-raf antagonist may include MLN2480, LY3009120, a MEK inhibitor such as trametinib (also known as MEKINIST®), or an EGFR inhibitor such as erlotinib (also known as TARCEVA®). Further descriptions of B-raf inhibitors may be found in Zambon, A. *et al.* *Bioorg. Med. Chem. Lett.* 22(2):789-92 (2012); Tentori, L., *et al.* *Trends Pharmacol. Sci.* 34(12):656-66 (2013); and Martin-Liberal, J. and Larkin, J. *Expert Opin. Pharmacother.* 15(9):1235-45 (2014), WO2007/002325, WO2007/002433, WO2009111278, WO2009111279, WO2009111277, WO2009111280 and U.S. Pat. No. 7,491,829.

[0117] In some embodiments, the B-raf antagonist is a selective B-raf antagonist of B-raf V600. In some embodiments, the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600E. In some embodiments, the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600E, B-raf V600K, and/or V600D. In some embodiments, the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600R. Techniques for generating and assaying B-raf antagonists that are selective for B-raf V600 have been described in the art; see, *e.g.*, Tsai, J., *et al.*, *Proc. Natl. Acad. Sci.* 105(8):3041-6 (2008).

[0118] In some embodiments, the cancer contains a BRAF V600E mutation, a BRAF wildtype, a KRAS wildtype, or an activating KRAS mutation. Methods for detecting the presence of such mutations may include, without limitation, PCR, Sanger sequencing, use of a mutation-specific antibody, and the like. Methods for determining whether a cancer expresses a B-raf V600 are known in the art, including without limitation the COBAS® 4800 B-raf V600 Mutation Test kit (Roche).

[0119] In some embodiments, the patient's cancer has been shown to express a B-raf biomarker. In some embodiments, B-raf biomarker is mutant B-raf. In some embodiments, mutant B-raf is B-raf V600. In some embodiments, B-raf V600 is B-raf V600E. In some embodiments, mutant B-raf is constitutively active.

[0120] In some embodiments, the cancer patient has progressed while receiving a B-raf antagonist therapy (i.e., the patient is "B-raf refractory"), or the patient has progressed within 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, or more after completing a B-raf antagonist-based therapy regimen.

[0121] In some embodiments, vemurafenib resistant cancer is meant that the cancer patient has progressed while receiving vemurafenib-based therapy (i.e., the patient is "vemurafenib refractory"). In some embodiments, cancer in the patient has progressed within 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, or more after completing a vemurafenib-based therapy regimen. In some embodiments, the cancer in the individual has progressed within 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 1 year, 2 years, 3 years, 4 years, or 5 years after completing a B-raf antagonist-based therapy regimen.

[0122] In some embodiments, the treatment results in a sustained response in the individual after cessation of the treatment.

[0123] In some embodiments, resistance to, e.g., B-raf inhibitor develops (is acquired) after treatment with B-raf antagonist. In other embodiments, the patient (e.g., the patient having B-raf resistant cancer) has not been previously treated with a B-raf antagonist.

[0124] In some embodiments, the patient is currently being treated with B-raf antagonist, such as a B-raf inhibitor. In some embodiments, the patient was previously treated with B-raf antagonist. In some embodiments, the patient was not previously treated with B-raf antagonist.

[0125] In some embodiments, the individual has colorectal cancer, melanoma, lung cancer, ovarian cancer, breast cancer, pancreatic cancer, hematological malignancy, bladder cancer, and/or renal cell carcinoma. In some embodiments, the individual has non-small cell lung cancer. The non-small cell lung cancer may be at early stage or at late stage. In some embodiments, the individual has small cell lung cancer. The small cell lung cancer may be at early stage or at late stage. In some embodiments, the individual has renal cell cancer. The renal cell cancer may be at early stage or at late stage. In some embodiments, the individual has colorectal cancer. The colorectal cancer may be at early stage or late stage. In some embodiments, the individual has ovarian cancer. The ovarian cancer may be at early stage or at late stage. In some embodiments, the individual has breast cancer. The breast cancer may be at early stage or at late stage. In some embodiments, the individual has pancreatic cancer. The pancreatic cancer may be at early stage or at late stage. In some embodiments, the individual has gastric carcinoma. The gastric carcinoma may be at early stage or at late stage. In some embodiments, the individual has bladder cancer. The bladder cancer may be at early stage or at late stage. In some embodiments, the individual has esophageal cancer. The esophageal cancer may be at early stage or at late stage. In some embodiments, the individual has mesothelioma. The mesothelioma may be at early stage or at late stage. In some embodiments, the individual has melanoma. The melanoma may be at early stage or at late stage. In some embodiments, the individual has head and neck cancer. The head and neck cancer may be at early stage or at late stage. In some embodiments, the individual has thyroid cancer. The thyroid cancer may be at early stage or at late stage. In some embodiments, the individual has sarcoma. The sarcoma may be at early stage or late stage. In some embodiments, the individual has prostate cancer. The prostate cancer may be at early stage or at late stage. In some embodiments, the individual has glioblastoma. The glioblastoma may be at early stage or at late stage. In some embodiments, the individual has cervical cancer. The cervical cancer may be at early stage or at late stage. In some embodiments, the individual has thymic carcinoma. The thymic carcinoma may be at early stage or at late stage. In some embodiments, the individual has leukemia. The leukemia may be at early stage or at late stage. In some embodiments, the individual has lymphomas. The lymphoma may be at early stage or at late stage. In some embodiments, the individual has myelomas. The myelomas may be at early stage or at late stage. In some embodiments, the individual has mycosis fungoides. The mycosis fungoides may be at early stage or at late stage. In some embodiments, the individual has merkel cell cancer. The merkel cell cancer may be at

early stage or at late stage. In some embodiments, the individual has hematologic malignancies. The hematological malignancies may be early stage or late stage. In some embodiments, the individual is a human. In some embodiments, the cancer is metastatic.

[0126] In some embodiments, the individual is a mammal, such as domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g., humans and non-human primates such as monkeys), rabbits, and rodents (e.g., mice and rats). In some embodiments, the individual treated is a human.

[0127] In another aspect, provided herein is a method of enhancing immune function in an individual having cancer comprising administering an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor.

[0128] In some embodiments, the CD8 T cells in the individual have enhanced priming, activation, proliferation and/or cytolytic activity relative to prior to the administration of the PD-1 pathway antagonist and the MEK inhibitor. In some embodiments, the CD8 T cell priming is characterized by elevated CD44 expression and/or enhanced cytolytic activity in CD8 T cells. In some embodiments, the CD8 T cell activation is characterized by an elevated frequency of γ -IFN $^+$ CD8 T cells. In some embodiments, the CD8 T cell is an antigen-specific T-cell. In some embodiments, the immune evasion by signaling through PD-L1 surface expression is inhibited.

[0129] In some embodiments, the cancer cells in the individual have elevated expression of MHC class I antigen expression relative to prior to the administration of the PD-1 pathway antagonist and the MEK inhibitor.

[0130] In some embodiments, the antigen presenting cells in the individual have enhanced maturation and activation relative prior to the administration of the PD-1 pathway antagonist and the MEK inhibitor. In some embodiments, wherein the antigen presenting cells are dendritic cells. In some embodiments, the maturation of the antigen presenting cells is characterized by increased frequency of CD83 $^+$ dendritic cells. In some embodiments, the activation of the antigen presenting cells is characterized by elevated expression of CD80 and CD86 on dendritic cells.

[0131] In some embodiments, the serum levels of cytokine IL-10 and/or chemokine IL-8, a human homolog of murine KC, in the individual are reduced relative prior to the administration of the anti-PD-L1 antibody and the MEK inhibitor.

[0132] In some embodiments, the cancer has elevated levels of T-cell infiltration.

[0133] In some embodiments, the combination therapy of the invention comprises administration of a PD-1 axis binding antagonist and a MEK inhibitor. The PD-1 axis binding antagonist and the MEK inhibitor may be administered in any suitable manner known in the art. For example, The PD-1 axis binding antagonist and the MEK inhibitor may be administered sequentially (at different times) or concurrently (at the same time).

[0134] In some embodiments, the MEK inhibitor is administered continuously. In some embodiments, the MEK inhibitor is administered intermittently. In some embodiments, the MEK inhibitor is administered before administration of the PD-1 axis binding antagonist. In some embodiments, the MEK inhibitor is administered simultaneously with administration of the PD-1 axis binding antagonist. In some embodiments, the MEK inhibitor is administered after administration of the PD-1 axis binding antagonist.

[0135] In some embodiments, provided is a method for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor, further comprising administering an additional therapy. The additional therapy may be radiation therapy, surgery (*e.g.*, lumpectomy and a mastectomy), chemotherapy, gene therapy, DNA therapy, viral therapy, RNA therapy, immunotherapy, bone marrow transplantation, nanotherapy, monoclonal antibody therapy, or a combination of the foregoing. The additional therapy may be in the form of adjuvant or neoadjuvant therapy. In some embodiments, the additional therapy is the administration of small molecule enzymatic inhibitor or anti-metastatic agent. In some embodiments, the additional therapy is the administration of side-effect limiting agents (*e.g.*, agents intended to lessen the occurrence and/or severity of side effects of treatment, such as anti-nausea agents, etc.). In some embodiments, the additional therapy is radiation therapy. In some embodiments, the additional therapy is surgery. In some embodiments, the additional therapy is a combination of radiation therapy and surgery. In some embodiments, the additional therapy is gamma irradiation. In some embodiments, the additional therapy is therapy targeting PI3K/AKT/mTOR pathway, HSP90 inhibitor, tubulin inhibitor, apoptosis inhibitor, and/or chemopreventative agent. The additional therapy may be one or more of the chemotherapeutic agents described hereabove.

[0136] The PD-1 axis binding antagonist and the MEK inhibitor may be administered by the same route of administration or by different routes of administration. In some embodiments, the PD-1 axis binding antagonist is administered intravenously, intramuscularly, subcutaneously, topically, orally, transdermally, intraperitoneally, intraorbitally, by implantation, by inhalation,

intrathecally, intraventricularly, or intranasally. In some embodiments, the MEK inhibitor is administered intravenously, intramuscularly, subcutaneously, topically, orally, transdermally, intraperitoneally, intraorbitally, by implantation, by inhalation, intrathecally, intraventricularly, or intranasally. An effective amount of the PD-1 axis binding antagonist and the MEK inhibitor may be administered for prevention or treatment of disease. The appropriate dosage of the PD-1 axis binding antagonist and/or the MEK inhibitor may be determined based on the type of disease to be treated, the type of the PD-1 axis binding antagonist and the MEK inhibitor, the severity and course of the disease, the clinical condition of the individual, the individual's clinical history and response to the treatment, and the discretion of the attending physician.

[0137] Any of the PD-1 axis binding antagonists and the MEK inhibitors known in the art or described below may be used in the methods.

PD-1 axis binding antagonists

[0138] Provided herein is a method for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor. For example, a PD-1 axis binding antagonist includes a PD-1 binding antagonist, a PD-L1 binding antagonist and a PD-L2 binding antagonist. Alternative names for "PD-1" include CD279 and SLEB2. Alternative names for "PD-L1" include B7-H1, B7-4, CD274, and B7-H. Alternative names for "PD-L2" include B7-DC, Btdc, and CD273. In some embodiments, PD-1, PD-L1, and PD-L2 are human PD-1, PD-L1 and PD-L2.

[0139] In some embodiments, the PD-1 binding antagonist is a molecule that inhibits the binding of PD-1 to its ligand binding partners. In a specific aspect the PD-1 ligand binding partners are PD-L1 and/or PD-L2. In another embodiment, a PD-L1 binding antagonist is a molecule that inhibits the binding of PD-L1 to its binding partners. In a specific aspect, PD-L1 binding partners are PD-1 and/or B7-1. In another embodiment, the PD-L2 binding antagonist is a molecule that inhibits the binding of PD-L2 to its binding partners. In a specific aspect, a PD-L2 binding partner is PD-1. The antagonist may be an antibody, an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide.

[0140] In some embodiment, the PD-1 binding antagonist is an anti-PD-1 antibody (e.g., a human antibody, a humanized antibody, or a chimeric antibody). In some embodiments, the anti-PD-1 antibody is selected from the group consisting of MDX-1106 (nivolumab, OPDIVO[®]), Merck 3745 (MK-3475, pembrolizumab, KEYTRUDA[®]), CT- 011 (pidilizumab), MEDI-0680

(AMP-514), PDR001, REGN2810, BGB-108, and BGB-A317. In some embodiments, the PD-1 binding antagonist is an immunoadhesin (e.g., an immunoadhesin comprising an extracellular or PD-1 binding portion of PD-L1 or PD-L2 fused to a constant region (e.g., an Fc region of an immunoglobulin sequence). In some embodiments, the PD-1 binding antagonist is AMP-224. In some embodiments, the PD-L1 binding antagonist is anti-PD-L1 antibody. In some embodiments, the anti-PD-L1 binding antagonist is selected from the group consisting of YW243.55.S70, MPDL3280A (atezolizumab), MEDI4736 (durvalumab), MDX-1105, and MSB0010718C (avelumab). MDX-1105, also known as BMS-936559, is an anti-PD-L1 antibody described in WO2007/005874. Antibody YW243.55.S70 (heavy and light chain variable region sequences shown in SEQ ID Nos: 20 and 21, respectively) is an anti-PD-L1 described in WO 2010/077634 A1. MEDI4736 is an anti-PD-L1 antibody described in WO2011/066389 and US2013/034559. MDX-1106, also known as nivolumab, MDX-1106-04, ONO-4538, BMS-936558, or OPDIVO®, is an anti-PD-1 antibody described in WO2006/121168. Merck 3745, also known as MK-3475, pembrolizumab, lambrolizumab, KEYTRUDA®, or SCH-900475, is an anti-PD-1 antibody described in WO2009/114335. CT-011, also known as hBAT or hBAT-1, is an anti-PD-1 antibody described in WO2009/101611. AMP-224, also known as B7-DCIg, is a PD-L2-Fc fusion soluble receptor described in WO2010/027827 and WO2011/066342.

[0141] In some embodiments, the anti-PD-1 antibody is MDX-1106. Alternative names for “MDX-1106” include MDX-1106-04, ONO-4538, BMS-936558 or nivolumab. In some embodiments, the anti-PD-1 antibody is Nivolumab (CAS Registry Number: 946414-94-4). In a still further embodiment, provided is an isolated anti-PD-1 antibody comprising a heavy chain variable region comprising the heavy chain variable region amino acid sequence from SEQ ID NO:22 and/or a light chain variable region comprising the light chain variable region amino acid sequence from SEQ ID NO:23. In a still further embodiment, provided is an isolated anti-PD-1 antibody comprising a heavy chain and/or a light chain sequence, wherein:

(a) the heavy chain sequence has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the heavy chain sequence:

QVQLVESGGVVQPGRSRLLDCKASGITFSNSGMHWVRQAPGKGLEWVAVIWY
DGSKRYYADSVKGRFTISRDNSKNTLFLQMNSLRAEDTAVYYCATNDDYWGQGTLVT
VSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPALV

QSSGLYSLSSVVTVPSQLGTKYTCNVDHKPSNTKVDKRVESKYGPPCPCPAPEFLGG
PSVFLPPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTPREEQF
NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQ
EEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDK
SRWQEGNVFSCSVMHEALHNHYTQKSLSLGK (SEQ ID NO:22), or

(b) the light chain sequences has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the light chain sequence:

EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASN RAT
GIPARFSGSGSGTDFLTISLEPEDFAVYYCQQSSNWPRTFGQGTKVEIKRTVAAPSVFI
FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSL
SSTLTL SKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:23).

[0142] Examples of anti-PD-L1 antibodies useful for the methods of this invention, and methods for making thereof are described in PCT patent application WO 2010/077634 A1, which is incorporated herein by reference.

[0143] In some embodiments, the PD-1 axis binding antagonist is an anti-PD-L1 antibody. In some embodiments, the anti-PD-L1 antibody is capable of inhibiting binding between PD-L1 and PD-1 and/or between PD-L1 and B7-1. In some embodiments, the anti-PD-L1 antibody is a monoclonal antibody. In some embodiments, the anti-PD-L1 antibody is an antibody fragment selected from the group consisting of Fab, Fab'-SH, Fv, scFv, and (Fab')₂ fragments. In some embodiments, the anti-PD-L1 antibody is a humanized antibody. In some embodiments, the anti-PD-L1 antibody is a human antibody.

[0144] The anti-PD-L1 antibodies useful in this invention, including compositions containing such antibodies, such as those described in WO 2010/077634 A1, may be used in combination with a MEK inhibitor to treat cancer. In some embodiments, the anti-PD-L1 antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:20 and a light chain variable region comprising the amino acid sequence of SEQ ID NO:21.

[0145] In one embodiment, the anti-PD-L1 antibody contains a heavy chain variable region polypeptide comprising an HVR-H1, HVR-H2 and HVR-H3 sequence, wherein:

- (a) the HVR-H1 sequence is GFTFSX₁SWIH (SEQ ID NO:1);
- (b) the HVR-H2 sequence is AWIX₂PYGGSX₃YYADSVKG (SEQ ID NO:2);
- (c) the HVR-H3 sequence is RHWPGGF DY (SEQ ID NO:3);

further wherein: X_1 is D or G; X_2 is S or L; X_3 is T or S.

[0146] In one specific aspect, X_1 is D; X_2 is S and X_3 is T. In another aspect, the polypeptide further comprises variable region heavy chain framework sequences juxtaposed between the HVRs according to the formula: (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a further aspect, the framework sequences are VH subgroup III consensus framework. In a still further aspect, at least one of the framework sequences is the following:

HC-FR1 is EVQLVESGGGLVQPGGSLRLSCAAS	(SEQ ID NO:4)
HC-FR2 is WVRQAPGKGLEWV	(SEQ ID NO:5)
HC-FR3 is RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR	(SEQ ID NO:6)
HC-FR4 is WGQGTLVTVSA	(SEQ ID NO:7).

[0147] In a still further aspect, the heavy chain polypeptide is further combined with a variable region light chain comprising an HVR-L1, HVR-L2 and HVR-L3, wherein:

(a) the HVR-L1 sequence is RASQX ₄ X ₅ X ₆ TX ₇ X ₈ A	(SEQ ID NO:8);
(b) the HVR-L2 sequence is SASX ₉ LX ₁₀ S,	(SEQ ID NO:9);
(c) the HVR-L3 sequence is QQX ₁₁ X ₁₂ X ₁₃ X ₁₄ PX ₁₅ T	(SEQ ID NO:10);

further wherein: X_4 is D or V; X_5 is V or I; X_6 is S or N; X_7 is A or F; X_8 is V or L; X_9 is F or T; X_{10} is Y or A; X_{11} is Y, G, F, or S; X_{12} is L, Y, F or W; X_{13} is Y, N, A, T, G, F or I; X_{14} is H, V, P, T or I; X_{15} is A, W, R, P or T.

[0148] In a still further aspect, X_4 is D; X_5 is V; X_6 is S; X_7 is A; X_8 is V; X_9 is F; X_{10} is Y; X_{11} is Y; X_{12} is L; X_{13} is Y; X_{14} is H; X_{15} is A. In a still further aspect, the light chain further comprises variable region light chain framework sequences juxtaposed between the HVRs according to the formula: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In a still further aspect, the framework sequences are derived from human consensus framework sequences. In a still further aspect, the framework sequences are VL kappa I consensus framework. In a still further aspect, at least one of the framework sequence is the following:

LC-FR1 is DIQMTQSPSSLSASVGDRVTITC	(SEQ ID NO:11)
LC-FR2 is WYQQKPGKAPKLLIY	(SEQ ID NO:12)
LC-FR3 is GPSPRFSGSGSGTDFLTISLQPEDFATYYC	(SEQ ID NO:13)

LC-FR4 is FGQGTKVEIKR (SEQ ID NO:14).

[0149] In another embodiment, provided is an isolated anti-PD-L1 antibody or antigen binding fragment comprising a heavy chain and a light chain variable region sequence, wherein:

- (a) the heavy chain comprises and HVR-H1, HVR-H2 and HVR-H3, wherein further:
 - (i) the HVR-H1 sequence is GFTFSX₁SWIH; (SEQ ID NO:1)
 - (ii) the HVR-H2 sequence is AWIX₂PYGGSX₃YYADSVKG (SEQ ID NO:2)
 - (iii) the HVR-H3 sequence is RHWPGGF DY, and (SEQ ID NO:3)
- (b) the light chain comprises and HVR-L1, HVR-L2 and HVR-L3, wherein further:
 - (i) the HVR-L1 sequence is RASQX₄X₅X₆TX₇X₈A (SEQ ID NO:8)
 - (ii) the HVR-L2 sequence is SASX₉LX₁₀S; and (SEQ ID NO:9)
 - (iii) the HVR-L3 sequence is QQX₁₁X₁₂X₁₃X₁₄PX₁₅T; (SEQ ID NO:10)

Further wherein: X₁ is D or G; X₂ is S or L; X₃ is T or S; X₄ is D or V; X₅ is V or I; X₆ is S or N; X₇ is A or F; X₈ is V or L; X₉ is F or T; X₁₀ is Y or A; X₁₁ is Y, G, F, or S; X₁₂ is L, Y, F or W; X₁₃ is Y, N, A, T, G, F or I; X₁₄ is H, V, P, T or I; X₁₅ is A, W, R, P or T.

[0150] In a specific aspect, X₁ is D; X₂ is S and X₃ is T. In another aspect, X₄ is D; X₅ is V; X₆ is S; X₇ is A; X₈ is V; X₉ is F; X₁₀ is Y; X₁₁ is Y; X₁₂ is L; X₁₃ is Y; X₁₄ is H; X₁₅ is A. In yet another aspect, X₁ is D; X₂ is S and X₃ is T, X₄ is D; X₅ is V; X₆ is S; X₇ is A; X₈ is V; X₉ is F; X₁₀ is Y; X₁₁ is Y; X₁₂ is L; X₁₃ is Y; X₁₄ is H and X₁₅ is A.

[0151] In a further aspect, the heavy chain variable region comprises one or more framework sequences juxtaposed between the HVRs as: (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one or more framework sequences juxtaposed between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In a still further aspect, the framework sequences are derived from human consensus framework sequences. In a still further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1	EVQLVESGGGLVQPGGSLRLSCAAS	(SEQ ID NO:4)
HC-FR2	WVRQAPGKGLEWV	(SEQ ID NO:5)
HC-FR3	RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR	(SEQ ID NO:6)

HC-FR4 WGQGTLVTVSA (SEQ ID NO:7).

[0152] In a still further aspect, the light chain framework sequences are derived from a Kabat kappa I, II, II or IV subgroup sequence. In a still further aspect, the light chain framework sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

LC-FR1	DIQMTQSPSSLSASVGDRVITIC	(SEQ ID NO:11)
LC-FR2	WYQQKPGKAPKLLIY	(SEQ ID NO:12)
LC-FR3	GVPSRFSGSGSGTDFTLTSSLQPEDFATYYC	(SEQ ID NO:13)
LC-FR4	FGQQGKVEIKR	(SEQ ID NO:14).

[0153] In a still further specific aspect, the antibody further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect the minimal effector function results from an “effector-less Fc mutation” or aglycosylation. In still a further embodiment, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

[0154] In yet another embodiment, provided is an anti-PD-L1 antibody comprising a heavy chain and a light chain variable region sequence, wherein:

- (a) the heavy chain further comprises an HVR-H1, HVR-H2 and an HVR-H3 sequence having at least 85% sequence identity to GFTFSDSWIH (SEQ ID NO:15), AWISPYGGSTYYADSVKG (SEQ ID NO:16) and RHWPGGFDY (SEQ ID NO:3), respectively, or
- (b) the light chain further comprises an HVR-L1, HVR-L2 and an HVR-L3 sequence having at least 85% sequence identity to RASQDVSTAVA (SEQ ID NO:17), SASFLYS (SEQ ID NO:18) and QQYLYHPAT (SEQ ID NO:19), respectively.

[0155] In a specific aspect, the sequence identity is 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. In another aspect, the heavy chain variable

region comprises one or more framework sequences juxtaposed between the HVRs as: (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one or more framework sequences juxtaposed between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a still further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1	EVQLVESGGGLVQPGGSLRLSCAAS	(SEQ ID NO:4)
HC-FR2	WVRQAPGKGLEWW	(SEQ ID NO:5)
HC-FR3	RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR	(SEQ ID NO:6)
HC-FR4	WGQGTLVTVSA	(SEQ ID NO:7).

[0156] In a still further aspect, the light chain framework sequences are derived from a Kabat kappa I, II, III or IV subgroup sequence. In a still further aspect, the light chain framework sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

LC-FR1	DIQMTQSPSSLSASVGDRVTITC	(SEQ ID NO:11)
LC-FR2	WYQQKPGKAPKLLIY	(SEQ ID NO:12)
LC-FR3	GVPSRFGSGSGTDFTLTISLQPEDFATYYC	(SEQ ID NO:13)
LC-FR4	FGQGTKVEIKR	(SEQ ID NO:14).

[0157] In a still further specific aspect, the antibody further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect the minimal effector function results from an

“effector-less Fc mutation” or aglycosylation. In still a further embodiment, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

[0158] In a still further embodiment, provided is an isolated anti-PD-L1 antibody comprising a heavy chain and a light chain variable region sequence, wherein:

(a) the heavy chain sequence has at least 85% sequence identity to the heavy chain sequence: EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWVRQAPGKGLEWVAWIS PYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGF DYWG QGTLVTVSA (SEQ ID NO:20), or

(b) the light chain sequence has at least 85% sequence identity to the light chain sequence: DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIY SASF LYSGVPSRFSGSQSGTDFLTISLQPEDFATYYCQQYLYHPATFGQGTKVEIKR (SEQ ID NO:21).

[0159] In a specific aspect, the sequence identity is 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. In another aspect, the heavy chain variable region comprises one or more framework sequences juxtaposed between the HVRs as: (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one or more framework sequences juxtaposed between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1	EVQLVESGGGLVQPGGSLRLSCAAS	(SEQ ID NO:4)
HC-FR2	WVRQAPGKGLEWV	(SEQ ID NO:5)
HC-FR3	RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR	(SEQ ID NO:6)
HC-FR4	WGQGTLVTVSA	(SEQ ID NO:7).

[0160] In a still further aspect, the light chain framework sequences are derived from a Kabat kappa I, II, III or IV subgroup sequence. In a still further aspect, the light chain framework

sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

LC-FR1	DIQMTQSPSSLSASVGDRVTTTC	(SEQ ID NO:11)
LC-FR2	WYQQKPGKAPKLLIY	(SEQ ID NO:12)
LC-FR3	GVPSRFSGSGSGTDFLTISLQPEDFATYYC	(SEQ ID NO:13)
LC-FR4	FGQGTKVEIKR	(SEQ ID NO:14).

[0161] In a still further specific aspect, the antibody further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic cells. In a still further specific aspect the minimal effector function results from an “effector-less Fc mutation” or aglycosylation. In still a further embodiment, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

[0162] In another further embodiment, provided is an isolated anti-PD-L1 antibody comprising a heavy chain and a light chain variable region sequence, wherein:

(a) the heavy chain sequence has at least 85% sequence identity to the heavy chain sequence: EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWRQAPGKGLEWVAWIS PYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFYWG QGTLTVVSS (SEQ ID NO:24), or

(b) the light chain sequence has at least 85% sequence identity to the light chain sequence: DIQMTQSPSSLSASVGDRVTTTCRASQDVSTAVAWYQQKPGKAPKLLIY SASFLYSGVPSRFSGSGSGTDFLTISLQPEDFATYYCQQYLYHPATFGQGTKVEIKR (SEQ ID NO:21).

[0163] In a still further embodiment, provided is an isolated anti-PDL1 antibody comprising a heavy chain and a light chain variable region sequence, wherein:

(a) the heavy chain sequence has at least 85% sequence identity to the heavy chain sequence: EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWRQAPGKGLEWVAWI

SPYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDYW
GQGTLVTVSSASTK (SEQ ID NO:28), or

(b) the light chain sequence has at least 85% sequence identity to the light chain sequence:

DIQMTQSPSSLSASVGDRVITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLYSGVPS
RFSGSGSGTDFTLTISSLQPEDFATYYCQQYLYHPATFGQGTKVEIKR (SEQ ID NO:21).

[0164] In a specific aspect, the sequence identity is 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. In another aspect, the heavy chain variable region comprises one or more framework sequences juxtaposed between the HVRs as: (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one or more framework sequences juxtaposed between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1	EVQLVESGGLVQPGGSLRLSCAAS	(SEQ ID NO:4)
HC-FR2	WVRQAPGKGLEWV	(SEQ ID NO:5)
HC-FR3	RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR	(SEQ ID NO:6)
HC-FR4	WGQGTLVTVSS	(SEQ ID NO:25).

[0165] In a still further aspect, the light chain framework sequences are derived from a Kabat kappa I, II, III or IV subgroup sequence. In a still further aspect, the light chain framework sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

LC-FR1	DIQMTQSPSSLSASVGDRVITC	(SEQ ID NO:11)
LC-FR2	WYQQKPGKAPKLLIY	(SEQ ID NO:12)
LC-FR3	GVPSRFSGSGTDFTLTISSLQPEDFATYYC	(SEQ ID NO:13)
LC-FR4	FGQGTKVEIKR	(SEQ ID NO:14).

[0166] In a still further specific aspect, the antibody further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic cells. In a still further specific aspect the minimal effector function results from an “effector-less Fc mutation” or aglycosylation. In still a further embodiment, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

[0167] In yet another embodiment, the anti-PD-1 antibody is MPDL3280A (atezolizumab). In a still further embodiment, provided is an isolated anti-PD-1 antibody comprising a heavy chain variable region comprising the heavy chain variable region amino acid sequence from SEQ ID NO:24 and/or a light chain variable region comprising the light chain variable region amino acid sequence from SEQ ID NO:25. In a still further embodiment, provided is an isolated anti-PD-1 antibody comprising a heavy chain and/or a light chain sequence, wherein:

(a) the heavy chain sequence has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the heavy chain sequence:

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWVRQAPGKGLEWVAWISPYGGST
YYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDYWGQGTLVT
VSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL
QSSGLYSLSSVVTVPSQLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEL
LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPR
EEQYASTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT
LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSK
LTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:26), or

(b) the light chain sequences has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the light chain sequence:

DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLIYSASFLYSGVPS

RFSGSGSGTDFLTISLQPEDFATYYCQQYLYHPATFGQGTKVEIKRTVAAPSVFIFPPS
DEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTL
TLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:27).

[0168] In a still further embodiment, the invention provides for compositions comprising any of the above described anti-PD-L1 antibodies in combination with at least one pharmaceutically-acceptable carrier.

[0169] In a still further embodiment, provided is an isolated nucleic acid encoding a light chain or a heavy chain variable region sequence of an anti-PD-L1 antibody, wherein:

- (a) the heavy chain further comprises and HVR-H1, HVR-H2 and an HVR-H3 sequence having at least 85% sequence identity to GFTFSDSWIH (SEQ ID NO:15), AWISPYGGSTYYADSVKG (SEQ ID NO:16) and RHWPGGF DY (SEQ ID NO:3), respectively, and
- (b) the light chain further comprises an HVR-L1, HVR-L2 and an HVR-L3 sequence having at least 85% sequence identity to RASQDVSTAVA (SEQ ID NO:17), SASFLYS (SEQ ID NO:18) and QQYLYHPAT (SEQ ID NO:19), respectively.

[0170] In a specific aspect, the sequence identity is 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. In aspect, the heavy chain variable region comprises one or more framework sequences juxtaposed between the HVRs as: (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one or more framework sequences juxtaposed between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1	EVQLVESGGGLVQPGGSLRLSCAAS	(SEQ ID NO:4)
HC-FR2	WVRQAPGKGLEWV	(SEQ ID NO:5)
HC-FR3	RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR	(SEQ ID NO:6)
HC-FR4	WGQGTLVTVSA	(SEQ ID NO:7).

[0171] In a still further aspect, the light chain framework sequences are derived from a Kabat kappa I, II, III or IV subgroup sequence. In a still further aspect, the light chain framework sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

LC-FR1	DIQMTQSPSSLSASVGDRVITIC	(SEQ ID NO:11)
LC-FR2	WYQQKPGKAPKLLIY	(SEQ ID NO:12)
LC-FR3	GVPSRFSGSGSGTDFTLTSSLQPEDFATYYC	(SEQ ID NO:13)
LC-FR4	FGQGTKVEIKR	(SEQ ID NO:14).

[0172] In a still further specific aspect, the antibody described herein (such as an anti-PD-1 antibody, an anti-PD-L1 antibody, or an anti-PD-L2 antibody) further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, and IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, and IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic cells. In a still further specific aspect the minimal effector function results from an “effector-less Fc mutation” or aglycosylation. In still a further aspect, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

[0173] In a still further aspect, provided herein are nucleic acids encoding any of the antibodies described herein. In some embodiments, the nucleic acid further comprises a vector suitable for expression of the nucleic acid encoding any of the previously described anti-PD-L1, anti-PD-1, or anti-PD-L2 antibodies. In a still further specific aspect, the vector further comprises a host cell suitable for expression of the nucleic acid. In a still further specific aspect, the host cell is a eukaryotic cell or a prokaryotic cell. In a still further specific aspect, the eukaryotic cell is a mammalian cell, such as Chinese Hamster Ovary (CHO).

[0174] The antibody or antigen binding fragment thereof, may be made using methods known in the art, for example, by a process comprising culturing a host cell containing nucleic acid encoding any of the previously described anti-PD-L1, anti-PD-1, or anti-PD-L2 antibodies or

antigen-binding fragment in a form suitable for expression, under conditions suitable to produce such antibody or fragment, and recovering the antibody or fragment.

[0175] In a still further embodiment, the invention provides for a composition comprising an anti-PD-L1, an anti-PD-1, or an anti-PD-L2 antibody or antigen binding fragment thereof as provided herein and at least one pharmaceutically acceptable carrier. In some embodiments, the anti-PD-L1, anti-PD-1, or anti-PD-L2 antibody or antigen binding fragment thereof administered to the individual is a composition comprising one or more pharmaceutically acceptable carrier. Any of the pharmaceutically acceptable carrier described herein or known in the art may be used.


MEK inhibitors

[0176] The invention provides methods for treating cancer or slowing progression of cancer in an individual comprising administering an effective amount of a PD-1 pathway antagonist and a MEK inhibitor. Any known MEK inhibitors are intended, such as the MEK inhibitor compounds described in PCT patent applications WO 03/077914 A1, WO 2005/121142 A1, WO 2007/044515 A1, WO 2008/024725 A1 and WO 2009/085983 A1, the content of which are incorporated herein by reference. The MEK inhibitor administered may be in a pharmaceutical composition or formulation. In some embodiments, the pharmaceutical composition or formulation comprises one or more MEK inhibitors described herein and a pharmaceutically acceptable carrier or excipient.

[0177] In some embodiments, the MEK inhibitor is a competitive inhibitor of MEK. In some embodiments, the MEK inhibitor is more selective against an activating KRAS mutation. In some embodiments, the MEK inhibitor is an allosteric inhibitor of MEK. In some embodiments, the MEK inhibitor is more selective against an activating B-raf mutation (e.g., B-raf V600E mutation). In some embodiments, the MEK inhibitor binds and inhibits the activity of MEK1 and/or MEK2 (such as human MEK1 and/or human MEK2).

[0178] In some embodiments, the MEK inhibitor is a compound selected from the group consisting of GDC-0973 (also known as “Cobimetinib” or “XL518”), G-38963, G02443714 (also known as “AS703206”), G02442104 (also known as “GSK-1120212”), and G00039805 (also known as “AZD-6244”), or a pharmaceutically acceptable salt or solvate thereof.

[0179] In some embodiments, the MEK inhibitor is a compound of formula (I),



or a pharmaceutically acceptable salt or solvate thereof, wherein A, X, R¹, R², R³, R⁴, R⁵, R⁶, and R⁷ are as defined in Group A, Group B, Group C, or Group D:

Group A:

A is arylene optionally substituted with one, two, three or four groups selected from R¹⁰, R¹², R¹⁴, R¹⁶, and R¹⁹ where R¹⁰, R¹², R¹⁴ and R¹⁶ are independently hydrogen, alkyl, alkenyl, alkynyl, halo, haloalkoxy, hydroxy, alkoxy, amino, alkylamino, dialkylamino, haloalkyl, -NHS(O)₂R⁸, -CN, -C(O)R⁸, -C(O)OR⁸, -C(O)NR⁸R^{8'} and -NR⁸C(O)R^{8'} and where R¹⁹ is hydrogen, alkyl, or alkenyl;

X is alkyl, halo, haloalkyl, or haloalkoxy;

R¹, R², R³, R⁴, R⁵ and R⁶ are independently hydrogen, halo, nitro, -NR⁸R^{8'}, -OR⁸, -NHS(O)₂R⁸, -CN, -S(O)_mR⁸, -S(O)₂NR⁸R^{8'}, -C(O)R⁸, -C(O)OR⁸, -C(O)NR⁸R^{8'}, -NR⁸C(O)OR^{8'}, -NR⁸C(O)NR⁸R^{8''}, -NR⁸C(O)OR^{8'}, -NR⁸C(O)R^{8'}, -CH₂N(R²⁵)(NR^{25a}R^{25b}), -CH₂NR²⁵C(=NH)(NR^{25a}R^{25b}), -CH₂NR²⁵C(=NH)(N(R^{25a})(NO₂)), -CH₂NR²⁵C(=NH)(N(R^{25a})(CN)), -CH₂NR²⁵C(=NH)(R²⁵), -CH₂NR²⁵C(NR^{25a}R^{25b})=CH(NO₂), alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, or heterocycloalkyl; where the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two, three, four, five, six or seven groups independently selected from halo, alkyl, haloalkyl, nitro, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted arylalkyl, optionally substituted heteroaryl, -OR⁸, -NR⁸R^{8'}, -NR⁸S(O)₂R⁹, -CN, -S(O)_mR⁹, -C(O)R⁸, -C(O)OR⁸, -C(O)NR⁸R^{8'}, -NR⁸C(O)NR⁸R^{8''}, -NR⁸C(O)OR^{8'} and -NR⁸C(O)R^{8'}; or one of R¹ and R² together with the carbon to which they are attached, R³ and R⁴ together with the carbon to which they are attached, and R⁵ and R⁶ together with the carbon to which they are attached form C(O) or C(=NOH);

m is 0, 1, or 2;

R⁷ is hydrogen, halo or alkyl;

each R^8 , $R^{8''}$ and $R^{8''''}$ is independently selected from hydrogen, hydroxy, optionally substituted alkoxy, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl; where the alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two, three, four, or five groups independently selected from alkyl, halo, hydroxy, hydroxyalkyl, optionally substituted alkoxy, alkoxyalkyl, haloalkyl, carboxy, alkoxy carbonyl, alkenyloxy carbonyl, optionally substituted cycloalkyl, optionally substituted cycloalkyloxy carbonyl, optionally substituted aryl, optionally substituted aryloxy, optionally substituted aryloxy carbonyl, optionally substituted arylalkyl, optionally substituted arylalkyloxy, optionally substituted arylalkyloxy carbonyl, nitro, cyano, optionally substituted heterocycloalkyl, optionally substituted heteroaryl, $-S(O)R^{31}$ (where n is 0, 1, or 2 and R^{31} is optionally substituted alkyl, optionally substituted aryl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), $-NR^{34}SO_2R^{34a}$ (where R^{34} is hydrogen or alkyl and R^{34a} is alkyl, alkenyl, cycloalkyl, aryl, heteroaryl, or heterocycloalkyl), $-SO_2NR^{35}R^{35a}$ (where R^{35} is hydrogen or alkyl and R^{35a} is alkyl, alkenyl, cycloalkyl, aryl, heteroaryl, or heterocycloalkyl), $-NR^{32}C(O)R^{32a}$ (where R^{32} is hydrogen or alkyl and R^{32a} is alkyl, alkenyl, alkoxy, or cycloalkyl), $-NR^{30}R^{30'}$ (where R^{30} and $R^{30'}$ are independently hydrogen, alkyl, or hydroxyalkyl), and $-C(O)NR^{33}R^{33a}$ (where R^{33} is hydrogen or alkyl and R^{33a} is alkyl, alkenyl, alkynyl, or cycloalkyl); and

each R^9 is independently selected from alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl; where the alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two, three, four, or five groups selected from halo, hydroxy, alkyl, haloalkyl, haloalkoxy, amino, alkylamino, and dialkylamino;

Group B:

A is heteroarylene optionally substituted with one, two, three, or four groups selected from R^{10} , R^{12} , R^{14} , R^{16} and R^{19} where R^{10} , R^{12} , R^{14} and R^{16} are independently hydrogen, alkyl, alkenyl, alkynyl, halo, haloalkoxy, hydroxy, alkoxy, cyano, amino, alkylamino, dialkylamino, haloalkyl, alkylsulfonylamino, alkylcarbonyl, alkenylcarbonyl, alkoxy carbonyl, alkenyloxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, or alkylcarbonylamino; where R^{19} is hydrogen, alkyl, or alkenyl;

and where each alkyl and alkenyl, either alone or as part of another group within R^{10} , R^{12} , R^{14} , R^{16} , and R^{19} , is independently optionally substituted with halo, hydroxy, or alkoxy; X is alkyl, halo, haloalkyl, or haloalkoxy; R^1 , R^2 , R^3 , R^4 , R^3 and R^6 are independently hydrogen, halo, nitro, $-NR^8R^{8'}$, $-OR^8$, $-NHS(O)2R^8$, $-CN$, $-S(O)_mR^8$, $-S(O)_2NR^8R^{8'}$, $-C(O)R^8$, $-C(O)OR^8$, $-C(O)NR^8R^{8'}$, $-NR^8C(O)OR^{8'}$, $-NR^8C(O)NR^{8'}$, $-NR^8C(O)OR^{8'}$, $-NR^8C(O)R^{8'}$, $-CH_2N(R^{25})(NR^{25a}R^{25b})$, $-CH_2NR^{25}C(=NH)(NR^{25a}R^{25b})$, $-CH_2NR^{25}C(=NH)(N(R^{25a})(NO_2))$, $-CH_2NR^{25}C(NH)(N(R^{25a})(CN))$, $-CH_2NR^{25}C(=NH)(R^{25})$, $-CH_2NR^{25}C(NR^{25a}R^{25b})=CH(NO_2)$, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, or heterocycloalkyl, where the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two, three, four, five, six or seven groups independently selected from halo, alkyl, haloalkyl, nitro, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted arylalkyl, optionally substituted heteroaryl, $-OR^8$, $-NR^8R^{8'}$, $-NR^8S(O)_2R^9$, $-CN$, $-S(O)_mR^9$, $-C(O)R^8$, $-C(O)OR^8$, $-C(O)NR^8R^{8'}$, $-NR^8C(O)NR^{8'}$, $-NR^8C(O)OR^{8'}$ and $-NR^8C(O)R^{8'}$; or one of R^1 and R^2 together with the carbon to which they are attached, R^3 and R^4 together with the carbon to which they are attached, and R^5 and R^6 together with the carbon to which they are attached form $C(O)$ or $C(=NOH)$;

m is 1 or 2;


R^7 is hydrogen, halo or alkyl; and

each R^8 , $R^{8'}$ and $R^{8''}$ is independently selected from hydrogen, hydroxy, optionally substituted alkoxy, alkyl, haloalkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl, where the alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two three, four, or five groups independently selected from alkyl, halo, hydroxy, hydroxyalkyl, optionally substituted alkoxy, alkoxyalkyl, haloalkyl, carboxy, carboxy ester, nitro, cyano, $-S(O)_nR^{31}$ (where n is 0, 1, or 2 and R^{31} is optionally substituted alkyl, optionally substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), $-NR^{36}S(O)_2R^{36a}$ (where R^{36} is hydrogen, alkyl, or alkenyl and R^{36a} is alkyl, alkenyl, optionally substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted

heteroaryl), $-\text{S}(\text{O})_2\text{NR}^{37}\text{R}^{37a}$ (where R^{37} is hydrogen, alkyl, or alkenyl and R^{37a} is alkyl, alkenyl, optionally substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted arylalkyl, optionally substituted aryloxy, optionally substituted arylalkyloxy, optionally substituted heteroaryl, $-\text{NHC}(\text{O})\text{R}^{32}$ (where R^{32} is alkyl, alkenyl, alkoxy, or cycloalkyl) and $-\text{NR}^{30}\text{R}^{30'}$ (where R^{30} and $\text{R}^{30'}$ are independently hydrogen, alkyl, or hydroxyalkyl), and $-\text{C}(\text{O})\text{NHR}^{33}$ (where R^{33} is alkyl, alkenyl, alkynyl, or cycloalkyl);

Group C:

A is



(a)

where R^{10} is hydrogen, alkyl, alkenyl, alkynyl, halo, haloalkoxy, hydroxy, alkoxy, amino, alkylamino, dialkylamino, haloalkyl, $-\text{NHS}(\text{O})_2\text{R}^8$, $-\text{CN}$, $-\text{C}(\text{O})\text{R}^8$, $-\text{C}(\text{O})\text{OR}^8$, $-\text{C}(\text{O})\text{NR}^8\text{R}^{8'}$ and $-\text{NR}^8\text{C}(\text{O})\text{R}^{8'}$;

R^{10a} is hydrogen, alkyl, or alkenyl;

Y^1 is $=\text{CH-}$ or $=\text{N-}$;

X is alkyl, halo, haloalkyl, or haloalkoxy;

$\text{R}^1, \text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5$ and R^6 are independently hydrogen, halo, nitro, $-\text{NR}^8\text{R}^{8'}$, $-\text{OR}^8$, $-\text{NHS}(\text{O})_2\text{R}^8$, $-\text{CN}$, $-\text{S}(\text{O})_m\text{R}^8$, $-\text{S}(\text{O})_2\text{NR}^8\text{R}^{8'}$, $-\text{C}(\text{O})\text{R}^8$, $-\text{C}(\text{O})\text{OR}^8$, $-\text{C}(\text{O})\text{NR}^8\text{R}^{8'}$, $-\text{NR}^8\text{C}(\text{O})\text{OR}^{8''}$, $-\text{NR}^8\text{C}(\text{O})\text{OR}^{8'}$, $-\text{NR}^8\text{C}(\text{O})\text{R}^{8'}$, $-\text{CH}_2\text{N}(\text{R}^{25})\text{NR}^{25a}\text{R}^{25b}$, $-\text{CH}_2\text{NR}^{25}\text{C}(\text{=NH})(\text{NR}^{25a}\text{R}^{25b})$, $-\text{CH}_2\text{NR}^{25}\text{C}(\text{=NH})(\text{N}(\text{R}^{25a})(\text{NO}_2))$, $-\text{CH}_2\text{NR}^{25}\text{C}(\text{=NH})(\text{N}(\text{R}^{25a})(\text{CN}))$, $-\text{CH}_2\text{NR}^{25}\text{C}(\text{=NH})(\text{R}^{25})$, $-\text{CH}_2\text{NR}^{25}\text{C}(\text{NR}^{25a}\text{R}^{25b})=\text{CH}(\text{NO}_2)$, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, or heterocycloalkyl, where the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two, three, four, five, six or seven groups independently selected from halo, alkyl, haloalkyl, nitro, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted arylalkyl, optionally substituted heteroaryl, $-\text{OR}^8$, $-\text{NR}^8\text{R}^{8'}$, $-\text{NR}^8\text{S}(\text{O})_2\text{R}^9$, $-\text{CN}$, $-\text{S}(\text{O})_m\text{R}^9$, $-\text{C}(\text{O})\text{R}^8$, $-\text{C}(\text{O})\text{OR}^8$, $-\text{C}(\text{O})\text{NR}^8\text{R}^{8'}$, $-\text{NR}^8\text{C}(\text{O})\text{NR}^{8''}\text{R}^{8''}$,

-NR⁸C(O)OR^{8'} and -NR⁸C(O)R^{8'}; or one of R¹ and R² together with the carbon to which they are attached, R³ and R⁴ together with the carbon to which they are attached, and R⁵ and R⁶ together with the carbon to which they are attached form C(O) or C(NOH);


m is 1 or 2;

R⁷ is hydrogen, halo or alkyl; and


each R⁸, R^{8'} and R^{8''} is independently selected from hydrogen, hydroxy, optionally substituted alkoxy, alkyl, haloalkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl, where the alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two three, four, or five groups independently selected from alkyl, halo, hydroxy, hydroxyalkyl, optionally substituted alkoxy, alkoxyalkyl, haloalkyl, carboxy, carboxy ester, nitro, cyano, -S(O)_nR³¹ (where n is 0, 1, or 2 and R³¹ is optionally substituted alkyl, optionally substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), -NR³⁶S(O)₂R^{36a} (where R³⁶ is hydrogen, alkyl, or alkenyl and R^{36a} is alkyl, alkenyl, optionally substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), -S(O)₂NR³⁷R^{37a} (where R³⁷ is hydrogen, alkyl, or alkenyl and R^{37a} is alkyl, alkenyl, optionally substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted arylalkyl, optionally substituted aryloxy, optionally substituted arylalkyloxy, optionally substituted heteroaryl, -NHC(O)R³² (where R³² is alkyl, alkenyl, alkoxy, or cycloalkyl) and -NR³⁰R^{30'} (where R³⁰ and R^{30'} are independently hydrogen, alkyl, or hydroxyalkyl), and -C(O)NHR³³ (where R³³ is alkyl, alkenyl, alkynyl, or cycloalkyl); or

Group D:

A is



(b) or



(c)

R⁴⁰ and R^{40a} are independently hydrogen or alkyl;

X is alkyl, halo, haloalkyl, or haloalkoxy;

R¹, R², R³, R⁴, R⁵ and R⁶ are independently hydrogen, halo, nitro, -NR⁸R^{8'}, -OR⁸, -NHS(O)R⁸, -CN, -S(O)_mR⁸, -S(O)₂NR⁸R^{8'}, -C(O)R⁸, -C(O)OR⁸, -C(O)NR⁸R^{8'}, -NR⁸C(O)OR⁸, -NR⁸C(O)NR⁸R^{8''}, -NR⁸C(O)OR^{8'}, -NR⁸C(O)R^{8'}, -CH₂N(R²⁵)(NR^{25a}R^{25b}), -CH₂NR²⁵C(=NH)(NR^{25a}R^{25b}), -CH₂NR²⁵C(=NH)(N(R^{25a})(NO₂)), -CH₂NR²⁵C(=NH)(N(R^{25a})(CN)), -CH₂NR²⁵C(=NH)(R²⁵), -CH₂NR²⁵C(NR^{25a}R^{25b})=CH(NO₂), alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, or heterocycloalkyl, where the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two, three, four, five, six or seven groups independently selected from halo, alkyl, haloalkyl, nitro, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted arylalkyl, optionally substituted heteroaryl, -OR⁸, -NR⁸R^{8'}, -NR⁸S(O)₂R⁹, -CN, -S(O)_mR⁹, -C(O)R⁸, -C(O)OR⁸, -C(O)NR⁸R^{8'}, -NR⁸C(O)NR⁸R^{8''}, -NR⁸C(O)OR^{8'} and -NR⁸C(O)R^{8'}; or one of R¹ and R² together with the carbon to which they are attached, R³ and R⁴ together with the carbon to which they are attached, and R⁵ and R⁶ together with the carbon to which they are attached form C(O) or C(NOH);


m is 1 or 2;

R⁷ is hydrogen, halo or alkyl; and

R⁸, R^{8'} and R^{8''} are independently selected from hydrogen, hydroxy, optionally substituted alkoxy, alkyl, haloalkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl, where the alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl are independently optionally substituted with one, two three, four, or five groups independently selected from alkyl, halo, hydroxy, hydroxyalkyl, optionally substituted alkoxy, alkoxyalkyl, haloalkyl, carboxy, carboxy ester, nitro, cyano, -S(O)_nR³¹ (where n is 0, 1, or 2 and R³¹ is optionally substituted alkyl, optionally


substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), -NR³⁶S(O)₂R^{36a} (where R³⁶ is hydrogen, alkyl, or alkenyl and R^{36a} is alkyl, alkenyl, optionally substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), -S(O)₂NR³⁷R^{37a} (where R³⁷ is hydrogen, alkyl, or alkenyl and R^{37a} is alkyl, alkenyl, optionally substituted aryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), optionally substituted heterocycloalkyl, or optionally substituted heteroaryl), -NHC(O)R³² (where R³² is alkyl, alkenyl, alkoxy, or cycloalkyl) and -NR³⁰R^{30'} (where R³⁰ and R^{30'} are independently hydrogen, alkyl, or hydroxyalkyl), and -C(O)NHR³³ (where R³³ is alkyl, alkenyl, alkynyl, or cycloalkyl).


[0180] In some variations, the MEK inhibitor compound of the formula (I) is a compound of the Group A, having the formula I(a) or I(b):





or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined for the formula (I), Group A, or as defined in WO 2007/044515 A1, incorporated herein by reference.

[0181] In some variations, the MEK inhibitor compound of the formula (I) is a compound of the Group B, having the formula I(c), I(d), I(e), I(f), I(g), I(h), I(i), I(j), I(k), I(m), I(n), I(o), I(p), I(q), I(r), I(s), I(u), I(v), I(w), I(x), I(cc) or I(dd):











or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined for the formula (I), Group B, or as defined in WO 2007/044515 A1, incorporated herein by reference.

[0182] In some variations, the MEK inhibitor compound of the formula (I) is a compound of the Group C, having the formula I(y) or I(z):



or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined for the formula (I), Group C, or as defined in WO 2007/044515 A1, incorporated herein by reference.

[0183] In some variations, the MEK inhibitor compound of the formula (I) is a compound of the Group D, having the formula I(aa) or I(bb):




or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined for the formula (I), Group D, or as defined in WO 2007/044515 A1, incorporated herein by reference.

[0184] In some embodiments, the MEK inhibitor compound of the formula (I) is a compound selected from the compound Nos. 1-362 as listed in WO 2007/044515 A1, Table 1 on pages 71-144 (herein collectively referred to as the Formula I Species), or a pharmaceutically acceptable salt or solvate thereof.

[0185] Also embraced are any variations of formula (I) as described in WO 2007/044515 A1, which is incorporated herein by reference. Compounds of the formula (I) or any variations thereof can be synthesized using methods known in the art, for example, the synthetic methods described in WO 2007/044515 A1, incorporated herein by reference.

[0186] Unless defined otherwise herein, the terms used in describing compounds of the formula (I) should be understood to have the same meaning as defined in WO 2007/044515 A1.

[0187] In some embodiments, the MEK inhibitor is a compound of formula (II):



(II)

or a pharmaceutically acceptable salt or solvate thereof, wherein:

Z¹ is CR¹ or N;

Z² is CR² or N;


Z³ is CR³ or N;

Z^4 is CR^4 or N ;

where one or two of Z^1 , Z^2 , Z^3 , and Z^4 are N ;

R^1 , R^2 , R^3 and R^4 are independently selected from H, halo, CN, CF_3 , $-OCF_3$, $-NO_2$,

$-(CR^{14}R^{15})_nC(=Y)R^{11}$, $-(CR^{14}R^{15})_nC(=Y)OR^{11}$, $-(CR^{14}R^{15})_nC(=Y)NR^{11}R^{12}$,
 $-(CR^{14}R^{15})_nNR^{11}R^{12}$, $-(CR^{14}R^{15})_nOR^{11}$, $-(CR^{14}R^{15})_nSR^{11}$, $-(CR^{14}R^{15})_nNR^{12}C(=Y)R^{11}$,
 $-(CR^{14}R^{15})_nNR^{12}C(=Y)OR^{11}$, $-(CR^{14}R^{15})_nNR^{13}C(=Y)NR^{11}R^{12}$, $-(CR^{14}R^{15})_nNR^{12}SO_2R^{11}$,
 $-(CR^{14}R^{15})_nOC(=Y)R^{11}$, $-(CR^{14}R^{15})_nOC(=Y)OR^{11}$, $-(CR^{14}R^{15})_nOC(=Y)NR^{11}R^{12}$,
 $-(CR^{14}R^{15})_nOS(O)_2(OR^{11})$, $-(CR^{14}R^{15})_nOP(=Y)(OR^{11})(OR^{12})$, $-(CR^{14}R^{15})_nOP(OR^{11})(OR^{12})$,
 $-(CR^{14}R^{15})_nS(O)R^{11}$, $-(CR^{14}R^{15})_nS(O)_2R^{11}$, $-(CR^{14}R^{15})_nS(O)_2NR^{11}R^{12}$, $-(CR^{14}R^{15})_nS(O)(OR^{11})$,
 $-(CR^{14}R^{15})_nS(O)_2(OR^{11})$, $-(CR^{14}R^{15})_nSC(=Y)R^{11}$, $-(CR^{14}R^{15})_nSC(=Y)OR^{11}$,
 $-(CR^{14}R^{15})_nSC(=Y)NR^{11}R^{12}$, C_1-C_{12} alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, carbocyclyl,
heterocyclyl, aryl, and heteroaryl;



R^5 and R^6 are independently selected from H or C_1-C_{12} alkyl;

X^1 is selected from R^{11} , $-OR^{11}$, $-NR^{11}R^{12}$, $-S(O)R^{11}$, and $-S(O)_2R^{11}$; when X^1 is R^{11} or $-OR^{11}$, R^{11} or $-OR^{11}$ of X^1 and $-R^5$ are optionally taken together with the nitrogen atom to which they are attached to form a 4-7 membered saturated or unsaturated ring having 0-2 additional heteroatoms selected from O, S and N, wherein said ring is optionally substituted with one or more groups selected from halo, CN, CF_3 , $-OCF_3$, $-NO_2$, oxo, $-Si(C_1-C_6\text{ alkyl})$,
 $-(CR^{19}R^{20})_nC(=Y')R^{16}$, $-(CR^{19}R^{20})_nC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nC(=Y')NR^{16}R^{17}$,
 $-(CR^{19}R^{20})_nNR^{16}R^{17}$, $-(CR^{19}R^{20})_nOR^{16}$, $-(CR^{19}R^{20})_nSR^{16}$, $-(CR^{19}R^{20})_nNR^{16}C(=Y')R^{17}$,
 $-(CR^{19}R^{20})_nNR^{16}C(=Y')OR^{17}$, $-(CR^{19}R^{20})_nNR^{18}C(=Y')NR^{16}R^{17}$, $-(CR^{19}R^{20})_nNR^{17}SO_2R^{16}$,
 $-(CR^{19}R^{20})_nOC(=Y')R^{16}$, $-(CR^{19}R^{20})_nOC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nOC(=Y')NR^{16}R^{17}$,
 $-(CR^{19}R^{20})_nOS(O)_2(OR^{16})$, $-(CR^{19}R^{20})_nOP(=Y')(OR^{16})(OR^{17})$, $-(CR^{19}R^{20})_nOP(OR^{16})(OR^{17})$,
 $-(CR^{19}R^{20})_nS(O)R^{16}$, $-(CR^{19}R^{20})_nS(O)_2R^{16}$, $-(CR^{19}R^{20})_nS(O)_2NR^{16}R^{17}$, $-(CR^{19}R^{20})_nS(O)(OR^{16})$,
 $-(CR^{19}R^{20})_nS(O)_2(OR^{16})$, $-(CR^{19}R^{20})_nSC(=Y')R^{16}$, $-(CR^{19}R^{20})_nSC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nSC(=Y')NR^{16}R^{17}$, and R^{21} ;

X^2 is selected from carbocyclyl, heterocyclyl, aryl, and heteroaryl;

R^{11} , R^{12} and R^{13} are independently H, C_1-C_{12} alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl, or R^{11} and R^{12} together with the nitrogen to which they are attached form a 3-8 membered saturated, unsaturated or aromatic ring having 0-2 heteroatoms selected from O, S and N, wherein said ring is optionally substituted with one or more groups selected from halo, CN, CF_3 , $-OCF_3$, $-NO_2$, C_1-C_6 alkyl, $-OH$, $-SH$, $-O(C_1-C_6$ alkyl), $-S(C_1-C_6$ alkyl), $-NH_2$, $-NH(C_1-C_6$ alkyl), $-N(C_1-C_6$ alkyl)₂, $-SO_2(C_1-C_6$ alkyl), $-CO_2H$, $-CO_2(C_1-C_6$ alkyl), $-C(O)NH_2$, $-C(O)NH(C_1-C_6$ alkyl), $-C(O)N(C_1-C_6$ alkyl)₂, $-N(C_1-C_6$ alkyl) $C(O)(C_1-C_6$ alkyl), $-NHC(O)(C_1-C_6$ alkyl), $-NHSO_2(C_1-C_6$ alkyl), $-N(C_1-C_6$ alkyl) $SO_2(C_1-C_6$ alkyl), $-SO_2NH_2$, $-SO_2NH(C_1-C_6$ alkyl), $-SO_2N(C_1-C_6$ alkyl)₂, $-OC(O)NH_2$, $-OC(O)NH(C_1-C_6$ alkyl), $-OC(O)N(C_1-C_6$ alkyl)₂, $-OC(O)O(C_1-C_6$ alkyl), $-NHC(O)NH(C_1-C_6$ alkyl), $-NHC(O)N(C_1-C_6$ alkyl)₂, $-N(C_1-C_6$ alkyl) $C(O)NH(C_1-C_6$ alkyl), $-N(C_1-C_6$ alkyl) $C(O)N(C_1-C_6$ alkyl)₂, $-NHC(O)NH(C_1-C_6$ alkyl), $-NHC(O)N(C_1-C_6$ alkyl)₂, $-NHC(O)O(C_1-C_6$ alkyl), and $-N(C_1-C_6$ alkyl) $C(O)O(C_1-C_6$ alkyl);

R^{14} and R^{15} are independently selected from H, C_1-C_{12} alkyl, aryl, carbocyclyl, heterocyclyl, and heteroaryl;

m and n are independently selected from 0, 1, 2, 3, 4, 5, or 6;

Y is independently O, NR^{11} , or S;

wherein each said alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl and heteroaryl of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , X^1 , X^2 , R^{11} , R^{12} , R^{13} , R^{14} , and R^{15} is independently optionally substituted with one or more groups independently selected from halo, CN, CF_3 , $-OCF_3$, $-NO_2$, oxo, $-Si(C_1-C_6$ alkyl), $-(CR^{19}R^{20})_nC(=Y')R^{16}$, $-(CR^{19}R^{20})_nC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nC(=Y')NR^{16}R^{17}$, $-(CR^{19}R^{20})_nNR^{16}R^{17}$, $-(CR^{19}R^{20})_nOR^{16}$, $-(CR^{19}R^{20})_nSR^{16}$, $-(CR^{19}R^{20})_nNR^{16}C(=Y')R^{17}$, $-(CR^{19}R^{20})_nNR^{16}C(=Y')OR^{17}$, $-(CR^{19}R^{20})_nNR^{18}C(=Y')NR^{16}R^{17}$, $-(CR^{19}R^{20})_nNR^{17}SO_2R^{16}$, $-(CR^{19}R^{20})_nOC(=Y')R^{16}$, $-(CR^{19}R^{20})_nOC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nOC(=Y')NR^{16}R^{17}$, $-(CR^{19}R^{20})_nOS(O)_2(OR^{16})$, $-(CR^{19}R^{20})_nOP(=Y')(OR^{16})(OR^{17})$, $-(CR^{19}R^{20})_nOP(OR^{16})(OR^{17})$, $-(CR^{19}R^{20})_nS(O)R^{16}$, $-(CR^{19}R^{20})_nS(O)_2R^{16}$, $-(CR^{19}R^{20})_nS(O)_2NR^{16}R^{17}$, $-(CR^{19}R^{20})_nS(O)(OR^{16})$, $-(CR^{19}R^{20})_nS(O)_2(OR^{16})$, $-(CR^{19}R^{20})_nSC(=Y')R^{16}$, $-(CR^{19}R^{20})_nSC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nSC(=Y')NR^{16}R^{17}$, and R^{21} ;

each R^{16} , R^{17} and R^{18} is independently H, C_1-C_{12} alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl, wherein said alkyl, alkenyl, alkynyl, carbocyclyl,

heterocyclyl, aryl, or heteroaryl is optionally substituted with one or more groups selected from halo, oxo, CN, $-\text{OCF}_3$, CF_3 , $-\text{NO}_2$, $\text{C}_1\text{-C}_6$ alkyl, $-\text{OH}$, $-\text{SH}$, $-\text{O}(\text{C}_1\text{-C}_6$ alkyl), $-\text{S}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NH}_2$, $-\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{SO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{CO}_2\text{H}$, $-\text{CO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHSO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{SO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{SO}_2\text{NH}_2$, $-\text{SO}_2\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{SO}_2\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{OC}(\text{O})\text{NH}_2$, $-\text{OC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{OC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{OC}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{NHC}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl), and $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl);

or R^{16} and R^{17} together with the nitrogen to which they are attached form a 3-8 membered saturated, unsaturated or aromatic ring having 0-2 heteroatoms selected from O, S and N, wherein said ring is optionally substituted with one or more groups selected from halo, CN, $-\text{OCF}_3$, CF_3 , $-\text{NO}_2$, $\text{C}_1\text{-C}_6$ alkyl, $-\text{OH}$, $-\text{SH}$, $-\text{O}(\text{C}_1\text{-C}_6$ alkyl), $-\text{S}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NH}_2$, $-\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{SO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{CO}_2\text{H}$, $-\text{CO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHSO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{SO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{SO}_2\text{NH}_2$, $-\text{SO}_2\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{SO}_2\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{OC}(\text{O})\text{NH}_2$, $-\text{OC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{OC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{OC}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{NHC}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl), and $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl);

R^{19} and R^{20} are independently selected from H, $\text{C}_1\text{-C}_{12}$ alkyl, $-(\text{CH}_2)_n$ -aryl, $-(\text{CH}_2)_n$ -carbocyclyl, $-(\text{CH}_2)_n$ -heterocyclyl, and $-(\text{CH}_2)_n$ -heteroaryl;


R^{21} is $\text{C}_1\text{-C}_{12}$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, $\text{C}_2\text{-C}_8$ alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl, wherein each member of R^{21} is optionally substituted with one or more groups selected from halo, CN, $-\text{OCF}_3$, CF_3 , $-\text{NO}_2$, $\text{C}_1\text{-C}_6$ alkyl, $-\text{OH}$, $-\text{SH}$, $-\text{O}(\text{C}_1\text{-C}_6$ alkyl), $-\text{S}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NH}_2$, $-\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{SO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{CO}_2\text{H}$, $-\text{CO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHSO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{SO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{SO}_2\text{NH}_2$, $-\text{SO}_2\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{SO}_2\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{OC}(\text{O})\text{NH}_2$, $-\text{OC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl),


$-\text{OC(O)N(C}_1\text{-C}_6\text{ alkyl})_2$, $-\text{OC(O)O(C}_1\text{-C}_6\text{ alkyl)}$, $-\text{NHC(O)NH(C}_1\text{-C}_6\text{ alkyl)}$, $-\text{NHC(O)N(C}_1\text{-C}_6\text{ alkyl})_2$, $-\text{N(C}_1\text{-C}_6\text{ alkyl)C(O)NH(C}_1\text{-C}_6\text{ alkyl)}$, $-\text{N(C}_1\text{-C}_6\text{ alkyl)C(O)N(C}_1\text{-C}_6\text{ alkyl})_2$, $-\text{NHC(O)NH(C}_1\text{-C}_6\text{ alkyl)}$, $-\text{NHC(O)N(C}_1\text{-C}_6\text{ alkyl})_2$, $-\text{NHC(O)O(C}_1\text{-C}_6\text{ alkyl)}$, and $-\text{N(C}_1\text{-C}_6\text{ alkyl)C(O)O(C}_1\text{-C}_6\text{ alkyl)}$;

each Y' is independently O , NR^{22} , or S ; and

R^{22} is H or $\text{C}_1\text{-C}_{12}$ alkyl.

[0188] In some variations, the MEK inhibitor compound of the formula (II) is a compound of the formula (II-1-a), (II-1-b), (II-1-c), (II-1-d), (II-1-e), (II-1-f), (II-1-g), (II-1-h), (II-1-i), (II-2-a), (II-2-b), (II-2-c), (II-2-d), (II-2-e), (II-2-f), (II-2-g), (II-2-h), (II-2-i), (II-3-a), (II-3-b), (II-3-c), (II-3-d), (II-3-e), (II-3-f), (II-3-g), (II-3-h), or (II-3-i):






or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined for the formula (II) or as defined in WO 2008/024725 A1, incorporated herein by reference.

[0189] In some embodiments, the MEK inhibitor compound of the formula (II) is a compound selected from the compounds of Examples 5-18, 20-102, 105-109, 111-118, 120-133, 136-149 and 151-160 in WO 2008/024725 A1 (herein collectively referred to as the Formula II Species), or a pharmaceutically acceptable salt or solvate thereof. These compounds exhibited an IC₅₀ of

less than 10 μ M in the assay described either in Example 8a or 8b (MEK activity assays). Most of these compounds exhibited an IC_{50} of less than 5 μ M. See page 62 in WO 2008/024725 A1.

[0190] Also embraced are MEK inhibitor compounds (and/or solvates and salts thereof) described in WO 2008/024725 A1, which is incorporated herein by reference, for example, aza-benzofuran compounds of the formula (II) (designated as formula I in WO 2008/024725 A1, e.g., on page 3) and variations thereof as described in WO 2008/024725 A1. Compounds of formula (II) can be synthesized using methods known in the art, for example, the synthetic methods described in WO 2008/024725 A1, incorporated herein by reference.

[0191] In some embodiments, the MEK inhibitor is a compound of formula (III):



(III)

or a pharmaceutically acceptable salt or solvate thereof, wherein:

Z^1 is CR^1 or N ;

R^1 is H , C_1 - C_3 alkyl, halo, CF_3 , CHF_2 , CN , OR^A or $NR^A R^A$;

R^1' is H , C_1 - C_3 alkyl, halo, CF_3 , CHF_2 , CN , OR^A , or $NR^A R^A$;

wherein each R^A is independently H or C_1 - C_3 alkyl;

Z^2 is CR^2 or N ;

Z^3 is CR^3 or N ; provided that only one of Z^1 , Z^2 and Z^3 can be N at the same time;

R^2 and R^3 are independently selected from H , halo, CN , CF_3 , $-OCF_3$, $-NO_2$,

$-(CR^{14}R^{15})_nC(=Y')R^{11}$, $-(CR^{14}R^{15})_nC(=Y')OR^{11}$, $-(CR^{14}R^{15})_nC(=Y')NR^{11}R^{12}$,

$-(CR^{14}R^{15})_nNR^{11}R^{12}$, $-(CR^{14}R^{15})_nOR^{11}$, $-(CR^{14}R^{15})_nSR^{11}$, $-(CR^{14}R^{15})_nNR^{12}C(=Y')R^{11}$,

$-(CR^{14}R^{15})_nNR^{12}C(=Y')OR^{11}$, $-(CR^{14}R^{15})_nNR^{13}C(=Y')NR^{11}R^{12}$, $-(CR^{14}R^{15})_nNR^{12}SO_2R^{11}$,

$-(CR^{14}R^{15})_nOC(=Y')R^{11}$, $-(CR^{14}R^{15})_nOC(=Y')OR^{11}$, $-(CR^{14}R^{15})_nOC(=Y')NR^{11}R^{12}$,

$-(CR^{14}R^{15})_nOS(O)_2(OR^{11})$, $-(CR^{14}R^{15})_nOP(=Y')(OR^{11})(OR^{12})$, $-(CR^{14}R^{15})_nOP(OR^{11})(OR^{12})$,


$-(CR^{14}R^{15})_nS(O)R^{11}$, $-(CR^{14}R^{15})_nS(O)_2R^{11}$, $-(CR^{14}R^{15})_nS(O)_2NR^{11}R^{12}$, $-(CR^{14}R^{15})_nS(O)(OR^{11})$,

$-(CR^{14}R^{15})_nS(O)_2(OR^{11})$, $-(CR^{14}R^{15})_nSC(=Y')R^{11}$, $-(CR^{14}R^{15})_nSC(=Y')OR^{11}$,

$-(CR^{14}R^{15})_nSC(=Y')NR^{11}R^{12}$, C₁–C₁₂ alkyl, C₂–C₈ alkenyl, C₂–C₈ alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl;

R⁴ is H, C₁–C₆ alkyl or C₃–C₄ carbocyclyl;

Y is W-C(O)- or W';



R⁵ is H or C₁–C₁₂ alkyl;

X¹ is selected from R^{11'} and –OR^{11'}; when X¹ is R^{11'}, X¹ is optionally taken together with R⁵ and the nitrogen atom to which they are bound to form a 4–7 membered saturated or unsaturated ring having 0–2 additional heteroatoms selected from O, S and N, wherein said ring is optionally substituted with one or more groups selected from halo, CN, CF₃, –OCF₃, –NO₂, oxo, $-(CR^{19}R^{20})_nC(=Y')R^{16}$, $-(CR^{19}R^{20})_nC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nC(=Y')NR^{16}R^{17}$, $-(CR^{19}R^{20})_nNR^{16}R^{17}$, $-(CR^{19}R^{20})_nOR^{16}$, $-(CR^{19}R^{20})_nSR^{16}$, $-(CR^{19}R^{20})_nNR^{16}C(=Y')R^{17}$, $-(CR^{19}R^{20})_nNR^{16}C(=Y')OR^{17}$, $-(CR^{19}R^{20})_nNR^{18}C(=Y')NR^{16}R^{17}$, $-(CR^{19}R^{20})_nNR^{17}SO_2R^{16}$, $-(CR^{19}R^{20})_nOC(=Y')R^{16}$, $-(CR^{19}R^{20})_nOC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nOC(=Y')NR^{16}R^{17}$, $-(CR^{19}R^{20})_nOS(O)_2(OR^{16})$, $-(CR^{19}R^{20})_nOP(=Y')(OR^{16})(OR^{17})$, $-(CR^{19}R^{20})_nOP(OR^{16})(OR^{17})$, $-(CR^{19}R^{20})_nS(O)R^{16}$, $-(CR^{19}R^{20})_nS(O)_2R^{16}$, $-(CR^{19}R^{20})_nS(O)_2NR^{16}R^{17}$, $-(CR^{19}R^{20})_nS(O)(OR^{16})$, $-(CR^{19}R^{20})_nS(O)_2(OR^{16})$, $-(CR^{19}R^{20})_nSC(=Y')R^{16}$, $-(CR^{19}R^{20})_nSC(=Y')OR^{16}$, $-(CR^{19}R^{20})_nSC(=Y')NR^{16}R^{17}$, and R²¹;


each R^{11'} is independently H, C₁–C₁₂ alkyl, C₂–C₈ alkenyl, C₂–C₈ alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl;

R¹¹, R¹² and R¹³ are independently H, C₁–C₁₂ alkyl, C₂–C₈ alkenyl, C₂–C₈ alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl,

or R¹¹ and R¹² together with the nitrogen to which they are attached form a 3–8 membered saturated, unsaturated or aromatic ring having 0–2 heteroatoms selected from O, S and N, wherein said ring is optionally substituted with one or more groups selected from halo, CN, CF₃, –OCF₃, –NO₂, C₁–C₆ alkyl, –OH, –SH, –O(C₁–C₆ alkyl), –S(C₁–C₆ alkyl), –NH₂, –NH(C₁–C₆ alkyl), –N(C₁–C₆ alkyl)₂, –SO₂(C₁–C₆ alkyl), –CO₂H, –CO₂(C₁–C₆ alkyl), –C(O)NH₂, –C(O)NH(C₁–C₆ alkyl), –C(O)N(C₁–C₆ alkyl)₂, –N(C₁–C₆ alkyl)C(O)(C₁–C₆ alkyl), –NHC(O)(C₁–C₆ alkyl), –NHSO₂(C₁–C₆ alkyl), –N(C₁–C₆ alkyl)SO₂(C₁–C₆ alkyl), –SO₂NH₂,

—SO₂NH(C₁—C₆ alkyl), —SO₂N(C₁—C₆ alkyl)₂, —OC(O)NH₂, —OC(O)NH(C₁—C₆ alkyl), —OC(O)N(C₁—C₆ alkyl)₂, —OC(O)O(C₁—C₆ alkyl), —NHC(O)NH(C₁—C₆ alkyl), —NHC(O)N(C₁—C₆ alkyl)₂, —N(C₁—C₆ alkyl)C(O)NH(C₁—C₆ alkyl), —N(C₁—C₆ alkyl)C(O)N(C₁—C₆ alkyl)₂, —NHC(O)NH(C₁—C₆ alkyl), —NHC(O)N(C₁—C₆ alkyl)₂, —NHC(O)O(C₁—C₆ alkyl), and —N(C₁—C₆ alkyl)C(O)O(C₁—C₆ alkyl);

R¹⁴ and R¹⁵ are independently selected from H, C₁—C₁₂ alkyl, aryl, carbocyclyl, heterocyclyl, and heteroaryl;




each R⁷ is independently selected from H, halo, CN, CF₃, —OCF₃, —NO₂, —(CR¹⁴R¹⁵)_nC(=Y')R¹¹, —(CR¹⁴R¹⁵)_nC(=Y')OR¹¹, —(CR¹⁴R¹⁵)_nC(=Y')NR¹¹R¹², —(CR¹⁴R¹⁵)_nNR¹¹R¹², —(CR¹⁴R¹⁵)_nOR¹¹, —(CR¹⁴R¹⁵)_nSR¹¹, —(CR¹⁴R¹⁵)_nNR¹²C(=Y')R¹¹, —(CR¹⁴R¹⁵)_nNR¹²C(=Y')OR¹¹, —(CR¹⁴R¹⁵)_nNR¹³C(=Y')NR¹¹R¹², —(CR¹⁴R¹⁵)_nNR¹²SO₂R¹¹, —(CR¹⁴R¹⁵)_nOC(=Y')R¹¹, —(CR¹⁴R¹⁵)_nOC(=Y')OR¹¹, —(CR¹⁴R¹⁵)_nOC(=Y')NR¹¹R¹², —(CR¹⁴R¹⁵)_nOS(O)₂(OR¹¹), —(CR¹⁴R¹⁵)_nOP(=Y')(OR¹¹)(OR¹²), —(CR¹⁴R¹⁵)_nOP(OR¹¹)(OR¹²), —(CR¹⁴R¹⁵)_nS(O)R¹¹, —(CR¹⁴R¹⁵)_nS(O)₂R¹¹, —(CR¹⁴R¹⁵)_nS(O)₂NR¹¹R¹², —(CR¹⁴R¹⁵)_nS(O)(OR¹¹), —(CR¹⁴R¹⁵)_nS(O)₂(OR¹¹), —(CR¹⁴R¹⁵)_nSC(=Y')R¹¹, —(CR¹⁴R¹⁵)_nSC(=Y')OR¹¹, —(CR¹⁴R¹⁵)_nSC(=Y')NR¹¹R¹², C₁—C₁₂ alkyl, C₂—C₈ alkenyl, C₂—C₈ alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl;

each R⁸ is independently selected from C₁–C₁₂ alkyl, aryl, carbocyclyl, heterocyclyl, and heteroaryl;

R⁹ is selected from H, -(CR¹⁴R¹⁵)_nC(=Y')R¹¹, -(CR¹⁴R¹⁵)_nC(=Y')OR¹¹, -(CR¹⁴R¹⁵)_nC(=Y')NR¹¹R¹², -(CR¹⁴R¹⁵)_qNR¹¹R¹², -(CR¹⁴R¹⁵)_qOR¹¹, -(CR¹⁴R¹⁵)_qSR¹¹, -(CR¹⁴R¹⁵)_qNR¹²C(=Y')R¹¹, -(CR¹⁴R¹⁵)_qNR¹²C(=Y')OR¹¹, -(CR¹⁴R¹⁵)_qNR¹³C(=Y')NR¹¹R¹², -(CR¹⁴R¹⁵)_qNR¹²SO₂R¹¹, -(CR¹⁴R¹⁵)_qOC(=Y')R¹¹, -(CR¹⁴R¹⁵)_qOC(=Y')OR¹¹, -(CR¹⁴R¹⁵)_qOC(=Y')NR¹¹R¹², -(CR¹⁴R¹⁵)_qOS(O)₂(OR¹¹), -(CR¹⁴R¹⁵)_qOP(=Y')(OR¹¹)(OR¹²), -(CR¹⁴R¹⁵)_qOP(OR¹¹)(OR¹²), -(CR¹⁴R¹⁵)_nS(O)R¹¹, -(CR¹⁴R¹⁵)_nS(O)₂R¹¹, -(CR¹⁴R¹⁵)_nS(O)₂NR¹¹R¹², C₁–C₁₂ alkyl, C₂–C₈ alkenyl, C₂–C₈ alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl;

R¹⁰ is H, C₁–C₆ alkyl or C₃–C₄ carbocyclyl;



R⁶ is H, halo, C₁–C₆ alkyl, C₂–C₈ alkenyl, C₂–C₈ alkynyl, carbocyclyl, heteroaryl, heterocyclyl, -OCF₃, -NO₂, -Si(C₁–C₆ alkyl), -(CR¹⁹R²⁰)_nNR¹⁶R¹⁷, -(CR¹⁹R²⁰)_nOR¹⁶, or -(CR¹⁹R²⁰)_nSR¹⁶;

R^{6'} is H, halo, C₁–C₆ alkyl, carbocyclyl, CF₃, -OCF₃, -NO₂, -Si(C₁–C₆ alkyl), -(CR¹⁹R²⁰)_nNR¹⁶R¹⁷, -(CR¹⁹R²⁰)_nOR¹⁶, -(CR¹⁹R²⁰)_nSR¹⁶, C₂–C₈ alkenyl, C₂–C₈ alkynyl, heterocyclyl, aryl, or heteroaryl;

p is 0, 1, 2 or 3;

n is 0, 1, 2 or 3;

q is 2 or 3;

wherein each said alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl and heteroaryl of R¹, R², R³, R⁴, R⁵, R⁶, R^{6'}, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R^{11'}, R¹², R¹³, R¹⁴, R¹⁵ and R^A is independently optionally substituted with one or more groups independently selected from halo, CN, CF₃, -OCF₃, -NO₂, oxo, -Si(C₁–C₆ alkyl), -(CR¹⁹R²⁰)_nC(=Y')R¹⁶, -(CR¹⁹R²⁰)_nC(=Y')OR¹⁶, -(CR¹⁹R²⁰)_nC(=Y')NR¹⁶R¹⁷, -(CR¹⁹R²⁰)_nNR¹⁶R¹⁷, -(CR¹⁹R²⁰)_nOR¹⁶, -(CR¹⁹R²⁰)_nSR¹⁶, -(CR¹⁹R²⁰)_nNR¹⁶C(=Y')R¹⁷, -(CR¹⁹R²⁰)_nNR¹⁶C(=Y')OR¹⁷, -(CR¹⁹R²⁰)_nNR¹⁸C(=Y')NR¹⁶R¹⁷, -(CR¹⁹R²⁰)_nNR¹⁷SO₂R¹⁶, -(CR¹⁹R²⁰)_nOC(=Y')R¹⁶, -(CR¹⁹R²⁰)_nOC(=Y')OR¹⁶, -(CR¹⁹R²⁰)_nOC(=Y')NR¹⁶R¹⁷, -(CR¹⁹R²⁰)_nOS(O)₂(OR¹⁶), -(CR¹⁹R²⁰)_nOP(=Y')(OR¹⁶)(OR¹⁷),

-(CR¹⁹R²⁰)_nOP(OR¹⁶)(OR¹⁷), -(CR¹⁹R²⁰)_nS(O)R¹⁶, -(CR¹⁹R²⁰)_nS(O)₂R¹⁶,
 -(CR¹⁹R²⁰)_nS(O)₂NR¹⁶R¹⁷, -(CR¹⁹R²⁰)_nS(O)(OR¹⁶), -(CR¹⁹R²⁰)_nS(O)₂(OR¹⁶),
 -(CR¹⁹R²⁰)_nSC(=Y')R¹⁶, -(CR¹⁹R²⁰)_nSC(=Y')OR¹⁶, -(CR¹⁹R²⁰)_nSC(=Y')NR¹⁶R¹⁷, and R²¹;
 each R¹⁶, R¹⁷ and R¹⁸ is independently H, C₁-C₁₂ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl, wherein said alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with one or more groups selected from halo, CN, -OCF₃, CF₃, -NO₂, C₁-C₆ alkyl, -OH, -SH, -O(C₁-C₆ alkyl), -S(C₁-C₆ alkyl), -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -SO₂(C₁-C₆ alkyl), -CO₂H, -CO₂(C₁-C₆ alkyl), -C(O)NH₂, -C(O)NH(C₁-C₆ alkyl), -C(O)N(C₁-C₆ alkyl)₂, -N(C₁-C₆ alkyl)C(O)(C₁-C₆ alkyl), -NHC(O)(C₁-C₆ alkyl), -NHSO₂(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)SO₂(C₁-C₆ alkyl), -SO₂NH₂, -SO₂NH(C₁-C₆ alkyl), -SO₂N(C₁-C₆ alkyl)₂, -OC(O)NH₂, -OC(O)NH(C₁-C₆ alkyl), -OC(O)N(C₁-C₆ alkyl)₂, -OC(O)O(C₁-C₆ alkyl), -NHC(O)NH(C₁-C₆ alkyl), -NHC(O)N(C₁-C₆ alkyl)₂, -N(C₁-C₆ alkyl)C(O)NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)C(O)N(C₁-C₆ alkyl)₂, -NHC(O)NH(C₁-C₆ alkyl), -NHC(O)N(C₁-C₆ alkyl)₂, -NHC(O)O(C₁-C₆ alkyl), and -N(C₁-C₆ alkyl)C(O)O(C₁-C₆ alkyl);



or R¹⁶ and R¹⁷ together with the nitrogen to which they are attached form a 3-8 membered saturated, unsaturated or aromatic ring having 0-2 heteroatoms selected from O, S and N, wherein said ring is optionally substituted with one or more groups selected from halo, CN, -OCF₃, CF₃, -NO₂, C₁-C₆ alkyl, -OH, -SH, -O(C₁-C₆ alkyl), -S(C₁-C₆ alkyl), -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -SO₂(C₁-C₆ alkyl), -CO₂H, -CO₂(C₁-C₆ alkyl), -C(O)NH₂, -C(O)NH(C₁-C₆ alkyl), -C(O)N(C₁-C₆ alkyl)₂, -N(C₁-C₆ alkyl)C(O)(C₁-C₆ alkyl), -NHC(O)(C₁-C₆ alkyl), -NHSO₂(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)SO₂(C₁-C₆ alkyl), -SO₂NH₂, -SO₂NH(C₁-C₆ alkyl), -SO₂N(C₁-C₆ alkyl)₂, -OC(O)NH₂, -OC(O)NH(C₁-C₆ alkyl), -OC(O)N(C₁-C₆ alkyl)₂, -OC(O)O(C₁-C₆ alkyl), -NHC(O)NH(C₁-C₆ alkyl), -NHC(O)N(C₁-C₆ alkyl)₂, -N(C₁-C₆ alkyl)C(O)NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)C(O)N(C₁-C₆ alkyl)₂, -NHC(O)NH(C₁-C₆ alkyl), -NHC(O)N(C₁-C₆ alkyl)₂, -NHC(O)O(C₁-C₆ alkyl), and -N(C₁-C₆ alkyl)C(O)O(C₁-C₆ alkyl);

R¹⁹ and R²⁰ are independently selected from H, C₁-C₁₂ alkyl, -(CH₂)_n-aryl, -(CH₂)_n-carbocyclyl, -(CH₂)_n-heterocyclyl, and -(CH₂)_n-heteroaryl;

R²¹ is C₁-C₁₂ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl, wherein each member of R²¹ is optionally substituted with one or more groups

selected from halo, oxo, CN, $-\text{OCF}_3$, CF_3 , $-\text{NO}_2$, $\text{C}_1\text{-C}_6$ alkyl, $-\text{OH}$, $-\text{SH}$, $-\text{O}(\text{C}_1\text{-C}_6$ alkyl), $-\text{S}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NH}_2$, $-\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{SO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{CO}_2\text{H}$, $-\text{CO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHSO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{SO}_2(\text{C}_1\text{-C}_6$ alkyl), $-\text{SO}_2\text{NH}_2$, $-\text{SO}_2\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{SO}_2\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{OC}(\text{O})\text{NH}_2$, $-\text{OC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{OC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{OC}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$ alkyl), $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_6$ alkyl)₂, $-\text{NHC}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl), and $-\text{N}(\text{C}_1\text{-C}_6$ alkyl) $\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_6$ alkyl); each Y' is independently O, NR²², or S; and R²² is H or C₁-C₁₂ alkyl.

[0192] In some variations, the MEK inhibitor compound of the formula (III) has the formula (III-a) or (III-b):



or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined for the formula (III) or as defined in WO 2009/085983 A1, incorporated herein by reference.

[0193] In some embodiments, the MEK inhibitor compound of the formula (III) is a compound selected from the compounds listed in Table 1, or a pharmaceutically acceptable salt or solvate thereof.

Table 1

Compound No.	Chemical Name	Structure
(III)-5	5-(2-Fluoro-4-iodophenylamino)-imidazo[1,5-a]pyridine-6-carboxylic acid (2-hydroxyethoxy)-amide	

Compound No.	Chemical Name	Structure
(III)-6	5-(2-Fluoro-4-iodo-phenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid ((<i>R</i>)-2,3-dihydroxy-propoxy)-amide	
(III)-7	5-(2-Fluoro-4-iodo-phenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid ((<i>S</i>)-2-hydroxy-propoxy)-amide	
(III)-8	5-(4-Bromo-2-fluorophenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid (2-hydroxyethoxy)-amide	
(III)-9	5-(4-Bromo-2-fluoro-phenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid ((<i>S</i>)-2-hydroxy-propoxy)-amide	
(III)-10	5-(4-Bromo-2-fluoro-phenylamino)-8-fluoro-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid ((<i>S</i>)-2-hydroxy-propoxy)-amide	
(III)-11	8-Fluoro-5-(2-fluoro-4-iodo-phenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid (2-hydroxy-ethoxy)-amide	
(III)-12	8-Fluoro-5-(2-fluoro-4-iodo-phenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid ((<i>R</i>)-2,3-dihydroxy-propoxy)-amide	


Compound No.	Chemical Name	Structure
(III)-13	8-Fluoro-5-(2-fluoro-4-iodophenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid ((S)-2-hydroxy-propoxy)-amide	
(III)-14	5-(2-Fluoro-methanesulfanyl-phenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid (2-hydroxy-ethoxy)-amide	
(III)-15	5-(2-Fluoro-4-iodophenylamino)-imidazo[1,5- <i>a</i>]pyrazine-6-carboxylic acid (2-hydroxy-ethoxy)-amide	
(III)-16	5-(2-Fluoro-4-iodophenylamino)-imidazo[1,5- <i>a</i>]pyrazine-6-carboxylic acid ((S)-2-hydroxy-propoxy)-amide	
(III)-17	5-(4-Cyclopropyl-2-fluoro-phenylamino)-imidazo[1,5- <i>a</i>]pyridine-6-carboxylic acid (2-hydroxy-ethoxy)-amide	
(III)-18	(R)- <i>N</i> -(2,3-Dihydroxypropoxy)-5-(2-fluoro-4-iodophenylamino)imidazo[1,5- <i>a</i>]pyrazine-6-carboxamide	
(III)-19	<i>N</i> -Ethoxy-5-(2-fluoro-4-iodophenylamino)imidazo[1,5- <i>a</i>]pyrazine-6-carboxamide	

Compound No.	Chemical Name	Structure
(III)-20	<i>N</i> -(Cyclopropylmethoxy)-5-(2-fluoro-4-iodophenylamino)imidazo[1,5- <i>a</i>]pyrazine-6-carboxamide	
(III)-21	5-(2-Fluoro-4-iodophenylamino)- <i>N</i> -methylimidazo[1,5- <i>a</i>]pyrazine-6-carboxamide	
(III)-22	5-(4-Bromo-2-fluorophenylamino)- <i>N</i> -(2-hydroxy-ethoxy)imidazo[1,5- <i>a</i>]pyrazine-6-carboxamide	
(III)-23	(S)-5-(4-Bromo-2-fluorophenylamino)- <i>N</i> -(2-hydroxy-propoxy)imidazo[1,5- <i>a</i>]pyrazine-6-carboxamide	
(III)-24	(R)-5-(4-Bromo-2-fluorophenylamino)- <i>N</i> -(2,3-dihydroxy-propoxy)imidazo[1,5- <i>a</i>]pyrazine-6-carboxamide	
(III)-25	5-(4-Bromo-2-fluorophenylamino)- <i>N</i> -(cyclopropylmethoxy)imidazo[1,5- <i>a</i>]pyrazine-6-carboxamide	

[0194] Compounds in Table 1 correspond to Examples 5-25 in WO 2009/085983 A1. Compounds (III)-5 – (III)-20 and (III)-22 – (III)-24 exhibited an IC₅₀ of less than 0.5 μ M in the assay described in Example 8b (MEK activity assay). Some of these compounds exhibited an IC₅₀ of less than 0.1 μ M. Compounds (III)-21 and (III)-25 exhibited an IC₅₀ of less than 10 μ M. See page 49 in WO 2009/085983 A1.

[0195] Also embraced are MEK inhibitor compounds (and/or solvates and salts thereof) described in WO 2009/085983 A1, which is incorporated herein by reference, for example, imidazopyridine compounds of the formula (III) (designated as formula I in WO 2009/085983 A1, e.g., on page 3) and variations thereof as described in WO 2009/085983 A1. Compounds of formula (III) can be synthesized using methods known in the art, for example, the synthetic methods described in WO 2009/085983 A1, incorporated herein by reference.


[0196] In some embodiments, the MEK inhibitor is a compound of formula (IV),




IV

or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined in WO 03/077914 A1 for the formula I on pages 4-9 or any applicable variations described in WO 03/077914 A1, incorporated herein by reference.

[0197] In some variations, the MEK inhibitor compound of the formula (IV) is a compound of the formula (IV-a), (IV-b), (IV-c), or (IV-d):





or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined in WO 03/077914 A1 for the formulae II, III, IIIa and IIIb, respectively on pages 10-13 or any applicable variations described in WO 03/077914 A1, incorporated herein by reference.

[0198] In some embodiments, the MEK inhibitor compound of the formula (IV) is a compound selected from the group consisting of:

7-Fluoro-6-(4-bromo-2-methyl-phenylamino)-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid cyclopropylmethoxy-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2,3-dihydroxy-propoxy)-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-pyran-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide;

[6-(5-Amino-[1,3,4]oxadiazol-2-yl)-4-fluoro-1H-benzimidazol-5-yl]-(4-bromo-2-methyl-phenyl)-amine;

1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazol-5-yl]-2-hydroxy-ethanone;

1-[6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazol-5-yl]-2-methoxy ethanone;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-1,1-dimethyl-ethoxy)-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3-(tetrahydro-furan-2-ylmethyl)-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide;

6-(4-Bromo-2-chloro-phenylamino)-7-fluoro-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide;


6-(Bromo-2-fluoro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide; and

6-(2,4-Dichloro-phenylamino)-7-fluoro-3-methyl-3H-benzoimidazole-5-carboxylic acid (2-hydroxy-ethoxy)-amide;

or a pharmaceutically acceptable salt or solvate thereof.

[0199] Also embraced are any variations of formula (IV) as described in WO 03/077914 A1, which is incorporated herein by reference. Compounds of the formula (IV) or any variations thereof can be synthesized using methods known in the art, for example, the synthetic methods described in WO 03/077914 A1, incorporated herein by reference.

[0200] In some embodiments, the MEK inhibitor is a compound of formula (V),




V

or a pharmaceutically acceptable salt or solvate thereof, wherein the variables are as defined in WO 2005/121142 A1 for the formula [I] on pages 6-10 or any applicable variations described in WO 2005/121142 A1, incorporated herein by reference.

[0201] Also embraced are any variations of formula (V) as described in WO 2005/121142 A1, such as the individual MEK inhibitor compounds described in WO 2005/121142 A1, e.g., Examples 1-1 to 1-343 in Table 1, Examples 2-1 and 2-2 in Table 2, Examples 3-1 to 3-9 in Table 3, Examples 4-1 to 4-148 in Table 4. Compounds of the formula (V) or any variations thereof can be synthesized using methods known in the art, for example, the synthetic methods described in WO 2005/121142 A1, incorporated herein by reference.

[0202] In some embodiments, the MEK inhibitor is a compound of formula (VI),



VI


or a pharmaceutically acceptable salt or ester thereof, wherein:

R1 is selected from the group consisting of bromo, iodo, ethynyl, cycloalkyl, alkoxy, azetidinyl, acetyl, heterocycyl, cyano, straight-chained alkyl and branched-chain alkyl;

R2 is selected from the group consisting of hydrogen, chlorine, fluorine, and alkyl;

R3 is selected from the group consisting of hydrogen, chlorine, and fluorine;

R4 is selected from the group consisting of hydrogen, optionally substituted aryl, alkyl, and cycloalkyl;



wherein R6 is selected from the group consisting of hydroxyl, alkoxy, cycloalkyl, optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl;

R7 and R8 are independently selected from the group consisting of hydrogen and optionally substituted alkyl;

or R6 and R7 can together form a cycloalkyl group and R8 is hydrogen.

[0203] In some variations, the MEK inhibitor compound is of the formula (VI), or a pharmaceutically acceptable salt or ester thereof, wherein the variables are as defined in WO 2007/096259 A1 for the formula I or any applicable variations described on pages 4-10 in WO 2007/096259 A1, incorporated herein by reference. Further embraced MEK inhibitors are compounds described in Examples 1-182 in WO 2007/096259 A1, incorporated herein by reference.

[0204] In some embodiments, the MEK inhibitor compound of the formula (VI) is a compound selected from the group consisting of:

(2S,3S)-N-(4-Bromo-phenyl)-2-[(R)-4-(4-methoxy-phenyl)-2,5-dioxo-imidazolidin-1-yl]-3-phenyl-butyramide;

(2S,3S)-N-(4-iodo-phenyl)-2-[(R)-4-(4-methoxy-phenyl)-2,5-dioxo-imidazolidin-1-yl]-3-phenyl-butyramide;

(2S,3S)-N-(2-Fluoro-4-iodo-phenyl)-2-[(R)-4-[4-(2-hydroxy-ethoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-3-phenyl-butyramide;

(2S,3S)-N-(4-Ethynyl-2-fluoro-phenyl)-2-[(R)-4-[4-(2-hydroxy-ethoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-3-phenyl-butyramide;

(2R,3S)-N-(4-Ethynyl-2-fluoro-phenyl)-2-[(R)-4-[4-(2-hydroxy-ethoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-3-phenyl-butyramide;

(2S,3S)-N-(2-Chloro-4-iodo-phenyl)-2-[(R)-4-[4-(2-hydroxy-ethoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-3-phenyl-butyramide;

(2S,3S)-2-[(R)-4-[4-(2-Hydroxy-ethoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-N-(4-iodo-2-methyl-phenyl)-3-phenyl-butyramide;

(2S,3S)-N-(2-Chloro-4-iodo-phenyl)-2-[(R)-4-[4-((R)-2,3-dihydroxy-propoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-3-phenyl-butyramide;

(2S,3S)-N-(2-Chloro-4-iodo-phenyl)-2-[(R)-4-[4-((S)-2,3-di hydroxy-propoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-3-phenyl-butyramide;

(2S,3S)-2-[(R)-2,5-Dioxo-4-[4-(2-oxo-2-pyrrolidin-1-yl-ethoxy)-phenyl]-imidazolidin-1-yl]-N-(2-fluoro-4-iodo-phenyl)-3-phenyl-butyramide;

(2S,3S)-2-((R)-2,5-Dioxo-4-thiophen-3-yl-imidazolidin-1-yl)-N-(4-iodo-phenyl)-3-phenyl-butyramide;

(S)-2-[(R)-4-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-2,5-dioxo-imidazolidin-1-yl]-N-(2-fluoro-4-iodo-phenyl)-3-phenyl-propionamide;

(S)-2-[(R)-4-(4-Acetylamino-phenyl)-2,5-dioxo-imidazolidin-1-yl]-N-(2-fluoro-4-iodo-phenyl)-3-phenyl-propionamide;

(4-[(R)-1-[(1S,2S)-1-(2-Fluoro-4-iodo-phenylcarbamoyl)-2-phenyl-propyl]-2,5-dioxo-imidazolidin-4-yl]-phenoxy-methyl)-phosphonic acid dimethyl ester;

(2S,3S)-N-(2-Fluoro-4-iodo-phenyl)-2-((R)-4-isopropyl-2,5-dioxo-imidazolidin-1-yl)-3-phenyl-butyramide;

(S)-N-(2-Fluoro-4-iodo-phenyl)-2-[(R)-4-[4-(2-hydroxy-ethoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-3-methyl-butyramide;

(S)-N-(2-Fluoro-4-iodo-phenyl)-2-[(R)-4-(4-methoxy-phenyl)-2,5-dioxo-imidazolidin-1-yl]-3-o-tolyl-propionamide;


(S)-N-(2-Fluoro-4-iodo-phenyl)-2-[(R)-4-(4-methoxy-phenyl)-2,5-dioxo-imidazolidin-1-yl]-3-m-tolyl-propionamide;

(S)-N-(2-Fluoro-4-iodo-phenyl)-2-[(R)-4-(4-methoxy-phenyl)-2,5-dioxo-imidazolidin-1-yl]-3-p-tolyl-propionamide; and

(S)-N-(4-Cyclopropyl-2-fluoro-phenyl)-3-(4-fluoro-phenyl)-2-[(R)-4-[4-(2-hydroxy-1-hydroxymethyl-ethoxy)-phenyl]-2,5-dioxo-imidazolidin-1-yl]-propionamide ;

or a pharmaceutically acceptable salt or ester thereof.

[0205] In some embodiments, the MEK inhibitor is a compound of formula (VII),



VII

or a pharmaceutically acceptable salt or ester thereof, wherein:

R1 is selected from the group consisting of halogen, ethynyl, and cycloalkyl;

R2 is selected from the group consisting of hydrogen and CH(R3)(R4);

R3 is selected from the group consisting of lower alkyl, lower alkoxy, optionally substituted aryl, and optionally substituted heteroaryl;

R4 is selected from the group consisting of hydrogen and lower alkyl;

R5 is hydrogen or, taken together with R2 and the carbon to which R2 and R5 are attached, forms lower cycloalkyl; and

R6 is selected from the group consisting of hydrogen, lower alkyl, lower cycloalkyl, optionally substituted aryl, and optionally substituted heteroaryl.

[0206] In some variations, the MEK inhibitor compound is of the formula (VI), or a pharmaceutically acceptable salt or ester thereof, wherein the variables are as defined in WO 2009/021887 A1 for the formula I or any applicable variations described on pages 4-5 in WO 2009/021887 A1, incorporated herein by reference. Further embraced MEK inhibitors are compounds described in Examples 1-21 in 2009/021887 A1, incorporated herein by reference.

[0207] In some embodiments, the MEK inhibitor compound of the formula (VI) is a compound selected from the group consisting of:

(R)-5-[4-(2-Hydroxy-ethoxy)-phenyl]-3-[(S)-1-(6-iodo-1*H*-benzoimidazol-2-yl)-2-phenyl-ethyl]-imidazolidine-2,4-dione;

(R)-5-[4-(2-Hydroxy-ethoxy)-phenyl]-3-(5-iodo-1*H*-benzoimidazol-2-ylmethyl)-imidazolidine-2,4-dione;

(R)-5-[4-(2-Hydroxy-ethoxy)-phenyl]-3-[(S)-1-(5-*ido*-1*H*-benzoimidazol-2-yl)-2-methyl-propyl]-imidazolidine-2,4-dione;

(R)-5-[4-(2-Hydroxy-ethoxy)-phenyl]-3-[(1*R*,2*R*)-1-(5-*ido*-1*H*-benzoimidazol-2-yl)-2-methoxy-propyl]-imidazolidine-2,4-dione;

3-[(S)-1-(5-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-ethyl]-imidazolidine-2,4-dione; compound with trifluoro-acetic acid;

(R)-3-[(S)-2-(4-Fluoro-phenyl)-1-(5-*ido*-1*H*-benzoimidazol-2-yl)-ethyl]-5-[4-(2-hydroxy-ethoxy)-phenyl]-imidazolidine-2,4-dione;

(R)-5-[4-(2-Hydroxy-ethoxy)-phenyl]-3-[(S)-1-(5-*ido*-1*H*-benzoimidazol-2-yl)-2-(4-methoxy-phenyl)-ethyl]-imidazolidine-2,4-dione;

(R)-5-[4-(2-Hydroxy-ethoxy)-phenyl]-3-[(S)-1-(5-*ido*-1*H*-benzoimidazol-2-yl)-2-thiophen-2-yl-ethyl]-imidazolidine-2,4-dione;

(R)-3-[(1*S*,2*S*)-1-(6-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-5-phenyl-imidazolidine-2,4-dione;

(R)-3-[(1*S*,2*S*)-1-(6-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-5-(4-methoxy-phenyl)-imidazolidine-2,4-dione;

(R)-5-[4-(2-Hydroxy-ethoxy)-phenyl]-3-[(1*S*,2*S*)-1-(6-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-imidazolidine-2,4-dione;

(R)-3-[(1*S*,2*S*)-1-(6-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-5-[4-(2-methoxy-ethoxy)-phenyl]-imidazolidine-2,4-dione;

2-(4-{(R)-1-[(1*S*,2*S*)-1-(6-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-2,5-dioxo-imidazolidin-4-yl}-phenoxy)-*N,N*-dimethyl-acetamide;

N,N-*Bis*-(2-hydroxy-ethyl)-2-(4-{(R)-1-[(1*S*,2*S*)-1-(6-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-2,5-dioxo-imidazolidin-4-yl}-phenoxy)-acetamide;

(R)-3-[(1*S*,2*S*)-1-(5-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-5-isopropyl-imidazolidine-2,4-dione;

(R)-5-Cyclohexyl-3-[(1*S*,2*S*)-1-(5-*ido*-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-imidazolidine-2,4-dione;

(R)-5-[4-(2-Hydroxy-ethoxy)-phenyl]-3-[1-(5-*ido*-1*H*-benzoimidazol-2-yl)-cyclopropyl]-imidazolidine-2,4-dione;

(R)-3-[(1*S*,2*S*)-1-(6-Bromo-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-5-[4-(2-hydroxy-ethoxy)-phenyl]-imidazolidine-2,4-dione;


(R)-3-[(S)-1-(5-Cyclopropyl-1*H*-benzoimidazol-2-yl)-2-phenyl-ethyl]-5-[4-(2-hydroxy-ethoxy)-phenyl]-imidazolidine-2,4-dione;

(R)-3-[(S)-1-(5-Ethynyl-1*H*-benzimidazol-2-yl)-2-phenyl-ethyl]-5-[4-(2-hydroxyethoxy)-phenyl]-imidazolidine-2,4-dione; and


(R)-3-[(1S,2S)-1-(5-Ethynyl-1*H*-benzoimidazol-2-yl)-2-phenyl-propyl]-5-[4-(2-hydroxy-ethoxy)-phenyl]-imidazolidine-2,4-dione;

or a pharmaceutically acceptable salt or solvate thereof.


[0208] In some embodiments, the MEK inhibitor is a compound selected from the group consisting of GDC-0973 (Methanone, [3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]phenyl][3-hydroxy-3-(2S)-2-piperidinyl-1-azetidinyl]-), G-38963, G02443714, G02442104, and G00039805, or a pharmaceutically acceptable salt or solvate thereof.




GDC-0973




G-38963



G02443714



G02442104



G00039805

IV Kits

[0209] In another aspect, provided herein is a kit comprising a PD-L1 axis binding antagonist and/or a MEK inhibitor for treating or delaying progression of a cancer in an individual or for enhancing immune function of an individual having cancer. In some embodiments, the kit comprises a PD-1 axis binding antagonist and a package insert comprising instructions for using

the PD-1 axis binding antagonist in combination with a MEK inhibitor to treat or delay progression of cancer in an individual or to enhance immune function of an individual having cancer. In some embodiments, the kit comprises a MEK inhibitor and a package insert comprising instructions for using the MEK inhibitor in combination with a PD-1 axis binding antagonist to treat or delay progression of cancer in an individual or to enhance immune function of an individual having cancer. In some embodiments, the kit comprises a PD-1 axis binding antagonist and a MEK inhibitor, and a package insert comprising instructions for using the PD-1 axis binding antagonist and the MEK inhibitor to treat or delay progression of cancer in an individual or to enhance immune function of an individual having cancer. Any of the PD-1 axis binding antagonists and/or MEK inhibitors described herein may be included in the kits.

[0210] In some embodiments, the kit comprises a container containing one or more of the PD-1 axis binding antagonists and MEK inhibitors described herein. Suitable containers include, for example, bottles, vials (e.g., dual chamber vials), syringes (such as single or dual chamber syringes) and test tubes. The container may be formed from a variety of materials such as glass or plastic. In some embodiments, the kit may comprise a label (e.g., on or associated with the container) or a package insert. The label or the package insert may indicate that the compound contained therein may be useful or intended for treating or delaying progression of cancer in an individual or for enhancing immune function of an individual having cancer. The kit may further comprise other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

EXAMPLES

[0211] The invention can be further understood by reference to the following examples, which are provided by way of illustration and are not meant to be limiting.

Example 1: Combination treatment with an anti-PDL1 antibody and a MEK inhibitor causes sustained tumor regression in Vemurafenib-progressing tumors

[0212] While B-raf inhibition (such as by treatment with Vemurafenib) is effective in eliciting short-term tumor regression, resistance is frequently observed. This Example describes the finding that treatment with a combination of a PD-1 axis binding antagonist and a MEK inhibitor induces sustained tumor regression and increased progression-free survival in animals with Vemurafenib-progressing tumors. Moreover, treatment with a combination of a PD-1 axis

binding antagonist and a MEK inhibitor was surprisingly superior to treatment with either agent individually.

Materials and Methods

Mouse model

[0213] A melanoma GEM model *B-raf*^{V600E}; *PTEN*^{fl/fl}; *TyCreER* was used. *B-raf*^{V600E} and *TyCreER* alleles were as described in Dankort, D., *et al.* *Nat. Genet.* 41(5):544-52 (2009). The *PTEN* conditional allele was as described in Lesche, R. *et al.* *genesis* 32:148-9 (2002).

Tumor initiation

[0214] Tumors were initiated by application of tamoxifen as described in Dankort, D., *et al.* *Nat. Genet.* 41(5):544-52 (2009). Animals were enrolled into the study once their tumors reached a size greater than or equal to 400 mm³.

Treatments

[0215] Prior to beginning treatment, each mouse received a biopsy of a melanoma tumor. After the biopsy, mice were allowed to recover for up to one week prior to receiving treatment. Mice were assigned into initial treatment groups (n=20), and treatment commenced at day 0.

[0216] For Vemurafenib treatment, mice were given either MCT, 200µL, PO, qd; or PLX-4032 (Vemurafenib), 50 mg/kg PO, BID (volume not to exceed 300µL). When animals in any group reached ~2000mm³, the tumors were biopsied a second time. After the biopsy procedure, mice recovered for up to one week prior to receiving further therapeutic treatment. The animals were then re-assigned to the following treatment groups: GDC-0973 (Cobimetinib), 7.5 mg/kg PO, qd (volume not to exceed 300µL) + Ragweed Control (IgG2a), 10mg/kg IP, three times weekly; GDC-0973, 7.5 mg/kg PO, qd (volume not to exceed 300µL) + anti-PDL1 (IgG1-WT), 10mg/kg IP, three times weekly; or MCT, 200uL, PO, qd + anti-PDL1 (IgG1-WT), 10mg/kg IP, three times weekly. IP dose volume did not exceed 300 µL.

[0217] Mice were weighed and tumors measured at least once a week until study termination. Mice received treatment each day until a mean tumor volume of 2500 mm³ was achieved. Mice were then euthanized, and melanoma tumors were collected for histology and assessing molecular changes. Mice were perfused under anesthesia at euthanasia.

[0218] Throughout the study, mice were monitored for clinical appearance (body condition, coat appearance, posture, labored breathing, etc.) at least 2 times a week, with increasing frequency, up to daily, depending on severity of adverse clinical signs observed. Moribund animals were euthanized. Mice with a body condition score < 2 were euthanized.

Results

[0219] Upon tumor induction, the melanoma GEM model $B\text{-}raf^{V600E};PTEN^{fl/fl};TyCreER$ causes tumors that show an initial regression in size upon treatment with the B-raf inhibitor Vemurafenib. After this initial regression, the tumors display steady re-growth, thereby modeling resistance to B-raf inhibition in Vemurafenib-progressing tumors.

[0220] This model was used to test the efficacy of PD-1 axis binding antagonists and MEK inhibitors as a 2nd line therapy for Vemurafenib-progressing tumors. As shown in **FIG. 1**, after first line treatment with Vemurafenib, animals were treated with an antibody against PD-L1, a MEK inhibitor (Cobimetinib), or both. Treatment with anti-PD-L1 alone showed no effect on tumor growth. Treatment with Cobimetinib caused an initial tumor regression, but this response was not sustained and tumor re-growth was observed. Combination treatment with anti-PD-L1 and Cobimetinib, however, caused regression in every tumor, and this regression was sustained. In addition, intratumoral GR1 levels were significantly reduced, and a signature of T cell activation was observed (e.g., increased CD8, PRF1, and MHC I). Importantly, treatment with anti-PD-L1 and Cobimetinib led to increased progression-free survival (PFS).

[0221] **FIG. 2** shows individual animal responses following crossover from Vemurafenib to anti-PD-L1, Cobimetinib, or combination treatment.

[0222] These results demonstrate that combined treatment with a PD-1 axis binding antagonist and a MEK inhibitor leads to dramatic, sustained tumor regression in Vemurafenib-progressing tumors, as compared to treatment with each agent alone. Moreover, these results demonstrate the superior efficacy of combined PD-1 axis/MEK inhibition as a 2nd line treatment for tumors resistant to B-raf inhibition.

[0223] All patents, patent applications, documents, and articles cited herein are herein incorporated by reference in their entireties.

CLAIMS

What is claimed is:

1. A method for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor, wherein the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist.
2. The method of claim 1, further comprising diagnosing the individual as having a cancer that is resistant to a B-raf antagonist, wherein the diagnosing occurs prior to administering the effective amount of the PD-1 axis binding antagonist and the MEK inhibitor.
3. The method of claim 1 or claim 2, wherein the individual has not been previously treated with a B-raf antagonist.
4. The method of claim 1 or claim 2, wherein the individual has been previously treated with a B-raf antagonist.
5. A method for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a PD-1 axis binding antagonist and a MEK inhibitor, wherein the individual has been previously treated with a B-raf antagonist for cancer.
6. The method of claim 4 or 5, wherein the cancer in the individual has progressed within 1 month, 6 months, 1 year, or 5 years after completing a B-raf antagonist-based therapy regimen.
7. The method of any one of claims 1-6, wherein the B-raf antagonist is a small molecule inhibitor, an antibody, a peptide, a peptidomimetic, an aptamer or a polynucleotide.
8. The method of claim 7, wherein the B-raf antagonist is dabrafenib, vemurafenib, GSK 2118436, RAF265, XL281, ARQ736, BAY73-4506, sorafenib, PLX4720, PLX-3603, GSK2118436, GDC-0879, or N-(3-(5-(4-chlorophenyl)-1H-pyrrolo[2,3-b]pyridine-3-carbonyl)-2,4-difluorophenyl)propane-1-sulfonamide.
9. The method of claim 7, wherein the B-raf antagonist is a selective B-raf antagonist of B-raf V600.
10. The method of claim 7, wherein the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600E.
11. The method of claim 7, wherein the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600E, B-raf V600K, and/or V600D.

12. The method of claim 7, wherein the selective B-raf antagonist of B-raf V600 is a selective antagonist of B-raf V600R.
13. The method of any one of claims 1-12, wherein the cancer contains a BRAF V600E mutation, a BRAF wildtype, a KRAS wildtype, or an activating KRAS mutation.
14. The method of any one of claims 1-13, wherein the treatment results in a sustained response in the individual after cessation of the treatment.
15. The method of any one of claims 1-14, wherein the individual has colorectal cancer, melanoma, lung cancer, ovarian cancer, breast cancer, pancreatic cancer, hematological malignancy, bladder cancer, and/or renal cell carcinoma.
16. The method of claim 15, wherein the cancer is metastatic.
17. The method of any one of the preceding claims, wherein the PD-1 axis binding antagonist is selected from the group consisting of a PD-1 binding antagonist, a PD-L1 binding antagonist and a PD-L2 binding antagonist.
18. The method of claim 17, wherein the PD-1 axis binding antagonist is a PD-1 binding antagonist.
19. The method of claim 18, wherein the PD-1 binding antagonist inhibits the binding of PD-1 to its ligand binding partners.
20. The method of claim 19, wherein the PD-1 binding antagonist inhibits the binding of PD-1 to PD-L1, PD-1 to PD-L2, or PD-1 to both PD-L1 and PD-L2.
21. The method of any one of claims 18-20, wherein the PD-1 binding antagonist is an antibody.
22. The method of any one of claims 18-20, wherein the PD-1 binding antagonist is nivolumab, pembrolizumab, pidilizumab, MEDI-0680, PDR001, REGN2810, BGB-108, BGB-A317, or AMP-224.
23. The method of claim 17, wherein the PD-1 axis binding antagonist is a PD-L1 binding antagonist.
24. The method of claim 23, wherein the PD-L1 binding antagonist inhibits the binding of PD-L1 to PD-1, PD-L1 to B7-1, or PD-L1 to both PD-1 and B7-1.
25. The method of claim 23 or claim 24, wherein the PD-L1 binding antagonist is an anti-PD-L1 antibody.
26. The method of claim 25, wherein the anti-PD-L1 antibody is a monoclonal antibody.
27. The method of claim 25, wherein the anti-PD-L1 antibody is an antibody fragment

selected from the group consisting of Fab, Fab'-SH, Fv, scFv, and (Fab')₂ fragments.

28. The method of any one of claims 25-27, wherein the anti-PD-L1 antibody is a humanized antibody or a human antibody.

29. The method of any one of claim 25, wherein the PD-L1 binding antagonist is selected from the group consisting of: YW243.55.S70, atezolizumab, durvalumab, MDX-1105, and avelumab.

30. The method of any one of claims 25-28, wherein the antibody comprises a heavy chain comprising HVR-H1 sequence of SEQ ID NO:15, HVR-H2 sequence of SEQ ID NO:16, and HVR-H3 sequence of SEQ ID NO:3; and a light chain comprising HVR-L1 sequence of SEQ ID NO:17, HVR-L2 sequence of SEQ ID NO:18, and HVR-L3 sequence of SEQ ID NO:19.

31. The method of any one of claims 25-28, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:24 or SEQ ID NO:28 and a light chain variable region comprising the amino acid sequence of SEQ ID NO:21.

32. The method of claim 17, wherein the PD-1 axis binding antagonist is a PD-L2 binding antagonist.

33. The method of claim 32, wherein the PD-L2 binding antagonist is an antibody.

34. The method of claim 32, wherein the PD-L2 binding antagonist is an immunoadhesin.

35. The method of any one of claims 1-34, wherein the MEK inhibitor is a competitive inhibitor of MEK.

36. The method of any one of claims 1-34, wherein the MEK inhibitor is more selective against an activating KRAS mutation.

37. The method of any one of claims 1-34, wherein the MEK inhibitor is an allosteric inhibitor of MEK.

38. The method of any one of claims 1-34, wherein the MEK inhibitor is more selective against an activating BRAF mutation.

39. The method of any one of claims 1-34, wherein the MEK inhibitor is a compound of the formula (I), (II), (III), (IV), (V), (VI) or (VII), or a pharmaceutically acceptable salt or solvate thereof.

40. The method of any one of claims 1-34, wherein the MEK inhibitor is selected from the group consisting of G02442104, G-38963, G02443714, G00039805 and GDC-0973, or a pharmaceutically acceptable salt or solvate thereof.

41. The method of claim 40, wherein the MEK inhibitor is G02443714, G02442104 or

G00039805.

42. The method of any one of claims 1-41, wherein the MEK inhibitor is administered continuously.
43. The method of any one of claims 1-41, wherein the MEK inhibitor is administered intermittently.
44. The method of any one of claims 1-41, wherein the MEK inhibitor is administered before the PD-1 axis binding antagonist.
45. The method of any one of claims 1-41, wherein the MEK inhibitor is administered simultaneous with the PD-1 axis binding antagonist.
46. The method of any one of claims 1-41, wherein the MEK inhibitor is administered after the PD-1 axis binding antagonist.
47. The method of any one of claims 1-46, wherein the PD-1 axis binding antagonist and/or the MEK inhibitor is administered intravenously, intramuscularly, subcutaneously, topically, orally, transdermally, intraperitoneally, intraorbitally, by implantation, by inhalation, intrathecally, intraventricularly, or intranasally.
48. A kit comprising a PD-1 axis binding antagonist and a package insert comprising instructions for using the PD-1 axis binding antagonist in combination with a MEK inhibitor to treat or delay progression of cancer in an individual, wherein the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist.
49. A kit comprising a PD-1 axis binding antagonist and a MEK inhibitor, and a package insert comprising instructions for using the PD-1 axis binding antagonist and the MEK inhibitor to treat or delay progression of cancer in an individual, wherein the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist.
50. A kit comprising a MEK inhibitor and a package insert comprising instructions for using the MEK inhibitor in combination with a PD-1 axis binding antagonist to treat or delay progression of cancer in an individual, wherein the individual has cancer or is at risk of developing cancer that is resistant to a B-raf antagonist.
51. A kit comprising a PD-1 axis binding antagonist and a package insert comprising instructions for using the PD-1 axis binding antagonist in combination with a MEK inhibitor to treat or delay progression of cancer in an individual, wherein the individual has been previously treated with a B-raf antagonist for cancer.
52. A kit comprising a PD-1 axis binding antagonist and a MEK inhibitor, and a package

insert comprising instructions for using the PD-1 axis binding antagonist and the MEK inhibitor to treat or delay progression of cancer in an individual, wherein the individual has been previously treated with a B-raf antagonist for cancer.

53. A kit comprising a MEK inhibitor and a package insert comprising instructions for using the MEK inhibitor in combination with a PD-1 axis binding antagonist to treat or delay progression of cancer in an individual, wherein the individual has been previously treated with a B-raf antagonist for cancer.




FIG. 1




FIG. 2

146392027540SEQLI ST

SEQUENCE LISTING

<110> GENENTECH, INC.
JUNTTILA, Meissa

<120> METHODS OF TREATING CANCER USING PD-1
AXIS BINDING ANTAGONISTS AND MEK INHIBITORS

<130> 146392027540

<140> Not Yet Assigned
<141> Concurrently Herewith

<150> US 62/024, 988
<151> 2014-07-15

<160> 28

<170> FastSEQ for Windows Version 4.0

<210> 1
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<220>
<221> VARIANT
<222> 6
<223> Xaa = Asp or Glu

<400> 1
Gly Phe Thr Phe Ser Xaa Ser Trp Ile His
1 5 10

<210> 2
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<220>
<221> VARIANT
<222> 4
<223> Xaa = Ser or Leu

<220>
<221> VARIANT
<222> 10
<223> Xaa = Thr or Ser

<400> 2
Ala Trp Ile Xaa Pro Tyr Gly Gly Ser Xaa Tyr Tyr Ala Asp Ser Val
1 5 10 15
Lys Glu

<210> 3
<211> 9
<212> PRT

146392027540SEQLIST

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 3

Arg His Trp Pro Gly Gly Phe Asp Tyr
1 5

<210> 4

<211> 25

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 4

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> 5

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 5

Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
1 5 10

<210> 6

<211> 32

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 6

Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr Leu Gln
1 5 10 15
Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
20 25 30

<210> 7

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 7

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
1 5 10

<210> 8

<211> 11

<212> PRT
 <213> Arti fi ci al Sequence

<220>
 <223> Synthetic Construct

<220>
 <221> VARI ANT
 <222> 5
 <223> Xaa = Asp or Val

<220>
 <221> VARI ANT
 <222> 6
 <223> Xaa = Val or Ile

<220>
 <221> VARI ANT
 <222> 7
 <223> Xaa = Ser or Asn

<220>
 <221> VARI ANT
 <222> 9
 <223> Xaa = Ala or Phe

<220>
 <221> VARI ANT
 <222> 10
 <223> Xaa = Val or Leu

<400> 8
 Arg Ala Ser Gln Xaa Xaa Xaa Thr Xaa Xaa Ala
 1 5 10

<210> 9
 <211> 7
 <212> PRT
 <213> Arti fi ci al Sequence

<220>
 <223> Synthetic Construct

<220>
 <221> VARI ANT
 <222> 4
 <223> Xaa = Phe or Thr

<220>
 <221> VARI ANT
 <222> 6
 <223> Xaa = Tyr or Ala

<400> 9
 Ser Ala Ser Xaa Leu Xaa Ser
 1 5

<210> 10
 <211> 9
 <212> PRT
 <213> Arti fi ci al Sequence

<220>
 <223> Synthetic Construct

<220>

146392027540SEQLI ST

<221> VARI ANT

<222> 3

<223> Xaa = Tyr, Gl y, Phe or Ser

<220>

<221> VARI ANT

<222> 4

<223> Xaa = Leu, Tyr, Phe or Trp

<220>

<221> VARI ANT

<222> 5

<223> Xaa = Tyr, Asn, Ala, Thr, Gl y, Phe or Ile

<220>

<221> VARI ANT

<222> 6

<223> Xaa = His, Val, Pro, Thr or Ile

<220>

<221> VARI ANT

<222> 8

<223> Xaa = Ala, Trp, Arg, Pro or Thr

<400> 10

Gl n Gl n Xaa Xaa Xaa Xaa Pro Xaa Thr

1

5

<210> 11

<211> 23

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 11

Asp Ile Gl n Met Thr Gl n Ser Pro Ser Ser Leu Ser Ala Ser Val Gl y

1

5

10

15

Asp Arg Val Thr Ile Thr Cys

20

<210> 12

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 12

Trp Tyr Gl n Gl n Lys Pro Gl y Lys Ala Pro Lys Leu Leu Ile Tyr

1

5

10

15

<210> 13

<211> 32

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 13

Gl y Val Pro Ser Arg Phe Ser Gl y Ser Gl y Ser Gl y Thr Asp Phe Thr

1

5

10

15

146392027540SEQLI ST
Leu Thr Ile Ser Ser Leu Glu Pro Glu Asp Phe Ala Thr Tyr Tyr Cys
20 25 30

<210> 14
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<400> 14
Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
1 5 10

<210> 15
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<400> 15
Gly Phe Thr Phe Ser Asp Ser Trp Ile His
1 5 10

<210> 16
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<400> 16
Ala Trp Ile Ser Pro Tyr Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
1 5 10 15
Lys Gly

<210> 17
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<400> 17
Arg Ala Ser Gln Asp Val Ser Thr Ala Val Ala
1 5 10

<210> 18
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

<400> 18
Ser Ala Ser Phe Leu Tyr Ser

<210> 19
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 19
 Gln Gln Tyr Leu Tyr His Pro Ala Thr
 1 5

<210> 20
 <211> 118
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 20
 Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Ser
 20 25 30
 Trp Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Trp Ile Ser Pro Tyr Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Arg His Trp Pro Gly Gly Phe Asp Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ala
 115

<210> 21
 <211> 108
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 21
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Ser Thr Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Leu Tyr His Pro Ala
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 100 105

<210> 22
 <211> 440
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 22
 Gln Val Gln Leu Val Glu Ser Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Asp Cys Lys Ala Ser Gly Ile Thr Phe Ser Asn Ser
 20 25 30
 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Val Ile Trp Tyr Asp Gly Ser Lys Arg Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Phe
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Gln Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Thr Asn Asp Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 100 105 110
 Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser
 115 120 125
 Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp
 130 135 140
 Tyr Phe Pro Gln Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr
 145 150 155 160
 Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr
 165 170 175
 Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys
 180 185 190
 Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp
 195 200 205
 Lys Arg Val Gln Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala
 210 215 220
 Pro Gln Phe Leu Gln Gln Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
 225 230 235 240
 Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Gln Val Thr Cys Val Val
 245 250 255
 Val Asp Val Ser Gln Gln Asp Pro Gln Val Gln Phe Asn Trp Tyr Val
 260 265 270
 Asp Gln Val Gln Val His Asn Ala Lys Thr Lys Pro Arg Gln Gln Gln
 275 280 285
 Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
 290 295 300
 Asp Trp Leu Asn Gln Lys Gln Tyr Lys Cys Lys Val Ser Asn Lys Gln
 305 310 315 320
 Leu Pro Ser Ser Ile Gln Lys Thr Ile Ser Lys Ala Lys Gln Gln Pro
 325 330 335
 Arg Gln Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Gln Gln Met Thr
 340 345 350
 Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
 355 360 365
 Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
 370 375 380
 Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gln Ser Phe Phe Leu Tyr
 385 390 395 400
 Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
 405 410 415
 Ser Cys Ser Val Met His Gln Ala Leu His Asn His Tyr Thr Gln Lys
 420 425 430
 Ser Leu Ser Leu Ser Leu Gln Lys
 435 440

146392027540SEQLI ST

<211> 214

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 23

Gl u Ile Val Leu Thr Gl n Ser Pro Al a Thr Leu Ser Leu Ser Pro Gl y
1 5 10 15
Gl u Arg Al a Thr Leu Ser Cys Arg Al a Ser Gl n Ser Val Ser Ser Tyr
20 25 30
Leu Al a Trp Tyr Gl n Gl n Lys Pro Gl y Gl n Al a Pro Arg Leu Leu Ile
35 40 45
Tyr Asp Al a Ser Asn Arg Al a Thr Gl y Ile Pro Al a Arg Phe Ser Gl y
50 55 60
Ser Gl y Ser Gl y Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gl u Pro
65 70 75 80
Gl u Asp Phe Al a Val Tyr Tyr Cys Gl n Gl n Ser Ser Asn Trp Pro Arg
85 90 95
Thr Phe Gl y Gl n Gl y Thr Lys Val Gl u Ile Lys Arg Thr Val Al a Al a
100 105 110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Gl u Gl n Leu Lys Ser Gl y
115 120 125
Thr Al a Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Gl u Al a
130 135 140
Lys Val Gl n Trp Lys Val Asp Asn Al a Leu Gl n Ser Gl y Asn Ser Gl n
145 150 155 160
Gl u Ser Val Thr Gl u Gl n Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
165 170 175
Ser Thr Leu Thr Leu Ser Lys Al a Asp Tyr Gl u Lys His Lys Val Tyr
180 185 190
Al a Cys Gl u Val Thr His Gl n Gl y Leu Ser Ser Pro Val Thr Lys Ser
195 200 205
Phe Asn Arg Gl y Gl u Cys
210

<210> 24

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 24

Gl u Val Gl n Leu Val Gl u Ser Gl y Gl y Gl y Leu Val Gl n Pro Gl y Gl y
1 5 10 15
Ser Leu Arg Leu Ser Cys Al a Al a Ser Gl y Phe Thr Phe Ser Asp Ser
20 25 30
Trp Ile His Trp Val Arg Gl n Al a Pro Gl y Lys Gl y Leu Gl u Trp Val
35 40 45
Al a Trp Ile Ser Pro Tyr Gl y Gl y Ser Thr Tyr Tyr Al a Asp Ser Val
50 55 60
Lys Gl y Arg Phe Thr Ile Ser Al a Asp Thr Ser Lys Asn Thr Al a Tyr
65 70 75 80
Leu Gl n Met Asn Ser Leu Arg Al a Gl u Asp Thr Al a Val Tyr Tyr Cys
85 90 95
Al a Arg Arg His Trp Pro Gl y Gl y Phe Asp Tyr Trp Gl y Gl n Gl y Thr
100 105 110
Leu Val Thr Val Ser Ser
115

<210> 25

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 25

Trp Gl y Gl n Gl y Thr Leu Val Thr Val Ser Ser
1 5 10

<210> 26

<211> 448

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 26

 Gl u Val Gl n Leu Val Gl u Ser Gl y Gl y Leu Val Gl n Pro Gl y Gl y
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Al a Al a Ser Gl y Phe Thr Phe Ser Asp Ser
 20 25 30
 Trp Ile His Trp Val Arg Gl n Al a Pro Gl y Lys Gl y Leu Gl u Trp Val
 35 40 45
 Al a Trp Ile Ser Pro Tyr Gl y Gl y Ser Thr Tyr Tyr Al a Asp Ser Val
 50 55 60
 Lys Gl y Arg Phe Thr Ile Ser Al a Asp Thr Ser Lys Asn Thr Al a Tyr
 65 70 75 80
 Leu Gl n Met Asn Ser Leu Arg Al a Gl u Asp Thr Al a Val Tyr Tyr Cys
 85 90 95
 Al a Arg Arg His Trp Pro Gl y Gl y Phe Asp Tyr Trp Gl y Gl n Gl y Thr
 100 105 110
 Leu Val Thr Val Ser Ser Al a Ser Thr Lys Gl y Pro Ser Val Phe Pro
 115 120 125
 Leu Al a Pro Ser Ser Lys Ser Thr Ser Gl y Gl y Thr Al a Al a Leu Gl y
 130 135 140
 Cys Leu Val Lys Asp Tyr Phe Pro Gl u Pro Val Thr Val Ser Trp Asn
 145 150 155 160
 Ser Gl y Al a Leu Thr Ser Gl y Val His Thr Phe Pro Al a Val Leu Gl n
 165 170 175
 Ser Ser Gl y Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 180 185 190
 Ser Leu Gl y Thr Gl n Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
 195 200 205
 Asn Thr Lys Val Asp Lys Lys Val Gl u Pro Lys Ser Cys Asp Lys Thr
 210 215 220
 His Thr Cys Pro Pro Cys Pro Al a Pro Gl u Leu Leu Gl y Gl y Pro Ser
 225 230 235 240
 Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
 245 250 255
 Thr Pro Gl u Val Thr Cys Val Val Val Asp Val Ser His Gl u Asp Pro
 260 265 270
 Gl u Val Lys Phe Asn Trp Tyr Val Asp Gl y Val Gl u Val His Asn Al a
 275 280 285
 Lys Thr Lys Pro Arg Gl u Gl u Gl n Tyr Al a Ser Thr Tyr Arg Val Val
 290 295 300
 Ser Val Leu Thr Val Leu His Gl n Asp Trp Leu Asn Gl y Lys Gl u Tyr
 305 310 315 320
 Lys Cys Lys Val Ser Asn Lys Al a Leu Pro Al a Pro Ile Gl u Lys Thr
 325 330 335
 Ile Ser Lys Al a Lys Gl y Gl n Pro Arg Gl u Pro Gl n Val Tyr Thr Leu
 340 345 350
 Pro Pro Ser Arg Gl u Gl u Met Thr Lys Asn Gl n Val Ser Leu Thr Cys
 355 360 365
 Leu Val Lys Gl y Phe Tyr Pro Ser Asp Ile Al a Val Gl u Trp Gl u Ser
 370 375 380
 Asn Gl y Gl n Pro Gl u Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp

146392027540SEQLIST

385		390		395		400									
Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser
				405				410					415		
Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glut	Ala
				420				425				430			
Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys
				435			440				445				

<210> 27

<211> 214

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 27

<210> 28

<211> 122

<212> PRT

<213> Artifi ci al Sequence

<220>

<223> Synthetic Construct

<400> 28

Gl u	Val	Gl n	Leu	Val	Gl u	Ser	Gl y	Gl y	Gl y	Leu	Val	Gl n	Pro	Gl y	Gl y
1					5			10						15	
Ser	Leu	Arg	Leu	Ser	Cys	Al a	Al a	Ser	Gl y	Phe	Thr	Phe	Ser	Asp	Ser
					20			25					30		
Trp	Ile	His	Trp	Val	Arg	Gl n	Al a	Pro	Gl y	Lys	Gl y	Leu	Gl u	Trp	Val
						35		40			45				
Al a	Trp	Ile	Ser	Pro	Tyr	Gl y	Gl y	Ser	Thr	Tyr	Tyr	Al a	Asp	Ser	Val
						50		55			60				
Lys	Gl y	Arg	Phe	Thr	Ile	Ser	Al a	Asp	Thr	Ser	Lys	Asn	Thr	Al a	Tyr
					70				75					80	
Leu	Gl n	Met	Asn	Ser	Leu	Arg	Al a	Gl u	Asp	Thr	Al a	Val	Tyr	Tyr	Cys

146392027540SEQLI ST

Ala Arg Arg His Trp Pro Gly Gly Phe Asp Tyr Trp Gly Gln Gly Thr
100 85 90 105 95 110
Leu Val Thr Val Ser Ser Ala Ser Thr Lys
115 120