




(86) Date de dépôt PCT/PCT Filing Date: 2016/02/05
(87) Date publication PCT/PCT Publication Date: 2016/08/11
(85) Entrée phase nationale/National Entry: 2017/07/25
(86) N° demande PCT/PCT Application No.: US 2016/016842
(87) N° publication PCT/PCT Publication No.: 2016/127097
(30) Priorité/Priority: 2015/02/06 (US62/113,300)

(51) Cl.Int./Int.Cl. *A61K 38/17* (2006.01),
A61K 38/22 (2006.01), *A61K 38/47* (2006.01),
C07K 14/50 (2006.01)
(71) Demandeur/Applicant:
THE REGENTS OF THE UNIVERSITY OF CALIFORNIA,
US
(72) Inventeur/Inventor:
DUBAL, DENA, US
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PROCEDES ET COMPOSITIONS POUR AMELIORER LA FONCTION COGNITIVE
(54) Title: METHODS AND COMPOSITIONS FOR IMPROVED COGNITION

FIG. 1A



(57) Abrégé/Abstract:

Provided herein are klotho polypeptide compositions and methods for improving cognitive function in an individual comprising treatment of with klotho polypeptides.



- (51) International Patent Classification:
A61K 38/17 (2006.01) *A61K 38/47* (2006.01)
A61K 38/22 (2006.01) *C07K 14/50* (2006.01)

(21) International Application Number:
PCT/US2016/016842

(22) International Filing Date:
5 February 2016 (05.02.2016)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/113,300 6 February 2015 (06.02.2015) US

(71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 1111 Franklin St., 5th Floor, Oakland, California 94607 (US).

(72) Inventor: DUBAL, Dena; 1111 Franklin St., 5th Floor, Oakland, California 94607 (US).

(74) Agents: HINSCH, Matthew E. et al.; Kilpatrick Townsend and Stockton LLP, Two Embarcadero Center, Eighth Floor, San Francisco, California 94111 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:
— with international search report (Art. 21(3))

[Continued on next page]

(54) Title: METHODS AND COMPOSITIONS FOR IMPROVED COGNITION

(57) Abstract: Provided herein are klotho polypeptide compositions and methods for improving cognitive function in an individual comprising treatment of with klotho polypeptides.

FIG. 1A



WO 2016/127097 A1

— *with sequence listing part of description (Rule 5.2(a))*

METHODS AND COMPOSITIONS FOR IMPROVED COGNITION

5 CROSS-REFERENCE TO RELATED PATENT APPLICATIONS

[0001] The present application claims benefit of priority to US Provisional Patent Application No. 62/113,300, filed February 6, 2015, which is incorporated by reference.

BACKGROUND OF THE INVENTION

[0002] Brain health is one of the biggest biomedical challenges with few if any effective
10 medical treatments. Cognition is a highly valued and central manifestation of brain health that is impaired or becomes disrupted in normal aging, numerous neurodegenerative, neurologic, and psychiatric diseases, childhood developmental syndromes, traumatic brain injury, and stress. Cognition is also disrupted by jet lag, medication side effects, and certain medical treatments, such as those for cancer. Thus, the potential to enhance cognition or
15 counter cognitive dysfunction is of enormous relevance across the human lifespan in health and disease.

BRIEF SUMMARY OF THE INVENTION

[0003] Provided herein are methods and compositions for improving cognition through
20 systemic administration of klotho or a protein comprising klotho or a functional fragment thereof. In some embodiments, the method comprises improving cognition in an individual comprising administering to the individual an effective amount of a protein comprising a Klotho polypeptide or a functional variant or fragment thereof, thereby improving cognitive function in the individual. In some embodiments, the administering is systemic, peripheral,
25 or nasal. In some embodiments, the protein is the Klotho polypeptide or a fragment thereof. In some embodiments, the administering is oral, mucosal, or carried out by injection. In some embodiments, the injection is intravenous, intraperitoneal, subcutaneous, or intramuscular. In some embodiments, the administration is by infusion, *e.g.*, continuous infusion using a reservoir or osmotic minipump.

30 [0004] In some embodiments, the individual is a human. In some embodiments, the human has at least normal cognitive function and the administering results in improved cognitive function compared to before the administering. In some embodiments, the human is 50 years

of age or older (*e.g.*, at least 60, 65, 70, 75, 80, 85, 90, 100, or older). In some embodiments, the individual is experiencing age-related cognitive decline. In some embodiments, the administering results in a reduced rate of cognitive decline, *e.g.*, so that short term memory does not decline as quickly as expected based on pre-treatment or age-cohort decline. In
 5 some embodiments, the individual is less than 50 years of age.

[0005] In some embodiments, the individual has a neurodegenerative disease, *e.g.*, Alzheimer's disease, Parkinson's disease, Huntington's disease, frontotemporal dementia, progressive supranuclear palsy, corticobasalar degeneration, mild cognitive impairment, vascular dementia, Lewy body dementia, multiple system atrophy, amyotrophic lateral
 10 sclerosis, prion disorder, and HIV-related dementia. In some embodiments, the individual has a mental or mood disorder, *e.g.*, depression, schizophrenia, attention deficit/ hyperactivity disorder, autism spectrum disorder, intellectual disability, a mood disorder, or a psychotic disorder. In some embodiments, the individual has a condition selected from traumatic brain injury, stroke, multiple sclerosis, neuroautoimmune disease, epilepsy, delirium, and a
 15 paraneoplastic disorder. In some embodiments, the individual has a condition selected from X-linked mental disorder, Down's syndrome, Angelman's syndrome, and Rett's syndrome. In some embodiments, the individual has a condition selected from phenylketonuria, Lesch-Nyhan, galactosemia, and adrenoleukodystrophy. In some embodiments, the individual is receiving radiation treatment or chemotherapy for cancer. In some embodiments, the
 20 individual is experiencing or is expected to experience (*e.g.*, in about 2 hours to 2 weeks, about 12-48 hours, or about 24 hours) stress, pain, sleep deprivation, or jet lag.

[0006] In some embodiments, the human has impaired motor function the administering results in improved motor function compared to before the administering.

[0007] In some embodiments, the klotho polypeptide has at least 75% identity (*e.g.*, at least
 25 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identity) to soluble human klotho (amino acids 34-979 of SEQ ID NO:1). In some embodiments, the klotho polypeptide is a functional fragment comprising a polypeptide with at least 75% identity (*e.g.*, at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identity) to the KL1 domain of human klotho. In some embodiments, the klotho polypeptide is a functional fragment comprising a
 30 polypeptide with at least 75% identity (*e.g.*, at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identity) to the KL2 domain of human klotho. In some embodiments, the klotho

polypeptide retains at least 40, 50, 60, 70, 80, 90, or 100% of the level of at least one activity of soluble human klotho.

[0008] In some embodiments, the klotho polypeptide is administered to the individual at a dose of 0.1-50,000 $\mu\text{g}/\text{kg}$ body weight (*e.g.*, 0.5-10 $\mu\text{g}/\text{kg}$, 0.5-500 $\mu\text{g}/\text{kg}$, 1-2000 $\mu\text{g}/\text{kg}$, 1-250 $\mu\text{g}/\text{kg}$, 5-250 $\mu\text{g}/\text{kg}$, 10-100 $\mu\text{g}/\text{kg}$, 1000-20,000 $\mu\text{g}/\text{kg}$, or about 10, 25, 50, 100, or 1000 $\mu\text{g}/\text{kg}$). In some embodiments, the klotho polypeptide is administered daily, twice per week, weekly, or every two weeks. In some embodiments, the klotho polypeptide is administered prior to (*e.g.*, 2, 6, 12, 24, or 48 hours) or in response to an event requiring heightened cognition, *e.g.*, stress, jet lag, sleep deprivation, or anticipated taxing mental task.

[0009] In some embodiments, the individual is tested for cognitive ability prior to the administering. In some embodiments, the individual is tested for cognitive ability after the administering. In some embodiments, the individual is tested for cognitive ability before and after the administering, or multiple times during the course of treatment. In some embodiments, the individual is tested for semantic, episodic, procedural, priming, and/or working memory. In some embodiments, the individual is tested for language ability, executive function, visuospatial function, or dementia. In some embodiments, the dose or frequency of administration of klotho polypeptide is increased when cognitive ability does not significantly increase.

[0010] In some embodiments, a method of improving motor function is provided. In some embodiments, a method is provided for improving motor function in an individual in need thereof, the method comprising administering to the individual an effective amount of a protein comprising a Klotho polypeptide or a functional fragment thereof, wherein the administering is systemic or peripheral, thereby improving motor function in the individual compared to before the administering. In some embodiments, the klotho polypeptide has at least 75% identity (*e.g.*, at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identity) to soluble human klotho (amino acids 34-979 of SEQ ID NO:1). In some embodiments, the klotho polypeptide is a functional fragment comprising a polypeptide with at least 75% identity (*e.g.*, at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identity) to the KL1 domain of human klotho. In some embodiments, the klotho polypeptide is a functional fragment comprising a polypeptide with at least 75% identity (*e.g.*, at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identity) to the KL2 domain of human klotho. In some embodiments, the klotho polypeptide retains at least 40, 50, 60, 70, 80, 90,

or 100% of the level of at least one activity of soluble human klotho. In some embodiments, the klotho polypeptide is administered to the individual at a dose of 0.1-50,000 µg/kg body weight (*e.g.*, 0.5-10 µg/kg, 0.5-500 µg/kg, 1-2000 µg/kg, 1-250 µg/kg, 5-250 µg/kg, 10-100 µg/kg, 1000-20,000 µg/kg, or about 10, 25, 50, 100, or 1000 µg/kg). In some embodiments, the klotho polypeptide is administered daily, twice per week, weekly, or every two weeks.

[0011] In some embodiments, the individual with impaired motor function has stroke to the brain or spinal cord (ischemic or hemorrhagic), neurodegenerative disease (Parkinson's disease, Lewy body dementia, multiple system atrophy, amyotrophic lateral sclerosis, prion disorder, Huntington's disease, supranuclear palsy), Parkinsonism, traumatic brain injury, neuroinfectious brain lesions, multiple sclerosis and related autoimmune and demyelinating disease, spinal cord lesions (compressive, infectious, toxic or metabolic, autoimmune, oncologic), brain tumor, epilepsy, paraneoplastic disorder, neurodevelopmental disorder (mitochondrial, autosomal genetic), muscle disease (polymyositis, dermatomyositis, inclusion body myositis, infectious, endocrine, metabolic, toxic, congenital myopathy, congenital muscular dystrophy, hereditary), neuropathies (Guillain-Barre syndrome, axonal and demyelinating, diabetic, toxic, metabolic, infectious, critical illness, entrapment), tick paralysis, myasthenia gravis, and spinal muscular atrophy.

BRIEF DESCRIPTION OF THE DRAWINGS

[0012] FIG. 1A-J. Klotho delivery enhances cognition in mice. Mice were tested in cognitive tasks following i.p. delivery of vehicle (Veh) or recombinant mouse klotho (KL) (10 µg/kg) (age 4 months). Data are mean ± SEM.

[0013] FIG. 1A. Mice were tested in the Y maze 18 hrs after treatment with vehicle or klotho. Percent alternations among arms during 4 minutes of exploration of a Y maze are shown (n=4 male mice/group).

[0014] FIG. 1B-J. Mice were tested in the Morris water maze after daily treatment with vehicle or klotho (two independent cohorts shown: Cohort 1 and Cohort 2).

[0015] FIG. 1B. Spatial learning curves when the platform is hidden are shown in Cohort 1. Veh or KL was administered 4h prior to testing. Data represent the daily average of total distance traveled on Days 1-4 to reach the hidden platform. Day 0 represents distance traveled on the first trial of Day 1. On Day 5, when the platform was visible, Veh- and klotho-treated mice located it equally well. (n=4 male mice/ group)

[0016] FIG. 1C. Veh- and klotho-treated mice swam at equal speeds in Cohort 1, as measured by the average velocity from hidden training on Day 4.

[0017] FIG. 1D. Spatial learning curves when the platform is hidden are shown in an independent cohort of mice, Cohort 2. Veh or KL was administered 18h prior to testing. Data represent the daily average of distance traveled on Days 1–4 to reach the hidden platform. (n=7–10 mice/group; males and females included)

[0018] FIG. 1E-J. Results of probe trials to assess spatial memory when the platform was removed 1hr and 24hrs after completion of hidden-platform training in Veh- and klotho-treated mice in Cohort 1.

[0019] FIG. 1E. Duration at the target center in a 1hr probe shows time spent at the original platform location.

[0020] FIG. 1F. Target platform crossings in a 1hr probe shows the frequency of crossings over the original platform location.

[0021] FIG. 1G. Percent time spent in the target quadrant, compared to the average time spent in other quadrants, in a 1hr probe. *p<0.05 (t-test)

[0022] FIG. 1H. Duration at the target center in a 24hr probe.

[0023] FIG. 1I. Target platform crossings in a 24hr probe.

[0024] FIG. 1J. Percent time spent in the target quadrant in a 24hr probe.

[0025] FIG. 2A-B. Acute delivery of klotho enhances cognition in aged mice. Mice were tested for spatial and working memory in the large Y-Maze following i.p. delivery of vehicle (Veh) or recombinant mouse klotho (KL) (10 µg/kg) 24 hours prior to training (n=8–9 mice per experimental group, sex-balanced groups, age 18 months, from NIH colony of aging mice). 18 hours after training, mice underwent testing and duration of time spent in the novel arm and the familiar arm was measured during exploration of the maze. (FIG. 2A) Novel arm preference is shown as a ratio time spent in the novel compared to the familiar arm throughout indicated times of exploration and (FIG. 2B) at 3 minutes. *p<0.05 (t-test). Data are mean ± SEM.

[0026] FIG. 3. Acute delivery of klotho induces long-lasting cognitive enhancement. Mice were tested for spatial and working memory in the large Y-Maze at 16 days following the last i.p. delivery of vehicle (Veh) or recombinant mouse klotho (KL) (0.5 or 2.5 µg/kg) that was

given daily for 5 days (n=8–10 mice per experimental group, sex-balanced groups, age 5–5.5 months of age). Duration of time spent in the novel arm and the familiar arm was measured during exploration of the maze 18 hours after training. Novel arm preference is shown as a ratio of time spent in the novel compared to the familiar arm at 5 min of exploration.

5 **p=0.01 (Bonferroni-Holm test). Data are mean $\pm$ SEM.

[0027] FIG. 4A-B. Acute delivery of klotho improves cognitive deficits in transgenic hSYN mice that express the human α -synuclein protein. Mice were tested for spatial and working memory in the large Y-Maze following i.p. delivery of vehicle (Veh) or recombinant mouse klotho (KL) (2.5 μ g/kg) at 22 hours prior to training and then 14 hours prior to testing
10 (n=6–9 male mice per group, age 2.5–6 months of age). Duration of time and number of entries in the novel and familiar arms were measured during exploration of the maze 18 hours after training. Data are mean $\pm$ SEM.

[0028] FIG. 4A. Novel arm preference is shown as ratio of time spent in the novel compared to the familiar arm at 5 min of exploration. Two-way ANOVA: hSYN effect
15 p=0.026, KL effect p=0.07; *p<0.05, #p=0.07 (Bonferroni-Holm test)

[0029] FIG. 4B. Novel arm preference is shown as ratio of entries into the novel compared to the familiar arm at 5 min of exploration. Two-way ANOVA: hSYN effect p=0.07, KL effect p=0.03; #p=0.07 (t-test).

[0030] FIG. 5. Acute delivery of klotho improves early motor deficits in transgenic hSYN
20 mice that express the human α -synuclein protein. Mice were tested for motor function on the rotarod task following i.p. delivery of vehicle (Veh) or recombinant mouse klotho (KL) (2.5 μ g/kg) at approximately 17 hours prior to testing (n=6–10 male mice per experimental group, age 2.5–6 months of age). Time spent on the spinning rod without falling is depicted in seconds (s). In hSYN mice, KL-treatment improves early motor function as shown by
25 increased duration of the average time spent on the spinning rod during Trials 1 through 3. Two-way ANOVA: hSYN effect 0.0025; **p<0.01, *p<0.05 (Bonferroni-Holm test). Data are mean $\pm$ SEM.

[0031] FIG. 6. Acute delivery of klotho does not alter the overall activity of mice. Mice were tested for level of overall movements and total activity in the open field task at 16 hours
30 following i.p. delivery of vehicle (Veh) or recombinant mouse klotho (KL) (2.5 μ g/kg) (n=8

mice per experimental group; sex-balanced groups; age 5 months). Total movements over 5 minutes are depicted. Data are mean $\pm$ SEM.

DETAILED DESCRIPTION OF THE INVENTION

I. Introduction

5 [0032] Klotho is a relatively large protein, approximately 130 kD in its secreted form, and is not expected to cross the blood-brain barrier. The results provided herein, however, show that Klotho exerts a positive effect on cognition within hours of systemic administration and long after its half-life elimination. These results are highly unexpected, and provide the advantage of safer, more convenient therapies compared to administration directly to the
10 brain or into the cerebrospinal fluid.

II. Definitions

[0033] Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by a person of ordinary skill in the art. See, *e.g.*, Lackie, DICTIONARY OF CELL AND MOLECULAR BIOLOGY, Elsevier (4th ed. 2007); Sambrook *et al.*,
15 MOLECULAR CLONING, A LABORATORY MANUAL, Cold Springs Harbor Press (Cold Springs Harbor, NY 1989). Any methods, devices and materials similar or equivalent to those described herein can be used in the practice of this invention. The following definitions are provided to facilitate understanding of certain terms used frequently herein and are not meant to limit the scope of the present disclosure.

20 [0034] The terms “klotho” or “klotho polypeptide” refer to soluble klotho polypeptide, and functional variants and fragments thereof, unless otherwise stated. Soluble klotho is any form of klotho that circulates in fluid (*e.g.*, serum, cerebrospinal fluid, etc.), and that does not include a transmembrane or intracellular component. Klotho can be cleaved from its transmembrane form and released into fluid, or otherwise secreted or shed from a cell.
25 Klotho RNA can also be alternatively spliced and directly secreted into the surrounding fluid (*i.e.*, without forming a transmembrane protein). Both forms are encompassed in the terms soluble klotho polypeptide, klotho polypeptide, and klotho.

[0035] As used herein, the terms “systemic” or “peripheral” refer to administration by a route that does not involve direct injection (or other administration) into the cerebrospinal
30 fluid (CSF) or central nervous system (CNS). That is, systemic and peripheral administration encompasses administration to the “blood” side of the blood-brain barrier. Examples of

systemic and peripheral routes include oral and mucosal, intravenous, intraperitoneal, intramuscular, and subcutaneous injection, and intravenous drip.

[0036] The terms “cognition,” “cognitive ability,” “cognitive function,” and like terms refer to a collection of mental tasks and functions, including but not limited to: memory (e.g., semantic, episodic, procedural, priming, or working); orientation; language; problem solving; visual perception, construction, and integration; planning; organizational skills; selective attention; inhibitory control; and ability to mentally manipulate information.

[0037] The terms “improved cognition,” “increased cognitive ability,” “improved cognitive function,” and like terms refer to an improvement in cognition under a given condition (e.g. treatment with klotho) compared to cognition absent the condition (e.g., absent treatment with klotho). For an individual experiencing cognitive decline, an improvement in cognition might be a reduction in the rate of cognitive decline (*i.e.*, an improvement compared to the absence of treatment), but not an actual improvement in cognitive ability. An increase in cognitive ability can also be an increase in brain activity in a specified area, e.g., as determined by MRI, or an inhibition of brain activity that results in better overall brain function. An increase in cognitive ability can also be improvement in a cognitive performance test as described in more detail herein. An improvement or increase in cognitive ability can be in any one cognitive aspect or function, or any combination of individual cognitive functions.

[0038] An individual in need of improved cognitive function refers to individuals with age-related cognitive decline; a neurodegenerative disease; a mental or mood disorder; traumatic brain injury; developmental delay; genetic disorder resulting in reduced cognitive ability; brain injury due to stroke, brain cancer, MS, epilepsy, radiation or chemotherapy; etc. An individual in need of improved cognitive function can also include individuals that desire increased mental function to fight the effects of stress, sleep deprivation, jet lag, or pain, or to heighten ability for a particular task. A more complete and specific list of such individuals is included in the “Cognitive conditions and disorders” section herein.

[0039] The words “protein”, “peptide”, and “polypeptide” are used interchangeably to denote an amino acid polymer or a set of two or more interacting or bound amino acid polymers. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as

to naturally occurring amino acid polymers, those containing modified residues, and non-naturally occurring amino acid polymer.

[0040] The term “amino acid” refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function similarly to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, *e.g.*, hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, *e.g.*, an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, *e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs may have modified R groups (*e.g.*, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions similarly to a naturally occurring amino acid.

[0041] Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

[0042] “Conservatively modified variants” applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical or associated, *e.g.*, naturally contiguous, sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode most proteins. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to another of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are “silent variations,” which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes silent variations of the nucleic acid. One of skill will recognize that in certain contexts each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can

be modified to yield a functionally identical molecule. Accordingly, often silent variations of a nucleic acid which encodes a polypeptide is implicit in a described sequence with respect to the expression product, but not with respect to actual probe sequences.

[0043] As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a “conservatively modified variant” where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention. The following amino acids are typically conservative substitutions for one another: 1) Alanine (A), Glycine (G); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); 7) Serine (S), Threonine (T); and 8) Cysteine (C), Methionine (M) (*see, e.g., Creighton, Proteins* (1984)).

[0044] The terms “identical” or “percent identity,” in the context of two or more nucleic acids, or two or more polypeptides, refer to two or more sequences or subsequences that are the same or have a specified percentage of nucleotides, or amino acids, that are the same (*i.e.,* about 60% identity, preferably 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or higher identity over a specified region, when compared and aligned for maximum correspondence over a comparison window or designated region) as measured using a BLAST or BLAST 2.0 sequence comparison algorithms with default parameters, or by manual alignment and visual inspection. *See e.g.,* the NCBI web site at ncbi.nlm.nih.gov/BLAST. Such sequences are then said to be “substantially identical.” This definition also refers to, or may be applied to, the complement of a nucleotide test sequence. The definition also includes sequences that have deletions and/or additions, as well as those that have substitutions. As described below, the algorithms can account for gaps and the like. Typically, identity exists over a region comprising an antibody epitope, or a sequence that is at least about 25 amino acids or nucleotides in length, or over a region that is 50-100 amino acids or nucleotides in length, or over the entire length of the reference sequence.

[0045] The term “recombinant” when used with reference, *e.g.*, to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, recombina
5 tant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all.

[0046] The term “heterologous” when used with reference to portions of a protein or nucleic acid indicates that the protein or nucleic acid comprises two or more subsequences
10 that are not found in the same relationship to each other in nature. For instance, the protein or nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, *e.g.*, a promoter from one source and a coding region from another source, or functional chimeric protein.

[0047] The terms “agonist,” “activator,” “inducer” and like terms refer to an agent that
15 increases activity or expression (*e.g.*, of klotho or a klotho signaling pathway) as compared to a control. Agonists are agents that, *e.g.*, stimulate, increase, activate, enhance activation, sensitize or upregulate the activity of klotho or a klotho signaling pathway. The expression or activity can be increased 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% 100% or more than that in a control. In certain instances, the activation is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-
20 fold, 10-fold, or more in comparison to a control.

[0048] The terms “inhibitor,” “repressor” or “antagonist” or “downregulator” interchangeably refer to a substance that results in a detectably lower expression or activity level as compared to a control. The inhibited expression or activity can be 10%, 20%, 30%,
40%, 50%, 60%, 70%, 80%, 90% or less than that in a control. In certain instances, the
25 inhibition is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, or more in comparison to a control.

[0049] A “control” sample or value refers to a sample that serves as a reference, usually a known reference, for comparison to a test sample. For example, a test sample can be taken from a test condition, *e.g.*, in the presence of a test compound, and compared to samples from
30 known conditions, *e.g.*, in the absence of the test compound (negative control), or in the presence of a known compound (positive control). A control can also represent an average value gathered from a number of tests or results. One of skill in the art will recognize that

controls can be designed for assessment of any number of parameters. For example, a control can be devised to compare therapeutic benefit based on pharmacological data (*e.g.*, half-life) or therapeutic measures (*e.g.*, comparison of benefit and/or side effects). Controls can be designed for *in vitro* applications. One of skill in the art will understand which controls are valuable in a given situation and be able to analyze data based on comparisons to control values. Controls are also valuable for determining the significance of data. For example, if values for a given parameter are widely variant in controls, variation in test samples will not be considered as significant.

[0050] A “label” or a “detectable moiety” is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, chemical, or other physical means. For example, useful labels include ^{32}P , fluorescent dyes, electron-dense reagents, enzymes (*e.g.*, as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins or other entities which can be made detectable, *e.g.*, by incorporating a radiolabel into a peptide or antibody specifically reactive with a target peptide. Any method known for conjugating a protein to the label may be employed, *e.g.*, using methods described in Hermanson, Bioconjugate Techniques 1996, Academic Press, Inc., San Diego.

[0051] A “labeled” molecule (*e.g.*, klotho polypeptide) is one that is bound, either covalently, through a linker or a chemical bond, or noncovalently, through ionic, van der Waals, electrostatic, or hydrogen bonds to a label such that the presence of the molecule may be detected by detecting the presence of the label bound to the molecule.

[0052] The term “diagnosis” refers to a relative probability that a disorder is present in an individual. Similarly, the term “prognosis” refers to a relative probability that a certain future outcome may occur in the individual. For example, in the context of the present disclosure, prognosis can refer to the likelihood that an individual suffer cognitive decline, or the likely severity of the disease (*e.g.*, severity of symptoms, rate of functional decline, etc.). The terms are not intended to be absolute, as will be appreciated by any one of skill in the field of medical diagnostics.

[0053] A “biological sample” can be obtained from a patient, *e.g.*, a biopsy, from an animal, such as an animal model, or from cultured cells, *e.g.*, a cell line or cells removed from a patient and grown in culture for observation. Biological samples include tissues and bodily fluids, *e.g.*, cerebrospinal fluid (CSF), blood, blood fractions, lymph, saliva, urine, feces, etc.

[0054] The terms “therapy,” “treatment,” and “amelioration” refer to any reduction in the severity of symptoms (cognitive decline), or improvement in cognitive function, or where motor function is affected, an improvement in motor function. As used herein, the terms “treat” and “prevent” are not intended to be absolute terms. Treatment and prevention can refer to any delay in cognitive decline, amelioration of symptoms (*e.g.*, confusion, delirium), etc. Treatment and prevention can be complete or partial, such that cognition is better than would be expected without treatment (*e.g.*, compared to cognition in the same individual before treatment or compared to cognition in similar non-treated individuals). The effect of treatment can be compared to an individual or pool of individuals not receiving the treatment, or to the same patient prior to treatment or at a different time during treatment. In some aspects, cognition is improved by at least 1%, as compared, *e.g.*, to the individual before administration or to a control individual not undergoing treatment. In some embodiments, cognition is improved by at least 2, 3, 5, 7, 10, 15, 20, 25%, 50%, 75%, 80%, or 90%, or more, determined using tests of cognition, molecular proxies, or structural changes associated with brain function. In some aspects, motor function is improved by at least 1%, as compared, *e.g.*, to the individual before administration or to a control individual not undergoing treatment. In some embodiments, motor function is improved by at least 2, 3, 5, 7, 10, 15, 20, 25%, 50%, 75%, 80%, or 90%, or more, determined using tests of motor function.

[0055] The terms “effective amount,” “effective dose,” “therapeutically effective amount,” etc. refer to that amount of the therapeutic agent sufficient to ameliorate a disorder, as described above. For example, for the given parameter, a therapeutically effective amount will show an increase or decrease of therapeutic effect at least 1%, 2%, 5%, 10%, 15%, 20%, 25%, 40%, 50%, 60%, 75%, 80%, 90%, or at least 100%. Therapeutic efficacy can also be expressed as “-fold” increase or decrease. For example, a therapeutically effective amount can have at least a 1.2-fold, 1.5-fold, 2-fold, 5-fold, or more effect over a control.

[0056] As used herein, the term “pharmaceutically acceptable” is used synonymously with physiologically acceptable and pharmacologically acceptable. A pharmaceutical composition will generally comprise agents for buffering and preservation in storage, and can include buffers and carriers for appropriate delivery, depending on the route of administration.

[0057] The terms “dose” and “dosage” are used interchangeably herein. A dose refers to the amount of active ingredient given to an individual at each administration. For the present

invention, the dose refers to the amount of Klotho polypeptide. The dose will vary depending on a number of factors, including frequency of administration; size and tolerance of the individual; type and severity of the condition; risk of side effects; and the route of administration. One of skill in the art will recognize that the dose can be modified depending on the above factors or based on therapeutic progress. The term “dosage form” refers to the particular format of the pharmaceutical, and depends on the route of administration. For example, a dosage form can be in a liquid, *e.g.*, a saline solution for injection.

[0058] “Subject,” “patient,” “individual” and like terms are used interchangeably and refer to, except where indicated, mammals such as humans and non-human primates, as well as dogs, horses, pigs, mice, rats, and other mammalian species. The term does not necessarily indicate that the subject has been diagnosed with a particular disease, but typically refers to an individual under medical supervision. A patient can be an individual that is seeking treatment, monitoring, adjustment or modification of an existing therapeutic regimen, etc.

III. Klotho

[0059] Klotho is a pleiotropic protein and an aging regulator that circulates throughout the body and brain (Imura *et al.* (2004) *FEBS Letters* 565:143; Kurosu *et al.* (2005) *Science* 309:1829). Human Klotho is described in GenBank Accession No. NC_000013 and Uniprot Accession No. Q9UEF7. A number of species homologs exist, including mouse and rat Klotho which share 86% and 85% identity with the human Klotho polypeptide, which is shown as SEQ ID NO:1. It exists in a transmembrane form that can be cleaved such that the extracellular portion (amino acids 34-979) is released as a hormone (Shiraki-ika *et al.* (1998) *FEBS Letters* 424:6). Klotho also has a splice variant that results in a 549 amino acid secreted form of the protein that is also functional (Wang and Sun (2009) *Ageing Res. Rev.* 8:43). Both cleaved and secreted klotho are soluble and functional in the body, but have a sequence variation at the C-terminal end due to the splice variation. Amino acids 535-549 are DTTLSQFTDLNVYLW (SEQ ID NO:2) for cleaved, soluble human Klotho and SQLTKPISSLTKPYH (SEQ ID NO:3) for spliced, soluble human Klotho. Full length soluble Klotho includes two conserved domains (KL1 and KL2) with homology to beta glycosidase proteins. The conserved beta-glucosidase/ 6-phospho-beta-glucosidase/ beta-galactosidase motif spans 62-497 in the human protein and 64-499 in the mouse. The conserved KL1 sequence is described in Chateau *et al.* (2010) *Aging* 2:567 and Matsumura *et al.* (1998) *Biochem Biophys Res Commun*, and comprises amino acids 34-549 of the human Klotho protein, with the glycosyl hydrolase consensus region spanning amino acids 57-506 of

the human Klotho protein (59-508 of the mouse). Klotho does not have beta-glycosidase activity, but shows some beta-glucuronidase activity.

[0060] Klotho suppresses insulin and wnt signaling, regulates ion channels and their transport, and promotes function of FGF23. *See, e.g., Chang et al. (2005) Science 310:490; Imura et al. (2007) Science 316:1615; Kurosu (2005); Liu et al. (2007) Science 317:803; and Urakawa et al. (2006) Nature 444:770).*

[0061] In mice, transgenic overexpression of klotho extends lifespan and associates with better cognitive functions in the normal and diseased brain (Kurosu (2005); Dubal *et al.* (2014) *Cell Reports* 7:1065; Dubal *et al.* (2015) *J. Neuroscience*). In humans, a single allele of the KL-VS variant of the *KLOTHO* gene, which increases secreted klotho promotes longevity (Arking *et al.*, 2002; Arking *et al.*, 2005; Invidia *et al.*, 2010) and also associates with better baseline cognitive functions in aging populations. *See, e.g., Arking et al. (2002) PNAS 99:856; Arking et al. (2005) Circ. Res. 96:412; Dubal et al (2014) Cell Reports 7:1065; Yokoyama et al. (2015) Ann. Clin. Translational Neurology 2:215.*

[0062] Klotho polypeptides that can be used for administration include species homologs (*e.g.*, non-human primate, mouse, rat), allelic variants, functional fragments, and functional variants of the wild type sequence that retain cognition improving activity. Examples include secreted Klotho, fragments comprising the KL1 domain, fragments comprising the KL2 domain, fragments comprising the KL1 and KL2 domains, variants comprising the KL1 domain with at least one (*e.g.*, 1-20, 5-50, 25-100) non-conserved amino acid in the KL1 domain substituted with a different amino acid or deleted, variants comprising the KL2 domain with at least one non-conserved amino acid in the KL2 domain substituted with a different amino acid.

[0063] Functional fragments of the Klotho polypeptide that can be used as described herein include the extracellular domain (*e.g.*, corresponding to or substantially identical or similar to amino acids 34-979 of human Klotho), secreted Klotho (*e.g.*, corresponding to or substantially identical or similar to 549 amino acid form), a KL1 domain (*e.g.*, corresponding to or substantially identical or similar to amino acids 34-549 of human Klotho), a glycosyl hydrolase consensus sequence (*e.g.*, corresponding to or substantially identical or similar to amino acids 57-506 of human Klotho), or a beta-glucosidase/6-phospho-beta-glucosidase/beta-galactosidase consensus sequence (*e.g.*, corresponding to or substantially identical or similar to amino acids 62-497 of human Klotho). In some embodiments, the

Klotho polypeptide comprises or is substantially identical or similar to amino acids 34-549 of human Klotho. In some embodiments, the Klotho polypeptide is part of a larger fusion protein. In some embodiments, the fusion protein comprises the Klotho polypeptide as described herein and further comprises no more than 100, 75, 50, or 30 additional amino acids. In some embodiments, the Klotho polypeptide is not fused to a Fibroblast growth factor (FGF). In some embodiments, the Klotho polypeptide comprises (e.g., is fused to) an affinity tag (e.g., a histidine tag) or a conjugate to increase stability or half-life *in vivo*.

[0064] A functional variant or fragment of Klotho is a variant or fragment that retains any klotho activity, e.g., at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of the level of any activity of soluble klotho. Soluble klotho activities include those described above, and include binding FGF-23, binding to FGFR1c, beta-glucuronidase activity, suppression of wnt signaling, suppression of insulin signaling, suppression of TGF-beta 1 activity, increasing GluN2B expression and/or synaptic localization, c-fos induction. Additional Klotho activities include causing changes in magnetic resonance imaging (MRI) brain scans, e.g., functional MRI, electroencephalograph (EEG), and transcranial magnetic and electrical stimulation (TMS and TES); and improved performance in neuropsychologic testing and cognitive ability.

IV. Administration of Klotho

[0065] Provided herein are methods of improving cognition and/or motor function in an individual comprising administering Klotho to the individual. In some embodiments, the method of treatment comprises administering to an individual an effective amount of a Klotho polypeptide (or functional variant or fragment thereof). In some embodiments, the treatment is prophylactic, e.g., for individual expecting stress (e.g., jet lag, military performance) or to prevent cognitive decline associated with aging. In some embodiments, the individual has been diagnosed with a cognitive disorder. In some embodiments, the individual is receiving or has received therapy for a cognitive disorder or for a condition that is related to cognitive function (e.g., cognitive decline in response to chemotherapy).

[0066] In some embodiments, the method further comprises monitoring the individual for cognitive ability, either through a molecular proxy (e.g., changes NMDA receptor or c-fos activation, or GluN2B levels in the brain), changes in MRI brain scans (e.g., functional MRI), changes in EEG, changes in TMS and TES, changes in neuropsychologic test scores, or tests of cognitive ability (e.g., for learning, short or long term memory, executive functions,

language ability, and visuospatial function). In some embodiments, the individual is monitored using more than one of the above tests in any combination. In some embodiments, the dose of the Klotho polypeptide for each administration is determined based on the therapeutic progress of the individual, *e.g.*, where a higher dose is administered if the individual is not responding sufficiently to therapy.

[0067] In some embodiments, the Klotho polypeptide is administered in a pharmaceutical composition with a physiologically (*i.e.*, pharmaceutically) acceptable carrier. The term “carrier” refers to a typically inert substance used as a diluent or vehicle for a diagnostic or therapeutic agent. The term also encompasses a typically inert substance that imparts cohesive qualities to the composition. Physiologically acceptable carriers can be liquid, *e.g.*, physiological saline, phosphate buffer, normal buffered saline (135-150 mM NaCl), water, buffered water, 0.4% saline, 0.3% glycine, glycoproteins to provide enhanced stability (*e.g.*, albumin, lipoprotein, globulin, etc.), and the like. Since physiologically acceptable carriers are determined in part by the particular composition being administered as well as by the particular method used to administer the composition, there are a wide variety of suitable formulations of pharmaceutical compositions of the present invention (*See, e.g.*, Remington's Pharmaceutical Sciences, 17th ed., 1989).

[0068] The presently described compositions can be sterilized by conventional, well-known sterilization techniques or may be produced under sterile conditions. Aqueous solutions can be packaged for use or filtered under aseptic conditions and lyophilized, the lyophilized preparation being combined with a sterile aqueous solution prior to administration. The compositions can contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents, and the like, *e.g.*, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, and triethanolamine oleate. Sugars can also be included for stabilizing the compositions, such as a stabilizer for lyophilized antibody compositions.

[0069] Dosage forms can be prepared for mucosal (*e.g.*, nasal, sublingual, vaginal, buccal, or rectal), parenteral (*e.g.*, subcutaneous, intravenous, intramuscular, or intraarterial injection, either bolus or infusion), oral, or transdermal administration to a patient. Examples of dosage forms include, but are not limited to: dispersions; suppositories; ointments; cataplasms (poultices); pastes; powders; dressings; creams; plasters; solutions; patches; aerosols (*e.g.*,

nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (*e.g.*, aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; and sterile solids (*e.g.*, crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

[0070] Injectable compositions can comprise a solution of the Klotho polypeptide suspended in an acceptable carrier, such as an aqueous carrier. Any of a variety of aqueous carriers can be used, *e.g.*, water, buffered water, 0.4% saline, 0.9% isotonic saline, 0.3% glycine, 5% dextrose, and the like, and may include glycoproteins for enhanced stability, such as albumin, lipoprotein, globulin, etc. In some embodiments, normal buffered saline (135-150 mM NaCl) is used. The compositions can contain pharmaceutically acceptable auxiliary substances to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents, *e.g.*, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, etc.

[0071] Formulations suitable for parenteral administration, such as, for example, by intraarticular (in the joints), intravenous, intramuscular, intradermal, intraperitoneal, and subcutaneous routes, include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. Injection solutions and suspensions can also be prepared from sterile powders, granules, and tablets. In some embodiments, the composition is administered by intravenous infusion, topically, intraperitoneally, intravesically, or intrathecally. The Klotho polypeptide formulation can be provided in unit-dose or multi-dose sealed containers, such as ampoules and vials.

[0072] The Klotho polypeptide composition, alone or in combination with other suitable components, can be made into aerosol formulations (“nebulized”) to be administered via inhalation. Aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, and nitrogen.

[0073] The pharmaceutical preparation can be packaged or prepared in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component, *e.g.*, according to the dose of Klotho polypeptide. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation. The composition can, if desired, also contain other compatible therapeutic agents. In some embodiments, the Klotho polypeptide composition can be formulated in a kit for administration.

[0074] In some embodiments, a pharmaceutical composition comprising a klotho polypeptide is administered orally. In some embodiments, a pharmaceutical composition comprising a klotho polypeptide is administered mucosally, *e.g.*, nasally. In some embodiments, a pharmaceutical composition comprising a klotho polypeptide is administered by injection, *e.g.*, subcutaneous, intraperitoneal, intravenous, or intramuscular. In some embodiments, a pharmaceutical composition comprising a klotho polypeptide is administered by infusion, *e.g.*, using a reservoir or osmotic minipump.

[0075] An example of administration of a pharmaceutical composition includes storing the Klotho polypeptide at 10 mg/ml in sterile isotonic aqueous saline solution at 4°C, and diluting it in an appropriate solution for injection prior to administration to the patient. In some embodiments, the Klotho polypeptide composition can be administered by intravenous infusion over the course of 0.25-2 hours. In some embodiments, the administration procedure is via bolus injection.

[0076] In therapeutic use, the Klotho polypeptide can be administered at the initial dosage of about 0.1 µg/kg to about 1000 µg/kg daily and adjusted over time. A daily dose range of about 1 µg/kg to about 500 µg/kg, or about 10 µg/kg to about 100 µg/kg, or about 30 µg/kg to about 50 µg/kg can be used. The dosage is varied depending upon the requirements of the patient, the severity of the condition being treated, and the route of administration. For example, for injection of Klotho polypeptide, the effective dose is typically in the range of 10-100 µg/kg, while for direct delivery to the central nervous system (CNS), the effective dosage is lower, *e.g.*, 5-30 µg/kg. For oral administration, the effective dose is higher, *e.g.*, in the range of 50-10,000 µg/kg (*e.g.*, 100µg/kg-2mg/kg). The dose is chosen in order to provide effective therapy for the patient. The dose may be repeated at an appropriate frequency which may be in the range of once or twice per day, once or twice per week to once every three months, depending on the pharmacokinetics of the Klotho polypeptide

composition (*e.g.*, half-life in the circulation) and the pharmacodynamic response (*e.g.*, the duration of the therapeutic effect).

[0077] Administration can be periodic. Depending on the route of administration, the dose can be administered, *e.g.*, once every 1, 3, 5, 7, 10, 14, 21, or 28 days or longer (*e.g.*, once every 2, 3, 4, or 6 months). In some cases, administration is more frequent, *e.g.*, 2 or 3 times per day. The patient can be monitored to adjust the dosage and frequency of administration depending on therapeutic progress and any adverse side effects, as will be recognized by one of skill in the art.

[0078] Dosages can be empirically determined considering the type and severity of cognitive condition diagnosed in a particular patient. The dose administered to a patient, in the context of the present disclosure, should be sufficient to affect a beneficial therapeutic response in the patient over time. The size of the dose will also be determined by the existence, nature, and extent of any adverse side-effects that accompany the administration of any particular composition in a particular patient, as will be recognized by the skilled practitioner.

[0079] In some embodiments, the Klotho polypeptide composition is administered to an (*e.g.*, human) individual having at least normal cognitive function. As described herein, it has been surprisingly discovered that not only can Klotho improve cognition in individuals with impaired cognition, Klotho can also improve cognition of individuals with at least normal cognition. Thus in some embodiments, the individual receiving the Klotho polypeptide composition begins initially with at least normal cognition and following administration of the Klotho polypeptide composition attains improved cognition compared to the initial level of cognition. The level of cognition of an individual can be determined as is known in the art. Normal cognitive functions are determined by scores from sets of cognitive tests that are compiled into global cognitive scores, as described in Dubal DB et al. (2014) *Cell Reports* 7:1065-1076. Such cognition tests include tests of executive function and working memory such as Trails A and Trails B (Dubal DB et al. (2014) *Cell Reports* 7:1065-1076). In some embodiments, administration of Klotho results in an improvement of cognition (whether initially at least normal or impaired), by at least 5%, 10%, 20% or more.

In some embodiments, administration results in improved motor function. In some embodiments, the Klotho polypeptide composition is administered to an (*e.g.*, human) individual having impaired motor function. For example, in some embodiments, the

individual has stroke to the brain or spinal cord (ischemic or hemorrhagic), neurodegenerative disease (Parkinson's disease, Lewy body dementia, multiple system atrophy, amyotrophic lateral sclerosis, prion disorder, Huntington's disease, supranuclear palsy), Parkinsonism, traumatic brain injury, neuroinfectious brain lesions, multiple sclerosis
 5 and related autoimmune and demyelinating disease, spinal cord lesions (compressive, infectious, toxic or metabolic, autoimmune, oncologic), brain tumor, epilepsy, paraneoplastic disorder, neurodevelopmental disorder (mitochondrial, autosomal genetic), muscle disease (polymyositis, dermatomyositis, inclusion body myositis, infectious, endocrine, metabolic, toxic, congenital myopathy, congenital muscular dystrophy, hereditary), neuropathies
 10 (Guillain-Barre syndrome, axonal and demyelinating, diabetic, toxic, metabolic, infectious, critical illness, entrapment), tick paralysis, myasthenia gravis, and spinal muscular atrophy. Changes in motor function can be assayed as known in the art. Exemplary motor function assays include but are not limited to electromyogram and nerve conduction studies, direct or device-assisted clinical testing of strength, tone, and muscle bulk, reflex examination,
 15 coordination examination, and gait analysis. Assays for testing etiologies causing deficits of motor function include but are not limited to magnetic resonance imaging of the central nervous system, muscle biopsy, nerve biopsy, and laboratory studies.

[0080] Thus in some embodiments, additional administration is dependent on patient progress, *e.g.*, the patient is monitored between administrations. For example, after the first
 20 administration or round of administrations, the patient can be monitored for cognitive ability or for side effects, *e.g.*, weakness, dizziness, nausea, etc.

[0081] In some embodiments, the individual has a chronic condition, so that klotho is administered over an indefinite period, *e.g.*, for the lifetime of the patient. In such cases, administration is typically periodic. Diseases that are considered long-term or chronic
 25 include, but are not limited to Alzheimer's disease, Parkinson's disease, Huntington's disease, and cognitive decline associated with hypertension and heart disease.

[0082] In some embodiments, the Klotho polypeptide is linked to a stabilizing moiety such as PEG, glycosylation, or a liposome or other nanocarrier. US Patent Nos. 4,732,863 and 7892554 and Chattopadhyay *et al.* (2010) *Mol Pharm* 7:2194 describe methods for attaching
 30 a polypeptide to PEG, PEG derivatives, and nanoparticles (*e.g.*, liposomes). Liposomes containing phosphatidyl-ethanolamine (PE) can be prepared by established procedures as described herein. The inclusion of PE provides an active functional site on the liposomal

surface for attachment. In some embodiments, the Klotho polypeptide is linked to an affinity tag, *e.g.*, a histidine tag (*e.g.*, 4-16 histidine residues), streptavidin, or an antibody target.

[0083] The Klotho polypeptide can also be formulated as a sustained-release preparation (*e.g.*, in a semi-permeable matrices of solid hydrophobic polymers (*e.g.*, polyesters, hydrogels (for example, poly (2-hydroxyethyl-methacrylate), or poly (vinylalcohol)), polylactides. The Klotho polypeptide can be entrapped in a nanoparticle prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin microcapsules and poly- (methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nanoparticles and nanocapsules) or in macroemulsions.

[0084] In some embodiments, the Klotho polypeptide is labeled, *e.g.*, for tracking in the body or *ex vivo*. The Klotho polypeptide can be labeled any diagnostic agent known in the art, as provided, for example, in the following references: Armstrong *et al.*, *Diagnostic Imaging*, 5th Ed., Blackwell Publishing (2004); Torchilin, V. P., Ed., *Targeted Delivery of Imaging Agents*, CRC Press (1995); Vallabhajosula, S., *Molecular Imaging: Radiopharmaceuticals for PET and SPECT*, Springer (2009). The diagnostic agent can be detected by a variety of ways, including as an agent providing and/or enhancing a detectable signal. Detectable signals include, but are not limited to, gamma-emitting, radioactive, echogenic, optical, fluorescent, absorptive, magnetic, or tomography signals. Techniques for imaging the diagnostic agent can include, but are not limited to, single photon emission computed tomography (SPECT), magnetic resonance imaging (MRI), optical imaging, positron emission tomography (PET), computed tomography (CT), x-ray imaging, gamma ray imaging, and the like. The terms “detectable agent,” “detectable moiety,” “label,” “imaging agent,” and like terms are used synonymously herein.

[0085] In some embodiments, the label can include optical agents such as fluorescent agents, phosphorescent agents, chemiluminescent agents, and the like. Numerous agents (*e.g.*, dyes, probes, labels, or indicators) are known in the art and can be used in the present invention. (*See, e.g.*, Invitrogen, *The Handbook—A Guide to Fluorescent Probes and Labeling Technologies*, Tenth Edition (2005)). Fluorescent agents can include a variety of organic and/or inorganic small molecules or a variety of fluorescent proteins and derivatives thereof. For example, fluorescent agents can include but are not limited to cyanines, phthalocyanines, porphyrins, indocyanines, rhodamines, phenoxazines, phenylxanthenes,

phenothiazines, phenoselenazines, fluoresceins, benzoporphyrins, squaraines, dipyrrolo
pyrimidones, tetracenes, quinolines, pyrazines, corrins, croconiums, acridones,
phenanthridines, rhodamines, acridines, anthraquinones, chalcogenopyrylium analogues,
chlorins, naphthalocyanines, methine dyes, indolenium dyes, azo compounds, azulenes,
5 azaazulenes, triphenyl methane dyes, indoles, benzoindoles, indocarbocyanines,
benzoindocarbocyanines, and BODIPY™ derivatives. Fluorescent dyes are discussed, for
example, in U.S. Pat. No. 4,452,720, U.S. Pat. No. 5,227,487, and U.S. Pat. No. 5,543,295.

[0086] The label can also be a radioisotope, *e.g.*, radionuclides that emit gamma rays,
positrons, beta and alpha particles, and X-rays. Suitable radionuclides include but are not
10 limited

to ^{225}Ac , ^{72}As , ^{211}At , ^{11}B , ^{128}Ba , ^{212}Bi , ^{75}Br , ^{77}Br , ^{14}C , ^{109}Cd , ^{62}Cu , ^{64}Cu , ^{67}Cu , ^{18}F , ^{67}Ga , ^{68}Ga ,
 ^3H ,

^{166}Ho , ^{123}I , ^{124}I , ^{125}I , ^{130}I , ^{131}I , ^{111}In , ^{177}Lu , ^{13}N , ^{15}O , ^{32}P , ^{33}P , ^{212}Pb , ^{103}Pd , ^{186}Re , ^{188}Re , ^{47}Sc , ^{153}Sm , ^{89}Sr , $^{99\text{m}}\text{Tc}$, ^{88}Y and ^{90}Y . In some embodiments, radioactive agents can include ^{111}In -

15 DTPA, $^{99\text{m}}\text{Tc}(\text{CO})_3\text{-DTPA}$, $^{99\text{m}}\text{Tc}(\text{CO})_3\text{-ENPy2}$, $^{62/64/67}\text{Cu-TETA}$, $^{99\text{m}}\text{Tc}(\text{CO})_3\text{-IDA}$,
and $^{99\text{m}}\text{Tc}(\text{CO})_3$ triamines (cyclic or linear). In some embodiments, the agents can include
DOTA and its various analogs with ^{111}In , ^{177}Lu , ^{153}Sm , $^{88/90}\text{Y}$, $^{62/64/67}\text{Cu}$, or $^{67/68}\text{Ga}$. In some
embodiments, a nanoparticle can be labeled by incorporation of lipids attached to chelates,
such as DTPA-lipid, as provided in the following references: Phillips *et al.*, *Wiley*

20 *Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*, 1(1): 69-83 (2008);
Torchilin, V.P. & Weissig, V., Eds. *Liposomes 2nd Ed.*: Oxford Univ. Press (2003);
Elbayoumi, T.A. & Torchilin, V.P., *Eur. J. Nucl. Med. Mol. Imaging* 33:1196–1205 (2006);
Mougin-Degraef, M. *et al.*, *Int'l J. Pharmaceutics* 344:110-117 (2007).

[0087] In some embodiments, the diagnostic agent can be associated with a secondary
25 binding ligand or to an enzyme (an enzyme tag) that will generate a colored product upon
contact with a chromogenic substrate. Examples of suitable enzymes include urease, alkaline
phosphatase, (horseradish) hydrogen peroxidase and glucose oxidase. Secondary binding
ligands include, *e.g.*, biotin and avidin or streptavidin compounds as known in the art.

V. Cognitive conditions and disorders

30 **[0088]** Klotho polypeptides (and functional variants and fragments thereof) can be
administered to improve cognition for a number of conditions and situations. This includes

treatment of individuals with lower than normal or declining cognitive ability, or prophylactic treatment of individuals in need of improved or increased cognitive ability.

[0089] Klotho polypeptides (and functional variants and fragments thereof) can be used to prevent or reduce cognitive decline associated with aging, *e.g.* in individuals 50 years of age or older, or upon initial signs of cognitive decline.

[0090] Klotho polypeptides (and functional variants and fragments thereof) can also be used to treat individuals with age-related, non-age related, or disease related conditions including, but not limited to:

Neurodegenerative diseases and dementia: Alzheimer's disease, Parkinson's disease, Huntington's disease, frontotemporal dementia, progressive supranuclear palsy, corticobasalar degeneration, mild cognitive impairment, vascular dementia, Lewy body dementia, amyotrophic lateral sclerosis, prion disorder, HIV-related dementia;

Mental or mood disorders: depression, schizophrenia, attention deficit/ hyperactivity disorder, autism spectrum disorder, intellectual disability, a mood disorder, and a psychotic disorder;

Childhood neurodevelopmental syndromes and brain tumors: X-linked mental disability or retardation, astrocytoma, ependymoma, medulloblastoma, oligodendroglioma;

Genetic syndromes affecting learning: Down's syndrome, Angelman's syndrome, Rett's syndrome;

Metabolic disorders affecting cognition: phenylketonuria, Lesch-Nyhan, galactosemia, and adrenoleukodystrophy;

Cognitive decline associated with chemotherapy and/or radiation therapy; and

Additional conditions and disorders: pain-associated cognitive effects, traumatic brain injury, stroke, multiple sclerosis, neuroautoimmune disease, epilepsy, delirium, paraneoplastic disorder, developmental delay, and leukodystrophies.

[0091] Klotho polypeptides (and functional variants and fragments thereof) can be also be administered to provide increased cognition for individuals desiring improved cognition, *e.g.*, individuals exposed to stress, sleep deprivation, or jet lag, or for individuals requiring superior cognitive function, such as surgeons, air-traffic controllers, and military personnel. In such cases, the klotho polypeptide composition can be administered 2-24 hours before the

desired effect, which can last about 3-5 days for working memory and about 2 weeks for spatial memory.

[0092] Cognitive ability can be measured using any method known in the art, *e.g.*, for testing memory, language ability, executive functions, visuospatial function, dementia, or multi-parameter neuropsychological abilities. In some embodiments, Klotho administration results in at least a 1%, 2%, 5%, 7%, 10%, 15%, 20%, 30%, 50%, or greater improvement in score on a standard cognitive ability test (*e.g.*, measured 1-3 days after administration). In some embodiments, the testing is carried out more than once for an individual, *e.g.*, one or more time over the course of treatment with Klotho.

[0093] For example, standard tests for memory and learning can be applied, *e.g.*, to determine semantic, episodic, procedural, priming, and/or working (*i.e.*, short term) memory. Common tests include Cambridge prospective memory test (CAMPROMPT), memory assessment scales (MAS), Rey auditory verbal learning test, Rivermead behavioral memory test, Test of memory and learning (TOMAL), Wechsler memory scale (WMS), and Test of memory malingering (TOMM). Tests for language functions include, *e.g.*, Boston Diagnostic Aphasia Examination (BDAE), Comprehensive aphasia test (CAT), and Multilingual aphasia examination (MAE).

[0094] Executive function (*e.g.*, problem solving, planning, organization, inhibitory control) can be tested using Behavioral assessment of dysexecutive syndrome (BADS), CNS vital signs (Brief Core Battery), Controlled oral word association test (COWAT), Delis-Kaplan Executive Function System (D-KEFS), Digit vigilance test, Kaplan Baycrest neurocognitive assessment (KBNA), Hayling and Brixton tests, Tests of variables of attention (TOVA), Wisconsin card sorting test (WCST), or Test of everyday attention (TEA). Visuospatial ability (*e.g.*, visual perception, construction and integration) can be tested using the Clock Test, Hooper visual organization task (VOT), or Rey-Osterrieth complex figure tests. Dementia can be quantified using the clinical dementia rating or dementia rating scale.

[0095] Multi-parameter tests for neuropsychological function (*e.g.*, cognitive function) include but are not limited to the Barcelona neuropsychological test (BNT), Cambridge neuropsychological test automated battery (CANTAB), Cognistat, Cognitive assessment screening instrument (CASI), Cognitive function scanner (CFS), Dean-Woodcock neuropsychology assessment system (DWNAS), General practical assessment of cognition

(GPCOG) Mini mental state examination (MMSE), NEPSY, or the CDR computerized assessment system.

[0096] Alternatively, cognition can be determined using structural or molecular proxies for cognitive activity, *e.g.*, compared over time to detect changes. Cognitive changes can be detected, *e.g.*, by observing changes to brain structure, connectivity, activation, inhibition, or synaptic plasticity, *e.g.*, by MRI, fMRI, EEG, TMS and TES, and/or any combination of these. In some embodiments, brain activity is observed. In some embodiments, Klotho administration results in a 1.5-fold, 2-fold, 5-fold, 7-fold, 10-fold, or greater increase in brain activity (*e.g.*, measured 1-3 days after administration). Molecular proxies for improved cognition include, but are not limited to: increased levels of GluN2B, increased GluN2B synaptic localization, increased NMDA receptor activation, and/or increased c-fos activation in the brain. These measures are particularly relevant to cognition. Such method can include, *e.g.*, obtaining a sample of neuronal tissue or CSF from an individual and using standard assays to determine gene expression or activation.

[0097] Similarly, in mice and other non-human animals, cognitive ability can be tested with measures of executive function (working memory, attention, processing speed, set shifting), visiospatial learning and memory, object memory, pattern recognition, fear memory, passive avoidance memory, habituation, and novel object recognition, for example. Common tests include but are not limited to the Morris water maze, Barnes maze, radial arm water maze, y-maze, T-maze, and open field habituation. Brain imaging techniques are similarly applicable.

VI. Examples

Example 1:

A. Materials and Methods

[0098] *Mice.* All mice were on a C57BL/6J background and were kept on a 12-h light/dark cycle with *ad libitum* access to food (Picolab Rodent Diet 20, Labdiet) and water. The standard housing group was five mice per cage except for single housing during water maze studies. All cognitive and behavioral studies were carried out during the light cycle. All studies were conducted in a blinded manner.

[0099] *Treatments.* Vehicle or klotho (recombinant mouse klotho, amino acids 35-982, with a C-terminal His tag, R&D Systems) was injected intraperitoneally (i.p.) before behavioral testing of mice.

[0100] *Morris water maze cognitive behavior test.* Water maze studies were carried out as described in Dubal *et al.* (2014) *Cell Reports* 7:1065; Dubal *et al.* (2015) *Journal of Neuroscience* 35:2358. Mice were treated with either vehicle or klotho (10µg/kg) i.p. 4h prior to testing daily for 5 days. Briefly, mice were tested in a pool (diameter, 122 cm) with white, opaque water (21°± 1°C). A square, 14-cm² platform was 2 cm below the surface. Before hidden platform training, mice underwent two pre-training trials by swimming through a channel to mount a hidden platform. Over the course of hidden platform training, the platform location stayed consistent while the drop location was varied between trials. Mice underwent two training sessions, consisting of two trials of 60 s each, daily for four days. For the probe trial testing, the platform was removed and the mice were allowed to swim for 60 s. After 1h and 24h probe trials, mice were tested for their ability to find a visible platform marked with a cue (15-cm pole on the platform) over two sessions. As part of the studies, swim velocities were also recorded.

[0101] *Y-maze cognitive behavior test.* Y-maze studies were carried out as described in Dubal *et al.* (2014) *Cell Reports* 7:1065. Mice were treated with either vehicle or klotho (10µg/kg) i.p. 18h prior to testing. Briefly, mice were acclimated to the room 30 min prior to testing. Then, mice were placed in one arm of the Y-maze (three identical arms, 120° apart) and explored for 4 min. Arm entries were recorded and an alternation was noted any time the mouse entered each of the three arms in successive arm entries; chance alternation was 22%. The apparatus was cleaned with 70% alcohol between testing sessions. Percent alternation was calculated from recorded data.

B. Systemic klotho delivery enhances working memory in young mice

[0102] To investigate whether therapeutic delivery of klotho can enhance cognition, we injected mice with recombinant klotho (i.p.) and 18h later, tested working memory in the Y-maze. Compared to vehicle-treated mice, klotho-treated mice showed more alternations, indicating superior working memory (FIG. 1A). Thus, systemic klotho delivery enhanced working memory, a process that involves frontal cortical brain regions.

C. Systemic klotho delivery enhances spatial learning in young mice

[0103] We next tested whether klotho treatment enhances spatial learning in the Morris water maze. We injected mice with recombinant klotho (i.p.) 4h prior to testing daily for five days in a group of mice (Cohort 1). Compared to vehicle-treated mice, klotho-treated mice performed better in spatial learning of the hidden platform location (FIG. 1B). Both groups swam at equal speeds (FIG. 1C). In an independent group of mice (Cohort 2), we injected recombinant klotho (i.p.) 18h prior to testing daily. Again, compared to vehicle-treated mice, klotho-treated mice performed better in spatial learning (FIG. 1D). Thus, systemic klotho delivery enhanced spatial learning, a process that involves frontal cortical and hippocampal brain regions.

D. Systemic klotho delivery enhances spatial memory in young mice

[0104] To determine whether klotho treatment can enhance spatial memory, we removed the platform and performed probe testing following hidden training in the Morris water maze. In probe trials of Cohort 1, spatial memory retention was assessed by measuring the affinity of mice for the target area. In probe testing at 1h, klotho-treated mice spent more time at the target center (FIG. 1E), showed increased frequency of crossing the target (FIG. 1F), and increased time in the target quadrant (FIG. 1G), compared to vehicle-treated controls. In probe testing at 24h, klotho-treated mice continued to show more center duration (FIG. 1H), crossing frequency (FIG. 1I), and percent time in the target quadrant (FIG. 1J), compared to vehicle-treated mice. Thus, systemic klotho delivery enhanced spatial memory, a process that engages hippocampal brain regions.

[0105] The data show that systemic delivery of klotho enhances normal cognition in mice. Klotho treatment improved learning and memory in multiple tests and measures including working memory, spatial learning, and spatial memory. These findings provide a direct therapeutic application for boosting cognitive functions, a therapy that is relevant, but not limited to, cognitive enhancement of the normal brain and cognitive dysfunction due to normal aging, numerous neurodegenerative, neurologic, and psychiatric diseases, childhood developmental syndromes, traumatic brain injury, and stress.

Example 2

[0106] Methods.

[0107] Mice. All mice were on a C57BL/6J background and were kept on a 12-h light/dark cycle with ad libitum access to food (Picolab Rodent Diet 20, Labdiet) and water. The standard housing group was five mice per cage except for single housing during water maze studies. All motor, cognitive and behavioral studies were carried out during the light cycle.

5 All studies were conducted in a blinded manner.

[0108] Aged mice (18 months) were obtained from the NIH aging colony and were used in experiments. Transgenic mice that model human neurodegenerative diseases related to α -synuclein toxicity were utilized in experiments; these mice express full length human α -synuclein from the mouse Thy-1 promoter (Rockenstein E, *et al.* (2002) *J Neurosci Res* 10 68:568–578) and exhibit motor, cognitive and behavioral deficits. Increased expression of the human α -synuclein protein contributes to several neurodegenerative diseases in the human condition including, but not limited to, Parkinson's disease (PD), Alzheimer's disease (AD), Lewy body dementia (LBD), and multiple system atrophy (MSA).

[0109] Treatments. Vehicle or α -klotho (recombinant mouse α -klotho, amino acids 35-982, 15 with a C-terminal His tag, R&D Systems) was injected intraperitoneally (i.p.) before behavioral testing of mice as indicated. All animal studies were approved by the Institutional Animal Care and Use Committee of the University of California, San Francisco and conducted in compliance with NIH guidelines.

[0110] Cognitive and Motor Behavior

20 **[0090]** Morris water maze. Water maze studies were carried out as described (Zarei M, *et al.* (2013) *J Neurol Neurosurg Psychiatry* 84:875-881; Dubal DB *et al.* (2014) *Cell Reports* 7:1065-1076). Mice were treated with either vehicle or klotho (10 μ g/kg) i.p. 4h prior to testing daily for 5 days. Briefly, mice were tested in a pool (diameter, 122 cm) with white, opaque water (21 $^{\circ}$ $\pm$ 1 $^{\circ}$ C). A square, 14-cm² platform was 2 cm below the surface. Before 25 hidden platform training, mice underwent two pre-training trials by swimming through a channel to mount a hidden platform. Over the course of hidden platform training, the platform location stayed consistent while the drop location was varied between trials. Mice underwent two training sessions, consisting of two trials of 60 s each, daily for four days. For the probe trial testing, the platform was removed and the mice were allowed to swim for 60 s. 30 After 1h and 24h probe trials, mice were tested for their ability to find a visible platform marked with a cue (15-cm pole on the platform) over two sessions. As part of the studies, swim velocities were also recorded.

[0090] Y-maze. Y-maze studies were carried out as described (Dubal DB et al. (2014) *Cell Reports* 7:1065-1076). Mice were treated with either vehicle or klotho (10µg/kg) i.p. 18h prior to testing. Briefly, mice were acclimated to the room 30 min prior to testing. Then, mice were placed in one arm of the Y-maze (three identical arms, 120° apart) and explored for 4 min. Arm entries were recorded and an alternation was noted any time the mouse entered each of the three arms in successive arm entries; chance alternation was 22%. The apparatus was cleaned with 70% alcohol between testing sessions. Percent alternations was calculated from recorded data.

[0111] Large Y-maze. Large Y-maze studies were performed as described (Dellu F, *et al.* (1992) *Brain Res* 588:132-139) with minor modifications. The large Y-maze apparatus consists of three identical arms, 120° apart, with a distinct and different visual cue at the back end of two arms; the third arm without a visual cue is the start arm. Prior to training or testing, mice were acclimated to the dimly lit room for 60 minutes. During training, one arm with a visual cue was blocked with a solid divider and mice were placed at the end of the start arm. Mice were allowed to explore the open arm for five minutes and then returned to their cages. The open arm used in training (the familiar arm) was counterbalanced throughout the cohort testing. 16 hours after training, mice underwent testing. During testing, the divider was removed. Mice were placed in the start arm and allowed to explore the arms with novel and familiar visual cues (novel arm and familiar arm) for 5 minutes. Number of entries and duration of time spent in the novel and familiar arms was recorded and measured. The ratio of novel to familiar arm entries or duration was calculated to assess spatial and working memory. The apparatus was cleaned with 70% alcohol between sessions.

[0112] Rota Rod. Mice were acclimated to the room 60 minutes prior to each session. Five mice were simultaneously put on the rotating rod (Rota Rod, Med Associates Inc, VT) at a constant speed of 16 rpm for a maximum of 300 seconds. Latency to fall was recorded in each trial. Three trials were performed consecutively with a 10 minute rest between trials. The apparatus was cleaned with 70% alcohol between testing sessions.

[0113] Open field. Total activity in the open field was measured as described (Dubal DB, *et al.* (2015) *J Neurosci* 35:2358-2371) with an automated Flex-Field/Open Field Photobeam Activity System (San Diego Instruments, San Diego, CA). Mice were acclimated to the testing room for 30 minute and then tested in a clear plastic chamber (41 x 30 cm) for 5 min,

with two photobeam arrays measuring movements. The apparatus was cleaned with 70% alcohol between testing sessions.

[0114] Results.

[0115] Systemic klotho delivery enhances cognition in aged mice. To determine whether therapeutic delivery of klotho can enhance cognition in aged mice, we injected mice with recombinant klotho (i.p., 10µg/kg) once and then trained them for a task to measure working and spatial memory in the large Y-maze 24h later. Then, 18h following training (42h after klotho delivery), mice underwent testing. At baseline, and without klotho treatment, the aged mice did not show novel arm preference, indicating a cognitive deficit. Compared to vehicle-treated mice, klotho-treated mice showed a persistent preference for the novel arm largely throughout exploration (FIG. 2A,B), indicating better cognition that involves frontal and hippocampal brain regions. These data show that klotho enhances cognition in mice with age-induced cognitive deficits. These data suggest that therapeutic delivery of klotho enhances cognition in the aging brain.

[0116] Systemic klotho delivery enhances cognition in a long-lasting manner. To determine whether therapeutic delivery of klotho can induce long-lasting cognitive enhancement, in a manner that extends beyond its half-life of 7 hours (Hu MC, *et al.* (2015) *J Am Soc Nephrol.*), we tested young mice approximately two weeks after 5 days of daily klotho treatment and cognitive testing (i.p., 0.5 or 2.5 µg/kg). As expected, vehicle-treated mice did not show preference for the novel arm, indicating a loss of memory for the visual cue after two weeks. In contrast, klotho-treated mice showed a clear preference for the novel arm of the maze – even two weeks after the last treatment and in a dose-dependent manner (FIG. 3). These data indicate that klotho-mediated cognitive enhancement in normal, young mice was long-lasting and extended at least two weeks beyond its half-life. These data suggest that therapeutic delivery of klotho induces organizational changes in the brain that enhance neural function.

[0117] Systemic klotho delivery improves cognitive dysfunction in mice that model neurodegenerative disease. We next tested whether acute, therapeutic delivery of klotho can improve existing cognitive deficits in a mouse model of neurodegenerative disease. In this well-characterized model of disease, mice express human α-synuclein (hSYN), a pathogenic protein that causes cognitive and motor deficits and contributes to Parkinson's disease, Alzheimer's disease, Lewy body dementia, and multiple system atrophy. In these mice,

cognitive function was assessed in the large-Y maze. This task is optimal since it utilizes the mouse's natural tendency to explore, avoids testing-induced stressors, and is not affected by motor difficulties that limit other tests (such as inability to swim appropriately in a water maze). Compared to nontransgenic (NTG) mice, hSYN mice showed decreased preference for the novel arm, as shown by decreased duration (FIG. 4A). Remarkably, klotho treatment of hSYN mice (i.p., 2.5 µg/kg,) increased novel arm preference (FIG. 4A,B), indicating acute and therapeutic cognitive improvement. There was an overall effect of klotho treatment in enhancing memory as indicated by increasing novel arm preference (Two-way ANOVA KL effect $p < 0.05$) (FIG. 4B). These data suggest that therapeutic delivery of klotho can reverse or improve cognitive deficits caused by α -synuclein toxicity in neurodegenerative diseases. Of note, klotho-mediated cognitive enhancement could contribute to improved motor learning.

[0118] Systemic klotho delivery improves early motor dysfunction in mice that model neurodegenerative disease. Since α -synuclein toxicity induces deficits in motor function, a major clinical problem in neurodegenerative diseases, we tested whether therapeutic delivery of klotho can improve this key measure in hSYN mice. Motor function was assessed on a spinning rod, the Rota rod. Compared to NTG mice, hSYN mice showed decreased motor function (FIG. 5). Remarkably, klotho treatment (i.p., 2.5 µg/kg) of hSYN mice acutely increased motor function, as shown by longer latency on the spinning rod (FIG. 5). These data indicate that klotho treatment acutely improves early motor function in hSYN mice. These data suggest that therapeutic delivery of klotho can improve deficits related to motor problems caused by α -synuclein toxicity in neurodegenerative diseases.

[0119] Systemic klotho delivery does not alter total movements or activity of mice. To examine whether klotho delivery alters total movements and activity of mice, we examined mouse exploration of an open field. Total movements did not differ between Veh- or KL-treated (i.p., 2.5 µg/kg) mice (FIG. 6).. These data suggest that the therapeutic effects of klotho are specific to cognitive and motor functions and are not influenced by non-specific actions such as hyperactivity.

[0120] Discussion

[0121] Our data show that systemic delivery of klotho, which does not cross the blood brain barrier, enhances normal cognition in young mice in a manner that is long-lasting, improves cognitive deficits in normal young mice and in aging mice; it also improves

cognitive and motor deficits in transgenic mice that model major neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, Lewy body dementia, and multiple system atrophy.

[0122] These findings provide a direct therapeutic application for boosting cognitive functions in the normal brain and improving brain function in aging and neurodegenerative disease. This therapeutic application is also relevant, but not limited to, cognitive dysfunction due to numerous neurologic, and psychiatric diseases, childhood developmental syndromes, traumatic brain injury, and stress.

[0123] Our data show that:

1. Systemic administration of mouse recombinant klotho forms enhance normal cognition across the lifespan from young (2–7 mos) to aged (18mos) male and female mice.
2. Systemic klotho therapy is effective in enhancing cognition when given from 4 hours to 16 days prior to testing. The cognitive-enhancing effects last for at least two weeks following extensive cognitive training.
3. Systemic doses of mouse klotho between 0.5 µg/kg and 10 µg/kg enhance cognition in conditions tested.
4. Systemic therapy with recombinant klotho enhances cognition in a mouse model of neurodegenerative disease relevant to, but not limited to, Alzheimer's disease, Parkinson's disease, Lewy body dementia, and multiple system atrophy.
5. In addition to improving cognitive deficits in a mouse model of neurodegenerative disease, klotho therapy also improved early motor function.
6. The therapeutic effects of klotho on enhancing cognition extend far beyond its half-life of approximately 7 hours.
7. The therapeutic effects of klotho appear specific to cognitive and motor functions and are not influenced by non-specific actions such as hyperactivity.

[0124] The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, databases, internet sources, patents, patent applications, and accession numbers cited herein are hereby incorporated by reference in their entireties for all purposes.

VII. Informal sequence listing

SEQ ID NO:1: Human Klotho Protein

	10	20	30	40	50
5	MPASAPRRRP	RPPPPSLSL	LVLLGLGGR	LRAEPGDGAQ	TWARFSRPPA
	60	70	80	90	100
	PEAAGLFQGT	FPDGFLWAVG	SAAYQTEGGW	QQHGKGASIW	DTFTHHPLAP
	110	120	130	140	150
	PGDSRNASLP	LGAPSPLQPA	TGDVASDSYN	NVFRDTEALR	ELGVTHYRFS
10	160	170	180	190	200
	ISWARVLPNG	SAGVPNREGL	RYYRLLERL	RELGVQPVVT	LYHWDLPQRL
	210	220	230	240	250
	QDAYGGWANR	ALADHFRDYA	ELCFRHFGGQ	VKYWITIDNP	YVVAWHGYAT
	260	270	280	290	300
15	GRLAPGIRGS	PRLGYLVAHN	LLLAHAKVWH	LYNTSFRPTQ	GGQVSIALSS
	310	320	330	340	350
	HWINPRRMTD	HSIKECQKSL	DFVLGWFAKP	VFIDGDYPES	MKNNLSSILP
	360	370	380	390	400
	DFTESEKKFI	KGTADFFALC	FGPTLSFQLL	DPHMKFRQLE	SPNLRQLLSW
20	410	420	430	440	450
	IDLEFNHPQI	FIVENGWFVS	GTTKRDDAKY	MYYLKKFIME	TLKAIKLDGV
	460	470	480	490	500
	DVIGYTAWSL	MDGFEWHRGY	SIRRGIFYVD	FLSQDKMLLP	KSSALFYQKL
	510	520	530	540	550
25	IEKNGFPPLP	ENQPLEGTFP	CDFAWGVVDN	YIQVDTTLSQ	FTDLNVYLWD
	560	570	580	590	600
	VHHSKRLIKV	DGVVTKKRKS	YCVDFAAIQP	QIALLOEMHV	THFRFSLDWA
	610	620	630	640	650
	LILPLGNQSQ	VNHTILQYYR	CMASELVRVN	ITPVVALWQP	MAPNQGLPRL
30	660	670	680	690	700
	LARQGAWENP	YTALAFAEYA	RLCFQELGHH	VKLWITMNEP	YTRNMTYSAG
	710	720	730	740	750
	HNLLKAHALA	WHVYNEKFRH	AQNGKISIAL	QADWIEPACP	FSQKDKEVAE
	760	770	780	790	800
35	RVLEFDIGWL	AEPIFGSGDY	PWVMRDWLNQ	RNNFLLPYFT	EDEKKLIQGT
	810	820	830	840	850
	FDFLALSHYT	TILVDSEKED	PIKYNDYLEV	QEMTDITWLN	SPSQVAVVPW
	860	870	880	890	900
	GLRKVLNWLK	FKYGDLPYI	ISNGIDDGLH	AEDDQLRVYY	MQNYINEALK

WO 2016/127097

PCT/US2016/016842

910920930940950

AHILDGINLC GYFAYSFNDR TAPRFGLYRY AADQFEPKAS MKHYRKIIDS

9609709809901000

NGFPGPETLE RFCPEEFTVC TECSFFHTRK SLLAFIAFLF FASIISLSLI

51010

FYYSKKGRRS YK

1 WHAT IS CLAIMED IS:

2 1. A method for improving cognitive function in an individual in need
3 thereof comprising
4 administering to the individual an effective amount of a protein comprising a
5 Klotho polypeptide or a functional fragment thereof, wherein the administering is systemic or
6 peripheral,
7 thereby improving cognitive function in the individual.

8
9 2. The method of claim 1, wherein the protein is a Klotho polypeptide or
10 a functional fragment thereof.

11 3. The method of claim 1, wherein the administering is oral, mucosal, or
12 carried out by injection.

13 4. The method of claim 3, wherein the injection is intravenous,
14 intraperitoneal, subcutaneous, or intramuscular.

15 5. The method of any one of the proceeding claims, wherein the
16 individual is a human.

17 6. The method of claim 5, wherein the human has at least normal
18 cognitive function and the administering results in improved cognitive function compared to
19 before the administering.

20 7. The method of claim 5, wherein the human is 50 years of age or older.

21 8. The method of claim 7, wherein the human has age related cognitive
22 decline.

23 9. The method of claim 5, wherein the human is less than 50 years of age.

24 10. The method of any one of the proceeding claims, wherein the
25 individual is a human having a neurodegenerative disease.

26 11. The method of claim 10, wherein the neurodegenerative disease is
27 selected from the group consisting of: Alzheimer's disease, Parkinson's disease, Huntington's
28 disease, frontotemporal dementia, progressive supranuclear palsy, corticobasalar

29 degeneration, mild cognitive impairment, vascular dementia, Lewy body dementia, multiple
30 system atrophy, amyotrophic lateral sclerosis, prion disorder, and HIV-related dementia.

31 12. The method of any one of the foregoing claims, wherein the individual
32 is a human having a condition selected from the group consisting of: depression,
33 schizophrenia, attention deficit/ hyperactivity disorder, autism spectrum disorder, intellectual
34 disability, a mood disorder, and a psychotic disorder.

35 13. The method of any one of the foregoing claims, wherein the individual
36 is a human having a condition selected from the group consisting of traumatic brain injury,
37 stroke, multiple sclerosis, neuroautoimmune disease, epilepsy, delirium, and a paraneoplastic
38 disorder.

39 14. The method of any one of the foregoing claims, wherein the individual
40 is a human having a condition selected from the group consisting of: an X-linked mental
41 disorder, Down's syndrome, Angelman's syndrome, Rett's syndrome, phenylketonuria,
42 Lesch-Nyhan, galactosemia, and adrenoleukodystrophy.

43 15. The method of any one of the foregoing claims, wherein the individual
44 is a human having a condition selected from astrocytoma, ependymoma, medulloblastoma,
45 and oligodendroglioma.

46 16. The method of any one of the foregoing claims, wherein the individual
47 is a human receiving radiation treatment or chemotherapy for cancer.

48 17. The method of any one of the foregoing claims, wherein the individual
49 is a human that is experiencing, or will experience within 24 hours, sleep deprivation or jet
50 lag.

51 18. The method of any one of the foregoing claims, wherein the Klotho
52 polypeptide has at least 85% identity to amino acids 34-534 of SEQ ID NO:1.

53 19. The method of any one of the foregoing claims, wherein the Klotho
54 protein is a functional fragment comprising a KL1 domain.

55 20. The method of any one of the foregoing claims, wherein the Klotho
56 protein is a functional fragment comprising KL1 and KL2 domains.

1 21. The method of any one of the foregoing claims, wherein the Klotho
2 protein has at least 85% identity to amino acids 34-979 of SEQ ID NO:1.

3 22. The method of any one of the foregoing claims, wherein the effective
4 amount is 1 ug to 1000 ug per kg body weight of the individual.

5 23. The method of any one of the foregoing claims, wherein the Klotho
6 polypeptide or a functional fragment thereof is administered more than once as part of a
7 course of treatment.

8 24. The method of claim 23, wherein the Klotho polypeptide or a
9 functional fragment thereof is administered once every 1-7 days.

10 25. The method of any one of the foregoing claims, further comprising
11 testing the cognitive function of the individual after administering.

12 26. The method of claim 25, further comprising testing the cognitive
13 function of the individual prior to administering, and comparing the cognitive function of the
14 individual prior to and after administering.

15 27. The method of claim 25 or 26, wherein cognitive function is
16 determined by testing the individual for semantic, episodic, procedural, priming, and/or
17 working memory.

18 28. The method of claim 25 or 26, wherein cognitive function is
19 determined by testing the individual for language ability, executive function, visuospatial
20 function, or dementia.

21 29. A method for improving motor function in an individual in need
22 thereof comprising
23 administering to the individual an effective amount of a protein comprising a
24 Klotho polypeptide or a functional fragment thereof, wherein the administering is systemic or
25 peripheral,
26 thereby improving motor function in the individual compared to before the
27 administering.

FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D










FIG. 1E**FIG. 1F****FIG. 1G****FIG. 1H****FIG. 1I****FIG. 1J**

FIG. 2A




FIG. 2B





FIG. 3 Long Lasting Effect

FIG. 4A




FIG. 4B





FIG. 5


FIG. 6

FIG. 1A

