



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2548323 C 2012/12/04

(11)(21) **2 548 323**

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**


(86) Date de dépôt PCT/PCT Filing Date: 2004/12/17
(87) Date publication PCT/PCT Publication Date: 2005/07/14
(45) Date de délivrance/Issue Date: 2012/12/04
(85) Entrée phase nationale/National Entry: 2006/06/06
(86) N° demande PCT/PCT Application No.: EP 2004/014399
(87) N° publication PCT/PCT Publication No.: 2005/063750
(30) Priorités/Priorities: 2003/12/23 (DE103 60 835.4);
2004/09/24 (DE10 2004 046 530.4)

(51) Cl.Int./Int.Cl. *C07D 473/04* (2006.01),
A61K 31/5025 (2006.01), *A61K 31/522* (2006.01),
A61P 3/10 (2006.01), *A61P 5/50* (2006.01),
C07D 235/00 (2006.01), *C07D 237/00* (2006.01),
C07D 487/04 (2006.01)

(72) Inventeurs/Inventors:
HIMMELSBACH, FRANK, DE;
LANGKOPF, ELKE, DE;
ECKHARDT, MATTHIAS, DE;
HAUEL, NORBERT, DE;
THOMAS, LEO, DE;
TADAYYON, MOHAMMAD, DE

(73) Propriétaire/Owner:

(54) Titre : COMPOSES IMIDAZOL BICYCLIQUES, LEUR FABRICATION ET LEUR UTILISATION COMME MEDICAMENTS
(54) Title: BICYCLIC IMIDAZOLE DERIVATIVES, THE PREPARATION THEREOF AND THEIR USE AS PHARMACEUTICAL COMPOSITIONS



(57) Abrégé/Abstract:


The present invention relates to bicyclic imidazole compounds of general formula (see formula I), wherein R¹ to R³ and A are defined as in claims 1 to 8, the tautomers, the enantiomers, the stereoisomers, the mixtures thereof and the salts thereof, which have valuable pharmacological properties, particularly an inhibiting effect on the activity of the enzyme dipeptidylpeptidase-IV (DPP-IV).

(73) Propriétaires(suite)/Owners(continued):BOEHRINGER INGELHEIM INTERNATIONAL GMBH, DE

(74) Agent: FETHERSTONHAUGH & CO.

Abstract

5 The present invention relates to bicyclic imidazole compounds of general formula



wherein R^1 to R^3 and A are defined as in claims 1 to 8, the tautomers, the
10 enantiomers, the stereoisomers, the mixtures thereof and the salts thereof, which
have valuable pharmacological properties, particularly an inhibiting effect on the
activity of the enzyme dipeptidylpeptidase-IV (DPP-IV).


15

20

25771-1192

Bicyclic imidazole derivatives, the preparation thereof and their use as pharmaceutical compositions

5 The present invention relates to new bicyclic imidazole compounds of general formula



10 the tautomers, the enantiomers, the stereoisomers, the mixtures thereof and the salts thereof, particularly the physiologically acceptable salts thereof with inorganic or organic acids which have valuable pharmacological properties, particularly an inhibiting effect on the activity of the enzyme dipeptidylpeptidase-IV (DPP-IV), the preparation thereof, the use thereof for preventing or treating illnesses or conditions
15 connected with an increased DPP-IV activity or capable of being prevented or alleviated by reducing the DPP-IV activity, particularly type I or type II diabetes mellitus, the pharmaceutical compositions containing a compound of general formula (I) or a physiologically acceptable salt thereof and processes for the preparation thereof.

20

In the above formula I

R^1 denotes a pyridinyl, phenylpyridinyl, (pyridinylphenyl)carbonyl, quinoliny, phenylquinoliny, isoquinoliny, phenylisoquinoliny or phenanthridinyl group substituted by the groups R^{10} to R^{12} , while the nitrogen atom of the above-mentioned groups is substituted by an oxygen atom, and

R^{10} denotes a hydrogen atom, a fluorine, chlorine, bromine or iodine atom, a C_{1-4} -alkyl, hydroxy or C_{1-4} -alkyloxy group,

a nitro, amino, C₁₋₃-alkylamino, di-(C₁₋₃-alkyl)amino, pyrrolidin-1-yl, piperidin-1-yl or morpholin-4-yl group,

a C₁₋₃-alkyl-carbonylamino or N-(C₁₋₃-alkyl)-C₁₋₃-alkyl-carbonylamino group,

a C₁₋₃-alkylsulphonylamino or N-(C₁₋₃-alkyl)-C₁₋₃-alkyl-sulphonylamino group,

5 a C₁₋₃-alkyl-carbonyl group,

a cyano, aminocarbonyl, (C₁₋₃-alkylamino)carbonyl, [di-(C₁₋₃-alkyl)-amino]carbonyl, pyrrolidin-1-ylcarbonyl, piperidin-1-ylcarbonyl or morpholin-4-ylcarbonyl group,

a methyl or methoxy group substituted by 1 to 3 fluorine atoms,

10 a C₁₋₃-alkylsulphonyl, C₁₋₃-alkylsulphinyl or C₁₋₃-alkylsulphonyl group,

a C₂₋₄-alkenyl or C₂₋₄-alkynyl group,

a C₃₋₄-alkenyloxy or C₃₋₄-alkynyoxy group,

a C₃₋₆-cycloalkyl or C₃₋₆-cycloalkyloxy group,

a C₃₋₆-cycloalkyl-C₁₋₃-alkyl or C₃₋₆-cycloalkyl-C₁₋₃-alkyloxy group or

15 an aryl, aryloxy, aryl-C₁₋₃-alkyl or aryl-C₁₋₃-alkyloxy group,

R¹¹ and R¹², which may be identical or different, represent a hydrogen atom, a fluorine, chlorine or bromine atom or a methyl, difluoromethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy or cyano group,

20 or a pyridazinyl, phenylpyridazinyl, (pyridazinylphenyl)carbonyl, pyrimidinyl, phenylpyrimidinyl, (pyrimidinylphenyl)carbonyl, pyrazinyl, phenylpyrazinyl, (pyrazinylphenyl)carbonyl, quinolinyl, phenylquinolinyl, quinazolinyl, phenylquinazolinyl, phthalazinyl, phenylphthalazinyl, quinoxalinyl, phenylquinoxalinyl,

25 naphthyridinyl or phenylnaphthyridinyl group substituted by the groups R¹⁰ to R¹², while at least one nitrogen atom of the above-mentioned groups is substituted by an oxygen atom, and R¹⁰ to R¹² are as hereinbefore defined,

R² denotes a 2-methyl-2-propen-1-yl, 2-chloro-2-propen-1-yl or 3-bromo-2-propen-1-yl group,

a 1-buten-1-yl, 3-methyl-1-buten-1-yl, 3-methyl-2-buten-1-yl, 2-buten-1-yl, 2-methyl-2-buten-1-yl or 2,3-dimethyl-2-buten-1-yl group,

a 2-butyn-1-yl group,

5

a 1-cyclopenten-1-ylmethyl group or

a benzyl, 2-fluorobenzyl, 2-chlorobenzyl, 2-bromobenzyl or 2-cyanobenzyl group,

10 R^3 denotes a 3-aminopiperidin-1-yl, 3-amino-azepan-1-yl, piperazin-1-yl or [1,4]-diazepan-1-yl group

or an amino group substituted by the groups R^4 and R^5 wherein

15 R^4 denotes a methyl or ethyl group and

R^5 denotes a 2-aminoethyl group, while the ethyl moiety of the 2-aminoethyl group may be substituted by one or two methyl groups,

and A denotes a $-CO-N(R^6)-$ group, while the nitrogen atom of this group is linked to

20 the imidazo ring of general formula I, and

R^6 denotes a hydrogen atom, a C_{1-4} -alkyl, C_{3-6} -cycloalkyl or aryl group,

a $-CH=CH-$ group substituted by R^6 , where R^6 is as hereinbefore defined,

25

a $-C(R^7)=N-$ group, where the nitrogen atom of this group is linked to the imidazo ring of general formula I, and

R^7 denotes a hydrogen atom, a C_{1-4} -alkyl, C_{3-6} -cycloalkyl or aryl group,

30

or an $-N=C(R^7)$ group, while the carbon atom of this group is linked to the imidazo ring of general formula I, and R^7 is as hereinbefore defined,

while by the aryl groups mentioned in the definition of the above groups is meant a phenyl group substituted by R^{10} and R^{11} and R^{10} and R^{11} are as hereinbefore defined,

5

and the above-mentioned alkyl and alkenyl groups may be straight-chain or branched,

the tautomers, the enantiomers, the diastereomers, the mixtures thereof and the salts
10 thereof.

Preferred compounds of general formula I are those wherein

R^1 denotes a pyridinyl, phenylpyridinyl, (pyridinylphenyl)carbonyl, quinolinyl,
15 phenylquinolinyl, isoquinolinyl, phenylisoquinolinyl or phenanthridinyl group substituted by the groups R^{10} and R^{11} , while the nitrogen atom of the above-mentioned groups is substituted by an oxygen atom, and

R^{10} and R^{11} , which may be identical or different, denote a hydrogen atom, a
20 fluorine, chlorine or bromine atom or a methyl, difluoromethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy or cyano group,

or a pyrimidinyl, phenylpyrimidinyl, (pyrimidinylphenyl)carbonyl, quinazolinyl, phenylquinazolinyl, quinoxaliny, phenylquinoxaliny or naphthyridinyl group
25 substituted by the groups R^{10} and R^{11} , while at least one nitrogen atom of the above-mentioned groups is substituted by an oxygen atom, and R^{10} and R^{11} are as hereinbefore defined,

R^2 denotes a 2-butyn-1-yl group,

30

R^3 denotes a 3-aminopiperidin-1-yl, piperazin-1-yl or [1,4]-diazepan-1-yl group, or

an amino group substituted by the groups R⁴ and R⁵ wherein

R⁴ denotes a methyl or ethyl group and

5 R⁵ denotes a 2-aminoethyl group, while the ethyl moiety of the 2-aminoethyl group may be substituted by one or two methyl groups,

and A denotes a -CO-N(R⁶) group, while the nitrogen atom of this group is linked to the imidazo ring of general formula I, and

10 R⁶ denotes a methyl, ethyl, propyl, isopropyl, cyclopropyl or phenyl group,

or an -N=C(R⁷)- group, while the carbon atom of this group is linked to the imidazo ring of general formula I, and

15 R⁷ denotes a hydrogen atom or a methyl group,

the tautomers, the enantiomers, the diastereomers, the mixtures thereof and the salts thereof.

20 Particularly preferred are those compounds of general formula I wherein

R¹ denotes a phenylpyridinyl, quinolinyl, isoquinolinyl or phenanthridinyl group substituted by the groups R¹⁰ and R¹¹, while the nitrogen atom of the above-mentioned groups is substituted by an oxygen atom, and

25

R¹⁰ denotes a hydrogen atom or a methyl, methoxy or cyano group and

R¹¹ denotes a hydrogen atom or a methyl group,

or a phenylpyrimidinyl, quinazolinyl, quinoxalinyl or naphthyridinyl group substituted 30 by the groups R¹⁰ and R¹¹, while at least one nitrogen atom of the above-mentioned groups is substituted by an oxygen atom, and R¹⁰ and R¹¹ are as hereinbefore defined,

R² denotes a 2-butyn-1-yl group,

R³ denotes a 3-aminopiperidin-1-yl, piperazin-1-yl or [1,4]-diazepan-1-yl group, or

5

an amino group substituted by the groups R⁴ and R⁵ wherein

R⁴ denotes a methyl group and

10 R⁵ denotes a 2-aminoethyl group, while the ethyl moiety of the 2-aminoethyl group may be substituted by one or two methyl groups,

and A denotes a -CO-N(R⁶)- group, while the nitrogen atom of this group is linked to the imidazo ring of general formula I, and

15

R⁶ denotes a methyl, ethyl, isopropyl, cyclopropyl or phenyl group,

or a -N=C(R⁷)- group, while the carbon atom of this group is linked to the imidazo ring of general formula I, and

20

R⁷ denotes a hydrogen atom or a methyl group,

the tautomers, the enantiomers, the diastereomers, the mixtures thereof and the salts thereof.

25

A first sub-group comprises those compounds of the above formula I wherein R¹, R² and A are as hereinbefore defined and R³ denotes a 3-aminopiperidin-1-yl group, the tautomers, the enantiomers, the diastereomers, the mixtures thereof and the salts thereof.

30

A second sub-group comprises those compounds of the above formula I wherein R¹, R² and A are as hereinbefore defined and R³ denotes a piperazin-1-yl group, the

tautomers, the enantiomers, the diastereomers, the mixtures thereof and the salts thereof.

A third sub-group comprises those compounds of the above formula I wherein R¹, R² 5 and A are as hereinbefore defined and R³ denotes a [1,4]-diazepan-1-yl group, the tautomers, the enantiomers, the diastereomers, the mixtures thereof and the salts thereof.

Most particularly preferred are those compounds of general formula I wherein

10

R¹ denotes a quinolinyl, isoquinolinyl, methylisoquinolinyl or phenanthridinyl group, while the nitrogen atom of the above-mentioned groups is substituted by an oxygen atom,

15

a quinazolinyl or methylquinazolinyl group, while the nitrogen atom of the above-mentioned groups is substituted by an oxygen atom,

or a quinoxalanyl group wherein both nitrogen atoms are substituted by oxygen atoms,

20

R² denotes a 2-butyn-1-yl group,

R³ denotes a 3-aminopiperidin-1-yl or a piperazin-1-yl group

25

and A denotes a -CO-N(R⁶)- group, while the nitrogen atom of this group is linked to the imidazo ring of general formula I, and R⁶ denotes a methyl group,

or a -N=C(R⁷)-group, while the carbon atom of this group is linked to the imidazo ring of general formula I, and R⁷ denotes a hydrogen atom,

30

the tautomers, the enantiomers, the diastereomers, the mixtures thereof and the salts thereof;

particular mention should be made of the following compounds of general formula I:

(a) 1-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,
5

(b) 1-[(1-oxy-quinolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,
10

(c) 1-[(3-methyl-2-oxy-isoquinolin-1-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,
15

(d) 1-[(5-oxy-phenanthridin-6-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,
20

(e) 1-[(3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,
25

(f) 2-((R)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(1-oxy-quinolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,
30

(g) 2-((R)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,
35

(h) 2-(piperazin-1-yl)-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,
40

(i) 2-((R)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,
45


(j) 1-[(2-oxy-isoquinolin-3-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine and
50

(k) 2-((*R*)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-(2-oxy-isoquinolin-3-yl)methyl]-3,5-dihydro-imidazo[4,5-*d*]pyridazin-4-one

as well as the tautomers, enantiomers, diastereomers, mixtures thereof and the salts
5 thereof.

According to the invention the compounds of general formula I are obtained by methods known *per se*, for example by the following methods:

10 a) reacting a compound of general formula



wherein


15 R¹, R² and A are as hereinbefore defined and
z¹ denotes a leaving group such as a halogen atom, a substituted hydroxy,
mercapto, sulphinyl, sulphonyl or sulphonyloxy group such as a chlorine or bromine
atom, a methanesulphonyl or methanesulphonyloxy group,
with R³-H, the enantiomers or the salts thereof, where R³ is as hereinbefore defined.

20 The reaction is expediently carried out in a solvent such as isopropanol, butanol, tetrahydrofuran, dioxane, dimethylformamide, dimethylsulphoxide, ethyleneglycol monomethylether, ethyleneglycol diethylether, N-methyl-pyrrolidin-2-one or sulpholane, optionally in the presence of an inorganic or tertiary organic base, e.g. 25 sodium carbonate, potassium carbonate or potassium hydroxide, a tertiary organic base, e.g. triethylamine, or in the presence of N-ethyl-diisopropylamine (Hünig base), while these organic bases may simultaneously also serve as solvent, and optionally in the presence of a reaction accelerator such as an alkali metal halide or a palladium-based catalyst at temperatures between -20 and 180°C, but preferably at

temperatures between -10 and 120°C. The reaction may, however, also be carried out without solvent or in an excess of the amino compound R³-H.

b) deprotecting a compound of general formula

5



wherein R¹, R² and A are as hereinbefore defined and R^{3'} denotes one of the groups given as a definition of R³ hereinbefore, wherein the amino or imino group is

10 protected by a protecting group such as a tert.-butyloxycarbonyl, benzyloxycarbonyl, formyl or trifluoroacetyl group, while for the amino function the phthalyl group is an additional possibility.

The tert.-butyloxycarbonyl group is preferably cleaved by treating with an acid such

15 as trifluoroacetic acid or hydrochloric acid or by treating with bromotrimethylsilane or iodotrimethylsilane, optionally using a solvent such as methylene chloride, ethyl acetate, dioxane, methanol, isopropanol or diethyl ether at temperatures between 0 and 80°C.

20 However, the benzyloxycarbonyl group is cleaved, for example, hydrogenolytically, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a suitable solvent such as methanol, ethanol, ethyl acetate or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid at temperatures between 0 and 100°C, but preferably at temperatures between 20 and 60°C, and at a 25 hydrogen pressure of 1 to 7 bar, but preferably 3 to 5 bar.

The formyl and trifluoroacetyl group are cleaved, for example, hydrolytically in an aqueous solvent, e.g. in water, isopropanol/water, acetic acid/water, tetrahydrofuran/water or dioxane/water, in the presence of an acid such as

30 trifluoroacetic acid, hydrochloric acid or sulphuric acid or in the presence of an alkali

metal base such as sodium hydroxide or potassium hydroxide at temperatures between 0 and 120°C, preferably at temperatures between 10 and 100°C.

A phthalyl group is preferably cleaved in the presence of hydrazine or a primary 5 amine such as methylamine, ethylamine or n-butylamine in a solvent such as methanol, ethanol, isopropanol, toluene/water or dioxane at temperatures between 20 and 50°C.

In the reactions described hereinbefore, any reactive groups present such as amino, 10 alkylamino or imino groups may be protected during the reaction by conventional protecting groups which are cleaved again after the reaction.

For example, a protecting group for an amino, alkylamino or imino group may be a 15 formyl, acetyl, trifluoroacetyl, ethoxycarbonyl, tert-butoxycarbonyl, benzyloxycarbonyl, benzyl, methoxybenzyl or 2,4-dimethoxybenzyl group and additionally, for the amino group, a phthalyl group.

Any protecting group used is optionally subsequently cleaved for example by 20 hydrolysis in an aqueous solvent, e.g. in water, isopropanol/water, acetic acid/water, tetrahydrofuran/water or dioxane/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulphuric acid or in the presence of an alkali metal base such as sodium hydroxide or potassium hydroxide or aprotically, e.g. in the presence of iodotrimethylsilane, at temperatures between 0 and 120°C, 25 preferably at temperatures between 10 and 100°C.

25 However, a benzyl, methoxybenzyl or benzyloxycarbonyl group is cleaved, for example, hydrogenolytically, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a suitable solvent such as methanol, ethanol, ethyl acetate or glacial acetic acid optionally with the addition of an acid such as hydrochloric acid at 30 temperatures between 0 and 100°C, but preferably at temperatures between 20 and 60°C, and at a hydrogen pressure of 1 to 7 bar, but preferably from 3 to 5 bar.

However, a 2,4-dimethoxybenzyl group is preferably cleaved in trifluoroacetic acid in the presence of anisole.

A tert.-butyl or tert.-butyloxycarbonyl group is preferably cleaved by treating with an

5 acid such as trifluoroacetic acid or hydrochloric acid or by treating with iodotrimethylsilane optionally using a solvent such as methylene chloride, dioxane, methanol or diethyl ether.

A trifluoroacetyl group is preferably cleaved by treating with an acid such as

10 hydrochloric acid optionally in the presence of a solvent such as acetic acid at temperatures between 50 and 120°C or by treating with sodium hydroxide solution, optionally in the presence of a solvent such as tetrahydrofuran at temperatures between 0 and 50°C.

15 A phthalyl group is preferably cleaved in the presence of hydrazine or a primary amine such as methylamine, ethylamine or n-butylamine in a solvent such as methanol, ethanol, isopropanol, toluene/water or dioxane at temperatures between 20 and 50°C.

20 Moreover, the compounds of general formula I obtained may be resolved into their enantiomers and/or diastereomers, as mentioned hereinbefore. Thus, for example, cis/trans mixtures may be resolved into their cis and trans isomers, and compounds with at least one optically active carbon atom may be separated into their enantiomers.

25 Thus, for example, the *cis/trans* mixtures obtained may be separated by chromatography into their *cis* and *trans* isomers, the compounds of general formula I obtained which occur as racemates may be separated by methods known *per se* (cf. Allinger N. L. and Eliel E. L. in "Topics in Stereochemistry", Vol. 6, Wiley Interscience, 30 1971) into their optical enantiomers and compounds of general formula I with at least 2 asymmetric carbon atoms may be resolved into their diastereomers on the basis of their physical-chemical differences using methods known *per se*, e.g. by

chromatography and/or fractional crystallisation, and, if these compounds are obtained in racemic form, they may subsequently be resolved into the enantiomers as mentioned above.

- 5 The enantiomers are preferably separated by column separation on chiral phases or by recrystallisation from an optically active solvent or by reacting with an optically active substance which forms salts or derivatives such as e.g. esters or amides with the racemic compound, particularly acids and the activated derivatives or alcohols thereof, and separating the diastereomeric mixture of salts or derivatives thus obtained, e.g. on the basis of their differences in solubility, whilst the free antipodes 10 may be released from the pure diastereomeric salts or derivatives by the action of suitable agents. Optically active acids in common use are e.g. the D- and L-forms of tartaric acid or dibenzoyltartaric acid, di-o-tolyltartaric acid, malic acid, mandelic acid, camphorsulphonic acid, glutamic acid, aspartic acid or quinic acid. An optically active 15 alcohol may be, for example, (+) or (-)-menthol and an optically active acyl group in amides, for example, may be a (+)- or (-)-menthyloxycarbonyl.

Furthermore, the compounds of formula I may be converted into the salts thereof, particularly for pharmaceutical use into the physiologically acceptable salts with 20 inorganic or organic acids. Acids which may be used for this purpose include for example hydrochloric acid, hydrobromic acid, sulphuric acid, methanesulphonic acid, phosphoric acid, fumaric acid, succinic acid, lactic acid, citric acid, tartaric acid or maleic acid.

- 25 The compounds of general formulae II and III used as starting compounds are either known from the literature or may be prepared by methods known from the literature (see Examples I to X).

As already mentioned hereinbefore, the compounds of general formula I according to 30 the invention and the physiologically acceptable salts thereof have valuable pharmaceutical properties, particularly an inhibiting effect on the enzyme DPP-IV.

The biological properties of the new compounds were investigated as follows:

The ability of the substances and their corresponding salts to inhibit the DPP-IV activity can be demonstrated in an experiment in which an extract of the human colon 5 carcinoma cell line Caco-2 is used as the DPP IV source. The differentiation of the cells in order to induce the DPP-IV expression was carried out in accordance with the description by Reiher *et al.* in an article entitled "Increased expression of intestinal cell line Caco-2" , which appeared in Proc. Natl. Acad. Sci. Vol. 90, pp. 5757-5761 10 (1993). The cell extract was obtained from cells solubilised in a buffer (10mM Tris HCl, 0.15 M NaCl, 0.04 t.i.u. aprotinin, 0.5% Nonidet-P40, pH 8.0) by centrifugation at 35,000 g for 30 minutes at 4°C (to remove cell debris).

The DPP-IV assay was carried out as follows:

15 50 µl of substrate solution (AFC; AFC is amido-4-trifluoromethylcoumarin), final concentration 100 µM, were placed in black microtitre plates. 20 µl of assay buffer (final concentrations 50 mM Tris HCl pH 7.8, 50 mM NaCl, 1 % DMSO) was pipetted in. The reaction was started by the addition of 30 µl of solubilised Caco-2 protein 20 (final concentration 0.14 µg of protein per well). The test substances under investigation were typically added prediluted to 20 µl, while the volume of assay buffer was then reduced accordingly. The reaction was carried out at ambient temperature, the incubation period was 60 minutes. Then the fluorescence was measured in a Victor 1420 Multilabel Counter, with the excitation wavelength at 405 nm and the emission wavelength at 535 nm. Dummy values (corresponding to 0 % 25 activity) were obtained in mixtures with no Caco-2 protein (volume replaced by assay buffer), control values (corresponding to 100 % activity) were obtained in mixtures without any added substance. The potency of the test substances in question, expressed as IC₅₀ values, were calculated from dosage/activity curves consisting of 11 measured points in each case. The following results were obtained:

Compound (Example no.)	DPP IV inhibition IC ₅₀ [nM]
1	2
1(1)	1
1(2)	4
1(3)	6
1(4)	2

The compounds prepared according to the invention are well tolerated as no toxic side effects could be detected in rats after the oral administration of 10 mg/kg of the compound of Example 1, for example.

In view of their ability to inhibit DPP-IV activity, the compounds of general formula I according to the invention and the corresponding pharmaceutically acceptable salts thereof are suitable for influencing any conditions or diseases which can be affected by the inhibition of the DPP-IV activity. It is therefore to be expected that the compounds according to the invention will be suitable for the prevention or treatment of diseases or conditions such as type I and type II diabetes mellitus, pre-diabetes, reduced glucose tolerance or changes in the fasting blood sugar, diabetic complications (e.g. retinopathy, nephropathy or neuropathies), metabolic acidosis or ketosis, reactive hypoglycaemia, insulin resistance, dyslipidaemias of various origins, arthritis, atherosclerosis and related diseases, obesity, allograft transplantation and osteoporosis caused by calcitonin. In addition, these substances are suitable for preventing B-cell degeneration such as e.g. apoptosis or necrosis of pancreatic B-cells. The substances are also suitable for improving or restoring the function of pancreatic cells and additionally increasing the size and number of pancreatic B-cells. Additionally, on the basis of the role of the glucagon-like peptides such as e.g. GLP-1 and GLP-2 and their link with DPP-IV inhibition, it is expected that the compounds according to the invention will be suitable for achieving, *inter alia*, a sedative or tranquillising effect, as well as having a favourable effect on catabolic

states after operations or hormonal stress responses or possibly reducing mortality and morbidity after myocardial infarct. Moreover, they are suitable for treating any conditions connected with the effects mentioned above and mediated by GLP-1 or GLP-2. The compounds according to the invention may also be used as diuretics or 5 antihypertensives and are suitable for preventing and treating acute kidney failure. The compounds according to the invention may also be used to treat inflammatory complaints of the respiratory tract. They are also suitable for preventing and treating chronic inflammatory bowel diseases such as e.g. irritable bowel syndrome (IBS), Crohn's disease or ulcerative colitis and also pancreatitis. It is also expected that they 10 can be used for all kinds of injury or damage to the gastrointestinal tract such as may occur in colitis and enteritis, for example. Moreover, it is expected that DPP-IV inhibitors and hence the compounds according to the invention can be used to treat infertility or to improve fertility in humans or mammals, particularly if the infertility is connected with insulin resistance or with polycystic ovary syndrome. On the other 15 hand these substances are suitable for influencing sperm motility and are thus suitable for use as male contraceptives. In addition, the substances are suitable for treating growth hormone deficiencies connected with restricted growth, and may reasonably be used for all indications for which growth hormone may be used. The compounds according to the invention are also suitable, on the basis of their 20 inhibitory effect on DPP-IV, for treating various autoimmune diseases such as e.g. rheumatoid arthritis, multiple sclerosis, thyroiditis and Basedow's disease, etc. They may also be used to treat viral diseases and also, for example, in HIV infections, for stimulating blood production, in benign prostatic hyperplasia, gingivitis, as well as for the treatment of neuronal defects and neurodegenerative diseases such as 25 Alzheimer's disease, for example. The compounds described may also be used for the treatment of tumours, particularly for modifying tumour invasion and also metastasis; examples here are their use in treating T-cell lymphomas, acute lymphoblastic leukaemia, cell-based pancreatic carcinomas, basal cell carcinomas or breast cancers. Other indications are stroke, ischaemia of various origins, 30 Parkinson's disease and migraine. In addition, further indications include follicular and epidermal hyperkeratoses, increased keratinocyte proliferation, psoriasis,

encephalomyelitis, glomerulonephritis, lipodystrophies, as well as psychosomatic, depressive and neuropsychiatric diseases of all kinds.

The compounds according to the invention may also be used in conjunction with
5 other active substances. Suitable therapeutic agents for such combinations include
for example antidiabetic agents such as metformin, sulphonylureas (e.g.
glibenclamid, tolbutamide, glimepiride), nateglinide, repaglinide, thiazolidinediones
(e.g. rosiglitazone, pioglitazone), PPAR-gamma agonists (e.g. GI 262570) and
antagonists, PPAR-gamma/alpha modulators (e.g. KRP 297), PPAR-
10 gamma/alpha/delta modulators, AMPK activators, ACC1 and ACC2 inhibitors, DGAT
inhibitors, SMT3 receptor agonists, 11 β -HSD inhibitors, FGF19 agonists or mimetics,
alpha-glucosidase inhibitors (e.g. acarbose, voglibose), other DPPIV inhibitors,
alpha2 antagonists, insulin and insulin analogues, GLP-1 and GLP-1 analogues (e.g.
exendin-4) or amylin. Also, SGLT2 inhibitors such as T-1095 or KGT-1251 (869682),
15 inhibitors of protein tyrosine phosphatase 1, substances which influence deregulated
glucose production in the liver, such as e.g. inhibitors of glucose-6-phosphatase, or
fructose-1,6-bisphosphatase, glycogen phosphorylase, glucagon receptor
antagonists and inhibitors of phosphoenol pyruvate carboxykinase, glycogen
synthase kinase or pyruvate dehydrokinase, lipid lowering agents, such as HMG-
20 CoA-reductase inhibitors (e.g. simvastatin, atorvastatin), fibrates (e.g. bezafibrate,
fenofibrate), nicotinic acid and its derivatives, PPAR-alpha agonists, PPAR-delta
agonists, ACAT inhibitors (e.g. avasimibe) or cholesterol absorption inhibitors such
as for example ezetimibe, bile acid-binding substances such as for example
cholestyramine, inhibitors of ileac bile acid transport, HDL-raising compounds such
25 as for example inhibitors of CETP or regulators of ABC1 or LX α antagonists,
LXR β agonists or LX α /beta regulators or active substances for the treatment
of obesity, such as e.g. sibutramine or tetrahydrolipostatin, dextroamphetamine, axokine,
antagonists of the cannabinoid1 receptor, MCH-1 receptor antagonists, MC4 receptor
agonists, NPY5 or NPY2 antagonists or β_3 -agonists such as SB-418790 or AD-9677
30 as well as agonists of the 5HT2c receptor.

It is also possible to combine the compounds with drugs for treating high blood pressure such as e.g. All antagonists or ACE inhibitors, diuretics, β -blockers, Ca-antagonists, etc., or combinations thereof.

- 5 The dosage required to achieve such an effect is expediently, by intravenous route, 1 to 100 mg, preferably 1 to 30 mg, and by oral route 1 to 1000 mg, preferably 1 to 100 mg, in each case 1 to 4 times a day. For this purpose, the compounds of formula I prepared according to the invention, optionally combined with other active substances, may be incorporated together with one or more inert conventional carriers and/or diluents, e.g. with corn starch, lactose, glucose, microcrystalline cellulose, magnesium stearate, polyvinylpyrrolidone, citric acid, tartaric acid, water, water/ethanol, water/glycerol, water/sorbitol, water/polyethylene glycol, propylene glycol, cetylstearyl alcohol, carboxymethylcellulose or fatty substances such as hard fat or suitable mixtures thereof into conventional galenic preparations such as plain or coated tablets, capsules, powders, suspensions or suppositories.
- 10
- 15

The Examples that follow are intended to illustrate the invention:

Preparation of the starting compounds:

Example I

1-[(4-Methyl-3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(R)-3-(tert.-

5 butyloxycarbonylamino)-piperidin-1-yl]-xanthine

A mixture of 300 mg of 3-methyl-7-(2-butyn-1-yl)-8-[(R)-3-(tert.-butyloxycarbonyl-amino)-piperidin-1-yl]-xanthine, 151 mg of 2-chloromethyl-4-methyl-quinazoline-3-oxide and 220 mg of potassium carbonate in 50 ml of acetonitrile is heated for seven minutes in the microwave at 170°C. Then the acetonitrile is distilled off and the flask 10 residue is chromatographed through a silica gel column with ethyl acetate/methanol (100:0 to 90:10) as eluant.

Yield: 121 mg (29 % of theory)

R_f value: 0.60 (silica gel, ethyl acetate/methanol = 9:1)

Mass spectrum (ESI $^+$): m/z = 589 [M+H] $^+$

15

Example II

3-Methyl-7-(2-butyn-1-yl)-8-[(R)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-xanthine

11.00 g of (R)-3-tert.-butyloxycarbonylamino-piperidine are added to 15.00 g of 3-methyl-7-(2-butyn-1-yl)-8-bromo-xanthine and 16.00 g of potassium carbonate in 100 ml of dimethylsulphoxide and the thick, light beige suspension is stirred for four hours with a mechanical stirrer at approx. 114°C. Then another 900 mg of (R)-3-tert.-butyloxycarbonylamino-piperidine, dissolved in 10 ml of dimethylsulphoxide, are added to the reaction mixture and this is stirred for a further two hours at 114°C. After 25 cooling to ambient temperature the reaction mixture is diluted with copious amounts of water. The precipitate formed is thoroughly triturated until no more clumps are left, and suction filtered. The light solid is again suspended with water, suction filtered, washed with water and diethyl ether and dried in the circulating air dryer at 60°C.

Yield: 19.73 g (94 % of theory)

30 R_f value: 0.64 (silica gel, ethyl acetate)

Mass spectrum (ESI $^+$): m/z = 417 [M+H] $^+$

The following compounds are obtained analogously to Example II :

(1) 2-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-3-(2-butyn-1-yl)-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

5 (carried out in N,N-dimethylformamide at 80°C)

R_f value: 0.55 (silica gel, ethyl acetate)

Mass spectrum (ESI $^+$): m/z = 387 [M+H] $^+$

Example III

10 3-methyl-7-(2-butyn-1-yl)-8-bromo-xanthine

17.06 g of 1-bromo-2-butyne are added to 30.17 g of 3-methyl-8-bromo-xanthine and 27.00 ml of Hünig base in 370 ml of N,N-dimethylformamide. The reaction mixture is stirred for two hours at ambient temperature, then another 1 ml of 1-bromo-2-butyne is added and stirring is continued for another hour at ambient temperature. For 15 working up the reaction mixture is diluted with approx. 300 ml of water. The light precipitate formed is suction filtered and washed with water. The filter cake is washed with a little ethanol and diethyl ether and dried in the circulating air dryer at 60°C.

Yield: 30.50 g (84 % of theory)

R_f value: 0.24 (silica gel, methylene chloride/methanol = 95:5)

20 Mass spectrum (ESI $^+$): m/z = 297, 299 [M+H] $^+$

Example IV

1-[(1-oxy-quinolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-xanthine

25 Prepared by heating 450 mg of 3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-xanthine, 245 mg of 2-chloromethyl-quinoline-1-oxide and 800 mg of potassium carbonate in 5 ml of N,N-dimethylformamide to 80°C.

Yield: 622 mg (100 % of theory)

R_f value: 0.26 (silica gel, ethyl acetate)

30 Mass spectrum (ESI $^+$): m/z = 574 [M+H] $^+$

The following compounds are obtained analogously to Example IV:

(1) 1-[(3-methyl-2-oxy-isoquinolin-1-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-xanthine

*R*_f value: 0.17 (silica gel, ethyl acetate)

5 Mass spectrum (ESI⁺): m/z = 588 [M+H]⁺

(2) 1-[(5-oxy-phenanthridin-6-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-xanthine

*R*_f value: 0.47 (silica gel, ethyl acetate)

10 Mass spectrum (ESI⁺): m/z = 624 [M+H]⁺

(3) 1-[(3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-xanthine

*R*_f value: 0.29 (silica gel, ethyl acetate)

15 Mass spectrum (ESI⁺): m/z = 575 [M+H]⁺

(4) 1-[(1,4-dioxy-quinoxalin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-xanthine

*R*_f value: 0.53 (silica gel, ethyl acetate)

20 Mass spectrum (ESI⁺): m/z = 591 [M+H]⁺

(5) 2-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-3-(2-butyn-1-yl)-5-[(1-oxy-quinolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

Mass spectrum (ESI⁺): m/z = 544 [M+H]⁺

25

(6) 2-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-3-(2-butyn-1-yl)-5-[(3-methyl-2-oxy-isoquinolin-1-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

Mass spectrum (ESI⁺): m/z = 558 [M+H]⁺

30 (7) 2-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-3-(2-butyn-1-yl)-5-[(5-oxy-phenanthridin-6-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

Mass spectrum (ESI⁺): m/z = 594 [M+H]⁺

(8) 2-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-3-(2-butyn-1-yl)-5-[(3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

Mass spectrum (ESI⁺): m/z = 545 [M+H]⁺

5

(9) 2-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-3-(2-butyn-1-yl)-5-[(1,4-dioxy-quinoxalin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

Mass spectrum (ESI⁺): m/z = 561 [M+H]⁺

10 (10) 2-bromo-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

R_f value: 0.30 (silica gel, ethyl acetate)

Mass spectrum (ESI⁺): m/z = 439, 441 [M+H]⁺

15 (11) 2-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

R_f value: 0.40 (silica gel, ethyl acetate/methanol = 9:1)

Mass spectrum (ESI⁺): m/z = 559 [M+H]⁺

20 (12) 1-[(2-oxy-isoquinolin-3-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-xanthine

R_f value: 0.35 (silica gel, methylene chloride/methanol = 95:5)

Mass spectrum (ESI⁺): m/z = 574 [M+H]⁺

25 (13) 2-[(*R*)-3-(tert.-butyloxycarbonylamino)-piperidin-1-yl]-3-(2-butyn-1-yl)-5-(2-oxy-isoquinolin-3-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

R_f value: 0.10 (silica gel, ethyl acetate/methanol = 98:2)

Mass spectrum (ESI⁺): m/z = 544 [M+H]⁺

30

Example V1-Chloromethyl-3-methyl-isoquinoline-2-oxide

A solution of 300 mg of 1-chloromethyl-3-methyl-isoquinoline in 3 ml of methylene chloride is combined with 390 mg of 3-chloroperoxybenzoic acid and stirred overnight

5 at ambient temperature. The reaction mixture is diluted with some methylene chloride and extracted with saturated sodium hydrogen carbonate solution. The combined organic phases are dried over magnesium sulphate and evaporated down. The solid, yellowish crude product is triturated with tert.-butylmethylether, suction filtered, washed with tert.-butylmethylether and dried.

10 Yield: 285 mg (88 % of theory)

R_f value: 0.31 (silica gel, ethyl acetate/petroleum ether = 3:2)

Mass spectrum (ESI $^+$): m/z = 208, 210 [M+H] $^+$

The following compound is obtained analogously to Example V:

15

(1) 6-chloromethyl-phenanthridine-5-oxide

R_f value: 0.66 (silica gel, ethyl acetate/petroleum ether = 3:2)

Mass spectrum (ESI $^+$): m/z = 244, 246 [M+H] $^+$

20

Example VI2-bromomethyl-quinazoline-3-oxide

A solution of 1.00 g of 2-methyl-quinazoline-3-oxide in 30 ml of glacial acetic acid is combined dropwise with a solution of 0.48 ml of bromine in 10 ml of glacial acetic acid and stirred overnight at ambient temperature. Then the reaction mixture is

25 stirred for another two hours at 80°C. The glacial acetic acid is largely distilled off and the residue is stirred with saturated sodium hydrogen carbonate solution. The lumpy precipitate formed is taken up in ethyl acetate. The ethyl acetate is distilled off again and the fine precipitate is suction filtered, washed with ethanol and tert.-butylmethylether and dried. The crude product is purified by chromatography through a silica gel column with ethyl acetate as eluant .

30 Yield: 654 mg (44 % of theory)

R_f value: 0.52 (silica gel, ethyl acetate)

Mass spectrum (ESI⁺): m/z = 239, 241 [M+H]⁺

Example VII

2-bromo-3-(2-butyn-1-yl)-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

5 0.31 ml of hydrazine hydrate (99%), dissolved in 1 ml of ethanol, are added dropwise at ambient temperature to a solution of 1.80 g of methyl 2-bromo-3-(2-butyn-1-yl)-5-formyl-3H-imidazole-4-carboxylate in 25 ml of ethanol. Five minutes later 1.5 ml of concentrated acetic acid are added and the mixture is refluxed for 30 minutes. After cooling the precipitated solid is suction filtered, washed with 10 ml of ethanol and 20 ml of diethyl ether and dried.

10 Yield: 1.25 g of (74 % of theory)

Mass spectrum (ESI⁺): m/z = 267, 269 [M+H]⁺

1H-NMR spectrum (d6-DMSO): δ = 1.80 (s, 3H); 5.28 (s, 2H); 8.38 (s, 1H); 12.99 (s, 1H) ppm

15

Example VIII

Methyl 2-bromo-3-(2-butyn-1-yl)-5-formyl-3H-imidazole-4-carboxylate

43 ml of a 1M solution of diisobutyl-aluminium hydride in tetrahydrofuran are added dropwise to a solution of 13.5 g of dimethyl 2-bromo-1-(2-butyn-1-yl)-1H-imidazole-20 4,5-dicarboxylate in 220 ml of tetrahydrofuran under an argon atmosphere at -70°C within 20 minutes. The mixture is stirred for a further four hours at -70°C, then 20 ml of a mixture of 1M hydrochloric acid and tetrahydrofuran are added dropwise. After heating to ambient temperature approx. 200 ml of water are added and the mixture is extracted three times with 70 ml of ethyl acetate. The combined extracts are dried 25 and evaporated down. The crude product thus obtained is purified by column chromatography through silica gel with petroleum ether/ethyl acetate (80:20 to 50:50) as eluant.

Yield: 6.40 g (52% of theory)

Mass spectrum (ESI⁺): m/z = 285, 287 [M+H]⁺

30 1H-NMR spectrum (d6-DMSO): δ = 1.80 (s, 3H); 3.93 (s, 3H); 5.11 (s, 2H); 10.12 (s, 1H) ppm

Example IXDimethyl 2-bromo-1-(2-butyn-1-yl)-1H-imidazole-4,5-dicarboxylate

A solution of 15.0 g of dimethyl 2-bromo-imidazole-4,5-dicarboxylate, 5.15 ml of 1-

5 bromo-2-butyne and 50 ml of N,N-diisopropylethylamine in 280 ml of tetrahydrofuran is refluxed for one hour. The mixture is concentrated by evaporation, the residue is combined with approx. 100 ml of water and extracted three times with 70 ml of ethyl acetate. The extracts are washed with 50 ml of water, dried and evaporated down.

10 The crude product thus obtained is purified by column chromatography through silica gel using methylene chloride/ethanol (100:0 to 98:2) as eluant .

Yield: 13.50 g (75 % of theory)

R_f value: 0.82 (silica gel, methylene chloride/ethanol = 9:1)

Mass spectrum (ESI $^+$): m/z = 315, 317 [M+H] $^+$

15

Example X3-chloromethyl-isoquinoline-2-oxide

Prepared by treating 3-chloromethyl-isoquinoline with 35 % hydrogen peroxide solution in glacial acetic acid at 70°C.

20 R_f value: 0.30 (silica gel, ethyl acetate/methanol = 98:2)


Mass spectrum (ESI $^+$): m/z = 194, 196 [M+H] $^+$

Preparation of the final compounds:

Example 1

1-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-

5 piperidin-1-yl)-xanthine



10 A mixture of 121 mg of 1-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(R)-3-(tert.-butyloxycarbonylamo)-piperidin-1-yl]-xanthine and 0.59 ml of trifluoroacetic acid in 4 ml of methylene chloride is stirred for one hour at ambient temperature. For working up the reaction mixture is diluted with methylene chloride and water, made alkaline with 1 N sodium hydroxide solution and extracted with 15 methylene chloride. The organic phase is washed with saturated sodium chloride solution, dried over magnesium sulphate and evaporated down. A brownish solid remains.

Yield: 84 mg (84 % of theory)


R_f value: 0.50 (reversed phase ready-made TLC plate (E. Merck), acetonitrile/water/

20 trifluoroacetic acid = 50:50:1)

Mass spectrum (ESI $^+$): m/z = 489 [M+H] $^+$

The following compounds are obtained analogously to Example 1:

25 (1) 1-[(1-oxy-quinolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine




(carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

R_f value: 0.53 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI⁺): m/z = 474 [M+H]⁺

5

(2) 1-[(3-methyl-2-oxy-isoquinolin-1-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine




(carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

10 R_f value: 0.39 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI⁺): m/z = 488 [M+H]⁺

(3) 1-[(5-oxy-phenanthridin-6-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-

15 piperidin-1-yl)-xanthine




(carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

R_f value: 0.47 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

20 Mass spectrum (ESI⁺): m/z = 524 [M+H]⁺

(4) 1-[(3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((*R*)-3-amino-piperidin-1-yl)-xanthine




(carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

5 R_f value: 0.41 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI⁺): m/z = 475 [M+H]⁺

(5) 1-[(1,4-dioxy-quinoxalin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((*R*)-3-amino-

10 piperidin-1-yl)-xanthine




(carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

R_f value: 0.55 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

15 Mass spectrum (ESI⁺): m/z = 491 [M+H]⁺

(6) 2-((*R*)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(1-oxy-quinolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one




20 (carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

R_f value: 0.33 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI⁺): m/z = 444 [M+H]⁺


(7) 2-((*R*)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(3-methyl-2-oxy-isoquinolin-1-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one



5 R_f value: 0.27 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI $^+$): m/z = 458 [M+H] $^+$

10 (8) 2-((*R*)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(5-oxy-phenanthridin-6-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one




(carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

R_f value: 0.24 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

15 Mass spectrum (ESI $^+$): m/z = 494 [M+H] $^+$


(9) 2-((*R*)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one



20 R_f value: 0.24 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI $^+$): m/z = 445 [M+H] $^+$


(10) 2-((*R*)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(1,4-dioxy-quinoxalin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one



5 R_f value: 0.26 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI $^+$): m/z = 461 [M+H] $^+$

10 (11) 2-((*R*)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one




R_f value: 0.60 (reversed phase ready-made TLC plate (E. Merck), acetonitrile/water/trifluoroacetic acid = 50:50:1)

Mass spectrum (ESI $^+$): m/z = 459 [M+H] $^+$

15

(12) 1-[(2-oxy-isoquinolin-3-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((*R*)-3-amino-piperidin-1-yl)-xanthine




(carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

20 R_f value: 0.48 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI $^+$): m/z = 474 [M+H] $^+$

(13) 2-((R)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-(2-oxy-isoquinolin-3-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one




(carried out with isopropanolic hydrochloric acid (5-6 M) in methylene chloride)

5 R_f value: 0.38 (silica gel, methylene chloride/methanol/conc. aqueous ammonia = 90:10:1)

Mass spectrum (ESI⁺): m/z = 444 [M+H]⁺

10 Example 2

2-(Piperazin-1-yl)-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one



A mixture of 250 mg of 2-bromo-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one and 250 mg of piperazine in 5 ml of N,N-dimethylformamide is heated for five minutes in the microwave at 200°C. Then the solvent is distilled off *in vacuo* and the flask residue is dissolved in methylene chloride. The solution is washed with water and saturated sodium chloride solution and evaporated down *in vacuo*. The crude product is purified through a silica gel column with methylene chloride/methanol/conc. methanolic ammonia (99:0.9:0.1 to 80:18:2) as eluant.

Yield: 35 mg (14 % of theory)

R_f value: 0.60 (silica gel, methylene chloride/methanol/conc. methanolic ammonia = 90:9:1)

25 Mass spectrum (ESI⁺): m/z = 445 [M+H]⁺

Example 3Coated tablets containing 75 mg of active substance

5

1 tablet core contains:

	active substance	75.0 mg
	calcium phosphate	93.0 mg
	corn starch	35.5 mg
10	polyvinylpyrrolidone	10.0 mg
	hydroxypropylmethylcellulose	15.0 mg
	magnesium stearate	<u>1.5 mg</u>
		230.0 mg

15 Preparation:

The active substance is mixed with calcium phosphate, corn starch, polyvinylpyrrolidone, hydroxypropylmethylcellulose and half the specified amount of magnesium stearate. Blanks about 13 mm in diameter are produced in a tablet-making machine and these are then rubbed through a screen with a mesh size of 1.5 mm using a suitable machine and mixed with the rest of the magnesium stearate. This granulate is compressed in a tablet-making machine to form tablets of the desired shape.

weight of core: 230 mg

die: 9 mm, convex

25 The tablet cores thus produced are coated with a film consisting essentially of hydroxypropylmethylcellulose. The finished film-coated tablets are polished with beeswax.

Weight of coated tablet: 245 mg.

Example 4Tablets containing 100 mg of active substance

5

Composition:

1 tablet contains:

active substance	100.0 mg
lactose	80.0 mg
10 corn starch	34.0 mg
polyvinylpyrrolidone	4.0 mg
magnesium stearate	<u>2.0 mg</u>
	220.0 mg

15 Method of Preparation:

The active substance, lactose and starch are mixed together and uniformly moistened with an aqueous solution of the polyvinylpyrrolidone. After the moist composition has been screened (2.0 mm mesh size) and dried in a rack-type drier at 50°C it is

20 screened again (1.5 mm mesh size) and the lubricant is added. The finished mixture is compressed to form tablets.

Weight of tablet: 220 mg

Diameter: 10 mm, biplanar, faceted on both sides and notched on one side.

25

Example 5Tablets containing 150 mg of active substance

5 Composition:

1 tablet contains:

active substance	150.0 mg
powdered lactose	89.0 mg
corn starch	40.0 mg
10 colloidal silica	10.0 mg
polyvinylpyrrolidone	10.0 mg
magnesium stearate	<u>1.0 mg</u>
	300.0 mg

15 Preparation:

The active substance mixed with lactose, corn starch and silica is moistened with a 20% aqueous polyvinylpyrrolidone solution and passed through a screen with a mesh size of 1.5 mm. The granules, dried at 45°C, are passed through the same screen

20 again and mixed with the specified amount of magnesium stearate. Tablets are pressed from the mixture.

Weight of tablet: 300 mg

die: 10 mm, flat

Example 6Hard gelatine capsules containing 150 mg of active substance

5 1 capsule contains:

active substance	150.0 mg
corn starch (dried)	approx. 180.0 mg
lactose (powdered)	approx. 87.0 mg
magnesium stearate	<u>3.0 mg</u>

10 approx. 420.0 mg

Preparation:

The active substance is mixed with the excipients, passed through a screen with a

15 mesh size of 0.75 mm and homogeneously mixed using a suitable apparatus. The finished mixture is packed into size 1 hard gelatine capsules.

Capsule filling: approx. 320 mg

Capsule shell: size 1 hard gelatine capsule.

20 Example 7Suppositories containing 150 mg of active substance

1 suppository contains:

25 active substance	150.0 mg
polyethyleneglycol 1500	550.0 mg
polyethyleneglycol 6000	460.0 mg
polyoxyethylene sorbitan monostearate	<u>840.0 mg</u>
	2,000.0 mg

Preparation:

After the suppository mass has been melted the active substance is homogeneously distributed therein and the melt is poured into chilled moulds.

5

Example 8Suspension containing 50 mg of active substance

10 100 ml of suspension contain:

active substance	1.00 g
carboxymethylcellulose-Na-salt	0.10 g
methyl p-hydroxybenzoate	0.05 g
propyl p-hydroxybenzoate	0.01 g
15 glucose	10.00 g
glycerol	5.00 g
70% sorbitol solution	20.00 g
flavouring	0.30 g
dist. water	ad 100 ml

20

Preparation:

The distilled water is heated to 70°C. The methyl and propyl p-hydroxybenzoates together with the glycerol and sodium salt of carboxymethylcellulose are dissolved

25 therein with stirring. The solution is cooled to ambient temperature and the active substance is added and homogeneously dispersed therein with stirring. After the sugar, the sorbitol solution and the flavouring have been added and dissolved, the suspension is evacuated with stirring to eliminate air.

5 ml of suspension contain 50 mg of active substance.

30

Example 9Ampoules containing 10 mg active substance

5 Composition:

active substance	10.0 mg
0.01 N hydrochloric acid q.s.	
double-distilled water	ad 2.0 ml

10 Preparation:

The active substance is dissolved in the necessary amount of 0.01 N HCl, made isotonic with common salt, filtered sterile and transferred into 2 ml ampoules.

15 Example 10Ampoules containing 50 mg of active substance

Composition:

20	active substance	50.0 mg
	0.01 N hydrochloric acid q.s.	
	double-distilled water	ad 10.0 ml

25 Preparation:


The active substance is dissolved in the necessary amount of 0.01 N HCl, made isotonic with common salt, filtered sterile and transferred into 10 ml ampoules.

25771-1192

38

CLAIMS:

1. A compound of general formula I



wherein

5 R^1 denotes a pyridinyl, phenylpyridinyl, (pyridinylphenyl)carbonyl, quinolinyl, phenylquinolinyl, isoquinolinyl, phenylisoquinolinyl or phenanthridinyl group substituted by the groups R^{10} to R^{12} , while the nitrogen atom of the above-mentioned heteroaryl groups is substituted by an oxygen atom, and

10 R^{10} denotes hydrogen, fluorine, chlorine, bromine, iodine, C_{1-4} -alkyl, hydroxy, C_{1-4} -alkyloxy, nitro, amino, C_{1-3} -alkylamino, di-(C_{1-3} -alkyl)amino, pyrrolidin-1-yl, piperidin-1-yl, morpholin-4-yl, C_{1-3} -alkyl-carbonylamino, N -(C_{1-3} -alkyl)- C_{1-3} -alkyl-carbonylamino, C_{1-3} -alkylsulphonylamino, N -(C_{1-3} -alkyl)- C_{1-3} -alkyl-sulphonylamino, C_{1-3} -alkyl-carbonyl, cyano, aminocarbonyl, (C_{1-3} -alkylamino)carbonyl, [di-(C_{1-3} -alkyl)-amino]carbonyl, pyrrolidin-1-ylcarbonyl, piperidin-1-ylcarbonyl, 15 morpholin-4-ylcarbonyl, methyl substituted by 1 to 3 fluorine atoms, methoxy substituted by 1 to 3 fluorine atoms, C_{1-3} -alkylsulphanyl, C_{1-3} -alkylsulphanyl, C_{1-3} -alkylsulphonyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, C_{3-4} -alkenyloxy, C_{3-4} -alkynyoxy, C_{3-6} -cycloalkyl, C_{3-6} -cycloalkyloxy, C_{3-6} -cycloalkyl- C_{1-3} -alkyl, C_{3-6} -cycloalkyl- C_{1-3} -alkyloxy, aryl, aryloxy, aryl- C_{1-3} -alkyl or aryl- C_{1-3} -alkyloxy,

20 R^{11} and R^{12} , which may be identical or different, represent hydrogen, fluorine, chlorine, bromine, methyl, difluoromethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy or cyano,

25771-1192

39

or R^1 denotes a pyridazinyl, phenylpyridazinyl, (pyridazinylphenyl)carbonyl, pyrimidinyl, phenylpyrimidinyl, (pyrimidinylphenyl)carbonyl, pyrazinyl, phenylpyrazinyl, (pyrazinylphenyl)carbonyl, quinolinyl, phenylquinolinyl, quinazolinyl, phenylquinazolinyl, phthalazinyl, phenylphthalazinyl, quinoxaliny, phenylquinoxaliny,

5 naphthyridinyl or phenylnaphthyridinyl group substituted by the groups R^{10} to R^{12} , while at least one nitrogen atom of the above-mentioned heteroaryl groups is substituted by an oxygen atom, and R^{10} to R^{12} are as hereinbefore defined,

R^2 denotes 2-methyl-2-propen-1-yl, 2-chloro-2-propen-1-yl, 3-bromo-2-propen-1-yl, 1-buten-1-yl, 3-methyl-1-buten-1-yl, 3-methyl-2-buten-1-yl, 2-buten-1-yl,

10 2-methyl-2-buten-1-yl, 2,3-dimethyl-2-buten-1-yl, 2-butyn-1-yl, 1-cyclopenten-1-ylmethyl, benzyl, 2-fluorobenzyl, 2-chlorobenzyl, 2-bromobenzyl- or 2-cyanobenzyl,

R^3 denotes 3-aminopiperidin-1-yl, 3-amino-azepan-1-yl, piperazin-1-yl, [1,4]-diazepan-1-yl, or an amino group substituted by the groups R^4 and R^5 wherein

15 R^4 denotes a methyl or ethyl group and

R^5 denotes a 2-aminoethyl group, while the ethyl moiety of the 2-aminoethyl group may be substituted by one or two methyl groups,

and A denotes a $-CO-N(R^6)-$ group, while the nitrogen atom of this group is linked to the imidazo ring of general formula I, and

20 R^6 denotes a hydrogen atom, a C_{1-4} -alkyl, C_{3-6} -cycloalkyl or aryl group,

a $-CH=CH-$ group substituted by R^6 , where R^6 is as hereinbefore defined,

a $-C(R^7)=N-$ group, where the nitrogen atom of this group is linked to the imidazo ring of general formula I, and

R^7 denotes a hydrogen atom, a C_{1-4} -alkyl, C_{3-6} -cycloalkyl or aryl group,

25771-1192

40

or an $-N=C(R^7)$ group, while the carbon atom of this group is linked to the imidazo ring of general formula I, and R^7 is as hereinbefore defined,

while by the aryl groups mentioned in the definition of the above groups is meant a phenyl group substituted by R^{10} and R^{11} and R^{10} and R^{11} are as hereinbefore defined,

5 and the above-mentioned alkyl and alkenyl groups may be straight-chain or branched,

or a tautomer, enantiomer, diastereomer, or mixture thereof or a salt thereof.

2. A compound of general formula I according to claim 1, wherein

R^1 denotes a pyridinyl, phenylpyridinyl, (pyridinylphenyl)carbonyl, quinolinyl, 10 phenylquinolinyl, isoquinolinyl, phenylisoquinolinyl or phenanthridinyl group substituted by the groups R^{10} and R^{11} , while the nitrogen atom of the above-mentioned heteroaryl groups is substituted by an oxygen atom, and

R^{10} and R^{11} , which may be identical or different, denote hydrogen, fluorine, chlorine, bromine, methyl, difluoromethyl, trifluoromethyl, methoxy, 15 difluoromethoxy, trifluoromethoxy or cyano,

or R^1 denotes a pyrimidinyl, phenylpyrimidinyl, (pyrimidinylphenyl)carbonyl, quinazolinyl, phenylquinazolinyl, quinoxalinyl, phenylquinoxalinyl or naphthyridinyl group substituted by the groups R^{10} and R^{11} , while at least one nitrogen atom of the above-mentioned heteroaryl groups is substituted by an oxygen atom, and

20 R^{10} and R^{11} are as hereinbefore defined,

R^2 denotes a 2-butyn-1-yl group,

R^3 denotes 3-aminopiperidin-1-yl, piperazin-1-yl, [1,4]-diazepan-1-yl or an amino group substituted by the groups R^4 and R^5 wherein

25771-1192

41

R^4 denotes a methyl or ethyl group and

R^5 denotes a 2-aminoethyl group, while the ethyl moiety of the 2-aminoethyl group may be substituted by one or two methyl groups, and A denotes a $-CO-N(R^6)$ group, while the nitrogen atom of this group is linked to

5 the imidazo ring of general formula I, and

R^6 denotes a methyl, ethyl, propyl, isopropyl, cyclopropyl or phenyl group,

or an $-N=C(R^7)$ - group, while the carbon atom of this group is linked to the imidazo ring of general formula I, and

10 R^7 denotes a hydrogen atom or a methyl group,

or a tautomer, enantiomer, diastereomer, or mixture thereof or a salt thereof.

3. A compound of general formula I according to claim 1, wherein

R^1 denotes a phenylpyridinyl, quinolinyl, isoquinolinyl or phenanthridinyl group substituted by the groups R^{10} and R^{11} , while the nitrogen atom of the above-
15 mentioned heteroaryl groups is substituted by an oxygen atom, and

R^{10} denotes hydrogen, methyl, methoxy or cyano and

R^{11} denotes hydrogen or methyl,

or R^1 denotes a phenylpyrimidinyl, quinazolinyl, quinoxalinyl or naphthyridinyl group substituted by the groups R^{10} and R^{11} , while at least one nitrogen atom of the above-
20 mentioned heteroaryl groups is substituted by an oxygen atom, and R^{10} and R^{11} are as hereinbefore defined,

R^2 denotes a 2-butyn-1-yl group,

25771-1192

42

R^3 denotes 3-aminopiperidin-1-yl, piperazin-1-yl, [1,4]-diazepan-1-yl, or

an amino group substituted by the groups R^4 and R^5 wherein

R^4 denotes a methyl group and

R^5 denotes a 2-aminoethyl group, while the ethyl moiety of the

5 2-aminoethyl group may be substituted by one or two methyl groups,

and A denotes a $-CO-N(R^6)-$ group, while the nitrogen atom of this group is linked to the imidazo ring of general formula I, and

R^6 denotes a methyl, ethyl, isopropyl, cyclopropyl or phenyl group,

or a $-N=C(R^7)-$ group, while the carbon atom of this group is linked to the imidazo ring

10 of general formula I, and

R^7 denotes a hydrogen atom or a methyl group,

or a tautomer, enantiomer, diastereomer, or mixture thereof or a salt thereof.

4. A compound according to any one of claims 1 to 3, wherein

R^1 , R^2 and A are defined as in claims 1 to 3 and R^3 denotes a 3-aminopiperidin-1-yl

15 group, or a tautomer, enantiomer, diastereomer, or mixture thereof or a salt thereof.

5. A compound according to any one of claims 1 to 3, wherein

R^1 , R^2 and A are defined as in any one of claims 1 to 3 and R^3 denotes a

piperazin-1-yl group, or a tautomer, enantiomer, diastereomer, or mixture thereof or a salt thereof.

20 6. A compound according to any one of claims 1 to 3, wherein

R^1 , R^2 and A are defined as in any one of claims 1 to 3 and R^3 denotes a

[1,4]-diazepan-1-yl group, or a tautomer, enantiomer, diastereomer, or mixture thereof or a salt thereof.

25771-1192

43

7. A compound of general formula I according to claim 1, wherein

R¹ denotes a quinolinyl, isoquinolinyl, methylisoquinolinyl or phenanthridinyl group, while the nitrogen atom of the above-mentioned heteroaryl groups is substituted by an oxygen atom, or

5 R¹ denotes a quinazolinyl or methylquinazolinyl group, while the nitrogen atom of the above-mentioned heteroaryl groups is substituted by an oxygen atom,

or R¹ denotes a quinoxaliny group wherein both nitrogen atoms are substituted by oxygen atoms,

R² denotes a 2-butyn-1-yl group,

10 R³ denotes a 3-aminopiperidin-1-yl or a piperazin-1-yl group

and A denotes a -CO-N(R⁶)- group, while the nitrogen atom of this group is linked to the imidazo ring of general formula I, and R⁶ denotes a methyl group,

or a -N=C(R⁷)- group, while the carbon atom of this group is linked to the imidazo ring of general formula I, and R⁷ denotes a hydrogen atom,

15 or a tautomer, enantiomer, diastereomer, or a mixture thereof or a salt thereof.

8. The following compounds of general formula I according to claim 1:

(a) 1-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,

20 (b) 1-[(1-oxy-quinolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,

(c) 1-[(3-methyl-2-oxy-isoquinolin-1-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,

25771-1192

44

(d) 1-[(5-oxy-phenanthridin-6-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,

(e) 1-[(3-oxy-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine,

5 (f) 2-((R)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(1-oxy-quinolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,

(g) 2-((R)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,

(h) 2-(piperazin-1-yl)-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,

10 (i) 2-((R)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-[(4-methyl-3-oxy-quinazolin-2-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one,

(j) 1-[(2-oxy-isoquinolin-3-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((R)-3-amino-piperidin-1-yl)-xanthine and

15 (k) 2-((R)-3-amino-piperidin-1-yl)-3-(2-butyn-1-yl)-5-(2-oxy-isoquinolin-3-yl)methyl]-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one

or a tautomer, enantiomer, diastereomer, or a mixture thereof or a salt thereof.

9. A physiologically acceptable salt of a compound according to any one of claims 1 to 8 with an inorganic acid or organic acid.

20 10. A pharmaceutical composition containing a compound according to any one of claims 1 to 8 or a physiologically acceptable salt thereof, together with one or more inert carriers and/or diluents.

25771-1192

45


11. The pharmaceutical composition of claim 10, wherein the physiologically acceptable salt of the compound is with an inorganic acid or an organic acid.

12. Use of a compound according to any one of claims 1 to 9 or a tautomer, enantiomer, diastereomer, or a mixture thereof or a salt thereof, for preparing a pharmaceutical composition which is suitable for treating type I diabetes, type II diabetes, mellitus, arthritis, obesity, allograft transplantation or calcitonin-induced osteoporosis.

13. Process for preparing a pharmaceutical composition according to claim 10 or 11, wherein a compound according to any one of claims 1 to 9 is incorporated in one or more inert carriers and/or diluents by a non-chemical method.

14. Process for preparing a compound of general formula I according to any one of claims 1 to 9, wherein

a) a compound of general formula



15

wherein


R^1 , R^2 and A are defined as mentioned in claims 1 to 8 and

Z^1 denotes a leaving group,

is reacted with R^3 -H, an enantiomer or the salt thereof, where R^3 is as hereinbefore defined, or

25771-1192

b) a compound of general formula



wherein R^1 , R^2 and A are defined as mentioned in claims 1 to 8, and R^3 denotes one of the groups mentioned in the definition of R^3 hereinbefore wherein the amino or

5 imino group is protected by a protecting group, is deprotected,

and/or

any protecting groups used during the reaction are then cleaved and/or

the compound of general formula I thus obtained is resolved into an enantiomer and/or diastereomer and/or

10 the compound of formula I thus obtained is converted into a salt.

15. The process of claim 14, wherein the salt obtained is a physiologically acceptable salt with an inorganic acid or an organic acid.

16. The process of claim 14 or 15, wherein the leaving group is a halogen, substituted hydroxy, mercapto, sulphinyl, sulphonyl or sulphonoxy group.

15 17. The pharmaceutical composition of claim 10 or 11, for use in the
treatment of type I diabetes, type II diabetes, mellitus, arthritis, obesity,
allograft transplantation or calcitonin-induced osteoporosis.

18. Use of a compound as defined in any one of claims 1 to 9, or a
tautomer, enantiomer, diastereomer or mixture thereof or a salt thereof in the
20 treatment of type I diabetes, type II diabetes, mellitus, arthritis, obesity,
allograft transplantation or calcitonin-induced osteoporosis.

