



- (72) CHEN, HONG, US
(72) FREIMER, NELSON B., US
(71) MILLENNIUM PHARMACEUTICALS, INC., US
(71) THE REGENTS OF THE UNIVERSITY OF CALIFORNIA, US
(51) Int.Cl.⁶ C07H 17/00, C12Q 1/68, A61K 35/14, C12P 21/06, A01N 43/04,
C12N 15/00, C07K 14/00
(30) 1998/03/16 (60/078,044) US
(30) 1998/06/05 (60/088,312) US
(30) 1998/10/28 (60/106,056) US
(30) 1999/01/22 (09/236,134) US
(54) **PROCEDES ET COMPOSITIONS POUR DIAGNOSTIQUER ET
TRAITER LES TROUBLES LIES AU CHROMOSOME-18P**
(54) **METHODS AND COMPOSITIONS FOR DIAGNOSING AND
TREATING CHROMOSOME-18P RELATED DISORDERS**

(57) La présente invention concerne le gène mammifère HKNG1, un gène associé au trouble affectif bipolaire (BAD) chez les humains. Cette invention concerne en particulier des procédés d'évaluation diagnostique, de tests génétiques et de pronostic des troubles neuropsychiatriques liés au HKNG1, notamment la schizophrénie, les troubles déficitaires de l'attention, les troubles schizo-affectifs, les troubles affectifs bipolaires ou les troubles affectifs unipolaires.

(57) The present invention relates to the mammalian HKNG1 gene, a gene associated with bipolar affective disorder (BAD) in humans. The invention relates, in particular, to methods for the diagnostic evaluation, genetic testing and prognosis of HKNG1 neuropsychiatric disorders including schizophrenia, attention deficit disorder, a schizo-affective disorder, a bipolar affective disorder or a unipolar affective disorder.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07H 17/00, C07K 14/00, A01N 43/04, A61K 35/14, C12Q 1/68, C12P 21/06, C12N 15/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/47535 (43) International Publication Date: 23 September 1999 (23.09.99)</p>
<p>(21) International Application Number: PCT/US99/05606 (22) International Filing Date: 16 March 1999 (16.03.99) (30) Priority Data: 60/078,044 16 March 1998 (16.03.98) US 60/088,312 5 June 1998 (05.06.98) US 60/106,056 28 October 1998 (28.10.98) US 09/236,134 22 January 1999 (22.01.99) US (71) Applicants (for all designated States except US): MILLENNIUM PHARMACEUTICALS, INC. [US/US]; 238 Main Street, Cambridge, MA 02142-1017 (US). THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 300 Lakeside Drive, Oakland, CA 94612 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CHEN, Hong [CN/US]; 46 Aspinwal Avenue #2, Brookline, MA 02146 (US). FREIMER, Nelson, B. [US/US]; 630 29th Street, San Francisco, CA 94131 (US). (74) Agents: CORUZZI, Laura, A. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>

(54) Title: METHODS AND COMPOSITIONS FOR DIAGNOSING AND TREATING CHROMOSOME-18p RELATED DISORDERS

M K I K A E K W 1
 24
 28
 32
 36
 40
 44
 48
 52
 56
 60
 64
 68
 72
 76
 80
 84
 88
 92
 96
 100
 104
 108
 112
 116
 120
 124
 128
 132
 136
 140
 144
 148
 152
 156
 160
 164
 168
 172
 176
 180
 184
 188
 192
 196
 200
 204
 208
 212
 216
 220
 224
 228
 232
 236
 240
 244
 248
 252
 256
 260
 264
 268
 272
 276
 280
 284
 288
 292
 296
 300
 304
 308
 312
 316
 320
 324
 328
 332
 336
 340
 344
 348
 352
 356
 360
 364
 368
 372
 376
 380
 384
 388
 392
 396
 400
 404
 408
 412
 416
 420
 424
 428
 432
 436
 440
 444
 448
 452
 456
 460
 464
 468
 472
 476
 480
 484
 488
 492
 496
 500
 504
 508
 512
 516
 520
 524
 528
 532
 536
 540
 544
 548
 552
 556
 560
 564
 568
 572
 576
 580
 584
 588
 592
 596
 600
 604
 608
 612
 616
 620
 624
 628
 632
 636
 640
 644
 648
 652
 656
 660
 664
 668
 672
 676
 680
 684
 688
 692
 696
 700
 704
 708
 712
 716
 720
 724
 728
 732
 736
 740
 744
 748
 752
 756
 760
 764
 768
 772
 776
 780
 784
 788
 792
 796
 800
 804
 808
 812
 816
 820
 824
 828
 832
 836
 840
 844
 848
 852
 856
 860
 864
 868
 872
 876
 880
 884
 888
 892
 896
 900
 904
 908
 912
 916
 920
 924
 928
 932
 936
 940
 944
 948
 952
 956
 960
 964
 968
 972
 976
 980
 984
 988
 992
 996
 1000

984
 988
 992
 996
 1000
 1004
 1008
 1012
 1016
 1020
 1024
 1028
 1032
 1036
 1040
 1044
 1048
 1052
 1056
 1060
 1064
 1068
 1072
 1076
 1080
 1084
 1088
 1092
 1096
 1100
 1104
 1108
 1112
 1116
 1120
 1124
 1128
 1132
 1136
 1140
 1144
 1148
 1152
 1156
 1160
 1164
 1168
 1172
 1176
 1180
 1184
 1188
 1192
 1196
 1200
 1204
 1208
 1212
 1216
 1220
 1224
 1228
 1232
 1236
 1240
 1244
 1248
 1252
 1256
 1260
 1264
 1268
 1272
 1276
 1280
 1284
 1288
 1292
 1296
 1300
 1304
 1308
 1312
 1316
 1320
 1324
 1328
 1332
 1336
 1340
 1344
 1348
 1352
 1356
 1360
 1364
 1368
 1372
 1376
 1380
 1384
 1388
 1392
 1396
 1400
 1404
 1408
 1412
 1416
 1420
 1424
 1428
 1432
 1436
 1440
 1444
 1448
 1452
 1456
 1460
 1464
 1468
 1472
 1476
 1480
 1484
 1488
 1492
 1496
 1500
 1504
 1508
 1512
 1516
 1520
 1524
 1528
 1532
 1536
 1540
 1544
 1548
 1552
 1556
 1560
 1564
 1568
 1572
 1576
 1580
 1584
 1588
 1592
 1596
 1600
 1604
 1608
 1612
 1616
 1620
 1624
 1628
 1632
 1636
 1640
 1644
 1648
 1652
 1656
 1660
 1664
 1668
 1672
 1676
 1680
 1684
 1688
 1692
 1696
 1700
 1704
 1708
 1712
 1716
 1720
 1724
 1728
 1732
 1736
 1740
 1744
 1748
 1752
 1756
 1760
 1764
 1768
 1772
 1776
 1780
 1784
 1788
 1792
 1796
 1800
 1804
 1808
 1812
 1816
 1820
 1824
 1828
 1832
 1836
 1840
 1844
 1848
 1852
 1856
 1860
 1864
 1868
 1872
 1876
 1880
 1884
 1888
 1892
 1896
 1900
 1904
 1908
 1912
 1916
 1920
 1924
 1928
 1932
 1936
 1940
 1944
 1948
 1952
 1956
 1960
 1964
 1968
 1972
 1976
 1980
 1984
 1988
 1992
 1996
 2000

(57) Abstract

The present invention relates to the mammalian *HKNG1* gene, a gene associated with bipolar affective disorder (BAD) in humans. The invention relates, in particular, to methods for the diagnostic evaluation, genetic testing and prognosis of *HKNG1* neuropsychiatric disorders including schizophrenia, attention deficit disorder, a schizoaffective disorder, a bipolar affective disorder or a unipolar affective disorder.

**METHODS AND COMPOSITIONS FOR DIAGNOSING AND TREATING
CHROMOSOME-18p RELATED DISORDERS**

5

This is a continuation-in-part of U.S. application no. 09/236,134, filed on January 22, 1999 which claims priority under 35 U.S.C. § 119(e)(1) of provisional application no. 60/078,044 filed on March 16, 1998, of provisional application no. 60/088,312 filed on June 5, 1998, and of
10 provisional application no. 60/106,056 filed on October 28, 1998, each of which is incorporated herein by reference in its entirety.

1. INTRODUCTION

The present invention relates, first, to the *HKNG1* gene,
15 shown herein to be associated with central nervous system-related disorders, e.g., neuropsychiatric disorders, in particular, bipolar affective disorder and schizophrenia and with myopia-related disorders. The invention includes recombinant DNA molecules and cloning vectors comprising
20 sequences of the *HKNG1* gene, and host cells and non-human host organisms engineered to contain such DNA molecules and cloning vectors. The present invention further relates to *HKNG1* gene products, and to antibodies directed against such
25 *HKNG1* gene products. The present invention also relates to methods of using the *HKNG1* gene and gene product, including drug screening assays, and diagnostic and therapeutic methods for the treatment of *HKNG1*-mediated disorders, including
HKNG1-mediated neuropsychiatric disorders such as bipolar affective disorder, as well as *HKNG1*-mediated myopia disorders such as early-onset autosomal dominant myopia.

30

2. BACKGROUND OF THE INVENTION

There are only a few psychiatric disorders in which clinical manifestations of the disorder can be correlated with demonstrable defects in the structure and/or function of the nervous system. Well-known examples of such disorders include Huntington's disease, which can be traced to a mutation in a single gene and in which neurons in the striatum degenerate, and Parkinson's disease, in which dopaminergic neurons in the nigro-striatal pathway degenerate. The vast majority of psychiatric disorders, however, presumably involve subtle and/or undetectable changes, at the cellular and/or molecular levels, in nervous system structure and function. This lack of detectable neurological defects distinguishes "neuropsychiatric" disorders, such as schizophrenia, attention deficit disorders, schizoaffective disorder, bipolar affective disorders, or unipolar affective disorder, from neurological disorders, in which anatomical or biochemical pathologies are manifest. Hence, identification of the causative defects and the neuropathologies of neuropsychiatric disorders are needed in order to enable clinicians to evaluate and prescribe appropriate courses of treatment to cure or ameliorate the symptoms of these disorders.

One of the most prevalent and potentially devastating of neuropsychiatric disorders is bipolar affective disorder (BAD), also known as bipolar mood disorder (BP) or manic-depressive illness, which is characterized by episodes of elevated mood (mania) and depression (Goodwin, et al., 1990, *Manic Depressive Illness*, Oxford University Press, New York). The most severe and clinically distinctive forms of BAD are BP-I (severe bipolar affective (mood) disorder), which affects 2-3 million people in the United States, and SAD-M (schizoaffective disorder manic type). They are characterized by at least one full episode of mania, with or without episodes of major depression (defined by lowered mood, or depression, with associated disturbances in rhythmic

behaviors such as sleeping, eating, and sexual activity). BP-I often co-segregates in families with more etiologically heterogeneous syndromes, such as with a unipolar affective disorder such as unipolar major depressive disorder (MDD), which is a more broadly defined phenotype (Freimer and Reus, 5 1992, in *The Molecular and Genetic Basis of Neurological Disease*, Rosenberg, et al., eds., Butterworths, New York, pp. 951-965; McInnes and Freimer, 1995, *Curr. Opin. Genet. Develop.*, 5, 376-381). BP-I and SAD-M are severe mood disorders that are frequently difficult to distinguish from one another on a cross-sectional basis, follow similar 10 clinical courses, and segregate together in family studies (Rosenthal, et al., 1980, *Arch. General Psychiat.* 37, 804-810; Levinson and Levitt, 1987, *Am. J. Psychiat.* 144, 415-426; Goodwin, et al., 1990, *Manic Depressive Illness*, Oxford University Press, New York). Hence, methods for 15 distinguishing neuropsychiatric disorders such as these are needed in order to effectively diagnose and treat afflicted individuals.

Currently, individuals are typically evaluated for BAD using the criteria set forth in the most current version of the American Psychiatric Association's Diagnostic and 20 Statistical Manual of Mental Disorders (DSM). While many drugs have been used to treat individuals diagnosed with BAD, including lithium salts, carbamazepine and valproic acid, none of the currently available drugs are adequate. For example, drug treatments are effective in only approximately 60-70% of individuals diagnosed with BP-I. Moreover, it is 25 currently impossible to predict which drug treatments will be effective in, for example, particular BP-I affected individuals. Commonly, upon diagnosis, affected individuals are prescribed one drug after another until one is found to be effective. Early prescription of an effective drug treatment, therefore, is critical for several reasons, 30 including the avoidance of extremely dangerous manic episodes, the risk of progressive deterioration if effective

treatments are not found, and the risk of substantial side effects of current treatments.

The existence of a genetic component for BAD is strongly supported by segregation analyses and twin studies (Bertelson, et al., 1977, Br. J. Psychiat. 130, 330-351; 5 Freimer and Reus, 1992, in *The Molecular and Genetic Basis of Neurological Disease*, Rosenberg, et al., eds., Butterworths, New York, pp. 951-965; Pauls, et al., 1992, Arch. Gen. Psychiat. 49, 703-708). Efforts to identify the chromosomal location of genes that might be involved in BP-I, however, 10 have yielded disappointing results in that reports of linkage between BP-I and markers on chromosomes X and 11 could not be independently replicated nor confirmed in the re-analyses of the original pedigrees, indicating that with BAD linkage studies, even extremely high lod scores at a single locus, can be false positives (Baron, et al., 1987, Nature 326, 289- 15 292; Egeland, et al., 1987, Nature 325, 783-787; Kelsoe, et al., 1989, Nature 342, 238-243; Baron, et al., 1993, Nature Genet. 3, 49-55).

Recent investigations have suggested possible localization of BAD genes on chromosomes 18p and 21q, but in 20 both cases the proposed candidate region is not well defined and no unequivocal support exists for either location (Berrettini, et al., 1994, Proc. Natl. Acad. Sci. USA 91, 5918-5921; Murray, et al., 1994, Science 265, 2049-2054; Pauls, et al., 1995, Am. J. Hum. Genet. 57, 636-643; Maier, et al., 1995, Psych. Res. 59, 7-15; Straub, et al., 1994, 25 Nature Genet. 8, 291-296).

Mapping genes for common diseases believed to be caused by multiple genes, such as BAD, may be complicated by the typically imprecise definition of phenotypes, by etiologic heterogeneity, and by uncertainty about the mode of genetic transmission of the disease trait. With neuropsychiatric 30 disorders there is even greater ambiguity in distinguishing

individuals who likely carry an affected genotype from those who are genetically unaffected. For example, one can define an affected phenotype for BAD by including one or more of the broad grouping of diagnostic classifications that constitute the mood disorders: BP-I, SAD-M, MDD, and bipolar affective
5 (mood) disorder with hypomania and major depression (BP-II).

Thus, one of the greatest difficulties facing psychiatric geneticists is uncertainty regarding the validity of phenotype designations, since clinical diagnoses are based solely on clinical observation and subjective reports. Also, with complex traits such as neuropsychiatric disorders, it is
10 difficult to genetically map the trait-causing genes because: (1) neuropsychiatric disorder phenotypes do not exhibit classic Mendelian recessive or dominant inheritance patterns attributable to a single genetic locus, (2) there may be incomplete penetrance, *i.e.*, individuals who inherit a
15 predisposing allele may not manifest disease; (3) a phenocopy phenomenon may occur, *i.e.*, individuals who do not inherit a predisposing allele may nevertheless develop disease due to environmental or random causes; (4) genetic heterogeneity may exist, in which case mutations in any one of several genes may result in identical phenotypes.

20 Despite these difficulties, however, identification of the chromosomal location, sequence and function of genes and gene products responsible for causing neuropsychiatric disorders such as bipolar affective disorders is of great importance for genetic counseling, diagnosis and treatment of individuals in affected families.

25

3. SUMMARY OF THE INVENTION

The present invention relates, first, to the discovery, identification, and characterization of novel nucleic acid molecules that are associated with central nervous system-related disorders and processes, *e.g.*, human neuropsychiatric
30 disorders, such as schizophrenia, attention deficit disorder, schizoaffective disorder, dysthymic disorder, major

depressive disorder, and bipolar affective disorder (BAD) including severe bipolar affective (mood) disorder (BP-I), bipolar affective (mood) disorder with hypomania and major depression (BP-II). The invention further relates to the discovery, identification, and characterization of proteins
5 encoded by such nucleic acid molecules, or by degenerate, e.g., allelic or homologous, variants thereof. The invention further relates to the discovery, identification, and characterization of novel nucleic acid molecules that are associated with human myopia or nearsightedness, such as
10 early-onset, autosomal dominant myopia, as well as to the discovery, identification, and characterization of proteins encoded by such nucleic acid molecules or by degenerate variants thereof.

In particular, the nucleic acid molecules of the present invention represent, first, nucleic acid sequences
15 corresponding to the gene referred to herein as *HKNG1*. As demonstrated in the Examples presented below in Sections 6, 8 and 14, the *HKNG1* gene is associated with human CNS-related disorders, e.g., neuropsychiatric disorders, in particular
BAD. The *HKNG1* gene is associated with other human neuropsychiatric disorders as well, such as schizophrenia.
20 As demonstrated in the Example presented below in Section 14, the *HKNG1* gene is also associated with human myopia, such as early-onset autosomal dominant myopia.

In addition to the positive correlation between mutations within the *HKNG1* gene and individuals exhibiting
25 symptoms of BAD, described in Section 6 and 8, the present invention is further based, in part, on Applicants' discovery of novel *HKNG1* cDNA sequences. Applicants' discovery of such cDNA sequences has led to the elucidation of the *HKNG1* genomic (that is, upstream untranslated, intron/exon, and downstream untranslated) structure, and to the discovery of
30 full-length and alternately spliced *HKNG1* variants and the polypeptides encoded by such variants. These discoveries are

described in Sections 7 and 10, below. Applicants' discovery of such cDNA sequences has also led to the elucidation of novel mammalian (e.g., guinea pig and bovine) *HKNG1* sequences, and to the discovery of novel allelic variants and polymorphisms of such sequences. These discoveries are
5 described in Section 10 below.

The invention encompasses nucleic acid molecules which comprise the following nucleotide sequences: (a) nucleotide sequences (e.g., SEQ ID NOS: 1, 3, 5, 6, 36, and 37) that comprise a human *HKNG1* gene and/or encode a human *HKNG1* gene
10 product (e.g., SEQ ID NO: 2; SEQ ID NO: 4), as well as allelic variants, homologs and orthologs thereof, including nucleotide sequences (e.g., SEQ ID NOS: 38, 40, 42, 44, and 46-48) that encode non-human *HKNG1* gene products (e.g., SEQ ID NOS: 39, 41, 43, 45, and 49); (b) nucleotide sequences
15 comprising the novel *HKNG1* sequences disclosed herein that encode mutants of the *HKNG1* gene product in which sequences encoding all or a part of one or more of the *HKNG1* domains is deleted or altered, or fragments thereof; (c) nucleotide sequences that encode fusion proteins comprising a *HKNG1* gene
20 product (e.g., SEQ ID NO: 2; SEQ ID NO: 4), or a portion thereof fused to a heterologous polypeptide; and (d) nucleotide sequences within the *HKNG1* gene, as well as chromosome 18p nucleotide sequences flanking the *HKNG1* gene, which can be utilized, e.g., as primers, in the methods of the invention for identifying and diagnosing individuals at
25 risk for or exhibiting an *HKNG1*-mediated disorder, such as BAD or schizophrenia, or for diagnosing individuals at risk for or exhibiting a form of myopia such as early-onset autosomal dominant myopia. The nucleic acid molecules of (a) through (d), above, can include, but are not limited to, CDNA, genomic DNA, and RNA sequences.

30

The invention also encompasses the expression products of the nucleic acid molecules listed above; i.e., peptides, proteins, glycoproteins and/or polypeptides that are encoded by the above *HKNG1* nucleic acid molecules.

5 The compositions of the present invention further encompass agonists and antagonists of the *HKNG1* gene product, including small molecules (such as small organic molecules), and macromolecules (including antibodies), as well as nucleotide sequences that can be used to inhibit *HKNG1* gene expression (e.g., antisense and ribozyme molecules, and gene
10 or regulatory sequence replacement constructs) or to enhance *HKNG1* gene expression (e.g., expression constructs that place the *HKNG1* gene under the control of a strong promoter system).

The compositions of the present invention further
15 include cloning vectors and expression vectors containing the nucleic acid molecules of the invention, as well as hosts which have been transformed with such nucleic acid molecules, including cells genetically engineered to contain the nucleic acid molecules of the invention, and/or cells genetically
20 engineered to express the nucleic acid molecules of the invention. In addition to host cells and cell lines, hosts also include transgenic non-human animals (or progeny thereof), particularly non-human mammals, that have been engineered to express an *HKNG1* transgene, or "knock-outs" that have been engineered to not express *HKNG1*.

Transgenic non-human animals of the invention include
25 animals engineered to express an *HKNG1* transgene at higher or lower levels than normal, wild-type animals. The transgenic animals of the invention also include animals engineered to express a mutant variant or polymorphism of an *HKNG1*
transgene which is associated with *HKNG1*-mediated disorder,
30 for example an *HKNG1*-mediated neuropsychiatric disorders, such as BAD and schizophrenia, or, alternatively, a myopia

disorder such as early-onset autosomal dominant myopia. The transgenic animals of the invention further include the progeny of such genetically engineered animals.

The invention further relates to methods for the treatment of *HKNG1*-mediated disorders in a subject, such as
5 *HKNG1*-mediated neuropsychiatric disorders and *HKNG1*-mediated myopia disorders, wherein such methods comprise administering a compound which modulates the expression of a *HKNG1* gene and/or the synthesis or activity of a *HKNG1* gene product so symptoms of the disorder are ameliorated.

10 The invention further relates to methods for the treatment of *HKNG1*-mediated disorders in a subject, such as *HKNG1*-mediated neuropsychiatric disorders and *HKNG1*-mediated myopia disorders, resulting from *HKNG1* gene mutations or aberrant levels of *HKNG1* expression or activity, wherein such
15 methods comprise supplying the subject with a nucleic acid molecule encoding an unimpaired *HKNG1* gene product such that an unimpaired *HKNG1* gene product is expressed and symptoms of the disorder are ameliorated.

The invention further relates to methods for the treatment of *HKNG1*-mediated disorders in a subject, such as
20 *HKNG1*-mediated neuropsychiatric disorders and *HKNG1*-mediated myopia disorders, resulting from *HKNG1* gene mutations or from aberrant levels of expression or activity, wherein such methods comprise supplying the subject with a cell comprising a nucleic acid molecule that encodes an unimpaired *HKNG1* gene
25 product such that the cell expresses the unimpaired *HKNG1* gene product and symptoms of the disorder are ameliorated.

The invention also encompasses pharmaceutical formulations and methods for treating *HKNG1*-mediated disorders, including neuropsychiatric disorders, such as BAD and schizophrenia, and myopia disorders, such as early-onset
30 autosomal dominant myopia, involving *HKNG1* gene.

In addition, the present invention is directed to methods that utilize the *HKNG1* nucleic acid sequences, chromosome 18p nucleotide sequences flanking the *HKNG1* human gene and/or *HKNG1* gene product sequences for mapping the
5 chromosome 18p region, and for the diagnostic evaluation, genetic testing and prognosis of a *HKNG1*-mediated disorder, such as a *HKNG1*-mediated neuropsychiatric disorder or a *HKNG1*-mediated myopia disorder. For example, in one embodiment, the invention relates to methods for diagnosing
10 *HKNG1*-mediated disorders, wherein such methods comprise measuring *HKNG1* gene expression in a patient sample, or detecting a *HKNG1* polymorphism or mutation in the genome of a mammal, including a human, suspected of exhibiting such a disorder. In one embodiment, nucleic acid molecules encoding *HKNG1* can be used as diagnostic hybridization probes or as
15 primers for diagnostic PCR analysis for the identification of *HKNG1* gene mutations, allelic variations and regulatory defects in the *HKNG1* gene which correlate with neuropsychiatric disorders such as BAD or schizophrenia.

The invention still further relates to methods for
20 identifying compounds which modulate the expression of the *HKNG1* gene and/or the synthesis or activity of the *HKNG1* gene products, including therapeutic compounds, which reduce or eliminate the symptoms of *HKNG1*-mediated disorders, including *HKNG1*-mediated neuropsychiatric disorders such as BAD and schizophrenia. In particular, cellular and non-cellular
25 assays are described that can be used to identify compounds that interact with the *HKNG1* gene product, e.g., modulate the activity of the *HKNG1* and/or bind to the *HKNG1* gene product. Such cell-based assays of the invention utilize cells, cell lines, or engineered cells or cell lines that express the
30 *HKNG1* gene product.

In one embodiment, such methods comprise contacting a compound to a cell that expresses a *HKNG1* gene, measuring the level of *HKNG1* gene expression, gene product expression or gene product activity, and comparing this level to the level of *HKNG1* gene expression, gene product expression or gene product activity produced by the cell in the absence of the compound, such that if the level obtained in the presence of the compound differs from that obtained in its absence, a compound that modulates the expression of the *HKNG1* gene and/or the synthesis or activity of the *HKNG1* gene products has been identified.

In another embodiment, such methods comprise contacting a compound to a cell that expresses a *HKNG1* gene and also comprises a reporter construct whose transcription is dependent, at least in part, on *HKNG1* expression or activity. In such an embodiment, the level of reporter transcription is measured and compared to the level of reporter transcription in the cell in the absence of the compound. If the level of reporter transcription obtained in the presence of the compound differs from that obtained in its absence, a compound that modulates expression of *HKNG1* or genes involved in *HKNG1*-related pathways or signal transduction has been identified.

In yet another embodiment, such methods comprise administering a compound to a host, such as a transgenic animal, that expresses an *HKNG1* transgene or a mutant *HKNG1* transgene associated with an *HKNG1*-mediated disorder such as a neuropsychiatric disorder (e.g., BAD or schizophrenia), or to an animal, e.g., a knock-out animal, that does not express *HKNG1*, and measuring the level of *HKNG1* gene expression, gene product expression, gene product activity, or symptoms of an *HKNG1*-mediated disorder such as an *HKNG1*-mediated neuropsychiatric disorder (e.g., BAD or schizophrenia). The

measured level is compared to the level obtained in a host that is not exposed to the compound, such that if the level obtained when the host is exposed to the compound differs from that obtained in a host not exposed to the compound, a compound modulates the expression of the mammalian *HKNG1* gene
5 and/or the synthesis or activity of the mammalian *HKNG1* gene products, and/or the symptoms of an *HKNG1*-mediated disorder such as a neuropsychiatric disorder (e.g., BAD or schizophrenia), has been identified.

The present invention still further relates to
10 pharmacogenomic and pharmacogenetic methods for selecting an effective drug to administer to an individual having a *HKNG1*-mediated disorder. Such methods are based on the detection of genetic polymorphisms in the *HKNG1* gene or variations in *HKNG1* gene expression due to, e.g., altered methylation,
15 differential splicing, or post-translational modification of the *HKNG1* gene product which can affect the safety and efficacy of a therapeutic agent.

As briefly discussed above, the present invention is based, in part, on the genetic and physical mapping of the *HKNG1* gene to a specific portion of the short arm of human
20 chromosome 18 that is associated with human neuropsychiatric disorders, in particular, bipolar affective disorder. These results are described in the Example presented, below, in Section 6. The invention is also based on the elucidation of the *HKGN1* nucleotide sequence, amino acid sequence and
expression pattern, as described in the Example presented,
25 below, in Section 7. The invention is further based on the identification of specific mutations and/or polymorphisms within the *HKNG1* gene which positively correlate with neuropsychiatric disorders, in particular, BAD, as described in the Example presented below in Section 8. These mutations
include a point mutation discovered in an individual affected
30 by BAD which is absent from the corresponding wild-type

nucleic acid derived from non-affected individuals and linkage disequilibrium of three markers showing cosegregation with a population of individuals with BAD. This mutation is a single base mutation which results in a mutant *HKNG1* gene product comprising substitution of a lysine residue for the wild-type glutamic acid residue at *HKNG1* amino acid position 202 of the polypeptide of SEQ ID NO:2 or the *HKNG1* amino acid residue 184 of the polypeptide of SEQ ID NO:4. These mutations further include the mutations discovered in schizophrenic and BAD patients that are detailed in FIGS. 5A-5B.

3.1. DEFINITIONS

As used herein, the following terms shall have the abbreviations indicated.

15 BAC, bacterial artificial chromosomes
 BAD, bipolar affective disorder(s)
 BP, bipolar mood disorder
 BP-I, severe bipolar affective (mood) disorder
 BP-II, bipolar affective (mood) disorder with
 hypomania and major depression
 20 bp, base pair(s)
 EST, expressed sequence tag
HKNG1, Hong Kong new gene 1
 lod, logarithm of odds
 MDD, unipolar major depressive disorder
 ROS, reactive oxygen species
 25 RT-PCR, reverse transcriptase PCR
 SSCP, single-stranded conformational polymorphism
 SAD-M, schizoaffective disorder manic type
 STS, sequence tagged site
 YAC, yeast artificial chromosome

"*HKNG1*-mediated disorders" include disorders involving
 30 an aberrant level of *HKNG1* gene expression, gene product

synthesis and/or gene product activity relative to levels found in clinically normal individuals, and/or relative to levels found in a population whose level represents a baseline, average *HKNG1* level. While not wishing to be bound by any particular mechanism, it is to be understood that
5 disorder symptoms can, for example, be caused, either directly or indirectly, by such aberrant levels. Alternatively, it is to be understood that such aberrant levels can, either directly or indirectly, ameliorate disorder symptoms, (e.g., as in instances wherein aberrant
10 levels of *HKNG1* suppress the disorder symptoms caused by mutations within a second gene).

HNKG1-mediated disorders include, for example, central nervous system (CNS) disorders. CNS disorders include, but are not limited to cognitive and neurodegenerative disorders such as Alzheimer's disease, senile dementia, Huntington's
15 disease, amyotrophic lateral sclerosis, and Parkinson's disease, as well as Gilles de la Tourette's syndrome, autonomic function disorders such as hypertension and sleep disorders, and neuropsychiatric disorders that include, but are not limited to schizophrenia, schizoaffective disorder, attention deficit disorder, dysthymic disorder, major
20 depressive disorder, mania, obsessive-compulsive disorder, psychoactive substance use disorders, anxiety, panic disorder, as well as bipolar affective disorder, e.g., severe bipolar affective (mood) disorder (BP-I), bipolar affective (mood) disorder with hypomania and major depression (BP-II). Further CNS-related disorders include, for example, those
25 listed in the American Psychiatric Association's Diagnostic and Statistical manual of Mental Disorders (DSM), the most current version of which is incorporated herein by reference in its entirety.

"*HKNG1*-mediated processes" include processes dependent and/or responsive, either directly or indirectly, to levels
30 of *HKNG1* gene expression, gene product synthesis and/or gene

product activity. Such processes can include, but are not limited to, developmental, cognitive and autonomic neural and neurological processes, such as, for example, pain, appetite, long term memory and short term memory.

5

4. BRIEF DESCRIPTION OF THE FIGURES

FIG. 1A-1B. The nucleotide sequence of human *HKNG1* cDNA (SEQ ID NO: 1) is depicted on the bottom line. The top line depicts the full length amino acid sequence of human *HKNG1* polypeptide (SEQ ID NO: 2) encoded by the human *HKNG1* cDNA sequence. The nucleotide sequence encoding SEQ ID NO:2 corresponds to SEQ ID NO:5.

FIG. 2A-2B. Nucleotide sequence of an alternately spliced human *HKNG1* variant, referred to as *HKNG1-V1* (SEQ ID NO: 3) is depicted on the bottom line. The derived amino acid sequence of the human *HKNG1* gene product (SEQ ID NO: 4) encoded by this alternately spliced cDNA variant is depicted on the top line. The nucleotide sequence encoding SEQ ID NO:4 corresponds to SEQ ID NO:6

FIG. 3A-3R. Genomic sequence of the human *HKNG1* gene (SEQ ID NO. 7). Exons are in bold and the 3' and 5' UTRs (untranslated regions) are underlined.

FIG. 4. Summary of in situ hybridization analysis of *HKNG1* mRNA distribution in normal human brain tissue.

FIGS. 5A-B. *HKNG1* polymorphisms relative to the *HKNG1* wild-type sequence. These polymorphisms were isolated from a collection of schizophrenic patients of mixed ethnicity from the United States (FIG. 5A) and from the San Francisco BAD collection (FIG. 5B).

FIGS. 6A-B. Nucleotide sequence of the RT-PCR products for *HKNG1-V2* (FIG. 6A; SEQ ID NO:36) and *HKNG1-V3* (FIG. 6B; SEQ ID NO:37).

FIG 7. The cDNA sequence (SEQ ID NO:38) and the predicted amino acid sequence (SEQ ID NO:39) of the guinea pig *HKNG1* ortholog *gphkng1815*.

FIG. 8. The cDNA sequence (SEQ ID NO:40) and the predicted amino acid sequence (SEQ ID NO:41) of *gphkng 7b*, an allelic variant of the guinea pig *HKNG1* ortholog *gphkng1815*.

FIG. 9. The cDNA sequence (SEQ ID NO:42) and the predicted amino acid sequence (SEQ ID NO:43) of *gphkng 7c*, an allelic variant of the guinea pig *HKNG1* ortholog *gphkng1815*.

FIG. 10. The cDNA sequence (SEQ ID NO:44) and the predicted amino acid sequence (SEQ ID NO:45) of *gphkng 7d*, an allelic variant of the guinea pig *HKNG1* ortholog *gphkng1815*.

FIG. 11. The cDNA sequence (SEQ ID NO:46) and the predicted amino acid sequence (SEQ ID NO:49) of the allelic variant *bhkng1* of the bovine *HKNG1* ortholog.

FIG. 12. The cDNA sequence (SEQ ID NO:47) and the predicted amino acid sequence (SEQ ID NO:49) of the allelic variant *bhkng2* of the bovine *HKNG1* homologue.

FIG. 13. The cDNA sequence (SEQ ID NO:48) and the predicted amino acid sequence (SEQ ID NO:49) of the allelic variant *bhkng3* of the bovine *HKNG1* homologue.

FIG. 14A-B. Alignments of the guinea pig *HKNG1* cDNA (FIG. 14A) and predicted amino acid (FIG. 14B) sequences for *gphkng1815*, *gphkng 7b*, *gphkng7c*, and *gphkng 7d*.

FIG. 15. Alignments of the cDNA sequences of the bovine *HKNG1* allelic variants *bhkng1*, *bhkng2*, and *bhkng3*.

FIG. 16. Alignments of the human (*hkng_aa*), bovine (*bhkng1_aa*) and guinea pig (*gphkng1815_aa*) *HKNG1* amino acid sequences.

FIG. 17. Alignments of the human *HKNG1* protein sequences; top line: the mature secreted *HKNG1* protein sequence (SEQ ID NO:51); second line: immature *HKNG1* protein form 1 (IPF1; SEQ ID NO:2); third line: immature *HKNG1* protein form 2 (IPF2; SEQ ID NO:64); bottom line: immature *HKNG1* protein form 3 (IPF3; SEQ ID NO:4).

FIG. 18. The nucleotide sequence of human *HKNG1* splice variant *HKNG1Δ7* cDNA (SEQ ID NO: 65) is depicted on the

bottom line. The top line depicts the full length amino acid sequence of human HKNG1 Δ 7 polypeptide (SEQ ID NO: 66) encoded by the human HKNG1 Δ 7 cDNA sequence.

5. DETAILED DESCRIPTION OF THE INVENTION

5

5.1. THE HKNG1 GENE

HKNG1 nucleic acid molecules are described in the section. Unless otherwise stated, the term "HKNG1 nucleic acid" refers collectively to the sequences described herein.

A human HKNG1 cDNA sequence (SEQ ID NO: 1) encoding the full length amino acid sequence (SEQ ID NO: 2) of the HKNG1 polypeptide is shown in FIG. 1A-1B. The human HKNG1 gene encodes a secreted polypeptide of 495 amino acid residues, as shown in FIG. 1A-1B, and SEQ ID NO: 2. The nucleotide sequence of the portion of the cDNA corresponding to the coding sequence for HKNG1 (SEQ ID NO:2) is depicted as SEQ ID NO:5.

The HKNG1 sequences of the invention also include splice variants of the HKNG1 sequences described herein. For example, an alternately spliced human HKNG1 cDNA sequence, referred to as HKNG1-V1 (SEQ ID NO: 3) encoding a human HKNG1 variant gene product (*i.e.*, the HKNG1-V1 gene product) is shown in FIG. 2A-2B. This splice variant of a human HKNG1 gene encodes a secreted polypeptide of 477 amino acid residues, as shown in FIG. 2A-2B, and SEQ ID NO:4. The nucleotide sequence of the portion of the cDNA corresponding to the coding sequence for HKNG1 (SEQ ID NO:4) is depicted in SEQ ID NO:6.

Another alternately spliced human HKNG1 cDNA sequence (SEQ ID NO:65), referred to as HKNG1 Δ 7, encodes a second HKNG1 variant gene product (the HKNG1 Δ 7 gene product) which is depicted in FIG. 18. This splice variant of the human

30

HKNG1 gene encodes the variant polypeptide shown in FIG. 18 (SEQ ID NO:66).

The genomic structure of the human *HKNG1* gene has been elucidated and is depicted in FIG. 3A-3R, with the *HKNG1* exons indicated in bold type, and the 5'- and 3'-untranslated regions indicated by underlining. The wild-type genomic sequence of the *HKNG1* gene is depicted in FIG. 3A-3R and SEQ ID NO:7.

Non-human homologues or orthologs mammalian orthologs, e.g., of the human *HKNG1* sequences discussed above are also provided. Specifically, a guinea pig cDNA sequence (SEQ ID NO:38), referred to herein as *gphkng1815*, encoding the full length amino acid sequence (SEQ ID NO:39) of a guinea pig *HKNG1* ortholog is shown in FIG. 7. This guinea pig cDNA sequence encodes a gene product of 466 amino acid residues, as shown in FIG. 7 and in SEQ ID NO:39.

Allelic variants of this guinea pig *HKNG1* ortholog, referred to as *gphkng 7b*, *gphkng 7c*, and *gphkng 7d* (SEQ ID NOS:40, 42, and 44, respectively), are shown in FIGS. 8-10, respectively. The allelic variants *gphkng7b*, *gphkng7c*, and *gphkng7d* each encode variants of the guinea pig *gphkng1815* *HKNG1* gene product which contain deletions of 16, 92, and 93 amino acids, respectively, as shown in FIGS. 8-10, in SEQ ID NOS:41, 43, and 45, respectively, and in the sequence alignment in FIG. 14B.

Bovine *HKNG1* ortholog cDNA sequences (SEQ ID NOS: 46-48), referred to herein as *bhkng1*, *bhkng2*, and *bhkng3*, and each encoding the same bovine ortholog gene product are shown in FIGS. 11-13, respectively. The bovine *HKNG1* allelic variants encode the same gene product, *i.e.*, a 465 amino acid protein, as shown in FIGS. 11-13 and in SEQ ID NO:49.

The *HKNG1* gene nucleic acid molecules of the invention include: (a) nucleotide sequences and fragments thereof

(e.g., SEQ ID NOS: 1, 3, 5, 6, 7, 36, 37, and 65) that encode a *HKNG1* gene product (e.g., SEQ ID NOS: 2, 4 and 66), as well as homologues, orthologs and allelic variants of such sequences and fragments thereof (e.g., SEQ ID NOS:38, 40, 42, 44, and 46-48) which encode homologue or ortholog *HKNG1* gene products (e.g., SEQ ID NOS:39, 41, 43, 45, and 49); (b) nucleotide sequences that encode one or more functional domains of a *HKNG1* gene product including, but not limited to, nucleic acid sequences that encode a signal sequence domain, or one or more clusterin domains as described in Section 5.2 below; (c) nucleotide sequences that comprise *HKNG1* gene sequences of upstream untranslated regions, intronic regions, and/or downstream untranslated regions or fragments thereof of the *HKNG1* nucleotide sequences in (a) above; (d) nucleotide sequences comprising the novel *HKNG1* sequences disclosed herein that encode mutants of the *HKNG1* gene product in which all or a part of one or more of the domains is deleted or altered, as well as fragments thereof; (e) nucleotide sequences that encode fusion proteins comprising a *HKNG1* gene product (e.g., SEQ ID NO: 2, 4, 39, 41, 43, 45, 49 and 65), or a portion thereof fused to a heterologous polypeptide; and (f) nucleotide sequences (e.g., primers) within the *HKNG1* gene, and chromosome 18p nucleotide sequences flanking the *HKNG1* gene which can be utilized as part of the methods of the invention for identifying and diagnosing individuals at risk for or exhibiting an *HKNG1*-mediated disorder, such as BAD, or myopia.

The *HKNG1* nucleotide sequences of the invention further include nucleotide sequences corresponding to the nucleotide sequences of (a)-(f) above wherein one or more of the exons, or fragments thereof, have been deleted. In one preferred embodiment, the *HKNG1* nucleotide sequence of the invention is

a sequence wherein the exon corresponding to exon 7 of SEQ ID NO:7, or a fragment thereof, has been deleted.

The *HKNG1* nucleotide sequences of the invention also include nucleotide sequences that have at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or more nucleotide sequence identity to the *HKNG1* nucleotide sequences of (a)-(f) above. The *HKNG1* nucleotide sequences of the invention further include nucleotide sequences that encode polypeptides having at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or higher amino acid sequence identity to the polypeptides encoded by the *HKNG1* nucleotide sequences of (a)-(f), e.g., SEQ ID NOS: 2, 4, 39, 41, 43, 45, 49, and 66 above.

To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical overlapping positions/total # of positions x 100%). In one embodiment, the two sequences are the same length.

The determination of percent identity between two sequences can also be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-2268, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA*

90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al. (1990) *J. Mol. Biol.* 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 5 12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to a protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as 10 described in Altschul et al. (1997) *Nucleic Acids Res.* 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (*Id.*). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be 15 used (see <http://www.ncbi.nlm.nih.gov>). Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, (1988) *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is 20 part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

The percent identity between two sequences can be determined using techniques similar to those described above, 25 with or without allowing gaps. In calculating percent identity, typically only exact matches are counted.

The *HKNG1* nucleotide sequences of the invention further include: (a) any nucleotide sequence that hybridizes to a *HKNG1* nucleic acid molecule of the invention under stringent conditions, e.g., hybridization to filter-bound DNA in 6x 30 sodium chloride/sodium citrate (SSC) at about 45°C followed

by one or more washes in 0.2xSSC/0.1% SDS at about 50-65°C, or (b) under highly stringent conditions, e.g., hybridization to filter-bound nucleic acid in 6xSSC at about 45°C followed by one or more washes in 0.1xSSC/0.2% SDS at about 68°C, or under other hybridization conditions which are apparent to those of skill in the art (see, for example, Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at pp. 6.3.1-6.3.6 and 2.10.3). Preferably the *HKNG1* nucleic acid molecule that hybridizes to the nucleotide sequence of (a) and (b), above, is one that comprises the complement of a nucleic acid molecule that encodes a *HKNG1* gene product. In a preferred embodiment, nucleic acid molecules comprising the nucleotide sequences of (a) and (b), above, encode gene products, e.g., gene products functionally equivalent to an *HKNG1* gene product.

Functionally equivalent *HKNG1* gene products include naturally occurring *HKNG1* gene products present in the same or different species. In one embodiment, *HKNG1* gene sequences in non-human species map to chromosome regions syntenic to the human 18p chromosome location within which human *HKNG1* lies. Functionally equivalent *HKNG1* gene products also include gene products that retain at least one of the biological activities of the *HKNG1* gene products, and/or which are recognized by and bind to antibodies (polyclonal or monoclonal) directed against the *HKNG1* gene products.

Among the nucleic acid molecules of the invention are deoxyoligonucleotides ("oligos") which hybridize under highly stringent or stringent conditions to the *HKNG1* nucleic acid molecules described above. In general, for probes between 14 and 70 nucleotides in length the melting temperature (TM) is calculated using the formula:

$$T_m(^{\circ}\text{C}) = 81.5 + 16.6(\log[\text{monovalent cations (molar)}]) + 0.41 (\%$$

G+C)-(500/N) where N is the length of the probe. If the hybridization is carried out in a solution containing formamide, the melting temperature is calculated using the equation $T_m(^{\circ}\text{C})=81.5+16.6(\log[\text{monovalent cations (molar)}])+0.41(\% \text{ G+C})-(0.61\% \text{ formamide})-(500/N)$ where N is
5 the length of the probe. In general, hybridization is carried out at about 20-25 degrees below T_m (for DNA-DNA hybrids) or 10-15 degrees below T_m (for RNA-DNA hybrids).

Exemplary highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for about 14-base oligos), 48°C (for about 17-base oligos), 55°C
10 (for about 20-base oligos), and 60°C (for about 23-base oligos).

These nucleic acid molecules may encode or act as antisense molecules, useful, for example, in *HKNG1* gene regulation, and/or as antisense primers in amplification reactions of *HKNG1* gene nucleic acid sequences. Further,
15 such sequences may be used as part of ribozyme and/or triple helix sequences, also useful for *HKNG1* gene regulation. Still further, such molecules may be used as components of diagnostic methods whereby, for example, the presence of a particular *HKNG1* allele involved in a *HKNG1*-related disorder,
20 e.g., a neuropsychiatric disorder, such as BAD, may be detected.

Fragments of the *HKNG1* nucleic acid molecules can be at least 10 nucleotides in length. In alternative embodiments, the fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 1000, 1500, 2000, 2500, 3000, 3500,
25 4000, 4500, 5000, or more contiguous nucleotides in length. Alternatively, the fragments can comprise sequences that encode at least 10, 20, 30, 40, 50, 100, 150, 200, 250, 300, 350, 400, 450 or more contiguous amino acid residues of the *HKNG1* gene products. Fragments of the *HKNG1* nucleic acid molecules can also refer to *HKNG1* exons or introns, and,
30 further, can refer to portions of *HKNG1* coding regions that

encode domains (e.g., clusterin domains) of *HKNG1* gene products.

The *HKNG1* nucleotide sequences of the invention can be readily obtained, for example, by standard sequencing and the sequence provided herein.

As will be appreciated by those skilled in the art, DNA sequence polymorphisms of a *HKNG1* gene will exist within a population of individual organisms (e.g., within a human population). Such polymorphisms may exist, for example, among individuals within a population due to natural allelic variation. Such polymorphisms include ones that lead to changes in amino acid sequence. An allele is one of a group of genes which occur alternatively at a given genetic locus.

As used herein, the phrase "allelic variant" refers to a nucleotide sequence which occurs at a given locus or to a gene product encoded by that nucleotide sequence. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of a given gene. Alternative alleles can be identified by sequencing the gene of interest in a number of different individuals. This can be readily carried out by using hybridization probes to identify the same genetic locus in a variety of individuals. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a polypeptide of the invention. The term can further include nucleic acid molecules comprising upstream and/or exon/intron sequences and structure.

With respect to *HKNG1* allelic variants, any and all such nucleotide variations and resulting amino acid polymorphisms or variations that are the result of natural allelic variation of the *HKNG1* gene are intended to be within the scope of the present invention. Such allelic variants include, but are not limited to, ones that do not alter the functional activity of the *HKNG1* gene product. Variants include, but are not limited to, variants comprising the

polymorphisms summarized in FIGS. 5A-B and a variant which encodes a full length *HKNG1* polypeptide comprising a substitution of a lysine amino acid at amino acid residue 202 of the *HKNG1* polypeptide shown in FIG. 1A-1B and SEQ ID NO:2 or the *HKNG1* amino acid residue 184 of the polypeptide of SEQ ID NO:4.

With respect to the cloning of additional allelic variants of the human *HKNG1* gene and homologues and orthologs from other species (e.g., guinea pig, cow, mouse), the isolated *HKNG1* gene sequences disclosed herein may be labeled and used to screen a cDNA library constructed from mRNA obtained from appropriate cells or tissues (e.g., brain or retinal tissues) derived from the organism (e.g., guinea pig, cow, and mouse) of interest. The hybridization conditions used should generally be of a lower stringency when the cDNA library is derived from an organism different from the type of organism from which the labeled sequence was derived, and can routinely be determined based on, e.g., relative relatedness of the target and reference organisms.

Alternatively, the labeled fragment may be used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. Appropriate stringency conditions are well known to those of skill in the art as discussed above, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook, et al., 1989, *Molecular Cloning, A Laboratory Manual, Second Edition*, Cold Spring Harbor Press, N.Y.; and Ausubel, et al., 1989-1999, *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y., both of which are incorporated herein by reference in their entirety.

30

Further, a *HKNG1* gene allelic variant may be isolated from, for example, human nucleic acid, by performing PCR using two degenerate oligonucleotide primer pools designed on the basis of amino acid sequences within the *HKNG1* gene product disclosed herein. The template for the reaction may be cDNA obtained by reverse transcription of mRNA prepared from, for example, human or non-human cell lines or tissue known or suspected to express a wild type or mutant *HKNG1* gene allele (such as, for example, brain cells, including brain cells from individuals having BAD). In one embodiment, the allelic variant is isolated from an individual who has a *HKNG1*-mediated disorder. Such variants are described in the examples below.

The PCR product may be subcloned and sequenced to ensure that the amplified sequences represent the sequences of a *HKNG1* gene nucleic acid sequence. The PCR fragment may then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment may be labeled and used to screen a bacteriophage cDNA library. Alternatively, the labeled fragment may be used to isolate genomic clones via the screening of a genomic library.

PCR technology may also be utilized to isolate full length cDNA sequences. For example, RNA may be isolated, following standard procedures, from an appropriate cellular or tissue source (*i.e.*, one known, or suspected, to express the *HKNG1* gene, such as, for example, brain tissue samples obtained through biopsy or post-mortem). A reverse transcription reaction may be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment may easily

be isolated. For a review of cloning strategies that may be used, see e.g., Sambrook et al., 1989, *supra*, or Ausubel et al., *supra*.

A cDNA of an allelic, e.g., mutant, variant of the *HKNG1* gene may be isolated, for example, by using PCR, a technique that is well known to those of skill in the art. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying a mutant *HKNG1* allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant *HKNG1* allele to that of the normal *HKNG1* allele, the mutation(s) responsible for the loss or alteration of function of the mutant *HKNG1* gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant *HKNG1* allele, or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant *HKNG1* allele. An unimpaired *HKNG1* gene or any suitable fragment thereof may then be labeled and used as a probe to identify the corresponding mutant *HKNG1* allele in such libraries. Clones containing the mutant *HKNG1* gene sequences may then be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant *HKNG1* allele in an individual suspected of or known to carry such a

mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against the normal *HKNG1* gene product, as described, below, in Section 5.3. (For screening techniques, see, for example, Harlow and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.)

In cases where a *HKNG1* mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), a polyclonal set of anti-*HKNG1* gene product antibodies are likely to cross-react with the mutant *HKNG1* gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

HKNG1 mutations or polymorphisms can further be detected using PCR amplification techniques. Primers can routinely be designed to amplify overlapping regions of the whole *HKNG1* sequence including the promoter regulating region. In one embodiment, primers are designed to cover the exon-intron boundaries such that, coding regions can be scanned for mutations. Exemplary primers for analyzing *HKNG1* exons are provided in Table 1, of Section 5.6, below.

The invention also includes nucleic acid molecules, preferably DNA molecules, that are the complements of the nucleotide sequences of the preceding paragraphs.

In certain embodiments, the nucleic acid molecules of the invention are present as part of nucleic acid molecules comprising nucleic acid sequences that do not contain heterologous (e.g., cloning vector or expression vector) sequences. In other embodiments, the nucleic acid molecules of the invention further comprise vector sequences, e.g., cloning vectors or expression vectors.

5.2. PROTEIN PRODUCTS OF THE HKNG1 GENE

HKNG1 gene products or peptide fragments thereof, can be prepared for a variety of uses. For example, such gene products, or peptide fragments thereof, can be used for the generation of antibodies, in diagnostic assays, or for the identification of other cellular or extracellular gene products involved in the regulation of *HKNG1*-mediated disorders, e.g., neuropsychiatric disorders, such as BAD.

The *HKNG1* gene products of the invention include, but are not limited to, human *HKNG1* gene products, e.g., polypeptides comprising the amino acid sequences depicted in FIGS. 1A-1B, 2A-2B, 17, and 18 (*i.e.*, SEQ ID NOS:2, 4, 51, and 66). The *HKNG1* gene products of the invention also include non-human, e.g., mammalian (such as bovine or guinea pig), *HKNG1* gene products. These include, but are not limited to, polypeptides comprising the amino acid sequences depicted in FIGS. 7-13 (*i.e.*, SEQ ID NOS:39, 41, 43, 45, and 49).

HKNG1 gene product, sometimes referred to herein as an "*HKNG1* protein" or "*HKNG1* polypeptide," includes those gene products encoded by the *HKNG1* gene sequences depicted in FIGS. 1A-1B, 2A-2B, 7-13, 17, and 18, as well as gene products encoded by other human allelic variants and non-human variants of *HKNG1* that can be identified by the methods herein described. Among such *HKNG1* gene product variants are gene products comprising *HKNG1* amino acid residues encoded by the polymorphisms depicted in FIGS. 5A and 5B. Such gene product variants also include a variant of the *HKNG1* gene product depicted in FIG. 1 (SEQ ID NO:2) wherein the amino acid residue Lys202 is mutated to a glutamic acid residue. Such *HKNG1* gene product variants also include a variant of the *HKNG1* gene product depicted in FIG. 2 (SEQ ID NO:4)

wherein the amino acid residue Lys184 is mutated to a glutamic acid residue.

In addition, *HKNG1* gene products may include proteins that represent functionally equivalent gene products. Functionally equivalent gene products may include, for
5 example, gene products encoded by one of the *HKNG1* nucleic acid molecules described in Section 5.1, above. In preferred embodiments, such functionally equivalent *HKNG1* gene products are naturally occurring gene products. Functionally
10 equivalent *HKNG1* gene products also include gene products that retain at least one of the biological activities of the *HKNG1* gene products described above, and/or which are recognized by and bind to antibodies (polyclonal or
monoclonal) directed against *HKNG1* gene products.

Equivalent *HKNG1* gene product may contain deletions,
15 including internal deletions, additions, including additions yielding fusion proteins, or substitutions of amino acid residues within and/or adjacent to the amino acid sequence encoded by the *HKNG1* gene sequences described, above, in
Section 5.1. Generally, deletions will be deletions of single amino acid residues, or deletions of no more than
20 about 2, 3, 4, 5, 10 or 20 amino acid residues, either contiguous or non-contiguous. Generally, additions or substitutions, other than additions that yield fusion proteins, will be additions or substitutions of single amino
acid residues, or additions or substitutions of no more than
25 about 2, 3, 4, 5, 10 or 20 amino acid residues, either contiguous or non-contiguous. Preferably, these modifications result in a "silent" change, in that the change produces a *HKNG1* gene product with the same activity as the
HKNG1 gene product depicted in FIG. 1A-1B, 2A-2B, 7-13, or
17.

Amino acid substitutions may be made on the basis of
30 similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues

involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged
5 (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

Alternatively, where alteration of function is desired, addition(s), deletion(s) or non-conservative alterations can produce altered, including reduced-activity, *HKNG1* gene
10 products. Such alterations can, for example, alter one or more of the biological functions of the *HKNG1* gene product. Further, such alterations can be selected so as to generate *HKNG1* gene products that are better suited for expression, scale up, etc. in the host cells chosen. For example,
15 cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges.

As another example, altered *HKNG1* gene products can be engineered that correspond to variants of the *HKNG1* gene product associated with *HKNG1*-mediated neuropsychiatric disorders such as BAD. Such altered *HKNG1* gene products
20 include, but are not limited to, *HKNG1* proteins or peptides comprising substitution of a lysine residue for the wild-type glutamic acid residue at *HKNG1* amino acid position 202 in FIG. 1A-1B (SEQ ID NO:2) or amino acid position 184 (SEQ ID NO:4) in FIG. 2A-2B.

HKNG1 protein fragments and/or *HKNG1* peptides comprise at least as many contiguous amino acid residues as necessary to represent an epitope fragment (that is to be recognized by an antibody directed to the *HKNG1* protein). For example, such protein fragments or peptides comprise at least about 8 contiguous *HKNG1* amino acid residues from a full length *HKNG1*
25 protein. In alternate embodiments, the *HKNG1* protein fragments and peptides of the invention can comprise about
30

10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450 or more contiguous amino acid residues of a HKNG1 protein.

Peptides and/or proteins corresponding to one or more domains of the *HKNG1* protein as well as fusion proteins in
5 which a *HKNG1* protein, or a portion of a *HKNG1* protein such as a truncated *HKNG1* protein or peptide or a *HKNG1* protein domain, is fused to an unrelated protein are also within the scope of this invention. Such proteins and peptides can be designed on the basis of the *HKNG1* nucleotide sequence
10 disclosed in Section 5.1, above, and/or on the basis of the *HKNG1* amino acid sequence disclosed in the Section. Fusion proteins include, but are not limited to, IgFc fusions which stabilize the *HKNG1* protein or peptide and prolong half life *in vivo*; or fusions to any amino acid sequence that allows
15 the fusion protein to be anchored to the cell membrane; or fusions to an enzyme, fluorescent protein, luminescent protein, or a flag epitope protein or peptide which provides a marker function.

The *HKNG1* protein, the *HKNG1* protein sequences described above can include a domain which comprises a signal sequence
20 that targets the *HKNG1* gene product for secretion. As used herein, a signal sequence includes a peptide of at least about 15 or 20 amino acid residues in length which occurs at the N-terminus of secretory and membrane-bound proteins and which contains at least about 70% hydrophobic amino acid residues such as alanine, leucine, isoleucine, phenylalanine,
25 proline, tyrosine, tryptophan, or valine. In a preferred embodiment, a signal sequence contains at least about 10 to 40 amino acid residues, preferably about 19-34 amino acid residues, and has at least about 60-80%, more preferably 65-75%, and more preferably at least about 70% hydrophobic residues. A signal sequence serves to direct a protein
30 containing such a sequence to a lipid bilayer.

In one embodiment, a *HNKNG1* protein contains a signal sequence at about amino acids 1 to 49 of SEQ ID NO:2. In another embodiment, a *HKNG1* protein contains a signal sequence at about amino acids 30-49 of SEQ ID NO:2. In yet another embodiment, a *HKNG1* protein contains a signal sequence at about amino acid residues 1 to 31 of SEQ ID NO:4. In yet another embodiment, a *HKNG1* protein contains a signal sequence at about amino acids 12-31 of SEQ ID NO:4. The signal sequence is cleaved during processing of the mature protein.

A signal sequence of a polypeptide of the invention can be used to facilitate secretion and isolation of the secreted protein or other proteins of interest. Signal sequences are typically characterized by a core of hydrophobic amino acids which are generally cleaved from the mature protein during secretion in one or more cleavage events. Such signal peptides contain processing sites that allow cleavage of the signal sequence from the mature proteins as they pass through the secretory pathway. Thus, the invention pertains to the described *HKNG1* polypeptides having a signal sequence (that is, "immature" polypeptides), as well as to the *HKNG1* signal sequences themselves and to the *HNKNG1* polypeptides in the absence of a signal sequence (*i.e.*, the "mature" *HKNG1* cleavage products). It is to be understood that *HKNG1* polypeptides of the invention can further comprise polypeptides comprising any signal sequence having characteristics as described above and a mature *HKNG1* polypeptide sequence.

In one embodiment, a nucleic acid sequence encoding a signal sequence of the invention can be operably linked in an expression vector to a protein of interest, such as a protein which is ordinarily not secreted or is otherwise difficult to isolate. The signal sequence directs secretion of the protein, such as from a eukaryotic host into which the expression vector is transformed, and the signal sequence is

subsequently or concurrently cleaved. The protein can then be readily purified from the extracellular medium by art recognized methods. Alternatively, the signal sequence can be linked to the protein of interest using a sequence which facilitates purification, such as with a GST domain.

5 The HKNG1 protein sequences described above can also include one or more domains which comprise a clusterin domain, *i.e.*, domains which are identical to or substantially homologous to (*i.e.*, 65%, 75%, 80%, 85%, 90%, 95% or more identical to) the domain corresponding to amino acid residues 134 to 160 or amino acid residues 334 to 362 of SEQ ID NO:2, 10 or to the domain corresponding to amino acid residues 105-131 or amino acid residues 305-333 of SEQ ID No:39, or to the domain corresponding to amino acid residues 105-131 or amino acid residues 304-332 of SEQ ID NO:49. Preferably, such domains comprise cysteine amino acid residues at positions corresponding to conserved cysteine residues of the clusterin 15 domains of SEQ ID NOS: 2, 39 or 49.

In particular, HKNG1 protein sequences described above can also include one or more domains which comprise a conserved cysteine domain. Such a domain corresponds, for example, to the domain of cysteines corresponding to Cys134, Cys145, Cys148, Cys158 and Cys160; or to Cys 334, Cys344, 20 Cys351, Cys354, and Cys362 of SEQ ID NO:2. In an alternative embodiment, a conserved cystein domain corresponds to one or more of the domains of SEQ ID NO:39 which comprises Cys105, Cys116, Cys119, Cys124, and Cys131; or Cys305, Cys315, Cys322, Cys325, and Cys333. In yet another alternative 25 embodiment, a conserved cysteine domain corresponds to one or more of the domains of SEQ ID NO:49 which comprises Cys105, Cys116, Cys119, Cys124, and Cys131; or Cys314, Cys321, Cys324, and Cys332.

Finally, the HKNG1 proteins of the invention also include HKNG1 protein sequences wherein domains encoded by one or more exons of the cDNA sequence, or fragments thereof, 30 have been deleted. In one particularly preferred embodiment,

the HKNG1 proteins of the invention are proteins in which the domain(s) corresponding the those domains encoded by exon 7 of SEQ ID NO:7, or fragments thereof, have been deleted.

The HKNG1 polypeptides of the invention can further comprise posttranslational modifications, including, but not limited to glycosylations, acetylations, and myrisalations.

The HKNG1 gene products, peptide fragments thereof and fusion proteins thereof, may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the HKNG1 gene products, polypeptides, peptides, fusion peptide and fusion polypeptides of the invention by expressing nucleic acid containing HKNG1 gene sequences are described herein. Methods that are well known to those skilled in the art can be used to construct expression vectors containing HKNG1 gene product coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. See, for example, the techniques described in Sambrook, et al., 1989, *supra*, and Ausubel, et al., 1989, *supra*. Alternatively, RNA capable of encoding HKNG1 gene product sequences may be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis", 1984, Gait, ed., IRL Press, Oxford.

A variety of host-expression vector systems may be utilized to express the HKNG1 gene product coding sequences of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells that may, when transformed or transfected with the appropriate nucleotide coding sequences, exhibit the HKNG1 gene product of the invention *in situ*. These include but are not limited to microorganisms such as bacteria (e.g., *E.*

coli, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing *HKNG1* gene product coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast
5 expression vectors containing the *HKNG1* gene product coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the *HKNG1* gene product coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g.,
10 cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing *HKNG1* gene product coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian
15 cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the *HKNG1* gene product being expressed. For example,
20 when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of *HKNG1* gene product or for raising antibodies to *HKNG1* gene product, for example, vectors that direct the expression of high levels of fusion protein products that are readily purified
25 may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which the *HKNG1* gene product coding sequence may be ligated individually into the vector in frame with the *lacZ* coding region so that a fusion protein is
30 produced; pIN vectors (Inouye and Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke and Schuster, 1989, J. Biol.

Chem. 264:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by
5 elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, *Autographa californica*, nuclear polyhidrosis virus (AcNPV) is used as a vector to express
10 foreign genes. The virus grows in *Spodoptera frugiperda* cells. The *HKNG1* gene product coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).
15 Successful insertion of *HKNG1* gene product coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect
20 *Spodoptera frugiperda* cells in which the inserted gene is expressed. (e.g., see Smith, et al., 1983, J. Virol. 46:584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the *HKNG1* gene
25 product coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-
30 essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable

of expressing *HKNG1* gene product in infected hosts. (e.g., See Logan and Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted *HKNG1* gene product coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire *HKNG1* gene, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of the *HKNG1* gene coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner, et al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such

mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, and WI38.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the *HKNG1* gene product may be
5 engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and
10 a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and
15 grow to form foci that in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the *HKNG1* gene product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the *HKNG1* gene product.

20 A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine
25 phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk^- , $hgprt^-$ or $aprt^-$ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: *dhfr*, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc.
30 Natl. Acad. Sci. USA 78:1527); *gpt*, which confers resistance to mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl.

Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hygro, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147).

5 Alternatively, the expression characteristics of an endogenous *HKNG1* gene within a cell line or microorganism may be modified by inserting a heterologous DNA regulatory element into the genome of a stable cell line or cloned microorganism such that the inserted regulatory element is operatively linked with the endogenous *HKNG1* gene. For
10 example, an endogenous *HKNG1* gene which is normally "transcriptionally silent", i.e., an *HKNG1* gene which is normally not expressed, or is expressed only at very low levels in a cell line or microorganism, may be activated by inserting a regulatory element which is capable of promoting the expression of a normally expressed gene product in that
15 cell line or microorganism. Alternatively, a transcriptionally silent, endogenous *HKNG1* gene may be activated by insertion of a promiscuous regulatory element that works across cell types.

A heterologous regulatory element may be inserted into a
20 stable cell line or cloned microorganism, such that it is operatively linked with an endogenous *HKNG1* gene, using techniques, such as targeted homologous recombination, which are well known to those of skill in the art, and described e.g., in Chappel, U.S. Patent No. 5,272,071; PCT publication No. WO 91/06667, published May 16, 1991.

25 Alternatively, any fusion protein may be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht, et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-
30 8976). In this system, the gene of interest is subcloned

into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺-nitriloacetic acid-agarose columns and
5 histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

The *HKNG1* gene products can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, sheep, cows, and non-human primates, e.g.,
10 baboons, monkeys, and chimpanzees may be used to generate *HKNG1* transgenic animals. The term "transgenic," as used herein, refers to animals expressing *HKNG1* gene sequences from a different species (e.g., mice expressing huma *HKNG1* gene sequences), as well as animals that have been
15 genetically engineered to overexpress endogenous (i.e., same species) *HKNG1* sequences or animals that have been genetically engineered to no longer express endogenous *HKNG1* gene sequences (i.e., "knock-out" animals), and their progeny.

20 Any technique known in the art may be used to introduce a *HKNG1* gene transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten, et al., 1985,
25 Proc. Natl. Acad. Sci., USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson, et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol. Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723) (For a review of such techniques,
30 see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115, 171-229)

Any technique known in the art may be used to produce transgenic animal clones containing a *HKNG1* transgene, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal or adult cells induced to quiescence (Campbell, et al., 1996, Nature 380:64-66; Wilmut, et al., Nature 385:810-813).

The present invention provides for transgenic animals that carry a *HKNG1* transgene in all their cells, as well as animals that carry the transgene in some, but not all their cells, i.e., mosaic animals. The transgene may be integrated as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko, et al., 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the *HKNG1* transgene be integrated into the chromosomal site of the endogenous *HKNG1* gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous *HKNG1* gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous *HKNG1* gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous *HKNG1* gene in only that cell type, by following, for example, the teaching of Gu, et al. (Gu, et al., 1994, Science 265, 103-106). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant *HKNG1* gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques that include but are not limited to Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and RT-PCR (reverse transcriptase PCR). Samples of *HKNG1* gene-expressing tissue, may also be evaluated immunocytochemically using antibodies specific for the *HKNG1* transgene product.

HKNG1 proteins can be used, e.g., to treat CNS-related disorders, e.g., neuropsychiatric disorders. Such *HKNG1* gene products include but are not limited to soluble derivatives such as peptides or polypeptides corresponding to one or more domains of the *HKNG1* gene product, particularly *HKNG1* gene products, that are modified such that they are deleted for one or more hydrophobic domains. Alternatively, antibodies to the *HKNG1* protein or anti-idiotypic antibodies that mimic the *HKNG1* gene product (including Fab fragments), antagonists or agonists can be used to treat neuropsychiatric disorders involving *HKNG1*. In yet another approach, nucleotide constructs encoding such *HKNG1* gene products can be used to genetically engineer host cells to express such *HKNG1* gene products in vivo; these genetically engineered cells can function as "bioreactors" in the body delivering a continuous supply of *HKNG1* gene product, *HKNG1* peptides, soluble *HKNG1* polypeptides.

5.3. ANTIBODIES TO *HKNG1* GENE PRODUCTS

Described herein are methods for the production of antibodies capable of specifically recognizing one or more

HKNG1 gene product epitopes or epitopes of conserved variants or peptide fragments of the *HKNG1* gene products. Further, antibodies that specifically recognize mutant forms of *HKNG1*, are encompassed by the invention. The terms "specifically
5 bind" and "specifically recognize" refer to antibodies that bind to *HKNG1* gene product epitopes at a higher affinity than they bind to non-*HKNG1* (e.g., random) epitopes.

Such antibodies may include, but are not limited to, polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies,
10 Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above, including the polyclonal and monoclonal antibodies described in Section 12 below. Such antibodies may be used, for example, in the
15 detection of a *HKNG1* gene product in an biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal levels of *HKNG1* gene products, and/or for the presence of abnormal forms of such gene products. Such
antibodies may also be utilized in conjunction with, for
20 example, compound screening schemes, as described, below, in Section 5.8, for the evaluation of the effect of test compounds on *HKNG1* gene product levels and/or activity. Additionally, such antibodies can be used in conjunction with the gene therapy techniques described, below, in Section
5.9.2 to, for example, evaluate the normal and/or engineered
25 *HKNG1*-expressing cells prior to their introduction into the patient.

Anti-*HKNG1* gene product antibodies may additionally be used in methods for inhibiting abnormal *HKNG1* gene product activity. Thus, such antibodies may, therefore, be utilized
30 as part of treatment methods for a *HKNG1*-mediated neuropsychiatric disorder, such as BAD or schizophrenia.

For the production of antibodies against a *HKNG1* gene product, various host animals may be immunized by injection with a *HKNG1* gene product, or a portion thereof. Such host animals may include, but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.

Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen, such as a *HKNG1* gene product, or an antigenic functional derivative thereof. For the production of polyclonal antibodies, host animals such as those described above, may be immunized by injection with *HKNG1* gene product supplemented with adjuvants as also described above.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique that provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, *Nature* 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, *Immunology Today* 4:72; Cole et al., 1983, *Proc. Natl. Acad. Sci. USA* 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, *Monoclonal Antibodies And Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated in

vitro or *in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison, et al., 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger, et al., 1984, Nature 312:604-608; Takeda, et al., 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. (See, e.g., Cabilly et al., U.S. Patent No. 4,816,567; and Boss et al., U.S. Patent No. 4,816,397, which are incorporated herein by reference in their entirety.)

In addition, techniques have been developed for the production of humanized antibodies. (See, e.g., Queen, U.S. Patent No. 5,585,089, which is incorporated herein by reference in its entirety.) An immunoglobulin light or heavy chain variable region consists of a "framework" region interrupted by three hypervariable regions, referred to as complementarily determining regions (CDRs). The extent of the framework region and CDRs have been precisely defined (see, "Sequences of Proteins of Immunological Interest", Kabat, E. et al., U.S. Department of Health and Human Services (1983)). Briefly, humanized antibodies are antibody molecules from non-human species having one or more CDRs from the non-human species and a framework region from a human immunoglobulin molecule.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242:423-426; Huston, et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward, et al., 1989, Nature 334:544-546) can be adapted to produce single chain

antibodies against *HKNG1* gene products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

5 Antibody fragments that recognize specific epitopes may be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragments, which can be produced by pepsin digestion of the antibody molecule and the Fab fragments, which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be
10 constructed (Huse, et al., 1989, Science 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

5.4. USES OF *HKNG1* GENE SEQUENCES GENE PRODUCTS, AND ANTIBODIES

15 Described herein are various applications of *HKNG1* gene sequences, *HKNG1* gene products, including peptide fragments and fusion proteins thereof, and of antibodies directed against *HKNG1* gene products and peptide fragments thereof. Such applications include, for example, mapping of chromosome
20 18p, prognostic and diagnostic evaluation of *HKNG1*-mediated disorders, including CNS-related disorders, e.g., neuropsychiatric disorders, such as BAD or schizophrenia, modulation of *HKNG1*-related processes, and the identification of subjects with a predisposition to such disorders, as
25 described, below, in Section 5.5.

Additionally, such applications include methods for the treatment of a *HKNG1*-mediated disorders, such as BAD or schizophrenia, as described, below, in Section 5.9, and for the identification of compounds that modulate the expression of the *HKNG1* gene and/or the synthesis or activity of the
30 *HKNG1* gene product, as described below, in Section 5.8. Such compounds can include, for example, other cellular products

that are involved in such processes as mood regulation and in *HKNG1*-mediated disorders, e.g., neuropsychiatric disorders such as BAD or schizophrenia. These compounds can be used, for example, in the amelioration of *HKNG1*-mediated disorders and for the modulation of *HKNG1*-mediated processes.

Uses of the *HKNG1* gene sequences, *HKNG1* gene products, including peptide fragments and fusion proteins thereof, and of antibodies directed against *HKNG1* gene products and/or peptide fragments thereof also include prognostic and diagnostic evaluation of a *HKNG1*-mediated myopia disorder such as early-onset autosomal dominant myopia, methods for the treatment of a *HKNG1*-mediated myopia disorder, and for the identification of compound that modulate the expression of the *HKNG1* gene and/or the synthesis or activity of the *HKNG1* gene product and could therefore be used in the amelioration of a *HKNG1*-mediated myopia such as early-onset autosomal dominant myopia. Indeed, such methods are substantially identical to the methods described, below, in Sections 5.5, 5.8, and 5.9 for the diagnosis and treatment of *HKNG1*-mediated disorders.

20 **5.5. DIAGNOSIS OF *HKNG1*-MEDIATED DISORDERS**

A variety of methods can be employed for the diagnostic and prognostic evaluation of *HKNG1*-mediated disorders, e.g., neuropsychiatric disorders and for the identification of subjects having a predisposition to such disorders.

Such methods may, for example, utilize reagents such as the *HKNG1* gene nucleotide sequences described in Sections 5.1, and antibodies directed against *HKNG1* gene products, including peptide fragments thereof, as described, above, in Section 5.3. Specifically, such reagents may be used, for example, for:

(1) the detection of the presence of *HKNG1* gene mutations, or the detection of either over- or under-expression of *HKNG1* gene relative to wild-type *HKNG1* levels of expression;

5 (2) the detection of over- or under-abundance of *HKNG1* gene product relative to wild-type abundance of *HKNG1* gene product; and

(3) the detection of an aberrant level of *HKNG1* gene product activity relative to wild-type *HKNG1* gene product activity levels.

10 *HKNG1* gene nucleotide sequences can, for example, be used to diagnose a *HKNG1*-mediated neuropsychiatric disorder using, for example, the techniques for *HKNG1* mutation/polymorphism detection described above in Section 5.1, and in Section 5.6 below.

15 Mutations at a number of different genetic loci may lead to phenotypes related to neuropsychiatric disorders. Ideally, the treatment of patients suffering from such neuropsychiatric disorder will be designed to target the particular genetic loci containing the mutation mediating the disorder. Genetic polymorphisms have been linked to
20 differences in drug effectiveness. Thus, identification of alterations in the *HKNG1* gene, protein or gene flanking regions, can be utilized in pharmacogenetic methods to optimize therapeutic drug treatments.

In one embodiment of the present invention, therefore, alterations, *i.e.*, polymorphisms, in the *HKNG1* gene or
25 protein encoded by genes comprising such polymorphisms, are associated with a drug or drugs' efficacy, tolerance, or toxicity, and may be used in pharmacogenomic methods to optimize therapeutic drug treatments, including therapeutic drug treatments for one of the disorders described herein, *e.g.*, *HKNG1*-mediated disorders such as schizophrenia and BAD.
30 Such polymorphisms can be used, for example, to refine the

design of drugs by decreasing the incidence of adverse events in drug tolerance studies, e.g., by identifying patient subpopulations of individuals who respond or do not respond to a particular drug therapy in efficacy studies, wherein the subpopulations have a *HKNG1* polymorphism associated with drug
5 responsiveness or unresponsiveness. The pharmacogenomic methods of the present invention can also provide tools to identify new drug targets for designing drugs and to optimize the use of already existing drugs, e.g., to increase the response rate to a drug and/or to identify and exclude non-
10 responders from certain drug treatments (e.g., individuals having a particular *HKNG1* polymorphism associated with unresponsiveness or inferior responsiveness to the drug treatment) or to decrease the undersirable side effects of certain drug treatments and/or to identify and exclude individuals with marked susceptibility to such side effects
15 (e.g., individuals having a particular *HKNG1* polymorphism associated with an undersirable side effect to the drug treatment).

In an embodiment of the present invention, polymorphisms in the *HKNG1* gene sequence or flanking this sequence, or
20 variations in *HKNG1* gene expression, or activity, e.g., variations due to altered methylation, differential splicing, or post-translational modification of the *HKNG1* gene product, may be utilized to identify an individual having a disease or condition resulting from a *HKNG1*-mediated disorder and thus define the most effective and safest drug treatment. Assays
25 such as those described herein may be used to identify such polymorphisms or variations in *HKNG1* gene expression or activity. Once a polymorphism in the *HKNG1* gene or in a flanking sequence in linkage disequilibrium with a disorder-causing allele, or a variation in *HKNG1* gene expression has
30 been identified in an individual, an appropriate drug treatment can be prescribed to the individual.

For the detection of *HKNG1* gene mutations or polymorphisms, any nucleated cell can be used as a starting source for genomic nucleic acid. For the detection of *HKNG1* gene expression or *HKNG1* gene products, any cell type or
5 tissue in which the *HKNG1* gene is expressed may be utilized.

Nucleic acid-based detection techniques are described, below, in Section 5.6. Peptide detection techniques are described, below, in Section 5.7.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits. The
10 invention therefore also encompasses kits for detecting the presence of a polypeptide or nucleic acid of the invention in a biological sample (*i.e.*, a test sample). Such kits can be used, *e.g.*, to determine if a subject is suffering from or is at increased risk of developing a disorder associated with a
15 disorder-causing allele, or aberrant expression or activity of a polypeptide of the invention (*e.g.*, a CNS disorder, including a neuropsychiatric disorder such as BAD or schizophrenia). For example, the kit can comprise a labeled compound or agent capable of detecting the polypeptide or mRNA or DNA or *HKNG1* gene sequences, *e.g.*, encoding the
20 polypeptide in a biological sample. The kit can further comprise a means for determining the amount of the polypeptide or mRNA in the sample (*e.g.*, an antibody which binds the polypeptide or an oligonucleotide probe which binds to DNA or mRNA encoding the polypeptide). Kits can also include instructions for observing that the tested subject is
25 suffering from or is at risk of developing a disorder associated with aberrant expression of the polypeptide if the amount of the polypeptide or mRNA encoding the polypeptide is above or below a normal level, or if the DNA correlates with presence of a *HKNG1* allele that causes a disorder.

For antibody-based kits, the kit can comprise, for
30 example: (1) a first antibody (*e.g.*, attached to a solid

support) which binds to a polypeptide of the invention; and, optionally, (2) a second, different antibody which binds to either the polypeptide or to the first antibody and is conjugated to a detectable agent.

For oligonucleotide-based kits, the kit can comprise, 5 for example: (1) an oligonucleotide (e.g., a detectably labeled oligonucleotide) which hybridizes to a nucleic acid sequence encoding a polypeptide of the invention, or (2) a pair of primers, such as the primers recited in Table 1, useful for amplifying a nucleic acid molecule encoding a polypeptide of the invention.

10 The kit can also comprise, for example, one or more buffering agents, preservatives, or protein stabilizing agents. The kit can also comprise components necessary for detecting the detectable agent (e.g., an enzyme or a substrate). The kit can also contain a control sample or a 15 series of control samples which can be assayed and compared to the test sample. Each component of the kit is usually enclosed within an individual container and all of the various containers are within a single package along with instructions for observing whether the tested subject is suffering from or is at risk of developing a disorder 20 associated with polymorphisms that correlate with alleles that cause *HKNG1*-related disorders, and/or aberrant levels of *HKNG1* mRNA, polypeptides or activity.

5.6. DETECTION OF *HKNG1* NUCLEIC ACID MOLECULES

25 A variety of methods can be employed to screen for the presence of *HKNG1* gene-specific mutations or polymorphisms (including polymorphisms flanking *HKNG1* gene) and to detect and/or assay levels of *HKNG1* nucleic acid sequences.

Mutations or polymorphisms within or flanking the *HKNG1* gene can be detected by utilizing a number of techniques. 30 Nucleic acid from any nucleated cell can be used as the starting point for such assay techniques, and may be isolated

according to standard nucleic acid preparation procedures that are well known to those of skill in the art.

HKNG1 nucleic acid sequences may be used in hybridization or amplification assays of biological samples to detect abnormalities involving *HKNG1* gene structure,
5 including point mutations, insertions, deletions, inversions, translocations and chromosomal rearrangements. Such assays may include, but are not limited to, Southern analyses, single-stranded conformational polymorphism analyses (SSCP), and PCR analyses.

10 Diagnostic methods for the detection of *HKNG1* gene-specific mutations or polymorphisms can involve for example, contacting and incubating nucleic acids obtained from a sample, e.g., derived from a patient sample or other appropriate cellular source with one or more labeled nucleic acid reagents including recombinant DNA molecules, cloned
15 genes or degenerate variants thereof, such as described in Section 5.1, above, under conditions favorable for the specific annealing of these reagents to their complementary sequences within or flanking the *HKNG1* gene. The diagnostic methods of the present invention further encompass contacting and incubating nucleic acids for the detection of single
20 nucleotide mutations or polymorphisms of the *HKNG1* gene. Preferably, these nucleic acid reagent sequences within the *HKNG1* gene, or chromosome 18p nucleotide sequences flanking the *HKNG1* gene are 15 to 30 nucleotides in length.

After incubation, all non-annealed nucleic acids are
25 removed from the nucleic acid:*HKNG1* molecule hybrid. The presence of nucleic acids that have hybridized, if any such molecules exist, is then detected. Using such a detection scheme, the nucleic acid from the cell type or tissue of interest can be immobilized, for example, to a solid support such as a membrane, or a plastic surface such as that on a
30 microtiter plate or polystyrene beads. In this case, after incubation, non-annealed, labeled nucleic acid reagents of

the type described in Section 5.1 are easily removed. Detection of the remaining, annealed, labeled *HKNG1* nucleic acid reagents is accomplished using standard techniques well-known to those in the art. The *HKNG1* gene sequences to which
5 the nucleic acid reagents have annealed can be compared to the annealing pattern expected from a normal *HKNG1* gene sequence in order to determine whether a *HKNG1* gene mutation is present.

In a preferred embodiment, *HKNG1* mutations or polymorphisms can be detected by using a microassay of *HKNG1*
10 nucleic acid sequences immobilized to a substrate or "gene chip" (see, e.g. Cronin, et al., 1996, Human Mutation 7:244-255).

Alternative diagnostic methods for the detection of *HKNG1* gene-specific nucleic acid molecules (or *HKNG1* flanking
15 sequences), in patient samples or other appropriate cell sources, may involve their amplification, e.g., by PCR (the experimental embodiment set forth in Mullis, 1987, U.S. Patent No. 4,683,202), followed by the analysis of the amplified molecules using techniques well known to those of skill in the art, such as, for example, those listed above.
20 The resulting amplified sequences can be compared to those that would be expected if the nucleic acid being amplified contained only normal copies of the *HKNG1* gene in order to determine whether a *HKNG1* gene mutation or polymorphism in linkage disequilibrium with a disease-causing *HKNG1* allele
25 exists.

Among those *HKNG1* nucleic acid sequences which are preferred for such amplification-related diagnostic screening analyses are oligonucleotide primers which amplify *HKNG1* exon
sequences. The sequences of such oligonucleotide primers are, therefore, preferably derived from *HKNG1* intron
30 sequences so that the entire exon, or coding region, can be analyzed as discussed below. Primer pairs useful for

amplification of *HKNG1* exons are preferably derived from adjacent introns. Appropriate primer pairs can be chosen such that each of the eleven *HKNG1* exons are amplified.

Primers for the amplification of *HKNG1* exons can be routinely
5 designed by one of ordinary skill in the art by utilizing the
exon and intron sequences of *HKNG1* shown in Figure 3A-3R.

As an example, and not by way of limitation, Table 1,
below, lists primers and primer pairs which can be utilized
for the amplification of each of the human *HKGN1* exons one
through eleven. In this table, a primer pair is listed for
10 each exon which consists of a forward primer derived from
intron sequence upstream of the exon to be amplified, and a
reverse primer derived from intron sequence downstream of the
exon to be amplified. For exons greater than about 300 base
pairs in length, i.e., exons 4 and 7, two primer pairs are
listed (marked 4a, 4b, 7a and 7b). Each of the primer pairs
15 can be utilized, therefore, as part of a standard PCR
reaction to amplify an individual *HKNG1* exon (or portion
thereof). Primer sequences are depicted in a 5' to 3'
orientation.

20

25

30

TABLE 1

	Primer Sequence		
1	cgggggttggtttccacc	(SEQ ID NO:8)	forward
	gcgaggagagaaatctggg	(SEQ ID NO:9)	reverse
5			
2	tgctcactactttgcagtgttc	(SEQ ID NO:10)	forward
	tgagatcgtgtcactgcattct	(SEQ ID NO:11)	reverse
3	gtaaactctcaaaatggtgggtaataag	(SEQ ID NO:12)	forward
	ctaactcttcttctatcattactc	(SEQ ID NO:13)	reverse
10			
4A	tgtttattgtgtgtctgctgtg	(SEQ ID NO:14)	forward
	ggacaaccaacatgcaaacag	(SEQ ID NO:15)	reverse
4B	cccaggtgttttcaattgatgc	(SEQ ID NO:16)	forward
	agcagttttgtccttccaagtg	(SEQ ID NO:17)	reverse
15			
5	gtgttttgtaatctgatcagatctc	(SEQ ID NO:18)	forward
	gcagtatttctgggtccagatc	(SEQ ID NO:19)	reverse
6	ggtgcacatagatcatgaaatgg	(SEQ ID NO:20)	forward
	taagctgaaataggtgccttaag	(SEQ ID NO:21)	reverse
20			
7A	tttattccatttctgtcccctac	(SEQ ID NO:22)	forward
	aaggctcagttaggtctgtatc	(SEQ ID NO:23)	reverse
7B	caggagttttaacgtcttcagac	(SEQ ID NO:24)	forward
	gactcagaaatgtctaccatttc	(SEQ ID NO:25)	reverse
25			
8	tgtctccacttcttcaaagtgc	(SEQ ID NO:26)	forward
	caaaatgtacctgagaacttaaag	(SEQ ID NO:27)	reverse
30			

	Primer Sequence		
9	cacctccaagtttcatggac	(SEQ ID NO:28)	forward
	caaggatgacacgtgtcatttc	(SEQ ID NO:29)	reverse
5	10	gaatgtgtattgggatttagtaaac	(SEQ ID NO:30) forward
		ttgagaattaactattcctgtcaac	(SEQ ID NO:31) reverse
	11	ccatcctggacttttactcc	(SEQ ID NO:32) forward
		ctttcctgcaactgtgtttattg	(SEQ ID NO:33) reverse

10

Each primer pair above can be used to generate an amplified sequence of about 300 base pairs. This is especially desirable in instances in which sequence analysis is performed using SSCP gel electrophoretic procedures, in that such procedures work optimally using sequences of about 15 300 base pairs or less.

Additional *HKNG1* nucleic acid sequences which are preferred for such amplification-related analyses are those which will detect the presence of an *HKNG1* polymorphism which differs from the *HKNG1* sequence depicted in FIG. 3A-3R. Such polymorphisms include ones which represent mutations 20 associated with an *HKNG1*-mediated neuropsychiatric disorder, such as BAD or schizophrenia. For example, a single base mutation identified in the Example presented in Section 8, below, results in a mutant *HKNG1* gene product comprising substitution of a lysine residue for the wild-type glutamic 25 acid residue at amino acid position 202 of the *HKNG1* amino acid sequence shown in FIG. 1A-1B (SEQ ID NO:2) or amino acid position 184 of the *HKNG1* amino acid sequence shown in FIG. 2A-2B (SEQ ID NO:4). Such polymorphisms also include ones that correlate with the presence of a *HKNG1*-mediated neuropsychiatric disorder, e.g., polymorphisms that are in 30 linkage disequilibrium with disorder-causing *HKNG1* alleles.

Amplification techniques are well known to those of skill in the art and can routinely be utilized in connection with primers such as those listed in Table 1 above. In general, hybridization conditions can be as follows. In general, for probes between 14 and 70 nucleotides in length the melting temperature T_m is calculated using the formula: $T_m(^{\circ}\text{C})=81.5+16.6(\log[\text{monovalent cations}])+0.41(\% \text{G+C})-(500/N)$ where N is the length of the probe. If the hybridization is carried out in a solution containing formamide, the melting temperature is calculated using the equation $T_m(^{\circ}\text{C})=81.5+16.6(\log[\text{monovalent cations}])+0.41(\% \text{G+C})-(0.61\% \text{ formamide})-(500/N)$ where N is the length of the probe.

Additionally, well-known genotyping techniques can be performed to identify individuals carrying *HKNG1* gene mutations. Such techniques include, for example, the use of restriction fragment length polymorphisms (RFLPs), which involve sequence variations in one of the recognition sites for the specific restriction enzyme used.

Further, improved methods for analyzing DNA polymorphisms, which can be utilized for the identification of *HKNG1* gene-specific mutations, have been described that capitalize on the presence of variable numbers of short, tandemly repeated DNA sequences between the restriction enzyme sites. For example, Weber (U.S. Pat. No. 5,075,217) describes a DNA marker based on length polymorphisms in blocks of $(\text{dC-dA})_n-(\text{dG-dT})_n$ short tandem repeats. The average separation of $(\text{dC-dA})_n-(\text{dG-dT})_n$ blocks is estimated to be 30,000-60,000 bp. Markers that are so closely spaced exhibit a high frequency co-inheritance, and are extremely useful in the identification of genetic mutations, such as, for example, mutations within the *HKNG1* gene, and the diagnosis of diseases and disorders related to *HKNG1* mutations.

Also, Caskey et al. (U.S. Pat.No. 5,364,759) describe a DNA profiling assay for detecting short tri and tetra

nucleotide repeat sequences. The process includes extracting the DNA of interest, such as the *HKNG1* gene, amplifying the extracted DNA, and labelling the repeat sequences to form a genotypic map of the individual's DNA.

5 Other methods well known in the art may be used to identify single nucleotide polymorphisms (SNPs), including biallelic SNPs or biallelic markers which have two alleles, both of which are present at a fairly high frequency in a population. Conventional techniques for detecting SNPs include, e.g., conventional dot blot analysis, single
10 stranded conformational polymorphism (SSCP) analysis (see, e.g., Orita et al., 1989, *Proc. Natl. Acad. Sci. USA* 86:2766-2770), denaturing gradient gel electrophoresis (DGGE), heteroduplex analysis, mismatch cleavage detection, and other routine techniques well known in the art (see, e.g., Sheffield et al., 1989, *Proc. Natl. Acad. Sci.* 86:5855-5892;
15 Grompe, 1993, *Nature Genetics* 5:111-117). Alternative, preferred methods of detecting and mapping SNPs involve microsequencing techniques wherein an SNP site in a target DNA is detected by a single nucleotide primer extension reaction (see, e.g., Goelet et al., PCT Publication No.
20 WO92/15712; Mundy, U.S. Patent No. 4,656,127; Vary and Diamond, U.S. Patent No. 4,851,331; Cohen et al., PCT Publication No. WO91/02087; Chee et al., PCT Publication No. WO95/11995; Landegren et al., 1988, *Science* 241:1077-1080; Nicerson et al., 1990, *Proc. Natl. Acad. Sci. U.S.A.* 87:8923-8927; Pastinen et al., 1997, *Genome Res.* 7:606-614; Pastinen
25 et al., 1996, *Clin. Chem.* 42:1391-1397; Jalanko et al., 1992, *Clin. Chem.* 38:39-43; Shumaker et al., 1996, *Hum. Mutation* 7:346-354; Caskey et al., PCT Publication No. WO 95/00669).

The level of *HKNG1* gene expression can also be assayed. For example, RNA from a cell type or tissue known, or
30 suspected, to express the *HKNG1* gene, such as brain, may be

isolated and tested utilizing hybridization or PCR techniques such as are described, above. The isolated cells can be derived from cell culture or from a patient. The analysis of cells taken from culture may be a necessary step in the assessment of cells to be used as part of a cell-based gene therapy technique or, alternatively, to test the effect of compounds on the expression of the *HKNG1* gene. Such analyses may reveal both quantitative and qualitative aspects of the expression pattern of the *HKNG1* gene, including activation or inactivation of *HKNG1* gene expression.

10 In one embodiment of such a detection scheme, a cDNA molecule is synthesized from an RNA molecule of interest (e.g., by reverse transcription of the RNA molecule into cDNA). A sequence within the cDNA is then used as the template for a nucleic acid amplification reaction, such as a PCR amplification reaction, or the like. The nucleic acid reagents used as synthesis initiation reagents (e.g., primers) in the reverse transcription and nucleic acid amplification steps of this method are chosen from among the *HKNG1* gene nucleic acid reagents described in Section 5.1. The preferred lengths of such nucleic acid reagents are at least 9-30 nucleotides. For detection of the amplified product, the nucleic acid amplification may be performed using radioactively or non-radioactively labeled nucleotides. Alternatively, enough amplified product may be made such that the product may be visualized by standard ethidium bromide staining or by utilizing any other suitable nucleic acid staining method.

25 Additionally, it is possible to perform such *HKNG1* gene expression assays "in situ", i.e., directly upon tissue sections (fixed and/or frozen) of patient tissue obtained from biopsies or resections, such that no nucleic acid purification is necessary. Nucleic acid reagents such as those described in Section 5.1 may be used as probes and/or primers for such in situ procedures (see, for example, Nuovo,

G.J., 1992, "PCR In Situ Hybridization: Protocols And Applications", Raven Press, NY).

Alternatively, if a sufficient quantity of the appropriate cells can be obtained, standard Northern analysis can be performed to determine the level of mRNA expression of
5 the *HKNG1* gene.

5.7. DETECTION OF *HKNG1* GENE PRODUCTS

Antibodies directed against unimpaired or mutant *HKNG1* gene products or conserved variants or peptide fragments
10 thereof, which are discussed, above, in Section 5.3, may also be used as diagnostics and prognostics for a *HKNG1*-mediated disorder, e.g., a neuropsychiatric disorder such as BAD or schizophrenia. Such methods may be used to detect abnormalities in the level of *HKNG1* gene product synthesis or
15 expression, or abnormalities in the structure, temporal expression, and/or physical location of *HKNG1* gene product.

The antibodies and immunoassay methods described herein have, for example, important *in vitro* applications in assessing the efficacy of treatments for *HKNG1*-mediated disorders.

Antibodies, or fragments of antibodies, such as those
20 described below, may be used to screen potentially therapeutic compounds *in vitro* to determine their effects on *HKNG1* gene expression and *HKNG1* gene product production. The compounds that have beneficial effects on a *HKNG1*-mediated disorder, such as BAD or schizophrenia.

In vitro immunoassays may also be used, for example, to
25 assess the efficacy of cell-based gene therapy for a *HKNG1*-mediated disorder, e.g., a neuropsychiatric disorder, such as BAD schizophrenia. Antibodies directed against *HKNG1* gene products may be used *in vitro* to determine, for example, the level of *HKNG1* gene expression achieved in cells genetically
30 engineered to produce *HKNG1* gene product. In the case of

intracellular *HKNG1* gene products, such an assessment is done, preferably, using cell lysates or extracts. Such analysis will allow for a determination of the number of transformed cells necessary to achieve therapeutic efficacy *in vivo*, as well as optimization of the gene replacement
5 protocol.

The tissue or cell type to be analyzed will generally include those that are known, or suspected, to express the *HKNG1* gene. The protein isolation methods employed herein may, for example, be such as those described in Harlow and
10 Lane (1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). The isolated cells can be derived from cell culture or from a patient. The analysis of cells taken from culture may be a necessary step in the assessment of cells to be used as part of a cell-based gene therapy technique or, alternatively, to
15 test the effect of compounds on the expression of the *HKNG1* gene.

Preferred diagnostic methods for the detection of *HKNG1* gene products, conserved variants or peptide fragments thereof, may involve, for example, immunoassays wherein the
20 *HKNG1* gene products or conserved variants or peptide fragments are detected by their interaction with an anti-*HKNG1* gene product-specific antibody.

For example, antibodies, or fragments of antibodies, such as those described, above, in Section 5.3, may be used to quantitatively or qualitatively detect the presence of
25 *HKNG1* gene products or conserved variants or peptide fragments thereof. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody (see below, this Section) coupled with light microscopic, flow cytometric, or fluorimetric detection. Such techniques are especially preferred for *HKNG1* gene
30 products that are expressed on the cell surface.

The antibodies (or fragments thereof) useful in the present invention may, additionally, be employed histologically, as in immunofluorescence or immunoelectron microscopy, for *in situ* detection of *HKNG1* gene products, conserved variants or peptide fragments thereof. *In situ* 5 detection may be accomplished by removing a histological specimen from a patient, and applying thereto a labeled antibody that binds to an rTs polypeptide. The antibody (or fragment) is preferably applied by overlaying the labeled antibody (or fragment) onto a biological sample. Through the use of such a procedure, it is possible to determine not only 10 the presence of the *HKNG1* gene product, conserved variants or peptide fragments, but also its distribution in the examined tissue. Using the present invention, those of ordinary skill will readily recognize that any of a wide variety of histological methods (such as staining procedures) can be 15 modified in order to achieve *in situ* detection of a *HKNG1* gene product.

Immunoassays for *HKNG1* gene products, conserved variants, or peptide fragments thereof will typically comprise incubating a sample, such as a biological fluid, a tissue extract, freshly harvested cells, or lysates of cells 20 in the presence of a detectably labeled antibody capable of identifying *HKNG1* gene product, conserved variants or peptide fragments thereof, and detecting the bound antibody by any of a number of techniques well-known in the art.

The biological sample may be brought in contact with and immobilized onto a solid phase support or carrier, such as 25 nitrocellulose, that is capable of immobilizing cells, cell particles or soluble proteins. The support may then be washed with suitable buffers followed by treatment with the detectably labeled *HKNG1* gene product specific antibody. The solid phase support may then be washed with the buffer a 30 second time to remove unbound antibody. The amount of bound

label on the solid support may then be detected by conventional means.

By "solid phase support or carrier" is intended any support capable of binding an antigen or an antibody. Well-known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to an antigen or antibody. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Preferred supports include polystyrene beads. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

One of the ways in which the *HKNG1* gene product-specific antibody can be detectably labeled is by linking the same to an enzyme, such as for use in an enzyme immunoassay (EIA) (Voller, A., "The Enzyme Linked Immunosorbent Assay (ELISA)", 1978, Diagnostic Horizons 2:1-7, Microbiological Associates Quarterly Publication, Walkersville, MD); Voller, A. et al., 1978, J. Clin. Pathol. 31:507-520; Butler, J.E., 1981, Meth. Enzymol. 73:482-523; Maggio, E. (ed.), 1980, Enzyme Immunoassay, CRC Press, Boca Raton, FL; Ishikawa, E. et al., (eds.), 1981, Enzyme Immunoassay, Kagaku Shoin, Tokyo). The enzyme which is bound to the antibody will react with an appropriate substrate, preferably a chromogenic substrate, in such a manner as to produce a chemical moiety that can be detected, for example, by spectrophotometric, fluorimetric or by visual means. Enzymes that can be used to detectably label the antibody include, but are not limited to, malate

dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, α -glycerophosphate, dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, β -galactosidase, ribonuclease, urease, catalase, 5 glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. The detection can be accomplished by colorimetric methods that employ a chromogenic substrate for the enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

10 Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it is possible to detect *HKNG1* gene products through the use of a radioimmunoassay (RIA) (see, for example, Weintraub, B., Principles of Radioimmunoassays, Seventh Training Course on 15 Radioligand Assay Techniques, The Endocrine Society, March, 1986). The radioactive isotope can be detected by such means as the use of a gamma counter or a scintillation counter or by autoradiography.

It is also possible to label the antibody with a fluorescent compound. When the fluorescently labeled 20 antibody is exposed to light of the proper wave length, its presence can then be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and 25 fluorescamine.

The antibody can also be detectably labeled using fluorescence emitting metals such as ^{152}Eu , or others of the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or 30 ethylenediaminetetraacetic acid (EDTA).

The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly
5 useful chemiluminescent labeling compounds are luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt and oxalate ester.

Likewise, a bioluminescent compound may be used to label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in
10 which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

15

**5.8. SCREENING ASSAYS FOR COMPOUNDS
THAT MODULATE *HKNG1* GENE ACTIVITY**

The following assays are designed to identify compounds that bind to a *HKNG1* gene product, compounds that bind to proteins, or portions of proteins that interact with a *HKNG1*
20 gene product, compounds that modulate, e.g., interfere with, the interaction of a *HKNG1* gene product with proteins and compounds that modulate the activity of the *HKNG1* gene (i.e., modulate the level of *HKNG1* gene expression and/or modulate the level of *HKNG1* gene product activity). Assays may
25 additionally be utilized that identify compounds that bind to *HKNG1* gene regulatory sequences (e.g., promoter sequences; see e.g., Platt, 1994, J. Biol. Chem. 269, 28558-28562), and that can modulate the level of *HKNG1* gene expression. Such compounds may include, but are not limited to, small organic molecules, such as ones that are able to cross the blood-
30 brain barrier, gain to and/or entry into an appropriate cell

and affect expression of the *HKNG1* gene or some other gene involved in a *HKNG1* regulatory pathway.

Methods for the identification of such proteins are described, below, in Section 5.8.2. Such proteins may be involved in the control and/or regulation of mood. Further, among these compounds are compounds that affect the level of *HKNG1* gene expression and/or *HKNG1* gene product activity and that can be used in the therapeutic treatment of *HKNG1*-mediated disorders, e.g., neuropsychiatric disorders such as BAD and schizophrenia as described, below, in Section 5.9.

Compounds may include, but are not limited to, peptides such as, for example, soluble peptides, including but not limited to, Ig-tailed fusion peptides, and members of random peptide libraries; (see, e.g., Lam, et al., 1991, Nature 354:82-84; Houghten, et al., 1991, Nature 354:84-86), and combinatorial chemistry-derived molecular library made of D- and/or L- configuration amino acids, phosphopeptides (including, but not limited to members of random or partially degenerate, directed phosphopeptide libraries; see, e.g., Songyang, et al., 1993, Cell 72:767-778), antibodies (including, but not limited to, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')₂ and FAb expression library fragments, and epitope-binding fragments thereof), and small organic or inorganic molecules.

Such compounds may further comprise compounds, in particular drugs or members of classes or families of drugs, known to ameliorate the symptoms of a *HKNG1*-mediated disorder, e.g., a neuropsychiatric disorder such as BAD or schizophrenia.

Such compounds include families of antidepressants such as lithium salts, carbamazepine, valproic acid, lysergic acid diethylamide (LSD), *p*-chlorophenylalanine, *p*-propyldopacetamide dithiocarbamate derivatives e.g., FLA 63;

anti-anxiety drugs, e.g., diazepam; monoamine oxidase (MAO) inhibitors, e.g., iproniazid, clorgyline, phenelzine and isocarboxazid; biogenic amine uptake blockers, e.g., tricyclic antidepressants such as desipramine, imipramine and 5 amitriptyline; serotonin reuptake inhibitors e.g., fluoxetine; antipsychotic drugs such as phenothiazine derivatives (e.g., chlorpromazine (thorazine) and trifluopromazine)), butyrophenones (e.g., haloperidol (Haldol)), thioxanthene derivatives (e.g., chlorprothixene), 10 and dibenzodiazepines (e.g., clozapine); benzodiazepines; dopaminergic agonists and antagonists e.g., L-DOPA, cocaine, amphetamine, α -methyl-tyrosine, reserpine, tetrabenazine, benzotropine, pargyline; noradrenergic agonists and antagonists e.g., clonidine, phenoxybenzamine, phentolamine, tropolone.

15 Compounds identified via assays such as those described herein may be useful, for example, in elaborating the biological function of the *HKNG1* gene product and for ameliorating *HKNG1*-mediated neuropsychiatric disorders, such as BAD and schizophrenia. Assays for testing the 20 effectiveness of compounds identified by, for example, techniques such as those described in Sections 5.8.1 - 5.8.3, are discussed, below, in Section 5.8.4.

5.8.1. IN VITRO SCREENING ASSAYS FOR COMPOUNDS THAT BIND TO THE *HKNG1* GENE PRODUCT

25 *In vitro* systems may be designed to identify compounds capable of binding the *HKNG1* gene products of the invention. Compounds identified may be useful, for example, in modulating the activity of unimpaired and/or mutant *HKNG1* gene products, may be useful in elaborating the biological function of the *HKNG1* gene product, may be utilized in 30 screens for identifying compounds that disrupt normal *HKNG1*

gene product interactions, or may in themselves disrupt such interactions.

The principle of the assays used to identify compounds that bind to the *HKNG1* gene product involves preparing a reaction mixture of the *HKNG1* gene product and the test
5 compound under conditions and for a time sufficient to allow the two components to interact and bind, thus forming a complex that can be removed and/or detected in the reaction mixture. These assays can be conducted in a variety of ways. For example, one method to conduct such an assay involves
10 anchoring a *HKNG1* gene product or a test substance onto a solid support and detecting *HKNG1* gene product/test compound complexes formed on the solid support at the end of the reaction. In one embodiment of such a method, the *HKNG1* gene product may be anchored onto a solid support, and the test compound, which is not anchored, may be labeled, either
15 directly or indirectly.

In practice, microtiter plates are conveniently utilized as the solid support. The anchored component may be immobilized by non-covalent or covalent attachments. Non-covalent attachment may be accomplished by simply coating the solid surface with a solution of the protein and drying.
20 Alternatively, an immobilized antibody, preferably a monoclonal antibody, specific for the protein to be immobilized may be used to anchor the protein to the solid surface. The surfaces may be prepared in advance and stored.

In order to conduct the assay, the non-immobilized component is added to the coated surface containing the
25 anchored component. After the reaction is complete, unreacted components are removed (e.g., by washing) under conditions such that any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. Where the previously non-immobilized component is
30 pre-labeled, the detection of label immobilized on the

surface indicates that complexes were formed. Where the previously non-immobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the previously non-immobilized component (the antibody, in turn, may be directly labeled or indirectly labeled with a labeled anti-Ig antibody).

Alternatively, a reaction can be conducted in a liquid phase, the reaction products separated from unreacted components, and complexes detected; e.g., using an immobilized antibody specific for *HKNG1* gene product or the test compound to anchor any complexes formed in solution, and a labeled antibody specific for the other component of the possible complex to detect anchored complexes.

5.8.2. ASSAYS FOR PROTEINS THAT INTERACT WITH *HKNG1* GENE PRODUCTS

Any method suitable for detecting protein-protein interactions may be employed for identifying *HKNG1* gene product-protein interactions.

Among the traditional methods that may be employed are co-immunoprecipitation, cross-linking and co-purification through gradients or chromatographic columns. Utilizing procedures such as these allows for the identification of proteins, including intracellular proteins, that interact with *HKNG1* gene products. Once isolated, such a protein can be identified and can be used in conjunction with standard techniques, to identify proteins it interacts with. For example, at least a portion of the amino acid sequence of a protein that interacts with the *HKNG1* gene product can be ascertained using techniques well known to those of skill in the art, such as via the Edman degradation technique (see, e.g., Creighton, 1983, "Proteins: Structures and Molecular Principles," W.H. Freeman & Co., N.Y., pp.34-49). The amino acid sequence obtained may be used as a guide for the

generation of oligonucleotide mixtures that can be used to screen for gene sequences encoding such proteins. Screening made be accomplished, for example, by standard hybridization or PCR techniques. Techniques for the generation of oligonucleotide mixtures and the screening are well-known.

5 (See, e.g., Ausubel, *supra*, and 1990, "PCR Protocols: A Guide to Methods and Applications," Innis, *et al.*, eds. Academic Press, Inc., New York).

10 Additionally, methods may be employed that result in the simultaneous identification of genes that encode a protein which interacts with a *HKNG1* gene product. These methods include, for example, probing expression libraries with labeled *HKNG1* gene product, using *HKNG1* gene product in a manner similar to the well known technique of antibody probing of λ gt11 libraries.

15 One method that detects protein interactions *in vivo*, the two-hybrid system, is described in detail for illustration only and not by way of limitation. One version of this system has been described (Chien, *et al.*, 1991, Proc. Natl. Acad. Sci. USA, 88:9578-9582) and is commercially available from Clontech (Palo Alto, CA).

20 Briefly, utilizing such a system, plasmids are constructed that encode two hybrid proteins: one consists of the DNA-binding domain of a transcription activator protein fused to the *HKNG1* gene product and the other consists of the transcription activator protein's activation domain fused to an unknown protein that is encoded by a cDNA that has been recombined into this plasmid as part of a cDNA library. The DNA-binding domain fusion plasmid and the cDNA library are transformed into a strain of the yeast *Saccharomyces cerevisiae* that contains a reporter gene (e.g., HBS or *lacZ*) whose regulatory region contains the transcription activator's binding site. Either hybrid protein alone cannot activate transcription of the reporter gene: the DNA-binding domain hybrid cannot because it does not provide activation

25
30

function and the activation domain hybrid cannot because it cannot localize to the activator's binding sites.

Interaction of the two hybrid proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is detected by an assay for the reporter
5 gene product.

The two-hybrid system or related methodologies may be used to screen activation domain libraries for proteins that interact with the "bait" gene product. By way of example, and not by way of limitation, *HKNG1* gene products may be used
10 as the bait gene product. Total genomic or cDNA sequences are fused to the DNA encoding an activation domain. This library and a plasmid encoding a hybrid of a bait *HKNG1* gene product fused to the DNA-binding domain are co-transformed into a yeast reporter strain, and the resulting transformants are screened for those that express the reporter gene. For
15 example, a bait *HKNG1* gene sequence, such as the open reading frame of the *HKNG1* gene, can be cloned into a vector such that it is translationally fused to the DNA encoding the DNA-binding domain of the GAL4 protein. These colonies are purified and the library plasmids responsible for reporter gene expression are isolated. DNA sequencing is then used to
20 identify the proteins encoded by the library plasmids.

A cDNA library of the cell line from which proteins that interact with bait *HKNG1* gene product are to be detected can be made using methods routinely practiced in the art. According to the particular system described herein, for
25 example, the cDNA fragments can be inserted into a vector such that they are translationally fused to the transcriptional activation domain of GAL4. Such a library can be co-transformed along with the bait *HKNG1* gene-GAL4 fusion plasmid into a yeast strain that contains a lacZ gene driven by a promoter that contains GAL4 activation sequence. A cDNA encoded protein, fused to a GAL4 transcriptional
30 activation domain that interacts with bait *HKNG1* gene product

will reconstitute an active GAL4 protein and thereby drive expression of the HIS3 gene. Colonies that express HIS3 can be detected by their growth on petri dishes containing semi-solid agar based media lacking histidine. The cDNA can then be purified from these strains, and used to produce and
5 isolate the bait *HKNG1* gene product-interacting protein using techniques routinely practiced in the art.

5.8.3. ASSAYS FOR COMPOUNDS THAT INTERFERE WITH OR POTENTIATE *HKNG1* GENE PRODUCT MACROMOLECULE INTERACTION

10 The *HKNG1* gene products may, *in vivo*, interact with one or more macromolecules, including intracellular macromolecules, such as proteins. Such macromolecules may include, but are not limited to, nucleic acid molecules and those proteins identified via methods such as those
15 described, above, in Sections 5.8.1 - 5.8.2. For purposes of this discussion, the macromolecules are referred to herein as "binding partners". Compounds that disrupt *HKNG1* gene product binding to a binding partner may be useful in regulating the activity of the *HKNG1* gene product, especially mutant *HKNG1* gene products. Such compounds may include, but
20 are not limited to molecules such as peptides, and the like, as described, for example, in Section 5.8.2 above.

The basic principle of an assay system used to identify compounds that interfere with or potentiate the interaction between the *HKNG1* gene product and a binding partner or
25 partners involves preparing a reaction mixture containing the *HKNG1* gene product and the binding partner under conditions and for a time sufficient to allow the two to interact and bind, thus forming a complex. In order to test a compound for inhibitory activity, the reaction mixture is prepared in the presence and absence of the test compound. The test
30 compound may be initially included in the reaction mixture, or may be added at a time subsequent to the addition of *HKNG1*

gene product and its binding partner. Control reaction mixtures are incubated without the test compound or with a compound which is known not to block complex formation. The formation of any complexes between the *HKNG1* gene product and the binding partner is then detected. The formation of a complex in the control reaction, but not in the reaction mixture containing the test compound, indicates that the compound interferes with the interaction of the *HKNG1* gene product and the binding partner. Additionally, complex formation within reaction mixtures containing the test compound and normal *HKNG1* gene product may also be compared to complex formation within reaction mixtures containing the test compound and a mutant *HKNG1* gene product. This comparison may be important in those cases wherein it is desirable to identify compounds that disrupt interactions of mutant but not normal *HKNG1* gene product.

In order to test a compound for potentiating activity, the reaction mixture is prepared in the presence and absence of the test compound. The test compound may be initially included in the reaction mixture, or may be added at a time subsequent to the addition of *HKNG1* gene product and its binding partner. Control reaction mixtures are incubated without the test compound or with a compound which is known not to block complex formation. The formation of any complexes between the *HKNG1* gene product and the binding partner is then detected. Increased formation of a complex in the reaction mixture containing the test compound, but not in the control reaction, indicates that the compound enhances and therefore potentiates the interaction of the *HKNG1* gene product and the binding partner. Additionally, complex formation within reaction mixtures containing the test compound and normal *HKNG1* gene product may also be compared to complex formation within reaction mixtures containing the test compound and a mutant *HKNG1* gene product. This comparison may be important in those cases wherein it is

desirable to identify compounds that enhance interactions of mutant but not normal *HKNG1* gene product.

In alternative embodiments, the above assays may be performed using a reaction mixture containing the *HKNG1* gene product, a binding partner, and a third which disrupts or enhances *HKNG1* gene product binding to the binding partner. The reaction mixture is prepared and incubated in the presence and absence of the test compound, as described above, and the formation of any complexes between the *HKNG1* gene product and the binding partner is detected. In this embodiment, the formation of a complex in the reaction mixture containing the test compound, but not in the control reaction, indicates that the test compound interferes with the ability of the second compound to disrupt *HKNG1* gene product binding to its binding partner.

The assays for compounds that interfere with or potentiate the interaction of the *HKNG1* gene products and binding partners can be conducted in a heterogeneous or homogeneous format. Heterogeneous assays involve anchoring either the *HKNG1* gene product or the binding partner onto a solid support and detecting complexes formed on the solid support at the end of the reaction. In homogeneous assays, the entire reaction is carried out in a liquid phase. In either approach, the order of addition of reactants can be varied to obtain different information about the compounds being tested. For example, test compounds that interfere with or potentiate the interaction between the *HKNG1* gene products and the binding partners, e.g., by competition, can be identified by conducting the reaction in the presence of the test substance; i.e., by adding the test substance to the reaction mixture prior to or simultaneously with the *HKNG1* gene product and interactive intracellular binding partner. Alternatively, test compounds that disrupt preformed complexes, e.g., compounds with higher binding constants that

displace one of the components from the complex, can be tested by adding the test compound to the reaction mixture after complexes have been formed. The various formats are described briefly below.

5 In a heterogeneous assay system, either the *HKNG1* gene product or the interactive binding partner, is anchored onto a solid surface, while the non-anchored species is labeled, either directly or indirectly. In practice, microtiter plates are conveniently utilized. The anchored species may be immobilized by non-covalent or covalent attachments. Non-covalent attachment may be accomplished simply by coating the
10 solid surface with a solution of the *HKNG1* gene product or binding partner and drying. Alternatively, an immobilized antibody specific for the species to be anchored may be used to anchor the species to the solid surface. The surfaces may be prepared in advance and stored.

15 In order to conduct the assay, the partner of the immobilized species is exposed to the coated surface with or without the test compound. After the reaction is complete, unreacted components are removed (e.g., by washing) and any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid
20 surface can be accomplished in a number of ways. Where the non-immobilized species is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the non-immobilized species is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody
25 specific for the initially non-immobilized species (the antibody, in turn, may be directly labeled or indirectly labeled with a labeled anti-Ig antibody). Depending upon the order of addition of reaction components, test compounds that inhibit complex formation or that disrupt preformed complexes can be detected.

30 Alternatively, the reaction can be conducted in a liquid phase in the presence or absence of the test compound, the

reaction products separated from unreacted components, and complexes detected; e.g., using an immobilized antibody specific for one of the binding components to anchor any complexes formed in solution, and a labeled antibody specific for the other partner to detect anchored complexes. Again,
5 depending upon the order of addition of reactants to the liquid phase, test compounds that inhibit complex formation or that disrupt preformed complexes can be identified.

In an alternate embodiment of the invention, a homogeneous assay can be used. In this approach, a preformed complex of the *HKNG1* gene product and the interactive binding
10 partner is prepared in which either the *HKNG1* gene product or its binding partners is labeled, but the signal generated by the label is quenched due to complex formation (see, e.g., U.S. Patent No. 4,109,496 by Rubenstein which utilizes this approach for immunoassays). The addition of a test substance
15 that competes with and displaces one of the species from the preformed complex will result in the generation of a signal above background. In this way, test substances that disrupt *HKNG1* gene product/binding partner interaction can be identified.

In another embodiment of the invention, these same
20 techniques can be employed using peptide fragments that correspond to the binding domains of the *HKNG1* product and/or the binding partner (in cases where the binding partner is a protein), in place of one or both of the full length proteins. Any number of methods routinely practiced in the art can be used to identify and isolate the binding sites.
25 These methods include, but are not limited to, mutagenesis of the gene encoding one of the proteins and screening for disruption of binding in a co-immunoprecipitation assay. Compensating mutations in the gene encoding the second species in the complex can then be selected. Sequence analysis of the genes encoding the respective proteins will
30 reveal the mutations that correspond to the region of the

protein involved in interactive binding. Alternatively, one protein can be anchored to a solid surface using methods described in this Section above, and allowed to interact with and bind to its labeled binding partner, which has been treated with a proteolytic enzyme, such as trypsin. After
5 washing, a short, labeled peptide comprising the binding domain may remain associated with the solid material, which can be isolated and identified by amino acid sequencing. Also, once the gene coding for the segments is engineered to express peptide fragments of the protein, it can then be tested for binding activity and purified or synthesized.
10 For example, and not by way of limitation, a *HKNG1* gene product can be anchored to a solid material as described, above, in this Section by making a *GST-HKNG1* fusion protein and allowing it to bind to glutathione agarose beads. The binding partner can be labeled with a radioactive isotope, such as ³⁵S, and cleaved with a proteolytic enzyme such as
15 trypsin. Cleavage products can then be added to the anchored *GST-HKNG1* fusion protein and allowed to bind. After washing away unbound peptides, labeled bound material, representing the binding partner binding domain, can be eluted, purified, and analyzed for amino acid sequence by well-known methods.
20 Peptides so identified can be produced synthetically or produced using recombinant DNA technology.

5.8.4. ASSAYS FOR IDENTIFICATION OF COMPOUNDS THAT AMELIORATE A *HKNG1*-MEDIATED DISORDER

Compounds, including but not limited to binding
25 compounds identified via assay techniques such as those described, above, in Sections 5.8.1 - 5.8.4, can be tested for the ability to ameliorate symptoms of a *HKNG1*-mediated disorder, e.g., a CNS-related disorder, such as a neuropsychiatric disorder, including schizophrenia and bipolar affective (mood) disorders, including severe bipolar
30 affective (mood) disorder (BP-I), bipolar affective (mood)

disorder with hypomania and major depression (BP-II), and myopia disorders

It should be noted that the assays described herein can identify compounds that affect *HKNG1* activity by either affecting *HKNG1* gene expression or by affecting the level of *HKNG1* gene product activity. For example, compounds may be identified that are involved in another step in the pathway in which the *HKNG1* gene and/or *HKNG1* gene product is involved and, by affecting this same pathway may modulate the effect of *HKNG1* on the development of a *HKNG1*-mediated disorder.

Such compounds can be used, e.g., as part of a therapeutic method for the treatment of the disorder.

Described below are cell-based and animal model-based assays for the identification of compounds exhibiting such an ability to ameliorate symptoms of a *HKNG1*-mediated disorder, e.g., neuropsychiatric disorder, such as BAD or schizophrenia.

First, cell-based systems can be used to identify compounds that may act to ameliorate symptoms of a *HKNG1*-mediated disorder. Such cell systems can include, for example, recombinant or non-recombinant cell, such as cell lines, that express the *HKNG1* gene.

In utilizing such cell systems, cells that express *HKNG1* may be exposed to a compound suspected of exhibiting an ability to ameliorate symptoms of a *HKNG1*-mediated disorder, e.g., a neuropsychiatric disorder, such as BAD or schizophrenia, at a sufficient concentration and for a sufficient time to elicit such an amelioration of such symptoms in the exposed cells. After exposure, the cells can be assayed to measure alterations in the expression of the *HKNG1* gene, e.g., by assaying cell lysates for *HKNG1* mRNA transcripts (e.g., by Northern analysis) or for *HKNG1* gene products expressed by the cell; compounds that modulate

expression of the *HKNG1* gene are good candidates as therapeutics.

In addition, animal-based systems or models for a *HKNG1*-mediated disorder, e.g., neuropsychiatric disorder, for example, transgenic mice containing a human or altered form of *HKNG1* gene, may be used to identify compounds capable of ameliorating symptoms of the disorder. Such animal models may be used as test substrates for the identification of drugs, pharmaceuticals, therapies and interventions. For example, animal models may be exposed to a compound suspected of exhibiting an ability to ameliorate symptoms, at a sufficient concentration and for a sufficient time to elicit such an amelioration of symptoms of a *HKNG1*-mediated disorder. The response of the animals to the exposure may be monitored by assessing the reversal of the symptoms of the disorder.

With regard to intervention, any treatments that reverse any aspect of symptoms of a *HKNG1*-mediated disorder, should be considered as candidates for human therapeutic intervention in such disorders. Dosages of test agents may be determined by deriving dose-response curves, as discussed in Section 5.10.1, below.

5.9. COMPOUNDS AND METHODS FOR THE TREATMENT OF *HKNG1*-MEDIATED DISORDERS

Described below are methods and compositions whereby a *HKNG1*-mediated disorder described herein, e.g., a *HKNG1*-mediated neuropsychiatric disorder, such as BAD or schizophrenia, may be treated. For example, such methods can comprise administering compounds which modulate the expression of a mammalian *HKNG1* gene and/or the synthesis or activity of a mammalian *HKNG1* gene product (e.g., a recombinant *HKNG1* gene product) so symptoms of the disorder are ameliorated.

Alternatively, in those instances whereby the *HKNG1*-mediated disorders result from *HKNG1* gene mutations, such methods can comprise supplying the subject with a nucleic acid molecule encoding an unimpaired *HKNG1* gene product such
5 that an unimpaired *HKNG1* gene product is expressed and symptoms of the disorder are ameliorated.

In another embodiment of methods for the treatment of *HKNG1*-mediated disorders resulting from *HKNG1* gene mutations, such methods can comprise supplying the subject with a cell comprising a nucleic acid molecule that encodes an unimpaired
10 *HKNG1* gene product such that the cell expresses the unimpaired *HKNG1* gene product and symptoms of the disorder are ameliorated.

In cases in which a loss of normal *HKNG1* gene product function results in the development of a *HKNG1*-mediated
15 disorder an increase in *HKNG1* gene product activity would facilitate progress towards an asymptomatic state in individuals exhibiting a deficient level of *HKNG1* gene expression and/or *HKNG1* gene product activity. Methods for enhancing the expression or synthesis of *HKNG1* can include,
20 for example, methods such as those described below, in Section 5.9.2.

Alternatively, symptoms of *HKNG1*-mediated neuropsychiatric disorders, may be ameliorated by administering a compound that decreases the level of *HKNG1*
25 gene expression and/or *HKNG1* gene product activity. Methods for inhibiting or reducing the level of *HKNG1* gene product synthesis or expression can include, for example, methods such as those described in Section 5.9.1.

In one embodiment of treatment methods, the compounds administered comprise compounds, in particular drugs, which
30 ameliorate the symptoms of a disorder described herein as a neuropsychiatric disorder, such as BAD or schizophrenia.

Such compounds include drugs within the families of antidepressants such as lithium salts, carbamazepine, valproic acid, lysergic acid diethylamide (LSD), *p*-chlorophenylalanine, *p*-propyldopacetamide dithiocarbamate derivatives e.g., FLA 63; anti-anxiety drugs, e.g., diazepam; 5 monoamine oxidase (MAO) inhibitors, e.g., iproniazid, clorgyline, phenelzine and isocarboxazid; biogenic amine uptake blockers, e.g., tricyclic antidepressants such as desipramine, imipramine and amitriptyline; serotonin reuptake inhibitors e.g., fluoxetine; antipsychotic drugs such as 10 phenothiazine derivatives (e.g., chlorpromazine (thorazine) and trifluopromazine)), butyrophenones (e.g., haloperidol (Haldol)), thioxanthene derivatives (e.g., chlorprothixene), and dibenzodiazepines (e.g., clozapine); benzodiazepines; dopaminergic agonists and antagonists e.g., L-DOPA, cocaine, 15 amphetamine, α -methyl-tyrosine, reserpine, tetrabenazine, benzotropine, pargyline; noradrenergic agonists and antagonists e.g., clonidine, phenoxybenzamine, phentolamine, tropolone.

In another embodiment, symptoms of a disorder described herein, e.g., a *HKNG1*-mediated neuropsychiatric disorder such 20 as BAD or schizophrenia, may be ameliorated by *HKNG1* protein therapy methods, e.g., decreasing or increasing the level and/or of *HKNG1*-activity using the *HKNG1* protein, fusion protein, and peptide sequences described in Section 5.2, above, or by the administration of proteins or protein 25 fragments (e.g., peptides) which interact with an *HKNG1* gene or gene product and thereby inhibit or potentiate its activity.

Such protein therapy may include, for example, the administration of a functional *HKNG1* protein or fragments of an *HKNG1* protein (e.g., peptides) which represent functional 30 *HKNG1* domains.

In one embodiment, HKNG1 fragments or peptides representing a functional HKNG1 binding domain are administered to an individual such that the peptides bind to an HKNG1 binding protein, e.g., an HKNG1 receptor. Such fragments or peptides may serve to inhibit HKNG1 activity in an individual by competing with, and thereby inhibiting, binding of HKNG1 to the binding protein, thereby ameliorating symptoms of a disorder described herein. Alternatively, such fragments or peptides may enhance HKNG1 activity in an individual by mimicking the function of HKNG1 *in vivo*, thereby ameliorating the symptoms of a disorder described herein.

The proteins and peptides which may be used in the methods of the invention include synthetic (e.g., recombinant or chemically synthesized) proteins and peptides, as well as naturally occurring proteins and peptides. The proteins and peptides may have both naturally occurring and non-naturally occurring amino acid residues (e.g., D-amino acid residues) and/or one or more non-peptide bonds (e.g., imino, ester, hydrazide, semicarbazide, and azo bonds). The proteins or peptides may also contain additional chemical groups (*i.e.*, functional groups) present at the amino and/or carboxy termini, such that, for example, the stability, bioavailability, and/or inhibitory activity of the peptide is enhanced. Exemplary functional groups include hydrophobic groups (e.g. carbobenzoxy, dansyl, and t-butyloxycarbonyl groups), an acetyl group, a 9-fluorenylmethoxy-carbonyl group, and macromolecular carrier groups (e.g., lipid-fatty acid conjugates, polyethylene glycol, or carbohydrates) including peptide groups.

5.9.1. INHIBITORY ANTISENSE, RIBOZYME AND TRIPLE HELIX APPROACHES

In another embodiment, symptoms of HKNG1-mediated neuropsychiatric disorders may be ameliorated by decreasing

the level of *HKNG1* gene expression and/or *HKNG1* gene product activity by using *HKNG1* gene sequences in conjunction with well-known antisense, gene "knock-out," ribozyme and/or triple helix methods to decrease the level of *HKNG1* gene
5 expression. Among the compounds that may exhibit the ability to modulate the activity, expression or synthesis of the *HKNG1* gene, including the ability to ameliorate the symptoms of a *HKNG1*-mediated neuropsychiatric disorder, such as BAD or schizophrenia, are antisense, ribozyme, and triple helix molecules. Such molecules may be designed to reduce or
10 inhibit either unimpaired, or if appropriate, mutant target gene activity. Techniques for the production and use of such molecules are well known to those of skill in the art.

Antisense RNA and DNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense approaches involve
15 the design of oligonucleotides that are complementary to a target gene mRNA. The antisense oligonucleotides will bind to the complementary target gene mRNA transcripts and prevent translation. Absolute complementarity, although preferred, is not required.

A sequence "complementary" to a portion of an RNA, as
20 referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity
25 and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized
30 complex.

In one embodiment, oligonucleotides complementary to non-coding regions of the *HKNG1* gene could be used in an antisense approach to inhibit translation of endogenous *HKNG1* mRNA. Antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

Regardless of the choice of target sequence, it is preferred that *in vitro* studies are first performed to quantitate the ability of the antisense oligonucleotide to inhibit gene expression. It is preferred that these studies utilize controls that distinguish between antisense gene inhibition and nonspecific biological effects of oligonucleotides. It is also preferred that these studies compare levels of the target RNA or protein with that of an internal control RNA or protein. Additionally, it is envisioned that results obtained using the antisense oligonucleotide are compared with those obtained using a control oligonucleotide. It is preferred that the control oligonucleotide is of approximately the same length as the test oligonucleotide and that the nucleotide sequence of the oligonucleotide differs from the antisense sequence no more than is necessary to prevent specific hybridization to the target sequence.

The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger, et al., 1989,

Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre, et al., 1987, Proc. Natl. Acad. Sci. U.S.A. 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. 5 WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents (see, e.g., Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents (see, e.g., Zon, 1988, Pharm. Res. 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, 10 transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 15 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 20 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil- 25 5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, 30 and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an
5 alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier, et al.,
10 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue, et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue, et al., 1987, FEBS Lett. 215:327-330).

Oligonucleotides of the invention may be synthesized by
15 standard methods known in the art, e.g., by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein, et al. (1988, Nucl. Acids Res. 16:3209),
20 methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin, et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

While antisense nucleotides complementary to the target gene coding region sequence could be used, those complementary to the transcribed, untranslated region are
25 most preferred.

Antisense molecules should be delivered to cells that express the target gene *in vivo*. A number of methods have been developed for delivering antisense DNA or RNA to cells; e.g., antisense molecules can be injected directly into the tissue site, or modified antisense molecules, designed to
30 target the desired cells (e.g., antisense linked to peptides or antibodies that specifically bind receptors or antigens

expressed on the target cell surface) can be administered systemically.

A preferred approach to achieve intracellular concentrations of the antisense sufficient to suppress translation of endogenous mRNAs utilizes a recombinant DNA
5 construct in which the antisense oligonucleotide is placed under the control of a strong pol III or pol II promoter. The use of such a construct to transfect target cells in the patient will result in the transcription of sufficient amounts of single stranded RNAs that will form complementary base pairs with the endogenous target gene transcripts and
10 thereby prevent translation of the target gene mRNA. For example, a vector can be introduced e.g., such that it is taken up by a cell and directs the transcription of an antisense RNA. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be
15 constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the antisense RNA can be by any promoter known in the art to act in mammalian, preferably human cells. Such promoters can be inducible or
20 constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, Nature 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, Cell 22:787-797), the herpes thymidine kinase promoter
25 (Wagner, et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster, et al., 1982, Nature 296:39-42), etc. Any type of plasmid, cosmid, YAC or viral vector can be used to prepare the recombinant DNA construct which can be introduced directly into the tissue site. Alternatively, viral vectors
30 can be used that selectively infect the desired tissue, in

which case administration may be accomplished by another route (e.g., systemically).

Ribozyme molecules designed to catalytically cleave target gene mRNA transcripts can also be used to prevent translation of target gene mRNA and, therefore, expression of target gene product. (See, e.g., PCT International Publication WO90/11364, published October 4, 1990; Sarver, et al., 1990, Science 247, 1222-1225).

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. (For a review, see Rossi, 1994, Current Biology 4:469-471). The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage event. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA, and must include the well known catalytic sequence responsible for mRNA cleavage. For this sequence, see, e.g., U.S. Patent No. 5,093,246, which is incorporated herein by reference in its entirety.

While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy target gene mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Myers, 1995, *Molecular Biology and Biotechnology: A Comprehensive Desk Reference*, VCH Publishers, New York, (see especially Figure 4, page 833) and in Haseloff and Gerlach, 1988, Nature, 334:585-591, which is incorporated herein by reference in its entirety.

Preferably the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the

target gene mRNA, i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as
5 the one that occurs naturally in *Tetrahymena thermophila* (known as the IVS, or L-19 IVS RNA) and that has been extensively described by Thomas Cech and collaborators (Zaug, et al., 1984, *Science*, 224:574-578; Zaug and Cech, 1986, *Science*, 231:470-475; Zaug, et al., 1986, *Nature*, 324:429-
10 433; published International patent application No. WO 88/04300 by University Patents Inc.; Been and Cech, 1986, *Cell*, 47:207-216). The Cech-type ribozymes have an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes which
15 target eight base-pair active site sequences that are present in the target gene.

As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (e.g., for improved stability, targeting, etc.) and should be delivered to cells that express the target gene *in vivo*. A preferred method of
20 delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous target gene messages and inhibit translation. Because
25 ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Endogenous target gene expression can also be reduced by inactivating or "knocking out" the target gene or its promoter using targeted homologous recombination (e.g., see Smithies, et al., 1985, *Nature* 317:230-234; Thomas and
30 Capecchi, 1987, *Cell* 51:503-512; Thompson, et al., 1989, *Cell* 5:313-321; each of which is incorporated by reference herein

in its entirety). For example, a mutant, non-functional target gene (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous target gene (either the coding regions or regulatory regions of the target gene) can be used, with or without a selectable marker and/or a
5 negative selectable marker, to transfect cells that express the target gene *in vivo*. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the target gene. Such approaches are particularly suited in the agricultural field where modifications to ES (embryonic stem) cells can be used to generate animal offspring with an
10 inactive target gene (e.g., see Thomas and Capecchi, 1987 and Thompson, 1989, *supra*). However this approach can be adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors.

15 Alternatively, endogenous target gene expression can be reduced by targeting deoxyribonucleotide sequences complementary to the regulatory region of the target gene (*i.e.*, the target gene promoter and/or enhancers) to form triple helical structures that prevent transcription of the target gene in target cells in the body. (See generally,
20 Helene, 1991, *Anticancer Drug Des.*, 6(6):569-584; Helene, et al., 1992, *Ann. N.Y. Acad. Sci.*, 660:27-36; and Maher, 1992, *Bioassays* 14(12):807-815).

Nucleic acid molecules to be used in triplex helix formation for the inhibition of transcription should be
25 single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences may be pyrimidine-based, which
30 will result in TAT and CGC⁺ triplets across the three associated strands of the resulting triple helix. The

pyrimidine-rich molecules provide base complementarily to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules may be chosen that are purine-rich, for example, contain a stretch of G residues. These molecules
5 will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

Alternatively, the potential sequences that can be targeted for triple helix formation may be increased by
10 creating a so called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

15 In instances wherein the antisense, ribozyme, and/or triple helix molecules described herein are utilized to inhibit mutant gene expression, it is possible that the technique may so efficiently reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by normal target gene alleles that
20 the possibility may arise wherein the concentration of normal target gene product present may be lower than is necessary for a normal phenotype. In such cases, to ensure that substantially normal levels of target gene activity are maintained, therefore, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal target
25 gene activity may, be introduced into cells via gene therapy methods such as those described, below, in Section 5.9.2 that do not contain sequences susceptible to whatever antisense, ribozyme, or triple helix treatments are being utilized. Alternatively, in instances whereby the target gene encodes an extracellular protein, it may be preferable to co-
30 administer normal target gene protein in order to maintain the requisite level of target gene activity.

Anti-sense RNA and DNA, ribozyme, and triple helix molecules of the invention may be prepared by any method known in the art for the synthesis of DNA and RNA molecules, as discussed above. These include techniques for chemically synthesizing oligodeoxyribonucleotides and
5 oligoribonucleotides well known in the art such as for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a
10 wide variety of vectors that incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

15

5.9.2. GENE REPLACEMENT THERAPY

HKNG1 gene nucleic acid sequences, described above in Section 5.1, can be utilized for transferring recombinant *HKNG1* nucleic acid sequences to cells and expressing said sequences in recipient cells. Such techniques can be used,
20 for example, in marking cells or for the treatment of a *HKNG1*-mediated neuropsychiatric disorder. Such treatment can be in the form of gene replacement therapy. Specifically, one or more copies of a normal *HKNG1* gene or a portion of the *HKNG1* gene that directs the production of a *HKNG1* gene
25 product exhibiting normal *HKNG1* gene function, may be inserted into the appropriate cells within a patient, using vectors that include, but are not limited to adenovirus, adeno-associated virus, and retrovirus vectors, in addition to other particles that introduce DNA into cells, such as liposomes.

30 Because the *HKNG1* gene is expressed in the brain, such gene replacement therapy techniques should be capable of

delivering *HKNG1* gene sequences to these cell types within patients. Thus, in one embodiment, techniques that are well known to those of skill in the art (see, e.g., PCT Publication No. WO89/10134, published April 25, 1988) can be used to enable *HKNG1* gene sequences to cross the blood-brain barrier readily and to deliver the sequences to cells in the brain. With respect to delivery that is capable of crossing the blood-brain barrier, viral vectors such as, for example, those described above, are preferable.

In another embodiment, techniques for delivery involve direct administration, e.g., by stereotactic delivery of such *HKNG1* gene sequences to the site of the cells in which the *HKNG1* gene sequences are to be expressed.

Additional methods that may be utilized to increase the overall level of *HKNG1* gene expression and/or *HKNG1* gene product activity include using targeted homologous recombination methods, discussed in Section 5.2, above, to modify the expression characteristics of an endogenous *HKNG1* gene in a cell or microorganism by inserting a heterologous DNA regulatory element such that the inserted regulatory element is operatively linked with the endogenous *HKNG1* gene in question. Targeted homologous recombination can thus be used to activate transcription of an endogenous *HKNG1* gene that is "transcriptionally silent", i.e., is not normally expressed or is normally expressed at very low levels, or to enhance the expression of an endogenous *HKNG1* gene that is normally expressed.

Further, the overall level of *HKNG1* gene expression and/or *HKNG1* gene product activity may be increased by the introduction of appropriate *HKNG1*-expressing cells, preferably autologous cells, into a patient at positions and in numbers that are sufficient to ameliorate the symptoms of

a *HKNG1*-mediated neuropsychiatric disorder. Such cells may be either recombinant or non-recombinant.

Among the cells that can be administered to increase the overall level of *HKNG1* gene expression in a patient are
5 normal cells, preferably brain cells, that express the *HKNG1* gene. Alternatively, cells, preferably autologous cells, can be engineered to express *HKNG1* gene sequences, and may then be introduced into a patient in positions appropriate for the amelioration of the symptoms of a *HKNG1*-mediated
10 neuropsychiatric disorder. Alternately, cells that express an unimpaired *HKNG1* gene and that are from a MHC matched individual can be utilized, and may include, for example, brain cells. The expression of the *HKNG1* gene sequences is controlled by the appropriate gene regulatory sequences to allow such expression in the necessary cell types. Such gene
15 regulatory sequences are well known to the skilled artisan. Such cell-based gene therapy techniques are well known to those skilled in the art, see, e.g., Anderson, U.S. Patent No. 5,399,349.

When the cells to be administered are non-autologous cells, they can be administered using well known techniques
20 that prevent a host immune response against the introduced cells from developing. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

25 Additionally, compounds, such as those identified via techniques such as those described, above, in Section 5.8, that are capable of modulating *HKNG1* gene product activity can be administered using standard techniques that are well known to those of skill in the art. In instances in which the compounds to be administered are to involve an
30 interaction with brain cells, the administration techniques

should include well known ones that allow for a crossing of the blood-brain barrier.

**5.10. PHARMACEUTICAL PREPARATIONS
AND METHODS OF ADMINISTRATION**

5 The compounds that are determined to affect *HKNG1* gene expression or gene product activity can be administered to a patient at therapeutically effective doses to treat or ameliorate a *HKNG1*-mediated disorder or modulate a *HKNG1*-related process described herein. A therapeutically effective dose refers to that amount of the compound
10 sufficient to result in amelioration of symptoms of such a disorder.

5.10.1. EFFECTIVE DOSE

Toxicity and therapeutic efficacy of such compounds can
15 be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD_{50} (the dose lethal to 50% of the population) and the ED_{50} (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as
20 the ratio LD_{50}/ED_{50} . Compounds that exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

25 The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage may vary within this range depending upon the dosage form
30 employed and the route of administration utilized. For any compound used in the method of the invention, the

therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC_{50} (*i.e.*, the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

As defined herein, a therapeutically effective amount of antibody, protein, or polypeptide (*i.e.*, an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight.

The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a protein, polypeptide, or antibody can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with antibody, protein, or polypeptide in the range of between about 0.1 to 20 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. It will also be appreciated that the effective dosage of antibody, protein, or polypeptide used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may result and become apparent from the results of diagnostic assays as described herein.

5.10.2. FORMULATIONS AND USE

Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers or excipients.

5 Thus, the compounds and their physiologically acceptable salts and solvates may be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral rectal or topical administration.

10 For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, 15 microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form 20 of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or 25 hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer 30 salts, flavoring, coloring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

5 For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane,
10 dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as
15 lactose or starch.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The
20 compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

25 The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In certain embodiments, it may be desirable to administer the pharmaceutical compositions of the invention
30 locally to the area in need of treatment. This may be

achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, 5 non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. In one embodiment, administration can be by direct injection at the site (or former site) of a malignant tumor or neoplastic or pre-neoplastic tissue.

10 For topical application, the compounds may be combined with a carrier so that an effective dosage is delivered, based on the desired activity

A topical formulation for treatment of some of the eye disorders discussed *infra* (e.g., myopia) consists of an effective amount of the compounds in a ophthalmologically acceptable excipient such as buffered saline, mineral oil, 15 vegetable oils such as corn or arachis oil, petroleum jelly, Miglyol 182, alcohol solutions, or liposomes or liposome-like products. Any of these compositions may also include preservatives, antioxidants, antibiotics, immunosuppressants, and other biologically or pharmaceutically effective agents 20 which do not exert a detrimental effect on the compound.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the 25 compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

The compositions may, if desired, be presented in a pack 30 or dispenser device that may contain one or more unit dosage forms containing the active ingredient. The pack may for

example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

5 6. **EXAMPLE: THE *HKNG1* GENE OF CHROMOSOME 18 IS ASSOCIATED WITH THE NEUROPSYCHIATRIC DISORDER BAD**

In the Example presented in this Section, studies are described that define a narrow interval of approximately 27 kb on the short arm of human chromosome 18 which is associated with the neuropsychiatric disorder BAD. The
10 interval is demonstrated to lie within the gene referred to herein as the *HKNG1* gene.

6.1. **MATERIALS AND METHODS**

6.1.1. **LINKAGE DISEQUILIBRIUM**

15 Linkage disequilibrium (LD) studies were performed using DNA from a population sample of neuropsychiatric disorder (BP-I) patients. The population sample and LD techniques were as described in Escamilla *et al.*, 1996, *Am J. Med. Genet.* 67:244-253. The present LD study took advantage of
20 the additional population sample collection and the additional physical markers identified via the physical mapping techniques described below.

6.1.2. **YEAST ARTIFICIAL CHROMOSOME (YAC) MAPPING**

For physical mapping, yeast artificial chromosomes (YACs) containing human sequences were mapped to the region
25 being analyzed based on publicly available maps (Cohen *et al.*, 1993, *C.R. Acad. Sci.* 316:1484-1488). The YACs were then ordered and contig reconstructed by performing standard short tag sequence (STS)-content mapping with microsatellite markers and non-polymorphic STSs available from databases
30 that surround the genetically defined candidate region.

6.1.3. BACTERIAL ARTIFICIAL CHROMOSOME (BAC) MAPPING

STSS from the short arm of human chromosome 18 were used to screen a human BAC library (Research Genetics, Huntsville, AL). The ends of the BACs were cloned or directly sequenced. The end sequences were used to amplify the next overlapping BACs. From each BAC, additional microsatellites were identified. Specifically, random sheared libraries were prepared from overlapping BACs within the defined genetic interval. BAC DNA was sheared with a nebulizer (CIS-US Inc., Bedford, MA). Fragments in the size range of 600 to 1,000 bp were utilized for the sublibrary production. Microsatellite sequences from the sublibraries were identified by corresponding microsatellite probes. Sequences around such repeats were obtained to enable development of PCR primers for genomic DNA.

6.1.4. RADIATION HYBRID (RH) MAPPING

Standard RH mapping techniques were applied to a Stanford G3 RH mapping panel (Research Genetics, Huntsville, AL) to order all microsatellite markers and non-polymorphic STSS in the region being analyzed.

6.1.5. SAMPLE SEQUENCING

Random sheared libraries were made from all the BACs within the defined genetic region. Approximately 9,000 subclones within the approximately 340 kb region containing the BAD interval were sequenced with vector primers in order to achieve an 8-fold sequence coverage of the region. All sequences were processed through an automated sequence analysis pipeline that assessed quality, removed vector sequences and masked repetitive sequences. The resulting sequences were then compared to public DNA and protein databases using BLAST algorithms (Altschul, et al., 1990, J. Molec. Biol., 215:403-410).

All sequences were contiged using Sequencher 3.0 (Gene Code Corp.) and PHRED and PHRAP (Phill Green, Washington University) into a single DNA fragment of 340 kb.

6.2. RESULTS

5 Genetic regions involved in bipolar affective disorder (BAD) human genes had previously been reported to map to portions of the long (18q) and short (18p) arms of human chromosome 18 (Freimer *et al.*, 1996, *Neuropsychiat. Genet.* 67:254-263; Freimer *et al.*, 1996, *Nature Genetics* 12:436-441; 10 and McInnis *et al.*, *Proc. Natl. Acad. Scie. U.S.A.* 93:13060-13065).

High resolution physical mapping using YAC, BAC and RH techniques. In order to provide the precise order of genetic markers necessary for linkage and LD mapping, and to guide new microsatellite marker development for finer mapping, a 15 high resolution physical map of the 18p candidate region was developed using YAC, BAC and RH techniques.

For such physical mapping, first, YACs were mapped to the chromosome 18 region being analyzed. Using the mapped YAC contig as a framework, the region from publicly available markers spanning the 18p region were also mapped and contiged 20 with BACs. Sublibraries from the contiged BACs were constructed, from which microsatellite marker sequences were identified and sequenced.

To ensure development of an accurate physical map, the radiation hybrid (RH) mapping technique was independently applied to the region being analyzed. RH was used to order 25 all microsatellite markers and non-polymorphic STSs in the region. Thus, the high resolution physical map ultimately constructed was obtained using data from RH mapping and STS-content mapping.

Linkage Disequilibrium. Prior to attempting to identify gene sequences, studies were performed to further narrow the 30 neuropsychiatric disorder region. Specifically, a linkage disequilibrium (LD) analysis was performed using population

samples and techniques as described in Section 6.1, above, which took advantage of the additional physical markers identified via the physical mapping techniques described below.

Initial LD analysis narrowed the interval which
5 associates with BAD disorders to a 340 kb region of 18p. BAC clones within this newly identified neuropsychiatric disorder region were analyzed to identify specific genes within the region. A combination of sample sequencing, cDNA selection and transcription mapping analyses were used to arrange
10 sequences into tentative transcription units, that is, tentatively delineating the coding sequences of genes within this genomic region of interest.

Subsequent LD analyses further narrowed the BAD region of 18p to a narrow interval of approximately 27 kb. This was accomplished by identifying the maximum haplotype shared among affected individuals using additional markers.
15 Statistical analysis of the entire 18p candidate region indicated that the 27 kb haplotype was significantly elevated in frequency among affected Costa Rican individuals (LOD = 2.2; $p = 0.0005$).

This newly identified narrow interval was found to map completely within one of the transcription units identified
20 as described above. The gene corresponding to this transcription unit is referred to herein as the *HKNG1* gene. Thus, the results of the mapping analyses presented in this Section demonstrate that the *HKNG1* gene of human chromosome 18 is associated the neuropsychiatric disorder BAD.

Analysis of the BAD interval indicated that the 27 kb
25 BAD disease-associated chromosomal interval identified in the linkage disequilibrium studies is contained within an approximately 60 kb genomic region which contains a sequence referred to as GS4642 or rod photoreceptor protein (RPP) gene (Shimizu-Matsumoto, A. et al., 1997, Invest. Ophthalmol. Vis. Sci. 38:2576-2585).
30

7. **EXAMPLE: SEQUENCE AND CHARACTERIZATION
OF THE *HKNG1* GENE**

As demonstrated in the Example presented in Section 6, above, the *HKNG1* gene is involved in the neuropsychiatric disorder BAD. The results presented in this Section further characterize the *HKNG1* gene and gene product. In particular, isolation of additional cDNA clones and analyses of genomic and cDNA sequences have revealed both the full length *HKNG1* amino acid sequence and the *HKNG1* genomic intron/exon structure. In particular, the nucleotide and predicted amino acid sequence of the *HKNG1* gene identified by these analyses disclose new *HKNG1* exon sequences, including new *HKNG1* protein coding sequence, discovered herein. Further, the expression of *HKNG1* in human tissue, especially neural tissue, is characterized by Northern and in situ hybridization analysis. The results presented herein are consistent with the *HKNG1* gene being a gene which mediates neuropsychiatric disorders such as BAD.

7.1. **MATERIALS AND METHODS**

***HKNG1* cDNA Clone Isolation:** Hybridization of a human brain and kidney cDNA library was performed according to standard techniques and identified a full-length *HKNG1* cDNA clone. In addition, a *HKNG1* cDNA derived from a splice variant was isolated, as described in Section 7.2, below.

Northern Blot Analysis: Standard RNA isolation techniques and Northern blotting procedures were followed. The *HKNG1* probe utilized corresponds to the complementary sequence of base pairs 1367 to 1578 of the full length *HKNG1* cDNA sequence (SEQ ID NO. 1). Clontech multiple tissue northern blots were probed. In particular, Clontech human I, human II, human III, human fetal II, human brain II and human brain III blots were utilized for this study.

In Situ Hybridization Analysis: Standard in situ hybridization techniques were utilized. The *HKNG1* probe utilized corresponds to the complementary sequence of base pairs 910 to 1422 of the full length *HKNG1* cDNA sequence (SEQ ID NO. 1). Brains for in situ hybridization analysis were obtained from McLean Hospital (The Harvard Brain Tissue Resource Center, Belmont, MA 02178).

Other techniques: The remaining techniques described in Section 7.2, below, were performed according to standard techniques or as discussed in Section 6.1, above.

10

7.2. RESULTS

7.2.1. *HKNG1* Nucleotide and Amino Acid Sequence

A human brain cDNA library was screened and a full-length clone of *HKNG1* was isolated from this library, as described above. By comparing the isolated cDNA sequence to sequences in the public databases, a clone was identified which had been previously identified as GS4642, or rod photoreceptor protein (RPP) gene (GenBank Accession No. D63813; Shimizu-Matsumoto, A. et al., 1997, Invest. Ophthalmol. Vis. Sci. 38:2576-2585). Although Shimizu-Matsumoto et al. refer to GS4642 as a full-length cDNA sequence, the isolated *HKNG1* cDNA extends approximately 200 bp beyond the 5'end of the identified GS4642 clone.

Importantly, the *HKNG1* clone isolated herein reveals that, contrary to the amino acid sequence described in Shimizu-Matsumoto et al., the full length *HKNG1* amino acid sequence contains an additional 29 amino acid residues N-terminal to what had previously been identified as the full-length RPP (SEQ ID NO:64). The full-length *HKNG1* nucleotide sequence (SEQ ID NO: 1) and the derived amino acid sequence of the full-length *HKNG1* polypeptide (SEQ ID NO: 2) encoded by this sequence are depicted in FIG. 1A-1B.

The full-length *HKNG1* polypeptide was found to contain two clusterin similarity domains: clusterin similarity

domain 1 which corresponds to amino acid residues 134 to amino acid residue 160, and clusterin similarity domain 2 which corresponds to amino acid residue 334 to amino acid residue 362. Such clusterin domains are typically characterized by five shared cysteine residues. In clusterin
5 domain 1, these shared cysteine residues correspond to Cys 134, Cys145, Cys148, Cys158, and Cys 160. The shared cysteine residues in clusterin domain 2 correspond to the residues Cys334, Cys344, Cys351, Cys354, and Cys362.

Full-length *HKNG1* cDNA sequence was compared with the genomic contig completed by random sheared library
10 sequencing. Exon-intron boundaries were identified manually by aligning the two sequences in Sequencher 3.0 and by observing the conservative splicing sites where the alignments ended. This sequence comparison revealed that the additional cDNA sequence discovered through isolation of the full-length *HKNG1* cDNA clone actually belongs within three
15 *HKNG1* exons.

Prior to the isolation and analysis of *HKNG1* cDNA described herein, nine exons were predicted to be present within the corresponding genomic sequence. As discovered herein, however, the *HKNG1* gene, in contrast, actually
20 contains 13 exons, with the new cDNA containing sequence which corresponds to a new exon 1, exon 2 and a 5' extension of what had previously been designated exon 1. Splice variants, discussed in Section 9 below, also exist which comprise additional exons 2' and 2". The genomic sequence and intron/exon structure of the *HKNG1* gene is shown in FIG.
25 3A-3R.

The breakdown of exons was confirmed by the perfect alignment of the cDNA sequence with the genomic sequence and by observation of expected splicing sites flanking each of the additional, newly discovered exons.

HKNG1 nucleotide sequence was used to search databases
30 of partial sequences of cDNA clones. This search identified

a partial cDNA sequence derived from IMAGE clone R61493 having similarity to the human *HKNG1* sequence. IMAGE clone R61493 was obtained and consists of a cDNA insert, the Lafmid BA vector backbone, and DNA originating from the oligo dT primer and Hind III adaptors used in cDNA library
5 construction. The Lafmid BA vector nucleotide sequence is available at the URL http://image.rzpd.de/lafmida_seq.html and descriptions of the oligo dT primer and Hind III adaptors are available in the GENBANK record corresponding to accession number R61493.

10 The sequence of the cDNA insert revealed that the insert was derived from an alternatively spliced *HKNG1* mRNA variant, referred to herein as *HKNG1-V1*. In particular, this *HKNG1* variant is deleted for exon 3 of the full length 13 exon *HKNG1* sequence. The nucleotide sequence of this *HKNG1* variant (SEQ ID NO:3) is depicted in FIG. 2A-B. The amino
15 acid sequence encoded by the *HKNG1* variant (SEQ ID NO:3) is also shown in FIG. 2A-B.

Preferably therefore, the nucleic acids of the invention include nucleic acid molecules comprising the nucleotide sequence of *HKNG1-V1* or encoding the polypeptide encoded by
20 *HKNG1-V1* in the absence of heterologous sequences (e.g., cloning vector sequences such as Lafmid BA; oligo dT primer, and Hind III adaptor).

7.2.2. HKNG1 GENE EXPRESSION

HKNG1 gene expression was examined by Northern blot
25 analysis in various human tissues. A transcript of approximately 2 kb was detected in fetal brain, lung and kidney, and in adult brain, kidney, pancreas, prostate, testis, ovary, stomach, thyroid, spinal cord, lymph node and trachea. An approximately 1.5 kb transcript was also seen in trachea. In addition, a larger transcript of approximately 5
30 kb was detected in all adult neural regions tested (that is, cerebellum, cortex, medulla, spinal cord, occipital pole,

frontal lobe, temporal, putamen, amygdala, caudate nucleus, corpus callosum, hippocampus, whole brain, substantia nigra, subthalamic nucleus and thalamus). Once again, this is in direct contrast to previous Northern analysis of the *RPP* gene, which reported that expression was limited to the retina (Shimizu-Matsumoto, A. et al., 1997, Invest. Ophthalmol. Vis. Sci. 38:2576-2585).

Analysis of *HKNG1* the tissue distribution was extended through an in situ hybridization analysis. In particular, the *HKNG1* mRNA distribution in normal human brain tissue was analyzed. The results of this analysis are depicted in FIG. 4. As summarized in FIG. 4, *HKNG1* is expressed throughout the brain, with transcripts being localized to neuronal and grey matter cell types.

Finally, expression of *HKNG1* in recombinant cells demonstrates that the *HKNG1* gene encodes a secreted polypeptide(s).

8. A MISSENSE MUTATION WITHIN *HKNG1* CORRELATES WITH BAD

The Example presented in Section 6, above, shows that the BAD disorder maps to an interval completely contained within the *HKNG1* gene of the short arm of human chromosome 18. The Example presented in Section 7, above, characterizes the *HKNG1* gene and gene products. The results presented in this Example further these studies by identifying a mutation within the coding region of a *HKNG1* allele of an individual exhibiting a BAD disorder.

Thus, the results described herein demonstrate a positive correlation between a mutation which encodes a non-wild-type *HKNG1* polypeptide and the appearance of the neuropsychiatric disorder BAD. The results presented herein, coupled with the results presented in Section 6, above,

identify *HKNG1* as a gene which mediates neuropsychiatric disorders such as BAD.

8.1. MATERIALS AND METHODS

5 Pairs of PCR primers that flank each exon (see TABLE 1, above) were made and used to PCR amplify genomic DNA isolated from BAD affected and normal individuals. The amplified PCR products were analyzed using SSCP gel electrophoresis or by DNA sequencing. The DNA sequences and SSCP patterns of the affected and controls were compared and variations were
10 further analyzed.

8.2. RESULTS

In order to more definitively show that the *HKNG1* gene mediates neuropsychiatric disorders, in particular BAD, a study was conducted to explore whether a *HKNG1* mutation that
15 correlates with BAD could be identified.

First, exon scanning was performed on all eleven exons of the *HKNG1* gene using chromosomes isolated from three affected and one normal individual from the Costa Rican population utilized for the LD studies discussed in Section 6, above. No obvious mutations correlating with BAD were
20 found through this analysis.

Next, *HKNG1* intron and 3'-untranslated regions within the 27 kb BAD interval were scanned by sscp and/or sequencing for all variants among three affected and one normal individual from the same population. Approximately 60
25 variants were identified after scanning approximately two-thirds of the 27 kb genomic interval, which can be genotyped and analyzed by haplotype sharing and LD analyses, as described above, in order to identify ones which correlate with bipolar affective disorder. Fig. 5 lists selected variants identified through this study.

30 Exon scanning using chromosomal DNA from the general population of Costa Rica, however, successfully identified a

HKNG1 missense mutation in an individual affected with BAD who did not share the common diseased haplotype identified by the LD analysis provided above. In particular, exon scanning was done on exons 1-11 of *HKNG1* nucleic acid from 129 individuals from the general population affected with BAD.

This analysis identified a point mutation in the coding region of exon 7 not seen in non-bipolar affected disorder individuals. Specifically, the guanine corresponding to nucleotide residue 604 of SEQ ID NO:1 (or nucleotide residue 550 of SEQ ID NO:3) had mutated to an adenine. *HKNG1* protein expressed from this mutated *HKNG1* allele comprises the substitution of a lysine residue at amino acid residue 202 of SEQ ID NO:2 (or amino acid residue 184 of SEQ ID NO:4) in place of the wild-type glutamic acid residue.

Additional *HKNG1* polymorphisms relative to the *HKNG1* wild-type sequence, and which, therefore, represent *HKNG1* alleles, were identified through sequence analysis of the *HKNG1* alleles within a collection of schizophrenic patients of mixed ethnicity from the United States and within a BAD collection from the San Francisco area. These variants are depicted in FIGS. 5A and 5B, respectively. Statistical analysis indicated that there were significantly more variants in the collection of schizophrenic patients of mixed ethnicity from the United States and the San Francisco BAD and Costa Rican BAD samples than in a collection of 242 controls ($p < 0.05$).

9. **EXAMPLE: IDENTIFICATION OF ADDITIONAL HKNG1 SPLICE VARIANTS**

This example describes the isolation and identification of three novel splice variants of the human gene *HKNG1*. First, a novel *HKNG1* clone was isolated from a human retinal cDNA library. This clone, which completely lacks exon 7 of the full length *HKNG1* cDNA sequence, is referred to herein as *HKNG1Δ7*. Because the deletion of exon 7 from the full length

HKNG1 sequence leads to an immediate frameshift, the clone *HKNG1Δ7* encodes a truncated form of the *HKNG1* protein. The *HKNG1Δ7* cDNA sequence (SEQ ID NO:65) is depicted in FIG. 18 along with the predicted amino acid sequence (SEQ ID NO:66) of the *HKNG1Δ7* gene product it encodes.

Two other novel splice variants, referred to herein as *HKNG1-V2* and *HKNG1-V3*, were isolated and identified by using RT-PCR analysis to isolate additional *HKNG1* sequences. The following primer sequences were used:

10 5'-AGTTGCGTCCCTCTCTGTTG-3' (SEQ ID NO:67)
5'-GCTTCATGTTCCCGCTGTTA-3' (SEQ ID NO:68)

These splice variants included additional exons between exons 2 and 3 of the full length *HKNG1* sequence (SEQ ID NO:1).

15 The RT-PCR product derived from *HKNG1-V2* includes a novel exon referred to as "exon 2'", whereas the RT-PCR product derived from *HKNG1-V3* includes a novel exon referred to as "exon 2"". The sequence of these novel exons are provided in Table 2 below. The nucleotide sequence of the *HKNG1-V2* RT-PCR product containing novel exon 2' is depicted in FIG. 6A (SEQ ID NO:36), whereas the *HKNG1-V3* RT-PCR product containing novel exon 2" is depicted in FIG. 6B (SEQ ID NO:37). Both exon 2' and 2" are part of the 5'-untranslated region of the *HKNG1* cDNA.

TABLE 2

25	Exon 2'	5'-TTCCCTCCCTTTGGAACGCAGCGTGGGCACC (SEQ ID NO:34) TGCAACGCAGAGACCACTGTATCCCCGGTGCAGA ATGTAATGAGTGCCTGATACATTTGCCGAATAAA CTATTCCAAGGGTTGAACTTGCTGGAAGCAAGAG AAGCACTATTCTGG-3'
----	---------	--

30

Exon 2"	5'-ATGGAGTCTTGCTCTCGTTGCCAGACTGGA	(SEQ DI NO:35)
	GTGCACTGCTGCGATCTCAGCTCACTGCAACCTC	
	TACCTCCCAGGTTCAAGCGATTCTCCTGCCTCAG	
	CCTCTCGAGTGGCTGGGACTATAG-3'	

5

10. EXAMPLE: IDENTIFICATION OF HKNG1 ORTHOLOGS

This example describes the isolation and characterization of genes in other mammalian species which are orthologs to human *HKNG1*. Specifically, both guinea pig and bovine *HKNG1* sequences are described.

10

10.1. GUINEA PIG HKNG1 ORTHOLOGS

A guinea pig *HKNG1* ortholog, referred to as *gphkng1815*, was isolated using RT-PCR. The cDNA sequence (SEQ ID NO:38) and predicted amino acid sequence (SEQ ID NO:39) are depicted in FIG. 7. Both the nucleotide and the predicted amino acid sequence of *gphkng 1815* are similar to the human *HKNG1* nucleotide and amino acid sequences. Specifically, the program ALIGNv2.0 identified a 71.5% nucleotide sequence identity and a 62.8% amino acid sequence identity using standard parameters (Scoring Matrix: PAM120; GAP penalties: -12/-4).

20

Like the human *HKNG1* polypeptide, the predicted *gphkng 1815* polypeptide also contains two clusterin similarity domains, which correspond to amino acid residues 105 to 131 (clusterin domain 1), and amino acid residues 305-333 (clusterin domain 2), respectively. Both of these domains contain the five conserved cysteine residues typically associated with clusterin domains. Specifically, these conserved cysteines correspond to Cys105, Cys116, Cys119, Cys124 and Cys131 (clusterin similarity domain 1) and Cys305, Cys315, Cys322, Cys325, and Cys333 (clusterin similarity domain 1) of the *gphkng 1815* polypeptide sequence.

25

Three allelic variants of *gphkng 1815*, referred to as *gphkng 7b*, *gphkng 7c*, and *gphkng 7d*, respectively, were also

30

identified by RT-PCR. Their nucleotide [SEQ ID NO:40 (*gphkng 7b*), SEQ ID NO:42 (*gphkng 7c*), and SEQ ID NO:44 (*gphkng 7d*)] and amino acid [SEQ ID NO:41 (*gphkng 7b*), SEQ ID NO:43 (*gphkng 7c*), and SEQ ID NO:45 (*gphkng 7d*)] sequences
5 are depicted in FIGS. 8 through 10, respectively. Each of these three allelic variants contains a deletion within a region homologous to exon 7 of human *HKNG1*. The allelic variants retain the open reading frame of the gene, however, each allelic variant contains a deletion, relative to *gphkng 1815*, of 16, 92, and 93 amino acid residues, respectively.

10 An alignment of the predicted amino acid sequences of *gphkng1815*, *gphkng 7b*, *gphkng 7c*, and *gphkng7d* is shown in FIG. 14. An alignment of the predicted amino acid sequences of the human *HKNG1* gene product, the guinea pig *HKNG1* ortholog *gphkng1815*, and the bovine *HKNG1* ortholog described
15 in Subsection 10.2 below are shown in FIG. 16.

10.2. BOVINE HKNG1 ORTHOLOGS

Bovine orthologs of *HKNG1* were also cloned by screening a cDNA library made from pooled bovine retinal tissue using a
20 nucleotide sequence that corresponded to the complementary sequence of base pairs 910-1422 of the full length human *HKNG1* cDNA sequence (SEQ ID NO:1) as a probe. Three independent bovine cDNA species, referred to as *bhkng1*, *bhkng2*, and *bhkng3* (SEQ ID NOS: 46 to 48, respectively) were isolated. Each of these allelic variants contains several single nucleotide polymorphisms (SNPs). None of the SNPs
25 results in an altered predicted amino acid sequence. Thus all three bovine cDNAs encodes the same predicted amino acid sequence (SEQ ID NO:49). These SNPs apparently reflect the natural allelic variation of the pooled cDNA library from which the sequences were isolated. Each of the three bovine *HKNG1* allelic variants is depicted in FIGS. 11 to 13,
30 respectively, along with the predicted amino acid sequence which they encode.

The predicted bovine HKNG1 polypeptide also contains two clusterin similarity domains, corresponding to amino acid residues 105-131 and amino acid residues 304-332, respectively, of SEQ ID NO:49. Clusterin domain 1 contains the five shared cysteine amino acid residues typically
 5 associated with this type of domain: Cys105, Cys116, Cys119, Cys124, and Cys131. Clusterin domain 2 of the bovine HKNG1 polypeptide contains four conserved cystein residues: Cys314, Cys321, Cys324, and Cys332.

11. EXPRESSION OF HUMAN HKNG1 GENE PRODUCT

10 Described in this example is the construction of expression vectors and the successful expression of recombinant human HKNG1 sequences. Expression vectors are described both for native HKNG1 and for various HKNG1 fusion proteins.

15 11.1. EXPRESSION OF HUMAN HKNG1:FLAG

A human HKNG1 flag epitope-tagged protein (HKNG1:flag) vector was constructed by PCR followed by ligation into an vector for expression in HEK 293T cells. The full open-reading frame of the full length HKNG1 cDNA sequence (SEQ ID
 20 NO:5) was PCR amplified using the following primer sequences:

5' primer 5'-TTTTTCTGAATTCGCCACCATGAAAATTA (SEQ ID NO:52)
 AAGCAGAGAAAAACG-3'

3' primer 5'-TTTTTGTCGACTTATCACTTGTCGTCGTC (SEQ ID NO:53)
 GTCCTTGTAGTCCCAGGTTTAAATGTTCTT
 TAAATGC-3'

the 5' primer incorporating a Kozak sequence upstream of and
 25 including the upstream initiator methionine and the 3' primer including the nucleotide sequence encoding the flag epitope DYKDDDDK (SEQ ID NO:50) followed by a termination codon.

The sequenced DNA construct was transiently transfected into HEK 293T cells in 150 mm plates using Lipofectamine (GIBCO/BRL) according to the manufacturer's protocol.
 30 Seventy-two hours post-transfection, the serum-free conditioned medium (OptiMEM, GIBCO/BRL) was harvested and

spun and the remaining monolayer of cells was lysed using 2 mL of lysis buffer [50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 0.05% SDS with "Complete" protease cocktail (Boehringer Mannheim) diluted according to manufacturers instructions]. Insoluble material was pelleted before
5 preparation of SDS-PAGE samples.

Conditioned medium was electroblotted onto a PVDF membrane (Novex) after separation by SDS-PAGE on 4-20% gradient gels and probed with an M2 anti-flag polyclonal antibody (1:500, Sigma) followed by horseradish peroxidase (HRP) conjugated sheep anti-mouse antibody (1:5000,
10 Amersham), developed using chemiluminescent reagents (Renaissance, Dupont), and exposed to autoradiography film (Biomax MR2 film, Kodak). Flag immunoreactivity appeared as a doublet of bands that migrated by SDS-PAGE between 60 and 95 kDa as determined by Multimark molecular weight markers (Novex), demonstrating secretion of the HKNG1:Flag protein.
15 The double band indicates at least two different species with different mobilities on SDS-PAGE. Such doublets most commonly arise with posttranslational modifications to the protein, such as glycosylation and/or proteolysis. Treatment of the PNGase F (Oxford Glycosciences) according to the manufacturer's directions resulted in a single band of
20 increased mobility, indicating that two original bands contain N-linked carbohydrate. When run in the absence of a reducing agent, the relative mobility of the immunoreactive bands was greater than 100 kDa relative to the same markers, indicating that HKNG1:flag fusion proteins may be a disulfide linked dimer or higher oligomer.

25

11.2. EXPRESSION OF HUMAN HKNG1-V1:FLAG

A human HKNG1-V1 flag epitope-tagged protein (HKNG1-V1:flag) vector was also constructed by PCR followed by ligation into an expression vector, pMET stop. The full
30 length open-reading frame of the HKNG1-V1 cDNA sequence (SEQ

ID NO:6) was PCR amplified using the following primer sequences:

5' primer 5'-TTTTTCTGAATTCACCATGAGGACCTGGG (SEQ ID NO:54)
ACTACAGTAAC-3'

5 3' primer 5'-TTTTTGTCGACTTATCACTTGTCGTCGTC (SEQ ID NO:53)
GTCCTGTAGTCCCAGGTTTAAATGTTTCCT
TAAATGC-3'

The 5' primer incorporated a Kozak sequence upstream of and including the upstream initiator methionine. The 3' primer included the nucleotide sequence encoding the flag epitope DYKDDDDK (SEQ ID NO:50) followed by a termination codon.

10 The sequenced DNA construct was transiently transfected into HEK 293T cells in 150 mm plates using Lipofectamine (GIBCO/BRL) according to the manufacturer's protocol. Seventy-two hours post-transfection, the serum-free conditioned medium (OptiMEM, GIBCO/BRL) was harvested and spun and the remaining monolayer of cells was lysed using
15 2 mL of lysis buffer [50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 0.05% SDS with "Complete" protease cocktail (Boehringer Mannheim) diluted according to manufacturers instructions]. Insoluble material was pelleted before preparation of SDS-PAGE samples.

20 Conditioned medium was electroblotted onto a PVDF membrane (Novex) after separation by SDS-PAGE on 4-20% gradient gels and probed with an M2 anti-flag polyclonal antibody (1:500, Sigma) followed by horseradish peroxidase (HRP) conjugated sheep anti-mouse antibody (1:5000, Amersham), developed using chemiluminescent reagents (Renaissance, Dupont), and exposed to autoradiography film
25 (Biomax MR2 film, Kodak). Flag immunoreactivity appeared as a doublet of bands that migrated by SDS-PAGE between 60 and 95 kDa as determined by Multimark molecular weight markers (Novex), demonstrating secretion of the HKNG1:Flag protein. When run in the absence of reducing agent, the relative mobility of the immunoreactive bands was greater than 100 kDa
30 relative to the same markers, suggesting that the HKNG1-

V1:flag fusion protein may be a disulfide linked dimer or higher oligomer.

11.3. EXPRESSION OF HUMAN HKNG1:Fc

5 A human HKNG1/hIgG1Fc fusion protein vector was constructed by PCR. The full-length open-reading frame of the full length HKNG1 cDNA (SEQ ID NO:5) was PCR amplified using the following primer sequences:

5' primer 5'-TTTTTCTCTCGAGACCATGAAAATTAAAG (SEQ ID NO:55)
CAGAGAAAACG-3'

10 3' primer 5'-TTTTTGGATCCGCTGCTGCCAGGTTTAA (SEQ ID NO:56)
AAATGTTTCCTTAAAATGC-3'

The 5' primer incorporated a Kozak sequence before the upstream methionine to the amino acid residue before the stop codon. The 3' PCR primer contained a 3 alanine linker at the junction of HKNG1 and the human IgG1 Fc domain, which starts
15 at residues DPE. The genomic sequence of the human IgG1 Fc domain was ligated along with the PCR product into a pCDM8 vector (Invitrogen, Carlsbad CA) for transient expression.

The sequenced DNA construct was transiently transfected into HEK 293T cells in 150 mm plates using Lipofectamine (GIBCO/BRL) according to the manufacturer's protocol.
20 Seventy-two hours post-transfection, the serum-free conditioned medium (OptiMEM, GIBCO/BRL) was harvested and spun and the remaining monolayer of cells was lysed using 2 mL of lysis buffer [50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 0.05% SDS with "Complete" protease cocktail (Boehringer Mannheim) diluted according to manufacturers
25 instructions]. Insoluble material was pelleted before preparation of SDS-PAGE samples.

Conditioned medium was electroblotted onto a PVDF membrane (Novex) after separation by SDS-PAGE on 4-20% gradient gels and probed with an M2 anti-Fc polyclonal antibody (1:500, Jackson ImmunoResearch Laboratories, Inc.)
30 followed by horseradish peroxidase (HRP) conjugated sheep anti-mouse antibody (1:5000, Amersham), developed using

chemiluminescent reagents (Renaissance, Dupont), and exposed to autoradiography film (Biomax MR2 film, Kodak). Human IgG1 Fc immunoreactivity appeared as a doublet of bands that migrated by SDS-PAGE between 148 and 60 kDa standards of the Multimark molecular weight markers (Novex), demonstrating
 5 secretion of the HKNG1:Fc fusion protein.

11.4. EXPRESSION OF HUMAN HKNG1-V1:Fc

A human HKNG1-V1/hIgG1Fc fusion protein (HKNG1-V1:Fc) vector was also constructed by PCR. The full-length open reading frame of HKNG1-V1 cDNA (SEQ ID NO:6) was PCR
 10 amplified using the following primer sequences:

5' primer 5'-TTTTTCTCTCGAGACCATGAGGACCTGGG (SEQ ID NO:57)
 ACTACAGTAAC-3'

3' primer 5'-TTTTTGGATCCGCTGCTGCCAGGTTTAA (SEQ ID NO:56)
 AAATGTTCTTAAATGC-3'

15 The 5' primer incorporated a Kozak sequence before the upstream methionine to the amino acid residue before the stop codon. The 3' PCR primer contained a 3 alanine linker at the junction of HKNG1-V1 and the human IgG1 Fc domain, which starts at residues DPE. The genomic sequence of the human IgG1 Fc domain was ligated along with the PCR product into a
 20 pCDM8 vector for transient expression.

The sequenced DNA construct was transiently transfected into HEK 293T cells in 150 mm plates using Lipofectamine (GIBCO/BRL) according to the manufacturer's protocol. Seventy-two hours post-transfection, the serum-free conditioned medium (OptiMEM, GIBCO/BRL) was harvested and
 25 spun and the remaining monolayer of cells was lysed using 2 mL of lysis buffer [50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 0.05% SDS with "Complete" protease cocktail (Boehringer Mannheim) diluted according to manufacturers instructions]. Insoluble material was pelleted before preparation of SDS-PAGE samples.

30 Conditioned medium was electroblotted onto a PVDF membrane (Novex) after separation by SDS-PAGE on 4-20%

gradient gels and probed with an anti-human Fc polyclonal antibody (1:500, Jackson ImmunoResearch Laboratories, Inc.) followed by horseradish peroxidase (HRP) conjugated sheep anti-mouse antibody (1:5000, Amersham), developed using chemiluminescent reagents (Renaissance, Dupont), and exposed
 5 to autoradiography film (Biomax MR2 film, Kodak). Human IgG1 Fc immunoreactivity appeared as a doublet of bands that migrated by SDS-PAGE between 148 and 60 kDa standards of the Multimark molecular weight markers (Novex) centered approximately between 125 and 150 kDa, demonstrating secretion mediated by the HKNG1 signal peptide.

10

11.5. EXPRESSION OF HUMAN HKNG1 Δ 7:Fc

A human HKNG1 Δ 7:hIgG1Fc fusion protein vector was also constructed by PCR. The sequence of the HKNG1 Δ 7 splice variant was amplified by PCR amplification using Exons 1 through 6 of the full length HKNG1 cDNA sequence (SEQ ID
 15 NO:1) as a template with the following primer sequences:

5' primer 5'-TTTTTCTGAATTCACCATGAAGCCGCCAC (SEQ ID NO:58)
 TCTTGGTG-3'

3' primer 5'-TTTTTGGATCCGCTGCGGCCTCCGTG (SEQ ID NO:59)
 GTCAGGAGCTTATTTTTCACAGAGGACCAGCT
 AG-3'

20 The 5' primer incorporated a Kozak sequence upstream of and including the upstream initiator methionine. The 3' primer included the first 17 (coding) nucleotides of exon 8 followed by nucleotides encoding a 3 alanine linker.

The genomic sequence of the human IgG1 Fc domain was ligated along with the PCR product into a pCDM8 vector for
 25 transient expression.

The sequenced DNA construct was transiently transfected into HEK 293T cells in 150 mm plates using Lipofectamine (GIBCO/BRL) according to the manufacturer's protocol. Seventy-two hours post-transfection, the serum-free conditioned medium (OptiMEM, GIBCO/BRL) was harvested and
 30 spun and the remaining monolayer of cells was lysed using

2 mL of lysis buffer [50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 0.05% SDS with "Complete" protease cocktail (Boehringer Mannheim) diluted according to manufacturers instructions]. Insoluble material was pelleted before preparation of SDS-PAGE samples.

5 Conditioned medium was electroblotted onto a PVDF membrane (Novex) after separation by SDS-PAGE on 4-20% gradient gels and probed with an anti-human Fc polyclonal antibody (1:500, Jackson ImmunoResearch Laboratories) followed by horseradish peroxidase (HRP) conjugated sheep anti-mouse antibody (1:5000, Amersham), developed using
 10 chemiluminescent reagents (Renaissance, Dupont), and exposed to autoradiography film (Biomax MR2 film, Kodak). Human IgG1 Fc immunoreactivity appeared as a band that migrated by SDS-PAGE between 42 and 60 kDa relative to Multimark molecular weight markers (Novex) centered approximately between 36.5 and 55.4 kDa relative to Mark 12 molecular weight markers
 15 (Novex).

11.6. EXPRESSION OF NATIVE HUMAN HKNG1

A human HKNG1 expression vector was constructed by PCR amplification of the human HKNG1 cDNA sequence (SEQ ID NO:1) followed by ligation into an expression vector, pcDNA3.1
 20 (Invitrogen, Carlsbad CA). The full open-reading frame of the HKNG1 cDNA sequence (SEQ ID NO:5) was PCR amplified using the following primer sequences:

5' primer 5'-TTTTTCTCTCGAGGACTACAGGACACAGC (SEQ ID NO:60)
 TAAATCC-3'

25 3' primer 5'-TTTTTGGATCCTTATCACCAGGTTTAA (SEQ ID NO:61)
 ATGTCCTTAAAATGC-3'

The 5' primer incorporated a Kozak sequence upstream of and including the upstream initiator methionine. The 3' primer included a tandem pair of termination codons.

The sequenced DNA construct was transiently transfected into HEK 293T cells in 150 mm plates using Lipofectamine
 30 (GIBCO/BRL) according to the manufacturer's protocol. Seventy-two hours post-transfection, the serum-free

conditioned medium (OptiMEM, GIBCO/BRL) was harvested and spun and the remaining monolayer of cells was lysed using 2 mL of lysis buffer [50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 0.05% SDS with "Complete" protease cocktail (Boehringer Mannheim) diluted according to manufacturers instructions]. Insoluble material was pelleted before preparation of SDS-PAGE samples.

Conditioned medium was electroblotted onto a PVDF membrane (Novex) after separation by SDS-PAGE on 4-20% gradient gels and probed with an anti-HKNG1 polyclonal antibody (#84, 1:500) followed by horseradish peroxidase (HRP) conjugated sheep anti-mouse antibody (1:5000, Amersham), developed using chemiluminescent reagents (Renaissance, Dupont), and exposed to autoradiography film (Biomax MR2 film, Kodak). HKNG1 immunoreactivity appeared as a doublet of bands that migrated by SDS-PAGE between 60 and 95 kDa as determined by Multimark molecular weight markers (Novex).

11.7. EXPRESSION OF NATIVE HUMAN HKNG1-V1

A human HKNG1-V1 expression vector was also constructed by PCR amplification of the human HKNG1-V1 cDNA sequence (SEQ ID NO:3) followed by ligation into an expression vector, pcDNA3.1. The full open-reading frame of the HKNG1 cDNA sequence (SEQ ID NO:6) was PCR amplified using the following primer sequences:

5' primer 5'-TTTTTCTGAATTCACCATGAAGCCGCCAC (SEQ ID NO:62)
TCTTGGTG-3'

5' primer 5'-TTTTTCTCTCGAGACCATGAGGACCTGGG (SEQ ID NO:63)
ACTACAGTAAC-3'

3' primer 5'-TTTTTGGATCCTTATCACCAGGTTTAA (SEQ ID NO:61)
ATGTTTCCTTAAAATGC-3'

The 5' primer incorporated a Kozak sequence upstream of and including the upstream initiator methionine. The 3' primer included a tandem pair of termination codons.

The sequenced DNA construct was transiently transfected into HEK 293T cells in 150 mm plates using Lipofectamine

(GIBCO/BRL) according to the manufacturer's protocol. Seventy-two hours post-transfection, the serum-free conditioned medium (OptiMEM, GIBCO/BRL) was harvested and spun and the remaining monolayer of cells was lysed using 2 mL of lysis buffer [50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 0.05% SDS with "Complete" protease cocktail (Boehringer Mannheim) diluted according to manufacturers instructions]. Insoluble material was pelleted before preparation of SDS-PAGE samples.

Conditioned medium was electroblotted onto a PVDF membrane (Novex) after separation by SDS-PAGE on 4-20% gradient gels and probed with an M2 anti-HKNG1 polyclonal antibody (#84, 1:500) followed by horseradish peroxidase (HRP) conjugated sheep anti-mouse antibody (1:5000, Amersham), developed using chemiluminescent reagents (Renaissance, Dupont), and exposed to autoradiography film (Biomax MR2 film, Kodak). HKNG immunoreactivity appeared as a doublet of bands that migrated by SDS-PAGE between 70 and 95 kDa as determined by Multimark molecular weight markers (Novex), demonstrating secretion mediated by the HKNG1 signal peptide.

11.8. EXPRESSION OF HUMAN HKNG:AP FUSION PROTEINS

Expression vectors were also constructed for human HKNG1 alkaline phosphatase C-terminal fusion protein (HKNG1:AP), human HKNG1-V1 alkaline phosphatase C-terminal fusion protein (HKNG1-V1:AP), and human HKNG1 alkaline phosphatase N-terminal fusion protein (AP:HKNG1).

The expression vector for human HKNG1:AP was constructed by PCR amplification followed by ligation into a vector for suitable for expression in HEK 293T cells. The full-length open-reading frame of human HKNG1 (SEQ ID NO:5) was PCR amplified using a 5' primer incorporating an EcoRI restriction site followed by a Kozak sequence prior to the upstream initiator methionine. The 3' primer included a XhoI restriction site immediately following the final codon of

HKNG1. Thus, the open reading frame of the construct includes the *HKNG1* signal peptide and the full *HKNG1* sequence followed by the full sequence of human placental alkaline phosphatase.

5 The expression vector for human *HKNG1-V1:AP* was constructed by PCR amplification followed by ligation into *pMEAP3* vector. The full length open reading frame of human *HKNG1-V1* (SEQ ID NO:6) was PCR amplified using a 5' primer incorporating an *EcoRI* restriction site followed by a Kozak sequence prior to the upstream initiator methionine. The 3'
10 primer included a *XhoI* restriction site immediately following the final codon of *HKNG1-V1*. Thus, the open reading frame of the construct includes the *HKNG1-V1* signal and the full length *HKNG1-V1* sequence followed by the full sequence of human placental alkaline phosphatase.

The expression vector for human *AP:HKNG1* was constructed
15 by PCR amplification followed by ligation into the *AP-Tag3* vector reported by Cheng and Flanagan, 1994, *Cell* 79:157-168. The full-length open-reading frame of human *HKNG1* (SEQ ID NO:5) was PCR amplified using a 5' primer incorporating a *BamHI* restriction site prior to the nucleotides encoding the first amino acids (*i.e.*, *APT*) of the mature *HKNG* protein, and
20 a 3' primer that included a *XhoI* restriction site immediately following the termination codon of *HKNG1*. Thus, the open reading frame of the complete construct includes the *AP* signal peptide and the full sequence of human placental alkaline phosphatase, followed by the full *HKNG1* sequence.

The sequenced DNA constructs were transiently
25 transfected in HEK 293T cells in 150 mM plates using Lipofectamine (GIBCO/BRL) according to the manufacturer's protocol. 72 hours post-transfection, the serum-free conditioned media (OptiMEM, Gibco/BRL) were harvested, spun and filtered. Alkaline phosphatase activity in the conditioned media was quantitated using an enzymatic assay
30 kit (Phospha-Light, Tropix) according to the manufacturer's instructions. When alkaline phosphatase fusion protein

concentrations below 2 nM were observed, conditioned medium was concentrated by centrifugation using a 30 kDa cut-off membrane. Conditioned medium samples before and after concentration were analyzed by SDS-PAGE followed by Western blot using anti-human alkaline phosphatase antibodies (1:250, 5 Genzyme) and chemiluminiscence detection. A band at 140 kDa was observed in concentrated supernatant of HKNG1:AP, HKNG1-V1:AP, and AP:HKNG1 transfections. Conditioned medium samples were adjusted to 10% fetal calf serum and stored at 4°C.

10 **11.9. PURIFICATION OF FLAG-TAGGED HKNG1 PROTEINS**

The secreted flag-tagged proteins described in subsections 12.1 and 12.2 above were isolated by a one step purification scheme utilizing the affinity of the flag epitope to M2 anti-flag antibodies. The conditioned media was passed over an M2-biotin (Sigma)/streptavidin Poros 15 column (2.1 x 30 mm, PE Biosystems). The column was then washed with PBS, pH 7.4, and flag-tagged protein was eluted with 200 mM glycine, pH 3.0. Fraction was neutralized with 1.0 M Tris pH 8.0. Eluted fractions with 280 nm absorbance greater than background were then analyzed on SDS-PAGE gels and by Western blot. The fractions containing flag tagged 20 protein were pooled and dialyzed in 8000 MWCO dialysis tubing against 2 changes of 4L PBS, pH 7.4 at 4°C with constant stirring. The buffered exchanged material was then sterile filtered (0.2 µm, Millipore) and frozen at -80°C.

25 **11.10. PURIFICATION OF HKNG1 Fc FUSION PROTEINS**

The secreted Fc fusion proteins described in Subsections 12.3-12.5 above were isolated by a one step purification scheme utilizing the affinity of the human IgG1 Fc domain to Protein A. The conditioned media was passed over a POROS A column (4.6 x 100 mm, PerSeptive Biosystems); the column was then washed with PBS, pH 7.4 and eluted with 200 mM glycine, 30 pH 3.0. Fractions were neutralized with 1.0 M Tris pH 8.0.

A constant flow rate of 7 ml/min was maintained throughout the procedure. Eluted fractions with 280 nm absorbance greater than background were then analyzed on SDS-PAGE gels and by Western blot. The fractions containing Fc fusion protein were pooled and dialyzed in 8000 MWCO dialysis tubing
 5 against 2 changes of 4L PBS, pH 7.4 at 4°C with constant stirring. The buffered exchanged material was then sterile filtered (0.2 µm, Millipore) and frozen at -80°C.

12. PRODUCTION OF ANTI-HKNG1 ANTIBODIES

10 Described in the example presented in this Section is the production and characterization of polyclonal and monoclonal antibodies directed against HKNG1 proteins.

12.1. PRODUCTION OF POLYCLONAL ANTIBODIES

15 Polyclonal antisera were raised in rabbits against each of the three peptides listed in Table 3 below. Each of the peptides was derived from the HKNG1 amino acid sequence (SEQ ID NO:2) by standard techniques (see, in particular, Harlow&Lane, 1988, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, the contents of which is incorporated herein by reference in its entirety). Each of
 20 the peptides is also represented in the HKNG1-V1 polypeptide sequence (SEQ ID NO:4). Antisera was subsequently affinity purified using the peptide immunogens.

TABLE 3

25	Antibody	Peptide/Immunogen	a.a. residues (SEQ ID NO:2)
	Antibody 84	APTWKDKTAISENLK	50-64
	Antibody 85	KAIEDLPKQDK	304-314
	Antibody 86	KALQHFKEHFKTW	483-495

30 12.2. PRODUCTION OF MONOCLONAL ANTIBODIES

Monoclonal antibodies were raised in mice by standard techniques (see, Harlow & Lane, *supra*) against the HKNG-Fc fusion protein described in Section 11.3 above. Wells were screened by ELISA for binding to the HKNG-Fc fusion protein. Those wells reacting with the Fc protein were identified by ELISA for binding to an irrelevant Fc fusion protein and discarded. HKNG-Fc specific wells were tested for their ability to immunoprecipitate HKNG-Fc and subjected to isotype analysis by standard techniques (Harlow & Lane, *supra*), and eight wells were selected for subcloning. The isotype of the subcloned monoclonal antibodies was confirmed and is presented in Table 4 below.

Based on Western blotting, immunoprecipitation and immunostaining data discussed in SubSection 12.3 below, two monoclonal antibodies (3D17 and 4N6) were selected for large scale production.

TABLE 4

Clone	Isotype
1F24	2a
1J18	2a
2020	1
3D17	1
3D24	2a
4N6	1
4O16	2b
10C6	2a

12.3. WESTERN BLOTTING AND IMMUNOPRECIPITATION OF RECOMBINANT HKNG PROTEIN

The polyclonal antisera and all eight monoclonal antibodies described in subsection 12.1 and 12.2 above were tested for their ability to recognize recombinant HKNG1 proteins on Western blots using standard techniques (see, in

particular, Harlow & Lane, 1988, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press). Polyclonal antisera 84 and 85 and monoclonal antibodies 3D17 and 4N6 were able to recognize all forms of the mature (*i.e.*,
 5 secreted) recombinant HKNG proteins tested (*i.e.*, HKNG1:Fc, HKNG1:flag, AP:HKNG1, and native HKNG1) in Western blots.

Table 5 indicates the ability of each monoclonal antibody to immunoprecipitate recombinant HKNG1, as assessed by Western blotting of immunoprecipitates with the polyclonal antisera 84 and 85. None of the polyclonal antisera were
 10 able to immunoprecipitate recombinant HKNG1 proteins. All eight monoclonal antibodies immunoprecipitated HKNG1:Fc. Immunoprecipitation of the other recombinant HKNG1 proteins was variable.

15

TABLE 5

Monoclonal Antibody	Protein			
	HKNG1:Fc	HKNG1:flag	AP:HKNG1	HKNG1 (native)
IF24	+	+	+	-/+
20 1J18	+	-	-/+	+/+
2020	+	-	+	-
3D17	+/+	+/+	-	+/+
3D24	+	-	-	-
4N6	+	+	+	+
25 4016	+	-	-	+/+
10C6	+	-	-	+

13. CONFIRMATION OF THE HKNG N-TERMINUS AND DISULFIDE BOND STRUCTURE

30 The experiments described in this section provide data identifying the N-terminus of the mature secreted human HKNG

protein. The experiments also provide data identifying the disulfide bond linkages between cysteine amino acid residues in the mature, secreted protein.

Specifically, mature, secreted HKNG:flag, HKNG, and HKNG:Fc recombinant proteins were produced and purified as
5 described in Section 11 above. The mature recombinant proteins were digested with trypsin, and the tryptic fragments were identified and sequenced using reverse-phase liquid chromatography coupled with electrospray ionization tandem mass spectrometry (LC/MS/MS). The N-terminus of all mature secreted proteins tested was unambiguously identified
10 as APTWKDKT, which corresponds to the amino acid sequence starting at alanine 50 of the HKNG1 amino acid sequence (FIG. 1; SEQ ID NO:2) or alanine 32 of the HKNG1-V1 amino acid sequence (FIG. 2; SEQ ID NO:4). Thus, although the cDNA sequences of HKNG1 and HKNG1-V1 encode distinct amino acid sequences, the mature secreted proteins produced by these two
15 splice variants of the human *HKNG1* gene are identical, since the alternative splicing that gives rise to HKNG1-V1 (*i.e.*, the deletion of exon 3) affects the amino acid sequence of the proteolytically cleaved signal peptide. The amino acid sequence of the mature secreted HKNG1 protein is shown in
20 FIG. 17 (SEQ ID NO:51)

The mature secreted HKNG protein is also distinct from the RPP amino acid sequence disclosed by Shimizu-Matsumo *et al.* (1997, *Invest. Ophthalmol. Vis. Sci.* 38:2576-2585). In particular, amino acid residues 1 to 20 of the RPP amino acid sequence disclosed in Figure 3 of Shimizu-Matsumo *et al.*,
25 *supra*, correspond to the cleaved signal peptide of HKNG1-V1. The amino acid sequence of the mature secreted form of the *HKNG1* gene product is depicted in FIG. 17 (SEQ ID NO:51).

Disulfide bond linkages for 8 of the 13 cysteine
30 residues in the mature, secreted HKNG protein were also identified from LC/MS/MS of peptides recovered from tryptic

digestion of the unreduced protein. In particular, the following disulfide bonded pairs of cysteines were identified (numbering refers to the HKNG1 protein shown in FIG. 1; SEQ ID NO:2):

- 5 Cys 134 to Cys 145;
 Cys 148 to Cys 153;
 Cys 160 to Cys 334; and
 Cys 354 to Cys 362.

14. **EXAMPLE: LOCALIZATION OF HKNG mRNA
 AND PROTEIN EXPRESSION**

10 This example describes experiments wherein the HKNG gene product is shown to be expressed in human brain and retinal tissue. Specifically, *in situ* hybridization experiments performed using standard techniques with a probe that corresponded to the complementary sequence of base pairs 910-1422 of the full length *HKNG1* cDNA sequence (SEQ ID NO:1)
 15 detected HKNG messenger RNA in the photoreceptor layer (outer nuclear layer) of human retina in eyes obtained from the New England Eye Bank.

The polyclonal antisera and all eight monoclonal antibodies described in Section 12 above were tested for
 20 immunostaining of human retina. Polyclonal antiserum 85 and monoclonal antibodies 1F24, 4N6 and 4O16 showed immunostaining of HKNG protein in the photoreceptor layer and adjacent layers of the retina. The immunostaining in these tissues with polyclonal antiserum was blocked by 85 peptide immunogen, but not by the other two peptide immunogens (*i.e.*,
 25 84 and 86), confirming that the immunostaining was due to HKNG protein expressed in the photoreceptor layer.

The same antibodies were then used to localize HKNG protein by immunostaining in sections of human and monkey brain. HKNG protein was observed in cortical neurons in the frontal cortex. The majority of pyramidal neurons in layers
 30 IV-V were immunoreactive for HKNG protein. A subpopulation of neurons was also labeled in layers I-III. HKNG

Both the full length *HKNG1* cDNA sequence shown in FIG. 1 (SEQ ID NO:1) and the alternatively spliced *HKNG1-V1* cDNA sequence shown in FIG. 2 (SEQ ID NO:3) encode predicted proteins that have methionines in close proximity to their predicted initiator methionines. The predicted protein sequence encoded by the full length *HKNG1* cDNA sequence has a second methionine at amino acid residue number 30 of the amino acid sequence depicted in FIG. 1 (SEQ ID NO:2). Thus, although FIG. 1 indicates that the full length *HKNG1* cDNA encodes the first immature form of the *HKNG1* protein depicted in FIG. 1 (referred to herein as IPF1), the full length *HKNG1* cDNA may additionally encode a second immature protein form (referred to herein as IPF2), whose sequence (SEQ ID NO:64) is provided on the third line of the protein alignment depicted in FIG. 17. IPF2 is initiated at methionine 30 of the IPF1 protein sequence, and is identical to the RPP polypeptide sequence taught by Shimizu-Matsumoto et al (1997, *Invest. Ophthalmol. Vis. Sci.* 38:2576-2585). Likewise, the alternatively spliced *HKNG1-V1* cDNA sequence encodes the predicted immature protein form, referred to herein as IPF3, depicted in FIG. 2 (SEQ ID NO:4). However, the *HKNG1-V1* cDNA may also encoded another immature protein form, identical to IPF 2, that is initiated at methionine 12 of the IPF3 protein sequence. FIG. 17 illustrates an alignment of the three immature *HKNG1* protein sequences IPF1 (second row), IPF2 (third row), and IPF3 (bottom row). As explained in Section 13 above, the mature *HKNG1* gene product secreted by cells expressing the *HKNG1* constructs described in Section 11, above, is in fact the same cleaved product (SEQ ID NO:51), regardless of the immature *HKNG1* protein (IPF1, IPF2, or IPF3) from which it is produced. An alignment of the mature secreted *HKNG1* protein is therefore also depicted in FIG. 17 (top row).

Modified *HKNG1:flag* and *HKNG1-V1:flag* expression vectors were constructed as described in Sections 12.1 and 12.2,

respectively. However, the nucleotide sequence of full length HKNG1 was modified, using standard site directed mutagenesis techniques, so as to introduce an additional base pair between the upstream methionine (*i.e.*, met 1 in SEQ ID NO:2) and the downstream methionine (*i.e.*, met 30 in SEQ ID NO:2). The nucleotide sequence of HKNG1-V1 was likewise modified, using standard site directed mutagenesis techniques, to introduce an additional base between its upstream methionine (*i.e.*, met 1 in SEQ ID NO:4) and downstream methionine (*i.e.*, met 12 in SEQ ID NO:4). Thus, in both modified constructs, the C-terminal flag epitope tag was no longer in the same reading frame as the upstream methionine but was in frame with the downstream methionine. Consequently, exclusive translation initiation at the first methionine of a construct would lead to the production of non-flag immunoreactive proteins. However, exclusive translation initiation at the second methionine of a construct would lead to the production of flag immunoreactive proteins.

Unmodified HKNG1:flag, unmodified HKNG1-V1:flag, modified HKNG1:flag, and modified HKNG1-V1:flag constructs were transfected into cells, and their resulting gene products were harvested, blotted onto a PVDF membrane, and probed with an M2 anti-flag polyclonal antibody, and developed according to the methods described in Sections 12.1 and 12.2 above.

Flag immunoreactivity was detected in all four samples. The unmodified HKNG1:flag and HKNG1-V1:flag expression vectors produced amounts of mature secreted HKNG1:flag protein consistent with the levels detected in Sections 12.1 and 12.2 above. Further, the flag immunoreactive band detected for the modified HKNG1:flag construct was indistinguishable in intensity from the band detected for the unmodified HKNG1:flag construct, indicating that the immature HKNG1 protein produced by full length HKNG1 cDNA is

predominantly IPF2, while IPF1 is produced by full length *HKNG1* cDNA in relatively minor amounts.

The flag immunoreactive band from the modified *HKNG1-V1:flag* construct had dramatically reduced intensity relative to the band from the unmodified *HKNG1-V1:flag* construct.
 5 Thus, *HKNG1-V1* produces primarily the immature *HKNG1* protein IPF3, while the immature *HKNG1* protein IPF2 is produced by *HKNG1-V1* in relatively minor amounts. These results are summarized below in Table 6.

10

TABLE 6

Construct	Immature Protein	Prominence
<i>HKNG1</i>	IPF1 (SEQ ID NO:2)	Minor
	IPF2 (SEQ ID NO:64)	Predominant
<i>HKNG1-V1</i>	IPF2 (SEQ ID NO:64)	Minor
	IPF3 (SEQ ID NO:4)	Predominant

15

Thus, the *HKNG1* gene products of the invention include gene products corresponding to the immature protein forms IPF1 and IPF3. However, preferably the *HKNG1* gene products of the invention do not include amino acid sequences
 20 consisting of the IPF2 sequence (SEQ ID NO:64).

16. REFERENCES CITED

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the
 25 invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying
 30 drawings.

All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

5

10

15

20

25

30

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a HKNG1 gene product comprising:
 - 5 (a) the amino acid sequence of SEQ ID NO:2;
 - (b) the amino acid sequence of SEQ ID NO:4;
 - (c) the amino acid sequence of SEQ ID NO:39;
 - (d) the amino acid sequence of SEQ ID NO:41;
 - (e) the amino acid sequence of SEQ ID NO:43;
 - (f) the amino acid sequence of SEQ ID NO:45;
 - 10 (g) the amino acid sequence of SEQ ID NO:49; or
 - (h) the amino acid sequence of SEQ ID NO:66.

2. The isolate nucleic acid molecule of Claim 1, wherein the isolate nucleic acid molecule comprises:
 - 15 (a) the nucleotide sequence of SEQ ID NO:1;
 - (b) the nucleotide sequence of SEQ ID NO:3;
 - (c) the nucleotide sequence of SEQ ID NO:7;
 - (d) the nucleotide sequence of SEQ ID NO:34; or
 - (e) the nucleotide sequence of SEQ ID NO:35.

3. The isolated nucleic acid molecule of Claim 1, wherein the isolated nucleic acid molecule comprises:
 - 20 (a) the nucleotide sequence of SEQ ID NO:38;
 - (b) the nucleotide sequence of SEQ ID NO:40;
 - (c) the nucleotide sequence of SEQ ID NO:42; or
 - (d) the nucleotide sequence of SEQ ID NO:44.

4. The isolated nucleic acid molecule of Claim 1, wherein the isolated nucleic acid molecule comprises:
 - 25 (a) the nucleotide sequence of SEQ ID NO:46;
 - (b) the nucleotide sequence of SEQ ID NO:47; or
 - (c) the nucleotide sequence of SEQ ID NO:48.

5. An isolated nucleic acid molecule consisting of a nucleotide sequence that encodes a mature HKNG1 protein having the amino acid sequence of SEQ ID NO:51.

6. An isolated nucleic acid molecule which hybridizes to the complement of the nucleic acid molecule of any one of claims 1-5 under highly stringent conditions comprising washing in 0.1xSSC/0.1% SDS at 68°C.

5 7. An isolated nucleic acid molecule which hybridizes to the complement of the nucleic acid molecule of any one of claims 1-5 under stringent conditions comprising washing in 0.2xSSC/0.1% SDS at 50-65°C.

8. The isolated nucleic acid molecule of Claim 6 or 7,
10 wherein said isolated nucleic acid molecule encodes a functionally equivalent HKNG1 gene product.

9. A vector comprising the nucleotide sequence of any one of Claims 1-5.

15 10. An expression vector comprising the nucleotide sequence of any one of Claims 1-5 operatively associated with a regulatory nucleotide sequence controlling the expression of the nucleotide sequence in a host cell.

11. A host cell genetically engineered to contain the
20 nucleotide sequence of any one of Claims 1-5.

12. A host cell genetically engineered to express the nucleotide sequence of any one of Claims 1-5 operatively associated with a regulatory nucleotide sequence controlling expression of the nucleotide sequence in said host cell.
25

13. An isolated polypeptide comprising the amino acid sequence of a HKNG1 gene product having:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:4;
- (c) the amino acid sequence of SEQ ID NO:39;
- 30 (d) the amino acid sequence of SEQ ID NO:41;
- (e) the amino acid sequence of SEQ ID NO:43;

- (f) the amino acid sequence of SEQ ID NO:45; or
- (g) the amino acid sequence of SEQ ID NO:49;
- (h) the amino acid sequence of SEQ ID NO:66.

14. An isolated polypeptide consisting of a mature
5 HKNG1 gene product having the amino acid sequence of SEQ ID
NO:51.

15. An isolated polypeptide comprising an amino acid
sequence encoded by the isolated nucleic acid molecule of
Claim 6 or 7.

10

16. An antibody which selectively binds to the HKNG1
gene product of any one of Claims 13 or 14.

17. A method for treating a HKNG1-mediated disorder in
an individual comprising administering to the individual a
15 compound which modulates the expression of an HKNG1 gene in
the individual.

18. The method of Claim 17, wherein the compound
inhibits or potentiates the expression of an HKNG1 gene in
the individual.

20

19. The method of Claim 17, wherein the compound is a
small molecule.

20. The method of Claim 17, wherein the HKNG1-mediated
disorder is a neuropsychiatric disorder.

25

21. The method of Claim 17, wherein the
neuropsychiatric disorder is bipolar affective disorder or
schizophrenia.

22. The method of Claim 17, wherein the HKNG1 gene
30 encodes a HKNG1 gene product comprising:

- (a) the amino acid sequence of SEQ ID NO:2;

- (b) the amino acid sequence of SEQ ID NO:4;
(c) the amino acid sequence of SEQ ID NO:39;
(d) the amino acid sequence of SEQ ID NO:41;
(e) the amino acid sequence of SEQ ID NO:43;
(f) the amino acid sequence of SEQ ID NO:45;
5 (g) the amino acid sequence of SEQ ID NO:49;
(h) the amino acid sequence of SEQ ID NO:51;
(i) the amino acid sequence of SEQ ID NO:64; or
(j) the amino acid sequence of SEQ ID NO:66.
23. The method of Claim 17, wherein the individual is a
10 mammal.
24. The method of Claim 23, wherein the mammal is a
human.
25. A method for treating a HKNG1-mediated disorder in
15 an individual comprising administering to the individual a
compound which modulates the expression or activity of a
HKNG1 gene product in the individual.
26. The method of Claim 25, wherein the compound
inhibits or potentiates the expression or activity of a HKNG1
20 gene product in the individual.
27. The method of Claim 25, wherein the compound is a
small molecule.
28. The method of Claim 25, wherein the HKNG1-mediated
25 disorder is a neuropsychiatric disorder.
29. The method of Claim 28, wherein the
neuropsychiatric disorder is bipolar affective disorder or
schizophrenia.
30. The method of Claim 25, wherein the HKNG1 gene
30 product comprises:

- (a) the amino acid sequence of SEQ ID NO:2;
(b) the amino acid sequence of SEQ ID NO:4;
(c) the amino acid sequence of SEQ ID NO:39;
(d) the amino acid sequence of SEQ ID NO:41;
(e) the amino acid sequence of SEQ ID NO:43;
5 (f) the amino acid sequence of SEQ ID NO:45;
(g) the amino acid sequence of SEQ ID NO:49;
(h) the amino acid sequence of SEQ ID NO:51;
(i) the amino acid sequence of SEQ ID NO:64; or
(j) the amino acid sequence of SEQ ID NO:66.
- 10 31. The method of Claim 25, wherein the individual is a mammal.
32. The method of Claim 31, wherein the mammal is a human.
- 15 33. A method for identifying a compound which modulates expression of an HKNG1 gene comprising:
- (a) contacting a test compound to a cell that expresses an HKNG1 gene;
(b) measuring a level of HKNG1 gene expression in the cell;
20 (c) comparing the level of HKNG1 gene expression in the cell in the presence of the test compound to a level of HKNG1 gene expression in the cell in the absence of the test compound,
- wherein if the level of HKNG1 gene expression in the cell in the presence of the test compound differs from the level of
25 expression of the HKNG1 gene in the cell in the absence of the test compound, a compound that modulates expression of an HKNG1 gene is identified.
34. The method of Claim 33, wherein the HKNG1 gene encodes an HKNG1 gene product comprising:
- 30 (a) the amino acid sequence of SEQ ID NO:2;
(b) the amino acid sequence of SEQ ID NO:4;

- (c) the amino acid sequence of SEQ ID NO:39;
- (d) the amino acid sequence of SEQ ID NO:41;
- (e) the amino acid sequence of SEQ ID NO:43;
- (f) the amino acid sequence of SEQ ID NO:45;
- (g) the amino acid sequence of SEQ ID NO:49;
- 5 (h) the amino acid sequence of SEQ ID NO:51;
- (i) the amino acid sequence of SEQ ID NO:64; or
- (j) the amino acid sequence of SEQ ID NO:66.

35. The method of Claim 34, wherein the HKNG1 gene comprises:

- 10 (a) the nucleotide sequence of SEQ ID NO:1;
- (a) the nucleotide sequence of SEQ ID NO:3;
- (a) the nucleotide sequence of SEQ ID NO:5;
- (a) the nucleotide sequence of SEQ ID NO:6;
- (a) the nucleotide sequence of SEQ ID NO:34;
- (a) the nucleotide sequence of SEQ ID NO:35;
- 15 (a) the nucleotide sequence of SEQ ID NO:38;
- (a) the nucleotide sequence of SEQ ID NO:40;
- (a) the nucleotide sequence of SEQ ID NO:42;
- (a) the nucleotide sequence of SEQ ID NO:44;
- (a) the nucleotide sequence of SEQ ID NO:46;
- (a) the nucleotide sequence of SEQ ID NO:47;
- 20 (a) the nucleotide sequence of SEQ ID NO:48; or
- (a) the nucleotide sequence of SEQ ID NO:65.

36. A method for identifying a compound which modulates expression or activity of an HKNG1 gene product comprising:

- 25 (a) contacting a test compound to a cell that expresses an HKNG1 gene product;
- (b) measuring a level of HKNG1 gene product expression or activity in the cell;
- (c) comparing the level of HKNG1 gene product expression or activity in the cell in the presence of the test compound to a level of HKNG1 gene product expression or activity in the cell in the absence of the test compound,
- 30

wherein if the level of HKNG1 gene product expression or activity in the cell in the presence of the test compound differs from the level of HKNG1 gene product expression or activity in the cell in the absence of the test compound, a compound that modulates expression or activity of an HKNG1 gene product is identified.

37. The method of Claim 36, wherein the HKNG1 gene product comprises:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:4;
- 10 (c) the amino acid sequence of SEQ ID NO:39;
- (d) the amino acid sequence of SEQ ID NO:41;
- (e) the amino acid sequence of SEQ ID NO:43;
- (f) the amino acid sequence of SEQ ID NO:45;
- (g) the amino acid sequence of SEQ ID NO:49;
- (h) the amino acid sequence of SEQ ID NO:51; or
- 15 (i) the amino acid sequence of SEQ ID NO:64.

38. A method for identifying an individual having or at risk of developing a HKNG1-mediated disorder comprising the step of detecting the presence or absence of a polymorphism that correlates with an HKNG1 allele associated with the disorder, wherein presence of the polymorphism indicates that the individual has or is at risk of developing the HKNG1-mediated disorder.

39. The method of Claim 38, wherein the mutation results in production of a protein comprising an amino acid sequence that is different from the amino acid sequence of SEQ ID NO:2 or 4.

40. The method of Claim 39, wherein the mutation results in the substitution of a lysine for a glutamic acid at amino acid residue 202 of SEQ ID NO:2.

41. The method of Claim 39, wherein the mutation results in the substitution of a lysine for a glutamic acid at amino acid residue 184 of SEQ ID NO:4.

42. The method of Claim 36, wherein the method
5 comprises the step of analyzing the sequence of the coding region of the human *HKNG1* gene by preparing and sequencing cDNA comprising a sequence that hybridizes under stringent conditions to the complement of a nucleotide sequence which encodes the polypeptide sequence depicted in SEQ ID NO:2.

10

15

20

25

30

TGGGTCACCTGCGAGGCCCCGGGGGGGGGGTGGTTTCCACCCCTGGAGGTGGCTGACACCCCTGTGGCCCTGGGCTGACTTC
 CAGGCGGTGGCACGAGGCGCTCCAGGGGGCAGCACTCAAGGGCATCTTAGGAATGACAGAGTGGGTCCTCTCTGTTG
 CCAGGCTGGAGTTCAGTGGCATGTTCTTAGGTCACGAGGCGCTCAAAATTCCTGGGTTCAAGTGAACCTCCCACTCAGC
 OOCATGAGGAOCTGGGACTACAGGACACAGCTAAATCCCTGACAGGG ATG AAA ATT AAA GCA GAG AAA AAC 8
 E G P S R S W W Q L H W G D I A N N S G 24
 GAA GGT OCT TCC AGA AGC TGG TGG CAA CTT CAC TGG GGA GAT-ATT GCA AAT AAC AGC GGG 28
 N M K P P L L V F I V C L L W L K D S H 84
 AAC ATG AAG CCG OCA CTC TTG GTG TTT ATT GTG TGT CTG CTG TGG TTG AAA GAC AGT CAC 48
 C A P T H K D K T A I S E N L K S F S E 144
 TGC GCA CCC ACT TGG AAG GAC AAA ACT OCT ATC AGT GAA AAC CTG AAG AGT TTT TCT GAG 68
 V G E I D A D E E V K K A L T G I K Q M 204
 GTG GGG GAG ATA GAT GCA GAT GAA GAG GTG AAG AAG GCT TTG ACT GGT ATT AAG CAA ATG 88
 K I H H E R K E K E H T N L H S T L K K 264
 AAA ATC ATG ATG GAA AGA AAA GAG AAG GAA CAC ACC AAT CTA ATG AGC ACC CTG AAG AAA 108
 C R E E K Q E A L K L L H E V Q E H L E 324
 TGC AGA GAA GAA AAG CAG GAG GCC CTG AAA CTT CTG AAT GAA GTT CAA GAA CAT CTG GAG 128
 E E E R L C R E S L A D S W G E C R S C 384
 GAA GAA GAA AGG CTA TGC CCG GAG TCT TTG GCA GAT TCC TGG GGT GAA TGC AGG TCT TGC 444
 L E N N C M R I Y T T C Q P S W S S V K 504
 CTG GAA AAT AAC TGC ATG AGA ATT TAT ACA ACC TGC CAA OCT ACC TGG TCC TCT GTG AAA 168
 N K I E R F F R K I Y Q F L F P F H E D 188
 AAT AAG ATT GAA CCG TTT TTC AGG AAG ATA TAT CAA TTT CTA TTT OCT TTC CAT GAA GAT 564
 N E K D L P I S E K L I E E D A Q L T Q 208
 AAT GAA AAA GAT CTC CCC ATC AGT GAA AAG CTC AAT GAG GAA GAT GCA CAA TTG ACC CAA 624
 H E D V F S Q L T V D V N S L F N R S F 228
 ATG GAG GAT GTG TTC ACC CAG TTG ACT GTG GAT GTG AAT TCT CTC TTT AAC AGG AGT TTT 684
 N V F R Q K Q Q E F D Q T F Q S H F I S 248
 AAC GTC TTC AGA CAG ATG CAG CAA GAG TTT GAC CAG ACT TTT CAA TCA CAT TTC ATA TCA 744
 D T D L T E P Y F F F A F S K E P M T K 268
 GAT ACA GAC CTA ACT GAG OCT TAC TTT TTT CCA OCT TTC TCT AAA GAG CCG ATG ACA AAA 804
 A D L E Q C W D I P N F F Q L F C N F S 288
 GCA GAT CTT GAG CAA TGT TGG GAC ATT CCC AAC TTC TTC CAG CTG TTT TGT AAT TTC AGT 864
 V S I Y E S V S E T I T K H L K A I E D 308
 GTC TCT ATT TAT GAA AGT GTC AGT GAA ACA ATT ACT AAG ATG CTG AAG GCA ATA GAA GAT 924
 L P K Q D K A P D H G G L I S K M L P G 328

FIG. 1A

TTA OCA AAA CAA GAC AAA OCT OCT GAC CAC GGA GGC CTG ATT TCA AAG ATG TTA OCT GGG 984
Q D R G L C G E L D Q N L S R C F K F H 348
CAG GAC AGA GGA CTG TGT GGG GAA CTT GAC CAG AAT TTG TCA AGA TGT TTC AAA TTT CAT 1044
E K C Q K C Q A H L S E D C P D V P A L 368
GAA AAA TGC CAA AAA TGT CAG GGT CAC CTA TCT GAA GAC TGT OCT GAT GTA OCT GGT CTG 1104
H T E L D E A I R L V N V S N Q Q Y G Q 388
CAC ACA GAA TTA GAC GAG GGG ATC AGG TTG GTC AAT GTA TOC AAT CAG CAG TAT GGC CAG 1164
I L Q M T R K H L E D T A Y L V E K M R 408
ATT CTC CAG ATG ACC CGG AAG CAC TTG GAG CAC ACG GCC TAT CTG GTG GAG AAG ATG AGA 1224
G Q F G W V S E L A N Q A P E T E I I F 428
GGG CAA TTT GGC TGG GTG TCT GAA CTG GCA AAC CAG GCC OCA GAA ACA GAG ATC ATC TTT 1284
N S I Q V V P R I H E G N I S K Q D E T 448
AAT TCA ATA CAG GTA GTT OCA AGG ATT CAT GAA GGA AAT ATT TOC AAA CAA GAT GAA ACA 1344
K H T D L S I L P S S N F T L K I P L E 468
ATG ATG ACA GAC TTA ACC ATT CTG OCT TOC TCT AAT TTC ACA CTC AAG ATC OCT CTT GAA 1404
E S A E S S N F I G Y V V A K A L Q H F 488
GAA AGT GCT GAG AGT TCT AAC TTC ATT GGC TAC GTA CTG GCA AAA GGT CTA CAG CAT TTT 1464
K E H F K T H 496
AAG GAA CAT TTT AAA ACC TGG TAA 1468

GAGATCAATGCATCCTATATOCAGTAGTAGAATTATOTCTTCATCTGGGACCTGGAAATCCTGAAATRAAAAAGGA
TAAATGCANAAACACAGTTGCAGGAAGTATGTTAGCCTATATACTATGAGTACTCTTAGTCTTACTATGTGAAATGGC
TEAGCTATTAATCTCAATGTGAGTTAAAATGAAAATTOCTOCTTAAAAATCAAAOCTAATAATGTATTACATTTCAAG
GTACTTAGTACTTCTTTGTATATGAAATAACTAAATCACTA

FIG. 1B

TGCGTCACTGACAGGGGGGGGGGGGGGGGGGGTGGTTTCCACCCCTGGAGGTTGCTGACACCCCTGTGCGCTGGGCTGACCTC
 CAGCGGGTGGCACAGAGGGCTCCAGGGGGGCGCACTCCAGGGCCATCTTAGGAATGACAGAGTTGGGTCCCTCTGGGTTG
 CCAGGGCTGGAGTTCCAGTGGCCATGTTCCATAGCTCACTGAAGGCTCAATTCCTGGGTTCAAGTGAACCCCTCCCTCAGC

 H R T W D Y S N S G N M K P P L L V F 19
 CCC ATG AGG ACC TGG GAC TAC AGT AAC AGC GGG AAC ATG AAG CCG CCA CTC TTG GTG TTT 57

 I V C L L W L K D S H C A P T N K D K T 39
 ATT GTG TGT CTG CTG TGG TTG AAA GAC AGT CAC TGC GCA CCC ACT TGG AAG GAC AAA ACT 117

 A I S E N L K S F S E V G E I D A D E E 59
 GCT ATC AGT GAA AAC CTG AAG AGT TTT TCT GAG GTG GGG GAG ATA GAT GCA GAT GAA GAG 177

 V K K A L T G I K Q H K I H M E R K E K 79
 GTG AAG AAG GCT TTG ACT GGT ATT AAG CAA ATG AAA ATC ATG ATG GAA AGA AAA GAG AAG 237

 E H T N L H S T L K K C R E E K Q E A L 99
 GAA CAC ACC AAT CTA ATG AGC ACC CTG AAG AAA TCC AGA GAA GAA AAG CAG GAG GGC CTG 297

 K L L N E V Q E H L E E E E R L C R E S 119
 AAA CTT CTG AAT GAA GTT CAA GAA CAT CTG GAG GAA GAA GAA AGG CTA TGC CCG GAG TCT 357

 L A D S W G E C R S C L E N N C M R I Y 139
 TTG GCA GAT TCC TGG GGT GAA TGC AGG TCT TCC CTG GAA AAT AAC TGC ATG AGA ATT TAT 417

 T T C Q P S W S S V K N K I E R F F R K 159
 ACA ACC TGC CAA CCT AGC TGG TCC TCT GTG AAA AAT AAG ATT GAA CCG TTT TTC AGG AAG 477

 I Y Q F L F P F H E D N E K D L P I S E 179
 ATA TAT CAA TTT CTA TTT CCT TTC CAT GAA GAT AAT GAA AAA GAT CTC CCC ATC AGT GAA 537

 K L I E E D A Q L T Q M E D V F S Q L T 199
 AAG CTC AAT GAG GAA GAT GCA CAA TTG ACC CAA ATG GAG GAT GTG TTC AGC CAG TTG ACT 597

 V D V N S L F N R S F N V F R Q H Q Q E 219
 GTG GAT GTG AAT TCT CTC TTT AAC AGG AGT TTT AAC GTC TTC AGA CAG ATG CAG CAA GAG 657

 F D Q T F Q S H F I S D T D L T E P Y F 239
 TTT GAC CAG ACT TTT CAA TCA CAT TTC ATA TCA GAT ACA GAC CTA ACT GAG CCT TAC TTT 717

 F P A F S K E P H T K A D L E Q C W D I 259
 TTT CCA GCT TTC TCT AAA GAG CCG ATG ACA AAA GCA GAT CTT GAG CAA TGT TGG GAC ATT 777

 P H F F Q L F C H F S V S I Y E S V S E 279
 CCC AAC TTC TTC CAG CTG TTT TGT AAT TTC AGT GTC TCT ATT TAT GAA AGT GTC AGT GAA 837

 T I T K H L K A I E D L P K Q D K A P D 299
 ACA ATT ACT AAG ATG CTG AAG CCA ATA GAA GAT TTA CCA AAA CAA GAC AAA GCT CCT GAC 897

 H G G L I S K H L P G Q D R G L C G E L 319
 CAC GGA GGC CTG ATT TCA AAG ATG TTA CCT GGG CAG GAC AGA GGA CTG TGT GGG GAA CTT 957

 D Q N L S R C F K F H E K C Q K C Q A H 339

FIG. 2A

WO 99/47535

4/57

PCT/US99/05606

GAC CAG AAT TTG TCA AGA TGT TTC AAA TTT CAT GAA AAA TGC CAA AAA TGT CAG GCT CAC 1017
 L S E D C P D V P A L H T E L D E A I R 359
 CTA TCT GAA GAC TGT OCT GAT GTA OCT GCT CTG CAC ACA GAA TTA GAC GAG GGG ATC AGG 1077
 L V N V S N Q Q Y G Q I L Q M T R K H L 379
 TTG GTC AAT GTA TCC AAT CAG CAG TAT GGC CAG AAT CTC CAG ATG ACC GGG AAG CAC TTG 1137
 E D T A Y L V E K M R G Q F G H V S E L 399
 GAG GAC ACC GGC TAT CTG GTG GAG AAG ATG ACA GGG CAA TTT GGC TGG GTG TCT GAA CTG 1197
 A N Q A P E T E I I F N S I Q V V P R I 419
 GCA AAC CAG GGC CCA GAA ACA GAG ATC ATC TTT AAT TCA ATA CAG GTA GTT CCA AGG ATT 1257
 H E G N I S K Q D E T H H T D L S I L P 439
 CAT GAA GGA AAT ATT TCC AAA CAA GAT GAA ACA ATG ATG ACA GAC TTA AGC ATT CTG OCT 1317
 S S N F T L K I P L E E S A E S S N F I 459
 TCC TCT AAT TTC ACA CTC AAG ATC OCT CTT GAA GAA AGT GCT GAG AGT TCT AAC TTC ATT 1377
 G Y V V A K A L Q H F K E H F K T W 478
 GGC TAC GTA GTG GCA AAA GCT CTA CAG CAT TTT AAG GAA CAT TTT AAA ACC TGG TAA 1434

GAAGATCTAATGCATCCTATATCCAGTAAAGTAGAATTTATCTCTTCATCTGGGACCTGGAAATCCTGAAATAAAAAGGA
 TAATGCCAATAAACACAGTTCCAGGAAGTATGTTAGCTATATACTATGAAGTACTCTTAGTTTACTTATGTTGAATGCC
 TTAGCTATTAATACTCAATTGAGTTRAAATGAAATTCCTCCTTAAAAATCAAAAGTAAATATGTATTACATTTCTAG
 GTACATTAGTAGTTCTTTCTATATGAAATAAATACTAAATCAOCTA

FIG. 2B

ACATTTTAAGCTACTTATAGTCCTTGGAAATAGCAACAAATATCTTAGTTATTGGACTATTATAACCTTAGTCATCTTATTACTGCTTG
 ATTATGAGACACTCTCCCTGCTAATCCTTAGAACATCTTGGTTCTTGGTACTTGACTTTTAGCCCCCTGACATATAGTTGATGTCAGA
 GTGTCTGGCATTTCAGTAGTGCTCTATTTTACAAATCCCAGTAAACTGCTCCACTGTGGCTTGTTTATGTGTTAATACTGCTTGTTTTC
 TGTATAAATTATTTTTGCTTTGGAGTAAGATATCATCATTTTGCATAGCTACAAATCTGAAGTTAAAGAAAATTTTAAAAATGTAAT
 TGTGGGAAAATAACAAATAGATCTGCTGAGATGGAGGCTTTGACTAATGTTTAAATAACAGGCAACAAACAAAGAGGCAGGATATTTT
 GGTCACAACCTAAACCTAAATTAATCCTCATACAAAGCCCCATTAAGATAAATGCTCAAATTCCTGGGAACATTTCACTTGCTTTGCCAG
 CAATTTTACCCTTCAGAGGGTGTGGATCTAATCAGGGGAACAACTACCCTGGGCTTAATTCATTAAACAGGGACTAATTTGTCAAAG
 CGGCAGTACTAGCTGAAGTGATGGGTATGGAAGCATTCACTGTGAGGATTTTGTCTGAGGTGCCTGGCACAGGGTAGGGGAACTCACCCA
 GGCTGCAAGATGCTAACAGTTCAAGTCAAGGTCTTAGTGTGGACTAAGGTGCAGTCAGGATGGGAACAGGTGCAACTTGGGCCAACAT
 CAGTATGAAGGGCCTGATCTGAGGGCAGGGGAAGGAGGGGGCATTCTGGGAAGCAAGAGTCTCTGGTATCCTGTGACCAGAGTCTTGG
 CCCAAGGATCAACGTATGAATTAAGTAGAAATACCAGAAACAAAGAAAGTTGGCAGAACTAGGAGAAGCAGAGTCTCAGCCAACTGG
 ACTGGGCTCAGCCTTGGCTACTGGCCCGGCAGATGATAGAAGAGAAAACCAGGAACCCAGGCTGAAGCCCAGTGGTTGGGCTGGCCACA
 CACCATGCATAGCCTTAAAGGGGTGGCCTAAGGGCATGGTCCGCTCCAAAAAAGGAAAGGGGGCCCCAGAATATTTCTGAATCCCCTC
 ACTGCCAGGGGAAGAACCCTCAATTCACTCAATAGTGCATTTCTCCTGCTTCTCAATAGGCTAATACTCTAGAGAATATGGGGACAAGGG
 GAGGAGGGTCTAGTGGAAACAGGTCTAAACTGGCGTTTGAATTTAAGATAAGTTAATCATACATTGGCTGGGTGAGCCATGTCTCTTAG
 TCTTTACAAAAGTAGAACACAAAAAATTCATGGAAATCTACAGACACCTATTTGCAGATGAGGAAACACGGCTATGAAGATTGGGAA
 GATTGGGAAGAACTGGCCAGGTGTGGTGTCTCACGCCTGTAATCCAGCACTTTGGGAGGCCGAGGCTGGTGGATCACTTGAGGTGAGGA
 GTTGGAGACCAGCCTGGGCAACATAGTAAAACCCTGTCTCTACTCAAATTACAAAAATCAGCAGGGCGTTGTGGTGGCCACCTGTAATC
 CCAGCTATGCAGGAGGCTGAGGCAGGACAATCACTTGAACCTGGTAGGCGGAGGTTGCAGTGAGCCAAAATCACGCCACTGTACTCCAG
 CCTGGGTGACAGAGCAAGACTTTGTTTAAAAAAGGGGAAGAACTAAAAATGTAATTTTCAAGGGGCTATCACAAATGGT
 CCCAATAAAGAGAAAGCAGGACTCATGTTAAGAAACCCATGAGATGTGTATGGACCTCATGGAAGAGCTCTTGCTTTCTAATGATCTA
 CGTAACAGATGAAAAGCAGAGCATAGGGCTAAGGATGAAAATACAACAGTAATAAGGTATTAATATATTATTAAGAAAGCTAATGCTCC
 ACATAAGCAGAGGACATTAAGGGACTTTTTTTCTTAAGGATATCTTAATGTTTAAATGAGAAGACATAGAAAGGGATAGGTCCAAC
 TCTTGGGATTGTTGCAGGTGGTTCCATCGGAAGCACTCTGAGTCTGAGATTTGTATGCAGAAAATTAATTTGAATGTGCTTTTCAGA
 TCACCAGGTGGGGAGGGAGGAAACCAGGACTGGGCAGAGAGAGGCTGGGCTGTAACCAAGTCACAACAAAGGTGTGAGCTGGTCCCA
 TGGTGAATTCGGACCTAGGATGGCTGATCCCAAGGCATTCCAACTGGGGCAAGGAAGTTGTGCTTTAAACTTCTCATTGACTGTCA
 GTCACTGGGCATGAGCAGTCCCAGGAAGGGGGGATGACCTTGAGCAAGGTGGATGTCTTCAGCCAAGGGCAA YCACTGGGAAGGAGAA
 CCCAGCTATGAACTGTGAGCTGCCAACACTCCCAGCATCTGAGAGGATGAGGGCTTCAATTCCTAAGGGCAGGGGCTCCAAGGGCAGGGG
 TACGGATGGTGGAACTCTGGGCAGTACCTTGTGGCTTCCACTACAGTCCACCCCTTGACCCTTAGTTCCACTGGCTTTTTTTTTTTTT
 TTCPTTCTGAGACAGTCTCACTCTGTCAACCAGGCTGGAGTGGGTGGCACAGATCTCGGCTCGCTGCAACCTCCGCTCCAGGTTCA
 AGCAATCTTGAACCTCCTGAGTAGCTGGGACTACAGATGTGTGCCACCACACCAGCTAATTTTTTGTATTTTTTAGTAGAGACGGGT
 TTTACCGTGTTAGCCAGATTGGTCTGATCTCCTGACCTCATGATCGGCTGCTTTGGCCCTCCAAAGTGTGGGATTACAGGTGTGAG
 CCACCGCACACAGCCAGATCCACTGGCTTCTATATAATTTCTGGGTGAAGCTAATTCAGGATTCGTATGGACCTGTCTTCCCGAGGGAA
 ACTTGTAAAAGGAAAGTTAGAGGGGACAACTATAGCCCCCTGCCACAGCAGCTGCTGTGAGGACAAAATGGTGTCTCTCATTTCCTT
 AACCACCTGACCTAGATTCCCCTAACCTTAGTGGGCACCTCTGTGGATGGAAGTGGTGGCTCACYKGGKGGRRMRYCMRRWYWCWY
 YCCCTGAGTGGTCTGAGCTCCAGTTACCAGGCOCTTCTCAGGCTGTGGCTGTGCACTTACCTCCCCAGCCATCCCCACTTTTTTTT
 CTTGAGACTGGGTCTTGTCTGTCAACCAGGCTGAAATGCAGTGGCATAACCTCAGCTCACTGCAGCCTTGATCTCCCAAGCTCAAGCC
 ATCTTCTCACCTCTGCCTCCCAAGTGGCTGGGACTACAGGCACATGCCACCATGCCAGCTAATATTTTTTATTTTTTATTTTTTTGTA
 GCAATGGGATTTTGGCATGTTTCCAGGCTGGGCTTGAATCCTAAGCTCAAGCTATCCTCCCACCTCTGCTTCCCAAGTGTCTGGGAT
 TACAGGCTGAGTCACTGCATCTGGCCACATTTATTCCTTTTAAACGTTAAAATGGAATGCAGGATCACTGAGAGACAGGTGAGTGATT
 ACCAGGGTGCCAAACATAACCTTCTCCTCCTTCTCCTGAGCTCTACCTCCTCCTGATGATCAGGACAATCATGTATGATGACTCCTTTC
 CTTGACTGCTGCTCTCTCAGAAGGAACCCATTGTGTTGGGTGAGAACACATCATTTGAAATTTAGTAAGACTCTTGTGTGCCTATGGT
 AGAAGCATCCCTCTCTGGGGCCAAGATCTTTAAATGCACAGAGTCCAAAGTCGTGGGAACCAAGCAGAAATTA AAAAGGAGATGACT
 GGGATTATGGTAAGAACTGTTTCCACCCTTGATTTGCTGCACCCATGTGTTCTACCTAGGAGATAGCACACCATATACTGGTTATTCA
 TTGGATTACATGCTGCATCCCGGAGAATGGGCACTGCATTCTCACTGGTCAATCATGTGAGAGCCTGGCTGCAGAGGCTTCCCATTGC
 TCTGTGAGTGTGTTATAGGGTCAAGGATTTCAATGGTCAATGTGCCCACTGCTGCACCTCCATTCTTGTAAAATGGGTCCCTCTGGTTCAA
 TGTGATGCCATGTGGGATCTTGTGTCAATAGAAATAAATACTCAGATGTTCTGGCTGAAGCTTTACAAGCAGAAAAGGCCAACCGATGAC

FIG. 3A

ATGGATCTAAGAAGAGTTATTGATTTTCAATTTGTCCAACCTTAATTCCTGTTTTGAAGACAGAAGTGATGACTTCCAAGCTCTTTATAT
 GTTGAACCCAACCCCATATTATTTTCAATTAGCAATTGCATATAGCAATGGTACATTGCATTTATAGAAATATAAATTGATGTTTGCCTG
 TGTATCTTTTTTCTATTATGTTGCTGAATTCATTTCTTAGTTCTAGGAATTTTTCAAATACATCCCTTAGGATATCTGTATACATAA
 TCATGTCATCTGCACATAGGGACAGTTTTATTCTTTTTCTAGTCTGTATTCTTATTTCCCTTTCTTGCCTTATTCAGTGGCTAGAA
 CTTGCAGCACTATATTAATAAAGAGTGGTAAAAGTGAACATTCCTTTCTTTGTTGCTGATCTTGGGGGAAAGTATTCAGTCTTTCACC
 ATTGAGCATAAATGTTAGCTGTAGGTGTTTTAAATCTTTATCCAGTTGACGAAGTTACCCTTTATTCCAATTTTTCTGAGAGTTTATATC
 AATAATGTTTAAATTTTGTCAAATTTTTTGTGCATGTATTGATATGATTATGTGGTTTTTCTTCTTTAGTTACTGCAGTGGGTTGCATT
 GATTGATTTCTATTATTGAACCAGCCTGCATTCCTGGAATAAACCCCATTTGGTCATGATGTATAAATCTTTTTTTTTATATTGCTGAAT
 TCTATTTGCTAATATTTTGTAAAGGATTTTGCATCTGTGTTTCATGAGGGATCTGGGCTGGTAGGTTTTTTTTCCCCCTGCAATGTCTC
 TGTCTGGTTTTGGTATTAAGGTAATTTTTTTTTKTTTTKTTTTTGAGATGGAGTCTCGCTCTGCTCACCCAGGCTGGAGTGCAGTGGCAC
 GATCTTGGCTCACTGCAACCTCCACCTCCCAGGTTAAGCGATTCTCCTGCCTCAGGCTCCTGAGTAGCTGGGACTACAGGTCACACCA
 CCACGCCCCGACTAATTTGGTATTAAGGTAATATTATCATCATAAAAATGAACTGGGAAGTGTGCCCTCTTCTTGTATTTCTTTTTTTTT
 TTTGAGACAGTCTTGTGTTGCCAGGCTGGAGTACAGTGGTACGATCATGGCTCACTGCAGCCTCAAACCTCCCAGGCTCAAGTGTATCT
 TCCTGCCTCAGCCTTCCCAGTACAGGGCAGGCTACCACATCTGGCCAATTTTTAAATTTTTCTTTTGTAGAGAGGGGCTCTCACTATGT
 TGCCCAGAGGATCTCAAGCAATTCACCTACCTTGGCCCCCTCTTCTTGTATTTTATGGAAGAATTATGGTGTCAATTCCTTCTTGAAGT
 TCTGTTAGAATTCCTCAGTGAAGCTGTATGGGCTTGAAGATTACTTTTTTTTTTTTTTTTTGAGATGGAATTCACTCTTGTGCCCC
 AGGCTGTAGTGCAGTGGTGTGACCTCTGCTCACTACAACCTCTGCCTCCCACGTTCAAGTATTCCCCTGCCTTACTCAGCCTCTGGAG
 GAGCTGGGATTACAGGCACCCGCCACCATGCCCGGCTAATTTTTTGTATTTTAGTAGAGACGGGGTTTACCATGTTGACCAGACTGG
 TCTGAACTCCTGACCTCAAGTGATCCACCCGCTCGGCTCTCAAAGTGTGGGATTACAGGCATGAGCCACCGCGCCAGCTGAAGA
 TTTCTTTTTGGGGAGTTTTAAATTATACAATCAATTTGCTTAATAGGTATAAGCTATTCAAGTTATCTATTTATACTGGATGAGTTGC
 AATAGTTTGTGGTTTTATGAGTTTATATGGTCCATTCATCTGAGGTATAAAATTTAYTTGTGTAGTATTGTTGGTAGTATTCCCCTGTT
 ATCTTTTTTATGTTACATGGTATATGGTGACAGTCCGTTTAAATTCCTAGTATTAGTAACTGGCTCTCTCTCTCTCTCTCTCTCTCT
 CTCTCTCTCTGGTCACTTTCCAGAGGTTGTCAATTTTGTGACTTTTTTCCCCAAAGAATCAGCTCTTGTTCATGGATTTCT
 GCTTTCTGTTTTCAACTTCATTGATTTCTGCTGTTATTATTTCTCTCCTTCTGTTGGTGTGAGTTTGTTTGCTTTTCTTTTTCTA
 CATATTCGATGTGAAATCTTACATTATTCACTCGGGACTTTCTTCTTTTTGATGTATGCATTTAGTATTCTAAATTTACTTCTKAGT
 ACTGCATACTGCTTGAACATATGCTGACAAATATTAATATATTGTTTTTAAATCTTTATTCAGTTCAGTGTATTTTTAAATTTCCCTC
 TCTGCCCTCTTTTTGATTTGTTATTTAGAATGTGTTGTTATTTTCCGAGTATTTACATTTTCCCTCTTATCTTTCTGCATTGATTCCAT
 CGTAGTCAGAGTGCATGCTCTGTACAGTTTCAGTCTTTCAAATTTATGAGCTTTGTTTAAATGGATCTGGATAACAGTTTATCTTGGCA
 TATATATATATATACACACACATATGTATGTGGGCGCTTGAAGAAGAAAGCGTATCTGCTGTTTGGTGAATGTTTGGAGTGTCTATAA
 GGGGTGATTAGATACTGTTGGTTGATGATGTCATTGAGGGTCCGATAAACCTACTGATTTAAATTTATTTAGTCTGTCAATTTATTCAGA
 GAGAGAGGTGTTGAACTCTGCAATGTGAATGTGGATTTGTCAATTTCTCCTTTCAGTCTATTAGTTTTTTCTTACATATTTTACAA
 CTCTGTTGTTTGGTGCATACACATTTATGCACCAAATTTAGGATTGCTATAACTTCTTGGTGGATTGACCCTTTTACATATATAATGT
 CTTTTCTGTCCCTGOTAATTTGGTTGCTCTGAAGTCTATGTTATCTCAATATAAATAGACAACCTCTGCTTTCTTTTTGATTAATGTTT
 ACATGATACATCTTTTTCTATTTCTTTACTTTCAACTTACTTATATTATTATGTTTGAAGTGAAGCTTCTTGTAGACAGCATGTAGTAGG
 TCATATATGTACATAGATATATATATTTTTTTGAGATGGTGTACTCTGTCAACCAGGCTGGAGTACAGTAGTCTCACTGCAACCTCTG
 CCTCCTGGGTCAAGTGTCTGTCCKCAGCKCCCCAGTAGCTGGGATTACAGGCACGCCACCATGCCCAGCTAATTTTTGTATTT
 TTAGTAGAGACGGGTTAAACCATGATGGACAGGCTGGTCTCGAACTCCCGACCTCCAGCGATTAGCCCACCTTGGCCCTCCCAAAGTGTCT
 GGCATTACAGGTGTGAGCCACCGTGCCTGGTTAATATTTTTTAAATCCACTCAGTCTTTGTCTTCTACTGGTGTACATAGACATTOGCAT
 GTAATGTAATGTTGATATGTAAGAGCTTGAATCTGTTATGTTTTGCTTTCTCTATGTTTTCTCAATTTTTAAATTTCTCTGTTTTCTT
 TTTTTCTGCTTCATATTGGCTAATGAACACTTTGAATCATTCOATTTTGAATTTACCTATAGTGTTTTTTAGTGTCTCTTTGCATAGC
 TTTTTAGGGTTACTTTAAGTATTTCAATTATATGTACATAACTTATCACAGTATATTGGTATOGTTATTTTACCAGTTCAAGTAAAGT
 ATGGAAATGTTTCCCTCTACATTCCTTTACCTCATTTATAATATAAATGTCTTAGGTATTTCTTGTACATACATTTTAAACCGGATGA
 GTGTTATTTTTGATTTAGCTATCAAATAATCCAAAAACTCAAGAAAAAAGGAAAGCTTACTATATTGACCCATATTTTCATTACCA
 TGTGTTTTCTTCCCTCTTATGCCCCATAGTTCCCTTCTTCTATTGTTTTCGTTTTAGAGAACTTCCCTAGCCATTCTATTGGGGTAGATCT
 CCTAGTGACAAATTCCTTAGCTTTCTTTCTCTGTGAATGTCTTTATTTCCCTCTTGTTCCTGGAGGACATTCCTCACTGGATATAGG
 ATTCTTGGCTATTGGGCTTTTTCTTTTTGGCACTTTTGTAAAGTGTGCAGCCTGCTGTCAAATAAAAATTAATAAATAAATAAATAAATAA
 GTTTTTCTTTGCTACGTTTCATGAAAGTATAATTCAGTGAATGAGGAGGACACCCATCTCTATAATCTGGAGGCCCATGCTCACCTCTG

FIG. 3C

AATAGTACATTTGCAGAGAAATTGGGGAAATCAAAGTCTGTTGAGACCAGCAAGATAAATAAGGCCAAAAGGATACAAAACCATATCCAA
AGAGAAATGGTTTAAAGGAACTAAGGCTGTTCTCCTAAAAGAAAATAGTTGGAGACATGTGACCTCCAAAGAAACAGGACTTTTCT
ATGGGCTCCAAGGGTTTCTATGAGAGAATGATAAAGGAGAGATTTAGCTTAGTCTCAGGAAGACTTTTCAACAACCAAACTGCC
AAAGATGGACTGCCCTGCCTAAGGATTGTGTTCTGACATTAAGGGTATGGAGGTATGGGTTAGATGAATATTTTACCAAATGCCATAG
ATATTTAGGCTATTGATGTTGTAATATCATACTAGGCAACTCCACTTCAATATGAGTCTCTATGATGTAAAATGAAATAGGATGTGTT
TCGATAGAGAGTTGCAGATTTCAATTTGATGTTAGCGACCACACAAAATTAATTTCCCTACATAAGAACATGTTATTACTCTAGTTGAT
GATGACTGCTTATGGGAAATGTGCTGCTTTGTTAGGAATCTTGCCATAATATATGTATAATTCAAGATGGTATTATAAAGTGACATATA
TGATTTTAAACATTTGCACCTAAAATAACACTTATTCTGTACCATGMASTGTCTAGGAGCTTCTACATATTCATTATTATCTTTATTTT
ACAAGACAGGAACTAAGGCATGGAGAGATTGAGTAATTTGTGCAATATTACCTACCTAGTAAGTGGTAAAGGAAAGATTGGAACCCAT
TCTGGCTCCAGGATCCAGGCTCAAAGCCAATATACTATCCACCACCCCAACTCTTTAGTTGATCAATTTGTCAAATATTTTACAGTT
ATTTATCTGTAAATTAAGGGGATAATTGCCAGTCAATAAATGTGTCCCCTCAAAGGTTACATACTTAACCAATGGTGTACTGCGGCT
CAGAACATTTTGGAACTACGATTTTGGTGGCAACCAAAAACCTCCAGTACATTCCTCTGAACATTCCTCCAGAGGCAAGTCTTCTCC
ATGGAGACTGGGCTTCATTTTGAATTAGCCTGAAGTTGTTGAGGTCAAATCTGATGAAAAGAGCGGCTGGGGAAGCTGGATATTTT
CGTTGCTGATTTAAACAGTAAATGCCACCTAAATGAGAAGGCTACTTCTTTGAAATGTTTGTAAACTGGCTTTGAAGGTACTTCTTT
AAAAAGAAGCACAAGAAAGACGGTACTGGCAACAGCCTCACTGGAATACGTCTCTAATCATCAAGGCAACCCACACTCATTTGGATG
TGTGCATCCGGTATGTTATTATTTTAAAGTTATGTGCCACAAAGATGCATTCCTTTGCTATACAAAAGAGCTGTTGTTAAATTTATAA
AGATATAAAAAGGGGAAAGGAGAAGGCACCAATGGAAGATTCTTAGGCATTAAGTGTCTCAGACAGCATAGATCTTCATTAGATGACGT
CAGGGAGAAGAGACACAGACTTTGCCATCTCAGGTAGAAGTATCAAAGTCAACAGCCTCCTAGTAAGACAGACCTGGGTTTGAAGCTCT
GCACAGCCATTTCTAGCTGGTCTGGGAAAATTAATTTCTTGAAGCCTCAGTGTCTTTATTTGTAAAGTAAGTGGAAATTATATTACCT
TGTCAGGATGTTGTCAGAATTAGAAATAATTTAAAGAGGTCCAGCAAGAGCAGGTCAATCAAGGGAAGATGTTAAAAATAACAACAGGT
GAAATGTACTCCCAAAGATAAAGTGGATACATAGATGAATCTTCTCACACACAGAGTATAATAACCTCAGAAAAATATTGCCCTAGAG
TAAACATGCCTCCCAAGCCAAGTTCATCATCCAGGAATACGGAGAGGATGTTTGGGATATGGGGGGCATGAAATTTTACAATTGTAGG
GCCCTTTAACAAGGGTAGACTTGCAAGTTGCACTGMCCTTCTGCCCCCTCTGCTGCTACCTGTTCCAGCATCCAGAGTTTGTGAACCTG
GGGCAAGGACAGCACCCCTGGCATGGGCAGGCCACTNGGCGACTCTCTCAGGGCTGCTGCAGCTGTGTCACTGTCCCCACAGGGAGN
CTGACATCCAGCCATGACCATCGCATTAAAGCCCAGCAGTCAGGGCAGGGGAGCAACTGCTCAGAGGCACCTTTGACCCACTACTTTTTT
CCCCCTGCTTTATCTGCCAGAGCGAGGCTCTCTTCTAATGTGTACAAGGCGTTCTACCTATGACTCGTGGTCTGCCATAGAAAT
GCTTTTTTTTTTTTAACTGAATTAAGTTGCCAAGTTTGAAAAATCAGAATTCACATAAGATCCCTATTTCTGTCTTCTTTTGA AAAA
CTGAATGTTCTTTCCACAGTGAGCCACATTCCTTCTGACGACCATCACCGTTCACTGGAGTAGAGAGGGCTCTGCTGGCTTCAGAT
COGGACCGCAGGTCTCTGCAGGCCCCGCCACCCGGCGTCACTGCAAGTCCCGCCACCCGGCGTCTGCAGCCCCCGCCACCCCGG
CCTCACCTGCAGGCCCCGCCACCCGGCGTCTGCAGGCCCCGCCACCCGGCGTCACTGCAAGGCCCCGCCACCCGGCGTCTGCAGG
CCCCCCCCCGGCGTCACTGCAAGGCCCCGCCACCCGGCGTCTGCAGGCCCCGCCACCCGGCGTCTGCAGGCCCCCGGCGTCTGCAGG
CGTTGGTTTCCACCTGGAGTTGCTGACACCTGTGCCCTGGCTGACTTCCAGCCGTTGGCACAGGCGCTCCAGGGGCGAGCACTC
AGGGCATCTTAGGATGACAGTTGAGARCATCCTCCGGGCCCCAGATTTCTCTCCTCGCCGCTTTGCCCATTTCTCCGGAGAGCCAG
AGAAAGCCGCTCCCAAGTCCAAGGCCGAGCTCCGAGAGCCCCGGCCCCCTCCGGCCGACAGAACAAAGCCATTGTTCTTGGCCGGGA
AGGTAGAAATACTGTGGGCTGCTTCAGAGGCTGCCGAGCAAACTCAGGCAATCTCCTGGGCTGTTCCAATAAGTTTATTCTCTTTTTC
AAAACAGGAGGAGGAGGTAGAGGCGGGGAGACACACCATCCCTGCAAACTACTGGCAAAACTAAGCCGAGCCGGGTGTGGTGGCTCA
CGCCTGTAATCTCAACACTTTGGGAGGCCGAGGGGGCCGATCACTTGAGGTCAGGAGTTGGAGGCCAGCCTGGCCGGCATGGTGA AAC
ACAAAATTAGTCGATTGTGGTGGTGCATGCTTGTAAATCCCATCTACTTGGGAGGCTGAGGCAGGAGAATCGCTTGAACCCGGGAGGCG
GAGGTGCACTGAGCCGAGATTGGGCCACTGCACTCCAGCCTGGACAACACAAGTGAGATTCTGTCTCAAATAAATAAATAAATAAAC
CCAAGCAGAAAAGAATCACTCTGAAAACGATCACATCTAATGCTCATACAGTTTATGGAATTTATCAGCCCAACTTGATAAA
ATCAGTATTTGAGGAACTGTGGATAAGCCCCCTGATTTCAATCCCATTTGTGCCAGGTCTGGTTAACTGAGGTTAACGAAGTAAAGA
GCTGCAGACACTATTAACCTACTACCTTAAACCGATTACTCTAGCTTAGCCTACTTTCCAGTACAGATTTTACCAGTGGACAACATGAT
GCTTTATCTTGTTTTTCTCTCCCTGGGACTTTTCTCCAGACATTGAAAACAGAAATACTAATAAGGCCACTTTTACCTGCCTGATGCAA
GAACAGAAATTTCAAACCTCAACATTAATGCAACTCCTCAGTCCCAGCAATGGCGGGTGGAAAAGTTTCTAAAATAATGCAGCAGCACA
ATTATCCGGAAAGAGATGAGATACTGTTACCTAATAAAAATGCCATAAATAGAGAATGATGAACTACCATGGGAAATGAATGCATAGAAG
AGGACATGCTGGAATGTGGGACAGTAAAATCACTTAAACTTTGCGTGACCTTGAAGAAAGTCAGGATGATCTGTTTTTCCAGGTCCCT
CAAACAGTGAGATGTGCTGTTTCCCAAGTCTTCTCTCCAGTGTAAAGGGTCTGAATTTAGACCTTTGTGAGTCTTCTTCTTTGGA

FIG. 3D

CAATAGGAAAATAAATTATTGAAATAGAGGAAGAGACAGGTAATAATAGAGGTATACACAAGTAGAATGGGGCAATAAATGGCGCATT
TCGCACCATCAAGAGTGCCCATGTAACAGAGATAAGTAAATGCATCTTGAGCTGAACACTGAAGGATAAGAAACAAAGGGGAGAAAGAC
CTAGAAGGGGCAATATACAGCAAGGAGGGAAAATAAATACTACTGTGCATTTCATGCCAGTGTAGCATTTAGGACATCTGGAAGCTAGAGG
TGGAGTGGAAAAGGAGAGAGTGATAGGAGCTGGGGTCAGAGAGTTTCAGGGTGGGGAAAGGTCTTGCAGGACCTTGTAGGTAATTGTAAA
GCATTTGGATTTTATTCTGAGGGTCACTGGGGTGTCAATTAGAGACTTTTGAGCAAAGAGGTACATGCTCTGACTGAACTTTATTCTGTG
AACAATCAGAATCAACTAGATGGATTTAAGTATGGGTATACCATGAAAGAAAATTACTTAAGATCCTTGCTACTCAAAGTATGAGCCAG
GACCAGCTACACTGGCATMAGCTGGGAACCTGTGTAGAAATGCAGAATCCCAAGTCCCCGAGACAAACTGAATCAGAACCTGCACCTTAA
CAAGATCCCAGGTGGCCCATTTGTATGGTAGAGTTAAGAAGCATTGGTTTAAAAGATCCCTCTTGATAGGAGCATGGAAGATACATTT
GAGACAGAATAGACAAGTCAGAGACAGGTGGGAAGGGCCTAAAACAGGGCAGAAGTAGGGAGGTAATGAGGAGACAAATACAAAGGAA
GAAAATGCACAGCACAGTGTAGACAATTCTAAATACTTAAAAAAATTTTTTTTGAATAATGATAGATTCACAGGAGGTTGCAAAGAA
ATGCGTAGGGAGAACAATGCACCCCTTACCCAGCCTCCTCCATCATTAAACATCTTATGCAACTATATTATAATATCGAAAACAATCAA
GTGACATTGCTACAACCCATAGAGCTTATTAGATTTCACCAGTTATTAGATGCACTCGTGTGTGTATGCATATAGCTCTGTGTAAT
TTTATCATATGTGAAGCTTTGCTACCACAATCAAGATATTCAAGCCATTAGCAGAAGATTTTCTGGTGTACCTCCTTATAGCCACAG
CATTCTCCATCATTAAACCCCTGGGAACAATACTGTTCATCTCTATAATTTCTATTTCACGAACATTTTGTAGATGGGTACATG
CAGTGTGTATCTTTTGGGATGGTAACAGAGCAAGACAGGATCTCACTCTGTCAACCAGGCTGGAGTGCAGTGTCTGTATCTTGGCTCA
TTGCAGCCTCCACCTCCTGGGCTCAGGTGATCCTTCCACCCAGCCTCCTGAGTAGCTGGGACTACAGACACACGCCACCTCACCTGGC
TAATTTTTTGTATTTTTATAATGATGGGGTTTCACCATTTTGCCTAGGCTAGTCTAGAACTCCTGGGCTCAAGTGATCCAACCGCCTTG
GCCTCCAAAATGCTGGGAGTACAGGCATGAGCCACCACCTCCACCAGCTTTTTCATTACATCTTCTTTGAAGTTCATCCAAGTTGTG
TGTATCAATACTTCACTCCTTCCAGTTGCTGAGTAGTATTCCATGGCTTGGAGGTGCTAGAGTMTATTTCATCACATTCACCCATTGAA
GGMCATTTGGGTGGCTTCCAAGTTTCCAGTTTGGGCTATTATGAACAAAGTFACTATGAACATTCATATACAATGGATACTTTTTGTA
TGAATGAATGGAATAGAATGGATAGGATTTAGTGATCAGCTATGTGGGATGAAGAGTGGCATAAGTAGTAAAAAGTAAACCCCTCAATGCA
ATGTGCAGCCAGCAAGTACCACAAAAGAGTTTATTTGTTTCATACATATATTTCTATATATACATACACACTTTATTAATAACCA
AATAGTATCCTTTTCAAATGAAAACAGTAATTTAACATAAACTATGAACTTAAAATCTAAAGTAAAACCTTGACAACAGTGATGCAGAAT
TTTTTGCTCCTTAGCTCAGTTAGGTCTGTGTTCTTATCTTATGACCAGGAAGAAGTACTAGGTACCCTGACATCAAAGAATGAGTGGCATAG
AATTTATTAAGCAAAAAGGAAAGCTCTCAGGAAAGAGTGGGGTCTGAAAGCAGGTGCTGGTGGCCCTTCGTAGTTGAATACAAGGG
CTTCTATATAAAACCTGATGGGGCCGAGTTCCCTGTTCGTATAAGGCATGAATTCCTGGTGGCTCCACOGCCCTCCCCAGTGGGTATG
TGGGACCTTCGTCCACTAGGGACATGTTTAGACAAGCTCCCTGTGCAOGTTCCCTTATCTGCACAAAACATGGGTTGGAGGTTCTCCGG
GGACCTTCCTTTACTTTCTGCCATAAGCAAGCTGGCTAACTCCTTTCAACAATACTAAAGACATACAGACAATGGTTCTCAGTACAAT
CATTTTAAATATTTAAGTAAACTTAAAATGGTGTGTTGTTTTGATTTGACATTTTAAAAGATATCCTGTCTTAAAAATCTGTGTTTTT
AGTTGTTTTGGGCTCCTATTCTACAATGTGCTATTACTATTAAGCATTCTTGTATCATGGCATTCTCCTCAAATAAGTTTTTAAATTA
AATTTGAAGAAGGAACATTCTGTACAGTCAAGGAAAGTGTCAAAAATGAAAATGAGGGCAGGGTGTGGTGGCTCACGCCGTGAATCTCCG
CACTTTGGGAGCCCTAGGTGGGTGGATGCTTGAOCCTAAGAATTTGAGACCAGCCTGGGCAATATGGTATAACCCCTGTGTGTACAAAA
AATACAAAATTAGCCAGGTGTGGTGGCCCAAGCCTGTAGTCCCAOCTACTTGGGAAGTTAGGGTGGGAAATCCTAGGTGACAGAATGA
GACCTTGTCTCAAAAAAAAAGAAAAAAGAAAATGATAAAGGATACATATCAGGAAAACATGCATGGTATTTTGTATCATCTACTTTA
GAGTAATTCAGTATAGTGGTTTTTTTGTGTTGTTGTTTTGTTTTATTTTTGAGAAAGGGTCTTGGCTGTCAACCAGGCTGGAGTGCAGTG
GTACGATCTTGGCTCACTGCAACCTCOGCTACCAGGTTCAAGCCATCCTCCCAACTCAGCCTCCAGAGTAGCTGGGACTACAGGTGTG
CGOCACCATGTCCAGATAATTTTGTATTTTTTGTAGAGATGGGATTTTGCATGTGCTTGAATGCTGGCTCAAGCAATCCACCCCTC
CTCAGCCTCCCAAAGTGTGGGATGTCAGGCGTGAGCCACCACACCCAGCCCCAGTGTAGTGTGTTTTCTTTTTCTTTTTTATTCTATG
TTTTAATGAATTTACAOGTTACCCAAATGTTCCCTAGTTTTTCTGCCCTTCCAAGATCACTCTGGAAGAATATTTAAGAATATACCAAAT
AAGAATATGCAAGTCTCCCTAAGGGTGGCAGGAAGAACACCCCTCCCCAGATGGTATTTAGCGCCTCTGGCTGGGAACGGCTTCCC
CATGCTCCTAGGTCAAGGCTCTCTTGGCATGACACTACCACCACAGTGCAGACCCACAACAGGGAGAAGGACGGCCACAGTCCCTCA
ATCCCCCTTTTCCAAGATGTGCACAGCCTGACTCCTAACTCCCCACCCTGACTCTAGGGGAAAACAGCACAGGGCAGGAAACGATTT
TCCATGTCAACAACCTTTCTCTGAGGGAACTACTGGCCACCTCCCTCTTAGGACCAGCCCATCGTCCACAACGTGGAAAGTCCAGCTTC
CGTTCAAATOGGAGTTCTTTCTTCATGACATTTCTTTGCAAAGTCCCGGAACCCACAGCTCTGAGACTCTGGCTGTCCCCAACCCACC
CCATCTTCTTGTCTCACCCTGGTCAGGAGAAGCCAAAACATCAGTCAOCTTCCCAGTAATCAAGCCTGGCTTTCTCACCCAGGGCT
CGCCCCAGAACAACCACGGCTTCTTTCAAGTGTAGCCAAAAGGCTATTGGAGTCTTCTCAAATGAAAGAGATTTTATCAAAGGCTTGG
GAAGAAAAGAAAAGAGGATTATATAATAAAACGTAAAACAACAACATATACACACAACAAAATAAACGTGAGATATGATTCTCCC

FIG. 3F

GGAGTGTTTAGAGCAGGAATGTTCTTGGGCATCTGCCTTCCCCACCAGCACCCCCACAAGGCAAGGCCAGTTCACCCCTCAGTGCTCA
CTACTTTGCAGTGTTTCATAGAATATTTGTAATAATTTTAGGGCGCTCCCTAAAAATTTCTTTTCTTTTCTTTTCTTTTCTTTAGAGTTGGC
TCCCTCTCGGTTGCCAGGCTGGAGTTCAGTGGCATGTTTCATAGCTCACTGAAGCCTCAAAATTCCTGGGTTCAGTGACCCCTCCTACCTC
AGCCCCATGAGGACCTGGGACTACAGGTATGCACCGCTATACCCGTCTATCTTTTATTTATTTATTTATTTAGAGACAGAGTCTAGCTC
TGTCACCCAGGCCAGAATGCAGTGACACGATCTCAGCTCACTGCAACTTCTGCCTCCCAGATTTAAGGGTTTCTCTTGCCTCAGCCTCC
CTACTAGCTGGGATTACAGGCTTGCACCACCTACGTCCGGCTAATTTTGTATTTTATAGTAGAGATGTGGTTTCACCATGTTGGCCAGG
CAGGTCTCGAGCTCCTGACCTCAAGTGATCCACCCGGGCTGGCCTCCCAAAGTGCTGGGATTACAGGCGTGAGCCACTACGCCAGCCT
ATTTTATTTTATAATTTTGTTTTAGACAAGGTCTAGCTCTGTGCTGGGCTGGAGTGTAGTGGTGCAATCAAGATTCAGTGGCGCCCT
GATCTCCTGGGTTGAGTGAGCCTTAGCCTCCTGTTTAGCTGGTACTACAGGTGCATGCCACCACCTAGCTAATTTTTTAAAAATTTT
TGTAGAGACGGGTCTCACCCCTGGTGTCCAGGCTGGTCTCAAACCTCCTGGGCTCCAGTGATGCTCCACATTTGGCGTCCCAAAGTGCTG
GGATTATAGGAGTGAACACTACTGTGCCAGTCTTTTTAAAAATTTTCAAGAGATTGGGGTCTTGCTATATTGCCAGGCTGGTCTCCAC
TCCCTGGTGTAAAGGATCCTCCACCTCAGCCTCCTGAGTAGCTGGGATGACATTACAGGCACACACTGCCACCCTGGCTCTAAAAC
TTCTTCTGTGCCATTTGTGCACTTCACCCAATTGCCCTCTTTGTAGTAATTAATTAGGATCTAGGGTGAAGAAAAAGTCAACAGCTATAT
ATAGTCCCTCAAAGTTTGTACGTATCTGAGCAGTCATCAGTTGCACAGTGCCAGAGGGATGAACGCGTCCCGCCACCTAAAAAGCATT
AGTGACCATCAGGGAACCGTCAGATGCATGCCAGACTAAAGCAGAGTGAGGCTGTGCTGGGTCTGTCTGTGGCTGCCCGTGTCTC
ACTTCCCTGTCTTGTCTGTGCTTTGGGAGGTTGACCCTGAGTTGGCATCTCAGGGTCTCAGTCTGCTGGTTTCCCTGSGTTCCCTTG
AAGGCTACTGCTCCCACAAGGCAACCACGGTCCCGCTCTGGCTCTCACTGAGCTCCAGAATCATTGTTTCTCCCTTACCCAAGTGA
GAATAATTATGTTTTATTCAGAACCCCTGACAAATGAAGAGGCCATAAAAAACCCCTAGGTATTATCCGATCTTGGTGATCAGGGAGGTG
TTTGTTTTGTTTTTTAATGCAGACACATAGTTTTAAAAATTTACTTCTACTGTAAGAAAAGTCAATTAATTCACAATTTTGA
TTAAAACAAACAACAACAACAACCTTCTGTGACATTTGGCTAACAAAGTGGTTCAATATTAAGCTTTGTCCACCAGGTGCAGTGGC
TCATGCCCTGTAGTCTCAGTCTTTAGGAGGCTGAGGTGGGAGGATCACTTGAGGCCAGGAGGTCGAGGCTGCAGTGAACCATGATCTCA
CTACTACACTCCAGCCTGGGCAACAGAGTGAGACTCTGTCTCTAACAAACAACAACAATAAGTATAGTTCTTTCAAGCATGGCAGA
CAATCTGTCTCCTTTGGCCTGGGTCTCTCACTGCCTTTTAGATAAAAACTGGCAATAACCAAAGAGTTTTTATAAGGCCCTGTGATCT
ATTTATAAGACATGCATATAATTTACTTGACCATTATAATACCATTATAATAATCTAAATCTATTTCTTTATCGTCCAATAATCCACA
GAGTCAGCACACAAGGATTCTTTTTTCCATATATAGGCTGAGTATTCCTTATCTTACATGCGTGACGCCAAAGTGTTCAGGTTCTGGA
TGTTTTGGGATTTTGAATATTTGCATATACACAATGAGATATCTTGGGGATAGAACCCTACATCTAACACAAAAATTCATTTATGTTTC
ATATACACCTTATACAGTACCTGAAGGTAAATTTACACAATATTTTTAATAATTTTCCACATAAAAAACAAGTTTGTATACATTGAAC
CATCAGGAAGCAAGGTGTCCCTGTCTCAGCCACCACAAGGACACTCTGTAGTGTCTTTTATTCTGATTCCGAATTTATACGCTACT
GACAAGCAATCATTTTCTTACACTTATTACACAAGAGCACTTAGTAAAAATATGACATATATATCTGGCATGCTCAGAAAAGCTATT
TTGCAGCAGAAAGGAGCTGGGAGGGTCTTTTTTCCCTTGGGACACGGAAATAATTTGTGATTATGTGCCTGCATTTTACTGTGAC
CCCATCACATGAGGTTAAGTGTAGAATTTTCCACTTGTCTCTGTGCTTAAAAAGTTTAGATTGGCCAGGCATGGTGGCTCATGGCTG
CAATCCCATCACTTTAGGAGGCCAAAGCAAGTGGTCAATTTGAGGTCAGGAGTCAAACCAGCCTGGCCAACATGGTGAACCCCTGTCT
CTACTAAAAATAAAAAAGTTAGCCTGGCATGTTGGTGCATGCTTGTAAATCCCAGCTACTCGGGAGGCCGAGCCAGGAGAATCTCTTGA
CCTGGGAGGCAGAGGTTGCAGAGACAGAGATCACTCCATTGCACTCCAGCCTGGGTGACAAAGCGAGACTCTGTCTCAAAAAA
AAAAAAGGTTAGATTTTGGAGCATTTTGGATTTTGGATTTTGCATTAAGTGTGTTCAAGCTGAAAAGAAAATCCGATTTGCTCAGGA
CAAACCTTAAACAACAAGTGAATATTTCCAATACTATATATATGCTCCTGTTTATATTTCCCTTAATTAATTTGGACTTGGAAACA
GCCAATTATGGATTAGAGGATGAGACTTAAATGTTACTGTACAAGGATAGAACGATTCATTCCCTCTATGTTATCAAATACTTATGGTA
TTTTTCCCATCCTGCTGTATGCAGATCCAAGAACCAATTTAAACACATTTGCCGGGTCATAATAATGTGGCCAGAATTTAAAGAAA
AACTTGATTTTTAATTTATGATGATTTGCTTGTTTAGTCTACCGATTTCTATTTGCTTTAGCTTACTCAAAAAATAAAGCGGGCACTT
CGAAGACTCAATAGTCTTCCATTCAATGTTGGCCTTTATAATGCACGGGCCAGATGCAATACATCTGGCGGTCTGCTTGGGTTGGCCAC
TGGATTGAAGGAGGCAGAGAAGTCTGGGATGATTTCCCAAATGTCTGGATCTGGTGACAGGGAGATATGGCAGGGCGAGCTTAGGGGAAA
AAGCTGGGTTAGGAACGTTGAAACTGAAATCCCTGAGGSYTKTGCCGACAGAGAGACAGCCGGTAGAAGGTTGTCTTTGCCTGTCTGT
GGTCCAGGTAACCTTCATCGAAAGAGAGTTTTCAGGCAGTAGAAATAAGAGCACCCAGGACAAAGCCCCAGGGAAGAGAAACATCTGACG
GAGGACAGAGGAAGAAGGTCAGGAATGAGACTGAGCAGGTGTCTGTGCTGACACCAGAGCCTGACACATAGTACGTAGTAGACT
CAGCAAATACCGTAAACAGAGATGAATCCAAGGCTGGGGAGGTTGGCTCACGCCGTGAATCCCCACACCTTGAGAGCCCTAAGTGGGAGG
ATCTCTTGAAGTCCAGGAGTTGAGAGACCAGCCTGGGAACATGGTGAACCTTCCCTCTAAAAAATAAAAAATTAACATTAAAAAAGA
GATGAATCCATAACCTGGCTGCTGGAGCCAACATGGGTTGGGTGAGCCCCTTACCAGCAGCTAATCAAAAAATTTGCCCTGGAAATCT

FIG. 3G

GAGGCTCCTGTCTACGTCTTGGCTGCTCCTCCCAGATCACCTTCTGGCCGGTCCCAAGTCCACTTCCCGTGTCTCTTGTCTCCCTTCCT
CCTGGTCTCCCTCACACTTTCCCTTCCCTACTCCCCTTCCCTCTGTGGCCCTGGCTCAGCCCAGCACAGGGAGAGCCCTGTGCCACCTAT
TACAGCTCACCTGCACCTTTGCATCTTTCAGAAAGGAGCACCTACAAGATAACCCACCCCCACCTTTTTTTTTTTTTTTTAGTAGTA
CAGATTGCCTCTCATAGCATAATTGGGCTTCATTATTATCCTTAAAGACCCCTTCTCTGTGGGGATTGGGATGGATAAAAATAAAGAAG
ATCGAGAGGTTGAAGAACCCATCCTGTTTTGCCAGTGAGAAGGGGATAGAATTAAGGATTAGGAGGGCTCAGGCATGGTGGCTCCAG
NGTGTCTATCCCAGCTACTCAGGAGGCTGAGGCGGGAGGATCACTTGAGCCCAGGAGTTGGAGACTATAGAGCACTATGATTACACCTGT
GAATAGCCACTGCACTCTAGCCTGGGCAACATATCAAGACCCCTGTTCCTAGGGACAAAAATATNNTTAAATAAATTAAAAATTAAGGG
AAAGGTAACCACATCCTGCTACAAAANAAAAGAAGNTGGAGAGGTANGANGAGGACCAAGAGCTAATGGCATCATTTACACAAAAAGAGA
TGCTTTAAAATCAGTTGCTCATCCAATTCCACAAGGACAATAAGTAAGAAAGAGGATAGAAAGTCAACCGTGGATTGGTGCATCATTGG
CTTCTTGATGACTTTAGCAACAAAAATCTTGTGGTAGTGAGAGTTAGACCCTGGTGGACTGGGTAGGGGGTTCTGGATCATGAGCA
AAGGCTGTGCCAGCCAATGGCCCCACTACACTCTGCCCCGGCCTTCTCATCTCAAAAAATGGCATCCCCATCCAAAGCTCAAGTC
AAGAATCCAGCAGCCACCTTTGATTCTGCACTTCCCCTCACCTCACAGTCCAGTCCCATCTCCAAAATAAGTTCCAAATYTCACCACCT
CTCATCTCCAAAGAGGACHATTATCTCTTCTGGTGATTAACAACAGCTTCTAACTGGSTTCCCTTCTACCTTGGCTTCCCATAGT
CCATCTCTCAGGACAACAACAGTGGCCCTTTAAAACCAGTGCATTATTGTTGCCCTTGGGAAATCCTCCACAATTATCCAGTCTTG
CTTCAAAAAATGTATGTATTTCTGACTTTTACCCTGCCCTACTTACAGGATATGCACATTTCTGATCTCCAGCCAATATCACACTTCT
TCTCTCACTGCACTCTGCCACACTTGGCCAAGTTTGTTCCTCACTCCTCTTGCACTTGTCTCAGATCTCAGAAGAGGGCGTCTCCTTGT
CTTTCAGGCCAGCCGGCTTACACATGTGCCACGTGCGCCCCCTCGCTCAGAAGGGATCTGTACTCGGTTTGGATCTATTGTTGCCATCT
TGAAACTCTTAATACTCTTTGAACACGGGGCCCGTATTTTCATTTTGCACCTGGGTCTGAAAATTTGTGTAGCTGGCTCTACTTTCAGGG
ATTGTATCAGAAGTCTCCTCCTCAAAGAGGCCTTCCCTCGGCCACTTATCCTCAAGTAGCTCCTCCCCCTCTAAGTTACTGGCTATCCCA
TCATTCACCTAATTTTCTTATAACAGTTGTATGCTTTTATACATCTGGCTTCTATATTTATTGTTGTTGTTCCAGTCTCCCTCC
CTTTGGACCCAGCTGTGGCACCTGCAACCCAGAGACCCTGTATCCCGGTGCAGAAATGTAATGAGTGCCTGATACATTTGCCGAATA
AATATTCCAAAGGTTGAACCTGCTGGAAAGCAGAGAGCACTATCTGGTAAAATGGAATTTAAATGTACTTGATATTTATATA
ATCCTAATCAATAATTAATTTGTGTAGTGCTGATCTAAACAGATAAATTTCTGGCTTCATGATGATGGTGAAGTGGAAATATAATTTCT
CATTTGTATTCAAACCTAGATCTTTTTCATGAAAGGATTGAAGTCTAGATTCAATGCCTACTTTTGTACTTATGTTATATGAAACTA
AAACAATTATTTATTGTTATTTTTTGGAGATGGAGTCTTCTCTGGTGGCCAGACTGGAGTGCCTGCTGCAATCTCAGCTCACTGCA
ACCTGCACTCCCAAGTTCAAGCBAATCTCCTGCCCTCAGCCCTCTGAGTGGCTGGGACTATAGGTGGCTGCCACCACACCAGCTAAT
TTGTATTTTTAGTAAAGATGGCTTTCACCATGTTGGCCAGCTGGTCTTGAACCTCTGACCCAAAGTGATCTGCCCTGCCCTCCGCTCC
CAAAGTGTGGATTACAGGCATGAGCCACTGTGCCCTGGCAATAATTTTAGTTAGTCTGAATTTTTTTTTTTTTTTTGGAGATGGAGTCTC
GCTCTGTTGCCAGGCTGGAGTGCAGTGACGCTATCTCAGCTCACAGAACTCCGCTCCTAGGTTTAAGCAATCCTCCTGTCTCAGC
CTCCGAGTAGCCAAGATTACAGGCACCTGCCACCACCCCCAGCTAATTTTTGTATTTTTTAGTAGAGATGGGGTTTACCATGTTGACC
AGGCTGGTCTCAAACCTCTGACCCAAAGTGATGTGTCTGCCCTCAGCCCTCCAAAATGCTGGGATTACAGGCTGAGCCACTGTGCCCTGGC
CTAGTCTGAATTTTTTAAAAGGTTATTGGTCTACCTTCCAATGACATTGCCTCTGTGTGGCTCAATAAAACATTTTCATTTATAATA
ACTAATTTGACCTGCTCAGCAATCTCTAAGCAAGATAGAGTAGCTGTAATTTCTTCAATTTTACAGGTCATGTCAAATCATTTCGTACATT
CCAGCTATGTACGAGAGCTTGGTGAGAATATGTGAATAATAATCACAGAACTTCAGAGCTGGGAGTAACAGCTGGAAATATTTCTTCCA
ATAATTCATTTTTTATGAGAGGACGATGAGGTCCAAAGTGGACAGGACCATGAGACAATCGTGTGCCAAGGAAGTTGATGCAATTTGAC
CTCTAAGTCACTGATCTTTATGTCCATCGGTCCTTCCAGCAAGTGAGTTAGCCAACTTTGCCCTGCAAAGGAGGAATTTTTAATTG
AGGATTTACACTCTGCTTCTAAAATTTTGTCTTATTATTGTGAATAATTTCTTTAAGTTTATTAAATGAATGGCTGAATAAATGGACAT
AAGGAAGAAGGAAGGGAGGAAGGAAGGGAGGGAGGKAAGGAAGGGAGGGAAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGA
AAGAAGAGAGGAAGGAAGGAAGGATAAGTCTGATGACAGCTGCTATTATATTCTACGTGGATAAATTTATTAGATCTTTATACTTTAT
CTTTGTTTTACTTCTCTTATGCATATTCTCCTCAACTTTTTTTCAGTGGGCCAGAGGAGGAGGACTGCCCTCTTGTGACTGTGGAAAGGA
CTTCTACCAGGCTAACACCCCTGGCCTCTCACCTCCCATTTCTCACCCCTGCAAAGCAGAGTGTCTATTGATTGATGTTCTTAGTCTGT
GGATCTCAGTTGAGGAGAACTCGTTAGAGATTTGCCCTCTTCTGTCTTTTTGAGACCTTACTGGTGCAAGACAGCAAATCCTAGCTGG
TGTCTACAGGACACATGCACCTTFAAGTTACATAACTGCAGGGACCCTGTCAATGTTATCCTGGAGCTGGTCTATATAAGACACAGCC
TGAGCAGTATATAGGCTTCCCTAGTCTGCTCCTGGCCAAATGTCCCAGTTGGAAGCCCAGAGGTTGTCTGGCTATGCCAGTGGCAGGATG
GGCAAGTCTAACTCAAGGGTGACATATTAGCAAGACCTTTATGGCCATGCATCTAAGATGCTCTGTCCAAGCCTGAACTTAGCAACAAT
AAACCTGACATTTTGAATCCATCTGATTCCCTCTATTTCCAGTTGATGCCACATGCATCCTCTTCCCATCTTCTTAATTAAGATGAC
TTTCTTCTAAATCTCCTTAATTAATCAAGCAGCTATCTACAATATTTGTAATCCCTTAAATCTTGAGCATAATGATGTCATAATTAT

FIG. 3H

GAAAGTGMCCGGWTTACATGAAGTATTGCTTAATCTTAAGAACAAAATGGCAGCTGTGAAAACAGATGAAGTAATTAGAGGAAGAGCC
 TTTTGGAAAGCTTCGAGATATTTCAAAGTAATTAGTACTAGTTAGCAATAAAGTTCTGTTCTGAGAAATTGCTCTTAAAGGAGGAACA
 TGGATTAAGAAAAAATCTGCTACTAGGAAGTAAGCCATCTTCCTATGTGTGTGATTGGTTTTGCTTCTGAAAACCTGGTCCGTTTT
 CAACAAAATTTGGGTCTGTTGAAAAAGAACACGCAGATGCCAGCCTTGATGTCAAACGGGCCCCAACTTGGACAGTGGTAAACTAATGA
 GCAATGGTGCACAGAGTCAAGGTAAAAGCTGGACAATTTCCCTATGACCAACTTTTCCAGGACTCTGCTCTGCTCTTCCCTGAGAAAAATA
 CCCAAAGTGTGCTTCCATTTGGCCCAACCATGCATCTTTCAGGATAGGMCACATCTGTTTATAGGTGTGGATTGTAGTTGCTCATA
 AGTGACATTAGGCTGTTTAAAAATAATAATAGTTCGAGTTTTGCTATGAGCTGATCTGTTTTCCAAGAGAGCTAAGAGTTTCCAGCTAA
 AAGAGGGAATTAGTGGGTAATCAAGGCAGCTGACATGGGGTGTGGCTGGGCCTTGAATGTGTGTCACTCTCTGTGCCAGGCAGAGCAA
 AGATAAACTCCAGACTGCATGTTGCTCAGAGACCAGGACCAACGTCATAGGGCGCCTAAAAGGCAGGTGGCCAGTTCAGAATTGTCAA
 GGTCTGACCTGCTTGGACAAGTGTGAGTACATAGTAAGGATGGATTGGCTAGTCTCTCAAACTTGCAAACAGGGCGCAGGTGATCTT
 GAGATTTCAAGTGCAGGAGAGACCCATCGTGTAGATTCCAGAGTTGGCTATCATGACTAACAGCTGTCTAAGTTGTTTTAAATGAATC
 ATTAAGGGCTACATTTTCAGTTCAGCTAATCAAGTAGCAAATTACGGTGGGTCTAAAATACTTATCTATTGCATTATGTATATGCTAGA
 CTTTATCACTTTAGTTGGTTATATCGCTTCATATACTAACAGTCAAAAAATGCCAAACGAGAAAAACAACAAAACAAAATGCCACATGA
 CTGTGTAATAACACTTTTCAAACGTTTTATCTAAGAGTTTACTCACTTTACATTTGTGGCTTATAGTATTTTCAATCTAAGAGACTAA
 TTTTGGCTTACATAGGAACTACATATTTTAAATGAAAATTAATAAATATTTTTAAGGTTTTAATGAGTCCATCAAAACACATTTGT
 ATATAGGAAGGTAGCCCAAGGTCACTGTTGCCAATTGTGTACACAGCCTGCCCTMTAGTGTTTTCTTCTAAACAGCACCAAATTTTAGA
 TCATAGTTGTAATCTCAAAATGTGGGTTAATAGGATTAACACTGTGTCAAAATTGATAGGACACAGCTAAATCCCTGACACGGG
TGAAAATTAAGCAGAGAAAAACGAAGGTCCCTCCAGAAAGCTGGTGGCAACTTCACTGGGGAGATATTGCAAAGTTAGTGGTAAATACA
 CTATATTAAGGTTTTGTTTTGTAATAGAGTAATGATAGAAGAAGAGTTAGTTGAAATGATGTATGTAATGTGATAACTGCATAA
 TTACTAGTACAGTTGCTAGTTTACGACTGTATTAAGAGACATTCCAAATGTTGATCAAATAATGGAGGTTTCTGTGGTTGTTTTCTTT
 TTAATAAGTAAATATACGTAAGCAGATAAATATCCCTTTGTGGGAGTTAAATAATCTAACTTATTTTATAGTTTAACTTTATTA
 AAGCATAAGACTATTCTAACTTATTTAACTTTTCTTAGTAAAGTTTTAACCTCTGTATTTAGAATATTTGTAACATAATGTGTATCGAAT
 TAACTCAAAGGGAAATTCATTAAGTGAAGAAAAAATTTAACTGTGCACTATTCACATAGCATAATGGGTTTTATAAGGAGTATGA
 GAAAAATGTGTGTGGTTGGTTTTGCTTTCTTAAAAATAATAGCGAACCCAGTAGGTAAAACTCACTTGAGAACATAGACTTTTGGAG
 GGAAATGCCAGGTGTGGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGGGGGGGATCACCTGAGGTGAGTAGTTGAGAG
 CCAGCCTGACCCACATGGAGAACTCCATCTCTACTAAAAATACAAAATTAACCGGGCTTGGTGGCGCATGCCATAATCCCAGCTACT
 TGGGAAGGCTGAGGCAGGAGAATCACTTGAACCTGGGAGGTGGAGGTGCGGTGGGCCGAGATCACGCCATTGCACTCCAGCCTGGGCA
 ACAAGAGCAAACTCCGTCTCAAAAAAAAAAAAAAAAAAAGAATTTTGGAGGGAAAAAATCCCTCTAACAGATTGGAATTAATTCT
 GTGTTTCGAGATGTTTACAAAATGAAGCTTGGACTCTGAGAGGATGTGATCTATCCTCTCCATTGCATTGAGTTTCAAGTACTTCACAT
 GCGGGCTTTTTTAACTGTGCTGAAGTTTAAACCAATAGGGACTAGAATTTGTTTGTTTTTTAACTTACATTTCAAGCTTCCCTATG
 TCTCAGGCACATTAGCATAAGTTGTCTAAAGTCATAAGGAAAAATGACAGAAAAATGCTTTGGAGCCCCAGGTGTTTTCAATTGATGC
 CAACAGAACTAACCAATGGAAGACATTTGATGCGGGTTATTTTTCTTTGCACTAACAGCGGGAACATGAAAGCCGCCACTCTGGT
 GTTATGTTGTGCTGCTGTGGTTGAAAGACAGTCACTGCCACCCACTTGGAAAGGACAAAACCTGCTATCAGTGAAAACCTGAAAGGTA
 CGTTTGGTTTTCTTACCTGTGCTGTGCTGTTTGCATGTTGGTTGTCTGCTGGCGTTTATAGTGAAGTCCAGTTGAGAGATAACCATA
 TTCGCTGTTTTACGGTGAACGTTCTCAAGGCCCTTAAACCAGGTCACTCTGACGCCAAACATCTGGGTAAAAATAGAAAATCCAAAT
 CACGTCTCTGCAGGCCCTTCCAGATGTTTGTATCATGTAGATACAACTTGCCAGTTTTTTCACTGCATTTTTTTGTATCATCC
 AGATGGTTGGTGTCACTCAGCACAGCTCTAATGAACAGTGAATACTTTCTAGCATTTGAAAAATTTAAACCATTAGAGTAATCTGT
 GCAATTGTTCTTAACTAGTGAAGAAATGGGTTATAATTACGTTGAATCTGGTTGTTCTGTGGCCATTAACCTTGAACCTTTGCTTGGTG
 ATATATACTTTGGGTACTTAATATATAGAAGAACAATTAGCTAAAATGCAGCTGATTTGGGCTCTGTAATAATCAGAGTCAAGAATGA
 GCTCCTCAGTAGGCCAAGTTGGCTATTTTGAACAGGGAATGACAATGAATTTTAACTTACTAAGGGCTTATTAAGGTGTATAAGACA
 CGTCCATTGAGTTATTAAGGAAGCTCGTATTACATGGGATACTTTCTAGGTCTCGTGCCTCCTTATTAGGTAAGCTGAAAGAAA
 GAGAAATGCTGACTGTGTTTGGAGTCCCCAGCTGGGCCTTAATATAAATATGAAGAAAATGCAAAATTTCTCTAATATAAACACA
 CTTGAGTCTTAAATGAAAGAAAAAATGGATAAATGAAAACAGGGCCTGAGCAAGTGACAAGAATGAGGTTCAAGTGAAGTCTATTTGT
 TTAGGCGCTCACAAGTGAGGAGTAGAAGGTATGGTCCGTGTGGCAGCTGTGTCCATGTGGCAGCTGACAGCTAATTCATTATGATCTGC
 TTTCAGAAATAGACCTATAAGAGAACAATTAAGCCTCTCTTTTGGAGACATGAAAGGTTGGTGAACCTGGTGTTTTTGTAATCTGATCA
 GATCTCAAAGAAAAATGCCACATGTCTTTTAGGTTTTCTGAGGTGGGGAGATAGATCCAGATGAAGAGGTGAAGAGGCTTTGAC
 TGGTATTAAGCAATGAAAATCATGATGGAAGAAAGGAAAGGAAACACCAATCTAATGAGCACCCCTGAAGAAATCCAGAGAGAAA

FIG. 31

AGCAGGTACAGTCATTGAAAATAATGTCTGTTCTTACACAGATCTGGACCAGAAATACTGCACCTGTTAGTGGGATTGATGAATTACTT
 ATTTTCCTTAGTAATAAATTTTCATGGGTAGCTGCTTTTATTTGAGGAAAAGTTTAAGGGAAGCTTCAGATTTCCCTGAAGAACATATTT
 CGTGTAGGATAGGCTTCTGCAAGACTCCAACCCGGAATCTGGGGGATTCATCTCTGTTAAGTGTGCTTTCTCAAAAATAGATTATTC
 TTGGTCTCTTCTGAGTTAGGATATTGAGTCAAAAGTATTTGAAGAGTTTTTTTTTTTACTAGATCAGTGGTCTCCAGAGTTTTTGTMTT
 TTGTTTTTTGTTTGTCTGTTTTTGGACAGAGTCTCGCTCTGTCAACCAGGCTGGAGTTGATCCCGCTCATTGCAACCTCCACCTCC
 TGGGTTCAAGTGATTCTCCTGTCTCAGCTCCCTAGTAGCTGGGATTACAGGCTCCTACCACCAGCCTGGCTAATTTTTTGTATTTTGA
 GAAGAGACGGGTTTACCATTGCTGGCCAGGCTGGTCCCGAACTCTGGGGCTCAAGTGATCCACCTGCCTCAGCTCCCAAAGTGCTGG
 AATTACAGGCATGGACCACCGTGCCTGGCCAGAGATTTTTGGTCTCTCATTCCTATGACTAAAAAATTTGTTACCCTCACTCCTAAA
 TATATGCATATTCATTTACTCATGAATTAGATACATGAATTGCTACCATTGATATCTCAAGGCACAATATGTATTTAAGGTGAGATTCA
 TCATTAGCGAGTGTGGATATAAGTCCACATTTCAAATAATCTTCTAGATATTTTGAAACTTTTAGCCGACTTGCCAGATCTGATTAGAT
 CACCATAGTTTTCCCTGTCACTTGGCCAATAAAGAGCTCATAATGATCAAGTGTGAGCTCTGCCATTTGCTTTTGGTCCGCTTGAGCT
 TAAATTATTCATTTTTAAAATCTGCCAAGTTTTTTTTTTTTTCAAAGAATCTTGTAAAGCCTCCTGTCCATTTAGTGAAGGTACTTTA
 GTTAAAATAGATAATAAATCCATCAGTCTACCTGAGTCTCTTACATGGCAACTCATTACAATTGGGTGCATGTGAACAGAGCAAGG
 GAACTATAGTTGATTCTTCTGGAATGTAGAGGATCCCCCTTTCCCAAGGTCATCACATACAGTTGGGCACACACAGTATCTGACATAT
 GCATCTCAAGAGAGTACCATGTATATCCAATAATGCATCAGCCTAATCACTTTTTCAAATCAAATAGCTTTATTTAACAGCTATAGCT
 TGAACTACATATTTATCCATGGAGAATACATATTATTTCAAATGTCTTTGGAAGATGTAAAAAATTTGTTTATATGCCACAGTATAAA
 GTTCAGTAAATTTCTAAATFATAGACATTGAATAGCTTGCAGTTAATGACATTAATAATTAACATCACTCAAACAATGACTTTTTT
 TAAAAAGGTTATCTTCAAMCATTTCCCTTAAATCAAAGAGGAAATTAAACTGTAACAAAAATAATTTGGAAAATATTTTCAATTTTA
 ATGTTGAGAGTAAATTAATTTTAAATKTATTTTTTATTTTTTGA AAAATGTTAAGTTGTAAAAATACATATAACAAAATTTACCATCATA
 ACCATTTTTAAGTGTAAACGTTCAAGTAGTGTAAATACATTCATACTGTTGTGCAACCAATCTCCAGAAATTTTTTCACTCTGCAAAAAC
 TGAAGTCTATACATATTAACAATGCCCAATCCCCCACCAGTCAAGTTTTAATTTAAAAATACAAGTGGAAAGTTCTAATATTT
 TCTATCTATCCCTCTATCTATAAAGTGGGGGCCACTGAATTCAGATTGCTGCTTGCATCTTTTTACTTCTGAGCATCATGGCCTCTG
 GGAGTCCGTTAAGCAACTGGAGCCGGGTAGTGTGACAGGCTGACCCCAAAGCTGTGTGTCAGCGTCAACGGACTGGTTGATGTTGCAGC
 CTCACCTACTGCCCTGAGTCAAGTCAAGGTTCTGGCAAGGAAAGGAGAATGCCTGACCAGCAGCTGCAAACCTTCTCCCTTTTGGCAGC
 AATCAAAGATTTTGAGGAAATCTAAAATAGCTCCTCATCAGGAAAATGTGGAAGCCCCCTCCAGCTGGGATCTTCCCTGGTGGGCTGT
 GAGCTGGCCATCTGGGAATAGAGACACTAGATAGCACTCATACTCTTCAAAAACACATTATCACATGGAAATGTTTTGAACATCTG
 GGTAAACCACTACTTTTCATTTTATAGCTAAGAAAAGTGGGTTGAGATGTTGTTAATTAACATGTTACTCCAACACTGTAATGAATG
 AACTGAGATAAAGTCAAGCAGATGTGTGCACGGGGGACCCAGTGAATTTCTGCTTTTCTCACTTCCCTGAACCTCCTGGCAAGGAGGACA
 GGTATAACAGCTTTAACAAGAATATTCCACTTTGGGTGGGTCAAGTAAGCAAAATGTGGATTTCACTTCTGGCCCTGAAGAATCCAAGCA
 ACTAGTAGAATTTTTGTTTATTTCTTAAAAATCTTATGTACAAAATTCATTGAATTATACTCTTAAGTTTGAGGCACTCAATTAGAAA
 GTTAATCGGAAAAAAAATCTGTTAACCCTGAGTATCCCTCCCTAAAATTAATTAAGCCTAGAATAAAGGTCAGTTAGACAAAT
 ATGAATTGGCAAAATATGGTGTAGCAACCTAGTCTCCAGTATTGAGCCCCACCATTCTCAAGAGTACTGCTCAGTGGTGGACCCAGC
 ATCCTCACTGTCCCTTCCCTCCACCCCTCCTTATTAATATTTAGTGGACTATCTGAACTTATTAAGTAGGAAACCCCTAGAGAAGGTT
 AGAGTCACTTGACCTCCAAATCAGGTTTTATTTGTATGTGTTTTAATGAAATGGGGTCTGCTATGTTGCTCAGCTGGTCTTGAACCT
 CCTGGGCTCAAGGGATCCTCCTGCCTCACTTCCCGAGTAGCTGGGATCACAGGCACTAGCCACCATGCCCTGGCTCAATGCCAGGTTAAT
 ATAGCCCTTTTGATAAACTGTCAACTATAGGAATAGAGTTATAAGCGTGAATCTGCCAGTTGGTACAATGCTTAGCAAGAAACGGAAAGG
 CGTCGATAGGATATTCCTTAGGAATGTTTACTAGACAGAGGCTACTTCTTCCATGGCAATGTTTCACTTCCAAAATTTGGGACCTGTG
 ATTTGGTAACTGTTTTTGTCTGCTTCTGGCCAGTGAATGGAAGGAAGCCTGAGAGATACTAGTTATTATACTGGACTAGTTATAATA
 ACAGATGCTTGCCTATGATAATGGATACTAGGTATAATAATAGATGCCCTGCTTGTTTAGCTCATTAAATGCAAAGACCTTGAGAAGT
 AGATACTATTATTCCTATTATCTTATTTGCAAATGAGGAGACTAAGGCTTATATGTATTAAAGTAATTTGCCCAAGGGTACACAGOCAC
 TGTAGTTTGGAAATGGGAATATTAGGATTTTGGCTTATGAGGACAATGAGCAGAATATGTAAAATTTGGGACTGATTGAGAAAATCCTGG
 AGGTATTGTTACTTGCCTTGGAGAAAACAATTTTTTTTTTTTTTTTGGAGACAGAGTCTTACTCTTGTGCCCAGGCTAAAGGACAATG
 GCACGATCTTGGCTCACTGCAACCTCCGCTCCTGGGTTCAAGCGATTCTCCTGCTTCAGCCTCTGAAGTGGCTGGGATTACAGGCACC
 CACCATCATGACCAGCTAATTTTTGTATTTCTAGCAGAGACAGGGTTTTACTATGTTGGCCAGGCTGTTCTCAAATCCTGACATCAGG
 TGATCCACCCGCTCCAGCCTCCAAAATGCTGGAATTACAGTGTGAGCCACTGCACCCTGCCGAAAAACAACCCTTTAAGATGTTA
 GATCCAGCCAAGTGAAGTGGCTCATGCCGCAATCCCAAGCACTTTGGGAGGTC AACCTGGGCAGATCACTTGAGGCCAGGAGTTGGA
 GNTCAGCCTGGGNAANTGGTGNAACTCCGTCTCTANTANAACATACAAAATTTNGCCCGCATGGTGGCAGCACCTGTACTCCACG

FIG. 3J

TACAGGGCAAAGAAGAGGGTCCAGGAAAGCAGCTGGGAGAACTGACTTCTGGTCACCAAAGGGGATGGGTGCCTTACATGCCATTCT
 ATCAAACAGTGCCTTCACTGTTTTTAAACTATGGACTTTGCAATTTATCTCAAAATAAAAAGTTTCATTTTTAAATGCTGAGGATTTAAT
 ATGACAGAAAATCATCAGGTTGTAAATAGTAATACATGTTTCCCTAATGTCAAACACTCTATTGGGAACCGCCAATTTTCTGTTGGATA
 GACTTCTCTTTTACACATTTTTATATGGATTGTTAATTTCTCTAGGGGAAAAAAGTTCTCAAACTTGATTGGCTTTAGATATTTTCT
 AAATCTTTGACCCCTGTTTATAACAGTATATGCATCTCCACACACACATACTCGCACACATATGTGTGTATATATATGTGTGTGTG
 TGTGTGTGTGTATATACATATATATGAGAAATGCAAAAAAGAATAGTAATAAAATAACCACCTATCACCCACTTTAAGAAACAGACAT
 TTCTAATATCTTTGAAACTTCTTCCCAATTATAGCTTTAAAAATTAATTATTAAGAGTTTTTTAAATAACAGAAAAGTCCAAGAGAAA
 AAGTGGTTCACAATCACCTATTTACTTAATCCTATTGACATCAGAAATACTAATGATATAAGACAAATGATTTTTAAAGTAATCAAATA
 TATAAAGAACAAAATAAATGAAAGCTGCCCTCTCCTACCTTATCAACTCCCTCTTCTAAAAGATAGTTATTAATAATTTCTCATGACT
 CCTCTAGAAAATAAAATTACATGCATTAATATATGTGTGTATATACTACTAATAAATTTCTAGTAATGAGATTCTTGGATTCAAGAGT
 GTGCAATTTTTAATAGCTGTTTCAAGTTGTCCAGGAAATTTATGCACCAACGTGCATTTCTGTGTCTAAATATAGGAAAAGGGCCAGGG
 GCGGTGGCTCATGCCTGTAATCCCAGCACTTTGGGAGGCCGAGGGCGGTGGATCATTGAGGTCAGGAGTTCAAGAAAACCGGCTGGAC
 AACATGGCGCAACCCCATCTCTACTAAAAGTACAAAGATTAGCTGGGCTTGGTGGCTCTCACCTGTAATCCCAGCTACTTGGGAGCCTG
 AGGCAGGAGAATCACTTGAACCCGGGAGGCAGAGGTTGCAGTGAGCCAAGATCCCGCCACTGCATTTAGCCTGGGCAACAAGCAAGAC
 TCTGTCTCAAAAATAAATAAATTAATAACATACATATAGGAAAAGATTTTGAAGCACTGGTAAGAAAAGCTGCGGCATTGTC
 TCCACTTCTCAAAGTGCAAACTCTTATGACACTAACGTGTAATGTTATGTTCCCTGTAGCTCCTGACCACGGAGGCCTGATTTCAA
 GATGTTACCTGGGCAGGACAGAGGACTGTGTGGGAACTTGACCAGAAATTTGTCAAGATGTTTCAAATTTTCAAAAAATGCCAAAAT
 GTCAGGCTCACCTATCTGAAGGTAATAATGCTATTTTTGTTTTTATTCTACTTTAAGTTCTCAGGTACATTTTGTATAAAGTTTCG
 GTGCCACAAAAGAAATAGCACTCGAATATAAATTTTTCTTTTTAATTCTCAGCAAGGAAAGTTACTTCTATAGAAGGGTGGCCCTTAC
 AGATGGAGCAATGGTGTAGCGTGCATTTGCCAAGGGAGGGGAAGGGGTTCTTAACCCCTGACAATGCACTGGCCCTGCTGCTGTGTGGT
 TCCCCTATTGGCTAGGGTTAGACCGCACAGGCTAGACTAATCCCATTGGCTAATTTAAAGAGAGTGACGAGGTGAGTGGTCTGGAGGG
 AAAATGGTTATGACAGAGCATGTAATCGGAATGAATCAGGGCCGAGCGTGAATCGGAATGAATCAGGGCCGAGCATGTAATCGGAAT
 GAATCAGGGTGGAGCGTGAATCGAAAAGGTTGCTTTACGAGGAAATTAAGTTTAAAGTAGAAGGCAAAGAATTGAACATACTGACA
 TACTGATTTCTTTGGAAAGAAATTTAGAACTCACATCTAACAAATTTTTAGGGTTCTTTAGTATTTCTGGACAGAGGACAAAATCTCATT
 CTCACAAGCATAGTGGATTCAATTTGCTTTCCCTCCAAGCACTTTTTTGCAGGCTCATTTCCATCTGGGGGCGTTCAATGTAGGTTTATAA
 ACTGGTGTMTTGTMTTGTMTTATGAGACAGAGTCTTGCTCTGTTGCCAGGCTGGNGTGGCACAACTCTCGGCTCACTGCAACCTCC
 ACCTCTCGGGTTCAAGCAATTTCTCTGCTCAGCCTGCCAAGTAGCTGGGATTACAGGCATGTGCCACCACGCCCCGCTAATTTTTTTTT
 GTATTTTTAGTAGGGACGAGGGTTTACCATATTTGGCCAGGCTGGTCTCGAACTCCTGACCTTGTGATCCGCCCACCTCGGCCCTCCAA
 AGTGTCTGGGATTACAGGCATGAACCACCGTGCCTGGCCTGGTTTATAAACTTTTTATTATTCCAAAGTATGTCATTCTTTCATTCTTT
 AATCCCTAATGTTCTTGTGATTTTTTTTTATGATTAATGACCAAACACTATTGTGTGCAAAAAGAAAACCTTGAGCAAATTAGCGCAA
 CTCTTCTCTTACCGCAAGCAAAAAGAACCCTGCCCCCAACCATGAAGAAGAACCTTTCAATCTGTAAATCAAGTGTTTAGACAAGTG
 AATATTTTTTTGAAAGTGGCATTTGGCTTTTCCCATTTGGTGGGTTAATGAACTAATTAGCATTAAATAGGGAAAAGTGGCTTCTCTC
 CCAAGCCCCAGGAATCCTTTTCCCTCCCTTTCTAGTTCCCTTCCCCAGGAAGGAAATCATTCTCCCTTTCCCTCCATCCCTCCCTCATT
 CCTTTCCCTTCTCCAGACTAAAGTCACTCCTCCAACCCACCAGGGCCAAATTACAACTTTTCTTACATAAAAACAAGAGCTTTTGATTC
 CTATGCTTCTGCATTTTATCTCACTAAAGCCCTAAGGGAAAGGAAATTTCAAAGTGTGACTAATGGCTTACAGTAGGAAATTTGGAAGAT
 ACAGAAGGGACAGAAATCAACATGTCAAGTAAATTTTACAACACTAGCTAGAGATTTGGGGCAAGTCATTTATGCTGTCTAGGCCCTCAGT
 TGAGTAATTTGTAAATAAAGGACCCAAAGATAATCTTTGGGTTCTAACAAAATTTCTCTGTAAAACAGTGGTCCCCAGCCTTCTGGCACC
 AGGGACTAGATTCCTGGAAGACAATTTTTCCAAAGATGGTGGGGCAGGGGGCACGTTTGGGGATGATCATCAGGCATTATTTCTCCTAAG
 GAGCGCTCAACCTAGACCCCTTTGCATGCACAGTTTACAATAGGGTTTGTGCTCCCGTGAGAATGGAATGCCCTCCGCTGATCTGACAGCA
 GGGGGGGCTCAGGCAGTCATGCTGCTCACCTGCGCTCACCTCCTGCTGTACAGCTCOGTTCCCTAAGAGGCTACAGGCTGATATGGGT
 COGTGGCCCAGGGTTGGGGACCCCTGCTATAAAGGAAGTTCAGAAAATCAGATTATAATTTCTGATTTTTATAAATCAGAATTTATAA
 AATTCAGATTATAATTTACTACCAAGTAATAGCTCTTTTGGCCCTTAACTTCCCACAGTGAAGACCACTGGAGTAATTTATATCAACGCA
 AAGAACAAAAGCATGGTCAAGTGAAGTCTGCCCCCTCCCTTGGCTTTCTCTCCTCAATCTAACAGTGAGCAAGTTGCAACAAATCGC
 GCCGTTCAAGAAAAGGGAGGATGGAATTTGTTACAACCGTTTCTGTGCCCCAGGCTGGAGTGCAAGTGGGGGATCTTCCCTCACTGAAA
 CCTCTACCTCCTGAGTTCAAGCGATTCTGCTGCCCTCAGCCTCCTGAGTAGCTGGGATTACAGGCACCGCCACCATACTGGCTGATTT
 TTGTATTTTTAGTAGAGATGGGGTTTACCATATTTGGCCAGGCTGGTCTCGAACTCCTGACCTCGKATCCTCCACCTCAGCCTCCCA
 AAGCGCTGGGATTACAGGTGTGAGCCATCGCCCTGGCCAAACAATTTGTTACAATGTTAACAACATAATATCCTAAACATATTTGGCTT

FIG. 3M

TTAAAGTATCATTAGATACACCACAATACTAATAAAGGTTACCTTTGGGTTTTAAGATTAAAGATGATTTTTAAAAATACTTCTTTCTG
 TATTTTCCAAACTCTTAACCATAAACATAAGATATTCCTTGACTTAGGATAGGATTATGTCACAACCCATCATAAGTTTGAAAAATCAT
 AAGTTGAACCATTTGTAATTTGGGGACCATATGTACATGTATGCATATATGATATTAATAATTATTAGACGTCTTTAAAAATTGACTTTT
 TAACATATTACTTTTATTTAATCACCTTGCTCAAGGAGCCTGTAAATTACATATTAATATTCTCCATTATGAAATAAGTCTTTCCATTG
 TGCAAATTAATGCATTGCAGAGGTTCTAAACATCTATATGCTTTGCAACTCGAAAGGAGTAAGTTTCCCTTTCTAATTTTTTTATTCAA
 TTAAATAAAAAAATGAGTTTAATAGAGTCTATTAAATTAGATCATTATTCGGAGTGGTTAGTAAACCTGTTTAGAGTCGACAACACTCC
 CTTTCTCTCTTTTTTTTTTTTTTTTTTTTTTTGTGCCAGAGTCTCGCTCTGTGCGCCGAGGCTGGAGTGCAATGGCACGATCTCGGCTCACT
 GCAACCTCCACTTCCCAGGTTCAAGTGATTCTCCTGCCTCAGCCTCTCGAGTAGCTGGGATTACAGGCAACCGCCACCATGCCAGCTA
 CTTTGTGTATTTTATGATAGAGATGGGTTTCAACATGTTGGTTAGGCTGGTGGCGAACTCCTGACCTCAAGTGATTGCTGCCTCTG
 CCTCCCAAAGTGCTGGGATTACAGGCTGAGCCACCATGCCAGCCCCCTTCTCCTTTTTAAATATCACCAGCCTGGGTTCTTTGTCT
 TTTTGTTTTGTTTYGTTTTTGTTTTTTGTTTTTTTGAGACGGAGTCTTGCTCCGTNGCCAGGCTGGAGGGCAGTGGCACAATCTTGGC
 TCAGTGCAACCTCCGCCTTCTGGGTTCAATGCCATTCTCCTGCCTCAGCCTCCTGAGTAGCTGGGACTACAGGCGCCCGCCACCATGCC
 GGCTAAATTTTTGTATTTTAGTAGAGACGGGTTTCAAGTGTAGCCAGGATGGTCTCGATCTCCTGACCTTGTGATCCACCTGCCT
 CGGCCTCCCAAAGTCCCTGGGATTACAGGCTGAGCCACCATCCCTGGCCTCCAGCCTGGGTTCTTATTGACACTGAATTTCTCAAGTTAG
 TTGGGCTAGTGAGGAAGTCAGGTTACACGGGCCACAGAACAAGAACAAGGATGTTCTTCTCTCTCTCTTCCACTTCATTCTCTGTCA
 GCCTCTCCCGACCTCAGTAGTTGGTCTTTCTCCCCCTTCTTTTGAAGCAGAGTCCATTATACAAATGGACTTGTTTACTTCTCCACA
 TCCCTCTGTGCAAAATTTCTGCCATGGACACCTTACCCACCTTAGAATGTATATTAGACAATTTTGACATCTAGAATGTCTTGTG
 GGCAGAAAAGCGTTTGGAAAGCGTTGCTCCAGGTAGCTCTGATTACAACTGGACCTTTTCGCGGGGTTACCTAGAGCAGTTGAGAGTG
 CTCTTTCTCCTGGCCAGGTGCAGTTGCTCATGGCTGTAATCCCAGCACTCTGGAAGGCCGAGGCGGGGGGATCACCTGCGGTCAGGAGT
 TTGAGACCAGCCTGGCCAACATGGCGAAACCCCGTTCTACTAAAAATACAAAAATTAGCCAGATATGGTGGTATGAACCTGTAATCCCA
 GCTACTCAGGAGGCTGAGGCAAGAGAATTGCTTGAACCTGGGAGGCAGAGGTTGCAGTGAGCTGAGATCAAGCCTCCAGCCTGGGCCCTC
 AGAGCGAGACTCTGTCTTGAATAATAATAATAATAAACAGATAAAATAAAATTTAAAAAATAAAAAAGGAGTGCTCTCTCTCCTG
 AACTGCTGACTCGAGGACTCTCTCAGCCTGTTTTATCATTGGAAGAGGAAATAATATATCTGCTTCGTACACATCTTTAGAAGTTTAA
 ATAAATGTCTGAAATATCAATGATTCTCATTATTCAAATATTGTTTTTTAAGTCACAGTTGCAAGGTTATATACAGAAGCATAGGTT
 TTTATAACAGAAAAATAGACACTTAATATACTGACCTCTTACAAAAATAGTCCCTGCTCAAGCATCCCATCTATGTATCATTAMCATCTA
 TTTCTTTCTACCCAGCTAAAAATAGTTTATTAATAATCCTTGAATGTCACAAGTNGAATACAGAATAAATCAGATAAATACATTAAAAATGC
 ACCTGATAATCAATATGCACCAGATAATGGACACAGTATACATCAGATAATACAGTACAAATTCATGAAAGTTTAGTGTGCAAAGGT
 AAAATGTAAGAATGTCCTAATGTGCTCCCATGCTGCTTAAAACTGTTATTATAAATTGCTTTTTTATTATAAATATATAAAGAATGATG
 TAATAGCCAGCCATGGTGGCTCATCCCTGTAATCCAGGTCTTTGGGAGGCTGAGGCAGGTGAATCACTTGAGGTTAGGAGTTTGAGA
 CCAGCCTGGCCAACATGGTGAACCCCGTCTCTACTAAAAATATAAAAAATTAGCCAGGTGTGGTGGTACGCACCTGTAGTCTCAGCTAC
 TCCGGAGGCTGAGGCAGGAGATCCCTTGAACCCAGAGCCGAGGTTGCAGTGGTCAAGATCAAGCAACTGCCTCCAGCCTAGGTTG
 ACAGAGCGAGACTTTGTCTCAGGAAAAAATAAATTTCTCAGTCACCTAGATTGAGAAATAGAACATTACCAAACAGATAAAGCCCCA
 CTGTGTTCCCATCCACATCACATTCACTTTATCTCCTCAAAGGAAAGTGCTATTTTGAATTTAGTATTAATTTTCCCTTGCATTTCT
 TCCTACTCATATCATGTGCCTATATACATATAATATATACAAATGCCGATATCATACATAGCAATGTTTACATTTGATTTTTCATT
 GTCATGTAGAATTTTTAACTTAAAAACATGCTTCATACAGCCGGGTGTGGTGGCTCATGCCTGTAATCCCAGCATTTTGGGAGGCCA
 AGGCAGGCGGATCGACGAGGTCAGGAGTTGAGACCAGCCTGACCAACATGGTGAACCCCATCTCTATTAAAAATACAAAAAATA
 TTAGCTGGTCATGGTGGCGGCTGCTGTAATCCCAGCTACTCAGGAGGCTGAGGCAGGAGAAATGTTTGAACCCAGGAGGCAGAGGTTG
 CAGTGAGCCGAGATCGCACCATTTGCCTCCAGCCTGGGTGACAGAGGAGACTCCATCTCAAAAAAATAAAGCTTCATACAAA
 CATGAAACGGGCACATGTCTGGCTGGGTGGCTCATGCCTGTAATCCCAGCACTTTGGGAGGCCAAGGCGGGCAATCACTTAAGG
 CCAGGAGTTGAGACCAGCCTGGTCAGCATGGTGAACCCCGTCTCTACTAAAAATACAAAAATTAGCCAGGATGGTGGCATGCGCCT
 GTAGTCCCAGCTACTGGGAGGCTGAGGCACAAGTATCACTTGATCCCAGGAAGCAGAGGTTGCAGTGAGCCAAGATTGTGTCACTGCA
 CTCCTGCCTGGGTAACAGAGTGATACTCTGTCTCAACAAACAAACAAAAAACAAGAAAAGAAAAAGAAAATGGGCACAT
 GTCAAATGTTAATTTGACTATGTAACCTTATTAATGAAGGAACCAGCAGGTTGTTAGAGCTGGGTCAAAGAAGTATAAGAGAGACTGGAG
 TGCTTACAGTCAAGCAGAGACAGAATGCTGAAAGGTTATGAAATTAGATATGTTAGTTAATTTGAAAGGGCAACTAACTGTAAATC
 TTGCCATTATCTTTCTATCAGACCAATAAATTTACATCTCTACTAGACAAACATTTGCCACTTTTCAATCCATAATCTATGGGTAAT
 TTCATGGAGTCTGGCCCTAATCAACAGTAAATAGTAAAGCCAAACAAAGGATCTCTTCCCTAGACCTTGAAGTGATCTTTGGGTGGACCC
 CTTAGACAATAATTTAGTATGACATTGAGAGGACACCGAAGCCTGGGCAOCATAGTGAGACCCCGCTCTACAAAAAATTTAAAAATTAG

FIG. 3N

CCGGGCATGGTGGTGTGAGCCTGTAGTCCTAGCTACTCAGGAGGCTAAGGTGGAATACCACTTGAGCCCAGGAGTTCGAGGCTGTAGT
 GAGCTATGATCATGCCATTGCACTCCAGCCTGGGTAACAGAGCGAGAACCCTGTCTTGAAAAAAGAAAAAGAAAAAGAAAAAGAAACA
 AAAGGAAATGCAGCCATTTTTTTTTTGCCTTATTTCCAAGTTCTGGATAATTTTTCTTTTTTAACAATATAAATATTACTTATGTA
 TTCTTTTGCAATATGGCTTTTCACTCAGTGTAGTTTGCAAGGGTTAGCCATGTGAATGCATGCTGCTCTAGTTCATTAATTCAGTGT
 GTATGTTGGTCTATGTAGGCATATCACAATWTATYCATTCOCTAGCTGAAGTACATTTGCTTTCAAGGTATGCTATTATAAACAAATC
 TCATACCTTTAATCAAATAATAATTTGTCTCTTCAATCAGCTNTGATTTACTTTGTTTCAANACNAAGCACACAACATAATTANAAT
 TTCATTAAGTATAAATAAATAAATTTTCCAAAACATCACAATCTTTTNTNNTNCACTATTTACTATAACACTTTNGGTCTNAATTTAA
 AGCGGCTTCACTATATGTGGTTCTTTTCTCTCTTCCCATACTAATTAAGTACTGGTACTGGACATATACATCCAAAATCAAATAGTARTGTC
 CTTTTAAGGGATAAATGGGATGTGATGTAGAAGGGGCATAGTAGGGACTTCATCTGTTTGGCAAATTTTTCTTAATATAGGTGGTA
 GGCATGTGGAATTTATAACAAAAGTTCTGTCTCCAGCCAGTTTCTGTTACATAAAAACCATATAATTAACAGTTAAACTGGATCTGGTT
 TGACACAGATGTAGACGATATTAATAATTAAGTACTCCAGAACAACAGGCATAACTAAAACACTACCACAGGCAAAGGGGAAAATAGAGAATG
 TAAGGGCTGGGACTTAAGCCCATGTTGCCACCTCCAAGTTTCATGGACTTTTCTCTTCCACATTACTTTCTTCTCTGCTAGACTGT
 OCTGATGTACCTGCTCTGCACACAGAATTAGACGAGGCGATCAGGTTGGTCAATGTATCCAATCAGCAGTATGGCCAGATTCTCCAGAT
 GACCCCGAAGCACTTGGAGGACACCCGCTATCTGGTGGAGAGATGAGAGGGCAATTTGGCTGGGTGTCTGAACTGGCAAACCAGGCCC
 CAGAAAACAGAGATCATCTTTAATTCATACAGGTAAAGGAGAGACCCAAGAGCAGATACGGAAATGACACGTGCATACCTTGATTTAC
 TGTTAATTTACTTATGAATGTGTCTGAATTTGAAAACAAGCTGTAGGAGGTATTCATATTTCCATTGTGATTGCCTTCAGGCTGACTT
 GATTTAACGTAGTTCATGGTCTTTAGAAAACAAGAAAGTCCATAAAGAAAAATCAATTTAAAACACAAAATACTTTCTAATCTAGAAATG
 GCTATTTCTGCTTAGAGTTATAGGGCTATACTGATAGAGGTAACCTTGAAGAAATATGGCCAATGTAGGTTTTAGGAGAGAAGACTTA
 CAAATAAAGCAATTTGAGTTCAAAATTTGACTCTGAAACTTACCAGCTGAGTAAGCTTGGGAAAAGTACCTCAACCATTCTAGGCCTCAG
 TGTTCCACCTGTAAAATGGTAACAATCATAGCTATCTTAACTGTACACCTATAAAGTGATTAGTATAGATTTCTTATACAAAACAAGA
 GCTCTGTAATTTATAGCTCTTATTAGTTGCTGACACAATAAAGCCACTGAGTTATCTTGAGAATTAACATTTATATGTTACTCGTCA
 ATAAAAATACATTTGCCAGCTGGGCGCAGTGGCTTATGCCTGTAATCCCAGCACTTTGGGAGGCTGAGGTGGGTGGATCACTTGAGGTCA
 GGAGTTTGAGACCAGCCTGGCTAATGTGGCGAAACCCCGTCTCTACCAAAAACATAAAAAATTAGCCAAGTGTGATGGCACACACTTGT
 AATCCCAGCTACTCAGGAGGCTGAGGCAGGAGAATCACTTGAACCCGGAAGGCAGAGGTTGCAGTGAGCTGAGATCGTGCCACTGCACT
 CCAGCCTGGGCGACAGAAGGAGACTCTGTCTCAAAAAAACAATAAARACATATTGCCATCTTAAATCCACCTATACCATGACTC
 CCAGATTCAGTCAATAACTTTTTGCATAACATGCAAGTGAATTTCTTCTTAAAGACATCCCCCTCCAACACACACACATTACCTTAAT
 CTACAAATGCGCCAGGCTAGTGATTCCTGATGAGGCTGGTTTTGAGGGTTCCCAAAAAGACTTGGATACAAAATTAAGTGGGCGAGCA
 ATTGAAGATGCAATATTCTGTGTGTAGTATGTTAGGTTATGTTGGTGGCCCTATCCAGATCCCTGGGGATCCCTTTTACCAGCTCCCCT
 GGTGCTGGTGTGCTGCTAAGTCTTATCTCTGAACTTTCTCCCAAGATTGCCCTTGGAGCACTTATGCCCCAGAGCTTCTGTCAGG
 ATCAGGCTGAGGCTAACAGTCATCTGAAGCCATATCCTTGCTTAGCTTCTTCACTTCTCTAGTTTGTCTTCTCATCCCCTTAAAAGT
 TGCACCTGAGAGCATTCTTTATAAACCCTTCTGTGAGAATCTCAGGCACTGCTTCTAGGAAATTAGACTTATGGCATTCTATAATCCA
 GCATTTCCCTCTTTTTTCAAACCTACAAGCTGTGGATCATGCCCTGATTTGAGAAATAAGTTTTAGAAAGTCACAGCAAGCTCATTAAAA
 ACAAAATTAAAAACCATACAAAATAGAAATAGGACAAAGTAGAAAATATTAGCATGCATTGCATTTCATAGTCATATGCACATCATG
 GAATTTCAATTTCCATTTTGTATGTGTATATGTGTGTAACATATATACACATATGTAGACATACGTGTGTGTTTTGAATCATGATGTCA
 AGTGTATTCACTTACGACACCACAGTCAAAGGGTTTTGAAAGCCACTGTTCCAATCCCTGCCAGCTCTCTGATTCTATAACTCTATTA
 GATTACACTTGAGGAAGGTAATAAATTCATATATTTGATCATCCTGGCATATATAGACTTTTAGTTTAAAGGAAAAAGTCTTGTA
 TTGAAGAATAAAAATGAAGAAAAATTTTAGCAGTGTCTTCAACCTTTAGAAATCTACAGTCAATATTTAGTTGTTTTTACCATTGTCA
 GTATTTTCTATTCTGTGCTTTGATTTACTTCCATTCTAGTGTCTCTTGAGTAACATAACAGATTTATCTAAAATTTCTTTATGCTGATAA
 CAAAGGCACTTCTATATAAAAACCTCCACATAAAAATAAATTTATGGTTTTCAATTATACATTTTTATAACAATTTATACCCTTAAGAG
 CATTACTGGGTGTCAGGCAATGTTCTAAGACTTTTTCCATATATCAGATCATTAAATACCCTCAATGACCCATAAGGGGAAGTAGAAT
 TCTTTCCCCAGTTTTTCAAATGAGGCACAGAGGAGGTTAAGCAACTGTCTGAGCTCACACAGCTAGTAAATGGTAGAACTAGAATTCA
 AACTCAAGCAGTATTTCTCTAGAATCAGTGAACGTAACCCTTTGCTAAAAGTGCCTGTGAAGTTACTTTTCTCAAAACAGCTCCTATTT
 CACCATGTAAAGAAAAGTACAAAACCCATAAAAATAGCAAGTGTGTAAGAGAAGCCTTATGAAAGAAATATACAAATCCAGCAAGTGAAA
 ACGGTTGTGGTCCCTGGTTGTATAATAGTTACATGGGTGTTGACTTTACAATTTTAAACCAACATAAATACTTTATGCAGTTTTTA
 TGTATGTTATACTCACAGAAAGAGAAGGGAAAAATTTTTAAATCATTTCTTAAAGGTTACATCAAGTTGGTATCAGTTCAGTTCATT
 TAAATGATTCAAATCAAAGTCTGTGCATTTGAGAATTCATTAAGAGAGTAACATACATGTTATTCATTAAGAGTAACATAAATTTGCA
 TTGATTTCTGCCAAAATCACACCTACAACCATAAAATTTGTAATTTCTAGGAAAACCTCAGTACAAAACCTGGTGAATGCAATAAAGTTT

FIG. 30

GTGGCACAGACAGTAATACTCAGCAAACATCCCACCTCCTCTCATATTTTCCAGCTCCCTTGTGGTTAAACGTTGCCATGTGGCAA
 GTTCTGGCCAGTGAAGCGTGAGCAAACTGAAAAGGGTCTTTGTAGATTGAGACAGTGAAGAGCCTATGTGTGCTCATCTATTCTCTT
 TTTCTGCTGAGGGCACAAAGAAAGTCCTGAAATCATGTGCTACAGCTATGAGATAATGTGCCTTTGCCTACCAGGCTTCTCAGTGTTTA
 CTGGTGTGGAGCCCTGTAAATGGACACATAACATGAACAAGAAATAAATCTTTGTTCATGAAGCCCTAGGAATGCCAGGACTAATCTG
 TTACCTCAGCACAAACCAGGCCTATCCTGACTAAGGTGGTATTAAATTACTATTGAATGTGTATTGGGATTTAGTAAACTTCTACTGT
 ATAATCCTTCTTCTGTAGGTAGTTCCAAGGATTCATGAAGGAAATATTTCCAACAAGATGAACAATGATGACAGACTTAAGCATTCT
 GCCTCCTCTAATTTACACTCAAGATCCCTCTTGAAGAAAGTGTGAGAGTTCTAACTTCATTGGCTACGTAGTGGCAAAAGCTCTAC
 AGCATTTTAAGGAACATTTTAAACCTGGTAAGCAGAGTGCCTGGTTAGGAATGCCTTGTGACAGGAATAGTTAATTCTCAAAGGGA
 AAAACAAAACCTGTTTCAAATACCTGGAAAACATGTTTAACTCATTAATAAAGACATGAAAACAAAACAAGATGGCATTCTCGCCTA
 TCAGATTTGCAAATTAATAAAAAACCAGGAAATCCTGATAGGAATGTGATGAAATGGGAATTCATATATCATGTATTGGTGGGAAC
 ATAATGGTTTTGCAATTTTGAAGCTATTGATTATGCATATGAAGAGCCATAAAATTTCCTTTTGATATAATAATTCCACTTCOGAA
 ATCAATCCTAAGGRATAAATCTAAATTTGATGAAMAKTCTCCCTCCAAGATCTAGATTTGCAGCATTATTTAAATATTAAGTTGGCC
 GGGCCAGTGGCTCATGCCGTAAATCCCAGCACTTTGGGAGGCTGAGGCGGGCGGATCACGAGGTCAGGAGATGAGACCATCCTGGAT
 AACACGGAGAACTGCGTCTCTACTAAAATAAAAAAATTAGCCGGGCATGGTGGCGGGCGCCTGTAGTCCCAGCTACTCGGGAGGCTG
 AGGCAGGAGAAATGGCGTGAACCCGGGAGGCAGARCTTGCAAGTGAAGAGATCGCGCCACTGCACTCCAGCCTGGGCGACAGAGCAAGA
 CTCTGTTTAAAAAATAAAAAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAA
 TGTGTAACAAGAGAATGATTAAGTAKATTATGACTAAATACACTCAATACATTTTATGAAACGTTAAAAATATTCAAAAATTTAA
 ATAATGACTTGCTAACTACTTTAACAAGAGCTTTATTATCAGCTAGTCTTGGAGGTAATAGTATTATCATGATTTTTCAGAAAAAGATC
 CTGAGGCTCAGTGTCCAAGGTCCAATGAACTACTCAGGTCCGAGGTGGTAGAGCAGCATGTGGAGCCAGTTCTCTCTCCGACTCCATCA
 TCACACTGCACGGCTTCTGTAAAGATATTTGCTCAAAAAATGCGAGATATAAAAAATCTGGGTAATATGATCAACCTTAAAGAATAATT
 ACATTTTAAATTTATTCATGAGACCTTGTAGTAGGTCACCATCAATGTGTAATTAAGCCAGATGTGACAGGATTTGTTGCCTCTCCCTT
 TACTTCTGAATTTTGGAGGCCTTTTTTTTTTCTAGTTGTATCAGTCAGCCAACCAATATCTTTTTAGCATCTACTAAGTTTAGATACG
 GAACTGGTACTCTGAAAGAGAAAATGAGAAAATTTGACAAGATCCTGTCCCAAGGAGCTTCCATCCAACAGGGGCACAAGACAGATA
 GATAGACACACACACACACACACACACACACACACACACTATAAAGCAAGGCAAGATTTAGAGAGTGCACAGGAGTGGG
 CTCTGGGAGTTCAGGGGAGGTCGTTACATTTCTGGTAGGGAAGATACTTCTGAGCTCAGTATATTCCCTTCTCACTGTCCCTCTATC
 CCCTCTCTCCCTCTCCCTCTCTTTTCTCTCTCTCTCCCTCTCCCTCTCTCCCTCTCTCCCTCTCTCCCTCTCTCCCTCTCTCCCTCT
 TTTTTTTTTGAGACAGAGTCTTGTCTGTCAACCCATACTGGAGTACAGTGGCACGATCTCGGCTCAGTCAACCTCGGCCCTCCAGGT
 TCAAATGATTTCTGTGCCTCAGCCTCCTGAGTAGCTGGGATTACAGGCGCACACCACCATGCCTGGCTAATTTTTGTGTTTTTAGTAG
 AGACAGGGTTTCAACATGTTGGCCAGGCTGGTCTTGAACCTCCTGACCTCAAGTAATCCACCCACCTTGGCCTCCCAAAGTGTGGGATC
 ACAGGCATGAGCCACCACACTGGCCTCCTCTCCCTTTCTTAAAAATACATCAATTAATTAATATATAAATGTAGATACACACACAGGC
 AGAATCAAAGTGTATAGGTTGGAGAGGAGACTGTTCCAAAAGGGGGGATGGCATGGGCAAAATACGGCAAGAAAGAGTAGAGCATCTAGG
 TACTGAGGGTGTGGAAAGTCTGCTAAAAATACGGCAAGAAAGAGTAGAGCATCTAGGTAAGGGTGTGGAAAGTCTGCTAAAG
 TGGTCCCTCCCACTGTGGGGCCTTTGAGTTTCCCTGTGCCAGGTTACCTGCCCTCTGTGAGTTTGTGTTCTTTTGGTTGCAAGCA
 ACCAAGACCAGCTCAGCTAAAGAAATGGATGGATACCGACTCATGAGTCAGAGGGGAAGCTGGAGTCTATGCCAGAGCCAGGCAGA
 AACGGTTCAGGTCTAGAGTCTGGGAGGAGGAAACCGATGGACAGCTGCTTCAGGGCCCAGCGCTCAGGGTGAAGCAGCTGCAGTTGTTT
 TTAGTCTCAGATCACTCTGCTCAAGATGTGACTTGCCAGGAGGAATCTGGCTGGCCCAGCTGGGACATGTGTGTCTACCTCTAGACCA
 GGAGAGAGGAGAGTCTTGGTGTGACAGTCCCCATGTAGTACCCCTTTGTTAGGTTACTGAGTCATCAACAGATCTCAGTTCAAATAGTC
 ACTTCTCAGGGGCAATATAACCTCTTCTACCCATAAACTAGGGGCAACATACCCCTCTCTCCCTTTTACACATGACCATAACACCATG
 TAGCACTCAACTCTTGAAGTGTGACATTTACCCATGTGACTCTTTATGAACGTTCACTCCATCCCGAGACCTACAGTCCATGAGGGTA
 CCACCGTCTAGGGTTTTGCTCTTCTCTTTGTGACTGGGGACTTAGGACTCTGCCCTGGCACAGGGCAAACCCCTCAATATTTGTTGAAT
 AAATTAATTAATAAACACGTGTAAATGAATATCAGTAGACTACAACAAGAGTAACAGTAGGCGAAGGTGGAAGGCAAAGGTGGGAAGAG
 GTCAGGGCTCTGAGTGTGGCTGTGGAGTCTGAGGTTCACTCTACAGCGCTGGTGAGACACGATAGGTTTTAGAGAAAGGAAGCCTCA
 TGCTGGTGCCTCAGTGGTACTGACTATGCATTTGTAGCCAAATCAAAGTATTTCCATAAAGTCACTATCTCTTCCAGTTGTTGGG
 ACTTCCAATGGCAATGGGAATTAAGATACTGAGTAATTTGGGAGATCAAGCAAATTTACTAACAAGGCACACGAAGTGAATTTTTCAC
 AGGCAATGTTAATGTTTTTCTTTTTATGTAGTTTTAAATTTCTAAAAGTAACAAAATCACAACCTACCAAACATTTAGAGGACAAAAT
 TATCCATAATCCACCATCTTAACACAACCACTATTATCATTTGTTTTCTTATTTCACATTTTCTACCTATTTTCTTAGATTYCCAAGA
 AATAGAATTAATGTTTTAGAGGTTATTAACATCTTATTGTTCTGGATATATATATATATATATATAGCTATATATAGCTAAATTTAATAACAG

FIG. 3P

TGGTAGTAGCAACAATAAAAAATAATATTATCAAGTAACTGATTCATAATTGACTCTCAAAAACGTTAATTTTCTGCTTTCCTTTACCT
AAGTTTACCTACATGTTGAATTTGTAAAGGGAAGGTTTTCTAGACCAATAATTTTCAAATATTTTGTCTCTCATACTTCCCTCAAAGG
AAACTGAAAAAGTTGCAACATACTTGCATGTCATTTTTCTATATAAGTTGAAAGAATAGCAAATTGTTATTTTCCCACGCATCGTAAAG
ATTAGCAGGTCAATCCCTCTTTAAAATGTACCAAATGGAATCTAAATATCATCGCAATTTGACCCAGCATCATCCATTTAAACAAATATA
CAAGTTTTCTTTAACAATGAGAAATTTATCTCATTACATTTTCTCCCTAAACTCTTATTTCAATCTACATTCCTAAGAATTTTATCC
TAATGTAGTATATTTTATGCTTAAATATCTTTTGTGATCAACACAATTTTGATCATTMTTAAATTTTAAAAATTAAGAACATCCTGT
GACATCAAATTCAGGTATGAAATATTTATCTAGATTGGGTGATCATTATAATTATTTTGTACATAATTGATCAAATAACATAAA
TATACTACAAATTTCTATGACTACTAAACATATAAAAAGTAAAATTTTAAACAAATATATCTCTTAATGAGAAGGAAGAGCTTTTATAC
TCCAATAAGTTAACGTATCCACTAATAATTATTTCTTCCCTAGAACAAGACAGGATTAAGCATCATGACCGTCCCTATTGGGGGATG
TTTTTATAGATGCAAGCACTGTGGCACCTACTGGTATAAATGCACCTGCTGATTGGAATGTTCTTTCCCCAGATCTTCCCTGCTGGTT
TCTTCCAGTATTCAGGTCTCAGCTCAAATGTGACTTCCCTCAATGAGGCCTCCTGGTGATCAGATCTAAAGCACCCCTCTACACAATCAC
TGTTTAGTGCTATACCCATTAATTTACTATCATCACACTTGTCACTATCTGCAGATGTCTTGTMTGGTTACTTTTGTNGTGTTTGTAC
TGCCAGAATATCAGTTCTATGAAGAAAAGGGCCTTGTCTATTTTGTACACTTATAGANATGATGNAGGNACGACATACAAATGGCCAATG
GGCATATGGAAAAACGCTTGACTTCAAGAGTACTNATGGNTATNACCAACATTTATGGAGTAACTACTTTGAAAAGAACCATTCTGTCT
TTACTATCAAGCCAAGATACTCAAGGAAGGCAGCAGAAGTGGAAGCTCCATGTGGGCAGAGGAGCCTAGTCTTGAGATGTGATTTAGCT
GGTATTTGGGTGAAACAAATAAACCAGCCTCAAATAACACAAGGGGCCGGGTGCAGTGGCTCACGCCTGTATCCCAGCACTTTGGGAG
GCTCGAGGCAGGCAGATTACTTCAGGTGAGGAGTTCGAGACCAGCCTGGCTAACATGGTGAACCTCCAT

FIG. 3R

7853-138

HKNG 1 mRNA expression in normal brain

Brain Regions	Gray Matter	White Matter	Neuron	Astrocytes	Oligodendrocytes
Frontal cortex(1)	+++	.	++	.	.
Motor cortex(2)	+++	.	++	.	.
Parietal cortex(3)	+++	.	++	.	.
Occipital cortex(4)	+++	.	++	.	.
Hippocampal formation(5)					
CA1	+++	.	++	.	.
CA2	+++	.	++	.	.
CA3	na	na	na	na	na
CA4	+++	.	++	.	.
Dentate gyrus	++	.	+	.	.
subiculum	+++	.	++	.	.
parahippocampal gyri	+++	.	++	.	.
Caudate/Putamen(6)	+/-	.	+/-	.	.
GPI/GPe/Putamen(7)					
GPI	+	.	+	.	.
GPe	+	.	+	.	.
Putamen	+/-	.	+/-	.	.
Amygdala(8)	++	.	+	.	.
Thalamus(9)medial	++	.	+	.	.
Substantia nigra level(10)					
SNc(substantia nigra pars compacta)	++	.	++	.	.
SNr(substantia nigra pars reticulata)	+	.	+	.	.
Red Nucleus	+	.	+	.	.
3rd cranial nerve nuclei	+	.	+	.	.
superior colliculus	+	.	+	.	.
Upper pons(11)					
Locus ceruleus	+	.	+	.	.
pontine nuclei	+++	.	++	.	.
Lower pons(12)					
locus ceruleus	+	.	+	.	.
pontine nuclei	+++	.	++	.	.
raphy nucleus(midline)	++	.	+	.	.
Medulla(13)					
Inferior olivary nucleus	++	.	+	.	.
12th cranial nerve nuclei	+	.	+	.	.
nucleus ambiguus(multipolar lower motor neurons)	+	.	+	.	.
Cerebellum(14)					
Purkinje cells	++	.	++	.	.
Granular layer	+	.	+/-	.	.
Molecular layer	+	.	+	.	.
Temporal pole(15)	+++	.	++	.	.
Cingulate cortex(16)	+++	.	++	.	.
Anterior thalamus(17)					
Subthalamic nucleus	?	.	?	.	.
Ventral anterior N. (VA),Ventral lateral N.(VL)	++	.	++	.	.
Hippocampal formation(18)					
CA1	na	na	na	na	na
CA2	na	na	na	na	na
CA3	+++	.	++	.	.
CA4	+++	.	++	.	.
subiculum	+++	.	++	.	.
parahippocampal gyri	+++	.	++	.	.
cervical cord (rostral portion)					
anterior motor nuclei	++	.	+	.	.
sensory nuclei group	++	.	+	.	.

FIG. 4

pedigree	Affected Individuals	Phenotype	a. a. change	exon	comment	nt change	nt position
30124	3010189	scz	R331T	8	3 of 4 affected individuals	AGA -> ACT	51,641 51,642
	3010185	scz					
	3010184	scz					
30105	3010027	scz	I23T	3	the only affected individual	ATT -> ACT	35,044
31102	3110017	major depr	E202K	7	all three affected individual (also seen once in Costa Rica)	GAA -> AAA	45,487
	3110014	scz					
	3110003	scz					
30120	3010155	scz	E202K	7	one of the affected individuals	GAA -> AAA	45,487
30126	3010203	scz	intronic	10	3 of 4 affected individuals	insertion: GAATGCCCTGGTTAG 21 base pairs 3' of exon 10	after 63,417
	3010210	scz					
	3010204	scz					
30140	3011486	scz	intronic	6	one of the two affected individuals	A -> T (24bp downstream of exon 6)	43,450
	3210041	scz			two of the three affected individuals		
32301	3210051	scz					

FIG. 5A

pedigree	Affected Individuals	Phenotype	a.a. change	exon	comment	nt change	nt position
30120	3010155	scz	L34L	4	one of the two affected individuals	CTC -> CTA	36,307
32200	3210104	scz	L34L	4	both affected individuals	CTC -> CTA	36,307
	3210009	scz					
31109	3110013	scz	I23T	3	one of the two affected individuals	ATT -> ACT	35,044

FIG. 5B

a.a. change	exon	nt change	position
non-coding 5'-UTR	1	G->C (35 bp upstream from 3' endo of exon 1)	15,385
L42L (silent)	4	CTG -> CTA	36,331
V123G	6	GTT -> GGT	43,184
non-coding (intronic)	6	A -> T (24 bp downstream from exon 6)	43,350
V30I	7	GTC -> ATC	45,571

FIG. 5C

AGTTGCGTCCCTCTCTGTTGCCAGGCTGGAGTTCAGTGGCATGTTTCATAGCTC
ACTGAAGCCTCAAATTCNTGGGTTCAAGTGACCCTCCTACCTCAGCCCCATGA
GGACCTGGGACTACAGTTCCTCCCTTTGGAACGCAGCGTGGGCACCTGCAA
CGCAGAGACCACTGTATCTCCGGTGCAGAATGTAATGAGTGCCTGATACATT
TGCCGAATAAACTATTCCAAGGGTTGAACTTGCTGGAAGCAANAGAAGCACT
ATTCTGGTAACAGCGGGAACATGAAGCCGCCACTCTTGGTGTTTATTGTGTGT
CTGCTGTGGTTGAAAGACAGTCACTGCGCACCCACTTGGAAGGACAAAACCTG
CTATCAGTGAAAACCTGAAGAGTTTTTCTGA

FIG. 6A

AGTTGCGTCCCTCTCTGTTGCCAGGCTGGAGTTCAGTGGCATGTTCTTAGCTC
ACTGAAGCCTCAAATTCCTGGGTTCAAGTGACCCTCCCACCTCAGCCCCATGA
GGACCTGGGACTACAGATGGAGTCTTGCTCTCGTTGCCAGACTGGAGTGCA
CTGCTGCGATCTCAGCTCACTGCAACCTCTACCTCCCAGGTTCAAGCGATTCT
CCTGCCTCAGCCTCTCGAGTGGCTGGGACTATAGTAACAGCGGGAACATGAA
GCCGCCACTCTTGGTGTTTATTGTGTGTCCGCTGTGGTTGAAAGACAGTCACT
GCGCACCCACTTGGAAGGACAAAACCTGCTATCAGTGAAAACCTGAAGAGTTT
TTCT

FIG. 6B

CTTGGAGTCAACTGAGTGTGGACTGAAACTTCCAAAACTGACATGAGGAGTCACTGGAGAATCATGATCAAGGAGCTA 79

CACACTCTGACTTAACTTTATTCTGTGGACAATGAGAGACAACCTGCAAGGATTAACAGTGAGAAC ATG AAG CTG 153

P L L M F P V C L L W L K D C H C A P T 23
CCA CTT TTG ATG TTT CCC GTG TGT CTG CTA TGG TTG AAA GAC TGT CAT TGT GCA CCT ACT 213

W K D K T A I S E N A N S F S E A G E I 43
TGG AAG GAC AAA ACT GCC ATC AGT GAA AAC GCG AAC AGT TTT TCT GAG GCT GGG GAG ATA 273

D V D G E V K I A L I G I K Q M K I M M 63
GAC GTA GAT GGA GAG GTG AAG ATA GCT TTG ATT GGC ATT AAA CAG ATG AAA ATC ATG ATG 333

E R R E E E H S K L M K T L K K C K E E 83
GAA AGG AGA GAG GAA GAA CAC AGC AAA CTA ATG AAA ACC TTG AAG AAG TGC AAA GAA GAA 393

K Q E A L K L M N E V H E H L E E E E S 103
AAG CAG GAG GCC CTG AAA CTT ATG AAT GAA GTT CAT GAA CAC CTG GAG GAG GAA GAA AGC 453

L C Q V S L A D S W D E C R A C L E S N 123
TTA TGC CAG GTT TCT CTG GCA GAT TCC TGG GAT GAA TGC AGG GCT TGC CTG GAA AGT AAC 513

C M R F D T T C Q P A W S S V K N M V E 143
TGC ATG AGG TTT GAT ACC ACC TGC CAA CCT GCA TGG TCC TCT GTG AAA AAT ATG GTG GAA 573

Q F F R K I Y Q F L F P L Q E N D R S G 163
CAG TTT TTC AGG AAG ATC TAT CAG TTT CTG TTT CCT CTC CAG GAA AAT GAC AGA AGT GGC 633

P V S K G V T E E D A Q V S H I E H V F 183
CCT GTC AGC AAA GGG GTC ACT GAG GAA GAT GCG CAG GTG TCA CAC ATA GAG CAT GTG TTC 693

S Q L S A D V T S L F N R S L Y V F K Q 203
AGC CAG CTG AGC GCA GAT GTG ACA TCT CTC TTC AAC AGA AGC CTT TAC GTC TTC AAA CAG 753

L R R E F D Q A F Q S Y F T S G T D V T 223
CTG CGG CGA GAA TTT GAC CAG GCT TTT CAG TCA TAT TTC ACA TCG GGG ACT GAC GTT ACA 813

E P F F F P S L S K E P A Y R A D A E P 243
GAG CCT TTC TTT TTT CCA TCT TTG TCC AAG GAG CCA GCC TAC AGA GCA GAT GCT GAG CCA 873

S W A I P N V F Q L L C N L S F S V Y Q 263
AGC TGG GCC ATT CCC AAT GTC TTC CAG CTG CTC TGC AAC TTG AGT TTC TCA GTT TAT CAA 933

S V S E K L I T T L R A T E D P P K Q D 283
AGT GTC AGT GAA AAA CTC ATC ACA ACC CTG CGT GCC ACA GAG GAC CCT CCA AAA CAA GAC 993

K D S N Q G G P I S K I L P E Q D R G S 303
AAA GAC TCC AAC CAG GGA GGC CCG ATT TCA AAG ATA CTA CCT GAG CAA GAC AGA GGC TCA 1053

D G K L G Q N L S D C V N F R K R C Q K 323
GAT GGG AAA CTT GGC CAG AAT TTG TCT GAT TGC GTT AAT TTT CGC AAG AGA TGC CAG AAA 1113

C Q D Y L S D D C P N V P E L Y R E L N 343
TGC CAG GAT TAT CTA TCT GAT GAC TGC CCT AAT GTG CCT GAA CTA TAC AGA GAA CTC AAT 1173

E A L R L V S R S N Q Q Y D Q V V Q M T 363
GAG GCC CTC CGA CTG GTC AGT AGA TCC AAT CAG CAA TAC GAC CAG GTG GTG CAG ATG ACC 1233

Q Y H L E D T T L L M E K M R E Q F G W 383
CAG TAT CAC CTG GAA GAC ACC ACG CTT CTG ATG GAG AAG ATG AGA GAG CAG TTT GGC TGG 1293

V S E L A Y Q S P G A E D I F N P V K V 403
GTT TCT GAA CTG GCA TAC CAG TCC CCA GGA GCT GAG GAC ATC TTT AAT CCA GTG AAA GTA 1353

FIG. 7A

WO 99/47535

PCT/US99/05606

30/57

M	V	A	L	S	A	H	E	G	N	S	S	D	Q	D	D	T	V	V	P	423
ATG	GTA	GCC	CTA	AGT	GCT	CAT	GAA	GGA	AAT	TCT	TCT	GAT	CAA	GAT	GAC	ACA	GTG	GTT	CCT	1413
S	S	L	L	P	S	S	N	F	T	L	S	S	P	L	E	K	S	A	G	443
TCA	AGC	CTC	CTG	CCT	TCC	TCT	AAC	TTC	ACA	CTC	AGC	AGC	CCT	CTT	GAA	AAG	AGT	GCT	GGC	1473
N	A	N	F	I	D	H	V	V	E	K	V	L	Q	H	F	K	E	H	F	463
AAC	GCT	AAC	TTC	ATT	GAT	CAC	GTG	GTA	GAG	AAG	GTT	CTT	CAG	CAC	TTT	AAG	GAG	CAC	TTT	1533
K	T	W	*																	
AAA	ACT	TGG	TAA																	467
																				1545
GAAGATTTAGTCCATCCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGAGAATTCTGAAAATACAAAGCAGGC																				1624
TAACACAATGAACACAGCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGGTTTACTTTGTTGAATGGAAGTTT																				1703
AATAGCTATTCAAATTGAGTTAATATAAAAATTTCTTCCTAAAAAGTAAAATGTACATATGTAGAATATGATGCATTAG																				1782
TTCTTTGTATACTAAATAAATACTGAGTCCCCT																				1815

FIG. 7B

CTTGGAGTCAACTGAGTGTGGACTGAAACTTCCAAAACTGACATGAGGAGTCACTGGAGAATCATGATCAAGGAGCTA 79

CACACTCTGACTTAACTTTATTCTGTGGACAATGAGAGACAACCTGCAAGGATTAACAGTGAGAAC ATG AAG CTG 3
153

P L L M F P V C L L W L K D C H C A P T 23
CCA CTT TTG ATG TTT CCC GTG TGT CTG CTA TGG TTG AAA GAC TGT CAT TGT GCA CCT ACT 213

W K D K T A I S E N A N S F S E A G E I 43
TGG AAG GAC AAA ACT GCC ATC AGT GAA AAC GCG AAC AGT TTT TCT GAG GCT GGG GAG ATA 273

D V D G E V K I A L I G I K Q M K I M M 63
GAC GTA GAT GGA GAG GTG AAG ATA GCT TTG ATT GGC ATT AAA CAG ATG AAA ATC ATG ATG 333

E R R E E E H S K L M K T L K K C K E E 83
GAA AGG AGA GAG GAA GAA CAC AGC AAA CTA ATG AAA ACC TTG AAG AAG TGC AAA GAA GAA 393

K Q E A L K L M N E V H E H L E E E E S 103
AAG CAG GAG GCC CTG AAA CTT ATG AAT GAA GTT CAT GAA CAC CTG GAG GAG GAA GAA AGC 453

L C Q V S L A D S W D E C R A C L E S N 123
TTA TGC CAG GTT TCT CTG GCA GAT TCC TGG GAT GAA TGC AGG GCT TGC CTG GAA AGT AAC 513

C M R F D T T C Q P A W S S V K N M E N 143
TGC ATG AGG TTT GAT ACC ACC TGC CAA CCT GCA TGG TCC TCT GTG AAA AAT ATG GAA AAT 573

D R S G P V S K G V T E E D A Q V S H I 163
GAC AGA AGT GGC CCT GTC AGC AAA GGG GTC ACT GAG GAA GAT GCG CAG GTG TCA CAC ATA 633

E H V F S Q L S A D V T S L F N R S L Y 183
GAG CAT GTG TTC AGC CAG CTG AGC GCA GAT GTG ACA TCT CTC TTC AAC AGA AGC CTT TAC 693

V F K Q L R R E F D Q A F Q S Y F T S G 203
GTC TTC AAA CAG CTG CGG CGA GAA TTT GAC CAG GCT TTT CAG TCA TAT TTC ACA TCG GGG 753

T D V T E P F F F P S L S K E P A Y R A 223
ACT GAC GTT ACA GAG CCT TTC TTT TTT CCA TCT TTG TCC AAG GAG CCA GCC TAC AGA GCA 813

D A E P S W A I P N V F Q L L C N L S F 243
GAT GCT GAG CCA AGC TGG GCC ATT CCC AAT GTC TTC CAG CTG CTC TGC AAC TTG AGT TTC 873

S V Y Q S V S E K L I T T L R A T E D P 263
TCA GTT TAT CAA AGT GTC AGT GAA AAA CTC ATC ACA ACC CTG CGT GCC ACA GAG GAC CCT 933

P K Q D K D S N Q G G P I S K I L P E Q 283
CCA AAA CAA GAC AAA GAC TCC AAC CAG GGA GGC CCG ATT TCA AAG ATA CTA CCT GAG CAA 993

D R G S D G K L G Q N L S D C V N F R K 303
GAC AGA GGC TCA GAT GGG AAA CTT GGC CAG AAT TTG TCT GAT TGC GTT AAT TTT CGC AAG 1053

R C Q K C Q D Y L S D D C P N V P E L Y 323
AGA TGC CAG AAA TGC CAG GAT TAT CTA TCT GAT GAC TGC CCT AAT GTG CCT GAA CTA TAC 1113

R E L N E A L R L V S R S N Q Q Y D Q V 343
AGA GAA CTC AAT GAG GCC CTC CGA CTG GTC AGT AGA TCC AAT CAG CAA TAC GAC CAG GTG 1173

V Q M T Q Y H L E D T T L L M E K M R E 363
GTG CAG ATG ACC CAG TAT CAC CTG GAA GAC ACC ACG CTT CTG ATG GAG AAG ATG AGA GAG 1233

Q F G W V S E L A Y Q S P G A E D I F N 383
CAG TTT GGC TGG GTT TCT GAA CTG GCA TAC CAG TCC CCA GGA GCT GAG GAC ATC TTT AAT 1293

P V K V M V A L S A H E G N S S D Q D D 403
CCA GTG AAA GTA ATG GTA GCC CTA AGT GCT CAT GAA GGA AAT TCT TCT GAT CAA GAT GAC 1353

FIG. 8A

WO 99/47535

PCT/US99/05606

32/57

T	V	V	P	S	S	L	L	P	S	S	N	F	T	L	S	S	P	L	E	423
ACA	GTG	GTT	CCT	TCA	AGC	CTC	CTG	CCT	TCC	TCT	AAC	TTC	ACA	CTC	AGC	AGC	CCT	CTT	GAA	1413
K	S	A	G	N	A	N	F	I	D	H	V	V	E	K	V	L	Q	H	F	443
AAG	AGT	GCT	GGC	AAC	GCT	AAC	TTC	ATT	GAT	CAC	GTG	GTA	GAG	AAG	GTT	CTT	CAG	CAC	TTT	1473
K	E	H	F	K	T	W	*													451
AAG	GAG	CAC	TTT	AAA	ACT	TGG	TAA													1497
GAAGATTTAGTCCATCCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGAGAATTCTGAAAATACAAAGCAGGC	1576																			
TAACACAATGAACACAGCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGGTTTACTTTGTTGAATGGAAGTTT	1655																			
AATAGCTATTCAAATTGAGTTAATATAAAAATTTCTTCCTAAAAAGTAAAATGTACATATGTAGAATATGATGCATTAG	1734																			
TTCTTTGTATACTAAATAAATACTGAGTCCCCT	1767																			

FIG. 8B

CTTGGAGTCAACTGAGTGTGGACTGAAACTTCCAAAACTGACATGAGGAGTCACTGGAGAATCATGATCAAGGAGCTA 79
 CACACTCTGACTTAACTTTATTCTGTGGACAATGAGAGACAACCTGCAAGGATTAACAGTGAAGAAC ATG AAG CTG 153
 P L L M F P V C L L W L K D C H C A P T 23
 CCA CTT TTG ATG TTT CCC GTG TGT CTG CTA TGG TTG AAA GAC TGT CAT TGT GCA CCT ACT 213
 W K D K T A I S E N A N S F S E A G E I 43
 TGG AAG GAC AAA ACT GCC ATC AGT GAA AAC GCG AAC AGT TTT TCT GAG GCT GGG GAG ATA 273
 D V D G E V K I A L I G I K Q M K I M M 63
 GAC GTA GAT GGA GAG GTG AAG ATA GCT TTG ATT GGC ATT AAA CAG ATG AAA ATC ATG ATG 333
 E R R E E E H S K L M K T L K K C K E E 83
 GAA AGG AGA GAG GAA GAA CAC AGC AAA CTA ATG AAA ACC TTG AAG AAG TGC AAA GAA GAA 393
 K Q E A L K L M N E V H E H L E E E E S 103
 AAG CAG GAG GCC CTG AAA CTT ATG AAT GAA GTT CAT GAA CAC CTG GAG GAG GAA GAA AGC 453
 L C Q V S L A D S W D E C R A C L E S N 123
 TTA TGC CAG GTT TCT CTG GCA GAT TCC TGG GAT GAA TGC AGG GCT TGC CTG GAA AGT AAC 513
 C M R F D T T C Q P A W S S V K N M E P 143
 TGC ATG AGG TTT GAT ACC ACC TGC CAA CCT GCA TGG TCC TCT GTG AAA AAT ATG GAG CCA 573
 A Y R A D A E P S W A I P N V F Q L L C 163
 GCC TAC AGA GCA GAT GCT GAG CCA AGC TGG GCC ATT CCC AAT GTC TTC CAG CTG CTC TGC 633
 N L S F S V Y Q S V S E K L I T T L R A 183
 AAC TTG AGT TTC TCA GTT TAT CAA AGT GTC AGT GAA AAA CTC ATC ACA ACC CTG CGT GCC 693
 T E D P P K Q D K D S N Q G G P I S K I 203
 ACA GAG GAC CCT CCA AAA CAA GAC AAA GAC TCC AAC CAG GGA GGC CCG ATT TCA AAG ATA 753
 L P E Q D R G S D G K L G Q N L S D C V 223
 CTA CCT GAG CAA GAC AGA GGC TCA GAT GGG AAA CTT GGC CAG AAT TTG TCT GAT TGC GTT 813
 N F R K R C Q K C Q D Y L S D D C P N V 243
 AAT TTT CGC AAG AGA TGC CAG AAA TGC CAG GAT TAT CTA TCT GAT GAC TGC CCT AAT GTG 873
 P E L Y R E L N E A L R L V S R S N Q Q 263
 CCT GAA CTA TAC AGA GAA CTC AAT GAG GCC CTC CGA CTG GTC AGT AGA TCC AAT CAG CAA 933
 Y D Q V V Q M T Q Y H L E D T T L L M E 283
 TAC GAC CAG GTG GTG CAG ATG ACC CAG TAT CAC CTG GAA GAC ACC ACG CTT CTG ATG GAG 993
 K M R E Q F G W V S E L A Y Q S P G A E 303
 AAG ATG AGA GAG CAG TTT GGC TGG GTT TCT GAA CTG GCA TAC CAG TCC CCA GGA GCT GAG 1053
 D I F N P V K V M V A L S A H E G N S S 323
 GAC ATC TTT AAT CCA GTG AAA GTA ATG GTA GCC CTA AGT GCT CAT GAA GGA AAT TCT TCT 1113
 D Q D D T V V P S S L L P S S N F T L S 343
 GAT CAA GAT GAC ACA GTG GTT CCT TCA AGC CTC CTG CCT TCC TCT AAC TTC ACA CTC AGC 1173
 S P L E K S A G N A N F I D H V V E K V 363
 AGC CCT CTT GAA AAG AGT GCT GGC AAC GCT AAC TTC ATT GAT CAC GTG GTA GAG AAG GTT 1233
 L Q H F K E H F K T W * 375
 CTT CAG CAC TTT AAG GAG CAC TTT AAA ACT TGG TAA 1269
 GAAGATTTAGTCCATCCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGAGAATTCTGAAAATACAAAGCAGGC 1348

FIG. 9A

WO 99/47535

PCT/US99/05606

34/57

TAACACAATGAACACAGCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGGTTTACTTTGTTGAATGGAAGTTT 1427
AATAGCTATTCAAATTGAGTTAATATAAAAATTTCTTCCTAAAAAGTAAAATGTACATATGTAGAATATGATGCATTAG 1506
TTCTTTGTATACTAAATAAATACTGAGTCCCCT 1539

FIG. 9B

CTTGGAGTCAACTGAGTGTGGACTGAAACTTCCAAAACTGACATGAGGAGTCACTGGAGAATCATGATCAAGGAGCTA	79				
CACACTCTGACTTAACTTTATTCTGTGGACAATGAGAGACAACTGCAAGGATTAACAGTGAGAAC	153				
<table border="0" style="width: 100%;"> <tr> <td style="text-align: right;">M K L</td> <td style="text-align: right;">3</td> </tr> <tr> <td>ATG AAG CTG</td> <td>153</td> </tr> </table>	M K L	3	ATG AAG CTG	153	
M K L	3				
ATG AAG CTG	153				
P L L M F P V C L L W L K D C H C A P T	23				
CCA CTT TTG ATG TTT CCC GTG TGT CTG CTA TGG TTG AAA GAC TGT CAT TGT GCA CCT ACT	213				
W K D K T A I S E N A N S F S E A G E I	43				
TGG AAG GAC AAA ACT GCC ATC AGT GAA AAC GCG AAC AGT TTT TCT GAG GCT GGG GAG ATA	273				
D V D G E V K I A L I G I K Q M K I M M	63				
GAC GTA GAT GGA GAG GTG AAG ATA GCT TTG ATT GGC ATT AAA CAG ATG AAA ATC ATG ATG	333				
E R R E E E H S K L M K T L K K C K E E	83				
GAA AGG AGA GAG GAA GAA CAC AGC AAA CTA ATG AAA ACC TTG AAG AAG TGC AAA GAA GAA	393				
K Q E A L K L M N E V H E H L E E E E S	103				
AAG CAG GAG GCC CTG AAA CTT ATG AAT GAA GTT CAT GAA CAC CTG GAG GAG GAA GAA AGC	453				
L C Q V S L A D S W D E C R A C L E S N	123				
TTA TGC CAG GTT TCT CTG GCA GAT TCC TGG GAT GAA TGC AGG GCT TGC CTG GAA AGT AAC	513				
C M R F D T T C Q P A W S S V K N M P A	143				
TGC ATG AGG TTT GAT ACC ACC TGC CAA CCT GCA TGG TCC TCT GTG AAA AAT ATG CCA GCC	573				
Y R A D A E P S W A I P N V F Q L L C N	163				
TAC AGA GCA GAT GCT GAG CCA AGC TGG GCC ATT CCC AAT GTC TTC CAG CTG CTC TGC AAC	633				
L S F S V Y Q S V S E K L I T T L R A T	183				
TTG AGT TTC TCA GTT TAT CAA AGT GTC AGT GAA AAA CTC ATC ACA ACC CTG CGT GCC ACA	693				
E D P P K Q D K D S N Q G G P I S K I L	203				
GAG GAC CCT CCA AAA CAA GAC AAA GAC TCC AAC CAG GGA GGC CCG ATT TCA AAG ATA CTA	753				
P E Q D R G S D G K L G Q N L S D C V N	223				
CCT GAG CAA GAC AGA GGC TCA GAT GGG AAA CTT GGC CAG AAT TTG TCT GAT TGC GTT AAT	813				
F R K R C Q K C Q D Y L S D D C P N V P	243				
TTT CGC AAG AGA TGC CAG AAA TGC CAG GAT TAT CTA TCT GAT GAC TGC CCT AAT GTG CCT	873				
E L Y R E L N E A L R L V S R S N Q Q Y	263				
GAA CTA TAC AGA GAA CTC AAT GAG GCC CTC CGA CTG GTC AGT AGA TCC AAT CAG CAA TAC	933				
D Q V V Q M T Q Y H L E D T T L L M E K	283				
GAC CAG GTG GTG CAG ATG ACC CAG TAT CAC CTG GAA GAC ACC ACG CTT CTG ATG GAG AAG	993				
M R E Q F G W V S E L A Y Q S P G A E D	303				
ATG AGA GAG CAG TTT GGC TGG GTT TCT GAA CTG GCA TAC CAG TCC CCA GGA GCT GAG GAC	1053				
I F N P V K V M V A L S A H E G N S S D	323				
ATC TTT AAT CCA GTG AAA GTA ATG GTA GCC CTA AGT GCT CAT GAA GGA AAT TCT TCT GAT	1113				
Q D D T V V P S S L L P S S N F T L S S	343				
CAA GAT GAC ACA GTG GTT CCT TCA AGC CTC CTG CCT TCC TCT AAC TTC ACA CTC AGC AGC	1173				
P L E K S A G N A N F I D H V V E K V L	363				
CCT CTT GAA AAG AGT GCT GGC AAC GCT AAC TTC ATT GAT CAC GTG GTA GAG AAG GTT CTT	1233				
Q H F K E H F K T W *	374				
CAG CAC TTT AAG GAG CAC TTT AAA ACT TGG TAA	1266				
GAAGATTTAGTCCATCCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGAGAATTCTGAAAATACAAAGCAGGC	1345				

FIG. 10A

WO 99/47535

36/57

PCT/US99/05606

TAACACAATGAACACAGCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGGTTTACTTTGTTGAATGGAAGTTT 1424
AATAGCTATTCAAATTGAGTTAATATAAAAAATTTCTTCCTAAAAAGTAAATGTACATATGTAGAATATGATGCATTAG 1503
TTCCTTGTATACTAAATAAATACTGAGTCCCCT 1536

FIG. 10B

WO 99/47535

38/57

PCT/US99/05606

CAC GAA GGA AAT TTC TCC AAA CAA GAT GAA AAG ATG ATA GAC ATA AGC ATT CTG CCT TCC 1329
S N F T L T I P L E E S A E S S D F I S 448
TCT AAT TTC ACA CTC ACC ATC CCT CTT GAA GAA AGT GCT GAG AGT TCC GAC TTC ATT AGC 1389
Y M L A K A V Q H F K E H F K S W * 466
TAC ATG CTG GCC AAA GCT GTA CAG CAT TTT AAG GAA CAT TTT AAA TCT TGG TAA 1443
GCAGAGTATTTGATTAGGGACGTTTGCTGATAGGAATAGATGGTTCTTAAAAGGGAAAAATGACAAAACCTAGCTTTTGA 1522
ATACCTTGAAAACGTATTCAACCTCATTAAATAATCAAAGGCATGAAAACCTAAGACAAGTTAGCAGTTTTTACCTATTGA 1601
ATTTTCAAATTAATAAAAAAAAAAATCCTGATAGAATGCAATGAAATGAGAATTCTTATATGTGATTGCCAGAAACAACTG 1680
GTTTTGTCTTTTGAAGTTATTCAATTATACATATCAAGAGTCATCAAATTTCTTTTAAATATAATAATTCCACTTC 1759
TGGAATCAATCCAAAGGAGTAAATCTAAAATTGAATTGAAGTTCCACCCCAAGATCAATATTTGCAAATTATTTAAAA 1838
TAGTAAACTGTTAAAAACTGAATGTCATCTGAATGTCTAAAAACCAGAAATGGTTAAAAGCTGTGGCTAAATATGCTCC 1917
AAATATCTTATAAAACCATTAATAAATATTATAAAATTTAAATCATGACATGACATCTGCTGGAACAAGAGTTTATTCT 1996
AAGCCTATCTATAAGGCAAATATTATTACTATCTTCCAGAAAAGAACTTGAGACTCAGGGTCCAAGTGTTAGTTG 2075
CTCAGTCATGTCTGACTCTTTGGGACCCCTTGGACTGTAGCCCACCAGGCTCCTCTGTCCGTGGGATTCTTCAGACAGG 2154
AATACTGGGGCAGGTTGCTATTTCTTCTCCAGGAAATCTTCCCTATCCAGGGATGGAACCCAGGTCTCCTGCATTGCA 2233
GGTAGATGCTTTACTATCTGAGCAACCAAATGAATTACTCAAGTCAGTAGGGGGTAGAGGCAAATTTTAACTTAGTTTT 2312
CTCTGAATCATAATTGCCACATTAACCTGGTTCTGTTGGGACATTTGGTTGAAAAAATAAAGTGAAAAATGAGTATA 2391
AAACTCTATAAATGTAATGATCAAAACGAAAAAATCTACAATCTGCATTAAAAATAAAAAGGGTTGGCAGG 2464

FIG. 11B

WO 99/47535

40/57

PCT/US99/05606

P	G	V	H	E	G	N	F	S	K	Q	D	E	K	M	I	D	I	S	I	425	
CCA	GGT	GTT	CAC	GAA	GGA	AAT	TTC	TCC	AAA	CAA	GAT	GAA	AAG	ATG	ATA	GAC	ATA	AGC	ATT	1332	
L	P	S	S	N	F	T	L	T	I	P	L	E	E	S	A	E	S	S	D	445	
CTG	CCT	TCC	TCT	AAT	TTC	ACA	CTC	ACC	ATC	CCT	CTT	GAA	GAA	AGT	GCT	GAG	AGT	TCC	GAC	1392	
F	I	S	Y	M	L	A	K	A	V	Q	H	F	K	E	H	F	K	S	W	465	
TTC	ATT	AGC	TAC	ATG	CTG	GCC	AAA	GCT	GTA	CAG	CAT	TTT	AAG	GAA	CAT	TTT	AAA	TCT	TGG	1452	
*																					
TAA																					466
																					1455
GCAGAGTATTTGATTAGGGACGTTTGCTGATAGGAATAGATGGTTCTTAAAAGGGAAAAATGACAAAAGTACTAGCTTTTGA																					1534
ATACCTTGAAAACGTATTCAACCTCATTAATAATCAAAGGCATGAAAACCTAAGACAAGTTAGCAGTTTTTACCTATTGA																					1613
ATTTTCAAATTAATAAAAAAAAAAATCCTGATAGAATGCAATGAAATGAGAATTCTTATATGTGATTGCCAGAAACAACTG																					1692
GTTTTGTCTTTTTGAAAAGTTATTCAATTATAACATATCAAGAGTCATCAAATTTCTTTTTAATAATAAATCCACTTC																					1771
TGGAATCAATCCAAAGGAGTAAATCTAAAATTGAATTGAAGTTCCACCCCAAGATCAATATTGCAAATTATTTAAAA																					1850
TAGTAACTGTTAAAACTGAATGTCATCTGAATGTCTAAAAACCAGAAATGGTTAAAAGCTGTGGCTAAATATGCTCC																					1929
AAATATCTTATAAAACCATTAAAAATATTTATAAAATTTAAATCATGACATGACATCTGCTGGAACAAGAGTTTATTCT																					2008
AAGCCTATCTATAAGGCAAATATTATTACTATCTTCCAGAAAAGAACTTGAGACTCAGGGTCCAAGTGTTAGTTG																					2087
CTCAGTCATGTCTGACTCTTTGAGACCCCTTGACTGTGGCCACCAGGCTCCTCTGTCCATGGGATTCTTCAGACAAG																					2166
AATACTGGAGCAGGTTGCTATTTCTTCTCCAGGAAATCTTCCCTATCCAGGGATGGAACCCAGGTCTCCTGCATTGCA																					2245
GGTAGATGCTTTACTATCTGAGCAACCAAATGAATTACTCAAGTCAGTAGGGGGTAGAGGCAAATTTTAACTTAGTTTT																					2324
CTCTGAATCATAATTGCCACATTAACTGGTTCCTGTTGGGACATTTGGTTGAAAAAATAAAGTGAAAAATGAGTATA																					2403
AAACTCTATAAATGTAATGATCAAACGAAAAAATACTACAATCTGCATTAAAAATAAAAAGGGTTGGCAGGAATTAC																					2482
GGTTGGAATGGATGATTTTTTTTAACTTTTCATCTTTTGATATTTTACAATTTTCTATAATGAATAAATAATTTTGA																					2561
GATTTCAAATTAGAAGATATGTTGCTAAAATAGCTAGGTAAATGTAGATTGAACACTGTATCAATGTGTTCTCATCTTT																					2640
AACTTTAGTATAAGTACTTCTATTCCATGGTAATCCTACAGTAAGACGAAATGTAAATCTGTTCCGGTCTACAGGAAAA																					2719
ACAACTAAATGACATTTTACAGACGTACATTACCATCTCTGTTAGGATAATCTTCTGAATTAATGGCACAAATTAGAACTGT																					2798
ACATAGTATTTCTCCTTTGGTAAAATGGTCAATCTTAAAGAAGCATTAAATGTTAATTTCTAAGTTATTACTCATAAGGGA																					2877
CCTGTAGGTAGGTCCCTATCAATGTATAATTAAGCTGGGTATTTCTAGATTGCTGCTCCTCTCCCTTTATCTCTGAATG																					2956
TTGGAGAGGTTGTTGGTCAATCAACCAATATCTTTTTAGCATCTTCTAAGTGAAGGC																					3016

FIG. 12B

GTGAAGGTCCTTACAGAAGCTGGTGGCAACCTCGTTGGTGAGAGCCTGCAGTTAGTGTCACGGCGGAAAC	M K	2
ATG AAG		76
P P I L V F I V Y L L Q L R D C Q C A P		22
CCG CCA ATC TTG GTG TTT ATC GTG TAT CTG CTG CAG CTG AGA GAC TGT CAG TGT GCG CCT		136
T G K D R T S I R E D P K G F S K A G E		42
ACA GGG AAG GAC CGA ACT TCC ATC CGT GAA GAC CCG AAG GGT TTT TCC AAG GCT GGG GAG		196
I D V D E E V K K A L I G M K Q M K I L		62
ATA GAC GTA GAT GAA GAG GTG AAG AAG GCT TTG ATT GGC ATG AAG CAG ATG AAA ATC CTG		256
M E R R E E E H S K L M R T L K K C R E		82
ATG GAA AGA AGA GAG GAG GAA CAT AGC AAA CTA ATG AGA ACC CTG AAG AAA TGC AGA GAA		316
E K Q E A L K L M N E V Q E H L E E E E		102
GAA AAG CAG GAG GCC CTG AAG CTT ATG AAT GAA GTT CAA GAA CAT CTA GAA GAG GAA GAA		376
R L C Q V S L M G S W D E C K S C L E S		122
AGG CTA TGC CAG GTG TCT CTG ATG GGT TCC TGG GAC GAA TGC AAA TCT TGC CTG GAA AGT		436
D C M R F Y T T C Q S S W S S M K S T I		142
GAC TGC ATG AGA TTT TAT ACA ACC TGC CAA AGC AGT TGG TCC TCT ATG AAA TCC ACG ATT		496
E R V F R K I Y Q F L F P F H E D D E K		162
GAA CGG GTT TTC CGG AAG ATA TAT CAG TTT CTC TTT CCT TTC CAT GAA GAC GAT GAA AAA		556
E L P V G E K F T E E D V Q L M Q I E N		182
GAG CTT CCT GTT GGT GAG AAG TTC ACT GAG GAA GAT GTA CAG CTG ATG CAG ATA GAG AAT		616
V F S Q L T V D V G F L Y N M S F H V F		202
GTG TTC AGC CAG CTG ACC GTG GAC GTG GGA TTT CTC TAT AAC ATG AGC TTT CAC GTC TTC		676
K Q M Q Q E F D L A F Q S Y F M S D T D		222
AAA CAG ATG CAG CAA GAA TTT GAC CTG GCT TTT CAA TCA TAC TTT ATG TCA GAC ACA GAC		736
S M E P Y F F P A F S K E P A K K A H P		242
TCC ATG GAG CCT TAC TTT TTT CCA GCT TTT TCC AAA GAG CCA GCA AAA AAA GCA CAT CCT		796
M Q S W D I P S F F Q L F C N F S L S V		262
ATG CAG AGT TGG GAC ATT CCC AGC TTC TTC CAG CTG TTT TGT AAT TTC AGC CTC TCT GTT		856
Y Q S V S A T V T E M L K A I E D L S K		282
TAT CAA AGT GTC AGC GCA ACA GTT ACA GAG ATG CTG AAG GCC ATT GAG GAC TTA TCC AAA		916
Q D K D S A H G G P S S T T W P V R G R		302
CAA GAC AAA GAT TCT GCC CAC GGT GGA CCG AGT TCC ACG ACG TGG CCT GTG CGG GGC AGA		976
G L C G E P G Q N S S E C L Q F H A R C		322
GGG CTG TGT GGA GAA CCT GGC CAG AAC TCG TCC GAA TGT CTC CAA TTT CAT GCA AGA TGC		1036
Q K C Q D Y L W A D C P A V P E L Y T K		342
CAG AAA TGT CAG GAT TAC CTA TGG GCA GAC TGC CCT GCT GTT CCT GAA CTA TAC ACA AAG		1096
A D E A L E L V N I S N Q Q Y A Q V L Q		362
GCG GAT GAG GCC CTT GAG TTG GTC AAC ATA TCC AAT CAG CAG TAT GCC CAG GTA CTC CAG		1156
M T Q H H L E D T T Y L M E K M R E Q F		382
ATG ACC CAG CAT CAC TTG GAG GAC ACC ACG TAT CTG ATG GAG AAG ATG AGA GAG CAG TTT		1216
G W V T E L A S Q T P G S E N I F S F I		402
GGT TGG GTA ACA GAG CTG GCC AGC CAG ACC CCA GGA AGC GAG AAC ATC TTC AGT TTC ATA		1276
K V V P G V H E G N F S K Q D E K M I D		422

FIG. 13A

AAG GTA GTT CCA GGT GTT CAC GAA GGA AAT TTC TCC AAA CAA GAT GAA AAG ATG ATA GAC	1336
I S I L P S S N F T L T I P L E E S A E	442
ATA AGC ATT CTG CCT TCC TCT AAT TTC ACA CTC ACC ATC CCT CTT GAA GAA AGT GCT GAG	1396
S S D F I S Y M L A K A V Q H F K E H F	462
AGT TCC GAC TTC ATT AGC TAC ATG CTG GCC AAA GCT GTA CAG CAT TTT AAG GAA CAT TTT	1456
K S W *	
AAA TCT TGG TAA	466
	1468
GCAGAGTATTTGATTAGGGACGTTTGCTGATAGGAATAGATGGTTCTTAAAAGGGAAAAATGACAAAAGCTTTTGA	1547
ATACCTTGAAAACGTATTCAACCTCATTAAATAATCAAAGGCATGAAAAGTAAAGACAAGTTAGCAGTTTTTACCTATTGA	1626
ATTTTCAAATTAATAAAAAAATCCTGATAGAATGCAATGAAATGAGAATTCCTTATATGTGATTGCCAGAAACAACTGG	1705
TTTTGTCTTTTTGAAAAGTTATTCAATTATACATATCAAGAGTCATCAAATTTCTTTTTAATATAATAATTCCACTTCT	1784
GGAATCAATCCAAAGGAGTAAATCTAAAATTGAATTGAAGTCCCACCCCAAGATCAATATTTGCAAATTATTTAAAAT	1863
AGTAAACTGTTAAAAGTGAATGTCATCTGAATGTCTAAAACCAGAAATGGTTAAAAGCTGTGGCTAAATATGCTCCA	1942
AATATCTTATAAAAACCATTAAAAATATTTATAAAATTTAAATCATGACATGACATCTGCTGGAACAAGAGTTTATTCTA	2021
AGCCTATCTATAAGGCAAATATTATTACTATCTTCCAGAAAAGAACTTGAGACTCAGGGTCCAAGTGTTAGTTGC	2100
TCAGTCATGCTGACTCTTTGAGACCCCTTGGACTGTAGCCCACCAGGCTCCTCTGTCCATGGGATTCTTCAGACAAGA	2179
ATACTGGAGCAGGTTGCTATTTCTTCTCCAGGAAATCTTCCCTATCCAGGGATGGAACCCAGGTCTCCTGCATTGCAG	2258
GTAGATGCTTTACTATCTGAGCAACCAAATGAATTACTCAAGTCAGTAGGGGGTAGAGGCAAATTTTAACTTAGTTTTC	2337
TCTGAATCATAATTGCCACATTAAACTGGTTCTGTGGGACATTTGGTTGAAAAAATAAAGTGAAAAATGAGTATAA	2416
AACTCTATAAATGTAATGATCAAACGAAAAAATCTACAATCTGCATTAAAAATAAAAAGGGTTGGCAGG	2488

FIG. 13B

WO 99/47535
7853-138

43/57

PCT/US99/05606

	C T T G G A G T C A A C T G A G T G T G G A C T G A A A C T T C C A A A A A C T				Majority
	10	20	30	40	
1	C T T G G A G T C A A C T G A G T G T G G A C T G A A A C T T C C A A A A A C T				gphkng1815-1.
1	C T T G G A G T C A A C T G A G T G T G G A C T G A A A C T T C C A A A A A C T				gp7b-1.
1	C T T G G A G T C A A C T G A G T G T G G A C T G A A A C T T C C A A A A A C T				gp7c-1.
1	C T T G G A G T C A A C T G A G T G T G G A C T G A A A C T T C C A A A A A C T				gp7d-1.
	G A C A T G A G G A G T C A C T G G A G A A T C A T G A T C A A G G A G C T A C				Majority
	50	60	70	80	
41	G A C A T G A G G A G T C A C T G G A G A A T C A T G A T C A A G G A G C T A C				gphkng1815-1.
41	G A C A T G A G G A G T C A C T G G A G A A T C A T G A T C A A G G A G C T A C				gp7b-1.
41	G A C A T G A G G A G T C A C T G G A G A A T C A T G A T C A A G G A G C T A C				gp7c-1.
41	G A C A T G A G G A G T C A C T G G A G A A T C A T G A T C A A G G A G C T A C				gp7d-1.
	A C A C T C T G A C T T A A C T T T A T T C T G T G G A C A A T G A G A G A C A				Majority
	90	100	110	120	
81	A C A C T C T G A C T T A A C T T T A T T C T G T G G A C A A T G A G A G A C A				gphkng1815-1.
81	A C A C T C T G A C T T A A C T T T A T T C T G T G G A C A A T G A G A G A C A				gp7b-1.
81	A C A C T C T G A C T T A A C T T T A T T C T G T G G A C A A T G A G A G A C A				gp7c-1.
81	A C A C T C T G A C T T A A C T T T A T T C T G T G G A C A A T G A G A G A C A				gp7d-1.
	A C T G C A A G G A T T A A C A G T G A G A A C A T G A A G C T G C C A C T T T				Majority
	130	140	150	160	
121	A C T G C A A G G A T T A A C A G T G A G A A C A T G A A G C T G C C A C T T T				gphkng1815-1.
121	A C T G C A A G G A T T A A C A G T G A G A A C A T G A A G C T G C C A C T T T				gp7b-1.
121	A C T G C A A G G A T T A A C A G T G A G A A C A T G A A G C T G C C A C T T T				gp7c-1.
121	A C T G C A A G G A T T A A C A G T G A G A A C A T G A A G C T G C C A C T T T				gp7d-1.
	T G A T G T T T C C C G T G T G T C T G C T A T G G T T G A A A G A C T G T C A				Majority
	170	180	190	200	
161	T G A T G T T T C C C G T G T G T C T G C T A T G G T T G A A A G A C T G T C A				gphkng1815-1.
161	T G A T G T T T C C C G T G T G T C T G C T A T G G T T G A A A G A C T G T C A				gp7b-1.
161	T G A T G T T T C C C G T G T G T C T G C T A T G G T T G A A A G A C T G T C A				gp7c-1.
161	T G A T G T T T C C C G T G T G T C T G C T A T G G T T G A A A G A C T G T C A				gp7d-1.
	T T G T G C A C C T A C T T G G A A G G A C A A A A C T G C C A T C A G T G A A				Majority
	210	220	230	240	
201	T T G T G C A C C T A C T T G G A A G G A C A A A A C T G C C A T C A G T G A A				gphkng1815-1.
201	T T G T G C A C C T A C T T G G A A G G A C A A A A C T G C C A T C A G T G A A				gp7b-1.
201	T T G T G C A C C T A C T T G G A A G G A C A A A A C T G C C A T C A G T G A A				gp7c-1.
201	T T G T G C A C C T A C T T G G A A G G A C A A A A C T G C C A T C A G T G A A				gp7d-1.
	A A C G C G A A C A G T T T T T C T G A G G C T G G G G A G A T A G A C G T A G				Majority
	250	260	270	280	
241	A A C G C G A A C A G T T T T T C T G A G G C T G G G G A G A T A G A C G T A G				gphkng1815-1.
241	A A C G C G A A C A G T T T T T C T G A G G C T G G G G A G A T A G A C G T A G				gp7b-1.
241	A A C G C G A A C A G T T T T T C T G A G G C T G G G G A G A T A G A C G T A G				gp7c-1.
241	A A C G C G A A C A G T T T T T C T G A G G C T G G G G A G A T A G A C G T A G				gp7d-1.

FIG. 14A

A T G G A G A G G T G A A G A T A G C T T T G A T T G G C A T T A A A C A G A T Majority
 290 300 310 320
 281 **A T G G A G A G G T G A A G A T A G C T T T G A T T G G C A T T A A A C A G A T** gphkng1815-
 281 **A T G G A G A G G T G A A G A T A G C T T T G A T T G G C A T T A A A C A G A T** gp7b-1.
 281 **A T G G A G A G G T G A A G A T A G C T T T G A T T G G C A T T A A A C A G A T** gp7c-1.
 281 **A T G G A G A G G T G A A G A T A G C T T T G A T T G G C A T T A A A C A G A T** gp7d-1.

G A A A A T C A T G A T G G A A A G G A G A G A G G A A G A A C A C A G C A A A Majority
 330 340 350 360
 321 **G A A A A T C A T G A T G G A A A G G A G A G A G G A A G A A C A C A G C A A A** gphkng1815-
 321 **G A A A A T C A T G A T G G A A A G G A G A G A G G A A G A A C A C A G C A A A** gp7b-1.
 321 **G A A A A T C A T G A T G G A A A G G A G A G A G G A A G A A C A C A G C A A A** gp7c-1.
 321 **G A A A A T C A T G A T G G A A A G G A G A G A G G A A G A A C A C A G C A A A** gp7d-1.

C T A A T G A A A A C C T T G A A G A A G T G C A A A G A A G A A A A G C A G G Majority
 370 380 390 400
 361 **C T A A T G A A A A C C T T G A A G A A G T G C A A A G A A G A A A A G C A G G** gphkng1815-
 361 **C T A A T G A A A A C C T T G A A G A A G T G C A A A G A A G A A A A G C A G G** gp7b-1.
 361 **C T A A T G A A A A C C T T G A A G A A G T G C A A A G A A G A A A A G C A G G** gp7c-1.
 361 **C T A A T G A A A A C C T T G A A G A A G T G C A A A G A A G A A A A G C A G G** gp7d-1.

A G G C C C T G A A A C T T A T G A A T G A A G T T C A T G A A C A C C T G G A Majority
 410 420 430 440
 401 **A G G C C C T G A A A C T T A T G A A T G A A G T T C A T G A A C A C C T G G A** gphkng1815-
 401 **A G G C C C T G A A A C T T A T G A A T G A A G T T C A T G A A C A C C T G G A** gp7b-1.
 401 **A G G C C C T G A A A C T T A T G A A T G A A G T T C A T G A A C A C C T G G A** gp7c-1.
 401 **A G G C C C T G A A A C T T A T G A A T G A A G T T C A T G A A C A C C T G G A** gp7d-1.

G G A G G A A G A A A G C T T A T G C C A G G T T T C T C T G G C A G A T T C C Majority
 450 460 470 480
 441 **G G A G G A A G A A A G C T T A T G C C A G G T T T C T C T G G C A G A T T C C** gphkng1815-
 441 **G G A G G A A G A A A G C T T A T G C C A G G T T T C T C T G G C A G A T T C C** gp7b-1.
 441 **G G A G G A A G A A A G C T T A T G C C A G G T T T C T C T G G C A G A T T C C** gp7c-1.
 441 **G G A G G A A G A A A G C T T A T G C C A G G T T T C T C T G G C A G A T T C C** gp7d-1.

T G G G A T G A A T G C A G G G C T T G C C T G G A A A G T A A C T G C A T G A Majority
 490 500 510 520
 481 **T G G G A T G A A T G C A G G G C T T G C C T G G A A A G T A A C T G C A T G A** gphkng1815-
 481 **T G G G A T G A A T G C A G G G C T T G C C T G G A A A G T A A C T G C A T G A** gp7b-1.
 481 **T G G G A T G A A T G C A G G G C T T G C C T G G A A A G T A A C T G C A T G A** gp7c-1.
 481 **T G G G A T G A A T G C A G G G C T T G C C T G G A A A G T A A C T G C A T G A** gp7d-1.

G G T T T G A T A C C A C C T G C C A A C C T G C A T G G T C C T C T G T G A A Majority
 530 540 550 560
 521 **G G T T T G A T A C C A C C T G C C A A C C T G C A T G G T C C T C T G T G A A** gphkng1815-
 521 **G G T T T G A T A C C A C C T G C C A A C C T G C A T G G T C C T C T G T G A A** gp7b-1.
 521 **G G T T T G A T A C C A C C T G C C A A C C T G C A T G G T C C T C T G T G A A** gp7c-1.
 521 **G G T T T G A T A C C A C C T G C C A A C C T G C A T G G T C C T C T G T G A A** gp7d-1.

FIG. 14B




FIG. 14C




FIG. 14D

A T T A T C T A T C T G A T G A C T G C C C T A A T G T G C C T G A A C T A T A Majority
 1130 1140 1150 1160
 1121 A T T A T C T A T C T G A T G A C T G C C C T A A T G T G C C T G A A C T A T A gphkng1815-1.
 1073 A T T A T C T A T C T G A T G A C T G C C C T A A T G T G C C T G A A C T A T A gp7b-1.
 845 A T T A T C T A T C T G A T G A C T G C C C T A A T G T G C C T G A A C T A T A gp7c-1.
 842 A T T A T C T A T C T G A T G A C T G C C C T A A T G T G C C T G A A C T A T A gp7d-1.

C A G A G A A C T C A A T G A G G C C C T C C G A C T G G T C A G T A G A T C C Majority
 1170 1180 1190 1200
 1161 C A G A G A A C T C A A T G A G G C C C T C C G A C T G G T C A G T A G A T C C gphkng1815-1.
 1113 C A G A G A A C T C A A T G A G G C C C T C C G A C T G G T C A G T A G A T C C gp7b-1.
 885 C A G A G A A C T C A A T G A G G C C C T C C G A C T G G T C A G T A G A T C C gp7c-1.
 882 C A G A G A A C T C A A T G A G G C C C T C C G A C T G G T C A G T A G A T C C gp7d-1.

A A T C A G C A A T A C G A C C A G G T G G T G C A G A T G A C C C A G T A T C Majority
 1210 1220 1230 1240
 1201 A A T C A G C A A T A C G A C C A G G T G G T G C A G A T G A C C C A G T A T C gphkng1815-1.
 1153 A A T C A G C A A T A C G A C C A G G T G G T G C A G A T G A C C C A G T A T C gp7b-1.
 925 A A T C A G C A A T A C G A C C A G G T G G T G C A G A T G A C C C A G T A T C gp7c-1.
 922 A A T C A G C A A T A C G A C C A G G T G G T G C A G A T G A C C C A G T A T C gp7d-1.

A C C T G G A A G A C A C C A C G C T T C T G A T G G A G A A G A T G A G A G A Majority
 1250 1260 1270 1280
 1241 A C C T G G A A G A C A C C A C G C T T C T G A T G G A G A A G A T G A G A G A gphkng1815-1.
 1193 A C C T G G A A G A C A C C A C G C T T C T G A T G G A G A A G A T G A G A G A gp7b-1.
 965 A C C T G G A A G A C A C C A C G C T T C T G A T G G A G A A G A T G A G A G A gp7c-1.
 962 A C C T G G A A G A C A C C A C G C T T C T G A T G G A G A A G A T G A G A G A gp7d-1.

G C A G T T T G G C T G G G T T T C T G A A C T G G C A T A C C A G T C C C C A Majority
 1290 1300 1310 1320
 1281 G C A G T T T G G C T G G G T T T C T G A A C T G G C A T A C C A G T C C C C A gphkng1815-1.
 1233 G C A G T T T G G C T G G G T T T C T G A A C T G G C A T A C C A G T C C C C A gp7b-1.
 1005 G C A G T T T G G C T G G G T T T C T G A A C T G G C A T A C C A G T C C C C A gp7c-1.
 1002 G C A G T T T G G C T G G G T T T C T G A A C T G G C A T A C C A G T C C C C A gp7d-1.

G G A G C T G A G G A C A T C T T T A A T C C A G T G A A A G T A A T G G T A G Majority
 1330 1340 1350 1360
 1321 G G A G C T G A G G A C A T C T T T A A T C C A G T G A A A G T A A T G G T A G gphkng1815-1.
 1273 G G A G C T G A G G A C A T C T T T A A T C C A G T G A A A G T A A T G G T A G gp7b-1.
 1045 G G A G C T G A G G A C A T C T T T A A T C C A G T G A A A G T A A T G G T A G gp7c-1.
 1042 G G A G C T G A G G A C A T C T T T A A T C C A G T G A A A G T A A T G G T A G gp7d-1.

C C C T A A G T G C T C A T G A A G G A A A T T C T T C T G A T C A A G A T G A Majority
 1370 1380 1390 1400
 1361 C C C T A A G T G C T C A T G A A G G A A A T T C T T C T G A T C A A G A T G A gphkng1815-1.
 1313 C C C T A A G T G C T C A T G A A G G A A A T T C T T C T G A T C A A G A T G A gp7b-1.
 1085 C C C T A A G T G C T C A T G A A G G A A A T T C T T C T G A T C A A G A T G A gp7c-1.
 1082 C C C T A A G T G C T C A T G A A G G A A A T T C T T C T G A T C A A G A T G A gp7d-1.

FIG. 14E

CACAGTGGTTCCTTCAAGCCTCCTGCCTTCCTCTAACTTC Majority
 1410 1420 1430 1440
 1401 CACAGTGGTTCCTTCAAGCCTCCTGCCTTCCTCTAACTTC gphkng1815-1.
 1353 CACAGTGGTTCCTTCAAGCCTCCTGCCTTCCTCTAACTTC gp7b-1.
 1125 CACAGTGGTTCCTTCAAGCCTCCTGCCTTCCTCTAACTTC gp7c-1.
 1122 CACAGTGGTTCCTTCAAGCCTCCTGCCTTCCTCTAACTTC gp7d-1.

ACACTCAGCAGCCCTCTTGAAAAGAGTGCTGGCAACGCTA Majority
 1450 1460 1470 1480
 1441 ACACTCAGCAGCCCTCTTGAAAAGAGTGCTGGCAACGCTA gphkng1815-1.
 1393 ACACTCAGCAGCCCTCTTGAAAAGAGTGCTGGCAACGCTA gp7b-1.
 1165 ACACTCAGCAGCCCTCTTGAAAAGAGTGCTGGCAACGCTA gp7c-1.
 1162 ACACTCAGCAGCCCTCTTGAAAAGAGTGCTGGCAACGCTA gp7d-1.

ACTTCATTGATCACGTGGTAGAGAAGGTTCTTCAGCACTT Majority
 1490 1500 1510 1520
 1481 ACTTCATTGATCACGTGGTAGAGAAGGTTCTTCAGCACTT gphkng1815-1.
 1433 ACTTCATTGATCACGTGGTAGAGAAGGTTCTTCAGCACTT gp7b-1.
 1205 ACTTCATTGATCACGTGGTAGAGAAGGTTCTTCAGCACTT gp7c-1.
 1202 ACTTCATTGATCACGTGGTAGAGAAGGTTCTTCAGCACTT gp7d-1.

TAAGGAGCACTTTAAAACCTTGGTAAGAAGATTTAGTCCAT Majority
 1530 1540 1550 1560
 1521 TAAGGAGCACTTTAAAACCTTGGTAAGAAGATTTAGTCCAT gphkng1815-1.
 1473 TAAGGAGCACTTTAAAACCTTGGTAAGAAGATTTAGTCCAT gp7b-1.
 1245 TAAGGAGCACTTTAAAACCTTGGTAAGAAGATTTAGTCCAT gp7c-1.
 1242 TAAGGAGCACTTTAAAACCTTGGTAAGAAGATTTAGTCCAT gp7d-1.

CCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGA Majority
 1570 1580 1590 1600
 1561 CCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGA gphkng1815-1.
 1513 CCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGA gp7b-1.
 1285 CCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGA gp7c-1.
 1282 CCTATAATCAGCAAGAATTACACCTTCGGCCAAGACCTGA gp7d-1.

GAAATTCGAAAATACAAAGCAGGCTAACACAATGAACACA Majority
 1610 1620 1630 1640
 1601 GAAATTCGAAAATACAAAGCAGGCTAACACAATGAACACA gphkng1815-1.
 1553 GAAATTCGAAAATACAAAGCAGGCTAACACAATGAACACA gp7b-1.
 1325 GAAATTCGAAAATACAAAGCAGGCTAACACAATGAACACA gp7c-1.
 1322 GAAATTCGAAAATACAAAGCAGGCTAACACAATGAACACA gp7d-1.

GCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGG Majority
 1650 1660 1670 1680
 1641 GCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGG gphkng1815-1.
 1593 GCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGG gp7b-1.
 1365 GCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGG gp7c-1.
 1362 GCTGCATGAAAGTTAGGTATATATTAGGAAGCACTATTGG gp7d-1.

FIG. 14F

TTTACTTTGTTGAATGGAAAGTTTAAATAGCTATTCAAATTG Majority
 1690 1700 1710 1720
 1681 **TTTACTTTGTTGAATGGAAAGTTTAAATAGCTATTCAAATTG** gphkng1815-1.
 1633 **TTTACTTTGTTGAATGGAAAGTTTAAATAGCTATTCAAATTG** gp7b-1.
 1405 **TTTACTTTGTTGAATGGAAAGTTTAAATAGCTATTCAAATTG** gp7c-1.
 1402 **TTTACTTTGTTGAATGGAAAGTTTAAATAGCTATTCAAATTG** gp7d-1.

AGTTAATAATAAAAATTTCTTCCTAAAAAGTAAAAATGTACA Majority
 1730 1740 1750 1760
 1721 **AGTTAATAATAAAAATTTCTTCCTAAAAAGTAAAAATGTACA** gphkng1815-1.
 1673 **AGTTAATAATAAAAATTTCTTCCTAAAAAGTAAAAATGTACA** gp7b-1.
 1445 **AGTTAATAATAAAAATTTCTTCCTAAAAAGTAAAAATGTACA** gp7c-1.
 1442 **AGTTAATAATAAAAATTTCTTCCTAAAAAGTAAAAATGTACA** gp7d-1.

TATGTAGAAATATGATGCATTAGTTCTTTGTATACTAAATA Majority
 1770 1780 1790 1800
 1761 **TATGTAGAAATATGATGCATTAGTTCTTTGTATACTAAATA** gphkng1815-1.
 1713 **TATGTAGAAATATGATGCATTAGTTCTTTGTATACTAAATA** gp7b-1.
 1485 **TATGTAGAAATATGATGCATTAGTTCTTTGTATACTAAATA** gp7c-1.
 1482 **TATGTAGAAATATGATGCATTAGTTCTTTGTATACTAAATA** gp7d-1.

AA TACTGAGTCCCCT Majority
 1810
 1801 **AA TACTGAGTCCCCT** gphkng1815-1.
 1753 **AA TACTGAGTCCCCT** gp7b-1.
 1525 **AA TACTGAGTCCCCT** gp7c-1.
 1522 **AA TACTGAGTCCCCT** gp7d-1.

FIG. 14G


```

1
gphkng1815_aa_ 80
gb7b_aa_
gb7c_aa_
gb7d_aa_
81
gphkng1815_aa_ 160
gb7b_aa_
gb7c_aa_
gb7d_aa_
161
gphkng1815_aa_ 240
gb7b_aa_
gb7c_aa_
gb7d_aa_
241
gphkng1815_aa_ 320
gb7b_aa_
gb7c_aa_
gb7d_aa_
321
gphkng1815_aa_ 400
gb7b_aa_
gb7c_aa_
gb7d_aa_
401
gphkng1815_aa_ 466
gb7b_aa_
gb7c_aa_
gb7d_aa_

```

MKLPLLMPVCLLMLKCHCAPTWKDTAISENANSFSEAGEIDVDGEVKIALIGIKQMKIMMEREEHHSKLMKTLKKC
MKLPLLMPVCLLMLKCHCAPTWKDTAISENANSFSEAGEIDVDGEVKIALIGIKQMKIMMEREEHHSKLMKTLKKC
MKLPLLMPVCLLMLKCHCAPTWKDTAISENANSFSEAGEIDVDGEVKIALIGIKQMKIMMEREEHHSKLMKTLKKC
MKLPLLMPVCLLMLKCHCAPTWKDTAISENANSFSEAGEIDVDGEVKIALIGIKQMKIMMEREEHHSKLMKTLKKC
81
KEEKQEAALKMNEVHEHLEEEESLCOVSLADSNDECRALESNCVRFDTTCQPAMSSVKMVEQFFRKYQFLFPLQEND
KEEKQEAALKMNEVHEHLEEEESLCOVSLADSNDECRALESNCVRFDTTCQPAMSSVKM
KEEKQEAALKMNEVHEHLEEEESLCOVSLADSNDECRALESNCVRFDTTCQPAMSSVKM
KEEKQEAALKMNEVHEHLEEEESLCOVSLADSNDECRALESNCVRFDTTCQPAMSSVKM
161
RSGPVSQGVTEEDAQVSHIEHVFSQLSADVTSLFNRSLYVFKQLRREFDQAFQSYFTSGIDVTEPFFPPLSKEPAYRAD
RSGPVSQGVTEEDAQVSHIEHVFSQLSADVTSLFNRSLYVFKQLRREFDQAFQSYFTSGIDVTEPFFPPLSKEPAYRAD
.....MEPAYRAD
.....MPAYRAD
241
AEPSSWAI PNVFQLLCNLSFSVYQSVSEKLIITLRATEDPPKQDKDSNQGPIISKILPEQDRGSDGKLGQNLSDCVNFRKR
AEPSSWAI PNVFQLLCNLSFSVYQSVSEKLIITLRATEDPPKQDKDSNQGPIISKILPEQDRGSDGKLGQNLSDCVNFRKR
AEPSSWAI PNVFQLLCNLSFSVYQSVSEKLIITLRATEDPPKQDKDSNQGPIISKILPEQDRGSDGKLGQNLSDCVNFRKR
AEPSSWAI PNVFQLLCNLSFSVYQSVSEKLIITLRATEDPPKQDKDSNQGPIISKILPEQDRGSDGKLGQNLSDCVNFRKR
321
CQKQDYLSDDCPNVPPELYRELNEALRLVRSNQQYDQVVMQTYHLEDITLLMEKMRQFGWVSELAYQSPGAEDIFNP
CQKQDYLSDDCPNVPPELYRELNEALRLVRSNQQYDQVVMQTYHLEDITLLMEKMRQFGWVSELAYQSPGAEDIFNP
CQKQDYLSDDCPNVPPELYRELNEALRLVRSNQQYDQVVMQTYHLEDITLLMEKMRQFGWVSELAYQSPGAEDIFNP
CQKQDYLSDDCPNVPPELYRELNEALRLVRSNQQYDQVVMQTYHLEDITLLMEKMRQFGWVSELAYQSPGAEDIFNP
401
VKVMVALSAHEGNSDQDDTVVPSLLPSSNFTLSSPLEKSAGNANFIDHVVEKVLQHFKEHFKTW
VKVMVALSAHEGNSDQDDTVVPSLLPSSNFTLSSPLEKSAGNANFIDHVVEKVLQHFKEHFKTW
VKVMVALSAHEGNSDQDDTVVPSLLPSSNFTLSSPLEKSAGNANFIDHVVEKVLQHFKEHFKTW
VKVMVALSAHEGNSDQDDTVVPSLLPSSNFTLSSPLEKSAGNANFIDHVVEKVLQHFKEHFKTW

FIG. 14H

1 80

bhkng1 -----GCAACCTCGTTGGTGAGAGCCTGCAGTTAGTGTACGGCGGAAACATGAAGCCGC
 bhkng2 -----CAGAAGCTGGTGGCAACCTCGTTGGTGAGAGCCTGCAGTTAGTGTACGGCGGAAACATGAAGCCGC
 bhkng3 GTGAAGGTCCTTACAGAAGCTGGTGGCAACCTCGTTGGTGAGAGCCTGCAGTTAGTGTACGGCGGAAACATGAAGCCGC
 81 160

bhkng1 CACTCTTGGTGTATTATTGTGTATCTGCTGCGGCTGAGAGACTGTGAGTGTGCGCCTACAGGGAAGGACCGAACTTCCATC
 bhkng2 CACTCTTGGTGTATTATTGTGTATCTGCTGCGGCTGAGAGACTGTGAGTGTGCGCCTACAGGGAAGGACCGAACTTCCATC
 bhkng3 CAATCTTGGTGTATTATTGTGTATCTGCTGCGGCTGAGAGACTGTGAGTGTGCGCCTACAGGGAAGGACCGAACTTCCATC
 161 240

bhkng1 CGTGAAGACCCGAAGGGTTTTTCCAAGGCTGGGGAGATAGACGTAGATGAAGAGGTGAAGAAGGCTTTGATTGGCATGAA
 bhkng2 CGTGAAGACCCGAAGGGTTTTTCCAAGGCTGGGGAGATAGACGTAGATGAAGAGGTGAAGAAGGCTTTGATTGGCATGAA
 bhkng3 CGTGAAGACCCGAAGGGTTTTTCCAAGGCTGGGGAGATAGACGTAGATGAAGAGGTGAAGAAGGCTTTGATTGGCATGAA
 241 320

bhkng1 GCAGATGAAAATCCTGATGGAAGAAGAGAGAGGAGGAACATAGCAAATAATGAGAACACTGAAGAAATGCAGAGAAGAAA
 bhkng2 GCAGATGAAAATCCTGATGGAAGAAGAGAGAGGAGGAACATAGCAAATAATGAGAACACTGAAGAAATGCAGAGAAGAAA
 bhkng3 GCAGATGAAAATCCTGATGGAAGAAGAGAGAGGAGGAACATAGCAAATAATGAGAACCCTGAAGAAATGCAGAGAAGAAA
 321 400

bhkng1 AGCAGGAGGCCCTGAAGCTTATGAATGAAGTTCAAGAACATCTAGAAGAGGAAGAAAGGCTATGCCAGGTGTCTCTGATG
 bhkng2 AGCAGGAGGCCCTGAAGCTTATGAATGAAGTTCAAGAACATCTAGAAGAGGAAGAAAGGCTATGCCAGGTGTCTCTGATG
 bhkng3 AGCAGGAGGCCCTGAAGCTTATGAATGAAGTTCAAGAACATCTAGAAGAGGAAGAAAGGCTATGCCAGGTGTCTCTGATG
 401 480

bhkng1 GGTTCTGGGACGAATGCAAATCTGCCTGGAAAGTGAAGTGCATGAGATTTTATAACAACCTGCCAAAGCAGTTGGTCCTC
 bhkng2 GGTTCTGGGACGAATGCAAATCTGCCTGGAAAGTGAAGTGCATGAGATTTTATAACAACCTGCCAAAGCAGTTGGTCCTC
 bhkng3 GGTTCTGGGACGAATGCAAATCTGCCTGGAAAGTGAAGTGCATGAGATTTTATAACAACCTGCCAAAGCAGTTGGTCCTC
 481 560

bhkng1 TATGAAATCCACGATTGAACGGGTTTTCCGGAAGATATATCAGTTTCTCTTCTTCCATGAAGACGATGAAAAAGAGC
 bhkng2 TATGAAATCCACGATTGAACGGGTTTTCCGGAAGATATATCAGTTTCTCTTCTTCCATGAAGACGATGAAAAAGAGC
 bhkng3 TATGAAATCCACGATTGAACGGGTTTTCCGGAAGATATATCAGTTTCTCTTCTTCCATGAAGACGATGAAAAAGAGC
 561 640

bhkng1 TTCCTGTTGGTGAGAAGTTCACTGAGGAAGATGTACAGCTGATGCAGATAGAGAATGTGTTTCCAGCCAGCTGACCGTGGAT
 bhkng2 TTCCTGTTGGTGAGAAGTTCACTGAGGAAGATGTACAGCTGATGCAGATAGAGAATGTGTTTCCAGCCAGCTGACCGTGGAT
 bhkng3 TTCCTGTTGGTGAGAAGTTCACTGAGGAAGATGTACAGCTGATGCAGATAGAGAATGTGTTTCCAGCCAGCTGACCGTGGAT
 641 720

bhkng1 GTGGGATTTCTCTATAACATGAGCTTTACAGTCTTCAAACAGATGCAGCAAGAATTTGACCTGGCTTTTCAATCATACTT
 bhkng2 GTGGGATTTCTCTATAACATGAGCTTTACAGTCTTCAAACAGATGCAGCAAGAATTTGACCTGGCTTTTCAATCATACTT
 bhkng3 GTGGGATTTCTCTATAACATGAGCTTTACAGTCTTCAAACAGATGCAGCAAGAATTTGACCTGGCTTTTCAATCATACTT
 721 800

bhkng1 TATGTCAGACACAGACTCCATGGAGCCTTACTTTTTTCCAGCTTTTTTCCAAAGAGCCAGCAAAAAAGCACATCCTATGC
 bhkng2 TATGTCAGACACAGACTCCATGGAGCCTTACTTTTTTCCAGCTTTTTTCCAAAGAGCCAGCAAAAAAGCACATCCTATGC
 bhkng3 TATGTCAGACACAGACTCCATGGAGCCTTACTTTTTTCCAGCTTTTTTCCAAAGAGCCAGCAAAAAAGCACATCCTATGC
 801 880

bhkng1 AGAGTTGGGACATTCCCAGCTTCTTCCAGCTGTTTTGTAATTTCCAGCCTCTCTGTTTATCAAAGTGTGAGCGCAACAGTT
 bhkng2 AGAGTTGGGACATTCCCAGCTTCTTCCAGCTGTTTTGTAATTTCCAGCCTCTCTGTTTATCAAAGTGTGAGCGCAACAGTT
 bhkng3 AGAGTTGGGACATTCCCAGCTTCTTCCAGCTGTTTTGTAATTTCCAGCCTCTCTGTTTATCAAAGTGTGAGCGCAACAGTT

FIG. 15A

881 960
bhkng1 ACAGAGATGCTGAAGGCCATTGAGGACTTATCCAAACAAGACAAAGATTCTGCCACGGTGGACCGAGTTCCACGACGTG
bhkng2 ACAGAGATGCTGAAGGCCATTGAGGACTTATCCAAACAAGACAAAGATTCTGCCACGGTGGACCGAGTTCCACGACGTG
bhkng3 ACAGAGATGCTGAAGGCCATTGAGGACTTATCCAAACAAGACAAAGATTCTGCCACGGTGGACCGAGTTCCACGACGTG
961 1040
bhkng1 GCCTGTGCGGGCAGAGGGCTGTGTGGAGAACCTGGCCAGAACTCGTCCGAATGTCTCCAATTTATGCAAGATGCCAGA
bhkng2 GCCTGTGCGGGCAGAGGGCTGTGTGGAGAACCTGGCCAGAACTCGTCCGAATGTCTCCAATTTATGCAAGATGCCAGA
bhkng3 GCCTGTGCGGGCAGAGGGCTGTGTGGAGAACCTGGCCAGAACTCGTCCGAATGTCTCCAATTTATGCAAGATGCCAGA
1041 1120
bhkng1 AATGTCAGGATTACCTATGGGCAGACTGCCCTGCTGTTCTGAACTATACACAAAGGCGGATGAGGCCCTTGAGTTGGTC
bhkng2 AATGTCAGGATTACCTATGGGCAGACTGCCCTGCTGTTCTGAACTATACACAAAGGCGGATGAGGCCCTTGAGTTGGTC
bhkng3 AATGTCAGGATTACCTATGGGCAGACTGCCCTGCTGTTCTGAACTATACACAAAGGCGGATGAGGCCCTTGAGTTGGTC
1121 1200
bhkng1 AACATATCCAATCAGCAGTATGCCCAGGTA CTCCAGATGACCCAGCATCACTTGGAGGACACCACGTATCTGATGGAGAA
bhkng2 AACATATCCAATCAGCAGTATGCCCAGGTA CTCCAGATGACCCAGCATCACTTGGAGGACACCACGTATCTGATGGAGAA
bhkng3 AACATATCCAATCAGCAGTATGCCCAGGTA CTCCAGATGACCCAGCATCACTTGGAGGACACCACGTATCTGATGGAGAA
1201 1280
bhkng1 GATGAGAGAGCAGTTTGGTTGGGTAACAGAGCTGGCCAGCCAGACCCAGGAAGCGAGAACATCTTCAGTTTCATAAAGG
bhkng2 GATGAGAGAGCAGTTTGGTTGGGTAACAGAGCTGGCCAGCCAGACCCAGGAAGCGAGAACATCTTCAGTTTCATAAAGG
bhkng3 GATGAGAGAGCAGTTTGGTTGGGTAACAGAGCTGGCCAGCCAGACCCAGGAAGCGAGAACATCTTCAGTTTCATAAAGG
1281 1360
bhkng1 TAGTTCAGGTGTTACGAAGGAAATTTCTCAAACAAGATGAAAAGATGATAGACATAAGCATTCTGCCTTCCTCTAAT
bhkng2 TAGTTCAGGTGTTACGAAGGAAATTTCTCAAACAAGATGAAAAGATGATAGACATAAGCATTCTGCCTTCCTCTAAT
bhkng3 TAGTTCAGGTGTTACGAAGGAAATTTCTCAAACAAGATGAAAAGATGATAGACATAAGCATTCTGCCTTCCTCTAAT
1361 1440
bhkng1 TTCACACTCACCATCCCTCTTGAAGAAAGTGCTGAGAGTCCGACTTCATTAGCTACATGCTGGCCAAAGCTGTACAGCA
bhkng2 TTCACACTCACCATCCCTCTTGAAGAAAGTGCTGAGAGTCCGACTTCATTAGCTACATGCTGGCCAAAGCTGTACAGCA
bhkng3 TTCACACTCACCATCCCTCTTGAAGAAAGTGCTGAGAGTCCGACTTCATTAGCTACATGCTGGCCAAAGCTGTACAGCA
1441 1520
bhkng1 TTTAAGGAACATTTTAAATCTTGGTAAGCAGAGTATTGATTAGGGACGTTTGCTGATAGGAATAGATGGTTCTTAAAA
bhkng2 TTTAAGGAACATTTTAAATCTTGGTAAGCAGAGTATTGATTAGGGACGTTTGCTGATAGGAATAGATGGTTCTTAAAA
bhkng3 TTTAAGGAACATTTTAAATCTTGGTAAGCAGAGTATTGATTAGGGACGTTTGCTGATAGGAATAGATGGTTCTTAAAA
1521 1600
bhkng1 GGGAAAAATGACAAAACCTAGCTTTTGAATACCTTGAAAACGTATTCAACCTCATTATAATCAAAGGCATGAAAACCTAAG
bhkng2 GGGAAAAATGACAAAACCTAGCTTTTGAATACCTTGAAAACGTATTCAACCTCATTATAATCAAAGGCATGAAAACCTAAG
bhkng3 GGGAAAAATGACAAAACCTAGCTTTTGAATACCTTGAAAACGTATTCAACCTCATTATAATCAAAGGCATGAAAACCTAAG
1601 1680
bhkng1 ACAAGTTAGCAGTTTTTACCTATTGAATTTCAAATTA AAAAAAAAAAATCCTGATAGAATGCAATGAAATGAGAATTCTT
bhkng2 ACAAGTTAGCAGTTTTTACCTATTGAATTTCAAATTA AAAAAAAAAAATCCTGATAGAATGCAATGAAATGAGAATTCTT
bhkng3 ACAAGTTAGCAGTTTTTACCTATTGAATTTCAAATTA AAAAAAAAAA . TCCTGATAGAATGCAATGAAATGAGAATTCTT
1681 1760
bhkng1 ATATGTGATTGCCAGAAACAACTGGTTTTGTCTTTTGA AAAAGTTATTCAATTATACATATCAAGAGTCATCAAATTC
bhkng2 ATATGTGATTGCCAGAAACAACTGGTTTTGTCTTTTGA AAAAGTTATTCAATTATACATATCAAGAGTCATCAAATTC
bhkng3 ATATGTGATTGCCAGAAACAACTGGTTTTGTCTTTTGA AAAAGTTATTCAATTATACATATCAAGAGTCATCAAATTC

FIG. 15B

1761 1840
bhkng1 TTTTAAATAATAAATCCACTTCTGGAATCAATCCAAAGGAGTAAATCTAAAATGAATTGAAGTTCCCACCCCAAGAT
bhkng2 TTTTAAATAATAAATCCACTTCTGGAATCAATCCAAAGGAGTAAATCTAAAATGAATTGAAGTTCCCACCCCAAGAT
bhkng3 TTTTAAATAATAAATCCACTTCTGGAATCAATCCAAAGGAGTAAATCTAAAATGAATTGAAGTTCCCACCCCAAGAT
1841 1920
bhkng1 CAATATTTGCAAATTATTTAAAATAGTAAACTGTAAAACTGAATGTCATCTGAATGCTAAAAACCAGAAATGGTTAA
bhkng2 CAATATTTGCAAATTATTTAAAATAGTAAACTGTAAAACTGAATGTCATCTGAATGCTAAAAACCAGAAATGGTTAA
bhkng3 CAATATTTGCAAATTATTTAAAATAGTAAACTGTAAAACTGAATGTCATCTGAATGCTAAAAACCAGAAATGGTTAA
1921 2000
bhkng1 AAGCTGTGGCTAAATATGCTCCAAATATCTTATAAAACCATTAAAAATTTTATAAAATTTAAATCATGACATGACATCT
bhkng2 AAGCTGTGGCTAAATATGCTCCAAATATCTTATAAAACCATTAAAAATTTTATAAAATTTAAATCATGACATGACATCT
bhkng3 AAGCTGTGGCTAAATATGCTCCAAATATCTTATAAAACCATTAAAAATTTTATAAAATTTAAATCATGACATGACATCT
2001 2080
bhkng1 GCTGGAACAAGAGTTTATTCTAAGCCTATCTATAAGGCAAATATTATTACTATCTTCCAGAAAAGAACTTGAGACT
bhkng2 GCTGGAACAAGAGTTTATTCTAAGCCTATCTATAAGGCAAATATTATTACTATCTTCCAGAAAAGAACTTGAGACT
bhkng3 GCTGGAACAAGAGTTTATTCTAAGCCTATCTATAAGGCAAATATTATTACTATCTTCCAGAAAAGAACTTGAGACT
2081 2160
bhkng1 CAGGGTCCAAGTGTAGTTGCTCAGTCATGTCTGACTCTTTGGGACCCCTGGACTGTAGCCCACCAGGCTCCTCTGTCC
bhkng2 CAGGGTCCAAGTGTAGTTGCTCAGTCATGTCTGACTCTTTGGGACCCCTGGACTGTAGCCCACCAGGCTCCTCTGTCC
bhkng3 CAGGGTCCAAGTGTAGTTGCTCAGTCATGTCTGACTCTTTGGGACCCCTGGACTGTAGCCCACCAGGCTCCTCTGTCC
2161 2240
bhkng1 GTGGGATTCTTCAGACAGGAATACTGGGGCAGGTTGCTATTTCCCTTCTCCAGGAAATCTTCCCTATCCAGGGATGGAACC
bhkng2 ATGGGATTCTTCAGACAAGAATACTGGAGCAGGTTGCTATTTCCCTTCTCCAGGAAATCTTCCCTATCCAGGGATGGAACC
bhkng3 ATGGGATTCTTCAGACAAGAATACTGGAGCAGGTTGCTATTTCCCTTCTCCAGGAAATCTTCCCTATCCAGGGATGGAACC
2241 2320
bhkng1 CAGGTCTCCTGCATTGCAGGTAGATGCTTTACTATCTGAGCAACCAAATGAATTAATCAAGTCAGTAGGGGGTAGAGGCA
bhkng2 CAGGTCTCCTGCATTGCAGGTAGATGCTTTACTATCTGAGCAACCAAATGAATTAATCAAGTCAGTAGGGGGTAGAGGCA
bhkng3 CAGGTCTCCTGCATTGCAGGTAGATGCTTTACTATCTGAGCAACCAAATGAATTAATCAAGTCAGTAGGGGGTAGAGGCA
2321 2400
bhkng1 AATTTAACTTAGTTTTCTCTGAATCATAATGCCACATTAACTGGTTCCCTGTTGGGACATTTGGTTGAAAAAATAAA
bhkng2 AATTTAACTTAGTTTTCTCTGAATCATAATGCCACATTAACTGGTTCCCTGTTGGGACATTTGGTTGAAAAAATAAA
bhkng3 AATTTAACTTAGTTTTCTCTGAATCATAATGCCACATTAACTGGTTCCCTGTTGGGACATTTGGTTGAAAAAATAAA
2401 2480
bhkng1 GTGAAAAATGAGTATAAACTCTATAAATGTAATGATCAAAACGAAAAAAATCTACAATCTGCATTAAAAATAAAAAGG
bhkng2 GTGAAAAATGAGTATAAACTCTATAAATGTAATGATCAAAACGAAAAAAATCTACAATCTGCATTAAAAATAAAAAGG
bhkng3 GTGAAAAATGAGTATAAACTCTATAAATGTAATGATCAAAACGAAAAAAATCTACAATCTGCATTAAAAATAAAAAGG
2481 2560
bhkng1 GTGGCAGG.....
bhkng2 GTGGCAGGAATTACGGTTGGAAATGGATGATTTTTTTTAACTTTTCATCTTTTGATATTTTACAATTTTCTATAATGA
bhkng3 GTGGCAGG.....
2561 2640
bhkng1
bhkng2 ATAAATAATTTTGAGATTTCAAATTAGAAGATATGTTGCTAAAATAGCTAGGTAATGTAGATTGAACACTGTATCAATG
bhkng3
2641 2720
bhkng1
bhkng2 TGTTCTCATCTTTAACTTTAGTATAAGTACTTCTATTCCATGGTAATCCTACAGTAAGACGAAATGTAATCTGTTCCG
bhkng3
2721 2800
bhkng1
bhkng2 TCTACAGGAAAAACAATAATGACATTTAGACGTAACATTACCATCTCTGTTAGGATAATCTTCTGAATTAATGGCACA
bhkng3
2801 2880
bhkng1
bhkng2 ATTAGAAGTACATAGTATTCTCCTTTGGTAAAATGGTCAATCTTAAAGAAGCATTAAATGTTAATTCTAAGTTATTAC
bhkng3
2881 2960
bhkng1
bhkng2 TCATAAGGACCTTGTTAGGTAGGTCCTATCAATGTATAATTAAGCTGGGTATTTCTAGATTGCTGCTCCTCTCCCTTTAT
bhkng3
2961 3029
bhkng1
bhkng2 CTCTGAATGTTGGAGAGGTTGTTGGTCATCAATCAACCAATATCTTTTAGCATCTTCTAAGTGAAGGC
bhkng3

FIG. 15C


```

1
hmhknng_aa      80
bhknng1_aa     80
gphknng1815_aa 80
-----MKPPLLVFIVYLLRLRDCQCAPTGKDRTSIREDPKGFSKAGEIDVDEEVKK
81
hmhknng_aa      160
bhknng1_aa     160
gphknng1815_aa 160
-----MKLPLLMPVCLLWLKDKCHCAPTWKDKTAISENANSFSEAGEIDVDGEVKI
ALTGIKQMKIMMERKEKEHTNLMSTLKKCREEKQEALKLLNEVQEHLEEEERLCRESLADSWGECRSCLENNCMRIYTTTC
ALIGMKQMKILMERREEEHSKLMRTLKKCREEKQEALKLMNEVQEHLEEEERLCQVSLMGSWDECKSCLESDCMRFYTTTC
ALIGIKQMKIMMERREEEHSKLMKTLKKCKEEKQEALKLMNEVHEHLEEEESLCQVSLADSWDECRACLESNMRFDTTTC
161
hmhknng_aa      240
bhknng1_aa     240
gphknng1815_aa 240
QPSWSSVKNKIERFFRKIYQFLPFFHEDNEKDLPISSEKLIIEEDAQLTQMEDVFSQTLVDVNSLNFNRSFNVFRMQQEFDQ
QSSWSSMKSTIERVFRKIYQFLPFFHEDDEKELPVGEKFTEDVQLMQIENVFSQTLVDVGFLYNMSFHVFKMQQEFDL
QPAWSSVKNMVEQFFRKIYQFLPFLQE.NDRSGPVSKGVTEEDAQVSHIEHVFSQLSADVTSLFNRSLYVFKQLRREFDQ
241
hmhknng_aa      320
bhknng1_aa     320
gphknng1815_aa 320
TFQSHFISDTDLTEPYFFPAFSKEPMTKADLEQCWDIPNFFQLFCNFSVSIYESVSETITKMLKAIEDLPKQDKAPDHGG
AFQSYFMSDIDSMEPYFFPAFSKEPAKKAHPMQSWDIPSPFFQLFCNFSLSVYQSVSATVTEMLKAIEDLSKQDKDSAHGG
AFQSYFTSGTDVTEPFFPFLSKEPAYRADAEP SWAIPNVFQLLNLFSVYQSVSEKLITTLRATEDPPKQDKDSNQQG
321
hmhknng_aa      400
bhknng1_aa     400
gphknng1815_aa 400
LISKMLPGQDRGLCGELDONLSRCFKFHEKQKQCAHLSQDCPDVPALHTELDEAIRLVNVSNOQYQGQILQMTKHLLEDT
PSSTTWPVRGRGLCGEPGQNSSECLQFHARCQKQDYLDWADCPAVPELYTKADEALELVNISNOQYAQVLQMTQHLEDT
PISKILPEQDRGSDGKLGQNLSDCVNFRKRCQKQDYLDSDDCPNVPELYRELNEALRLVSRSNQOYDQVQMTQYHLEDT
401
hmhknng_aa      480
bhknng1_aa     480
gphknng1815_aa 480
AYLVEKMRGQFGWVSELANQAPETEIIFNSIQVVPRI..HEGNISKQDETMMDLSILPSSNFTLKIPLLESAESSNFIG
TYLMEKMREQFGWVTELASQTPGSENIFFSPIKVVPGV..HEGNFSKQDE.KMIDISILPSSNFTLTIPLEESAESSDFIS
TLLMEKMREQFGWVSELAYQSPGAEDIFNPVKVMVALSAHEGNSSDQDD.TVVPSSLLPSSNFTLSSPLEKSAGNANFID
481
hmhknng_aa      497
bhknng1_aa     497
gphknng1815_aa 497
YVAKALQHFKEHFKTW
YMLAKAVQHFKEHFKSW
HVVEKVLQHFKEHFKTW

```

FIG. 16

matureHKNG -----APTWKDKTAIS
 HKNG1-V1-IPF3 -----MRTWDYSNSGNMKPPLLVFIVCLLWLKDSHCAPTWKDKTAIS
 HKNG1/1-V1-IPF2 -----MKPPLLVFIVCLLWLKDSHCAPTWKDKTAIS
 HKNG1-IPF1 MKIKAENEKNEGPSRSWWQLHWGDIANNSGNMKPPLLVFIVCLLWLKDSHCAPTWKDKTAIS

matureHKNG ENLKSFSEVGEIDADEEVKKALTGIKQMKIMMERKEKEHTNLMSTLKKCREEKQEALKLL
 HKNG1-V1-IPF3 ENLKSFSEVGEIDADEEVKKALTGIKQMKIMMERKEKEHTNLMSTLKKCREEKQEALKLL
 HKNG1/1-V1-IPF2 ENLKSFSEVGEIDADEEVKKALTGIKQMKIMMERKEKEHTNLMSTLKKCREEKQEALKLL
 HKNG1-IPF1 ENLKSFSEVGEIDADEEVKKALTGIKQMKIMMERKEKEHTNLMSTLKKCREEKQEALKLL

matureHKNG NEVQEHLEEEERLCRESLADSWGECRSCLENNCMRIYTTTCQPSWSSVKNKIERFFRKIYQ
 HKNG1-V1-IPF3 NEVQEHLEEEERLCRESLADSWGECRSCLENNCMRIYTTTCQPSWSSVKNKIERFFRKIYQ
 HKNG1/1-V1-IPF2 NEVQEHLEEEERLCRESLADSWGECRSCLENNCMRIYTTTCQPSWSSVKNKIERFFRKIYQ
 HKNG1-IPF1 NEVQEHLEEEERLCRESLADSWGECRSCLENNCMRIYTTTCQPSWSSVKNKIERFFRKIYQ

matureHKNG FLFPFHEDNEKDLPISEKLI EEDAQLTQMEDVFSQLTVDVNSLNFNRSFNVFRQMQQEFDQ
 HKNG1-V1-IPF3 FLFPFHEDNEKDLPISEKLI EEDAQLTQMEDVFSQLTVDVNSLNFNRSFNVFRQMQQEFDQ
 HKNG1/1-V1-IPF2 FLFPFHEDNEKDLPISEKLI EEDAQLTQMEDVFSQLTVDVNSLNFNRSFNVFRQMQQEFDQ
 HKNG1-IPF1 FLFPFHEDNEKDLPISEKLI EEDAQLTQMEDVFSQLTVDVNSLNFNRSFNVFRQMQQEFDQ

matureHKNG TFQSHFISDIDLTEPYFFPAFSKEPMTKADLEQCWDIPNFFQLFCNFSVSIYESVSETIT
 HKNG1-V1-IPF3 TFQSHFISDIDLTEPYFFPAFSKEPMTKADLEQCWDIPNFFQLFCNFSVSIYESVSETIT
 HKNG1/1-V1-IPF2 TFQSHFISDIDLTEPYFFPAFSKEPMTKADLEQCWDIPNFFQLFCNFSVSIYESVSETIT
 HKNG1-IPF1 TFQSHFISDIDLTEPYFFPAFSKEPMTKADLEQCWDIPNFFQLFCNFSVSIYESVSETIT

matureHKNG KMLKAI EDLPKQDKAPDHGGLISKMLPGQDRGLCGELDQNL SRCFKFHEKCQKCQAHLSE
 HKNG1-V1-IPF3 KMLKAI EDLPKQDKAPDHGGLISKMLPGQDRGLCGELDQNL SRCFKFHEKCQKCQAHLSE
 HKNG1/1-V1-IPF2 KMLKAI EDLPKQDKAPDHGGLISKMLPGQDRGLCGELDQNL SRCFKFHEKCQKCQAHLSE
 HKNG1-IPF1 KMLKAI EDLPKQDKAPDHGGLISKMLPGQDRGLCGELDQNL SRCFKFHEKCQKCQAHLSE

matureHKNG DCPDVPALHTELDEAIRLVNVSNOQYGOILOMTRKHLEDTAYLVEKMRGQFGWVSELANQ
 HKNG1-V1-IPF3 DCPDVPALHTELDEAIRLVNVSNOQYGOILOMTRKHLEDTAYLVEKMRGQFGWVSELANQ
 HKNG1/1-V1-IPF2 DCPDVPALHTELDEAIRLVNVSNOQYGOILOMTRKHLEDTAYLVEKMRGQFGWVSELANQ
 HKNG1-IPF1 DCPDVPALHTELDEAIRLVNVSNOQYGOILOMTRKHLEDTAYLVEKMRGQFGWVSELANQ

matureHKNG APETEIIFN SIQVV PRIHEGNISKQDETMMTDLSILPSSNFTLKI PLEESAESSNFIGYV
 HKNG1-V1-IPF3 APETEIIFN SIQVV PRIHEGNISKQDETMMTDLSILPSSNFTLKI PLEESAESSNFIGYV
 HKNG1/1-V1-IPF2 APETEIIFN SIQVV PRIHEGNISKQDETMMTDLSILPSSNFTLKI PLEESAESSNFIGYV
 HKNG1-IPF1 APETEIIFN SIQVV PRIHEGNISKQDETMMTDLSILPSSNFTLKI PLEESAESSNFIGYV

matureHKNG VAKALQHFKEHFKTW
 HKNG1-V1-IPF3 VAKALQHFKEHFKTW
 HKNG1/1-V1-IPF2 VAKALQHFKEHFKTW
 HKNG1-IPF1 VAKALQHFKEHFKTW

FIG. 17

R H L Q A R A A G L V S T L E V A D T 19
 TG CGT CAC CTG CAG GCC CGG GCC GCG GGG TTG GTT TCC ACC CTG GAG GTT GCT GAC ACC 57

L C P R L T S S R W H R R L Q G A A L K 39
 CTG TGC CCT CGG CTG ACT TCC AGC CGG TGG CAC AGA CGC CTC CAG GGG GCA GCA CTC AAG 117

R I L G M T E L R P S L L P G W S S V A 59
 CGC ATC TTA GGA ATG ACA GAG TTG CGT CCC TCT CTG TTG CCA GGC TGG AGT TCA GTG GCA 177

C S * L T E A S N S W V Q V T L P P Q P 79
 TGT TCT TAG CTC ACT GAA GCC TCA AAT TCC TGG GTT CAA GTG ACC CTC CCA CCT CAG CCC 237

H E D L G L Q D T A K S L T R M K I K A 99
 CAT GAG GAC CTG GGA CTA CAG GAC ACA GCT AAA TCC CTG ACA CGG ATG AAA ATT AAA GCA 297

E K N E G P S R S W W Q L H W G D I A N 119
 GAG AAA AAC GAA GGT CCT TCC AGA AGC TGG TGG CAA CTT CAC TGG GGA GAT ATT GCA AAT 357

N S G N M K P P L L V F I V C L L W L K 139
 AAC AGC GGG AAC ATG AAG CCG CCA CTC TTG GTG TTT ATT GTG TGT CTG CTG TGG TTG AAA 417

D S H C A P T W K D K T A I S E N L K S 159
 GAC AGT CAC TGC GCA CCC ACT TGG AAG GAC AAA ACT GCT ATC AGT GAA AAC CTG AAG AGT 477

F S E V G E I D A D E E V K K A L T G I 179
 TTT TCT GAG GTG GGG GAG ATA GAT GCA GAT GAA GAG GTG AAG AAG GCT TTG ACT GGT ATT 537

K Q M K I M M E R K E K E H T N L M S T 199
 AAG CAA ATG AAA ATC ATG ATG GAA AGA AAA GAG AAG GAA CAC ACC AAT CTA ATG AGC ACC 597

L K K C R E E K Q E A L K L L N E V Q E 219
 CTG AAG AAA TGC AGA GAA GAA AAG CAG GAG GCC CTG AAA CTT CTG AAT GAA GTT CAA GAA 657

H L E E E E R L C R E S L A D S W G E C 239
 CAT CTG GAG GAA GAA GAA AGG CTA TGC CGG GAG TCT TTG GCA GAT TCC TGG GGT GAA TGC 717

R S C L E N N C M R I Y T T C Q P S W S 259
 AGG TCT TGC CTG GAA AAT AAC TGC ATG AGA ATT TAT ACA ACC TGC CAA CCT AGC TGG TCC 777

S V K N K L L T T E A * F Q R C Y L G R 279
 TCT GTG AAA AAT AAG CTC CTG ACC ACG GAG GCC TGA TTT CAA AGA TGT TAC CTG GGC AGG 837

T E D C V G N L T R I C Q D V S N F M K 299
 ACA GAG GAC TGT GTG GGG AAC TTG ACC AGA ATT TGT CAA GAT GTT TCA AAT TTC ATG AAA 897

N A K N V R L T Y L K T V L M Y L L C T 319
 AAT GCC AAA AAT GTC AGG CTC ACC TAT CTG AAG ACT GTC CTG ATG TAC CTG CTC TGC ACA 957

Q N * T R R S G W S M Y P I S S M A R F 339
 CAG AAT TAG ACG AGG CGA TCA GGT TGG TCA ATG TAT CCA ATC AGC AGT ATG GCC AGA TTC 1017

S R * P G S T W R T P P I W W R R * E G 359
 TCC AGA TGA CCC GGA AGC ACT TGG AGG ACA CCG CCT ATC TGG TGG AGA AGA TGA GAG GGC 1077

N L A G C L N W Q T R P Q K Q R S S L I 379
 AAT TTG GCT GGG TGT CTG AAC TGG CAA ACC AGG CCC CAG AAA CAG AGA TCA TCT TTA ATT 1137

Q Y R * F Q G F M K E I F P N K M K Q * 399
 CAA TAC AGG TAG TTC CAA GGA TTC ATG AAG GAA ATA TTT CCA AAC AAG ATG AAA CAA TGA 1197

* Q T * A F C L P L I S H S R S L L K K 419
 TGA CAG ACT TAA GCA TTC TGC CTT CCT CTA ATT TCA CAC TCA AGA TCC CTC TTG AAG AAA 1257

V L R V L T S L A T * W Q K L Y S I L R 439
 GTG CTG AGA GTT CTA ACT TCA TTG GCT ACG TAG TGG CAA AAG CTC TAC AGC ATT TTA AGG 1317

FIG. 18A

WO 99/47535

57/57

PCT/US99/05606

N	I	L	K	P	G	K	K	I	*	C	I	L	Y	P	V	S	R	I	I	459
AAC	ATT	TTA	AAA	CCT	GGT	AAG	AAG	ATC	TAA	TGC	ATC	CTA	TAT	CCA	GTA	AGT	AGA	ATT	ATC	1377
S	S	S	G	T	W	K	S	*	N	K	K	G	*	C	N	K	H	S	C	479
TCT	TCA	TCT	GGG	ACC	TGG	AAA	TCC	TGA	AAT	AAA	AAA	GGA	TAA	TGC	AAT	AAA	CAC	AGT	TGC	1437
R	K	V	C	*	L	Y	T	M	K	Y	S	*	F	T	Y	V	E	W	L	499
AGG	AAA	GTA	TGT	TAG	CTA	TAT	ACT	ATG	AAG	TAC	TCT	TAG	TTT	ACT	TAT	GTT	GAA	TGG	CTT	1497
S	Y	*	Y	S	N	*	V	K	M	K	I	P	P	*	K	I	K	R	N	519
AGC	TAT	TAA	TAC	TCA	AAT	TGA	GTT	AAA	ATG	AAA	ATT	CCT	CCT	TAA	AAA	ATC	AAA	CGT	AAT	1557
M	Y	Y	I	S	W	Y	I	S	S	S	L	Y	I	E	*	I	L	N	H	539
ATG	TAT	TAC	ATT	TCA	TGG	TAC	ATT	AGT	AGT	TCT	TTG	TAT	ATT	GAA	TAA	ATA	CTA	AAT	CAC	1617
L																				540
CTA																				1620

FIG. 18B