

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3768846 T5**

(12) **Rettet oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **C 12 N 15/85 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2024-08-19**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-06-21**
- (86) Europæisk ansøgning nr.: **19715345.5**
- (86) Europæisk indleveringsdag: **2019-03-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2021-01-27**
- (86) International ansøgning nr.: **US2019023209**
- (87) Internationalt publikationsnr.: **WO2019183248**
- (30) Prioritet: **2018-03-21 US 201862645892 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Aldevron, L.L.C., 4055 41st Avenue, South Fargo, ND 58104, USA**
- (72) Opfinder: **WILLIAMS, James A., 6420 Pueblo Ct., Lincoln, Nebraska 68516, USA**
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **VIRALE OG IKKE-VIRALE NANOPLASMIDVEKTORER MED FORBEDRET PRODUKTION**
- (56) Fremdragne publikationer:
US-B1- 6 709 844
JIAMIAO LU ET AL: "A 5' Noncoding Exon Containing Engineered Intron Enhances Transgene Expression from Recombinant AAV Vectors in vivo", HUMAN GENE THERAPY, vol. 28, no. 1, 1 January 2017 (2017-01-01), pages 125-134, XP055587331, US ISSN: 1043-0342, DOI: 10.1089/hum.2016.140 cited in the application
WILLIAMS J A ET AL: "Plasmid DNA vaccine vector design: Impact on efficacy, safety and upstream production", BIOTECHNOLOGY ADVANCES, ELSEVIER PUBLISHING, BARKING, GB, vol. 27, no. 4, 1 July 2009 (2009-07-01), pages 353-370, XP026127885, ISSN: 0734-9750, DOI: 10.1016/J.BIOTECHADV.2009.02.003 [retrieved on 2009-02-20]
WO-A1-2014/035457
WO-A1-2014/077863
US-A1- 2015 275 221

DESCRIPTION

FIELD OF THE INVENTION

[0001] The present invention relates to methods for improving viral and non-viral vector manufacturing yield and quality, reducing transfection associated toxicity, improving transposition from non-viral transposon vectors, improving packaging titers from viral vectors, improving expression of viral and non-viral vector encoded genes, and for eliminating viral vector and non-viral vector mediated antibiotic selection marker gene transfer.

BACKGROUND OF THE INVENTION

[0002] *E. coli* plasmids have long been an important source of recombinant DNA molecules used by researchers and by industry. Today, plasmid DNA is becoming increasingly important as the next generation of biotechnology products (e.g. gene medicines and DNA vaccines) make their way into clinical trials, and eventually into the pharmaceutical marketplace. Plasmid DNA vaccines may find application as preventive vaccines for viral, bacterial, or parasitic diseases; immunizing agents for the preparation of hyper immune globulin products; therapeutic vaccines for infectious diseases; or as cancer vaccines. Plasmids are also utilized in gene therapy or gene replacement applications, wherein the desired gene product is expressed from the plasmid after administration to a patient. Plasmids are also utilized in non-viral transposon vectors for gene therapy or gene replacement applications, wherein the desired gene product is expressed from the genome after transposition from the plasmid and genome integration. Plasmids are also utilized in viral vectors for gene therapy or gene replacement applications, wherein the desired gene product is packaged in a transducing virus particle after transfection of a production cell line, and is then expressed from the virus in a target cell after viral transduction.

[0003] Non-viral and viral vector plasmids typically contain a pMB1, ColE1 or pBR322 derived replication origin. Common high copy number derivatives have mutations affecting copy number regulation, such as ROP (Repressor of primer gene) deletion, with a second site mutation that increases copy number (e.g. pMB1 pUC G to A point mutation, or ColE1 pMM1). Higher temperature (42°C) can be employed to induce selective plasmid amplification with pUC and pMM1 replication origins.

[0004] Carnes, AE and Williams, JA, 2011 US Patent 7943377 describe methods for fed-batch fermentation, in which plasmid-containing *E. coli* cells were grown at a reduced temperature during part of the fed-batch phase, during which growth rate was restricted, followed by a temperature upshift and continued growth at elevated temperature in order to accumulate plasmid; the temperature shift at restricted growth rate improved plasmid yield and purity. Other fermentation processes for plasmid production are described in Carnes A.E. 2005 BioProcess Intl 3:36-44.

[0005] The art teaches that one of the limitations of application of plasmid therapies and plasmid vaccines is regulatory agency (e.g. Food and Drug Administration, European Medicines Agency) safety concerns regarding: 1) plasmid transfer and replication in endogenous bacterial flora, or 2) plasmid encoded selection marker expression in human cells, or endogenous bacterial flora. Additionally, regulatory agency guidance's recommend removal of all non-essential sequences in a vector. Plasmids containing a pMB1, ColE1 or pBR322 derived replication origin can replicate promiscuously in *E. coli* hosts. This presents a safety concern that a plasmid therapeutic gene or antigen will be transferred to, and replicated within, a patient's endogenous flora. Ideally, a therapeutic or vaccine plasmid would be replication incompetent in endogenous *E. coli* strains. This requires replacement of the pMB1, ColE1 or pBR322 derived replication origin with a conditional replication origin that requires a specialized cell line for propagation. As well, regulatory agencies, such as the EMEA and FDA, are concerned with utilization of antibiotic resistance or alternative protein markers in gene therapy and gene vaccine vectors, due to concerns that the gene (antibiotic resistance marker or protein marker) may be expressed in a patient's cells. Ideally, plasmid therapies and plasmid vaccines would: 1) be replication incompetent in endogenous *E. coli* strains, 2) not encode a protein-based selection marker and 3) be minimized to eliminate all non-essential sequences.

[0006] The art further teaches that one of the limitations of application of plasmid vectors is that transgene expression duration from plasmid vectors is reduced due to promoter inactivation mediated by the bacterial region (i.e. the region encoding the bacterial replication origin and selectable marker) of the vector (Chen ZY, He CY, Meuse L, Kay MA. 2004. Gene Ther 11:856-864; Suzuki M, Kasai K, Saeki Y. 2006. J Virol 80:3293-3300). This results in short duration transgene expression. A strategy to improve transgene expression duration is to remove the bacterial region of the plasmid. For example, minicircle vectors have been developed which do not contain a bacterial region. Removal of the bacterial region in minicircle vectors improved transgene expression duration (Chen *et al.*, *Supra*, 2004). In minicircle vectors, the eukaryotic region polyadenylation signal is covalently linked to the eukaryotic region promoter through a short spacer typically less than 200 bp comprised of the recombined attachment sites. This linkage (spacer region) can tolerate a much longer spacer sequence since while long spacers ≥ 1 kb in length resulted in transgene expression silencing *in vivo*, shorter spacers ≤ 500 bp exhibited similar transgene expression patterns to conventional minicircle DNA vectors (Lu J, Zhang F, Xu S, Fire AZ, Kay MA. 2012. Mol Ther. 20:2111-9).

[0007] Williams, 2014. DNA plasmids with improved expression. World Patent Application WO2014035457 disclose minimized Nanoplasmid™ vectors utilize RNA-OUT antibiotic-free selection and replace the large 1000 bp pUC replication origin with a novel, 300 bp, R6K origin. Reduction of the spacer region linking the 5' and 3' ends of the transgene expression cassette to < 500 bp with R6K origin-RNA-OUT backbones improved expression duration to that of conventional minicircle DNA vectors as expected from the teachings of Lu *et al.*, *Supra*, 2012.

[0008] The 1.1 kb pFAR4 vector pUC-origin tRNA antibiotic free selection spacer has improved expression duration compared to a 2.2 kb pUC origin-kanR antibiotic selection marker spacer region (Quiviger, M, Arfi A, Mansard D, Delacotte L, Pastor M, Scherman D, Marie C. 2014. Gene Therapy 21: 1001-1007). This teaches that improved expression duration can be obtained with some bacterial regions up to 1.1 kb.

[0009] Expression level improvement compared to plasmid vectors is also observed with some spacer regions ≤ 1.1 kb. For example, pVAX1 derivatives with the 2 kb bacterial backbone reduced to 1.2, 1.1 or 0.7 kb show ≥ 2 -fold improved expression compared to the parent pVAX1 vector (Table 1). NTC8685 derivatives with the 1.5 kb bacterial backbone reduced to 0.9 kb, 466 bp or 281 bp (Nanoplasmid™ vectors) show ≥ 2 -fold improved expression compared to the parent NTC8685 vector (Table 2).

[0010] This teaches that improved expression level can be obtained with short bacterial regions up to 1.2 kb.

Table 1: pVAX1 mammalian expression vector spacer region (SR) derivatives expression level

pVAX1 Vector derivatives	Bacterial Backbone (bp)	Selection Marker	Selection	In Vitro Expression	In vivo Expression
pVAX1	2 kb	KanR	Kan	1X (baseline)	1X (baseline)
pVAX1-A2 ^a	1.2 kb	RNA-OUT	Sucrose	4x	ND
pVAX1/MINI ^b	0.7 kb	RNAI (pUC origin)	murA translation	2x	ND
pFAR4 ^c	1.1 kb	tRNA (amber)	thyA amber mutation suppression	ND	>5x ID EP

pVAX1 Vector derivatives	Bacterial Backbone (bp)	Selection Marker	Selection	In Vitro Expression	In vivo Expression
Minicircle ^d	50 bp	NA, Minicircle	NA, Minicircle	1.4-1.6x (equimolar dose)	1.3x IM (equimolar dose)

^a Games et al., (2010) J. Gene Medicine 12:818-31

^b Ribeiro et al., (2011) Cell Reprogram 14:130

^c Marie et al., (2010) J. Gene Medicine 12:323-32

^d Wang et al., (2014) J Virology 88: 1924-34

Table 2: NTC8685 mammalian expression vector spacer region (SR) derivatives expression level

NTC8685 Vector derivatives ^a	Replication origin	Bacterial backbone (bp)	Selection Marker	In vitro Expression (A549)	In vivo Expression (ID + EP)
NTC8685	pUC origin	1.5 kb	RNA-OUT	1X (baseline)	1X (baseline)
NTC8385-Min	Minimalized pUC origin	0.9 kb	RNA-OUT	2.1x	1.4-1.9 x
NTC9385R Nanoplasmid™	R6K origin	466 bp	RNA-OUT	2.5x	5.3-6.3x
NTC9385C Nanoplasmid™	ColE2 origin	281 bp	RNA-OUT	1.1x	2.7-3.1x

^a Williams, *Supra*, 2014

[0011] Various investigators have identified that minicircle vectors are superior to plasmid vectors for production of AAV vectors (improved transducing unit titers - Table 3) and transposon vectors (increased transposition - Table 3). The improved performance due to improved expression duration with short backbone minicircle vectors should also be observed with short bacterial backbone plasmid vectors up to 1.1 kb.

Table 3: Minicircle applications with various viral and non-viral vector platforms

Vector platform	pUC Plasmid (>1.5 kb pUC origin-antibiotic marker spacer region)	Minicircle (MC) retrofit (<400 bp spacer region)	Result-performance	Result-antibiotic marker gene transfer
self-complementary (sc) AAV	pAAV-scGFP	MC.AAVscGFP	Up to 30-fold improved Transducing units ^a	Plasmid backbone antibiotic resistance marker packaged in up to 26.1% viral particles ^a potential for genome integration in transduced cells
single-stranded (ss) AAV	AAV2-ssGFP	MC.AAV-ssGFP	Up to 3-fold improved Transducing units ^a	Plasmid backbone antibiotic resistance marker packaged in up to 2.9% ^a or 3% ^b of viral particles potential for genome integration in transduced cells
Sleeping Beauty Transposon	SB puroR reporter plasmid	SB puroR reporter MC	2-fold increased transposition rate into established cell lines ^c	Plasmid backbone antibiotic resistance potential for genome integration in T-cells
	SB CD19-CAR	MC: SB CD19 CAR	4-fold increased yield of CD 19-CAR T-cells; reduced transfection associated toxicity ^d	
	SB	MC: SB (218 bp MC Backbone)	6-7-fold increased transposition into Human Hematopoietic Stem cells; reduced transfection associated toxicity ^e	

^a Schnödt et al., (2016) Mol Ther - Nucleic Acids 5 e355. This citation, along with Chadeuf et al., (2005) Molecular Therapy 12:744-53 and Gray, 2017. WO2017066579 also reports that AAV helper plasmid antibiotic resistance markers are packaged into viral particles, demonstrating need to remove antibiotic markers from AAV helper plasmids as well as the AAV vector. Interestingly, no improvement in viral titer compared to plasmid is observed with linear Doggybone™ DNA single-stranded (ss) AAV vector retrofits (Karbowniczek et al., 2017. Cell and Gene Therapy Insights 731-8). This suggests that short backbone minicircle vector improvement compared to >1.5 kb pUC-antibiotic marker plasmid backbone vectors require a short backbone circular vector rather than a linear vector such as Doggybone™ vectors.

^b Gray, *Supra*, 2017 describes adding a large spacer region to move selection marker away from ITRs. Resultant vector, pAAV CMV GFP is 11 kb compared to 5-6 kb for standard plasmid AAV vector so will have reduced transducing units compared to parental

^c Sharma et al., (2013) Molecular Therapy Nucleic Acids 2:e74

^d Nucleofection transfection: Monjezi et al., (2017) Leukemia 31: 186-94

^e Nucleofection transfection: Holstein et al., (2018) Molecular Therapy epub Jan 17, 2018

[0012] Indeed 2-fold improved sleeping beauty transposition into human cells was reported with the 1.1 kb bacterial backbone pFAR4 SB transposon vector/SB100x transposase vector combination compared to a 2.8 kb bacterial backbone pT2 plasmid SB transposon vector/SB 100x transposase vector combination (Pastor, M, Johnen S, Harmening N, Quiviger M, Pailloux J, Kropp, M, Walter P, Ivics Z, Izsvak Z, Thumann G, Scherman D. 2018 Molecular Therapy 11: 57-67).

[0013] However, viral vectors such as AAV, Lentiviral and Retroviral vectors, and transposon vectors contained structured DNA sequences at their termini. For example, Sleeping Beauty transposon vectors contain flanking IR/DR sequences, AAV vectors contain flanking ITRs, and Lentiviral and Retroviral vectors contain flanking LTRs.

[0014] The close proximity of the pUC origin to a structured DNA sequence results in aberrant replication termination, resulting in replication intermediates which unacceptably reduce plasmid quality (Levy J. 2004. US Patent 6709844). Levy teaches that replication intermediates form when any high copy replication origin is < 1kb from a structured DNA sequence such as an enhancer, LTR or IRES, but not when the high copy replication origin is >1.5 kb away. Since the pUC origin itself is 1 kb, there is no configuration to make a <1.1 kb bacterial region AAV, Lentiviral, Retroviral or transposon vector containing the pUC origin which is not predicted to produce replication intermediates.

[0015] Lu J, Williams JA, Luke J, Zhang F, Chu K, and Kay MA. 2017. Human Gene Therapy 28: 125-34 disclose antibiotic free Mini-Intronic Plasmid (MIP) AAV vectors and suggest that MIP intron AAV vectors could have the vector backbone removed to create a short backbone AAV vector. Attempts to create a minicircle like 6 or 10 bp spacer region in Mini-Intronic Plasmid AAV vectors were toxic (see Table 7, footnote e) presumably due to creation of a long palindrome by such close juxtaposition of

the AAV ITRs. While MIP vectors with longer spacer regions < 1kb can be made, a drawback of the MIP intron strategy is that it requires cloning of a replication and selection encoding intron into the eukaryotic region, which is not possible or desired with many vectors.

[0016] A drawback of the minicircle strategy to create short bacterial region AAV, Lentiviral, Retroviral or transposon vectors, is that methods to manufacture minicircle vectors are expensive and not easily scalable. For minicircle vectors, *E. coli*-based manufacturing systems have been developed in which, after plasmid production, the bacterial region and the eukaryotic region are separated and circularized into a minicircle (eukaryotic region) and a bacterial region circle via the action of phage recombinases on recognition sequences in the plasmid. In some methods, a restriction enzyme is then utilized to digest the bacterial region circle at a unique site to eliminate this difficult to remove contaminant. These production procedures are very inefficient. For example, optimal manufacture of minicircle vectors yields only 5 mg of minicircle per liter culture (Kay MA, He CY, Chen ZY. 2010. Nat Biotechnol 28:1287-1289).

[0017] Methods for high yield manufacture of pFAR vectors have not been reported; this system utilizes a plasmid borne suppressor tRNA gene to complement a TAG amber nonsense mutation of the thyA gene to complement thymidine auxotrophy and allow cell growth on minimal media (Marie *et al.*, *Supra*, 2010).

[0018] A solution is needed to develop AAV, Lentiviral, Retroviral or transposon vector that contain short spacer regions preferably less than 1000 bp that can be efficiently manufactured without replication intermediates or poor production. These vectors should not encode a protein-based selection marker and should be minimized to eliminate all non-essential sequences.

SUMMARY OF THE INVENTION

[0019] In a first aspect, the present invention provides a method for improving the replication of a covalently closed circular plasmid comprising the following steps:

1. a. providing a covalently closed circular plasmid comprising:
 1. i. a Pol I-dependent origin of replication, and
 2. ii. an insert comprising a structured DNA sequence selected from the group consisting of inverted repeat sequence, direct repeat sequence, homopolymeric repeat sequence, or eukaryotic origin of replication, wherein the structured DNA sequence is located at a distance of less than 1000 bp from the Pol I-dependent origin of replication in the direction of replication; and
2. b. modifying the covalently closed circular plasmid of a. such that the Pol I-dependent origin of replication is replaced with an R6K gamma replication origin whereby the resultant R6K gamma replication origin covalently closed circular plasmid has improved replication selected from the group consisting of reduced production of replication intermediates and increased plasmid copy number relative to the covalently closed circular plasmid of a, wherein the R6K gamma replication origin does not require Pol I, and wherein the structured DNA sequence is located at a distance of less than 1000 bp from the R6K gamma replication origin in the direction of replication.

[0020] In a second aspect, the present invention provides a method for improving the replication of a covalently closed circular plasmid comprising the following steps:

1. a. providing a covalently closed circular plasmid comprising:
 1. i. a bacterial replication-selection region comprising a Pol I-dependent origin of replication and an antibiotic selectable marker, and
 2. ii. an insert comprising a structured DNA sequence selected from the group consisting of inverted repeat sequence, direct repeat sequence, homopolymeric repeat sequence, or eukaryotic origin of replication, wherein the structured DNA sequence is located at a distance of less than 1000 bp from the Pol I-dependent origin of replication in the direction of replication; and
2. b. modifying the covalently closed circular plasmid of a. such that the antibiotic selectable marker is replaced with an RNA selectable marker and the Pol I-dependent origin of replication is replaced with an R6K gamma replication origin, whereby the resultant R6K gamma replication origin covalently closed circular plasmid has improved replication selected from the group consisting of reduced production of replication intermediates and increased plasmid copy number relative to the covalently closed circular plasmid of a, wherein the R6K gamma replication origin does not require Pol I, and wherein the structured DNA sequence is located at a distance of less than 1000 bp from the R6K gamma replication origin in the direction of replication.

[0021] In an embodiment, the Pol I-dependent origin of replication may be selected from the group consisting of pUC origin, pMB1 origin, and ColE1 origin.

[0022] In an embodiment, said R6K gamma replication origin may be an R6K gamma replication origin with at least 95% sequence identity to a sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 18.

[0023] In an embodiment, said structured DNA sequence may be:

1. a. a homopolymeric repeat, and wherein the homopolymeric repeat is a polyA repeat;
2. b. a eukaryotic origin of replication, and wherein the eukaryotic origin of replication is a SV40 origin of replication;
3. c. a direct repeat sequence, and wherein the direct repeat sequence is selected from the group consisting of viral LTR, a Lentiviral LTR, or a Retro viral LTR; or
4. d. an inverted repeat sequence, and wherein the inverted repeat sequence is selected from the group consisting of a transposon IR/DR repeat, a Sleeping Beauty transposon IR/DR repeat, an AAV ITR, a transposon ITR, or a PiggyBac transposon ITR.

[0024] In an embodiment, said RNA selectable marker may be an RNA-IN regulating RNA-OUT functional variant with at least 95% sequence identity to a sequence selected from the group consisting of SEQ ID NO: 5 and SEQ ID NO: 7.

[0025] In an embodiment, said RNA selectable marker may be an RNA-OUT RNA selectable marker that encodes an RNA-IN regulating RNA-OUT RNA with at least 95% sequence identity to SEQ ID NO: 6.

[0026] In an embodiment, said bacterial replication-selection region comprising a Pol I-dependent origin of replication and an antibiotic selectable marker may be replaced with a Pol III-dependent R6K origin-RNA-OUT RNA selectable marker bacterial replication-selection region with at least 95% sequence identity to a sequence selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, and SEQ ID NO: 17.

[0027] The Pol III-dependent replication origin plasmids provided by the invention have surprisingly improved manufacturing quality and yield than the parent pMB1, ColE1 or pBR322 derived replication origin expression plasmid vector.

[0028] Further objects and advantages of the invention will become apparent from a consideration of the drawings and ensuing description.

BRIEF DESCRIPTION OF THE FIGURES

[0029]

FIGS. 1A-1F depict the R6K origin (FIGS. 1A, 1E, and 1F), RNA-OUT selectable marker (FIG. 1B), and 14 and 3 CpG R6K-RNA-OUT bacterial backbones (FIGS. 1C and 1D);

FIGS. 2A-2B depict a Pol I-dependent pUC origin Sleeping Beauty transposon vector (FIG. 2A) and a Pol III-dependent R6K origin Sleeping Beauty transposon vector (FIG. 2B);

FIGS. 3A-3C depict Pol I-dependent pUC origin AAV vectors (FIGS. 3A and 3B) and a Pol III-dependent R6K origin AAV vectors (FIG. 3C); and

FIGS. 4A-4F depict Pol I-dependent pUC origin A60 polyA repeat encoding mRNA vectors (FIGS. 4A-4B), a Pol III-dependent R6K origin A60 polyA repeat encoding mRNA vector (FIG. 4C), Pol I-dependent pUC origin A99 polyA repeat encoding mRNA vectors (FIGS. 4D-4E), and a Pol III-dependent R6K origin A99 polyA repeat encoding mRNA vector (FIG. 4F).

Table 1: pVAX1 mammalian expression vector spacer region (SR) derivatives expression level

Table 2: NTC8685 mammalian expression vector spacer region (SR) derivatives expression level

Table 3: Minicircle applications with various viral and non-viral vector platforms

Table 4: pNTC multiple cloning site flanked R6K Origin-RNA-OUT selection marker vectors

Table 5: SV40 origin Lentiviral vectors: pUC versus R6K origin shake flask production yields/quality

Table 6: Sleeping Beauty Transposon vectors: pUC versus R6K origin shake flask production yields/quality

Table 7: AAV vectors: pUC versus R6K origin shake flask production yields/quality

Table 8: mRNA vectors: pUC versus R6K origin DH5 α HyperGRO fermentation yields/quality

Table 9: AAV helper vectors: pUC versus R6K origin plasmid production yields/quality

SEQ ID NO:1: R6K gamma origin

SEQ ID NO:2: 1 CpG R6K gamma origin

SEQ ID NO:3: CpG free R6K gamma origin

SEQ ID NO:4: Extended R6K gamma origin

SEQ ID NO:5: RNA-OUT Selectable Marker

SEQ ID NO:6: RNA-OUT antisense repressor RNA

SEQ ID NO:7: 2 CpG RNA-OUT Selectable Marker

SEQ ID NO:8: R6K gamma origin-RNA-OUT bacterial region flanked by NheI and KpnI restriction sites

SEQ ID NO:9: 1 CpG R6K gamma origin-2 CpG RNA-OUT bacterial region flanked by NheI and KpnI restriction sites

SEQ ID NO:10: pNTC-NP1 polylinker trpA R6K-RNA-OUT polylinker cloning cassette: EcoRI/HindIII

SEQ ID NO:11: pNTC-NP2 polylinker trpA R6K-RNA-OUT polylinker cloning cassette: EcoRI/HindIII

SEQ ID NO:12: pNTC-NP3 polylinker trpA R6K-RNA-OUT polylinker cloning cassette: EcoRI/HindIII

SEQ ID NO:13: pNTC-NP4 polylinker trpA R6K-RNA-OUT polylinker cloning cassette: EcoRI/HindIII

SEQ ID NO:14: pNTC-NP5 polylinker trpA R6K-RNA-OUT polylinker cloning cassette: KasI/HindIII

SEQ ID NO:15: pNTC-NP6 polylinker trpA R6K-RNA-OUT polylinker cloning cassette: EcoRI/SacI

SEQ ID NO:16: pNTC-NP7 polylinker trpA R6K-RNA-OUT polylinker cloning cassette: BssHII-BssHII

SEQ ID NO:17: pNTC-3xCpG NP1 polylinker R6K-RNA-OUT polylinker cloning cassette: HindIII-EcoRI

SEQ ID NO:18: R6K gamma origin (7 iteron)

SEQ ID NO:19: R6K gamma origin 22 bp iteron repeat

SEQ ID NO:20: R6K gamma origin 22 bp iteron repeat

SEQ ID NO:21: R6K gamma origin 22 bp iteron repeat

SEQ ID NO:22: R6K gamma origin 22 bp iteron repeat

SEQ ID NO:23: R6K gamma origin 22 bp iteron repeat

Definition of Terms

[0030]

AAV vector: Adeno-associated virus vector, an episomal viral vector. Includes self-complementary (sc) Adeno-associated virus vectors (scAAV) and single-stranded

(ss) Adeno-associated virus vectors (ssAAV)

AF: Antibiotic-free

amp: Ampicillin

ampR: Ampicillin Resistance gene

Antibiotic selectable marker: A gene that confers resistance to an antibiotic, e.g. ampicillin resistance gene, kanamycin resistance gene, chloramphenicol resistance gene, tetracycline resistance gene

Approximately: As used herein, the term "approximately" or "about," as applied to one or more values of interest, refers to a value that is the same or similar to a stated reference value

Bacterial region: Region of a plasmid vector required for propagation and selection in the bacterial host

bp: basepairs

ccc: Covalently Closed Circular

cl: Lambda repressor

clTs857: Lambda repressor further incorporating a C to T (Ala to Thr) mutation that confers temperature sensitivity. clTs857 is a functional repressor at 28-30°C but is mostly inactive at 37-42°C. Also called cl857

Cat^R: Chloramphenicol resistance gene

cmv: Cytomegalovirus

dem methylation: *E. coli* methyltransferase that methylated the sequences CC(AT)GG at the C5 position of the second cytosine

DNA replicon: A genetic element that can replicate under its own control; examples include plasmids, cosmids, bacterial artificial chromosomes (BACs), bacteriophages, viral vectors and hybrids thereof

E. coli: *Escherichia coli*, a gram negative bacteria

EGFP: Enhanced green fluorescent protein

EP: Electroporation

Eukaryotic expression vector: A vector for expression of mRNA, protein antigens, protein therapeutics, shRNA, RNA or microRNA genes in a target eukaryotic organism using RNA Polymerase I, II or III promoters

Eukaryotic region: The region of a plasmid that encodes eukaryotic sequences and/or sequences required for plasmid function in the target organism. This includes the region of a plasmid vector required for expression of one or more transgenes in the target organism including RNA Pol II enhancers, promoters, transgenes and polyA sequences. This also includes the region of a plasmid vector required for expression of one or more transgenes in the target organism using RNA Pol I or RNA Pol III promoters, RNA Pol I or RNA Pol III expressed transgenes or RNAs. The eukaryotic region may optionally include other functional sequences, such as eukaryotic transcriptional terminators, supercoiling-induced DNA duplex destabilized (SIDD) structures, S/MARs, boundary elements, etc. In a Lentiviral or Retroviral vector, the eukaryotic region contains flanking direct repeat LTRs, in a AAV vector the eukaryotic region contains flanking inverted terminal repeats, while in a Transposon vector the eukaryotic region contains flanking transposon inverted terminal repeats or IR/DR termini (e.g. Sleeping Beauty). In genome integration vectors, the eukaryotic region may encode homology arms to direct targeted integration

Exon: A nucleotide sequence encoded by a gene that is transcribed and present within a mature mRNA product after RNA splicing to remove introns has been completed

Expression vector: A vector for expression of mRNA, protein antigens, protein therapeutics, shRNA, RNA or microRNA genes in a target organism.

g: Gram, kg for kilogram

gene of interest: gene to be expressed in the target organism. Includes mRNA genes that encode protein or peptide antigens, protein or peptide therapeutics, and mRNA, shRNA, RNA or microRNA that encode RNA therapeutics, and mRNA, shRNA, RNA or microRNA that encode RNA vaccines, etc.

Hr(s): Hour(s)

ID: Intradermal

IM: Intramuscular

immune response: Antigen reactive cellular (e.g. antigen reactive T cells) or antibody (e.g. antigen reactive IgG) responses

Intron: A nucleotide sequence encoded by a gene that is transcribed and subsequently removed from a mature mRNA product by RNA splicing

IR/DR: Inverted Repeats which are each Directly Repeated twice. For example, Sleeping Beauty transposon IR/DR repeats

Iteron: Directly repeated DNA sequences in a origin of replication that are required for replication initiation. R6K origin iteron repeats are 22 bp

ITR: Inverted Terminal Repeat

kan: Kanamycin

kanR: Kanamycin Resistance gene

Kd: Kilodalton

kozak sequence: Optimized consensus DNA sequence gccRccATG (R = G or A) immediately upstream of an ATG start codon that ensures efficient translation initiation. A Sall site (GTCGAC) immediately upstream of the ATG start codon (GTCGACATG) is an effective kozak sequence

Lentiviral vector: Integrative viral vector that can infect dividing and non-dividing cells. Also call Lentiviral transfer plasmid. Plasmid encodes Lentiviral LTR flanked expression unit. Transfer plasmid is transfected into production cells along with Lentiviral envelope and packaging plasmids required to make viral particles

Lentiviral envelope vector: Plasmid encoding envelope glycoprotein

Lentiviral packaging vector: One or two plasmids that express gag, pol and Rev functions required to package the Lentiviral transfer vector

minicircle: Covalently closed circular plasmid derivatives in which the bacterial region has been removed from the parent plasmid by *in vivo* or *in vitro* site-specific recombination or *in vitro* restriction digestion/ligation. Minicircle vectors are replication incompetent in bacterial cells **mRNA:** Messenger RNA

mSEAP: Murine secreted alkaline phosphatase

NA: Not Applicable

Nanoplasmid™ vector: Vector combining an RNA selectable marker with a R6K, ColE2 or ColE2 related replication origin. For example, NTC9385C, NTC9685C, NTC9385R, NTC9685R vectors and modifications described in Williams, *Supra*, 2014

NTC8385: NTC8385, NTC8485 and NTC8685 plasmids are antibiotic-free pUC origin vectors that contain a short RNA (RNA-OUT) selectable marker instead of an antibiotic resistance marker such as kanR. The creation and application of these RNA-OUT based antibiotic-free vectors are described in Williams, JA 2008 World Patent Application WO2008153733

NTC8485: NTC8485 is an antibiotic-free pUC origin vector that contains a short RNA (RNA-OUT) selectable marker instead of an antibiotic resistance marker such as kanR. The creation and application of NTC8485 is described in Williams, JA 2010 US Patent Application 20100184158 **NTC8685:** NTC8685 is an antibiotic-free pUC origin vector that contains a short RNA (RNA-OUT) selectable marker instead of an antibiotic resistance marker such as kanR. The creation and application of NTC8685 is described in Williams, *Supra*, 2010

NTC9385R: The NTC9385R Nanoplasmid™ vector described in Williams, *Supra*, 2014 has a spacer region encoded NheI- trpA terminator-R6K origin RNA-OUT -KpnI bacterial region (SEQ ID NO:8) linked through the flanking NheI and KpnI sites to the eukaryotic region.

OD₆₀₀: optical density at 600 nm

PAS: Primosomal assembly site. Priming of DNA synthesis on a single stranded DNA ssi site. ØX174 type PAS: DNA hairpin sequence that binds *priA*, which, in turn, recruits the remaining proteins to form the preprimosome [*priB*, *dnaT*, recruits *dnaB* (delivered by *dnaC*)], which then also recruits primase (*dnaG*), which then, finally, makes a short RNA substrate for DNA polymerase I. ABC type PAS: DNA hairpin binds *dnaA*, recruits *dnaB* (delivered by *dnaC*) which then also recruits primase (*dnaG*), which then, finally, makes a short RNA substrate for DNA polymerase I. For example, the R6K plasmid CpG free *ssiA* primosomal assembly site or alternative ØX174 type or ABC type primosomal assembly sites

PAS-BH: Primosomal assembly site on the heavy (leading) strand

PAS-BH region: pBR322 origin region between ROP and PAS-BL (approximately pBR322 2067-2351)

PAS-BL: Primosomal assembly site on the light (lagging) strand

PBS: Phosphate buffered Saline

PCR: Polymerase Chain Reaction

pDNA: Plasmid DNA

PiggyBac Transposon: PB transposon. A transposon system that integrates an ITR flanked PB transposon into the genome by a simple cut and paste mechanism mediated by PB transposase. The transposon vector typically contains a promoter-transgene-polyA expression cassette between the PB ITRs which is excised and integrated into the genome

pINT pR pL vector: The pINT pR pL att_{HK022} integration expression vector is described in Luke et al., 2011 Mol Biotechnol 47:43. The target gene to be expressed is cloned downstream of the pL promoter. The vector encodes the temperature inducible cI857 repressor, allowing heat inducible target gene expression

P_L promoter: Lambda promoter left. P_L is a strong promoter that is repressed by the cI repressor binding to OL1, OL2 and OL3 repressor binding sites. The temperature sensitive cI857 repressor allows control of gene expression by heat induction since at 30°C the cI857 repressor is functional and it represses gene expression, but at 37-42 °C the repressor is inactivated so expression of the gene ensues

P_L (OL1 G to T) promoter: Lambda promoter left. P_L is a strong promoter that is repressed by the cI repressor binding to OL1, OL2 and OL3 repressor binding sites. The temperature sensitive cI857 repressor allows control of gene expression by heat induction since at 30°C the cI857 repressor is functional and it represses gene expression, but at 37-42 °C the repressor is inactivated so expression of the gene ensues. The cI repressor binding to OL1 is reduced by the OL1 G to T mutation resulting in increased promoter activity at 30°C and 37-42 °C as described in Williams, *Supra*, 2014.

Plasmid: An extra chromosomal DNA molecule separate from the chromosomal DNA which is capable of replicating independently from the chromosomal DNA

Plasmid copy number: the number of copies of a plasmid per cell. Increases in plasmid copy number increase plasmid production yield

Pol: Polymerase

Pol I: *Escherichia coli* DNA Polymerase I

Pol I dependent origin of replication: A replication origin that requires Pol I, for example the pMB1, ColE1 or pBR322 or derivatives such as the high copy pUC origin. For these origins the RNAlI primer forms an RNA: DNA R-loop that is cleaved by RNase H to create a primer for DNA pol I directed DNA synthesis. DNA synthesis then converts to DNA pol III. Numerous additional Pol I dependent replication origins are known in the art, many of which are summarized in del Solar et al., 1998 Microbiol. Mol. Biol. Rev 62:434-464

Pol III: *Escherichia coli* DNA Polymerase III

Pol III dependent origin of replication: A replication origin that doesn't require Pol I, for example the rep protein dependent R6K gamma replication origin. Numerous additional Pol III dependent replication origins are known in the art, many of which are summarized in del Solar et al., *Supra*, 1998

polyA: Polyadenylation signal or site. Polyadenylation is the addition of a poly(A) tail to an RNA molecule. The polyadenylation signal contains the sequence motif recognized by the RNA cleavage complex. Most human polyadenylation signals contain an AAUAAA motif and conserved sequences 5' and 3' to it. Commonly utilized polyA signals are derived from the rabbit β globin, bovine growth hormone, SV40 early, or SV40 late polyA signals

pUC origin: pBR322-derived replication origin, with G to A transition that increases copy number at elevated temperature and deletion of the ROP negative regulator

pUC free: Plasmid that does not contain the pUC origin. Non-replicative fragments of the pUC origin may be included, for example the RNAI selectable marker

pUC plasmid: Plasmid containing the pUC origin

R6K plasmid: NTC9385R, NTC9685R, NTC9385R2-01, NTC9385R2-02, NTC9385R2a-01, NTC9385R2a-02, NTC9385R2b-01, NTC9385R2b-02, NTC9385Ra-01, NTC9385Ra-02, NTC9385RaF, and NTC9385RbF vectors as well as modifications and alternative vectors containing a R6K replication origin that were described in Williams, *Supra*, 2014. Alternative R6K vectors known in the art including, but not limited to, pCOR vectors (Gencell), pCpGfree vectors (Invivogen), and CpG free University of Oxford vectors including pGM169

R6K replication origin: a region which is specifically recognized by the R6K Rep protein to initiate DNA replication. Includes but not limited to R6K gamma replication origin sequence disclosed as SEQ ID NO: 1, SEQ ID NO:2 SEQ ID NO:4, and SEQ ID NO: 18. Also includes CpG free versions (e.g. SEQ ID NO:3) as described in Drocourt et al., United States Patent 7244609

R6K replication origin-RNA-OUT bacterial region: Contains a R6K replication origin for propagation and the RNA-OUT selectable marker (e.g. SEQ ID NO:8; SEQ ID NO:9; SEQ ID NO:10; SEQ ID NO:11; SEQ ID NO:12; SEQ ID NO:13; SEQ ID NO: 14; SEQ ID NO:15; SEQ ID NO:16; SEQ ID NO:17)

Rep: Replication

Replication intermediates: Linear DNA fragments resulting from premature termination of plasmid replication

Rep protein dependent plasmid: A plasmid in which replication is dependent on a replication (Rep) protein provided in Trans. For example, R6K replication origin, ColE2-P9 replication origin and ColE2 related replication origin plasmids in which the Rep protein is expressed from the host strain genome. Numerous additional Rep protein dependent plasmids are known in the art, many of which are summarized in del Solar *et al.*, *Supra*, 1998

Retroviral vector: Integrative viral vector that can infect dividing cells. Also call transfer plasmid. Plasmid encodes Retroviral LTR flanked expression unit. Transfer plasmid is transfected into production cells along with envelope and packaging plasmids required to make viral particles

Retroviral envelope vector: Plasmid encoding envelope glycoprotein

Retroviral packaging vector: Plasmid that encodes Retroviral gag, pol genes required to package the Retroviral transfer vector

RNA-IN: Insertion sequence 10 (IS10) encoded RNA-IN, an RNA complementary and antisense to a portion of RNA RNA-OUT. When RNA-IN is cloned in the untranslated leader of a mRNA, annealing of RNA-IN to RNA-OUT reduces translation of the gene encoded downstream of RNA-IN

RNA-IN regulated selectable marker: A genomically expressed RNA-IN regulated selectable marker. In the presence of plasmid borne RNA-OUT antisense repressor RNA (SEQ ID NO:6), expression of a protein encoded downstream of RNA-IN is repressed. An RNA-IN regulated selectable marker is configured such that RNA-IN regulates either 1) a protein that is lethal or toxic to said cell per se or by generating a toxic substance (e.g. SacB), or 2) a repressor protein that is lethal or toxic to said bacterial cell by repressing the transcription of a gene that is essential for growth of said cell (e.g. murA essential gene regulated by RNA-IN tetR repressor gene). For example, genomically expressed RNA-IN-SacB cell lines for RNA-OUT plasmid selection/propagation are described in Williams, *Supra*, 2008. Alternative selection markers described in the art may be substituted for SacB

RNA-OUT: Insertion sequence 10 (IS10) encoded RNA-OUT, an antisense RNA that hybridizes to, and reduces translation of, the transposon gene expressed downstream of RNA-IN. The sequence of the RNA-OUT RNA (SEQ ID NO:6) and complementary RNA-IN SacB genomically expressed RNA-IN-SacB cell lines can be modified to incorporate alternative functional RNA-IN/RNA-OUT binding pairs such as those described in Mutalik *et al.*, 2012 Nat Chem Biol 8:447, including, but not limited to, the RNA-OUT A08/RNA-IN S49 pair, the RNA-OUT A08/RNA-IN S08 pair, and CpG free modifications of RNA-OUT A08 that modify the CG in the RNA-OUT 5' TTGGC sequence to a non-CpG sequence. An example of a CpG free RNA-OUT selection marker, in which the two CpG motifs in the RNA-OUT RNA (one of which is present in the RNA-IN complementary region) are removed, was described in Williams 2015. Replicative minicircle vectors with improved expression. US Patent Application US 2015/0275221. A multitude of alternative substitutions to remove the two CpG motifs (mutating each CpG to either CpA, CpC, CpT, ApG, GpG, or TpG) may be utilized to make a CpG free RNA-OUT

RNA-OUT Selectable marker: An RNA-OUT selectable marker DNA fragment including *E. coli* transcription promoter and terminator sequences flanking an RNA-OUT RNA. An RNA-OUT selectable marker, utilizing the RNA-OUT promoter and terminator sequences, that is flanked by DraIII and KpnI restriction enzyme sites, and designer genomically expressed RNA-IN-SacB cell lines for RNA-OUT plasmid propagation, are described in Williams, *Supra*, 2008. The RNA-OUT promoter and terminator sequences in SEQ ID NO: 5 that flank the RNA-OUT RNA (SEQ ID NO:6; FIG. 1B) may be replaced with heterologous promoter and terminator sequences. For example, the RNA-OUT promoter may be substituted with a CpG free promoter known in the art, for example the I-EC2K promoter or the P5/6 5/6 or P5/6 6/6 promoters described in Williams, *Supra*, 2008. A 2 CpG RNA-OUT selectable marker in which the two CpG motifs in the RNA-OUT promoter are removed is given as SEQ ID NO: 7. An example of a CpG free RNA-OUT transcription unit, in which the two CpG motifs in the RNA-OUT RNA (one of which is present in the RNA-IN complementary region) and the two CpG motifs in the RNA-OUT promoter are removed was described in Williams, *Supra*, 2015. Vectors incorporating CpG free RNA-OUT selectable marker may be selected for sucrose resistance using the RNA-IN-SacB cell lines for RNA-OUT plasmid propagation described in Williams, *Supra*, 2008. Alternatively, the RNA-IN sequence in these cell lines can be modified to incorporate the 1 bp change needed to perfectly match the CpG free RNA-OUT region complementary to RNA-IN.

RNA polymerase II promoter: Promoter that recruits RNA Polymerase II to synthesize mRNAs, most small nuclear RNAs and microRNAs. For example, constitutive promoters such as the human or murine CMV promoter, elongation factor 1 (EF1) promoter, the chicken β -actin promoter, the β -actin promoter from other species, the elongation factor-1 α (EF1 α) promoter, the phosphoglycerokinase (PGK) promoter, the Rous sarcoma virus (RSV) promoter, the human serum albumin (SA) promoter, the spleen focus-forming virus (SFFV) promoter, the α -1 antitrypsin (AAT) promoter, the thyroxine binding globulin (TBG) promoter, the cytochrome P450 2E1 (CYP2E1) promoter, etc. The vectors may also utilize combination promoters such as the chicken β -actin/CMV enhancer (CAG) promoter, the human or murine CMV-derived enhancer elements combined with the elongation factor 1 α (EF1 α) promoters, CpG free versions of the human or murine CMV-derived enhancer elements combined with the elongation factor 1 α (EF1 α) promoters, the albumin promoter combined with an α -fetoprotein MERII enhancer, etc., or the diversity of tissue specific or inducible promoters known in the art such as the muscle specific promoters muscle creatine kinase (MCK), and C5-12 or the liver-specific promoter apolipoprotein A-I (ApoA1), etc.

RNA polymerase III promoter: Promoter that recruits RNA Polymerase III to synthesize tRNAs, 5S ribosomal RNA, and other small RNAs. For example, Class I promoters such as the 5s rRNA promoter, Class II promoter such as tRNA promoters, Class III promoters such as the U6 small nuclear RNA promoter or the H1 nuclear RNase P promoter, etc.

RNA selectable marker: An RNA selectable marker is a plasmid borne expressed non-translated RNA that regulates a chromosomally expressed target gene to afford selection. This may be a plasmid borne nonsense suppressing tRNA that regulates a nonsense suppressible selectable chromosomal target as described by Crouzet J and Soubrier F 2005 US Patent 6,977,174. This may also be a plasmid borne antisense repressor RNA, a non limiting list includes RNA-OUT that represses RNA-IN regulated targets (Williams, *Supra*, 2008), pMB1 plasmid origin encoded RNAI that represses RNAII regulated targets (Grabherr R, Pfaffenzeller I. 2006 US patent application US20060063232; Cranenburgh RM. 2009; US Patent 7,611,883), IncB plasmid pMU720 origin encoded RNAI that represses RNA II regulated targets (Wilson IW, Siemerling KR, Praszkiel J, Pittard AJ. 1997. J Bacterial 179:742-53), ParB locus Sok of plasmid R1 that represses Hok regulated targets, Flm locus FlmB of F plasmid that represses flmA regulated targets (Morsey MA, 1999 US patent US5922583). An RNA selectable marker may be another natural antisense repressor RNAs known in the art such as those described in Wagner EGH, Altuvia S, Romby P. 2002. Adv Genet 46:361-98 and Franch T, and Gerdes K. 2000. Current Opin Microbiol 3:159-64. An RNA selectable marker may also be an engineered repressor RNAs such as synthetic small RNAs expressed SgrS, MicC or MicF scaffolds as

described in Na D, Yoo SM, Chung H, Park H, Park JH, Lee SY. 2013. Nat Biotechnol 31:170-4. An RNA selectable marker may also be an engineered repressor RNA as part of a selectable marker that represses a target RNA fused to a target gene to be regulated such as SacB as described in Williams, *Supra*, 2015

ROP: Repressor of primer

RSM: RNA selectable marker

SacB: Structural gene encoding *Bacillus subtilis* levansucrase. Expression of SacB in gram negative bacteria is toxic in the presence of sucrose

SD: Standard deviation

SEAP: Secreted alkaline phosphatase

Selectable marker: A selectable marker, for example a kanamycin resistance gene or an RNA selectable marker

Selection marker: A selectable marker, for example a kanamycin resistance gene or an RNA selectable marker

SIDD: supercoiling-induced DNA duplex destabilized (SIDD) structures. These sites, when incorporated into a vector, may alter the susceptibility of other sequences within the vector to be destabilized. This can alter function. For example, addition of a SIDD site to an expression vector may reduce the helical destabilization of a promoter. This may increase or decrease promoter activity, depending on the promoter since some promoters have increased expression with promoter helical destabilization, while others will have reduced expression with promoter helical destabilization

shRNA: Short hairpin RNA

S/MAR: Scaffold/matrix attached region. Eukaryotic sequences that mediate DNA attachment to the nuclear matrix

Sleeping Beauty Transposon: SB transposon. A transposon system that integrates an IR/DR flanked SB transposon into the genome by a simple cut and paste mechanism mediated by SB transposase. The transposon vector typically contains a promoter-transgene-polyA expression cassette between the IR/DRs which is excised and integrated into the genome

Spacer region: As used herein, spacer region is the region linking the 5' and 3' ends of the eukaryotic region sequences. The eukaryotic region 5' and 3' ends are typically separated by the bacterial replication origin and bacterial selectable marker in plasmid vectors (bacterial region) so many spacer regions consist of the bacterial region. In Pol III dependent origin of replication vectors of the invention, this spacer region preferably is less than 1000 bp

SR: Spacer region.

ssi: Single stranded initiation sequences

Structured DNA sequence: As used herein, a DNA sequence that is capable of forming replication inhibiting secondary structures (Mirkin and Mirkin, 2007. Microbiology and Molecular Biology Reviews 71: 13-35). This includes but is not limited to inverted repeats, palindromes, direct repeats, IR/DRs, homopolymeric repeats or repeat containing eukaryotic promoter enhancers, or repeat containing eukaryotic origin of replications.

SV40 origin: Simian Virus 40 genomic DNA that contains the origin of replication

SV40 enhancer: Simian Virus 40 genomic DNA that contains the 72 bp and optionally the 21 bp enhancer repeats

target antigen: Immunogenic protein or peptide epitope, or combination of proteins and epitopes, against which an immune response can be mounted. Target antigens may be derived from a pathogen for infectious disease or allergy applications or derived from a host organism for applications such as cancer, allergy, or autoimmune diseases. Target antigens are well defined in the art. Some examples are described in Williams, *Supra*, 2008

TE buffer: A solution containing approximately 10mM Tris pH 8 and 1 mM EDTA

TetR: Tetracycline resistance gene

ToI2 Transposon: A transposon system that integrates an ITR flanked ToI2 transposon into the genome by a simple cut and paste mechanism mediated by ToI2 transposase. The transposon vector typically contains a promoter-transgene-polyA expression cassette between the ToI2 ITRs which is excised and integrated into the genome

Transcription terminator: Bacterial: A DNA sequence that marks the end of a gene or operon for transcription. This may be an intrinsic transcription terminator or a Rho-dependent transcriptional terminator. For an intrinsic terminator, such as the trpA terminator, a hairpin structure forms within the transcript that disrupts the mRNA-DNA-RNA polymerase ternary complex. Alternatively, Rho-dependent transcriptional terminators require Rho factor, an RNA helicase protein complex, to disrupt the nascent mRNA-DNA-RNA polymerase ternary complex. Eukaryotic: PolyA signals are not 'terminators', instead internal cleavage at PolyA sites leaves an uncapped 5' end on the 3'UTR RNA for nuclease digestion. Nuclease catches up to RNA Pol II and causes termination. Termination can be promoted within a short region of the poly A site by introduction of RNA Pol II pause sites (eukaryotic transcription terminator). Pausing of RNA Pol II allows the nuclease introduced into the 3' UTR mRNA after PolyA cleavage to catch up to RNA Pol II at the pause site. A nonlimiting list of eukaryotic transcription terminators known in the art include the C2x4 and the gastrin terminator. Eukaryotic transcription terminators may elevate mRNA levels by enhancing proper 3'-end processing of mRNA

transfection: Method to deliver nucleic acids into cells [e.g. poly(lactide-co-glycolide) (PLGA), ISCOMs, liposomes, niosomes, virosomes, block copolymers, Pluronic block copolymers, chitosan, and other biodegradable polymers, microparticles, microspheres, calcium phosphate nanoparticles, nanoparticles, nanocapsules, nanospheres, poloxamine nanospheres, electroporation, nucleofection, piezoelectric permeabilization, sonoporation, iontophoresis, ultrasound, SQZ high speed cell deformation mediated membrane disruption, corona plasma, plasma facilitated delivery, tissue tolerable plasma, laser microporation, shock wave energy, magnetic fields, contactless magnetopermeabilization, gene gun, microneedles, microdermabrasion, hydrodynamic delivery, high pressure tail vein injection, etc] as known in the art

Transgene: Gene of interest that is cloned into a vector for expression in a target organism

Transposase vector: A vector which encodes a transposase

Transposon vector: A vector which encodes a transposon which is a substrate for transposase mediated gene integration

ts: Temperature sensitive

µg: Microgram

µl: Microliter

UTR: Untranslated region of a mRNA (5' or 3' to the coding region)

Vector: A gene delivery vehicle, including viral (e.g. Alphavirus, Poxvirus, Lentivirus, Retrovirus, Adenovirus, Adenovirus related virus, etc.) and non-viral (e.g. plasmid, MIDGE, transcriptionally active PCR fragment, minicircles, bacteriophage, etc.) vectors. These are well known in the art

Vector backbone: Eukaryotic region and bacterial region of a vector, without the transgene or target antigen coding region

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0031] The current technology relates generally to short < 1kb bacterial region plasmid DNA vector methods that improve plasmid manufacture yield and quality, reduce transfection associated toxicity, and increase transgene expression. The current technology can be practiced to improve expression and manufacturing of vectors such as non-viral vectors (transposon vector, transposase vector, Sleeping Beauty transposon vector, PiggyBac transposon vector, PiggyBac transposase vector, expression vector, etc.) and viral vectors (e.g. AAV vector, AAV rep cap vector, AAV helper vector, Ad helper vector, Lentivirus vector, Lentiviral envelope vector, Lentiviral packaging vector, Retroviral vector, Retroviral envelope vector, Retroviral packaging vector, etc.).

[0032] Improved plasmid expression is defined herein as improved transgene expression level and/or expression duration *in vitro* or *in vivo* compared to a transgene encoding plasmid containing a bacterial region encoding the pUC replication origin.

[0033] The methods of plasmid modification of the present current technology have been surprisingly found to provide a solution to provide short spacer region vectors containing structured DNA sequences with efficient high yield manufacture.

[0034] As used herein, the term "sequence identity" refers to the degree of identity between any given query sequence, e.g. SEQ ID NO: 2, and a subject sequence. A subject sequence may, for example, have at least 90 percent, at least 95 percent, or at least 99 percent sequence identity to a given query sequence. To determine percent sequence identity, a query sequence (e.g. a nucleic acid sequence) is aligned to one or more subject sequences using any suitable sequence alignment program that is well known in the art, for instance, the computer program ClustalW (version 1.83, default parameters), which allows alignments of nucleic acid sequences to be carried out across their entire length (global alignment). Chema et al., 2003 Nucleic Acids Res., 31:3497-500. In a preferred method, the sequence alignment program (e.g. ClustalW) calculates the best match between a query and one or more subject sequences, and aligns them so that identities, similarities, and differences can be determined. Gaps of one or more nucleotides can be inserted into a query sequence, a subject sequence, or both, to maximize sequence alignments. For fast pair-wise alignments of nucleic acid sequences, suitable default parameters can be selected that are appropriate for the particular alignment program. The output is a sequence alignment that reflects the relationship between sequences. To further determine percent identity of a subject nucleic acid sequence to a query sequence, the sequences are aligned using the alignment program, the number of identical matches in the alignment is divided by the length of the query sequence, and the result is multiplied by 100. It is noted that the percent identity value can be rounded to the nearest tenth. For example, 78.11, 78.12, 78.13, and 78.14 are rounded down to 78.1, while 78.15, 78.16, 78.17, 78.18, and 78.19 are rounded up to 78.2.

[0035] Turning now to the drawings, FIGS. 1A-1F show annotated maps of: FIG. 1A) R6K origin with the locations of the 22 bp iteron repeats, DnaA boxes 1 and 2, and the regions included in the SEQ ID NO: 1, 2, 3, and 4 R6K origins; FIG. 1B) SEQ ID NO: 5 RNA-OUT selectable marker with the locations of the RNA-OUT promoter -35 and -10 elements, SEQ ID NO: 6 RNA OUT antisense RNA with RNA-IN complementary homology region and RNA-OUT terminator 3' hairpin; FIG. 1C) 14 CpG R6K-RNA-OUT bacterial backbone composed of SEQ ID NO: 1 R6K replication origin and SEQ ID NO: 5 RNA-OUT selectable marker including the trpA bacterial terminator upstream of the R6K origin and flanked by NheI and KpnI cloning sites; FIG. 1D) 3 CpG R6K-RNA-OUT bacterial backbone composed of SEQ ID NO: 2 1x CpG R6K replication origin and SEQ ID NO: 7 2x CpG RNA-OUT selectable marker flanked by NheI and KpnI cloning sites; FIG. 1E) R6K origin from SEQ ID NO: 1, with locations of the 6 iterons highlighted. The individual 22 bp iteron repeat sequences are shown below the origin map; and FIG. 1F) R6K origin from SEQ ID NO: 18, with locations of the 7 iterons highlighted. The individual 22 bp iteron repeat sequences are shown below the origin map. In this example 7 iteron vector iteron 5 has been tandemly duplication; however, a 7 iteron vector of the invention can be obtained by tandem duplication of any of iterons 1, 2, 3, 4, 5 or 6.

[0036] FIGS. 2A-2B show annotated maps of: FIG. 2A) Pol I-dependent pUC origin-Kanamycin selection Sleeping Beauty transposon vector pUC57-Kan SB1 (see Table 6); and FIG. 2B) Pol III-dependent R6K origin-RNA-OUT antibiotic free selection Sleeping Beauty transposon vector NTC9 SB1 (see Table 6). The locations of the left and right Sleeping Beauty IR/DR relative to the bacterial backbone replication origins and selection markers are shown.

[0037] FIGS. 3A-3C show annotated maps of: FIG. 3A) Pol I-dependent pUC origin- Ampicillin selection AAV vector pAAV (see Table 7); FIG. 3B) Pol I-dependent pUC origin-RNA-OUT antibiotic free selection AAV vector NTC8-AAV (see Table 7); and FIG. 3C) Pol III-dependent R6K origin-RNA-OUT antibiotic free selection AAV vector NTC9-AAV (see Table 7). The locations of the left and right AAV ITRs relative to the bacterial backbone replication origins and selection markers are shown.

[0038] FIGS. 4A-4F show annotated maps of: FIG. 4A) Pol I-dependent pUC origin- Ampicillin selection A60 polyA repeat encoding mRNA vector pGEM4Z T7 A60 pA (see Table 8); FIG. 4B) Pol I-dependent pUC origin-RNA-OUT antibiotic free selection A60 polyA repeat encoding mRNA vector NTC8-T7 A60 pA (see Table 8); FIG. 4C) Pol III-dependent R6K origin-RNA-OUT antibiotic free selection A60 polyA repeat encoding mRNA vector NTC9-T7 A60 pA (see Table 8); FIG. 4D) Pol I-dependent pUC origin- Ampicillin selection A99 polyA repeat encoding mRNA vector pT3/T7 A99 pA (see Table 8); FIG. 4E) Pol I-dependent pUC origin-kanR selection A99 polyA repeat encoding mRNA vector NTC7-T7 A99 pA (see Table 8); and FIG. 4F) Pol III-dependent R6K origin-RNA-OUT antibiotic free selection A99 polyA repeat encoding mRNA vector NTC9-T7 A99 pA (see Table 8). The location of the A60 or A99 polyA repeat relative to the bacterial backbone replication origins and selection markers are shown.

[0039] In a first aspect, the present invention provides a method for improving the replication of a covalently closed circular plasmid comprising the following steps:

1. a. providing a covalently closed circular plasmid comprising:
 - iii. a Pol I-dependent origin of replication, and
 - iv. an insert comprising a structured DNA sequence selected from the group consisting of inverted repeat sequence, direct repeat sequence, homopolymeric repeat sequence, or eukaryotic origin of replication, wherein the structured DNA sequence is located at a distance of less than 1000 bp from the Pol I-dependent origin of replication in the direction of replication; and
2. b. modifying the covalently closed circular plasmid of a. such that the Pol I-dependent origin of replication is replaced with an R6K gamma replication origin whereby the resultant R6K gamma replication origin covalently closed circular plasmid has improved replication selected from the group consisting of reduced production of replication intermediates and increased plasmid copy number relative to the covalently closed circular plasmid of a, wherein the R6K gamma replication origin does not require Pol I, and wherein the structured DNA sequence is located at a distance of less than 1000 bp from the R6K gamma replication origin in the direction of replication.

[0040] In an embodiment said Pol I-dependent origin of replication is selected from the group consisting of: pUC origin, pMB 1 origin, and ColE1 origin. In a further

embodiment said R6K gamma replication origin is an R6K gamma replication origin with at least 95% sequence identity to a sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 18. In a further embodiment said structured DNA sequence is selected from the group consisting of polyA repeat, SV40 origin of replication, viral LTR, Lentiviral LTR, Retroviral LTR, transposon IR/DR repeat, Sleeping Beauty transposon IR/DR repeat, AAV ITR, transposon ITR, and PiggyBac transposon ITR.

[0041] In a second aspect, the present invention provides a method for improving the replication of a covalently closed circular plasmid comprising the following steps:

1. a. providing a covalently closed circular plasmid comprising:
 - iii. a bacterial replication-selection region comprising a Pol I-dependent origin of replication and an antibiotic selectable marker, and
 - iv. an insert comprising a structured DNA sequence selected from the group consisting of inverted repeat sequence, direct repeat sequence, homopolymeric repeat sequence, or eukaryotic origin of replication, wherein the structured DNA sequence is located at a distance of less than 1000 bp from the Pol I-dependent origin of replication in the direction of replication; and
2. b. modifying the covalently closed circular plasmid of a. such that the antibiotic selectable marker is replaced with an RNA selectable marker and the Pol I-dependent origin of replication is replaced with an R6K gamma replication origin, whereby the resultant R6K gamma replication origin covalently closed circular plasmid has improved replication selected from the group consisting of reduced production of replication intermediates and increased plasmid copy number relative to the covalently closed circular plasmid of a, wherein the R6K gamma replication origin does not require Pol I, and wherein the structured DNA sequence is located at a distance of less than 1000 bp from the R6K gamma replication origin in the direction of replication.

[0042] In an embodiment said Pol I-dependent origin of replication is selected from the group consisting of: pUC origin, pMB1 origin, and ColE1 origin. In a further embodiment said R6K gamma replication origin is an R6K gamma replication origin with at least 95% sequence identity to a sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 18. In a further embodiment said RNA selectable marker is an RNA-IN regulating RNA-OUT functional variant with at least 95% sequence identity to a sequence selected from the group consisting of SEQ ID NO: 5, and SEQ ID NO: 7. In a further embodiment said RNA selectable marker is an RNA-OUT RNA selectable marker that encodes an RNA-IN regulating RNA-OUT RNA with at least 95% sequence identity to SEQ ID NO: 6. In a further embodiment said bacterial replication-selection region comprising a Pol I-dependent origin of replication and an antibiotic selectable marker is replaced with a Pol III-dependent R6K origin-RNA-OUT RNA selectable marker bacterial replication-selection region with at least 95% sequence identity to a sequence selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, and SEQ ID NO: 17. In a further embodiment said structured DNA sequence is selected from the group consisting of polyA repeat, SV40 origin of replication, viral LTR, Lentiviral LTR, Retroviral LTR, transposon IR/DR repeat, Sleeping Beauty transposon IR/DR repeat, AAV ITR, transposon ITR, and PiggyBac transposon ITR.

EXAMPLES

[0043] The methods of the current technology are further illustrated by the following examples. These are provided by way of illustration and are not intended in any way to limit the scope of the disclosure.

Example 1: pUC, and R6K replication origin plasmid replication and production

[0044] pUC origin vector replication and production background: The vast majority of therapeutic plasmids use the pUC origin which is a high copy derivative of the pMB1 origin (closely related to the ColE1 origin). For pMB 1 replication, plasmid DNA synthesis is unidirectional and does not require a plasmid borne initiator protein. The pUC origin is a copy up derivative of the pMB 1 origin that deletes the accessory ROP (rom) protein and has an additional temperature sensitive mutation that destabilizes the RNAI/RNAlI interaction. Shifting of a culture containing these origins from 30 to 42°C leads to an increase in plasmid copy number. pUC plasmids can be produced in a multitude of *E. coli* cell lines.

[0045] RNA-OUT antibiotic free selectable marker background: Antibiotic-free selection is performed in *E. coli* strains containing phage lambda attachment site chromosomally integrated *pCAH63-CAT RNA-IN-SacB (P516 616)* as described in Williams, *Supra*, 2008. SacB (*Bacillus subtilis* levansucrase) is a counterselectable marker which is lethal to *E. coli* cells in the presence of sucrose. Translation of SacB from the RNA-IN-SacB transcript is inhibited by plasmid encoded RNA-OUT (Fig. 1B). This facilitates plasmid selection in the presence of sucrose, by inhibition of SacB mediated lethality.

[0046] R6K origin vector replication and production background: The R6K gamma plasmid replication origin requires a single plasmid replication protein π that binds as a replication initiating monomer to multiple repeated 'iteron' sites (seven core repeats containing TGAGNG consensus) and as a replication inhibiting dimer to repressive sites (TGAGNG) and to iterons with reduced affinity. Replication requires multiple host factors including IHF, DnaA, and primosomal assembly proteins DnaB, DnaC, DnaG (Abhyankar et al., 2003 J Biol Chem 278:45476-45484). The R6K core origin contains binding sites for DnaA and IHF that affect plasmid replication since π , IHF and DnaA interact to initiate replication.

[0047] Different versions of the R6K gamma replication origin have been utilized in various eukaryotic expression vectors, for example pCOR vectors (Soubrier et al., 1999, Gene Therapy 6: 1482-88) and a CpG free version in pCpGfree vectors (Invivogen, San Diego CA), and pGM169 (University of Oxford). Incorporation of the R6K replication origin per se does not improve transgene expression levels compared to an optimized pUC origin vector (Soubrier et al., *Supra*, 1999). However, use of a conditional replication origin such as R6K gamma that requires a specialized cell line for propagation adds a safety margin since the vector will not replicate if transferred to a patient's endogenous flora.

[0048] A highly minimized 6 iteron R6K gamma derived replication origin (SEQ ID NO:1; Fig. 1E) that contains core sequences required for replication (including the DnaA box and stb 1-3 sites; Wu et al., 1995. J Bacterial. 177: 6338-6345), but with the upstream π dimer repressor binding sites and downstream π promoter deleted (by removing one copy of the iterons) was described in Williams, *Supra*, 2014. This R6K origin contains 6 tandem direct repeat iterons (Fig. 1E). The NTC9385R Nanoplasmid™ vector including this minimized R6K origin and the RNA-OUT AF selectable marker in the spacer region, was described in Williams, *Supra*, 2014.

[0049] Typical R6K production strains express from the genome the π protein derivative PIR116 that contains a P106L substitution that increases copy number (by reducing π dimerization; π monomers activate while π dimers repress). Fermentation results with pCOR (Soubrier et al., *Supra*, 1999) and pCpG plasmids (Hebel HL, Cai Y, Davies LA, Hyde SC, Pringle IA, Gill DR. 2008. Mol Ther 16: S110) were low, around 100 mg/L in PIR116 cell lines.

[0050] Mutagenesis of the pir-116 replication protein and selection for increased copy number has been used to make new production strains. For example, the TEX2pir42 strain contains a combination of P106L and P42L. The P42L mutation interferes with DNA looping replication repression. The TEX2pir42 cell line improved copy number and fermentation yield with pCOR plasmids with reported yields of 205 mg/L (Soubrier F. 2004. World Patent Application WO2004033664).

[0051] Other combinations of π copy number mutants that improve copy number include 'P42L and P113S' and 'P42L, P106L and F107S' (Abhyankar et al., 2004. J Biol Chem 279:6711-6719).

[0052] Williams, *Supra*, 2014 describes host strains expressing phage HK022 attachment site integrated pL promoter heat inducible π P42L, P106L and F107S high copy mutant replication (Rep) protein for selection and propagation of R6K origin Nanoplasmid™ vectors. This is an additional Nanoplasmid™ safety factor since R6K origin vectors can only replicate within the engineered Rep protein-expressing *E. coli* host strain.

[0053] RNA-OUT selectable marker-R6K plasmid propagation and fermentations described in Williams, *Supra*, 2014 were performed using heat inducible 'P42L, P106L and F107S' π copy number mutant cell lines such as DH5 α host strain NTC711772 = DH5 α dcm- att λ ::P_{5/6} 6/6-RNA-IN-SacB, catR; att_{HK022}::pL (OL1-G to T) P42L-P106L-F107S (P3-), SpecR StrepR. Production yields up to 695 mg/L were reported.

[0054] Additional cell lines were created and disclosed herein including:

NTC821601 DH5 α att λ ::P_{5/6} 6/6-RNA-IN- SacB, catR; att_{HK022}::pL (OL1-G to T) P42L-P106L-F107S (P3-), SpecR StrepR = dcm+ version of NTC711772

NTC940211 DH5 α att λ ::P_{5/6} 6/6-RNA-IN- SacB, catR; att_{HK022}::pL (OL1-G to T) P42L-P106L-F107S

P113S (P3-), SpecR StrepR = high copy substitution of P106L for P106L combined with P113S to create a quadruple copy number increasing mutant rep protein derivative of NTC821601

NTC1050811 DH5 α att λ ::P_{5/6} 6/6-RNA-IN- SacB, catR; att_{HK022}::pL (OL1-G to T) P42L-P106L-F107S

P113S (P3-), SpecR StrepR; att₈₀₈::pARA-CI857ts, tetR = pARA-CI857ts derivative of NTC940211. This strain contains a phage ϕ 80 attachment site chromosomally integrated copy of an arabinose inducible CI857ts gene. Addition of arabinose to plates or media (e.g. to 0.2-0.4% final concentration) induces pARA mediated CI857ts repressor expression which reduces copy number at 30° C through CI857ts mediated downregulation of the Rep protein expressing pL promoter [*i.e.* additional CI857ts mediates more effective downregulation of the pL (OL1-G to T) promoter at 30°C]. Copy number induction after temperature shift to 37-42°C is not impaired since the CI857ts repressor is inactivated at these elevated temperatures. A dcm- derivative (NTC1050811 dcm-) is used in cases where dcm methylation is undesirable.

NTC1011641; Stb14 att λ ::P_{5/6} 6/6-RNA-IN- SacB, catR; att_{HK022}::pL P42L-P106L-F107S (P3-) SpecR StrepR = Stb14 version of NTC661135 (XL1Blue- dcm- att λ ::P_{5/6} 6/6-RNA-IN- SacB, catR; att_{HK022}::pR pL P42L-P106L-F107S (P3-) SpecR StrepR described in Williams, *Supra*, 2014

[0055] Nanoplasmid™ production yields are improved with the quadruple mutant heat inducible pL (OL1-G to T) P42L-P106L-F107S **P113S** (P3-) compared to the triple mutant heat inducible pL (OL1-G to T) P42L-P106L-F107S (P3-) described in Williams, *Supra*, 2014. Yields in excess of 2 g/L Nanoplasmid™ have been obtained with the quadruple mutant NTC1050811 cell line (e.g. 2240 mg/L with NTC9 T7 A99 pA, **Table 8**)

[0056] Use of a conditional replication origin such as these R6K origins that requires a specialized cell line for propagation adds a safety margin since the vector will not replicate if transferred to a patient's endogenous flora.

Example 2: pUC and R6K origin vector production

[0057] Shake flask production: Shake flask production was performed using proprietary Plasmid+ shake culture medium. The seed cultures were started from glycerol stocks or colonies and streaked onto LB medium agar plates containing 50 μ g/mL antibiotic (for ampR or kanR selection plasmids) or 6% sucrose (for RNA-OUT selection plasmids). The plates were grown at 30-32°C; cells were resuspended in media and used to provide approximately 2.5 OD₆₀₀ inoculums for the 500 mL Plasmid+ shake flasks that contained 50 μ g/mL antibiotic for ampR or kanR selection plasmids or 0.5% sucrose to select for RNA-OUT plasmids. Flasks were grown with shaking to saturation at the growth temperatures as indicated in **Tables 5, 6, 7, and 9**.

[0058] Fermentation production: Fermentations were performed using proprietary fed-batch media (NTC3019, HyperGRO media) in New Brunswick BioFlo 110 bioreactors as described (Carnes and Williams, *Supra*, 2011). The seed cultures were started from glycerol stocks or colonies and streaked onto LB medium agar plates containing 50 μ g/mL antibiotic (for ampR or kanR selection plasmids) or 6% sucrose (for RNA-OUT selection plasmids). The plates were grown at 30-32°C; cells were resuspended in media and used to provide approximately 0.1% inoculums for the fermentations that contained 50 μ g/mL antibiotic for ampR or kanR selection plasmids or 0.5% sucrose for RNA-OUT plasmids. HyperGRO temperature shifts were as indicated in **Tables 8 and 9**.

[0059] Production hosts: pUC origin AmpR or KanR plasmid fermentations were performed in *E. coli* strain DH5 α [F- Φ 80*lacZ* Δ M15 Δ (*lacZ*Y Δ -argF) U169 *recA1 endA1 hsdR17* (rK-, mK+) *phoA supE44* λ - *thi-1 gyrA96 relA1*] (Invitrogen, Carlsbad CA) or Stb14.

[0060] Antibiotic-free pUC origin RNA-OUT plasmid fermentations were performed in *E. coli* strain DH5 α containing phage lambda attachment site chromosomally integrated *pCAH63-CAT RNA-IN SacB (P5/6 6/6)* as described in Williams, *Supra*, 2008. The production strain is NTC4862 = DH5 α att λ ::P_{5/6} 6/6-RNA-IN- SacB, catR.

[0061] Antibiotic-free R6K gamma origin RNA-OUT plasmid propagation and fermentations were performed using *E. coli* RNA-OUT selection hosts further encoding phage HK022 attachment site integrated pL promoter heat inducible π copy number mutant cell line lines, methods for the creation of which are described in Williams, *Supra*, 2014.

Production strains:

[0062]

pUC origin-AmpR or KanR antibiotic selection hosts

DH5 α

Stb14

pUC origin-RNA-OUT sucrose selection hosts

NTC4862 DH5α attλ::P_{5/6 6/6}-RNA-IN- SacB, catR

NTC1011592 Stbl4 attλ::P_{5/6 6/6}-RNA-IN- SacB, catR

R6K origin-RNA-OUT sucrose selection Nanoplasmid™ hosts

NTC1050811 DH5α attλ::P_{5/6 6/6}-RNA-IN- SacB, catR; att_{HK022}::pL (OL1-G to T) P42L-P106L-F107S P113S (P3-), SpecR StrepR; att_{φ80}::pARA-Cl857ts, tetR

NTC1011641 Stbl4 attλ::P_{5/6 6/6}-RNA-IN- SacB, catR; att_{HK022}::pL P42L-P106L-F107S (P3-) SpecR StrepR

[0063] Analytical Methods: Culture samples were taken at key points and at regular intervals during all fermentations. Samples were analyzed immediately for biomass (OD₆₀₀) and for plasmid yield. Plasmid yield was determined by quantification of plasmid obtained from Qiagen Spin Miniprep Kit preparations as described (Carnes and Williams, *Supra*, 2011). Briefly, cells were alkaline lysed, clarified, plasmid was column purified, and eluted prior to quantification. Plasmid quality was determined by agarose gel electrophoresis analysis (AGE) and was performed on 0.8-1% Tris/acetate/EDTA (TAE) gels as described in Carnes and Williams, *Supra*, 2011.

Example 3: pUC and R6K origin structured vector construction and manufacturing

[0064] The R6K gamma origin (SEQ ID NO:1; Fig. 1E)-RNA-OUT (SEQ ID NO:5; Fig. 1B) bacterial replication-selection region (SEQ ID NO:8; Fig. 1C) was cloned into the polylinker region of a variety of pUC57 based vectors to create the pNTC-NP1, pNTC-NP2, pNTC-NP3, pNTC-NP4, pNTC-NP5, pNTC-NP6, pNTC-NP7, vectors. Each vector has different flanking restriction sites that can be used to retrofit a target vector to R6K replication-RNA-OUT selection. The 5' and 3' polylinker sequences flanking the R6K-RNA-OUT insert in the pNTC-NP 1-7 vectors are shown in Table 4. A pUC57 based version of the 1 CpG R6K gamma origin- 2 CpG RNA-OUT bacterial replication-selection region (SEQ ID NO:9; Fig. 1D) was also created (pNTC-3xCpG NP1) and is shown in Table 4.

[0065] The R6K gamma origin (SEQ ID NO:1) is an engineered 6 iteron R6K origin (Fig. 1E). A pUC57 based version of a 7 iteron R6K gamma origin (SEQ ID NO:18; Fig. 1F)-RNA-OUT (SEQ ID NO:5; Fig. 1B) bacterial replication-selection region was also created and used to construct and evaluate the utility of additional iterons on manufacturing. Similarly, high quality, high yield manufacture was obtained with vectors differing only by containing either the SEQ ID NO:18 seven iteron R6K gamma origin or the six iteron R6K gamma origin (SEQ ID NO:1). For example, the following harvest production yields were obtained in 30-42°C 10 hr ramp temperature shift HyperGRO fermentations:

SEQ ID NO:1 6 iteron 3203 bp R6K origin vector: a biomass of 120 OD₆₀₀; plasmid titer of 1363 mg/L; plasmid specific yield of 11.3 mg plasmid/L/OD₆₀₀

SEQ ID NO:18 7 iteron 3225 bp R6K origin vector: a biomass of 137 OD₆₀₀; plasmid titer of 1503 mg/L; plasmid specific yield of 11.0 mg plasmid/L/OD₆₀₀

[0066] The 7 iteron R6K gamma origin in SEQ ID NO:18 is a tandem duplication of iteron 5 (Fig. 1F; SEQ ID NO:18) but the 7 iteron R6K gamma origin vectors of the invention can be tandem duplications of any of the iterons given as SEQ ID NO:19; SEQ ID NO:20; SEQ ID NO:21; SEQ ID NO:22; SEQ ID NO:23 (Fig. 1E), or random combinations of SEQ ID NO:19; SEQ ID NO:20; SEQ ID NO:21; SEQ ID NO:22; SEQ ID NO:23 into 7 iteron R6K origin compositions, or iteron repeat variants that retain the TGAGNG consensus. Additional iteron derivatives (e.g. 8, 9 or 10 iteron vectors) are also contemplated for practice of the invention.

Table 4: pNTC multiple cloning site flanked R6K Origin-RNA-OUT selection marker vectors

Vector	R6K 5' flanking restriction sites	trpA term	R6K origin	Linker site	RNA OUT Selection marker	RNA-OUT 3' flanking restriction site
pNTC-NP1 (SEQ ID NO: 10)	EcoRI, SacI, KpnI, NruI, NsiI, XmaIII, NotI, <i>NheI</i>	Yes	SEQ ID NO: 1	DraIII ^a	SEQ ID NO: 5	<i>NheI</i> BamHI, XmaI, ApaI, SalI, HincII, PstI, StuI, AatI, SphI, <i>HindIII</i> (in R6K)
pNTC-NP2 (SEQ ID NO: 11)	EcoRI, SacI, KpnI, NruI, NsiI, XmaIII, NotI, <i>NheI</i>	Yes	SEQ ID NO: 1	DraIII ^a	SEQ ID NO: 5	SpeI, XmaI, SspI BamHI, XmaI, ApaI, SalI, HincII, PstI, StuI, AatI, SphI, <i>HindIII</i> (in R6K)
pNTC-NP3 (SEQ ID NO: 12)	EcoRI, <i>SacI</i> , <i>KpnI</i> , NruI, NsiI, XmaIII, NotI, <i>NheI</i>	Yes	SEQ ID NO: 1	DraIII ^a	SEQ ID NO: 5	<i>KpnI</i> , <i>SacI</i> BamHI, XmaI, ApaI, SalI, HincII, PstI, StuI, AatI, SphI, <i>HindIII</i> (in R6K)
pNTC-NP4 (SEQ ID NO: 13)	<i>NheI</i> , XmaIII, NotI, NsiI, NruI, <i>KpnI</i> , <i>SacI</i> BamHI, XmaI, ApaI, SalI, HincII, SfcI, PstI, StuI, AatI, SphI, <i>HindIII</i> (in R6K)	Yes	SEQ ID NO: 1	DraIII ^a	SEQ ID NO: 5	EcoRI, <i>SacI</i> , <i>KpnI</i>
pNTC-NP5 (SEQ ID NO: 14)	KasI, <i>NheI</i>	Yes	SEQ ID NO: 1	DraIII ^a	SEQ ID NO: 5	<i>KpnI</i> AflIII PstI, AatI, SphI, <i>HindIII</i> (in R6K)
pNTC-NP6 (SEQ ID NO: 15)	EcoRI, PstI, EcoRV, BstXI, NotI, <i>NheI</i>	Yes	SEQ ID NO: 1	DraIII ^a	SEQ ID NO: 5	<i>KpnI</i> , ApaI, PvuI, SalI, <i>SacI</i>
pNTC-NP7 (SEQ ID NO: 16)	BssHII PacI <i>NheI</i>	Yes	SEQ ID NO: 1	DraIII ^a	SEQ ID NO: 5	<i>KpnI</i> PacI BssHII
pNTC-3x CpG NP1 (SEQ ID NO: 17)	XhoI, XbaI, ApaI, SalI, HincII, PstI, StuI, AatI, SphI, <i>HindIII</i> (in R6K)	No	SEQ ID NO: 2	BsrGI	SEQ ID NO: 7	EcoRI, <i>SacI</i> , <i>KpnI</i> , NruI, NsiI, XmaIII, NotI, <i>NheI</i> , <i>KpnI</i>

^a Non-palindromic unique 3 bp NNN sticky end DraIII site (CACNNNGTG) separating R6K and RNA-OUT of sequence CACGTTGTG can be used to assemble R6K and RNA-OUT from separate pNTC vectors in directional multi-fragment ligation reactions

[0067] Viral and non-viral vector pUC origin-antibiotic selection bacterial backbone retrofits to R6K-RNA-OUT were performed by:

1. 1) selecting restriction sites that flank the pUC origin and antibiotic selection marker region in the target viral and non-viral vector;
2. 2) Identifying a pNTC-NP compatible polylinker -R6K-RNA-OUT polylinker cassette (either pNTC-NP1, 2, 3, 4, 5, 6, or 7; Table 4);

3. 3) Excising the pUC origin antibiotic selection marker region and replacing with the selected R6K origin RNA-OUT region using the selected restriction digestion approach and standard ligase mediated cloning.

[0068] In some cases, the R6K origin and RNA-OUT units were assembled in multi-fragment ligations from separate restriction fragments using the non-palindromic DralIII linker site (see **Table 4**). In the case of the fd6 Ad helper retrofit (**Table 9**), a 3-fragment ligation was performed using a short 500 bp synthetic gene DralIII RNA-OUT-Ad helper-AvrII to link RNA-OUT to a unique AvrII site in the fd6 Ad helper eukaryotic region in a 12 kb AvrII-SalI restriction fragment, and to a SalI-R6K origin-DralIII fragment from pNTC-NP4.

[0069] Example vector maps and vector characteristics of the original pUC origin- antibiotic selection marker vector and the retrofitted R6K origin-RNA-OUT antibiotic free selection marker vectors are shown for Sleeping Beauty (**Fig 2; Table 6**), AAV (**Fig. 3; Table 7**) and mRNA (**Fig. 4; Table 8**) vectors. The vector characteristics of the original pUC origin- antibiotic selection marker vector and the retrofitted R6K origin-RNA-OUT antibiotic free selection marker vectors are shown for AAV helper vectors (**Table 8**). The vector characteristics of pUC origin- RNA-OUT antibiotic free selection marker vector and the retrofitted R6K origin-RNA-OUT antibiotic free selection marker vectors are shown for Lentiviral vectors (**Table 5**) and AAV vectors (**Table 7**).

[0070] In all cases, the bacterial backbone size was < 1 kb in the R6K origin-RNA-OUT antibiotic free selection marker retrofitted vectors (460-610 bp). This is well below the 1.1 kb bacterial backbone size limit required to improve vector expression level (**Tables 1-2**) and duration (Quiviger *et al.*, *Supra*, 2014). In all cases, the original pUC origin-antibiotic selection bacterial backbone prior to retrofit was > 1.2 kb (2340-2750 bp) as were the pUC origin-RNA-OUT retrofits (1210-1500 bp). Thus, these AAV, AAV helper, Sleeping Beauty, and Lentiviral R6K origin-RNA-OUT antibiotic free selection marker retrofit vectors meet the short spacer region requirement for improved expression level and duration compared to the original pUC origin- antibiotic selection marker vector. Additionally, these AAV, AAV helper, Sleeping Beauty, and Lentiviral R6K origin-RNA-OUT antibiotic free selection marker retrofit vectors have no chance of antibiotic marker gene transfer by transduction (AAV, Lentiviral vectors) or transposition (Sleeping Beauty vectors) due to removal of the KanR or ampR antibiotic resistance selection marker in the parent vector. Additionally, the vectors of the current technology do not require the complicated difficult to scale expensive additional manufacturing steps required to remove the large bacterial region between the eukaryotic polyA and promoter with minicircle vectors (Kay *et al.*, *Supra*, 2010).

[0071] However, in a Lentiviral vector the eukaryotic region contains flanking direct repeat LTRs, in an AAV vector the eukaryotic region contains flanking inverted terminal repeats, while in a Sleeping Beauty transposon vector the eukaryotic region contains flanking transposon IR/DR termini. These flanking sequences are all structured DNA sequences.

[0072] Levy, *Supra*, 2004 teaches that replication intermediates form when any high copy number prokaryotic origin of replication is < 1kb from a structured DNA sequence such as an enhancer, LTR or IRES, but not when the high copy replication origin is >1.5 kb. Consistent with this, Replication intermediates were formed in all pUC origin- RNA-OUT marker vectors in which the pUC origin was <1 kb from a Lentiviral vector LTR (**Table 5**: 400 bp) or a pUC origin- antibiotic resistance marker vector in which the pUC origin was <1 kb from a Sleeping Beauty IR/DR (**Table 6**: 280 bp). For AAV and mRNA vectors, the original pUC origin- antibiotic selection marker vectors have the pUC origin 0 bp from an ITR (AAV vector; **Table 7**) or 170 bp from an A99 repeat (mRNA vector, **Table 8**) which may make a replication intermediate that is too small to detect on an agarose gel. However, in these cases production yields were very low, indicative of low plasmid copy number due to replication blockage. By contrast, as expected, in the case where the original pUC origin- antibiotic selection marker vector pUC origin was > 1.5 kb from a structured DNA sequence (A60 repeat), high plasmid production yields were obtained (**Table 8**: mRNA vector pGEM4Z T7 A60).

[0073] Williams, *Supra*, 2017 reported that pUC origin vector production yield is improved with a PAS-BH extended pUC origin when the pUC origin is >1.5 kb from a homopolymeric A64C31 repeat. However, production yields were low when a PAS-BH extended pUC origin is orientated <400 bp from the A64C31 repeat (**Table 8**, see footnotes d and e). This teaches that addition of a PAS-BH primosomal assembly site does not overcome the poor pUC origin directed replication of closely positioned structured DNA sequences.

[0074] Since the pUC origin itself is 1 kb, there is no configuration to make a <1.1 kb bacterial region AAV, Lentiviral, Retroviral or transposon vector containing the pUC origin which is not predicted to produce replication intermediates as seen above and predicted by Levy, *Supra*, 2004 and poor plasmid yields as reported herein.

[0075] Surprisingly, replication intermediates were not observed in any R6K origin-RNA-OUT antibiotic free selection marker retrofitted vectors, include those in which the R6K origin was <1 kb from a Lentiviral vector LTR (**Table 5**: 400 bp) or a Sleeping Beauty IR/DR (**Table 6**: <40 bp). Further, for AAV vectors, while the original pUC origin- antibiotic selection marker vectors with the pUC origin 0 bp from the ITR had very poor production yields, the two R6K origin-RNA-OUT antibiotic free selection marker retrofitted vectors with the R6K origin 40 bp from the ITR had much higher production yields (**Table 7**). This improved production is specific to R6K and not RNA-OUT, since the two AAV pUC-RNA-OUT retrofits with the pUC origin 50 bp from the ITR had equally poor plasmid production yields as the original pUC antibiotic marker vector (**Table 7**); as well the direct comparison of pUC-RNA-OUT with R6K-RNA-OUT retrofits positioned 400 bp from an LTR repeat in a Lentiviral backbone showed replication intermediates with all three pUC-RNA-OUT backbones but none of the three R6K-RNA-OUT backbones (**Table 5**).

[0076] This surprising improvement in plasmid copy number (plasmid production yields) and quality (eliminated replication intermediates) with the R6K origin vector implies that the R6K origin can replicate through a structured DNA sequence more effectively than the pUC origin. While Levy, *Supra*, 2004 teaches that replication intermediates form when any high copy number prokaryotic origin of replication is < 1kb from a structured DNA sequence such as an enhancer, LTR or IRES, but not when the high copy replication origin is >1.5 kb away, the examples provided by Levy, *Supra*, 2004 were all with pUC origin plasmids.

[0077] A fundamental difference between these replication origins is that the pUC origin is a Pol I dependent origin of replication while the R6K origin is a Pol III dependent origin of replication. With the pUC origin the RNAII primer forms an RNA: DNA R-loop that is cleaved by RNase H to create a primer for DNA Pol I directed DNA synthesis during initial leading strand synthesis. DNA synthesis then converts from slow DNA Pol I to the highly processive DNA Pol III from 400 bp to up to 1.3 kb downstream of the origin (Allen *et al.*, 2011. *Nucleic Acids Research* 39:7020-33). The R6K gamma replication origin rep protein interacts with dnaB helicase and dnaG primase which creates short RNA primers for DNA Pol III replication without requirement for DNA Pol I (Abhyankar *et al.*, *Supra*, 2003). The pUC origin DNA Pol I replication zone of up to 1.3 kb from the origin corresponds closely with the Levy, *Supra*, 2004 defined upper limit of replication intermediate formation (between 1 and 1.5 kb from the origin). We propose that the observed surprisingly improved replication of structured DNA when in close proximity to the R6K but not the pUC origin is due to an unexpected improvement of replication of structured DNA sequences by DNA Pol III compared to DNA Pol I.

[0078] The vector methods and compositions disclosed herein demonstrate that a Pol III-dependent origin of replication such as the R6K origin can be used to replicate structured DNA sequences which are poorly replicated by a Pol I-dependent origin of replication such as the pUC origin.

[0079] These results demonstrate the vectors of the invention are useful for improving viral and non-viral vector manufacturing yield and quality.

Example 4: Improved performance of R6K origin structured vector

[0080] The vectors of the invention are additionally useful for eliminating antibiotic resistance marker gene transfer by viral and non-viral vectors; reducing transfection

associated toxicity; improving transposition from non-viral transposon vectors; improving packaging titers from viral vectors; improving expression of viral and non-viral vector encoded transgenes, etc.

[0081] As an example, R6K origin third generation lentiviral vectors [4 vectors: **Table 5** transfer plasmid, gag pol packaging plasmid; env plasmid; REV plasmid (not shown) with R6K origin and < 1 kb bacterial backbone] of the invention showed reduced toxicity and improved viral packaging titers compared to pUC origin vector comparators with >1.5 kb bacterial backbone. Transfection of Lenti-X 293 T cell line (Takara Bio Mountain View, CA) with **Table 5** R6K origin third generation lentiviral vectors with <1 kb bacterial backbone or original pUC origin-antibiotic selection marker vector with >1.5 kb bacterial backbone control in 24 well plates using Lipofectamine 3000 (Thermo Fisher Scientific, Waltham, MA) as recommended by the manufacturer resulted in higher titer lentivirus production (>1.5 kb bacterial backbone pUC origin control vectors: $1.00x \pm 0.32$; < 1 kb bacterial backbone R6K origin vectors $1.45x \pm 0.42$) as measured using the Lenti-X p24 Rapid Titer Kit (Takara Bio Mountain View, CA). Transfection of Lenti-X 293 T cell line in 24 well plates with **Table 5** third generation lentiviral vectors (>1.5 kb bacterial backbone pUC origin control or < 1 kb bacterial backbone R6K origin) using Calcium Phosphate transfection as described (Marino MP, Luce MJ, Reiser J. 2003, Methods Mol Biol 229:43-55) resulted in higher titer lentivirus production (>1.5 kb bacterial backbone pUC origin control: $1.00x \pm 0.30$; < 1 kb bacterial backbone R6K origin vectors: $1.32x \pm 0.19$) as measured using the Lenti-X p24 Rapid Titer Kit (Takara Bio Mountain View, CA). Significantly, Calcium Phosphate transfection of the >1.5 kb bacterial backbone pUC origin third generation Lentiviral vectors resulted in extensive transfection associated toxicity (>80% cell death) compared to low toxicity with the R6K origin < 1 kb bacterial backbone R6K origin third generation Lentiviral vectors in this 24 well plate transfection. This reduced transfection associated toxicity should result in dramatically improved viral titers in larger manufacturing scale transfections. These results demonstrate the < 1 kb bacterial backbone R6K origin Nanoplasmid vectors of the invention reduce transfection associated toxicity and improve packaging titers from viral vectors compared to >1.5 kb bacterial backbone vectors.

Summary

[0082] While the above description contains many examples, these should not be construed as limitations on the scope of the disclosure, but rather should be viewed as an exemplification of preferred embodiments thereof. Many other variations are possible.

[0083] For example, in the vectors of the current technology various orientations of the Pol III dependent replication origin, and the RNA selectable marker, may be utilized. For example, any of the eight orientations of the Pol III dependent replication origin, and the RNA selectable marker in vectors of the current technology may be used (i.e. ← Pol III replication origin RSM→; ← Pol III replication origin ← RSM; Pol III replication origin - RSM →; Pol III replication origin → ← RSM; ← RSM Pol III replication origin →; ← RSM ← Pol III replication origin; RSM - Pol III replication origin →; RSM → ← Pol III replication origin).

[0084] Further, a variety of RNA selectable markers known in the art may be substituted for RNA-OUT.

[0085] Further, an antibiotic resistance maker may be substituted for RNA-OUT, for example in the case where a simple retrofit of the pUC origin to the R6K origin is desired to improve plasmid production yield and/or quality.

[0086] Thus, the reader will see that the improved Pol III dependent replication origin vectors of the current technology provide for an approach to reduce transfection associated toxicity, improve transposition from non-viral transposon vectors, improve packaging titers from viral vectors, improve expression of viral and non-viral vector encoded genes, and eliminate viral vector and non-viral vector mediated antibiotic selection marker gene transfer (i.e. through incorporation of a bacterial region preferably less than 1000 bp) while dramatically improving manufacture compared to alternative vectors such as pUC plasmids and minicircles.

Table 5: Lentiviral vectors: pUC origin-RNA-OUT versus R6K origin-RNA-OUT shake flask production yields/quality

Lentiviral Vector	Replication origin	Vector size	Transfer plasmid bacterial backbone (LTR spacing) ^a	Replication origin spacing from SV40 Ori ^b	Replication origin spacing from LTR ^c	Cell line ^d	Production culture ^e	Harvest OD ₆₀₀	Harvest spec yield (mg/L/OD ₆₀₀)	Harvest yield (mg/L)	Production quality ^f
Transfer plasmid 1 with SV40 Ori	pUC origin	9.9 kb	1210 bp (1210 bp)	<50 bp	400 bp	DH5α	30 to 37C	7.5	3.4	26	CCC Monomer + Replicate Intermediate
	R6K origin	9.2 kb	460 bp (460 bp)	<50 bp	400 bp	DH5α	30 to 37C	5.0	4.7	24	CCC Monomer
Transfer plasmid 2 with SV40 Ori	pUC origin	7.3 kb	1210 bp (1210 bp)	<50 bp	400 bp	DH5α	30 to 37C	9.0	3.9	35	CCC Monomer + Replicate Intermediate
	R6K origin	6.5 kb	460 bp (460 bp)	<50 bp	400 bp	DH5α	30 to 37C	8.4	7.2	61	CCC Monomer
Transfer plasmid 3 with SV40 Ori	pUC origin	8.3 kb	1210 bp (1210 bp)	<50 bp	400 bp	DH5α	30 to 37C	16.8	3.0	51	CCC Monomer + Replicate Intermediate
	R6K origin	7.5 kb	460 bp (460 bp)	<50 bp	400 bp	DH5α	30 to 37C	11.9	7.7	92	CCC Monomer
Gag pol packaging plasmid	pUC origin	8.4 kb	1210 bp	NA	NA	DH5α	30 to 37C	10.2	4.3	44	CCC Monomer
	R6K origin	7.5 kb	460 bp	NA	NA	DH5α	30 to 37C	9.7	7.6	73	CCC Monomer
Envelope plasmid	pUC origin	5.4 kb	1210 bp	NA	NA	DH5α	30 to 37C	17.0	5.7	96	CCC Monomer
	R6K origin	4.4 kb	460 bp	NA	NA	DH5α	30 to 37C	10.8	6.0	65	CCC Monomer

^aAll vectors are antibiotic free RNA-OUT selection retrofits. Original pUC origin-antibiotic selection marker vector bacterial backbone size were all > 1.5 kb ^b Distance in bp from 3' end of replication origin to SV40 origin in direction of replication ^c Distance in bp from 3' end of replication origin to nearest LTR in direction of replication ^d DH5α = NTC4862 for pUC origin, NTC1050811 for R6K origin ^e 500 mL Plasmid+ Shake Flask Culture temperature shifted from 30 to 37C ^f Harvest plasmid preparations with detectable replication intermediates indicated. Levy, *Supra*, 2004 teach that replication intermediates form when

any high copy number prokaryotic origin of replication is less than 1000 bp from a promoter/enhancer or LTR promoter or DNA repeat or a complex secondary structure such as an IRES, while replication intermediates do not form if the origin of replication is a distance greater than about 1.5 kb.

Table 6: Sleeping Beauty Transposon and transposase vector: pUC versus R6K origin shake flask production yields/quality

Replication origin/ vector	Bacterial selection	Vector IR/DR Configuration	Vector size	bacterial backbone size (IR/DR spacing)	Replication origin spacing from IR/DR ^a	Cell line ^b	Production culture ^c	Harvest OD ₆₀₀	Harvest spec yield (mg/L/OD ₆₀₀)	Harvest yield (mg/L)	Production quality ^d
Transposon vector pUC origin (pUC57-kan-SB1: see Fig. 2)	KanR	IR DR(R) <pUC<kanR IR DR(L) Promoter> Transgene>	5.5 kb	2600 bp (2600 bp)	280 bp	DH5α	30 to 37C	1.9	2.1	4	CCC Monomer +Replicate Intermediate
Transposon vector R6K origin (NTC9-SB1: see Fig 2)	RNA-OUT AF	IR DR(R) <R6K R-OUT > IR DR(L) Promoter> Transgene>	3.4 kb	475 bp (475 bp)	<40 bp	DH5α	30 to 37C	11.2	1.0	11	CCC Monomer
Transposon vector R6K origin (NTC9-SB2)	RNA-OUT AF	IR DR(R) <R6K R-OUT > IR DR(L) Promoter> Transgene>	3.6 kb	475 bp (475 bp)	<40 bp	DH5α	30 to 37C	4.1	6.2	25	CCC Monomer
Transposase vector pUC origin	KanR	Not Applicable	4.7 kb	2600 bp (NA)	Not Applicable	DH5α	30 to 37C	21.6	1.8	39	CCC Monomer
Transposase vector R6K origin	RNA-OUT AF	Not Applicable	2.5 kb	475 bp (NA)	Not Applicable	DH5α	30 to 37C	10.9	4.7	51	CCC Monomer

^a Distance in bp from 3' end of replication origin to nearest Sleeping Beauty IR/DR in direction of replication

^b DH5α = DH5α for pUC57-kanR pUC origin, NTC1050811 for R6K origin

^c 500 mL Plasmid+ Shake Flask Culture temperature shifted from 30 to 37C

^d Harvest plasmid preparations with detectable replication intermediates indicated. Levy, *Supra*, 2004 teach that replication intermediates form when any high copy number prokaryotic origin of replication is less than 1000 bp from a promoter/enhancer or LTR promoter or DNA repeat or a complex secondary structure such as an IRES, while replication intermediates do not form if the origin of replication is a distance greater than about 1.5 kb.

Table 7: AAV vectors: pUC versus R6K origin shake flask production yields/quality

AAV Vector	Replication origin	Bacterial selection	Vector Configuration	Vector size	bacterial backbone size (ITR spacing) ^e	Bacterial selection	Replication origin spacing from ITR ^a	Cell line ^b	Production culture ^c	Harvest OD ₆₀₀	Harvest spec yield (mg/L/OD ₆₀₀)	Harvest yield (mg/L)	Production quality ^d
Backbone 1 ^e	pUC origin (pAAV: see Fig 3)	AmpR	ITR <F1Ori-AmpR>pUC> ITR CMV-transgene-pA	5.4 kb	2600 bp (2600 bp)	AmpR	0 bp	Stb14	30-37C shake	17.8	0.4	8	CCC Monomer
	pUC origin (NTC8-AAV: see Fig 3)	RNA-OUT AF	ITR <pUC R-OUT> ITR CMV-transgene-pA	4.0 kb	1240 bp (1240 bp)	RNA-OUT AF	50 bp	Stb14	30-37C shake	10.4	0.7	7	CCC Monomer
	R6K origin (NTC9-AAV: see Fig 3)	RNA-OUT AF	ITR <R6K R-OUT> ITR CMV-transgene-pA	3.3 kb	490 bp (490 bp)	RNA-OUT AF	40 bp	Stb14	30-37C shake	11.8	3.5	42	CCC Monomer
	R6K origin	RNA-OUT AF	ITR <R6K R-OUT> Rep> Cap> ITR CMV-transgene-pA	7.7 kb	490 bp (490 bp)	RNA-OUT AF	40 bp	Stb14	30-37C shake	11.0	2.1	23	CCC Monomer
Backbone 2	pUC origin (NTC8)	RNA-OUT AF	ITR <pUC R-OUT> ITR CAG-transgene-WPRE-pA	5.5 kb	1250 bp (NA)	RNA-OUT AF	50 bp	Stb14	30-37C shake	11.7	0.4	5	CCC Monomer
	R6K origin (NTC9)	RNA-OUT AF	ITR <R6K R-OUT> ITR CAG-transgene-WPRE-pA	4.8 kb	490 bp (NA)	RNA-OUT AF	40 bp	Stb14	30-37C shake	11.1	1.6	18	CCC Monomer

^a Distance in bp from 3' end of replication origin from nearest AAV ITR in direction of replication

^b Stb14 = Stb14 for pUC origin-ampR, NTC1011592 for pUC origin-RNA-OUT, NTC1011641 for R6K origin-RNA-OUT

^c Plasmid+ Shake Flask Culture temperature shifted from 30 to 37C

^d High quality CCC monomer with no detectable replication intermediate and low amounts of recombination mediated ITR deletion. May not see replication intermediates on gel with pUC origin since <50 from DNA structure, but probably formed since plasmid yield lower than R6K origin

^e A R6K origin-RNA-OUT MIP intron version of this vector with 6 bp ITR spacing was unclonable, presumably due to toxicity of juxtaposing the 2 AAV ITR immediately adjacent to each other. A R6K origin-RNA-OUT MIP intron version in a second AAV vector backbone (10 bp ITR spacing) was also unclonable

Table 8: mRNA vector: pUC versus R6K origin DH5α^f HyperGRO^g Fermentation Yields/quality

polyA mRNA expression vector	Replication origin	Selection	Vector Configuration	Vector size	Bacterial backbone size	Replication origin bp from polyA	Harvest OD ₆₀₀	Harvest spec yield (mg/L/OD ₆₀₀)	Harvest spec yield (mg/gWCW)	Harvest yield (mg/L)	Harvest yield (g) ^b	Production quality
pT3/T7 A99 pA ^h mRNA expression vector and retrofit derivatives	pUC origin (pT3/T7 A99 pA)	AmpR	A99<pUC<AmpR T7> Transgene	7.2 kb	2660 bp	170 bp	113	3.8	1.6	430 ^a	4.3	CCC Monomer ^l
	pUC origin + PAS-BH ^d (NTC7 T7 A99 pA)	kanR	A99<KanR pUC>PAS-BH> T7> Transgene	7.2 kb	2660 bp	5000 bp	140	8.9	4.1	1250 ^a	12.5	CCC Monomer
	R6K origin (NTC9 T7 A99 pA)	RNA-OUT AF	A99<R-OUT R6K> T7> Transgene	5.2 kb	610 bp	4600 bp	147	15.3	6.7	2240 ^a	22.4	CCC Monomer
pGEM4Z A60 pA ^c mRNA expression vector and retrofit derivatives	pUC origin (pGEM4Z T7 A60 pA)	AmpR	A60 AmpR> pUC> T7> Transgene	6.8 kb	2500 bp	4400 bp	112	8.3	4.4	929	9.3	CCC Monomer
	pUC origin + PAS-BH ^e (NTC8 T7 A60 pA)	RNA-OUT AF	A60<R-OUT pUC> PAS-BH> T7> Transgene	5.9 kb	1500 bp	4600 bp	146	11.5	4.9	1673 ^a	16.7	CCC Monomer
	R6K origin (NTC9 T7 A60 pA)	RNA-OUT AF	A60<R-OUT R6K> T7> Transgene	4.9 kb	460 bp	4300 bp	113	13.1	5.0	1483 ^a	14.8	CCC Monomer

^a PolyA region correct by sequencing ^b gram plasmid/10 L fermentor ^c See Fig. 4 for vector maps
^d KanR-pUC+ PASBH vector backbone described in Williams 2017 WO2017025447. This pUC+ PASBH backbone dramatically improved plasmid yield when oriented >1.5 kb away from polyA64-polyC31 compared to a pUC origin oriented <200 bp away from the polyA64-polyC31 repeat or a pUC + PASBH origin oriented <400 bp away from the polyA64-polyC31 repeat. (Williams, *Supra*, 2017, P1140-K2 versus P1140 and P1140-K1, Table 1). Thus PAS-BH does not dramatically improve pUC origin replication into a repeat structure <400 bp from the pUC origin.
^e RNA-OUT-pUC+ PASBH vector backbone described in Williams, *Supra*, 2017. This pUC+ PASBH backbone dramatically improved plasmid yield when oriented >1.5 kb away from polyA64-polyC31 compared to a pUC origin oriented <200 bp away from the polyA64-polyC31 repeat or a pUC + PASBH origin oriented <400 bp away from the polyA64-polyC31 repeat. (Williams, *Supra*, 2017, Table 1: P1140-AF2 versus P1140 and P1140-AF1). Thus PAS-BH does not dramatically improve pUC origin replication into a repeat structure <400 bp from the pUC origin.
^f DH5 α = NTC4862 for NTC8; NTC1050811 for NTC9: NTC1050811 DH5 α att_x::P_{5/6 6/6}-RNA-IN- SacB, catR; att_{HK022}::pL (OL1-G to T) P42L-P106I-F107S P113S (P3-), SpecR StrepR; att₈₀₀::pARA-CI857ts, TetR. Arabinose induces pARA-CI857ts expression which reduces copy number through CI857ts mediated downregulation of the Rep protein expressing pL promoter
^g HyperGRO fermentation 30-42°C temperature shift at 55 OD₆₀₀. For NTC9 vectors fermentation media contained 0.2% arabinose
^h Elango et al., 2005. Biochemical and Biophysical Research Communications 330:958-66 teach poly A100 mRNA vectors are unstable in DH5 α and XL1Blue and cannot be produced in mg quantities. See Fig. 4 for vector maps
ⁱ May not see replication intermediates on gel with pUC origin since <170 bp from DNA structure, but probably formed since plasmid yield lower than with R6K origin vectors

Table 9: AAV helper vectors: pUC versus R6K origin plasmid production yields/quality

AAV Helper Vector	Replication origin	Selection	Vector size	bacterial backbone size	Cell line ^b	Production culture ^a	Harvest OD ₆₀₀	Harvest spec yield (mg/L/OD ₆₀₀)	Harvest yield (mg/L)	Production quality
AAV2 Rep/AAV9 Cap (Rep Cap helper)	pUC origin	RNA-OUT AF	5.7 kb	1220 bp	DH5 α	30 to 37C ^a	16.5	2.6	43	CCC Monomer
	R6K origin	RNA-OUT AF	4.9 kb	470 bp	DH5 α	30 to 37C ^a	17.5	5.5	96	CCC Monomer
Fd6 (Ad Helper)	pUC origin	ampR	15.4 kb	2750 bp	Stb14 ^c	30C ^{a, c}	20	0.8 ^d	16 ^d	CCC Monomer and <u>deletion productions</u>
	R6K origin	RNA-OUT AF	12.9 kb	540 bp	DH5 α	30 to 37C ^a	17.8	1.3	23	CCC Monomer
pHelper Rep Cap NP (Ad Helper + Rep Cap single helper plasmid)	R6K origin	RNA-OUT AF	14.2 kb	470 bp	DH5 α	30 to 37C ^a	6.8	4.8	33	CCC Monomer
pHelper (Ad Helper)	pUC origin	ampR	11.6 kb	2340 bp	DH5 α	30-42C HyperGRO shift	129	14.2	1833	CCC Monomer
pHelper-NP (Ad Helper)	R6K origin	RNA-OUT AF	9.8 kb	460 bp	DH5 α	30-42C HyperGRO RAMP ^e	84.5	18.1	1530	CCC Monomer

^a 500 mL Plasmid+ Shake Flask Culture;
^b DH5 α = NTC1050811 for NTC9: NTC1050811 DH5 α att_x::P_{5/6 6/6}-RNA-IN- SacB, catR; att_{HK022}::pL (OL1-G to T) P42L-P106I-F107S P113S (P3-), SpecR StrepR; att₈₀₀::pARA-CI857ts, TetR. Arabinose induces pARA-CI857ts expression which reduces copy number through CI857ts mediated downregulation of the Rep protein expressing pL promoter. Retrofits were obtained using Sucrose + 0.2% Arabinose plates to reduce copy number ^c Fd6 Stb14 best cell line, Fd6/Stb13 see deletions, Fd6/DH5 α and Fd6/XL1Blue cell lines are unstable, used 30C throughout since plasmid lost in 30-37C process
^d Actual yield lower due to high gDNA in minipreps
^e HyperGRO fermentation, inoculated from sucrose + 0.2% arabinose culture grown at 30°C, arabinose added to batch medium to 0.2% final concentration. 30-42°C ramp temperature shift

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US7246693B](#) [\[0004\]](#)
- [WO2014035457A](#) [\[0007\]](#)
- [WO2017086579A](#) [\[0011\]](#)
- [US670844B](#) [\[0014\]](#)
- [WO2008159733A](#) [\[0030\]](#)
- [US20100184159A](#) [\[0030\]](#)
- [US7246693B](#) [\[0030\]](#)
- [US20150275221A](#) [\[0030\]](#)
- [US6977174B](#) [\[0030\]](#)
- [US2009063232A](#) [\[0030\]](#)
- [US7611893B](#) [\[0030\]](#)
- [US622283A](#) [\[0030\]](#)
- [WO200403864A](#) [\[0050\]](#)
- [WO2017026447A](#) [\[0086\]](#)

Non-patent literature cited in the description

- CARNES A.E. *BioProcess Intl*, 2005, vol. 3, 36-44 [\[0004\]](#)
- CHEN ZYHE CYMEUSE LKAY MA *Gene Ther*, 2004, vol. 11, 856-864 [\[0006\]](#)
- SUZUKI MKASAI KSAEKI Y.J *J Virol*, 2006, vol. 80, 3293-3300 [\[0008\]](#)
- LU JZHANG FXU SFIRE AZKAY MA. *Mol Ther*, 2012, vol. 20, 2111-9 [\[0009\]](#)
- QUIVIGER, MARFI AMANSARD DDELACOTTE LPASTOR MSCHERMAN DMARIE C. *Gene Therapy*, 2014, vol. 21, 1001-1007 [\[0008\]](#)
- GAMES et al. *J. Gene Medicine*, 2010, vol. 12, 818-31 [\[0010\]](#)
- RIBEIRO et al. *Cell Reprogram*, 2011, vol. 14, 130- [\[0010\]](#)
- MARIE et al. *J. Gene Medicine*, 2010, vol. 12, 323-32 [\[0010\]](#)
- WANG et al. *J Virology*, 2014, vol. 88, 1924-34 [\[0010\]](#)
- SCHNÖDT et al. *Mol Ther - Nucleic Acids*, 2016, vol. 5, e355- [\[0011\]](#)
- CHADEUF et al. *Molecular Therapy*, 2005, vol. 12, 744-53 [\[0011\]](#)
- KARBOWNICKEZ et al. *Cell and Gene Therapy Insights*, 2017, 731-8 [\[0011\]](#)
- SHARMA et al. *Molecular Therapy Nucleic Acids*, 2013, e74- [\[0011\]](#)
- MONJEZI et al. *Leukemia*, 2017, vol. 31, 186-94 [\[0011\]](#)
- HOLSTEIN et al. *Molecular Therapy*, 2018, [\[0011\]](#)
- PASTOR, MJOHNEN SHARMENING NQUIVIGER MPAILLOUX JKROPP, MWALTER PIVICS ZIZSVAK ZTHUMANN G. *Molecular Therapy*, 2018, vol. 11, 57-67 [\[0012\]](#)
- LU JWILLIAMS JALUKE JZHANG FCHU KKAY MA. *Human Gene Therapy*, 2017, vol. 28, 125-34 [\[0015\]](#)
- KAY MAHE CYCHEN ZY. *Nat Biotechnol*, 2010, vol. 28, 1287-1289 [\[0015\]](#)
- LUKE et al. *Mol Biotechnol*, 2011, vol. 47, 43- [\[0030\]](#)
- SOLAR et al. *Microbiol. Mol. Biol. Rev*, 1998, vol. 62, 434-464 [\[0030\]](#)
- MUTALIK et al. *Nat Chem Biol*, 2012, vol. 8, 447- [\[0030\]](#)
- WILSON IWSIEMERING KRPRASZKIER JPITTARD AJ. *J Bacterial*, 1997, vol. 179, 742-53 [\[0030\]](#)
- WAGNER EGHALTUVIA SROMBY P. *Adv Genet*, 2002, vol. 46, 361-98 [\[0030\]](#)
- FRANCH TGERDES K. *Current Opin Microbiol*, 2000, vol. 3, 159-64 [\[0030\]](#)
- NA DYOY SMCHUNG HPARK HPARK JHLEE SY. *Nat Biotechnol*, 2013, vol. 31, 170-4 [\[0030\]](#)
- MIRKINMIRKIN. *Microbiology and Molecular Biology Reviews*, 2007, vol. 71, 13-35 [\[0030\]](#)
- CHEMA et al. *Nucleic Acids Res.*, 2003, vol. 31, 3497-500 [\[0034\]](#)
- ABHYANKAR et al. *J Biol Chem*, 2003, vol. 278, 45476-45484 [\[0036\]](#)
- SOUBRIER et al. *Gene Therapy*, 1999, vol. 6, 1482-88 [\[0047\]](#)
- WU et al. *J Bacterial.*, 1995, vol. 177, 6338-6345 [\[0049\]](#)
- HEBEL HLCAL YDAVIES LAHYDE SCPRINGLE IAGILL DR. *Mol Ther*, 2008, vol. 16, S110- [\[0049\]](#)
- ABHYANKAR et al. *J Biol Chem*, 2004, vol. 279, 6711-6719 [\[0051\]](#)
- ALLEN et al. *Nucleic Acids Research*, 2011, vol. 39, 7020-33 [\[0077\]](#)
- MARINO MPLUCE MJREISER J. *Methods Mol Biol*, 2003, vol. 229, 43-55 [\[0081\]](#)
- ELANGO et al. *Biochemical and Biophysical Research Communications*, 2005, vol. 330, 958-66 [\[0086\]](#)

Patentkrav

1. Fremgangsmåde til forbedring af replikationen af et kovalent lukket cirkulært plasmid omfattende følgende trin:

- a. tilvejebringelse af et kovalent lukket cirkulært plasmid omfattende:
- 5 i. et Pol I-afhængigt replikationsorigin, og
- ii. et insert omfattende en struktureret DNA-sekvens valgt fra gruppen bestående af inverteret gentagelsessekvens, direkte gentagelsessekvens, homopolymer gentagelsessekvens eller eukaryotisk replikationsorigin, hvor den strukturerede DNA-sekvens
- 10 er placeret i en afstand på mindre end 1000 bp fra det Pol I-afhængige replikationsorigin i replikationsretningen; og
- b. modificering af det kovalent lukkede cirkulære plasmid af a. således at det Pol I-afhængige replikationsorigin erstattes med et R6K-gamma-replikationsorigin, hvorved det resulterende R6K-gammareplikations-origin
- 15 kovalent lukkede cirkulære plasmid har forbedret replikation valgt fra gruppen bestående af reduceret produktion af replikationsmellemprodukter og øget plasmidkopiantal i forhold til det kovalent lukkede cirkulære plasmid af a, hvor R6K-gamma-replikationsoriginet ikke kræver Pol I, og hvor den strukturerede DNA-sekvens er placeret i en afstand på mindre
- 20 end 1000 bp fra R6K gamma-replikationsoriginet i replikationsretningen.

2. Fremgangsmåden ifølge krav 1, hvor det Pol I-afhængige replikationsorigin er valgt fra gruppen bestående af pUC-origin, pMB1-origin og ColE1-origin.

25 **3.** Fremgangsmåden ifølge krav 1, hvor R6K-gamma-replikationsstartstedet er et R6K-gamma-replikationsorigin med mindst 95% sekvensidentitet med en sekvens valgt fra gruppen bestående af SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 og SEQ ID NO: 18.

30 **4.** Fremgangsmåden ifølge krav 1, hvor den strukturerede DNA-sekvens er:

a. en homopolymer gentagelse, og hvor den homopolymere gentagelse er en polyA-gentagelse;

b. et eukaryotisk replikationsorigin, og hvor det eukaryotiske replikationsorigin er et SV40-replikationsorigin;

7. Fremgangsmåden ifølge krav 5, hvor R6K-gamma-replikationsoriginet er et R6K-gamma-replikationsorigin med mindst 95% sekvensidentitet med en sekvens valgt fra gruppen bestående af SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 og SEQ ID NO: 18.

5

8. Fremgangsmåden ifølge krav 5, hvor den RNA-selektive markør er en RNA-IN regulerende RNA-OUT funktionel variant med mindst 95% sekvensidentitet med en sekvens valgt fra gruppen bestående af SEQ ID NO: 5 og SEQ ID NO: 7.

10 **9.** Fremgangsmåden ifølge krav 5, hvor den RNA-selektive markør er en RNA-OUT RNA-selektiv markør, der koder for et RNA-IN regulerende RNA-OUT-RNA med mindst 95% sekvensidentitet med SEQ ID NO: 6.

10. Fremgangsmåden ifølge krav 5, hvor den bakterielle replikations-
15 selektionsregion omfattende et Pol I-afhængigt replikationsorigin og en antibiotisk selektiv markør er erstattet med en Pol III-afhængig R6K origin-RNA-OUT RNA-selektiv markør bakteriel replikations-selektionsregion med mindst 95% sekvensidentitet med en sekvens valgt fra gruppen bestående af SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13
20 SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16 og SEQ ID NO: 17.

11. Fremgangsmåden ifølge krav 5, hvor den strukturerede DNA-sekvens er:

- a. en homopolymer gentagelse, og hvor den homopolymere gentagelse er en polyA-gentagelse;
- 25 b. et eukaryotisk replikationsorigin, og hvor det eukaryotiske replikationsorigin er et SV40-replikationsorigin;
- c. en direkte gentagelsessekvens, og hvor den direkte gentagelsessekvens er valgt fra gruppen bestående af viral LTR, en lentiviral LTR eller en retroviral LTR; eller
- 30 d. en inverteret gentagelsessekvens, og hvor den inverterede gentagelsessekvens er valgt fra gruppen bestående af en transposon IR/DR-gentagelse, en Sleeping Beauty transposon IR/DR-gentagelse, en AAV ITR, en transposon ITR eller en PiggyBac transposon ITR.

35

DRAWINGS

FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D




FIG. 4E





FIG. 4F



SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

