

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
16 June 2005 (16.06.2005)

PCT

(10) International Publication Number
WO 2005/054479 A1

- (51) International Patent Classification⁷: C12N 15/82, (74) Agents: GAAL, Jozsef, Christopher et al.; Intellectual Property Dept., Syngenta Limited, Jealott's Hill International Research Centre, P.O. Box 3538, Bracknell, Berks RG42 6YA (GB).
C12Q 1/68, A01H 5/00
- (21) International Application Number: PCT/EP2004/012662
- (22) International Filing Date: 9 November 2004 (09.11.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/526,112 1 December 2003 (01.12.2003) US
- (71) Applicant (for all designated States except US): SYNGENTA PARTICIPATIONS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): NEGROTTO, David, Vincent [US/US]; Syngenta Biotechnology, Inc., 3054 Cornwallis Road, Research Triangle Park, North Carolina, NC 27709 (US). SHOTKOSKI, Frank, Arthur [US/US]; Syngenta Biotechnology, Inc., 3054 Cornwallis Road, Research Triangle Park, North Carolina, NC 27709 (US). YU, Wenjin [CA/US]; Syngenta Biotechnology, Inc., 3054 Cornwallis Road, RTP, North Carolina, NC 27709 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2005/054479 A1

(54) Title: INSECT RESISTANT COTTON PLANTS AND METHODS OF DETECTING THE SAME

(57) Abstract: The present application relates an insect resistant transgenic cotton plant. In particular, it relates to a specific event, designated COT202. The application also relates to polynucleotides which are characteristic of the COT202 event, plants comprising said polynucleotides, and methods of detecting the COT202 event.

INSECT RESISTANT COTTON PLANTS AND METHODS OF DETECTING THE
SAME

- 5 The present invention relates to genetic engineering of plants and in particular to an insect resistant transgenic cotton plant. Specifically, the invention relates to a cotton plant designated COT202 which comprises a VIP3A gene. It also relates to methods of detecting material derived from the plant.
- 10 Plant pests are a major factor in the loss of the world's important agricultural crops. About \$8 billion is lost every year in the U.S. due to infestations of plants by non-mammalian pests including insects. In addition to losses in field crops, insect pests are also a burden to vegetable and fruit growers, to producers of ornamental flowers, and to home gardeners.
- 15 Insect pests are mainly controlled by intensive applications of chemical pesticides, which are active through inhibition of insect growth, prevention of insect feeding or reproduction, or cause death. Good control of insect pests can thus be reached, but these chemicals can sometimes also affect other, beneficial insects. Another problem resulting from the wide use of chemical pesticides is the appearance of resistant insect varieties. This has been
- 20 partially alleviated by various resistance management practices, but there is an increasing need for alternative pest control agents. Biological pest control agents, such as *Bacillus thuringiensis* strains expressing pesticidal toxins like δ -endotoxins, have also been applied to crop plants with satisfactory results, offering an alternative or compliment to chemical pesticides. The genes coding for some of these δ -endotoxins have been isolated and their
- 25 expression in heterologous hosts has been shown to provide another tool for the control of economically important insect pests. In particular, the expression of insecticidal toxins such as *Bacillus thuringiensis* δ -endotoxins in transgenic plants, has provided efficient protection against selected insect pests, and transgenic plants expressing such toxins have been commercialised, allowing farmers to reduce applications of chemical insect control
- 30 agents.

Recently, a new family of insecticidal proteins produced by *Bacillus* sp. during the vegetative stages of growth (vegetative insecticidal proteins (VIPs)) has been identified.

U.S. Patents 5,877,012, 6,107,279, and 6,137,033 describe *vip3A* toxin genes isolated from *Bacillus* species. The VIP3A toxins possess insecticidal activity against a wide spectrum of lepidopteran insects including but not limited to fall armyworm, *Spodoptera frugiperda*, black cutworm, *Agrotis ipsilon*, sugarcane borer, *Diatraea saccharalis*, and lesser cornstalk borer, *Elasmopalpus lignosellus*, and when expressed in transgenic plants, for example cotton, confer protection on the plant from insect feeding damage.

The cotton family, genus *Gossypium*, a member of the Malvaceae, consists of 39 species, of which *Gossypium hirsutum* is the most commonly cultivated species. Three other species are also cultivated: *G. arboreum*, *G. barbadense*, and *G. herbaceum*. These cultivated species are grown primarily for the seed hairs that are made into textiles. Cotton is suitable as a textile fibre because the mature dry hairs twist in such a way that fine strong threads can be spun from them. Other products, such as cottonseed oil, cake, and cotton linters are by-products of fibre production.

Damage to cotton crops by insect pests throughout the world results in a significant yield loss each year. Effective control of these pests to minimise yield loss is of great economic importance. Examples of insect pests of cotton include Beet armyworm (*Spodoptera exigua*), Boll weevil (*Anthonomus grandis grandis*), Cabbage looper (*Trichoplusia ni*), Clouded plant bug (*Neurocolpus nubilus*), Cotton aphid (*Aphis gossypii*), Cotton bollworm (*Heliocoverpa zea*), Cutworms (*Feltia subterranea*, *Peridroma saucia*, *Agrotis ipsilon*), European corn borer (*Ostrinia nubilalis*), Fall armyworm (*Spodoptera frugiperda*), Seedling thrips (*Frankliniella* spp.), Soybean looper (*Pseudoplusia includens*), Stink bugs (*Nezara viridula*, *Acrosternum hilare*, *Euschistus servus*), Tarnished plant bug (*Lygus lineolaris*), Tobacco budworm (*Heliothis virescens*) and Whiteflies (*Trialeurodes abutilonea*, *Bemisia tabaci*).

Transformation and regeneration of cotton plants is now a well-established procedure, typically based on *Agrobacterium tumefaciens* mediated transfer of foreign DNA into cotton plant parts and regeneration of said plant parts in tissue culture into fully fertile, transgenic cotton plants.

There exists a requirement to generate a cotton plant that is insect resistant so that yield loss through damage to cotton crops by insect pests is reduced. An insect resistant cotton plant could reduce the need to apply chemical pesticides, which may be detrimental to other, beneficial insects and the environment. Further, it is desirable to provide an insect resistant plant that comprises a VIP gene, as an alternative to transgenic plants comprising crystal proteins from *Bacillus thuringiensis*. This may be of use in insect resistance management.

Therefore, the present invention relates to an insect resistant transgenic cotton event, designated COT202. It also relates to methods of detecting plant material derived therefrom. "COT202 event" in the context of this application refers to the original insecticidal transgenic cotton plant described herein. "Insecticidal" as used herein refers to any inhibitory effect on an insect, including but not limited to reduced feeding, retarded growth, reduced fecundity, paralysis or death. "Fecundity" comprises all aspects related to reproduction such as reproductive ability, reproductive frequency and number of offspring. Also embraced by this invention is any plant material derived from the COT202 event, including seeds.

The COT202 event exhibits a novel genotype comprising at least one expression cassette. The cassette comprises a suitable promoter for expression in plants operably linked to a gene that encodes a VIP3A insecticidal toxin, useful in controlling a wide spectrum of lepidopteran insect pests, and a suitable polyadenylation signal. Suitable promoters may be isolated from, *inter alia*, plants. Numerous plant promoters have been isolated and characterised including constitutive, switchable and/or tissue specific promoters. Suitable promoters may be selected from the following, non-limiting group: CaMV35S, FMV35S, Ubiquitin, Act2, NOS, OCS, Cestrum yellow leaf curl virus promoter, Patatin, E9, alcA/alcR switch, GST switch, RMS switch, oleosin, Gelvin, ribulose bisphosphate carboxylase-oxygenase small sub-unit, actin 7, MR7 promoter (maize), Gos 9 (rice), GOS2 promoters, MasOcs (or super promoter), RolD promoter (*Agrobacterium rhizogenes*), SuperMAS promoter, and Suc2 promoter (*Arabidopsis*). In one embodiment of the present invention, the promoter is the Ubiquitin promoter, UBQ3, from *Arabidopsis thaliana*. Additional elements such as enhancer sequences may also be incorporated into the expression cassette in order to boost levels of gene expression, for example transcriptional

or translational enhancers, such as tobacco etch virus (TEV) translation activator, CaMV35S enhancer, and FMV35S enhancer. Alternatively it may be desirable to include a targeting sequence, for example, to direct transportation of the VIP3A toxin to a particular cellular compartment. For example if it is desired to provide the protein outside
5 of the cell then an extracellular targeting sequence may be ligated to the polynucleotide encoding the VIP protein. Other examples of targeting include targeting to a specific intracellular organelle or compartment, for example to the endoplasmic reticulum using a 'KDEL' retention sequence. Numerous polyadenylation signals have been isolated and characterised. Examples of suitable polyadenylation signals functional in plants include
10 that from the nopaline synthase gene (nos) of *Agrobacterium tumefaciens*, from the proteinase inhibitor II gene and from the alpha-tubulin gene (EP-A 652,286). In one embodiment of the present invention, the polyadenylation signal is that from the nos gene of *Agrobacterium tumefaciens*.

15 According to the invention, the polynucleotide encoding the VIP3A protein may also be codon-optimised or otherwise altered to enhance for example, transcription once it is incorporated into plant material. Such codon optimisation may also be used to alter the predicted secondary structure of the RNA transcript produced in any transformed cell, or to destroy cryptic RNA instability elements present in the unaltered transcript, thereby
20 increasing the stability and/or availability of the transcript in the transformed cell (Abler and Green (1996) Plant Molecular Biology (32) pp.63-78).

In a precursor to the COT202 event, a second cassette is present that comprises a gene which, when expressed, can be used as a selectable marker. Numerous selectable markers
25 have been characterised, including some that confer tolerance to antibiotics and others that confer tolerance to herbicides. Examples of suitable selectable marker genes include those that confer tolerance to hygromycin, kanamycin or gentamycin. Further suitable selectable markers include genes that confer resistance to herbicides such as glyphosate-based herbicides or resistance to toxins such as eutypine. Other forms of selection are also
30 available such as hormone based selection systems such as the Multi Auto Transformation (MAT) system of Hiroyrasu Ebinuma *et al.* (1997) PNAS Vol. 94 pp.2117-2121; visual selection systems which use the known green fluorescence protein, β glucuronidase; and any other selection system such as mannose isomerase (PositechTM), xylose isomerase and

2-deoxyglucose (2-DOG). In one embodiment of the present invention, the selectable marker gene is one that confers tolerance to hygromycin. This second expression cassette is useful for selecting transformants during and following plant transformation.

Optionally, it may be segregated away from the COT202 event precursor after
5 transformation to leave the COT202 event itself. The COT202 event *per se* does not comprise a selectable marker cassette. Further expression cassettes are optionally comprised in the COT202 event. For example these may provide other desirable benefits such as herbicide resistance.

10 The expression cassettes may be introduced into the plant on the same or different plasmids. If the expression cassettes are present on the same plasmid and introduced into the plant via an *Agrobacterium*-mediated transformation method, they may be present within the same or different T-DNA regions. In one embodiment of the present invention, two expression cassettes are present on different T-DNA regions within the same plasmid.

15 According to the first aspect of the invention, there is provided a polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 1. In one embodiment said polynucleotide comprises at least 18 contiguous nucleotides from SEQ ID NO: 1. In a further embodiment said polynucleotide comprises
20 at least 20 contiguous nucleotides from SEQ ID NO: 1. In a still further embodiment said polynucleotide comprises at least 22 contiguous nucleotides from SEQ ID NO: 1. In a further embodiment said polynucleotide comprises at least 23 contiguous nucleotides from SEQ ID NO: 1. In yet a further embodiment said polynucleotide comprises at least 24 contiguous nucleotides from SEQ ID NO: 1. In a further embodiment said polynucleotide
25 comprises at least 25 contiguous nucleotides from SEQ ID NO: 1. In a still further embodiment there is provided a polynucleotide comprising the sequence of SEQ ID NO: 1.

In a further aspect of the invention, there is provided a polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 2. In one
30 embodiment said polynucleotide comprises at least 18 contiguous nucleotides from SEQ ID NO: 2. In a further embodiment said polynucleotide comprises at least 20 contiguous nucleotides from SEQ ID NO: 2. In a still further embodiment said polynucleotide comprises at least 22 contiguous nucleotides from SEQ ID NO: 2. In a further

embodiment said polynucleotide comprises at least 23 contiguous nucleotides from SEQ ID NO: 2. In yet a further embodiment said polynucleotide comprises at least 24 contiguous nucleotides from SEQ ID NO: 2. In a further embodiment said polynucleotide comprises at least 25 contiguous nucleotides from SEQ ID NO: 2. In a still further
5 embodiment there is provided a polynucleotide comprising the sequence of SEQ ID NO: 2.

In a further aspect of the present invention there is provided a polynucleotide as described above further comprising the sequence of SEQ ID NO: 7. In a further aspect of the present invention there is provided a polynucleotide as described above further comprising the
10 sequence of SEQ ID NO: 8.

In another aspect of the present invention there is provided a plant comprising a polynucleotide which comprises at least 17 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. In one embodiment said plant comprises at least 18 contiguous
15 nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. In a further embodiment said plant comprises at least 20 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. In a further embodiment said plant comprises at least 22 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. In a further embodiment said plant comprises at least 23 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. In a still further
20 embodiment said plant comprises at least 24 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. In a further embodiment said plant comprises at least 25 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2. In yet a further embodiment said plant comprises the sequence of SEQ ID NO: 1 and / or SEQ ID NO: 2. In a further embodiment, said plant additionally comprises the sequence of SEQ ID NO: 7. In a further
25 embodiment still, said plant additionally comprises the sequence of SEQ ID NO: 8. In one embodiment of the present invention, said plant is a cotton plant. In a further embodiment, said plant is an insecticidal cotton plant which is the COT202 event, or a plant derived therefrom.

30 The skilled man is familiar with plant transformation methods. In particular, two principal techniques have been characterised across a wide range of plant species: transformation by *Agrobacterium* and transformation by direct DNA transfer.

Agrobacterium-mediated transformation is a commonly used method for transformation of dicotyledonous plants. The foreign DNA to be introduced into the plant is cloned into a binary vector in between left and right border consensus sequences. This is the T-DNA region. The binary vector is transferred into an *Agrobacterium* cell, which is subsequently used to infect plant tissue. The T-DNA region of the vector comprising the foreign DNA is inserted into the plant genome. The marker gene cassette and trait gene cassette may be present on the same T-DNA region, different T-DNA regions in the same vector, or even different T-DNA regions in different vectors. In one embodiment of the present invention, the cassettes are present on different T-DNA regions in the same vector.

10

Alternatively, direct DNA transfer can be used to introduce the DNA directly into a plant cell. One suitable method of direct transfer may be bombardment of plant cells with a vector comprising the DNA for insertion using a particle gun (particle-mediated biolistic transformation); another established method, 'whiskers', involves coating the DNA onto silicon carbide fibres onto which cells are impaled. Other methods for transforming plant cells include protoplast transformation (optionally in the presence of polyethylene glycols); sonication of plant tissues, cells or protoplasts in a medium comprising the polynucleotide or vector; micro-insertion of the polynucleotide or vector into plant material (optionally employing the known silicon carbide "whiskers" technique), electroporation and the like.

20

Following transformation, transgenic plants must be regenerated from the transformed plant tissue, and progeny possessing the foreign DNA selected using an appropriate marker such as resistance to hygromycin. The skilled man is familiar with the composition of suitable regeneration media. The selectable marker can be segregated away from transgenic events by conventional plant breeding methods, thus resulting in, for example, the COT202 event.

25

A plant of the invention, as described herein, has an insecticidal effect on insects from one or more species from the group comprising *Heliothis* sp., *Helicoverpa* sp. and *Spodoptera* sp. which may infest it. "Infest" as used herein refers to attack, colonisation, feeding or damage in any way by one or more insects. Thus, for example, the plant of the present invention will provide a self-defence mechanism against infestation by pest insects such as *Helicoverpa zea* (cotton boll worm). As a result, a reduced number of insecticide sprays

30

are required during the cultivation of said plant compared to a non-transgenic cotton plant of the same variety and yield loss through insect pests is kept at a minimal level.

The present invention is not limited to the COT202 event itself, but is further extended to include any plant material derived therefrom, including seeds in so far as they contain at least one of the present inventive polynucleotides. The present invention includes, but is not limited to plants that are derived from a breeding cross with the COT202 event or a derivative therefrom by conventional breeding or other methods. The invention also includes plant material derived from the COT202 event that may comprise additional, modified or fewer polynucleotide sequences compared to the COT202 event or exhibit other phenotypic characteristics. For example it may be desirable to transform plant material derived from the COT202 event to generate a new event that possesses an additional trait, such as a second insect resistance gene. This process is known as gene stacking. The second insect resistance gene may encode, for example insecticidal lectins, insecticidal protease inhibitors and insecticidal proteins derived from species of the *Bacillus thuringiensis*, *Xenorhabdus nematophilus*, or *Photorhabdus luminescens*. In one aspect, the second insect resistance gene encodes an insecticidal gene from *Bacillus thuringiensis*. Preferably, the second insect resistance gene encodes a Cry gene from the bacterium *Bacillus thuringiensis*, which Cry gene produces a toxin with a different mode of action or binding site in the insect gut to VIP for the control of different insect species. The Cry gene may, for example, be Cry1Ab.

The present invention further provides plant material derived from the COT202 event which possesses an additional trait such as herbicide resistance, nematode resistance or fungal resistance. In one embodiment, said additional trait is herbicide resistance. The herbicide resistance trait may be provided, for example, by a herbicide degradation enzyme, or a target-site specific resistant enzyme. In a further embodiment, said herbicide resistance trait provides resistance to a herbicide which comprises glyphosate acid or an agriculturally acceptable salt thereof. In a further embodiment still, said herbicide resistance trait is provided by a gene encoding EPSP synthase or a mutant thereof.

The present invention further provides a method of controlling insects comprising providing the COT202 event or plant material derived from the COT202 event at a locus

where said insects feed. The invention yet further provides a method of controlling insects comprising providing the COT202 event or plant material derived from the COT202 event at a locus where said insects feed, and applying other agrochemicals to said plant material such as herbicides, fungicides and other insecticidal compounds including other
5 insecticidal proteins. Examples of possible insecticidal compounds include insecticidal lectins, insecticidal protease inhibitors and insecticidal proteins derived from species of the *Bacillus thuringiensis*, *Xenorhabdus nematophilus*, or *Photorhabdus luminescens*. Examples of possible chemicals include pyrethroids, carbamates, imidacloprid, organochlorines, and macromolecules such as spinosad, abamectin or emamectin.

10

According to yet a further aspect of the present invention, there is provided a method of detecting the COT202 event or plant material derived from the COT202 transgenic event comprising obtaining a sample for analysis; extracting DNA from the sample; providing a pair of primers designed to bind to a polynucleotide comprising at least 17 contiguous
15 nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2; amplifying the region which lies between the sites at which the primers bind; and detecting the presence of the amplification product. Suitable pairs of primers for use in this method of detection can be designed using parameters well known to those skilled in the art of molecular biology now that SEQ ID NOs 1 and 2 are made available. For example, one or both primers of the pair may be
20 designed to be vector-specific, trait gene specific, promoter specific, and/or specific to the sequence of the junction between the inserted DNA and the genomic DNA. Preferably one of the primers is designed to be specific to the inserted sequence, and the other primer specific to the genomic DNA upstream or downstream of the insertion site. In one embodiment, the sequence of said primers is depicted as SEQ ID NO: 3 and SEQ ID NO:
25 4.

In an embodiment of the present invention, the region amplified by said method (the 'amplicon') is between 100 and 1000 base pairs in length. In a further embodiment the amplicon is between 100 and 400 base pairs in length. In a still further embodiment the
30 amplicon is 181 base pairs in length. In a further embodiment the amplicon is produced using the above method in conjunction with the primers of the sequence of SEQ ID NO: 3 and SEQ ID NO: 4, and is 181 base pairs in length. These primers are specific for the COT202 event.

Alternative primers which may be used in combination to detect the COT202 event include SEQ ID NOs 13 and 14 which are specific for the COT202 event and produce an 86 bp amplicon, and SEQ ID NOs 5 and 6 which are specific for the VIP gene and produce a
5 556bp amplicon.

There are many amplification methods that may be used in accordance with this aspect of the invention. The underlying principle, a known technique to those skilled in the art, is the polymerase chain reaction (PCR). The amplification product from a PCR reaction may
10 be visualised by staining with ethidium bromide and excitation with UV light, typically after size separation using agarose gel electrophoresis.

An embodiment of the present invention employs variations of the PCR principle such as TaqMan™. This involves labelling at least one of the primers involved in the
15 amplification process with a fluorescent dye. When unbound, the primer adopts a conformation such that no fluorescence can be detected. However, when the primer is bound to a piece of DNA, the conformation changes and fluorescence can be detected. In this way, the amplification process can be monitored in real-time, the intensity of fluorescence corresponding directly to the level of amplification. Suitable primers for use
20 in TaqMan™ PCR are depicted as SEQ ID NOs 13 to 15. These may be used in conjunction with internal control primers such as those depicted as SEQ ID NOs 10 to 12. TaqMan™ analysis may be useful for example, for detecting the presence of the COT202 event in a background of wild type cotton, or for detecting the adventitious presence of COT202 in other germplasm. Further embodiments of the present invention include, but
25 are not limited to, RACE PCR.

A further embodiment of the present invention involves the use of multiplex PCR for distinguishing between homozygous COT202 plant material and heterozygous COT202 plant material. This is known to those skilled in the art as zygosity testing, and involves
30 the use of three PCR primers which bind to specific parts of the cotton genome and / or inserted DNA. The presence or absence of each of two amplification products of particular sizes indicates whether the test sample is heterozygous or homozygous for COT202. Suitable primers for use in such a zygosity test are depicted as SEQ ID NOs 16 to 18.

In another aspect of the invention there is provided a method of detecting plant material derived from the COT202 event comprising obtaining a sample for analysis; providing a probe designed to bind to the complement of a polynucleotide which comprises at least 17 contiguous nucleotides of SEQ ID NO: 1 and / or SEQ ID NO: 2 when said polynucleotide is single stranded; hybridising said probe with the sample; and detecting whether the probe has hybridised. In one embodiment, said probe comprises the sequence of SEQ ID NO: 1 and / or SEQ ID NO: 2. In an embodiment of the present invention there is provided a method of detecting plant material derived from the COT202 event using a probe comprising SED ID NO: 7 or SEQ ID NO: 8. In one embodiment, said probe comprises SEQ ID NO: 7. In a further embodiment, said probe consists of SEQ ID NO: 7. In one embodiment, said probe comprises SEQ ID NO: 8. In a further embodiment, said probe consists of SEQ ID NO: 8. The probe may be, for example, a PCR product or restriction digestion fragment. In a further embodiment, the probe as described herein may be tagged with a fluorescent, radioactive, enzymatic or other suitable label to enable hybridisation to be detected. The skilled man will know how to design suitable probes, now that he has the benefit of the present disclosure.

In a further embodiment of the present invention, there is provided a method of hybridising a probe to the sample under stringent conditions and detecting whether the probe has hybridised. Stringent hybridisation conditions are well known to the skilled man and comprise, for example: hybridisation at a temperature of about 60°C in a solution containing 6 x SSC, 0.01% SDS and 0.25% skimmed milk powder, followed by rinsing at the same temperature in a solution containing 1 x SSC and 0.1% SDS. More stringent hybridisation conditions may comprise: hybridisation at a temperature of about 65°C in a solution containing 6 x SSC, 0.01% SDS and 0.25% skimmed milk powder, followed by rinsing at the same temperature in a solution containing 0.2 x SSC and 0.1% SDS.

Suitable techniques for detecting plant material derived from the COT202 event based on the hybridisation principle include, but are not limited to Southern Blots, Northern Blots and *in-situ* hybridisation. The skilled man is familiar with techniques such as these. Typically, they involve incubating a probe with a sample, washing to remove unbound probe, and detecting whether the probe has hybridised. Said detection method is

dependent on the type of tag attached to the probe – for example, a radioactively labelled probe can be detected by exposure to and development of x-ray film. Alternatively, an enzymatically labelled probe may be detected by conversion of a substrate to effect a colour change.

5

In a further aspect of the invention there is provided a method of detecting plant material derived from the COT202 event comprising obtaining a sample for analysis; providing an antibody or binding protein designed to bind to a VIP protein contained within a plant comprising at least 17 contiguous nucleotides from SEQ ID NO: 1 and / or SEQ ID NO: 2;
10 incubating said antibody or binding protein with the sample; and detecting whether the antibody or binding protein has bound. In one embodiment of the present invention said VIP protein comprises the sequence of SEQ ID NO: 9.

Suitable methods of detecting plant material derived from the COT202 event based on said
15 antibody binding include, but are not limited to Western Blots, Enzyme-Linked ImmunoSorbent Assays (ELISA) and SELDI mass spectrometry. The skilled man is familiar with these immunological techniques. Typical steps include incubating a sample with an antibody that binds to the VIP protein, washing to remove unbound antibody, and detecting whether the antibody has bound. Many such detection methods are based on
20 enzymatic reactions - for example the antibody may be tagged with an enzyme such as horse radish peroxidase, and on application of a suitable substrate, a colour change detected. Suitable antibodies may be monoclonal or polyclonal.

In another aspect of the invention there is provided a method of detecting plant material
25 derived from the COT202 event comprising obtaining a sample for analysis; making a protein extract of the sample; providing a test strip designed to detect the presence of a VIP protein present within the sample; incubating the test strip with the sample; and detecting whether VIP protein is present. In one embodiment of the present invention said VIP protein comprises the sequence of SEQ ID NO: 9.

30

An alternative antibody-based detection method for COT202 uses of dipsticks or test strips. Typical steps include incubating a test strip with a sample and observing the presence or absence of coloured bands on the test strip. The coloured bands are indicative

of the presence of a protein in the sample. Such dipstick or test strip tests are protein specific, and may be used for rapid testing of samples in the field.

In a further aspect of the present invention there is provided a method of detecting plant material derived from the COT202 event comprising obtaining a sample for analysis;
5 subjecting one or more insects of the species *Spodoptera frugiperda* (susceptible to VIP3A) to the sample; subjecting one or more insects of species *Ostrinia nubilalis* (not susceptible to VIP3A) to the sample as a control; detecting whether the sample has an insecticidal effect on insects from each species; and comparing the results with an
10 authentic COT202 bioassay profile. The results are compared against an authentic COT202 bioassay profile that is produced using insects of the same condition (including insect age and culture conditions) which have been subjected to the same dose and type of COT202 plant material (including plant age, plant variety and tissue type) and where the insecticidal effect is detected the same length of time after subjecting the insects to the
15 COT202 sample. Detection of an insecticidal effect may be, for example, an assessment of insect mortality, or of the growth stage of the insects. *Spodoptera frugiperda* is a positive control for COT202 as it is susceptible to a suitable dose of VIP3A, while *Ostrinia nubilalis* is a negative control for COT202 as it is not susceptible to a suitable dose of VIP3A. Alternative insect species that are either susceptible or not susceptible to VIP3A
20 may be substituted in an assay as described above as appropriate, provided that the results are compared against an authentic profile generated using the same insect species.

In one embodiment of the invention, the method of detecting plant material derived from the COT202 event includes but is not limited to leaf-feeding bioassays in which a leaf or
25 other suitable plant part from the COT202 event or any plant material derived from the COT202 event, is infested with one or more pest insects. Detection may be through assessment of damage to the leaf or plant part after set time periods, assessment of mortality or another insecticidal effect on the insects. Alternative plant parts which may be used for such bioassays include bolls and squares. Such bioassays may, for example, be
30 carried out in the laboratory, field, or glasshouse, and may be subject to natural or artificial insect infestation.

In another aspect of the invention, there is provided a kit of parts comprising a means for detecting the presence in a sample of plant material derived from the COT202 event.

Preferably, said kit of parts comprises a means for detecting the presence in a sample of a polynucleotide comprising at least 17 contiguous nucleotides from the sequence of SEQ ID NO: 1 and / or SEQ ID NO: 2, or a protein encoded by a polynucleotide as described above, or a VIP protein. In an embodiment of the present invention, said kit of parts may comprise DNA amplification-detection technology such as PCR or TaqMan™. In a further embodiment of the present invention, said kit of parts may comprise probe hybridisation-detection technology such as Southern Blots, Northern Blots or *in-situ* Hybridisation. In another embodiment of the present invention, said kit of parts may comprise antibody binding-detection technology such as Western Blots, ELISA's, SELDI mass spectrometry or test strips. In a further embodiment of the present invention, said kit of parts may comprise insect bioassay-detection technology such as leaf feeding bioassays or mortality bioassays. Each of these detection technologies may be used as described above. In a further embodiment of the present invention, said kit of parts may comprise any combination of the afore-mentioned detection technologies. In a still further embodiment, said kit of parts may comprise in the form of instructions one or more of the methods described above.

According to the present invention, there is provided the use of one or more of the polynucleotides of the invention as described above for detecting the COT202 event. In one embodiment, said polynucleotides may be used in a method for detecting the COT202 event as described above.

EXAMPLES

The invention will be further apparent from the following non-limiting examples in conjunction with the associated sequence listings as described below:

SEQ ID NO 1: Polynucleotide sequence which extends across the junction where the 5' end of the COT202 insert is inserted into the cotton genome in event COT202.

- SEQ ID NO 2: Polynucleotide sequence which extends across the junction where the 3' end of the COT202 insert is inserted into the cotton genome in event COT202.
- 5 SEQ ID NOs 3 - 6: Polynucleotide sequences suitable for use as primers in the detection of the COT202 event.
- SEQ ID NOs 7 - 8: Polynucleotide sequences suitable for use as probes in the detection of the COT202 event.
- SEQ ID NO 9: Amino acid sequence of the VIP3A toxin protein.
- 10 SEQ ID NOs 10 - 15: Polynucleotide sequences suitable for use as TaqMan™ primers in the detection of the COT202 event.
- SEQ ID NOs 16 - 18: Polynucleotide sequences suitable for use as primers in the detection of the COT202 event via zygosity testing.

Example 1: Cloning and Transformation

15 *1.1 Vector cloning*

Standard gene cloning techniques of restriction digestion and ligation of fragments from in-house vectors were used to construct the transformation vector, pNOV103. The vector included a selectable marker cassette comprising a Ubiquitin (UBQ3) promoter, the UBQ3 intron, a gene sequence which encodes a protein conferring resistance to hygromycin, and a nos polyadenylation sequence. The vector also included the expression cassette of the target gene, which cassette comprised a Ubiquitin (UBQ3) promoter, the UBQ3 intron, a sequence encoding the VIP3A gene that had been codon optimised for expression in maize, and a nos polyadenylation sequence. The selectable marker cassette and VIP3A containing cassette were cloned between the left and right border sequences within different T-DNA regions of vector pNOV103. The vector also comprised a gene conferring resistance to an antibiotic, kanamycin, for prokaryotic selection.

20

25

The vector was transformed into *Agrobacterium tumefaciens* strain GV3101 using standard *Agrobacterium* transformation techniques, and transformed cells selected through their resistance to kanamycin.

30

1.2 Plant transformation

The COT202 event was produced by *Agrobacterium*-mediated transformation of *Gossypium hirsutum* L. cv Coker 312.

5 Coker 312 seeds were sown in the glasshouse. Tender petioles were cut from 3 to 5 weeks old plants, and sterilized by immersion in 70% ethanol. The petioles were then immersed in a 5% Clorox + 2ml/L Tween 20 solution for 20 minutes. Petioles were washed 3 times in ddH₂O. The ends of petioles were cut off, and petioles transferred to petiole pre-culture medium (4.3g/L MS salts, 200X B5 vitamins, 30g/L glucose, 2.4g/L phytoigel, pH 7.0) and allowed to pre-culture in the light at 30°C for 3 days.

10

2 ml cultures of *Agrobacterium* containing the pNOV103 construct were grown overnight in appropriate antibiotics and then diluted with liquid MMS1 medium (4.3g/L MS salts, 200X B5 vitamins, 0.05mg/L 2,4-D, 0.1mg/L kinetin, 30g/L glucose, pH 6.5) to an OD₆₆₀ of between 0.1 and 0.2.

15

The ends were cut off the petioles and placed in 3 to 5ml of bacterial solution in a sterile petri dish. Once in the solution, the petioles were cut lengthwise and then cut into 2cm sections. After the petiole explants had soaked in bacterial solution for 5 to 10 minutes, they were transferred to co-culture plates, and allowed to co-culture at 24°C for 48 hours 20 under low light intensity. Co-cultured explants were transferred to MMS1 medium (recipe as for MMS1 liquid medium, additionally with 2.4g/L phytoigel) containing 500mg/L Cefotaxime and 10mg/L Hygromycin, and incubated at 30 °C under a light cycle of 16 hours light and 8 hours dark. Explants were transferred to fresh medium after 2 weeks, and every 4 to 6 weeks thereafter until callus was formed.

25

Once calli were the size of a garden pea, they were removed from the explants and transferred to fresh MMS1 medium containing 500mg/L Cefotaxime and 10mg/L Hygromycin, and maintained in tissue culture by subculturing every 4 weeks as appropriate.

30

1.5g callus tissue was broken up thoroughly and placed in a 50 ml Erlenmeyer flask containing 10 ml of liquid MMS2 medium (4.3g/L MS salts, 200X B5 vitamins, 1.9g/L KNO₃, 30 g/L glucose, pH 6.5). The suspended callus was shaken at 100 rpm in the light

at 30°C until small white slightly round cell clusters were visible. These clusters indicate that the tissue is embryogenic. The suspension culture cells were rinsed 3 times in MMS2 liquid medium, resuspended and plated onto solid MMS2 medium (recipe as per liquid MMS2 medium, additionally with 2.4g/L phytoigel). Once plated, excess liquid MMS2 medium was removed, and the plates incubated at 30°C in the light. Plates were checked for somatic embryo development each week. Somatic embryos formed within 1 to 2 months. Somatic embryos were transferred to EG (embryoid germination) medium (10X EG stock (consisting of 1x 10L pack of Musashige and Skoog Modified Basal Salt Mixture (Sigma), 19g KNO₃, 50ml 200X B5 vitamins, water to 1L), 1g/L glutamine, 0.5g/L asparagine, recipe), and sub-cultured to fresh EG medium every 3 to 4 weeks.

Once somatic embryos turned green and were larger than 2cm, they were plated root down in EG medium. At all stages of regeneration, growing plantlets were prevented from reaching the lids or sides of their containers. Germinated embryos with 1 to 2 true leaves were transferred to EG medium in 175ml Greiners. Strong plantlets with true leaves were transferred to sterile peat plugs expanded with dH₂O in 175ml Greiners and transferred to a growth cabinet under conditions of 14 hours daylight at 30°C and 10 hours darkness at 20°C. Thereafter, plantlets were transplanted into pots and grown in the glasshouse.

1.3 Identification and selection of transgenics

Putative transgenic plants were screened by PCR for the presence of the VIP3A gene. Positive events were identified and screened using insect bioassays for insecticidal activity against Fall Armyworm (*Spodoptera frugiperda*) (see Example 7). Insecticidal lines were characterized for copy number by TaqMan™ analysis (see Example 2). T1 seed from several events were observed in a field trial for insect resistance and agronomic quality. Two events, COT202 and COT203, were chosen based on having a single copy of the transgene, good protein expression as identified by ELISA (see Example 4), good insecticidal activity against Cotton Boll Worm (*Helicoverpa zea*) and field performance. The hygromycin selectable marker cassette was segregated away using conventional plant breeding to result in the COT202 event and the COT203 event.

1.4 Verification of sequence of COT202

Genomic DNA was isolated from the COT202 event. This was used in the sequencing of the junctions of the DNA insertion site with the cotton genomic DNA in the COT202 event, using standard DNA sequencing techniques.

5 **Example 2: COT202 event specific Detection via TaqMan™**

2.1 *DNA extraction*

DNA was extracted from leaf tissue using the Wizard™ Magnetic 96 DNA Plant System (Promega, #FF3760), according to the manufacturers instructions, with an additional step at the beginning of the protocol: following grinding of the leaf material, 0.9ml Cotton
 10 Extraction Buffer (0.2M Tris pH 8.0, 50mM EDTA, 0.25M NaCl, 0.1% v/v 2-mercaptoethanol, 2.5% w/v polyvinyl-pyrrolidone) was added to each well, the plant tissue resuspended and the plate centrifuged at 4,000 rpm (2755g) for 10 minutes. After aspirating and discarding the supernatant; 300ul Lysis Buffer A (Promega) was added and the manufacturers protocol was followed from this point. This procedure resulted in
 15 approximately 85ul of purified genomic DNA at a concentration of approximately 10ng/ul.

2.2 *TaqMan™ PCR reactions*

TaqMan™ PCR reactions were setup using a standard reaction mix comprising:

20 5ul 2x Jumpstart Master Mix for Q-PCR (Sigma, #P2893), supplemented with
 15mM MgCl₂ and 200nM Strata-ROX
 0.2ul 50x FAM primer / probe mix
 0.2ul 50x VIC primer / probe mix
 1.6ul Water.

25 50x primer / probe mixes comprised 45ul of each primer at a concentration of 1mM, 50ul of the probe at a concentration of 100uM and 860ul nuclease free water, and were stored in an amber tube at 4°C. Examples of suitable primer / probe sequence combinations which were used are:

<u>Primer Name</u>	<u>Primer Sequence 5'-3'</u>	<u>SEQ ID</u>
GhCHI2b-F Forward	GGTCCCTGGATACGGTGTCA	SEQ ID NO: 10

GhCHI2b-R TTGAGGGTTGGATCCTTTGC SEQ ID NO: 11
Reverse

GhCHI2bNEW-VIC CACCAACATCATCAATGGTGGCATCG SEQ ID NO: 12
Probe (5' label = VIC, 3' label = TAMRA)

COT202-F GGAATGTGGCGAATGGTGAT SEQ ID NO: 13
Forward

COT202-R TGTCGTTTCCCGCCTTCA SEQ ID NO: 14
Reverse

COT202-FAM CAAATTGCCCATTTTCATTCATCCAAA SEQ ID NO: 15
Probe AGC
 (5' label = FAM, 3' label = TAMRA)

7ul of master mix was dispensed into each well of a 384-well TaqMan™ assay plate. 3ul DNA template was added to the appropriate wells. 3ul of copy control dilution series was added to specific wells as a control. The reactions were run in an ABI7900HT (Applied Biosystems) using the following cycling conditions:

	Step	Temperature	Time
	1	50°C	2 min
	2	95°C	10 min
10	3	95°C	15 sec
	4	60°C	1 min
	5	Goto step 3, repeat 40 times	

Data was analysed using SDS2.0 version A, software (Applied Biosystems).

Example 3: COT202 Detection via Multiplex PCR Zygoty Test*3.1 Genomic DNA extraction*

Genomic DNA from COT202 was extracted as described in Example 2.1.

5 *3.2 Multiplex PCR*

PCR primers were designed to bind to cotton genomic DNA sequence upstream of the site at which the COT202 cassette inserted (SEQ ID NO: 16); the cotton genomic DNA sequence downstream of the site at which the COT202 cassette inserted (SEQ ID NO: 17); and the COT202 cassette sequence itself (SEQ ID NO: 18). A 25ul PCR reaction was set
10 up for each sample to be tested as follows:

1x JumpState ReadyMix REDTaq PCR (Sigma P-1107)

0.5uM primer 1 (SEQ ID NO: 16)

0.5uM primer 2 (SEQ ID NO: 17)

15

0.5uM primer 3 (SEQ ID NO: 18)

0.2% BSA

20 ng genomic DNA

ddH₂O to 25ul

20 The PCR reactions were heated in a thermocycler at 94°C for 5 minutes, followed by 30 cycles as follows: 94°C for 30 seconds, 55°C for 45 seconds, 72°C for 1 minute. The reaction was completed by heating at 72°C for 5 minutes.

3.3 Analysis

25 PCR reactions were run on an agarose gel, and DNA bands visualised under UV light after staining with ethidium bromide. The presence of 2 bands indicated that the sample was from a COT202 heterozygote plant; 1 band of 181bp in size indicated that the sample was from a COT202 homozygote plant; and 1 band of approximately 400bp in size indicated that the sample was from a homozygote wild type cotton plant.

30

3.4 COT202 Detection via Standard PCR

As an alternative to the multiplex PCR, the COT202 event can be detected in a simple PCR reaction using the primers depicted as SEQ ID NO: 3 and 4, SEQ ID NO: 13 and 14, or

SEQ ID NO: 16 and 18. The composition of the PCR reaction mixture is the same as described in example 3.2 above. The PCR reactions are heated in a thermocycler at 94°C for 5 minutes, followed by 30 cycles as follows: 94°C for 30 seconds, 55°C for 30 seconds, 72°C for 20 seconds. The reaction is completed by heating at 72°C for 5 minutes. A DNA
5 fragment of 181bp, 86bp or 181bp in size respectively indicates the presence of the COT202 event.

Example 4: COT202 Detection via Southern Blot

4.1 DNA extraction for use in Southern Blotting

10 Approximately 2 to 3g fresh weight of young leaf tissue was ground in a chilled mortar and pestle to a fine powder and added to 15 ml of ice-cold Nuclei extraction buffer (0.35M glucose, 0.1M Tris-HCl pH8, 50mM Na₂EDTA, 2% Polyvinyl-pyrrolidone-10, 0.1% ascorbic acid, 0.2% B-mercaptoethanol) in a labelled tube. The sample was incubated on ice for 15-20 minutes. The tube was mixed gently and centrifuged at 2700g for 20 minutes
15 at 4°C. The supernatant was discarded and 8ml of nuclei lysis buffer (0.14M sorbitol, 0.22M Tris-Cl pH8, 0.8M NaCl, 0.22M Na₂EDTA, 0.8%w/v CTAB, 1% Sarkosyl, 1% Polyvinyl-pyrrolidone-10, 0.1% ascorbic acid, 0.2% B-mercaptoethanol, 5µg/ml proteinase K) was added. After mixing, the tubes were incubated at 65°C for 30 minutes. 10ml chloroform was added, and the tube mixed gently by inversion until an emulsion formed
20 followed by centrifugation at 4600rpm for 10 minutes at room temperature.

The aqueous layer was removed into a new tube containing 10µl RNase A (10mg sigma R4642), and the tube incubated for 30 minutes at 37°C. The chloroform and centrifugation steps were repeated once. The aqueous layer was removed into a new tube containing
25 10ml propan-2-ol. After approximately 15 minutes incubation at room temperature, a gelatinous precipitate was observed in the middle of the tube. The tube was mixed gently to precipitate out the DNA. The DNA was spooled out using a sterile loop into a falcon tube containing 70% ethanol. The DNA was air-dried to remove the ethanol and resuspended in 200-400µl TE.

30

4.2 Alternative method for DNA extraction

2-3 young cotton leaves (approximately 1g fresh weight) are ground to a paste in a mortar and pestle at room temperature, with 2ml of grinding buffer (100mM NaOAc pH 4.8,

50mM EDTA pH8.0, 500mM NaCl, 2% PVP (10,000 MW), 1.4 % SDS) and a little sand. The ground tissue is transferred to a 15ml falcon tube, and the remnants in the mortar rinsed with 1 ml of grinding buffer into the tube. The sample is incubated at 65°C for 15 minutes, shaking occasionally. 4ml 10M ammonium acetate is added, and the sample
5 mixed well and incubated at 65°C for 10 minutes to precipitate proteins. The samples are incubated at room temperature at 4600 rpm for 10 minutes. The aqueous phase is transferred to a fresh 15ml tube.

0.6 volumes of cold isopropanol are added and the sample is incubated at room
10 temperature for approximately 30 minutes. After mixing by slowly inverting the tube several times, the DNA is spooled out and dissolved in 500ul TE. 10ul of 10mg/ml RNase are added and incubated for 15 minutes at room temperature. Following extraction with 500ul of phenol: chloroform: isoamyl alcohol (25:24:1), the sample is mixed gently and centrifuged at 13000rpm for 5 min.

15 The supernatant is transferred to a fresh tube using a fine Pasteur pipette and re-extracted with chloroform: isoamyl alcohol (24:1) as above. The supernatant is transferred to fresh tubes, 1/10 volume 3M NaOAc (pH4.8) added and mixed, and then one volume cold isopropanol is added. The sample may be incubated at room temperature for up to 30
20 minutes to precipitate the DNA. The DNA is spooled out and resuspended in 70% ethanol. The DNA is air-dried to remove the ethanol and resuspended in 200ul water.

4.3 *Restriction enzyme digests*

The DNA was quantified using a spectrophotometer and running out on a gel. Suitable
25 enzyme digests were prepared using 5ug DNA per digest in a total volume of 40ul. Digests included *HindIII*, *XmaI*, *BamHI*, *NheI*, and *SacI*, both alone and in combination. In particular, a *HindIII* and *XmaI* double digest was used to detect the intactness of the VIP3A gene; a *NheI* digest was used to detect VIP3A locus number. Digests were incubated for 6 hours at the appropriate temperature for each enzyme.

30

4.4 *Gel electrophoresis*

Bromophenol blue loading dye was added to each sample from 4.2 above, and each sample loaded on a 0.8% TBE agarose gel. The gel was run at 50 volts overnight.

After running, the gel was washed in 0.25M HCl for 10 minutes to depurinate the DNA, incubated in denaturing solution (0.5M NaOH, 1.5M NaCl) with gentle agitation for 30 minutes, rinsed with distilled water and then incubated in neutralising solution (0.5M Tris, 1.5M NaCl) for 30 minutes.

A Southern Blot was prepared as follows: A glass plate was placed over a tray containing 20X SSC and a strip of 3M paper was placed onto the glass plate such that both ends dipped into the 20X SSC solution (to act as a wick). A piece of 3M paper the same size as the gel was placed on the wick, and the gel placed on this. Strips of nescofilm were laid around the edges of the gel to form a seal. A Hybond membrane was placed on top of the gel, followed by two further pieces of 3M paper. Throughout the assembly of the blot, care was taken to ensure that no air bubbles were trapped between the membrane, gel and 3M paper. A 5cm-10cm stack of absorbent paper towels was placed on top of the 3M paper and held in place with a weight.

The DNA was allowed to transfer to the Hybond membrane overnight. After transfer the Southern Blot stack was disassembled and the DNA was bound to the membrane via UV cross-linking.

4.5 Hybridisation

A suitable DNA probe was prepared by PCR or restriction digest of binary plasmid. 25ng probe DNA in 45ul TE was boiled for 5 minutes, placed on ice for 5 minutes then transferred to a Rediprime II (Amersham Pharmacia Biotech, #RPN1633) tube. After addition of 5ul P32-labelled dCTP to the Rediprime tube, the probe was incubated at 37°C for 1 hour. The probe was purified by centrifugation through a microspin G-50 column (Amersham Pharmacia Biotech, #27-5330-01) according to the manufacturers instructions to remove unincorporated dNTPs. The activity of the probe was measured roughly by comparing the amount of radioactive component remaining in the column to the amount in the sample tube, with a ratio of at least 50:50 being acceptable. The Hybond membrane was pre-hybridised by wetting with 40 ml pre-warmed Rapid-Hyb buffer (Amersham-Pharmacia), at 65°C for 30 minutes. The labelled probe was boiled for 5 minutes, and placed on ice for 5 minutes. An appropriate amount of probe (1 million counts per 1ml

pre-hybridisation buffer) was added to the pre-hybridisation buffer and hybridisation occurred at 65°C overnight. The following day, the hybridisation buffer was discarded, and following a rinse with 50ml 2xSSC/1%SDS solution the membrane washed in 150ml 2xSSC/1%SDS solution at 65°C for 30-45 minutes. This process was repeated twice with 5 0.1xSSC/1%SDS solution .The membrane was exposed to a phosphor screen or X-ray film to detect where the probe had bound.

Example 5: COT202 Detection via ELISA

5.1 Protein extraction

10 Cotton tissue for analysis was harvested and frozen at -70°C. Fresh tissue was ground to a fine powder and weighed into a labelled polypropylene tube. Extraction buffer (100mM Tris, 100mM Sodium Borate, 5mM MgCl, 0.05% Tween 20, 0.2% Sodium Ascorbate, Water, pH 7.8, 1mM AEBSEF, 0.001mM Leupeptin) was added to the sample in a ratio of 2:1 (volume extraction buffer : sample fresh weight) for fresh tissue or 30:1 (volume 15 extraction buffer : sample dry weight) for lyophilised tissue. The sample was vortexed and homogenised using a Brinkman PT 10/35 Polytron equipped with a PTA 10TS foam-reducing generator, until the mixture became liquefied. Extracts were centrifuged at 10,000 x g for 15 minutes. The protein extract supernatant was stored at 2-8°C.

20 *5.2 ELISA protocol*

The ELISA procedure used standard techniques as follows. A 96-well plate was soaked in ethanol for 2 hours, and air-dried. The plate was coated with 50ul goat anti-VIP3A antibody per well and incubated overnight at 2-8°C. After washing three times with 1X ELISA wash solution (100mM Tris, 0.5% Tween-20, 75mM NaCl, pH8.5), the plate was 25 dried briefly by tapping upside down on a paper towel. 150ul blocking solution (10mM NaPO₄, 140mM NaCl, 1% BSA, 0.02% Sodium Azide, titrated to pH7.4 with monobasic NaPi and dibasic NaPi) was added to each well followed by incubation at room temperature for 45 minutes. The plate was washed 3 times as described above.

30 VIP3A standards and protein extract samples were applied to appropriate wells of the plate in triplicate, 50ul total volume per well. The plate was incubated at 2-8°C for 1 hour 30 minutes, followed by room temperature for a further 30 minutes. The plate was washed three times with ELISA wash solution, and then incubated at 35-39°C for 1 hour with 50ul

rabbit anti-VIP3A antibody per well. The plate was washed three times with ELISA wash solution, and incubated at room temperature for 30 minutes with 50ul donkey anti-rabbit alkaline phosphatase per well. Following a further three washes with ELISA wash solution, 50ul phosphatase substrate solution was added per well and the plate incubated
5 for 30 minutes at room temperature. The reaction was stopped by addition of 50ul 3M NaOH per well. The absorbance of the solution in each well was measured at 405nm using a Ceres 900C multiwell plate reader and the results analysed using KC3 Curve fitting software (Bio-Tek Instruments Inc.). The concentration of VIP3A in the samples was calculated by reference to the VIP3A protein standards.

10

Example 6: COT202 detection via DipStick

6.1 Protein extraction

A piece of leaf tissue approximately 2 cm² was placed in a tube containing extraction buffer. A plastic stirrer was used to extract protein from the tissue, by cutting into and
15 mascerating the tissue.

6.2 Dipstick test

A test strip was placed into the tube and incubated for 5 to 10 minutes for the result to develop. The test strip comprised a first band at which anti-VIP3A antibody was bound,
20 and a second band at which a control antibody was bound. After incubation, a double red line in the result window of the test strip indicated that VIP3A was present. The lower line indicated the presence of VIP3A protein while the upper line was a control indicating that the assay was working correctly.

Example 7: COT202 Detection via Insect Bioassay

7.1 Leaf biosassays

Leaf assays were performed on Fall Army Worm (*Spodoptera frugiperda*), Cotton Boll Worm (*Helicoverpa zea*) and Tobacco Budworm (*Heliothis virescens*) as follows:
30 Pads were soaked with 300ul to 500ul distilled water and placed into Gelman dishes. Leaf pieces measuring between approximately 0.5 square inches and 0.75 square inches were excised from cotton plants 8 to 12 inches in height, and placed on the pads. Between 8 and 10 insect larvae were placed in each dish and a lid fitted. The dishes were incubated at

28°C. On the third and sixth days after infestation, damage to the leaf in each dish was scored and compared with the control plants.

7.2 *Boll bioassays*

- 5 Four absorbent pads were saturated with water and placed inside a large plastic cup. Three extra thick glass filters, each soaked with 100ul distilled water, were placed in a smaller plastic cup, which was then seated inside the larger cup. A 1.25 inch long boll was excised, immersed in 10mg/ml to 20mg/ml Nystatin and placed on the filters in the small cup. 50 insect larvae were placed on the square or boll and a lid attached to the larger cup.
- 10 The squares or bolls were re-infested with 50 more larvae after 7 days.

The experiment was incubated at room temperature for approximately 3 weeks. The bolls were then cut open to determine damage. Damage to the boll was compared to the control samples.

15

7.3 *Lyophilised leaf bioassays*

Bioassays using freeze-dried leaf tissue were performed on *Heliothis virescens* as follows:

- 20 Terminal leaves were snap-frozen on dry-ice at time of picking and lyophilised overnight. The freeze dried tissue was ground in a mortar and pestle to a fine powder and resuspended in 0.2% agar solution to make an 8% (0.08g/ml) suspension of leaf powder. The suspension was overlaid on top of artificial insect diet in 96-well plates and left to dry. A single neonate insect larva was introduced into each well and the plates sealed. The plates
- 25 were incubated at 28°C. On the sixth day after infestation, larval mortality was scored and compared with control samples.

Example 8: COT202 Field Trials

8.1 *Field trial design*

- 30 The efficacy of the COT202 event against *Heliothis virescens* (Tobacco Bud Worm) and *Helicoverpa zea* (Cotton Boll Worm) was tested by conducting field trials at three locations in the US, namely Leland (MS), Quitman (GA) and Beasley (TX). Trials in each location were set up using a randomised complete block design, with four entry plots comprising four rows of 40 feet in length and four repetitions per trial. Seed was planted

to obtain a plant stand of approximately 3 plants per foot of row length. Each field trial included non transgenic Coker 312 plants for control purposes, and two other transgenic events designated event A and event B for comparison purposes.

5 8.2 *Field trial assessment*

An assessment of the natural insect populations was made at each trial location at the first white flower stage, approximately 80 days after planting. Where insect pressure was below the US economic threshold of 10% damage, artificial infestation of Cotton Boll Worm and Tobacco Bud Worm was made. The artificial infestation method was designed
10 to obtain a rate of 10 eggs per foot per insect species. Assessment of damage to cotton squares and bolls was made by visual inspection of 50 fruiting forms per plot at 5 – 7 days after artificial infestation. When relying on a natural infestation, damage ratings were made when the non transgenic Coker 312 control plants showed fruiting form damage above the economic threshold level of 10% in all control plots.

15

8.3 *Field trial results*

The results presented below show percentage damage to cotton squares and bolls at each field trial location, for each plant category. The data below represents an average of 200 fruiting forms (squares or bolls) per event per trial.

20

Location: Leland, MS

	Square Damage	Boll damage
Control	77	No data
COT202 event	6.6	No data
Event A	3.6	No data
Event B	29.6	No data

25

Location: Quitman, GA

	Square Damage	Boll Damage
Control	80	52
COT202 event	2	0
Event A	1.5	0
Event B	24.5	2

5 Location: Beasley, TX, USA

	Square Damage	Boll Damage
Control	24.4	13.2
COT202 event	4	2
Event A	2	1.2
Event C	5.2	3.2

CLAIMS

1. A polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 1.
5
2. A polynucleotide according to claim 1 comprising at least 18 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 1.
3. A polynucleotide according to claim 1 comprising at least 20 contiguous nucleotides
10 from the 26-nucleotide sequence of SEQ ID NO: 1.
4. A polynucleotide according to claim 1 comprising the sequence of SEQ ID NO: 1.
5. A polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide
15 sequence of SEQ ID NO: 2.
6. A polynucleotide according to claim 5 comprising at least 18 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 2.
- 20 7. A polynucleotide according to claim 5 comprising at least 20 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 2.
8. A polynucleotide according to claim 5 comprising the sequence of SEQ ID NO: 2.
- 25 9. A polynucleotide according to any of the preceding claims comprising the sequence of SEQ ID NO: 8.
10. An insect resistant plant comprising a VIP3A protein and a polynucleotide according to any of claims 1 to 9.
- 30 11. A plant according to claim 10 which is a cotton plant.

12. An insecticidal cotton plant according to claim 11 which is derived from the COT202 event.
13. A method of detecting plant material derived from the COT202 event comprising:
- 5 (a) obtaining a sample for analysis;
- (b) providing DNA from the sample;
- (c) providing a pair of primers designed to bind to a polynucleotide as claimed in claims 1 to 9 when said polynucleotide is single stranded;
- (d) amplifying the region which lies between the sites at which the primers bind; and
- 10 (e) detecting the presence of the amplification product;
- whereby the presence of the amplification product is indicative that the sample is derived from the COT202 event.
14. A method according to claim 13 wherein the first primer has the sequence of SEQ ID NO: 3 and the second primer has the sequence of SEQ ID NO: 4.
15. A method of detecting plant material derived from the COT202 event comprising:
- (a) obtaining a sample for analysis;
- (b) providing a probe designed to bind to the complement of a polynucleotide as
- 20 claimed in claims 1 to 9 when said polynucleotide is single stranded;
- (c) hybridising said probe with the sample; and
- (d) detecting whether the probe has hybridised;
- whereby the hybridisation of the probe is indicative that the sample is derived from the COT202 event.
- 25
16. A method according to claim 15 wherein the sequence of the probe is selected from the group comprising SEQ ID NO: 7 and SEQ ID NO: 8.
17. A method according to claims 15 or 16 wherein the probe hybridises to the sample
- 30 under stringent hybridisation conditions.
18. A method of detecting plant material derived from the COT202 event comprising:
- (a) obtaining a sample for analysis;

- (b) providing an antibody designed to bind to a VIP protein contained within a plant according to claims 10 to 12;
- (c) incubating said antibody with the sample; and
- (d) detecting whether the antibody has bound;
- 5 whereby the presence of antibody which has bound is indicative that the sample is derived from the COT202 event.
19. A method of detecting plant material derived from the COT202 event comprising:
- (a) obtaining a sample for analysis;
- 10 (b) making a protein extract of the sample;
- (c) providing a test strip designed to detect the presence of a VIP protein present within the sample;
- (d) incubating the test strip with the sample; and
- (e) detecting whether VIP protein is present;
- 15 wherein the presence of VIP protein is indicative that the sample is derived from the COT202 event.
20. A method according to claim 18 or 19 wherein the VIP protein has the sequence of SEQ ID NO: 9.
- 20
21. A method of detecting plant material derived from the COT202 event comprising:
- (a) obtaining a sample for analysis;
- (b) subjecting one or more insects of the species *Spodoptera frugiperda* to the sample;
- (c) subjecting one or more insects of species *Ostrinia nubilalis* to the sample as a
- 25 control;
- (d) detecting whether the sample has an insecticidal effect on insects from each species; and
- (e) comparing the results with an authentic COT202 bioassay profile.
- 30 22. A kit of parts comprising a means for detecting the presence in a sample of plant material derived from the COT202 event.

23. A kit of parts according to claim 22 comprising a means for detecting the presence in a sample of a polynucleotide according to claims 1 to 9, or a protein encoded by a polynucleotide according to claims 1 to 9, or a VIP protein.

5 24. A kit of parts according to claims 22 or 23 comprising in the form of instructions one or more of the methods according to claims 13 to 21.

25. Use of the nucleotide sequences according to any one of claims 1 to 9 in a method of detecting the COT202 event.

SEQUENCE LISTING

<110> SYNGENTA PARTICIPATIONS AG

<120> INSECT RESISTANT COTTON PLANTS AND METHODS OF DETECTING THE SAME
<130> 70325
<160> 18
<170> PatentIn version 3.1
<210> 1
<211> 26
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<400> 1
aacaaacaca aaatcttttc accagt 26

<210> 2
<211> 26
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<400> 2
ttcccgcctt cagatcttct gcaaca 26

<210> 3
<211> 23
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<400> 3
gggtgccatc gggtagtcca taa 23

<210> 4
<211> 24
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<400> 4
tctatgttac tagatcggga attg 24

<210> 5
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<400> 5
 gatcggggtc aggaaggtct 20

<210> 6
 <211> 20
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> COT202 nucleotide motif

<400> 6
 cagcatcatg aacgagcact 20

<210> 7
 <211> 290
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> COT202 nucleotide motif

<400> 7
 gacaaggaca gcttgagcga ggtgatctac ggcgacatgg acaagctgct gtgtccggac 60
 cagagcgagc aaatctacta caccaacaac atcgtgttcc cgaacgagta cgtgatcacc 120
 aagatcgact tcaccaagaa gatgaagacc ctgcgctacg aggtgaccgc caacttctac 180
 gacagcagca ccggcgagat cgacctgaac aagaagaagg tggagagcag cgaggccgag 240
 taccgcaccc tgagcgcgaa cgacgacggc gtctacatgc cactgggcgt 290

<210> 8
 <211> 4382
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> COT202 nucleotide motif

<400> 8
 gatcttgagc caagtctcat aaacgccatt gtggaagaaa gtcttgagtt ggtggtaatg 60
 taacagagta gtaagaacag agaagagaga gagtgtgaga tacatgaatt gtcgggcaac 120
 aaaaatcctg aacatcttat tttagcaaag agaaagagtt ccgagtctgt agcagaagag 180
 tgaggagaaa ttaagctct tggacttctg aattgttccg cctcttgaat acttcttcaa 240
 tcctcatata ttcttcttct atgttaoctg aaaaccggca ttaaatctcg cgggtttatt 300
 ccggttcaac attttttttg ttttgagtta ttatctgggc ttaataacgc aggcctgaaa 360
 taaattcaag gcccaactgt tttttttttt aagaagttgc tgtaaaaaa aaaaaaaggg 420
 aattaacaac aacaacaaaa aaagataaag aaaataataa caattacttt aattgtagac 480
 taaaaaaaca tagatcttat catgaaaaaa agagaaaaa aataaaaaact tggatcaaaa 540

aaaaacatac agatccttcta attattaact tttcttaaaa attaggtcct ttttcccaac 600
aattagggttt agagttttgg aattaaacca aaaagattgt tctaaaaaat actcaaattt 660
ggtagataag tttccttatt ttaattagtc aatggtagat actttttttt cttttcttta 720
ttagagtaga ttagaatcct ttatgccaaag tattgataaa ttaaatcaag aagataaact 780
atcataatca acatgaaatt aaaagaaaaa tctcatatat agtattagta ttctctatat 840
atattatgat tgcttattct taatgggttg ggttaacca gacatagtct taatggaaag 900
aatctttttt gaactttttc cttattgatt aaattcttct atagaaaaga aagaaattat 960
ttgaggaaaa gtatatacaa aaagaaaaat agaaaaatgt cagtgaagca gatgtaatgg 1020
atgacctaact ccaaccacca ccataggatg tttctacttg agtcggtcct ttaaaaacgc 1080
acggtggaaa atatgacacg tatcatatga ttccttcctt tagtttcgtg ataataatcc 1140
tcaactgata tcttcctttt tttgttttgg ctaaagatat tttattctca ttaatagaaa 1200
agacggtttt gggcttttgg tttgcgatat aaagaagacc ttcgtgtgga agataataat 1260
tcatectttc gtctttttct gactcttcaa tctctcccaa agcctaaagc gatctctgca 1320
aatctctcgc gactctctct ttcaaggat attttctgat tctttttggt tttgattcgt 1380
atctgatctc caatttttgt tatgtggatt attgaatcct ttgtataaat tgcttttgac 1440
aatattgttc gtttcgtcaa tccagcttct aaattttgtc ctgattacta agatatcgat 1500
tcgtagtggt tacatctgtg taatttcttg cttgattgtg aaattaggat tttcaaggac 1560
gatctattca atttttgtgt tttctttggt cgattctctc tgtttttaggt ttcttatggt 1620
tagatccggt tctcttttgg gttgttttga tttctcttac ggcttttgat ttggtatatg 1680
ttegctgatt ggtttctact tgttctattg ttttatttca ggtggatcca ccatgaacaa 1740
gaacaacacc aagctgagca cccgcgcct gccgagcttc atcgactact tcaacggcat 1800
ctacggcttc gccaccggca tcaaggacat catgaacatg atcttcaaga ccgacaccgg 1860
cggcgacctg accctggacg agatcctgaa gaaccagcag ctgctgaacg acatcagcgg 1920
caagctggac ggcgtgaacg gcagcctgaa cgacctgat gccagggca acctgaacac 1980
cgagctgagc aaggagatcc ttaagatcgc caacgagcag aaccaggtgc tgaacgacgt 2040
gaacaacaag ctggacgcca tcaacacat gctgcgcgtg tacctgccga agatcaccag 2100
catgctgagc gacgtgatga agcagaacta cgccctgagc ctgcagatcg agtacctgag 2160
caagcagctg caggagatca gcgacaagct ggacatcctc aacgtgaacg tcctgatcaa 2220
cagcaccctg accgagatca ccccgcccta ccagcgcctc aagtacgtga acgagaagtt 2280
cgaagagctg accttcgcca ccgagaccag cagcaaggtg aagaaggacg gcagcccggc 2340

cgacatcctg gacgagctga ccgagctgac cgagctggcg aagagcgtga ccaagaacga 2400
cgtggacggc ttcgagttct acctgaacac cttccacgac gtgatgggtg gcaacaacct 2460
gttcggccgc agcgccctga agaccgccag cgagctgatc accaaggaga acgtgaagac 2520
cagcggcagc gaggtgggca acgtgtacaa cttcctgatc gtgctgaccg ccctgcaggc 2580
ccaggccttc ctgaccctga ccacctgtcg caagctgctg ggcctggccg acatcgacta 2640
caccagcatc atgaacgagc acttgaacaa ggagaaggag gagttccgcg tgaacatcct 2700
gccgaccctg agcaacacct tcagcaacc gaactacgcc aaggatgaagg gcagcgacga 2760
ggacgccaag atgatcgtgg aggctaagcc gggccacgcg ttgatcggct tcgagatcag 2820
caacgacagc atcacctgac tgaagggtga cgaggccaag ctgaagcaga actaccaggt 2880
ggacaaggac agcttgagcg aggtgatcta cggcgacatg gacaagctgc tgtgtccgga 2940
ccagagcgag caaatctact acaccaaaa catcgtgttc ccgaacgagt acgtgatcac 3000
caagatcgac ttcaccaaga agatgaagac cctgcgctac gaggtgaccg ccaacttcta 3060
cgacagcagc accggcgaga tcgacctgaa caagaagaag gtggagagca gcgaggccga 3120
gtaccgcacc ctgagcgcgga acgacgacgg cgtctacatg ccaactggcg tgatcagcga 3180
gaccttcctg accccgatca acggctttgg cctgcaggcc gacgagaaca gccgcctgat 3240
caccctgacc tgtaagagct acctgcgca gctgctgcta gccaccgacc tgagcaaaaa 3300
ggagaccaag ctgatcgtgc caccgagcgg cttcatcagc aacatcgtgg agaacggcag 3360
catcgaggag gacaacctgg agccgtggaa ggccaacaac aagaacgcct acgtggacca 3420
caccggcggc gtgaacggca ccaaggccct gtacgtgcac aaggacggcg gcatcagcca 3480
gttcatcggc gacaagctga agccgaagac cgagtacgtg atccagtaca ccgtgaaggg 3540
caagccatcg attcacctga aggacgagaa caccggctac atccactacg aggacaccaa 3600
caacaacctg gaggactacc agaccatcaa caagcgcttc accaccggca ccgacctgaa 3660
ggcgctgtac ctgatcctga agagccagaa cggcgacgag gcctggggcg acaacttcat 3720
catcctggag atcagcccga gcgagaagct gctgagcccg gagctgatca acaccaaaaa 3780
ctggaccagc accggcagca ccaacatcag cggcaacacc ctgaccctgt accagggcgg 3840
ccgcggcatc ctgaagcaga acctgcagct ggacagcttc agcacctacc gcgtgtactt 3900
cagcgtgagc ggcgacgcca acgtgcgcat ccgcaactcc cgcgaggtgc tgttcgagaa 3960
gaggtacatg agcggcgcca aggacgtgag cgagatgttc accaccaagt tcgagaagga 4020
caacttctac atcgagctga gccagggcaa caacctgtac ggcgggcccga tcgtgcactt 4080
ctacgacgtg agcatcaagt aggagctcta gatccccgga atttccccga tcgttcaaac 4140

atttggcaat aaagtttctt aagattgaat cctgttgccg gtcttgcgat gattatcata 4200
 taatttctgt tgaattacgt taagcatgta ataattaaca tgtaatgcat gacgttatatt 4260
 atgagatggg tttttatgat tagagtcccc caattataca tttaatacgc gatagaaaaac 4320
 aaaatatagc gcgcaaaacta ggataaatta tcgcgcgccg tgatcatctat gttactagat 4380
 cg 4382

<210> 9
 <211> 789
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> VIP3A protein motif

<400> 9

Met Asn Lys Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe
 1 5 10 15
 Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
 20 25 30
 Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu
 35 40 45
 Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys
 50 55 60
 Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
 65 70 75 80
 Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
 85 90 95
 Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
 100 105 110
 Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
 115 120 125
 Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
 130 135 140
 Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val
 145 150 155 160
 Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile
 165 170 175
 Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr
 180 185 190
 Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Leu Asp Glu
 195 200 205

Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val
 210 215 220
 Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly
 225 230 235 240
 Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile
 245 250 255
 Thr Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr
 260 265 270
 Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Gln Ala Phe Leu Thr
 275 280 285
 Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr
 290 295 300
 Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val
 305 310 315 320
 Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala
 325 330 335
 Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys
 340 345 350
 Pro Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser Ile Thr
 355 360 365
 Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp
 370 375 380
 Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu
 385 390 395 400
 Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe
 405 410 415
 Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys
 420 425 430
 Thr Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly
 435 440 445
 Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr
 450 455 460
 Arg Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu Gly Val
 465 470 475 480
 Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala
 485 490 495
 Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg
 500 505 510
 Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile
 515 520 525

Val Pro Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Ser Ile
 530 535 540

Glu Glu Asp Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn Ala Tyr
 545 550 555 560

Val Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His
 565 570 575

Lys Asp Gly Gly Ile Ser Gln Phe Ile Gly Asp Lys Leu Lys Pro Lys
 580 585 590

Thr Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser Ile His
 595 600 605

Leu Lys Asp Glu Asn Thr Gly Tyr Ile His Tyr Glu Asp Thr Asn Asn
 610 615 620

Asn Leu Glu Asp Tyr Gln Thr Ile Asn Lys Arg Phe Thr Thr Gly Thr
 625 630 635 640

Asp Leu Lys Gly Val Tyr Leu Ile Leu Lys Ser Gln Asn Gly Asp Glu
 645 650 655

Ala Trp Gly Asp Asn Phe Ile Ile Leu Glu Ile Ser Pro Ser Glu Lys
 660 665 670

Leu Leu Ser Pro Glu Leu Ile Asn Thr Asn Asn Trp Thr Ser Thr Gly
 675 680 685

Ser Thr Asn Ile Ser Gly Asn Thr Leu Thr Leu Tyr Gln Gly Gly Arg
 690 695 700

Gly Ile Leu Lys Gln Asn Leu Gln Leu Asp Ser Phe Ser Thr Tyr Arg
 705 710 715 720

Val Tyr Phe Ser Val Ser Gly Asp Ala Asn Val Arg Ile Arg Asn Ser
 725 730 735

Arg Glu Val Leu Phe Glu Lys Arg Tyr Met Ser Gly Ala Lys Asp Val
 740 745 750

Ser Glu Met Phe Thr Thr Lys Phe Glu Lys Asp Asn Phe Tyr Ile Glu
 755 760 765

Leu Ser Gln Gly Asn Asn Leu Tyr Gly Gly Pro Ile Val His Phe Tyr
 770 775 780

Asp Val Ser Ile Lys
 785

- <210> 10
- <211> 20
- <212> DNA
- <213> Artificial Sequence
- <220>
- <223> COT202 nucleotide motif

<400> 10
ggtcacctgga tacggtgtca 20

<210> 11
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<400> 11
ttgagggttg gatcctttgc 20

<210> 12
<211> 26
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<220>
<221> modified_base
<222> (1)..(1)
<223> VIC label at 5' end

<220>
<221> modified_base
<222> (26)..(26)
<223> TAMRA label at 3' end

<400> 12
caccaacatc atcaatggtg gcatcg 26

<210> 13
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<400> 13
ggaatgtggc gaatggtgat 20

<210> 14
<211> 18
<212> DNA
<213> Artificial Sequence
<220>
<223> COT202 nucleotide motif

<400> 14
tgtcgtttcc cgccttca 18

<210> 15
<211> 29

<212> DNA
 <213> Artificial Sequence
 <220>
 <223> COT202 nucleotide motif

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> FAM label at 5' end

<220>
 <221> modified_base
 <222> (29)..(29)
 <223> TAMRA label at 3' end

<400> 15
 caaattgccc atttcattca tccaaaagc 29

<210> 16
 <211> 23
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> COT202 nucleotide motif

<400> 16
 ggtgtccatc gggtagtcca taa 23

<210> 17
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> COT202 nucleotide motif

<400> 17
 tgagtaggag atgtaagttg gcgc 24

<210> 18
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> COT202 nucleotide motif

<400> 18
 tctatgttac tagatcggga attg 24

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/012662

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/82 C12Q1/68 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N C12Q A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, Sequence Search, EMBL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! 25 November 2001 (2001-11-25), "Mus musculus clone RP23-182G17, LOW-PASS SEQUENCE SAMPLING." XP002318755 retrieved from EBI accession no. EM_PRO:AC101211 Database accession no. AC101211 Positions 51663 - 51683</p> <p style="text-align: center;">----- -/--</p>	1-3

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

° Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

25 February 2005

Date of mailing of the international search report

14/03/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bucka, A

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/EP2004/012662

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! 28 August 2002 (2002-08-28), "Mus musculus BAC clone RP23-307E9 from chromosome 7, complete sequence." XP002318982 retrieved from EBI accession no. EM_PRO:AC131669 Database accession no. AC131669 Positions 14695 - 14715 -----	1-3
X	ANONYMOUS: "Application for licence for dealings involving an intentional release into the environment DIR 036/2003 Title: Breeding and pre-commercial evaluation of transgenic cotton expressing a vegetative insecticidal protein (VIP) and a herbicide tolerance gene" INTERNET ARTICLE, 'Online! October 2003 (2003-10), XP002318972 AUSTRALIA Retrieved from the Internet: URL:http://www.ogtr.gov.au/pdf/ir/dir036fi nalramp.pdf> 'retrieved on 2005-02-22! the whole document -----	1-25
P,X	WO 2004/039986 A (SYNGENTA PARTICIPATIONS AG; ELLIS, DANIEL, MURRAY; NEGROTTI, DAVID, VI) 13 May 2004 (2004-05-13) the whole document -----	13,15-25
X	WO 96/10083 A (CIBA-GEIGY AG) 4 April 1996 (1996-04-04) the whole document -----	10-25
Y	page 12 -----	1-9,14, 16
X	WO 02/078437 A (SYNGENTA PARTICIPATIONS AG; MILES, PAUL; KRAMER, VANCE; SHEN, ZHICHENG) 10 October 2002 (2002-10-10) the whole document -----	10-25
Y	page 43 -----	1-9
X	RICE W C: "Specific primers for the detection of Vip3A insecticidal gene within a Bacillus thuringiensis collection" LETTERS IN APPLIED MICROBIOLOGY, vol. 28, no. 5, May 1999 (1999-05), pages 378-382, XP002318971 ISSN: 0266-8254 the whole document -----	13,15, 17,22-25
Y	page 379, left-hand column; figures 1,2 -----	1-9,14, 16
	----- -/-	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/012662

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>SELVAPANDIYAN A ET AL: "Toxicity analysis of N- and C- terminus-deleted vegetative insecticidal protein from Bacillus thuringiensis" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, WASHINGTON,DC, US, vol. 67, no. 12, December 2002 (2002-12), pages 5855-5858, XP002251845 ISSN: 0099-2240 page 5855, right-hand column page 5857, right-hand column</p>	<p>13,15, 17,22-25</p>
X	<p>YU C-G ET AL: "THE BACILLUS THURINGIENSIS VEGETATIVE INSECTICIDAL PROTEIN VIP3A Lyses MIDGUT EPITHELIM CELLS OF SUSCEPTIBLE INSECTS" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, WASHINGTON,DC, US, vol. 63, no. 2, February 1997 (1997-02), pages 532-536, XP000673006 ISSN: 0099-2240 figure 2</p>	<p>18-20, 22-25</p>
A	<p>WO 03/013224 A (BAYER BIOSCIENCE N.V) 20 February 2003 (2003-02-20) page 6 - page 8 page 13 - page 17</p>	<p>1-25</p>

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/012662

Patent document cited in search report	A	Publication date	Patent family member(s)	Publication date
WO 2004039986	A	13-05-2004	WO 2004039986 A1	13-05-2004
WO 9610083	A	04-04-1996	US 5849870 A	15-12-1998
			AT 256743 T	15-01-2004
			AU 692934 B2	18-06-1998
			AU 3743395 A	19-04-1996
			BG 101384 A	31-10-1997
			BR 9509099 A	30-09-1997
			CA 2199049 A1	04-04-1996
			CN 1160420 A	24-09-1997
			CZ 9700908 A3	16-02-2000
			DE 69532333 D1	29-01-2004
			DE 69532333 T2	07-10-2004
			DK 792363 T3	26-04-2004
			WO 9610083 A1	04-04-1996
			EP 1382611 A2	21-01-2004
			EP 0792363 A1	03-09-1997
			ES 2213162 T3	16-08-2004
			HU 77449 A2	28-04-1998
			JP 10506532 T	30-06-1998
			PT 792363 T	31-05-2004
			RU 2196824 C2	20-01-2003
			SI 792363 T1	30-04-2004
			TR 960263 A2	21-06-1996
			US 6107279 A	22-08-2000
			US 6137033 A	24-10-2000
			US 6066783 A	23-05-2000
			US 5888801 A	30-03-1999
			US 5990383 A	23-11-1999
			US 5872212 A	16-02-1999
			US 5889174 A	30-03-1999
			US 5770696 A	23-06-1998
			US 5840868 A	24-11-1998
			US 5866326 A	02-02-1999
			US 5877012 A	02-03-1999
			US 2002078473 A1	20-06-2002
			TW 496896 B	01-08-2002
			ZA 9508121 A	29-04-1996
WO 02078437	A	10-10-2002	BR 0208589 A	23-03-2004
			CA 2442594 A1	10-10-2002
			EP 1377157 A2	07-01-2004
			HU 0303738 A2	01-03-2004
			WO 02078437 A2	10-10-2002
			ZA 200307266 A	03-05-2004
WO 03013224	A	20-02-2003	US 2003097687 A1	22-05-2003
			BR 0211726 A	21-09-2004
			WO 03013224 A2	20-02-2003
			EP 1417312 A2	12-05-2004
			US 2005005334 A1	06-01-2005