



(86) **Date de dépôt PCT/PCT Filing Date:** 2018/01/29  
 (87) **Date publication PCT/PCT Publication Date:** 2018/08/02  
 (45) **Date de délivrance/Issue Date:** 2022/11/29  
 (85) **Entrée phase nationale/National Entry:** 2019/07/24  
 (86) **N° demande PCT/PCT Application No.:** US 2018/015779  
 (87) **N° publication PCT/PCT Publication No.:** 2018/140894  
 (30) **Priorité/Priority:** 2017/01/30 (US62/452,220)

(51) **Cl.Int./Int.Cl. A61K 31/137** (2006.01),  
**A61K 47/12** (2006.01), **A61P 9/00** (2006.01),  
**A61P 9/12** (2006.01)  
 (72) **Inventeurs/Inventors:**  
 YADAV, VIVEK, US;  
 GARAPATI, SRIRAMYA, US;  
 HINGORANI, TUSHAR, US;  
 ILITCHEV, IOURI V., US;  
 AKASAPU, PREM SAGAR, US;  
 SOPPIMATH, KUMARESH, US;  
 PURI, NAVNEET, US  
 (73) **Propriétaire/Owner:**  
 NEVAKAR INJECTABLES INC., US  
 (74) **Agent:** LAVERY, DE BILLY, LLP

(54) **Titre : COMPOSITIONS DE NOREPINEPHRINE ET PROCEDES ASSOCIES**  
 (54) **Title: NOREPINEPHRINE COMPOSITIONS AND METHODS THEREFOR**

(57) **Abrégé/Abstract:**

The inventive subject matter is directed to compositions and methods for ready-to-inject norepinephrine compositions with improved stability. Most preferably, compositions presented herein are substantially antioxidant free and exhibit less than 10% isomerization of R-norepinephrine and exhibit less than 5% degradation of total norepinephrine.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(10) International Publication Number  
**WO 2018/140894 A1**

(43) International Publication Date  
02 August 2018 (02.08.2018)

## (51) International Patent Classification:

*A61K 31/137* (2006.01)      *A61P 9/00* (2006.01)  
*A61K 47/12* (2006.01)      *A61P 9/12* (2006.01)

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
KM, ML, MR, NE, SN, TD, TG).

## (21) International Application Number:

PCT/US2018/015779

## Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

## (22) International Filing Date:

29 January 2018 (29.01.2018)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

62/452,220      30 January 2017 (30.01.2017)      US

(71) Applicant: NEVAKAR, INC [US/US]; 1019 ROUTE 202/206, BLDG. K, BRIDGEWATER, New Jersey 08807 (US).

(72) Inventors: YADAV, Vivek; 1000 American Pacific Drive, Apartment 2711, Henderson, Nevada 89074 (US). GARA-PATI, Sriramya; 13 Fleet St., North Brunswick, New Jersey 08902 (US). HINGORANI, Tushar; 138 Bonney Ct, Bridgewater, New Jersey 08807 (US). ILITCHEV, Iouri V.; 21 Reed Rd., Hillsborough, New Jersey 08844 (US). AKASAPU, Prem Sagar; 906 Rivendell Way, Edison, New Jersey 08817 (US). SOPPIMATH, Kumaresh; 97 Autumn Ln, Skillman, New Jersey 08558 (US). PURI, Navneet; 40 Still Hollow Road, Lebanon, New Jersey 08833 (US).

(74) Agent: FESSENMAIER, Martin; Umberg Zipser LLP, 1920 Main Street, Suite 750, Irvine, California 92614 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: NOREPINEPHRINE COMPOSITIONS AND METHODS THEREFOR

(57) Abstract: The inventive subject matter is directed to compositions and methods for ready-to-inject norepinephrine compositions with improved stability. Most preferably, compositions presented herein are substantially antioxidant free and exhibit less than 10% isomerization of *R*-norepinephrine and exhibit less than 5% degradation of total norepinephrine.



WO 2018/140894 A1

## NOREPINEPHRINE COMPOSITIONS AND METHODS THEREFOR

[0001]

### **Field of the Invention**

[0002] The field of the invention is pharmaceutical compositions comprising norepinephrine, especially as it relates to storage stable, ready-to-inject, antioxidant free compositions, and method of manufacturing such compositions.

### **Background**

[0003] The background description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0004] Where a definition or use of a term in a reference is inconsistent or contrary to the definition of that term provided herein, the definition of that term provided herein applies and the definition of that term in the reference does not apply.

[0005] Norepinephrine is often used during CPR (cardio-pulmonary resuscitation), and in the treatment of cardiac arrest and profound hypotension. Norepinephrine is also used for blood pressure control in certain acute hypotensive states, including for example sympathectomy, poliomyelitis, pheochromocytectomy, spinal anesthesia, myocardial infarction, blood transfusion, and septicemia.

[0006] Currently, norepinephrine is marketed as Levophed<sup>®</sup>, which is a concentrated 4 mg per 4 mL norepinephrine bitartrate formulation to be administered by intravenous infusion following dilution with dextrose or dextrose and sodium chloride injection. Norepinephrine is also marketed by Baxter which supplies as a norepinephrine concentrate that is free of sodium metabisulfite and packaged under nitrogen. Unfortunately, most, if not all diluted commercially available norepinephrine formulations lack storage and should therefore be discarded within one day after reconstitution when stored at room temperature. Consequently,

risk for microbial contamination and dilution errors is present. In addition, Levophed also contains sodium metabisulphite as an antioxidant, and carries a warning label that sulfite may cause allergic type reactions including anaphylactic shock and life threatening or less severe asthmatic episodes in susceptible people. **Table 1** depicts ingredients of currently marketed norepinephrine compositions.

**Table 1:** Composition of currently marketed Norepinephrine Bitartrate Products.

Ingredient	Levophed® (Hospira)	Norepinephrine Bitartrate (Baxter)
Norepinephrine Bitartrate equivalent to Norepinephrine Base	1 mg/mL	1 mg/mL
Sodium Chloride	Isotonic	Isotonic
Sodium Metabisulphite	0.2 mg/mL	-
pH	3 – 4.5	3 – 4.5
Water for injection	q.s. 1 mL	q.s. 1 mL

[0007] Stability of Levophed® and Norepinephrine bitartrate injection (Baxter), in normal saline solutions is presented in **Table 2** and **Table 3** where norepinephrine was diluted to a concentration of 16 µg/ml. Stability was assessed in 250 ml saline at accelerated (*i.e.*, 40 ± 2 °C and 75 ± 5 % relative humidity, duration as indicated) and long term stability (*i.e.*, 25 ± 2 °C and 60 ± 5 % relative humidity, duration as indicated) storage conditions.

**Table 2:** Stability study of Levophed® diluted in 0.9% Saline (Hospira) at 16 µg/mL.

Storage Condition	25 ± 2 °C/60 ± 5 % RH				40 ± 2 °C/75 ± 5 % RH			
	Initial	1 Month	2 Month	3 Month	Initial	1 Month	2 Month	3 Month
Assay	97.3	98.9	97.9	91.9	98.8	96.5	80.2	71.9
Total Impurities	0.05	-	0.71	8.08	0.03	1.96	5.29	9.73

**Table 3.** Stability study of Norepinephrine bitartrate injection [Baxter] diluted in 0.9% Saline (Hospira) at 16 µg/mL

Storage Condition	25 ± 2 °C/60 ± 5 % RH				40 ± 2 °C/75 ± 5 % RH			
	Initial	1 Month	2 Month	3 Month	Initial	1 Month	2 Month	3 Month
Assay	99.9	99.7	97.0	92.2	99.4	91.5	82.9	77.6
Total Impurities	0.08	1.73	2.68	10.17	0.10	2.34	4.46	6.71

[0008] As can be seen from the results, the norepinephrine at ready-to-inject concentrations underwent significant degradation. Oxidative degradation could possibly be reduced or even prevented by addition of effective amounts of sodium metabisulphite to the ready-to-inject norepinephrine solution. However, the quantities of sodium metabisulphite that would be administered by injection of 250 ml of the ready-to-inject solution would be substantial and detrimental to the patient. To avoid issues associated with sodium metabisulphite, efforts have been made to provide norepinephrine formulations with a non-sulfite anti-oxidant. For example US 2016/0058715 teaches a ready-to-inject dosage form of norepinephrine that uses butylated hydroxyl anisole as an anti-oxidant. While generally deemed safe for topical and cosmetic use, butylated hydroxyl anisole was shown to produce some renal and hepatic damage (e.g., *Int J Toxicol.* 2002;21 Suppl 2:19-94).

[0009] In other attempts to provide ready-to-administer norepinephrine formulations with increased storage stability and reduced risk of human error, the pH on the injectable solution was reduced to between 3.2 and 3.6 with 40-200 µg/ml norepinephrine as is described in WO 2015/128418. While such formulations exhibited reduced degradation as compared to higher pH formulations, significant discomfort can occur at the injection site. Worse yet, at the pH used, norepinephrine isomerized relatively quickly from the active R (-) isomer to the inactive S (+) isomer. Isomerization is also encountered at exposure of norepinephrine to higher temperatures.

[0010] Therefore, there is a need for improved stable, low concentration, ready-to-inject and antioxidant free norepinephrine formulations, and methods of manufacturing and storing the same.

### **Summary of the Invention**

[0011] The inventive subject matter is directed to antioxidant free sterilizable/autoclavable ready-to-inject norepinephrine compositions having improved stability and a physiologically acceptable pH.

[0012] In one aspect of the inventive subject matter, the inventors contemplate a ready to ready-to-inject norepinephrine composition that comprises an aqueous acidic buffer having a pH range of between 3.7 and 4.3, wherein the aqueous buffer further comprises a chelating agent and a pharmaceutically acceptable salt. Most typically, the chelating agent is present in an amount of between 1 µg/ml and 100 µg/ml, and the pharmaceutically acceptable salt is present in an amount of between 0.6 wt% and 1.2 wt%. Norepinephrine (typically enantiomerically pure (*i.e.*, at least 98%) *R*-isomer) is dissolved at a concentration that is suitable for administration to a patient in need thereof. In further preferred aspects, the ready-to-administer norepinephrine composition is substantially free of antioxidants, and the ready-to-administer norepinephrine composition is formulated such that after storage over at least three months equal or less than 10% of the *R*-isomer form will isomerize to the *S*-isomer and such that equal or less than 5% of the total norepinephrine will degrade to degradation products. As used herein, reference to the term norepinephrine should be interpreted broadly to include pharmaceutically acceptable salts and prodrugs thereof.

[0013] Therefore, the inventors also contemplate a method of preparing a ready-to-inject norepinephrine composition that includes a step of formulating a liquid parenteral composition that contains in an aqueous acidic buffer norepinephrine as an *R*-isomer such that (a) the formulation exhibits less than 10% of isomerization of the *R*-isomer to an *S*-isomer after three months of storage of the liquid composition, and (b) the formulation exhibits equal or less than 5% degradation of total norepinephrine after three months of storage of the liquid composition. The aqueous acidic buffer will typically have a pH range of between 3.7 and 4.3, and the aqueous buffer will further comprise a chelating agent and a pharmaceutically acceptable salt. In such methods, the total norepinephrine is present in the liquid parenteral composition at a concentration of between 10 µg/ml and 100 µg/ml, and the ready-to-inject norepinephrine composition is substantially free of antioxidants.

[0014] Viewed from a different perspective, the inventors also contemplate a method of preparing a ready-to-inject norepinephrine composition that includes the steps of preparing an aqueous acidic buffer at a pH range of between 3.7 and 4.3, wherein the aqueous buffer also

includes a chelating agent and a pharmaceutically acceptable salt. Preferably, the chelating agent is present in an amount of between 1 µg/ml and 100 µg/ml, and tonicity is adjusted with the pharmaceutically acceptable salt (*e.g.*, NaCl). In a further step, norepinephrine (preferably enantiomerically pure *R*-isomer) is dissolved at a concentration suitable for administration to a patient in need thereof, and the ready-to-administer norepinephrine composition is formulated such that after storage over at least three months equal or less than 10% of the *R*-isomer form will isomerize to the *S*-isomer and such that equal or less than 5% of the total norepinephrine will degrade to degradation products. As before, it is generally preferred that the ready-to-administer norepinephrine composition is substantially free of antioxidants. In yet another step, the composition is autoclaved to sterility.

[0015] Most typically, but not necessarily, the aqueous acidic buffer is a citrate buffer and/or preferably has a concentration of between 5 mM and 20 mM. Furthermore, preferred aqueous acidic buffers will have a pH of between 3.8 and 4.2. With respect to the chelating agent it is contemplated that such agents are a bicarboxylic acid (*e.g.*, optionally hydroxylated, tartrate), a tricarboxylic acid (*e.g.*, aconitic acid, trimesic acid, citric acid), and/or an aminopolycarboxylic acid (*e.g.*, EDTA, EGTA, etc.), and that such chelating agents are present at low concentrations, preferably between 1 µg/ml and 10 µg/ml, or between 10 µg/ml and 100 µg/ml. The norepinephrine is typically present at a concentration of between 10 µg/ml and 100 µg/ml, for example, at a concentration of 16 µg/ml (+/- 10%), 32 µg/ml (+/- 10%), or 64 µg/ml (+/- 10%). Contemplated methods may also include a step of autoclaving the compositions.

[0016] With respect to stability it is contemplated that the storage condition is over at least three months at 40 °C and 75% (+/- 5) relative humidity, that equal or less than 6% of the *R*-isomer form will isomerize to the *S*-isomer, and/or that equal or less than 3.5% of the total norepinephrine will degrade to degradation products.

[0017] Where desired, contemplated compositions have a dissolved oxygen concentration of equal or less than 1 ppm (*e.g.*, by formulating the liquid parenteral composition using deoxygenated water), and/or or by packaging the composition together with a (preferably metal free) oxygen scavenger. Packaging may further make use of a container that is configured (*e.g.*, aluminized or otherwise treated) to reduce light-mediated oxidation of the norepinephrine.

**[0017a]** The inventive subject matter is also directed to the use of the ready-to-inject composition as defined herein for treating hypotension in a subject. The inventive subject matter is also directed to the ready-to-inject composition as defined herein for treating hypotension in a subject.

**[0017b]** In an embodiment, the ready-to-inject composition is for administration at an initial dose per minute and at a second dose per minute, and wherein the initial dose per minute is greater than the second dose per minute. In an embodiment, the initial dose per minute is a dose of between 8 and 12  $\mu\text{g}/\text{min}$ . In an embodiment, the second dose per minute is a dose of between 2 and 4  $\mu\text{g}/\text{min}$ . In an embodiment, the initial dose per minute is at least twice the second dose per minute.

### **Detailed Description of the Invention**

[0018] The inventive subject matter is directed to stable aqueous pharmaceutical preparations of norepinephrine (and pharmaceutically acceptable salts thereof) in a ready-to-inject form that are sterile and preferably substantially free of antioxidants. Most preferably, stability of such compositions is characterized by low (oxidative and photo-induced) degradation as well as low isomerization.

[0019] More specifically, the inventors have discovered that formulations can be prepared that will exhibit less than 8%, more typically less than 6%, even more typically less than 4%, and most typically less than 3% of degradation as determined by HPLC-UV, and that will exhibit less than 10%, more typically less than 8%, even more typically less than 6%, and most typically less than 4% of isomerization from *R*- to *S*-configuration as determined by HPLC-UV. Most notably, such formulations were found to be stable over extended periods without antioxidants (*e.g.*, at least 1 month, or at least two months, or at least three months), even at elevated storage temperatures (*e.g.*, accelerated storage conditions such as 40 °C and 75% relative humidity (+/- 5%)). Even more remarkable, such formulations could also be subjected to thermal sterilization, and particularly sterilizing to sterility (*e.g.*, over at least 5 min, or at least 10 min, or at least 15 min at 121 °C), without substantial increase (*i.e.*, >1.5%, or >1.0%, or >0.7%) of the *S*-isomer of norepinephrine.

[0020] Additionally, it should be appreciated that contemplated formulations can be filled in a polymer bag (*e.g.*, polypropylene) or other container that may subsequently be placed into a secondary container together with an oxygen scavenger, and especially a metal-free oxygen scavenger. Most typically, at least one of the polymer bag and the secondary container may be impervious to light in general or light of a wavelength that promotes photo-initiated degradation. For example, containers may be metalized (*e.g.*, aluminized) or combined or coated with carbonaceous materials or other dye(s). If desired, contemplated formulations are sufficiently stable to also allow filling into containers using a blow-fill-seal (BFS) process.

[0021] Therefore, contemplated norepinephrine formulations of the inventive subject matter can advantageously be provided in a ready-to-inject form to thereby avoid the inconvenience associated with diluting concentrated small volume norepinephrine parenteral formulations into diluents prior to infusion. Thus, the ready-to-inject formulations also eliminate microbial contamination risks and calculation errors associated with dilution. Most typically, contemplated formulations will be available in a range of concentrations commonly required

by medical practitioners for emergency restoration of blood pressure, for example in cases of acute hypotension. Consequently, norepinephrine will typically be present in formulations at a concentration of between 10 µg/ml and 100 µg/ml, including concentration of 16 µg/ml (+/- 10%), 32 µg/ml (+/- 10%), and 64 µg/ml (+/- 10%).

**[0022]** As will be readily appreciated, the norepinephrine for preparation of contemplated formulations is preferably (*R*)-Norepinephrine, or enantiomerically pure (*i.e.*, at least 98% *R*-isomer) norepinephrine. However, in less preferred aspects, isomeric purity can also be between 95-98%, or even between 90-95%. Of course, it should also be appreciated that the norepinephrine may be a salt of any suitable and pharmaceutically acceptable form, including mineral salts (*e.g.*, HCl salt) and organic salts (*e.g.*, bitartrate). Similarly, where desired, the norepinephrine may also be used in any suitable prodrug form (*e.g.*, β,3-dihydroxytyrosine, L-dihydroxyphenylserine, etc.).

**[0023]** Suitable buffers are generally buffers that stabilize the pH of the contemplated liquid formulations in an acidic pH range and will therefore include glycine buffers, citrate buffers, citrate/phosphate buffers, acetate buffers, etc. However, the inventors have further discovered that where the norepinephrine is provided as the norepinephrine bitartrate salt, a buffer can advantageously be omitted and the pH can be adjusted with suitable acid and/or base as is well known in the art. Notably, the bitartrate appeared to act as a weak buffer in the stability range for the norepinephrine as is shown in more detail below. Most typically the pH of the formulation will be less than 5.0 and more typically less than 4.5, and most typically less than 4.3, but higher than 3.0, more typically higher than 3.5, and most typically higher than 3.7. For example, suitable buffers will have a pH in the range of between 3.7 and 4.3, or between 3.7 and 4.0, or between 3.8 and 4.1, or between 3.9 and 4.2, or between 4.0 and 4.2. Notably, such pH range provided remarkable stability for low concentrations of norepinephrine, especially when in combination with a chelator and a salt. While not limiting to the inventive subject matter, the buffer strength is typically relatively low, for example, equal or less than 100 mM, and more typically equal or less than 50 mM, and most typically between 5 mM and 20 mM (*e.g.*, 10 mM).

**[0024]** Moreover, in further contemplated aspects, the formulation will also include one or more chelating agents, and particularly metal ion chelators. For example, suitable chelators include various bicarboxylic acids, tricarboxylic acids, and aminopolycarboxylic acids such as ethylenediaminetetraacetic acid (EDTA), ethylene glycol-bis(β-aminoethyl ether)-

N,N,N',N'-tetraacetic acid (EGTA), and penta(carboxymethyl)diethylenetriamine (DTPA), and salts and hydrates thereof. While not limiting to the inventive subject matter, it is contemplated that the metal ion chelators will slow down both the baseline and metal ion-stimulated autoxidation of norepinephrine. Remarkably, the inventors unexpectedly observed that the desirable effect of the chelators was observable at relatively low concentrations of the chelators. For example, reduction of the baseline and metal ion-stimulated autoxidation of norepinephrine was observed at chelator concentrations of between 1 µg/ml and 10 µg/ml, and between 10 µg/ml and 100 µg/ml. Interestingly, the chelators, and especially the aminopolycarboxylic acids retained stabilizing effect despite the relatively low pH favoring protonated forms of the chelators.

[0025] With respect to suitable salts it is contemplated that the salt is a pharmaceutically acceptable salt that can be used to increase tonicity. Therefore, pharmaceutically acceptable salts are contemplated, and especially NaCl, at a concentration of at least 0.6 wt%, or at least 0.7 wt%, or at least 0.8 wt%, or at least 0.9 wt%. For example, suitable salt concentrations are between 0.6 wt% and 1.2 wt%. Depending on the particular salt concentration, additional tonicity agents may be added and suitable tonicity agents include glycerol, thioglycerol, mannitol, lactose, and dextrose. The amount of tonicity adjusting agent used can be adjusted to obtain osmolality of the formulations in the range of 260 to 340 mOsm/kg. An osmometer can be used to check and adjust the amount of tonicity adjusting agent to be added to obtain the desired osmolality.

[0026] It should further be appreciated that contemplated compositions are substantially free of antioxidants (*i.e.*, do not include antioxidants in an amount effective to reduce degradation of total norepinephrine by at least 1% when stored over a period of at least three months at 25 °C. Indeed, the inventors unexpectedly discovered that some formulations with antioxidants (particularly with ascorbic acid) had decreased stability. Notably, contemplated formulations were stable as described in more detail below, even in the absence of effective quantities of antioxidants, especially where deoxygenated solvents (*e.g.*, typically water and/or buffer) were employed. Deoxygenation (*i.e.*, reduction of molecular dissolved oxygen) can be achieved in numerous manners, including sparging with inert gases (*e.g.*, helium, various freons, argon, xenon), agitation under vacuum, and/or using enzymatic systems that deplete a solution of dissolved oxygen (see *e.g.*, US 9187779). Additionally, or alternatively, ingress of molecular oxygen into the formulation can also be reduced by co-packaging a container

with the formulation in a secondary container that includes an oxygen scavenger, and especially a metal-free oxygen scavenger (*e.g.*, GLS100, Ageless®, Pharmakeep®, all commercially available from Mitsubishi Gas Chemical America).

[0027] With respect to the sterilization of contemplated formulations it should be appreciated that contemplated formulations may be sterilized using all known manners of sterilization, including filtration through 0.22 micron filters, heat sterilization, autoclaving, radiation (*e.g.*, gamma, electron beam, microwave). Unexpectedly, and as shown in more detail below, the inventors have also discovered that contemplated formulations were heat stable and did not undergo significant isomerization, even under conditions of sterilization (exposure to high-pressure saturated steam) at 121 °C for at least 5, or at least 10, or at least 15 minutes.

[0028] Based on the unexpected heat stability, the formulations contemplated herein can also be filtered through a 0.22 micron filter, and filled in to a polyethylene, polypropylene or low-density polyethylene containers in a blow-fill-seal (BFS) process. BFS is a form of advanced aseptic manufacturing wherein the container is formed, filled, and sealed in one continuous, automated system not requiring human intervention. The process begins with the extrusion of plastic granules in the form of a hot hollow pipe of molten plastic called a parison. The next step is the blow molding of the container with an open top through which the container is filled, all while the plastic remains hot and in a molten state. Once filled, the container is hermetically sealed and cooled. The blow-fill seal process can take several seconds, and contemplated ready-to-inject compositions advantageously are formulated to withstand the temperature and pressure requirements without substantial degradation of norepinephrine (*e.g.*, less than 5 wt%, less than 3 wt%, less than 2 wt%, less than 1 wt% degradation).

[0029] Once the norepinephrine formulations are filled in large volume polymeric, semi-permeable infusion containers (*e.g.*, BFS container or flexible IV bags), the containers can optionally be layered or covered with a secondary packaging system including an aluminum pouch or other oxygen scavenger. For example, the BFS containers can further be sealed in an oxygen and moisture barrier blister packaging. The blister packaging can comprise one or more layers, and the one or more layers can include aluminum foil or other oxygen absorber having an Oxygen Transmission Rate (OTR) between 0.0005 to 5.00 cc/100 in<sup>2</sup>/24hrs. Additionally or alternatively, one or more oxygen absorbers (metal or metal free, organic material) can be incorporated into any portion of the BFS container, the secondary packaging system, or between the two (*e.g.*, between the BFS container and the multi-layer packaging)

such that the oxygen absorber removes at least a portion of oxygen from the air surrounding said oxygen-sensitive drug. A beneficial feature of the oxygen absorber is the absorbance and removal of oxygen present in the primary packaging and in the liquid drug itself. Notably, it was found that the oxygen absorber also removed residual headspace oxygen in the primary packaging and also dissolved oxygen in the liquid over time, thereby further improving stability of norepinephrine.

[0030] The following examples are provided for illustrative purposes only and should not be interpreted as limiting the present invention.

### **Examples**

[0031] The following examples illustrate some of the experiments leading to the formulations according to the inventive subject matter, however, should not be construed to limit the scope of the claims in any way.

[0032] Stability and Isomerization: The ionization behavior of norepinephrine in aqueous solution is complex. Common with other o-hydroquinone systems, norepinephrine in aqueous solution is susceptible to oxidation to form the corresponding o-quinone, which can then also undergo various secondary reactions, which also becomes more prevalent as the pH becomes more alkaline. Norepinephrine may further isomerize to the pharmacologically less active *S*-enantiomer at low pH values, corresponding to protonation of the hydroxyl group at the benzylic chiral center. Therefore, to prevent norepinephrine cyclization reactions pH values less than 6.0 are desired. A pH range of 3.0 to 6.2 was screened to determine pH of optimum stability. Composition of norepinephrine bitartrate equivalent to 16 µg/mL norepinephrine base at various pH values were prepared are described below, with **Table 4** listing compositions of norepinephrine bitartrate in citrate buffer (10 mM),

For preparation of the solutions, about 90% of the final quantity of water was collected in a glass media bottle. Nitrogen (N<sub>2</sub>) gas was purged for about thirty minutes to reduce the dissolved oxygen levels. Sodium chloride was added and the solution was stirred until a homogenous solution was obtained. Citric acid was added and the solution was stirred until a homogenous solution was obtained. The pH of the bulk solutions was adjusted to pH 3.0, 3.4, 3.8, 4.2, 4.6, 5.0, 5.4, 5.8, and 6.2, respectively for each formulation composition using sufficient quantity of 10% w/v sodium hydroxide or 10% w/v hydrochloric acid. Norepinephrine bitartrate was added and the solution was stirred for approximately 10

minutes until a clear solution was formed. Solutions were made up to volume with water. The solutions were filled into 10 mL glass vials, overlaid with nitrogen, stoppered, and sealed. The stability was studied at 4 °C, 25 °C, and 60 °C by assay. Samples were observed visually for precipitation and change in color for a period of 7 days. Data are presented in **Table 5**.

**Table 4:** Compositions of Norepinephrine Bitartrate for pH dependent stability in Citrate Buffer (10 mM).

Ingredients	Concentration (mg/mL)								
	I	II	III	IV	V	VI	VII	VIII	IX
Norepinephrine Bitartrate equivalent to Norepinephrine base	0.016	0.016	0.016	0.016	0.016	0.016	0.016	0.016	0.016
Sodium Chloride	9	9	9	9	9	9	9	9	9
Citric acid	1.92	1.92	1.92	1.92	1.92	1.92	1.92	1.92	1.92
Sodium Citrate	2.94	2.94	2.94	2.94	2.94	2.94	2.94	2.94	2.94
HCl/NaOH (q.s. pH)	3.0	3.4	3.8	4.2	4.6	5.0	5.4	5.8	6.2
Water for Injection (q.s. mL)	1	1	1	1	1	1	1	1	1

**Table 5** :Effect of pH on stability of Norepinephrine Bitartrate in citrate buffer.

Temperature	Formulation	Assay T <sub>0</sub>	Assay T <sub>7</sub>	pH	Color	Precipitation
4° C	I	96.4	96.5	3.0	No	No
	II	98.0	97.5	3.4	No	No
	III	99.0	98.5	3.8	No	No
	IV	99.1	98.4	4.2	No	No
	V	98.1	98.6	4.6	No	No
	VI	98.4	98.1	5.0	No	No
	VII	97.1	96.6	5.4	No	No
	VIII	97.8	97.5	5.8	No	No
	IX	91.5	91.2	6.2	No	No
25° C	I	96.4	96.4	3.0	No	No
	II	98.0	97.5	3.4	No	No
	III	99.0	97.9	3.8	No	No
	IV	99.1	97.7	4.2	No	No
	V	98.1	97.3	4.6	No	No
	VI	98.4	97.3	5.0	No	No
	VII	97.1	95.9	5.4	No	No
	VIII	97.8	94.5	5.8	No	No
	IX	91.5	80.4	6.2	Yes	No
60° C	I	96.4	95.2	3.0	No	No
	II	98.0	95.0	3.4	No	No
	III	99.0	95.2	3.8	No	No
	IV	99.1	93.2	4.2	No	No
	V	98.1	88.9	4.6	No	No
	VI	98.4	77.4	5.0	Yes	No
	VII	97.1	46.8	5.4	Yes	No
	VIII	97.8	NT	5.8	Yes	No
	IX	91.5	NT	6.2	Yes	No

[0033] No change in physical appearance was observed in the solutions stored at 4 °C. In the solutions stored at 25 °C, a change in color was observed at pH 6.2. Red brown color was observed in solutions stored at or above pH 5.0 at 60 °C. Oxidation and color formation are very common with norepinephrine in unfavorable conditions and the speed of the reaction and the nature of the final products are dependent on the catalysts (*e.g.*, metal ion impurities) and buffers employed. A pH range from 3.0 to 4.5 was selected for further testing.

[0034] Stability of Norepinephrine in selected pH ranges and formulations: The formulations for the next experiments are shown in **Table 6** below, involving three different compositions of norepinephrine bitartrate at three different pH (3.5, 4.0, 4.5, and 5.0) values. Lab scale

batches were prepared and subjected to lab scale stability tests at accelerated (40°C/75%RH) and long term stability (25°C/60% RH) storage conditions. The test results from the stability studies are presented in **Table-7 – Table 10**, with CCS indicating Clear colorless solution; ND indicating Not Detected; NR indicating Not Reported (<0.05%); and NT indicating Not Tested.

**Table 6:** Formulation composition selected for further development activities and optimization

Ingredient	Quantity (mg/mL)			
	X	XI	XII	XIII
<b>Formulation</b>	X	XI	XII	XIII
<b>Norepinephrine Bitartrate</b>	0.016	0.016	0.016	0.016
<b>Edetate Sodium</b>	0.10	0.10	0.10	0.10
<b>Sodium chloride</b>	9	9	9	9
<b>HCl/NaOH</b>	q.s. pH 3.5	q.s. pH 4.0	q.s. pH 4.5	q.s. pH 5.0
<b>Water for Injection Q.S.</b>	1 mL	1 mL	1 mL	1 mL
<b>Dissolved Oxygen (ppm)</b>	< 1	< 1	< 1	< 1

**Table 7:** Stability study of Formulation X - Norepinephrine Bitartrate Injection (16 µg/ml) filled in glass vial (pH 3.5).

Storage Condition	25 ± 2 °C/60 ± 5 % RH							40 ± 2 °C/75 ± 5 % RH					
	Initial	1 Month	2 Month	3 Month	4 Month	5 Month	6 Month	1 Month	2 Month	3 Month	4 Month	5 Month	6 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	3.50	3.65	3.59	3.56	3.58	3.54	3.48	3.66	3.61	3.59	3.64	3.60	3.59
Assay	101.4	99.6	97.1	97.1	101.0	102.3	102.2	99.5	97.0	98.7	100.4	101.7	101.4
S-form	NT	NT	NT	NT	1.8	2.2	2.2	NT	NT	NT	7.6	8.1	9.8
Total	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

**Table 8:** Stability study of Formulation XI- Norepinephrine Bitartrate Injection (16 µg/ml) filled in glass vial (pH 4.0).

Storage Condition	25 ± 2 °C/60 ± 5 % RH					40 ± 2 °C/75 ± 5 % RH			
	Initial	1 Month	2 Month	3 Month	6 Month	1 Month	2 Month	3 Month	6 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	4.02	3.96	3.98	3.97	3.91	4.01	3.99	4.02	4.03
Assay	101.3	98.7	95.5	99.2	100.5	98.6	95.3	97.1	97.5
S-form	NT	NT	NT	NT	1.7	NT	NT	NT	7.8
Total Impurities	0.1	ND	0.06	ND	0.80	ND	0.06	0.1	0.79

**Table 9:** Stability study of Formulation XII - Norepinephrine Bitartrate Injection (16 µg/ml) filled in glass vial (pH 4.5).

Storage Condition	25 ± 2 °C/60 ± 5 % RH					40 ± 2 °C/75 ± 5 % RH			
	Initial	1 Month	2 Month	3 Month	6 Month	1 Month	2 Month	3 Month	6 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	4.50	4.35	4.36	4.32	4.33	4.33	4.40	4.39	4.29
Assay	100.1	98.9	95.5	98.2	97.9	97.1	92.5	93.7	77.2
S-form	NT	NT	NT	NT	1.2	NT	NT	NT	2.9
Total Impurities	ND	0.32	0.79	0.52	3.41	1.18	0.38	5.59	10.38

**Table 10:** Stability study of Formulation XIII - Norepinephrine Bitartrate Injection (16 µg/ml) filled in glass vial (pH 5.0).

Storage Condition	25 ± 2 °C/60 ± 5 % RH					40 ± 2 °C/75 ± 5 % RH			
	Initial	1 Month	2 Month	3 Month	4 Month	1 Month	2 Month	3 Month	4 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	4.99	4.62	4.51	4.57	4.51	4.87	4.81	4.83	4.53
Assay	102.7	100.5	95.6	99.2	100.4	98.3	89.8	87.0	72.3
S-form	NT	NT	NT	NT	1.2	NT	NT	NT	3.0
Total Impurities	ND	0.75	0.81	0.48	1.29	0.94	2.4	5.39	14.91

[0035] Based on the above considerations, the effect of different levels of EDTA on stability of norepinephrine was determined. Three batches at concentrations of 16 µg/mL, 32 µg/mL, and 64 µg/mL were made with EDTA concentrations of 100 µg/mL: Formulation XIV (16 µg/mL), Formulation XV (32 µg/mL), Formulation XVI (64 µg/mL). Two additional batches were made at 10 µg/mL EDTA Formulation XVII and 1 µg/mL EDTA (Formulation XVIII) at 64 µg/mL Norepinephrine. The composition of the batches is specified in Table. The drug product was compounded as described earlier and packaged in 250 mL in polypropylene bags. This was further packed into aluminum overwrap with an oxygen scavenger (GLS 100, Mitsubishi Gas Chemicals). The batches were then stored at room temperature and accelerated temperature conditions.

**Table 11:** Formulation composition selected with different level of EDTA concentrations.

Ingredient	Quantity (mg/mL)				
	XIV	XV	XVI	XVII	XVIII
Formulation Number	XIV	XV	XVI	XVII	XVIII
Norepinephrine Bitartrate	0.016	0.032	0.064	0.064	0.064
Edetate Sodium	0.10	0.10	0.10	0.010	0.0010
Sodium chloride	9	9	9	9	9
Hydrochloric Acid/ Sodium Hydroxide	q.s. pH 4.0	q.s. pH 4.0	q.s. pH 4.0	q.s. pH 4.0	q.s. pH 4.0
Water for Injection	q.s. 1 mL	q.s. 1 mL	q.s. 1 mL	q.s. 1 mL	q.s. 1 mL

[0036] The resultant stability data on these formulations are presented in **Table 12 – Table 16** (CCS- Clear colorless solution; ND – Not Detected). The results of the stability studies at different amounts of EDTA at pH 4.0 indicates that both 0.01%, 0.001% of EDTA significantly prevented the degradation rate of norepinephrine in terms of known and unknown impurities. Moreover, with respect to isomerization from the *R*-isomer to the *S*-isomer it was notably observed that the amount of EDTA had substantially no influence on racemization or enantiomer formation during stability and after autoclaving.

**Table 12:** Stability study of Formulation XIV - Norepinephrine bitartrate injection (16  $\mu\text{g/mL}$ ); pH 4.0 at 100  $\mu\text{g/mL}$  EDTA.

Storage Condition	25 $\pm$ 2 °C/60 $\pm$ 5 % RH				40 $\pm$ 2 °C/75 $\pm$ 5 % RH		
	Initial	1 Month	2 Month	3 Month	1 Month	2 Month	3 Month
Time Point	Initial	1 Month	2 Month	3 Month	1 Month	2 Month	3 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	3.99	3.96	4.08	4.08	4.02	4.08	4.08
Assay	98.5	100.4	100.1	99.7	100.3	100.0	99.5
S-form	0.9	1.1	1.4	1.3	1.9	2.9	4.2
Total Impurities	0.05	ND	ND	ND	ND	0.10	0.38

**Table 13:** Stability study of Formulation XV - Norepinephrine bitartrate injection (32  $\mu\text{g/mL}$ ); pH 4.0 at 100  $\mu\text{g/mL}$  EDTA.

Storage Condition	25 $\pm$ 2 °C/60 $\pm$ 5 % RH				40 $\pm$ 2 °C/75 $\pm$ 5 % RH		
	Initial	1 Month	2 Month	3 Month	1 Month	2 Month	3 Month
Time Point	Initial	1 Month	2 Month	3 Month	1 Month	2 Month	3 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	4.01	3.99	4.08	4.08	4.02	4.08	4.08
Assay	101.0	102.9	97.1	100.7	102.9	99.4	100.6
S-form	0.9	1.1	1.3	1.4	1.9	3.0	4.1
Total Impurities	0.06	ND	ND	ND	ND	ND	0.14

**Table 14:** Stability study of Formulation XVI - Norepinephrine bitartrate injection (64  $\mu\text{g/mL}$ ); pH 4.0 at 100  $\mu\text{g/mL}$  EDTA.

Storage Condition	25 $\pm$ 2 °C/60 $\pm$ 5 % RH				40 $\pm$ 2 °C/75 $\pm$ 5 % RH		
	Initial	1 Month	2 Month	3 Month	1 Month	2 Month	3 Month
Time Point	Initial	1 Month	2 Month	3 Month	1 Month	2 Month	3 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	4.00	3.99	4.08	4.08	3.98	4.07	4.07
Assay	98.4	103.2	98.7	100.2	104.6	99.3	99.8
S-form	0.9	1.1	1.3	1.3	2.0	3.2	4.2
Total Impurities	0.06	ND	0.12	ND	ND	ND	ND

**Table 15:** Stability study of Formulation XVII - Norepinephrine bitartrate injection (64 µg/mL); pH 4.0 at 10 µg/mL EDTA.

Storage Condition	25 ± 2 °C/60 ± 5 % RH				40 ± 2 °C/75 ± 5 % RH		
	Initial	1 Month	2 Month	3 Month	1 Month	2 Month	3 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	4.00	3.98	4.06	4.06	3.99	4.05	4.05
Assay	102.7	105.7	103.4	104.3	107.8	103.6	103.9
S-form	0.9	1.1	1.2	1.5	2.0	3.3	4.3
Total	0.06	ND	ND	ND	ND	0.26	ND

**Table 16:** Stability study of Formulation XVIII - Norepinephrine bitartrate injection (64 µg/mL); pH 4.0 at 1 µg/mL EDTA.

Storage Condition	25 ± 2 °C/60 ± 5 % RH				40 ± 2 °C/75 ± 5 % RH		
	Initial	1 Month	2 Month	3 Month	1 Month	2 Month	3 Month
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	4.00	3.98	4.07	4.07	4.02	4.06	4.06
Assay	98.7	102.6	100.4	100.4	105.0	99.9	99.2
S-form	0.9	1.1	1.3	1.4	2.0	3.2	4.3
Total	0.06	ND	ND	ND	ND	ND	ND

[0037] Sterilization and Stability: The volume for ready-to-inject formulations is 250 mL and as such classifies as a large volume parenteral (LVP). To achieve a desired or required sterility assurance level of  $10^{-6}$  for a LVP terminal sterilization via heat it is typically required. The inventors therefore investigated whether or not contemplated formulations could be terminally sterilized via autoclaving.

**[0038]** Formulations at a concentration 16 µg/mL and 64 µg/mL (Formulation XVII) Norepinephrine base were prepared substantially as shown in **Table 11** above and packaged in secondary packaging of aluminum overwrap with an oxygen scavenger and shipped for terminal sterilization. The secondary packaging was removed and the bags were terminally sterilized using steam sterilizer (Fedegari, Model # FOB3) with an air over-pressure (AOP) sterilization cycle. The terminal sterilization was performed at 121 °C for 5, 10, and 15 min. Post completion of sterilization temperature, the bags underwent spontaneous cooling to 95 °C and forced cooling to 70 °C. The total exposure time and calculated F<sub>0</sub> values were 11.09, 17.04, and 22.42 for 5 min, 10 min, and 15 min cycles respectively. The bags were then analyzed for assay, impurities, and S-isoforn, and the results are shown in **Table 17** and **Table 18**.

**Table 17:** Stability study of Norepinephrine bitartrate injection (16 µg/mL) filled in 100 mL PP bags (pH 4.0); 10 µg/mL EDTA; terminally sterilized.

Time Point	Initial	5 Min			10 Min			15 Min		
Bag Number		1	2	3	1	2	3	1	2	3
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	3.76	3.85	3.78	3.77	3.76	3.76	3.78	3.76	3.75	3.76
Dissolved Oxygen	0.63	4.93	4.86	4.89	0.75	0.48	0.55	0.65	0.78	0.77
Assay	103.1	103.1	103.1	103.1	103.1	103.0	103.1	103.1	103.2	103.1
S-Form	1.0	3.0	3.0	3.0	3.8	3.7	3.7	4.3	4.3	4.3
Total	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

**Table 18:** Stability study of Norepinephrine bitartrate injection (64 µg/ml) filled in 100 mL PP bags (pH 4.0); 10 µg/mL EDTA terminally sterilized.

Time Point	Initial	5 Min			10 Min			15 Min		
Bag Number		1	2	3	1	2	3	1	2	3
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	3.74	3.74	3.75	3.73	3.74	3.74	3.76	3.74	3.73	3.74
Dissolved Oxygen	0.69	5.15	5.03	5.00	0.52	0.59	0.75	0.69	0.80	0.74
Assay	101.2	102.2	101.2	101.5	101.7	101.2	101.3	101.2	101.3	102.2
S-Form	1.0	3.0	3.0	3.0	3.7	3.7	3.7	4.3	4.3	4.3
Total	ND	ND	ND	ND	ND	ND	ND	0.1	ND	ND

[0039] As can be seen from the data, the *S*-isoform appears to increase proportionally to time during the terminal sterilization cycle. No increase in reportable impurities was observed.

[0040] Test method - Determination of norepinephrine and degradation products: Separation of Norepinephrine and related compounds was performed using a gradient HPLC method with UV detection. Pentofluorophenylpropyl terminated silica was used as a stationary phase for chromatographic analysis. The mobile phase was prepared by mixing water and methanol, with both solvents containing formic acid. Related compounds were defined by their relative retention times (RRT) based on the NE peak retention time. Quantitation of related compounds was accomplished by comparing the corresponding peak area from a sample solution chromatogram to that of the NE peak from a Reference Standard (RS) solution of a known concentration. Relative Response Factors (RRF) were used to correct for chemical structure effects on the responses of the identified impurities. Chromatography was performed using parameters and methods as shown in **Table 19**.

HPLC	Waters Alliance e2695		
Column	Supelco Discovery HS F-5 Column, 3 $\mu$ m, 4.6x150 mm		
Column Temperature	35 °C		
Sample Temperature	Ambient		
Injection volume	85.0 $\mu$ l.		
Flow Rate	0.8 mL/min		
Detection	Spectrum: 200-600 nm, resolution 1.2 nm Single channel: 280 nm, resolution 4.8 nm PDA Filter Time Constant: Normal Sampling rate: 5 points/sec		
Solution A	0.1% Formic acid in Water		
Solution B	0.1% Formic acid in Methanol		
Mobile Phase	Time (mins)	%Solution A	%Solution B
	0	100	0
	3	100	0
	6	93	7
	8	93	7
	15	88	12
	30	2	98
	35	2	98
	36	100	0
	40	100	0

Table 19

**[0041]** Test Method - Identification, Assay and Enantiomeric Purity of Norepinephrine: Identification and quantification of *S*-norepinephrine and *R*-norepinephrine was performed using an HPLC method with UV detection. HPLC-UV was used to separate and quantitate the amount of (*R*)- and (*S*)-enantiomers of norepinephrine (NE) present in the drug product with the NE concentrations of 16, 32 and 64  $\mu$ g/ml. The comparison of the sum of (*R*)- and (*S*)-peak responses in a sample chromatogram versus a reference standard chromatogram gives the total amount of NE. The (*S*)-enantiomer was quantitated based on its peak response as the percentage of the total peak response of both enantiomers.

**[0042]** More specifically, determination of R- and S-enantiomers of norepinephrine in the drug product solution was performed using an isocratic reverse-phase HPLC method with UV detection. Separation was achieved by using a protein-based column with functional chiral selectors. The chiral selector is cellobiohydrolase (CBH), a stable enzyme that has been immobilized onto spherical silica particles. This enzyme preferentially separates compounds containing one or more basic nitrogen groups together with one or more hydrogen-accepting or hydrogen-donating groups. Chromatography was performed using parameters and methods as shown in **Table 20**.

HPLC	Agilent 1260 Infinity
Column	Daicel Chiralpak CBH™ column, 5 µm, 4.0x100 mm
Column Temperature	27°C ± 2°C
Sample Temperature	Ambient
Injection volume	20.0 µL for 16 mcg/mL, 10.0 µL for 32 mcg/mL, 5.0 µL for 64 mcg/mL.
Flow Rate	0.9 mL/min
Detection	Single channel: 280 nm, resolution 4.8 nm Spectrum: 200-600 nm, resolution 1.2 nm
Mobile Phase:	Buffer/IPA 95:5 v/v Buffer: Sodium Phosphate, Disodium Edetate, pH 6.0
Run Time	8 min

Table 20

[0043] While contemplated formulations can be administered following various protocols, the inventors contemplate that administration of the formulations, especially administration for treatment of hypotension, will follow a protocol that comprises at least two distinct steps, with an accelerated administration followed by a maintenance administration as exemplarily described in Table 21 below.

Presentation (mg/mL)	Concentration (µg/mL)	Initial Dose		Maintenance Dose	
		Dose per Minute (µg/min)	Flow Rate (mL/min)	Dose per Minute (µg/min)	Flow Rate (mL/min)
16 µg/mL (4 mg in 250 mL)	16	8-12	0.500-0.750	2-4	0.125-0.250
32 µg/mL (8 mg in 250 mL)	32		0.250-0.375		0.062-0.125
64 µg/mL (16 mg in 250 mL)	64		0.125-0.187		0.031-0.062

Table 21

[0044] As used in the description herein and throughout the claims that follow, the meaning of “a,” “an,” and “the” includes plural reference unless the context clearly dictates otherwise.

Also, as used in the description herein, the meaning of “in” includes “in” and “on” unless the context clearly dictates otherwise.

[0045] In some embodiments, the numbers expressing quantities of ingredients, properties such as concentration, reaction conditions, and so forth, used to describe and claim certain embodiments of the invention are to be understood as being modified in some instances by the term “about.” Accordingly, in some embodiments, the numerical parameters set forth in the written description and attached claims are approximations that can vary depending upon the desired properties sought to be obtained by a particular embodiment. In some embodiments, the numerical parameters should be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of some embodiments of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as practicable. The numerical values presented in some embodiments of the invention may contain certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

[0046] It should be apparent, however, to those skilled in the art that many more modifications besides those already described are possible without departing from the inventive concepts herein. The inventive subject matter, therefore, is not to be restricted except in the spirit of the disclosure. One skilled in the art will recognize many methods and materials similar or equivalent to those described herein, which could be used in the practice of the present invention. Indeed, the present invention is in no way limited to the methods and materials described.

[0047] Moreover, in interpreting the disclosure all terms should be interpreted in the broadest possible manner consistent with the context. In particular the terms “comprises” and “comprising” should be interpreted as referring to the elements, components, or steps in a non-exclusive manner, indicating that the referenced elements, components, or steps can be present, or utilized, or combined with other elements, components, or steps that are not expressly referenced.

What is claimed is:

1. A ready-to-inject norepinephrine composition, comprising:
  - an aqueous acidic solution having a pH range of between 3.7 and 4.3, wherein the aqueous acidic solution comprises a metal ion chelating agent and a pharmaceutically acceptable salt;
  - wherein the aqueous acidic solution comprises a buffer, or wherein the aqueous acidic solution is free of buffer when the norepinephrine is norepinephrine bitartrate;
  - wherein the metal ion chelating agent is present in an amount of between 1  $\mu\text{g/ml}$  and 100  $\mu\text{g/ml}$ , and wherein the pharmaceutically acceptable salt is present in an amount of between 0.6 wt% and 1.2 wt%;
  - norepinephrine dissolved at a concentration of between 10  $\mu\text{g/ml}$  and 100  $\mu\text{g/ml}$ , wherein the norepinephrine is an *R*-isomer;
  - wherein the ready-to-inject norepinephrine composition is free of antioxidants;
  - and
  - wherein after storage at 25 °C and 60% relative humidity over at least three months equal or less than 10% of the *R*-isomer form isomerize to the *S*-isomer and equal or less than 5% of the total norepinephrine degrade to degradation products.
2. The composition of claim 1, wherein the aqueous acidic solution comprises a buffer having a concentration of between 5 mM and 20 mM.
3. The composition of claim 1, wherein the norepinephrine is norepinephrine bitartrate.
4. The composition of any one of claims 1 to 3, wherein the aqueous acidic solution has a pH of between 4.0 and 4.2.
5. The composition of any one of claims 1 to 4, wherein the metal ion chelating agent is a bicarboxylic acid, a tricarboxylic acid, or an aminopolycarboxylic acid.
6. The composition of claim 5, wherein the metal ion chelating agent is ethylenediaminetetraacetic acid (EDTA).

7. The composition of any one of claims 1 to 6, wherein the metal ion chelating agent is present at a concentration of between 10 µg/ml and 100 µg/ml.
8. The composition of any one of claims 1 to 7, wherein the pharmaceutically acceptable salt is sodium chloride.
9. The composition of any one of claims 1 to 8, wherein the composition is heat sterilized.
10. The composition of any one of claims 1 to 9, wherein the norepinephrine is present at a concentration of between 16 µg/ml and 64 µg/ml.
11. The composition of any one of claims 1 to 9, wherein the norepinephrine is present at a concentration of 16 µg/ml +/- 10%, 32 µg/ml +/- 10%, or 64 µg/ml +/- 10%.
12. The composition of any one of claims 1 to 11, wherein equal or less than 6% of the *R*-isomer form isomerize to the *S*-isomer.
13. The composition of any one of claims 1 to 12, wherein equal or less than 3.5% of the total norepinephrine degrade to degradation products.
14. The composition of any one of claims 1 to 13, wherein the composition has a dissolved oxygen concentration of equal or less than 1 ppm.
15. The composition of claim 1, wherein the aqueous acidic solution has a pH of between 3.7 and 4.0.
16. A method of preparing a ready-to-inject norepinephrine composition, comprising:
  - formulating a liquid parenteral composition that contains in an aqueous acidic solution norepinephrine as an *R*-isomer wherein:
    - (a) the formulation exhibits less than 10% of isomerization of the *R*-isomer to an *S*-isomer after three months of storage of the liquid composition at 25 °C and 60% relative humidity, and
    - (b) the formulation exhibits equal or less than 5% degradation of total norepinephrine after three months of storage of the liquid composition at 25 °C and 60% relative humidity;

wherein the aqueous acidic solution comprises a buffer, or wherein the aqueous acidic solution is free of buffer when the norepinephrine is norepinephrine bitartrate;

wherein the aqueous acidic solution has a pH range of between 3.7 and 4.3, and wherein the aqueous acidic solution comprises a metal ion chelating agent and a pharmaceutically acceptable salt;

wherein the total norepinephrine is present in the liquid parenteral composition at a concentration of between 10 µg/ml and 100 µg/ml; and

wherein the ready-to-inject norepinephrine composition is free of antioxidants.

17. The method of claim 16, wherein the norepinephrine is present at a concentration of 16 µg/ml +/- 10%, 32 µg/ml +/- 10%, or 64 µg/ml +/- 10%.

18. The method of any one of claims 16 and 17, wherein the aqueous acidic solution comprises a citrate buffer at a concentration of between 5 mM and 20 mM.

19. The method of any one of claims 16-18, wherein the norepinephrine is norepinephrine bitartrate.

20. The method of any one of claims 16-19, wherein the aqueous acidic solution has a pH of between 4.0 and 4.2.

21. The method of any one of claims 16-20, wherein the metal ion chelating agent is a bicarboxylic acid, a tricarboxylic acid, or an aminopolycarboxylic acid.

22. The method of claim 21, wherein the metal ion chelating agent is ethylenediaminetetraacetic acid (EDTA).

23. The method of any one of claims 16-22, wherein the metal ion chelating agent is present at a concentration of between 10 µg/ml and 100 µg/ml.

24. The method of any one of claims 16-23, wherein the pharmaceutically acceptable salt is sodium chloride.

25. The method of any one of claims 16-24, wherein the step of formulating the liquid parenteral composition uses deoxygenated water having dissolved oxygen at a concentration of equal or less than 1 ppm.

26. The method of any one of claims 16-25, further comprising a step of heat sterilizing the composition.
27. The method of any one of claims 16-26, further comprising a step of packaging the composition in a container, and placing the container in a second container that includes a metal-free oxygen scavenger.
28. The method of claim 27, wherein the container and/or the second container is configured to reduce light-mediated oxidation of the norepinephrine.
29. A method of preparing a ready-to-inject norepinephrine composition, comprising:  
preparing an aqueous solution having a pH range of between 3.7 and 4.3,  
wherein the aqueous solution comprises a metal ion chelating agent  
and a pharmaceutically acceptable salt;  
wherein the metal ion chelating agent is present in an amount of between 1  
 $\mu\text{g/ml}$  and 100  $\mu\text{g/ml}$ ; and adjusting tonicity with the pharmaceutically  
acceptable salt;  
dissolving norepinephrine at a concentration suitable for administration to a  
patient in need thereof, wherein the norepinephrine is an *R*-isomer;  
wherein the aqueous solution comprises a buffer, or wherein the aqueous  
solution is free of buffer when the norepinephrine is norepinephrine  
bitartrate;  
wherein after storage at 25 °C and 60% relative humidity over at least three  
months equal or less than 10% of the *R*-isomer form isomerize to the  
*S*-isomer and equal or less than 5% of the total norepinephrine degrade  
to degradation products;  
wherein the ready-to-inject norepinephrine composition is free of antioxidants;  
and  
heat sterilizing the composition.
30. The method of claim 29, wherein the norepinephrine is present at a concentration  
of between 10  $\mu\text{g/ml}$  and 100  $\mu\text{g/ml}$ .
31. The method of claim 29 or 30, wherein the norepinephrine is present at a  
concentration of 16  $\mu\text{g/ml}$  +/- 10%, 32  $\mu\text{g/ml}$  +/- 10%, or 64  $\mu\text{g/ml}$  +/- 10%.

32. The method of any one of claims 29-31, wherein the aqueous solution comprises a citrate buffer.
33. The method of claim 32, wherein the citrate buffer has a concentration of between 5 mM and 20 mM.
34. The method of any one of claims 29-33, wherein the aqueous solution has a pH of between 4.0 and 4.2.
35. The method of any one of claims 29-34, wherein the metal ion chelating agent is a bicarboxylic acid, a tricarboxylic acid, or an aminopolycarboxylic acid.
36. The method of claim 35, wherein the metal ion chelating agent is ethylenediaminetetraacetic acid (EDTA).
37. The method of any one of claims 29-36, wherein the metal ion chelating agent is present at a concentration of between 10 µg/ml and 100 µg/ml.
38. The method of any one of claims 29-37, wherein the pharmaceutically acceptable salt is sodium chloride.
39. The method of any one of claims 29-38, wherein the step of preparing the aqueous solution uses deoxygenated water having dissolved oxygen at a concentration of equal or less than 1 ppm.
40. The method of any one of claims 29-39, wherein the step of heat sterilizing the composition comprises autoclaving the composition for at least 5 minutes.
41. The method of any one of claims 29-40, further comprising a step of packaging the composition in a container, and placing the container in a second container that includes a metal-free oxygen scavenger.
42. The method of claim 41, wherein the container and/or the second container is configured to reduce light-mediated oxidation of the norepinephrine.

43. Use of the ready-to-inject composition as defined in any one of claims 1-15 for treating hypotension in a subject.
44. The use of claim 43, wherein the ready-to-inject composition is for administration at an initial dose per minute and at a second dose per minute, and wherein the initial dose per minute is greater than the second dose per minute.
45. The use of claim 44, wherein the initial dose per minute is a dose of between 8 and 12  $\mu\text{g}/\text{min}$ .
46. The use of claim 44 or 45, wherein the second dose per minute is a dose of between 2 and 4  $\mu\text{g}/\text{min}$ .
47. The use of claim 44, wherein the initial dose per minute is at least twice the second dose per minute.
48. The ready-to-inject composition as defined in any one of claims 1-15 for use in treating hypotension in a subject.
49. The ready-to-inject composition for use of claim 48, wherein the ready-to-inject composition is for administration at an initial dose per minute and at a second dose per minute, and wherein the initial dose per minute is greater than the second dose per minute.
50. The ready-to-inject composition for use of claim 49, wherein the initial dose per minute is a dose of between 8 and 12  $\mu\text{g}/\text{min}$ .
51. The ready-to-inject composition for use of claim 49 or 50, wherein the second dose per minute is a dose of between 2 and 4  $\mu\text{g}/\text{min}$ .
52. The ready-to-inject composition for use of claim 49, wherein the initial dose per minute is at least twice the second dose per minute.