



HU000026671T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 026 671**(13) **T2**

EURÓPAI SZABADALOM

SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 12 823510**(22) A bejelentés napja: **2012. 08. 17.**

(96) Az európai bejelentés bejelentési száma:

EP 20120823510

(97) Az európai bejelentés közzétételi adatai:

EP 2746265 A1 **2014. 06. 25.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2746265 B1 **2015. 11. 18.**(51) Int. Cl.: **C07D 235/06** (2006.01)**A61K 313/43** (2006.01)**A61K 314/16** (2006.01)**A61K 3141/84** (2006.01)**A61K 314/23** (2006.01)**A61K 314/28** (2006.01)**A61K 314/33** (2006.01)**A61K 3144/39** (2006.01)**A61K 314/54** (2006.01)**A61K 314/96** (2006.01)**A61K 3153/77** (2006.01)**A61P 1/02** (2006.01)**A61P 1/04** (2006.01)**A61P 1/18** (2006.01)**A61P 9/10** (2006.01)**A61P 11/06** (2006.01)**A61P 13/10** (2006.01)**C07D 231/56** (2006.01)**C07D 263/56** (2006.01)**C07D 307/79** (2006.01)**C07D 403/04** (2006.01)**C07D 405/04** (2006.01)**C07D 405/12** (2006.01)**C07D 409/12** (2006.01)

(86) A nemzetközi (PCT) bejelentési szám:

PCT/JP 12/070902

(87) A nemzetközi közzétételi szám:

WO 13024898

(30) Elsőbbségi adatok:

2011179134**2011. 08. 18.****JP**

(72) Feltaláló(k):

OTSU, Hironori, Matsubara-shi Osaka 580-0015 (JP)

(73) Jogosult(ak):

Nippon Shinyaku Co., Ltd., Kyoto-shi Kyoto
601-8550 (JP)

(74) Képviselő:

SBGK Szabadalmi Ügyvivői Iroda, Budapest(54) **Heterociklusos származékok, mint mikroszomális prosztaglandin E szintetáz inhibitorok (mPGEs)**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 746 265 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
18.11.2015 Bulletin 2015/47

(21) Application number: **12823510.8**

(22) Date of filing: **17.08.2012**

(51) Int Cl.:
C07D 235/06 ^(2006.01) **A61K 31/343** ^(2006.01)
A61K 31/416 ^(2006.01) **A61K 31/423** ^(2006.01)
A61K 31/428 ^(2006.01) **A61K 31/433** ^(2006.01)
A61K 31/454 ^(2006.01) **A61K 31/496** ^(2006.01)
A61P 1/02 ^(2006.01) **A61P 1/04** ^(2006.01)
A61P 1/18 ^(2006.01) **A61P 9/10** ^(2006.01)
A61P 11/06 ^(2006.01) **A61P 13/10** ^(2006.01)
C07D 231/56 ^(2006.01) **C07D 263/56** ^(2006.01)
C07D 307/79 ^(2006.01) **C07D 403/04** ^(2006.01)
C07D 405/04 ^(2006.01) **C07D 405/12** ^(2006.01)
C07D 409/12 ^(2006.01) **A61K 31/4184** ^(2006.01)
A61K 31/4439 ^(2006.01) **A61K 31/5377** ^(2006.01)

(86) International application number:
PCT/JP2012/070902

(87) International publication number:
WO 2013/024898 (21.02.2013 Gazette 2013/08)

(54) **HETEROCYCLIC DERIVATIVE AS MICROSOMAL PROSTAGLANDIN E SYNTHASE (mPGEs)
INHIBITOR**

HETEROCYCLISCHES DERIVAT ALS MICROSOMAL PROSTAGLANDIN E SYNTHASE (mPGEs)
HEMMER

DÉRIVÉ HÉTÉROCYCLIQUE COMME INHIBITEUR DE LA SYNTHASE DE LA PROSTAGLANDINE
E MICROSOmale (mPGEs).

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **18.08.2011 JP 2011179134**

(43) Date of publication of application:
25.06.2014 Bulletin 2014/26

(73) Proprietor: **Nippon Shinyaku Co., Ltd.
Kyoto-shi
Kyoto 601-8550 (JP)**

(72) Inventor: **OTSU, Hironori
Matsubara-shi
Osaka 580-0015 (JP)**

(74) Representative: **Vossius & Partner
Patentanwälte Rechtsanwälte mbB
Siebertstrasse 3
81675 München (DE)**

(56) References cited:
**WO-A1-2008/071944 WO-A1-2009/103778
WO-A1-2011/048004 WO-A2-2006/099379**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[Field of the Invention]

5 **[0001]** The present invention relates to novel heterocyclic derivatives.

[Background Art]

10 **[0002]** Prostaglandins (PG) are produced abundantly in the site of inflammation and involved in progress of inflammation. Prostaglandin production is initiated with the release of arachidonic acid from membrane glycerophospholipid by phospholipase A2, and the arachidonic acid is then converted into prostaglandin H2 (PGH2) by cyclooxygenase (COX). PGH2 is converted into prostaglandins including prostaglandin E2 (PGE2), prostaglandin F2 α (PGF2 α), prostaglandin D2 (PGD2), prostaglandin 12 (PGI2), and thromboxane A2 (TXA2). These prostaglandins are known to have various physiological or pathophysiological activities, including inflammation inducing effect. Especially, PGE2 is known as an inflammatory inducer in acute and chronic inflammations and further known to induce pyrexia and hyperpathia. Non-steroidal anti-inflammatory drugs (NSAIDs) and selective COX-2 inhibitors have anti-inflammatory effect via reduced production of PGE2 based on their COX-1 and/or COX-2 inhibitory effect. PGE2 synthase (PGES) catalyzes the final step of the synthetic pathway of PGE2, which is an inflammatory mediator. To date, three subtypes of PGES, microsomal prostaglandin E synthase-1 (mPGES-1) [for example, Non-Patent Document 1], mPGES-2 [for example, Non-Patent Document 2] and cytosolic prostaglandin synthase (cPGES) [for example, Non-Patent Document 3] are known. Among these, mPGES-1, in the same manner as COX-2, is primarily induced during inflammation and plays a major part in PGE2 production in inflammatory lesion. On the other hand, cPGES is constitutively expressed PGES and coupled to COX-1 to play a part in basal PGE2 production [for example, Non-Patent Document 4]. As to mPGES-2, it is a subject of controversy as there is a report that it can be coupled to both COX isoforms. The studies in mPGES-1 deficient mouse suggest that mPGES-1 contributes to pathological progress in various inflammation models, such as acetic acid writhing model (e.g., Non-Patent Document 5), arthritis model (e.g., Non-Patent Documents 5 and 6), multiple sclerosis model (e.g., Non-Patent Document 7), fever model (e.g., Non-Patent Document 8). Also, mPGES-1 inhibitors specifically inhibit COX-2-dependent PGE2 production, and therefore, they are expected to reduce various side effects, compared with NSAIDs or COX-2 inhibitors. It is believed that elevated risk of cardiovascular events by COX-2 inhibitors is attributed to enhanced coagulation system and vasoconstriction via inhibited COX-2-dependent PGI2 production (e.g., Non-Patent Document 9). In contrast, mPGES-1 inhibitors are believed not to increase the risk of cardiovascular events, which is a problem with COX-2 inhibitors, since they do not inhibit PGI2 production (e.g., Non-Patent Document 10). mPGES-1 inhibitors are expected to serve as a safe anti-inflammatory agent by inhibiting only PGE2 production, which participates in inflammation. Thus, a pharmaceutical agent which is able to inhibit mPGES-1, and reduce PGE2 production is useful in the treatment or prevention of a disease, such as an inflammatory disease in which mPGES-1 participates. mPGES-1 inhibitors have been disclosed in the patent applications by NovaSAID AB (Patent Documents 1 and 2) and Boehringer Ingelheim International GmbH (Patent Document 3).

[Prior Art Documents]

[Patent Documents]

[0003]

[Patent Document 1] WO2009/103778

[Patent Document 2] US2010/0324086

[Patent Document 3] WO2011/048004

[Non-Patent Documents]

[0004]

[Non-Patent Document 1] Jakobsson et al., Proc. Natl. Acad. Sci. USA, 1999, 96, 7220-7225

[Non-Patent Document 2] Tanikawa et al., Biochem. Biophys. Res. Commun., 2002, 291, 884-889

[Non-Patent Document 3] Tanioka et al., J. Biol. Chem., 2000, 275, 32775-32782

[Non-Patent Document 4] Murakami et al., J. Biol. Chem., 2000, 275, 32783-32792

[Non-Patent Document 5] Kamei et al., J. Biol. Chem., 2004, 279, 33684-33695

[Non-Patent Document 6] Kojima et al., J. Immunol., 2008, 108, 3861-3868

- [Non-Patent Document 7] Kimura et al., Proc. Natl. Acad. Sci. USA, 2000, 106, 21807-21812
- [Non-Patent Document 8] Engblom et al., Nat. Neurosci., 2003, 6, 1137-1138
- [Non-Patent Document 9] Foudi et al., Cardiovasc. Res., 2009, 81, 269-277
- [Non-Patent Document 10] Cheng et al., J. Clin. Invest., 2006, 116, 1391-1399
- 5 [Non-Patent Document 11] J. Biol. Chem., 2004, 279(13), 12647-12658
- [Non-Patent Document 12] Biomed. Pharmacother., 2011, 65(1), 77-84
- [Non-Patent Document 13] Acta. Med. Okayama, 2008, 62(6), 373-378
- [Non-Patent Document 14] Proc. Natl. Acad. Sci. USA, 2009, 106, 21807-21812
- [Non-Patent Document 15] J. Biol. Chem., 279(32), 33684-33695
- 10 [Non-Patent Document 16] J. Immunol., 2008, 180, 8361-8368
- [Non-Patent Document 17] Glia, 2011, 59, 208-218
- [Non-Patent Document 18] J. Burn. Care. Res., 2011, 32(1), 79-90
- [Non-Patent Document 19] J. Biol. Chem., 2003, 278(21), 19396-19405
- [Non-Patent Document 20] Oncogene, 2012, 31(24), 2943-2952
- 15 [Non-Patent Document 21] Cancer Res., 2008, 68(9), 3251-3259
- [Non-Patent Document 22] Proc. Natl. Acad. Sci. USA, 2006, 103, 14507-14512
- [Non-Patent Document 23] Proc. Natl. Acad. Sci. USA, 2006, 103(31), 11790-11795
- [Non-Patent Document 24] J. Rheumatol., 2005, 32(5), 887-895
- [Non-Patent Document 25] J. Pharmacol. Exp. Ther., 2008, 326, 754-763
- 20 [Non-Patent Document 26] Kidney Int., 2011, 79(1), 77-88
- [Non-Patent Document 27] J. Immunol., 2012, 188, 4093-4102
- [Non-Patent Document 28] Am. J. Physiol. Lung Cell Mol. Physiol., 2004, 287, L981-L991
- [Non-Patent Document 29] Am. J. Physiol. Lung Cell Mol. Physiol., 2005, 288, L1010-L1016
- [Non-Patent Document 30] J. Immunol., 2012, 188, 4093-4102
- 25 [Non-Patent Document 31] Oncogene, 2012, 31(24), 2943-2952
- [Non-Patent Document 32] Arthritis Res. Ther., 2011, 13, R6

[Summary of the Invention]

30 [Problem to be solved by the invention]


[0005] The present invention provides a novel heterocycle derivative or a pharmaceutically acceptable salt thereof. Also, the invention provides with a pharmaceutical composition containing such heterocycle derivative or a pharmaceutically acceptable salt thereof as an active ingredient.

35 [Means for solving the problem]

[0006] The present invention is based on the inventors' discovery that the novel heterocycle derivative or a pharmaceutically acceptable salt thereof, as described below, has an excellent mPGES-1 inhibiting activity.




40 **[0007]** The present invention provides a heterocycle derivative represented by the general formula [1] or its tautomer (hereinafter referred to as "compound of the invention"), or a pharmaceutically acceptable salt thereof.

[Chemical Formula 1]



wherein ring A is a group represented by the general formulae [2], [3] or [4]:

[Chemical Formula 2]



wherein

X¹ is NH, N-alkyl, or O;

A¹ is hydrogen or alkyl;

A² is

i) hydrogen;

ii) halogen;

iii) alkyl optionally substituted with one to three groups selected from the group consisting of halogen, amino, monoalkylamino, dialkylamino, carbamoyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, a saturated cyclic aminocarbonyl, alkoxy, alkoxyalkoxy and alkylcarbonyloxy;

iv) cycloalkyl optionally substituted with alkyl optionally substituted with one to three halogens;

v) alkoxy;

vi) a saturated heterocycle group optionally substituted with alkyl, alkyloxycarbonyl, alkylcarbonyl or oxo;


vii) alkylthio;

viii) alkylsulfonyl;

ix) alkylsulfinyl;

x) a group represented by the general formula [5]:

[Chemical Formula 3]



wherein R^3 and R^4 are the same or different group selected from

- a) hydrogen,
- b) alkyl optionally substituted with a group selected from the group consisting of monoalkylamino, dialkylamino, a saturated cyclic amino optionally substituted with alkyl, a saturated heterocycle group optionally substituted with alkyl, alkoxy, hydroxycarbonyl, hydroxyl, alkyloxycarbonyl and alkylthio, or
- c) cycloalkyl; or

xi) a saturated cyclic amino optionally substituted with alkyl, amino, monoalkylamino, dialkylamino, alkoxy or hydroxyl;

R^1 is phenyl, benzyl, naphthyl, cycloalkyl, cycloalkylmethyl, heteroaryl, heteroaryl-methyl, 1,2,3,4-tetrahydronaphthalen-5-yl, 1,2,3,4-tetrahydronaphthalen-6-yl, 2,3-dihydro-1H-inden-4-yl, 2,3-dihydro-1H-inden-5-yl, 1,2-dihydrocyclobutabenzen-3-yl, 1,2-dihydrocyclobutabenzen-4-yl or alkyl,

wherein said phenyl, benzyl, cycloalkyl, cycloalkylmethyl, heteroaryl and heteroaryl-methyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen,
- ii) alkyl optionally substituted with one to three groups selected from the group consisting of halogen, hydroxy and phenyl,
- iii) alkoxy,
- iv) hydroxy, and
- v) cyano;

R^2 is phenyl or pyridyl,

wherein said phenyl and pyridyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen,
- ii) alkylsulfonyl,
- iii) alkoxy optionally substituted with one to three halogens or alkoxy;
- iv) alkynyl optionally substituted with alkoxyalkyl or cycloalkyl, and
- v) alkyl optionally substituted with one to three groups selected from the group consisting of alkoxy, alkoxyalkoxy, cycloalkyl, phenyl and halogen.

[0008] A preferred embodiment of the invention is any one of the following (A) to (C), or a pharmaceutically acceptable salt thereof:

(A) a compound of the invention or a pharmaceutically acceptable salt thereof, wherein the ring A is a group of formula [4] and X^1 is NH;

(B) a compound of the invention or a pharmaceutically acceptable salt thereof, wherein R^1 is phenyl, 1,2,3,4-tetrahydronaphthalen-5-yl, 1,2,3,4-tetrahydronaphthalen-6-yl, 2,3-dihydro-1H-inden-4-yl, 2,3-dihydro-1H-inden-5-yl,

1,2-dihydrocyclobutabenzen-3-yl, or

1,2-dihydrocyclobutabenzen-4-yl, and said phenyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen,
- ii) alkyl optionally substituted with one to three halogens,
- iii) alkoxy, and
- iv) cyano

; or

(C) a compound of the invention a pharmaceutically acceptable salt thereof, wherein R² is phenyl and said phenyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen
- ii) alkylsulfonyl,
- iii) alkoxy optionally substituted with alkoxy,
- iv) alkynyl optionally substituted with alkoxyalkyl or cycloalkyl, and
- v) alkyl optionally substituted with one to three groups selected from the group consisting of halogen, alkoxy, alkoxyalkoxy, cycloalkyl and phenyl

[0009] A more preferred embodiment of the invention is a compound of the invention or a pharmaceutically acceptable salt thereof, wherein

the ring A is a group of formula [4],

X¹ is NH,

A² is

- i) hydrogen,
- ii) alkyl optionally substituted with a group selected from the group consisting of halogen, monoalkylamino, dialkylamino, monoalkylaminocarbonyl, dialkylaminocarbonyl, a saturated cyclic aminocarbonyl, alkoxy, alkoxyalkoxy and alkylcarbonyloxy,
- iii) cycloalkyl optionally substituted with alkyl optionally substituted with one to three halogens,
- iv) alkoxy,
- v) a saturated heterocyclic group optionally substituted with alkyl or alkyloxycarbonyl,
- vi) alkylthio,
- vii) alkylsulfonyl,
- viii) alkylsulfinyl,
- ix) amino substituted with alkyl wherein said alkyl is optionally substituted with a group selected from the group consisting of monoalkylamino, dialkylamino, a saturated cyclic amino optionally substituted with alkyl, tetrahydrofuryl, morpholino, alkoxy, hydroxycarbonyl, hydroxyl and alkylthio,
- x) amino substituted with cycloalkyl or xi) a saturated cyclic amino optionally substituted with alkyl, dialkylamino, alkoxy or hydroxyl, and

R¹ is

- i) phenyl optionally substituted with one to three groups selected from the group consisting of halogen, alkyl optionally substituted with one to three halogens, alkoxy and cyano,
- ii) 1,2,3,4-tetrahydronaphthalen-5-yl,
- iii) 2,3-dihydro-1H-inden-5-yl,
- iv) benzyl optionally substituted with halogen or alkyl optionally substituted with one to three halogens,
- v) cycloalkyl,
- vi) cycloalkylmethyl,
- vii) naphthyl,
- viii) pyridylmethyl optionally substituted with alkyl optionally substituted one to three halogens,
- ix) thienyl,
- x) thienylmethyl,
- xi) benzothiazolyl,
- xii) benzothiadiazolyl,

- xiii) indolyl or
- xiv) alkyl, and

R² is phenyl or pyridyl

wherein said phenyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen,
- ii) alkylsulfonyl,
- iii) alkoxy optionally substituted with alkoxy,
- iv) alkynyl optionally substituted with alkoxyalkyl or cycloalkyl, and
- v) alkyl optionally substituted with one to three groups selected from the group consisting of halogen, alkoxy, alkoxy-alkoxy, cycloalkyl and phenyl, and

said pyridyl is optionally substituted with halogen

[0010] A further more preferred embodiment of the invention is a compound of the invention or a pharmaceutically acceptable salt thereof, wherein

the ring A is a group of formula [4],

X¹ is NH,

A² is alkyl substituted with alkoxy, dialkylamino, tetrahydrofuryl, tetrahydrofurylmethyl, alkoxyalkylamino, or cycloalkyl optionally substituted with alkyl optionally substituted with one to three halogens,

R¹ is phenyl substituted with one halogen and one methyl, and R² is phenyl optionally substituted with one trifluoromethyl or two halogens

[0011] A particularly preferred embodiment of the invention is any one of the following (1)-(239), or its tautomer or a pharmaceutically acceptable salt thereof:

- (1) N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (2) N-cyclohexyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (3) N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (4) N-[(1-hydroxycyclohexyl)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (5) N-[2-(trifluoromethyl)benzyl]-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (6) N-cyclohexyl-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (7) N-(3-chloro-2-methylphenyl)-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (8) N-cyclohexyl-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-indazole-7-carboxamide,
- (9) N-[2-(trifluoromethyl)benzyl]-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-indazole-7-carboxamide,
- (10) N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (11) 2-methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (12) N-cyclohexyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (13) N-(3-chloro-2-methylphenyl)-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (14) N-cyclopentyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (15) N-cyclobutyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (16) N-(3-chloro-2-methylphenyl)-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (17) N-cyclohexyl-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (18) 2-ethyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (19) N-cyclohexyl-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (20) 2-(methoxymethyl)-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (21) 2-(methoxymethyl)-N-(2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (22) 2-(methoxymethyl)-N-(4-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-

carboxamide,

(23) N-(2-chlorobenzyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(24) 2-(methoxymethyl)-N-(4-methylbenzyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(25) N-(4,4-difluorocyclohexyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(26) N-(4-tert-butylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(27) 2-(methoxymethyl)-N-[4-(trifluoromethyl)phenyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(28) N-(2,4-dimethylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(29) N-(2-chloro-4-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(30) N-(3,4-dimethylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(31) N-(3-chloro-4-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(32) N-(2,3-dihydro-1H-inden-5-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(33) 2-(methoxymethyl)-N-(5,6,7,8-tetrahydronaphthalen-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(34) N-(2-fluorophenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(35) 2-(methoxymethyl)-N-(2-methoxyphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(36) 2-(methoxymethyl)-N-(4-methoxyphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(37) N-(3-bromo-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(38) N-(3-chloro-2-methylbenzyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(39) N-(2,6-difluorophenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(40) N-(3-cyano-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(41) 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-N-({[3-(trifluoromethyl)pyridin-2-yl]methyl})-1H-benzimidazole-4-carboxamide,

(42) N-(2-chloro-6-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(43) 2-(2-amino-2-oxoethyl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(44) 2-(2-amino-2-oxoethyl)-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(45) N-(3-chloro-2-methylphenyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(46) N-cyclohexyl-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(47) 1-methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(48) N-(3-chloro-2-methylphenyl)-1-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(49) N-cyclohexyl-1-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(50) 1-ethyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(51) N-(3-chloro-2-methylphenyl)-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide,

(52) 2-methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide,

boxamide,

(53) N-(3-chloro-2-methylphenyl)-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide,

(54) N-(3-chloro-2-methylphenyl)-2-ethoxy-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(55) 2-ethoxy-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(56) N-(3-chloro-2-methylphenyl)-2-(1-chloro-2-methylpropan-2-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(57) N-(3-chloro-2-methylphenyl)-2-[(dimethylamino)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(58) N-(3-chloro-2-methylphenyl)-2-(2-methylpropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(59) 2-(2-methylpropyl)-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(60) tert-butyl 3-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}azetidine-1-carboxylate,

(61) N-(3-chloro-2-methylphenyl)-2-[(methylamino)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(62) 4-{[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}methyl acetate,

(63) N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(64) 2-[[[2(R)-tetrahydrofuran-2-yl]-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(65) N-(3-chloro-2-methylphenyl)-2-[(2S)-tetrahydrofuran-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(66) 2-[[[2(S)-tetrahydrofuran-2-yl]-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(67) 2-(1-acetylazetidine-3-yl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(68) tert-butyl (2S)-2-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}pyrrolidine-1-carboxylate,

(69) tert-butyl (2R)-2-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}pyrrolidine-1-carboxylate,

(70) N-(3-chloro-2-methylphenyl)-2-[(2S)-pyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(71) N-(3-chloro-2-methylphenyl)-2-[(2S)-1-methylpyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(72) 2-[[[2(S)-1-acetylpyrrolidin-2-yl]-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(73) N-(3-chloro-2-methylphenyl)-2-[(2-methoxyethoxy)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(74) N-(3-chloro-2-methylphenyl)-2-(1-methoxy-2-methylpropan-2-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(75) 2-tert-butyl-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(76) 2-tert-butyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-N-[[3-(trifluoromethyl)pyridin-2-yl]methyl]-1H-benzimidazole-4-carboxamide,

(77) N-(3-chloro-2-methylphenyl)-2-(2-ethoxyethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(78) N-(3-chloro-2-methylphenyl)-2-(ethoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(79) 2-(ethoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-N-[[3-(trifluoromethyl)pyridin-2-yl]methyl]-1H-benzimidazole-4-carboxamide,

(80) N-(3-chloro-2-methylphenyl)-2-(2-methoxyethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(81) N-(3-chloro-2-methylphenyl)-2-(2,2-dimethylpropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benz-

imidazole-4-carboxamide,

(82) N-(3-chloro-2-methylphenyl)-2-cyclopropyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(83) N-(3-chloro-2-methylphenyl)-2-(2-methylpentan-2-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(84) N-(3-chloro-2-methylphenyl)-2-(1-methylcyclopropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(85) 2-tert-butyl-N-(3-chloro-4-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(86) 2-tert-butyl-N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(87) 2-tert-butyl-N-(3-chloro-4-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(88) N-(3-chloro-2-methylphenyl)-2-[1-(trifluoromethyl)cyclopropyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(89) N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(90) N-(2-chlorobenzyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(91) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(92) 6-({[2-chloro-4-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-methoxymethyl-1H-benzimidazole-4-carboxamide,

(93) 6-({[2-chloro-5-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(94) N-(3-chloro-2-methylphenyl)-6-({[2-chlorophenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(95) N-(3-chloro-2-methylphenyl)-6-({[2-chloropyridin-3-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(96) 6-({[2-bromophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(97) N-(3-chloro-2-methylphenyl)-6-({[2,6-dichlorophenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(98) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(99) 6-({[2-chloro-3-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(100) 6-({[2-chloro-3,6-difluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(101) 6-({[2-bromo-6-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(102) 6-({[2-bromo-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(103) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-6-methylphenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(104) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-4-methylphenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(105) 6-({[5-bromo-2-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(106) 6-({[2-bromo-5-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(107) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-5-methylphenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(108) N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[5-methyl-2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(109) 6-({[2,5-bis(trifluoromethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(110) 6-({[2,4-bis(trifluoromethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

imidazole-4-carboxamide,

(111) N-(3-chloro-2-methylphenyl)-6-({[5-fluoro-2-(trifluoromethyl)phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(112) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-6-(trifluoromethyl)phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(113) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-5-[2-(propan-2-yloxy)ethoxy]phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(114) 6-({[2-chloro-5-(2-ethoxyethoxy)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(115) 6-({[2-chloro-5-(3-methoxypropyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(116) 6-({[5-(3-tert-butoxyprop-1-yn-1-yl)-2-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(117) 6-({[5-(3-tert-butoxypropyl)-2-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(118) 6-({[2-chloro-5-(3-hydroxy-3-methylbutyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(119) 6-({[2-chloro-5-(ethoxymethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(120) 6-({[2-chloro-5-[(2-ethoxyethoxy)methyl]phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(121) 6-({[2-chloro-5-(2-cyclopropylethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(122) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-5-(2-phenylethyl)phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(123) N-(3-chloro-2-methylphenyl)-2-cyclopentyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(124) N-(3-chloro-2-methylphenyl)-2-cyclopentyl-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(125) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-cyclopentyl-1H-benzimidazole-4-carboxamide,

(126) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(127) N-(3-chloro-2-methylphenyl)-6-({[2,6-dichlorophenyl]carbonyl}amino)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(128) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(129) N-(3-chloro-2-methylphenyl)-2-[(2S)-5-oxopyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(130) N-(3-chloro-2-methylphenyl)-2-[(2R)-5-oxopyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(131) N-(3-chloro-2-methylphenyl)-2-[2-oxo-2-(pyrrolizin-1-yl)ethyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(132) N-(3-chloro-2-methylphenyl)-2-[2-(dimethylamino)-2-oxoethyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(133) N-(3-chloro-2-methylphenyl)-2-[2-(methylamino)-2-oxoethyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(134) 2-chloro-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(135) N-(3-chloro-2-methylphenyl)-2-[(2-methoxyethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(136) N-(3-chloro-2-methylphenyl)-2-[(2-hydroxyethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(137) N-(3-chloro-2-methylphenyl)-2-(methylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(138) N-(3-chloro-2-methylphenyl)-2-(ethylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(139) N-(3-chloro-2-methylphenyl)-2-[(2,2-dimethylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-

1H-benzimidazole-4-carboxamide,

(140) N-(3-chloro-2-methylphenyl)-2-(cyclopentylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(141) N-(3-chloro-2-methylphenyl)-2-(piperidin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(142) N-(3-chloro-2-methylphenyl)-2-(4-methylpiperazin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(143) 2-bis(2-hydroxyethyl)amino-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(144) N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(145) N-(3-chloro-2-methylphenyl)-2-([2-(morpholin-4-yl)ethyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(146) N-(3-chloro-2-methylphenyl)-2-([2-(dimethylamino)ethyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(147) N-(3-chloro-2-methylphenyl)-2-(3-hydroxyazetidin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(148) N-(3-chloro-2-methylphenyl)-2-[(3S)-3-(dimethylamino)pyrrolizin-1-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(149) N-(3-chloro-2-methylphenyl)-2-[(3S)-3-hydroxypyrrolizin-1-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(150) N-(3-chloro-2-methylphenyl)-2-([2-(diethylamino)ethyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(151) N-(3-chloro-2-methylphenyl)-2-([2-(pyrrolizin-1-yl)ethyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(152) N-(3-chloro-2-methylphenyl)-2-([3-(dimethylamino)propyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(153) N-(3-chloro-2-methylphenyl)-2-([3-(dimethylamino)-2,2-dimethylpropyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(154) N-(3-chloro-2-methylphenyl)-2-([2-(dipropylamino)ethyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(155) N-(3-chloro-2-methylphenyl)-2-(morpholin-4-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(156) 2-amino-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(157) N-(3-chloro-2-methylphenyl)-2-([3-(hydroxy-2,2-dimethylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(158) N-(3-chloro-2-methylphenyl)-2-([3-methoxy-2,2-dimethylpropyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(159) tert-butyl N-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-2-yl}glycinate,

(160) N-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-2-yl}glycine,

(161) N-(3-chloro-2-methylphenyl)-2-([3-(hydroxy-2,2-dimethylpropyl)amino]-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(162) N-(3-chloro-2-methylphenyl)-2-([3-methoxy-2,2-dimethylpropyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(163) N-(3-chloro-2-methylphenyl)-2-(pyrrolizin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(164) 2-(azetidin-1-yl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(165) N-(3-chloro-2-methylphenyl)-2-(3-methoxyazetidin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(166) N-(3-chloro-2-methylphenyl)-2-([2-(hydroxy-2-methylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(167) N-(3-chloro-2-methylphenyl)-2-([(2S)-tetrahydrofuran-2-ylmethyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(168) N-(3-chloro-2-methylphenyl)-2-([(2R)-tetrahydrofuran-2-ylmethyl]amino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

onyl}amino)-1H-benzimidazole-4-carboxamide,

(169) N-(3-chloro-2-methylphenyl)-2-(((2S)-1-hydroxy-3-methylbuta n-2-yl)amino)-6-({[2-

(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(170) N-(3-chloro-2-methylphenyl)-2-(((2R)-1-hydroxy-3-methylbuta n-2-yl)amino)-6-({[2-

(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(171) N-(3-chloro-2-methylphenyl)-2-(((2S)-1-hydroxy-3,3-dimethyl butan-2-yl)amino)-6-({[2-(trifluoromethyl)phe-
nyl]carbonyl}a mino)-1H-benzimidazole-4-carboxamide,

(172) N-(3-chloro-2-methylphenyl)-2-[(3-methoxy-2,2-dimethylpropy l)(methyl)amino]-6-({[2-(trifluoromethyl)phe-
nyl]carbonyl}am ino)-1H-benzimidazole-4-carboxamide,

(173) N-(3-chloro-2-methylphenyl)-2-[(3-methoxypropyl)amino]-6-({ [2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-
benzimidazole -4-carboxamide,

(174) N-(3-chloro-2-methylphenyl)-2-[[2-(propan-2-yloxy)ethyl]ami no]-6-({[2-(trifluoromethyl)phenyl]carbonyl}ami-
no)-1H-benzi midazole-4-carboxamide,

(175) 2-[(2-tert-butoxyethyl)amino]-N-(3-chloro-2-methylphenyl)-6 -({[2-(trifluoromethyl)phenyl]carbonyl}amino)-
1H-benzimidaz ole-4-carboxamide,

(176) N-(3-chloro-2-methylphenyl)-2-[(2-methoxy-2-methylpropyl)am ino]-6-({[2-(trifluoromethyl)phenyl]carbon-
yl}amino)-1H-benz imidazole-4-carboxamide,

(177) N-(3-chloro-2-methylphenyl)-2-[[2-(methylsulfanyl)ethyl]ami no]-6-({[2-(trifluoromethyl)phenyl]carbonyl}ami-
no)-1H-benzi midazole-4-carboxamide,

(178) N-(3-chloro-2-methylphenyl)-2-(methylsulfanyl)-6-({[2-(trif luoromethyl)phenyl]carbonyl}amino)-1H-benzimi-
dazole-4-carbo xamide,

(179) N-(3-chloro-2-methylphenyl)-2-(methylsulfonyl)-6-({[2-(trif luoromethyl)phenyl]carbonyl}amino)-1H-benzimi-
dazole-4-carbo xamide,

(180) N-(3-chloro-2-methylphenyl)-2-(methylsulfinyl)-6-({[2-(trif luoromethyl)phenyl]carbonyl}amino)-1H-benzimi-
dazole-4-carbo xamide,

(181) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimida-
zole-4-carboxam ide,

(182) N-(3-chloro-2-methylphenyl)-6-[[[(2,6-dichlorophenyl)carbony l]amino]-2-(dimethylamino)-1H-benzimidazole-
4-carboxamide,

(183) N-(3-chloro-2-methylphenyl)-6-[[[(2,4-dichlorophenyl)carbony 1]amino]-2-(dimethylamino)-1H-benzimida-
zole-4-carboxamide,

(184) N-(3-chloro-2-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbony 1]amino]-2-(dimethylamino)-1H-benzimida-
zole-4-carboxamide,

(185) 6-[[[(2-bromo-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-m ethylphenyl)-2-(dimethylamino)-1H-benzimi-
dazole-4-carboxami de,

(186) 6-[[[(2-bromo-6-chlorophenyl)carbonyl]amino]-N-(3-chloro-2-m ethylphenyl)-2-(dimethylamino)-1H-benzimi-
dazole-4-carboxami de,

(187) 6-[[[2-chloro-5-(cyclopropylethynyl)phenyl]carbonyl]amino)-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-
1H-benzimidaz ole-4-carboxamide,

(188) N-(3-chloro-2-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbony l]amino]-2-[(3-hydroxy-2,2-dimethylpropyl)ami-
no]-1H-benzimi dazole-4-carboxamide,

(189) N-(3-chloro-2-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbony l]amino]-2-[(3-methoxy-2,2-dimethylpropyl)ami-
no]-1H-benzimi dazole-4-carboxamide,

(190) N-(3-chloro-2-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbony l]amino]-2-[(2-hydroxy-2-methylpropyl)amino]-
1H-benzimidazo le-4-carboxamide,

(191) N-(3-chloro-2-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbony l]amino]-2-[(2-methoxy-2-methylpropyl)amino]-
1H-benzimidazo le-4-carboxamide,

(192) N-(3-chloro-2-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbony l]amino]-2-[[2-(propan-2-yloxy)ethyl]amino]-
1H-benzimidazol e-4-carboxamide,

(193) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2 -methylphenyl)-2-[[2-(propan-2-yloxy)ethyl]ami-
no]-1H-benzim idazole-4-carboxamide,

(194) 2-[(2-tert-butoxyethyl)amino]-6-[[[(2-chloro-6-fluorophenyl) carbonyl]amino]-N-(3-chloro-2-methylphenyl)-1H-
benzimidazol e-4-carboxamide,

(195) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2 -methylphenyl)-2-[(3-methoxy-2,2-dimethylpro-
pyl)amino]-1H-b enzimidazole-4-carboxamide,

(196) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2 -methylphenyl)-2-[(2-methoxy-2-methylpro-
pyl)amino]-1H-benzi midazole-4-carboxamide,

(197) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2 -methylphenyl)-2-[[[(2S)-tetrahydrofuran-2-ylme-

thyl]amino}-1 H-benzimidazole-4-carboxamide,
 (198) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-[[2-(2R)-tetrahydrofuran-2-ylmethyl]amino]-1 H-benzimidazole-4-carboxamide,
 (199) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide,
 (200) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-[[2-(2S)-1-hydroxy-3-methylbutan-2-yl]amino]-1H-benzimidazole-4-carboxamide,
 (201) N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (202) N-(4-tert-butylphenyl)-2-(dimethylamino)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (203) N-(2,3-dihydro-1H-inden-5-yl)-2-(dimethylamino)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (204) 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,
 (205) N-(3-chloro-4-methylphenyl)-6-[[[(2,6-dichlorophenyl)carbonyl]amino]-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,
 (206) N-(3-chloro-4-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbonyl]amino]-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,
 (207) N-(3-chloro-2-methylphenyl)-2-cyclopropyl-6-[[[(2,5-dichlorophenyl)carbonyl]amino]-1H-benzimidazole-4-carboxamide,
 (208) N-(3-chloro-4-methylphenyl)-2-cyclopropyl-6-[[[(2,5-dichlorophenyl)carbonyl]amino]-1H-benzimidazole-4-carboxamide,
 (209) N-(3-chloro-2-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbonyl]amino]-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxamide,
 (210) N-(3-chloro-4-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbonyl]amino]-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxamide,
 (211) N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-[[2-(methylsulfonyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (212) N-(3-chloro-4-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbonyl]amino]-2-(2-methoxyethyl)-1H-benzimidazole-4-carboxamide,
 (213) 2-(methoxymethyl)-N-phenyl-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (214) 2-(methoxymethyl)-N-propyl-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (215) 2-(methoxymethyl)-N-(pyridin-3-yl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (216) N-benzyl-2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (217) N-(cyclohexylmethyl)-2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (218) 2-(methoxymethyl)-N-(naphthalen-1-yl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (219) 2-(methoxymethyl)-N-(thiophen-3-yl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (220) N-(2,1,3-benzothiadiazol-4-yl)-2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (221) N-(1,1-dioxide-1-benzothiophen-6-yl)-2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (222) 2-(methoxymethyl)-N-(thiophen-2-ylmethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (223) N-(1H-indol-5-yl)-2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (224) N-(1,3-benzothiazol-2-yl)-2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (225) N-(2,2-dimethylpropyl)-2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,
 (226) 2-(methoxymethyl)-N-(thiophen-2-yl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-

carboxamide,

(227) N-(5-chloro-1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(228) N-(2-benzylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(229) 2-(methoxymethyl)-N-(quinolin-8-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(230) N-(cycloheptylmethyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(231) N-(1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(232) N-(6-chloro-1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(233) N-[3-chloro-2-(hydroxymethyl)phenyl]-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(234) N-(3-chloro-2-methylphenyl)-6-({[3-fluoropyridin-2-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(235) N-(3-chloro-2-methylphenyl)-6-({[3-chloropyridin-4-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(236) N-(3-chloro-2-methylphenyl)-6-({[3,5-dichloropyridin-4-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(237) 6-({[5-butoxy-2-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(238) 6-({[2-chloro-5-(2,2-difluoroethoxy)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(239) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-5-(4,4,4-trifluorobutoxy)phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide.

DETAILED DESCRIPTION OF THE INVENTION

[0012] Detail description of the terms used in the present specification is provided as follows.

[0013] Examples of "halogen" include fluorine, chlorine, bromine and iodine.

[0014] Examples of "alkyl" include straight or branched alkyl having 1 to 8 carbon atoms, such as, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, n-hexyl, isohexyl, n-heptyl, isoheptyl, n-octyl. Among which alkyl having 1 to 6 carbon atoms is preferred, and alkyl having 1 to 3 carbon atoms is more preferred.

[0015] The alkyl moiety of "monoalkylamino", "dialkylamino", "monoalkyl aminocarbonyl", "dialkylaminocarbonyl", "alkylcarbonyloxy", "alkyloxycarbonyl", "alkylcarbonyl", "alkylthio", "alkylsulfonyl", "alkylsulfinyl", "alkoxyalkyl", "monohaloalkyl", "dihaloalkyl", "trihaloalkyl" and "alkoxyalkylamino" is as defined above for "alkyl".

[0016] Examples of "alkoxy" include straight or branched alkoxy having 1 to 8 carbon atoms, such as, for example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, t-butoxy, n-pentyloxy, n-hexyloxy, n-heptyloxy, n-octyloxy.

[0017] The alkoxy moiety of "alkoxyalkoxy", "alkoxyalkyl" and "alkoxyalkylamino" is as defined above for "alkoxy".

[0018] Examples of "heteroaryl" include monocyclic or bicyclic aromatic rings having 1 to 3 hetero atoms selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom. Specific examples include furyl (e.g., 2-furyl, 3-furyl), thienyl (e.g., 2-thienyl, 3-thienyl), pyrrolyl (e.g., 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl), imidazolyl (e.g., 1-imidazolyl, 2-imidazolyl, 4-imidazolyl), pyrazolyl (e.g., 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl), triazolyl (e.g., 1,2,4-triazol-1-yl, 1,2,4-triazol-3-yl, 1,2,4-triazol-4-yl), tetrazolyl (e.g., 1-tetrazolyl, 2-tetrazolyl, 5-tetrazolyl), oxazolyl (e.g., 2-oxazolyl, 4-oxazolyl, 5-oxazolyl), isoxazolyl (e.g., 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl), oxadiazolyl (e.g., 1,3,4-oxadiazol-2-yl), thiazolyl (e.g., 2-thiazolyl, 4-thiazolyl, 5-thiazolyl), thiadiazolyl (e.g., 1,3,4-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,3-thiadiazolyl), isothiazolyl (e.g., 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl), pyridyl (e.g., 2-pyridyl, 3-pyridyl, 4-pyridyl), pyridazinyl (e.g., 3-pyridazinyl, 4-pyridazinyl), pyrimidinyl (e.g., 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl), pyrazinyl (e.g., 2-pyrazinyl), benzothiadiazolyl (e.g., 1,2,3-benzothiadiazol-4-yl, 1,2,3-benzothiadiazol-5-yl, 2,1,3-benzothiadiazol-4-yl, 2,1,3-benzothiadiazol-5-yl), benzothiazolyl (e.g., benzothiazol-2-yl, benzothiazol-4-yl, benzothiazol-5-yl, benzothiazol-6-yl, benzothiazol-7-yl), indolyl (e.g., indol-3-yl and indol-4-yl, indol-5-yl, indol-6-yl, indol-7-yl), benzothiophenyl (e.g., 1-benzothiophen-2-yl, 1-benzothiophen-3-yl, 1-benzothiophen-4-yl, 1-benzothiophen-5-yl, 1-benzothiophen-6-yl, 1-benzothiophen-7-yl), 1,1-dioxo-1-benzothiophenyl (e.g., 1,1-dioxo-1-benzothiophen-2-yl, 1,1-dioxo-1-benzothiophen-3-yl, 1,1-dioxo-1-benzothiophen-4-yl, 1,1-dioxo-1-benzothiophen-5-yl, 1,1-dioxo-1-benzothiophen-6-yl, 1,1-dioxo-1-benzothiophen-7-yl), quinolyl (quinolin-2-yl, quinolin-3-yl, quinolin-4-yl, quinolin-5-yl, quinolin-6-yl, quinolin-7-yl, quinolin-8-yl) and

1,3-benzoxazol-2-yl.

[0019] The heteroaryl moiety of "heteroaryl methyl" is as defined above for "heteroaryl".

[0020] Examples of "a saturated cyclic amino" include 4- to 7-membered saturated cyclic amino groups having one or two nitrogen atoms, said ring optionally having one oxygen or sulfur atom and optionally substituted with oxo. Specific examples include 1-azetidiny, 1-pyrrolidiny, 1-imidazolidiny, piperidino, 1-piperaziny, 1-tetrahydropyrimidiny, 4-morpholino, 4-thiomorpholino, 1-homopiperaziny, and 2-oxo-oxazolidin-3-yl.

[0021] The saturated cyclic amino moiety of "a saturated cyclic aminocarbonyl" is as defined above for "a saturated cyclic amino".

[0022] Examples of "a saturated heterocycle group" include 4- to 6-membered saturated heterocycle group having one nitrogen or oxygen atom in the ring. Specific examples include 2-pyrrolidiny, 3-pyrrolidiny, 2-piperidiny, 3-piperidiny, 4-piperidiny, 2-oxetany, 3-oxetany, 2-tetrahydrofurany, and 3-tetrahydrofurany.

[0023] Examples of "cycloalkyl" include cycloalkyl having 3 to 8 carbon atoms. Specific examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

[0024] The cycloalkyl moiety of "cycloalkyl methyl" is as defined above for "cycloalkyl".

[0025] Examples of "naphthyl" include 1-naphthyl and 2-naphthyl.

[0026] Examples of "pyridyl" include 2-pyridyl, 3-pyridyl and 4-pyridyl.

[0027] Examples of "alkynyl" include straight or branched alkynyl having 2 to 6 carbon atoms. Specific examples include ethynyl, 1-propynyl, 1-butynyl, 1-pentynyl, 2-propynyl, 2-butynyl, 3-butynyl, 1-methyl-2-propynyl, 2-pentynyl, 3-pentynyl and 4-pentynyl.

[Mode for Carrying Out the Invention]

[0028] The compound of the invention can be prepared according to the following procedures, working examples or procedures known in the art. If the starting material has a substituent group that may interfere with a reaction during the process, it may be protected with an appropriate protecting group according to known method before subjecting to the reaction.

[0029] The following abbreviations can be used herein to simplify the description.

p-: para-,

t-: tert-,

s-: sec-,

m-: meta-,

THF: tetrahydrofuran,

DMF: N,N-dimethylformamide,

DMA: N,N-dimethylacetamide,

HBTU: O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate,

HATU: O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate,

TFA: trifluoroacetic acid,

DME: ethylene glycol dimethyl ether,

NMP: N-methyl-2-pyrrolidone,

DMSO: dimethyl sulfoxide,


MeOH: methanol,

EtOH: ethanol.

Process 1

[0030]

[Formula 4]




wherein ring A, R¹ and R² are as defined above.

[0031] This reaction is the condensation of Compound [6] with Compound [7] and can be performed according to a method for condensation reaction known per se. A compound of the invention can be synthesized by the reaction of a carboxylic acid Compound [6] or its reactive derivative with an amine derivative [7]. Examples of such reactive derivative of Compound [6] include those commonly used in an amide condensation formation, such as, for example, acid halides (e.g., acid chloride, acid bromide), mixed anhydrides, imidazolides, active amides. When using Compound [6], the reaction can be performed using a condensing agent at a temperature in the range from -20°C to 100°C in the presence or absence of a base. Examples of the condensing agent which may be used for this reaction include 1,1'-oxalyldiimidazole,

1-ethyl-3-(3-dimethylaminopropyl)carbodiimide, dicyclohexylcarbodiimide, diethyl cyanophosphonate, HBTU, HATU, 1H-benzotriazol-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate. Examples of the base which can be used for this reaction include organic bases such as triethylamine, N,N-diisopropylethylamine, N,N-dimethylaniline, pyridine, 1,8-diazabicyclo[5,4,0]-7-undecene. Any solvent may be used so long as it does not interfere with the reaction, and examples of such solvent include ethers such as THF, 1,4-dioxane and diethyl ether, amides such as DMF and DMA, nitriles such as acetonitrile and propionitrile, hydrocarbons such as benzene and toluene, halogenated hydrocarbons such as chloroform and methylene chloride, and mixed solvents thereof. Also, an additive agent can be used if necessary. Examples of such additive agent which can be used include 1-hydroxybenzotriazol and 1-hydroxy-7-aza-benzotriazol. Preferred reaction time is generally within the range from 10 minutes to 24 hours, but it should vary depending on the starting material, the reaction temperature, etc. Preferred amounts of Compound [7] and the condensing agent to be used are within the range from 1 to 3 moles for one mole of Compound [6]. Preferred amount of the base to be used is within the range from 1 Eq to 10 Eq, preferably from 1 Eq to 4 Eq, to Compound [6].

[0032] For example, Compound [6] can be prepared according to the process as described below.

[Formula 5]



wherein ring A and R² are as defined above, X represents halogen and R⁵ represents alkyl.

Step 1-A

[0033] This reaction is a condensation reaction of Compound [8A] with Compound [9A] to synthesize Compound [10] according to Process 1 as described above.

Step 1-B


[0034] The reaction is a coupling reaction of Compound [8B] with Compound [9B] using palladium catalyst, and the reaction is performed according to a method known per se. The solvent which can be used is not limited so long as it does not interfere with a reaction, and examples of such solvent include hydrocarbons such as toluene and xylene, ethers such as 1,4-dioxane and THF, amides such as DMF, DMA and NMP, and a mixed solvent thereof. The reaction is performed in the presence of a base at a temperature in the range from 20°C to 200°C, optionally using microwave. Examples of the palladium catalyst which can be used include tris(dibenzylideneacetone)(chloroform)dipalladium (0), tris(dibenzylideneacetone)dipalladium (0) and palladium acetate (II). Suitable amount of such palladium catalyst is within the range from 0.001 mol to 0.3 mol, to 1 mol of aryl halide. Examples of the ligand for the palladium catalyst which can be used include 1,1'-bis(diphenylphosphino)ferrocene, 4,5-bis(diphenylphosphino)-9,9'-dimethylxanthene, 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl, (±)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, 2-(di-*t*-butylphosphino)biphenyl, bis[2-(diphenylphosphino)phenyl] ether, tri-*t*-butylphosphine, etc. Examples of the base which can be used include sodium *t*-butoxide, tripotassium phosphate and cesium carbonate. The amount of the base to be used is within the range from 1 Eq to 10 Eq, preferably within the range from 1 Eq to 4 Eq, to Compound [8B]. Appropriate reaction time is generally within the range from 10 minutes to 24 hours, but it should vary depending on the starting material, the reaction temperature, etc.

Step 2

[0035] Compound [6] can be prepared by hydrolyzing Compound [10] according to a known method. The reaction is usually performed in the presence of acid or base in a suitable solvent. Examples of the acid used in the hydrolysis include inorganic acids such as hydrochloric acid and sulfuric acid, and examples of the base include inorganic bases such as sodium hydroxide and potassium hydroxide. Examples of reaction solvent include alcohols such as MeOH and EtOH, ethers such as THF and dioxane, water, and mixed solvents thereof. The reaction is performed at a temperature within the range from 0°C to 100°C, and the reaction time is usually within the range from 30 minutes to 24 hours.

[0036] For example, Compound [8A] can be prepared by the following process.

[Formula 6]




wherein ring A and R^5 are as defined above.

[0037] This reaction is a reduction reaction of Compound [11] to aromatic amine, and the reaction can be carried out with a conventional method. For example, the reaction is achieved by catalytic hydrogen reduction of Compound [11] using a catalyst such as Raney nickel, palladium, rhodium, platinum, etc., in a suitable solvent under hydrogen gas atmosphere, hydride reduction using lithium aluminum hydride, etc., iron reduction using reduced iron reagent and ammonium chloride etc., or zinc reduction using zinc dust and acetic acid, etc. In addition, there are also a method using sulfides such as hydrosulfite sodium and a reducing method by ammonium formate, hydrazine, etc. with a metal catalyst such as palladium on carbon. Selection of the solvent depends on the kind of compound or reagent to be used, and the solvent may be used alone or as a mixture thereof. Examples of such solvent include toluene, THF, 1,4-dioxane, 1,2-dimethoxyethane, ethyl acetate, acetone, acetonitrile, DMF, or alcohols such as MeOH, EtOH and tert-butanol, and water. Although the reaction temperature depends on the kind of compound and reagent to be used, it is usually within the range from 0°C to 300°C, preferably within the range from 20°C to 150°C.

[0038] When the ring A of Compound [8A] is a group represented by the general formula [2] or [3], the compound can be prepared according to the method described in WO2008/65508. When the ring A of Compound [8A] is a group represented by the general formula [4], the compound can be prepared according to a method described in literature (e.g., EP2226315, J. Org. Chem., 1960, 25,942, etc.). Also, the compound may be prepared according to the following process.

[Formula 7]



wherein X^1 , R^5 and A^2 are as defined above, and X^2 represents NH_2 or OH .

Step 1

[0039] This reaction is an acylation reaction of Compound [12] with Compound [13] or its reactive derivative and can be performed according to method known per se as acylation reaction. Examples of the reactive derivative of Compound [13] include those generally used for acylation reaction and include acid halides (e.g., acid chloride, acid bromide), mixed acid anhydrides, imidazolides, active amides, etc. When using Compound [13], the reaction can be conducted using a condensing agent in the presence or absence of a base at temperature within a range from -20°C to 100°C. Examples of the condensing agent which can be used for this reaction include 1,1'-oxalyldiimidazole, 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide, dicyclohexylcarbodiimide, diethyl cyanophosphonate, HBTU, HATU and 1H-benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate. Examples of the base which can be used for this reaction include organic bases such as triethylamine, N,N-diisopropylethylamine, N,N-dimethylaniline, pyridine, 1,8-diazabicyclo[5,4,0]-7-undecene. The solvent to be used is not limited so long as it does not interfere with a reaction, and examples of such solvent include ethers such as THF, 1,4-dioxane and diethyl ether, amides such as DMF and DMA, nitriles such as acetonitrile and propionitrile, hydrocarbons such as benzene and toluene, halogenated hydrocarbons such as chloroform and methylene chloride, and a mixed solvent thereof. Also, an additive agent can be used if necessary. Examples of such additive agent which can be used include 1-hydroxybenzotriazol and 1-hydroxy-7-aza-benzotriazol. Preferred amount of the base to be used is within the range from 1 Eq to 10 Eq, preferably 1 Eq to 4 Eq, to Compound [13]. Preferred reaction time is generally within the range from 10 minutes to 24 hours, but it should vary depending on the starting material, the reaction temperature, etc. Preferred amounts of Compound [13] and the condensing agent to be used are within the range from 1 to 3 moles for one mole of Compound [12].

Step 2

[0040] This reaction is intramolecular cyclization of Compound [14] using an acid catalyst and can be performed according to a method known per se. The reaction is performed in a suitable solvent or in the absence of a solvent, and examples of the acid which can be used include hydrochloric acid, p-toluenesulfonic acid, acetic acid, pyridinium p-toluenesulfonate, polyphosphoric acid, phosphoryl chloride, etc. The reaction can be carried out usually at a temperature within the range from 0°C to 200°C.

The solvent which can be used is not limited so long as it does not interfere with a reaction, and examples of such solvent includes hydrocarbons such as toluene and xylene, alcohols such as MeOH and EtOH, ethers such as 1,4-dioxane and THF, amides such as DMF and DMA, halogenated hydrocarbon such as chloroform and dichloromethane, acetonitrile, or a mixed solvent thereof. The reaction time is generally within the range from 30 minutes to 48 hours although it should vary depending on the starting material and the reaction temperature.

Step 3

[0041] This reaction is a nitration reaction of Compound [15] and can be performed according to a known method. Generally, a nitrating agent, such as nitric acid, mixed acid, metal nitrate, acetyl nitrate, dinitrogen pentaoxide, nitronium salt (e.g., nitronium tetrafluoro borate, nitronium trifluoromethanesulfonate), etc., is used. The reaction solvent is not limited so long as it does not interfere with a reaction, and example of such solvent include halogenated solvents such as dichloromethane and chloroform, pentane, TFA, sulfolane, acetonitrile, etc.


The reaction is carried out under neutral or acidic condition. When the reaction is carried out under acidic condition, examples of the acid to be used include sulfuric acid, nitric acid, acetic acid, and acetic anhydride. Also, only such acid may be used as a reaction solvent without using of the reaction solvent as mentioned above. The reaction temperature is generally within the range from -20°C to room temperature although it should vary depending on the compound and reagent to be used. Preferred reaction time is generally within the range from 30 minutes to 24 hours although it should vary depending on the starting material and the reaction temperature.

Step 4

[0042] This reaction is a reduction reaction of Compound [16] to an aromatic amine and can be carried out to prepare Compound [8A'] according to the similar process of the preparation of Compound [8A].

[0043] Compound [8B] can be prepared according to the method described in literature (e.g., J. Med. Chem., 1999, 42, 5020, WO2008/65508). When the ring A of Compound [8B] is a group represented by the general formula [4], the compound also can be prepared by the following process.

[Formula 8]



15 wherein A^2 , X , X^1 , X^2 and R^5 are as defined above.

Step 1

20 **[0044]** This reaction is a condensation reaction of Compound [17] with Compound [13] or its reactive derivative and can be carried out to prepare Compound [18] according to the similar procedure of Step 1 in the process for Compound [8A'].


Step 2

25 **[0045]** This reaction is an intramolecular cyclization reaction of acyl group and amino group using an acid catalyst and can be carried out to prepare Compound [8B'] according to the procedure of Step 2 in the above-mentioned process of the preparation of Compound [8A'].


[0046] For example, Compound [17] can be prepared according to the following process.

[Formula 9]

30 $X^2 = -NH_2$



45 $X^2 = -OH$



wherein X and R^5 are as defined above.

Step 1-1

[0047] This reaction is a halogenation reaction of Compound [19] and can be carried out according to a method known per se as halogenation reaction. Examples of halogenating agent which can be used include N-bromosuccinimide, N-iodosuccinimide, bromine, iodine, etc., and generally, the reaction can be carried out at a temperature within the range from 0°C to 200°C. The solvent which can be used is not limited so long as it does not interfere with a reaction, and examples of such solvent include hydrocarbons, such as toluene and xylene, ethers such as 1, 4-dioxane and THF, amides such as DMF and DMA, halogenated hydrocarbons such as chloroform and dichloromethane, acetonitrile, and a mixed solvent thereof. Also a suitable base may be added if necessary, and examples of such base which can be used include pyridine, N, N-diisopropylethylamine, etc. The amount of the base to be used, for example, is within the range from 1 Eq to 10 Eq, preferably within the range from 1 Eq - 4 Eq, to Compound [19]. Preferred reaction time is generally within the range from 30 minutes to 24 hours although it should vary depending on the starting material and the reaction temperature.

Step 1-2

[0048] This reaction is a reduction reaction of Compound [20] to an aromatic amine, and can be performed to prepare Compound [17A] according to the similar process of the above-mentioned Compound [8A].


Step 2-1

[0049] This reaction is a halogenation reaction of Compound [21] and can be performed to prepare Compound [17B] according to the similar procedure of Step 1-1 in the process of the preparation of Compound [17].

Process 2

[0050]

[Formula 10]




wherein ring A, R¹ and R² are as defined above.

[0051] This reaction is a condensation reaction of Compound [22A] with Compound [9A] and can be performed according to Process 1 to prepare a compound of the invention.

[0052] Also, a compound of the invention may be prepared by the following procedures.

[Formula 11]




15 wherein ring A, R¹, R² and X are as defined above.

[0053] This reaction is a coupling reaction of Compound [22B] with Compound [9B], and can be performed to prepare a compound of the invention according to the similar procedure of Step 1-B in the process of Compound [6].

[0054] Compound [22A] can be prepared according to the following process, for example.

20 [Formula 12]



wherein ring A and R¹ are as defined above.

Step 1

35 **[0055]** This reaction is a hydrolysis reaction of Compound [11] as starting material and can be performed to prepare Compound [23] according to the similar procedure of Step 2 in the process of the preparation of Compound [6].

Step 2


40 **[0056]** This reaction is a condensation reaction of Compound [23] with Compound [7] and can be performed to prepare Compound [24] according to Process 1.

Step 3

45 **[0057]** This reaction is a reduction reaction of Compound [24] to an aromatic amine and can be performed to prepare Compound [22A] according to the process of the preparation of Compound [8A].

[0058] For example, Compound [22B] can also be prepared by the following process.

[Formula 13]




wherein ring A, R¹ and X are as defined above.

[0059] This reaction is a condensation reaction of Compound [25] with Compound [7], and can be performed to prepare Compound [22B] according to Process 1. Compound [25] can be prepared according to a method described in literature (e.g., EP234872B1; US6387938B1; Bioorg. Med. Chem., 1999, 7, 2271; J. Med. Chem., 1999, 42, 5020), and can also be prepared by hydrolyzing Compound [8B] according to the similar procedure of Step 2 in the process of the preparation of Compound [6].

Process 3

[0060] When the ring A is a group represented by the general formula [4] and A² is alkylthio, alkoxy or a group represented by the general formula [5], the compound can be prepared according to the following process.

[Formula 14]




wherein R¹, R² and X¹ are as defined above, and A⁰ represents halogen, mesylate, tosylate, etc., and A³ represents alkylthio, alkoxy or a group represented by the general formula [5].

[0061] This reaction is a nucleophilic substitution reaction of Compound [26] by amines, metal alkoxy or substituted thiol and can be carried out according to known method. The reaction is carried out in a suitable solvent or in the absence of a solvent, using an excessive amount of reagents, or in the presence of a base. Examples of suitable base to be used include pyridine, triethylamine, N,N-diisopropylethylamine, potassium carbonate, sodium hydrogencarbonate, etc. The solvent to be used is not limited so long as it does not interfere with a reaction, and examples of such solvent include ethers such as THF and diethyl ether, amides such as DMF and DMA, nitriles such as acetonitrile and propionitrile, hydrocarbons such as benzene and toluene, alcohols such as MeOH and EtOH, water, or a mixed solvent thereof. Also, an excessive amount of the amine may be used in the reaction instead of the solvent. The reaction is generally carried out at a temperature within the range from 0°C to 200°C although it should depend on the compound and reagents to be used. The amount of the base is, for example, within the range from 1 Eq to 10 Eq, preferably within the range from 1 Eq to 4 Eq, to Compound [26]. Preferred reaction time is generally within the range from 30 minutes to 24 hours although it should vary depending on the starting material and the reaction temperature.

[0062] Compound [26] can be prepared, for example, according to the following process.

[Formula 15]



wherein X^1 , R^1 , R^2 , R^5 and A^0 are as defined above.

Step 1

[0063] This reaction is a nitration reaction of Compound [27] as a starting material and can be performed according to the similar procedure of Step 3 in the process for the preparation of Compound [8A']. Compound [27] can be prepared according to a method described in literature (e.g., WO2006/116412; J. Med. Chem., 1993, 36, 2182).

Step 2

[0064] This reaction is a reduction reaction of Compound [28] to an aromatic amine and can be performed to prepare Compound [29] according to the similar procedure of the process for the preparation of Compound [8A].

Step 3

[0065] This reaction is a condensation reaction of Compound [29] with Compound [9A] and can be performed to prepare Compound [30] according to the similar procedure in Process 1.

Step 4

[0066] This reaction is a hydrolysis reaction of Compound [30] and can be performed to prepare Compound [31] according to similar procedure in Step 2 in the preparation of Compound [6].


Step 5

[0067] This reaction is a condensation reaction of Compound [31] with Compound [7] and can be performed to prepare Compound [26] according to similar procedure in Process 1.

Process 4

[0068] The compound also can be prepared by the following process in case where the ring A is a group represented by the general formula [4] and A is alkylsulfinyl, alkylsulfonyl.

[Formula 16]



wherein R^1 , R^2 , and X^1 are as defined above, A^4 represents alkylthio and A^5 represents alkylsulfinyl, alkylsulfonyl.

[0069] This reaction is an oxidation reaction of Compound [32] as a starting material and can be carried out according to a conventional method. For example, the reaction is achieved in a suitable solvent using a peroxide such as potassium permanganate, metachloro perbenzoic acid and oxone permonosulfate. The solvent should be selected according to the kind of starting material and not limited so long as it does not interfere with a reaction, and examples of such solvent include dichloromethane, chloroform, dichloroethane, THF, 1,4-dioxane, DME, toluene, MeOH, etc., and such solvent may be used alone or as a mixed solvent. The amount of the oxidizing agent is within the range from 0.5 Eq to 10 Eq, preferably within the range from 0.9 Eq to 3 Eq, to the starting material. The reaction temperature is usually within the range from -20°C to 80°C , preferably within the range from 0°C to 50°C although it should depend on the kinds of the compound and reagent to be used. Preferred reaction temperature is usually within the range from 30 minutes to 24 hours although it depends on the kind of starting material to be used and the reaction temperature.

[0070] The compound of the invention may be used as a pharmaceutical as it is, and also a pharmaceutically acceptable salt thereof formed according to known method may be used. Examples of such salt include inorganic salt of acids such as hydrochloric acid, hydrobromic acid, sulfuric acid and phosphoric acid, and organic salt of acids such as acetic acid, citric acid, tartaric acid, maleic acid, succinic acid, fumaric acid, p-toluenesulfonic acid, benzenesulfonic acid, and methanesulfonic acid.


[0071] For example, a hydrochloride of a compound of the invention can be formed by dissolving the compound of the invention in a solution of hydrogen chloride in alcohol, ethyl acetate or diethyl ether.

[0072] The compound of the invention may have an asymmetrical carbon, and each of such optical isomers and a mixture thereof are within the scope of the present invention. Such optical isomer can be prepared by optical resolution from a racemic mixture as obtained in the following working examples, according to known method using an optically active acid such as tartaric acid, dibenzoyltartaric acid, mandelic acid, 10-camphor sulfonic acid, etc., or by the use of an optically active compound previously prepared as a starting material. Alternatively, such compound may be prepared by optical resolution using a chiral column or asymmetric synthesis.

[0073] Some of the compounds of the invention may exist as tautomers, and each of such tautomers and a mixture thereof are within the scope of the invention.

[0074] For example, the heterocycle derivative represented by the general formula [1] (i.e., heterocycle derivative represented by the following general formula [1X]), wherein the ring A is a group represented by the general formula [4] and X^1 is NH, may form a heterocycle derivative represented by the following general formula [1XA].

[Formula 17]



wherein R¹, R², and A² are as defined above.

[0075] As shown in the test examples as described below, the compound of the invention or a pharmaceutically acceptable salt thereof has mPGES-1 inhibiting activity. Also, since the compound of the invention or a pharmaceutically acceptable salt thereof has mPGES-1 inhibiting activity, it has PGE2 inhibitory effect, analgesic action and anti-inflammatory effect.

[0076] Therefore, the compound of the invention or a pharmaceutically acceptable salt thereof can be used as a preventing or treating agent for diseases that involve mPGES-1, diseases associated with PGE2, and diseases on which effectiveness is expected based on analgesic or anti-inflammatory action.

[0077] Examples of disease which can apply a compound of the invention or a pharmaceutically acceptable salt thereof include inflammatory bowel disease (e.g., see Non-Patent Document 11), irritable bowel syndrome, migraine, headache, low back pain, spinal stenosis, herniated disk, temporomandibular joint disorders, cervical syndrome, cervical spondylosis, endometriosis (e.g., see Non-Patent Document 12), adenomyosis, preterm labour and delivery, threatened premature delivery, dysmenorrhea, overactive bladder, nocturia (e.g., see Non-Patent Document 13), interstitial cystitis, neurodegenerative disease such as Alzheimer's disease, multiple sclerosis (e.g., see Non-Patent Document 14), psoriasis, rheumatoid arthritis (e.g., see Non-Patent Document 15 and Non-Patent Document 16), rheumatic fever, fibromyalgia, neuralgia (e.g., see Non-Patent Document 17), complex regional pain syndrome, fascial dysfunction, viral infections such as influenza, common cold, zoster and AIDS, bacterial infection, mycosis, burn (e.g., see Non-Patent Document 18), inflammation and pain after operation, injury and dental extraction, malignant tumors such as colon cancer, breast cancer, lung cancer, prostatic cancer, etc. (e.g., see Non-Patent Document 19, Non-Patent Document 20, and Non-Patent Document 21), atherosclerosis (e.g., see Non-Patent Document 22), stroke (e.g., see Non-Patent Document 23), gout, arthritis, osteoarthritis (e.g., see Non-Patent Document 24 and Non-Patent Document 25), juvenile arthritis, ankylosing spondylitis, tenosynovitis, ligament ossification, systemic lupus erythematosus, vasculitis, pancreatitis, nephritis (e.g., see Non-Patent Document 26), conjunctivitis, iritis, scleritis, uveitis, wound therapy, dermatitis, eczema, osteoporosis, asthma (e.g., see Non-Patent Document 27), chronic obstructive pulmonary disease (e.g., see Non-Patent Document 28), pulmonary fibrosis (e.g., see Non-Patent Document 29), allergic disease (e.g., see Non-Patent Document 30), familial adenomatous polyposis (e.g., see Non-Patent Document 31), scleroderma (e.g., see Non-Patent Document 32), bursitis, leiomyoma of uterus, prostatitis, and a pain from cancer.

[0078] When administered as a pharmaceutical, a compound of the invention or a pharmaceutically acceptable salt thereof is administered as it is or as a pharmaceutical composition containing, for example, 0.001% to 99.5%, preferably 0.1% to 90%, in a pharmaceutically acceptable non-toxic and inactive carrier to a mammal including human.

[0079] In the pharmaceutical composition, a diluent in the form of a solid, a semi-solid or a liquid, a bulking agent, and one or more of other formulation additives can be used as a carrier. Preferably, the pharmaceutical composition of the invention is administered in a unit dosage form. The pharmaceutical composition may be administered by intra-tissue administration, oral administration, intravenous administration, local administration such as dermal administration, ocular instillation, intraperitoneal administration, intrathoracic administration, etc., or transrectal administration. Of course, the composition should be administered in a dosage form suitable for these administration routes.

[0080] The dosage as a medicament should be adjusted preferably in consideration of conditions of the patient such as age, body weight, nature and severity of the disease, route of administration, the compound of the invention to be administered, whether such compound is a salt or not, and the kind of such salt. For oral administration, a daily dosage of the compound of the invention or a pharmaceutically acceptable salt thereof as an active ingredient for adult is generally within the range from 0.01 mg to 5 g, preferably 1 mg to 500 mg for adult human. However, a lower dosage under said range may be sufficient in some cases, or a higher dosage over the said range may be needed in other cases. Generally, a daily dosage is administered once in a day or may be administered in several divisions in a day. Alternatively, a daily

dosage can be administered intravenously by prompt administration or continuous infusion over 24 hours.

[Example]

5 **[0081]** Although the present invention is further described in detail in the following Reference Examples, Examples, Test Examples and Formulation Examples, present invention is not limited thereto.

[0082] The measurement conditions for high-performance liquid chromatography mass spectrometer; LCMS are as follows.

Analytical instrument: ACUITY UPLC MS/PDA System (Waters) Mass spectrometer: Waters 3100 MS detector

10 Photodiode array detector: ACUITY PDA detector (210-400 nm)

Column: Acuity BEH C₁₈, 1.7 μm, 2.1x50mm

Flow rate: 0.5 mL/min

Column temperature: 40°C

Solvent:

15

A liquid: 0.1% formic acid/H₂O (v/v)

B liquid: 0.1% formic acid / acetonitrile (v/v)

Gradient condition:

20

Method A:

0.0-2.5min; %A liquid / %B liquid = 90/10 → 10/90

2.5-3.0min; %A liquid / %B liquid = 10/90

25

3.0-3.5min; %A liquid / %B liquid = 0/100

Method B:

0.0-2.5min; %A liquid / %B liquid = 50/50 → 10/90

30

2.5-3.0min; %A liquid / %B liquid = 10/90

3.0-3.5min; %A liquid / %B liquid = 0/100

[0083] The values [MS(m/z)] (MS: mass spectrometry) observed in the mass spectrometry are expressed in m/z, and the retention times are expressed in Rt (minute).

35

Reference Example 1

2-Chloro-5-(ethoxymethyl)benzoic acid

40 **[0084]** The mixture of methyl 5-(bromomethyl)-2-chlorobenzoate (prepared as described in WO2010/132999) (0.2g), potassium carbonate (0.21g), EtOH (3 mL) and THF (3 mL) was stirred at 80°C for 4 hours. The reaction mixture was cooled to room temperature, water was then added, and the mixture was extracted with ethyl acetate. The organic layer was separated and washed with brine, and dried over magnesium sulfate and concentrated in vacuo. The residue was purified on column chromatography to obtain ethyl 2-chloro-5-(ethoxymethyl)benzoate (0.15g). This was dissolved in THF-MeOH-H₂O (3:3:2, 8 mL), and lithium hydroxide hydrate (0.15g) was added, and the mixture was stirred at room temperature for 14 hours. The reaction mixture was acidified with 1N hydrochloric acid and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, and concentrated in vacuo to obtain the titled compound (0.12g) as pale yellow oil.

45

50 Reference Example 2

2-Chloro-5-[(2-ethoxyethoxy)methyl]benzoic acid

[0085] The titled compound was obtained according to the procedure as described in Reference Example 1, using 2-ethoxyethanol instead of EtOH.

55

Reference Example 3

2-Chloro-5-(cyclopropylethynyl)benzoic acid

- 5 **[0086]** To a solution of ethyl 5-bromo-2-chlorobenzoate (1.9g) in DMF (12 mL), toluene (0.3 mL), cyclopropylacetylene (714 mg), copper iodide (275 mg), dichloro bis(triphenylphosphine)palladium (1.01g) and triethylamine (20.1 mL) were added, and the mixture was degassed, stirred at 100°C under argon atmosphere for 8 hours later, and then the reaction mixture was cooled to room temperature and diluted with ethyl acetate. The insoluble material was filtered off on celite. The mother liquor was washed with brine, dried over magnesium sulfate and concentrated. By purification on column chromatography, ethyl 2-chloro-5-(cyclopropylethynyl)benzoate (1.86g) was obtained as brown oil.
- 10 **[0087]** This was dissolved in MeOH-THF (1:1, 60 mL), 10% aqueous sodium hydroxide solution (20 mL) was added, and the solution was stirred at room temperature for 2 hours. The solvent was removed under reduced pressure, water was added to the residue, and the mixture was separated with diethyl ether.
- 15 **[0088]** The aqueous layer was separated, acidified with 1N hydrochloric acid and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate and concentrated in vacuo. The residual solid was washed with n-hexane, and the titled compound (1.02g) was obtained as beige powder.

Reference Example 4

20 2-Chloro-5-(2-cyclopropylethyl)benzoic acid

- [0089]** To a solution of 2-chloro-5-(cyclopropylethynyl)benzoic acid (200 mg) in ethyl acetate (20 mL) was added 10% palladium on carbon (40 mg), and the mixture was stirred under hydrogen atmosphere (0.2MPa), for 14 hours later, then was filtered off on celite, and the mother liquor was concentrated in vacuo. The residue was purified on silica gel column chromatography to obtain the titled compound (190 mg) as yellow oil.
- 25

Reference Example 5

30 2-Chloro-5-(2-phenylethyl)benzoic acid

[Step 1]

2-chloro-5-(phenylethynyl)benzoic acid

- 35 **[0090]** According to the procedure as described in Reference Example 3 using phenylacetylene instead of cyclopropylacetylene, 2-chloro-5-(phenylethynyl)benzoic acid was obtained as yellow powder.

[Step 2]

40 2-Chloro-5-(2-phenylethyl)benzoic acid

- [0091]** The titled compound was obtained as pale yellow powder, according to the procedure as described in Reference Example 4, using 2-chloro-5-(phenylethynyl)benzoic acid instead of 2-chloro-5-(cyclopropylethynyl)benzoic acid.

45 Reference Example 6

5-(3-tert-Butoxy-1-propynyl)-2-chlorobenzoic acid

- 50 **[0092]** The titled compound was obtained as white powder according to the procedure as described in Reference Example 3, using tert-butylpropargyl ether was used instead of cyclopropylacetylene.

Reference Example 7

55 5-(3-tert-Butoxypropyl)-2-chlorobenzoic acid

- [0093]** The titled compound was obtained as a pale yellow oil according to the procedure as described in Reference Example 4, using 5-(3-tert-butoxy-1-propynyl)-2-chlorobenzoic acid instead of 2-chloro-5-(cyclopropylethynyl)benzoic acid.

Reference Example 8

2-Chloro-5-(3-hydroxy-3-methylbutyl)benzoic acid

[Step 1]

2-Chloro-5-(3-hydroxy-3-methyl-1-butynyl)benzoic acid

[0094] The titled compound was obtained as white powder according to the procedure as described in Reference Example 3, using 2-methyl-3-butyn-2-ol instead of cyclopropylacetylene.

[Step 2]

2-Chloro-5-(3-hydroxy-3-methylbutyl)benzoic acid

[0095] The titled compound was obtained as a pale yellow oil according to the procedure as described in Reference Example 4, using 2-chloro-5-(3-hydroxy-3-methyl-1-butynyl)benzoic acid instead of 2-chloro-5-(cyclopropylethynyl)benzoic acid.

Reference Example 9

3-Methoxy-N,2,2-trimethylpropan-1-amine

[Step 1]

tert-Butyl (3-hydroxy-2,2-dimethylpropyl) carbamate

[0096] A solution of 3-amino-2,2-dimethyl-1-propanol (6.98g) and sodium carbonate (7.18g) in 1,4-dioxane-H₂O (1:1, 240 mL) was stirred under ice-cooling, and di-tert-butyl dicarbonate (14.77g) was added, and the mixture was stirred for 5 hours. Ethyl acetate was added, and the organic layer was separated, washed sequentially with water and brine, dried over magnesium sulfate and concentrated in vacuo to obtain the titled compound (12.7g) as a colorless solid.

[Step 2]

3-Methoxy-N,2,2-trimethylpropan-1-amine

[0097] A solution of 60% sodium hydride (1.89g) in DMF (60 mL) was stirred under ice-cooling, a solution of tert-butyl (3-hydroxy-2,2-dimethylpropyl) carbamate (3.84g) in DMF (45 mL) was dropped slowly over 5 minutes. Methyl iodide (10.73g) was added and the mixture was stirred at room temperature for 3 hours. Diethyl ether was added to the reaction mixture. The organic layer was separated, washed sequentially with water and brine, dried over magnesium sulfate and concentrated in vacuo. The residue was purified on column chromatography to obtain tert-butyl (3-methoxy-2,2-dimethylpropyl)methyl carbamate (3.1g) as a colorless oil. This was stirred under ice-cooling, TFA (8 mL) was added, and the mixture was stirred at room temperature for 1 hour later. The mixture was stirred under ice-cooling, and raised the pH to about pH 9 with 1N aqueous sodium hydroxide solution, and extracted with diethyl ether. The diethyl ether layer was dried and concentrated in vacuo with magnesium sulfate to obtain the titled compound (2.6g) as pale yellow oil.

Reference Example 10

5-Butoxy-2-chlorobenzoic acid

[0098] To a solution of 2-chloro-5-fluorobenzoic acid (50 mg) and 1-butanol (263 μ L) in THF (0.5 mg) and DMF (3 mL) was added potassium t-butoxide (329 mg) and the solution was reacted at 120°C for 10 minutes in a microwave synthesizer (Biotage, Initiator). The reaction mixture was cooled and acidified by addition of water and 2M hydrochloric acid, extracted with diethyl ether, dried over anhydrous sodium sulfate. The solvent was removed in vacuo, and the residue was purified on column chromatography to obtain the titled compound (17 mg) as white powder.

Reference Example 11

2-Chloro-5-(2,2-difluoroethoxy)benzoic acid

5 [Step 1]

Methyl 2-chloro-5-(2,2-difluoroethoxy)benzoate

10 **[0099]** To a suspension of methyl 2-chloro-5-hydroxybenzoate (50 mg) and potassium carbonate (111 mg) in acetone (1 mL) was added 2-iodo-1,1-difluoroethane (35 μ L), and the suspension was reacted at 120°C for 10 minutes in a microwave synthesizer (Biotage, Initiator). After reaction at 130°C for additional 15 minutes, the reaction mixture was directly purified on silica gel column chromatography to obtain the titled compound (55 mg) as yellow oil.

15 [Step 2]

2-Chloro-5-(2,2-difluoroethoxy)benzoic acid

20 **[0100]** To a solution of methyl 2-chloro-5-(2,2-difluoroethoxy) benzoate (55 mg) in MeOH (1 mL) was added 1M aqueous lithium hydroxide (0.88 mL), and the solution was stirred at room temperature for 12 hours. The reaction mixture was stirred under ice-cooling, 2M hydrochloric acid was added slowly to acidify to pH 2-3, and the precipitation solid was filtered. The solid was washed sequentially with a small amount of water and diethyl ether, dried under reduced pressure to obtain the titled compound (28 mg) as white powder.

25 Reference Example 12

2-Chloro-5-(4,4,4-trifluorobutoxy)benzoic acid

[Step 1]

30 Methyl 2-chloro-5-(4,4,4-trifluorobutoxy)benzoate

[0101] To a suspension of methyl 2-chloro-5-hydroxy benzoate (50 mg) and potassium carbonate (111 mg) in acetone (1 mL) was added 1-bromo-4,4,4-trifluorobutane (50 μ L) and the suspension was reacted at 120°C for 10 minutes in a microwave synthesizer (Biotage, Initiator). The reaction mixture was directly purified on silica gel column chromatography to obtain the titled compound (74 mg) as colorless oil.

[Step 2]

40 2-Chloro-5-(4,4,4-trifluorobutoxy)benzoic acid

[0102] To a solution of methyl 2-chloro-5-(4,4,4-trifluorobutoxy)benzoate (74 mg) in MeOH (1 mL) was added 1M aqueous lithium hydroxide solution (0.99 mL) and the reaction mixture was stirred at room temperature for 12 hours. The reaction mixture was stirred under ice-cooling, and 2M hydrochloric acid was added slowly to acidify to pH 2-3. The precipitated solid was filtered, washed with water, and dried under reduced pressure to obtain the titled compound (49 mg) as a colorless powder.

Example 1

50 N-[2-(Trifluoromethyl)benzyl]-6-([2-(trifluoromethyl)phenyl]carbonylamino)-1H-benzimidazol-4-carboxamide

[Step 1]

Methyl 1H-benzimidazole-4-carboxylate

55 **[0103]** To a suspension of methyl 2-amino-3-nitrobenzoate (1.0 g) in formic acid (> 87%) was added 5% palladium on carbon (100 mg) and the suspension was heated with stirring at 100°C for 23 hours. After the reaction mixture was cooled to room temperature, the catalyst was filtered off on celite, and the mother liquor was concentrated in vacuo. The residual solid was washed with diethyl ether under stirring, the solid was collected by filtration to obtain the titled compound

(872 mg) as a colorless powder.

[Step 2]

5 Methyl 6-nitro-1H-benzimidazole-4-carboxylate

[0104] Methyl 1H-benzimidazole-4-carboxylate (315 mg) was dissolved in conc. sulfuric acid (3 mL), and the solution was stirred on ice bath. A small amount of potassium nitrate (199 mg) was added portion-wise, the solution was stirred at room temperature for 4 hours. The reaction mixture was poured into ice, alkalified with 3N aqueous sodium hydroxide solution under stirring on ice bath, and extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate. The titled compound (334 mg) was obtained as colorless powder after distilling off a solvent under reduced pressure.

[Step 3]

15 Methyl 6-amino-1H-benzimidazole-4-carboxylate

[0105] Methyl 6-nitro-1H-benzimidazole-4-carboxylate (304 mg) was suspended in MeOH (10 mL), the suspension was added with 5% palladium on carbon (30 mg), and stirred under hydrogen (1 atm) atmosphere. The catalyst was filtered off on celite and washed with MeOH, and the mother liquor was concentrated in vacuo to obtain the titled compound (269 mg) as pale yellow powder.
MS(ESI+) m/z 192(M+H)⁺

[Step 4]

25 methyl 6-([2-(Trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate

[0106] To a solution of methyl 6-amino-1H-benzimidazole-4-carboxylate (266 mg) in dehydrated THF (12 mL), N,N-diisopropylethylamine (284 μ L) was added, and the solution was stirred under ice-cooling. To this solution, 2-(trifluoromethyl) benzoyl chloride (284 μ L) in dehydrated THF (5 mL) was added slowly dropwise, and stirred at the same temperature for 3 hours. After ice water was added, THF was removed under reduced pressure, and aqueous saturated sodium bicarbonate was added, and the mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and the solvent was removed under reduced pressure. The residue was purified on NH silica gel column chromatography (ethyl acetate $\rightarrow$ ethyl acetate/MeOH) to obtain the titled compound (309 mg) as pale yellow powder.

[Step 5]

40 6-([2-(Trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid

[0107] Methyl 6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate (356 mg) was dissolved in MeOH (10 mL), and 1N aqueous sodium hydroxide solution (2 mL) was added, and the reaction mixture was stirred at room temperature overnight. After MeOH was removed under reduced pressure, water was added to the residue, and the solution was acidified with 1N hydrochloric acid to pH 5 under ice-cooling. The precipitate was collected by filtration to obtain the titled compound (276 mg) as slightly brown powder.

[Step 6]

50 N-[2-(Trifluoromethyl)benzyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0108] To a solution of 6-([2-(Trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid (50 mg) in DMF (2 mL) were added N'-(ethylcarbonimidoyl)-N,N-dimethyl-1,3-propanediamine hydrochloride (33 mg), 1-hydroxy-benzotriazole (23 mg) and triethylamine (24 μ L), and 2-(trifluoromethyl)benzylamine (24 μ L) was added under ice-cooling, and the solution was stirred at room temperature overnight. The mixture was poured into ice water and extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was washed with n-hexane/ethyl acetate (1:1), and collected by filtration to obtain the titled compound (61 mg) as colorless powder. MS(ESI+)m/z 507 (M+H)⁺

Example 2

N-Cyclohexyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

- 5 **[0109]** The titled compound was obtained as colorless powder according to the procedure as described in Step 6 of Example 1 using cyclohexylamine instead of 2-(trifluoromethyl)benzylamine. MS(ESI⁺)m/z 431 (M+H)⁺

Example 3

- 10 N-(3-Chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

- [0110]** To a solution of 6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid (example 1, Step 5) (40 mg) in DMF (2 mL), HBTU (53 mg), triethylamine (20 μ L) were added. Under ice-cooling, 3-chloro-2-methylaniline (17 μ L) was added and the solution was stirred at room temperature for 24 hours. The reaction mixture was poured into ice water, extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate and removed the solvent under reduced pressure. The residue was washed with n-hexane/ethyl acetate (1:1) and collected by filtration to obtain the titled compound (33 mg) as colorless powder. MS(ESI⁺)m/z 473 (M+H)⁺

20 Example 5

N-[(1-Hydroxycyclohexyl)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

- 25 **[0111]** The titled compound was obtained as colorless powder according to the procedure as described in Step 6 of Example 1, using 1-aminomethyl-1-cyclohexanol instead of 2-(trifluoromethyl)benzylamine. MS(ESI⁺)m/z 461 (M+H)⁺

Example 6

- 30 N-[2-(Trifluoromethyl)benzyl]-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide

[Step 1]

Ethyl 5-bromo-2,3-dihydro-1-benzofuran-7-carboxylate

- 35 **[0112]** To a solution of 5-bromo-2,3-dihydro-1-benzofuran-7-carboxylic acid (1.0g) in EtOH (15 mL), sulfuric acid (0.5 mL) was added, and the solution was heated at reflux for 30 hours. The reaction mixture was cooled to room temperature, and EtOH was removed. The residue was added with water, and the solution was extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with saturated aqueous sodium bicarbonate and brine, dried over anhydrous magnesium sulfate. The solvent was removed under reduced pressure to obtain the titled compound (1.07g) as pale yellow powder.

[Step 2]

- 45 Ethyl

5-({[2-(Trifluoromethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxylate

- 50 **[0113]** 1,4-dioxane (20 ml) was added to ethyl 5-bromo-2,3-dihydro-1-benzofuran-7-carboxylate (600 mg), 2-(trifluoromethyl)benzamide (501 mg), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (Xantphos) (96 mg), cesium carbonate (1.01g) and tris(dibenzylideneacetone)dipalladium (114 mg). After degassing, the mixture was stirred at 100°C under argon atmosphere for 24 hours. The reaction mixture was filtered off on celite, the solvent was removed under reduced pressure. The resultant residue was purified on silica gel column chromatography to obtain the titled compound (220 mg) as slightly yellow powder.

55

[Step 3]

5-([2-(Trifluoromethyl)phenyl]carbonyl)amino)-2,3-dihydro-1-benzofuran-7-carboxylic acid

- 5 **[0114]** To a suspension of ethyl 5-([2-(trifluoromethyl)phenyl]carbonyl)amino)-2,3-dihydro-1-benzofuran-7-carboxylate (220 mg) obtained in Step 2 in EtOH (5 mL), 1N aqueous sodium hydroxide solution (1 mL) was added and the mixture was stirred at 80°C for 4 hours. The reaction mixture was cooled to room temperature, and EtOH was removed under reduced pressure. The residue was added with water, and acidified to pH 3 with 1N hydrochloric acid under ice-cooling and extracted with ethyl acetate. The organic layer was washed sequentially with water and brine, dried over
10 anhydrous magnesium sulfate, and removed the solvent under reduced pressure to obtain the titled compound (195 mg) as slightly yellow powder.

[Step 4]

- 15 N-[2-(Trifluoromethyl)benzyl]-5-([2-(trifluoromethyl)phenyl]carbonyl)amino)-2,3-dihydro-1-benzofuran-7-carboxamide

- [0115]** To a solution of 5-([2-(trifluoromethyl)phenyl]carbonyl)amino)-2,3-dihydro-1-benzofuran-7-carboxylic acid (35 mg) in DMF (2 mL), N'-(ethylcarbonimidoyl)-N,N-dimethyl-1,3-propanediamine hydrochloride (23 mg), 1-hydroxybenzotriazol (23 mg) and triethylamine (17 µL) were added. Under ice-cooling, The mixture was added 2-(trifluoromethyl)benzylamine (15 µL), and stirred at room temperature overnight. The reaction mixture was poured into ice water, and extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate and removed the solvent under reduced pressure. The residue was washed with n-hexane/ethyl acetate (1:1) and collected by filtration to obtain the titled compound (43 mg) as colorless powder.
20 MS(ESI+) m/z 509 (M+H)⁺, Rt = 2.31 minutes (method A)

25

Example 7

N-Cyclohexyl-5-([2-(trifluoromethyl)phenyl]carbonyl)amino)-2,3-dihydro-1-benzofuran-7-carboxamide

- 30 **[0116]** The titled compound was obtained as colorless powder according to the procedure as described in Step 4 of Example 6 using cyclohexylamine instead of 2-(trifluoromethyl)benzylamine.
MS(ESI+) m/z 433 (M+H)⁺, Rt= 2.25 minutes (method A)

Example 8

35

N-(3-Chloro-2-methylphenyl)-5-([2-(trifluoromethyl)phenyl] carbonyl)amino)-2,3-dihydro-1-benzofuran-7-carboxamide

- [0117]** The titled compound was obtained as colorless powder according to the procedure as described in Example 3, using 5-([2-(trifluoromethyl)phenyl]carbonyl)amino)-2,3-dihydro-1-benzofuran-7-carboxylic acid instead of 6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid. MS(ESI+) m/z 475 (M+H)⁺, Rt= 2.47 minutes (method A)
40

Example 9

- 45 N-Cyclohexyl-5-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-indazole-7-carboxamide

[Step 1]

Methyl 5-bromo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-indazole-7-carboxylate

50

- [0118]** Under argon atmosphere, a solution of methyl 5-bromo-1H-indazole-7-carboxylate (as prepared according to WO2008/65508) (658 mg) in DMF (20 mL) was stirred under ice-cooling. 60% Sodium hydride (124 mg) was added slowly, and the mixture was stirred for 30 minutes at same temperature. 2-(Trimethylsilyl)ethoxymethyl chloride (544 µL) was added dropwise slowly using a syringe, and the mixture was stirred at room temperature for 4 hours. The mixture
55 was poured into ice water and extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on silica gel column chromatography to obtain the titled compound (696 mg) as slightly yellow oil.

[Step 2]

Methyl

5 5-({[(benzyloxy)carbonyl]amino}-1-{{2-(trimethylsilyl)ethoxy}methyl})-1H-indazole-7-carboxylate

[0119] The titled compound was obtained as slightly yellow oil from methyl 5-bromo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-indazole-7-carboxylate prepared in Step 1, according to the procedure as described in Step 2 of Example 6, using benzyl carbamate instead of 2-(trifluoromethyl)benzamide.

10

[Step 3]

Methyl

15 5-amino-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-indazole-7-carboxylate

[0120] To a solution of methyl 5-({[(benzyloxy)carbonyl]amino}-1-{{2-(trimethylsilyl)ethoxy}methyl})-1H-indazole-7-carboxylate (364 mg) in MeOH (10 mL) was added 5% palladium on carbon (55 mg), and the reaction mixture was stirred overnight under hydrogen atmosphere at an ordinary pressure. The reaction mixture was filtered off on celite and washed with MeOH. The mother liquor was concentrated under reduced pressure to obtain the titled compound (234 mg) as green powder.

20

[Step 4]

25 Methyl

5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-indazole-7-carboxylate

[0121] The titled compound (323 mg) was obtained as colorless powder according to the procedure as described in Step 4 of Example 1, using methyl 5-amino-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-indazole-7-carboxylate (231 mg).

30

[Step 5]

5-({[2-(Trifluoromethyl)phenyl]carbonyl}amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-indazole-7-carboxylic acid

35

[0122] The titled compound (291 mg) was obtained as colorless powder according to the procedure as described in Step 5 of Example 1, using methyl 5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-indazole-7-carboxylate (321 mg).

40

[Step 6]

N-Cyclohexyl-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-indazole-7-carboxamide

[0123] To a solution of 5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-indazole-7-carboxylic acid (obtained in Step 5) (48 mg) in DMF (2 mL) were added HBTU (46 mg) and triethylamine (17 μ L), and the mixture was stirred at room temperature, and added with cyclohexylamine (12.5 μ L) and stirred at same temperature overnight. The reaction mixture was poured into ice water, extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with saturated aqueous sodium bicarbonate and brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue is triturated in n-hexane/ethyl acetate (10:1), and collected by filtration to obtain 59 mg of N-cyclohexyl-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1-{{2-(trimethylsilyl)ethoxymethyl}-indazole-7-carboxamide as colorless powder. This was dissolved in MeOH (2 mL), 6N hydrochloric acid (2 mL) was added, and the solution was stirred on oil bath at 60-80°C for 10 hours. The solvent was removed, and ice water was added to the residue. The precipitate was collected by filtration and dried to obtain the titled compound (33 mg) as colorless powder.

45

50

55 MS(ESI+) m/z 431 (M+H)⁺, Rt= 1.91 minutes (method A)

Elemental Analysis for C₂₂H₂₁F₃N₄O₂+0.1H₂O

Calcd. (%) C:61.13 H:4.94 N:12.96

Fond. (%) C:60.93 H:4.91 N:12.71

Example 10

N-[2-(Trifluoromethyl)benzyl]-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-indazole-7-carboxamide

[0124] The titled compound was obtained as colorless powder according to the procedure as described in Step 6 of Example 9, using 2-(trifluoromethyl)benzylamine instead of cyclohexylamine.
MS(ESI+) m/z 507 (M+H)⁺, Rt= 2.05 minutes (method A)

Example 11

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 2-amino-3-[(methoxyacetyl)amino]benzoate

[0125] To a solution of methyl 2, 3-diaminobenzoate (3.0g) in THF (50 mL) was added N,N-diisopropylethylamine (4.0 mL). The solution was stirred under ice-cooling, and added with methoxyacetyl chloride (1.81 mL) in THF (10 mL) dropwise slowly and stirred for 3 hours at same temperature. The reaction mixture was added with saturated aqueous sodium bicarbonate, and removed THF under reduced pressure. The residue was added with saturated aqueous sodium bicarbonate, and extracted with ethyl acetate, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on silica gel column chromatography to obtain the titled compound (3.60g) as pale yellow powder.

[Step 2]

Methyl 2-(methoxymethyl)-1H-benzimidazole-4-carboxylate

[0126] Methyl 2-amino-3-[(methoxyacetyl)amino]benzoate (280 mg) was dissolved in acetic acid (6 mL), and the solution was heated at 100°C for 0.5 hour with stirring. After cooling the reaction mixture to room temperature, acetic acid was removed. The residue was added with saturated aqueous sodium bicarbonate under ice-cooling, and extracted with ethyl acetate. The organic layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate and removed the solvent under reduced pressure to obtain the titled compound (241 mg) as pale yellow powder.

[Step 3]

Methyl 2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxylate

[0127] The titled compound (274 mg) was obtained as colorless powder according to the procedure as described in Step 2 of Example 1, using methyl 2-(methoxymethyl)-1H-benzimidazole-4-carboxylate (240 mg) instead of methyl 1H-benzimidazole-4-carboxylate.

[Step 4]

Methyl 6-amino-2-(methoxymethyl)-1H-benzimidazole-4-carboxylate

[0128] The titled compound (202 mg) was obtained as yellow powder according to the procedure as described in Step 3 of Example 1, using methyl 2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxylate (271 mg) instead of methyl 6-nitro-1H-benzimidazole-4-carboxylate.

[Step 5]

Methyl

2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate

[0129] The titled compound (281 mg) was obtained as colorless powder according to the procedure as described in

EP 2 746 265 B1

Step 4 of Example 1, using methyl 6-amino-2-(methoxymethyl)-1H-benzimidazole-4-carboxylate (200 mg) instead of methyl 6-amino-1H-benzimidazole-4-carboxylate.

[Step 6]

2-(Methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid

[0130] The titled compound (263 mg) was obtained as colorless powder according to the procedure as described in Step 5 of Example 1, using methyl 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate (278 mg) instead of methyl 6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate.

[Step 7]

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0131] To a solution of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid (153 mg) in DMF (5 mL), were added HBTU (178 mg) and triethylamine (65 μ L). The mixture was added with 3-chloro-2-methylaniline (56 μ L) under stirring at room temperature, and the mixture was stirred overnight. The reaction mixture was poured into saturated aqueous sodium bicarbonate, extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on silica gel column chromatography to obtain N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (165 mg) as colorless powder.

[Step 8]

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0132] N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (165 mg) obtained in Step 7 was suspended in EtOH (3 mL). The suspension was added with 1M hydrochloric acid (0.32 mL) and stirred to obtain a homogeneous solution, and removed the solvent under reduced pressure. The residue was triturated in ethyl acetate, collected by filtration, and dried to obtain the titled compound (167 mg) as colorless powder.

MS(ESI+) m/z 517 (M+H)⁺, Rt= 2.41 minutes (method A)

Elemental Analysis for C₂₅H₂₀ClF₃N₄O₃•HCl+1.0H₂O

Calcd. (%) C:52.55 H:4.06 N:9.81

Found. (%) C:52.35 H:3.96 N:9.89

Example 12

2-Methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0133] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 11, using acetyl chloride instead of methoxyacetyl chloride, and 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 521 (M+H)⁺,

Rt= 1.74 minutes (method A)

Elemental Analysis for C₂₅H₁₈F₆N₄O₂•HCl+1.5H₂O

Calcd. (%) C:51.42 H:3.80 N:9.60

Fond. (%) C:51.68 H:3.81 N:9.45

Example 13

N-Cyclohexyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

5 **[0134]** The titled compound was obtained as pale yellow powder according to the procedure as described in Example 11, using acetyl chloride instead of methoxyacetyl chloride, cyclohexylamine instead of 3-chloro-2-methylaniline. MS(ESI+) m/z 445 (M+H)⁺, Rt= 1.52 minutes (method A)

Example 14

10

N-(3-Chloro-2-methylphenyl)-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

15 **[0135]** The titled compound was obtained as pale yellow powder according to the procedure as described in Example 11, using acetyl chloride instead of methoxyacetyl chloride. MS(ESI+) m/z 487 (M+H)⁺, Rt= 2.10 minutes (method A)

Example 15

20

N-Cyclopentyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0136] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using acetyl chloride instead of methoxyacetyl chloride, and cyclopentylamine instead of 3-chloro-2-methylaniline.
25 MS(ESI+) m/z 431 (M+H)⁺, Rt= 1.39 minutes (method A)

Example 16

30

N-Cyclobutyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0137] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using acetyl chloride instead of methoxyacetyl chloride, and cyclobutylamine instead of 3-chloro-2-methylaniline.
35 MS(ESI+) m/z 417 (M+H)⁺, Rt= 1.29 minutes (method A)

Example 17

40

N-(3-Chloro-2-methylphenyl)-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0138] The titled compound was obtained as white powder according to the procedure as described in Step 1 to Step 7 of Example 11, using propionyl chloride instead of methoxyacetyl chloride.
MS(ESI+) m/z 501 (M+H)⁺, Rt= 2.39 minutes (method A)

Example 18

45

N-Cyclohexyl-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0139] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using propionyl chloride instead of methoxyacetyl chloride, and cyclohexylamine instead of 3-chloro-2-methylaniline.
50 MS(ESI+) m/z 459 (M+H)⁺, Rt= 1.75 minutes (method A)

Example 19

55

2-Ethyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0140] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using propionyl chloride instead of methoxyacetyl chloride, and 2-(trifluoromethyl)benzylamine

instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 535 (M+H)⁺, Rt= 1.96 minutes (method A)

Example 20

N-Cyclohexyl-2-(methoxymethyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0141] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using cyclohexylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 475 (M+H)⁺, Rt= 1.97 minutes (method A)

Example 21

2-(Methoxymethyl)-N-[2-(trifluoromethyl)benzyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0142] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 551 (M+H)⁺, Rt= 2.12 minutes (method A)

Example 22

2-(Methoxymethyl)-N-(2-methylphenyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0143] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using ortho-toluidine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 483 (M+H)⁺, Rt= 1.06 minutes (method B)

Example 23

2-(Methoxymethyl)-N-(4-methylphenyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0144] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using para-toluidine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 483 (M+H)⁺, Rt= 1.06 minutes (method B)

Example 24

N-(2-Chlorobenzyl)-2-(methoxymethyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0145] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 11, using 2-chlorobenzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 517 (M+H)⁺, Rt= 2.04 minutes (method A)

Example 25

2-(Methoxymethyl)-N-(4-methylbenzyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0146] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 4-methylbenzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 497 (M+H)⁺, Rt= 1.99 minutes (method A)

EP 2 746 265 B1

Example 26

N-(4,4-Difluorocyclohexyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0147] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 4,4-difluorocyclohexylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 511 (M+H)⁺, Rt= 1.88 minutes (method A)

Example 27

N-(4-tert-Butylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0148] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 4-tert-butylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 525 (M+H)⁺, Rt= 1.63 minutes (method B)

Example 28

2-(Methoxymethyl)-N-[4-(trifluoromethyl)phenyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0149] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 1-amino-4-(trifluoromethyl)benzene instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 537 (M+H)⁺, Rt= 1.39 minutes (method B)

Example 29

N-(2,4-Dimethylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0150] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2,4-dimethylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 497 (M+H)⁺, Rt= 1.24 minutes (method B)

Example 30

N-(2-Chloro-4-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0151] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-chloro-4-methylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 517 (M+H)⁺, Rt= 1.42 minutes (method B)

Example 31

N-(3,4-Dimethylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0152] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 3,4-dimethylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 497 (M+H)⁺, Rt= 1.22 minutes (method B)

EP 2 746 265 B1

Example 32

N-(3-Chloro-4-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0153] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 3-chloro-4-methylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 517 (M+H)⁺, Rt= 1.46 minutes (method B)

Example 33

N-(2,3-Dihydro-1H-inden-5-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0154] The titled compound was obtained as gray powder according to the procedure as described in Example 11, using 5-aminoindan instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 509 (M+H)⁺, Rt= 2.36 minutes (method A)

Example 34

2-(Methoxymethyl)-N-(5,6,7,8-tetrahydronaphthalen-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0155] The titled compound was obtained as white powder according to the procedure as described in Example 11, using 5,6,7,8-tetrahydro-1-naphthylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 523 (M+H)⁺, Rt= 2.44 minutes (method A)

Example 35

N-(2-Fluorophenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0156] A stirring solution of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid in THF (2 mL) was stirred under ice-cooling, added with catalytic amount of DMF, and then oxalyl chloride (7.7 μ L) was added. One hour later, the reaction mixture was added with additional oxalyl chloride (7.7 μ L) and stirred for additional one hour, and removed volatile elements under reduced pressure. The residue was dissolved in THF (2 mL), and 2-fluoroaniline (8.8 μ L) and triethylamine (11 μ L) were added sequentially, and the mixture was stirred at room temperature for 1 hour and added with saturated aqueous sodium bicarbonate, and extracted with ethyl acetate.

[0157] The ethyl acetate layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated. The residue was triturated in n-hexane/diethyl ether to obtain the titled compound (21 mg) as colorless powder.

MS(ESI+) m/z 487 (M+H)⁺, Rt= 2.14 minutes (method A)

Example 36

2-(Methoxymethyl)-N-(2-methoxyphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0158] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using ortho-anisidine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 499 (M+H)⁺, Rt= 2.07 minutes (method A)

Example 37

2-(Methoxymethyl)-N-(4-methoxyphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0159] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using para-anisidine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 499 (M+H)⁺, Rt= 1.96 minutes (method A)

Example 38

5 N-(3-Bromo-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0160] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using 3-bromo-2-methylaniline instead of 3-chloro-2-methylaniline.

10 MS(ESI+) m/z 563 (M+H)⁺, Rt= 2.45 minutes (method A)

Example 39

15 N-(3-Chloro-2-methylbenzyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0161] The titled compound was obtained as colorless powder according to the procedure as described in Step 1 to Step 7 of Example 11, using 3-chloro-2-methylbenzylamine instead of 3-chloro-2-methylaniline.

20 MS(ESI+) m/z 531 (M+H)⁺, Rt= 2.19 minutes (method A)

Example 40

25 N-(2,6-Difluorophenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0162] The titled compound was obtained as colorless powder according to the procedure as described in Example 35, using 2,6-difluoroaniline instead of 2-fluoroaniline.

MS(ESI+) m/z 505 (M+H)⁺, Rt= 1.94 minutes (method A)

30 Example 41

N-(3-Cyano-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

35 **[0163]** To a solution of N-(3-bromo-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (Example 38) (85 mg) in DMF (0.5 mL), was added zinc cyanide (60% content, 60 mg). After degassing, the mixture was added with tetrakis(triphenylphosphine)palladium (88 mg), and stirred at 100°C for 24 hours. The reaction mixture was added with ice water, and extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on preparative thin-layer chromatography to obtain the titled compound (14 mg) as colorless powder.

40 MS(ESI+) m/z 508 (M+H)⁺, Rt= 1.02 minutes (method B)

Example 42

45 2-(Methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-N-({[3-(trifluoromethyl)pyridin-2-yl]methyl})-1H-benzimidazole-4-carboxamide dihydrochloride

[0164] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-(aminomethyl)-3-(trifluoromethyl)pyridine instead of 3-chloro-2-methylaniline.

50 MS(ESI+) m/z 552 (M+H)⁺, Rt= 1.86 minutes (method A)

Example 43

55 N-(2-Chloro-6-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0165] The titled compound was obtained as colorless powder according to the procedure as described in Example

EP 2 746 265 B1

11, using 2-chloro-6-methylaniline instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 517 (M+H)⁺, Rt= 2.35 minutes (method A)

Example 44

2-(2-Amino-2-oxoethyl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 2-(2-amino-2-oxoethyl)-6-nitro-1H-benzimidazole-4-carboxylate

[0166] Under ice-cooling, potassium nitrate (117 mg) was added slowly portion wise to a stirring solution of methyl 2-(cyanomethyl)-1H-benzimidazole-4-carboxylate (prepared as described in EP1479681) (226 mg) in conc. sulfuric acid (5 mL), and the mixture was stirred at room temperature for 5 hours. The reaction mixture was poured into ice, and alkalinized with 3N aqueous sodium hydroxide solution, and extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate and removed the solvent under reduced pressure. The residue was washed with MeOH, filtered and dried to obtain the titled compound (243 mg).

[Step 2]

Methyl

6-amino-2-(2-amino-2-oxoethyl)-1H-benzimidazole-4-carboxylate

[0167] The titled compound was obtained as yellow powder according to the procedure as described in Step 4 of Example 11, using methyl 2-(2-amino-2-oxoethyl)-6-nitro-1H-benzimidazole-4-carboxylate instead of methyl 2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxylate.

[Step 3]

Methyl

2-(2-amino-2-oxoethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate

[0168] The titled compound was obtained as orange powder according to the procedure as described in Step 4 of Example 1, using methyl 6-amino-2-(2-amino-2-oxoethyl)-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate.

[Step 4]

2-(2-Amino-2-oxoethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid

[0169] The titled compound was obtained as brown powder according to the procedure as described in Step 5 of Example 1, using methyl 2-(2-amino-2-oxoethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate instead of methyl 6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate.

[Step 5]

2-(2-Amino-2-oxoethyl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0170] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 7 of Example 11, using 2-(2-amino-2-oxoethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid.

MS(ESI+) m/z 530 (M+H)⁺, Rt= 1.86 minutes (method A)

Example 45

2-(2-Amino-2-oxoethyl)-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0171] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 7 of Example 11, using 2-(2-amino-2-oxoethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid, and 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 564 (M+H)⁺, Rt= 1.72 minutes (method A)

Example 46

N-(3-Chloro-2-methylphenyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[Step 1]

Methyl 1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate

[0172] To a solution of methyl 6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate (Example 1, Step 4) (189 mg) in THF (5 mL), NaH (60%, 25 mg) was added slowly under ice-cooling, and the mixture was stirred at same temperature for 30 minutes and additionally at room temperature for 1 hour. The reaction mixture was further stirred under ice-cooling, and was added with methyl iodide (81 μ L) slowly, and stirred at room temperature overnight. The reaction mixture was added with saturated aqueous ammonium chloride, and extracted with ethyl acetate. The organic layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on column chromatography to obtain the titled compound (78 mg) as colorless powder.

[Step 2]

1-Methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid

[0173] The titled compound was obtained as colorless powder according to the procedure as described in Step 5 of Example 1, using methyl 1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate instead of methyl 6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate.

[Step 3]

N-(3-Chloro-2-methylphenyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0174] The titled compound was obtained as colorless powder according to the procedure as described in Step 7 of Example 11, using 1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid.

MS(ESI+) m/z 487 (M+H)⁺, Rt= 1.44 minutes (method B)

Example 47

N-Cyclohexyl-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0175] The titled compound was obtained as colorless powder according to the procedure as described in Example 46, using cyclohexylamine instead of 3-chloro-2-methylaniline.

MS (ESI+) m/z 445 (M+H)⁺

Example 48

1-Methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0176] The titled compound was obtained as colorless powder according to the procedure as described in Example 46, using 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 521 (M+H)⁺, Rt= 1.06 minutes (method B)

Example 49

N-(3-Chloro-2-methylphenyl)-1-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0177] The titled compound was obtained as white powder according to the procedure as described in Example 46, using ethyl iodide instead of methyl iodide.

MS(ESI+) m/z 501 (M+H)⁺, Rt= 2.53 minutes (method A)

Example 50

N-Cyclohexyl-1-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0178] The titled compound was obtained as colorless powder according to the procedure as described in Example 46, using ethyl iodide instead of methyl iodide, and cyclohexylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 459 (M+H)⁺, Rt= 2.14 minutes (method A)

Example 51

1-Ethyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0179] The titled compound was obtained as colorless powder according to the procedure as described in Example 46, using ethyl iodide instead of methyl iodide, and 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 535 (M+H)⁺, Rt= 2.28 minutes (method A)

Example 52

N-(3-Chloro-2-methylphenyl)-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide

[Step 1]

6-Bromo-2-methyl-1,3-benzoxazole-4-carboxylic acid

[0180] To a solution of 5-bromo-3-hydroxyanthranilic acid (prepared as described in Eur. J. Med. Chem., 1999, 34, 729) (400 mg) in xylene (30 mL), were added acetyl chloride (135 μ L), triethylamine (265 μ L) and pyridinium p-toluenesulfonate (130 mg), and the mixture was heated at reflux for 8 hours. The reaction mixture was cooled to room temperature, diluted with ethyl acetate, washed sequentially with water and brine, and dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on silica gel column chromatography to obtain the titled compound (471 mg).

[Step 2]

6-Bromo-N-(3-chloro-2-methylphenyl)-2-methyl-1,3-benzoxazole-4-carboxamide

[0181] To a solution of 6-bromo-2-methyl-1,3-benzoxazole-4-carboxylic acid (136 mg) in DMF (2 mL), were added HBTU (242 mg) and triethylamine (110 μ L). The mixture was added with 3-chloro-2-methylaniline (76 μ L) and stirred at room temperature over night. The reaction mixture was poured into ice water and extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on silica gel column chromatography to obtain the titled compound (53 mg) as white powder.

[Step 3]

N-(3-Chloro-2-methylphenyl)-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide

5 **[0182]** The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 6, using 6-bromo-N-(3-chloro-2-methylphenyl)-2-methyl-1,3-benzoxazole-4-carboxamide instead of ethyl 5-bromo-2,3-dihydro-1-benzofuran-7-carboxylate.

MS(ESI) m/z 488 (M+H)⁺, Rt= 1.96 minutes (method B)

10 Example 53

2-Methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide

[Step 1]

15

6-Bromo-2-methyl-N-[2-(trifluoromethyl)benzyl]-1,3-benzoxazole-4-carboxamide

[0183] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 52, using 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.

20

[Step 2]

2-Methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide

25 **[0184]** The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 6, using 6-bromo-2-methyl-N-[2-(trifluoromethyl)benzyl]-1,3-benzoxazole-4-carboxamide instead of ethyl 5-bromo-2,3-dihydro-1-benzofuran-7-carboxylate.

MS(ESI) m/z 522 (M+H)⁺, Rt= 2.49 minutes (method A)

30 Example 54

N-(3-Chloro-2-methylphenyl)-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide

[Step 1]

35

6-Bromo-2-ethyl-1,3-benzoxazole-4-carboxylic acid

[0185] The titled compound was obtained according to the procedure as described in Step 1 of Example 52, using propionyl chloride instead of acetyl chloride.

40

[Step 2]

6-Bromo-N-(3-chloro-2-methylphenyl)-2-ethyl-1,3-benzoxazole-4-carboxamide

45 **[0186]** The titled compound was obtained according to the procedure as described in Step 2 of Example 52, using 6-bromo-2-ethyl-1,3-benzoxazole-4-carboxylic acid instead of 6-bromo-2-methyl-1,3-benzoxazole-4-carboxylic acid.

[Step 3]

50 N-(3-Chloro-2-methylphenyl)-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1,3-benzoxazole-4-carboxamide

[0187] The titled compound was obtained as powder according to the procedure as described in Step 2 of Example 6, using 6-bromo-N-(3-chloro-2-methylphenyl)-2-ethyl-1,3-benzoxazole-4-carboxamide instead of ethyl 5-bromo-2,3-dihydro-1-benzofuran-7-carboxylate.

55

MS(ESI) m/z 502 (M+H)⁺, Rt= 2.29 minutes (method B)

Example 55

N-(3-Chloro-2-methylphenyl)-2-ethoxy-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[Step 1]

Methyl 2-ethoxy-6-nitro-1H-benzimidazole-4-carboxylate

[0188] To a stirring solution of methyl 2-ethoxy-1H-benzimidazole-4-carboxylate (prepared as described in J. Med. Chem., 1993, 36, 2182) (500 mg) in conc. sulfuric acid (5 mL), potassium nitrate (275 mg) was added portion-wise under ice-cooling, and the mixture was stirred at room temperature for 2.5 hours. The reaction mixture was poured into ice, alkalified with aqueous sodium hydroxide solution, and extracted with ethyl acetate. The ethyl acetate layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure to obtain the titled compound (595 mg) as white powder.

[Step 2]

Methyl 6-amino-2-ethoxy-1H-benzimidazole-4-carboxylate

[0189] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 3 of Example 1, using methyl 2-ethoxy-6-nitro-1H-benzimidazole-4-carboxylate instead of methyl 6-nitro-1H-benzimidazole-4-carboxylate.

[Step 3]

Methyl

2-ethoxy-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate

[0190] The titled compound was obtained as colorless powder according to the procedure as described in Step 4 of Example 1, using methyl 6-amino-2-ethoxy-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate.

[Step 4]

2-Ethoxy-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid

[0191] The titled compound was obtained as colorless powder according to the procedure as described in Step 5 of Example 1, using methyl 2-ethoxy-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate instead of methyl 6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate.

[Step 5]

N-(3-Chloro-2-methylphenyl)-2-ethoxy-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0192] The titled compound was obtained as white powder according to the procedure as described in Step 7 of Example 11, using 2-ethoxy-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid. MS(ESI+) m/z 517 (M+H)⁺, Rt= 1.72 minutes (method B)

Example 56

2-Ethoxy-N-[2-(trifluoromethyl)benzyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0193] The titled compound was obtained as white powder according to the procedure as described in Example 55,

EP 2 746 265 B1

using 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 551 (M+H)⁺, Rt= 1.32 minutes (method B)

Example 57

N-(3-Chloro-2-methylphenyl)-2-(1-chloro-2-methylpropan-2-yl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0194] The titled compound was obtained as white powder according to the procedure as described in Example 11, using 3-chloro-2,2-dimethylpropionyl chloride instead of methoxyacetyl chloride.
MS(ESI+) m/z 563 (M+H)⁺, Rt= 2.75 minutes (method A)

Example 58

N-(3-Chloro-2-methylphenyl)-2-[(dimethylamino)methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl

2-(chloromethyl)-6-nitro-1H-benzimidazole-4-carboxylate

[0196] The titled compound was obtained as slightly yellow powder according to the procedure as described in Step 2 of Example 1, using methyl 2-(chloromethyl)-1H-benzimidazole-4-carboxylate (prepared as described in WO2003/106430) instead of methyl 1H-benzimidazole-4-carboxylate.

[Step 2]

Methyl

2-[(dimethylamino)methyl]-6-nitro-1H-benzimidazole-4-carboxylate

[0197] To a solution of methyl 2-(chloromethyl)-6-nitro-1H-benzimidazole-4-carboxylate (72 mg) in acetonitrile (3 mL), was added 2M dimethylamine/MeOH solution (1.3 mL), and the mixture was stirred at 80°C overnight. The solvent was removed under reduced pressure, and the residue was purified on silica gel chromatography to obtain the titled compound (68 mg) as slightly yellow solid.

[Step 3]

Methyl

6-amino-2-[(dimethylamino)methyl]-1H-benzimidazole-4-carboxylate

[0198] The titled compound was obtained as yellow amorphous according to the procedure as described in Step 3 of Example 1, using methyl 2-[(dimethylamino)methyl]-6-nitro-1H-benzimidazole-4-carboxylate instead of methyl 6-nitro-1H-benzimidazole-4-carboxylate.

[Step 4]

Methyl

2-[(dimethylamino)methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate

[0199] The titled compound was obtained as colorless powder according to the procedure as described in Step 4 of Example 1, using methyl 6-amino-2-[(dimethylamino)methyl]-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate.

[Step 5]

2-[(Dimethylamino)methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazole-4-carboxylic acid

5 **[0200]** The titled compound was obtained as yellow amorphous according to the procedure as described in Step 5 of Example 1, using methyl 2-[(dimethylamino)methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazole-4-carboxylate instead of methyl 6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazole-4-carboxylate.

[Step 6]

10

N-(3-Chloro-2-methylphenyl)-2-[(dimethylamino)methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazole-4-carboxamide hydrochloride

15 **[0201]** The titled compound was obtained as white powder according to the procedure as described in Step 7 of Example 11, using 2-[(dimethylamino)methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazole-4-carboxylic acid.
MS(ESI+) m/z 530 (M+H)⁺, Rt= 1.54 minutes (method A)

20 Example 59

N-(3-Chloro-2-methylphenyl)-2-(2-methylpropyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazole-4-carboxamide hydrochloride

25 **[0202]** The titled compound was obtained as white powder according to the procedure as described in Example 11, using isovaleryl chloride instead of methoxyacetyl chloride. MS(ESI+) m/z 529 (M+H)⁺, Rt= 1.95 minutes (method B)

Example 60

30 2-(2-Methylpropyl)-N-[2-(trifluoromethyl)benzyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazole-4-carboxamide hydrochloride

35 **[0203]** The titled compound was obtained as white powder according to the procedure as described in Example 11, using isovaleryl chloride instead of methoxyacetyl chloride, and 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 563 (M+H)⁺, Rt= 1.20 minutes (method B)

Example 61

40 tert-Butyl

3-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino-1H-benzimidazol-2-yl}azetidine-1-carboxylate

45 **[0204]** The titled compound was obtained as pale yellow powder according to the procedures as described in Steps 1 to 7 of Example 11, using 1-(tert-butoxycarbonyl)azetidine-3-carbonyl chloride (prepared as described in US6020368) instead of methoxyacetyl chloride.
MS(ESI+) m/z 628 (M+H)⁺, Rt= 1.87 minutes (method B)

50 Example 62

55

EP 2 746 265 B1

N-(3-Chloro-2-methylphenyl)-2-[(methylamino)methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide dihydrochloride

[Step 1]

Methyl

2-([[(tert-butoxycarbonyl)(methyl)amino]methyl]-6-nitro-1H-benzimidazole-4-carboxylate

[0205] To a solution of methyl 2-(chloromethyl)-6-nitro-1H-benzimidazole-4-carboxylate (Example 58, Step 1) (72 mg) in acetonitrile (2 mL), was added 40% methylamine/MeOH solution (0.27 mL), and the mixture was stirred at 80°C for 1 hour. The solvent was removed under reduced pressure. The residue was added with water, and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated. The obtained residue (68 mg) was dissolved in anhydrous THF (2 mL), triethylamine (93 µL) was added to the solution. Di-tert-butyl dicarbonate (141 µL) was added to the solution with stirring under ice-cooling for 3 hours. The reaction mixture was added with water, and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over anhydrous magnesium and concentrated under reduced pressure. The residue was purified on column chromatography to obtain the titled compound (54 mg) as slightly yellow amorphous.

[Step 2]

Methyl

6-amino-2-([[(tert-butoxycarbonyl)(methyl)amino]methyl]-1H-benzimidazole-4-carboxylate

[0206] The titled compound was obtained as yellow amorphous according to the procedure as described in Step 3 of Example 1, using methyl 2-([[(tert-butoxycarbonyl)(methyl)amino]methyl]-6-nitro-1H-benzimidazole-4-carboxylate instead of methyl 6-nitro-1H-benzimidazole-4-carboxylate.

[Step 3]

Methyl

2-([[(tert-butoxycarbonyl)(methyl)amino]methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate

[0207] The titled compound was obtained as slightly yellow amorphous according to the procedure as described in Step 4 of Example 1, using methyl 6-amino-2-([[(tert-butoxycarbonyl)(methyl)amino]methyl]-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate.

[Step 4]

2-([[(tert-Butoxycarbonyl)(methyl)amino]methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid

[0208] The titled compound was obtained as yellow amorphous according to the procedure as described in Step 5 of Example 1, using methyl 2-([[(tert-butoxycarbonyl)(methyl)amino]methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate instead of methyl 6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate.

[Step 5]

N-(3-Chloro-2-methylphenyl)-2-[(methylamino)methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide dihydrochloride

[0209] 2-([[(tert-Butoxycarbonyl)(methyl)amino]methyl]-N-(3-chloro-2-methylphenyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide (9 mg) was obtained according to the procedure as described in Step 7 of Example 11, using 2-([[(tert-butoxycarbonyl)(methyl)amino]methyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid instead of methyl 6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate.

yl}amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid. This was dissolved in methylene chloride (1 mL). The residue was added with TFA (0.5 mL) under ice-cooling, and stirred for 0.5 hour. The reaction mixture was added with ice and then saturated aqueous sodium bicarbonate, and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain N-(3-chloro-2-methylphenyl)-2-[(methylamino)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (5 mg) as slightly yellow powder. This was dissolved in MeOH (0.5 mL), and treated with 1M hydrochloric acid (2 Eq), and concentrated to obtain the titled compound (5 mg) as slightly yellow powder. MS(ESI+) m/z 516 (M+H)⁺, Rt= 1.55 minutes (method A)

Example 63

{4-[(3-Chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}methyl acetate

[Step 1]

Methyl

2-[(acetyloxy)methyl]-6-nitro-1H-benzimidazole-4-carboxylate

[0210] To a solution of methyl 2-(chloromethyl)-6-nitro-1H-benzimidazole-4-carboxylate (Example 58, Step 1) (99 mg) in DMF (1 mL), was added sodium acetate (17 mg), and the mixture was stirred at 50°C for 6 hours. The reaction mixture was added with ice water, and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over anhydrous magnesium sulfate and concentrated. The residue was purified on column chromatography to obtain the titled compound (65 mg) as colorless powder.

[Step 2]

Methyl

2-[(acetyloxy)methyl]-6-amino-1H-benzimidazole-4-carboxylate

[0211] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 3 of Example 1, using methyl 2-[(acetyloxy)methyl]-6-nitro-1H-benzimidazole-4-carboxylate instead of methyl 6-nitro-1H-benzimidazole-4-carboxylate.

[Step 3]

Methyl 2-[(acetyloxy)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate

[0212] The titled compound was obtained as pale yellow amorphous according to the procedure as described in Step 4 of Example 1, using methyl 2-[(acetyloxy)methyl]-6-amino-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate.

[Step 4]

2-(Hydroxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid

[0213] The titled compound was obtained as colorless powder according to the procedure as described in Step 5 of Example 1, using methyl 2-[(acetyloxy)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate instead of methyl 6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate.

[Step 5]

{4-[(3-Chloro-2-methylphenyl)carbamoyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazol-2-yl}methyl acetate

[0214] A mixture of 2-(hydroxymethyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid (50 mg) and N,N-diisopropylethylamine (49 μ L) in methylene chloride-THF (1:1)(2 mL) was stirred under ice-cooling, added with acetyl chloride (19 μ L), and stirred at room temperature overnight. The reaction mixture was poured into a cold 10% aqueous citric acid solution, and extracted with ethyl acetate. The organic layer was separated, and washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was dissolved in DMF (3 mL), and added with HBTU (65 mg) and triethylamine (24 μ L), and stirred at room temperature. The mixture was added with 3-chloro-2-methylaniline (16 μ L) and stirred overnight. The reaction mixture was poured into ice water, and extracted with ethyl acetate. The organic layer was washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure.

The obtained residue was purified on column chromatography to obtain the titled compound (6 mg) as colorless powder. MS(ESI+) m/z 545 (M+H)⁺, Rt= 2.41 minutes (method A)

Example 64

N-(3-Chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0215] The titled compound was obtained as white powder according to the procedure as described in Example 11, using (2R)-tetrahydro-2-furancarboxyl chloride (prepared as described in WO2006/79642) instead of methoxyacetyl chloride.

MS(ESI+) m/z 543 (M+H)⁺, Rt= 2.56 minutes (method A)

Elemental Analysis for C₂₇H₂₂ClF₃N₄O₃•HCl+1.5H₂O

Calcd. (%) C:53.48 H:4.32 N:9.24

Fond. (%) C:53.26 H:4.17 N:9.19

Example 65

2-[(2R)-Tetrahydrofuran-2-yl]-N-[2-(trifluoromethyl)benzyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0216] The titled compound was obtained as white powder according to the procedure as described in Example 11, using (2R)-tetrahydro-2-furancarboxyl chloride (prepared as described in WO2006/79642) instead of methoxyacetyl chloride, and 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 577 (M+H)⁺, Rt= 2.25 minutes (method A)

Example 66

N-(3-Chloro-2-methylphenyl)-2-[(2S)-tetrahydrofuran-2-yl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0217] The titled compound was obtained as white powder according to the procedure as described in Example 11, using (2S)-tetrahydro-2-furancarboxyl chloride (prepared as described in WO2006/79642) instead of methoxyacetyl chloride. MS(ESI+) m/z 543 (M+H)⁺, Rt= 2.56 minutes (method A)

Example 67

2-[(2S)-Tetrahydrofuran-2-yl]-N-[2-(trifluoromethyl)benzyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0218] The titled compound was obtained as white powder according to the procedure as described in Example 11, using (2S)-tetrahydro-2-furancarboxyl chloride (prepared as described in WO2006/79642) instead of methoxyacetyl chloride, and 2-(trifluoromethyl)benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 577 (M+H)⁺, Rt= 2.25 minutes (method A)

Example 68

2-(1-Acetylazetidin-3-yl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0219] To a solution of tert-butyl 3-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-2-anyl}azetidine-1-carboxylate (Example 61) (50 mg) in methylene chloride (2 mL), was added thioanisole (50 μ L). The mixture was added with TFA (1 mL) under ice-cooling, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was poured into saturated aqueous sodium bicarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was triturated in n-hexane/ethyl acetate, and the precipitate was collected by filtration and dried to obtain 2-(azetidin-3-yl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (38 mg). This was dissolved in THF (1 mL), and added with pyridine (23 μ L), and the mixture was stirred under ice-cooling. The mixture was added dropwise with acetyl chloride (7.7 μ L), and stirred at room temperature for 2 hours. The reaction mixture was poured into ice water, and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on column chromatography to obtain 2-(1-acetylazetidin-3-yl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (7.5 mg). This was dissolved in MeOH (0.5 mL), treated with 1M hydrogen chloride/MeOH, and concentrated to obtain the titled compound (5.5 mg) as slightly gray powder. MS(ESI+) m/z 570 (M+H)⁺, Rt= 1.98 minutes (method A)

Example 69

tert-Butyl

(2S)-2-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}pyrrolidine-1-carboxylate

[0220] The titled compound was obtained as white powder according to the procedures as described in Steps 1 to 7 of Example 11, using tert-butyl (2S)-2-(chlorocarbonyl)pyrrolidine-1-carboxylate (prepared as described in Tetrahedron Asymmetry, 2007, 18, 2011) instead of methoxyacetyl chloride. MS(ESI+) m/z 642 (M+H)⁺, Rt= 2.77 minutes (method A)

Example 70

tert-Butyl

(2R)-2-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}pyrrolidine-1-carboxylate

[0221] The titled compound was obtained as white powder according to the procedures as described in Steps 1 to 7 of Example 11, using tert-butyl (2R)-2-(chlorocarbonyl)pyrrolidine-1-carboxylate (prepared as described in Tetrahedron Asymmetry, 2007, 18, 2011) instead of methoxyacetyl chloride. MS(ESI+) m/z 642 (M+H)⁺, Rt= 2.77 minutes (method A)

Example 71

N-(3-Chloro-2-methylphenyl)-2-[(2S)-pyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide dihydrochloride

[0222] To a solution of tert-butyl (2S)-2-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-2-yl}pyrrolidine-1-carboxylate (26 mg) (Example 69) in ethyl acetate (2 mL), was added 4N hydrogen chloride/ethyl acetate solution (2 mL), and the mixture was stirred at room temperature for 2 hours. N-hexane was added to the reaction mixture, the precipitates was collected, and dried to obtain the titled compound (20 mg) as pale yellow powder. MS(ESI+) m/z 542 (M+H)⁺, Rt= 1.65 minutes (method A)

Example 72

N-(3-Chloro-2-methylphenyl)-2-[(2S)-1-methylpyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0223] To a solution of N-(3-chloro-2-methylphenyl)-2-[(2S)-pyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (Example 71) (25 mg) in MeOH (0.4 mL), was added 38% aqueous formaldehyde solution (7.2 μ L), and the mixture was stirred at room temperature overnight. The mixture was added with 2-picoline borane (10 mg), and stirred for 3 hours, removed the solvent under reduced pressure. The residue was purified on column chromatography to obtain the titled compound (11 mg) as white powder.

MS(ESI+) m/z 556 (M+H)⁺, Rt= 1.67 minutes (method A)

Example 73

2-[(2S)-1-Acetylpyrrolidin-2-yl]-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0224] To a solution of N-(3-chloro-2-methylphenyl)-2-[(2S)-pyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (Example 71) (26 mg) in THF solution (0.5 mL), was added pyridine (12 μ L), and the mixture was stirred under ice-cooling. The mixture was added with acetyl chloride (5.1 μ L), and stirred at room temperature for 1 hour. The reaction mixture was added with ice water, and the mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and removed the solvent under reduced pressure. The residue was purified on column chromatography to obtain the titled compound (15 mg) as white powder.

MS(ESI+) m/z 584 (M+H)⁺, Rt= 2.26 minutes (method A)

Example 74

N-(3-Chloro-2-methylphenyl)-2-[(2-methoxyethoxy)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0225] The titled compound was obtained as white powder according to the procedure as described in Example 11, using 2-(methoxyethoxy)acetyl chloride instead of methoxyacetyl chloride.

MS(ESI+) m/z 561 (M+H)⁺, Rt= 2.43 minutes (method A)

Example 75

N-(3-Chloro-2-methylphenyl)-2-(1-methoxy-2-methylpropan-2-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0226] The titled compound was obtained as white powder according to the procedure as described in Example 11, using 3-methoxy-2,2-dimethylpropanoyl chloride (prepared as described in Bull. Chem. Soc. Jpn., 2001, 74, 1695) instead of methoxyacetyl chloride.

MS(ESI+) m/z 559 (M+H)⁺, Rt= 2.82 minutes (method A)

Example 76

2-tert-Butyl-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0227] The titled compound was obtained as white powder according to the procedure as described in Example 11, using pivaloyl chloride instead of methoxyacetyl chloride. MS(ESI+) m/z 529 (M+H)⁺, Rt= 3.03 minutes (method A)

Example 77

2-tert-Butyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-N-({[3-(trifluoromethyl)pyridin-2-yl]methyl})-1H-benzimidazole-4-carboxamide dihydrochloride

[0228] The titled compound was obtained as white powder according to the procedure as described in Example 11, using pivaloyl chloride instead of methoxyacetyl chloride, and 2-(aminomethyl)-3-(trifluoromethyl)pyridine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 564 (M+H)⁺, Rt= 2.39 minutes (method A)

Example 78

N-(3-Chloro-2-methylphenyl)-2-(2-ethoxyethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0229] The titled compound was obtained as white powder according to the procedure as described in Example 11, using 3-ethoxy-propionylchloride (prepared as described in EP1803350) instead of methoxyacetyl chloride.

MS(ESI+) m/z 545 (M+H)⁺, Rt= 2.69 minutes (method A)

Example 79

N-(3-Chloro-2-methylphenyl)-2-(ethoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0230] The titled compound was obtained as white powder according to the procedure as described in Example 11, using ethoxyacetyl chloride (prepared as described in US2004/39038) instead of methoxyacetyl chloride.

MS(ESI+) m/z 531 (M+H)⁺, Rt= 2.77 minutes (method A)

Example 80

2-(Ethoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-N-({[3-(trifluoromethyl)pyridin-2-yl]methyl})-1H-benzimidazole-4-carboxamide dihydrochloride

[0231] The titled compound was obtained as white powder according to the procedure as described in Example 11, using ethoxyacetyl chloride (prepared as described in US2004/39038) instead of methoxyacetyl chloride, and 2-(aminomethyl)-3-(trifluoromethyl)pyridine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 566 (M+H)⁺, Rt= 2.18 minutes (method A)

Example 81

N-(3-Chloro-2-methylphenyl)-2-(2-methoxyethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0232] The titled compound was obtained as white powder according to the procedure as described in Example 11, using methoxypropionyl chloride instead of methoxyacetyl chloride.

MS(ESI+) m/z 531 (M+H)⁺, Rt= 2.58 minutes (method A)

Example 82

N-(3-Chloro-2-methylphenyl)-2-(2,2-dimethylpropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0233] The titled compound was obtained as white powder according to the procedure as described in Example 11, using tert-butylacetyl chloride instead of methoxyacetyl chloride.

MS(ESI+) m/z 543 (M+H)⁺, Rt= 3.25 minutes (method A)

Example 83

N-(3-Chloro-2-methylphenyl)-2-cyclopropyl-6-({[2-(trifluoro methyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0234] The titled compound was obtained as white powder according to the procedure as described in Example 11, using cyclopropanecarbonyl chloride instead of methoxyacetyl chloride.

MS(ESI+) m/z 513 (M+H)⁺, Rt= 2.91 minutes (method A)

Example 84

N-(3-chloro-2-methylphenyl)-2-(2-methylpentan-2-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0235] The titled compound was obtained as white powder according to the procedure as described in Example 11, using 2,2-dimethylpentanoyl chloride (prepared as described in J. Am. Chem. Soc., 1974, 96, 1518) instead of methoxyacetyl chloride. MS(ESI+) m/z 557 (M+H)⁺, Rt= 3.16 minutes (method B)

Example 85

N-(3-Chloro-2-methylphenyl)-2-(1-methylcyclopropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0236] The titled compound was obtained as white powder according to the procedure as described in Example 11, using 1-methyl-cyclopropanecarbonyl chloride (prepared as described in WO2009/68512) instead of methoxyacetyl chloride.

MS(ESI+) m/z 527 (M+H)⁺, Rt= 3.16 minutes (method A)

Elemental Analysis for C₂₇H₂₂ClF₃N₄O₂•HCl+1.0H₂O

Calcd.(%) C:55.78 H:4.33 N:9.64

Fond.(%) C:55.48 H:3.94 N:9.63

Example 86

2-tert-Butyl-N-(3-chloro-4-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0237] The titled compound was obtained as white powder according to the procedure as described in Example 11, using pivaloyl chloride instead of methoxyacetyl chloride, and 3-chloro-4-methylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 529 (M+H)⁺, Rt= 2.83 minutes (method B)

Example 87

2-tert-Butyl-N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 6-amino-2-tert-butyl-1H-benzimidazole-4-carboxylate

[0238] The title compound was obtained as white powder according to the procedures as described in step 1 to step 4 of Example 11, using pivaloyl chloride instead of methoxyacetyl chloride in Step 1.

[Step 2]

2-tert-Butyl-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid

[0239] The title compound was obtained according to the procedures as described in step 4 and step 5 of Example

1, using methyl 6-amino-2-tert-butyl-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate, and 2,5-dichlorobenzoyl chloride instead of 2-(trifluoromethyl)benzoyl chloride.

[Step 3]

2-tert-Butyl-N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0240] The title compound was obtained according to the procedures as described in step 7 and step 8 of Example 11, using 2-tert-Butyl-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid. MS(ESI+) m/z 529 (M+H)⁺, Rt= 2.97 minutes (method B)

Example 88

2-tert-Butyl-N-(3-chloro-4-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0241] The titled compound was obtained as white powder according to the procedures as described in step 7 and step 8 of Example 11, using 2-tert-Butyl-6-({[2,5-dichlorophenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid, and 3-chloro-4-methylaniline instead of 3-chloro-2-methylaniline. MS(ESI+) m/z 529 (M+H)⁺, Rt= 3.05 minutes (method B)

Example 89

N-(3-Chloro-2-methylphenyl)-2-[1-(trifluoromethyl)cyclopropyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0242] The titled compound was obtained as white powder according to the procedures as described in Steps 1 to 7 of Example 11, using 1-(trifluoromethyl)cyclopropanecarbonyl chloride (prepared as described in WO2005/23773) instead of methoxyacetyl chloride. MS(ESI+) m/z 581 (M+H)⁺, Rt= 3.15 minutes (method A)

Example 90

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[Step 1]

2-(Methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid

[0243] The titled compound was obtained according to the procedure as described in Example 46, using methyl 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate instead of methyl 6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate.

[Step 2]

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0244] The titled compound was obtained as colorless powder according to the procedure as described in Step 7 of Example 11, using 2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid. MS(ESI+) m/z 531 (M+H)⁺, Rt= 2.58 minutes (method A)

Example 91

N-(2-Chlorobenzyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0245] The titled compound was obtained as colorless powder according to the procedure as described in Step 7 of Example 11, using 2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid, and 2-chloro benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 531 (M+H)⁺, Rt= 2.25 minutes (method A)

Example 92

6-({[2-Chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

6-Amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide

[0246] To a suspension of methyl 2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxylate (Example 11, Step 3) (4.8g) in MeOH (120 mL), were added lithium hydroxide hydrate (4.56g) and water (54 mL). The mixture was stirred at room temperature for 16 hours. Under ice-cooling, 1N hydrochloric acid was added slowly to the reaction mixture to adjust the pH to about 2. The precipitate was collected by filtration and dried to obtain 2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxylic acid (4.6g). To the solution of 2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxylic acid (4.6g) in DMF (90 mL), HBTU (8.23g), triethylamine (2.19g) and 3-chloro-2-methylaniline (3.07g) were added sequentially, and the solution was stirred for 18 hours. The reaction mixture was poured into saturated aqueous sodium bicarbonate and stirred for 1 hour. The precipitate was collected by filtration and washed sequentially with water and ethyl acetate to obtain N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxamide (7.2g) as pale yellow powder. To the solution of N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxamide (7.2g) in MeOH-THF (1:1, 120 mL), were added 1% platinum + 0.1% copper-activated carbon (Degussa type CF105 R/W) (1.4g). The mixture was stirred under hydrogen atmosphere (3 atm) for 4 hours. The catalyst was filtered off, and the mother liquid was concentrated. The residual solid was washed with ethyl acetate to obtain 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide (5.1g) as pale yellow powder.

[Step 2]

6-({[2-Chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0247] To a solution of 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide (40 mg) in THF (1 mL), was added N,N-diisopropylethylamine (24 μ L), and the mixture was stirred under ice-cooling. The mixture was added with 2-chloro-6-fluorobenzoyl chloride (24 mg), and stirred for 3 hours. The reaction mixture was added with saturated aqueous sodium bicarbonate, and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide (50 mg). This was dissolved in ethyl acetate (2 mL), and treated with 4N hydrogen chloride/ethyl acetate (1.2 Eq). The precipitate was collected by filtration, and dried to obtain the titled compound (40 mg) as white powder.

MS(ESI+) m/z 501 (M+H)⁺, Rt= 2.39 minutes (method A)

Example 93

6-({[2-Chloro-4-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-methoxymethyl-1H-benzimidazole-4-carboxamide hydrochloride

[0248] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 2 of Example 92, using 2-chloro-4-fluorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

EP 2 746 265 B1

MS(ESI+) m/z 501 (M+H)⁺, Rt= 2.40 minutes (method A)

Example 94

5 6-[[[(2-Chloro-5-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0249] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2-chloro-5-fluorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

10 MS(ESI+) m/z 501 (M+H)⁺, Rt= 2.41 minutes (method A)

Example 95

15 N-(3-Chloro-2-methylphenyl)-6-[[[(2-chlorophenyl)carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0250] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2-chlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

20 MS(ESI+) m/z 483 (M+H)⁺, Rt= 2.36 minutes (method A)

Example 96

25 N-(3-Chloro-2-methylphenyl)-6-[[[(2-chloropyridin-3-yl)carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0251] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2-chloronicotinoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

30 MS(ESI+) m/z 484 (M+H)⁺, Rt= 2.16 minutes (method A)

Example 97

35 6-[[[(2-Bromophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0252] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2-bromobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 527 (M+H)⁺, Rt= 2.39 minutes (method A)

Example 98

40 N-(3-Chloro-2-methylphenyl)-6-[[[(2,6-dichlorophenyl)carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0253] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2,6-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

45 MS(ESI+) m/z 517 (M+H)⁺, Rt= 2.46 minutes (method A)

Example 99

50 N-(3-Chloro-2-methylphenyl)-6-[[[(2,5-dichlorophenyl)carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0254] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2,5-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

55 MS(ESI+) m/z 517 (M+H)⁺, Rt= 2.54 minutes (method A)

Example 100

6-[[[2-Chloro-3-fluorophenyl]carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0255] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2-chloro-3-fluorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 501 (M+H)⁺, Rt= 2.43 minutes (method A)

Example 101

6-[[[2-Chloro-3,6-difluorophenyl]carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0256] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2-chloro-3,6-difluorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 519 (M+H)⁺, Rt= 2.46 minutes (method A)

Example 102

6-[[[2-Bromo-6-chlorophenyl]carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0257] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 2-chloro-6-bromobenzoyl chloride (prepared as described in WO2008/124575) instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 561 (M+H)⁺, Rt= 2.50 minutes (method A)

Example 103

6-[[[2-Bromo-6-fluorophenyl]carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0258] To a solution of 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide (50 mg) in DMF (1.5 mL), were added HBTU (66 mg), N,N-diisopropylethylamine (30 μ L) and 2-bromo-6-fluorobenzoic acid (38 mg) in turn, and the mixture was stirred at room temperature for 24 hours. The reaction mixture was poured into saturated aqueous sodium bicarbonate, and extracted with ethyl acetate. The organic layer was separated, washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain 6-[[[2-bromo-6-fluorophenyl]carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide (46 mg). This was dissolved in ethyl acetate (2 mL), and treated with 4N hydrogen chloride/ethyl acetate (1.2 Eq). The precipitate was collected by filtration, and dried to obtain the titled compound (24 mg) as white powder.

MS(ESI+) m/z 545 (M+H)⁺, Rt= 2.44 minutes (method A)

Example 104

N-(3-Chloro-2-methylphenyl)-6-[[[2-chloro-6-methylphenyl]carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0259] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-6-methylbenzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 497 (M+H)⁺, Rt= 2.44 minutes (method A)

EP 2 746 265 B1

Example 105

N-(3-Chloro-2-methylphenyl)-6-[[2-(chloro-4-methylphenyl)carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0260] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-4-methylbenzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 497 (M+H)⁺, Rt= 2.47 minutes (method A)

Example 106

6-[[5-Bromo-2-chlorophenyl]carbonyl]amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0261] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 5-bromo-2-chlorobenzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 561 (M+H)⁺, Rt= 2.62 minutes (method A)

Example 107

6-[[2-Bromo-5-chlorophenyl]carbonyl]amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0262] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-bromo-5-chlorobenzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 561 (M+H)⁺, Rt= 2.63 minutes (method A)

Example 108

N-(3-Chloro-2-methylphenyl)-6-[[2-(chloro-5-methylphenyl)carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0263] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-5-methylbenzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 497 (M+H)⁺, Rt= 2.54 minutes (method A)

Example 109

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-6-[[5-methyl-2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0264] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 5-methyl-2-(trifluoromethyl)benzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 531 (M+H)⁺, Rt= 2.59 minutes (method A)

Example 110

6-[[2,5-Bis(trifluoromethyl)phenyl]carbonyl]amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0265] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2,5-bis(trifluoromethyl)benzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 585 (M+H)⁺, Rt= 2.76 minutes (method A)

Example 111

6-({[2,4-Bis(trifluoromethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0266] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2,4-bis(trifluoromethyl)benzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 585 (M+H)⁺, Rt= 2.80 minutes (method A)

Example 112

N-(3-Chloro-2-methylphenyl)-6-({[5-fluoro-2-(trifluoromethyl)phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0267] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 5-fluoro-2-(trifluoromethyl)benzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 535 (M+H)⁺, Rt= 2.68 minutes (method A)

Example 113

N-(3-Chloro-2-methylphenyl)-6-({[2-chloro-6-(trifluoromethyl)phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0268] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-6-(trifluoromethyl)benzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 551 (M+H)⁺, Rt= 2.76 minutes (method A)

Example 114

N-(3-Chloro-2-methylphenyl)-6-({[2-chloro-5-[2-(propan-2-yl oxy)ethoxy]phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0269] Under argon atmosphere, potassium tert-butoxide (168 mg) was added to a solution of 2-isopropoxyethanol (172 μ L) in NMP (1.5 mL). The mixture was stirred at room temperature for 10 minutes, added with 2-chloro-5-fluorobenzoic acid (87 mg), and then stirred at 130°C for 4 hours. The reaction mixture was cooled on ice bath, and added with water, and then acidified with 1N hydrochloric acid and extracted with ethyl acetate. The organic layer was separated, washed with brine, and dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to obtain crude 2-chloro-5-[2-(propan-2-yloxy)ethoxy]benzoic acid. This was dissolved in DMF (2 mL), added with HATU (61 mg), N,N-diisopropylethylamine (29 μ L) and 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide (50 mg) in this order, and stirred at room temperature for 12 hours. The reaction mixture was poured into saturated aqueous sodium bicarbonate, and extracted with ethyl acetate. The organic layer was separated, washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain N-(3-chloro-2-methylphenyl)-6-({[2-chloro-5-[2-(propan-2-yl oxy)ethoxy]phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide (41 mg) as white powder. This was dissolved in ethyl acetate (2 mL), and treated with 4N hydrogen chloride/ethyl acetate (1.2 Eq). The precipitate was collected by filtration, and dried to obtain the titled compound (28 mg) as white powder.

MS(ESI+) m/z 585 (M+H)⁺, Rt= 3.14 minutes (method A)

Example 115

6-({[2-Chloro-5-(2-ethoxyethoxy)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0270] The titled compound was obtained as white powder according to the procedure as described in Example 114, using 2-ethoxyethanol instead of 2-isopropoxy ethanol.

MS(ESI+) m/z 571 (M+H)⁺, Rt= 2.71 minutes (method A)

Example 116

6-([2-Chloro-5-(3-methoxypropyl)phenyl]carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0271] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 103, using 2-chloro-5-(3-methoxypropyl)benzoic acid (prepared as described in Bioorg. Med. Chem. Lett., 2010, 20, 2204) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 555 (M+H)⁺, Rt= 2.73 minutes (method A)

Example 117

6-([5-(3-tert-Butoxy-1-propynyl)-2-chlorophenyl]carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0272] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 103, using 5-(3-tert-butoxy-1-propynyl)-2-chlorobenzoic acid (Reference Example 6) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 593 (M+H)⁺, Rt= 3.17 minutes (method A)

Example 118

6-([5-(3-tert-Butoxypropyl)-2-chlorophenyl]carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0273] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 5-(3-tert-butoxypropyl)-2-chlorobenzoic acid (Reference Example 7) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 597 (M+H)⁺, Rt= 3.28 minutes (method A)

Example 119

6-([2-Chloro-5-(3-hydroxy-3-methylbutyl)phenyl]carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0274] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 103, using 2-chloro-5-(3-hydroxy-3-methylbutyl)benzoic acid (Reference Example 8) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 569 (M+H)⁺, Rt= 2.64 minutes (method A)

Example 120

6-([2-Chloro-5-(ethoxymethyl)phenyl]carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide

[0275] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-5-(ethoxymethyl)benzoic acid (Reference Example 1) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 541 (M+H)⁺, Rt= 2.52 minutes (method A)

Example 121

6-([2-Chloro-5-[(2-ethoxyethoxy)methyl]phenyl]carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0276] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-5-[(2-ethoxyethoxy)methyl]benzoic acid (Reference Example 2) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 585 (M+H)⁺, Rt= 2.51 minutes (method A)

Example 122

6-([2-Chloro-5-(2-cyclopropylethyl)phenyl]carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide

[0277] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-5-(2-cyclopropylethyl)benzoic acid (Reference Example 4) instead of 2-bromo-6-fluorobenzoic acid. MS(ESI+) m/z 551 (M+H)⁺, Rt= 2.92 minutes (method A)

Example 123

N-(3-Chloro-2-methylphenyl)-6-([2-chloro-5-(2-phenylethyl)phenyl]carbonyl)amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0278] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-5-(2-phenylethyl)benzoic acid (Reference Example 5) instead of 2-bromo-6-fluorobenzoic acid. MS(ESI+) m/z 587 (M+H)⁺, Rt= 2.98 minutes (method A)

Example 124

N-(3-Chloro-2-methylphenyl)-2-cyclopentyl-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 2-cyclopentyl-6-nitro-1H-benzimidazole-4-carboxylate

[0279] The titled compound was obtained as pale yellow powder according to the procedure as described in Steps 1 to 3 of Example 11, using cyclopentanecarbonyl chloride instead of methoxyacetyl chloride.

[Step 2]

6-Amino-N-(3-chloro-2-methylphenyl)-2-cyclopentyl-1H-benzimidazole-4-carboxamide

[0280] The titled compound was obtained as yellow powder according to the procedure as described in Step 1 of Example 92, using methyl 2-cyclopentyl-6-nitro-1H-benzimidazole-4-carboxylate instead of methyl 2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxylate.

[Step 3]

N-(3-Chloro-2-methylphenyl)-2-cyclopentyl-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0281] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 6-amino-N-(3-chloro-2-methylphenyl)-2-cyclopentyl-1H-benzimidazole-4-carboxamide instead of 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide, and 2-(trifluoromethyl)benzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride. MS(ESI+) m/z 541 (M+H)⁺, Rt= 3.21 minutes (method A)

Example 125

N-(3-Chloro-2-methylphenyl)-2-cyclopentyl-6-([2,5-dichlorophenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0282] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 6-amino-N-(3-chloro-2-methylphenyl)-2-cyclopentyl-1H-benzimidazole-4-carboxamide instead of 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide, and 2,5-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 543 (M+H)⁺, Rt= 3.32 minutes (method A)

Example 126

5 6-[[2-Chloro-6-fluorophenyl]carbonyl]amino}-N-(3-chloro-2-methylphenyl)-2-cyclopentyl-1H-benzimidazole-4-carboxamide hydrochloride

10 **[0283]** The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 6-amino-N-(3-chloro-2-methylphenyl)-2-cyclopentyl-1H-benzimidazole-4-carboxamide instead of 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide.
MS(ESI+) m/z 525 (M+H)⁺, Rt= 3.24 minutes (method A)

Example 127

15 6-[[2-Chloro-6-fluorophenyl]carbonyl]amino}-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

20 Methyl

6-nitro-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxylate

25 **[0284]** The titled compound was obtained as pale yellow powder according to the procedure as described in Steps 1 to 3 of Example 11, using (2R)-tetrahydro-2-furancarboxyl chloride (prepared as described in WO2006/79642) instead of methoxyacetyl chloride.

[Step 2]

30 6-Amino-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide

[0285] The titled compound was obtained as yellow powder according to the procedure as described in Step 1 of Example 92, using methyl 6-nitro-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxylate instead of methyl 2-(methoxymethyl)-6-nitro-1H-benzimidazole-4-carboxylate.

35 [Step 3]

40 6-[[2-Chloro-6-fluorophenyl]carbonyl]amino}-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide hydrochloride

[0286] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 6-amino-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide instead of 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide.
MS(ESI+) m/z 527 (M+H)⁺, Rt= 2.75 minutes (method A)

45 Example 128

50 N-(3-Chloro-2-methylphenyl)-6-[[2,6-dichlorophenyl]carbonyl]amino}-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide hydrochloride

[0287] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 2 of Example 92, using 6-amino-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide instead of 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide, and 2,6-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.
MS(ESI+) m/z 543 (M+H)⁺, Rt= 2.80 minutes (method A)

Example 129

N-(3-Chloro-2-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide hydrochloride

[0288] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 92, using 6-amino-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide instead of 6-amino-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide, and 2,5-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 543 (M+H)⁺, Rt= 2.86 minutes (method A)

Example 130

N-(3-Chloro-2-methylphenyl)-2-[(2S)-5-oxopyrrolidin-2-yl]-6-([(2-(trifluoromethyl)phenyl)carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0289] The titled compound was obtained as white powder according to the procedure as described in Example 11, using (S)-(-)-2-pyrrolidone 5-carbonyl chloride (prepared as described in Tetrahedron Lett., 1997, 38, 2259) instead of methoxyacetyl chloride.

MS(ESI+) m/z 556 (M+H)⁺, Rt= 2.03 minutes (method A)

Example 131

N-(3-Chloro-2-methylphenyl)-2-[(2R)-5-oxopyrrolidin-2-yl]-6-([(2-(trifluoromethyl)phenyl)carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0290] The titled compound was obtained as white powder according to the procedure as described in Example 11, using (R)-(+)-2-pyrrolidone-5-carbonyl chloride (prepared as described in Tetrahedron Lett., 1997, 38, 2259) instead of methoxyacetyl chloride.

MS(ESI+) m/z 556 (M+H)⁺, Rt= 2.03 minutes (method A)

Example 132

N-(3-Chloro-2-methylphenyl)-2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([(2-(trifluoromethyl)phenyl)carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 2-(2-ethoxy-2-oxoethyl)-1H-benzimidazole-4-carboxylate

[0291] A solution of methyl 2,3-diaminobenzoate (680 mg) and ethyl 3-amino-3-ethoxyacrylate hydrochloride (880 mg) in EtOH (12 mL) was stirred at 60°C for 2 hours. EtOH was removed under reduced pressure, and the residue was purified on column chromatography to obtain the titled compound (1.14g) as a pale yellow solid.

[Step 2]

Methyl 2-(2-ethoxy-2-oxoethyl)-6-([(2-(trifluoromethyl)phenyl)carbonyl]amino)-1H-benzimidazole-4-carboxylate

[0292] The titled compound was obtained as pale yellow powder according to the procedures as described in Steps 3 to 5 of Example 11, using methyl 2-(2-ethoxy-2-oxoethyl)-1H-benzimidazole-4-carboxylate instead of methyl 1H-benzimidazole-4-carboxylate as a starting material.

[Step 3]

Methyl 2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([(2-(trifluoromethyl)phenyl)carbonyl]amino)-1H-benzimidazole-4-carboxylate

[0293] Pyrrolidine (1 mL) was added to methyl 2-(2-ethoxy-2-oxoethyl)-6-([(2-(trifluoromethyl)phenyl)carbonyl]amino)-

1H-benzimidazole-4-carboxylate (80 mg), and the solution was stirred at 70°C for 30 minutes. The reaction mixture was diluted with ethyl acetate, and washed with brine. The ethyl acetate layer was concentrated under reduced pressure. The residue was purified on column chromatography to obtain the titled compound (74 mg) as white powder.

5 [Step 4]

2-[2-Oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid

10 **[0294]** A solution of methyl 2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate (74 mg), lithium hydroxide hydrate (66 mg) in MeOH-water (1:1, 2 mL) was stirred at 50°C for 30 minutes. The reaction mixture was removed the solvent under reduced pressure, added with water, and to neutralize by 1N hydrochloric acid under ice-cooling. The precipitate was collected by filtration, washed with diethyl ether, and dried under reduced pressure to obtain the titled compound (60 mg) as pale yellow powder.

15 [Step 5]

N-(3-Chloro-2-methylphenyl)-2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

20 **[0295]** To a solution of 2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid (30 mg) and HATU (30 mg) in DMF (0.5 mL), were added N,N-diisopropylethylamine (28 µL) and 3-chloro-2-methylaniline (10 µL), and the mixture was stirred at room temperature for 15 minutes. The reaction mixture was diluted with ethyl acetate, washed with brine, and the organic layer was concentrated under reduced pressure. The residue was purified on column chromatography to obtain pale yellow powder (28 mg). This was suspended in MeOH (1 mL), and 2N hydrogen chloride/EtOH solution (1 Eq) was added to obtain a homogeneous solution, which was then concentrated under reduced pressure. The residue was triturated in diethyl ether and washed. The precipitate was collected by filtration and dried under reduced pressure to obtain the titled compound (20 mg) as white powder. MS(ESI+) m/z 584 (M+H)⁺, Rt= 2.24 minutes (method A)

30 Example 133

N-(3-Chloro-2-methylphenyl)-2-[2-(dimethylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

35 [Step 1]

Methyl 2-[2-(dimethylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate

40 **[0296]** The titled compound was obtained as pale yellow powder according to the procedure as described in Step 3 of Example 132, using 40% aqueous dimethylamine solution and THF instead of pyrrolidine.

[Step 2]

45 2-[2-(Dimethylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid

[0297] The titled compound was obtained as gray powder according to the procedure as described in Step 4 of Example 132, using methyl 2-[2-(dimethylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate instead of methyl 2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate.

[Step 3]

55 N-(3-Chloro-2-methylphenyl)-2-[2-(dimethylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0298] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 5 of Example 132, using 2-[2-(dimethylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid instead of methyl 2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate.

EP 2 746 265 B1

zole-4-carboxylic acid instead of 2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid. MS(ESI+) m/z 558 (M+H)⁺, Rt= 2.12 minutes (method A)

Example 134

N-(3-Chloro-2-methylphenyl)-2-[2-(methylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 2-[2-(methylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate

[0299] The titled compound was obtained as white powder according to the procedure as described in Step 3 of Example 132, using 40% aqueous methylamine solution and THF instead of pyrrolidine.

[Step 2]

2-[2-(Methylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid

[0300] The titled compound was obtained as brown powder according to the procedure as described in Step 4 of Example 132, using methyl 2-[2-(methylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate instead of methyl 2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate.

[Step 3]

N-(3-Chloro-2-methylphenyl)-2-[2-(methylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0301] The titled compound was obtained as pale red powder according to the procedure as described in Step 5 of Example 132, using 2-[2-(methylamino)-2-oxoethyl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid instead of 2-[2-oxo-2-(pyrrolidin-1-yl)ethyl]-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid. MS(ESI+) m/z 544 (M+H)⁺, Rt= 1.98 minutes (method A)

Example 135

2-Chloro-N-(3-chloro-2-methylphenyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[Step 1]

Methyl 2-chloro-1H-benzimidazole-4-carboxylate

[0302] A solution of methyl 2-oxo 2,3-dihydro-1H-benzimidazole-4-carboxylate (prepared as described in US2009/186879) (2.0g) in phosphoryl chloride (18 mL) was stirred at 120°C for 2.5 hours. Excessive phosphoryl chloride was removed under reduced pressure. The residue was added with saturated aqueous sodium bicarbonate under ice-cooling, and extracted with ethyl acetate. The ethyl acetate layer was dried over anhydrous magnesium sulfate and concentrated. The residue was purified on column chromatography to obtain the titled compound (1.5g) as white powder.

[Step 2]

Methyl 2-chloro-6-nitro-1H-benzimidazole-4-carboxylate

[0303] The titled compound (0.98g) was obtained as pale yellow powder according to the procedure as described in Step 2 of Example 1, using methyl 2-chloro-1H-benzimidazole-4-carboxylate (1.46g) instead of methyl-1H-benzimidazole-4-carboxylate.

[Step 3]

Methyl 6-amino-2-chloro-1H-benzimidazole-4-carboxylate

[0304] To a solution of methyl 2-chloro-6-nitro-1H-benzimidazole-4-carboxylate (0.87g) in MeOH (100 mL), was added 1% platinum + 0.1% copper-activated carbon (Degussa type CF105 R/W) (0.19g), and the mixture was stirred under hydrogen atmosphere (0.2 MPa) for 4 hours. The reaction mixture was filtered off through celite and washed with MeOH. The mother liquid was concentrated to obtain the titled compound (0.78g) as yellow powder.

[Step 4]

Methyl

2-chloro-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate

[0305] The titled compound (0.69g) was obtained as colorless powder according to the procedure as described in Step 4 of Example 1, using methyl 6-amino-2-chloro-1H-benzimidazole-4-carboxylate (0.78g) instead of methyl 6-amino-1H-benzimidazole-4-carboxylate.

[Step 5]

2-Chloro-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid

[0306] The titled compound (620 mg) was obtained as white powder according to the procedure as described in Step 5 of Example 1, using methyl 2-chloro-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate (688 mg) instead of methyl 6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylate.

[Step 6]

2-Chloro-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0307] Oxalyl chloride (1.42 mL) was added dropwise to a stirring solution of 2-chloro-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid (1.27g) in THF (13 mL) under ice-cooling. The mixture was added with DMF (five drops), and stirred at room temperature for 2 hours. From the reaction mixture was removed the solvent and excessive oxalyl chloride under reduced pressure. The residue was added with toluene, and concentrated under reduced pressure, dried, and then dissolved in THF (10 mL). The solution was dropped slowly to a solution of 3-chloro-2-methylaniline (0.42 mL) and N,N-diisopropylethylamine (2.3 mL) in THF (10 mL) under ice-cooling for 1 hour. The reaction mixture was diluted with ethyl acetate, washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain the titled compound (0.58g) as white powder.

MS(ESI⁺) m/z 507 (M+H)⁺, Rt= 2.56 minutes (method A)

Example 136

N-(3-Chloro-2-methylphenyl)-2-[(2-methoxyethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0308] 2-Methoxyethylamine (2 mL) was added to 2-chloro-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (40 mg, Example 135). The mixture was stirred at 150 °C in a sealed pressure-proof stainless steel container for 17 hours. The reaction mixture was washed with brine and concentrated under reduced pressure. The residue was purified on column chromatography to obtain N-(3-chloro-2-methylphenyl)-2-[(2-methoxyethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (35 mg) as white powder. This was dissolved in MeOH (1 mL), was added with 2N hydrogen chloride/EtOH solution (1 Eq), stirred and concentrated. The residue was triturated in diethyl ether, and the precipitate was collected by filtration and dried to obtain the titled compound (28 mg) as white powder.

MS(ESI⁺) m/z 546 (M+H)⁺, Rt= 1.80 minutes (method A)

Example 137

N-(3-Chloro-2-methylphenyl)-2-[(2-hydroxyethyl)amino]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0309] The titled compound was obtained as brown powder according to the procedure as described in Example 136, using 2-aminoethanol instead of 2-methoxyethylamine.

MS(ESI+) m/z 532 (M+H)⁺, Rt= 1.58 minutes (method A)

Example 138

N-(3-Chloro-2-methylphenyl)-2-(methylamino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0310] The titled compound was obtained as white powder according to the procedure as described in Example 136, using 40% aqueous methylamine solution instead of 2-methoxyethylamine.

MS(ESI+) m/z 502 (M+H)⁺, Rt= 1.69 minutes (method A)

Example 139

N-(3-Chloro-2-methylphenyl)-2-(ethylamino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0311] The titled compound was obtained as white powder according to the procedure as described in Example 136, using 70% aqueous ethylamine solution instead of 2-methoxyethylamine.

MS(ESI+) m/z 516 (M+H)⁺, Rt= 1.80 minutes (method A)

Example 140

N-(3-Chloro-2-methylphenyl)-2-[(2,2-dimethylpropyl)amino]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0312] The titled compound was obtained as white powder according to the procedure as described in Example 136, using neopentylamine instead of 2-methoxyethylamine.

MS(ESI+) m/z 558 (M+H)⁺, Rt= 2.22 minutes (method A)

Elemental Analysis for C₂₈H₂₇Cl₃N₅O₂•HCl+0.5H₂O

Calcd.(%) C:55.73 H:4.84 N:11.61

Found.(%) C:55.99 H:4.75 N:11.69

Example 141

N-(3-Chloro-2-methylphenyl)-2-(cyclopentylamino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0313] The titled compound was obtained as white powder according to the procedure as described in Example 136, using cyclopentylamine instead of 2-methoxyethylamine.

MS(ESI+) m/z 556 (M+H)⁺, Rt= 2.10 minutes (method A)

Example 142

N-(3-Chloro-2-methylphenyl)-2-(piperidin-1-yl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0314] The titled compound was obtained as white powder according to the procedure as described in Example 136, using piperidine instead of 2-methoxyethylamine.

MS(ESI+) m/z 556 (M+H)⁺, Rt= 2.54 minutes (method A)

EP 2 746 265 B1

Example 143

N-(3-Chloro-2-methylphenyl)-2-(4-methylpiperazin-1-yl)-6-({ [2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide dihydrochloride

[0315] The titled compound was obtained as white powder according to the procedure as described in Example 136, using 1-methylpiperazine instead of 2-methoxyethylamine.

MS(ESI+) m/z 571 (M+H)⁺, Rt= 1.63 minutes (method A)

Example 144

2-[Bis(2-hydroxyethyl)amino]-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0316] The titled compound was obtained as white powder according to the procedure as described in Example 136, using diethanolamine instead of 2-methoxyethylamine.

MS(ESI+) m/z 576 (M+H)⁺, Rt= 1.66 minutes (method A)

Example 145

N-(3-Chloro-2-methylphenyl)-2-(dimethylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0317] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 50% aqueous dimethylamine solution instead of 2-methoxyethylamine.

MS(ESI+) m/z 516 (M+H)⁺, Rt= 2.13 minutes (method A)

Elemental Analysis for C₂₅H₂₁ClF₃N₅O₂•HCl+0.5H₂O

Calcd.(%) C:53.49 H:4.13 N:12.48

Found.(%) C:53.44 H:4.17 N:12.39

Example 146

N-(3-Chloro-2-methylphenyl)-2-{[2-(morpholin-4-yl)ethyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0318] The titled compound was obtained as pink powder according to the procedure as described in Example 136, using 4-(2-aminoethyl)morpholine instead of 2-methoxyethylamine.

MS(ESI+) m/z 601 (M+H)⁺, Rt= 1.53 minutes (method A)

Example 147

N-(3-Chloro-2-methylphenyl)-2-{[2-(dimethylamino)ethyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0319] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using N,N-dimethylethylenediamine instead of 2-methoxyethylamine.

MS(ESI+) m/z 559 (M+H)⁺, Rt= 1.51 minutes (method A)

Example 148

N-(3-Chloro-2-methylphenyl)-2-(3-hydroxyazetidin-1-yl)-6-({ [2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0320] The titled compound was obtained as white powder according to the procedure as described in Example 136, using 3-hydroxyazetidine hydrochloride, N,N-diisopropylethylamine and EtOH instead of 2-methoxyethylamine.

MS(ESI+) m/z 544 (M+H)⁺, Rt= 1.77 minutes (method A)

EP 2 746 265 B1

Example 149

N-(3-Chloro-2-methylphenyl)-2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide dihydrochloride

[0321] The titled compound was obtained as brown powder according to the procedure as described in Example 136, using (S)-(-)-3-dimethylaminopyrrolidine, N,N-diisopropylethylamine and EtOH instead of 2-methoxyethylamine. MS(ESI+) m/z 585 (M+H)⁺, Rt= 1.53 minutes (method A)

Example 150

N-(3-Chloro-2-methylphenyl)-2-[(3S)-3-hydroxypyrrolidin-1-yl]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0322] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using (S)-3-hydroxypyrrolidine, N,N-diisopropylethylamine and EtOH instead of 2-methoxyethylamine. MS(ESI+) m/z 558 (M+H)⁺, Rt= 1.75 minutes (method A)

Example 151

N-(3-Chloro-2-methylphenyl)-2-([2-(diethylamino)ethyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0323] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using N,N-diethylethylenediamine instead of 2-methoxyethylamine. MS(ESI+) m/z 587 (M+H)⁺, Rt= 1.57 minutes (method A)

Example 152

N-(3-Chloro-2-methylphenyl)-2-([2-(pyrrolidin-1-yl)ethyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0324] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 1-(2-aminoethyl)pyrrolidine instead of 2-methoxyethylamine. MS(ESI+) m/z 585 (M+H)⁺, Rt= 1.59 minutes (method A)

Example 153

N-(3-Chloro-2-methylphenyl)-2-([3-(dimethylamino)propyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0325] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using N,N-dimethyl-1,3-propanediamine instead of 2-methoxyethylamine. MS(ESI+) m/z 573 (M+H)⁺, Rt= 1.43 minutes (method A)

Example 154

N-(3-Chloro-2-methylphenyl)-2-([3-(dimethylamino)-2,2-dimethylpropyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0326] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using N,N,2,2-tetramethyl-1,3-propanediamine instead of 2-methoxyethylamine. MS(ESI+) m/z 601 (M+H)⁺, Rt= 1.56 minutes (method A)

Example 155

N-(3-Chloro-2-methylphenyl)-2-([2-(dipropan-2-ylamino)ethyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0327] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using N,N-diisopropylethylenediamine instead of 2-methoxyethylamine.

MS(ESI+) m/z 615 (M+H)⁺, Rt= 1.69 minutes (method A)

Example 156

N-(3-Chloro-2-methylphenyl)-2-(morpholin-4-yl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0328] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136 using morpholine instead of 2-methoxyethylamine.

MS(ESI+) m/z 558 (M+H)⁺, Rt= 2.25 minutes (method A)

Example 157

2-Amino-N-(3-chloro-2-methylphenyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0329] According to the procedure as described in Example 136, N-(3-chloro-2-methylphenyl)-2-[(4-methoxybenzyl)amino]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide (97 mg) was obtained as yellow powder, using 4-methoxybenzylamine instead of 2-methoxyethylamine. This was dissolved in TFA (2 mL), and the solution was stirred at 70°C for 2 hours, and removed TFA under reduced pressure. The residue was added with saturated aqueous sodium bicarbonate, and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain 2-amino-N-(3-chloro-2-methylphenyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide (35 mg). This was suspended in MeOH (3 mL), added with 2N hydrogen chloride/MeOH solution (0.3 mL), and removed MeOH under reduced pressure. The residue was triturated in 2-propanol/diethyl ether, and the precipitate was collected by filtration and dried to obtain the titled compound (21 mg) as yellow powder.

MS(ESI+) m/z 488 (M+H)⁺, Rt= 1.62 minutes (method A)

Example 158

N-(3-Chloro-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0330] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 3-amino-2,2-dimethyl-1-propanol instead of 2-methoxyethylamine.

MS(ESI+) m/z 574 (M+H)⁺, Rt= 1.81 minutes (method A)

Example 159

N-(3-Chloro-2-methylphenyl)-2-[(3-methyloxetan-3-yl)methyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0331] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 1-(3-methyloxetan-3-yl)methanamine (prepared as described in US2009/76062) instead of 2-methoxyethylamine.

MS(ESI+) m/z 572 (M+H)⁺, Rt= 1.66 minutes (method A)

Example 160

tert-Butyl

5 N-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-([2-(trifluoro methyl)phenyl]carbonyl)amino}-1H-benzimidazol-2-yl}glycinate hydrochloride

10 **[0332]** The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using glycine tert-butyl ester instead of 2-methoxyethylamine. MS(ESI+) m/z 602 (M+H)⁺, Rt= 2.42 minutes (method A)

Example 161

15 N-{4-[(3-Chloro-2-methylphenyl)carbamoyl]-6-([2-(trifluoro methyl)phenyl]carbonyl)amino}-1H-benzimidazol-2-yl} glycine hydrochloride

20 **[0333]** TFA (1 mL) was added to a solution of tert-butyl N-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-([2-(trifluoro methyl)phenyl]carbonyl)amino}-1H-benzimidazol-2-yl}glycinate (95 mg) in methylene chloride (2 mL), and the mixture was stirred for 1 hour at room temperature, and concentrated. The residue was neutralized with 1N aqueous NaOH under ice-cooling and extracted with EtOAc. The ethyl acetate layer was dried over anhydrous magnesium sulfate and concentrated. The residue was purified on column chromatography to obtain N-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-([2-(trifluoro methyl)phenyl]carbonyl)amino}-1H-benzimidazol-2-yl}glycine (50 mg) as white powder. N-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-([2-(trifluoro methyl)phenyl]carbonyl)amino}-1H-benzimidazol-2-yl}glycine (16 mg) was suspended in MeOH. The suspension was added with 2N hydrogen chloride/EtOH solution (1 Eq) to obtain a homogeneous solution. The reaction mixture was removed the solvent under reduced pressure. The residue was triturated in diethyl ether, and the precipitate was collected by filtration and dried to obtain the titled compound (12 mg) as white powder. MS(ESI+) m/z 546 (M+H)⁺, Rt= 1.74 minutes (method A)

Example 162

30 N-(3-Chloro-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1-methyl-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

35 **[0334]** Sodium hydroxide powder (6.2 mg) was added to a solution of N-(3-chloro-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide (80 mg) in DMF (2 mL), and the solution was stirred at room temperature for 30 minutes, added with methyl iodide (8.7 μ L), and stirred for 17 hours. The reaction mixture was diluted with ethyl acetate, washed with brine, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain N-(3-chloro-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1-methyl-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide (20 mg) as white powder. This was suspended in MeOH, and added with 2N hydrogen chloride/EtOH (1 Eq) and stirred to obtain the solution and removed the solvent under reduced pressure. The residue was washed with diethyl ether, and the precipitate was collected by filtration and dried to obtain the titled compound (15 mg) as white powder. MS(ESI+) m/z 588 (M+H)⁺, Rt= 2.06 minutes (method A)

Example 163

N-(3-Chloro-2-methylphenyl)-2-[(3-methoxy-2,2-dimethylpropyl)amino]-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

50 **[0335]** The titled compound was obtained as white powder according to the procedure as described in Example 136, using 3-methoxy-2,2-dimethyl propylamine (prepared as described in WO2007/28051) instead of 2-methoxyethylamine. MS(ESI+) m/z 588 (M+H)⁺, Rt= 2.11 minutes (method A)
 Elemental Analysis for C₂₉H₂₉ClF₃N₅O₃•HCl+0.5H₂O
 Calcd.(%) C:54.98 H:4.93 N:11.06
 55 Found. (%) C:54.94 H:4.63 N:11.09

Example 164

N-(3-Chloro-2-methylphenyl)-2-(pyrrolidin-1-yl)-6-({[2-(tri fluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carb oxamide hydrochloride

[0336] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using pyrrolidine instead of 2-methoxyethylamine.

MS(ESI+) m/z 542 (M+H)⁺, Rt= 2.15 minutes (method A)

Elemental Analysis for C₂₇H₂₃ClF₃N₅O₂•HCl+1.0H₂O

Calcd.(%) C:54.37 H:4.39 N:11.74

Fond.(%) C:54.34 H:4.66 N:12.26

Example 165

2-(Azetidin-1-yl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carbox amide hydrochloride

[0337] A solution of 2-chloro-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (80 mg, Example 135), azetidine hydrochloride (60 mg) and N, N-diisopropylethylamine (208 μL) in EtOH (1.8 mL) was reacted at 120°C for 20 minutes in a microwaves reactor (Biotage, Initiator). The reaction mixture was concentrated under reduced pressure, and the residue was purified on column chromatography to obtain 2-(azetidin-1-yl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carbox amide (27 mg). This was dissolved in MeOH (1 mL), and added with 2N hydrogen chloride/MeOH solution (1 Eq), and stirred and concentrated. The residue was triturated in diethyl ether, and the precipitate was collected by filtration and dried to obtain the titled compound (21 mg) as white powder.

MS(ESI+) m/z 528 (M+H)⁺, Rt= 2.09 minutes (method A)

Example 166

N-(3-Chloro-2-methylphenyl)-2-(3-methoxyazetidin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole -4-carboxamide hydrochloride

[0338] The titled compound was obtained as white powder according to the procedure as described in Example 136, using 3-methoxy azetidine hydrochloride, N,N-diisopropylethylamine and EtOH instead of 2-methoxyethylamine.

MS(ESI+) m/z 558 (M+H)⁺, Rt= 2.20 minutes (method A)

Example 167

N-(3-Chloro-2-methylphenyl)-2-[(2-hydroxy-2-methylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0339] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 1-amino-2-methyl-2-propanol (prepared as described in WO2009/57827) and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 560 (M+H)⁺, Rt= 1.87 minutes (method A)

Example 168

N-(3-Chloro-2-methylphenyl)-2-[(2S)-tetrahydrofuran-2-ylmethyl]amino-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0340] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using (S)-(+)-tetrahydrofurfurylamine and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 572 (M+H)⁺, Rt= 2.04 minutes (method A)

Example 169

N-(3-Chloro-2-methylphenyl)-2-[[2-(2R)-tetrahydrofuran-2-ylmethyl]amino]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0341] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using (R)-(-)-tetrahydrofurfurylamine and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 572 (M+H)⁺, Rt= 2.04 minutes (method A)

Elemental Analysis for C₂₈H₂₅Cl₃N₅O₃·HCl+0.7H₂O.

Calcd. (%) C:54.15 H:4.45 N:11.28

Found. (%) C:54.16 H:4.57 N:11.19

Example 170

N-(3-Chloro-2-methylphenyl)-2-[[2-(2S)-1-hydroxy-3-methylbutan-2-yl]amino]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0342] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using (S)-(+)-2-amino-3-methyl-1-butanol and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 574 (M+H)⁺, Rt= 1.98 minutes (method A)

Example 171

N-(3-Chloro-2-methylphenyl)-2-[[2-(2R)-1-hydroxy-3-methylbutan-2-yl]amino]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0343] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using (R)-(-)-2-amino-3-methyl-1-butanol and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 574 (M+H)⁺, Rt= 1.98 minutes (method A)

Example 172

N-(3-Chloro-2-methylphenyl)-2-[[2-(2S)-1-hydroxy-3,3-dimethylbutan-2-yl]amino]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0344] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using (S)-tert-leucinol and THF instead of 2-methoxyethylamine. MS(ESI+) m/z 588 (M+H)⁺, Rt= 2.20 minutes (method A)

Example 173

N-(3-Chloro-2-methylphenyl)-2-[[3-methoxy-2,2-dimethylpropyl(methyl)amino]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0345] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 3-methoxy-N,2,2-trimethylpropan-1-amine (Reference Example 9) and THF instead of 2-methoxyethylamine. MS(ESI+) m/z 602 (M+H)⁺, Rt= 2.71 minutes (method A)

Example 174

N-(3-Chloro-2-methylphenyl)-2-[[3-methoxypropyl]amino]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0346] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 3-methoxypropylamine and THF instead of 2-methoxyethylamine. MS(ESI+) m/z 560 (M+H)⁺, Rt= 2.10 minutes (method A)

Example 175

N-(3-Chloro-2-methylphenyl)-2-([2-(propan-2-yloxy)ethyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0347] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 2-aminoethyl isopropyl ether and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 574 (M+H)⁺, Rt= 2.29 minutes (method A)

Example 176

2-([2-tert-Butoxyethyl]amino)-N-(3-chloro-2-methylphenyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0348] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using O-tert-butyl-2-aminoethanol and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 588 (M+H)⁺, Rt= 2.42 minutes (method A)

Example 177

N-(3-Chloro-2-methylphenyl)-2-([2-methoxy-2-methylpropyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0349] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 2-methoxy-2-methylpropylamine and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 574 (M+H)⁺, Rt= 2.30 minutes (method A)

Example 178

N-(3-Chloro-2-methylphenyl)-2-([2-(methylsulfanyl)ethyl]amino)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0350] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 136, using 2-(methylthio)ethylamine and THF instead of 2-methoxyethylamine.

MS(ESI+) m/z 562 (M+H)⁺, Rt= 2.51 minutes (method A)

Example 179

N-(3-Chloro-2-methylphenyl)-2-(methylsulfanyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[Step 1]

Methyl 2-(methylsulfanyl)-1H-benzimidazole-4-carboxylate

[0351] To a solution of methyl 2-sulfanyl-1H-benzimidazole-4-carboxylate (prepared as described in WO2003/106430) (1.0g) in DMF (15 mL), were added potassium carbonate (700 mg) and methyl iodide (0.32 mL), and the mixture was stirred at room temperature for 3 days. The reaction mixture was diluted with ethyl acetate, washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain the titled compound (860 mg) as white powder.

[Step 2]

Methyl 2-(methylsulfanyl)-6-nitro-1H-benzimidazole-4-carboxylate

[0352] The titled compound (840 mg) was obtained as pale yellow powder according to the procedure as described in Step 2 of Example 1, using methyl 2-(methylsulfanyl)-1H-benzimidazole-4-carboxylate (860 mg) instead of methyl-1H-benzimidazole-4-carboxylate.

[Step 3]

Methyl 6-amino-2-(methylsulfanyl)-1H-benzimidazole-4-carboxylate

[0353] The titled compound (380 mg) was obtained as pale yellow powder according to the procedure as described in Step 3 of Example 1, using methyl 2-(methylsulfanyl)-6-nitro-1H-benzimidazole-4-carboxylate (840 mg) instead of methyl 6-nitro-1H-benzimidazole-4-carboxylate.

[Step 4]

Methyl 2-(methylsulfanyl)-6-([2-(trifluoromethyl)phenyl]carbonyl) amino)-1H-benzimidazole-4-carboxylate

[0354] The titled compound (250 mg) was obtained as white powder according to the procedure as described in Step 4 of Example 1, using methyl 6-amino-2-(methylsulfanyl)-1H-benzimidazole-4-carboxylate (150 mg) instead of methyl 6-amino-1H-benzimidazole-4-carboxylate.

[Step 5]

2-(Methylsulfanyl)-6-([2-(trifluoromethyl)phenyl]carbonyl) amino)-1H-benzimidazole-4-carboxylic acid

[0355] The titled compound (240 mg) was obtained as pale yellow powder according to the procedure as described in Step 5 of Example 1, using methyl 2-(methylsulfanyl)-6-([2-(trifluoromethyl)phenyl]carbonyl) amino)-1H-benzimidazole-4-carboxylate (250 mg) instead of methyl 6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylate.

[Step 6]

N-(3-Chloro-2-methylphenyl)-2-(methylsulfanyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0356] The titled compound (35 mg) was obtained as white powder according to the procedure as described in Step 6 of Example 135, using 2-(methylsulfanyl)-6-([2-(trifluoromethyl)phenyl]carbonyl) amino)-1H-benzimidazole-4-carboxylic acid (45 mg) instead of 2-chloro-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxylic acid. MS(ESI+) m/z 519 (M+H)⁺, Rt= 2.79 minutes (method A)

Example 180

N-(3-Chloro-2-methylphenyl)-2-(methylsulfonyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0357] To a solution of N-(3-chloro-2-methylphenyl)-2-(methylsulfanyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide (33 mg) in MeOH (5 mL), was added an aqueous solution (0.5 mL) of Oxone monopersulfate compound (75 mg), and the mixture was stirred at room temperature for 17 hours. Additionally the mixture was added with an aqueous solution (0.5 mL) of Oxone monopersulfate compound (50 mg), and stirred for 3 hours. The reaction mixture was diluted with ethyl acetate and washed with saturated aqueous sodium bicarbonate. The organic layer was separated, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain titled compound (11 mg) as white powder. MS(ESI+) m/z 551 (M+H)⁺, Rt= 2.54 minutes (method A)

Example 181

N-(3-Chloro-2-methylphenyl)-2-(methylsulfinyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide

[0358] To a solution of N-(3-chloro-2-methylphenyl)-2-(methylsulfanyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide (26 mg) in THF (3 mL), was added an aqueous solution of m-chloroperbenzoic acid (75%, 12 mg) (1 mL), and the mixture was stirred at room temperature for 3 hours. The reaction mixture was diluted with ethyl acetate, washed sequentially with saturated aqueous sodium bicarbonate and brine, dried over anhydrous

magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to yield the titled compound (10 mg) as pale yellow powder.

MS(ESI+) m/z 535 (M+H)⁺, Rt= 2.42 minutes (method A)

5 Example 182

6-[[[(2-Chloro-6-fluorophenyl)carbonyl]amino}-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

10 [Step 1]

Methyl 2-(dimethylamino)-1H-benzimidazole-4-carboxylate

15 **[0359]** (Dichloromethylene)dimethyliminium chloride (4.93g) was added to a solution of methyl 2,3-diaminobenzoate (4.2g) in methylene chloride (75 mL), and the solution was stirred on oil bath at 50°C for 1 hour. The reaction mixture was added with saturated aqueous sodium bicarbonate was added, and extracted with ethyl acetate. The organic layer was separated, washed with brine and concentrated under reduced pressure. The residue was purified on column chromatography to obtain the titled compound (4.7g) as a pale yellow solid.

20 [Step 2]

Methyl

2-(dimethylamino)-6-nitro-1H-benzimidazole-4-carboxylate

25

[0360] The titled compound (1.8g) was obtained as orange powder according to the procedure as described in Step 2 of Example 1, using methyl 2-(dimethylamino)-1H-benzimidazole-4-carboxylate (4.57g) instead of methyl 1H-benzimidazole-4-carboxylate.

30 [Step 3]

2-(Dimethylamino)-6-nitro-1H-benzimidazole-4-carboxylic acid

35 **[0361]** A solution of methyl 2-(dimethylamino)-6-nitro-1H-benzimidazole-4-carboxylate (1.5g) and lithium hydroxide hydrate (2.4g) in THF-H₂O (1:1, 100 mL) was stirred at 50°C for 2 hours. THF was removed under reduced pressure, and 1N hydrochloric acid was added to the residue with stirring under ice-cooling to adjust the pH to 5. The precipitate was collected by filtration and dried under reduced pressure to obtain the titled compound (1.38g) as yellow solid.

[Step 4]

40

N-(3-Chloro-2-methylphenyl)-2-(dimethylamino)-6-nitro-1H-benzimidazole-4-carboxamide

45 **[0362]** To a solution of 2-(dimethylamino)-6-nitro-1H-benzimidazole-4-carboxylic acid (520 mg) and HBTU (1.03g) in DMF (6 mL), were added N,N-diisopropylethylamine (942 μ L) and 3-chloro-2-methylaniline (323 μ L), and the mixture was stirred at room temperature for 15 hours. The reaction mixture was diluted with the ethyl acetate-THF (1:1), washed with brine, and the organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was washed with MeOH to obtain the titled compound (650 mg) as a yellow solid.

[Step 5]

50

6-Amino-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide

55 **[0363]** To a suspension of N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-6-nitro-1H-benzimidazole-4-carboxamide (580 mg) in MeOH-THF (1:1, 60 mL), was added 2N hydrogen chloride/EtOH (1 mL). The mixture was added with 1% platinum + 0.1% copper-activated carbon (Degussa type CF105 R/W) (150 mg) and stirred vigorously under hydrogen atmosphere (0.2MPa) for 3 hours. The reaction mixture was filtered off through celite, and the mother mixture was concentrated. The residue was added with saturated aqueous sodium bicarbonate, and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under

reduced pressure. The residue was purified on column chromatography to obtain the titled compound (265 mg) as brown powder.

[Step 6]

6-[[2-chloro-6-fluorophenyl]carbonyl]amino-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0364] To a solution of 6-amino-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide (30 mg) in THF (1 mL), was added N,N-diisopropylethylamine (22 μ L). The mixture was stirred under ice-cooling, and added slowly with 2-chloro-6-fluorobenzoyl chloride (13 μ L), and stirred for 2 hours. The reaction mixture was diluted with ethyl acetate, washed sequentially with saturated aqueous sodium bicarbonate and brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified on column chromatography to obtain 6-[[2-chloro-6-fluorophenyl]carbonyl]amino-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide (30 mg) as pale yellow powder. This was suspended in MeOH (1 mL), and was added with 1N hydrogen chloride/MeOH (1 Eq) to obtain a homogeneous solution, and removed MeOH under reduced pressure. The residue was triturated in n-hexane/ethyl acetate (1:1), and the precipitate was collected by filtration and dried under reduced pressure to obtain the titled compound (23 mg) as pale yellow powder.

MS(ESI+) m/z 500 (M+H)⁺, Rt= 2.13 minutes (method A)

Example 183

N-(3-Chloro-2-methylphenyl)-6-[[2,6-dichlorophenyl]carbonyl]amino-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0365] The titled compound was obtained as white powder according to the procedure as described in Step 6 of Example 182, using 2,6-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 516 (M+H)⁺, Rt= 2.19 minutes (method A)

Example 184

N-(3-Chloro-2-methylphenyl)-6-[[2,4-dichlorophenyl]carbonyl]amino-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0366] The titled compound was obtained as white powder according to the procedure as described in Step 6 of Example 182, using 2,4-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 516 (M+H)⁺, Rt= 2.31 minutes (method A)

Example 185

N-(3-Chloro-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0367] The titled compound was obtained as white powder according to the procedure as described in Step 6 of Example 182, using 2,5-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 516 (M+H)⁺, Rt= 2.31 minutes (method A)

Elemental Analysis for C₂₄H₂₀Cl₃N₅O₂•HCl•0.5H₂O

Calcd.(%) C:51.27 H:3.94 N:12.46

Fond.(%) C:51.10 H:4.02 N:12.45

Example 186

6-[[2-Bromo-6-fluorophenyl]carbonyl]amino-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0368] The titled compound was obtained as gray powder according to the procedure as described in Step 6 of Example 182, using 2-bromo-6-fluorobenzoyl chloride (prepared as described in WO2007/144327) instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 544 (M+H)⁺, Rt= 2.23 minutes (method A)

Example 187

5 6-([(2-Bromo-6-chlorophenyl)carbonyl]amino)-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0369] The titled compound was obtained as light brown powder according to the procedure as described in Step 6 of Example 182, using 2-bromo-6-chlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

10 MS(ESI+) m/z 560 (M+H)⁺, Rt= 2.29 minutes (method A)

Example 188

15 6-([(2-Chloro-5-(cyclopropylethynyl)phenyl)carbonyl]amino)-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0370] To a solution of 2-chloro-5-(cyclopropylethynyl)benzoic acid (33 mg, Reference Example 3), 6-amino-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide (50 mg) and HATU (72 mg) in DMF (1 mL), was added N,N-diisopropylethylamine (66 μ L), and the mixture was stirred at room temperature for 17 hours. The reaction mixture was diluted with ethyl acetate, and washed sequentially with saturated aqueous sodium bicarbonate and brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain 6-([(2-chloro-5-(cyclopropylethynyl)phenyl)carbonyl]amino)-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide (74 mg). This was suspended in MeOH (1 mL), and was added with 2N hydrogen chloride/EtOH solution (1 Eq) to obtain a homogeneous solution, and removed the solvent under reduced pressure. The residue was triturated in ethyl acetate, and the precipitated was collected by filtration and dried under reduced pressure to obtain the titled compound (58 mg) as pale yellow powder.

MS(ESI+) m/z 546 (M+H)⁺, Rt= 2.60 minutes (method A)

Example 189

30 N-(3-Chloro-2-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

35 Methyl

2-chloro-6-([(2,5-dichlorophenyl)carbonyl]amino)-1H-benzimidazole-4-carboxylate

40 **[0371]** The titled compound (1.68g) was obtained as colorless amorphous according to the procedure as described in Step 4 of Example 1, using methyl 6-amino-2-chloro-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate, and 2,5-dichlorobenzoyl chloride instead of 2-(trifluoromethyl)benzoyl chloride.

[Step 2]

45 2-Chloro-6-([(2,5-dichlorophenyl)carbonyl]amino)-1H-benzimidazole-4-carboxylic acid

[0372] The titled compound (1.41g) was obtained as white powder according to the procedure as described in Step 5 of Example 1, using methyl 2-chloro-6-([(2,5-dichlorophenyl)carbonyl]amino)-1H-benzimidazole-4-carboxylate (1.68g) instead of methyl 6-([(2-(trifluoromethyl)phenyl)carbonyl]amino)-1H-benzimidazole-4-carboxylate.

[Step 3]

55 2-Chloro-N-(3-chloro-2-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-1H-benzimidazole-4-carboxamide

[0373] The titled compound was obtained as a white solid according to the procedure as described in Step 6 of Example 135, using 2-chloro-6-([(2,5-dichlorophenyl)carbonyl]amino)-1H-benzimidazole-4-carboxylic acid instead of 2-chloro-6-([(2-(trifluoromethyl)phenyl)carbonyl]amino)-1H-benzimidazole-4-carboxylic acid.

[Step 4]

N-(3-Chloro-2-methylphenyl)-6-[[{(2,5-dichlorophenyl)carbonyl}amino]-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide hydrochloride

[0374] THF (1 mL) was added to 2-chloro-N-(3-chloro-2-methylphenyl)-6-[[{(2,5-dichlorophenyl)carbonyl}amino]-1H-benzimidazole-4-carboxamide (50 mg) and 3-amino-2,2-dimethyl-1-propanol (400 mg). The mixture was stirred at 150°C in a sealed pressure-proof stainless steel container for 8 hours. The reaction mixture was diluted with ethyl acetate, washed with brine and concentrated under reduced pressure. The residue was purified on column chromatography to obtain N-(3-chloro-2-methylphenyl)-6-[[{(2,5-dichlorophenyl)carbonyl}amino]-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide (50 mg) as pale yellow powder. This was dissolved in MeOH (1 mL), and added with 2N Hydrogen chloride/EtOH solution (1 Eq), and stirred and concentrated. The residue was triturated in diethyl ether, and the precipitate was collected by filtration and dried to obtain the titled compound (40 mg) as white powder.

MS(ESI+) m/z 574 (M+H)⁺, Rt= 2.06 minutes (method A)

Example 190

N-(3-Chloro-2-methylphenyl)-6-[[{(2,5-dichlorophenyl)carbonyl}amino]-2-[(3-methoxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide hydrochloride

[0375] The titled compound was obtained as white powder according to the procedure as described in Example 189, using 3-methoxy-2,2-dimethylpropylamine instead of 3-amino-2,2-dimethyl-1-propanol.

MS(ESI+) m/z 588 (M+H)⁺, Rt= 2.41 minutes (method A)

Example 191

N-(3-Chloro-2-methylphenyl)-6-[[{(2,5-dichlorophenyl)carbonyl}amino]-2-[(2-hydroxy-2-methylpropyl)amino]-1H-benzimidazole-4-carboxamide hydrochloride

[0376] The titled compound was obtained as white powder according to the procedure as described in Example 189, using 2-hydroxy-2-methylpropylamine instead of 3-amino-2,2-dimethyl-1-propanol.

MS(ESI+) m/z 560 (M+H)⁺, Rt= 2.12 minutes (method A)

Example 192

N-(3-Chloro-2-methylphenyl)-6-[[{(2,5-dichlorophenyl)carbonyl}amino]-2-[(2-methoxy-2-methylpropyl)amino]-1H-benzimidazole-4-carboxamide hydrochloride

[0377] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 189, using 2-methoxy-2-methylpropylamine instead of 3-amino-2,2-dimethyl-1-propanol.

MS(ESI+) m/z 574 (M+H)⁺, Rt= 2.48 minutes (method A)

Example 193

N-(3-Chloro-2-methylphenyl)-6-[[{(2,5-dichlorophenyl)carbonyl}amino]-2-[[2-(propan-2-yloxy)ethyl]amino]-1H-benzimidazole-4-carboxamide hydrochloride

[0378] The titled compound was obtained as milky white powder according to the procedure as described in Example 189, using 2-aminoethyl isopropyl ether instead of 3-amino-2,2-dimethyl-1-propanol.

MS(ESI+) m/z 574 (M+H)⁺, Rt= 2.60 minutes (method A)

Elemental Analysis for C₂₇H₂₆Cl₃N₅O₃•HCl•0.1H₂O

Calcd.(%) C:52.89 H:4.47 N:11.42

Fond.(%) C:52.55 H:4.10 N:11.36

Example 194

6-[[[(2-Chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-[[2-(propan-2-yloxy)ethyl]amino]-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

2-Chloro-6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-1H-benzimidazole-4-carboxamide

[0379] The titled compound was obtained as a white solid according to the procedures as described in Steps 1 to 3 of Example 189, using 2-chloro-6-fluorobenzoyl chloride instead of 2,5-dichlorobenzoyl chloride in Step 1.

[Step 2]

6-[[[(2-Chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-[[2-(propan-2-yloxy)-ethyl]amino]-1H-benzimidazole-4-carboxamide hydrochloride

[0380] THF (1 mL) was added to 2-chloro-6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-1H-benzimidazole-4-carboxamide (50 mg) and 2-aminoethyl isopropyl ether (32 mg). The mixture was stirred at 150°C in a sealed pressure-proof stainless steel container for 17 hours. The reaction mixture was diluted with ethyl acetate and washed with brine and concentrated under reduced pressure. The residue was purified on column chromatography to obtain 6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-[[2-(propan-2-yloxy)ethyl]amino]-1H-benzimidazole-4-carboxamide (41 mg) as pale yellow powder. This was dissolved in MeOH (1 mL), and added with 2N hydrogen chloride/EtOH solution (1 Eq), and stirred and concentrated. The residue was triturated in diethyl ether, and the precipitate was collected by filtration and dried to obtain the titled compound (35 mg) as white powder. MS(ESI+) m/z 558 (M+H)⁺, Rt= 2.41 minutes (method A)

Example 195

2-[[2-(tert-Butoxyethyl)amino]-6-[[[(2-chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0381] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 194, using O-tert-butyl 2-aminoethanol instead of 2-aminoethyl isopropyl ether.

MS(ESI+) m/z 572 (M+H)⁺, Rt= 2.49 minutes (method A)

Example 196

6-[[[(2-Chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-[(3-methoxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide hydrochloride

[0382] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 194, using 3-methoxy-2,2-dimethylpropylamine instead of 2-aminoethyl isopropyl ether.

MS(ESI+) m/z 572 (M+H)⁺, Rt= 2.51 minutes (method A)

Example 197

6-[[[(2-Chloro-6-fluorophenyl)carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-[(2-methoxy-2-methylpropyl)amino]-1H-benzimidazole-4-carboxamide hydrochloride

[0383] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 194, using 2-methoxy-2-methylpropylamine instead of 2-aminoethyl isopropyl ether.

MS(ESI+) m/z 558 (M+H)⁺, Rt= 2.34 minutes (method A)

Example 198

6-(((2-Chloro-6-fluorophenyl)-carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(((2S)-tetrahydrofuran-2-ylmethyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0384] The titled compound was obtained as pale pink powder according to the procedure as described in Step 2 of Example 194, using (S)-(+)-tetrahydrofurfurylamine instead of 2-aminoethyl isopropyl ether.

MS(ESI+) m/z 556 (M+H)⁺, Rt= 2.28 minutes (method A)

Example 199

6-(((2-Chloro-6-fluorophenyl)-carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(((2R)-tetrahydrofuran-2-ylmethyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0385] The titled compound was obtained as pale pink powder according to the procedure as described in Step 2 of Example 194, using (R)-(-)-tetrahydrofurfurylamine instead of 2-aminoethyl isopropyl ether.

MS(ESI+) m/z 556 (M+H)⁺, Rt= 2.28 minutes (method A)

Example 200

6-(((2-Chloro-6-fluorophenyl)carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-((3-hydroxy-2,2-dimethylpropyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0386] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 2 of Example 194, using 3-amino-2,2-dimethyl-1-propanol instead of 2-aminoethyl isopropyl ether.

MS(ESI+) m/z 558 (M+H)⁺, Rt= 2.22 minutes (method A)

Example 201

6-(((2-Chloro-6-fluorophenyl)-carbonyl)amino)-N-(3-chloro-2-methylphenyl)-2-(((2S)-1-hydroxy-3-methylbutan-2-yl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0387] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 2 of Example 194, using (S)-(+)-2-amino-3-methyl-1-butanol instead of 2-aminoethyl isopropyl ether.

MS(ESI+) m/z 558 (M+H)⁺, Rt= 2.31 minutes (method A)

Example 202

N-(3-Chloro-4-methylphenyl)-2-(dimethylamino)-6-((2-(trifluoromethyl)phenyl)carbonyl)amino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

2-(dimethylamino)-6-((2-(trifluoromethyl)phenyl)carbonyl)amino)-1H-benzimidazole-4-carboxylic acid

[0388] To a stirring solution of methyl 6-amino-2-chloro-1H-benzimidazole-4-carboxylate (400 mg, Example 135, Step 3) and triethylamine (209 μ L) in THF (18 mL), 2-(trifluoromethyl)benzoyl chloride (316 μ L) was dropped slowly under ice-cooling. After 1 hour, the reaction mixture was diluted with ethyl acetate, washed sequentially with water and brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to obtain crude methyl 2-chloro-6-((2-(trifluoromethyl)phenyl)carbonyl)amino)-1H-benzimidazole-4-carboxylate. This was dissolved in THF (15 mL) and MeOH (7 mL), and added with 1N aqueous lithium hydroxide solution (9mL), and stirred at room temperature overnight. The reaction mixture was removing THF and MeOH under reduced pressure. The residue was added with water, and neutralized with 1N hydrochloric acid under ice-cooling. The precipitation was collected by filtration to obtain crude 2-chloro-6-((2-(trifluoromethyl)phenyl)carbonyl)amino)-1H-benzimidazole-4-carboxylic acid. This was dissolved in THF (10 mL), and added with 40% aqueous dimethylamine solution (7 mL), and stirred at 120°C for 14 hours. The reaction mixture was cooled to room temperature, removed THF under reduced pressure. The residue was added with water, and adjusted to PH 3-4 with 1N hydrochloric acid under ice-cooling. The precipitate was collected by filtration, washed with diethyl ether, and dried under reduced pressure to obtain the titled compound (589 mg) as slightly yellow solid.

[Step 2]

N-(3-Chloro-4-methylphenyl)-2-(dimethylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0389] To a solution of 2-(dimethylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid (50 mg) and HATU (82 mg) in DMF (1 mL), were added N,N-diisopropylethylamine (88 μ L) and 3-chloro-4-methylaniline (26 μ L), and the mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted with ethyl acetate, washed sequentially with saturated aqueous sodium bicarbonate and brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column chromatography to obtain N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (65 mg) as white powder. This was suspended in MeOH (1.5 mL), and added with 2N hydrogen chloride/MeOH (1 Eq), stirred and concentrated. The residue was triturated in diethyl ether, and the precipitate was collected by filtration and dried to obtain the titled compound (40 mg) as white powder.

MS(ESI+) m/z 516 (M+H)⁺, Rt= 2.20 minutes (method A)

Example 203

N-(4-tert-Buthylphenyl)-2-(dimethylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0390] The titled compound was obtained as pale yellow powder according to the procedure as described in Step 2 of Example 202, using 4-tert-butylaniline instead of 3-chloro-4-methylaniline.

MS(ESI+) m/z 524 (M+H)⁺, Rt= 2.30 minutes (method A)

Example 204

N-(2,3-Dihydro-1H-inden-5-yl)-2-(dimethylamino)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0391] The titled compound was obtained as gray powder according to the procedure as described in Step 2 of Example 202, using 5-aminoindan instead of 3-chloro-4-methylaniline.

MS(ESI+) m/z 508 (M+H)⁺, Rt= 2.11 minutes (method A)

Example 205

6-({[2-Chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

6-Amino-N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide

[0392] N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-6-nitro-1H-benzimidazole-4-carboxamide (520 mg) was obtained as yellow powder according to the procedure as described in Step 4 of Example 182, using 3-chloro-4-methylaniline instead of 3-chloro-2-methylaniline. This was then reacted according to the procedure as described in Step 5 of Example 182, to obtain the titled compound (358 mg) as yellow powder.

[Step 2]

6-({[2-Chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0393] To a solution of 6-amino-N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide (50 mg) in THF (1.5 mL), was added N,N-diisopropylethylamine (37 μ L) and the mixture was stirred under ice-cooling. The mixture was added slowly with 2-chloro-6-fluorobenzoyl chloride (24 μ L), and stirred for 2 hours. The reaction mixture was diluted with ethyl acetate, washed sequentially with saturated aqueous sodium bicarbonate and brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified on column

chromatography to obtain 6-[[2-chloro-6-fluorophenyl]carbonyl]amino-N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide (30 mg) as pale yellow powder. This was then suspended in MeOH (1 mL), and added with 2N hydrogen chloride/MeOH (1 Eq) to obtain a homogeneous solution, and MeOH was removed under reduced pressure. The residue was triturated in n-hexane/ethyl acetate (1:1), and the precipitate was collected by filtration and dried to obtain the titled compound (24 mg) as pale purple powder.

MS(ESI+) m/z 500 (M+H)⁺, Rt= 2.20 minutes (method A)

Example 206

N-(3-Chloro-4-methylphenyl)-6-[[2,6-dichlorophenyl]carbonyl]amino-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0394] The titled compound was obtained as pale purple powder according to the procedure as described in Step 2 of Example 205, using 2,6-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 516 (M+H)⁺, Rt= 2.26 minutes (method A)

Example 207

N-(3-Chloro-4-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino-2-(dimethylamino)-1H-benzimidazole-4-carboxamide hydrochloride

[0395] The titled compound was obtained as white powder according to the procedure as described in Step 2 of Example 205, using 2,5-dichlorobenzoyl chloride instead of 2-chloro-6-fluorobenzoyl chloride.

MS(ESI+) m/z 516 (M+H)⁺, Rt= 2.39 minutes (method A)

Elemental Analysis for C₂₄H₂₀Cl₃N₅O₂•HCl

Calcd.(%) C:52.10 H:3.83 N:12.66

Fond.(%) C:51.96 H:3.61 N:12.76

Example 208

N-(3-Chloro-2-methylphenyl)-2-cyclopropyl-6-[[2,5-dichlorophenyl]carbonyl]amino-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 6-amino-2-cyclopropyl-1H-benzimidazole-4-carboxylate

[0396] The title compound was obtained as white powder according to the procedures as described in step 1 to step 4 of Example 11, using cyclopropanecarbonyl chloride instead of methoxyacetyl chloride in Step 1.

[Step 2]

2-Cyclopropyl-6-[[2,5-dichlorophenyl]carbonyl]amino-1H-benzimidazole-4-carboxylic acid

[0397] The title compound was obtained according to the procedures as described in step 4 and step 5 of Example 11, using methyl 6-amino-2-cyclopropyl-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate, and 2,5-dichlorobenzoyl chloride instead of 2-(trifluoromethyl)benzoyl chloride.

[Step 3]

N-(3-Chloro-2-methylphenyl)-2-cyclopropyl-6-[[2,5-dichlorophenyl]carbonyl]amino-1H-benzimidazole-4-carboxamide hydrochloride

[0398] The title compound was obtained according to the procedures as described in step 7 and step 8 of Example 11, using 2-cyclopropyl-6-[[2,5-dichlorophenyl]carbonyl]amino-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino-1H-benzimidazole-4-carboxylic acid.

MS(ESI+) m/z 513 (M+H)⁺, Rt= 1.88 minutes (method A)

Example 209

N-(3-Chloro-4-methylphenyl)-2-cyclopropyl-6-[[2,5-dichlorophenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0399] The titled compound was obtained as white powder according to the procedures as described in step 7 and step 8 of Example 11, using 2-cyclopropyl-6-[[2,5-dichlorophenyl]carbonyl]amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxylic acid, and 3-chloro-4-methylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 513 (M+H)⁺, Rt= 1.85 minutes (method A)

Example 210

N-(3-Chloro-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]aminol-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 6-amino-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxylate

[0400] The title compound was obtained as white powder according to the procedures as described in step 1 to step 4 of Example 11, using 1-methyl-cyclopropanecarbonyl chloride (prepared as described in WO2009/68512) instead of methoxyacetyl chloride in Step 1.

[Step 2]

2-(1-Methylcyclopropyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-1H-benzimidazole-4-carboxylic acid

[0401] The title compound was obtained according to the procedures as described in step 4 and step 5 of Example 1, using methyl 6-amino-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate, and 2,5-dichlorobenzoyl chloride instead of 2-(trifluoromethyl)benzoyl chloride.

[Step 3]

N-(3-Chloro-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0402] The title compound was obtained according to the procedures as described in step 7 and step 8 of Example 11, using 2-(1-methylcyclopropyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxylic acid.

MS(ESI+) m/z 527 (M+H)⁺, Rt= 2.12 minutes (method A)

Example 211

N-(3-Chloro-4-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0403] The titled compound was obtained as white powder according to the procedures as described in step 7 and step 8 of Example 11, using 2-(1-methylcyclopropyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-1H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxylic acid, and 3-chloro-4-methylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 527 (M+H)⁺, Rt= 2.18 minutes (method A)

Example 212

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(methylsulfonyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0404] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 103, using 2-(methylsulfonyl)benzoic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 527 (M+H)⁺, Rt= 2.08 minutes (method A)

Example 213

N-(3-Chloro-4-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(2-methoxyethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[Step 1]

Methyl 6-amino-2-(2-methoxyethyl)-1H-benzimidazole-4-carboxylate

[0405] The title compound was obtained as white powder according to the procedures as described in step 1 to step 4 of Example 11, using methoxypropionyl chloride instead of methoxyacetyl chloride in Step 1.

[Step 2]

2-(2-Methoxyethyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-1 H-benzimidazole-4-carboxylic acid

[0406] The title compound was obtained according to the procedures as described in step 4 and step 5 of Example 1, using methyl 6-amino-2-(2-methoxyethyl)-1H-benzimidazole-4-carboxylate instead of methyl 6-amino-1H-benzimidazole-4-carboxylate, and 2,5-dichlorobenzoyl chloride instead of 2-(trifluoromethyl)benzoyl chloride.

[Step 3]

N-(3-Chloro-4-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(2-methoxyethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0407] The title compound was obtained according to the procedures as described in step 7 and step 8 of Example 11, using 2-(2-methoxyethyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-1 H-benzimidazole-4-carboxylic acid instead of 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxylic acid.

MS(ESI+) m/z 531 (M+H)⁺, Rt= 2.43 minutes (method A)

Example 214

2-(Methoxymethyl)-N-phenyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0408] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using aniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 469 (M+H)⁺, Rt= 2.01 minutes (method A)

Example 215

2-(Methoxymethyl)-N-propyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0409] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using n-propylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 435 (M+H)⁺, Rt= 1.61 minutes (method A)

Example 216

2-(Methoxymethyl)-N-(pyridin-3-yl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0410] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 3-aminopyridine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 470 (M+H)⁺, Rt= 1.25 minutes (method A)

Example 217

N-Benzyl-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0411] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using benzylamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 483 (M+H)⁺, Rt= 1.83 minutes (method A)

Example 218

N-(Cyclohexylmethyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0412] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using cyclohexanemethylamine instead of 3-chloro-2-methylaniline. MS(ESI+) m/z 489 (M+H)⁺, Rt= 2.11 minutes (method A)

Example 219

2-(Methoxymethyl)-N-(naphthalen-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0413] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 1-naphthylamine instead of 3-chloro-2-methylaniline. MS(ESI+) m/z 519 (M+H)⁺, Rt= 2.29 minutes (method A)

Example 220

2-(Methoxymethyl)-N-(thiophen-3-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0414] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 3-aminothiophene instead of 3-chloro-2-methylaniline. MS(ESI+) m/z 475 (M+H)⁺, Rt= 1.92 minutes (method A)

Example 221

N-(2,1,3-Benzothiadiazol-4-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0415] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 4-amino-2,1,3-benzothiadiazole instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 527 (M+H)⁺, Rt= 2.16 minutes (method A)

Example 222

N-(1,1-Dioxide-1-benzothiophen-6-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0416] The titled compound was obtained as colorless powder according to the procedure as described in Example

EP 2 746 265 B1

11, using 6-aminobenzo[b]thiophene 1,1-dioxide instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 557 (M+H)⁺, Rt= 1.82 minutes (method A)

Example 223

2-(Methoxymethyl)-N-(thiophen-2-ylmethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0417] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-thiophene methylamine instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 489 (M+H)⁺, Rt= 1.77 minutes (method A)

Example 224

N-(1H-Indol-5-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0418] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 5-aminoindole instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 508 (M+H)⁺, Rt= 1.75 minutes (method A)

Example 225

N-(1,3-Benzothiazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0419] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-aminobenzothiazole instead of 3-chloro-2-methylaniline. MS(ESI+) m/z 526 (M+H)⁺, Rt= 2.20 minutes (method A)

Example 226

N-(2,2-Dimethylpropyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0420] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using neopentyl amine instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 463 (M+H)⁺, Rt= 1.95 minutes (method A)

Example 227

2-(Methoxymethyl)-N-(thiophen-2-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0421] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-aminothiophene instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 475 (M+H)⁺, Rt= 1.94 minutes (method A)

Example 228

N-(5-Chloro-1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0422] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-amino-5-chlorobenzoxazole instead of 3-chloro-2-methylaniline.
MS(ESI+) m/z 544 (M+H)⁺, Rt= 2.09 minutes (method A)

Example 229

N-(2-Benzylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0423] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-benzylaniline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 559 (M+H)⁺, Rt= 2.34 minutes (method A)

Example 230

2-(Methoxymethyl)-N-(quinolin-8-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0424] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 11, using 8-aminoquinoline instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 520 (M+H)⁺, Rt= 2.03 minutes (method A)

Example 231

N-(Cycloheptylmethyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0425] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 11, using cycloheptanemethyleneamine instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 503 (M+H)⁺, Rt= 2.23 minutes (method A)

Example 232

N-(1,3-Benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0426] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-aminobenzoxazole instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 510 (M+H)⁺, Rt= 1.80 minutes (method A)

Example 233

N-(6-Chloro-1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide hydrochloride

[0427] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using 2-amino-6-chlorobenzoxazole instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 544 (M+H)⁺, Rt= 2.09 minutes (method A)

Example 234

N-3-Chloro-2-(hydroxymethyl)phenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide

[0428] The titled compound was obtained as colorless powder according to the procedure as described in Example 11, using (2-amino-6-chlorophenyl)methanol (prepared as described in J. Med. Chem., 2005, 48, 2080) instead of 3-chloro-2-methylaniline.

MS(ESI+) m/z 533 (M+H)⁺, Rt= 2.08 minutes (method A)

Example 235

N-(3-Chloro-4-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide methanesulfonate

[0429] To a suspension of N-(3-chloro-4-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide (200 mg) (obtained in Example 207) in MeOH (2 ml), was added methanesulfonic acid (1 Eq). The mixture was stirred at room temperature for 30 minutes, and filtered off through glass filter to collect crystals, which were dried under reduced pressure to obtain the titled compound (180 mg) as white powder.

Elemental Analysis for $C_{24}H_{20}Cl_3N_5O_2 \cdot CH_3SO_3H + 1.5H_2O$

Calcd.(%) C:46.92 H:4.25 N:10.94

Fond.(%) C:46.85 H:4.04 N:10.89

Example 236

N-(3-Chloro-4-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide 4-methylbenzenesulfonate

[0430] The titled compound was obtained as white powder according to the procedure as described in Example 235, using p-toluenesulfonic acid monohydrate (1 Eq) instead of methanesulfonic acid.

Elemental Analysis for $C_{24}H_{20}Cl_3N_5O_2 \cdot C_7H_8O_3S + 1.2H_2O$

Calcd.(%) C:52.40 H:4.31 N:9.86

Found.(%) C:52.24 H:4.28 N:9.95

Example 237

N-(3-Chloro-4-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide sulfate

[0431] The titled compound was obtained as white powder according to the procedure as described in Example 235, using sulfuric acid (1 Eq) instead of methanesulfonic acid.

Elemental Analysis for $C_{24}H_{20}Cl_3N_5O_2 \cdot H_2SO_4 + 1.5H_2O$

Calcd.(%) C:44.91 H:3.93 N:10.91

Found.(%) C:44.71 H:3.81 N:10.85

Example 238

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide methanesulfonate

[0432] Methanesulfonic acid (1 Eq) was added to a suspension of N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide (250 mg) (obtained in Example 11) in MeOH (2 ml). After stirring for 30 minutes at room temperature, MeOH was removed under reduced pressure to obtain the titled compound as white powder.

Elemental Analysis for $C_{25}H_{20}ClF_3N_4O_3 \cdot CH_3SO_3H + 1.0H_2O$

Calcd.(%) C:49.49 H:4.15 N:8.88

Fond.(%) C:49.20 H:4.14 N:8.78

Example 239

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-6-([2-(trifluoromethyl)phenyl]carbonyl)amino)-1H-benzimidazole-4-carboxamide 4-methylbenzenesulfonate

[0433] The titled compound was obtained as white powder according to the procedure as described in Example 238, using p-toluenesulfonic acid monohydrate (1 Eq) instead of methanesulfonic acid.

Elemental Analysis for $C_{25}H_{20}ClF_3N_4O_3 \cdot C_7H_8O_3S$

Calcd.(%) C:55.78 H:4.10 N:8.13

Fond.(%) C:55.62 H:4.12 N:8.16

Example 240

N-(3-Chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide sulfate

[0434] The titled compound was obtained as white powder according to the procedure as described in Example 238, using sulfuric acid (1 Eq) instead of methanesulfonic acid.

Elemental Analysis for $C_{25}H_{20}ClF_3N_4O_3 \cdot H_2SO_4 + 0.5H_2O$

Calcd. (%) C:48.12 H:3.72 N:8.98

Found. (%) C:48.34 H:3.57 N:8.99

Example 241

N-(3-Chloro-2-methylphenyl)-2-(1-methylcyclopropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide methanesulfonate

[0435] Methanesulfonic acid (1 Eq) was added to a suspension of N-(3-chloro-2-methylphenyl)-2-(1-methylcyclopropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide (250 mg) (obtained in Example 85) in MeOH (2 ml). After stirring at room temperature for 30 minutes, MeOH was removed under reduced pressure to obtain the titled compound as white powder.

Elemental Analysis for $C_{27}H_{22}ClF_3N_4O_2 \cdot CH_3SO_3H + 0.5H_2O$

Calcd. (%) C:53.21 H:4.31 N:8.86

Fond. (%) C:52.97 H:4.14 N:8.90

Example 242

N-(3-Chloro-2-methylphenyl)-2-(1-methylcyclopropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide 4-methylbenzenesulfonate

[0436] The titled compound was obtained as white powder according to the procedure as described in Example 241, using p-toluenesulfonic acid monohydrate (1 Eq) instead of methanesulfonic acid.

Elemental Analysis for $C_{27}H_{22}ClF_3N_4O_2 \cdot C_7H_8O_3S$

Calcd. (%) C:58.41 H:4.32 N:8.01

Fond. (%) C:58.13 H:4.50 N:8.00

Example 243

N-(3-Chloro-2-methylphenyl)-2-(1-methylcyclopropyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide sulfate

[0437] The titled compound was obtained as white powder according to the procedure as described in Example 241, using sulfuric acid (1 Eq) instead of methanesulfonic acid.

Elemental Analysis for $C_{27}H_{22}ClF_3N_4O_2 \cdot H_2SO_4 + 0.6H_2O$

Calcd. (%) C:51.00 H:3.99 N:8.81

Fond. (%) C:51.35 H:4.39 N:8.85

Example 244

N-(3-Chloro-2-methylphenyl)-6-({[3-fluoropyridin-2-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0438] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 3-fluoropyridine-2-carboxylic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 468 (M+H)⁺, Rt= 2.17 minutes (method A)

EP 2 746 265 B1

Example 245

N-(3-Chloro-2-methylphenyl)-6-[[3-chloropyridin-4-yl]carbonylamino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0439] The titled compound was obtained as pale yellow powder according to the procedure as described in Example 103, using 3-chloropicolinic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 484 (M+H)⁺, Rt= 2.08 minutes (method A)

Example 246

N-(3-Chloro-2-methylphenyl)-6-[[3,5-dichloropyridin-4-yl]carbonylamino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0440] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 3,5-dichloroisonicotinic acid instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 518 (M+H)⁺, Rt= 2.24 minutes (method A)

Example 247

6-[[5-Butoxy-2-chlorophenyl]carbonylamino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0441] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 5-butoxy-2-chlorobenzoic acid (prepared in Reference Example 10) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 555 (M+H)⁺, Rt= 2.67 minutes (method A)

Example 248

6-[[2-Chloro-5-(2,2-difluoroethoxy)phenyl]carbonylamino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0442] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-5-(2,2-difluoroethoxy)benzoic acid (prepared in Reference Example 11) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 563 (M+H)⁺, Rt= 1.66 minutes (method A)

Example 249

N-(3-Chloro-2-methylphenyl)-6-[[2-chloro-5-(4,4,4-trifluorobutoxy)phenyl]carbonylamino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide hydrochloride

[0443] The titled compound was obtained as white powder according to the procedure as described in Example 103, using 2-chloro-5-(4,4,4-trifluorobutoxy)benzoic acid (prepared in Reference Example 12) instead of 2-bromo-6-fluorobenzoic acid.

MS(ESI+) m/z 609 (M+H)⁺, Rt= 1.99 minutes (method A)

[0444] The chemical structures of the compounds of Examples 1 to 249 are shown in the following Tables 1 to 14.

EP 2 746 265 B1

[Table 1]

Example	Structure	Example	Structure	Example	Structure
1		7		13	
2		8		14	
3		9		15	
		10		16	
5		11		17	
6		12		18	

[Table 2]

Example	Structure	Example	Structure	Example	Structure
19		25		31	
20		26		32	
21		27		33	
22		28		34	
23		29		35	
24		30		36	

[Table 3]

Example	Structure	Example	Structure	Example	Structure
37		43		49	
38		44		50	
39		45		51	
40		46		52	
41		47		53	
42		48		54	

[Table 4]

Example	Structure	Example	Structure	Example	Structure
55		61		67	
56		62		68	
57		63		69	
58		64		70	
59		65		71	
60		66		72	



















[Table 5]

Example	Structure	Example	Structure	Example	Structure
73		79		85	
74		80		86	
75		81		87	
76		82		88	
77		83		89	
78		84		90	



















[Table 6]

Example	Structure	Example	Structure	Example	Structure
91		97		103	
92		98		104	
93		99		105	
94		100		106	
95		101		107	
96		102		108	













[Table 7]

Example	Structure	Example	Structure	Example	Structure
109		115		121	
110		116		122	
111		117		123	
112		118		124	
113		119		125	
114		120		126	








[Table 8]

Example	Structure	Example	Structure	Example	Structure
127		133		139	
128		134		140	
129		135		141	
130		136		142	
131		137		143	
132		138		144	



















[Table 9]

Example	Structure	Example	Structure	Example	Structure
145		151		157	
146		152		158	
147		153		159	
148		154		160	

(continued)

Structure	Structure	Exam ple	Structure	Exam ple	Structure	Exam ple
		149		162		150
		161		156		



















[Table 10]

Example	Structure	Example	Structure	Example	Structure
163		169		175	
164		170		176	
165		171		177	
166		172		178	
167		173		179	
168		174		180	



















[Table 11]

Example	Structure	Example	Structure	Example	Structure
181		187		193	
182		188		194	
183		189		195	
184		190		196	
185		191		197	
186		192		198	
















[Table 12]

Example	Structure	Example	Structure	Example	Structure
199		205		211	
200		206		212	
201		207		213	
202		208		214	
203		209		215	
204		210		216	

[Table 13]

Example	Structure	Example	Structure	Example	Structure
217		223		229	
218		224		230	
219		225		231	
220		226		232	
221		227		233	
222		228		234	

[Table 14]

Example	Structure	Example	Structure	Example	Structure
235		241		247	
236		242		248	
237		243		249	
238		244			
239		245			
240		246			

Test Example 1: mPGES-1 inhibiting activity

[0445] mPGES-1 microsome fractions were prepared from CHO-K1 cells transiently transfected with plasmid encoding the human mPGES-1cDNA. Microsomes were diluted with potassium phosphate buffer containing reduced glutathione (pH7.4), and DMSO containing test compound or DMSO alone, which was each 1% of DMSO final concentration, was

EP 2 746 265 B1

added and incubated at 4°C for 20 minutes. Then, the enzymatic reactions were initiated by the addition of PGH2 substrate (final concentration 1 μ M) and incubated at 4°C for 60 seconds. The reaction was terminated by addition of the citrate solution (final citrate concentration 50mM) containing ferric chloride (final concentration 1 mg/mL). PGE2 production in the enzyme reaction aliquot was measured using HTRF kit (Cisbio International, catalogue #62P2APEC). The solution free of test compound was used as positive control, and the solution free of test compound and microsome sample was used as negative control. 100% activity was defined as PGE2 production in the positive control minus PGE2 production in negative control. IC50 value was calculated by standard method.

Test Example 2: Inhibition of PGE2 and PGF2 α production in A549 cell

[0446] Human A549 cells were seeded on to a 96-well plate at 2x10⁴ cells/100 μ L per well and allowed to incubate overnight. After removing the culture medium and washed with phosphate buffered saline, the culture medium was replaced by RPMI culture medium with 3%FBS containing a DMSO solution containing test compound or DMSO alone, which was each 0.1% of DMSO final concentration. After incubation for 60 minutes, IL-1 β (5 ng/well) was added and incubated at 37°C for 24 hours. Then, PGE2 in the culture medium was measured using HTRF kit (Cisbio International, catalogue #62P2APEC), and PGF2 α in the culture medium was measured using EIA kit (Cayman Chemical Company, catalogue #516011). The solution free of test compound was used as positive control, and the solution free of test compound and IL-1 β was used as negative control. 100% activity was defined as PGE2 and PGF2 α production in the positive control minus PGE2 and PGF2 α production in the negative control. IC50 value was calculated by standard method.

[0447] The results of Test Example 1 and Test Example 2 are shown in Tables 15-18 ("-" means unadministered test).

[Table 15]

Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)
1	12.7	88.9	39	10.4	236.9
2	7.6	240.5	40	194.5	300.2
3	1.5	24.6	41	52.5	145.0
			42	28.5	1857.9
5	453.4	-	43	70.7	438.3
6	27.4	-	44	4.1	1101.2
7	105.1	341.6	45	4.9	2084.5
8	160.3	105.8	46	26.0	230.8
9	452.2	-	47	269.6	-
10	332.8	-	48	49.9	146.3
11	6.6	12.9	49	100.3	375.3
12	7.1	41	50	140.3	810.2
13	35.1	75.6	51	120.2	330.4
14	3.8	15.2	52	40.2	280.7
15	37.4	465.0	53	179.3	279.9
16	63.9	1303.5	54	23.2	225.3
17	9.8	21.6	55	2.7	64.3
18	21.6	110.1	56	7.5	85.0
19	9.4	47.8	57	0.5	16.8
20	5.6	98	58	5.4	157.0
21	8.3	69.4	59	2.1	12.9

EP 2 746 265 B1

(continued)

5	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)
	22	4.8	104.2	60	19.8	8.0
	23	7.2	115.7	61	3.2	66.7
10	24	10.5	750.6	62	11.0	51.6
	25	89.0	849.6	63	5.6	54.8
	26	171.5	871.5	64	2.3	20.1
	27	2.4	16.0	65	9.6	72.2
15	28	3.3	114.4	66	9.9	57.3
	29	3.5	85.5	67	11.0	56.4
	30	1.8	210.1	68	63.4	10000<
20	31	4.1	69.0	69	10.0	21.6
	32	5.2	71.7	70	4.6	146.9
	33	3.0	33.6	71	33.1	227.2
	34	4.9	27.6	72	9.8	60.0
25	35	20.3	204.6	73	18.6	76.4
	36	147.2	234.2	74	71.3	10.4
	37	23.3	265.1	75	31.7	19.7
30	38	6.6	17.4	76	0.2	5.5

[Table 16]

35	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)
	77	1.7	32.9	115	11.5	37.0
	78	96.8	176.8	116	9.7	251.5
40	79	15.0	284.6	117	583.8	-
	80	29.4	1251.4	118	7.3	5.3
	81	5.0	404.5	119	207.6	-
45	82	0.4	6.3	120	57.5	513.2
	83	9.8	16.5	121	6.3	432.4
	84	0.8	6.5	122	9.7	1000<
	85	0.5	3.7	123	8.7	99.8
50	86	1.0	5.1	124	2.2	3.2
	87	5.7	6.9	125	25.6	56.9
	88	15.1	9.2	126	5.6	13.1
55	89	1.2	2.1	127	6.3	144.1
	90	117.7	348.4	128	1	253.3
	91	327.3	-	129	59.4	177.6

EP 2 746 265 B1

(continued)

Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)
92	6.7	32.2	130	3.9	10000<
93	4.2	153.5	131	32.8	10000<
94	9.1	182.1	132	2.2	590.4
95	5.1	105.3	133	7.0	487.5
96	34.0	125.7	134	10.3	921.3
97	6.1	43.1	135	0.3	87.3
98	6.0	11.9	136	2.4	61.9
99	3.7	18.9	137	3.3	1106.7
100	33.3	122.2	138	2.2	87.4
101	77.8	117.9	139	1.9	72.8
102	1.6	49.1	140	0.2	33.0
103	2.8	59.6	141	0.4	36.5
104	2.9	71.1	142	0.4	82.7
105	8.9	324.6	143	5.3	87.5
106	4.5	63.8	144	9.7	2992.1
107	1.5	42.0	145	1.6	13.0
108	11.5	87.9	146	3.1	128.9
109	7.8	8.7	147	15.8	185.9
110	2.6	35.5	148	1.9	324.4
111	764.8	-	149	1.0	43.3
112	2.6	186.7	150	2.1	375.8
113	約 0.1	1.2	151	11.6	327.5
114	3.5	8.8	152	31.6	854.4

[Table 17]

Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)
153	22.3	617.8	191	3.9	49.5
154	20.0	64.4	192	9.9	18.1
155	100.6	208.7	193	0.4	18.9
156	1.6	24.6	194	0.4	33.8
157	3.9	10856.6	195	2.3	17.2
158	8.5	82.2	196	6.5	35.9
159	403.3	-	197	5.2	34.4
160	7.6	71.6	198	6.0	34.1

EP 2 746 265 B1

(continued)

Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)
161	3.8	10000<	199	9.7	43.2
162	23.8	52.2	200	4.6	107.6
163	5.9	23.2	201	0.5	253.0
164	0.3	13.3	202	4.9	5.9
165	1.3	7.4	203	1.4	5.2
166	3.2	29.0	204	1.0	12.2
167	1.0	217.3	205	3.7	25.6
168	3.2	23.9	206	3.2	74.5
169	2.4	88.6	207	3.6	6.2
170	3.2	17.8	208	0.7	13.0
171	4.7	97.8	209	1.0	10.3
172	4.5	134.4	210	4.7	13.8
173	1.1	60.8	211	2.5	8.5
174	2.0	117.2	212	42.3	305.4
175	3.6	143.9	213	0.9	141.6
176	0.3	138.4	214	5.2	60.7
177	8.3	45.2	215	140.4	772.5
178	1.2	12.8	216	203.6	-
179	3.6	20.2	217	26.9	316.7
180	1.1	773.6	218	5.3	42.8
181	1.1	705.0	219	2.0	26.0
182	1.6	38.4	220	8.6	-
183	1.9	6.1	221	2.2	-
184	1.0	138.2	222	661.1	-
185	4.5	5.6	223	68.2	-
186	6.1	13.9	224	26.5	-
187	3.6	12.7	225	15.6	-
188	456.9	1019.5	226	26.9	-
189	1.4	24.5	227	0.1	-
190	0.1	7.0	228	1.7	-

[Table 18]

Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)
229	2.3	39	244	56.2	104
230	1.9	100	245	25.7	166

EP 2 746 265 B1

(continued)

Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)	Example	mPGES-1 Inhibition IC50 (nM)	A549 Cell PGE2 Production Inhibition IC50 (nM)
231	1.9	150	246	35.9	132
232	7.7	125	247	68.4	55
233	2.5	195	248	201	-
234	97.4	198	249	404	-

Test Example 3: Inhibition of PGE2 production in human whole blood

[0448] Human whole blood assay was carried out according to Brideau et al. (Inflamm. Res., vol.45, p.68, 1996). Venous bloods from volunteers were collected into each tube (containing heparin). These volunteers did not have clinical appearance of inflammation and not take NSAID at least seven days before the blood collection. DMSO solution containing test compound or DMSO alone, which was each 0.25% of DMSO final concentration, was added to the blood, and incubated at 37°C for 20 minutes. LPS from E. coli (E. coli serotype 0111:B4 diluted with phosphate-buffered saline) was added at the final concentration of 100 µg/mL and incubated at 37°C for 24 hours. After the incubation, the blood was centrifuged at 2000 rpm for 5 minutes at 4°C, and PGE2 in the supernatant was measured using HTRF kit (Cisbio International, catalogue #62P2APEC). The solution free of test compound was used as positive control, and the solution free of test compound and LPS was used as negative control. 100% activity was defined as PGE2 production in the positive control minus PGE2 production in the negative control. IC50 value was calculated by standard method.

Test Example 4 : Inhibition of PGE2 production in mouse air pouch model

[0449] Air pouch was established in BALB/c mouse by the subcutaneous injection (twice at intervals of 3 days) of sterilized air into the back. 3 days after the second air injection, 0.5% zymosan was injected into the air pouch. 5 hours later, 1 mL of phosphate-buffered saline containing 10 µM indomethacin was injected into the air pouch to wash the interior thereof, and the white blood cells and PGE2 in the pouch fluid were measured. PGE2 was quantified using HTRF kit (Cisbio International, catalogue #62P2APEC). The test compound was suspended in 0.5% methylcellulose solution and administered orally 1 hour before the zymosan injection. The group which was administered only methylcellulose solution was used as control for comparison.

Test Example 5: Pharmacokinetic assay in rat

[0450] A 10mg/5mL liquid of a test compound was prepared by suspending the compound in 0.5% methylcellulose solution. The liquid was administered orally to female SD rats at fasting state at a dose of 10 mg/kg, and bloods were collected from the jugular vein 1, 2, 4 and 6 hours after the administration. The Blood samples were centrifuged to separate plasma. After deproteinization of the plasma, the concentration of the test compound in plasma was measured by HPLC.

Test Example 6: Evaluation of analgesic effect using mouse writhing test

[0451] Analgesic effect was evaluated from pain-related behavior induced by the intraperitoneal injection of 0.6% acetic acid solution (10mL/kg) into ddY mouse. The number of times of writhing in 20 minutes after the acetic acid injection was counted as pain-related behavior. The test compound was suspended in 0.5% methylcellulose solution and administered orally 1 hour before the acetic acid injection. The group which was administered only methylcellulose solution was used as control for comparison.

Test Example 7: Evaluation of analgesic effect using mouse formalin test

[0452] Analgesic effect was evaluated from pain-related behavior induced by the injection of 5% formalin solution (0.02 mL/mouse) into the right hind-foot pad of ICR mouse. The sum of the duration of licking in 30 minutes after the formalin injection was measured as pain-related behavior. The test compound was suspended in 0.5% methylcellulose solution and administered orally 1 hour before the formalin injection. The group which was administered only methylcellulose solution was used as control for comparison.

Test Example 8: Evaluation of antiinflammatory effect in rat carrageenan edema model

[0453] Antiinflammatory effect was evaluated from paw edema induced by the injection of 0.5% carrageenan solution (0.1 mL/rat) into the right hind-foot pad of SD rat. The volume of the paw was measured before and 3 hours later the carrageenan injection using a plethysmometer (Unicom Inc., catalogue #TK-105). Swelling volume was determined from the increase in the volume from the volume before the injection. The inhibiting rate was determined as the ratio of the average swelling volume in the group given the test compound to the average swelling in the control group. The test compound was suspended in 0.5% methylcellulose solution and administered orally 1 hour before the carrageenan injection. The group which was administered only methylcellulose solution was used as control for comparison.

[0454] The compound of the invention or a pharmaceutically acceptable salt thereof showed potent mPGES-1 inhibiting activity and inhibited PGE₂ production, as shown in Test Example 1 and Test Example 2. Thus, it is particularly useful as an agent for the treatment or prevention of inflammatory bowel disease, irritable bowel syndrome, migraine, headache, low back pain, spinal stenosis, herniated disk, temporomandibular joint disorder, cervical syndrome, cervical spondylosis, endometriosis, adenomyosis, preterm labour and delivery, threatened premature delivery, dysmenorrhea, overactive bladder, nocturia, interstitial cystitis, neurodegenerative disease, psoriasis, rheumatoid arthritis, rheumatic fever, fibromyalgia, neuralgia, complex regional pain syndrome, fascial dysfunction, viral infections, bacterial infection, mycosis, burn, inflammation and pain after operation, injury and dental extraction, malignant tumors, atherosclerosis, stroke, gout, arthritis, osteoarthritis, juvenile arthritis, ankylosing spondylitis, tenosynovitis, ligament ossification, systemic lupus erythematosus, vasculitis, pancreatitis, nephritis, conjunctivitis, iritis, scleritis, uveitis, wound therapy, dermatitis, eczema, osteoporosis, asthma, chronic obstructive pulmonary disease, pulmonary fibrosis, allergic disease, familial adenomatous polyposis, scleroderma, bursitis, leiomyoma of uterus, prostatitis, and a pain from cancer.

Formulation 1: 80 mg Tablets for oral administration

[0455]

Ingredient	Quantity (mg/tablet)
Compound of Example 1	5.0
Corn starch	46.6
Cellulose, crystalline	24.0
Methylcellulose	4.0
Magnesium stearate	0.4


[0456] The components are blended and compressed to form tablets.

[Industrial applicability]




[0457] The compound of the invention or a pharmaceutically acceptable salt thereof has mPGES-1 inhibiting activity and is particularly useful as an agent for the treatment or prevention of inflammatory bowel disease, irritable bowel syndrome, migraine, headache, low back pain, spinal stenosis, herniated disk, temporomandibular joint disorder, cervical syndrome, cervical spondylosis, endometriosis, adenomyosis, preterm labour and delivery, threatened premature delivery, dysmenorrhea, overactive bladder, nocturia, interstitial cystitis, neurodegenerative disease, psoriasis, rheumatoid arthritis, rheumatic fever, fibromyalgia, neuralgia, complex regional pain syndrome, fascial dysfunction, viral infections, bacterial infection, mycosis, burn, inflammation and pain after operation, injury and dental extraction, malignant tumors, atherosclerosis, stroke, gout, arthritis, osteoarthritis, juvenile arthritis, ankylosing spondylitis, tenosynovitis, ligament ossification, systemic lupus erythematosus, vasculitis, pancreatitis, nephritis, conjunctivitis, iritis, scleritis, uveitis, wound therapy, dermatitis, eczema, osteoporosis, asthma, chronic obstructive pulmonary disease, pulmonary fibrosis, allergic disease, familial adenomatous polyposis, scleroderma, bursitis, leiomyoma of uterus, prostatitis, and a pain from cancer.

Claims

1. A heterocyclic derivative represented by the general formula [1]:




or its tautomer or a pharmaceutically acceptable salt thereof, wherein ring A is a group represented by the general formulae [2], [3] or [4]:



wherein

X¹ is NH, N-alkyl, or O;
A¹ is hydrogen or alkyl;
A² is

- i) hydrogen;
- ii) halogen;
- iii) alkyl optionally substituted with one to three groups selected from the group consisting of halogen, amino, monoalkylamino, dialkylamino, carbamoyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, a saturated cyclic aminocarbonyl, alkoxy, alkoxyalkoxy and alkylcarbonyloxy;
- iv) cycloalkyl optionally substituted with alkyl optionally substituted with one to three halogens;
- v) alkoxy;
- vi) a saturated heterocycle group optionally substituted with alkyl, alkyloxycarbonyl, alkylcarbonyl or oxo;
- vii) alkylthio;
- viii) alkylsulfonyl;
- ix) alkylsulfinyl;
- x) a group of the general formula [5]:



wherein
R³ and R⁴ are the same or different group selected from

- a) hydrogen,
- b) alkyl optionally substituted with a group selected from the group consisting of monoalkylamino, dialkylamino, a saturated cyclic amino optionally substituted with alkyl, a saturated heterocycle group optionally substituted with alkyl, alkoxy, hydroxycarbonyl, hydroxyl, alkyloxycarbonyl and alkylthio, or
- c) cycloalkyl; or

xi) a saturated cyclic amino optionally substituted with alkyl, amino, monoalkylamino, dialkylamino, alkoxy or hydroxyl;

R¹ is phenyl, benzyl, naphthyl, cycloalkyl, cycloalkylmethyl, heteroaryl, heteroarylmethyl, 1,2,3,4-tetrahydronaphthalen-5-yl, 1,2,3,4-tetrahydronaphthalen-6-yl, 2,3-dihydro-1H-inden-4-yl, 2,3-dihydro-1H-inden-5-yl, 1,2-dihydrocyclobutabenzen-3-yl, 1,2-dihydrocyclobutabenzen-4-yl or alkyl, wherein said phenyl, benzyl, cycloalkyl, cycloalkylmethyl, heteroaryl and heteroarylmethyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen,
- ii) alkyl optionally substituted with one to three groups selected from the group consisting of halogen, hydroxy and phenyl,
- iii) alkoxy,
- iv) hydroxy, and
- v) cyano;

R² is phenyl or pyridyl, wherein said phenyl and pyridyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen,
- ii) alkylsulfonyl,
- iii) alkoxy optionally substituted with one to three halogens or alkoxy;
- iv) alkynyl optionally substituted with alkoxyalkyl or cycloalkyl, and
- v) alkyl optionally substituted with one to three groups selected from the group consisting of alkoxy, alkoxyalkoxy, cycloalkyl, phenyl and halogen.

2. The heterocyclic derivative according to claim 1, wherein the ring A is a group of formula [4] and X¹ is NH, or its tautomer or a pharmaceutically acceptable salt thereof.

3. The heterocyclic derivative according to claim 1, wherein R¹ is phenyl, 1,2,3,4-tetrahydronaphthalen-5-yl, 1,2,3,4-tetrahydronaphthalen-6-yl, 2,3-dihydro-1H-inden-4-yl, 2,3-dihydro-1H-inden-5-yl, 1,2-dihydrocyclobutabenzen-3-yl, or 1,2-dihydrocyclobutabenzen-4-yl, and said phenyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen,
- ii) alkyl optionally substituted with one to three halogens,
- iii) alkoxy, and
- iv) cyano,

or its tautomer or a pharmaceutically acceptable salt thereof.

4. The heterocyclic derivative according to claim 1, wherein R² is phenyl and said phenyl is optionally substituted with one to three groups selected from the group consisting of

- i) halogen
- ii) alkylsulfonyl,
- iii) alkoxy optionally substituted with alkoxy,
- iv) alkynyl optionally substituted with alkoxyalkyl or cycloalkyl, and
- v) alkyl optionally substituted with one to three groups selected from the group consisting of halogen, alkoxy, alkoxyalkoxy, cycloalkyl and phenyl,

or its tautomer or a pharmaceutically acceptable salt thereof.

5. The heterocyclic derivative according to claim 1,
wherein

the ring A is a group of formula [4],

X¹ is NH,

A² is

i) hydrogen,

ii) alkyl optionally substituted with a group selected from the group consisting of halogen, monoalkylamino, dialkylamino, monoalkylaminocarbonyl, dialkylaminocarbonyl, a saturated cyclic aminocarbonyl, alkoxy, alkoxy-alkoxy and alkylcarbonyloxy,

iii) cycloalkyl optionally substituted with alkyl optionally substituted with one to three halogens,

iv) alkoxy,

v) a saturated heterocyclic group optionally substituted with alkyl or alkyloxycarbonyl,

vi) alkylthio,

vii) alkylsulfonyl,

viii) alkylsulfinyl,

ix) amino substituted with alkyl wherein said alkyl is optionally substituted with a group selected from the group consisting of monoalkylamino, dialkylamino, a saturated cyclic amino optionally substituted with alkyl, tetrahydrofuryl, morpholino, alkoxy, hydroxycarbonyl, hydroxyl and alkylthio,

x) amino substituted with cycloalkyl or

xi) a saturated cyclic amino optionally substituted with alkyl, dialkylamino, alkoxy or hydroxyl, and

R¹ is

i) phenyl optionally substituted with one to three groups selected from the group consisting of halogen, alkyl optionally substituted with one to three halogens, alkoxy and cyano,

ii) 1,2,3,4-tetrahydronaphthalen-5-yl,

iii) 2,3-dihydro-1H-inden-5-yl,

iv) benzyl optionally substituted with halogen or alkyl optionally substituted with one to three halogens,

v) cycloalkyl,

vi) cycloalkylmethyl,

vii) naphthyl,

viii) pyridylmethyl optionally substituted with alkyl optionally substituted one to three halogens,

ix) thienyl,

x) thienylmethyl,

xi) benzothiazolyl,

xii) benzothiadiazolyl,

xiii) indolyl or

xiv) alkyl, and

R² is phenyl or pyridyl

wherein said phenyl is optionally substituted with one to three groups selected from the group consisting of

i) halogen,

ii) alkylsulfonyl,

iii) alkoxy optionally substituted with alkoxy,

iv) alkynyl optionally substituted with alkoxyalkyl or cycloalkyl, and

v) alkyl optionally substituted with one to three groups selected from the group consisting of halogen, alkoxy, alkoxyalkoxy, cycloalkyl and phenyl, and

said pyridyl is optionally substituted with halogen,

or its tautomer or a pharmaceutically acceptable salt thereof.

6. The heterocyclic derivative according to claim 1, wherein
the ring A is a group of formula [4],

X¹ is NH,

A² is alkyl substituted with alkoxy, dialkylamino, tetrahydrofuryl, tetrahydrofurylmethyl, alkoxyalkylamino, or cycloalkyl optionally substituted with alkyl optionally substituted with one to three halogens,

R¹ is phenyl substituted with one halogen and one methyl, and R² is phenyl optionally substituted with one trifluoromethyl or two halogens,

or its tautomer or a pharmaceutically acceptable salt thereof.

7. The heterocyclic derivative according to claim 1 selected from the following (1)-(239), or its tautomer or a pharmaceutically acceptable salt thereof:

- (1) N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (2) N-cyclohexyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (3) N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (4) N-[(1-hydroxycyclohexyl)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (5) N-[2-(trifluoromethyl)benzyl]-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (6) N-cyclohexyl-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (7) N-(3-chloro-2-methylphenyl)-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (8) N-cyclohexyl-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-indazole-7-carboxamide,
- (9) N-[2-(trifluoromethyl)benzyl]-5-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-indazole-7-carboxamide,
- (10) N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (11) 2-methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (12) N-cyclohexyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (13) N-(3-chloro-2-methylphenyl)-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (14) N-cyclopentyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (15) N-cyclobutyl-2-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (16) N-(3-chloro-2-methylphenyl)-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (17) N-cyclohexyl-2-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (18) 2-ethyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (19) N-cyclohexyl-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (20) 2-(methoxymethyl)-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (21) 2-(methoxymethyl)-N-(2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (22) 2-(methoxymethyl)-N-(4-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (23) N-(2-chlorobenzyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (24) 2-(methoxymethyl)-N-(4-methylbenzyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (25) N-(4,4-difluorocyclohexyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (26) N-(4-tert-butylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (27) 2-(methoxymethyl)-N-[4-(trifluoromethyl)phenyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (28) N-(2,4-dimethylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (29) N-(2-chloro-4-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

zimidazole-4-carboxamide,

(30) N-(3,4-dimethylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(31) N-(3-chloro-4-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(32) N-(2,3-dihydro-1H-inden-5-yl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(33) 2-(methoxymethyl)-N-(5,6,7,8-tetrahydronaphthalen-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(34) N-(2-fluorophenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(35) 2-(methoxymethyl)-N-(2-methoxyphenyl)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(36) 2-(methoxymethyl)-N-(4-methoxyphenyl)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(37) N-(3-bromo-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(38) N-(3-chloro-2-methylbenzyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(39) N-(2,6-difluorophenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(40) N-(3-cyano-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(41) 2-(methoxymethyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-N-[[3-(trifluoromethyl)pyridin-2-yl]methyl]-1H-benzimidazole-4-carboxamide,

(42) N-(2-chloro-6-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(43) 2-(2-amino-2-oxoethyl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl) phenyl] carbonyl }amino)-1H-benzimidazole-4-carboxamide,

(44) 2-(2-amino-2-oxoethyl)-N-[2-(trifluoromethyl)benzyl]-6-({[2 -(trifluoromethyl) phenyl] carbonyl }amino)-1H-benzimidazole-4-carboxamide,

(45) N-(3-chloro-2-methylphenyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(46) N-cyclohexyl-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(47) 1-methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(48) N-(3-chloro-2-methylphenyl)-1-ethyl-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(49) N-cyclohexyl-1-ethyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(50) 1-ethyl-N-[2-(trifluoromethyl)benzyl]-6-({[2(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(51) N-(3-chloro-2-methylphenyl)-2-methyl-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1,3-benzoxazole-4-carboxamide,

(52) 2-methyl-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1,3-benzoxazole-4-carboxamide,

(53) N-(3-chloro-2-methylphenyl)-2-ethyl-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1,3-benzoxazole-4-carboxamide,

(54) N-(3-chloro-2-methylphenyl)-2-ethoxy-6-({[2-(trifluoromethyl)phenyl] carbonyl }amino)-1H-benzimidazole-4-carboxamide,

(55) 2-ethoxy-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(56) N-(3-chloro-2-methylphenyl)-2-(1-chloro-2-methylpropan-2-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-H-benzimidazole-4-carboxamide,

(57) N-(3-chloro-2-methylphenyl)-2-[(dimethylamino)methyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(58) N-(3-chloro-2-methylphenyl)-2-(2-methylpropyl)-6-({[2-(trifluoromethyl) phenyl] carbonyl }amino)-1H-benzimidazole-4-carboxamide,

(59) 2-(2-methylpropyl)-N-[2-(trifluoromethyl)benzyl]-6-({[2-(trifluoromethyl) phenyl] carbonyl}amino)-1H-benz-

imidazole-4-carboxamide,

(60) tert-butyl 3-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino}-1H-benzimidazol-2-yl}azetidine-1-carboxylate,

(61) N-(3-chloro-2-methylphenyl)-2-[(methylamino)methyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(62) 4-{[(3-chloro-2-methylphenyl)carbamoyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazol-2-yl}methyl acetate,

(63) N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(64) 2-[[2-(2R)-tetrahydrofuran-2-yl]-N-[2-(trifluoromethyl)benzyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-H-benzimidazole-4-carboxamide,

(65) N-(3-chloro-2-methylphenyl)-2-[(2S)-tetrahydrofuran-2-yl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(66) 2-[[2-(2S)-tetrahydrofuran-2-yl]-N-[2-(trifluoromethyl)benzyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(67) 2-(1-acetylazetidin-3-yl)-N-(3-chloro-2-methylphenyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(68) tert-butyl (2S)-2-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazol-2-yl}pyrrolidine-1-carboxylate,

(69) tert-butyl (2R)-2-{4-[(3-chloro-2-methylphenyl)carbamoyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazol-2-yl}pyrrolidine-1-carboxylate,

(70) N-(3-chloro-2-methylphenyl)-2-[(2S)-pyrrolidin-2-yl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(71) N-(3-chloro-2-methylphenyl)-2-[(2S)-1-methylpyrrolidin-2-yl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(72) 2-[(2S)-1-acetylpyrrolidin-2-yl]-N-(3-chloro-2-methylphenyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(73) N-(3-chloro-2-methylphenyl)-2-[(2-methoxyethoxy)methyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(74) N-(3-chloro-2-methylphenyl)-2-(1-methoxy-2-methylpropan-2-yl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(75) 2-tert-butyl-N-(3-chloro-2-methylphenyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(76) 2-tert-butyl-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-N-[[3-(trifluoromethyl)pyridin-2-yl]methyl]-1H-benzimidazole-4-carboxamide,

(77) N-(3-chloro-2-methylphenyl)-2-(2-ethoxyethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(78) N-(3-chloro-2-methylphenyl)-2-(ethoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(79) 2-(ethoxymethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-N-[[3-(trifluoromethyl)pyridin-2-yl]methyl]-1H-benzimidazole-4-carboxamide,

(80) N-(3-chloro-2-methylphenyl)-2-(2-methoxyethyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(81) N-(3-chloro-2-methylphenyl)-2-(2,2-dimethylpropyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(82) N-(3-chloro-2-methylphenyl)-2-cyclopropyl-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(83) N-(3-chloro-2-methylphenyl)-2-(2-methylpentan-2-yl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(84) N-(3-chloro-2-methylphenyl)-2-(1-methylcyclopropyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(85) 2-tert-butyl-N-(3-chloro-4-methylphenyl)-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(86) 2-tert-butyl-N-(3-chloro-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(87) 2-tert-butyl-N-(3-chloro-4-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(88) N-(3-chloro-2-methylphenyl)-2-[1-(trifluoromethyl)cyclopropyl]-6-[[2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

yl}amino)-1H-benzimidazole-4-carboxamide,

(89) N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(90) N-(2-chlorobenzyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(91) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(92) 6-({[2-chloro-4-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-methoxymethyl-1H-benzimidazole-4-carboxamide,

(93) 6-({[2-chloro-5-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(94) N-(3-chloro-2-methylphenyl)-6-({[2-chlorophenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(95) N-(3-chloro-2-methylphenyl)-6-({[2-chloropyridin-3-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(96) 6-({[2-bromophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(97) N-(3-chloro-2-methylphenyl)-6-({[2,6-dichlorophenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(98) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(99) 6-({[2-chloro-3-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(100) 6-({[2-chloro-3,6-difluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(101) 6-({[2-bromo-6-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(102) 6-({[2-bromo-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(103) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-6-methylphenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(104) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-4-methylphenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(105) 6-({[5-bromo-2-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(106) 6-({[2-bromo-5-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(107) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-5-methylphenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(108) N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-({[5-methyl-2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(109) 6-({[2,5-bis(trifluoromethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(110) 6-({[2,4-bis(trifluoromethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(111) N-(3-chloro-2-methylphenyl)-6-({[5-fluoro-2-(trifluoromethyl)phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(112) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-6-(trifluoromethyl)phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(113) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-5-[2-(propan-2-yloxy)ethoxy]phenyl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(114) 6-({[2-chloro-5-(2-ethoxyethoxy)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(115) 6-({[2-chloro-5-(3-methoxypropyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(116) 6-({[5-(3-tert-butoxyprop-1-yn-1-yl)-2-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(117) 6-({[5-(3-tert-butoxypropyl)-2-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(meth-

oxymethyl)-1H-benzimidazole-4-carboxamide,

(118) 6-({[2-chloro-5-(3-hydroxy-3-methylbutyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(119) 6-({[2-chloro-5-(ethoxymethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(120) 6-({[2-chloro-5-[(2-ethoxyethoxy)methyl]phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(121) 6-({[2-chloro-5-(2-cyclopropylethyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(122) N-(3-chloro-2-methylphenyl)-6-({[2-chloro-5-(2-phenylethyl)phenyl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(123) N-(3-chloro-2-methylphenyl)-2-cyclopentyl-6-({[2-(trifluoromethyl)phenyl] carbonyl }amino)-1H-benzimidazole-4-carboxamide,

(124) N-(3-chloro-2-methylphenyl)-2-cyclopentyl-6-({[2,5-dichlorophenyl]carbonyl} amino)-1H-benzimidazole-4-carboxamide,

(125) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-cyclopentyl-1H-benzimidazole-4-carboxamide,

(126) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(127) N-(3-chloro-2-methylphenyl)-6-({[2,6-dichlorophenyl]carbonyl}amino)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(128) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(129) N-(3-chloro-2-methylphenyl)-2-[(2S)-5-oxopyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(130) N-(3-chloro-2-methylphenyl)-2-[(2R)-5-oxopyrrolidin-2-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(131) N-(3-chloro-2-methylphenyl)-2-[2-oxo-2-(pyrrolizin-1-yl)ethyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(132) N-(3-chloro-2-methylphenyl)-2-[2-(dimethylamino)-2-oxoethyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(133) N-(3-chloro-2-methylphenyl)-2-[2-(methylamino)-2-oxoethyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(134) 2-chloro-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl} amino)-1H-benzimidazole-4-carboxamide,

(135) N-(3-chloro-2-methylphenyl)-2-[(2-methoxyethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(136) N-(3-chloro-2-methylphenyl)-2-[(2-hydroxyethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(137) N-(3-chloro-2-methylphenyl)-2-(methylamino)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(138) N-(3-chloro-2-methylphenyl)-2-(ethylamino)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(139) N-(3-chloro-2-methylphenyl)-2-[(2,2-dimethylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(140) N-(3-chloro-2-methylphenyl)-2-(cyclopentylamino)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(141) N-(3-chloro-2-methylphenyl)-2-(piperidin-1-yl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(142) N-(3-chloro-2-methylphenyl)-2-(4-methylpiperazin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(143) 2-[bis(2-hydroxyethyl)amino]-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(144) N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(145) N-(3-chloro-2-methylphenyl)-2-[(2-(morpholin-4-yl)ethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(146) N-(3-chloro-2-methylphenyl)-2-[(2-(dimethylamino)ethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

yl}amino)-1H-benzimidazole-4-carboxamide,

(147) N-(3-chloro-2-methylphenyl)-2-(3-hydroxyazetidin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(148) N-(3-chloro-2-methylphenyl)-2-[(3S)-3-(dimethylamino)pyrrolizin-1-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(149) N-(3-chloro-2-methylphenyl)-2-[(3S)-3-hydroxypyrrolizin-1-yl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(150) N-(3-chloro-2-methylphenyl)-2-{{[2-(diethylamino)ethyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(151) N-(3-chloro-2-methylphenyl)-2-{{[2-(pyrrolizin-1-yl)ethyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(152) N-(3-chloro-2-methylphenyl)-2-{{[3-(dimethylamino)propyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(153) N-(3-chloro-2-methylphenyl)-2-{{[3-(dimethylamino)-2,2-dimethylpropyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(154) N-(3-chloro-2-methylphenyl)-2-{{[2-(dipropylamino)ethyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(155) N-(3-chloro-2-methylphenyl)-2-(morpholin-4-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(156) 2-amino-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(157) N-(3-chloro-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(158) N-(3-chloro-2-methylphenyl)-2-{{[3-methyloxetan-3-yl)methyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(159) tert-butyl N-[4-{{[3-chloro-2-methylphenyl]carbonyl}amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl]glycinate,

(160) N-[4-{{[3-chloro-2-methylphenyl]carbonyl}amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl]glycine,

(161) N-(3-chloro-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1-methyl-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(162) N-(3-chloro-2-methylphenyl)-2-[(3-methoxy-2,2-dimethylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(163) N-(3-chloro-2-methylphenyl)-2-(pyrrolizin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(164) 2-(azetidin-1-yl)-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(165) N-(3-chloro-2-methylphenyl)-2-(3-methoxyazetidin-1-yl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(166) N-(3-chloro-2-methylphenyl)-2-[(2-hydroxy-2-methylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(167) N-(3-chloro-2-methylphenyl)-2-{{[(2S)-tetrahydrofuran-2-ylmethyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(168) N-(3-chloro-2-methylphenyl)-2-{{[(2R)-tetrahydrofuran-2-ylmethyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(169) N-(3-chloro-2-methylphenyl)-2-{{[(2S)-1-hydroxy-3-methylbutan-2-yl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(170) N-(3-chloro-2-methylphenyl)-2-{{[(2R)-1-hydroxy-3-methylbutan-2-yl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(171) N-(3-chloro-2-methylphenyl)-2-{{[(2S)-1-hydroxy-3,3-dimethylbutan-2-yl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(172) N-(3-chloro-2-methylphenyl)-2-{{[3-methoxy-2,2-dimethylpropyl(methyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(173) N-(3-chloro-2-methylphenyl)-2-[(3-methoxypropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(174) N-(3-chloro-2-methylphenyl)-2-{{[2-(propan-2-yloxy)ethyl]amino}-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(175) 2-[(2-tert-butoxyethyl)amino]-N-(3-chloro-2-methylphenyl)-6-({[2-(trifluoromethyl)phenyl]carbonyl}ami-

no)-1H-benzimidazole-4-carboxamide,

(176) N-(3-chloro-2-methylphenyl)-2-[(2-methoxy-2-methylpropyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(177) N-(3-chloro-2-methylphenyl)-2-[(2-(methylsulfanyl)ethyl)amino]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(178) N-(3-chloro-2-methylphenyl)-2-(methylsulfanyl)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(179) N-(3-chloro-2-methylphenyl)-2-(methylsulfonyl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(180) N-(3-chloro-2-methylphenyl)-2-(methylsulfinyl)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(181) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,

(182) N-(3-chloro-2-methylphenyl)-6-({[2,6-dichlorophenyl]carbonyl}amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,

(183) N-(3-chloro-2-methylphenyl)-6-({[2,4-dichlorophenyl]carbonyl}amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,

(184) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,

(185) 6-({[2-bromo-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,

(186) 6-({[2-bromo-6-chlorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1 H-benzimidazole-4-carboxamide,

(187) 6-({[2-chloro-5-(cyclopropylethynyl)phenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,

(188) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(189) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-[(3-methoxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(190) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-[(2-hydroxy-2-methylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(191) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-[(2-methoxy-2-methylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(192) N-(3-chloro-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-[(2-(propan-2-yloxy)ethyl)amino]-1H-benzimidazole-4-carboxamide,

(193) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(2-(propan-2-yloxy)ethyl)amino]-1H-benzimidazole-4-carboxamide,

(194) 2-[(2-tert-butoxyethyl)amino]-6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-1H-benzimidazole-4-carboxamide,

(195) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(3-methoxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(196) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(2-methoxy-2-methylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(197) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(2S)-tetrahydrofuran-2-ylmethyl]amino]-1H-benzimidazole-4-carboxamide,

(198) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-ylmethyl]amino]-1H-benzimidazole-4-carboxamide,

(199) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(200) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-2-methylphenyl)-2-[(2S)-1-hydroxy-3-methylbutan-2-yl]amino]-1H-benzimidazole-4-carboxamide,

(201) N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(202) N-(4-tert-buthylphenyl)-2-(dimethylamino)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(203) N-(2,3-dihydro-1H-inden-5-yl)-2-(dimethylamino)-6-({[2-(trifluoromethyl) phenyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(204) 6-({[2-chloro-6-fluorophenyl]carbonyl}amino)-N-(3-chloro-4-methylphenyl)-2-(dimethylamino)-1H-benz-

imidazole-4-carboxamide,

(205) N-(3-chloro-4-methylphenyl)-6-([(2,6-dichlorophenyl)carbonyl]amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,

(206) N-(3-chloro-4-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-(dimethylamino)-1H-benzimidazole-4-carboxamide,

(207) N-(3-chloro-2-methylphenyl)-2-cyclopropyl-6-([(2,5-dichlorophenyl)carbonyl] amino)-1H-benzimidazole-4-carboxamide,

(208) N-(3-chloro-4-methylphenyl)-2-cyclopropyl-6-([(2,5-dichlorophenyl)carbonyl] amino)-1H-benzimidazole-4-carboxamide,

(209) N-(3-chloro-2-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxamide,

(210) N-(3-chloro-4-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-(1-methylcyclopropyl)-1H-benzimidazole-4-carboxamide,

(211) N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-6-([(2-(methylsulfonyl) phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(212) N-(3-chloro-4-methylphenyl)-6-([(2,5-dichlorophenyl)carbonyl]amino)-2-(2-methoxyethyl)-1 H-benzimidazole-4-carboxamide,

(213) 2-(methoxymethyl)-N-phenyl-6-([(2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(214) 2-(methoxymethyl)-N-propyl-6-([(2-(trifluoromethyl)phenyl]carbonyl]amino)-1 H-benzimidazole-4-carboxamide,

(215) 2-(methoxymethyl)-N-(pyridin-3-yl)-6-([(2-(trifluoromethyl)phenyl]carbonyl] amino)-1H-benzimidazole-4-carboxamide,

(216) N-benzyl-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(217) N-(cyclohexylmethyl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(218) 2-(methoxymethyl)-N-(naphthalen-1-yl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(219) 2-(methoxymethyl)-N-(thiophen-3-yl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(220) N-(2,1,3-benzothiadiazol-4-yl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl) phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(221) N-(1,1-dioxide-1-benzothiophen-6-yl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(222) 2-(methoxymethyl)-N-(thiophen-2-ylmethyl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(223) N-(1H-indol-5-yl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl]carbonyl] amino)-1H-benzimidazole-4-carboxamide,

(224) N-(1,3-benzothiazol-2-yl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(225) N-(2,2-dimethylpropyl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(226) 2-(methoxymethyl)-N-(thiophen-2-yl)-6-([(2-(trifluoromethyl)phenyl]carbonyl] amino)-1H-benzimidazole-4-carboxamide,

(227) N-(5-chloro-1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl) phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(228) N-(2-benzylphenyl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(229) 2-(methoxymethyl)-N-(quinolin-8-yl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(230) N-(cycloheptylmethyl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(231) N-(1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl] carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(232) N-(6-chloro-1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-([(2-(trifluoromethyl) phenyl]carbonyl]amino)-1H-benzimidazole-4-carboxamide,

(233) N-[3-chloro-2-(hydroxymethyl)phenyl]-2-(methoxymethyl)-6-([(2-(trifluoromethyl)phenyl]carbonyl]ami-

no)-1H-benzimidazole-4-carboxamide,

(234) N-(3-chloro-2-methylphenyl)-6-[[[3-fluoropyridin-2-yl]carbonyl]amino]-2-(methoxymethyl)-1 H-benzimidazole-4-carboxamide,

(235) N-(3-chloro-2-methylphenyl)-6-[[[3-chloropyridin-4-yl]carbonyl]amino]-2-(methoxymethyl)-1 H-benzimidazole-4-carboxamide,

(236) N-(3-chloro-2-methylphenyl)-6-[[[3,5-dichloropyridin-4-yl]carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(237) 6-[[[5-butoxy-2-chlorophenyl]carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(238) 6-[[[2-chloro-5-(2,2-difluoroethoxy)phenyl]carbonyl]amino]-N-(3-chloro-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide,

(239) N-(3-chloro-2-methylphenyl)-6-[[[2-chloro-5-(4,4,4-trifluorobutoxy)phenyl] carbonyl]amino]-2-(methoxymethyl)-1H-benzimidazole-4-carboxamide.

8. A pharmaceutical composition containing a compound selected from the group consisting of the heterocyclic derivative according to any one of claims 1 to 7, its tautomer and a pharmaceutically acceptable salt thereof as an active ingredient.

9. A compound selected from the group consisting of the heterocyclic derivative according to any one of claims 1 to 7, its tautomer and a pharmaceutically acceptable salt thereof for use as an mPGES inhibiting agent.

10. A compound selected from the group consisting of the heterocyclic derivative according to any one of claims 1 to 7, its tautomer and a pharmaceutically acceptable salt thereof, for use in the prevention or treatment of inflammatory bowel disease, irritable bowel syndrome, migraine, headache, low back pain, spinal stenosis, herniated disk, temporomandibular joint disorders, cervical syndrome, cervical spondylosis, endometriosis, adenomyosis, preterm labour and delivery, threatened premature delivery, dysmenorrhea, overactive bladder, nocturia, interstitial cystitis, neurodegenerative disease, psoriasis, rheumatoid arthritis, rheumatic fever, fibromyalgia, neuralgia, complex regional pain syndrome, fascial dysfunction, viral infections, bacterial infection, mycosis, burn, inflammation and pain after operation, injury and dental extraction, malignant tumors, atherosclerosis, stroke, gout, arthritis, osteoarthritis, juvenile arthritis, ankylosing spondylitis, tenosynovitis, ligament ossification, systemic lupus erythematosus, vasculitis, pancreatitis, nephritis, conjunctivitis, iritis, scleritis, uveitis, wound therapy, dermatitis, eczema, osteoporosis, asthma, chronic obstructive pulmonary disease, pulmonary fibrosis, allergic disease, familial adenomatous polyposis, scleroderma, bursitis, leiomyoma of uterus, or a pain from cancer.


11. The compound for use according to claim 10, wherein the neurodegenerative disease is Alzheimer's disease or multiple sclerosis.

12. The compound for use according to claim 10, wherein the viral infectious disease is influenza, common cold, zoster or AIDS.

13. The compound for use according to claim 10, wherein the malignant tumor is colon cancer, breast cancer, lung cancer or prostatic cancer.


Patentansprüche

1. Ein Heterocyclisches Derivat, dargestellt durch die allgemeine Formel [1]:




[1]


oder sein Tautomer oder ein pharmazeutisch verträgliches Salz davon, wobei Ring A eine Gruppe ist, dargestellt durch die allgemeinen Formeln [2], [3] oder [4]:



[2]



[3]



[4]

wobei

X¹ gleich NH, N-Alkyl oder O ist;
A¹ Wasserstoff oder Alkyl ist;
A²

i) Wasserstoff;

ii) Halogen;

iii) Alkyl, gegebenenfalls substituiert mit einer bis drei Gruppen, ausgewählt aus der Gruppe bestehend aus Halogen, Amino, Monoalkylamino, Dialkylamino, Carbamoyl, Monoalkylaminocarbonyl, Dialkylaminocarbonyl, einem gesättigten cyclischen Aminocarbonyl, Alkoxy, Alkoxyalkoxy und Alkylcarbonyloxy;

iv) Cycloalkyl, gegebenenfalls substituiert mit Alkyl, gegebenenfalls substituiert mit einem bis drei Halogenen;

v) Alkoxy;


vi) eine gesättigte Heterocyclusgruppe, gegebenenfalls substituiert mit Alkyl, Alkylloxycarbonyl, Alkylcarbonyl oder Oxo;

vii) Alkylthio;

viii) Alkylsulfonyl;

ix) Alkylsulfinyl;

x) eine Gruppe der allgemeinen Formel [5]:



[5]

wobei

R³ und R⁴ die gleiche oder verschiedene Gruppe sind, ausgewählt aus

- a) Wasserstoff,
- b) Alkyl, gegebenenfalls substituiert mit einer Gruppe, ausgewählt aus der Gruppe bestehend aus Monoalkylamino, Dialkylamino, einem gesättigten cyclischen Amino, gegebenenfalls substituiert mit Alkyl, einer gesättigten heterocyclischen Gruppe, gegebenenfalls substituiert mit Alkyl, Alkoxy, Hydroxycarbonyl, Hydroxyl, Alkyloxycarbonyl und Alkylthio oder
- c) Cycloalkyl; oder

xi) ein gesättigtes cyclisches Amin, gegebenenfalls substituiert mit Alkyl, Amino, Monoalkylamino, Dialkylamino, Alkoxy oder Hydroxyl ist;

R¹ Phenyl, Benzyl, Naphthyl, Cycloalkyl, Cycloalkylmethyl, Heteroaryl, Heteroaryl-methyl, 1,2,3,4-Tetrahydronaphthalen-5-yl, 1,2,3,4-Tetrahydronaphthalen-6-yl, 2,3-Dihydro-1H-inden-4-yl, 2,3-Dihydro-1H-inden-5-yl, 1,2-Dihydrocyclobutabenzol-3-yl, 1,2-Dihydrocyclobutabenzol-4-yl oder Alkyl ist, wobei das Phenyl, Benzyl, Cycloalkyl, Cycloalkylmethyl, Heteroaryl und Heteroaryl-methyl gegebenenfalls mit einer bis drei Gruppen substituiert ist, ausgewählt aus der Gruppe bestehend aus

- i) Halogen,
- ii) Alkyl, gegebenenfalls substituiert mit einer bis drei Gruppen, ausgewählt aus der Gruppe bestehend aus Halogen, Hydroxy und Phenyl,
- iii) Alkoxy,
- iv) Hydroxy und
- v) Cyano;

R² Phenyl oder Pyridyl ist, wobei das Phenyl und Pyridyl gegebenenfalls mit einer bis drei Gruppen substituiert ist, ausgewählt aus der Gruppe bestehend aus

- i) Halogen,
- ii) Alkylsulfonyl,
- iii) Alkoxy, gegebenenfalls substituiert mit einem bis drei Halogenen oder Alkoxy;
- iv) Alkynyl, gegebenenfalls substituiert mit Alkoxyalkyl oder Cycloalkyl und
- v) Alkyl, gegebenenfalls substituiert mit einer bis drei Gruppen, ausgewählt aus der Gruppe bestehend aus Alkoxy, Alkoxyalkoxy, Cycloalkyl, Phenyl und Halogen.

2. Das heterocyclische Derivat nach Anspruch 1, wobei Ring A eine Gruppe der Formel [4] ist und X¹ NH ist, oder sein Tautomer oder ein pharmazeutisch verträgliches Salz davon.

3. Das heterocyclische Derivat nach Anspruch 1, wobei R¹ Phenyl, 1,2,3,4-Tetrahydronaphthalen-5-yl, 1,2,3,4-Tetrahydronaphthalen-6-yl, 2,3-Dihydro-1H-inden-4-yl, 2,3-Dihydro-1H-inden-5-yl, 1,2-Dihydrocyclobutabenzol-3-yl oder 1,2-Dihydrocyclobutabenzol-4-yl ist und das Phenyl gegebenenfalls substituiert ist mit einer bis drei Gruppen, ausgewählt aus der Gruppe bestehend aus

- i) Halogen,
- ii) Alkyl, gegebenenfalls substituiert mit einem bis drei Halogenen,
- iii) Alkoxy und
- iv) Cyano,

oder sein Tautomer oder ein pharmazeutisch verträgliches Salz davon.

4. Das heterocyclische Derivat nach Anspruch 1, wobei R² Phenyl ist und das Phenyl gegebenenfalls mit einer bis drei Gruppen substituiert ist, ausgewählt aus der Gruppe bestehend aus

- i) Halogen,
- ii) Alkylsulfonyl,
- iii) Alkoxy, gegebenenfalls substituiert mit Alkoxy,
- iv) Alkynyl, gegebenenfalls substituiert mit Alkoxyalkyl oder Cycloalkyl und

v) Alkyl, gegebenenfalls substituiert mit einer bis drei Gruppen, ausgewählt aus der Gruppe bestehend aus Halogen, Alkoxy, Alkoxyalkoxy, Cycloalkyl und Phenyl,

oder sein Tautomer oder ein pharmazeutisch verträgliches Salz davon.

- 5
10
15
20
25
30
5. Das heterocyclische Derivat nach Anspruch 1, wobei
der Ring A eine Gruppe der Formel [4] ist,
X¹ NH ist,
A²

- i) Wasserstoff,
ii) Alkyl, gegebenenfalls substituiert mit einer Gruppe, ausgewählt aus der Gruppe bestehend aus Halogen, Monoalkylamino, Dialkylamino, Monoalkylaminocarbonyl, Dialkylaminocarbonyl, einem gesättigten cyclischen Aminocarbonyl, Alkoxy, Alkoxyalkoxy und Alkylcarbonyloxy,
iii) Cycloalkyl, gegebenenfalls substituiert mit Alkyl, gegebenenfalls substituiert mit einem bis drei Halogenen,
iv) Alkoxy,
v) eine gesättigte heterocyclische Gruppe, gegebenenfalls substituiert mit Alkyl oder Alkyloxycarbonyl,
vi) Alkylthio,
vii) Alkylsulfonyl,
viii) Alkylsulfinyl,
ix) Amino, substituiert mit Alkyl, wobei das Alkyl gegebenenfalls mit einer Gruppe substituiert ist, ausgewählt aus der Gruppe bestehend aus Monoalkylamino, Dialkylamino, einem gesättigten cyclischen Amino, gegebenenfalls substituiert mit Alkyl, Tetrahydrofuryl, Morpholino, Alkoxy, Hydroxycarbonyl, Hydroxyl und Alkylthio,
x) Amino, substituiert mit Cycloalkyl oder
xi) ein gesättigtes cyclisches Amino, gegebenenfalls substituiert mit Alkyl, Dialkylamino, Alkoxy oder Hydroxyl, ist, und

R¹

- i) Phenyl, gegebenenfalls substituiert mit einer bis drei Gruppen, ausgewählt aus der Gruppe bestehend aus Halogen, Alkyl, gegebenenfalls substituiert mit einem bis drei Halogenen, Alkoxy und Cyano,
ii) 1,2,3,4-Tetrahydronaphthalen-5-yl,
iii) 2,3-Dihydro-1H-inden-5-yl,
iv) Benzyl, gegebenenfalls substituiert mit Halogen oder Alkyl, gegebenenfalls substituiert mit einem oder drei Halogenen,
v) Cycloalkyl,
vi) Cycloalkylmethyl,
vii) Naphthyl,
viii) Pyridylmethyl, gegebenenfalls substituiert mit Alkyl, gegebenenfalls substituiert mit einem bis drei Halogenen,
ix) Thienyl,
x) Thienylmethyl,
xi) Benzothiazolyl,
xii) Benzothiadiazolyl,
xiii) Indolyl oder
xiv) Alkyl ist, und

R² Phenyl oder Pyridyl ist,

wobei das Phenyl gegebenenfalls mit einer bis drei Gruppen substituiert ist, ausgewählt aus der Gruppe bestehend aus

- i) Halogen,
ii) Alkylsulfonyl,
iii) Alkoxy, gegebenenfalls substituiert mit Alkoxy,
iv) Alkynyl, gegebenenfalls substituiert mit Alkoxyalkyl oder Cycloalkyl, und
v) Alkyl, gegebenenfalls substituiert mit einer bis drei Gruppen, ausgewählt aus der Gruppe bestehend aus Halogen, Alkoxy, Alkoxyalkoxy, Cycloalkyl und Phenyl

und das Pyridyl gegebenenfalls mit Halogen substituiert ist
oder sein Tautomer oder ein pharmazeutisch verträgliches Salz davon.

6. Das heterocyclische Derivat nach Anspruch 1, wobei

der Ring A eine Gruppe der Formel [4] ist,

X¹ NH ist,

A² Alkyl, substituiert mit Alkoxy, Dialkylamino, Tetrahydrofuryl, Tetrahydrofurylmethyl, Alkoxyalkylamino oder Cycloalkyl, gegebenenfalls substituiert mit Alkyl, gegebenenfalls substituiert mit einem bis drei Halogenen, ist R¹ Phenyl, substituiert mit einem Halogen und einem Methyl, ist und R² Phenyl, gegebenenfalls substituiert mit einem Trifluormethyl oder zwei Halogenen, ist,

oder sein Tautomer oder ein pharmazeutisch verträgliches Salz davon.

7. Das heterocyclische Derivat nach Anspruch 1, ausgewählt aus den folgenden (1)-(239), oder seine Tautomere oder ein pharmazeutisch verträgliches Salz davon:

(1) N-[2-(Trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1 H-benzimidazol-4-carboxamid,

(2) N-Cyclohexyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amine)-1H-benzimidazol-4-carboxamid,

(3) N-(3-Chlor-2-methylphenyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1 H-benzimidazol-4-carboxamid,

(4) N-[(1-Hydroxycyclohexyl)methyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(5) N-[2-(Trifluormethyl)benzyl]-5-({[2-(trifluormethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamid,

(6) N-Cyclohexyl-5-({[2-(trifluormethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamid,

(7) N-(3-Chlor-2-methylphenyl)-5-({[2-(trifluormethyl)phenyl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamid,

(8) N-Cyclohexyl-5-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-indazol-7-carboxamid,

(9) N-[2-(Trifluormethyl)benzyl]-5-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-indazol-7-carboxamid,

(10) N-(3-Chlor-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(11) 2-Methyl-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(12) N-Cyclohexyl-2-methyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(13) N-(3-Chlor-2-methylphenyl)-2-methyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(14) N-Cyclopentyl-2-methyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(15) N-Cyclobutyl-2-methyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(16) N-(3-Chlor-2-methylphenyl)-2-ethyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(17) N-Cyclohexyl-2-ethyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(18) 2-Ethyl-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(19) N-Cyclohexyl-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(20) 2-(Methoxymethyl)-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(21) 2-(Methoxymethyl)-N-(2-methylphenyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(22) 2-(Methoxymethyl)-N-(4-methylphenyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(23) N-(2-Chlorbenzyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(24) 2-(Methoxymethyl)-N-(4-methylbenzyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(25) N-(4,4-Difluorocyclohexyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(26) N-(4-tert-Buthylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(27) 2-(Methoxymethyl)-N-[4-(trifluormethyl)phenyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimi-

dazol-4-carboxamid,

(28) N-(2,4-Dimethylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(29) N-(2-Chlor-4-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(30) N-(3,4-Dimethylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(31) N-(3-Chlor-4-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(32) N-(2,3-Dihydro-1H-inden-5-yl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(33) 2-(Methoxymethyl)-N-(5,6,7,8-tetrahydronaphthalen-1-yl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(34) N-(2-Fluorphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(35) 2-(Methoxymethyl)-N-(2-methoxyphenyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(36) 2-(Methoxymethyl)-N-(4-methoxyphenyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(37) N-(3-Brom-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(38) N-(3-Chlor-2-methylbenzyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(39) N-(2,6-Difluorphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(40) N-(3-Cyano-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(41) 2-(Methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-N-{{[3-(trifluormethyl)pyridin-2-yl]methyl}-1H-benzimidazol-4-carboxamid,

(42) N-(2-Chlor-6-methylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(43) 2-(2-Amino-2-oxoethyl)-N-(3-chlor-2-methylphenyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(44) 2-(2-Amino-2-oxoethyl)-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(45) N-(3-Chlor-2-methylphenyl)-1-methyl-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(46) N-Cyclohexyl-1-methyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(47) 1-Methyl-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(48) N-(3-Chlor-2-methylphenyl)-1-ethyl-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(49) N-Cyclohexyl-1-ethyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(50) 1-Ethyl-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(51) N-(3-Chlor-2-methylphenyl)-2-methyl-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1,3-benzoxazol-4-carboxamid,

(52) 2-Methyl-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1,3-benzoxazol-4-carboxamid,

(53) N-(3-Chlor-2-methylphenyl)-2-ethyl-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1,3-benzoxazol-4-carboxamid,

(54) N-(3-Chlor-2-methylphenyl)-2-ethoxy-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(55) 2-Ethoxy-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(56) N-(3-Chlor-2-methylphenyl)-2-(1-chlor-2-methylpropan-2-yl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(57) N-(3-Chlor-2-methylphenyl)-2-[(dimethylamino)methyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-

benzimidazol-4-carboxamid,

(58) N-(3-Chlor-2-methylphenyl)-2-(2-methylpropyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(59) 2-(2-Methylpropyl)-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(60) tert-Butyl-3-{4-[(3-chlor-2-methylphenyl)carbamoyl]-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}azetidin-1-carboxylat,

(61) N-(3-Chlor-2-methylphenyl)-2-[(methylamino)methyl]-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(62) {4-[(3-Chlor-2-methylphenyl)carbamoyl]-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-2-yl}methylacetat,

(63) N-(3-Chlor-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(64) 2-[[2-(2R)-Tetrahydrofuran-2-yl]-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(65) N-(3-Chlor-2-methylphenyl)-2-[(2S)-tetrahydrofuran-2-yl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(66) 2-[[2-(2S)-Tetrahydrofuran-2-yl]-N-[2-(trifluormethyl)benzyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(67) 2-(1-Acetylazetidin-3-yl)-N-(3-chlor-2-methylphenyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(68) tert-Butyl-(2S)-2-{4-[(3-chlor-2-methylphenyl)carbamoyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}pyrrolidin-1-carboxylat,

(69) tert-Butyl-(2R)-2-{4-[(3-chlor-2-methylphenyl)carbamoyl]-6-({[2-(trifluoromethyl)phenyl]carbonyl}amino)-1H-benzimidazol-2-yl}pyrrolidin-1-carboxylat,

(70) N-(3-Chlor-2-methylphenyl)-2-[(2S)-pyrrolidin-2-yl]-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(71) N-(3-Chlor-2-methylphenyl)-2-[(2S)-1-methylpyrrolidin-2-yl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(72) 2-[(2S)-1-Acetylpyrrolidin-2-yl]-N-(3-chlor-2-methylphenyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(73) N-(3-Chlor-2-methylphenyl)-2-[(2-methoxyethoxy)methyl]-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(74) N-(3-Chlor-2-methylphenyl)-2-(1-methoxy-2-methylpropan-2-yl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(75) 2-tert-Butyl-N-(3-chlor-2-methylphenyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(76) 2-tert-Butyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-N-[[3-(trifluormethyl)pyridin-2-yl]methyl]-1H-benzimidazol-4-carboxamid,

(77) N-(3-Chlor-2-methylphenyl)-2-(2-ethoxyethyl)-6-({[2-(trifluoromethyl)phenyl] carbonyl}amino)-1 H-benzimidazol-4-carboxamid,

(78) N-(3-Chlor-2-methylphenyl)-2-(ethoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(79) 2-(Ethoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-N-[[3-(trifluormethyl)pyridin-2-yl]methyl]-1H-benzimidazol-4-carboxamid,

(80) N-(3-Chlor-2-methylphenyl)-2-(2-methoxyethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(81) N-(3-Chlor-2-methylphenyl)-2-(2,2-dimethylpropyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(82) N-(3-Chlor-2-methylphenyl)-2-cyclopropyl-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(83) N-(3-Chlor-2-methylphenyl)-2-(2-methylpentan-2-yl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(84) N-(3-Chlor-2-methylphenyl)-2-(1-methylcyclopropyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(85) 2-tert-Butyl-N-(3-chlor-4-methylphenyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(86) 2-tert-Butyl-N-(3-chlor-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl} amino)-1H-benzimidazol-4-carbo-

xamid,

(87) 2-tert-Butyl-N-(3-chlor-4-methylphenyl)-6-({(2,5-dichlorophenyl)carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(88) N-(3-Chlor-2-methylphenyl)-2-[1-(trifluormethyl)cyclopropyl]-6-({[2-(trifluormethyl)phenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(89) N-(3-Chlor-2-methylphenyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluormethyl)phenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(90) N-(2-Chlorbenzyl)-2-(methoxymethyl)-1-methyl-6-({[2-(trifluoromethyl) phenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(91) 6-({[2-Chlor-6-fluorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(92) 6-({[2-Chlor-4-fluorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-methoxymethyl-1H-benzimidazol-4-carboxamid,

(93) 6-({[2-Chlor-5-fluorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(94) N-(3-Chlor-2-methylphenyl)-6-({[2-chlorophenyl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(95) N-(3-Chlor-2-methylphenyl)-6-({[2-chlorpyridin-3-yl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(96) 6-({[2-Bromphenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(97) N-(3-Chlor-2-methylphenyl)-6-({[2,6-dichlorophenyl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(98) N-(3-Chlor-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(99) 6-({[2-Chlor-3-fluorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(100) 6-({[2-Chlor-3,6-difluorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl) -2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(101) 6-({[2-Brom-6-chlorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(102) 6-({[2-Brom-6-fluorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(103) N-(3-Chlor-2-methylphenyl)-6-({[2-chlor-6-methylphenyl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(104) N-(3-Chlor-2-methylphenyl)-6-({[2-chlor-4-methylphenyl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(105) 6-({[5-Brom-2-chlorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(106) 6-({[2-Brom-5-chlorophenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(107) N-(3-Chlor-2-methylphenyl)-6-({[2-chlor-5-methylphenyl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(108) N-(3-Chlor-2-methylphenyl)-2-(methoxymethyl)-6-({[5-methyl-2-(trifluormethyl)phenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(109) 6-({[2,5-bis(Trifluormethyl)phenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(110) 6-({[2,4-bis(Trifluormethyl)phenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(111) N-(3-Chlor-2-methylphenyl)-6-({[5-fluor-2-(trifluormethyl)phenyl] carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(112) N-(3-Chlor-2-methylphenyl)-6-({[2-chlor-6-(trifluormethyl)phenyl] carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(113) N-(3-Chlor-2-methylphenyl)-6-({[2-chlor-5-[2-(propan-2-yloxy)ethoxy] phenyl]carbonyl} amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(114) 6-({[2-Chlor-5-(2-ethoxyethoxy)phenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(115) 6-({[2-Chlor-5-(3-methoxypropyl)phenyl]carbonyl} amino)-N-(3-chlor-2-methylphenyl)-2-(methoxyme-

thyl)-1H-benzimidazol-4-carboxamid,
 (116) 6-([5-(3-tert-Butoxyprop-1-in-1-yl)-2-chlorophenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,
 (117) 6-([5-(3-tert-Butoxypropyl)-2-chlorophenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,
 5 (118) 6-([2-Chlor-5-(3-hydroxy-3-methylbutyl)phenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,
 (119) 6-([2-Chlor-5-(ethoxymethyl)phenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,
 10 (120) 6-([2-Chlor-5-[(2-ethoxyethoxy)methyl]phenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,
 (121) 6-([2-Chlor-5-(2-cyclopropylethyl)phenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,
 (122) N-(3-Chlor-2-methylphenyl)-6-([2-chlor-5-(2-phenylethyl)phenyl]carbonyl) amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,
 15 (123) N-(3-Chlor-2-methylphenyl)-2-cyclopentyl-6-([2-(trifluormethyl)phenyl] carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (124) N-(3-Chlor-2-methylphenyl)-2-cyclopentyl-6-([2,5-dichlorophenyl]carbonyl) amino)-1H-benzimidazol-4-carboxamid,
 20 (125) 6-([2-Chlor-6-fluorophenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-cyclopentyl-1H-benzimidazol-4-carboxamid,
 (126) 6-([2-Chlor-6-fluorophenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-yl]-1H-benzimidazol-4-carboxamid,
 (127) N-(3-Chlor-2-methylphenyl)-6-([2,6-dichlorophenyl]carbonyl)amino)-2-[(2R)-tetrahydrofuran-2-yl]-1 H-benzimidazol-4-carboxamid,
 25 (128) N-(3-Chlor-2-methylphenyl)-6-([2,5-dichlorophenyl]carbonyl)amino)-2-[(2R)-tetrahydrofuran-2-yl]-1 H-benzimidazol-4-carboxamid,
 (129) N-(3-Chlor-2-methylphenyl)-2-[(2S)-5-oxopyrrolidin-2-yl]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 30 (130) N-(3-Chlor-2-methylphenyl)-2-[(2R)-5-oxopyrrolidin-2-yl]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (131) N-(3-Chlor-2-methylphenyl)-2-[2-oxo-2-(pyrrolizin-1-yl)ethyl]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (132) N-(3-Chlor-2-methylphenyl)-2-[2-(dimethylamino)-2-oxoethyl]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 35 (133) N-(3-Chlor-2-methylphenyl)-2-[2-(methylamino)-2-oxoethyl]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (134) 2-Chlor-N-(3-chlor-2-methylphenyl)-6-([2-(trifluormethyl)phenyl]carbonyl) amino)-1H-benzimidazol-4-carboxamid,
 40 (135) N-(3-Chlor-2-methylphenyl)-2-[(2-methoxyethyl)amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (136) N-(3-Chlor-2-methylphenyl)-2-[(2-hydroxyethyl)amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (137) N-(3-Chlor-2-methylphenyl)-2-(methylamino)-6-([2-(trifluoromethyl)phenyl] carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 45 (138) N-(3-Chlor-2-methylphenyl)-2-(ethylamino)-6-([2-(trifluoromethyl)phenyl] carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (139) N-(3-Chlor-2-methylphenyl)-2-[(2,2-dimethylpropyl)amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 50 (140) N-(3-Chlor-2-methylphenyl)-2-(cyclopentylamino)-6-([2-(trifluormethyl) phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (141) N-(3-Chlor-2-methylphenyl)-2-(piperidin-1-yl)-6-([2-(trifluormethyl) phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (142) N-(3-Chlor-2-methylphenyl)-2-(4-methylpiperazin-1-yl)-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 55 (143) 2-bis(2-Hydroxyethyl)amino)-N-(3-chlor-2-methylphenyl)-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,
 (144) N-(3-Chlor-2-methylphenyl)-2-(dimethylamino)-6-([2-(trifluormethyl) phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

midazol-4-carboxamid,

(145) N-(3-Chlor-2-methylphenyl)-2-[[2-(morpholin-4-yl)ethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(146) N-(3-Chlor-2-methylphenyl)-2-[[2-(dimethylamino)ethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(147) N-(3-Chlor-2-methylphenyl)-2-(3-hydroxyazetidin-1-yl)-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(148) N-(3-Chlor-2-methylphenyl)-2-[(3S)-3-(dimethylamino)pyrrolizin-1-yl]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(149) N-(3-Chlor-2-methylphenyl)-2-[(3S)-3-hydroxypyrrolizin-1-yl]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(150) N-(3-Chlor-2-methylphenyl)-2-[[2-(diethyl amino)ethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(151) N-(3-Chlor-2-methylphenyl)-2-[[2-(pyrrolizin-1-yl)ethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(152) N-(3-Chlor-2-methylphenyl)-2-[[3-(dimethylamino)propyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(153) N-(3-Chlor-2-methylphenyl)-2-[[3-(dimethylamino)-2,2-dimethylpropyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(154) N-(3-Chlor-2-methylphenyl)-2-[2-(dipropan-2-ylamino)ethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(155) N-(3-Chlor-2-methylphenyl)-2-(morpholin-4-yl)-6-([2-(trifluoromethyl) phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(156) 2-Amino-N-(3-chlor-2-methylphenyl)-6-([2-(trifluormethyl)phenyl] carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(157) N-(3-Chlor-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(158) N-(3-Chlor-2-methylphenyl)-2-[[3-methyloxetan-3-yl)methyl]amino]-6-([2-(trifluormethyl)phenyl] carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(159) tert-Butyl-N-{4-[(3-chlor-2-methylphenyl)carbamoyl]-6-([2-(trifluormethyl) phenyl]carbonyl)amino)-1H-benzimidazol-2-yl}glycinat,

(160) N-{4-[(3-chlor-2-methylphenyl)carbamoyl]-6-([2-(trifluormethyl)phenyl] carbonyl)amino)-1H-benzimidazol-2-yl}glycine,

(161) N-(3-Chlor-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1-methyl-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(162) N-(3-Chlor-2-methylphenyl)-2-[(3-methoxy-2,2-dimethylpropyl)amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amine)-1H-benzimidazol-4-carboxamid,

(163) N-(3-Chlor-2-methylphenyl)-2-(pyrrolizin-1-yl)-6-([2-(trifluormethyl)phenyl] carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(164) 2-(Azetidin-1-yl)-N-(3-chlor-2-methylphenyl)-6-([2-(trifluormethyl)phenyl] carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(165) N-(3-Chlor-2-methylphenyl)-2-(3-methoxyazetidin-1-yl)-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(166) N-(3-Chlor-2-methylphenyl)-2-[(2-hydroxy-2-methylpropyl)amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(167) N-(3-Chlor-2-methylphenyl)-2-[(2S)-tetrahydrofuran-2-ylmethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(168) N-(3-Chlor-2-methylphenyl)-2-[(2R)-tetrahydrofuran-2-ylmethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(169) N-(3-Chlor-2-methylphenyl)-2-[(2S)-1-hydroxy-3-methylbutan-2-yl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(170) N-(3-Chlor-2-methylphenyl)-2-[(2R)-1-hydroxy-3-methylbutan-2-yl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(171) N-(3-Chlor-2-methylphenyl)-2-[(2S)-1-hydroxy-3,3-dimethylbutan-2-yl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(172) N-(3-Chlor-2-methylphenyl)-2-[(3-methoxy-2,2-dimethylpropyl)(methyl amino)-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(173) N-(3-Chlor-2-methylphenyl)-2-[(3-methoxypropyl)amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-

1H-benzimidazol-4-carboxamid,

(174) N-(3-Chlor-2-methylphenyl)-2-[[2-(propan-2-yloxy)ethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(175) 2-[(2-tert-Butoxyethyl)amino]-N-(3-chlor-2-methylphenyl)-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(176) N-(3-Chlor-2-methylphenyl)-2-[(2-methoxy-2-methylpropyl)amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(177) N-(3-Chlor-2-methylphenyl)-2-[[2-(methylsulfanyl)ethyl]amino]-6-([2-(trifluormethyl)phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(178) N-(3-Chlor-2-methylphenyl)-2-(methylsulfanyl)-6-([2-(trifluormethyl)phenyl] carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(179) N-(3-Chlor-2-methylphenyl)-2-(methylsulfonyl)-6-([2-(trifluormethyl) phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(180) N-(3-Chlor-2-methylphenyl)-2-(methylsulfiny)-6-([2-(trifluormethyl) phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(181) 6-[[2-Chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-(dimethylamino)-1 H-benzimidazol-4-carboxamid,

(182) N-(3-Chlor-2-methylphenyl)-6-[[2,6-dichlorophenyl]carbonyl]amino)-2-(dimethylamino)-1H-benzimidazol-4-carboxamid,

(183) N-(3-Chlor-2-methylphenyl)-6-[[2,4-dichlorophenyl]carbonyl]amino)-2-(dimethylamino)-1H-benzimidazol-4-carboxamid,

(184) N-(3-Chlor-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino -2-(dimethylamino)-1 H-benzimidazol-4-carboxamid,

(185) 6-[[2-Brom-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazol-4-carboxamid,

(186) 6-[[2-Brom-6-chlorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazol-4-carboxamid,

(187) 6-([2-Chlor-5-(cyclopropylethynyl)phenyl]carbonyl)amino)-N-(3-chlor-2-methylphenyl)-2-(dimethylamino)-1H-benzimidazol-4-carboxamid,

(188) N-(3-Chlor-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1H-benzimidazol-4-carboxamid,

(189) N-(3-Chlor-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-2-[(3-methoxy-2,2-dimethylpropyl)amino]-1H-benzimidazol-4-carboxamid,

(190) N-(3-Chlor-2-methylphenyl)-6-[[2, 5-dichlorophenyl]carbonyl]amino)-2-[(2-hydroxy-2-methylpropyl)amino]-1H-benzimidazol-4-carboxamid,

(191) N-(3-Chlor-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-2-[(2-methoxy-2-methylpropyl)amino]-1H-benzimidazol-4-carboxamid,

(192) N-(3-Chlor-2-methylphenyl)-6-[[2,5-dichlorophenyl]carbonyl]amino)-2-[[2-(propan-2-yloxy)ethyl]amino)-1H-benzimidazol-4-carboxamid,

(193) 6-[[2-Chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-[[2-(propan-2-yloxy)ethyl]amino)-1H-benzimidazol-4-carboxamid,

(194) 2-[(2-tert-Butoxyethyl)amino]-6-[[2-chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-1H-benzimidazol-4-carboxamid,

(195) 6-[[2-Chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-[(3-methoxy-2,2-dimethylpropyl)amino]-1H-benzimidazol-4-carboxamid,

(196) 6-[[2-Chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-[(2-methoxy-2-methylpropyl)amino]-1H-benzimidazol-4-carboxamid,

(197) 6-[[2-Chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-[[2S]-tetrahydrofuran-2-ylmethyl]amino)-1H-benzimidazol-4-carboxamid,

(198) 6-[[2-Chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-[[2R]-tetrahydrofuran-2-ylmethyl]amino)-1H-benzimidazol-4-carboxamid,

(199) 6-[[2-Chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-[(3-hydroxy-2,2-dimethylpropyl)amino]-1H-benzimidazol-4-carboxamid,

(200) 6-[[2-Chlor-6-fluorophenyl]carbonyl]amino)-N-(3-chlor-2-methylphenyl)-2-[[2S]-1-hydroxy-3-methylbutan-2-yl]amino)-1H-benzimidazol-4-carboxamid,

(201) N-(3-Chlor-4-methylphenyl)-2-(dimethylamino)-6-([2-(trifluormethyl) phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

(202) N-(4-tert-Buthylphenyl)-2-(dimethylamino)-6-([2-(trifluormethyl) phenyl]carbonyl)amino)-1H-benzimidazol-4-carboxamid,

zol-4-carboxamid,

(203) N-(2,3-Dihydro-1H-inden-5-yl)-2-(dimethylamino)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(204) 6-({[2-Chlor-6-fluorophenyl]carbonyl}amino)-N-(3-chlor-4-methylphenyl)-2-(dimethylamino)-1 H-benzimidazol-4-carboxamid,

(205) N-(3-Chlor-4-methylphenyl)-6-({[2,6-dichlorophenyl]carbonyl}amino)-2-(dimethylamino)-1 H-benzimidazol-4-carboxamid,

(206) N-(3-Chlor-4-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(dimethylamino)-1 H-benzimidazol-4-carboxamid,

(207) N-(3-Chlor-2-methylphenyl)-2-cyclopropyl-6-({[2,5-dichlorophenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(208) N-(3-Chlor-4-methylphenyl)-2-cyclopropyl-6-({[2,5-dichlorophenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(209) N-(3-Chlor-2-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(1-methylcyclopropyl)-1H-benzimidazol-4-carboxamid,

(210) N-(3-Chlor-4-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(1-methylcyclopropyl)-1H-benzimidazol-4-carboxamid,

(211) N-(3-Chlor-2-methylphenyl)-2-(methoxymethyl)-6-({[2-(methylsulfonyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(212) N-(3-Chlor-4-methylphenyl)-6-({[2,5-dichlorophenyl]carbonyl}amino)-2-(2-methoxyethyl)-1H-benzimidazol-4-carboxamid,

(213) 2-(Methoxymethyl)-N-phenyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1 H-benzimidazol-4-carboxamid,

(214) 2-(Methoxymethyl)-N-propyl-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(215) 2-(Methoxymethyl)-N-(pyridin-3-yl)-6-({[2-(trifluormethyl)phenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(216) N-Benzyl-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(217) N-(Cyclohexylmethyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(218) 2-(Methoxymethyl)-N-(naphthalen-1-yl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(219) 2-(Methoxymethyl)-N-(thiophen-3-yl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(220) N-(2,1,3-Benzothiadiazol-4-yl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(221) N-(1,1-Dioxid-1-benzothiophen-6-yl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(222) 2-(Methoxymethyl)-N-(thiophen-2-ylmethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(223) N-(1H-Indol-5-yl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(224) N-(1,3-Benzothiazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(225) N-(2,2-Dimethylpropyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(226) 2-(Methoxymethyl)-N-(thiophen-2-yl)-6-({[2-(trifluormethyl)phenyl]carbonyl} amino)-1H-benzimidazol-4-carboxamid,

(227) N-(5-Chlor-1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(228) N-(2-Benzylphenyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(229) 2-(Methoxymethyl)-N-(chinolin-8-yl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(230) N-(Cycloheptylmethyl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(231) N-(1,3-Benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl] carbonyl}amino)-1H-benzimi-

dazol-4-carboxamid,

(232) N-(6-Chlor-1,3-benzoxazol-2-yl)-2-(methoxymethyl)-6-({[2-(trifluormethyl) phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(233) N-[3-Chlor-2-(hydroxymethyl)phenyl]-2-(methoxymethyl)-6-({[2-(trifluormethyl)phenyl]carbonyl}amino)-1H-benzimidazol-4-carboxamid,

(234) N-(3-Chlor-2-methylphenyl)-6-({[3-fluorpyridin-2-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(235) N-(3-Chlor-2-methylphenyl)-6-({[3-chlorpyridin-4-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(236) N-(3-Chlor-2-methylphenyl)-6-({[3,5-dichlorpyridin-4-yl]carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(237) 6-({[5-Butoxy-2-chlorphenyl]carbonyl}amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(238) 6-({[2-Chlor-5-(2,2-difluorethoxy)phenyl]carbonyl}amino)-N-(3-chlor-2-methylphenyl)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid,

(239) N-(3-Chlor-2-methylphenyl)-6-({[2-chlor-5-(4,4,4-trifluorbutoxy)phenyl] carbonyl}amino)-2-(methoxymethyl)-1H-benzimidazol-4-carboxamid.

8. Ein Arzneimittel, enthaltend eine Verbindung, ausgewählt aus der Gruppe bestehend aus dem heterocyclischen Derivat nach einem der Ansprüche 1 bis 7, seinem Tautomer und einem pharmazeutisch verträglichen Salz davon als Wirkstoff.

9. Eine Verbindung, ausgewählt aus der Gruppe bestehend aus dem heterocyclischen Derivat nach einem der Ansprüche 1 bis 7, seinem Tautomer und einem pharmazeutisch verträglichen Salz davon zur Verwendung als ein mPGES-Inhibitor.

10. Eine Verbindung, ausgewählt aus der Gruppe bestehend aus dem heterocyclischen Derivat nach einem der Ansprüche 1 bis 7, seinem Tautomer und einem pharmazeutisch verträglichen Salz davon, zur Verwendung bei der Vorbeugung oder Behandlung von entzündlicher Darmerkrankung, Reizdarmsyndrom, Migräne, Kopfschmerzen, Lumbago, Spinalstenose, Bandscheibenvorfall, Kraniomandibulären Dysfunktionen, Zervikalsyndrom, Zervikalspondylose, Endometriose, Adenomyose, vorzeitiger Wehentätigkeit und Entbindung, drohender Frühgeburt, Dysmenorrhö, hyperaktiver Blase, Nykturie, interstitieller Zystitis, neurodegenerativer Erkrankung, Psoriasis, rheumatoider Arthritis, rheumatischem Fieber, Fibromyalgie, Neuralgie, komplexem regionalem Schmerzsyndrom, fasziärer Dysfunktion, Virusinfektionen, bakterieller Infektion, Mykose, Verbrennung, Entzündung und Schmerz nach einer Operation, Verletzung und Zahnextraktion, bösartigen Tumoren, Atherosklerose, Schlaganfall, Gicht, Arthritis, Osteoarthritis, juveniler Arthritis, Spondylitis ankylosans, Sehnenscheidenentzündung, Bänderverknöcherung, systemischem Lupus erythematoses, Vaskulitis, Pancreatitis, Nephritis, Konjunktivitis, Iritis, Skleritis, Uveitis, Wundtherapie, Dermatitis, Ekzem, Osteoporose, Asthma, chronisch obstruktiver Lungenerkrankung, Lungenfibrose, allergischer Erkrankung, Familiärer adenomatöser Polyposis, Skleroderma, Bursitis, Leiomyom des Uterus oder eines Schmerzes aufgrund von Krebs.


11. Die Verbindung zur Verwendung nach Anspruch 10, wobei die neurodegenerative Erkrankung Alzheimer-Krankheit oder multiple Sklerose ist.

12. Die Verbindung zur Verwendung nach Anspruch 10, wobei die Virusinfektionskrankheit Influenza, Erkältung, Zoster, oder AIDS ist.




13. Die Verbindung zur Verwendung nach Anspruch 10, wobei der bösartige Tumor Dickdarmkrebs, Brustkrebs, Lungenkrebs oder Prostatakrebs ist.

Revendications

1. Dérivé hétérocyclique représenté par la formule générale [1] :



ou son tautomère ou un sel pharmaceutiquement acceptable de celui-ci, dans laquelle le cycle A est un groupe représenté par la formule générale [2], [3] ou [4] :



dans lesquelles

X¹ est NH, N-alkyle ou O ;

A¹ est un atome d'hydrogène ou un groupe alkyle ;

A² est

i) un atome d'hydrogène ;

ii) un atome d'halogène ;

iii) un groupe alkyle facultativement substitué par un à trois groupes choisis dans le groupe constitué par un atome d'halogène, un groupe amino, un groupe monoalkylamino, un groupe dialkylamino, un groupe carbamoyle, un groupe monoalkylaminocarbonyle, un groupe dialkylaminocarbonyle, un groupe aminocarbonyle cyclique saturé, un groupe alcoxy, un groupe alcoxyalcoxy et un groupe alkylcarbonyloxy ;

iv) un groupe cycloalkyle facultativement substitué par un groupe alkyle facultativement substitué par un à trois atomes d'halogène ;

v) un groupe alcoxy ;


vi) un groupe hétérocyclique saturé facultativement substitué par un groupe alkyle, un groupe alkyloxycarbonyle, un groupe alkylcarbonyle ou un groupe oxo ;

vii) un groupe alkylthio ;

viii) un groupe alkylsulfonyl ;

ix) un groupe alkylsulfinyl ;

x) un groupe de la formule générale [5] :



dans laquelle R³ et R⁴ sont des groupes identiques ou différents choisis parmi

- a) un atome d'hydrogène,
- b) un groupe alkyle facultativement substitué par un groupe choisi dans le groupe constitué par un groupe monoalkylamino, un groupe dialkylamino, un groupe amino cyclique saturé facultativement substitué par un groupe alkyle, un groupe hétérocycle saturé facultativement substitué par un groupe alkyle, un groupe alcoxy, un groupe hydroxycarbonyl, un groupe hydroxyle, un groupe alkyloxycarbonyl et un groupe alkylthio, ou
- c) un groupe cycloalkyle ; ou

xi) un groupe amino cyclique saturé facultativement substitué par un groupe alkyle, un groupe amino, un groupe monoalkylamino, un groupe dialkylamino, un groupe alcoxy ou un groupe hydroxyle ;

R¹ est un groupe phényle, un groupe benzyle, un groupe naphthyle, un groupe cycloalkyle, un groupe cycloalkylméthyle, un groupe hétéroaryle, un groupe hétéroarylméthyle, un groupe 1,2,3,4-tétrahydronaphtalén-5-yle, un groupe 1,2,3,4-tétrahydronaphtalén-6-yle, un groupe 2,3-dihydro-1H-indén-4-yle, un groupe 2,3-dihydro-1H-indén-5-yle, un groupe 1,2-dihydrocyclobutabenzén-3-yle, un groupe 1,2-dihydrocyclobutabenzén-4-yle ou un groupe alkyle, où lesdits groupes phényle, benzyle, cycloalkyle, cycloalkylméthyle, hétéroaryle et hétéroarylméthyle sont facultativement substitués par un à trois groupes choisis dans le groupe constitué par

- i) un atome d'halogène,
- ii) un groupe alkyle facultativement substitué par un à trois groupes choisis dans le groupe constitué par un atome d'halogène, un groupe hydroxy et un groupe phényle,
- iii) un groupe alcoxy,
- iv) un groupe hydroxy, et
- v) un groupe cyano ;

R² est un groupe phényle ou un groupe pyridyle, où lesdits groupes phényle et pyridyle sont facultativement substitués par un à trois groupes choisis dans le groupe constitué par

- i) un atome d'halogène,
- ii) un groupe alkylsulfonyl,
- iii) un groupe alcoxy facultativement substitué par un à trois atomes d'halogène ou groupes alcoxy ;
- iv) un groupe alcynyle facultativement substitué par un groupe alcoxyalkyle ou un groupe cycloalkyle, et
- v) un groupe alkyle facultativement substitué par un à trois groupes choisis dans le groupe constitué par un groupe alcoxy, un groupe alcoxylalcoxy, un groupe cycloalkyle, un groupe phényle et un atome d'halogène.

2. Dérivé hétérocyclique selon la revendication 1, dans lequel le cycle A est un groupe de formule [4] et X¹ est NH, ou son tautomère ou un sel pharmaceutiquement acceptable de celui-ci.

3. Dérivé hétérocyclique selon la revendication 1, dans lequel R¹ est un groupe phényle, un groupe 1,2,3,4-tétrahydronaphtalén-5-yle, un groupe 1,2,3,4-tétrahydronaphtalén-6-yle, un groupe 2,3-dihydro-1H-indén-4-yle, un groupe 2,3-dihydro-1H-indén-5-yle, un groupe 1,2-dihydrocyclobutabenzén-3-yle, ou un groupe 1,2-dihydrocyclobutabenzén-4-yle, et ledit groupe phényle est facultativement substitué par un à trois groupes choisis dans le groupe constitué par

- i) un atome d'halogène,
- ii) un groupe alkyle facultativement substitué par un à trois atomes d'halogène,
- iii) un groupe alcoxy, et
- iv) un groupe cyano,

ou son tautomère ou un sel pharmaceutiquement acceptable de celui-ci.

4. Dérivé hétérocyclique selon la revendication 1, dans lequel R² est un groupe phényle et ledit groupe phényle est facultativement substitué par un à trois groupes choisis dans le groupe constitué par

- i) un atome d'halogène,

- ii) un groupe alkylsulfonyle,
- iii) un groupe alcoxy facultativement substitué par un groupe alcoxy,
- iv) un groupe alcynyle facultativement substitué par un groupe alcoxyalkyle ou un groupe cycloalkyle, et
- v) un groupe alkyle facultativement substitué par un à trois groupes choisis dans le groupe constitué par un atome d'halogène, un groupe alcoxy, un groupe alcoxyalcoxy, un groupe cycloalkyle et un groupe phényle,

ou son tautomère ou un sel pharmaceutiquement acceptable de celui-ci.

5. Dérivé hétérocyclique selon la revendication 1, dans lequel le cycle A est un groupe de formule [4],

X¹ est NH,

A² est

- i) un atome d'hydrogène,
- ii) un groupe alkyle facultativement substitué par un groupe choisi dans le groupe constitué par un atome d'halogène, un groupe monoalkylamino, un groupe dialkylamino, un groupe monoalkylaminocarbonyle, un groupe dialkylaminocarbonyle, un groupe aminocarbonyle cyclique saturé, un groupe alcoxy, un groupe alcoxyalcoxy et un groupe alkylcarbonyloxy,
- iii) un groupe cycloalkyle facultativement substitué par un groupe alkyle facultativement substitué par un à trois atomes d'halogène,
- iv) un groupe alcoxy,
- v) un groupe hétérocyclique saturé facultativement substitué par un groupe alkyle ou un groupe alkyloxycarbonyle,
- vi) un groupe alkylthio,
- vii) un groupe alkylsulfonyle,
- viii) un groupe alkylsulfinyle,
- ix) un groupe amino substitué par un groupe alkyle où ledit groupe alkyle est facultativement substitué par un groupe choisi dans le groupe constitué par un groupe monoalkylamino, un groupe dialkylamino, un groupe amino cyclique saturé facultativement substitué par un groupe alkyle, un groupe tétrahydrofuryle, un groupe morpholino, un groupe alcoxy, un groupe hydroxycarbonyle, un groupe hydroxyle et un groupe alkylthio,
- x) un groupe amino substitué par un groupe cycloalkyle ou
- xi) un groupe amino cyclique saturé facultativement substitué par un groupe alkyle, un groupe dialkylamino, un groupe alcoxy ou un groupe hydroxy,

R¹ est

- i) un groupe phényle facultativement substitué par un à trois groupes choisis dans le groupe constitué par un atome d'halogène, un groupe alkyle facultativement substitué par un à trois atomes d'halogène, un groupe alcoxy et un groupe cyano,
- ii) un groupe 1,2,3,4-tétrahydronaphtalén-5-yle,
- iii) un groupe 2,3-dihydro-1H-indén-5-yle,
- iv) un groupe benzyle facultativement substitué par un atome d'halogène ou un groupe alkyle facultativement substitué par un à trois atomes d'halogène,
- v) un groupe cycloalkyle,
- vi) un groupe cycloalkylméthyle,
- vii) un groupe naphthyle,
- viii) un groupe pyridylméthyle facultativement substitué par un groupe alkyle facultativement substitué une à trois atomes d'halogène,
- ix) un groupe thiényl,
- x) un groupe thiénylméthyle,
- xi) un groupe benzothiazolyle,
- xii) un groupe benzothiadiazolyle,
- xiii) un groupe indolyle ou
- xiv) un groupe alkyle, et

R² est un groupe phényle ou un groupe pyridyle,

où ledit groupe phényle est facultativement substitué par un à trois groupes choisis dans le groupe constitué par

- i) un atome d'halogène,

- ii) un groupe alkylsulfonyl,
- iii) un groupe alcoxy facultativement substitué par un groupe alcoxy,
- iv) un groupe alcynyle facultativement substitué par un groupe alcoxyalkyle ou un groupe cycloalkyle, et
- v) un groupe alkyle facultativement substitué par un à trois groupes choisis dans le groupe constitué par un atome d'halogène, un groupe alcoxy, un groupe alcoxyalcoxy, un groupe cycloalkyle et un groupe phényle, et

ledit groupe pyridyle est facultativement substitué par un atome d'halogène, ou son tautomère ou un sel pharmaceutiquement acceptable de celui-ci.

6. Dérivé hétérocyclique selon la revendication 1, dans lequel le cycle A est un groupe de formule [4],
 X¹ est NH,
 A² est un groupe alkyle substitué par un groupe alcoxy, un groupe dialkylamino, un groupe tétrahydrofuryle, un groupe tétrahydrofurylméthyle, un groupe alcoxyalkylamino, ou un groupe cycloalkyle facultativement substitué par un groupe alkyle facultativement substitué par un à trois atomes d'halogène,
 R¹ est un groupe phényle substitué par un atome d'halogène et un groupe méthyle, et
 R² est un groupe phényle facultativement substitué par un groupe trifluorométhyle ou deux atomes d'halogène, ou son tautomère ou un sel pharmaceutiquement acceptable de celui-ci.
7. Dérivé hétérocyclique selon la revendication 1 choisi parmi les composés (1) à (239) suivants ou son tautomère ou un sel pharmaceutiquement acceptable de celui-ci :

- (1) N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1 H-benzimidazole-4-carboxamide,
- (2) N-cyclohexyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (3) N-(3-chloro-2-méthylphényl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (4) N-[(1-hydroxycyclohexyl)méthyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (5) N-[2-(trifluorométhyl)benzyl]-5-({[2-(trifluorométhyl)phényl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (6) N-cyclohexyl-5-({[2-(trifluorométhyl)phényl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (7) N-(3-chloro-2-méthylphényl)-5-({[2-(trifluorométhyl)phényl]carbonyl}amino)-2,3-dihydro-1-benzofuran-7-carboxamide,
- (8) N-cyclohexyl-5-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-indazole-7-carboxamide,
- (9) N-[2-(trifluorométhyl)benzyl]-5-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1 H-indazole-7-carboxamide,
- (10) N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (11) 2-méthyl-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (12) N-cyclohexyl-2-méthyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (13) N-(3-chloro-2-méthylphényl)-2-méthyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (14) N-cyclopentyl-2-méthyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (15) N-cyclobutyle-2-méthyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (16) N-(3-chloro-2-méthylphényl)-2-éthyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (17) N-cyclohexyl-2-éthyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (18) 2-éthyl-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (19) N-cyclohexyl-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (20) 2-(méthoxyméthyl)-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (21) 2-(méthoxyméthyl)-N-(2-méthylphényl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (22) 2-(méthoxyméthyl)-N-(4-méthylphényl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,
- (23) N-(2-chlorobenzyl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-

4-carboxamide,

(24) 2-(méthoxyméthyl)-N-(4-méthylbenzyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(25) N-(4,4-difluorocyclohexyl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(26) N-(4-tert-butylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(27) 2-(méthoxyméthyl)-N-[4-(trifluorométhyl)phényl]-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(28) N-(2,4-diméthylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(29) N-(2-chloro-4-méthylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(30) N-(3,4-diméthylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(31) N-(3-chloro-4-méthylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(32) N-(2,3-dihydro-1H-indén-5-yl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(33) 2-(méthoxyméthyl)-N-(5,6,7,8-tétrahydronaphthalén-1-yl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(34) N-(2-fluorophényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(35) 2-(méthoxyméthyl)-N-(2-méthoxyphényl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(36) 2-(méthoxyméthyl)-N-(4-méthoxyphényl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(37) N-(3-bromo-2-méthylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(38) N-(3-chloro-2-méthylbenzyl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(39) N-(2,6-difluorophényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(40) N-(3-cyano-2-méthylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(41) 2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-N-([3-(trifluorométhyl)pyridin-2-yl]méthyl)-1H-benzimidazole-4-carboxamide,

(42) N-(2-chloro-6-méthylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(43) 2-(2-amino-2-oxoéthyl)-N-(3-chloro-2-méthylphényl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(44) 2-(2-amino-2-oxoéthyl)-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(45) N-(3-chloro-2-méthylphényl)-1-méthyl-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(46) N-cyclohexyl-1-méthyl-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(47) 1-méthyl-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(48) N-(3-chloro-2-méthylphényl)-1-éthyl-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(49) N-cyclohexyl-1-éthyl-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(50) 1-éthyl-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(51) N-(3-chloro-2-méthylphényl)-2-méthyl-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1,3-benzoxazole-4-carboxamide,

(52) 2-méthyl-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1,3-benzoxazole-4-carboxamide,

(53) N-(3-chloro-2-méthylphényl)-2-éthyl-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1,3-benzoxazole-4-

carboxamide,

(54) N-(3-chloro-2-méthylphényl)-2-éthoxy-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(55) 2-éthoxy-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(56) N-(3-chloro-2-méthylphényl)-2-(1-chloro-2-méthylpropan-2-yl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(57) N-(3-chloro-2-méthylphényl)-2-[(diméthylamino)méthyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(58) N-(3-chloro-2-méthylphényl)-2-(2-méthylpropyl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(59) 2-(2-méthylpropyl)-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(60) 3-{4-[(3-chloro-2-méthylphényl)carbamoyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazol-2-yl}azétidine-1-carboxylate de tert-butyle,

(61) N-(3-chloro-2-méthylphényl)-2-[(méthylamino)méthyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(62) {4-[(3-chloro-2-méthylphényl)carbamoyl]-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazol-2-yl}méthyl acétate,

(63) N-(3-chloro-2-méthylphényl)-2-[(2R)-tétrahydrofuran-2-yl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(64) 2-[[{(2R)-tétrahydrofuran-2-yl]-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(65) N-(3-chloro-2-méthylphényl)-2-[(2S)-tétrahydrofuran-2-yl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(66) 2-[[{(2S)-tétrahydrofuran-2-yl]-N-[2-(trifluorométhyl)benzyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(67) 2-(1-acétylazétidin-3-yl)-N-(3-chloro-2-méthylphényl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(68) (2S)-2-{4-[(3-chloro-2-méthylphényl)carbamoyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazol-2-yl}pyrrolidine-1-carboxylate de tert-butyle,

(69) (2R)-2-{4-[(3-chloro-2-méthylphényl)carbamoyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazol-2-yl}pyrrolidine-1-carboxylate de tert-butyle,

(70) N-(3-chloro-2-méthylphényl)-2-[(2S)-pyrrolidin-2-yl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(71) N-(3-chloro-2-méthylphényl)-2-[(2S)-1-méthylpyrrolidin-2-yl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(72) 2-[(2S)-1-acétylpyrrolidin-2-yl]-N-(3-chloro-2-méthylphényl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(73) N-(3-chloro-2-méthylphényl)-2-[(2-méthoxyéthoxy)méthyl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(74) N-(3-chloro-2-méthylphényl)-2-(1-méthoxy-2-méthylpropan-2-yl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(75) 2-tert-butyl-N-(3-chloro-2-méthylphényl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(76) 2-tert-butyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-N-[[3-(trifluorométhyl)pyridin-2-yl]méthyl]-1H-benzimidazole-4-carboxamide,

(77) N-(3-chloro-2-méthylphényl)-2-(2-éthoxyéthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(78) N-(3-chloro-2-méthylphényl)-2-(éthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(79) 2-(éthoxyméthyl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-N-[[3-(trifluorométhyl)pyridin-2-yl]méthyl]-1H-benzimidazole-4-carboxamide,

(80) N-(3-chloro-2-méthylphényl)-2-(2-méthoxyéthyl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(81) N-(3-chloro-2-méthylphényl)-2-(2,2-diméthylpropyl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(82) N-(3-chloro-2-méthylphényl)-2-cyclopropyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

zole-4-carboxamide,

(83) N-(3-chloro-2-méthylphényl)-2-(2-méthylpentan-2-yl)-6-([2-(trifluorométhyl) phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(84) N-(3-chloro-2-méthylphényl)-2-(1-méthylcyclopropyl)-6-([2-(trifluorométhyl) phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(85) 2-tert-butyl-N-(3-chloro-4-méthylphényl)-6-([2-(trifluorométhyl) phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(86) 2-tert-butyl-N-(3-chloro-2-méthylphényl)-6-([2,5-dichlorophényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(87) 2-tert-butyl-N-(3-chloro-4-méthylphényl)-6-([2,5-dichlorophényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(88) N-(3-chloro-2-méthylphényl)-2-[1-(trifluorométhyl)cyclopropyl]-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(89) N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1-méthyl-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(90) N-(2-chlorobenzyl)-2-(méthoxyméthyl)-1-méthyl-6-([2-(trifluorométhyl) phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(91) 6-([2-chloro-6-fluorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1 H-benzimidazole-4-carboxamide,

(92) 6-([2-chloro-4-fluorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-méthoxyméthyl-1H-benzimidazole-4-carboxamide,

(93) 6-([2-chloro-5-fluorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(94) N-(3-chloro-2-méthylphényl)-6-([2-chlorophényl]carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(95) N-(3-chloro-2-méthylphényl)-6-([2-chloropyridin-3-yl]carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(96) 6-([2-bromophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1 H-benzimidazole-4-carboxamide,

(97) N-(3-chloro-2-méthylphényl)-6-([2,6-dichlorophényl]carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(98) N-(3-chloro-2-méthylphényl)-6-([2,5-dichlorophényl]carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(99) 6-([2-chloro-3-fluorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(100) 6-([2-chloro-3,6-difluorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl) -2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(101) 6-([2-bromo-6-chlorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(102) 6-([2-bromo-6-fluorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(103) N-(3-chloro-2-méthylphényl)-6-([2-chloro-6-méthylphényl]carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(104) N-(3-chloro-2-méthylphényl)-6-([2-chloro-4-méthylphényl]carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(105) 6-([5-bromo-2-chlorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(106) 6-([2-bromo-5-chlorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(107) N-(3-chloro-2-méthylphényl)-6-([2-chloro-5-méthylphényl]carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(108) N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-6-([5-méthyl-2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(109) 6-([2,5-bis(trifluorométhyl)phényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(110) 6-([2,4-bis(trifluorométhyl)phényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(111) N-(3-chloro-2-méthylphényl)-6-([5-fluoro-2-(trifluorométhyl)phényl]carbonyl)amino)-2-(méthoxyméthyl)-

1H-benzimidazole-4-carboxamide,

(112) N-(3-chloro-2-méthylphényl)-6-([2-chloro-6-(trifluorométhyl)phényl] carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(113) N-(3-chloro-2-méthylphényl)-6-([2-chloro-5-[2-(propan-2-yloxy)éthoxy] phényl]carbonyl)amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(114) 6-([2-chloro-5-(2-éthoxyéthoxy)phényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(115) 6-([2-chloro-5-(3-méthoxypropyl)phényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(116) 6-([5-(3-tert-butoxyprop-1-yn-1-yl)-2-chlorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(117) 6-([5-(3-tert-butoxypropyl)-2-chlorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(118) 6-([2-chloro-5-(3-hydroxy-3-méthylbutyl)phényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(119) 6-([2-chloro-5-(éthoxyméthyl)phényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(120) 6-([2-chloro-5-[(2-éthoxyéthoxy)méthyl]phényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(121) 6-([2-chloro-5-(2-cyclopropyléthyl)phényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(122) N-(3-chloro-2-méthylphényl)-6-([2-chloro-5-(2-phényléthyl)phényl]carbonyl) amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(123) N-(3-chloro-2-méthylphényl)-2-cyclopentyl-6-([2-(trifluorométhyl)phényl] carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(124) N-(3-chloro-2-méthylphényl)-2-cyclopentyl-6-([2,5-dichlorophényl]carbonyl) amino)-1H-benzimidazole-4-carboxamide,

(125) 6-([2-chloro-6-fluorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-cyclopentyl-1H-benzimidazole-4-carboxamide,

(126) 6-([2-chloro-6-fluorophényl]carbonyl)amino)-N-(3-chloro-2-méthylphényl)-2-[(2R)-tétrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(127) N-(3-chloro-2-méthylphényl)-6-([2,6-dichlorophényl]carbonyl)amino)-2-[(2R)-tétrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(128) N-(3-chloro-2-méthylphényl)-6-([2,5-dichlorophényl]carbonyl)amino)-2-[(2R)-tétrahydrofuran-2-yl]-1H-benzimidazole-4-carboxamide,

(129) N-(3-chloro-2-méthylphényl)-2-[(2S)-5-oxopyrrolidin-2-yl]-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(130) N-(3-chloro-2-méthylphényl)-2-[(2R)-5-oxopyrrolidin-2-yl]-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(131) N-(3-chloro-2-méthylphényl)-2-[2-oxo-2-(pyrrolizin-1-yl)éthyl]-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(132) N-(3-chloro-2-méthylphényl)-2-[2-(diméthylamino)-2-oxoéthyl]-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(133) N-(3-chloro-2-méthylphényl)-2-[2-(méthylamino)-2-oxoéthyl]-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(134) 2-chloro-N-(3-chloro-2-méthylphényl)-6-([2-(trifluorométhyl)phényl]carbonyl) amino)-1H-benzimidazole-4-carboxamide,

(135) N-(3-chloro-2-méthylphényl)-2-[2-(méthoxyéthyl)amino]-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(136) N-(3-chloro-2-méthylphényl)-2-[2-(hydroxyéthyl)amino]-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(137) N-(3-chloro-2-méthylphényl)-2-(méthylamino)-6-([2-(trifluorométhyl)phényl] carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(138) N-(3-chloro-2-méthylphényl)-2-(éthylamino)-6-([2-(trifluorométhyl)phényl] carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(139) N-(3-chloro-2-méthylphényl)-2-[2,2-diméthylpropyl]amino)-6-([2-(trifluorométhyl)phényl]carbonyl)amino)-1H-benzimidazole-4-carboxamide,

(140) N-(3-chloro-2-méthylphényl)-2-(cyclopentylamino)-6-([2-(trifluorométhyl) phényl]carbonyl)amino)-1H-

benzimidazole-4-carboxamide,

(141) N-(3-chloro-2-méthylphényl)-2-(pipéridin-1-yl)-6-({[2-(trifluorométhyl) phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(142) N-(3-chloro-2-méthylphényl)-2-(4-méthylpipérazin-1-yl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(143) 2-[bis(2-hydroxyéthyl)amino]-N-(3-chloro-2-méthylphényl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(144) N-(3-chloro-2-méthylphényl)-2-(diméthylamino)-6-({[2-(trifluorométhyl) phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(145) N-(3-chloro-2-méthylphényl)-2-{[2-(morpholin-4-yl)éthyl]amino}-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(146) N-(3-chloro-2-méthylphényl)-2-{[2-(diméthylamino)éthyl]amino}-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(147) N-(3-chloro-2-méthylphényl)-2-(3-hydroxyazétidin-1-yl)-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(148) N-(3-chloro-2-méthylphényl)-2-[(3S)-3-(diméthylamino)pyrrolizin-1-yl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(149) N-(3-chloro-2-méthylphényl)-2-[(3S)-3-hydroxypyrrolizin-1-yl]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(150) N-(3-chloro-2-méthylphényl)-2-{[2-(diéthylamino)éthyl]amino}-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(151) N-(3-chloro-2-méthylphényl)-2-{[2-(pyrrolizin-1-yl)éthyl]amino}-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(152) N-(3-chloro-2-méthylphényl)-2-{[3-(diméthylamino)propyl]amino}-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(153) N-(3-chloro-2-méthylphényl)-2-[[3-(diméthylamino)-2,2-diméthylpropyl]amino]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(154) N-(3-chloro-2-méthylphényl)-2-{[2-(dipropan-2-ylamino)éthyl]amino}-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(155) N-(3-chloro-2-méthylphényl)-2-(morpholin-4-yl)-6-({[2-(trifluorométhyl) phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(156) 2-amino-N-(3-chloro-2-méthylphényl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(157) N-(3-chloro-2-méthylphényl)-2-[(3-hydroxy-2,2-diméthylpropyl)amino]-6-({[2-(trifluorométhyl)phényl]carbonyl }amino)-1H-benzimidazole-4-carboxamide,

(158) N-(3-chloro-2-méthylphényl)-2-[[3-méthyloxetan-3-yl)méthyl]amino]-6-({[2-(trifluorométhyl)phényl]carbonyl }amino)-1H-benzimidazole-4-carboxamide,

(159) N-{4-[(3-chloro-2-méthylphényl)carbamoyl]-6-({[2-(trifluorométhyl) phényl]carbonyl}amino)-1H-benzimidazol-2-yl}glycinate de tert-butyle,

(160) N-{4-[(3-chloro-2-méthylphényl)carbamoyl]-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazol-2-yl}glycine,

(161) N-(3-chloro-2-méthylphényl)-2-[(3-hydroxy-2,2-diméthylpropyl)amino]-1-méthyl-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(162) N-(3-chloro-2-méthylphényl)-2-[(3-méthoxy-2,2-diméthylpropyl)amino]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(163) N-(3-chloro-2-méthylphényl)-2-(pyrrolizin-1-yl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(164) 2-(azétidin-1-yl)-N-(3-chloro-2-méthylphényl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(165) N-(3-chloro-2-méthylphényl)-2-(3-méthoxyazétidin-1-yl)-6-({[2-(trifluorométhyl)phényl]carbonyl} amino)-1H-benzimidazole-4-carboxamide,

(166) N-(3-chloro-2-méthylphényl)-2-[(2-hydroxy-2-méthylpropyl)amino]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(167) N-(3-chloro-2-méthylphényl)-2-[[2S]-tétrahydrofuran-2-ylméthyl]amino]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(168) N-(3-chloro-2-méthylphényl)-2-[[2R]-tétrahydrofuran-2-ylméthyl]amino]-6-({[2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(169) N-(3-chloro-2-méthylphényl)-2-[[2S]-1-hydroxy-3-méthylbutan-2-yl]amino]-6-({[2-(trifluorométhyl)phé-

nyl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(170) N-(3-chloro-2-méthylphényl)-2-[[2R]-1-hydroxy-3-méthylbutan-2-yl]amino]-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(171) N-(3-chloro-2-méthylphényl)-2-[[2S]-1-hydroxy-3,3-diméthylbutan-2-yl]amino]-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(172) N-(3-chloro-2-méthylphényl)-2-[(3-méthoxy-2,2-diméthylpropyl)(méthyl) amino]-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(173) N-(3-chloro-2-méthylphényl)-2-[(3-méthoxypropyl)amino]-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(174) N-(3-chloro-2-méthylphényl)-2-[[2-(propan-2-yloxy)éthyl]amino]-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(175) 2-[(2-tert-butoxyéthyl)amino]-N-(3-chloro-2-méthylphényl)-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(176) N-(3-chloro-2-méthylphényl)-2-[(2-méthoxy-2-méthylpropyl)amino]-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(177) N-(3-chloro-2-méthylphényl)-2-[[2-(méthylsulfanyl)éthyl]amino]-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(178) N-(3-chloro-2-méthylphényl)-2-(méthylsulfanyl)-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(179) N-(3-chloro-2-méthylphényl)-2-(méthylsulfonyl)-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(180) N-(3-chloro-2-méthylphényl)-2-(méthylsulfinyl)-6-([2-(trifluorométhyl)phényl]carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(181) 6-[[2-chloro-6-fluorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-2-(diméthylamino)-1H-benzimidazole-4-carboxamide,

(182) N-(3-chloro-2-méthylphényl)-6-[[2,6-dichlorophényl]carbonyl]amino]-2-(diméthylamino)-1H-benzimidazole-4-carboxamide,

(183) N-(3-chloro-2-méthylphényl)-6-[[2,4-dichlorophényl]carbonyl]amino]-2-(diméthylamino)-1H-benzimidazole-4-carboxamide,

(184) N-(3-chloro-2-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino]-2-(diméthylamino)-1H-benzimidazole-4-carboxamide,

(185) 6-[[2-bromo-6-fluorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-2-(diméthylamino)-1H-benzimidazole-4-carboxamide,

(186) 6-[[2-bromo-6-chlorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-2-(diméthylamino)-1H-benzimidazole-4-carboxamide,

(187) 6-([2-chloro-5-(cyclopropyléthynyl)phényl]carbonyl}amino)-N-(3-chloro-2-méthylphényl)-2-(diméthylamino)-1H-benzimidazole-4-carboxamide,

(188) N-(3-chloro-2-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino]-2-[(3-hydroxy-2,2-diméthylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(189) N-(3-chloro-2-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino]-2-[(3-méthoxy-2,2-diméthylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(190) N-(3-chloro-2-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino]-2-[(2-hydroxy-2-méthylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(191) N-(3-chloro-2-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino]-2-[(2-méthoxy-2-méthylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(192) N-(3-chloro-2-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino]-2-[[2-(propan-2-yloxy)éthyl]amino]-1H-benzimidazole-4-carboxamide,

(193) 6-[[2-chloro-6-fluorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-2-[[2-(propan-2-yloxy)éthyl]amino]-1H-benzimidazole-4-carboxamide,

(194) 2-[(2-tert-butoxyéthyl)amino]-6-[[2-chloro-6-fluorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-1H-benzimidazole-4-carboxamide,

(195) 6-[[2-chloro-6-fluorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-2-[(3-méthoxy-2,2-diméthylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(196) 6-[[2-chloro-6-fluorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-2-[(2-méthoxy-2-méthylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(197) 6-[[2-chloro-6-fluorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-2-[[2S]-tétrahydrofuran-2-ylméthyl]amino]-1H-benzimidazole-4-carboxamide,

(198) 6-[[2-chloro-6-fluorophényl]carbonyl]amino]-N-(3-chloro-2-méthylphényl)-2-[[2R]-tétrahydrofuran-2-yl-

méthyl]amino}-1H-benzimidazole-4-carboxamide,

(199) 6-[[2-chloro-6-fluorophényl]carbonyl]amino}-N-(3-chloro-2-méthylphényl)-2-[(3-hydroxy-2,2-diméthylpropyl)amino]-1H-benzimidazole-4-carboxamide,

(200) 6-[[2-chloro-6-fluorophényl]carbonyl]amino}-N-(3-chloro-2-méthylphényl)-2-[(2S)-1-hydroxy-3-méthylbutan-2-yl]amino}-1H-benzimidazole-4-carboxamide,

(201) N-(3-chloro-4-méthylphényl)-2-(diméthylamino)-6-[[2-(trifluorométhyl) phényl]carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(202) N-(4-tert-butylphényl)-2-(diméthylamino)-6-[[2-(trifluorométhyl) phényl]carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(203) N-(2,3-dihydro-1H-indén-5-yl)-2-(diméthylamino)-6-[[2-(trifluorométhyl) phényl]carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(204) 6-[[2-chloro-6-fluorophényl]carbonyl]amino}-N-(3-chloro-4-méthylphényl)-2-(diméthylamino)-1H-benzimidazole-4-carboxamide,

(205) N-(3-chloro-4-méthylphényl)-6-[[2,6-dichlorophényl]carbonyl]amino}-2-(diméthylamino)-1 H-benzimidazole-4-carboxamide,

(206) N-(3-chloro-4-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino}-2-(diméthylamino)-1 H-benzimidazole-4-carboxamide,

(207) N-(3-chloro-2-méthylphényl)-2-cyclopropyl-6-[[2,5-dichlorophényl]carbonyl] amino}-1H-benzimidazole-4-carboxamide,

(208) N-(3-chloro-4-méthylphényl)-2-cyclopropyl-6-[[2,5-dichlorophényl]carbonyl] amino}-1H-benzimidazole-4-carboxamide,

(209) N-(3-chloro-2-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino}-2-(1-méthylcyclopropyl)-1 H-benzimidazole-4-carboxamide,

(210) N-(3-chloro-4-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino}-2-(1-méthylcyclopropyl)-1 H-benzimidazole-4-carboxamide,

(211) N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-6-[[2-(méthylsulfonyl) phényl]carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(212) N-(3-chloro-4-méthylphényl)-6-[[2,5-dichlorophényl]carbonyl]amino}-2-(2-méthoxyéthyl)-1H-benzimidazole-4-carboxamide,

(213) 2-(méthoxyméthyl)-N-phényl-6-[[2-(trifluorométhyl)phényl]carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(214) 2-(méthoxyméthyl)-N-propyl-6-[[2-(trifluorométhyl)phényl]carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(215) 2-(méthoxyméthyl)-N-(pyridin-3-yl)-6-[[2-(trifluorométhyl)phényl]carbonyl] amino}-1H-benzimidazole-4-carboxamide,

(216) N-benzyl-2-(méthoxyméthyl)-6-[[2-(trifluorométhyl)phényl]carbonyl]amino}-1 H-benzimidazole-4-carboxamide,

(217) N-(cyclohexylméthyl)-2-(méthoxyméthyl)-6-[[2-(trifluorométhyl)phényl] carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(218) 2-(méthoxyméthyl)-N-(naphthalén-1-yl)-6-[[2-(trifluorométhyl)phényl] carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(219) 2-(méthoxyméthyl)-N-(thiophén-3-yl)-6-[[2-(trifluorométhyl)phényl] carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(220) N-(2,1,3-benzothiadiazol-4-yl)-2-(méthoxyméthyl)-6-[[2-(trifluorométhyl) phényl]carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(221) N-(1,1-dioxide-1-benzothiophén-6-yl)-2-(méthoxyméthyl)-6-[[2-(trifluorométhyl)phényl]carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(222) 2-(méthoxyméthyl)-N-(thiophén-2-ylméthyl)-6-[[2-(trifluorométhyl)phényl] carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(223) N-(1H-indol-5-yl)-2-(méthoxyméthyl)-6-[[2-(trifluorométhyl)phényl]carbonyl] amino}-1H-benzimidazole-4-carboxamide,

(224) N-(1,3-benzothiazol-2-yl)-2-(méthoxyméthyl)-6-[[2-(trifluorométhyl)phényl] carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(225) N-(2,2-diméthylpropyl)-2-(méthoxyméthyl)-6-[[2-(trifluorométhyl)phényl] carbonyl]amino}-1H-benzimidazole-4-carboxamide,

(226) 2-(méthoxyméthyl)-N-(thiophén-2-yl)-6-[[2-(trifluorométhyl)phényl]carbonyl] amino}-1H-benzimidazole-4-carboxamide,

(227) N-(5-chloro-1,3-benzoxazol-2-yl)-2-(méthoxyméthyl)-6-[[2-(trifluorométhyl) phényl]carbonyl]amino}-1H-

benzimidazole-4-carboxamide,

(228) N-(2-benzylphényl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(229) 2-(méthoxyméthyl)-N-(quinolin-8-yl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(230) N-(cycloheptylméthyl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(231) N-(1,3-benzoxazol-2-yl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(232) N-(6-chloro-1,3-benzoxazol-2-yl)-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(233) N-[3-chloro-2-(hydroxyméthyl)phényl]-2-(méthoxyméthyl)-6-({[2-(trifluorométhyl)phényl] carbonyl}amino)-1H-benzimidazole-4-carboxamide,

(234) N-(3-chloro-2-méthylphényl)-6-({[3-fluoropyridin-2-yl] carbonyl}amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(235) N-(3-chloro-2-méthylphényl)-6-({[3-chloropyridin-4-yl] carbonyl}amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(236) N-(3-chloro-2-méthylphényl)-6-({[3,5-dichloropyridin-4-yl] carbonyl}amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(237) 6-({[5-butoxy-2-chlorophényl] carbonyl}amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(238) 6-({[2-chloro-5-(2,2-difluoroéthoxy)phényl] carbonyl}amino)-N-(3-chloro-2-méthylphényl)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide,

(239) N-(3-chloro-2-méthylphényl)-6-({[2-chloro-5-(4,4,4-trifluorobutoxy)phényl] carbonyl}amino)-2-(méthoxyméthyl)-1H-benzimidazole-4-carboxamide.

8. Composition pharmaceutique comprenant un composé choisi dans le groupe constitué par le dérivé hétérocyclique selon l'une quelconque des revendications 1 à 7, son tautomère et un sel pharmaceutiquement acceptable de celui-ci en tant que principe actif.

9. Composé choisi dans le groupe constitué par le dérivé hétérocyclique selon l'une quelconque des revendications 1 à 7, son tautomère et un sel pharmaceutiquement acceptable de celui-ci pour une utilisation en tant qu'agent inhibant mPGES.

10. Composé choisi dans le groupe constitué par le dérivé hétérocyclique selon l'une quelconque des revendications 1 à 7, son tautomère et un sel pharmaceutiquement acceptable de celui-ci, pour une utilisation dans la prévention ou le traitement d'une maladie intestinale inflammatoire, du syndrome du côlon irritable, de la migraine, des maux de tête, d'une douleur lombaire, de la sténose spinale, de l'hernie discale, des troubles de l'articulation temporo-mandibulaire, du syndrome cervical, de l'arthrose cervicale, de l'endométriose, de l'adénomyose, du travail et de l'accouchement prématurés, de l'accouchement prématuré dangereux pour la santé, de la dysménorrhée, de la vessie hyperactive, de la nycturie, de la cystite interstitielle, d'une maladie neurodégénérative, du psoriasis, de la polyarthrite rhumatoïde, du rhumatisme articulaire aigu, de la fibromyalgie, de la névralgie, du syndrome douloureux régional complexe, du dysfonctionnement facial, des infections virales, des infections bactériennes, des mycoses, des brûlures, de l'inflammation et d'une douleur après opération, de lésions et de l'extraction dentaire, de tumeurs malignes, de l'athérosclérose, de l'accident vasculaire cérébral, de la goutte, de l'arthrite, de l'arthrose, de l'arthrite juvénile, de la spondylarthrite ankylosante, de la ténosynovite, de l'ossification de ligaments, du lupus érythémateux aigu disséminé, de la vasculite, de la pancréatite, de la néphrite, de la conjonctivite, de l'irritation, de la sclérite, de l'uvéïte, du traitement de plaies, de la dermatite, de l'eczéma, de l'ostéoporose, de l'asthme, de la maladie pulmonaire obstructive chronique, de la fibrose pulmonaire, d'une maladie allergique, de la polypose adénomateuse familiale, de la sclérodermie, de la bursite, du fibrome de l'utérus, ou d'une douleur provenant d'un cancer.

11. Composé pour une utilisation selon la revendication 10, dans lequel la maladie neurodégénérative est la maladie d'Alzheimer ou la sclérose en plaques.

12. Composé pour une utilisation selon la revendication 10, dans lequel la maladie infectieuse virale est la grippe, le rhume, le zona ou le SIDA.

13. Composé pour une utilisation selon la revendication 10, dans lequel la tumeur maligne est le cancer du côlon, le

EP 2 746 265 B1

cancer du sein, le cancer du poumon ou le cancer de la prostate.

5

10

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2009103778 A [0003]
- US 20100324086 A [0003]
- WO 2011048004 A [0003]
- WO 200865508 A [0038] [0043] [0118]
- EP 2226315 A [0038]
- EP 234872 B1 [0059]
- US 6387938 B1 [0059]
- WO 2006116412 A [0063]
- WO 2010132999 A [0084]
- EP 1479681 A [0166]
- WO 2003106430 A [0196] [0351]
- US 6020368 A [0204]
- WO 200679642 A [0215] [0216] [0217] [0218] [0284]
- EP 1803350 A [0229]
- US 200439038 B [0230] [0231]
- WO 200968512 A [0236] [0400]
- WO 200523773 A [0242]
- WO 2008124575 A [0257]
- US 2009186879 A [0302]
- US 200976062 B [0331]
- WO 200728051 A [0335]
- WO 200957827 A [0339]
- WO 2007144327 A [0368]


Non-patent literature cited in the description

- JAKOBSSON et al. *Proc. Natl. Acad. Sci. USA*, 1999, vol. 96, 7220-7225 [0004]
- TANIKAWA et al. *Biochem. Biophys. Res. Commun.*, 2002, vol. 291, 884-889 [0004]
- TANIOKA et al. *J. Biol. Chem.*, 2000, vol. 275, 32775-32782 [0004]
- MURAKAMI et al. *J. Biol. Chem.*, 2000, vol. 275, 32783-32792 [0004]
- KAMEI et al. *J. Biol. Chem.*, 2004, vol. 279, 33684-33695 [0004]
- KOJIMA et al. *J. Immunol.*, 2008, vol. 108, 3861-3868 [0004]
- KIMURA et al. *Proc. Natl. Acad. Sci. USA*, 2000, vol. 106, 21807-21812 [0004]
- ENGBLOM et al. *Nat. Neurosci.*, 2003, vol. 6, 1137-1138 [0004]
- FOUDI et al. *Cardiovasc. Res.*, 2009, vol. 81, 269-277 [0004]
- CHENG et al. *J. Clin. Invest.*, 2006, vol. 116, 1391-1399 [0004]
- *J. Biol. Chem.*, 2004, vol. 279 (13), 12647-12658 [0004]
- *Biomed. Pharmacother.*, 2001, vol. 65 (1), 77-84 [0004]
- *Acta. Med. Okayama*, 2008, vol. 62 (6), 373-378 [0004]
- *Proc. Natl. Acad. Sci. USA*, 2009, vol. 106, 21807-21812 [0004]
- *J. Biol. Chem.*, vol. 279 (32), 33684-33695 [0004]
- *J. Immunol.*, 2008, vol. 180, 8361-8368 [0004]
- *Glia*, 2011, vol. 59, 208-218 [0004]
- J. BURN. *Care. Res.*, 2011, vol. 32 (1), 79-90 [0004]
- *J. Biol. Chem.*, 2003, vol. 278 (21), 19396-19405 [0004]
- *Oncogene*, 2012, vol. 31 (24), 2943-2952 [0004]
- *Cancer Res.*, 2008, vol. 68 (9), 3251-3259 [0004]
- *Proc. Natl. Acad. Sci. USA*, 2006, vol. 103, 14507-14512 [0004]
- *Proc. Natl. Acad. Sci. USA*, 2006, vol. 103, 11790-11795 [0004]
- *J. Rheumatol.*, 2005, vol. 32 (5), 887-895 [0004]
- *J. Pharmacol. Exp. Ther.*, 2008, vol. 326, 754-763 [0004]
- *Kidney Int.*, 2011, vol. 79 (1), 77-88 [0004]
- *J. Immunol.*, 2012, vol. 188, 4093-4102 [0004]
- *Am. J. Physiol. Lung Cell Mol. Physiol.*, 2004, vol. 287, L981-L991 [0004]
- *Am. J. Physiol. Lung Cell Mol. Physiol.*, 2005, vol. 288, L1010-L1016 [0004]
- *Arthritis Res. Ther.*, 2011, vol. 13, R6 [0004]
- *J. Org. Chem.*, 1960, vol. 25, 942 [0038]
- *J. Med. Chem.*, 1999, vol. 42, 5020 [0043] [0059]
- *Bioorg. Med. Chem.*, vol. 7, 2271 [0059]
- *J. Med. Chem.*, 1993, vol. 36, 2182 [0063] [0188]
- *Eur. J. Med. Chem.*, 1999, vol. 34, 729 [0180]
- *Tetrahedron Asymmetry*, 2007, vol. 18, 2011 [0220] [0221]
- *Bull. Chem. Soc. Jpn.*, 2001, vol. 74, 1695 [0226]
- *J. Am. Chem. Soc.*, 1974, vol. 96, 1518 [0235]
- *Bioorg. Med. Chem. Lett.*, 2010, vol. 20, 2204 [0271]
- *Tetrahedron Lett.*, 1997, vol. 38, 2259 [0289] [0290]
- *J. Med. Chem.*, 2005, vol. 48, 2080 [0428]
- BRIDEAU et al. *Inflamm. Res.*, 1996, vol. 45, 68 [0448]


Heterociklusos származékok, mint mikroszomális prosztaglandin E szintetáz inhibitorok
(mPGEs)

SZABADALMI IGÉNYPONTOK

1. Egy [1] általános képletű heterociklusos származék:



vagy annak tautomerje vagy gyógyászatilag elfogadható sója, ahol az A gyűrű valamely, [2], [3] vagy [4] általános képletű csoport:



ahol

X^1 jelentése NH-, N-alkilcsoport, vagy O atom;

A^1 hidrogénatom vagy alkilcsoport;

A^2 jelentése

i) hidrogénatom;

ii) halogénatom;


iii) adott esetben egy-három, a halogénatom, amino-, monoalkilamino-,

dialkilamino-, karbamoil-, monoalkilaminokarbonil-, dialkilaminokarbonil-,

telített ciklusos aminokarbonil-, alkoxi-, alkoxialkoxi- és alkilkarboniloxicsop

ortból választott szubsztituenssel helyettesített alkilcsoport;

- iv) adott esetben egy-három halogénatommal helyettesített alkilcsoporttal adott esetben helyettesített cikloalkilcsoport;
- v) alkoxycsoport;
- vi) adott esetben alkil-, alkiloxikarbonil-, alkilkarbonil- vagy oxocsoporttal helyettesített telített heterociklusos csoport;
- vii) alkiltiocsoport;
- viii) alkilszulfonilcsoport;
- ix) alkilszulfonilcsoport;
- x) egy [5] általános képletű csoport:



ahol

R^3 és R^4 azonos vagy különböző, az alábbiak közül kiválasztva:

- a) hidrogénatom,
- b) alkilcsoport, mely adott esetben valamely, a monoalkilamino-, dialkilamino-, adott esetben alkilcsoporttal helyettesített telített ciklusos aminocsoport, az alkil-, alkoxi-, hidroxikarbonil-, hidroxil-, alkiloxikarbonil és alkiltiocsoport csoportból választott szubsztituenssel adott esetben helyettesített telített heterociklusos csoport, vagy
- c) cikloalkilcsoport; vagy

- xi) adott esetben alkil-, amino-, monoalkilamino-, dialkilamino-, alkoxi- vagy hidroxilcsoporttal helyettesített telített ciklusos aminocsoport;

R^1 jelentése fenil-, benzil-, naftil-, cikloalkil-, cikloalkilmetil-, heteroaril-, heteroarilmetil-, 1,2,3,4-tetrahidronaftalin-5-il-, 1,2,3,4-tetrahidronaftalin-6-il-, 2,3-dihidro-1H-inden-4-il-, 2,3-dihidro-1H-inden-5-il-, 1,2-dihidrociklobutabenzil-3-il-, 1,2-dihidrociklobutabenzil-4-il- vagy alkilcsoport,

ahol az említett fenil-, benzil-, cikloalkil-, cikloalkilmetil-, heteroaril- és heteroarilmetilcsoport adott esetben egy-három, a következő csoportból választott helyettesítővel helyettesítve lehet:

- i) halogénatom,

ii) adott esetben egy-három, a halogénatom, hidroxil- és fenilcsoport csoportból választott szubsztituenssel helyettesítet alkilcsoport;

iii) alkoxics csoport;

iv) hidroxilcsoport; és

v) cianocsoport;

R^2 fenilcsoport vagy piridilcsoport,

ahol az említett fenilcsoport vagy piridilcsoport adott esetben egy-három, az alábbi csoportból választott szubsztituenssel lehet helyettesítve:

i) halogénatom,

ii) alkilszulfonilcsoport,

iii) alkoxics csoport, mely adott esetben egy-három halogénatommal vagy alkoxics csoporttal lehet helyettesítve;

iv) alkinilcsoport, mely adott esetben alkoxialkil- vagy cikloalkilcsoporttal lehet helyettesítve, és

v) alkilcsoport, mely adott esetben egy-három, a következő csoportból választott szubsztituenssel lehet helyettesítve: alkoxi, alkoxialkoxi-, cikloalkil-, fenilcsoport és halogénatom.

2. Az 1. igénypont szerinti heterociklusos származék, ahol az A gyűrű egy [4] általános képletű csoport, és X^1 jelentése NH, vagy e vegyület tautomerje vagy gyógyászatilag elfogadható sója.

3. Az 1. igénypont szerinti heterociklusos származék, ahol R^1 jelentése fenil-, 1,2,3,4-tetrahidronaftalin-5-il-, 1,2,3,4-tetrahidronaftalin-6-il-, 2,3-dihidro-1H-inden-4-il-, 2,3-dihidro-1H-inden-5-il-, 1,2-dihidrociklobutabenzil-3-il-, vagy 1,2-dihidrociklobutabenzil-4-il-csoport, és az említett fenilcsoport adott esetben az alábbi csoportból kiválasztott egy-három szubsztituenssel lehet helyettesítve:

i) halogénatom,

ii) adott esetben egy-három halogénatommal helyettesített alkilcsoport,

iii) alkoxics csoport, és

iv) cianocsoport,

vagy e vegyület tautomere vagy gyógyászatilag elfogadható sója.

4. Az 1. igénypont szerinti heterociklusos származék, ahol R^2 jelentése fenilcsoport, és az említett fenilcsoport adott esetben egy-három, az alábbi csoportból kiválasztott szubsztituenssel lehet helyettesítve:

- i) halogénatom,
- ii) alkilszulfonilcsoport,
- iii) adott esetben alkoxicsoporttal helyettesített alkoxicsoport,
- iv) adott esetben alkoxialkil- vagy ciklolakilcsoporttal helyettesített alkinilcsoport,
- és
- v) adott esetben egy-három, a halogénatom, alkoxi-, alkoxialkoxi-, cikloalkil- és fenilcsoport csoportból kiválasztott szubsztituenssel helyettesített alkilcsoport,

vagy e vegyület tautomere vagy gyógyászatilag elfogadható sója.

5. Az 1. igénypont szerinti heterociklusos származék,

ahol

az A gyűrű egy [4] általános képletű csoport,

X^1 jelentése NH,

A^2 jelentése

- i) hidrogénatom,
- ii) alkilcsoport, mely adott esetben a halogénatom, monoalkilamino-, dialkilamino-, monoalkilaminokarbonil-, dialkilaminokarbonil-, telített ciklusos aminokarbonil-, alkoxi, alkoxialkoxi- és alkilkarboniloxicsoport csoportból kiválasztott szubsztituenssel lehet helyettesítve,
- iii) adott esetben egy-három halogénatommal helyettesített alkilcsoporttal helyettesített cikloalkilcsoport,
- iv) alkoxicsoport,
- v) adott esetben alkil- vagy alkiloxikarbonilcsoporttal szubsztituált telített heterociklusos csoport,
- vi) alkiltiocsoport,
- vii) alkilszulfonilcsoport,
- viii) alkilszulfonilcsoport,
- ix) alkilcsoporttal helyettesített aminocsoport, ahol az említett alkilcsoport adott esetben egy, a monoalkilamino-, dialkilamino-, telített ciklusos aminocsoport, mely adott esetben alkil-, tetrahydrofuril-, morfolino-, alkoxi, hidroxikarbonil-, hidroxil- és alkiltiocsoporttal szubsztituált, csoportból választott szubsztituenssel lehet helyettesítve,
- x) cikloalkilcsoporttal helyettesített aminocsoport vagy

xi) telített ciklusos aminosocsoport, mely adott esetben alkil-, dialkilamino-, alkoxi- vagy hidroxilcsoporttal szubsztituált, és

R^1 jelentése

- i) adott esetben egy-három, a halogénatom, adott esetben egy-három halogénatommal szubsztituált alkilcsoport, alkoxi- és cianocsoportból választott csoporttal szubsztituált fenilcsoport,
- ii) 1,2,3,4-tetrahidronaftalin-5-il,
- iii) 2,3-dihidro-1H-inden-5-il,
- iv) benzilcsoport, mely adott esetben halogénatommal vagy adott esetben egy-három halogénatommal szubsztituált alkilcsoporttal lehet helyettesítve,
- v) cikloalkilcsoport,
- vi) cikloalkilmetilcsoport,
- vii) naftilcsoport,
- viii) piridilmetilcsoport, mely adott esetben alkilcsoporttal lehet helyettesítve, mely utóbbi adott esetben egy-három halogénatommal helyettesített,
- ix) tienilcsoport,
- x) tienilmetilcsoport,
- xi) benzotiazolilcsoport,
- xii) benzotiadiazolilcsoport,
- xiii) indolilcsoport vagy
- xiv) alkilcsoport, és

R^2 fenil- vagy piridilcsoport,

ahol az említett fenilcsoport adott esetben az alábbi csoportból kiválasztott egy-három helyettesítővel lehet helyettesítve

- i) halogénatom,
- ii) alkilszulfonilcsoport;
- iii) adott esetben alkoxicsocporttal helyettesített alkoxicsocport,
- iv) adott esetben alkoxialkil- vagy cikloalkilcsoporttal helyettesített alkinilcsoport, és
- v) adott esetben egy-három, a halogénatom, alkoxi-, alkoxialkoxi-, cikloalkil-, és fenilcsoport csoportból kiválasztott helyettesítővel szubsztituált alkilcsoport, és

az említett piridilcsoport adott esetben halogénatommal lehet helyettesítve,

vagy e vegyület tautomere vagy gyógyászatilag elfogadható sója.

6. Az 1. igénypont szerinti heterociklusos származék, ahol

az A gyűrű egy [4] általános képletű csoport,

X^1 jelentése NH ,

A^2 alkilcsoport, mely alkoxi-, dialkilamino-, tetrahydrofuril-, tetrahydrofurilmetil-, alkoxialkilamino-, vagy adott esetben alkilcsoporttal, mely adott esetben egy-három halogénatommal lehet helyettesítve, szubsztituált cikloalkilcsoporttal van helyettesítve,

R^1 egy halogénatommal és egy metilcsoporttal szubsztituált fenilcsoport, és R^2 adott esetben egy trifluormetilcsoporttal vagy két halogénatommal szubsztituált fenilcsoport,

vagy e vegyület tautomere vagy gyógyászatilag elfogadható sója.

7. Az 1. igénypont szerinti, a következő (1)-(239) közül választott heterociklusos származék, vagy annak tautomere vagy gyógyászatilag elfogadható sója:

- (1) N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (2) N-ciklohexil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (3) N-(3-klór-2-metilfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (4) N-[(1-hidroxikiklohexil)-metil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (5) N-[2-(trifluormetil)-benzil]-5-([2-(trifluormetil)-fenil]-karbonil)-amino)-2,3-dihidro-1-benzofurán-7-karboxamid,
- (6) N-ciklohexil-5-([2-(trifluormetil)-fenil]-karbonil)-amino)-2,3-dihidro-1-benzofurán-7-karboxamid,
- (7) N-(3-klór-2-metilfenil)-5-([2-(trifluormetil)-fenil]-karbonil)-amino)-2,3-dihidro-1-benzofurán-7-karboxamid,
- (8) N-ciklohexil-5-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-indazol-7-karboxamid,
- (9) N-[2-(trifluormetil)-benzil]-5-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-indazol-7-karboxamid,
- (10) N-(3-klór-2-metilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (11) 2-metil-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,

- (12) N-ciklohexil-2-metil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (13) N-(3-klór-2-metilfenil)-2-metil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (14) N-ciklopentil-2-metil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (15) N-ciklobutil-2-metil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (16) N-(3-klór-2-metilfenil)-2-etil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-H-benzimidazol-4-karboxamid,
- (17) N-ciklohexil-2-etil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (18) 2-etil-N-[2-(trifluormetil)-benzil]-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (19) N-ciklohexil-2-(metoximetil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-H-benzimidazol-4-karboxamid,
- (20) 2-(metoximetil)-N-[2-(trifluormetil)-benzil]-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (21) 2-(metoximetil)-N-(2-metilfenil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (22) 2-(metoximetil)-N-(4-metilfenil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (23) N-(2-klórbenzil)-2-(metoximetil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (24) 2-(metoximetil)-N-(4-metilbenzil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (25) N-(4,4-difluorciklohexil)-2-(metoximetil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (26) N-(4-terc-butilfenil)-2-(metoximetil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (27) 2-(metoximetil)-N-[4-(trifluormetil)-fenil]-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,

- (28) N-(2,4-dimetilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (29) N-(2-klór-4-metilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid
- (30) N-(3,4-dimetilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (31) N-(3-klór-4-metilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (32) N-(2,3-dihidro-1H-inden-5-il)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (33) 2-(metoximetil)-N-(5,6,7,8-tetrahidronaftalin-1-il)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (34) N-(2-fluorfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (35) 2-(metoximetil)-N-(2-metoxifenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (36) 2-(metoximetil)-N-(4-metoxifenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (37) N-(3-bróm-2-metilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (38) N-(3-klór-2-metilbenzil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (39) N-(2,6-difluorfenil)-2-(metoximetil)-6-([2-(metoxi-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (40) N-(3-ciano-2-metilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (41) 2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-N-([3-(trifluormetil)-piridin-2-il]-metil)-1H-benzimidazol-4-karboxamid,
- (42) N-(2-klór-6-metilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (43) 2-(2-amino-2-oxoetil)-N-(3-klór-2-metilfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,

- (44) 2-(2-amino-2-oxoetil)-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (45) N-(3-klór-2-metilfenil)-1-metil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (46) N-ciklohexil-1-metil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (47) 1-metil-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (48) N-(3-klór-2-metilfenil)-1-etil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (49) N-ciklohexil-1-etil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (50) 1-etil-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (51) N-(3-klór-2-metilfenil)-2-metil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1,3-benzoxazol-4-karboxamid,
- (52) 2-metil-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1,3-benzoxazol-4-karboxamid,
- (53) N-(3-klór-2-metilfenil)-2-etil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1,3-benzoxazol-4-karboxamid,
- (54) N-(3-klór-2-metilfenil)-2-etoxi-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (55) 2-etoxi-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (56) N-(3-klór-2-metilfenil)-2-(1-klór-2-metilpropan-2-il)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-H-benzimidazol-4-karboxamid,
- (57) N-(3-klór-2-metilfenil)-2-[(dimetilamino)-metil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (58) N-(3-klór-2-metilfenil)-2-(2-metilpropil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (59) 2-(2-metilpropil)-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-imidazol-4-karboxamid,

- (60) terc-butil-3-{4-[(3-klór-2-metilfenil)-karbamoil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino}-1H-benzimidazol-2-il}azetidin-1-karboxilát,
- (61) N-(3-klór-2-metilfenil)-2-[(metilamino)-metil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (62) {4-[(3-klór-2-metilfenil)-karbamoil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-2-il}-metil-acetát,
- (63) N-(3-klór-2-metilfenil)-2-[(2R)-tetrahydrofurán-2-il]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (64) 2-[(2R)-tetrahydrofurán-2-il]-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-H-benzimidazol-4-karboxamid,
- (65) N-(3-klór-2-metilfenil)-2-[(2S)-tetrahydrofurán-2-il]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (66) 2-[(2S)-tetrahydrofurán-2-il]-N-[2-(trifluormetil)-benzil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (67) 2-(1-acetilazetidin-3-il)-N-(3-klór-2-metilfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (68) terc-butyl (2S)-2-{4-[(3-klór-2-metilfenil)-karbamoil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-2-il}-pirrolidin-1-karboxilát,
- (69) terc-butyl (2R)-2-{4-[(3-klór-2-metilfenil)-karbamoil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-2-il}-pirrolidin-1-karboxilát,
- (70) N-(3-klór-2-metilfenil)-2-[(2S)-pirrolidin-2-il]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (71) N-(3-klór-2-metilfenil)-2-[(2S)-1-metilpirrolidin-2-il]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (72) 2-[(2S)-1-acetilpirrolidin-2-il]-N-(3-klór-2-metilfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (73) N-(3-klór-2-metilfenil)-2-[(2-metoxietoxi)-metil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (74) N-(3-klór-2-metilfenil)-2-(1-metoxi-2-metilpropan-2-il)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (75) 2-terc-butyl-N-(3-klór-2-metilfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,

- (76) 2-terc-butil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-N-({[3-(trifluormetil)-piridin-2-il]-metil})-1H-benzimidazol-4-karboxamid,
- (77) N-(3-klór-2-metilfenil)-2-(2-etoxietil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (78) N-(3-klór-2-metilfenil)-2-(etoximetil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (79) 2-(etoximetil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-N-({[3-(trifluormetil)-piridin-2-il]-metil})-1H-benzimidazol-4-karboxamid,
- (80) N-(3-klór-2-metilfenil)-2-(2-metoxietil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (81) N-(3-klór-2-metilfenil)-2-(2,2-dimetilpropil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (82) N-(3-klór-2-metilfenil)-2-ciklopropil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (83) N-(3-klór-2-metilfenil)-2-(2-metilpentan-2-il)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (84) N-(3-klór-2-metilfenil)-2-(1-metilciklopropil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (85) 2-terc-butil-N-(3-klór-4-metilfenil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (86) 2-terc-butil-N-(3-klór-2-metilfenil)-6-({[2,5-diklórfenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (87) 2-terc-butil-N-(3-klór-4-metilfenil)-6-({[2,5-diklórfenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (88) N-(3-klór-2-metilfenil)-2-[1-(trifluormetil)-ciklopropil]-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (89) N-(3-klór-2-metilfenil)-2-(metoximetil)-1-metil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (90) N-(2-klórbenzil)-2-(metoximetil)-1-metil-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (91) 6-({[2-klór-6-fluorfenil]-karbonil}-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-H-benzimidazol-4-karboxamid,

- (92) 6-[[(2-klór-4-fluorfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-metoximetil-1H-benzimidazol-4-karboxamid,
- (93) 6-[[(2-klór-5-fluorfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (94) N-(3-klór-2-metilfenil)-6-[[(2-klórfenil)-karbonil]-amino}-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (95) N-(3-klór-2-metilfenil)-6-[[(2-klórpíridin-3-il)-karbonil]-amino}-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (96) 6-[[(2-brómfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (97) N-(3-klór-2-metilfenil)-6-[[(2,6-diklórfenil)-karbonil]-amino}-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (98) N-(3-klór-2-metilfenil)-6-[[(2,5-diklórfenil)-karbonil]-amino}-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (99) 6-[[(2-klór-3-fluorfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (100) 6-[[(2-klór-3,6-difluorfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (101) 6-[[(2-bróm-6-klórfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (102) 6-[[(2-bróm-6-fluorfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (103) N-(3-klór-2-metilfenil)-6-[[(2-klór-6-metilfenil)-karbonil]-amino}-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (104) N-(3-klór-2-metilfenil)-6-[[(2-klór-4-metilfenil)-karbonil]-amino}-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (105) 6-[[(5-bróm-2-klórfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (106) 6-[[(2-bróm-5-klórfenil)-karbonil]-amino}-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (107) N-(3-klór-2-metilfenil)-6-[[(2-klór-5-metilfenil)-karbonil]-amino}-2-(metoximetil)-1H-benzimidazol-4-karboxamid,

- (108) N-(3-klór-2-metilfenil)-2-(metoximetil)-6-([5-metil-2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (109) 6-([2,5-bisz(trifluormetil)-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (110) 6-([2,4-bisz(trifluormetil)-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (111) N-(3-klór-2-metilfenil)-6-([5-fluor-2-(trifluormetil)-fenil]-karbonil)-amino)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (112) N-(3-klór-2-metilfenil)-6-([2-klór-6-(trifluormetil)-fenil]-karbonil)-amino)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (113) N-(3-klór-2-metilfenil)-6-([2-klór-5-[2-(propan-2-il-oxi)-etoxi]-fenil]-karbonil)-amino)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (114) 6-([2-klór-5-(2-etoxietoxi)-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (115) 6-([2-klór-5-(3-metoxipropil)-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (116) 6-([5-(3-terc-butoxiprop-1-in-1-il)-2-klór-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (117) 6-([5-(3-terc-butoxipropil)-2-klór-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (118) 6-([2-klór-5-(3-hidroxi-3-metilbutil)-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (119) 6-([2-klór-5-(etoximetil)-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (120) 6-([2-klór-5-[(2-etoxietoxi)-metil]-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (121) 6-([2-klór-5-(2-ciklopropiletil)-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (122) N-(3-klór-2-metilfenil)-6-([2-klór-5-(2-feniletil)-fenil]-karbonil)-amino)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (123) N-(3-klór-2-metilfenil)-2-ciklopentil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,

- (124) N-(3-klór-2-metilfenil)-2-ciklopentil-6-([(2,5-diklórfenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (125) 6-([(2-klór-6-fluorfenil)-karbonil]-amino)-N-(3-klór-2-metilfenil)-2-ciklopentil-1H-benzimidazol-4-karboxamid,
- (126) 6-([(2-klór-6-fluorfenil)-karbonil]-amino)-N-(3-klór-2-metilfenil)-2-[(2R)-tetrahydrofuran-2-il]-1H-benzimidazol-4-karboxamid,
- (127) N-(3-klór-2-metilfenil)-6-([(2,6-diklórfenil)-karbonil]-amino)-2-[(2R)-tetrahydrofuran-2-il]-1H-benzimidazol-4-karboxamid,
- (128) N-(3-klór-2-metilfenil)-6-([(2,5-diklórfenil)-karbonil]-amino)-2-[(2R)-tetrahydrofuran-2-il]-1H-benzimidazol-4-karboxamid,
- (129) N-(3-klór-2-metilfenil)-2-[(2S)-5-oxopirrolidin-2-il]-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (130) N-(3-klór-2-metilfenil)-2-[(2R)-5-oxopirrolidin-2-il]-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (131) N-(3-klór-2-metilfenil)-2-[2-oxo-2-(pirrolizin-1-il)-etil]-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (132) N-(3-klór-2-metilfenil)-2-[2-(dimetilamino)-2-oxoetil]-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (133) N-(3-klór-2-metilfenil)-2-[2-(metilamino)-2-oxoetil]-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (134) 2-klór-N-(3-klór-2-metilfenil)-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (135) N-(3-klór-2-metilfenil)-2-[(2-metoxietil)-amino]-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (136) N-(3-klór-2-metilfenil)-2-[(2-hidroxietil)-amino]-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (137) N-(3-klór-2-metilfenil)-2-(metilamino)-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (138) N-(3-klór-2-metilfenil)-2-(etilamino)-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (139) N-(3-klór-2-metilfenil)-2-[(2,2-dimetilpropil)-amino]-6-([(2-(trifluormetil)-fenil)-karbonil]-amino)-1H-benzimidazol-4-karboxamid,

- (140) N-(3-klór-2-metilfenil)-2-(ciklopentilamino)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (141) N-(3-klór-2-metilfenil)-2-(piperidin-1-il)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (142) N-(3-klór-2-metilfenil)-2-(4-metilpiperazin-1-il)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (143) 2-[bisz(2-hidroxietyl)-amino]-N-(3-klór-2-metilfenil)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (144) N-(3-klór-2-metilfenil)-2-(dimetilamino)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (145) N-(3-klór-2-metilfenil)-2-{{[2-(morfolin-4-il)-etyl]-amino}}-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (146) N-(3-klór-2-metilfenil)-2-{{[2-(dimetilamino)-etyl]-amino}}-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (147) N-(3-klór-2-metilfenil)-2-(3-hidroxiacetidin-1-il)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (148) N-(3-klór-2-metilfenil)-2-[(3S)-3-(dimetilamino)-pirrolizin-1-il]-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (149) N-(3-klór-2-metilfenil)-2-[(3S)-3-hidroxi-2-pirrolizin-1-il]-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (150) N-(3-klór-2-metilfenil)-2-{{[2-(diethylamino)-etyl]-amino}}-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (151) N-(3-klór-2-metilfenil)-2-{{[2-(pirrolizin-1-il)-etyl]-amino}}-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (152) N-(3-klór-2-metilfenil)-2-{{[3-(dimetilamino)-propil]-amino}}-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (153) N-(3-klór-2-metilfenil)-2-{{[3-(dimetilamino)-2,2-dimetilpropil]-amino}}-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (154) N-(3-klór-2-metilfenil)-2-{{[2-(dipropán-2-il-amino)-etyl]-amino}}-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,
- (155) N-(3-klór-2-metilfenil)-2-(morfolin-4-il)-6-({[2-(trifluormetil)-fenil]-karbonil}-amino)-1H-benzimidazol-4-karboxamid,

- (156) 2-amino-N-(3-klór-2-metilfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (157) N-(3-klór-2-metilfenil)-2-[(3-hidroxi-2,2-dimetilpropil)-amino]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (158) N-(3-klór-2-metilfenil)-2-[(3-metiloxetán-3-il)-metil]-amino)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (159) terc-butil N-{4-[(3-klór-2-metilfenil)-karbamoil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-2-il}-glicinát,
- (160) N-{4-[(3-klór-2-metilfenil)-karbamoil]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-2-il}-glicin,
- (161) N-(3-klór-2-metilfenil)-2-[(3-hidroxi-2,2-dimetilpropil)-amino]-1-metil-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (162) N-(3-klór-2-metilfenil)-2-[(3-metoxi-2,2-dimetilpropil)-amino]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (163) N-(3-klór-2-metilfenil)-2-(pirrolizin-1-il)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (164) 2-(azetidín-1-il)-N-(3-klór-2-metilfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-H-benzimidazol-4-karboxamid,
- (165) N-(3-klór-2-metilfenil)-2-(3-metoxiazetidín-1-il)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (166) N-(3-klór-2-metilfenil)-2-[(2-hidroxi-2-metilpropil)-amino]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (167) N-(3-klór-2-metilfenil)-2-[(2S)-tetrahydrofurán-2-il-metil]-amino)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (168) N-(3-klór-2-metilfenil)-2-[(2R)-tetrahydrofurán-2-il-metil]-amino)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (169) N-(3-klór-2-metilfenil)-2-[(2S)-1-hidroxi-3-metilbután-2-il]-amino)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (170) N-(3-klór-2-metilfenil)-2-[(2R)-1-hidroxi-3-metilbután-2-il]-amino)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (171) N-(3-klór-2-metilfenil)-2-[(2S)-1-hidroxi-3,3-dimetilbután-2-il]-amino)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,

- (172) N-(3-klór-2-metilfenil)-2-[(3-metoxi-2,2-dimetilpropil)-(metil)-amino]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (173) N-(3-klór-2-metilfenil)-2-[(3-metoxipropil)-amino]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (174) N-(3-klór-2-metilfenil)-2-([2-(propan-2-iloxi)-etil]-amino)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (175) 2-[(2-terc-butoxietyl)-amino]-N-(3-klór-2-metilfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (176) N-(3-klór-2-metilfenil)-2-[(2-metoxi-2-metilpropil)-amino]-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (177) N-(3-klór-2-metilfenil)-2-([2-(metilszulfanil)-etil]-amino)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (178) N-(3-klór-2-metilfenil)-2-(metilszulfanil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (179) N-(3-klór-2-metilfenil)-2-(metilszulfonil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (180) N-(3-klór-2-metilfenil)-2-(metilszulfenil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (181) 6-([(2-klór-6-fluorfenil)-karbonil]-amino)-N-(3-klór-2-metilfenil)-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (182) N-(3-klór-2-metilfenil)-6-([(2,6-diklórfenil)-karbonil]-amino)-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (183) N-(3-klór-2-metilfenil)-6-([(2,4-diklórfenil)-karbonil]-amino)-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (184) N-(3-klór-2-metilfenil)-6-([(2,5-diklórfenil)-karbonil]-amino)-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (185) 6-([(2-bróm-6-fluorfenil)-karbonil]-amino)-N-(3-klór-2-metilfenil)-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (186) 6-([(2-bróm-6-klórfenil)-karbonil]-amino)-N-(3-klór-2-metilfenil)-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (187) 6-([2-klór-5-(ciklopropiletinil)-fenil]-karbonil)-amino)-N-(3-klór-2-metilfenil)-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,

- (188) N-(3-klór-2-metilfenil)-6-[[[(2,5-diklórfenil)-karbonil]-amino]-2-[(3-hidroxi-2,2-dimetilpropil)-amino]-1H-benzimidazol-4-karboxamid,
- (189) N-(3-klór-2-metilfenil)-6-[[[(2,5-diklórfenil)-karbonil]-amino]-2-[(3-metoxi-2,2-dimetilpropil)-amino]-1H-benzimidazol-4-karboxamid,
- (190) N-(3-klór-2-metilfenil)-6-[[[(2,5-diklórfenil)-karbonil]-amino]-2-[(2-hidroxi-2-metilpropil)-amino]-1H-benzimidazol-4-karboxamid,
- (191) N-(3-klór-2-metilfenil)-6-[[[(2,5-diklórfenil)-karbonil]-amino]-2-[(2-metoxi-2-metilpropil)-amino]-1H-benzimidazol-4-karboxamid,
- (192) N-(3-klór-2-metilfenil)-6-[[[(2,5-diklórfenil)-karbonil]-amino]-2-[[2-(propan-2-iloxi)-etil]-amino]-1H-benzimidazol-4-karboxamid,
- (193) 6-[[[(2-klór-6-fluorfenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-[[2-(propan-2-iloxi)-etil]-amino]-1H-benzimidazol-4-karboxamid,
- (194) 2-[[2-(terc-butoxi)etil]-amino]-6-[[[(2-klór-6-fluorfenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-1H-benzimidazol-4-karboxamid,
- (195) 6-[[[(2-klór-6-fluorfenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-[(3-metoxi-2,2-dimetilpropil)-amino]-1H-benzimidazol-4-karboxamid,
- (196) 6-[[[(2-klór-6-fluorfenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-[(2-metoxi-2-metilpropil)-amino]-1H-benzimidazol-4-karboxamid,
- (197) 6-[[[(2-klór-6-fluorfenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-[[2-(2S)-tetrahydrofurán-2-il-metil]-amino]-1H-benzimidazol-4-karboxamid,
- (198) 6-[[[(2-klór-6-fluorfenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-[[2-(2R)-tetrahydrofurán-2-il-metil]-amino]-1H-benzimidazol-4-karboxamid,
- (199) 6-[[[(2-klór-6-fluorfenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-[(3-hidroxi-2,2-dimetilpropil)-amino]-1H-benzimidazol-4-karboxamid,
- (200) 6-[[[(2-klór-6-fluorfenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-[[2-(2S)-1-hidroxi-3-metilbután-2-il]-amino]-1H-benzimidazol-4-karboxamid,
- (201) N-(3-klór-4-metilfenil)-2-(dimetilamino)-6-[[2-(trifluorometil)-fenil]-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (202) N-(4-terc-butilfenil)-2-(dimetilamino)-6-[[2-(trifluorometil)-fenil]-karbonil]-amino)-1H-benzimidazol-4-karboxamid,
- (203) N-(2,3-dihidro-1H-inden-5-il)-2-(dimetilamino)-6-[[2-(trifluorometil)-fenil]-karbonil]-amino)-1H-benzimidazol-4-karboxamid,

- (204) 6-[[2-klór-6-fluorfenil]-karbonil]-amino}-N-(3-klór-4-metilfenil)-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (205) N-(3-klór-4-metilfenil)-6-[[2,6-diklórfenil]-karbonil]-amino}-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (206) N-(3-klór-4-metilfenil)-6-[[2,5-diklórfenil]-karbonil]-amino}-2-(dimetilamino)-1H-benzimidazol-4-karboxamid,
- (207) N-(3-klór-2-metilfenil)-2-ciklopropil-6-[[2,5-diklórfenil]-karbonil]-amino}-1H-benzimidazol-4-karboxamid,
- (208) N-(3-klór-4-metilfenil)-2-ciklopropil-6-[[2,5-diklórfenil]-karbonil]-amino}-1H-benzimidazol-4-karboxamid,
- (209) N-(3-klór-2-metilfenil)-6-[[2,5-diklórfenil]-karbonil]-amino}-2-(1-metilciklopropil)-1H-benzimidazol-4-karboxamid,
- (210) N-(3-klór-4-metilfenil)-6-[[2,5-diklórfenil]-karbonil]-amino}-2-(1-metilciklopropil)-1H-benzimidazol-4-karboxamid,
- (211) N-(3-klór-2-metilfenil)-2-(metoximetil)-6-([2-(metilszulfonil)-fenil]-karbonil)-amino}-1H-benzimidazol-4-karboxamid,
- (212) N-(3-klór-4-metilfenil)-6-[[2,5-diklórfenil]-karbonil]-amino}-2-(2-metoxietil)-1H-benzimidazol-4-karboxamid,
- (213) 2-(metoximetil)-N-fenil-6-([2-(trifluorometil)-fenil]-karbonil)-amino}-1H-benzimidazol-4-karboxamid,
- (214) 2-(metoximetil)-N-propil-6-([2-(trifluorometil)-fenil]-karbonil)-amino}-1H-benzimidazol-4-karboxamid,
- (215) 2-(metoximetil)-N-(piridin-3-il)-6-([2-(trifluorometil)-fenil]-karbonil)-amino}-1H-benzimidazol-4-karboxamid,
- (216) N-benzil-2-(metoximetil)-6-([2-(trifluorometil)-fenil]-karbonil)-amino}-1H-benzimidazol-4-karboxamid,
- (217) N-(ciklohexilmetil)-2-(metoximetil)-6-([2-(trifluorometil)-fenil]-karbonil)-amino}-1H-benzimidazol-4-karboxamid,
- (218) 2-(metoximetil)-N-(naftalin-1-il)-6-([2-(trifluorometil)-fenil]-karbonil)-amino}-1H-benzimidazol-4-karboxamid,
- (219) 2-(metoximetil)-N-(tiofén-3-il)-6-([2-(trifluorometil)-fenil]-karbonil)-amino}-1H-benzimidazol-4-karboxamid,

- (220) N-(2,1,3-benzotiadiazol-4-il)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (221) N-(1,1-dioxid-1-benzotiofén-6-il)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (222) 2-(metoximetil)-N-(tiofén-2-ilmetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (223) N-(1H-indol-5-il)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (224) N-(1,3-benzotiazol-2-il)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (225) N-(2,2-dimetilpropil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (226) 2-(metoximetil)-N-(tiofén-2-il)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (227) N-(5-klór-1,3-benzoxazol-2-il)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (228) N-(2-benzilfenil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (229) 2-(metoximetil)-N-(kinolin-8-il)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (230) N-(cikloheptilmetil)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (231) N-(1,3-benzoxazol-2-il)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (232) N-(6-klór-1,3-benzoxazol-2-il)-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (233) N-[3-klór-2-(hidroximetil)-fenil]-2-(metoximetil)-6-([2-(trifluormetil)-fenil]-karbonil)-amino)-1H-benzimidazol-4-karboxamid,
- (234) N-(3-klór-2-metilfenil)-6-[(3-fluorpiridin-2-il)-karbonil]-amino)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (235) N-(3-klór-2-metilfenil)-6-[(3-klórpíridin-4-il)-karbonil]-amino)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,

- (236) N-(3-klór-2-metilfenil)-6-[[[(3,5-diklórpíridin-4-il)-karbonil]-amino]-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (237) 6-[[[(5-butoxi-2-klór-fenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (238) 6-[[[(2-klór-5-(2,2-difluoretóxi)-fenil)-karbonil]-amino]-N-(3-klór-2-metilfenil)-2-(metoximetil)-1H-benzimidazol-4-karboxamid,
- (239) N-(3-klór-2-metilfenil)-6-[[[(2-klór-5-(4,4,4-trifluorbutoxi)-fenil)-karbonil]-amino]-2-(metoximetil)-1H-benzimidazol-4-karboxamid.

8. Gyógyszerkészítmény, mely az 1-7. igénypontok bármelyike szerinti heterociklusos származékok vagy azok tautomerei és gyógyászati lag elfogadható sói csoportból kiválasztott vegyületet tartalmaz hatóanyagként.

9. Az 1-7. igénypontok szerinti heterociklusos származékok, azok tautomerei és gyógyászati lag elfogadható sói csoportból kiválasztott vegyület alkalmazása mPGES gátló szerként.


10. Az 1-7. igénypontok szerinti heterociklusos származékok, azok tautomerei és gyógyászati lag elfogadható sói csoportból kiválasztott vegyület alkalmazása gyulladásos bélbetegségek, irritábilis bél-szindróma, migrén, fejfájás, derékfájás, spinal stenosis, porckorong sérv, temporomandibuláris ízületi betegségek, nyaki gerinc szindróma, nyaki gerinc spondilosis, endometriosis, adenomyosis, koraszülés és vajúdas, fenyegető koraszülés, dysmenorrhea, hiperaktív hólyag, nocturia, interstitial cystitis, neurodegeneratív betegségek, psoriasis, rheumatoid arthritis, reumás láz, fibromyalgia, neuralgia, komplex regionális fájdalom szindróma, fasciális zavar, vírus fertőzés, baktérium fertőzés, mycosis, égés, műtét utáni gyulladás és fájdalom, sérülés és foghúzás, rosszindulatú tumorok, atherosclerosis, stroke, kőszvény, arthritis, osteoarthritis, fiatalkori arthritis, Bechterew spondilitis, inhuvely gyulladás, ligament ossifikáció, szisztémás lupus erythematosus, vasculitis, pancreatitis, nephritis, conjunctivitis, iritis, scleritis, uveitis, sebgyógyítás, dermatitis, ekcéma, osteoporosis, asztma, krónikus obstruktív tüdőbetegségek, tüdő fibrózis, allergiás betegségek, familiáris adenomatosus polyposis, scleroderma, bursitis, uterus leiomyoma, vagy rákos fájdalom megelőzésére vagy kezelésére.

11. A 10. igénypont szerinti alkalmazásra szolgáló vegyület, ahol a neurodegeneratív betegség Alzheimer-kór vagy multiple sclerosis.

12. A 10. igénypont szerinti alkalmazásra szolgáló vegyület, ahol a vírusfertőzés az influenza, megfázás, övsömör vagy AIDS.

13. A 10. igénypont szerinti alkalmazásra szolgáló vegyület, ahol a rosszindulatú tumor vastagbél rák, mellrák, tüdőrák vagy prosztatara k.

A meghatalmazott



Derzi Katalin
szabadságn ügyvivő
SSGK Szabadságn Ügyvivői Iroda
H-1052 Budapest, Andrássy út 113.
Telefon: 461-1000 Fax: 461-4999
Email: derzi@ssgk.hu