

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199933723 B2**
(10) Patent No. **762314**

- (54) Title
Oligonucleotide probes for detecting enterobacteriaceae and
quinolone-resistant enterobacteriaceae
- (51)⁶ International Patent Classification(s)
C12Q 001/68
- (21) Application No: 199933723 (22) Application Date: 1999 . 03 . 30
- (87) WIPO No: WO99/50458
- (30) Priority Data
- | | | |
|-------------|----------------|--------------|
| (31) Number | (32) Date | (33) Country |
| 60/080375 | 1998 . 04 . 01 | US |
- (43) Publication Date : 1999 . 10 . 18
(43) Publication Journal Date : 1999 . 12 . 16
(44) Accepted Journal Date : 2003 . 06 . 19
- (71) Applicant(s)
The Government of the United States of America, as represented by
the Secretary, Department of Health and Human Services
- (72) Inventor(s)
Linda M. Weigel; Fred C. Tenover
- (74) Agent/Attorney
F.B. RICE and CO., 139 Rathdowne Street, CARLTON VIC 3053
- (56) Related Art
US 5645994
BACHOVAL ET AL, 1998, MICROBIAL DRUG RESISTANCE 4(4):271-276
OZEKI ET AL, 1997, J CLINICAL MICROBIOLOGY 35(9):2315-2319

33723/99



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68	A2	(11) International Publication Number: WO 99/50458 (43) International Publication Date: 7 October 1999 (07.10.99)
(21) International Application Number: PCT/US99/06963 (22) International Filing Date: 30 March 1999 (30.03.99) (30) Priority Data: 60/080,375 1 April 1998 (01.04.98) US (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Centers for Disease Control and Prevention, Office of Technology Transfer, Executive Park, Building 4, Suite 1103, Atlanta, GA 30329 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WEIGEL, Linda, M. [US/US]; 1736 Wilson's Crossing Drive, Decatur, GA 30033 (US). TENOVER, Fred, C. [US/US]; 2044 Breckenridge Drive, Atlanta, GA 30345 (US). (74) Agents: GREENE, Jamie, L. et al.; Jones & Askew, LLP, 37th floor, 191 Peachtree Street, N.E., Atlanta, GA 30303 (US).	(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published Without international search report and to be republished upon receipt of that report. IP AUSTRALIA 18 OCT 1999 RECEIVED	
(54) Title: OLIGONUCLEOTIDE PROBES FOR DETECTING ENTEROBACTERIACEAE AND QUINOLONE-RESISTANT ENTEROBACTERIACEAE (57) Abstract Oligonucleotide probes for detecting <i>Enterobacteriaceae</i> species. Unique <i>gyrA</i> coding regions permit the development of probes specific for eight different species: <i>Escherichia coli</i>, <i>Citrobacter freundii</i>, <i>Enterobacter aerogenes</i>, <i>Enterobacter cloacae</i>, <i>Klebsiella oxytoca</i>, <i>Klebsiella pneumoniae</i>, <i>Providencia stuartii</i> and <i>Serratia marcescens</i>. The invention thereby provides methods for the species-specific identification of these <i>Enterobacteriaceae</i> in a sample, and detection and diagnosis of <i>Enterobacteriaceae</i> infection in a subject. Further, nucleic acids are provided for determining quinolone-resistant status of these <i>Enterobacteriaceae</i>.		

5

10

**OLIGONUCLEOTIDE PROBES FOR
DETECTING *Enterobacteriaceae* AND
QUINOLONE-RESISTANT *Enterobacteriaceae***

15 This invention was made in the Centers for Disease Control and Prevention, an agency of the United States Government. The U.S. Government has certain rights in this invention.

Technical Field of the Invention

20 This invention relates in general to the field of diagnostic microbiology. In particular, the invention relates to the species-specific detection of *Enterobacteriaceae*.

Background of the Invention

25 *Enterobacteriaceae* is a family of closely related, Gram-negative organisms associated with gastrointestinal diseases and a wide range of opportunistic infections. They are leading causes of bacteremia and urinary tract infections and are associated with wound infections, pneumonia, meningitis, and various gastrointestinal disorders. (Farmer, J. J., III. *Enterobacteriaceae: Introduction and Identification*. in Murray, P. R., et al., *Manual of Clinical Microbiology*, Washington, D.C., ASM Press, 6th (32): 438-449 (1998)). Many of these infections are life threatening and are often nosocomial (hospital-acquired) infections. (Schaberg et al., *The Am. J. Med.*, 91:72s-75s (1991) and CDC NNIS System Report *Am. J. Infect. Control.*, 24:380-388 (1996)).

30 Conventional methods for isolation and identification of these organisms include growth on selective and/or differential media followed

by biochemical tests of the isolated organism. Total incubation times require 24-48 hours. Slow-growing or fastidious strains require extended incubation times. An additional 18-24 hours is required for susceptibility testing, usually by disk diffusion or broth dilution. More recently, the
5 identification of bacteria by direct hybridization of probes to bacterial genes or by detection of amplified genes has proven to be more time efficient.

Quinolones are broad-spectrum antibacterial agents effective in the treatment of a wide range of infections, particularly those caused by
10 Gram-negative pathogens. (Stein, *Clin. Infect. Diseases*, 23(Suppl 1):S19-24 (1996) and Maxwell, *J. Antimicrob. Chemother.*, 30:409-416 (1992)). For example, nalidixic acid is a first-generation quinolone. Ciprofloxacin is an example of a second generation quinolone, which is also a fluoroquinolone. Sparfloxacin is an example of a third generation
15 quinolone, which is also a fluoroquinolone. As used herein, the term "quinolone" is intended to include this entire spectrum of antibacterial agents, including the fluoroquinolones. This class of antibiotics has many advantages, including oral administration with therapeutic levels attained in most tissues and body fluids, and few drawbacks. As a result,
20 indiscriminate use has led to the currently increasing incidence of quinolone/fluoroquinolone resistance. Hooper, *Adv. Exptl. Medicine and Biology*, 390:49-57 (1995). Mechanisms of resistance to quinolones include alterations in DNA gyrase and/or topoisomerase IV and decreased intracellular accumulation of the antibiotic due to alterations in membrane
25 proteins. (Hooper et al., *Antimicrob. Agents Chemother.*, 36:1151-1154 (1992)).

The primary target of quinolones, including the fluoroquinolones, in Gram-negative bacteria is DNA gyrase, a type II topoisomerase required for DNA replication and transcription. (Cambau et
30 al., *Drugs*, 45(Suppl. 3):15-23 (1993) and Deguchi et al., *J. Antimicrob. Chemother.*, 40:543-549 (1997)). DNA gyrase, composed of two A subunits and two B subunits, is encoded by the *gyrA* and *gyrB* genes. Resistance to quinolones has been shown to be associated most frequently with alterations in *gyrA*. (Yoshida et al., *Antimicrob. Agents
35 Chemother.* 34:1271-1272 (1990)). These mutations are localized at the 5' end of the gene (nucleotides 199-318 in the *E. coli* gene sequence) in an area designated as the quinolone resistance-determining region, or QRDR,

located near the active site of the enzyme, Tyr-122. (Hooper, *Adv. Exptl. Medicine and Biology*, 390:49-57 (1995)).

Previous studies of fluoroquinolone-resistant strains of *Escherichia coli*, *Citrobacter freundii*, *Serratia marcescens* and
5 *Enterobacter cloacae* have revealed that codons 81, 83, and 87 of *gyrA* are the sites most frequently mutated in Gram-negative organisms. (Nishino et al., *FEMS Microbiology Letters*, 154:409-414 (1997), and Kim et al., *Antimicrob. Agents Chemother.*, 42:190-193 (1998)). However, the association of *gyrA* mutations with fluoroquinolone resistance in
10 *Enterobacter aerogenes*, *Klebsiella oxytoca*, and *Providencia stuartii* has not been established.

Previous publications have referred to the use of *gyrA* sequences to identify species within a single genus, such as Husmann et al., *J. Clin. Microbiol.*, 35(9):2398-2400 (1997) for *Campylobacters*, and
15 Guillemin et al., *Antimicrob. Agents Chemo.*, 39(9):2145-2149 (1995) for *Mycobacterium*. The complete gene sequences of DNA gyrase A has previously been published for *Escherichia coli* (Swanberg, et al., *J. Mol. Biol.*, 197:729-736 (1987)) and *Serratia marcescens* (Kim et al., *Antimicrob. Agents Chemother.*, 42:190-193 (1998)). Fragments of *gyrA*
20 including the QRDR have been published for *Enterobacter cloacae* (Deguchi, *J. Antimicrob. Chemother.* 40:543-549 (1997)) and *Citobacter freundii* (Nishino et al., *FEMS Microbiology Letters*, 154:409-414 (1997)). Additionally, the putative *gyrA* sequence for *Klebsiella pneumoniae* was published (Dimri et al., *Nucleic Acids Research*, 18:151-
25 156 (1990)), however, the present invention demonstrates that the most likely organism used in that work was *Klebsiella oxytoca*.

The prior art has not provided enough information about different *Enterobacteriaceae* to develop probes capable of distinguishing
30 between as many species as desirable, nor for determining the quinolone resistance-status of the species. It would be desirable to characterize additional *gyrA* genes and mutations from quinolone-resistant *Enterobacteriaceae* for species-specific identification and quinolone resistance determination using oligonucleotide probes.

Summary of the Invention

The present invention relates to oligonucleotide probes for detecting *Enterobacteriaceae* species. Unique *gyrA* coding regions permit the development of probes specific for identifying eight different species: *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii* and *Serratia marcescens*. The invention thereby provides methods for the species-specific identification of these *Enterobacteriaceae* in a sample, and detection and diagnosis of *Enterobacteriaceae* infection in a subject.

Furthermore, the described unique DNA sequences from the 5' end of *gyrA*, within or flanking the quinolone resistance-determining region, permit the development of probes specific for determining the quinolone-resistant status of eight different species: *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii* and *Serratia marcescens*. The invention thereby provides methods for the species-specific identification of these quinolone-resistant *Enterobacteriaceae*, and detection and diagnosis of quinolone-resistant *Enterobacteriaceae* infection in a subject.

Therefore, it is desirable to provide improved materials and methods for detecting and differentiating *Enterobacteriaceae* species and/or quinolone resistance in the clinical laboratory and research settings.

In a first aspect, the present invention provides an isolated nucleic acid probe, comprising 10 to 50 consecutive nucleotides of one of the *gyrA* genes of *Escherichia coli* (SEQ ID NO: 1), *Enterobacter aerogenes* (SEQ ID NO: 3), *Enterobacter cloacae* (SEQ ID NO: 4), *Klebsiella oxytoca* (SEQ ID NO: 5), *Klebsiella pneumoniae* (SEQ ID NO: 6), or *Providencia stuartii* (SEQ ID NO: 7), wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, but does not hybridize to any of the other *gyrA* genes set forth in SEQ ID NOs: 1-8, wherein the probe identifies an *Enterobacteriaceae* species as *Escherichia coli*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, or *Providencia stuartii*.

In a second aspect, the present invention provides an isolated nucleic acid probe, consisting essentially of 10 to 50 consecutive nucleotides of one of the *gyrA* genes of *Citrobacter freundii* (SEQ ID NO: 2), or *Serratia marcescens* (SEQ ID NO: 8), wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, but does not hybridize to any of the other *gyrA* genes set forth in SEQ ID NOs: 1-8, wherein the

probe identifies an *Enterobacteriaceae* species as *Citrobacter freundii* or *Serratia marcescens*.

In a third aspect, the present invention provides an isolated nucleic acid probe having a nucleic acid sequence of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, or SEQ ID NO: 24, or a complementary sequence thereof.

In a fourth aspect, the present invention provides an isolated nucleic acid probe, comprising 10 to 50 consecutive nucleotides of one of the *gyrA* genes of *Escherichia coli* (SEQ ID NO: 1), *Enterobacter aerogenes* (SEQ ID NO: 3), *Klebsiella oxytoca* (SEQ ID NO: 5), *Klebsiella pneumoniae* (SEQ ID NO: 6), or *Providencia stuartii* (SEQ ID NO: 7), wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, wherein a quinolone susceptible strain selectively hybridizes to the probe, and wherein a quinolone resistant strain has a one or more base pair mismatch with the probe.

In a fifth aspect, the present invention provides an isolated nucleic acid probe, consisting essentially of 10 to 50 consecutive nucleotides of one of the *gyrA* genes of *Citrobacter freundii* (SEQ ID NO: 2), *Enterobacter cloacae* (SEQ ID NO: 4), or *Serratia marcescens* (SEQ ID NO: 8), wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, wherein a quinolone susceptible strain selectively hybridizes to the probe, and wherein a quinolone resistant strain has a one or more base pair mismatch with the probe.

In a sixth aspect, the present invention provides an isolated nucleic acid probe having a nucleic acid sequence of SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32, or SEQ ID NO: 33, or a complementary sequence thereof.

In a seventh aspect, the present invention provides a method of identifying in a sample an *Enterobacteriaceae* species selected from the group consisting of *Escherichia coli* (SEQ ID NO: 1), *Citrobacter freundii* (SEQ ID NO: 2), *Enterobacter aerogenes* (SEQ ID NO: 3), *Enterobacter cloacae* (SEQ ID NO: 4), *Klebsiella oxytoca* (SEQ ID NO: 5), *Klebsiella pneumoniae* (SEQ ID NO: 6), *Providencia stuartii* (SEQ ID NO: 7), and *Serratia marcescens* (SEQ ID NO: 8), comprising combining the sample with the isolated nucleic acid probe according to any one of the first to third aspects, wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, but

does not hybridize to any of the other *gyrA* genes set forth in SEQ ID NOs: 1-8, wherein the presence of hybridization with the single *gyrA* gene identifies the *Enterobacteriaceae* species as *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*,
 5 *Providencia stuartii*, or *Serratia marcescens*.

In an eighth aspect, the present invention provides a method of determining in a sample the quinolone resistance of an *Enterobacteriaceae* species selected from the group consisting of *Escherichia coli* (SEQ ID NO: 1), *Citrobacter freundii* (SEQ ID NO: 2), *Enterobacter aerogenes* (SEQ ID NO: 3), *Enterobacter cloacae* (SEQ ID NO:
 10 4), *Klebsiella oxytoca* (SEQ ID NO: 5), *Klebsiella pneumoniae* (SEQ ID NO: 6), *Providencia stuartii* (SEQ ID NO: 7), and *Serratia marcescens* (SEQ ID NO: 8), comprising combining the sample with the isolated nucleic acid probe according to any one of the fourth to sixth aspects, wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 9-16, or a
 15 complementary sequence thereof, the presence of hybridization with the single *gyrA* gene indicating the quinolone resistance of the species, wherein a quinolone susceptible strain selectively hybridizes to the probe, and wherein a quinolone resistant strain has a one or more base pair mismatch with the probe.

These features and advantages of the present invention will become apparent
 20 after a review of the following detailed description of the disclosed embodiments and the appended claims.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of
 25 any other element, integer or step, or group of elements, integers or steps.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date
 30 of each claim of this application.

Brief Description of the Drawings

5 **Figures 1A and 1B** show the nucleic acid sequence (SEQ ID NOS:1-8) alignments for a portion of the *gyrA* gene in *Escherichia coli* (EC), *Citrobacter freundii* (CF), *Enterobacter aerogenes* (EA), *Enterobacter cloacae* (ECL), *Klebsiella oxytoca* (KO), *Klebsiella pneumoniae* (KP), *Providencia stuartii* (PS) and *Serratia marcescens* (SM).

10 **Figure 2** shows the DNA sequence (SEQ ID NOS:9-16) similarity of the quinolone resistance-determining region (QRDR) in



Escherichia coli, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii* and *Serratia marcescens*.

5 **Figure 3** shows the deduced amino acid sequences (SEQ ID NOS:36-43) of the QRDR for *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii*, and *Serratia marcescens*.

10 **Figures 4A and 4B** show the alterations in GyrA amino acid sequences and susceptibilities of quinolone resistant clinical isolates of *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii*, and *Serratia marcescens*.

15

Detailed Description of the Invention

20 The present invention provides a simple, rapid and useful method for differentiating *Enterobacteriaceae* species and determining their quinolone-resistance status. This invention provides materials and methods to apply the species-specific probes to isolated DNA from host samples for an *in vitro* diagnosis of *Enterobacteriaceae* infection.

25 The present invention provides the nucleic acid sequences of conserved and unique regions of the *gyrA* gene of the following species of the Family *Enterobacteriaceae*: *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii* and *Serratia marcescens*. The present invention provides the nucleic acid sequences of the quinolone resistance-determining region (QRDR) and surrounding regions of *gyrA* of each species listed above.

30 DNA sequence analyses revealed that *gyrA* is unique to each species and highly conserved within the species. However, the *gyrA* mutations resulting in amino acid substitutions which confer quinolone resistance vary in number, type, and position depending on the species. The invention demonstrates that these unique sequences can be used for
35 identification of enteric organisms (genus and species) as well as detection of quinolone resistance within a given species. In addition, comparisons of *Enterobacteriaceae gyrA* with *gyrA* sequences from bacteria not closely

related to *Enterobacteriaceae* species suggest that *gyrA* sequences are unique for all bacterial species and may be used for identification of any species.

The invention provides unique, isolated nucleic acids containing regions of specificity for eight different members of the Family *Enterobacteriaceae*. These nucleic acids are from the *gyrA* gene of the *Enterobacteriaceae* genome. In particular, the invention provides isolated nucleic acids from *Escherichia coli* (SEQ ID NO:1), *Citrobacter freundii* (SEQ ID NO:2), *Enterobacter aerogenes* (SEQ ID NO:3), *Enterobacter cloacae* (SEQ ID NO:4), *Klebsiella oxytoca* (SEQ ID NO:5), *Klebsiella pneumoniae* (SEQ ID NO:6), *Providencia stuartii* (SEQ ID NO:7) and *Serratia marcescens* (SEQ ID NO:8). These sequences can be used to identify and distinguish the respective species of *Enterobacteriaceae*. Figures 1A and 1B show the nucleic acids of SEQ ID NOS:1-8. The sequences correspond to nucleotides #25-613, based on the *E. coli gyrA* sequence numbers of Swanberg et al., *J. Mol. Biol.*, 197:729-736 (1987).

The invention also provides unique, isolated nucleic acids from the quinolone resistance-determining region of *Escherichia coli* (SEQ ID NO:9), *Citrobacter freundii* (SEQ ID NO:10), *Enterobacter aerogenes* (SEQ ID NO:11), *Enterobacter cloacae* (SEQ ID NO:12), *Klebsiella oxytoca* (SEQ ID NO:13), *Klebsiella pneumoniae* (SEQ ID NO:14), *Providencia stuartii* (SEQ ID NO:15) and *Serratia marcescens* (SEQ ID NO:16). These sequences can be used to determine the quinolone resistance status of each species. The QRDR nucleic acids are shown in Figure 2.

Furthermore, the invention provides specific examples of isolated nucleic acid probes derived from the above nucleic acid sequences which may be used as species-specific identifiers of *Escherichia coli* (SEQ ID NO:17), *Citrobacter freundii* (SEQ ID NO:18), *Enterobacter aerogenes* (SEQ ID NO:19), *Enterobacter cloacae* (SEQ ID NO:20), *Klebsiella oxytoca* (SEQ ID NO:21), *Klebsiella pneumoniae* (SEQ ID NO:22), *Providencia stuartii* (SEQ ID NO:23) and *Serratia marcescens* (SEQ ID NO:24).

The invention also provides specific examples of isolated nucleic acid probes derived from the QRDR of the above nucleic acid sequences which may be used as determinants of quinolone resistance for *Escherichia coli* (SEQ ID NOS:25 and 26), *Citrobacter freundii* (SEQ ID

NO:27), *Enterobacter aerogenes* (SEQ ID NO:28), *Enterobacter cloacae* (SEQ ID NO:29), *Klebsiella oxytoca* (SEQ ID NO:30), *Klebsiella pneumoniae* (SEQ ID NO:31), *Providencia stuartii* (SEQ ID NO:32) and *Serratia marcescens* (SEQ ID NO:33).

5 Such probes can be used to selectively hybridize with samples containing nucleic acids from species of *Enterobacteriaceae*. The probes can be incorporated into hybridization assays using polymerase chain reaction, ligase chain reaction, or oligonucleotide arrays on chips or membranes, for example. Additional probes can routinely be derived
10 from the sequences given in SEQ ID NOs:1-8, which are specific for identifying the respective species or for determining quinolone resistance. Therefore, the probes shown in SEQ ID NOs:17-24 and 25-33 are only provided as examples of the species-specific probes or quinolone resistance-determining probes, respectively, that can be derived from SEQ
15 ID NOs:1-8.

By "isolated" is meant nucleic acid free from at least some of the components with which it naturally occurs. By "selective" or "selectively" is meant a sequence that does not hybridize with other nucleic acids to prevent adequate determination of an *Enterobacteriaceae* species or
20 quinolone resistance, depending upon the intended result. As used herein to describe nucleic acids, the term "selectively hybridizes" excludes the occasional randomly hybridizing nucleic acids, and thus has the same meaning as "specifically hybridizing".

A hybridizing nucleic acid should have at least 70%
25 complementarity with the segment of the nucleic acid to which it hybridizes. The selectively hybridizing nucleic acids of the invention can have at least 70%, 80%, 85%, 90%, 95%, 97%, 98%, and 99% complementarity with the segment of the sequence to which it hybridizes. The exemplary probes shown in SEQ ID NOs:17-24 and 25-33 are
30 designed to have 100% hybridization with the target DNA.

The invention contemplates sequences, probes and primers which selectively hybridize to the complementary, or opposite, strand of nucleic acid as those specifically provided herein. Specific hybridization with nucleic acid can occur with minor modifications or substitutions in the
35 nucleic acid, so long as functional species-specific or quinolone resistance determining hybridization capability is maintained. By "probe" is meant a nucleic acid sequence that can be used as a probe or primer for selective

hybridization with complementary nucleic acid sequences for their detection or amplification, which probe can vary in length from about 5 to 100 nucleotides, or preferably from about 10 to 50 nucleotides, or most preferably about 25 nucleotides. The invention provides isolated nucleic acids that selectively hybridize with the species-specific nucleic acids under stringent conditions. *See generally*, Maniatis, et al., *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., (1982) latest edition.

Molecular biology techniques permit the rapid detection of hybridization, such as through confocal laser microscopy and high density oligonucleotide arrays and chips. See, Kozal et al., *Nat. Med.*, 2(7): 753-759 (1996), Schummer et al., *Biotech.*, 23:1087-1092 (1997) or Lockhart et al., *Nat. Biotech.* 14:1675-1680 (1996). Another example of a detection format is the use of controlled electric fields that permit the rapid determination of single base mismatches, as described in Sosnowski et al., *Proc. Natl. Acad. Sci. USA*, 94:1119-1123 (1997). The invention contemplates the use of the disclosed nucleic acid sequences and probes derived therefrom with these currently available techniques and those new techniques discovered in the future.

If used as primers, the invention provides compositions including at least two oligonucleotides (i.e., nucleic acids) that hybridize with different regions of DNA so as to amplify the desired region between the two primers. Depending on the length of the probe or primer, the target region can range between 70% complementary bases and full complementarity and still hybridize under stringent conditions. For example, for the purpose of diagnosing the presence of the *Enterobacteriaceae* in a clinical sample, the degree of complementarity between the nucleic acid (probe or primer) and the target sequence to which it hybridizes (e.g., *Enterobacteriaceae* DNA from a sample) is at least enough to distinguish hybridization with a non-target nucleic acid from other *Enterobacteriaceae*. The invention provides examples of nucleic acids having sequences unique to *Enterobacteriaceae* such that the degree of complementarity required to distinguish selectively hybridizing from nonselectively hybridizing nucleic acids under stringent conditions can be clearly determined for each nucleic acid.

Alternatively, the nucleic acid probes can be designed to have homology with nucleotide sequences present in more than one species of

Enterobacteriaceae. Such a nucleic acid probe can be used to selectively identify a group of *Enterobacteriaceae* species. Additionally, the invention provides that the nucleic acids can be used to differentiate *Enterobacteriaceae* species in general from other species. Such a determination is clinically significant, since therapies for these infections differ.

The invention further provides methods of using the nucleic acids to detect and identify the presence of *Enterobacteriaceae*, or particular species thereof. The methods involve the steps of obtaining a sample suspected of containing *Enterobacteriaceae*. The sample, such as blood, urine, lung lavage fluids, spinal fluid, bone marrow aspiration, vaginal mucosa, tissues, etc., may be taken from an individual, or taken from the environment. The *Enterobacteriaceae* cells in the sample can then be lysed, and the DNA released (or made accessible) for hybridization with oligonucleotide probes.

The DNA sample is preferably amplified prior to hybridization using primers derived from the *gyrA* regions of the *Enterobacteriaceae* DNA that are designed to amplify several species. Examples of such primers are shown below as GYRA6 (SEQ ID NO:34) and or GYRA631R (SEQ ID NO:35). Detection of and/or the determination of quinolone resistance in the target species of *Enterobacteriaceae* is achieved by hybridizing the amplified *gyrA* DNA with an *Enterobacteriaceae* species-specific probe that selectively hybridizes with the DNA. Detection of hybridization is indicative of the presence of the particular species of *Enterobacteriaceae* or quinolone resistance, depending upon the probe. In the case where the species of *Enterobacteriaceae* is known, for example through previous hybridization with a species-specific identifying probe of SEQ ID NOS:17-24, the lack of subsequent hybridization with a species-specific quinolone resistance-determining probe of SEQ ID NOS:25-33 is indicative of quinolone resistance in the sample.

Preferably, detection of nucleic acid hybridization can be facilitated by the use of reporter or detection moieties. For example, the species-specific probes can be labeled with digoxigenin, and a universal-*Enterobacteriaceae* species probe can be labeled with biotin and used in a streptavidin-coated microtiter plate assay. Other examples of detectable

moieties include radioactive labeling, enzyme labeling, and fluorescent labeling.

The invention further contemplates a kit containing one or more species-specific and/or quinolone resistance-determining probes, which can be used for the identification and/or quinolone resistance determination of particular *Enterobacteriaceae* species. Such a kit can also contain the appropriate reagents for hybridizing the probe to the sample and detecting bound probe. The invention may be further demonstrated by the following non-limiting examples.

Examples

Example 1

In this Example, the DNA sequence of the *gyrA* was determined for eight species of *Enterobacteriaceae*. Oligonucleotide primers were designed from conserved *gyrA* gene sequences flanking the QRDR and used to amplify and sequence the 5' region of *gyrA* from ATCC type strains and fluoroquinolone-resistant clinical isolates. The nucleotide and the inferred amino acid sequences were aligned and compared.

The QRDR sequences from 60 clinical isolates with decreased fluoroquinolone susceptibilities were analyzed for alterations associated with fluoroquinolone resistance. The primer sequences at the 3' and 5' ends have been removed leaving nucleotides #25-613, based on the *E. coli gyrA* sequence numbers of Swanberg et al., *J. Mol. Biol.*, 197:729-736 (1987). The organisms, abbreviations and ATCC type strain designation numbers are as follows.

EC = *Escherichia coli* (*E. coli*) ATCC 11775

CF = *Citrobacter freundii* (*C. freundii*) ATCC 8090

EA = *Enterobacter aerogenes* (*E. aerogenes*) ATCC 13048

ECL = *Enterobacter cloacae* (*E. cloacae*) ATCC 13047

KO = *Klebsiella oxytoca* (*K. oxytoca*) ATCC 13182

KP = *Klebsiella pneumoniae* (*K. pneumoniae*) ATCC 13883

PS = *Providencia stuartii* (*P. stuartii*) ATCC 29914

SM = *Serratia marcescens* (*S. marcescens*) ATCC 13880

Amplification of *gyrA*

Bacterial strains and determination of antibiotic susceptibility profiles.

- Type strains of *Enterobacteriaceae* were from American Type Culture Collection (ATCC). Fluoroquinolone resistant and susceptible clinical isolates were selected from the Intensive Care Antimicrobial Resistance Epidemiology (ICARE) study, collected from 39 hospitals across the U.S. between June, 1994 and April 1997 (Archibald et al., *CID*, 24(2):211-215 (1997)). ICARE isolates were screened to exclude duplicate strains from the same patient.
- Minimal inhibitory concentrations (MICs) were determined by the broth microdilution method with cation-adjusted Mueller-Hinton broth according to the methods of the National Committee for Clinical Laboratory Standards (NCCLS M7-A4 (1997)). Ciprofloxacin was purchased from Bayer Corporation (West Haven, CT), ofloxacin and nalidixic acid were from Sigma (St. Louis, MO) and sparfloxacin was from Rhône-Poulenc Rorer (Collegeville, PA).

Amplification of 5' region of *gyrA*.

- Oligonucleotide primers were designed based on homologous regions of *gyrA* sequences in *E. coli* (Swanberg et al., *J. Mol. Biol.*, 1987, 197:729-736) and *K. oxytoca* (published by Dimri et al., *Nuc. Acids Res.*, 1990, 18(1):151-156 as *K. pneumonia*), as follows:

GYRA6

- 5'-CGACCTTGCGAGAGAAAT-3' (SEQ ID NO:34)

GYRA631R

5'-GTTCCATCAGCCCTTCAA-3' (SEQ ID NO:35)

- Primer GYRA6 corresponds to nucleotides 6 to 23 and primer GYRA631R is complementary to nucleotides 610 to 631 of the *E. coli gyrA* sequence.

- DNA fragments were amplified from chromosomal DNA in cell lysates. Amplifications were carried out in a GeneAmp 9600 PCR System (Perkin-Elmer, Applied Biosystems Division, Foster City, CA) in 50 µl volume containing 50 pmol of each primer, 200 µM deoxynucleoside triphosphates, 10 ul cell lysate containing approximately 100 ng template

DNA, 1X reaction buffer with 1.5 mM MgCl₂ and 1 U native Taq polymerase (Perkin Elmer). An initial 4 minute period of denaturation at 94°C was followed by 30 cycles including: denaturation for 1 minute at 94°C, annealing for 30 seconds at 55°C, extending for 45 seconds at 72°C, followed by a final cycle of 72°C for 10 minutes. Amplification products were visualized by agarose gel electrophoresis and ethidium bromide staining to determine specificity and size of gene fragments. PCR products were purified on QIAquick spin columns (QIAGEN, Chatsworth, CA) and sequenced with the ABI Prism Dye Terminator or dRhodamine Terminator Cycle Sequencing Kit and an ABI 377 automated sequencer (Perkin Elmer). To eliminate errors due to amplification artifacts, the forward and reverse sequences of each QRDR were determined using products from independent PCR reactions. The GCG (Genetics Computer Group, Madison, WI) analyses programs were used for the construction of DNA and amino acid sequence alignments.

The resultant sequences of the *gyrA* regions for *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii* and *Serratia marcescens* are shown below in Table 1 and in Figures 1A-1B. The sequences provided correspond to nucleotide positions 25 to 613 of the *gyrA* regions for *Escherichia coli*.

TABLE 1**Gyrase A 5' Region Sequences***Escherichia coli*

ACACCGGT CAACATTGAG GAAGAGCTGA AGAGCTCCTA TCTGGATTAT
GCGATGTCGG TCATTGTTGG CCGTGCGCTG CCAGATGTCC GAGATGGCCT
GAAGCCGGTA CACCGTCGCG TACTTTACGC CATGAACGTA CTAGGCAATG
ACTGGAACAA AGCCTATAAA AAATCTGCCC GTGTCGTTGG TGACGTAATC
GGTAAATACC ATCCCATGG TGA CTGCGG GTTTATGACA CGATCGTCCG
TATGGCGCAG CCATTCTCGC TGC GTTACAT GCTGGTAGAC GGTCAGGGTA
ACTTCGGTTC CATCGACGGC GACTCTGCGG CGGCAATGCG TTATACGGAA
ATCCGTCTGG CGAAAATTGC CCATGAACTG ATGGCTGATC TCGAAAAAGA
GACGGTCGAT TTCGTTGATA ACTATGACGG TACGGAAAAA ATTCCGGACG
TCATGCCAAC CAAAATTCCT AACCTGCTGG TGAACGGTTC TTCCGGTATC

GCCGTAGGTA TGGCAACCAA CATCCCGCCG CACAACCTGA CGGAAGTCAT
CAACGGTTGT CTGGCGTATA TCGATGATGA AGACATCAGC A (SEQ ID
NO:1)

5 *Citrobacter freundii*

ACACCGGT CAACATTGAG GAAGAGCTGA AGAGCTCCTA TCTGGATTAT
GCGATGTCGG TCATTGTTGG CCGTGCGCTG CCAGACGTCC GAGATGGCCT
GAAGCCGGTT CACCGTCGCG TACTTTACGC CATGAACGTA TTGGGCAACG
ACTGGAATAA AGCCTATAAA AAATCTGCCC GTGTCGTTGG TGACGTAATC
10 GGTAATACC ACCCTCATGG TGATACCGCC GTTTACGACA CCATTGTTCC
TATGGCGCAG CCATTCTCCT TGC GTTACAT GCTGGTAGAT GGTCAGGGTA
ACTTTGGTTC TGTCGATGGC GACTCCGCAG CGGCGATGCG TTATACGGAA
ATCCGTATGT CGAAAATCGC CCATGAGCTG ATGGCTGACC TGGAAAAAGA
AACGGTTGAT TTCGTCGATA ACTACGACGG CACCGAACAA ATTCTGACG
15 TCATGCCGAC CAAAATTCCT AACCTGCTGG TGAACGGTTC GTCCGGTATC
GCGGTAGGTA TGGCGACCAA CATTCCGCCG CACAACCTGA CTGAAGTGAT
CAACGGCTGT CTGGCATATA TTGACGATGA AGACATCAGC A (SEQ ID
NO:2)

20 *Enterobacter aerogenes*

ACACGGGT CAACATTGAG GAAGAGCTGA AAAGCTCGTA TCTGGATTAT
GCGATGTCGG TCATTGTTGG CCGTGCGCTG CCGGATGTCC GAGATGGCCT
GAAGCCGGTA CACCGTCGCG TACTATACGC CATGAACGTA TTGGGCAATG
ACTGGAACAA AGCCTATAAA AAATCAGCCC GTGTCGTTGG CGACGTAATC
25 GGTAATACC ACCCGCATGG TGATACCGCC GTTTATGACA CCATCGTACG
TATGGCGCAG CCGTTCTCCT TGC GTTATAT GCTGGTCGAT GGCCAGGGTA
ACTTTGGTTC TGTCGATGGC GACTCCGCTG CAGCGATGCG TTATACGGAA
ATCCGTATGT CGAAGATCGC TCATGAGCTG ATGGCCGATC TCGAAAAAGA
GACGGTTGAT TTCGTCGACA ACTATGACGG CACGGAGAAA ATCCCTGACG
30 TCATGCCGAC AAAAATCCCT AACCTGCTGG TGAACGGTTC TTCCGGTATC
GCCGTAGGTA TGGCGACCAA CATTCCGCCG CATAACCTGA CGGAAGTTAT
CAACGGCTGC CTGGCATACG TTGATAACGA AGACATCAGC A (SEQ ID
NO:3)

35 *Enterobacter cloacae*

ACACCGGTTA ACATCGAGGA AGAGCTGAAG AGCTCCTATC TGGACTATGC
GATGTCGGTC ATTGTTGGCC GTGCGCTGCC GGACGTCCGC GATGGCCTGA
AGCCGGTACA CCGTCGCGTA CTATACGCCA TGAACGTATT GGGCAATGAC

TGGAATAAAG CCTACAAAAA ATCTGCCCCG GTCGTTGGTG ACGTAATCGG
TAAATACCAT CCCCATGGTG ATTCCGCGGT GTACGACACC ATCGTTCGTA
TGGCGCAGCC TTTCTCGCTG CGTTACATGC TGGTAGATGG TCAGGGTAAC
TTTGGTTCTA TCGACGGCGA CTCCGCCGCG GCAATGCGTT ATACGGAAAT
5 CCGTCTGGCG AAAATTGCCC ATGAGCTGAT GGCCGACCTG GAAAAAGAGA
CGGTTGATTT CGTTGATAAC TACGATGGCA CGGAAAAAAT TCCTGACGTC
ATGCCAACGA AGATCCCTAA CCTGCTGGTG AACGGTTCGT CCGGTATCGC
CGTAGGGATG GCGACCAACA TTCCGCCGCA CAACATCACC GAAGTGATCA
ACGGCTGCCT GGCCTATATC GACGATGAAG ACATCAGCA (SEQ ID NO:4)

10

Klebsiella oxytoca

ACACCGGT CAACATTGAG GAAGAGCTGA AGAGCTCCTA TCTGGATTAT
GCGATGTCGG TCATTGTTGG CCGTGCGCTG CCGGATGTCC GAGATGGCCT
GAAGCCGGTA CACCGTCGCG TACTATACGC CATGAACGTA TTGGGCAATG
15 ACTGGAACAA AGCCTATAAA AAATCTGCCC GTGTCGTGGG TGACGTCATC
GGTAAATACC ACCCTCATGG TGATACTGCC GTATACGACA CCATTGTACG
TATGGCGCAG CCATTCTCCC TGC GTTACAT GCTGGTAGAT GGCCAGGGTA
ACTTTGGTTC GGTCGACGGC GACTCCGCCG CAGCGATGCG TTATACGGAA
ATCCGTATGT CGAAGATCGC CCATGAACTG ATGGCCGACC TCGAAAAAGA
20 GACGGTGGAT TTCGTCGATA ACTATGACGG CACGGAGAAA ATCCCTGACG
TTATGCCGAC CAAAATCCCG AACCTGCTAG TCAACGGTTC GTCCGGTATC
GCGGTAGGTA TGGCGACTAA TATTCCGCCG CACAACCTGA CCGAAGTGAT
CAACGGCTGT CTGGCCTACG TTGAAAACGA AGACATCAGC A (SEQ ID
NO:5)

25

Klebsiella pneumoniae

ACACCGGT CAACATTGAG GAAGAGCTTA AGAACTCTTA TCTGGATTAT
GCGATGTCGG TCATTGTTGG CCGTGCGCTG CCGGATGTCC GAGATGGCCT
GAAGCCGGTA CACCGTCGCG TACTTTACGC CATGAACGTA TTGGGCAATG
30 ACTGGAACAA AGCCTATAAA AAATCAGCCC GTGTCGTTGG TGACGTAATC
GGTAAATACC ACCCGCACGG CGACTCCGCG GTATACGACA CCATCGTGCG
TATGGCGCAG CCGTTCTCGC TGC GTTACAT GCTGGTGGAC GGCCAGGGTA
ACTTTGGTTC CATCGACGGC GACTCCGCCG CGGCGATGCG TTATACCGAA
ATTCGTCTGG CGAAAATCGC TCATGAGCTG ATGGCCGATC TTGAAAAAGA
35 GACGGTCGAT TTCGTCGACA ACTATGACGG TACGGAGCGT ATCCGGACG
TCATGCCGAC CAAAATTCCT AACCTGCTGG TGAACGGCGC CTCCGGGATC
GCCGTAGGGA TGGCCACCAA CATACGCCA CATAACCTGA CGGAAGTGAT

TAACGGCTGT CTGGCGTATG TTGACGATGA AGACATCAGC A (SEQ ID NO:6)

Providencia stuartii

5 ACACCGGT CAATATCGAA GAAGAACTCA AAAGTTCGTA TTTGGATTAT
GCGATGTCCG TTATTGTCGG GCGCGCGCTT CCAGATGTTC GAGATGGACT
GAAGCCAGTACACCGCAGAG TACTGTTTGC GATGAATGTA TTGGGAAATG
ATTGGAATAA ACCCTATAAA AAATCTGCCC GTATAGTCGG GGACGTTATC
GGTAAATACC ATCCACATGG TGATAGCGCT GTTTATGAGA CAATCGTTTCG
10 TCTTGCTCAG CCTTTTCTA TCGGTTATAT GCTGGTAGAT GGTCAGGGGA
ACTTTGGTTC AGTTGACGGA GATTCCGCAG CTGCAATGCG TTATACGGAA
ATCCGTATGG CGAAAATTGC CCATGAAATG TTAGCGGATC TTGAAAAAGA
GACCGTTGAT TTCGTCCCAA ACTATGATGG TACAGAGCAA ATCCCTGAAG
TTATGCCTAC GAAAATCCCT AACCTATTGG TTAATGGTTC GTCAGGTATT
15 GCTGTTGGGA TGGCAACGAA CATTCTCCA CACAACCTAG GGAAGTGAT
CAGCGGTTGC CTGCTTATA TAGATGATGA AGATATTAGC A (SEQ ID NO:7)

Serratia marcescens

ACACCGGT AAACATCGAA GACGAGTTGA AAAACTCGTA TCTGGACTAT
20 GCGATGTCCG TTATTGTCGG ACGTGCCCTG CCAGATGTTC GTGATGGACT
GAAGCCGGTT CACCGCCGCG TTCTGTACGC GATGAGCGTA TTGGGTAACG
ACTGGAATAA ACCATACAAG AAATCGGCCC GTGTCGTCGG GGACGTGATC
GGTAAATATC ACCCGCACGG TGACAGCGCG GTTACGACA CTATCGTGCG
TATGGCTCAG CCGTTTTCAC TCGCTACAT GCTGGTGGAC GGTCAGGGTA
25 ACTTCGGTTC CGTCGACGGC GACTCCGCGG CGGCGATGCG TTATACCGAA
GTGCGCATGT CCAAGATTGC TCACGAACTG TTGGCGGATC TGGAAAAAGA
AACCGTCGAC TTCGTGCCTA ACTATGATGG CACCGAGCAG ATCCCGGCCG
TCATGCCGAC CAAGATCCCG AACCTGCTGG TCAACGGCTC GTCGGGCATC
GCCGTGGGCA TGGCTACCAA TATTCCGCCG CACAACCTGG CGGAAGTCGT
30 CAACGGCTGC CTGGCCTATA TCGACGATGA AAACATCAGC A (SEQ ID NO:8)

35 The QRDR sequences from positions 199 to 318 (relative to *E. coli*) are shown below in Table 2.

TABLE 2**Quinolone Resistance-Determining Region Sequences***Escherichia coli*

5 GCGG TGTCGTTGGT GACGTAATCG GTAAATACCA TCCCCATGGT
GACTCGGCCG TTTATGACAC GATCGTCCGT ATGGCGCAGC CATTCTCGCT
GCGTTACATG CTGGTAGACG GTCAG (SEQ ID NO:9)

Citrobacter freundii

10 GCGG TGTCGTTGGT GACGTAATCG GTAAATACCA CCCTCATGGT
GATACCGCCG TTTACGACAC CATTGTTTCGT ATGGCGCAGC CATTCTCCTT
GCGTTACATG CTGGTAGATG GTCAG (SEQ ID NO:10)

Enterobacter aerogenes

15 GC CCGTGTCGTT GGCGACGTAA TCGGTAAATA CCACCCGCAT
GGTGATACCG CCGTTTATGA CACCATCGTA CGTATGGCGC AGCCGTTCTC
CTTGCGTTAT ATGCTGGTCG ATGGCCAG (SEQ ID NO:11)

Enterobacter cloacae

20 GC CCGTGTCGTT GGTGACGTAA TCGGTAAATA CCATCCCCAT
GGTGATTCCG CCGGTGTACGA CACCATCGTT CGTATGGCGC AGCCTTTCTC
GCTGCGTTAC ATGCTGGTAG ATGGTCAG (SEQ ID NO:12)

Klebsiella oxytoca

25 GCGG TGTCGTTGGT GACGTAATCG GTAAATACCA TCCCCATGGT
CTGCCGTATA CGACACCATT GTACGTATGG CGCAGCCATT CTCCCTGCGT
TACATGCTGG TAGATGGCCA G (SEQ ID NO:13)

Klebsiella pneumoniae

30 GC CCGTGTCGTT GGTGACGTAA TCGGTAAATA CCACCCGCAC
GGCGACTCCG CCGTATACGA CACCATCGTG CGTATGGCGC AGCCGTTCTC
GCTGCGTTAC ATGCTGGTGG ACGGCCAG (SEQ ID NO:14)

Providencia stuartii

35 GCGG TGTCGTTGGT GACGTAATCG GTAAATACCA TCCCCATGGT
CGCTGTTTAT GAGACAATCG TTCGTCTTGC TCAGCCTTTT TCTATGCGTT
ATATGCTGGT AGATGGTCAG (SEQ ID NO:15)

Serratia marcescens

GCCCGTGTC GTCGGGGACG TGATCGGTAA ATATCACCCG CACGGTGACA
GCGCGGTTTA CGACACTATC GTGCGTATGG CTCAGCCGTT TTCACTGCGC
TACATGCTGG TGGACGGTCA G (SEQ ID NO:16)

5

Oligonucleotide primers GYRA6 and GYRA631R successfully amplified the expected 626 bp DNA fragment from *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii* and *Serratia marcescens* (Figs. 1A-1B). In additional experiments, amplification with GYRA6 and GYRA631 produced the expected GYRA fragment from *S. typhimurium* (data not shown).

The PCR products were sequenced and the 120 bp regions of *gyrA* known as the QRDR were analyzed. Alignment of the QRDR DNA sequences of the type strains revealed numerous nucleotide substitutions when compared with the *E. coli* sequence (Fig. 2). Eighty-seven of 120 nucleotides (72.5%) were conserved. Similarity to the *E. coli* sequence varied from 93.3% for *E. cloacae* to 80.8% for *P. stuartii* (Figs. 4A-4B). Significant diversity was noted when the *gyrA* QRDR sequences of two species from one genus were aligned. *E. aerogenes* and *E. cloacae* shared 90.5% identity and *K. pneumoniae* and *K. oxytoca* shared 89.3% identity in this region, less similarity than between several of the different genera.

The *gyrA* QRDR sequence of the *E. coli* type strain (ATCC 11775) was compared with the *E. coli* K12 *gyrA* sequence published by Swanberg and Wang (*J. Mol. Biol.* 197:729-736 (1997)) and 4 nucleotide differences were detected at positions 255 (C -> T), 267 (T -> C), 273 (C -> T), and 300 (T -> C).

When the QRDR sequence from the *K. pneumoniae* type strain was compared with the *gyrA* gene sequence from *K. pneumoniae* strain M5a1 published by Dimri and Das (*Nucleic Acids Research*, 18:151-156 (1990)), differences were detected in 15 of 120 nucleotides. Of these 15 nucleotides, only one resulted in an amino acid change. At nucleotide position 247 a T to A change altered the deduced amino acid from Ser-83 (ATCC type strain) to Thr (M5a1). When the M5a1 *gyrA* sequence was compared with that of the *K. oxytoca* type strain, only 4 nucleotide differences were detected. In addition, Ser was consistently found at position 83 in the fluoroquinolone-susceptible strains of *K. pneumoniae*

35

and Thr was consistently found at this position in the *K. oxytoca* strains (Figs. 4A and 4B). These data indicate that the Dimri and Das-sequence of the M5a1 strain most likely was from a strain of *K. oxytoca* and not *K. pneumoniae*.

5 In the sequence from the *S. marcescens* type strain (ATCC 13880), the QRDR was identical to the sequence published by Kim et al. (ATCC 14756)(*Antimicrob. Agents Chemother.*, 42:190-193 (1998)). One nucleotide difference was found in the flanking region (nt 321, T to C) with no change in amino acid sequence (data not shown). The *C. freundii*
10 QRDR sequence was identical to that of Nishino et al. (*FEMS Microbiology Letters*, 154:409-414 (1997)), however, an additional 393 nucleotides are presented herein.

The deduced amino acid sequences of the QRDR were highly conserved (Fig. 3). *E. cloacae*, *K. pneumoniae* and *S. marcescens* shared
15 identical amino acid sequences with *E. coli*. In *C. freundii*, *E. aerogenes* and *K. oxytoca*, one conservative substitution, Ser-83 to Thr was found. Only *P. stuartii* exhibited more than one amino acid substitution in this region. In this organism two conservative changes were detected, Val-69 to Ile and Asp-87 to Glu. In addition, the Leu-92 and Met-98 positions
20 were reversed when compared with the amino acid sequences of other members of the *Enterobacteriaceae* family included in this study. The Glu at position 87 is typical for *gyrA* in Gram-positive organisms (Tankovic et al., *Antimicrob. Agents Chemother.*, 40:2505-2510 (1996)), but has not previously been described for a Gram-negative organism.

25 After determining the DNA sequence of the QRDR from the quinolone-susceptible type strains, the 5' region of *gyrA* in ciprofloxacin-resistant and -susceptible clinical isolates was amplified, sequenced, and analyzed for mutations leading to amino acid changes associated with fluoroquinolone resistance (Figs. 4A and 4B). Comparisons of the
30 fluoroquinolone-susceptible type strain and the resistant clinical isolates of *E. coli* revealed single mutations in codon 83 in *gyrA* associated with low levels of resistance and double mutations (codons 83 and 87) with high levels of resistance (≥ 16 ug/ml ciprofloxacin) as previously described (Vila et al., *Antimicrob. Agents Chemother.*, 38:2477-2479
35 (1994) and Heisig et al., *Antimicrob. Agents Chemother.*, 37:696-701 (1993)). However, in all other species in this study, high levels of resistance were found in strains with single as well as double *gyrA*

mutations. MICs varied significantly among strains with the same mutation, confirming that factors other than *gyrA* are involved in determining the level of resistance to fluoroquinolones (Everett et al., *Antimicrob. Agents Chemother.*, 40:2380-2386 (1996) and Piddock, *Drugs*, 49 (Suppl):29-35 (1995)).

All clinical isolates of *C. freundii* with reduced susceptibility to fluoroquinolones were found to have Thr-83 to Ile mutations, resulting from C-to-T substitutions at nucleotide position 248. Two isolates also displayed alterations of Asp-87 to Gly. However, as noted for isolate *C. freundii* 9023 (Figs. 4A and 4B), the presence of a double mutation was not required for high-level resistance (MICs of 16 µg/ml ciprofloxacin). The nucleotide substitutions in codon 83 of *E. aerogenes gyrA* (Thr-83 to Ile) were identical to those of *C. freundii*. No double mutations were detected in *gyrA* from 7 strains of *E. aerogenes* with reduced levels of susceptibility to fluoroquinolones. However, MICs of isolates with the single mutation ranged from 2 - 16 µg/ml ciprofloxacin.

Clinical isolates of *E. cloacae* exhibited numerous substitutions resulting in Ser-83 changes to Phe, Tyr, or Ile with no single amino acid change associated with either low level or high level resistance. There was no alteration of Ser-83 in the clinical isolate *E. cloacae* 1524 which had a marginal decrease in susceptibility to the fluoroquinolones. However, Asp-87 was changed to Asn. This alteration, found as part of a double mutation in *E. cloacae* 1224, may contribute to high-level resistance if additional changes occur in the QRDR of *E. cloacae* 1524.

K. pneumoniae isolates exhibited either single or double mutations involving Ser-83 and Asp-87, and ciprofloxacin MICs ranged from 1 - 16 µg/ml. Again, double mutations were not required for high-level resistance and no specific mutation (Ser-83 to Phe or Tyr) was associated with low or high levels of fluoroquinolone resistance.

K. oxytoca mutations were confined to the Thr-83 codon and were consistent C-to-T substitutions in the second position resulting in amino acid change to Ile, similar to *C. freundii* and *E. aerogenes*. MICs associated with this alteration ranged from 0.5 - 16 µg/ml ciprofloxacin.

Changes in the QRDR of *P. stuartii gyrA* were also confined to codon 83, however, the nucleotide substitutions varied. The single nucleotide substitutions included A-to-C at the first position or C-to-G at the third position, both resulting in Ser-to-Arg mutations, or G-to-T in the

second position resulting in Ser-to-Ile mutations. MICs ranged from 2 to 16 µg/ml ciprofloxacin.

S. marcescens displayed the greatest diversity in mutations with Gly-81, Ser-83, or Asp-87 involved. No double mutations were
5 detected in the QRDR of *gyrA* from 6 fluoroquinolone-resistant clinical isolates. An unusual mutation of Gly-81 to Cys was found in two isolates. However, this mutation has been described in *E. coli* (Yoshida et al., *Antimicrob. Agents Chemother.*, 34:1271-1272 (1990)).

The data in this Example provides for the first time enough
10 comparative nucleic acid sequence data for the *gyrA* gene to enable one to prepare probes that will selectively hybridize to target nucleic acid to identify the species and/or quinolone resistance of *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii* and
15 *Serratia marcescens*.

Example 2

Development of Probes

Identification of *Enterobacteriaceae* Species

20 Oligonucleotide probes can be selected for species-specific identification of *Enterobacteriaceae* in or near the QRDR of *gyrA*. The region which includes the codons most often associated with fluoroquinolone resistance (nucleotides 239-263) was not used for the reason that if identification were based on one or more nucleotide changes,
25 the changes associated with resistance would interfere with identification. Each probe for identification was selected for maximum difference, and it is recognized that a smaller region within some probes could be used, based on single base changes. However, most of the probes have at least two nucleotide differences compared with the same region in other strains.
30 When there were variations, other than those associated with resistance, within the susceptible and/or the resistance strains for any given species, the position of the probe was shifted to a region which was completely conserved for all strains sequenced. For this reason, the probes were in the region 5' of the QRDR.

35

TABLE 3**Oligonucleotide probes for identification of *Enterobacteriaceae***

- 5 *E. coli* 5' ACT TTA CGC CAT GAA CGT ACT AGG C 3'
(SEQ ID NO:17) (144-168)
- C. freundii* 5' TGG GCA ACG ACT GGA ATA AAG CC 3'
(SEQ ID NO:18) (164-186)
- 10 *E. aerogenes* 5' TTA TAT GCT GGT CGA TGG CCA G 3'
(SEQ ID NO:19) (297-323)
- E. cloacae* 5' GCC GGA CGT CCG CGA TGG CCT 3'
(SEQ ID NO:20) (102-122)
- 15 *K. oxytoca* 5' GTA GAT GGC CAG GGT AAC TTT GGT TCG
GTC 3' (SEQ ID NO:21) (307-336)
- K. pneumoniae* 5' GTG CGT ATG GCG CAG CCG TTC TCG CTG 3'
20 (SEQ ID NO:22) (268-294)
- P. stuartii* 5' CGT CTT GCT CAG CCT TTT TCT ATG C 3'
(SEQ ID NO:23) (271-295)
- 25 *S. marcescens* 5' GGA ATA AAC CAT ACA AGA AA 3'
(SEQ ID NO:24) (176-195)

Note: Numbers in parentheses refer to base positions in *E. coli* sequence

30 **Fluoroquinolone resistance probes**

- Simultaneous identification of the species and mutations leading to resistance can be determined by using one of the above oligonucleotide probes in combination with the resistance probes set forth below. All oligonucleotide probes shown in Table 4 for quinolone resistance span the region containing the amino acid codons most frequently associated with resistance (nucleotides 239-263). Susceptible strains will hybridize to the resistance probe for that species and resistance
- 35

will be detected as one or more basepair mismatch with the susceptible strain sequence.

TABLE 4

Oligonucleotide probes for quinolone resistance in
Enterobacteriaceae

5	<i>E. coli</i>	5' ATG GTG ACT CGG CGG TTT ATG ACA C 3'
	(SEQ ID NO:25)	
10	OR	5' ATG GTG ACT CGG CGG TCT ATG ACA C 3'
	(SEQ ID NO:26)	
	<i>C. freundii</i>	5' ATG GTG ATA CCG CCG TTT ACG ACA C 3'
	(SEQ ID NO:27)	
15	<i>E. aerogenes</i>	5' ATG GTG ATA CCG CCG TTT ATG ACA C 3'
	(SEQ ID NO:28)	
	<i>E. cloacae</i>	5' ATG GTG ATT CCG CGG TGT ACG ACA C 3'
20	(SEQ ID NO:29)	
	<i>K. oxytoca</i>	5' ATG GTG ATA CTG CCG TAT ACG ACA C 3'
	(SEQ ID NO:30)	
25	<i>K. pneumoniae</i>	5' ACG GCG ACT CCG CGG TAT ACG ACA C 3'
	(SEQ ID NO:31)	
	<i>P. stuartii</i>	5' ATG GTG ATA GCG CTG TTT ATG AGA C 3'
	(SEQ ID NO:32)	
30	<i>S. marcescens</i>	5' ACG GTG ACA GCG CGG TTT ACG ACA C 3'
	(SEQ ID NO:33)	

EDITORIAL NOTE

APPLICATION NUMBER - 33723/99

The following Sequence Listing pages 1 to 8 are part of the description. The claims pages follow on pages "23" to "29".

SEQUENCE LISTING

<110> The Government of the United States of America . .

<120> Oligonucleotide Probes for Detecting Enterobacteriaceae
and Quinolone-Resistant Enterbacteriaceae

<130> 03063-0430wp

<140>

<141>

<150> 60/080375

<151> 1998-04-01

<160> 35

<170> PatentIn Ver. 2.0

<210> 1

<211> 589

<212> DNA

<213> Escherichia coli

<400> 1

```
acaccggtca acattgagga agagctgaag agctcctatc tggattatgc gatgtcggtc 60
attgttggcc gtgcgctgcc agatgtccga gatggcctga agccggtaca ccgtcgcgta 120
ctttacgcca tgaacgtact aggcaatgac tggaacaaaag cctataaaaa atctgcccgt 180
gtcgttgggtg acgtaatcgg taaataccat ccccatgggtg actcggcggt ttatgacacg 240
atcgtccgta tggcgagcc attctcgtg cgttacatgc tggtagacgg tcagggtaac 300
ttcggttcca tcgacggcga ctctcgccg gcaatgcgtt atacggaaat ccgtctggcg 360
aaaattgccc atgaactgat ggctgatctc gaaaaagaga cggtcgattt cgttgataac 420
tatgacggta cggaaaaaat tccggacgtc atgccaacca aaattcctaa cctgctgggtg 480
aacggttctt ccggtatcgc cgtaggtatg gcaaccaaca tcccgccgca caacctgacg 540
gaagtcacga acggttgctt ggcgtatata gatgatgaag acatcagca 589
```

<210> 2

<211> 589

<212> DNA

<213> Citrobacter freundii

<400> 2

```
acaccggtca acattgagga agagctgaag agctcctatc tggattatgc gatgtcggtc 60
attgttggcc gtgcgctgcc agacgtccga gatggcctga agccggttca ccgtcgcgta 120
ctttacgcca tgaacgtatt gggcaacgac tggaataaag cctataaaaa atctgcccgt 180
gtcgttgggtg acgtaatcgg taaataccac cctcatgggtg ataccgccgt ttacgacacc 240
attgttcgta tggcgagcc attctcctt cgttacatgc tggtagatgg tcagggtaac 300
tttggttctg tcgatggcga ctccgagcg gcgatgcgtt atacggaaat ccgtatgtcg 360
```

aaaatcgccc atgagctgat ggctgacctg gaaaaagaaa cggttgattt cgtcgataac 420
tacgacggca ccgaacaaat tcctgacgtc atgccgacca aaattccctaa cctgctgggtg 480
aacggttcgt ccggtatcgc ggtaggtatg gcgaccaaca ttccgccgca caacctgact 540
gaagtgatca acggctgtct ggcataatg gacgatgaag acatcagca 589

<210> 3

<211> 589

<212> DNA

<213> Enterobacter aerogenes

<400> 3

acacgggtca acattgagga agagctgaaa agctcgtatc tggattatgc gatgtcggtc 60
attgttgccc gtgcgctgcc ggatgtccga gatggcctga agccgggtaca ccgtcgcgta 120
ctatacgcca tgaacgtatt gggcaatgac tggaacaaaag cctataaaaa atcagcccgt 180
gtcgttgccg acgtaatcgg taaataccac ccgcatgggtg ataccgccgt ttatgacacc 240
atcgtacgta tggcgcagcc gttctccttg cgttatatgc tggtcgatgg ccagggtaac 300
tttggttctg tcgatggcga ctccgctgca gcgatgcgtt atacggaaat ccgtatgtcg 360
aaagtcgctc atgagctgat ggcgcatctc gaaaaagaga cggttgattt cgtcgacaac 420
tatgacggca cggagaaaaat ccctgacgtc atgccgacaa aaatccctaa cctgctgggtg 480
aacggttctt ccggtatcgc cgtaggtatg gcgaccaaca ttccgccgca taacctgacg 540
gaagttatca acggctgcct ggcatacgtt gataacgaag acatcagca 589

<210> 4

<211> 589

<212> DNA

<213> Enterobacter cloacae

<400> 4

acaccggtta acatcgagga agagctgaag agctcctatc tggactatgc gatgtcggtc 60
attgttgccc gtgcgctgcc ggacgtccgc gatggcctga agccgggtaca ccgtcgcgta 120
ctatacgcca tgaacgtatt gggcaatgac tggaataaaag cctacaaaaa atctgcccgt 180
gtcgttggtg acgtaatcgg taaataccat ccccatgggtg attccgcggt gtacgacacc 240
atcgttcgta tggcgcagcc tttctcgctg cgttacatgc tggtagatgg tcagggtaac 300
tttggttcta tcgacggcga ctccgccgcg gcaatgcgtt atacggaaat ccgtctggcg 360
aaaattgccc atgagctgat ggcgacctg gaaaaagaga cggttgattt cgttgataac 420
tacgatggca cggaaaaaat tcctgacgtc atgccaacga agatccctaa cctgctgggtg 480
aacggttcgt ccggtatcgc cgtagggatg gcgaccaaca ttccgccgca caacatcacc 540
gaagtgatca acggctgcct ggcctatctc gacgatgaag acatcagca 589

<210> 5

<211> 589

<212> DNA

<213> Klebsiella oxytoca

<400> 5

acaccgggtca acattgagga agagctgaag agctcctatc tggattatgc gatgtcggtc 60
attgttgccc gtgcgctgcc ggatgtccga gatggcctga agccgggtaca ccgtcgcgta 120
ctatacgcca tgaacgtatt gggcaatgac tggaacaaaag cctataaaaa atctgcccgt 180

gtcgtgggtg acgtcatcgg taaataccac cctcatggtg atactgccgt atacgacacc 240
attgtacgta tggcgagcc attctccctg cgttacatgc tggtagatgg ccagggtaac 300
tttggttcgg tcgacggcga ctccgccgca gcgatgcgtt atacggaaat ccgtatgtcg 360
aagatcgccc atgaactgat ggccgacctc gaaaaagaga cgggtggattt cgtcgataac 420
tatgacggca cggagaaaaa cctcgacgtt atgccgacca aaatcccgaa cctgctagtc 480
aacggttcgt ccggtatcgc ggtagggtatg gcgactaata ttccgccgca caacctgacc 540
gaagtgatca acggctgtct ggcctacgtt gaaaacgaag acatcagca 589

<210> 6

<211> 589

<212> DNA

<213> *Klebsiella pneumoniae*

<400> 6

acaccgggtca acattgagga agagcttaag aactcttattc tggattatgc gatgtcggtc 60
attgttggcc gtgcgctgcc ggatgtccga gatggcctga agccgggtaca ccgtcgcgta 120
ctttacgccca tgaacgtatt gggcaatgac tggacaaaag cctataaaaa atcagcccgt 180
gtcgttgggtg acgtaatcgg taaataccac ccgcacggcg actccgcggt atacgacacc 240
atcgtcgta tggcgagcc gttctcgctg cgttacatgc tggtagatgg ccagggtaac 300
tttggttcca tcgacggcga ctccgccgcg gcgatgcgtt ataccgaaat tcgtctggcg 360
aaaatcgctc atgagctgat ggccgatctt gaaaaagaga cggtcgattt cgtcgacaac 420
tatgacggta cggagcgtat tccggacgtc atgccgacca aaatccctaa cctgctgggt 480
aacggcgccct ccgggacgcg cgtagggtatg gccaccaaca taccgccaca taacctgacg 540
gaagtgatta acggctgtct ggcgtatggt gacgatgaag acatcagca 589

<210> 7

<211> 589

<212> DNA

<213> *Providencia stuartii*

<400> 7

acaccgggtca atacgaaga agaactcaaa agttcgtatt tggattatgc gatgtccggt 60
attgtcgggc gcgcgcttcc agatgttcga gatggactga agccagtaca ccgcagagta 120
ctgtttcga tgaatgtatt gggaaatgat tggaaataaac cctataaaaa atctgccgt 180
atagtcgggg acgttatcgg taaataccat ccacatgggtg atagcgctgt ttatgagaca 240
atcgttcgtc ttgctcagcc tttttctatg cgttatatgc tggtagatgg tcagggggaa 300
tttggttcag ttgacggaga ttccgcagct gcaatgcgtt atacggaaat ccgtatggcg 360
aaaattgccc atgaaatgtt agcggatctt gaaaaagaga cgtttgattt cgtcccaaac 420
tatgatggta cagagcaaat ccctgaagtt atgcctacga aaatccctaa cctattggtt 480
aatggttcgt caggtattgc tgttgggatg gcaacgaaca ttccctccaca caacctaggg 540
gaagtgatca gcggttcgct tgcttatata gatgatgaag atattagca 589

<210> 8

<211> 589

<212> DNA

<213> *Serratia marcescens*

<400> 8

acaccggtaa acatcgaaga cgagttgaaa aactcgtatc tggactatgc gatgtccgtt 60
attgtcggac gtgccctgcc agatgttcgt gatggactga agccggttca ccgccgcgtt 120
ctgtacgcga tgagcgtatt gggtaacgac tggataaac catacaagaa atcggcccggt 180
gtcgtcgggg acgtgatcgg taaatatcac ccgcacggtg acagcgcggt ttacgacact 240
atcgtgcgta tggctcagcc gttttcactg cgctacatgc tggtaggacgg tcagggtaac 300
ttcggttccg tcgacggcga ctcccgcgcg gcgatgcgtt ataccgaagt gcgcatgtcc 360
aagattgtc acgaactgtt ggcgatctg gaaaaagaaa ccgtcgactt cgtgcctaac 420
tatgatggca ccgagcagat ccggccgctc atgccgacca agatccgaa cctgctggtc 480
aacggctcgt cgggcatcgc cgtgggcatg gctaccaata ttccgcgcga caacctggcg 540
gaagtctca acggtgcct ggctatatc gacgatgaaa acatcagca 589

<210> 9

<211> 120

<212> DNA

<213> Escherichia coli

<400> 9

gcccgtgtcg ttggtgacgt aatcggtaaa taccatcccc atggtgactc ggcggtttat 60
gacacgatcg tccgtatggc gcagccattc tcgctgcgtt acatgctggt agacggtcag 120

<210> 10

<211> 120

<212> DNA

<213> Citrobacter freundii

<400> 10

gcccgtgtcg ttggtgacgt aatcggtaaa taccaccctc atggtgatac cgcggtttac 60
gacaccattg ttcgtatggc gcagccattc tccttgcggtt acatgctggt agatggtcag 120

<210> 11

<211> 120

<212> DNA

<213> Enterobacter aerogenes

<400> 11

gcccgtgtcg ttggcgacgt aatcggtaaa taccaccgcg atggtgatac cgcggtttat 60
gacaccatcg tacgtatggc gcagccgttc tccttgcggtt atatgctggt cgtatggcag 120

<210> 12

<211> 120

<212> DNA

<213> Enterobacter cloacae

<400> 12

gcccgtgtcg ttggtgacgt aatcggtaaa taccatcccc atggtgattc cgcggtgtac 60
gacaccatcg ttcgtatggc gcagccttcc tcgctgcgtt acatgctggt agatggtcag 120

<210> 13

<211> 120
<212> DNA
<213> *Klebsiella oxytoca*

<400> 13
gcccgtgtcg tgggtgacgt catcggtaaa taccaccctc atggtgatac tgccgtatac 60
gacaccattg tacgtatggc gcagccattc tccctgcggt acatgctggt agatggccag 120

<210> 14
<211> 120
<212> DNA
<213> *Klebsiella pneumoniae*

<400> 14
gcccgtgtcg ttggtgacgt aatcggtaaa taccaccgc acggcgactc cgcggtatac 60
gacaccatcg tgcgtatggc gcagccgttc tcgctgcggt acatgctggt ggacggccag 120

<210> 15
<211> 120
<212> DNA
<213> *Providencia stuartii*

<400> 15
gcccgtatag tcggggacgt tatcggtaaa taccatccac atggtgatag cgctgtttat 60
gagacaatcg ttcgtcttgc tcagcctttt tctatgcggt atatgctggt agatggtcag 120

<210> 16
<211> 120
<212> DNA
<213> *Serratia marcescens*

<400> 16
gcccgtgtcg tcggggacgt gatcggtaaa taccaccgc acggtgacag cgcggtttac 60
gacactatcg tgcgtatggc tcagccgttt tcaactgcgt acatgctggt ggacggtcag 120

<210> 17
<211> 25
<212> DNA
<213> *Escherichia coli*

<400> 17
actttacgcc atgaacgtac taggc 25

<210> 18
<211> 23
<212> DNA
<213> *Citrobacter freundii*

<400> 18
tgggcaacga ctggaataaa gcc 23

<210> 19
<211> 22
<212> DNA
<213> Enterobacter aerogenes

<400> 19
ttatatgctg gtcgatggcc ag 22

<210> 20
<211> 21
<212> DNA
<213> Enterobacter cloacae

<400> 20
gccggacgtc cgcgatggcc t 21

<210> 21
<211> 30
<212> DNA
<213> Klebsiella oxytoca

<400> 21
gtagatggcc agggtaactt tggttcggtc 30

<210> 22
<211> 27
<212> DNA
<213> Klebsiella pneumoniae

<400> 22
gtgcgtatgg cgcagccgtt ctcgctg 27

<210> 23
<211> 25
<212> DNA
<213> Providencia stuartii

<400> 23
cgtcttgctc agcctttttc tatgc 25

<210> 24
<211> 20
<212> DNA
<213> Serratia marcescens

<400> 24	
ggaataaacc atacaagaaa	20
<210> 25	
<211> 25	
<212> DNA	
<213> Escherichia coli	
<400> 25	
atggtgactc ggcggtttat gacac	25
<210> 26	
<211> 25	
<212> DNA	
<213> Escherichia coli	
<400> 26	
atggtgactc ggcggtttat gacac	25
<210> 27	
<211> 25	
<212> DNA	
<213> Citrobacter freundii	
<400> 27	
atggtgatac cgccgtttac gacac	25
<210> 28	
<211> 25	
<212> DNA	
<213> Enterobacter aerogenes	
<400> 28	
atggtgatac cgccgtttat gacac	25
<210> 29	
<211> 25	
<212> DNA	
<213> Enterobacter cloacae	
<400> 29	
atggtgattc cgccgtgtac gacac	25
<210> 30	
<211> 25	
<212> DNA	
<213> Klebsiella oxytoca	

<400> 30 atggtgatac tgccgtatac gacac	25
<210> 31 <211> 25 <212> DNA <213> <i>Klebsiella pneumoniae</i>	
<400> 31 acggcgactc cgcggtatac gacac	25
<210> 32 <211> 25 <212> DNA <213> <i>Providencia stuartii</i>	
<400> 32 atggtgatag cgctgtttat gagac	25
<210> 33 <211> 25 <212> DNA <213> <i>Serratia marcescens</i>	
<400> 33 acggtgacag cgcggtttac gacac	25
<210> 34 <211> 18 <212> DNA <213> <i>Enterobacter</i> sp.	
<400> 34 cgaccttgcg agagaaat	18
<210> 35 <211> 18 <212> DNA <213> <i>Enterobacter</i> sp.	
<400> 35 gttccatcag ccttcaa	18

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. An isolated nucleic acid probe, comprising 10 to 50 consecutive nucleotides of one of the *gyrA* genes of *Escherichia coli* (SEQ ID NO: 1), *Enterobacter aerogenes* (SEQ ID NO: 3), *Enterobacter cloacae* (SEQ ID NO: 4), *Klebsiella oxytoca* (SEQ ID NO: 5), *Klebsiella pneumoniae* (SEQ ID NO: 6), or *Providencia stuartii* (SEQ ID NO: 7), wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, but does not hybridize to any of the other *gyrA* genes set forth in SEQ ID NOs: 1-8, wherein the probe identifies an *Enterobacteriaceae* species as *Escherichia coli*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, or *Providencia stuartii*.
2. The isolated nucleic acid probe of claim 1, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Escherichia coli* (SEQ ID NO: 1), or a complementary sequence thereof.
3. The isolated nucleic acid probe of claim 1, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Enterobacter aerogenes* (SEQ ID NO: 3), or a complementary sequence thereof.
4. The isolated nucleic acid probe of claim 1, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Enterobacter cloacae* (SEQ ID NO: 4), or a complementary sequence thereof.
5. The isolated nucleic acid probe of claim 1, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Klebsiella oxytoca* (SEQ ID NO: 5), or a complementary sequence thereof.
6. The isolated nucleic acid probe of claim 1, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Klebsiella pneumoniae* (SEQ ID NO: 6), or a complementary sequence thereof.
7. The isolated nucleic acid probe of claim 1, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Providencia stuartii* (SEQ ID NO: 7), or a complementary sequence thereof.

8. An isolated nucleic acid probe, consisting essentially of 10 to 50 consecutive nucleotides of one of the *gyrA* genes of *Citrobacter freundii* (SEQ ID NO: 2), or *Serratia marcescens* (SEQ ID NO: 8), wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, but does not hybridize to any of the other *gyrA* genes set forth in SEQ ID NOs: 1-8, wherein the probe identifies an *Enterobacteriaceae* species as *Citrobacter freundii* or *Serratia marcescens*.
9. The isolated nucleic acid probe of claim 8, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Citrobacter freundii* (SEQ ID NO: 2), or a complementary sequence thereof.
10. The isolated nucleic acid probe of claim 8, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Serratia marcescens* (SEQ ID NO: 8), or a complementary sequence thereof.
11. An isolated nucleic acid probe having a nucleic acid sequence of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, or SEQ ID NO: 24, or a complementary sequence thereof.
12. An isolated nucleic acid probe, comprising 10 to 50 consecutive nucleotides of one of the *gyrA* genes of *Escherichia coli* (SEQ ID NO: 1), *Enterobacter aerogenes* (SEQ ID NO: 3), *Klebsiella oxytoca* (SEQ ID NO: 5), *Klebsiella pneumoniae* (SEQ ID NO: 6), or *Providencia stuartii* (SEQ ID NO: 7), wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, wherein a quinolone susceptible strain selectively hybridizes to the probe, and wherein a quinolone resistant strain has a one or more base pair mismatch with the probe.
13. The isolated nucleic acid probe of claim 12, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Escherichia coli* (SEQ ID NO: 1), or a complementary sequence thereof.
14. The isolated nucleic acid probe of claim 12, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Enterobacter aerogenes* (SEQ ID NO: 3), or a complementary sequence thereof.

15. The isolated nucleic acid probe of claim 12, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Klebsiella oxytoca* (SEQ ID NO: 5), or a complementary sequence thereof.
- 5 16. The isolated nucleic acid probe of claim 12, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Klebsiella pneumoniae* (SEQ ID NO: 6), or a complementary sequence thereof.
17. The isolated nucleic acid probe of claim 12, wherein the probe selectively
10 hybridizes under stringent conditions to the *gyrA* gene of *Providencia stuartii* (SEQ ID NO: 7), or a complementary sequence thereof.
18. An isolated nucleic acid probe, consisting essentially of 10 to 50 consecutive nucleotides of one of the *gyrA* genes of *Citrobacter freundii* (SEQ ID NO: 2),
15 *Enterobacter cloacae* (SEQ ID NO: 4), or *Serratia marcescens* (SEQ ID NO: 8), wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, wherein a quinolone susceptible strain selectively hybridizes to the probe, and wherein a quinolone resistant strain has a one or more base pair mismatch with the probe.
20
19. The isolated nucleic acid probe of claim 18, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Citrobacter freundii* (SEQ ID NO: 2), or a complementary sequence thereof.
- 25 20. The isolated nucleic acid probe of claim 18, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Enterobacter cloacae* (SEQ ID NO: 4), or a complementary sequence thereof.
21. The isolated nucleic acid probe of claim 18, wherein the probe selectively
30 hybridizes under stringent conditions to the *gyrA* gene of *Serratia marcescens* (SEQ ID NO: 8), or a complementary sequence thereof.
22. An isolated nucleic acid probe having a nucleic acid sequence of SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32, or SEQ ID NO: 33, or a complementary sequence thereof.

23. A method of identifying in a sample an *Enterobacteriaceae* species selected from the group consisting of *Escherichia coli* (SEQ ID NO: 1), *Citrobacter freundii* (SEQ ID NO: 2), *Enterobacter aerogenes* (SEQ ID NO: 3), *Enterobacter cloacae* (SEQ ID NO: 4), *Klebsiella oxytoca* (SEQ ID NO: 5), *Klebsiella pneumoniae* (SEQ ID NO: 6), *Providencia stuartii* (SEQ ID NO: 7), and *Serratia marcescens* (SEQ ID NO: 8), comprising combining the sample with the isolated nucleic acid probe according to any one of claims 1 to 11, wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 1-8, or a complementary sequence thereof, but does not hybridize to any of the other *gyrA* genes set forth in SEQ ID NOs: 1-8, wherein the presence of hybridization with the single *gyrA* gene identifies the *Enterobacteriaceae* species as *Escherichia coli*, *Citrobacter freundii*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Providencia stuartii*, or *Serratia marcescens*.
24. The method of claim 23, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Escherichia coli* (SEQ ID NO: 1), or a complementary sequence thereof, the presence of hybridization identifying *Escherichia coli* in the sample.
25. The method of claim 23, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Citrobacter freundii* (SEQ ID NO: 2), or a complementary sequence thereof, the presence of hybridization identifying *Citrobacter freundii* in the sample.
26. The method of claim 23, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Enterobacter aerogenes* (SEQ ID NO: 3), or a complementary sequence thereof, the presence of hybridization identifying *Enterobacter aerogenes* in the sample.
27. The method of claim 23, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Enterobacter cloacae* (SEQ ID NO: 4), or a complementary sequence thereof, the presence of hybridization identifying *Enterobacter cloacae* in the sample.

28. The method of claim 23, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Klebsiella oxytoca* (SEQ ID NO: 5), or a complementary sequence thereof, the presence of hybridization identifying *Klebsiella oxytoca* in the sample.

5

29. The method of claim 23, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Klebsiella pneumoniae* (SEQ ID NO: 6), or a complementary sequence thereof, the presence of hybridization identifying *Klebsiella pneumoniae* in the sample.

10

30. The method of claim 23, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Providencia stuartii* (SEQ ID NO: 7), or a complementary sequence thereof, the presence of hybridization identifying *Providencia stuartii* in the sample.

15

31. The method of claim 23, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Serratia marcescens* (SEQ ID NO: 8), or a complementary sequence thereof, the presence of hybridization identifying *Serratia marcescens* in the sample.

20

32. A method of determining in a sample the quinolone resistance of an *Enterobacteriaceae* species selected from the group consisting of *Escherichia coli* (SEQ ID NO: 1), *Citrobacter freundii* (SEQ ID NO: 2), *Enterobacter aerogenes* (SEQ ID NO: 3), *Enterobacter cloacae* (SEQ ID NO: 4), *Klebsiella oxytoca* (SEQ ID NO: 5),
 25 *Klebsiella pneumoniae* (SEQ ID NO: 6), *Providencia stuartii* (SEQ ID NO: 7), and *Serratia marcescens* (SEQ ID NO: 8), comprising combining the sample with the isolated nucleic acid probe according to any one of claims 12 to 22, wherein the probe selectively hybridizes under stringent conditions to a single *gyrA* gene set forth in SEQ ID NOs: 9-16, or a complementary sequence thereof, the presence of hybridization with
 30 the single *gyrA* gene indicating the quinolone resistance of the species, wherein a quinolone susceptible strain selectively hybridizes to the probe, and wherein a quinolone resistant strain has a one or more base pair mismatch with the probe.

35

33. The method of claim 32, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Escherichia coli* (SEQ ID NO: 9), or a complementary sequence thereof, the presence of hybridization indicating the quinolone resistance of the *Escherichia coli* in the sample.

5

34. The method of claim 32, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Citrobacter freundii* (SEQ ID NO: 10), or a complementary sequence thereof, the presence of hybridization indicating the quinolone resistance of the *Citrobacter freundii* in the sample.

10

35. The method of claim 32, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Enterobacter aerogenes* (SEQ ID NO: 11), or a complementary sequence thereof, the presence of hybridization indicating the quinolone resistance of the *Enterobacter aerogenes* in the sample.

15

36. The method of claim 32, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Enterobacter cloacae* (SEQ ID NO: 12), or a complementary sequence thereof, the presence of hybridization indicating the quinolone resistance of the *Enterobacter cloacae* in the sample.

20

37. The method of claim 32, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Klebsiella oxytoca* (SEQ ID NO: 13), or a complementary sequence thereof, the presence of hybridization indicating the quinolone resistance of the *Klebsiella oxytoca* in the sample.

25

38. The method of claim 32, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Klebsiella pneumoniae* (SEQ ID NO: 14), or a complementary sequence thereof, the presence of hybridization indicating the quinolone resistance of the *Klebsiella pneumoniae* in the sample.

30

39. The method of claim 32, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Providencia stuartii* (SEQ ID NO: 15), or a complementary sequence thereof, the presence of hybridization indicating the quinolone resistance of the *Providencia stuartii* in the sample.

35

40. The method of claim 32, wherein the probe selectively hybridizes under stringent conditions to the *gyrA* gene of *Serratia marcescens* (SEQ ID NO: 16), or a complementary sequence thereof, the presence of hybridization indicating the quinolone resistance of the *Serratia marcescens* in the sample.

5

Dated this twentieth day of March 2003

The Government of the United States of
America, represented by The Secretary of
the Department of Health and Human
Services

Patent Attorneys for the Applicant:

F B RICE & CO

2003

2003

1/6

EC-11775.DNA 25: ACACCGGTCAACATTGAGGAAGAGCTGAAGAGCTCCTATCTGGATTATGCGATGTCGGTC
CF-8090.DNA
EA-13048.DNAG.....A.....G.....
ECL13047.DNAT.....C.....C.....
KO-13182.DNA
KP-13883.DNAT.....A.....T.....
PS-29914.DNAT.....C.....A.....C.....A.....T.....G.....T.....C.....T.....
SM-13880.DNAA.....C.....A.....C.....T.....A.....A.....G.....C.....C.....T.....

EC-11775.DNA 85: ATTGTTGGCCGTGCGCTGCCAGATGTCGAGATGGCCTGAAGCCGGTACACCGTCGCGTA
CF-8090.DNAC.....T.....
EA-13048.DNAG.....
ECL13047.DNAG.....C.....C.....
KO-13182.DNAG.....
KP-13883.DNAG.....
PS-29914.DNAC.....G.....C.....T.....T.....A.....A.....CA.....A.....
SM-13880.DNAC.....A.....C.....T.....T.....A.....T.....C.....T.....

EC-11775.DNA 145: CTTTACGCCATGAACGTACTAGGCAATGACTGGAACAAAGCCTATAAAAAATCTGCCCGT
CF-8090.DNAT.G.....C.....T.....
EA-13048.DNAA.....T.G.....A.....
ECL13047.DNAA.....T.G.....T.....C.....
KO-13182.DNAA.....T.G.....
KP-13883.DNAT.G.....A.....
PS-29914.DNAG.TT..G.....T.....T.G..A.....T.....T.....C.....
SM-13880.DNAG.....G.....T.G..T..C.....T.....C.A..C..G.....G.....

EC-11775.DNA 205: GTCGTTGGTGACGTAATCGGTAAATACCATCCCCATGGTGACTCGGCGGTTTATGACACG
CF-8090.DNAC..T.....TA.C..C.....C.....C.....
EA-13048.DNAC.....C..G.....TA.C..C.....C.....C.....
ECL13047.DNAT.....C.....G..C.....C.....
KO-13182.DNAG.....C.....C..T.....TA.T..C..A..C.....C.....
KP-13883.DNAC.....C..G..C..C.....C.....A..C.....C.....
PS-29914.DNA A.A..C..G.....T.....A.....TAGC..T.....G..A.....
SM-13880.DNAC..G.....G.....T..C..G..C.....AGC.....C.....T.....

EC-11775.DNA 265: ATCGTCCGTATGGCGCAGCCATTCTCGCTGCGTTACATGCTGGTAGACGGTCAGGGTAAC
CF-8090.DNA ..T..T.....CT.....T.....
EA-13048.DNAA.....G.....CT.....T.....C..T..C.....
ECL13047.DNAT.....T.....T.....
KO-13182.DNA ..T..A.....C.....T..C.....
KP-13883.DNAG.....G.....G.....C.....
PS-29914.DNAT..C..T..T.....T..T..TA.....T.....T.....G..
SM-13880.DNAG.....T.....G..T..A.....C.....G.....

Fig - 1A

SUBSTITUTE SHEET (RULE 26)

2/6


EC-11775.DNA 325: TTCGGTTCCATCGACGGCGACTCTGCGGCGGCAATGCGTTATACGGAAATCCGTCTGGCG
CF-8090.DNA ..T....TG....T.....C..A....G.....A..T..
EA-13048.DNA ..T....TG....T.....C..T..A..G.....A..T..
ECL13047.DNA ..T....T.....C..C.....A..T..
KO-13182.DNA ..T....GG.....C..C..A..G.....A..T..
KP-13883.DNA ..T.....C..C.....G.....C.....T..
PS-29914.DNA ..T....AG.T....A..T..C..A..T.....A..
SM-13880.DNAG.....C.....G.....C..G..CA..T.C
** ***** * * * * *

EC-11775.DNA 385: AAAATTGCCCATGAACTGATGGCTGATCTCGAAAAAGAGACGGTCGATTTCGTTGATAAC
CF-8090.DNAC.....G.....C..G.....A....T.....C..
EA-13048.DNA ..G..C..T....G.....C.....T.....C..C..
ECL13047.DNAG.....C..C..G.....T.....
KO-13182.DNA ..G..C.....C..C.....G.....C..
KP-13883.DNAC..T....G.....C..T.....C..C..
PS-29914.DNAA..T..A..G....T.....C..T.....CCCA..
SM-13880.DNA ..G....T..C.....T...G.....G.....A..C....GCC..
** *

EC-11775.DNA 445: TATGACGGTACGGAAAAATTCGGACGTCATGCCAACCAAAATTCCTAACCTGCTGGTG
CF-8090.DNA ..C....C..C..C.....T.....G.....
EA-13048.DNAC....G....C..T.....G..A....C.....
ECL13047.DNA ..C..T..C.....T.....G..G..C.....
KO-13182.DNAC.....G....C..T....T....G.....C..G.....A..C
KP-13883.DNAGCGT.....G.....
PS-29914.DNAT....A..GC....C..T..A..T....T..G.....C.....AT...T
SM-13880.DNAT..C..C..GC.G..C....C.....G....G..C..G.....C
** *

EC-11775.DNA 505: AACGGTTCTTCCGGTATCGCCGTAGGTATGGCAACCAACATCCCGCCGCACAACTGACG
CF-8090.DNAG.....G.....T.....T.....
EA-13048.DNAG.....G.....T.....T.....
ECL13047.DNAG.....G.....G.....T.....A..C..C
KO-13182.DNAG.....G.....G.....T.....T.....C
KP-13883.DNACG.C....G.....G....C.....A....A..T.....
PS-29914.DNA ..T....G..A....T..T..T..G.....G....T..T..A.....AGG.
SM-13880.DNAC..G..G..C.....G..C....T....T.....G..
** *

EC-11775.DNA 565: GAAGTCATCAACGGTTGTCTGGCGTATATCGATGATGAAGACATCAGCA
CF-8090.DNAG.....C.....A....T..C..
EA-13048.DNAT.....C..C....A..CG.T..A.C.....
ECL13047.DNAG.....C..C....C.....C.....
KO-13182.DNAG.....C.....C..CG.T..AA.C.....
KP-13883.DNAG..T....C.....G.T..C.....
PS-29914.DNAG....G.....C..T..T....A.....T..T..
SM-13880.DNAG.....C..C....C.....C.....A..
***** * * * * * * * * * * * * * * * * *

 1B

SUBSTITUTE SHEET (RULE 26)

DNA Sequence Similarity of the QDR* in Enterobacteriaceae

	199	S83	258
<i>E. coli</i>	**	GCCCGTGTGTTGGTGACGTATCGGTAATACCATCCCATGGTGACTCGGCGTTTAT	
<i>C. freundii</i>		C..T..TA.C.C.....C	
<i>E. aerogenes</i>		C.....C.G.....TA.C.C.....	
<i>E. cloacae</i>		G.....C.....T.C.....G.C	
<i>K. oxytoca</i>		A.A.C.G.....T.....TA.T.C.A.C	
<i>K. pneumoniae</i>		A.A.C.G.....T.....TAGC.T.....	
<i>P. stuartii</i>		C.G.....G.....T.C.G.C.....AGC.....C	
<i>S. marcescens</i>		***** ** ** ***** ** ** ** ** ** - ** ** *	
<i>E. coli</i>		GACACGATCGTCCGTATGGCGCAGCCATTCTCGTGTACATGCTGTAGACGGTCAG	
<i>C. freundii</i>		C.T.T.....CT.....T.....	
<i>E. aerogenes</i>		C.....A.....G.....CT.....C.T.C.....	
<i>E. cloacae</i>		C.....T.....T.....T.....	
<i>K. oxytoca</i>		C.T.A.....C.....T.C.....	
<i>K. pneumoniae</i>		C.....G.....G.....G.....C.....	
<i>P. stuartii</i>		G.A.....T.C.T.T.....T.T.TA.....T.....	
<i>S. marcescens</i>		T.....G.....T.....G.T.A.....C.....G.....	
		** ** ** ** * ** ***** ** ** - ** ** ** ** - ** ** *	

* Yoshida et al., 1990. Antimicrob. Agents Chemother. 34:1271.

** Swanberg and Wang, 1987. J. Mol. Biol. 197:729.

Similarity of the Amino Acid Sequences of the QRDR
of Enterobacteriaceae

	67	83	87	106
<i>E. coli</i>	ARVVG	DVIGKY	PHGDS	SAVYDTIVRMAQPFSLRYMLVDGQ
<i>C. freundii</i>				
<i>E. aerogenes</i>			T.....	
<i>E. cloacae</i>			T.....	
<i>K. oxytoca</i>			T.....	
<i>K. pneumoniae</i>				
<i>P. stuartii</i>		I.....		E.....L.....M.....
<i>S. marcescens</i>				
	**	*****	*****	*****

5/6

Strain	MIC (µg/ml) ¹			Amino acid change ²		
	CIP	OFLX	SPAR	GyrA		
				81	83	87 (codon)
<i>C. freundii</i>						
ATCC 8090	≤0.12	≤0.25	≤0.008	Gly(GGT)	Thr(ACC)	Asp(GAC)
Cf 7377	2	8	>2	-	Ile(ATC)	-
Cf 0759	2	8	>2	-	Ile(ATC)	-
Cf 9085	4	8	2	-	Ile(ATC)	-
Cf 9417	8	8	>2	-	Ile(ATC)	-
Cf 1958	≥16	≥16	>2	-	Ile(ATC)	Gly(GGC)
Cf 5757	≥16	≥16	>2	-	Ile(ATC)	Gly(GGC)
Cf 9023	≥16	≥16	>2	-	Ile(ATC)	-
<i>E. aerogenes</i>						
ATCC 13048	0.5	1	0.25	Gly(GGT)	Thr(ACC)	Asp(GAC)
Ea 1747	≤0.12	<0.25	0.06	-	-	-
Ea 2786	2	4	2	-	Ile(ATC)	-
Ea 9032	4	4	2	-	Ile(ATC)	-
Ea 5593	8	8	>2	-	Ile(ATC)	-
Ea 9433	8	≥16	>2	-	Ile(ATC)	-
Ea 3521	8	≥16	>2	-	Ile(ATC)	-
Ea 5590	8	≥16	>2	-	Ile(ATC)	-
Ea 2775	≥16	32	>2	-	Ile(ATC)	-
<i>E. cloacae</i>						
ATCC 13047	≤0.12	≤0.25	0.03	Gly(GGT)	Ser(TCC)	Asp(GAC)
Ecl 1700	≤0.12	<0.25	0.06	-	-	-
Ecl 1524	0.25	0.5	0.12	-	-	Asn(AAC)
Ecl 1963	2	4	2	-	Phe(TTC)	-
Ecl 1286	2	4	1	-	Tyr(TAC)	-
Ecl 3529	4	8	>2	-	Thr(ACC)	-
Ecl 1544	8	8	>2	-	Tyr(TAC)	-
Ecl 1627	8	≥16	>2	-	Tyr(TAC)	-
Ecl 9028	≥16	8	>2	-	Ile(ATC)	-
Ecl 1224	≥16	8	>2	-	Phe(TTC)	Asn(AAC)
Ecl 1251	≥16	≥16	>2	-	Ile(ATC)	-
Ecl 63	≥16	≥16	>2	-	Phe(TTC)	-
Ecl 105	≥16	≥16	>2	-	Phe(TTC)	-
Ecl 1783	≥16	≥16	>2	-	Tyr(TAC)	-
Ecl 9030	≥16	≥16	>2	-	Tyr(TAC)	-
Ecl 9031	≥16	>32	>2	-	Ile(ATC)	-
<i>E. coli</i>						
ATCC 11775	≤0.12	≤0.25	0.03	Gly(GGT)	Ser(TCG)	Asp(GAC)
Ec 748	≤0.12	≤0.25	0.03	-	-	-
Ec 3535	1	2	2	-	Leu(TTG)	-
Ec 5524	2	8	2	-	Leu(TTG)	-
Ec 9419	≥16	32	>2	-	Leu(TTG)	Gly(GGA)
Ec 9421	≥16	32	>2	-	Leu(TTG)	Tyr(TAC)
Ec 9425	≥16	>32	>2	-	Leu(TTG)	Asn(AAC)



SUBSTITUTE SHEET (RULE 26)

6/6

Strain	MIC ($\mu\text{g/ml}$)			Amino acid change		
	CIP	OFLX	SPAR	81	GyrA 83	87 (codon)
<i>K. pneumoniae</i>						
ATCC 13883	0.5	≤ 0.25	0.06	Gly(GGC)	Ser(TCC)	Asp(GAC)
Kp 570	≤ 0.12	≤ 0.25	0.06	-	-	-
Kp 1961	≤ 0.12	≤ 0.25	0.03	-	-	-
Kp 2778	≤ 0.12	≤ 0.25	0.03	-	-	-
Kp 2790	≤ 0.12	≤ 0.25	0.03	-	-	-
Kp 1361	1	2	0.5	-	Phe(TTC)	-
Kp 1362	1	2	1	-	Phe(TTC)	-
Kp 1177	4	≥ 16	> 2	-	Phe(TTC)	-
Kp 682	≥ 16	≥ 16	> 2	-	Phe(TTC)	-
Kp 1768	≥ 16	32	> 2	-	Tyr(TAC)	-
Kp 1775	≥ 16	> 32	> 2	-	Phe(TTC)	Gly(GGC)
Kp 1565	≥ 16	> 32	> 2	-	Tyr(TAC)	Asn(AAC)
<i>K. oxytoca</i>						
ATCC 13182	≤ 0.12	≤ 0.25	0.12	Gly(GGT)	Thr(ACT)	Asp(GAC)
Ko 702	≤ 0.12	≤ 0.25	0.03	-	-	-
Ko 2538	≤ 0.12	≤ 0.25	0.06	-	-	-
Ko 2110	≤ 0.12	≤ 0.25	0.06	-	-	-
Ko 1199	≤ 0.12	≤ 0.25	0.06	-	-	-
Ko 57	0.5	0.05	0.25	-	Ile(ATT)	-
Ko 1577	4	4	2	-	Ile(ATT)	-
Ko 1817	8	≥ 16	> 2	-	Ile(ATT)	-
Ko 466	≥ 16	8	> 2	-	Ile(ATT)	-
Ko 32	≥ 16	≥ 16	> 2	-	Ile(ATT)	-
Ko 1578	≥ 16	≥ 16	> 2	-	Ile(ATT)	-
Ko 1612	≥ 16	≥ 16	> 2	-	Ile(ATT)	-
Ko 2777	≥ 16	> 32	> 2	-	Ile(ATT)	-
<i>P. stuartii</i>						
ATCC 29914	≤ 0.12	≤ 0.25	≤ 0.12	Gly(GGT)	Ser(AGC)	Glu(GAG)
Ps 1571	0.25	1	0.5	-	-	-
Ps 1284	2	4	2	-	Arg(CGC)	-
Ps 2469	8	≥ 16	> 2	-	Ile(ATC)	-
Ps 2783	≥ 16	≥ 16	2	-	Arg(AGG)	-
Ps 9428	≥ 16	≥ 16	> 2	-	Ile(ATC)	-
Ps 7375	≥ 16	32	> 2	-	Ile(ATC)	-
Ps 2468	≥ 16	32	> 2	-	Ile(ATC)	-
Ps 1773	≥ 16	> 32	> 2	-	Arg(AGG)	-
<i>S. marcescens</i>						
ATCC 13880	0.5	1	1	Gly(GGT)	Ser(AGC)	Asp(GAC)
Sm 1714	0.25	1	1	-	-	-
Sm 9745	2	4	2	-	Ile(ATC)	-
Sm 1221	4	8	> 2	Cys(TGT)	-	-
Sm 1969	4	≥ 16	> 2	-	-	Asn(AAC)
Sm 1570	8	8	> 2	Cys(TGT)	-	-
Sm 5591	8	≥ 16	> 2	-	Arg(AGA)	-
Sm 1568	≥ 16	≥ 16	> 2	-	Arg(CGC)	-

¹ CIP, ciprofloxacin; OFLX, ofloxacin; SPAR, sparfloxacin² -, identical to the ATCC type strain


SUBSTITUTE SHEET (RULE 26)