



(86) Date de dépôt PCT/PCT Filing Date: 1996/03/28

(87) Date publication PCT/PCT Publication Date: 1996/10/03

(45) Date de délivrance/Issue Date: 2001/12/18

(85) Entrée phase nationale/National Entry: 1997/08/25

(86) N° demande PCT/PCT Application No.: JP 96/00820

(87) N° publication PCT/PCT Publication No.: WO 96/30339

(30) Priorité/Priority: 1995/03/31 (PCT/JP95/00631) WO

(51) Cl.Int.⁶/Int.Cl.⁶ C07D 207/12, A61K 31/695, A61K 31/44, A61K 31/40, C07D 409/14, C07D 401/12, C07F 7/10, C07D 405/14, C07D 295/13, C07D 409/12, C07D 405/12


(72) Inventeur/Inventor:
ITO, Fumitaka, JP

(73) Propriétaire/Owner:
PFIZER INC., US

(74) Agent: SMART & BIGGAR

(54) Titre : COMPOSES D'ACIDE PYRROLIDINYL-HYDROXAMIQUE ET LEUR PROCEDE DE PRODUCTION

(54) Title: PYRROLIDINYL HYDROXAMIC ACID COMPOUNDS AND THEIR PRODUCTION PROCESS



(57) **Abrégé/Abstract:**

A compound of formula (I) and its pharmaceutically acceptable salt, wherein A is hydrogen, hydroxy or OY, wherein Y is a hydroxy protecting group; Ar is phenyl optionally substituted with one or more substituents selected from halo, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, CF₃, C₁-C₄ alkoxy-C₁-C₄ alkyloxy, and carboxy-C₁-C₄ alkyloxy; X is phenyl, naphthyl, biphenyl, indanyl, benzofuranyl, benzothiophenyl, 1-tetralone-6-yl, C₁-C₄ alkylenedioxy, pyridyl, furyl and thienyl, these groups optionally being substituted with up to three substituents selected from halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, NO₂, CF₃ and SO₂CH₃; and R is hydrogen, C₁-C₄ alkyl or a hydroxy protecting group. These compounds and pharmaceutical compositions containing them are useful as analgesic, anti-inflammatory, diuretic, anesthetic or neuroprotective agents, or an agent for stroke or treatment of functional bowel diseases such as abdominal pain, for the treatment of a mammalian subject, especially a human subject. Further, the present invention provides processes for producing the hydroxamic compounds of formula (I) and their intermediate compounds of formula (II).


**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 207/12, 295/12, 401/06, 405/06, A61K 31/40		A1	(11) International Publication Number: WO 96/30339
			(43) International Publication Date: 3 October 1996 (03.10.96)
(21) International Application Number: PCT/JP96/00820 (22) International Filing Date: 28 March 1996 (28.03.96) (30) Priority Data: PCT/JP95/00631 31 March 1995 (31.03.95) WO (34) Country for which the regional or international application was filed: JP (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017-5755 (US). (71) Applicant (for JP only): PFIZER PHARMACEUTICALS INC. [JP/JP]; Shinjuku Mitsui Building, 1-1, Nishi-shinjuku 2-chome, Shinjuku-ku, Tokyo 163-04 (JP). (72) Inventor; and (75) Inventor/Applicant (for US only): ITO, Fumitaka [JP/JP]; 2-71-1, Nashinoki, Taketoyo-cho, Chita-gun, Aichi-ken 470-23 (JP). (74) Agent: MURAKAMI, Kiyoshi; Yuasa and Hara, Section 206, New Ohtemachi Building, 2-1, Ohtemachi 2-chome, Chiyoda-ku, Tokyo 100 (JP).		(81) Designated States: AU, BG, BR, BY, CA, CN, CZ, HU, IS, JP, KR, KZ, LV, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	

(54) Title: PYRROLIDINYL HYDROXAMIC ACID COMPOUNDS AND THEIR PRODUCTION PROCESS**(57) Abstract**

A compound of formula (I) and its pharmaceutically acceptable salt, wherein A is hydrogen, hydroxy or OY, wherein Y is a hydroxy protecting group; Ar is phenyl optionally substituted with one or more substituents selected from halo, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, CF₃, C₁-C₄ alkoxy-C₁-C₄ alkyloxy, and carboxy-C₁-C₄ alkyloxy; X is phenyl, naphthyl, biphenyl, indanyl, benzofuranyl, benzothiophenyl, 1-tetralone-6-yl, C₁-C₄ alkylenedioxy, pyridyl, furyl and thienyl, these groups optionally being substituted with up to three substituents selected from halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, NO₂, CF₃ and SO₂CH₃; and R is hydrogen, C₁-C₄ alkyl or a hydroxy protecting group. These compounds and pharmaceutical compositions containing them are useful as analgesic, anti-inflammatory, diuretic, anesthetic or neuroprotective agents, or an agent for stroke or treatment of functional bowel diseases such as abdominal pain, for the treatment of a mammalian subject, especially a human subject. Further, the present invention provides processes for producing the hydroxamic compounds of formula (I) and their intermediate compounds of formula (II).



PYRROLIDINYL HYDROXAMIC ACID COMPOUNDS AND THEIR
PRODUCTION PROCESS

Technical Field

5 This invention relates to novel hydroxamic acid derivatives and their pharmaceutically acceptable salts, and to pharmaceutical compositions containing them. These compounds and compositions are useful as analgesic, antiinflammatory, diuretic, anesthetic or neuroprotective agents, or an agent for treatment of stroke or functional bowel diseases such as abdominal pain, for the treatment of a mammalian subject, especially a human subject.

10

Background Art

15 Opioid analgesics such as morphine are therapeutically useful, but their usage is strictly limited because of their side effects such as drug dependency. Thus, analgesics with high usefulness and reduced tendency to cause drug dependency are desired. Considerable pharmacological and biochemical studies have been carried out to discover the opioid peptides and opioid receptors, and the discovery of the subtype of opioid receptor such as μ , δ , κ at a peripheral nerve in a variety of species, including human, has made a beginning towards creating new analgesics. As it is thought that opioid analgesics such as morphine act as a μ -receptor agonist, separating the action based on a κ -receptor agonist from the action based on μ -receptor agonist has been investigated. Recently κ -selective agonists have been reported from the above viewpoint for example, EMD-60400: A. Barber et al., Naunyn-Schmied. Arch. Pharmacol., 345 (Suppl.): Abst 456. Some of them actually have been studied in clinical trials (Med. Res. Rev., 12, 525 (1992)).


20

25 However, even when a selective κ -receptor agonist is employed, use of high doses can give rise to side effects such as sedation. Therefore, it would be desired to provide compounds having better agonist activity toward opioid κ -receptor, and in particular compounds having only low sedative activity.

Brief Disclosure of the Invention

The present invention provides a compound of the following formula:

2



(I)

and the salts thereof, wherein

A is hydrogen, hydroxy or OY, wherein Y is a hydroxy-protecting group;

Ar is phenyl optionally substituted with one or more (preferably up to three) substituents selected from halo, hydroxyl, C₁-C₄ alkyl, C₁-C₄ alkoxy, CF₃, C₁-C₄ alkoxy-C₁-C₄ alkoxy and carboxy-C₁-C₄ alkoxy;

X is phenyl, naphthyl, biphenyl, indanyl, benzofuranyl, benzothiophenyl, 1-tetralone-6-yl, pyridyl, furyl or thienyl, each being optionally substituted with up to three substituents selected from C₁-C₄ alkylenedioxy, halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxyl, NO₂, CF₃ and SO₂CH₃; and

R is hydrogen, C₁-C₄ alkyl or a hydroxy-protecting group.

The hydroxamic acid derivatives of the present invention of formula (I), wherein A is hydrogen or hydroxyl and R is hydrogen or C₁-C₄ alkyl, exhibit significant agonist activity toward opioid κ -receptor. Therefore these κ agonists are particularly useful as an analgesic agent in mammals, especially humans. They are also useful as antiinflammatory, diuretic, anesthetic or neuroprotective

2a

agents, or an agent for treatment of stroke or functional bowel diseases such as abdominal pain, for the treatment of a mammalian subject, especially a human subject.

Accordingly, the present invention also provides a pharmaceutical composition useful as an analgesic, antiinflammatory, diuretic, anesthetic or neuroprotective agent, or an agent for treatment of stroke or functional bowel diseases such as abdominal pain, for the treatment of a mammalian subject, especially a human subject, which
10 comprises a therapeutically effective amount of a hydroxamic acid of formula (I), wherein A is hydrogen or hydroxyl and R is hydrogen or C₁-C₄ alkyl, or its pharmaceutically acceptable salt together with a pharmaceutically acceptable carrier.

The compounds of formula (I), wherein either or both of OY and OR represent a protected hydroxyl group, are useful as chemical intermediates to the κ agonist of

formula (I). Typical hydroxy protecting groups are benzyl, triphenylmethyl, tetrahydropyranyl, methoxymethyl and $R^1R^2R^3Si$, wherein R^1, R^2 and R^3 are each C_1 - C_6 alkyl or phenyl.

5 A preferred group of κ agonists compounds of the present invention consists of the compounds of formula (I), wherein A is hydrogen or hydroxy, Ar is phenyl, X is phenyl substituted with up to three substituents selected from chloro, methyl and CF_3 , more preferably 3,4-dichlorophenyl, and R is hydrogen. The preferred configuration of the carbon atom to which the group Ar is attached is (S).

Preferred individual compounds of the invention are:

10 2-(3,4-Dichlorophenyl)-N-hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]acetamide;

N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-(2,3,6-trichlorophenyl)acetamide;

15 N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-(4-trifluoromethylphenyl)acetamide;

N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-(2,4,6-trimethylphenyl)acetamide;

2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

20 2-(4-Bromophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(4-trifluoromethylphenyl)acetamide;

25 2-(4-Chlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

2-(2,3-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

2-(2,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

30 2-(2,5-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-

phenylethyl]acetamide;


2-(2,6-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

5 N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(2,3,6-trichlorophenyl)acetamide;

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide; and

2-(3,4-Dimethylphenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

10 Further, the present invention provides a compound of the formula:



and the salts thereof, wherein

A is hydrogen, hydroxy or OY, wherein Y is hydroxy protecting group;

15 Ar is phenyl optionally substituted with one or more substituents selected from halo, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, CF₃, C₁-C₄ alkoxy-C₁-C₄ alkyloxy, and carboxy-C₁-C₄ alkyloxy;

R is hydrogen, C₁-C₄ alkyl or a hydroxy protecting group.


20 These compounds of formula (II) can be used as intermediates to prepare the compounds of formula (I).

Further, the present invention provides processes for producing the hydroxamic compounds of formula (I) and their intermediate compounds of formula (II).

25 Detailed Disclosure of the Invention

The κ agonists of formula (I) of this invention can be prepared by a numbers of methods. For example, they can be readily prepared according to the procedure shown in Scheme (I).

Scheme 1



Thus, the κ agonists compounds of formula (I), wherein A is hydrogen or hydroxy and R is hydrogen, can be prepared by reaction of a compound of the formula (VI) with a carboxylic acid of the formula XCH_2COOH , followed by removal of the protecting group P, and the protecting group in A¹ if necessary. This is a conventional acylation reaction, which can be carried out using standard methods, well-known to

those skilled in the art. However, a convenient way of acylating a compound of formula (VI) with an acid of the formula XCH_2COOH comprises coupling the two compounds in the presence of a carbodiimide compound. An especially convenient carbodiimide compound is 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, which is sometimes referred to as water-soluble carbodiimide, or WSC. This reaction is carried out by contacting substantially equivalent amounts of the acid and amine with a small excess of the carbodiimide in an appropriate solvent at a temperature in the range from -30 to $100^\circ C$, usually from 0 to $30^\circ C$. Appropriate solvents are inactive aromatic hydrocarbons, ethers, halogenated hydrocarbons, especially dichloromethane. The reaction takes about 30 minutes to 24 hours, usually 30 minutes to 3 hours at room temperature. The product can be isolated and purified by standard techniques.

The protecting group P, and any protecting group in A^1 , is removed by the appropriate method for the particular protecting group chosen. Thus, a typical protecting group is benzyl. This can be removed by catalytic hydrogenation. Appropriate catalysts for hydrogenation are Pd/C, Pearlman's catalyst, Pd black, or Pd/BaSO₄, especially 10% Pd/C.

A further convenient protecting group for P and A^1 is the tetrahydropyranyl group (THP). This can be removed by acid-catalyzed hydrolysis. Appropriate acid catalysts are organic acid, inorganic acid, or Lewis acid such as AcOH, *p*-TsOH, HCl, Me₂AlCl etc., especially HCl.


The κ agonist compounds of formula (I), wherein R is a C₁-C₄ alkyl group, can be prepared by alkylation of the corresponding compounds of formula (I), wherein R is hydroxy. This alkylation can be carried out by standard methods. A particularly convenient method involves base catalyzed alkylation using alkyl halide in the presence of phase transfer catalyst such as tetra-n-butylammonium hydrogen sulfate. The intermediate hydroxylamine of the formula (VI) can be prepared from the alcohol (V), by treatment with methanesulfonyl chloride in the presence of a base such as triethylamine followed by addition of a protected hydroxylamine (NH₂OP).

The alcohol (V) is obtained from the appropriate ethanolamine compound (III) and the appropriate ethane compound of the formula (IV).

The compounds of formula (III) and (IV) are either known compounds, which can be made by the known methods, or they are analogs of known compounds, which can be prepared by methods analogous to the known methods.

The intermediate compounds of formula (II) wherein Ar is substituted phenyl can be prepared according to the procedures shown in the following Scheme 2.

Scheme 2



(Q is, for example, halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkoxy-C₁-C₄ alkyloxy or CF₃, n=1-5, preferably 1-3)

In the above Scheme 2, a compound (VII) can be reacted with a substituted-styrene oxide (VIII) to form a mixture of a compounds (IX) and (X). This reaction may be carried out in the absence or presence of a reaction inert solvent (e.g., methanol (MeOH), ethanol (EtOH), isopropylalcohol, tetrahydrofuran (THF), dioxane, dimethylformamide (DMF), dimethylsulfoxide (DMSO), methylene chloride (CH₂Cl₂), water, benzene, toluene, n-hexane, cyclohexane) at a temperature from -78 °C to reflux temperature of the solvent, preferably from 0 °C to 25 °C for 5 minutes to 48

hours preferably from 0.5 to 12 hours. A compound (II') can be prepared from the mixture of a compound (IX) and a compound (X) under the same conditions as already described in Scheme 1.

5 According to the above procedures, R, S configuration of compounds (IX) and (X) can be selectively determined. In addition, in the above procedures, 1-substitutedphenyl-1,2-ethanediol 2-tosylate can be used instead of the substituted-styrene oxide (VIII).

10 The compounds of formula (I) of this invention are basic, and therefore they will form acid-addition salts. All such salts are within the scope of this invention. However, it is necessary to use an acid addition salts which is pharmaceutically-acceptable for administration to a mammal. The acid-addition salts can be prepared by standard methods, e.g., by contacting the basic and acidic compounds in substantially equivalent proportions in water or an organic solvent such as methanol or ethanol, or a mixture thereof. The salts can be isolated by evaporation of the solvent. Typical salts which can be formed are the hydrochloride, nitrate, sulfate, 15 bisulfate, phosphate, acetate, lactate, citrate, tartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, *p*-toluenesulfonate, oxalate and pamoate (1,1'-methylene-bis-(2-hydroxy-3-naphtoate)) salts.

20 The compounds of formula (I) of this invention, wherein R is hydrogen, are acidic, and they will form base salts. All such salts are within the scope of this invention. However, it is necessary to use a base salt which is pharmaceutically-acceptable for administration to a mammal. The base salts can be prepared by standard methods, e.g., by contacting the acidic and basic compounds in substantially equivalent proportions in water or an organic solvent such as methanol or ethanol, or 25 a mixture thereof. The salts can be isolated by evaporation of the solvent. Typical base salts which can be formed are the sodium, potassium, calcium and magnesium salts, and also salts with ammonia and amines, such as ethylamine, diethylamine, triethylamine, cyclohexylamine, piperidine and morpholine salts.

30 Also included within the scope of this invention are bioprecursors (also called pro-drugs) of the κ agonist compounds of the formula (I). A bioprecursor of a kappa agonist of formula (I) is a chemical derivative thereof which is readily converted back

into the parent compound of formula (I) in biological systems. In particular, a bioprecursor of a κ agonist of formula (I) is converted back to the parent compound of formula (I) after the bioprecursor has been administered to, and absorbed by, a mammalian subject, e.g., a human subject. For example, it is possible to make a bioprecursor of a κ agonist of the invention of formula (I) in which one or both of A and OR is hydroxy groups by making an ester to the hydroxy group. When only one of A and OR is a hydroxy group, only mono-esters are possible. When both A and OR are hydroxy, mono- and di-esters (which can be the same or different) can be made. Typical esters are simple alkanoate esters, such as acetate, propionate, butyrate, etc. In addition, when A or OR is a hydroxy group, bioprecursors can be made by converting the hydroxy group to an acyloxymethyl derivative (e.g., a pivaloyloxymethyl derivative) by reaction with an acyloxymethyl halide (e.g., pivaloyloxymethyl chloride).

The κ agonists compounds of the present invention of formula (I) exhibit significant agonist activity toward opioid κ -receptor and are thus useful as analgesic, antiinflammatory, diuretic, anesthetic and neuroprotective agents, or an agent for treatment of stroke or functional bowel diseases such as abdominal pain, for the treatment of a mammalian subject, especially a human subject, for the treatment of mammals, especially humans in need of such agents.

The activity of the κ -agonists compounds of formula

9a

(I) of the present invention, is demonstrated by the opioid receptor activity. Such activity may be determined in homogenates from guinea pig whole brain, as described by Regina, A. et al. in J. Receptor Res. 12: 171-180, 1992. In summary, tissue homogenate is incubated at 25°C for 30 min in the presence of labelled ligand and test compounds. The μ -sites are labelled by 1 nM of (3H)-[D-Ala²,MePhe⁴,Gly-^o15]-enkephalin (DAMGO), the δ -sites by 1 nM of (3H)-[D-Pen²,5]-enkephalin (DPDPE) and the κ -sites by 0.5 nM (3H)-CI-977.

10 The non specific binding is measured by use of 1 μ M CI-977 (κ), 1 μ M (DAMGO) (μ), 1 μ M (DPDPE) (δ). Data are expressed as the IC₅₀ values obtained by a non-linear fitting program using the Cheng and Prusoff equation. Some compounds prepared in the Examples showed a low IC₅₀ value in the range of 0.01 to 100 nM.

The activity of the κ agonists compounds can also be demonstrated by the Formalin Test as described by Wheeler-Aceto, H. et al. in Psychopharmacology 104: 35-44, 1991. In this testing, male SD rats (80-100 g) are injected s.c. with a test compound dissolved in 0.1% methyl cellulose saline or vehicle. After 30 min., 50 ml of a 2% formalin are injected into a hind paw. The number of licking the injected paw per observation period is measured 15-30 min. after the injection of formalin and expressed as % inhibition compared to the respective vehicle group.

The activity of the κ agonists can also be demonstrated by the Rotarod Test as described by Hayes, A.G. et al. in Br. J. Pharmacol. 79: 731-736, 1983. In this testing, a group of 6-10 male SD rats (100-120 g) are selected for their ability to balance on a rotating rod (diameter 9 cm, rate of rotation 5 r.p.m.). The selected rats are then injected s.c. with a test compound dissolved in 0.1% methyl cellulose saline. The animals are tested again 30 min. after treatment; a rat falling off the bar more than twice within 150 seconds is considered to be showing motor impairment and the animal's performance (i.e., time on the rotarod) are recorded. The ED_{50} value, defined as the dose of the drug which halves the performance time is observed in the control group.

The κ agonists compounds of formula (I) of this invention can be administered via either the oral, parenteral or topical routes to mammals. In general, these compounds are most desirably administered to humans in doses ranging from 0.01 mg to 50 mg per day, although variations will necessarily occur depending upon the weight and condition of the subject being treated, the disease state being treated and the particular route of administration chosen. However, a dosage level that is in the range of from 0.01 mg to 1 mg per kg of body weight per day, single or divided dosage is most desirably employed in humans for the treatment of pain in a postoperative patient.

The compounds of the present invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the above routes previously indicated, and such administration can be carried out in single or multiple doses. More particularly, the novel therapeutic agents of the invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets,

capsules, lozenges, trochees, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various nontoxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging 5% to 70% by weight, preferably 10% to 50% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dipotassium phosphate and glycine may be employed along with various disintegrants such as starch and preferably corn, potato or tapioca starch, alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatine capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration, solutions of a compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH > 8) if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intra-articular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art. Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions

68277-1

12

of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

Examples and Preparations

The present invention is illustrated by the following examples and preparations.

5 However, it should be understood that the invention is not limited to the specific details of these examples and preparations. Melting points were taken with a Buchi micro melting point apparatus and uncorrected. Infrared Ray absorption spectra (IR) were measured by a Shimadzu infrared spectrometer (IR-470). ¹H and ¹³C nuclear magnetic resonance spectra (NMR) were measured in CDCl₃ by a JEOL* NMR
10 spectrometer (JNM-GX270, 270MHz) unless otherwise indicated and peak positions are expressed in parts per million (ppm) downfield from tetramethylsilane. The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad.

Preparation 1.

(S)-N-Benzylloxy-1-phenyl-2-pyrrolidinoethylamine

15 To a stirred solution of (R)-2-phenyl-2-pyrrolidinoethanol (E. Brown et al, Tetrahedron: Asymmetry, 1991, 2, 339; 4.78g, 25mmol) and triethylamine (3.95ml, 28mmol) in CH₂Cl₂ (50ml) was added methanesulfonyl chloride (2ml, 26mmol) dropwise at 0 °C (ice bath). After 3h stirring at 0 °C to room temperature (rt), the reaction mixture was washed with saturated NaHCO₃ aqueous solution, dried (Na₂SO₄),
20 and concentrated to give 5.88g of yellow solid and brown viscous oil mixture. To this mixture was added O-benzylhydroxylamine (this was prepared from O-benzylhydroxylamine hydrochloride 5.99g (37.5mmol) by basification) and ethanol (6ml) and the mixture was stirred at 80 °C for 1h. The solvent was evaporated to give 9.47g of white solid which was collected by filtration and washed with ethanol/ether
25 to afford 6.96g (83.7%) of hydrochloride salt of desired product as white crystalline, mp 161-162 °C.

¹H NMR (270MHz, CDCl₃) δ 7.44-7.25 (10H, m), 6.40 (1H, br.s), 4.68 (1H, d, J = 11.7Hz), 4.68-4.62 (1H, m), 4.63 (1H, d, J = 11.7Hz), 3.90-3.70 (1H, m), 3.60 (1H, dd, J = 7.7, 13.2Hz), 3.55-3.40 (1H, m), 3.05 (1H, dd, J = 5.5, 13.2Hz), 2.80-2.65
30 (1H, m), 2.65-2.45 (1H, m), 2.25-2.05 (2H, m), 2.05-1.80 (3H, m).

Anal. Calcd for C₁₉H₂₄N₂O·HCl : C, 68.56 ; H, 7.57; N, 8.42 ; Cl, 10.65.

* Trade-mark

Found : C, 68.36 ; H, 7.70 ; N, 8.39 ; Cl, 11.13.

This hydrochloride salt (80mg) was basified with ammonium hydroxide solution, extracted with CH_2Cl_2 , dried (Na_2SO_4), and concentrated to give 67mg of free amine derivative as a colorless oil.

¹H NMR (270MHz, CDCl_3) δ 7.46 - 7.12 (10H, m), 6.53 (1H, br.s), 4.53 (1H, d, J = 11.0Hz), 4.45 (1H, d, J = 11.4Hz), 4.20 (1H, dd, J = 3.7, 11.4Hz), 2.90 (1H, dd, J = 11.4, 12.5Hz), 2.70 - 2.60 (2H, m), 2.50 - 2.35 (2H, m), 2.28 (1H, dd, J = 4.0, 12.5Hz), 1.80 - 1.70 (4H, m).

IR(neat) : 3250 cm^{-1} .

$[\alpha]_D = +44.6$ (c = 0.67, MeOH).

Example 1

N-Benzoyloxy-2-(3,4-dichlorophenyl)-N-[1-(S)-phenyl-2-(1-pyrrolidiny)ethyl]acetamide

To a stirred solution of (S)-1-(2-O-benzylhydroxylamino-2-phenylethyl)pyrrolidine hydrochloride (2.88g, 8.65mmol) and 3,4-dichlorophenylacetic acid (2.05g, 10mmol) in CH_2Cl_2 (30ml) was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (2.30g, 12mmol) at room temperature. After 1hr stirring, the reaction mixture was washed with water and saturated NaHCO_3 aqueous solution, dried (Na_2SO_4), and concentrated to give 4.44g of pale brown viscous oil. To this oil was added methanol (2ml) and stood for 1hr. The white crystalline appeared was collected by filtration to give 1.60g of white powder. The filtrate was concentrated to afford 2.84g of oil and solid mixture, which was purified by column chromatography(silica gel; 100g, $\text{CH}_2\text{Cl}_2/\text{MeOH}$: 40/1) to give 0.82g of clear yellow viscous oil, which was crystallized by addition of methanol (0.2ml).

Total yield was 2.42g(57.9%). mp 88.5-90 °C .

¹H NMR (270MHz, CDCl_3) δ 7.46 - 7.21 (12H, m), 6.98 (1H, dd, J = 2.2, 8.4Hz), 5.80 - 5.65 (1H, m), 4.73 (1H, d, J = 10.3Hz), 4.43 (1H, d, J = 10.6Hz), 3.77 (1H, d, J = 15.8Hz), 3.61 - 3.51 (2H, m, including 1H, d, J = 15.4Hz at 3.54ppm), 2.75 - 2.60 (3H, m), 2.55 - 2.40 (2H, m), 1.80 - 1.50 (4H, m).

IR(Nujol) : 1670cm⁻¹.

Example 2

2-(3,4-Dichlorophenyl)-N-hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]acetamide

AsuspensionmixtureofN-benzyloxy-2-(3,4-dichlorophenyl)-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]acetamide (1.60g, 3.3mmol), 10% palladium on carbon (0.16g), and HCl gas saturated methanol (20ml) in methanol (20ml) was stirred under hydrogen atmosphere at room temperature for 13h. After removal of the catalyst by Celite filtration, the filtrate was concentrated to give 1.63g of violet colored viscous oil, which was basified with NH₄OH and extracted with CH₂Cl₂ (20ml x 3). The extract combined was dried (Na₂SO₄) and concentrated to afford a brown colored crystalline, which was collected by filtration and washed with ether/hexane to give 1.04g (80%) of pale yellow powder. mp 118 - 120 °C .

¹H NMR (270MHz, CDCl₃) δ 7.44 (1H, d, J = 1.8Hz), 7.37 - 7.24 (6H, m, including 1H, d, J = 8.4Hz at 7.36ppm), 7.17 (1H, dd, J = 1.8, 8.4Hz), 5.56 (1H, dd, J = 5.9, 10.3Hz), 3.90 (1H, d, J = 14.3Hz), 3.70 (1H, d, J = 13.9Hz), 3.31 (1H, dd, J = 10.6, 12.5Hz), 2.73 (1H, dd, J = 5.9, 12.5Hz), 2.60 - 2.45 (4H, m), 1.80 - 1.55 (4H, m).

IR(CH₂Cl₂) : 3450, 1650cm⁻¹.

MS m/z: 394 (M⁺+2, 0.48), 392(M⁺, 1.1), 211(4.8), 173(3.1), 149(12.9), 132(12.8), 99(28.8), 84(100).

925mg of this crystalline was dissolved in CH₂Cl₂ (10ml). To this solution was added HCl gas saturated ether (10ml) at room temperature. The mixture solution was concentrated to give a white crystalline, which was collected by filtration and washed with ether to afford 971mg of HCl salt as white powder.

mp 161-162 °C .

[α]_D = +119.8(c = 0.884, MeOH).

Anal. Calcd for C₂₀H₂₂Cl₂N₂O₂·HCl·0.5H₂O : C, 54.75 ; 5.51; N, 6.38

Found : C, 54.96 ; H, 5.49 ; N, 6.44.

Example 3

2-(3,4-Dichlorophenyl)-N-methoxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]acetamide

A mixture of 2-(3,4-Dichlorophenyl)-N-hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]acetamide (598mg, 1.5mmol), tetrabutylammonium hydrogen sulfate(10mg), NaOH 50% aqueous solution (1ml), and iodomethane (0.12ml, 2mmol) in toluene (4ml) was stirred at room temperature for 3h. The mixture was extracted with ethyl acetate (20ml x 2). The extract combined was washed with brine, dried (Na₂SO₄) and concentrated to afford 1.06g of brown viscous oil, which was purified by column chromatography (silica gel 60g, CH₂Cl₂/MeOH: 20/1) to give 304mg (49.8%) of yellow viscous oil.

¹H NMR (270MHz, CDCl₃) δ 7.41 - 7.26 (7H, m), 7.09 (1H, dd, J = 1.8, 8.1Hz), 5.70 - 5.60 (1H, m), 3.83 (1H, d, J = 15.4Hz), 3.65 (1H, d, J = 15.4Hz), 3.50 (3H, s), 3.50 (1H, dd, J = 9.9, 12.5Hz), 2.75 - 2.57 (3H, m, including 1H, dd, J = 4.8, 12.5Hz at 2.60ppm), 2.55 - 2.40 (2H, m), 1.70 (4H, m).

IR(neat) : 1670cm⁻¹.

304mg of this crystalline was dissolved in MeOH (5ml). To this solution was added HCl gas saturated ether (5ml) at room temperature. The mixture solution was concentrated to give a white crystalline, which was collected by filtration and washed with ether to afford 277mg of HCl salt as white powder.

mp 165-166 °C Anal. Calcd for C₂₁H₂₄Cl₂N₂O₂·HCl·0.5H₂O : C, 55.70 ; 5.79; N, 6.19

Found : C, 55.53 ; H, 5.80 ; N, 6.19.

Example 4N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-(2,3,6-trichlorophenyl)acetamide

This was prepared from (S)-1-(2-O-benzylhydroxylamino-2-phenylethyl)pyrrolidine in 68% yield according to a procedure similar to that described in Examples 2 and 3.

mp 217-218.5 °C (HCl salt)

¹H NMR (270MHz, free amine, CDCl₃) δ 7.44 - 7.20 (8H, m), 5.61 (1H, dd, J = 5.9, 10.6Hz), 4.36 (1H, d, J = 16.9Hz), 4.26 (1H, d, J = 17.2Hz), 3.40 (1H, dd,

16

$J = 10.6, 12.5\text{Hz}$), 2.80 (1H, dd, $J = 5.9, 12.5\text{Hz}$), 2.76 - 2.55 (4H, m), 1.90 - 1.70 (4H, m).

IR(neat, free amine) : 1650cm^{-1} .

Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{Cl}_3\text{N}_2\text{O}_2 \cdot \text{HCl} \cdot 0.5\text{H}_2\text{O}$: C, 50.76 ; 4.90; N, 5.92

5 Found : C, 50.58 ; H, 4.65 ; N, 5.83.

Example 5

N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-(4-trifluoromethylphenyl)acetamide

10 This was prepared from (S)-1-(2-O-benzylhydroxylamino-2-phenylethyl)pyrrolidine in 66.6% yield according to a procedure similar to that described in Examples 2 and 3.

mp $172.8\text{-}177^\circ\text{C}$ (HCl salt)

15 ^1H NMR (270MHz, free amine, CDCl_3) δ 7.55 (2H, d, $J = 8.4\text{Hz}$), 7.45 (2H, d, $J = 8.1\text{Hz}$), 7.40 - 7.20 (6H, m), 5.57 (1H, dd, $J = 5.9, 10.3\text{Hz}$), 4.00 (1H, d, $J = 13.9\text{Hz}$), 3.81 (1H, d, $J = 13.9\text{Hz}$), 3.30 (1H, dd, $J = 10.6, 12.5\text{Hz}$), 2.71 (1H, dd, $J = 5.9, 12.5\text{Hz}$), 2.60 - 2.40 (4H, m), 1.80 - 1.50 (4H, m).

IR(neat, free amine) : $3150, 1650\text{cm}^{-1}$.

Anal. Calcd for $\text{C}_{21}\text{H}_{23}\text{F}_3\text{N}_2\text{O}_2 \cdot \text{HCl} \cdot \text{H}_2\text{O}$: C, 56.44 ; 5.86; N, 6.27

Found : C, 56.16 ; H, 5.77 ; N, 6.76.

20

Example 6

N-Hydroxy-2-(1-naphthyl)-N-[1-(S)-Phenyl-2-(1-pyrrolidinyl)ethyl]acetamide

This was prepared from (S)-1-(2-O-benzylhydroxylamino-2-phenylethyl)pyrrolidine in 65.1% yield according to a procedure similar to that described in Examples 2 and 3.

25 mp $81.0\text{-}83.5^\circ\text{C}$ (HCl salt)

^1H NMR (270MHz, free amine, CDCl_3) δ 7.55 - 7.20 (13H, m), 5.59 (1H, dd, $J = 5.9, 10.3\text{Hz}$), 4.43 (1H, d, $J = 14.7\text{Hz}$), 4.10 (1H, d, $J = 15.0\text{Hz}$), 3.31 (1H, dd, $J = 11.0, 12.1\text{Hz}$), 2.65 (1H, dd, $J = 5.9, 12.5\text{Hz}$), 2.55 - 2.35 (4H, m), 1.60 - 1.35 (4H, m).

IR(neat, free amine) : 3150, 1650cm⁻¹.

Anal. Calcd for C₂₄H₂₆N₂O₂·HCl·1.2H₂O : C, 66.64 ; 6.85; N, 6.48

Found : C, 66.93 ; H, 6.50 ; N, 6.02.

Example 7

5 N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-(2,4,6-trimethylphenyl)acetamide

This was prepared from (S)-1-(2-O-benzylhydroxylamino-2-phenylethyl)pyrrolidine in 58.9% yield according to a procedure similar to that described in Examples 2 and 3.

10 mp 186 - 187.2 °C (HCl salt).

¹H NMR (270MHz, free amine, CDCl₃) δ 7.42 - 7.24 (6H, m), 6.82 (2H, s), 5.70 - 5.55 (1H, m), 3.86 (2H, br.s), 3.38 (1H, dd, J = 10.6, 12.1Hz), 2.74 (1H, dd, J = 5.9, 12.5Hz), 2.70 - 2.55 (4H, m), 2.22 (9H, s), 1.85 - 1.75 (4H, m).

IR(neat, free amine) : 3220, 1640cm⁻¹.

15 Anal. Calcd for C₂₃H₃₀N₂O₂·HCl·1.3H₂O : C, 64.79 ; 7.94; N, 6.57

Found : C, 64.51 ; H, 7.48 ; N, 6.31.

Example 8

N-Hydroxy-2-(4-pyridyl)-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]acetamide

20 This was prepared from (S)-1-(2-O-benzylhydroxylamino-2-phenylethyl)pyrrolidine in 67.9% yield according to a procedure similar to that described in Examples 2 and 3.

¹H NMR (270MHz, free amine, CDCl₃) δ 8.46 (2H, d, J = 5.9Hz), 7.40 - 7.18 (8H, m), 5.61 (1H, dd, J = 5.5, 10.6Hz), 3.91 (1H, d, J = 14.3Hz), 3.77 (1H, d, J = 13.9Hz), 3.33 (1H, dd, J = 11.0, 12.1Hz), 2.68 (1H, dd, J = 5.5, 12.5Hz), 2.57 - 2.40 (4H, m), 1.80 - 1.55 (4H, m).

25 IR(neat, free amine) : 1640cm⁻¹.

Example 9

2-(Benzo[b]furan-4-yl)-N-hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]acetamide

This was prepared from (*S*)-1-(2-*O*-benzylhydroxylamino-2-phenylethyl)pyrrolidine in 73.5% yield according to a procedure similar to that described in Examples 2 and 3.

¹H NMR (270MHz, CDCl₃) δ 7.59 (1H, d, J = 1.8Hz), 7.45 - 7.20 (9H, m), 6.98 (1H, br.s), 5.58 (1H, dd, J = 5.9, 10.6Hz), 4.24 (1H, d, J = 13.6Hz), 3.91 (1H, d, J = 13.6Hz), 3.28 (1H, dd, J = 11.3, 11.7Hz), 2.60 (1H, dd, J = 5.9, 12.5Hz), 2.45 - 2.30 (4H, m), 1.60 - 1.30 (4H, m).

IR(neat, free amine) : 1650cm⁻¹.

Preparation 2

1,4-Diiodo-2-(*S*)-(tetrahydropyranyloxy)butane

To a stirred solution of (*S*)-(-)-1,2,4-butanetriol (10.61g, 0.1mol) in pyridine (100ml) was added p-toluenesulfonyl chloride (38.13g, 0.2mol) by portions at 0 °C. After 2h stirring, the reaction mixture was poured into 10% HCl aqueous solution including ice and acidified to pH2. The mixture was extracted with ethyl acetate (150ml x 3). The extract combined was washed with brine, dried (Na₂SO₄), and concentrated to give 42.88g of colorless oil. A mixture of this crude ditosylate (42.88g, 0.1mol) and NaI (44.97g, 0.3mol) in acetone (300ml) was refluxed with stirring for 5h. The solid precipitated was removed by filtration and the filtrate was concentrated. The residue was dissolved in ethyl acetate and washed with Na₂S₂O₃ aqueous solution and brine. After dry (Na₂SO₄), the solvent was evaporated and the residue was purified by column chromatography (silica gel 250g, hexane/ethyl acetate: 10/1) to afford 24.81g of colorless oil. A mixture of this oil (24.81g, 76.1mmol), 3,4-dihydro-2H-pyran (21.9ml, 0.24mol), and pyridinium p-toluenesulfonate (125mg) in CH₂Cl₂ (100ml) was stirred at room temperature for 12h. The reaction mixture was diluted with CH₂Cl₂ (100ml), washed with NaHCO₃ aqueous solution, and dried (Na₂SO₄). Evaporation of the solvent gave 33.56g of pale yellow oil, which was purified by column chromatography (silica gel 250g, hexane/ethyl acetate: 20/1) to afford 28.75g (70.1% for 3 steps) of colorless oil.

¹H NMR (270MHz, CDCl₃) δ 4.80 - 4.75 (1H, m), 4.02 - 3.85 (1H, m), 3.70 - 3.17 (6H, m), 2.27 - 2.01 (2H, m), 1.90 - 1.55 (6H, m).

Preparation 32-(R)-Phenyl-2-(3-(S)-tetrahydropyranyloxy)pyrrolidin-1-yl)ethanol

A suspension mixture of 1,4-diiodo-2-(S)-(tetrahydropyranyloxy)-butane (12.50g, 30mmol), *R*-(-)-phenylglycinol (3.43g, 25mmol), and K₂CO₃ (6.91g, 50mmol) in ethanol (50ml) was refluxed with stirring for 6h. The white solid was removed by filtration and the filtrate was concentrated. The residue was diluted with NaHCO₃ aqueous solution (30ml) and extracted with CH₂Cl₂ (20ml x 3). After dry (Na₂SO₄), the solvent was evaporated to give 9.54g of clear yellow oil, which was purified by column chromatography (silica gel 150g, CH₂Cl₂/MeOH: 20/1) to afford 7.22g (99%) of colorless viscous oil.

¹H NMR (270MHz, CDCl₃) δ 7.37 - 7.27 (5H, m), 4.61 - 4.51 (1H, m), 4.40 - 4.28 (1H, m), 3.91 - 3.75 (3H, m), 3.55 - 3.42 (2H, m), 2.92 - 2.72 (1H, m), 2.70 - 2.57 (2H, m), 2.55 - 2.25 (2H, m), 2.20 - 1.95 (1H, m), 1.93 - 1.60 (3H, m), 1.60 - 1.45 (4H, m).

IR(neat) : 3450cm⁻¹.

Preparation 41-(S)-Phenyl-N-tetrahydropyranyloxy-2-(3-(S)-tetrahydropyranyloxy)pyrrolidin-1-yl)ethylamine

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydro-pyranyloxy)pyrrolidin-1-yl)ethanol and *O*-tetrahydropyranyl-hydroxylamine (R.N.Warrener and E.N.Cain, Angew. Chem. Int. Edit. 1966, 5, 511) in 42.5% yield as a brown oil according to a procedure similar to that described in Preparation 1.

¹H NMR (270MHz, CDCl₃) δ 7.45 - 7.25 (5H, m), 6.51 (1H, br.s), 4.80 - 4.73 (1H, m), 4.65 - 4.55 (1H, m), 4.45 - 4.33 (1H, m), 4.28 - 4.15 (1H, m), 4.00 - 3.75 (2H, m), 3.70 - 2.55 (9H, m), 2.30 - 2.05 (1H, m), 1.90 - 1.35 (12H, m).

Example 102-(3,4-Dichlorophenyl)-N-tetrahydropyranyloxy-N-[2-(3-(S)-tetrahydropyranyloxy)pyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 1-(*S*)-Phenyl-N-tetrahydropyranyloxy-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethylamine and 3,4-dichlorophenylacetic acid in 69.8% yield as a clear brown viscous oil according to a procedure similar to that described in Example 1.

5 ¹H NMR (270MHz, CDCl₃) δ 7.43 - 7.15 (7.4H, m), 6.98 - 6.91 (0.6H, m), 5.69 (0.4H, dd, J=4.0, 11.0Hz), 5.58 (0.6H, dd, J=4.8, 11.4Hz), 5.35 - 5.20 (1H, m), 4.65 - 4.53 (1H, m), 4.41 - 4.21 (1H, m), 4.15 - 3.80 (4H, m), 3.68 - 3.10 (4H, m), 3.03 - 2.80 (2H, m), 2.70 - 2.35 (3H, m), 2.20 - 1.10 (13H, m).
IR(neat) : 1660cm⁻¹.

10

Example 112-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(*S*)-hydroxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]acetamide

A mixture of 2-(3,4-Dichlorophenyl)-N-tetrahydropyranyloxy-N-[2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]acetamide (1.13g, 1.96mmol)
15 and HCl gas saturated MeOH (4ml) in MeOH (20ml) was stirred at room temperature for 7h. The solvent was evaporated. The residue was basified with saturated NaHCO₃ aqueous solution, extracted with CH₂Cl₂, and dried (Na₂SO₄). Evaporation of the solvent gave 0.80g of brown viscous oil, which was crystallized by adding ether and scratching. The crystalline was collected by filtration and washed with ether to afford
20 377mg(47.1%) of white powder.
mp 98.5 - 99.5°C.

¹H NMR (270MHz, CDCl₃) δ 7.45 - 7.20 (7H, m), 7.14 (1H, dd, J=1.8, 9.9Hz), 5.62 (1H, dd, J=5.5, 11.0Hz), 5.00 - 3.00 (2H, almost flat br.s), 4.35 - 4.25 (1H, m), 3.85 (1H, d, J=14.3 Hz), 3.73 (1H, d, J=13.9Hz), 3.38 (1H, dd, J=11.0, 12.5
25 Hz), 2.95 (1H, dt, J=5.1, 8.8 Hz), 2.73 (1H, d, J=10.6 Hz), 2.65 (1H, dd, J=5.5, 12.5Hz), 2.51 (1H, dd, J=5.5, 10.6Hz), 2.40 - 2.27 (1H, m), 2.22 - 2.07 (1H, m), 1.65 - 1.50 (1H, m).

IR(Nujol) : 3070, 1640cm⁻¹.

MS m/z: 412(M⁺+4, 10.3), 410(M⁺+2, 85.7), 408(M⁺, 100), 304(8.6), 149(50.2),
30 114(22.7), 112(24.2).

21

$[\alpha]_D = +102.9 (c = 0.516, \text{MeOH})$.

HCl salt: mp 65.5-67.0 °C.

Anal. Calcd for $\text{C}_{20}\text{H}_{22}\text{Cl}_2\text{N}_2\text{O}_3 \cdot \text{HCl} \cdot 0.5\text{H}_2\text{O}$: C, 52.82 ; H, 5.32; N, 6.16 .

Found : C, 53.09 ; H, 5.29 ; N, 6.17 .

5

Preparation 5

(R)-(-)-2-(4-Fluorophenyl)glycinol

This was prepared from 4-fluoro-D-a-phenylglycine in 88% yield according to the procedure of D.A.Evans (Organic Synthesis, 68, 77).

^1H NMR (270MHz, CDCl_3) δ 7.30 (2H, dd, $J=5.5, 8.4\text{Hz}$), 7.03 (2H, t, $J=8.4\text{Hz}$), 4.05 (1H, dd, $J=4.4, 8.1\text{Hz}$), 3.71 (1H, dd, $J=4.4, 10.6\text{Hz}$), 3.53 (1H, dd, $J=8.4, 10.6\text{Hz}$), 2.19 (3H, br.s).

IR(KBr) : 3350, 3280 cm^{-1} .

Preparation 6

2-(R)-(4-Fluorophenyl)-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol

This was prepared from (R)-(-)-2-(4-fluorophenyl)glycinol in 68.8% yield according to a procedure similar to that described in Preparation 3.

^1H NMR (270MHz, CDCl_3) δ 7.31-7.26 (2H, m), 7.03 (2H, dd, $J=8.4, 8.8\text{Hz}$), 4.65-4.51 (1H, m), 4.40-4.27 (1H, m), 3.90-3.75 (3H, m), 3.55-3.40 (2H, m), 2.90-2.70 (1H, m), 2.70-2.50 (2H, m), 2.50-2.35 (1H, m), 2.30-1.95 (2H, m), 1.95-1.60 (3H, m), 1.60-1.45 (4H, m).

IR(neat) : 3450 cm^{-1} .

Example 12

2-(3,4-Dichlorophenyl)-N-[1-(S)-(4-fluorophenyl)ethyl-2-(3-(S)-hydroxypyrrolidin-1-yl)]-N-hydroxyacetamide

25

This was prepared from 2-(R)-(4-fluorophenyl)-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 3,4-dichlorophenylacetic acid in 52.8% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

^1H NMR (270MHz, CDCl_3) δ 7.41 - 7.26 (4H, m), 7.12 (1H, dd, $J=1.8, 8.1\text{Hz}$),

6.99 (2H, dd, $J=8.4, 8.8\text{Hz}$), 5.60 (1H, dd, $J=5.1, 11.0\text{Hz}$), 4.35 - 4.25 (1H, m),
 3.82 (1H, d, $J=13.9\text{Hz}$), 3.72 (1H, d, $J=14.3\text{Hz}$), 3.71 (1H, s), 3.58 (1H, s), 3.35
 (1H, dd, $J=11.7, 12.1\text{Hz}$), 3.00 - 2.90 (1H, m), 2.73 (1H, br.d, $J=11.0\text{Hz}$), 2.58
 (1H, dd, $J=5.1, 12.5\text{Hz}$), 2.51 (1H, dd, $J=5.5, 10.6\text{Hz}$), 2.37-2.10 (2H, m), 1.65-
 1.55 (1H, m).

IR(neat) : 3200, 1640 cm^{-1} .

MS m/z : 426(M^+).

HCl salt: amorphous solid.

Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{Cl}_2\text{FN}_2\text{O}_3 \cdot \text{HCl} \cdot 0.7\text{H}_2\text{O}$: C, 50.43 ; H, 4.95; N, 5.88 .

Found : C, 50.80 ; H, 4.96 ; N, 5.45 .

Example 13

2-(4-Bromophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 4-bromophenylacetic acid in 44.6% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

^1H NMR (270MHz, CDCl_3) δ 7.50- 7.14 (9H, m), 5.61 (1H, dd, $J=5.1, 11.0\text{Hz}$),
 4.28-4.22 (1H, m), 3.90 (1H, d, $J=13.6\text{Hz}$), 3.70 (1H, d, $J=13.9\text{Hz}$), 3.33 (1H, dd,
 $J=11.0, 12.5\text{Hz}$), 2.92-2.82 (1H, m), 2.72-2.64 (2H, m), 2.50 (1H, dd, $J=5.5,$
 10.6Hz), 2.38-2.28 (1H, m), 2.20 (2H, br.s), 2.16-2.01 (1H, m), 1.60-1.50 (1H, m).

IR(neat) : 3200, 1630 cm^{-1} .

MS m/z : 418(M^+).

HCl salt: amorphous solid.

Anal. Calcd for $\text{C}_{20}\text{H}_{23}\text{BrN}_2\text{O}_3 \cdot \text{HCl} \cdot 0.5\text{H}_2\text{O}$: C, 51.68 ; H, 5.42; N, 6.03 .

Found : C, 51.75 ; H, 5.51 ; N, 5.71 .

Example 14

2-(3-Bromophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-

yl)ethanol and 3-bromophenylacetic acid in 29.8% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.51-7.15 (9H, m), 5.62 (1H, dd, J=5.5, 11.0Hz), 4.28-4.20 (1H, m), 3.94 (1H, d, J=13.9Hz), 3.70 (1H, d, J=13.6Hz), 3.35 (1H, dd, J=11.4, 12.5Hz), 2.92-2.83 (1H, m), 2.70-2.62 (2H, m), 2.51 (1H, dd, J=5.1, 10.6Hz), 2.42 (2H, br.s), 2.38-2.28 (1H, m), 2.18-2.03 (1H, m), 1.60-1.46 (1H, m). IR(neat) : 3200, 1630cm⁻¹.

MS m/z: 418(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₀H₂₃BrN₂O₃·HCl·H₂O : C, 50.70 ; H, 5.53; N, 5.91 .

Found : C, 50.57 ; H, 5.58 ; N, 5.90 .

Example 15

2-(4-Fluorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(R)-phenyl-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 4-fluorophenylacetic acid in 23.6% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.40-7.22 (7H, m), 7.10-6.95 (2H, m), 5.67-5.61 (1H, m), 4.34-4.22 (1H, m), 3.92 (1H, d, J=13.6Hz), 3.73 (1H, d, J=13.9Hz), 3.36 (1H, dd, J=10.6, 12.5Hz), 2.96-2.86 (1H, m), 2.76-2.62 (2H, m), 2.58-2.48 (1H, m), 2.40-2.28 (1H, m), 2.24-1.70 (3H, m), 1.64-1.48 (1H, m).

IR(neat) : 3400, 1630cm⁻¹.

MS m/z: 358(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₀H₂₃FN₂O₃·HCl·0.4H₂O : C, 59.74 ; H, 6.22; N, 6.97 .

Found : C, 59.81 ; H, 6.43 ; N, 6.88 .

Example 16

2-(3,4-Dimethoxyphenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 3,4-dimethoxyphenylacetic acid in 10.6% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.40-7.22 (5H, m), 6.95-6.78 (3H, m), 5.70-5.60 (1H, m), 4.25-4.15 (1H, m), 3.91 (1H, d, J=13.9Hz), 3.88 (3H, s), 3.87 (3H, s), 3.68 (1H, d, J=13.9Hz), 3.33 (1H, dd, J=11.4, 11.7Hz), 2.90-2.78 (1H, m), 2.74-2.60 (2H, m), 2.47 (1H, dd, J=5.1, 10.6Hz), 2.34-2.20 (1H, m), 2.14-1.98 (1H, m), 1.90 (2H, br.s), 1.50-1.36 (1H, m).

IR(neat) : 3400, 1640cm⁻¹.

MS m/z: 400(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₂H₂₈N₂O₅·HCl·2.7H₂O : C, 54.42 ; H, 7.14; N, 5.77 .

Found : C, 54.31 ; H, 6.77 ; N, 5.92 .

Example 17

N-Hydroxy-N-[2-(3-(*S*)-hydroxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]-2-(3-trifluoromethylphenyl)acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 3-trifluoromethylphenylacetic acid in 18.9% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.60-7.26 (9H, m), 5.75-5.65 (1H, m), 4.35-4.25 (1H, m), 3.99 (1H, d, J=14.3Hz), 3.86 (1H, d, J=14.3Hz), 3.54-3.38 (1H, m), 3.04-2.94 (1H, m), 2.84-2.40 (6H, m), 2.20-2.06 (1H, m), 1.70-1.55 (1H, m).

IR(neat) : 3350, 1630cm⁻¹.

MS m/z: 408(M+H)⁺.

HCl salt: amorphous solid.

Anal. Calcd for C₂₁H₂₃F₃N₂O₃·HCl·1.9H₂O : C, 54.70 ; H, 5.64; N, 6.08 . Found : C, 54.83 ; H, 5.97 ; N, 6.21 .

Example 18

N-Hydroxy-N-[2-(3-(*S*)-hydroxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]-2-(4-

trifluoromethylphenyl)acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 4-trifluoromethylphenylacetic acid in 35.4% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

5 ¹H NMR (270MHz, CDCl₃) δ 7.56 (2H, d, J=8.1Hz), 7.44 (2H, d, J=8.1Hz), 7.33-7.26 (5H, m), 5.65 (1H, dd, J=5.9, 11.0Hz), 4.35 - 4.20 (1H, m), 3.99 (1H, d, J=14.3Hz), 3.85 (1H, d, J=13.9Hz), 3.41 (1H, dd, J=12.1, 12.5Hz), 3.00-2.90 (1H, m), 2.82-2.02 (7H, m), 1.64-1.50 (1H, m).

IR(neat) : 3100, 1650cm⁻¹.

10 MS m/z: 408(M⁺).

HCl salt: mp 142.5-144.2 °C

Anal. Calcd for C₂₁H₂₃F₃N₂O₃·HCl·0.2H₂O : C, 56.24 ; H, 5.48; N, 6.25 .

Found : C, 56.27 ; H, 5.61 ; N, 6.08 .

Example 19

15 2-(4-Biphenyl)-N-hydroxy-N-[2-(3-(*S*)-hydroxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 4-biphenylacetic acid in 38.8% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

20 ¹H NMR (270MHz, CDCl₃) δ 7.60-7.26 (14H, m), 5.66 (1H, dd, J=5.1, 11.0Hz), 4.20-4.14 (1H, m), 4.04 (1H, d, J=13.6Hz), 3.76 (1H, d, J=13.2Hz), 3.35 (1H, dd, J=10.3, 13.6Hz), 2.90-2.80 (1H, m), 2.73-2.63 (2H, m), 2.55-2.45 (1H, m), 2.35-2.22 (1H, m), 2.10-1.96 (1H, m), 1.90 (2H, br.s), 1.50-1.35 (1H, m).

MS m/z: 417(M+H)⁺.

25 HCl salt: mp 163.8-165.5 °C

Anal. Calcd for C₂₆H₂₈N₂O₃·HCl·0.5H₂O : C, 67.60 ; H, 6.55; N, 6.06 .

Found : C, 67.77 ; H, 6.42 ; N, 5.76 .

Example 20

N-Hydroxy-N-[2-(3-(*S*)-hydroxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]-2-(4-

nitrophenyl)acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 4-nitrophenylacetic acid in 11.6% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 8.14 (2H, d, J=8.8Hz), 7.44 (2H, d, J=8.8Hz), 7.35-7.16 (5H, m), 5.74 (1H, dd, J=4.8, 10.3Hz), 4.46-4.38 (1H, m), 4.03 (1H, d, J=15.0Hz), 3.96 (1H, d, J=15.0Hz), 3.64-3.50 (1H, m), 3.20-3.10 (1H, m), 2.96 (1H, br.d, J=10.3Hz), 2.90-2.74 (3H, m), 2.66 (2H, br.s), 2.30-2.16 (1H, m), 1.84-1.70 (1H, m).

IR(neat) : 3400, 1630cm⁻¹.

MS m/z: 385(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₀H₂₃N₃O₅·HCl·1.5H₂O : C, 53.51 ; H, 6.06; N, 9.36 .

Found : C, 53.71 ; H, 6.01 ; N, 9.11 .

15

Example 21N-Hydroxy-N-[2-(3-(*S*)-hydroxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]-2-(3-nitrophenyl)acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 3-nitrophenylacetic acid in 11.6% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 8.17-8.08 (2H, m), 7.66-7.20 (7H, m), 5.64 (1H, dd, J=5.9, 11.0Hz), 4.38-4.30 (1H, m), 4.03 (1H, d, J=14.7Hz), 3.90 (1H, d, J=14.3Hz), 3.50-3.38 (1H, m), 3.06-2.94 (1H, m), 2.84-2.70 (2H, m), 2.66-2.56 (1H, m), 2.50-2.32 (1H, m), 2.20-2.04 (1H, m), 1.96 (2H, br.s), 1.70-1.50 (1H, m).

MS m/z: 386(M+H)⁺.

HCl salt: mp 154.3-155.5 °C.

Anal. Calcd for C₂₀H₂₃N₃O₅·HCl·0.3H₂O : C, 56.22 ; H, 5.80; N, 9.83 .

Found : C, 56.29 ; H, 5.80 ; N, 9.55 .

Example 22

2-(4-Chlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(R)-phenyl-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 4-chlorophenylacetic acid in 49.4% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.40-7.20 (9H, m), 5.65 (1H, dd, J=5.1, 11.0Hz), 5.00-3.30 (2H, wide spread br.s), 4.35-4.25 (1H, m), 3.86 (1H, d, J=13.9Hz), 3.74 (1H, d, J=13.9Hz), 3.40 (1H, dd, J=11.7, 12.1Hz), 3.02-2.90 (1H, m), 2.75 (1H, br.d, J=10.6Hz), 2.61 (1H, dd, J=5.1, 12.5Hz), 2.51 (1H, dd, J=5.1, 10.3Hz), 2.40-2.25 (1H, m), 2.23-2.08 (1H, m), 1.65-1.50 (1H, m).

IR(neat) : 3400, 1630cm⁻¹.

MS m/z: 374(M⁺).

HCl salt: mp 146.5-147.3 °C.

Anal. Calcd for C₂₀H₂₃ClN₂O₃·HCl·0.3H₂O : C, 57.64 ; H, 5.95; N, 6.72 .

Found : C, 57.87 ; H, 5.88 ; N, 6.78 .

Example 23

2-(3-Chlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(R)-phenyl-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 3-chlorophenylacetic acid in 29.6% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.34-7.20 (9H, m), 5.75-5.62 (1H, m), 4.35-4.25 (1H, m), 3.94 (1H, d, J=13.9Hz), 3.74 (1H, d, J=13.9Hz), 3.45 (1H, dd, J=9.5, 12.1Hz), 3.05-2.92 (1H, m), 2.80 (1H, br.d, J=10.6Hz), 2.77-2.30 (3H, m), 3.80-2.30 (2H, almost flat peak), 2.23-2.06 (1H, m), 1.68-1.54 (1H, m).

IR(neat) : 3350, 1630cm⁻¹.

MS m/z: 374(M⁺).

HCl salt: mp 113.2-114.3 °C.

Anal. Calcd for C₂₀H₂₃ClN₂O₃·HCl·0.4H₂O : C, 57.40 ; H, 5.97; N, 6.69 .

Found : C, 57.79 ; H, 5.84 ; N, 6.74 .

Example 242-(2-Chlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 2-chlorophenylacetic acid in 31.2% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.16 (9H, m), 5.85-5.70 (1H, m), 4.44-4.34 (1H, m), 4.14 (1H, d, J=16.1Hz), 3.91 (1H, d, J=16.1Hz), 3.68-3.48 (1H, m), 3.24-3.10 (1H, m), 2.98-2.40 (6H, m), 2.34-2.18 (1H, m), 1.86-1.70 (1H, m).

IR(neat) : 3400, 1640cm⁻¹.

MS m/z: 374(M⁺).

HCl salt: mp 146 °C.

Anal. Calcd for C₂₀H₂₃ClN₂O₃·HCl·H₂O : C, 55.95 ; H, 6.10; N, 6.52 .

Found : C, 56.18 ; H, 6.00 ; N, 6.55 .

Example 25N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(2,3,5-trichlorophenyl)acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 2,3,5-trichlorophenylacetic acid in 51.6% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.26 (6H, m), 7.14 (1H, d, J=2.2Hz), 5.70 (1H, dd, J=4.8, 11.0Hz), 4.48-4.30 (1H, m), 4.20-3.00 (2H, wide spread br.s), 4.06 (1H, d, J=16.5Hz), 3.90 (1H, d, J=16.1Hz), 3.50 (1H, dd, J=11.4, 12.1Hz), 3.20-3.10 (1H, m), 2.86 (1H, br.d, J=10.3Hz), 2.75-2.60 (2H, m), 2.55-2.35 (1H, m), 2.35-2.20 (1H, m), 1.85-1.70 (1H, m).

IR(neat) : 3400, 1640cm⁻¹.

MS m/z: 444(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₀H₂₁Cl₃N₂O₃·HCl·H₂O : C, 48.21 ; H, 4.86; N, 5.62 .

Found : C, 48.56 ; H, 5.17 ; N, 5.40 .

Example 26N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(2,4,6-trichlorophenyl)acetamide

5 This was prepared from 2-(R)-phenyl-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 2,4,6-trichlorophenylacetic acid in 14.0% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.50-7.26 (7H, m), 5.60 (1H, dd, J=4.8, 11.4Hz), 4.47-4.38 (1H, m), 4.19 (2H, s), 3.49 (1H, dd, J=11.7, 12.1Hz), 3.25-3.10 (1H, m), 2.84 (1H, br.d, J=9.5Hz), 2.75-2.60 (2H, m), 2.50-2.35 (2H, m), 2.35-2.20 (2H, m), 1.90-1.70 (1H, m).

1R(KBr) : 3450, 1640cm⁻¹.

MS m/z: 442(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₀H₂₁Cl₃N₂O₃·HCl·0.2H₂O : C, 49.65 ; H, 4.67; N, 5.79 .

15 Found : C, 49.42 ; H, 4.39 ; N, 5.96 .

Example 27N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(2,4,6-trimethylphenyl)acetamide

20 This was prepared from 2-(R)-phenyl-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 2,4,6-trimethylphenylacetic acid in 67.8% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.25 (5H, m), 6.81 (2H, s), 5.80-5.65 (1H, m), 4.40-4.30 (1H, m), 3.86 (2H, s), 3.49 (1H, dd, J=11.7, 13.2Hz), 3.20-3.10 (1H, m), 2.80 (1H, br.d, J=10.3Hz), 2.65-2.50 (2H, m), 2.35-2.25 (3H, m), 2.23 (3H, s), 2.18 (6H, s), 1.90-1.65 (1H, m), 1.65-1.50 (1H, m).

25 1R(neat) : 3250, 1630cm⁻¹.

MS m/z: 382(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₃H₃₀N₂O₃·HCl·0.2H₂O : C, 64.01 ; H, 7.57; N, 6.49 .

30 Found : C, 64.08 ; H, 7.85 ; N, 6.61 .

Example 282-(2,3-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(R)-phenyl-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 2,3-dichlorophenylacetic acid in 56% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.50-7.05 (8H, m), 5.69 (1H, dd, J=5.1, 11.4Hz), 5.00 - 3.00 (2H, almost flat br.s), 4.45-4.35 (1H, m), 4.10 (1H, d, J=16.1Hz), 3.92 (1H, d, J=16.1Hz), 3.48 (1H, dd, J=11.7, 12.1Hz), 3.20-3.10 (1H, m), 2.82 (1H, d, J=10.3Hz), 2.70-2.55 (2H, m), 2.45-2.20 (2H, m), 1.80-1.70 (1H, m).

IR(neat) : 3200, 1640cm⁻¹.

MS m/z: 408(M⁺).

HCl salt: mp 155.3-158.1 °C.

Anal. Calcd for C₂₀H₂₂Cl₂N₂O₃·HCl : C, 53.89 ; H, 5.20; N, 6.28 .

Found : C, 53.72 ; H, 5.24 ; N, 6.16 .

Example 292-(2,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(R)-phenyl-2-(3-(S)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 2,4-dichlorophenylacetic acid in 71.9% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.15 (8H, m), 5.69 (1H, dd, J=5.1, 11.4Hz), 6.50-4.50 (2H, almost flat br.s), 4.35-4.25 (1H, m), 4.00 (1H, d, J=16.1Hz), 3.86 (1H, d, J=16.1Hz), 3.47 (1H, dd, J=11.7, 12.1Hz), 3.20-3.10 (1H, m), 2.83 (1H, d, J=10.6Hz), 2.61 (2H, dd, J=5.5, 12.1Hz), 2.45-2.20 (2H, m), 1.80-1.65 (1H, m).

IR(neat) : 3200, 1635cm⁻¹.

MS m/z: 408(M⁺).

HCl salt: mp 149-151.5 °C.

Anal. Calcd for C₂₀H₂₂Cl₂N₂O₃·HCl·0.2H₂O : C, 53.46 ; H, 5.25; N, 6.23 . Found :

C, 53.46 ; H, 5.19 ; N, 6.19 .

Example 302-(2,5-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(*R*)-phenyl-2-(3-(*S*)-tetrahydropyranyloxypyrrolidin-1-yl)ethanol and 2,5-dichlorophenylacetic acid in 56.3% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.15 (8H, m), 5.69 (1H, dd, J=5.1, 11.0Hz), 5.60-4.50 (2H, almost flat br.s), 4.35-4.25 (1H, m), 4.03 (1H, d, J=16.1Hz), 3.86 (1H, d, J=16.1Hz), 3.47 (1H, t, J=11.7Hz), 3.20-3.10 (1H, m), 2.82 (1H, d, J=10.6Hz), 2.63 (2H, dd, J=5.1, 12.1Hz), 2.45-2.20 (2H, m), 1.85-1.70 (1H, m).

IR(neat) : 3200, 1635cm⁻¹.

MS m/z: 408(M⁺).

HCl salt: 157.5-158.2 °C.

Anal. Calcd for C₂₀H₂₂Cl₂N₂O₃·HCl·0.2H₂O : C, 53.46 ; H, 5.25; N, 6.23 .

Found : C, 53.35 ; H, 5.21 ; N, 6.14 .

Preparation 72-(3-(S)-Methoxymethyloxypyrrolidin-1-yl)-2-(R)-phenylethanol

To a stirred solution of (*S*)-(-)-butanetriol (10.61g, 0.1mol) in pyridine (50ml) was added p-toluenesulfonyl chloride (38.13g, 0.2mol) by portions at 0 °C (ice bath). After 1h stirring, the reaction mixture was poured into c-HCl aqueous solution including ice and acidified to pH2. The mixture was extracted with ethyl acetate (100ml x 3). The extract combined was washed with brine, dried (Na₂SO₄), and concentrated to give 36.22g of pale brown clear oil. To a stirred solution of this crude ditosylate (36.22g) and methylal(50 ml) in CH₂Cl₂ (50ml) was added P₂O₅ (20g). After 1h stirring, another 10g of P₂O₅ was added to the reaction mixture. After 2h stirring, the CH₂Cl₂ layer was separated. Residual dark brown solid was washed with CH₂Cl₂. The combined CH₂Cl₂ layer was washed with NaHCO₃ aqueous solution, dried (Na₂SO₄), and concentrated to give 38.51g of brown viscous oil. A mixture of this oil (38.51g, 84mmol), (*R*)-(-)-2-phenylglycinol (10.97g, 80mmol), and triethylamine (23mmol, 160mmol) in ethanol (40ml) was refluxed with stirring for 15h. The solvent

was evaporated and the residue was dissolved in CH_2Cl_2 (200ml), washed with NaHCO_3 aqueous solution and brine, dried (Na_2SO_4), and concentrated to give 28.43g of brown viscous oil. This oil was purified by column chromatography (silica gel 200g, CH_2Cl_2 /methanol: 40/1 to 20/1) to afford 9.74g (48.4%) of clear brown viscous oil.

5 ^1H NMR (270MHz, CDCl_3) δ 7.40 - 7.25 (5H, m), 4.62 (1H, d, $J=7.0\text{Hz}$), 4.58 (1H, d, $J=6.6\text{Hz}$), 4.26 - 4.18 (1H, m), 3.92 (1H, dd, $J=6.2, 11.0\text{Hz}$), 3.82 (1H, dd, $J=5.5, 11.0\text{Hz}$), 3.54 (2H, t, $J=5.9\text{Hz}$), 3.33 (3H, s), 2.93 (1H, br.s), 2.85 - 2.66 (3H, m), 2.56 - 2.47 (1H, m), 2.16 - 2.02 (1H, m), 1.88 - 1.77 (1H, m).

Example 31

10 2-(2,6-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-phenylethanol and 2,6-dichlorophenylacetic acid in 47.2% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

15 ^1H NMR (270MHz, CDCl_3) δ 7.50-7.25 (7H, m), 7.20-7.10 (1H, m), 5.71 (1H, dd, $J=5.1, 11.4\text{Hz}$), 5.40-3.70 (2H, almost flat br.s), 4.50-4.40 (1H, m), 4.25 (2H, s), 3.50 (1H, dd, $J=11.0, 12.5\text{Hz}$), 3.28-3.15 (1H, m), 2.87 (1H, d, $J=10.3\text{Hz}$), 2.75-2.55 (2H, m), 2.50-2.25 (2H, m), 1.90-1.70 (1H, m).

IR(KBr) : 3400, 1640 cm^{-1} .

20 MS m/z: 408(M^+).

HCl salt: mp 95.5-96.8 $^{\circ}\text{C}$.

Anal. Calcd for $\text{C}_{20}\text{H}_{22}\text{Cl}_2\text{N}_2\text{O}_3 \cdot \text{HCl} \cdot 0.5\text{H}_2\text{O}$: C, 52.82 ; H, 5.32; N, 6.16 .

Found : C, 52.61 ; H, 5.13 ; N, 6.10 .

Example 32

25 2-(3,5-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-phenylethanol and 3,5-dichlorophenylacetic acid in 47.8% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

33

¹H NMR (270MHz, CDCl₃) δ 7.45-7.15 (8H, m), 5.63 (1H, dd, J=5.5, 11.0Hz), 4.50-3.00 (2H, almost flat br.s), 4.40-4.28 (1H, m), 3.87 (1H, d, J=14.3Hz), 3.71 (1H, d, J=14.3Hz), 3.39 (1H, dd, J=11.4, 12.1Hz), 3.05-2.95(1H, m), 2.74 (1H, d, J=11.0Hz), 2.65 (1H, dd, J=5.5, 12.5Hz), 2.54 (1H, dd, J=5.5, 10.6Hz), 2.45-2.30 (1H, m), 2.25-2.10 (1H, m), 1.70-1.55 (1H, m).

IR(neat) : 3350, 1650cm⁻¹.

MS m/z: 408(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₀H₂₂Cl₂N₂O₃·HCl·2H₂O : C, 49.86 ; H, 5.65; N, 5.81 .

Found : C, 49.49 ; H, 5.53 ; N, 5.59 .

Example 33

N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(2,3,6-trichlorophenyl)acetamide

This was prepared from 2-(3-(S)-methoxymethoxypyrrolidin-1-yl)-2-(R)-phenylethanol and 2,3,6-trichlorophenylacetic acid in 46.7% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.20 (7H, m), 5.69 (1H, dd, J=4.8, 11.0Hz), 5.00-3.50 (2H, almost flat br.s), 4.50-4.40 (1H, m), 4.29 (2H, s), 3.49 (1H, t, J=11.7Hz), 3.25-3.15 (1H, m), 2.85 (1H, d, J=10.3Hz), 2.70-2.60 (2H, m), 2.45-2.20 (2H, m), 1.90-1.70 (1H, m).

IR(KBr) : 3400, 1640cm⁻¹.

MS m/z: 442(M⁺).

HCl salt: mp 102-103 °C.

Anal. Calcd for C₂₀H₂₁Cl₃N₂O₃·HCl·H₂O : C, 48.21 ; H, 4.86; N, 5.62 .

Found : C, 48.40 ; H, 4.64 ; N, 5.52 .

Example 34

2-(Benzo[b]furan-4-yl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(3-(S)-methoxymethoxypyrrolidin-1-yl)-2-(R)-

phenylethanol and 4-benzo[*b*]furanacetic acid in 57.5% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.64 (1H, d, J=2.2Hz), 7.50-7.25 (7H, m), 7.14 (1H, d, J=7.3Hz), 6.84 (1H, dd, J=0.7, 2.2Hz), 5.61 (1H, dd, J=5.5, 11.4Hz), 4.24 (1H, d, J=13.6Hz), 4.05-3.95 (1H, m), 3.91 (1H, d, J=13.2Hz), 3.31 (1H, dd, J=11.7, 12.1Hz), 2.75-2.65 (1H, m), 2.63-2.50 (2H, m), 2.30 (1H, dd, J=5.1, 10.3Hz), 2.20-2.10 (1H, m), 2.00-1.85 (1H, m).

IR(neat) : 3400, 1635cm⁻¹.

MS m/z: 380(M⁺).

10 HCl salt: amorphous solid.

Anal. Calcd for C₂₂H₂₄N₂O₄·HCl·1.1H₂O : C, 60.51 ; H, 6.28; N, 6.41 .

Found : C, 60.31 ; H, 5.98 ; N, 6.47 .

Example 35

15 N-Hydroxy-N-[2-(3-(*S*)-hydroxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]-2-(1-tetralon-6-yl)acetamide

This was prepared from 2-(3-(*S*)-methoxymethyloxypyrrolidin-1-yl)-2-(*R*)-phenylethanol and (1-tetralon-6-yl)acetic acid in 59.4% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

20 ¹H NMR (270MHz, CDCl₃) δ 7.96 (1H, d, J=8.1Hz), 7.40-7.18 (7H, m), 5.66 (1H, dd, J=5.5, 11.0Hz), 4.30-4.20 (1H, m), 3.94 (1H, d, J=14.3Hz), 3.81 (1H, d, J=13.9Hz), 3.80-2.00 (2H, almost flat br.s), 3.40 (1H, dd, J=11.7, 12.1Hz), 3.00-2.85 (3H, m), 2.80-2.50 (5H, m), 2.45-2.30 (1H, m), 2.20-2.05 (3H, m), 1.65-1.50 (1H, m).

IR(neat) : 3400, 1680, 1640cm⁻¹.

25 MS m/z: 408(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₄H₂₈N₂O₄·HCl·1.2H₂O : C, 61.78 ; H, 6.78; N, 6.00 .

Found : C, 61.60 ; H, 6.59 ; N, 6.35 .

Example 36

2-(3,4-Dimethylphenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-phenylethanol and 3,4-dimethylphenylacetic acid in 66.8% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.25 (5H, m), 7.20-7.00 (3H, m), 5.66 (1H, dd, J=5.1, 11.4Hz), 4.25-4.10 (1H, m), 3.87 (1H, d, J=13.9Hz), 3.67 (1H, d, J=13.9Hz), 3.37 (1H, dd, J=11.7, 12.1Hz), 3.00-2.85 (1H, m), 2.71 (1H, d, J=9.9Hz), 2.55 (1H, dd, J=5.5, 12.5Hz), 2.42 (1H, dd, J=5.1, 9.9Hz), 2.35-2.05 (9H, m, including each 3H, s, at 2.22 and 2.21ppm), 1.80-1.35 (2H, m).

IR(neat) : 3350, 1630cm⁻¹.

MS m/z: 368(M⁺).

HCl salt: amorphous solid.

Anal. Calcd for C₂₂H₂₈N₂O₃·HCl·1.8H₂O : C, 60.42 ; H, 7.51; N, 6.41 .

Found : C, 60.51 ; H, 7.71 ; N, 6.29 .

Example 37

2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(R)-phenylethyl]acetamide

This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(S)-phenylethanol and 3,4-dichlorophenylacetic acid in 32.8% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.25 (7H, m), 7.13 (1H, dd, J=1.5, 8.1Hz), 5.61 (1H, dd, J=5.5, 10.6Hz), 5.00-3.90 (2H, almost flat br.s), 4.45-4.35 (1H, m), 3.85 (1H, d, J=14.7Hz), 3.77 (1H, d, J=14.3Hz), 3.37 (1H, dd, J=11.0, 12.5Hz), 2.89 (1H, dd, J=4.7, 8.4Hz), 2.80-2.60 (3H, m), 2.45-2.35 (1H, m), 2.15-2.00 (1H, m), 1.80-1.65 (1H, m).

IR(KBr) : 3450, 3250, 1650cm⁻¹.

MS m/z: 408(M⁺).

mp 125.5-126.0 °C.

[α]_D = -95.4 ° (c=0.218, methanol).

Anal. Calcd for $C_{20}H_{22}Cl_2N_2O_3$: C, 58.69 ; H, 5.42; N, 6.84 .

Found : C, 58.51 ; H, 5.42 ; N, 6.70 .

Example 38

5 2-(3,4-Difluorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-phenylethanol and 3,4-difluorophenylacetic acid in 53.6% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

10 ^1H NMR (270MHz, $CDCl_3$) δ 7.40-7.25 (5H, m), 7.18-6.95 (3H, m), 5.65 (1H, dd, $J=5.5, 11.4Hz$), 5.00-3.90 (2H, almost flat br.s), 4.35-4.25 (1H, m), 3.82 (1H, d, $J=14.3Hz$), 3.74 (1H, d, $J=14.3Hz$), 3.40 (1H, dd, $J=10.6, 13.2Hz$), 2.95 (1H, dt, $J=4.4, 8.8Hz$), 2.75 (1H, d, $J=10.6Hz$), 2.61 (1H, dd, $J=5.1, 12.5Hz$), 2.51 (1H, dd, $J=5.5, 10.6Hz$), 2.40-2.10 (2H, m), 1.70-1.50 (1H, m).

IR(neat) : 3350, 3250, 1630 cm^{-1} .

15 MS m/z: 376(M^+).

HCl salt : amorphous solid.

Anal. Calcd for $C_{20}H_{22}F_2N_2O_3 \cdot HCl \cdot 0.5H_2O$: C, 56.94 ; H, 5.73; N, 6.64 .

Found : C, 57.21 ; H, 6.07 ; N, 6.63 .

Example 39

20 2-(Benzo[b]thiophen-4-yl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-phenylethanol and (benzo[b]thiophen-4-yl)acetic acid in 48.8% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

25 ^1H NMR (270MHz, $CDCl_3$) δ 7.79 (1H, d, $J=7.7Hz$), 7.66 (1H, d, $J=5.5Hz$), 7.50-7.20 (8H, m), 5.60 (1H, dd, $J=5.5, 11.4Hz$), 4.60-3.20 (2H, almost flat br.s), 4.32 (1H, d, $J=13.6Hz$), 4.01 (1H, d, $J=13.6Hz$), 4.00-3.90 (1H, m), 3.30 (1H, dd, $J=11.7, 12.1Hz$), 2.70-2.45 (3H, m), 2.28 (1H, dd, $J=5.1, 10.3Hz$), 2.20-2.10 (1H, m), 1.95-1.80 (1H, m), 1.20-1.05 (1H, m).

IR(neat) : 3400, 3200, 1630cm⁻¹.

MS m/z: 396(M⁺).

HCl salt : amorphous solid.

Anal. Calcd for C₂₂H₂₄N₂O₃S·HCl·0.5H₂O : C, 59.79 ; H, 5.93; N, 6.34 .

5 Found : C, 59.85 ; H, 6.09 ; N, 6.27 .

Example 40

N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(3,4-methylenedioxyphenyl)acetamide

10 This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-phenylethanol and 3,4-methylenedioxyphenylacetic acid in 59.7% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.25 (5H, m), 6.85-6.70 (3H, m), 5.92 (2H, s), 5.66 (1H, dd, J=5.5, 11.4Hz), 4.50-3.30 (2H, almost flat br.s), 4.30-4.20 (1H, m), 3.86 (1H, d, J=13.6Hz), 3.64 (1H, d, J=13.9Hz), 3.39 (1H, t, J=12.1Hz), 3.05-2.95 (1H, m), 2.72 (1H, d, J=10.3Hz), 2.59 (1H, dd, J=5.5, 12.5Hz), 2.48 (1H, dd, J=5.5, 10.3Hz), 2.35-2.10 (2H, m), 1.65-1.50 (1H, m).

15 IR(neat) : 3400, 3250, 1630cm⁻¹.

MS m/z: 384(M⁺).

HCl salt : amorphous solid.

20 Anal. Calcd for C₂₁H₂₄N₂O₅·HCl·1.4H₂O : C, 56.54 ; H, 6.28; N, 6.28 .

Found : C, 56.74 ; H, 6.38 ; N, 5.89 .

Example 41

2-(3,5-Difluorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

25 This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-phenylethanol and 3,4-difluorophenylacetic acid in 40.0% yield according to a procedure similar to that described in Preparation 4, Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.40-7.25 (5H, m), 6.82 (2H, d, J=8.1Hz), 6.72-6.64 (1H, m), 5.65 (1H, dd, J=5.1, 11.0Hz), 5.30-4.20 (2H, almost flat br.s), 4.40-4.30

38

(1H, m), 3.86 (1H, d, $J=14.3\text{Hz}$), 3.74 (1H, d, $J=14.3\text{Hz}$), 3.41 (1H, dd, $J=11.7, 12.1\text{Hz}$), 3.10-2.95 (1H, m), 2.76 (1H, d, $J=10.6\text{Hz}$), 2.61 (1H, dd, $J=5.1, 12.5\text{Hz}$), 2.52 (1H, dd, $J=5.5, 10.6\text{Hz}$), 2.40-2.10 (2H, m), 1.70-1.55 (1H, m).

IR(neat) : 3350, 3200, 1630 cm^{-1} .

5 MS m/z: 376(M^+).

HCl salt : amorphous solid.

Anal. Calcd for $\text{C}_{20}\text{H}_{22}\text{F}_2\text{N}_2\text{O}_3 \cdot \text{HCl} \cdot 0.5\text{H}_2\text{O}$: C, 56.94 ; H, 5.73; N, 6.64 .

Found : C, 57.01 ; H, 5.93 ; N, 6.45 .

Preparation 8

10 1-Benzyl-3-(R)-tetrahydropyranyloxypyrrolidine

To a stirred solution of (R)-(+)-1-benzyl-3-pyrrolidinol (5.00g, 28mmol) and D-camphor-10-sulfonic acid (6.97g, 30mmol) in CH_2Cl_2 (10ml) was added 3,4-dihydro-2H-pyran (20ml) at rt and the reaction mixture was stirred for 14h (in most cases, the reaction was completed after exothermic reaction subsided). The reaction mixture was
15 diluted with CH_2Cl_2 (100ml), washed with saturated NaHCO_3 aqueous solution, dried (Na_2SO_4), and concentrated to give brown oil. This was purified by column chromatography(silica gel:200g, $\text{CH}_2\text{Cl}_2/\text{MeOH}$:40/1 as eluent) to give 8.78g (97.6%) of desired compound as brown oil.

¹H NMR (270MHz, CDCl_3) δ 7.34-7.22 (5H, m), 4.61 (0.5H, dd, $J=2.9, 4.4\text{Hz}$),
20 4.54 (0.5H, dd, $J=2.9, 4.4\text{Hz}$), 4.42-4.31 (1H, m), 3.90-3.79 (1H, m), 3.67 (1H, d, $J=12.8\text{Hz}$), 3.59 (0.5H, d, $J=12.8\text{Hz}$), 3.58 (0.5H, d, $J=12.8\text{Hz}$), 3.50-3.40 (1H, m), 2.88 (0.5H, dd, $J=6.6, 10.3\text{Hz}$), 2.74-2.45 (3.5H, m), 2.25-2.05 (1H, m), 1.95-1.45 (7H, m).

Preparation 9

25 3-(R)-Tetrahydropyranyloxypyrrolidine

A mixture of 1-benzyl-3-(R)-tetrahydropyranyloxypyrrolidine (8.78g, 27.3mmol) and Pearlman's catalyst (3.50g) in MeOH (100ml) was stirred under hydrogen atmosphere at rt for 4h. After removal of the catalyst by Celite filtration, the filtrate was concentrated to give 5.74g of clear light brown oil. This was used for the next

reaction without purification.

¹H NMR (270MHz, CDCl₃) δ 4.62 (1H, br.s), 4.45-4.30 (1H, m), 3.90-3.80 (1H, m), 3.55-3.45 (1H, m), 3.20-2.80 (5H, m), 2.00-1.40 (8H, m).

Preparation 10

5 1-(S)-Phenyl-2-(3-(R)-tetrahydropyranyloxy pyrrolidin-1-yl)ethanol and 2-(R)-Phenyl-2-(3-(R)-tetrahydropyranyloxy pyrrolidin-1-yl)ethanol

10 A mixture of 3-(R)-tetrahydropyranyloxy pyrrolidine (1.43g, 8.32 mmol) and (S)-(-)-styrene oxide (1.00g, 8.32mmol) in EtOH (10ml) was refluxed with stirring for 1h. Evaporation of the solvent gave 3.098g of brown oil, which was purified by column chromatography(silica gel:100g, CH₂Cl₂/MeOH:40/1 to 15/1 as eluent) to afford 1.68g (69.3%) of clear light brown oil as about 2 to 1 mixture of title compounds in which 1-(S)-phenyl-2-(3-(R)-tetrahydro-pyranyloxy pyrrolidin-1-yl)ethanol was main.

15 ¹H NMR (270MHz, CDCl₃) δ 7.40-7.24 (5H, m), 4.72 and 4.68 (total 0.67H, app.each d, J=2.6Hz, OCH₂O), 4.63-4.55 (1H,m, PhCH₂OH and OCH₂O), 4.43-4.25 (1H, m, OCH₂CH₂N), 3.89-3.81 (1.67H,m), 3.52-3.46 (1.33H, m), 2.88-2.47 (5.33H, m), 2.15-1.90 (2H, m), 1.86-1.66 (3H, m), 1.58-1.51 (4H, m).

Example 42

20 2-(3,4-Dichlorophenyl)-N-[1-(S)-phenyl-2-(3-(R)-tetrahydropyranyloxy pyrrolidin-1-yl)ethyl]-N-tetrahydropyranyloxyacetamide

25 To a stirred solution of 1-(S)-phenyl-2-(3-(R)-tetrahydropyranyloxy pyrrolidin-1-yl)ethanol (1.67g, 5.73mmol) and Et₃N (0.96ml, 6.88mmol) in CH₂Cl₂ (20ml) was added dropwise mesyl chloride (0.53ml, 6.88mmol) at 0 °C. The reaction mixture was stirred at rt for 16h. The reaction mixture was washed with saturated NaHCO₃ aqueous solution and brine, dried(Na₂SO₄), and concentrated to give 2.02g of brown oil. This oil was used for next reaction without purification.

¹H NMR (270MHz, CDCl₃) δ 7.42-7.30 (5H, m), 4.94 (1H, dd, J=5.9, 8.1Hz, PhCH₂Cl), 4.60 and 4.52 (total 1H, each m, OCH₂O), 4.35-4.31 (1H, m, OCH₂CH₂N),

3.88-3.82 (1H, m), 3.48-3.45 (1H, m), 3.25-3.17 (1H, m), 3.02-2.69 (3H, m), 2.66-2.50 (3H, m), 1.88-1.67 (3H, m), 1.56-1.51 (4H, m).

A mixture of crude chloride derivative (2.02g, 5.73mmol) and *O*-(tetrahydropyranyl)hydroxylamine (0.806g, 6.88mmol) in EtOH (10ml) was refluxed with stirring for 0.5h. The reaction mixture was concentrated, diluted with CH₂Cl₂ (30ml), washed with saturated NaHCO₃ aqueous solution and brine, dried (Na₂SO₄), and concentrated to give 2.59g of brown oil. This oil was used for the next reaction without purification.

A mixture of the above crude amine derivative (2.59g, 5.73mmol), 3,4-dichlorophenylacetic acid (1.41g, 6.88mmol), and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (abbreviated as WSC, 1.32g, 6.88mmol) in CH₂Cl₂ (15ml) was stirred at rt for 0.5h. The reaction mixture was washed with saturated NaHCO₃ aqueous solution and brine, dried (Na₂SO₄), and concentrated to give 4.12g of brown oil. This oil was purified by column chromatography (silica gel:100g, CH₂Cl₂/MeOH:50/1 to 40/1 as eluent) to give 2.22g(67.1%) of pale yellow oil.

Example 43

2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(*R*)-hydroxypyrrolidin-1-yl)-1-(*S*)-phenylethyl]acetamide

A mixture of above amide derivative (2.20g, 3.81mmol) and HCl gas containing MeOH (10ml) was stirred at rt for 1h. The reaction mixture was concentrated, basified with NH₃ aqueous solution, and extracted with CH₂Cl₂ (30ml). The extract was washed with brine, dried (Na₂SO₄), and concentrated to give light brown powder. This was collected by filtration and washed with hexane to afford 1.117g (71.6%) of light brown powder.

¹H NMR (270MHz, CDCl₃) δ 7.41-7.28 (7H, m), 7.13 (1H, dd, J = 1.8, 8.4Hz), 5.61 (1H, dd, J = 5.5, 10.6Hz), 4.50-3.50 (2H, almost flat br.s), 4.40-4.35 (1H, m), 3.84 (1H, d, J = 14.7Hz), 3.77 (1H, d, J = 14.3Hz), 3.38 (1H, dd, J = 11.0, 12.1Hz), 2.94-2.85 (1H, m), 2.74-2.63 (3H, m), 2.44-2.35 (1H, m), 2.15-2.01 (1H, m), 1.80-1.65 (1H, m). IR(KBr) : 3250, 1650cm⁻¹.

MS m/z : 408(M⁺)

HCl salt : amorphous solid.

Anal. Calcd for C₂₀H₂₂Cl₂N₂O.HC·0.8H₂O: C, 52.20; H, 5.39; N, 6.09. Found: C, 52.22; H, 5.39; N, 6.12.

5

Example 44

2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(R)-hydroxypyrrolidin-1-yl)-1-(R)-phenylethyl]acetamide

This was prepared from 3-(R)-tetrahydropyranyloxypyrrolidine and (R)-(+)-styrene oxide in 33.3% yield according to the procedure similar to that described in Examples 3 to 5.

10

¹H NMR (270MHz, CDCl₃) δ 7.38 (1H, d, J=8.4Hz), 7.36-7.25 (6H, m), 7.13 (1H, dd, J = 1.8, 8.1Hz), 5.64 (1H, dd, J=5.1, 11.0Hz), 5.00-3.50 (2H, almost flat br.s), 4.35-4.25 (1H, m), 3.84 (1H, d, J=14.3Hz), 3.73 (1H, d, J=13.2Hz), 3.40 (1H, dd, J=11.4, 12.5Hz), 3.05-2.95 (1H, m), 2.74 (1H, br.d, J=10.3Hz), 2.62 (1H, dd, J=5.1, 12.5Hz), 2.51 (1H, dd, J=5.5, 10.6Hz), 2.40-2.25 (1H, m), 2.25-2.10 (1H, m), 1.70-1.55 (1H, m).

15

IR(KBr) : 3400, 3200, 1640cm⁻¹.

MS m/z : 408(M⁺)

HCl salt : amorphous solid

20

Anal. Calcd for C₂₀H₂₂Cl₂N₂O₃·HCl·0.5H₂O: C, 52.82; H, 5.32; N, 6.16. Found: C, 52.71; H, 5.59; N, 6.15.

Preparation 11

(S)-1-(3-Methylphenyl)-1,2-ethanediol

A mixture of 3-methylstyrene (1.69ml, 12.7mmol), and AD-mix-α (17.78g, 12.7mmol) in water (65ml) and *t*-BuOH (65ml) was stirred at 0 °C for 3.5h. To this reaction mixture was added Na₂SO₃ (20g) and the mixture was stirred at rt for 1h. The reaction mixture was extracted with ethyl acetate. The extract was washed with brine, dried (Na₂SO₄), and concentrated to give 2.07g of light brown oil, which was purified by column chromatography (silica gel: 110g, ethyl acetate/hexane:3/2) to afford

25

1.89g(98%) of desired product as light brown oil.

¹H NMR (270MHz, CDCl₃) δ 7.24 (1H, dd, J=7.3, 7.7Hz), 7.19-7.09 (3H, m), 4.77 (1H, dd, J=3.7, 8.1Hz), 3.74 (1H, dd, J=3.7, 11.4Hz), 3.65 (1H, dd, J=8.1, 11.4Hz), 2.82 (1H, br.s), 2.35 (3H, s), 1.77 (1H, br.s).

5

Preparation 12

(S)-1-(3-Methylphenyl)-1,2-ethanediol 2-tosylate

To a stirred solution of (S)-1-(3-methylphenyl)-1,2-ethanediol (1.78g, 11.7mmol) in pyridine (35ml) was added p-toluenesulfonyl chloride (2.46g, 12.9mmol), and 4-dimethylaminopyridine (1.58g, 12.9mmol) at 0 °C and the reaction mixture was stirred at 0 °C to rt for 17h. The reaction mixture was acidified with 6N HCl aqueous solution and extracted with CH₂Cl₂. The extract was washed with water and brine, dried (Na₂SO₄), and concentrated to give 3.02g of yellow oil, which was purified by column chromatography (silica gel: 100g, ethyl acetate/hexane:1/9 to 1/3) to afford 2.63g (73%) of desired product as light yellow oil. Its optical purity was 97% ee by HPLC employing a chiral stationary phase (chiral pak AS, Daicel Chemical Industries, eluted with n-hexane/EtOH:98/2; detection time: 55min for (R)-isomer 59min for (S)-isomer).

15

20

¹H NMR (270MHz, CDCl₃) δ 7.77 (2H, d, J=8.4Hz), 7.33 (2H, d, J=8.1Hz), 7.22 (1H, dd, J=7.7, 8.1Hz), 7.15-7.05 (3H, m), 4.94 (1H, ddd, J=2.9, 2.9, 8.4Hz), 4.15 (1H, dd, J=2.9, 10.3Hz), 4.04 (1H, dd, J=8.4, 10.3Hz), 2.54 (1H, br.d, J=2.9Hz), 2.45 (3H, s), 2.33 (3H, s), 1.58 (3H, s).

Preparation 13

2-(3-(S)-Methoxymethyloxypyrrolidin-1-yl)-1-(S)-(3-methylphenyl)-ethanol and 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-(3-methylphenyl)ethanol

25

A mixture of (S)-1-(3-methylphenyl)-1,2-ethanediol 2-tosylate (2.63g, 8.59mmol), (S)-3-methoxymethyloxypyrrolidine (1.24g, 9.45mmol), and K₂CO₃ (1.31g, 9.45mmol) in ethanol (25ml) was refluxed with stirring for 2h. After removal of the solvent by evaporation, the residue was diluted with water and extracted with CH₂Cl₂. The extract was washed with brine, dried (Na₂SO₄), and concentrated to give

2.11g of brown oil, which was purified by column chromatography (silica gel:110g, CH₂Cl₂/MeOH: 15/1 to 10/1) to afford 1.72g (76%) of 3 to 2 mixture of desired products as a light brown oil.

¹H NMR (270MHz, CDCl₃) δ 7.26-7.05 (4H, m), 4.68 (0.6H, dd, J=2.9, 10.6Hz, PhCH₂OH), 4.67 (0.6H, d, J=7.0Hz, OCH₂O), 4.63 (0.6H, d, J=6.6Hz, OCH₂O), 4.62 (0.4H, d, J=7.0Hz, OCH₂O), 4.59 (0.4H, d, J=7.0Hz, OCH₂O), 4.34-4.24 (0.6H, m, OCHCH₂N), 4.24-4.16 (0.4H, m, OCHCH₂N), 3.88 (0.4H, dd, J=6.2, 10.6Hz, CHCH₂OH), 3.79 (0.4H, dd, J=5.8, 11.0Hz, CHCH₂OH), 3.47 (0.4H, dd, J=5.8, 6.2Hz, NCHPh), 3.38 (1.8H, s), 3.33 (1.2H, s), 3.05-2.92 (1.2H, m), 2.82-2.40 (4H, m), 2.35 (3H, s), 2.25-1.50 (3H, m).

Example 45

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-1-(S)-(3-methylphenyl)ethyl]-N-tetrahydropyranyloxyacetamide

This was prepared from a mixture of 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-1-(S)-(3-methylphenyl)ethanol and 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-(3-methylphenyl)ethanol in 60% yield according to the procedure similar to that described in Example 4.

¹H NMR (270MHz, CDCl₃) δ 7.39 (0.5H, d, J=1.8Hz), 7.35 (0.5H, d, J=8.4Hz), 7.27-7.02 (5.5H, m), 6.96 (0.5H, dd, J=1.8, 8.4Hz), 5.65 (0.5H, dd, J=5.1, 11.4Hz, PhCHN), 5.52 (0.5H, dd, J=4.8, 11.0Hz, PhCHN), 5.30-5.20 (1H, m, NOCHO), 4.64 (0.5H, d, J=6.6Hz, OCH₂O), 4.63 (0.5H, d, J=7.0Hz, OCH₂O), 4.61 (0.5H, d, J=6.6Hz, OCH₂O), 4.60 (0.5H, d, J=6.6Hz, OCH₂O), 4.30-4.20 (0.5H, m, OCHCH₂N), 4.20-4.10 (0.5H, m, OCHCH₂N), 4.06-3.85 (3H, m), 3.56-3.36 (1.5H, m), 3.35 (1.5H, s, OMe), 3.34 (1.5H, s, OMe), 3.24-3.10 (0.5H, m), 3.01-2.80 (2H, m), 2.66-2.40 (3H, m), 2.34 (1.5H, s), 2.28 (1.5H, s), 2.15-1.15 (8H, m).

Example 46

2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-(3-methylphenyl)ethyl]acetamide

This was prepared from 2-(3,4-dichlorophenyl)-N-[2-(3-(*S*)-methoxymethoxy-pyrrolidin-1-yl)-1-(*S*)-(3-methylphenyl)ethyl]-N-tetrahydropyranyloxyacetamide in 77% yield according to the procedure similar to that described in Example 5

¹H NMR (270MHz, CDCl₃) δ 7.42-7.05 (7H, m), 5.59 (1H, dd, J=5.1, 11.0Hz, PhCHN), 4.35-4.25 (1H, m, CHOH), 3.85 (1H, d, J=14.3Hz, COCH₂Ph), 3.74 (1H, d, J=15.8Hz, COCH₂Ph), 3.50-2.50 (2H, almost flat br.s, OHx2), 3.38 (1H, dd, J=11.7, 12.1Hz), 3.00-2.90 (1H, m), 2.73 (1H, br.d, J=10.6Hz), 2.62 (1H, dd, J=5.1, 12.5Hz), 2.53 (1H, dd, J=5.5, 10.6Hz), 2.40-2.25 (4H, m, including 3H, s at 2.30ppm), 2.23-2.07 (1H, m), 1.65-1.55 (1H, m).

IR(neat) : 3350, 1650cm⁻¹.

MS m/z : 422(M⁺)

HCl salt : amorphous solid

Anal. Calcd for C₂₁H₂₄Cl₂N₂O₃·HCl·1.5H₂O: C, 51.81; H, 5.80; N, 5.75. Found: C, 51.85; H, 5.72; N, 5.47.

15

Example 47

N-[1-(*S*)-(4-Chlorophenyl)-2-(3-(*S*)-hydroxypyrrolidin-1-yl)ethyl]-2-(3,4-dichlorophenyl)-N-hydroxyacetamide

This was prepared from 4-chlorostyrene and 3-(*S*)-methoxymethoxypyrrolidine in 12% overall yield according to a procedure similar to that described in Examples 7 to 11.

20

¹H NMR (270MHz, CDCl₃) δ 7.40 (1H, d, J=2.2Hz), 7.36 (1H, d, J=8.4Hz), 7.30-7.20 (4H, m), 7.14 (1H, dd, J=2.2, 8.1Hz), 5.58 (1H, dd, J=5.1, 11.0Hz, PhCHN), 5.00-3.00 (2H, almost flat br.s, OHx2), 4.35-4.25 (1H, m, CHOH), 3.85 (1H, d, J=14.3Hz, COCH₂Ph), 3.72 (1H, d, J=13.9Hz, COCH₂Ph), 3.33 (1H, t, J=11.7Hz), 3.00-2.85 (1H, m), 2.74 (1H, br.d, J=10.3Hz), 2.65 (1H, dd, J=5.1, 12.5Hz), 2.60-2.45 (1H, m), 2.45-2.25 (1H, m), 2.25-2.05 (1H, m), 1.70-1.50 (1H, m).

25

HCl salt : amorphous solid

IR(KBr) : 3400, 3100, 1650cm⁻¹.

MS m/z : 443(M+H)⁺.

30

Anal. Calcd for C₂₀H₂₁Cl₃N₂O₃·HCl·0.7H₂O: C, 48.74; H, 4.79; N, 5.68. Found: C,

49.15; H, 5.21; N, 5.58.

Example 48

5 2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-(4-methoxyphenyl)ethyl]acetamide and 2-(3,4-dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(R)-(4-methoxyphenyl)-ethyl]acetamide

This was prepared from 4-methoxystyrene and 3-(S)-methoxymethyloxypyrrolidine in 5.2% overall yield according to a procedure similar to that described in Examples 7 to 11.

10 In this case racemization occurred at 1-position to afford the title compounds during the following reactions (mesylation, addition of THPONH₂, and acylation).

¹H NMR (270MHz, CDCl₃) δ 7.40-7.26 (4H, m), 7.12 (0.5H, dd, J=2.2, 8.4Hz), 7.11 (0.5H, dd, J=2.6, 8.4Hz), 6.84 (2H, d, J=8.4Hz), 5.70-5.60 (1H, m, PhCHN), 4.50-4.40 (0.5H, m, CHOH), 4.50-3.00 (2H, almost flat br.s, OHx2), 4.40-4.30 (0.5H, m, CHOH), 3.84 (1H, d, J=14.3Hz, COCH₂Ph), 3.79 (3H, s), 3.73 (1H, d, J=14.7Hz, COCH₂Ph), 3.65-3.40 (1H, m), 3.15-3.00 (1H, m), 2.90-2.40 (4H, m), 2.30-2.10 (1H, m), 1.90-1.78 (0.5H, m), 1.78-1.60 (0.5H, m).

HCl salt : amorphous solid

1R(KBr) : 3400, 3150, 1650cm⁻¹.

MS m/z : 438(M⁺)

20 Anal. Calcd for C₂₁H₂₄Cl₂N₂O₄·HCl·2.5H₂O: C, 48.43; H, 5.81; N, 5.38. Found: C, 48.21; H, 5.75; N, 5.35.

Example 49

2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-(4-trifluoromethylphenyl)ethyl]acetamide

25 This was prepared from 4-trifluoromethylstyrene and 3-(S)-methoxymethyloxypyrrolidine in 25.3% overall yield according to a procedure similar to that described in Examples 7 to 11.

¹H NMR (270MHz, CDCl₃) δ 7.60-7.35 (6H, m), 7.20-7.10 (1H, m), 5.65 (1H, dd, J=5.5, 11.0Hz, PhCHN), 4.40-4.30 (1H, m, CHOH), 3.90 (1H, d, J=13.9Hz,

COCH₂Ph), 3.73 (1H, d, J=12.5Hz, COCH₂Ph), 3.34 (1H, dd, J=11.0, 12.5Hz), 3.00-2.90 (1H, m), 2.75-2.65 (2H, m), 2.54 (1H, dd, J=5.1, 10.6Hz), 2.50-2.00 (4H, m), 1.70-1.55 (1H, m).

IR(neat) : 3400, 3250, 1635cm⁻¹.

5 MS m/z : 476(M⁺)

HCl salt : amorphous solid

Anal. Calcd for C₂₁H₂₁Cl₂F₃N₂O₃·HCl·2H₂O: C, 45.88; H, 4.77; N, 5.10. Found: C, 45.90; H, 4.83; N, 4.71.

Preparation 14

10 (S)-1-(4-Methylphenyl)-1,2-ethanediol 2-tosylate

This was prepared from 4-methylstyrene in 75% overall yield according to a procedure similar to that described in Examples 7 and 8. Optical purity was 98.3% ee by HPLC analysis.

15 ¹H NMR (270MHz, CDCl₃) δ 7.77 (2H, d, J=8.1Hz), 7.33 (2H, d, J=8.4Hz), 7.20 (2H, d, J=8.1Hz), 7.14 (2H, d, J=8.1Hz), 5.00-4.90 (1H, m), 4.13 (1H, dd, J=3.3, 10.3Hz), 4.03 (1H, d, J=8.4, 10.3Hz), 2.49 (1H, d, J=2.9Hz), 2.45 (3H, s), 1.57 (3H, s).

Preparation 15

(S)-4-Methylstyrene oxide

20 A mixture of (S)-1-(4-methylphenyl)-1,2-ethanediol 2-tosylate (4.13g, 13.5mmol) and 50% NaOH aqueous solution (5ml) in THF (25ml) was stirred at rt for 1h and at 50 °C for 2h. After cooling down to rt, the reaction mixture was diluted with water and extracted with ethyl acetate. The extract was washed with water and brine, dried (Na₂SO₄), and concentrated to give 1.59g (88%) of desired compound as pale brown oil. This oil was used for next reaction without purification.

25 ¹H NMR (270MHz, CDCl₃) δ 7.20-7.10 (4H, m), 3.83 (1H, dd, J=2.6, 4.0Hz), 3.13 (1H, dd, J=4.0, 5.5Hz), 2.80 (1H, dd, J=2.6, 5.5Hz), 2.34 (3H, s).

Preparation 16

2-(3-(S)-Methoxymethyloxypyrrolidin-1-yl)-1-(S)-(4-methylphenyl)-ethanol and 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-(4-methylphenyl)ethanol

A mixture of (S)-4-methylstyrene oxide (1.59g, 11.9mmol) and 3-(S)-methoxymethyloxypyrrolidine (1.55g, 11.9mmol) in isopropanol (25ml) was refluxed for 7h. The solvent was evaporated and the residue was purified by column chromatography (silica gel: 150g, CH₂Cl₂/MeOH: 50/1 to 15/1) to give 2.39g (76%) of desired products as a pale brown oil. This was 3 to 2 mixture of title compounds. ¹H NMR (270MHz, CDCl₃) δ 7.26 (1.2H, d, J=8.1Hz), 7.21-7.10 (2.8H, m), 4.75-4.55 (2.6H, m, including 0.6H, d, J=6.6Hz at 4.66ppm, 0.6H, d, J=7.0Hz at 4.63ppm, 0.4H, d, J=7.0Hz at 4.62ppm, 0.4H, d, J=7.0Hz at 4.58ppm), 4.35-4.23 (0.6H, m, OCHCH₂N), 4.23-4.15 (0.4H, m, OCHCH₂N), 3.87 (0.4H, dd, J=6.2, 10.6Hz, CHCH₂OH), 3.77 (0.4H, dd, J=5.9, 10.6Hz, CHCH₂OH), 3.49 (0.4H, dd, J=5.9, 6.2Hz, NCHPh), 3.38 (1.8H, s), 3.33 (1.2H, s), 3.05-2.90 (1.2H, m), 2.80-2.40 (5H, m), 2.34 (3H, s), 2.25-2.00 (1H, m), 1.95-1.75 (1H, m).

Example 50

2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-(4-methylphenyl)ethyl]acetamide

This was prepared from 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-1-(S)-(4-methylphenyl)ethanol and 2-(3-(S)-methoxymethyloxypyrrolidin-1-yl)-2-(R)-(4-methylphenyl)ethanol in 29.5% overall yield according to a procedure similar to that described in Examples 10 and 11.

¹H NMR (270MHz, CDCl₃) δ 7.40-7.30 (2H, m), 7.23 (2H, app.d, J=8.1Hz), 7.11 (3H, app.d, J=7.7Hz), 5.64 (1H, dd, J=5.1, 11.4Hz, PhCHN), 5.00-3.00 (2H, almost flat br.s, OHx2), 4.40-4.30 (1H, m, CHOH), 3.84 (1H, d, J=14.7Hz, COCH₂Ph), 3.73 (1H, d, J=14.3Hz, COCH₂Ph), 3.46 (1H, dd, J=11.4, 12.1Hz), 3.10-2.95 (1H, m), 2.83 (1H, br.d, J=11.0Hz), 2.75-2.40 (3H, m), 2.32 (3H, s), 2.25-2.10 (1H, m), 1.75-1.60 (1H, m).

HCl salt : amorphous solid

MS m/z : 422(M⁺)

IR(KBr) : 3420, 3180, 1650cm⁻¹.

Anal. Calcd for $C_{21}H_{24}Cl_2N_2O_3 \cdot HCl \cdot 0.5H_2O$: C, 53.80; H, 5.59; N, 5.98. Found: C, 53.51; H, 5.67; N, 6.04.

Preparation 17

(S)-1-(3-Methoxymethoxyphenyl)-1,2-ethanediol

5 This was prepared from 3-methoxymethoxystyrene (prepared by methoxymethylation of 3-hydroxystyrene in a standard manner) in quantitative yield according to a procedure similar to that described in Example 7.

^1H NMR (270MHz, $CDCl_3$) δ 7.25 (1H, dd, $J=7.7, 8.1$ Hz), 7.03 (1H, d, $J=1.8$ Hz), 6.98-6.92 (2H, m), 5.15 (2H, s, OCH_2OMe), 4.74 (1H, dd, $J=3.3, 8.1$ Hz, ArCH $\underline{O}H$), 3.71 (1H, br.d, $J=9.9$ Hz, CHCH $\underline{2}OH$), 3.65-3.55 (2H, m, including 1H, dd, $J=8.1, 11.0$ Hz at 3.61ppm, CHCH $\underline{2}OH$), 3.44 (3H, s, OCH_2OMe), 3.14 (1H, br.s, $\underline{O}H$).

Preparation 18

(S)-1-(3-Methoxymethoxyphenyl)-1,2-ethanediol 2-tosylate

15 This was prepared from (S)-1-(3-methoxymethoxyphenyl)-1,2-ethanediol in 64% yield according to a procedure similar to that described in Example 8. Its optical purity was 96%ee by HPLC.

^1H NMR (270MHz, $CDCl_3$) δ 7.77 (2H, d, $J=8.4$ Hz), 7.34 (2H, d, $J=8.1$ Hz), 7.25 (1H, dd, $J=7.7, 8.4$ Hz), 7.00-6.92 (3H, m), 5.15 (2H, s), 4.95 (1H, ddd, $J=3.3, 3.3, 8.4$ Hz, ArCH $\underline{O}H$), 4.15 (1H, dd, $J=3.3, 10.3$ Hz, CHCH $\underline{2}OTs$), 4.03 (1H, dd, $J=8.4, 10.3$ Hz, CHCH $\underline{2}OTs$), 3.46 (3H, s, OCH_2OMe), 2.65 (1H, d, $J=3.3$ Hz, ArCH $\underline{O}H$), 2.45 (3H, s, Ph $\underline{Me}$).

Example 51

2-(3,4-Dichlorophenyl)-N-[1-(S)-(3-methoxymethoxyphenyl)-2-(3-(S)-tetrahydropyranyloxy-pyrrolidin-1-yl)ethyl]-N-tetrahydropyranyloxy-acetamide

25 This was prepared from (S)-1-(3-methoxymethoxyphenyl)-1,2-ethanediol 2-tosylate in 52% overall yield according to the procedure similar to that described in Examples 9 and 10.

¹H NMR (270MHz, CDCl₃) δ 7.42-6.91 (7H, m), 5.65 (0.5H, dd, J=3.3, 9.9Hz, PhCHN), 5.54 (0.5H, dd, J=4.4, 11.0Hz, PhCHN), 5.35-5.25 (1H, m, NOCHO), 5.19 (0.5H, d, J=6.6Hz, OCH₂O), 5.15 (0.5H, d, J=6.6Hz, OCH₂O), 5.14 (0.5H, d, J=7.0Hz, OCH₂O), 5.10 (0.5H, d, J=7.0Hz, OCH₂O), 4.65-4.55 (1H, m, CHOCHO), 4.40-4.30 (0.5H, m, OCHCH₂N), 4.30-4.20 (0.5H, m, OCHCH₂N), 4.10-3.85 (4H, m, including 0.5H, d, J=16.5Hz at 4.06ppm, 0.5H, d, J=16.5Hz at 3.92ppm, and 1H, s at 3.92ppm, COCH₂Ph), 3.68-3.15 (6H, m, including each 1.5H, s, at 3.47 and 3.46ppm, OMe), 3.02-2.80 (2H, m), 2.66-2.35 (3H, m), 2.20-1.15 (14H, m).

10

Example 522-(3,4-Dichlorophenyl)-N-hydroxy-N-[1-(S)-(3-hydroxyphenyl)-2-(3-(S)-hydroxypyrrolidin-1-yl)ethyl]acetamide

This was prepared from 2-(3,4-dichlorophenyl)-N-[1-(S)-(3-methoxymethoxyphenyl)-2-(3-(S)-tetrahydropyranyloxy-pyrrolidin-1-yl)ethyl]-N-tetrahydropyranyloxyacetamide in 46% yield according to the procedure similar to that described in Example 11.

¹H NMR (270MHz, CDCl₃ and DMSO-d₆) δ 7.56 (1H, s, PhOH), 7.40 (1H, d, J=1.8Hz), 7.37 (1H, d, J=8.4Hz), 7.17 (1H, dd, J=1.8, 8.1Hz), 7.11 (1H, dd, J=7.7, 8.1Hz), 6.90-6.70 (3H, m), 5.56 (1H, dd, J=5.1, 10.6Hz, PhCHN), 4.30-4.20 (1H, m, CHOH), 3.90 (1H, d, J=15.0Hz, COCH₂Ph), 3.74 (1H, d, J=14.5Hz, COCH₂Ph), 4.50-2.50 (2H, almost flat br.s, OHx2), 3.32 (1H, dd, J=11.4, 11.7Hz), 3.00-2.85 (1H, m), 2.75-2.55 (3H, m, including 1H, dd, J=5.1, 11.0Hz), 2.40-2.30 (1H, m), 2.15-2.00 (1H, m), 1.80-1.60 (1H, m).

IR(KBr) : 3350, 3200, 1630cm⁻¹.


MS m/z : 424(M⁺)

Free amine : mp 151.6-153.1 °C

Anal. Calcd for C₂₀H₂₂Cl₂N₂O₄.7H₂O: C, 54.85; H, 5.39; N, 6.40. Found: C, 54.70; H, 4.99; N, 6.42.

The chemical structures of the compounds prepared in the Examples 1 to 52 are summarized in the following tables.

50

TABLE

(I)

	Ex.#	A	Ar	R	X
5	1	hydrogen	(S)-phenyl	benzyl	3,4-dichlorophenyl
	2	hydrogen	(S)-phenyl	hydrogen	3,4-dichlorophenyl
	3	hydrogen	(S)-phenyl	methyl	3,4-dichlorophenyl
	4	hydrogen	(S)-phenyl	hydrogen	2,3,6-trichlorophenyl
	5	hydrogen	(S)-phenyl	hydrogen	4-trifluoro-methylphenyl
10	6	hydrogen	(S)-phenyl	hydrogen	1-naphthyl
	7	hydrogen	(S)-phenyl	hydrogen	2,4,6-trimethylphenyl
	8	hydrogen	(S)-phenyl	hydrogen	4-pyridyl
	9	hydrogen	(S)-phenyl	hydrogen	benzo[<i>b</i>]furan-4-yl
	10	(S)-tetra-hydropyranyloxy	(S)-phenyl	tetrahydro-pyranyloxy	3,4-dichlorophenyl
15	11	(S)-hydroxy	(S)-phenyl	hydrogen	3,4-dichlorophenyl
	12	(S)-hydroxy	(S)-4-fluoro-phenyl	hydrogen	3,4-dichlorophenyl
	13	(S)-hydroxy	(S)-phenyl	hydrogen	4-bromophenyl
	14	(S)-hydroxy	(S)-phenyl	hydrogen	3-bromophenyl
	15	(S)-hydroxy	(S)-phenyl	hydrogen	4-fluorophenyl
20	16	(S)-hydroxy	(S)-phenyl	hydrogen	3,4-dimethoxyphenyl
	17	(S)-hydroxy	(S)-phenyl	hydrogen	3-trifluoromethyl-phenyl
	18	(S)-hydroxy	(S)-phenyl	hydrogen	4-trifluoromethyl-phenyl
	19	(S)-hydroxy	(S)-phenyl	hydrogen	4-biphenyl
	20	(S)-hydroxy	(S)-phenyl	hydrogen	4-nitrolphenyl

	Ex.#	A	Ar	R	X
5	21	(S)-hydroxy	(S)-phenyl	hydrogen	3-nitrophenyl
	22	(S)-hydroxy	(S)-phenyl	hydrogen	4-chlorophenyl
	23	(S)-hydroxy	(S)-phenyl	hydrogen	3-chlorophenyl
	24	(S)-hydroxy	(S)-phenyl	hydrogen	2-chlorophenyl
	25	(S)-hydroxy	(S)-phenyl	hydrogen	2,3,5-trichlorophenyl
10	26	(S)-hydroxy	(S)-phenyl	hydrogen	2,4,6-trichlorophenyl
	27	(S)-hydroxy	(S)-phenyl	hydrogen	2,4,6-trimethylphenyl
	28	(S)-hydroxy	(S)-phenyl	hydrogen	2,3-dichlorophenyl
	29	(S)-hydroxy	(S)-phenyl	hydrogen	2,4-dichlorophenyl
	30	(S)-hydroxy	(S)-phenyl	hydrogen	2,5-dichlorophenyl
15	31	(S)-hydroxy	(S)-phenyl	hydrogen	2,6-dichlorophenyl
	32	(S)-hydroxy	(S)-phenyl	hydrogen	3,5-dichlorophenyl
	33	(S)-hydroxy	(S)-phenyl	hydrogen	2,3,6-trichlorophenyl
	34	(S)-hydroxy	(S)-phenyl	hydrogen	benzo[<i>b</i>]furan-4-yl
	35	(S)-hydroxy	(S)-phenyl	hydrogen	1-tetralon-6-yl
20	36	(S)-hydroxy	(S)-phenyl	hydrogen	3,4-dimethylphenyl
	37	(S)-hydroxy	(R)-phenyl	hydrogen	3,4-dichlorophenyl
	38	(S)-hydroxy	(S)-phenyl	hydrogen	3,4-difluorophenyl
	39	(S)-hydroxy	(S)-phenyl	hydrogen	benzo[<i>b</i>]thiophen-4-yl
	40	(S)-hydroxy	(S)-phenyl	hydrogen	3,4-methylene-dioxyphenyl
25	41	(S)-hydroxy	(S)-phenyl	hydrogen	3,5-difluorophenyl
	42	(R)-tetrahydro-pyranyloxy	(S)-phenyl	tetrahydro-pyranyl	3,4-dichlorophenyl
	43	(R)-hydroxy	(S)-phenyl	hydrogen	3,4-dichlorophenyl
	44	(R)-hydroxy	(R)-phenyl	hydrogen	3,4-dichlorophenyl
	45	(S)-methoxymethyloxy	(S)-3-methyl-phenyl	tetrahydro-pyranyl	3,4-dichlorophenyl


Ex.#	A	Ar	R	X
46	(S)-hydroxy	(S)-3-methyl-phenyl	hydrogen	3,4-dichlorophenyl
47	(S)-hydroxy	(S)-4-chloro-phenyl	hydrogen	3,4-dichlorophenyl
48	(S)-hydroxy	(S)-4-methoxy-phenyl	hydrogen	3,4-dichlorophenyl
5 49	(S)-hydroxy	(S)-4-trifluoro-methylphenyl	hydrogen	3,4-dichlorophenyl
50	(S)-hydroxyl	(S)-4-methyl-phenyl	hydrogen	3,4-dichlorophenyl
51	(S)-tetrahydro-pyranyloxy	(S)-3-methoxy-methyloxy-phenyl	tetrahydro-pyranyl	3,4-dichlorophenyl
52	(S)-hydroxy	(S)-3-hydroxy-phenyl	hydrogen	3,4-dichlorophenyl

68277-1

53

CLAIMS:

1. A compound of the following formula:



(I)

(wherein:

A is hydrogen, hydroxy or OY, wherein Y is a hydroxy-protecting group selected from the group consisting of benzyl, triphenylmethyl, tetrahydropyranyl, methoxymethyl and $R^1R^2R^3Si$ in which R^1 , R^2 and R^3 are each C_1 - C_6 alkyl or phenyl;

Ar is phenyl optionally substituted with one or more substituents selected from halo, hydroxyl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, CF_3 , C_1 - C_4 alkoxy- C_1 - C_4 alkoxy and carboxy- C_1 - C_4 alkoxy;

X is phenyl, naphthyl, biphenyl, indanyl, benzofuranyl, benzothiophenyl, 1-tetralone-6-yl, pyridyl, furyl or thienyl, each being optionally substituted with up to three substituents selected from halo, C_1 - C_4 alkylenedioxy, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxyl, NO_2 , CF_3 and SO_2CH_3 ; and

R is hydrogen, C_1 - C_4 alkyl or a hydroxy-protecting group selected from the group consisting of benzyl, triphenylmethyl, tetrahydropyranyl, methoxymethyl and $R^1R^2R^3Si$ in which R^1 , R^2 and R^3 are each C_1 - C_6 alkyl or phenyl) or a salt thereof.

2. A compound according to claim 1, wherein A is hydrogen or hydroxyl, and R is hydrogen or C_1 - C_4 alkyl.

68277-1

54

3. A compound according to claim 1 or 2, wherein Ar is phenyl.

4. A compound according to claim 1, 2 or 3, wherein X is phenyl substituted with up to three substituents selected from
5 chloro, methyl and CF₃, and R is hydrogen.

5. A compound according to claim 1, 2, 3 or 4, wherein X is 3,4-dichlorophenyl.

6. 2-(3,4-Dichlorophenyl)-N-hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]acetamide.

10 7. N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-(2,3,6-trichlorophenyl)acetamide.

8. N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-(4-trifluoromethylphenyl)acetamide.

9. N-Hydroxy-N-[1-(S)-phenyl-2-(1-pyrrolidinyl)ethyl]-2-
15 (2,4,6-trimethylphenyl)acetamide.

10. 2-(3,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

11. 2-(4-Bromophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

20 12. N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(4-trifluoromethylphenyl)acetamide.

13. 2-(4-Chlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

14. 2-(2,3-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.
25

15. 2-(2,4-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

68277-1

55

16. 2-(2,5-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

17. 2-(2,6-Dichlorophenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

5 18. N-Hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-2-(2,3,6-trichlorophenyl)acetamide.

19. 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.


20. 2-(3,4-Dimethylphenyl)-N-hydroxy-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide.

21. A compound according to claim 1, wherein A is OY, and R is the hydroxy protecting group.

22. A pharmaceutical composition used as an analgesic, antiinflammatory, diuretic, anesthetic or neuroprotective agent, which comprises (1) an effective amount of a compound according to any one of claims 1 to 21 or a pharmaceutically acceptable salt thereof, and (2) a pharmaceutically acceptable inert carrier.

23. A pharmaceutical composition used as an agent for treatment of stroke or functional bowel disease, which comprises (1) an effective amount of a compound according to any one of claims 1 to 21 or a pharmaceutically acceptable salt thereof, and (2) a pharmaceutically acceptable inert carrier.

24. A compound of the formula:



68277-1

56

(wherein:



A is hydrogen, hydroxy or OY, wherein Y is a hydroxy-protecting group selected from the group consisting of benzyl, triphenylmethyl, tetrahydropyranyl, methoxymethyl and $R^1R^2R^3Si$ in which R^1 , R^2 and R^3 are each C_1 - C_6 alkyl or phenyl;

Ar is phenyl optionally substituted with one or more substituents selected from halo, hydroxyl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, CF_3 , C_1 - C_4 alkoxy- C_1 - C_4 alkoxy, and carboxy- C_1 - C_4 alkoxy; and

R is hydrogen, C_1 - C_4 alkyl, or a hydroxy-protecting group selected from the group consisting of benzyl, triphenylmethyl, tetrahydropyranyl, methoxymethyl and $R^1R^2R^3Si$ in which R^1 , R^2 and R^3 are each C_1 - C_6 alkyl or phenyl) or a salt thereof.

25. A process for producing a compound of the formula (II) as defined in claim 24, which comprises:


reacting an ethanol amine compound of the formula (III) with a hydroxylamine of the formula (IV):



(wherein Ar is as defined in claim 24, L is a leaving group and A^1 is hydrogen or a hydroxyl group protected by a hydroxy-protecting group selected from the group consisting of benzyl, triphenylmethyl, tetrahydropyranyl, methoxymethyl and $R^1R^2R^3Si$ in which R^1 , R^2 and R^3 are each C_1 - C_6 alkyl or phenyl) to obtain a compound of the formula (V):

68277-1

57



(V)


(wherein Ar and A¹ have the meanings given above), and

5 then reacting a compound of the formula (V) with methanesulfonyl chloride in the presence of a base followed by addition of a protected hydroxylamine and,


where required, removal of the protecting group.

26. A process for producing a compound of the formula
10 (II) as defined in claim 24, which comprises:

reacting a pyrrolidinyll compound of the formula (VII) with a substituted or unsubstituted phenyloxide of the formula (VIII):



(VII)





(VIII)

15 (wherein A is as defined in claim 24, Q is halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkoxy-C₁-C₄ alkoxy or CF₃ and n
20 is an integer of 1 to 5) to obtain a mixture of a compound of the formula (IX) and a compound of the formula (X);

68277-1

58



5 then reacting the obtained mixture with
methanesulfonyl chloride in the presence of a base followed by
addition of a protected hydroxylamine and,

where required, removal of the protecting group.

27. A process for producing a compound of the formula (I)
10 as defined in claim 1, which comprises:

reacting a compound of the formula (II) of claim 24
wherein R is the hydroxy protecting group and A and Ar are as
defined in claim 24 with a carboxylic acid of the formula XCH_2
COOH (wherein X is as defined in claim 1),

15 removing the protecting group from the obtained
compound, and,

where necessary, alkylation of the obtained compound.

SMART & BIGGAR

OTTAWA, CANADA

PATENT AGENTS

