

(19) **DANMARK**

(10) **DK/EP 3411468 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 C 3/08 (2006.01)** **C 12 C 3/12 (2006.01)** **C 12 H 1/16 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2023-06-26**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-06-07**
- (86) Europæisk ansøgning nr.: **17705043.2**
- (86) Europæisk indleveringsdag: **2017-02-03**
- (87) Den europæiske ansøgnings publiceringsdag: **2018-12-12**
- (86) International ansøgning nr.: **EP2017052450**
- (87) Internationalt publikationsnr.: **WO2017134263**
- (30) Prioritet: **2016-02-05 EP 16154513**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Heineken Supply Chain B.V., Burgemeester Smeetsweg 1, 2382 PH Zoeterwoude, Holland**
- (72) Opfinder: **BROUWER, Eric Richard, Burgemeester Smeetsweg 1, 2382 PH Zoeterwoude, Holland**
DEKONINCK, Tinne, Burgemeester Smeetsweg 1, 2382 PH Zoeterwoude, Holland
VANBENEDEN, Nele, Burgemeester Smeetsweg 1, 2382 PH Zoeterwoude, Holland
VAN VEEN, Marcel, Burgemeester Smeetsweg 1, 2382 PH Zoeterwoude, Holland
SCHOUTEN, Maria Elizabeth Wilhelmina, Burgemeester Smeetsweg 1, 2382 PH Zoeterwoude, Holland
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **Lysstabil humleekstrakt**
- (56) Fremdragne publikationer:
WO-A1-2006/104387
WO-A2-2011/054838
GB-A- 2 362 891
US-A- 4 767 640
US-A- 5 811 144
HUVAERE KEVIN ET AL: "Flavin-induced photodecomposition of sulfur-containing amino acids is decisive in the formation of beer lightstruck flavor", PHOTOCHEMICAL AND PHOTOBIOLOGICAL SCIENCES, ROYAL SOCIETY OF CHEMISTRY, CAMBRIDGE, GB, vol. 5, no. 10, 13 September 2006 (2006-09-13), pages 961-969, XP009108620, ISSN: 1474-905X, DOI: 10.1039/B609337J

DESCRIPTION

TECHNICAL FIELD OF THE INVENTION

[0001] The present invention relates to light-stable hop extracts and to the application of such hop extracts in beer. The invention further provides a method of preparing such a light-stable hop extract, said method comprising illuminating a pre-isomerised hop extract.

BACKGROUND OF THE INVENTION

[0002] Hops are the female flowers of the hop plant *Humulus lupulus*. They are used as an ingredient in beer, to which they impart a bitter, tangy flavour. Hops are usually dried in an oast house before they are used in the brewing process. In the brewing process the wort (sugar-rich liquid produced from malt) is boiled with hops before it is cooled down and yeast is added, to start fermentation.

[0003] Boiling the wort with hops results in the extraction of alpha acids (humulones, such as humulone, adhumulone, cohumulone, posthumulone and prehumulone) into the sweet wort, which under the influence of temperature (thermal isomerisation) are partially isomerised to the corresponding iso-alpha acids (isohumulones). These iso-alpha acids are responsible for the characteristic bitter taste of hopped beer. Typical alpha acid levels in the wort at the start of the wort boiling are below 0.01 wt % (below 100 ppm). Disadvantages of this traditional hopping approach to impart bitterness are the inefficient alpha acid extraction and isomerisation, typically resulting in iso-alpha acid yields below 40%.

[0004] Hop utilization can be improved significantly by performing the alpha acid isomerisation outside the brewing process. To this end pre-isomerised hop products have been developed. These pre-isomerised hop product are produced starting from hop extract.

[0005] Hop extracts are usually obtained via liquid or supercritical carbon dioxide extraction. Carbon dioxide hop extracts provide predominantly alpha acids (humulones) next to beta acids (lupulones), and they can be further fractionated to obtain alpha acid enriched hop extracts. Starting from these alpha acid containing extracts, the "off-line" pre-isomerisation of alpha acids outside the brewing (wort boiling) process can be achieved using alkali metal and/or alkaline earth metal based compounds via two approaches. The first approach applies a solvent-free reaction medium while in the other approach the transformation is performed after addition of solvents, either pure water or alternatively water mixed with an organic solvent.

[0006] It is known to isomerise alpha acid to iso-alpha acids by means of photolysis. AU 7540281, for instance, describes a process for the up-grading of the bitter substances from hops, and more particularly of practically pure alpha- and beta acids obtained from a highly

purified hop extract, particularly by treatment with liquid carbon dioxide, characterized in that the alpha and/or beta acids are subjected to photolysis by sunlight, for a predetermined duration, under conditions adapted to produce an optimum quantic yield under suitable temperature conditions, preferably close to 60°C, to obtain respectively iso-alpha acids and deoxy-alpha acids

[0007] US 4,767,640 describes a method of preparing light stable hop product that essentially consists of reduced isohumulones, and less than 0.5% unreduced isohumulones and non-isohumulones unstable products. Example 1 describes a process in which an isomerized hop extract was reduced, followed by acidification and separation of the aqueous layer and the oily layer. The oily extract so obtained contained 70% reduced isohumulone, 6.5% background material and less than 0.5% unreduced isohumulone. Example 3 describes how the oily extract was further purified to yield a "Post Purified Aqueous Phase" and a "Post Purified Oil Phase". Example 4 describes the chromatographic separation of the oily extract of Example 1 into three fractions. Example 5 describes an experiment in which the light stability of the aforementioned purified extracts ("Post Purified Aqueous Phase" and a "Post Purified Oil Phase") and of the three chromatographic fractions was evaluated. This was done by adding these materials to beer and irradiating for 2 hours with 600 foot candles of fluorescent light.

[0008] WO 93/02177 describes a process for producing a foam stabilizing and bittering agent for a malt beverage characterized by extracting hops under conditions which favour the separation and recovery of a fraction rich in adprehumulone, and subjecting the adprehumulone fraction to photo-isomerisation to produce iso-adprehumulone.

[0009] US 5,811,144 describes a process for the production of a beer wherein wort having a high riboflavin content is hopped to produce the desired beverage, the improvement comprising treating the wort with an effective amount of actinic radiation adapted to decompose riboflavin and having a wave length of from 300 to 700 nanometers; wherein, during said treatment with the actinic radiation, the wort is maintained at a temperature of from about 25 °C - 90 °C and is substantially anaerobic; whereby the riboflavin content is reduced to less than 0.2 ppm and the resulting beer has enhanced stability to light.

[0010] WO 2006/104387 relates to a composition that can be used as an additive in beverages or foodstuffs to prevent or reduce light induced flavour changes. The composition containing at least 10 µg of substituted pyrroles per kg of dry matter and at least 100 mg per kg of dry matter of a pyranone selected from the group consisting of maltol, 2,3-dihydro-3,5-dihydroxy-6-methyl-4H-pyran-4-one and combinations thereof.

[0011] WO 2011/054838 relates to a bottle, container or beverage glass for containing a hops based beverage, including a concentration between 10 µg/l and 10 mg/l of riboflavin, said bottle, container or beverage glass being at least partially transparent or translucent to visible light and having an optical filter characteristic preventing light transmission of wavelengths between 200 nm and 510 nm to a level preventing generation of more than a tasteable concentration of MBT in said beverage through photochemical reactions and photochemically

initiated auto-catalytic reactions involving said Riboflavin.

[0012] GB 2,362,891 relates to a method of providing a beverage comprising treating a malted beverage such as beer to modify the original flavour or aroma of the beverage as brewed by exposing the beverage to a light source when the beverage is dispensed for consumption.

[0013] It is well-known that the flavour quality of beer is compromised by exposure to light, a phenomenon which is generally referred to as 'lightstruck' or 'sunstruck' flavour, and which is triggered by light-induced degradation of iso-alpha acids.

[0014] The formal mechanism for formation of beer lightstruck flavour in model systems, composed of isohumulones, riboflavin, and cysteine, on exposure to visible light, has been suggested by Kuroiwa et al. already in 1963. Photo-excited riboflavin induces cleavage of isohumulones to a 4-methylpent-3-enyl radical, which undergoes decarbonylation to a 3-methylbut-2-enyl radical. Trapping of this stabilized allyl radical by a thiol radical derived from cysteine leads to formation of 3-methylbut-2-ene-1-thiol (3-MBT), the substance responsible for lightstruck flavour. 3-MBT has an extremely low flavour threshold level of around or below 1 ppt.

[0015] In order for the formation of 3-MBT to occur in beer besides iso-alpha acids, the reaction requires light energy in the 300-550 nm range of the spectrum, a photosensitizer (e.g. riboflavin, i.e. vitamin B2), and a sulphur source (e.g. sulphur containing amino acid).

[0016] The photolytic degradation of iso-alpha acids occurs as a consequence of the presence of an iso-3-hexenoyl side chain in the iso-alpha acid molecules. By modifying the molecular structure of the iso-alpha acids, notably by reducing the C=C and/or C=O bonds in the iso-3-hexenoyl side chain, substantial 3-MBT formation in beer can be prevented.

[0017] Reduced iso-alpha acid derivatives that are commercially available contain dihydro-, tetrahydro- and/or hexahydro-iso-alpha acids, and are usually added after the primary fermentation stage of the brewing process. The dihydro-iso-alpha acids (also called rho-iso-alpha acids) are obtained by the reduction of the carbonyl group in the aforementioned iso-3-hexenoyl chain to a hydroxyl group, generally using alkali metal borohydride as the reducing agent. The tetrahydro-iso-alpha acids are obtained via hydrogenation of the C=C bonds in the aforementioned iso-3-hexenoyl side chain and the isopentenyl side chain. The hexahydro-iso-alpha acids are produced by combining the aforementioned reduction and hydrogenation processes.


[0018] There is a need for a pre-isomerised hop extract that is light-stable, i.e. that does not substantially contribute to the formation of 3-MBT in light-exposed beer, but that has not been chemically reduced/hydrogenated.

SUMMARY OF THE INVENTION

[0019] The inventors have unexpectedly discovered that light-stability of pre-isomerised hop extracts can be improved substantially by illuminating such hop extracts with sunlight. Although the inventors do not wish to be bound by theory, it is believed that as a result of this illumination iso-alpha acids undergo the same type of photolytic degradation that is responsible for the formation of 3-MBT in beer. However, in the absence of important reactants (such as thiol-containing substances), the reactive intermediate degradation products formed during illumination that would otherwise participate in the formation of 3-MBT, are converted into other reaction products that do not adversely affect the flavour or flavour stability of beer.

[0020] The illumination of pre-isomerised hop extract in accordance with the present invention greatly improves the light-stability of the hop extract, but surprisingly it has not more than a limited impact on the bitterness and desirable flavour characteristics of the hop extract.


[0021] The inventors have further discovered that during illumination of pre-isomerised hop extract substantial quantities of the following hop derived substances (including all isomers) are formed:



wherein R = $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ or $-\text{CH}_2\text{CH}(\text{CH}_3)_2$.


[0022] These substances occur in conventional pre-isomerised hop extracts in not more than trace amounts. However, in the illuminated pre-isomerised hop extracts of the present invention they are present in significant concentration levels.

[0023] One aspect of the present invention relates to a light-stable hop extract comprising at least 1 g per kg of dry matter of one or more hop derived substances represented by the following formula (I):



wherein R = $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ or $-\text{CH}_2\text{CH}(\text{CH}_3)_2$.


[0024] The present invention also concerns a beer comprising light-stable hop extract, said beer containing at least 40 $\mu\text{g/l}$ riboflavin and at least 0.1 mg/l of one or more hop derived substances represented by the following formula (I):



wherein R = $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ or $-\text{CH}_2\text{CH}(\text{CH}_3)_2$.

DETAILED DESCRIPTION OF THE INVENTION

[0025] Accordingly, a first aspect of the present invention relates to a light-stable hop extract comprising at least 1 g per kg of dry matter of one or more hop derived substances represented by the following formula (I):



wherein R = $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ or $-\text{CH}_2\text{CH}(\text{CH}_3)_2$.

[0026] The term "hop derived substances" as used herein, unless indicated otherwise, refers to substituted oxycyclohexyl substances represented by aforementioned formula (I). The aforementioned hop derived substances can occur in different isomeric forms. All these isomeric forms, including tautomers and stereo-isomeric forms are encompassed by the above mentioned definition "hop derived substances".

[0027] The term "iso-alpha acids" as used herein refers to substances selected from the group of isohumulone, isoadhumulone, isocohumulone, pre-isohumulone, post-isohumulone and combinations thereof. The term "iso-alpha acids" encompasses different stereo-isomers (cis-iso-alpha acids and trans-iso-alpha acids).

[0028] The term "alpha acids" as used herein refers to substances selected from the group of

humulone, adhumulone, cohumulone, prehumulone, posthumulone and combinations thereof.

[0029] The term "beta acids" as used herein refers to substances selected from the group of lupulone, adlupulone, colupulone, prelupulone, and postlupulone and combinations thereof.

[0030] The term "isohumulone derived substance" as used herein refers to hop derived substances as defined above, wherein $R = -CH_2CH(CH_3)_2$.

[0031] The term "isocohumulone derived substance" as used herein refers to hop derived substances as defined above, wherein $R = -CH(CH_3)_2$.

[0032] The term "beer" as used herein encompasses both alcohol-containing and alcohol-free beer.

[0033] The concentration of the hop derived substances in a hop extract or in beer can suitably be determined by means of LC-MS, using the methodology that is described in this document.

[0034] The light-stable hop extract of the present invention typically contains the hop derived substances in a concentration of at least 3 g per kg of dry matter, more preferably in a concentration of at least 6 g per kg of dry matter and most preferably of at least 10 g per kg of dry matter. Typically, the hop extract contains not more than 350 g, preferably not more than 250 g of the hop derived substances per kg of dry matter.

[0035] According to a preferred embodiment of the present invention, the light-stable hop extract contains significant amounts of one or more hop derived substances according to formula (I) in which $R = -CH(CH_3)_2$. This hop derived substance is believed to originate from isocohumulone and is referred to herein as isocohumulone derived substance. Typically, the light-stable hop extract contains the isocohumulone derived substance in a concentration of at least 0.3 g per kg of dry matter, more preferably in a concentration of at least 1 g per kg of dry matter and most preferably of at least 3 g per kg of dry matter.

[0036] As explained below, the inventors have identified two isocohumulone derived substances in the light-stable hop extract of the present invention. These two isocohumulone derived substances, referred to as Co1 and Co2, are described in more detail below. Typically, the light-stable hop extract contains the isocohumulone derived substance Co2 in a concentration of at least 0.2 g per kg of dry matter, more preferably in a concentration of at least 0.7 g per kg of dry matter and most preferably of at least 2 g per kg of dry matter.

[0037] According to a further preferred embodiment of the present invention, the light-stable hop extract contains significant amounts of one or more hop derived substances according to formula (I) in which $R = -CH_2CH(CH_3)_2$. This hop derived substance is believed to originate from isohumulone and is referred to herein as isohumulone derived substance. Typically, the light-

stable hop extract contains the isohumulone derived substance in a concentration at least 0.5 g per kg of dry matter, more preferably in a concentration of at least 2 g per kg of dry matter and most preferably of at least 5 g per kg of dry matter.

[0038] The light-stable hop extract of the present invention typically contains at least 1%, more preferably at least 2% and most preferably at least 3% of iso-alpha acids by weight of dry matter.

[0039] According to a particularly preferred embodiment of the present invention the light-stable hop extract contains the hop derived substances in a concentration of at least 1%, more preferably at least 3%, even more preferably at least 5%, yet more preferably at least 10% and most preferably at least 20% by weight of the iso-alpha acids that are contained in the extract.

[0040] Typically, the light-stable hop extract contains at least 0.3%, more preferably at least 3% and most preferably at least 6% of the isocohumulone derived substance by weight of the iso-alpha acids that are contained in the extract.

[0041] The light-stable hop extract preferably contains at least 0.5%, more preferably at least 5% and most preferably at least 10% of the isohumulone derived substance by weight of the iso-alpha acids that are contained in the extract.

[0042] Alpha acids are typically contained in the light-stable hop extract in a concentration of 0-10%, more preferably of 0-3% and most preferably of 0.01-2% by weight of dry matter.

[0043] The light-stable hop extract of the present invention typically contains 0-10% beta acids by weight of dry matter, more preferably 0-5% beta acids by weight of dry matter.

[0044] The amount of reduced iso-alpha acids (dihydro-iso-alpha acids, tetrahydro-iso-alpha acids and hexahydro-iso-alpha acids) in the light-stable hop extract preferably does not exceed 10% by weight of the iso-alpha acids. Even more preferably the amount of reduced iso-alpha acids does not exceed 3% by weight of the iso-alpha acids.


[0045] The light-stable hop extract of the present invention typically comprises at least 20 mg/l of the hop derived substances. More preferably, the light-stable hop extract contains at least 80 mg/l of the hop derived substances, most preferably at least 150 mg/l of said hop derived substances.

[0046] The light-stable hop extract preferably is a hop extract that has been isolated from hop by means of extraction of with liquid or supercritical carbon dioxide.

[0047] The light-stable hop extract of the present invention preferably is a pre-isomerised hop extract, more preferably a pre-isomerised hop extract containing iso-alpha acids and alpha acids in a weight ratio that exceeds 10:1, most preferably in a weight ratio that exceeds 20:1.

[0048] The light-stable hop extract preferably has a water content of at least 30 wt.%, more preferably a water content of at least 50 wt.% and most preferably of at least 70 wt.%.

[0049] Another aspect of the present invention relates to a beer comprising light-stable hop extract, said beer containing at least 40 µg/l riboflavin and at least 0.1 mg/l of one or more hop derived substances represented by the following formula (I):



wherein R = -CH(CH₃)₂, -CH(CH₃)CH₂CH₃ or -CH₂CH(CH₃)₂.

[0050] As explained herein before, riboflavin plays an important role in the light-induced formation of 3-MBT in beer. Riboflavin is naturally present in beer and undergoes extremely rapid photolysis. As a matter of fact, in ordinary beer riboflavin is photolysed much more rapidly than iso-alpha acids. Thus, in light-struck beer riboflavin content is usually reduced to only a fraction of the original content even before a significant fraction of the iso-alpha acids has been photolysed.

[0051] The light-stable beer of the present invention differs from ordinary beer in that it contains at least 0.1 mg/l of the aforementioned one or more hop derived substances. This light-stable beer differs from light-struck beer in that it simultaneously contains at least 0.1 mg/l of these hop derived substances and at least 40 µg/l riboflavin.

[0052] The beer of the present invention typically contains at least 0.2 mg/l, more preferably at least 0.4 mg/l, and even more preferably at least 1 mg/l of the one or more hop derived substances. Typically the concentration of the one or more photolysed hop substances in the beer does not exceed 50 mg/l. More preferably said concentration does not exceed 40 mg/l, even more preferably it does not exceed 30 mg/l and most preferably it does not exceed 20 mg/l.

[0053] The light-stable beer preferably contains at least 0.03 mg/l, more preferably at least 0.1 mg/l and most preferably at least 0.3 mg/l of the isocohumulone derived substance.

[0054] The isocohumulone derived substance Co2 is preferably contained in the light-stable beer in a concentration of at least 0.02 mg/l, more preferably at least 0.07 mg/l and most preferably of at least 0.2 mg/l.

[0055] The light-stable beer preferably contains at least 0.05 mg/l, more preferably at least 0.2 mg/l and most preferably at least 0.5 mg/l of the isohumulone derived substance.

[0056] The riboflavin content of the light-stable beer preferably is at least 60 µg/l. Most preferably, the riboflavin content is in the range of 80 to 2,000 µg/l.

[0057] According to a particularly preferred embodiment, the one or more hop derived substances and riboflavin are present in the light-stable beer in a weight ratio that does not exceed 200: 1, more preferably in a weight ratio that is in the range of 1:1 to 100:1 and most preferably in the range of 2:1 to 50:1.

[0058] The light-stable beer typically contains the isocohumulone derived substance and riboflavin in a weight ratio that does not exceed 100:1, more preferably in a weight ratio of 1:2 to 50:1, most preferably of 1:1 to 40:1.

[0059] The light-stable beer typically contains the isohumulone derived substance and riboflavin in a weight ratio that does not exceed 160:1, more preferably in a weight ratio of 1:1 to 80:1, most preferably of 2:1 to 60:1.

[0060] The light-stable beer of the present invention typically contains at least 0.1 mg/l, more preferably at least 0.3 mg/l, even more preferably at least 0.6 mg/l and most preferably at least 1 mg/l of iso-alpha acids. Typically the level of iso-alpha acids in the beer will not exceed 30 mg/l.

[0061] The one or more hop derived substances are preferably contained in the light-stable beer in a concentration of at least 1%, more preferably of at least 3%, even more preferably at least 5% and most preferably of at least 10% by weight of the iso-alpha acids that are also contained in the beer.

[0062] The light-stable beer of the present invention typically contains less than 100 ng/l 3-methyl-2-butene-1-thiol (3-MBT). Even more preferably the 3-MBT content is less than 50 ng/l, most preferably less than 30 ng/l. The concentration of 3-MBT can suitably be determined by means of the method described by Hughes et al. (Hughes P. S., Burke S. and Meacham A. E. (1997) "Aspects of the lightstruck character of beer". Institute of Brewing, Proceedings of the 6th Central and South Africa Section, pp. 123-128).

[0063] A further aspect of the present invention relates to a process of preparing a beer as defined herein before, said process comprising introducing a light-stable hop extract as defined herein before.

[0064] The light-stable hop extract is preferably added to wort or beer before clarification, i.e. before removal of entrained solids and yeast from the fermented beer. Typically, clarification of fermented beer is achieved by means of filtration.

[0065] In the aforementioned process the light-stable hop extract is preferably introduced into the wort or beer in a concentration of at least 2 mg/l, more preferably in a concentration of 5-

200 mg/l, most preferably in a concentration of 10-150 mg/l.

[0066] Following addition of the light-stable hop extract to the wort, the wort is subjected to further processing steps, including:

- fermenting the wort with the help of brewer's yeast;
- clarifying the fermented beer; and
- packaging.

[0067] According to a particularly preferred embodiment, the pre-isomerised hop extract is illuminated with light having a maximum intensity at a wavelength in the range of 200-800 nm, more preferably in the range of 250-600 nm, most preferably of 300-500 nm.

[0068] The illumination intensity employed in the present process preferably exceeds 50 W/m². More preferably, said illumination intensity exceeds 100 W/m², most preferably it exceeds 200 W/m².

[0069] The pre-isomerised hop extract is typically illuminated with the aforementioned illumination intensity for at least 30 minutes, more preferably for at least 1 hour and most preferably for 2-48 hours.

[0070] During illumination the temperature of the hop extract typically remains within the range of 0-100°C, more preferably of 15-80°C.

[0071] The inventors have found that particularly good results can be obtained if the pre-isomerised hop extract is exposed to illumination after being diluted with water. Accordingly, in accordance with a particularly preferred embodiment, the present method comprises the steps of:

- diluting the pre-isomerised extract with water to produce an aqueous solution; and
- illuminating an aqueous solution of the pre-isomerised hop extract.

[0072] The aqueous diluted solution of the pre-isomerised hop extract typically has an iso-alpha acid content in the range of 0.2-200 g/l, more preferably of 0.5-70 g/l and most preferably 1-25 g/l when the illuminations starts.

[0073] The aqueous diluted solution of the pre-isomerised hop extract typically contains at least 80 wt.% water. Even more preferably said dilution contains at least 90 wt.% water. Most preferably, at least 95 wt.% water.

[0074] The invention is further illustrated by the following non-limiting examples.

EXAMPLES***Method of determining the concentration of hop derived substances***

[0075] The concentrations of 4 different hop derived substances represented by formula (I) can be determined by means of UPLC-MS using the procedure described below. These 4 hop derived substances include two isocohumulone derived substances (referred to as Co1 and Co2) and two isohumulone derived substance (referred to as N1 and N2). Typical retention times of these substances in the UPLC-MS method described below are listed below.

Substance	Retention time (min.)
Co1	6.87
Co2	7.99
N1	9.08
N2	9.99

[0076] The mass spectra of these hop derived substances are shown in Figures 1-4.

[0077] The concentration of each hop derived substance is determined on the basis of calibration curves for each of the hop derived substances. These hop derived substances can be isolated in highly pure form from illuminated pre-isomerised hop extract by means of preparative UPLC.

[0078] The calibration curves for the hop derived substances are determined using the detector conditions described below.

[0079] The equipment, materials and conditions described below can be used to determine the concentrations of the aforementioned hop derived substances.

Equipment:

[0080]

UPLC system:	Acquity UPLC (Waters) equipped with a BEH C18 column (1.7 µm, 2.1 mm × 150 mm; Waters product number: 186002353)
Detector:	XEVO TQ-S mass spectrometer (Waters)

Chemicals:

[0081]

Milli-Q® Ultrapure Water

Acetonitrile ≥ 99.9% (JT-Baker: 9853-02)

Formic acid ≥ 98% (Sigma-Aldrich product number: 33015)

UPLC Instrument conditions:**[0082]**

Column temperature: 60 °C

Injection volume: 1 µL

Sample temperature: 10 °C

Run time: 25 min

Eluent A = 1000 mL Ultrapure water + 1 mL formic acid

Eluent B = 1000 mL acetonitrile + 1 mL formic acid

Gradient:**[0083]**

Time (min)	Flow (mL/min)	%A	%B
0.0	0.25	95	5
10	0.25	59	41
11	0.25	95	5
25	0.25	95	5

Detector conditions:

[0084] Analytical method is set-up in negative scan mode (ES-) and Multiple Reaction Monitoring (MRM) modes.

MS settings:

[0085]

Polarity	ES-
Capillary (kV)	2.5
Source Temperature (°C)	150
Desolvation temperature (°C)	600
Cone gas flow (L/Hr)	150
Desolvation gas flow (L/Hr)	1000
Collision gas flow (mL/min)	0.22

Multiple Reaction Monitoring (MRM) modes:

[0086]

	Parent (m/z)	Daughter (m/z)	Coll (eV)
Co1 and Co2	267	195	20
N1 and N2	281	209	20

[0087] Samples are degassed prior to analysis.

Method of determining the concentration of riboflavin

[0088] The concentration of riboflavin in beer can be determined by means of UPLC-FLR using equipment, materials and conditions as described below.

Equipment:

[0089]

UPLC system:	Acquity UPLC (Waters) equipped with a BEH C18 column (1.7 µm, 2.1 mm × 150 mm; Waters product number: 186002353)
--------------	--

Detector:	Acquity Fluorescence (FLR) detector (Waters)
-----------	--

Chemicals:

[0090]

Milli-Q® Ultrapure Water

Acetonitrile $\geq 99.9\%$ (Sigma Aldrich product number: 34998)

Phosphoric acid $\geq 85.0\%$ (Sigma Aldrich product number: 30417)

Riboflavin (Sigma Aldrich product number: R-7649)

Sodium dihydrogen phosphate monohydrate (Merck product number: 106346)

pH buffer 2.7 (7,5 g sodium dihydrogen phosphate monohydrate in 200 mL Ultrapure water + 1 mL Phosphoric acid in 1000 mL Ultrapure water)

UPLC instrument conditions:

[0091]

Column temperature: 40 °C

Injection volume: 10 μ L

Sample temperature: 10 °C

Run time: 15 min

Eluens A = pH buffer 2.7

Eluens B = acetonitrile

Gradient:

[0092]

Time (min)	Flow (mL/min)	%A	%B
0.00	0.20	95	5
10.0	0.20	95	5
10.5	0.20	15	85
12.0	0.20	15	85
12.2	0.20	95	5
15.0	0.20	95	5

Detector conditions:**[0093]** $\lambda_{\text{ex}} = 444 \text{ nm}$ $\lambda_{\text{em}} = 525 \text{ nm}$ **[0094]** Samples are degassed prior to analysis.**Example 1**

[0095] A pre-isomerised hop extract (so-called 'Isohop' obtained from Barth-Haas Group) was used as a starting material. This hop extract had the following characteristics (according to the product description sheet from Barth-Haas Group):

- contains $30 \pm 0.5 \text{ wt\%}$ iso-alpha acids
- contains $< 0.7\%$ alpha acids
- contains $< 0.3\%$ beta acids

[0096] The pre-isomerised hop extract was diluted with 'Milli-Q® Ultrapure Water' (1:50 w/w). 16 mL of the diluted hop extract was introduced into a glass tube with screw cap (Duran glass, tube diameter is 16 mm, tube length is 160 mm). 30 of these glass tubes were simultaneously illuminated for a period of 32 hours inside a Suntest XLS+ device (ATLAS), using a Xenon lamp. During illumination, the following conditions were applicable:

- Power of Xenon lamp is 1700 W
- Tubes were horizontally positioned inside the Suntest XLS+ device

- Distance between glass tubes and lamp is 25 cm
- Amount of irradiance at sample level was set at 765 W/m²
- Temperature of diluted hop extract increases from ambient to circa 50°C (no cooling nor heating was intentionally applied during illumination)

[0097] Before and after illumination, the concentrations of iso-alpha acids and hop derived substances in the hop extract were determined. Determination of iso-alpha acids in the hop extract was performed according to the method described by the European Brewery Convention (EBC) in 2005: 'Method 7.8: Iso-alpha-, alpha- and beta-acids in hops, hop powder products and hop extracts (HPLC method)'. Determination of the concentration of the hop derived substances in the extract was performed according to the method described above. The results of these analyses are shown in Table 2:

Table 2

	Iso-alpha acids (mg/L)	Hop derived substances				
		Co1 (mg/L)	Co2 (mg/L)	N1 (mg/L)	N2 (mg/L)	Sum (mg/L)
Hop Extract	5808	<0.01	<0.01	<0.01	<0.01	<0.01
Light-Stable Hop Extract	339	65	264	174	372	875

Example 2

[0098] The illuminated and non-illuminated hop extracts of Example 1 were added to unhopped beer (300 mL unhopped beer in green bottle). A first beer (reference beer) was prepared by adding 20 mg/L of the non-illuminated hop extract. A second beer (light-stable beer) was prepared by adding 60 mg/L of the illuminated hop extract. The illuminated hop extract was applied in a higher concentration in order to compensate for the loss in bitter intensity that had occurred as a result of the illumination. The flavour profile of the beer containing the illuminated hop extract had a flavour profile that was very similar to that of the beer containing the non-illuminated hop extract.

[0099] Both beers were exposed to artificial sunlight by putting the bottles horizontally in the Suntest XLS+ device and illuminating them for a period of 24 hours. During illumination, the same conditions were applied as in Example 1.

[0100] Before and during exposure to the artificial sunlight the concentrations of the following substances were determined in both beers:

- the hop derived substances of formula (I),

- iso-alpha acids,
- riboflavin, and
- 3-MBT.

[0101] Determination of iso-alpha acids in beer was performed according to the method described by the American Society of Brewing Chemists (ASBC) in 2009 (Methods of Analysis, 14th Edition): 'Method Beer 23 (section E: Iso-alpha acids in beer by HPLC)'. The 3-MBT content was determined using the method described by Hughes et al. ('Aspects of the lightstruck character of beer'. Institute of Brewing, Proceedings of the 6th Central and South Africa Section (1997), pp. 123-128). Determination of the concentration of the hop derived substances and riboflavin in beer was performed according to the methods described above.

[0102] The results of these analyses are summarised in Table 3:

Table 3

	Iso-alpha acids (mg/L)	Hop derived substances					Riboflavin (µg/L)	MBT (ng/L)
		Co1 (mg/L)	Co2 (mg/L)	N1 (mg/L)	N2 (mg/L)	Sum (mg/L)		
Reference beer								
0h illumination	19.2	<0.01	<0.01	<0.01	<0.01	<0.01	318	7
2h illumination	18.8	0.01	0.02	0.03	0.04	0.11	16	309
6h illumination	17.8	0.05	0.08	0.11	0.15	0.39	2.8	660
24h illumination	16.0	0.19	0.28	0.42	0.54	1.42	0.3	2072
Light-stable beer								
0h illumination	4.5	0.53	2.32	1.65	3.54	8.04	313	8
2h illumination	4.4	0.56	2.29	1.66	3.45	7.95	15	74
6h illumination	4.2	0.61	2.20	1.77	3.29	7.87	2.0	137
24h illumination	3.6	0.86	2.19	2.18	3.18	8.41	0.4	506

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- [AU7540281](#) [0006]
- [US4767640A](#) [0007]
- [WO9302177A](#) [0008]
- [US5811144A](#) [0009]
- [WO2006104387A](#) [0010]
- [WO2011054838A](#) [0011]
- [GB2362891A](#) [0012]

Non-patent literature cited in the description

- **HUGHES P. S.BURKE S.MEACHAM A. E.**Aspects of the lightstruck character of beerProceedings of the 6th Central and South Africa Section, 1997, 123-128 [0062]
- **Method Beer 23** (section E: Iso-alpha acids in beer by HPLC)Methods of Analysis20090000 [0101]
- **HUGHES et al.**Aspects of the lightstruck character of beerProceedings of the 6th Central and South Africa Section, 1997, 123-128 [0101]

Patentkrav

1. Øl omfattende lysstabil humleekstrakt, hvilken øl indeholder mindst 40 µg/l riboflavin og mindst 0,1 mg/l af et eller flere humleafledte stoffer repræsenteret ved formel (I):



hvor R = -CH(CH₃)₂, -CH(CH₃)CH₂CH₃ eller -CH₂CH(CH₃)₂.

2. Øl ifølge krav 1, hvor øllet indeholder mindst 0,2 mg/l, fortrinsvis mindst 0,4 mg/l, mere fortrinsvis mindst 1 mg/l af de et eller flere humleafledte stoffer.

3. Øl ifølge krav 1 eller 2, hvor øllet ikke indeholder mere end 50 mg/l, fortrinsvis ikke mere end 40 mg/l, mere fortrinsvis ikke mere end 30 mg/l af de et eller flere humleafledte stoffer.

4. Øl ifølge et hvilket som helst af de foregående krav, hvor øllet indeholder mindst 60 µg/l, fortrinsvis 80-2.000 µg/l riboflavin.


5. Øl ifølge et hvilket som helst af de foregående krav, hvor de en eller flere humleafledte stoffer og riboflavin er til stede i et vægtforhold, der ikke overstiger 200:1, fortrinsvis i et vægtforhold, der er i området fra 1:1 til 100:1.

6. Øl ifølge et hvilket som helst af de foregående krav, hvor øllet indeholder mindst 0,1 mg/l, fortrinsvis mindst 0,3 mg/l, mere fortrinsvis mindst 0,6 mg/l

iso-alfa-syrer.

5 **7.** Øl ifølge krav 6, hvor øllet indeholder mindst 1%, mere foretrukket mindst 3% og mest foretrukket mindst 5 vægt% af iso-alfa-syrerne af de humleafledte stoffer.

8. Lysstabil humleekstrakt, der omfatter mindst 1 g pr. kg tørstof af et eller flere humleafledte stoffer repræsenteret ved følgende formel (I):



10 hvor R = $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ eller $-\text{CH}_2\text{CH}(\text{CH}_3)_2$.

15 **9.** Humleekstrakt ifølge krav 8, hvor humleekstrakten indeholder det ene eller flere humleafledte stoffer i en koncentration på mindst 3 g pr. kg tørstof, fortrinsvis i en koncentration på mindst 6 g pr. kg tørstof og mere fortrinsvis på mindst 10 g pr. kg tørstof.

10. Humleekstrakt ifølge krav 8 eller 9, hvor humleekstrakten indeholder mindst 1%, mere foretrukket mindst 3% og mest fortrinsvis mindst 5 vægt% af iso-alfasyrerne af den ene eller flere humleafledte stoffer.

20

11. Humleekstrakt ifølge et hvilket som helst af kravene 8-10, hvor humleekstrakten indeholder mindst 1 vægt% iso-alfasyrer af tørstof.

- 12.** Humleekstrakt ifølge et hvilket som helst af kravene 8-11, hvor humleeks-trakten indeholder alfa-syrer i en koncentration på 0-10 vægt-% tørstof, fortrinsvis 0-3 vægt-% tørstof og mere fortrinsvis 0,01-2 vægt-% tørstof.
- 5 **13.** Humleekstrakt ifølge et hvilket som helst af kravene 8-12, hvor humleeks-trakten indeholder beta-syrer i en koncentration på 0-10 vægt% tørstof, fortrinsvis 0-5 vægt% tørstof.
- 10 **14.** Humleekstrakt ifølge et hvilket som helst af kravene 8-13, hvor humleeks-trakten indeholder reducerede iso-alfa-syrer i en koncentration, der ikke overstiger 10 vægt-% af iso-alfa-syrerne, fortrinsvis ikke overstiger 3 vægt-% af iso-alfa-syrer; de reducerede iso-alfa-syrer er valgt blandt dihydro-iso-alfa-syrer, tetrahydro-iso-alfa-syrer, hexahydro-iso-alfa-syrer og kombinationer deraf.
- 15 **15.** Fremgangsmåde til fremstilling af en øl, hvilken fremgangsmåde omfatter at indføre en humleekstrakt ifølge et hvilket som helst af kravene 8-14.

DRAWINGS


Fig. 1

Fig. 2




Fig. 3




Fig. 4

