



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2535652 A1 2005/02/24

(21) **2 535 652**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2004/08/12
(87) Date publication PCT/PCT Publication Date: 2005/02/24
(85) Entrée phase nationale/National Entry: 2006/02/13
(86) N° demande PCT/PCT Application No.: GB 2004/003473
(87) N° publication PCT/PCT Publication No.: 2005/016909
(30) Priorités/Priorities: 2003/08/15 (US60/495,580);
2004/05/28 (US60/576,416)

(51) Cl.Int./Int.Cl. *C07D 333/38* (2006.01),
C07D 409/12 (2006.01), *A61K 31/38* (2006.01),
C07D 333/00 (2006.01), *C07D 307/00* (2006.01),
C07D 241/00 (2006.01), *A61P 35/00* (2006.01)

(71) Demandeur/Applicant:
ASTRAZENECA AB, SE

(72) Inventeurs/Inventors:
ASHWELL, SUSAN, US;
GERO, THOMAS, US;
IOANNIDIS, STOPHANOS, US;
JANETKA, JAMES, US;
LYNE, PAUL, US;
...

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : THIOPHENES SUBSTITUES ET LEURS UTILISATIONS
(54) Title: SUBSTITUTED THIOPHENES AND USES THEREOF

(57) **Abrégé/Abstract:**

This invention relates to novel compounds having the structural formula (I) and to their pharmaceutical salts, compositions and methods of use. These novel compounds provide a treatment or prophylaxis of cancer.



(72) Inventeurs(suite)/Inventors(continued): OZA, VIBHA, US; SPRINGER, STEPHANIE, US; SU, MEI, US; YU, DINGWEI, US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
24 February 2005 (24.02.2005)

PCT

(10) International Publication Number
WO 2005/016909 A1

(51) International Patent Classification⁷: **C07D 333/38**,
409/12, A61K 31/38, A61P 35/00 // (C07D 409/12,
333:00, 333:00) (C07D 409/12, 333:00, 307:00) (C07D
409/12, 333:00, 241:00)

(21) International Application Number:
PCT/GB2004/003473

(22) International Filing Date: 12 August 2004 (12.08.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/495,580 15 August 2003 (15.08.2003) US
60/576,416 28 May 2004 (28.05.2004) US

(71) Applicant (for all designated States except MG, US): **ASTRAZENECA AB** [SE/SE]; SE-151 85 Södertälje (SE).

(71) Applicant (for MG only): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ASHWELL, Susan** [GB/US]; AstraZeneca R & D Boston, 35 Gatehouse Drive, Waltham, MA 02451 (US). **GERO, Thomas** [US/US]; AstraZeneca R & D Boston, 35 Gatehouse Drive, Waltham, MA 02451 (US). **IOANNIDIS, Stophanos** [GR/US]; AstraZeneca R & D Boston, 35 Gatehouse Drive, Waltham, MA 02451 (US). **JANETKA, James** [US/US]; AstraZeneca R & D Boston, 35 Gatehouse Drive, Waltham, MA 02451 (US). **LYNE, Paul** [IE/US]; AstraZeneca R & D Boston, 35 Gatehouse Drive, Waltham, MA 02451 (US).

OZA, Vibha [IN/US]; AstraZeneca R & D Boston, 35 Gatehouse Drive, Waltham, MA 02451 (US). **SPRINGER, Stephanie** [US/US]; 37 Village Green Drive, North Andover, MA 01845 (US). **SU, Mei** [CN/US]; AstraZeneca R & D Boston, 35 Gatehouse Drive, Waltham, MA 02451 (US). **YU, Dingwei** [US/US]; AstraZeneca R & D Boston, 35 Gatehouse Drive, Waltham, MA 02451 (US).

(74) Agent: **GLOBAL INTELLECTUAL PROPERTY**; AstraZeneca AB, SE-151 85 Södertälje (SE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED THIOPHENES AND USES THEREOF

(57) Abstract: This invention relates to novel compounds having the structural formula (I) and to their pharmaceutical salts, compositions and methods of use. These novel compounds provide a treatment or prophylaxis of cancer.

WO 2005/016909 A1

- 1 -

SUBSTITUTED THIOPHENES AND USES THEREOF**Field of the invention**

5 The present invention relates to novel substituted thiophenes, their pharmaceutical compositions and methods of use. In addition, the present invention relates to therapeutic methods for the treatment and prevention of cancers.

Background of the invention

10

Chemotherapy and radiation exposure are currently the major options for the treatment of cancer, but the utility of both these approaches is severely limited by adverse effects on normal tissue, and the frequent development of tumor cell resistance. It is therefore desirable to improve the efficacy of such treatments in a way that does not increase the toxicity associated with them.

15 One way to achieve this is by the use of specific sensitizing agents such as those described herein.

 An individual cell replicates by making an exact copy of its chromosomes, and then segregating these into separate cells. This cycle of DNA replication, chromosome separation and division is regulated by mechanisms within the cell that maintain the order of the steps and ensure that each step is precisely carried out. Involved in these processes are the cell cycle checkpoints (Hartwell *et al.*, *Science*, Nov 3, 1989, 246(4930):629-34) where cells may arrest to ensure DNA repair mechanisms have time to operate prior to continuing through the cycle into mitosis. There are two such checkpoints in the cell cycle – the G1/S checkpoint that is regulated by p53 and the G2/M checkpoint that is monitored by the Ser/Thr kinase checkpoint kinase 1
20 (CHK1).
25

 The cell cycle arrest induced by these checkpoints is a mechanism by which cells can overcome the damage resulting from radio- or chemotherapy, their abrogation by novel agents should increase the sensitivity of tumor cells to DNA damaging therapies. Additionally, the tumor specific abrogation of the G1/S checkpoint by p53 mutations in the majority of tumors can
30 be exploited to provide tumor selective agents. One approach to the design of chemosensitizers


- 2 -

that abrogate the G2/M checkpoint is to develop inhibitors of the key G2/M regulatory kinase CHK1, and this approach has been shown to work in a number of proof of concept studies. (Koniaras *et al.*, *Oncogene*, 2001, 20:7453; Luo *et al.*, *Neoplasia*, 2001, 3:411; Busby *et al.*, *Cancer Res.*, 2000, 60:2108; Jackson *et al.*, *Cancer Res.*, 2000, 60:566).

5

Summary of the invention

Provided herein are novel compounds of structural formula (I) or a pharmaceutically acceptable salt thereof:



10

wherein:

15 R^1 and R^2 are at each occurrence independently selected from H, optionally substituted C_{1-6} alkyl, or optionally substituted heterocyclyl; with the proviso that R^1 and R^2 are not both H; or R^1 and R^2 and the N to which they are attached in combination form an optionally substituted heterocyclyl;

R^4 is selected from H, OH, optionally substituted carbocyclyl, optionally substituted heterocyclyl, or optionally substituted C_{1-6} alkyl;

20 R^5 is selected from H, optionally substituted carbocyclyl, or optionally substituted C_{1-6} alkyl.

The invention also encompasses stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts of compounds of formula I, pharmaceutical compositions and formulations containing them, methods of using them to treat diseases and

25


- 3 -

conditions either alone or in combination with other therapeutically-active compounds or substances, processes and intermediates used to prepare them, uses of them as medicaments, uses of them in the manufacture of medicaments and uses of them for diagnostic and analytic purposes.

5

Detailed Description of the Invention

Provided herein are novel compounds of structural formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor thereof:



wherein:

15 R^1 and R^2 are at each occurrence independently selected from H, optionally substituted C_{1-6} alkyl, or optionally substituted heterocyclyl; with the proviso that R^1 and R^2 are not both H; or R^1 and R^2 and the N to which they are attached in combination form an optionally substituted heterocyclyl;

R^4 is selected from H, OH, optionally substituted carbocyclyl, optionally substituted heterocyclyl, or optionally substituted C_{1-6} alkyl;

20 R^5 is selected from H, optionally substituted carbocyclyl, or optionally substituted C_{1-6} alkyl.

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R^1 is an optionally substituted heterocyclyl.

25

- 4 -

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R¹ is an optionally substituted heterocyclyl wherein 1,2, or 3 substituents is/are independently selected from

5 halogen, nitro, amino, cyano, trifluoromethyl, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, hydroxy, alkylhydroxy, carbonyl, -CH(OH)CH₃, -CH₂NH-alkyl-OH, alkyl-(OH)CH₃, -CH₂-phenyl-(OCH₃)₂, -Oalkyl, -OCH₃, -Ophenyl, -OCOalkyl, -NHCHO, -Nalkyl, -N-(alkyl)-CHO, -NH-CO-amino, -N-(alkyl)-CO-amino, -NH-COalkyl, -N-(alkyl)-COalkyl, -carboxy, -amidino, -CO-amino, -CO-alkyl, -CO₂alkyl, mercapto, -Salkyl, -SCH₂furanyl, -SO(alkyl), -SO₂(alkyl), -SO₂-

10 amino, -alkylsulfonylamino, phenyl, anisole, dimethoxyphenyl, trimethoxyphenyl, halophenyl, cycloalkyl, heterocyclyl, -alkyl-NH-cycloalkyl, -alkyl-NH- heterocyclyl, -alkyl-NH-alkyl-OH, -C(=O)OC(CH₃)₃, -N(CH₃)₂, -N(CH₂CH₃)₂, -alkyl-NH-alkyl- heterocyclyl, -alkyl-aryl, -methyl-phenyl, alkyl-polycyclyl, alkyl-amino, alkyl-hydroxy, -CH₂NH-alkyl-heterocyclyl, -CH₂NHCH₂CH(CH₃)₂, vicinal -O(alkyl)O-, vicinal -OC(haloalkyl)O-, vicinal -CH₂O(alkyl)O-,

15 vicinal -S(alkyl)S- and -O(alkyl)S-.

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R¹ is an optionally substituted heterocyclyl wherein 1,2, or 3 substituents is/are independently selected from: -OH,

20 C(=O)OC(CH₃)₃, NH₂, C₁₋₆alkyl, methoxybenzene, or dimethoxy benzene.

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R¹ is a heterocyclyl wherein heterocyclyl is selected from piperdiny, pyridiny, pyrrolidiny, pyraziny,

25 azepany, azetidiny, azabicycloziny, furany, thieny.

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R² is H.

- 5 -

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R⁴ is H.

One embodiment of the present invention provides compounds of formula (I) or a
5 pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R⁵ is H or an optionally substituted C₁₋₆alkyl.

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R⁵ is H or an
10 optionally substituted C₁₋₆alkyl wherein 1,2 or 3 substituents is/are independently selected from: NH₂, NHCH₃, N(CH₂CH₃)₂, N(CH₃)₂, OCH₃, OH, -C₁₋₆alkyl, morpholino, piperidinyl, pyrrolidinyl.

One embodiment of the present invention provides compounds of formula (I) or a
15 pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R⁵ is H or an optionally substituted C₁₋₃alkyl.

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein R⁵ is H or an
20 optionally substituted C₁₋₃alkyl wherein 1,2 or 3 substituents is/are independently selected from: NH₂, NHCH₃, N(CH₂CH₃)₂, N(CH₃)₂, OCH₃, OH, -C₁₋₆alkyl, morpholino, piperidinyl, pyrrolidinyl.

One embodiment of the present invention provides compounds of formula (I) or a
25 pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein

R¹ is an optionally substituted heterocyclyl;

R² is H;

R⁴ is H;

R⁵ is H or an optionally substituted C₁₋₆alkyl.

- 6 -

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein:

R^1 is an optionally substituted heterocyclyl wherein the substituent is selected from one or more of the following: $-NH_2$, C_{1-6} alkyl, $-C(=O)OC(CH_3)_3$,

5 R^2 is H;

R^4 is H;

R^5 is H or an optionally substituted C_{1-6} alkyl wherein the substituent is selected from one or more of the following: $-C_{1-6}$ alkyl, $-N(C_{1-3}alkyl)_2$.

10 One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein:

R^1 is an optionally substituted heterocyclyl wherein the substituent is selected from one or more of the following: $-NH_2$, C_{1-6} alkyl, $-C(=O)OC(CH_3)_3$,

R^2 is H;

15 R^4 is H;

R^5 is H or an optionally substituted C_{1-3} alkyl wherein 1,2 or 3 substituents is/are independently selected from: NH_2 , $NHCH_3$, $N(CH_2CH_3)_2$, $N(CH_3)_2$, OCH_3 , OH , $-C_{1-6}$ alkyl, morpholino, piperidinyl, pyrrolidinyl.

20 One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein:

R^1 is a heterocyclyl;

R^2 is H;

R^4 is H;

25 R^5 is H or a C_{1-6} alkyl.

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt or an *in vivo*-hydrolysable precursor wherein:

R^1 is a 6-membered heterocyclyl containing at least one N in the ring;

30 R^2 is H;

- 7 -

R⁴ is H;

R⁵ is a C₁₋₃alkyl.

One embodiment of the present invention provides compounds of formula (I) selected from the following:

tert-butyl 3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-ylthiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-ylthiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide;

tert-butyl 3-{[(2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-ylthiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-N-[(3R)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;

N-(3-[(4-aminopiperidin-1-yl)carbonyl]-5-{4-[2-(diethylamino)ethoxy]phenyl}-2-thienyl)urea;

2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[3-(hydroxymethyl)phenyl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-ylthiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-N-(2-aminoethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-pyridin-3-ylthiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(1-methylpiperidin-4-yl)thiophene-3-carboxamide;

- 8 -

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-1-methylazepan-3-yl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-[3-(hydroxymethyl)phenyl]thiophene-3-carboxamide;

5 2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-pyrrolidin-3-ylthiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-pyridin-3-ylthiophene-3-carboxamide;

10 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-1-methylpiperidin-3-yl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-pyrrolidin-3-ylthiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-ylmethyl]thiophene-3-carboxamide;

15 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]thiophene-3-carboxamide;

20 2-[(aminocarbonyl)amino]-N-[2-(dimethylamino)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-N-[2-(diethylamino)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;

25 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-yl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(piperidin-4-ylmethyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-pyrrolidin-3-ylthiophene-3-carboxamide;

30 2-[(aminocarbonyl)amino]-N-(1-ethylpiperidin-3-yl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

- 9 -

- 2-[(aminocarbonyl)amino]-N-[(3S)-1-ethylazepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(3-hydroxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;
- 5 2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;
- tert-butyl (3S)-3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)pyrrolidine-1-carboxylate;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-piperidin-3-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-(1-benzylpiperidin-4-yl)-5-(4-methoxyphenyl)thiophene-3-
- 10 carboxamide;
- tert-butyl 3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate;
- 2-[(aminocarbonyl)amino]-5-[4-(2-piperidin-1-ylethoxy)phenyl]-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide;
- 15 2-[(aminocarbonyl)amino]-5-[4-(2-piperidin-1-ylethoxy)phenyl]-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-azetidin-3-yl-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(2S)-pyrrolidin-2-ylmethyl]thiophene-3-carboxamide;
- 20 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-pyridin-4-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperazin-1-ylethyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperidin-1-ylethyl)thiophene-3-carboxamide;
- 25 2-[(aminocarbonyl)amino]-N-1-azabicyclo[2.2.2]oct-3-yl-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-(2-hydroxyethyl)-5-(4-hydroxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-(trans-4-hydroxycyclohexyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

- 10 -

2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperazin-1-ylethyl)thiophene-3-carboxamide;

5 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-pyridin-3-ylethyl)thiophene-3-carboxamide;

10 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-3-ylethyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(2-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;

15 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(tetrahydrofuran-2-ylmethyl)thiophene-3-carboxamide;

tert-butyl (3R)-3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-3-ylmethyl)thiophene-3-carboxamide; -

20 tert-butyl 3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)azetidine-1-carboxylate;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-4-ylmethyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(3-methoxypropyl)thiophene-3-carboxamide;

25 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[2-(2-thienyl)ethyl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-thienylmethyl)thiophene-3-carboxamide;

N-[3-(1,4-diazepan-1-ylcarbonyl)-5-(4-methoxyphenyl)-2-thienyl]urea;

2-[(aminocarbonyl)amino]-N-(2-methoxyethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

30 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-thienylmethyl)thiophene-3-carboxamide;

- 11 -

2-[(aminocarbonyl)amino]-N-{2-[(2-furylmethyl)thio]ethyl}-5-(4-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-[2-(2-thienyl)ethyl]thiophene-3-carboxamide;
N-(3-[(4-aminopiperidin-1-yl)carbonyl]-5-{3-[2-(diethylamino)ethoxy]phenyl}-2-thienyl)urea;

5 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-ylmethyl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(1,2,3,4-tetrahydroquinolin-3-yl)thiophene-3-carboxamide;

10 2-[(aminocarbonyl)amino]-N-(1,3-benzodioxol-5-ylmethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-N-(3-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-N-[2-(3,4-dimethoxyphenyl)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;

15 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(5-methyl-2-furyl)methyl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-2-ylmethyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-N-(4-fluorobenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

20 tert-butyl 4-({[2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate;

2-[(aminocarbonyl)amino]-N-(2-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-phenoxyethyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-2-ylethyl)thiophene-3-carboxamide;

25 tert-butyl 4-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate;

2-[(aminocarbonyl)amino]-N-(4-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide;

- 12 -

2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3R)-piperidin-3-yl]thiophene-3-carboxamide;

tert-butyl (3S)-3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

5 2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-{4-[2-(diethylamino)ethoxy]phenyl}thiophene-3-carboxamide;

tert-butyl (3R)-3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

N-[3-{[(3S)-3-aminoazepan-1-yl]carbonyl}-5-(4-methoxyphenyl)-2-thienyl]urea;

10 5-{4-[2-(diethylamino)ethoxy]phenyl}-2-{[(pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide;

5-{3-[2-(diethylamino)ethoxy]phenyl}-2-{[(pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide;

15 5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

20 N-(2-aminoethyl)-5-(4-methoxyphenyl)-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

25 5-(4-methoxyphenyl)-N-piperidin-4-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

tert-butyl 3-{[(5-{3-[2-(diethylamino)ethoxy]phenyl}-2-{[(pyrazin-2-ylamino)carbonyl]amino}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

- 13 -

5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino]carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide;

N-[3-(1,4-diazepan-1-ylcarbonyl)-5-(4-methoxyphenyl)-2-thienyl]-N'-pyrazin-2-ylurea;

N-[3-[(3-aminopyrrolidin-1-yl)carbonyl]-5-(4-methoxyphenyl)-2-thienyl]-N'-pyrazin-2-ylurea;

5 tert-butyl 4-[[5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino]carbonyl]amino]-3-thienyl]carbonyl]amino}piperidine-1-carboxylate;

tert-butyl 3-[[5-{4-[2-(diethylamino)ethoxy]phenyl}-2-[[pyrazin-2-ylamino]carbonyl]amino]-3-thienyl]carbonyl]amino}piperidine-1-carboxylate;

10 5-[4-(2-diethylamino-ethoxy)-phenyl]-2-(3-hydroxy-urea)-thiophene-3-carboxylic acid-(S)-piperidin-3-ylamide;

2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(3-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(2-hydroxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide;

15 2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-[2-(benzyloxy)phenyl]-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide.

20 One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt thereof for use as a medicament.

One embodiment of the present invention provides compounds of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment or prophylaxis of disorders associated with cancer.

25

One embodiment of the present invention provides a method for the treatment of cancer comprising administering to a human a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

- 14 -

One embodiment of the present invention provides a method for the treatment of breast cancer, colorectal cancer, ovarian cancer, lung (non small cell) cancer, malignant brain tumors, sarcomas, melanoma and lymphoma by administering a compound of formula I or a pharmaceutically acceptable salt thereof.

5

One embodiment the of present invention provides a method of treating cancer by administering to a human a compound of formula (I) or a pharmaceutically acceptable salt thereof and an anti-tumor agent.

10

One embodiment of the present invention provides a method of treating cancer by administering to a human a compound of formula (I) or a pharmaceutically acceptable salt thereof and a DNA damaging agent.

15

One embodiment of the present invention provides a method for the treatment of infections associated with cancer comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

20

One embodiment of the present invention provides a method for the prophylaxis treatment of infections associated with cancer comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

25


One embodiment of the present invention provides a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with at least one pharmaceutically acceptable carrier, diluent or excipient.

30

One embodiment of the present invention provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which comprises:


(a) the reaction of a 2-aminothiophene shown below as formula II

- 15 -




II

wherein the hydrogen at the 2-amino position is displaced to form an amide, shown as formula III below




III

wherein the methyl ester is converted to an amide utilizing the desired amine in conjunction with an aluminate organometallic complex, to give the product shown as formula IV below:



IV


wherein the amide is converted to various substituted secondary ureas by the reaction with various isocyanates to yield the product shown as formula V below:



V

- 16 -

One embodiment of the present invention provides the use of a compound of formula (VI) below or a pharmaceutically acceptable salt or an in vivo hydrolysable precursor in the manufacture of a compound of formula (I).



VI

The definitions set forth in this application are intended to clarify terms used throughout this application. The term "herein" means the entire application.

As used in this application, the term "optionally substituted," as used herein, means that substitution is optional and therefore it is possible for the designated atom to be unsubstituted. In the event a substitution is desired then such substitution means that any number of hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the normal valency of the designated atom is not exceeded, and that the substitution results in a stable compound. For example when a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced. Examples of such substituents are as follows:

halogen, nitro, amino, cyano, trifluoromethyl, C₁₋₆alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, hydroxy, alkylhydroxy, carbonyl, -CH(OH)CH₃, -CH₂NH-alkyl-OH, alkyl-(OH)CH₃, -CH₂-phenyl-(OCH₃)₂, -Oalkyl, -OCH₃, -Ophenyl, -OCOalkyl, -NHCHO, -N-(alkyl)-CHO, -Nalkyl, -NH-CO-amino, -N-(alkyl)-CO-amino, -NH-COalkyl, -N-(alkyl)-COalkyl, -carboxy, -amidino, -CO-amino, -CO-alkyl, -CO₂alkyl, mercapto, -Salkyl, -SCH₂furanyl, -SO(alkyl), -SO₂(alkyl), -SO₂-amino, -alkylsulfonylamino, phenyl, anisole, dimethoxyphenyl, trimethoxyphenyl, halophenyl, cycloalkyl, heterocyclyl, -alkyl-NH-cycloalkyl, -alkyl-NH-heterocyclyl, -alkyl-NH-alkyl-OH, -C(=O)OC(CH₃)₃, -N(CH₃)₂, -N(CH₂CH₃)₂, -alkyl-NH-alkyl-heterocyclyl, -alkyl-aryl, -methyl-phenyl, alkyl-polycyclyl, alkyl-amino, alkyl-hydroxy, -CH₂NH-alkyl-heterocyclyl, -CH₂NHCH₂CH(CH₃)₂.

- 17 -

If the selection is attached to a ring the substituents could also be selected from: vicinal -O(alkyl)O-, vicinal -OC(haloalkyl)O-, vicinal -CH₂O(alkyl)O-, vicinal -S(alkyl)S- and -O(alkyl)S-.

5

A variety of compounds in the present invention may exist in particular geometric or stereoisomeric forms. The present invention takes into account all such compounds, including cis- and trans isomers, R- and S- enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, as being covered within the scope of this invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention. The compounds herein described may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. When required, separation of the racemic material can be achieved by methods known in the art. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

30

- 18 -

As used herein, "alkyl" or "alkylene" used alone or as a suffix or prefix, is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having from 1 to 12 carbon atoms or if a specified number of carbon atoms is provided then that specific number would be intended. For example "C₁₋₆ alkyl" denotes alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms.

5 Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, pentyl, and hexyl. As used herein, "C₁₋₃ alkyl", whether a terminal substituent or an alkylene group linking two substituents, is understood to specifically include both branched and straight-chain methyl, ethyl, and propyl.

10 As used herein "alkylhydroxy" represents an alkyl group straight chain or branched as defined above with the indicated number of carbon atoms with one or more hydroxy groups attached. One such example of alkylhydroxy would be -CH₂OH.

15 As used herein, the term "carbocyclyl" is intended to include both alicyclic and aromatic ring structures wherein the closed ring is made of carbon atoms. These may include fused or bridged polycyclic systems. Carbocyclyls may have from 3 to 10 carbon atoms in their ring structure, and often have 3, 4, 5, and 6 carbons in the ring structure. For example, "C₃₋₆ carbocyclyl" denotes such groups as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopenta-diene or phenyl.

20

As used herein, the term "cycloalkyl" is intended to include saturated ring groups, having the specified number of carbon atoms. These may include fused or bridged polycyclic systems. Preferred cycloalkyls have from 3 to 10 carbon atoms in their ring structure, and more preferably have 3, 4, 5, and 6 carbons in the ring structure. For example, "C₃₋₆ cycloalkyl" denotes such
25 groups as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

As used herein, "alkenyl" or "alkenylene" is intended to include from 2 to 12 hydrocarbon atoms of either a straight or branched configuration with one or more carbon-carbon double bonds that may occur at any stable point along the chain. Examples of "C₃₋₆alkenyl" include, but

- 19 -

are not limited to, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 3-methyl-2-butenyl, 2-pentenyl, 3-pentenyl, hexenyl.

As used herein, "alkynyl" or "alkynylene" is intended to include from 2 to 12 hydrocarbon
5 chains of either a straight or branched configuration with one or more carbon-carbon triple bonds that may occur at any stable point along the chain. Examples of alkynyl include but are not limited to ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne.

As used herein, the term "alkylcycloalkyl" is intended to mean an alkyl attached to the
10 formula atom modified with a cycloalkyl. Examples of alkylcycloalkyl include, but are not limited to cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, cycloheptylmethyl, cyclopropylethyl, cyclopentylethyl, cyclohexylethyl, cycloheptylethyl, cyclopropylpropyl, cyclopentylpropyl, cyclohexylpropyl, cycloheptylpropyl.

As used herein, "cycloalkenyl" refers to ring-containing hydrocarbyl groups having at
15 least one carbon-carbon double bond in the ring, and having from 3 to 12 carbons atoms.

As used herein, "cycloalkynyl" refers to ring-containing hydrocarbyl groups having at
least one carbon-carbon triple bond in the ring, and having from 7 to 12 carbons atoms.

20

As used herein, the term "aralkyl" refers to an alkyl group substituted with an aryl group (an aromatic or heteroaromatic group).

As used herein, "aromatic" refers to hydrocarbyl groups having one or more
25 polyunsaturated carbon rings having aromatic character, (e.g., $4n + 2$ delocalized electrons) and comprising up to about 14 carbon atoms.

The term "aryl" as used herein includes 5-, 6- and 7-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, furan, imidazole,
30 isoxazole, nicotinic, isonicotinic, oxazole, phenyl, pyrazole, pyrazine, pyridazine, pyridine,

- 20 -

pyrimidine, thiazole, thiophene, triazole and the like. Those aryl groups having heteroatoms in the ring structure may also be referred to as "heteroaryl" or "heteroaromatics." The aromatic ring can be substituted at one or more ring positions with such substituents as described above. The term "aryl" also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings (the rings are "fused rings") wherein at least one of the rings is aromatic, for example, the other cyclic rings can be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls.

The terms ortho, meta and para apply to 1,2-, 1,3- and 1,4-disubstituted benzenes, respectively. For example, the names 1,2-dimethylbenzene and ortho-dimethylbenzene are synonymous.

As used herein, the term "heterocyclyl" or "heterocyclic" or "heterocycle" refers to a ring-containing monovalent and divalent structures having one or more heteroatoms, independently selected from N, O and S, as part of the ring structure and comprising from 3 to 20 atoms in the rings, more preferably 3- to 7- membered rings. Heterocyclic groups may be saturated or unsaturated, containing one or more double bonds, and heterocyclic groups may contain more than one ring as in the case of polycyclic systems. The heterocyclic rings described herein may be substituted on carbon or on a heteroatom atom if the resulting compound is stable. If specifically noted, nitrogen in the heterocyclyl may optionally be quaternized. It is understood that when the total number of S and O atoms in the heterocyclyl exceeds 1, then these heteroatoms are not adjacent to one another.

Examples of heterocyclyls include, but are not limited to, 1H-indazole, 2H, 6H-1, 5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidonyl, 4aH-carbazole, 4H-quinolizinyl, 6H-1, 2,5-thiadiazinyl, acridinyl, azabicyclo, azetidine, azepane, aziridine, azocinyl, benzimidazolyl, benzodioxol, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benzotriazolyl, benzotetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalonyl, carbazolyl, 4aH-carbazolyl, b-carbolinyl, chromanyl, chromenyl, cinnolinyl, diazepane, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dioxolane, furyl, 2,3-dihydrofuran, 2,5-dihydrofuran, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, homopiperidinyl, imidazolidine, imidazolidinyl, imidazolyl, 1H-indazolyl, indolenyl, indolinyl,

- 21 -

indoliziny, indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxirane, oxazolidinylperimidinyl, phenanthridinyl, phenanthrolinyl, phenarsazinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, pteridinyl, piperidonyl, 4-piperidonyl, purinyl, pyranyl, pyrrolidinyl, pyrroline, pyrrolidine, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, N-oxide-pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, pyrrolyl, pyridine, quinazolinyl, quinolinyl, 4H-quinoliziny, quinoxaliny, quinuclidinyl, carbolinyl, tetrahydrofuranyl, tetrahydroquinoline, tetrahydroisoquinolinyl, thiophane, thiotetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiopheneyl, thiirane, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, xanthenyl.

15

The terms “polycyclyl” or “polycyclic group” refer to two or more rings (for example, cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and /or heterocyclyls) in which two or more carbons are common to two adjoining rings, for example, the rings are “fused rings.” Rings that are joined through non-adjacent atoms are termed “bridged” rings. Each of the rings of the polycycle can be substituted with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, carbonyl, carboxyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like. Examples of such bridged heterocyclyls include quinuclidine, diazabicyclo[2.2.1]heptane and 7-oxabicyclo[2.2.1]heptane, substituted piperazine.


25

As used herein, the term “amine” or “amino” refers to groups of the general formula –NRR’, wherein R and R’ are each independently represented by but not limited to hydrogen, alkyl, cycloalkyl, alkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl. Example of the amino group include, but are not limited to NH₂, methylamine, ethylamine, dimethylamine, diethylamine, propylamine, benzylamine and the like.

30

- 22 -

As used herein, the term "amido" is art-recognized as an amino-substituted carbonyl and includes a moiety that can be represented by the general formula:



5 wherein R and R' are each independently represented by but not limited to hydrogen, alkyl, cycloalkyl, alkenyl, aryl, heteroaryl, heterocyclyl, aralkyl, or heteroaralkyl, or R and R' may form a ring.


As used herein, "alkoxy" or "alkyloxy" represents an alkyl group as defined above with
10 the indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, t-butoxy, n-pentoxo, isopentoxo, cyclopropylmethoxy, allyloxy and propargyloxy. Similarly, "alkylthio" or "thioalkoxy" represent an alkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge.

15

As used herein, the term "acyl" refers to groups of the of the general formula ---C(=O)---R , wherein R is hydrogen, hydrocarbonyl radical. Examples of acyl groups include, but are not limited to acetyl, propionyl, benzoyl, phenyl acetyl.

20

As used herein, the term "carbonyl" is art recognized and includes such moieties as can be represented by the general formula:




wherein X is a bond or represents an oxygen or sulfur, and R represents a hydrogen, an alkyl, an alkenyl, $\text{---(CH}_2\text{)}_m\text{---R''}$ or a pharmaceutically acceptable salt, R' represents a hydrogen, an alkyl, an
25 alkenyl or $\text{---(CH}_2\text{)}_m\text{---R''}$, where m is an integer less than or equal to ten, and R'' is alkyl, cycloalkyl, alkenyl, aryl, or heteroaryl. Where X is an oxygen and R and R' is not hydrogen, the formula represents an "ester". Where X is an oxygen, and R is as defined above, the moiety is

- 23 -


referred to herein as a carboxyl group, and particularly when R' is a hydrogen, the formula represents a "carboxylic acid." Where X is oxygen, and R' is a hydrogen, the formula represents a "formate." In general, where the oxygen atom of the above formula is replaced by sulfur, the formula represents a "thiolcarbonyl" group. Where X is a sulfur and R and R' is not hydrogen, the formula represents a "thiolester." Where X is sulfur and R is hydrogen, the formula represents a "thiolcarboxylic acid." Where X is sulfur and R' is hydrogen, the formula represents a "thiolformate." On the other hand, where X is a bond, and R is not a hydrogen, the above formula represents a "ketone" group. Where X is a bond, and R is hydrogen, the above formula is represents an "aldehyde" group.

As used herein, the term "sulfonylamino" is art-recognized and refers to a moiety that can be represented by the general formula:



wherein R and R' are each independently represented by but not limited to hydrogen, alkyl, cycloalkyl, alkenyl, aryl, heteroaryl, heterocyclyl, aralkyl, or heteroaralkyl.

As used herein, the term "sulfonyl" is art-recognized and refers to a moiety that can be represented by the general formula:



wherein R is represented by but not limited to hydrogen, alkyl, cycloalkyl, alkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl.

As used herein, "halo" or "halogen" refers to fluoro, chloro, bromo, and iodo.

"Counterion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate, tosylate, benzenesulfonate, and the like.

- 24 -

As used herein, "haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example $-C_vF_w$ where $v=1$ to 3 and $w=1$ to $(2v+1)$). Examples of haloalkyl include, but are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, pentachloroethyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, heptafluoropropyl, and heptachloropropyl. "Haloalkoxy" is intended to mean a haloalkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge; for example trifluoromethoxy, pentafluoroethoxy, 2,2,2-trifluoroethoxy, and the like. "Haloalkylthio" is intended to mean a haloalkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge.

As used herein, the phrase "protecting group" means temporary substituents which protect a potentially reactive functional group from undesired chemical transformations. Examples of such protecting groups include esters of carboxylic acids, silyl ethers of alcohols, and acetals and ketals of aldehydes and ketones respectively. The field of protecting group chemistry has been reviewed (Greene, T.W.; Wuts, P.G.M. *Protective Groups in Organic Synthesis*, 3rd ed.; Wiley: New York, 1999).

As used herein, "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional

- 25 -

non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, phosphoric, and the like; and the salts prepared from organic acids such as lactic, maleic, citric, benzoic, methanesulfonic, and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are used.

As used herein, "in vivo hydrolysable ester" means an in vivo hydrolysable (or cleavable) ester of a compound of the formula (I) that contains a carboxy or a hydroxy group. For example amino acid esters, C₁₋₆ alkoxymethyl esters like methoxymethyl; C₁₋₆alkanoyloxymethyl esters like pivaloyloxymethyl; C₃₋₈cycloalkoxycarbonyloxy C₁₋₆alkyl esters like 1-cyclohexylcarbonyloxyethyl, acetoxymethoxy, or phosphoramidic cyclic esters.

As used herein "stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The anti-cancer treatment defined herein may be applied as a sole therapy or may involve, in addition to the compound of the invention, conventional surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti-tumor agents:

(i) antiproliferative/antineoplastic/DNA damaging drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan and nitrosoureas);

- 26 -

antimetabolites (for example antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea and gemcitabine); antitumour antibiotics (for example anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and vinorelbine and taxoids like taxol and taxotere); topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan, irinotecan and camptothecin) and cytodifferentiating agents (for example All-trans retinoic acid, 13-cis retinoic acid and fenretinide);

(ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene and idoxifene), oestrogen receptor down regulators (for example fulvestrant), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and buserelin), progestogens (for example megestrol acetate), aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase such as finasteride;

(iii) agents which inhibit cancer cell invasion (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function);

(iv) inhibitors of growth factor function, for example such inhibitors include growth factor antibodies, growth factor receptor antibodies (for example the anti-erbb2 antibody trastuzumab [HerceptinTM] and the anti-erbb1 antibody cetuximab [C225]), farnesyl transferase inhibitors, tyrosine kinase inhibitors and serine/threonine kinase inhibitors, for example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), for example inhibitors of the platelet-derived growth factor family and for example inhibitors of the hepatocyte growth factor family;

(v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth factor, (for example the anti-vascular endothelial cell growth factor antibody bevacizumab [AvastinTM], compounds such as those disclosed in published International Patent Applications WO 97/22596, WO 97/30035, WO 97/32856 and WO 98/13354) and compounds

- 27 -

that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function and angiostatin);

(vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in International Patent Applications WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224,
5 WO 02/04434 and WO 02/08213;

(vii) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;

(viii) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug
10 therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; and

(ix) immunotherapy approaches, including for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as
15 interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines and approaches using anti-idiotypic antibodies.

Such conjoint treatment may be achieved by way of the simultaneous, sequential or
20 separate dosing of the individual components of the treatment. Such combination products employ the compounds of this invention.

Compounds of the present invention may be administered orally, parenteral, buccal, vaginal, rectal, inhalation, insufflation, sublingually, intramuscularly, subcutaneously, topically,
25 intranasally, intraperitoneally, intrathoracically, intravenously, epidurally, intrathecally, intracerebroventricularly and by injection into the joints.

The dosage will depend on the route of administration, the severity of the disease, age and weight of the patient and other factors normally considered by the attending physician, when determining the individual regimen and dosage level as the most appropriate for a particular
30 patient.

- 28 -

An effective amount of a compound of the present invention for use in therapy of infection is an amount sufficient to symptomatically relieve in a warm-blooded animal, particularly a human the symptoms of infection, to slow the progression of infection, or to reduce in patients with symptoms of infection the risk of getting worse.

5 For preparing pharmaceutical compositions from the compounds of this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories.

A solid carrier can be one or more substances, which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, or tablet disintegrating agents; it can
10 also be an encapsulating material.

In powders, the carrier is a finely divided solid, which is in a mixture with the finely divided active component. In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

For preparing suppository compositions, a low-melting wax such as a mixture of fatty
15 acid glycerides and cocoa butter is first melted and the active ingredient is dispersed therein by, for example, stirring. The molten homogeneous mixture is then poured into convenient sized molds and allowed to cool and solidify.

Suitable carriers include magnesium carbonate, magnesium stearate, talc, lactose, sugar, pectin, dextrin, starch, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, a low-
20 melting wax, cocoa butter, and the like.

Some of the compounds of the present invention are capable of forming salts with various inorganic and organic acids and bases and such salts are also within the scope of this invention. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, phosphoric, and the like; and the salts prepared from organic acids such as lactic,
25 maleic, citric, benzoic, methanesulfonic, trifluoroacetate and the like.

In one embodiment a compound of the formula (I) or a pharmaceutically acceptable salt thereof for the therapeutic treatment (including prophylactic treatment) of mammals including humans, it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

- 29 -

In addition to the compounds of the present invention, the pharmaceutical composition of this invention may also contain, or be co-administered (simultaneously or sequentially) with, one or more pharmacological agents of value in treating one or more disease conditions referred to herein.

5 The term composition is intended to include the formulation of the active component or a pharmaceutically acceptable salt with a pharmaceutically acceptable carrier. For example this invention may be formulated by means known in the art into the form of, for example, tablets, capsules, aqueous or oily solutions, suspensions, emulsions, creams, ointments, gels, nasal
10 sprays, suppositories, finely divided powders or aerosols or nebulisers for inhalation, and for parenteral use (including intravenous, intramuscular or infusion) sterile aqueous or oily solutions or suspensions or sterile emulsions.

 Liquid form compositions include solutions, suspensions, and emulsions. Sterile water or water-propylene glycol solutions of the active compounds may be mentioned as an example of liquid preparations suitable for parenteral administration. Liquid compositions can also be
15 formulated in solution in aqueous polyethylene glycol solution. Aqueous solutions for oral administration can be prepared by dissolving the active component in water and adding suitable colorants, flavoring agents, stabilizers, and thickening agents as desired. Aqueous suspensions for oral use can be made by dispersing the finely divided active component in water together with a
viscous material such as natural synthetic gums, resins, methyl cellulose, sodium carboxymethyl
20 cellulose, and other suspending agents known to the pharmaceutical formulation art.

 The pharmaceutical compositions can be in unit dosage form. In such form, the composition is divided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of the preparations, for example, packeted tablets, capsules, and powders in vials or ampoules.
25 The unit dosage form can also be a capsule, cachet, or tablet itself, or it can be the appropriate number of any of these packaged forms.

 Compounds of formula (I) have been shown to inhibit checkpoint kinase activity in vitro. Inhibitors of checkpoint kinase have been shown to allow cells to progress inappropriately to the metaphase of mitosis leading to apoptosis of effected cells, and to therefore have anti-
30 proliferative effects. Therefore it is believed that the compounds of formula (I) may be used for

- 30 -

the treatment of neoplastic disease. Hence compounds of formula (I) and their salts are expected to be active against neoplastic disease such as carcinoma of the brain, breast, ovary, lung, colon, prostate, skin or other tissues, as well as leukemias and lymphomas, tumors of the central and peripheral nervous system, and other tumor types such as melanoma, sarcomas, fibrosarcoma and osteosarcoma. In addition, compounds of formula (I) are also expected to be useful for the treatment other proliferative diseases. It is expected that the compounds of formula (I) would most likely be used in combination with a broad range of DNA damaging agents but could also be used as a single agent.

Generally, the compounds of formula (I) have been identified in one or both assays described below as having an IC_{50} value of 100 micromolar or less. For example compound of example 2 has an IC_{50} value of 10 nM.

Checkpoint Kinase 1 Assay: This in vitro assay measures the inhibition of CHK1 kinase by compounds. The kinase domain is expressed in baculovirus and purified by the GST tag. Purified protein and biotinylated peptide substrate (Cdc25C) is then used in a 384 well automated Scintillation Proximity Assay (SPA). Specifically, peptide, enzyme and reaction buffer are mixed and aliquoted into a 384 well plate containing dilution series of compounds and controls. Cold and hot ATP are then added to initiate the reaction. After 2 hours, a SPA bead slurry, CsCl₂ and EDTA are added to stop the reaction and capture the biotinylated peptide. Plates are then counted on a Topcount. Data is analyzed and IC_{50} s determined for individual compounds.

Abrogation Assay: This cellular assay measures the ability of CHK1 inhibitors to abrogate the DNA-damage induced G2/M checkpoint. Compounds active against the enzyme (< 2 μ M) are tested in the cellular assay. Briefly HT29 cells (colon cancer cell line, p53 null) are plated in 96 well plates on day 1. The following day, cells are treated with camptothecin for 2 hours to induce DNA damage. After 2 hours, camptothecin is removed and cells are treated for an additional 18 hours with test compound and nocodazole, a spindle poison that traps in cells in mitosis that abrogate the checkpoint. Cells are then fixed with formaldehyde, stained for the presence of phosphohistone H3, a specific marker for mitosis and labeled with Hoechst dye so that cell number can be measured. Plates are scanned using the Mitotic Index protocol on the Array Scan (Cellomics). As a positive control for abrogation, 4 mM caffeine is used.

- 31 -

Compounds are tested in a 12-point dose response in triplicate. Data is analyzed and EC50s determined for individual compounds.

The compounds of the present invention can be prepared in a number of ways well known to one skilled in the art of organic synthesis. The compounds of the present invention can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Such methods include, but are not limited to, those described below. All references cited herein are hereby incorporated in their entirety by reference.


The novel compounds of this invention may be prepared using the reactions and techniques described herein. The reactions are performed in solvents appropriate to the reagents and materials employed and are suitable for the transformations being effected. Also, in the description of the synthetic methods described below, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of the experiment and workup procedures, are chosen to be the conditions standard for that reaction, which should be readily recognized by one skilled in the art. It is understood by one skilled in the art of organic synthesis that the functionality present on various portions of the molecule must be compatible with the reagents and reactions proposed. Such restrictions to the substituents, which are not compatible with the reaction conditions, will be readily apparent to one skilled in the art and alternate methods must then be used.

The starting materials for the examples contained herein are either commercially available or are readily prepared by standard methods from known materials. For example the following reactions are illustrations but not limitations of the preparation of some of the starting materials and examples used herein.

General procedures for making the compounds of the invention is as follows:


- 32 -

The first of these procedures initiates from a common intermediate. This intermediate 2-aminothiophene core is produced by a one-pot Gewald synthesis from reaction of cyanomethylacetate with various benzaldehydes and elemental sulfur under basic conditions shown below in Scheme I.




Scheme I

If not commercially available, the corresponding benzaldehydes could be synthesized using some or all of the transformations described in Scheme II.



Scheme II


Compounds of Formula (I) can then be synthesized from the general synthetic methods described below in Schemes III-IX. The first general method shown in Scheme III involves a Weinreb amide formation from reaction of either the trichloroacetyl-protected or free urea with an amino aluminate organometallic complex.



Scheme III

- 33 -


An alternate method for the generation of similar compounds is described in Scheme IV. This general route utilizes the same starting 2-aminothiophene ester from Scheme III but amide bond formation is executed from the reaction of the corresponding carboxylic acids with various amines. A variety of coupling agents can be used to effect this transformation including EDCI, DIC, BOP, and HATU under standard coupling methods very familiar to those practicing and trained in the art of organic synthesis. Generation of the final primary urea is then performed using a simple two-step reaction with trichloroacetylisocyanate followed by cleavage with ammonia in methanol.

**Scheme IV**


As shown in the following Schemes V-VII, the amide product formed prior to urea generation in Scheme IV can be used as a common intermediate for the formation of various substituted ureas where R^4 is not hydrogen. Reaction with isocyanates, acyl azides (in particular pyrazine acyl azides), or carbonyldiimidazole and amines (in particular hydroxylamine hydrochloride), leads to the creation of various substituted secondary ureas where R^4 is selected from OH, optionally substituted carbocyclyl, optionally substituted heterocyclyl, or optionally substituted C_{1-6} alkyl.

20


- 34 -



Scheme V




Scheme VI



Scheme VII

- 10 An additional general process presented in this invention involves an improvement on the construction of general compounds with formula (I). The method, which employs a Suzuki Coupling of a 5-bromothiophene intermediate as its key transformation, is shown in Scheme VIII. This method allows for increased diversity much later in the synthesis and is amenable to parallel combinatorial methods of organic synthesis. Commercially available 2-amino-
- 15 thiophene-3-carboxylic acid methyl ester is protected as the trichloroacetyl urea, followed by selective bromination at the 5-position with bromine in acetic acid. Removal of the protecting group with ammonia in methanol, followed by Weinreb amidation yields the common intermediate. Standard Suzuki reaction conditions are then employed to finally generate the target compounds.

- 35 -




Scheme VIII

5

If the final compound generated from any of the above-mentioned methods or schemes has a protecting group on nitrogen (in particular a carbamate) or oxygen (in particular an ether or ester) present anywhere on the molecule standard methods of removal can be utilized to generate final compounds. Shown in Scheme IX is the general method used for the deprotection of a tert-butoxycarbonyl carbamate to yield the secondary amine product ($R^2 = \text{XNHR}^3$) as the corresponding hydrochloride or trifluoroacetate salt. Cleavage of methyl ethers to phenols (not shown) is affected by reaction with boron tribromide in methylene chloride. Both of these methods are very familiar with those trained in the art of organic synthesis.

10



15

Scheme IX

Examples:Table 1

IUPAC Name	Ex.	LCMS (M+1, ES or APCI detection)	¹ H NMR (d ₆ -DMSO unless stated otherwise)	Synthesis Method (Scheme)
tert-butyl 3-{[(2- [(aminocarbonyl)amino]-5-{4-[2- (diethylamino)ethoxy]phenyl}-3- thienyl)carbonyl]amino}piperidine-1- carboxylate trifluoroacetate	1	560	MeOD; 7.55(d, 2H), 7.45(s, 1H), 7.05(d, 2H), 4.35(m, 2H), 3.90(m, 3H), 3.60(m, 2H), 3.30(m, 4H), 2.95(dd, 2H), 2.10(m, 1H), 1.80(m, 1H), 1.50(m, 2H), 1.40(s, 9H), 1.35(t, 6H)	III
2-[(aminocarbonyl)amino]-5-{4-[2- (diethylamino)ethoxy]phenyl}-N- piperidin-3-ylthiophene-3-carboxamide trifluoroacetate	2	460		III
2-[(aminocarbonyl)amino]-5-{3-[2- (diethylamino)ethoxy]phenyl}-N- piperidin-3-ylthiophene-3-carboxamide trifluoroacetate	3	460		III
2-[(aminocarbonyl)amino]-5-(4- methoxyphenyl)-N-[(3S)-piperidin-3- yl]thiophene-3-carboxamide	4	375	1.76-1.49 (m, 2H), 1.93 (brs, 2H), 2.85 (t, 2H), 3.16 (s, 1H), 3.22 (brd, 1H), 3.31 (brd, 1H), 3.77 (s, 3H), 4.11 (brs, 1H), 6.97 (d, 2H, J=8.59 Hz), 7.45 (d, 2H, J=8.59 Hz), 7.58 (s, 1H), 8.02 (d, 1H, J=7.33 Hz), 8.65 (brs, 2H), 10.81 (s, 1H)	III
tert-butyl 3-{[(2- [(aminocarbonyl)amino]-5-{3-[2- (diethylamino)ethoxy]phenyl}-3- thienyl)carbonyl]amino}piperidine-1- carboxylate trifluoroacetate	5	560	MeOD; 7.65(d, 1H), 7.30(dd, 2H), 7.20(s, 1H), 6.90(d, 1H), 4.35(m, 2H), 3.90(m, 2H), 3.60(m, 2H), 3.30(m, 4H), 3.00(m, 2H), 2.10(m, 1H), 1.80(m, 1H), 1.50(m, 2H), 1.45(s, 9H), 1.40(t, 6H)	III
2-[(aminocarbonyl)amino]-5-{4-[2- (diethylamino)ethoxy]phenyl}-N- piperidin-4-ylthiophene-3-carboxamide	6	460	MeOD; 7.52(d, 2H), 7.49(s, 1H), 7.01(d, 2H), 4.35(m, 2H), 3.60(m, 2H), 3.48(m, 1H), 3.42(m, 2H), 3.33(q, 4H), 3.13(m, 2H), 2.14(m, 2H), 1.85(m, 2H), 1.35(t, 6H)	III
2-[(aminocarbonyl)amino]-N-[(3R)- azepan-3-yl]-5-(4- methoxyphenyl)thiophene-3- carboxamide	7	389	CDCl ₃ ; 11.30(br s, 1H), 7.50 (d, 2H), 7.05(d, 1H), 7.00 (s, 1H), 6.85(d, 2H), 5.15 (s, 2H), 4.20(m, 1H), 3.80(s, 3H), 3.10 (m, 1H), 3.00 (m, 2H), 2.80 (m, 1H), 1.80(m, 2H), 1.70(m, 3H), 1.50 (m, 1H)	III

- 37 -

N-(3-[(4-aminopiperidin-1-yl)carbonyl]-5-{4-[2-(diethylamino)ethoxy]phenyl}-2-thienyl)urea trifluoroacetate	8	460	MeOD; 7.43(d, 2H), 6.93(d, 2H), 6.92(s, 1H), 4.33(m, 2H), 4.27(dd, 2H), 3.52(dd, 2H), 3.32(m, 1H), 3.24(q, 4H), 3.02(m, 2H), 1.99(m, 1H), 1.95(m, 1H), 1.47(m, 2H), 1.27(t, 6H)	III
2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[3-(hydroxymethyl)phenyl]thiophene-3-carboxamide trifluoroacetate (salt)	9	483	MeOD; 7.61(m, 1H), 7.57(s, 1H), 7.49(m, 1H), 7.47(d, 2H), 7.23(t, 1H), 7.03(dd, 1H), 6.93(d, 2H), 4.52(s, 2H), 4.27(dd, 2H), 4.31(dd, 2H), 3.24(q, 4H), 1.27(t, 6H)	III
2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-ylthiophene-3-carboxamide trifluoroacetate	10	460	MeOD; 7.62(s, 1H), 7.29(dd, 1H), 7.24(m, 1H), 7.13(m, 1H), 6.87(m, 1H), 4.35(dd, 2H), 3.59(dd, 2H), 3.46(m, 1H), 3.40(m, 2H), 3.31(q, 4H), 3.10(m, 2H), 2.16(m, 2H), 1.83(m, 2H), 1.34(t, 6H)	III
2-[(aminocarbonyl)amino]-N-(2-aminoethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	11	335	MeOD; 7.38(d, 2H), 7.36(s, 1H), 6.82(d, 2H), 3.70(s, 3H), 3.57(dd, 2H), 3.10(dd, 2H),	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	12	375	1.70 (m, 2H), 1.99 (d, 2H, J=12.88 Hz), 3.01 (q, 2H, J=11.03 Hz), 3.35 (d, 2H, J=12.38 Hz), 3.76 (s, 3H), 4.01 (brs, 1H), 6.97 (d, 2H, J=8.59 Hz), 7.45 (d, 2H, J=8.84 Hz), 7.63 (s, 1H), 8.04 (d, 1H, J=7.07 Hz), 8.43 (d, 1H, J=9.10 Hz), 8.65 (d, 1H, J=8.59 Hz), 10.91 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-pyridin-3-ylthiophene-3-carboxamide trifluoroacetate	13	454	MeOD; 9.51(s, 1H), 8.75(d, 1H), 8.51(m, 1H), 7.97(s, 1H), 7.93(m, 1H), 7.35(t, 1H), 7.29(m, 2H), 6.93(m, 1H), 4.45(dd, 2H), 3.67(dd, 2H), 3.39(q, 4H), 1.42(t, 6H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(1-methylpiperidin-4-yl)thiophene-3-carboxamide	14	389	1.76 (q, 2H), 2.04 (d, 2H, J=12.38 Hz), 2.77 (brd, 3H), 3.08 (q, 2H), 3.48 (d, 2H, J=11.62 Hz), 3.76 (s, 3H), 3.96 (m, 1H), 6.97 (d, 2H), 7.45 (d, 2H, J=8.84 Hz), 7.63 (s, 1H), 8.07 (d, 1H, J=7.33 Hz), 9.62 (brs, 1H), 10.91 (s, 1H)	III

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-1-methylazepan-3-yl]thiophene-3-carboxamide hydrochloride	15	403	CDCl ₃ ; 11.35(br s, 1H), 7.50 (d, 2H), 7.15(d, 1H), 6.95 (s, 1H), 6.90(d, 2H), 5.30 (s, 2H), 4.20(m, 1H), 3.80(s, 3H), 2.85 (m, 1H), 2.80 (d, 1H), 2.60 (dd, 1H), 2.40(m, 1H), 2.45(s, 3H), 1.50-2.00 (m, 6H)	III
2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-[3-(hydroxymethyl)phenyl]thiophene-3-carboxamide trifluoroacetate (salt)	16	483	MeOD; 7.84(s, 1H), 7.75(m, 1H), 7.63(dd, 1H), 7.35(t, 2H), 7.32(m, 1H), 7.22(m, 1H), 7.16(dd, 1H), 6.92(m, 1H), 4.65(s, 2H), 4.41(dd, 2H), 3.63(dd, 2H), 3.37(q, 4H), 1.40(t, 6H)	III
2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-pyrrolidin-3-ylthiophene-3-carboxamide trifluoroacetate	17	446	MeOD; 7.46(d, 2H), 7.16(s, 1H), 6.95(d, 2H), 4.28(dd, 2H), 3.89(m, 3H), 3.70 (m, 2H), 3.53(dd, 2H), 3.28(q, 4H), 2.30(m, 1H), 2.06(m, 1H), 1.29(t, 6H)	III
2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-pyridin-3-ylthiophene-3-carboxamide trifluoroacetate	18	454	MeOD; 9.59(s, 1H), 8.75(d, 1H), 8.56(d, 1H), 8.00(m, 2H), 7.71(s, 1H), 7.59(d, 2H), 7.07(d, 2H), 4.41(dd, 2H), 3.65(dd, 2H), 3.38(q, 4H), 1.40(t, 6H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-1-methylpiperidin-3-yl]thiophene-3-carboxamide hydrochloride	19	389	CDCl ₃ ; 11.25(br s, 1H), 7.50 (d, 2H), 7.15(s, 1H), 6.85 (d, 2H), 6.90(m, 2H), 5.40 (s, 2H), 4.25(m, 1H), 3.80(s, 3H), 2.40-2.80 (m, 4H), 2.30(s, 3H), 2.20(m, 1H), 1.50-1.90 (m, 3H)	III
2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-pyrrolidin-3-ylthiophene-3-carboxamide trifluoroacetate	20	446	MeOD; 7.40(s, 1H), 7.35(dd, 1H), 7.27(dd, 1H), 7.20(s, 1H), 6.93(m, 1H), 4.42(dd, 2H), 4.01(m, 3H), 3.83 (m, 2H), 3.65(dd, 2H), 3.37(q, 4H), 2.44(m, 1H), 2.18(m, 1H), 1.40(t, 6H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-ylmethyl]thiophene-3-carboxamide	21	389	1.22 (dd, 1H, J ₁ =12.38 Hz, J ₂ =2.53 Hz), 1.57 (d, 1H, J=13.14 Hz), 1.78 (brt, 2H), 1.95 (brs, 1H), 2.60 (q, 1H, J=11.37), 2.78 (q, 1H, J=10.61 Hz), 3.09 (m, 1H), 3.25 (m, 3H), 3.77 (s, 3H), 6.97 (d, 2H, J=8.59 Hz), 7.44 (d, 2H, J=8.84 Hz), 7.58 (s, 1H), 8.24 (brs, 1H), 8.30 (brt, 1H), 8.57 (brs, 1H), 10.92 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide	22	361		III

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]thiophene-3-carboxamide	23	361		III
2-[(aminocarbonyl)amino]-N-[2-(dimethylamino)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide	24	363	MeOD; 7.37(d, 2H), 7.35(s, 1H), 6.80(d, 2H), 3.69(s, 3H), 3.64(m, 2H), 3.29(m, 2H), 2.91(s, 6H)	III
2-[(aminocarbonyl)amino]-N-[2-(diethylamino)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide	25	391		III
2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide hydrochloride	26	389	δ 10.9, s, 1H; δ 9.58, br s, 1H; δ 9.29, br s, 1H; δ 8.39, d, 1H; δ 7.82, s, 1H; δ 7.48, d, 2H; δ 6.96, d, 2H; δ 4.36, m, 1H; δ 3.77, s, 3H; δ 3.29, m, 1H; δ 3.20, m, 2H; δ 3.07, m, 1H; δ 1.98, m, 1H; δ 1.84, m, 4H; δ 1.59, m, 1H	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-yl]thiophene-3-carboxamide	27	375	1.52 (m, 2H), 1.76 (brs, 1H), 1.89 (brs, 1H), 2.61 (m, 2H), 2.99 (t, 1H), 3.16 (m, 1H), 3.76 (s, 3H), 3.95 (brs, 1H), 6.97 (d, 2H), 7.46 (d, 2H), 7.61 (s, 1H), 7.89 (d, 1H), 10.90 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(piperidin-4-ylmethyl)thiophene-3-carboxamide	28	389	1.06 (m, 2H), 1.62 (d, 3H, J=10.11 Hz), 2.95 (d, 2H, J=12.13 Hz), 3.11 (t, 2H, J=5.68 Hz), 3.76 (s, 3H), 6.90 (d, 1H, J=8.59 Hz), 6.96 (d, 2H, J=8.59 Hz), 7.44 (d, 2H, J=8.59 Hz), 7.61 (s, 1H), 8.14 (t, 1H, J=5.43 Hz), 10.99 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-pyrrolidin-3-ylthiophene-3-carboxamide	29	361	2.02 (brt, 2H, J=5.81 Hz), 2.22 (dd, 2H, J1=13.52 Hz, J2=6.69 Hz), 3.21 (dd, 1H, J1=13.77 Hz, J2=6.95 Hz), 3.68 (brs, 1H), 3.76 (s, 3H), 3.86 (brs, 1H), 6.96 (brm, 2H), 7.28 (brs, 1H), 7.50 (d, 2H, J=8.84 Hz), 8.16 (brs, 1H), 10.31 (s, 1H)	III
2-[(aminocarbonyl)amino]-N-(1-ethylpiperidin-3-yl)-5-(4-methoxyphenyl)thiophene-3-carboxamide hydrochloride	30	403	11.0(s, 1H), 7.80(d, 1H), 7.65(s, 1H), 7.45(d, 2H), 6.95(d, 2H), 6.90(br s, 2H), 3.90(m, 1H), 3.75(s, 3H), 2.85(dd, 2H), 2.30(m, 2H), 1.80(m, 3H), 1.70(m, 1H), 1.50(m, 1H), 13.0(m, 1H), 1.0(t, 3H)	III

- 40 -

2-[(aminocarbonyl)amino]-N-[(3S)-1-ethylazepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide hydrochloride	31	417	11.0(s, 1H), 7.75(d, 1H), 7.65(s, 1H), 7.45(d, 2H), 6.95(d, 2H), 6.90(br s, 2H), 4.05(m, 1H), 3.75(s, 3H), 2.75(m, 1H), 2.40-2.70(m, 5H), 1.85(m, 1H), 1.40-1.75(m, 5H), 0.95(t, 3H)	III
2-[(aminocarbonyl)amino]-5-(3-hydroxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	32	361		III
2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	33	361		III
2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	34	375		III
tert-butyl (3S)-3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)pyrrolidine-1-carboxylate	35	481		III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-piperidin-3-ylthiophene-3-carboxamide	36	375		III
2-[(aminocarbonyl)amino]-N-(1-benzylpiperidin-4-yl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	37	465	1.75 (m, 2H), 2.08 (d, 2H, J=13.39 Hz), 3.10 (m, 2H), 3.43 (d, 2H, J=11.62 Hz), 3.76 (s, 3H), 3.95 (m, 1H), 4.31 (d, 2H, J=4.55 Hz), 6.97 (d, 2H, J=8.59 Hz), 7.45 (d, 2H, J=8.59 Hz), 7.49 (brs, 5H, J=2.27 Hz), 7.61 (s, 1H), 8.05 (d, 1H, J=6.82 Hz), 9.65 (brs, 1H), 10.89 (s, 1H)	III
tert-butyl 3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate	38	475	1.39 (s, 9H), 1.46-1.97 (bm, 4H), 2.72 (t, 2H), 3.16 (s, 2H), 3.77 (s, 3H), 6.96 (d, 2H), 7.45 (d, 2H), 7.62 (s, 1H), 7.85 (brs, 1H), 8.77 (d, 1H), 10.95 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-[4-(2-piperidin-1-ylethoxy)phenyl]-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide	39	494		III
2-[(aminocarbonyl)amino]-5-[4-(2-piperidin-1-ylethoxy)phenyl]-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide	40	451		III

- 41 -

2-[(aminocarbonyl)amino]-N-azetidin-3-yl-5-(4-methoxyphenyl)thiophene-3-carboxamide	41	347		III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(2S)-pyrrolidin-2-ylmethyl]thiophene-3-carboxamide	42	375	1.71 (m, 1H), 1.92 (m, 2H), 2.04 (m, 1H), 3.33-3.11 (m, 2H), 3.62-3.45 (m, 2H), 3.76 (s, 3H), 6.97 (d, 2H, J=8.84 Hz), 7.45 (d, 2H, J=8.59 Hz), 7.52 (s, 1H), 7.94 (s, 1H), 8.47 (t, 1H, J=5.81 Hz), 8.52 (brs, 1H), 9.15 (brs, 1H), 10.79 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-pyridin-4-ylthiophene-3-carboxamide	43	369	3.78 (s, 3H), 7.01 (d, 2H), 7.52 (d, 2H), 7.79 (s, 1H), 8.15 (d, 2H), 8.70 (brs, 2H), 10.62 (s, 1H), 10.68 (brs, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperazin-1-ylethyl)thiophene-3-carboxamide	44	404	2.06 (m, 4H), 2.34 (brs, 2H), 2.42 (m, 2H), 2.67 (m, 4H), 3.76 (s, 3H), 6.97 (d, 2H, J=8.84 Hz), 7.44 (d, 2H, J=8.59 Hz), 7.55 (s, 1H), 8.10 (t, 1H, J=5.68 Hz), 10.95 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperidin-1-ylethyl)thiophene-3-carboxamide	45	403	1.37 (s, 2H), 1.48 (s, 4H), 2.37 (brs, 4H), 2.42 (t, 2H), 3.76 (s, 3H), 6.94 (brs, 1H), 6.97 (d, 2H), 7.44 (d, 2H), 7.56 (t, 2H), 8.11 (t, 1H), 10.96 (s, 1H)	III
2-[(aminocarbonyl)amino]-N-1-azabicyclo[2.2.2]oct-3-yl-5-(4-methoxyphenyl)thiophene-3-carboxamide	46	401		III
2-[(aminocarbonyl)amino]-N-(2-hydroxyethyl)-5-(4-hydroxyphenyl)thiophene-3-carboxamide	48	322	3.06(t,1H), 3.5(m,2H), 6.54(bs,1H), 6.78-6.97(m,4H), 7.33(m,2H),7.49(s,1H), 8.34(m,1H), 9.56(s,1H), 10.94(s,1H)	III
2-[(aminocarbonyl)amino]-N-(trans-4-hydroxycyclohexyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	49	390	1.10-1.45 (dq, 4H), 1.85 (t, 4H), 3.40 (brs, 1H), 3.71 (brm, 1H), 3.76 (s, 3H), 4.56 (brs, 1H), 6.90 (brs, 1H), 6.96 (d, 2H), 7.45 (d, 2H), 7.62 (s, 1H), 7.81 (d, 1H), 11.03 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide	50	383	3.17 (t, 2H, J=6.44 Hz), 3.64 (m, 2H), 6.77 (d, 2H, J=8.59 Hz), 7.31 (d, 2H, J=8.59 Hz), 7.45 (s, 1H), 7.97 (d, 2H, J=6.32 Hz), 8.27 (s, 1H), 8.83 (d, 2H, J=6.06 Hz), 10.79 (s, 1H)	III

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperazin-1-ylethyl)thiophene-3-carboxamide	52	525	1.34-1.58 (m, 2H), 1.75 (t, 2H, J=14.02 Hz), 1.98 (brs, 1H), 2.01 (d, 1H), 2.80 (m, 3H), 3.48 (d, 2H, J=8.59 Hz), 3.73 (s, 6H), 3.75 (s, 3H), 6.63 (dd, 2H, J1=8.34 Hz, J2=4.80 Hz), 6.90 (brs, 1H), 6.95 (d, 2H, J=8.59 Hz), 7.21 (t, 2H), 7.45 (d, 2H, J=8.84 Hz), 7.46 (s, 1H), 7.86 (d, 1H, J=7.58 Hz), 11.00 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide	54	397	3.12 (t, 2H), 3.62 (t, 2H), 3.76 (s, 3H), 6.97 (d, 2H), 7.42 (d, 2H), 7.49 (s, 1H), 7.85 (d, 2H), 8.28 (s, 1H), 8.78 (d, 2H), 10.84 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-pyridin-3-ylethyl)thiophene-3-carboxamide	55	383	3.17 (t, 2H, J=6.44 Hz), 3.64 (m, 2H), 6.77 (d, 2H, J=8.59 Hz), 7.31 (d, 2H, J=8.59 Hz), 7.45 (s, 1H), 7.97 (d, 2H, J=6.32 Hz), 8.27 (s, 1H), 8.83 (d, 2H, J=6.06 Hz), 10.79 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-3-ylethyl)thiophene-3-carboxamide	56	397	2.99 (t, 2H), 3.55 (m, 2H), 3.76 (s, 3H), 6.97 (d, 2H), 7.42 (d, 2H), 7.50 (s, 1H), 7.76 (m, 1H), 8.16 (d, 1H), 8.25 (s, 1H), 8.65 (s, 1H), 8.69 (s, 1H), 10.85 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)thiophene-3-carboxamide	58	431	1.44 (s, 12H), 1.63 (t, 2H), 1.95 (d, 2H), 3.76 (s, 3H), 4.34 (brs, 1H), 6.96 (d, 2H), 7.45 (d, 2H), 7.67 (s, 1H), 8.12 (m, 2H), 9.06 (bd, 1H), 10.92 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(2-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	59	375		III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(tetrahydrofuran-2-ylmethyl)thiophene-3-carboxamide	60	376	1.52(m,1H), 1.77(m,3H), 3.25(m,2H), 3.72(m,1H), 3.71(s,3H), 3.93(m,2H), 6.91(bd,4H,J=8.84Hz), 7.39(d,2H,J=8.59Hz), 7.59(s,1H), 8.17(bs,1H), 10.91(s,1H)	III
tert-butyl (3R)-3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate	62	475	1.38 (s, 9H), 1.47-1.99 (brm, 6H), 3.56 (brm, 2H), 3.74 (brm, 1H), 3.76 (s, 3H), 6.96 (d, 2H, J=8.59 Hz), 7.45 (d, 2H, J=8.84 Hz), 7.62 (s, 1H), 7.85 (brs, 1H), 8.77 (d, 1H, J=7.58 Hz), 10.96 (s, 1H)	III

- 43 -

2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-3-ylmethyl)thiophene-3-carboxamide	63	383	3.76 (s, 3H), 4.48 (d, 2H), 6.96 (d, 2H), 7.36 (dd, 1H), 7.44 (d, 2H), 7.61 (s, 1H), 7.72 (dt, 1H), 8.46 (bdd, 1H), 8.56 (bd, 1H), 8.77 (t, 1H)	III
tert-butyl 3-([(2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl)amino)azetidine-1-carboxylate	64	447	1.39 (s, 9H), 3.77 (s, 3H), 3.86 (m, 2H), 4.13 (t, 2H, J=7.83 Hz), 4.61 (m, 1H), 6.97 (d, 2H, J=8.84 Hz), 7.45 (d, 2H, J=8.59 Hz), 7.60 (s, 1H), 8.57 (d, 1H, J=6.82 Hz), 10.82 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-4-ylmethyl)thiophene-3-carboxamide	65	383		III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(3-methoxypropyl)thiophene-3-carboxamide	67	364	1.76 (m, 2H), 3.23 (s, 3H), 3.30 (t, 2H), 3.37 (t, 2H), 3.76 (s, 3H), 6.96 (d, 2H), 7.44 (d, 2H), 7.57 (s, 1H), 8.15 (brs, 1H), 10.97 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[2-(2-thienyl)ethyl]thiophene-3-carboxamide	68	402		III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-thienylmethyl)thiophene-3-carboxamide	69	388	3.76 (s, 3H), 4.62 (d, 2H), 6.96 (d, 1H), 6.96 (d, 2H), 7.03 (d, 1H), 7.39 (d, 1H), 7.43 (d, 2H), 7.60 (s, 1H), 8.79 (t, 1H), 10.92 (s, 1H)	III
N-[3-(1,4-diazepan-1-ylcarbonyl)-5-(4-methoxyphenyl)-2-thienyl]urea	70	375	7.50 d J=8.8 Hz 2H, 7.09 s 1H, 6.95 d J=8.8 Hz 2H, 3.86 s 3H, 1.23-1.35 m 6H.	III
2-[(aminocarbonyl)amino]-N-(2-methoxyethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	71	350	3.27 (s, 3H), 3.37-3.51 (m, 4H), 3.76 (s, 3H), 6.97 (d, 2H), 7.44 (d, 2H), 7.61 (s, 1H), 8.15-8.26 (m, 1H), 10.95 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-thienylmethyl)thiophene-3-carboxamide	72	374	4.61 (d, 2H, J=5.81 Hz), 6.77 (d, 2H, J=8.59 Hz), 6.96 (dd, 2H, J1=5.05 Hz, J2=3.54 Hz), 7.02 (d, 1H, J=3.03 Hz), 7.32 (d, 2H, J=8.34 Hz), 7.38 (d, 1H, J=5.05 Hz), 7.53 (s, 1H), 8.77 (t, 1H, J=5.81 Hz), 9.53 (s, 1H), 10.89 (s, 1H)	III

- 44 -

2-[(aminocarbonyl)amino]-N-{2-[(2-furylmethyl)thio]ethyl}-5-(4-methoxyphenyl)thiophene-3-carboxamide	74	432	2.61 (t, 2H), 2.65 (t, 2H), 3.75 (s, 3H), 3.81 (d, 2H), 6.29 (d, 1H, J=3.03 Hz), 6.38 (dd, 1H, J1=3.03 Hz, J2=1.77 Hz), 6.97 (d, 2H, J=8.84 Hz), 7.44 (d, 2H, J=8.84 Hz), 7.55 (s, 1H), 7.57 (brs, 1H), 8.31 (t, 1H, J=5.68 Hz), 9.10 (t, 1H, J=4.93 Hz), 10.92 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-[2-(2-thienyl)ethyl]thiophene-3-carboxamide	75	388	3.50(m,2H), 3.80(m,2H), 6.54(bs,1H), 6.78-7.34(m,7H), 7.49(s,1H), 8.34(m,1H), 9.56(s,1H), 10.94(s,1H)	III
N-(3-[(4-aminopiperidin-1-yl)carbonyl]-5-{3-[2-(diethylamino)ethoxy]phenyl}-2-thienyl)urea trifluoroacetate	76	460	MeOD; 7.24(dd, 1H), 7.16(dd, 1H), 7.08(s, 1H), 7.06(s, 1H), 6.83(dd, 1H), 4.34(m, 2H), 4.30(dd, 2H), 3.53(dd, 2H), 3.32(m, 1H), 3.27(q, 4H), 3.04(m, 2H), 1.99(m, 2H), 1.50(m, 2H), 1.28(t, 6H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-ylmethyl]thiophene-3-carboxamide	77	389	1.22 (dd, 1H, J1=12.38 Hz, J2=2.53 Hz), 1.57 (d, 1H, J=13.14 Hz), 1.78 (brt, 2H), 1.95 (brs, 1H), 2.60 (q, 1H, J=11.37), 2.78 (q, 1H, J=10.61 Hz), 3.09 (m, 1H), 3.25 (m, 3H), 3.77 (s, 3H), 6.97 (d, 2H, J=8.59 Hz), 7.44 (d, 2H, J=8.84 Hz), 7.58 (s, 1H), 8.24 (brs, 1H), 8.30 (brt, 1H), 8.57 (brs, 1H), 10.92 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(1,2,3,4-tetrahydroquinolin-3-yl)thiophene-3-carboxamide	78	423		III
2-[(aminocarbonyl)amino]-N-(1,3-benzodioxol-5-ylmethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	79	426	3.76 (s, 3H), 4.36 (d, 2H), 5.97 (s, 2H), 6.81 (d, 1H), 6.89 (d, 1H), 6.96 (d, 2H), 7.44 (d, 2H), 7.62 (s, 1H), 8.64 (t, 1H), 10.94 (s, 1H)	III
2-[(aminocarbonyl)amino]-N-(3-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	80	412	3.74(s,3H), 3.78(s,3H), 4.44(d,2H,J=5.56Hz), 6.82(m,1H), 6.91(m,2H), 6.98(d,2H,J=8.59Hz), 7.25(t, 2H, J=7.83), 7.46(d,2H,J=8.59Hz), 7.66(s,1H), 8.71(s,1H), 10.95(s,1H)	III

- 45 -

2-[(aminocarbonyl)amino]-N-[2-(3,4-dimethoxyphenyl)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide	81	456	3.73(s,3H), 3.74(s,3H), 3.77(s,3H), 4.40(d,2H,J=5.56Hz), 6.92(m,7H), 7.45(d,2H,J=8.84Hz), 7.66(s,1H), 8.64(bs,1H), 10.97(s,1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(5-methyl-2-furyl)methyl]thiophene-3-carboxamide	84	386	2.22 (s, 3H), 3.76 (s, 3H), 4.38 (d, 2H), 5.99 (s, 1H), 6.14 (s, 1H), 6.96 (d, 2H), 7.43 (d, 2H), 7.64 (s, 1H), 8.57 (t, 1H), 10.93 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-2-ylmethyl)thiophene-3-carboxamide	85	383	3.77 (s, 3H), 4.69 (d, 2H), 6.98 (d, 2H), 7.46 (d, 2H), 7.65 (t, 1H), 7.73 (t, 1H), 7.67 (s, 1H), 8.20 (t, 1H), 8.69 (d, 1H), 8.96 (t, 1H), 10.76 (s, 1H)	III
2-[(aminocarbonyl)amino]-N-(4-fluorobenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	86	400		III
tert-butyl 4-({[2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate	88	475		III
2-[(aminocarbonyl)amino]-N-(2-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	89	412	3.78(s,3H), 3.83(s,3H), 4.45(bd,2h,J=5.56Hz), 6.97(m,5H), 7.23(m,2H), 7.47(d,2H,J=8.84), 7.71(s,1H), 8.55(bt,1H), 10.97(s,1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-phenoxyethyl)thiophene-3-carboxamide	90	412	3.56-3.7 (m, 2H), 3.76 (s, 3H), 4.11 (t, 2H), 6.85-7.05 (m, 5H), 7.28 (t, 2H), 7.44 (d, 2H), 7.61 (s, 1H), 8.38 (brs, 1H), 10.93 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-2-ylethyl)thiophene-3-carboxamide	93	397	3.00 (t, 2H, J=7.45 Hz), 3.61 (m, 2H), 3.76 (s, 3H), 6.97 (d, 2H, J=8.84 Hz), 7.22 (dd, 1H, J1=6.95 Hz, J2=5.18 Hz), 7.27 (d, 1H, J=7.83 Hz), 7.43 (d, 2H, J=8.59 Hz), 7.54 (s, 1H), 7.70 (dt, 1H, J1=7.58 Hz, J2=1.77 Hz), 8.27 (t, 1H, J=5.56 Hz), 8.51 (d, 1H, J=4.55 Hz), 10.97 (s, 1H)	III
tert-butyl 4-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate	94	475		III

- 46 -

2-[(aminocarbonyl)amino]-N-(4-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	95	412	3.72 (s, 3H), 3.76 (s, 3H), 4.39 (d, 2H), 6.89 (d, 2H), 6.96 (d, 2H), 7.25 (d, 2H), 7.44 (d, 2H), 7.63 (s, 1H), 8.64 (t, 1H), 10.95 (s, 1H)	III
2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide trifluoroacetate	110	460	MeOD; 7.55(d, 2H), 7.45(s, 1H), 7.05(d, 2H), 4.35(dd, 2H), 4.25(m, 1H), 3.60(dd, 2H), 3.50(m, 1H), 3.30(m, 5H), 2.95(dd, 2H), 2.10(dd, 2H), 1.80(m, 2H), 1.35(t, 6H)	IV
2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3R)-piperidin-3-yl]thiophene-3-carboxamide trifluoroacetate	111	460	MeOD; 7.55(d, 2H), 7.45(s, 1H), 7.05(d, 2H), 4.35(dd, 2H), 4.25(m, 1H), 3.60(dd, 2H), 3.50(m, 1H), 3.30(m, 5H), 2.95(dd, 2H), 2.10(dd, 2H), 1.80(m, 2H), 1.35(t, 6H)	IV
tert-butyl (3S)-3-[[2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl]carbonyl]amino}piperidine-1-carboxylate trifluoroacetate	112	560	MeOD; 7.55(d, 2H), 7.45(s, 1H), 7.05(d, 2H), 4.35(dd, 2H), 3.60-3.90(m, 3H), 3.60(dd, 2H), 3.30(m, 4H), 2.95(m, 2H), 1.90(dd, 2H), 1.55(m, 2H), 1.45(s, 9H), 1.35(t, 6H)	IV
2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-{4-[2-(diethylamino)ethoxy]phenyl}thiophene-3-carboxamide hydrochloride	113	474	10.85(s, 1H), 9.50(br s, 1H), 9.10(br s, 1H), 8.95(br s, 1H), 8.20(d, 1H), 7.65(s, 1H), 7.50(d, 2H), 7.05(d, 2H), 6.95, (br s, 2H), 4.35(dd, 2H), 4.25(m, 1H), 3.50(dd, 2H), 3.10-3.40(m, 8H), 2.00(m, 1H), 1.80(m, 4H), 1.55(m, 1H), 1.25(t, 6H)	IV
tert-butyl (3R)-3-[[2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl]carbonyl]amino}piperidine-1-carboxylate trifluoroacetate	114	560	MeOD; 7.45(d, 2H), 7.35(s, 1H), 6.90(d, 2H), 4.25(dd, 2H), 3.60-3.90(m, 3H), 3.50(dd, 2H), 3.20(m, 5H), 2.85(s, 1H), 1.80(dd, 2H), 1.45(m, 2H), 1.35(s, 9H), 1.25(t, 6H)	IV
N-[3-[(3S)-3-aminoazepan-1-yl]carbonyl]-5-(4-methoxyphenyl)-2-thienyl]urea hydrochloride	115	389	9.65 (s, 1H), 8.25 (s, 3H), 7.50 (d, 2H), 7.10(s, 1H), 6.90(d, 2H), 6.75 (br s, 2H), 4.00 (m, 1H), 3.80(s, 3H), 3.40 (m, 4H), 2.0 (m, 1H), 1.00-1.80 (m, 5H)	IV
2-([(2,5-dimethoxyphenyl)amino]carbonyl)amino)-5-(4-methoxyphenyl)-N-piperidin-3-ylthiophene-3-carboxamide	116	511	11.28 s 1H, 9.65 bs 1H, 9.08 bs 1H, 8.43 s 1H, 7.90 s 1H, 7.46 d J = 8.7 Hz 2H, 6.83 d J = 8.8 Hz 2H, 6.65 d J = 8.8 Hz 1H, 6.43 - 6.47 m 1H, 4.35 bs 1H, 3.76 s 3H, 7.73 s 3H, 3.54 s 3H, 3.24 - 3.45 m 2H, 2.81 bs 2H, 2.44 bs 4H.	V

- 47 -

5-{4-[2-(diethylamino)ethoxy]phenyl}-2-[[pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide	120	524	1.21 (t, 6H), 2.00 (m, 1H), 2.20 (m, 1H), 3.18 (m, 4H), 3.43 (m, 6H), 4.31 (t, 2H), 4.53 (m, 1H), 7.05 (d, 2H), 7.52 (d, 2H), 7.70 (s, 1H), 8.30 (m, 3H), 8.78 (s, 1H), 8.85 (s, 1H), 8.98 (s, 1H), 9.32 (s, 1H), 10.90 (s, 1H)	VI
5-{3-[2-(diethylamino)ethoxy]phenyl}-2-[[pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide hydrochloride	121	524	1.25 (t, 6H), 2.02 (m, 1H), 2.20 (m, 1H), 3.22 (m, 4H), 3.43 (m, 6H), 4.34 (t, 2H), 4.53 (m, 1H), 6.92 (d, 1H), 7.12 (s, 1H), 7.28 (d, 1H), 7.39 (dxd, 1H), 7.84 (s, 1H), 8.32 (m, 3H), 8.78 (s, 1H), 8.85 (s, 1H), 8.98 (s, 1H), 9.30 (s, 1H), 10.92 (s, 1H)	VI
5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-yl-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide	122	538	0.95 (t, 6H), 1.52 (m, 2H), 1.62 (m, 4H), 2.52 (m, 4H), 2.78 (m, 4H), 3.00 (m, 12H), 3.18 (t, 2H), 4.05 (t, 2H), 6.68 (bs, 1H), 6.80 (d, 1H), 7.10 (m, 2H), 7.28 (m, 2H), 7.82 (s, 1H), 8.22 (s, 1H), 8.28 (s, 1H), 9.00 (s, 1H)	VI
N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide hydrochloride	123	467	δ 12.6, br s, 1H; δ 10.9, s, 1H; δ 9.55, br s, 1H; δ 9.24, br s, 1H; δ 8.88, s, 1H; δ 8.49, d, 1H; δ 8.35, dd, 1H; δ 8.29, d, 1H; δ 7.92, s, 1H; δ 7.54, d, 2H; δ 6.99, d, 2H; δ 4.42, m, 1H; δ 3.33, m, 1H; δ 3.23, m, 2H; δ 3.10, m, 1H; δ 2.02, m, 1H; δ 1.85, m, 4H; δ 1.62, m, 1H	VI
5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-yl-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide hydrochloride	124	538	0.95 (t, 6H), 1.52 (m, 2H), 1.62 (m, 4H), 2.52 (m, 4H), 2.78 (m, 4H), 3.00 (m, 12H), 3.18 (t, 2H), 4.05 (t, 2H), 6.68 (bs, 1H), 6.80 (d, 1H), 7.10 (m, 2H), 7.28 (m, 2H), 7.82 (s, 1H), 8.22 (s, 1H), 8.28 (s, 1H), 9.00 (s, 1H)	VI
N-(2-aminoethyl)-5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide	125	413	8.53 s 1H, 8.31-8.33 m 1H, 8.13 d J=2.8 Hz 1H, 7.42 d J=8.8 Hz 2H, 7.38 s 1H, 6.85 d J=8.8 Hz 2H, 3.72 s 3H.	VI
5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-yl-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide	126	538	0.95 (t, 6H), 1.52 (m, 2H), 1.79 (m, 2H), 2.42 (m, 4H), 2.70 (m, 4H), 3.08 (m, 2H), 3.98 (m, 3H), 6.95 (d, 2H), 7.48 (d, 2H), 7.70 (s, 1H), 8.28 (d, 2H), 8.95 (s, 1H)	VI







- 48 -

5-(4-methoxyphenyl)-N-piperidin-4-yl-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide	127	453		VI
tert-butyl 3-[[5-{3-[2-(diethylamino)ethoxy]phenyl}-2-[[pyrazin-2-ylamino)carbonyl]amino}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate trifluoroacetate	128	638	1.21 (t, 6H), 1.32 (s, 9H), 1.75 (m, 1H), 1.92 (m, 1H), 2.85 (m, 1H), 3.20 (m, 4H), 3.50 (t, 2H), 3.80 (m, 2H), 4.35(t,2H), 6.90 (d, 1H), 7.15 (s, 1H) 7.25 (d, 1H), 7.38 (dxd, 1H), 7.76 (s, 1H), 7.98 (bs, 1H), 8.30 (d, 2H), 8.90 (s, 1H), 9.25 (bs, 1H), 10.92 (s, 1H)	VI
5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-yl-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide	129	538	0.95 (t, 6H), 1.52 (m, 2H), 1.79 (m, 2H), 2.42 (m, 4H), 2.70 (m, 4H), 3.08 (m, 2H), 3.98 (m, 3H), 6.95 (d, 2H), 7.48 (d, 2H), 7.70 (s, 1H), 8.28 (d, 2H), 8.95 (s, 1H)	VI
5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide	130	439		VI
N-[3-(1,4-diazepan-1-ylcarbonyl)-5-(4-methoxyphenyl)-2-thienyl]-N'-pyrazin-2-ylurea	131	453	8.47 s 1H, 8.14-8.25 m 2H, 7.43 d J=8.8 Hz 2H, 7.06 s 1H, 7.85 d J=8.8 Hz 2H, 3.90 bs 2H, 3.79 t J=6.0 Hz 2H, 3.72 s 3H, 3.40 bs 2H, 2.10 bs 2H.	VI
N-[3-[(3-aminopyrrolidin-1-yl)carbonyl]-5-(4-methoxyphenyl)-2-thienyl]-N'-pyrazin-2-ylurea	132	439	11.97 bs 1H, 10.72 s 1H, 8.79 s 1H, 8.22-8.26 m 2H, 7.50 d J=8.6 Hz 2H, 8.04 bs 2H, 7.31 s 1H, 7.20-7.25 m 2H, 7.08 d J=8.3 Hz 1H, 7.00 t J=7.2 Hz 1H, 6.92 d J=8.6 Hz 2H 3.72 s 3H, 2.11-2.19 m 1H, 3.80 bs 2H, 3.64 bs 2H.	VI
tert-butyl 4-[[5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino)carbonyl]amino}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate	133	553		VI
tert-butyl 3-[[5-{4-[2-(diethylamino)ethoxy]phenyl}-2-[[pyrazin-2-ylamino)carbonyl]amino}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate	134	638	1.18 (t, 6H), 1.32 (s, 9H), 1.72 (m, 1H), 1.90 (m, 1H), 2.85 (m, 1H), 3.21 (m, 4H), 3.50 (t, 2H), 3.80 (m, 2H), 4.30 (t,2H), 7.05 (d, 2H), 7.52 (d, 2H), 7.76 (s, 1H), 7.92 (bs, 1H), 8.30 (d, 2H), 8.90 (s, 1H), 9.25 (bs, 1H), 10.90 (s, 1H)	VI








- 49 -

5-[4-(2-diethylamino-ethoxy)-phenyl]-2-(3-hydroxy-urea)-thiophene-3-carboxylic acid-(S)-piperidin-3-ylamide	136	476	δ 1.17 - 1.40 (m, 6 H) 1.54 - 1.81 (m, 2 H) 1.84 - 2.07 (m, 2 H) 2.76 - 2.99 (m, 2 H) 3.05 - 3.25 (m, 4 H) 3.41 - 3.61 (m, 4 H) 3.73 - 3.88 (m, 1 H) 4.29 - 4.42 (m, 1 H) 4.40 - 4.64 (m, 1 H) 5.86 (s, 1 H) 7.07 (d, $J=8.48$ Hz, 2 H) 7.55 (d, $J=8.48$ Hz, 2 H) 7.82 (s, 1 H) 9.44 (s, 1 H) 9.65 (s, 1 H);	VII
2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(3-methoxyphenyl)thiophene-3-carboxamide	137	389	10.90 (s, 1H), 9.21 (s, 1H), 9.01 (s, 1H), 8.29 (d, 1H), 7.93 (s, 1H), 7.29 (t, 1H), 7.11 (d, 2H), 7.03 (s, 1H), 6.83 (d, 1H), 4.33 (s, 1H), 3.81 (s, 3H), 3.20 (m, 4H), 2.00 (m, 1H), 1.84 (m, 4H), 1.56 (m, 1H).	VIII
2-[(aminocarbonyl)amino]-5-(2-hydroxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide	138	361	δ 10.85 (s, 1H), 10.13 (s, 1H), 9.29 (d, 2H), 8.31 (d, 1H), 7.92 (s, 1H), 7.63 (dd, 1H), 7.05 (td, 1H), 6.94 (dd, 1H), 6.82 (td, 1H), 6.91 (brs, 2H), 4.21 (brs, 1H), 3.26 (dd, 2H), 2.89 (m, 2H), 1.88-1.618 (m, 4H).	VIII
2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide	139	375	10.94 (s, 1H), 9.41 (s, 1H), 9.19 (s, 1H), 8.47 (d, 1H), 8.11 (s, 1H), 7.30 (t, 1H), 7.15 (d, 1H), 7.05 (br, 2H), 6.84 (d, 1H), 4.25 (s, 1H), 3.82 (s, 3H), 3.29 (d, 1H), 3.12 (d, 1H), 2.97 (m, 2H), 1.92 (d, 2H), 1.69 (m, 2H).	VIII
2-[(aminocarbonyl)amino]-5-[2-(benzyloxy)phenyl]-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide	140	451	δ 10.8 (s, 1H), 8.63 (brs, 2H), 8.04 (d, 1H), 7.79 (s, 1H), 7.63 (d, 1H), 7.55 (d, 2H), 7.38 (t, 2H), 7.31 (t, 1H), 7.23 (t, 1H), 7.20 (t, 1H), 7.02 (t, 1H), 6.95 (m, 2H), 5.28 (s, 2H), 4.15 (m, 1H), 3.24 (m, 1H), 2.85 (m, 2H), 1.90-1.58 (m, 5H).	VIII








Table 2

Structure	IUPAC Name	M.W. (g/mol)	Ex.
	tert-butyl 3-[[2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate trifluoroacetate	559.7	1
	2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-ylthiophene-3-carboxamide trifluoroacetate	459.6	2
	2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-ylthiophene-3-carboxamide trifluoroacetate	459.6	3
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide	374.5	4
	tert-butyl 3-[[2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate trifluoroacetate	559.7	5
	2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-ylthiophene-3-carboxamide	459.6	6








- 51 -

	2-[(aminocarbonyl)amino]-N-[(3R)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide	388.5	7
	N-(3-[(4-aminopiperidin-1-yl)carbonyl]-5-{4-[2-(diethylamino)ethoxy]phenyl}-2-thienyl)urea trifluoroacetate	459.6	8
	2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[3-(hydroxymethyl)phenyl]thiophene-3-carboxamide trifluoroacetate (salt)	482.6	9
	2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-ylthiophene-3-carboxamide trifluoroacetate	459.6	10
	2-[(aminocarbonyl)amino]-N-(2-aminoethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	334.4	11
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	374.5	12
	2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-pyridin-3-ylthiophene-3-carboxamide trifluoroacetate	453.6	13








- 52 -

	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(1-methylpiperidin-4-yl)thiophene-3-carboxamide	388.5	14
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-1-methylazepan-3-yl]thiophene-3-carboxamide hydrochloride	402.5	15
	2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-[3-(hydroxymethyl)phenyl]thiophene-3-carboxamide trifluoroacetate (salt)	482.6	16
	2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-pyrrolidin-3-ylthiophene-3-carboxamide trifluoroacetate	445.6	17
	2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-pyridin-3-ylthiophene-3-carboxamide trifluoroacetate	453.6	18
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-1-methylpiperidin-3-yl]thiophene-3-carboxamide hydrochloride	388.5	19
	2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-pyrrolidin-3-ylthiophene-3-carboxamide trifluoroacetate	445.6	20








- 53 -

	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-ylmethyl]thiophene-3-carboxamide	388.5	21
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide	360.4	22
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]thiophene-3-carboxamide	360.4	23
	2-[(aminocarbonyl)amino]-N-[2-(dimethylamino)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide	362.4	24
	2-[(aminocarbonyl)amino]-N-[2-(diethylamino)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide	390.5	25
	2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide hydrochloride	388.5	26
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-yl]thiophene-3-carboxamide	374.5	27








- 54 -

	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(piperidin-4-ylmethyl)thiophene-3-carboxamide	388.5	28
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-pyrrolidin-3-ylthiophene-3-carboxamide	360.4	29
	2-[(aminocarbonyl)amino]-N-(1-ethylpiperidin-3-yl)-5-(4-methoxyphenyl)thiophene-3-carboxamide hydrochloride	402.5	30
	2-[(aminocarbonyl)amino]-N-[(3S)-1-ethylazepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide hydrochloride	416.5	31
	2-[(aminocarbonyl)amino]-5-(3-hydroxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	360.4	32
	2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	360.4	33
	2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	374.5	34








- 55 -

	tert-butyl (3S)-3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)pyrrolidine-1-carboxylate	460.6	35
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-piperidin-3-ylthiophene-3-carboxamide	374.5	36
	2-[(aminocarbonyl)amino]-N-(1-benzylpiperidin-4-yl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	464.6	37
	tert-butyl 3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate	474.6	38
	2-[(aminocarbonyl)amino]-5-[4-(2-piperidin-1-ylethoxy)phenyl]-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide	493.6	39
	2-[(aminocarbonyl)amino]-5-[4-(2-piperidin-1-ylethoxy)phenyl]-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide	450.6	40
	2-[(aminocarbonyl)amino]-N-azetidin-3-yl-5-(4-methoxyphenyl)thiophene-3-carboxamide	346.4	41








- 56 -

	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(2S)-pyrrolidin-2-ylmethyl]thiophene-3-carboxamide	374.5	42
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-pyridin-4-ylthiophene-3-carboxamide	368.4	43
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperazin-1-ylethyl)thiophene-3-carboxamide	403.5	44
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperidin-1-ylethyl)thiophene-3-carboxamide	402.5	45
	2-[(aminocarbonyl)amino]-N-1-azabicyclo[2.2.2]oct-3-yl-5-(4-methoxyphenyl)thiophene-3-carboxamide	400.5	46
	2-[(aminocarbonyl)amino]-N-(2-hydroxyethyl)-5-(4-hydroxyphenyl)thiophene-3-carboxamide	321.4	48
	2-[(aminocarbonyl)amino]-N-(trans-4-hydroxycyclohexyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	389.5	49








- 57 -

	2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide	382.4	50
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperazin-1-ylethyl)thiophene-3-carboxamide	524.6	52
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide	396.5	54
	2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-pyridin-3-ylethyl)thiophene-3-carboxamide	382.4	55
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-3-ylethyl)thiophene-3-carboxamide	396.5	56
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)thiophene-3-carboxamide	430.6	58
	2-[(aminocarbonyl)amino]-5-(2-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide	374.5	59








- 58 -








	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(tetrahydrofuran-2-ylmethyl)thiophene-3-carboxamide	375.4	60
	tert-butyl (3R)-3-([2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl)amino)piperidine-1-carboxylate	474.6	62
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-3-ylmethyl)thiophene-3-carboxamide	382.4	63
	tert-butyl 3-([2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl)amino)azetidine-1-carboxylate	446.5	64
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-4-ylmethyl)thiophene-3-carboxamide	382.4	65
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(3-methoxypropyl)thiophene-3-carboxamide	363.4	67
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[2-(2-thienyl)ethyl]thiophene-3-carboxamide	401.5	68








- 59 -







	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-thienylmethyl)thiophene-3-carboxamide	387.5	69
	N-[3-(1,4-diazepan-1-ylcarbonyl)-5-(4-methoxyphenyl)-2-thienyl]urea	374.5	70
	2-[(aminocarbonyl)amino]-N-(2-methoxyethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	349.4	71
	2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-thienylmethyl)thiophene-3-carboxamide	373.5	72
	2-[(aminocarbonyl)amino]-N-{2-[(2-furylmethyl)thio]ethyl}-5-(4-methoxyphenyl)thiophene-3-carboxamide	431.5	74
	2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-[2-(2-thienyl)ethyl]thiophene-3-carboxamide	387.5	75
	N-(3-[(4-aminopiperidin-1-yl)carbonyl]-5-{3-[2-(diethylamino)ethoxy]phenyl}-2-thienyl)urea trifluoroacetate	459.6	76

- 60 -







	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-ylmethyl]thiophene-3-carboxamide	388.5	77
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(1,2,3,4-tetrahydroquinolin-3-yl)thiophene-3-carboxamide	422.5	78
	2-[(aminocarbonyl)amino]-N-(1,3-benzodioxol-5-ylmethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	425.5	79
	2-[(aminocarbonyl)amino]-N-(3-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	411.5	80
	2-[(aminocarbonyl)amino]-N-[2-(3,4-dimethoxyphenyl)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide	455.5	81
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(5-methyl-2-furyl)methyl]thiophene-3-carboxamide	385.4	84
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-2-ylmethyl)thiophene-3-carboxamide	382.4	85







	2-[(aminocarbonyl)amino]-N-(4-fluorobenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	399.4	86
	tert-butyl 4-([2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl)amino)piperidine-1-carboxylate	474.6	88
	2-[(aminocarbonyl)amino]-N-(2-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	411.5	89
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-phenoxyethyl)thiophene-3-carboxamide	411.5	90
	2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-2-ylethyl)thiophene-3-carboxamide	396.5	93
	tert-butyl 4-([2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl)amino)piperidine-1-carboxylate	474.6	94
	2-[(aminocarbonyl)amino]-N-(4-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide	411.5	95

	2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide trifluoroacetate	459.6	110
	2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3R)-piperidin-3-yl]thiophene-3-carboxamide trifluoroacetate	459.6	111
	tert-butyl (3S)-3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate trifluoroacetate	559.7	112
	2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-{4-[2-(diethylamino)ethoxy]phenyl}thiophene-3-carboxamide hydrochloride	473.6	113
	tert-butyl (3R)-3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate trifluoroacetate	559.7	114
	N-[3-[(3S)-3-aminoazepan-1-yl]carbonyl]-5-(4-methoxyphenyl)-2-thienyl]urea hydrochloride	388.5	115
	5-{4-[2-(diethylamino)ethoxy]phenyl}-2-[[[(pyrazin-2-ylamino)carbonyl]amino]-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide	523.7	120


	5-{3-[2-(diethylamino)ethoxy]phenyl}-2-[[pyrazin-2-ylamino]carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide hydrochloride	523.7	121
	5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-yl-2-[[pyrazin-2-ylamino]carbonyl]amino}thiophene-3-carboxamide	537.7	122
	N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino]carbonyl]amino}thiophene-3-carboxamide hydrochloride	466.6	123
	5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-yl-2-[[pyrazin-2-ylamino]carbonyl]amino}thiophene-3-carboxamide hydrochloride	537.7	124
	N-(2-aminoethyl)-5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino]carbonyl]amino}thiophene-3-carboxamide	412.5	125
	5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-yl-2-[[pyrazin-2-ylamino]carbonyl]amino}thiophene-3-carboxamide	537.7	126

- 64 -

	5-(4-methoxyphenyl)-N-piperidin-4-yl-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide	452.5	127
	tert-butyl 3-[[5-{3-[2-(diethylamino)ethoxy]phenyl}-2-[[pyrazin-2-ylamino)carbonyl]amino}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate trifluoroacetate	637.8	128
	5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-yl-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide	537.7	129
	5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide	438.5	130
	N-[3-(1,4-diazepan-1-ylcarbonyl)-5-(4-methoxyphenyl)-2-thienyl]-N'-pyrazin-2-ylurea	452.5	131
	N-[3-[(3-aminopyrrolidin-1-yl)carbonyl]-5-(4-methoxyphenyl)-2-thienyl]-N'-pyrazin-2-ylurea	438.5	132

	tert-butyl 4-[[[(5-(4-methoxyphenyl)-2-[(pyrazin-2-ylamino)carbonyl]amino)-3-thienyl)carbonyl]amino]piperidine-1-carboxylate	552.6	133
	tert-butyl 3-[[[(5-{4-[2-(diethylamino)ethoxy]phenyl}-2-[(pyrazin-2-ylamino)carbonyl]amino)-3-thienyl)carbonyl]amino]piperidine-1-carboxylate	637.8	134
	5-[4-(2-diethylamino-ethoxy)-phenyl]-2-(3-hydroxy-urea)-thiophene-3-carboxylic acid-(S)-piperidin-3-ylamide	475.2	136
	2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(3-methoxyphenyl)thiophene-3-carboxamide	388.5	137
	2-[(aminocarbonyl)amino]-5-(2-hydroxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide	360.4	138
	2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide	374.5	139

- 66 -

	2-[(aminocarbonyl)amino]-5-[2-(benzyloxy)phenyl]-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide	450.6	140
---	---	---------------------	-------------------

Example 4**2-[(Aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide**

(4-Methoxy-phenyl)-acetaldehyde. To a stirred solution of (4-Methoxy-phenyl)-acetic acid methyl ester (18.0 g, 100 mmol) in anhydrous toluene (200 mL) cooled to -78°C under N_2 was added diisobutylaluminum hydride (DIBAL, 1.0 M in toluene, 150 mL, 150 mmol) over a period of 10-15 minutes. The mixture was stirred at -78°C for an additional 2h. The reaction was quenched by the slow addition of MeOH, followed by the introduction of 10% Rochelle's Salt. The suspension was diluted with EtOAc and stirred at room temperature for 1h. The EtOAc layer was set aside and the aqueous layer was extracted with EtOAc(2x). The combined organic layers were combined and dried over Na_2SO_4 and filtered. The solution was concentrated under vacuum to yield 12.0 g (100%) of the title aldehyde as a yellow viscous semisolid, which was used in the next step without purification. LC/MS (APCI, ES, $\text{M}+\text{H}=151$).

2-Amino-5-(4-methoxy-phenyl)-thiophene-3-carboxylic acid methyl ester. To a solution of 4-methoxyphenylacetaldehyde (12.0g) in DMF (200mL) was added cyanomethyl acetate (8.9 mL, 100 mmol) and sulfur (3.2g, 100mmol), followed by diisopropylethylamine (Hunig's Base, 17.4 mL, 100 mmol). The resultant suspension immediately turned dark yellow to brown with an exotherm. The reaction mixture was stirred overnight at room temperature. The reaction was slowly added to water (~1L) while stirring. A tan precipitate formed and was filtered after an additional 30 minutes of stirring. The resultant solid was purified by column chromatography (SiO_2 , 10-20% EtOAc/ hexanes) to yield 15.3 g (58%) of the title compound as a light yellow

- 67 -

solid. ^1H NMR (d_6 -DMSO δ 7.41, br s, 2H; δ 7.37, d, 2H; δ 7.07, s, 1H; δ 6.90, d, 2H; δ 3.75, s, 3H; δ 3.72, s, 3H), LC/MS (APCI, ES, $\text{M}+\text{H}=264$).

Methyl 5-(4-methoxyphenyl)-2-([(trichloroacetyl)amino]carbonyl)amino)thiophene-3-

5 **carboxylate.** To a stirred solution of 2-Amino-5-(4-methoxy-phenyl)-thiophene-3-carboxylic acid methyl ester (7.15 g, 27.2 mmol) in anhydrous THF (150 mL) was added trichloroacetyl isocyanate (6.4 mL, 54 mmol) slowly over a period of 5 min. After the addition was complete, a precipitate formed and the reaction stirred for an additional 1h. The desired product was obtained by filtration to give 6.9 g (56%) an off-white solid. The product was used in the next step
10 without purification ^1H NMR (d_6 -DMSO δ 12.3, br s, 1H; δ 12.2, s, 1H; δ 7.46, d, 2H; δ 7.32, s, 1H; δ 6.85, d, 2H; δ 3.75, s, 3H; δ 3.66, s, 3H), LC/MS (APCI, ES, $\text{M}+\text{H}=451$).

***tert*-Butyl (3S)-3-([2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-**

thienyl]carbonyl)amino)piperidine-1-carboxylate. To a solution of methyl 5-(4-
15 methoxyphenyl)-2-([(trichloroacetyl)amino]carbonyl)amino)thiophene-3-carboxylate (1.0 g, 2.2 mmol) in dry THF (20 mL) was added a solution of $[\text{Me}_2\text{Al}-3\text{-Boc-(S)-3-aminopiperidine}]$ in THF (which was preformed by the addition of Me_3Al (2.0 M in hexanes, 2.2 mL, 4.4 mmol) to a solution of (S)-3-Amino-piperidine-1-carboxylic acid *tert*-butyl ester (0.89 g, 4.4 mmol) in 10 mL THF at -78°C followed by warming to room temperature for an additional 15 min). The
20 resulting orange-colored solution was stirred overnight at room temperature. The reaction mixture was cooled with ice and a 10% aqueous solution of Rochelle's salt was added slowly to quench the reaction. The resulting biphasic solution was warmed to room temperature and stirred for an additional 1h. The mixture was diluted with EtOAc and H_2O , the aqueous layer was extracted with EtOAc (3x) and the combined organic extracts were washed with H_2O , brine and
25 dried (Na_2SO_4). Evaporation gave a pale orange solid. Purification by column chromatography (SiO_2 , 50 % EtOAc/hexanes) gave 0.70 g (67%) of a light yellow solid. LC/MS (APCI, ES, $\text{M}+\text{H}=475$).

2-[(Aminocarbonyl)amino]-5-(4-methoxyphenyl)-*N*-[(3S)-piperidin-3-yl]thiophene-3-

30 **carboxamide; hydrochloride.** To a stirred solution of *tert*-butyl (3S)-3-([2-

- 68 -

[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl} amino)piperidine-1-carboxylate (0.70 g, 1.47 mmol) in anhydrous MeOH (5.0 mL) was added 4.0N HCl in 1, 4-dioxane (10 mL). A small amount of precipitate forms shortly and the reaction is stirred for an additional 4h at room temperature. The solvent was removed under vacuum. The residue was redissolved in methanol and concentrated under vacuum (2x) to yield 0.51 g (85%) of a light yellow solid. ¹H NMR (d₆-DMSO δ 10.9, s, 1H; δ 9.39, br s, 1H; δ 9.20, br s, 1H; δ 8.37, d, 1H; δ 7.88, s, 1H; δ 7.49, d, 2H; δ 6.96, d, 2H; δ 6.97, br s, 2H; δ 4.24, m, 1H; δ 3.77, s, 3H; δ 3.29, m, 1H; δ 3.11, m, 1H; δ 2.93, m, 2H; δ 1.91, m, 2H; δ 1.68, m, 2H), LC/MS (APCI, ES, M+H=6875).

Example 26

2-[(Aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide hydrochloride

(S)-3-Amino-azepane-1-carboxylic acid tert-butyl ester. (S)-Azepan-3-ylamine (5g; 43.8 mmol) was dissolved in 100 mL of anhydrous CH₂Cl₂ and cooled to -78 °C while stirring with a magnetic stirring bar. In another flask N-(tert-butoxycarbonyloxy)succinimide [Boc-OSu] (9.7 g; 45 mmol) was dissolved in 50 mL of anhydrous CH₂Cl₂. To the stirred solution of the amine was added the solution of the succinimide over a period of 10-15 minutes so as to keep the reaction mixture at -78 °C while stirring. After the addition was complete, the reaction was allowed to warm to room temperature and then stirred for an additional 4h or until the reaction was complete by TLC (Ninhydrin; R_f 0.3; 0.1:1:10 NH₄OH, MeOH; CH₂Cl₂). The reaction mixture was washed with 50 mL of H₂O. The aqueous layer was brought to a pH >13 by the addition of 6N NaOH and extracted with CH₂Cl₂ (3 x 100 mL). The organic layer was dried over Na₂CO₃, filtered, and concentrated in a vacuum to yield pure (S)-3-amino-azepane-1-carboxylic acid tert-butyl ester as a viscous oil (5.1 g, 54%). ¹H NMR (d₆-DMSO, d 3.4, m, 2H; d 2.89, m, 1H; d 2.71, m, 1H; d 2.54, m, 1H; d 1.54, m, 3H; d 1.34, m, 3H; d 1.27, s, 9H; d 1.12, m, 2H), LC/MS (APCI, ES, M+H=215).

- 69 -

***tert*-Butyl (3*S*)-3-([2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-**

thienyl]carbonyl}amino)azepane-1-carboxylate. To a solution of methyl 5-(4-

methoxyphenyl)-2-([[(trichloroacetyl)amino]carbonyl}amino)thiophene-3-carboxylate (1.36 g, 3 mmol) in anhydrous THF (20 mL) was added a solution of [Me₂Al-3-Boc-(*S*)-3-

5 aminohomopiperidine] in THF (preformed by the careful addition of Me₃Al (2.0 M in hexanes, 3.0 mL, 6.0 mmol) to a solution of (*S*)-3-amino-azepane-1-carboxylic acid *tert*-butyl ester in 10 mL of THF at -78 °C followed by warming to room temperature under nitrogen and stirring for an additional 15 min.). The resulting deep yellow/orange solution was stirred overnight at room

temperature. The reaction mixture was cooled with ice and a 10% aqueous solution of Rochelle's
10 salt was added slowly to quench the reaction. The resulting biphasic solution was warmed to room temperature and stirred for an additional 1h. The mixture was diluted with EtOAc and H₂O, the aqueous layer was extracted with EtOAc (3x) and the combined organic extracts were washed with H₂O, brine and dried (Na₂SO₄). Evaporation gave a pale orange solid. Purification by ISCO MPLC (SiO₂, 60-80% EtOAc/hexanes) gave 0.9 g (62%) of the title compound as a
15 white solid. ¹H NMR (d₆-DMSO, δ 11.0, s, 1H; δ 7.95, d, 0.5H; δ 7.81, d, 0.5H; δ 7.65, s, 0.5H; δ 7.56, s, 0.5H; δ 7.46, d, 2H; δ 6.97, d, 2H; δ 6.96, br s, 2H; δ 4.19, m, 0.5H; δ 4.11, m, 0.5H; δ 3.77, m, 3H; δ 3.65, m, 1H; δ 3.48, m, 1H; δ 3.20, m, 2H; δ 1.75, m, 3H; δ 1.58, m, 2H; δ 1.42, s, 4.5H; δ 1.39, m, 1H; δ 1.36, s, 4.5H), LC/MS (APCI, ES, M+H=489).

20 **2-[(Aminocarbonyl)amino]-*N*-[(3*S*)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide; hydrochloride.** To a stirred solution of *tert*-butyl (3*S*)-3-([2-

[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)azepane-1-carboxylate (0.9g, 1.8 mmol) in 1, 4-dioxane (10 mL) was added 4.0N HCl in 1, 4-dioxane (10 mL, 40 mmol). A precipitate forms shortly and the reaction is stirred for an additional 4h at room

25 temperature. Due to the hygroscopic nature of the salt form, the solvent was removed under vacuum. The residue was dissolved in methanol and concentrated under vacuum (2x) to yield and off-white solid. Recrystallization from using 2-propanol gave 0.45g (59%) of a white solid.

¹H NMR (d₆-DMSO, δ 10.9, s, 1H; δ 9.58, br s, 1H; δ 9.29, br s, 1H; δ 8.39, d, 1H; δ 7.82, s, 1H; δ 7.48, d, 2H; δ 6.96, d, 2H; δ 4.36, m, 1H; δ 3.77, s, 3H; δ 3.29, m, 1H; δ 3.20, m, 2H; δ 3.07, m, 1H; δ 1.98, m, 1H; δ 1.84, m, 4H; δ 1.59, m, 1H), LC/MS (APCI, ES, M+H=389).
30

- 70 -

Example 110

2-[(Aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide trifluoroacetate

To a stirred solution of *tert*-butyl (3S)-3-({[2-
 5 [(aminocarbonyl) amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl} amino)piperidine-1-carboxylate dissolved in a small amount of methanol was added 4.0 N HCl in dioxane. The solution was stirred for 1h at RT. The product was purified by Gilson (5%MeCN-H₂O→98%MeCN-H₂O) to yield 27 mg of the title compound as the trifluoroacetate salt. ¹H NMR (300 MHz, d₃-MeOD; δ 7.55(d, 2H), 7.45(s, 1H), 7.05(d, 2H), 4.35(dd, 2H), 4.25(m, 1H),
 10 3.60(dd, 2H), 3.50(m, 1H), 3.30(m, 5H), 2.95(dd, 2H), 2.10(dd, 2H), 1.80(m, 2H), 1.35(t, 6H)), LCMS, (ES, M+H=460).

Example 112

tert-Butyl (3S)-3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl]carbonyl]amino}piperidine-1-carboxylate trifluoroacetate

Methyl {4-[2-(diethylamino)ethoxy]phenyl}acetate To a solution of methyl (4-hydroxyphenyl)acetate (16.6 g, 10 mmol) in DMF (100 mL) was added 2-bromo-*N,N*-diethylethanamine hydrobromide (2.6 g, 10 mmol) and), Cs₂CO₃ (6.6g, 20 mmol). After 1h an
 20 additional equivalent of the bromide was added and then stirred overnight at room temperature. The reaction mixture was poured in to a large volume of cold water. The product was then isolated by filtration and purified by column chromatography (SiO₂, 10% MeOH/DCM) to give 15.6 g of the title compound as an off-white solid. LC/MS (APCI, ES, M+H=266).

{4-[2-(Diethylamino)ethoxy]phenyl}acetaldehyde To a stirred solution of methyl {4-[2-(diethylamino)ethoxy]phenyl}acetate (5.3 g, 20 mmol) in anhydrous toluene (100 mL)-cooled to -78 °C under N₂ was added diisobutylaluminum hydride (DIBAL, 1.0M in toluene, 100 mL, 100 mmol) over a period of 10-15 minutes. The mixture was stirred at -78 °C for an additional 2h. The reaction was quenched by the slow addition of MeOH, followed by the introduction of 10%
 30 Rochelle's Salt. The suspension was diluted with EtOAc and stirred at room temperature for 1h. The EtOAc layer was set aside and the aqueous layer was extracted with EtOAc(2x). The

- 71 -

combined organic layers were combined and dried over Na₂SO₄ and filtered. The solution was concentrated under vacuum to yield 4.7 g (100%) of the title aldehyde as a yellow viscous semisolid, which was used in the next step without purification. LC/MS (APCI, ES, M+H=236).

- 5 **Methyl 2-amino-5-{4-[2-(diethylamino)ethoxy]phenyl}thiophene-3-carboxylate** To a solution of {4-[2-(diethylamino)ethoxy]phenyl}acetaldehyde (4.7 g, 20mmol) in DMF (30 mL) was added cyanomethyl acetate (1.5 mL, 20 mmol) and sulfur (0.6 g, 20mmol), followed by diisopropylethylamine (Hunig's Base, 2.5 mL, 20 mmol). The resultant suspension immediately turned dark yellow to brown with an exotherm. The reaction mixture was stirred overnight at
10 room temperature. The reaction was slowly added to water (~200 mL) while stirring. A tan precipitate formed and was filtered after an additional 30 minutes of stirring. The resultant solid was purified by column chromatography (SiO₂, 5-10% MeOH/DCM/0.5% NH₄OH) to yield 2.4 g of the title compound as a light yellow solid. LC/MS (APCI, ES, M+H=349).
- 15 **2-Amino-5-{4-[2-(diethylamino)ethoxy]phenyl}thiophene-3-carboxylic acid** To a stirred solution of methyl 2-amino-5-{4-[2-(diethylamino)ethoxy]phenyl}thiophene-3-carboxylate (2.0 g, 5.7 mmol) in MeOH (50 mL) was added 6N NaOH (50 mL) and water (50 mL). The reaction was heated to reflux for 2h or until starting material was gone by TLC or LCMS. The solution was concentrated under vacuum to about half of the original volume. The pH of the resultant
20 cloudy mixture was adjusted to 3-5 by the careful addition of 6N HCl (~150 mL) while stirring. The gummy red precipitate was filtered and dried. Purification was achieved by triturating in boiling hexanes. The product (1.6 g) was isolated in pure form by filtration after cooling to room temperature and drying in a vacuum oven overnight. LC/MS (APCI, ES, M+H=335).
- 25 **tert-Butyl (3S)-3-{[(2-amino-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl) carbonyl] amino}piperidine-1-carboxylate** To a stirred solution of 2-amino-5-{4-[2-(diethylamino)ethoxy]phenyl}thiophene-3-carboxylic acid (100 mg, 0.3 mmol) in anhydrous DMF (2.0 mL) is added (S)-3-amino-azepane-1-carboxylic acid tert-butyl ester (60 mg, 0.3 mmol), EDCI (63 mg, 0.33 mmol), HOBt (61 mg, 0.45 mmol), and NMM (0.04 mL, 0.3 mmol).
30 The reaction mixture was stirred overnight at room temperature. The solution was diluted with

- 72 -

water and EtOAc. The organic layer was separated and set aside. The remaining aqueous layer was extracted with EtOAc (2x) and then the combined organic extracts were pooled and washed with brine. The resultant EtOAc solution was dried over Na₂SO₄, filtered, and concentrated under vacuum to yield a brown solid. Purification was performed by Gilson (5%MeCN-H₂O→98%MeCN-H₂O) to give 90 mg of an off-white solid. LC/MS (APCI, ES, M+H=517).

***tert*-Butyl (3*S*)-3-[[2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate** To a stirred solution of *tert*-butyl (3*S*)-3-[[2-amino-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl) carbonyl]amino}piperidine-1-carboxylate (90 mg, 0.17 mmol) in anhydrous THF (5.0 mL) was added trichloroacetyl isocyanate (0.09 mL, 0.7 mmol) slowly over a period of 5 min. After the addition was complete, a precipitate formed and the reaction stirred for an additional 1h. The reaction mixture was concentrated in a vacuum. The crude residue was dissolved in methanol and then charged with 2.0N NH₃ in methanol (0.35 mL). Purification by Gilson (5%MeCN-H₂O→98%MeCN-H₂O) gave the title (50 mg) as a tan solid. ¹H NMR (300 MHz, d₃-MeOD; δ 7.55(d, 2H), 7.45(s, 1H), 7.05(d, 2H), 4.35(dd, 2H), 3.60-3.90(m, 3H), 3.60(dd, 2H), 3.30(m, 4H), 2.95(m, 2H), 1.90(dd, 2H), 1.55(m, 2H), 1.45(s, 9H), 1.35(t, 6H)), (APCI, ES, M+H=560).

Example 123.....

N-[(3*S*)-Azepan-3-yl]-5-(4-methoxyphenyl)-2-[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide hydrochloride

2-Amino-5-(4-methoxy-phenyl)-thiophene-3-carboxylic acid. To a stirred solution of 2-amino-5-(4-methoxy-phenyl)-thiophene-3-carboxylic acid methyl ester (6.6 g, 26.6 mmol) in MeOH (200 mL) was added 6N NaOH (100 mL) and water (50 mL). The reaction was heated to reflux for 2h or until starting material was gone by TLC or LCMS. The solution was concentrated under vacuum to about half of the original volume. The pH of the resultant cloudy mixture was adjusted to 3-5 by the careful addition of 6N HCl (~150 mL) while stirring. The gummy red precipitate was filtered and dried. Purification was achieved by triturating in boiling hexanes. The product (6.0 g, 91%) was isolated in pure form by filtration after cooling to room

- 73 -

temperature and drying in a vacuum oven overnight. ^1H NMR (d_6 -DMSO δ 7.37, d, 2H; δ 7.11, s, 1H; δ 7.10, br s, 2H; δ 6.94, d, 2H), LC/MS (APCI, ES, $M+H=250$).

Pyrazine-2-carboxylic acid hydrazide. To a stirred solution of pyrazine-2-carboxylic acid methyl ester (11.1 g, 80 mmol) in 140 mL of EtOH was added hydrazine hydrate (15.6 mL, 320 mmol). The resultant solution was heated to reflux for 2h. The solvent was removed under reduced pressure and dried under high vacuum to yield the title amide (11.1 g, 100%) as a white solid. The product was used in subsequent steps without purification. ^1H NMR (d_6 -DMSO δ 10.1, br s, 1H; δ 9.12, d, 1H; δ 8.83, d, 1H; δ 8.70, dd, 1H; δ 4.64, br s, 2H), LC/MS (APCI, ES, $M+H=139$).

Pyrazine-2-carbonyl azide. Pyrazine-2-carboxylic acid hydrazide (11.1 g, 80 mmol) was dissolved in 140 mL of water and charged with 6N HCl (13.3 mL, 80 mmol) and cooled to 0 °C. To the stirred reaction mixture was added a solution of sodium nitrite (8.3 g, 120 mmol) in 80 mL of water was added slowly over a period of 15-30 minutes using an addition funnel. After the addition was complete the reaction was warmed to room temperature and stirred for an additional 5h. The solution was the neutralized by the careful addition of solid NaHCO_3 and then extracted with CHCl_3 (3x). The pooled organic fractions were dried over Na_2SO_4 , filtered, concentrated and dried under high vacuum overnight to yield 2.5 g (21%) the title acyl azide. The product was used in subsequent steps without purification. ^1H NMR (d_6 -DMSO δ 9.30, d, 1H; δ 9.03, d, 1H; δ 8.90, dd, 1H).

(S)-3-[[2-Amino-5-(4-methoxy-phenyl)-thiophene-3-carbonyl]-amino]-azepane-1carboxylic acid tert-butyl ester. To a stirred solution of 2-amino-5-(4-methoxy-phenyl)-thiophene-3-carboxylic acid (1.0 g, 4.0 mmol) in anhydrous DMF (20 mL) is added (S)-3-amino-azepane-1-carboxylic acid tert-butyl ester (1.03 g, 4.8 mmol), BOP (2.6 g, 6.0 mmol) and NMM (0.6 mL, 5 mmol). The reaction mixture was stirred overnight at room temperature. The solution was diluted with water and EtOAc. The organic layer was separated and set aside. The remaining aqueous layer was extracted with EtOAc (2x) and then the combined organic extracts were pooled and washed with brine. The resultant EtOAc solution was dried over Na_2SO_4 , filtered,

- 74 -

and concentrated under vacuum to yield a brown solid. Purification was performed by ISCO MPLC (SiO₂, 30-50% EtOAc/hexanes) to give 0.9 g (50%) of an off-white solid. ¹H NMR (d₆-DMSO δ 7.51, d, 0.5H; δ 7.46, s, 0.5H; δ 7.37, s, 0.5H; δ 7.36, d, 0.5H; δ 7.34, br s, 2H; δ 7.33, d, 2H; δ 6.93, d, 2H; δ 4.11, br s, 1H; δ 3.76, s, 3H; δ 3.61, dq, 1H; δ 3.47, m, 1H; δ 3.11, m, 2H; δ 1.73, m, 3H; δ 1.56, m, 2H; δ 1.42, s, 4.5H; δ 1.38, s+m, 5.5H;), LC/MS (APCI, ES, M+H=446).

***tert*-Butyl (3*S*)-3-[(5-(4-methoxyphenyl)-2-[(pyrazin-2-ylamino)carbonyl]amino)-3-thienyl)carbonyl]amino}azepane-1-carboxylate.** A solution of (S)-3-{[2-amino-5-(4-methoxyphenyl)-thiophene-3-carbonyl]-amino}-azepane-1carboxylic acid tert-butyl ester (0.76 g, 1.7 mmol) and pyrazine-2-carbonyl azide (0.5 g, 3.4 mmol) in 20 mL of anhydrous DME was refluxed for 2h. The solvent was removed under reduced pressure and the crude product was purified using ISCO MPLC (40-60% EtOAc/hexanes) to give the title 0.51 g (53%) compound as a light yellow solid. ¹H NMR (d₆-DMSO δ 12.5, br s, 0.5H; δ 12.4, br s, 0.5H; δ 10.90, s, 0.5H; δ 10.88, s, 0.5H; δ 8.93, s, 0.5H; δ 8.89, s, 0.5H; δ 8.33, d, 1H; δ 8.29, t, 1H; δ 8.05, d, 0.5H; δ 7.91, d, 0.5H; δ 7.74, s, 0.5H; δ 7.65, s, 0.5H; δ 7.52, dd, 2H; δ 7.00, d, 2H; δ 4.26, m, 0.5H; δ 4.17, m, 0.5H; δ 3.79, s, 3H; δ 3.69, m, 1H; δ 3.48, m, 1H; δ 3.21, m, 2H; δ 1.77, m, 3H; δ 1.61, m, 2H; δ 1.44, s, 4.5H; δ 1.38, s+m, 5.5H), LC/MS (APCI, ES, M+H=567).

***N*-[(3*S*)-Azepan-3-yl]-5-(4-methoxyphenyl)-2-[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide; hydrochloride.** To a stirred solution of *tert*-butyl (3*S*)-3-[(5-(4-methoxyphenyl)-2-[(pyrazin-2-ylamino)carbonyl]amino)-3-thienyl)carbonyl]amino}azepane-1-carboxylate (0.51 g, 0.9 mmol) in 10 mL of MeOH is added 10 mL (40 mmol) of 4.0 N HCl in dioxane. The solution was stirred at room temperature for 4h and then concentrated under vacuum. The residue was partially recrystallized by triturating in refluxing 2-propanol to yield the title compound are a light orange solid (0.30 g, 67%). ¹H NMR (d₆-DMSO δ 12.6, br s, 1H; δ 10.9, s, 1H; δ 9.55, br s, 1H; δ 9.24, br s, 1H; δ 8.88, s, 1H; δ 8.49, d, 1H; δ 8.35, dd, 1H; δ 8.29, d, 1H; δ 7.92, s, 1H; δ 7.54, d, 2H; δ 6.99, d, 2H; δ 4.42, m, 1H; δ 3.33, m, 1H; δ 3.23, m, 2H; δ 3.10, m, 1H; δ 2.02, m, 1H; δ 1.85, m, 4H; δ 1.62, m, 1H;), LC/MS (APCI, ES, M+H=467).

- 75 -

Example 136**2-[(Aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide trifluoroacetate**

5 *tert*-Butyl (3S)-3-[[5-{4-[2-(diethylamino)ethoxy]phenyl}-2-[(hydroxyamino)carbonyl]amino}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate To a solution of *tert*-butyl (3S)-3-[[2-amino-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl) carbonyl]amino}piperidine-1-carboxylate (115 mg, 0.22 mmol) in THF (640 μ L) was added 1,1'-carbonyl
 10 diimidazole (178 mg, 1.1 mmol). The resulting cloudy solution was stirred at rt for 1 h whereupon hydroxylamine hydrochloride (76.4 mg, 1.1 mmol) and Et₃N (100 μ L) were added and the resulting dark solution was stirred for 48 h at rt. The mixture was partitioned between EtOAc and H₂O and the organic layer was washed with H₂O, brine and dried (MgSO₄). Evaporation afforded a yellow residue. Purification by Gilson (5%MeCN-H₂O→98%MeCN-
 15 H₂O) gave the title hydroxy urea.

5-{4-[2-(Diethylamino)ethoxy]phenyl}-2-[(hydroxyamino)carbonyl]amino}-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide A stirred solution of *tert*-butyl (3S)-3-[[5-{4-[2-(diethylamino)ethoxy]phenyl}-2-[(hydroxyamino)carbonyl]amino}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate in dioxane was treated with 4.0N HCl solution in
 20 dioxane (3 mL) and the resulting cloudy mixture was stirred at rt for 1 h. Evaporation of the solvent gave 5-[4-(2-diethylamino-ethoxy)-phenyl]-2-(3-hydroxy-urea)-thiophene-3-carboxylic acid-(S)-piperidin-3-ylamide as the hydrochloride salt (8 mg). ¹H NMR (300 MHz, DMSO-d₆) δ ppm 1.17 - 1.40 (m, 6 H) 1.54 - 1.81 (m, 2 H) 1.84 - 2.07 (m, 2 H) 2.76 - 2.99 (m, 2 H) 3.05 -
 25 3.25 (m, 4 H) 3.41 - 3.61 (m, 4 H) 3.73 - 3.88 (m, 1 H) 4.29 - 4.42 (m, 1 H) 4.40 - 4.64 (m, 1 H) 5.86 (s, 1 H) 7.07 (d, *J*=8.48 Hz, 2 H) 7.55 (d, *J*=8.48 Hz, 2 H) 7.82 (s, 1 H) 9.44 (s, 1 H) 9.65 (s, 1 H); LC/MS (ES, M+H=476).

Example 137

30 **2-[(Aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(3-methoxyphenyl)thiophene-3-carboxamide**

- 76 -

***tert*-Butyl (3*S*)-3-[(2-[(aminocarbonyl)amino]-5-bromo-3-thienyl) carbonyl)amino]azepane-1-carboxylate**

To a solution of methyl 2-[(aminocarbonyl)amino]-5-bromothiophene-3-carboxylate (1 equiv) in
 5 dry THF (0.3 M) was added a solution of [Me₂Al-Boc-3-(*S*)-aminohomopiperidine] (2 equiv) in
 THF (1.0 M) (which was preformed by the addition of Me₃Al (2.0 M in hexanes) to a solution of
 Boc-3-(*S*)-homopiperidine in THF at -78 °C and the resulting yellow solution was warmed to
 room temperature and stirred for an additional 15 min) and the resulting deep yellow solution
 was stirred overnight at room temperature. The reaction mixture was cooled with ice and a 10%
 10 aqueous solution of Rochelle's salt was added slowly to quench the reaction. The resulting
 biphasic solution was warmed to room temperature and stirred for an additional 1h. The mixture
 was diluted with EtOAc and H₂O, the aqueous layer was extracted with EtOAc (3x) and the
 combined organic extracts were washed with H₂O, brine and dried (MgSO₄). Evaporation gave a
 pale orange solid. Purification by column chromatography (40-60% EtOAc/Hexanes) gave a
 15 white solid. LC/MS (ES, M+H=462).

***tert*-Butyl (3*S*)-3-([2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl)amino)azepane-1-carboxylate**

A flask was loaded with *tert*-butyl (3*S*)-3-
 20 [(2-[(aminocarbonyl)amino]-5-bromo-3-thienyl) carbonyl)amino]azepane-1-carboxylate (1.0
 mmol), 3-methoxyphenylboronic acid (1.5 mmol), Cs₂CO₃ (3.0 mmol) and Pd(PPh₃)₄ (0.05-0.1
 mmol) and was purged with nitrogen for 10 mins. Dioxane (4mL) and H₂O (1mL) were added
 under nitrogen atmosphere and the resulting mixture was heated to 90°C for 2-4h. The mixture
 was allowed to cool to RT and the mixture was filtered (0.45 uM or diatomaceous earth). The
 water layer was separated and remaining solvent was concentrated to dryness. The residue was
 25 purified by column chromatography (SiO₂) on an MPLC ISCO separation system (30-60%
 EtOAc/hexanes) to give an off-white solid. LCMS, (ES, M+H=489).

2-[(Aminocarbonyl)amino]-N-[(3*S*)-azepan-3-yl]-5-(3-methoxyphenyl)thiophene-3-carboxamide hydrochloride

To a stirred solution of *tert*-butyl (3*S*)-3-([2-
 30 [(aminocarbonyl)amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl)amino)azepane-1-carboxylate

- 77 -

dissolved in a small amount of methanol was added 4.0 N HCl in dioxane. The solution was stirred for 1h at RT. The product as the hydrochloride salt was obtained as an off-white solid after removal of the solvent and drying. ¹H NMR (d₆-DMSO δ 10.90 (s, 1H), 9.21 (s, 1H), 9.01 (s, 1H), 8.29 (d, 1H), 7.93 (s, 1H), 7.29 (t, 1H), 7.11 (d, 2H), 7.03 (s, 1H), 6.83 (d, 1H), 4.33 (s, 1H), 3.81 (s, 3H), 3.20 (m, 4H), 2.00 (m, 1H), 1.84 (m, 4H), 1.56 (m, 1H)), LCMS, (ES, M+H=389).

Example 139**2-[(Aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide**

Methyl 2-([(trichloroacetyl)amino]carbonyl)amino)thiophene-3-carboxylate. To a stirred solution of 2-amino-thiophene-3-carboxylic acid methyl ester (1 eq) in anhydrous THF (mL) was added trichloroacetyl isocyanate (1 eq) slowly over a period of 5 min. After the addition was complete, a precipitate formed and the reaction stirred for an additional 1h. The desired product was obtained by filtration to give methyl 2-([(trichloroacetyl)amino]carbonyl)amino)thiophene-3-carboxylate (99%) as an off-white solid. The product was used in the next step without any further purification. LC/MS (ES, M+H=345).

Methyl 5-bromo-2-([(trichloroacetyl)amino]carbonyl)amino)thiophene-3-carboxylate. To a stirred solution of methyl 2-([(trichloroacetyl)amino]carbonyl)amino)thiophene-3-carboxylate (1 eq) in glacial acetic acid (20 mL) was added a solution of bromine (1.3 eq) in glacial acetic acid (5 mL) slowly over a period of 5 min. After the addition was complete, the resulting dark solution was stirred for 30 mins at RT. The solvent was evaporated under vacuum and the residue was triturated with H₂O. The title compound was obtained by filtration (99%) as an off-white solid. The product was used in the next step without any further purification after drying for 2 days under P₂O₅. LC/MS (ES, M+H=425).

Methyl 2-[(aminocarbonyl)amino]-5-bromothiophene-3-carboxylate. A stirred solution of methyl 5-bromo-2-([(trichloroacetyl)amino]carbonyl)amino)thiophene-3-carboxylate (1 eq) in

- 78 -

anhydrous methanol was purged with dry ammonia for 20 mins. After stirring for extra 10 mins at RT, precipitation was observed and the product was isolated by filtration (100% yield). LC/MS (ES, M+H=280).

5 ***tert*-Butyl(3*S*)-3-[(2-[(aminocarbonyl)amino]-5-bromo-3-thienyl)carbonyl]amino]piperidine-1-carboxylate**

To a solution of methyl 2-[(aminocarbonyl)amino]-5-bromothiophene-3-carboxylate (1 equiv) in dry THF (0.3 M) was added a solution of [Me₂Al-Boc-3-(*S*)-aminopiperidine] (2 equiv) in THF (1.0M) (which was preformed by the addition of Me₃Al (2.0M in hexanes) to a solution of Boc-3-
10 (S)-piperidine in THF at -78°C and the resulting yellow solution was warmed to room temperature and stirred for an additional 15 min) and the resulting deep yellow solution was stirred overnight at room temperature. The reaction mixture was cooled with ice and a 10% aqueous solution of Rochelle's salt was added slowly to quench the reaction. The resulting biphasic solution was warmed to room temperature and stirred for an additional 1h. The mixture
15 was diluted with EtOAc and H₂O, the aqueous layer was extracted with EtOAc (3x) and the combined organic extracts were washed with H₂O, brine and dried (MgSO₄). Evaporation gave a pale orange solid. Purification by column chromatography (40-60% EtOAc/Hexanes) gave a white solid. ¹H NMR (d₆-DMSO, δ 10.9, s, 1H; δ 9.48, br s, 1H; δ 9.31, br s, 1H; δ 8.48, d, 1H; δ 8.10, s, 1H; δ 7.57, d, 2H; δ 7.38, t, 2H; δ 7.23, t, 1H; δ 7.01, br s, 2H; δ 4.26, m, 1H; δ
20 3.29, m, 1H; δ 3.11, m, 1H; δ 2.94, m, 2H; δ 1.91, m, 2H; δ 1.69, m, 2H), LC/MS (APCI, ES, M+H=345).

***tert*-Butyl (3*S*)-3-([2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl)amino]piperidine-1-carboxylate** A flask was loaded with *tert*-butyl(3*S*)-3-
25 [(2-[(aminocarbonyl)amino]-5-bromo-3-thienyl)carbonyl]amino] piperidine-1-carboxylate (1.0 mmol), 3-methoxyphenylboronic acid (1.5 mmol), Cs₂CO₃ (3.0 mmol) and Pd(PPh₃)₄ (0.05-0.1 mmol) and was purged with nitrogen for 10 mins. Dioxane (4 mL) and H₂O (1 mL) were added under nitrogen atmosphere and the resulting mixture was heated to 90 °C for 2-4h. The mixture was allowed to cool to RT and the mixture was filtered (0.45 uM or diatomaceous earth). The
30 water layer was separated and remaining solvent was concentrated to dryness. The residue was

- 79 -

purified by column chromatography (SiO₂) on an MPLC ISCO separation system (30-60% EtOAc/hexanes) to give an off-white solid. LCMS, (ES, M+H=475).

2-[(Aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-

- 5 **carboxamide hydrochloride** To a stirred solution of *tert*-butyl (3*S*)-3-({[2-[(aminocarbonyl) amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl} amino)piperidine-1-carboxylate dissolved in a small amount of methanol was added 4.0 N HCl in dioxane. The solution was stirred for 1h at RT. The product as the hydrochloride salt was obtained as an off-white solid after removal of the solvent and drying. ¹H NMR (d₆-DMSO δ 10.94 (s, 1H), 9.41 (s, 1H), 9.19 (s, 1H), 8.47 (d, 1H),
10 8.11 (s, 1H), 7.30 (t, 1H), 7.15 (d, 1H), 7.05 (br, 2H), 6.84 (d, 1H), 4.25 (s, 1H), 3.82 (s, 3H), 3.29 (d, 1H), 3.12 (d, 1H), 2.97 (m, 2H), 1.92 (d, 2H), 1.69 (m, 2H)), LCMS, (ES, M+H=375).

Other Examples

- 15 Examples **1-3**, **5-25**, and **27-109** were prepared in a similar fashion to that described for examples **4** and **26**.

Examples **111** and **113-115** were prepared in a similar fashion to that described for examples **110** and **112**.

20

Examples **116-119** were prepared according to the general **Scheme V**.

Examples **121-122** and **124-135** were prepared in a similar fashion to that described for examples **123** and **136**.

25


Examples **138** and **140** were prepared in a similar fashion to that described for examples **137** and **139**.

Claims

CLAIMS

- 80 -

1. A compound of formula (I) or a pharmaceutically acceptable salt or *in vivo*-hydrolysable precursors thereof:



5

wherein:

R^1 and R^2 are at each occurrence independently selected from H, optionally substituted C_{1-6} alkyl, or optionally substituted heterocyclyl; with the proviso that R^1 and R^2 are not both H; or R^1 and R^2 and the N to which they are attached in combination form an optionally substituted heterocyclyl;

10

R^4 is selected from H, OH, optionally substituted carbocyclyl, optionally substituted heterocyclyl, or optionally substituted C_{1-6} alkyl;

R^5 is selected from H, optionally substituted carbocyclyl, or optionally substituted C_{1-6} alkyl.

15

2. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof as recited in claim 1 wherein R^2 , R^4 , and R^5 have any of the meanings defined in claim 1 and

R^1 is an optionally substituted heterocyclyl.

20

3. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof as recited in claim 1 wherein R^2 , R^4 , and R^5 have any of the meanings defined in claim 1 and R^1 is an optionally substituted heterocyclyl wherein 1,2, or 3 substituents is/are independently selected from halogen, nitro, amino, cyano, trifluoromethyl, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, hydroxy, alkylhydroxy, carbonyl, $-CH(OH)CH_3$, $-CH_2NH$ -alkyl-OH,

25

- 81 -

alkyl-(OH)CH₃, -CH₂-phenyl-(OCH₃)₂, -Oalkyl, -OCH₃, -Ophenyl, -OCOalkyl, -NHCHO, -
 Nalkyl, -N-(alkyl)-CHO, -NH-CO-amino, -N-(alkyl)-CO-amino, -NH-COalkyl, -N-(alkyl)-
 COalkyl, -carboxy, -amidino, -CO-amino, -CO-alkyl, -CO₂alkyl, mercapto, -Salkyl, -
 SCH₂furanyl, -SO(alkyl), -SO₂(alkyl), -SO₂-amino, -alkylsulfonylamino, phenyl, anisole,
 5 dimethoxyphenyl, trimethoxyphenyl, halophenyl, cycloalkyl, heterocyclyl, -alkyl-NH-cycloalkyl,
 -alkyl-NH- heterocyclyl, -alkyl-NH-alkyl-OH, -C(=O)OC(CH₃)₃, -N(CH₃)₂, -N(CH₂CH₃)₂, -
 alkyl-NH-alkyl- heterocyclyl, -alkyl-aryl, -methyl-phenyl, alkyl-polycyclyl, alkyl-amino, alkyl-
 hydroxy, -CH₂NH-alkyl-heterocyclyl, -CH₂NHCH₂CH(CH₃)₂, vicinal -O(alkyl)O-, vicinal -
 OC(haloalkyl)O-, vicinal -CH₂O(alkyl)O-, vicinal -S(alkyl)S- and -O(alkyl)S-.

10

4. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable
 precursor thereof as recited in claim 1 wherein R², R⁴, and R⁵ have any of the meanings defined
 in claim 1 and R¹ is an optionally substituted heterocyclyl wherein 1,2, or 3 substituents is/are
 independently selected from: -OH, C(=O)OC(CH₃)₃, NH₂, C₁₋₆alkyl, methoxybenzene, or
 15 dimethoxy benzene.

5. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable
 precursor thereof as recited in claim 1 wherein R², R⁴, and R⁵ have any of the meanings defined
 in claim 1 and

20

R¹ is a heterocyclyl wherein heterocyclyl is selected from piperdinyl, pyridinyl,
 pyrrolidinyl, pyrazinyl, azepanyl, azetidiny, azabicycloziny, furanyl, thienyl.

6. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable
 precursor thereof as recited in claim 1 wherein R¹, R⁴, and R⁵ have any of the meanings defined
 25 in claim 1 and

R² is H.

7. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable
 precursor thereof as recited in claim 1 wherein R¹, R², and R⁵ have any of the meanings defined
 30 in claim 1 and

- 82 -

R⁴ is H.

8. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof as recited in claim 1 wherein R¹, R², and R⁴ have any of the meanings defined in claim 1 and

R⁵ is H or an optionally substituted C₁₋₆alkyl.

9. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof as recited in claim 1 wherein R¹, R², and R⁴ have any of the meanings defined in claim 1 and

R⁵ is H or an optionally substituted C₁₋₆alkyl wherein 1,2 or 3 substituents is/are independently selected from: NH₂, NHCH₃, N(CH₂CH₃)₂, N(CH₃)₂, OCH₃, OH, -C₁₋₆alkyl, morpholino, piperidinyl, pyrrolodiny.

10. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof as recited in claim 1 wherein R¹, R², and R⁴ have any of the meanings defined in claim 1 and

R⁵ is H or an optionally substituted C₁₋₃alkyl.

11. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof as recited in claim 1 wherein R¹, R², and R⁴ have any of the meanings defined in claim 1 and

R⁵ is H or an optionally substituted C₁₋₃alkyl wherein 1,2 or 3 substituents is/are independently selected from: NH₂, NHCH₃, N(CH₂CH₃)₂, N(CH₃)₂, OCH₃, OH, -C₁₋₆alkyl, morpholino, piperidinyl, pyrrolodiny.

12. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof, as recited in claim 1 wherein:

R¹ is an optionally substituted heterocyclyl;

R² is H;

- 83 -

R⁴ is H;

R⁵ is H or an optionally substituted C₁₋₆alkyl.

13. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof, as recited in claim 1 wherein:

R¹ is an optionally substituted heterocyclyl wherein the substituent is selected from one or more of the following: -NH₂, C₁₋₆alkyl, -C(=O)OC(CH₃)₃,

R² is H;

R⁴ is H;

10 R⁵ is H or an optionally substituted C₁₋₆alkyl wherein the substituent is selected from one or more of the following: -C₁₋₆alkyl, -N(C₁₋₃alkyl)₂.

14. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof, as recited in claim 1 wherein:

15 R¹ is an optionally substituted heterocyclyl wherein the substituent is selected from one or more of the following: -NH₂, C₁₋₆alkyl, -C(=O)OC(CH₃)₃,

R² is H;

R⁴ is H;

20 R⁵ is H or an optionally substituted C₁₋₃alkyl wherein 1,2 or 3 substituents is/are independently selected from: NH₂, NHCH₃, N(CH₂CH₃)₂, N(CH₃)₂, OCH₃, OH, -C₁₋₆alkyl, morpholino, piperidinyl, pyrrolodinyl.

15. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof, as recited in claim 1 wherein:

25 R¹ is a heterocyclyl;

R² is H;

R⁴ is H;

R⁵ is H or a C₁₋₆alkyl.

- 84 -

16. A compound of formula (I) or a pharmaceutically salt or an *in vivo*-hydrolysable precursor thereof, as recited in claim 1 wherein:

R^1 is a 6-membered heterocyclyl containing at least one N in the ring;

R^2 is H;

5 R^4 is H;

R^5 is a C_{1-3} alkyl.

17. A compound of formula (I) selected from:

- tert-butyl 3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;
 10 2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-ylthiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-ylthiophene-3-carboxamide;
 15 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide;
 tert-butyl 3-{[(2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;
 2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-ylthiophene-3-
 20 carboxamide;
 2-[(aminocarbonyl)amino]-N-[(3R)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;
 N-(3-[(4-aminopiperidin-1-yl)carbonyl]-5-{4-[2-(diethylamino)ethoxy]phenyl}-2-thienyl)urea;
 2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[3-(hydroxymethyl)phenyl]thiophene-3-carboxamide;
 25 2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-ylthiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-N-(2-aminoethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-pyridin-3-ylthiophene-3-
 30 carboxamide;

- 85 -

- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(1-methylpiperidin-4-yl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-1-methylazepan-3-yl]thiophene-3-carboxamide;
- 5 2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-[3-(hydroxymethyl)phenyl]thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-pyrrolidin-3-ylthiophene-3-carboxamide;
- 10 2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-pyridin-3-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-1-methylpiperidin-3-yl]thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-{3-[2-(diethylamino)ethoxy]phenyl}-N-pyrrolidin-3-ylthiophene-3-carboxamide;
- 15 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-ylmethyl]thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]thiophene-3-carboxamide;
- 20 2-[(aminocarbonyl)amino]-N-[2-(dimethylamino)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-[2-(diethylamino)ethyl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 25 2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-yl]thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(piperidin-4-ylmethyl)thiophene-3-carboxamide;
- 30 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-pyrrolidin-3-ylthiophene-3-carboxamide;

- 86 -

- 2-[(aminocarbonyl)amino]-N-(1-ethylpiperidin-3-yl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-[(3S)-1-ethylazepan-3-yl]-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 5 2-[(aminocarbonyl)amino]-5-(3-hydroxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;
- tert-butyl (3S)-3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)pyrrolidine-1-carboxylate;
- 10 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-piperidin-3-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-(1-benzylpiperidin-4-yl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- tert-butyl 3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate;
- 15 2-[(aminocarbonyl)amino]-5-[4-(2-piperidin-1-ylethoxy)phenyl]-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-[4-(2-piperidin-1-ylethoxy)phenyl]-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-azetidin-3-yl-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 20 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(2S)-pyrrolidin-2-ylmethyl]thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-pyridin-4-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperazin-1-ylethyl)thiophene-3-carboxamide;
- 25 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperidin-1-ylethyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-1-azabicyclo[2.2.2]oct-3-yl-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-N-(2-hydroxyethyl)-5-(4-hydroxyphenyl)thiophene-3-carboxamide;

- 87 -

- 2-[(aminocarbonyl)amino]-N-(trans-4-hydroxycyclohexyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide;
- 5 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-piperazin-1-ylethyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-4-ylethyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-pyridin-3-ylethyl)thiophene-3-carboxamide;
- 10 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-3-ylethyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)thiophene-3-carboxamide;
- 15 2-[(aminocarbonyl)amino]-5-(2-methoxyphenyl)-N-piperidin-4-ylthiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(tetrahydrofuran-2-ylmethyl)thiophene-3-carboxamide;
- tert-butyl (3R)-3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate;
- 20 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-3-ylmethyl)thiophene-3-carboxamide;
- tert-butyl 3-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)azetidine-1-carboxylate;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-4-ylmethyl)thiophene-3-carboxamide;
- 25 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(3-methoxypropyl)thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[2-(2-thienyl)ethyl]thiophene-3-carboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-thienylmethyl)thiophene-3-carboxamide;
- 30 N-[3-(1,4-diazepan-1-ylcarbonyl)-5-(4-methoxyphenyl)-2-thienyl]urea;

- 88 -

- 2-[(aminocarbonyl)amino]-N-(2-methoxyethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-(2-thienylmethyl)thiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-N-{2-[(2-furylmethyl)thio]ethyl}-5-(4-methoxyphenyl)thiophene-3-carboxamide;
 5 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-N-[2-(2-thienyl)ethyl]thiophene-3-carboxamide;
 N-(3-[(4-aminopiperidin-1-yl)carbonyl]-5-{3-[2-(diethylamino)ethoxy]phenyl}-2-thienyl)urea;
 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(3R)-piperidin-3-ylmethyl]thiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(1,2,3,4-tetrahydroquinolin-3-yl)thiophene-
 10 3-carboxamide;
 2-[(aminocarbonyl)amino]-N-(1,3-benzodioxol-5-ylmethyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-N-(3-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-N-[2-(3,4-dimethoxyphenyl)ethyl]-5-(4-methoxyphenyl)thiophene-3-
 15 carboxamide;
 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-[(5-methyl-2-furyl)methyl]thiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(pyridin-2-ylmethyl)thiophene-3-carboxamide;
 20 2-[(aminocarbonyl)amino]-N-(4-fluorobenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
 tert-butyl 4-({[2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate;
 2-[(aminocarbonyl)amino]-N-(2-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;
 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-phenoxyethyl)thiophene-3-carboxamide;
 25 2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-N-(2-pyridin-2-ylethyl)thiophene-3-carboxamide;
 tert-butyl 4-({[2-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thienyl]carbonyl}amino)piperidine-1-carboxylate;
 2-[(aminocarbonyl)amino]-N-(4-methoxybenzyl)-5-(4-methoxyphenyl)thiophene-3-carboxamide;

- 89 -

2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-N-[(3R)-piperidin-3-yl]thiophene-3-carboxamide;

5 tert-butyl (3S)-3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-{4-[2-(diethylamino)ethoxy]phenyl}thiophene-3-carboxamide;

10 tert-butyl (3R)-3-{[(2-[(aminocarbonyl)amino]-5-{4-[2-(diethylamino)ethoxy]phenyl}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

N-[3-{[(3S)-3-aminoazepan-1-yl]carbonyl}-5-(4-methoxyphenyl)-2-thienyl]urea;

5-{4-[2-(diethylamino)ethoxy]phenyl}-2-{[(pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide;

15 5-{3-[2-(diethylamino)ethoxy]phenyl}-2-{[(pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide;

5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

N-[(3S)-azepan-3-yl]-5-(4-methoxyphenyl)-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

20 5-{3-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

N-(2-aminoethyl)-5-(4-methoxyphenyl)-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

25 5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-3-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

5-(4-methoxyphenyl)-N-piperidin-4-yl-2-{[(pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

tert-butyl 3-{[(5-{3-[2-(diethylamino)ethoxy]phenyl}-2-{[(pyrazin-2-ylamino)carbonyl]amino}-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

- 90 -

5-{4-[2-(diethylamino)ethoxy]phenyl}-N-piperidin-4-yl-2-[[pyrazin-2-ylamino)carbonyl]amino}thiophene-3-carboxamide;

5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino)carbonyl]amino}-N-[(3S)-pyrrolidin-3-yl]thiophene-3-carboxamide;

5 N-[3-(1,4-diazepan-1-ylcarbonyl)-5-(4-methoxyphenyl)-2-thienyl]-N'-pyrazin-2-ylurea;

N-[3-[(3-aminopyrrolidin-1-yl)carbonyl]-5-(4-methoxyphenyl)-2-thienyl]-N'-pyrazin-2-ylurea;

tert-butyl 4-[[5-(4-methoxyphenyl)-2-[[pyrazin-2-ylamino)carbonyl]amino]-3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

tert-butyl 3-[[5-{4-[2-(diethylamino)ethoxy]phenyl}-2-[[pyrazin-2-ylamino)carbonyl]amino]-

10 3-thienyl)carbonyl]amino}piperidine-1-carboxylate;

5-[4-(2-diethylamino-ethoxy)-phenyl]-2-(3-hydroxy-urea)-thiophene-3-carboxylic acid-(S)-piperidin-3-ylamide;

2-[(aminocarbonyl)amino]-N-[(3S)-azepan-3-yl]-5-(3-methoxyphenyl)thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-(2-hydroxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-

15 carboxamide;

2-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide;

2-[(aminocarbonyl)amino]-5-[2-(benzyloxy)phenyl]-N-[(3S)-piperidin-3-yl]thiophene-3-carboxamide.

20

18. A compound of formula (I) or a pharmaceutically acceptable salt thereof as recited in any one of claims 1 to 17 for use as a medicament.

19. The use of a compound of formula (I) or a pharmaceutically acceptable salt thereof as recited
25 in any one of claims 1 to 17, in the manufacture of a medicament for the treatment or prophylaxis of disorders associated with cancer.

20. A method for the treatment of cancer comprising administering to a human a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as
30 defined in any one of claims 1 to 17.

- 91 -

21. A method for the treatment of breast cancer, colorectal cancer, ovarian cancer, lung (non small cell) cancer, malignant brain tumors, sarcomas, melanoma and lymphoma by administering a compound of formula I or a pharmaceutically acceptable salt thereof as defined in any one of
5 claims 1 to 17.

22. A method of treating cancer by administering to a human a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 17 and an anti-tumor agent.

10 23. A method of treating cancer by administering to a human a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 17 and a DNA damaging agent.

15 24. A method for the treatment of infections associated with cancer comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 17.


20 25. A method for the prophylaxis treatment of infections associated with cancer comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 17.

25 26. A pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 17 together with at least one pharmaceutically acceptable carrier, diluent or excipient.

30 27. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt or *in vivo*-hydrolysable precursors thereof as defined in any one of claims 1 to 17, which comprises:

- 92 -


(a) the reaction of a 2-aminothiophene shown below as Formula II



II


wherein the hydrogen at the 2-amino position is displaced to form an amide, shown as formula III

5 below



III

wherein the methyl ester is converted to an amide utilizing the desired amine in conjunction with an aluminate organometallic complex, to give the product shown as formula IV below:




10

IV

Wherein the amide is converted to various substituted secondary ureas by the reaction with various isocyanates to yield the product shown as formula V below:


15

- 93 -



V

28. The use of a compound of formula (VI) below or a pharmaceutically acceptable salt or an in vivo hydrolysable precursor in the manufacture of a compound of formula (I) as set forth in any one of claims 1-17.



VI