



(11) (21) (C) **2,006,418**  
(22) 1989/12/21  
(43) 1990/06/28  
(45) 2000/04/11

(72) Angst, Max, CH  
(72) Gugumus, François, CH  
(72) Rist, Gunther, CH  
(72) Vogt, Manfred, DE  
(72) Rody, Jean, CH  
(73) NOVARTIS AG, CH  
(51) Int.Cl.<sup>5</sup> A01N 25/00  
(30) 1988/12/28 (4825/88-8) CH  
(54) **PESTICIDE**  
(54) **PEST CONTROL**

(57) A composition for controlling harmful insects and representatives of the order Acarina, containing as biologically active components a) a substance that modifies the behaviour of the pests and b) at least one pesticidally active compound, wherein the biologically active components are contained in a flowable or viscous non-hardening matrix that is resistant to water and weather, is protected against light and is suitable for distribution in the form of droplets or droplet-like units or spots that adhere to a substrate, by means of which matrix the behaviour-modifying substance is protected against UV radiation and from which it is slowly released in a biologically effective amount over a prolonged period, and wherein the pesticidally active compound can be taken up in a pesticidally effective amount from the surface of the droplets or droplet-like units or spots by the pests to be controlled; as well as a method of controlling the said pests, especially plant-destructive insects.



Pest controlAbstract

A composition for controlling harmful insects and representatives of the order Acarina, containing as biologically active components a) a substance that modifies the behaviour of the pests and b) at least one pesticidally active compound, wherein the biologically active components are contained in a flowable or viscous non-hardening matrix that is resistant to water and weather, is protected against light and is suitable for distribution in the form of droplets or droplet-like units or spots that adhere to a substrate, by means of which matrix the behaviour-modifying substance is protected against UV radiation and from which it is slowly released in a biologically effective amount over a prolonged period, and wherein the pesticidally active compound can be taken up in a pesticidally effective amount from the surface of the droplets or droplet-like units or spots by the pests to be controlled; as well as a method of controlling the said pests, especially plant-destructive insects.

5-17394/=

## Pest control

The present invention relates to a novel pesticidal composition for controlling harmful insects and representatives of the order Acarina, especially those of the plant-destructive kind, wherein a substance that alters the behaviour of the pests is freed in a biologically effective amount.

In the control of plant pests, especially insects and spider mites, in agricultural crops, especially monocrops, the aim in general is to allow as little contact as possible of the pesticidal active ingredients with the crop plants to be protected and the locus thereof, and with the soil in which they are growing. The pests found mostly on the plants should themselves, however, be exposed as much as possible to the active ingredient, whether by contact action, stomach poison action or gaseous phase action. A further aim is to ensure that useful insects and warm-blooded animals are not significantly harmed by the pesticidal active ingredients. In order to achieve these aims, in accordance with the invention a composition and a method of control have been proposed with which, in principle, the behaviour of the pests to be controlled can be so specifically directed or selectively modified using signal substances that the pests are led to a source of pesticidal active ingredient that is provided in discrete regions within the crop area to be protected.

Of the behaviour-modifying substances, so far mainly pheromones have been used in practice, in traps, for the early recognition of pest attack, for controlling pest populations by interrupting or interfering with the normal pairing process, or for catching pests, mainly insects (cf. for example, GB-PS 1 548 920 and GB Patent Application 2 063 068). In general, behaviour-modifying substances, especially pheromones, must have a slow release over a relatively long period of time and need to be pro-

tected from the influences of light and weather, GB Patent Application 2 064 323, for example, describes sprayable, film-forming compositions containing UV stabilisers and antioxidants that are said to provide slow release of pheromones and be suitable for the control of insects (cf. also EP-PS 55 475). It is also known from PCT Publication WO 87/04591 published on 13 August 1987 for the control of spider mites, to spray a slow release substance that modifies the behaviour of pests onto the leaves of the plants to be treated, which substance is applied in combination with a miticide in the conventional form of a spray mixture. The distribution of flowable, copolymeric pheromone-containing insecticidal compositions in the form of limited regions in the form of droplets or spots that provide slow release of the pheromone is described in GB-PS 2 141 932. These copolymeric compositions can be applied to the leaves of plants to be protected in the form of thin films or droplets that harden or set.

It is not possible, however, to achieve the aims of the invention set forth at the beginning using the above-described compositions and method of application, either because with the known pesticide/attractant combinations the entire agricultural area to be protected has to be given a surface cover spray, or because the compositions applied in discrete regions do not release the behaviour-modifying substances for a sufficient length of time or do not ensure adequate efficiency in respect of actual kill of the pests using a pesticidal active ingredient. In addition, the known formulations, which can be applied in droplet form (cf. GB-PS 2 141 932), do not meet requirements in practice as regards UV protection even though they contain UV absorbers, antioxidants, etc. It has also been found that phytotoxic phenomena often occur using known compositions of this kind having a polymer-based formulation.

The disadvantages mentioned above have now been overcome by the compositions and methods of control according to the invention. The invention accordingly relates to a composition for controlling harmful insects and representatives of the order Acarina, containing as biologically active components a) a substance that modifies the behaviour of the pests and b) at least one pesticidally active compound, wherein this composition contains the biologically active components in a flowable or viscous non-hardening matrix that is resistant to water and weather, is protected against light and is suitable for distribution in the form of droplets or droplet-like units or spots that adhere to a substrate, by means of which matrix the behaviour-modifying substance is protected against UV radiation and from which it is slowly released in a biologically effective amount over a prolonged period, and the pesticidally active compound can be taken up in a pesticidally effective amount from the surface of the droplets or droplet-like units or spots by the pests to be controlled. The invention further relates to a method of controlling harmful, especially plant-destructive, insects and representatives of the order Acarina, which comprises applying to or distributing over the area to be protected against pest attack, or the crop area, the afore-described composition in the form of a predetermined number of discrete droplets or droplet-like units or spots, the behaviour-modifying substance being slowly released in a biologically effective amount over a prolonged period into the surrounding atmosphere.

The invention therefore provides a viscous, non-hardening composition for controlling harmful insects and representatives of the order Acarina that is resistant to water and weather and having a viscosity of at least about 20,000 cp which comprises:

a) about 0.01-30% by weight of a pheromone or kairomone;

- 3a -

b) about 0.1-10% of a pesticidal compound that is compatible with other constituents of said composition; and

c) about 51-98% of an ultraviolet-absorber having an absorption range of from 270 to 400 nm selected from the group consisting of 2-H-benzotriazoles, 2-hydroxyalkoxybenzophenones, oxalanilides, cinnamic acid ester and triazines;

wherein the pheromone or kairomone is slowly released in a biologically effective amount over a period of time; and

wherein said composition is distributable for direct application without dilution as droplets or spots.

It was surprisingly found that the composition of the invention is outstandingly suitable for pest control and at the same time has a good lasting action, is tolerated by plants and provides a high degree of preservation of useful insects. As a result of the use of the mentioned signal substances or attractants, the amount of insecticides, which are often toxic to warm-blooded animals and useful insects, required for the treatment of the crop areas to be protected can be restricted considerably. This novel procedure which is also advantageous from the ecological point of view belongs to those concepts that are based on integrated plant protection and have already received considerable attention. As mentioned, the formulations of the invention are distinguished especially by the fact that both the behaviour-modifying substance and the pesticidally active compound are present directly in a liquid or flowable UV absorbent. As a result, optimum and prolonged UV protection, especially of the sensitive behaviour-modifying substance, is achieved.

The compositions of the invention exhibit particular activity especially in the control of plant-attacking insects and representatives of the order Acarina. The compositions and methods of the invention are suitable

especially for controlling insects of the orders: Lepidoptera, Coleoptera, Homoptera, Heteroptera, Diptera, Thysanoptera, Orthoptera, Anoplura, Siphonaptera, Mallophaga, Thysanura, Isoptera, Psocoptera and Hymenoptera, and representatives of the order Acarina of the families: Ixodidae, Argasidae, Tetranychidae and Dermanyssidae.

The compositions of the invention are used preferably for controlling destructive phytophagous insects in forests, in plantations of ornamental and useful plants, especially in cotton crops (e.g. *Pectinophora gossypiella*, *Spodoptera littoralis*, *Anthonomus grandis* and *Heliothis virescens*), in maize crops, in forestry and in viniculture. The compositions are distinguished by a pronounced activity not only against adult insects but also against larvae, especially of phytophagous insect pests.

The compositions of the invention can also be used for controlling flies, e.g. *Musca domestica* and *Stomoxys calcitrans*, mosquitos, and undesired populations of other insects, e.g. *Blattella germanica*, as well as for controlling ectoparasitic insects, e.g. *Lucilia sericata*, that harm domestic animals and livestock, for example by treating livestock buildings and pastures.

The following species of insects, for example, can be controlled successfully with the composition of the invention:

*Acleris* spp.  
*Adoxophyes fasciata*  
*Adoxophyes orana*  
*Aegeria apiformis*  
*Agriotes* spp.  
*Agrotis ipsilon*  
*Agrotis segetum*  
*Amylois transitella*  
*Anthonomus grandis*  
*Aonidiella aurantii*  
*Aonidiella citrana*  
*Archips* spp.  
*Argyrotaenia* spp.  
*Autographa* spp.  
*Blattella germanica*  
*Busseola fusca*  
*Cadra cautella*  
*Chilo partellus*

Chilo suppressalis  
Choristoneura fumiferana  
Choristoneura murinana  
Choristoneura rosaceana  
Choristoneura spp.  
Cnephasia spp.  
Cochylis spp.  
Coleophora laricella  
Coleophora spp.  
Cossus cossus  
Crocidolomia binotalis  
Curculio caryae  
Cydia pomonella  
Dacus dorsalis  
Dacus oleae  
Dacus spp.  
Dasychira spp.  
Dendroctonus brevicomis  
Dendroctonus frontalis  
Dendroctonus ponderosae  
Dendrolimus pini  
Dendroctonus spp.  
Dermestes spp.  
Diabrotica balteata  
Diabrotica longicornis  
Diabrotica vigifera  
Diabrotica barberi  
Diabrotica undecimpunctata  
Diatraea grandiosella  
Diatraea saccharalis  
Earias biplaga  
Earias insulana  
Earias vittella  
Ephestia elutella  
Ephestia kuehniella  
Eucosma ponderosa  
Eucosma sonomana  
Eucosma spp.  
Eupoecilia ambiguella  
Euproctis similis xanthocampa  
Euxoa spp.  
Glossina morsitans morsitans  
Gnathotrichus spp.  
Grapholita funebrana  
Grapholita janthinana  
Grapholita molesta  
Grapholita prunivora  
Grapholita sp.  
Hedya nubiferana  
Heliothis armigera  
Heliothis spp.  
Heliothis virescens  
Heliothis zea  
Hylobius abietis

Hylotrupes bajulus  
Ips paraconfusus  
Ips spp.  
Ips typographus  
Keiferia lycopersicella  
Leptinotarsa decemlineata  
Leucoptera scitella  
Lobesia botrana  
Lymantria dispar  
Lymantria monacha  
Lyonetia clerkella  
Malacosoma spp.  
Mamestra brassicae  
Manduca sexta  
Musca domestica  
Neodiprion sertifer  
Neodiprion spp.  
Operophtera brumata  
Ostrinia nubilalis  
Pammene rhediella  
Pammene spp.  
Pandemis spp.  
Pandemis heparana  
Panolis flammea  
Pectinophora gossypiella  
Pectinophora spp.  
Periplaneta americana  
Pityogenes chalcographus  
Pityokteines spp.  
Planococcus citri  
Platypus flavicoruis  
Plutella xylostella  
Popillia japonica  
Prays citri  
Prays oleae  
Pseudaulacaspis pentagona  
Pseudococcus comstocki  
Quadraspidiotus perniciosus  
Scolytus multistriatus  
Scolytus scolytus  
Scolytus spp.  
Sesamia spp.  
Sitotroga spp.  
Sparganothis pilleriana  
Sparganothis spp.  
Spodoptera exempta  
Spodoptera exigua  
Spodoptera frugiperda  
Spodoptera littoralis  
Spodoptera litura  
Spodoptera spp.  
Stomoxys calcitrans  
Synanthedon formicaeformis  
Synanthedon spp

*Tetranychus urticae*  
*Thaumetopoea pityocampa*  
*Tortrix viridana*  
*Trichoplusia ni*  
*Trogoderma granarium*  
*Trogoderma* spp.  
*Trypodendron domesticum*  
*Trypodendron lineatum*  
*Vespula* spp.  
*Yponomeuta* spp.  
*Zeiraphera diniana*  
*Zeuzera pyrina*

The compositions of the invention also exhibit good activity against plant-destructive acarids (spider mites: e.g. of the Tetranychidae, Tarsonemidae, Eriophyidae, Tyroglyphidae and Glycyphagidae families). The compositions of the invention are suitable especially for controlling the following species of mites that attack fruit and vegetable crops:

*Tetranychus urticae*, *Tetranychus cinnabarinus*, *Panonychus ulmi*, *Bryobia rubrioculus*, *Panonychus citri*, *Eriophyes pyri*, *Eriophyes ribis*, *Eriophyes vitis*, *Tarsomemus pallidus*, *Phyllocoptes vitis* and *Phyllocoptura oleivora*.

The good pesticidal activity of the compositions of the invention corresponds to a killing rate (mortality) of at least 50-60 % of the mentioned pests.

Behaviour-modifying substances suitable for the invention are chiefly pheromones, but kairomones and attractants are also suitable. Such signal substances are effective even in extraordinarily low concentrations and can modify the behaviour of insects in a manner that renders possible their control. The proportion of behaviour-modifying substance in the compositions of the invention is preferably from 0.01 to 30 % by weight. In the case of pheromones, the proportions used are preferably from 0.01 to 2.0 % by weight, and in the case of kairomones and attractants, which are usually contained in compositions for controlling larval stages, the proportions used are from 0.5 to 30 % by weight.

Pheromones are sexual signal substances, produced in most cases by the adult females of pests, mainly insects, that attract individuals of the opposite sex of the same pest species. The natural pheromones and pheromone mixtures are volatile and their attracting effect can reach over long distances. In principle, all pheromones described in the literature are suitable for the purposes of the invention. The structure and composition of pheromones is known from the literature, c.f., e.g.: A.F. Kydoneus et al., "Insect Suppression with Controlled Release Pheromone Systems", CRC Press (1982); H. Arn et al., "List of Pheromones of Lepidoptera and Related Attractants", OILB (1986).

The following pheromones, for example, may be used within the scope of the present invention:

Z-5-Decenyl acetate  
Dodecanyl acetate  
Z-7-Dodecenyl acetate  
E-7-Dodecenyl acetate  
Z-8-Dodecenyl acetate  
E-8-Dodecenyl acetate  
Z-9-Dodecenyl acetate  
E-9-Dodecenyl acetate  
E-10-Dodecenyl acetate  
11-Dodecenyl acetate  
Z-9,11-Dodecadienyl acetate  
E-9,11-Dodecadienyl acetate  
Z-11-Tridecenyl acetate  
E-11-Tridecenyl acetate  
Tetradecanyl acetate  
E-7-Tetradecenyl acetate  
Z-8-Tetradecenyl acetate  
E-8-Tetradecenyl acetate  
Z-9-Tetradecenyl acetate  
E-9-Tetradecenyl acetate  
Z-10-Tetradecenyl acetate  
E-10-Tetradecenyl acetate  
Z-11-Tetradecenyl acetate  
E-11-Tetradecenyl acetate  
Z-12-Pentadecenyl acetate  
E-12-Pentadecenyl acetate  
Hexadecanyl acetate  
Z-7-Hexadecenyl acetate  
Z-11-Hexadecenyl acetate  
E-11-Hexadecenyl acetate  
Octadecanyl acetate  
E,Z-7,9-Dodecadienyl acetate  
Z,E-7,9-Dodecadienyl acetate

E,E-7,9-Dodecadienyl acetate  
Z,Z-7,9-Dodecadienyl acetate  
E,E-8,10-Dodecadienyl acetate  
E,Z-9,12-Dodecadienyl acetate  
E,Z-4,7-Tridecadienyl acetate  
4-methoxy-cinnamaldehyde  
β-Ionone  
Estragol  
Eugenol  
Indole  
8-Methyl-2-decyl-propanoate  
E,E-9,11-Tetradecadienyl acetate  
Z,Z-9,12-Tetradecadienyl acetate  
Z,Z-7,11-Hexadecadienyl acetate  
E,Z-7,11-Hexadecadienyl acetate  
Z,E-7,11-Hexadecadienyl acetate  
E,E-7,11-Hexadecadienyl acetate  
Z,E-3,13-Octadecadienyl acetate  
E,Z-3,13-Octadecadienyl acetate  
E,E-3,13-Octadecadienyl acetate

Ethanol  
Hexanol  
Heptanol  
Octanol  
Decanol  
Z-6-Nonenol  
E-6-Nonenol  
Dodecanol  
11-Dedecenol  
Z-7-Dedecenol  
E-7-Dedecenol  
Z-8-Dedecenol  
E-8-Dedecenol  
E-9-Dedecenol  
Z-9-Dedecenol  
E-9,11-Dodecadienol  
Z-9,11-Dodecadienol  
Z,E-5,7-Dodecadienol  
E,E-5,7-Dodecadienol  
E,E-8,10-Dodecadienol  
E,Z-8,10-Dodecadienol  
Z,Z-8,10-Dodecadienol  
Z,E-8,10-Dodecadienol  
E,Z-7,9-Dodecadienol  
Z,Z-7,9-Dodecadienol  
E-5-Tetradecenol  
Z-8-Tetradecenol  
Z-9-Tetradecenol  
E-9-Tetradecenol  
Z-10-Tetradecenol  
Z-11-Tetradecenol  
E-11-Tetradecenol  
Z-11-Hexadecenol

Z,E-9,11-Tetradecadienol  
 Z,E-9,12-Tetradecadienol  
 Z,Z-9,12-Tetradecadienol  
 Z,Z-10,12-Tetradecadienol  
 Z,Z-7,11-Hexadecadienol  
 Z,E-7,11-Hexadecadienol  
 (E)-14-Methyl-8-Hexadecen-1-ol  
 (Z)-14-Methyl-8-Hexadecen-1-ol  
 E,E-10,12-Hexadecadienol  
 E,Z-10,12-Hexadecadienol  
 Dodecanal  
 Z-9-Dodecenal  
 tetradecanal  
 Z-7-Tetradecenal  
 Z-9-Tetradecenal  
 Z-11-Tetradecenal  
 E-11-Tetradecenal  
 E-11,13-Tetradecadienal  
 E,E-8,10-Tetradecadienal  
 Z,E-9,11-Tetradecadienal  
 Z,E-9,12-Tetradecadienal

Hexadecanal  
 Z-8-Hexadecenal  
 Z-9-Hexadecenal  
 Z-10-Hexadecenal  
 E-10-Hexadecenal  
 Z-11-Hexadecenal  
 E-11-Hexadecenal  
 Z-12-Hexadecenal  
 Z-13-Hexadecenal  
 (Z)-14-Methyl-8-Hexadecenal  
 (E)-14-Methyl-8-Hexadecenal  
 Z,Z-7,11-Hexadecadienal  
 Z,E-7,11-Hexadecadienal  
 Z,E-9,11-Hexadecadienal  
 E,E-10,12-Hexadecadienal  
 E,Z-10,12-Hexadecadienal  
 Z,E-10,12-Hexadecadienal  
 Z,Z-10,12-Hexadecadienal  
 Z,Z-11,13-Hexadecadienal  
 Octadecanal  
 Z-11-Octadecenal  
 E-13-Octadecenal  
 Z-13-Octadecenal  
 Z-5-Decenyl-3-methyl-butanoate

Disparlure:	(+) cis-7,8-Epoxy-2-methyloctadecane
Seudenol:	3-Methyl-2-cyclohexen-1-ol
Sulcatol:	6-Methyl-5-hepten-2-ol
Ipsenol:	2-Methyl-6-methylene-7-octen-4-ol
Ipsdienol:	2-Methyl-6-methylene-2,7-octadien-4-ol
Grandlure I:	cis-2-Isopropenyl-1-methylcyclobutane ethanol
Grandlure II:	Z-3,3,-Dimethyl-1-cyclohexane ethanol

Grandlure III:	Z-3,3,-Dimethyl-1-cyclohexane acetaldehyde
Grandlure IV:	E-3,3,-Dimethyl-1-cyclohexane acetaldehyde
cis-2-Verbenol:	cis-4,6,6-Trimethylbicyclo[3.1.1]hept-3-en-2-ol
	cucurbitacin
	2-Methyl-3-buten-2-ol
	4-Methyl-3-heptanol
Cucurbitacin	
	2-Methyl-3-buten-2-ol
	4-Methyl-3-heptanol
$\alpha$ -Pinen:	2,6,6-Trimethylbicyclo[3.1.1]hept-2-ene
$\alpha$ -Caryophyllen:	4,11,11-Trimethyl-8-methylenebicyclo[7.2.0]undecane
	Z-9-Tricosene
$\alpha$ -Multistriatin	2(2-endo,4-endo)-5-Ethyl-2,4-dimethyl-6,8-dioxabicyclo- [3.2.1]octane
Methyleugenol:	1,2-Dimethoxy-4-(2-propenyl)phenol
Lineatin:	3,3,7-Trimethyl-2,9-dioxatricyclo[3.3.1.0]nonane
Chalcogran:	2-Ethyl-1,6-dioxaspiro[4.4]nonane
Frontalin:	1,5-Dimethyl-6,8-dioxabicyclo[3.2.1]octan
endo-Brevicommin:	endo-7-Ethyl-5-methyl-6,8-dioxabicyclo[3.2.1]octane
exo-Brevicommin:	exo-7-Ethyl-5-methyl-6,8-dioxabicyclo[3.2.1]octane
	(Z)-5-(1-Decenyl)dihydro-2(3H)-furanone
Farnesol	3,7-,11-Trimethyl-2,6,10-dodecatrien-1-ol
Nerolidol	3,7-,11-Trimethyl-1,6,10-dodecatrien-3-ol
	3-Methyl-6-(1-methylethenyl)-9-decen-1-ol-acetate
	(Z)-3-Methyl-6-(1-methylethenyl)-3,9-decadien-1-ol- acetate
	(E)-3,9-Methyl-6-(1-methylethenyl)-5,8-decadien- 1-ol-acetate
	3-Methylene-7-methyl-octen-1-ol-propionate
	(Z)-3,7-Dimethyl-2,7-octadien-1-ol-propionate
	(Z)-3,9-Dimethyl-6-(1-methylethenyl)-3,9-decadien- 1-ol-propionate

Kairomones, too, are signal substances of natural origin. They are produced by plants and consist chiefly of a mixture of a number of different volatile odorous substances. Kairomones are able to attract insects and representatives of the order Acarina. Depending on the concentration, however, they may also have a repelling effect. The activity of kairomones should be seen in conjunction with the fact that insects and acarines form a close living partnership with the relevant plants. Kairomones suitable for the purposes of the invention and methods of obtaining them are described, for example, in the following literature sources: Science, 154, 1392-93 (1966); P.A. Hedin "Bioregulators for Pest Control", American Chemical Society, Washington, 353-366 (1985).

So-called "attractants" are known chemical compounds that are relatively readily available and can exert behaviour-modifying effects on pests for the purposes of making these pests accessible and permitting substantial exposure thereof to the action of pesticides that are simultaneously present. Such attractants, which can also be used for the purposes of the invention, preferably in combination with pheromones and/or kairomones, are known, for example, from the following literature sources: G.R. Waller, "Allelochemicals: Role in Agriculture and Forestry", ACS, Washington, 431-438 (1987); Entomol. exp. appl 41, 11-16 (1986); Can. Entomol. 115, 1-5 (1983).

Conventional insecticidal and acaricidal active ingredients that are suitable and are known for controlling the relevant pests are used as pesticidally active compounds in the compositions and preparations of the invention. Obviously, the pesticidal active ingredients must be compatible with the other constituents of the flowable or viscous composition

of the invention and must be at least substantially soluble therein. For this reason preference is given in accordance with the invention to liquid pesticidally active compounds. However, the pesticidal active ingredients must not evaporate too quickly, but should be present at or on the surface of the applied droplets or droplet-like units or spots for a reasonable period of time so as to render possible the uptake of the active ingredient by the pest to be controlled. Accordingly, preferably pesticidal contact-active ingredients are present in the compositions of the invention. The compositions according to the invention generally contain from 0.1 to 10 % by weight, preferably from 1 to 5 % by weight, of the pesticidally active compound. Representatives of the following classes of pesticidal active ingredient are suitable for the purposes of the invention: carbamates, organophosphorus compounds, nitrophenols and derivatives thereof, formamidines, ureas, phenylbenzoylureas, pyrethroids, chlorinated hydrocarbons and *Bacillus thuringiensis* preparations.

Preferably, the compositions of the invention contain as classes of pesticidally active compounds carbamates, organophosphorus compounds, pyrethroids and insect-growth regulators.

Suitable carbamates are, for example:

CLOETOCARB	BENDIOCARB
BPMC	CARBOFURAN
METHOMYL	CARBOSULFAN
BUFENCARB	PIRIMICARB
DIOXACARB	ISOPROCARB
THIODICARB	BENFURACARB
FURATHIOCARB	ALDICARB
NITRILACARB	CARBARYL
METHOMYL	THIODICARB
OXAMYL	

Suitable organophosphorus compounds are, for example:

DIMETHOAT	PROFENOFOS	MALATHION
ZOLAPROFOS	AZAMETHIPHOS	CHLORPYRIFOS
AZINPHOS-METHYL	METHACRIFOS	CHLORPYRIFOS-M
SULPROFOS	CHLORFENVINPHOS	TRIAZOPHOS

METHYLPARATHION  
 PARATHION  
 FENITROTHION  
 CARBOFURAN  
 TRICHLORPHON  
 PHOXIM  
 ACEPHATE  
 METAMIDOPHOS  
 ISAZOPHOS

PHOSPHAMIDON  
 DICROTOPHOS  
 MONOCROTOPHOS  
 DICHLORVOS  
 DIMETILAN  
 DIAZINON  
 METHIDATION  
 JODFENPHOS  
 DEMETON-S-METHYL

ENDOSULFAN  
 PROPAPHOS  
 SULPROFOS  
 PIRIMIPHOS-M  
 PHOSALONE  
 VAMIDOTHION  
 CHLORMEPHOS  
 FORMOTHION  
 ETRIMIFOS  
 FONOFOS

Suitable pyrethroids are, for example:

BIORESMETHRIN  
 CYPERMETHRIN  
 CYHALOTHRIN  
 TEFLUTHRIN  
 PERMETHRIN  
 DECAMETHRIN  
 TRALOCYTRIN

FENPROPATHRIN  
 RESMETHRIN  
 PHENOTHRIN  
 FENVALERATE  
 FENPROPATHRIN  
 FLUVALINATE

Suitable insect-growth regulators that are used especially for controlling larval stages are, for example:

CHLORDIMEFORM  
 CYROMAZINE  
 CHLORFLUAZURON  
 BUPROFEZINE  
 DIFLUBENZURON  
 FLUFENOXURON

The follow acaricides, for example, are suitable in accordance with the invention for controlling representatives of the order Acarina:

CHLORBENZILAT  
 CHLORPROPHYLAT  
 BROMPROPHYLAT  
 AMITRAZ  
 HEXYTHIAZOX  
 ROTENON  
 DIPEL


The pesticidally active compound can be omitted from certain forms of the compositions of the invention. This is the case, for example, when extremely small larvae are to be attracted which become stuck to the matrix and therefore die without pesticidally active compound.

It is an essential feature of the present invention that the matrix of the compositions of the invention, in which the biologically active components are contained preferably in dissolved or emulsified form, consists for the most part of a liquid UV absorbent that is known per se. Liquid or flowable UV absorbers having an absorption range of from 270 to 400 nm have proved to be especially suitable for the purposes of the invention because on the one hand they ensure the necessary protection of the biological components contained in the composition against the influences of weather and light, and because on the other hand they have properties of permanent flowability. Consequently it is possible to impart to the compositions, if appropriate with the addition of certain viscosity-regulating substances, a consistency that makes them suitable for distribution in the form of droplets or droplet-like units or spots on a substrate, the substrate preferably being the leaves or other parts of the plants to be protected. These droplets, or droplet-like units or spots, which are each of an order of magnitude of from 1 to 100  $\mu\text{l}$ , preferably from 5 to 40  $\mu\text{l}$ , adhere to the relevant substrate and provide slow and controlled release over a prolonged time period of the substance that modifies the behaviour of the pests. On account of their differing biological, mobility and feeding behaviour, larval and adult stages of the pest are expediently controlled by producing droplets or droplet-like units of differing size on the attacked crop surfaces, these sizes preferably being from 1 to 20  $\mu\text{l}$  for larvae and from 30 to 100  $\mu\text{l}$  for adults. Generally, the slow or controlled release of the behaviour-modifying substance takes place over a period of up to 15 weeks, but at least up to 6 weeks. When the attracted pests come into contact with the permanently flowable or viscous composition, they inevitably take up from the surface thereof a pesticidally effective amount of the insecticide or acaricide contained in the composition. Such a controlled pesticide intake ("attract-and-kill" effect) is not provided, for example, by com-

positions that adhere to substrates, for example to parts of plants, in the form of a lacquer-like layer or in the form of solid and hard polymers.

For the compositions of the invention, there are used as a matrix base, that is to say as chief quantitative component of the matrix, liquid or flowable UV absorbers that are known per se, preferably having an absorption range of from 270 to 400 nm. Generally from 51 to 98 % by weight, preferably from 70 to 98 % by weight, of the compositions of the invention consists of the matrix, that is to say the UV absorber. It has proved advantageous for the UV absorbers themselves to have a viscosity of from 1000 to 40,000 cp. Classes and types of UV absorbers that are suitable in accordance with the invention include, for example, the following, provided they are liquid and meet the other conditions of the invention:

1. 2-(2-hydroxyphenyl)-benzotriazoles:



R<sub>1</sub> = H, Cl;


R<sub>2</sub> = H, alkyl, phenylalkyl, phenyl;

R<sub>3</sub> = alkyl, phenylalkyl, phenyl, -(CH<sub>2</sub>)<sub>2</sub>-COO-alkyl.

The following UV absorbers of that type are preferred:

Trade name	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
Tinuvin 109	Cl	-C(CH <sub>3</sub> ) <sub>3</sub>	50 % $-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{C}_8\text{H}_{17}$ 50 % $-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{CH}_2-\underset{\text{C}_2\text{H}_5}{\text{CH}}-\text{C}_4\text{H}_9$
Tinuvin 171	H	-C <sub>12</sub> H <sub>25</sub> -(iso- meric mixture)	-CH <sub>3</sub> or C <sub>2</sub> -C <sub>12</sub> alkyl
Tinuvin 1130	H	-C(CH <sub>3</sub> ) <sub>3</sub>	ca. 50 % $-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-(\text{CH}_2-\text{CH}_2-\text{O})_{300}-\text{H}$ ca. 38 % $\left[ \text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-(\text{CH}_2-\text{CH}_2-\text{O})_{150} \right]_2$ ca. 12 % polyethyleneglycol (EO 300)
"SL 874"	H	-C(CH <sub>3</sub> ) <sub>3</sub>	$-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-(\text{CH}_2)_8-\text{CH}=\text{CH}-(\text{CH}_2)_7-\text{CH}_3$


2. 2-hydroxy-4-alkoxybenzophenones:







R = alkyl (e.g. iso-C<sub>8</sub>H<sub>17</sub>, -C<sub>12</sub>H<sub>25</sub>)

R' = H, alkyl, alkoxy

3. Oxalanilides:



4. Cinnamic acid derivatives:R = alkyl (e.g. n-C<sub>4</sub>H<sub>9</sub>)R = alkyl (e.g. C<sub>4</sub>H<sub>9</sub>, -CH<sub>2</sub>-CH(C<sub>2</sub>H<sub>5</sub>)-C<sub>4</sub>H<sub>9</sub>)5. Triazine derivatives:R = C<sub>1</sub>-C<sub>18</sub>alkyl, -O-(C<sub>1</sub>-C<sub>18</sub>alkyl)R<sub>1</sub> = H, OH

n = 0, 1 or 2

If necessary, the flowable preparations of the invention may contain conventional viscosity-regulating thickeners in an amount of from 1 to 47 %, preferably from 1 to 20 %, based on the weight of the matrix. The following, for example, are suitable as organic thickeners: acrylic acid polymers of high molecular weight and relatively high viscosity neutralised with bases ("Carbople" types), polyvinylpyrrolidones, cellulose

gums, especially cellulose alkyl esters and alkyl ethers ("Blanose" types), liquid polyalkylene glycol block mixed polymers of ethylene oxide and propylene oxide ("Pluronic" types) and also polyethylene glycols having a molecular weight of more than 10,000. Of the inorganic thickeners, the following, for example, may be mentioned: precipitated or pyrogenic silicic acids, aluminium oxides and natural mineral fillers, especially calcite, types of talcum, kaolins, bentonites, montmorillonites, smectites and attapulgite, aluminium oxide/silicon dioxide ("Aerosil" types) and sodium-aluminium silicates. Quartz sand or solid crosslinked polymers in powder form may be incorporated in the preparations as additional fillers.

In order to adjust the viscosity of the flowable preparations of the invention, it may be expedient in certain cases to add an inert solvent or diluent. These solvents should be compatible with the other constituents of the preparation and should preferably not be excessively volatile. The following solvents, for example, are suitable: ethers and ethereal compounds that are not readily volatile, such as dipropyl ether, dibutyl ether, dioxane, dimethoxyethane and tetrahydrofuran; N,N-dialkylated carboxylic acid amides; aliphatic, aromatic and halogenated hydrocarbons, especially pentanes, hexanes, heptanes, octanes, hexadecane, toluene, xylenes, chlorohydrocarbons and chlorobenzenes, alcohols, such as ethanol, propanols, t-butanol and higher alcohols; nitriles, such as acetonitrile or propionitrile; and also ketones, for example methyl isopropyl ketone and methyl isobutyl ketone; alkyl esters of aliphatic carboxylic acids, such as propionic acid butyl ester, oxalic acid methyl ester, sebacic acid dibutyl ester and sebacic acid di-(2-ethylhexyl) ester. In general, only relatively small amounts of solvent, for example from 1 to 2 % by weight, are contained in the formulations of the invention.

If necessary, the compositions of the invention may contain further formulation auxiliaries, which are used to adapt the compositions to specific conditions or to protect the biological active ingredients contained therein against particular environmental conditions and influences. Such formulation auxiliaries may be substances that ensure an

additional UV protection, such as fine-particled carbon powder (e.g. carbon black), dyestuffs and colouring pigments (e.g. Sudan black, chromophthalic blue, Terasil blue, Cibacet yellow, titanium dioxide, zinc sulfate and zinc oxides), optical brighteners (e.g. Uvitex or Tinopal DMS), antioxidants (e.g. Tinuvin 765, butylhydroxytoluene or 2,6-di-tert-butyl-p-cresol) and in some cases also certain surface-active substances and emulsifiers (e.g. anionic surfactants: Na lauryl sulfate, dodecylbenzenesulfonic acid Ca salt, and non-ionic surfactants: fatty alcohol ethoxylates, alkylphenol ethoxylates, oleyl alcohol ethoxylates, ethylene oxide/propylene oxide block copolymers, fatty amine ethoxylates, silicone surfactants). Lime pigments and colouring pigments may be present in the compositions of the invention in an amount of from 1 to 20 % by weight, preferably from 3 to 10 % by weight, optical brighteners in an amount of from 0.1 to 2 % by weight, antioxidants in an amount of from 0.1 to 5 % by weight and surface-active substances and emulsifiers in an amount of from 0.1 to 1 % by weight.

Within the scope of the method of the invention for the control of plant-destructive insects and representatives of the order Acarina, the compositions of the invention can be distributed in the form of droplets or droplet-like units or spots over the agricultural crop area to be treated using conventional devices as known to the person skilled in the art. Automatic pipettes with appropriate metering means, for example, have proved suitable for smaller areas. Also suitable are arrangements with intermittent production of droplets controlled mechanically or by compressed air, which are suitable mainly for extensive surface areas. Within the scope of the method of the invention, the number of droplets or droplet-like units or spots distributed over the crop area to be treated may vary in accordance with the nature of the pests to be controlled and their stages of development. From 50 to 100,000,000 of said droplets or droplet-like units or spots may be distributed per 1000 m<sup>2</sup> of crop area to be protected, from 50 to 1000 droplets or droplet-like units of the compositions of the invention generally being used to control adults and from 500,000 to 100,000,000 to control larvae.

The following Examples illustrate the invention, the percentage values in the formulations relating to weight unless otherwise indicated.

Example 1: A flowable formulation having a viscosity of 25,000 cp is prepared by intimately mixing the following constituents:

Tinuvin 171 (cf. above data)	85.6 %
carbon black powder (density 1.8-1.9 g/cm <sup>3</sup> , particle size 20 nm, surface area 240 m <sup>2</sup> /g)	4.3 %
Aerosil COK 84 (finely particulate SiO <sub>2</sub> /Al <sub>2</sub> O <sub>3</sub> -84:16 % by wt.)	2.6 %
Furathiocarb	4.3 %
Codlemon	0.1 %
hexane	3.1 %

Droplets of this formulation, each of 100 µl, are applied to an aluminium sheet and each day are exposed alternately to UV radiation (UV lamp) at 35°C for 9 hours and to a temperature of 12°C in darkness for 15 hours. Droplets with different durations of treatment are then, immediately after that treatment, examined in a "wind tunnel" for insecticidal activity and their ability to attract *Cydia pomonella* males.

The wind tunnel used, as has already been described in principle in the literature (cf. J.R. Hiller and W.L. Roelofs, *J. Chem. Ecology*, 4, 187-198, 1978), consists essentially of a sealed tunnel of transparent material of rectangular cross-section. A droplet of the attractant/insecticide formulation as described above is arranged at one end of the tunnel. Short open glass tubes for receiving the insects (*Cydia pomonella* males) and a take-off pad are arranged at the opposite end of the tunnel. An attracting effect of the attractant contained in the droplet, which takes effect on the insect in the glass tube when a stream of air is produced, causes a state of excitation in the *Cydia pomonella* males, which is manifested initially by motorisation and buzzing of the moths' wings. The source of attractant (that is the droplet containing the active ingredient) is then specifically flown to from the take-off pad.

In the case of substances that do not have an attracting effect, the moth remains, unstimulated, in the open glass tube. The glass tube is occupied by one moth for every flight test. On average, 40 flight tests are carried out for each formulation to be tested. For the purposes of evaluation, the percentage of moths that fly in the direction of the droplet and touch ("contact") it is ascertained. The moths that touch the droplets are captured and their percentage mortality is ascertained after 24 hours. The results obtained, compared with control batches without attractant, are compiled in the following Table:

Action on C. pomonella	Duration of treatment of the droplets (in days)				
	0	3	7	14	21
% contact	95	85	95	95	80
% mortality	100	100	100	100	100

Example 2: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the preceding Example 1:

Tinuvin 171	85.5 %
carbon black powder (particle size 35 nm, surface area 1000 m <sup>2</sup> /g)	4.3 %
Sudan black B (dyestuff)	4.3 %
Furathiocarb	4.3 %
Codlemon	0.2 %
hexane	1.4 %

Droplets of 100 µl of this formulation are applied in the field to the leaves of apple trees and tested in the wind tunnel after 35 days. These droplets still exhibit a very good attracting effect and insecticidal activity in the wind tunnel. 80 % of the tested males of *Cydia pomonella* are attracted by such a droplet, fly to it and touch it. After 24 hours the mortality of these males is 100 %. The wind tunnel tests are carried out as indicated in Example 1.

Example 3: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	81.5 %
Aerosil COK 84	3.2 %
carbon black powder	4.1 %
Sudan black B	4.1 %
Furathiocarb	4.1 %
Codlemon	0.1 %
hexane	2.9 %

In a field test with the codling moth *Cydia pomonella*, a 1.2 ha orchard with about 1200 apple trees is treated with 8 droplets, each of 100 µl, of the above flowable formulation per tree (approximately 8000 droplets/ha). A second corresponding application with this formulation is carried out 41 days later. Figures for the attack in the treated orchard compared with an untreated control orchard are as follows:

Evaluation point of time (No. of days after first application)	% attack by <i>C. pomonella</i>	
	treated	untreated
46	0.6 %	7 %
87	0.8 %	22 %

Example 4: A viscous formulation is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	80.1 %
Tinuvin 765	1.6 %
Aerosil COK 84	3.2 %
carbon powder (particle size 20 nm)	4.0 %
Sudan black B	4.0 %
METHIDATHION	4.0 %

# 2006418

- 24 -

29-dodecenyl acetate	}	(pheromone mixture)	0.05 %
dodecanyl acetate			0.05 %
octadecanyl acetate			0.1 %
hexane			2.9 %

The prepared formulation is tested in a wind tunnel against *Eupoecilia ambiguella* males in accordance with the method described in Example 1. With droplets of the above formulation that had been treated for 7 days in the manner described in Example 1, evaluation indicated a mortality of from 75 to 100 %.

Example 5: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	81.2 %
Aerosil COK 84	3.2 %
carbon black powder	4.0 %
Sudan black B	4.0 %
Furathiocarb	4.5 %
hexane	3.0 %
pheromone	0.1 %

Composition of the pheromone used:

Proportions by weight:

(Z)9-tetradecenyl acetate	9
(Z)11-tetradecanyl acetate	1
(Z)9-tetradecenol	1
(Z)11-tetradecenol	0.2

In a field test with the moth *Adoxophyes reticulana*, 1 droplet of 100 µl of the formulation is placed in the middle of a paper sheet 16 x 16 cm coated with glue (polybutylene). The paper is secured around an approximately 5 cm thick branch of an apple tree about 2 m above the ground and the moths that have been attracted by the formulation droplet and caught

on the sticky paper are counted weekly. A rubber stopper treated with the same amount of pheromone, which is also secured to the middle of a sticky paper, is used as a control. 6 replicates were carried out per variant:

Evaluation point of time (No. of days after application)	Number of moths caught (period of capture 1 week, 6 traps)	
	formulation	rubber stopper (control)
14	67	7
70	17	6

Example 6: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	81.2 %
Aerosil COK 84	3.2 %
carbon black powder	4.0 %
Sudan black B	4.0 %
Furathiocarb	4.5 %
hexane	3.0 %
pheromone	0.1 %

Composition of the pheromone used:

Proportions by weight:

(Z)9-dodecenyl acetate	1
dodecanyl acetate	1
octadecanyl acetate	2

In a field test with the vine moth *Eupoecilia ambiguella* 1 droplet of 100 µl is placed in the middle of a paper sheet 16 x 16 cm coated with glue (polybutylene). The paper is secured around a vine at eye level, and the moths that have been attracted by the formulation droplet and caught on the sticky paper are counted weekly. A rubber stopper treated with the same amount of pheromone, which is also secured to the middle of a sticky paper, is used as a control. 6 replicates were carried out per variant:

Evaluation-point of time (No. of days after applica- tion)	Number of moths caught (period of capture 1 week, 6 traps)	
	formulation	rubber peg (control)
14	19	17

Example 7: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	83.6 %
Aerosil COK 84	3.3 %
carbon black powder	4.2 %
Sudan black B	4.2 %
Deltamethrin	4.2 %
hexane	0.4 %
pheromone	0.2 %

Composition of the pheromone used:

Proportions by weight:

Z,Z-7,11-hexadecadienyl acetate	1
Z,E-7,11-hexadecadienyl acetate	1

In a field test with the Pink Bollworm (*Pectinophora gossypiella*), an area of 0.75 ha of cotton is treated with droplets, each of 50  $\mu$ l, of the above formulation (approximately 10,000 droplets/ha). The action is ascertained as a percentage of the capsules attacked by larvae. Figures for the attack of the treated plot compared with an untreated control plot are as follows:

Evaluation point of time (No. of days after applica- tion)	% attack by <i>P. gossypiella</i>	
	treated	untreated (control)
30	0.14 %	3.3 %

Example 8: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	85.4 %
Aerosil COK 84	3.4 %
carbon black powder	4.3 %
Sudan black B	4.3 %
Deltamethrin	0.9 %
hexane	1.5 %
pheromone	0.2 %

Composition of the pheromone used:

Proportions by weight:

(+) <i>cis</i> -2-isopropenyl-1-methylcyclobutane ethanol	40
( <i>Z</i> )-3,3-dimethyl- $\Delta$ -1, $\beta$ -cyclohexane ethanol	30
( <i>Z</i> )-3,3-dimethyl- $\Delta$ -1, $\alpha$ -cyclohexane acetaldehyde	15
( <i>E</i> )-3,3-dimethyl- $\Delta$ -1, $\alpha$ -cyclohexane acetaldehyde	15

Droplets each of 100  $\mu$ l are placed on an aluminium sheet and exposed under a UV lamp for 3 days. Adult *Anthonomus grandis* are then brought into contact with the formulation for a short period, and the mortality is observed after 24 hours.

An analogous formulation without Deltamethrin is used as a control.

Exposure of the formulation (No. of days under UV)	% mortality of <i>A. grandis</i> 24 hours after exposure	
	treated	untreated (control)
3	100 %	5 %

# 2006418

- 28 -

Example 9: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	83.7 %
Aerosil COK 84	3.4 %
carbon black powder	4.3 %
Sudan black B	4.3 %
Methidathion	2.7 %
hexane	1.5 %
8-methyl-2-decanol propanoate (pheromone)	0.1 %

Droplets each of 100 µl are placed on an aluminium sheet and exposed under a UV lamp for 5 days. Adult *Diabrotica balteata* are then brought into contact with the formulation for a short period, and the mortality is observed after 24 hours.

An analogous formulation without Methidathion is used as a control.

Exposure of the formulation (No. of days under UV)	% mortality of <i>D. balteata</i> 24 hours after exposure	
	treated	untreated (control)
5	100 %	0 %

Example 10: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	85.4 %
Aerosil COK 84	3.4 %
carbon black powder	4.3 %
Sudan black B	4.3 %
Cypermethrin	0.9 %
hexane	1.5 %
estragol	0.1 %
4-methoxycinnamaldehyde (pheromone)	0.1 %

Droplets each of 100  $\mu$ l are placed on an aluminium sheet and exposed under a UV lamp for 4 days. In each case one leg of an adult *Diabrotica balteata* is then brought into contact with the formulation for a short period, and the mortality is observed after 24 hours. An analogous formulation without Cypermethrin is used as a control.

Exposure of the formulation (No. of days under UV)	% mortality of <i>D. balteata</i> 24 hours after exposure	
	treated	untreated (control)
4	100 %	0 %

Example 11: A formulation having a viscosity of 20,000 cp is prepared from the following constituents in accordance with the above Example 1:

Tinuvin 171	27.2 %
quartz sand (particle size < 0.7 mm)	66.6 %
Aerosil COK 84	1.1 %
carbon black powder	1.4 %
Sudan black B	1.4 %
Furathiocarb	1.4 %
hexane	0.9 %
E,E-8,10-dodecadienol (pheromone)	0.03 %

Droplets each of 100  $\mu$ l are placed on an aluminium sheet, exposed under a UV lamp for 49 days and then tested in the wind tunnel. These droplets still exhibit a very good attracting effect and insecticidal activity in the wind tunnel. 100 % of the tested males of *Cydia pomonella* are attracted by such a droplet, fly to it and touch it. After 24 hours the mortality of these males is 100 %. The wind tunnel tests are carried out as indicated in Example 1.

Examples 12 to 20: The following formulations are prepared as described in Example 1, and tested against *Cydia pomonella* according to the methods described in Examples 1 to 3, mortality rates of from 75 to 100 % being achieved:

Example

No.

12	-	Tinuvin 171 (see above data)	90.2 %
		Aerosil COK 84	4.1 %
		Furathiocarb	4.5 %
		Codlemon	0.1 %
		sebacic acid dibutyl ester	1.1 %
13	-	Tinuvin 1130 (see above data)	84.9 %
		Aerosil COK 84	4.2 %
		carbon powder	4.2 %
		Furathiocarb	4.2 %
		Codlemon	0.1 %
		ethanol	2.4 %
14	-	Tinuvin 1130	80.9 %
		Aerosil COK 84	3.6 %
		carbon powder	4.0 %
		Sudan black B	4.0 %
		Furathiocarb	4.0 %
		Codlemon	0.1 %
		ethanol	3.4 %
15	-	Tinuvin 171	81.0 %
		Aerosil COK 84	4.0 %
		carbon black (particle size 13 nm)	4.0 %
		Sudan black B	4.0 %
		Furathiocarb	4.0 %
		Codlemon	0.1 %
		sebacic acid dibutyl ester	2.9 %

16	-	Tinuvin 171	84.9 %
		Aerosil COK 84	3.4 %
		Furathiocarb	4.3 %
		Cibacetyyellow 2CG (dyestuff)	4.3 %
		Codlemon	0.1 %
		sebacic acid dibutyl ester	3.0 %
17	-	Tinuvin 171	89.7 %
		Aerosil COK 84	4.0 %
		Furathiocarb	4.0 %
		Erythrosin (dyestuff)	1.0 %
		Codlemon	0.1 %
		hexane	1.2 %
18	-	Tinuvin 171	80.2 %
		Tinuvin 765	1.6 %
		Aerosil COK 84	3.2 %
		carbon powder (particle size 20 nm)	4.0 %
		Sudan black B	4.0 %
		Furathiocarb	4.0 %
		Codlemon	0.1 %
		hexane	3.9 %
19	-	Tinuvin 109	80.2 %
		Tinuvin 765	1.6 %
		Aerosil COK 84	3.2 %
		carbon powder (particle size 20 nm)	4.0 %
		Sudan black B (dyestuff)	4.0 %
		Furathiocarb	4.0 %
		Codlemon	0.1 %
		hexane	2.9 %

# 2006418

- 32 -

20	-	Tinuvin 171	61.0 %
		Aerosil COK 84	10.0 %
		Furathiocarb	4.5 %
		Codlemon	4.5 %
		hexane	10.0 %

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A viscous, non-hardening composition for controlling harmful insects and representatives of the order Acarina that is resistant to water and weather and having a viscosity of at least about 20,000 cp which comprises:

a) about 0.01-30% by weight of a pheromone or kairomone;

b) about 0.1-10% of a pesticidal compound that is compatible with other constituents of said composition; and

c) about 51-98% of an ultraviolet-absorber having an absorption range of from 270 to 400 nm selected from the group consisting of 2-H-benzotriazoles, 2-hydroxyalkoxybenzophenones, oxalanilides, cinnamic acid ester and triazines;

wherein the pheromone or kairomone is slowly released in a biologically effective amount over a period of time; and

wherein said composition is distributable for direct application without dilution as droplets or spots.

2. A composition according to claim 1, wherein the composition additionally contains a viscosity-regulating thickener or solvent, antioxidant, dyestuff or colouring pigment, optical brightener, tackifiers or surface-active substance.

3. A composition according to any one of claims 1 or 2, additionally comprising from 1 to 47% by weight of a viscosity-regulating thickener or solvent.

4. A composition according to any one of claims 1 to 3, which has permanently adhesive properties.

21489-7899

- 34 -

5. A composition according to any one of claims 1 to 4, which contains as additional UV absorbent finely particulate carbon.
6. A composition according to any one of claims 1 to 5, wherein the slow or controlled release of the pheromone or kairomone takes place over a period of up to 15 weeks.
7. A composition according to claim 7, which contains the pheromone Codlemon.
8. A method of controlling harmful insects and representatives of the order Acarina, which comprises distributing within the area to be protected against pest attack a composition according to any one of the preceding claims 1 to 7 in the form of a predetermined number of discrete droplets or droplet-like units or spots, the pheromone or kairomone being slowly released in a biologically effective amount over a prolonged period into the surrounding atmosphere.
9. A method according to claim 8 which comprises distributing the composition over an agricultural crop area.
10. A method according to claim 9 which comprises distributing the composition uniformly over the crop area to be protected in the form of droplets or droplet-like units or spots that adhere to a substrate and are of the order of magnitude of from 1 to 100  $\mu$ l.
11. A method according to claim 9 or 10 which comprises distributing over the crop area to be protected from 50 to 100,000,000 droplets or droplet-like units or spots per 1000 m<sup>2</sup>.

12. A method according to any one of claims 9 to 11 which comprises applying droplets or droplet-like units or spots substantially to the plants to be protected.

FETHERSTONHAUGH & CO.  
OTTAWA, CANADA

PATENT AGENTS