

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 2831113 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 07 K 16/26 (2006.01)** **A 61 K 39/395 (2006.01)** **A 61 P 29/00 (2006.01)**
C 07 K 7/18 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2018-06-25**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-03-14**
- (86) Europæisk ansøgning nr.: **13715049.6**
- (86) Europæisk indleveringsdag: **2013-03-15**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-02-04**
- (86) International ansøgning nr.: **US2013031836**
- (87) Internationalt publikationsnr.: **WO2013148296**
- (30) Prioritet: **2012-03-28 US 201261616845 P** **2013-02-04 FR 1350953**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **SANOFI, 54 rue La Boétie, 75008 Paris, Frankrig**
- (72) Opfinder: **KOMINOS, Dorothea, 99 Rolling Hill Drive, Millington, NJ 07946, USA**
ZHANG, Jie, 6 Alexis Court, Bridgewater, NJ 08807, USA
PRITSKER, Alla, 142 Bramble Drive, Morganville, NJ 07751, USA
SUBRAMANIAN, Govindan, 28 Boice Lane, Belle Mead, NJ 08502, USA
CHEN, Xin, 6 Troy Drive, Short Hills, NJ 07078-1334, USA
Li, Han, 388 Twig Lane, Yardley, PA 19067, USA
DAVISON, Matthew, 109 Rebecca Road, Whitinsville, MA 01588, USA
BAURIN, Nicolas, 35 rue de la Bretoniere, 91180 Arpajon, Frankrig
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **ANTISTOFFER MOD BRADYKININ-B1-RECEPTORLIGANDER**
- (56) Fremdragne publikationer:
BEDI G S ET AL: "Monoclonal antibodies to bradykinin inhibit smooth muscle contractile action of bradykinin", BIOCHIMICA ET BIOPHYSICA ACTA - GENERAL SUBJECTS, ELSEVIER SCIENCE PUBLISHERS, NL, vol. 842, no. 1, 27 September 1985 (1985-09-27), pages 90-99, XP025852771, ISSN: 0304-4165, DOI: 10.1016/0304-4165(85)90298-3 [retrieved on 1985-09-27]
U. HILGENFELDT ET AL: "Strategy of Measuring Bradykinin and Kallidin and Their Concentration in Plasma and Urine", ANALYTICAL BIOCHEMISTRY, vol. 228, no. 1, 1 June 1995 (1995-06-01), pages 35-41, XP055064491, ISSN: 0003-2697, DOI: 10.1006/abio.1995.1311
CARRETERO O A ET AL: "Measurement of urinary kallikrein activity by kinin radioimmunoassay", BIOCHEMICAL PHARMACOLOGY, PERGAMON, OXFORD, GB, vol. 25, no. 20, 15 October 1976 (1976-10-15), pages 2265-2270, XP025509763, ISSN: 0006-2952, DOI: 10.1016/0006-2952(76)90008-3 [retrieved on 1976-10-15]

Fortsættes ...

GEPPETTI P ET AL: "Kallidin applied to the human nasal mucosa produces algescic response not blocked by capsaicin desensitization", REGULATORY PEPTIDES, ELSEVIER SCIENCE BV, NL, vol. 33, no. 3, 17 May 1991 (1991-05-17), pages 321-329, XP025208931, ISSN: 0167-0115, DOI: 10.1016/0167-0115(91)90234-8 [retrieved on 1991-05-17]

DESCRIPTION

BACKGROUND

[0001] The bradykinin B1 receptor has been implicated in pathogenesis of inflammatory disease and chronic pain. By modulating tissue inflammation and renal fibrosis, the B1 receptor has also been associated with pathogenesis of acute kidney injury as well as chronic kidney diseases which are the main causes of end-stage renal failure.

In humans, the major agonists of the bradykinin B1 receptor are the kinins. Kinins are bioactive peptides produced from the proteolytic cleavage of kininogen proteins. The major kinin agonists of bradykinin B1 receptor are the decapeptide Kallidin, and the nonapeptide des-Arg₁₀-Kallidin (formed by the proteolytic cleavage the c-terminal arginine form Kallidin). Therefore, agents that can inhibit the binding of Kallidin and des-Arg₁₀-Kallidin to the bradykinin B1 receptor have the potential to treat or prevent bradykinin B1 receptor-mediated pathologies.

[0002] Accordingly, there is a need in the art for novel agents that inhibit the binding of Kallidin and des-Arg₁₀-Kallidin to the bradykinin B1 receptor for use in the treatment of bradykinin B1 receptor-mediated human pathologies.

[0003] Bedi et al. (Biochimica et Biophysica Acta, vol. 842, no. 1, 1985, pages 90-99) discloses monoclonal antibodies that bind to bradykinin.

[0004] Hilgenfeldt et al. (Analytical Biochemistry, vol. 228, no. 1, 1995, pages 35-41) describes an antiserum specific for kallidin.

[0005] Carretero et al. (Biochemical Pharmacology, vol. 25, no. 20, 1976, pages 2265-2270) describes anti kallidin antisera.

[0006] Geppetti et al. (Regulatory Peptides, vol. 33, no. 3, 1991, pages 321-329) describes the analgesic effect of kallidin on human nasal mucose.

SUMMARY

[0007] The invention relates to an isolated monoclonal antibody or antigen binding fragment thereof that specifically binds to Kallidin or des-Arg₁₀-Kallidin but not to Bradykinin or des-Arg₉-Bradykinin, wherein the antibody or antigen binding fragment comprises:

1. i) a heavy chain variable domain comprising a heavy chain complementarity determining region 3 (HCDR3) amino acid sequence, a heavy variable complementary determining region 2 (HCDR2) amino acid sequence, and a heavy chain variable complementary determining region 1 (HCDR1) amino acid sequence selected from the group consisting of:
 1. a)

SEQ ID NO: 7 [X₁YX₂X₃DX₄HAMX₅Y], wherein

X₁ is Y, F or H,

X₂ is R, D, A, V, L, I, M, F, Y or W,

X₃ is Y, F, W or H,

X₄ is D, E or Y, and,

X₅ is D or E;

SEQ ID NO: 8 [YFX₁PX₂NGNTGYNQKFRG], wherein

X₁ is D, R, A, V, L, I, M, F, Y or W, and

X₂ is Y, D, E, N, or Q; and

SEQ ID NO: 9 [GYSFTDYX₁IY], wherein X₁ is N, W or Y;

respectively,

2. b)

SEQ ID NO: 63 [X₁EYDGX₂YX₃X₄LDX₅], wherein

X₁ is W or F,

X₂ is N or no amino acid;

X₃ is Y or S,

X₄ is D or P, and

X₅ is F or Y;

SEQ ID NO: 64 [WX₁DPENGDX₂X₃YAPKFQG], wherein

X₁ is I, or V,

X₂ is T, or S, and

X₃ is G, or D; and

SEQ ID NO: 65 [GFNIKDYX₁H], wherein X₁ is L, or M;

respectively,

3. c)

SEQ ID NO: 13;

SEQ ID NO: 14; and

SEQ ID NO: 15;

respectively,

4. d)

SEQ ID NO: 32;

SEQ ID NO: 33; and

SEQ ID NO: 34;

respectively,

5. e)

SEQ ID NO: 40;

SEQ ID NO: 41; and

SEQ ID NO: 42;

respectively,

6. f)

SEQ ID NO: 47;

SEQ ID NO: 48;

SEQ ID NO: 49;

respectively,

and

7. g)

SEQ ID NO: 55;

SEQ ID NO: 56;

SEQ ID NO: 57;

respectively; and

2. ii) a light chain variable domain comprising a light chain complementarity determining region 3 (LCDR3) amino acid sequence, a light chain complementarity determining region 2 (LCDR2) amino acid sequence, and a light chain complementarity determining region 1 (LCDR1) amino acid sequence selected from the group consisting of:

h)

SEQ ID NO: 10 [QQ X₁ X₂S X₃P X₄T], wherein

X₁ is Y, F or H,

X₂ is Y, F, H or W,

X_3 is Y, F, T or H, and,

X_4 is W, Y, F, H or L:

SEQ ID NO: 11 [WASTRX₁], wherein X_1 is E, D, Q or N; and

SEQ ID NO: 12 [KSSQSLL X_1 SSNQKN X_2 LA], wherein

X_1 is W, H, Y or F, and

X_2 is H or Y;

respectively,

i)

SEQ ID NO: 66 [QX₁X₂X₃SX₄PX₅T], wherein

X_1 is Q or N,

X_2 is Y, F, D or H,

X_3 is Y, F, H or W,

X_4 is Y, F, T or H, and

X_5 is W, Y, F, H or L;

SEQ ID NO: 67 [X₂ASTRX₂], wherein

X_1 is W or G, and

X_2 is E, D, Q or N; and

SEQ ID NO: 68 [KSSQSLLX₁X₂SX₃QX₄NX₅LA], wherein

X_1 is W, H, Y or F,

X_2 is S or G,

X_3 is N or D,

X_4 is K or R,

X_5 is H or Y;

respectively,

j)

SEQ ID NO: 69 [X₁QGTHFPYT], wherein X_1 is L or M;

SEQ ID NO: 36; and

SEQ ID NO: 70 [KSSQSLLYSNGX1TYLN], wherein X1 is K or E;

respectively,

k)

SEQ ID NO: 16;

SEQ ID NO: 17; and

SEQ ID NO: 18;

respectively,

l)

SEQ ID NO: 35;

SEQ ID NO: 36; and

SEQ ID NO: 37;

respectively,

m)

SEQ ID NO: 43;

SEQ ID NO: 17; and

SEQ ID NO: 44;

respectively,

n)

SEQ ID NO: 50;

SEQ ID NO: 51; and

SEQ ID NO: 52;

respectively,

and

o)

SEQ ID NO: 58;

SEQ ID NO: 59; and

SEQ ID NO: 60;

respectively.

[0008] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof of claim 1 comprises

1. a) a heavy chain variable domain comprising the HCDR3, HCDR2 and HCDR1 region amino sequences set forth in SEQ ID NOs: 13, 14, and 15, respectively, and one or more amino acid substitution at positions selected from the group consisting of H1, H5, H9, H11, H12, H16, H38, H40, H41, H43, H44, H66, H75, H79, H81, H82A, H83, H87, and H108, according to Kabat; and
2. b) a light chain variable domain comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in SEQ ID NOs: 16, 17, and 18, respectively, and one or more amino acid substitution at positions selected from the group consisting of L5, L9, L15, L18, L19, L21, L22, L43, L63, L78, L79, L83, L85, L100 and L104, according to Kabat.

[0009] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof of claim 1 comprises the heavy chain and light chain variable domain amino acid sequences set forth in SEQ ID NO: 24 and 30, respectively.

[0010] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof is in a pharmaceutical composition with one or more pharmaceutically acceptable carrier(s).

[0011] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof of claim 1 comprises:

a heavy chain variable region domain amino acid sequence, and a light chain variable domain amino acid sequence with at least 90% identity to an amino acid sequence selected from the group consisting of:

1. a) SEQ ID NOs: 19 and 26, respectively,
2. b) SEQ ID NOs: 20 and 27, respectively,
3. c) SEQ ID NOs: 21 and 28, respectively,
4. d) SEQ ID NOs: 22 and 28, respectively,
5. e) SEQ ID NOs: 23 and 29, respectively,
6. f) SEQ ID NOs: 24 and 30, respectively,
7. g) SEQ ID NOs: 25 and 31, respectively,
8. h) SEQ ID NOs: 38 and 39, respectively,
9. i) SEQ ID NOs: 45 and 46, respectively,
10. j) SEQ ID NOs: 53 and 54, respectively,
11. k) SEQ ID NOs: 61 and 62, respectively.

[0012] In an embodiment, the isolated monoclonal antibody or antigen binding fragment thereof of claim 1 comprises:

a heavy chain variable domain comprising an amino acid sequence, and a light chain variable domain amino acid sequence selected from the group consisting of:

1. a) SEQ ID NOs: 19 and 26, respectively,
2. b) SEQ ID NOs: 20 and 27, respectively,
3. c) SEQ ID NOs: 21 and 28, respectively,
4. d) SEQ ID NOs: 22 and 28, respectively,
5. e) SEQ ID NOs: 23 and 29, respectively,
6. f) SEQ ID NOs: 24 and 30, respectively,
7. g) SEQ ID NOs: 25 and 31, respectively,
8. h) SEQ ID NOs: 38 and 39, respectively,
9. i) SEQ ID NOs: 45 and 46, respectively,
10. j) SEQ ID NOs: 53 and 54, respectively,
11. k) SEQ ID NOs: 61 and 62, respectively.

[0013] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof comprises:

1. a) a heavy chain variable domain comprising the consensus HCDR3, HCDR2 and HCDR1 region amino sequences set forth in SEQ ID NOs: 7, 8, and 9, respectively; and
2. b) a light chain variable domain comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in SEQ ID NOs: 10, 11, and 12, respectively.

[0014] In a embodiment, the isolated monoclonal antibody or antigen binding fragment thereof comprises:

1. a) a heavy chain variable domain comprising the HCDR3, HCDR2 and HCDR1 region amino sequences set forth in SEQ ID NOs: 13, 14, and 15, respectively; and
2. b) a light chain variable domain comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in SEQ ID NOs: 16, 17, and 18, respectively.

[0015] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof comprises:

1. a) a heavy chain variable domain comprising the HCDR3, HCDR2 and HCDR1 region amino sequences set forth in SEQ ID NOs: 32, 33, and 34, respectively; and
2. b) a light chain variable domain comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in SEQ ID NOs: 35, 36, and 37, respectively.

[0016] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof that specifically binds to Kallidin or des-Arg₁₀-Kallidin but not to Bradykinin or des-Arg₉-Bradykinin of claim 1, wherein the antibody or antigen binding fragment comprises:

1. a) a heavy chain variable domain comprising the HCDR3, HCDR2 and HCDR1 region amino sequences set forth in SEQ ID NOs: 40, 41 and 42, respectively; and
2. b) a light chain variable domain comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in SEQ ID NOs: 43, 17, and 44, respectively.

[0017] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof comprises:

1. a) a heavy chain variable domain comprising the HCDR3, HCDR2 and HCDR1 region amino sequences set forth in SEQ ID NOs: 47, 48, and 49, respectively; and
2. b) a light chain variable domain comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in SEQ ID NOs: 50, 51, and 52, respectively.

[0018] In an embodiment of the invention, the isolated monoclonal antibody or antigen binding fragment thereof comprises:

1. a) a heavy chain variable domain comprising the HCDR3, HCDR2 and HCDR1 region amino sequences set forth in SEQ ID NOs: 55, 56, and 57, respectively; and
2. b) a light chain variable domain comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in SEQ ID NOs: 58, 59, and 60, respectively.

[0019] The present application further discloses nucleic acids encoding anti-Kallidin and des-Arg₁₀-Kallidin antibodies, recombinant expression vectors and host cells for making such antibodies, or fragments thereof. Methods of using antibodies, or fragments thereof, of the disclosure to detect Kallidin and des-Arg₁₀-Kallidin or to modulate Kallidin and des-Arg₁₀-Kallidin activity, either *in vitro* or *in vivo*, are also disclosed. The present application also discloses methods of making antibodies that specifically bind to des-Arg₉- Bradykinin and des-Arg₁₀-Kallidin-like peptide.

[0020] Accordingly, in one aspect the present disclosure provides an isolated monoclonal antibody or antigen binding fragment thereof that:

1. a) specifically binds to Kallidin or des-Arg₁₀-Kallidin but not to Bradykinin or des-Arg₉-Bradykinin;
2. b) specifically binds to Kallidin or des-Arg₁₀-Kallidin with a K_D of less than 1×10^{-10} M;
3. c) specifically binds to Kallidin or des-Arg₁₀-Kallidin with a K_{off} of less than 1×10^4 s⁻¹; or

4. d) specifically binds to Kallidin or des-Arg₁₀-Kallidin and inhibits binding to the bradykinin B1 receptor.

[0021] In one embodiment, the antibody or antigen binding fragment thereof binds to the N-terminal Lysine residue of Kallidin or des-Arg₁₀-Kallidin.

[0022] In another embodiment, the antibody or antigen binding fragment thereof inhibits the binding of Kallidin or des-Arg₁₀-Kallidin to a bradykinin-1 receptor.

[0023] In another embodiment, the antibody or antigen binding fragment thereof binds specifically to mouse Kallidin-like peptide (KLP).

[0024] In another embodiment, the antibody or antigen binding fragment thereof comprises a heavy chain variable domain comprising an HCDR3 amino acid sequence selected from the group consisting of:

1. a) SEQ ID NO: 7 [X₁Y X₂ X₃D X₄HAM X₅Y], wherein

X₁ is Y, F or H,

X₂ is R, D, A, V, L, I, M, F, Y or W,

X₃ is Y, F, W or H,

X₄ is D, E or Y, and,

X₅ is D or E;

2. b) SEQ ID NO: 63 [X₁EYDGX₂YX₃X₄LDX₅], wherein

X₁ is W or F,

X₂ is N or no amino acid;

X₃ is Y or S,

X₄ is D or P, and

X₅ is F or Y;

3. c) SEQ ID NO: 13;

4. d) SEQ ID NO: 32;

5. e) SEQ ID NO: 40;

6. f) SEQ ID NO: 47; and

7. g) SEQ ID NO: 55.

[0025] In another embodiment, the antibody or antigen binding fragment thereof comprises an HCDR2 amino acid sequence selected from the group consisting of:

1. a) SEQ ID NO: 8 [YFX₁PX₂NGNTGYNQKFRG], wherein

X₁ is D, R, A, V, L, I, M, F, Y or W, and

X₂ is Y, D, E, N, or Q;

2. b) SEQ ID NO: 64 [WX₁DPENGDX₂X₃YAPKFQG], wherein

X_1 is I, or V,

X_2 is T, or S, and

X_3 is G, or D;

3. c) SEQ ID NO: 14

4. d) SEQ ID NO: 33;

5. e) SEQ ID NO: 41;

6. f) SEQ ID NO: 48; and

7. g) SEQ ID NO: 56.

[0026] In another embodiment, the antibody or antigen binding fragment thereof comprises an HCDR1 amino acid sequence selected from the group consisting of:

1. a) SEQ ID NO: 9 [GYSFTDY X_1 IY], wherein X_1 is N, W or Y;

2. b) SEQ ID NO: 65 [GFNIKDY X_1 H], wherein X_1 is L, or M;

3. c) SEQ ID NO: 15;

4. d) SEQ ID NO: 34;

5. e) SEQ ID NO: 42;

6. f) SEQ ID NO: 49; and

7. g) SEQ ID NO: 57.

[0027] In another embodiment, the antibody or antigen binding fragment thereof comprises a light chain variable domain comprising an LCDR3 amino acid sequence selected from the group consisting of:

1. a) SEQ ID NO: 10 [QQ X_1 X_2 S X_3 P X_4 T], wherein

X_1 is Y, F or H,

X_2 is Y, F, H or W,

X_3 is Y, F, T or H, and,

X_4 is W, Y, F, H or L;

2. b) SEQ ID NO: 66 [Q X_1 X_2 X_3 S X_4 P X_5 T], wherein

X_1 is Q or N,

X_2 is Y, F, D or H,

X_3 is Y, F, H or W,

X_4 is Y, F, T or H, and

X_5 is W, Y, F, H or L;

3. c) SEQ ID NO: 69 [X_1 QGTHTFPYT], wherein X_1 is L or M;

4. d) SEQ ID NO: 16;

5. e) SEQ ID NO: 35;

6. f) SEQ ID NO: 43;

7. g) SEQ ID NO: 50; and

8. h) SEQ ID NO: 58.

[0028] In another embodiment, the antibody or antigen binding fragment thereof comprises an LCDR2 amino acid sequence selected from the group consisting of:

1. a) SEQ ID NO: 11 [WASTRX₁], wherein X₁ is E, D, Q or N;
2. b) SEQ ID NO: 67 [X₂ASTRX₂], wherein
X₁ is W or G, and
X₂ is E, D, Q or N;
3. c) SEQ ID NO: 17;
4. d) SEQ ID NO: 36;
5. e) SEQ ID NO: 51; and
6. f) SEQ ID NO: 59.

[0029] In another embodiment, the antibody or antigen binding fragment thereof comprises an LCDR1 amino acid sequence selected from the group consisting of:

- a) SEQ ID NO: 12 [KSSQSLL X₁SSNQKN X₂LA], wherein
X₁ is W, H, Y or F, and
X₂ is H or Y;
- b) SEQ ID NO: 68 [KSSQSLLX₁X₂SX₃QX₄NX₅LA], wherein
X₁ is W, H, Y or F,
X₂ is S or G,
X₃ is N or D,
X₄ is K or R,
X₅ is H or Y.
- c) SEQ ID NO: 70 [KSSQSLLYSNGX₁TYLN], wherein X₁ is K or E;
- b) SEQ ID NO: 18;
- c) SEQ ID NO: 37;
- d) SEQ ID NO: 44;
- e) SEQ ID NO: 52; and
- f) SEQ ID NO: 60.

[0030] In another embodiment, the antibody or antigen binding fragment comprises a light chain variable domain comprising an LCDR3 amino acid sequence selected from the group consisting of:

1. a) SEQ ID NO: 10 [QQ X₁ X₂S X₃P X₄T], wherein

- X_1 is Y, F or H,
 X_2 is Y, F, H or W,
 X_3 is Y, F, T or H, and,
 X_4 is W, Y, F, H or L:
2. b) SEQ ID NO: 66 [QX₁X₂X₃SX₄PX₅T], wherein

X_1 is Q or N,
 X_2 is Y, F, D or H,
 X_3 is Y, F, H or W,
 X_4 is Y, F, T or H, and
 X_5 is W, Y, F, H or L;
 3. c) SEQ ID NO: 69 [X₁QGTHFPYT], wherein X_1 is L or M;
 4. d) SEQ ID NO: 16;
 5. e) SEQ ID NO: 35;
 6. f) SEQ ID NO: 43;
 7. g) SEQ ID NO: 50; and
 8. h) SEQ ID NO: 58.

[0031] In another embodiment, the antibody or antigen binding fragment thereof comprises an LCDR2 amino acid sequence selected from the group consisting of:

1. a) SEQ ID NO: 11 [WASTRX₁], wherein X_1 is E, D, Q or N;
2. b) SEQ ID NO: 67 [X₂ASTRX₂], wherein

X_1 is W or G, and
 X_2 is E, D, Q or N;
3. c) SEQ ID NO: 17;
4. d) SEQ ID NO: 36;
5. e) SEQ ID NO: 51; and
6. f) SEQ ID NO: 59.

[0032] In another embodiment, the antibody or antigen binding fragment thereof comprises an LCDR1 amino acid sequence selected from the group consisting of:

- a) SEQ ID NO: 12 [KSSQSLL X₁SSNQKN X₂LA], wherein

X_1 is W, H, Y or F, and
 X_2 is H or Y;
- b) SEQ ID NO: 68 [KSSQSLLX₁X₂SX₃QX₄NX₅LA], wherein

X_1 is W, H, Y or F,
 X_2 is S or G,
 X_3 is N or D,

X₄ is K or R,

X₅ is H or Y.

c) SEQ ID NO: 70 [KSSQSLLYSNGX₁TYLN], wherein X₁ is K or E;

b) SEQ ID NO: 18;

c) SEQ ID NO: 37;

d) SEQ ID NO: 44;

e) SEQ ID NO: 52; and

f) SEQ ID NO: 60.

[0033] In another embodiment, the antibody or antigen binding fragment thereof comprises a heavy chain variable region comprising the HCDR3, HCDR2 and HCDR1 region amino sequences set forth in SEQ ID NOs 13, 14, and 15, respectively, and one or more amino acid substitutions at positions selected from the group consisting of H1, H5, H9, H11, H12, H16, H38, H40, H41, H43, H44, H66, H75, H79, H81, H82A, H83, H87, and H108 according to Kabat.

[0034] In another embodiment, the antibody or antigen binding fragment thereof comprises a light chain variable region comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in SEQ ID NOs 16, 17, and 18, respectively, and one or more amino acid substitution at positions selected from the group consisting of L5, L9, L15, L18, L19, L21, L22, L43, L63, L78, L79, L83, L85, L100 and L104, according to Kabat.

[0035] In another embodiment, the antibody or antigen binding fragment thereof comprises a heavy chain variable region amino acid sequence with at least 90% identity to an amino acid sequence selected from the group consisting of SEQ ID NOs: 19, 20, 21, 22, 24, 25, 38, 45, 53, and 61.

[0036] In another embodiment, the antibody or antigen binding fragment thereof comprises a light chain variable domain amino acid sequence with at least 90% identity to an amino acid sequence selected from the group consisting of SEQ ID NOs: 26, 27, 28, 29, 29, 30, 31, 39, 46, 54, and 62.

[0037] In another embodiment, the antibody or antigen binding fragment thereof comprises a light chain variable region amino acid sequence with at least 90% identity to an amino acid sequence selected from the group consisting of SEQ ID NOs: 26, 27, 28, 29, 29, 30, 31, 39, 46, 54, and 62.

[0038] In another embodiment, the antibody or antigen binding fragment thereof comprises a heavy chain variable domain comprising an amino acid sequence selected from the group consisting of: SEQ ID NO: 19, 20, 21, 22, 24, 25, 38, 45, 53, and 61.

[0039] In another embodiment, the antibody or antigen binding fragment thereof comprises a light chain variable domain amino acid sequence selected from the group consisting of: SEQ ID NO: 26,

27, 28, 29, 30, 31, 39, 46, 54, and 62.

[0040] In another embodiment, the antibody or antigen binding fragment thereof comprises an amino acid sequence selected from the group consisting of: SEQ ID NO: 26, 27, 28, 29, 30, 31, 39, 46, 54, and 62.

[0041] In another embodiment, the antibody or antigen binding fragment thereof comprises the heavy chain and light chain variable region amino acid sequences set forth in SEQ ID NO: 19 and 26, SEQ ID NO: 20 and 27, SEQ ID NO: 21 and 28; SEQ ID NO: 22 and 28; SEQ ID NO: 23 and 29; SEQ ID NO: 24 and 30; SEQ ID NO: 25 and 31; SEQ ID NO: 38 and 39, SEQ ID NO: 45 and 46, SEQ ID NO: 53 and 54, or SEQ ID NO: 61 and 62, respectively.

[0042] In another aspect, the disclosure provides an antibody, or antigen binding fragment thereof, that specifically binds to Kallidin and des-Arg₁₀-Kallidin, wherein the antibody, or antigen binding fragment thereof, competes for binding to Kallidin and des-Arg₁₀-Kallidin with an antibody comprising the heavy chain and light chain variable region amino acid sequences set forth in SEQ ID NO: 19 and 26, SEQ ID NO: 38 and 39, SEQ ID NO: 45 and 46, SEQ ID NO: 53 and 54, or SEQ ID NO: 61 and 62, respectively.

[0043] In another aspect, the disclosure provides an isolated monoclonal antibody or antigen binding fragment thereof that competes for binding to Kallidin or des-Arg₁₀-Kallidin with the antibody of any one of the preceding claims, and does not bind to Bradykinin or desArg₉-Bradykinin.

[0044] In another aspect, the disclosure provides an isolated monoclonal antibody or antigen binding fragment thereof that specifically binds to a conformational epitope of kallidin (KD) or desArg₁₀-Kallidin (DAKD) which adopts a Pro4 kink conformation comprising a type II tight turn at Proline 4 of the KD or DAKD). In one embodiment, the Pro 4 kink conformation of KD or DAKD further comprises amino acid repeats of a sigmoid shape which align the hydrophobic side chains of the amino acids in a spatially stacking mode. In another embodiment, the antibody or antigen binding fragment thereof comprises (a) specifically binds Kallidin or des-Arg₁₀-Kallidin but not to Bradykinin or des-Arg₉-Bradykinin; b) specifically binds to Kallidin or des-Arg₁₀-Kallidin with a KD of less than 1×10^{-10} M; c) specifically binds to Kallidin or des-Arg₁₀-Kallidin with a K_{off} of less than 1×10^4 s⁻¹; or d) specifically binds to Kallidin or des-Arg₁₀-Kallidin and inhibits binding to the bradykinin B1 receptor.

[0045] In another aspect, the antibody or antigen binding fragment is conjugated to a diagnostic or therapeutic agent.

[0046] In another aspect, the disclosure provides isolated nucleic acid encoding the amino acid sequence of the antibody, or antigen binding fragment thereof, of the disclosure.

[0047] In another aspect, the disclosure provides recombinant expression vector comprising the nucleic acid of the disclosure.

[0048] In another aspect, the disclosure provides a host cell comprising the recombinant expression vector of the disclosure.

[0049] In another aspect, the disclosure provides a method of producing an antibody that binds specifically to Kallidin and des-Arg₁₀-Kallidin, comprising culturing the host cell of the disclosure under conditions such that an antibody that binds specifically to Kallidin and des-Arg₁₀-Kallidin is produced by the host cell.

[0050] In another aspect, the disclosure provides a pharmaceutical composition comprising the antibody, or antigen binding fragment thereof, of the disclosure and one or more pharmaceutically acceptable carriers.

[0051] In another aspect, the disclosure provides a method for treating a disease or disorder Kallidin or des-Arg₁₀-Kallidin-associated disease or disorder, the method comprising administering to a subject in need of thereof the pharmaceutical composition of the disclosure.

[0052] In one embodiment, the disease or disorder is chronic pain.

[0053] In another aspect, the disclosure provides a method of generating an antibody that specifically binds to des-Arg₉- Bradykinin and des-Arg₁₀-Kallidin-like peptide comprising: immunizing an animal with an immunogen comprising a peptide, wherein the peptide consists of the amino acid sequence set forth in SEQ ID No.11, and wherein the amino terminal arginine of the peptide is indirectly coupled to a carrier moiety through a linker moiety, such that an antibody that specifically binds to des-Arg₉-Bradykinin, des-Arg₁₀-Kallidin and des-Arg₁₀-Kallidin-like peptide is produced by the immune system of the animal.

[0054] In another embodiment, the method further comprises isolating from the animal, the antibody, a nucleic acid encoding the antibody, or an immune cell expressing the antibody.

[0055] In one embodiment, the carrier moiety is a protein. In another embodiment, the protein is Keyhole limpet hemocyanin (KLH). In another embodiment, wherein the linker moiety comprises [Gly-Gly-Gly]_n, wherein n is at least 1.

BRIEF DESCRIPTION OF THE DRAWINGS

[0056]

Figure 1 depicts the results of ELISA assays demonstrating binding of EE1 antibody to kinin peptides.

Figure 2 depicts the results of differential scanning calorimetry measurements of antibody F151.

Figure 3 depicts amino acid sequence alignments of the variable regions of the murine and humanized F151 antibody. All identical residues are listed in the alignment, while homologous residues are identified by "+" sign and non-homologous residues are left blank.

Figure 4 depicts an electron density map of the antigen binding site of the F151 antibody/kallidin complex.

Figure 5 depicts an electron density map of the antigen binding site of the F151 antibody/des-Arg10-kallidin complex.

Figure 6 depicts a ribbon and stick representation of the Fv subunit of F151 bound to kallidin.

Figure 7 depicts an amino acid sequence alignment of the light chain variable regions of exemplary murine anti-kallidin antibodies of the disclosure. Amino acid residues that interact with kallidin are marked with asterisks.

Figure 8 depicts an amino acid sequence alignment of the heavy chain variable regions of exemplary murine anti-kallidin antibodies of the disclosure. Amino acid residues that interact with kallidin are marked with asterisks.

Figure 9 depicts the results of *in vivo* experiments determining the effect of EE1 antibody on formalin-induced acute inflammatory pain.

Figure 10 depicts the results of *in vivo* experiments determining the effect of EE1 antibody on CFA-induced mechanical hypersensitivity.

Figure 11 depicts the results of *in vivo* experiments determining the effect of EE1 antibody on CFA-induced thermal hypersensitivity.

Figure 12 depicts the results of *in vivo* experiments determining the effect of EE1 antibody on CCI-induced mechanical hypersensitivity.

Figure 13 depicts the results of *in vivo* experiments determining the effect of EE1 antibody on CCI-induced thermal hypersensitivity.

Figure 14 depicts schematic maps of VL and VH expression constructs for generating humanized F151 variant HC3a/LC3a with the restriction DNA endonuclease sites presented as deduced sequences in bold and underlined. Panel A depicts the light chain and Panel B depicts the heavy chain.

Figure 15 depicts an alignment of the F151 heavy chain (A) and light chain (B) amino acid sequences with the closest human germline amino acid sequences.

Figure 16 depicts an alignment of the F151 heavy chain (A) and light chain (B) with a heavy chain locus (1-08 & 1-18) and light chain (V IV-B3) locus of the VH1 sub-family. CDR regions and Vernier regions are indicated in boldface and humanizing mutations are underlined.

Figure 17 depicts (A) the secondary and (B) tertiary structure of the main chain polypeptide backbone conformation of kallidin (KD) as bound to F151 antibody which comprises a type II tight turn at Proline 4 (C).

DETAILED DESCRIPTION

[0057] The present disclosure provides antibodies that specifically bind to Kallidin and des-Arg₁₀-Kallidin and prevent binding to the bradykinin B1 receptor. Such antibodies are particularly useful for treating Kallidin and des-Arg₁₀-Kallidin -associated disease or disorders (e.g., pain). The disclosure also provides pharmaceutical compositions, as well as nucleic acids encoding anti-Kallidin and des-Arg₁₀-Kallidin antibodies, recombinant expression vectors and host cells for making such antibodies, or fragments thereof. Methods of using antibodies of the disclosure to detect Kallidin and des-Arg₁₀-Kallidin or to modulate Kallidin and des-Arg₁₀-Kallidin activity, either *in vitro* or *in vivo*, are also encompassed by the disclosure.

I. Definitions

[0058] In order that the present disclosure may be more readily understood, certain terms are first defined.

[0059] As used herein, the term "Kallidin" refers to a peptide comprising or consisting of the amino acid sequence KRPPGFSPFR (SEQ ID NO. 1).

[0060] As used herein, the term "des-Arg₁₀-Kallidin" refers to a peptide comprising or consisting of the amino acid sequence KRPPGFSPF (SEQ ID NO. 2).

[0061] As used herein, the term "mouse Kallidin" or "Kallidin-like peptide" refers to a peptide comprising or consisting of the amino acid sequence RRPPGFSPFR (SEQ ID NO. 3)

[0062] As used herein, the term "mouse des-Arg₁₀-Kallidin" or "des-Arg₁₀Kallidin-like peptide" refers to a peptide comprising or consisting of the amino acid sequence RRPPGFSPF (SEQ ID NO. 4).

[0063] As used herein, the term "Bradykinin" refers to a peptide comprising or consisting of the amino acid sequence RPPGFSPFR (SEQ ID NO. 5).

[0064] As used herein, the term "des-Arg₉-Bradykinin" refers to a peptide comprising or consisting of the amino acid sequence RPPGFSPF (SEQ ID NO. 6).

[0065] As used herein, the term "antibody" refers to immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, as well as multimers thereof (e.g., IgM). Each heavy chain comprises a heavy chain variable region (abbreviated V_H or V_H) and a heavy chain constant region (C_H or C_H). The heavy chain constant region comprises three domains, C_H1, C_H2 and C_H3. Each light chain comprises a light chain variable region (abbreviated V_L) and a light chain constant region (C_L or C_L). The light chain constant region comprises one domain (C_L1). The V_H and V_L regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed with

regions that are more conserved, termed framework regions (FR). Each V_H and V_L is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

[0066] As used herein, the term "antigen-binding fragment" of an antibody includes any naturally occurring, enzymatically obtainable, synthetic, or genetically engineered polypeptide or glycoprotein that specifically binds an antigen to form a complex. Antigen-binding fragments of an antibody may be derived, e.g., from full antibody molecules using any suitable standard techniques such as proteolytic digestion or recombinant genetic engineering techniques involving the manipulation and expression of DNA encoding antibody variable and optionally constant domains. Non-limiting examples of antigen-binding portions include: (i) Fab fragments; (ii) $F(ab')_2$ fragments; (iii) Fd fragments; (iv) Fv fragments; (v) single-chain Fv (scFv) molecules; (vi) dAb fragments; and (vii) minimal recognition units consisting of the amino acid residues that mimic the hypervariable region of an antibody (e.g., an isolated complementarity determining region (CDR)). Other engineered molecules, such as diabodies, triabodies, tetrabodies and minibodies, are also encompassed within the expression "antigen-binding fragment."

[0067] As used herein, the term "CDR" or "complementarity determining region" means the noncontiguous antigen combining sites found within the variable region of both heavy and light chain polypeptides. These particular regions have been described by Kabat et al., J. Biol. Chem. 252, 6609-6616 (1977) and Kabat et al., Sequences of protein of immunological interest. (1991), and by Chothia et al., J. Mol. Biol. 196:901-917 (1987) and by MacCallum et al., J. Mol. Biol. 262:732-745 (1996) where the definitions include overlapping or subsets of amino acid residues when compared against each other. The amino acid residues which encompass the CDRs as defined by each of the above cited references are set forth for comparison. In an embodiment of the disclosure, the term "CDR" is a CDR as defined by Kabat, based on sequence comparisons.

[0068] As used herein the term "framework (FR) amino acid residues" refers to those amino acids in the framework region of an Ig chain. The term "framework region" or "FR region" as used herein, includes the amino acid residues that are part of the variable region, but are not part of the CDRs (e.g., using the Kabat definition of CDRs). Therefore, a variable region framework is between about 100-120 amino acids in length but includes only those amino acids outside of the CDRs.

[0069] As used herein, the term "specifically binds to" refers to the ability of an antibody or an antigen-binding fragment thereof to bind to an antigen with an K_d of at least about 1×10^{-6} M, 1×10^{-7} M, 1×10^{-8} M, 1×10^{-9} M, 1×10^{-10} M, 1×10^{-11} M, 1×10^{-12} M, or more. The term also encompasses refers to the ability of an antibody or an antigen-binding fragment thereof to bind to an antigen with an affinity that is at least two-fold greater than its affinity for a nonspecific antigen. It shall be understood, however, that an antibody, or an antigen-binding fragment thereof, is capable of specifically binding to two or more antigens which are related in sequence (e.g., Kallidin or des-Arg10-Kallidin and mouse Kallidin or des-Arg10-Kallidin).

[0070] As used herein, the term "antigen" refers to the binding site or epitope recognized by an antibody or antigen binding fragment thereof.

[0071] As used herein, the term "vector" is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid," which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. The terms, "plasmid" and "vector" may be used interchangeably. However, the disclosure is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

[0072] As used herein, the term "host cell" is intended to refer to a cell into which a recombinant expression vector has been introduced. It should be understood that this term is intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein.

[0073] As used herein, the term "treat," "treating," and "treatment" refer to therapeutic or preventative measures described herein. The methods of "treatment" employ administration to a subject, an antibody or antigen binding fragment of the present disclosure, for example, a subject having a Kallidin and des-Arg₁₀-Kallidin-associated disease or disorder (e.g. an inflammatory disease) or predisposed to having such a disease or disorder, in order to prevent, cure, delay, reduce the severity of, or ameliorate one or more symptoms of the disease or disorder or recurring disease or disorder, or in order to prolong the survival of a subject beyond that expected in the absence of such treatment.

[0074] As used herein, the term "Kallidin or des-Arg₁₀-Kallidin -associated disease or disorder" includes disease states and/or symptoms associated with a disease state, where altered levels or activity of Kallidin or des-Arg₁₀-Kallidin are found. Exemplary Kallidin or des-Arg₁₀-Kallidin-associated diseases or disorders include, but are not limited to, pain and fibrosis.

[0075] As used herein, the term "effective amount" refers to that amount of an antibody or an antigen binding fragment thereof that binds Kallidin or des-Arg₁₀-Kallidin, which is sufficient to effect treatment, prognosis or diagnosis of a Kallidin or des-Arg₁₀-Kallidin-associated disease or disorder, as described herein, when administered to a subject. A therapeutically effective amount will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. The dosages for administration can range from, for example, about 1 ng to about 10,000 mg, about 1 ug to about 5,000 mg, about 1 mg to about

1,000 mg, about 10 mg to about 100 mg, of an antibody or antigen binding fragment thereof, according to the disclosure. Dosage regimens may be adjusted to provide the optimum therapeutic response. An effective amount is also one in which any toxic or detrimental effects (i.e., side effects) of an antibody or antigen binding fragment thereof are minimized or outweighed by the beneficial effects.

[0076] As used herein, the term "subject" includes any human or non-human animal.

[0077] As used herein, the term "epitope" refers to an antigenic determinant that interacts with a specific antigen binding site in the variable region of an antibody molecule known as a paratope. A single antigen may have more than one epitope. Thus, different antibodies may bind to different areas on an antigen and may have different biological effects. Epitopes may be either conformational or linear. A conformational epitope is produced by spatially juxtaposed amino acids from different segments of the linear polypeptide chain. A linear epitope is one produced by adjacent amino acid residues in a polypeptide chain.

[0078] It is noted here that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise.

II. Anti-Kallidin or des-Arg₁₀-Kallidin Antibodies

[0079] In one aspect the disclosure provides antibodies, or antigen binding fragments thereof, that specifically bind to Kallidin or des-Arg₁₀-Kallidin. Exemplary VH, VL and CDR amino acid sequences of the antibodies of the disclosure are set forth in Table 1.

Table 1. VH, VL and CDR amino acid sequences of exemplary anti-Kallidin or des-Arg₁₀-Kallidin antibodies.

Antibody Clone	Sequence	SEQ ID NO.
F151 HCDR3 consensus	X ₁ YX ₂ X ₃ DX ₄ HAMX ₅ Y	7
	where:	
	X ₁ is Y, F or H;	
	X ₂ is R, D, A, V, L, I, M, F, Y or W;	
	X ₃ is Y, F, W or H;	
	X ₄ is D, E or Y; and	
	X ₅ is D or E.	
F151 HCDR2 consensus	YFX ₁ PX ₂ NGNTGYNQKFRG	8
	where:	
	X ₁ is D, R, A, V, L, I, M, F, Y or W; and X ₂ is Y, D, E, N, or Q.	

Antibody Clone	Sequence	SEQ ID NO.
F151 HCDR1 consensus	GYSFTDYX ₁ IY	9
	Where X ₁ is N, W or Y.	
F151 LCDR3 consensus	QQX ₁ X ₂ SX ₃ PX ₄ T	10
	where:	
	X ₁ is Y, F or H;	
	X ₂ is Y, F, H or W;	
	X ₃ is Y, F, T or H; and	
	X ₄ is W, Y, F, H or L.	
F151 LCDR2 consensus	WASTRX ₁	11
	where X ₁ is E, D, Q or N.	
F151 LCDR1 consensus	KSSQSLLX ₁ SSNQKNX ₂ LA	12
	where:	
	X ₁ is W, H, Y or F; and	
	X ₂ is H or Y.	
F151 HCDR3	YYRYDDHAMDY	13
F151 HCDR2	YFDPYNGNTGYNQKFRG	14
F151 HCDR1	GYSFTDYNIY	15
F151 LCDR3	QQYYSPWT	16
F151 LCDR2	WASTRES	17
F151 LCDR1	KSSQSLLYSSNQKNYLA	18
F151 VH	EIQLVQSGPELVKPGTSVKVSCASGYSTFDYNIYWVKQSHCK SLEWIGYFDPYNGNTGYNQKFRGKATLTVDKSSSTAFMHLSSL TSDDSAVYYCANYRYDDHAMDYWGQGSTVIVSS	19
F151 Humanized HC1	EIQLVQSGPEVKKPGASVKVSCASGYSTFDYNIYWVKQSPGK SLEWIGYFDPYNGNTGYNQKFRGKATLTVDKSSSTAFMHLSSL TSEDSAVYYCANYRYDDHAMDYWGQGSTVIVSS	20
F151 Humanized HC2a	QIQLVQSGAEVKKPGASVKVSCASGYSTFDYNIYWVKQSPGK GLEWIGYFDPYNGNTGYNQKFRGKATLTVDKSSSTAFMHLSSL TSEESAVYYCANYRYDDHAMDYWGQGSTVIVSS	21
F151 Humanized HC2b	QIQLVQSGAEVKKPGASVKVSCASGYSTFDYNIYWVKQSPGK GLEWIGYFDPYNGNTGYNEKFRGKATLTVDKSSSTAFMHLSSL TSEESAVYYCANYRYDDHAMDYWGQGSTVIVSS	22

Antibody Clone	Sequence	SEQ ID NO.
F151 Humanized HC2c	QIQLVQSGAEVKKPGASVKVSCKASGYSFTDYNIIYWKQSPGK GLEWIGYFDYPYNGNTGYNQKFRGRKATLTVDKSSSTA VMHLGSK TSEESAVYYCANYRYDDHAMDYWGQGTSTVTVSS	23
F151 Humanized HC3a	QIQLVQSGAEVKKPGASVKVSCKASGYSFTDYNIIYWRQAPGQ GLEWIGYFDYPYNGNTGYNQKFRGRATLTVDKSTSTAYMELRSL RSDDTAVYYCANYRYDDHAMDYWGQGTSTVTVSS	24
F151 Humanized HC3b	QVQLVQSGAEVKKPGASVKVSCKASGYSFTDYNIIYWRQAPGQ GLEWMGYFDYPYNGNTGYNQKFRGRVTMTDTSTSTAYMELRSL RSDDTAVYYCANYRYDDHAMDYWGQGTSTVTVSS	25
F151 VL	DIVMSQSPSSSLAVSVGEKVTMSCKSSQSLLYSSNQKNYLAWYQ QKPGQSPKPLIYWASTRESGVPDRFTGSGSGTDFTLTISSVKA EDLAIYYCQYYSYPTWTFGGGTKLEIK	26
F151 Humanized LC1	DIVMSQSPSSSLAASVGDRTVTMSCKSSQSLLYSSNQKNYLAWYQ QKPGKSPKPLIYWASTRESGVPDRFTGSGSGTDFTLTISSVQA EDLAIYYCQYYSYPTWTFGGGTKLEIK	27
F151	DIVMTQSPSSSLASVGDRTVTISCKSSQSLLYSSNQKNYLAWYQ QKPGKSPKPLIYWASTRESGVPDRFSGSGSGTDFTLTISSVQA	28
Humanized LC2a	EDLAIYYCQYYSYPTWTFGGGTKLEIK	
F151 Humanized LC2b	DIVMTQSPSSSLASVGDRTVTISCKSSQSLLYSSNQKNYLAWYQ QKPGKSPKPLIYWASTRESGVPDRFSGSGSGTDFTLTISSVQA EDKAIYYCQYYSYPTWTFGGGTKLEIK	29
F151 Humanized LC3a	DIVMTQSPDSSLAVSLGERATINCKSSQSLLYSSNQKNYLAWYQ QKPGQPPKPLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQA EDVAVYYCQYYSYPTWTFGGGTKVEIK	30
F151 Humanized LC3b	DIVMTQSPDSSLAVSLGERATINCKSSQSLLYSSNQKNYLAWYQ QKPGQPPKLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQA EDVAVYYCQYYSYPTWTFGGGTKVEIK	31
B21 HCDR3	WEYDGYDLDY	32
B21 HCDR2	WIDPENGDTGYARKFQG	33
B21 HCDR1	GFNIKDYLLH	34
B21 LCDR3	LQGTHFPYT	35
B21 LCDR2	LVSKLDS	36
B21 LCDR1	KSSQSLLYSNGKTYLN	37

Antibody Clone	Sequence	SEQ ID NO.
B21 VH	EVQLQQSGAELVRSGASVKLSCTASGFNIKDYLLHWVKORPEQ GLEWIGWIDPENGDTGYARKFQGKATMTADTSSNTVYLHLSSL TSEDVAVYYFNAWEYDGYDLDYWGQGTSTVTVSS	38
B21 VL	DVMTQTPTLTLSVTIGQPASISCKSSQSLLYSNGKTYLNWLLQ RPGQSPKRLIYLVSKLDSGVPDRFTGSGSGTDFTLKIIRVEAE DLGVYYCLQGTHFPYTFGGGKLEIK	39
C63 HCDR3	EDYGGDY	40
C63 HCDR2	EIRSKSNNYATHYAESVKG	41
C63 HCDR1	GFTFSNYWMN	42
C63 LCDR3	QQYYSYPYT	43
C63 LCDR2	WASTRES	17
C63 LCDR1	KSSQSLLYSSDQRNYLA	44
C63 VH	EVKLEESGGGLVQPGGSMKLSCTVASGFTFSNYWMNWVRSPEK GLEWVAEIRSKSNNYATHYAESVKGRTISRDDSKSSVYLQMN NLRAEDTGIYYCIGEDYGGDYWGQGTSTVTVSS	45
C63 VL	DIVMSQSPSSLAVSVGEKVTMSCKSSQSLLYSSDQRNYLAWYQ QRSGQSPKLLIYWASTRESGVPDRFTGSGSGTDFTLTISSVKA EDLAVYYCQQYYSYPYTFGGGKLEIK	46
I22 HCDR3	FEYDGNYSPLDF	47
I22 HCDR2	WVDPENGDSYAPKFQ	48
I22 HCDR1	GFNIKDYMH	49
I22 LCDR3	QNDHSYPLT	50
I22 LCDR2	GASTRES	51
I22 LCDR1	KSSQSLLNSGNQKNYLA	52
I22 VH	EVQLQQSGAELVRSGASVKLSCTASGFNIKDYMHVVKORPEQ GLEWIGWVDPENGDSYAPKFQKATMTADTSSNTVYLQFSSL TSEDVAVYYCNAFEYDGNYSPLDFWGQGTSTVTVSS	53
I22 VL	DIVMTQSPSSLVSVSAGEKVTMSCKSSQSLNSGNQKNYLA QKPGQPPKLLIYGASTRESGVPDRFTGSGSGTDFTLTISSVQA EDLAVYYCQNDHSYPLTFGAGTKLEIK	54
I54 HCDR3	FEYDGNYSPLDF	55
I54 HCDR2	WVDPENGDSYAPKFQG	56
I54 HCDR1	GFNIKDYMH	57
I54 LCDR3	MQGTHFPYT	58
I54 LCDR2	LVSKLDS	59

Antibody Clone	Sequence	SEQ ID NO.
154 LCDR1	KSSQSLLYSNGETYLN	60
154 VH	EVQLQQSGAELVRSGLASVKLSCTASGFNIKDYIMHWVKORPEQ GLEWIGWVDPENGSDYAPKFQKATMTADTSSNTVYLQFSSSL TSEDYAVYYCNAFEYDGNYSPLDFWGQGTSTVTVSS	61
154 VL	DVVMITQTPLTSLSPICQPASISCKSSQSLLYSNGETYLNWLLQ RPGQSPKRLIYLVSKLDSGVDPDRFTGSRSGTDFTLKISRVESE DLGVYYCMQGTIHFYPTFGGGTKLEIK	62
B21/I22/I54 HCDR3 consensus	X_1 EYD GX_2 Y X_3 X_4 L DX_5	63
	where:	
	X_1 is W or F;	
	X_2 is N or no amino acid;	
	X_3 is Y or S;	
	X_4 is D or P; and	
	X_5 is F or Y.	
B21/I22/I54 HCDR2 consensus	W X_1 DPENG DGX_2 X_3 YAPKFQG	64
	where:	
	X_1 is I, or V;	
	X_2 is T, or S; and	
	X_3 is G, or D.	
B21/I22/I54 HCDR1 consensus	GFNIKDY YX_1 H	65
	where X_1 is L, or M.	
F151/C63/I22 LCDR3 consensus	Q X_1 X_2 X_3 S X_4 P X_5 T	66
	where:	
	X_1 is Q or N;	
	X_2 is Y, F, D or H;	
	X_3 is Y, F, H or W;	
	X_4 is Y, F, T or H; and	
	X_5 is W, Y, F, H or L.	
F151/C63/I22 LCDR2 consensus	X_2 ASTR X_2	67
	where:	
	X_1 is W or G; and	
	X_2 is E, D, Q or N	

Antibody Clone	Sequence	SEQ ID NO.
F151/C63/I22 LCDR1 consensus	KSSQSLLX ₁ X ₂ SX ₃ QX ₄ NX ₅ LA	68
	where:	
	X ₁ is W, H, Y or F;	
	X ₂ is S or G;	
	X ₃ is N or D;	
	X ₄ is K or R;	
	X ₅ is H or Y.	
B21/I54 LCDR3 consensus	X ₁ QGTTFPYT	69
	where:	
	X ₁ is L or M;	
B21/I54 LCDR2 both identical	LVSKLDS	36
B21/I54 LCDR1	KSSQSLLYSNGX ₁ TYLN	70
consensus	where:	
	X ₁ is K or E;	

[0080] In certain embodiments, the antibody, or antigen binding fragment thereof, comprises one or more CDR region amino acid sequence selected from the group consisting of SEQ ID NO: 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 32, 33, 34, 35, 36, 37, 40, 41, 42, 43, 44, 47, 48, 49, 50, 51, 52, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, and 70.

[0081] In other embodiments, the antibody, or antigen binding fragment thereof, comprises HCDR3, HCDR2 and HCDR1 region amino acid sequences selected from the group consisting of:

1. a) SEQ ID NO: 7, 8, and 9;
2. b) SEQ ID NO: 13, 14, and 15;
3. c) SEQ ID NO: 32, 33, and 34;
4. d) SEQ ID NO: 40, 41, and 42;
5. e) SEQ ID NO: 47, 48, and 49;
6. f) SEQ ID NO: 55, 56, and 57; and
7. g) SEQ ID NO: 63, 64, and 65, respectively.

[0082] In other embodiments, the antibody, or antigen binding fragment thereof, comprises the LCDR3, LCDR2 and LCDR1 region amino acid sequences selected from the group consisting of:

1. a) SEQ ID NO: 10, 11, and 12;
2. b) SEQ ID NO: 16, 17, and 18;

3. c) SEQ ID NO: 35, 36, and 37;
4. d) SEQ ID NO: 43, 17, and 44;
5. e) SEQ ID NO: 50, 51, and 52;
6. f) SEQ ID NO: 58, 59, and 60;
7. g) SEQ ID NO: 66, 67, and 68; and
8. h) SEQ ID NO: 69, 25, and 70, respectively.

[0083] In other embodiments, the antibody, or antigen binding fragment thereof, comprises the HCDR3, HCDR2, HCDR1, LCDR3, LCDR2 and LCDR1 region amino acid sequences selected from the group consisting of:

1. a) SEQ ID NO: 7, 8, 9, 10, 11, and 12;
2. b) SEQ ID NO: 13, 14, 15, 16, 17, and 18;
3. c) SEQ ID NO: 32, 33, 34, 35, 36 and 37;
4. d) SEQ ID NO: 40, 41, 42, 43, 17, and 44;
5. e) SEQ ID NO: 47, 48, 49, 50, 51, and 52; and
6. f) SEQ ID NO: 55, 56, 57, 58, 59, and 60, respectively

[0084] In other embodiment, the disclosure provides humanized antibodies, or antigen binding fragments thereof, comprising one or more CDR regions (or conservatively modified variants thereof) from the murine antibodies disclosed herein. Any method of humanization can be employed to generate the humanized antibodies of the disclosure. Suitable methods are disclosed herein and specifically exemplified in Example 4.

[0085] In a one particular embodiment, the humanized antibody, or antigen binding fragment thereof comprises:

a heavy chain variable region comprising the HCDR3, HCDR2 and HCDR1 region amino sequences set forth in 13, 14, and 15, respectively, and one or more amino acid substitution at positions selected from the group consisting of H1, H5, H9, H11, H12, H16, H38, H40, H41, H43, H44, H66, H75, H79, H81, H82A, H83, H87, and H108; and/or

a light chain variable region comprising the LCDR3, LCDR2 and LCDR1 region amino sequences set forth in 16, 17, and 18, respectively, and one or more amino acid substitution at positions selected from the group consisting of L5, L9, L15, L18, L19, L21, L22, L43, L63, L78, L79, L83, L85, L100 and L104 (according to the Kabat numbering convention).

[0086] In other embodiments, the antibody, or antigen binding fragment thereof, comprises the VH region amino acid sequences set forth in SEQ ID NO: 19, 20, 21, 22, 24, 25, 38, 45, 53, and/or 61.

[0087] In other embodiments, the antibody, or antigen binding fragment thereof, comprises the VL region amino acid sequences set forth in SEQ ID NO: 26, 27, 28, 29, 30, 31, 39, 46, 54, and/or

62.

[0088] In other embodiments, the antibody, or antigen binding fragment thereof, comprises the VH and VL region amino acid sequences selected from the group consisting of: SEQ ID NO: 19 and 26, SEQ ID NO: 20 and 27, SEQ ID NO: 21 and 28; SEQ ID NO: 22 and 28; SEQ ID NO: 23 and 29; SEQ ID NO: 24 and 30; SEQ ID NO: 25 and 31; SEQ ID NO: 38 and 39, SEQ ID NO: 45 and 46, SEQ ID NO: 53 and 54, or SEQ ID NO: 61 and 62, respectively.

[0089] In certain embodiments, the antibody, or antigen binding fragment thereof, comprises one or more CDR region amino acid sequence selected from the group consisting of SEQ ID NO: 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 32, 33, 34, 35, 36, 37, 40, 41, 42, 43, 44, 47, 48, 49, 50, 51, 52, 55, 56, 57, 58, 59 and 60, wherein the one or more CDR region amino acid sequences comprises at least one or more conservative amino acid substitutions.

[0090] The present disclosure also encompasses "conservative amino acid substitutions" in the CDR amino acid sequences (e.g., SEQ ID NOs: , 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 32, 33, 34, 35, 36, 37, 40, 41, 42, 43, 44, 47, 48, 49, 50, 51, 52, 55, 56, 57, 58, 59 and 60) of the antibodies of the disclosure, i.e., amino acid sequence modifications which do not abrogate the binding of the antibody to the antigen, e.g., Kallidin or des-Arg10-Kallidin. Conservative amino acid substitutions include the substitution of an amino acid in one class by an amino acid of the same class, where a class is defined by common physicochemical amino acid side chain properties and high substitution frequencies in homologous proteins found in nature, as determined, for example, by a standard Dayhoff frequency exchange matrix or BLOSUM matrix. Six general classes of amino acid side chains have been categorized and include: Class I (Cys); Class II (Ser, Thr, Pro, Ala, Gly); Class III (Asn, Asp, Gln, Glu); Class IV (His, Arg, Lys); Class V (Ile, Leu, Val, Met); and Class VI (Phe, Tyr, Trp). For example, substitution of an Asp for another class III residue such as Asn, Gln, or Glu, is a conservative substitution. Thus, a predicted nonessential amino acid residue in an anti-Kallidin or des-Arg10-Kallidin antibody is preferably replaced with another amino acid residue from the same class. Methods of identifying amino acid conservative substitutions which do not eliminate antigen binding are well-known in the art (see, e.g., Brummell et al., *Biochem.* 32:1180-1187 (1993); Kobayashi et al. *Protein Eng.* 12(10):879-884 (1999); and Burks et al. *Proc. Natl. Acad. Sci. USA* 94:412-417 (1997)).

[0091] In another embodiment, the present disclosure provides anti-Kallidin or des-Arg10-Kallidin antibodies, or antigen binding fragment thereof, that comprise a VH and/or VL region amino acid sequence with about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, identity to the VH region amino acid sequence set forth in SEQ ID NO: 19, 20, 21, 22, 24, 25, 38, 45, 53, or 61, or the VL region amino acid sequence set forth in SEQ ID NO: 26, 27, 28, 29, 30, 31, 39, 46, 54, or 62, respectively.

[0092] In another embodiment, the present disclosure provides anti-Kallidin or des-Arg10-Kallidin antibodies that bind to the same epitope and/or cross compete with an antibody, or antigen binding fragment thereof comprising the VH and VL region amino acid sequences set forth in SEQ ID NO: 19 and 25, SEQ ID NO: 38 and 39, SEQ ID NO: 45 and 46, SEQ ID NO: 53 and 54, or SEQ ID NO: 61 and 62, respectively. Such antibodies can be identified using routine competition binding assays including, for example, surface plasmon resonance (SPR)-based competition assays.

[0093] In certain embodiments, the antibodies of the disclosure bind a conformational epitope of kallidin (KD) or desArg10-Kallidin (DAKD) which adopts a "Pro4 kink" conformation. As depicted in Figure 17, a hallmark of the "Pro 4 kink" conformation is a type II tight turn in the main chain polypeptide backbone of KD or DAKD at Proline 4. As known to those of skill in the art, a type II tight turn conformation comprises three residues (X1-X2-X3) with the carbonyl of residue X1 forming a hydrogen bond with the amide N of residue X3, which is typically a glycine (see Richardson JS. "The anatomy and taxonomy of protein structure." *Adv Protein Chem.* 1981 ;34:167-339, which is incorporated by reference herein). Accordingly, in certain embodiments, a type II tight turn conformation is formed by the Pro3-Pro4-Gly5 motif of KD or DADK. In more specific embodiments, the "Pro 4 kink" conformation is further defined by all or substantially all of the remaining amino acids of KD (1-2 and 6-9) or DAKD adopting repeats of a sigmoid shape which align the hydrophobic side chains in a spatially stacking mode.

III. Modified Anti-Kallidin or des-Arg₁₀-Kallidin antibodies

[0094] In certain embodiments, anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure may comprise one or more modifications. Modified forms of anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure can be made using any techniques known in the art.

i) Reducing Immunogenicity

[0095] In certain embodiments, anti-Kallidin or des-Arg10-Kallidin antibodies, or antigen binding fragments thereof, of the disclosure are modified to reduce their immunogenicity using art-recognized techniques. For example, antibodies, or fragments thereof, can be chimericized, humanized, and/or deimmunized.

[0096] In one embodiment, an antibody, or antigen binding fragments thereof, of the disclosure may be chimeric. A chimeric antibody is an antibody in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies, or fragments thereof, are known in the art. See, e.g., Morrison, *Science* 229:1202 (1985); Oi et al., *BioTechniques* 4:214 (1986); Gillies et al., *J. Immunol. Methods* 125:191-202 (1989); U.S. Pat. Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entireties. Techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci.* 81:851-855 (1984); Neuberger et al., *Nature* 312:604-608 (1984); Takeda et al., *Nature* 314:452-454 (1985)) may be employed for the synthesis of said molecules. For example, a genetic sequence encoding a binding specificity of a mouse anti-Kallidin or des-Arg10-Kallidin antibody molecule may be fused together with a sequence from a human antibody molecule of appropriate biological activity. As used herein, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region, e.g., humanized antibodies.

[0097] In another embodiment, an antibody, or antigen binding fragment thereof, of the disclosure is humanized. Humanized antibodies, have a binding specificity comprising one or more complementarity determining regions (CDRs) from a non-human antibody and framework regions from a human antibody molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Pat. No. 5,585,089; Riechmann et al., *Nature* 332:323 (1988), which are incorporated herein by reference in their entireties.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Pat. Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, *Molecular Immunology* 28(4/5):489-498 (1991); Studnicka et al., *Protein Engineering* 7(6):805-814 (1994); Roguska. et al., *PNAS* 91:969-973 (1994)), and chain shuffling (U.S. Pat. No. 5,565,332).

[0098] In a particular embodiment, a humanization method is employed that is based on the impact of the molecular flexibility of the antibody during and at immune recognition (see WO2009/032661, which is incorporated herein by reference in its entirety). Protein flexibility is related to the molecular motion of the protein molecule. Protein flexibility is the ability of a whole protein, a part of a protein or a single amino acid residue to adopt an ensemble of conformations which differ significantly from each other. Information about protein flexibility can be obtained by performing protein X-ray crystallography experiments (see, for example, Kundu et al. 2002, *Biophys J* 83:723-732.), nuclear magnetic resonance experiments (see, for example, Freedberg et al., *J Am Chem Soc* 1998, 120(31):7916-7923) or by running molecular dynamics (MD) simulations. An MD simulation of a protein is done on a computer and allows one to determine the motion of all protein atoms over a period of time by calculating the physical interactions of the atoms with each other. The output of a MD simulation is the trajectory of the studied protein over the period of time of the simulation. The trajectory is an ensemble of protein conformations, also called snapshots, which are periodically sampled over the period of the simulation, e.g. every 1 picosecond (ps). It is by analyzing the ensemble of snapshots that one can quantify the flexibility of the protein amino acid residues. Thus, a flexible residue is one which adopts an ensemble of different conformations in the context of the polypeptide within which that residue resides. MD methods are known in the art, see, e.g., Brooks et al. "Proteins: A Theoretical Perspective of Dynamics, Structure and Thermodynamics" (Wiley, New York, 1988). Several software enable MD simulations, such as Amber (see Case et al. (2005) *J Comp Chem* 26:1668-1688), Charmm (see Brooks et al. (1983) *J Comp Chem* 4:187-217; and MacKerell et al. (1998) in "The Encyclopedia of Computational Chemistry" vol. 1:271-177, Schleyer et al., eds. Chichester: John Wiley & Sons) or Impact (see Rizzo et al. *J Am Chem Soc*; 2000; 122(51):12898-12900.)

[0099] Most protein complexes share a relatively large and planar buried surface and it has been shown that flexibility of binding partners provides the origin for their plasticity, enabling them to conformationally adapt to each other (*Structure* (2000) 8, R137-R142). As such, examples of "induced fit" have been shown to play a dominant role in protein-protein interfaces. In addition, there is a steadily increasing body of data showing that proteins actually bind ligands of diverse shapes sizes and composition (*Protein Science* (2002) 11:184-187) and that the conformational

diversity appears to be an essential component of the ability to recognize different partners (Science (2003) 299, 1362-1367). Flexible residues are involved in the binding of protein-protein partners (Structure (2006) 14, 683-693).

[0100] The flexible residues can adopt a variety of conformations that provide an ensemble of interaction areas that are likely to be recognized by memory B cells and to trigger an immunogenic response. Thus, an antibody can be humanized by modifying a number of residues from the framework so that the ensemble of conformations and of recognition areas displayed by the modified antibody resemble as much as possible those adopted by a human antibody. That can be achieved by modifying a limited number of residues by: (1) building a homology model of the parent mAb and running an MD simulation; (2) analyzing the flexible residues and identification of the most flexible residues of a non-human antibody molecule, as well as identifying residues or motifs likely to be a source of heterogeneity or of degradation reaction; (3) identifying a human antibody which displays the most similar ensemble of recognition areas as the parent antibody; (4) determining the flexible residues to be mutated, residues or motifs likely to be a source of heterogeneity and degradation are also mutated; and (5) checking for the presence of known T cell or B cell epitopes. The flexible residues can be found using an MD calculation as taught herein using an implicit solvent model, which accounts for the interaction of the water solvent with the protein atoms over the period of time of the simulation.

[0101] Once the set of flexible residues has been identified within the variable light and heavy chains, a set of human heavy and light chain variable region frameworks that closely resemble that of the antibody of interest are identified. That can be done, for example, using a BLAST search on the set of flexible residues against a database of antibody human germ line sequence. It can also be done by comparing the dynamics of the parent mAb with the dynamics of a library of germ line canonical structures. The CDR residues and neighboring residues are excluded from the search to ensure high affinity for the antigen is preserved. Flexible residues then are replaced.

[0102] When several human residues show similar homologies, the selection is driven also by the nature of the residues that are likely to affect the solution behavior of the humanized antibody. For instance, polar residues will be preferred in exposed flexible loops over hydrophobic residues. Residues which are a potential source of instability and heterogeneity are also mutated even if there are found in the CDRs. That will include exposed methionines as sulfoxide formation can result from oxygen radicals, proteolytic cleavage of acid labile bonds such as those of the Asp-Pro dipeptide (Drug Dev Res (2004) 61:137-154), deamidation sites found with an exposed asparagine residue followed by a small amino acid, such as Gly, Ser, Ala, His, Asn or Cys (J Chromatog (2006) 837:35-43) and N-glycosylation sites, such as the Asn-X-Ser/Thr site. Typically, exposed methionines will be substituted by a Leu, exposed asparagines will be replaced by a glutamine or by an aspartate, or the subsequent residue will be changed. For the glycosylation site (Asn-X-Ser/Thr), either the Asn or the Ser/Thr residue will be changed.

[0103] The resulting composite antibody sequence is checked for the presence of known B cell or linear T-cell epitopes. A search is performed, for example, with the publicly available Immune Epitope Data Base (IEDB) (PLoS Biol (2005) 3(3)e91). If a known epitope is found within the composite sequence, another set of human sequences is retrieved and substituted. Thus, unlike the resurfacing method of U.S. Pat. No. 5,639,641, both B-cell-mediated and T-cell-mediated

immunogenic responses are addressed by the method. The method also avoids the issue of loss of activity that is sometimes observed with CDR grafting (U.S. Pat. No. 5,530,101). In addition, stability and solubility issues also are considered in the engineering and selection process, resulting in an antibody that is optimized for low immunogenicity, high antigen affinity and improved biophysical properties.

[0104] In some embodiments, de-immunization can be used to decrease the immunogenicity of an antibody, or antigen binding fragment thereof. As used herein, the term "de-immunization" includes alteration of an antibody, or antigen binding fragment thereof, to modify T cell epitopes (see, e.g., WO9852976A1, WO0034317A2). For example, VH and VL sequences from the starting antibody may be analyzed and a human T cell epitope "map" may be generated from each V region showing the location of epitopes in relation to complementarity-determining regions (CDRs) and other key residues within the sequence. Individual T cell epitopes from the T cell epitope map are analyzed in order to identify alternative amino acid substitutions with a low risk of altering activity of the final antibody. A range of alternative VH and VL sequences are designed comprising combinations of amino acid substitutions and these sequences are subsequently incorporated into a range of Kallidin or des-Arg10-Kallidin-specific antibodies or fragments thereof for use in the diagnostic and treatment methods disclosed herein, which are then tested for function. Typically, between 12 and 24 variant antibodies are generated and tested. Complete heavy and light chain genes comprising modified V and human C regions are then cloned into expression vectors and the subsequent plasmids introduced into cell lines for the production of whole antibody. The antibodies are then compared in appropriate biochemical and biological assays, and the optimal variant is identified.

ii) Effector Functions and Fc Modifications

[0105] Anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure may comprise an antibody constant region (e.g. an IgG constant region e.g., a human IgG constant region, e.g., a human IgG1 or IgG4 constant region) which mediates one or more effector functions. For example, binding of the C1 component of complement to an antibody constant region may activate the complement system. Activation of complement is important in the opsonisation and lysis of cell pathogens. The activation of complement also stimulates the inflammatory response and may also be involved in autoimmune hypersensitivity. Further, antibodies bind to receptors on various cells via the Fc region, with a Fc receptor binding site on the antibody Fc region binding to a Fc receptor (FcR) on a cell. There are a number of Fc receptors which are specific for different classes of antibody, including IgG (gamma receptors), IgE (epsilon receptors), IgA (alpha receptors) and IgM (mu receptors). Binding of antibody to Fc receptors on cell surfaces triggers a number of important and diverse biological responses including engulfment and destruction of antibody-coated particles, clearance of immune complexes, lysis of antibody-coated target cells by killer cells (called antibody-dependent cell-mediated cytotoxicity, or ADCC), release of inflammatory mediators, placental transfer and control of immunoglobulin production. In preferred embodiments, the antibodies, or fragments thereof, of the disclosure bind to an Fc-gamma receptor. In alternative embodiments, anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure may comprise a constant region which is devoid of one or more effector functions (e.g., ADCC activity) and/or is unable to bind Fc receptor.

[0106] Certain embodiments of the disclosure include anti-Kallidin or des-Arg10-Kallidin antibodies in which at least one amino acid in one or more of the constant region domains has been deleted or otherwise altered so as to provide desired biochemical characteristics such as reduced or enhanced effector functions, the ability to non-covalently dimerize, increased ability to localize at the site of a tumor, reduced serum half-life, or increased serum half-life when compared with a whole, unaltered antibody of approximately the same immunogenicity. For example, certain antibodies, or fragments thereof, for use in the diagnostic and treatment methods described herein are domain deleted antibodies which comprise a polypeptide chain similar to an immunoglobulin heavy chain, but which lack at least a portion of one or more heavy chain domains. For instance, in certain antibodies, one entire domain of the constant region of the modified antibody will be deleted, for example, all or part of the CH2 domain will be deleted.

[0107] In certain other embodiments, anti-Kallidin or des-Arg10-Kallidin antibodies comprise constant regions derived from different antibody isotypes (e.g., constant regions from two or more of a human IgG1, IgG2, IgG3, or IgG4). In other embodiments, anti-Kallidin or des-Arg10-Kallidin antibodies comprises a chimeric hinge (i.e., a hinge comprising hinge portions derived from hinge domains of different antibody isotypes, e.g., an upper hinge domain from an IgG4 molecule and an IgG1 middle hinge domain). In one embodiment, an anti-Kallidin or des-Arg10-Kallidin antibodies comprises an Fc region or portion thereof from a human IgG4 molecule and a Ser228Pro mutation (EU numbering) in the core hinge region of the molecule.

[0108] In certain anti-Kallidin or des-Arg10-Kallidin antibodies, the Fc portion may be mutated to increase or decrease effector function using techniques known in the art. For example, the deletion or inactivation (through point mutations or other means) of a constant region domain may reduce Fc receptor binding of the circulating modified antibody thereby increasing tumor localization. In other cases it may be that constant region modifications consistent with the instant disclosure moderate complement binding and thus reduce the serum half life and nonspecific association of a conjugated cytotoxin. Yet other modifications of the constant region may be used to modify disulfide linkages or oligosaccharide moieties that allow for enhanced localization due to increased antigen specificity or flexibility. The resulting physiological profile, bioavailability and other biochemical effects of the modifications, such as tumor localization, biodistribution and serum half-life, may easily be measured and quantified using well know immunological techniques without undue experimentation.

[0109] In certain embodiments, an Fc domain employed in an antibody of the disclosure is an Fc variant. As used herein, the term "Fc variant" refers to an Fc domain having at least one amino acid substitution relative to the wild-type Fc domain from which said Fc domain is derived. For example, wherein the Fc domain is derived from a human IgG1 antibody, the Fc variant of said human IgG1 Fc domain comprises at least one amino acid substitution relative to said Fc domain.

[0110] The amino acid substitution(s) of an Fc variant may be located at any position (i.e., any EU convention amino acid position) within the Fc domain. In one embodiment, the Fc variant comprises a substitution at an amino acid position located in a hinge domain or portion thereof. In another embodiment, the Fc variant comprises a substitution at an amino acid position located in a CH2 domain or portion thereof. In another embodiment, the Fc variant comprises a substitution at

an amino acid position located in a CH3 domain or portion thereof. In another embodiment, the Fc variant comprises a substitution at an amino acid position located in a CH4 domain or portion thereof.

[0111] The antibodies of the disclosure may employ any art-recognized Fc variant which is known to impart an improvement (e.g., reduction or enhancement) in effector function and/or FcR binding. Said Fc variants may include, for example, any one of the amino acid substitutions disclosed in International PCT Publications WO88/07089A1, WO96/14339A1, WO98/05787A1, WO98/23289A1, WO99/51642A1, WO99/58572A1, WO00/09560A2, WO00/32767A1, WO00/42072A2, WO02/44215A2, WO02/060919A2, WO03/074569A2, WO04/016750A2, WO04/029207A2, WO04/035752A2, WO04/063351A2, WO04/074455A2, WO04/099249A2, WO05/040217A2, WO05/070963A1, WO05/077981A2, WO05/092925A2, WO05/123780A2, WO06/019447A1, WO06/047350A2, and WO06/085967A2 or U.S. Pat. Nos. 5,648,260; 5,739,277; 5,834,250; 5,869,046; 6,096,871; 6,121,022; 6,194,551; 6,242,195; 6,277,375; 6,528,624; 6,538,124; 6,737,056; 6,821,505; 6,998,253; and 7,083,784, each of which is incorporated by reference herein. In one exemplary embodiment, an antibody of the disclosure may comprise an Fc variant comprising an amino acid substitution at EU position 268 (e.g., H268D or H268E). In another exemplary embodiment, an antibody of the disclosure may comprise an amino acid substitution at EU position 239 (e.g., S239D or S239E) and/or EU position 332 (e.g., I332D or I332Q).

[0112] In certain embodiments, an antibody of the disclosure may comprise an Fc variant comprising an amino acid substitution which alters the antigen-independent effector functions of the antibody, in particular the circulating half-life of the antibody. Such antibodies exhibit either increased or decreased binding to FcRn when compared to antibodies lacking these substitutions, therefore, have an increased or decreased half-life in serum, respectively. Fc variants with improved affinity for FcRn are anticipated to have longer serum half-lives, and such molecules have useful applications in methods of treating mammals where long half-life of the administered antibody is desired, e.g., to treat a chronic disease or disorder. In contrast, Fc variants with decreased FcRn binding affinity are expected to have shorter half-lives, and such molecules are also useful, for example, for administration to a mammal where a shortened circulation time may be advantageous, e.g. for in vivo diagnostic imaging or in situations where the starting antibody has toxic side effects when present in the circulation for prolonged periods. Fc variants with decreased FcRn binding affinity are also less likely to cross the placenta and, thus, are also useful in the treatment of diseases or disorders in pregnant women. In addition, other applications in which reduced FcRn binding affinity may be desired include those applications in which localization the brain, kidney, and/or liver is desired. In one exemplary embodiment, the altered antibodies of the disclosure exhibit reduced transport across the epithelium of kidney glomeruli from the vasculature. In another embodiment, the altered antibodies of the disclosure exhibit reduced transport across the blood brain barrier (BBB) from the brain, into the vascular space. In one embodiment, an antibody with altered FcRn binding comprises an Fc domain having one or more amino acid substitutions within the "FcRn binding loop" of an Fc domain. The FcRn binding loop is comprised of amino acid residues 280-299 (according to EU numbering). Exemplary amino acid substitutions which altered FcRn binding activity are disclosed in International PCT Publication No. WO05/047327 which is incorporated by reference herein. In certain exemplary embodiments, the antibodies, or fragments thereof, of the disclosure comprise an Fc domain having one or more of the following substitutions: V284E, H285E, N286D, K290E and S304D (EU numbering).

[0113] In other embodiments, antibodies, for use in the diagnostic and treatment methods described herein have a constant region, e.g., an IgG1 or IgG4 heavy chain constant region, which is altered to reduce or eliminate glycosylation. For example, an antibody of the disclosure may also comprise an Fc variant comprising an amino acid substitution which alters the glycosylation of the antibody. For example, said Fc variant may have reduced glycosylation (e.g., N- or O-linked glycosylation). In exemplary embodiments, the Fc variant comprises reduced glycosylation of the N-linked glycan normally found at amino acid position 297 (EU numbering). In another embodiment, the antibody has an amino acid substitution near or within a glycosylation motif, for example, an N-linked glycosylation motif that contains the amino acid sequence NXT or NXS. In a particular embodiment, the antibody comprises an Fc variant with an amino acid substitution at amino acid position 228 or 299 (EU numbering). In more particular embodiments, the antibody comprises an IgG1 or IgG4 constant region comprising an S228P and a T299A mutation (EU numbering).

[0114] Exemplary amino acid substitutions which confer reduce or altered glycosylation are disclosed in International PCT Publication No. WO05/018572, which is incorporated by reference herein. In preferred embodiments, the antibodies, or fragments thereof, of the disclosure are modified to eliminate glycosylation. Such antibodies, or fragments thereof, may be referred to as "agly" antibodies, or fragments thereof, (e.g. "agly" antibodies). While not being bound by theory, it is believed that "agly" antibodies, or fragments thereof, may have an improved safety and stability profile in vivo. Exemplary agly antibodies, or fragments thereof, comprise an aglycosylated Fc region of an IgG4 antibody which is devoid of Fc-effector function thereby eliminating the potential for Fc mediated toxicity to the normal vital organs that express Kallidin or des-Arg10-Kallidin. In yet other embodiments, antibodies, or fragments thereof, of the disclosure comprise an altered glycan. For example, the antibody may have a reduced number of fucose residues on an N-glycan at Asn297 of the Fc region, i.e., is afucosylated. In another embodiment, the antibody may have an altered number of sialic acid residues on the N-glycan at Asn297 of the Fc region.

iii) Covalent Attachment

[0115] Anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure may be modified, e.g., by the covalent attachment of a molecule to the antibody such that covalent attachment does not prevent the antibody from specifically binding to its cognate epitope. For example, but not by way of limitation, the antibodies, or fragments thereof, of the disclosure may be modified by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, etc. Additionally, the derivative may contain one or more non-classical amino acids.

[0116] Antibodies, or fragments thereof, of the disclosure may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalent and non-covalent conjugations) to polypeptides or other compositions. For example, anti-Kallidin or des-Arg10-Kallidin antibodies may be recombinantly fused or conjugated to molecules useful as

labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO 92/08495; WO 91/14438; WO 89/12624; U.S. Pat. No. 5,314,995; and EP 396,387.

[0117] Anti-Kallidin or des-Arg10-Kallidin antibodies may be fused to heterologous polypeptides to increase the in vivo half life or for use in immunoassays using methods known in the art. For example, in one embodiment, PEG can be conjugated to the anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure to increase their half-life in vivo. Leong, S. R., *et al.*, Cytokine 16:106 (2001); Adv. in Drug Deliv. Rev. 54:531 (2002); or Weir *et al.*, Biochem. Soc. Transactions 30:512 (2002).

[0118] Moreover, anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure can be fused to marker sequences, such as a peptide to facilitate their purification or detection. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, Calif., 91311), among others, many of which are commercially available. As described in Gentz *et al.*, Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, Cell 37:767 (1984)) and the "flag" tag.

[0119] Anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure may be used in nonconjugated form or may be conjugated to at least one of a variety of molecules, e.g., to improve the therapeutic properties of the molecule, to facilitate target detection, or for imaging or therapy of the patient. Anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure can be labeled or conjugated either before or after purification, when purification is performed. In particular, anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure may be conjugated to therapeutic agents, prodrugs, peptides, proteins, enzymes, viruses, lipids, biological response modifiers, pharmaceutical agents, or PEG.

[0120] The present disclosure further encompasses anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure conjugated to a diagnostic or therapeutic agent. The anti-Kallidin or des-Arg10-Kallidin antibodies can be used diagnostically to, for example, monitor the development or progression of an immune cell disorder (e.g., CLL) as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment and/or prevention regimen. Detection can be facilitated by coupling the anti-Kallidin or des-Arg10-Kallidin antibodies to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. See, for example, U.S. Pat. No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present disclosure. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include

luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{111}In or ^{99}Tc .

[0121] Anti-Kallidin or des-Arg10-Kallidin antibodies for use in the diagnostic and treatment methods disclosed herein may be conjugated to cytotoxins (such as radioisotopes, cytotoxic drugs, or toxins) therapeutic agents, cytostatic agents, biological toxins, prodrugs, peptides, proteins, enzymes, viruses, lipids, biological response modifiers, pharmaceutical agents, immunologically active ligands (e.g., lymphokines or other antibodies wherein the resulting molecule binds to both the neoplastic cell and an effector cell such as a T cell), or PEG.

[0122] In another embodiment, an anti-Kallidin or des-Arg10-Kallidin antibody for use in the diagnostic and treatment methods disclosed herein can be conjugated to a molecule that decreases tumor cell growth. In other embodiments, the disclosed compositions may comprise antibodies, or fragments thereof, coupled to drugs or prodrugs. Still other embodiments of the present disclosure comprise the use of antibodies, or fragments thereof, conjugated to specific biotoxins or their cytotoxic fragments such as ricin, gelonin, *Pseudomonas* exotoxin or diphtheria toxin. The selection of which conjugated or unconjugated antibody to use will depend on the type and stage of cancer, use of adjunct treatment (e.g., chemotherapy or external radiation) and patient condition. It will be appreciated that one skilled in the art could readily make such a selection in view of the teachings herein.

[0123] It will be appreciated that, in previous studies, anti-tumor antibodies labeled with isotopes have been used successfully to destroy tumor cells in animal models, and in some cases in humans. Exemplary radioisotopes include: ^{90}Y , ^{125}I , ^{131}I , ^{123}I , ^{111}In , ^{105}Rh , ^{153}Sm , ^{67}Cu , ^{67}Ga , ^{166}Ho , ^{177}Lu , ^{186}Re and ^{188}Re . The radionuclides act by producing ionizing radiation which causes multiple strand breaks in nuclear DNA, leading to cell death. The isotopes used to produce therapeutic conjugates typically produce high energy alpha- or beta-particles which have a short path length. Such radionuclides kill cells to which they are in close proximity, for example neoplastic cells to which the conjugate has attached or has entered. They have little or no effect on non-localized cells. Radionuclides are essentially non-immunogenic.

IV. Expression of Anti-Kallidin or des-Arg10-Kallidin Antibodies, or Antigen Binding Fragments Thereof

[0124] Following manipulation of the isolated genetic material to provide anti-Kallidin or des-Arg10-Kallidin antibodies of the disclosure as set forth above, the genes are typically inserted in an expression vector for introduction into host cells that may be used to produce the desired quantity of the claimed antibodies, or fragments thereof.

[0125] The term "vector" or "expression vector" is used herein for the purposes of the specification and claims, to mean vectors used in accordance with the present disclosure as a vehicle for introducing into and expressing a desired gene in a cell. As known to those skilled in the art, such vectors may easily be selected from the group consisting of plasmids, phages, viruses and retroviruses. In general, vectors compatible with the instant disclosure will comprise a selection marker, appropriate restriction sites to facilitate cloning of the desired gene and the ability to enter

and/or replicate in eukaryotic or prokaryotic cells.

[0126] Numerous expression vector systems may be employed for the purposes of this disclosure. For example, one class of vector utilizes DNA elements which are derived from animal viruses such as bovine papilloma virus, polyoma virus, adenovirus, vaccinia virus, baculovirus, retroviruses (RSV, MMTV or MOMLV) or SV40 virus. Others involve the use of polycistronic systems with internal ribosome binding sites. Additionally, cells which have integrated the DNA into their chromosomes may be selected by introducing one or more markers which allow selection of transfected host cells. The marker may provide for prototrophy to an auxotrophic host, biocide resistance (e.g., antibiotics) or resistance to heavy metals such as copper. The selectable marker gene can either be directly linked to the DNA sequences to be expressed, or introduced into the same cell by cotransformation. Additional elements may also be needed for optimal synthesis of mRNA. These elements may include signal sequences, splice signals, as well as transcriptional promoters, enhancers, and termination signals. In particularly preferred embodiments the cloned variable region genes are inserted into an expression vector along with the heavy and light chain constant region genes (preferably human) synthetic as discussed above.

[0127] In other preferred embodiments the anti-Kallidin or des-Arg1 0-Kallidin antibodies, or fragments thereof, of the disclosure may be expressed using polycistronic constructs. In such expression systems, multiple gene products of interest such as heavy and light chains of antibodies may be produced from a single polycistronic construct. These systems advantageously use an internal ribosome entry site (IRES) to provide relatively high levels of polypeptides of the disclosure in eukaryotic host cells. Compatible IRES sequences are disclosed in U.S. Pat. No. 6,193,980, which is incorporated by reference herein. Those skilled in the art will appreciate that such expression systems may be used to effectively produce the full range of polypeptides disclosed in the instant application.

[0128] More generally, once a vector or DNA sequence encoding an antibody, or fragment thereof, has been prepared, the expression vector may be introduced into an appropriate host cell. That is, the host cells may be transformed. Introduction of the plasmid into the host cell can be accomplished by various techniques well known to those of skill in the art. These include, but are not limited to, transfection (including electrophoresis and electroporation), protoplast fusion, calcium phosphate precipitation, cell fusion with enveloped DNA, microinjection, and infection with intact virus. See, Ridgway, A. A. G. "Mammalian Expression Vectors" Chapter 24.2, pp. 470-472 Vectors, Rodriguez and Denhardt, Eds. (Butterworths, Boston, Mass. 1988). Most preferably, plasmid introduction into the host is via electroporation. The transformed cells are grown under conditions appropriate to the production of the light chains and heavy chains, and assayed for heavy and/or light chain protein synthesis. Exemplary assay techniques include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), or fluorescence-activated cell sorter analysis (FACS), immunohistochemistry and the like.

[0129] As used herein, the term "transformation" shall be used in a broad sense to refer to the introduction of DNA into a recipient host cell that changes the genotype and consequently results in a change in the recipient cell.

[0130] Along those same lines, "host cells" refers to cells that have been transformed with vectors

constructed using recombinant DNA techniques and encoding at least one heterologous gene. In descriptions of processes for isolation of polypeptides from recombinant hosts, the terms "cell" and "cell culture" are used interchangeably to denote the source of antibody unless it is clearly specified otherwise. In other words, recovery of polypeptide from the "cells" may mean either from spun down whole cells, or from the cell culture containing both the medium and the suspended cells.

[0131] In one embodiment, the host cell line used for antibody expression is of mammalian origin; those skilled in the art can determine particular host cell lines which are best suited for the desired gene product to be expressed therein. Exemplary host cell lines include, but are not limited to, DG44 and DUXB11 (Chinese Hamster Ovary lines, DHFR minus), HELA (human cervical carcinoma), CVI (monkey kidney line), COS (a derivative of CVI with SV40 T antigen), R1610 (Chinese hamster fibroblast) BALBC/3T3 (mouse fibroblast), HAK (hamster kidney line), SP2/O (mouse myeloma), BFA-1c1BPT (bovine endothelial cells), RAJI (human lymphocyte), 293 (human kidney). In one embodiment, the cell line provides for altered glycosylation, e.g., afucosylation, of the antibody expressed therefrom (e.g., PER.C6.RTM. (Crucell) or FUT8-knockout CHO cell lines (Potelligent.RTM. Cells) (Biowa, Princeton, N.J.)). In one embodiment NS0 cells may be used. CHO cells are particularly preferred. Host cell lines are typically available from commercial services, the American Tissue Culture Collection or from published literature.

[0132] *In vitro* production allows scale-up to give large amounts of the desired polypeptides. Techniques for mammalian cell cultivation under tissue culture conditions are known in the art and include homogeneous suspension culture, e.g. in an airlift reactor or in a continuous stirrer reactor, or immobilized or entrapped cell culture, e.g. in hollow fibers, microcapsules, on agarose microbeads or ceramic cartridges. If necessary and/or desired, the solutions of polypeptides can be purified by the customary chromatography methods, for example gel filtration, ion-exchange chromatography, chromatography over DEAE-cellulose and/or (immuno-)affinity chromatography.

[0133] Genes encoding the anti-Kallidin or des-Arg10-Kallidin antibodies, or fragments thereof, of the disclosure can also be expressed non-mammalian cells such as bacteria or yeast or plant cells. In this regard it will be appreciated that various unicellular non-mammalian microorganisms such as bacteria can also be transformed; i.e. those capable of being grown in cultures or fermentation. Bacteria, which are susceptible to transformation, include members of the enterobacteriaceae, such as strains of *Escherichia coli* or *Salmonella*; Bacillaceae, such as *Bacillus subtilis*; *Pneumococcus*; *Streptococcus*, and *Haemophilus influenzae*. It will further be appreciated that, when expressed in bacteria, the polypeptides can become part of inclusion bodies. The polypeptides must be isolated, purified and then assembled into functional molecules.

[0134] In addition to prokaryotes, eukaryotic microbes may also be used. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among eukaryotic microorganisms although a number of other strains are commonly available. For expression in *Saccharomyces*, the plasmid YRp7, for example, (Stinchcomb et al., *Nature*, 282:39 (1979); Kingsman et al., *Gene*, 7:141 (1979); Tschemper et al., *Gene*, 10:157 (1980)) is commonly used. This plasmid already contains the TRP1 gene which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example ATCC No. 44076 or PEP4-1 (Jones, *Genetics*, 85:12 (1977)). The presence of the *trp1* lesion as a characteristic of the yeast host cell genome then provides an effective environment for detecting transformation by growth in the

absence of tryptophan.

V. Pharmaceutical Formulations and Methods of Administration of Anti-Kallidin or des-Arg10-Kallidin Antibodies.

[0135] In another aspect, the disclosure provides pharmaceutical compositions comprising an anti-Kallidin or des-Arg10-Kallidin antibody, or fragment thereof.

[0136] Methods of preparing and administering antibodies, or fragments thereof, of the disclosure to a subject are well known to or are readily determined by those skilled in the art. The route of administration of the antibodies, or fragments thereof, of the disclosure may be oral, parenteral, by inhalation or topical. The term parenteral as used herein includes intravenous, intraarterial, intraperitoneal, intramuscular, subcutaneous, rectal or vaginal administration. The intravenous, intraarterial, subcutaneous and intramuscular forms of parenteral administration are generally preferred. While all these forms of administration are clearly contemplated as being within the scope of the disclosure, a form for administration would be a solution for injection, in particular for intravenous or intraarterial injection or drip. Usually, a suitable pharmaceutical composition for injection may comprise a buffer (e.g. acetate, phosphate or citrate buffer), a surfactant (e.g. polysorbate), optionally a stabilizer agent (e.g. human albumin), etc. However, in other methods compatible with the teachings herein, the polypeptides can be delivered directly to the site of the adverse cellular population thereby increasing the exposure of the diseased tissue to the therapeutic agent.

[0137] Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. In the subject disclosure, pharmaceutically acceptable carriers include, but are not limited to, 0.01-0.1M and preferably 0.05M phosphate buffer or 0.8% saline. Other common parenteral vehicles include sodium phosphate solutions, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers, such as those based on Ringer's dextrose, and the like. Preservatives and other additives may also be present such as for example, antimicrobials, antioxidants, chelating agents, and inert gases and the like. More particularly, pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In such cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage and will preferably be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic

acid, thimerosal and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols, such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

[0138] In any case, sterile injectable solutions can be prepared by incorporating an active compound (e.g., an antibody by itself or in combination with other active agents) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated herein, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle, which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying, which yields a powder of an active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The preparations for injections are processed, filled into containers such as ampoules, bags, bottles, syringes or vials, and sealed under aseptic conditions according to methods known in the art. Further, the preparations may be packaged and sold in the form of a kit such as those described in U.S. Publication No. 2002/0102208 which is incorporated herein by reference. Such articles of manufacture will preferably have labels or package inserts indicating that the associated compositions are useful for treating a subject suffering from, or predisposed to autoimmune or neoplastic disorders.

[0139] Effective doses of the stabilized antibodies, or fragments thereof, of the present disclosure, for the treatment of the above described conditions vary depending upon many different factors, including means of administration, target site, physiological state of the patient, whether the patient is human or an animal, other medications administered, and whether treatment is prophylactic or therapeutic. Usually, the patient is a human, but non-human mammals including transgenic mammals can also be treated. Treatment dosages may be titrated using routine methods known to those of skill in the art to optimize safety and efficacy.

[0140] For passive immunization with an antibody of the disclosure, the dosage may range, e.g., from about 0.0001 to 100 mg/kg, and more usually 0.01 to 5 mg/kg (e.g., 0.02 mg/kg, 0.25 mg/kg, 0.5 mg/kg, 0.75 mg/kg, 1 mg/kg, 2 mg/kg, etc.), of the host body weight. For example dosages can be 1 mg/kg body weight or 10 mg/kg body weight or within the range of 1-10 mg/kg, preferably at least 1 mg/kg. Doses intermediate in the above ranges are also intended to be within the scope of the disclosure.

[0141] Subjects can be administered such doses daily, on alternative days, weekly or according to any other schedule determined by empirical analysis. An exemplary treatment entails administration in multiple dosages over a prolonged period, for example, of at least six months. Additional exemplary treatment regimes entail administration once per every two weeks or once a month or once every 3 to 6 months. Exemplary dosage schedules include 1-10 mg/kg or 15 mg/kg on consecutive days, 30 mg/kg on alternate days or 60 mg/kg weekly. In some methods, two or more monoclonal antibodies with different binding specificities are administered simultaneously, in which case the dosage of each antibody administered may fall within the ranges indicated.

[0142] Antibodies, or fragments thereof, of the disclosure can be administered on multiple

occasions. Intervals between single dosages can be, e.g., daily, weekly, monthly or yearly. Intervals can also be irregular as indicated by measuring blood levels of polypeptide or target molecule in the patient. In some methods, dosage is adjusted to achieve a certain plasma antibody or toxin concentration, e.g., 1-1000 ug/ml or 25-300 ug/ml. Alternatively, antibodies, or fragments thereof, can be administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, humanized antibodies show the longest half-life, followed by chimeric antibodies and nonhuman antibodies. In one embodiment, the antibodies, or fragments thereof, of the disclosure can be administered in unconjugated form. In another embodiment, the antibodies of the disclosure can be administered multiple times in conjugated form. In still another embodiment, the antibodies, or fragments thereof, of the disclosure can be administered in unconjugated form, then in conjugated form, or vice versa.

[0143] The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, compositions containing the present antibodies or a cocktail thereof are administered to a patient not already in the disease state to enhance the patient's resistance. Such an amount is defined to be a "prophylactic effective dose." In this use, the precise amounts again depend upon the patient's state of health and general immunity, but generally range from 0.1 to 25 mg per dose, especially 0.5 to 2.5 mg per dose. A relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives.

[0144] In therapeutic applications, a relatively high dosage (e.g., from about 1 to 400 mg/kg of antibody per dose, with dosages of from 5 to 25 mg being more commonly used for radioimmunoconjugates and higher doses for cytotoxin-drug conjugated molecules) at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

[0145] In one embodiment, a subject can be treated with a nucleic acid molecule encoding a polypeptide of the disclosure (e.g., in a vector). Doses for nucleic acids encoding polypeptides range from about 10 ng to 1 g, 100 ng to 100 mg, 1 ug to 10 mg, or 30-300 ug DNA per patient. Doses for infectious viral vectors vary from 10-100, or more, virions per dose.

[0146] Therapeutic agents can be administered by parenteral, topical, intravenous, oral, subcutaneous, intraarterial, intracranial, intraperitoneal, intranasal or intramuscular means for prophylactic and/or therapeutic treatment. Intramuscular injection or intravenous infusion are preferred for administration of a antibody of the disclosure. In some methods, therapeutic antibodies, or fragments thereof, are injected directly into the cranium. In some methods, antibodies, or fragments thereof, are administered as a sustained release composition or device, such as a Medipad™ device.

[0147] Agents of the disclosure can optionally be administered in combination with other agents that are effective in treating the disorder or condition in need of treatment (e.g., prophylactic or therapeutic). Preferred additional agents are those which are art recognized and are standardly administered for a particular disorder.

[0148] Effective single treatment dosages (i.e., therapeutically effective amounts) of ^{90}Y -labeled antibodies of the disclosure range from between about 5 and about 75 mCi, more preferably between about 10 and about 40 mCi. Effective single treatment non-marrow ablative dosages of ^{131}I -labeled antibodies range from between about 5 and about 70 mCi, more preferably between about 5 and about 40 mCi. Effective single treatment ablative dosages (i.e., may require autologous bone marrow transplantation) of ^{131}I -labeled antibodies range from between about 30 and about 600 mCi, more preferably between about 50 and less than about 500 mCi. In conjunction with a chimeric modified antibody, owing to the longer circulating half life vis-a-vis murine antibodies, an effective single treatment non-marrow ablative dosages of iodine-131 labeled chimeric antibodies range from between about 5 and about 40 mCi, more preferably less than about 30 mCi. Imaging criteria for, e.g., the ^{111}In label, are typically less than about 5 mCi.

[0149] While a great deal of clinical experience has been gained with ^{131}I and ^{90}Y , other radiolabels are known in the art and have been used for similar purposes. Still other radioisotopes are used for imaging. For example, additional radioisotopes which are compatible with the scope of the instant disclosure include, but are not limited to, ^{123}I , ^{125}I , ^{32}P , ^{57}Co , ^{64}Cu , ^{67}Cu , ^{77}Br , ^{81}Rb , ^{81}Kr , ^{87}Sr , ^{113}In , ^{127}Cs , ^{129}Cs , ^{132}I , ^{197}Hg , ^{203}Pb , ^{206}Bi , ^{177}Lu , ^{186}Re , ^{212}Pb , ^{212}Bi , ^{47}Sc , ^{105}Rh , ^{109}Pd , ^{153}Sm , ^{188}Re , ^{199}Au , ^{225}Ac , ^{211}A ^{213}Bi . In this respect alpha, gamma and beta emitters are all compatible within the instant disclosure. Further, in view of the instant disclosure it is submitted that one skilled in the art could readily determine which radionuclides are compatible with a selected course of treatment without undue experimentation. To this end, additional radionuclides which have already been used in clinical diagnosis include ^{125}I , ^{123}I , ^{99}Tc , ^{43}K , ^{52}Fe , ^{67}Ga , ^{68}Ga , as well as ^{111}In . Antibodies have also been labeled with a variety of radionuclides for potential use in targeted immunotherapy (Peirersz et al. *Immunol. Cell Biol.* 65: 111-125 (1987)). These radionuclides include ^{188}Re and ^{186}Re as well as ^{199}Au and ^{67}Cu to a lesser extent. U.S. Pat. No. 5,460,785 provides additional data regarding such radioisotopes and is incorporated herein by reference.

[0150] As previously discussed, the antibodies, or fragments thereof, of the disclosure, can be administered in a pharmaceutically effective amount for the in vivo treatment of mammalian disorders. In this regard, it will be appreciated that the disclosed antibodies, or fragments thereof, will be formulated so as to facilitate administration and promote stability of the active agent. Preferably, pharmaceutical compositions in accordance with the present disclosure comprise a pharmaceutically acceptable, non-toxic, sterile carrier such as physiological saline, non-toxic buffers, preservatives and the like. For the purposes of the instant application, a pharmaceutically effective amount of a antibody of the disclosure, conjugated or unconjugated to a therapeutic agent, shall be held to mean an amount sufficient to achieve effective binding to a target and to achieve a benefit, e.g., to ameliorate symptoms of a disease or disorder or to detect a substance or a cell. In the case of tumor cells, the polypeptide will be preferably be capable of interacting with selected immunoreactive antigens on neoplastic or immunoreactive cells and provide for an increase in the death of those cells. Of course, the pharmaceutical compositions of the present disclosure may be administered in single or multiple doses to provide for a pharmaceutically effective amount of the polypeptide.

[0151] In keeping with the scope of the present disclosure, the antibodies of the disclosure may be

administered to a human or other animal in accordance with the aforementioned methods of treatment in an amount sufficient to produce a therapeutic or prophylactic effect. The polypeptides of the disclosure can be administered to such human or other animal in a conventional dosage form prepared by combining the antibody of the disclosure with a conventional pharmaceutically acceptable carrier or diluent according to known techniques. It will be recognized by one of skill in the art that the form and character of the pharmaceutically acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. Those skilled in the art will further appreciate that a cocktail comprising one or more species of polypeptides according to the present disclosure may prove to be particularly effective.

VI. Methods of Treating Kallidin or des-Arg10-Kallidin-Associated Disease or Disorders

[0152] The anti-Kallidin or des-Arg10-Kallidin antibodies, or fragments thereof, of the disclosure are useful for antagonizing Kallidin or des-Arg10-Kallidin activity. Accordingly, in another aspect, the disclosure provides methods for treating Kallidin or des-Arg10-Kallidin-associated diseases or disorders by administering to a subject in need of thereof a pharmaceutical composition comprising one or more anti-Kallidin or des-Arg10-Kallidin antibody, or antigen binding fragment thereof of the disclosure.

[0153] Kallidin or des-Arg10-Kallidin-associated diseases or disorders amenable to treatment include, without limitation, pathophysiologic conditions such as inflammation, trauma, burns, shock, allergy, acute or chronic pain, and fibrosis, e.g., renal fibrosis. In certain exemplary, embodiments, antibodies of the disclosure may be issued to treat renal fibrosis and associated acute kidney injury as well as chronic kidney diseases which are the main causes of end-stage renal failure.

[0154] One skilled in the art would be able, by routine experimentation, to determine what an effective, non-toxic amount of antibody (or additional therapeutic agent) would be for the purpose of treating a Kallidin or des-Arg10-Kallidin-associated disease or disorder. For example, a therapeutically active amount of a polypeptide may vary according to factors such as the disease stage (e.g., stage I versus stage IV), age, sex, medical complications (e.g., immunosuppressed conditions or diseases) and weight of the subject, and the ability of the antibody to elicit a desired response in the subject. The dosage regimen may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily, or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation. Generally, however, an effective dosage is expected to be in the range of about 0.05 to 100 milligrams per kilogram body weight per day and more preferably from about 0.5 to 10, milligrams per kilogram body weight per day.

VII. Exemplification

[0155] The disclosure further provides the following examples which should not be construed as further limiting.

[0156] Furthermore, in accordance with the present disclosure there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (herein "Sambrook et al., 1989"); *DNA Cloning: A Practical Approach*, Volumes I and II (D.N. Glover ed. 1985); *Oligonucleotide Synthesis* (M.J. Gait ed. 1984); *Nucleic Acid Hybridization* [B.D. Hames & S.J. Higgins eds. (1985)]; *Transcription And Translation* [B.D. Hames & S.J. Higgins, eds. (1984)]; *Animal Cell Culture* [R.I. Freshney, ed. (1986)]; *Immobilized Cells And Enzymes* [IRL Press, (1986)]; B. Perbal, *A Practical Guide To Molecular Cloning* (1984); F.M. Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994).

Example 1: Hybridoma Production: Immunization of Mice with Kallidin Peptide Conjugated to KLH and Antibody Generation against Human BKR1 Ligands

[0157] The objective was to develop cross-reactive antibodies against Kallidin (KD; SEQ ID NO:1) and des-arg-Kallidin (DAKD; SEQ ID NO:2) that would inhibit these ligands binding to the human BKR1. Generally, immunization of mice with KLH conjugated KD through additional cysteines on either the C- or N- terminus of the peptide was used to obtain mouse splenocytes for fusion with mouse myeloma cell lines as a fusion partner to produce the hybridomas.

[0158] Briefly, the immunization protocol was as follows: BALB/c Mice (8-20 week-old naïve female) were immunized intraperitoneally with a mixture of even amounts of KLH-KD and KD-KLH in phosphate buffered saline (PBS) as an antigen total of 100 ug per mouse mixed at 1:1 ratio of Sigma Adjuvant System (Sigma cat #6322) in a total volume of 200 µl per mouse (day 0). On day 21, mice were boosted with a mixture of even amounts of KLH-KD and KD-KLH in PBS as an antigen total of 50 ug per mouse mixed at 1:1 ratio of Sigma Adjuvant System (Sigma cat #6322) in a total volume of 200 µl per mouse. On day 30, blood samples were harvested for KD specific antibody titer evaluation. On day 51, mice were boosted for fusion with a mixture of even amounts of KLH-KD and KD-KLH in PBS as an antigen total of 50 ug per mouse mixed at 1:1 ratio of Sigma Adjuvant System (Sigma cat #6322) in a total volume of 200 µl per mouse. At day 55 mice were sacrificed by CO₂ chamber, blood was collected through the cardiac puncture and spleen was harvested for hybridoma production.

[0159] Hybridomas were made by fusing mouse myeloma cells that are deficient in adenosine phosphoribosyltransferase (APRT) with spleen cells from mice immunized with specific antigens. A selection system using HAT (hypoxanthine, azaserine, and thymidine) medium eliminates all but the fusion cells that are APRT⁺. Successful hybridomas must also retain the immunoglobulin (Igh) heavy chain, one of the immunoglobulin light chain loci and secrete functional antibody.

[0160] Hybridoma Production Medium (IMDM) was made by combining the following: 500ml Iscove's Modified Dulbecco's Medium (HyClone SH30259.01), 50ml fetal bovine serum (HyClone SH30070.03), 5ml L-glutamine (Gibco Invitrogen cat # 25030), 5ml non-essential amino acids (Gibco Invitrogen cat # 11140050), 5ml sodium pyruvate (Gibco Invitrogen cat # 11360070), 5ml

0.1% penicillin-streptomycin (Gibco Invitrogen cat # 15140148). The medium was filtered before use. Expansion medium was made by combining the following: 1000ml serum free medium (Gibco Hybridoma SFM # 12045), 100ml 10% HyClone SuperLow IgG Defined FBS # SH30898.03 and 10ml penicillin/streptomycin. Freezing medium was 45ml heat inactivated FBS (HyClone SH30070.03) and 5ml DMSO, filter sterilized. Other materials included the following : HAT (50x) was obtained from Sigma-Aldrich (# HO262); Hybridoma Fusion and Cloning Supplement (50X) (Roche Diagnostics 11 363 735 001); Trypan Blue Stain 0.4% (Invitrogen cat # 15250-061 or T10282); PEG 1500 in 75 mM Hepes 50% w/v (Roche cat # 783641(10783641001)). All the reagents except HAT and Hybridoma Fusion and cloning supplement were used at 37°C.

Table 2. Peptide Reagents Used in Immunization and Screening

Peptide No.	SEQ ID NO.	Peptide Sequence	Peptide Name	Alternative Name
1	5	RPPGFSPFR	bradykinin	BK
2	70	biotin-RPPGFSPFR		b-BK
3	71	RPPGFSPFR-biotin		BK-b
4	72	KLH-RPPGFSPFR		KLH-BK
5	73	RPPGFSPFR-KLH		BK-KLH
6	1	KRPPGFSPFR	kallidin	KD
7	74	biotin-KRPPGFSPFR		b-KD
8	75	KRPPGFSPFR-biotin		KD-b
9	76	KLH-KRPPGFSPFR		KLH-KD
10	77	KRPPGFSPFR-KLH		KD-KLH
11	6	RPPGFSPF	desArg9bradykinin	DABK
12	78	biotin-RPPGFSPF		b-DABK
13	79	RPPGFSPF-biotin		DABK-b
14	80	KLH-RPPGFSPF		KLHDABK
15	81	RPPGFSPF-KLH		DABK-KLH
16	2	KRPPGFSPF	desArg10kallidin	DAKD
17	82	biotin-KRPPGFSPF		b-DAKD
18	83	KRPPGFSPF-biotin		DAKD-b
19	84	KLH-KRPPGFSPF		KLH-DAKD
20	85	KRPPGFSPF-KLH		DAKD-KLH
21	3	RRPPGFSPFR	Kallidin like peptide	KLP
		biotin-		

Peptide No.	SEQ ID NO.	Peptide Sequence	Peptide Name	Alternative Name
22	86	RRPPGFSPFR		b-KLP
23	87	RRPPGFSPFR-biotin		KLP-b
24	88	KLH-RRPPGFSPFR		KLH-KLP
25	89	RRPPGFSPFR-KLH		KLP-KLH
26	90	RRPPGFSPF	desArg10kallidin like peptide	DAKLP
27	91	biotin-RRPPGFSPF		b-DAKLP
28	92	RRPPGFSPF-biotin		DAKLP-b
29	93	KLH-RRPPGFSPF		KLH-DAKLP
30	94	RRPPGFSPF-KLH		DAKLP-b
31	95	RPPGF	bradykinin1-5	BK15
32	96	biotin-RPPGF		b-BK15

[0161] Briefly, three or four days before the fusion, the mouse was boosted with an antigen of interest either intraperitoneally or intravenously. On the day of the fusion, the mouse was sacrificed in CO₂ chamber, blood was collected by cardiac puncture and the spleen was taken out and placed into 10 ml of serum free IMDM in a Petri dish. Fusion partner cells myeloma: FO (ATCC ref CRL-1646)/ X63 Ag8.653 (ATCC ref CRL1580) were grown at a log phase, then split one day before the fusion (1:2 and 1:5), and collected into 20 ml centrifuge tubes, spun and resuspended the pellet in 10ml IMDM. The pellet was washed two times with serum free IMDM medium. All the centrifugations are performed at 1570 rpm for 5 min. Final resuspension was in 10ml serum free IMDM. The connective tissue was dissected away from the spleen. The spleen was injected with 1 ml of serum free IMDM preheated to 37°C by 1 ml syringe and 25-gauge needle. Splenocytes are squeezed out of the fibroelastic coat by forceps and washed two times in 10 ml of serum free IMDM (including initial spin) and were resuspended in 10ml serum free IMDM. Cells were counted in Countess Automated Cell Counter.

[0162] Fusion partner cells and splenocytes were combined in one 50ml tube at ratio of 1:2 to 1:10 (by cell number) and spun down at 970 rpm for 10 min (slow spin) to form a loose pellet. After the "slow" spin, supernatant was taken out with the precaution not to disturb the pellet, but minimize the amount of liquid over the cells in order not to dilute PEG 1500. The last remaining medium was reserved and added back after the PEG is added (below). Preheated PEG 1500 (37°C, total 1 ml) was added drop by drop to the cell pellet over 1 minute period of time and cells were mixed after every drop of PEG was added. Pellet was incubated with PEG for another 1 minute followed by addition of 10 ml of serum-free IMDM medium over 1 minute, so that the first 1 ml out of 10 is added over 30 sec. Cells underwent slow spin at 970 rpm for 10 min and supernatant decanted.

Into (2) 100ml troughs, the following was added: 70ml IMDM with 10% FBS, 2ml HAT and 2ml Hybridoma and Fusion Cloning Supplement. Cells were resuspended in 10ml IMDM with 10% FBS and split into (2) 50ml tubes (5ml cells/tube) and 25ml IMDM with 10% FBS was added. The resulting 30ml was transferred to the troughs containing 70ml HBSS/HAT/cloning supplement and 200ul cells/well were pipetted into (10) 96-well plates. Fusion was ready for screening by ELISA (50ul) about 10 to 14 days later, or when medium in the wells turns yellow. After the primary screening, positive clones are selected, numbered and moved to a 24-well plate in 500 ul per well of IMDM with 10% FBS. Hybridoma supernatants were screened by ELISA on streptavidin plated coated with N- and C-term biotinylated peptides (see below).

Example 2: Characterization and Selection of Hybridomas Expressing Antibodies Against Human BKR1 Ligands

[0163] Hybridoma supernatants were screened by ELISA on streptavidin plated coated with N- and C-term biotinylated peptides (see e.g., those set forth in Table 2) and then antibody binding kinetics were determined for confirmed positive hybridoma clones.

[0164] The ability of the antibodies in hybridoma supernatants to bind to BKR1 ligand peptide was evaluated with an ELISA assay. DAKD-biotin or KD-biotin peptides was coated on a 96-well SA plate in phosphate buffered saline (PBS) buffer for an hour at room temperature at 5 ug/ml, and the nonspecific binding sites were blocked with 1% bovine serum albumin (BSA) in PBS buffer. This plate was used to perform primary and secondary screening of the crude hybridoma supernatants. Hybridoma supernatants were added to the plates for binding to the coated KD or DAKD peptides. After 1 hour incubation, the plate was washed and bound antibodies were detected using horseradish peroxidase (HRP) conjugated secondary antibody (HRP-goat antimouse IgG (H+L): Jackson ImmunoResearch Labs # 115-035-166) and developed using 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) substrate (Roche diagnostics # 11 204 521 001). Data was analyzed using Excel. The antibodies showing positive signals (2 fold higher than 1:10000 serum dilution ELISA signal) were selected and re-screened in duplicates for confirmation. Confirmed positive hybridoma clones were selected and subjected to binding dissociation rate ranking by Biacore.

[0165] For antibody binding kinetics, the instruments used were the BIACORE 2000 or BIACORE 3000 (GE Healthcare), designed for biomolecular interaction analysis (BIA) in real time. The sensor chip used was SA chip (GE Healthcare) with streptavidin covalently immobilized on a carboxymethylated dextran matrix. Each sensor chip has four parallel flow cells (Fc). Every biotinylated BKR1 or BKR2 ligand peptides were immobilized to one of the flow cells 2 to 4 (Fc2 to Fc4) in the SA chip for binding dissociation rate screening and selectivity screening. Flow cell 1 (Fc1) was reserved and immobilized with a random peptide (biotinylated at one terminus) with equal or close peptide length in comparison to the testing ligand peptides as the negative control. In screening assays, cell culture supernatants of the hybridoma clones selected through primary screening of transiently expressed humanized variants were injected over immobilized peptides. Hybridoma cell culture media was also injected over the chip surface as blank to establish a baseline. After subtracting signals of Fc1 and blank buffer runs, the dissociation rate of the antibodies from the supernatants to each peptide was analyzed and ranked using BIAevaluation

software. Only the antibody clones that demonstrated superior ($k_d < 10^{-4}$ 1/s) binding dissociation rate were selected for subcloning and further characterization. In kinetics analysis, the corresponding biotin-peptides identified in screening for the testing antibody were immobilized in Fc2 to Fc4 while Fc1 with a random peptide used as reference cell. Each purified antibody selected from screenings were made into a series of two fold dilutions in running buffer (1 x HBS-EP buffer, GE Healthcare) between 0.1 to 10 nM. Binding association rate, dissociation rate and the overall affinity were calculated in BIAevaluation. Antibody binding kinetics for each antibody was always confirmed in triplicate assays using Biacore.

[0166] A total of 8 mice were immunized with mixed KLH-KD/KD-KLH and KLH-DAKD/DAKD-KLH and the spleens were fused using the above protocols. After primary screening of about 7680 hybridoma clones in ELISA with DAKD-biotin and KD-biotin, only 76 clones were confirmed positive and selected for binding dissociation rate ranking in Biacore 3000/2000 over the immobilized DAKD-biotin and KD-biotin on Streptavidin (SA) chips. Among those, 8 hybridoma clones with binding dissociation rate $\leq 10^{-4}$ were subcloned, sequenced, purified and further characterized (see Table 3).

Table 3 : Immunization Results with KLH-KD/KD-KLH and KLH-DAKD/DAKD-KLH

Ligand	Peptide Used in Assay	Assay	Clone ID							
			B21	C63	F151	F306	I2	I8	I22**	I54**
DAKD	b-DAKD	ELISA	+	+	-	-	+	+	+	+
		Biacore (KD,M)	-	-	-	-	-	-	NS	NS
	DAKD-b	ELISA	+	+	+	+	+	+	+	+
		Biacore (KD,M)	4.15 E-11	1.42 E-10	1.60 E-10	1.60 E-10	1.10 E-10	6.25 E-10	NTD	NTD
	DAKD (50nM)	FLIPR	22-25 ₁	-	6.9	6.9			9.4	8.1
		(nM)	9							
BK	b-DABK	ELISA	+/-	+/-	-	-	+	+	+	+
		Biacore (KD,M)	-	-	-	-	-	-	NS	NS
	DABK-b	ELISA	+	+/-	-	-	+	+	+	+
		Biacore (KD,M)	-*	-	-	-	-	-	NS	NS
	K	FLIPR								

DAI	DAB									
-----	-----	--	--	--	--	--	--	--	--	--

		(nM)								
BK	b-BK	ELISA	-	-	-	-	-	-	-	-
		Biacore (KD,M)	-	-	-	-	-	-	NS	NS
	BK-b	ELISA	+	-	-	-	+	+	+	+
		Biacore (KD,M)	8.57 E-09	-	-	-	-	-	NS	NS
	BK	FLIPR (nM)	-	-	-	-				
KD	b-KD	ELISA	+	-	-	-	-	-	-	-
		Biacore (KD,M)	-	-	-	-	-	-	NS	NS
	KD-b	ELISA	+	+	+	+	+	+	+	+
		Biacore (KD,M)	8.4 E-11	2.21 E-10	2.9 E-11	2.9 E-11	7.8 E-11	7.53 E-10	NTD	NTD
	KD (15nM)/V)	FLIPR (nM)	12	-	3.0	3.0	7.3	8.3	25	30

NA = not applicable, negative in ELISA

NS = nonspecific binding

NTD = not to be determined

-* = residual binding (low RU in Biacore)

Based on results seen in Table 3, five clones with unique sequences were selected for kinetic studies. These antibodies were highly selective for DAKD-biotin, KD-biotin, DAKLP-biotin and KLP-biotin binding (see Table 4). They do not bind to other kinin peptides or to peptides biotinylated at the N-terminus.

Table 4. Summary of Kinetics of Selected anti-DAKD/KD Antibody Candidates

Antibody	DAKD-b		KD-b	
	koff	KD	koff	KD
C63	9.36E-05	1.42E-10	1.00E-04	2.21E-10
B21	9.89E-05	4.15E-11	2.04E-04	8.40E-11

	DAKD-b		KD-b	
Antibody	koff	KD	koff	KD
F151	1.36E-04	1.62E-10	2.00E-05	2.88E-11
I22	3.19E-04	2.17E-10	2.10E-05	4.40E-12
I54	3.06E-05	9.53E-12	3.88E-05	1.12E-11
	DAKL-b		KLP-b	
	koff	KD	koff	KD
C63	n/b	n/b	n/b	n/b
B21	2.30E-04	1.34E-10	1.12E-04	1.92E-10
F151	6.58E-05	2.12E-10	≤1.0E-06	≈1.66E-11
I22	≤1.0E-06	≈1.83E-12	1.03E-05	1.82E-12
I54	5.66E-05	1.17E-11	6.04E-05	9.56E-12
n/b = no binding				

[0167] Additional immunization were performed with an array of immunogens (see list of peptides, Table 2) for generating antibodies blocking the rodent BKR1 ligands, DABK and DAKD as well as antibodies with other binding specificities against different member of kinin family of peptides. Table 5 lists the heavy and light sequences of the antibodies generated.

Table 5 . Heavy and Light Chain Sequences of Antibodies

Antibody	Isotype	SEQ ID NO.	Heavy Chain Sequence
B21	IgG1/k	97	LPEFQVKLEESGAEIVRSASVVKLSCTASGFNIKDYLLHWVKORPEQGLEWIGWIDPENGDT GYARKEFGKATMTADTSSNTVYLHLSSLTSEDVAVYYFNAWEYDGYDLDYWGQGTSTVTVSS AKTTPPSVYGSS
C63		98	LPEFQVKLEESGGGLVQPGGGMKLSCTASGFNFKDYLLHWVKORPEQGLEWVAEIRSKSN NYATHYAESVKGRTISRDDSKSSVYLQMNRLRAEDTGIYYCIGEDYGGDYWGQGTSTVTV SSAKTTPPSVYGSS
F151		99	LPEFEVQLEESGPELVKPGTSVKVSCASGYSFTDYNLYWVKSHGKSLWIGYFDPYNGNT GYNQKFRGKATLTVDKSSSTAFMHLSSLTSDDSAVYYCANYYRYDDHMDYWGQGTSTVTVSS AKTTPPSVYGSS
I22		100	LPEFEVKLEESGAEIVRSASVVKLSCTASGFNIKDYLLHWVKORPEQGLEWIGWIDPENGDS DYAPKFGKATMTADTSSNTVYLQFSSLTSEDVAVYYCNAFEYDGNYSPLDFWGQGTSTVTVSS SAKTTPPSVYGSS
I54		101	LPEFEVKLEESGAEIVRSASVVKLSCTASGFNIKDYLLHWVKORPEQGLEWIGWIDPENGDS DYAPKFGKATMTADTSSNTVYLQFSSLTSEDVAVYYCNAFEYDGNYSPLDFWGQGTSTVTVSS SAKTTPPSVYGSS
B21		38	EVQLQQSGAEIVRSASVVKLSCTASGFNIKDYLLHWVKORPEQGLEWIGWIDPENGDTGYAR KFGKATMTADTSSNTVYLHLSSLTSEDVAVYYFNAWEYDGYDLDYWGQGTSTVTVSSAKTT PPS
C63		45	EVKLEESGGGLVQPGGGMKLSCTASGFNFKDYLLHWVKORPEQGLEWVAEIRSKSNNYATHY AESVKGRTISRDDSKSSVYLQMNRLRAEDTGIYYCIGEDYGGDYWGQGTSTVTVSSAKTTP S

F151	mIgG1/K	19	ETQLQQSGPELVKPGTSVKVSC <u>CK</u> ASGYSTFDYNIYVVKQSHGKSLEWIGYFDPYNGNTGYNQ KFRGKATLTVDKSSSTAFMHLSSLTSDDSAVYYC <u>AN</u> YYRYDDHAMDYWGOGTSVTVSSAKTT PPS
I22		53	EVQLQQSGAELVRSAGASVKLSCTASGFNIRKDYIMHWVKQRPEQGLEWIGWVDPENGDSFYAP KFQGGKATMTADTSSNTVYLVQFSLSLTSEDYAVYYCNAFEYDGNYSPLDFWGGOGTSVTVSSAKT TPPS
154		61	EVQLQQSGAELVRSAGASVKLSCTASGFNIRKDYIMHWVKQRPEQGLEWIGWVDPENGDSFYAP KFQGGKATMTADTSSNTVYLVQFSLSLTSEDYAVYYCNAFEYDGNYSPLDFWGGOGTSVTVSSAKT TPPS
Antibody	Isotype	SEQ ID NO.	LightChainSequence

B21	IgG1/k	102	ELDIVMTQTTLTSLVTIGQPASIS <u>CKSSQSLLYS</u> NGRTYLNWLLORPGQSPKRLIYLVSKLD SGVPDRFTGSGSGTDFTLKIRVEAEDLGVIY <u>CCLOGTHFPYTFGGG</u> KLEIKRADAAPT VSI FPPSKLELY
C63		103	ELDIVLTQSPSSLAVSVGEKVTMS <u>CKSSQSLLYSSDQRNYLAWYQQRSGQSPKLLIYWASTR</u> <u>ESGVPDRFTGSGSGTDFTLT</u> ISSVKAEDLAVYY <u>CQQYYSYPYTFGGG</u> KLEIKRADAAPT VSI IFPPSKLELY
F151		104	ELDIVMTQTPSSLAVSVGEKVTMS <u>CKSSQSLLYTSNQKNYLAWYQQRSGQSPKPLIYWASTR</u> <u>ESGVPDRFTGSGSGTDFTLT</u> ISSVKAEDLAIYY <u>CQQYYSYPWTFGGG</u> KLEIKRADAAPT VSI IFPPSKLELY
I22		105	ELDIVITQTTLTSLVPIGQPASIS <u>CKSRQSLLYSNGETYLNWLLORPGQSPKRLIYLVSKLD</u> SGVPDRFTGSRSGTDFTLKISRVEAEDLGVIY <u>CCMQGTHFPYTFGGG</u> KLEIKRADAAPT VSI FPPSKLELY
154		106	ELDIVITQSTTLTSLVPIGQPASIS <u>CKSSQSLLYSNGETYLNWLLORPGQSPKRLIYLVSKLD</u> SGVPDRFTGSRSGTDFTLKISRVEAEDLGVIY <u>CCMQGTHFPYTFGGG</u> KLEIKRADAAPT VSI FPPSKLELY
B21	mIgG1/K	39	DVVMQTPTLTLTSLVTIGQPASIS <u>CKSSQSLLYS</u> NGRTYLNWLLORPGQSPKRLIYLVSKLD <u>SG</u> VPDRFTGSGSGTDFTLTKIRVEAEDLGVIY <u>CCLOGTHFPYTFGGG</u> KLEIKRADAAPT
C63		46	DIVMSQSPSSLAVSVGEKVTMS <u>CKSSQSLLYSSDQRNYLAWYQQRSGQSPKLLIYWASTRES</u> <u>GVPDRFTGSGSGTDFTLT</u> ISSVKAEDLAVYY <u>CQQYYSYPYTFGGG</u> KLEIKRADAAPT
F151		26	DIVMSQSPSSLAVSVGEKVTMS <u>CKSSQSLLYSSNQKNYLAWYQQRSGQSPKPLIYWASTRES</u> <u>GVPDRFTGSGSGTDFTLT</u> ISSVKAEDLAIYY <u>CQQYYSYPWTFGGG</u> KLEIKRADAAPT
I22		54	DIVMTQSPSSLVSAAGEKVTMS <u>CKSSQSLLNSGNQKNYLAWYQQRSGQSPKPLIYGASTRES</u> <u>GVPDRFTGSGSGTDFTLT</u> ISSVQAEDLAVYY <u>CQNDHSYPLTFGAGT</u> KLEIKRADAAPT
154		62	DVVMQTPTLTLTSLVPIGQPASIS <u>CKSSQSLLYS</u> NGRTYLNWLLORPGQSPKRLIYLVSKLD <u>SG</u> VPDRFTGSRSGTDFTLKISRVEAEDLGVIY <u>CCMQGTHFPYTFGGG</u> KLEIKRADAAPT
Single underscore = CDR region; double underscore = signature amino acids for identifying CDRs			

EXAMPLE 3: Generation of Surrogate Antibody for Murine Animal Studies

[0168] A surrogate antibody to be used in murine animal studies needed to be able to bind and

neutralize rodent BKR1 ligands, DABK and DAKLP (mouse equivalent of DAKD). In order to generate the required surrogate antibody, mice were first immunized with DABK and/or DAKD with KLH directly conjugated to the N-terminals of the peptides. Biotin-DABK/biotin-DAKD (biotinylation directly on N-terminus of the peptide) positive hybridoma clones from ELISA screening were selected for scaling up and purification. The antibodies listed in Family 7 (see Table 12) that demonstrated high binding affinities to biotin-DABK, biotin-DAKLP and biotin-DAKD were selected based on Biacore direct binding assay (Table 10). However, these Family 7 antibodies showed no binding to the native, unmodified DABK and DAKD peptides in competitive ELISA, and lacked neutralizing functionality in a calcium influx assay with Functional Drug Screening System (FDSS) (Hamamatsu Photonics K.K., Japan). Moreover, the biotin-DABK and biotin-DAKD completely lost bioactivity in the FDSS assay in comparison to the native, unmodified DABK and DAKD peptides (data not shown).

[0169] It was hypothesized that the direct N-terminus conjugation of KLH and biotin prevented the native conformation of DABK and DAKD to form. With the aim to restore the native conformation in KLH- and biotin- conjugated peptides, linkers were designed and added to the N-terminus of DABK and/or DAKD with the intention to "cushion" the KLH/biotin conjugation effects on peptide conformation. Poly-glycine linkers were first attempted and tested because of their simple, nonpolar and neutral properties based on modeling results. The FDSS assay results indicated that the gly-gly-gly (3G) linker was the best according to its ability to restore the bioactivities of KLH and biotin conjugated DABK and DAKD peptides (data not shown). Therefore, KLH-3G-DABK was chosen to immunize mice. And biotin-3G-DABK and biotin-3G-KD were used in binding based screening assays (ELISA and Biacore). Several DABK/DAKD specific antibodies (Family 3, see Table 13) were identified in this new round of surrogate antibody hybridoma selection. EE1 was selected as the lead surrogate antibody based on its superior binding affinity and neutralization activity against native DABK/DAKD and lack of cross-reactivity to other peptides (see Tables 6 - 12)

[0170] Antibodies with different specificities were generated when using the different immunogens listed in Table 13. Family 4 antibodies were specific to the BKR2 receptor ligands, BK and KD. Family 5 antibodies specifically bind to the C terminus of BK and DABK. Family 6 antibodies bind BK, DABK and DAKD but do not bind to KD.

[0171] Additional linkers were evaluated for binding to the surrogate EE1 antibody for their ability to fit into the DABK/DAKD binding pocket in EE1, including longer poly-glycine linkers, poly-alanine linkers and preexisting linkers such as polyethylene glycol (PEG2) linker and aminohexanoic acid (Ahx) linker (a 6-carbon inert linker). All linker peptides were custom synthesized by Abgent (Can Diego, CA). All tested biotinylated peptides with linkers (biotin-linker-DABK/DAKD) bound well to EE1, indicating that any inert N-terminus linkers helped DABK and DAKD peptides to retain their native bioactive conformation when conjugated with biotin and other molecules. In contrast, no binding or poor binding to EE1 was observed with biotin-DABK and biotin-DAKD, peptides that have direct N-terminal biotin conjugation (see Figure 1).

[0172] The binding kinetics of generated antibodies are summarized in Tables 5-11. Then, all antibodies generated were sorted into families and their binding specificities are summarized below in Table 12. Table 13 provides the heavy and light chain sequences of antibodies that were placed into family 1 and family 2 based on their binding specificity (see Table 12).

Table 6. Summary of Antibody Kinetics to b-3G-DABK and b-3G-DAKD Peptides

Clone	b-3G-DABK			b-3G-DAKD		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DD20	1.5E+06	2.3E-04	1.6E-10	4.7E+05	4.0E-04	8.8E-10
UR11	2.0E+05	3.0E-04	1.5E-09	3.0E+05	1.6E-03	5.2E-09
DD7	2.3E+05	6.0E-04	2.7E-09	2.1E+05	1.4E-03	6.6E-09
EE1	4.4E+05	1.2E-04	2.8E-10	4.4E+05	2.0E-04	4.5E-10
EE36	4.3E+03	5.3E-04	1.2E-07	n/b	n/b	n/b
UR29	n/b*	n/b	n/b	n/b	n/b	n/b
JK3	3.44E+05	3.91E-05	1.14E-10	3.18E+05	5.07E-05	1.60E-10
LR4	n/b	n/b	n/b	n/b	n/b	n/b
LR16	n/b	n/b	n/b	n/b	n/b	n/b
LR6	n/b	n/b	n/b	n/b	n/b	n/b
LR12	n/b	n/b	n/b	n/b	n/b	n/b
NR1	n/b	n/b	n/b	n/b	n/b	n/b
NR15	n/b	n/b	n/b	n/b	n/b	n/b

n/b* = non-specific binding, n/b = no binding

Table 7. Summary of Antibody Kinetics to b-3G-DAKLP and b-3G-BK Peptides

Clone	b-3G-DAKLP			b-3G-BK		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DD20	3.1E+05	6.1E-04	2.0E-09	3.0E+05	7.1E-04	2.3E-09
UR11	n/b	n/b	n/b	n/b	n/b	n/b
DD7	1.7E+05	2.1E-03	1.2E-08	1.3E+05	8.8E-04	6.8E-09
EE1	4.2E+05	2.9E-04	6.8E-10	2.5E+05	2.6E-03	1.1E-08
EE36	n/b	n/b	n/b	n/b	n/b	n/b
UR29	n/b	n/b	n/b	n/b	n/b	n/b
JK3	ND	ND	ND	n/b	n/b	n/b
LR4	n/b	n/b	n/b	n/b	n/b	n/b
LR16	n/b	n/b	n/b	n/b	n/b	n/b
LR6	n/b	n/b	n/b	n/b	n/b	n/b
LR12	n/b	n/b	n/b	n/b	n/b	n/b
NR1	n/b	n/b	n/b	n/b	n/b	n/b
NR15	n/b	n/b	n/b	n/b	n/b	n/b

n/b = no binding; ND = not determined

Table 8. Summary of Antibody Kinetics to b-3G-KLP and b-3G-KD Peptides

Clone	b-3G-KLP			b-3G-KD		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DD20	3.8E+05	5.7E-04	1.5E-09	n/b	n/b	n/b
UR11	n/b	n/b	n/b	1.2E+05	1.3E-03	1.1E-08
DD7	1.5E+05	2.1E-03	1.5E-08	2.2E+05	1.8E-03	8.4E-09
EE1	4.0E+05	2.1E-03	5.3E-09	n/b	n/b	n/b
EE36	n/b	n/b	n/b	n/b	n/b	n/b
UR29	n/b	n/b	n/b	n/b	n/b	n/b
JK3	ND	ND	ND	n/b	n/b	n/b
LR4	n/b	n/b	n/b	n/b	n/b	n/b
LR16	n/b	n/b	n/b	n/b	n/b	n/b
LR6	n/b	n/b	n/b	n/b	n/b	n/b
LR12	n/b	n/b	n/b	n/b	n/b	n/b
NR1	n/b	n/b	n/b	n/b	n/b	n/b
NR15	n/b	n/b	n/b	n/b	n/b	n/b

n/b = no binding; ND = not determined

Table 9. Summary of Antibody Kinetics to DABK-b and DAKLP-b Peptides

Clone	DABK-b			DAKLP-b		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DD20	n/b	n/b	n/b	n/b	n/b	n/b
UR11	n/b	n/b	n/b	n/b	n/b	n/b
DD7	n/b	n/b	n/b	n/b	n/b	n/b
EE1	n/b	n/b	n/b	n/b	n/b	n/b
EE36	n/b	n/b	n/b	n/b	n/b	n/b
UR29	1.5E+06	5.8E-05	3.9E-11	3.0E+06	2.1E-03	6.8E-10
JK3	n/b	n/b	n/b	n/b	n/b	n/b
LR4	n/b	n/b	n/b	n/b	n/b	n/b
LR16	n/b	n/b	n/b	n/b	n/b	n/b
LR6	n/b	n/b	n/b	n/b	n/b	n/b
LR12	n/b	n/b	n/b	n/b	n/b	n/b
NR1	n/b	n/b	n/b	n/b	n/b	n/b
NR15	n/b	n/b	n/b	n/b	n/b	n/b

n/b = no binding; ND = not determined

Table 10. Summary of Antibody Kinetics to BK-b and b-BK Peptides

Clone	BK-b			b-BK		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DD20	n/b	n/b	n/b	n/b	n/b	n/b

Clone	BK-b			b-BK		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
UR11	n/b	n/b	n/b	n/b	n/b	n/b
DD7	n/b	n/b	n/b	n/b	n/b	n/b
EE1	n/b	n/b	n/b	n/b	n/b	n/b
EE36	n/b	n/b	n/b	n/b	n/b	n/b
UR29	1.5E+06	1.0E-04	7.2E-11	n/b	n/b	n/b
JK3	n/b	n/b	n/b	n/b	n/b	n/b
LR4	n/b	n/b	n/b	n/b	n/b	n/b
LR16	n/b	n/b	n/b	n/b	n/b	n/b
LR6	n/b	n/b	n/b	n/b	n/b	n/b
LR12	n/b	n/b	n/b	n/b	n/b	n/b
NR1	n/b	n/b	n/b	8.69E+04	8.77E-04	1.01E-08
NR15	n/b	n/b	n/b	2.95E+05	1.09E-03	3.68E-09

n/b = no binding; ND = not determined

Table 11. Summary of Antibody Kinetics to b-DABK and b-DAKD Peptides

Clone	b-DABK			b-DAKD		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DD20	n/b	n/b	n/b	n/b	n/b	n/b
UR11	n/b	n/b	n/b	n/b	n/b	n/b
DD7	n/b	n/b	n/b	n/b	n/b	n/b
EE1	n/b	n/b	n/b	n/b	n/b	n/b
EE36	n/b	n/b	n/b	n/b	n/b	n/b
UR29	n/b	n/b	n/b	n/b	n/b	n/b
JK3	n/b	n/b	n/b	n/b	n/b	n/b
LR4	1.48E+05	1.04E-03	7.15E-09	3.27E+05	7.63E-04	2.36E-09
LR16	4.34E+05	4.38E-05	1.01E-10	2.07E+05	3.39E-03	1.65E-08
LR6	n/b	n/b	n/b	n/b	n/b	n/b
LR12	2.91E+05	5.40E-04	3.63E-09	n/b	n/b	n/b
NR1	n/b	n/b	n/b	n/b	n/b	n/b
NR15	n/b	n/b	n/b	n/b	n/b	n/b

n/b = no binding; ND = not determined

Table 12. Summary of Antibody Kinetics to b-DAKLP and b-KD Peptides

Clone	b-DAKLP			b-KD		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DD20	n/b	n/b	n/b	n/b	n/b	n/b
UR11	n/b	n/b	n/b	n/b	n/b	n/b

Clone	b-DAKLP			b-KD		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DD7	n/b	n/b	n/b	n/b	n/b	n/b
EE1	n/b	n/b	n/b	n/b	n/b	n/b
EE36	n/b	n/b	n/b	n/b	n/b	n/b
UR29	n/b	n/b	n/b	n/b	n/b	n/b
JK3	n/b	n/b	n/b	n/b	n/b	n/b
LR4	1.84E+05	2.58E-04	1.40E-09	n/b	n/b	n/b
LR16	2.34E+05	1.11E-04	4.74E-10	n/b	n/b	n/b
LR6	6.80E+05	4.01E-04	7.45E-10	n/b	n/b	n/b
LR12	n/b	n/b	n/b	n/b	n/b	n/b
NR1	n/b	n/b	n/b	1.66E+05	5.81E-03	3.56E-08
NR15	n/b	n/b	n/b	7.66E+05	5.66E-03	7.41E-09

n/b = no binding; ND = not determined

Table 13. Summary of Anti-kinin peptide antibody generation

Antibody Families	Immunogens	Representative antibodies	Binding Specificity
Family 1	KD-KLH + KLH-KD or DAKD-KLH + KLH-DAKD	F151, B21, I22, I54	N-terminus of DAKD (DAKLP) and KD (KLP)
Family 2	KD-KLH + KLH-KD or DAKD-KLH + KLH-DAKD	C63	N-terminus of DAKD and KD
Family 3	KLH-3G-DABK	EE1, DD20, JK3	C-terminus of DABK and DAKD (DAKLP)
Family 4	KLH-BK and BSA-BK	NR15, NR1	C-terminus of BK and KD
Family 5	KLH-BK	UR29	N-terminus of BK and DABK
Family 6	KLH-BK	UR11	BK, DABK and DAKD
Family 7	KLH-DABK or KLH-DAKD	LR4, LR6, LR12 and LR16	no binding with native peptides

Example 4: Characterization of des-arg-Kinin Ligand Depletion using Calcium Mobilization

[0173] A functional assay was used to further characterize the seven families of generated antibodies. The Bradykinin B1 Receptor signaling is Gq coupled, therefore receptor activation can be monitored using Gq activation of IP3 and downstream calcium mobilization. HEK mBKR1 (recombinant mouse bradykinin B1 receptor) cells or MRC5 (endogenous expression of bradykinin B2 receptors; (ATCC CCL-171)) were used to measure calcium mobilization.

[0174] Briefly, the mouse Bdkrb1 gene (sequence provided below) was amplified from mouse lung

cDNA (Biochain, Cat# C1334152) using PCR primers 804_cGWY_F: 5'-AAAAGCAGGCTTAGGAGCGGCCGCCATGGCGTCCCAGGCCTCGCTG-3' (SEQ ID NO: 107) and 804_cGWY_R: 5'-CAAGAAAGCTGGGTTCGGATCCTTATAAAGTTCCCAGAACCCCTGGTC-3' (SEQ ID NO: 108) and Pfu Polymerase (Agilent Technologies, Cat# 600264) and cloned into pDONR201 using BP clonase enzyme mix (Invitrogen, Cat# 11789-020). In parallel, the pEAK8 expression vector (EDGE Biosystems) was modified by inserting a N-terminal HA tag (GCATACCCATACGACGTCCCAGACTACGCT, GenBank SEQ ID NO:109 CY100443) into pEAK8 linearized with EcoRI and HindIII (vector pEAK8-nHA) and subsequent insertion of the Gateway cassette B (Invitrogen, Cat# 11828-029) into pEAK8_nHA digested with EcoRI and NotI and blunt-ended with Klenow polymerase (NEB, cat# M0210S) resulting in vector pEAK8_nHA_DEST. Next mouse Bdkrb1 was subcloned into pEAK8_nHA_DEST using LR clonase (Invitrogen, Cat# 11791-100). 293-PSC cells were then transfected with pEAK8-Bdkrb1 plasmid using Fugene 6 transfection reagent. The cells were put under antibiotic (puromycin) selection 24 hours after transfection, and selection was maintained to generate a stable cell line. Presence of the Bdkrb1 gene in the resultant stable cell lines was confirmed using real time RT-PCR, and by agarose gel electrophoresis. Cell surface expression of the Bradykinin B1 receptor was performed by using an antibody against the N-terminal-HA tag (Covance, Cat # MMS-101P) on the Bradykinin B1R on a FACS instrument. Functional activity of the Bradykinin B1 receptor was demonstrated in calcium mobilization assay with selective agonists.

Bdkrb1 gene subcloned into cells:

```
ATGGCGTCCCAGGCCTCGCTGAAGCTACAGCCTTCTAACCAGCCAGCAGGCCCTCCCAACATCACCTC
CTGCCAGGGCGCCCGGAAGCCTGGGATCTGCTGTGTCGGGTGCTGCCAGGGTTTGTCTACTGTCTGT
TCTTTGGCCTCCTGGGGAACCTTTAGTCTCTCTCTCTCTCTCTTTTGCCTTGGCGACGATGGTGGCAGCAG
CGGCGGCAGCGCCTAACCATAGCAGAAATCTACCTGGCTAACTTGGCAGCTTCTGATCTGGTGTGTGTGCT
GGGCTGCCCTTCTGGGAGAGAACGTTGGGAACCGTTTCAACTGGCCCTTTGGAAGTGACCTCTGCCGGG
TGGTCAGCGGGGTTCATCAAGGCCAACCTGTTCATCAGCATCTTCTGGTGGTGGCCATCAGTCAGGACCGC
TACAGGTGTGCTGGTATACCCCATGACCAGCTGGGGGAACCGCGCGCGACGGCAAGCCCAAGTGACCTGCCCT
GCTCATCTGGGTAGCTGGGGGCCCTCTTGAGCACCCCCACGTTCTTCTGCGTTCCGTCAAAGTCGTCCCTG
ATCTGAACATCTCTGCCCTGCATCCTGCTTTTCCCCACGAAGCTTGGCACTTTGTAAGGATGGTGGAGTTG
AACGTTTTGGGTTTCTCTCTCCATTGGCTGCCATCTCTACTTCACTTTACATCCTGGCCTCCCTGAG
AGGACAGAAGGAGGCCAGCAGAACCCGGTGTGGGGGACCCAAGGACAGCAAGACAATGGGGCTGATCCTCA
CACTGGTAGCCTCCTTCTGGTCTGCTGGGCCCCCTTACCACCTTCTTTGCCCTTCTGGATTTCCTGGTCCAG
GTGAGAGTGATCCAGGACTGCTTCTGGAAGGAGCTCACAGACCTGGGCCCTGCAGCTGGCCAACTTCTTTGC
TTTTGTCAACAGCTGCCTGAACCACTGATTTATGTCTTTGCAGGCCGGCTCTTTAAGACCAGGGTTCTGG
GAACCTTATAA (GenBank NM_007539; SEQ ID NO:110)
```

[0175] HEK mBKR1 or MRC5 cells were plated into 384 well clear bottom plates in growth medium, and allowed to attach overnight. Then growth media was removed, cells were washed in assay buffer (HBSS, 20mM HEPES, 2.5 mM probenecid), then dye-loaded with 0.5 uM Fluo-4AM, a cell permeable calcium sensing dye, with 0.04% Pluronic Acid for 1 hr at 37C. The AM ester is cleaved, and the calcium dye is retained in the cytoplasm. After 1 hr, the cells were washed to remove excess dye, and 20ul of residual buffer remained on the cells. Treatments were added as 2x solutions on the Functional Drug Screening System from Hamamatsu (FDSS), and the calcium mobilization was monitored kinetically for at least 4 minutes. B1R or B2R receptor activation results in G α_q mediated activation of phospholipase C and IP3 mediated calcium mobilization. The Fluo-4 dye chelates the released calcium, and a robust change in fluorescence is observed. The results were exported as max-min relative fluorescence units to normalize for differences between cell density or dye loading across the plate.

[0176] Ligand potency was determined each day by running concentration response curves of

ligand, and an approximate EC70-80 concentration of ligand was selected for incubation with antibodies. An EC80 concentration was selected because it is on the linear range of the detection curve and there was ample window to see a decrease with antagonists or ligand depleting antibodies. Dose response curve of antibodies were allowed to bind a EC80 concentration ligand, and the extent of ligand depletion was monitored using change in fluorescence. Results were normalized to buffer and EC80 ligand response, and an EC50 for ligand depletion was calculated. The results were then reported as molar ratio which corresponds to the Antibody concentration that reduces depletes 50% of the ligand response (i.e., EC50 of Ab) divided by the ligand concentration used. The theoretical max should be 0.5 because one unit of antibody should be able to deplete 2 units of ligand, but we have seen lower values in practice but that may be a reflection of the insensitivity of the detection method for low ligand concentrations, rather than a stoichiometric constraint for the antibody. The results of these experiments are set forth in Tables 14-16.

[0177] All family 1 and family 2 antibodies (see Table 13) demonstrated superior binding kinetics by Biacore (Table 3) and neutralization activity as measured by calcium mobilization against DAKD and KD peptides (Tables 14 and 15). The antibodies were further analyzed for their thermal stability and sequence suitability for humanization. F151 was advanced for humanization because it was thermally stable, there were no problematic residues in the CDR regions and it was cross-reactive to the mouse ligand KLP and DAKLP.

Table 14: Characterization of des-arg-Kinin Ligand Depletion using Calcium Mobilization in HEK mBKR1 cells

Family	Antibody	Depletion of DABK			Depletion of DAKD			Depletion of DAKLP		
		DABK (Mean Molar Ratio)	SD Molar Ratio	n	DAKD (Mean Molar Ratio)	SD Molar Ratio	n	DAKLP (Mean Molar Ratio)	SD Molar Ratio	n
1	F151	IA100		5	0.08	0.04	7	0.15	0.04	4
1	B21	IA100		1	0.15	0.04	3	0.67		1
1	122	IA100		1	0.07	0.02	3	0.21		1
1	154	IA100		1	0.15	0.05	3	0.35		1
2	C63	IA100		1	0.08	0.02	3	5.85		1
3	EE1	1.03	0.52	5	0.86	0.52	3	0.57	0.36	4
3	DD20	3.45	1.34	3	1.82	0.76	3	1.31	0.86	3
	DD7	2.18	0.45	3	4.22	0.95	3	5.34	1.22	2
3	JK3	1.86	0.03	2	ND			1.44	0.03	2
4	MBK3	ND			ND			ND		
4	NR15	ND			ND			ND		
4	NR1	ND			ND			ND		
5	UR29	0.60	0.12	5	IA200		3	IA300		4
6	UR11	6.99	1.61	3	19.65	14.95	3	11.09	3.13	2
7	LR4	IA100		1	IA400		1	IA400		1

		Depletion of DABK			Depletion of DAKD			Depletion of DAKLP		
Family	Antibody	DABK (Mean Molar Ratio)	SD Molar Ratio	n	DAKD (Mean Molar Ratio)	SD Molar Ratio	n	DAKLP (Mean Molar Ratio)	SD Molar Ratio	n
7	LR6	IA100		1	IA100		1	ND		
7	LR12	IA100		1	IA100		1	ND		
7	LR16	IA100		1	IA100		1	ND		

Antibodies were pre-incubated with a set concentration of ligand, usually an EC70-80 for activating calcium mobilization at the Bradykinin B1 Receptor. The antibody-ligand mixture was added to HEK mBKR1 cells pre-loaded with a calcium sensing dye (Fluo-4AM or Fluo-8AM) on the Hamamatsu FDSS6000 instrument, and calcium mobilization was monitored. Data was exported as a max-min relative fluorescence of the biological response, and IC50 for ligand depletion was calculated using sigmoidal curve fit in Graph Pad Prism V4.03. Data reported as molar ratio for ligand depletion by the antibody to standardize the different concentration of ligand that was used for the various experiments.

Molar Ratio for ligand depletion = [IC50 of Antibody]/ [Ligand]

SD = Standard Deviation; ND = not determined; IA100 = Inactive at 100nM; IA200 = Inactive at 200nM; IA300 = Inactive at 300nM; IA400 = Inactive at 400nM

Table 15: Characterization of Kinin Ligand Depletion using Calcium Mobilization in MRC5 Fetal Lung Fibroblasts cells

		Depletion of BK			Depletion of KD			Depletion of KLP		
Family	Antibody	BK (Mean Molar Ratio)	SD Molar Ratio	n	KD (Mean Molar Ratio)	SD Molar Ratio	n	KLP (Mean Molar Ratio)	SD Molar Ratio	n
1	F151	IA100		5	0.14	0.05	5	0.15	0.02	3
1	B21	IA100		1	0.33		1	ND		
1	I22	IA100		1	0.22		1	ND		
1	I54	IA100		1	0.30		1	ND		
2	C63	IA100		1	0.23		1	ND		
3	EE1	IA300		4	IA300		4	IA150		1
3	DD20	IA600		4	IA600		5	IA150		1
	DD7	7.11	3.62	3	17.37	12.11	3	4.27		1
3	JK3	IA300		2	IA300		2	ND		
4	MBK3	22.11	14.10	9	3.46	2.64	6	9.45		1
4	NR15	15.26	11.51	5	4.34	2.55	5	11.18		1
4	NR1	39.31		1	42.15		1	32.58		1
5	UR29	1.15	0.86	5	0.30	0.08	2	0.41		1
6	UR11	5.41	0.80	2	25.21	4.54	2	1.53		1

7	LR4	IA100		1	IA100		1	ND		
7	LR6	IA100		1	IA100		1	ND		
7	LR12	IA100		1	IA100		1	ND		
7	LR16	IA100		1	IA100		1	ND		
Antibodies were pre-incubated with a set concentration of ligand, usually an EC70-80 for activating calcium mobilization at the Bradykinin B2 Receptor. The antibody-ligand mixture was added to MRC5 Fetal Lung Fibroblasts (ATCC CCL-171) pre-loaded with a calcium sensing dye (Fluo-4AM or Fluo-8AM) on the Hamamatsu FDSS6000 instrument, and calcium mobilization was monitored. Data was exported as a max-min relative fluorescence of the biological response, and IC50 for ligand depletion was calculated using sigmoidal curve fit in Graph Pad Prism V4.03. Data reported as molar ratio for ligand depletion by the antibody to standardize the different concentration of ligand that was used for the various experiments.										
Molar Ratio for ligand depletion = [IC50 of Antibody]/ [Ligand]										
SD = Standard Deviation; ND = not determined; IA100 = Inactive at 100nM; IA150 = Inactive at 150nM; IA300 = Inactive at 300nM; IA400 = Inactive at 400nM; IA600 = Inactive at 600nM										

Example 5: Engineering of F151: Humanization, Stabilization and Mutation of Unwanted Sequence Motifs

1. HUMANIZATION

[0178] The humanization protocol used has been described in PCT/US08/74381 (US20110027266), herein incorporated by reference in its entirety. The variable light (VL) and variable heavy (VH) sequences of murine F151 were used to build a homology model of anti-DAKD/KD F151 LC and HC in Molecular Operating Environment (MOE; v. 2009.10; Chemical Computing Group). The following templates were used: light chain framework - 1SBS (93% identity in the framework regions), heavy chain framework - 2VXT (84% identity in the framework regions), L1 - 1LVE (93% identity), L2 - 1EEU (100% identity), L3 - 2R56 (93% identity), H1 - 1NJ9 (95% identity), H2 - 2VXU (76% identity) and H3 - 1HIL (49% identity). Templates were available at the RCSB Protein Data Bank found on the world wide web at rcsb.org, a website managed by Rutgers and the University of California San Diego (Berman, H.M; Westbrook J.; Feng. Z.; Gilliland, G.; Bhat, T.N.; Weissig, H.; Shindyalov, I.N.; Bourne, P.E. The Protein Data Bank, Nucleic Acids Research, 2000, 28, 235-242.). The homology model was subsequently energy minimized using the standard procedures implemented in MOE. A molecular dynamics (MD) simulation of the minimized 3D homology model of the murine F151 was subsequently performed, with constraints on the protein backbone at 500 K temperature for 1.1 nanoseconds (ns) in Generalized Born implicit solvent. Ten diverse conformations were extracted from this first MD run every 100 picoseconds (ps) for the last 1 ns. These diverse conformations were then each submitted to a MD simulation, with no constraints on the protein backbone and at 300 K temperature, for 2.3 ns. For each of the 10 MD runs, the last 2,000 snapshots, one every ps, from the MD trajectory were then

used to calculate, for each murine F151 amino acid, its root mean square deviations (rmsd) compared to a reference medoid position. By comparing the average rmsd on the 10 separate MD runs of a given amino-acid to the overall average rmsd of all F151 murine amino-acids, one decides if the amino-acid is flexible enough, as seen during the MD to be considered as likely to interact with T-cell receptors and responsible for activation of the immune response. 62 amino-acids were identified as flexible in the murine F151 antibody, excluding the CDR and its immediate 5 Å vicinity.

[0179] The motion of the 28 most flexible murine F151 amino acids, during the 20 ns (10 x 2 ns), were then compared to the motion of the corresponding flexible amino-acids of 49 human germline homology models, for each of which were run the 10 x 2 ns MD simulations. The 49 human germline models were built by systematically combining the 7 most common human germline light chains (vk1, vk2, vk3, vk4, vlamba1, vlamba2, vlamba3) and 7 most common human germline heavy chains (vh1a, vh1 b, vh2, vh3, vh4, vh5, vh6). The vk1-vh1b human germline antibody showed 0.80 4D similarity of its flexible amino-acids compared to the flexible amino-acids of the murine F151 antibody; the vk1-vh1b germline antibody was therefore used to humanize F151 antibody focusing on the flexible amino-acids. For the pair wise amino-acid association between murine F151 vk1-vh1b amino-acids, the 2 sequences were aligned based on the optimal 3D superposition of the alpha carbons of the 2 corresponding homology models (see Figure 15 for an alignment of F151 LC and F151 HC with vk1 and vh1b, respectively).

2. STABILIZATION

[0180] Two approaches were used to improve the stability of the antibody.

a) KNOWLEDGE-BASED APPROACH

[0181] The amino acids of the light and heavy chains with low frequency of occurrence vs. their respective canonical sequences, excluding the CDRs, were proposed to be mutated into the most frequently found amino acids ($\Delta\Delta G_{th} > 0.5$ kcal/mol; E. Monsellier, H. Bedouelle. J. Mol. Biol. 362, 2006, p. 580-593). This first list of consensus mutations for the light chain (LC) and heavy chain (HC) was restricted to the amino acids found in the closest human germline (vk1-vh1b). Suggested changes in the immediate vicinity of the CDRs (5 Angstroms "Vernier" zone, J. Mol. Biol. 224, 1992, p. 487-499) were removed from consideration. This resulted in 5 stabilizing mutations in the LC (see Table 19) and 4 stabilizing mutations in the HC (see Table 20). Other criteria were taken into account to consider these mutations for potentially stabilizing the anti-DAKD/KD F151 antibody. These criteria were a favorable change of hydropathy at the surface or a molecular mechanics based predicted stabilization of the mutant. Also, additional stabilizing mutations reported to be successful in the literature (E. Monsellier & H. Bedouelle, J. Mol. Biol., 362, 2006, p. 580-593; B.J. Steipe et al. J. Mol. Biol, 1994, 240, 188-192) were considered (see Tables 16-22). One of these changes was incorporated as a stabilizing mutation (D89E) in sequences HC2a, HC2b and HC2c below. Another suggested change (Q62E) was incorporated in variant HC2b.

b) 3D AND MD-BASED APPROACHES

[0182] 3D and MD- based approaches have been previously reported (Seco J, Luque FJ, Barril X., J Med Chem. 2009 Apr 23;52(8):2363-71; Malin Jonsson et al., J. Phys. Chem. B 2003, 107, 5511-5518). Hydrophobic regions of the antibody were explicitly identified by analyzing the molecular dynamics simulation of the Fab in a binary solvent (20% isopropanol in water, 20 ns production simulation). Lysine mutations were then introduced in the vicinity of these regions as an attempt to prevent the aggregation. Additional analysis using a hydrophobic surface map within Schrodinger's maestro software (v. 8.5.207) was completed. Using a combination of these two techniques, 2 Lys mutations, 1 in the heavy chain and 1 in the light chain, are suggested.

3. HUMANIZATION BY GRAFTING

[0183] Humanization using grafting grafting techniques has previously been reported (Peter T. Jones, Paul H. Dear, Jefferson Foote, Michael S. Neuberger & Greg Winter Nature, 1986, 321, 522-525). The humanization process which was used started by identifying the closest human germlines to anti-DAKD/KD light and heavy chains. This is done by performing a BLAST search vs. all the human germlines which were systematically enumerated (all possible combinations of the V & J domains for the kappa and lambda chains; V, D and J domains for the heavy chains).

[0184] The following closest human germlines were identified with 83% and 62% sequence identity to anti-DAKD/KD F151 light chains (LC) and heavy chains (HC), respectively (see Figure 16). Using the internal VBASE germline, the light chain is found to be close to V IV-B3 (-83% identity) locus and the heavy chain close to 1-08 & 1-18 (-62% identity) locus of the VH1 subfamily. CDR regions (as defined by MOE), and Vernier regions (as defined in Foote & Winter, J. Mol. Biol., 1992, 224, 487-499) are indicated in boldface The humanizing mutations in underlining were obtained by performing a pairwise comparison of the 2 aligned sequences, excluding the CDR & Vernier zone residues as defined above. In another variant of the humanization, only the CDRs were excluded in the comparison.

4. MUTATION OF UNWANTED SEQUENCE MOTIFS

[0185] The following motifs of sequences were considered: Asp-Pro (acid labile bond), Asn-X-Ser/Thr (glycosylation, X=any amino-acid but Pro), Asp-Gly/Ser/Thr (succinimide/iso-asp formation in flexible regions), Asn-Gly/His/Ser/Ala/Cys (exposed deamidation sites), and Met (oxidation in exposed areas). Among other criteria, the VL & VH domains of murine F151 was selected from other murine antibodies because murine F151 did not have exposed unwanted sequence motifs, but they are introduced in some humanized variants.

[0186] LC3a, LC3b, HC3a and HC3b each have potentially problematic succinimide sites that were identified. These sites were not modified in the proposed sequences as the residues involved are potentially involved in H-bond network (visual inspection of the homology model). These positions are also found in a number of other antibody structures. Additionally, in both HC3a and HC3b, a

strict humanization by grafting would include a substitution of Ser115 to Met. This Methionine is exposed. A substitution to Leucine at this position is suggested as a humanizing mutation as it is a common residue among many close human germline sequences.

[0187] The resulting humanized sequences were blasted for sequence similarity against the International Epitope Database (IEDB) database (found on the world wide web at immuneepitope.com; version June 2009; Vita R, Zarebski L, Greenbaum JA, Emami H, Hoof I, Salimi N, Damle R, Sette A, Peters B. The immune epitope database 2.0. Nucleic Acids Res. 2010 Jan;38(Database issue):D854-62. Epub 2009 Nov 11) to ensure that none of the sequences contain any known human B- or T-cell epitopes (sequence identity of 70% used as cut-off for the results obtained through BLAST search and considering only the results from human species).

5. ORIGINAL SEQUENCES OF MURINE F151 VARIABLE DOMAINS

[0188] CDRs are highlighted in bold and Vernier regions (as defined in Foote & Winter, J. Mol. Biol., 1992, 224, 487-499) are underlined.

Light Chain (SEQ ID NO:26)

DIVMSQSPSS	LAVSVGEKVTMSC KSSQSLLYSSNQKNYLA
WYQQKPGQSP	KPLIY WASTRES GVDPDRFTGSGSGTDFTLT
ISSVKAEDLA	IYYC QQYYSPWT FGGGTKLEIK

Germinality index = 83% with Z46615_1_V_X67858_1_J [V IV-B3]

Heavy Chain (SEQ ID NO:19) :

EQQLQQSGPELVKPGTSVKVSCKAS GYSFTD YNIYWVKQS
HGKSLEWIGY FDPYNGNTGYNQKFRG KATLTVDKSSSTAF
MHLSSLTSDDSAVYYC ANYRYDDHAMDY WGQGTSTVSS

Germinality index = 62% with Z12316_1_VX97051_4_D_X97051_5_J [VH1 1-18]

6. ENGINEERED SEQUENCES

a) Background

[0189] 5 versions for the light chain (LC1, LC2a, LC2b, LC3a, and LC3b) and 5 versions of the heavy chain (HC1, HC2a, HC2b, HC3a, and HC3b) were proposed.

[0190] LC1 contains 5 humanizing mutations identified using the 4D humanization protocol. LC2a introduced an additional 5 stabilizing mutations. LC2b added 1 Lysine mutations to help prevent aggregation. LC3a contains 15 mutations derived from grafting to the closest human germline sequence and retaining the murine CDR and Vernier zone residues. LC3b contained 16 mutations derived from CDR-grafting with one additional humanizing mutation.

[0191] HC1 has 6 humanizing mutations identified by the in-house protocol. HC2a introduced 5 additional stabilizing mutations while HC2b contains 6 additional stabilizing mutations as compared to HC1. HC2c contains 1 Lys mutation, in addition to the stabilizing mutations of HC2a, to help prevent aggregation. HC3a contains 19 mutations derived from grafting to the closest human germline sequence and retaining the murine CDR and Vernier zone residues. HC3b contains 25 mutations derived from CDR grafting.

[0192] 6 combinations in total were proposed (summarized in Table 16):

- LC1 x HC1 (mutations addressing humanization only)
- LC2a x HC2a (mutations addressing humanization and stabilization)
- LC2a x HC2b (mutations addressing humanization and stabilization)
- LC2b x HC2c (mutations addressing humanization, stabilization and "anti-aggregation")
- LC3a x HC3a (mutations addressing mostly humanization by grafting + Vernier)
- LC3b x HC3b (mutations addressing humanization by grafting)

Table 16: Summary of the 6 LCxHC combinations proposed

	(LC1) Humanizing	(LC2a) Humanizing + stabilizing	(LC2b) Humanizing + stabilizing + anti-aggregation	(LC3a) Grafting With Vernier Regions	(LC3b) Grafting
(HC1) Humanizing	X				
(HC2a) Humanizing + stabilizing		X			
(HC2b) Humanizing + stabilizing		X			
(HC2c) Humanizing + stabilizing + "anti-aggregation"			X		
(HC3a) grafting				X	
(HC3b) grafting					X

Table 17: Mutations of the 5 LC variants of the anti-DAKD/KD F151 antibody

Light Chain Sequential numbering	Light Chain Kabat Numbering	(LC1) Humanizing mutations	(LC2a) Humanizing + stabilizing mutations	(LC2b) Humanizing + stabilizing mutations + anti-aggregation mutations	(LC3a) Grafting CDRs + Vernier residues	(LC3b) Grafting CDRs only
Ser5	Ser5		Thr	Thr	Thr	Thr
Ser9	Ser9				Asp	Asp
Ala12	Ala12		Ser	Ser		
Val13	Val13	Ala	Ala	Ala		
Val15	Val15				Leu	Leu
Glu17	Glu17	Asp	Asp	Asp		
Lys18	Lys18	Arg	Arg	Arg	Arg	Arg
Val19	Val19				Ala	Ala
Met21	Met21		Ile	Ile	Ile	Ile
Ser22	Ser22				Asn	Asn
Gln48	Gln42	Lys	Lys	Lys		
Ser49	Ser43				Pro	Pro
Pro52	Pro46					Leu
Thr69	Thr63		Ser	Ser	Ser	Ser
Val84	Val78				Leu	Leu
Lys85	Lys79	Gln	Gln	Gln	Gln	Gln
Leu89	Leu83			Lys	Val	Val
Ile91	Ile85		Thr	Thr	Val	Val
Gly106	Gly100				Gln	Gln
Leu110	Leu104				Val	Val
Mutations:		5	10	11	15	16

Table 18: Mutations of the 6 HC variants of the anti-DAKD/KD F151 antibody

Heavy Chain Sequential numbering	Heavy Chain Kabat numbering	(HC1) Humanizing mutations	(HC2a) Humanizing +stabilizing mutations	(HC2b) Humanizing stabilizing mutations	(HC2c) humanizing +stabilizing mutations + antiaggregation mutations	(HC3a) Grafting CDRs + Vernier residue	(HC3b) Grafting CDRs only
Glu1	Glu1		Gln	Gln	Gln	Gln	Gln
Ile2	Ile2						Val
Gln5	Gln5	Val	Val	Val	Val	Val	Val
Pro9	Pro9		Ala	Ala	Ala	Ala	Ala

Leu11	Leu11	Val	Val	Val	Val	Val	Val
Val12	Val12	Lys	Lys	Lys	Lys	Lys	Lys
Thr16	Thr16	Ala	Ala	Ala	Ala	Ala	Ala
Lys38	Lys38					Arg	Arg
Ser40	Ser40					Ala	Ala
His41	His41	Pro	Pro	Pro	Pro	Pro	Pro
Lys43	Lys43					Gln	Gln
Ser44	Ser44		Gly	Gly	Gly	Gly	Gly
Ile48	Ile48						Met
Gln62	Gln61			Glu			
Lys67	Lys66					Arg	Arg
Ala68	Ala67						Val
Leu70	Leu69						Met
Val72	Val71						Thr
Lys74	Lys73						Thr
Ser76	Ser75					Thr	Thr

Phe80	Phe79		Tyr	Tyr	Tyr	Tyr	Tyr
His82	His81					Glu	Glu
Ser84	Ser82A					Arg	Arg
Leu 86	Leu82C				Lys		
Thr87	Thr83					Arg	Arg
Asp89	Asp85	Glu	Glu	Glu	Glu		
Asp90	Asp86		Glu	Glu	Glu		
Ser91	Ser87					Thr	Thr
Ser115	Ser108					Leu	Leu
Mutations:		6	11	12	12	19	25

a) Engineered light chain sequences:

[0193] No potentially problematic known T-cell or B-cell epitopes were found in all the variants proposed.

LC1 (SEQ ID NO:27), humanizing mutations are underlined, CDRs and vernier zones are in bold:

DIVMTQSPSSSLAASVGDRVTMSCKSSQSLLYSSNQNLYLA
 WYQQKPGKSPKPLIYWASTRESGVPDRFSGSGSTDFTLT
 ISSVQAEDLAIIYYCQQYYSPWTFGGGKLEIK

LC2a (SEQ ID NO:28), humanizing mutations are underlined, CDRs and vernier zones are in bold, stabilization mutations are in italics (T at position 5, S at position 12, I at position 21, S at position 69, T at position 91 shown below) :

DIVMTQSPSSSLAASVGDRVTISCKSSQSLLYSSNQNLYLA
 WYQQKPGKSPKPLIYWASTRESGVPDRFSGSGSTDFTLT
 ISSVQAEDLAIIYYCQQYYSPWTFGGGKLEIK

LC2b (SEQ ID NO:29) humanizing mutations are underlined, CDRs and vernier zones are in bold, stabilization mutations are in italics (T at position 5, S at position 12, I at position 21, S at position 69, T at position 91 shown below) and an anti-aggregation mutation is K at position 89:

DIVMTQSPSSSLAASVGDRVTISCKSSQSLLYSSNQNLYLA
 WYQQKPGKSPKPLIYWASTRESGVPDRFSGSGSTDFTLT
 ISSVQAEDKAIIYYCQQYYSPWTFGGGKLEIK

LC3a (SEQ ID NO:30), grafted mutations shown in underline and CDRs and vernier zones shown in bold:

DIVMTQSPDLSAVSLGERATINCKSSQSLLYSSNQNLYLA
 WYQQKPGQPPKPLIYWASTRESGVPDRFSGSGSTDFTLT
 ISSLQAEDVAIIYYCQQYYSPWTFGGGKLEIK

LC3b (SEQ ID NO:31), grafted mutations shown in underline and CDRs and vernier zones shown in bold:

DIVMTQSPDLSAVSLGERATINCKSSQSLLYSSNQNLYLA
 WYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSTDFTLT
 ISSLQAEDVAIIYYCQQYYSPWTFGGGKLEIK

[0194] Note that L at position 52 is a vernier residue that is mutated to human.

c) Engineered Heavy chain sequences

[0195]

HC1 (SEQ ID NO:20), humanizing mutations are underlined, CDRs and vernier zones are in bold:

EIQLVQSGPEVKKPGASVKVSCKASGYSFTDYNIIYWKQS
 PGKSLIEWIGYFDPYNGNTGYNQKFRGKATLTVDKSSSTAF
 MHLSSLTSED~~SA~~VYYCANYRYDDHAMDYWGQGTSTVTVSS

HC2a (SEQ ID NO:21), humanizing mutations are underlined, CDRs and vernier zones are in bold, stabilization mutations are in italics (Q at position 1, A at position 9, G at position 44, Y at position 80 and E at position 90 shown below):

QIQLVQSGAEVKKPGASVKVSCKASGYSFTDYNIIYWKQS
 PGKGLEWIGYFDPYNGNTGYNQKFRGKATLTVDKSSSTAY
 MHLSSLTSE~~ES~~VYYCANYRYDDHAMDYWGQGTSTVTVSS

HC2b (SEQ ID NO:22), humanizing mutations are underlined, CDRs and vernier zones are in bold, stabilization mutations are in italics (Q at position 1, A at position 9, G at position 44, E at position 62, Y at position 80 and E at position 90 shown below):

QIQLVQSGAEVKKPGASVKVSCCKASGYSFTDYNLYWVKQS
 PGKGLEWIGYFDPYNGNTGYNQKFRGKATLTVDKSSSTAY
 MHLSSLTSEESAVYYCANYRYDDHAMDYWGQGTSTVTVSS

No human epitopes were identified for sequence HC2b in IEDB database.

HC2c (SEQ ID NO:23), humanizing mutations are underlined, CDRs and vernier zones are in bold, stabilization mutations are in italics (Q at position 1, A at position 9, G at position 44, Y at position 80 and E at position 90 shown below) and an anti-aggregation mutation at K at position 86:

QIQLVQSGAEVKKPGASVKVSCCKASGYSFTDYNLYWVKQS
 PGKGLEWIGYFDPYNGNTGYNQKFRGKATLTVDKSSSTAY
 MHLSSLTSEESAVYYCANYRYDDHAMDYWGQGTSTVTVSS

HC3a (SEQ ID NO:24), grafted mutations shown in underline and CDRs and vernier zones shown in bold:

QIQLVQSGAEVKKPGASVKVSCCKASGYSFTDYNLYWVRQA
 PGQGLEWIGYFDPYNGNTGYNQKFRGRATLTVDKSTSTAY
 MELRSLRSDDTAVYYCANYRYDDHAMDYWGQGTSTVTVSS

LC3b (SEQ ID NO:25), grafted mutations shown in underline and CDRs and vernier zones shown in bold:

QVQLVQSGAEVKKPGASVKVSCCKASGYSFTDYNLYWVRQA
 PGQGLEWIGYFDPYNGNTGYNQKFRGRVTMTDTSTSTAY MELRSLRSDDTAVYYCANYRYDDHAMDYWGQGTSTVTVSS

[0196] Note that the following Vernier Residue are mutated to human: V at position 2, M at position 48, V at position 68, M at position 70 and T at position 74.

[0197] No human epitopes were identified for sequence HC3b in IEDB database.

HC3b germinality index = 83% with Z12316_1_V_J00235_1_D_U42590_1_J [1-18/DP-14].

Table 19: Stabilizing Changes Proposed in Light Chain

Residue	Proposed Change	Calculated Gth	Accept Change
Ser-5	Thr	2.32286	Yes
Ala-12	Ser	0.75228	Yes
Met-21	Ile	0.768959	Yes
Pro-52	Leu	1.70059	No - Vernier region
Thr-69	Ser	1.10843	Yes
Lys-86	Glu	2.00115	No - changed to Gln during humanization
Ile-91	Thr	1.27255	Yes

Table 20: Stabilizing Changes Proposed in Heavy Chain

Residue	Proposed Change	Calculated Gth	Accept Change
Glu-1	Gln	0.562423	Yes
Ile-2	Val	2.15882	No - Vernier region
Pro-9	Ala	0.505324	Yes
Thr-16	Ala	1.50552	Already changed to Ala in humanization

Residue	Proposed Change	Calculated Gth	Accept Change
Val-20	Leu	2.21586	No - not in germline sequence
Ser-40	Arg	1.03643	No - not in germline sequence
His-41	Pro	1.67738	Already changed to Pro in humanization
Ser-44	Gly	1.5068	Yes
Gln-62	Glu	0.74934	No - not in germline sequence
Arg-65	Lys	2.32314	No - not in germline sequence
Phe-80	Tyr	1.30935	Yes
His-82	Gln	2.24674	No - not in germline sequence
Asp-89	Glu	1.65409	Already changed to Glu in humanization
Asn-98	Arg	3.65643	No - Vernier region

Table 21: Combinations of stabilizing mutations evaluated

Combination*	Additional changes suggested	Accept Change
L1 (46->P & 48->Q)	K48->Q	No - K48 humanizing mutation
L2 (51->K)	None - already K51	None
L3 (80->T)	None - already T80	None
L4 (82->S)	None - already S82	None
L5 (90->A, 91->T)	None - already A90, T91 suggested above (Table 1)	None
H1 (15->G)	None - already G15	None
H2 (62->E, 63->K, 64->F)	Q62->E, already K63 and F64	Yes - considered in HC2b
H3 (87->T, 88->S, 89->D)	D89->E, already T87 and S88	Yes - potential salt bridges with K63 and K43
S1 (L1 & L5)	K48->Q	No - K48 humanizing mutation
S2 (H1 & H3)	D89->E	No (see H3)

*Note: Sequential numbering used to refer to residues

Table 22: Potential Stabilizing Mutations

Light Chain Residue*	Additional changes suggested	Accept Change
15->L	V15->L	No - V15 in Vk1 germline
96->Q	None - already Q96	None
38->Y	None - already Y38	None
112->I	None - already I112	None
69->S	G69->S	No - G69 is in Vernier Region
21->I	M21->I	Already changed (see Table

Light Chain Residue*	Additional changes suggested	Accept Change
		19)
*Note: Sequential numbering used to refer to residues		

Example 6: Characterization of Humanization Variants

[0198] Based on the in silico modeling presented in Table 16, the variable region of the light chain (VL) and heavy chain (VH) DNA of humanized F151 were codon optimized for HEK293 expression and gene synthesized by GeneArt (subsidiary of Life Technologies). The synthesized DNA fragments were cloned into the constant region of the light chain (CL) encoding vectors, pFF0362 (A. Human Kappa LC vector) at ApaLI/BsiWI sites and the constant regions of the heavy chain (CH1, CH2 and CH3) encoding vectors, pFF0363 (B. Human IgG1 HC vector) at ApaLI/ApaI sites respectively. The resulted plasmids pFF0460 containing the full sequence of LC and pFF0466 containing the full length of HC of humanized F151 variants were co-transfected and transiently expressed in FreeStyle™ 293 Expression System (Invitrogen/Life Technologies, catalog no. K9000-01).

[0199] The six humanized variants shown in Table 16 were characterized by various parameters such as binding kinetics (discussed above) as well as chemical and physical properties such as thermostability that are routinely used in the art.

[0200] The characterization was done in two tiers. Tier I included differential scanning calorimetry (DSC) shown in Table 24 and Figure 2. Briefly, for the DCS experiments, the antibodies were dialyzed against phosphate-buffered saline solution. Antibody concentrations were measured by UV absorbance. The antibodies were diluted to 1 mg/mL using PBS. Scans were performed using a Calorimetry Sciences Corporation N-DSC II instrument using a 0.3268 mL capillary cell with PBS in the reference cell. The scan rate was 2°C/min and the samples were scanned from 20°C to 100°C.

[0201] All variants, except for HC3b/LC3b showed comparable binding affinities to the parental antibody. Variant HC3a/LC3a was selected over the other variants based on other physiochemical properties such as SEC data, stability and lack of aggregation (see Tables 23-25).

Table 23. Comparison of Kinetics of the Humanized F151 Variants

	HC1/LC1			HC2a/LC2a		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DAKD-b	4.16E+05	6.00E-06	1.45E-11	6.66E+05	1.22E-05	1.83E-11
KD-b	4.24E+05	1.74E-07	3.94E-13	7.03E+05	6.12E-06	8.71E-12
DAKLP-b	5.00E+05	7.96E-06	1.60E-11	4.10E+05	5.67E-06	1.38E-11
KLP-b	4.81E+05	2.67E-06	5.54E-12	6.15E+05	2.68E-05	4.34E-11

	HC2b/LC2a			HC2c/LC2b		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DAKD-b	4.17E+05	1.05E-05	2.57E-11	4.81E+05	4.34E-05	9.01E-11
KD-b	3.75E+05	1.66E-06	4.72E-12	5.64E+05	9.08E-06	1.74E-11
DAKLP-b	4.46E+05	1.30E-05	2.97E-11	9.03E+05	1.10E-05	1.21E-11
KLP-b	4.01E+05	2.20E-06	5.76E-12	5.16E+05	1.02E-05	1.98E-11
	HC3a/LC3a			HC3b/LC3b		
	Ka(1/Ms)	Kd(1/s)	KD(M)	Ka(1/Ms)	Kd(1/s)	KD(M)
DAKD-b	5.06E+05	1.28E-05	2.53E-11	3.85E+05	5.15E-05	1.35E-10
KD-b	4.27E+05	2.95E-06	6.78E-12	2.51E+05	3.02E-06	1.44E-11
DAKLP-b	4.65E+05	1.42E-05	3.05E-11	7.04E+04	2.76E-03	4.05E-08
KLP-b	5.02E+05	5.43E-07	1.06E-12	5.39E+05	2.72E-04	5.26E-10
For comparison: Ka(1/Ms) of mF151 was 7.84E+05 for DAKD-b, 8.30E+05 for KD-b, 1.81E+06 for DAKLP-b, and 1.12E+06 for KLP-b						

Table 24. Tier 1 comparison of humanization variants

Variant	Test	Protein Conc. (mg/ml)	Purity		Stability	Functional Potency Assay		Ligand Affinity
	Method	UV Spectroscopy (A280)	1D-gel (reducing & non-reducing)	SEC	DSC (Tm °C) (Tm of parental F151=73°C)	FDSS Assay	FDSS Assay without preincubation	Biacore Binding
HC1/LC1		1.69	No Ag/Deg	≈5% Ag	81.6(M) 70.0(m)	nM potency at DAKD, sub nM potency at KD, comparable among 5 variants	Active at DAKD and KD comparable among 5 variants	Kon 10E5, koffless than or equal to 10E-6 comparable among 5 variants
HC2a/LC2a		2.01	No Ag/Deg	No Ag	75.0(M) 82.5(m)			
HC2b/LC2a		1.83	No Ag/Deg	No Ag	74.6(M) 83.0(m)			
HC2c/LC2b		0.68	No Ag/Deg	No Ag	71.0(M) 64.0(m) 82.5(m) unstable			
HC3a/LC3a		1.71	No Ag/Deg	No Ag	82.3(M) 71.8(m) Most Stable			
HC3b/LC3b		1.47	No Ag/Deg	No Ag	79.4(M) 71.4(m)	Lost potency to KLP & DAKLP		Decreased affinity to KLP & DAKLP

Abbreviations: Ag=aggregation, Deg=degradation

Table 25. Tier 2 comparison of humanization variants

	Thermostability			Stability	Intactness Confirmation (LC, HC)	N-terminal sequence confirmation
Variant	1D-gel	SEC	Biacore	SEC	LC-MS	N-terminal sequencing
HC1/ LC1	No Ag/Deg	No Ag/Deg	45C slightly faster off rate than 4C	No Ag/Deg	LC(+1 Da off) HC(+1 Da off) G0 dominant	N terminal of LC & HC intact
HC2a/ LC2a	No Ag/Deg	No Ag/Deg	45C slightly faster off rate than 4C	No Ag/Deg	LC(spot on) HC(spot on) G0 dominant	N terminal of LC & HC intact
HC2b/ LC2a	No Ag/Deg	No Ag/Deg	45C slightly faster off rate than 4C	No Ag/Deg	LC(+1 Da off) HC(+1 Da off) G0 dominant	N terminal of LC & HC intact
HC2c/ LC2b	X	X	X	X	X	X
HC3a/ LC3a	No Ag/Deg	No Ag/Deg	45C slightly faster off rate than 4C	No Ag/Deg	LC(-2Da off) HC(-2Da off) G0 dominant	N terminal of LC & HC intact
HC3b/ LC3b	X	X	X	X	X	X
Thermostability = Incubation at 4°C (control) and 45°C for 3 days; 1D-gel was under nonreducing conditions; Stability = 2 cycles of freeze/thaw; LC-MS = reduced and reduced/deg lcosylation Abbreviations: Ag=aggregation, Deg=degradation, X = no data presented						

Table 26. Comparison of Parental F151 and humanized variant F151 (HC3a/LC3a)

Parental F151		
	Variable Heavy Chain	Variable Light Chain
Gene	GAGATCCAGCTGCAGCAGTCTGGACCTGAGCTG GTGAAGCCTGGGACITCAGTGAAGGTGTCTGC AAGGCTTCTGGTTACTCAITCACTGACTACAAC ATCTACTGGGTGAAACAGAGCCATGGAAGAGC CTTGAGTGGATTGGATATTTGATCCTTACAAT GGTAATACTGGCTACAACCAGAAAGTTCAGGGGC AAGGCCACATTGACTGTTGACAAGTCCCTCCAGC ACAGCCITCATGCATCTCAGCAGCCTGACATCT GATGACTCTGCAGTCTATTACTGTGCAAACTAC TATAGGTATGACGACCATGCTATGGACTATTGG GGTCAAGGAACCTCAGTCACCGTCTCCTCA (SEQ ID NO:127)	GACATTGTGATGTCACAGTCTCCATCCT CCCTAGCTGTGTCAGTTGGAGAGAAGGI TACTATGAGCTGCAAGTCCAGTCAGAGC CTTTATATAGTAGCAATCAAAAGAACT ACTTGGCCTGGTACCAGCAGAAACCAGG GCAGTCTCCTAAACCGCTGATTTACTGG GCATCCACTAGGGAATCTGGGGTCCCTG ATCGTTTCACAGGCAGTGGATCTGGGAC AGATTTCACTCTCACCATCAGCAGTGTG AAGGCTGAAGACCTGGCAATTTATTACT GTCAGCAATATTATAGCIATCCGTGGAC GTTCGGTGGAGGCAACCAAGCTGGAAATC AAA (SEQ ID NO:128)
	EIQLQQSGPELVKPGTSVKVSCKASGYFETDYN IYVVKQSHGKSLEWIGYFDPYNGNTGYNQKFRG KATLTIVDKSSSTAFMHLSSLTSDSAVYYCANY YVDYDIAKDVVSCGCTGTTGCG	DIVMSQSPSSLAIVSVGEKVTMSCKSSQSL LYSSNQKNYLAWYQQKPGQSPKPLIYWA STRESGVPRFTGSGSGTDFTLTISIVK RDDIATYVGGQWVGRFTSGSGTKELEIK

Protein	<u>IK</u> <u>IDDHAMD</u> <u>YWGQ</u> <u>GISV</u> <u>IVSS</u> (SEQ ID NO:26)	<u>AED</u> <u>LA</u> <u>TI</u> <u>TC</u> <u>QQ</u> <u>YIS</u> <u>IPW</u> <u>IFG</u> <u>Q</u> <u>TK</u> <u>LE</u> <u>IK</u> (SEQ ID NO:19)
---------	--	--

Humanized F151 (HC3a/LC3a)		
Gene	CAGATTTCAGCTGGTGCTGAGTCTGGCGCCGAAGTG AAGAAACCTGGCGCCAGCGTGAAGGTGTCTTGC AAGGCCAGCGGCTACAGCTTCACCGACTACAAC ATCTACTGGGTCCGACAGGCTCCAGGCCAGGGA CTGGAATGGATCGGCTACTTCGACCCCTACAAC GGCAACACCGGCTACAACCAGAAAGTTCGGGGC AGAGCCACCCTGACCGTGEACAAGAGCACCAGC ACCGCCTACATGGAAGTGGGAGCCTGAGAAGC GACGACACCGCGGTGTACTACTGCGCCAACACTAC TACAGATACGACGACCACGCCATGGACTACTGG GGCCAGGACACCCTGGTCACCGTGTCTCT (SEQ ID NO:129)	GACATCGTGATGACCCAGAGCCCCGACA GCCTGGCCGTGTCTCTGGGCGAGCGGGC CACCATCAACTGCAAGAGCAGCCAGAGC CTGCTGTACTCTAGCAACCAGAAGAAGT ACCTGGCCTGGTATCAGCAGAAGCCCCGG CCAGCCCCCAAGCCCCCTGATCTACTGG GCCAGCACCCGCGAGAGCGGCGTGCCCG ATAGATTTTCCGGCAGCGGCTCCGGCAC CGACTTCACCCCTGACCAATCAGCAGCCTG CAGGCCGAGGACGTGGCCGTGTACTACT GCCAGCAGTACTACAGCTACCCCTGGAC CTTCGGCCAGGGCACCAAGGTGGAAATC AAG (SEQ ID NO:130)
Protein	<u>Q</u> <u>I</u> <u>Q</u> <u>L</u> <u>V</u> <u>Q</u> <u>S</u> <u>G</u> <u>A</u> <u>E</u> <u>V</u> <u>K</u> <u>K</u> <u>P</u> <u>G</u> <u>A</u> <u>S</u> <u>V</u> <u>K</u> <u>V</u> <u>S</u> <u>C</u> <u>K</u> <u>A</u> <u>S</u> <u>G</u> <u>Y</u> <u>S</u> <u>F</u> <u>T</u> <u>D</u> <u>Y</u> <u>N</u> <u>I</u> <u>Y</u> <u>W</u> <u>V</u> <u>R</u> <u>Q</u> <u>A</u> <u>P</u> <u>G</u> <u>Q</u> <u>G</u> <u>L</u> <u>E</u> <u>W</u> <u>I</u> <u>G</u> <u>Y</u> <u>F</u> <u>D</u> <u>P</u> <u>Y</u> <u>N</u> <u>G</u> <u>N</u> <u>T</u> <u>G</u> <u>Y</u> <u>N</u> <u>Q</u> <u>K</u> <u>F</u> <u>R</u> <u>G</u> <u>R</u> <u>A</u> <u>T</u> <u>L</u> <u>T</u> <u>V</u> <u>D</u> <u>K</u> <u>S</u> <u>T</u> <u>S</u> <u>T</u> <u>A</u> <u>Y</u> <u>M</u> <u>E</u> <u>L</u> <u>R</u> <u>S</u> <u>L</u> <u>R</u> <u>S</u> <u>D</u> <u>D</u> <u>T</u> <u>A</u> <u>V</u> <u>Y</u> <u>Y</u> <u>C</u> <u>A</u> <u>N</u> <u>Y</u> <u>Y</u> <u>R</u> <u>Y</u> <u>D</u> <u>D</u> <u>H</u> <u>A</u> <u>M</u> <u>D</u> <u>Y</u> <u>W</u> <u>G</u> <u>Q</u> <u>I</u> <u>L</u> <u>V</u> <u>T</u> <u>V</u> <u>S</u> (SEQ ID NO:24)	<u>D</u> <u>I</u> <u>V</u> <u>M</u> <u>T</u> <u>Q</u> <u>S</u> <u>P</u> <u>D</u> <u>S</u> <u>L</u> <u>A</u> <u>V</u> <u>S</u> <u>L</u> <u>G</u> <u>E</u> <u>R</u> <u>A</u> <u>T</u> <u>I</u> <u>N</u> <u>C</u> <u>K</u> <u>S</u> <u>S</u> <u>Q</u> <u>S</u> <u>L</u> <u>L</u> <u>Y</u> <u>S</u> <u>S</u> <u>N</u> <u>Q</u> <u>K</u> <u>N</u> <u>Y</u> <u>L</u> <u>A</u> <u>W</u> <u>Y</u> <u>Q</u> <u>K</u> <u>P</u> <u>G</u> <u>Q</u> <u>P</u> <u>P</u> <u>K</u> <u>P</u> <u>L</u> <u>I</u> <u>Y</u> <u>W</u> <u>A</u> <u>S</u> <u>T</u> <u>R</u> <u>E</u> <u>S</u> <u>G</u> <u>V</u> <u>P</u> <u>D</u> <u>R</u> <u>F</u> <u>S</u> <u>G</u> <u>S</u> <u>G</u> <u>T</u> <u>D</u> <u>F</u> <u>I</u> <u>L</u> <u>T</u> <u>I</u> <u>S</u> <u>S</u> <u>L</u> <u>Q</u> <u>A</u> <u>E</u> <u>D</u> <u>V</u> <u>A</u> <u>V</u> <u>Y</u> <u>Y</u> <u>C</u> <u>Q</u> <u>Q</u> <u>Y</u> <u>S</u> <u>Y</u> <u>P</u> <u>W</u> <u>I</u> <u>F</u> <u>G</u> <u>Q</u> <u>G</u> <u>T</u> <u>K</u> <u>V</u> <u>E</u> <u>I</u> <u>K</u> (SEQ ID NO:30)
Single underscore = CDR region; double underscore = signature amino acids for identifying CDRs		

[0202] For alignment of light and heavy chains of parental F151 to humanized F151 variant (HC3a/LC3a), see Figure 3.

Example 7: Crystal Structure of Humanized Antibody F151 against BRK1 Ligand Kallidan and des-arg¹⁰-Kallidin

[0203] The crystal structures of humanized F151 (HC3a/LC3a) Fab bound to kallidan or des-arg¹⁰-kallidin was determined and the molecular interactions analyzed.

[0204] Kallidin powder was purchased from Phoenix Pharmaceuticals (Cat. No. 009-37). For Fab protein generation, the DNA of heavy chain (HC) VH region from humanized F151 HC3a was cloned into 6XHis tagged CH1 vector pFF0366. The light chain (LC) plasmid used here was the same as of the original F151 LC3a used in F151 humanization (see Example 5). The two plasmids were co-transfected into free style HEK293 cells for Fab expression. The Fab protein was purified using cobalt-resin, buffer exchanged to 50 mM MES pH6.0, 50mM NaCl before being concentrated to about 9 mg/mL. Purified F151 Fab protein was mixed with kallidin in a molar ratio of 1:2 and set up for crystallization screening. Crystallization screening was done with a wide range of conditions. The best crystal was observed under condition B10, B12 and G10 of Hampton Research screening kit PEG/ION HT. The crystals were cryo-protected with 20% glycerol in well buffer and frozen for diffraction data collection. The X-ray diffraction data for both complexes were collected at Canadian

Light Source, beamline CMCF-08ID. The Rmerge for the F151-KD complex is 8.9% and I/s(I) = 20.2, while those for the F151 -DAKD are 7.7% and 18.5, respectively. The F151-KD structure was solved by molecular replacement in Phaser using Fab coordinates from PDB entry 3QOS, treating the V_L - V_H and C_L - C_H1 domains as independent units. The structure was refined in autoBuster at 2.07 Å resolution in space group $P2_12_12_1$ to an Rfactor of 0.205 and an Rfree of 0.228. The F151 -DAKD structure was solved using the F151-KD coordinates. The structure was refined in autoBuster at 1.86 Å resolution in space group $P212121$ to an Rfactor of 0.232 and an Rfree of 0.238.

[0205] The electron density maps shown in Figures 4 and 5 depict the binding of kallidin (KD) and Des-Arg¹⁰-kallidin (DAKD) to the F151 Fab and unambiguously determine the positions of each amino acid. For kallidin, the electron density for the extreme C-terminal residue Arg¹⁰ is not present. This is in agreement with the observation that DAKD, which is missing the C-terminal residue arginine (shown in Table 27 below), binds equally well to F151 as KD. The IC₅₀ values of F151 in the neutralization FDSS cellular assay towards KD and DAKD are 0.12 nM and 0.09 nM, respectively. In both cases the electron density is weaker towards the C-termini of the peptides. Since Phe⁹ in KD has slightly better electron density than that in DAKD, it is possible that the presence of the additional arginine at the C-terminus of KD stabilizes the C-terminus of this peptide when binding to F151 although this arginine itself is not stable enough to be observed by X-ray. Since the two structures are essentially identical (rms between KD and DAKD is 0.139 for C atoms and 0.328 for all atoms), all the following discussions are based on the F151-KD structure.

Table 27. A selected list of kinin peptides

Peptide name	SEQ ID NO:	Sequence									
		1	2	3	4	5	6	7	8	9	10
KD (Kallidin)	1	Lys	-Arg	-Pro	-Pro	-Gly	-Phe	-Ser	-Pro	-Phe	-Arg
DAKD (des-Arg ¹⁰ -Kallidin)	2	Lys	-Arg	-Pro	-Pro	-Gly	-Phe	-Ser	-Pro	-Phe	
KLP (rodent KD ortholog)	3	Arg	-Arg	-Pro	-Pro	-Gly	-Phe	-Ser	-Pro	-Phe	-Arg
BK (Bradykinin)	5		Arg	-Pro	-Pro	-Gly	-Phe	-Ser	-Pro	-Phe	-Arg

[0206] KD is bound with its N-terminus buried in the interface between Fv subunits of the light and heavy chains, as shown in Figure 6. The interface between light and heavy chains are packed with aromatic amino acids, including Tyr-L42, Tyr-L93, Tyr-L100, Trp_L102, Phe-L104 and Tyr-H35, Trp-H47, Tyr-H50, Tyr-H99, Trp-H110, stabilizing each other through stacking and hydrophobic interactions. Residues from each of the CDR's of light and heavy chains contribute to the binding. The residues along the light and heavy chains that are involved in interactions with KD as mapped on the CDRs are shown in Figures 7 and 8. CDR H3 of the heavy chain is the longest loop and the one most frequently used in the interactions with KD, forming a side cover for KD. The loop was stabilized mostly through interactions with the other two CDRs, H1 and H2 of the heavy chain, namely, Salt bridge between Asp-H101 and Arg-H52 (stabilizing H1 and H3), arene-H interaction between Tyr-H102 and Tyr-H54 (stabilizing H2 and H3), H-bond between Asp-H108 and Tyr-H35 and H-bond between His-H105 and Tyr-L55 (stabilizing H3 and L2).

Comparing the KD interacting residues among the antibodies generated, it can be seen that there is similarity among the antibodies, and some were more related in use of particular amino acids for KD-interaction than others. For example, in the light chain F151, C63 and I22 use more similar amino acids in their CDRs to bind KD, while B21 and I54 were more similar. In the heavy chain, F151 and C63 were surprisingly unique from each other and from B21, I22 and I54. The latter three appear to form a group in similarity. C63 is particularly interesting in its heavy chain, that the loop length in H2 and H3 are more different from all others. Considering the Fab as a whole, B21 and I54 were most closely related.

[0207] In the crystal structure, we found that KD is involved in systematic hydrogen bond and hydrophobic interactions with the Fab. The N-terminus of KD is buried in the Fab and harbors more intensive interactions, while the C-terminus is essentially solvent exposed. Except for the first 4 residues (Lys-Arg-Pro-Pro), the other residues of KD gradually extend into the bulk solvent. The amidinium group of Lys1 sidechain is anchored by Glu-L61 (L: light chain) through salt bridges, while the amino terminal amino group Lys 1 forms a salt bridge with Asp-H108 (H: heavy chain). The amidinium group of Lys1 sidechain also hangs over the aromatic ring of Tyr-L55, involved in cation interactions. Such intensive interactions involved with Lys1 tightly anchor the amino terminus of KD in the Fab. This also accounts for the importance of Lys1 in the binding of KD to F151. Without it (i.e. bradykinin), no detectable binding to hF151 or F151 can be measured. Like Lys1, Arg2 interacts with the Fab through a salt bridge. The guanidium group of Arg2 interacts with the sidechain of Asp-H104. The sidechain of Arg2 is also H-bonded with the mainchain carbonyl oxygen of Arg-H101. Also, the mainchain oxygen of Pro8 is H-bonded to the sidechain of Arg-H101. Tyr-H102 is half-way intercalated into Phe8 and Pro9, involving in hydrophobic interactions with KD. In addition to direct interaction, numerous water-mediated H-bonds between KD and the Fab are also seen. It is also interesting to notice that tyrosine residues are most frequently used in the interaction compared to other amino acids; 9 out of the 16 residues marked with asterisks in Figures 7 and 8 are tyrosines. All the residues in F151 surrounding KD appear to play a role in ligand binding, except for Asn-H33, which is close to Phe6 sidechain but incompatible in polarity and lack of other important interactions. Substitution with aromatic/hydrophobic residues, such Trp or Tyr to interaction with Phe8 appears to be a quick pick if affinity maturation is considered. These two aromatic amino acids are in fact seen in other antibodies (Trp in C63, and Tyr in B21, I22, I54). Table 28 below provides a detailed analysis of 16 KD-interacting amino acid residues marked in Figures 7 and 8 and sets forth functional substitutions that can be made in the CDR regions that should not disrupt antigen binding.

Table 28. A list of amino acid residues found around the KD binding pocket, and their roles in KD binding and potential functional substitutions (light chain residues in grey-colored cells and heavy chain residues in unshaded cells)

Residue	Role in KD Binding or CDR Stabilization	Functional Substitution
Tyr-L31	• Edge-on hydrophobic interactions with Pro4	• His to add an H-bond with amide N of Pro5
	• Along with Tyr-L38 and Tyr-L98 form three orthogonal planes that surround the 90deg turn of KD at Pro4	• other aromatic a.a, such as Trp, and Phe
	• hydrophobic stacking with Pro4	• His to add an H-bond with

Residue	Role in KD Binding or CDR Stabilization	Functional Substitution
Tyr-L38	Along with Tyr-L31 and Tyr-L98 form three orthogonal planes that surround the 90deg turn of KD at Pro4	carbonyl O of Arg2
Tyr-L55	cation- interaction with amidinium ion of Lys1 sidechain	Trp to pair with H105 mutations of Gln, Asn, Glu or Asp (maintaining H-bond); Other variants of His_H105 are Tyr and Ser.
	H-bond with His-H105; Tyr-L55 --His-H105 pairing adds stabilization between L2 and H3 loops	Other aromatic a.a., such as Trp, His and Phe
Glu-L61	Forming key salt bridges with Lys1 sidechain	Asp, Gln, ASN (Asp is seen in B21 and I54 already, Figure 7)
Tyr-L97	H-bond with amide N of Trp-L56	Aromatic a.a. such as Phe or His (too tight space for Trp)
	Forming pocket for Arg1 extended sidechain	
Tyr-L98	Along with Tyr-L31 and Tyr-L38 form three orthogonal planes that surround the 90deg turn of KD at Pro4	Other aromatic a.a, such as Phe, Trp or His
Tyr-L100	Forming pocket surface for Pro3	Aromatic a.a., such as Phe (better hydrophobic interactions with Pro3)
	Part of the aromatic a.a. cluster interface between L/H chains, further including Tyr-L42, Tyr-L93, Trp_L102, Phe-L104 and Tyr-H35, Trp-H47, Tyr-H50, Tyr-H99, Trp-H110	Other variants seen are Thr (in B21 and I54) and His (in I22)
	Partial stacking with Tyr-H50	
Trp-L102	Part of the aromatic a.a. cluster interface between L/H chains	Other aromatic a.a, such as Tyr (in C63 and B21), Phe and His
	Stacking with Trp-H47	Other hydrophobic residue, such as L (in I22)
Asn-H33	Close to Phe6 sidechain but incompatible in polarity, no other roles seen either; can be a target for affinity maturation	Replace with aromatic/hydrophobic a.a., such as Trp (seen in C63) or Tyr (seen in B21, I22, I54)
Asp-H52	Salt bridge with Arg-H101, stabilizing H1 and H3 loops	Mutate as a pair with Arg-H101 to reversely charged a.a., such Arg-H52/Asp-H101, or a pair of hydrophobic a.a. (Leu, Ile, Val,

Residue	Role in KD Binding or CDR Stabilization	Functional Substitution
		Met, Phe, Tyr, Trp, Ala) to form a cluster with Phe6 of KD
Tyr-H54	• Close to Pro8 but no specific interactions	• Mutated to negatively charged a.a. to stabilize Arg-H101, such D or E (seen in B21, I22 and I54) or N or Q, also provide H-bond with carbonyl O of Pro8 (A Lys in C63, which can be charge-reversed to Glu)
	• Close to Arg-H101, but no charge interactions	
Tyr-H99	• Part of aromatic interface between H and L chains	• Mutate to small aromatic a.a. except W, such as Phe and His
	• H-bond with Asn-H33	
	• Tight space	
Arg-H101	• H-bond with amide of Pro8	• Mutate as a pair with Asp-H52 to reversely charged a.a., such Arg-H52/Asp-H101, or a pair of hydrophobic a.a. to form a cluster with Phe6 of KD
	• supported by Asp-H52 (salt bridge)	
Tyr-H102	• Half-way intercalating into Phe9 and Pro8, hydrophobic interactions with KD	• Phe can be better, Trp or His may be OK too
Asp-H104	• Key residue to salt-bridge with Arg2	• Glu to maintain salt bridge with Arg2
		• Bigger a.a to fill the gap from Pro3, such as Tyr as seen in B21, I22 and I54
Asp-H108	• Key residue to salt bridge with N-term -NH ₃ ⁺ of KD	• Glu
	• H-bond with Tyr-H33, stabilizing H3 loop	
	• Conserved residue! Not in CDR	

[0208] Analysis of the conformational epitope of kallidin (KD) or desArg10-Kallidin (DAKD) revealed that it adopts a "Pro4 kink" conformation. As depicted in Figure 17, a hallmark of the "Pro 4 kink" conformation is a type II tight turn in the main chain polypeptide backbone of KD or DAKD at Proline 4 (see Richardson JS. "The anatomy and taxonomy of protein structure." Adv Protein Chem. 1981 ;34:167-339, which is incorporated by reference herein). The "Pro 4 kink" conformation may further defined by all or substantially all of the remaining amino acids of KD (1-2 and 6-9) or DAKD adopting repeats of a sigmoid shape which align the hydrophobic side chains in a spatially stacking mode.

Example 8: In vivo Pharmacology of anti-BKR1-Ligand Antibodies in Pain Models

[0209] The examples of the present disclosure illustrate the in vivo efficacy of anti-BKR1 receptor-ligand antibodies in different preclinical models of acute and chronic pain according to modified procedures described in (a) Saddi GM and Abbott FV., Pain (2000), 89:53-63; (b) Chen et al., Molecular Pain (2010), 2:6-13 and (c) Bennett GJ and Xie YK., Pain (1988), 33:87-107.

Animals

[0210] Experiments were carried out using adult male OF1 mice (20-30 gr) for formalin studies and adult male C57Bl/6J mice (25-30 gr) for both CFA and CCI studies. The mice were kept in a controlled temperature room under a 12-h light-dark cycle. Food and water were provided *ad libitum*. For all of the experiments, mice were acclimatized to the laboratory room for at least 2 hours before testing. No randomization was performed for the studies. Experimenters performing the behavioral tests were not blind to treatment; however they were not aware of the study hypothesis. All procedures have been approved by the "Comité d'Expérimentation pour la Protection de l'Animal de Laboratoire" (Animal Care and Use Committee) of sanofi-aventis recherche & développement and were carried out in accordance with French legislation (Decree n°87-848 - 19 October 1987 - and decision -19 April 1988) implementing European directive 86/609/EEC.

A. Formalin-induced acute inflammatory pain

[0211] The formalin test was used to measure nociceptive and inflammatory pain. Indeed, intraplantar injection of formalin induces an initial acute nociceptive behavioral response (0-12 minutes), followed by a second inflammatory-mediated response (15-45 minutes), which is attributed to spinal cord excitability.

[0212] Formaldehyde (37%, Sigma) was diluted in saline (v/v) to obtain a 2.5% formaldehyde concentration (i.e. $\approx$ 6.25% formalin concentration). Mice were gently restrained and 20 μ L of this solution was injected subcutaneously into the dorsal part of one hind paw. Behavioral responses were scored immediately after formalin injection, then at 3 minutes intervals over 45 minutes as follows: (0): normal weight bearing of the injected paw; (1): injected paw resting lightly on floor; (2): lifting-elevation of the injected paw; (3): licking or biting the injected paw. Group sizes were 11-12 male OF1 mice.

[0213] Scores were plotted versus time and areas under the curves (AUC) were calculated from the mean scores ($\pm$ SEM) for both the early (0-12 min) and the late (15-45 min) phases. Reversal of pain-like behaviors was expressed as change in AUC in %.

[0214] EE1 antibody inhibited the pain-like behavior in the late phase of the formalin test in male OF1 mice. EE1 antibody, when administered intravenously 48 hours before intraplantar injection of formalin, showed a dose dependent reversal of the pain-like behavior only in the late phase with a

Minimal Effective Dose (MED)=2.5 mg/kg, as depicted in Figure 9. Indeed, when administered at 2.5, 10 and 30 mg/kg, EE1 reversed the late phase by $35\pm5\%$, $33\pm5\%$ and $45\pm7\%$, respectively, as depicted in Table 29.

[0215] In contrast, F151 weakly inhibits the pain-like behavior in the late phase of the formalin test when administered 48 hours before intraplantar injection of formalin. Indeed, when administered at 2.5 and 10 mg/kg, F151 reversed the late phase by $15\pm7\%$ and $21\pm5\%$, respectively, as depicted in Table 29.

Table 29. Effect of EE1 and F151 antibodies on formalin-induced pain-like behavior in male OF1 mice

Group	Dose (mg/kg, i.v.)	A.U.C. $\pm$ SEM (15-45 min)	Reversal of pain-like behavior (in %) $\pm$ SEM (15-45 min)
Isotype-control 1B7.11 (EE1)	30	63.6 ± 2.9	0 ± 5
EE1	2.5	41.6 ± 3.4 (***)	35 ± 5
	10	42.9 ± 2.9 (***)	33 ± 5
	30	36.5 ± 4.3 (***)	45 ± 7
Isotype-control 1B7.11 (F151)	10	57.3 ± 3	0 ± 5
F151	2.5	48.9 ± 3.8 (NS)	15 ± 7
	10	45.0 ± 2.8 (*)	21 ± 5
*, $p<0.05$; ***, $p<0.001$: Student's t-test versus adequate control was used.			
NS: non significant			

B. CFA (Complete Freund's adjuvant)-induced chronic inflammatory pain

[0216] Chronic inflammation was induced under brief anesthesia (Isoflurane, 3%) by an intraplantar administration of 25 μ L of Complete Freund's Adjuvant (CFA) containing 1 μ g/ μ L heat-killed Mycobacterium tuberculosis in mineral oil and mannide monooleate (Sigma). Group sizes were 8 male C57Bl/6 mice.

[0217] EE1 antibody was administered intravenously 22 hours after intraplantar CFA injection at 2.5 and 30 mg/kg and mechanical and thermal hypersensitivities were assessed at Day 1 (D1), Day 4 (D4) and Day 7 (D7) post-CFA intraplantar administration.

B1. Mechanical Hypersensitivity

[0218] Mechanical hypersensitivity was assessed by measuring the Frequency of withdrawal Response (FR, in %) following 10 applications of a 0.6 g Von Frey filament (Bioseb, France) onto the plantar surface of the injected paw.

To investigate the efficacy of EE1 antibody on pain-like behavior, we calculated the reversal of mechanical hypersensitivity (in %) as follows:

Percent reversals were calculated as $(\text{Mean FR-isotype-control}_{\text{postdose}} - \text{FR-ipsi}_{\text{postdose}}) / (\text{Mean FR-isotype-control}_{\text{postdose}} - \text{Mean FR-sham}_{\text{postdose}})$ for each mouse.

[0219] At D1, D4 and D7 after intraplantar injection of CFA, a significant increase of FR to the Von Frey filaments was observed in the isotype-control 1 B7.11-treated group in comparison with the naive group, demonstrating the development of mechanical hypersensitivity. EE1 antibody, when administered intravenously 22 hours after intraplantar CFA, was able to significantly decrease this FR at the different times studied compared with that obtained in the isotype-control 1 B7.11-treated group. (Figure 10).

[0220] Reversal of mechanical hypersensitivity was $41 \pm 8\%$ and $22 \pm 8\%$ at D1, $36 \pm 9\%$ and $32 \pm 9\%$ at D4 and $27 \pm 10\%$ and $50 \pm 9\%$ at D7 for a 2.5 mg/kg and 30 mg/kg intravenous administration of EE1 antibody, respectively (Table 30).

Table 30. Effect of EE1 antibody on CFA-induced mechanical hypersensitivity in male C57Bl/6 mice

Group	Dose (mg/kg, i.v.)	Day 1 post-CFA FR (%) % effect		Day 4 post-CFA FR (%) % effect		Day 7 post-CFA FR (%) % effect	
Naive	n.a.	41.3 $\pm$ 4.4	100 $\pm$ 11	43.8 $\pm$ 2.6	100 $\pm$ 8	45 $\pm$ 5	100 $\pm$ 13
Isotype-control 1B7.11	30	81.3 $\pm$ 4.4	0 $\pm$ 11	78.8 $\pm$ 3	0 $\pm$ 8	82.5 $\pm$ 2.5	0 $\pm$ 7
EE1	2.5	65 $\pm$ 3.3 (**)	41 $\pm$ 8	66.3 $\pm$ 3.2 (*)	36 $\pm$ 9	72.5 $\pm$ 3.7 (*)	27 $\pm$ 10
	30	72.5 $\pm$ 3.1 (*)	22 $\pm$ 8	67.5 $\pm$ 3.1 (*)	32 $\pm$ 9	63.8 $\pm$ 3.2 (***)	50 $\pm$ 9
FR: Frequency of Response (in %) $\pm$ SEM, % effect $\pm$ SEM, n.a. not applicable							
*, p<0.05, **, p<0.01 and ***, p<0.001, Two-Way ANOVA with time as repeated measure followed by Dunnett's test for factor group for each level of factor time							

B2. Thermal Hypersensitivity

[0221] For thermal hypersensitivity, measures of Paw Withdrawal Latencies (PWL, in seconds) in response to a radiant heat using a plantar apparatus (IITC, Woodland Hills, USA) were assessed.

[0222] To investigate the efficacy of EE1 antibody on pain-like behavior, we calculated the reversal of thermal hypersensitivity (in %) as follows:

Percent reversals were calculated as $(PWL_{\text{postdose}} - \text{Mean isotype-control}_{\text{postdose}}) / (\text{Mean isotype-control}_{\text{predose}} - \text{Mean isotype-control}_{\text{postdose}})$ for each mouse.

[0223] Thermal hypersensitivities were not different between all groups at baseline, before intraplantar injection of CFA (data not shown).

[0224] At D1, D4 and D7 after intraplantar injection of CFA, a significant decrease in paw withdrawal latency of the injected paw was observed in isotype-control 1B7.11 -treated group of mice, demonstrating that CFA induced a thermal hypersensitivity (data not shown).

[0225] EE1 antibody, administered intravenously 22 hours after intraplantar CFA injection (i.e. on Day 1 post-intraplantar CFA injection), and was not able to increase the Paw Withdrawal Latency at D1, whatever the dose tested (Figure 11). However, EE1 significantly increased the Paw Withdrawal Latency at D4 and this effect was also present at D7 (Figure 11).

[0226] Reversal of thermal hypersensitivity was $41 \pm 15\%$ and $58 \pm 21\%$ at D4 and $46 \pm 10\%$ and $52 \pm 17\%$ at D7 for a 2.5 mg/kg and 30 mg/kg intravenous administration of EE1, respectively (Table 31).

Table 31. Effect of EE1 antibody on CFA-induced thermal hypersensitivity in male C57B1/6 mice

Group	Dose (mg/kg, i.v.)	Day 1 post-CFA PWL (sec) % effect		Day 4 post-CFA PWL (sec) % effect		Day 7 post-CFA PWL (sec) % effect	
Isotype-control 1B7.11	30	3.3±0.3	n.a.	4.1±0.1	n.a.	3.6±0.3	n.a.
EE1	2.5	3.1±0.3	-7±10	5.2±0.4	41±15	5.0±0.3 (*)	46±10
	30	2.6±0.2	-20±5	5.7±0.6 (*)	58±21	5.2±0.6 (*)	52±17
PWL: Paw withdrawal latency ± SEM, % effect ± SEM, n.a. not applicable							
*, p<0.05, Two-Way ANOVA with time as repeated measure followed by Dunnett's test for factor group for each level of factor time							

C. CCI (Chronic Constriction Injury)-induced neuropathic-like pain (Bennett's model)

[0227] CCI model was used as a model of peripheral nerve injury. Briefly, mice were anesthetized with Isoflurane (3%), and the right sciatic nerve was exposed at mid thigh level through a small

incision. Three loose ligatures of 6.0 chromic gut (Ethicon) at 1 mm space were placed around the sciatic nerve. The surgical procedure was completed by closing the muscles and skin. The day of CCI surgery was considered as Day 0. Group sizes were 6-10 male C57Bl/6 mice.

[0228] EE1 antibody was administered intravenously at 2.5 and 30 mg/kg on Day 11 post surgery and mechanical and thermal hypersensitivities were assessed on Day 12 (D12), Day 14 (D14) and Day 18 (D18) post-surgery which corresponded to Day 1 (D1), Day 3 (D3) and Day 7 (D7) posttreatment.

C1. Mechanical Hypersensitivity

[0229] Mechanical hypersensitivity was assessed by measuring hind paw withdrawal thresholds (on both injured [i.e. Ipsi] and non-injured [i.e. Contra] paws) to an increasing pressure (in g) stimulus using a Dynamic Plantar Aesthesiometer (Ugo-Basile, Italy); a steel rod was applied to the hind paws of the mice with an increasing force (5 grams in 10 seconds).

[0230] To investigate the efficacy of EE1 antibody on pain-like behavior, we determined the reversal of mechanical hypersensitivity as follows: percent reversals were calculated as $(\text{Ipsi}_{\text{postdose}} - \text{Ipsi}_{\text{predose}}) / (\text{Contra}_{\text{predose}} - \text{Ipsi}_{\text{predose}})$ for each mouse.

[0231] Following surgery, operated mice developed a robust sensitization to mechanical stimulus on the injured paw, whereas the non-injured paw was not affected. At Day 11, the mechanical sensitization on the injured paw reached a plateau (data not shown).

[0232] EE1 antibody, administered intravenously on Day 11 demonstrated a slight tendency to reverse CCI-induced mechanical hypersensitivity on D12, D14 and D18 with $15.2 \pm 4.9\%$ and $15.2 \pm 5.7\%$ on D12, $26.8 \pm 5.7\%$ and $25.7 \pm 4.5\%$ on D14 and $30.3 \pm 7.1\%$ and $20.8 \pm 5.9\%$ on D18, at 2.5 and 30 mg/kg respectively (Figure 12 and Table 32).

Table 32. Effect of EE1 antibody on CCI-induced mechanical hypersensitivity in male C57Bl/6 mice

Group	Dose (mg/kg, i.v.)	Day 12 post-CCI % effect $\pm$ SEM	Day 14 post-CCI % effect $\pm$ SEM	Day 18 post-CCI % effect $\pm$ SEM
Isotype-control 1B7.11	30	0.2 ± 3.0	1.8 ± 4.3	18.1 ± 6.6
EE1	2.5	15.2 ± 4.9	26.8 ± 5.7 (**)	30.3 ± 7.1
	30	15.2 ± 5.7	25.7 ± 4.5 (*)	20.8 ± 5.9

*, $p < 0.05$, and **, $p < 0.01$ Two-Way ANOVA with time as repeated measure followed by Dunnett's test for factor group for each level of factor time (statistics performed on Delta ipsi values)

C2. Thermal Hypersensitivity

[0233] For thermal hypersensitivity, measures of Paw Withdrawal Latencies (in seconds) in response to a radiant heat using a plantar apparatus (IITC, Woodland Hills, USA) were assessed on the injected hind paw.

[0234] To investigate the efficacy of EE1 antibody on pain-like behavior, we calculated the reversal of thermal hypersensitivity (in %) as follows:

Percent reversals were calculated as $(\text{Ipsi}_{\text{postdose}} - \text{Mean isotype-control}_{\text{postdose}}) / (\text{Mean naive}_{\text{postdose}} - \text{Mean isotype-control}_{\text{postdose}})$ for each mouse.

[0235] Following surgery, operated mice developed a robust sensitization to thermal stimulus on the injured paw, whereas the non-injured paw was not affected. At Day 11, the thermal sensitization on the injured paw reached a plateau (data not shown).

[0236] EE1 antibody, administered intravenously on Day 11 did not significantly increase Paw Withdrawal Latency of the injured paw on D12, even if a trend was observed. However, from D14, EE1 antibody significantly increased the Paw Withdrawal (Figure 13).

Reversal of thermal hypersensitivity was $41 \pm 16\%$ and $56 \pm 24\%$ at D12 and $51 \pm 16\%$ and $98 \pm 48\%$ at D14 and $78 \pm 19\%$ and $84 \pm 22\%$ at D18, for a 2.5 mg/kg and 30 mg/kg intravenous administration of EE1 antibody, respectively (Table 33).

Table 33. Effect of EE1 antibody on CCI-induced thermal hypersensitivity in male C57Bl/6 mice Kinetic evaluation

Group	Dose (mg/kg, i.v.)	Day 12 post-CCI PWL (sec) % effect		Day 14 post-CCI PWL (sec) % effect		Day 18 post-CCI PWL (sec) % effect	
naive	n.a.	6.3±0.5	100±19	6.2±0.4	100±14	6.1±0.5	100±17
Isotype-control 1 B7.11	30	3.8±0.2	0±7	3.4±0.2	0±5	3.4±0.2	0±7
EE1	2.5	4.8±0.4	41±16	4.8±0.5	51±16	5.5±0.5 (*)	78±19
	30	5.2±0.6	56±24	6.1±1.3(**)	98±48	5.7±0.6 (**)	84±22
PWL: Paw withdrawal latency ± SEM, % effect ± SEM, n.a. non applicable							
*, p<0.05, and **,p<0.01 Two-Way ANOVA with time as repeated measure followed by Dunnett's test for factor group for each level of factor time							

SEQUENCE LISTING

[0237]

<110> SANOFI

<120> ANTIBODIES TO BRADYKININ B1 RECEPTOR LIGANDS

<130> 543895 SA9-029PC

<140> New Application

<141> Concurrently Herewith

<150> 61/616,845

<151> 2012-03-28

<150> FR 1350953

<151> 2013-02-04

<160> 137

<170> PatentIn version 3.5

<210> 1

<211> 10

<212> PRT

<213> Homo sapiens

<400> 1

Lys	Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe	Arg
1				5					10

<210> 2

<211> 9

<212> PRT

<213> Homo sapiens

<400> 2

Lys	Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe
1				5				

<210> 3

<211> 10

<212> PRT

<213> Mus musculus

<400> 3

Arg	Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe	Arg
1				5					10

<210> 4

<211> 9

<212> PRT

<213> Mus musculus

<400> 4

Arg Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 5

<211> 9

<212> PRT

<213> Unknown

<220>

<221> source

<223> /note="Description of Unknown: Bradykinin peptide"

<400> 5

Arg Pro Pro Gly Phe Ser Pro Phe Arg
1 5

<210> 6

<211> 8

<212> PRT

<213> Unknown

<220>

<221> source

<223> /note="Description of Unknown: Des-Arg9-Bradykinin peptide"

<400> 6

Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 7

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (1)..(1)

<223> /replace="Phe" or "His"

<220>

<221> misc_feature

<222> (1)..(1)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<220>

<221> VARIANT

<222> (3)..(3)

<223> /replace="Asp" or "Ala" or "Val" or "Leu" or "Ile" or "Met" or "Phe" or "Tyr" or "Trp"

<220>

<221> VARIANT

<222> (4)..(4)

<223> /replace="Phe" or "Trp" or "His"

<220>

<221> misc_feature

<222> (3)..(4)

<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

<220>

<221> VARIANT

<222> (6)..(6)

<223> /replace="Glu" or "Tyr"

<220>

<221> misc_feature

<222> (6)..(6)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<220>

<221> VARIANT

<222> (10)..(10)

<223> /replace="Glu"

<220>

<221> misc_feature

<222> (10)..(10)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<400> 7

Tyr	Tyr	Arg	Tyr	Asp	Asp	His	Ala	Met	Asp	Tyr
1				5					10	

<210> 8

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (3)..(3)

<223> /replace="Arg" or "Ala" or "Val" or "Leu" or "Ile" or "Met" or "Phe" or "Tyr" or "Trp"

<220>

<221> misc_feature

<222> (3)..(3)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<220>

<221> VARIANT

<222> (5)..(5)

<223> /replace="Asp" or "Glu" or "Asn" or "Gln"

<220>

<221> misc_feature

<222> (5)..(5)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<400> 8

Tyr	Phe	Asp	Pro	Tyr	Asn	Gly	Asn	Thr	Gly	Tyr	Asn	Gln	Lys	Phe	Arg
1				5					10					15	

Gly

<210> 9

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (8)..(8)

<223> /replace="Trp" or "Tyr"

<220>

<221> misc_feature

<222> (8)..(8)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<400> 9

Gly	Tyr	Ser	Phe	Thr	Asp	Tyr	Asn	Ile	Tyr
1				5					10

<210> 10
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>
 <221> VARIANT
 <222> (3)..(3)
 <223> /replace="Phe" or "His"

<220>
 <221> VARIANT
 <222> (4)..(4)
 <223> /replace="Phe" or "His" or "Trp"

<220>
 <221> misc_feature
 <222> (3)..(4)
 <223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

<220>
 <221> VARIANT
 <222> (6)..(6)
 <223> /replace="Phe" or "Thr" or "His"

<220>
 <221> misc_feature
 <222> (6)..(6)
 <223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<220>
 <221> VARIANT
 <222> (8)..(8)
 <223> /replace="Tyr" or "Phe" or "His" or "Leu"

<220>
 <221> misc_feature
 <222> (8)..(8)
 <223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<400> 10
 Gln Gln Tyr Tyr Ser Tyr Pro Trp Thr
 1 5

<210> 11
 <211> 6
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>
 <221> VARIANT
 <222> (6)..(6)
 <223> /replace="Asp" or "Gln" or "Asn"

<220>
 <221> misc_feature
 <222> (6)..(6)
 <223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<400> 11
 Trp Ala Ser Thr Arg Glu
 1 5

<210> 12
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>
 <221> VARIANT
 <222> (8)..(8)
 <223> /replace="His" or "Tyr" or "Phe"

<220>
 <221> misc_feature
 <222> (8)..(8)
 <223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<220>
 <221> VARIANT
 <222> (15)..(15)
 <223> /replace="Tyr"

<220>

<221> misc_feature

<222> (15)..(15)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<400> 12

Lys	Ser	Ser	Gln	Ser	Leu	Leu	Trp	Ser	Ser	Asn	Gln	Lys	Asn	His	Leu
1				5					10					15	

Ala

<210> 13

<211> 11

<212> PRT

<213> Mus musculus

<400> 13

Tyr	Tyr	Arg	Tyr	Asp	Asp	His	Ala	Met	Asp	Tyr
1			5						10	

<210> 14

<211> 17

<212> PRT

<213> Mus musculus

<400> 14

Tyr	Phe	Asp	Pro	Tyr	Asn	Gly	Asn	Thr	Gly	Tyr	Asn	Gln	Lys	Phe	Arg
1				5					10					15	

Gly

<210> 15

<211> 10

<212> PRT

<213> Mus musculus

<400> 15

Gly	Tyr	Ser	Phe	Thr	Asp	Tyr	Asn	Ile	Tyr
1			5						10

<210> 16

<211> 9

<212> PRT

<213> Mus musculus

<400> 16

Gln	Gln	Tyr	Tyr	Ser	Tyr	Pro	Trp	Thr
1			5					

<210> 17

<211> 7

<212> PRT

<213> Mus musculus

<400> 17

Trp Ala Ser Thr Arg Glu Ser
 1 5

<210> 18

<211> 17

<212> PRT

<213> Mus musculus

<400> 18

Lys Ser Ser Gln Ser Leu Leu Tyr Ser Ser Asn Gln Lys Asn Tyr Leu
 1 5 10 15

Ala

<210> 19

<211> 120

<212> PRT

<213> Mus musculus

<400> 19

Glu Ile Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Thr
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
 20 25 30

Asn Ile Tyr Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
 35 40 45

Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr Asn Gln Lys Phe
 50 55 60

Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Phe
 65 70 75 80

Met His Leu Ser Ser Leu Thr Ser Asp Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
 100 105 110

Gly Thr Ser Val Thr Val Ser Ser
 115 120

<210> 20

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 20

Glu Ile Gln Leu Val Gln Ser Gly Pro Glu Val Lys Lys Pro Gly Ala

```

1             5             10             15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
      20             25             30

Asn Ile Tyr Trp Val Lys Gln Ser Pro Gly Lys Ser Leu Glu Trp Ile
      35             40             45

Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr Asn Gln Lys Phe
      50             55             60

Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Phe
      65             70             75             80

Met His Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
      85             90             95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
      100            105            110

Gly Thr Ser Val Thr Val Ser Ser
      115            120

<210> 21
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 21
Gln Ile Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1             5             10             15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
      20             25             30

Asn Ile Tyr Trp Val Lys Gln Ser Pro Gly Lys Gly Leu Glu Trp Ile
      35             40             45

Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr Asn Gln Lys Phe
      50             55             60

Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
      65             70             75             80

Met His Leu Ser Ser Leu Thr Ser Glu Glu Ser Ala Val Tyr Tyr Cys
      85             90             95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
      100            105            110

Gly Thr Ser Val Thr Val Ser Ser
      115            120

```

<210> 22

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 22

Gln	Ile	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala
1				5					10					15	

Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Asp	Tyr
			20					25					30		

Asn	Ile	Tyr	Trp	Val	Lys	Gln	Ser	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Ile
		35					40					45			

Gly	Tyr	Phe	Asp	Pro	Tyr	Asn	Gly	Asn	Thr	Gly	Tyr	Asn	Glu	Lys	Phe
	50					55					60				

Arg	Gly	Lys	Ala	Thr	Leu	Thr	Val	Asp	Lys	Ser	Ser	Ser	Thr	Ala	Tyr
65					70					75				80	

Met	His	Leu	Ser	Ser	Leu	Thr	Ser	Glu	Glu	Ser	Ala	Val	Tyr	Tyr	Cys
			85						90					95	

Ala	Asn	Tyr	Tyr	Arg	Tyr	Asp	Asp	His	Ala	Met	Asp	Tyr	Trp	Gly	Gln
			100					105						110	

Gly	Thr	Ser	Val	Thr	Val	Ser	Ser
		115				120	

<210> 23

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 23

Gln	Ile	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala
1				5					10					15	

Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Asp	Tyr
			20					25					30		

Asn	Ile	Tyr	Trp	Val	Lys	Gln	Ser	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Ile
		35					40					45			

Gly	Tyr	Phe	Asp	Pro	Tyr	Asn	Gly	Asn	Thr	Gly	Tyr	Asn	Gln	Lys	Phe
	50					55					60				

Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met His Leu Ser Ser Lys Thr Ser Glu Glu Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Ser Val Thr Val Ser Ser
115 120

<210> 24

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 24

Gln Ile Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
20 25 30

Asn Ile Tyr Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr Asn Gln Lys Phe
50 55 60

Arg Gly Arg Ala Thr Leu Thr Val Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 25

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 25

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
 20 25 30

Asn Ile Tyr Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr Asn Gln Lys Phe
 50 55 60

Arg Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
 100 105 110

Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 26

<211> 113

<212> PRT

<213> Mus musculus

<400> 26

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
 1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45

Ser Pro Lys Pro Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Ile Tyr Tyr Cys Gln Gln
 85 90 95

Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
 100 105 110

Lys

<210> 27

<211> 113

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 27

Asp	Ile	Val	Met	Ser	Gln	Ser	Pro	Ser	Ser	Leu	Ala	Ala	Ser	Val	Gly
1				5					10					15	

Asp	Arg	Val	Thr	Met	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser
			20					25					30		

Ser	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys
		35					40					45			

Ser	Pro	Lys	Pro	Leu	Ile	Tyr	Trp	Ala	Ser	Thr	Arg	Glu	Ser	Gly	Val
	50					55					60				

Pro	Asp	Arg	Phe	Thr	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
65					70					75					80

Ile	Ser	Ser	Val	Gln	Ala	Glu	Asp	Leu	Ala	Ile	Tyr	Tyr	Cys	Gln	Gln
			85						90					95	

Tyr	Tyr	Ser	Tyr	Pro	Trp	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile
			100					105					110		

Lys

<210> 28

<211> 113

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 28

Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5					10					15	

Asp	Arg	Val	Thr	Ile	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser
			20					25					30		

Ser	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys
		35					40					45			

Ser	Pro	Lys	Pro	Leu	Ile	Tyr	Trp	Ala	Ser	Thr	Arg	Glu	Ser	Gly	Val
	50					55					60				

Pro	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
65					70					75					80

Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Thr Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
100 105 110

Lys

<210> 29

<211> 113

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 29

Asp Ile Val Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
35 40 45

Ser Pro Lys Pro Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Gln Ala Glu Asp Lys Ala Thr Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
100 105 110

Lys

<210> 30

<211> 113

<212> **PRT**

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 30

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Pro Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
100 105 110

Lys

<210> 31

<211> 113

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 31

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
100 105 110

Lys

<210> 32

<211> 11

<212> PRT

<213> Mus musculus

<400> 32

Trp Glu Tyr Asp Gly Tyr Tyr Asp Leu Asp Tyr
1 5 10

<210> 33

<211> 17

<212> PRT

<213> Mus musculus

<400> 33

Trp Ile Asp Pro Glu Asn Gly Asp Thr Gly Tyr Ala Arg Lys Phe Gln
1 5 10 15

Gly

<210> 34

<211> 10

<212> PRT

<213> Mus musculus

<400> 34

Gly Phe Asn Ile Lys Asp Tyr Tyr Leu His
1 5 10

<210> 35

<211> 9

<212> PRT

<213> Mus musculus

<400> 35

Leu Gln Gly Thr His Phe Pro Tyr Thr
1 5

<210> 36

<211> 7

<212> PRT

<213> Mus musculus

<400> 36

Leu Val Ser Lys Leu Asp Ser
1 5

<210> 37

<211> 16

<212> PRT

<213> Mus musculus

<400> 37

Lys Ser Ser Gln Ser Leu Leu Tyr Ser Asn Gly Lys Thr Tyr Leu Asn
1 5 10 15

<210> 38

<211> 120

<212> PRT

<213> Mus musculus

<400> 38

```

Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Ser Gly Ala
1           5           10           15

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Tyr
          20           25           30

Tyr Leu His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
          35           40           45

Gly Trp Ile Asp Pro Glu Asn Gly Asp Thr Gly Tyr Ala Arg Lys Phe
          50           55           60

Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser Asn Thr Val Tyr
65           70           75           80

Leu His Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Phe
          85           90           95

Asn Ala Trp Glu Tyr Asp Gly Tyr Tyr Asp Leu Asp Tyr Trp Gly Gln
          100          105          110

Gly Thr Ser Val Thr Val Ser Ser
          115          120

```

<210> 39

<211> 112

<212> PRT

<213> Mus musculus

<400> 39

```

Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val Thr Ile Gly
1           5           10           15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
          20           25           30

Asn Gly Lys Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser
          35           40           45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro
          50           55           60

Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65           70           75           80

Ile Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Leu Gln Gly
          85           90           95

Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
          100          105          110

```

<210> 40

<211> 7

<212> PRT

<213> Mus musculus

<400> 40

Glu	Asp	Tyr	Gly	Gly	Asp	Tyr
1				5		

<210> 41

<211> 19

<212> PRT

<213> Mus musculus

<400> 41

Glu	Ile	Arg	Ser	Lys	Ser	Asn	Asn	Tyr	Ala	Thr	His	Tyr	Ala	Glu	Ser
1				5				10						15	

Val Lys Gly

<210> 42

<211> 10

<212> PRT

<213> Mus musculus

<400> 42

Gly	Phe	Thr	Phe	Ser	Asn	Tyr	Trp	Met	Asn
1				5				10	

<210> 43

<211> 9

<212> PRT

<213> Mus musculus

<400> 43

Gln	Gln	Tyr	Tyr	Ser	Tyr	Pro	Tyr	Thr
1				5				

<210> 44

<211> 17

<212> PRT

<213> Mus musculus

<400> 44

Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser	Ser	Asp	Gln	Arg	Asn	Tyr	Leu
1				5				10						15	

Ala

<210> 45

<211> 118

<212> PRT

<213> Mus musculus

<400> 45

Glu	Val	Lys	Leu	Glu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1				5				10						15	

Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asn Tyr
20 25 30

Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
35 40 45

Ala Glu Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr His Tyr Ala Glu
50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
65 70 75 80

Val Tyr Leu Gln Met Asn Asn Leu Arg Ala Glu Asp Thr Gly Ile Tyr
85 90 95

Tyr Cys Ile Gly Glu Asp Tyr Gly Gly Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Ser Val Thr Val Ser Ser
115

<210> 46

<211> 113

<212> PRT

<213> Mus musculus

<400> 46

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asp Gln Arg Asn Tyr Leu Ala Trp Tyr Gln Gln Arg Ser Gly Gln
35 40 45

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
100 105 110

Lys

<210> 47

<211> 12

<212> PRT

<213> Mus musculus

<400> 47

Phe	Glu	Tyr	Asp	Gly	Asn	Tyr	Ser	Pro	Leu	Asp	Phe
1				5					10		

<210> 48

<211> 16

<212> PRT

<213> Mus musculus

<400> 48

Trp	Val	Asp	Pro	Glu	Asn	Gly	Asp	Ser	Asp	Tyr	Ala	Pro	Lys	Phe	Gln
1				5				10					15		

<210> 49

<211> 10

<212> PRT

<213> Mus musculus

<400> 49

Gly	Phe	Asn	Ile	Lys	Asp	Tyr	Tyr	Met	His
1				5				10	

<210> 50

<211> 9

<212> PRT

<213> Mus musculus

<400> 50

Gln	Asn	Asp	His	Ser	Tyr	Pro	Leu	Thr
1				5				

<210> 51

<211> 7

<212> PRT

<213> Mus musculus

<400> 51

Gly	Ala	Ser	Thr	Arg	Glu	Ser
1				5		

<210> 52

<211> 17

<212> PRT

<213> Mus musculus

<400> 52

Lys	Ser	Ser	Gln	Ser	Leu	Leu	Asn	Ser	Gly	Asn	Gln	Lys	Asn	Tyr	Leu
1				5					10					15	

Ala

<210> 53

<211> 121

<212> PRT

<213> Mus musculus

<400> 53

Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Ser Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

Gly Trp Val Asp Pro Glu Asn Gly Asp Ser Asp Tyr Ala Pro Lys Phe
50 55 60

Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser Asn Thr Val Tyr
65 70 75 80

Leu Gln Phe Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Asn Ala Phe Glu Tyr Asp Gly Asn Tyr Ser Pro Leu Asp Phe Trp Gly
100 105 110

Gln Gly Thr Ser Val Thr Val Ser Ser
115 120

<210> 54

<211> 113

<212> PRT

<213> Mus musculus

<400> 54

Asp Ile Val Met Thr Gln Ser Pro Ser Ser Leu Ser Val Ser Ala Gly
1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser
20 25 30

Gly Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Leu Leu Ile Tyr Gly Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn
85 90 95

Asp His Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu
100 105 110

Lys

<210> 55

<211> 12

<212> PRT

<213> Mus musculus

<400> 55

Phe	Glu	Tyr	Asp	Gly	Asn	Tyr	Ser	Pro	Leu	Asp	Phe
1				5					10		

<210> 56

<211> 17

<212> PRT

<213> Mus musculus

<400> 56

Trp	Val	Asp	Pro	Glu	Asn	Gly	Asp	Ser	Asp	Tyr	Ala	Pro	Lys	Phe	Gln
1				5					10					15	

Gly

<210> 57

<211> 10

<212> PRT

<213> Mus musculus

<400> 57

Gly	Phe	Asn	Ile	Lys	Asp	Tyr	Tyr	Met	His
1				5				10	

<210> 58

<211> 9

<212> PRT

<213> Mus musculus

<400> 58

Met	Gln	Gly	Thr	His	Phe	Pro	Tyr	Thr
1				5				

<210> 59

<211> 7

<212> PRT

<213> Mus musculus

<400> 59

Leu	Val	Ser	Lys	Leu	Asp	Ser
1				5		

<210> 60

<211> 16

<212> PRT

<213> Mus musculus

<400> 60

Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser	Asn	Gly	Glu	Thr	Tyr	Leu	Asn
1				5					10					15	

<210> 61

<211> 121

<212> PRT

<213> Mus musculus

<400> 61

```

Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Ser Gly Ala
1           5           10           15

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Tyr
          20           25           30

Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
          35           40           45

Gly Trp Val Asp Pro Glu Asn Gly Asp Ser Asp Tyr Ala Pro Lys Phe
          50           55           60

Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser Asn Thr Val Tyr
65           70           75           80

Leu Gln Phe Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
          85           90           95

Asn Ala Phe Glu Tyr Asp Gly Asn Tyr Ser Pro Leu Asp Phe Trp Gly
          100          105          110

Gln Gly Thr Ser Val Thr Val Ser Ser
          115          120

```

<210> 62

<211> 112

<212> PRT

<213> Mus musculus

<400> 62

```

Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val Pro Ile Gly
1           5           10           15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
          20           25           30

Asn Gly Glu Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser
          35           40           45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro
          50           55           60

Asp Arg Phe Thr Gly Ser Arg Ser Gly Thr Asp Phe Thr Leu Lys Ile
65           70           75           80

Ser Arg Val Glu Ser Glu Asp Leu Gly Val Tyr Tyr Cys Met Gln Gly
          85           90           95

Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
          100          105          110

```

<210> 63

<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>
<221> VARIANT
<222> (1)..(1)
<223> /replace="Phe"

<220>
<221> misc_feature
<222> (1)..(1)
<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<220>
<221> VARIANT
<222> (6)..(6)
<223> /replace=" "

<220>
<221> misc_feature
<222> (6)..(6)
<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<220>
<221> VARIANT
<222> (8)..(8)
<223> /replace="Ser"

<220>
<221> VARIANT
<222> (9)..(9)
<223> /replace="Pro"

<220>
<221> misc_feature
<222> (8)..(9)
<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

<220>
<221> VARIANT
<222> (12)..(12)

<223> /replace="Tyr"

<220>

<221> misc_feature

<222> (12)..(12)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<400> 63

Trp	Glu	Tyr	Asp	Gly	Asn	Tyr	Tyr	Asp	Leu	Asp	Phe
1				5					10		

<210> 64

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (2) .. (2)

<223> /replace="Val"

<220>

<221> misc_feature

<222> (2)..(2)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<220>

<221> VARIANT

<222> (9)..(9)

<223> /replace="Ser"

<220>

<221> misc_feature

<222> (9)..(9)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<220>

<221> VARIANT

<222> (10)..(10)

<223> /replace="Asp"

<220>

<221> misc_feature

<222> (10)..(10)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<400> 64

Trp	Ile	Asp	Pro	Glu	Asn	Gly	Asp	Thr	Gly	Tyr	Ala	Pro	Lys	Phe	Gln
1				5					10					15	

Gly

<210> 65

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (9)..(9)

<223> /replace="Met"

<220>

<221> misc_feature

<222> (9)..(9)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<400> 65

Gly	Phe	Asn	Ile	Lys	Asp	Tyr	Tyr	Leu	His
1				5				10	

<210> 66

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (2)..(2)

<223> /replace="Asn"

<220>

<221> VARIANT

<222> (3)..(3)

<223> /replace="Phe" or "Asp" or "His"

<220>

<221> misc_feature

<222> (2)..(3)

<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

<220>

<221> VARIANT

<222> (4)..(4)

<223> /replace="Phe" or "His" or "Trp"

<220>

<221> misc_feature

<222> (4)..(4)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<220>

<221> VARIANT

<222> (6)..(6)

<223> /replace="Phe" or "Thr" or "His"

<220>

<221> misc_feature

<222> (6)..(6)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<220>

<221> VARIANT

<222> (8)..(8)

<223> /replace="Tyr" or "Phe" or "His" or "Leu"

<220>

<221> misc_feature

<222> (8)..(8)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<400> 66

Gln Gln Tyr Tyr Ser Tyr Pro Trp Thr
1 5

<210> 67

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (1)..(1)

<223> /replace="Gly"

<220>

<221> misc_feature

<222> (1)..(1)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<220>

<221> VARIANT

<222> (6)..(6)

<223> /replace="Asp" or "Gln" or "Asn"

<220>

<221> misc_feature

<222> (6)..(6)

<223> /note="Residue given in the sequence has no preference with respect to those in the annotations for said position"

<400> 67

Trp	Ala	Ser	Thr	Arg	Glu
1				5	

<210> 68

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (8)..(8)

<223> /replace="His" or "Tyr" or "Phe"

<220>

<221> VARIANT

<222> (9)..(9)

<223> /replace="Gly"

<220>

<221> misc_feature

<222> (8)..(9)

<223> /note="Residues given in the sequence have no preference with respect to those in the annotations for said positions"

<220>

<221> VARIANT

<222> (11)..(11)

<223> /replace="Asp"

<220>

<221> misc_feature

<222> (11)..(11)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<220>

<221> VARIANT

<222> (13)..(13)

<223> /replace="Arg"

<220>

<221> misc_feature

<222> (13)..(13)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<220>

<221> VARIANT

<222> (15)..(15)

<223> /replace="Tyr"

<220>

<221> misc_feature

<222> (15)..(15)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<400> 68

Lys	Ser	Ser	Gln	Ser	Leu	Leu	Trp	Ser	Ser	Asn	Gln	Lys	Asn	His	Leu
1				5					10					15	

Ala

<210> 69

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (1)..(1)

<223> /replace="Met"

<220>

<221> misc_feature

<222> (1)..(1)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<400> 69

Leu	Gln	Gly	Thr	His	Phe	Pro	Tyr	Thr
1				5				

<210> 70

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic consensus peptide"

<220>

<221> VARIANT

<222> (12)..(12)

<223> /replace="Glu"

<220>

<221> misc_feature

<222> (12)..(12)

<223> /note="Residue given in the sequence has no preference with respect to that in the annotation for said position"

<400> 70

Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser	Asn	Gly	Lys	Thr	Tyr	Leu	Asn
1				5					10					15	

<210> 71

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term biotin"

<400> 71

Arg Pro Pro Gly Phe Ser Pro Phe Arg
1 5

<210> 72

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term KLH"

<400> 72

Arg Pro Pro Gly Phe Ser Pro Phe Arg
1 5

<210> 73

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term KLH"

<400> 73

Arg Pro Pro Gly Phe Ser Pro Phe Arg
1 5

<210> 74

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term biotin"

<400> 74

Lys	Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe	Arg
1				5					10

<210> 75

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term biotin"

<400> 75

Lys	Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe	Arg
1				5					10

<210> 76

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term KLH"

<400> 76

Lys	Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe	Arg
1				5					10

<210> 77

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term KLH"

<400> 77

Lys	Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe	Arg
1				5					10

<210> 78

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term biotin"

<400> 78

Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe
1			5				

<210> 79

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term biotin"

<400> 79

Arg	Pro	Pro	Gly	Phe	Ser	Pro	Phe
1			5				

<210> 80

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term KLH"

<400> 80

Arg Pro Pro Gly Phe Ser Pro Phe

1

5

<210> 81

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term KLH"

<400> 81

Arg Pro Pro Gly Phe Ser Pro Phe

1

5

<210> 82

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term biotin"

<400> 82

Lys Arg Pro Pro Gly Phe Ser Pro Phe

1

5

<210> 83

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term biotin"

<400> 83

Lys Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 84

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term KLH"

<400> 84

Lys Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 85

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term KLH"

<400> 85

Lys Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 86

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term biotin"

<400> 86

Arg Arg Pro Pro Gly Phe Ser Pro Phe Arg
1 10

1 5 10

<210> 87
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>
 <221> source
 <223> /note="C-term biotin"

<400> 87
 Arg Arg Pro Pro Gly Phe Ser Pro Phe Arg
 1 5 10

<210> 88
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>
 <221> source
 <223> /note="N-term KLH"

<400> 88
 Arg Arg Pro Pro Gly Phe Ser Pro Phe Arg
 1 5 10

<210> 89
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>
 <221> source
 <223> /note="C-term KLH"

<400> 89
 Arg Arg Pro Pro Gly Phe Ser Pro Phe Arg
 1 5 10

<210> 90
<211> 9
<212> PRT
<213> Mus musculus

<400> 90
Arg Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 91
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>
<221> source
<223> /note="N-term biotin"

<400> 91
Arg Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 92
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>
<221> source
<223> /note="C-term biotin"

<400> 92
Arg Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 93
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term KLH"

<400> 93

Arg Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 94

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="C-term KLH"

<400> 94

Arg Arg Pro Pro Gly Phe Ser Pro Phe
1 5

<210> 95

<211> 5

<212> PRT

<213> Unknown

<220>

<221> source

<223> /note="Description of Unknown: Bradykinin 1-5 peptide"

<400> 95

Arg Pro Pro Gly Phe
1 5

<210> 96

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>

<221> source

<223> /note="N-term biotin"

<400> 96

Arg Pro Pro Gly Phe

1 5
 <210> 97
 <211> 136
 <212> PRT
 <213> Mus musculus

<400> 97
 Leu Pro Glu Phe Gln Val Lys Leu Glu Glu Ser Gly Ala Glu Leu Val
 1 5 10 15
 Arg Ser Gly Ala Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn
 20 25 30
 Ile Lys Asp Tyr Tyr Leu His Trp Val Lys Gln Arg Pro Glu Gln Gly
 35 40 45
 Leu Glu Trp Ile Gly Trp Ile Asp Pro Glu Asn Gly Asp Thr Gly Tyr
 50 55 60
 Ala Arg Lys Phe Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser
 65 70 75 80
 Asn Thr Val Tyr Leu His Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala
 85 90 95
 Val Tyr Tyr Phe Asn Ala Trp Glu Tyr Asp Gly Tyr Tyr Asp Leu Asp
 100 105 110
 Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr Thr
 115 120 125
 Pro Pro Ser Val Tyr Gly Ser Ser
 130 135

<210> 98
 <211> 134
 <212> PRT
 <213> Mus musculus

<400> 98
 Leu Pro Glu Phe Gln Val Gln Leu Glu Glu Ser Gly Gly Gly Leu Val
 1 5 10 15
 Gln Pro Gly Gly Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr
 20 25 30
 Phe Ser Asn Tyr Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly
 35 40 45
 Leu Glu Trp Val Ala Glu Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr
 50 55 60
 His Tyr Ala Glu Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp
 65 70 75 80
 Ser Lys Ser Ser Val Tyr Leu Gln Met Asn Asn Leu Arg Ala Glu Asp
 85 90 95

Thr Gly Ile Tyr Tyr Cys Ile Gly Glu Asp Tyr Gly Gly Asp Tyr Trp
 100 105 110

Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr Thr Pro Pro
 115 120 125

Ser Val Tyr Gly Ser Ser
 130

<210> 99

<211> 136

<212> PRT

<213> Mus musculus

<400> 99

Leu Pro Glu Phe Glu Val Gln Leu Glu Glu Ser Gly Pro Glu Leu Val
 1 5 10 15

Lys Pro Gly Thr Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser
 20 25 30

Phe Thr Asp Tyr Asn Ile Tyr Trp Val Lys Gln Ser His Gly Lys Ser
 35 40 45

Leu Glu Trp Ile Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr
 50 55 60

Asn Gln Lys Phe Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser
 65 70 75 80

Ser Thr Ala Phe Met His Leu Ser Ser Leu Thr Ser Asp Asp Ser Ala
 85 90 95

Val Tyr Tyr Cys Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp
 100 105 110

Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr Thr
 115 120 125

Pro Pro Ser Val Tyr Gly Ser Ser
 130 135

<210> 100

<211> 137

<212> PRT

<213> Mus musculus

<400> 100

Leu Pro Glu Phe Glu Val Lys Leu Gln Glu Ser Gly Ala Glu Leu Val
 1 5 10 15

Arg Ser Gly Ala Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn
 20 25 30

Ile Lys Asp Tyr Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly
 35 40 45

Leu Glu Trp Ile Gly Trp Val Asp Pro Glu Asn Gly Asp Ser Asp Tyr
50 55 60

Ala Pro Lys Phe Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser
65 70 75 80

Asn Thr Val Tyr Leu Gln Phe Ser Ser Leu Thr Ser Glu Asp Thr Ala
85 90 95

Val Tyr Tyr Cys Asn Ala Phe Glu Tyr Asp Gly Asn Tyr Ser Ser Leu
100 105 110

Asp Phe Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr
115 120 125

Thr Pro Pro Ser Val Tyr Gly Ser Ser
130 135

<210> 101

<211> 137

<212> PRT

<213> Mus musculus

<400> 101

Leu Pro Glu Phe Glu Val Lys Leu Glu Gln Ser Gly Ala Glu Leu Val
1 5 10 15

Arg Ser Gly Ala Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn
20 25 30

Ile Lys Asp Tyr Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly
35 40 45

Leu Glu Trp Ile Gly Trp Val Asp Pro Glu Asn Gly Asp Ser Asp Tyr
50 55 60

Ala Pro Lys Phe Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser
65 70 75 80

Asn Thr Val Tyr Leu Gln Phe Ser Ser Leu Thr Ser Glu Asp Thr Ala
85 90 95

Val Tyr Tyr Cys Asn Ala Phe Glu Tyr Asp Gly Asn Tyr Ser Pro Leu
100 105 110

Asp Phe Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr
115 120 125

Thr Pro Pro Ser Val Tyr Gly Ser Ser
130 135

<210> 102

<211> 133

<212> PRT

<213> Mus musculus

<400> 102

Glu Leu Asp Ile Val Met Thr Gln Thr Thr Leu Thr Leu Ser Val Thr
 1 5 10 15

Ile Gly Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu
 20 25 30

Tyr Ser Asn Gly Lys Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly
 35 40 45

Gln Ser Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly
 50 55 60

Val Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu
 65 70 75 80

Lys Ile Ile Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Leu
 85 90 95

Gln Gly Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu
 100 105 110

Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro Ser
 115 120 125

Lys Leu Glu Leu Tyr
 130

<210> 103

<211> 134

<212> PRT

<213> Mus musculus

<400> 103

Glu Leu Asp Ile Val Leu Thr Gln Ser Pro Ser Ser Leu Ala Val Ser
 1 5 10 15

Val Gly Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu
 20 25 30

Tyr Ser Ser Asp Gln Arg Asn Tyr Leu Ala Trp Tyr Gln Gln Arg Ser
 35 40 45

Gly Gln Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser
 50 55 60

Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr
 65 70 75 80

Leu Thr Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys
 85 90 95

Gln Gln Tyr Tyr Ser Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu
 100 105 110

Glu Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro
 115 120 125

Ser Lys Leu Glu Leu Tyr
130

<210> 104

<211> 134

<212> PRT

<213> Mus musculus

<400> 104

Glu Leu Asp Ile Val Met Thr Gln Thr Pro Ser Ser Leu Ala Val Ser
1 5 10 15

Val Gly Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu
20 25 30

Tyr Thr Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro
35 40 45

Gly Gln Ser Pro Lys Pro Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser
50 55 60

Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr
65 70 75 80

Leu Thr Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Ile Tyr Tyr Cys
85 90 95

Gln Gln Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu
100 105 110

Glu Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro
115 120 125

Ser Lys Leu Glu Leu Tyr
130

<210> 105

<211> 133

<212> PRT

<213> Mus musculus

<400> 105

Glu Leu Asp Ile Val Ile Thr Gln Thr Thr Leu Ser Leu Ser Val Pro
1 5 10 15

Ile Gly Gln Pro Ala Ser Ile Ser Cys Lys Ser Arg Gln Ser Leu Leu
20 25 30

Tyr Ser Asn Gly Glu Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly
35 40 45

Gln Ser Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly
50 55 60

Val Pro Asp Arg Phe Thr Gly Ser Arg Ser Gly Thr Asp Phe Thr Leu
65 70 75 80

Lys Ile Ser Arg Val Glu Ser Glu Asp Leu Gly Val Tyr Tyr Cys Met
85 90 95

Gln Gly Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu
100 105 110

Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro Ser
115 120 125

Lys Leu Glu Leu Tyr
130

<210> 106

<211> 133

<212> PRT

<213> Mus musculus

<400> 106

Glu Leu Asp Ile Val Ile Thr Gln Ser Thr Leu Thr Leu Ser Val Pro
1 5 10 15

Ile Gly Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu
20 25 30

Tyr Ser Asn Gly Glu Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly
35 40 45

Gln Ser Pro Lys Arg Gln Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly
50 55 60

Val Pro Asp Arg Phe Thr Gly Ser Arg Ser Gly Thr Asp Phe Thr Leu
65 70 75 80

Lys Ile Ser Arg Val Glu Ser Glu Asp Leu Gly Val Tyr Tyr Cys Met
85 90 95

Gln Gly Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu
100 105 110

Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro Ser
115 120 125

Lys Leu Glu Leu Tyr
130

<210> 107

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic primer"

<400> 107

aaaagcaggc ttaggagcgg ccgccatggc gtcccaggcc tcgctg

46

<210> 108

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic primer"

<400> 108

caagaaagct gggtcggatc cttataaagt tcccagaacc ctggtc 46

<210> 109

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic oligonucleotide"

<400> 109

gcatacccat acgacgtccc agactacgct 30

<210> 110

<211> 1005

<212> DNA

<213> Mus musculus

<400> 110

atggcggtccc aggcctcgct gaagctacag ccttctaacc aaagccagca ggcccctccc	60
aacatcacct cctgcgaggg cgccccggaa gcctgggatc tgctgtgtcg ggtgctgcca	120
gggtttgtca tcaactgtctg tttctttggc ctctctggga accttttagt cctgtccttc	180
ttccttttgc cttggcgacg atggtggcag cagcggcggc agcgccaac catagcagaa	240
atctacctgg ctaactggc agcttctgat ctggtgtttg tgctgggcct gcccttctgg	300
gcagagaacg ttgggaaccg tttcaactgg ccctttggaa gtgacctctg ccgggtggtc	360
agcggggtca tcaaggccaa cctgttcac agcatcttcc tgggtggtgg catcagtcag	420
gaccgctaca ggttgctggt ataccccatg accagctggg ggaaccggcg gcgacggcaa	480
gcccagtgta cctgcctgct catctgggta gctgggggcc tcttgagcac cccacgttc	540
cttctgcgtt ccgtcaaagt cgtccctgat ctgaacatct ctgcctgcat cctgcttttc	600
ccccacgaag cttggcactt tgtaaggatg gtggagtga acgttttggg tttcctcctc	660
ccattggctg ccatcctcta cttcaacttt cacatcctgg cctccctgag aggacagaag	720
gaggccagca gaaccgggtg tgggggaccc aaggacagca agacaatggg gctgatcctc	780
aaactggtag cctccttctt ggtctgctgg gcccttacc acttctttgc cttcctggat	840
ttcctggtcc aggtgagagt gatccaggac tgcctctgga aggagctcac agacctgggc	900
cttctctctc cctctctctc cctctctctc cctctctctc cctctctctc cctctctctc	960

```

ctgcagctgg ccaacttttt tgcttttgc aacagctgcc tgaacccact gatttatgtc      960
tttgcaggcc ggctcttttaa gaccagggtt ctgggaactt tataa                      1005

```

<210> 111

<211> 111

<212> PRT

<213> Mus musculus

<400> 111

```

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1           5           10           15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
          20           25           30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
          35           40           45

Ser Pro Lys Pro Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
          50           55           60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65           70           75           80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Ile Tyr Tyr Cys Gln Gln
          85           90           95

Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu
          100          105          110

```

<210> 112

<211> 105

<212> PRT

<213> Homo sapiens

<400> 112

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10           15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr
          20           25           30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
          35           40           45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
          50           55           60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65           70           75           80

Glu Asp Leu Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Ser Thr Pro Pro
          85           90           95

Thr Phe Gly Gln Gly Thr Lys Val Glu
          100          105

```

<210> 113

<211> 120

<212> PRT

<213> Mus musculus

<400> 113

```

Glu Ile Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Thr
1           5           10           15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
           20           25           30

Asn Ile Tyr Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
           35           40           45

Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr Asn Gln Lys Phe
50           55           60

Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Phe
65           70           75           80

Met His Leu Ser Ser Leu Thr Ser Asp Asp Ser Ala Val Tyr Tyr Cys
           85           90           95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
100          105          110

Gly Thr Ser Val Thr Val Ser Ser
115          120

```

<210> 114

<211> 123

<212> PRT

<213> Homo sapiens

<400> 114

```

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1           5           10           15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
           20           25           30

Tyr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35           40           45

Gly Trp Ile Asn Pro Asn Ser Gly Gly Thr Asn Tyr Ala Gln Lys Phe
50           55           60

Gln Gly Arg Val Thr Met Thr Arg Asp Lys Ser Ser Ser Thr Ala Tyr
65           70           75           80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
           85           90           95

Ala Arg Trp Gly Tyr Asp Tyr Asp Val Phe Tyr Tyr Ala Met Asp Tyr
100          105          110

```

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 115

<211> 113

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 115

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Ser
20 25 30

Ser Asn Asn Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Thr Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
100 105 110

Lys

<210> 116

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 116

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Gly Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Glv Trp Ile Ser Ala Trp Asn Glv Asn Thr Asn Trp Ala Gln Lys Leu

```

-----
50          55          60
Gln Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
65          70          75          80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85          90          95

Ala Arg Glu Tyr Ser Ser Ser Ser Asp Ala Phe Asp Ile Trp Gly Gln
100        105        110

Gly Thr Met Val Thr Val Ser Ser
115        120

<210> 117
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>
<221> source
<223> /note="N-term biotin"

<400> 117
Arg Pro Pro Gly Phe Ser Pro Phe Arg
1          5

<210> 118
<211> 127
<212> PRT
<213> Mus musculus

<400> 118
Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Ser Gly Ala
1          5          10        15

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20          25          30

Tyr Leu His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35          40          45

Gly Trp Ile Asp Pro Glu Asn Gly Asp Thr Gly Tyr Ala Arg Lys Phe
50          55          60

Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser Asn Thr Val Tyr
65          70          75          80

Leu His Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Phe
85          90          95

```

Asn Ala Trp Glu Tyr Asp Gly Tyr Tyr Asp Leu Asp Tyr Trp Gly Gln
 100 105 110

Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr Thr Pro Pro Ser
 115 120 125

<210> 119

<211> 125

<212> PRT

<213> Mus musculus

<400> 119

Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
 35 40 45

Ala Glu Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr His Tyr Ala Glu
 50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
 65 70 75 80

Val Tyr Leu Gln Met Asn Asn Leu Arg Ala Glu Asp Thr Gly Ile Tyr
 85 90 95

Tyr Cys Ile Gly Glu Asp Tyr Gly Gly Asp Tyr Trp Gly Gln Gly Thr
 100 105 110

Ser Val Thr Val Ser Ser Ala Lys Thr Thr Pro Pro Ser
 115 120 125

<210> 120

<211> 127

<212> PRT

<213> Mus musculus

<400> 120

Glu Ile Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Thr
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
 20 25 30

Asn Ile Tyr Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
 35 40 45

Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr Asn Gln Lys Phe
 50 55 60

Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Phe
 65 70 75 80

Met His Leu Ser Ser Leu Thr Ser Asp Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr Thr Pro Pro Ser
115 120 125

<210> 121

<211> 128

<212> PRT

<213> Mus musculus

<400> 121

Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Ser Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

Gly Trp Val Asp Pro Glu Asn Gly Asp Ser Asp Tyr Ala Pro Lys Phe
50 55 60

Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser Asn Thr Val Tyr

65 70 75 80

Leu Gln Phe Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Asn Ala Phe Glu Tyr Asp Gly Asn Tyr Ser Pro Leu Asp Phe Trp Gly
100 105 110

Gln Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr Thr Pro Pro Ser
115 120 125

<210> 122

<211> 128

<212> PRT

<213> Mus musculus

<400> 122

Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Ser Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

Gly Trp Val Asp Pro Glu Asn Gly Asp Ser Asp Tyr Ala Pro Lys Phe
50 55 60

50 55 60
 Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser Asn Thr Val Tyr
 65 70 75 80

Leu Gln Phe Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Asn Ala Phe Glu Tyr Asp Gly Asn Tyr Ser Pro Leu Asp Phe Trp Gly
 100 105 110

Gln Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr Thr Pro Pro Ser
 115 120 125

<210> 123

<211> 119

<212> PRT

<213> Mus musculus

<400> 123

Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val Thr Ile Gly
 1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

Asn Gly Lys Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser
 35 40 45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro
 50 55 60

Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80

Ile Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Leu Gln Gly
 85 90 95

Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

Arg Ala Asp Ala Ala Pro Thr
 115

<210> 124

<211> 120

<212> PRT

<213> Mus musculus

<400> 124

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
 1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

Ser Asp Gln Arg Asn Tyr Leu Ala Trp Tyr Gln Gln Arg Ser Gly Gln
 35 40 45

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
100 105 110

Lys Arg Ala Asp Ala Ala Pro Thr

115 120

<210> 125

<211> 120

<212> PRT

<213> Mus musculus

<400> 125

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Ser Pro Lys Pro Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Ile Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
100 105 110

Lys Arg Ala Asp Ala Ala Pro Thr
115 120

<210> 126

<211> 120

<212> PRT

<213> Mus musculus

<400> 126

Asp Ile Val Met Thr Gln Ser Pro Ser Ser Leu Ser Val Ser Ala Gly
1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser
20 25 30

Gly Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Leu Leu Ile Tyr Gly Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn
85 90 95

Asp His Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu
100 105 110

Lys Arg Ala Asp Ala Ala Pro Thr
115 120

<210> 127

<211> 360

<212> DNA

<213> Mus musculus

<400> 127

gagatccagc tgcagcagtc tggacctgag ctggtgaagc ctgggacttc agtgaagggtg	60
tcctgcaagg cttctgggta ctcattcact gactacaaca tctactgggt gaaacagagc	120
catggaaaga gccttgagtg gattggatat ttgatacctt acaatggtaa tactggctac	180
aaccagaagt tcaggggcaa ggccacattg actgttgaca agtcctccag cacagccttc	240
atgcatctca gcagcctgac atctgatgac tctgcagtct attactgtgc aaactactat	300
aggtatgacg accatgctat ggactattgg ggtcaaggaa cctcagtcac cgtctcctca	360

<210> 128

<211> 339

<212> DNA

<213> Mus musculus

<400> 128

gacattgtga tgtcacagtc tccatcctcc ctagctgtgt cagttggaga gaaggttact	60
atgagctgca agtccagtca gagcctttta tatagtagca atcaaaagaa ctacttggcc	120
tggtaccagc agaaaccagg gcagtctcct aaaccgctga tttactgggc atccactagg	180
gaatctgggg tccttgatcg cttcacaggc agtggatctg ggacagattt cactctcacc	240
atcagcagtg tgaaggetga agacctggca atttattact gtcagcaata ttatagctat	300
ccgtggacgt tcggtggagg caccaagctg gaaatcaaa	339

<210> 129

<211> 360

<212> DNA

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polynucleotide"

<400> 129

```

cagattcagc tgggtgcagtc tggcgccgaa gtgaagaaac ctggcgccag cgtgaagggtg      60
tcctgcaagg ccagcggcta cagcttcacc gactacaaca tctactgggt ccgacaggct      120
ccaggccagg gactggaatg gatcggctac ttcgaccctt acaacggcaa caccggctac      180
aaccagaagt tccggggcag agccaccctg accgtggaca agagcaccag caccgcctac      240
atggaactgc ggagcctgag aagcgacgac accgccgtgt actactgccc caactactac      300
agatacgacg accacgccat ggactactgg ggccagggca ccctggtcac cgtgtcctct      360

```

<210> 130

<211> 339

<212> DNA

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polynucleotide"

<400> 130

```

gacatcgtga tgaccagag ccccgacagc ctggccgtgt ctctgggcca gggggccacc      60
atcaactgca agagcagcca gagcctgctg tactctagca accagaagaa ctacctggcc      120
tggtatcagc agaagcccg ccagccccc aagcccctga tctactgggc cagcaccgcg      180
gagagcggcg tgcccgatag attttcgggc agcggctccg gcaccgactt caccctgacc      240
atcagcagcc tgcaggccga ggacgtggcc gtgtactact gccagcagta ctacagctac      300
ccctggacct tcggccaggg caccaagggtg gaaatcaag      339

```

<210> 131

<211> 119

<212> PRT

<213> Mus musculus

<400> 131

```

Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val Pro Ile Gly
1           5           10           15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20           25           30

Asn Gly Glu Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser
35           40           45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro
50           55           60

Asp Arg Phe Thr Gly Ser Arg Ser Gly Thr Asp Phe Thr Leu Lys Ile
65           70           75           80

```

Ser Arg Val Glu Ser Glu Asp Leu Gly Val Tyr Tyr Cys Met Gln Gly
85 90 95

Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

Arg Ala Asp Ala Ala Pro Thr
115

<210> 132

<211> 4

<212> PRT

<213> Homo sapiens

<400> 132

Lys Arg Pro Pro
1

<210> 133

<211> 6

<212> PRT

<213> Homo sapiens

<400> 133

Arg Pro Pro Gly Phe Ser
1 5

<210> 134

<211> 220

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 134

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Pro Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
100 105 110

```

      ---          ---          ---

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
    115                      120                      125

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
    130                      135                      140

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
    145                      150                      155                      160

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
                      165                      170                      175

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
                      180                      185                      190

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
    195                      200                      205

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
    210                      215                      220

<210> 135
<211> 229
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 135
Gln Ile Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1          5          10          15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
          20          25          30

Asn Ile Tyr Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Ile
          35          40          45

Gly Tyr Phe Asp Pro Tyr Asn Gly Asn Thr Gly Tyr Asn Gln Lys Phe
          50          55          60

Arg Gly Arg Ala Thr Leu Thr Val Asp Lys Ser Thr Ser Thr Ala Tyr
          65          70          75          80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
          85          90          95

Ala Asn Tyr Tyr Arg Tyr Asp Asp His Ala Met Asp Tyr Trp Gly Gln
          100          105          110

Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
          115          120          125

Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala

```

Phe Phe Leu Ala Phe Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
130 135 140

Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
145 150 155 160

Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
165 170 175

Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
180 185 190

Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
195 200 205

Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys His
210 215 220

His His His His His
225

<210> 136

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 136

Ala Ser Thr Lys Gly Pro
1 5

<210> 137

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic 6xHis tag"

<400> 137

His His His His His His
1 5

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the

references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US5807715A \[0096\]](#)
- [US4816567A \[0096\]](#)
- [US4816397A \[0096\]](#)
- [US5585089A \[0097\] \[0097\]](#)
- [EP239400A \[0097\]](#)
- [WO9109967A \[0097\]](#)
- [US5225539A \[0097\]](#)
- [US5530101A \[0097\] \[0103\]](#)
- [EP592106A \[0097\]](#)
- [EP519596A \[0097\]](#)
- [US5565332A \[0097\]](#)
- [WO2009032661A \[0098\]](#)
- [US5639641A \[0103\]](#)
- [WO9852976A1 \[0104\]](#)
- [WO0034317A2 \[0104\]](#)
- [WO8807089A1 \[0111\]](#)
- [WO9614339A1 \[0111\]](#)
- [WO9805787A1 \[0111\]](#)
- [WO9823289A1 \[0111\]](#)
- [WO9951642A1 \[0111\]](#)
- [WO9958572A1 \[0111\]](#)
- [WO0009560A2 \[0111\]](#)
- [WO0032767A1 \[0111\]](#)
- [WO0042072A2 \[0111\]](#)
- [WO0244215A2 \[0111\]](#)
- [WO02060919A2 \[0111\]](#)
- [WO03074569A2 \[0111\]](#)
- [WO04016750A2 \[0111\]](#)
- [WO04029207A2 \[0111\]](#)
- [WO04035752A2 \[0111\]](#)
- [WO04063351A2 \[0111\]](#)
- [WO04074455A2 \[0111\]](#)
- [WO04099249A2 \[0111\]](#)
- [WO05040217A2 \[0111\]](#)
- [WO05070963A1 \[0111\]](#)
- [WO05077981A2 \[0111\]](#)
- [WO05092925A2 \[0111\]](#)
- [WO05123780A2 \[0111\]](#)
- [WO06019447A1 \[0111\]](#)
- [WO06047350A2 \[0111\]](#)

- [WO06085967A2 \[0111\]](#)
- [US5648260A \[0111\]](#)
- [US5739277A \[0111\]](#)
- [US5834250A \[0111\]](#)
- [US5869046A \[0111\]](#)
- [US6096871A \[0111\]](#)
- [US6121022A \[0111\]](#)
- [US6194551B \[0111\]](#)
- [US6242195B \[0111\]](#)
- [US6277375B \[0111\]](#)
- [US6528624B \[0111\]](#)
- [US6538124B \[0111\]](#)
- [US6737056B \[0111\]](#)
- [US6821505B \[0111\]](#)
- [US6998253B \[0111\]](#)
- [US7083784B \[0111\]](#)
- [WO05047327A \[0112\]](#)
- [WO05013572A \[0114\]](#)
- [WO9208495A \[0116\]](#)
- [WO9114438A \[0116\]](#)
- [WO8912624A \[0116\]](#)
- [US5314995A \[0116\]](#)
- [EP396387A \[0116\]](#)
- [US4741900A \[0120\]](#)
- [US6193980B \[0127\]](#)
- [US20020102208A \[0138\]](#)
- [US5460785A \[0149\]](#)
- [US0874381W \[0178\]](#)
- [US20110027266A \[0178\]](#)
- [FR61616845 \[0237\]](#)
- [FR1350953 \[0237\]](#)

Non-patent literature cited in the description

- **BEDI et al.**Biochimica et Biophysica Acta, 1985, vol. 842, 190-99 [0003]
- **HILGENFELDT et al.**Analytical Biochemistry, 1995, vol. 228, 135-41 [0004]
- **CARRETERO et al.**Biochemical Pharmacology, 1976, vol. 25, 202265-2270 [0005]
- **GEPPETTI et al.**Regulatory Peptides, 1991, vol. 33, 3321-329 [0006]
- **KABAT et al.**J. Biol. Chem., 1977, vol. 252, 6609-6616 [0067]
- **KABAT et al.**Sequences of protein of immunological interest, 1991, [0067]
- **CHOTHIA et al.**J. Mol. Biol., 1987, vol. 196, 901-917 [0067]
- **MACCALLUM et al.**J. Mol. Biol., 1996, vol. 262, 732-745 [0067]

- **BRUMMELL et al.**Biochem., 1993, vol. 32, 1180-1187 [0090]
- **KOBAYASHI et al.**Protein Eng., 1999, vol. 12, 10 [0090]
- **BURKS et al.**Proc. Natl. Acad. Sci. USA, 1997, vol. 94, 412-417 [0090]
- **RICHARDSON JS**The anatomy and taxonomy of protein structureAdv Protein Chem., 1981, vol. 34, 167-339 [0093] [0208]
- **MORRISON**Science, 1985, vol. 229, 1202- [0096]
- **Ol et al.**BioTechniques, 1986, vol. 4, 214- [0096]
- **GILLIES et al.**J. Immunol. Methods, 1989, vol. 125, 191-202 [0096]
- **MORRISON et al.**Proc. Natl. Acad. Sci., 1984, vol. 81, 851-855 [0096]
- **NEUBERGER et al.**Nature, 1984, vol. 312, 604-608 [0096]
- **TAKEDA et al.**Nature, 1985, vol. 314, 452-454 [0096]
- **RIECHMANN et al.**Nature, 1988, vol. 332, 323- [0097]
- **PADLAN**Molecular Immunology, 1991, vol. 28, 4/5489-498 [0097]
- **STUDNICKA et al.**Protein Engineering, 1994, vol. 7, 6805-814 [0097]
- **ROGUSKA et al.**PNAS, 1994, vol. 91, 969-973 [0097]
- **KUNDU et al.**Biophys J, 2002, vol. 83, 723-732 [0098]
- **FREEDBERG et al.**J Am Chem Soc, 1998, vol. 120, 317916-7923 [0098]
- **BROOKS et al.**Proteins: A Theoretical Perspective of Dynamics, Structure and ThermodynamicsWiley19880000 [0098]
- **CASE et al.**J Comp Chem, 2005, vol. 26, 1668-1688 [0098]
- **BROOKS et al.**J Comp Chem, 1983, vol. 4, 187-217 [0098]
- **MACKERELL et al.**The Encyclopedia of Computational ChemistryJohn Wiley & Sons19980000vol. 1, 271-177 [0098]
- **RIZZO et al.**J Am Chem Soc, 2000, vol. 122, 5112898-12900 [0098]
- Protein Science, 2002, vol. 11, 184-187 [0099]
- Science, 2003, vol. 299, 1362-1367 [0099]
- Structure, 2006, vol. 14, 683-693 [0099]
- Drug Dev Res, 2004, vol. 61, 137-154 [0102]
- J Chromatog, 2006, vol. 837, 35-43 [0102]
- PLoS Biol, 2005, vol. 3, 3e91- [0103]
- **GENTZ et al.**Proc. Natl. Acad. Sci. USA, 1989, vol. 86, 821-824 [0118]
- **WILSON et al.**Cell, 1984, vol. 37, 767- [0118]
- **RIDGWAY, A. A. G.**Mammalian Expression VectorsButterworths19880000470-472 [0128]
- **STINCHCOMB et al.**Nature, 1979, vol. 282, 39- [0134]
- **KINGSMAN et al.**Gene, 1979, vol. 7, 141- [0134]
- **TSCHEMPER et al.**Gene, 1980, vol. 10, 157- [0134]
- **JONES**Genetics, 1977, vol. 85, 12- [0134]
- **PEIRERSZ et al.**Immunol. Cell Biol., 1987, vol. 65, 111-125 [0149]
- **SAMBROOKFRITSCHMANIATIS**Molecular Cloning: A Laboratory ManualCold Spring Harbor Laboratory Press19890000 [0156]
- DNA Cloning: A Practical Approach19850000vol. I and II, [0156]
- Oligonucleotide Synthesis19840000 [0156]
- Nucleic Acid Hybridization19850000 [0156]
- Transcription And Translation19840000 [0156]
- Animal Cell Culture19860000 [0156]
- Immobilized Cells And EnzymesIRL Press19860000 [0156]

- **B. PERBALA** Practical Guide To Molecular Cloning, 1984, [0156]
- Current Protocols in Molecular Biology John Wiley & Sons, Inc. 19940000 [0156]
- **BERMAN, H.MWESTBROOK J.FENG. Z.GILLILAND, G.BHAT, T.N.WEISSIG, H.SHINDYALOV, I.N.BOURNE, P.E.**The Protein Data Bank, Nucleic Acids Research, 2000, vol. 28, 235-242 [0178]
- **E. MONSELLIERH. BEDOUELLEJ.** Mol. Biol., 2006, vol. 362, 580-593 [0181] [0181]
- J. Mol. Biol., 1992, vol. 224, 487-499 [0181]
- **B.J. STEIPE et al.**J. Mol. Biol, 1994, vol. 240, 188-192 [0181]
- **SECO JLUQUE FJBARRIL X.**J Med Chem., 2009, vol. 52, 82363-71 [0182]
- **MALIN JONSSON et al.**J. Phys. Chem. B, 2003, vol. 107, 5511-5518 [0182]
- **PETER T. JONESPAUL H. DEARJEFFERSON FOOTEMICHAEL SNEUBERGERGREG WINTER**Nature, 1986, vol. 321, 522-525 [0183]
- **FOOTEWINTERJ.** Mol. Biol., 1992, vol. 224, 487-499 [0184] [0188]
- **VITA RZAREBSKI LGREENBAUM JAEMAMI HHOOF ISALIMI NDAMLE RSETTE APETERS B.**The immune epitope database 2.0. Nucleic Acids Res., 2010, vol. 38, D854-62 [0187]
- **SADDI GMABBOTT FV.**Pain, 2000, vol. 89, 53-63 [0209]
- **CHEN et al.**Molecular Pain, 2010, vol. 2, 6-13 [0209]
- **BENNETT GJXIE YK.**Pain, 1988, vol. 33, 87-107 [0209]

Patentkrav

1. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, hvor antistoffet eller det antigenbindende fragment omfatter:
- i) et variabelt domæne af tung kæde, der omfatter en aminosyresekvens for komplementaritetsbestemmende region 3 af tung kæde (HCDR3), en aminosyresekvens for komplementaritetsbestemmende region 2 af tung kæde (HCDR2) og en aminosyresekvens for komplementaritetsbestemmende region 1 af tung kæde (HCDR1) valgt fra gruppen, der består af:
- a) henholdsvis
- SEQ ID NO: 7 [X₁YX₂X₃DX₄HAMX₅Y], hvor
- X₁ er Y, F eller H,
- X₂ er R, D, A, V, L, I, M, F, Y eller W,
- X₃ er Y, F, W eller H,
- X₄ er D, E eller Y, og
- X₅ er D eller E;
- SEQ ID NO: 8 [YFX₁PX₂NGNTGYNQKFRG], hvor
- X₁ er D, R, A, V, L, I, M, F, Y eller W, og
- X₂ er Y, D, E, N eller Q; og
- SEQ ID NO: 9 [GYSFTDYX₁IY], hvor X₁ er N, W eller Y;
- b) henholdsvis
- SEQ ID NO: 63 [X₁EYDGX₂YX₃X₄LDX₅], hvor
- X₁ er W eller F,
- X₂ er N eller ingen aminosyre;
- X₃ er Y eller S,
- X₄ er D eller P, og
- X₅ er F eller Y;
- SEQ ID NO: 64 [WX₁DPENGDX₂X₃YAPKFQG], hvor
- X₁ er I eller V,
- X₂ er T eller S, og
- X₃ er G eller D;
- SEQ ID NO: 65 [GFNIKDYYX₁H], hvor X₁ er L eller M;
- c) henholdsvis
- SEQ ID NO: 13;

SEQ ID NO: 14; og
SEQ ID NO: 15;
d) henholdsvis
SEQ ID NO: 32;
5 SEQ ID NO: 33; og
SEQ ID NO: 34;
e) henholdsvis
SEQ ID NO: 40;
SEQ ID NO: 41; og
10 SEQ ID NO: 42;
f) henholdsvis
SEQ ID NO: 47;
SEQ ID NO: 48; og
SEQ ID NO: 49;
15 og
g) henholdsvis
SEQ ID NO: 55;
SEQ ID NO: 56; og
SEQ ID NO: 57;
20 og
ii) et variabelt domæne af let kæde, der omfatter en
aminosyresekvens for komplementaritetsbestemmende region 3 af
let kæde (LCDR3), en aminosyresekvens for
komplementaritetsbestemmende region 2 af let kæde (LCDR2) og
25 en aminosyresekvens for komplementaritetsbestemmende region 1
af let kæde (LCDR1) valgt fra gruppen, der består af:
h) henholdsvis
SEQ ID NO: 10 [QQX₁X₂SX₃PX₄T], hvor
X₁ er Y, F eller H,
30 X₂ er Y, F, H eller W,
X₃ er Y, F, T eller H, og
X₄ er W, Y, F, H eller L;
SEQ ID NO: 11 [WASTRX₁], hvor X₁ er E, D, Q eller N; og
SEQ ID NO: 12 [KSSQSLLX₁SSNQKNX₂LA], hvor
35 X₁ er W, H, Y eller F, og
X₂ er H eller Y;
i) henholdsvis
SEQ ID NO: 66 [QX₁X₂X₃SX₄PX₅T], hvor

X₁ er Q eller N,
X₂ er Y, F, D eller H,
X₃ er Y, F, H eller W,
X₄ er Y, F, T eller H, og
5 X₅ er W, Y, F, H eller L;
SEQ ID NO: 67 [X₂ASTRX₂], hvor
X₁ er W eller G, og
X₂ er E, D, Q eller N; og
SEQ ID NO: 68 [KSSQSLLX₁X₂SX₃QX₄NX₅LA], hvor
10 X₁ er W, H, Y eller F,
X₂ er S eller G,
X₃ er N eller D,
X₄ er K eller R,
X₅ er H eller Y;
15 j) henholdsvis
SEQ ID NO: 69 [X₁QGTHFPYT], hvor X₁ er L eller M;
SEQ ID NO: 36; og
SEQ ID NO: 70 [KSSQSLLYSNGX₁TYLN], hvor X₁ er K eller E;
k) henholdsvis
20 SEQ ID NO: 16;
SEQ ID NO: 17; og
SEQ ID NO: 18;
l) henholdsvis
SEQ ID NO: 35;
25 SEQ ID NO: 36; og
SEQ ID NO: 37;
m) henholdsvis
SEQ ID NO: 43;
SEQ ID NO: 17; og
30 SEQ ID NO: 44;
n) henholdsvis
SEQ ID NO: 50;
SEQ ID NO: 51; og
SEQ ID NO: 52;
35 og
o) henholdsvis
SEQ ID NO: 58;
SEQ ID NO: 59; og

SEQ ID NO: 60.

2. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det antigenbindende fragment omfatter:

a) et variabelt domæne af tung kæde, der omfatter HCDR3-, HCDR2- og HCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 13, 14 og 15 og en eller flere aminosyresubstitutioner i positioner valgt fra gruppen, der består af H1, H5, H9, H11, H12, H16, H38, H40, H41, H43, H44, H66, H75, H79, H81, H82A, H83, H87 og H108 ifølge Kabat; og

b) et variabelt domæne af let kæde, der omfatter LCDR3-, LCDR2- og LCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 16, 17 og 18 og en eller flere aminosyresubstitutioner i positioner valgt fra gruppen, der består af L5, L9, L15, L18, L19, L21, L22, L43, L63, L78, L79, L83, L85, L100 og L104 ifølge Kabat.

3. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det antigenbindende fragment deraf omfatter aminosyresekvenserne for variabelt domæne af tung kæde og let kæde ifølge henholdsvis SEQ ID NO: 24 og 30.

4. Farmaceutisk sammensætning, der omfatter antistoffet eller det antigenbindende fragment deraf ifølge krav 1 og et eller flere farmaceutisk acceptable bærestoffer.

5. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det antigenbindende fragment omfatter:
en aminosyresekvens for variabelt regiondomæne af tung kæde og

en aminosyresekvens for variabelt domæne af let kæde med mindst 90 % identitet med en aminosyresekvens valgt fra gruppen, der består af:

- a) henholdsvis SEQ ID NO: 19 og 26,
- 5 b) henholdsvis SEQ ID NO: 20 og 27,
- c) henholdsvis SEQ ID NO: 21 og 28,
- d) henholdsvis SEQ ID NO: 22 og 28,
- e) henholdsvis SEQ ID NO: 23 og 29,
- f) henholdsvis SEQ ID NO: 24 og 30,
- 10 g) henholdsvis SEQ ID NO: 25 og 31,
- h) henholdsvis SEQ ID NO: 38 og 39,
- i) henholdsvis SEQ ID NO: 45 og 46,
- j) henholdsvis SEQ ID NO: 53 og 54,
- k) henholdsvis SEQ ID NO: 61 og 62.

15

6. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det

20 antigenbindende fragment omfatter:

et variabelt domæne af tung kæde, der omfatter en aminosyresekvens, og et variabelt domæne af let kæde, der omfatter en aminosyresekvens valgt fra gruppen, der består af:

- a) henholdsvis SEQ ID NO: 19 og 26,
- 25 b) henholdsvis SEQ ID NO: 20 og 27,
- c) henholdsvis SEQ ID NO: 21 og 28,
- d) henholdsvis SEQ ID NO: 22 og 28,
- e) henholdsvis SEQ ID NO: 23 og 29,
- f) henholdsvis SEQ ID NO: 24 og 30,
- 30 g) henholdsvis SEQ ID NO: 25 og 31,
- h) henholdsvis SEQ ID NO: 38 og 39,
- i) henholdsvis SEQ ID NO: 45 og 46,
- j) henholdsvis SEQ ID NO: 53 og 54,
- k) henholdsvis SEQ ID NO: 61 og 62.

35

7. Farmaceutisk sammensætning, der omfatter antistoffet eller det antigenbindende fragment deraf ifølge krav 2 og et eller flere farmaceutisk acceptable bærestoffer.

8. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det antigenbindende fragment omfatter:

a) et variabelt domæne af tung kæde, der omfatter konsensus-HCDR3-, HCDR2- og HCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 7, 8 og 9; og

b) et variabelt domæne af let kæde, der omfatter LCDR3-, LCDR2- og LCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 10, 11 og 12.

9. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 8, hvor antistoffet eller det antigenbindende fragment omfatter:

a) et variabelt domæne af tung kæde, der omfatter HCDR3-, HCDR2- og HCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 13, 14 og 15; og

b) et variabelt domæne af let kæde, der omfatter LCDR3-, LCDR2- og LCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 16, 17 og 18.

10. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det antigenbindende fragment omfatter:

a) et variabelt domæne af tung kæde, der omfatter HCDR3-, HCDR2- og HCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 32, 33 og 34; og

b) et variabelt domæne af let kæde, der omfatter LCDR3-, LCDR2- og LCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 35, 36 og 37.

11. Isoleret monoklonalt antistof eller antigenbindende

fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det antigenbindende fragment omfatter:

- 5 a) et variabelt domæne af tung kæde, der omfatter HCDR3-, HCDR2- og HCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 40, 41 og 42; og
b) et variabelt domæne af let kæde, der omfatter LCDR3-, LCDR2- og LCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID
10 NO: 43, 17 og 44.

12. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det antigenbindende fragment omfatter:

- 15 a) et variabelt domæne af tung kæde, der omfatter HCDR3-, HCDR2- og HCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 47, 48 og 49; og
20 b) et variabelt domæne af let kæde, der omfatter LCDR3-, LCDR2- og LCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 50, 51 og 52.

13. Isoleret monoklonalt antistof eller antigenbindende fragment deraf, der binder specifikt til kallidin eller des-Arg₁₀-kallidin, men ikke til bradykinin eller des-Arg₉-bradykinin, ifølge krav 1, hvor antistoffet eller det antigenbindende fragment omfatter:

- 25 a) et variabelt domæne af tung kæde, der omfatter HCDR3-, HCDR2- og HCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 55, 56 og 57; og
30 b) et variabelt domæne af let kæde, der omfatter LCDR3-, LCDR2- og LCDR1-aminosyresekvenserne ifølge henholdsvis SEQ ID NO: 58, 59 og 60.

DRAWINGS




Figure 1




Figure 2

Heavy Chain Alignment

Parental:	<u>ETQLQQSGPELVKPGTSVKVSCCKASGYSFTDYNLYWVKQSHKKSLEWIGYFDFPYNGNTGY</u> <u>+IQL QSG E+ KPG SVKVSCKASGYSFTDYNLYWV+Q+ G+ LEWIGYFDFPYNGNTGY</u>
Humanized:	<u>QIQLVQSGAEVKKPGASVKVSCKASGYSFTDYNLYWVROAPGQGLEWIGYFDFPYNGNTGY</u>
Parental:	<u>NQKFRGKATLTVDKSSSTAFMHLSSLTSDDSAVYYCANYRYDDHAMDYWGQGTSTVTVSS</u> <u>NQKFRG+ATLTVDKS+STA+M L SL SDD+AVYYCANYRYDDHAMDYWGQGT VTVSS</u>
Humanized:	<u>NQKFRGRATLTVDKSTSTAYMELRSLRSDDTAVYYCANYRYDDHAMDYWGQGLTVTVSS</u>

Light Chain Alignment

Parental:	<u>DIVMSQSPSSLAVSVGEKVTMSCKSSQSLLYSSNQKNYLAWYQQKPGQSPKPLIYWASTR</u> <u>DIVM+QSP SLAVS+GE+ T++CKSSQSLLYSSNQKNYLAWYQQKPGQ PKPLIYWASTR</u>
Humanized:	<u>DIVMTQSPDSLAVSLGERATINCKSSQSLLYSSNQKNYLAWYQQKPGQPKPLIYWASIR</u>
Parental:	<u>ESGVPDRFTGSGSGTDFTLTISSVKAEDLAIYYCQYYSYSPWTFGGGTKLEIK</u> <u>ESGVPDRF+GSGSGTDFTLTISS++AED+A+YYCQYYSYSPWTFG GTK+ELK</u>
Humanized:	<u>ESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCQYYSYSPWTFGGGTKVEIK</u>

Figure 3




Figure 4




Figure 5




Figure 6

Heavy Chain

H1

H2

	10	20	30	40	50	60	70	80
hF151 :	QI	LVQSGAE	LVK	CSVKVSC	KASGYST	LVWVQ	AG	LEWIG
F151 :	EI	LVQSGPE	LVK	CSVKVSC	KASGYST	LVWVQ	AG	LEWIG
C063 :	EV	LVQSGPE	LVK	CSVKVSC	KASGYST	LVWVQ	AG	LEWIG
B021 :	EV	LVQSGPE	LVK	CSVKVSC	KASGYST	LVWVQ	AG	LEWIG
I022 :	EV	LVQSGPE	LVK	CSVKVSC	KASGYST	LVWVQ	AG	LEWIG
I054 :	EV	LVQSGPE	LVK	CSVKVSC	KASGYST	LVWVQ	AG	LEWIG

E3

	90	100	110	120	130	140	150	160
hF151 :	AMER	LVRSDD	LVY	LVY	LVY	LVY	LVY	LVY
F151 :	AMER	LVRSDD	LVY	LVY	LVY	LVY	LVY	LVY
C063 :	AMER	LVRSDD	LVY	LVY	LVY	LVY	LVY	LVY
B021 :	AMER	LVRSDD	LVY	LVY	LVY	LVY	LVY	LVY
I022 :	AMER	LVRSDD	LVY	LVY	LVY	LVY	LVY	LVY
I054 :	AMER	LVRSDD	LVY	LVY	LVY	LVY	LVY	LVY

E3

	170	180	190	200	210	220	230
hF151 :	TVSN	SGAL	TS	GVH	TF	PA	V
F151 :	TVSN	SGAL	TS	GVH	TF	PA	V
C063 :	TVSN	SGAL	TS	GVH	TF	PA	V
B021 :	TVSN	SGAL	TS	GVH	TF	PA	V
I022 :	TVSN	SGAL	TS	GVH	TF	PA	V
I054 :	TVSN	SGAL	TS	GVH	TF	PA	V

Figure 8




Figure 9




Figure 10





Figure 11






Figure 13

