

A&A Ref: 149945

PUBLICATION PARTICULARS AND ABSTRACT  
(Section 32(3)(a) - Regulations 22(1)(g) and 31)

21	01	PATENT APPLICATION NO	22	LODGING DATE	43	ACCEPTANCE DATE
----	----	-----------------------	----	--------------	----	-----------------

2004 2133

7 April 2004

14-12-2004

51	INTERNATIONAL CLASSIFICATION	NOT FOR PUBLICATION
----	------------------------------	---------------------

C07D

CLASSIFIED BY: ISA

71	FULL NAME(S) OF APPLICANT(S)
----	------------------------------

Institute of Pharmacology and Toxicology Academy of Military Medical Sciences p.l.a China  
Zhejiang Xianju Pharmaceutical Co. Ltd.

72	FULL NAME(S) OF INVENTOR(S)
----	-----------------------------

ZHONG, Bohua  
WANG, Yaping  
QIN, Boyi

GONG, Zehui  
LIU, Yongshao

EARLIEST PRIORITY CLAIMED	COUNTRY		NUMBER	DATE	
	33	CN	31	01142149.5	32 14 September 2001

NOTE: The country must be indicated by its International Abbreviation - see schedule 4 of the Regulations

54	TITLE OF INVENTION
----	--------------------

New oripavine derivatives and their uses as medicines


57	ABSTRACT (NOT MORE THAN 150 WORDS)
----	------------------------------------

NUMBER OF SHEETS	38
------------------	----

The sheet(s) containing the abstract is/are attached.

If no classification is furnished, Form P.9 should accompany this form.  
The figure of the drawing to which the abstract refers is attached.

**Abstract:** The present invention disclosed oripavine derivatives of formula (I) and their nontoxic pharmaceutical acceptable salts. In the formula (I), R<sup>1</sup> is hydrogenatom and methyl, R<sup>2</sup> is methyl, cyclopropyl, cyclobutyl or allyl, R<sup>3</sup> is thiophenethyl, cycloalkyl methyl wherein cycloalkyl have 3 to 6 carbon atoms. These compounds can be used in preparing analgesics or abstinence agents of opium habit-forming drugs.



## Description


### New oripavine derivatives and their uses as pharmaceuticals

#### Field of the Invention

The present invention relates to novel oripavine derivatives, their non-toxic pharmaceutically acceptable salts, and their uses in the preparation of analgesics or abstinence agents of opium habit-forming drugs.


#### Background art

The Great Britain patent 1136214 disclosed compounds represented by the following formula:



wherein R<sup>1</sup> is hydrogen or methyl, R<sup>2</sup> is cyclopropyl methyl or allyl, R<sup>3</sup> is alkyl, phenyl or phenyl alkyl. These compounds have potent central analgesic activity and opiate antagonist activity.

Among them, buprenorphine (where R<sup>1</sup> is hydrogen, R<sup>2</sup> is cyclopropyl methyl, R<sup>3</sup> is tert-butyl) has good analgesic activity and little dependence with potency 25-30 times as that of morphine and less dependence. It has now been widely used as analgesics and anti-addiction. But buprenorphine only have moderate efficacy (40% that of morphine) and low bioavailability. So it could not be administered orally.



## Buprenorphine

U.S. patent 3931189 disclosed buprenorphine analogues wherein  $R^2$  is aromatic heterocyclic alkyl; Chinese patent CN1168377A disclosed buprenorphine analogues wherein  $R^3$  is cyclobutyl or cyclopropyl.

Although the prior arts have given these teachings, it is still of great need for new drugs for the treatment of pain and addiction.


### DETAILED DESCRIPTION OF THE INVENTION

The inventor of the present invention has discovered that the oripavine derivatives represented by the formula (I) or their pharmaceutically acceptable salts not only have strong analgesic potency and analgesic efficacy, but also exhibit good oral bioavailability, long acting time, low small animal body dependency. As a result, the invention has been completed.

Therefore, the present invention at one aspect aims to provide novel oripavine derivatives represented by the formula (I) and their non-toxic pharmaceutically acceptable salts.

The present invention at another aspect relates to a pharmaceutical composition comprising as active ingredient oripavine derivatives represented by the formula (I) and their non-toxic pharmaceutically acceptable salts and pharmaceutically acceptable carriers.


The present invention relates to oripavine derivatives represented by the formula (I):




(I)

wherein  $R^1$  is hydrogen or methyl,  $R^2$  is methyl, cyclopropyl methyl, cyclobutyl methyl or allyl,  $R^3$  is thiophenylethyl or cycloalkyl methyl where the cycloalkyl has 3 to 6 carbon atoms, and their non-toxic pharmaceutically acceptable salts.


The compounds of the formula I are preferably selected from the compounds represented by the following formulae:




1




I 2




I 3




I 4



I 5




I 6



I 7


or their non-toxic pharmaceutically acceptable salts.


The oripavine derivatives according to the present invention can be prepared by the following reaction schemes:



Starting material thebaine (II) is reacted with methyl vinyl ketone by Diels-Alder reaction to obtain a compound of formula III with a yield of 80-90%; the resulting compound of formula III is hydrogenated in the presence of palladium catalyst to obtain a compound of formula IV, the resulting compound of formula IV is reacted thiophenylethyl bromide by Grignard addition reaction to obtain a compound of formula V; the compound of formula V is reacted with cyanobromide to give the compound of formula VI substituted by N-cyano group, the compound of formula VI is hydrolyzed with potassium hydroxide to obtain a compound of formula VII, which is a key intermediate. Finally the compound of formula VII is subjected alkylation reaction on nitrogen atom to yield the target compound.

The compounds of the invention wherein  $R^3$  is cycloalkyl methyl where the cycloalkyl has 3 to 6 carbon atoms can be prepared by the following schemes:





the intermediate of formula VI obtained according to the first scheme is reacted with cyanobromide to obtain a compound of formula V' substituted by N-cyano group, the resulting compound of formula V' is hydrolyzed with potassium hydroxide to obtain a compound of VI'; the resulting compound of formula VI' is reacted with cyclopropyl methyl bromide to obtain an intermediate of formula VII'; the intermediate of formula VII' is subjected to Grignard addition reaction and then demethoxylated to yield target compound of formula I.

When a compound has sufficient acidic strength or basic capability to form a stable a nontoxic acidic or basic salt, it is appropriate that the compound is administered in the form of a salt. Examples of pharmaceutically acceptable salts are organic addition salts formed with acids, these acids form physiologically acceptable anions, such as tosylates, methyl sulfonates,

acetates, citrates, malonates, tartrates, succinates, benzoates, ascorbates, alpjha-keto-glutaric acid salts, maleates, fumarates, benzenesulfonates, and alpha-glycerin phosphates. These acids also form suitable inorganic salts, including hydrobromides, hydrochlorides, sulfates, nitrates, bicarbonates and carbonates.

The pharmaceutically acceptable salts can be obtained by common methods in the art, for example, a physiologically acceptable anion can be formed by reacting a compound with sufficient basic strength such as an amine with a suitable acid.

The compounds according to the invention can be administered in the form of pharmaceutical compositions comprising the compounds and appropriate carriers. These pharmaceutical compositions can be prepared by various processes and contain common carriers in the art. The guidelines for these processes and components have been taught in Remington's Pharmaceutical Sciences, edited by E.W. Martin (Mark Publ. Co., 15<sup>th</sup> Ed., 1975 ). For the necessity for the completeness of the invention, the reference document is introduced for reference. The compounds and the pharmaceutical compositions according to the invention can be administered non-enterogastrically (such as intravenously, intraperitoneally, intramuscularly), locally, transdermally, orally or rectally.

#### Brief Description of the Drawings

Figure 1 is comparative curve of administering compound I<sub>1</sub> orally or by subcutaneous injection.

### Mode of carrying out the invention

Following examples further describes the present invention, but not limit the invention at any way.

#### **Example 1**

##### **Preparation of 7 $\alpha$ -acetyl-6,14-endoetheno tetrahydro thebaine (III)**

49.8 g of thebaine (II) and 140 ml of methyl vinyl ketone were heated and refluxed for 1 hour. Excess methyl vinyl ketone was removed by distillation under reduced pressure. To the residue was added 60 ml of methanol and the mixture was heated to dissolve the residue. The resulting solution was cooled, the precipitated solid was filtered, and washed with cooled methanol and dried to give 56.3 g of a compound of formula III with a melting point of 118-120°C, the yield is 89%.

#### **Example 2**

##### **Preparation of 7 $\alpha$ -acetyl-6,14-endoethano tetrahydro thebaine (IV)**

The mixture of 19.2 g of the compound of formula III, 4 g of 10% Pd-C and 200 ml of absolute ethanol were hydrogenated at 50-60°C, 40-50 Kg/cm<sup>-1</sup> of hydrogen pressure for 8-12 hours. Upon the completion of the reaction, the catalyst was filtered off and the filtrate was concentrated and cooled. The precipitated solid was filtered, and washed with cooled ethanol and dried to give 15.4 g of a compound of formula IV with a melting point of 135-137°C, the yield is 80%.

#### **Example 3**

##### **Preparation of 7 $\alpha$ -[(S)-1-hydroxy-1-methyl-3-(2-thiophenyl)-propyl]-6,14-endoethano tetrahydro thebaine (V)**

The Grignard reagent was prepared by the reaction of 6.38 g (0.03 mol) of 2-thiophen-2-yl ethyl bromide and 4.9 g (0.2 mol) of magnesium in 100ml

of ether. To the Grignard reagent solution was added drop-wise a solution of 4.9 g (0.013 mol) IV in 100ml of dried benzene. The mixture was heated and refluxed for 3 hours. Then the mixture was cooled to room temperature, and saturated ammonium chloride solution was added, extracted with ether, washed with water and then dried over  $MgSO_4$ . The solvent was removed under reduced pressure. The residue was recrystallized with methanol to give 4.54 g of a compound of formula V with a melting point of 183-185°C, the yield is 62%. By elemental analysis,  $C_{29}H_{37}NO_4S$  has a theoretic value (%): C 70.30, H 7.47, N 2.83, experimental value (%) was C 70.18, H 7.56, N 2.74.

#### **Example 4**

#### **Preparation of N-cyano-7a-acetyl-6,14- endoethano tetrahydro-nor-thebaine (V')**

1.6 g of cyanobromide was dissolved in 50 ml of chloroform, to the solution was added 5 g of the compound of formula IV, after refluxing for 12 hours, the solvent was removed under reduced pressure, the residue was recrystallized with absolute ethanol to give 3.2 g of a compound of formula V' with a melting point of 198-200°C, the yield is 62.3%.

#### **Example 5**

#### **Preparation of N-cyano-7a-[(S)-1-hydroxy-1-methyl-3-(thiophen-2-yl)-propyl]- 6,14- endoethano tetrahydro-nor-thebaine (VI)**

4.07 g ( 0.0385 mol) of cyanobromide was dissolved in 18 ml of dry methylene chloride; to the refluxing solution was added 4.54 g(0.0092 mol) of V in 18 ml of methylene chloride, after 4 hours of reaction, the solvent was removed by distillation. The residue was recrystallized with absolute ethanol to give 4.23 g of a compound of formula VI with a melting point of 171-173 °C, the yield is 91.1%. By elemental analysis,  $C_{29}H_{34}N_2O_4S$  has theoretic

value (%): C 68.77, H 6.72, N 5.53, and experimental value (%) was: C 68.81, H 6.72, N 5.40.

### **Example 6**

#### **Preparation of 7 $\alpha$ -acetyl-6,14-endoethano tetrahydro nor-thebaine (VI')**

To 45 ml of 2 N hydrochloric acid was added 3.82 g of the compound of formula V', the mixture was refluxed for 2 hours and then cooled to 0°C; to the cooled solution was added dropwise 0.78 g of sodium nitrite. The reaction was continued until there was no gas released, the reaction mixture was neutralized with ammonia to pH8-9 and extracted with chloroform, dried with MgSO<sub>4</sub>. The solvent was removed under reduced pressure and the residue was recrystallized with methanol to obtain 1.46 g of a compound of formula VI' with a melting point of >300°C, the yield is 40.8%.

### **Example 7**

#### **Preparation of 7 $\alpha$ -[(S)-1-hydroxy-1-methyl-3-(2-thiophenyl)-propyl]-6,14-endoethano tetrahydro-nor-oripavine hydrochloride (VII)**

To 4 g of the compound of formula VI was added 50 ml of diethylene glycol and 10 g of KOH. The mixture was stirred under N<sub>2</sub> at 190-200°C for 1 hour. After finishing the reaction, the reaction mixture was poured into ice-water, and saturated ammonium chloride solution were added to adjust the pH 8-9, The produced solid precipitate was collected and recrystallized with methanol to yield 2.9 g of a compound of formula VII with a melting point of 268-270°C, the yield is 72%. By elemental analysis, C<sub>27</sub>N<sub>33</sub>NO<sub>4</sub>S.HCl.H<sub>2</sub>O has a theoretic value (%) C 62.18, H 6.91, N 2.69; an experimental value (%) C 62.30, H 6.87, N 2.38.

### **Example 8**

## **Preparation of N-cyclopropyl-7 $\alpha$ -acetyl-6,14- endoethano tetrahydro nor-thebaine (VII')**

To 275 ml of DMF was added 11.82 g of the compound of formula VI', 5.43 ml of cyclopropyl methyl bromide and 6.4 g of sodium hydrocarbonate. The mixture was heated stirring under N<sub>2</sub> at 70°C for 16 hours. The solid was filtered off, the filtrated was removed under reduced pressure, the residue was extracted with methylene chloride, dried with MgSO<sub>4</sub>. The solvent was removed under reduced pressure. The residue was recrystallized with methanol to give 8.2g of a compound of formula VII' with a melting point of 104-106°C, the yield is 60.74%.

### **Example 9**

#### **Preparation of N-cyclopropyl-7 $\alpha$ -[(S)-1-hydroxy-1-methyl-2-cyclopropyl-ethyl]- 6,14- endoethano tetrahydro-nor-thebaine (VIII)**

The Grignard reagent was prepared by the reaction of 1.2 g of cyclopropyl methyl bromide and 0.6 g of magnesium in 20ml of ether. To the Grignard reagent solution was added drop-wise a solution of 0.88 g of the compound of formula VII' in 30ml of dried ether. The mixture was heated and refluxed for 4 hours. Then the mixture was cooled to room temperature, and saturated ammonium chloride solution was added, the organic part was separated and the water part was extracted with ether 2 times, the organic solution was combined and dried with Na<sub>2</sub>SO<sub>4</sub>. The solvent was removed under reduced pressure and the residue was chromatographed on a silica gel column, the product obtained translated to hydrochloride with HCl-ether to give 0.55 g of a compound of formula VIII which was used for the next step without further purification.

### **Example 10**

## Preparation of N-cyclopropylmethyl-7a-[(S)-1-hydroxy-1-methyl-3-(2-thiophenyl)-propyl]-6,14-endoethano tetrahydro-nororipavine hydrochloride (I<sub>1</sub>)

To 210 ml of DMF was added 7 g (0.015 mol) of the compound of formula VII, 4.1 g (0.03 mol) of cyclopropyl methyl bromide, 3.95 g (0.047 mol) of sodium hydrocarbonate. The mixture was heated stirring under N<sub>2</sub> at 70°C for 16 hours. The solid was filtered off, the solvent of the filtrate was removed under reduced pressure, the residue was extracted with methylene chloride, dried with MgSO<sub>4</sub>. The solvent was removed under reduced pressure. The residue was chromatographed on a silica gel column, recrystallized with methanol to give 3.79 g of the compound I<sub>1</sub> in the form of base in 48.4% yield, MP170-172°C. Take 1.4 g of the product and dissolved with absolute ethanol, to the solution was added dropwise the ether solution of hydrochloride. The precipitated solid was collected and recrystallized with absolute ethanol to give 1.21 g of the compound I<sub>1</sub>, melting pointing of 255-257 °C. By elemental analysis, C<sub>31</sub>H<sub>38</sub>NO<sub>4</sub>S.2HCl.0.5 H<sub>2</sub>O has theoretic values (%): C 65.61 H 7.17 N 2.47 S 5.64, and an experimental value (%) of 65.53 H 7.18 N 2.07 S 5.41. IR: 3406 cm<sup>-1</sup> (w), ν<sub>as</sub> C<sub>19</sub>-OH; 3224 cm<sup>-1</sup> (w), ν<sub>as</sub> C<sub>3</sub>-OH; 2989, 2926 cm<sup>-1</sup> (m), γ, C<sub>1,2</sub>-H, ; 1634, 1609 cm<sup>-1</sup> (w), δ, ph-H, thiophenyl; 1080, 1023 cm<sup>-1</sup> (s), δ, C-N, C-O; 1023 cm<sup>-1</sup> (s), δ, cyclopropyl methyl; <sup>1</sup>HNMR: δ (ppm, CDCl<sub>4</sub>): 7.12-6.90 (dd, 2H, Ar-H); 6.85-6.50 (ddd, 3H, thiophenyl; 5.17 (s, 1H, 3-OH); 3.56 (s, 1H, 5β-H); 3.55 (s, 3H, 6-OCH<sub>3</sub>); 3.00-1.61 (m, 15H, C<sub>20,21,15,16,9,10</sub>, C<sub>1',2'</sub>, C<sub>19</sub>-OH); 1.40 (m, 3H, 19-CH<sub>3</sub>); 1.10-0.09 (m, 11H, C<sub>17,18,7,8</sub>, C<sub>3',4'</sub>); <sup>13</sup>CNMR: δ (ppm, CDCl<sub>4</sub>): 119.471 (1C, C<sub>1</sub>); 116.511 (1C, C<sub>2</sub>); 117.352 (1C, C<sub>3</sub>); 145.547 (1C, C<sub>4</sub>); 97.370 (1C, C<sub>5</sub>); 80.506 (1C, C<sub>6</sub>); 47.188 (1C, C<sub>7</sub>); 31.599 (1C, C<sub>8</sub>); 58.254 (1C, C<sub>9</sub>); 43.636 (1C, C<sub>10</sub>); 123.944 (1C, C<sub>11</sub>); 132.211 (1C, C<sub>12</sub>); 45.686 (1C, C<sub>13</sub>); 35.880 (1C, C<sub>14</sub>); 35.607 (1C, C<sub>15</sub>); 75.771 (1C, C<sub>16</sub>); 17.726 (1C, C<sub>17</sub>); 29.763 (1C, C<sub>18</sub>); 75.771 (1C, C<sub>19</sub>);

43.454 ( 1C,C<sub>20</sub>); 23.494 ( 1C, C<sub>21</sub>); 23.236 ( 1C, C<sub>19</sub>-CH<sub>3</sub>); 52.759 ( 1C, C<sub>6</sub>-OCH<sub>3</sub>); 146.004 ( 1C, C<sub>22</sub>); 122.734 ( 1C, C<sub>23</sub>); 128.047 ( 1C, C<sub>24</sub>); 126.666 ( 1C, C<sub>25</sub>); 59.787 ( 1C, C<sub>1</sub>); 9.393 ( 1C, C<sub>2</sub>); 3.974, 3.484 ( 2C, C<sub>3',4'</sub>)

### **Example 11**

#### **Preparation of N-cyclobutyl-methyl-7 $\alpha$ -[(S)-1-hydroxy-1-methyl-3-(2-thiophenyl)-propyl]-6,14-endoethano tetrahydro-nororipavine hydrochloride (I<sub>2</sub>)**

To 25 ml of DMF was added 1 g (0.002 mol) of the compound of formula VII, 0.64g (0.0043mol) of cyclobutyl methyl bromide, 0.53 g of sodium hydrocarbonate, and 0.1 g of NaI. The mixture was heated stirring under N<sub>2</sub> at 70°C for 16 hours. The solid was filtered off, the solvent of the filtrate was removed under reduced pressure, the residue was extracted with methylene chloride, dried with MgSO<sub>4</sub>. The solvent was removed under reduced pressure. The residue was chromatographed on a silica gel column. The product collected was dissolved with absolute ethanol, to the solution was added dropwise the ether solution of hydrochloride. The precipitated solid was collected and recrystallized with methanol to give 0.59 g of I<sub>2</sub> in 50.4% yield, MP 241-3 °C. By elemental analysis, C<sub>32</sub>H<sub>40</sub>NO<sub>4</sub>S.HCl.1.5 H<sub>2</sub>O has a theoretic value (%): C 64.27 H 7.36 N 2.34 S 5.35, experimental value (%): C 65.10 H 7.41 N 2.06 S 5.01

### **Example 12**

#### **Preparation of N-allyl-methyl-7 $\alpha$ -[(S)-1-hydroxy-1-methyl-3-(2-thiophenyl)-propyl]-6,14-endoethano tetrahydro-nor-oripavine hydrochloride (I<sub>3</sub>)**

To 205 ml of DMF was added 1.5 g (0.003 mol) of the compound of formula VII, 0.60 g (0.004mol) of allyl bromide and 0.53 g of sodium hydrocarbonate. The mixture was heated stirring under N<sub>2</sub> at 70°C for 16 hours. The solid was filtered off, the solvent of the filtrate was removed under

reduced pressure, the residue was extracted with methylene chloride, dried with  $MgSO_4$ . The solvent was removed under reduced pressure. The residue was chromatographed on a silica gel column. The product collected was dissolved with absolute ethanol, to the solution was added dropwise the ether solution of hydrochloride. The precipitated solid was collected and recrystallized with methanol to give 0.35 g of the compound of  $I_3$  with a melting point of 228-30°C, the yield is 22%. By elemental analysis,  $C_{30}H_{36}NO_4S.HCl$  has theoretic value (%): C 66.23 H 7.00 N 2.58 S 5.89; experimental value (%): C 66.40 H 7.14 N 2.36 S 5.80.

### **Example 13**

#### **Preparation of $7\alpha$ -[(S)-1-hydroxy-1-methyl-3-(2-thiophenyl)-propyl]-6,14-endoethano tetrahydro-oripavine hydrochloride ( $I_4$ )**

To 50 ml of diethylene glycol was added 4 g of the compound of formula V and 10 g of KOH . The mixture was stirred under  $N_2$  at 190-200 °C for 1 hour and then was poured into ice-water. Saturated ammonium chloride solution were added to adjust pH 8-9. The precipitated solid was collected and recrystallized with methanol give 2.9 g of base of  $I_4$  in 72 yield, MP 268-270°C. The base was translated to salt of hydrochloride with HCl-ether to give 3.1 g of a compound of formula  $I_4$  with a melting point of >300°C. By element analysis,  $C_{28}H_{34}NO_4S.HCl$  has theoretic value (%) C 64.86 H 6.95 N 2.70 S 6.18; experimental value (%): C 64.42 H 7.22 N 2.54 S 5.20.

### **Example 14**

#### **Preparation of N-cyclopropylmethyl- $7\alpha$ -[(S)-1-hydroxy-1-methyl-2-cyclopropyl-ethyl]-6,14-endoethano tetrahydro-nor-oripavine hydrochloride ( $I_5$ )**

To 8 ml of diethylene glycol was added 1.8 g of KOH . The mixture was heated under N<sub>2</sub> at 205-6°C until the temprature stable. Then 0.55 g of the compound of formula VIII was added to the reaction mixture, stirred at the same temperature for 2 hours. The reaction mixture was poured into ice-water. Saturated ammonium chloride solution was added to adjust pH 8-9. The precipitated solid was collected and chromatographed on silica gel column. The product collected and made to salt of hydrochloride with HCl-ether, 0.16 g of a compound I<sub>5</sub> was obtained in 30% yield, melting pointing of 185-190°C. By element analysis, C<sub>29</sub>H<sub>40</sub>NO<sub>4</sub>. HCl has theroretic value (%): C 69.25, H 7.96, N 2.79, S 6.37; experimental value (%): C 69.24 , H 7.72 , N 2.45, S 6.68.

According to similar methods to prepare the compound I<sub>5</sub> , N-cyclopropylmethyl-7 $\alpha$ -[(S)-1-hydroxy-1-methyl-2-cyclopentyl-ethyl]-6,14-endoethano tetrahydro-nor-oriavine hydrochloride ( I<sub>6</sub> ) , with a melting point of 270°C, decomposed, was prepared by reaction of cyclopropyl methyl chloride compound via Grinard addition reaction. Also N-cyclopropylmethyl-7 $\alpha$ -[(S)-1-hydroxy-1-methyl-cyclohexyl-ethyl]-6,14-endoethano tetrahydro-nor-oriavine hydrochloride ( I<sub>7</sub> ) , with a melting point of 241-6 °C, decomposed, was prepared by reaction of cyclohexyl methyl chloride compound.

### **Experiment of Pharmacological Activity Evaluation**

Pharmacological effect of the compounds according to the invention was measured by the methods of acetic acid writhing test, 55°C hot plate test, rat 55°C hot water bath test and so on.

#### **1.Antinociceptive tests**

##### **1.1. Methods**

###### **(1) mice 55°C hot plate test (sc/po)**

Female mice weighing 18 - 22 g were used in the test. The latency was measured as the period from placing the animals on the 55°C plate to the appearance of response to the hot nociceptive stimulus (licking hindpaws, jumping and flicking of paws) before and after administration of drugs (sc/po), respectively. Each group had 8 mice. The cut off time was 60 sec. The results were expressed as possible maximal analgesic percentage (PMAP),

$$PMAP = \frac{Latency \ after \ administration - Latency \ before \ administration}{60 - Latency \ before \ administration} \times 100$$

ED<sub>50</sub> was calculated with Logit program.

### (2) Rat hot water bath test

Male and female (1 : 1) Wistar rats weighing 180 - 200 g were used. The latency was defined as a period from insertion of the tip part of the rat tail into 55°C hot water to withdrawal of tail from it before and after administration of drugs, respectively. Each group had 8 mice. The results were expressed as PMAP and the computational method was same as that as mentioned above. The longest latency was defined as 15 s. And then ED<sub>50</sub> was calculated with Logit program.

### (3) Acetic acid writhing test

Male and female (1 : 1) mice weighing 18 - 22 g were used. Acetic acid (0.6 %, 0.4 ml per mouse, ip) was administered to each animal. 5 min later, the frequency of writhing in the following 15 min was counted. The drugs or normal saline were administrated by sc 30 min and by po 60 min prior to injection of acetic acid, respectively. ED<sub>50</sub> was calculated with Logit program.

$$\text{analgesic percentage} = \frac{\text{writhing number in normal saline} - \text{writhing number in drug}}{\text{writhing number in normal saline}} \times 100$$

## 1.2 Results

**Table 1 Antinociceptive activity of compounds**

Compounds	acetic acid writhing test		hot plate test	
	ED <sub>50</sub> ( $\mu\text{g}\cdot\text{kg}^{-1}$ )	Max.efficacy (%)	ED <sub>50</sub> ( $\mu\text{g}\cdot\text{kg}^{-1}$ )	Max.efficacy (%)
I <sub>1</sub>	116	100.0	860	81.7
I <sub>2</sub>	Not determined	Not determined	159	71.2
I <sub>4</sub>	Not determined	Not determined	15	100

In the mice hot plate test, analgesic dose-response curves of compound of compound I<sub>1</sub> was familiar with that of buprenorphine, Max. efficacy of which was less than 100%, which produced a dose-dependent partial agonist properties. Efficacy of compound of compound I<sub>1</sub> was stronger than that of buprenorphine, especially in serious analgesic models.

In the mice acetic acid writhing test, the Max. analgesic efficacy of compound I<sub>1</sub> reached 100% while that of buprenorphine was 92.5%. In the hot plate test, the Max. analgesic efficacy of compound I<sub>1</sub> was 81.7% while that of buprenorphine was 40.3%(Tab 1). In rat 55 °C hot water bath test and rat formaldehyde test, efficacy and potency of compound I<sub>1</sub> was stronger than that of buprenorphine (Tab 2). In the rhesus monkey tail flick test, the latency of tail flick was prolonged along with increasing of dose when given compound I<sub>1</sub> 16- $\mu\text{g}\cdot\text{kg}^{-1}$  im or 6-24 $\mu\text{g}\cdot\text{kg}^{-1}$  po, but the effects of the latter was weaker than that of former (Tab 4). From above results, compound I<sub>1</sub> showed preferable antinociceptive activity.

**Table 2 Antinociceptive activity of compounds**

Compounds	writhing test		hot plate test	
	ED <sub>50</sub> ( $\mu\text{g}\cdot\text{kg}^{-1}$ )	Max.efficacy (%)	ED <sub>50</sub> ( $\mu\text{g}\cdot\text{kg}^{-1}$ )	Max.efficacy (%)
I <sub>5</sub>	6.39	100.0	200	71.2
I <sub>6</sub>	55.9	100	50	54.2
I <sub>7</sub>	Not determined	Not determined	159.1	94
Bup.	7.3	90	1100	40

In the mice hot plate test, efficacy and potency of compound I<sub>5</sub>- I<sub>7</sub> was stronger than that of buprenorphine, Max. analgesic efficacy of which were less than 100%, producing a dose-dependent partial agonist properties.

**Table 3 Comparison of Antinociceptive effects of compounds I<sub>1</sub> and buprenorphine in rats**

Compounds	hot water bath test		formaldehyde test	
	ED <sub>50</sub> (mg·kg <sup>-1</sup> )	Max.efficacy (%)	ED <sub>50</sub> (mg·kg <sup>-1</sup> )	Max. efficacy P(%)
buprenorphine	8.75	64.7	0.34	97.7
I <sub>1</sub>	3.48	79.3	0.15	100.0

**Table 4 Analgesic effect of compound I<sub>1</sub> in the rhesus monkey tail flick test**

Dose μg·kg <sup>-1</sup>	Pathway	of	Latency tail flick (X±SD, sec)	Dose μg·kg <sup>-1</sup>	Pathway	of	latency (X±SD, sec)
0	Im		3.5±0.2	0	po		4.0±0.7
1	Im		4.8±2.1	6	po		5.1±1.8
2	Im		6.5±3.6	12	po		6.0±1.9
4	Im		7.9±2.5*	24	po		6.7±2.3*
6	Im		9.2±3.1*	48	po		5.7±0.5*

\* P<0.05, Comparison with 0 μg·kg<sup>-1</sup> group; n=5

In mice hot plate test, analgesic effects of the compound I<sub>1</sub> administered by po and sc, was compared (see figure 1). The ED<sub>50</sub> value po was 2.5 times of that of sc, but the Max. efficacy of them was almost identical. In the same test, the ED<sub>50</sub> value of buprenorphine po was 12.4 times of that of sc. In other analgesic models, the ratio of ED<sub>50</sub> value po/ ED<sub>50</sub> value sc of compound I<sub>1</sub> was less than that of buprenorphine (Tab 4). The above results

indicated that bioavailability of compound I<sub>1</sub> was higher than that of buprenorphine, the range of effective dose of compound I<sub>1</sub> was 1-3mg·kg<sup>-1</sup> in two analgesic models.

## 2. Physical dependent test

### 2.1. Methods and animals

Male Swiss mice, weighing 18 - 22 g, pretreated with morphine (24 mg·kg<sup>-1</sup>, s.c.) or buprenorphine (3.6 mg·kg<sup>-1</sup>, s.c.) or compound I<sub>1</sub> (5.0 mg·kg<sup>-1</sup>, s.c.) respectively 3 times daily for 14d, naloxone(10 mg·kg<sup>-1</sup>, i.p.) was injected 4 h after the last administration of drugs, The number of jumping was immediately observed within a period of 15 min and loss of body weight were marked 60 min after administration of naloxone.

Wister rats, male, weighing 180 - 200 g, were used. Morphine, buprenorphine and compound I<sub>1</sub> were administrated as mentioned above. 4 h after the last administration of drugs, all subjects were injected naloxone(5 mg·kg<sup>-1</sup> s.c.). The frequency of gasps, ptosis, shakes, teeth chatter and yawns was immediately observed for 15 min after injection of naloxone. The total score for abstinence signs was calculated as the sum of the scores for all individual signs of the withdrawal reaction. Loss of body weight was recorded for 60 min after injection of naloxone.

### 2.2. Results

There was a significant increase in the number of total abstinence signs in morphine-treated group (30×Tid×7d, 30×Tid×14d) compared with saline group. Statistical evaluation of these data showed no significant differences in the number of abstinence syndrome in compound I<sub>1</sub>-treated mice compared with saline group. These data indicated that compound I<sub>1</sub> has low potency of dependence.

**Table 5 comparison of the analgesic effect of compound by s.c. and p.o. administration in mice.**

Methods of test	Compound	ED <sub>50</sub> (mg/kg)		
		S.C	P.O	P.O/S.C
Acetic Writhing	Buprenorphine	0.02	0.37	17.6
	I <sub>1</sub>	0.08	0.64	8.2
Hot plate	Buprenorphine	1.01	12.52	12.4
	I <sub>1</sub>	0.57	3.10	5.4
tail-flick	Buprenorphine	8.75	---	Not detectable
	I <sub>1</sub>	1.75	2.61	1.5

\* < 30%

**Table 6 comparison of the experimental results on the physical dependence in mice**

Compound	Dose (mg · kg <sup>-1</sup> )	Percent of jumping (%)	Number jumping	of loss weight (g)
Saline	—	20	2.1 ± 3.0	0.4 ± 0.4
Morphine	24.0 × Tid × 14d	100	49.5 ± 56.0**	1.6 ± 0.2**
Buprenorphine	3.6 × Tid × 14d	80	14.6 ± 27.4*	0.8 ± 0.2
I <sub>1</sub>	5.0 × Tid × 14d	0	0 ± 0	0.5 ± 0.2

## 2. Psychological dependence experiment

### (1) Conditioned Place Preference testing in mice

Swiss mice, male, weighted 18-22g. Animals were immediately confined for 40 min to one compartment after injection of compound I<sub>1</sub>, and to the other compartment after injection of saline. Animals that had been injected with drugs were confined to one of the end compartments A.M, and

to the other of the end compartments after the injection of saline P.M. This tendentious conditioning cycle was performed for 5d. On day 6, preference state, after placing the animals in the neutral middle compartment and allowing them free access to each compartment. The time spent in drug-paired compartment was measured.

The results are shown in Table 7. As seen in Table 7, morphine ( $10 \text{ mg} \cdot \text{kg}^{-1}$ ) and buprenorphine ( $0.3 \text{ mg} \cdot \text{kg}^{-1}$ ) and compound  $I_1$  ( $3$  and  $10 \text{ mg} \cdot \text{kg}^{-1}$ ) induced significant Conditioned Place Preference.

**Table 7 comparison of the experimental results on the physical dependence in rats**

Compound	Dose ( $\text{mg} \cdot \text{kg}^{-1}$ )	Score of withdrawal loss weight	
		symptoms(%)	(g)
Saline	—	$1.8 \pm 1.3$	$1.4 \pm 0.6$
Morphine	$30 \times \text{Tid} \times 7\text{d}$	$8.0 \pm 2.2^{**}$	$12.0 \pm 2.6^{**}$
Buprenorphine	$5 \times \text{Tid} \times 7\text{d}$	$4.0 \pm 2.0$	$4.2 \pm 2.5$
$I_1$	$3 \times \text{Tid} \times 7\text{d}$	$2.2 \pm 1.3$	$0.4 \pm 0.6$
Saline	—	$1.8 \pm 1.3$	$2.8 \pm 1.3$
Morphine	$30 \times \text{Tid} \times 14\text{d}$	$6.4 \pm 2.2^{**}$	$4.2 \pm 1.2$
Buprenorphine	$5 \times \text{Tid} \times 14\text{d}$	$3.6 \pm 1.1$	$4.2 \pm 1.3$
$I_1$	$3 \times \text{Tid} \times 14\text{d}$	$2.8 \pm 0.8$	$2.0 \pm 1.9$

**Table 8 Experimental results of conditional place preference in mice**

Compound	Dose (mg/kg)	Time spent in drug-paired place (min)
Saline	---	316.2 ± 119.6
Morphine	10	566.6 ± 131.1**
Buprenorphine	0.3	547.5 ± 175.0**
$I_1$	1	415.3 ± 119.2
	3	452.3 ± 66.6 *
	10	474.0 ± 136.4*

## (2) Self –administration testing

The subjects were male Wistar rats, 350-400g and rhesus monkey, 4-6 kg at the start of the experiments. Anesthesia was induced by administration of sodium pentobarbital ( $40 \text{ mg} \cdot \text{kg}^{-1}$ , i.p.). While anesthetized, rats were implanted with guide cannulas, one end of guide cannulas was implanted right atrium, and the other end was connected with self-administration installation. Penicillin G was administered immediately after surgery.

Self-administration sessions were began 3 days after surgery. Rats were trained to self-administration compound  $I_1$  at a dose of  $0.1 \text{ mg} \cdot \text{kg}^{-1}$  infusion on FR1 schedule. Daily sessions were 6 h. In duration, the number of infusions was recorded, while animals were deprived food.

The results indicated that compound  $I_1$  ( $0.05, 0.08, 0.1 \text{ mg} \cdot \text{kg}^{-1}$ ) couldn't induce self-administration in rat in continuous 35-40 days session (daily injections  $< 10$ ). Opiate receptor partial agonist buprenorphine also couldn't induce self-administration. Compound  $I_1$  ( $0.025-0.05 \text{ mg} \cdot \text{kg}^{-1}$ /inject) couldn't induce the development of self-administration in rhesus monkey. These data indicate that the potency of dependence of compound  $I_1$  is weak.

## 3. Substitution test

## **(1) Methods and animals**

Methods and animals were as similar to self-administration test. Heroin was substituted by compound I<sub>1</sub> after development of stable self-administration induced by heroin in rat. Step number and rating was recorded.

Stable self-administration was developed by morphine (0.25 mg·kg<sup>-1</sup>/inject) in rhesus monkey. Compound I<sub>1</sub> (0.025 mg·kg<sup>-1</sup>/inject), morphine (0.25 mg·kg<sup>-1</sup> mg·kg<sup>-1</sup>/inject), buprenorphine (0.05 mg·kg<sup>-1</sup>/inject), morphine (0.25 mg·kg<sup>-1</sup>/inject) and compound I<sub>1</sub> (0.05 mg·kg<sup>-1</sup>/inject) was substituted by turns.

## **(2) Results**

Self-administration could continue induced by compound I<sub>1</sub> in heroin dependent rat. There were not increases of step number of 3 rats, but increased in other 3 rats. Step numbers were decreased in most of rat compared with the treatment of heroin.

Compound I<sub>1</sub> (0.025 mg·kg<sup>-1</sup>/inject) could continue self-administration in the place of morphine (0.25 mg/kg/inject), so did buprenorphine. Compound I<sub>1</sub> (0.05 mg·kg<sup>-1</sup>/inject) could not maintain this test in replace of morphine (0.25 mg·kg<sup>-1</sup>/inject) or buprenorphine (0.05 mg·kg<sup>-1</sup>/inject). Results inferred that compound I<sub>1</sub> had potency of addict, was as similar as buprenorphine, and was weaker than morphine.

## **4. Na<sup>+</sup> index Determination**

### **(1) Methods**

The subjects were male wistar rats, weighing 180 - 200 g. Rats were sacrificed by decapitation. The brain of rats without cerebellum was used to make membrane preparation containing opiate receptors. The brain was homogenized in 50 mmol·L<sup>-1</sup>tris-HCL solutions and centrifuged (20,000 rev/min, 20 min). This procedure was repeated twice more. Each assay contained 0.5 mg membrane protein, 5 nmol · L<sup>-1</sup> <sup>3</sup>H-naloxone and different concentrations of drugs ( 1 - 100000 nmol · L<sup>-1</sup>). Nonspecific binding was

determined in the presence of  $10 \mu\text{mol} \cdot \text{L}^{-1}$  naloxone. The  $\text{IC}_{50}$  was calculated with Logit method. The  $\text{Na}^+$  index was determined by the ratio of the  $\text{IC}_{50}$  in the presence of  $100 \text{ nmol} \cdot \text{L}^{-1}$   $\text{NaCl}$  to those in absence of it.

## (2) Results

Compound  $\text{I}_1$  could inhibit  $^3\text{H}$ -naloxone binding with opiate receptors as similar to morphine and buprenorphine. The affinity of formula  $\text{I}_1$  to opiate receptors was higher 100 times than that of morphine.  $\text{Na}^+$  index is 0.42, which is character of typical antagonist.

**Table 9 Influence of  $\text{Na}^+$  in competitive inhibition of drugs on binding affinity with opiate receptors of rat brain**


Compound	$\text{IC}_{50} (\text{nmol} \cdot \text{L}^{-1})$		$\text{Na}^+$ index
	Free of $\text{Na}^+$	Presence of $\text{Na}^+$	
Morphine	18	170	9.44
Buprenorphine	0.45	0.30	0.67
$\text{I}_1$	0.19	0.08	0.42

In all, we have described this invention in detail by using preferred embodiments of the invention. Obviously, it is allowed to improve and transform the invention at the premise of undeviating from privilege of the invention.

## Claims

We claim:


1. An oripavine derivative represented by formula (I),




(I)

wherein R<sup>1</sup> is hydrogen or methyl, R<sup>2</sup> is methyl, cyclopropyl methyl, cyclobutyl methyl or allyl, R<sup>3</sup> is thiophenyl ethyl or cycloalkyl methyl where the cycloalkyl has 3 to 6 carbon atoms, or a non-toxic pharmaceutically acceptable salt thereof.


2. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 1, characterized in that R<sup>3</sup> is thiophenyl ethyl.
3. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 1, characterized in that R<sup>2</sup> is cyclopropyl methyl, R<sup>3</sup> is cycloalkyl methyl where the cycloalkyl has 3 to 6 carbon atoms.
4. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 2, characterized in that the compound is represented by the following formula:




5. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 2, characterized in that the compound is represented by the following formula:




6. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 2, characterized in that the compound is represented by the following formula:




7. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 2, characterized in that the compound is represented by the following formula:




8. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 3, characterized in that the compound is represented by the following formula:



9. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 3, characterized in that the compound is represented by the following formula:



10. The oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to claim 3, characterized in that the compound is represented by the following formula:



11. A pharmaceutical composition comprising as active ingredient oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to any one of claims 1 to 10 and pharmaceutical acceptable carriers.

12. Use of oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to any one of claims 1 to 10 for the preparation of analgesics or abstinence agents of opium habit-forming drugs.

13. Use of oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to any one of claims 1 to 10 in the manufacture of a preparation for treating pain or drug addiction.
14. A substance or composition for use in a method of analgesia or as an abstinence agent of opium habit-forming drugs, said substance or composition comprising an oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to any one of claims 1 to 10, and said method comprising administering said substance or composition.
15. A substance or composition for use in a method of treating pain or drug addiction, said substance or composition comprising oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to any one of claims 1 to 10, and said method comprising administering said substance or composition.
16. A compound according to claim 1, substantially as herein described and illustrated.
17. A composition according to claim 11, substantially as herein described and illustrated.
18. Use according to claim 12, substantially as herein described and illustrated.
19. A substance or composition for use in a method of treatment according to claim 13, substantially as herein described and illustrated.

20. A new compound, a new composition, a new use of an oripavine derivative or non-toxic pharmaceutically acceptable salt thereof according to any one of claims 1 to 10, or a substance or composition for a new use in a method of treatment, substantially as herein described.

Fig. 1

