



US00PP16118P3

(12) **United States Plant Patent**  
**Hofmann**

(10) **Patent No.:** **US PP16,118 P3**

(45) **Date of Patent:** **Nov. 15, 2005**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED  
'FISUPNIC LIPINK'**

(50) Latin Name: *Impatiens hawkeri*  
Varietal Denomination: **Fisupnic Lipink**

(75) Inventor: **Birgit C. Hofmann**, Bendorf (DE)

(73) Assignee: **Florfis AG**, Binningen (SE)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 56 days.

(21) Appl. No.: **10/829,217**

(22) Filed: **Apr. 22, 2004**

(65) **Prior Publication Data**

US 2004/0216202 P1 Oct. 28, 2004

(30) **Foreign Application Priority Data**

Apr. 22, 2003 (CH) ..... 03-1996

(51) **Int. Cl.<sup>7</sup>** ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./318**

(58) **Field of Search** ..... **Plt./318**

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

PP10,109 P \* 11/1997 Trees ..... Plt./318

PP11,706 P \* 12/2000 Klemm ..... Plt./318

PP13,180 P2 \* 11/2002 Strope ..... Plt./138

PP13,579 P2 \* 2/2003 Hofmann ..... Plt./138

PP13,714 P2 \* 4/2003 Hofmann ..... Plt./318

2002/0174469 P1 \* 11/2002 Strope ..... Plt./318

2004/0216202 P1 \* 10/2004 Hofmann ..... Plt./318

**OTHER PUBLICATIONS**

UPOV ROM GIITM Computer Database, GTI Jouve  
Retrieval Software 2004/06 Citation(s) for 'Fisupnic  
Lipink'.\*

\* cited by examiner

*Primary Examiner*—Howard J. Locker

*Assistant Examiner*—W. C. Haas

(74) *Attorney, Agent, or Firm*—Foley & Lardner, LLP

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea *Impatiens* plant  
named 'Fisupnic Lipink', and particularly characterized by  
the combined features of light pink to nearly white flowers  
with purple eyes; large, round, flat flowers; uniform, deep  
green foliage; moderately tall, rounded plant habit; and early  
to medium flowering response.

**1 Drawing Sheet**

**1**

Botanical description: *Impatiens hawkeri*.  
Variety denomination: 'Fisupnic Lipink'.

**BACKGROUND OF THE INVENTION**

The present invention comprises a new and distinct cul-  
tivar of New Guinea *Impatiens* plant, botanically known as  
*Impatiens hawkeri*, and hereinafter referred to by the cultivar  
name 'Fisupnic Lipink'.

'Fisupnic Lipink' is a product of a planned breeding  
program and originated from a hybridization made by the  
inventor, Birgit C. Hofmann, in a controlled breeding pro-  
gram in Hillscheid, Germany, in 2000.

The female parent was the variety 'Malita' (unpatented),  
characterized by carmine and orange mixed flower color,  
flowers with a purple center, medium to light green foliage,  
and fairly compact plant habit.

The male parent of 'Fisupnic Lipink' was 'Fisimp 131'  
(U.S. Plant Pat. No. 13,714), characterized by round, rose-  
red flowers, medium green foliage, medium sized, and round  
and tight plant habit.

'Fisupnic Lipink' was discovered and selected as one  
flowering plant within the progeny of the stated cross by the  
inventor in April, 2001 in a greenhouse in Galdar, Gran  
Canaria, Spain.

The first act of asexual reproduction of 'Fisupnic Lipink'  
was accomplished when vegetative cuttings were taken from

**2**

the initial selection in July, 2001 in a controlled environment  
in Galdar, Gran Canaria, Spain, by, or under the supervision  
of the inventor.

Horticultural examination of plants grown from these  
cuttings initiated in the spring of 2002 in Hillscheid,  
Germany, and continuing thereafter, has demonstrated that  
the combination of characteristics as herein disclosed for  
'Fisupnic Lipink' are firmly fixed and are retained through  
successive generations of asexual reproduction. The new  
cultivar reproduces true to type.

'Fisupnic Lipink' has not been observed under all possible  
environmental conditions. The phenotype may vary signifi-  
cantly with variations in environment such as temperature,  
light intensity and day length, without, however, any varia-  
tion in genotype. The following observations, measurements,  
and comparisons describe plants grown in  
Hillscheid, Federal Republic of Germany, under greenhouse  
conditions which approximate those generally used in com-  
mercial practice.

**BRIEF SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and  
are determined to be basic characteristics of 'Fisupnic  
Lipink', which in combination distinguish this *Impatiens* as  
a new and distinct cultivar:

1. light pink to nearly white flowers with purple eyes;
2. large, round, flat flowers;

3. uniform, deep green foliage;
4. moderately tall, rounded plant habit; and
5. early to medium flowering response.

Of the many commercial cultivars known to the inventor, the most similar in comparison to 'Fisupnic Lipink' is the patented variety 'Fisnics Light Pink' (U.S. Plant Pat. No. 13,579).

In comparison to 'Fisnics Light Pink', 'Fisupnic Lipink' has a more variable pink flower color, and much taller, more upright plant habit.

#### BRIEF DESCRIPTION OF THE DRAWING

The accompanying photographic drawing shows typical flower and foliage characteristics of 'Fisupnic Lipink' with colors being as true as possible with illustrations of this type.

The photographic drawing shows a side view of a typical flowering plant of 'Fisupnic Lipink'.

#### DETAILED BOTANICAL DESCRIPTION

In the following description color references are made to The Royal Horticultural Society Colour Chart (R.H.S.). The color values were determined indoors from plants growing in a green-house in May 2003, Hillscheid, Germany.

The description is based on plants which were planted as rooted cuttings in 12 cm pots in late February 2003, and then grown in the greenhouse at a minimum temperature of 16° C. Most observations and measurements were made after the beginning of flowering in mid May, when the plants were about 12 weeks old.

#### PLANT

General appearance and form:

*Plant habit.*—Tall and upright, surface of foliage canopy rounded, and well-branched; growth is indeterminate, though weak after beginning of flowering.

*Height.*—17.9 cm.

*Width.*—31.3 cm.

*Number of branches.*—Approximately 11.5 per plant.

*Length of branches.*—12 to 14 cm.

*Internode length.*—5.5 to 7.0 cm.

*Diameter of branches.*—8 to 9 mm.

*Stem color.*—Brown, RHS 178 C.

*Propagation.*—Terminal shoot tips for cuttings.

*Rooting.*—Roots initiate in about 18 days at 22° C., from sticking to transplanting.

*Cultivation time.*—It takes about 9 to 10 weeks of growing time to produce a marketable flowering plant in a 12 cm pot.

Foliage:

*Leaf arrangement.*—Primarily in whorls.

*Shape of leaf.*—Elliptic, with acute base and acuminate tip.

*Texture.*—Surface slightly glossy, smooth or faintly rippled.

*Margin.*—Slightly serrated, ciliated.

*Leaf length.*—13.3 cm.

*Leaf width.*—4.2 cm.

*Upper surface, main color.*—Medium to dark green, uniform; mature leaves between RHS 137 A and 139 A, young leaves near RHS 137 A.

*Veins on upper surface color.*—Dark red, RHS 183 C, near the base.

*Lower surface color.*—Light green, about RHS 139 C (both young and mature leaves).

*Veins on lower surface color.*—RHS 183 A.

*Petiole.*—2.2 cm in length, 3 mm in diameter.

*Petiole color.*—Upper side dark-red, RHS 185 A, lower side RHS 185 A.

#### INFLORESCENCE

*Flowering response.*—9 weeks after planting of rooted cuttings.

*Flowering season.*—Generally indeterminate, mainly from March to October, depending on light intensity.

Flower:

*Number of flowers per node.*—6 to 8, in various stages of development.

*Form of corolla.*—Single-type, 5 petals.

*Shape of corolla.*—Large, nearly round, with the petals overlapping, almost flat.

*Diameter of corolla.*—Average length: 75 mm. Average width: 75 mm. Depth of corolla: 10 mm.

*Shape of petals.*—Cordate, only weakly lobed at the top end, base attenuate. Top petal: 32 mm long, 52 mm wide. Lateral petals: 35 mm long, 40 mm wide. Lower petals 37 mm long, 48 mm wide. Texture: Smooth, velvety. Aspect: Flat.

*Color (general tonality from a distance of three meters).*—Light pink to nearly white, somewhat variable ground color, with a small purple eye.

*Color of upper surface.*—Light pink RHS 65 A to RHS 69 D, nearly white.

*Color of eye zone.*—Red-purple, RHS N 66 B.

*Color of lower surface.*—Mainly pale light pink, RHS 69 C.

*Spur.*—Downwardly curved, 6.7 mm long, 3 mm in diameter at the flower end; color deep pink, RHS 63 B.

*Peduncle color.*—Pink, RHS 63 C.

*Peduncle length.*—Approximately 50 mm.

*Flower bud.*—Ovoid shape, 24 mm in length, 16 mm in width; color RHS 69 B.

Reproductive organs:

*Stamens.*—5 in number, fused, upper surface color is RHS N57 C.

*Anthers.*—Fused, hooded.

*Pollen.*—Whitish-yellow, about RHS 8 D.

*Style and stigma.*—Five in number, very short, yellowish, about RHS 150 D.

*Ovary.*—5-celled, 5 mm long, surface green, RHS 139 B.

Disease/pest resistance/susceptibility: No observations to date.

I claim:

1. A new and distinct New Guinea *Impatiens* Plant named 'Fisupnic Lipink' as described and illustrated herein.

\* \* \* \* \*



UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : PP 16,118 P3  
DATED : November 15, 2005  
INVENTOR(S) : Birgit C. Hofmann

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:


Title page.

Item [73], Assignee, should read:

-- **Florfis AG**, Binningen (CH) --.

Signed and Sealed this

Eleventh Day of April, 2006

A handwritten signature in black ink on a light gray dotted background. The signature reads "Jon W. Dudas" in a cursive, stylized script. The "J" is large and loops around the "on". The "W" is written with two distinct peaks. The "D" is large and loops around the "udas".

JON W. DUDAS

*Director of the United States Patent and Trademark Office*