

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2002242676 B2**

(54) Title
Bilayer pharmaceutical tablet comprising telmisartan and a diuretic and preparation thereof

(51) International Patent Classification(s)
A61K 9/26 (2006.01) **A61K 31/4965** (2006.01)
A61K 9/20 (2006.01) **A61K 31/54** (2006.01)
A61K 9/24 (2006.01) **A61K 31/549** (2006.01)
A61K 31/341 (2006.01) **A61K 31/635** (2006.01)
A61K 31/402 (2006.01) **A61K 45/06** (2006.01)
A61K 31/4035 (2006.01) **A61K 47/04** (2006.01)
A61K 31/404 (2006.01) **A61K 47/16** (2006.01)
A61K 31/415 (2006.01) **A61K 47/26** (2006.01)
A61K 31/4184 (2006.01) **A61P 13/00** (2006.01)
A61K 31/495 (2006.01) **G01N 27/414** (2006.01)

(21) Application No: **2002242676** (22) Date of Filing: **2002.01.16**

(87) WIPO No: **WO03/059327**

(43) Publication Date: **2003.07.30**

(43) Publication Journal Date: **2003.09.04**

(44) Accepted Journal Date: **2008.05.29**

(71) Applicant(s)
Boehringer Ingelheim Pharma GmbH & Co. KG

(72) Inventor(s)
Friendl, Thomas;Schepky, Gottfried

(74) Agent / Attorney
Davies Collison Cave, Level 3 303 Coronation Drive Milton Business Centre, QLD, 4064

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 July 2003 (24.07.2003)

PCT

(10) International Publication Number
WO 03/059327 A1

- (51) International Patent Classification⁷: **A61K 9/20**, 31/635, 31/54, 31/495, 31/415
- (21) International Application Number: PCT/EP02/00395
- (22) International Filing Date: 16 January 2002 (16.01.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (71) Applicant (*for all designated States except US*): **BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG** [DE/DE]; 55216 Ingelheim am Rhein (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **FRIEDL, Thomas** [DE/DE]; Dr. Hans-Liebherr-Strasse 48/4, 88416 Ochsenhausen (DE). **SCHEPKY, Gottfried** [DE/DE]; Händelstrasse 26, 79312 Emmendingen (DE).
- (74) Agent: **BARZ, Peter**; Kaiserplatz 2, 80803 München (DE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 03/059327 A1

(54) Title: BILAYER PHARMACEUTICAL TABLET COMPRISING TELMISARTAN AND A DIURETIC AND PREPARATION THEREOF

(57) Abstract: A bilayer pharmaceutical tablet comprises a first layer formulated for immediate release of the angiotensin II receptor antagonist telmisartan from a dissolving tablet matrix which contains telmisartan in substantially amorphous form, and a second layer formulated for immediate release of a diuretic like hydrochlorothiazide from a fast disintegrating tablet matrix. A method of producing the bilayer tablet is also disclosed.

BILAYER PHARMACEUTICAL TABLET COMPRISING TELMISARTAN AND A DIURETIC AND PREPARATION THEREOF


Field of invention

The present invention relates to a bilayer pharmaceutical tablet formulation comprising the angiotensin II receptor antagonist telmisartan in combination with a diuretic such as hydrochlorothiazide (HCTZ). The present invention also provides a method of producing said bilayer tablet.

Background of the invention

INN Telmisartan is an angiotensin II receptor antagonist developed for the treatment of hypertension and other medical indications as disclosed in EP-A-502314.


Its chemical name is 4'-[2-n-propyl-4-methyl-6-(1-methylbenzimidazol-2-yl)-benzimidazol-1-ylmethyl]-biphenyl-2-carboxylic acid having the following structure:



Telmisartan is generally manufactured and supplied in the free acid form. It is characterized by its very poor solubility in aqueous systems at the physiological pH range of the gastro-intestinal tract of between pH 1 to 7. As disclosed in WO 00/43370, crystalline telmisartan exists in two polymorphic forms having different melting points. Under the influence of heat and humidity, the lower melting polymorph B transforms irreversibly into the higher melting polymorph A.

Hydrochlorothiazide (HCTZ) is a thiazide diuretic which is orally administered in the treatment of edema and hypertension.

The chemical name of HCTZ is 6-chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-sulfonamide-1,1-dioxide having the following structure



Combination therapy of telmisartan with a diuretic like HCTZ is expected to show synergistic therapeutic efficacy in the treatment of hypertension.

One or more embodiments of the present invention may provide a fixed dose combination drug comprising telmisartan and a diuretic such as HCTC, said combination drug displaying the required fast dissolution and immediate drug release profile combined with adequate stability.

Generally, a fixed-dose combination of drugs intended for immediate release is prepared by either making a powder mixture or a co-granulate of the two active ingredients with the necessary excipients, normally keeping the basic formulation of the corresponding mono-drug preparation and simply adding the second drug component.

With a combination of telmisartan and HCTZ, this approach was not feasible due to the incompatibility of HCTZ with basic compounds such as, e.g., meglumine (N-methyl-D-glucamine) which is a component of conventional telmisartan formulations, and the reduced dissolution rate of HCTZ from a dissolving matrix as compared with dissolution from a disintegrating tablet.

Several galenical approaches to overcome the incompatibility problem have been investigated. A classical approach is to coat the HCTZ particles in a fluidized-bed granulator with a polymer solution containing water soluble polymers like hydroxypropylcellulose, hydroxypropylmethylcellulose or polyvinylpyrrolidone, thereby

reducing the contact surface area of the HCTZ particles with the telmisartan formulation during mixing and compressing. Yet, by these means it was not possible to reduce the contact area of HCTZ with the telmisartan formulation in a compressed tablet to a degree sufficient to achieve the desired prolonged shelf life.

Furthermore, the dissolution rate of HCTZ from tablets comprising coated HCTZ in a telmisartan formulation was further reduced due to the gel-forming properties of the polymer.

Another approach was to produce separate film-coated tablets for telmisartan and HCTZ in such a size and shape that these could be filled into a capsule. By dividing the doses into two to four single small tablets for telmisartan and into one or two small tablets for HCTZ, a capsule of size 1 to 0 long could be filled. Yet, with this approach the drug dissolution rate of telmisartan was reduced compared to the single entities due to a lag-time effect of the large capsule shells. Furthermore, with regard to patients' compliance a zero long capsule is not deemed reliable.

Summary of the invention

In accordance with the present invention, it has now been found that the above-described problems associated with conventional approaches in the preparation of a fixed dose combination drug comprising telmisartan and a diuretic could be overcome by means of a bilayer pharmaceutical tablet comprising a first layer containing telmisartan in substantially amorphous form in a dissolving tablet matrix, and a second layer containing a diuretic in a disintegrating tablet matrix.

The bilayer tablet according to the present invention provides a largely pH-independent dissolution of the poorly water-soluble telmisartan, thereby facilitating dissolution of the drug at a physiological pH level, and also provides for immediate release of the diuretic from the fast disintegrating matrix. At the same time, the bilayer tablet structure overcomes the stability problem caused by the incompatibility of diuretics like HCTZ with basic constituents of the telmisartan formulation.

In a further aspect, the present invention relates to an improvement in bilayer tableting technology and provides a method of producing a bilayer pharmaceutical tablet comprising the steps of:

- (i) providing a first tablet layer composition by
 - a) preparing an aqueous solution of telmisartan, at least one basic agent and, optionally, a solubilizer and/or a crystallization retarder;
 - b) spray-drying said aqueous solution to obtain a spray-dried granulate;
 - c) mixing said spray-dried granulate with a water-soluble diluent to obtain a premix;
 - d) mixing said premix with a lubricant to obtain a final blend for the first tablet layer;
 - e) optionally, adding other excipients and/or adjuvants in any of steps a) to d);
- (ii) providing a second tablet layer composition by
 - f) mixing and/or granulating a diuretic with the constituents of a disintegrating tablet matrix and, optionally, further excipients and/or adjuvants;
 - g) admixing a lubricant to obtain a final blend for the second tablet layer;
- (iii) introducing the first or the second tablet layer composition in a tablet press;
- (iv) compressing said tablet layer composition to form a tablet layer;
- (v) introducing the other tablet layer composition into the tablet press; and
- (vi) compressing both tablet layer compositions to form a bilayer tablet.

Definitions

As used herein, the term "substantially amorphous" refers to a product comprising amorphous constituents in a proportion of at least 90%, preferably at least 95%, as determined by X-ray powder diffraction measurement.

The term "dissolving tablet matrix" refers to a pharmaceutical tablet base formulation having immediate release (fast dissolution) characteristics that readily dissolves in a physiological aqueous medium.

The term "diuretic" refers to thiazide and thiazide-analogue diuretics like hydrochlorothiazide (HCTZ), clopamide, xipamide or chlorotalidone, and any other diuretic suitable in the treatment of hypertension like, e.g., furosemide and piretanide, and combinations thereof with amiloride and triamteren.

The term "disintegrating tablet matrix" refers to a pharmaceutical tablet base formulation having immediate release characteristics that readily swells and disintegrates in a physiological aqueous medium.

Description of the preferred embodiments

The bilayer tablet according to the present invention comprises a first layer containing telmisartan in substantially amorphous form in a dissolving tablet matrix, and a second layer containing a diuretic in a disintegrating tablet matrix.

The active ingredient telmisartan is generally supplied in its free acid form, although pharmaceutically acceptable salts may also be used. Since during subsequent processing telmisartan is normally dissolved and transformed into a substantially amorphous form, its initial crystal morphology and particle size are of little importance for the physical and biopharmaceutical properties of the bilayer tablet formulation obtained. It is however preferred to remove agglomerates from the starting material, e.g. by sieving, in order to facilitate wetting and dissolution during further processing.

Substantially amorphous telmisartan may be produced by any suitable method known to those skilled in the art, for instance, by freeze drying of aqueous solutions, coating of carrier particles in a fluidized bed, and solvent deposition on sugar pellets or other carriers. Preferably, however, the substantially amorphous telmisartan is prepared by the specific spray-drying method described hereinafter.

The other active ingredient, i.e. the diuretic, is usually employed as a fine-crystalline powder, optionally in fine-milled, peg-milled or micronized form. For instance, the particle size distribution of hydrochlorothiazide, as determined by the method of laser light scattering in a dry dispersion system (Sympatec Helos/Rodos, focal length 100 mm) is preferably as follows:

- d₁₀ : ≤ 20 μm, preferably 2 to 10 μm
- d₅₀ : 5 to 50 μm, preferably 10 to 30 μm
- d₉₀ : 20 to 100 μm, preferably 40 to 80 μm

The bilayer tablet according to the present invention generally contains 10 to 160 mg, preferably 20 to 80 mg, of telmisartan and 6.25 to 50 mg, preferably 12.5 to 25 mg, of diuretic. Presently preferred forms are bilayer tablets comprising 40/12.5 mg, 80/12.5 mg and 80/25 mg of telmisartan and HCTZ, respectively.

The first tablet layer contains telmisartan in substantially amorphous form dispersed in a dissolving tablet matrix having immediate release (fast dissolution) characteristics. The dissolving tablet matrix may have acidic, neutral or basic properties, although a basic tablet matrix is preferred.

In such preferred embodiments, the dissolving matrix comprises a basic agent, a water-soluble diluent and, optionally, other excipients and adjuvants.

Specific examples of suitable basic agents are alkali metal hydroxides such as NaOH and KOH; basic amino acids such as arginine and lysine; and meglumine (N-methyl-D-glucamine), NaOH and meglumine being preferred.

Specific examples of suitable water-soluble diluents are carbohydrates such as monosaccharides like glucose; oligosaccharides like sucrose, anhydrous lactose and lactose monohydrate; and sugar alcohols like sorbitol, mannitol, dulcitol, ribitol and xylitol. Sorbitol is a preferred diluent.

The other excipients and/or adjuvants are, for instance, selected from binders, carriers, fillers, lubricants, flow control agents, crystallization retarders, solubilizers, coloring agents, pH control agents, surfactants and emulsifiers, specific examples of which are given below in connection with the second tablet layer composition. The excipients and/or adjuvants for the first tablet layer composition are preferably chosen such that a non-acidic, fast dissolving tablet matrix is obtained.

The first tablet layer composition generally comprises 3 to 50 wt.%, preferably 5 to 35 wt.%, of active ingredient; 0.25 to 20 wt.%, preferably 0.40 to 15 wt.%, of basic agent; and 30 to 95 wt.%, preferably 60 to 80 wt.% of water-soluble diluent.

Other (optional) constituents may, for instance, be chosen from one or more of the following excipients and/or adjuvants in the amounts indicated:

- 10 to 30 wt.%, preferably 15 to 25 wt.%, of binders, carriers and fillers, thereby replacing the water-soluble diluent;
- 0.1 to 5 wt.%, preferably 0.5 to 3 wt.%, of lubricants;

0.1 to 5 wt.%, preferably 0.3 to 2 wt.%, of flow control agents;
1 to 10 wt.%, preferably 2 to 8 wt.%, of crystallization retarders;
1 to 10 wt.%, preferably 2 to 8 wt.%, of solubilizers;
0.05 to 1.5 wt.%, preferably 0.1 to 0.8 wt.%, of coloring agents;
0.5 to 10 wt.%, preferably 2 to 8 wt.%, of pH control agents;
0.01 to 5 wt.%, preferably 0.05 to 1 wt.%, of surfactants and emulsifiers.

The second tablet layer composition contains a diuretic in a fast disintegrating tablet matrix. In a preferred embodiment, the disintegrating tablet matrix comprises a filler, a binder, a disintegrant and, optionally, other excipients and adjuvants.

The filler is preferably selected from anhydrous lactose, spray-dried lactose and lactose monohydrate.

The binder is selected from the group of dry binders and/or the group of wet granulation binders, depending on the manufacturing process chosen for the second tablet layer. Suitable dry binders are, e.g., cellulose powder and microcrystalline cellulose. Specific examples of wet granulation binders are corn starch, polyvinyl pyrrolidone (Povidon), vinylpyrrolidone-vinylacetate copolymer (Copovidone) and cellulose derivatives like hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose and hydroxypropylmethylcellulose.

Suitable disintegrants are, e.g., sodium starch glycolate, Croscopovidon, Croscarmellose, sodium carboxymethylcellulose and dried corn starch, sodium starch glycolate being preferred.

The other excipients and adjuvants, if used, are preferably selected from diluents and carriers such as cellulose powder, microcrystalline cellulose, cellulose derivatives like hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose and hydroxypropylmethylcellulose, dibasic calcium phosphate, corn starch, pregelatinized starch, polyvinyl pyrrolidone (Povidone) etc.; lubricants such as stearic acid, magnesium stearate, sodium stearyl fumarate, glycerol tribehenate, etc.; flow control agents such as colloidal silica, talc, etc.; crystallization retarders such as Povidone, etc.; solubilizers such as Pluronic, Povidone, etc.; coloring agents, including dyes and pigments such as Iron Oxide Red or Yellow, titanium dioxide, talc, etc.; pH control agents such as citric acid, tartaric acid, fumaric acid, sodium citrate, dibasic calcium phosphate, dibasic

sodium phosphate, etc.; surfactants and emulsifiers such as Pluronic, polyethylene glycols, sodium carboxymethyl cellulose, polyethoxylated and hydrogenated castor oil, etc.; and mixtures of two or more of these excipients and/or adjuvants.

The second tablet layer composition generally comprises 1.5 to 35 wt.%, preferably 2 to 15 wt.%, of active ingredient; 25 to 75 wt.%, preferably 35 to 65 wt.%, of filler; 10 to 40 wt.%, preferably 15 to 35 wt.%, of dry binder; 0.5 to 5 wt.%, preferably 1 to 4 wt.%, of wet granulation binder; and 1 to 10 wt.%, preferably 2 to 8 wt.%, of disintegrant. The other excipients and adjuvants are generally employed in the same amount as in the first tablet layer composition.

For preparing the bilayer tablet according to the present invention, the first and second tablet layer compositions may be compressed in the usual manner in a bilayer tablet press, e.g. a high-speed rotary press in a bilayer tableting mode. However, care should be taken not to employ an excessive compression force for the first tablet layer. Preferably, the ratio of the compression force applied during compression of the first tablet layer to the compression force applied during compression of both the first and second tablet layers is in the range of from 1:10 to 1:2. For instance, the first tablet layer may be compressed at moderate force of 4 to 8 kN, whereas the main compression of first plus second layer is performed at a force of 10 to 20 kN.

During bilayer tablet compression adequate bond formation between the two layers is achieved by virtue of distance attraction forces (intermolecular forces) and mechanical interlocking between the particles.

The bilayer tablets obtained release the active ingredients rapidly and in a largely pH-independent fashion, with complete release occurring within less than 60 min and release of the major fraction occurring within less than 15 min. The dissolution/-disintegration kinetics of the bilayer tablet may be controlled in different ways. For instance, both layers may dissolve/disintegrate simultaneously. Preferably, however, the second tablet layer containing the diuretic disintegrates first whereas the first tablet layer containing telmisartan dissolves in parallel or subsequently.

In accordance with the present invention, a substantially increased dissolution rate of the active ingredients and, in particular, of telmisartan is achieved. Normally, at least 70% and typically at least 90% of the drug load are dissolved after 30 min.

The bilayer tablets of the present invention tend to be slightly hygroscopic and are therefore preferably packaged using a moisture-proof packaging material such as aluminium foil blister packs, or polypropylene tubes and HDPE bottles which preferably contain a desiccant.

For optimum dissolution/disintegration and drug release properties, a specific method of producing the bilayer tablet according to the present invention has been developed which method comprises

- (i) providing a first tablet layer composition by
 - a) preparing an aqueous solution of telmisartan, at least one basic agent and, optionally, a solubilizer and/or a crystallization retarder;
 - b) spray-drying said aqueous solution to obtain a spray-dried granulate;
 - c) mixing said spray-dried granulate with a water-soluble diluent to obtain a premix;
 - d) mixing said premix with a lubricant to obtain a final blend for the first layer;
 - e) optionally, adding other excipients and/or adjuvants in any of steps a) to d);
- (ii) providing a second tablet layer composition by
 - f) mixing and /or granulating a diuretic with the constituents of a disintegrating tablet matrix and, optionally, further excipients and/or adjuvants;
 - g) admixing a lubricant to obtain a final blend for the second tablet layer;
- (iii) introducing the first or the second tablet layer composition into a tablet press;
- (iv) compressing said tablet layer composition to form a tablet layer;
- (v) introducing the other tablet layer composition into the tablet press; and
- (vi) compressing both tablet layer compositions to form a bilayer tablet.

In a preferred embodiment of this method, an aqueous alkaline solution of telmisartan is prepared by dissolving the active ingredient in purified water with the help of one or more basic agents like sodium hydroxide and meglumine. Optionally, a solubilizer and/or a recrystallization retarder may be added. The dry matter content of the starting aqueous solution is generally 10 to 40 wt.%, preferably 20 to 30 wt.%.

The aqueous solution is then spray-dried at room temperature or preferably at increased temperatures of, for instance, between 50 and 100°C in a co-current or countercurrent spray-drier at a spray pressure of, for instance, 1 to 4 bar. Generally

speaking, the spray-drying conditions are preferably chosen in such a manner that a spray-dried granulate having a residual humidity of ≤ 5 wt.%, preferably ≤ 3.5 wt.%, is obtained in the separation cyclone. To that end, the outlet air temperature of the spray-drier is preferably kept at a value of between about 80 and 90°C while the other process parameters such as spray pressure, spraying rate, inlet air temperature, etc. are adjusted accordingly.

The spray-dried granulate obtained is preferably a fine powder having the following particle size distribution:

- d_{10} : $\leq 20 \mu\text{m}$, preferably $\leq 10 \mu\text{m}$
- d_{50} : $\leq 80 \mu\text{m}$, preferably 20 to 55 μm
- d_{90} : $\leq 350 \mu\text{m}$, preferably 50 to 150 μm

After spray-drying, the active ingredient (telmisartan) as well as the excipients contained in the spray-dried granulate are in a substantially amorphous state with no crystallinity being detectable. From a physical point of view, the spray-dried granulate is a solidified solution or glass having a glass transition temperature T_g of preferably $> 50^\circ\text{C}$, more preferably $> 80^\circ\text{C}$.

Based on 100 parts by weight of active ingredient (telmisartan), the spray-dried granulate preferably contains 5 to 200 parts by weight of basic agent and, optionally, solubilizer and/or crystallization retarder.

The water-soluble diluent is generally employed in an amount of 30 to 95 wt.%, preferably 60 to 80 wt.%, based on the weight of the first tablet layer composition.

The lubricant is generally added to the premix in an amount of 0.1 to 5 wt.%, preferably 0.3 to 2 wt.%, based on the weight of the first tablet layer composition.

Mixing is carried out in two stages, i.e. in a first mixing step the spray-dried granulate and the diluent are admixed using, e.g., a high-shear mixer or a free-fall blender, and in a second mixing step the lubricant is blended with the premix, preferably also under conditions of high shear. The method of the invention is however not limited to these mixing procedures and, generally, alternative mixing procedures may be employed in steps c), d), and also in the subsequent steps f) and g), such as, e.g., container mixing with intermediate screening.

For direct compression, the second tablet layer composition may be prepared by dry-mixing the constituent components, e.g. by means of a high-intensity mixer or a free-fall blender. Alternatively and preferably, the second tablet layer composition is prepared using a wet granulation technique wherein an aqueous solution of a wet granulation binder is added to a premix and subsequently the wet granulate obtained is dried, e.g. in a fluidized-bed dryer or drying chamber. The dried mixture is screened and then a lubricant is admixed, e.g. using a tumbling mixer or free-fall blender, whereafter the composition is ready for compression.

For production of the bilayer tablet according to the present invention, the first and second tablet layer compositions are compressed in a bilayer tablet press, e.g. a rotary press in the bilayer tableting mode, in the manner described above. In order to avoid any cross-contamination between the first and second tablet layers (which could lead to decomposition of HTCZ), any granulate residues have to be carefully removed during tableting by intense suction of the die table within the tableting chamber.

In order to further illustrate the present invention, the following non-limiting examples are given.

Example 1

Constituents		mg / 1.684 mg SD granulate	volatile constituent	kg / batch
(01)	Telmisartan	1.000		45.000
(02)	Sodium hydroxide	0.084		3.780
(03)	Povidone K 25	0.300		13.500
(04)	Meglumine	0.300		13.500
(05)	Purified water		5.000	(225.000)
		1.684	5.000	75.780

Manufacturing:**1. Spray solution**

225.000 kg of purified water are measured into a suitable stainless steel vessel at a temperature of between 20-40°C. In sequence, 3.780 of kg sodium hydroxide, 45.000 kg of telmisartan (mixture of polymorph A and B), 13.500 kg of Povidone K 25 and 13.500 kg of meglumine are dissolved in the purified water under intensive stirring until a virtually clear, slightly yellowish, alkaline solution is obtained.

2. Spray drying

The solution is sprayed into a suitable spray dryer, e.g. a Niro P 6.3 equipped with Schlick atomizing nozzles of 1.0 mm diameter, with a flow-through heating coil connected upstream of the dryer, and dried to give a white to off-white fine granulate. The spray mode is counter-current at a spray-pressure of about 3 bar, an inlet air temperature of about 125°C and a spray rate of about 11 kg/h, thus resulting in an outlet air temperature of about 85°C. The temperature of the flow through heating coil water bath is set at a temperature of about 80°C.

3. Protective Screening

The dry granulate powder is screened through a screen of 0.5 mm mesh size, e.g. using a Vibra Sieve machine.

The resulting amorphous telmisartan spray-dried granulate may be further processed to telmisartan mono-tablets or the first layer of the said bilayer tablet composition.

Example 2

	Constituents	mg/tablet 1st layer	mg/SD granulate	mg/tablet 2nd layer
(01)	Telmisartan SD granulate	67.360		
	consisting of (02) to (06):			
(02)	Telmisartan		40.000	
(03)	Sodium hydroxide		3.360	
(04)	Polyvidone (Kollidon 25)		12.000	
(05)	Meglumine		12.000	
(06)	Purified water		264.000*	
(07)	Sorbitol P/6	168.640		
(08)	Magnesium stearate, screened	4.000		1.000
(09)	Hydrochlorothiazide			12.500
(10)	Microcrystalline cellulose (Avicel PH 101)			64.000
(11)	Red iron oxide			0.330
(12)	Sodium starch glycolate			4.000
(13)	Lactose monohydrate fine, screened			112.170
(14)	Maize starch, dried at 45 °C			6.000
		240.000	67.360	200.000

* 200 mg in SD granulate, 64 mg in granulation liquid of HCTZ granulate

Manufacturing:

1. Final blend A

168.640 kg of sorbitol are mixed with 67.360 kg of telmisartan spray dried granulate in a suitable high shear mixer, e.g. Diosna P 600, for 4 minutes using both impeller and chopper. Next 4.0 kg of magnesium stearate are added to the resulting pre-mix and admixed in the high shear mixer for further 30 seconds.

2. Final blend B

9.000 kg of purified water of about 70°C are transferred to a suitable mixing vessel, 6.000 kg of maize starch, dried at 45°C, are suspended in the water. This suspension is stirred into 55.000 kg of purified water of about 90°C using e.g. an Ekato stirrer.

Next, 112.170 kg of lactose monohydrate, 12.500 kg of hydrochlorothiazide, 64.000 kg of microcrystalline cellulose (Avicel PH 101), 0.330 kg of red iron oxide and 4.000 kg of sodium starch glycolate are mixed in a suitable high shear granulator, e.g. Diosna P 600, until homogeneous, and moistened with 70.000 kg of the above-prepared aqueous granulating liquid.

Process parameters for wet granulation:

Process step	Duration (min)	Impeller (setting)	Chopper (setting)
Pre-mixing	3	1	1
Moistening	2	1	1
Wet mixing	4	2	2
Emptying	About 0.5	1	0

After moistening, the resulting wet granulate is dried in a suitable fluid bed dryer, e.g. Glatt WSG 120 at an inlet air temperature of 100°C, an inlet air flow of 2000-3000 m³/h until a product temperature of about 55°C is reached.

The dry granulate is screened to reduce the particle size using a suitable screening machine, e.g. a Comil screen machine equipped with a rasp screen of 2 mm mesh size. Finally 1.000 kg of prescreened magnesium stearate are admixed to the screened granulate material and mixed in a suitable tumbling mixer, e.g. a Lerner rotating spike mixer, for 100 revolutions at a speed of 8-10 rpm.

3. Bilayer tablet compression

Using a suitable rotary tablet press, 240 kg of the final blend (A) and 200 kg of the final blend (B) are compressed into bilayer tablets. The target weight for the first layer is 240 mg, the target weight for the second layer is 200 mg.

Process parameters for tableting:

Tablet press	Fette 3090	
Tabletting speed	100.000 (80.000 – 120.000) tabl./h	
Stirrer blade speed:	1st layer about 30 rpm	2nd layer about 75 rpm
Compression force	5 (4 – 6) KN	12 (10 – 14) KN

As a rule, the tablet hardness is adjusted by variation of the main compression force of the second layer.

The resulting bilayer tablets have the following characteristics:

Shape / diameter	oval, both faces convex / 14 x 6.8 mm
Colour	first layer: white to off-white second layer: red
Weight	440 mg (total) 240 mg (layer 1: with telmisartan) 200 mg (layer 2: with hydrochlorothiazide)
Thickness	about 5.2 mm
Hardness	about 120 N
Disintegration time	NMT 15 min (total)

Example 3

	Constituents	mg/tablet 1st layer	mg/SD granulate	mg/tablet 2nd layer
(01)	Telmisartan SD granulate	67.360		
	consisting of (02) to (06):			
(02)	Telmisartan		40.000	
(03)	Sodium hydroxide		3.360	
(04)	Polyvidone (Kollidon 25)		12.000	
(05)	Meglumine		12.000	
(06)	Purified water		(200.000)	
(07)	Sorbitol P/6	168.640		
(08)	Magnesium stearate, screened	4.000		1.000
(09)	Hydrochlorothiazide			25.000
(10)	Microcrystalline cellulose (Avicel PH 101)			64.000
(11)	Yellow iron oxide			0.330
(12)	Sodium starch glycolate			4.000
(13)	Lactose monohydrate fine, screened			105.67
		240.000	67.360	200.000

Manufacturing:

Manufacturing is carried out as in Example 2. Instead of the wet granulation process described in Example 2, the second layer composition is manufactured by dry mixing of (09) to (13) in a suitable free fall blender, e.g. a 1 m³ container mixer, for 200 revolutions at a speed of 10 rpm. Then, (08) is admixed to the main mixture for further 50 revolutions in the container mixer. In order to achieve a homogenous distribution of the color pigment, an additional premix with yellow iron oxide and a portion of the microcrystalline cellulose, e.g. 2.000 kg, which is screened through an 0.8 mm mesh screen manually before transfer to the main mixture, may be performed. The resulting bilayer tablets display virtually the same physical characteristics as described in example 2, except for the color.

Example 4

Composition of Telmisartan/Hydrochlorothiazide Bilayer Tablets (mg per tablet):

Ingredient	40/12.5 mg	80/12.5 mg
Telmisartan layer		
Telmisartan	40.000	80.000
Sodium hydroxide	3.360	6.720
Povidone	12.000	24.000
Meglumine	12.000	24.000
Purified water*	(200.000)	(400.000)
Sorbitol	168.640	337.280
Magnesium stearate	4.000	8.000
Total telmisartan layer	240.000	480.000
Hydrochlorothiazide layer		
Hydrochlorothiazide	12.500	12.500
Lactose monohydrate	112.170	112.170
Microcrystalline Cellulose	64.000	64.000
Corn starch	6.000	6.000
Red iron oxide	0.330	0.330
Sodium starch glycolate	4.000	4.000
Purified water*	(64.000)	(64.000)
Magnesium stearate	1.000	1.000
Total HCTZ layer	200.000	200.000
Total tablet weight	440.000	680.000

*Does not appear in final product

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A bilayer pharmaceutical tablet comprising a first layer containing telmisartan in substantially amorphous form in a dissolving tablet matrix, and a second layer containing a diuretic in a disintegrating tablet matrix.
2. A bilayer pharmaceutical tablet as claimed in claim 1 wherein the diuretic is selected from at least one of hydrochlorothiazide, furosemide, chlorotalidone, piretanide and amiloride.
3. A bilayer pharmaceutical tablet as claimed in claim 2 wherein the diuretic is hydrochlorothiazide.
4. A bilayer pharmaceutical tablet as claimed in any one of claims 1-3 wherein the dissolving tablet matrix has immediate release characteristics.
5. A bilayer pharmaceutical tablet as claimed in any one of claims 1-4 wherein the dissolving tablet matrix comprises a basic agent, a water-soluble diluent and, optionally, other excipients and adjuvants.
6. A bilayer pharmaceutical tablet as claimed in claim 5 where the basic agent is selected from alkali metal hydroxides, basic amino acids and meglumine.
7. A bilayer pharmaceutical tablet as claimed in claims 5 or 6 wherein the water-soluble diluent is selected from carbohydrates such as monosaccharides like glucose; oligosaccharides like sucrose and lactose; and sugar alcohols like sorbitol, mannitol, dulcitol, ribitol and xylitol.
8. A bilayer pharmaceutical tablet as claimed in any one of claims 5-7 wherein the other excipients and adjuvants are selected from binders, carriers, fillers, lubricants, flow control agents, crystallization retarders, solubilizers, coloring agents, pH control agents, surfactants and emulsifiers.

9. A bilayer pharmaceutical tablet as claimed in any one of claims 1-8 wherein the first tablet layer has been produced by spray-drying an aqueous solution comprising telmisartan and a basic agent to obtain a spray-dried granulate, mixing said spray-dried granulate with a water-soluble diluent to obtain a premix, mixing said premix with a lubricant to obtain a final blend and compressing the final blend to form the first tablet layer.
10. A bilayer pharmaceutical tablet as claimed in any one of claims 1-10 wherein the disintegrating tablet matrix comprises a filler, a binder, a disintegrant and, optionally, other excipients and adjuvants.
11. A bilayer pharmaceutical tablet as claimed in claim 10 wherein the other excipients and adjuvants are selected from carriers, diluents, lubricants, flow control agents, solubilizers, coloring agents, pH control agents, surfactants and emulsifiers.
12. A bilayer pharmaceutical tablet as claimed in any one of claims 1-11 containing 10 to 160 mg, preferably 20 to 80 mg, of telmisartan and 6.25 to 50 mg, preferably 12.5 to 25 mg, of diuretic.
13. A bilayer pharmaceutical tablet as claimed in any one of claims 1-12 packaged in a moisture proof packaging material such as aluminium foil blister packs, or polypropylene tubes and HDPE bottles.
14. A method of producing a bilayer pharmaceutical tablet comprising the steps of:
- (i) providing a first tablet layer composition by
 - a) preparing an aqueous solution of telmisartan, at least one basic agent and, optionally, a solubilizer and/or a crystallization retarder;
 - b) spray-drying said aqueous solution to obtain a spray-dried granulate;
 - c) mixing said spray-dried granulate with a water-soluble diluent to obtain a premix;
 - d) mixing said premix with a lubricant to obtain a final blend for the first tablet layer;

- e) optionally, adding other excipients and/or adjuvants in any of steps a) to d);
- (ii) providing a second tablet layer composition by
- f) mixing and/or granulating a diuretic with the constituents of a disintegrating tablet matrix and, optionally, further excipients and/or adjuvants;
- g) admixing a lubricant to obtain a final blend for the second tablet layer;
- (iii) introducing the first or the second tablet layer composition into a tablet press;
- (iv) compressing said tablet layer composition to form a tablet layer;
- (v) introducing the other tablet layer composition into the tablet press; and
- (vi) compressing both tablet layer compositions to form a bilayer tablet.

15. A method as claimed in claim 14 wherein spray-drying in step b) is carried out under conditions so as to obtain a spray-dried granulate having a residual humidity of ≤ 5 wt.%, preferably ≤ 3.5 wt.%.

16. A method as claimed in claims 14 or 15 wherein spray-drying in step b) is carried out at an outlet air temperature of the spray-drier of between about 80 and 90°C.

17. A method as claimed in any one of claims 14-16 wherein mixing in any of steps c), d), f) and g) is carried out in a high shear mixer or a free-fall blender.

18. A method as claimed in any one of claims 14-17 wherein mixing in step f) is carried out under conditions of dry-mixing or, preferably, under wet granulation conditions.

19. A method as claimed in any one of claims 14-18 wherein the ratio of the compression force applied during compression of the first tablet layer to the compression force applied during compression of both the first and second tablet layers is in the range of from 1:10 to 1:2.

2002242676 08 Apr 2008

- 21 -

20. A bilayer pharmaceutical tablet prepared by a method according to any one of claims 14-19.

21. A bilayer pharmaceutical tablet according to claim 1 or 20, or a method according to claim 14, substantially as hereinbefore described and/or exemplified.