



US00PP22746P2

(12) **United States Plant Patent**
Graff

(10) **Patent No.:** **US PP22,746 P2**

(45) **Date of Patent:** **May 22, 2012**

(54) **HIBISCUS PLANT NAMED 'ARIONICUS'**

(50) Latin Name: *Hibiscus rosa-sinensis*
Varietal Denomination: **Arionicus**

(75) Inventor: **Poul Graff**, Sabro (DK)

(73) Assignee: **Graff Breeding A/S**, Sabro (DK)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/927,926**

(22) Filed: **Nov. 30, 2010**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./257**

(58) **Field of Classification Search** **Plt./257**
See application file for complete search history.

Primary Examiner — Annette Para

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Hibiscus* plant named 'Arionicus', characterized by its upright to somewhat outwardly spreading plant form; dense and bushy plant habit; glossy dark green-colored leaves; freely, uniform and continuous flowering habit; large red-colored flowers with dark red purple-colored centers; good flower longevity; and good garden performance.

2 Drawing Sheets

1

Botanical designation: *Hibiscus rosa-sinensis*.
Cultivar denomination: 'ARIONICUS'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Hibiscus* plant, botanically known as *Hibiscus rosa-sinensis* and hereinafter referred to by the name 'Arionicus'.

The new *Hibiscus* plant is a product of a planned breeding program conducted by the Inventor in Sabro, Denmark. The objective of the breeding program is to create new strong *Hibiscus* plants with attractive and long-lasting flowers.

The new *Hibiscus* plant originated from a cross-pollination in December, 2007 in Sabro, Denmark of *Hibiscus rosa-sinensis* 'Camaro', not patented, as the female, or seed, parent with a proprietary selection of *Hibiscus rosa-sinensis* identified as code number GB 2006-4715, not patented, as the male, or pollen, parent. The new *Hibiscus* plant was discovered and selected by the Inventor as a flowering plant within the progeny of the stated cross-pollination in a controlled greenhouse environment in Sabro, Denmark in July, 2008.

Asexual reproduction of the new *Hibiscus* plant by vegetative terminal cuttings in a controlled greenhouse environment in Sabro, Denmark since September, 2008, has shown that the unique features of this new *Hibiscus* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Hibiscus* have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Arionicus'. These characteristics in combination distinguish 'Arionicus' as a new and distinct cultivar of *Hibiscus* plant:

1. Upright to somewhat outwardly spreading plant form.
2. Dense and bushy plant habit.
3. Glossy dark green-colored leaves.

2

4. Freely, uniform and continuous flowering habit.

5. Large red-colored flowers with dark red purple-colored centers.

6. Good flower longevity.

7. Good garden performance.

Plants of the new *Hibiscus* can be compared to plants of the female parent, 'Camaro'. Plants of the new *Hibiscus* differ primarily from plants of 'Camaro' in flower size and longevity as plants of the new *Hibiscus* have larger and longer-lasting flowers than plants of 'Camaro'. In addition, leaves of plants of the new *Hibiscus* have crenate margins whereas leaves of plants 'Camaro' have serrate margins.

Plants of the new *Hibiscus* can be compared to plants of the male parent selection. Plants of the new *Hibiscus* differ primarily from plants of the male parent selection in flower size and color as plants of the male parent selection have small dark pink-colored flowers.

Plants of the new *Hibiscus* can be compared to plants of the *Hibiscus rosa-sinensis* 'Cairo Red', not patented. In side-by-side comparisons conducted in Sabro, Denmark, plants of the new *Hibiscus* differed from plants of 'Cairo Red' in the following characteristics:

1. Flowers of plants of the new *Hibiscus* were brighter red in color than flowers of plants of 'Cairo Red'.
2. Flowers of plants of the new *Hibiscus* were longer lasting than flowers of plants of 'Cairo Red'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Hibiscus* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Hibiscus* plant.

The photograph on the first sheet comprises a side perspective view of typical flowering plants of 'Arionicus' grown in a container.

The photograph on the second sheet are close-up views of typical developing flower buds and a fully opened flower of 'Arionicus'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown during the summer in 13-cm containers in a glass-covered greenhouse in Sabro, Denmark and under conditions which closely approximate commercial *Hibiscus* production. During the production of the plants, day temperatures ranged from 20° C. to 25° C., night temperatures ranged from 19° C. to 21° C. and light levels ranged from 40 to 50 klux. Plants were pinched one time about seven weeks after planting and were 22 weeks old when the photographs and the description were taken. In the description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used. Botanical classification: *Hibiscus rosa-sinensis* 'Arionicus'. Parentage:

Female, or seed, parent.—*Hibiscus rosa-sinensis* 'Camaro', not patented.

Male or pollen parent.—Proprietary selection of *Hibiscus rosa-sinensis* identified as code number GB 2006-4715, not patented.

Propagation:

Type.—By vegetative terminal cuttings.

Time to initiate roots, summer.—About three weeks at temperatures of 24° C.

Time to initiate roots, winter.—About four weeks at temperatures of 24° C.

Time to produce a rooted young plant, summer.—About seven weeks at temperatures of 24° C.

Time to produce a rooted young plant, winter.—About eight weeks at temperatures of 24° C.

Root description.—Medium in thickness, fleshy; color, close to 158A.

Rooting habit.—Freely branching; dense.

Plant description:

Plant form and growth habit.—Upright to somewhat outwardly spreading plant form; moderately vigorous growth habit.

Branching habit.—Freely branching, usually about four to six lateral branches develop; pinching enhances lateral branch development; dense and bushy plant habit.

Plant height.—About 35 cm to 55 cm.

Plant diameter (area of spread).—About 30 cm to 50 cm.

Lateral branch description:

Length.—About 15 cm to 25 cm.

Diameter.—About 3 mm to 6 mm.

Internode length.—About 1 cm to 5 cm.

Strength.—Strong.

Texture.—Woody.

Color.—Close to N199A and 197A.

Foliage description:

Arrangement.—Alternate, simple.

Length.—About 8 cm to 10 cm.

Width.—About 5 cm to 7 cm.

Shape.—Broadly cordate.

Apex.—Cuspidate to acuminate.

Base.—Cordate.

Margin.—Crenate.

Texture, upper and lower surfaces.—Smooth, glabrous.

Luster, upper surface.—Glossy.

Luster, lower surface.—Slightly glossy.

Venation pattern.—Pinnate; arcuate.

Color.—Developing leaves, upper surface: Close to 146B. Developing leaves, lower surface: Close to 143B. Fully expanded leaves, upper surface: Close to 147A; venation, close to 138B. Fully expanded leaves, lower surface: Close to 137B; venation, close to 138B.

Petiole.—Length: About 3 cm to 5 cm. Diameter: About 2 mm to 3 mm. Texture, upper surface: Pubescent. Texture, lower surface: Smooth, glabrous. Color, upper and lower surfaces: Close to N199A.

Flower description:

Flower shape and arrangement.—Rounded flowers with imbricate petals arranged singly at terminal leaf axils; uniform, continuous and freely flowering habit with numerous flower buds and/or open flowers per plant at one time; flowers face mostly upright to outwardly.

Fragrance.—None detected.

Flower longevity.—Good flower longevity, flowers last for about two to three days; flowers persistent.

Natural flowering season.—Usually spring and summer or during periods of warm weather; plants flower year-round in the greenhouse; plants begin flowering about 13 to 15 weeks after pinching.

Flower diameter.—About 15 cm to 19 cm.

Flower length (height).—About 8 cm to 10 cm.

Flower bud.—Rate of opening: Flowers buds open in about two days. Length: About 5 cm to 7 cm. Diameter: About 1.5 cm to 2.5 cm. Shape: Ovate to lanceolate. Color: Close to N34A.

Petals.—Arrangement: Corolla consists of five imbricate petals in a single whorl. Length: About 10 cm to 12 cm. Width: About 8 cm to 9 cm. Shape: Roughly spatulate. Apex: Rounded. Base: Attenuate. Margin: Entire; undulate. Texture, upper surface: Glabrous; velvety. Texture, lower surface: Glabrous; satiny. Color: When opening, upper surface: Close to N44A or 45B; at the base, close to 59A. When opening, lower surface: Close to 45A. Fully opened, upper surface: Close to N44A or 45B; at the base, close to 59A. Fully opened, lower surface: Close to 45B; with development, color becomes closer to 45C.

Sepals.—Appearance: Five sepals fused into a campanulate-shaped calyx. Length: About 2.5 cm to 3 cm. Width: About 1.3 cm. Shape: Lanceolate. Apex: Acuminate. Margin: Entire. Texture, upper surface: Rough. Texture, lower surface: Smooth. Color, upper surface: Close to 144A to 144C. Color, lower surface: Close to 144B.

Peduncles.—Length: About 2 cm to 4 cm. Diameter: About 2 mm to 3 mm. Angle: Mostly upright. Strength: Strong. Texture: Smooth, glabrous. Color: Close to 144A.

Reproductive organs.—Androecium: Stamen number: Numerous, about 100. Filament length: About 2.5 mm. Filament color: Close to 55A. Anther shape: Rounded to ovate. Anther length: About 1 mm to 2 mm. Anther color: Close to 22B. Amount of pollen: Moderate to abundant. Pollen color: Close to 17A. Gynoecium: Pistil length: About 9 cm. Style length: About 8 cm. Style texture: Smooth, waxy. Style color:

Towards the apex, close to 55A; mid-section, close to 46B; towards the base, close to 53A. Stigma appearance: Five-parted, rounded. Stigma color: Close to 46B. Ovary color: Close to 145A and 2C.

Seeds.—Quantity produced per flower: None to ten. Length: About 3 mm to 4 mm. Diameter: About 2 mm to 3 mm. Color: Close to N200A.

Garden performance: Plants of the new *Hibiscus* have been observed to have good garden performance and to tolerate rain, wind and temperatures ranging from about 1° C. to about 30° C.

Pathogen/pest resistance: Plants of the new *Hibiscus* have not been observed to be resistant to pathogens and pests common to *Hibiscus*.

It is claimed:

1. A new and distinct *Hibiscus* plant named 'Arionicus' as illustrated and described.

* * * * *



