



US 20130288929A1

(19) **United States**(12) **Patent Application Publication**
Hansen et al.(10) **Pub. No.: US 2013/0288929 A1**(43) **Pub. Date: Oct. 31, 2013**(54) **METHOD FOR MAKING AN ENRICHED LIBRARY**(86) PCT No.: **PCT/EP2011/065117**


§ 371 (c)(1),

(2), (4) Date: **Sep. 21, 2012**(75) Inventors: **Nils Jakob Vest Hansen**, Copenhagen (DK); **Allan Beck Christensen**, Copenhagen (DK); **Leif Kongskov Larsen**, Copenhagen (DK); **Frank Abildgaard Sløk**, Copenhagen (DK); **Lars Kolster Petersen**, Copenhagen (DK); **Judith Rasmussen-Dietvorst**, Copenhagen (DK); **Peter Blakskjær**, Copenhagen (DK); **Tara Heitner Hansen**, Copenhagen (DK); **Johan Holmkvist**, Copenhagen (DK)(30) **Foreign Application Priority Data**

Sep. 27, 2010 (EP) 10180435.9


Publication Classification(51) **Int. Cl.**
C12N 15/10 (2006.01)(52) **U.S. Cl.**
CPC **C12N 15/1075** (2013.01)
USPC **506/26**(73) Assignee: **Vipergen**, Copenhagen (DK)(21) Appl. No.: **13/636,668**(22) PCT Filed: **Sep. 1, 2011**(57) **ABSTRACT**

A method for making an enriched library comprising specific nucleic acid sequence information allowing to identifying at least one binding entity that binds to at least one target wherein the specific binding entity has been present in an in vitro display library.




Illustrative example of the principle in ECC

Figure 1



Illustrative example of the principle in ECC

Figure 2



Illustrative example of the principle in ECC.using emulsion PCR.

Figure 3





Figure 4

The Low range DNA marker (Fermentas) was loaded in the lane M. In lane 1, PCR product derived from the binding reaction without pre-incubation with biotin was loaded. In lane 2, PCR product derived from the binding reaction with pre-incubation with biotin was loaded.

Figure 5

Condition	Spiked (1:1000)	Preincubation w biotin	# Total counts	# Biotin_yR counts	% Biotin_yR	Fold enrichment
A	Yes	No	6339	2385	37.62	376
A	Yes	No	7032	2250	32.00	320
B	Yes	Yes	6897	21	0.30	3
B	Yes	Yes	7195	52	0.72	7
C	No	No	5750	1	0.02	0
C	No	No	6425	3	0.05	0

Results from DNA sequencing

Figure 6




Figure 7

Sample	Library	BSB_yr_spike	Target	Target conc (nM)	Target pre-incubated w/ 1 uM BSB	Emulsion force (rpm)	Dissociation time (min)	Total number of reads	Higest counts	2nd higest counts	Counts for BSB_yr	Frequency (BSB_yr)	Fold enrichment
1	Lib01651	0.000005	CA II	10		5500	2	7808	54	24	54	0.0069	1383
2	Lib01651	0.000005	CA II	10	1	5500	2	7970	17	15	0	<0.0001	NA
3	Lib01651	0.000005	CA II	10		5500	30	10470	15	14	0	<0.0001	NA
4	Lib01651	0.000005	CA II	10	1	5500	30	7053	18	15	0	<0.0001	NA
5	Lib01651	0.000005	CA II	10		NA	NA	15864	15	12	0	<0.0001	NA

Results from DNA sequencing

METHOD FOR MAKING AN ENRICHED LIBRARY

FIELD OF THE INVENTION

[0001] The present invention relates to a method for making an enriched library comprising specific nucleic acid sequence information allowing to identifying at least one binding entity that binds to at least one target wherein the specific binding entity has been present in an in vitro display library.

BACKGROUND

[0002] Display technologies have been developed to combine information storage and amplification capabilities of nucleic acids with the functional activities of other compound. Display technologies rely on an association between a functional binding entity (i.e. phenotype) and a nucleic acid sequence informative (genotype) about the structure of the binding entity. Note: Nucleic acid aptamer technology is considered a display technology although a special case as the pheno- and genotype consist of the same molecule (DNA or RNA).

[0003] An advantage of such methods is that very large libraries can be constructed and probed for a desired activity of the functional binding entities. Library members having the desired activity can then be partitioned from library members not having the desired activity, thus creating an enriched library with a higher fraction of members having the desired activity. This process is called selection or enrichment. Some display technologies allows for rounds of selections, where the enriched library from one round is amplified and used to prepare a new enriched display library and used in a next round of selection and so forth. The structures of the library members in the enriched library can then be identified by their cognate nucleic acid sequence, thus allowing identification even from minute amounts of material.

[0004] Herein relevant libraries may according to the art be termed “in vitro display libraries”.

[0005] The term “in vitro display library” shall herein be understood according to the art—i.e. as a library comprising numerous different binding entities wherein each binding entity is attached to a nucleic acid molecule and the nucleic acid molecule comprises specific nucleic acid sequence information allowing to identify the binding entity—i.e. once one knows the specific nucleic acid sequence information of the nucleic acid molecule one directly knows the structure of the specific binding entity attached to the nucleic acid molecule—the structure of the binding entity (i.e. phenotype) attached to the nucleic acid molecule (genotype) is herein termed B-structure.

[0006] The prior art describes a number of different methods to make such in vitro display libraries—herein suitable examples include e.g. EP1809743B1 (Viperger), EP1402024B1 (Nuevolution), EP1423400B1 (David Liu), Nature Chem. Biol. (2009), 5:647-654 (Clark), WO 00/23458 (Harbury), Nature Methods (2006), 3(7), 561-570, 2006 (Miller), Nat. Biotechnol. 2004; 22, 568-574 (Melkko), Nature. (1990); 346(6287), 818-822 (Ellington), or Proc Natl Acad Sci USA (1997). 94 (23): 12297-302 (Roberts), WO06053571A2 (Rasmussen).

[0007] As described in e.g. above mentioned prior art—one can today make in vitro display libraries comprising very many (e.g. 10^{15}) specific binding entities (e.g. 10^{15} different chemical compounds).

[0008] In view of this—it is evident that it would be very interesting to be able to improve the selection/enrichment step of such libraries to make an enriched library—e.g. to more efficient be able to identify the structure of a specific binding entity (e.g. a chemical compound) that binds to a target of interest (e.g. a medical important receptor molecule).

[0009] In FIG. 3 herein is shown an example of the in vitro display technology as described in EP1809743B1 (Viperger)—as can be seen in this FIG. 3—the selection step of this example is performed by immobilizing the target (e.g. a receptor) to a solid surface (e.g. a bead or a glass plate).

[0010] Without being limited to theory—to our knowledge, the example in FIG. 3 herein may be seen as an example of herein relevant in vitro display technology prior art (e.g. above mentioned prior art)—i.e. to our knowledge the selection for suitable binding entities present within in vitro display libraries are in the prior art generally done by immobilizing the target (e.g. a receptor) to a solid support (e.g. a glass plate, a column, a bead, a nitrocellulose filter, a cell etc) before or after the display library binding event. Non-binders and low affinity binders are typically washed away, whereas the population enriched for binders are recovered from the solid support.

[0011] In prior art in vitro compartmentalization (IVC) have been described employed in technologies utilizing phenotype and genotype linkage for interrogating libraries. These prior art technologies can be divided into two groups: a) IVC utilized for facilitating establishing correct phenotype and genotype linkage, which allows for selection of function (e.g. specific target binding) later (post compartment disruption), and b) IVC for facilitating establishing correct phenotype and genotype linkage based on an activity of the phenotype inside the compartment, i.e. in a compartment a gene is transcribed and translated and the resulting protein's function inside the compartment is used directly or indirectly for sorting, survival or amplification.

[0012] In other words herein relevant so-called IVC prior art technologies—may be described as a: group a)—wherein the phenotype activity is interrogated AFTER the compartmentalized step; or group b)—wherein the phenotype activity is interrogated DURING the compartmentalized step.

[0013] Examples of IVC prior art belonging to group a):

[0014] Bertschinger et al, (2007) Protein Engineering, Design & Selection vol. 20 no. 2 pp. 57-68;

[0015] Miller O J, Bernath K, Agresti J J, Amitai G, Kelly B T, Mastrobattista E, Taly V, Magdassi S, Tawfik D S, Griffiths A D. Directed evolution by in vitro compartmentalization. Nat Methods. 2006 July; 3(7):561-70;

[0016] Doi, N. and Yanagawa, H. (1999) FEBS Lett., 457, 227-230; and Yonezawa, M., Doi, N., Kawahashi, Y., Higashinakagawa, T. and Yanagawa, H. (2003) Nucleic Acids Res., 31, e118.

[0017] Examples of IVC prior art belonging to group b):

[0018] Tawfik, D. S. and Griffiths, A. D. (1998) Man-made cell-like compartments for molecular evolution. Nat. Biotechnol., 16, 652-656;

[0019] Ghadessy, F. J., Ong, J. L. and Holliger, P. (2001) Proc. Natl. Acad. Sci. USA, 98, 4552-4557;

- [0020] Tay Y, Ho C, Droge P, Ghadessy F J. Selection of bacteriophage lambda integrases with altered recombination specificity by in vitro compartmentalization. *Nucleic Acids Res.* 2010 March; 38(4):e25. Epub 2009 Dec. 4;
- [0021] Zheng Y, Roberts R J. Selection of restriction endonucleases using artificial cells. *Nucleic Acids Res.* 2007; 35(11):e83. Epub 2007;
- [0022] Mastrobattista E, Taly V, Chanudet E, Treacy P, Kelly B T, Griffiths A D. High-throughput screening of enzyme libraries: in vitro evolution of a beta-galactosidase by fluorescence-activated sorting of double emulsions. *Chem. Biol.* 2005 December; 12(12): 1291-300;
- [0023] Levy M, Griswold K E, Ellington A D. Direct selection of trans-acting ligase ribozymes by in vitro compartmentalization. *RNA.* 2005 October; 11(10):1555-62. Epub 2005 Aug. 30;
- [0024] Sepp A, Choo Y. Cell-free selection of zinc finger DNA-binding proteins using in vitro compartmentalization. *J Mol Biol.* 2005 Nov. 25; 354(2):212-9. Epub 2005 Oct. 3;
- [0025] Bernath K, Magdassi S, Tawfik D S. Directed evolution of protein inhibitors of DNA-nucleases by in vitro compartmentalization (IVC) and nano-droplet delivery. *J Mol Biol.* 2005 Feb. 4; 345(5):1015-26. Epub 2004 Dec. 7.
- [0026] Examples of further IVC prior art may be found in:
- [0027] Bertschinger et al, (2004) *Protein Engineering, Design & Selection* vol. 20 no. 2 pp. 699-707;
- [0028] Chen Yu et al, (November 2008) *Nucleic Acid Research*, Vol. 36, Nr. 19, Pages: Article No. E128;
- [0029] Hansen et al. *J. Am. Chem. Soc.*, 2009, 131 (3), pp 1322-1327.

SUMMARY OF THE INVENTION

[0030] The problem to be solved by the present invention may be seen as to provide an improved in vitro display based method in order to make an enriched library comprising at least one binding entity (e.g. a chemical compound) that binds to a target of interest (e.g. a medical relevant receptor).

[0031] In many cases, most notably in the development of therapeutics, two parameters for a binding entity (drug) are especially important, namely the potency (affinity) and the off rate (dissociative half-life of drug:target complex). The present invention provides an improved solution for in vitro display methods to enrich for both these important binding parameters.

[0032] In other cases, the on-rate characteristic for a binding identity is desired. The present invention provides an improved solution for in vitro display methods to enrich for on-rate characteristic for a binding identity.

[0033] The solution may be seen as based on that:

(i): making an in vitro display library of binding entities (i.e. phenotype) attached to nucleic acid molecules (genotype)—this step may be made according to known prior art techniques for making such in vitro display libraries;

(ii): making structures comprising target (i.e. phenotype) attached to a nucleic acid molecule (genotype)—this step may be made according to known prior art techniques for making such structures; and

wherein the method as described herein may be seen as characterized by that:

(iii): the binding step is performed in solution (e.g. under aqueous conditions);

(iv): there is used a suitable in vitro compartmentalization system (e.g. a water-in-oil emulsion system) creating more individual compartments than target molecules;

(v): fuse co-compartmentalized target and binding entity genotypes;

(vi): de-compartmentalize; to get an enrichment of fused genotypes (positive binders in the in vitro display library will have a higher propensity for being fused than non-binders); and

(vii): optionally e.g. purify and/or preferential amplify the fused genotypes.

[0034] Based on the detailed description herein and the common general knowledge—the skilled person may perform the steps (iii) to steps (vi) in a number of different ways.

[0035] Step (vii) is an optional step—as described herein once one has obtained the enriched library of step (vi) one may use this library in different ways according to art—e.g. the enriched library may be considered as an enriched in vitro display library that e.g. can be used in a second round of selection/enrichment or one may identify the structure of a specific binding entity of interest directly from the enriched library of step (vi).

[0036] As discussed above—herein relevant so-called IVC prior art technologies—may be described as a:

group a)—wherein the phenotype activity is interrogated AFTER the compartmentalized step; or

group b)—wherein the phenotype activity is interrogated DURING the compartmentalized step.

[0037] As evident from above and as further discussed herein—the method as described herein is conceptionally different from such so-called IVC prior art technologies—e.g. due to that the phenotype activity is interrogated in step (iii) of first aspect, which is BEFORE the compartmentalized step (iv) of first aspect.

[0038] A simple way to explain the principle of the novel method as described herein, is that non-binders in the display library is randomly distributed in the compartments and therefore co-compartmentalize with the target in a random fashion, with a frequency depending on the ratio between the number of compartments and the number of target molecules. In contrast, binders, due to the binding activity, will co-compartmentalize together with target molecules—independently of the ratio between the number of compartments and the number of target molecules. Consequently, enrichment of a binder is achieved when the ratio between the number of compartments and the number of target molecules is larger than 1—the higher ratio the higher enrichment.

[0039] In FIGS. 1 and 2 herein are provided illustrative examples of the novel method as described herein.

[0040] In working examples 1 and 2 herein are provided an example with herein relevant numbers of e.g. binding entities and target of interest—as can be seen in the conclusion of the examples 1 and 2—by using the method as described herein one may get e.g. an 1000 times enrichment of binders in the library.

[0041] Accordingly, a first aspect of the invention relates to a method a method for making an enriched library comprising specific nucleic acid sequence information allowing to identifying at least one binding entity that binds to at least one target wherein the specific binding entity has been present in an in vitro display library and wherein the method comprises the steps of:

(i): making an in vitro display library of at least 100 different binding entities (B_n ($n=100$ or more)), wherein each binding

entity is attached to a nucleic acid molecule and the nucleic acid molecule comprises specific nucleic acid sequence information allowing to identify the binding entity—i.e. once one knows the specific nucleic acid sequence information of the nucleic acid molecule one directly knows the structure of the specific binding entity attached to the nucleic acid molecule—the structure of the binding entity (i.e. phenotype) attached to the nucleic acid molecule (genotype) is herein termed B-structure;

(ii): making nucleic acid molecules with at least one target T_n ($n=1$ or more) attached to a nucleic acid molecule and the nucleic acid molecule comprises specific nucleic acid sequence information allowing to identify the specific target, wherein the target is capable of binding to at least one of the binding entities present in the library of step (i)—the structure of the target (i.e. phenotype) attached to the nucleic acid molecule (genotype) is herein termed T-structure;

and wherein the method is characterized by that:

(iii): mixing a solution comprising X (X is a number greater than 10^4) numbers of B-structures of the library of step (i) with a solution comprising Y (Y is a number greater than 10^2) numbers of T-structures of step (ii) under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do and wherein one gets binding of at least one of the binding entities to at least one target thereby creating a complex comprising a B-structure bound to a T-structure (herein termed $B_{BoundTo}$ T-structure);

(iv): applying an in vitro compartmentalization system—under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do—wherein the compartmentalization system comprises at least 2 times more individual compartments than the Y number of T-structures present in step (iii) under conditions wherein the B-structures, T-structures and $B_{BoundTo}$ T-structures enter randomly into the individual compartments; and

(v): fusing the nucleic acid molecules of a B-structure and a T-structure which are both present within the same individual compartment—i.e. fusing the nucleic acid molecule of the B-structure to the nucleic acid molecule of the T-structure—this structure is herein termed BT_{Fused} -structure and the BT_{Fused} -structure comprises the specific nucleic acid sequence information allowing to identify the binding entity of step (i) and the specific nucleic acid sequence information allowing to identify the specific target of step (ii); and

(vi): combining the content of the individual compartments of step (v) under conditions wherein there is no fusing of the nucleic acid molecules of a B-structure and a T-structure—i.e. there is not created any new BT_{Fused} -structure not already created in step (v)—in order to get a library of BT_{Fused} -structures, wherein the library is an enriched library of species of BT_{Fused} -structures originating from binding pairs of target and binder entity when compared to BT_{Fused} -structures originating from nonbinding pairs of target and binder entity.

[0042] The method of the first aspect as described herein may be termed Enrichment by Co-Compartmentalization (ECC).

[0043] Advantageous in ECC method as described herein is that enrichment for important binding characteristics can be

optimized for in isolation—because ECC is a homogenous assay—target is not immobilization to a solid support. Prior art methods are heterogenous—rely on target immobilization to a solid support (e.g. beads, columns, cells, plastic, filters etc). Heterogenous assays are notoriously more difficult to control than homogenous assay due e.g. avidity effects, density of coating, and interference of the solid support itself with the assay.

[0044] As discussed above, in herein relevant in vitro display technology prior art (e.g. above mentioned prior art)—selection for suitable binding entities present within in vitro display libraries are in the prior art generally done by immobilizing the target (e.g. a receptor) to a solid support (e.g. a glass plate, a column, a bead, a nitrocellulose filter, a cell etc) before or after the display library binding event. Non-binders and low affinity binders are typically washed away, whereas the population enriched for binders are recovered from the solid support.

[0045] Accordingly, as understood by the skilled person, when there above is said that herein relevant prior art methods “rely on target immobilization to a solid support” it is understood by the skilled person in the way that the selection for suitable binding entities relies on this immobilization of target to a solid support as an essential element to get the selection for suitable binding entities.

[0046] As evident to the skilled person, the method of the first aspect is not such a prior art method that rely on target immobilization to a solid support, since the selection of the binding entities is based on the separation of the $B_{BoundTo}$ T-structures into the individual compartments as required in step (iv) of the first aspect.

[0047] In line of above and as understood by the skilled person—in the method of the first aspect one could theoretically image a situation, wherein the target T-structure of step (ii) would e.g. comprise a bead. It could theoretically be a T-structure, wherein the target is bound to a bead and the nucleic acid molecule that comprises the specific nucleic acid sequence information allowing identifying the specific target of the T-structure of step (ii) is then also bound to the bead.

[0048] As evident to the skilled person—such a special T-structure comprising a bead will not change the fact that the method of the first aspect is not a method, wherein the selection for suitable binding entities relies on this immobilization of target to a solid support.

[0049] In line of above and as understood by the skilled person in the present context, the method of the first aspect may be seen as a method which implies that the $B_{BoundTo}$ T-structures (i.e. the target-binding entity complexes) remain suspended in solution in the individual/separated compartments of step (iv) of the first aspect.

[0050] ECC allows optimizing for major binding characteristic for binding of binding entity to target in isolation. For example potency (affinity), association rate (on rate) or dissociative half-life of binding entity and target (off rate).

[0051] Affinity based selection is achieved by using equilibrium conditions and controlled by the target concentration in the mixing step (binding step), i.e. 90% of the molecules of a binding entity in the display library having a K_d equal to 10 times smaller than the target concentration are target bound, whereas 50% of the molecules of a binding entity having a K_d equal to the target concentration are, and 10% of the molecules of a binding entity having a K_d 10 times smaller than

the target concentration are. Consequently, enrichment for affinity is easily controlled by the target concentration in the mixing step.

[0052] A separate aspect of the invention relates to an enriched library of step (vi) of the first aspect and which is obtainable by the method of the first aspect or herein related embodiments of the first aspect.

[0053] Embodiments of the present invention are described below, by way of examples only.

DRAWINGS

[0054] FIG. 1: Illustrative example of the principle of the principle of the method as described herein.

[0055] FIG. 2: Illustrative example of the principle of the principle of the method as described herein—it is an illustrative example wherein emulsion PCR is used in the fusion step (v) of the first aspect.

[0056] FIG. 3: Herein is shown an example of the in vitro display technology as described in EP1809743B1 (Vipergen)—as can be seen in this FIG. 3—the selection step of this example is performed by immobilizing the target (e.g. a receptor) to a solid surface (e.g. a bead or a glass plate).

[0057] FIGS. 4-7: These figures are further discussed in working examples herein.

DETAILED DESCRIPTION OF THE INVENTION

In Vitro Display Library—Step (i) of First Aspect

[0058] The term “in vitro display library” shall be understood according to the art—i.e. as a library comprising numerous different binding entities wherein each binding entity is attached to a nucleic acid molecule and the nucleic acid molecule comprises specific nucleic acid sequence information allowing to identify the binding entity—i.e. once one knows the specific nucleic acid sequence information of the nucleic acid molecule one directly knows the structure of the specific binding entity attached to the nucleic acid molecule—the structure of the binding entity (i.e. phenotype) attached to the nucleic acid molecule (genotype) is herein termed B-structure.

[0059] As discussed herein—the prior art describes a number of different methods to make such in vitro display libraries—i.e. an in vitro display library of step (i).

[0060] Said in other words, it is today routine work for the skilled person to properly make a structure of the binding entity (i.e. phenotype) attached to the nucleic acid molecule (genotype)—i.e. what is herein termed a “B-structure”.

[0061] As known in the art—binding entity (i.e. phenotype) may be attached to the nucleic acid molecule (genotype) by e.g. a covalent binding or e.g. a high affinity non-covalent binding.

[0062] It may herein be preferred that the binding entity (i.e. phenotype) is attached to the nucleic acid molecule (genotype) by a covalent binding.

[0063] An in vitro display library of step (i) comprises a number of different B-structures—i.e. in line of above it is routine work for the skilled person to make an in vitro display library of step (i).

[0064] Herein suitable examples include e.g. EP1809743B1 (Vipergen), EP1402024B1 (Nuevolution), EP1423400B1 (David Liu), Nature Chem. Biol. (2009), 5:647-654 (Clark), WO 00/23458 (Harbury), Nature Methods (2006), 3(7), 561-570, 2006 (Miller), Nat. Biotechnol. 2004;

22, 568-574 (Melkko), Nature. (1990); 346(6287), 818-822 (Ellington), or Proc Natl Acad Sci USA (1997). 94 (23): 12297-302 (Roberts).

[0065] Said in other words, the in vitro display library of step (i) of first aspect may be made in a numbers of ways as described in the prior art.

[0066] Without being limited to theory—herein suitable examples of in vitro display library technologies include DNA Encoded Chemical Library technologies, Aptamer technologies, RNA/DNA display technologies such as CIS display, Ribosome display, mRNA display or bead display system (using nucleic acids for encoding).

[0067] As described in the prior art (see e.g. EP1809743B1 (Vipergen))—the nucleic acid molecule of the B-structure may e.g. be PNA, LNA, RNA, DNA or combinations thereof. Preferably, the nucleic acid molecule of the B-structure is DNA.

[0068] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the binding entity (phenotype) in the B-structure may be a double stranded nucleic acid molecule.

[0069] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the binding entity (phenotype) in the B-structure may be at least 0% double stranded (i.e. single stranded), may be at least 10% double stranded, at least 20% double stranded, at least 30% double stranded, at least 40% double stranded, at least 50% double stranded, at least 60% double stranded, at least 70% double stranded, at least 80% double stranded, at least 90% double stranded, or 100% double stranded.

[0070] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the binding entity (phenotype) in the B-structure may contain a PCR priming site or a fraction hereof.

[0071] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the binding entity (phenotype) in the B-structure may contain 2 PCR priming sites or fractions hereof.

[0072] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the binding entity (phenotype) in the B-structure may contain at least 3 PCR priming sites or fractions hereof.

[0073] In some embodiments of the present invention a fraction of a PCR priming site comprises at least 5 nucleotides, at least 6 nucleotides, at least 7 nucleotides, at least 8 nucleotides, at least 9 nucleotides, at least 10 nucleotides, at least 11 nucleotides, at least 12 nucleotides, at least 13 nucleotides, at least 14 nucleotides, at least 15 nucleotides, at least 16 nucleotides, at least 17 nucleotides, at least 18 nucleotides, at least 19 nucleotides, or at least 20 nucleotides.

[0074] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to the binding entity (phenotype) in the B-structure may contain a single stranded overhang reverse complement to a single stranded overhang of the genotype of the B structure.

[0075] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to the binding entity (phenotype) in the B-structure may contain a single stranded overhang reverse complement to a single stranded overhang of the genotype of the B structure. The overhang may preferentially be 1 nucleotide, 2 nucleotides, 3 nucleotides, 4 nucleotides, 5 nucleotides, 6 nucleotides, 7 nucleotides, 8 nucleotides, 9 nucleotides, or 10 nucleotides long.

Binding Entity

[0076] The Binding entity may any suitable binding entity of interest.

[0077] Step (i) of first aspect reads “at least 100 different binding entities (B_n ($n=100$ or more))”.

[0078] In practice, there may many times be many more different binding entities present in the library of step (i)—such as e.g. at least 10^4 , at least 10^5 or at least 10^6 different binding entities—i.e. where $n=$ at least 10^4 , $n=$ at least 10^5 or $n=$ at least 10^6 .

[0079] Accordingly, in a theoretical situation, wherein the library comprises exactly 10^4 different binding entities—one may herein express this as B_n ($n=10^4$) or B_{10^4} .

[0080] Without being limited to theory it may be difficult to make a library with more than 10^{20} different binding entities.

[0081] Suitable examples may be wherein the binding entity is at least one binding entity selected from the group consisting of: a protein, a polypeptide, a nucleic acid and a chemical compound (preferably a small chemical compound with an average molecular weight MW below 10000 dalton, more preferably an average molecular weight MW below 5000 dalton, even more preferably an average molecular weight MW below 1000 dalton).

[0082] Suitable examples of a herein relevant binding entity (such as e.g. a chemical compound) may be found in the prior art—see e.g. EP1809743B1 (Viperger), EP1402024B1 (Nuevolution), EP1423400B1 (David Liu), Nature Chem. Biol. (2009), 5:647-654 (Clark), WO 00/23458 (Harbury), Nature Methods (2006), 3(7), 561-570, 2006 (Miller), Nat. Biotechnol. 2004; 22, 568-574 (Melkko), Nature. (1990); 346(6287), 818-822 (Ellington), or Proc Natl Acad Sci USA (1997). 94 (23): 12297-302 (Roberts).

[0083] In short, the skilled person is aware of numerous different possible binding entities that could be of interest in the present context.

Step (ii) of First Aspect

[0084] As discussed herein—the target shall be capable of binding to at least one of the binding entities present in the library of step (i)—otherwise it is not a suitable target that can be used to identify a specific binding entity that binds to at least one target.

[0085] In line of above—it is today routine work for the skilled person to properly attach a target (i.e. phenotype) to a nucleic acid molecule (genotype) and thereby make a structure of the target (i.e. phenotype) attached to the nucleic acid molecule (genotype)—i.e. what is herein termed “T-structure”.

[0086] Said in other words, one may make herein relevant “T-structure” based on e.g. the same prior art literature discussed above for making the in vitro display library of step (i).

[0087] As known in the art—target (i.e. phenotype) may be attached to the nucleic acid molecule (genotype) by e.g. a covalent binding or e.g. a high affinity non-covalent binding.

[0088] It may herein be preferred that the target (i.e. phenotype) is attached to the nucleic acid molecule (genotype) by a covalent binding.

[0089] Step (i) of first aspect reads “at least one target T_n ($n=1$ or more)”.

[0090] As discussed herein—an advantage of the method as described herein is that one in an efficient and rapid way can simultaneous screen for binding entities that could bind to e.g. two or more targets.

[0091] For instance—the targets could be two different receptor molecules and the method as described herein could then simultaneous identify one binding entity that binds to one of the receptors and another binding entity that binds to the other receptor.

[0092] In the example above (with two different e.g. receptor targets) we would have a situation, wherein the target T_n ($n=2$) or alternatively expressed T_2 .

[0093] In line of above—it may be relevant to have at least two different targets in step (ii) [i.e. T_n ($n=2$ or more)], or to at least three different targets in step (ii) [i.e. T_n ($n=3$ or more)], or to have at least ten different targets in step (ii) [i.e. T_n ($n=10$ or more)], or to at least hundred different targets in step (ii) [i.e. T_n ($n=100$ or more)].

[0094] Without being limited to theory it may be difficult to have than 100.000 different targets in step (ii)—i.e. more than 100.000 different T-structures.

[0095] As described in the prior art (see e.g. EP1809743B1 (Viperger))—the nucleic acid molecule of the T-structure may e.g. be PNA, LNA, RNA, DNA or combinations thereof. Preferably, the nucleic acid molecule of the T-structure is DNA.

[0096] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may be at least 5 nucleotides long, at least 10 nucleotides long, at least 20 nucleotides long, at least 30 nucleotides long, at least 40 nucleotides long, at least 50 nucleotides long, at least 60 nucleotides long, at least 70 nucleotides long, at least 80 nucleotides long, at least 90 nucleotides long, at least 100 nucleotides long, at least 200 nucleotides long, at least 300 nucleotides long, at least 400 nucleotides long, or at least 500 nucleotides long.

[0097] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may be a double stranded nucleic acid molecule.

[0098] In a preferred embodiment of the present invention the double stranded nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may be at least 5 base pairs long, at least 10 base pairs long, at least 20 base pairs long, at least 30 base pairs long, at least 40 base pairs long, at least 50 base pairs long, at least 60 base pairs long, at least 70 base pairs long, at least 80 base pairs long, at least 90 base pairs long, at least 100 base pairs long, at least 200 base pairs long, at least 300 base pairs long, at least 400 base pairs long, or at least 500 base pairs long.

[0099] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may be at least 0% double stranded (i.e. single stranded), may be at least 10% double stranded, at least 20% double stranded, at least 30% double stranded, at least 40% double stranded, at least 50% double stranded, at least 60% double stranded, at least 70% double stranded, at least 80% double stranded, at least 90% double stranded, or 100% double stranded.

[0100] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain a PCR priming site or a fraction hereof.

[0101] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain 2 PCR priming sites or fractions hereof.

[0102] In a preferred embodiment of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain at least 3 PCR priming sites or fractions hereof.

[0103] In some embodiments of the present invention a fraction of a PCR priming site comprises at least 5 nucleotides, at least 6 nucleotides, at least 7 nucleotides, at least 8 nucleotides, at least 9 nucleotides, at least 10 nucleotides, at least 11 nucleotides, at least 12 nucleotides, at least 13 nucleotides, at least 14 nucleotides, at least 15 nucleotides, at least 16 nucleotides, at least 17 nucleotides, at least 18 nucleotides, at least 19 nucleotides, or at least 20 nucleotides.

[0104] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain a single stranded overhang reverse complement to a single stranded overhang of the genotype of the B structure.

[0105] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain a single stranded overhang reverse complement to a single stranded overhang of the genotype of the B structure. The overhang may preferentially be 1 nucleotide, 2 nucleotides, 3 nucleotides, 4 nucleotides, 5 nucleotides, 6 nucleotides, 7 nucleotides, 8 nucleotides, 9 nucleotides, or 10 nucleotides long.

[0106] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain a unique sequence specific for each target molecule (Unique Molecule Identifier—UMI).

[0107] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain a unique sequence specific for each target molecule (Unique Molecule Identifier—UMI) consisting of at least 16 Ns (N=A, C, G, or T), at least 17 Ns, at least 18 Ns, at least 19 Ns, at least 20 Ns, at least 21 Ns, at least 22 Ns, at least 23 Ns, at least 24 Ns, at least 25 Ns, at least 26 Ns, at least 27 Ns, at least 28 Ns, at least 29 Ns, or at least 30 Ns.

[0108] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain a unique sequence specific for each target molecule (Unique Molecule Identifier—UMI) consisting of a continuous sequence.

[0109] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to the target (phenotype) in the T-structure may contain a unique sequence specific for each target molecule (Unique Molecule Identifier—UMI) consisting of a discontinuous sequence.

[0110] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to a first target (phenotype) in the T-structure may contain a first sequence different from a second target's second genotype sequence (allowing multiplexing).

[0111] In some embodiments of the present invention the nucleic acid molecule (genotype) attached to a first target (phenotype) in the T-structure may contain a first sequence different from a second target's second genotype sequence (allowing multiplexing), wherein the first and second target genotype comprise different PCR priming sites.

Target

[0112] The target may be any suitable target of interest.

[0113] In a preferred embodiment of the present invention—specific enriching methods for the enrichment facilitating identification of binding entities with desired characteristics include but are not limited to: enrichment on nucleic acid attached target molecules. In this approach the target molecules is e.g. DNA, RNA, protein, carbohydrate, organic or inorganic molecule.

[0114] As known in the art—a suitable target could e.g. be a receptor molecule present in e.g. the human body and one would be interested in identifying a binding entity (e.g. a chemical compound) that can bind to the receptor.

[0115] In accordance with the prior art—suitable examples may be wherein the target is DNA, RNA, protein, carbohydrate, organic or inorganic molecule or fragments hereof.

[0116] In accordance with the prior art—suitable examples may be wherein the target is an autoantigen, a bacterial protein, a blood protein, a cell adhesion protein, a cytokine, a cytoskeleton protein, a DNA-binding protein, a developmental protein, an engineered protein, an enzyme, an extracellular matrix protein, a GTP-binding protein regulator, a glycoprotein, a growth factor, a heat shock protein, a lipoprotein, a membrane protein, a metalloprotein, a motor protein, a phosphoprotein, a prion, a protein complex, a protein domain, a RNA-binding protein, a receptor, a recombinant protein, a seed storage protein, a structural protein, a transcription coregulator protein, a transport protein, a viral protein or fragments hereof.

[0117] In short, the skilled person is aware of numerous different possible targets than could be of interest in the present context.

Step (iii) of First Aspect:

[0118] In the illustrative example of FIG. 1 herein—this step (iii) corresponds to the step “1 Binding”.

[0119] As discussed above—step (iii) reads:

“mixing a solution comprising X (X is a number greater than 10^4) numbers of B-structures of the library of step (i))”

[0120] The term “X” in relation to numbers of B-structures shall be understood as the total numbers of B-structures of the library of step (i).

[0121] For instance—if the library comprises 100 different binding entities (B_n ($n=100$)) and there are 100 copies of each of the 100 different B-structures then the number “X” is equal to $100 \times 100 = 10^4$.

[0122] In practice the number X may many times be higher—for instance, if the library comprises 10^6 different binding entities [B_n ($n=10^6$)] and there are 10^4 copies of each of the 10^6 different B-structures then the number “X” is equal to $10^6 \times 10^4 = 10^{10}$.

[0123] As discussed above—step (iii) reads:

“a solution comprising Y (Y is a number greater than 10^2) numbers of T-structures of step (ii))”

[0124] The term “Y” in relation to numbers of T-structures shall be understood as the total numbers of T-structures of the library of step (ii).

[0125] For instance—if there is only one target in step (ii) [T_n ($n=1$)] and there are 10^2 copies of each of the T-structure then the number “Y” is equal to $1 \times 10^2 = 10^2$.

[0126] As discussed above—one could have e.g. 2 different targets (e.g. two different receptor molecules)—in this case there would be two targets in step (ii) [T_n ($n=2$)] and if there would be 10^2 copies of each of the two different T-structures then the number “Y” would be equal to $2 \times 10^2 = 2 \times 10^2 = 200$.

[0127] In practice one may many times have significant more copies of a relevant T-structure—the reason for this is that one preferably wants to have numerous copies of a relevant T-structure in order to increase the probability for that the target on a T-structure bind to the binding entity of a B-structure.

[0128] Accordingly, in a preferred embodiment there are at least 10^0 , at least 10^1 , at least 10^2 , at least 10^3 , at least 10^4 , at least 10^5 , at least 10^6 , at least 10^7 at least 10^8 , at least 10^9 , at least 10^{10} , at least 10^{11} , at least 10^{12} , at least 10^{13} , at least 10^{14} , at least 10^{15} or at least 10^{16} copies of a T-structure of interest.

[0129] Advantageous in ECC method as described herein is that enrichment for important binding characteristics can be optimized for in isolation—because ECC is a homogenous assay—target is not immobilization to a solid support. Prior art methods are heterogeneous—rely on target immobilization to a solid support (e.g. beads, columns, cells, plastic, filters etc). Heterogeneous assays are notoriously more difficult to control than homogenous assay due e.g. avidity effects, density of coating, and interference of the solid support itself with the assay.

[0130] ECC allows optimizing for major binding characteristic for binding of binding entity to target in isolation. For example potency (affinity), association rate (on rate) or dissociative half-life of binding entity and target (off rate).

[0131] Affinity based selection is achieved in step (iii) e.g. by using equilibrium conditions and controlled by the target concentration in the mixing step (binding step), i.e. 90% of the molecules of a binding entity in the display library having a K_d equal to 10 times smaller than the target concentration are target bound, whereas 50% of the molecules of a binding entity having a K_d equal to the target concentration are, and 10% of the molecules of a binding entity having a K_d 10 times smaller than the target concentration are. Consequently, enrichment for affinity is easily controlled by the target concentration in the mixing step.

[0132] In a preferred embodiment of the present invention—the concentration of T-structures in the “mixing step (iii)” is at least 10^{-15} M, at least 10^{-14} M, at least 10^{-13} M, at least 10^{-12} M, at least 10^{-11} M, at least 10^{-10} M, at least 10^{-9} M, at least 10^{-8} M, at least 10^{-7} M, at least 10^{-6} M, at least 10^{-5} M, at least 10^{-4} M, or at least 10^{-3} M.

[0133] Alternatively, association rate based selection is achieved by controlling the time allowed for the mixing step (iii)—accordingly, the “mixing step” may be performed for a time period shorter than the time needed to reach binding equilibrium conditions.

[0134] Step (iii) further reads:

“under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do and wherein one gets binding of at least one of the binding entities to at least one target thereby creating a complex comprising a B-structure bound to a T-structure (herein termed $B_{BoundTo}$ T-structure)”

[0135] The term “binds more efficiently” shall be understood according to common practice e.g. higher affinity, faster on rate, or slower dissociation rate.

[0136] As known to the skilled person—in the present context it is routine work for the skilled person to perform step (iii) under conditions, wherein one gets this “binds more efficiently” effect.

[0137] For instance—one may easily obtain this “binds more efficiently” effect by e.g. using B and T-structures genotypes that essentially do not bind (e.g. by hybridization base pairing) under the binding conditions of step (iii)—as evident to the skilled person this could e.g. be obtained by using e.g. double stranded DNA with none or very small single stranded base-pairing overlap as genotypes for the B and T-structures.

[0138] It would be routine work for the skilled person to optimize the binding conditions of step (iii) in order to get the required “binds more efficiently” effect of step (iii).

[0139] As known to the skilled person—herein relevant optimization parameters may e.g. be ionic strength, temperature etc.

[0140] Accordingly, under any practical herein relevant circumstance—the skilled person would not be in any reasonable doubt if he (after e.g. proper routine adjustment of the binding conditions) would work under binding conditions of step (iii) or not.

[0141] In a preferred embodiment, step (iii) is performed under binding conditions, wherein a B-structure containing a binding entity capable of binding to a target molecule, binds 10 fold (more preferably 100 fold, even more preferably 1000 fold) more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do.

Step (iii-b)—Dilution Step—Preferred Embodiment:

[0142] In the illustrative example of FIG. 1 herein—this optional step (iii-b) corresponds to the step “2 Dilution”.

[0143] The mixing step (iii) may preferably be followed by a dilution step—this is herein termed step (iii-b) and is performed before the step (iv) of the first aspect.

[0144] Accordingly, in a preferred embodiment the method of the first aspect comprises an additional step (iii-b) that is performed before the step (iv) of the first aspect, comprising: (iii-b): diluting the solution of step (iii) at least 2 fold under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do.

[0145] The dilution solution introduced and the conditions (e.g. temperature) in the dilution step (iii-b) may be different from the binding conditions of the mixing step (iii)—but the above described effects shall be maintained in dilution step (iii-b).

[0146] It may be preferred in step (iii-b) to have a diluting the solution of step (iii) at least 10^2 fold, or have a diluting the solution of step (iii) at least 10^3 fold, or have a diluting the solution of step (iii) at least 10^4 fold, or have a diluting the solution of step (iii) at least 10^5 fold, or have a diluting the solution of step (iii) at least 10^6 fold, or have a diluting the solution of step (iii) at least 10^7 fold, or have a diluting the solution of step (iii) at least 10^8 fold or have a diluting the solution of step (iii) at least 10^9 fold.

[0147] An advantage of this diluting step is that enrichment can be performed based on dissociative half-life of the BT-structures and easily controlled by the degree of dilution and the incubation time. When the mixing solution of step (iii) is diluted binding of binding entity and target is a less likely event to happen whereas the “un-binding event”—the off rate (the dissociative half-life) is independent of the dilution. Consequently, in a very dilute solution (T-structure concentration $\ll K_d$) essentially only dissociation will take place. Therefore, enrichment for dissociative half-life of the

BT-structures is conveniently controlled by the degree of dilution and the incubation time.

[0148] The dissociative half-life together with the affinity is of greatest importance in the usability of a binding entity. Most notable, for development of effective new drugs where high affinity and long dissociative half-life are critical parameters for pharmacological effect (Nature Reviews Drug Discovery (2006) 5, 730-739, (Copeland). Hence, the method of the present new invention permits enrichment for these two parameters in an unprecedented effective and controllable manner. Moreover, the two parameters can be controlled independently of each other.

Step (iv) of First Aspect:

[0149] A simple way to view this step is that the binding of target with binding entity of step (iii) is “transformed” into co-compartmentalization of B-structures and T-structures.

[0150] The conditions of this step (iv) shall be “under binding conditions” that gives an effect corresponding to the effect in step (iii)—see above.

[0151] In the illustrative example of FIG. 1 herein—this step (iv) corresponds to the step “3 Emulsion w/o”.

[0152] Step (iv) of first aspect further reads:

“wherein the compartmentalization system comprises at least 2 times more individual compartments than the Y number of T-structures present in step (iii)”

[0153] This may herein be seen as an essential step of the method as described herein—i.e. it is essential to have “at least 2 times more individual compartments than the Y number of T-structures present in step (iii)”.

[0154] In the FIG. 1 herein—the in vitro compartmentalization system may be e.g. a water-in-oil emulsion system—as further discussed below herein suitable water-in-oil emulsion systems are well known in the art.

[0155] In the hypothetical theoretical illustrative example in FIG. 1 there is only one target in step (ii) [T_n ($n=1$)] and there are 3 copies of each of the T-structure—i.e. the number “Y” is 3.

[0156] Accordingly, in this theoretical illustrative example of FIG. 1 there should be at least $(2 \times 3) = 6$ individual compartments (e.g. oil droplets) in the in vitro compartmentalization system—please note that in FIG. 1 herein are there less than 30 individual compartments (i.e. FIG. 1 is just an illustration of some of the elements of the method as described herein).

[0157] As discussed above—in practice there may be for instance at least 10^4 copies of a T-structure of interest—i.e. Y could in this case be 10^4 and there should in this case be at least $(2 \times 10^4) = 2 \times 10^4$ individual compartments (e.g. oil droplets) in the in vitro compartmentalization system.

[0158] An advantage of having this “at least 2 times more individual compartments than the Y number of T-structures” is that non-binders in the display library is randomly distributed in the compartments and therefore co-compartmentalize with the target in a random fashion, with a frequency depending on the ratio between the number of compartments and the number of target molecules (in this case 1 out of 10), whereas binders, due to the binding activity, will co-compartmentalize together with target molecules independently of the ratio between the number of compartments and the number of target molecules (in the ideal case 1 out of 1). Consequently, in this case binders will be enriched 2 fold when compared to non-binders.

[0159] In line of above one may say one gets an even better enrichment if there is relatively more individual compartments in the in vitro compartmentalization system—accordingly, in a preferred embodiment there is “at least 10 times more individual compartments than the Y number of T-structures present in step (iii)”, more preferably there is “at least 100 times more individual compartments than the Y number of T-structures present in step (iii)”, more preferably there is “at least 10 000 times more individual compartments than the Y number of T-structures present in step (iii)”, more preferably there is “at least 100 000 times more individual compartments than the Y number of T-structures present in step (iii)”, more preferably there is “at least 1 000 000 times more individual compartments than the Y number of T-structures present in step (iii)”.

[0160] In a preferred embodiment of the present invention the number of compartments is larger than 2, 5, 10, 50, 100, 1000, 5000, 10 000, 50 000, 100 000, 500 000, 1 000 000, 5 000 000, or 10 000 000 times the Y number of T-structures of step (iii).

[0161] Step (iv) of first aspect further reads:

“under conditions wherein the B-structures, T-structures and BT-structures enter randomly into the individual compartments”

[0162] This should understood as the skilled person would understand it in the present context—relating to that in order to get the herein advantageous enrichment one needs to have conditions wherein the B-structures, T-structures and BT-structures enter randomly into the individual compartments.

[0163] Said in other words—the propensity for any B-structures, T-structures and BT-structures for being compartmentalized in any given compartment is dependent on the volume of said compartment and the total volume.

[0164] Said in other words—if virtually all the B-structures, T-structures and BT-structures would only enter into one specific individual compartment one will obviously not get the advantageous enrichment as discussed herein.

[0165] As discussed below—if one e.g. uses a suitable water-in-oil emulsion system as the in vitro compartmentalization system one can easily identify conditions, wherein the B-structures, T-structures and BT-structures enter randomly into the individual compartments (e.g. an individual oil droplets)—in fact it would be quite difficult to identify conditions, where it is not randomly—i.e. wherein virtually all the B-structures, T-structures and BT-structures would only enter into one specific individual compartment (e.g. an individual oil droplets).

[0166] In a preferred embodiment of step (iv)—there is at least square root 10 (3.16) times more individual compartments than the X number of B-structures in step (iii).

[0167] A Poisson distribution is assumed to describe the distribution of B-structures in the compartments. This implies that all compartments are of equal volumes. The probability that a compartment has $n=0, 1, 2$ or more B-structures molecules can be calculated using the formula:

$$f(n, \lambda) = \frac{e^{-\lambda} \lambda^n}{n!}$$

where λ is the ratio between X number of B-structures and number of compartments. When λ is square root 10 (3.16) less than 5% (4.1%) of the compartments will have more than one B-structures. This means that the effect of this is to

lowering the enrichment of positive binding entity with less than 5%, which is insignificant in most cases.

[0168] As discussed above and as understood by the skilled person in the present context, the method of the first aspect may be seen as a method which implies that the $B_{BoundTo}T$ -structures (i.e. the target-binding entity complexes) remain suspended in solution in the individual/separated compartments of step (iv) of the first aspect.

[0169] Accordingly, just to make it 100% clear, one may express this as that the method of the first aspect and herein relevant embodiments of this method, is a method wherein the $B_{BoundTo}T$ -structures remain suspended in solution in the individual compartments of step (iv) of the first aspect.

[0170] Said in other words, the method does not rely on target immobilization on a solid support as for herein relevant prior art methods as discussed above.

In Vitro Compartmentalization System

[0171] As discussed above, a herein suitable in vitro compartmentalization system may e.g. be a water-in-oil emulsion system.

[0172] In case wherein the in vitro compartmentalization system is a water-in-oil emulsion system—one may say that the “applying an in vitro compartmentalization system . . . to the solution of step (iii)” of step (iv) may be expressed as “adding the solution of step (iii) to an water-in-oil emulsion system”.

[0173] Herein suitable water-in-oil emulsion systems are described in the art e.g. Nat Methods. 2006 July; 3(7):545-50 (Williams et al), Nat Methods. 2006 July; 3(7):551-9 (Diehl et al), Proc Natl Acad Sci USA. 2003 Jul. 22; 100(15):8817-22 (Dressman et al), J Biotechnol. 2003 Apr. 24; 102(2):117-24. (Nakano et al), or Biomacromolecules. 6, 1824-1828 (2005) (Musyanovych et al).

[0174] As evident to the skilled person—in the case of using emulsions as the compartmentalization system and in analogy with similar size distributions, the compartment volume distribution is modeled as a log-normal distribution, also called a Galton distribution. By assuming a log-normal distribution and performing measurements of the actual droplet sizes the expected value (mean) and the standard deviation can be calculated for a specific experiment. According to this distribution, 95% of the compartment volumes will be within L logarithmic units from the mean (log) volume, where L is 1.96 times the standard deviation of the log-volumes.

[0175] In a preferred embodiment of the present invention the average compartments size, the variation, and the standard deviation is taken into account when analyzing the data.

[0176] In a preferred embodiment of the present invention compartments with a volume, larger than 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 times the average compartment size are removed from the experiment.

[0177] In a preferred embodiment of the present invention compartments with a volume, smaller than $\frac{1}{100}$, $\frac{1}{90}$, $\frac{1}{80}$, $\frac{1}{70}$, $\frac{1}{60}$, $\frac{1}{50}$, $\frac{1}{40}$, $\frac{1}{30}$, $\frac{1}{20}$, $\frac{1}{10}$, $\frac{1}{9}$, $\frac{1}{8}$, $\frac{1}{7}$, $\frac{1}{6}$, $\frac{1}{5}$, $\frac{1}{4}$, $\frac{1}{3}$, or $\frac{1}{2}$, times the average compartment size are removed from the experiment.

[0178] As evident to the skilled person—several technologies may be employed to exclude compartments from the experiment based on the volume of the compartments, for example by FACS sorting, equilibrium centrifugation, filtration, or microfluidic systems etc.

[0179] Alternatively, the in vitro compartmentalization system may be agarose droplet microfluidics. (Lab Chip.

2010 Sep. 13. [Epub ahead of print] Agarose droplet microfluidics for highly parallel and efficient single molecule emulsion PCR. Leng X, Zhang W, Wang C, Cui L, Yang C J).

[0180] Alternatively, the in vitro compartmentalization system may simply be to disperse the solution into e.g. “a micro titer plate” and of step (iii) is simply randomly putted into the individual wells (i.e. the individual compartments) of the micro titer plate—as known this may today be done rapidly and efficient by e.g. a suitable robot machine or an open well system. An example of such a system based on a high-density array of nanoliter PCR assays, functionally equivalent to a microtiter plate, the nanoplate system makes possible up to 3,072 simultaneous PCR reactions in a device, the size of a standard microscope slide (Methods Mol Biol. 2009; 496:161-74. (Brennan et al)).

[0181] Another example is a silicone device presenting a large array of micrometer-sized cavities, which can be used it to tightly enclose volumes of solution, as low as femtoliters, over long periods of time. The microchip insures that the chambers are uniform and precisely positioned (Nat Biotechnol. 2005 March; 23(3):361-5 (Rondelez et al)).

[0182] As evident to the skilled person—microfluidic devices can be employed in the in vitro compartmentalization system (For review see e.g. Angew Chem Int Ed Engl. 2010 Aug. 9; 49(34):5846-68 (Theberge et al)).

[0183] In a preferred embodiment of the present invention a suitable average compartments volume is less than 10^{-6} liter, less than 10^{-7} liter, less than 10^{-8} liter, less than 10^{-9} liter, less than 10^{-10} liter, less than 10^{-11} liter, less than 10^{-12} liter, less than 10^{-13} liter, less than 10^{-14} liter, less than 10^{-16} liter, less than 10^{-16} liter, less than 10^{-17} liter, less than 10^{-18} liter, less than 10^{-19} liter, less than 10^{-20} liter, less than 10^{-21} liter, or less than 10^{-22} liter.

[0184] As evident to the skilled person—the compartment volume cannot be infinitely small as the compartment should be larger than the molecules compartmentalized.

[0185] In short, the skilled person is aware of numerous different in vitro compartmentalization systems than could be of interest in the present context.

Step (v) of First Aspect

[0186] Step (v) of first aspect reads:

“fusing the nucleic acid molecules of a B-structure and a T-structure which are both present within the same individual compartment—i.e. fusing the nucleic acid molecule of the B-structure to the nucleic acid molecule of the T-structure”

[0187] A simple way to view this step is that the co-compartmentalization of B-structures and T-structures origination of (iv) is “transformed” into fused cognate genotypes.

[0188] In the present context “fusing the nucleic acid molecules of a B-structure and a T-structure” shall be understood as joining the genetic information carried by the two genotypes in the compartment.

[0189] As known to the skilled person, this can be accomplished in several ways such as e.g.:

a) information transfer by e.g. overlap PCR or overlap genome extension (overlap PCR without outer primers)—one strand originating from one genotype acts as a primer and uses a strand originating from the other genotype as a template;

b) information joining catalyzed by an enzyme forming an amplifiable facilitating bond—e.g. by a DNA ligase where a phosphodiester bond between at least one of the strands from each genome is form;

c) information joining catalyzed by an enzyme forming a non-amplifiable facilitating bond—e.g. having a moiety on each genome thus upon induction an enzyme capable of linking the two moieties between at least one of the strands from each genotype is form;

d) information joining not catalyzed by an enzyme—e.g. having an inducible chemical reactive group on each genome thus upon induction a chemical bond between at least one of the strands from each genotype is form; and

e) information joining transient—e.g. having an affinity tag (these may be different or identical) on each genome thus by providing an agent with affinity for both tags a ternary complex containing both genotype is formed.

[0190] In the illustrative example of FIG. 1 herein—this step (v) corresponds to fusing the nucleic acid molecules of the B-structure and the T-structure present in the individual compartment number three from the left.

[0191] A herein very important advantage is that during this step one may say that the binding between the binding entity and target is no longer relevant—i.e. when one here performs the fusion of the nucleic acid molecules step one can do it under conditions, wherein one does not have to worry about this binding entity to target binding and spatial arrangements. A simple way to view this step is that the binding of target with binding entity origination from step (i) is now transformed into co-compartmentalization of B-structures and T-structures.

[0192] This may be seen as a very big advantage of the method as described herein.

[0193] For instance, if one wants to make the fusion of the nucleic acid molecules of the B-structure and the T-structure by hybridization of overlapping base pairing regions one can in this step (v) adequately change e.g. the temperature to get the relevant base pairing hybridization without being worried if the binding between the binding entity and target could be destroyed.

[0194] Accordingly, the step (v) may be performed under conditions, wherein there is essentially no binding of any of the binding entities of step (i) to any of the target(s) of step (ii).

[0195] As already discussed above—the fusing of the nucleic acid molecules of the B-structure and the T-structure may be done in different ways than e.g. by hybridization of overlapping base pairing regions.

[0196] For instance, if a e.g. a ligase enzyme is used in step (v) to get the fusing nucleic acid molecules—then one does not need to have any base pairing overlapping regions between the nucleic acid molecules (genotype) of the B-structures step (i) and the nucleic acid molecules (genotype) of the T-structures of step (ii).

[0197] As evident to the skilled person—if e.g. a ligase or a polymerase enzyme is used—this ligase enzyme should preferably have been added to the solution of step (iii) or during the optional diluting step (iii-b) in order to properly be present in the relevant individual compartments of step (v).

[0198] As evident to the skilled person—several different enzymatic reactions may be employed to fuse the genome—a large number enzymatic reactions have been reported in the literature e.g. Kabanov et al., *Biochimica et Biophysica Acta*, 996 (1989) 147-152, Salon et al. *Biochemistry* 1992, 31, 8072-8079, Anarbaev et al, *Biochimica et Biophysica Acta* 1384 1998, 315-324, Ong et al., (2006). *J. Mol. Biol.* 361: 537-50, Ghadessy, F. J. Ong, J. L. and Holliger, P. (2001). *Proc. Natl. Acad. Sci. USA* 98: 4552-4557, *Protein Engineering, Design & Selection* vol. 17 no. 3 pp. 201-204, 2004, Levy


et al, *RNA* (2005), 11:1555-1562, Turner et al., *Nucleic Acids Res.* 2008 August; 36(13): e82.

[0199] In a preferred embodiment of the present invention the co-compartmentalized genotypes are fused by an enzyme.

[0200] As evident to the skilled person—despite the concentration of genotypes in the solution before compartmentalization may be very low e.g. picomolar-micromolar range the concentration of genotypes in a compartment with co-compartmentalized genotypes may be high e.g. when the compartment volume is in the femtoliter (10^{-15} liters) range the concentration of the genotypes are in the nanomolar range (10^{-9} M), when the compartment volume is in the attoliter (10^{-18} liters) range the concentration of the genotypes are in the micromolar (10^{-6} M) range, or when the compartment volume is in the zeptoliter (10^{-21} liters) range the concentration of the genotypes are in the millimolar (10^{-3} M) range. Consequently, the genotype concentration in a compartment may be controlled for facilitating enzymatic reactions and even traditional chemical reactions.

[0201] As evident to the skilled person—if e.g. inducible chemical cross-linking in emulsion is used—neither the inducer nor other reagents may be mandatory before compartmentalization as these may be “delivered” later to the formed compartments. For example inducers may be light, temperature or a chemical activator delivered through the continuous phase. Such embodiments may be advantageous when small compartments are desired.






[0202] In a preferred embodiment of the present new invention a chemical cross-linking is performed by



[0203] The nucleophilic substitution reaction can essentially be performed as described by:




[0204] Z. J. Gartner, et al. J. Am. Chem. Soc. 2001, 123, 6961.

[0205] In a preferred embodiment of the present new invention a chemical cross-linking is performed by



[0206] The nucleophilic aromatic substitution reaction can essentially be performed as described by: Clark et al, Nature Chemical Biology 5, 647-654 (2009)

[0207] In a preferred embodiment of the present new invention a chemical cross-linking is performed by





[0208] The nucleophilic substitution reaction can essentially be performed as described by:

[0209] Z. J. Gartner, et al. J. Am. Chem. Soc. 2001, 123, 6961.

[0210] In a preferred embodiment of the present new invention a chemical cross-linking is performed by

Reductive Amination






[0211] The reductive amination can essentially be performed as described by:

[0212] Z. J. Gartner, et al. Angew. Chem. Int. Ed. 2002, 41, 1796.

[0213] In a preferred embodiment of the present new invention a chemical cross-linking is performed by

Amine Acylation




[0214] The Amine acylation can essentially be performed as described by:

[0215] Z. J. Gartner, et al. Angew. Chem. Int. Ed. 2002, 41, 1796.

[0216] In a preferred embodiment of the present new invention a chemical cross-linking is performed by

Phosphoramidate formation





[0217] The Phosphoramidate formation can essentially be performed as described by:

[0218] Luther A, et al. Nature 1998, 396:245-248.

[0219] In a preferred embodiment of the present new invention a chemical cross-linking is performed by

Aldol Condensation




[0220] The Aldol condensation reaction can essentially be performed as described by:


[0221] Zhuo Tang et al. Angew. Chem. Int. Ed. 2007, 46, 7297-7300

[0222] In a preferred embodiment of the present new invention a chemical cross-linking is performed by


Cycloaddition Crosslinks
Diels-Alder Cycloaddition




-continued



1,3-Dipolar Cycloaddition



Huisgen Cycloaddition



[0223] The Cycloaddition reactions can essentially be performed as described by:

[0224] Buller et al. Bioorganic & Medicinal Chemistry Letters 18 (2008) 5926-5931

[0225] Fujimoto K, J Am Chem Soc 2000, 122:5646-5647.


[0226] Gartner Z. J. et al. Angew Chem Int Ed Engl 2002, 41:1796-1800.

[0227] Gartner Z. J. et al. Angew Chem Int Ed Engl 2003, 42:1370-1375.

[0228] Poulin-Kerstien A. T. et al. J Am Chem Soc 2003, 125:15811-15821.

[0229] In a preferred embodiment of the present new invention a chemical cross-linking is performed by

Disulfide Crosslinks

Tr = Trityl = C(Ph)₃


[0230] The Disulfide crosslink can essentially be performed as described by:

[0231] Mays, J. R. et al. Tetrahedron Lett. 2007, 48, 4579.

[0232] Theodoropoulos, D. et al. Journal of Medicinal Chemistry, 1985, vol. 28, 10, p. 1536-1539

[0233] Lorenz, Katrin B. et al. Journal of Organic Chemistry, 2004, vol. 69, 11 p. 3917-3927

[0234] In a preferred embodiment of the present new invention a chemical cross-linking is performed by




[0235] The urea crosslink can essentially be performed as described by:

[0236] EP1809743B1 (Vipergen)

[0237] In a preferred embodiment of the present new invention a chemical cross-linking is performed by

Olefination Crosslinks

Wittig Olefination




[0238] The Wittig olefination reaction can be performed as described by:

[0239] Gartner Z. J. et al. Angew Chem Int Ed Engl 2002, 41:1796-1800

[0240] In a preferred embodiment of the present new invention a chemical cross-linking is performed by

Olefination Crosslinks
Wittig Olefination




[0241] The Wittig olefination reaction can essentially be performed as described by:


[0242] Gartner Z. J. et al. Angew Chem Int Ed Engl 2002, 41:1796-1800.

[0243] In a preferred embodiment of the present new invention a chemical cross-linking is performed by


Transition metal catalysed crosslink
Heck Coupling




Sonogashira Coupling




-continued



Suzuki Coupling



Alken-Alkyn Coupling



[0244] Transition metal catalysed reactions can essentially be performed as described by:

[0245] Czapinski J. L. et al. J Am Chem Soc 2001, 123: 8618-8619.


[0246] Gartner Z. J. et al. Angew Chem Int Ed Engl 2002, 41: 1796-1800

[0247] Calderone C. T. et al. Angew Chem Int Ed Engl 2005, 44: 1-5

[0248] Kanan M. W. et al. Nature 431, 545-549, 2004

[0249] In a preferred embodiment of the present new invention a chemical cross-linking is performed by

Photo Crosslinking



[0250] Photo crosslinking can essentially be performed as described by:

[0251] Quamrul, A. et al. Bioorganic & Medicinal Chemistry Letters 18 (2008) 5923-5925

[0252] Weber, T. et al. Journal of the American Chemical Society; 117; 11; (1995); 3084-3095

[0253] Chee, G. et al. Bioorganic and Medicinal Chemistry; 18; 2; (2010); 830-838

[0254] Nassal, M. Journal of the American Chemical Society; 106; (1984); 7540-7545

[0255] Pandurangi, R. S. et al. Bioorganic Chemistry; 25; 2; (1997); 77-87

[0256] Patent; KENT STATE UNIVERSITY; LELAND STANFORD JUNIOR UNIVERSITY; US2010/29952; (2010); (A1)

[0257] In short, the skilled person is aware of numerous different ways of fusing the nucleic acid molecules of a

B-structure and a T-structure which are both present within the same individual compartment.

[0258] As understood by the skilled person in the present context—it is preferred that there is no herein significant fusion of the nucleic acid molecules of a B-structure and a T-structure in the steps (iii) and (iv) of the method of the first aspect—said in other words, there is preferably no significant creation of the BT_{Fused} -structures before step (v).

Step (Vi) of First Aspect

[0259] Step (vi) reads:

“combining the content of the individual compartments of step (v).”

[0260] It is evident that the “combining the content of the individual compartments of step (v)” is done in a suitable way depending on the in vitro compartmentalization system used in step (iv).

[0261] For instance, if the in vitro compartmentalization system used in step (iv) is a micro titer plate like format (see above) the content of the individual wells are simply combined (put together).

[0262] For instance, if the in vitro compartmentalization system used in step (iv) is a suitable water-in-oil emulsion—the individual oil compartments may simply be disrupted by e.g. centrifugation, increase the temperature or by adding a suitable organic solvent.

[0263] In short, in the present context it is routine work for the skilled person to combine the content of the individual compartments of step (v).

[0264] Step (vi) further reads:

“... under conditions wherein there is no fusing of the nucleic acid molecules of a B-structure and a T-structure—i.e. there is not created any new BT_{Fused} -structure not already created in step (v)—in order to get a library of BT_{Fused} -structures.”

[0265] As discussed above—one may say that the enrichment for the herein wanted BT_{Fused} -structures has already been obtained in the steps above—accordingly, as evident to the skilled person one is in this step not interested in creating more “new” BT_{Fused} -structures as such.

[0266] It is routine work for the skilled person to perform step (vi) under such conditions—for instance, if a ligase has been used in step (v) to obtain the wanted BT_{Fused} -structures, this ligase could be inactivated (e.g. by properly raising the temperature) before step (vi) is performed.

[0267] In view of the discussions herein—it is evident for the skilled person that this step (vi) will result in a library of BT_{Fused} -structures.

[0268] As evident to the skilled person—this library of BT_{Fused} -structures may be described as an enriched library of species of BT_{Fused} -structures, comprising nucleic acid sequence information allowing to identify the binding entity and the target—i.e. the sequence information of step (i) and (ii), originating from binding pairs of target and binder entity when compared to BT_{Fused} -structures originating from non-binding pairs of target and binder entity.

Optional Step (Vii)—i.e. Subsequently Use of the Enriched Library of Step (Iv) of First Aspect.

[0269] As discussed above—step (vii) is an optional step.

[0270] As described herein once one has obtained the enriched library of step (vi) one may use this library in different ways according to art—e.g. the enriched library may be considered as an enriched in vitro display library that e.g. can be used in a second round of selection/enrichment or one may

identify the structure of a specific binding entity of interest directly from the enriched library of step (vi).

[0271] Accordingly, an embodiment of the invention relates to the method as described herein, wherein there is an extra step (vii) comprising use the enriched library of step (vi) to identify at least one individual binding entity that binds to at least one target of interest.

Purification of Fused Genotypes:

[0272] In a suitable embodiment of the present invention the fused genotypes (i.e. the BT_{Fused} -structures) may be purified.

[0273] The skilled person in the art can routinely identify numerous different strategies in order to purify the fused genotypes, without being limited for example by: agarose gel electrophoresis, polyacrylamid gel electrophoresis, spun-columns, enzymatic treatment, HPLC purified, affinity purified or capillary electrophoresis (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press)

[0274] In a preferred embodiment of the present invention the fused genotypes are purified post compartmentalization e.g. by gel purification or enzymatic degradation of undesired nucleic acid species. For a skilled person in the art it is evident to design such procedures. For example in the case of using overlap PCR for genotype fusing the size of the genotypes may be chosen to facilitate gel purification e.g. the length of the display library genotype could be chosen to around 250 bp and the length of the target genotype could be around 100 bp and the overlap region to be around 20 bases, the resulting fused genotypes will then be around 330 bp which are easily separated and purified from the original un-fused species by standard agarose gels electrophoresis or polyacrylamide gel electrophoresis. Furthermore, unused primers and ssDNA originating from primer extension using un-fused genotypes as templates may conveniently be degraded enzymatically e.g. by ExoSAP-IT (Amersham Biosciences). Another example in the case of using ligase or chemical crosslinking or transient linking for genotype fusing the size of the genotypes may be chosen to facilitate gel purification e.g. the length of the display library genotype could be chosen to around 250 bp and the length of the target genotype could be around 100 bp, the resulting fused genotypes will then be around 350 bp which are easily separated and purified from the original un-fused species by standard agarose gels electrophoresis or polyacrylamide gel electrophoresis

[0275] In a preferred embodiment of the present invention the fused genotypes is gel purified.

[0276] In short, the skilled person is aware of numerous different ways of purifying fused nucleic acid molecules of a B-structure and a T-structure which was both present within the same individual compartment.

Polish Fused Genotypes:

[0277] In a preferred embodiment of the present invention the fused genotype may be polished (in cases where the genotypes are not fused by an approach compatible with DNA amplification), i.e. to form an amplifiable bond between the two genotypes in the fused genomes—an amplifiable bond is a phosphodiester bond (or alike) between a 3' end of one genotype with a 5' end of the other genotype in the fused genotype.

[0278] The skilled person in the art can routinely identify numerous different strategies in order to purify the fused genotypes, for example without being limited: enzymatically (e.g. *E. coli* DNA Ligase, Taq DNA Ligase, 9° NTM DNA Ligase, T4 DNA Ligase, T4 RNA Ligase 1 (ssRNA Ligase), T4 RNA Ligase 2 (dsRNA Ligase), T4 RNA Ligase 2, truncated) or chemically.

[0279] As evident to the skilled person—correct phosphodiester bond (or alike) formation between cognate genotypes (genotypes originating from the same compartment) post compartmentalization is easily controlled because these are pseudo-intramolecular reaction, thus, independent of concentration of the genotypes. In contrast, incorrect phosphodiester bond (or alike) formation between non-cognate genotypes is an intermolecular reaction, thus, dependent on the concentration of the genotypes.

[0280] In a preferred embodiment of the present invention a DNA ligase is used for polishing.

[0281] In short, the skilled person is aware of numerous different ways of polishing fused nucleic acid molecules of a B-structure and a T-structure which were both present within the same individual compartment.

Removal or Inactivation of the Target Attached to the Nucleic Acid

[0282] In a preferred embodiment of the present invention the target attached to the fused genotypes may be removed or inactivated.

[0283] The skilled person in the art can routinely identify numerous different strategies in order to remove or inactivate the target attached to the fused genotypes, for example without being limited: heat, protease treatment, 6 M Guanidinium chloride, or linker cleavage in case the target was attached by a cleavable linker,

[0284] In a preferred embodiment of the present invention the target attached to the fused genotypes is removed or inactivated by heat protease treatment, or 6 M Guanidinium chloride.

[0285] In a preferred embodiment of the present invention the target attached to the fused genotypes is removed proteinase K treatment.

[0286] In a preferred embodiment of the present invention the target attached to the fused genotypes is removed displacement by primer extension.

[0287] In short, the skilled person is aware of numerous different ways of removing or destroying the target attached to the fused nucleic acid molecules of a B-structure and a T-structure which were both present within the same individual compartment.

Next Round of ECC:

[0288] In a preferred embodiment of the present invention the fused genotype may be subjected to a next round of ECC, i.e. in a next round the fused genotypes will be fused with the new target's genotype—the new target may be the same or a different type as the previous target.

[0289] The skilled person in the art can appreciate that the enriched library of fused genotypes of step (iv) is an in vitro display library.

[0290] In a preferred embodiment of the present invention the earlier round of ECC target attached to the nucleic acid is removed or destroyed prior to a next round of ECC.

[0291] In a preferred embodiment of the present invention the target in a next round of ECC is the same as in an earlier round of ECC.

[0292] In a preferred embodiment of the present invention the target in a next round of ECC is not the same as in an earlier round of ECC.

[0293] In a preferred embodiment of the present invention the genotype of the target in a next round of ECC is fused to a free terminus end of the original genotype for the binding entity.

[0294] In a preferred embodiment of the present invention the genotype of the target in a next round of ECC is fused to a free terminus end of a target genotype from an earlier round of ECC.

Traditional Selection/Enrichment Methods:

[0295] In a preferred embodiment of the present invention the fused genotype may be subjected to a round of prior art known traditional selection/enrichment methods.

[0296] In a preferred embodiment of the present invention the earlier round of ECC target attached to the nucleic acid is removed or destroyed prior to a round of traditional selection/enrichment methods.

[0297] The skilled person in the art can routinely identify numerous different strategies in order to perform a round of a round of traditional selection/enrichment methods, for example without being limited: EP1809743B1 (Viperger), EP1402024B1 (Nuevolution), EP1423400B1 (David Liu), Nature Chem. Biol. (2009), 5:647-654 (Clark), WO 00/23458 (Harbury), Nature Methods (2006), 3(7), 561-570, 2006 (Miller), Nat. Biotechnol. 2004; 22, 568-574 (Melkko), Nature. (1990); 346(6287), 0 818-822 (Ellington), or Proc Natl Acad Sci USA (1997). 94 (23): 12297-302 (Roberts).

[0298] In short, the skilled person is aware of numerous different ways to perform a round of traditional selection/enrichment methods of the library of fused nucleic acid molecules of a B-structure and a T-structure which was both present within the same individual compartment.

Amplification of Fused Genotypes:

[0299] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes may be amplified—i.e. the BT_{Fused} -structures present in the enriched library of step (vi) may be amplified.

[0300] The skilled person in the art can routinely identify numerous different strategies in order to amplify the nucleic acid in the fused genotypes, for example without being limited: PCR (U.S. Pat. No. 4,683,202; Mullis), Emulsion PCR (Nakano et al., J Biotechnol. 2003; 102(2):117-24), Digital PCR (Vogelstein, B; Kinzler K W (1999). "Digital PCR". Proc Natl Acad Sci USA. 96 (16): 9236-41), NASBA (Compton J. Nucleic acid sequence-based amplification. Nature. 1991; 350(6313):91-2), or Rolling Circle Amplification (American Journal of Pathology. 2001; 159:63-69)

[0301] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the de-compartmentalization step.

[0302] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the compartmentalization step performed by PCR.

[0303] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the compartmentalization step performed by PCR,

where the forward PCR priming site is in the B-structure genotype and the backward priming site is in the T-structure genotype.

[0304] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the compartmentalization step performed by PCR where the forward PCR priming site is in the B-structure genotypes and a part of the backward priming site is in the first T-structure genotypes and the remaining part is in the second T-structure genotype.

[0305] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the compartmentalization step performed by PCR where the forward PCR priming is in the first T-structure genotypes site and part of the backward priming site is in the B-structure genotypes and the remaining part is in the second T-structure genotype.

[0306] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the compartmentalization step performed by PCR where part of the forward PCR priming is in the first T-structure genotypes site and the remaining part of the forward PCR priming is in the B-structure genotypes and a part of the backward priming site is in the B-structure genotypes and the remaining part is in the second T-structure genotype.

[0307] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the compartmentalization step performed by PCR where the forward PCR priming site is in the B-structure genotypes and 30-70% of the backward priming site is in the first T-structure genotypes and the remaining 30-70% is in the second T-structure genotype.

[0308] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the compartmentalization step performed by PCR where the forward PCR priming is in the first T-structure genotypes site and 30-70% of the backward priming site is in the B-structure genotypes and the remaining 30-70% is in the second T-structure genotype.

[0309] In a preferred embodiment of the present invention the nucleic acid in the fused genotypes is amplified subsequently to the compartmentalization step performed by PCR where 30-70% of the forward PCR priming is in the first T-structure genotypes site and the remaining 30-70% of the forward PCR priming is in the B-structure genotypes and 30-70% of the backward priming site is in the B-structure genotypes and the remaining 30-70% is in the second T-structure genotype.

[0310] In short, the skilled person is aware of numerous different ways of amplifying the nucleic acid in fused nucleic acid molecules of a B-structure and a T-structure which was both present within the same individual compartment.

Translation:

[0311] In a preferred embodiment of the present invention the nucleic acid of fused genotypes may be amplified and subjected to a translation process where the library of enriched fused genotypes is translated into a new enriched in vitro display library.

[0312] The skilled person in the art can routinely identify numerous different strategies in order to amplify and subject the fused genotypes to a translation process, for example without being limited: EP1809743B1 (Viperger), EP1423400B1 (David Liu), WO 00/23458 (Harbury), Nature

Methods (2006), 3(7), 561-570, 2006 (Miller), Nature. (1990); 346(6287), 818-822 (Ellington), or Proc Natl Acad Sci USA (1997). 94 (23): 12297-302 (Roberts).

[0313] In short, the skilled person is aware of numerous different ways to amplify and perform a translation process of nucleic acid molecules of a B-structure and a T-structure which was both present within the same individual compartment.

Analysis for Identities and Composition:

[0314] In a preferred embodiment of the present invention the nucleic acid of the fused genotype may be analyzed for identities and composition.

[0315] The skilled person in the art can routinely identify numerous different strategies in order to analyzed for identities of the nucleic acid of the fused genotypes, for example without being limited: sequencing (for review: Metzker, Michael L. (2010). "Sequencing technologies—the next generation". Nat Rev Genet 11 (1): 31-46.), DNA hybridization technologies (Science 270 (5235): 467-470), restriction enzyme digest, PCR, methods in EP1809743B1 (Vipergen), EP1402024B1 (Nuevolution), EP1423400B1 (David Liu), Nature Chem. Biol. (2009), 5:647-654 (Clark), WO 00/23458 (Harbury), Nature Methods (2006), 3(7), 561-570, 2006 (Miller), Nat. Biotechnol. 2004; 22,568-574 (Melkko), Nature. (1990); 346(6287), 818-822 (Ellington), or Proc Natl Acad Sci USA (1997). 94 (23): 12297-302 (Roberts), WO06053571A2 (Rasmussen).

[0316] In a preferred embodiment of the present invention the nucleic acid of the fused genotype may be analyzed for identities and composition by DNA sequencing.

[0317] In a preferred embodiment of the present invention the nucleic acid of the fused genotype may be analyzed for identities and composition by DNA sequencing using the 454 technology (Margulies M, Egholm M, Altman W E, et al (September 2005). "Genome sequencing in microfabricated high-density picolitre reactors". Nature 437 (7057): 376-80).

[0318] In short, the skilled person is aware of numerous different ways for analysis for identities and composition of the nucleic acid of the fused genotype of a B-structure and a T-structure which was both present within the same individual compartment.

A Separate Independent Aspect of the Invention

[0319] A separate independent aspect of the invention is described below.

[0320] As understood by the skilled person—the method of this separate independent aspect of the invention uses the same basic technical principles as described above for the first aspect of the invention and thereto related embodiments.

[0321] In line of this and as understood by the skilled person—specific preferred embodiments of the first aspect of the invention (such as e.g. that the in vitro compartmentalization system of step (iv) is a water-in-oil emulsion system) may also be corresponding preferred embodiments of this separate independent aspect of the invention.

[0322] Accordingly, a separate independent aspect of the invention relates to a method for making an enriched library comprising specific nucleic acid sequence information allowing to identifying at least one binding entity that binds to at least one target wherein the specific binding entity has been present in an in vitro display library and wherein the method comprises the steps of:

(i): making an in vitro display library of at least 100 different binding entities (B_n , ($n=100$ or more)), wherein each binding entity is attached to a nucleic acid molecule and the nucleic acid molecule comprises specific nucleic acid sequence information allowing to identify the binding entity—i.e. once one knows the specific nucleic acid sequence information of the nucleic acid molecule one directly knows the structure of the specific binding entity attached to the nucleic acid molecule—the structure of the binding entity (i.e. phenotype) attached to the nucleic acid molecule (genotype) is herein termed B-structure;

(ii): making structures with one target T attached to an enzyme capable of fusing two DNA molecules, wherein the target is capable of binding to at least one of the binding entities present in the library of step (i)—the structure of the target attached to the enzyme capable of fusing two DNA molecules is herein termed T-structure;

and wherein the method is characterized by that:

(iiia): mixing a solution comprising X (X is a number greater than 10^4) numbers of B-structures of the library of step (i) with a solution comprising Y (Y is a number greater than 10^2) numbers of T-structures of step (ii) under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do and wherein one gets binding of at least one of the binding entities to at least one target thereby creating a complex comprising a B-structure bound to a T-structure (herein termed $B_{BoundTo}$ T-structure);

(iiib): mixing to the solution of step (iiia) a solution comprising at least 2 times more nucleic acid molecules than the Y number of T-structures present in step (iiia), wherein the nucleic acid molecules comprise specific nucleic acid sequence information allowing to identify the specific target (herein termed Target-DNA);

(iv): applying an in vitro compartmentalization system—under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do—wherein the compartmentalization system comprises at least 2 times more individual compartments than the Y number of T-structures present in step (iii) under conditions wherein the B-structures, T-structures, $B_{BoundTo}$ T-structures and Target-DNA enter randomly into the individual compartments; and

(v): fusing the nucleic acid molecules of a B-structure and a Target-DNA which are both present within the same individual compartment—this structure is herein termed BT_{Fused} -structure and the BT_{Fused} -structure comprises the specific nucleic acid sequence information allowing to identify the binding entity of step (i) and the specific nucleic acid sequence information allowing to identify the specific target of step (ii); and

(vi): combining the content of the individual compartments of step (v) under conditions wherein there is no fusing of the nucleic acid molecules of a B-structure and a T-structure—i.e. there is not created any new BT_{Fused} -structure not already created in step (v)—in order to get a library of BT_{Fused} -structures, wherein the library is an enriched library of species of BT_{Fused} -structures originating from binding pairs of target and binder entity when compared to BT_{Fused} -structures originating from nonbinding pairs of target and binder entity.

[0323] As known to the skilled person—suitable examples of an enzyme capable of fusing two DNA molecules are e.g. a ligase or a polymerase.

[0324] In line of above discussion of the first aspect of the invention and herein relates embodiment to this—it may be preferred that the herein relevant nucleic acid molecules are DNA molecules and in line of this it may be preferred that the ligase or polymerase is a DNA ligase or a DNA polymerase.

[0325] As understood by the skilled person in the present context—the fusing of the nucleic acid molecules of a B-structure and a Target-DNA of step (v) of this separate independent aspect of the invention is done by the enzyme capable of fusing two DNA molecules (e.g. a ligase or a polymerase) as present in the T-structure of step (ii) of this separate independent aspect of the invention.

[0326] Contrary to the method of the first aspect as discussed herein (wherein there may be more than one different target T present)—there is only one target T present in this separate independent aspect of the invention.

sequence information allowing to identify the specific target is a PCR amplifiable sequence, since the BT_{Fused}-structure of step (v) can then be PCR amplified.

EXAMPLES

Example 1

Enrichment by Co-Compartmentalization Using Overlap ePCR for Genotype-Genotype Fusion

Spiking Experiment

[0329] For overview see FIG. 2.

Methods

DNA Oligonucleotides Used

[0330] Continuous strand of DNA analogue to the yoctoreactor [Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327]:

```
CGCTAAtggtccctggcagtcctccTTAGCGgaccGACTCcTgctcGAAGACAACGGTgttttacAC
CGTTGTCTTCgagcTgtACCTGCgcAAGTGCgttttacGCACTTgcGCAGGTacTgtGCAT
CgacAAGACCgttttacGGTCTTgtcGATGCacTgGAGTCggtcCTGTTTCGATCTTGGGCG
TAT
vip1461:
ATACGCCCAAGATCGAACAG
vip2501:
x-TGGTCCCTGGCAGTCTCC (x = 5'-biotin-TEG)
vip2504:
CTGTTTCGATCTTGGGCGTATGAGAAGAGCCAGAAACGTGGCTTCAGGCACCAA
GGAAGAC
vip2512:
GCCTTGCCAGCCCGCTCAGGCAAGTCTTACAGCCGATCAGTCTTCCTTGGTGCC
TGAAG
vip2502:
CTGTTTCGATCTTGGGCGTAT
vip2500:
x-GCCTTGCCAGCCCGCTCAG (x = 5' carboxyl)
vip157:
GCCTTGCCAGCCCGCTCAG
vip660:
TGGTCCCTGGCAGTCT
vip1481:
GAACAGGACCGA
vip1471:
CTGTTTCGATCTTGGGCGTAT
```

[0327] Accordingly, the specific nucleic acid sequence information allowing identifying the specific target of the nucleic acid molecules of step (iiib) of this separate independent aspect of the invention may simply be a herein relevant characterizing single sequence.

[0328] In line of above discussion of the first aspect of the invention it is preferred that this specific nucleic acid

Preparation of Yoctoreactor Library

[0331] The library is constructed according to Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327) with the following modification: The splint oligonucleotide vip1481 and the oligonucleotide vip1471 are used for introducing the backward priming site.

Preparation of Known Target Binder (Biotin) Attached to Encoding DNA

[0332] A continuous stranded DNA analogue to the yoc-toreactor library sequences is used for PCR using the vip1461 primer and the vip2501 primer, which has a 5'-biotin. Thus, a 5'-biotin is introduced in the yoc-toreactor DNA analogue.

Protocol

[0333] PCR mixture:

[0334] 50 μ L 2 $\times$ PCR mastermix (40 mM Tris-HCl, 20 mM $(\text{NH}_4)_2\text{SO}_4$, 20 mM KCl, 16 mM MgSO_4 , 0.2% Triton X-100, 0.2 mg/mL BSA, 0.4 mM each of dATP, dTTP, dGTP, and dCTP, pH 8.8 @ 25° C.)

10 μ L 5M Betaine (final conc. 0.5M)

1 μ L 50 μ M vip2501 (final conc 0.5 μ M)

1 μ L 50 μ M vip1461 (final conc. 0.5 μ M)

1 μ L (10^7 molecules) of continuous stranded DNA analogue to the yoc-toreactor

1 μ L (2 u/ μ L) Vent (exo-) polymerase

36 μ L water

[0335] The mixture is subjected to thermal cycling by applying the following program in a PCR machine:

92 degrees 2 min, 25 cycles of (92 degrees 30 seconds, 72 degrees 1 min), 72 degrees 2 min), 72 degrees for 2 min.

[0336] The 185 bp DNA fragment is purified by PAGE purification according to standard procedure (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press) and ethanol precipitated

Preparation of Target DNA (TD)

[0337] The 99-mer target DNA is prepared in a one-step overlapping PCR protocol and subsequently purified on a 10% TBE-PAGE native gel

Protocol

PCR Mixture:

[0338] 50 μ L 2 $\times$ PCR mastermix (40 mM Tris-HCl, 20 mM $(\text{NH}_4)_2\text{SO}_4$, 20 mM KCl, 16 mM MgSO_4 , 0.2% Triton X-100, 0.2 mg/mL BSA, 0.4 mM each of dATP, dTTP, dGTP, and dCTP, pH 8.8 @ 25° C.)

10 μ L 5M Betaine (final conc. 0.5M)

1 μ L 50 μ M vip2500 (final conc 0.5 μ M)

1 μ L 50 μ M vip2502 (final conc. 0.5 μ M)

1 μ L 20 μ M vip2504 (final concentration 0.2 μ M)

1 μ L 20 μ M vip2512 (final concentration 0.2 μ M)

1 μ L (2 u/ μ L) Vent (exo-) polymerase

35 μ L water

[0339] The mixture is subjected to thermal cycling by applying the following program in a PCR machine:

92 degrees 2 min, 25 cycles of (92 degrees 30 seconds, 72 degrees 1 min), 72 degrees for 2 min.

[0340] The 99 bp DNA fragment is purified by PAGE purification according to standard procedure (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press). and ethanol precipitated

DNA—Target Conjugation

Materials

[0341] 1-Ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC, 100 mM stock freshly prepared, Aldrich E6383)

N-Hydroxysulfosuccinimide (s-NHS, 200 mM stock, Aldrich 56485)

Morpholinoethanesulfonic acid, MES buffer pH 6.0, 500 mM beta-Mercaptoethanol, 500 mM stock

PCR product with 5' carboxyl group on lower strand (see above)

Streptavidin protein (AbCam 78833) (Target protein)

Protocol

[0342] The carboxyl modified oligonucleotide (ssDNA) or PCR product with terminal carboxyl can be pre-activated using EDC/s-NHS system prior to reaction with target protein (see refs for examples of activation of various types of carboxyls). Exposure of target protein to EDC may render it inactive, e.g. by chemically modifying tyrosine or cysteine residues. Thus, before mixing the activated DNA-carboxyl with target protein, residual EDC optionally may be quenched by addition of e.g. beta-mercaptoethanol to a final concentration of 20 mM.

Example Pre-Activation Mixture

Final Conc

[0343]

10 μ L DNA (ssDNA or dsDNA)	
10 μ L 500 mM MES, pH 6	100 mM
2.5 μ L 100 mM EDC	5 mM
2.5 μ L 200 mM s-NHS	10 mM
25 μ L water	
Total 50 μ L	

[0344] Carboxylic acid activation is allowed to incubate at 20 C for 15-30 min.

[0345] Optional: To quench residual EDC, 2 μ L of 500 mM beta-mercaptoethanol in water was added (final conc. 20 mM).

[0346] Subsequently, the s-NHS-activated ester should be used immediately.

Test of Pre-Activation

[0347] An aliquot of the preactivation mixture (25 pmoles ssDNA) can be diluted with water and a 1% phenethylamine in MeCN (primary amine that quenches the activated ester). This mixture can be allowed to react for 15 min, followed by precipitated using EtOH. After dissolution in 100 mM triethylammonium acetate (TEAA, pH 7), the DNA product can be analyzed by RP-HPLC using a gradient of MeCN in 100 mM TEAA. Shift from initial retention time to higher retention time indicates 1) transformation of carboxyl->NHS ester and 2) subsequent reaction with amine.

Reaction with Target Protein

[0348] Check enzyme shipment buffer composition for primary amines. This protocol should tolerate the presence of e.g. DTT or EDTA in the protein stock solution, but primary

amines must be removed e.g. by dialysis. Primary amines will quench the activated ester thus abolishing DNA—protein cross-link.

[0349] Otherwise, mix pre-activation mixture and enzyme as concentrated as possible to drive chemical reactions.

[0350] This should be allowed to incubate for 1-2 h at 20°C (possibly overnight in cases of slow reaction) then purify DNA—protein complex by e.g. electrophoresis.

Association Mixture

[0351] A diverse YoctoReactor library consisting of 10^6 different molecules and a total of 10^9 molecules i.e. potential ligands coupled to double stranded (ds) DNA, is spiked with 10^6 biotin molecules coupled to ds DNA (known target binder). The spiked library is mixed with 10^7 molecules streptavidin coupled to dsDNA (target attached to DNA). The molecules in the mixtures are allowed to associate in a total volume of 3 μ l Binding Buffer (PBS, 0.05% tween20, 0.2% BSA for 1 hour at room temperature to reach equilibrium. The concentration of streptavidin (the target) is around 6 μ M which is more than 100 fold more than the reported K_d of the biotin-streptavidin complex of $\sim 10^{-14}$ mol/L, which means that practical all biotin will be streptavidin bound at equilibrium.

PCR Mixture

[0352] 67 mM Tris-HCl (pH 8.8), 16.6 mM NH_4SO_4 , 6.7 mM MgCl_2 , 10 mM 2-mercaptoethanol, 1 mM of each dNTP, 7.5 μ M of each primer (vip157 and vip660), 45 units of Taq polymerase in a total volume of 610 μ l

Emulsion PCR

[0353] Two mL of an emulsion consisting of approx. 5×10^9 compartments per ml is prepared by a method similar to the method described by Dressman et al., 2003.

[0354] The DNA fragments coupled to the target (streptavidin) or ligands (non-binding or binding), respectively, have overlapping regions resulting in the potential assembly of the three fragments to combined fragments i.e. two types of combined fragment may be generated through assembly PCR per mixture; fragment (A) signifies that Streptavidin and Biotin have been present in the same compartment and fragment (B) signifies that Streptavidin and a random library molecule (not biotin linked) have been present in the same compartment. The two types of fragments can be differentiated through sequencing or restriction site digestion.

[0355] One mL and 500 μ l (1.5 mL) continuous phase is prepared by dissolving 4.5% (vol/vol) Span80 in mineral oil, followed by 0.40% (vol/vol) Tween80 and 0.05% (vol/vol) Triton X-100 under constant stirring (1,400 rpm) in a 5 ml round bottom Cryo vial, using a magnetic stirring bar with a pivot ring. The continuous phase is split into two times 600 μ l in separate 5 ml round bottom Cryo vials.

[0356] The aqueous phase is made by adding 597 μ l PCR mixture to the 3 μ l association mixture. Three hundred (300) μ l of the aqueous phases is gradually added (10 μ l every 15 s) to each of the two continuous phases under constant stirring (1400 rpm) using a magnetic stirring bar with a pivot ring. After addition of the aqueous phases, the stirring is continued for 30 min.

[0357] The emulsions are aliquoted into approx. twenty wells of a 96-well PCR plate, each containing 100 μ l. The amplification program comprises of 30 cycles with the fol-

lowing steps: initial denaturation at 92°C. for 2 min; 20 cycles consisting of dsDNA denaturation at 92°C. for 30 s, primer annealing and extension at 72°C. for 2 min and 30 s; and final elongation at 72°C. for 2 min.

Breaking the Emulsion

[0358] The DNA fragments from the emulsion PCR are rescued by pooling the emulsions and centrifuging at 13,000 g for 5 min at 25°C. The oil phase is discarded. Residual mineral oil and surfactants are removed from the emulsion by performing the following extraction twice: add 1 ml of water-saturated diethyl ether, vortex the tube, and dispose of the upper (solvent) phase.

Anticipated Results

[0359] Assuming; equal size spherical compartments, random distribution of molecules and complex in compartments, 100% association of Streptavidin-Biotin complex, no dissociation of Streptavidin-Biotin complex, assuming none or few binders with adequate binding affinity for streptavidin in the yR library, no bias of the PCR reaction, and 100% reaction efficiency in “primer extension” (only one priming site present—no co-compartmentalization) during the PCR cycling and 10 000 fold amplification when both priming sites present (co-compartmentalization).

[0360] After emulsion PCR the theoretical amounts of the different DNA species are:

Streptavidin fragment (100 bp): 10^7 molecules

Streptavidin fragment (100 nt): 20 cycles times 10^7 molecules = 2×10^8 molecules

YoctoReactor fragment (250 bp): 10^9 molecules

YoctoReactor fragment (250 nt): 20 cycles times 10^9 molecules = 2×10^{10} molecules (all original Biotin fragment assumed converted to fused species see below)

[0361] Fused genotype—known binder: Fragment A (Streptavidin-Biotin DNA fragment) (330 bp): $10\,000 \times 10^6$ molecules = 10^{10} (PCR amplification times number of biotin fragments)—the probability under the above mentioned assumption for co-compartmentalize the Biotin fragment with target DNA is 1.

[0362] Fused genotype—non-binder: Fragment B (random co-compartmentalized fragments) (330 bp): $10\,000 \times 10^{-3} \times 10^9$ molecules = 10^{10} (PCR amplification times number of initial yR library molecules)—the probability under the above mentioned assumption for co-compartmentalize a non-binder is $\# \text{target molecules} / \# \text{compartments} = 10^7 / 10^{10} = 10^{-3}$

[0363] Consequently, after this process 50% of the 330 bp fragments contain DNA origination from biotin. Moreover, the 330 bp fragment constitutes about 15 ng and constitutes most of the total double stranded DNA. The single stranded DNA is conveniently removed by ExoSAP-IT (Amersham Biosciences) according to manufactures instructions and the 330 bp fragment is conveniently PAGE purified by standard procedure.

Analysis of Enrichment by DNA Sequencing Using 454 Sequencing Technology

[0364] The 454 sequencing priming sites is introduced by PCR using primers with terminal A and B sequences. The resulting fragment is PAGE purified and submitted for 454 DNA sequencing using manufactures protocol.

[0365] The DNA sequences are analyzed and the frequency of the biotin genotype calculated.

Conclusions

[0366] As can be understood from above—by using the method as described herein one gets an 1000 times enrichment—It is expected from the above calculation that the biotin genotype will be observed with a high frequency ~1 out of 2 whereas each of the assumed non-binding yotoreactor library members will be observed ~1 out of 2 million on average. Consequently, the binder has been enriched 1000 fold over each of the non-binders. This will demonstrate the feasibility of the present new invention.

Example 2

Enrichment by Co-Compartmentalization Using eLigation for Genotype-Genotype fusion

Spiking Experiment

[0367] For overview see FIG. 1.

Methods

DNA Oligonucleotides Used

[0368]

vip1481:	GAACAGGACCGA
vip1471:	CTGTTTCGATCTTGGGCGTAT
vip2513:	ACGCCCAAGATCGAACAG

[0369] Biotin-modified continuous one-stranded DNA analogue to the yotoreactor:

X-

CGCTAATgggtccctggcaggtctccTTAGCGgaccGACTCcTgctcGAAGACAACGGTgttttacAC

CGTTGTCTTCgagcTgtACCTGCgcAAGTGCgttttacGCACCTTgcGCAGGTacTgtGCAT

CgacAAGACCgttttacGGTCTTgtcGATGCacTgGAGTCggtcCTGTTTCGATCTTGGGCG

TAT (x = 5'-biotin-TEG)

[0370] The biotin-modified continuous one-stranded DNA analogue to the yotoreactor may be assembled by smaller oligonucleotides essentially as described in Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327) using a 5'-biotin TEG-modified oligonucleotide in the 5'-position

vip2514:

x-

GCCTTGCCAGCCCGCTCAGGGGAAGGACGTTGGTGTAGAAGCGTTCACCT

GGTGAAGTAT (x = 5' carboxyl)

vip2515:

ACTTCCACCAAGTGAACGCT

vip157:

GCCTTGCCAGCCCGCTCAG

vip660:

TGGTCCCTGGCAGTCT

Preparation of Yotoreactor Library

[0371] The library is constructed according to Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327) with the following modifications:

[0372] 1) The splint oligonucleotide vip1481 and the oligonucleotide vip1471 are used for introducing the backward priming site

[0373] 2) The oligonucleotide vip2513 is used for the dismantling of the yotoreactor by primer extension

Preparation of Known Target Binder (Biotin) Attached to Encoding DNA

[0374] The known target binder (biotin) attached to double-stranded encoding DNA is made by dismantling the biotin-modified continuous one-stranded DNA analogue to the yotoreactor by primer extension using vip2513. (Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327). The primer is chosen, so a 3'-overhang of 2 nt is made. The 3'-overhang will facilitate the subsequent ligation.

[0375] The double stranded DNA fragment is purified by PAGE purification according to standard procedure (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press) and ethanol precipitated.

Preparation of Target DNA (TD)

Protocol

[0376] The 61-mer target DNA is prepared by primer extension and subsequently purified on a 10% TBE-PAGE native gel. The primer is chosen, so a 2 nucleotide 3'-over-

hang, complementary to the 3'-overhang of the yotoreactor 3'-overhang, is made. Furthermore, ligation is enabled by phosphorylating the primer.

Example:

Phosphorylation of Primer

[0377] 2 µL (200 pmol) vip2515

20 µL 10x buffer 0 (50 mM Tris-HCl (pH 7.5 at 37° C.), 10 mM MgCl₂,

100 mM NaCl, 0.1 mg/ml BSA)

2 µl 100 mM ATP (final concentration 2 mM)

10 µL T4 Polynucleotide Kinase (100 u)

[0378] 66 µL Water, nuclease-free

[0379] The phosphorylation reaction is incubated @37 degrees C. for 30 minutes, and the kinase is inactivated by incubation @75 degrees C. for 10 minutes

[0380] The DNA is precipitated by ethanol precipitation, washed in 70% ethanol, and resuspended in 10 μ L TE buffer.

Primer Extension:

[0381] 10 μ L phosphorylated vip2515 (200 pmoles)
10 μ L 10 $\times$ buffer 0 (New England Biolabs)
2 μ L 10 mM dNTP mix (final concentration 0.2 mM of dATP, dCTP, dGTP, and dTTP)

77 μ L water

1 μ L (5 u) Klenow (exo-, 5 u/ μ L)

[0382] The reaction is allowed to proceed for 15 minutes, and the double stranded DNA purified by extraction from a 15% acrylamide gel according to standard procedure (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press). and ethanol precipitated

DNA—Target Conjugation

Materials

[0383] 1-Ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC, 100 mM stock freshly prepared, Aldrich E6383)

N-Hydroxysulfosuccinimide (s-NHS, 200 mM stock, Aldrich 56485)

Morpholinoethanesulfonic acid, MES buffer pH 6.0, 500 mM beta-Mercaptoethanol, 500 mM stock

PCR product with 5' carboxyl group on lower strand (see above)

Streptavidin protein (AbCam 78833) (Target protein)

Protocol

[0384] The carboxyl modified oligonucleotide (ssDNA) or PCR product with terminal carboxyl can be pre-activated using EDC/s-NHS system prior to reaction with target protein (see refs for examples of activation of various types of carboxyls). Exposure of target protein to EDC may render it inactive, e.g. by chemically modifying tyrosine or cysteine residues. Thus, before mixing the activated DNA-carboxyl with target protein, residual EDC optionally may be quenched by addition of e.g. beta-mercaptoethanol to a final concentration of 20 mM.

Example Pre-Activation Mixture

Final Conc

[0385]

10 μ L DNA (ssDNA or dsDNA)	
10 μ L 500 mM MES, pH 6	100 mM
2.5 μ L 100 mM EDC	5 mM
2.5 μ L 200 mM s-NHS	10 mM
25 μ L water	
Total 50 μ L	

[0386] Carboxylic acid activation is allowed to incubate at 20 C for 15-30 min.

[0387] Optional: To quench residual EDC, 2 μ L of 500 mM beta-mercaptoethanol in water was added (final conc. 20 mM).

[0388] Subsequently, the s-NHS-activated ester should be used immediately.

Test of Pre-Activation

[0389] An aliquot of the preactivation mixture (25 pmoles ssDNA) can be diluted with water and a 1% phenethylamine in MeCN (primary amine that quenches the activated ester). This mixture can be allowed to react for 15 min, followed by precipitated using EtOH. After dissolution in 100 mM triethylammonium acetate (TEAA, pH 7), the DNA product can be analyzed by RP-HPLC using a gradient of MeCN in 100 mM TEAA. Shift from initial retention time to higher retention time indicates 1) transformation of carboxyl->NHS ester and 2) subsequent reaction with amine.

Reaction with Target Protein

[0390] Check enzyme shipment buffer composition for primary amines. This protocol should tolerate the presence of e.g. DTT or EDTA in the protein stock solution, but primary amines must be removed e.g. by dialysis. Primary amines will quench the activated ester thus abolishing DNA—protein cross-link.

[0391] Otherwise, mix pre-activation mixture and enzyme as concentrated as possible to drive chemical reactions.

[0392] This should be allowed to incubate for 1-2 h at 20 C (possibly overnight in cases of slow reaction) then purify DNA—protein complex by e.g. electrophoresis.

Association Mixture

[0393] A diverse YoctoReactor library consisting of 10^6 different molecules and a total of 10^9 molecules i.e. potential ligands coupled to double stranded (ds) DNA, is spiked with 10^6 biotin molecules coupled to ds DNA (known target binder). The spiked library is mixed with 10^7 molecules streptavidin coupled to dsDNA (target attached to DNA). The molecules in the mixtures are allowed to associate in a total volume of 3 μ L Binding Buffer (PBS, 0.05% tween20, 0.2% BSA for 1 hour at room temperature to reach equilibrium. The concentration of streptavidin (the target) is around 6 μ M which is more that 100 fold more than the reported K_d of the biotin-streptavidin complex of 10^{-14} mol/L, which means that practically all biotin will be streptavidin bound at equilibrium.

Ligation Mixture

[0394] 1 $\times$ Tag ligation buffer (20 mM Tris-HCl, 25 mM potassium acetate, 10 mM Magnesium Acetate, 1 mM NAD, 10 mM Dithiothreitol 0.1% Triton X-100 pH 7.6 @ 25° C.) is added 2 μ L (40 u/ μ L) Taq DNA ligase in a total volume of 610 μ L.

[0395] 67 mM Tris-HCl (pH 8.8), 16.6 mM NH_4SO_4 , 6.7 mM MgCl_2 , 10 mM 2-mercaptoethanol, 1 mM of each dNTP, 7.5 μ M of each primer (vip157 and vip660), 45 units of Taq polymerase in a total volume of 610 μ L

Ligation in Emulsion

[0396] Two mL of an emulsion consisting of approx. 5×10^9 compartments per ml is prepared by a method similar to the method described by Dressman et al., 2003.

[0397] The DNA fragments coupled to the target (streptavidin) or ligands (non-binding or binding), respectively, are able to be ligated on one strand. i.e. two types of combined fragment may be generated through ligation; fragment (A) signifies that Streptavidin and Biotin have been present in the

same compartment and fragment (B) signifies that Streptavidin and a random library molecule (not biotin linked) have been present in the same compartment. The two types of fragments can be differentiated through sequencing or restriction site digestion.

[0398] One mL and 500 μ L (1.5 mL) continuous phase is prepared by dissolving 4.5% (vol/vol) Span80 in mineral oil, followed by 0.40% (vol/vol) Tween80 and 0.05% (vol/vol) Triton X-100 under constant stirring (1,400 rpm) in a 5 mL round bottom Cryo vial, using a magnetic stirring bar with a pivot ring. The continuous phase is split into two times 600 μ L in separate 5 mL round bottom Cryo vials and is kept ice-cold. The aqueous phase is made by adding 597 μ L ice-cold ligation mixture to the 3 μ L association mixture. Three hundred (300) μ L of the aqueous phase is gradually added (10 μ L every 15 s) to each of the two continuous phases under constant stirring (1400 rpm) using a magnetic stirring bar with a pivot ring. After addition of the aqueous phase, the stirring is continued for 30 min under ice-cold conditions.

[0399] The emulsions are heated to 45 degrees C. and allowed to ligate for one hour.

Breaking the Emulsion

[0400] The ligation mixtures are added 30 μ L 500 mM EDTA each and vortexed briefly. The DNA fragments are rescued by pooling the emulsions and centrifuging at 13,000 g for 5 min at 25° C. The oil phase is discarded. Residual mineral oil and surfactants are removed from the emulsion by performing the following extraction twice: add 1 mL of water-saturated diethyl ether, vortex the tube, and dispose of the upper (solvent) phase.

[0401] The DNA is concentrated by precipitation, is fractionated by size on denaturing 10% polyacrylamide gels and the ligated fragments isolated by PAGE purification according to standard procedure (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press), ethanol precipitated and resuspended in 10 μ L TE buffer.

Amplification of Ligated DNA

[0402] Finally, the ligated fragments are amplified by PCR

Example:

PCR Mixture:

[0403] 10 μ L purified ligated DNA
50 μ L 2 $\times$ PCR mastermix (40 mM Tris-HCl, 20 mM (NH₄)₂SO₄, 20 mM KCl, 16 mM MgSO₄, 0.2% Triton X-100, 0.2 mg/mL BSA, 0.4 mM each of dATP, dTTP, dGTP, and dCTP, pH 8.8 @ 25° C.)

10 μ L 5M Betaine (final conc. 0.5M)

1 μ L 50 μ M vip167 (final conc 0.5 μ M)

1 μ L 50 μ M vip660 (final conc. 0.5 μ M)

1 μ L (2 u/ μ L) Vent (exo-) polymerase

27 μ L water

[0404] The mixture is subjected to thermal cycling by applying the following program in a PCR machine:

92 degrees C., 2 min, 25 cycles of (92 degrees C. 30 seconds, 72 degrees C. 1 min), 72 degrees C. for 2 min.

[0405] The resulting library of DNA fragments is sequenced, and the enrichment for the binding fragment calculated.

Anticipated Results

[0406] Assuming; equal size spherical compartments, random distribution of molecules and complex in compartments, 100% association of Streptavidin-Biotin complex, no dissociation of Streptavidin-Biotin complex, assuming none or few binders with adequate binding affinity for streptavidin in the yR library, no bias of the ligation reaction or the following PCR reaction, —no co-compartmentalization) during the ligation and 100 reaction efficiency when the library and target DNA fragments are co-compartmentalized.

[0407] After emulsion ligation, theoretical amounts of the ligated DNA species are: (all original Biotin fragment assumed converted to fused species see below)

[0408] Ligated genotype—known binder: Fragment A (Streptavidin-Biotin DNA fragment) (250 bp): 10⁶ molecules—the probability under the above mentioned assumption for co-compartmentalize the Biotin fragment with target DNA is 1.

[0409] Fused genotype—non-binder: Fragment B (random co-compartmentalized fragments) (250 bp): 10⁻³×10⁹ molecules=10⁶ (Probability for random co-compartmentalization times number initial yR library molecules)—the probability under the above mentioned assumption for co-compartmentalize a non-binder is #target molecules/# compartments=10⁷/10¹⁰=10⁻³

[0410] The final PCR amplification is expected to be of same efficiency for the two types of molecules, and Consequently, after this process 50% of the 330 by fragments contain DNA originating from the biotin-streptavidin binding.

Analysis of Enrichment by DNA Sequencing Using 454 Sequencing Technology

[0411] The 454 sequencing priming sites is introduced by PCR using primers with terminal A and B sequences. The resulting fragment is PAGE purified and submitted for 454 DNA sequencing using manufactures protocol.

[0412] The DNA sequences are analyzed and the frequency of the biotin genotype calculated.

Conclusions

[0413] As can be understood from above—by using the method as described herein one gets an 1000 times enrichment—It is expected from the above calculation that the biotin genotype will be observed with a high frequency ~1 out 2 whereas each of the assumed non-binding ycoreactor library members will be observed ~1 out of 2 million on average. Consequently, the binder has been enriched 1000 fold over each of the non-binders. This will demonstrate the feasibility of the present new invention.

Examples Below

[0414] All the Examples below were made based on the technical information disclosed above (e.g. in the working examples above) plus based on the common general knowledge of the skilled person.

Example 3

Enrichment by Co-Compartmentalization Using Overlap ePCR for Genotype-Genotype Fusion

[0415] The fundamental principle of ECC, co-compartmentalization of binding partners and fusion of their attached

DNA as a result hereof, was demonstrated, using biotin and streptavidin (SA) as the binding partners. Biotin DNA conjugate (yR_biotin) was subjected to ECC using SA conjugated to DNA (SA_TD001) as the target. As a negative control ECC was run in parallel using SA_TD001 preincubated with biotin as the targets. For overview see FIG. 2.

Methods

DNA Oligonucleotides Applied:

[0416] Applied for continuous strand of DNA analogue to the yoctoreactor (Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327):

CGCTAAtggtccctggcaggtctccTTAGCGgaccGACTCcTgctcGAAGACAACGGTgttttacAC

CGTTGTCTTCgagcTgtACCTGCgcAAGTGCgttttacGCACTTgcGCAGGTacTgtGCAT

CgacAAGACCgttttacGGTCTTgtcGATGCacTgGAGTCggtcCTGTTCGATCTTGGGCG

TAT

vip1481:
GAACAGGACCGA

vip1471:
CTGTTCGATCTTGGGCGTAT

Applied for yR_biotin
vip1461:
ATACGCCCAAGATCGAACAG

vip2501:
x-TGGTCCCTGGCAGTCTCC (x = biotin-TEG)

Applied for e PCR
vip157:
GCCTTGCCAGCCCGCTCAG

vip660:
TGGTCCCTGGCAGTCT

Applied for TD001
vip2500:
x-GCCTTGCCAGCCCGCTCAG (x = carboxyl modification)

vip2502:
CTGTTCGATCTTGGGCGTAT

vip2512:
GCCTTGCCAGCCCGCTCAGGCAAGTCTTACAGCCGATCAGTCTTCCTTGGTGCC

TGAAG

vip2507:
CTGTTCGATCTTGGGCGTATTGTTTTAGCTGCCCCAACTCCTTCAGGCACCAAG

GAAGAC

Applied for Rescue PCR
vip2549:
GCAAGTCTTACAGCCGATCA

vip660:
TGGTCCCTGGCAGTCT

Preparation of yR

[0417] Continuous strand of DNA analogue to the yoctoreactor was constructed as described in example 1

Preparation of yR_Biotin

[0418] Preparation of known target binder (biotin) attached to yR was constructed as described in example 1

Preparation of Target DNA (TD001)

[0419] The TD001 was prepared as described in example 1 except the oligo vip2507 being applied instead of Vip2504.

Materials

[0420] MOPS 3-(N-Morpholino) propanesulfonic acid (Sigma-Aldrich)

Silicone polyether/cyclopentasiloxane (Dow Corning, DC5225C)

Cyclopentasiloxane/trimethylsiloxysilicate (Dow Corning, DC749)

[0421] AR20 silicone oil (Sigma-Aldrich)

1-Ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC, 100 mM stock freshly prepared, Aldrich E6383)

N-Hydroxysulfosuccinimide (s-NHS, 200 mM stock, Aldrich 56485)

Morpholinoethanesulfonic acid, MES buffer pH 6.0, 500 mM beta-Mercaptoethanol, 500 mM stock

PCR product with 5' carboxyl group on lower strand (see above)

Streptavidin protein (AbCam 78833) (Target protein)
Slide-A-lyzer mini (Pierce)

TissueLyzer II (Qiagen)

DNA—Target Conjugation

Pre-Activation

[0422] Pre-activation was done by mixing 5.4 μ l TD001 [9.3 μ M] with 1 μ l MOPS pH 6 [1 M], 1 μ l EDC [50 mM], 1 μ l s-NHS [100 mM] and 1.6 μ l water.

[0423] Carboxylic acid activation is allowed to incubate at 20° C. for 30 min.

[0424] To quench residual EDC, 1 μ l of 250 mM beta-mercaptoethanol in water was added.

Reaction with Target Protein

[0425] Prior to conjugation, SA was dialyzed 2 times 30 min against Dialysis Buffer (10 mM MOPS (pH 8), 50 mM NaCl) using Slide-A-Lyzer mini dialysis device according to manufactures instructions (Pierce).

[0426] 5 μ l of dialyzed SA [58 μ M] was added to 1.6 μ l MOPS pH 6 [1 M], 1.6 μ l NaCl in water [1 M] and 11 μ l TD001 [4.6 μ M]. The reaction was allowed for 2 hours at 20° C.

[0427] To quench the reaction, 2 μ l Tris pH 8 [1 M] was added. The SA_TD001 conjugate was isolated from reactants by PAGE from a 6% TBE gel that was run for 40 min at 200V.

[0428] The bands are extracted 3 times in 500 μ l Extraction Buffer (50 mM Tris pH8, 150 mM NaCl, 0.1% Tween20) at 4° C. (30 min/o.n./30 min).

[0429] Residual gel was removed by filtration, and the samples concentrated in a Microcon YM30 device according to manufactures instructions (Millipore). The concentration of the conjugate was estimated to be 0.38 μ M by measuring the DNA concentration using Picogreen according to manufactures instructions (Molecular Probes).

Association Reactions (Binding Reaction)

[0430] Prior to yR_biotin and SA_TD001 binding, 6e8 molecules SA_TD001 molecules/ μ l in a total volume of 50 μ l Association Buffer (10 mM Tris-HCl (pH 7.8), 0.05% Triton-X100.) was incubated with or without 1 μ M biotin (6e11 molecules biotin/ μ l) for 30 min at 20° C.

[0431] The following binding reactions were made in Association Buffer:

[0432] 1) 3e8 yR_biotin molecules/ μ l and 3e8 SA_TD001 molecules/ μ l in a total volume of 50 μ l, using SA_TD001 that had not been pre-incubated with biotin

[0433] 2) 3e8 yR_biotin molecules/ μ l and 3e8 SA_TD001 molecules/ μ l in a total volume of 50 μ l, using SA_TD001 that had been pre-incubated with biotin.

[0434] The binding reaction was incubated for 1 h at 20° C. and hereafter diluted to a concentration of 3e6 molecules/ μ l of yR_biotin and 3e6 molecules/ μ l SA_TD001 in Association Buffer.

Emulsion PCR (ePCR)

[0435] Assembly of yR and TD001 and amplification of yR_TD001 fusion molecule in emulsion using PCR.

Continuous Phase

[0436] Continuous phase was prepared as described by (Turner and Hurles, Nat Protoc. 2009; 4(12): 1771-1783). 1200 μ l continuous phase was made per reaction
480 μ l Silicone polyether/cyclopentasiloxane (DC5225C)

360 μ l Cyclopentasiloxane/trimethylsiloxysilicate (DC749)

[0437] 360 μ l AR20 silicone oil

PCR Aqueous Phase

[0438] 600 μ l PCR aq. was made per reaction:

60 μ l Pfu buffer (10 $\times$)

12 μ l BSA (50 mg/ml)

12 μ l dNTP (10 mM)

3 μ l Vip157 (100 μ M)

3 μ l Vip660 (100 μ M)

[0439] 4 μ l Pfu-turbo (2.5 u/ μ l)

446 μ l water

60 μ l template the resulting concentration of yR_biotin and SA_TD001 are 3e5 molecules/ μ l of each

Emulsification

[0440] In a 2 ml Eppendorf tube 1000 μ l continuous phase and 500 μ l PCR phase, and a 5 mm steel bead were added per reaction

[0441] The reaction was emulsified by mixing for 8 min at 30 Hz in a Tissuelyser II at 20° C. 100 μ l emulsion was added per PCR tube and the mixture was subjected to thermal cycling by applying the following program in a PCR machine:

92° C. for 2 min, 30 cycles of (92° C. 30 seconds, 55° C. for 1 min and 72° C. for 1.5 min), 72° C. for 5 min.

Recovery of DNA

[0442] The emulsion was broken by adding 100 μ l 1-butanol per PCR tube. The contents of 8 PCR tubes per condition were pooled and 600 μ l NaCl in water [4 M] was added. The content was mixed by vortexing for 10 sec at max speed, and the organic phase was removed after centrifugation at 14000 g for 1 min. Another 800 μ l 1-butanol was added to the pooled PCR product and the vortexing and centrifugation step was repeated. The extraction with 1-butanol was repeated one more time.

[0443] The DNA was further purified by PCR purification columns (Macherey-Nagel) according to manufactures instructions. Elute with 50 μ l elution buffer per condition

[0444] The eluted DNA was diluted 20 fold in Dilution Buffer (10 mM Tris (pH 7.8), 20 mM NaCl, 0.1% Triton-X100) prior to the rescue PCR.

Rescue PCR

[0445] Amplification of yR_TD001 fusion molecule

PCR Mixture Per Reaction:

[0446] 50 μ L 2 $\times$ PCR mastermix (40 mM Tris-HCl, 20 mM (NH₄)₂SO₄, 20 mM KCl, 16 mM MgSO₄, 0.2% Triton X-100, 0.2 mg/mL BSA, 0.4 mM each of dATP, dTTP, dGTP, and dCTP, pH 8.8 at 25° C.)

1 μ L 50 μ M vip2549 (final conc 0.5 μ M)

1 μ L 50 μ M vip660 (final conc. 0.5 μ M)

1 μ L (2 u/ μ L) Vent (exo-) polymerase

10 μ L template

37 μ L water

[0447] The mixture was subjected to thermal cycling by applying the following program in a PCR machine:

92° C. for 2 min, 20 cycles of (92° C. 30 seconds, 55° C. for 1 min and 72° C. for 1.5 min), 72° C. for 5 min.

Results

[0448] The fusion product yR_TD001 has a predicted length of 245 bp. The DNA products were visualized on a 10% TBE PAGE run for 40 min. at 200V. The results showed the presence of a band with the expected size, if the association reaction was performed without pre-incubation with biotin (lane 1) and the absence of a band if association reaction was performed on molecules that had been pre-incubated with biotin (lane 2), see FIG. 4.

Conclusion

[0449] This result demonstrated co-compartmentalization of binding partners and fusion of their attached DNA as a result hereof.

Example 4

Enrichment by Co-Compartmentalization Using Overlap ePCR for Genotype-Genotype Fusion

Spiking Experiment

[0450] ECC was demonstrated by enriching for yR_biotin that was spiked into a diverse yR library using SA_TD001 as the target. As a negative control ECC was run in parallel using SA_TD001 preincubated with biotin as the targets.

Methods

DNA Oligonucleotides Applied

[0451] Applied for continuous strand of DNA analogue to the ycoreactor (Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327):

CGCTAAtgggtccctggcagtcctccTTAGCGgaccGACTCcTgctcGAAGACAACGGTggttttacAC

CGTTGTCTTCgagcTgtACCTGCGcAAGTGCggttttacGCACCTTgcGCAGGTacTgtGCAT

CgacAAGACCGgttttacGGTCTTgtcGATGCacTgGAGTCggtcCTGTTCGATCTTGGGCG

TAT

vip1481:
GAACAGGACCGA

vip1471:
CTGTTTCGATCTTGGGCGTAT

Applied for yR_biotin
vip1461:
ATACGCCCAAGATCGAACAG

vip2501:
x-TGGTCCTGGCAGTCTCC (x = biotin-TEG)

Applied for e PCR
vip157:
GCCTTGCCAGCCCGCTCAG

vip660:
TGGTCCCTGGCAGTCT

Applied for TD001
vip2500:
x-GCCTTGCCAGCCCGCTCAG (x = carboxyl modification)

vip2502:
CTGTTTCGATCTTGGGCGTAT

vip2512:
GCCTTGCCAGCCCGCTCAGGCAAGTCTTACAGCCGATCAGTCTTCCTTGGTGCC

TGAAG

vip2507:
CTGTTTCGATCTTGGGCGTATTGTTTTAGCTGCCCAACTCCTTCAGGCACCAAG

GAAGAC

Applied for Rescue PCR
vip2549:
GCAAGTCTTACAGCCGATCA

-continued

vip660:
TGGTCCCTGGCAGTCT

Applied for PCR of a yR diverse library
vip341:
TGGTCCCTGGCAGTCTCC

vip1461:
ATACGCCCAAGATCGAACAG

Applied for 454 PCR
vip2593:
CCTATCCCCTGTGTGCCTTGGCAGTCTCAGGTCTTCTTGGTGCTGAAG

vip2465:
CCATCTCATCCCTGCGTGTCTCCGACTCAGAGGTGGTCCCTGGCAGTCTCC

vip2467:
CCATCTCATCCCTGCGTGTCTCCGACTCAGATCGTGGTCCCTGGCAGTCTCC

vip2468:
CCATCTCATCCCTGCGTGTCTCCGACTCAGATGCTGGTCCCTGGCAGTCTCC

vip2469:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCACTGGTCCCTGGCAGTCTCC

vip2470:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCAGATGGTCCCTGGCAGTCTCC

vip2471:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCCATTGGTCCCTGGCAGTCTCC

Preparation of yR_Biotin

[0452] yR_biotin was prepared as described in example 1

Preparation of a Diverse Yoctoreactor Library

[0453] The yR library was essentially constructed as described by (Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327). The diverse yR library was PCR amplified by the following method;

PCR Mixture Per Reaction:

[0454] 50 μ L 2 $\times$ PCR mastermix (40 mM Tris-HCl, 20 mM (NH₄)₂SO₄, 20 mM KCl, 16 mM MgSO₄, 0.2% Triton X-100, 0.2 mg/mL BSA, 0.4 mM each of dATP, dTTP, dGTP, and dCTP, pH 8.8 at 25° C.)

10 μ L 5M Betaine (final conc. 0.5M)

1 μ L 50 μ M vip341 (final conc 0.5 μ M)

1 μ L 50 μ M vip1461 (final conc. 0.5 μ M)

1 μ L (10⁸ molecules) of continuous stranded DNA analogue to the yoctoreactor

1 μ L (2 u/ μ L) Vent (exo-) polymerase

36 μ L water

[0455] The mixture was subjected to thermal cycling by applying the following program in a PCR machine:

92° C. for 2 min, 15 cycles of (92° C. for 30 seconds, 72° C. for 1 min), 72° C. for 2 min.

[0456] The 185 bp DNA fragment was purified by PAGE purification according to standard procedure (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press) and ethanol precipitated

Association Reactions (Binding Reactions)

[0457] Binding reactions was performed as described in example 3, with the following changes.

[0458] Prior to yR_biotin and SA_TD001 binding, 6e8 molecules SA_TD001 molecules/ μ L in a total volume of 50 μ L association buffer was incubated with or without 1 μ M biotin (6e11 molecules biotin/ μ L) for 30 min at 20° C.

[0459] The following association reactions were made in Association Buffer:

[0460] 1) 3e7 yR_biotin molecules/ μ L and 3e8 SA_TD001 molecules/ μ L in a total volume of 50 μ L, using SA_TD001 that had not been pre-incubated with biotin

[0461] 2) 3e7 yR_biotin molecules/ μ L and 3e8 SA_TD001 molecules/ μ L in a total volume of 50 μ L, using SA_TD001 that had been pre-incubated with biotin.

[0462] Binding reactions were incubated for 1 h at 20° C.

[0463] Hereafter the following conditions were setup:

[0464] A) Without biotin pre-incubation: A 1000 fold dilution of binding reaction (1) to a concentration of 3e4 molecules/ μ L of yR_biotin and 3e5 molecules/ μ L SA_TD001 in Association Buffer containing 3e7 molecules yR library/ μ L (final concentration).

[0465] B) With biotin pre-incubation: A 1000 fold dilution of association reaction (2) to a concentration of 3e4 molecules/ μ L of yR_biotin and 3e5 molecules/ μ L SA_TD001 in Association Buffer containing 3e7 molecules yR library/ μ L (final concentration).

[0466] C) Without yR-biotin: Association Buffer containing a 3e7 yR library molecules/ μ L and 3e5 molecules/ μ L SA_TD001 in Association Buffer was made

Emulsion PCR

[0467] ePCR was performed as described in example 3, but performed in duplicate and with 40 PCR cycles

Recovery of DNA

[0468] Breaking of emulsions was performed as described in example 3, but performed by pooling the emulsions from 16 PCR tubes and eluting with 100 μ L elution buffer per condition

Rescue PCR

[0469] Rescue PCR was performed as described in example 3, but with the following thermal profile during PCR amplification; 92° C. for 2 min, 20 cycles of (92° C. for 30 seconds, 72° C. for 1.5 min), 72° C. for 5 min.

Preparation for 454-Sequencing

[0470] PCR protocol for amplification of yR_TD001 fusion molecules hereby including 454 sequence tags into the sequences leading to a predicted size of 309 bp. For each condition and duplicate a unique forward primer was applied.

PCR Mixture Per Reaction:

[0471] 50 μ L 2 $\times$ PCR mastermix (40 mM Tris-HCl, 20 mM (NH₄)₂SO₄, 20 mM KCl, 16 mM MgSO₄, 0.2% Triton X-100, 0.2 mg/mL BSA, 0.4 mM each of dATP, dTTP, dGTP, and dCTP, pH 8.8 at 25° C.)

10 μ L 5M Betaine (final conc. 0.5M)

1 μ L 50 μ M vip2593 (final conc 0.5 μ M)

2 μ L 25 μ M vip2465, vip2467, vip2468, vip2469, vip2470 or vip2471 (final conc. 0.5 μ M)

5 μ L Template from each condition and duplicate

1 μ L (2 u/ μ L) Vent (exo-) polymerase

31 μ L water

[0472] The DNA was purified on PCR purification columns (Macherey-Nagel) according to manufactures instructions and the DNA concentrations were determined using a spectrophotometer (Eppendorf). The concentrations were adjusted upon comparative visual inspection of the products on 10% TBE gels that were run for 40 min at 200V. The DNA products were pooled so that all DNA products were represented by similar amounts of DNA. The pooled DNA was run on a 10% TBE PAGE gel and a DNA fragment of approx. 309 bp was purified by PAGE purification according to standard procedure (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press), and ethanol precipitated.

454-Sequencing

[0473] 454-sequencing was performed as described by (Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327).

Results

[0474] The sequencing results showed, see FIG. 5, that although the yR_biotin was spiked into the yR library in a 1000 fold lower concentration than the yR library the percentages of yR_biotin counts were:

37.6% and 32% for the duplicates in condition (A) i.e. enrichment without pre-incubation of SA_TD001 with biotin. 0.30% and 0.72% for the duplicates in condition (B) i.e. enrichment with pre-incubation of SA_TD001 with biotin. 0.02% and 0.05% for the duplicates in condition (C) i.e. enrichment without yR_biotin included in the sample.

[0475] Consequently, more than 300 fold enrichment of yR_biotin was observed using SA_TD001 as the target. In contrast, the negative control target, SA_TD001 preincubated with biotin, provided a 3-7 fold enrichment.

CONCLUSION

[0476] ECC was demonstrated by enriching for yR_biotin that was spiked into a diverse yR library using SA_TD001 as the target.

Example 5

Enrichment by Co-Compartmentalization Using eLigation for Genotype-Genotype Fusion

[0477] The fundamental principle of ECC, co-compartmentalization of binding partners and fusion of their attached DNA as a result hereof, was demonstrated, using desthiobiotin (desBio) and streptavidin (SA) as the binding partners. Desthiobiotin DNA conjugate (yR_desBio) was subjected to ECC using SA conjugated to DNA (SA_TD002) as the target. As a negative control ECC was run in parallel using SA_TD002 preincubated with biotin as the targets.

[0478] For overview see FIG. 1.

Methods

[0479] DNA oligonucleotides used are described in example 2. In addition, the following were applied:

Applied for yR labeled with desthiobiotin (desBio_yR):
vip2815:
x-TGGTCCCTGGCAGTCTCC (x = desthiobiotin)

vip2535:
CACACGATGGCAATGCATTCTTCGCTGCCATTCTG

Applied for rescue PCR:
vip660:
TGGTCCCTGGCAGTCT

vip2824:
CGATGTCTGAGGTGGAAGT

Applied for 'Scavenger DNA':
vip2554:
GGCAAGTGATTGTCCATGTGCATGAGAAGAGGCCACATT

vip2555:
CACATGGACAATCACTTGCC

vip2556:
AATGTGGGCCTCTTCTCATG

Applied for TD002
vip2528:
TCCACATCCTCCAGTTCA

vip2529:
ACTTCCACCTCAGGACATCGAGCTGGAGCTTGCTGTTAGC

vip2530:
AGGTTTCGCTCCCTCCTTAAGTCAGGAGGATGTGACACCAA

vip2531:
CGATGTCTGAGGTGGAAGTTGAACCTGGAGGATGTGGACA

vip2532:
CTTAAGGAGGGAGCGAACCTGCTAACAGCAAGCTCCAGCT

-continued

vip2558:
x-TTGGTGTACATCCTCCTGA
(x = C6-amino modification)

Preparation of yR Labeled with Desthiobiotin (desBio_yR)

[0480] A 5'-desthiobiotin was introduced in the yR analogue by using primers vip2815 and vip2535 in PCR with a continuous stranded DNA analogue to the yotoreactor library sequences as template DNA (Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327). To create a 2 bp overhang suitable for ligation to target DNA conjugated with streptavidin, the PCR product was digested with BseMI.

Preparation of Target DNA (TD002)

Protocol

[0481] TD002 (98 bp double stranded DNA with a GA nucleotide overhang and 5' carboxyl group on the lower strand) was assembled by ligation of phosphorylated oligonucleotides vip2528, vip2529, vip2530, vip2531 and vip2532 and the non-phosphorylated oligonucleotide vip2558. Phosphorylation with T4 Polynucleotide Kinase and ligation with T4 DNA ligase was performed according to manufactures instructions (Fermentas). The double stranded DNA fragment was purified by PAGE purification according to standard procedure (Molecular Cloning: A Laboratory Manual (3-Volume Set), 3rd Edition, 2001-01 by Joseph Sambrook, David W. Russell, Publisher: Cold Spring Harbor Laboratory Press) and precipitated with ethanol. Prior to ligation vip2559 was modified to have a 5' carboxyl group. Thus, simple C6-amino modification was interchanged to a carboxylic acid by treatment with disuccinimidylsuberate (DSS, C8-di-NHS ester, Pierce #21580). The oligonucleotide was treated with 40 mM DSS in HEPBS buffer pH 9 in a water—NMP 1:1 mixture over night followed by treatment with LiOH to hydrolyse the remaining NHS ester. After neutralization and precipitation, the crude carboxy modified oligonucleotide was used without further modification.

Preparation of Target DNA Conjugated with Streptavidin (SA_TD002)

[0482] The SA_TD002 was prepared as described in example 3.

Association Reactions (Binding Reaction)

Materials

1 M Tris-HCl, pH 7.5

4 M NaCl

[0483] 10% triton X-100
10 μ M biotin

SA_TD002

[0484] desBio_yR

Protocol

[0485] In a total volume of 0.5 μ L Binding Buffer (10 mM Tris-HCl (pH7.5), 50 mM NaCl, 0.1% triton), 3E8 desBio_yR molecules were mixed with 1.4E9 molecules SA_TD002 in the presence or absence of 1 μ M biotin (inhibitor). Asso-

ciation of the molecules was allowed by incubating the binding mixtures for 1 hour on ice.

Dissociation Reactions (Dilution)

Materials

Standard Ligation Buffer:

50 mM Tris-HCl, pH 7.5

50 mM NaCl

0.1% Triton X-100

0.75 μ M BSA

9 mM KCL

4.5% Glycerol

0.2 mM EDTA

1 mM DTT

2 mM ATP

[0486] 1 μ M T4 DNA ligase (Fermentas)

0.01 μ M 'Scavenger DNA' (40-mer nicked dsDNA fragment):

40-mer dsDNA fragment containing a single nick was prepared by assembly of oligonucleotides vip2554, phosphorylated vip2555 and vip2556.

[0487] Continuous phase was prepared as described in example 3

[0488] 2 mL micro tubes with screw cap

Protocol

[0489] A volume of 0.12 μ L was transferred from the binding mixture to the lid of a 2 mL Eppendorf tube containing 600 μ L aqueous phase containing 1 μ M T4 DNA ligase (standard ligation buffer). The dissociation reaction was initiated by mixing the binding mixture with the aqueous phase by inverting the tubes twice followed by vortexing the tubes for 10 seconds. After a short spin in the microcentrifuge, 500 μ L of the mixture was transferred to an ice-cold 2 mL micro tube containing 1 mL continuous phase and left on ice for the remaining time to finally obtain a dissociation time of 2 minutes.

Emulsification

Materials

Induction Buffer:

50 mM Tris-HCl, pH 7.5

50 mM NaCl

0.1% Triton X-100

1.5 μ M BSA

10 mM KCL

5% Glycerol

0.2 mM EDTA

1 mM DTT

2 mM ATP

135 mM MgCl2

Protocol

[0490] The dissociation reactions were terminated exactly 2 min. after initiation by mixing the continuous phase (1 mL) and the aqueous phase (0.5 mL) by emulsification for 3×20 seconds at 5500 rpm (with 10 seconds pause in between the 20 seconds runs) on the Precellys24 (Bertin Technologies). In parallel, induction-emulsions containing magnesium but no ligase for the activation of T4 DNA ligase were prepared by emulsification for 3×20 seconds at 5500 rpm of 1 mL continuous phase and 0.5 mL aqueous phase containing 135 mM MgCl₂ (Induction Buffer).

Ligation in Emulsion

Protocol

[0491] A volume of 150 µL induction-emulsion containing MgCl₂ was added per emulsion and mixed by rotation for one hour at RT to activate T4 DNA ligase. Ligation of desBio_yR and SA_TD002 in emulsion was allowed by incubating the emulsions (1650 µL) for 16 hours in a thermo block at 16° C. and 300 rpm.

Emulsion Breaking and DNA Recovery

Materials

[0492] 1-butanol

Isopropanol

[0493] 100% ethanol

100 bp no-limits DNA

PCR clean-up kit (NucleoSpin ExtractII, Macherey-Nagel)

10% triton X-100

Protocol

[0494] The ligation reaction was stopped by incubating the tubes for 30 minutes at 65° C. followed by a short spin in the microcentrifuge. For breaking of the emulsions, half of the volume of each emulsion was transferred to a clean 2 mL eppendorf tube. To each tube 850 µL 1-butanol plus 15 ng 100 bp no-limits DNA [10 ng/µL] was added and mixed by thoroughly vortexing for 10 seconds. The tubes were centrifuged for 1 min at 14,000×g and the supernatant was discarded. Residual silicone oil and surfactants were removed from the emulsion by repeating the 1-butanol extraction once more with the addition of 1 volume of 1-butanol. The recovered water-phases of the previously splitted emulsions were pooled into one tube and the DNA fragments (desBio_yR_TD002_SA fusion molecules) were rescued by purification using a PCR clean-up kit (NucleoSpin ExtractII, Macherey-Nagel) according to the supplier's recommendations. The DNA was eluted into EB buffer (5 mM Tris/HCl, pH 8.5) containing 0.1% triton X-100. Prior to qPCR analysis, the eluted DNA was diluted 10 fold in Dilution Buffer (10 mM Tris-HCl pH 7.5, 10 mM NaCl, 0.05% Triton-X100).

Amplification and Analysis of Ligated DNA (Rescue PCR)

Materials

2×PCR Mastermix (pH 8.8 at 25° C.):

40 mM Tris-HCl

20 mM (NH₄)₂SO₄

20 mM KCl

16 mM MgSO₄,

0.2% Triton X-100,

[0495] 0.2 mg/mL BSA

0.4 mM of each nucleotide suitable for hot-start PCR dATP, dTTP, dGTP, and dCTP (CleanAmp, TriLink)

Protocol

[0496] To analyze the number of desBio_yR_TD002_SA fusion molecules present in the different emulsions, qPCR with primers vip660 and vip2824 and using 5 µL of the 10 fold diluted purified recovered DNA samples as a template was run. For the standard curve 3E8, 3E7, 3E6, 3E5 or 3E4 copies of pre-ligated yR_TD002 (control template) were added per qPCR reaction.

qPCR Mixture Per Reaction:

5 µL template

10 µL 2×PCR mastermix

2 µL 5M Betaine (final conc. 0.5 M)

0.4 µL SyBR Green (final conc. 2.5E-5%)

0.2 µL 100 µM vip660 (final conc. 1 µM)

0.2 µL 100 µM vip2824 (final conc 1 µM)

0.2 µL (2 u/µL) Vent (exo-) polymerase (Fermentas)

2 µL water

[0497] The mixture was subjected to thermal cycling by applying the following program in the qPCR machine:

92° C. for 10 min

[0498] 30 cycles of 95° C. 30 seconds and 72° C. for 2 min 30 seconds

72° C. for 5 min

Results

[0499] In order to calculate the number of fusion molecules present in the different qPCR reactions, the standard curve was defined. The calculated numbers of desBio_yR_TD002_SA fusion molecules were translated into a column chart to visualize the difference between the signal obtained in association reactions incubated in presence or absence of 1 µM biotin (FIG. 6). The results show the presence of a significantly higher number of fusion molecules if the association reaction was performed in the absence of biotin compared to if the association reaction was performed in the presence of the inhibitor biotin.

Conclusion

[0500] This result demonstrated co-compartmentalization of desBio_yR and SA_TD002 molecules originating from the desthiobiotin-streptavidin binding and ligation of their attached DNA as a result hereof.

Example 6

Enrichment by Co-Compartmentalization Using
eLigation for Genotype-Genotype Fusion

Spiking Experiment

[0501] ECC was demonstrated by enriching for N-benzyl-4-sulfamoyl-benzamide conjugated to yR DNA (BSB_yR) that was spiked into a diverse yR library using human Carbonic anhydrase II as the target. As a negative control ECC was run in parallel using target preincubated with BSB as the target. Furthermore, dissociation time dependent enrichment was demonstrated in the same system.

[0502] For overview see FIG. 1.

Methods

[0503] DNA oligonucleotides used are described in example 2 and 5. In addition, the following were applied:

Used for yR labeled with BSB (BSB_yR):

vip2260:
ATGAAAGACGTGGCCATTGC

vip2724_vip2607:
CTGACATGGTCCCTGGCAGTCTCCTGTCAGGACCGACTCCXGCTCGAAGA

C
(x = dT-C6-amino modification)

vip2970:
CTATCGGTTTTACCGATAGGTCTTCGAGCTGTACCTGCGC

vip2973:
AGCTAGGTTTTACCTAGCTGCGCAGGTACTGTGCATCGAC

vip2980:
CTATCGGTTTTACCGATAGGTGATGCACTGGAGTCGGTC

Used for TD003:
vip2536:
CTTATGCTGGCAGTTTCA

vip2529:
ACTTCCACCTCAGGACATCGAGCTGGAGCTTGCTGTTAGC

vip2538:
AGGTTGCTCCCTCCTTAAGCCAGCAGTGGTAATTCGACA

vip2996:
CGATGTCCTGAGGTGGAAGTTGAACTGCCAGCATAAGGA

vip2532:
CTTAAGGAGGAGCGAACCTGCTAACAGCAAGCTCCAGCT

vip2559:
x-TGTCGAATTACCACTGCTGG
(x = C6-amino modification)

Used for 454-sequencing:
Vip3018:
CCTATCCCTGTGTGCTTGGCAGTCTCAGCGATGTCCTGAGGTGAAGT

vip2459:
CCATCTCATCCCTGCGTGTCTCCGACTCAGAACCTGGTCCCTGGCAGTCT

CC

vip2460:
CCATCTCATCCCTGCGTGTCTCCGACTCAGAAGGTGGTCCCTGGCAGTCT

CC

-continued

vip2461:
CCATCTCATCCCTGCGTGTCTCCGACTCAGACACTGGTCCCTGGCAGTCT

CC

vip2462:
CCATCTCATCCCTGCGTGTCTCCGACTCAGACTGTGGTCCCTGGCAGTCT

CC

vip2463:
CCATCTCATCCCTGCGTGTCTCCGACTCAGAGAGTGGTCCCTGGCAGTCT

CC

vip2464:
CCATCTCATCCCTGCGTGTCTCCGACTCAGAGCATGGTCCCTGGCAGTCT

CC

vip2465:
CCATCTCATCCCTGCGTGTCTCCGACTCAGAGGTTGGTCCCTGGCAGTCT

CC

vip2466:
CCATCTCATCCCTGCGTGTCTCCGACTCAGAGTCTGGTCCCTGGCAGTCT

CC

vip2467:
CCATCTCATCCCTGCGTGTCTCCGACTCAGATCGTGGTCCCTGGCAGTCT

CC

vip2468:
CCATCTCATCCCTGCGTGTCTCCGACTCAGATGCTGGTCCCTGGCAGTCT

CC

vip2469:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCACTTGGTCCCTGGCAGTCT

CC

vip2470:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCAGATGGTCCCTGGCAGTCT

CC

vip2471:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCCATTGGTCCCTGGCAGTCT

CC

vip2472:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCCTATGGTCCCTGGCAGTCT

CC

vip2473:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCGAATGGTCCCTGGCAGTCT

CC

vip2474:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCTACTGGTCCCTGGCAGTCT

CC

vip2475:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCTCATGGTCCCTGGCAGTCT

CC

-continued

vip2476:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCTGTTGGTCCCTGGCAGTCT

CC

vip2477:
CCATCTCATCCCTGCGTGTCTCCGACTCAGCTTGTGGTCCCTGGCAGTCT

CC

vip2478:
CCATCTCATCCCTGCGTGTCTCCGACTCAGGAATGGTCCCTGGCAGTCT

CC

vip2479:
CCATCTCATCCCTGCGTGTCTCCGACTCAGGACATGGTCCCTGGCAGTCT

CC

vip2480:
CCATCTCATCCCTGCGTGTCTCCGACTCAGGAGTTGGTCCCTGGCAGTCT

CC

vip2481:
CCATCTCATCCCTGCGTGTCTCCGACTCAGGATGTGGTCCCTGGCAGTCT

CC

vip2482:
CCATCTCATCCCTGCGTGTCTCCGACTCAGGCAATGGTCCCTGGCAGTCT

CC

vip2483:
CCATCTCATCCCTGCGTGTCTCCGACTCAGGTCTTGGTCCCTGGCAGTCT

CC

vip2484:
CCATCTCATCCCTGCGTGTCTCCGACTCAGGTGATGGTCCCTGGCAGTCT

CC

vip2485:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTACGTGGTCCCTGGCAGTCT

CC

vip2486:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTAGCTGGTCCCTGGCAGTCT

CC

vip2487:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTCAGTGGTCCCTGGCAGTCT

CC

vip2488:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTCCATGGTCCCTGGCAGTCT

CC

vip2489:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTCGTGGTCCCTGGCAGTCT

CC

vip2490:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTCTCTGGTCCCTGGCAGTCT

CC

-continued

vip2491:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTGACTGGTCCCTGGCAGTCT

CC

vip2492:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTGTGTGGTCCCTGGCAGTCT

CC

vip2493:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTTCTGGTCCCTGGCAGTCT

CC

vip2494:
CCATCTCATCCCTGCGTGTCTCCGACTCAGTTGGTGGTCCCTGGCAGTCT

CC

Preparation of Yoctoreactor Library

[0504] The library was constructed according to (Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327) but in the tetramer format instead of the trimer format.

Preparation of yR Labeled with BSB (BSB_yR)

Materials

[0505] 10 mM Fmoc-NH-PEG(12)-CO₂H
100 mM 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methyl-morpholinium chloride (DMT-MM)
200 mM N-(2-Hydroxyethyl)piperazine-N'-(4-butanesulfonic acid) pH 9 (HEPBS)
N-Methyl-2-pyrrolidone (NMP)
0.5 M piperidine in NMP


10 mM Fmoc-L-Phenylglycine (Fmoc-L-Phe)

[0506] 10 mM 4-carboxybenzenesulfonamide
0.1 M acetonitrile/triethylammonium acetate (pH 7)


BSB: BSB was synthesized according to (Drabovich et al., *Anal. Chem.* 2009, 81, 490-494)

Protocol

[0507] BSB_yR was prepared by ligation. Position 2 oligonucleotide vip2970, position 3 oligonucleotide vip2973 and position 4 oligonucleotide vip2980 were prepared for ligation by phosphorylation with T4 Polynucleotide Kinase performed according to manufactures instructions (Fermentas). BSB labeling of position 1 oligonucleotide vip2724_vip2607 was synthesized according to the following reaction scheme:



-continued



[0508] Amino-PEG12 derived oligonucleotide was synthesized from a 51 mer oligonucleotide (vip2724_vip2607) [5 nmol] with internally modified dT (amino-C6-dT) that was coupled with Fmoc-NH-PEG(12)-CO₂H [10 mM] in a solution of 100 mM DMT-MM, 200 mM HEPBS pH 9, in 200 μ L NMP:water 1:1. After 1 hour the mixture was ethanol precipitated and dissolved in 100 μ L water. A volume of 100 μ L 0.5 M piperidine in NMP was added and incubated at 25° C. for 2 hr. The amino-PEG12-oligonucleotide was isolated by ethanol precipitation and used without further purification in the next coupling.

[0509] BSB-PEG12 labeled position 1 conjugate was synthesized by conjugating Fmoc-L-Phg to a 51 mer amino modified oligonucleotide in which the primary amine was linked through a PEG12 linker on an internally modified dT. The amino-PEG12 derived oligonucleotide was coupled with Fmoc-L-Phg [10 mM] in a solution of 100 mM DMT-MM and 200 mM HEPBS pH 9 in 200 μ L NMP:water 1:1. After 1 hour the mixture was ethanol precipitated and dissolved in 100 μ L water. 100 μ L 0.5 M piperidine in NMP was added and incubated at 25° C. for 2 hr. Final coupling of 4-carboxybenzenesulfonamide was made by treatment of a solution of the L-Phg-PEG12 derived oligonucleotide with 4-carboxybenzenesulfonamide [10 mM] in a solution of 100 mM DMT-MM and 200 mM HEPBS pH 9 in 200 μ L NMP:water 1:1. After incubation at 25° C. for 1 hour the crude oligonucleotide conjugate was isolated by ethanol precipitation and purified by reverse phase HPLC on a C-18 Waters XBridge column with acetonitrile/triethylammonium acetate (pH 7, 0.1 M) gradient 6-50% acetonitrile over 20 min. Appropriate fractions were collected and evaporated in vacuo and resulted into 1680 pmol of vip2724_vip2607 labeled with BSB. Equivalent amounts of the stem complimentary position 1 oligonucleotide vip2724_vip2607 conjugated with BSB, position 2, position 3 and position 4 oligonucleotides were mixed and ligated to form the yR (Hansen et al. *J. Am. Chem. Soc.*, 2009, 131 (3), pp 1322-1327). Ligation with T4 DNA ligase was performed according to manufactures instructions (Fermentas). The BSB_yR was un-folded and made double

stranded via primer extension using phosphorylated vip2260 in a Klenow (exo-) driven reaction performed according to manufactures instructions (Fermentas).

Preparation of Target DNA (TD003)

[0510] TD003 was prepared similar to TD002 as described in example 5.

Conjugation of Carbonic Anhydrase II to TD003 (CAII_TD003)

[0511] Recombinant human CAII (RnD systems; 2184CA)

Protocol

[0512] Conjugation of TD003 to CAII was done similarly as described in example 3. Pre-activation mixture with target DNA for conjugation with CAII. Pre-activation was done by mixing 21 μ L TD003 [4.7 μ M] with 3 μ L MOPS pH 6 [1 M], 3 μ L EDC [50 mM], and 3 μ L s-NHS [100 mM].

[0513] Carboxylic acid activation was allowed to incubate at 20° C. for 30 min. The buffer was removed by using a G25 Illustra column according to manufactures instructions (GE Healthcare).

[0514] Prior to conjugation, the protein was dialyzed 2 $\times$ 30 min against a Dialysis Buffer at 4° C. using Slide-A-Lyzer mini dialysis device according to manufactures instructions (Pierce).

[0515] For the conjugation reaction 1 μ L MOPS [1M] pH 8.0, 1 μ L NaCl [1 M] and 1 μ L water was added to 9 μ L of dialyzed CAII protein. Approx. 35 μ L activated DNA was added to this mixture. The reaction was incubated at 4° C. for 20 h.

[0516] The conjugation reaction was quenched by adding Tris (pH 8) to a final concentration of 50 mM. The CAII_TD003 conjugate was isolated from reactants by PAGE from a 6% TBE gel as described in example 3. The concentration of the conjugate was estimated to be 0.21 μ M by measuring the DNA concentration using Picogreen according to manufactures instructions (Molecular Probes).

Association Reactions (Binding Reaction)

Materials

1 M Tris-HCl, pH 7.5

4 M NaCl

[0517] 10% triton

10 μ M BSB

Protocol

[0518] In a total volume of 1.5 μ L Binding Buffer (10 mM Tris-HCl (pH7.5), 50 mM NaCl, 0.1% triton X-100), 6.5E10 library molecules and 3.3E5 BSB_yR molecules (YoctoReactor library consisting of 1E12 molecules spiked 1 to 200 000 with 5E6 BSB_yR molecules) were mixed with 9E9 molecules CAII_TD003 in the presence or absence of 1 μ M BSB (inhibitor). Association of the molecules was allowed by incubating the binding mixtures for 1 hour on ice.

Dissociation Reactions (Dilution)

Materials

[0519] Standard ligation buffer:

50 mM Tris-HCl, pH 7.5

50 mM NaCl

0.1% Triton X-100

0.75 μ M BSA

9 mM KCL

4.5% Glycerol

0.2 mM EDTA

1 mM DTT

2 mM ATP

[0520] 1 μ M T4 DNA ligase (Fermentas)

0.01 μ M 'Scavenger' DNA

[0521] Continuous phase was prepared as described in example 3

2 mL micro tubes with screw cap

Protocol

[0522] A volume of 0.12 μ L was transferred from the binding mixture to the lid of a 2 mL Eppendorf tube containing 600 μ L aqueous phase containing 1 μ M T4 DNA ligase (standard ligation buffer). The dissociation reaction was initiated by mixing the binding mixture with the aqueous phase by inverting the tubes twice followed by vortexing the tubes thoroughly for 10 seconds. After a short spin on the micro centrifuge, 500 μ L of the mixture was transferred to an ice-cold 2 mL tube containing 1 mL continuous phase and left on ice for the remaining time to finally obtain dissociation times of 2 or 30 minutes.

Emulsification

Materials

[0523] Induction buffer:

50 mM Tris-HCl, pH 7.5

50 mM NaCl

0.1% Triton X-100

1.5 μ M BSA

10 mM KCL

5% Glycerol

0.2 mM EDTA

1 mM DTT

2 mM ATP

135 mM $MgCl_2$

Protocol

[0524] The dissociation reaction was terminated exactly 2 min or 30 min after initiation by mixing the continuous phase

(1 mL) and the aqueous phase (0.5 mL) by emulsification for 3 $\times$ 20 seconds at 5500 rpm (with 10 seconds pause in between the seconds runs) on the Precellys 24 (Bertin Technologies). In parallel, induction-emulsions containing magnesium but no ligase for the activation of T4 DNA ligase were prepared by emulsification for 3 $\times$ 20 seconds at 5500 rpm of 1 mL continuous phase and 0.5 mL aqueous phase containing 135 mM $MgCl_2$ (induction buffer).

Ligation in Emulsion

Protocol

[0525] A volume of 150 μ L induction-emulsion containing $MgCl_2$ was added per emulsion and mixed by rotation for one hour at RT to activate T4 DNA ligase. Ligation was allowed by incubating the emulsions (1650 μ L) for 16 hours in a thermo block at 16° C. and 300 rpm.

Emulsion Breaking and DNA Recovery

Materials

[0526] 1-butanol

Isopropanol

[0527] 100% ethanol

100 bp no-limits DNA

TissueLyser II (Qiagen)

[0528] PCR clean-up kit (NucleoSpin ExtractII, Macherey-Nagel)

10% triton X-100

Protocol

[0529] The ligation reaction was stopped by incubating the tubes for 30 minutes at 65° C. followed by a short spin on the micro centrifuge. For breaking of the emulsions, 300 μ L 1-butanol, 150 μ L isopropanol, 50 μ L ethanol and 20 ng 100 bp no-limits DNA [10 ng/ μ L] was added per emulsion and mixed on the TissueLyser II (Qiagen) for 1 min at 15 Hz. Subsequently the tubes were rotated for 1 hour at RT, centrifuged for 2 min at 14,000 $\times$ g and the supernatant was discarded. Residual silicone oil and surfactants were removed from the emulsion by performing the following extraction twice: addition of 1 volume of 1-butanol, mixing for 1 min at 15 Hz on the TissueLyser, and discarding the upper phase. The DNA fragments were rescued by purification using a PCR clean-up kit (NucleoSpin ExtractII, Macherey-Nagel) according to the supplier's recommendations. The DNA was eluted into EB buffer (5 mM Tris/HCl, pH 8.5) containing 0.1% triton.

Amplification of Ligated DNA (Rescue PCR)

Materials

2 $\times$ PCR Mastermix (pH 8.8 at 25° C.):

40 mM Tris-HCl

20 mM $(NH_4)_2SO_4$

20 mM KCl

16 mM $MgSO_4$,

0.2% Triton X-100,

[0530] 0.2 mg/mL BSA

0.4 mM of each nucleotide suitable for hot-start PCR dATP, dTTP, dGTP, and dCTP (CleanAmp, TriLink)

Protocol

[0531] The ligated fragments were amplified by PCR using 10 μ L of the purified recovered DNA samples as a template in a total volume of 100 μ L.

PCR mixture:

10 μ L template DNA

50 μ L 2 $\times$ PCR Mastermix

[0532] 10 μ L 5 M Betaine (final conc. 0.5 M)

1 μ L 100 μ M vip660 (final conc. 1 μ M)

1 μ L 100 μ M vip2824 (final conc. 1 μ M)

1 μ L (2 u/ μ L) Vent (exo-) polymerase (Fermentas)

27 μ L water

[0533] The mixture was subjected to thermal cycling by applying the following program in a PCR machine:

10 min at 95° C.

[0534] 32 cycles of 30 sec at 95° C. and 2 min 30 sec at 72° C.

2 min at 72° C.

Preparation for 454-Sequencing

Protocol

[0535] Samples for 454-sequencing were prepared as described for the yR_TD001 fusion molecules in example 4. 454-Sequencing tags were included in the PCR for amplification of yR_TD003 fusion molecules using unique forward primers vip2459 to vip2494.

DNA Sequencing

Protocol

[0536] 454-Sequencing was performed as described by (Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327). The DNA sequences were analyzed and the frequency of the BSB genotype calculated.

Results

[0537] The sequencing results (FIG. 7) showed that the BSB_yR was successfully enriched for (about 1300 fold) by CA II using a dissociation time of two minutes. In contrast, no enrichment of BSB_DNA was observed when using a dissociation time of 30 minutes or when CA II was preincubated with BSB prior to the binding step.

Conclusion

[0538] ECC was demonstrated by enriching in dissociation time dependent fashion for BSB_yR that was spiked into a diverse yR library using CAII as the target.

REFERENCE LIST

- [0539]** 1: EP1809743B1 (Vipergen)
[0540] 2: EP1402024B1 (Nuevolution)
[0541] 3: EP1423400B1 (David Liu)
[0542] 4: Nature Chem. Biol. (2009), 5:647-654 (Clark)
[0543] 5: WO 00/23458 (Harbury)
[0544] 6: Nature Methods (2006), 3(7), 561-570 7: 2006 (Miller)
[0545] 7: Nat. Biotechnol. 2004; 22,568-574 (Melkko)
[0546] 8: Nature. (1990); 346(6287), 818-822 (Ellington)
[0547] 9: Proc Natl Acad Sci USA (1997). 94 (23): 12297-302 (Roberts)
[0548] 10: WO06053571A2 (Rasmussen)
[0549] 11: Bertschinger et al, (2007) Protein Engineering, Design & Selection vol. 20 no. 2 pp. 57-68
[0550] 12: Miller O J, Bernath K, Agresti J J, Amitai G, Kelly B T, Mastrobattista E, Taly V, Magdassi S, Tawfik D S, Griffiths A D. Directed evolution by in vitro compartmentalization. Nat Methods. 2006 July; 3(7):561-70
[0551] 13: Doi, N. and Yanagawa, H. (1999) FEBS Lett., 457,227-230
[0552] 14: Yonezawa, M., Doi, N., Kawahashi, Y., Higashinakagawa, T. and Yanagawa, H. (2003) Nucleic Acids Res., 31, e118.
[0553] 15: Tawfik, D. S. and Griffiths, A. D. (1998) Man-made cell-like compartments for molecular evolution. Nat. Biotechnol., 16,652-656
[0554] 16: Ghadessy, F. J., Ong, J. L. and Holliger, P. (2001) Proc. Natl. Acad. Sci. USA, 98, 4552-4557;
[0555] 17: Tay Y, Ho C, Droge P, Ghadessy F J. Selection of bacteriophage lambda integrases with altered recombination specificity by in vitro compartmentalization. Nucleic Acids Res. 2010 March; 38(4):e25. Epub 2009 Dec. 4
[0556] 18: Zheng Y, Roberts R J. Selection of restriction endonucleases using artificial cells. Nucleic Acids Res. 2007; 35(11):e83. Epub 2007
[0557] 19: Mastrobattista E, Taly V, Chanudet E, Treacy P, Kelly B T, Griffiths A D. High-throughput screening of enzyme libraries: in vitro evolution of a beta-galactosidase by fluorescence-activated sorting of double emulsions. Chem Biol. 2005 December; 12(12):1291-300
[0558] 20: Levy M, Griswold K E, Ellington A D. Direct selection of trans-acting ligase ribozymes by in vitro compartmentalization. RNA. 2005 October; 11(10):1555-62. Epub 2005 Aug. 30;
[0559] 21: Sepp A, Choo Y. Cell-free selection of zinc finger DNA-binding proteins using in vitro compartmentalization. J Mol Biol. 2005 Nov. 25; 354(2):212-9. Epub 2005 Oct. 3;
[0560] 22: Bernath K, Magdassi S, Tawfik D S. Directed evolution of protein inhibitors of DNA-nucleases by in vitro compartmentalization (IVC) and nano-droplet delivery. J Mol Biol. 2005 Feb. 4; 345(5):1015-26. Epub 2004 Dec. 7.
[0561] 23: Bertschinger et al, (2004) Protein Engineering, Design & Selection vol. 17 no. 9 pp. 699-707
[0562] 24: Chen Yu et al, (November 2008) Nucleic Acid Research, Vol. 36, Nr. 19, Pages: Article No. E128
[0563] 25: Hansen et al. J. Am. Chem. Soc., 2009, 131 (3), pp 1322-1327
[0564] 26: Drabovich et al., Anal. Chem. 2009, 81, 490-494
[0565] 27: Turner and Hurles, Nat Protoc. 2009; 4(12): 1771-1783

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 78

<210> SEQ ID NO 1
<211> LENGTH: 191
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..191
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

<400> SEQUENCE: 1

```
cgctaagtgt ccttgccagt ctccttagcg gaccgactcc tgctcgaaga caacgggtgtt    60
ttacaccggt gtcttcgagc tgtacctgcg caagtgcgtt ttacgcactt gcgcaggtac    120
tgtgcatcga caagaccgtt ttacggtctt gtcgatgcac tggagtcggt cctgttcgat    180
cttgggcgta t                                                                191
```

<210> SEQ ID NO 2
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

<400> SEQUENCE: 2

```
atacgcccaa gatcgaacag                                                                20
```

<210> SEQ ID NO 3
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..18
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

<400> SEQUENCE: 3

```
tggtccctgg cagtctcc                                                                18
```

<210> SEQ ID NO 4
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..60
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

<400> SEQUENCE: 4

```
ctgttcgatc ttgggcgtat gagaagagcc agaaacgtgg cttcaggcac caaggaagac    60
```

<210> SEQ ID NO 5
<211> LENGTH: 59

-continued

```
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..59
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="Artificial Sequence"

<400> SEQUENCE: 5

gccttgccag cccgctcagg caagtcttac agccgatcag tcttccttgg tgctgaag      59

<210> SEQ ID NO 6
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="Artificial Sequence"

<400> SEQUENCE: 6

ctgttcgatc ttgggcgtat      20

<210> SEQ ID NO 7
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..19
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="Artificial Sequence"

<400> SEQUENCE: 7

gccttgccag cccgctcag      19

<210> SEQ ID NO 8
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..19
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="Artificial Sequence"

<400> SEQUENCE: 8

gccttgccag cccgctcag      19

<210> SEQ ID NO 9
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..16
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="Artificial Sequence"

<400> SEQUENCE: 9

tggtccctgg cagtct      16
```

-continued

<210> SEQ ID NO 10
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..12
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

<400> SEQUENCE: 10

gaacaggacc ga 12

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

<400> SEQUENCE: 11

ctgttcgatac ttgggcgtat 20

<210> SEQ ID NO 12
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..18
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

<400> SEQUENCE: 12

acgcccaaga tcgaacag 18

<210> SEQ ID NO 13
<211> LENGTH: 61
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..61
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

<400> SEQUENCE: 13

gccttgccag cccgctcagg ggaaggacgt tgggtgtagaa gcgttcactt ggtggaagta 60
t 61

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="Artificial Sequence"

-continued

<400> SEQUENCE: 14

acttcaccca agtgaacgct

20

<210> SEQ ID NO 15

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: artificial sequences

<220> FEATURE:

<221> NAME/KEY: source

<222> LOCATION: 1..60

<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 15

ctgttcgatc ttgggcgat tggttttagct gcccacactc cttcaggcac caaggaagac

60

<210> SEQ ID NO 16

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: artificial sequences

<220> FEATURE:

<221> NAME/KEY: source

<222> LOCATION: 1..20

<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 16

gcaagtctta cagccgatca

20

<210> SEQ ID NO 17

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: artificial sequences

<220> FEATURE:

<221> NAME/KEY: source

<222> LOCATION: 1..18

<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 17

tggtccctgg cagtctcc

18

<210> SEQ ID NO 18

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: artificial sequences

<220> FEATURE:

<221> NAME/KEY: source

<222> LOCATION: 1..50

<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 18

cctatccct gtgtgccttg gcagtctcag gtcttccttg gtgcctgaag

50

<210> SEQ ID NO 19

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: artificial sequences

<220> FEATURE:

<221> NAME/KEY: source

<222> LOCATION: 1..52

-continued

<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 19

ccatctcatc cctgcgtgtc tccgactcag aggttggtcc ctggcagtct cc 52

<210> SEQ ID NO 20
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="PRIMER"
/organism="artificial sequences"

<400> SEQUENCE: 20

ccatctcatc cctgcgtgtc tccgactcag atcgttggtcc ctggcagtct cc 52

<210> SEQ ID NO 21
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 21

ccatctcatc cctgcgtgtc tccgactcag atcgttggtcc ctggcagtct cc 52

<210> SEQ ID NO 22
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 22

ccatctcatc cctgcgtgtc tccgactcag cacttggtcc ctggcagtct cc 52

<210> SEQ ID NO 23
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 23

ccatctcatc cctgcgtgtc tccgactcag cagatgggtcc ctggcagtct cc 52

<210> SEQ ID NO 24
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences

-continued

```
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 24

ccatctcatc cctgcgtgtc tccgaactcag ccattggtcc ctggcagtct cc          52

<210> SEQ ID NO 25
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..18
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 25

tggtccctgg cagtctcc          18

<210> SEQ ID NO 26
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..36
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 26

caccacgatg gcaatgcatt cttcgctgcc attctg          36

<210> SEQ ID NO 27
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 27

cgatgtcctg aggtggaagt          20

<210> SEQ ID NO 28
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 28

ggcaagtgat tgtccatgtg catgagaaga ggccccacatt          40

<210> SEQ ID NO 29
```

-continued

```
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
    /note="Primer"
    /organism="artificial sequences"

<400> SEQUENCE: 29

cacatggaca atcacttgcc                                20

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
    /note="Primer"
    /organism="artificial sequences"

<400> SEQUENCE: 30

aatgtgggcc tcttctcatg                                20

<210> SEQ ID NO 31
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..18
<223> OTHER INFORMATION: /mol_type="DNA"
    /note="Primer"
    /organism="artificial sequences"

<400> SEQUENCE: 31

tccacatcct ccagttca                                  18

<210> SEQ ID NO 32
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
    /note="Primer"
    /organism="artificial sequences"

<400> SEQUENCE: 32

acttccacct caggacatcg agctggagct tgctgtagc          40

<210> SEQ ID NO 33
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
    /note="Primer"
    /organism="artificial sequences"

<400> SEQUENCE: 33

aggttcgctc cctccttaag tcaggaggat gtgacaccaa          40
```

-continued

<210> SEQ ID NO 34
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 34

cgatgtcctg aggtggaagt tgaactggag gatgtggaca 40

<210> SEQ ID NO 35
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 35

cttaaggagg gagcgaacct gctaacagca agtccagct 40

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 36

ttggtgtcac atcctcctga 20

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 37

atgaaagacg tggccattgc 20

<210> SEQ ID NO 38
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..50
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"
<220> FEATURE:

-continued

<221> NAME/KEY: misc_feature
<222> LOCATION: 41

<400> SEQUENCE: 38

ctgacatggt ccctggcagt ctctgtcag gaccgactcc gctcgaagac 50

<210> SEQ ID NO 39
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 39

ctatcggttt taccgatagg tcttcgagct gtacctgcgc 40

<210> SEQ ID NO 40
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 40

agctaggttt tacctagctg cgcaggtact gtgcatcgac 40

<210> SEQ ID NO 41
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 41

ctatcggttt taccgatagg tcgatgcact ggagtcggtc 40

<210> SEQ ID NO 42
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..18
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 42

cttatgctgg cagtttca 18

<210> SEQ ID NO 43
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:

-continued

```
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 43

acttccacct caggacatcg agctggagct tgctgtagc

40

<210> SEQ ID NO 44
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 44

aggttcgctc cctccttaag ccagcagtg taattcgaca

40

<210> SEQ ID NO 45
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 45

cgatgtcctg aggtggaagt tgaaactgcc agcataagga

40

<210> SEQ ID NO 46
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..40
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 46

cttaaggagg gagcgaacct gctaacagca agtccagct

40

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..20
<223> OTHER INFORMATION: /mol_type="DNA"
      /note="Primer"
      /organism="artificial sequences"

<400> SEQUENCE: 47

tgtcgaatta ccactgctgg

20

<210> SEQ ID NO 48
<211> LENGTH: 50
```

-continued

<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..50
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 48

cctatccct gtgtgccttg gcagtcctcag cgatgtcctg aggtggaagt 50

<210> SEQ ID NO 49
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 49

ccatctcatc cctgcgtgtc tccgactcag aacctgggcc ctggcagtct cc 52

<210> SEQ ID NO 50
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 50

ccatctcatc cctgcgtgtc tccgactcag aagtggtgcc ctggcagtct cc 52

<210> SEQ ID NO 51
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 51

ccatctcatc cctgcgtgtc tccgactcag aactgggtcc ctggcagtct cc 52

<210> SEQ ID NO 52
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 52

ccatctcatc cctgcgtgtc tccgactcag actgtgggcc ctggcagtct cc 52

-continued

<210> SEQ ID NO 53
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 53

ccatctcatc cctgcgtgtc tccgactcag agagtgtgtcc ctggcagtct cc 52

<210> SEQ ID NO 54
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 54

ccatctcatc cctgcgtgtc tccgactcag agcatgtgtcc ctggcagtct cc 52

<210> SEQ ID NO 55
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 55

ccatctcatc cctgcgtgtc tccgactcag agtctgtgtcc ctggcagtct cc 52

<210> SEQ ID NO 56
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 56

ccatctcatc cctgcgtgtc tccgactcag cctatgtgtcc ctggcagtct cc 52

<210> SEQ ID NO 57
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 57

-continued

ccatctcatc cctgcgtgtc tccgactcag cgaatgggcc ctggcagtct cc 52

<210> SEQ ID NO 58
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 58

ccatctcatc cctgcgtgtc tccgactcag ctactgggcc ctggcagtct cc 52

<210> SEQ ID NO 59
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 59

ccatctcatc cctgcgtgtc tccgactcag ctcatgggcc ctggcagtct cc 52

<210> SEQ ID NO 60
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 60

ccatctcatc cctgcgtgtc tccgactcag ctgttgggcc ctggcagtct cc 52

<210> SEQ ID NO 61
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 61

ccatctcatc cctgcgtgtc tccgactcag cttgtgggcc ctggcagtct cc 52

<210> SEQ ID NO 62
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"

-continued

```
/organism="artificial sequences"

<400> SEQUENCE: 62

ccatctcatc cctgcgtgtc tccgactcag gaactggtec ctggcagtct cc          52

<210> SEQ ID NO 63
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
                        /note="Primer"
                        /organism="artificial sequences"

<400> SEQUENCE: 63

ccatctcatc cctgcgtgtc tccgactcag gacatggtec ctggcagtct cc          52

<210> SEQ ID NO 64
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
                        /note="Primer"
                        /organism="artificial sequences"

<400> SEQUENCE: 64

ccatctcatc cctgcgtgtc tccgactcag gagttggtec ctggcagtct cc          52

<210> SEQ ID NO 65
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
                        /note="Primer"
                        /organism="artificial sequences"

<400> SEQUENCE: 65

ccatctcatc cctgcgtgtc tccgactcag gatgtggtec ctggcagtct cc          52

<210> SEQ ID NO 66
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
                        /note="Primer"
                        /organism="artificial sequences"

<400> SEQUENCE: 66

ccatctcatc cctgcgtgtc tccgactcag gcaatggtec ctggcagtct cc          52

<210> SEQ ID NO 67
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
```

-continued

<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 67

ccatctcatc cctgcgtgtc tccgactcag gtcttggtcc ctggcagtct cc 52

<210> SEQ ID NO 68
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Pprimer"
/organism="artificial sequences"

<400> SEQUENCE: 68

ccatctcatc cctgcgtgtc tccgactcag gtgatgggtcc ctggcagtct cc 52

<210> SEQ ID NO 69
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 69

ccatctcatc cctgcgtgtc tccgactcag tacgtgggtcc ctggcagtct cc 52

<210> SEQ ID NO 70
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 70

ccatctcatc cctgcgtgtc tccgactcag tagctgggtcc ctggcagtct cc 52

<210> SEQ ID NO 71
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 71

ccatctcatc cctgcgtgtc tccgactcag tcagtgggtcc ctggcagtct cc 52

<210> SEQ ID NO 72
<211> LENGTH: 52
<212> TYPE: DNA

-continued

<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 72

ccatctcatc cctgcgtgtc tccgactcag tccatgggcc ctggcagtct cc 52

<210> SEQ ID NO 73
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 73

ccatctcatc cctgcgtgtc tccgactcag tcgttggtcc ctggcagtct cc 52

<210> SEQ ID NO 74
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 74

ccatctcatc cctgcgtgtc tccgactcag tctctgggcc ctggcagtct cc 52

<210> SEQ ID NO 75
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 75

ccatctcatc cctgcgtgtc tccgactcag tgactgggcc ctggcagtct cc 52

<210> SEQ ID NO 76
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
/note="Primer"
/organism="artificial sequences"

<400> SEQUENCE: 76

ccatctcatc cctgcgtgtc tccgactcag tgttggtgcc ctggcagtct cc 52

-continued

```

<210> SEQ ID NO 77
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
                        /note="Primer"
                        /organism="artificial sequences"

<400> SEQUENCE: 77

ccatctcatc cctgcgtgtc tccgactcag ttctgtgtcc ctggcagtct cc           52

<210> SEQ ID NO 78
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: artificial sequences
<220> FEATURE:
<221> NAME/KEY: source
<222> LOCATION: 1..52
<223> OTHER INFORMATION: /mol_type="DNA"
                        /note="Primer"
                        /organism="artificial sequences"

<400> SEQUENCE: 78

ccatctcatc cctgcgtgtc tccgactcag ttggtgtgtcc ctggcagtct cc           52

```

1. A method for making an enriched library comprising specific nucleic acid sequence information allowing to identifying at least one binding entity that binds to at least one target wherein the specific binding entity has been present in an in vitro display library and wherein the method comprises the steps of:

- (i): making an in vitro display library of at least 100 different binding entities, wherein each binding entity is attached to a nucleic acid molecule and the nucleic acid molecule comprises specific nucleic acid sequence information allowing to identify the binding entity once one knows the specific nucleic acid sequence information of the nucleic acid molecule one directly knows the structure of the specific binding entity attached to the nucleic acid molecule—the structure of the binding entity, attached to the nucleic acid molecule, is herein termed B-structure;
- (ii): making nucleic acid molecules with at least one target T_n attached to a nucleic acid molecule and the nucleic acid molecule comprises specific nucleic acid sequence information allowing to identify the specific target, wherein the target is capable of binding to at least one of the binding entities present in the library of step (i)—the structure of the target attached to the nucleic acid molecule is herein termed T-structure;

and wherein the method is characterized by that:

- (iii): mixing a solution comprising X, wherein X is a number greater than 10^4 , numbers of B-structures of the library of step (i) with a solution comprising Y, wherein Y is a number greater than 10^2 , numbers of T-structures of step (ii) under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do and wherein one gets binding of at least

one of the binding entities to at least one target thereby creating a complex comprising a B-structure bound to a T-structure, herein termed $B_{BoundTo}$ T-structure;

- (iv): applying an in vitro compartmentalization system—under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do—wherein the compartmentalization system comprises at least 2 times more individual compartments than the Y number of T-structures present in step (iii) under conditions wherein the B-structures, T-structures and $B_{BoundTo}$ T-structures enter randomly into the individual compartments; and
- (v): fusing the nucleic acid molecules of a B-structure and a T-structure which are both present within the same individual compartment—i.e. fusing the nucleic acid molecule of the B-structure to the nucleic acid molecule of the T-structure—this structure is herein termed BT_{Fused} -structure and the BT_{Fused} -structure comprises the specific nucleic acid sequence information allowing to identify the binding entity of step (i) and the specific nucleic acid sequence information allowing to identify the specific target of step (ii); and
- (vi): combining the content of the individual compartments of step (v) under conditions wherein there is no fusing of the nucleic acid molecules of a B-structure and a T-structure—i.e. there is not created any new BT_{Fused} -structure not already created in step (v)—in order to get a library of BT_{Fused} -structures, wherein the library is an enriched library of species of BT_{Fused} -structures originating from binding pairs of target and binder entity when compared to BT_{Fused} -structures originating from nonbinding pairs of target and binder entity; and

wherein the method is further characterized by at least one of (a) or (b) below:

(a) $B_{BoundTo}$ T-structures remain suspended in solution in the individual compartments of step (iv) of the first aspect;

(b) the method does not rely on target immobilization on a solid support.

2. The method of claim 1, wherein the binding entity of step (i) is attached to the nucleic acid molecule by a covalent binding and wherein the target of step (ii) is attached to the nucleic acid molecule by a covalent binding and wherein the nucleic acid molecule of the B-structure is DNA and the nucleic acid molecule of the T-structure is DNA.

3. The method of claim 2, wherein the DNA nucleic acid molecule in the B-structure is a double stranded nucleic acid molecule and wherein the DNA nucleic acid molecule in the T-structure is a double stranded nucleic acid molecule.

4. The method of claim 1, wherein the nucleic acid molecule attached to the binding entity in the B-structure contains a PCR priming site and wherein the nucleic acid molecule attached to the target in the T-structure contains a PCR priming site.

5. The method of claim 1, wherein the in vitro library of step (i) comprises at least 10^5 different binding entities and wherein the binding entities of step (i) are chemical compounds with an average molecular weight MW below 5000 dalton.

6. The method of claim 1, wherein there is at least two different targets in step (ii).

7. The method of claim 1, wherein at least one target is a protein.

8. The method of claim 1, wherein there in step (iii) is at least 10^5 copies of a T-structure of interest, "Y" is at least 10^5 and wherein the concentration of T-structures in the "mixing step (iii)" is at least 10^{-9} M.

9. The method of claim 1, wherein step (iii) is performed under binding conditions, wherein a B-structure containing a binding entity capable of binding to a target molecule, binds 100 fold more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do.

10. The method of claim 1, wherein said method comprises an additional step (iii-b) that is performed before the step (iv), comprising:

(iii-b): diluting the solution of step (iii) at least 100 fold under binding conditions, i.e. conditions where a B-structure containing a binding entity capable of binding to a target molecule, binds more efficiently to the corresponding T-structure, than a B-structure containing a binding entity not capable of binding to the same target do.

11. The method of claim 1, wherein there in step (iv) is at least 100 times more individual compartments than the Y number of T-structures present in step (iii) and wherein there in step (iv) there is at least the square root of 10, 3.16, times more individual compartments than the X number of B-structures in step (iii).

12. The method of claim 1, wherein the in vitro compartmentalization system of step (iv) is a water-in-oil emulsion system and wherein the average compartments volume is less than 10^{-12} liter.

13. The method of claim 1, wherein the fusing of the nucleic acid molecules of a B-structure and a T-structure which are both present within the same individual compartment of step (v) is:

(a): done by a DNA ligase where a phosphodiester bond between a 3'-OH and a 5'-phosphate groups is formed; or
(b): by using a DNA polymerase using emulsion PCR.

14. The method of claim 1, wherein the in vitro compartmentalization system of step (iv) is a water-in-oil emulsion and wherein the content of the individual compartments of step (v) is combined in step (vi) by a method, wherein the oil compartments are disrupted.

15. The method of claim 1, wherein there is an extra step (vii), wherein the BT_{Fused} -structures present in the enriched library of step (vi) is amplified by PCR and thereafter subjected to DNA sequencing to identify at least one individual binding entity that binds to at least one target of interest.

16. The method of claim 1, wherein there is an extra step (vii) comprising use the enriched library of step (vi) to identify at least one individual binding entity that binds to at least one target of interest.

17-18. (canceled)

* * * * *