



(86) Date de dépôt PCT/PCT Filing Date: 1998/09/25
(87) Date publication PCT/PCT Publication Date: 1999/04/01
(45) Date de délivrance/Issue Date: 2005/05/17
(85) Entrée phase nationale/National Entry: 2000/03/24
(86) N° demande PCT/PCT Application No.: US 1998/020208
(87) N° publication PCT/PCT Publication No.: 1999/015703
(30) Priorité/Priority: 1997/09/26 (60/060,170) US


(51) Cl.Int.⁶/Int.Cl.⁶ C12Q 1/68, A61K 48/00, C12N 15/85,
C12N 15/63, C12P 19/34, C12N 15/11, C07H 21/04,
C12N 15/00

(72) Inventeurs/Inventors:
WELLSTEIN, ANTON, US;
CZUBAYKO, FRANK, US

(73) Propriétaire/Owner:
GEORGETOWN UNIVERSITY, US

(74) Agent: SMART & BIGGAR

(54) Titre : INHIBITION DE LA PROTEINE DE FIXATION DU FGF AU MOYEN DE RIBOZYMES
(54) Title: INHIBITION OF AN FGF-BINDING PROTEIN USING RIBOZYMES



(57) Abrégé/Abstract:

Increased expression of a secreted FGF-binding protein (FGF-BP) occurs in certain autoimmune and malignant disease conditions. It is found, for example, that tumor secretions of FGF-BP results in mobilization and activation of locally-stored FGFs that can serve as an angiogenic switch molecule. Furthermore, it has been found that in an animal model of multiple sclerosis (MS), the exacerbation of the disease is accompanied by increased FGF-BP. Using ribozymes, it is possible to cause cleavage of the FGF-BP mRNA. Hence, administration of ribozymes which cleave the FGF-BP mRNA in sufficient amounts to inhibit disease processes triggered by FGF-BP is appropriate.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, C12P 19/34, C12N 15/00, 15/11, 15/63, 15/85, C07H 21/04, A61K 48/00	A1	(11) International Publication Number: WO 99/15703 (43) International Publication Date: 1 April 1999 (01.04.99)
(21) International Application Number: PCT/US98/20208 (22) International Filing Date: 25 September 1998 (25.09.98) (30) Priority Data: 60/060,170 26 September 1997 (26.09.97) US (71) Applicant: GEORGETOWN UNIVERSITY [US/US]; Office of Technology Transfer, 177 Building D, 4000 Reservoir Road, N.W., Washington, DC 20007 (US). (72) Inventors: WELLSTEIN, Anton; 4406 Volta Place, Washing- ton, DC 20007 (US). CZUBAYKO, Frank; 5009 Sentinel Drive #57, Bethesda, MD 20816 (US). (74) Agent: HENDRICKS, Glenna; P.O. Box 2509, Fairfax, VA 22031-2509 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: INHIBITION OF AN FGF-BINDING PROTEIN USING RIBOZYMES (57) Abstract Increased expression of a secreted FGF-binding protein (FGF-BP) occurs in certain autoimmune and malignant disease conditions. It is found, for example, that tumor secretions of FGF-BP results in mobilization and activation of locally-stored FGFs that can serve as an angiogenic switch molecule. Furthermore, it has been found that in an animal model of multiple sclerosis (MS), the exacerbation of the disease is accompanied by increased FGF-BP. Using ribozymes, it is possible to cause cleavage of the FGF-BP mRNA. Hence, administration of ribozymes which cleave the FGF-BP mRNA in sufficient amounts to inhibit disease processes triggered by FGF-BP is appropriate.		

WO 99/15703

PCT/US98/20208

Title: INHIBITION OF AN FGF-BINDING PROTEIN USING RIBOZYMES

Field of the Invention:

5 This invention relates to use of ribozymes for inhibition of fibroblast growth factor binding protein (FGF-BP). The increase of FGF-BP is accompanied by various pathological changes, including the disease phenomena seen in autoimmune disease, e.g., multiple sclerosis and in growth and metastatic spread of cancer.

Background of the Invention:

10 Fibroblast growth factor (FGF) is a potentially active growth factor found in large amounts in the nervous system and, to a somewhat lesser extent, in the rest of the body. Because of the blood-brain barrier, the two reservoirs, including the components such as the FGF and FGF-binding protein, remain separated. However, in certain instances of infection or
15 trauma there is cross-over between the two reservoirs.

Tumor growth and, ultimately, metastasis is a complex process regulated in part by factors controlling cellular proliferation and death as well as tumor angiogenesis. The driving factors which regulate angiogenesis and tumor growth
20 need to be understood to control the process of growth in malignancies. It is known that tumor cells and their normal stroma express a multitude of candidate angiogenic factors. Very few specific inhibitors have been generated to assess which of these gene products are only innocent bystanders and
25 which contribute significantly to tumor angiogenesis and metastasis.

Developmental expression of the retinoid-regulated FGF-BP gene is prominent in the skin and intestine during the perinatal phase and is down-modulated in the adult. The gene
30 is, however, up-regulated in various cancers such as carcinogen-induced skin tumor, in squamous cell carcinoma, in breast cancers and in some colon cancer cell lines and tumor samples. FGF-BP is also up-regulated in autoimmune responses of the

61181-89

2

nervous system such as in the case of multiple sclerosis.

Tumor-angiogenesis, a process whereby factors stimulating the ingrowth of blood vessels into the tumor are secreted into the local tumor milieu by cancer and stroma cells, also plays a critical role by regulating the balance between cell proliferation and cell death and by providing a route for distant spread. Both clinical and laboratory evidence suggest the spread of malignant cells from a localized tumor is directly related to the number of microvessels in the primary tumor. Of the multitude of factors secreted by tumor and stroma cells which are potentially angiogenic, two have been confirmed as angiogenic factors which are rate-limiting in in vivo tumor models. The importance of one of these, vascular endothelial growth factor/vascular permeability factors (VEGF/VPF), was previously demonstrated through functional knockout through use of blocking antibodies. A critical role for the other factor, pleiotrophin (PTN), has been shown in angiogenesis and metastasis associated with melanoma using a hammerhead-ribozyme PTN mRNA depletion strategy.

Summary of the Invention:

It has now been discovered that increased expression of a secreted FGF-binding protein (FGF-BP) occurs in certain autoimmune and malignant disease conditions. It is found, for example, that tumor secretions of FGF-BP results in mobilization and activation of locally-stored FGFs that can serve as an angiogenic switch molecule. Furthermore, it has been found that in an animal model of multiple sclerosis (MS), the exacerbation of the disease is accompanied by increased FGF-BP. Using ribozymes, it is possible to cause cleavage of the FGF-BP mRNA. Hence, administration of ribozymes which cleave the FGF-BP mRNA in

61181-89

2a

sufficient amounts to inhibit disease processes triggered by FGF-BP is appropriate.

Thus in one aspect the present invention provides a method of inhibiting expression of fibroblast growth factor binding protein (FGF-BP) in a cell in vitro by administration of FGF-BP ribozymes which cleave FGF-BP mRNA, wherein said ribozymes are hammerhead ribozymes as depicted in (FIG. 1) having a target cleavage site selected from the group consisting of 258, 332 and 697 nucleotides downstream of the translation initiation site of FGF-BP mRNA, and wherein said ribozymes comprise antisense sequences 5' and 3' to the ribozyme catalytic center that direct the ribozymes to said target cleavage site, such that expression of FGF-BP is inhibited.

In another aspect, the present invention provides a composition comprising: (a) ribozymes which specifically cleave FGF-BP mRNA, wherein said ribozymes are hammerhead ribozymes as depicted in (FIG. 1) having a target cleavage site selected from the group consisting of 258, 332 and 697 nucleotides downstream of the translation initiation site of FGF-BP mRNA, and wherein said ribozymes comprise antisense sequences 5' and 3' to the ribozyme catalytic center that direct the ribozymes to said target cleavage site; and (b) a pharmaceutically acceptable carrier.

In another aspect, the present invention provides use in the preparation of a medicament for reducing tumor growth in a subject by reducing the expression of FGF-BP mRNA in said tumor of the composition of the invention.

In another aspect, the present invention provides Use for reducing tumor growth in a subject by reducing the expression of FGF-BP mRNA in said tumor of the composition of the invention.

61181-89

2b

In another aspect, the present invention provides a commercial package comprising the composition of the present invention together with instructions for use for reducing tumor growth in a subject.

5 Brief Description of the Figures:

Figure 1 shows the ribozyme target sites, expression vector and in vitro cleavage of FGF-BP mRNA.

61181-89

3

Description of the Invention:

It has now been found that expression of FGF-binding protein (FGF-BP) causes mobilization and activation of locally stored FGFs that can serve as switch molecules. The FGF-BP expression can be controlled by targeting with ribozymes, thereby reducing the FGF-BP release of biologically active bFGF from cells. Furthermore, the growth and angiogenesis of tumors can be decreased in parallel with the reduction of FGF-BP. Additionally, it now is seen that autoimmune responses can be modulated by reducing FGF-BP release using the methods of the invention.

Materials and Methods:Generation of constructs:

The FGF-BP expression vector containing the open reading frame (ORF) of FGF-BP from -42 to +707 (with A in the translation initiation codon being +1, Genbank #M60047) was previously described (Czubayko, et al., J. Biol. Chem. 269: pp 28243-28248 (1994)). The FGF-BP-targeted ribozymes were designed to cleave 258, 332, 697 or 753 downstream of the translation initiation site (Fig. 1). The PCR product was purified using the ion exchange columns and reagents purchased from Qiagen* (Chatsworth, CA). The product of the PCR reaction was digested in HindIII and XbaI and cloned into pRc/CMV expression vector purchased from Invitrogen. All of the ribozymes exemplified have 8 nucleotides of 5' and 3' flanking antisense sequence stretches to direct them to the predicted cleavage site. Synthetic sense and antisense oligonucleotides containing the catalytic center and the flanking regions of each of the ribozymes were annealed and ligated as described earlier. (See Czubayko, et al., J. Biol. Chem. 269, 21358 to 21363 (1994).)

The mRNA sequence giving rise to FGF-BP as recorded in the gene bank is of the formula:

*Trade-mark

WO 99/15703

PCT/US98/20208

4

	1	ctctacctga	cacagctgca	gcctgcaatt	cactcccact	gcctgggatt
		gcactggatc				
<u>5</u>	61	cgtgtgctca	gaacaagggtg	aacgcccagc	tgcagccatg	aagatctgta
		gcctcacct			↑	
	121	gctctccttc	ctcctactgg	ctgctcaggt	gctcctgggtg	gaggggaaaa
		aaaaagtga				
<u>10</u>	181	gaatggactt	cacagcaaag	tgggtctcaga	acaaaaggac	actctgggca
		acaccagat				
	241	taagcagaaa	agcaggcccg	ggaacaaagg	caagtttgtc	accaaagacc
<u>15</u>		aagccaactg				
	301	cagatgggct	gctactgagc	aggaggaggg	catctctctc	aaggttgagt
		gcactcaatt				
<u>20</u>	361	ggaccatgaa	ttttcctgtg	tctttgctgg	caatccaacc	tcatgcctaa
		agctcaagga				
	421	tgagagagtc	tattggaac	aagttgcccg	gaatctgcgc	tcacagaaag
		acatctgtag				
<u>25</u>	481	atattccaag	acagctgtga	aaaccagagt	gtgcagaaag	gattttccag
		aatccagtct				
	541	taagctagtc	agctccactc	tatttgaggaa	cacaaagccc	aggaaggaga
<u>30</u>		aaacagagat				
	601	gtccccccagg	gagcacatca	agggcaaaga	gaccaccccc	tctagcctag
		cagtgaacca				
<u>35</u>	661	gaccatggcc	accaaagctc	ccgagtgtgt	ggaggacca	gatatggcaa
		accagaggaa				
	721	gactgccctg	gagttctgtg	gagagacttg	gagctctctc	tgcacattct
		tcctcagcat				
<u>40</u>	781	agtgcaggac	acgtcatgct	aatgaggtca	aaagagaacg	ggttccttta
		agagatgtca				
	841	tgtcgtaagt	ccctctgtat	actttaaaagc	tctctacagt	cccccaaaa
<u>45</u>		tatgaacttt				
	901	tgtgcttagt	gagtgcacg	aaatatttaa	acaagttttg	tattttttgc
		ttttgtgttt				
<u>50</u>	961	tggaatttgc	cttatttttc	ttggatgcga	tgttcagagg	ctgtttcctg
		cagcatgtat				
	1021	ttccatggcc	cacacagcta	tgtgtttgag	cagcgaagag	tctttgagct
		gaatgagcca				
<u>55</u>	1081	gagtgataat	ttcagtgcaa	cgaactttct	gctgaattaa	tggttaataaa
		actctgggtg				

61181-89

5

1141 tttttcaaaa aaaaaaaaaaaa aaa

wherein † identifies the beginning of the coding sequence. (The identification of the sequence chosen as a target sequence relates to the number attributed to the position with relation to the beginning of the coding sequence, and not with relation to the number appearing on the right hand side of the listings.)

Cell lines, transfections and growth assays:

SW-13, LS174T and ME-180 cells (obtained from American Type Culture Collection (ATCC), Rockville, Maryland) were cultured in IMEM with 10% fetal calf serum. Cells were transfected at 50% confluency for 5 hours with 20 µg of plasmid DNA mixed with 140 µl of LIPOFECTAMINE™ in serum-free OPTIMEM™ medium (Life Technologies) at 37°C with 5% CO₂. The transfection medium was then replaced with fresh medium and, 36 hours later, G418 at 50 µg/ml (SW-13, LS174T) or at 1000 µg/ml (ME-180) was added to select stably transfected cells for another 6-8 weeks. Studies of anchorage-independent growth of transfected SW-13 cells were carried out as described previously by Fang, et al. (J. Biol. Chem 267, 25889-25897 (1992)).

Detection of FGF-BP MRNA by Northern blot:

Total RNAs from cell lines or tumor tissues were isolated, blotted and quantitated using a random primed FGF-BP cDNA probe. FGF-BP transcript was quantitated after phosphorimaging and probing for loading with GAPDH. (See Liaudet-Coopman, et al., J. Biol. Chem. 271, 21303-21308 (1996).)

Staining for basic FGF:

Cells were plated overnight, washed, three times with PBS, fixed in 3.7% formaldehyde/0.1% Triton* X-10 for 10 minutes, washed again in PBS and then incubated for 20 minutes at room temperature with a 1:2000 dilution of rabbit anti-bFGF antibody in PBS. After a further triplicate wash in PBS, bound antibody was detected by fluorescence of rhodamine-labeled mouse anti-rabbit IgG (Boehringer Mannheim).

Detection of FGF-BP protein:

Serum-free media conditioned overnight by the different cell lines at approximately 80% confluency were harvested and

*Trade-mark

proteins present in 500 μ l of undiluted, and PBS-diluted (1:3, 1:10 and 1:30) conditioned media were immobilized by micro-filtration onto nitrocellulose membrane with a dotblot apparatus (Biorad). After washing in Tris-buffered saline, (TBS) and blocking of free sites with 5% skim milk, affinity-purified rabbit IgG raised against a human FGF-PB/GST fusion protein was used for detection of the immobilized antigen with an ECL detection assay (Amersham).

Ribozyme sites were numbered according to their cleavage sites relative to the translation start site in the FGF-BP mRNA. Ribozyme target sites were of the formulas:

gtgcactc 258 aattggac (Seq. 2)

agagagtc 332 tattggaa (Seq. 3)

gacacgtc 697 atgctaata (Seq. 4)

cgtaagtc 753 cctctgta (Seq. 5)

The ribozyme directed against Seq. 5 (a non-translational region) were not effective.

Ribozyme expression vectors were generated and evaluated for in vitro catalytic activity. Ribozyme expression was under the control of a CMV promotor. Each of the individual ribozymes, as well as a multimerized construct containing all of the ribozymes in tandem, were found to cleave the FGF-BP mRNA at the predicted cleavage sites. After in vitro run-off transcription using T7 polymerase, ³²P-labeled FGF-BP transcripts were incubated with unlabeled ribozyme transcripts for different times and the resulting products were separated on a polyacrylamide gel. The expected sizes of the intact FGF-BP transcript (812 nts) and the cleavage products (491 and 321 nts) obtained with Rz258 were noted of particular interest.

While the ribozymes tested were 16 bases in length, ribozymes of 8 to 30 bases targeting any of the transcriptional mRNA would be suggested as useful, with ribozymes of 16 to 20 bases are preferred.

To assess the in vivo efficacy of the ribozyme constructs, the targeted FGF-BP was expressed (alone or in combination with different ribozyme constructs) by stable transfection of FGF-

BP-negative SW-13 cells. More specifically, SW-13 cells were stably transfected with an expression vector for FGF-BP (5 μ g) and the different ribozyme expression vectors or the empty vector (20 μ g). (The Rz753 is targeted to the 3'-untranslated region not included in the expression vector, and could serve as an additional negative control.) It was found that co-transfection of ribozymes Rz258, Rz697 or Rz332 was effective in reducing transcript levels of the transfected gene. A multimerized ribozyme construct containing all ribozymes in tandem had similar efficacy.

In a bioassay to study effects of ribozyme-targeting, it was found that soft agar colony formation of the FGF-BP-transfected SW-13 cells was prevented by co-expression of Rz258, Rz697 or Rz 332. In addition, bFGF release for FGF-BP-transfected cells was reduced to background levels and tumorigenicity in nude mice was reversed upon co-transfection of these ribozyme constructs. It was also shown that Rz753 served as an additional negative control, since its target sequence in the 3'-UTR is not contained in the FGF-BP expression vector. Indeed, co-transfection with expression vector Rz753 did not affect the in vitro or in vivo phenotype of the FGF-BP-transfected SW-13 cells.

To evaluate the FGF-BP as a potential angiogenic switch for squamous cell carcinoma (SCC), endogenous FGF-BP in human ME-180 cells was reduced to varying residual levels using Rz332 and Rz697. These two ribozymes were selected because the target site of Rz258 is very close to Rz332 and both ribozymes had been equally effective in the previous studies. As shown on Northern blot, Rz332 and Rz697 decreased FGF-BP mRNA significantly by 57% and 20%, respectively ($p < 0.001$). Quantitative slot blots revealed a similar reduction of the FGF-BP protein.

Proliferation of the ME-180 cells was not altered by the reduction of FGF-BP, suggesting that FGF-BP is not limiting for in vitro phenotype. In addition to FGF-BP, ME-180 cells also produce bFGF, but no aFGF. Since FGF-BP can release bFGF from the extracellular matrix, the conditioned media from the ME-180

61181-89

8

cell lines were concentrated and partially purified by heparin-affinity chromatography. Peak activities of FGF-BP elute at 1 M NaCl, whereas bFGF elutes at salt concentration of ≥ 1.5 M NaCl from heparin-Sepharose*. In a bioassay, aliquots of the 1M and 2M elutes of ME-180 cells stimulated colony formation of SW-13 cells. FGF-BP stimulates colony formation of these cells by releasing their endogenous bFGF stored in the extracellular matrix. Addition of a neutralizing antibody to bFGF inhibited the stimulation by the 1M and 2M NaCl fractions, whereas a control anti-aFGF antibody showed no effect. The reduction of FGF-BP decreased the amount of stimulating activity of the 1M as well as the 2M fractions. Again, the treatment with the neutralizing antibodies confirmed that the stimulation was due to bFGF. This demonstrated that biologically active bFGF is released into the media of ME-180 cells (the 2 M NaCl fraction) and that this release is dependent on the production of FGF-BP. In parallel with the release of bFGF into media of FGF-BP depleted cells, cellular bFGF was increased.

One million each of different ME-180 cell lines in 0.1 ml of serum-free media were injected subcutaneously into female athymic nude mice and the tumor size measured. The tumors were surgically removed when they reached a size of 50 to 70 mm² and subsequently analyzed for FGF-BP mRNA levels with Northern analysis. Tumor growth of the ME-180 cells was reduced in parallel with the reduced levels of the FGF-BP mRNA. Even the rather small reduction of endogenous FGF-BP mRNA by only 20% using the R2697 group resulted in a 35% reduction of the size of subcutaneous tumors one month after the injection of tumor cells. A further reduction of FGF-BP mRNA resulted in a further (81%) reduced tumor growth.

To assess whether a selection against the FGF-BP-depleted subpopulations of cells occurred in vivo, FGF-BP mRNA levels were analyzed in six different tumors in each of the three groups. For this, tumors were allowed to grow to a size of 50-70 mm². The relative mRNA levels in the tumors after in vivo growth correlated perfectly with the native mRNA levels found

*Trade-mark

in the cultured tumor cells prior to their inoculation into animals. Obviously, no in vivo selection occurs. This suggests that the overall stimulus for the tumor cells towards the tumor stroma drives extension of the stroma and no individual, high-producer tumor cell will have a selection advantage in this setting.

Reduction of FGF-BP did not alter the well-differentiated morphology of the ME-180 tumors. However, a significant reduction of the number of microvessels was observed in the FGF-BP depletion tumors after highlighting endothelial cell lining with an anti-CD31 antibody.

A strong developmental regulation of FGF-BP in the intestine was observed, and the gene product expression in colon cancer was found in cell lines and samples. To assess whether the findings on the role of FGF-BP in SCC cells may also be valid for FGF-BP-positive colon cancer cells, LS174T cells were used as a model. From these highly tumorigenic cells several derivative cell lines were stably transfected with the empty control vector or with a ribozyme expression vector for Rz332. In the in vitro characterization of the cells, it was found that ribozyme transfection reduced the FGF-BP mRNA levels by 70% and the amount of secreted FGF-BP protein by 60%, but that this did not affect proliferation of the cells in culture. In the study, the LS174T cells were stably transfected with the empty vector or the Rz332 vector and analyzed for FGF-BP mRNA and secreted protein. The tumor xenograft studies in these animals showed a significant reduction of the growth (by 60%, $p < 0.001$) and of tumor angiogenesis by $23.1 \pm 11.3\%$ ($n=4$ blinded observers, $p < 0.05$) of the FGF-BP-reduced colon cancer cells. These results have been reproduced in a second independent study with a further ribozyme-transfected and FGF-BP-reduced LS174T derivative cell line (LS174T/Rz332-1).

It was previously known that FGF-BP is expressed in SCC cell lines from different organs and that at very high levels in almost all SCC tumor tissue specimen from patients. This gene product also was found in some colon cancer samples and

cell lines. In contrast, normal adult tissues did not appear to express FGF-BP mRNA, as evidenced by Northern blotting. It was observed that FGF-BP expression is up-regulated transcriptionally by the phorbol ester TPA in cultured cells and increased dramatically in the final stages of carcinogen-induced skin tumors in mice as well as in carcinogen-treated skin grafted onto SCID mice.

The genetic approach of ribozyme-targeting to down-regulate spontaneous expression of FGF-BP mRNA as taught herein is an appropriate means of decreasing tumor growth and angiogenesis. The ribozymes of the invention may be administered using viral or non-viral transfer vectors. Synthetic ribozymes may also be administered. The ribozymes in pharmaceutically acceptable carrier may be administered in any manner that will result in contacting the malignant cells, including, for example, intravenous, intra-arterial, intra-ocular, intranasal, transvaginal, topical, intrathecal, intraperitoneal and subcutaneous administration. Compositions containing the ribozymes may be administered directly into the tumor or into the arterial blood supply to the tumor.

The ribozymes of the invention may be administered in means known for the administration of oligonucleotides such as in viral expression vectors or in liposomes.

The antisense oligonucleotides can be phosphorothioated and then encapsulated in liposomes such as those disclosed in WO 95/11670. The preferred liposomes for this purpose are multilamellar having a solute entrapped in its aqueous compartments. The following is an example of means for making such liposomal compositions.

Distearoyl phosphatidylcholine (DSPC), cholesterol and dimethyldioctadecylammonium bromide (DDAB) liposomes (50 mole percent DSPC, 40 mole percent cholesterol and 10 mole percent DDAB) are prepared by dissolving 3.59 mg DSPC, 1.54 mg cholesterol and .631 mg DDAB in methanol in a 100 ml round-bottom flask. HEPES buffer (10 mM HEPES, 150 mM NaCl) is added to the flask and the sample is roto-evaporated to remove organic solvent. An additional 5 ml HEPES buffer is added to

WO 99/15703

PCT/US98/20208

11

the flask, which is then heated to 65 °C. The dried film is hydrated over night with 4 mg oligonucleotides of the invention in 4 ml HEPES buffer under refrigeration. The film is dispersed by centrifugation at 20,000 rpm for 20 minutes followed by sonication. The liposomes may then be frozen and rethawed several times to produce multilamellar vesicles.

5

SEQUENCE LISTING

10 <110> WELLSTEIN, ANTON
 CZUBAYKO, FRANK
 <120> INHIBITION OF AN FGF-BINDING PROTEIN USING RIBOZYMES
 <130> 23521.0076
 <140> 09/160,588
 <141> 1998-09-25
 <160> 5
 <170> PatentIn Ver. 2.1

20 <210> 1
 <211> 1163
 <212> DNA
 <213> Homo sapiens
 <400> 1
 ctctacctga cacagctgca gcctgcaatt cactcccact gcctgggatt gcactggatc 60
 cgtgtgctca gaacaagggtg aacgcccagc tgcagccatg aagatctgta gcctcaccct 120
 gctctccttc ctctactgg ctgctcaggt gctcctgggtg gaggggaaaa aaaaagtga 180
 gaatggactt cacagcaaag tgggtctcaga acaaaaggac actctgggca acaccagat 240
 taagcagaaa agcaggcccg ggaacaaagg caagtttgtc accaaagacc aagccaactg 300
 cagatgggct gctactgagc aggaggagg catctctctc aaggttgagt gcactcaatt 360
 ggacatgaa ttttcctgtg tctttgctgg caatccaacc tcatgcctaa agctcaagga 420
 tgagagagtc tattggaaac aagttgcccg gaatctgctc tcacagaaag acatctgtag 480
 atattccaag acagctgtga aaaccagagt gtgcagaaag gattttccag aatccagtct 540
 taagctagtc agctccactc tatttgggaa cacaaagccc aggaaggaga aaacagagat 600
 gtccccccagg gagcacatca agggcaaaga gaccacccc tctagcctag cagtgaccca 660
 30 gaccatggcc accaaagctc ccgagtgtgt ggaggaccca gatatggcaa accagaggaa 720
 gactgccctg gagttctgtg gagagacttg gagctctctc tgcacattct tcctcagcat 780
 agtgcaggac acgtcatgct aatgaggtca aaagagaacg ggttccttta agagatgtca 840
 tgtcgtaagt ccctctgtat actttaagc tctctacagt cccccaaaa tatgaacttt 900
 tgtgcttagt gagtgcaacg aaatatttaa acaagttttg tatttttttg ttttgtgttt 960
 tggaatttgc cttatttttc ttggatgcga tgttcagagg ctgtttcctg cagcatgtat 1020
 ttccatggcc cacacagcta tgtgtttgag cagcgaagag tctttgagct gaatgagcca 1080
 gagtgataat ttcagtgcaa cgaactttct gctgaattaa tggtaataaa actctgggtg 1140
 tttttcaaaa aaaaaaaaaa aaa 1163

40 <210> 2
 <211> 16
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Ribozyme
 target site
 <400> 2
 gtgcactcaa ttggac 16

50 <210> 3
 <211> 16
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Ribozyme
 target site
 <400> 3
 agagagtcta ttggaa 16

60 <210> 4
 <211> 16
 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Ribozyme
target site

<400> 4

gacacgtcat gctaat

16

10

<210> 5

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Ribozyme
target site

<400> 5

cgtaagtccc tctgta

16

61181-89

14

CLAIMS:

1. A method of inhibiting expression of fibroblast growth factor binding protein (FGF-BP) in a cell in vitro by administration of FGF-BP ribozymes which cleave FGF-BP mRNA,
5 wherein said ribozymes are hammerhead ribozymes as depicted in (FIG. 1) having a target cleavage site selected from the group consisting of 258, 332 and 697 nucleotides downstream of the translation initiation site of FGF-BP mRNA, and wherein said ribozymes comprise antisense
10 sequences 5' and 3' to the ribozyme catalytic center that direct the ribozymes to said target cleavage site, such that expression of FGF-BP is inhibited.
2. A method of claim 1 wherein the ribozymes are administered in the form of liposomes.
- 15 3. A method of claim 1 wherein the ribozymes are administered in viral vectors.
4. The method of any one of claims 1 to 3, wherein said ribozymes are in a pharmaceutically acceptable carrier.
5. The method of any one of claims 1 to 4, wherein
20 said FGF-BP is encoded by the DNA sequence of SEQ ID NO: 1.
6. The method of any one of claims 1 to 5, wherein said 5' and 3' antisense sequences together are 8 to 30 nucleotides long.
7. The method of any one of claims 1 to 6, wherein
25 said antisense sequences are selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 5.
8. A composition comprising: (a) ribozymes which specifically cleave FGF-BP mRNA, wherein said ribozymes are

61181-89

15

hammerhead ribozymes as depicted in (FIG. 1) having a target cleavage site selected from the group consisting of 258, 332 and 697 nucleotides downstream of the translation initiation site of FGF-BP mRNA, and wherein said ribozymes
5 comprise antisense sequences 5' and 3' to the ribozyme catalytic center that direct the ribozymes to said target cleavage site; and (b) a pharmaceutically acceptable carrier.

9. The composition of claim 8, wherein said ribozymes
10 are comprised in liposomes.

10. The composition of claim 8 or 9, wherein said FGF-BP is encoded by the DNA sequence of SEQ ID NO: 1.

11. The composition of any one of claims 8 to 10, wherein said 5' and 3' antisense sequences are together at
15 least 8 to 30 nucleotides long.

12. The composition of claim 11, wherein said antisense sequences are selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 5.

20 13. Use in the preparation of a medicament for reducing tumor growth in a subject by reducing the expression of FGF-BP mRNA in said tumor of the composition of any one of claims 8 to 12.

14. The use of claim 13 wherein said medicament is for
25 direct injection into said tumor.

15. Use for reducing tumor growth in a subject by reducing the expression of FGF-BP mRNA in said tumor of the composition of any one of claims 8 to 12.

61181-89

16

16. The use of claim 15, wherein said composition is formulated for direct injection into said tumor.

17. A commercial package comprising the composition of any one of claims 8 to 12 together with instructions for use
5 for reducing tumor growth in a subject.

SMART & BIGGAR

OTTAWA, CANADA

PATENT AGENTS




FIG. 1

FGF-BP mRNA ribozyme target sites

