

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2008 (15.05.2008)

PCT

(10) International Publication Number
WO 2008/055877 A1

(51) International Patent Classification:
C07D 231/44 (2006.01) *A61K 31/415* (2006.01)
A01N 43/56 (2006.01) *A61P 33/00* (2006.01)
A01N 43/00 (2006.01)

(74) Common Representative: BASF SE; 67056 Ludwigshafen (DE).

(21) International Application Number:
PCT/EP2007/061891

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
5 November 2007 (05.11.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/865,178 10 November 2006 (10.11.2006) US
60/913,617 24 April 2007 (24.04.2007) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): BASF SE [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SUKOPP, Martin** [DE/DE]; Richard-Wagner-Str. 33, 68165 Mannheim (DE). **KUHN, Oliver** [DE/LU]; 17, Rue du Pont, L-06581 Rosport (LU). **GRÖNING, Carsten** [DE/DE]; Normannenstr.29, 68259 Mannheim (DE). **KEIL, Michael** [DE/DE]; Fontanestr.4, 67251 Freinsheim (DE). **LONGLET, Jon J.** [US/US]; 3116 Nashville Avenue, Nederland, TX 77627 (US).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(54) Title: PROCESS FOR THE SULFINYLATION OF A PYRAZOLE DERIVATIVE

(57) Abstract: The present invention relates to a process for the sulfinylation of a pyrazole derivative, characterized in that 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3- carbonitrile (II) is reacted with a sulfinylating agent selected from trifluoromethylsulfonic acid, trifluoromethylsulfonic acid anhydride, and a trifluoromethylsulfinate alkaline or alkaline earth metal salt and mixtures of the acid and/or the salt(s), in the presence of at least one amine acid complex wherein the amine(s) are selected from tertiary amines and the acid(s) are selected from hydrofluoric, hydrochloric, hydrobromic and hydroiodic acid and sulfonic acid derivatives, and with the addition of a halogenating agent.

WO 2008/055877 A1

Process for the sulfinylation of a pyrazole derivative

The present invention relates to a novel process for the sulfinylation of a pyrazole derivative, characterized in that 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile (II) is reacted with a sulfinylating agent selected from trifluoromethylsulfonic acid, trifluoromethylsulfonic acid anhydride, and a trifluoromethylsulfinate alkaline or alkaline earth metal salt and mixtures of the acid and/or the salt(s), in the presence of at least one amine acid complex wherein the amine(s) are selected from tertiary amines and the acid(s) are selected from hydrofluoric, hydrochloric, hydrobromic and hydroiodic acid and sulfonic acid derivatives, and with the addition of a halogenating agent.

The sulfinylation of a pyrazole-type compound refers to the substitution of a hydrogen atom on a pyrazole heterocycle carbon atom by an RS(=O)- group.

The direct sulfinylation of various organic molecules (not including pyrazole derivatives) employing a mixture of P(O)Cl₃ and CF₃S(O)ONa has been described in T. Billard, A. Greiner, B. R. Langlois, Tetrahedron 55 (1999), p. 7243 – 7250. Likewise, C. Wakselman, M. Tordeux, C. Freslon, L. Saint-Jalmes, Synlett 2001, p. 550-552 teaches that direct sulfinylation of aromatic compounds takes place by CF₃S(O)ONa or CF₃S(O)OK in the presence of triflic acid (CF₃S(O)₂OH).

Processes for the direct sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile (II) have been described in EP-A 668 269, EP-A 1 331 222, CN-A 1374298, and in Y.Huilong, M. Zengeng, W. Shujuan, J. Hebei University of Science of Technology, Vol. 25(2), Sum 69 (2004), Serial no. 1008-1542 (2004) 02-0018-03.

In EP 668 269, the sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile (II) with trifluoromethylsulfonic acid CF₃S(O)OH and its derivatives CF₃S(O)Cl, CF₃S(O)ONa, CF₃S(O)N(CH₃)₂, or CF₃S(O)N(CH₂CH₃)₂ has been described. As chlorinating agent, phosgene, chloroformates, PCl₅ and SOCl₂ are mentioned. It is described that a reagent (“ compound C”) chosen from the group consisting of the tosylates, hydrochlorides and mesylates of a primary, secondary, or tertiary amine, preferably of dimethylamine, of pyridine, of trimethylamine, of diethylamine or of isopropylamine or gaseous hydrogen chloride, optionally in the presence of an equimolar amount of para-toluenesulfonic acid, may be added to complete the reaction. Examples are given for the following combinations of reactants:

CF₃S(O)Cl, dimethylamine p-tosylate;

CF₃S(O)Cl, pyridine hydrochloride salt;

CF₃S(O)N(CH₃)₂, p-toluolsulfonic acid, hydrochloric acid;

CF₃S(O)Cl, dimethylamine p-tosylate, hydrochloric acid; and
CF₃S(O)ONa, dimethylamine p-tosylate, SOCl₂.

The reactions carried out with CF₃S(O)Cl as the sulfinylating agent give the highest
yield of the final product.

5

The process described in CN-A 1374298 has been developed to overcome certain
shortcomings of the process described in EP 668 269. CN-A 1374298 cites that
CF₃S(O)Cl is extremely unstable, CF₃S(O)N(CH₃)₂ and CF₃SOOH are relatively difficult
to prepare, that the reactivity CF₃S(O)ONa is not high, and that the yield in the
10 sulfinylation reaction is correspondingly relatively low. CN-A 1374298 describes the
sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-
carbonitrile (II) with the potassium salt of trifluoromethylsulfonic acid, CF₃S(O)OK, or
mixtures of the potassium and the sodium salt of trifluoromethylsulfonic acid,
CF₃S(O)OK with CF₃S(O)ONa, wherein the sulfinylating agent is combined with POCl₃,
15 PCl₃, SOCl₂, COCl₂, or trichloromethylchloromethanoate. Optionally, the amine acid
complex dimethylamine p-tosylate can be added to complete the reaction.

Examples are given for the following combinations of reactants:

CF₃S(O)OK; dimethylamine p-tosylate; POCl₃; and

CF₃S(O)OK/Na; dimethylamine p-tosylate; SOCl₂.

20

Huilong et al. describes the reaction of 5-amino-1-[2,6-dichloro-4-
(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile (II) with the sodium salt of
trifluoromethylsulfonic acid (CF₃S(O)ONa), dimethylamine p-tosylate, and SOCl₂, with
addition of catalytic amounts of DMF (dimethylformamide).

25

In EP-A 1 331 222, 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-
carbonitrile (II) is sulfinylated using N-trifluoromethylsulfinylsuccinimide as the
sulfinylating agent in the presence of triethylamine and without the addition of a
chlorinating agent. The intermediate N-trifluoromethylsulfinylamino-pyrazole is isolated
30 and under the conditions of a Thia-Fries rearrangement transformed into the final
product 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-
pyrazole-3-carbonitrile.

Thus, the sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-
35 pyrazole-3-carbonitrile (II) to the final product 5-amino-1-[2,6-dichloro-4-
(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile (common
name: fipronil) has received considerable attention in the literature, the focus being on
the optimization of the sulfinylating agent.

40 However, and also as cited in the recent review article “ research progress on synthe-
sis of fipronil and its main intermediates” , Chinese Journal of Pesticides, 2004, Vol.

43, no.12, 529-531, the sulfinylation of the pyrazole intermediate was still found not to be generally suitable for a large scale industrial production.

5 The reaction product of the present sulfinylation is fipronil, which is a market insecticide of considerable importance. Generally, technical manufacturing processes of pesticides have to fulfill high requirements with regard to yield and purity of the product, for reasons of profitability, but also, most important, in order to avoid the presence of potentially noxious side products. This is of special relevance for fipronil as it is also used in animal health products and therefore also comes into contact with companion animals.

10

It is also a legal requirement for technical manufacturing processes to avoid exposure of the plant employees but also of the environment to reagents which can have adverse effects to the health of the employees or environment. Therefore, it is desirable to have a technical manufacturing process at hand which avoids the use of toxic reagents such as $\text{CF}_3\text{S}(\text{O})\text{Cl}$.

15

Furthermore, during the scale up of a process from the laboratory scale to a technical scale, problems may arise that were not as such or to the respective extend foreseeable in the laboratory.

20

- For example, the filling and/or dissolution of voluminous starting materials may take much longer on a big scale than in a small vessel, with the effect that the kinetic of the reaction is significantly changed, and thereby the conversion and the product spectrum.

25

- Another example that can be mentioned is the appearance of side products that are, due to solubility or texture, difficult to separate from the desired main product on a large scale. Problems with extraction, filtration and clogging of filters may occur. Insoluble starting materials or reaction (side) products may also challenge the agitation, heat dissipation or pumping thus leading to inhomogenous reaction mixtures.

30

- Yet another challenge is the control of the course of the reaction temperature in large-scale processes. The temperature rates generally are lower which may have an influence on the side product spectrum. As high reaction temperatures and/or aggressive reaction media may cause corrosion, and also because of economic reasons, moderate reaction conditions (low temperatures) are preferred.

35

- Hygroscopic properties of solids can complicate reactions that favorably are conducted under essentially water free conditions. For example, when the process as defined above is conducted using an amine acid complex wherein the acid is H_2SO_4 – and not an acid as defined for the inventive process – the yield of the reaction is extremely poor.

40

- Nonreactive catalysts – such as tertiary amines, e.g. triethylamine acids – are preferably used in the inventive process in order to avoid side reactions. Secondary or primary amines can react with the sulfinylating agent.

- With a view to facilitating the work-up, reagents are preferably used that can be removed by distillation. Solids are removed by washing with acidic or alkaline solvents. It is not advantageous to employ reagents that have phase transfer catalytic properties which may hinder a phase separation during work-up.

5

Against this background and facing the fact that one of the essential starting materials for the current industrial production of fipronil is CF_3Br (see e.g. WO 01/30760) which exhibits high environmental toxicity and is scheduled for production phase out by the Montreal Protocol on Substances that Deplete the Ozone Layer (it may then be used only as a feedstock material), it was an object of the present invention to develop a new, large scale industrial process for the manufacturing of fipronil which gives fipronil in high purity and yield while avoiding the use of dangerous reagents and avoiding problems with the technical reaction control.

10

Accordingly, the process defined at the outset was found. The obtained product fipronil is suitable for use as a pesticide for agricultural uses as well as non-crop uses for combating pests. Also, the obtained fipronil is suitable for use in the animal health field for combating animal health pests and parasites, especially for the long lasting protection against fleas and ticks on mammals.

20

Thus, the current invention also pertains to pesticidal or parasitocidal composition containing 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile as prepared by the inventive process.

25

Likewise, the present invention relates to a method for the control of insects, acarids or nematodes by contacting the insect, acarid or nematode or their food supply, habitat, breeding ground or their locus with a pesticidally effective amount of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by the inventive process as well as to a method of protecting growing plants from attack or infestation by insects, acarids or nematodes by applying to the foliage or the seeds of the plants, or to the soil or water in which they are growing, a pesticidally effective amount of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by the inventive process. According to these methods, 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile is usually applied in an amount of from 5 g/ha to 2000 g/ha.

30

35

Moreover, the present invention relates to a method for treating, controlling, preventing or protecting animals against infestation or infection by parasites which comprises orally, topically or parenterally administering or applying to the animals or their habitat a parasitocidally effective amount of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-

40

(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by the inventive process or its veterinarily acceptable enantiomers or salts.

Also, the present invention relates to a process for the preparation of a composition for
5 treating, controlling, preventing or protecting animals against infestation or infection by
parasites which comprises admixing 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-
4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by the inventive process or
its enantiomers or veterinarily acceptable salts with a veterinarily acceptable carrier.
The composition may either be a concentrate or contains 5-amino-1-[2,6-dichloro-4-
10 (trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile in a parasiti-
cidally effective amount.

While there are examples given for certain amine acid complexes added at the begin-
ning or during the course of the sulfinylation reaction, there is no teaching as to the
15 crucial importance of the specific nature of the amine acid complex with regard to reac-
tion control or to the yield and / or the purity of the final product fipronil.

In none of the prior art documents, mention is made of the favorable combination of the
key features of the present invention, namely employing
20 a) trifluoromethylsulfinic acid, trifluoromethylsulfinic acid anhydride, or a trifluoromethyl-
sulfinate alkaline or alkaline earth metal salt, or mixtures of the acid and/or the salt(s)
as sulfinylating agents and
b) at least one amine acid complex wherein the amine(s) are selected from tertiary
amines and the acid(s) are selected from hydrofluoric, hydrochloric, hydrobromic and
25 hydroiodic acid and sulfonic acid derivatives.


In EP-A1 668 269, the sulfinylating agents $\text{CF}_3\text{S}(\text{O})\text{Cl}$, $\text{CF}_3\text{S}(\text{O})\text{N}(\text{CH}_3)_2$,
 $\text{CF}_3\text{S}(\text{O})\text{N}(\text{CH}_2\text{CH}_3)_2$, $\text{CF}_3\text{S}(\text{O})\text{OH}$ and $\text{CF}_3\text{S}(\text{O})\text{ONa}$, together with phosgene or SOCl_2
or $\text{ClCO}_2\text{C}_2\text{H}_5$ are described as being equivalent alternatives in processes leading to
30 fipronil and its close analgon 5-amino-1-[2,6-dichloro-4-(pentafluorosulfanyl)phenyl]-4-
(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile. One example is given for the use of
 $\text{CF}_3\text{S}(\text{O})\text{ONa}$, but not together with tertiary amines as part of the amine acid complex.

Also, in EP-A1 668 269, the following list of preferred amines is given: tosylates, hydro-
35 chlorides or mesylates of dimethylamine, pyridine, trimethylamine, diethylamine, iso-
propylamine. An example is given for pyridine hydrochloride as amine acid complex,
however used together with $\text{CF}_3\text{S}(\text{O})\text{Cl}$ as the sulfinylating agent.

The novel subject of the present invention thus is the sulfinylation of 5-amino-1-[2,6-
40 dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile (II) with a sulfinylating
agent selected from trifluoromethylsulfinic acid, trifluoromethylsulfinic acid anhydride

and a trifluoromethylsulfinate alkaline or alkaline earth metal salt and mixtures of the acid and/or the salt(s), in the presence of at least one amine acid complex wherein the amine(s) are selected from tertiary amines and the acid(s) are selected from hydrofluoric, hydrochloric, hydrobromic and hydroiodic acid and sulfonic acid derivatives, and
 5 with the addition of a halogenating agent.

A reaction scheme may be depicted as follows:



The sulfinylating agent is preferably selected from trifluoromethylsulfonic acid, trifluoro-
 10 methylsulfonic acid anhydride, trifluoromethylsulfinate sodium salt, trifluoromethylsulfi-
 nate potassium salt, and mixtures of these.

According to a preferred embodiment of the present invention, trifluoromethylsulfinate
 sodium salt is used as the sulfinylating agent.

15

According to another preferred embodiment of the present invention, trifluoromethyl-
 sulfinate potassium salt is used as the sulfinylating agent.

According to yet another preferred embodiment of the present invention, trifluoro-
 20 methylsulfonic acid is used as the sulfinylating agent.

According to yet another preferred embodiment of the present invention, trifluoro-
 methylsulfonic acid anhydride is used as the sulfinylating agent.

25 According to a preferred embodiment of the present invention, a mixture of the
 trifluoromethylsulfinate sodium and potassium salts, in a mixing ratio of from 0,01 :
 99,99 weight % to 50 : 50 weight % is used as the sulfinylating agent.

30 Preferably, 1.0 to 1.35 molar equivalents, most preferably 1.2 molar equivalents, of the
 sulfinylating agent relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-
 pyrazole-3-carbonitrile are used.

In a preferred embodiment, the sulfinylating agent is dried before its use until it is essentially water-free. "Water free" means that the content of water in the solid does not exceed 5 ppm to 100 ppm.

- 5 The halogenating agent is selected from thionylchloride, thionylbromide, phosphorochloride, oxalylchloride, phosgen, triphosgen ((CCl₃)₂C(=O)), chloroformates, phosphorpentachloride, phosphortrichloride, trichloromethylchloromethanoat, and xylenesulfonic acid chloride.
- 10 According to a preferred embodiment of the present invention, a chlorinating agent is used as the halogenating agent. Preferably, thionylchloride or phosphorochloride are used as the chlorinating agent.


15 According to another preferred embodiment of the present invention, phosphorochloride is used as the chlorinating agent.

Most preferably, thionylchloride is used as the chlorinating agent.

20 Preferably, 1.15 to 1.35 molar equivalents, most preferably about 1.2 molar equivalents, of the halogenating agent relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile are used.

25 We have found that the choice of the amine acid complex plays a key role in the sulfonylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile. Critical properties that influence the sulfonylation reaction are steric (bulk) properties and molecular weight, and also pKs value and solubility.

30 The sulfonylation reaction of the present invention is a one-pot synthesis of a two-step reaction. The first step involves the addition of the CF₃S(O)- group to the amino group of the pyrazole ring. In a second step, fipronil is formed via a Thia-Fries rearrangement:



The amine acid complex has two functions in this two-step reaction: (1) when sulfinylates are used as sulfinylating agents, it catalyzes the activation of the sulfinylate with the halogenating agent via intermediate formation of sulfinic acid. For this, catalytic amounts of 0,01 to 1,0 molar equivalents of amine acid complex relative to the pyrazole compound II are needed. (2) It accelerates the Thia-Fries rearrangement and has a significant influence on selectivity. With a view to obtain high yields and high purity, overall amounts of above 1 molar equivalents of amine acid complex relative to the pyrazole compound II are advantageously used for step two.

Preferred are amine acid complexes exhibiting low or essentially no hygroscopicity, as the sulfinylating process of the present invention advantageously is conducted in the essential absence of water (i.e. below 5 to 100 ppm of water).

Preferred tertiary amines are

alkyl amines such as trimethylamine, triethylamine, tripropylamine, triisopropylamine, tributylamine, dimethyl ethyl amine, diethyl methylamine, dimethyl n-propyl amine, diisopropyl ethyl amine, DBU (1,8-diazabicyclo[5.4.0]undec-7-ene), DBN (1,5-diazabicyclo[4.3.0]non-5-ene), methyl morpholine, ethyl morpholine, N,N-dimethylaniline, methyl piperidine, methyl pyrrolidine, or methyldibenzylamine; or aromatic amines such as pyridine, DMAP (dimethylaminopyridine), collidine, lutidine, pyrimidine, pyrazine, or piperazine.

In a further preferred embodiment, the tertiary amine is selected from triethylamine, tripropylamine, triisopropylamine, dimethyl ethyl amine, diethyl methylamine, dimethyl n-propyl amine, diisopropylethylamine, DBU (1,8-diazabicyclo[5.4.0]undec-7-ene), DBN (1,5-diazabicyclo[4.3.0]non-5-ene), methyl morpholine, ethyl morpholine, N,N-dimethylaniline, methyl piperidine, methyl pyrrolidine, and methyldibenzylamine, DMAP (dimethylaminopyridine), collidine, lutidine, pyrimidine, pyrazine, and piperazine.

In another preferred embodiment, the tertiary amine is selected from trimethylamine, triethylamine, dimethyl ethyl amine, diethyl methylamine, dimethyl n-propyl amine, methyl morpholine, N,N-dimethylaniline, methyl piperidine, methyl pyrrolidine, methyldibenzylamine, and pyridine.

Especially preferred are trimethylamine, triethylamine, dimethyl ethyl amine, dimethyl n-propyl amine, or pyridine. Very preferred are triethylamine and pyridine. In a very preferred embodiment, the amine is triethylamine. Also, in a very preferred embodiment the amine is pyridine. Also, in a very preferred embodiment the amine is trimethylamine. Also, in a very preferred embodiment at least one of the alkyl groups attached to the nitrogen atom of the amine is a methyl group. Also, in a very preferred

embodiment the nitrogen atom of the amine group is sp^3 hybridized, i.e. it does not form a double bond to a neighbour atom.

5 Preferred acids of the amine acid complex for use in the present invention are hydrochloric acid, hydrofluoric acid, hydrobromic acid, hydroiodic acid or sulfonic acid derivatives such as aromatic sulfonic acids, e.g. p-toluenesulfonic acid, benzenesulfonic acid, 4-ethyl benzenesulfonic acid, 4-chlorobenzenesulfonic acid, xylene sulfonic acid, 2,3-dimethylbenzene sulfonic acid, 2,4-dimethylbenzene sulfonic acid, 2,5-dimethylbenzene sulfonic acid, 2,6-dimethylbenzene sulfonic acid,
 10 1-naphthalenesulfonic acid, 2-naphthalenesulfonic acid, mixtures of two or more of the isomers of dimethylbenzene sulfonic acids, or mesitylene sulfonic acid; or alkyl sulfonic acids, e.g. methane sulfonic acid or camphor sulfonic acid; or haloalkylsulfonic acids, e.g. trifluoromethylsulfonic acid.

15 More preferred acids of the amine acid complex for use in the present invention are hydrochloric acid, hydrofluoric acid, or sulfonic acid derivatives such as aromatic sulfonic acids, e.g. p-toluenesulfonic acid, benzenesulfonic acid, xylene sulfonic acid, mesitylene sulfonic acid; or alkyl sulfonic acids, e.g. methanesulfonic acid; or haloalkylsulfonic acids, e.g. trifluoromethylsulfonic acid.

20 Especially preferred are acids with a pKs-value of below 2.

From the choice of amine acid complexes as used in the present invention, those with a pKa below 6, preferably 5, and above 10 are preferred.

25 Most preferred acids are hydrochloric acid, p-toluenesulfonic acid, xylene sulfonic acid, or benzene sulfonic acid.

The most preferred acid with regard to optimized yield is hydrochloride acid.

30 Preferred amine acid complexes Q are listed in Table 1 which follows.

Table 1

No.	amine	Acid
Q-1	$N(CH_3)_3$	HF
Q-2	$N(CH_3)_3$	HCl
Q-3	$N(CH_3)_3$	HBr
Q-4	$N(CH_3)_3$	HI
Q-5	$N(CH_3)_3$	p-toluene sulfonic acid
Q-6	$N(CH_3)_3$	benzene sulfonic acid

No.	amine	Acid
Q-7	$N(CH_3)_3$	xylene sulfonic acid
Q-8	$N(CH_3)_3$	methane sulfonic acid
Q-9	$N(CH_3)_3$	trifluoromethyl sulfonic acid
Q-10	$N(CH_3)_3$	mesitylene sulfonic acid
Q-11	$N(CH_2CH_3)_3$	HF
Q-12	$N(CH_2CH_3)_3$	HCl
Q-13	$N(CH_2CH_3)_3$	HBr
Q-14	$N(CH_2CH_3)_3$	HI
Q-15	$N(CH_2CH_3)_3$	p-toluene sulfonic acid
Q-16	$N(CH_2CH_3)_3$	benzene sulfonic acid
Q-17	$N(CH_2CH_3)_3$	xylene sulfonic acid
Q-18	$N(CH_2CH_3)_3$	methane sulfonic acid
Q-19	$N(CH_2CH_3)_3$	trifluoromethyl sulfonic acid
Q-20	$N(CH_2CH_3)_3$	mesitylene sulfonic acid
Q-21	$N(CH_2CH_2CH_3)_3$	HF
Q-22	$N(CH_2CH_2CH_3)_3$	HCl
Q-23	$N(CH_2CH_2CH_3)_3$	HBr
Q-24	$N(CH_2CH_2CH_3)_3$	HI
Q-25	$N(CH_2CH_2CH_3)_3$	p-toluene sulfonic acid
Q-26	$N(CH_2CH_2CH_3)_3$	benzene sulfonic acid
Q-27	$N(CH_2CH_2CH_3)_3$	xylene sulfonic acid
Q-28	$N(CH_2CH_2CH_3)_3$	methane sulfonic acid
Q-29	$N(CH_2CH_2CH_3)_3$	trifluoromethyl sulfonic acid
Q-30	$N(CH_2CH_2CH_3)_3$	mesitylene sulfonic acid
Q-31	$N(CH_2CH_2CH_2CH_3)_3$	HF
Q-32	$N(CH_2CH_2CH_2CH_3)_3$	HCl
Q-33	$N(CH_2CH_2CH_2CH_3)_3$	HBr
Q-34	$N(CH_2CH_2CH_2CH_3)_3$	HI
Q-35	$N(CH_2CH_2CH_2CH_3)_3$	p-toluene sulfonic acid
Q-36	$N(CH_2CH_2CH_2CH_3)_3$	benzene sulfonic acid
Q-37	$N(CH_2CH_2CH_2CH_3)_3$	xylene sulfonic acid
Q-38	$N(CH_2CH_2CH_2CH_3)_3$	methane sulfonic acid
Q-39	$N(CH_2CH_2CH_2CH_3)_3$	trifluoromethyl sulfonic acid
Q-40	$N(CH_2CH_2CH_2CH_3)_3$	mesitylene sulfonic acid
Q-41	$N[CH(CH_3)_2]_3$	HF
Q-42	$N[CH(CH_3)_2]_3$	HCl
Q-43	$N[CH(CH_3)_2]_3$	HBr
Q-44	$N[CH(CH_3)_2]_3$	HI

No.	amine	Acid
Q-45	$N[CH(CH_3)_2]_3$	p-toluene sulfonic acid
Q-46	$N[CH(CH_3)_2]_3$	benzene sulfonic acid
Q-47	$N[CH(CH_3)_2]_3$	xylene sulfonic acid
Q-48	$N[CH(CH_3)_2]_3$	methane sulfonic acid
Q-49	$N[CH(CH_3)_2]_3$	trifluoromethyl sulfonic acid
Q-50	$N[CH(CH_3)_2]_3$	mesitylene sulfonic acid
Q-51	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	HF
Q-52	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	HCl
Q-53	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	HBr
Q-54	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	HI
Q-55	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	p-toluene sulfonic acid
Q-56	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	benzene sulfonic acid
Q-57	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	xylene sulfonic acid
Q-58	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	methane sulfonic acid
Q-59	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	trifluoromethyl sulfonic acid
Q-60	$N(CH_2CH_3)[(CH(CH_3)_2)_2]$	mesitylene sulfonic acid
Q-61	DBU	HF
Q-62	DBU	HCl
Q-63	DBU	HBr
Q-64	DBU	HI
Q-65	DBU	p-toluene sulfonic acid
Q-66	DBU	benzene sulfonic acid
Q-67	DBU	xylene sulfonic acid
Q-68	DBU	methane sulfonic acid
Q-69	DBU	trifluoromethyl sulfonic acid
Q-70	DBU	mesitylene sulfonic acid
Q-71	DBN	HF
Q-72	DBN	HCl
Q-73	DBN	HBr
Q-74	DBN	HI
Q-75	DBN	p-toluene sulfonic acid
Q-76	DBN	benzene sulfonic acid
Q-77	DBN	xylene sulfonic acid
Q-78	DBN	methane sulfonic acid
Q-79	DBN	trifluoromethyl sulfonic acid
Q-80	DBN	mesitylene sulfonic acid
Q-81	methyl morpholine	HF
Q-82	methyl morpholine	HCl

No.	amine	Acid
Q-83	methyl morpholine	HBr
Q-84	methyl morpholine	HI
Q-85	methyl morpholine	p-toluene sulfonic acid
Q-86	methyl morpholine	benzene sulfonic acid
Q-87	methyl morpholine	xylene sulfonic acid
Q-88	methyl morpholine	methane sulfonic acid
Q-89	methyl morpholine	trifluoromethyl sulfonic acid
Q-90	methyl morpholine	mesitylene sulfonic acid
Q-91	ethyl morpholine	HF
Q-92	ethyl morpholine	HCl
Q-93	ethyl morpholine	HBr
Q-94	ethyl morpholine	HI
Q-95	ethyl morpholine	p-toluene sulfonic acid
Q-96	ethyl morpholine	benzene sulfonic acid
Q-97	ethyl morpholine	xylene sulfonic acid
Q-98	ethyl morpholine	methane sulfonic acid
Q-99	ethyl morpholine	trifluoromethyl sulfonic acid
Q-100	ethyl morpholine	mesitylene sulfonic acid
Q-101	N,N-dimethylaniline	HF
Q-102	N,N-dimethylaniline	HCl
Q-103	N,N-dimethylaniline	HBr
Q-104	N,N-dimethylaniline	HI
Q-105	N,N-dimethylaniline	p-toluene sulfonic acid
Q-106	N,N-dimethylaniline	benzene sulfonic acid
Q-107	N,N-dimethylaniline	xylene sulfonic acid
Q-108	N,N-dimethylaniline	methane sulfonic acid
Q-109	N,N-dimethylaniline	trifluoromethyl sulfonic acid
Q-110	N,N-dimethylaniline	mesitylene sulfonic acid
Q-111	methyl piperidine	HF
Q-112	methyl piperidine	HCl
Q-113	methyl piperidine	HBr
Q-114	methyl piperidine	HI
Q-115	methyl piperidine	p-toluene sulfonic acid
Q-116	methyl piperidine	benzene sulfonic acid
Q-117	methyl piperidine	xylene sulfonic acid
Q-118	methyl piperidine	methane sulfonic acid
Q-119	methyl piperidine	trifluoromethyl sulfonic acid
Q-120	methyl piperidine	mesitylene sulfonic acid

No.	amine	Acid
Q-121	methyl pyrrolidine	HF
Q-122	methyl pyrrolidine	HCl
Q-123	methyl pyrrolidine	HBr
Q-124	methyl pyrrolidine	HI
Q-125	methyl pyrrolidine	p-toluene sulfonic acid
Q-126	methyl pyrrolidine	benzene sulfonic acid
Q-127	methyl pyrrolidine	xylene sulfonic acid
Q-128	methyl pyrrolidine	methane sulfonic acid
Q-129	methyl pyrrolidine	trifluoromethyl sulfonic acid
Q-130	methyl pyrrolidine	mesitylene sulfonic acid
Q-131	methyl dibenzyl amine	HF
Q-132	methyl dibenzyl amine	HCl
Q-133	methyl dibenzyl amine	HBr
Q-134	methyl dibenzyl amine	HI
Q-135	methyl dibenzyl amine	p-toluene sulfonic acid
Q-136	methyl dibenzyl amine	benzene sulfonic acid
Q-137	methyl dibenzyl amine	xylene sulfonic acid
Q-138	methyl dibenzyl amine	methane sulfonic acid
Q-139	methyl dibenzyl amine	trifluoromethyl sulfonic acid
Q-140	methyl dibenzyl amine	mesitylene sulfonic acid
Q-141	pyridine	HF
Q-142	pyridine	HCl
Q-143	pyridine	HBr
Q-144	pyridine	HI
Q-145	pyridine	p-toluene sulfonic acid
Q-146	pyridine	benzene sulfonic acid
Q-147	pyridine	xylene sulfonic acid
Q-148	pyridine	methane sulfonic acid
Q-149	pyridine	trifluoromethyl sulfonic acid
Q-150	pyridine	mesitylene sulfonic acid
Q-151	DMAP	HF
Q-152	DMAP	HCl
Q-153	DMAP	HBr
Q-154	DMAP	HI
Q-155	DMAP	p-toluene sulfonic acid
Q-156	DMAP	benzene sulfonic acid
Q-157	DMAP	xylene sulfonic acid
Q-158	DMAP	methane sulfonic acid

No.	amine	Acid
Q-159	DMAP	trifluoromethyl sulfonic acid
Q-160	DMAP	mesitylene sulfonic acid
Q-161	collidine	HF
Q-162	collidine	HCl
Q-163	collidine	HBr
Q-164	collidine	HI
Q-165	collidine	p-toluene sulfonic acid
Q-166	collidine	benzene sulfonic acid
Q-167	collidine	xylene sulfonic acid
Q-168	collidine	methane sulfonic acid
Q-169	collidine	trifluoromethyl sulfonic acid
Q-170	collidine	mesitylene sulfonic acid
Q-171	lutidine	HF
Q-172	lutidine	HCl
Q-173	lutidine	HBr
Q-174	lutidine	HI
Q-175	lutidine	p-toluene sulfonic acid
Q-176	lutidine	benzene sulfonic acid
Q-177	lutidine	xylene sulfonic acid
Q-178	lutidine	methane sulfonic acid
Q-179	lutidine	trifluoromethyl sulfonic acid
Q-180	lutidine	mesitylene sulfonic acid
Q-181	pyrimidine	HF
Q-182	pyrimidine	HCl
Q-183	pyrimidine	HBr
Q-184	pyrimidine	HI
Q-185	pyrimidine	p-toluene sulfonic acid
Q-186	pyrimidine	benzene sulfonic acid
Q-187	pyrimidine	xylene sulfonic acid
Q-188	pyrimidine	methane sulfonic acid
Q-189	pyrimidine	trifluoromethyl sulfonic acid
Q-190	pyrimidine	mesitylene sulfonic acid
Q-191	pyrazine	HF
Q-192	pyrazine	HCl
Q-193	pyrazine	HBr
Q-194	pyrazine	HI
Q-195	pyrazine	p-toluene sulfonic acid
Q-196	pyrazine	benzene sulfonic acid

No.	amine	Acid
Q-197	pyrazine	xylene sulfonic acid
Q-198	pyrazine	methane sulfonic acid
Q-199	pyrazine	trifluoromethyl sulfonic acid
Q-200	pyrazine	mesitylene sulfonic acid
Q-201	piperazine	HF
Q-202	piperazine	HCl
Q-203	piperazine	HBr
Q-204	piperazine	HI
Q-205	piperazine	p-toluene sulfonic acid
Q-206	piperazine	benzene sulfonic acid
Q-207	piperazine	xylene sulfonic acid
Q-208	piperazine	methane sulfonic acid
Q-209	piperazine	trifluoromethyl sulfonic acid
Q-210	piperazine	mesitylene sulfonic acid
Q-211	$N(CH_3)_2CH_2CH_3$	HF
Q-212	$N(CH_3)_2CH_2CH_3$	HCl
Q-213	$N(CH_3)_2CH_2CH_3$	HBr
Q-214	$N(CH_3)_2CH_2CH_3$	HI
Q-215	$N(CH_3)_2CH_2CH_3$	p-toluene sulfonic acid
Q-216	$N(CH_3)_2CH_2CH_3$	benzene sulfonic acid
Q-217	$N(CH_3)_2CH_2CH_3$	xylene sulfonic acid
Q-218	$N(CH_3)_2CH_2CH_3$	methane sulfonic acid
Q-219	$N(CH_3)_2CH_2CH_3$	trifluoromethyl sulfonic acid
Q-220	$N(CH_3)_2CH_2CH_3$	mesitylene sulfonic acid
Q-221	$N(CH_3)_2CH_2CH_2CH_3$	HF
Q-222	$N(CH_3)_2CH_2CH_2CH_3$	HCl
Q-223	$N(CH_3)_2CH_2CH_2CH_3$	HBr
Q-224	$N(CH_3)_2CH_2CH_2CH_3$	HI
Q-225	$N(CH_3)_2CH_2CH_2CH_3$	p-toluene sulfonic acid
Q-226	$N(CH_3)_2CH_2CH_2CH_3$	benzene sulfonic acid
Q-227	$N(CH_3)_2CH_2CH_2CH_3$	xylene sulfonic acid
Q-228	$N(CH_3)_2CH_2CH_2CH_3$	methane sulfonic acid
Q-229	$N(CH_3)_2CH_2CH_2CH_3$	trifluoromethyl sulfonic acid
Q-230	$N(CH_3)_2CH_2CH_2CH_3$	mesitylene sulfonic acid
Q-231	$NCH_3(CH_2CH_3)_2$	HF
Q-232	$NCH_3(CH_2CH_3)_2$	HCl
Q-233	$NCH_3(CH_2CH_3)_2$	HBr
Q-234	$NCH_3(CH_2CH_3)_2$	HI

No.	amine	Acid
Q-235	$\text{NCH}_3(\text{CH}_2\text{CH}_3)_2$	p-toluene sulfonic acid
Q-236	$\text{NCH}_3(\text{CH}_2\text{CH}_3)_2$	benzene sulfonic acid
Q-237	$\text{NCH}_3(\text{CH}_2\text{CH}_3)_2$	xylene sulfonic acid
Q-238	$\text{NCH}_3(\text{CH}_2\text{CH}_3)_2$	methane sulfonic acid
Q-239	$\text{NCH}_3(\text{CH}_2\text{CH}_3)_2$	trifluoromethyl sulfonic acid
Q-240	$\text{NCH}_3(\text{CH}_2\text{CH}_3)_2$	mesitylene sulfonic acid

Amine acid complexes wherein the acid is hydrochloric acid or hydrobromic acid, especially hydrochloric acid, are preferred.

- 5 Also, amine acid complexes wherein the acid is p-toluene sulfonic acid, benzene sulfonic acid, or xylene sulfonic acid are preferred.

10 Amine acid complexes Q-2, Q-5, Q-12, Q-17, Q-22, Q-32, Q-42, Q-52, Q-142, Q-145, Q-147, Q-162, Q-165, Q-172, Q-175, Q-212, Q-215, Q-222, Q-225, Q-232, and Q-235 are preferred.

15 Even more preferred are amine acid complexes Q-2, Q-5, Q-6, Q-7, Q-12, Q-82, Q-85, Q-86, Q-87, Q-102, Q-105, Q-106, Q-107, Q-112, Q-115, Q-116, Q-117, Q-122, Q-125, Q-126, Q-127, Q-132, Q-135, Q-136, Q-137, Q-142, Q-145, Q-146, Q-147, Q-212, Q-215, Q-216, Q-217, Q-222, Q-225, Q-226, Q-227, Q-232, Q-235, Q-236, and Q-237.

20 Most preferred are amine acid complexes Q-2, Q-5, Q-6, Q-7, Q-12, Q-142, Q-145, Q-146, Q-147, Q-212, Q-215, Q-216, Q-217, Q-222, Q-225, Q-226, Q-227, Q-232, Q-235, Q-236, and Q-237.

With regard to their use in the inventive process, the combinations of sulfinylating agent and amine acid complex given in the tables below are especially preferred.

Table 2

25 Trifluoromethylsulfinate sodium salt is used as the sulfinylating agent, and the amine acid complex in each case is a row of table 1.

Table 3

30 Trifluoromethylsulfinate potassium salt is used as the sulfinylating agent, and the amine acid complex in each case is a row of table 1.

Table 4

Trifluoromethylsulfinic acid is used as the sulfinylating agent, and the amine acid complex in each case is a row of table 1.

5 Table 5

Trifluoromethylsulfinic acid anhydride is used as the sulfinylating agent, and the amine acid complex in each case is a row of table 1.

Table 6

10 A mixture of trifluoromethylsulfinate sodium and potassium salt in a mixing ratio of from 0,01 : 99,99 weight % to 50 : 50 weight % is used as the sulfinylating agent, and the amine acid complex in each case is a row of table 1.

Also, in a further embodiment of the present invention, a Lewis acid such as AlCl_3 ,
15 FeCl_3 , CaCl_2 , ZnCl_2 , BF_3 , TiCl_4 , or ZrCl_4 can be used in exchange for the protonic acids cited above.

It can be advantageous to add the amine acid complex in two portions, one portion for step 1 and one portion after the addition of the pyrazole of formula II.

20

It can be advantageous to use two different amine acid complexes during the course of the reaction. For example, the first amine acid complex can be added in step 1 in amounts of 0,2 to 1 molar equivalents relative to pyrazole II, catalyzing the activation of the sulfinylate with the halogenating agent. After addition of the pyrazole of formula II,
25 in step 2, the Thia-Fries rearrangement, a second amine acid complex different from the first one is added, in amounts of 0,2 to 1 molar equivalents relative to pyrazole II.

Preferably, 1.4 to 2.2 molar equivalents, most preferably 1.5 to 1.8 molar equivalents, of the amine acid complex according to the present invention relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile are used.

30

When the sulfinylating agent is trifluoromethylsulfinic acid or a mixture containing trifluoromethylsulfinic acid, the molar amount of amine acid complex which is molar equivalent to the molar amount of trifluoromethylsulfinic acid is preferably generated in situ by addition of amine, and the remaining molar amount necessary to obtain the required 1.4 to 2.2 molar equivalents is added as amine acid complex.

35

In a preferred embodiment, the amine acid complex is dried before its use until it is essentially water-free. "Water free" means that the content of water in the solid does not exceed 5 ppm to 100 ppm.

40

Further additives can advantageously be added to the reaction mixture, such as potassium fluoride, pentafluorophenol, dimethylformamide, or 2,4-dinitrophenol. These additives are preferably added to the reaction mixture or solution or suspension of starting materials, respectively, before or at the reaction start. Most preferably, the additives are added at a low temperature of 5°C to 10°C.

In a preferred embodiment, 0.1 to 1.5 molar equivalents of potassium fluoride relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile are added to the reaction mixture or solution or suspension of starting materials, respectively, at 5°C to 10°C at or before the reaction start.

It is advantageous to add pentafluorophenol, dimethylformamide, or 2,4-dinitrophenol in catalytic amounts or in 0.10 molar equivalents relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile.

In a preferred embodiment, the additive is dried before its use until it is essentially water-free. "Water free" means that the content of water in the solid does not exceed 5 ppm to 100 ppm.

The reaction can be conducted in an inert organic solvent, preferably selected from

- aliphatic, alicyclic or aromatic, optionally halogenated hydrocarbons such as aromatic organic hydrocarbons, e.g. toluene, xylene, trifluoromethylbenzene, benzene, nitrobenzene, monochlorobenzene, dichlorobenzene, and ethylbenzene, preferably toluene and xylene, most preferably toluene; or aliphatic or alicyclic, optionally halogenated hydrocarbons such as hexane, cyclohexane, benzene, 1,2 dichloroethane, dichloromethane, trichloromethane (chloroform), tetrachlorocarbon, preferably 1,2 dichloroethane, dichloromethane, trichloromethane; and
- ethers, e.g. diethylether, dioxane, tetrahydrofuran, 2-methyl-tetrahydrofuran or ethylene glycol dimethyl- or diethylether; and
- ketons, e.g. acetone or butanon ; and
- nitriles, e.g. acetonitrile or propionitrile ; and
- amides, e.g. dimethylformamid, DMI (1,3-Dimethyl-2-Imidazolidinon), dimethylacetamid, N-methylformanilid, N-methylpyrrolidon or hexamethylphosphoric acid triamide; and
- sulfoxides, e.g. dimethylsulfoxide.

In a preferred embodiment, solvents which are essentially water free are used. "Water free" means that the content of water in the solvent does not exceed 5 ppm to 100 ppm. The most preferred solvent is water free toluene.

The reaction is carried out under an inert gas atmosphere, such as under an argon or a nitrogen atmosphere.

5 In a preferred embodiment, a total of 3.0 to 8.0 molar equivalents, more preferably 4.0 to 7.5 molar equivalents, and most preferably 4.5 to 6.5 molar equivalents of the solvent relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile are used. This relatively high concentration of starting materials maximizes the conversion to the sulfinamid intermediate.

10 In cases where the starting materials are dissolved and/or suspended, respectively, separately before their combination, about 25% to 40 % of the solvent are employed for dissolving and/ or suspending the 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile.

15 Generally, the sequence of addition of the starting materials 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile, amine acid complex, the sulfinylating agent and the chlorinating agent generally can be freely chosen.

20 Preferably, the respective starting materials are dissolved or suspended, respectively, in the reaction solvent before addition to the reaction mixture.

The halogenating agent is preferably not added to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile in the absence of the amine acid complex or the sulfinylating agent in the reaction mixture. In a preferred embodiment,
25 the halogenating agent is dissolved in the solvent and added to a reaction mixture containing 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile, amine acid complex and the sulfinylating agent, all dissolved or suspended, respectively, in the solvent.

30 In a preferred embodiment, 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile (II) is combined with a mixture containing the sulfinylating agent, the amine acid complex and the halogenating agent. In this case, it can be advantageous to include a first portion (equaling about 1 molar equivalent relative to compound II) of halogenating agent in the mixture containing the sulfinylating agent, the amine acid complex and the halogenating agent and then to add the second portion (equaling
35 about 0,1 to 0,2 molar equivalents relative to compound II) after addition of the compound II and stirring for approximately 30 to 60 minutes and shortly before rising the temperature to 30 to 50°C.

40 When the sulfinylating agent is trifluoromethylsulfinic acid, it is preferred to simultaneously add the trifluoromethylsulfinic acid and the halogenating agent to a solution or

suspension of the amine acid complex, followed by addition of a solution of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile to the reaction mixture.

5 In another preferred embodiment, a dissolved or suspended mixture of the sulfinylating agent, the amine acid complex and the halogenating agent in the solvent (preferably toluene) is cooled to about 3°C to 10°C, and a solution of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile in the solvent (preferably toluene) which has been heated to about 90°C to 110°C is combined with the cooled mixture.

10

In a preferred embodiment, after combination of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile, the sulfinylating agent, the amine acid complex, and the chlorinating agent, the temperature is raised to 30°C to 55 °C within 5 to 60 minutes.

15

It is also preferred to hold the reaction temperature initially at -20°C to 10°C for 5 to 60 minutes, preferably 20 to 40 minutes, followed by rising the temperature to 30°C to 55 °C at a rate of 5°C / min to 45°C / min. Preferably, in order to obtain high purity products, the reaction mixture is raised to a temperature not above 35°C. When the

20 sulfinylating agent is or contains CF₃S(O)OH, the initial reaction temperature preferably is -20°C to 5°C, in case of trifluoromethylsulfinate alkaline or alkaline earth metal salts, the initial reaction temperature preferably is -5°C to 10°C.

25

The reaction time depends upon the reaction temperature, the temperature control during the process, and the different reagents and solvents. The skilled artisan will be able to determine the appropriate reaction time in order to achieve the desired yield and purity. Typically, the reaction time will be about 5 to 15 hours, preferably 10 to 15 hours.

30

In a further preferred embodiment, the reaction is carried out in a pressure vessel at a pressure of 1, 013 bar (1 atm) to about 4 bar.

35

After completion of the reaction, fipronil can be isolated by employing conventional methods such as quenching the reaction with hydrogen carbonates, such as NaHCO₃, carbonates such as NaCO₃, or hydroxides, such as NaOH, extracting fipronil with an unpolar organic solvent such as ethylacetate or methyl-tert.-butylether, washing the extract, e.g. with hydrogen carbonates such as NaHCO₃, concentrating the extract, e.g. in vacuo, crystallization of fipronil, and the like. The isolated fipronil can be purified by a technique such as chromatography, recrystallization and the like, if necessary.

40

The crystallization of the final product 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile is typically conducted from a solution in a nonpolar, inert, preferably aromatic solvent with nonre-

5 particularly from a solution in benzene, ethylbenzene, monochlorobenzene, monofluorobenzene, 1,2-dichlorobenzene, 1,3-dichlorobenzene, 1,4-dichlorobenzene, toluene, o-xylene, m-xylene, p-xylene, styrene, i-propyl benzene, n-propyl benzene, 2-chlorotoluene, 3-chlorotoluene, 4-chlorotoluene, tert.-butyl benzene, sec.-butyl benzene, iso-butyl benzene, n-butyl benzene, 1,3-diisopropylbenzene, 1,4-

10 diisopropylbenzene, 2-nitrotoluene, 3-nitrotoluene, 4-nitrotoluene, nitrobenzene, benzonitrile, mesitylene, trifluoromethyl benzene, 1,2-dichloroethane, acetonitrile, dimethylsulfoxide, tetrahydrofuran, acetone, alcohols such as methanol, ethanol, n-propanol, isopropanol, n-butanol, isobutanol, 2-butanol, or tert-butanol, or mixtures thereof, preferably from a solution in monochlorobenzene, dichlorobenzene, ethylben-

15 zene or toluene.

Preferably, the crystallization is done from monochlorobenzene.

20 Preferably, the crystallization is done from dichlorobenzene.

Preferably, the crystallization is done from ethylbenzene.

Preferably, the crystallization is done from toluene.

25 It can be advantageous to add about 1 to 30 percent of a polar solvent such as ketons, amides, alcohols, esters or ethers, preferably esters, ketons or ethers, such as acetone methyl ethyl ketone, pentan-2-one, diethylketone, 4-methyl-2-pentanone, 3-methylbutan-2-one,

30 tert-butyl-methyl-ketone, cyclohexanone, methylacetate, ethylacetate, isopropylacetate, N-butylacetate, isobutylacetate, diethylcarbonate, 2-butoxyethylacetate, dimethylformamide, dimethylacetamide, dimethylsulfoxide, nitromethane, nitroethane, water, ethanol, methanol, propan-1-ol, propan-2-ol, butan-1-ol, butan-2-ol, tert-butanol, 2-methyl-propan-1-ol, 2-methyl-propan-2-ol, pentan-3-ol, 2-methyl butan-1-ol, 3-methyl butan-1-ol, 1,2-ethanediol, 1,3-propandiol, 1,2-propandiol, cyclohexanol, dioxane, tet-


35 rahydrofurane, diethylether, methyl tert.-butyl ether, 2-methyl tetrahydrofuran, acetonitrile, propionitrile, or mixtures thereof.

In another embodiment, fipronil is crystallized from water, optionally with the addition of about 1 to 30 percent of a polar organic solvent.

40

Purification of the crude product can also be achieved via filtration over charcoal or silica or washing with water.

- 5 When obtained according to the inventive process, the obtained product fipronil in the crude reaction mixture before crystallization contains less than 3,0 weight %, calculated without solvent, of compound F, a typical, biologically active side product of fipronil syntheses.



Compound F

- 10 After purification of the crude product by suitable methods such as washing and (re-) crystallizing, fipronil obtained by the inventive process contains less than 1.0 weight% of compound F.

- 15 Besides, the obtained product fipronil contains less than 10 ppm of compound D which is a common side product of the currently large scale industrial process as described e.g. in WO 01/30760, even after purification . Fipronil when prepared by the inventive process in an inert atmosphere contains less than 200 ppm of compounds containing sulfur in its oxidation state (IV). It also is free of compound E which typically can appear as side product of the current industrial process.

- 20 Furthermore, the obtained product fipronil is also free of trifluoroacetic acid, which is a reagent used in the current industrial process.

- 25 Moreover, when a chlorinating agent is used as the halogenating agent, the obtained fipronil product is practically free of bromine, meaning that it does not contain more than 5 to 20 ppm of bromine.

Examples

- 30 HPLCs were taken on a Hewlett Packard HP 1200, Chemstation, equipped with a J'Sphere ODS-H80, 4 μ m, 4.6 x 250 mm (YMC) column, eluent A: 90 wt.-% water + 10 wt.-% acetonitrile, eluent B: 10 wt.-% water + 90 wt.-% acetonitrile, flow: 0.85 ml/min, detection: 235 nm,

gradient:	time [min]	0	2	17	25	35
	A [%]	60	60	25	0	0
	B [%]	40	40	75	100	100.

- 5 Yields given below are in mol percent of the obtained purified crystalline product after work-up. Purity is given in weight percent of the obtained solid.

10 Example 1: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with triethylamine hydrochloride, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene

15 Within a 3-neck, 50 mL round bottom flask equipped with a magnetic stirrer bar and a thermometer were placed vacuum dried sodium trifluoromethylsulfinate (4.29 g, 27.5 mmol), vacuum dried triethylamine hydrochloride (5.16 g, 37.5 mmol), and 13 mL anhydrous toluene (6.5 molar equivalents relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile) under an argon atmosphere. After cooling to 0 °C to 5 °C with an ice bath, thionylchloride (3.57 g, 30 mmol) was added slowly while keeping the reaction temperature below 5 °C. After stirring for another 30 min, vacuum dried 5-amino-1-(2,6-dichloro-4-trifluoromethyl-phenyl)-1H-pyrazole-3-carbonitrile (8.03 g, 25 mmol, 99 % purity) was added at 5 °C, and the reaction mixture was heated to 50 °C within 5 min by a preheated water bath. The temperature of 50 °C was kept for another 6 hours before quenching the reaction with 50 mL of saturated NaHCO₃ solution.

25 The resulting suspension was diluted with 30 mL of ethylacetate. After phase separation the organic layer was washed once with saturated NaHCO₃ solution and concentrated under reduced pressure until dryness. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (8.06 g, 70 % yield, 94 % purity by quantitative HPLC). ¹H-NMR (Bruker DRX-500, d⁶-DMSO): δ [ppm]: 8.33 (s), 7.57 (s).

30 Example 2: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with triethylamine hydrochloride, potassium trifluoromethylsulfinate and thionylchloride

35 Within a 3-neck, 50 mL round bottom flask equipped with a magnetic stirrer bar and a thermometer were placed vacuum dried potassium trifluoromethylsulfinate (4.73 g, 27.5 mmol), vacuum dried triethylamine hydrochloride (5.16 g, 37.5 mmol), thionylchloride (3.57 g, 30 mmol), and 15 mL anhydrous toluene under an argon atmosphere. The reaction is further conducted as described above for example 1.

The crude product was crystallized from refluxing chlorobenzene (100 g) affording the title compound as a white crystalline powder (7.44 g, 66 % yield, 96 % purity by quantitative HPLC). ¹H-NMR (Bruker DRX-500, d⁶-DMSO): δ [ppm]: 8.33 (s), 7.57 (s).

5 Example 3: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile via addition of a mixture of sodium trifluoromethylsulfinate, triethylamine hydrochloride and thionylchloride to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile

10 Within a 3-neck, 50 mL round bottom flask equipped with a magnetic stirrer bar and a thermometer were placed vacuum dried sodium trifluoromethylsulfinate (4.29 g, 27.5 mmol), vacuum dried triethylamine hydrochloride (5.16 g, 37.5 mmol), and 10 g anhydrous toluene under an argon atmosphere. After cooling to 0 - 5 °C with an ice bath thionylchloride (3.57 g, 30 mmol) was added while keeping the reaction temperature
15 below 5 °C. After stirring for another 30 min the cooled sulfinic acid solution was added at once to a stirred suspension of vacuum dried 5-amino-1-(2,6-dichloro-4-trifluoromethyl-phenyl)-1H-pyrazole-3-carbonitrile (8.03 g, 25 mmol, 99 % purity) in 5 g toluene with a temperature of 50 °C. The temperature of 50 °C was kept for another 5 hours before quenching the reaction with 50 mL of saturated NaHCO₃ solution.

20 The resulting suspension was diluted with 30 mL of ethylacetate. After phase separation the organic layer was washed once with saturated NaHCO₃ solution and concentrated under reduced pressure until dryness. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (7.25 g, 63 % yield, 94 % purity by quantitative HPLC). ¹H-NMR (Bruker DRX-500, d⁶-
25 DMSO): δ [ppm]: 8.33 (s), 7.57 (s).

Example 4: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with triethylamine hydrochloride, trifluoromethylsulfonic acid and thionylchloride

30 Within a 3-neck, 50 mL round bottom flask equipped with a magnetic stirrer bar and a thermometer were placed trifluoromethylsulfonic acid (3.69 g, 27.5 mmol), anhydrous triethylamine (2.78 g, 27.5 mmol), and 10 g anhydrous toluene under an argon atmosphere. After cooling to 0 - 5 °C with an ice bath thionylchloride (3.57 g, 30 mmol) was
35 added while keeping the reaction temperature below 5 °C. After stirring for another 30 min, the cooled sulfinic acid solution was added at once to a stirred suspension of vacuum dried 5-amino-1-(2,6-dichloro-4-trifluoromethyl-phenyl)-1H-pyrazole-3-carbonitrile (8.03 g, 25 mmol, 99 % purity) and vacuum dried triethylamine hydrochloride (1.38 g, 10.0 mmol) in 5 g anhydrous toluene in a second 50 mL round bottom flask equipped
40 with a magnetic stirrer bar and a thermometer having a temperature of 50 °C inside.

The temperature of 50 °C was kept for another 6 hours before quenching the reaction with 50 mL of saturated NaHCO₃ solution.

The resulting suspension was diluted with 30 mL of ethylacetate. After phase separation the organic layer was washed once with saturated NaHCO₃ solution and concentrated under reduced pressure until dryness. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder. ¹H-NMR (Bruker DRX-500, d⁶-DMSO): δ [ppm]: 8.33 (s), 7.57 (s).

Example 5: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with triethylamine hydrochloride, sodium trifluoromethylsulfinate and thionylchloride under addition of potassium fluoride

An oven dried 100 mL 3 neck round bottom flask equipped with a magnetic stir bar, thermocouple, condenser, N₂ inlet, and rubber septum, was charged with vacuum dried potassium fluoride (1.53 g, 26.1 mmol), anhydrous toluene (20.1 g), vacuum dried sodium trifluoromethylsulfinate (4.53 g, 29.0 mmol), and thionylchloride (3.76 g, 31.6 mmol) under nitrogen. The solution was cooled to 0°C and triethylamine hydrochloride (5.44g, 39.5 mmol) was added slowly, controlling the temperature to less than 10 °C. Vacuum dried 5-amino-1-(2,6-dichloro-4-trifluoromethyl-phenyl)-1H-pyrazole-3-carbonitrile (8.48 g, 26.1 mmol, 99 % pure) was added at 0°C. The solution was then quickly warmed to 50 °C, in less than 5 minutes, using a hot water bath. The reaction was allowed to stir for 5 hrs at 50 °C before quenching with 50 ml of saturated aqueous NaHCO₃. The resulting suspension was diluted with 30 mL of ethylacetate and then allowed to phase separate. The aqueous phase was extracted with 30 mL ethyl acetate and the combined organic phases were washed with 30 ml of saturated aqueous NaCO₃. The organic phase was concentrated under reduced pressure until dryness, affording the crude title compound (11.6 g, 67 % yield, 68.5 % purity). ¹H-NMR (Bruker DRX-500, d⁶-DMSO): δ [ppm]: 8.33 (s), 7.57 (s).

Example 6: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with triethylamine hydrochloride, sodium trifluoromethylsulfinate and phosphoroychloride

Within a 3-neck, 50 mL round bottom flask equipped with a magnetic stirrer bar and a thermometer were placed sodium trifluoromethylsulfinate (8.84 g, 55.0 mmol) and 40 mL dried toluene (6.8 molar equivalents relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile) under a nitrogen atmosphere. After cooling to 0°C to 5 °C with an ice bath, phosphoroychloride (9.20 g, 60.0 mmol) was added slowly while keeping the reaction temperature below 5 °C. After complete addition of phosphoroychloride, triethylamine hydrochloride (10.32g, 75.0 mmol) was added at 5°C. After stirring for another 30 min, vacuum dried 5-amino-1-(2,6-dichloro-4-

trifluoromethyl-phenyl)-1H-pyrazole-3-carbonitrile (16.06 g, 50 mmol, 99 % purity) was added at 5 °C, and the reaction mixture was heated to 50 °C within 5 min by a pre-heated water bath. The temperature of 50 °C was kept for another 6 hours before quenching the reaction with 100 g of 10% NaHCO₃ solution. The resulting suspension
5 was diluted with 100 mL of ethylacetate. After phase separation, the organic layer was analyzed by quantitative HPLC (57.3% yield).

Example 7: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with pyridine tosylate, sodium trifluoromethylsulfinate and
10 thionylchloride, in 6.5 molar equivalents of toluene

The preparation procedure was conducted as described above for example 1. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (75 % yield, 94 % purity by quantitative HPLC).
15

Example 8: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with trimethylamine hydrochloride, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene

The preparation procedure was conducted as described above for example 1. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (71 % yield, 97 % purity by quantitative HPLC).
20

Example 9: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with pyridine hydrochloride, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene
25

The preparation procedure was conducted as described above for example 1. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (67 % yield, 95 % purity by quantitative HPLC).
30

Example 10: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with triethylamine hydrochloride, potassium trifluoromethylsulfinate and thionylchloride, in 4.5 molar equivalents of toluene
35

Within a 3-neck, 50 mL round bottom flask equipped with a magnetic stirrer bar and a thermometer were placed vacuum dried potassium trifluoromethylsulfinate (5.16 g, 30 mmol), vacuum dried triethylamine hydrochloride (5.16 g, 37.5 mmol), and 10.4 g anhydrous toluene (4.5 molar equivalents relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile) under an argon atmosphere. After cooling to 0 °C to 5 °C with external cooling, thionylchloride (3.57 g, 30 mmol) was
40

added within 15 min while keeping the reaction temperature below 5 °C. After stirring for another 30 min, vacuum dried 5-amino-1-(2,6-dichloro-4-trifluoromethyl-phenyl)-1H-pyrazole-3-carbonitrile (8.03 g, 25 mmol, 99 % purity) was added at 5 °C, and the reaction mixture was kept at 5 °C for 60 min and then heated to 35 °C within 45 min. The temperature of 35 °C was kept for 3 hours before adding 4.6 g of toluene. After another 7 hours at 35 °C the reaction was quenched with 20 g of sodium hydroxide solution (10 wt.%).

The resulting suspension was diluted with 30 mL of ethylacetate. After phase separation the organic layer was washed once with sodium hydroxide solution (10 wt.%). After phase separation, the organic layer was analyzed by quantitative HPLC (80.4 % yield). The crude product was crystallized from a mixture of ethylacetate and toluene affording the title compound as a white crystalline powder (7.98 g, 73 % yield, 98 % purity by quantitative HPLC).

Example 11: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with thionylchloride, triethylamine hydrochloride and dosage of potassium trifluoromethylsulfinate

Within a 750 mL reactor with a mechanical stirrer and a thermometer were placed vacuum dried triethylamine hydrochloride (51.1 g, 368 mmol), 147 g anhydrous toluene (6.5 molar equivalents relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile), and thionylchloride (35.7 g, 294 mmol) under an argon atmosphere. After cooling to 0°C to 5 °C with external cooling, vacuum dried potassium trifluoromethylsulfinate (50.4 g, 296 mmol) was added in three equal portions every 10 min while keeping the reaction temperature below 5 °C. After stirring for another 30 min, vacuum dried 5-amino-1-(2,6-dichloro-4-trifluoromethyl-phenyl)-1H-pyrazole-3-carbonitrile (79.5 g, 245 mmol, 99 % purity) was added at 5 °C, and the reaction mixture was kept at 5 °C for 60 min and then heated to 35 °C within 45 min. The temperature of 35 °C was kept for another 10 hours before quenching the reaction with 200 g of sodium hydroxide solution (10 wt.%).

The resulting suspension was diluted with 176 mL of ethylacetate. After phase separation the organic layer was washed once with sodium hydroxide solution (10 wt.%). After phase separation, the organic layer was analyzed by quantitative HPLC (79 % yield). The content of compound F was below 2.9 weight percent in the crude mixture (without solvent). The product was crystallized from a mixture of ethylacetate and toluene affording the title compound as a white crystalline powder (77.1 g, 75 % yield, 98 % purity by quantitative HPLC).

Example 12:

Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with trimethylamine tosylate, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene

5

The preparation procedure was conducted as described above for example 1. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (76 % yield, 96 % purity by quantitative HPLC).

10 Example 13:

Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with pyridine hydrochloride, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene

15 The preparation procedure was conducted as described above for example 1. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (67 % yield, 95 % purity by quantitative HPLC).

20 Example 14: Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with trimethylamine tosylate, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene

The preparation procedure was conducted as described above for example 11. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (73 % yield, 98 % purity by quantitative HPLC).

25 Example 15:

Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with triethylamine hydrochloride, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene

30

The preparation procedure was conducted as described above for example 11. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (75 % yield, 98 % purity by quantitative HPLC).

35

Example 16:

Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with pyridine tosylate, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene

40

The preparation procedure was conducted as described above for example 11. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (72 % yield, 98 % purity by quantitative HPLC).

5 Example 17:

Sulfinylation of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile with triethylamine tosylate, sodium trifluoromethylsulfinate and thionylchloride, in 6.5 molar equivalents of toluene

- 10 The preparation procedure was conducted as described above for example 1. The crude product was crystallized from refluxing toluene (100 g) affording the title compound as a white crystalline powder (44 % yield, 76 % purity by quantitative HPLC). No formation of insoluble material (as with diethylamine tosylate as the amine acid complex, compare example C4 of Table 3) was observed.

15

Comparative Examples

In order to demonstrate the advantages of the inventive process, the following examples are conducted employing the preparation procedure given above for example 1.

20

Table 2

No.	Sulfinylating agent	Amine acide complex	Chlorinating agent
C1	CF ₃ S(O)ONa	dimethylamine hydrochloride	SOCl ₂
C2	CF ₃ S(O)ONa	diethylamine hydrochloride	SOCl ₂
C3	CF ₃ S(O)ONa	diethylamine tosylate	SOCl ₂

- 25 In these experiments, it is demonstrated that the inventive process gives higher yields and/or higher purities as compared to the sulfinylation processes described in the prior art. The results are summarized in Table 3 which follows.

Table 3

Example no.	Sulfinylating Agent	Amine Acid Complex	Chlorinating Agent	Yield Crystalline Fipronil [mol%]	Purity of Fipronil [weight%]	Comments
Exp.8	CF ₃ S(O)ONa	trimethylamine hydrochloride	SOCl ₂	71	97	
C1	CF ₃ S(O)ONa	dimethylamine hydrochloride	SOCl ₂	57	84	Disadvantageously, an insoluble material was formed that remained throughout the work-up and recrystallization
Exp.1	CF ₃ S(O)ONa	triethylamine hydrochloride	SOCl ₂	70	94	
C2	CF ₃ S(O)ONa	diethylamine hydrochloride	SOCl ₂	55	84	Disadvantageously, an insoluble material was formed that remained throughout the work-up and recrystallization
Exp. 12	CF ₃ S(O)ONa	trimethylamine tosylate	SOCl ₂	76	96	
C3	CF ₃ S(O)ONa	dimethylamine tosylate	SOCl ₂	73	97	
Exp. 17	CF ₃ S(O)ONa	triethylamine tosylate	SOCl ₂	44	76	
C4	CF ₃ S(O)ONa	diethylamine tosylate	SOCl ₂	65	95	Disadvantageously, an insoluble material was formed that remained throughout the work-up and recrystallization

Further comparative experiments are conducted employing the following preparation procedure:

To a 750 ml reactor with a mechanical stirrer and a thermometer containing the amine acid complex (1.5 eq.), 147 g anhydrous toluene (6.5 eq.), and 44.8 g CF_3SOCl (1.2 eq.) under an argon atmosphere at 0 °C was added vacuum dried 5-amino-1-(2,6-dichloro-4-trifluoromethyl-phenyl)-1H-pyrazole-3-carbonitrile (1 eq., 79.5 g, 245 mmol, 99 % purity). The reaction mixture was kept at 5 °C for 60 min and then heated to 35 °C within 45 min. The temperature of 35 °C was kept for another 10 hours before quenching the reaction with 200 g of sodium hydroxide solution (10 wt.%). The resulting suspension was diluted with 176 ml of ethylacetate. After phase separation the organic layer was washed once with sodium hydroxide solution (10 wt.%). After phase separation, the organic layer was analyzed by quantitative HPLC. The product was crystallized from a mixture of ethylacetate and toluene affording the title compound as a white crystalline powder.

No.	Sulfinylating agent	Amine acide complex	Chlorinating agent
Ca	$\text{CF}_3\text{S(O)Cl}$	pyridine hydrochloride	-
Cb	$\text{CF}_3\text{S(O)Cl}$	trimethylamine hydrochloride	-

These experiments Ca and Cb where compared to examples 18 and 19, which were conducted as described above for example 11, employing potassium trifluoromethylsulfinate / pyridine hydrochloride / SOCl_2 and potassium trifluoromethylsulfinate / trimethylamine hydrochloride / SOCl_2 as reagents.


In these experiments, it is demonstrated that the inventive process gives higher yields and/or higher purities as compared to the sulfinylation processes described in the prior art.

Claims:

1. Process for the sulfinylation of a pyrazole derivative, characterized in that 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile (II) is reacted with a sulfinylating agent selected from trifluoromethylsulfonic acid, trifluoromethylsulfonic acid anhydride, and a trifluoromethylsulfinate alkaline or alkaline earth metal salt and mixtures of the acid and/or the salt(s), in the presence of at least one amine acid complex wherein the amine(s) are selected from tertiary amines and the acid(s) are selected from hydrofluoric, hydrochloric, hydrobromic and hydroiodic acid and sulfonic acid derivatives, and with the addition of a halogenating agent.
2. Process according to claim 1 wherein the halogenating agent is selected from thionylchloride, thionylbromide, phosphoroxychloride, oxalylchloride, phosgen, triphosgen ((CCl₃)₂C(=O)), chloroformates, phosphorpentachloride, phosphortrichloride, trichloromethylchloromethanoat, and xylenesulfonic acid chloride.
3. Process according to claims 1 or 2 wherein the halogenating agent is a chlorinating agent selected from thionylchloride and phosphoroxychloride.
4. Process according to any of claims 1 to 3 wherein the amine of the amine acid complex is selected from the alkyl amines trimethylamine, triethylamine, tripropylamine, triisopropylamine, tributylamine, dimethyl ethyl amine, diethyl methylamine, dimethyl n-propyl amine, diisopropyl ethyl amine, DBU (1,8-diazabicyclo[5.4.0]undec-7-ene), DBN (1,5-diazabicyclo[4.3.0]non-5-ene), methyl morpholine, ethyl morpholine, N,N-dimethyl aniline, methyl piperidine, methyl pyrrolidine, methyl dibenzyl amine, and the aromatic amines pyridine, DMAP (dimethylamino pyridine), collidine, lutidine, pyrimidine, pyrazine, and piperazine.
5. Process according to any of claims 1 to 4 wherein the amine of the amine acid complex is selected from trimethylamine, triethylamine, dimethyl ethyl amine, dimethyl n-propyl amine or pyridine, preferably from trimethylamine, triethylamine and pyridine.
6. Process according to any of claims 1 to 5 wherein the acid of the amine acid complex is selected from hydrofluoric acid, hydrochloric acid, p-toluenesulfonic acid, benzenesulfonic acid, xylene sulfonic acid, methanesulfonic acid and trifluoromethylsulfonic acid, preferably from hydrochloric acid and p-toluenesulfonic acid.

7. Process according to any of claims 1 to 6 wherein the sulfinylating agent is selected from trifluoromethylsulfinic acid, trifluoromethylsulfinate sodium salt, trifluoromethylsulfinate potassium salt, and mixtures of these.
- 5 8. Process according to any of claims 1 to 7 wherein the reaction is conducted in an organic solvent selected from toluene, benzene, xylene, trifluoromethylbenzene, monochlorobenzene, dichlorobenzene, and ethylbenzene.
- 10 9. Process according to any of claims 1 to 8 wherein 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile is combined with a reaction mixture of the sulfinylating agent, the amine acid complex and the halogenating agent.
- 15 10. Process according to any of claims 1 to 9 wherein 1.4 to 2.2 molar equivalents of the amine acid complex relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile are used.
- 20 11. Process according to any of claims 1 to 10 wherein 1.15 to 1.35 molar equivalents of the halogenating agent relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile are used.
- 25 12. Process according to any of claims 1 to 11 wherein 1.0 to 1.35 molar equivalents of the sulfinylating agent relative to 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile are used.
- 30 13. Process according to any of claims 1 to 12 wherein, after combination of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile, the sulfinylating agent, amine acid complex, and the halogenating agent, the temperature is raised to 30°C to 55 °C, preferably 30°C to 39°C, within 5 to 60 minutes.
- 35 14. 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile, prepared by the process as defined in any of the claims 1 to 13.
15. 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile, prepared by a process as defined in any of the claims 3 to 13, which contains less than 20 ppm of bromine.

16. 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile, prepared by a process as defined in any of the claims 1 to 13, or according to claim 15, which contains less than 10 ppm of the compound D



- 5
17. 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile, prepared by a process as defined in any of the claims 1 to 13, or according to claims 15 or 16, and in an inert atmosphere, which contains less than 200 ppm of compounds containing sulfur in its oxidation state (IV).
- 10
18. Pesticidal or parasitocidal composition containing 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile as defined in any of the claims 14 to 17.
- 15
19. Use of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by a process as defined in any of the claims 1 to 13, or as defined in any of the claims 14 to 17 for combating pests.
- 20
20. Use of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by a process as defined in any of the claims 1 to 13, or as defined in any of the claims 14 to 17 for combating animal health pests and parasites.
- 25
21. A method for the control of insects, acarids or nematodes by contacting the insect, acarid or nematode or their food supply, habitat, breeding ground or their locus with a pesticidally effective amount of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by a process as defined in any of the claims 1 to 13, or as defined in any of the
- 30
- claims 14 to 17.
22. A method of protecting growing plants from attack or infestation by insects, acarids or nematodes by applying to the foliage or the seeds of the plants, or to the soil or water in which they are growing, a pesticidally effective amount of

5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by a process as defined in any of the claims 1 to 13, or as defined in any of the claims 14 to 17.

- 5 23. A method as claimed in claims 21 or 22, wherein 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile is applied in an amount of from 5 g/ha to 2000 g/ha.
- 10 24. A method for treating, controlling, preventing or protecting animals against infestation or infection by parasites which comprises orally, topically or parenterally administering or applying to the animals or their habitat a parasitically effective amount of 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by a process as defined in any of the claims 1 to 13, or as defined in any of the claims 14 to 17 or its veterinarily acceptable enantiomers or salts.
- 15 25. A process for the preparation of a composition for treating, controlling, preventing or protecting animals against infestation or infection by parasites which comprises admixing 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile prepared by a process as defined in any of the claims 1 to 13, or as defined in any of the claims 14 to 17, or its veterinarily acceptable enantiomers or salts with a veterinarily acceptable carrier.
- 20 26. The process according to claim 25 wherein 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)-pyrazole-3-carbonitrile or its veterinarily acceptable enantiomer or salt is present in a parasitically effective amount.
- 25

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2007/061891

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07D231/44 A01N43/56 A01N43/00 A61K31/415 A61P33/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 C07D A01N A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
 EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 295 117 A (MAY & BAKER LTD [GB] RHONE POULENC AGRICULTURE [GB]) 14 December 1988 (1988-12-14) page 3, compound 42;claims 11-13	14-26
X	US 5 618 945 A (CASADO MICHEL [FR] ET AL) 8 April 1997 (1997-04-08) column 4, lines 10-33; example 1 & EP 0 668 269 A (RHONE POULENC AGROCHIMIE [FR] AVENTIS CROPSCHIENCE SA [FR]) 23 August 1995 (1995-08-23) cited in the application	1-26
A	CN 1 374 298 A (JIANGSU PROV INST OF PESTICIDE [CN] INT EPODOC Caesar accession number) 16 October 2002 (2002-10-16) cited in the application the whole document	1-26

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

19 March 2008

Date of mailing of the international search report

31/03/2008

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

De Jong, Bart

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2007/061891

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 19-21, 23, 24 are (at least partially) directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2007/061891

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0295117	A	14-12-1988	AT 191479 T 15-04-2000
			AT 307118 T 15-11-2005
			AU 618266 B2 19-12-1991
			AU 1755488 A 15-12-1988
			BR 8803258 A 31-01-1989
			CA 1330089 C 07-06-1994
			CN 88103601 A 28-12-1988
			CZ 8804052 A3 16-12-1998
			DD 281744 B5 20-02-1997
			DD 281744 A5 22-08-1990
			DE 3856402 D1 11-05-2000
			DE 3856402 T2 19-10-2000
			DE 3856585 T2 24-08-2006
			DK 314088 A 13-12-1988
			EG 19113 A 30-11-1994
			ES 2144390 T3 16-06-2000
			ES 2251806 T3 01-05-2006
			FI 882735 A 13-12-1988
			GR 3033663 T3 31-10-2000
			HK 1005289 A1 09-02-2001
			HU 48875 A2 28-07-1989
			HU 9500470 A3 30-10-1995
			IE 20020481 A1 11-02-2004
			IL 86492 A 08-07-1993
			JP 2669538 B2 29-10-1997
			JP 63316771 A 26-12-1988
			MA 21292 A1 31-12-1988
			MX 11842 A 01-12-1993
			NL 350001 I1 01-12-2000
			NO 882551 A 13-12-1988
			NZ 224979 A 26-05-1992
			OA 8880 A 31-10-1989
			PH 26895 A 03-12-1992
			PL 272998 A1 06-03-1989
			PT 87697 A 31-05-1989
			RO 100612 B1 07-07-1992
			RO 106496 B1 31-05-1993
			SG 63529 A1 30-03-1999
			SK 405288 A3 06-05-1998
			RU 2051909 C1 10-01-1996
			TR 23696 A 15-06-1990
ZA 8804179 A 22-02-1989			
ZW 7388 A1 21-12-1988			
US 5618945	A	08-04-1997	AT 218127 T 15-06-2002
			AU 678876 B2 12-06-1997
			AU 1238395 A 31-08-1995
			BR 9500858 A 24-10-1995
			CA 2143135 A1 23-08-1995
			CN 1110682 A 25-10-1995
			CZ 9500461 A3 18-10-1995
			DE 69526815 D1 04-07-2002
			DE 69526815 T2 14-11-2002
			DK 668269 T3 17-06-2002
			EP 0668269 A1 23-08-1995
			ES 2173932 T3 01-11-2002
			FI 950800 A 23-08-1995
			FR 2716453 A1 25-08-1995

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2007/061891

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5618945 A		HU 69333 A2	28-09-1995
		IL 112708 A	17-05-2005
		JP 3963965 B2	22-08-2007
		JP 7278106 A	24-10-1995
		JP 2005187478 A	14-07-2005
		JP 2007153906 A	21-06-2007
		PL 307393 A1	04-09-1995
		PT 668269 T	30-09-2002
		RO 113639 B1	30-09-1998
		RU 2136665 C1	10-09-1999
		SK 23495 A3	13-09-1995
		ZA 9501468 A	16-02-1996
	<hr/>		
EP 0668269 A	23-08-1995	AT 218127 T	15-06-2002
		AU 678876 B2	12-06-1997
		AU 1238395 A	31-08-1995
		BR 9500858 A	24-10-1995
		CA 2143135 A1	23-08-1995
		CN 1110682 A	25-10-1995
		CZ 9500461 A3	18-10-1995
		DE 69526815 D1	04-07-2002
		DE 69526815 T2	14-11-2002
		DK 668269 T3	17-06-2002
		ES 2173932 T3	01-11-2002
		FI 950800 A	23-08-1995
		FR 2716453 A1	25-08-1995
		HU 69333 A2	28-09-1995
		IL 112708 A	17-05-2005
		JP 3963965 B2	22-08-2007
		JP 7278106 A	24-10-1995
		JP 2005187478 A	14-07-2005
		JP 2007153906 A	21-06-2007
		PL 307393 A1	04-09-1995
		PT 668269 T	30-09-2002
		RO 113639 B1	30-09-1998
		RU 2136665 C1	10-09-1999
SK 23495 A3	13-09-1995		
US 5618945 A	08-04-1997		
ZA 9501468 A	16-02-1996		
<hr/>			
CN 1374298 A	16-10-2002	NONE	