



(51) International Patent Classification:

A01N 43/90 (2006.01)

A01P 7/04 (2006.01)

A01P 1/00 (2006.01)

C07D 487/22 (2006.01)

A01P 3/00 (2006.01)

(21) International Application Number:

PCT/CA2020/050197

(22) International Filing Date:

14 February 2020 (14.02.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/806,084

15 February 2019 (15.02.2019)

US

(71) Applicant: SUNCOR ENERGY INC. [CA/CA]; Suncor Energy Centre, West Tower, PO Box 2844, 150 - 6th Avenue SW, Calgary, Alberta T2P 3E3 (CA).

(72) Inventors: FEFER, Michael; 2489 North Sheridan Way, Mississauga, Ontario L5K 1A8 (CA). LIU, Jun; 2489 North Sheridan Way, Mississauga, Ontario L5K 1A8 (CA). TERAZONO, Yuichi; 2489 North Sheridan Way, Mississauga, Ontario L5K 1A8 (CA). NG, Kenneth; 2489 North Sheridan Way, Mississauga, Ontario L5K 1A8 (CA). SHEN, Youqing; Apt 102, Bldg5-1, 655 XI XI Street, Hangzhou Shi, Zhejiang 310088 (CN).

(74) Agent: ROBIC; 630 Rene-Levesque Blvd. West, 20th Floor, Montreal, Québec H3B 1S6 (CA).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,


SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE

(54) Title: THE USE OF PROTOPORPHYRIN IX DERIVATIVES TO IMPROVE THE HEALTH OF PLANTS



(I)

(57) Abstract: Provided herein are compounds of general Formula (I) or agriculturally acceptable salts thereof. The compounds of Formula (I) can be used to improve the health of plants. For example, the compounds of Formula (I) can be used to inhibit a microbial pathogen of a plant, or to increase resistance of a plant to one or more abiotic stress.

RELATED APPLICATION

[001] This application claims priority to United-States provisional application No. 62/806,084 filed on February 15, 2019, the content of which is incorporated herein by reference in its entirety for all purposes.

FIELD


[002] The technical field generally relates to photosensitizer tetrapyrrole compounds and their use for promoting the health of plants. More particularly, the technical field relates to modified protoporphyrin IX (PP IX) compounds and use thereof for photodynamic inhibition of microbial pathogens, such as fungal or bacterial pathogens, in plants. The modified PP IX compounds can also be used for increasing abiotic stress resistance or tolerance in plants and/or as insecticides to protect plants from plant pests.

BACKGROUND

[003] Photodynamic inhibition of microbial pathogens involves exposing a photosensitive agent to light in order to generate reactive oxygen species (ROS), such as singlet oxygen, which can have detrimental effects on the microbial pathogens. Existing photodynamic inhibition techniques and applications have various shortcomings.

SUMMARY

[004] In one aspect of the present description, there is provided a compound of Formula I:



Formula I

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

Z^1 and Z^2 are each independently OR^1 or NR^2R^3 ;

each R^1 , R^2 and R^3 is independently H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, or substituted alkynyl, wherein:

if Z^1 and Z^2 are both OR^1 then at least one R^1 is not H,

if Z^1 and Z^2 are both NR^2R^3 then at least one R^3 is not H, and

if one of Z^1 and Z^2 is OR^1 and the other one of Z^1 and Z^2 is NR^2R^3 , then at least one of R^1 and R^3 is not H;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

----- is a single bond or a double bond;

===== is a single bond or a double bond; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted aryl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more -X, $-R^B$, $-O^-$, $=O$, $-OR^B$, $-SR^B$, $-S^-$, $-NR^B_2$, $Si(R^C)_3$, $-N^+R^B_3$, $-NR^B-(Alk)-NR^B_2$, $-NR^B-(Alk)-N^+R^B_3$, $-NR^B-(Alk)-OR^B$, $-NR^B-(Alk)-OP(=O)(OR^B)(O^-)$, $-NR^B-(Alk)-OP(=O)(OR^B)_2$, $-NR^B-(Alk)-Si(R^C)_3$, $-NR^B-(Alk)-SR^B$, $-O-(Alk)-NR^B_2$, $-O-(Alk)-N^+R^B_3$, $-O-(Alk)-OR^B$, $-O-(Alk)-OP(=O)(OR^B)(O^-)$, $-O-(Alk)-OP(=O)(OR^B)_2$, $-O-(Alk)-Si(R^C)_3$, $-O-(Alk)-SR^B$, $=NR^B$, $-CX_3$, $-CN$, $-OCN$, $-SCN$, $-N=C=O$, $-NCS$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-NHC(=O)R^B$, $-OC(=O)R^B$, $-NHC(=O)NR^B_2$, $-S(=O)_2-$, $-S(=O)_2OH$, $-S(=O)_2R^B$, $-OS(=O)_2OR^B$, $-S(=O)_2NR^B_2$, $-S(=O)R^B$, $-OP(=O)(OR^B)(O^-)$, $-OP(=O)(OR^B)_2$, $-P(=O)(OR^B)_2$, $-P(=O)(O^-)_2$, $-P(=O)(OH)_2$, $-P(O)(OR^B)(O^-)$, $-C(=O)R^B$, $-C(=O)X$, $-C(S)R^B$, $-C(O)OR^B$, $-C(O)O^-$, $-C(S)OR^B$, $-C(OSR^B)$, $-C(S)SR^B$, $-C(O)NR^B_2$, $-C(S)NR^B_2$ or $-C(=NR^B)NR^B_2$;

each X is independently a halogen: F, Cl, Br or I;


each R^B is independently H, alkyl, aryl, arylalkyl, a heterocycle, an alkyloxy group such as poly(ethyleneoxy), PEG or poly(methyleneoxy), a capped poly(ethyleneoxy), capped PEG or capped polymethyleneoxy, or a protecting group;

the capped poly(ethyleneoxy), capped PEG and capped poly(methyleneoxy) groups being each independently capped with alkyl, aryl, arylalkyl, alkenyl, alkynyl, CO(alkyl), CO(aryl), CO(arylalkyl), CO(alkenyl) or CO(alkynyl);

each R^C is independently alkyl, aryl, arylalkyl, O(alkyl), O(aryl), O(arylalkyl), or O(tri-substituted silyl);

each tri-substituted silyl is independently substituted with three functional groups selected from alkyl, alkenyl, alkynyl, aryl and arylalkyl; and each Alk is independently alkylene, alkenylene, or alkynylene.

[005] In another aspect of the present description, there is provided a compound of Formula I:



Formula I

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z¹ and Z² is OR¹; and

the other one of Z¹ and Z² is NR²R³, NR²-(CH₂)_n-NR⁴R⁵, NR²-(CH₂)_n-N⁺R⁴R⁵R⁶ Y⁻, NR²-(CH₂)_n-O(PO₃H)⁻ W⁺, NR²-(CH₂)_n-Si(R⁷)₃, NR²-(CH₂)_n-SR⁸, NR²-(CH₂)_n-NR⁴-(CH₂)_p-NR⁹R¹⁰, NR²-(CH₂)_n-NR⁴-(CH₂)_p-N⁺R⁹R¹⁰R¹¹ Y⁻, NR²-(CH₂)_n-NR⁴-(CH₂)_p-O(PO₃H)⁻ W⁺, NR²-(CH₂)_n-NR⁴-(CH₂)_p-Si(R⁷)₃, NR²-(CH₂)_n-NR⁴-(CH₂)_p-SR⁸, O(CH₂)_n-NR⁴R⁵, O(CH₂)_n-N⁺R⁴R⁵R⁶ Y⁻, O(CH₂)_n-O(PO₃H)⁻ W⁺, O(CH₂)_n-Si(R⁷)₃, O(CH₂)_n-SR⁸, O(CH₂)_n-NR⁴-(CH₂)_p-NR⁹R¹⁰, O(CH₂)_n-NR⁴-(CH₂)_p-N⁺R⁹R¹⁰R¹¹ Y⁻, O(CH₂)_n-NR⁴-(CH₂)_p-O(PO₃H)⁻ W⁺ or O(CH₂)_n-NR⁴-(CH₂)_p-Si(R⁷)₃;

or

Z¹ is NR²R³, NR²-(CH₂)_n-NR⁴R⁵, NR²-(CH₂)_n-N⁺R⁴R⁵R⁶ Y⁻, NR²-(CH₂)_n-O(PO₃H)⁻ W⁺, NR²-(CH₂)_n-Si(R⁷)₃, NR²-(CH₂)_n-SR⁸, NR²-(CH₂)_n-NR⁴-(CH₂)_p-NR⁹R¹⁰, NR²-(CH₂)_n-NR⁴-(CH₂)_p-N⁺R⁹R¹⁰R¹¹ Y⁻, NR²-(CH₂)_n-NR⁴-(CH₂)_p-O(PO₃H)⁻ W⁺, NR²-(CH₂)_n-NR⁴-(CH₂)_p-Si(R⁷)₃, NR²-(CH₂)_n-NR⁴-(CH₂)_p-SR⁸, O(CH₂)_n-NR⁴R⁵, O(CH₂)_n-N⁺R⁴R⁵R⁶ Y⁻, O(CH₂)_n-O(PO₃H)⁻ W⁺, O(CH₂)_n-Si(R⁷)₃, O(CH₂)_n-SR⁸, O(CH₂)_n-NR⁴-(CH₂)_p-NR⁹R¹⁰, O(CH₂)_n-NR⁴-(CH₂)_p-N⁺R⁹R¹⁰R¹¹ Y⁻, O(CH₂)_n-NR⁴-(CH₂)_p-O(PO₃H)⁻ W⁺ or O(CH₂)_n-NR⁴-(CH₂)_p-Si(R⁷)₃; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

R^3 is alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

each R^4 , R^6 , R^8 , R^9 , R^{10} and R^{11} is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^5 is alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^7 is alkyl, O(alkyl) or O(tri-substituted silyl);

R^{13} is H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

W^+ is an agriculturally acceptable cation;

Y^- is an agriculturally acceptable anion;

n is an integer selected from 1 to 16;

p is an integer selected from 1 to 16;

m is an integer selected from 1 to 100;

q is an integer selected from 0 to 16;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

===== is a single bond or a double bond;


===== is a single bond or a double bond; and

M is 2H or a metal species,

wherein each substituted alkyl, substituted aryl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[006] In yet another aspect of the present description, there is provided a compound of Formula I-B1:

5



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is NR^2R^3 ; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2R^3$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

R^3 is alkyl or substituted alkyl;


each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[007] In yet another aspect of the present description, there is provided a compound of Formula I-B1:

6



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-NR^4R^5$ or $O-(CH_2)_n-NR^4R^5$; and
the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2)_n-NR^4R^5$ or $O-(CH_2)_n-NR^4R^5$; and

$Z^2 = Z^1$;

R^5 is alkyl, substituted alkyl or $-(CH_2)_p-NR^9R^{10}$;

each R^1 , R^2 , R^4 , R^9 and R^{10} is, independently, H, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

p is an integer selected from 1 to 16;


each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is $2H$ or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[008] In yet another aspect of the present description, there is provided a compound of Formula I-B1:

7



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-Si(R^7)_3$, $O-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $O-(CH_2)_n-SR^8$; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2)_n-Si(R^7)_3$, $O-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $O-(CH_2)_n-SR^8$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

R^7 is alkyl, O(alkyl) or O(trisubstituted silyl);

R^8 is H, alkyl, substituted alkyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^{13} is H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

n is an integer selected from 1 to 16;

m is an integer selected from 1 to 100;

q is an integer selected from 0 to 16;


each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is $2H$ or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[009] In yet another aspect of the present description, there is provided a compound of Formula I-B1:

8



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)}_2$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)}_2$, $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$; and the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = \text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)}_2$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)}_2$, $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$; and $Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

W^+ is an agriculturally acceptable cation;


each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[010] In yet another aspect of the present description, there is provided a compound of Formula I-B1:

9



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-NR^4R^5R^{6+} Y^-$ or $O-(CH_2)_n-NR^4R^5R^{6+} Y^-$; and the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2)_n-NR^4R^5R^{6+} Y^-$ or $O-(CH_2)_n-NR^4R^5R^{6+} Y^-$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

each R^4 , R^5 and R^6 is, independently, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

Y^- is an agriculturally acceptable anion;


each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[011] In yet another aspect of the present description, there is provided a compound of Formula I-B1:

10



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2CH_2O)_m-R^{13}$ or $O-(CH_2CH_2O)_m-R^{13}$; and the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2CH_2O)_m-R^{13}$ or $O-(CH_2CH_2O)_m-R^{13}$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

R^{13} is H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

m is an integer selected from 1 to 100;


each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is $2H$ or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[012] In yet another aspect of the present description, there is provided a compound of Formula I-B1:

11



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and

the other one of Z^1 and Z^2 is OR^1 ;

or

Z^1 is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[013] The present description also provides a composition for use in promoting the health of a plant, the composition comprising at least one compound as defined herein, or an agriculturally acceptable salt thereof, and a carrier fluid.

[014] The present description also provides a method for promoting the health of a plant, comprising: applying to the plant a compound as defined herein or an agriculturally acceptable salt thereof, or a composition as defined herein; and exposing the plant to light. Promoting the health of the plant can include at least one of preventing or inhibiting growth of a microbial pathogen of the plant (e.g., a fungus or a bacterial pathogen), increasing

resistance of the plant to one or more abiotic stress, and controlling a pest of the plant (e.g., a noxious insect or corresponding larva).

DETAILED DESCRIPTION

[015] Some microbial pathogens, such as Gram-negative bacteria and certain types of fungi have a cellular membrane that is difficult to penetrate. More specifically, these microbial pathogens sometimes have an impermeable outer cell membrane that contains endotoxins and can block small molecules such as antibiotics, dyes and detergents, thereby protecting the sensitive inner membrane and cell wall. It can therefore be challenging to use photodynamic therapy to inhibit growth of certain microbial pathogens in plants because the photosensitizer compounds tend to not achieve good penetration inside the cell wall. It can also be challenging to increase resistance of plants to damage caused by abiotic stresses.

[016] In some scenarios, photodynamic inhibition of microbial pathogens that are present on plants can be achieved by applying a photosensitizer compound. The photosensitizer compound reacts to light by generating reactive oxygen species (ROS). In other scenarios, a photosensitizer compound can be used to increase resistance of plants to damage caused by one or more abiotic stress.

[017] Some of the compounds of the present description are photosensitizer compounds that can be derived from the protoporphyrin IX (hereinafter “PP IX”) scaffold. The compounds derived from PP IX can also be referred to as “modified PP IX”. Some of the compounds of the present description are photosensitizer compounds that have a scaffold similar to that of PP IX, but that are not necessarily derived from PP IX. In some scenarios, these compounds can be used to improve the health of plants. That is, depending on the compound and/or mode of application, the compounds of the present description can be applied to plants to photodynamically inhibit microbial pathogens on plants, to increase resistance of plants to damage caused by one or more abiotic stress and/or to be used as insecticides.

Definitions

[018] Unless stated otherwise, the following terms and phrases as used herein are intended to have the following meanings.

[019] When trade names are used herein, it is intended to independently include the tradename product and the active ingredient(s) of the tradename product.

[020] As used herein, the phrase “a compound of Formula I” means a compound of Formula I or an agriculturally acceptable salt thereof. With respect to isolatable intermediates, the phrase “a compound of Formula (number)” means a compound of that formula and salts thereof, and optionally agriculturally acceptable salts thereof.

[021] The term “Alkyl”, as used herein, means a hydrocarbon containing primary, secondary, tertiary or cyclic carbon atoms. For example, and without being limiting, an alkyl group can have 1 to 20 carbon atoms (*i.e.*, C₁-C₂₀ alkyl), 1 to 8 carbon atoms (*i.e.*, C₁-C₈ alkyl), 1 to 6 carbon atoms (*i.e.*, C₁-C₆ alkyl) or 1 to 4 carbon atoms (*i.e.*, C₁-C₄ alkyl). Examples of suitable alkyl groups include, but are not limited to, methyl (Me, -CH₃), ethyl (Et, -CH₂CH₃), 1-propyl (*n*-Pr, *n*-propyl, -CH₂CH₂CH₃), 2-propyl (*i*-Pr, *i*-propyl, -CH(CH₃)₂), 1-butyl (*n*-Bu, *n*-butyl, -CH₂CH₂CH₂CH₃), 2-methyl-1-propyl (*i*-Bu, *i*-butyl, -CH₂CH(CH₃)₂), 2-butyl (*s*-Bu, *s*-butyl, -CH(CH₃)CH₂CH₃), 2-methyl-2-propyl (*t*-Bu, *t*-butyl, -C(CH₃)₃), 1-pentyl (*n*-pentyl, -CH₂CH₂CH₂CH₂CH₃), 2-pentyl (-CH(CH₃)CH₂CH₂CH₃), 3-pentyl (-CH(CH₂CH₃)₂), 2-methyl-2-butyl (-C(CH₃)₂CH₂CH₃), 3-methyl-2-butyl (-CH(CH₃)CH(CH₃)₂), 3-methyl-1-butyl (-CH₂CH₂CH(CH₃)₂), 2-methyl-1-butyl (-CH₂CH(CH₃)CH₂CH₃), 1-hexyl (-CH₂CH₂CH₂CH₂CH₂CH₃), 2-hexyl (-CH(CH₃)CH₂CH₂CH₂CH₃), 3-hexyl (-CH(CH₂CH₃)(CH₂CH₂CH₃)), 2-methyl-2-pentyl (-C(CH₃)₂CH₂CH₂CH₃), 3-methyl-2-pentyl (-CH(CH₃)CH(CH₃)CH₂CH₃), 4-methyl-2-pentyl (-CH(CH₃)CH₂CH(CH₃)₂), 3-methyl-3-pentyl (-C(CH₃)(CH₂CH₃)₂), 2-methyl-3-pentyl (-CH(CH₂CH₃)CH(CH₃)₂), 2,3-dimethyl-2-butyl (-C(CH₃)₂CH(CH₃)₂), 3,3-dimethyl-2-butyl (-CH(CH₃)C(CH₃)₃), and octyl (-CH₂)₇CH₃.

[022] The term “Alkenyl”, as used herein, means a hydrocarbon containing primary, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, *i.e.* a carbon-carbon *sp*² double bond. For example, and without being limiting, an alkenyl group can have 2 to 20 carbon atoms (*i.e.*, C₂-C₂₀ alkenyl), 2 to 8 carbon atoms (*i.e.*, C₂-C₈ alkenyl), 2 to 6 carbon atoms (*i.e.*, C₂-C₆ alkenyl) or 2 to 4 carbon atoms (*i.e.*, C₂-C₄ alkenyl). Examples of suitable alkenyl groups include, but are not limited to, ethylene or vinyl (-CH=CH₂), allyl (-CH₂CH=CH₂), cyclopentenyl (-C₅H₇), and 5-hexenyl (-CH₂CH₂CH₂CH₂CH=CH₂).

[023] The term “Alkynyl”, as used herein, means a hydrocarbon containing primary, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, *i.e.* a carbon-carbon, *sp* triple bond. For example, and without being limiting, an alkynyl group can have 2 to 20 carbon atoms (*i.e.*, C₂-C₂₀ alkynyl), 2 to 8 carbon atoms (*i.e.*, C₂-C₈ alkynyl), 2 to 6 carbon atoms (*i.e.*, C₂-C₆ alkynyl) or 2 to 4 carbon atoms (*i.e.*, C₂-C₄ alkynyl). Examples of

suitable alkynyl groups include, but are not limited to, acetylenic ($\text{-C}\equiv\text{CH}$) and propargyl ($\text{-CH}_2\text{C}\equiv\text{CH}$).

[024] The term “Alkoxy”, as used herein, is interchangeable with the term “O(Alkyl)”, in which an “Alkyl” group as defined above is attached to the parent molecule via an oxygen atom. For example, and without being limiting, the alkyl portion of an O(Alkyl) group can have 1 to 20 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{20}$ alkyl), 1 to 8 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_8$ alkyl), 1 to 6 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_6$ alkyl) or 1 to 4 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_4$ alkyl). Examples of suitable Alkoxy or O(Alkyl) groups include, but are not limited to, methoxy (-OCH_3 or -OMe), ethoxy ($\text{-OCH}_2\text{CH}_3$ or -OEt) and *t*-butoxy ($\text{-O-C(CH}_3)_3$ or -OtBu). Similarly, “O(alkenyl)”, “O(alkynyl)” and the corresponding substituted groups will be understood by a person skilled in the art.

[025] The term “Acyl”, as used herein, is meant to encompass several functional moieties such as “C=O(Alkyl)”, “C=O(Alkenyl)”, “C=O(Alkynyl)” and their corresponding substituted groups, in which an “Alkyl”, “Alkenyl” and “Alkynyl” groups are as defined above and attached to an O, N, S of a parent molecule via a C=O group. For example, and without being limiting, the alkyl portion of a C=O(Alkyl) group can have 1 to 20 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{20}$ alkyl), 1 to 8 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_8$ alkyl), 1 to 6 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_6$ alkyl) or 1 to 4 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_4$ alkyl). Examples of suitable Acyl groups include, but are not limited to, formyl (*i.e.*, a carboxyaldehyde group), acetyl, trifluoroacetyl, propionyl, and butanoyl. A person skilled in the art will understand that a corresponding definition applies for “C=O(Alkenyl)” and “C=O(Alkynyl)” moieties. In the present description, “C=O(Alkyl)”, “C=O(Alkenyl)”, “C=O(Alkynyl)” can also be written as “CO(Alkyl)”, “CO(Alkenyl)” and “CO(Alkynyl)”, respectively.

[026] The term “Alkylene”, as used herein, means a saturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. For example, and without being limiting, an alkylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, 1 to 6 carbon atoms or 1 to 4 carbon atoms. Typical alkylene radicals include, but are not limited to, methylene ($\text{-CH}_2\text{-}$), 1,1-ethyl ($\text{-CH(CH}_3\text{)-}$), 1,2-ethyl ($\text{-CH}_2\text{CH}_2\text{-}$), 1,1-propyl ($\text{-CH(CH}_2\text{CH}_3\text{)-}$), 1,2-propyl ($\text{-CH}_2\text{CH(CH}_3\text{)-}$), 1,3-propyl ($\text{-CH}_2\text{CH}_2\text{CH}_2\text{-}$) and 1,4-butyl ($\text{-CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-}$).

[027] The term “Alkenylene”, as used herein, means an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the

removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkene. For example, and without being limiting, an alkenylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, 1 to 6 carbon atoms or 1 to 4 carbon atoms. Typical alkenylene radicals include, but are not limited to, 1,2-ethylene ($-\text{CH}=\text{CH}-$).

[028] The term “Alkynylene”, as used herein, means an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkyne. For example, and without being limiting, an alkynylene group can have 2 to 20 carbon atoms, 2 to 10 carbon atoms, 2 to 6 carbon atoms or 2 to 4 carbon atoms. Typical alkynylene radicals include, but are not limited to, acetylene ($-\text{C}\equiv\text{C}-$), propargyl ($-\text{CH}_2\text{C}\equiv\text{C}-$), and 4-pentynyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{C}-$).

[029] The term “Aryl”, as used herein, means an aromatic hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. For example, and without being limiting, an aryl group can have 6 to 20 carbon atoms, 6 to 14 carbon atoms, or 6 to 10 carbon atoms. Typical aryl groups include, but are not limited to, radicals derived from benzene (e.g., phenyl), substituted benzene, naphthalene, anthracene and biphenyl.

[030] The term “Arylalkyl”, as used herein, means an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with an aryl radical. Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like. For example, and without being limiting, the arylalkyl group can include 7 to 20 carbon atoms, e.g., the alkyl moiety is 1 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

[031] The term “Arylalkenyl”, as used herein, means an acyclic alkenyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but also an sp^2 carbon atom, is replaced with an aryl radical. The aryl portion of the arylalkenyl can include, for example, any of the aryl groups described herein, and the alkenyl portion of the arylalkenyl can include, for example, any of the alkenyl groups described herein. The arylalkenyl group can include 8 to 20 carbon atoms, e.g., the alkenyl moiety is 2 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

[032] The term “Arylalkynyl”, as used herein, means an acyclic alkynyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but also an sp carbon atom, is replaced with an aryl radical. The aryl portion of the arylalkynyl can include, for example, any of the aryl groups disclosed herein, and the alkynyl portion of the arylalkynyl can include, for example, any of the alkynyl groups disclosed herein. For example, and without being limiting, the arylalkynyl group can include 8 to 20 carbon atoms, *e.g.*, the alkynyl moiety is 2 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

[033] The term “heterocycle”, as used herein, means a group including a covalently closed ring wherein at least one atom forming the ring is a heteroatom. For example, and without being limiting, heterocyclic rings can be formed by three, four, five, six, seven, eight, nine, or more than nine atoms. Any number of those atoms can be heteroatoms (*i.e.*, a heterocyclic ring can include one, two, three, four, five, six, seven, eight, nine, or more than nine heteroatoms). In heterocyclic rings including two or more heteroatoms, those two or more heteroatoms can be the same or different from one another. Heterocycles can be substituted. Binding to a heterocycle can be at a heteroatom or via a carbon atom. It should also be understood that in the present description, the term “heterocycle” also encompasses “heteroaryl” groups.

[034] The term “protecting group”, as used herein, means a moiety of a compound that masks or alters the properties of a functional group or the properties of the compound as a whole. The chemical substructure of a protecting group can greatly vary. One function of a protecting group is to serve as an intermediate in the synthesis of the parental active substance. Chemical protecting groups and strategies for protection/deprotection are well known in the art. See: "Protective Groups in Organic Chemistry", Theodora W. Greene (John Wiley & Sons, Inc., New York, 1991).

[035] The term “substituted”, as used herein in reference to alkyl, alkylene, alkoxy, alkenyl, alkynyl, alkenylene, aryl, alkynylene, etc., for example “substituted alkyl”, “substituted alkylene”, “substituted alkoxy” – “or substituted O(Alkyl)”, “substituted alkenyl”, “substituted alkynyl”, “substituted alkenylene”, “substituted aryl” and “substituted alkynylene”, unless otherwise indicated, means alkyl, alkylene, alkoxy, alkenyl, alkynyl, alkenylene, aryl and alkynylene, respectively, in which one or more hydrogen atoms are each independently replaced with a non-hydrogen substituent.

[036] Typical non-hydrogen substituents include, but are not limited to, $-X$, $-R^B$, $-O^-$, $=O$, $-OR^B$, $-SR^B$, $-S^-$, $-NR^{B_2}$, $Si(R^C)_3$, $-N^+R^{B_3}$, $-NR^b-(Alk)-NR^{B_2}$, $-NR^B-(Alk)-N^+R^{B_3}$, $-NR^B-(Alk)-OR^B$, $-NR^B-(Alk)-OP(=O)(OR^B)(O^-)$, $-NR^B-(Alk)-OP(=O)(OR^B)_2$, $-NR^B-(Alk)-Si(R^C)_3$, $-NR^B-(Alk)-SR^B$, $-O-(Alk)-NR^{B_2}$, $-O-(Alk)-N^+R^{B_3}$, $-O-(Alk)-OR^B$, $-O-(Alk)-OP(=O)(OR^B)(O^-)$, $-O-(Alk)-OP(=O)(OR^B)_2$, $-O-(Alk)-Si(R^C)_3$, $-O-(Alk)-SR^B$, $=NR^B$, $-CX_3$, $-CN$, $-OCN$, $-SCN$, $-N=C=O$, $-NCS$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-NHC(=O)R^B$, $-OC(=O)R^B$, $-NHC(=O)NR^{B_2}$, $-S(=O)_2^-$, $-S(=O)_2OH$, $-S(=O)_2R^B$, $-OS(=O)_2OR^B$, $-S(=O)_2NR^{B_2}$, $-S(=O)R^B$, $-OP(=O)(OR^B)(O^-)$, $-OP(=O)(OR^B)_2$, $-P(=O)(OR^B)_2$, $-P(=O)(O^-)_2$, $-P(=O)(OH)_2$, $-P(O)(OR^B)(O^-)$, $-C(=O)R^B$, $-C(=O)X$, $-C(S)R^B$, $-C(O)OR^B$, $-C(O)O^-$, $-C(S)OR^B$, $-C(O)SR^B$, $-C(S)SR^B$, $-C(O)NR^{B_2}$, $-C(S)NR^{B_2}$ or $-C(=NR^B)NR^{B_2}$ where each X is independently a halogen: F, Cl, Br, or I; each R^B is independently H, alkyl, aryl, arylalkyl, a heterocycle, an alkyloxy group such as poly(ethyleneoxy), PEG or poly(methyleneoxy), or a protecting group; each R^C is independently alkyl, O(alkyl) or O(tri-substituted silyl); and each Alk is independently alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene or substituted alkynylene. Unless otherwise indicated, when the term "substituted" is used in conjunction with groups such as arylalkyl, which have two or more moieties capable of substitution, the substituents can be attached to the aryl moiety, the alkyl moiety, or both.

[037] It should also be understood that the term "tri-substituted silyl" refers to a silyl group that is independently substituted with three functional groups selected from alkyl, alkenyl, alkynyl, aryl and arylalkyl. Non-limiting examples of tri-substituted silyl groups include trimethylsilyl and dimethylphenylsilyl.

[038] The term "PEG" or "poly(ethylene glycol)", as used herein, is meant to encompass any water-soluble poly(ethylene oxide). Typically, substantially all, or all monomeric subunits are ethylene oxide subunits, though the PEG can contain distinct end capping moieties or functional groups. PEG chains of the present description can include one of the following structures: $-(CH_2CH_2O)_m-$ or $-(CH_2CH_2O)_{m-1}CH_2CH_2-$, depending on if the terminal oxygen has been displaced, where m is an integer, optionally selected from 1 to 100, 1 to 50, 1 to 30, 5 to 30, 5 to 20 or 5 to 15. The PEG can be capped with an "end capping group" that is generally a non-reactive carbon-containing group attached to a terminal oxygen or other terminal atom of the PEG. Non-limiting examples of end capping groups can include alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl),

CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl).

[039] The term “natural amino acid”, as used herein refers to the twenty natural amino acids. More specifically, the natural amino acid can be selected from the group consisting of: Alanine (Ala), Glycine (Gly), Isoleucine (Ile), Leucine (Leu), Proline (Pro), Valine (Val), Phenylalanine (Phe), Tryptophan (Trp), Tyrosine (Tyr), Aspartic acid (Asp), Glutamic acid (Glu), Arginine (Arg), Histidine (His), Lysine (Lys), Serine (Ser), Threonine (Thr), Cysteine (Cys), Methionine (Met), Asparagine (Asn) and Glutamine (Gln). The natural amino acids (except for Glycine) bear an asymmetric carbon (the alpha-carbon) that is of (S) configuration (also referred to as L-amino acids). It should be understood that the expression “natural amino acid attached to the compound by its amino group bonded to the alpha-carbon”, as used herein, means that a hydrogen of the amino group bonded to the alpha-carbon is removed and replaced by a bond between the amino acid and the rest of the compound.

[040] A person skilled in the art will recognize that substituents and other moieties of the compounds of the present description should be selected in order to provide an agriculturally useful compound which can be formulated into an acceptably stable agricultural composition that can be applied to plants. The definitions and substituents for various genus and subgenus of the compounds of the present description are described and illustrated herein. It should be understood by a person skilled in the art that any combination of the definitions and substituents described herein should not result in an inoperable species or compound. It should also be understood that the phrase “inoperable species or compound” means compound structures that violate relevant scientific principles (such as, for example, a carbon atom connecting to more than four covalent bonds) or compounds too unstable to permit isolation and formulation into agriculturally acceptable compositions.

[041] Selected substituents of the compounds of the present description can be present to a recursive degree. In this context, “recursive substituent” means that a substituent may recite another instance of itself. Because of the recursive nature of such substituents, theoretically, a large number of compounds can be present in any given implementation. For example, R^x includes a R^y substituent. R^y can be R. R can be W^β . W^β can be W^4 and W^4 can be R or include substituents including R^y . A person skilled in the art of organic chemistry understands that the total number of such substituents is to be reasonably limited by the desired properties of the compound intended. Such properties include, by way of example and not limitation, physical properties such as molecular weight, solubility or log P, application properties such

as activity against the intended target, possibility of application to plants, and practical properties such as ease of synthesis. Typically, each recursive substituent can independently occur 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0, times in a given implementation. For example, each recursive substituent can independently occur 3 or fewer times in a given embodiment. Recursive substituents are an intended aspect of the compounds of the present description. A person skilled in the art of organic chemistry understands the versatility of such substituents.

[042] The term “agriculturally acceptable salt”, as used herein, refers to salts that exhibit pesticidal activity (*i.e.*, that are active against one or more biotic stress) or that can improve resistance of a plant to one or more abiotic stresss. The term also refers to salts that are or can be converted in plants, water or soil to a compound or salt that exhibits pesticidal activity or that can improve resistance of a plant to one or more abiotic stress. The “agriculturally acceptable salt” can be an agriculturally acceptable cation or agriculturally acceptable anion. Non-limiting examples of agriculturally acceptable cations can include cations derived from alkali or alkaline earth metals and cations derived from ammonia and amines. For example, agriculturally acceptable cations can include sodium, potassium, magnesium, alkylammonium and ammonium cations. Non-limiting examples of agriculturally acceptable anions can include halide, phosphate, alkylsulfate and carboxylate anions. For example, agriculturally acceptable anions can include chloride, bromide, methylsulfate, ethylsulfate, acetate, lactate, dimethyl phosphate or polyalkoxylated phosphate anions.

[043] The term “optionally substituted”, as used herein in reference to a particular moiety of the compounds of the present description, means a moiety wherein all substituents are hydrogen or wherein one or more of the hydrogens of the moiety can be replaced by substituents such as those listed under the definition of the term “substituted” or as otherwise indicated.

[044] It will be understood that all enantiomers, diastereomers, and racemic mixtures, tautomers, polymorphs, and pseudopolymorphs of compounds within the scope of the formulae and compositions described herein and their agriculturally acceptable salts thereof, are embraced by the present description. All mixtures of such enantiomers and diastereomers are also within the scope of the present description.

[045] A compound of the present description and its agriculturally acceptable salts may exist as different polymorphs or pseudopolymorphs. As used herein, crystalline polymorphism

means the ability of a crystalline compound to exist in different crystal structures. The crystalline polymorphism may result from differences in crystal packing (packing polymorphism) or differences in packing between different conformers of the same molecule (conformational polymorphism). As used herein, crystalline pseudopolymorphism means the ability of a hydrate or solvate of a compound to exist in different crystal structures. Pseudopolymorphs of the compounds of the present description may exist due to differences in crystal packing (packing pseudopolymorphism) or due to differences in packing between different conformers of the same molecule (conformational pseudopolymorphism). The description and depiction of the compounds of the present description is intended to include all polymorphs and pseudopolymorphs of the compounds and their agriculturally acceptable salts.

[046] A compound of the present description and its agriculturally acceptable salts may also exist as an amorphous solid. As used herein, an amorphous solid is a solid in which there is no long-range order of the positions of the atoms in the solid. The description and depiction of the compounds of the present description is intended to include all amorphous forms of the compounds and their agriculturally acceptable salts.

[047] The modifier “about” used in connection with a quantity is inclusive of the stated value and has the meaning dictated by the context. For example, the modifier “about” can include the degree of error associated with the measurement of the quantity.

[048] For agricultural use (*i.e.*, for application to plants), salts of the compounds of the present description are agriculturally acceptable salts. However, salts which are not agriculturally acceptable can also find use, for example, in the preparation or purification of an agriculturally acceptable compound. All salts, whether or not they are agriculturally acceptable salts, are therefore to be understood as within the scope of the present description.

[049] It will be understood that the compounds described herein can be in their un-ionized, ionized, as well as zwitterionic form, and in combinations with various amounts of water (*e.g.*, stoichiometric amounts of water) such as in hydrates.

[050] Whenever a compound described herein is substituted with more than one of the same designated group, *e.g.*, “R¹” or “R²”, then it will be understood that the groups may be the same or different, *i.e.*, each group is independently selected. For example, in the

expression “Si(OR⁷)₃ with each R⁷ being independently alkyl or aryl”, it is understood that each R⁷ can independently be selected from alkyl groups and aryl groups. Si(OR⁷)₃ therefore includes both symmetrical groups where all three R⁷ are the same and asymmetrical groups where at least one R⁷ group is different from the other two R⁷ groups, or where each R⁷ group is different. It is also understood that this applies to all R^q or Z^q groups defined herein (e.g., q being selected from 1 to 17, from a to f or from A to C). A group “Z¹” will be understood to be necessarily the same as another group “Z²” only when it is explicitly stated that “Z¹ = Z²”.

[051] The compounds described herein can also exist as tautomeric forms in certain cases. Although only one delocalized resonance structure will typically be depicted, all such forms are contemplated within the scope of the present description. For example, various tautomers can exist for the tetrapyrrole ring systems described herein, and all their possible tautomeric forms are within the scope of the present description.

[052] The term “growing medium”, as used herein, refers to any soil (of any composition) or soil-free (e.g., hydroponic) medium that is suitable for growing and cultivating a plant. The growing medium can further include any naturally occurring and/or synthetic substance(s) that are suitable for growing and cultivating the plant. The phrase “any surface of the growing medium” or “a surface of the growing medium”, as used herein, refers to a surface that is directly exposed to natural and/or simulated light and/or weather.

[053] The term “applying”, as used herein, refers to contacting a surface of the growing medium with at least one compound of the present description (e.g., combinations, compositions, solutions, emulsions including at least one compound of the present description), by any means known in the art (e.g., pouring, root bathing, soil drenching, drip irrigation, etc.), or contacting an area that is beneath the surface of the growing medium with at least one compound of the present description (e.g., by soil injection), or any combination thereof, or directly contacting the plant with at least one compound of the present description (e.g., spraying).

[054] The term “crop plant”, as used herein, refers to a non-woody plant, which is grown, tended to, and harvested in a cycle of one year or less as source of foodstuffs and/or energy. Non-limiting examples of crop plants include sugar cane, wheat, rice, corn (maize), potatoes, sugar beets, barley, sweet potatoes, cassava, soybeans, tomatoes, and legumes (beans and peas). The crop plant can be a monocot or a dicot.

[055] The term “woody plant”, as used herein, refers to a woody perennial plant having a single stem or trunk, and bearing lateral branches at some distance from the ground (e.g., a tree). The woody plant can be a deciduous tree, an evergreen tree (e.g., a coniferous) or a shrub. Non-limiting examples of woody plants include maple trees, citrus trees, apple trees, pear trees, oak trees, ash trees, pine trees, and spruce trees.

[056] The term “turf grass”, as used herein, refers to a cultivated grass that provides groundcover, for example a turf or lawn that is periodically cut or mowed to maintain a consistent height. Grasses belong to the *Poaceae* family, which is subdivided into six subfamilies, three of which include common turf grasses: the *Festucoideae* subfamily of cool-season turf grasses; and the *Panicoideae* and *Eragrostoideae* subfamilies of warm-season turf grasses. A limited number of species are in widespread use as turf grasses, generally meeting the criteria of forming uniform soil coverage and tolerating mowing and traffic. In general, turf grasses have a compressed crown that facilitates mowing without cutting off the growing point. In the present context, the term “turf grass” includes areas in which one or more grass species are cultivated to form relatively uniform soil coverage, including blends that are a combination of different cultivars of the same species, or mixtures that are a combination of different species and/or cultivars.

[057] Non-limiting examples of turf grasses include: bluegrasses (e.g., Kentucky bluegrass), bentgrasses (e.g., creeping bentgrass), Redtop, fescues (e.g., red fescue), ryegrasses (e.g., annual ryegrass), wheatgrasses (e.g., crested wheatgrass), beachgrass, Brome grasses (e.g., Arizona Brome), cattails (e.g., sand cattail), Alkaligrass (*Puccinellia distans*), crested dog's-tail (*Cynosurus cristatus*), bermudagrass (*Cynodon* spp. such as *Cynodon dactylon*), hybrid bermudagrass (e.g., tifdwarf bermudagrass), Zoysiagrasses (e.g., *Zoysia japonica*), St. Augustinegrass (e.g., Bitter Blue St. Augustinegrass), Centipedegrass (*Eremochloa ophiuroides*), Carpetgrass (*Axonopus fissifolius*), Bahiagrass (*Paspalum notatum*), Kikuyugrass (*Pennisetum clandestinum*), Buffalograss (*Buchloe dactyloids*), Seashore paspalum (*Paspalum vaginatum*), Blue Grama (*Bouteloua gracilis*), Black Grama (*Bouteloua eriopoda*), Sideoats Grama (*Bouteloua curtipendula*), Sporobolus spp. (e.g., Alkali Sacaton), Sand Dropseed (*Sporobolus cryptandrus*), Prairie Dropseed (*Sporobolus heterolepis*), Hordeum spp. (e.g., California Barley), Common Barley, Meadow Barley, Alopecurus spp. (e.g., Creeping Foxtail and Meadow Foxtail), Stipa spp. (e.g., Needle & Thread), Elymus spp. (e.g., Blue Wildrye), Buffelgrass (*Cenchrus ciliaris*), Big Quaking Grass (*Briza maxima*), Big Bluestem (*Andropogon gerardii*), Little Bluestem (*Schizachyruim*

scoparium), Sand Bluestem (*Andropogon hallii*), Deergrass (*Muhlenbergia rigens*), Eastern Gamagrass (*Tripsacum dactyloides*), Galleta (*Hilaria jamesii*), Tufted Hairgrass (*Deschampsia caespitosa*), Indian Rice Grass (*Oryzopsis hymenoides*), Indian Grass (*Sorghastrum nutans*), Sand Lovegrass (*Eragrostis trichodes*); Weeping Lovegrass (*Eragrostis curvula*), California Melic (*Melica californica*), Prairie Junegrass (*Koeleria pyramidata*), Prairie Sandreed (*Calamovilfa longifolia*), Redtop (*Agrostis alba*), Reed Canarygrass (*Phalaris arundinacea*), Sloughgrass (*Spartina pectinata*), Green Sprangletop (*Leptochloa dubia*), Bottlebush Squirreltail (*Sitanion hystrix*), Panicum Switchgrass (*virgatum*), and Purple Threeawn (*Aristida purpurea*).

[058] The phrase “promoting the health of a plant”, as used herein, includes at least one of controlling a disease, condition, or injury caused by a pest of a plant and increasing abiotic stress resistance or tolerance in a plant. In other words, the phrase “promoting the health of a plant” includes at least one of “controlling infection of a plant by one or more biotic agent”, “controlling infestation of a plant by one or more insect” and “increasing resistance of a plant to one or more abiotic stress”.

[059] The phrase “controlling infection of a plant by a biotic agent”, as used herein, means to diminish, ameliorate, or stabilize the infection and/or any other existing unwanted condition or side effect that is caused by the association of a microbial pathogen or infestation of an insect on the plant. The microbial pathogen can include fungi, bacteria (gram positive or gram negative), viruses, viroids, virus-like organisms, phytoplasma, etc.

[060] The term “abiotic stress”, as used herein, refers to environmental conditions that negatively impact growth, development, yield and yield quality of crop and other plants. below optimum levels. Non-limiting examples of abiotic stresses include, for example: photooxidative conditions, drought (water deficit), excessive watering (flooding, and submergence), extreme temperatures (chilling, freezing and heat), extreme levels of light (high and low), radiation (UV-B and UV-A), salinity due to excessive Na⁺ (sodicity), chemical factors (e.g., pH), mineral (metal and metalloid) toxicity, deficiency or excess of essential nutrients, gaseous pollutants (ozone, sulfur dioxide), wind, mechanical factors, and other stressors.

[061] As used herein, the term “increasing stress resistance” (and the like) refers to an increase in the ability of a plant to survive or thrive in stress conditions. Enhanced resistance or tolerance can be specific for a particular stressor, e.g., drought, excess water, nutrient

deficiency, salt, cold, shade or heat, or multiple stressors. In some scenarios, increased resistance to one or more abiotic stresses can be exemplified by the reduction in degradation of quality of the plant, as compared to an untreated plant subjected to the same stress. In other scenarios, increased resistance to one or more abiotic stress can be exemplified by maintained or improved plant quality, as compared to an untreated plant subjected to the same stress.

Photosensitizer compounds

[062] As discussed above, photosensitizer compounds can be used to enable photodynamic inhibition of biotic agents (*i.e.*, microbial pathogens and/or insects) that are present on plants. The photosensitizer compounds react to light by generating reactive oxygen species (ROS).


[063] Depending on the type of ROS generated, photosensitizers can be classified into two classes, namely Type I photosensitizers and Type II photosensitizers. On the one hand, Type I photosensitizers form short lived free radicals through electron abstraction or transfer from a substrate when excited at an appropriate wavelength in the presence of oxygen. On the other hand, Type II photosensitizers form a highly reactive oxygen state known as “singlet oxygen”, also referred to herein as “reactive singlet oxygen species”. Singlet oxygen species are generally relatively long lived and can have a large radius of action.

[064] It should be understood that the photosensitizer compound can be metallated or non-metallated. When metallated, as can be the case for various nitrogen-bearing macrocyclic compounds that are complexed with a metal, the metal can be selected to generate either a Type I or a Type II photosensitizer in response to light exposure. For example, when porphyrin photosensitizer compounds are metallated with copper, the ROS that are generated are typically Type I photosensitizers. When the same porphyrin photosensitizer compounds are metallated with magnesium, the ROS that are generated are typically Type II photosensitizers. Both Type I and Type II photosensitizers can be used to enable photodynamic inhibition of biotic agents that are present on plants or to protect a plant from abiotic stress.

[065] It should be understood that the term “singlet oxygen photosensitizer”, as used herein, refers to a compound that produces reactive singlet oxygen species when excited by light. In

other words, the term “singlet oxygen photosensitizer” refers to a photosensitizer in which the Type II process defined above is dominant compared to the Type I process.


[066] Protoporphyrin IX (PP IX) is an organic compound, which is one of the most common porphyrins in nature. PP IX is a deeply colored pigment that is not very soluble even in basic water. PP IX is encountered in nature in the form of its iron complexes. When complexed with ferrous iron, the molecule is called heme. Other iron complexes have also been synthesized, for example with Fe(III) or Fe(IV). PP IX is a largely planar tetrapyrrole having a 20-carbon atom macrocyclic ring, each pyrrole being linked to two other pyrroles of the macrocyclic ring by a one-carbon bridge. It will be understood that by “tetrapyrrole”, it is meant four pyrrole-like rings. As used herein, a “pyrrole-like” ring is a five-atom ring with four carbon atoms and one nitrogen atom. In the depiction of PP IX below, the carbons of the macrocyclic ring are numbered from 1 to 20. In the chemical structure of PP IX, two carboxylic acid-bearing moieties are provided at the C13 ($\text{CH}_2\text{CH}_2\text{COOH}$) and C17 ($\text{CH}_2\text{CH}_2\text{COOH}$) positions.



Protoporphyrin IX (PP IX)

[067] The compounds of the present description include photosensitizer compounds based on or similar to the PP IX scaffold above, that are of general Formula I represented below, or an agriculturally acceptable salt thereof. In what follows, the term “photosensitizer compound” refers to one or more compounds of Formula I. In other words, the term “photosensitizer compound” can refer to one compound of Formula I or to a combination or mixture of two or more compounds of Formula I.

26



Formula I

[068] In one aspect, there is provided a compound of Formula I, or an agriculturally acceptable salt thereof, wherein:

Z^1 and Z^2 are each independently OR^1 or NR^2R^3 ;

each R^1 , R^2 and R^3 is independently H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, or substituted alkynyl, wherein:

if Z^1 and Z^2 are both OR^1 then at least one R^1 is not H,

if Z^1 and Z^2 are both NR^2R^3 then at least one R^3 is not H, and

if one of Z^1 and Z^2 is OR^1 and the other one of Z^1 and Z^2 is NR^2R^3 , then at least one of R^1 and R^3 is not H;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

----- is a single bond or a double bond;

===== is a single bond or a double bond; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted aryl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more -X, $-R^B$, $-O^-$, $=O$, $-OR^B$, $-SR^B$, $-S^-$, $-NR^B_2$, $Si(R^C)_3$, $-N^+R^B_3$, $-NR^B-(Alk)-NR^B_2$, $-NR^B-(Alk)-N^+R^B_3$, $-NR^B-(Alk)-OR^B$, $-NR^B-(Alk)-OP(=O)(OR^B)(O^-)$, $-NR^B-(Alk)-OP(=O)(OR^B)_2$, $-NR^B-(Alk)-Si(R^C)_3$, $-NR^B-(Alk)-SR^B$, $-O-(Alk)-NR^B_2$, $-O-(Alk)-N^+R^B_3$, $-O-(Alk)-OR^B$, $-O-(Alk)-OP(=O)(OR^B)(O^-)$, $-O-(Alk)-OP(=O)(OR^B)_2$, $-O-(Alk)-Si(R^C)_3$, $-O-(Alk)-SR^B$, $=NR^B$, $-CX_3$, $-CN$, $-OCN$, $-SCN$, $-N=C=O$, $-NCS$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-NHC(=O)R^B$, $-OC(=O)R^B$, $-NHC(=O)NR^B_2$, $-S(=O)_2-$, $-S(=O)_2OH$, $-S(=O)_2R^B$, $-OS(=O)_2OR^B$, $-S(=O)_2NR^B_2$, $-S(=O)R^B$, $-OP(=O)(OR^B)(O^-)$, $-OP(=O)(OR^B)_2$, $-P(=O)(OR^B)_2$, $-P(=O)(O^-)_2$, $-P(=O)(OH)_2$, $-P(O)(OR^B)(O^-)$, $-C(=O)R^B$,

$-C(=O)X$, $-C(S)R^B$, $-C(O)OR^B$, $-C(O)O^-$, $-C(S)OR^B$, $-C(O)SR^B$, $-C(S)SR^B$, $-C(O)NR^{B_2}$, $-C(S)NR^{B_2}$ or $-C(=NR^B)NR^{B_2}$;

each X is independently a halogen: F, Cl, Br or I;

each R^B is independently H, alkyl, aryl, arylalkyl, a heterocycle, an alkyloxy group such as poly(ethyleneoxy), PEG or poly(methyleneoxy), a capped poly(ethyleneoxy), capped PEG or capped polymethyleneoxy, or a protecting group;

the capped poly(ethyleneoxy), capped PEG and capped poly(methyleneoxy) groups being each independently capped with alkyl, aryl, arylalkyl, alkenyl, alkynyl, CO(alkyl), CO(aryl), CO(arylalkyl), CO(alkenyl) or CO(alkynyl);

each R^C is independently alkyl, aryl, arylalkyl, O(alkyl), O(aryl), O(arylalkyl), or O(tri-substituted silyl);

each tri-substituted silyl is independently substituted with three functional groups selected from alkyl, alkenyl, alkynyl, aryl and arylalkyl; and

each Alk is independently alkylene, alkenylene, or alkynylene.

[069] In some implementations, the compound of Formula I is such that:

one of Z^1 and Z^2 is OR^1 ; and

the other one of Z^1 and Z^2 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$, $NR^2-(CH_2)_n-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-N^+R^9R^{10}R^{11} Y^-$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-Si(R^7)_3$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-SR^8$, $O(CH_2)_n-NR^4R^5$, $O(CH_2)_n-N^+R^4R^5R^6 Y^-$, $O(CH_2)_n-O(PO_3H)^- W^+$, $O(CH_2)_n-Si(R^7)_3$, $O(CH_2)_n-SR^8$, $O(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$, $O(CH_2)_n-NR^4-(CH_2)_p-N^+R^9R^{10}R^{11} Y^-$, $O(CH_2)_n-NR^4-(CH_2)_p-O(PO_3H)^- W^+$ or $O(CH_2)_n-NR^4-(CH_2)_p-Si(R^7)_3$;

or

Z^1 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$, $NR^2-(CH_2)_n-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-N^+R^9R^{10}R^{11} Y^-$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-Si(R^7)_3$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-SR^8$, $O(CH_2)_n-NR^4R^5$, $O(CH_2)_n-N^+R^4R^5R^6 Y^-$, $O(CH_2)_n-O(PO_3H)^- W^+$, $O(CH_2)_n-Si(R^7)_3$, $O(CH_2)_n-SR^8$, $O(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$, $O(CH_2)_n-NR^4-(CH_2)_p-N^+R^9R^{10}R^{11} Y^-$, $O(CH_2)_n-NR^4-(CH_2)_p-O(PO_3H)^- W^+$ or $O(CH_2)_n-NR^4-(CH_2)_p-Si(R^7)_3$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

R^3 is alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

each R^4 , R^6 , R^8 , R^9 , R^{10} and R^{11} is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^5 is alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^7 is alkyl, O(alkyl) or O(tri-substituted silyl);

R^{13} is H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

W^+ is an agriculturally acceptable cation;

Y^- is an agriculturally acceptable anion;

n is an integer selected from 1 to 16;

p is an integer selected from 1 to 16;

m is an integer selected from 1 to 100;

q is an integer selected from 0 to 16;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

----- is a single bond or a double bond;

===== is a single bond or a double bond; and

M is $2H$ or a metal species,

wherein each substituted alkyl, substituted aryl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[070] In some implementations, $Z^1 = Z^2 = NR^2R^3$. In other implementations, Z^1 is NR^2R^3 and Z^2 is OH, or Z^1 is OH and Z^2 is NR^2R^3 . R^3 can for example be alkyl or substituted alkyl.

[071] In some implementations, ----- is a double bond and/or ===== is a double bond. More specifically: in some scenarios, ----- is a double bond and ===== is a double bond. In other scenarios, ----- is a double bond and ===== is a single bond. In yet other scenarios, ----- is a single bond and ===== is a double bond. In yet other scenarios, ----- is a single bond and ===== is a single bond.

[072] In some implementations, each R^a , R^b , R^c , R^d , R^e and R^f is, independently, alkyl or alkenyl. In a non-limiting example, R^a , R^c , R^e and R^f are methyl while R^b and R^d are vinyl.

[073] In some implementations, M is 2H. In some implementations, M is a metal species selected from the group consisting of Mg, Zn, Pd, Sn, Al, Pt, Si, Ge, Ga, In, Cu, Co, Fe and Mn. It should be understood that when a metal species is mentioned without its degree of oxidation, all suitable oxidation states of the metal species are to be considered, as would be understood by a person skilled in the art. In other implementations, M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III). In yet other implementations, M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II). In yet other implementations, M is a metal species selected from the group consisting of Cu(II), Co(III), Fe(III) and Mn(III).

[074] In some implementations, each R^1 , R^2 , R^4 , R^6 , R^8 , R^9 , R^{10} and R^{11} is, independently, H, alkyl or substituted alkyl. In some implementations, each R^3 and R^5 is, independently, alkyl or substituted alkyl. In some implementations, R^{13} is H, alkyl, substituted alkyl, CO(alkyl) or CO(substituted alkyl).

[075] In some implementations, the compound of Formula I is selected such that at least one of the following is true: R^1 is H, R^2 is H, R^3 is alkyl, R^4 is H or alkyl, R^5 is alkyl, R^6 is alkyl, R^7 is O(tri-substituted silyl), R^8 is H or alkyl, R^9 is alkyl, R^{10} is alkyl, R^{11} is alkyl and R^{13} is H, alkyl, alkenyl, CO(alkyl) or CO(alkenyl).

[076] In some implementations, W^+ is selected from the group consisting of sodium, potassium, magnesium and ammonium cations. In some implementations, Y^- is selected from the group consisting of chloride, bromide, phosphate, dimethylphosphate, methylsulfate, ethylsulfate, acetate and lactate.

[077] In some implementations, n is an integer selected from 1 to 16, or from 1 to 12, or from 1 to 8, or from 1 to 6, or from 1 to 4, or from 2 to 4. Similarly, in some implementations, p is an integer selected from 1 to 16, or from 1 to 12, or from 1 to 8, or from 1 to 6, or from 1 to 4, or from 2 to 4. Regarding the PEG moieties, m is an integer that can be selected from 1 to 100, or from 1 to 80, or from 1 to 60, or from 1 to 50, or from 1 to 30, or from 1 to 20, or from 1 to 10, or from 5 to 30, or from 5 to 20, or from 5 to 10. Still regarding PEG moieties, q is an integer that can be selected from 0 to 16, or from 0 to 8, or from 0 to 4, or from 0 to 2. In some implementations, q = 1. In other implementations, 1 = 0.

[078] In some implementations, Z^1 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$, $NR^2-(CH_2)_n-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$,


$\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-NR}^9\text{R}^{10}$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-N}^+\text{R}^9\text{R}^{10}\text{R}^{11} \text{Y}^-$,
 $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-O(PO}_3\text{H)}^- \text{W}^+$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-Si(R}^7\text{)}_3$,
 $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-SR}^8$, $\text{O(CH}_2\text{)}_n\text{-NR}^4\text{R}^5$, $\text{O(CH}_2\text{)}_n\text{-N}^+\text{R}^4\text{R}^5\text{R}^6 \text{Y}^-$,
 $\text{O(CH}_2\text{)}_n\text{-O(PO}_3\text{H)}^- \text{W}^+$, $\text{O(CH}_2\text{)}_n\text{-Si(R}^7\text{)}_3$, $\text{O(CH}_2\text{)}_n\text{-SR}^8$, $\text{O(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-NR}^9\text{R}^{10}$,
 $\text{O(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-N}^+\text{R}^9\text{R}^{10}\text{R}^{11} \text{Y}^-$, $\text{O(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-O(PO}_3\text{H)}^- \text{W}^+$ or
 $\text{O(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-Si(R}^7\text{)}_3$; and $\text{Z}^2 = \text{Z}^1$.

[079] In some implementations, one of Z^1 and Z^2 is NR^2R^3 , $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{R}^5$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-N}^+\text{R}^4\text{R}^5\text{R}^6 \text{Y}^-$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-O(PO}_3\text{H)}^- \text{W}^+$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-Si(R}^7\text{)}_3$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-SR}^8$ or $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-NR}^9\text{R}^{10}$; and the other one of Z^1 and Z^2 is OR^1 ; or Z^1 is NR^2R^3 , $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{R}^5$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-N}^+\text{R}^4\text{R}^5\text{R}^6 \text{Y}^-$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-O(PO}_3\text{H)}^- \text{W}^+$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-Si(R}^7\text{)}_3$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-SR}^8$ or $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-NR}^9\text{R}^{10}$; and $\text{Z}^2 = \text{Z}^1$.

[080] In some implementations, one of Z^1 and Z^2 is NR^2R^3 , $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{R}^5$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-N}^+\text{R}^4\text{R}^5\text{R}^6 \text{Y}^-$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-O(PO}_3\text{H)}^- \text{W}^+$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-Si(R}^7\text{)}_3$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-SR}^8$ or $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-NR}^9\text{R}^{10}$; and the other one of Z^1 and Z^2 is OR^1 .

[081] In some implementations, Z^1 is NR^2R^3 , $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{R}^5$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-N}^+\text{R}^4\text{R}^5\text{R}^6 \text{Y}^-$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-O(PO}_3\text{H)}^- \text{W}^+$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-Si(R}^7\text{)}_3$, $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-SR}^8$ or $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{-(CH}_2\text{)}_p\text{-NR}^9\text{R}^{10}$; and $\text{Z}^2 = \text{Z}^1$.

[082] In another aspect, there is provided a compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant,
 wherein:

one of Z^1 and Z^2 is NR^2R^3 ; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2R^3$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

R^3 is alkyl or substituted alkyl;


each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and


M is 2H or a metal species,


wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[083] In some implementations, R^1 is H, R^2 is H and/or R^3 is alkyl. R^3 can for example be a (C_1-C_{12}) alkyl, a (C_1-C_8) alkyl or a (C_1-C_4) alkyl. In some implementations, one of Z^1 and Z^2 is NR^2R^3 ; and the other one of Z^1 and Z^2 is OR^1 . In other implementations, $Z^1 = NR^2R^3$; and $Z^2 = Z^1$.

[084] In some implementations, the compound of Formula I-B1 is:














or an agriculturally acceptable salt thereof.

[085] In some implementations, the compound of Formula I-B1 is:





34




or an agriculturally acceptable salt thereof.

[086] In some implementations, the compound of Formula I-B1 is:





[087] In another aspect, there is provided a compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-NR^4R^5$ or $O-(CH_2)_n-NR^4R^5$; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2)_n-NR^4R^5$ or $O-(CH_2)_n-NR^4R^5$; and

$Z^2 = Z^1$;

R^5 is alkyl, substituted alkyl or $-(CH_2)_p-NR^9R^{10}$;

each R^1 , R^2 , R^4 , R^9 and R^{10} is, independently, H, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

p is an integer selected from 1 to 16;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is $2H$ or a metal species,


wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[088] In some implementations, R^1 is H, R^2 is H and/or R^4 is H or alkyl. In some implementations, R^4 is H and R^5 is alkyl. In some implementations, R^4 and R^5 are alkyl. R^4 and/or R^5 can for example each independently be a (C_1-C_{12}) alkyl, a (C_1-C_8) alkyl or a (C_1-C_4) alkyl. In some implementations, R^5 is $-(CH_2)_p-NR^9R^{10}$. In some implementations, R^9 and R^{10} are alkyl, or R^9 is H and R^{10} is alkyl. R^9 and/or R^{10} can for example each independently be a (C_1-C_{12}) alkyl, a (C_1-C_8) alkyl or a (C_1-C_4) alkyl.

[089] In some implementations, n is an integer selected from 1 to 16, or from 1 to 12, or from 1 to 8, or from 1 to 6, or from 1 to 4, or from 2 to 4. In some implementations, p is an integer selected from 1 to 16, or from 1 to 12, or from 1 to 8, or from 1 to 6, or from 1 to 4, or from 2 to 4.


[090] In some implementations, one of Z^1 and Z^2 is $NR^2-(CH_2)_n-NR^4R^5$; and the other one of Z^1 and Z^2 is OR^1 . In other implementations, $Z^1 = NR^2-(CH_2)_n-NR^4R^5$; and $Z^2 = Z^1$.

[091] In some implementations, the compound of Formula I-B1 is:




or an agriculturally acceptable salt thereof.

[092] In some implementations, the compound of Formula I-B1 is:



or an agriculturally acceptable salt thereof.


[093] In some implementations, the compound of Formula I-B1 is:



or an agriculturally acceptable salt thereof.

[094] In another aspect, there is provided a compound of Formula I-B1:

41



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-Si(R^7)_3$, $O-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $O-(CH_2)_n-SR^8$; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2)_n-Si(R^7)_3$, $O-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $O-(CH_2)_n-SR^8$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

R^7 is alkyl, O(alkyl) or O(trisubstituted silyl);

R^8 is H, alkyl, substituted alkyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^{13} is H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

n is an integer selected from 1 to 16;

m is an integer selected from 1 to 100;


q is an integer selected from 0 to 16;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is $2H$ or a metal species,


wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

43




or an agriculturally acceptable salt thereof,
wherein m is an integer selected from 4 to 15.

[098] In some implementations, the compound of Formula I-B1 is:





44



or an agriculturally acceptable salt thereof.


[099] In some implementations, the compound of Formula I-B1 is:





or an agriculturally acceptable salt thereof.

[0100] In another aspect, there is provided a compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR_2-(CH_2)_n-OP=O(OH)_2$ or $O-(CH_2)_n-OP=O(OH)_2$, $NR_2-(CH_2)_n-OP=O(OH)O^- W^+$ or $O-(CH_2)_n-OP=O(OH)O^- W^+$; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR_2-(CH_2)_n-OP=O(OH)_2$ or $O-(CH_2)_n-OP=O(OH)_2$, $NR_2-(CH_2)_n-OP=O(OH)O^- W^+$ or $O-(CH_2)_n-OP=O(OH)O^- W^+$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

W^+ is an agriculturally acceptable cation;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and


M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .


[0101] In some implementations, R^1 is H and/or R^2 is H. In some implementations, n is an integer selected from 1 to 16, or from 1 to 12, or from 1 to 8, or from 1 to 6, or from 1 to 4, or from 2 to 4. W^+ can be selected from the group consisting of sodium, potassium, magnesium and ammonium cations.

[0102] In some implementations, one of Z^1 and Z^2 is $NR_2-(CH_2)_n-OP=O(OH)_2$ or $O-(CH_2)_n-OP=O(OH)_2$, $NR_2-(CH_2)_n-OP=O(OH)O^- W^+$ or $O-(CH_2)_n-OP=O(OH)O^- W^+$; and the other one of Z^1 and Z^2 is OR^1 . In other implementations, $Z^1 = NR_2-(CH_2)_n-OP=O(OH)_2$ or $O-(CH_2)_n-OP=O(OH)_2$, $NR_2-(CH_2)_n-OP=O(OH)O^- W^+$ or $O-(CH_2)_n-OP=O(OH)O^- W^+$; and $Z^2 = Z^1$.

[0103] In some implementations, the compound of Formula I-B1 is:




[0104] In some implementations, the compound of Formula I-B1 is:



or an agriculturally acceptable salt thereof, wherein W^+ is a cation that can be selected from the group consisting of: of sodium, potassium, magnesium, ammonium and alkylammonium cations. For example, W^+ can be an *n*-octylammonium cation.

[0105] In another aspect, there is provided a compound of Formula I-B1:

49



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-NR^4R^5R^{6+} Y^-$ or $O-(CH_2)_n-NR^4R^5R^{6+} Y^-$; and the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2)_n-NR^4R^5R^{6+} Y^-$ or $O-(CH_2)_n-NR^4R^5R^{6+} Y^-$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

each R^4 , R^5 and R^6 is, independently, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

Y^- is an agriculturally acceptable anion;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and


M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

[0106] In some implementations, R^1 is H and/or R^2 is H. In some implementations, n is an integer selected from 1 to 16, or from 1 to 12, or from 1 to 8, or from 1 to 6, or from 1 to 4, or from 2 to 4. In some implementations, R^4 , R^5 and R^6 are alkyl and optionally $R^4 = R^5 = R^6$. In some implementations, Y^- is selected from the group consisting of chloride, bromide, phosphate, dimethylphosphate, methylsulfate, ethylsulfate, acetate and lactate.

[0107] In some implementations, one of Z^1 and Z^2 is $NR^2-(CH_2)_n-NR^4R^5R^{6+} Y^-$ or $O-(CH_2)_n-NR^4R^5R^{6+} Y^-$; and the other one of Z^1 and Z^2 is OR^1 . In other implementations, $Z^1 = NR^2-(CH_2)_n-NR^4R^5R^{6+} Y^-$ or $O-(CH_2)_n-NR^4R^5R^{6+} Y^-$; and $Z^2 = Z^1$.


[0108] In some implementations, the compound of Formula I-B1 is:



wherein Y^- is an agriculturally acceptable anion that can be selected from the group consisting of: chloride, bromide, phosphate, dimethylphosphate, methylsulfate, ethylsulfate, acetate, citrate, tartrate and lactate.

[0109] In another aspect, there is provided a compound of Formula I-B1:

51



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2CH_2O)_m-R^{13}$ or $O-(CH_2CH_2O)_m-R^{13}$; and the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2CH_2O)_m-R^{13}$ or $O-(CH_2CH_2O)_m-R^{13}$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

R^{13} is H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

m is an integer selected from 1 to 100;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and


M is $2H$ or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .


[0110] In some implementations, R^1 is H and/or R^{12} is H. In some implementations, m is an integer selected from 5 to 100, or from 5 to 80, or from 5 to 50, or from 5 to 20, or from 5 to 10. In some implementations, R^{13} is H, alkyl, alkenyl, CO(alkyl) or CO(alkenyl).

[0111] In some implementations, one of Z^1 and Z^2 is $NR^2-(CH_2CH_2O)_m-R^{13}$ or $O-(CH_2CH_2O)_m-R^{13}$; and the other one of Z^1 and Z^2 is OR^1 . In other implementations, $Z^1 = NR^2-(CH_2CH_2O)_m-R^{13}$ or $O-(CH_2CH_2O)_m-R^{13}$; and $Z^2 = Z^1$.

[0112] In some implementations, the compound of Formula I-B1 is:



53



or an agriculturally acceptable salt thereof,

wherein m is an integer selected from 1 to 100; and


R^{14} is alkyl, alkenyl, alkynyl, substituted alkyl, substituted alkenyl or substituted alkynyl.

[0113] In some implementations, m is an integer selected from 5 to 20 and $(CH_2CH_2O)_mCOR^{14}$ is PEG₆₀₀-oleate, PEG₄₀₀-oleate, PEG₆₀₀-allyl or PEG₄₀₀-allyl.

[0114] In some implementations, R^{13} is H. In some implementations, $(CH_2CH_2O)_mR^{13}$ is PEG₆₀₀ or PEG₄₀₀.


[0115] In some implementations, the compound of Formula I-B1 is:

54



or an agriculturally acceptable salt thereof.

[0116] In yet another aspect of the present description, a compound of Formula I-B1 is provided:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and

the other one of Z^1 and Z^2 is OR^1 ;

or

Z^1 is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .


[0117] In some implementations, one of Z^1 and Z^2 is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and the other one of Z^1 and Z^2 is OR^1 .

[0118] In other implementations, Z^1 is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and $Z^2 = Z^1$.

[0119] In some implementations, Z^1 is one of the natural amino acids and Z^2 is OH; Z^2 is one of the natural amino acids and Z^1 is OH; or Z^1 is one of the natural amino acids and $Z^2 = Z^1$.


[0120] In some implementations, Z^1 is Glycine, L-Alanine or L-Valine and Z^2 is OH; Z^2 is Glycine, L-Alanine or L-Valine and Z^1 is OH; or Z^1 is Glycine, L-Alanine or L-Valine and $Z^2 = Z^1$.

[0121] In some implementations, the compound of Formula I-B1 is:



or an agriculturally acceptable salt thereof.


[0122] In some implementations, the compound of Formula I-B1 is:



or an agriculturally acceptable salt thereof.

[0123] In some implementations, the compound of Formula I-B1 is:


58



or an agriculturally acceptable salt thereof.

[0124] In some implementations, the compound of Formula I-B1 is:

59




or an agriculturally acceptable salt thereof, wherein R^{17} is the side chain of one of the twenty natural amino acids.

Synthesis of the photosensitizer compounds


[0125] In another aspect, there is provided a method of manufacture of the compounds described herein, comprising reacting one of the compounds of Formula II, or a salt thereof, with a corresponding amine or an alcohol.

[0126] For example, a compound of Formula II can first be reacted with a coupling agent (e.g., DCC, EDC etc.) before being reacted with an amine, as shown in the scheme below, to give a mixture of mono and bis substituted amide compounds.

60



[0127] In another example, the compound of Formula II can be esterified using esterification methods known in the art.



[0128] Examples of such reactions can be found in the Examples section.

Chelating agent

[0129] In some implementations, the photosensitizer compound can be applied to a plant in combination with a chelating agent (also referred to herein as a permeabilizing agent). In some scenarios, the photosensitizer compound reacts to light by generating ROS, while the chelating agent can increase the overall impact of suppression of the growth of the microbial pathogen, for example by increasing the permeability of the outer membrane of the microbial pathogen to the photosensitizer. It should be understood that the term “chelating agent”, as used herein, refers generally to a compound that can form several chelating bonds to one or several metals or ions.

[0130] In some implementations, the chelating agent can include at least one carboxylic group, at least one hydroxyl group, at least one phenol group and/or at least one amino group or an agriculturally acceptable salt thereof. In some implementations, the chelating agent can include an aminocarboxylic acid compound or an agriculturally acceptable salt thereof. The aminocarboxylic acid or agriculturally acceptable salt thereof can include an amino polycarboxylic acid or an agriculturally acceptable salt thereof. For example, the amino polycarboxylic acid can include two amino groups and two alkylcarboxyl groups bound to each amino group. The alkylcarboxyl groups can be methylcarboxyl groups.

[0131] In some implementations, the chelating agent is selected from the group consisting of: an aminopolycarboxylic acid, an aromatic or aliphatic carboxylic acid, an amino acid, a phosphonic acid, and a hydroxycarboxylic acid or an agriculturally acceptable salt thereof.

[0132] In some implementations, the methods and compositions described herein include one or more aminopolycarboxylic acid chelating agents. Examples of aminopolycarboxylic acid chelating agents include, without limitation, ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), hydroxyethylenediaminetriacetic acid (HEDTA), and ethylenediaminedisuccinate (EDDS), cyclohexanediaminetetraacetic acid (CDTA), N-(2-hydroxyethyl)ethylenediaminetriacetic acid (EDTA-OH) glycol ether diaminetetraacetic acid (GEDTA), alanine diacetic acid (ADA), alkoyl ethylene diamine triacetic acids (e.g., lauroyl ethylene diamine triacetic acids (LED3A)), aspartic acid diacetic acid (ASDA), aspartic acid monoacetic acid, diamino cyclohexane tetraacetic acid (CDTA), 1,2-diaminopropanetetraacetic acid (DPTA-OH), 1,3-diamino-2-propanoltetraacetic acid (DTPA), diethylene triamine pentam ethylene phosphonic acid (DTPMP), diglycolic acid, dipicolinic acid (DPA), ethanolamine diacetic acid, ethanol diglycine (EDG), ethylenediaminediglutaric

acid (EDDG), ethylenediaminedi(hydroxyphenylacetic acid (EDDHA), ethylenediaminedipropionic acid (EDDP), ethylenediaminedisuccinate (EDDS), ethylenediaminemonosuccinic acid (EDMS), ethylenediaminetetraacetic acid (EDTA), ethylenediaminetetrapropionic acid (EDTP), and ethyleneglycolaminoethylestertetraacetic acid (EGTA) and agriculturally acceptable salts (for example, the sodium salts, calcium salts and/or potassium salts) thereof.

[0133] One non-limiting example of chelating agent is ethylenediaminetetraacetic acid (EDTA) or an agriculturally acceptable salt thereof. The aminocarboxylate salt can for example be a sodium or calcium salt.

[0134] Another non-limiting example of chelating agent is polyaspartic acid or an agriculturally acceptable salt thereof (*i.e.*, a polyaspartate), such as sodium polyaspartate. The molecular weight of the polyaspartate salt can for example be between 2,000 and 3,000.

[0135] The chelating agent can thus be a polymeric compound, which can include aspartate units, carboxylic groups, and other features found in polyaspartates. The polyaspartate can be a co-polymer that has alpha and beta linkages, which may be in various proportions (*e.g.*, 30% alpha, 70% beta, randomly distributed along the polymer chain). One non-limiting example of a sodium polyaspartate is Baypure® DS 100.

[0136] Other non-limiting examples of chelating agents include EDDS (ethylenediamine-N,N'-disuccinic acid), IDS (iminodisuccinic acid (N-1,2-dicarboxyethyl)-D,L-aspartic acid), isopropylamine, triethanolamine, triethylamine, ammonium hydroxide, tetrabutylammonium hydroxide, hexamine, GLDA (L-glutamic acid N,N-diacetic acid), or agriculturally acceptable salts thereof. The chelating agent can be metallated or non-metallated. In some implementations, IDS can be used as a tetrasodium salt of IDS (*e.g.*, tetrasodium iminodisuccinate), which can be Baypure® CX100. In some implementations, EDDS can be used as a trisodium salt of EDDS. In some implementations, GLDA can be used as a tetrasodium salt of GLDA.

[0137] In some implementations, the chelating agent can include one or more amino acid chelating agents. Examples of amino acid chelating agents include, without limitation, alanine, arginine, asparagine, aspartic acid, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, proline, serine, threonine, tyrosine, valine, or salts (for example, the sodium salts, calcium salts and/or potassium salts) and combinations thereof.

[0138] In some implementations, the chelating agent can include one or more aromatic or aliphatic carboxylic acid chelating agents. Examples of aromatic or aliphatic carboxylic acid chelating agents include, without limitation, oxalic acid, succinic acid, pyruvic acid malic, acid, malonic acid, salicylic acid, and anthranilic acid, and salts (for example, the sodium salts, calcium salts and/or potassium salts) thereof. In some implementations, the methods and compositions described herein include one or more polyphenol chelating agents. One non-limiting example of a polyphenol chelating agent is tannins such as tannic acid.

[0139] In some implementations, the chelating agent can include one or more hydroxycarboxylic acid chelating agents. Examples of the hydroxycarboxylic acid type chelating agents include, without limitation, malic acid, citric acid, glycolic acid, heptonic acid, tartaric acid and salts (for example, the sodium salts, calcium salts and/or potassium salts) thereof.

[0140] It will be understood that the one or more chelating agents can be provided as the free acid, as an agriculturally acceptable salt, or as combinations thereof. In some implementations, each of one or more the chelating agent(s) is applied as the free acid. In other implementations, the chelating agent(s) can be applied as a salt. Exemplary salts include sodium salts, potassium salts, calcium salts, ammonium salts, amine salts, amide salts, and combinations thereof. In still other implementations, when more than one chelating agent is present, at least one of the chelating agents is applied as a free acid, and at least one of the chelating agents is applied as a salt.

Additives and adjuvants

[0141] In some implementations, the photosensitizer compound can be applied to a plant in combination with one or more agriculturally suitable adjuvants. Each of the one or more agriculturally suitable adjuvants can be independently selected from the group consisting of one or more activator adjuvants (e.g., one or more surfactants; e.g., one or more oil adjuvants, e.g., one or more penetrants) and one or more utility adjuvants (e.g., one or more wetting or spreading agents; one or more humectants; one or more emulsifiers; one or more drift control agents; one or more thickening agents; one or more deposition agents; one or more water conditioners; one or more buffers; one or more anti-foaming agents; one or more UV blockers; one or more antioxidants; one or more fertilizers, nutrients, and/or micronutrients; and/or one or more herbicide safeners). Exemplary adjuvants are provided in Hazen, J.L. Weed Technology 14: 773-784 (2000), which is incorporated by reference in its entirety.

[0142] In some implementations, the photosensitizer compound can be applied to a plant in combination with oil. The oil can be selected from the group consisting of a mineral oil (e.g., paraffinic oil), a vegetable oil, an essential oil, and a mixture thereof. In some scenarios, combining the photosensitizer compound with an oil can improve solubility of the photosensitizer compound when in contact with the plant. The oil can be added with the photosensitizer compound, or separately, in the presence or absence of a carrier fluid such as water.

[0143] Non-limiting examples of vegetable oils include oils that contain medium chain triglycerides (MCT), or oil extracted from nuts. Other non-limiting examples of vegetable oils include coconut oil, canola oil, soybean oil, rapeseed oil, sunflower oil, safflower oil, peanut oil, cottonseed oil, palm oil, rice bran oil or mixtures thereof. Non-limiting examples of mineral oils include paraffinic oils, branched paraffinic oils, naphthenic oils, aromatic oils or mixtures thereof.

[0144] Non-limiting examples of paraffinic oils include various grades of poly-alpha-olefin (PAO). For example, the paraffinic oil can include HT60™, HT100™, High Flash Jet, LSRD™, and N65DW™. The paraffinic oil can include a paraffin having a number of carbon atoms ranging from about 12 to about 50, or from about 16 to 35. In some scenarios, the paraffin can have an average number of carbon atoms of 23. In some implementations, the oil can have a paraffin content of at least 80 wt%, or at least 90 wt%, or at least 99 wt%.

[0145] The photosensitizer compound and the oil can be added sequentially or simultaneously. When added simultaneously, the nitrogen-bearing macrocyclic compound and the oil can be added as part of the same composition or as part of two separate compositions. In some implementations, the nitrogen-bearing macrocyclic compound and the oil can be combined in an oil-in-water emulsion. That is, the combination can include the nitrogen-bearing macrocyclic compound combined with the oil and water so that the combination is formulated as an oil-in-water emulsion. The oil-in-water emulsion can also include other additives such as a chelating agent, a surfactant, or combinations thereof.

[0146] As used herein, the term “oil-in-water emulsion” refers to a mixture in which one of the oil (e.g., the paraffinic oil) and water is dispersed as droplets in the other (e.g., the water). In some implementations, an oil-in-water emulsion is prepared by a process that includes combining the paraffinic oil, water, and any other components and the paraffinic oil and applying shear until the emulsion is obtained. In other implementations, an oil-in-water

emulsion is prepared by a process that includes combining the paraffinic oil, water, and any other components in the mixing tank and spraying through the nozzle of a spray gun.

[0147] In some implementations, the photosensitizer compound is part of a composition that includes a carrier fluid. A suitable carrier fluid can allow obtaining a stable solution, suspension and/or emulsion of the components of the composition in the carrier fluid. In some implementations, the carrier fluid is water. In other implementations, the carrier fluid is a mixture of water and other solvents or oils that are non-miscible or only partially soluble in water.

[0148] In some implementations, a combination of photosensitizer compound and oil can be used to inhibit growth of a microbial pathogen in a plant. The combination can be an oil-in-water emulsion, where the surfactant is selected such that the photosensitizer compound is maintained in dispersion in the oil-in-water emulsion for delivery to the plant.

[0149] The combination can include a surfactant (also referred to as an emulsifier). The surfactant can be selected from the group consisting of an ethoxylated alcohol, a polymeric surfactant, a fatty acid ester, a poly(ethylene glycol), an ethoxylated alkyl alcohol, a monoglyceride, an alkyl monoglyceride, an amphipathic glycoside, and a mixture thereof. For example, the fatty acid ester can be a sorbitan fatty acid ester. The surfactant can include a plant derived glycoside such as a saponin. The surfactant can be present as an adjuvant to aid coverage of plant foliage. The surfactant can be an acceptable polysorbate type surfactant (e.g., Tween 80), a nonionic surfactant blend (e.g., Altox™ 3273), or another suitable surfactant.

[0150] In some implementations, the poly(ethylene glycol) can include a poly(ethylene glycol) of Formula $R^{15}-O-(CH_2CH_2O)_f-R^{16}$, wherein: each R^{15} and R^{16} is each, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, CO(alkyl) or CO(substituted alkyl); and f is an integer selected from 1 to 100; wherein the substituted alkyl groups are, independently, substituted with one or more F, Cl, Br, I, hydroxy, alkenyl, CN and N_3 .

Compositions including a photosensitizer compound

[0151] It should also be understood that the photosensitizer compounds and the other agents (e.g., chelating agent, oil, surfactant, etc.) can be provided to a plant separately or together as part of the same composition. In some implementations, the components of the compositions can be packaged in a concentrated form, without carrier fluid, and the carrier

fluid (e.g., water) can be added to form the composition directly by the operator that can then apply the composition to plants.

[0152] When the components are provided as part of a single composition, the composition can be provided to have certain concentrations and relative proportions of components. For example, the composition can have between about 100 nM and about 50 mM, between about 5 micromolar and about 10 mM, between about 1 micromolar and about 1000 micromolar, between about 5 micromolar and about 200 micromolar of the photosensitizer compound, between about 10 micromolar and about 150 micromolar of the nitrogen-bearing macrocyclic compound, between about 25 micromolar and about 100 micromolar of the nitrogen-bearing macrocyclic compound, or between about 50 micromolar and about 75 micromolar of the photosensitizer compound.

[0153] For example, and without being limiting, the composition can also include between about 2 micromolar and about 10,000 micromolar of the chelating agent, between about 5 micromolar and about 5,000 micromolar of the chelating agent, between about 10 micromolar and about 1,000 micromolar of the chelating agent, between about 25 micromolar and about 500 micromolar of the chelating agent, or between about 50 micromolar and about 100 micromolar of the chelating agent.

[0154] For example, and without being limiting, the relative proportion, by weight, of the nitrogen-bearing macrocyclic compound and the chelating agent in the composition can be between about 50:1 and about 1:1000, between about 20:1 and about 1:500, between about 10:1 and about 1:100, or between about 1:1 and about 1:10.

[0155] For example, and without being limiting, the photosensitizer compound and the oil can be applied in a relative proportion, by weight, between about 50:1 and about 1:1000, between about 20:1 and about 1:500, between about 10:1 and about 1:100, or between about 1:1 and about 1:10.

[0156] The composition including the photosensitizer compound can be applied to plants in various ways. For example, the composition can be prepared to include the photosensitizer compound, a chelating agent as well as a delivery fluid, such as water or a water-oil emulsion. The composition can be applied to the plant by spraying, misting, sprinkling, pouring, or any other suitable method. The anti-microbial composition can be applied to the foliage, roots

and/or stem of the plant. Other additives can also be included in the anti-microbial composition, and other application methods can also be performed.

[0157] The plants on which the composition is applied can be outdoors or indoors (e.g., greenhouse) where they are exposed to natural sunlight, or in an indoor location where they are exposed to artificial light. The exposure to the incident light is provided such that the photosensitizer compound can generate ROS that, in turn, facilitate disruption of microbial growth.

[0158] In some implementations, the photosensitizer compound can be used to treat seeds or seedlings. In some scenarios, the treatment of seeds or seedlings can stimulate germination and growth, and/or can increase resistance of the plant to abiotic stresses. In some implementations, the seeds or seedlings can be treated with the photosensitizer compound prior to being planted into a growing medium. In some implementations, the seeds or seedlings can be treated with the photosensitizer compound after being planted into a growing medium.

[0159] The photosensitizer compound can be directly surface-coated onto the seeds, applied to seedlings roots or seedlings leafs (foliar application on seedlings). In some implementations, a solution or emulsion containing the photosensitizer compound can be directly sprayed onto the seeds or seedlings. In some implementations, the seeds or seedlings can be dipped into a solution or emulsion containing the photosensitizer compound. In some implementations, the root of the seedling can be dipped into a solution or emulsion containing the photosensitizer compound. In some implementations, the seeds can be placed into a container, and a solution containing the photosensitizer compound can be introduced into the container. The container can then be shaken for an appropriate period (e.g., between about 1 minute to several minutes) such that the solution contacts the seeds. The shaken seeds can then be dried (e.g., air dried) prior to being planted.

[0160] The photosensitizer compound can be applied once, twice, or more than twice to seeds or seedlings, using various modes of application. For example, the seeds can be treated after having been planted into a growing medium. In another example, the seeds and/or seedlings can be treated prior to having been planted and after having been planted (e.g., in furrow treatment and/or foliar application). In yet another example, the seed can be treated prior to having been planted and/or after having been planted, and the ensuing seedling can be further treated (e.g., root treatment and/or foliar treatment).

Microbial pathogens

[0161] The microbial pathogens to which the composition including the photosensitizer compound can be applied include fungal and bacterial pathogens. In such case, the composition can be referred to as an “anti-microbial composition”.

[0162] The fungal pathogens to which the anti-microbial composition can be applied include *Alternaria solani*, which can infect plants such as tomatoes and potatoes; *Botrytis cinerea*, which can infect grapes, as well as soft fruits and bulb crops; or *Sclerotinia homoeocarpa*, which can commonly infect turfgrasses. Other fungal pathogens in the *Alternaria*, *Botrytis* or *Sclerotinia* genera can also receive application of the anti-microbial composition. The anti-microbial composition can be applied to plants that are affected or susceptible to pathogens that cause various plant diseases, e.g., *Colletotrichum*, *Fusarium*, *Puccinia*, *Erysiphaceae*, *Cercospora*, *Rhizoctonia*, *Bipolaris*, *Microdochium*, *Venturia inaequalis*, *Monilinia fructicola*, *Gymnosporangium juniperi-virginianae*, *Plasmodiophora brassicae*, *Ustilago zaeae*, *Phytophthora*, *Pythium*, *Fusarium oxysporum*, *Phytophthora infestans*, *Taphrina deformans*, *Powdery Mildew*, *Phragmidium spp.*, or other fungal pathogens.

[0163] The bacterial pathogens to which the anti-microbial composition can be applied include gram-negative bacteria, such as *Erwinia amylovora*, or other bacterial pathogens in the genus *Erwinia* that can infect woody plants. *E. amylovora* causes fire blight on various plants, including pears, apples, and other *Rosaceae* crops. The anti-microbial composition can be applied to plants that are affected or susceptible to pathogens that cause various plant diseases, e.g., *Pseudomonas*, *Xanthomonas*, *Agrobacterium*, *Curtobacterium*, *Streptomyces*, *E. Coli*, *Xylella fastidiosa* (which causes Olive Quick Decline Syndrome (OQDS) disease), or other bacterial pathogens.

[0164] It is also noted that the anti-microbial compositions described herein can have various inhibitory effects on the microbial pathogens depending on the type of plant and pathogen as well as the state of microbial infection. While herein it is described that the anti-microbial composition can inhibit microbial pathogen growth on a plant, such expressions should not be limiting but should be understood to include suppression of microbial pathogens, prevention against microbial pathogens, killing of microbial pathogens or generally increase toxicity toward microbial pathogens.

Abiotic stresses

[0165] As mentioned above, in some implementations, the photosensitizer compounds and compositions of the present description can be used to increase tolerance of plants to one or more abiotic stresses such as photooxidative conditions, drought (water deficit), excessive watering (flooding, and submergence), extreme temperatures (chilling, freezing and heat), extreme levels of light (high and low), radiation (UV-B and UV-A), salinity due to excessive Na^+ (sodicity), chemical factors (e.g., pH), mineral (metal and metalloid) toxicity, deficiency or excess of essential nutrients, gaseous pollutants (ozone, sulfur dioxide), wind, mechanical factors, and other stressors.

Cold hardiness

[0166] When the abiotic stress is cold stress, application of the photosensitizer compound, alone or in combination with additives such as an oil, a surfactant and/or a chelating agent, can improve cold hardiness of the plant. That is, application of the photosensitizer compound can allow the plant to withstand temperature conditions that are colder than would typically be experienced in the plant's optimal or native growing conditions. Various types of cold stress are possible, such as unexpected frost (for example an early fall frost when healthy crop, fruit, grain, seeds or leaves are still present on the plant, or a late spring frost that occurs after spring plant growth has begun), a cooler than average growing season, colder than native winter conditions, minimal winter snow cover, ice accumulation, etc.

[0167] It should be noted that what constitutes a cold stress condition for one plant may not be a cold stress condition for another plant. With reference to the USDA zone map, a cold stress condition for a zone 9 plant may in fact be a native growing condition for a zone 8 plant. Likewise, the depth of snow cover required for survival of one type of plant may not be required for a second type of plant. It is therefore understood that various types of cold stress are possible, depending on the type of plant in question.

[0168] The photosensitizer compound, compositions or combinations described herein may be used to protect plants, including woody plants, non-woody plants and turfgrasses, from frost injury. The frost can be an early frost, for example before harvest, after harvest and before dormancy. The frost can be a late frost, for example after budding. The cold damage can also be winter kill induced by winter temperatures, which may result in a loss of viable branches or shoots and lead to plant mortality. Plants treated by the photosensitizer

compound, compositions or combinations described herein can be frost or cold sensitive plants, in that they are naturally susceptible to frost, freezing or cold damage or injury in economically or aesthetically significant amounts.

[0169] Increasing resistance to cold stress can be exemplified by a delayed onset of dormancy. Plant dormancy can be triggered by a drop in temperature, *e.g.*, the onset of cold stress. By increasing resistance of the plant to cold stress, dormancy of the plant can be delayed until triggered by a further drop in temperature.

[0170] The photosensitizer compound, compositions or combinations described herein can be used periodically (*e.g.*, at 2 or 3-week intervals starting with spring at breaking the dormancy) and/or by applying one or more treatments (*e.g.*, 2 in the fall), to provide a response in reducing or delaying the dormancy period of certain plants.

[0171] As used herein, the term “reducing dormancy period” refers to a plant that has a reduced dormancy period or extended growing period relative to a control, *e.g.*, a non-treated plant.

[0172] In some implementations, the harvesting step may be carried out one week, one month, two months or more after the last application of the photosensitizer compound, compositions or combinations described herein, with the active agent still being effective to reduce the effects of cold stress on the plant during the intervening period.

[0173] In some scenarios, resistance to cold stress includes resistance to early or late frost, or winter damage. In some scenarios, the photosensitizer compound, compositions or combinations described herein can be used to protect early growth from cold during fluctuations in temperature (*e.g.*, in early spring). In some scenarios, the photosensitizer compound, compositions or combinations described herein can be used to protect plants from cold during the cold months (*e.g.*, in winter).

[0174] In some scenarios, the photosensitizer compound, compositions or combinations described herein can be applied by soil drenching and/or foliar application (*e.g.*, sprayed until run-off) at the onset or prior to exposure to the low temperature (*e.g.*, fall when the trees have full healthy and vigorous foliage). In some scenarios, the photosensitizer compound, compositions or combinations described herein can be applied by soil drenching and/or foliar application (*e.g.*, sprayed until run-off) during late fall and winter (*e.g.*, for warm climates). In some scenarios, the photosensitizer compound, compositions or combinations described

herein can be applied by soil drenching in the late fall following by a foliar application (e.g., sprayed until run-off) in the winter in order to reach maximum hardiness.

[0175] In some scenarios, the photosensitizer compound, compositions or combinations described herein can be applied 1-4 times at a 1 to 6-month interval (e.g., every 2 to 3 months). Further treatments may be applied in the spring and/or during the growing season to improve resistance to subsequent cold stress conditions.

Heat hardiness

[0176] When the abiotic stress is heat stress, application of the photosensitizer compound, compositions or combinations described herein can improve tolerance to high temperatures during the growing season. That is, application of the photosensitizer compound, compositions or combinations described herein can allow the plant to withstand temperature conditions that are higher than would typically be experienced in the plant's optimal or native growing conditions. Heat stress can have various causes, such as lack of shade for plants that typically require shaded growing conditions, or higher than normal soil and air temperatures.

[0177] It should be noted that what constitutes a heat stress condition for one plant may not be a heat stress condition for another plant.

Photooxidative hardiness

[0178] When the abiotic stress is photooxidative stress, application of the photosensitizer compound, compositions or combinations described herein can improve tolerance to stressful light condition during periods of increased generation of reactive oxygen species. That is, application of the photosensitizer compound, compositions or combinations described herein can allow the plant to withstand light exposure conditions (e.g., ultraviolet irradiation conditions) that are higher than would typically be experienced in the plant's optimal or native growing conditions. Photooxidative stress can have various causes, such as high light conditions or certain types of lighting that induce formation of free radicals.

[0179] It should be noted that what constitutes a photooxidative stress condition for one plant may not be a photooxidative stress condition for another plant.

Shade hardiness

[0180] Shade stress, or “low light (LL) stress” can be a problem that influences plant growth and quality. When the abiotic stress is shade stress, application of the photosensitizer compound, compositions or combinations described herein can improve shade hardiness of the plant. That is, application of the photosensitizer compound, compositions or combinations described herein can allow the plant to withstand shady conditions for plants whose optimal or native growing conditions typically require partial or full sun exposure. Various types of shade stress are possible, such as a prolonged period of cloudy weather, excessive growth of adjacent plants or trees that cast shade onto the plant, or lack of availability of a sunny planting location.

[0181] Shade can be a periodic problem. For example, during certain months of the year, a structure situated near a plant may cast a shadow on the plant, causing a shade stress. As the earth moves over the course of a year, the structure may no longer cast the shadow on the plant for another series of months and then the situation can be repeated during the next annual cycle. In such instances, the photosensitizer compound, compositions or combinations described herein can be applied to the plant prior to onset of the period of shade stress and can also be applied during the period of shade stress. The damage to the plant that would typically result on account of the period of shade stress can be prevented or reduced.

[0182] Shade conditions are not considered to be an abiotic stress condition for many types of plants, as some plants have a requirement for shade as part of their optimal growing conditions. It should also be noted that what constitutes a shade stress condition for one plant may not be a shade stress condition for another plant.

Drought hardiness

[0183] Drought can be defined as the absence of rainfall or irrigation for a period of time sufficient to deplete soil moisture and injure plants. Drought stress results when water loss from the plant exceeds the ability of the plant's roots to absorb water and/or when the plant's water content is reduced enough to interfere with normal plant processes. The severity of the effect of a drought condition may vary between plants, as the plant's need for water may vary by plant type, plant phenological stage, plant age, root depth, soil quality, etc.

[0184] The photosensitizer compound, compositions or combinations described herein can be applied to a plant prior to onset of a drought and/or during a drought. Application of the photosensitizer compound, compositions or combinations described herein can increase the resistance of the plant to the drought stress. Increasing resistance can include maintaining or increasing a quality of the plant as compared to an untreated plant subjected to the same drought stress. Increasing resistance can include reducing the degradation in quality of the plant, as compared to an untreated plant subjected to the same drought stress. If plants do not receive adequate rainfall or irrigation, the resulting drought stress can reduce growth more than all other environmental stresses combined.

[0185] It should also be noted that what constitutes a drought stress condition for one plant may not be a drought stress condition for another plant.

Prevention of salt damage

[0186] Salts can be naturally present in the growing environment of a plant. Salinity stress refers to osmotic forces exerted on a plant when the plant is growing in a saline soil or under other excessively saline conditions. For example, plants growing near a body of salt water can be exposed to salt present in the air or in water used to water the plants. In another example, salt applied to road, sidewalk and driveway surfaces during the winter for improved driving conditions can be transferred and/or leach into the soil of plants growing in the proximity. Such increased salt content in a growing environment of the plant can result in salinity stress, which can damage the plant.

[0187] Application of the photosensitizer compound, compositions or combinations described herein to the plant can increase the plant's resistance to the salinity stress and prevent or reduce a deterioration in quality of the plant which would occur if untreated. The combination can be applied prior to or during the period of salinity stress.

[0188] It should also be noted that what constitutes a salt stress condition for one plant may not be a salt stress condition for another plant.

Transplant shock hardness

[0189] A plant that is subjected to transplanting from one growing environment to another, e.g., from a pot to flower bed or garden, can be subjected to transplant shock stress as a result of exposure to new environmental conditions such as wind, direct sun, or new soil

conditions. Application of the photosensitizer compound, compositions or combinations described herein to the roots of the plant can reduce the impact to the plant caused by the transplanting. In some scenarios, stunting of plant growth and/or development of a transplanted plant can be reduced or prevented by application of the photosensitizer compound, compositions or combinations described herein.

[0190] It should be noted that what constitutes a transplant shock stress condition for one plant may not be a transplant shock stress condition for another plant.

Excess water or flooding hardiness

[0191] Although plants require a certain volume of water for healthy plant growth and development, the exposure of a plant to excess volumes of water (“water stress”) can damage the plant. Application of the photosensitizer compound, compositions or combinations described herein to a plant prior to the onset of an excess water condition can increase the plant’s resistance to the water stress. The photosensitizer compound, compositions or combinations described herein can be applied during the water stress, however, dilution of the photosensitizer compound, compositions or combinations described herein may occur on account of the excess water. Accordingly, pre-treatment in advance of a period of excess water can be more effective.

[0192] It should be noted that what constitutes an excess water stress condition for one plant may not be an excess water stress condition for another plant.

Insecticide activity

[0193] In some implementations, the compounds and combinations of the present description can be used to protect the plant from a plant pest. It should be understood that the term “plant pest” or “pest”, as used herein, refers to insects and/or their larvae, which are known to or have the potential to cause damage to the plant. In some implementations, the compounds and combinations of the present description can induce photoinduced mortality in plant pests.

[0194] In some implementations, the insect pests are selected from the order of Hemiptera (groups of aphids, whiteflies, scales, mealybugs, stink bugs), Coleoptera (groups of beetles), Lepidoptera (groups of butterflies, moths), Diptera (groups of flies), Thysanoptera (group of

thrips), Orthoptera (group of grasshoppers, locusts), Hymenoptera (groups of wasps, ants), Blattodea (groups of cockroaches and termites) and mite pests (spider mites).

[0195] Non-limiting examples of insect pests include: larvae of the order Lepidoptera, such as armyworms, (e.g., beet armyworm (*Spodoptera exigua*)), cutworms, loopers, (e.g., cabbage looper (*Trichoplusia ni*)) and heliothines, in the family Noctuidae (e.g., fall armyworm (*Spodoptera fugiperda* J. E. Smith)), beet armyworm (*Spodoptera exigua* Hubner), black cutworm (*Agrotis ipsilon* Hufnagel), and tobacco budworm (*Heliothis virescens* Fabricius); borers, casebearers, webworms, coneworms, cabbageworms and skeletonizers from the family Pyralidae (e.g., European corn borer (*Ostrinia nubilalis* Hubner)), navel orangeworm (*Amyelois transitella* Walker), corn root webworm (*Crambus caliginosellus* Clemens), and sod webworms (Pyralidae: Crambinae) such as sod webworm (*Herpetogramma licarsisalis* Walker), leafrollers, budworms, seed worms, and fruit worms in the family Tortricidae (e.g., codling moth (*Cydia pomonella* Linnaeus)), grape berry moth (*Endopiza viteana* Clemens), oriental fruit moth (*Grapholita molesta* Busck) and many other economically important *Lepidoptera* (e.g., diamondback moth (*Plutella xylostella* Linnaeus)), pink bollworm (*Pectinophora gossypiella* Saunders), and gypsy moth (*Lymantria dispar* Linnaeus); foliar feeding larvae and adults of the order Coleoptera including weevils from the families Anthribidae, Bruchidae, and Curculionidae (e.g., boll weevil (*Anthonomus grandis* Boheman)), rice water weevil (*Lissorhoptrus oryzophilus* Kuschel), granary weevil (*Sitophilus granarius* Linnaeus), rice weevil (*Sitophilus oryzae* Linnaeus), annual bluegrass weevil (*Listronotus maculicollis* Dietz), bluegrass billbug (*Sphenophorus parvulus* Gyllenhal), hunting billbug (*Sphenophorus venatus* vestitus), Denver billbug (*Sphenophorus cicatristriatus* Fahraeus), flea beetles, cucumber beetles, rootworms, leaf beetles, Colorado potato beetles (*Leptinotarsa decemlineata*), and leafminers in the family Chrysomelidae, western corn rootworm (*Diabrotica virgifera virgifera* LeConte); chafers and other beetles from the family Scarabaeidae (e.g., Japanese beetle (*Popillia japonica* Newman)), oriental beetle (*Anomala orientalis* Waterhouse), northern masked chafer (*Cyclocephala borealis* Arrow), southern masked chafer (*Cyclocephala immaculate* Olivier), black turfgrass *ataenius* (*Ataenius spretulus* Haldeman), green June beetle (*Cotinis nitida* Linnaeus), Asiatic garden beetle (*Maladera castanea* Arrow), May/June beetles (*Phyllophaga* spp.) and European chafer (*Rhizotrogus majalis* Razoumowsky)); carpet beetles from the family Dermestidae; wireworms from the family Elateridae; bark beetles from the family Scolytidae; flour beetles from the family Tenebrionidae; adults and nymphs of the order Orthoptera including grasshoppers, locusts, and crickets (e.g., migratory grasshoppers (e.g., *Melanoplus*

sanguinipes Fabricius, *M. differentialis* Thomas)), American grasshoppers (e.g., *Schistocerca americana* Drury), desert locust (*Schistocerca gregaria* Forskal), migratory locust (*Locusta migratoria* Linnaeus), bush locust (*Zonocerus* spp.); adults and larvae of the order Diptera including leafminers, midges, fruit flies (Tephritidae), fruit flies (e.g., *Oscinella frit* Linnaeus), soil maggots; adults and nymphs of the orders Hemiptera and Homoptera such as plant bugs from the family Miridae, leafhoppers (e.g., *Empoasca* spp.) from the family Cicadellidae; planthoppers from the families Fulgoroidae and Delphacidae (e.g., corn plant hopper (*Peregrinus maidis*)); treehoppers from the family Membracidae; chinch bugs (e.g., hairy chinch bug (*Blissus leucopterus* hirtus Montandon) and southern chinch bug (*Blissus insularis* Barber) and other seed bugs from the family Lygaeidae; spittlebugs from the family Cercopidae; squash bugs from the family Coreidae; red bugs and cotton stainers from the family Pyrrhocoridae; mealybugs from the family Pseudococcidae (e.g. *Planicoccus citri* Risso), cicadas from the family Cicadidae; psyllids from the family Psyllidae (e.g. Citrus psyllid (*Diaphorina citri*)), whiteflies from the family Aleyrodidae (silverleaf whitefly (*Bemisia argentifolii*)); aphids from the family Aphididae, such as cotton melon aphid (*Aphis gossypii*), pea aphid (*Acyrtosiphon pisum* Harris), cowpea aphid (*Aphis craccivora* Koch), black bean aphid (*Aphis fabae* Scopoli), melon or cotton aphid (*Aphis gossypii* Glover), apple aphid (*Aphis pomi* De Geer), spirea aphid (*Aphis spiraecola* Patch), foxglove aphid (*Aulacorthum solani* Kaltenbach), strawberry aphid (*Chaetosiphon fragaefolii* Cockerell), Russian wheat aphid (*Diuraphis noxia* Kurdjumov/Mordvilko), rosy apple aphid (*Dysaphis plantaginea* Paaserini), woolly apple aphid (*Eriosoma lanigerum* Hausmann), mealy plum aphid (*Hyalopterus pruni* Geoffroy), turnip aphid (*Lipaphis erysimi* Kaltenbach), cereal aphid (*Metopolophium dirrhodum* Walker), potato aphid (*Macrosipum euphorbiae* Thomas), peach-potato and green peach aphid (*Myzus persicae* Sulzer), lettuce aphid (*Nasonovia ribisnigri* Mosley), root aphids and gall aphids, corn leaf aphid (*Rhopalosiphum maidis* Fitch), bird cherry-oat aphid (*Rhopalosiphum padi* Linnaeus), greenbug (*Schizaphis graminum* Rondani), English grain aphid (*Sitobion avenae* Fabricius), spotted alfalfa aphid (*Therioaphis maculata* Buckton), black citrus aphid (*Toxoptera aurantii* Boyer de Fonscolombe), brown citrus aphid (*Toxoptera citricida* Kirkaldy) and green peach aphid (*Myzus persicae*); phylloxera from the family Phylloxeridae; mealybugs from the family Pseudococcidae; scales from the families Coccidae, Diaspididae, and Margarodidae; lace bugs from the family Tingidae; stink bugs from the family Pentatomidae; adults and immatures of the order Thysanoptera including onion thrips (*Thrips tabaci* Lindeman), flower thrips (*Frankliniella* spp.), and other foliar feeding thrips. Agronomic pests also include invertebrate arthropods such as mites from the family Tetranychidae: twospotted spider mite (e.g. *Tetranychus urticae* Koch), flat mite from

family Rutacea (e.g., citrus flat mite (*Brevipalpus lewisi* McGregor); rust and bud mites from the family Eriophyidae and other foliar feeding mites. Economically important agricultural pests nematodes (e.g., root knot nematodes in the genus *Meloidogyne*, lesion nematodes in the genus *Pratylenchus*, and stubby root nematodes in the genus *Trichodorus*) and members of the classes Nematoda, Cestoda, Trematoda, and Acanthocephala from orders of Strongylida, Ascaridida, Oxyurida, Rhabditida, Spirurida, and Enoplida.

[0196] The photosensitizer compounds of the present description can be applied to the plant before, at or after infestation of the plant by the insect pests.

[0197] In some implementations, the photosensitizer compounds of the present description can be used as insecticides by applying them to insects (i.e., without applying the photosensitizers to a plant). The present description therefore also provides a method for controlling insect population. The method includes applying to the insects a photosensitizer compound of the present description; and exposing the insects to light to activate the photosensitizer compound and generate reactive oxygen species.

[0198] It should be understood that applying the photosensitizer to the insects can include indirectly applying the photosensitizer to the insect pests (e.g., by applying the photosensitizer to a food source that is then fed to the insects) and/or directly applying the photosensitizer to the insect pests (e.g., by directly contacting the insects with the photosensitizer, such as by spraying a composition including the photosensitizer onto the insects).

Types of plants

[0199] The photosensitizer compounds and compositions of the present description can be used for various types of plants. The plant can be a non-woody crop plant, a woody plant or a turfgrass. The plant can be selected from the group consisting of a crop plant, a fruit plant, a vegetable plant, a legume plant, a cereal plant, a fodder plant, an oil seed plant, a field plant, a garden plant, a green-house plant, a house plant, a flower plant, a lawn plant, a turfgrass, a tree such as a fruit-bearing tree, and other plants that may be affected by microbial pathogens and/or one or more abiotic stress. Some of the compounds of the present description can display a certain degree of toxicity against a variety of noxious plant pests, in the absence or presence of light.

[0200] In some implementations, the plant is a crop plant selected from the group consisting of sugar cane, wheat, rice, corn (maize), potatoes, sugar beets, barley, sweet potatoes, cassava, soybeans, tomatoes, and legumes (beans and peas).

[0201] In other implementations, the plant is a tree selected from the group consisting of deciduous trees and evergreen trees. Examples of trees include, without limitation, maple trees, fruit trees such as citrus trees, apple trees, and pear trees, an oak tree, an ash tree, a pine tree, and a spruce tree.

[0202] In yet other implementations, the plant is a shrub.

[0203] In yet other implementations, the plant is a fruit or nut plant. Non-limiting examples of such plants include: acerola (barbados cherry), atemoya, carambola (star fruit), rambutan, almonds, apricots, cherries, nectarines, peaches, pistachio, apples, avocados, bananas, plantains, figs, grapes, mango, olives, papaya, pears, pineapple, plums, strawberries, grapefruit, lemons, limes, oranges (e.g., navel and Valencia), tangelos, tangerines, mandarins and plants from the berry and small fruits plant group.

[0204] In other implementations, the plant is a vegetable plant. Non-limiting examples of such plants include: asparagus, bean, beets, broccoli, Chinese broccoli, broccoli raab, brussels sprouts, cabbage, cauliflower, Chinese cabbage (e.g., bok choy and mapa), Chinese mustard cabbage (gai choy), cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens, celery, chayote, Chinese waxgourd, citron melon, cucumber, gherkin, hyotan, cucuzza, hechima, Chinese okra, balsam apple, balsam pear, bitter melon, Chinese cucumber, true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey galls, mango melon, Persian melon, pumpkin, summer squash, winter squash, watermelon, dasheen (taro), eggplant, ginger, ginseng, herbs and spices (e.g., curly leaf basil, lemon balm, cilantro, Mexican oregano, mint), Japanese radish (daikon), lettuce, okra, peppers, potatoes, radishes, sweet potatoes, Chinese artichoke (Japanese artichoke), corn and tomatoes.

[0205] In other implementations, the plant is a flowering plant, such as roses, flowering shrubs or ornamentals. Non-limiting examples of such plants include: flowering and foliage plants including roses and other flowering shrubs, foliage ornamentals & bedding plants, fruit-bearing trees such as apple, cherry, peach, and pear trees, non-fruit-bearing trees, shade

trees, ornamental trees, and shrubs (e.g., conifers, deciduous and broadleaf evergreens & woody ornamentals).

[0206] In some implementations, the plant is a houseplant. Non-limiting examples of such plants include: chrysanthemum, dieffenbachia, dracaena, ferns, gardenias, geranium, jade plant, palms, philodendron, and schefflera.

[0207] In some implementations, the plant is a plant grown in a greenhouse. Non-limiting examples of such plants include: ageratum, crown of thorns, dieffenbachia, dogwood, dracaena, ferns, ficus, holly, lisianthus, magnolia, orchid, palms, petunia, poinsettia, schefflera, sunflower, aglaonema, aster, azaleas, begonias, browallia, camellias, carnation, celosia, chrysanthemum, coleus, cosmos, crepe myrtle, dusty miller, easter lilies, fuchsia, gardenias, gerbera, helichrysum, hibiscus foliage, hydrangea, impatiens, jade plant, marigold, new guinea, impatiens, nicotonia, philodendron, portulaca, reiger begonias, snapdragon, and zinnias.

Synergistic effect of the combinations

[0208] In some scenarios, the combinations can exhibit a synergistic response for inhibiting growth of microbial pathogens in plants. It should be understood that the terms “synergy” or “synergistic”, as used herein, refer to the interaction of two or more components of a combination (or composition) so that their combined effect is greater than the sum of their individual effects. This may include, in the context of the present description, the action of two or more of the nitrogen-bearing macrocyclic compounds, the oil, and the chelating agent. In some scenarios, the nitrogen-bearing macrocyclic compound and the oil can be present in synergistically effective amounts. In some scenarios, the nitrogen-bearing macrocyclic compound and the chelating agent can be present in synergistically effective amounts. In some scenarios, the oil and the chelating agent can be present in synergistically effective amounts. In some scenarios, the nitrogen-bearing macrocyclic compound, the oil and the chelating agent can be present in synergistically effective amounts.


[0209] In some scenarios, the approach as set out in S. R. Colby, “Calculating synergistic and antagonistic responses of herbicide combinations”, Weeds 15, 20-22 (1967), can be used to evaluate synergy. Expected efficacy, E, may be expressed as: $E = X + Y(100 - X)/100$, where X is the efficacy, expressed in % of the untreated control, of a first component of a combination, and Y is the efficacy, expressed in % of the untreated control, of a second

component of the combination. The two components are said to be present in synergistically effective amounts when the observed efficacy is higher than the expected efficacy.

EXAMPLES

Compounds

[0210] PP IX-di-DMAE and PP IX-mono-DMAE




[0211] Protoporphyrin IX (500 mg, 0.89 mmol) and N-Hydroxysuccinimide (NHS, 245 mg, 2.136 mmol) were dissolved in 100 mL of DMF at 0°C. 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC, 409 mg, 2.136 mmol) in 10 mL DMF was then added to the reaction mixture and stirred at room temperature for 6 h. 2-Dimethylaminoethylamine (0.188 mg, 2.136 mmol) was then added and the solution was stirred overnight at room temperature. DMF was removed in vacuo and the residue was dissolved in 25 mL of methanol. The product was precipitated in cold ether (yield 80%).

[0212] The product contained about 50% di-substituted compound and 50% mono-substituted compound. When the reaction was performed a second time using the first product as starting material, the ratio was brought up to 80% di-substituted compound and 20% mono-substituted compound.

[0213] PP IX-mono-DMAE amide: MS-ESI (HRMS) m/z calculated for C₃₈ H₄₄ N₆ O₃ (M+H⁺): 633.35. Found: 633.087.

[0214] PP IX-di-DMAE amide: MS-ESI (HRMS) m/z calculated for C₄₂ H₅₄ N₈ O₂ (M): 703.44 Found: 703.611.


[0215] PP IX mono-PEG₆₀₀



[0216] Protoporphyrin IX (500 mg, 0.89 mmol) and N-Hydroxysuccinimide (NHS, 307 mg, 2.67 mmol) were dissolved in DMF (50 ml). The solution of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimidehydrochloride (EDC, 512 mg, 2.67 mmol) in DMF (10 ml) was then added to the reaction mixture and allowed to stirred overnight at room temperature. Methoxyl polyethylene glycol 600 (mPEG-OH600, 640 mg, 1.06 mmol) was added to the solution and stirred overnight at room temperature. The mixture was concentrated to afford the product. The crude product was purified by flash column chromatography over silica gel, eluting with PE/ME (5:1 to 2:1) to give the product.

[0217] Mass spectrometry and ¹H NMR analyses confirmed that a mixture of both PP IX mono-PEG₆₀₀ was obtained.


[0218] PP IX di-PEG₆₀₀



[0219] Protoporphyrin IX (500mg, 0.89mmol) and N-Hydroxysuccinimide (NHS, 610 mg, 5.2 mmol) were dissolved in DMF (50 ml) stirring at 0 °C for 1 h. A solution of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimidehydrochloride (EDC, 1012.0 mg, 5.2 mmol) in DMF (10 ml) was then added to the solution. The mixture was stirred overnight at room temperature. PEG600 (1290.0 mg, 2.2 mmol) was then added to the solution and further stirred overnight at room temperature. The mixture was concentrated to afford the crude product. The crude product was purified by flash column chromatography over silica gel, eluting with PE/ME (5:1 to 2:1) to give the product.

[0220] Mass spectrometry and ¹H NMR analyses confirmed that PP IX di-PEG₆₀₀ was obtained as the major product.

[0221] PP IX-mono-L-Alanine




[0222] A mixture of Protoporphyrin IX (500 mg, 0.89 mmol) and N-Hydroxysuccinimide (307 mg, 2.67 mmol) in DMF (50 ml) was stirred at 0°C for 1 h. A solution of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimidehydrochloride (512 mg, 2.67 mmol) in DMF (10 ml) was then added to the reaction mixture and allowed to stir overnight at room temperature.

[0223] To a solution of L-Alanine in water, cooled to 0°C, was added triethylamine and di-tert-butyl dicarbonate. The reaction mixture was stirred at room temperature for 2 h. The pH was adjusted to about 6 by progressively adding a 10% w/v aqueous citric acid solution. The combined aqueous layers were extracted with ethyl acetate and the combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The product obtained (200 mg) was added to the previously obtained EDC-activated reaction mixture.

[0224] The mixture was adjusted to pH 9 with 10% HCl and concentrated to afford a crude product. The crude product was dissolved in methanol (50 ml) and filtered. The filtrate was slowly dripped into methyl tert-butyl ether (200 ml) at 0°C and filtered to obtain PP IX-mono-L-Analine as a mixture containing mainly the two L-Alanine mono-substituted PP IX compounds.

[0225] Mass spectrometry and ¹H NMR analyses confirmed that PP IX-mono-L-Alanine was obtained as the major product.

[0226] PP IX-mono-L-Valine



[0227] A mixture of Protoporphyrin IX (500 mg, 0.89 mmol) and N-Hydroxysuccinimide (307 mg, 2.67 mmol) in DMF (50 ml) was stirred at 0°C for 1 h. A solution of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimidehydrochloride (512 mg, 2.67 mmol) in DMF (10 ml) was then added to the reaction mixture and allowed to stir overnight at room temperature.


[0228] To a solution of L-Valine in water, cooled to 0°C, was added triethylamine and di-tert-butyl dicarbonate. The reaction mixture was stirred at room temperature for 2 h. The pH was adjusted to about 6 by progressively adding a 10% w/v aqueous citric acid solution. The combined aqueous layers were extracted with ethyl acetate and the combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The product obtained (200 mg) was added to the previously obtained EDC-activated reaction mixture.

[0229] The mixture was adjusted to pH 9 with 10% HCl and concentrated to afford a crude product. The crude product was dissolved in methanol (50 ml) and filtered. The filtrate was

slowly dripped into methyl tert-butyl ether (200 ml) at 0°C and filtered to obtain PP IX-mono-L-Valine as a mixture containing mainly the two L-Valine mono-substituted PP IX compounds.

[0230] Mass spectrometry and ^1H NMR analyses confirmed that PP IX-mono-L-Valine was obtained as the major product.

[0231] PP IX-mono-glycine



[0232] A mixture of Protoporphyrin IX (500 mg, 0.89 mmol) and N-Hydroxysuccinimide (307 mg, 2.67 mmol) in DMF (50 ml) was stirred at 0°C for 1 h. A solution of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimidehydrochloride (512 mg, 2.67 mmol) in DMF (10 ml) was then added to the reaction mixture and allowed to stir overnight at room temperature.

[0233] To a solution of Glycine in water, cooled to 0°C, was added triethylamine and di-tert-butyl dicarbonate. The reaction mixture was stirred at room temperature for 2 h. The pH was adjusted to about 6 by progressively adding a 10% w/v aqueous citric acid solution. The combined aqueous layers were extracted with ethyl acetate and the combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The product obtained (200 mg) was added to the previously obtained EDC-activated reaction mixture.

[0234] The mixture was adjusted to pH 9 with 10% HCl and concentrated to afford a crude product. The crude product was dissolved in methanol (50 ml) and filtered. The filtrate was slowly dripped into methyl tert-butyl ether (200 ml) at 0°C and filtered to obtain PP IX-mono-Glycine as a mixture containing mainly the two glycine mono-substituted PP IX compounds.

[0235] Mass spectrometry and ^1H NMR analyses confirmed that PP IX-mono-L-Glycine was obtained as the major product.

Biological activity

Example 1

[0236] In this example, control of the gram-negative bacterial plant pathogen *Pseudomonas syringae* pv. *tabaci* with PP IX and modified PP IX was assessed, with and without chelating agents. Treatments were prepared in Phosphate Buffered Saline (PBS) in 96 well plates at desired concentrations. A bacterial suspension was inoculated into the PBS and incubated at 28°C in the dark for 30 minutes. After 30 minutes, the 96 well plate was placed under illumination for 1 hour (at 21°C). Following illumination, bacterial suspensions were serially diluted and 10 μL of each dilution is spread uniformly on Tryptic Soy Agar (TSA) plates and placed in the dark in an incubator at 28°C for 48 hours. After 48 hours, bacterial colonies were counted, and the results were log transformed (log colony forming units (CFU)/mL). The relative inactivation was determined by taking the difference between logCFU(PBS control) and logCFU(treatments). Sample Illumination was provided by LED lights (Heliospectra RX30) emitting about 1000 $\mu\text{mol}/\text{m}^2/\text{s}$ photosynthetically active radiation (PAR). The results are summarized in Table 1 below.

Table 1: Effect of 10 μM PP IX and derivatives on *Pseudomonas syringae*

Compound	log CFU/ml
PBS (control)	8.7
10 μM PP IX disodium salt	7.4
10 μM (PP IX-mono-DMAE : PP IX-di-DMAE – 50:50)	8.7
10 μM (PP IX-mono-DMAE : PP IX-di-DMAE – 20:80)	5.5
10 μM PP IX disodium salt + 5mM NaEDTA	3.8
10 μM (PP IX-mono-DMAE : PP IX-di-DMAE – 50:50) + 5mM NaEDTA	3.1
10 μM (PP IX-mono-DMAE : PP IX-di-DMAE – 20:80) + 5mM NaEDTA	0.0

Example 2

[0237] In this example, control of dollar spot fungus (*Sclerotinia homoeocarpa*) with PP IX and modified PP IX was assessed. Treatments were prepared in Phosphate Buffered Saline (PBS) in 24 well plates (in duplicates for light vs. dark incubation) at desired concentrations.

Then, a 5mm diameter plug of a *Sclerotinia homoeocarpa* isolate (3 isolates total tested) was inoculated into the PBS and incubated at 21°C in the dark for 2 hours. After 2 hours, one of the 24 well plates (with isolates in triplicate) was left in the dark and one 24 well plate was placed under illumination for 1 hour (all at 21°C). Following illumination, fungal plugs were removed from PBS, blotted dry on sterile filter paper and transferred to non-amended Potato Dextrose Agar (PDA). Radial growth of the fungus was monitored daily until the growth of *S. homoeocarpa* reaches the edge of the Petri-dish. Illumination was provided by LED lights emitting about 1000 $\mu\text{mol}/\text{m}^2/\text{s}$ photosynthetically active radiation (PAR). The results are summarized at Tables 2A and 2B below.

Table 2A: Results in Dark (no light exposure)

Treatment ¹	Mean Radial Growth ^{2,3}	% Inhibition ⁴
PBS (control)	10.1	-
31 μM Protoporphyrin IX disodium salt + 0.21% BrijO10:ArlasolveDMI (1:4)	10.5	-3.8
31 μM (PP IX-mono-DMAE : PP IX-di-DMAE – 20:80) + 0.21% BrijO10:ArlasolveDMI (1:4)	0.0	100.0
31 μM (PP IX-mono-DMAE : PP IX-di-DMAE – 50:50) + 0.21% BrijO10:ArlasolveDMI (1:4)	0.0	100.0
0.21% BrijO10:ArlasolveDMI (1:4)	10.3	-1.6

Notes on above table:

¹ Treatments were prepared in Phosphate Buffered Saline (PBS), incubated on shaker (200 rpm) for 2 hours in the dark, then kept in dark for 1 hour with no shaking.

² Means were calculated based on 3 fungal isolates replicated 3 times, with 2 measurements per replicate (18 total measurements)

³ Means represent growth that occurred between 24 and 48 hours of incubation at 21°C

⁴ % Inhibition calculated relative to non-amended control

Table 2B: Results in Light (exposed to light for 1 hour)

Treatment ¹	Mean Radial Growth ^{2,3}	% Inhibition ⁴
PBS (control)	10.2	-
31 μM Protoporphyrin IX disodium salt + 0.21% BrijO10:ArlasolveDMI (1:4)	0.0	100.0
31 μM (PP IX-mono-DMAE : PP IX-di-DMAE – 20:80) + 0.21% BrijO10:ArlasolveDMI (1:4)	0.0	100.0

31 μ M (PP IX-mono-DMAE : PP IX-di-DMAE – 50:50) + 0.21% BrijO10:ArlasolveDMI (1:4)	0.0	100.0
0.21% BrijO10:ArlasolveDMI (1:4)	6.1	40.2

Notes on above table:

¹ Treatments were prepared in Phosphate Buffered Saline (PBS), incubated on shaker (200 rpm) for 2 hours in the dark, then exposed to light (Helios, 1000 PAR) for 1 hour.

² Means were calculated based on 3 fungal isolates replicated 3 times, with 2 measurements per replicate (18 total measurements)

³ Means represent growth that occurred between 24 and 48 hours of incubation at 21°C

⁴% Inhibition calculated relative to non-amended control

Example 3

[0238] Control of the fungal plant pathogen *Colletotrichum orbiculare* ATC20767 (Cgm) on the host plant *Nicotiana benthamiana* following treatment with modified PP IX compounds was assessed. Treatments were applied to *N. benthamiana* plants approximately 2 h prior to inoculation with a spore suspension of Cgm. Plants were then exposed to light for a 24-hour period followed by dark incubation until disease symptoms were evident on the water treated control plants. Once disease symptoms were evident, lesions were counted, and leaf area measured in order to determine the number of lesions/cm² leaf area. Four replicate plants were used per treatment and plants were randomized under the light source. Illumination is provided by LED lights emitting about 180 μ mol/m²/s photosynthetically active radiation (PAR). The results are shown below.

Table 3: Effect of modified PP IX compounds on *Colletotrichum orbiculare*.


Treatment	% inhibition
untreated control	0
0.05% (PP IX-mono-DMAE : PP IX-di-DMAE – 20:80)	93
0.05% (PP IX-mono-DMAE : PP IX-di-DMAE – 50:50)	89
0.05% PP IX-mono-PEG ₆₀₀	56
0.05% PP IX-mono-L-valine	50
0.05% PP IX-mono-glycine	35

[0239] All publications, patents, and patent documents cited herein above are incorporated by reference herein, as though individually incorporated by reference. The compounds, compositions, methods and uses described herein have been described with reference to various embodiments and techniques. However, one skilled in the art will understand that

many variations and modifications can be made while remaining within the spirit and scope of the appended claims.

CLAIMS

1. A compound of Formula I:



Formula I

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

Z^1 and Z^2 are each independently OR^1 or NR^2R^3 ;

each R^1 , R^2 and R^3 is independently H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, or substituted alkynyl, wherein:

if Z^1 and Z^2 are both OR^1 then at least one R^1 is not H,

if Z^1 and Z^2 are both NR^2R^3 then at least one R^3 is not H, and

if one of Z^1 and Z^2 is OR^1 and the other one of Z^1 and Z^2 is NR^2R^3 , then at least one of R^1 and R^3 is not H;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

----- is a single bond or a double bond;

===== is a single bond or a double bond; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted aryl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more -X, $-R^B$, $-O^-$, $=O$, $-OR^B$, $-SR^B$, $-S^-$, $-NR^B_2$, $Si(R^C)_3$, $-N^+R^B_3$, $-NR^B-(Alk)-NR^B_2$, $-NR^B-(Alk)-N^+R^B_3$, $-NR^B-(Alk)-OR^B$, $-NR^B-(Alk)-OP(=O)(OR^B)(O^-)$, $-NR^B-(Alk)-OP(=O)(OR^B)_2$, $-NR^B-(Alk)-Si(R^C)_3$, $-NR^B-(Alk)-SR^B$, $-O-(Alk)-NR^B_2$, $-O-(Alk)-N^+R^B_3$, $-O-(Alk)-OR^B$, $-O-(Alk)-OP(=O)(OR^B)(O^-)$, $-O-(Alk)-OP(=O)(OR^B)_2$, $-O-(Alk)-Si(R^C)_3$, $-O-(Alk)-SR^B$, $=NR^B$, $-CX_3$, $-CN$, $-OCN$, $-SCN$, $-N=C=O$, $-NCS$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-NHC(=O)R^B$, $-OC(=O)R^B$, $-NHC(=O)NR^B_2$, $-S(=O)_2^-$, $-S(=O)_2OH$, $-S(=O)_2R^B$, $-OS(=O)_2OR^B$, $-S(=O)_2NR^B_2$, $-S(=O)R^B$, $-OP(=O)(OR^B)(O^-)$,

-OP(=O)(OR^B)₂, -P(=O)(OR^B)₂, -P(=O)(O⁻)₂, -P(=O)(OH)₂, -P(O)(OR^B)(O⁻), -C(=O)R^B, -C(=O)X, -C(S)R^B, -C(O)OR^B, -C(O)O⁻, -C(S)OR^B, -C(O)SR^B, -C(S)SR^B, -C(O)NR^B₂, -C(S)NR^B₂ or -C(=NR^B)NR^B₂;

each X is independently a halogen: F, Cl, Br or I;

each R^B is independently H, alkyl, aryl, arylalkyl, a heterocycle, an alkyloxy group such as poly(ethyleneoxy), PEG or poly(methyleneoxy), a capped poly(ethyleneoxy), capped PEG or capped polymethyleneoxy, or a protecting group;

the capped poly(ethyleneoxy), capped PEG and capped poly(methyleneoxy) groups being each independently capped with alkyl, aryl, arylalkyl, alkenyl, alkynyl, CO(alkyl), CO(aryl), CO(arylalkyl), CO(alkenyl) or CO(alkynyl);

each R^C is independently alkyl, aryl, arylalkyl, O(alkyl), O(aryl), O(arylalkyl), or O(tri-substituted silyl);

each tri-substituted silyl is independently substituted with three functional groups selected from alkyl, alkenyl, alkynyl, aryl and arylalkyl; and

each Alk is independently alkylene, alkenylene, or alkynylene.

2. The compound of claim 1, wherein Z¹ = Z² = NR²R³.

3. The compound of claim 1, wherein:

Z¹ is NR²R³ and Z² is OH, or Z¹ is OH and Z² is NR²R³.

4. The compound of any one of claims 1 to 3, wherein R³ is alkyl or substituted alkyl.

5. The compound of any one of claims 1 to 4, or an agriculturally acceptable salt thereof, wherein:

===== is a double bond.

6. The compound of any one of claims 1 to 5, or an agriculturally acceptable salt thereof, wherein:

===== is a double bond.

7. The compound of any one of claims 1 to 6, or an agriculturally acceptable salt thereof, wherein:

R^a, R^c, R^e and R^f are methyl; and


R^b and R^d are vinyl.

8. The compound of any one of claims 1 to 7, or an agriculturally acceptable salt thereof, wherein M is 2H.

9. The compound of any one of claims 1 to 7, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

10. The compound of any one of claims 1 to 7, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II).

11. A compound of Formula I:



Formula I

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z¹ and Z² is OR¹; and

the other one of Z¹ and Z² is NR²R³, NR²-(CH₂)_n-NR⁴R⁵, NR²-(CH₂)_n-N⁺R⁴R⁵R⁶ Y⁻, NR²-(CH₂)_n-O(PO₃H)⁻ W⁺, NR²-(CH₂)_n-Si(R⁷)₃, NR²-(CH₂)_n-SR⁸, NR²-(CH₂)_n-NR⁴-(CH₂)_p-NR⁹R¹⁰, NR²-(CH₂)_n-NR⁴-(CH₂)_p-N⁺R⁹R¹⁰R¹¹ Y⁻, NR²-(CH₂)_n-NR⁴-(CH₂)_p-O(PO₃H)⁻ W⁺, NR²-(CH₂)_n-NR⁴-(CH₂)_p-Si(R⁷)₃, NR²-(CH₂)_n-NR⁴-(CH₂)_p-SR⁸, O(CH₂)_n-NR⁴R⁵, O(CH₂)_n-N⁺R⁴R⁵R⁶ Y⁻, O(CH₂)_n-O(PO₃H)⁻ W⁺, O(CH₂)_n-Si(R⁷)₃, O(CH₂)_n-SR⁸, O(CH₂)_n-NR⁴-(CH₂)_p-NR⁹R¹⁰, O(CH₂)_n-NR⁴-(CH₂)_p-N⁺R⁹R¹⁰R¹¹ Y⁻, O(CH₂)_n-NR⁴-(CH₂)_p-O(PO₃H)⁻ W⁺ or O(CH₂)_n-NR⁴-(CH₂)_p-Si(R⁷)₃;

or

Z^1 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$, $NR^2-(CH_2)_n-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-N^+R^9R^{10}R^{11} Y^-$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-Si(R^7)_3$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-SR^8$, $O(CH_2)_n-NR^4R^5$, $O(CH_2)_n-N^+R^4R^5R^6 Y^-$, $O(CH_2)_n-O(PO_3H)^- W^+$, $O(CH_2)_n-Si(R^7)_3$, $O(CH_2)_n-SR^8$, $O(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$, $O(CH_2)_n-NR^4-(CH_2)_p-N^+R^9R^{10}R^{11} Y^-$, $O(CH_2)_n-NR^4-(CH_2)_p-O(PO_3H)^- W^+$ or $O(CH_2)_n-NR^4-(CH_2)_p-Si(R^7)_3$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

R^3 is alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

each R^4 , R^6 , R^8 , R^9 , R^{10} and R^{11} is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^5 is alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^7 is alkyl, O(alkyl) or O(tri-substituted silyl);

R^{13} is H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

W^+ is an agriculturally acceptable cation;

Y^- is an agriculturally acceptable anion;

n is an integer selected from 1 to 16;

p is an integer selected from 1 to 16;

m is an integer selected from 1 to 100;

q is an integer selected from 0 to 16;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl;

----- is a single bond or a double bond;

===== is a single bond or a double bond; and

M is 2H or a metal species,

wherein each substituted alkyl, substituted aryl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

12. The compound of claim 11, or an agriculturally acceptable salt thereof, wherein:
===== is a double bond.

13. The compound of claims 11 or 12, or an agriculturally acceptable salt thereof,
wherein:
===== is a double bond.

14. The compound of any one of claims 11 to 13, or an agriculturally acceptable salt
thereof, wherein:
each R^a, R^b, R^c, R^d, R^e and R^f is, independently, alkyl or alkenyl.

15. The compound of claim 14, or an agriculturally acceptable salt thereof, wherein:
R^a, R^c, R^e and R^f are methyl; and
R^b and R^d are vinyl.

16. The compound of any one of claims 11 to 15, or an agriculturally acceptable salt
thereof, wherein M is 2H.

17. The compound of any one of claims 11 to 15, or an agriculturally acceptable salt
thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II),
Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

18. The compound of any one of claims 11 to 15, or an agriculturally acceptable salt
thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II),
Fe(II) and Mn(II).

19. The compound of any one of claims 11 to 18, or an agriculturally acceptable salt
thereof, wherein R¹ is H.

20. The compound of any one of claims 11 to 19, or an agriculturally acceptable salt
thereof, wherein R² is H.

21. The compound of any one of claims 11 to 20, or an agriculturally acceptable salt
thereof, wherein R³ is alkyl.

22. The compound of any one of claims 11 to 21, or an agriculturally acceptable salt thereof, wherein R^4 is H or alkyl.

23. The compound of any one of claims 11 to 22, or an agriculturally acceptable salt thereof, wherein R^5 is alkyl.

24. The compound of any one of claims 11 to 23, or an agriculturally acceptable salt thereof, wherein R^6 is alkyl.

25. The compound of any one of claims 11 to 24, or an agriculturally acceptable salt thereof, wherein R^7 is O(tri-substituted silyl).

26. The compound of any one of claims 11 to 25, or an agriculturally acceptable salt thereof, wherein R^8 is H or alkyl.

27. The compound of any one of claims 11 to 26, or an agriculturally acceptable salt thereof, wherein R^9 is alkyl.

28. The compound of any one of claims 11 to 27, or an agriculturally acceptable salt thereof, wherein R^{10} is alkyl.

29. The compound of any one of claims 11 to 28, or an agriculturally acceptable salt thereof, wherein R^{11} is alkyl.

30. The compound of any one of claims 11 to 29, or an agriculturally acceptable salt thereof, wherein R^{13} is H, alkyl, alkenyl, CO(alkyl) or CO(alkenyl).

31. The compound of any one of claims 11 to 30, or an agriculturally acceptable salt thereof, wherein W^+ is selected from the group consisting of sodium, potassium, magnesium and ammonium cations.

32. The compound of any one of claims 11 to 31, or an agriculturally acceptable salt thereof, wherein Y^- is selected from the group consisting of chloride, bromide, phosphate, dimethylphosphate, methylsulfate, ethylsulfate, acetate and lactate.

33. The compound of any one of claims 11 to 32, or an agriculturally acceptable salt thereof, wherein n is an integer selected from 2 to 4.

34. The compound of any one of claims 11 to 33, or an agriculturally acceptable salt thereof, wherein p is an integer selected from 2 to 4.

35. The compound of any one of claims 11 to 34, or an agriculturally acceptable salt thereof, wherein m is an integer selected from 1 to 20.

36. The compound of any one of claims 11 to 35, or an agriculturally acceptable salt thereof, wherein the substituted alkyl, substituted aryl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more F, Cl, Br and I.

37. The compound of any one of claims 11 to 36, or an agriculturally acceptable salt thereof, wherein:

Z^1 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$,
 $NR^2-(CH_2)_n-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$,
 $NR^2-(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-N^+R^9R^{10}R^{11} Y^-$,
 $NR^2-(CH_2)_n-NR^4-(CH_2)_p-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-NR^4-(CH_2)_p-Si(R^7)_3$,
 $NR^2-(CH_2)_n-NR^4-(CH_2)_p-SR^8$, $O(CH_2)_n-NR^4R^5$, $O(CH_2)_n-N^+R^4R^5R^6 Y^-$,
 $O(CH_2)_n-O(PO_3H)^- W^+$, $O(CH_2)_n-Si(R^7)_3$, $O(CH_2)_n-SR^8$, $O(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$,
 $O(CH_2)_n-NR^4-(CH_2)_p-N^+R^9R^{10}R^{11} Y^-$, $O(CH_2)_n-NR^4-(CH_2)_p-O(PO_3H)^- W^+$ or
 $O(CH_2)_n-NR^4-(CH_2)_p-Si(R^7)_3$; and
 $Z^2 = Z^1$.


38. The compound of any one of claims 11 to 36, or an agriculturally acceptable salt thereof, wherein:

one of Z^1 and Z^2 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$, $NR^2-(CH_2)_n-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $NR^2-(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$; and
the other one of Z^1 and Z^2 is OR^1 ;
or
 Z^1 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$, $NR^2-(CH_2)_n-O(PO_3H)^- W^+$,
 $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $NR^2-(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$; and
 $Z^2 = Z^1$.

39. The compound of claim 38, or an agriculturally acceptable salt thereof, wherein:
 one of Z^1 and Z^2 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$, $NR^2-(CH_2)_n-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $NR^2-(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$; and
 the other one of Z^1 and Z^2 is OR^1 .

40. The compound of claim 38, or an agriculturally acceptable salt thereof, wherein:
 Z^1 is NR^2R^3 , $NR^2-(CH_2)_n-NR^4R^5$, $NR^2-(CH_2)_n-N^+R^4R^5R^6 Y^-$, $NR^2-(CH_2)_n-O(PO_3H)^- W^+$, $NR^2-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $NR^2-(CH_2)_n-NR^4-(CH_2)_p-NR^9R^{10}$; and
 $Z^2 = Z^1$.

41. A compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant,
 wherein:

one of Z^1 and Z^2 is NR^2R^3 ; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2R^3$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

R^3 is alkyl or substituted alkyl;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N₃.

42. The compound of claim 41, or an agriculturally acceptable salt thereof, wherein: each R^a, R^b, R^c, R^d, R^e and R^f is, independently, alkyl or alkenyl.

43. The compound of claim 41 or 42, or an agriculturally acceptable salt thereof, wherein:

R^a, R^c, R^e and R^f are methyl; and

R^b and R^d are vinyl.

44. The compound of any one of claims 41 to 43, or an agriculturally acceptable salt thereof, wherein M is 2H.

45. The compound of any one of claims 41 to 43, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

46. The compound of any one of claims 41 to 43, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II).

47. The compound of any one of claims 41 to 46, or an agriculturally acceptable salt thereof, wherein R¹ is H.

48. The compound of any one of claims 41 to 47, or an agriculturally acceptable salt thereof, wherein R² is H.

49. The compound of any one of claims 41 to 48, or an agriculturally acceptable salt thereof, wherein R³ is alkyl.

50. The compound of claim 49, or an agriculturally acceptable salt thereof, wherein R³ is (C₁-C₁₂)alkyl.

51. The compound of any one of claims 41 to 50, or an agriculturally acceptable salt thereof, wherein:

one of Z^1 and Z^2 is NR^2R^3 ; and


the other one of Z^1 and Z^2 is OR^1 .


52. The compound of any one of claims 41 to 50, or an agriculturally acceptable salt thereof, wherein:

$Z^1 = NR^2R^3$; and

$Z^2 = Z^1$.

53. The compound of claim 41, that is:












or an agriculturally acceptable salt thereof.

54. The compound of claim 41, that is:

100







101




or an agriculturally acceptable salt thereof.

55. The compound of claim 41, that is:






103



or an agriculturally acceptable salt thereof.

56. A compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant,
wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-NR^4R^5$ or $O-(CH_2)_n-NR^4R^5$; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2)_n-NR^4R^5$ or $O-(CH_2)_n-NR^4R^5$; and

$Z^2 = Z^1$;

R^5 is alkyl, substituted alkyl or $-(CH_2)_p-NR^9R^{10}$;

each R^1 , R^2 , R^4 , R^9 and R^{10} is, independently, H, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

p is an integer selected from 1 to 16;

each R^a, R^b, R^c, R^d, R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N₃.

57. The compound of claim 56, or an agriculturally acceptable salt thereof, wherein: each R^a, R^b, R^c, R^d, R^e and R^f is, independently, alkyl or alkenyl.

58. The compound of claim 56 or 57, or an agriculturally acceptable salt thereof, wherein:

R^a, R^c, R^e and R^f are methyl; and

R^b and R^d are vinyl.

59. The compound of any one of claims 56 to 58, or an agriculturally acceptable salt thereof, wherein M is 2H.

60. The compound of any one of claims 56 to 58, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

61. The compound of any one of claims 56 to 58, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II).

62. The compound of any one of claims 56 to 61, or an agriculturally acceptable salt thereof, wherein R¹ is H.

63. The compound of any one of claims 56 to 62, or an agriculturally acceptable salt thereof, wherein R² is H.

64. The compound of any one of claims 56 to 63, or an agriculturally acceptable salt thereof, wherein R⁴ is H or alkyl.

65. The compound of any one of claims 56 to 64, or an agriculturally acceptable salt thereof, wherein n is an integer selected from 2 to 4.

66. The compound of any one of claims 56 to 65, or an agriculturally acceptable salt thereof, wherein R^5 is alkyl.

67. The compound of any one of claims 56 to 65, or an agriculturally acceptable salt thereof, wherein R^5 is $-(CH_2)_p-NR^9R^{10}$.

68. The compound of claim 67, or an agriculturally acceptable salt thereof, wherein R^9 is alkyl.

69. The compound of claim 67 or 68, or an agriculturally acceptable salt thereof, wherein R^{10} is alkyl.

70. The compound of any one of claims 67 to 69, or an agriculturally acceptable salt thereof, wherein p is an integer selected from 2 to 4.

71. The compound of any one of claims 56 to 70, or an agriculturally acceptable salt thereof, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-NR^4R^5$; and

the other one of Z^1 and Z^2 is OR^1 .


72. The compound of any one of claims 56 to 70, or an agriculturally acceptable salt thereof, wherein:

$Z^1 = NR^2-(CH_2)_n-NR^4R^5$; and

$Z^2 = Z^1$.

73. The compound of claim 56, that is:


106



or an agriculturally acceptable salt thereof.

74. The compound of claim 56, that is:







107



or an agriculturally acceptable salt thereof.

75. The compound of claim 56, that is:


108



or an agriculturally acceptable salt thereof.

109

76. A compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $NR^2-(CH_2)_n-Si(R^7)_3$, $O-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $O-(CH_2)_n-SR^8$; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = NR^2-(CH_2)_n-Si(R^7)_3$, $O-(CH_2)_n-Si(R^7)_3$, $NR^2-(CH_2)_n-SR^8$ or $O-(CH_2)_n-SR^8$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

R^7 is alkyl, O(alkyl) or O(trisubstituted silyl);

R^8 is H, alkyl, substituted alkyl or $-(CH_2)_q-(CH_2CH_2O)_m-R^{13}$;

R^{13} is H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

n is an integer selected from 1 to 16;

m is an integer selected from 1 to 100;

q is an integer selected from 0 to 16;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

77. The compound of claim 76, or an agriculturally acceptable salt thereof, wherein: each R^a , R^b , R^c , R^d , R^e and R^f is, independently, alkyl or alkenyl.

78. The compound of claim 76 or 77, or an agriculturally acceptable salt thereof, wherein:

R^a , R^c , R^e and R^f are methyl; and

R^b and R^d are vinyl.

79. The compound of any one of claims 76 to 78, or an agriculturally acceptable salt thereof, wherein M is 2H.

80. The compound of any one of claims 76 to 78, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

81. The compound of any one of claims 76 to 78, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II).

82. The compound of any one of claims 76 to 81, or an agriculturally acceptable salt thereof, wherein R^1 is H.


83. The compound of any one of claims 76 to 82, or an agriculturally acceptable salt thereof, wherein R^2 is H.

84. The compound of any one of claims 76 to 83, or an agriculturally acceptable salt thereof, wherein n is an integer selected from 2 to 4.

85. The compound of any one of claims 76 to 84, or an agriculturally acceptable salt thereof, wherein R^7 is O(tri-substituted silyl).


86. The compound of claims 76 to 84, or an agriculturally acceptable salt thereof, wherein R^8 is $-(CH_2CH_2O)_m-R^{13}$.

112




or an agriculturally acceptable salt thereof,
wherein m is an integer selected from 4 to 15.

92. The compound of claim 76, that is:





113




or an agriculturally acceptable salt thereof.

93. The compound of claim 76, that is:



94. A compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)}_2$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)}_2$, $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$; and

the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = \text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)}_2$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)}_2$, $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

W^+ is an agriculturally acceptable cation;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

95. The compound of claim 94, or an agriculturally acceptable salt thereof, wherein:

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, alkyl or alkenyl.

115

96. The compound of claim 94 or 95, or an agriculturally acceptable salt thereof, wherein:

R^a , R^c , R^e and R^f are methyl; and

R^b and R^d are vinyl.

97. The compound of any one of claims 94 to 96, or an agriculturally acceptable salt thereof, wherein M is 2H.

98. The compound of any one of claims 94 to 96, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

99. The compound of any one of claims 94 to 96, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II).

100. The compound of any one of claims 94 to 99, or an agriculturally acceptable salt thereof, wherein R^1 is H.

101. The compound of any one of claims 94 to 100, or an agriculturally acceptable salt thereof, wherein R^2 is H.

102. The compound of any one of claims 94 to 101, or an agriculturally acceptable salt thereof, wherein n is an integer selected from 2 to 4.

103. The compound of any one of claims 94 to 102, or an agriculturally acceptable salt thereof, wherein W^+ is selected from the group consisting of sodium, potassium, magnesium and ammonium cations.

104. The compound of claim 103, or an agriculturally acceptable salt thereof, wherein W^+ is an ammonium cation.

105. The compound of any one of claims 94 to 104, or an agriculturally acceptable salt thereof, wherein:


116

one of Z^1 and Z^2 is $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)}_2$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)}_2$,
 $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$; and
 the other one of Z^1 and Z^2 is OR^1 .

106. The compound of any one of claims 94 to 104, or an agriculturally acceptable salt thereof, wherein:


$Z^1 = \text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)}_2$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)}_2$,
 $\text{NR}_2-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$ or $\text{O}-(\text{CH}_2)_n\text{-OP=O(OH)O}^- \text{W}^+$; and
 $Z^2 = Z^1$.

107. The compound of claim 94, that is:



117

108. A compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is $\text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{R}^5\text{R}^{6+} \text{Y}^-$ or $\text{O-(CH}_2\text{)}_n\text{-NR}^4\text{R}^5\text{R}^{6+} \text{Y}^-$; and the other one of Z^1 and Z^2 is OR^1 ;

or

$Z^1 = \text{NR}^2\text{-(CH}_2\text{)}_n\text{-NR}^4\text{R}^5\text{R}^{6+} \text{Y}^-$ or $\text{O-(CH}_2\text{)}_n\text{-NR}^4\text{R}^5\text{R}^{6+} \text{Y}^-$; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

each R^4 , R^5 and R^6 is, independently, alkyl or substituted alkyl;

n is an integer selected from 1 to 16;

Y^- is an agriculturally acceptable anion;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

109. The compound of claim 108, or an agriculturally acceptable salt thereof, wherein:

R^a , R^c , R^e and R^f are methyl; and

R^b and R^d are vinyl.

118

110. The compound of any one of claims 108 to 109, or an agriculturally acceptable salt thereof, wherein M is 2H.

111. The compound of any one of claims 108 to 109, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

112. The compound of any one of claims 108 to 109, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II).

113. The compound of any one of claims 108 to 112, or an agriculturally acceptable salt thereof, wherein R¹ is H.

114. The compound of any one of claims 108 to 113, or an agriculturally acceptable salt thereof, wherein R² is H.

115. The compound of any one of claims 108 to 114, or an agriculturally acceptable salt thereof, wherein R⁴, R⁵ and R⁶ are alkyl and R⁴ = R⁵ = R⁶.

116. The compound of any one of claims 108 to 115, or an agriculturally acceptable salt thereof, wherein Y⁻ is selected from the group consisting of chloride, bromide, phosphate, dimethylphosphate, methylsulfate, ethylsulfate, acetate, citrate, tartrate and lactate.

117. The compound of claim 116, wherein Y⁻ is acetate.

118. The compound of any one of claims 108 to 117, wherein n is an integer selected from 2 to 4.

119. The compound of any one of claims 108 to 118, or an agriculturally acceptable salt thereof, wherein:

one of Z¹ and Z² is NR²-(CH₂)_n-NR⁴R⁵R⁶⁺ Y⁻ or O-(CH₂)_n-NR⁴R⁵R⁶⁺ Y⁻; and
the other one of Z¹ and Z² is OR¹.


119

120. The compound of any one of claims 108 to 118, or an agriculturally acceptable salt thereof, wherein:

$Z^1 = \text{NR}^2-(\text{CH}_2)_n-\text{NR}^4\text{R}^5\text{R}^{6+} \text{Y}^-$ or $\text{O}-(\text{CH}_2)_n-\text{NR}^4\text{R}^5\text{R}^{6+} \text{Y}^-$; and

$Z^2 = Z^1$.


121. The compound of claim 108, that is:



wherein Y^- is an agriculturally acceptable anion selected from the group consisting of: chloride, bromide, phosphate, dimethylphosphate, methylsulfate, ethylsulfate, acetate, citrate, tartrate and lactate.

120

122. A compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z¹ and Z² is NR²-(CH₂CH₂O)_m-R¹³ or O-(CH₂CH₂O)_m-R¹³; and the other one of Z¹ and Z² is OR¹;

or

Z¹ = NR²-(CH₂CH₂O)_m-R¹³ or O-(CH₂CH₂O)_m-R¹³; and

Z² = Z¹;

each R¹ and R² is, independently, H, alkyl or substituted alkyl;

R¹³ is H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CO(alkyl), CO(substituted alkyl), CO(alkenyl), CO(substituted alkenyl), CO(alkynyl) or CO(substituted alkynyl);

m is an integer selected from 1 to 100;

each R^a, R^b, R^c, R^d, R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N₃.

123. The compound of claim 122, or an agriculturally acceptable salt thereof, wherein:

R^a, R^c, R^e and R^f are methyl; and

R^b and R^d are vinyl.

121

124. The compound of any one of claims 122 or 123, or an agriculturally acceptable salt thereof, wherein M is 2H.

125. The compound of any one of claims 122 or 123, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

126. The compound of any one of claims 122 or 123, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II).

127. The compound of any one of claims 122 to 126, or an agriculturally acceptable salt thereof, wherein R¹ is H.

128. The compound of any one of claims 122 to 127, or an agriculturally acceptable salt thereof, wherein R² is H.

129. The compound of any one of claims 122 to 128, or an agriculturally acceptable salt thereof, wherein R¹³ is alkyl, alkenyl, CO(alkyl) or CO(alkenyl).

130. The compound of any one of claims 122 to 129, or an agriculturally acceptable salt thereof, wherein m is an integer selected from 5 to 20.


131. The compound of any one of claims 122 to 130, or an agriculturally acceptable salt thereof, wherein:


one of Z¹ and Z² is NR²-(CH₂CH₂O)_m-R¹³ or O-(CH₂CH₂O)_m-R¹³; and
the other one of Z¹ and Z² is OR¹.

132. The compound of any one of claims 122 to 130, or an agriculturally acceptable salt thereof, wherein:

Z¹ = NR²-(CH₂CH₂O)_m-R¹³ or O-(CH₂CH₂O)_m-R¹³; and
Z² = Z¹.

133. The compound of claim 122, that is:





or an agriculturally acceptable salt thereof,

wherein m is an integer selected from 1 to 100; and

R^{14} is alkyl, alkenyl, alkynyl, substituted alkyl, substituted alkenyl or substituted alkynyl.


134. The compound of claim 133, or an agriculturally acceptable salt thereof, wherein m is an integer selected from 5 to 20.

135. The compound of claim 133, or an agriculturally acceptable salt thereof, wherein $(CH_2CH_2O)_mCOR^{14}$ is PEG₆₀₀-oleate, PEG₄₀₀-oleate, PEG₆₀₀-allyl or PEG₄₀₀-allyl.

136. The compound of any one of claims 122 to 128, or an agriculturally acceptable salt thereof, wherein R^{13} is H.


137. The compound of claim 136, or an agriculturally acceptable salt thereof, wherein $(CH_2CH_2O)_mR^{13}$ is PEG₆₀₀ or PEG₄₀₀.

138. The compound of claim 122, that is:



125

139. A compound of Formula I-B1:



Formula I-B1

or an agriculturally acceptable salt thereof, for use in promoting the health of a plant, wherein:

one of Z^1 and Z^2 is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and

the other one of Z^1 and Z^2 is OR^1 ;

or

Z^1 is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and

$Z^2 = Z^1$;

each R^1 and R^2 is, independently, H, alkyl or substituted alkyl;

each R^a , R^b , R^c , R^d , R^e and R^f is, independently, H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl; and

M is 2H or a metal species,

wherein the substituted alkyl, substituted alkenyl and substituted alkynyl groups are, independently, substituted with one or more OH, F, Cl, Br, I, CN and N_3 .

140. The compound of claim 139, or an agriculturally acceptable salt thereof, wherein:

R^a , R^c , R^e and R^f are methyl; and

R^b and R^d are vinyl.

141. The compound of any one of claims 139 or 140, or an agriculturally acceptable salt thereof, wherein M is 2H.

126

142. The compound of any one of claims 139 or 140, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Mg(II), Zn(II), Pd(II), Sn(IV), Al(III), Pt(II), Si(IV), Ge(IV), Ga(III) and In(III).

143. The compound of any one of claims 139 or 140, or an agriculturally acceptable salt thereof, wherein M is a metal species selected from the group consisting of Cu(II), Co(II), Fe(II) and Mn(II).

144. The compound of any one of claims 139 to 143, or an agriculturally acceptable salt thereof, wherein R¹ is H.

145. The compound of any one of claims 139 to 144, or an agriculturally acceptable salt thereof, wherein R² is H.

146. The compound of any one of claims 139 to 145, or an agriculturally acceptable salt thereof, wherein:

one of Z¹ and Z² is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and
the other one of Z¹ and Z² is OR¹.


147. The compound of any one of claims 139 to 145, or an agriculturally acceptable salt thereof, wherein:

Z¹ is a natural amino acid attached to the compound by its amino group bonded to the alpha carbon; and
Z² = Z¹.

148. The compound of any one of claims 139 to 145, or an agriculturally acceptable salt thereof, wherein:

Z¹ is Glycine, L-Alanine, or L-Valine and Z² is OH;
Z² is Glycine, L-Alanine, or L-Valine and Z¹ is OH; or
Z¹ is Glycine, L-Alanine or L-Valine and Z² = Z¹.


149. The compound of claim 139, that is:




or an agriculturally acceptable salt thereof.

150. The compound of claim 139, that is:

128





or an agriculturally acceptable salt thereof.

151. A composition for use in promoting the health of a plant, the composition comprising:

at least one compound as defined in any one of claims 1 to 150, or an agriculturally acceptable salt thereof; and
a carrier fluid.

152. The composition of claim 151, further comprising a chelating agent.

153. The composition of claim 152, wherein the chelating agent comprises at least one carboxylic acid group or an agriculturally acceptable salt thereof.

154. The composition of claim 152 or 153, wherein the chelating agent comprises at least two carboxylic acid groups or agriculturally acceptable salts thereof.

155. The composition of any one of claims 152 to 154, wherein the chelating agent comprises at least one amino group or an agriculturally acceptable salt thereof.

156. The composition of any one of claims 152 to 155, wherein the chelating agent comprises an amino-carboxylic acid compound or an agriculturally acceptable salt thereof.

157. The composition of claim 156, wherein the amino-carboxylic acid compound comprises an amino polycarboxylic acid compound or an agriculturally acceptable salt thereof.

158. The composition of any one of claims 152 to 157, wherein the chelating agent is selected from the group consisting of ethylenediaminetetraacetic acid (EDTA) or an agriculturally acceptable salt thereof, ethylenediamine-N,N'-disuccinic acid (EDDS) or an agriculturally acceptable salt thereof, iminodisuccinic acid (IDS) or an agriculturally acceptable salt thereof, and mixtures thereof.

159. The composition of any one of claims 152 to 157, wherein the chelating agent comprises a polyaspartic acid or an agriculturally acceptable salt thereof.

160. The composition of any one of claims 152 to 159, wherein the chelating agent is metallated.

161. The composition of any one of claims 152 to 159, wherein the chelating agent is metal-free.

162. The composition of any one of claims 151 to 161, further comprising a surfactant.

163. The composition of claim 162, wherein the surfactant is selected from the group consisting of an ethoxylated alcohol, a polymeric surfactant, a fatty acid ester, a poly(ethylene glycol), an ethoxylated alkyl alcohol, a monoglyceride, an alkyl monoglyceride and a mixture thereof.

164. The composition of claim 162 or 163, wherein the surfactant comprises a poly(ethylene glycol) of Formula $R^{15}-O-(CH_2CH_2O)_f-R^{16}$,

wherein:

each R^{15} and R^{16} is each, independently, H, alkyl, substituted alkyl, aryl, substituted aryl, CO(alkyl) or CO(substituted alkyl); and

f is an integer selected from 1 to 100;

wherein the substituted alkyl groups are, independently, substituted with one or more F, Cl, Br, I, hydroxy, alkenyl, CN and N_3 .

131

165. The composition of claim 164, wherein the substituted alkyl groups are methyl, vinyl or CO(methyl).

166. The composition of any one of claims 151 to 165, further comprising an oil selected from the group consisting of a mineral oil, an essential oil, a vegetable oil and a mixture thereof.

167. The composition of claim 166, wherein the oil comprises a vegetable oil selected from the group consisting of coconut oil, canola oil, soybean oil, rapeseed oil, sunflower oil, safflower oil, peanut oil, cottonseed oil, palm oil, rice bran oil and mixtures thereof.

168. The composition of claim 166 or 167, wherein the oil comprises a mineral oil selected from the group consisting of a paraffinic oil, a branched paraffinic oil, naphthenic oil, an aromatic oil and mixtures thereof.

169. The composition of any one of claims 166 to 168, wherein the oil comprises a poly-alpha-olefin (PAO).

170. The composition of any one of claims 151 to 169, wherein the carrier fluid comprises water.

171. The composition of any one of claims 151 to 170, wherein the carrier fluid comprises an oil-in-water emulsion.

172. A method for promoting the health of a plant, comprising:
applying to the plant the compound as defined in any one of claims 1 to 150 or an agriculturally acceptable salt thereof, or the composition as defined in any one of claims 151 to 171; and
exposing the plant to light.

173. The method of claim 172, wherein promoting the health of the plant comprises preventing or inhibiting growth of a microbial pathogen of the plant.

174. The method of claim 172, wherein promoting the health of the plant comprises inhibiting growth of a microbial pathogen of the plant.

175. The method of claim 173 or 174, wherein the microbial pathogen comprises a fungal pathogen.

176. The method of claim 175, wherein the fungal pathogen comprises at least one of *Botrytis cinerea*, *Alternaria solani* and *Sclerotinia homoeocarpa*.

177. The method of claim 173 or 174, wherein the microbial pathogen comprises a bacterial pathogen.

178. The method of claim 177, wherein the bacterial pathogen comprises Gram negative bacteria.

179. The method of claim 178, wherein the bacterial pathogen comprises at least one of *Erwinia amylovora*, *Xanthomonas axonopodis*, and *E. Coli*.

180. The method of claim 173 or 174, wherein the microbial pathogen comprises a virus, a viroid, a virus-like organism or a phytoplasma.

181. The method of claim 180, wherein the microbial pathogen comprises a virus.

182. The method of claim 172, wherein promoting the health of the plant comprises increasing resistance of the plant to one or more abiotic stress.

183. The method of claim 182, wherein the one or more abiotic stress is selected from the group consisting of cold stress, heat stress, water stress, transplant shock stress, low light stress, photooxidative stress, drought stress and salinity stress.

184. The method of claim 182 or 183, wherein applying the compound to the plant is performed at or before onset of the one or more abiotic stress.

185. The method of claim 182 or 183, wherein applying the compound to the plant is performed after onset of the abiotic stress.

133

186. The method of claim 172, wherein promoting the health of the plant comprises controlling a pest of the plant.

187. The method of claim 186, wherein the pest of the plant is selected from the group consisting of insects and insect larvae.

188. The method of claim 186 or 187, wherein applying the compound to the plant is performed at or before contamination of the plant by the pest.

189. The method of claim 186 or 187, wherein applying the compound to the plant is performed after contamination of the plant by the pest.

190. The method of any one of claims 172 to 189, wherein the plant is a non-woody crop plant, a woody plant or a turfgrass.

191. The method of claim 190, wherein the plant is a woody plant.

192. The method of claim 191, wherein the woody plant is a tree.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA2020/050197

A. CLASSIFICATION OF SUBJECT MATTER

IPC: *A01N 43/90* (2006.01), *A01P 1/00* (2006.01), *A01P 3/00* (2006.01), *A01P 7/04* (2006.01),*C07D 487/22* (2006.01)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: A01N 43/90 (2006.01), *A01P 1/00* (2006.01), *A01P 3/00* (2006.01), *A01P 7/04* (2006.01),
C07D 487/22 (2006.01)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used)

STN (structure searching with and without control terms relating to plants, plant growth)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2010080819, ECKERT et al., 15 July 2010 (15-07-2010)	1-191
A	JUNG et al., Toxic Tetrapyrrole Accumulation in Protoporphyrinogen IX Oxidase-Overexpressing Transgenic Rice Plants, <i>Plant Mol. Biol.</i> (2008), 67, p. 535-546.	1-191

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* "A" "D" "E" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance document cited by the applicant in the international application earlier application or patent but published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	"T" "X" "Y" "&"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
---	--	--------------------------	--

Date of the actual completion of the international search
2 March 2020 (02-03-2020)Date of mailing of the international search report
20 April 2020 (20-04-2020)Name and mailing address of the ISA/CA
Canadian Intellectual Property Office
Place du Portage I, C114 - 1st Floor, Box PCT
50 Victoria Street
Gatineau, Quebec K1A 0C9
Facsimile No.: 819-953-2476

Authorized officer

Daniel Black (819) 639-7540

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2020/050197

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
WO2010080819A1	15 July 2010 (15-07-2010)	AU2010203698A1	21 July 2011 (21-07-2011)
		AU2010203698A2	22 September 2011 (22-09-2011)
		AU2010203698B2	21 July 2016 (21-07-2016)
		AU2016204543A1	14 July 2016 (14-07-2016)
		AU2016204543B2	18 January 2018 (18-01-2018)
		CA2749082A1	15 July 2010 (15-07-2010)
		CN102405053A	04 April 2012 (04-04-2012)
		CN102405053B	29 July 2015 (29-07-2015)
		EP2381953A1	02 November 2011 (02-11-2011)
		EP2381953A4	29 June 2016 (29-06-2016)
		US2010184681A1	22 July 2010 (22-07-2010)
		US8303962B2	06 November 2012 (06-11-2012)
		US2011039761A1	17 February 2011 (17-02-2011)
		US8754039B2	17 June 2014 (17-06-2014)
		US2013108575A1	02 May 2013 (02-05-2013)
		US9072793B2	07 July 2015 (07-07-2015)
		US2014349917A1	27 November 2014 (27-11-2014)
		US9597407B2	21 March 2017 (21-03-2017)
		US2010184683A1	22 July 2010 (22-07-2010)
		US2010184684A1	22 July 2010 (22-07-2010)
		US2011039762A1	17 February 2011 (17-02-2011)
		US2011039763A1	17 February 2011 (17-02-2011)
		US2017266306A1	21 September 2017 (21-09-2017)
		WO2010080836A2	15 July 2010 (15-07-2010)
		WO2010080836A9	17 July 2014 (17-07-2014)