

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

08 November 2018 (08.11.2018)



(10) International Publication Number

WO 2018/204559 A1

(51) International Patent Classification:

C11D 1/37 (2006.01) C11D 11/00 (2006.01)

C11D 1/83 (2006.01) C11D 1/22 (2006.01)

C11D 3/20 (2006.01) C11D 1/29 (2006.01)

MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/US2018/030752

Published:

— with international search report (Art. 21(3))

(22) International Filing Date:

03 May 2018 (03.05.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

17169631.3 05 May 2017 (05.05.2017) EP

18154155.8 30 January 2018 (30.01.2018) EP

(71) Applicant: THE PROCTER & GAMBLE COMPANY

[US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).

(72) Inventors: BOUTIQUE, Jean-Pol; NV Procter & Gamble

Services Company SA, Temselaan 100, 1853 Strombeek-Bever (BE). DKIDAK, Aicha; NV Procter & Gamble Services Company SA, Temselaan 100, 1853 Strombeek-Bever (BE). MAHIEU, Bill, Karel; NV Procter & Gamble Services Company SA, Temselaan 100, 1853 Strombeek-Bever (BE). STENGER, Patrick, Christopher; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). GONZALES, Denis, Alfred; NV Procter & Gamble Services Company SA, Temselaan 100, 1853 Strombeek-Bever (BE).

(74) Agent: KREBS, Jay, A.; c/o The Procter & Gamble Com-

pany, Global IP Services, One Procter & Gamble Plaza, C9, Cincinnati, OH 45202 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

(54) Title: LAUNDRY DETERGENT COMPOSITIONS WITH IMPROVED GREASE REMOVAL

(57) Abstract: The need for a laundry detergent composition which provides improved grease removal both with and without pretreatment, while also providing improved viscosity, is met using a composition having a pH of from 6.0 to 8.9 and comprising anionic and nonionic surfactants and C4-C10 alkyl branched alcohol.



WO 2018/204559 A1

LAUNDRY DETERGENT COMPOSITIONS WITH IMPROVED GREASE REMOVAL

FIELD OF THE INVENTION

Laundry detergent compositions, especially liquid laundry detergent compositions for cleaning
5 grease stains.

BACKGROUND OF THE INVENTION

Laundry detergent compositions are formulated to provide good cleaning to fabrics: To keep white
fabrics white, and to keep coloured fabrics bright. The laundry detergent compositions are also
10 typically formulated to remove stains. A stain is a local discoloration that can be clearly
distinguished on the fabric it is found upon. Since they result in a discoloration that strongly
contrasts with the unstained fabrics, they are particularly noticeable on fabrics.

Grease-containing stains remain challenging to remove, especially at low temperatures. Various
organic solvents, including terpenes and terpene-like compounds, are rather well-known for use in
15 hard surface cleaners for their grease removal ability, as disclosed in EP0137616A1. Such cleaners
often contain 10%, or more, of a solvent such as d-limonene, together with a surfactant, especially
nonionic surfactants which are also well-known for their grease removal performance. EP0040882
also discloses hard surface cleaners comprising a mixture of benzyl alcohol, terpenes, surfactants
and other deterative ingredients. High composition pH also improves grease cleaning. However,
20 high pH can also damage fabrics and fabric colour, especially for delicate fabrics, and especially
during direct application, for instance during pretreatment.

Moreover, laundry detergent compositions are typically significantly diluted during the laundry
wash. Even pretreating grease-containing stains remains challenging, since the grease is typically
heavily impregnated into the fabric and the time for the laundry detergent composition to penetrate
25 the stain is short, especially considering the hydrophobic nature, and hence water-repellency, of
such grease-containing stains.

While the use of relatively high concentrations of solvents in heavy duty liquid laundry detergents
improves stain penetration during pretreatment, higher solvent levels have disadvantages such as
damaging delicate fabrics and affecting color retention on fabrics. Moreover, solvents typically
30 lower the viscosity of liquid laundry detergent compositions.

As such, a need remains for laundry compositions, especially liquid laundry detergent compositions which are suitable for use in pretreating, which provide improved removal of grease-containing stains while not damaging delicate fabrics. A further need remains for such compositions which provide improved removal of grease-containing stains, while not requiring high levels of the solvent, even during washing without pretreatment.

US4561991A relates to stain removal compositions which comprise mixtures of grease-cutting solvents and polyamines for removing stains comprising a mixture of grease and particulate matter from fabrics. WO 2012/117004 A2 and US 2012/0222348 A1 relate to C5-15-alkanols in foam inhibitors for biodiesel fuel or diesel fuel-biodiesel blends. WO 94/11331 A1 relates to the use of an alkoxylate of 2-propyl-1-heptanol for degreasing. EP 0666308 A2 relates to multi-purpose aqueous cleaning compositions comprising hydrogen peroxide, a 2-alkyl alkanol, a hydrophobic surfactant having an HLB below 14 and an anionic surfactant. US 3,856,695 relates to solvent based detergent compositions which consist essentially of 10-80% liquid aliphatic chlorinated hydrocarbon solvents, 10-40% liquid aliphatic ketones and/or liquid aliphatic esters, 6 to 50% water soluble liquid alkanols having two to five carbon atoms and 1-10% surface active agents. WO 98104761 relates to emulsions for cleaning, the emulsions containing water, one or more anionic surfactants from a group of selected molecular weight carboxylic acid salts, and one or more organic solvents so that the compositions have low conductivity and low viscosity. WO 03/006734 A2 relates to aqueous textile fibre degreasing agents essentially consisting of at least one amine oxide, at least one hexanediol, at least one secondary, branched or cyclic hexanol, at least one aminosurfactant, and water, and optionally at least one formulation additive. DE4021265 A1 relates to the use of saturated secondary alcohols with branching in the alpha-position to the OH and with molecular weight of 210-300 as additives to washing compositions. EP3170884 A1 relates to the use of an alcohol in a liquid cleaning composition comprising surfactant, for cleaning hydrophobic stains from surfaces and/or for providing suds longevity, wherein the alcohol is selected from the group consisting of: C4-C6 linear mono-alcohols, branched C4-C10 mono-alcohols having one or more C 1-C4 branching groups, alkyl mono-glycerols, and mixtures thereof.

SUMMARY OF THE INVENTION

The present invention relates to a laundry detergent composition comprising: deterative surfactant, wherein the deterative surfactant comprises a combination of anionic and nonionic surfactant;

solvent selected from the group consisting of: C4-C10 alkyl branched alcohols, and mixtures thereof, wherein “C4-C10 alkyl” refers to the primary alkyl chain, and refers to the weight average alkyl chain length, and wherein the composition has a pH of from 6.0 to 8.9.

The present invention further relates to a method of laundering fabric, preferably a stained fabric, comprising the steps of: providing a laundry detergent composition according to any preceding claim; diluting the laundry detergent composition to provide a wash liquor having a total surfactant concentration of greater than 300 ppm; and washing fabric in the wash liquor.

The present invention further relates to the use of solvent selected from the group consisting of: C4-C10 alkyl branched alcohols, and mixtures thereof, in a laundry detergent composition, for removing stains comprising natural oil and/or natural grease from fabrics.

DETAILED DESCRIPTION OF THE INVENTION

Laundry detergent compositions comprising C4-C10 alkyl branched alcohols, and mixtures thereof, have been found to provide improved removal of grease-containing stains, both via pre-treatment and also through the wash. Such grease removal benefits can be achieved using relatively low levels of the solvent. In addition, the solvents improve the viscosity of liquid laundry detergent compositions.

Unless otherwise noted, all component or composition levels are in reference to the active portion of that component or composition, and are exclusive of impurities, for example, residual solvents or by-products, which may be present in commercially available sources of such components or compositions.

All percentages and ratios are calculated by weight unless otherwise indicated. All percentages and ratios are calculated based on the total composition unless otherwise indicated.

All measurements are performed at 25°C unless otherwise specified.

As used herein, the articles including “a” and “an” when used in a claim, are understood to mean one or more of what is claimed or described.

Detergent composition:

As used herein, “liquid detergent composition” refers to liquid detergent composition which is fluid, and preferably capable of wetting and cleaning a fabric, e.g., clothing in a domestic washing machine. As used herein, “laundry detergent composition” refers to compositions suitable for washing clothes. The composition can include solids or gases in suitably subdivided form, but the overall composition excludes product forms which are non-fluid overall, such as tablets or

granules. The liquid laundry detergent composition preferably has a density in the range from 0.9 to 1.3 grams per cubic centimeter, more specifically from 1.00 to 1.10 grams per cubic centimeter, excluding any solid additives but including any bubbles, if present.

Aqueous liquid laundry detergent compositions are preferred. For such aqueous liquid laundry detergent compositions, the water content can be present at a level of from 5 % to 99 %, preferably from 15 % to 90 %, more preferably from 25 % to 80 % by weight of the liquid detergent composition.

The pH range of the detergent composition is from 6.0 to 8.9, preferably from pH 7 to 8.8.

Deterstive Surfactants

Deterstive surfactant as used herein means surfactants or mixtures of surfactants that provide cleaning, stain removing, or laundering benefit to soiled material. Suitable deterstive surfactants can be: anionic surfactant, nonionic surfactant, zwitterionic surfactant, and combinations thereof. The deterstive surfactant comprises a combination of anionic and nonionic surfactant.

The laundry composition can comprises deterstive surfactant at a level of from 1 wt% to 70 wt%, preferably from 10 wt% to 50 wt%, more preferably from 15 wt% to 35 wt%.

Suitable anionic surfactants can be selected from the group consisting of: alkyl sulphates, alkyl ethoxy sulphates, alkyl sulphonates, alkyl benzene sulphonates, fatty acids and their salts, and mixtures thereof. However, by nature, every anionic surfactant known in the art of detergent compositions may be used, such as disclosed in "Surfactant Science Series", Vol. 7, edited by W. M. Linfield, Marcel Dekker. However, the composition preferably comprises at least a sulphonic acid surfactant, such as a linear alkyl benzene sulphonic acid, but water-soluble salt forms may also be used. Alkyl ethoxy sulphates, or mixtures thereof, are also preferred.

Anionic sulfonate or sulfonic acid surfactants suitable for use herein include the acid and salt forms of linear or branched alkylbenzene sulfonates, alkyl ester sulfonates, alkane sulfonates, alkyl sulfonated polycarboxylic acids, and mixtures thereof. Suitable anionic sulfonate or sulfonic acid surfactants include: C5-C20 alkylbenzene sulfonates, more preferably C10-C16 alkylbenzene sulfonates, more preferably C11-C13 alkylbenzene sulfonates, C5-C20 alkyl ester sulfonates, C6-C22 primary or secondary alkane sulfonates, C5-C20 sulfonated polycarboxylic acids, and any mixtures thereof, but preferably C11-C13 alkylbenzene sulfonates. The aforementioned surfactants can vary widely in their 2-phenyl isomer content.

Anionic sulphate salts suitable for use in the compositions of the invention include the primary and secondary alkyl sulphates, having a linear or branched alkyl or alkenyl moiety having from 9

to 22 carbon atoms or more preferably 12 to 18 carbon atoms. Also useful are beta-branched alkyl sulphate surfactants or mixtures of commercial available materials, having a weight average (of the surfactant or the mixture) branching degree of at least 50%.

Mid-chain branched alkyl sulphates or sulfonates are also suitable anionic surfactants for use in the compositions of the invention. Preferred are the C5-C22, preferably C10-C20 mid-chain branched alkyl primary sulphates. When mixtures are used, a suitable average total number of carbon atoms for the alkyl moieties is preferably within the range of from greater than 14.5 to 17.5. Preferred mono-methyl-branched primary alkyl sulphates are selected from the group consisting of the 3-methyl to 13-methyl pentadecanol sulphates, the corresponding hexadecanol sulphates, and mixtures thereof. Dimethyl derivatives or other biodegradable alkyl sulphates having light branching can similarly be used.

Other suitable anionic surfactants for use herein include fatty methyl ester sulphonates and/or alkyl alkoxyated sulphates such as alkyl ethoxy sulphates (AES) and/or alkyl polyalkoxyated carboxylates (AEC). When used, the alkyl alkoxyated sulphate surfactant is preferably a blend of one or more alkyl ethoxyated sulphates. Suitable alkyl alkoxyated sulphates include C10-C18 alkyl ethoxylate, more preferably C12-C15 alkyl ethoxylate, with a degree of ethoxylation of from 1 to 5, preferably from 2 to 3.

The anionic surfactants are typically present in the form of their salts with alkanolamines or alkali metals such as sodium and potassium.

For improved stability and grease cleaning, the liquid detergent composition can comprise linear alkyl benzene sulfonate surfactant and alkyl alkoxyated sulphate surfactant, such that the ratio of linear alkyl benzene sulfonate surfactant to alkyl alkoxyated sulphate surfactant is from 0.1 to 5, preferably from 0.25 to 3, more preferably from 0.6 to 1.1.

The liquid detergent composition can comprise nonionic surfactant. The level of nonionic surfactant in the liquid detergent composition can be present at a level of less than 10 wt%, preferably less than 5 wt%, more preferably less than 1 wt%, most preferably less than 0.5 wt %.

Suitable nonionic surfactants include, but are not limited to C12-C18 alkyl ethoxylates ("AE") including the so-called narrow peaked alkyl ethoxylates and C6-C12 alkyl phenol alkoxyates (especially ethoxylates and mixed ethoxy/propoxy), block alkylene oxide condensate of C6-C12 alkyl phenols, alkylene oxide condensates of C8-C22 alkanols and ethylene oxide/propylene oxide block polymers (Pluronic - BASF Corp.), as well as semi polar nonionics (e.g., amine oxides and

phosphine oxides) can be used in the present compositions. An extensive disclosure of these types of surfactants is found in U.S. Pat. 3,929,678, Laughlin et al., issued December 30, 1975.

Alkylpolysaccharides such as disclosed in U.S. Pat. 4,565,647 Llenado are also useful nonionic surfactants in the compositions of the invention.


5 Also suitable are alkyl polyglucoside surfactants.

Nonionic surfactants of use include those of the formula $R_1(OC_2H_4)_nOH$, wherein R_1 is a C10-C16 alkyl group or a C8-C12 alkyl phenyl group, and n is from preferably 3 to 80. In some embodiments, the nonionic surfactants may be condensation products of C12-C15 alcohols with from 5 to 20 moles of ethylene oxide per mole of alcohol, e.g., C12-C13 alcohol condensed with
10 6.5 moles of ethylene oxide per mole of alcohol

Suitable amine oxide surfactants are amine oxides having the following formula : $R_1R_2R_3NO$ wherein R_1 is an hydrocarbon chain comprising from 1 to 30 carbon atoms, preferably from 6 to 20, more preferably from 8 to 16 and wherein R_2 and R_3 are independently saturated or unsaturated, substituted or unsubstituted, linear or branched hydrocarbon chains comprising from 1 to 4 carbon
15 atoms, preferably from 1 to 3 carbon atoms, and more preferably are methyl groups. R_1 may be a saturated or unsaturated, substituted or unsubstituted linear or branched hydrocarbon chain.

Suitable amine oxides for use herein are for instance preferably C_{12} - C_{14} dimethyl amine oxide, commercially available from Albright & Wilson, C_{12} - C_{14} amine oxides commercially available under the trade name Genaminox® LA from Clariant or AROMOX® DMC from AKZO Nobel.

20 Additional suitable nonionic surfactants include polyhydroxy fatty acid amides of the formula:



wherein R is a C9-17 alkyl or alkenyl, R_1 is a methyl group and Z is glycidyl derived from a reduced sugar or alkoxyated derivative thereof. Examples are N-methyl N-1-deoxyglucityl cocoamide and N-methyl N-1-deoxyglucityl oleamide. Processes for making polyhydroxy fatty
25 acid amides are known and can be found in Wilson, U.S. Patent 2,965,576 and Schwartz, U.S. Patent 2,703,798.

The liquid detergent composition can comprise a zwitterion. The zwitterion can be present at a level of from 0.1 wt% to 5 wt%, preferably from 0.2 wt% to 2 wt%, more preferably from 0.4 wt% to 1 wt %.

Suitable amphoteric or zwitterionic deterative surfactants include those which are known for use in hair care or other personal care cleansing. Non-limiting examples of suitable zwitterionic or amphoteric surfactants are described in U.S. Pat. Nos. 5,104,646 (Bolich Jr. et al.), 5,106,609 (Bolich Jr. et al.). Suitable amphoteric deterative surfactants include those surfactants broadly described as derivatives of aliphatic secondary and tertiary amines in which the aliphatic radical can be straight or branched chain and wherein one of the aliphatic substituents contains from 8 to 18 carbon atoms and one contains an anionic group such as carboxy, sulfonate, sulfate, phosphate, or phosphonate. Suitable amphoteric deterative surfactants for use in the present invention include, but are not limited to: cocoamphoacetate, cocoamphodiacetate, lauroamphoacetate, lauroamphodiacetate, and mixtures thereof.

Preferably surfactants comprising saturated alkyl chains are used.

Solvent

Suitable solvents are selected from the group consisting of: C4-C10 alkyl branched alcohols, and mixtures thereof. Such solvents have been found to improve the grease removal in detergent compositions when used during pretreatment of fabrics. Even more surprisingly, detergent compositions comprising such solvents have been found to improve grease removal even when during washing without pretreatment, and even when the solvents are present at low levels. Detergent compositions comprising such solvents are particularly effective at removing oils or grease which are at least partially naturally derived, such as animal fat, vegetable fat, and mixtures thereof, from fabrics. In addition, the solvents can improve the rheological profile of liquid detergent compositions.

“C4-C10 alkyl” refers to the primary alkyl chain, and refers to the weight average alkyl chain length. Preferred C4-C10 alkyl branched alcohols are C4-C10 primary mono-alcohols having one or more C1-C4 branching groups, preferably selected from the group consisting of: methyl butanol, ethyl butanol, methyl pentanol, ethyl pentanol, methyl hexanol, ethyl hexanol, propyl hexanol, dimethyl hexanol, trimethyl hexanol, methyl heptanol, ethyl heptanol, propyl heptanol, dimethyl heptanol, trimethyl heptanol, methyl octanol, ethyl octanol, propyl octanol, butyl octanol, dimethyl octanol, trimethyl octanol, methyl nonanol, ethyl nonanol, propyl nonanol, butyl nonanol, dimethyl nonanol, trimethyl nonanol and mixtures thereof.

More preferred C4-C10 alkyl branched alcohols are C6-C8 alkyl branched alcohols, preferably selected from the group consisting of: methyl hexanol, ethyl hexanol, propyl hexanol, dimethyl hexanol, trimethyl hexanol, methyl heptanol, ethyl heptanol, propyl heptanol, dimethyl heptanol, trimethyl heptanol, methyl octanol, ethyl octanol, propyl octanol, butyl octanol, dimethyl octanol, trimethyl octanol, and mixtures thereof.

Even more preferred are C6-C7 alkyl branched alcohols, and mixtures thereof. Most preferably, the solvent is selected from the group consisting of: 2-ethylhexanol, 3,5,5-trimethyl-1-hexanol, 2-propylheptanol, and mixtures thereof.

The laundry detergent composition can comprise the solvent at a level of from 0.1 wt% to 10wt%, preferably from 0.3 wt% to 7.5 wt%, preferably from 0.5 wt% to 5 wt% of the solvent.

For the present invention, alkanolamines, such as monoethanolamine and triethanolamine, are considered as alkali agents, or neutralising agents for anionic surfactants and the like, but not as solvents.

Optional Ingredients

- 5 The detergent composition may additionally comprise one or more of the following optional ingredients: external structurant or thickener, enzymes, enzyme stabilizers, cleaning polymers, bleaching systems, optical brighteners, hueing dyes, particulate material, perfume and other odour control agents, hydrotropes, suds suppressors, fabric care benefit agents, pH adjusting agents, dye transfer inhibiting agents, preservatives, non-fabric substantive dyes and mixtures thereof. In more
10 preferred embodiments, the laundry detergent composition does not comprise a bleach.

External structurant or thickener: Preferred external structurants and thickeners are those that do not rely on charge – charge interactions for providing a structuring benefit. As such, particularly preferred external structurants are uncharged external structurants, such as those selected from the group consisting of: non-polymeric crystalline, hydroxyl functional structurants, such as
15 hydrogenated castor oil; microfibrillated cellulose; uncharged hydroxyethyl cellulose; uncharged hydrophobically modified hydroxyethyl cellulose; hydrophobically modified ethoxylated urethanes; hydrophobically modified non-ionic polyols; and mixtures thereof.

Suitable non-polymeric crystalline, hydroxyl functional structurants are known in the art, and generally comprise a crystallizable glyceride which can be pre-emulsified to aid dispersion into the
20 final liquid detergent composition. A non-limiting example of such a pre-emulsified external structuring system comprises: (a) crystallizable glyceride(s); (b) anionic surfactant; and (c) water and optionally, non-aminofunctional organic solvents. Each of these components is discussed in detail below. The preferred non-polymeric crystalline, hydroxy-functional structurant comprises a

crystallizable glyceride, preferably hydrogenated castor oil or "HCO".

Suitable polymeric structurants include naturally derived and/or synthetic polymeric structurants. Examples of naturally derived polymeric structurants of use in the present invention include: microfibrillated cellulose, hydroxyethyl cellulose, hydrophobically modified hydroxyethyl
5 cellulose, carboxymethyl cellulose, polysaccharide derivatives and mixtures thereof. Non-limiting examples of microfibrillated cellulose are described in WO 2009/101545 A1. Suitable polysaccharide derivatives include: pectine, alginate, arabinogalactan (gum Arabic), carrageenan, gellan gum, xanthan gum, guar gum and mixtures thereof.

Examples of synthetic polymeric structurants or thickeners of use in the present invention include:

10 polycarboxylates, hydrophobically modified ethoxylated urethanes (HEUr), hydrophobically modified non-ionic polyols and mixtures thereof.

Preferably the polycarboxylate polymer is a polyacrylate, polymethacrylate or mixtures thereof.

In another preferred embodiment, the polyacrylate is a copolymer of unsaturated mono- or di-carbonic acid and 1-30C alkyl ester of the (meth) acrylic acid. Such copolymers are available from

15 Noveon inc under the tradename Carbopol Aqua 30. Suitable polyacrylates include alkali swellable emulsion (ASE) thickeners and hydrophobically modified alkali sellable emulsion (HASE) thickeners. Suitable hydrophobically modified alkali swellable emulsions (HASE) are sold under the various brand names by Lubrizol Corporation, Clariant, Akzo Nobel, Coatex, 3V Sigma, SEPPIC, Ashland and BASF. Particularly suited, are Novethix L10 and Novethix HC200
20 (Lubrizol), Crystasense Sapphire (Clariant), Alcoguard 5800 (Akzo Nobel), Rheosolve 637 and Rheosolve 650 (Coatex), Polygel W30 (3V Sigma), Capigel98 (SEPPIC), Jaypol AT4 (Ashland), Rheovis AT120, Salcare SC80 and Luvigel FIT (BASF)."

The branched solvents of the present invention have been surprisingly found to synergistically improve both the viscosity and yield strength provided by the external structurant or thickener,
25 while maintaining the dispersibility of the liquid detergent composition. In addition, the branched alcohols have surprisingly been found to result in structuring in addition to thickening, when combined with polymeric thickeners.

Preferably, the aqueous liquid detergent composition has a viscosity of 50 to 5,000, preferably 75 to 1,000, more preferably 100 to 500 mPa.s, when measured at a shear rate of 100 s⁻¹, at a
30 temperature of 20°C. For improved phase stability, and also improved stability of suspended ingredients, the aqueous liquid detergent composition has a viscosity of 50 to 250,000, preferably 5,000 to 125,000, more preferably 10,000 to 35,000 mPa.s, when measured at a shear rate of 0.05 s⁻¹, at a temperature of 20°C.

Cleaning polymers: Suitable cleaning polymers provide for broad-range soil cleaning of surfaces and fabrics and/or suspension of the soils. Any suitable cleaning polymer may be of use. Useful cleaning polymers are described in USPN 2009/0124528A1. Non-limiting examples of useful categories of cleaning polymers include: amphiphilic alkoxyated grease cleaning polymers; clay
5 soil cleaning polymers; soil release polymers; and soil suspending polymers. The detergent composition may comprise amphiphilic alkoxyated grease cleaning polymers, which may have balanced hydrophilic and hydrophobic properties such that they remove grease particles from fabrics and surfaces. The amphiphilic alkoxyated grease cleaning polymers may comprise a core structure and a plurality of alkoxyate groups attached to that core structure. These may comprise
10 alkoxyated polyalkyleneimines, for example. Such compounds may comprise, but are not limited to, ethoxylated polyethyleneimine, ethoxylated hexamethylene diamine, and sulfated versions thereof. Polypropoxylated derivatives may also be included. A wide variety of amines and polyalkyleneimines can be alkoxyated to various degrees. A useful example is 600g/mol polyethyleneimine core ethoxylated to 20 EO groups per NH and is available from BASF. The
15 alkoxyated polyalkyleneimines may have an inner polyethylene oxide block and an outer polypropylene oxide block. Other suitable cleaning polymers include polyester based soil release polymers, such as SRA300, supplied by Clariant. The detergent compositions may comprise from 0.1% to 10%, preferably, from 0.1% to 8%, more preferably from 0.1% to 6%, by weight of the detergent composition, of alkoxyated polyamines.

20 Polymer Deposition Aid: The laundry detergent composition can comprise from 0.1% to 7%, more preferably from 0.2% to 3%, of a polymer deposition aid. As used herein, "polymer deposition aid" refers to any cationic polymer or combination of cationic polymers that significantly enhance deposition of a fabric care benefit agent onto the fabric during laundering. Suitable polymer deposition aids can comprise a cationic polysaccharide and/or a copolymer. "Fabric care benefit
25 agent" as used herein refers to any material that can provide fabric care benefits. Non-limiting examples of fabric care benefit agents include: silicone derivatives, oily sugar derivatives, dispersible polyolefins, polymer latexes, cationic surfactants and combinations thereof. Preferably, the deposition aid is a cationic or amphoteric polymer. The cationic charge density of the polymer preferably ranges from 0.05 milliequivalents/g to 6 milliequivalents/g. The charge density is
30 calculated by dividing the number of net charge per repeating unit by the molecular weight of the repeating unit. In one embodiment, the charge density varies from 0.1 milliequivalents/g to 3 milliequivalents/g. The positive charges could be on the backbone of the polymers or the side chains of polymers.

Organic builder and/or chelant: The laundry detergent composition can comprise from 0.6% to 10%, preferably from 2 to 7% by weight of one or more organic builder and/or chelants. Suitable organic builders and/or chelants are selected from the group consisting of: MEA citrate, citric acid, aminoalkylenepoly(alkylene phosphonates), alkali metal ethane 1-hydroxy disphosphonates, and
5 nitrilotrimethylene, phosphonates, diethylene triamine penta (methylene phosphonic acid) (DTPMP), ethylene diamine tetra(methylene phosphonic acid) (DDTMP), hexamethylene diamine tetra(methylene phosphonic acid), hydroxy- ethylene 1,1 diphosphonic acid (HEDP), hydroxyethane dimethylene phosphonic acid, ethylene di-amine di-succinic acid (EDDS), ethylene diamine tetraacetic acid (EDTA), hydroxyethylethylenediamine triacetate (HEDTA),
10 nitrilotriacetate (NTA), methylglycinediacetate (MGDA), iminodisuccinate (IDS), hydroxyethyliminodisuccinate (HIDS), hydroxyethyliminodiacetate (HEIDA), glycine diacetate (GLDA), diethylene triamine pentaacetic acid (DTPA), catechol sulfonates such as TironTM and mixtures thereof.

Enzymes: Suitable enzymes provide cleaning performance and/or fabric care benefits. Examples
15 of suitable enzymes include, but are not limited to, hemicellulases, peroxidases, proteases, cellulases, xylanases, lipases, phospholipases, esterases, cutinases, pectinases, keratanases, reductases, oxidases, phenoloxidas, lipoxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases, hyaluronidase, chondroitinase, laccase, and known amylases, or combinations thereof. A preferred enzyme combination comprises a
20 cocktail of conventional deterative enzymes such as protease, lipase, cutinase and/or cellulase in conjunction with amylase. Deterative enzymes are described in greater detail in U.S. Patent No. 6,579,839.

Enzyme stabiliser: Enzymes can be stabilized using any known stabilizer system such as calcium and/or magnesium compounds, boron compounds and substituted boric acids, aromatic borate
25 esters, peptides and peptide derivatives, polyols, low molecular weight carboxylates, relatively hydrophobic organic compounds [e.g. certain esters, diacyl glycol ethers, alcohols or alcohol alkoxylates], alkyl ether carboxylate in addition to a calcium ion source, benzamidine hypochlorite, lower aliphatic alcohols and carboxylic acids, N,N-bis(carboxymethyl) serine salts; (meth)acrylic acid-(meth)acrylic acid ester copolymer and PEG; lignin compound, polyamide
30 oligomer, glycolic acid or its salts; poly hexa methylene bi guanide or N,N-bis-3-amino-propyl-dodecyl amine or salt; and mixtures thereof.

Hueing dyes: The detergent composition may comprise fabric hueing agent (sometimes referred to as shading, bluing, or whitening agents). Typically the hueing agent provides a blue or violet

shade to fabric. Hueing agents can be used either alone or in combination to create a specific shade of hueing and/or to shade different fabric types. This may be provided for example by mixing a red and green-blue dye to yield a blue or violet shade. Hueing agents may be selected from any known chemical class of dye, including but not limited to acridine, anthraquinone (including polycyclic quinones), azine, azo (e.g., monoazo, disazo, trisazo, tetrakisazo, polyazo), including premetallized azo, benzodifurane and benzodifuranone, carotenoid, coumarin, cyanine, diazahemicyanine, diphenylmethane, formazan, hemicyanine, indigoids, methane, naphthalimides, naphthoquinone, nitro and nitroso, oxazine, phthalocyanine, pyrazoles, stilbene, styryl, triarylmethane, triphenylmethane, xanthenes and combinations thereof.

Optical brighteners: The detergent composition may comprise, based on the total detergent composition weight, from 0.005 to 2 %, preferably 0.01 to 0.1 % of a fluorescent agent (optical brightener). Fluorescent agents are well known and many fluorescent agents are available commercially. Usually, these fluorescent agents are supplied and used in the form of their alkali metal salts, for example, the sodium salts. Preferred classes of fluorescent agent are: Di-styryl biphenyl compounds, e.g. Tinopal (Trade Mark) CBS-X, Di-amine stilbene di-sulphonic acid compounds, e.g. Tinopal DMS pure Xtra and Blankophor (Trade Mark) HRH, and Pyrazoline compounds, e.g. Blankophor SN. Preferred fluorescers are: sodium 2-(4-styryl-3-sulfophenyl)-2H-naphthol[1,2-d]trazole, disodium 4,4'-bis{[(4-anilino-6-(N methyl-N-2 hydroxyethyl) amino 1,3,5-triazin-2-yl)]amino}stilbene-2-2' disulfonate, disodium 4,4'-bis{[(4-anilino-6-morpholino-1,3,5-triazin-2-yl)]annino} stilbene-2-2' disulfonate, and disodium 4,4'-bis(2-sulfoslyryl)biphenyl.

Hydrotrope: The detergent composition may comprise, based on the total detergent composition weight, from 0 to 30%, preferably from 0.5 to 5%, more preferably from 1.0 to 3.0%, which can prevent liquid crystal formation. The addition of the hydrotrope thus aids the clarity/transparency of the composition. Suitable hydrotropes comprise but are not limited to urea, salts of benzene sulphonate, toluene sulphonate, xylene sulphonate or cumene sulphonate. Preferably, the hydrotrope is selected from the group consisting of propylene glycol, xylene sulfonate, ethanol, and urea to provide optimum performance.

Particles: The composition can also comprise particles, especially when the composition further comprises a structurant or thickener. The composition may comprise, based on the total composition weight, from 0.02% to 10%, preferably from 0.1% to 4%, more preferably from 0.25% to 2.5% of particles. Said particles include beads, pearlescent agents, microcapsules, and mixtures thereof.

Microcapsules: Suitable capsules are typically formed by at least partially, preferably fully, surrounding a benefit agent with a wall material. Preferably, the capsule is a perfume capsule, wherein said benefit agent comprises one or more perfume raw materials. The capsule wall material may comprise: melamine, polyacrylamide, silicones, silica, polystyrene, polyurea, polyurethanes, polyacrylate based materials, polyacrylate esters based materials, gelatin, styrene malic anhydride, polyamides, aromatic alcohols, polyvinyl alcohol, resorcinol-based materials, poly-isocyanate-based materials, acetals (such as 1,3,5-triol-benzene-gluteraldehyde and 1,3,5-triol-benzene melamine), starch, cellulose acetate phthalate and mixtures thereof. Preferably, the capsule wall comprises melamine and/or a polyacrylate based material. The perfume capsule may be coated with a deposition aid, a cationic polymer, a non-ionic polymer, an anionic polymer, or mixtures thereof. Preferably, the perfume capsules have a volume weighted mean particle size from 0.1 microns to 100 microns, preferably from 0.5 microns to 60 microns. Especially where the composition comprises capsules having a shell formed at least partially from formaldehyde, the composition can additionally comprise one or more formaldehyde scavengers.

Process of making the laundry detergent composition:

The laundry detergent compositions can be made using any suitable process known to the skilled person. Typically, the ingredients are blended together in any suitable order. Preferably, the deterative surfactants are added as part of a concentrated premix, to which are added the other optional ingredients. Preferably, the solvent is added either last, or if an external structurant is added, immediately before the external structurant, with the external structurant being added as the last ingredient.

Method of laundering fabrics:

The laundry detergent compositions of the present invention can be used to launder fabrics. In such methods, the laundry detergent composition is diluted to provide a wash liquor having a total surfactant concentration of greater than 300 ppm, preferably from 400 ppm to 2,500 ppm, more preferably from 600 ppm to 1000 ppm. The fabric is then washed in the wash liquor, and preferably rinsed.

The method of the present invention is particularly suited for removing stains, especially removing stains comprising natural oil and/or grease, such as animal fat, vegetable fat, and mixtures thereof.

METHODS:A) pH measurement:

The pH is measured, at 25°C, using a Santarius PT-10P pH meter with gel-filled probe (such as the Toledo probe, part number 52 000 100), calibrated according to the instructions manual. The pH is measured in a 10% dilution in demineralised water (i.e. 1 part laundry detergent composition and 9 parts demineralised water).

B) Method of measuring viscosity:

The viscosity is measured using an AR 2000 rheometer from TA instruments using a cone and plate geometry with a 40 mm diameter and an angle of 1°. The viscosity at the different shear rates is measured via a logarithmic shear rate sweep from 0.1 s⁻¹ to 1200 s⁻¹ in 3 minutes time at 20°C. Low shear viscosity is measured at a continuous shear rate of 0.05 s⁻¹.

EXAMPLES:

A following laundry detergent compositions were prepared by simple mixing, comprising 5 wt% of different solvents, except for comparative example C, which comprised no solvent:

	Ex. 1	Ex. 2	Ex. 3	Ex. A*	Ex. B*	Ex. C*
	wt%	wt%	wt%	wt%	wt%	wt%
Linear C10-C12 alkyl benzene sulfonate	4.8	4.8	4.8	4.8	4.8	4.8
C12-C14 alkyl ethoxylated (EO3) sulfate	2.2	2.2	2.2	2.2	2.2	2.2
C45 EO7 ethoxylated nonionic surfactant	3.6	3.6	3.6	3.6	3.6	3.6
Topped kernel fatty acid	2.5	2.5	2.5	2.5	2.5	2.5
Citric acid	1.7	1.7	1.7	1.7	1.7	1.7
2-ethylhexanol	5	-	-	-	-	-
3,5,5-trimethyl-1-hexanol	-	5	-	-	-	-
2-propylheptanol	-	-	5	-	-	-
1,2 propanediol	-	-	-	5	-	-
1-butanol	-	-	-	-	5	-
Sodium hydroxide	to pH 8.4	to pH 8.4	to pH 8.4	to pH 8.4	to pH 8.4	to pH 8.4
Water	to 100%	to 100%	to 100%	to 100%	to 100%	to 100%

*comparative

100 ml of each composition was tested in a full-scale wash test to compare stain removal performance of the compositions of the present invention, versus the comparative compositions.

- 5 Knitted cotton swatches (30cm x 30 cm) comprising fatty stains, including cooked beef, bacon grease and make-up stains were ordered from Warwick Equest.

The stains were pretreated by applying 1ml of the composition and scrubbing 10 times with an Ariel pretreat dosing cap (10 vertical strokes). The treated stains were then left for 5 minutes before placing in a Miele W527 washing machine with 3Kg of mixed load (13 cotton and 10 poly-cotton fabrics, 30 cm x 30 cm) and the remaining laundry detergent composition (to 100 ml) was dosed via a dosing ball. The wash test was run with a cotton short cycle at 30°C and 2.5mMol water hardness.

The wash test was repeated 4 times, with 2 replicates in each load, resulting in 8 test replicates per composition.

- 15 The fabric stain removal was graded using a Hunter Colorquest by measuring the amount of stain that is left after washing compared to an initial unwashed set. The results of the stain removal data were averaged for each stain and each product to give the overall stain removal index (SRI) grade vs. the reference (comparative example C, comprising no solvent).

- 20 The response was measured at 460 nm and used to calculate the soil removal delta in % (ΔSR) according to the following formula:

Formula 1: for calculating the soil removal delta in % (ΔSR)

$$\Delta SRI = \frac{R_{before} - R_{after}}{R_{before} - R_{clean}} \times 100\%$$

25

R_{before} = Remission before washing

R_{after} = Remission after washing

R_{clean} = Remission from clean white cotton

- 30 The results are shown on the tables below (versus Ex. C, comprising no solvent):

	Ex. C	Ex. 1	Ex. 2	Ex. 3
Stain	Δ SRI (%)	Change in Δ SRI (%)	Change in Δ SRI (%)	Change in Δ SRI (%)
Cooked beef	60.3	+12.9	+12.5	+9.1
Bacon grease	60.5	+12.8	+15.1	+11.3
Make up	68.9	+18.7	+18.7	+17.6

	Ex. C	Ex. A	Ex. B	Ex. 3
Stain	SRI (%)	Change in Δ SRI (%)	Change in Δ SRI (%)	Change in Δ SRI (%)
Cooked beef	66.3	-2.1	+4.4	+17.1
Bacon grease	52.3	-0.9	-1.1	+26.4
Make up	66.8	-7.9	-14.1	+21.9

As can be seen from the above data, pretreating and washing with laundry detergent compositions of the present invention, comprising the branched alcohol, resulted in improved removal of greasy stains.

A following laundry detergent compositions were prepared by simple mixing, comprising 2 wt% of different solvents:

	Ex. 4	Ex. D*	Ex. E*
	wt%	wt%	wt%
Linear C10-C12 alkyl benzene sulfonate	4.8	4.8	4.8
C12-C14 alkyl ethoxylated (EO3) sulfate	2.2	2.2	2.2
C45 EO7 ethoxylated nonionic surfactant	3.6	3.6	3.6
Topped kernel fatty acid	2.5	2.5	2.5
Citric acid	1.7	1.7	1.7
2-propylheptanol	2	-	-
1,2-propanediol	-	2	-
1-butanol	-	-	2
Sodium hydroxide	to pH 8.4	to pH 8.4	to pH 8.4
Water	to 100%	to 100%	to 100%

*comparative

The stain removal test was repeated, using example D (2 wt% propanediol) as reference, with the following results:

	Ex. D*	Ex. 4	Ex. E*
Stain	SRI (%)	Δ SRI (%)	Δ SRI (%)
Cooked beef	71.5	+14.0	+1.0
Bacon grease	53.4	+17.6	-0.6
Make up	63.1	+25.0	-1.8

- 5 As can be seen from the above data, laundry detergent compositions of the present invention, comprising the branched alcohol, resulted in improved removal of greasy stains, even when the solvent is added at 2wt%.

10 The stain removal test was repeated using the same conditions, except that no pretreatment step was used, with example D (2 wt% propanediol) as reference. The test used knitted cotton swatches (30cm x 30 cm) comprising fatty stains, including cooked beef, burnt butter, bacon grease and make-up stains. All other test settings remained the same.

The results are given in the table below:

	Ex. D*	Ex. 4	Ex. E*
Stain	SRI (%)	Δ SRI (%)	Δ SRI (%)
Cooked beef	47.5	+14.0	+3.3
Burnt butter	57.1	+14.9	+2.2
Bacon grease	46.3	+11.9	+0.3
Make up	41.5	+33.2	+1.2

15

As can be seen from the above data, laundry detergent compositions of the present invention, comprising the branched alcohol, resulted in improved removal of greasy stains, even when the solvent is added at 2wt% and even when no pretreatment was done on the stained fabrics.

- 20 The following laundry detergent compositions were prepared by simple mixing, with example F being comparative and comprising no solvent, and examples 10 to 12 being of the invention,

comprising various levels of 2-propylheptanol and surfactant:

	Ex. F *	Ex. 10	Ex. 11	Ex. 12
	wt%	wt%	wt%	wt%
Linear C10-C12 alkyl benzene sulfonate	4.8	3.9	4.4	4.6
C12-C14 alkyl ethoxylated (EO3) sulfate	2.2	1.8	2.0	2.1
C45 EO7 ethoxylated nonionic surfactant	3.6	2.9	3.2	3.4
Topped kernel fatty acid	2.5	2.1	2.3	2.4
Citric acid	1.7	1.7	1.7	1.7
2-propylheptanol	-	2	1	0.5
Sodium hydroxide	to pH 8.4	to pH 8.4	to pH 8.4	to pH 8.4
Water	to 100%	to 100%	to 100%	to 100%

The stain removal test was repeated using the same conditions, with no pretreatment step, using comparative example F (0 wt% 2-propylheptanol) as the reference. The test used knitted cotton swatches (30cm x 30 cm) comprising fatty stains, including cooked beef, burnt butter, bacon grease and make-up stains. All other test settings remained the same as with the earlier described test.

The results are given in the table below:

	Ex. F *	Ex. 10	Ex. 11	Ex. 12
Stain	SRI (%)	Δ SRI (%)	Δ SRI (%)	Δ SRI (%)
Cooked beef	48.2	+14.7	+7.2	+2.5
Burnt butter	64.1	+10.7	+4.9	+0.1
Bacon grease	49.9	+7.5	+3.6	-0.2
Make up	44.4	+21.7	+16.6	+3.9

As can be seen from the above data, laundry detergent compositions of the present invention, comprising the branched alcohol, resulted in improved removal of greasy stains, even when the solvent is added down to 0.5wt%, even at reduced surfactant levels, and even when the detergent composition is added as part of the main wash, with no pretreat step.

Examples 13 to 18 are non-limiting embodiments of the present invention. Percentages are by weight unless otherwise specified.

	Ex. 13	Ex. 14	Ex. 15	Ex. 16	Ex. 17	Ex. 18
	wt%	wt%	wt%	wt%	wt%	wt%
C12-14 EO3 alkyl ethoxylated sulfate	9.8	4.0	0.6	7.1	9.1	3.4
C12-15 linear alkylbenzene sulfonic acid	14.5	9.2	5.9	10.8	6.9	4.6
C12-15 EO8 ethoxylated alcohol	2.0	4.0	3.0	7.0	6	4.6
Dodecydimethylamine-N-oxide	0	0.5	0	0	0	0
Citric Acid	4.8	2.8	1.9	2.8	3.3	2.2
C12-18 Fatty Acid	3.3	1.7	1.2	4.7	4.9	1.6
Sodium Cumene Sulfonate	0	1.7	0.2	0	0	0
Zwitterionic polyamine ¹	0.7	0.7	0.3	0	0	0
Diethylenetriamine penta(methylene phosphonic acid), Sodium salt (DTPMP)	0	0.5	0.2	0	0.5	0.2
1-hydroxyethane 1,1-diphosphonic acid (HEDP)	0.6	0	0	2.0	0	0
Mannanase ²	0.003	0.002	0	0	0.0002	0.002
Amylase ³	0.013	0.004	0.0016	0	0.02	0.005
Protease ⁴	0.039	0.02	0.018	0	0.06	0.008
Cellulase ⁵	0	0	0	0	0.006	0
Pectate Lyase ⁶	0.005	0.002	0	0	0.001	0
Lipase ⁷	0	0	0	0	0.010	0
PEG-PVAc Polymer ⁸	1.9	1.3	0.9	0	0	0
Di-ethoxylated poly (1,2 propylene terephthalate) short block soil release polymer ⁹	0	0	0	0	0.55	0
Ethoxylated Polyethylenimine ¹⁰	0	0	0	1.2	0	0
Brightener 49	0.08	0.05	0.05	0.24	0	0.05
Bis azo or azo thiophene hueing dye ¹¹	0	0	0.02	0	0	0
Hydrogenated castor oil	1	0.30	0.44	0	0	0
2-propylheptanol	5	-	-	6	1	0.5
2-ethylhexanol	-	2	-	-	-	1
3,5,5-trimethyl-1-hexanol	-	-	3	-	4	-
1, 2 propanediol	8.3	1.3	1.0	9.2	6.1	3.6

Ethanol	0	0	0.5	0	1.9	0
Glycerine	0	0	0	0.5	0.3	1.0
Sodium formate	0	0.03	0.3	0	0	0
Calcium Chloride	0.03	0.01	0.006	0	0	0
Boric acid	0	0	0	0	1.1	1.8
Monoethanolamine	8.8	0.24	0.35	6.1	0	0
Triethanolamine	0	0	0	4.1	0	0
Sodium hydroxide	to pH 7.4	to pH 7.9	to pH 8.0	to pH 8.8	to pH 8.5	to pH 8.3
Acticide MBS2550	0.003	0.001	0.001	0.01	0.03	0.02
Silicone suds suppressor	0	0.003	0.003	0	0	0.02
Perfume microcapsules	0	0.25	0	0	0	0
Perfume	1.5	0.9	0.6	2.0	1.0	0.8
Dye	0.009	0.005	0.004	0	0.005	0.002
Water	to 100%	to 100%	to 100%	to 100%	to 100%	to 100%

¹ Zwitterionic ethoxylated quaternized sulfated hexamethylene diamine, supplied by BASF, Germany

² Mannanase enzyme originating from *Bacillus* sp. I633 available from Novozymes, Denmark

³ Termamyl® Ultra, available from Novozymes, Denmark

5 ⁴ Protease enzyme from *Bacillus Amyloliquefaciens* as described in EP 0 130 756 B1 published January 9, 1985

⁵ Carezyme® available from Novozymes, Denmark

⁶ Pectawash® 20L, supplied by Novozymes, Denmark

⁷ Lipex®, supplied by Novozymes, Denmark

10 ⁸ Polyvinyl acetate grafted polyethylene oxide copolymer having a polyethylene oxide backbone and multiple polyvinyl acetate side chains, supplied by BASF, Germany.

⁹ TexCare® SRN-100, supplied by Clariant, Germany

¹⁰ Polyethyleneimine (MW = 600) with 20 ethoxylate groups per –NH, supplied by BASF

¹¹ Supplied by Milliken, USA

15

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as “40 mm” is intended to mean “about 40 mm”.

CLAIMS

What is claimed is

1. A laundry detergent composition comprising:
 - a) deterative surfactant, wherein the deterative surfactant comprises a combination of anionic and nonionic surfactant;
 - b) solvent selected from the group consisting of: C4-C10 alkyl branched alcohols, and mixtures thereof, wherein "C4-C10 alkyl" refers to the primary alkyl chain, and refers to the weight average alkyl chain length; andwherein the composition has a pH of from 6.0 to 8.9.
2. The laundry detergent composition according to claim 1, wherein the solvent is selected from the group consisting of: C4-C10 primary mono-alcohols having one or more C1-C4 branching groups, preferably selected from the group consisting of: methyl butanol, ethyl butanol, methyl pentanol, ethyl pentanol, methyl hexanol, ethyl hexanol, propyl hexanol, dimethyl hexanol, trimethyl hexanol, methyl heptanol, ethyl heptanol, propyl heptanol, dimethyl heptanol, trimethyl heptanol, methyl octanol, ethyl octanol, propyl octanol, butyl octanol, dimethyl octanol, trimethyl octanol, methyl nonanol, ethyl nonanol, propyl nonanol, butyl nonanol, dimethyl nonanol, trimethyl nonanol and mixtures thereof.
3. The laundry detergent composition according to claim 2, wherein the solvent is selected from the group consisting of: C6-C8 alkyl branched alcohols, preferably selected from the group consisting of: methyl hexanol, ethyl hexanol, propyl hexanol, dimethyl hexanol, trimethyl hexanol, methyl heptanol, ethyl heptanol, propyl heptanol, dimethyl heptanol, trimethyl heptanol, methyl octanol, ethyl octanol, propyl octanol, butyl octanol, dimethyl octanol, trimethyl octanol, and mixtures thereof, more preferably 2-ethylhexanol, 3,5,5-trimethyl-1-hexanol, 2-propylheptanol, and mixtures thereof.
4. The laundry detergent composition according to any preceding claims, wherein the composition comprises solvent at a level of from 0.1 wt% to 10wt%, preferably from 0.3 wt% to 7.5 wt%, preferably from 0.5 wt% to 5 wt% of the solvent.

5. The laundry detergent composition according to any preceding claims, wherein the deterative surfactant is selected from the group consisting of: anionic surfactant, non-ionic surfactant, and mixtures thereof.
6. The laundry detergent composition according to claim 5, wherein the deterative surfactant comprises anionic surfactant, wherein the anionic surfactant comprises linear alkyl benzene sulfonate and alkyl alkoxyated sulfate and the ratio of linear alkyl benzene sulfonate surfactant to alkyl alkoxyated sulphate surfactant is from 0.1 to 5, preferably from 0.25 to 3, more preferably from 0.6 to 1.1.
7. The laundry detergent composition according to any preceding claims, wherein the laundry composition comprises deterative surfactant at a level of from 1 wt% to 70 wt%, preferably from 10 wt% to 50 wt%, more preferably from 15 wt% to 35 wt%.
8. The laundry detergent composition according to any preceding claim, wherein the liquid laundry detergent composition comprises an external structurant or thickener selected from the group consisting of: non-polymeric crystalline, hydroxyl functional structurants; microfibrillated cellulose; uncharged hydroxyethyl cellulose; uncharged hydrophobically modified hydroxyethyl cellulose; hydrophobically modified ethoxylated urethanes; polyacrylates; hydrophobically modified non-ionic polyols; and mixtures thereof.
9. The laundry detergent composition according to claim 8, wherein the laundry composition further comprises particles, preferably selected from the group consisting of: beads, pearlescent agents, microcapsules, and mixtures thereof, more preferably microcapsules.
10. The laundry detergent composition according to any preceding claims, wherein the laundry composition comprises enzymes, chelants, additional non-aminofunctional organic solvents, and mixtures thereof.
11. A method of laundering fabric, preferably a stained fabric, comprising the steps of:
 - a) providing a laundry detergent composition according to any preceding claim;
 - b) diluting the laundry detergent composition to provide a wash liquor having a total surfactant concentration of greater than 300 ppm; and
 - c) washing fabric in the wash liquor.

12. The use of solvent selected from the group consisting of: C4-C10 alkyl branched alcohols, and mixtures thereof, in a laundry detergent composition for laundering fabrics, for removing stains comprising natural oil and/or natural grease from fabrics.
13. The use according to claim 12, wherein the oil or grease are at least partially naturally derived.
14. The use according to any of claims 12 or 13, wherein the oil or grease comprise: animal fat, vegetable fat, and mixtures thereof.
15. The use according to any of claims 12 to 14, wherein the laundry detergent composition is diluted before applying to the fabric.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2018/030752

A. CLASSIFICATION OF SUBJECT MATTER

INV. C11D1/37 C11D1/83 C11D3/20 C11D11/00
ADD. C11D1/22 C11D1/29

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 40 21 265 A1 (KREUSSLER CHEM FAB [DE]) 9 January 1992 (1992-01-09) cited in the application example 1	1-3,5,7, 10-15
X	----- EP 0 666 308 A2 (PROCTER & GAMBLE [US]) 9 August 1995 (1995-08-09) cited in the application page 1, lines 3-5, 50-52 page 2, lines 45-50 page 4, lines 12-17 page 5, lines 27-30, 43-50 examples claims ----- -/-	1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

31 July 2018

Date of mailing of the international search report

13/08/2018

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Bertran Nadal, Josep

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2018/030752

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2012/252716 A1 (BARNABAS FREDDY ARTHUR [US] ET AL) 4 October 2012 (2012-10-04) paragraphs [0006], [0016], [0041] examples claims -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2018/030752

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 4021265	A1	09-01-1992	NONE

EP 0666308	A2	09-08-1995	AU 701779 B2 04-02-1999
		BR 9506715 A 09-09-1997	
		CA 2182672 A1 10-08-1995	
		CN 1144533 A 05-03-1997	
		EG 20508 A 30-06-1999	
		EP 0666308 A2 09-08-1995	
		FI 963078 A 01-10-1996	
		JP H09508655 A 02-09-1997	
		WO 9521233 A1 10-08-1995	

US 2012252716	A1	04-10-2012	CA 2829638 A1 04-10-2012
		CN 103459576 A 18-12-2013	
		EP 2691503 A1 05-02-2014	
		JP 5805845 B2 10-11-2015	
		JP 2014510849 A 01-05-2014	
		RU 2013138306 A 10-05-2015	
		US 2012252716 A1 04-10-2012	
		WO 2012135411 A1 04-10-2012	
