

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2014289342 B2

- (54) Title
Synthetic vaccines against Streptococcus pneumoniae type 1
- (51) International Patent Classification(s)
C07H 15/04 (2006.01) **A61P 31/04** (2006.01)
A61K 31/7028 (2006.01) **C07K 19/00** (2006.01)
- (21) Application No: **2014289342** (22) Date of Filing: **2014.07.07**
- (87) WIPO No: **WO15/004041**
- (30) Priority Data
- | | | |
|----------------------------------|--------------------------------|---------------------------|
| (31) Number
13196568.3 | (32) Date
2013.12.10 | (33) Country
EP |
| 13183826.0 | 2013.09.10 | EP |
| 13175447.5 | 2013.07.07 | EP |
- (43) Publication Date: **2015.01.15**
(44) Accepted Journal Date: **2018.05.31**
- (71) Applicant(s)
Max-Planck-Gesellschaft zur Forderung der Wissenschaften e.V.
- (72) Inventor(s)
Seeberger, Peter H.;Pereira, Claney Lebev;Anish, Chakkumkal;Schumann, Benjamin
- (74) Agent / Attorney
Cullens Pty Ltd, GPO Box 1074, Brisbane, QLD, 4001, AU
- (56) Related Art
**XIANGYANG WU ET AL, CHEMISTRY - A EUROPEAN JOURNAL, (2010-03-15),
vol. 16, no. 11, doi:10.1002/chem.200902460, ISSN 0947-6539, pages 3476 - 3488**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2015/004041 A1

(43) International Publication Date
15 January 2015 (15.01.2015)

WIPO | PCT

(51) International Patent Classification:
C07H 15/04 (2006.01) *A61K 31/7028* (2006.01)
C07K 19/00 (2006.01) *A61P 31/04* (2006.01)

(21) International Application Number:
PCT/EP2014/064407

(22) International Filing Date:
7 July 2014 (07.07.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
13175447.5 7 July 2013 (07.07.2013) EP
13183826.0 10 September 2013 (10.09.2013) EP
13196568.3 10 December 2013 (10.12.2013) EP

(71) Applicant: MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V. [DE/DE]; Hofgartenstrasse 8, 80539 Munich (DE).

(72) Inventors: SEEBERGER, Peter H.; Rudolf-Breitscheid-Strasse 46, 14532 Kleinmachnow (DE). PEREIRA, Claney Lebev; Hortensiengasse 66, 12203 Berlin (DE). ANISH, Chakkumkal; Dilgesstrasse 26, 12249 Berlin (DE). SCHUMANN, Benjamin; Hauptstrasse 63, 10827 Berlin (DE).

(74) Agent: ARTH, Hans-Lothar; ABK Patent Attorneys, Jasmundweg 9, 14052 Berlin (DE).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: SYNTHETIC VACCINES AGAINST STREPTOCOCCUS PNEUMONIAE TYPE 1



(57) Abstract: The present invention relates to the total synthesis of saccharide structures contained in the capsular polysaccharide of *Streptococcus pneumoniae* type 1, to glycoconjugates containing said saccharide structures obtained by total synthesis and to use of such glycoconjugates and pharmaceutical compositions thereof in the immunization against diseases associated with bacteria containing said saccharide structures in their capsular polysaccharide, and more specifically associated with *Streptococcus pneumoniae*.

WO 2015/004041 A1

Synthetic vaccines against *Streptococcus pneumoniae* type 1

5

Field of the invention

The present invention relates to the total synthesis of saccharide structures contained in the capsular polysaccharide of *Streptococcus pneumoniae* type 1, to glycoconjugates 10 containing said saccharide structures obtained by total synthesis and to the use of such glycoconjugates and pharmaceutical compositions thereof in the immunization against diseases associated with bacteria, and more specifically against diseases associated with *Streptococcus pneumoniae*.

15 **Background of the invention**

Gram-positive encapsulated bacterium *Streptococcus pneumoniae* (pneumococcus) is a major cause of morbidity and mortality worldwide. They colonize the upper respiratory tract and cause invasive pneumococcal diseases such as meningitis, bacteremia and bactermic pneumonia, and non-invasive pneumococcal diseases 20 including acute otitis media and pneumonia. These diseases are prevalent in young children, the elderly and immunocompromised individuals of all ages. In developing countries *Streptococcus pneumoniae* related diseases cause an estimated 1.2 million deaths annually of young children.

25 Structurally, three distinguished layers can be seen on the bacterial surface: plasma membrane, cell wall and capsule. The cell wall consists of a peptidoglycan backbone anchoring the cell wall polysaccharide (C-polysaccharide) and the capsular polysaccharide (CPS). The C-polysaccharide is a structure common to all the pneumococcal serotypes, whereas CPS is specific to each of the 90 known serotypes 30 and is the main virulence factor.

Out of the 90 serotypes the most common and prevalent serotypes found in the world are shown in figure 1. This distribution varies also based on geography and age difference. Thus, a vaccine comprising glycoconjugates containing an immunogenic 35 carrier and saccharide structures derived from the capsular polysaccharide of the most common and prevalent *Streptococcus pneumoniae* serotypes would provide immunization against a high percentage of the diseases caused by this class of Gram-positive bacteria.

Several poly-valent pneumococcal vaccines were manufactured up to present. The commercially available 23-valent pneumococcal polysaccharide vaccine (PPV), contains purified capsular polysaccharide (CPS) antigens of 23 serotypes. However, this vaccine is not effective in the case of infants and young children. The currently marketed pneumococcal conjugate vaccine (PCV), PCV-7 (PrevnarTM) contains saccharides of capsular antigens of serotype 4, 6B, 9V, 14, 18C, 19F and 23F individually conjugated to diphtheria CRM₁₉₇ and is effective in infants.

The currently marketed vaccines are effective in North America and Europe for individuals of a particular age. The manufacturing process for these vaccines is complex and results in a higher price. Therefore, the vaccine is unaffordable in most developing countries. In one aspect, the present invention advantageously provides affordable synthetic saccharide vaccines that contain most of the prevalent serotypes of the developing world.

Streptococcus pneumoniae type 1 (SP1) is one of the most prevalent *S. pneumoniae* serotypes. *Streptococcus pneumoniae* type 1 capsular polysaccharide is a linear polymer having as a repeating unit: [→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp-(1→].

Synthetic saccharide structures derived from [→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp-(1→] trisaccharide repeating unit of *Streptococcus pneumoniae* type 1 capsular polysaccharide were already reported. However, the method developed by Bundle (*Chem. Eur. J.* **2010**, *16*, 3476.) provides α-methoxy saccharides, which are not suitable for conjugation to an immunogenic carrier.

In one aspect, the present invention advantageously provides an improved synthetic route to access saccharide structures functionalized with a linker, said saccharide structures being derived from [→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp-(1→] trisaccharide repeating unit of *Streptococcus pneumoniae* type 1 capsular polysaccharide. Said saccharide structures have the advantage of being functionalized with a linker thus, being suitable to be conjugated to an immunogenic carrier. Therefore, in one aspect the present invention advantageously provides glycoconjugates and pharmaceutical compositions containing said glycoconjugates for immunization against diseases associated with bacteria containing in their capsular polysaccharide one of the following structures: α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp; α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp; α-D-GalAp-(1→3)-α-D-GalAp; α-D-

GalAp; α -2,4,6-trideoxy-4-amino-D-GalNAc; α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc; α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc; α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp.

The pharmaceutical compositions comprising the saccharides of general formula (I) and/or the intermediates of general formula (II) and/or the glycoconjugates according to the present invention are for use in immunization against diseases associated with bacteria, and especially associated with *Streptococcus pneumoniae*, said diseases including pneumonia, meningitis, otitis media, bacteremia and acute exacerbation of chronic bronchitis, sinusitis, arthritis and conjunctivitis

Further advantageous features, aspects and details of the invention are evident from the dependent claims, the description, the figures, and the examples of the present application.

Definitions

The term "linker" as used herein encompasses molecular fragments capable of connecting the reducing-end monosaccharide of a saccharide with an immunogenic carrier or a solid support, optionally by binding to at least one interconnecting molecule. Thus, the function of the linker *per se* or together with the interconnecting molecule is to establish, keep and/or bridge a special distance between the reducing-end monosaccharide and an immunogenic carrier or a solid support. More specifically, one extremity of the linker is connected to the exocyclic oxygen atom at the anomeric center of the reducing-end monosaccharide and the other extremity is connected via the sulfur atom with the interconnecting molecule, or directly with the immunogenic carrier or the solid support.

As used herein, the term "interconnecting molecule" refers to a bifunctional molecule containing functional group X and functional group Y, wherein functional group X is capable of reacting with the terminal thiol group on the linker A and the functional group Y is capable of binding to an immunogenic carrier or to a solid support. Figure 2 displays examples of commercially available interconnecting molecules, but does not restrict the interconnecting molecules that can be used according to the present invention to the examples displayed herein.

The term "adjuvant" as used herein refers to an immunological adjuvant i.e. a material used in a vaccine composition that modifies or augments the effects of said vaccine by enhancing the immune response to a given antigen contained in the vaccine without being antigenically related to it. For the person skilled in the art,

5 classically recognized examples of adjuvants include:

- mineral-containing compositions, including calcium salts and aluminum salts (or mixtures thereof). Calcium salts include calcium phosphate. Aluminum salts include hydroxides, phosphates, sulfates, etc., with the salts taking any suitable form (e.g. gel, crystalline, amorphous, etc.). Adsorption to these salts is preferred. The mineral

10 containing compositions may also be formulated as a particle of metal salt. The adjuvants known as aluminum hydroxide and aluminum phosphate may be also used. The invention can use any of the "hydroxide" or "phosphate" adjuvants that are in general use as adjuvants. The adjuvants known as "aluminium hydroxide" are typically aluminium oxyhydroxide salts, which are usually at least partially crystalline.

15 The adjuvants known as "aluminium phosphate" are typically aluminium hydroxyphosphates, often also containing a small amount of sulfate (i. e. aluminium hydroxyphosphate sulfate). They may be obtained by precipitation, and the reaction conditions and concentrations during precipitation influence the degree of substitution of phosphate for hydroxyl in the salt. Mixtures of both an aluminium hydroxide and an

20 aluminium phosphate can be employed in the formulation according to the present invention;

- saponins, which are a heterologous group of sterol glycosides and triterpenoid glycosides that are found in the bark, leaves, stems, roots and even flowers of a wide range of plant species. Saponin from the bark of the *Quillaia saponaria*, Molina tree

25 have been widely studied as adjuvants. Saponin can also be commercially obtained from *Smilax ornata* (sarsaparilla), *Gypsophila paniculata* (brides veil), and *Saponaria officinalis* (soap root). Saponin adjuvant formulations include purified formulations, such as QS21, as well as lipid formulations, such as ISCOMs. Saponin compositions have been purified using HPLC and RP-HPLC. Specific purified fractions using these

30 techniques have been identified, including QS7, QS 17, QS 18, QS2 1, QH-A, QH-B and QH-C. Saponin formulations may also comprise a sterol, such as cholesterol. Combinations of saponins and sterols can be used to form unique particles called immunostimulating complexes (ISCOMs). ISCOMs generally include a phospholipid such as phosphatidylethanolamine or phosphatidylcholine. Any known

35 saponin can be used in ISCOMs. Preferably, the ISCOM includes one or more of QuilA, QHA & QHC;

- microparticles (i.e. a particle of 100 nm to 150 pm in diameter, more preferably 200 nm to 30 pm in diameter, or 500 nm to 10 pm in diameter) formed from materials that are biodegradable and non-toxic. Such non-toxic and biodegradable materials

include, but are not restricted to poly(α -hydroxy acid), polyhydroxybutyric acid, polyorthoester, polyanhydride, polycaprolactone;

- CD1d ligands, such as an α -glycosylceramide, phytosphingosine-containing α -glycosylceramides, OCH, KRN7000 [(2S,3S,4R)-1-O-(α -D-galactopyranosyl)-2-(N-

5 hexacosanoylamino)-1,3,4-octadecanetriol], CRONY- 101, 3"-sulfo-galactosylceramide;

- immunostimulatory oligonucleotides, such CpG motif containing ones (a dinucleotide sequence containing an unmethylated cytosine residue linked by a phosphate bond to a guanosine residue), or CpI motif containing ones (a dinucleotide

10 sequence containing cytosine linked to inosine), or a double-stranded RNA, or an oligonucleotide containing a palindromic sequence, or an oligonucleotide containing a poly(dG) sequence. Immunostimulatory oligonucleotides can include nucleotide modifications/analogs such as phosphorothioate modifications and can be double-stranded or (except for RNA) single-stranded;

15 - compounds containing lipids linked to a phosphate-containing acyclic backbone, such as the TLR4 antagonist E5564;

- oil emulsions (e.g. Freund's adjuvant).

Theoretically, each molecule or substance that is able to favor or amplify a particular 20 situation in the cascade of immunological events, ultimately leading to a more pronounced immunological response, can be defined as an adjuvant.

In principle, through the use of adjuvants in vaccine formulations, one can

- direct and optimize immune responses that are appropriate or desirable for the vaccine;

25 - enable mucosal delivery of vaccines, i.e. administration that results in contact of the vaccine with a mucosal surface such as buccal or gastric or lung epithelium and the associated lymphoid tissue;

- promote cell-mediated immune responses;

- enhance the immunogenicity of weaker immunogens, such as highly purified or 30 recombinant antigens;

- reduce the amount of antigen or the frequency of immunization required to provide protective immunity; and

- improve the efficacy of vaccines in individuals with reduced or weakened immune responses, such as newborns, the aged, and immunocompromised vaccine 35 recipients.

Although little is known about their mode of action, it is currently believed that adjuvants augment immune responses by one of the following mechanisms:

- increasing the biological or immunologic half-life of antigens;

- improving antigen delivery to antigen-presenting cells (APCs), as well as antigen processing and presentation by the APCs e.g., by enabling antigen to cross endosomal membranes into the cytosol after ingestion of antigen-adjuvant complexes by APC;

- 5 - mimicking danger inducing signals from stressed or damaged cells, which serve to initiate an immune response;
- inducing the production of immunomodulatory cytokines;
- biasing the immune response towards a specific subset of the immune system; and
- blocking the rapid dispersal of the antigen challenge.

10

Saccharides are known by the person skilled in the art as TI-2 (T cell independent-2) antigens and poor immunogens. Therefore, to produce a saccharide-based vaccine, said saccharides are conjugated to an immunogenic carrier to provide a glycoconjugate, which presents an increased immunogenicity in comparison with the


15 saccharide. In this context the term "immunogenic carrier" is defined as a structure, which is conjugated to the saccharide to form a glycoconjugate that presents an increased immunity in comparison with the saccharide *per se*. Thus, the conjugation of the saccharides to the immunogenic carrier has as effect the stimulation of the immune response against said saccharide, without inducing an immune response

20 against the said immunogenic carrier.

As used herein the term "**O-glycosidic bond**" refers to the covalent bond connecting the anomeric carbon of sugar fragments S1, S2 and S3 (i.e. the carbon C-1) to sugar fragments S1, S2, S3 or to -O-A-SH fragment through an **oxygen atom**.

25 In case the anomeric carbon of sugar fragments S1, S2 and S3 is connected to -O-A-SH fragment, the **oxygen atom** is the terminal oxygen atom of the fragment -O-A-SH. In case the anomeric carbon of sugar fragments S1, S2 and S3 is connected to another sugar fragment S1, S2, S3, then the **oxygen atom** is the oxygen atom at position 3 of sugar fragment S1, or the oxygen atom at position 3 of sugar fragment S2, or the oxygen atom at position 4 of sugar fragment S3. In other words, the saccharides according to the invention do not contain -O-O- bonds or sugar fragments connected or bound to each other *via* their anomeric or C-1 carbons.

Thus, the present invention relates to saccharides of general formula (I):






(I)

5

wherein A is a linker;

M, N and P represent independently of each other one of the following sugar fragments:



10

wherein the sugar fragments S1, S2, S3 are connected to each other and to $—\text{O}—\text{A}—\text{SH}$ fragment via O-glycosidic bonds, each sugar fragment S1, S2, and S3 is not more than once present in the general formula (I), sugar fragment S1 cannot be simultaneously connected to $—\text{O}—\text{A}—\text{SH}$ and sugar fragment S3, sugar fragment S3

15 cannot be simultaneously connected to $—\text{O}—\text{A}—\text{SH}$ and sugar fragment S2, and sugar fragment S2 cannot be simultaneously connected to $—\text{O}—\text{A}—\text{SH}$ and sugar fragment S1, and n_1 , n_2 and n_3 are integers selected from 0 and 1, wherein at least one of the integers n_1 , n_2 and n_3 is 1

and pharmaceutically acceptable salts of these saccharides.

20

Each sugar fragment S1, S2, and S3 is not more than once present in the general formula (I) means that each of M, N and P has to represent one of S1, S2 and S3 and none of the sugar fragments S1, S2 and S3 can be selected twice. Thus, if M is S1, then N can only be selected from S2 and S3 but cannot represent S1, and if M is


25 S1 and N is S2, P can only be S3.

The term “cannot be simultaneously connected to” refers to a direct connection. “Direct connection” means that, for instance when the sugar fragment S1 is directly connected to the fragment $—\text{O}—\text{A}—\text{SH}$, then the sugar fragment S1 is linked through 30 its anomeric carbon atom to the fragment $—\text{O}—\text{A}—\text{SH}$ and not indirectly via another sugar fragment (e.g. S3) to the fragment $—\text{O}—\text{A}—\text{SH}$. Each sugar fragment S1 or S2

or S3 can be connected through two positions, which are indicated by dotted lines. One position is the anomeric carbon C-1, which can be linked to the oxygen of the fragment $-O-A-SH$ or to an oxygen atom with a dotted line of another sugar fragment. The oxygen with a dotted line of each sugar fragment S1 or S2 or S3 5 cannot be linked to the fragment $-O-A-SH$ and can only be linked to a hydrogen atom (in case of the terminal sugar fragment) or to an anomeric carbon C-1 of another sugar fragment. However, when linking the sugar fragments and the $-O-A-SH$ fragment together, the exceptions as disclosed herein have to be taken into consideration.

10

Thus, under the scope of the present invention are falling saccharides of general formula (Ia)




Ia

15 wherein


$n_1 = n_2 = n_3 = 1$, or $n_1 = n_2 = 1$ and $n_3 = 0$, or $n_2 = n_3 = 1$ and $n_1 = 0$, or $n_1 = 1$ and $n_2 = n_3 = 0$, or $n_2 = 1$ and $n_1 = n_3 = 0$;

and saccharides of general formula (Ib)

**Ib**

wherein

- n1 = n2 = n3 = 1, or n1 = n2 = 1 and n3 = 0, or n2 = n3 = 1 and n1 = 0, or n1 = 1
 5 and n2 = n3 = 0, or n3 = 1 and n1 = n2 = 0;
 and sacharides of general formula (**Ic**)


**Ic**

wherein

n1 = n2 = n3 = 1, or n1 = n2 = 1 and n3 = 0, or n2 = n3 = 1 and n1 = 0, or n1 = 1 and n2 = n3 = 0, or n3 = 1 and n1 = n2 = 0

5 and pharmaceutically acceptable salts of these saccharides.

In other words, the present invention relates to saccharides of general formula (I):



wherein A is a linker;

15 P represents S1, N represents S3, M represents S2;

or

P represents S3, N represents S2, M represents S1;

or



P represents S2, N represents S1, M represents S3;

20 and


n1 = n2 = n3 = 1, or n1 = n2 = 1 and n3 = 0, or n2 = n3 = 1 and n1 = 0, or n1 = 1 and n2 = n3 = 0, or n2 = 1 and n1 = n3 = 0, or n3 = 1 and n1 = n2 = 0

wherein S1, S2 and S3 are sugar fragments defined as

25



S3 =



and the sugar fragments S1, S2, S3 are connected to the each other and to $-O-A-SH$ fragment via O-glycosidic bonds,
and pharmaceutically acceptable salts of these saccharides.


5

Preferably not more than one of n1, n2 and n3 is 0 or n1 = n2 = n3 = 1, and even more preferred n1 = n2 = n3 = 1. Also saccharides of general formula (I) comprising the sugar S1 are preferred. Thus, it is preferred that general formula (I) represents the following sugars: H-(S1)-(S3)-(S2)-O-A-SH, H-(S2)-(S1)-(S3)-O-A-SH,



10 H-(S3)-(S2)-(S1)-O-A-SH, H-(S2)-(S1)-O-A-SH, H-(S3)-(S2)-O-A-SH, H-(S1)-(S3)-O-A-SH, H-(S1)-O-A-SH, H-(S2)-O-A-SH and more preferred H-(S1)-(S3)-(S2)-O-A-SH, H-(S2)-(S1)-(S3)-O-A-SH, H-(S3)-(S2)-(S1)-O-A-SH, H-(S2)-(S1)-O-A-SH, H-(S1)-(S3)-O-A-SH, and H-(S1)-O-A-SH.

15

Especially preferred saccharides of general formula (I) are:



H-S1-S3-S2-O-A-SH



A is defined as a linker and is part of the fragment –O–A–SH. Thus, the linker A is bound to an oxygen atom and to an SH-group, while the oxygen atom and the SH-group are bound to different carbon atoms of the linker A. It is preferred that at least 5 two carbon atoms of the linker are between the oxygen atom and the SH-group, like –O–C–C–SH.

The linker A preferably contains between 2 and 20 carbon atoms (including the carbon atoms of optional side chains), more preferably between 2 and 18, more

preferably between 2 and 16, more preferably between 2 and 14, more preferably between 2 and 12, and still more preferably between 2 and 10 carbon atoms.

The shortest atom chain between the oxygen (i.e. the oxygen of $-\text{O}-\text{A}-\text{SH}$) and the SH-group consists preferably of 2 to 14 atoms, more preferably of 2 to 12 atoms, more preferably of 2 to 10 atoms, more preferably of 2 to 8 atoms. In case the shortest chain (which is the shortest possible connection between the oxygen and the SH-group) consists of 2 to 5 atoms, these are preferably carbon atoms. In case the shortest chain consists of 4 to 8 atoms, the chain may contain 1, 2, 3, or 4 heteroatoms selected from O, N and S. In case the shortest chain consists of 9 to 14 atoms, the chain may contain 1, 2, 3, 4, 5, or 6 heteroatoms selected from O, N and S. It is preferred that the shortest chain contains 0, 1, or 2 sulphur atoms and/or 0, 1, or 2 nitrogen atoms and/or 0, 1, 2, or 3 oxygen atoms. In case more than 4 oxygen atoms are present, preferably no other heteroatoms are present.

15

It is also preferred that the linker A, or the shortest chain is fully or partially fluorinated. The linker A may contain a 4-membered or a 5-membered or a 6-membered saturated carbocycle or a 6-membered partly unsaturated (and not aromatic) carbocycle or a 4-membered or a 5-membered or a 6-membered saturated oxygen heterocycle or a 4-membered or a 5-membered or a 6-membered saturated nitrogen heterocycle.


















The linker A may also contain amide ($-\text{NH}-\text{CO}-$, $-\text{CO}-\text{NH}-$) and/or urea ($-\text{NH}-\text{CO}-\text{NH}-$) residues and preferably only one amide or urea residue. The linker 25 may also contain substituents and preferably one substituent such as R^1 or two substituents such as R^1 and R^2 , which have the meanings as defined herein and which are preferably selected from $-\text{OCH}_3$, $-\text{OC}_2\text{H}_5$, $-\text{CH}_3$, $-\text{C}_2\text{H}_5$, $-\text{CH}_2\text{F}$, $-\text{CF}_2\text{H}$, $-\text{CF}_3$, $-\text{C}(\text{O})-\text{NH}_2$, $-\text{NHAc}$, $-\text{NH}(\text{CH}_3)$, $-\text{NH}(\text{C}_2\text{H}_5)$, $-\text{N}(\text{CH}_3)_2$, $-\text{N}(\text{C}_2\text{H}_5)_2$, $-\text{NH}-\text{C}(\text{O})-\text{CH}_3$, and $-\text{C}(\text{O})-\text{CH}_3$.

30

In case the linker is fluorinated, more than two substituents $-\text{F}$ are preferred. In case an oxygen heterocycle is present each carbon atom of the oxygen heterocycle may be substituted by a hydroxy group ($-\text{OH}$). Thus, a 5-membered oxygen heterocycle may contain one or two hydroxy groups and a 6-membered oxygen heterocycle may contain one or two or preferably 3 hydroxy groups.

The linker A, also defined herein as $-\text{A}^a-\text{A}^b-\text{A}^c-\text{A}^d-$ or $-\text{A}^a-\text{A}^b-\text{A}^d-$ contains preferably 1, 2, 3 or 4 and more preferably 1, 2, or 3 of the following fragments:

$-(CH_2)_{o1}-$, $-(CR^1R^2)_{o1}-$, $-(CH_2)_{o3}-(CH_2-CH_2-O)_{o2}-(CH_2)_{o1}-$, $-(CH_2)_{o1}-S-(CH_2)_{o4}-$,
 $-(CH_2)_{o1}-O-(CH_2)_{o4}-$, $-(CH_2)_{o1}-NH-(CH_2)_{o4}-$, $-(CH_2)_{o1}-NAC-(CH_2)_{o4}-$,
 $-(CH_2)_{o1}-C(O)-(CH_2)_{o4}-$, $-(CH_2)_{p1}-$, $-(CR^7R^8)_{p1}-$, $-(CH_2-CH_2-O)_{p1}-$, $-O-$, $-S-$,
 $-NH-$, $-C(O)-$, $-NH-C(O)-NH-$, $-NH-C(O)-(CH_2)_{p2}-$, $-C(O)-NH-(CH_2)_{p2}-$,
5 $-NH-C(O)-C_2H_4-C(O)-NH-$, $-C(O)-NH-$, $-C(O)-NH-(CH_2-CH_2-O)_{p1}-C_2H_4-$,
 $-C(O)-NH-(CH_2-CH_2-O)_{p1}-$, $-(CH_2)_{q1}-$, $-(CR^{16}R^{17})_{q1}-$, $-(CH_2)_{q1}-NH-C(O)-$,
 $-(CH_2)_{q1}-C(O)-NH-$,



wherein not two heteroatoms of the above-mentioned residues are linked together, such as a group $-\text{S}-$ is not linked to a group $-\text{NH}-\text{C}(\text{O})-\text{NH}-$;

p_2 , o_2 , o_3 are integers selected independently of each other from: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10;

5 p_1 , q_1 , o_1 , o_4 are integers selected independently of each other from: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10;

and

R^1 , R^2 , R^7 , R^8 , R^{16} , and R^{17} represent independently of each other $-\text{H}$, $-\text{OCH}_3$, $-\text{OC}_2\text{H}_5$, $-\text{CH}_3$, $-\text{C}_2\text{H}_5$, $-\text{C}_3\text{H}_7$, $-\text{CH}(\text{CH}_3)_2$, $-\text{F}$, $-\text{CH}_2\text{F}$, $-\text{CF}_2\text{H}$, $-\text{CF}_3$,

10 $-\text{C}(\text{O})-\text{NH}_2$, $-\text{SCH}_3$, $-\text{SC}_2\text{H}_5$, $-\text{NHAc}$, $-\text{NH}(\text{CH}_3)$, $-\text{NH}(\text{C}_2\text{H}_5)$, $-\text{N}(\text{CH}_3)_2$, $-\text{N}(\text{C}_2\text{H}_5)_2$, $-\text{NH}-\text{C}(\text{O})-\text{CH}_3$, and $-\text{C}(\text{O})-\text{CH}_3$.


The linker $-\text{A}-$ according to the present invention represents:


$-\text{A}^a-\text{A}^b-\text{A}^c-\text{A}^d-$ or $-\text{A}^a-\text{A}^b-\text{A}^d-$ or $-\text{A}^a-\text{A}^d-$ or $-\text{A}^a-$;

15 wherein

A^a represents $-(\text{CH}_2)_{o_1}-$, $-(\text{CR}^1\text{R}^2)_{o_1}-$, $-(\text{CR}^1\text{R}^2)_{o_1}-(\text{CR}^3\text{R}^4)_{o_2}-$,
 $-(\text{CR}^1\text{R}^2)_{o_1}-(\text{CR}^3\text{R}^4)_{o_2}-(\text{CR}^5\text{R}^6)_{o_3}-$, $-(\text{CR}^1\text{R}^2)_{o_1}-(\text{CR}^3\text{R}^4)_{o_3}-(\text{CR}^5\text{R}^6)_{o_1}-$,
 $-(\text{CR}^1\text{R}^2)_{o_1}-(\text{CR}^3\text{R}^4)_{o_2}-\text{O}-(\text{CR}^5\text{R}^6)_{o_4}-$, $-(\text{CR}^1\text{R}^2)_{o_1}-(\text{CR}^3\text{R}^4)_{o_2}-\text{S}-(\text{CR}^5\text{R}^6)_{o_4}-$,
 $-(\text{CR}^1\text{R}^2)_{o_1}-\text{S}-(\text{CR}^3\text{R}^4)_{o_4}-$, $-(\text{CR}^1\text{R}^2)_{o_1}-\text{O}-(\text{CR}^3\text{R}^4)_{o_4}-$,
 $-(\text{CR}^1\text{R}^2)_{o_1}-\text{NH}-(\text{CR}^3\text{R}^4)_{o_4}-$, $-(\text{CR}^1\text{R}^2)_{o_1}-\text{NH}-(\text{CR}^5\text{R}^6)_{o_3}-(\text{CR}^3\text{R}^4)_{o_4}-$,
 $-(\text{CR}^1\text{R}^2)_{o_1}-\text{C}(\text{O})-(\text{CR}^3\text{R}^4)_{o_4}-$, $-(\text{CR}^1\text{R}^2)_{o_1}-\text{C}(\text{O})-(\text{CR}^5\text{R}^6)_{o_3}-(\text{CR}^3\text{R}^4)_{o_4}-$,

25






















A^b represents $-(CH_2)_{p1}-$, $-(CR^7R^8)_{p1}-$, $-(CH_2-CH_2-O)_{p1}-$,
 $-(CR^7R^8)_{p1}-S-(CR^9R^{10})_{p2}-$, $-(CR^7R^8)_{p1}-O-(CR^9R^{10})_{p2}-$, $-(CR^7R^8)_{p1}-NH-(CR^9R^{10})_{p2}-$,
 $-O-$, $-S-$, $-NH-$, $-C(O)-$, $-NH-C(O)-NH-$, $-NH-C(O)-(CH_2)_{p2}-$,
5 $-C(O)-NH-(CH_2)_{p2}-$, $-NH-C(O)-C_2H_4-C(O)-NH-$,
 $-(CR^7R^8)_{p1}-(CR^9R^{10})_{p2}-(CH_2-CH_2-O)_{p3}-$, $-(CR^7R^8)_{p1}-(CH_2-CH_2-O)_{p2}-(CR^9R^{10})_{p3}-$

$-(CH_2-CH_2-O)_{p1}-(CR^7R^8)_{p2}-(CR^9R^{10})_{p3}-$,
 $-(CR^7R^8)_{p1}-(CR^9R^{10})_{p2}-(CR^{11}R^{12})_{p3}$,
 $-C(O)-NH-CH(R^{18})-C(O)-NH-CH(R^{19})-$,
 $-C(O)-NH-(CH_2-CH_2-O)_{p1}-$,


$-C(O)-NR^{15}-$,
 $-C(O)-NH-CH(R^{18})-$,
 $-C(O)-NH-(CH_2-CH_2-O)_{p1}-C_2H_4-$,









A^c represents $-(CH_2)_{q1}-$, $-(CR^{16}R^{17})_{q1}-$, $-(CH_2)_{q1}-NH-C(O)-$, $-(CH_2)_{q1}-C(O)-NH-$,





5 A^d represents $-(CH_2)_{m1}-$, $-(CR^{13}R^{14})_{m1}-$, $-CH_2-S-(CH_2)_{m1}-$, $-CH_2-O-(CH_2)_{m1}-$,
 $-CH_2-C(O)-(CH_2)_{m1}-$, $-(CH_2-CH_2-O)_{m1}-(CR^{13}R^{14})_{m2}-$, $-(CH_2)_{m1}-(CR^{13}R^{14})_{m2}-$,
 $-(CR^{13}R^{14})_{m2}-(CH_2)_{m1}-$, $-(CR^{13}R^{14})_{m1}-(CH_2)_{m2}-$, $-(CH_2)_{m2}-(O-CH_2-CH_2)_{m1}-$,



R^1 - R^{14} , R^{16} and R^{17} represent independently of each other -H, -OCH₃, -OC₂H₅, -OC₃H₇, -cyclo-C₃H₅, -cyclo-C₄H₇, -cyclo-C₅H₉, -cyclo-C₆H₁₁, -cyclo-C₇H₁₃, -cyclo-C₈H₁₅, -CH₂-cyclo-C₆H₁₁, -C(cyclo-C₆H₁₁)₃, -CH₃, -C₂H₅, -C₃H₇, -CH(CH₃)₂, -C₄H₉, -CH₂-CH(CH₃)₂, -CH(CH₃)-C₂H₅, -C(CH₃)₃, -C₅H₁₁, -CH(CH₃)-C₃H₇, -CH₂-CH(CH₃)-C₂H₅, -CH(CH₃)-CH(CH₃)₂, -C(CH₃)₂-C₂H₅, -CH₂-C(CH₃)₃, -CH(C₂H₅)₂, -C₂H₄-CH(CH₃)₂, -C₆H₁₃, -C₃H₆-CH(CH₃)₂, -C₂H₄-CH(CH₃)-C₂H₅, -CH(CH₃)-C₄H₉, -CH₂-CH(CH₃)-C₃H₇, -CH(CH₃)-CH₂-CH(CH₃)₂, -CH(CH₃)-CH(CH₃)-C₂H₅, -CH₂-CH(CH₃)-CH(CH₃)₂, -CH₂-C(CH₃)₂-C₂H₅, -C(CH₃)₂-C₃H₇, -C(CH₃)₂-CH(CH₃)₂, -C₂H₄-C(CH₃)₃, -CH(CH₃)-C(CH₃)₃, -C₇H₁₅, -C₈H₁₇, -C₆H₄-OCH₃, -CH₂-CH₂-OCH₃, -CH₂-OCH₃, -CH₂-C₆H₄-OCH₃, -F, -CH₂F, -CF₂H, -CF₃, -C(O)-NHR¹⁵, -C(O)-NHR²², -C(O)-NHR²³, -C(O)-NHR²⁴, -C(O)-NHR²⁵, -SCH₃, -SC₂H₅, -NR¹⁵R²², -NHR¹⁵, -NHR²², -NHR²³, -NHR²⁴, -NHR²⁵, -NH-C(O)-R¹⁵, -NH-C(O)-R²², -NH-C(O)-R²³, -NH-C(O)-R²⁴, -NH-C(O)-R²⁵, -C(O)-R¹⁵, -C(O)-R²², -C(O)-R²³, -C(O)-R²⁴, -C(O)-R²⁵.


R^{15} , R^{22} , R^{23} , R^{24} and R^{25} represents : -CH₃, -C₂H₅, -C₃H₇, -CH(CH₃)₂, -C₄H₉, -CH₂-CH(CH₃)₂, -CH(CH₃)-C₂H₅, -C(CH₃)₃, -C₅H₁₁, -CH(CH₃)-C₃H₇, -CH₂-CH(CH₃)-C₂H₅, -CH(CH₃)-CH(CH₃)₂, -C(CH₃)₂-C₂H₅, -CH₂-C(CH₃)₃, -CH(C₂H₅)₂, -C₂H₄-CH(CH₃)₂, -C₆H₁₃, -C₃H₆-CH(CH₃)₂, -C₂H₄-CH(CH₃)-C₂H₅, -CH(CH₃)-C₄H₉, -CH₂-CH(CH₃)-C₃H₇, -CH(CH₃)-CH₂-CH(CH₃)₂, -CH(CH₃)-CH(CH₃)-C₂H₅, -CH₂-CH(CH₃)-CH(CH₃)₂, -CH₂-C(CH₃)₂-C₂H₅, -C(CH₃)₂-C₃H₇, -C(CH₃)₂-CH(CH₃)₂, -C₂H₄-C(CH₃)₃;

25 R^{18} , R^{19} , R^{20} and R^{21} represent independently of each other -CH₂-OCH₃, -CH₂-SCH₃, -CH₂-SeCH₃, -C₂H₄-OCH₃, -C₂H₄-SCH₃, -C₂H₄-SeCH₃, -H -CH₃, -C₂H₅, -C₃H₇, -CH(CH₃)(OCH₃), -CH(CH₃)₂, -CH₂-CH(CH₃)₂, -CH₂-Ph, -CH₂-CH(CH₃)(C₂H₅), -CH₂-C(O)-NH₂, -C₂H₄-C(O)-NH₂, -CH₂-p-C₆H₄-OCH₃;

30 p2, p3, o2, o3 are integers selected independently of each other from: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10; p1, q1, o1, o4, m1, m2 are integers selected independently of each other from: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10.

35 Preferably the linker A represents the residue -A^a-.

-A^a- is preferably a linear carbon chain or a saturated carbocycle selected from




Also preferably the linker A represents $-(CH_2)_{o1}-A^b-A^c-A^d-$, $-(CH_2)_{o1}-A^b-A^d-$, $-(CH_2)_{o1}-A^d-$, $-(CR^1R^2)_{o1}-A^b-A^c-A^d-$, $-(CR^1R^2)_{o1}-A^b-A^d-$, or $-(CR^1R^2)_{o1}-A^d-$, wherein A^b , A^c and A^d have the meanings as defined herein and R^1 and R^2 have the

5 meanings as defined herein and preferably represent independently of each other $-H$, $-OCH_3$, $-OC_2H_5$, $-CH_3$, $-C_2H_5$, $-C_3H_7$, $-CH(CH_3)_2$, $-F$, $-CH_2F$, $-CF_2H$, $-CF_3$, $-C(O)-NH_2$, $-SCH_3$, $-SC_2H_5$, $-NHAc$, $-NH(CH_3)$, $-NH(C_2H_5)$, $-N(CH_3)_2$, $-N(C_2H_5)_2$, $-NH-C(O)-CH_3$, and $-C(O)-CH_3$.


10 A^b represents in general and especially in the afore-mentioned general formula $-(CH_2)_{o1}-A^b-A^c-A^d-$, $-(CH_2)_{o1}-A^b-A^d-$, $-(CR^1R^2)_{o1}-A^b-A^c-A^d-$, $-(CR^1R^2)_{o1}-A^b-A^d-$ preferably one of the following residues: $-O-$, $-S-$, $-NH-$, $-C(O)-$, $-NH-C(O)-NH-$, $-NH-C(O)-(CH_2)_{p2}-$, $-C(O)-NH-(CH_2)_{p2}-$, $-NH-C(O)-C_2H_4-C(O)-NH-$, $-C(O)-NR^{15}-$, $-C(O)-NH-CH(R^{18})-$, $-C(O)-NH-(CH_2-CH_2-O)_{p1}-C_2H_4-$, $-C(O)-NH-(CH_2-CH_2-O)_{p1}-$, wherein the substituents R^{15} and R^{18} have the meanings as defined herein and preferably $-CH_3$, $-C_2H_5$, or $-C_3H_7$.

15 A^c represents in general and especially in the afore-mentioned general formula $-(CH_2)_{o1}-A^b-A^c-A^d-$ and $-(CR^1R^2)_{o1}-A^b-A^c-A^d-$ preferably one of the following residues: $-(CH_2)_{q1}-$,



20 A^d represents in general and especially in the afore-mentioned general formula $-(CH_2)_{o1}-A^b-A^c-A^d-$, $-(CH_2)_{o1}-A^b-A^d-$, $-(CH_2)_{o1}-A^d-$, $-(CR^1R^2)_{o1}-A^b-A^c-A^d-$,

$-(CR^1R^2)_{o1}-A^b-A^d-$, and $-(CR^1R^2)_{o1}-A^d-$ preferably one of the following residues:





- 5 The function of the linker alone or together with the interconnecting molecule is to covalently connect the reducing-end of the saccharides to an immunogenic carrier or to a solid support. An interconnecting molecule according to the present invention refers to a bifunctional molecule containing functional group X and functional group Y, wherein functional group X is capable of reacting with the terminal thiol group on
 10 the linker A and the functional group Y is capable of binding to an immunogenic carrier or to a solid support. Thus, the present invention refers to saccharides of the formula (I), wherein the linker *per se* or together with the interconnecting molecule is capable of establishing, keeping and/or bridging a special distance between the reducing-end monosaccharide and an immunogenic carrier or a solid support.

15


In a preferred embodiment according to the present invention the linker $-A-$ represents $-A^a-A^b-A^c-A^d-$. Preferably, the fragment $-A^a-A^b-A^c-A^d-$ is selected from the following fragments:


20

24





Another preferred embodiment is directed to saccharides of general formula (I), wherein the linker $-A-$ represents $-A^a-A^b-A^d-$. Preferably, fragment $-A^a-A^b-A^d-$ is selected from:










Preferably the linker A represents $-A^a-A^d-$, and more preferably fragment $-A^a-A^d-$ is selected from



Preferably, the linker $-A-$ represents $-A^a-$, wherein $-A^a-$ is selected from:






- 5 $-(CH_2)_{o1}-$, $-(CR^1R^2)_{o1}-$, $-(CR^1R^2)_{o1}-(CR^3R^4)_{o2}-(CR^5R^6)_{o3}-$, $-(CR^1R^2)_{o3}-(CH_2-CH_2-O)_{o2}-(CR^3R^4)_{o1}-$,
 $-(CH_2)_{o1}-(CR^3R^4)_{o2}-S-(CH_2)_{o4}-$, $-(CH_2)_{o1}-(CR^3R^4)_{o2}-NH-(CH_2)_{o4}-$, $-(CR^1R^2)_{o1}-(CR^3R^4)_{o4}-$, $-(CR^1R^2)_{o1}-O-(CR^3R^4)_{o4}-$,
 $-(CR^1R^2)_{o1}-NH-(CR^3R^4)_{o4}$, or $-(CR^1R^2)_{o1}-C(O)-(CR^3R^4)_{o4}$.

In a preferred embodiment, substituents R^1-R^{14} , R^{16} and R^{17} are selected from:

- 10 $-H$, $-OCH_3$, $-OC_2H_5$, $-CH_3$, $-C_2H_5$, $-C_3H_7$, $-F$, $-CH_2F$, $-CF_2H$, $-CF_3$,
 $-CH(CH_3)_2$, $-C_4H_9$, $-CH_2-CH(CH_3)_2$, $-CH(CH_3)-C_2H_5$, $-C(CH_3)_3$,
 $-CH_2-CH_2-OCH_3$, $-CH_2-OCH_3$, $-SCH_3$, $-SC_2H_5$, $-NR^{15}R^{22}$, $-NHR^{15}$,
 $-NHR^{22}$, $-NHR^{23}$, $-NHR^{24}$, $-NHR^{25}$, $-NH-C(O)-R^{15}$, $-NH-C(O)-R^{22}$,
 $-NH-C(O)-R^{23}$, $-NH-C(O)-R^{24}$, $-NH-C(O)-R^{25}$.

15

- Even more preferred are linkers, wherein $-A-$ represents $-A^a-$. Preferably, fragment $-A^a-$ has the meaning: $-(CH_2)_{o1}-$, $-(CR^1R^2)_{o1}-$, $-(CH_2)_{o1}-(CR^3R^4)_{o2}-(CH_2)_{o3}-$,
 $-(CH_2-CH_2-O)_{o2}-(CH_2)_{o1}-$, $-(CH_2-CH_2-O)_{o2}-(CR^1R^2)-(CH_2)_{o1}-$,
 $-(CH_2)_{o1}-(CR^3R^4)_{o2}-S-(CH_2)_{o4}-$, $-(CR^1R^2)_{o1}-S-(CH_2)_{o4}-$, $-(CH_2)_{o1}-(CR^3R^4)_{o2}-O-(CH_2)_{o4}-$,
20 $-(CR^1R^2)_{o1}-O-(CH_2)_{o4}$, or $-(CR^1R^2)_{o1}-C(O)-(CR^3R^4)_{o4}$.



wherein

o1 and o4 are integers selected from 1, 2, 3, 4, 5, 6;

o2 and o3 are integers selected from 0, 1, 2, 3, 4, 5, 6;

and, substituents R¹-R⁶ are selected from:

- 5 -H, -OCH₃, -OC₂H₅, -CH₃, -C₂H₅, -C₃H₇, -F, -CH₂F, -CF₂H, -CF₃,
-CH(CH₃)₂, -C₄H₉, -CH₂-CH(CH₃)₂, -CH(CH₃)-C₂H₅, -C(CH₃)₃,
-CH₂-CH₂-OCH₃, -CH₂-OCH₃, -SCH₃, -SC₂H₅, -NHR¹⁵, -NHR²², -NHR²³,
-NHR²⁴, -NHR²⁵, -NH-C(O)-R¹⁵, -NH-C(O)-R²², -NH-C(O)-R²³,
-NH-C(O)-R²⁴, -NH-C(O)-R²⁵.

10

The linker A according to the current invention can be easily accessed by the person skilled in the art following procedures described in the literature.

Further, the present invention refers to synthetic saccharides of the formula (I),
15 wherein the linker is a molecular fragment capable of connecting the reducing-end monosaccharide of saccharides of general formula (I) via the thiol group with an immunogenic carrier or a solid support, optionally by binding to at least one further interconnecting molecule. More specifically, one extremity of the linker is connected to the exocyclic oxygen atom at the anomeric center of the reducing-end
20 monosaccharide and the other extremity is connected via the sulfur atom with the interconnecting molecule, or directly with the immunogenic carrier or the solid support.


The compounds of the present invention bear basic and/or acidic substituents and
25 they may form salts with organic or inorganic acids or bases. Examples of suitable acids for such acid addition salt formation are hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, acetic acid, citric acid, oxalic acid, malonic acid, salicylic acid, *p*-aminosalicylic acid, malic acid, fumaric acid, succinic acid, ascorbic acid, maleic acid, sulfonic acid, phosphonic acid, perchloric acid, nitric acid, formic acid, propionic acid, gluconic acid, lactic acid, tartaric acid, hydroxymaleic acid, pyruvic acid, phenylacetic acid, benzoic acid, *p*-aminobenzoic acid, *p*-hydroxybenzoic acid, methanesulfonic acid, ethanesulfonic acid, nitrous acid, hydroxyethanesulfonic acid, ethylenesulfonic acid, *p*-toluenesulfonic acid, naphthylsulfonic acid, sulfanilic acid, camphorsulfonic acid, china acid, mandelic acid, *o*-methylmandelic acid,
30 35 hydrogen-benzenesulfonic acid, picric acid, adipic acid, d-*o*-tolyltartaric acid, tartronic acid, (*o*, *m*, *p*)-toluic acid, naphthylamine sulfonic acid, and other mineral or carboxylic acids well known to those skilled in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner.

- Examples of suitable inorganic or organic bases are, for example, NaOH, KOH, NH₄OH, tetraalkylammonium hydroxide, lysine or arginine and the like. Salts may be prepared in a conventional manner using methods well known in the art, for example
- 5 by treatment of a solution of the compound of the general formula (I) with a solution of an acid, selected out of the group mentioned above.

Further, it is also possible that the compounds of the present invention bear simultaneously basic and acid groups. Further, it may also occur that these basic and




10 acid groups appear to be in close vicinity to one another enabling an intramolecular proton transfer from the acidic group to the basic group. Therefore, in a preferred embodiment of the present invention the compound of the formula (I) may be zwitterionic, bearing at least e.g. one O^- and one NH_3^+ group.

- 15 Thus, the present invention relates to saccharides of general formula (I):



wherein A is a linker;

- 20 M, N and P represent independently of each other one of the following sugar fragments:




- 25 wherein each sugar fragment S1, S2, and S3 is not more than once present in the general formula (I), and if sugar fragment S1 is present then its anomeric carbon can be linked only to $—\text{O}—\text{A}—\text{SH}$ or to the oxygen atom at position 4 of sugar fragment S3, and if sugar fragment S2 is present, then its anomeric carbon is linked only to $—\text{O}—\text{A}—\text{SH}$ or to the oxygen atom at position 3 of sugar fragment S1, and if sugar fragment S3 is present then its anomeric carbon can be linked only to $—\text{O}—\text{A}—\text{SH}$ or to
- 30 the oxygen atom at position 3 of sugar fragment S2;

n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1 and pharmaceutically acceptable salts of these saccharides.

5 The same connections as disclosed herein for the saccharides of general formula (I) apply also to the disulfides (i.e. the dimeric saccharides) of general formula (II) named herein as intermediates.

Thus, included under the scope of the present invention are trisaccharides of general formula: $H-(S1)-(S3)-(S2)-O-A-SH$, $H-(S2)-(S1)-(S3)-O-A-SH$,
 10 $H-(S3)-(S2)-(S1)-O-A-SH$; disaccharide of general formula: $H-(S1)-(S3)-O-A-SH$, $H-(S3)-(S2)-O-A-SH$, $H-(S2)-(S1)-O-A-SH$, and monosaccharides of general formula: $H-(S1)-O-A-SH$, $H-(S3)-O-A-SH$, wherein A is defined as a linker.

15 A preferred embodiment of the present application is directed to saccharides of general formula (I)



20 wherein A is a linker defined as above,

P represents S1,

N represents S3,


M represents S2,

sugar fragments S1, S2, S3 are connected to each other and to $-O-A-SH$

25 fragment via O-glycosidic bonds, sugar fragment S2 cannot be connected to the sugar fragment S1, and

n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1 and pharmaceutically acceptable salts of these saccharides.

30 In other words, preferred saccharides of the present invention are saccharides of general formula (I)



35 wherein A is a linker defined as above,

P represents S1,

N represents S3,

M represents S2,

sugar fragments S1, S2, S3 are connected to the each other and to $-\text{O}-\text{A}-\text{SH}$ fragment *via* O-glycosidic bonds and

n1 = n2 = n3 = 1, or n1 = n2 = 1 and n3 = 0, or n1 = 1 and n2 = n3 = 0, or n1 = 0

5 and n2 = n3 = 1, or n1 = n2 = 0 and n3 = 1.

In yet another preferred embodiment of the present invention, the compound according to the general formula (I) is selected from the group comprising or consisting of:

10

2-mercaptoproethyl O-(2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl)-(1 \rightarrow 4)-O-(α -D-galactopyranosyluronate)-(1 \rightarrow 3)-O-(α -D-galactopyranosyluronate);

2-mercaptoproethyl O-(α -D-galactopyranosyluronate)-(1 \rightarrow 3)-O-(2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl)-(1 \rightarrow 4)-O-(α -D-galactopyranosyluronate);

2-mercaptoproethyl O-(α -D-galactopyranosyluronate)-(1 \rightarrow 3)-O-(α -D-galactopyranosyluronate)-(1 \rightarrow 3)-O-(2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranoside);

2-mercaptoproethyl O-(α -D-galactopyranosyluronate)-(1 \rightarrow 3)-O-(α -D-galactopyranosyluronate);

2-mercaptoproethyl O-(2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl)-(1 \rightarrow 4)-O-(α -D-galactopyranosyluronate);


2-mercaptoproethyl O-(α -D-galactopyranosyluronate);

2-mercaptoproethyl O-(α -D-galactopyranosyluronate)-(1 \rightarrow 3)-O-(2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranoside);

2-mercaptoproethyl O-(2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranoside)

Chemical synthesis


15 Another aspect of the present invention relates to the synthesis of saccharides of general formula (I):




wherein A is a linker;

M, N and P represent independently of each other one of the following sugar


5 fragments:



S1



S2




S3

wherein sugar fragments S1, S2, S3 are connected to each other and to $-\text{O}-\text{A}-\text{SH}$ fragment via O-glycosidic bonds, each sugar fragment S1, S2, and S3 is

10 not more than once present in the general formula (I), sugar fragment S1 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{SH}$ and sugar fragment S3, sugar fragment S3 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{SH}$ and sugar fragment S2, sugar fragment S2 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{SH}$ and sugar fragment S1, and n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the

15 integers n1, n2 and n3 is 1 and comprising the steps:


A1) Reacting the compound **2** of the formula:



2


20 wherein $\text{P}^1 - \text{P}^3$ represent protecting groups,

with the compound **3** of the formula:




3

wherein P^4 represents a protecting group, in order to obtain compound **4** of general formula:

**4**


5 wherein P^1 - P^4 and A are defined as above;

and
performing removal of protecting groups P^1 - P^4 on compound **4** to afford monosaccharide disulfide **5** of general formula:

**5**

10


wherein A is defined as above, and wherein monosaccharide disulfide **5** is further treated with a reducing agent to afford monosaccharide **6** of general formula:

**6 (H-S2-O-A-SH)**

15 wherein A is defined as above;

or

performing selective deprotection on compound **4** to afford compound **7** of general formula




7

wherein P^5 is a protecting group and P^1 , P^3 , P^4 and A are defined as above.

or

5


A2) Reacting compound 8 of general formula



8

wherein P^6 and P^7 represent protecting groups, with compound 3 to afford compound 9

10 of general formula




9


wherein P^6 , P^7 and A are defined as above;

and

15 performing conversion of the azido group to acetamido group and removal of the protecting groups P^4 , P^6 and P^7 on compound 9 to afford monosaccharide disulfide 10 of general formula:

**10**

wherein A is defined as above, and wherein monosaccharide disulfide **10** is further treated with a reducing agent to afford monosaccharide **11** of general formula:




5

11 (H-S1-O-A-SH)

wherein A is defined as above;

or

10 performing selective deprotection on compound **9** to afford compound **12** of general formula:


**12**

wherein P⁴, P⁷ and A are defined as above.

or


15

A3) Reacting compound **13** of general formula



13

wherein $P^8 - P^{11}$ represent protecting groups, with compound **3** to afford compound **14** of general formula:




5

14

wherein P^4 , $P^8 - P^{11}$ are defined as above

and

10 performing selective deprotection of compound **14** to afford compound **15** of general formula:




15

wherein P^4 , P^8 , P^9 , P^{11} and A are defined as above.

15 and


B1) Reacting compound **7** with compound **13** to afford compound **16** of general formula:

**16**

wherein P^1 , P^3 - P^5 , P^8 - P^{11} and A are defined as above;


and

- 5 performing removal of protecting groups P^1 , P^3 - P^5 , P^8 - P^{11} on compound **16** to afford disaccharide disulfide **17** of general formula:

**17**

wherein A is defined as above and wherein disaccharide disulfide **17** is further treated

- 10 with a reducing agent to afford disaccharide **18** of general formula:

**18 (H-S3-S2-O-A-SH)**


wherein A is defined as above;

5

or

performing selective removal of protecting group P^{10} on compound **16** to afford compound **19** of general formula:

10


**19**

wherein P^1 , P^3 - P^5 , P^8 , P^9 , P^{11} and A are defined as above.

15

or

B2) Reacting compound **15** with compound **8** to afford compound **20** of general formula:




wherein P^4 , P^6 - P^9 , P^{11} and A are defined as above


5 and

performing conversion of the azido group to acetamido group and removal of the protecting groups P^4 , P^6 - P^9 , P^{11} on compound **20** to afford disaccharide disulfide **21** of general formula:

10



wherein A is defined as above and wherein disaccharide disulfide **21** is treated with a reducing agent to afford disaccharide **22** of general formula:



22 (H-S1-S3-O-A-SH)


wherein A is defined as above;

5

or

performing selective removal of protecting group P^6 on compound **20** to afford compound **23** of general formula:

10




23

wherein P^4 , P^7 - P^9 , P^{11} and A are defined as above;

or

15

B3) Reacting compound **12** with compound **2** to afford compound **24** of general formula:




24

wherein P^1 - P^4 , P^7 and A are defined as above,


and

- 5 performing conversion of the azido group to acetamido group and removal of protecting groups P^1 - P^4 and P^7 on compound 24 to afford disaccharide disulfide 25 of general formula:



25

- 10 wherein A is defined as above, and wherein disaccharide disulfide 25 is further treated with a reducing agent to afford disaccharide 26 of general formula:




26 (H-S2-S1-O-A-SH)

wherein A is defined as above;

or

performing selective deprotection on compound **24** to afford compound **27** of general formula:



5


27

wherein P^{12} is a protecting group and P^1 , P^3 , P^4 , P^7 and A are defined as above.

and

10

C1) Reacting compound **19** with compound **8** to afford compound **28** of general formula:



15


28

wherein P^1 , P^3 - P^9 , P^{11} and A are defined as above;

and

5

wherein protecting group P^6 is replaced with protecting group P^{13} in order to obtain compound **29** of the following chemical formula:



10


wherein P^1 , P^3 - P^5 , P^7 - P^9 , P^{11} , P^{13} and A are defined as above;

15

and

conversion of compound **29** to trisaccharide disulfide **30** by conversion of the azido group in the acetamido group and cleavage of the protecting group P^1 , P^3 - P^5 , P^7 - P^9 , P^{11} , P^{13} , wherein compound **30** is of general formula:


20



and wherein A is defined as above;

and

- 5 conversion of trisaccharide disulfide **30** to trisaccharide **31** by treatment with a reducing agent, wherein compound **31** is of general formula:




31 (H-S1-S3-S2-O-A-SH)

and wherein A is defined as above.

- 10 **or**

- C2) Reacting compound **23** with compound **2** to afford compound **32** of general formula:




32

wherein $P^1 - P^4$, $P^7 - P^9$, P^{11} and A are defined as above;

and

- 5 conversion of compound 32 to trisaccharide disulfide 33 by conversion of the azido group to the acetamido group and cleavage of the protecting group $P^1 - P^4$, $P^7 - P^9$, P^{11} , wherein compound 33 is of general formula:




33

wherein A is defined as above;

and

conversion of trisaccharide disulfide **33** to trisaccharide **34** by treatment with a reducing agent, wherein compound **34** is of general formula:




5

34 (H-S2-S1-S3-O-A-SH)

wherein A is defined as above;

or

- 10 C3) Reacting compound **27** with compound **13** to afford compound **35** of general formula:




wherein P^1 , P^3 , P^4 , P^7 - P^{11} and A are defined as above;

and

conversion of compound **35** to trisaccharide disulfide **36** by conversion of the azido group to the acetamido group and cleavage of the protecting group P^1 , P^3 , P^4 , P^7 - P^{11} ,


5 wherein compound **36** is of general formula:



wherein A is defined as above;

and

10 conversion of trisaccharide disulfide **36** to trisaccharide **37** by treatment with a reducing agent, wherein compound **37** is of general formula:



37 (H-S3-S2-S1-O-A-SH)

wherein A is defined as above.

The term “protecting group” as used herein refers to commonly used groups in organic synthesis, preferably used for protection of amines, hydroxyl groups, thiols, imines, carbonyls, carboxyls or other common functional groups, and particularly preferred for amines, hydroxyl groups, thiols and carboxyls.

5

More specifically, P¹, P², P⁵, P⁶, P⁸-P¹⁰, P¹² and P¹³ preferably are suitable protecting groups for hydroxyl groups, more preferably different suitable protecting groups for hydroxyl groups capable of being removed subsequently one after another by a suitable sequence of deprotection reactions. Therefore, protecting groups for

10 hydroxyl groups, namely P¹, P², P⁵, P⁶, P⁸-P¹⁰, P¹² and P¹³ may be selected from the group consisting of or comprising: acetyl, benzyl, isopropylidene, benzylidene, benzoyl, *p*-methoxybenzyl, *p*-methoxybenzylidene, *p*-methoxyphenyl, *p*-bromobenzyl, *p*-bromobenzylidene, *p*-nitrophenyl, allyl, acetyl, isopropyl, *p*-bromobenzyl dimethoxytrityl, trityl, 2-naphthylmethyl, pyvaloyl, triisopropylsilyl, *tert*-
15 butyldimethylsilyl, *tert*-butyldiphenylsilyl, *tert*-butylmethoxphenylsilyl, triethylsilyl, trimethylsilyl, 2-trimethylsilylethoxymethyl, 9-fluorenylmethoxycarbonyl, benzyloxymethyl, methyloxymethyl, *tert*-butyloxymethyl, methoxyethyloxymethyl, levulinoyl.

20 More specifically, in a preferred embodiment of the present invention protecting groups P¹, P⁵, P⁸, P⁹ and P¹² are benzyl, P² is benzylidene, P⁶ is levulinoyl, P¹⁰ is 9-fluorenylmethoxycarbonyl and P¹³ is benzyloxymethyl.

25 Amines are generally protected as carbamates. Therefore, protecting group P⁷ may be selected from the group consisting of or comprising *tert*-butyloxy carbonyl, 9-fluorenylmethoxy carbonyl, allyloxy carbonyl, 2,2,2-trichloroethoxy carbonyl, benzyloxy carbonyl. In a preferred embodiment of the present invention P⁷ is benzyloxy carbonyl.


30 Carboxylic acids are generally protected as esters. Therefore protecting groups P³ and P¹¹ may be selected from the group consisting of or comprising methyl, ethyl, allyl, isopropyl, *tert*-butyl, phenyl, benzyl, *p*-methoxybenzyl. In a preferred embodiment of the present invention protecting groups P³ and P¹¹ are methyl.

35 Protecting groups for hydroxyl groups may serve as well as protecting groups for thiols. Therefore, preferred protecting groups for thiols groups are benzyl, benzoyl, 4-O-*p*-methoxybenzyl, allyl, acetyl, methylsulfonylethoxycarbonyl, levulinyl, dimethoxytrityl, 2-naphthylmethyl, triisopropylsilyl, *tert*-butyldimethylsilyl, *tert*-

butyldiphenylsilyl, 2-trimethylsilylethoxymethyl. Specifically, in a preferred embodiment of the present invention protecting group P⁴ is a benzyl group.

5 The protecting groups employed in the synthesis of saccharides of general formula (I) can be differentiated in permanent protecting groups and temporary protecting groups. Permanent protecting groups are protecting groups that are stable during the entire synthesis and that can be efficiently removed at the late stage of the synthesis. Such permanent groups include, but they are not restricted to benzyl, benzylidene, benzoate, acetate, alkyl esters. The temporary protecting groups are generally 10 orthogonal protecting groups that can be selectively removed at different levels of the synthesis to free hydroxyl groups for subsequent introduction of different substituents, including monosaccharides or other protecting groups. The ingenious choice of protecting groups allows expedient access to a library of saccharides of general formula (I) functionalized with a thiol group for subsequent conjugation to a 15 carrier immunogen or a solid support.

A preferred embodiment of the present invention is directed to the synthesis of saccharides of general formula (I)



wherein A is a linker defined as above,

P represents S1,

25 N represents S3,

M represents S2,

sugar fragments S1, S2, S3 are connected to the each other and to —O—A—SH fragment *via* O-glycosidic bonds, sugar fragment S2 cannot be connected to sugar fragment S1, and

30 n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1


and

comprising the steps:

35


A1) Reacting the compound 2 of the formula:

50




wherein $P^1 - P^3$ represent protecting groups,
with the compound **3** of the formula

5



wherein P^4 represents a protecting group, in order to obtain compound **4** of general formula:

10




wherein P^1-P^4 and A are defined as above;

and


15

performing removal of protecting groups P^1-P^4 on compound **4** to afford monosaccharide disulfide **5** of general formula:



20

wherein A is defined as above, and wherein monosaccharide disulfide **5** is further treated with a reducing agent to afford monosaccharide **6** of general formula:




6 (H-S2-O-A-SH)

5 wherein A is defined as above;

or

10 performing selective deprotection on compound **4** to afford compound **7** of general formula




7

wherein P⁵ is a protecting group and P¹, P³, P⁴ and A are defined as above.


15 or

A2) Reacting compound **8** of general formula



8


20 wherein P⁶ and P⁷ represent protecting groups, with compound **3** to afford compound **9** of general formula



wherein P^6 , P^7 and A are defined as above;


5 and

performing conversion of the azido group to acetamido group and removal of the protecting groups P^4 , P^6 and P^7 on compound 9 to afford monosaccharide disulfide 10 of general formula:



10

wherein A is defined as above and, wherein monosaccharide disulfide 10 is further treated with a reducing agent to afford monosaccharide 11 of general formula:




15

11 (H-S1-O-A-SH)

wherein A is defined as above.


or

A3) Reacting compound **13** of general formula



wherein $P^8 - P^{11}$ represent protecting groups, with compound **3** to afford compound **14**

5 of general formula:




14

wherein $P^4, P^8 - P^{11}$ are defined as above

10 and

performing selective deprotection of compound **14** to afford compound **15** of general formula:




15

wherein P^4, P^8, P^9, P^{11} and A are defined as above.

and

B1) Reacting compound **7** with compound **13** to afford compound **16** of general formula:




5

wherein P^1 , P^3 - P^5 , P^8 - P^{11} and A are defined as above;

and


10

performing removal of protecting groups P^1 , P^3 - P^5 , P^8 - P^{11} on compound **16** to afford disaccharide disulfide **17** of general formula:



15

wherein A is defined as above and wherein disaccharide disulfide **17** is further treated with a reducing agent to afford disaccharide **18** of general formula:

**18 (H-S3-S2-O-A-SH)**


wherein A is defined as above;

5

or

performing selective removal of protecting group P^{10} on compound **16** to afford compound **19** of general formula:

10


**19**

wherein P^1 , P^3 - P^5 , P^8 , P^9 , P^{11} and A are defined as above.

15

or

B2) Reacting compound **15** with compound **8** to afford compound **20** of general formula:




wherein P^4 , P^6 - P^9 , P^{11} and A are defined as above;


5 and

performing conversion of the azido group to acetamido group and removal of the protecting groups P^4 , P^6 - P^9 , P^{11} on compound **20** to afford disaccharide disulfide **21** of general formula:

10




wherein A is defined as above and wherein disaccharide disulfide **21** is treated with a
15 reducing agent to afford disaccharide **22** of general formula:

**22 (H-S1-S3-O-A-SH)**

wherein A is defined as above.

5 and

C1) Reacting compound **19** with compound **8** to afford compound **28** of general formula:



10


28

wherein P¹, P³-P⁹, P¹¹ and A are defined as above;

and

wherein protecting group P^6 is replaced with protecting group P^{13} in order to obtain compound **29** of the following chemical formula:


5

**29**

10 wherein P^1 , P^3 - P^5 , P^7 - P^9 , P^{11} , P^{13} and A are defined as above;

and


15 conversion of compound **29** to trisaccharide disulfide **30** by conversion of the azido group to the acetamido group and cleavage of the protecting group P^1 , P^3 - P^5 , P^7 - P^9 , P^{11} , P^{13} , wherein compound **30** is of general formula:



and wherein A is defined as above;

5 and

conversion of trisaccharide disulfide **30** to trisaccharide **31** by treatment with a reducing agent, wherein compound **31** is of general formula:



10

and wherein A is defined as above.

The conversion of the saccharide disulfides **36**, **33**, **30**, **25**, **21**, **17**, **10** and **5** to the saccharides **37**, **34**, **31**, **26**, **22**, **18**, **11**, and **6** respectively is performed in presence of a reducing agent. Known reducing agent for the person skilled in the art include, but they are not restricted to: mercaptoethanol, ditriotheritol, tris(2-carboxyethyl)phosphine, magnesium/methanol, sodium/ammonia followed by ammonium chloride/chlorhydric acid. Preferably the conversion of the saccharide disulfides to the corresponding saccharides is carried out with tris(2-carboxyethyl)phosphine.

- 10 It is preferred that the reaction between compounds **2** and **3**, compounds **2** and **12**, compounds **5*** and **11***, compounds **19*** and **21***, compounds **2** and **23** is performed in presence of (dimethylthio)methylsulfonium trifluoromethanesulfonate (DMTST) and 2,4,6-tri-*tert*-butylpyridine (TTBPy) in a mixture of non-polar solvent and polar aprotic solvent. In addition activated molecular sieve (MS) such as 3Å molecular sieve, 4Å molecular sieves or 3Å acid washed molecular sieves can be used. The reaction temperature is between –20 °C and room temperature, preferably the temperature is between –10 °C and room temperature, more preferably the temperature is between –5 °C and room temperature and most preferably the temperature is between 0 °C and room temperature. Preferred polar aprotic solvents are tetrahydrofuran, diethyl ether and dioxane. Preferred non-polar solvents are toluene, halogenated solvents such as chloroform and methylene chloride. Preferred mixtures of non-polar and polar aprotic solvent are: methylene chloride / tetrahydrofuran, methylene chloride / diethyl ether, toluene / diethyl ether, toluene/ tetrahydrofuran.
- 15
- 20
- 25 It is also preferred that the reaction of compounds **13** and **7**, compounds **8** and **3**, compounds **12*** and **9***, compounds **13*** and **2***, compounds **2*** and **21***, compounds **2*** and **11***, compounds **19** and **8**, compounds **27** and **13** and compounds **8** and **15**, is performed in a non polar solvent or a mixture of non polar and polar aprotic solvents in presence of silyl triflate. Examples of silyl triflate include, but are not restricted to trimethylsilyl trifluoromethanesulfonate, *tert*-butyl dimethyl trifluoromethanesulfonate, triisopropyl trifluoromethanesulfonate. Suitable non-polar solvents are toluene, chloroform and methylene chloride. Preferred polar aprotic solvents are tetrahydrofuran, diethyl ether and dioxane. The reaction temperature is between –20 °C and +20 °C, preferably the temperature is between –10 °C and +10 °C and more preferably the temperature is between –5 °C and +5 °C and most preferably about 0 °C. Preferably, activated molecular sieve (MS) such as 3Å molecular sieve, 4Å molecular sieves or 3Å acid washed molecular sieves can be used.
- 30
- 35

Preferably the replacement of protecting group P^6 on compound **28** with protecting group P^{13} to obtain compound **29** is performed in two steps, first involving the reaction of compound **28** with hydrazine or a hydrazinium salt in a solvent or mixture of solvents and second by treatment of the product obtained after the first step with 5 $BnOCH_2SCy$, DMTST and TTBPy in a non-polar solvent. For the first step, hydrazinium salts of weak acids are preferred such as hydrazinium acetate or hydrazinium propionate. Suitable solvents for this reaction are non-polar solvents, such as methylene chloride, polar solvents such as pyridine and acetic acid, and mixtures thereof. The second step is preferably conducted in presence of activated 10 molecular sieves as the one mentioned above at a temperature preferably between -10 °C and 20 °C and most preferably between 0 °C and 10 °C. Suitable non-polar solvents are mentioned above. It was found that the replacement of protecting group P^6 with protecting group P^{13} is essential for avoiding side-reactions during the cleavage of the permanent protecting group.

15

The conversion of compound **4** to **5**, **16** to **17**, **13*** to **27*** and **20*** to **22*** requires cleavage of the protecting group, and more specifically of the permanent protecting groups. The cleavage of said protecting groups involves first cleavage of the base-labile protecting groups by treatment with a base in a mixture of polar aprotic and 20 polar protic solvent; and second cleavage of the protecting groups sensitive to hydrogenation by exposure to sodium and ammonia in a mixture of polar protic and polar aprotic solvents. For the first step suitable aprotic solvents include tetrahydrofuran, ethyl acetate, acetone, *N,N*-dimethylformamide, acetonitrile and *N,N*-dimethylsulfoxide, which are mixed with a suitable protic solvent such as water 25 and alcohols including methanol, ethanol, propanol, isopropanol or *tert*-butanol.

The basic cleavage of the protecting groups is preferably performed at room temperature comprised between 0 °C and room temperature. Appropriate base for performing first step include lithium hydroxide, sodium hydroxide and potassium hydroxide. The cleavage of the protecting groups sensitive to hydrogenation is 30 conducted by exposure to sodium and ammonia in a mixture of polar protic and polar aprotic solvents at a temperature comprised between -78 °C and room temperature. Optionally, lithium can be used as equivalent of sodium during the cleavage of the protecting groups sensitive to hydrogenation.

35 Previous to the above mentioned cleavage of permanent protecting groups, the conversion of compounds **9** to **10**, **20** to **21**, **24** to **25**, **29** to **30**, **32** to **33**, **36** to **37**, and **16*** to **18*** involves conversion of the azido group in the acetamido group, which is preferably performed in presence of thioacetic acid and pyridine. An alternative method is to conduct conversion of the azido group in the acetamido group in two

steps: first chemoselective reduction of the azido group, and then acetylation. The chemoselective reduction can be carried out using by hydrogenolysis on Pd/C in presence of ammonia, ammonium acetate, triphenylphosphine or pyridine. The acetylation can be accomplished using acetyl chloride or acetic anhydride in presence of a base.

The saccharides I, 5, 6, 10, 11, 17, 18, 21, 22, 25, 26, 30, 31, 33, 34, 36, 37, 27*, 26*, 24*, 22* and 18* bear basic and/or acidic substituents and they may form salts with organic or inorganic acids or bases. Examples of suitable acids for such acid addition salt formation are hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, acetic acid, citric acid, oxalic acid, malonic acid, salicylic acid, *p*-aminosalicylic acid, malic acid, fumaric acid, succinic acid, ascorbic acid, maleic acid, sulfonic acid, phosphonic acid, perchloric acid, nitric acid, formic acid, propionic acid, gluconic acid, lactic acid, tartaric acid, hydroxymaleic acid, pyruvic acid, phenylacetic acid, benzoic acid, *p*-aminobenzoic acid, *p*-hydroxybenzoic acid, methanesulfonic acid, ethanesulfonic acid, nitrous acid, hydroxyethanesulfonic acid, ethylenesulfonic acid, *p*-toluenesulfonic acid, naphthylsulfonic acid, sulfanilic acid, camphorsulfonic acid, china acid, mandelic acid, *o*-methylmandelic acid, hydrogen-benzenesulfonic acid, picric acid, adipic acid, *d*-*o*-tolyltartaric acid, tartronic acid, (*o*, *m*, *p*)-toluic acid, naphthylamine sulfonic acid, and other mineral or carboxylic acids well known to those skilled in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner.

The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their corresponding salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the salts are otherwise equivalent to their corresponding free base forms for purposes of this invention.

Examples for suitable inorganic or organic bases are, for example, NaOH, KOH, NH₄OH, tetraalkylammonium hydroxide, lysine or arginine and the like. Salts may be prepared in a conventional manner using methods well known in the art, for example by treatment of a solution of the compound of the general formula (I) with a solution of an acid, selected out of the group mentioned above.

Further, it is also possible that the compounds of the present invention bear simultaneously basic and acid groups. Further, it may also occur that these basic and acid groups appear to be in close vicinity to one another enabling an intramolecular

proton transfer from the acidic group the basic group. Therefore, in a preferred embodiment of the present invention the compound of the formula (I) may be zwitterionic, bearing at least e.g. one $-\text{O}^-$ and one $-\text{NH}_3^+$ group.

- 5 This invention includes within its scope stoichiometric solvates including hydrates as well as compounds containing variable amounts of water that may be produced by processes such as lyophilisation.

10 Thus, the synthesis of the saccharides of the general formula (I) may further comprise step D:


D) preparing a salt of the compound of general formula (I) or preparing a lyophilisate of the compound of general formula (I) or of the salt of the compound of general formula (I).

15 In another preferred embodiment, the synthesis of intermediates of general formula (II) may further comprise step D:

D) preparing a salt of the compound of general formula (II) or preparing a lyophilisate of the compound of general formula (II) or of the salt of the compound of general formula (II).

20

In a preferred embodiment, the synthesis of saccharides of the general formula (I)



25 wherein A is a linker defined as above,

P represents S1,

N represents S3,

M represents S2,

sugar fragments S1, S2, S3 are connected to each other and to $-\text{O}-\text{A}-\text{SH}$

30 fragment via O-glycosidic bonds, sugar fragment S2 cannot be connected to sugar fragment S1, and

n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1

and

35


may further comprise step D)

- D) preparing a salt of the compound of general formula (I) or preparing a lyophilisate of the compound of general formula (I) or of the salt of the compound of general formula (I).

5




Intermediates

- 10 Another aspect of the present invention is directed to intermediates of general formula (II):



- 15 wherein A is a linker;

M, N and P represent independently of each other one of the following fragments:




- 20 wherein sugar fragments S1, S2, S3 are connected to each other and to $-\text{O}-\text{A}-\text{S}-$ fragment via O-glycosidic bonds, each sugar fragment S1, S2, and S3 is not more than once present in the fragment $\text{H}-(\text{P})_{n3}-(\text{N})_{n2}-(\text{M})_{n1}-\text{O}-\text{A}-\text{S}-$, sugar fragment S1 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{S}-$ and sugar fragment S3, sugar fragment S3 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{S}-$ and sugar fragment S2, sugar fragment S2 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{S}-$ and sugar fragment S1, and

n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1, and pharmaceutically acceptable salts of these saccharides.

30


Thus, under the scope of the present invention are falling intermediates of general formula (IIa)



10 wherein

n1 = n2 = n3 = 1, or n1 = n2 = 1 and n3 = 0, or n2 = n3 = 1 and n1 = 0, or n1 = 1 and n2 = n3 = 0, or n2 = 1 and n1 = n3 = 0;

and intermediates of general formula (IIb)




IIb

5

Wherein

10 $n_1 = n_2 = n_3 = 1$, or $n_1 = n_2 = 1$ and $n_3 = 0$, or $n_2 = n_3 = 1$ and $n_1 = 0$, or $n_1 = 1$ and $n_2 = n_3 = 0$, or $n_3 = 1$ and $n_1 = n_2 = 0$;

and intermediates of general formula (IIc)



IIIc


Wherein

- 5 $n_1 = n_2 = n_3 = 1$, or $n_1 = n_2 = 1$ and $n_3 = 0$, or $n_2 = n_3 = 1$ and $n_1 = 0$, or $n_1 = 1$ and $n_2 = n_3 = 0$, or $n_3 = 1$ and $n_1 = n_2 = 0$

and pharmaceutically acceptable salts of these saccharides.

10

In other words, the present invention relates to intermediates of general formula (II):



15

wherein A is a linker;

P represents S1, N represents S3, M represents S2;

or

P represents S3, N represents S2, M represents S1;

or

5


P represents S2, N represents S1, M represents S3, and

wherein


$n_1 = n_2 = n_3 = 1$, or $n_1 = n_2 = 1$ and $n_3 = 0$, or $n_2 = n_3 = 1$ and $n_1 = 0$, or $n_1 = 1$ and $n_2 = n_3 = 0$, or $n_2 = 1$ and $n_1 = n_3 = 0$, or $n_3 = 1$ and $n_1 = n_2 = 0$;

10 wherein


S1 =



S2 =




S3 =



and the sugar fragments S1, S2, S3 are connected to each other and to
 $-O-A-S-$ fragment via O-glycosidic bonds,
and pharmaceutically acceptable salts of these saccharides.

15

Preferred are intermediates of general formula (II)



wherein

20 A is a linker defined as above,

P represents S1,

N represents S3,

M represents S2,


sugar fragments S1, S2, S3 are connected to the each other and to -O-A-S- fragment *via* O-glycosidic bonds and sugar fragment S2 cannot be connected to

5 sugar fragment S1, and

n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1 and

pharmaceutically acceptable salts of these saccharides.

10 More preferred are intermediates of general formula (II)



wherein A is a linker defined as above,

P represents S1,

15 N represents S3,

M represents S2,

sugar fragments S1, S2, S3 are connected to the each other and to -O-A-S- fragment *via* O-glycosidic bonds and

wherein n1 = n2 = n3 = 1 or wherein n1 = n2 = 1 and n3 = 0 or wherein n1 = 1

20 and n2 = n3 = 0 or wherein n1 = 0 and n2 = n3 = 1 or wherein n1 = n2 = 0 and n3 = 1.

Even more preferred are the intermediates:

H-(S1)-(S3)-(S2)-O-A-S-S-A-O-(S2)-(S3)-(S1)-H,

25 H-(S2)-(S1)-(S3)-O-A-S-S-A-O-(S3)-(S1)-(S2)-H,

H-(S3)-(S2)-(S1)-O-A-S-S-A-O-(S1)-(S2)-(S3)-H,

H-(S1)-(S3)-O-A-S-S-A-O-(S3)-(S1)-H,

H-(S3)-(S2)-O-A-S-S-A-O-(S2)-(S3)-H,

H-(S2)-(S1)-O-A-S-S-A-O-(S1)-(S2)-H, H-(S1)-O-A-S-S-A-O-(S1)-H, and

30 H-(S3)-O-A-S-S-A-O-(S3)-H. Particularly preferred are intermediates of general formula (II) comprising the sugar fragment S1, such as:

H-(S1)-(S3)-(S2)-O-A-S-S-A-O-(S2)-(S3)-(S1)-H,

H-(S2)-(S1)-(S3)-O-A-S-S-A-O-(S3)-(S1)-(S2)-H,

H-(S3)-(S2)-(S1)-O-A-S-S-A-O-(S1)-(S2)-(S3)-H,

35 H-(S1)-(S3)-O-A-S-S-A-O-(S3)-(S1)-H,

H-(S2)-(S1)-O-A-S-S-A-O-(S1)-(S2)-H, and H-(S1)-O-A-S-S-A-O-(S1)-H.

Glycoconjugates

Another aspect of the present invention refers to a glycoconjugate obtained by reacting 5 a saccharide of general formula (I), (Ia), (Ib) and (Ic) with an immunogenic carrier. Said glycoconjugate proved to be efficient as a vaccine for immunization against diseases associated with bacteria. Hence, glycoconjugates comprising a saccharide of general formula (I), (Ia), (Ib) and (Ic) covalently linked to an immunogenic carrier are useful for raising a protective immune response in a human 10 and/or animal host and therefore are useful for the prevention and/or treatment of diseases associated with bacteria.

Saccharides are known by the person skilled in the art as generally TI-2 (T cell independent-2) antigens and poor immunogens. TI-2 antigens are antigens, which 15 are recognized only by mature B cells through the cross linking of surface exposed immunoglobulin receptors. Without T cell help, no immunological memory is generated and neither isotype switching from IgM to other IgG subclasses, nor B cells affinity maturation occurs. Moreover, saccharides are known poor 20 immunogens in humans due to the structural homology to human glycolipids and glycoproteins. Due to their poor immunogenic properties, saccharides manifest poor ability to produce both antibody production by B cells, as well as the formation of memory cells, features which are essential for the production of potent vaccines.

Therefore, to produce a potent saccharide-based vaccine, the saccharides of general 25 formula (I), (Ia), (Ib) and (Ic) are conjugated to an immunogenic carrier to provide glycoconjugates, which present increased immunogenicity in comparison with the saccharide.

In this context the term "immunogenic carrier" is defined as a structure, which is 30 conjugated to the saccharide to form a glycoconjugate that presents an increased immunity in comparison with the saccharide *per se*. Thus, the conjugation of the saccharides of general formula (I), (Ia), (Ib) and (Ic) to the immunogenic carrier has as effect the stimulation of the immune response against the saccharide of general formula (I), (Ia), (Ib) or (Ic) without inducing an immune response against the said 35 immunogenic carrier.

Preferred immunogenic carriers are carrier proteins or glycosphingolipid with immunomodulatory properties. For the person skilled in the art, a carrier protein is a protein selected from the group comprising or consisting of: a diphtheria toxoid, a

mutated diphtheria toxoid, a modified diphtheria toxoid, a mutated and modified diphtheria toxoid, a tetanus toxoid, a modified tetanus toxoid, a mutated tetanus toxoid, outer membrane protein (OMP), bovine serum albumin (BSA), keyhole limpet hemocyanine (KLH) or cholera toxin (CT). The term "toxoid" as used herein refers to a bacterial toxin (usually an exotoxin), whose toxicity has been inactivated or suppressed either by chemical (formalin) or heat treatment, while other properties, typically immunogenicity, are maintained. A mutated toxoid as used herein is a recombinant bacterial toxin, which has been amended to be less toxic or even non-toxic by amending the wild-type amino acid sequence. Such a mutation could be a substitution of one or more amino acids. Such a mutated toxoid presents on its surface a functionality that can react with the functional group Y of the interconnecting molecule to provide a modified toxoid. Said functionality is known to the person skilled in the art and includes, but is not restricted to the primary amino functionality of a lysine residue that can react with activated esters, an isocyanate group or an aldehyde in presence of a reducing agent, to the carboxylate functionality of a glutamate or aspartate residue that can be activated by carbodiimides or to the thiol functionality of a cysteine residue. Activated esters include N-(γ-maleimidobutyryloxy) sulfosuccinimide ester (sulfo-GMBS), succinimidyl (4-iodoacetyl) aminobenzoate (sulfo-SIAB), succinimidyl-3-(bromoacetamido)propionate (SBAP), disuccinimidyl glutarate (DSG), 2-pyridyldithiol-tetraoxatetradecane-N-hydroxysuccinimide (PEG-4-SPDP) (see **Figure 2**). The cysteine residue on the carrier protein can be converted to the corresponding dehydroalanine that can be further reacted with a suitable interconnecting molecule to provide modified carrier protein having on their surface the functional group X of the interconnecting molecule. In this case the functional group Y on the interconnecting molecule might be a thiol group and the group X might be an alkene. Such interconnecting molecules include allylmercaptan. After reaction with such interconnecting molecule, the carrier protein is converted to a modified carrier protein presenting the vinyl group X of the interconnecting molecule, which is suitable to react with the saccharides of general formula (I), (Ia), (Ib) and (Ic).

It is especially preferred that the saccharides of general formula (I), (Ia), (Ib) and (Ic) and preferably saccharides 37, 34, 31, 26, 22, 18, 11 and 6 are conjugated to the non-toxic mutated diphtheria toxin CRM₁₉₇ presenting as a functionality a primary amine functionality of a lysine residue.

CRM₁₉₇ like wild-type diphtheria toxin is a single polypeptide chain of 535 amino acids (58 kD) consisting of two subunits linked by disulfide bridges having a single

amino acid substitution of glutamic acid for glycine. It is utilized as a carrier protein in a number of approved conjugate vaccines for diseases such as Prevnar.

Thus, in a preferred embodiment of the present invention the carrier protein presents 5 on its surface primary amino functionalities of lysine residues that are able to react with the functional group Y of the interconnecting molecule to provide modified carrier protein having on their surface said functional group X of the interconnecting molecule, which is able to react with the terminal thiol group of the linker of the 10 compounds of general formula (I). Said functional group X of the interconnecting molecules is selected of the group comprising or consisting of maleimide; α -iodoacetyl; α -bromoacetyl; *N*-hydroxysuccinimide ester (NHS), 2-pyridylthiols, thiol and vinyl (see **Figure 3**).

Preferably, the saccharide of general formula (I), (Ia), (Ib) or (Ic) is conjugated to the 15 non-toxic mutated diphtheria toxin CRM₁₉₇, which is modified by maleimide. In yet another preferred embodiment, the saccharide of general formula (I), (Ia), (Ib) or (Ic) is conjugated to the non-toxic mutated diphtheria toxin CRM₁₉₇, which is modified by vinyl. In the most preferred embodiment, the saccharide of general formula (I), (Ia), (Ib) or (Ic) is conjugated to the non-toxic mutated diphtheria toxin CRM₁₉₇, which is 20 modified by α -bromoacetamide.

In another embodiment, said immunogenic carrier is preferably a glycosphingolipid with immunomodulatory properties, and more preferably (2S,3S,4R)-1-(α -D-galactopyranosyl)-2-hexacosanoylaminoctadecane-3,4-diol. The term 25 glycosphingolipid with immunomodulatory properties, as used herein, refers to a suitable glycosphingolipid capable of stimulating the immune system's response to a target antigen, but which does not in itself confer immunity as defined above.

Glycosphingolipids as used herein are compounds containing a carbohydrate moiety 30 α -linked to a sphingolipid. Preferably, the carbohydrate moiety is a hexopyranose and most preferably is α -D-galactopyranose. For the person skilled in the art, sphingolipids are a class of lipids containing a C18 amino alcohol connected via an amide bond to a fatty acid. The C18 amino alcohol is preferably mono-, di- or polysubstituted with hydroxyl groups. Especially preferred, the C18 amino alcohol is 35 phytosphingosine. The fatty acid is preferably a monocarboxylic acid having a saturated alkyl chain of a number of carbons ranging from 16 to 28 and more preferably from 18 to 26. Glycosphingolipids with immunomodulatory properties include, but they are not restricted to (2S,3S,4R)-1-(α -D-galactopyranosyl)-2-hexacosanoylaminoctadecane-3,4-diol, which can stimulate natural killer (NK)

activity and cytokine production by natural killer T (NKT) cells and exhibits potent antitumor activity *in vivo* (*Proc. Natl Acad. Sci. USA*, **1998**, 95, 5690).

5 The conjugates of the saccharides of general formula (I), (Ia), (Ib) or (Ic) with a glycosphingolipid with immunomodulatory properties have the advantage of being heat stable. To be suitable for conjugation, on the glycosphingolipid with immunomodulatory properties a functionality is introduced. Said functionality is prone to react directly with the terminal thiol group of the linker of the saccharides of general formula (I), (Ia), (Ib) or (Ic) to provide glycoconjugates of the saccharides of 10 general formula (I), (Ia), (Ib) or (Ic), or with the functional group Y of the interconnecting molecule to provide the modified glycosphingolipid with immunomodulatory properties.

15 Preferably, said functionality is introduced at the C6 of the carbohydrate moiety of the glycosphingolipid with immunomodulatory properties. Thus, the glycosphingolipid with immunomodulatory properties is functionalized with a functionality, which is prone of reacting with thiol group, activated ester, isocyanate group, aldehyde, vinyl, amino group and azido group to provide directly the glycoconjugate of the saccharides of general formula (I) or the modified glycosphingolipid with 20 immunomodulatory properties presenting the functional group X of the interconnecting molecule.

25 Preferably, the functionality introduced at the C6 of the carbohydrate moiety of the glycosphingolipid with immunomodulatory properties is selected from the group comprising or containing an amine, a thiol, an alcohol, a carboxylic acid, a vinyl, maleimide, α -iodoacetyl, α -bromoacetyl, *N*-hydroxysuccinimide ester (NHS), 2-pyridyldithiols.

30 Said functional group X of the interconnecting molecules is selected of the group comprising or consisting of maleimide, α -iodoacetyl, α -bromoacetyl, *N*-hydroxysuccinimide ester (NHS), 2-pyridyldithiols, thiol and vinyl.

35 As used herein, the term “interconnecting molecule” refers to a bifunctional molecule containing functional group X and functional group Y, wherein functional group X is capable of reacting with the terminal thiol group on the linker A and the functional group Y is capable of binding to an immunogenic carrier or to a solid support.

It was found that the glycoconjugates obtained by reacting the saccharides of general formula (I), (Ia), (Ib) and (Ic) with an immunogenic carrier are suitable to elicit an

immune response in an animal, and therefore are useful as a vaccine in immunization against diseases associated with bacteria containing in their capsular polysaccharide a saccharide structure selected from:

α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp
α-D-GalAp-(1→3)-α-D-GalAp
α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp

5

Preferably, the bacterium containing in the capsular polysaccharide one of the above mentioned saccharide structures is *Streptococcus pneumoniae* type 1.

10 In a preferred embodiment, the glycoconjugates obtained by reacting the saccharides of general formula (I), (Ia), (Ib) and (Ic) with an immunogenic carrier are useful as a vaccine for immunization against diseases associated with bacteria, wherein said diseases include pneumonia, meningitis, otitis media, bacteremia and acute exacerbation of chronic bronchitis, sinusitis, arthritis and conjunctivitis

15 One aspect of the present invention relates to pharmaceutical compositions, especially vaccines comprising at least one glycoconjugate obtained by reacting any saccharide of general formula (I) with an immunogenic carrier, and/or one saccharide of general formula (I) and/or a intermediate of general formula (II), together with at least one pharmaceutically acceptable cryoprotectant, lyoprotectant, excipient and/or diluent.

20 Said vaccine may be prepared in the form of a suspension or may be lyophilized. The suspension form may be stored frozen. In the lyophilized form, it is preferable to add one or more stabilizers. Optionally, one or more adjuvants may be added as well. Any conventional stabilizers and adjuvants may be included in a vaccine according to this invention.

25 The term "adjuvant" as used herein refers to an immunological adjuvant i.e. a material used in a vaccine composition that modifies or augments the effects of said vaccine by enhancing the immune response to a given antigen contained in the vaccine without being antigenically related to it. For the persons skilled in the art,

classically recognized examples immunological adjuvants include but are not restricted to oil emulsions (e.g. Freund's adjuvant), saponins, aluminium or calcium salts (e.g. alum), non-ionic block polymer surfactants, and many others.

5 Vaccination can be performed at any age. The vaccine may be administered subcutaneously, by spray, by injection, orally, intraocularly, intratracheally or nasally.

Another aspect of the present invention relates to pharmaceutical formulations and pharmaceutical compositions containing the vaccine as an active ingredient, together
10 with at least one pharmaceutically acceptable carrier, excipient, solvent and/or diluents.

Further preferred, the pharmaceutical composition is formulated in the form of a lyophilisate or liquid buffer solution.

15

The vaccine can also be administered in form of its pharmaceutically active salt optionally using substantially nontoxic pharmaceutically acceptable carrier, excipients, adjuvants or diluents. The vaccine of the present invention is prepared in a conventional solid or liquid carrier or diluents and a conventional pharmaceutically-
20 made adjuvant at suitable dosage level in a known way. The preferred preparations and formulations are in administrable form, which is suitable for oral application. These administrable forms, for example, include pills, tablets, film tablets, coated tablets, capsules, powders and deposits. Other than oral administratable forms are also possible. The inventive vaccine may be administered by any appropriate
25 means, including but not limited to inhalation; injection (intravenous, intraperitoneal, intramuscular, subcutaneous); by absorption through epithelial or mucocutaneous linings (oral mucosa, rectal and vaginal epithelial linings, nasopharyngial mucosa, intestinal mucosa); orally, rectally, transdermally, topically, intradermally, intragastrically, intracutaneously, intravaginally, intravasally, intranasally,
30 intrabuccally, percutaneously, sublingually, or any other means available within the pharmaceutical arts.

The vaccine of the present invention, containing the glycoconjugate obtained by reacting the saccharide of general formula (I) with an immunogenic carrier or
35 pharmaceutically acceptable salts thereof as an active ingredient will typically be administered in admixture with suitable carrier materials suitably selected with respect to the intended form of administration, i.e. oral tablets, capsules (either solid-filled, semi-solid filled or liquid filled), powders for constitution, oral gels, elixirs, dispersible granules, syrups, suspensions, and the like, and consistent with

conventional pharmaceutical practices. For example, for oral administration in the form of tablets or capsules, the active ingredient may be combined with any oral nontoxic pharmaceutically acceptable inert carrier, such as lactose, starch, sucrose, cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, talc, mannitol, 5 ethyl alcohol (liquid forms) and the like. Moreover, when desired or needed, suitable binders, lubricants, disintegrating agents and coloring agents may also be incorporated in the mixture.

10 Suitable binders include starch, gelatin, natural sugars, corn sweeteners, natural and synthetic gums such as acacia, sodium alginate, carboxymethyl-cellulose, polyethylene glycol and waxes. Among the lubricants that may be mentioned for use in these dosage forms, boric acid, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrants include starch, methylcellulose, guar gum and the like. Sweetening and flavoring agents and preservatives may also be included 15 where appropriate. Some of the terms noted above, namely disintegrants, diluents, lubricants, binders and the like, are discussed in more detail below.

20 Additionally, the vaccine of the present invention may be formulated in sustained release form to provide the rate controlled release of any one or more of the components or active ingredients to optimize the therapeutic effects. Suitable dosage forms for sustained release include layered tablets containing layers of varying disintegration rates or controlled release polymeric matrices impregnated with the active components and shaped in tablet form or capsules containing such impregnated or encapsulated porous polymeric matrices.

25 30 Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injections or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

35 Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier such as inert compressed gas, e.g. nitrogen.

For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides such as cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein by stirring or similar mixing. The molten homogeneous

mixture is then poured into convenient sized molds, allowed to cool and thereby solidifies.

Also included are solid form preparations that are intended to be converted, shortly 5 before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The vaccine of the present invention may also be deliverable transdermally. The transdermal compositions may take the form of creams, lotions, aerosols and/or 10 emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

The term capsule refers to a special container or enclosure made of methyl cellulose, polyvinyl alcohols, or denatured gelatins or starch for holding or containing 15 compositions comprising the active ingredients. Hard shell capsules are typically made of blends of relatively high gel strength bone and pork skin gelatins. The capsule itself may contain small amounts of dyes, opaques, plasticizers and preservatives.

20 Tablet means compressed or molded solid dosage form containing the active ingredients with suitable diluents. The tablet can be prepared by compression of mixtures or granulations obtained by wet granulation, dry granulation or by compaction well known to a person skilled in the art.

25 Oral gels refer to the active ingredients dispersed or solubilized in a hydrophilic semi-solid matrix.

Powders for constitution refer to powder blends containing the active ingredients and suitable diluents which can be suspended in water or juices.

30 Suitable diluents are substances that usually make up the major portion of the composition or dosage form. Suitable diluents include sugars such as lactose, sucrose, mannitol and sorbitol, starches derived from wheat, corn, rice and potato, and celluloses such as microcrystalline cellulose. The amount of diluents in the 35 composition can range from about 5 to about 95% by weight of the total composition, preferably from about 25 to about 75%, more preferably from about 30 to about 60% by weight, and most preferably from about 40 to 50% by weight.

The term disintegrants refers to materials added to the composition to help it break apart (disintegrate) and release the medicaments. Suitable disintegrants include starches, "cold water soluble" modified starches such as sodium carboxymethyl starch, natural and synthetic gums such as locust bean, karaya, guar, tragacanth and agar, cellulose derivatives such as methylcellulose and sodium carboxymethylcellulose, microcrystalline celluloses and cross-linked microcrystalline celluloses such as sodium croscarmellose, alginates such as alginic acid and sodium alginate, clays such as bentonites, and effervescent mixtures. The amount of disintegrant in the composition can range from about 1 to about 40% by weight of the composition, preferably 2 to about 30% by weight of the composition, more preferably from about 3 to 20% by weight of the composition, and most preferably from about 5 to about 10% by weight.

Binders characterize substances that bind or "glue" powders together and make them cohesive by forming granules, thus serving as the "adhesive" in the formulation. Binders add cohesive strength already available in the diluents or bulking agent. Suitable binders include sugars such as sucrose, starches derived from wheat, corn rice and potato; natural gums such as acacia, gelatin and tragacanth; derivatives of seaweed such as alginic acid, sodium alginate and ammonium calcium alginate; cellulosic materials such as methylcellulose and sodium carboxymethylcellulose and hydroxypropyl-methylcellulose; polyvinylpyrrolidone; and inorganics such as magnesium aluminum silicate. The amount of binder in the composition can range from about 1 to 30% by weight of the composition, preferably from about 2 to about 20% by weight of the composition, more preferably from about 3 to about 10% by weight, even more preferably from about 3 to about 6% by weight.

Lubricant refers to a substance added to the dosage form to enable the tablet, granules, etc. after it has been compressed, to release from the mold or die by reducing friction or wear. Suitable lubricants include metallic stearates such as magnesium stearate, calcium stearate or potassium stearate; stearic acid; high melting point waxes; and water soluble lubricants such as sodium chloride, sodium benzoate, sodium acetate, sodium oleate, polyethylene glycols and D, L-leucine. Lubricants are usually added at the very last step before compression, since they must be present on the surfaces of the granules and in between them and the parts of the tablet press. The amount of lubricant in the composition can range from about 0.05 to about 15% by weight of the composition, preferably 0.2 to about 5% by weight of the composition, more preferably from about 0.3 to about 3%, and most preferably from about 0.3 to about 1.5% by weight of the composition.

Glidants are materials that prevent caking and improve the flow characteristics of granulations, so that flow is smooth and uniform. Suitable glidants include silicon dioxide and talc. The amount of glidant in the composition can range from about 0.01 to 10% by weight of the composition, preferably 0.1% to about 7% by weight of 5 the total composition, more preferably from about 0.2 to 5% by weight, and most preferably from about 0.5 to about 2% by weight.

Coloring agents are excipients that provide coloration to the composition or the dosage form. Such excipients can include food grade dyes and food grade dyes 10 adsorbed onto a suitable adsorbent such as clay or aluminum oxide. The amount of the colouring agent can vary from about 0.01 to 10% by weight of the composition, preferably from about 0.05 to 6% by weight, more preferably from about 0.1 to about 4% by weight of the composition, and most preferably from about 0.1 to about 1%.

15 Techniques for the formulation and administration of the vaccine of the present invention may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton PA. A suitable vaccine composition comprising at least one glycoconjugate of the present invention and/or pharmaceutically acceptable salts thereof may be a solution of one glycoconjugate obtained by reacting any saccharide 20 of general formula (I) with an immunogenic carrier in a suitable liquid pharmaceutical carrier or any other formulation such as tablets, pills, film tablets, coated tablets, dragees, capsules, powders and deposits, gels, syrups, slurries, suspensions, emulsions, and the like.

25 A therapeutically effective dosage of one glycoconjugate obtained by reacting any saccharide of general formula (I) with an immunogenic carrier refers to that amount of the compound that results in an at least a partial immunization against a disease. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical, pharmacological, and toxicological procedures in cell cultures or 30 experimental animals. The dose ratio between toxic and therapeutic effect is the therapeutic index. The actual amount of the composition administered will be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgement of the prescribing physician.

35 Another preferred embodiment of the present invention is directed to pharmaceutical composition comprising the glycoconjugate obtained by reacting any saccharide of general formula (I) with an immunogenic carrier, and/or the saccharide of general formula (I), and/or the intermediate of general formula (II) together with at least one 40 one pharmaceutically acceptable cryoprotectant, lyoprotectant, excipient and/or

diluent. Said pharmaceutical composition are useful in immunization against diseases associated with *Streptococcus pneumoniae* bacteria.

Streptococcus pneumoniae bacteria referred herein include the following serotypes

- 5 *Streptococcus pneumoniae* type 1, *Streptococcus pneumoniae* type 4, *Streptococcus pneumoniae* type 9V, *Streptococcus pneumoniae* type 2, *Streptococcus pneumoniae* type 19F, *Streptococcus pneumoniae* type 3, *Streptococcus pneumoniae* type 19A, *Streptococcus pneumoniae* type 12F, *Streptococcus pneumoniae* type 31, *Streptococcus pneumoniae* type 7F, *Streptococcus pneumoniae* type 5, *Streptococcus pneumoniae* type 14, *Streptococcus pneumoniae* type 6A, *Streptococcus pneumoniae* type 6B, *Streptococcus pneumoniae* type 18C and *Streptococcus pneumoniae* type 23F.
- 10
- 15
- 20

A preferred embodiment of the present invention is directed to a pharmaceutical composition, especially a vaccine comprising the glycoconjugate obtained by reacting any saccharide of general formula (I) with an immunogenic carrier and/or the saccharide of general formula (I), and/or the intermediate of general formula (II) together with at least one one pharmaceutically acceptable cryoprotectant, lyoprotectant, excipient and/or diluents for use in immunization against diseases associated with *Streptococcus pneumoniae* type 1.

The saccharides derived from $[\rightarrow 3]-\alpha-2,4,6\text{-trideoxy-4-amino-D-GalNAc-(1\rightarrow 4)-\alpha-D-GalAp-(1\rightarrow 3)-\alpha-D-GalAp-(1\rightarrow 3)-\alpha-D-GalAp]$ are functionalized with suitable linker, which allows their conjugation to an immunogenic carrier, as defined herein to provide glycoconjugates. Said glycoconjugates proved to be efficient as a vaccine for immunization against diseases associated with bacteria, and in particularly against diseases associated with *Streptococcus pneumoniae*, and particularly against *Streptococcus pneumoniae* type 1. Said diseases include pneumonia, meningitis, otitis media, bacteremia and acute exacerbation of chronic bronchitis, sinusitis, arthritis and conjunctivitis.

Yet another aspect of the present invention refers to saccharide of general formula (I) and/or intermediate of general formula (II) for use as a marker in immunological assays for diagnostics of diseases caused by bacteria containing in the capsular polysaccharide a saccharide structure selected from:

$\alpha-2,4,6\text{-trideoxy-4-amino-D-GalNAc-(1\rightarrow 4)-\alpha-D-GalAp-(1\rightarrow 3)-\alpha-D-GalAp}$
 $\alpha-2,4,6\text{-trideoxy-4-amino-D-GalNAc-(1\rightarrow 4)-\alpha-D-GalAp}$
 $\alpha-D-GalAp-(1\rightarrow 3)-\alpha-D-GalAp$

α -D-GalAp

α -2,4,6-trideoxy-4-amino-D-GalNAc

α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc

α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc

α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp

Such assays comprise, for instance, microarray and ELISA useful for diagnosis of diseases caused by bacteria containing or comprising the saccharides of the present invention, said diseases including pneumonia, meningitis, otitis media, bacteremia and acute exacerbation of chronic bronchitis, sinusitis, arthritis and conjunctivitis.


The saccharides of the present invention can be easily conjugated to solid supports for providing assays useful for diagnosis of diseases caused by bacteria containing or comprising the saccharides of the present invention. Said solid supports present

10 on their surface a functionality that is prone to react with the functional group Y of the interconnecting molecule to provide modified solid supports, presenting on their surface the functional group X of the interconnecting molecule, which are able to react with the thiol group of saccharides of general formula (I). Said solid supports include, but are not restricted to microarray slides, which present on their surface a
15 functionality that is prone to react with the functional group Y of the interconnecting molecule to provide modified microarray slides, presenting of their surface the functional group X of the interconnecting molecule. Preferably, the microarray slides present on their surface an amino group. Microarray slides presenting on their surface an amino group include, but are not restricted to amine-coated GAPS II
20 slides (Corning) or CodeLink NHS slides on which the amino functionality was introduced by incubation with Bovin Serum Albumin (BSA).

Microarray slides coated with the saccharides of general formula (I) were synthesized by conjugating the saccharides of general formula (I) to said modified microarray
25 slides, and incubated with rabbit anti-*Streptococcus pneumoniae* type 1 typing serum human pneumococcal serum 007sp in the presence or absence of native *Streptococcus pneumoniae* type 1 polysaccharide. The binding experiments show that both rabbit anti-*Streptococcus pneumoniae* type 1 typing serum and human pneumococcal serum 007sp bound to α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -
30 D-GalAp-(1 \rightarrow 3)- α -D-GalAp and α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp saccharide structures (see **Figures 4-8**). Moreover, the binding could be inhibited with the native *Streptococcus pneumoniae* type 1 polysaccharide, suggesting that the saccharides according to the present invention share epitopes that are recognized by the immune system (see **Figure 5** and **Figure 6**).

Definitions of the specific embodiments of the invention as claimed herein follow.


According to a first embodiment of the invention, there is provided a saccharide of general formula (I):



wherein A represents $(CH_2)_{01}$;


o1 is an integer selected from 1, 2, 3, 4, 5 and 6;

M, N and P represent independently of each other one of the following sugar fragments:



wherein the sugar fragments S1, S2, S3 are connected to each other and to $-O-A-SH$ fragment via O-glycosidic bonds, each sugar fragment S1, S2, and S3 is not more than once present in the general formula (I), sugar fragment S1 cannot be simultaneously connected to $-O-A-SH$ and sugar fragment S3, sugar fragment S3 cannot be simultaneously connected to $-O-A-SH$ and sugar fragment S2, and sugar fragment S2 cannot be simultaneously connected to $-O-A-SH$ and sugar fragment S1, and n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1 and pharmaceutically acceptable salts of these saccharides.




According to a second embodiment of the invention, there is provided a synthesis of the saccharide of the general formula (I):



wherein A represents $(\text{CH}_2)_{01}$;


o1 is an integer selected from 1, 2, 3, 4, 5 and 6;

M, N and P represent independently of each other one of the following sugar fragments:




wherein the sugar fragments S1, S2, S3 are connected to each other and to $-O-A-SH$ fragment via O-glycosidic bonds, each sugar fragment S1, S2, and S3 is not more than once present in the general formula (I), sugar fragment S1 cannot be simultaneously connected to $-O-A-SH$ and sugar fragment S3, sugar fragment S3 cannot be simultaneously connected to $-O-A-SH$ and sugar fragment S2, and sugar fragment S2 cannot be simultaneously connected to $-O-A-SH$ and sugar fragment S1, and n_1, n_2 and n_3 are integers selected from 0 and 1, wherein at least one of the integers n_1, n_2 and n_3 is 1 and comprising the steps:

A1) Reacting the compound **2** of the formula:




wherein $P^1 - P^3$ represent protecting groups, with the compound **3** of the formula



wherein P^4 represents a protecting group, in order to obtain compound **4** of general formula:

2014289342 30 Apr 2018




4

wherein P¹-P⁴ and A are defined as above;


and

performing removal of protecting groups P¹-P⁴ on compound 4 to afford monosaccharide disulfide 5 of general formula:



5

wherein A is defined as above, and wherein monosaccharide disulfide 5 is further treated with a reducing agent to afford monosaccharide 6 of general formula:




6 (H-S2-O-A-SH)

wherein A is defined as above;

or

performing selective deprotection on compound 4 to afford compound 7 of general formula

2014289342 30 Apr 2018




7

wherein P^5 is a protecting group and P^1 , P^3 , P^4 and A are defined as above.


or

A2) Reacting compound **8** of general formula



8

wherein P^6 and P^7 represent protecting groups, with compound **3** to afford compound **9** of general formula




9


wherein P^6 , P^7 and A are defined as above;

and

performing conversion of the azido group to acetamido group and removal of the protecting groups P^4 , P^6 and P^7 on compound **9** to afford monosaccharide disulfide **10** of general formula:



wherein A is defined as above, and wherein monosaccharide disulfide **10** is further treated with a reducing agent to afford monosaccharide **11** of general formula:




11 (H-S1-O-A-SH)

wherein A is defined as above;

or

performing selective deprotection on compound **9** to afford compound **12** of general formula:




12


wherein P⁴, P⁷ and A are defined as above.

or

2014289342 30 Apr 2018

A3) Reacting compound **13** of general formula**13**


wherein $P^8 - P^{11}$ represent protecting groups, with compound **3** to afford compound **14** of general formula:

**14**

wherein $P^4, P^8 - P^{11}$ are defined as above

and


performing selective deprotection of compound **14** to afford compound **15** of general formula:

**15**

wherein P^4, P^8, P^9, P^{11} and A are defined as above.

and


B1) Reacting compound **7** with compound **13** to afford compound **16** of general formula:



wherein P^1 , P^3 - P^5 , P^8 - P^{11} and A are defined as above;


and

performing removal of protecting groups P^1 , P^3 - P^5 , P^8 - P^{11} on compound **16** to afford disaccharide disulfide **17** of general formula:




wherein A is defined as above and wherein disaccharide disulfide **17** is further treated with a reducing agent to afford disaccharide **18** of general formula:

81h

**18 (H-S3-S2-O-A-SH)**


wherein A is defined as above;

or

performing selective removal of protecting group P¹⁰ on compound **16** to afford compound **19** of general formula:**19**wherein P¹, P³-P⁵, P⁸, P⁹, P¹¹ and A are defined as above.

or


B2) Reacting compound **15** with compound **8** to afford compound **20** of general formula:



20

wherein P^4 , P^6 - P^9 , P^{11} and A are defined as above
and


performing conversion of the azido group to acetamido group and removal of the protecting groups P^4 , P^6 - P^9 , P^{11} on compound **20** to afford disaccharide disulfide **21** of general formula:



21

wherein A is defined as above and wherein disaccharide disulfide **21** is treated with a reducing agent to afford disaccharide **22** of general formula:

2014289342 30 Apr 2018




22 (H-S1-S3-O-A-SH)

wherein A is defined as above;

or

performing selective removal of protecting group P⁶ on compound **20** to afford compound **23** of general formula:




23

wherein P⁴, P⁷-P⁹, P¹¹ and A are defined as above;

or

B3) Reacting compound **12** with compound **2** to afford compound **24** of general formula:


2014289342 30 Apr 2018

**24**


wherein P¹-P⁴, P⁷ and A are defined as above,

and

performing conversion of the azido group to acetamido group and removal of protecting groups P¹-P⁴ and P⁷ on compound **24** to afford disaccharide disulfide **25** of general formula:

**25**


wherein A is defined as above, and wherein disaccharide disulfide **25** is further treated with a reducing agent to afford disaccharide **26** of general formula:

**26 (H-S2-S1-O-A-SH)**

wherein A is defined as above;

or

performing selective deprotection on compound **24** to afford compound **27** of general formula:




27

wherein P^{12} is a protecting group and P^1 , P^3 , P^4 , P^7 and A are defined as above.

and

C1) Reacting compound **19** with compound **8** to afford compound **28** of general formula:




28

wherein P^1 , P^3 - P^9 , P^{11} and A are defined as above;

and


wherein protecting group P^6 is replaced with protecting group P^{13} in order to obtain compound **29** of the following chemical formula:



wherein P^1 , P^3-P^5 , P^7-P^9 , P^{11} , P^{13} and A are defined as above;

and


conversion of compound **29** to trisaccharide disulfide **30** by conversion of the azido group in the acetamido group and cleavage of the protecting group P^1 , P^3-P^5 , P^7-P^9 , P^{11} , P^{13} , wherein compound **30** is of general formula:



and wherein A is defined as above;

and


conversion of trisaccharide disulfide **30** to trisaccharide **31** by treatment with a reducing agent, wherein compound **31** is of general formula:



and wherein A is defined as above.

or

C2) Reacting compound **23** with compound **2** to afford compound **32** of general formula:




32

wherein $P^1 - P^4$, $P^7 - P^9$, P^{11} and A are defined as above;

and

conversion of compound 32 to trisaccharide disulfide 33 by conversion of the azido group in the acetamido group and cleavage of the protecting group $P^1 - P^4$, $P^7 - P^9$, P^{11} , wherein compound 33 is of general formula:




33

wherein A is defined as above;

and

conversion of trisaccharide disulfide **33** to trisaccharide **34** by treatment with a reducing agent, wherein compound **34** is of general formula:




34 (H-S2-S1-S3-O-A-SH)

wherein A is defined as above;

or

C3) Reacting compound **27** with compound **13** to afford compound **35** of general formula:




35

wherein P¹, P³, P⁴, P⁷-P¹¹ and A are defined as above;

and


conversion of compound **35** to trisaccharide disulfide **36** by conversion of the azido group in the acetamido group and cleavage of the protecting group P¹, P³, P⁴, P⁷-P¹¹, wherein compound **36** is of general formula:



wherein A is defined as above;


and

conversion of trisaccharide disulfide **36** to trisaccharide **37** by treatment with a reducing agent, wherein compound **37** is of general formula:



wherein A is defined as above.




According to a third embodiment of the invention, there is provided an intermediate of the general formula (II):



wherein A represents $(\text{CH}_2)_{o1}$;

$o1$ is an integer selected from 1, 2, 3, 4, 5 and 6;

M, N and P represent independently of each other one of the following fragments:



wherein the sugar fragments S1, S2, S3 are connected to each other and to $-\text{O}-\text{A}-\text{S}-$ fragment via O-glycosidic bonds, each sugar fragment S1, S2, and S3 is not more than once present in the fragment $\text{H}-(\text{P})_{n3}-(\text{N})_{n2}-(\text{M})_{n1}-\text{O}-\text{A}-\text{S}-$,

sugar fragment S1 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{S}-$ and sugar fragment S3, sugar fragment S3 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{S}-$ and sugar fragment S2, sugar fragment S2 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{S}-$ and sugar fragment S1, and

$n1$, $n2$ and $n3$ are integers selected from 0 and 1, wherein at least one of the integers $n1$, $n2$ and $n3$ is 1

and pharmaceutically acceptable salts of these saccharides.

According to a fourth embodiment of the invention, there is provided a glycoconjugate obtained by reacting a saccharide of general formula (I) according to the first embodiment with an immunogenic carrier.

According to a fifth embodiment of the invention, there is provided a pharmaceutical composition comprising the glycoconjugate according to the fourth embodiment and/or the saccharide according to the first embodiment and/or the intermediate according to the third embodiment together with at least one pharmaceutically acceptable cryoprotectant, lyoprotectant, excipient and/or diluent.

2014289342 30 Apr 2018

According to a sixth embodiment of the invention, there is provided a saccharide according to the first embodiment and/or a intermediate according to the third embodiment for use as marker in immunological assays for diagnostics of diseases caused by bacteria containing in their capsular polysaccharide a saccharide structure selected from:

α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp
α-D-GalAp-(1→3)-α-D-GalAp
α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp.

According to a seventh embodiment of the invention, there is provided a method of vaccinating a subject against diseases associated with bacteria containing in their capsular polysaccharide a saccharide structure selected from:

α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp
α-D-GalAp-(1→3)-α-D-GalAp
α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp,
comprising administering to said subject the glycoconjugate according to the fourth embodiment.

According to an eighth embodiment of the invention, there is provided a use of the glycoconjugate according to the fourth embodiment in the manufacture of a vaccine for immunization against diseases associated with bacteria containing in their capsular polysaccharide a saccharide structure selected from:

α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp
α-D-GalAp-(1→3)-α-D-GalAp
α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc

2014289342 30 Apr 2018

α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc
 α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc
 α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp.

According to a ninth embodiment of the invention, there is provided a method of immunizing a subject against diseases associated with *Streptococcus pneumoniae* bacteria, comprising administering to said subject the pharmaceutical composition according to the fifth embodiment.

According to a tenth embodiment of the invention, there is provided a use of the glycoconjugate according to the fourth embodiment and/or the saccharide according to the first embodiment and/or the intermediate according to the third embodiment in the manufacture of a medicament for immunization against diseases associated with *Streptococcus pneumoniae* bacteria.

According to an eleventh embodiment of the invention, there is provided a use of the saccharide according to the first embodiment and/or the intermediate according to the third embodiment as a marker in immunological assays for diagnostics of diseases caused by bacteria containing in their capsular polysaccharide a saccharide structure selected from:

α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp
 α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp
 α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp
 α -D-GalAp
 α -2,4,6-trideoxy-4-amino-D-GalNAc
 α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc
 α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc
 α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp.

The term 'comprise' and variants of the term such as 'comprises' or 'comprising' are used herein to denote the inclusion of a stated integer or stated integers but not to exclude any other integer or any other integers, unless in the context or usage an exclusive interpretation of the term is required.

30 Apr 2018

2014289342

Any reference to publications cited in this specification is not an admission that the disclosures constitute common general knowledge in Australia.

[text continues on page 82]

Description of the figures

- 5 **Figure 1** shows the global distribution of *Streptococcus pneumoniae* serotypes
- 10 **Figure 2** provides examples of commercially available interconnecting molecules according to the present invention.
- 15 **Figure 3** provides examples of functional group X of the interconnecting molecule according to the present invention.
- 20 **Figure 4** shows the printing pattern of the saccharides of general formula (I) on microarray slides
- 25 **Figure 5** shows the binding of human pneumococcal serum 007sp (pooled sera of 287 humans immunized with Pneumovax vaccine) to saccharides of general formula (I), which are coated on modified CodeLink NHS slides in presence and in absence of native *Streptococcus pneumoniae* type 1 polysaccharide.
- 30 **Figure 6** shows the binding of the rabbit anti-*Streptococcus pneumoniae* type 1 typing serum to saccharides of general formula (I), which are coated on modified amine-coated GAPS II slides (Corning) in presence and in absence of native *Streptococcus pneumoniae* type 1 polysaccharide.
- 35 **Figure 7** shows the binding of the rabbit anti-*Streptococcus pneumoniae* type 1 typing serum to saccharides of general formula (I), which are coated on modified amine-coated GAPS II slides (Corning).
- 40 **Figure 8** shows the binding of the rabbit anti-*Streptococcus pneumoniae* type 1 typing serum to saccharides of general formula (I), which are coated on modified CodeLink NHS slides.

35 The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those skilled in the art that the techniques disclosed in the examples, which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those skilled in the art should, in light of the present disclosure, appreciate that many changes can be made

in the specific embodiments, which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.


Further modifications and alternative embodiments of various aspects of the invention will be apparent to those skilled in the art in view of this description. Accordingly, this description is to be construed as illustrative only and is for the purpose of teaching those skilled in the art the general manner of carrying out the invention. It is to be understood that the forms of the invention shown and described herein are to be taken as examples of embodiments. Elements and materials may be substituted for those illustrated and described herein, parts and processes may be reversed, and certain features of the invention may be utilized independently, all as would be apparent to one skilled in the art after having the benefit of this description of the invention. Changes may be made in the elements described herein without departing from the spirit and scope of the invention as described in the following claims.

Examples

Chemical synthesis


General information for chemical synthesis. Commercial reagents were used without further purification except where noted. Solvents were dried and redistilled prior to use in the usual way. All reactions were performed in oven-dried glassware under an inert atmosphere unless noted otherwise. Analytical thin layer chromatography (TLC) was performed on Kieselgel 60 F254 glass plates precoated with a 0.25 mm thickness of silica gel. The TLC plates were visualized with UV light and by staining with Hanessian solution (ceric sulfate and ammonium molybdate in aqueous sulfuric acid) or sulfuric acid-ethanol solution. Column chromatography was performed on Fluka Kieselgel 60 (230-400 mesh). Optical rotations (OR) were measured with a Schmidt & Haensch UniPol L1000 polarimeter at a concentration (c) expressed in g/100 mL. ¹H and ¹³C NMR spectra were measured with a Varian 400-MR or Varian 600 spectrometer with Me₄Si as the internal standard. NMR chemical shifts (δ) were recorded in ppm and coupling constants (J) were reported in Hz. High-resolution mass spectra (HRMS) were recorded with an Agilent 6210 ESI-TOF mass spectrometer at the Freie Universität Berlin, Mass Spectrometry Core Facility.

Example 1: 4-(Benzylloxycarbonyl)amino-3-O-levulinoyl-4,6-dideoxy-D-galactal (1⁺):



To a stirred solution of 4-O-(benzyloxycarbonyl)amino-3-hydroxy-4,6-dideoxy-D-galactal (*Org. Lett.* **2010**, 12, 1624) (1.64 g, 6.21 mmol) (1.64 g, 6.21 mmol) in CH_2Cl_2 (40 ml) were added at 0 °C pyridine (0.501 ml, 6.21 mmol), levulinic acid (0.96 ml, 9.31 mmol), DMAP (0.152 g, 1.242 mmol) and EDC (1.205 ml, 6.83 mmol).
 5 The mixture was warmed to room temperature and stirred at that temperature. After 3 h, 0.5 eq. levulinic acid and 0.5 eq. EDC were added to drive the reaction to completion. After 5 h, the mixture was diluted with 100 ml DCM and washed with water (50 ml), sat. aq. NH_4Cl (50 ml), sat. aq. NaHCO_3 (50 ml) and brine (50 ml). The
 10 organic fraction was dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 1:1) to give ester **1*** (2.07 g, 5.73 mmol, 92%) as a clear oil. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{23}\text{NO}_6$ ($\text{M}+\text{Na}^+$) 384.1423 found 384.1415 *m/z*.

15 **Example 2:** Dibutyl [2-azido-4-(benzyloxycarbonyl)amino-3-O-levulinoyl-2,4,6-trideoxy-D-galactopyranosyl] phosphate (2*):




To a stirred solution of galactal **1*** (3.17 g, 8.77 mmol) in dry MeCN (44 ml) were added at -25°C ceric ammonium nitrate (14.42 g, 26.3 mmol) and sodium azide (0.86 g, 13.15 mmol). The reaction was stirred vigorously between -25 °C and -20 °C for 6 h. The mixture was diluted with cold Et_2O (50 ml). The organic layer was washed with cold water (3x30 ml), dried over Na_2SO_4 and concentrated. The residue was filtered through a plug of silica gel (EtOAc/hexanes/ Et_3N 1:1:0.01) to give the crude glycosyl nitrate as 4:1 *galacto/talo* mixture (2.01 g) as a slightly yellow oil.
 25 To the crude glycosyl nitrate (2.01 g) was added at room temperature a solution of cesium dibutyl phosphate (2.21 g, 6.45 mmol) in dry DMF (28 ml). The mixture was stirred at that temperature for 4.5 h, diluted with EtOAc (100 ml) and poured into water (100 ml). The organic phase was washed with water (5x50 ml) and the combined aqueous fractions were extracted with EtOAc (50 ml). The organic phase
 30 was dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 45:55 to 50:50) to give glycosyl phosphate **2*** (1.84

g, 3.00 mmol, 37%, 1:10 α/β) as a clear oil. HRMS (ESI) calcd for $C_{27}H_{41}N_4O_{10}P$ ($M+Na^+$) 635.2458 found 635.2422 m/z .

Example 3: Ethyl 2-O-benzyl-3,4-isopropylidene-1-thio- β -D-galactopyranoside

5 (3*):




To a stirred solution of ethyl 6-O-*tert*-butyldimethylsilyl-3,4-isopropylidene-1-thio- β -D-galactopyranoside (*Bioorg. Med. Chem.* **2001**, 9, 1395) (45.7 g, 121 mmol) in DMF (150 ml) and THF (75 ml) were added at 0 °C portionwise sodium hydride (60%, 7.24

10 g, 181 mmol) and then benzyl bromide (17.2 ml, 145 mmol). The mixture was stirred for 1 h at 0 °C, slowly warmed to room temperature and stirred for 16 h at that temperature. The reaction was quenched at 0 °C with sat. aq. NH4Cl (20 ml), diluted with water (200 ml) and EtOAc (150 ml) and stirred for 15 min at 0°C. After separation, the organic phase was washed with water (5x100 ml) and the combined 15 aqueous fractions were re-extracted with EtOAc (2x100 ml). The combined organic extracts were dried over Na2SO4 and concentrated to give the crude benzyl ether (61 g) as a yellow oil.

To a stirred solution of the crude benzyl ether (61 g) in THF (370 ml) was added at 0 °C tetrabutylammonium fluoride (1 M in THF, 166 ml, 166 mmol). The mixture was

20 warmed to room temperature and stirred for 1 h. The reaction was diluted with sat. aq. NaHCO3 (200 ml) and EtOAc (100 ml). After separation, the aqueous phase was extracted with EtOAc (3x100 ml), the combined organic fractions were dried over MgSO4 and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 0:1 to 1:3 to 1:1) to give alcohol 3* as a white solid. HRMS (ESI) 25 calcd for $C_{18}H_{26}O_5S$ ($M+Na^+$) 377.1398 found 377.1416 m/z .

Example 4: Methyl (ethyl 2-O-benzyl-1-thio- β -D-galactopyranosid)uronate (4*):



To a vigorously stirred solution of alcohol 4* (6.0 g, 16.93 mmol) in CH2Cl2 (50 ml)

30 and H2O (25 ml) were added at 0 °C TEMPO (0.53 g, 3.39 mmol) and BAIB (10.9 g,

33.9 mmol). The mixture was warmed to room temperature and stirred for 1 h at that temperature. The reaction was quenched with 10% aq. $\text{Na}_2\text{S}_2\text{O}_3$ (10 ml) and diluted with EtOAc (30 ml). After separation, the organic phase was washed with 10% $\text{Na}_2\text{S}_2\text{O}_3$ (4x20 ml). The aqueous phase was extracted with EtOAc (2x20 ml) and the combined organic fractions were dried over Na_2SO_4 and concentrated to give the crude acid (7.92 g) as yellow oil.

To a stirred solution of acetyl chloride (6.04 ml, 85 mmol) in MeOH (300 ml) was added dropwise at 0 °C a solution of the crude acid (7.92 g) in MeOH (40 ml). The mixture was warmed to room temperature, stirred for 2 h at that temperature and cooled to 0 °C. The reaction was quenched with sat. aq. NaHCO_3 (30 ml) and neutralized to pH 7 with solid NaHCO_3 . The volatiles were evaporated and the mixture was diluted with EtOAc (70 ml). After separation, the aqueous phase was extracted with EtOAc (5x50 ml). The combined organic fractions were dried over Na_2SO_4 and concentrated. Flash chromatography was performed (EtOAc/hexanes 2:3 to 1:1, then 1:0) to give the crude product, which was crystallized in methanol at -20 °C (5 ml/g crude product) to give diol **4*** (3.47 g, 10.13 mmol, 60%) as a white solid. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{22}\text{O}_6\text{S} (\text{M}+\text{Na})^+$ 365.1034 found 365.1058 *m/z*.


Example 5: Methyl (ethyl 2-O-benzyl-3,4-O-*endo*-benzylidene-1-thio- β -D-galactopyranosid)uronate (**5***) and Methyl (ethyl 2-O-benzyl-3,4-O-*exo*-benzylidene-1-thio- β -D-galactopyranosid)uronate (**6***):



To a stirred solution of diol **4*** (2.99 g, 8.73 mmol) in dry acetonitrile (29 ml) were added at room temperature benzaldehyde dimethyl acetal (6.57 ml, 43.6 mmol) and DL-camphorsulfonic acid (0.51 g, 2.18 mmol). The mixture was stirred at room temperature for 5 h and the reaction was quenched by addition of triethylamine (0.35 ml). The mixture was concentrated under reduced pressure to give a residue, which was filtered through a short plug of silica gel (hexanes/EtOAc 8:1 (2% Et_3N) to 1:1 (2% Et_3N)) to give a 1:1 mixture of benzylidene acetals **5*** (*endo*) and **6*** (*exo*) (3.46 g, 8.03 mmol, 92%). The isomers were separated by selective crystallization of exo-isomer **6*** from EtOAc/hexanes and chromatographic separation of the mother liquor (Biotage, flat gradient of 10% to 40% EtOAc in hexanes + 0.5% Et_3N). Analytical data for **5***: Clear oil. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{26}\text{O}_6\text{S} (\text{M}+\text{Na})^+$ 453.1348, found 453.1352

m/z. Analytical data for **6***: White solid. HRMS (ESI) calcd for $C_{23}H_{26}O_6S$ ($M+Na$)⁺ 453.1348, found 453.1338 *m/z*.


Example 6: Methyl (ethyl 2,3-O-benzyl-1-thio- β -D-galactopyranosyl)uronate (7*):



5


To a solution of acetal **6*** (162 mg, 0.38 mmol) and sodium cyanoborohydride (296 mg, 4.70 mmol) in THF (9.4 ml) was added at room temperature a solution of hydrogen chloride (1 M in Et₂O) until the evolution of gas ceased. After 10 min, sodium cyanoborohydride (296 mg, 4.70 mmol) was added, followed by the addition 10 of HCl. The reaction was stirred at room temperature, diluted with EtOAc (30 ml) and quenched with sat. aq. NaHCO₃ (30 ml). After separation, the organic layer was washed with sat. aq. NaHCO₃ (20 ml) and the aqueous layer was re-extracted with EtOAc (2x20 ml). The organic extracts were pooled, dried over MgSO₄ and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 1:1) to give alcohol **7*** (67.5 mg, 0.156 mmol, 42%) as a white solid. HRMS (ESI) calcd for $C_{23}H_{28}O_6S$ ($M+Na$)⁺ 455.1504 found 455.1511 *m/z*.

Example 7: Methyl (ethyl 2,3-O-benzyl-4-O-fluorenylmethoxycarbonyl-1-thio- β -D-galactopyranosyl)uronate (8*):




To a stirring solution of alcohol **7*** (160 mg, 0.370 mmol) in pyridine (1.2 ml) was added at 0 °C FmocCl (383 mg, 1.48 mmol). The mixture was warmed to room temperature and stirred for 3 h. The mixture was then diluted with EtOAc (50 ml) and 25 washed with 1 N HCl (2x30 ml) and sat. aq. NaHCO₃ (30 ml). The organic phase was dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 1:2) to give carbonate **8*** (217 mg, 0.331 mmol, 90%) as a white foam. HRMS (ESI) calcd for $C_{38}H_{38}O_8S$ ($M+Na$)⁺ 677.2185 found 677.2167 *m/z*.

Example 8: Dibutyl [methyl (2,3-O-benzyl-4-O-fluorenylmethoxycarbonyl- α / β -D-galactopyranosyl)uronate] phosphate (9 *):



Thioglycoside 8 * (200 mg, 0.305 mmol) was co-evaporated with dry toluene (2x30 ml), kept under high vacuum for 1 h and dissolved in dry CH₂Cl₂ (3 ml). Activated molecular sieves (3 Å-AW) were added and the solution was stirred for 15 min at room temperature. The solution was then cooled to 0 °C, treated with dibutyl phosphoric acid (128 mg, 0.611 mmol) and stirred for another 15 min. The mixture was then treated with NIS (89 mg, 0.397 mmol), warmed to room temperature and stirred for 3 h at that temperature. The reaction was diluted with CH₂Cl₂ (20 ml) and quenched with a 1:1 (v/v) mixture of 10% aq. Na₂S₂O₃ and sat. aq. NaHCO₃ (20 ml). The aqueous phase was extracted with CH₂Cl₂ (3x30 ml), the combined organic fractions were dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 1:1 to 2:1) to give glycosyl phosphate 9 * (218 mg, 0.272 mmol, 89%, 10:1 α / β as clear oil. Analytical data of 9 $^*\alpha$: HRMS (ESI) calcd for C₄₄H₅₁O₁₂P (M+Na)⁺ 825.3015 found 825.3020 m/z. Analytical data of 9 $^*\beta$: HRMS (ESI) calcd for C₄₄H₅₁O₁₂P (M+Na)⁺ 825.3015 found 825.2970 m/z.


Example 9: Methyl (2-O-benzyl-3,4-O-endo-benzylidene- α / β -D-galactopyranosyl)uronate-(1→1)-2-(benzylthio)ethanol (10 *):



Thioglycoside 5 * (102 mg, 0.237 mmol), 2-(benzylthio)ethanol 11 * (60 mg, 0.355 mmol) and TTBPy. (117 mg, 0.474 mmol) were co-evaporated with anh. toluene (3x10 ml) and kept under high vacuum for 30 min. The mixture was dissolved in THF (4.8 ml) and stirred over activated molecular sieves (3 Å) for 30 min at room temperature. The solution was cooled to 0 °C and treated with DMTST (92 mg, 0.355 mmol in 0.2 ml dry CH₂Cl₂). The reaction was warmed to room temperature and stirred for 2 h at that temperature. The reaction was quenched with a 1:1 (v/v)


MeOH/Et₃N mixture (0.1 ml) and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes/Et₃N 0:1:0.01 to 30:70:0.01 to 45:55:0.01) to give thioether **10*α** (59 mg, 0.110 mmol, 46%) as a clear oil, along with the corresponding β-isomer **10*β** (35 mg, 0.065 mmol, 27%). Analytical data for **10*α**: HRMS (ESI) 5 calcd for C₃₀H₃₂O₇S (M+Na)⁺ 559.1766 found 559.1731 m/z.

Example 10: Methyl (2,4-di-O-benzyl-α-D-galactopyranosid)uronate-(1→1)-2-(benzylthio)ethanol (12*):



10 To a stirred solution of acetal **10*α** (100.0 mg, 0.186 mmol) in dry THF (5.3 ml) was added first borane trimethylamine complex (57.4 mg, 0.745 mmol) and then aluminium chloride (149 mg, 1.118 mmol) at room temperature. The mixture was stirred for 4.5 h. The reaction was quenched by addition of water (10 ml) and 1 M aq. HCl (5 ml). The mixture was extracted with EtOAc (3x 10 ml) and the combined 15 organic fractions were dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 2:5 to 1:1) to give alcohol **12*** (70.0 mg, 0.13 mmol, 70%) as a clear oil. HRMS (ESI) calcd for C₃₀H₃₄O₇S (M+Na)⁺ 561.1923 found 561.1879 m/z.


20 **Example 11: Methyl (2,3-di-O-benzyl-α-D-galactopyranosyl)uronate-(1→3)-methyl (2,4-di-O-benzyl-α-D-galactopyranosyl)uronate-(1→3)-(2-(benzylthio)ethanol (13*):**



Alcohol **12*** (90 mg, 0.166 mmol) and glycosyl phosphate **9*** (208 mg, 0.259 mmol) were co-evaporated with dry toluene (3x10 ml) and kept under high vacuum for 1 h. The mixture was dissolved in dry CH_2Cl_2 (3.3 ml) and stirred over activated molecular sieves (3 Å-AW) for 30 min at room temperature. The solution was cooled to 0 °C 5 and treated dropwise with TBSOTf (0.133 mmol in 0.2 ml dry CH_2Cl_2). The solution was warmed to room temperature and stirred for 20 h. The reaction was diluted with CH_2Cl_2 (10 ml) and quenched with a 1:1 (v/v) MeOH/pyridine mixture (0.2 ml). The solution was filtered through Celite and concentrated. The crude product was filtered 10 through a short plug of silica gel (EtOAc/hexanes 1:1) to give the intermediate disaccharide mixture (150 mg, 0.133 mmol, 80%, 3:1 α/β) as a clear oil.

To a stirred solution of the carbonate mixture (150 mg) in CH_2Cl_2 (2.6 ml) was added at room temperature triethylamine (1.1 ml, 7.96 mmol). The reaction was stirred for 3 h at that temperature and co-evaporated with toluene (2x10 ml). The residue was purified by flash chromatography (EtOAc/hexanes 1:6 to 2:3 to 1:1) to give alcohol 15 **13*** (62 mg, 0.068 mmol, 51%) along with the corresponding β -anomer (20 mg, 0.022 mmol, 17%). HRMS (ESI) calcd for $\text{C}_{51}\text{H}_{56}\text{O}_{13}\text{S}$ ($\text{M}+\text{Na}$)⁺ 931.3339, found 931.3340 *m/z*.


**Example 12: 2-Azido-4-(benzyloxycarbonyl)amino-3-O-levulinoyl-2,4,6-trideoxy-
 α -D-galactopyranosyl-(1→4)-methyl (2,3-di-O-benzyl- α -D-galactopyranosyl)uronate-(1→3)-methyl (2,4-di-O-benzyl- α -D-galactopyranosyl)uronate-(1→1)-2-(benzylthio)ethanol (14*):**



Alcohol **13*** (65 mg, 0.062 mmol) and glycosyl phosphate **2*** (61 mg, 0.100 mmol) 25 were co-evaporated with dry toluene (3x10 ml) and kept under high vacuum for 30 min. The mixture was dissolved in CH_2Cl_2 (2.1 ml) and stirred over activated molecular sieves (4 Å-AW) for 1 h at room temperature. The solution was then

cooled to 0 °C and treated with TMSOTf (17 μ l, 0.093 mmol in 0.2 ml dry CH_2Cl_2). The mixture was allowed to stir for 3 h at 0 °C, when TLC (EtOAc/hexanes 2:3) indicated complete consumption of the acceptor. The reaction was quenched with a 1:1 (v/v) MeOH/NEt₃ mixture (0.5 ml), diluted with CH_2Cl_2 (20 ml) and filtered through 5 Celite. The crude product was purified by flash chromatography (EtOAc/hexanes 1:2 to 1:1) to give trisaccharide **14*** (69 mg, 0.053 mmol, 85%) as a clear oil. HRMS (ESI) calcd. for $\text{C}_{70}\text{H}_{78}\text{N}_4\text{O}_{19}\text{S} (\text{M}+\text{Na})^+$ 1333.4879 found 1333.4911 *m/z*.

10 **Example 13:** 2-Azido-4-(benzyloxycarbonyl)amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-methyl (2,3-di-O-benzyl- α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-methyl (2,4-di-O-benzyl- α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(benzylthio)ethanol (**15***):




15

To a stirred solution of levulinoyl ester **14*** (30 mg, 0.023 mmol) in dry CH_2Cl_2 (1.0 ml) was added at room temperature first a mixture of pyridine (56 μ l, 0.692 mmol) and acetic acid (37 μ l, 0.646 mmol), and then hydrazine hydrate (2 μ l, 0.041 mmol). 20 The mixture was allowed to stir for 4 h at room temperature, diluted with EtOAc (2 ml), quenched with acetone (0.1 ml) and poured into water (15 ml). The aqueous phase was extracted with EtOAc (4x10 ml), and the combined organic extracts were dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc/hexanes 0:1 to 1:2 to 2:3) to give alcohol 25 **15*** (28 mg, 0.023 mmol, 100%) as a clear oil. HRMS (ESI) calcd. for $\text{C}_{65}\text{H}_{72}\text{N}_4\text{O}_{17}\text{S} (\text{M}+\text{Na})^+$ 1235.4511 found 1235.4539 *m/z*.

Example 14: 2-Azido-4-(benzyloxycarbonyl)amino-3-O-benzyloxymethyl-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-methyl (2,3-di-O-benzyl- α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-methyl (2,4-di-O-benzyl- α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(benzylthio)ethanol (16*):

5



Alcohol 15* (8.6 mg, 7.1 μ mol), benzyloxymethyl thiocyclohexane (79 mg, 0.354 mmol) and TTBPy. (105 mg, 0.425 mmol) were coevaproated with dry toluene (3x10 ml) and kept under high vacuum for 30 min. The mixture was dissolved in dry CH_2Cl_2 (0.4 ml) and stirred over activated molecular sieves (3 \AA) for 30 min at room temperature. The mixture was cooled to 0 $^{\circ}\text{C}$ and DMTST (7.1 mg, 0.18 mmol in 0.1 ml CH_2Cl_2) was added dropwise over 45 min, while the reaction temperature was 10

10


15

kept below 10 $^{\circ}\text{C}$. The reaction was stirred for another 45 min, quenched by addition of a 1:1 (v/v) MeOH/Et₃N mixture and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 1:10 to 1:2) to give acetal 16* (6.0 mg, 4.5 μ mol, 64%) as a clear oil. HRMS (ESI) calcd for $\text{C}_{73}\text{H}_{80}\text{N}_4\text{O}_{18}\text{S}$ ($\text{M}+\text{Na}$)⁺ 1355.5086 found 1355.5071 *m/z*.

20


Example 15: 2-Acetamido-4-(benzyloxycarbonyl)amino-3-O-benzyloxymethyl-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-methyl (2,3-di-O-benzyl- α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-methyl (2,4-di-O-benzyl- α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(benzylthio)ethanol (17*):

25



To a stirred solution of azide **16*** (14.0 mg, 10.5 μ mol) in dry pyridine (0.35 ml) was added at 0 °C thioacetic acid (0.35 ml). The mixture was warmed to room temperature and stirred for 24 h at that temperature. The solution was co-evaporated with toluene (2x5 ml) and the residue was purified by flash chromatography (EtOAc/hexanes 1:10 to acetone/hexanes 1:7 to 1:5 to 1:3) to give acetamide **17*** (9.4 mg, 7.0 μ mol, 66%) as a white solid. HRMS (ESI) calcd for $C_{75}H_{84}N_2O_{19}S$ ($M+Na$)⁺ 1371.5281 found 1371.5314 *m/z*.


10 **Example 16:** 2,2'-Dithiobis[2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1→4)- α -D-galactopyranosyluronate-(1→3)- α -D-galactopyranosyluronate-(1→1)-1-ethanol] (**18***):



To a stirred solution of diester **17*** in THF (4.0 ml) and MeOH (0.8 ml) was added at 0 °C a 1 M solution of NaOH in water (1.5 ml). The reaction was slowly warmed to room temperature and stirred for 16 h. The reaction was diluted with EtOAc (5 ml) and acidified to pH 4 with 0.5 M aq. NaHSO₄. After separation, the aqueous fraction 5 was extracted with EtOAc (8x10 ml), the combined organic fractions were dried over Na₂SO₄ and concentrated to give the intermediate diacid as a white solid.


To a stirring solution of liquid ammonia (5 ml) was added at -78 °C a solution of the crude diacid in THF (1.5 ml). The mixture was treated with *t*BuOH (0.5 ml) and lumps 10 of freshly cut sodium (45 mg) were added until a deeply blue color persisted. The reaction was stirred at -78 °C for 45 min and quenched by addition of solid ammonium acetate (200 mg). The solution was warmed to room temperature under a stream of argon and co-evaporated with MeOH (2x10 ml) and water (2x5 ml). The residue was left under air for 16 h, purified by size exclusion chromatography (Sephadex G-25, 1:1 MeOH/5 mM aq. NH₄OAc) and lyophilized repeatedly to give 15 disulfide **18*** (1.4 mg, 1.65 µmol, 32%) as a white solid. HRMS (MALDI) calcd for C₄₄H₇₀N₄O₃₂S₂ (M-H⁺) 1229.3330 found 1229.3342 *m/z*.

Example 17: Methyl (ethyl 2,3-O-benzyl-4-O-levulinoyl-1-thio- β -D-galactopyranosyl)uronate (19*):



To a stirred solution of alcohol **7*** (94 mg, 0.217 mmol) in CH₂Cl₂ (1.9 mL) were added at room temperature levulinic acid (386 mg, 3.26 mmol), DCC (673 mg, 3.26 mmol) and pyridine (0.26 mL, 3.26 mmol). The mixture was stirred at that 25 temperature for 35 h, diluted with CH₂Cl₂ (5 mL) and filtered through Celite. The mixture was concentrated, the residue was dissolved in a minimal volume of CH₂Cl₂ (1-3 mL) and filtered through cotton wool. The same procedure was repeated 3 times. The residue was purified by flash chromatography (EtOAc/toluene 1:1) to give ester **19*** (91 mg, 0.171 mmol, 79%) as a slightly yellow oil. HRMS (ESI) calcd for 30 C₂₈H₃₄O₈S (M+Na)⁺ 553.1872 found 553.1872 *m/z*.

Example 18: Methyl (2,3-di-O-benzyl- α -D-galactopyranosid)uronate-(1→1)-6-(benzylthio)hexanol (20*):




Thioglycoside **19*** (87 mg, 0.164 mmol), 6-(benzylthio)hexanol **21*** (85 mg, 0.379 mmol) and TTBPY. (97 mg, 0.392 mmol) were co-evaporated with anh. toluene (3x10 ml) and kept under high vacuum for 30 min. The mixture was dissolved in Et₂O (2.5

5 ml) and CH₂Cl₂ (0.83 ml) and stirred over activated molecular sieves (3 Å) for 30 min at room temperature. The solution was cooled to 0 °C and treated with DMTST (63.5 mg, 0.246 mmol). The reaction was warmed to room temperature and stirred for 8 h at that temperature. The reaction was quenched with a 1:1 (v/v) mixture of MeOH and triethylamine (0.1 ml) and concentrated. The residue was purified by flash

10 chromatography (EtOAc/CH₂Cl₂/hexanes 0:0:1 to 1:2:1) to give the corresponding glycosides (60 mg) as an inseparable α/β mixture.

To a stirred solution of the glycoside mixture in dry CH₂Cl₂ (2.2 ml) were added at room temperature first a mixture of pyridine (195 µl, 2.411 mmol) and acetic acid (137 µl, 2.393 mmol), and then hydrazine hydrate (5.9 µl, 0.121 mmol). The mixture 15 was stirred for 2 h at that temperature, diluted with EtOAc (2 ml), quenched with acetone (0.1 ml) and poured into water (15 ml). The aqueous phase was extracted with EtOAc (4x10 ml), the combined organic extracts were dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 1:2) to give alcohol **20*** (29 mg, 0.049 mmol, 30% over two steps) as a clear oil, 20 along with the corresponding β-isomer (22 mg, 0.037 mmol, 22%). HRMS (ESI) calcd. for C₃₄H₄₂O₇S (M+Na)⁺ 617.2544 found 617.2542 m/z.

Example 19: 6,6'-Dithiobis[α-D-galactopyranosyluronate-(1→1)-1-hexanol] (22*):



To a stirred solution of ester **20*** (10 mg, 0.017 mmol) in THF (1.0 ml) and MeOH (0.5 ml) was added at 0 °C a 1 M solution of NaOH in water (0.8 ml). The reaction was slowly warmed to room temperature and stirred for 16 h. The reaction was diluted with EtOAc (5 ml) and water (5 ml) and acidified to pH 4 with 0.5 M aq. NaHSO₄.


5 After separation, the aqueous fraction was extracted with EtOAc (8x5 ml), the combined organic fractions were dried over Na₂SO₄ and concentrated to give the intermediate acid as a white solid.

To a stirred solution of liquid ammonia (8 ml) was added at -78 °C a solution of the crude diacid in THF (2 ml). The mixture was treated with *t*BuOH (0.4 ml) and lumps of

10 freshly cut sodium (45 mg) were added until a deeply blue color persisted. The reaction was stirred at -78 °C for 45 min and quenched by addition of solid ammonium acetate (100 mg). The solution was warmed to room temperature under a stream of argon and co-evaporated with MeOH (2x10 ml) and water (2x5 ml). The residue was left under air for 16 h, purified by size exclusion chromatography

15 (Sephadex G-25, 9:1 MeOH/5 mM aq. NH₄OAc) and lyophilized repeatedly to give disulfide **22*** (3.1 mg, 5.1 µmol, 60% over two steps) as a white solid. HRMS (MALDI) calcd for C₂₄H₄₂O₁₄S₂(M-H⁺) 617.1938 found 617.1954 *m/z*.


20 **Example 20: 2-Acetamido-4-(benzyloxycarbonyl)amino-3-O-levulinoyl-2,4,6-trideoxy- α -D-galactopyranosyl-(1→1)-6-(benzylthio)hexanol (23*):**



6-(Benzylthio)hexanol **21*** (29 mg, 0.171 mmol) and glycosyl phosphate **2*** (70 mg, 0.114 mmol) were co-evaproated with dry toluene (3x10 ml) and kept under high 25 vacuum for 30 min. The mixture was dissolved in CH₂Cl₂ (1.8 ml) and stirred over activated molecular sieves (4 Å-AW) for 1 h at room temperature. The solution was then cooled to 0 °C and treated with TMSOTf (31 µl, 0.171 mmol in 0.2 ml dry CH₂Cl₂). The mixture was stirred for 3 h at that temperature, quenched with a 1:1 (v/v) mixture of MeOH and triethylamine (0.5 ml), diluted with CH₂Cl₂ (20 ml) and 30 filtered through Celite. The residue was purified by flash chromatography (EtOAc/hexanes 2:3 to 3:2) to give the corresponding glycosides (57 mg) as an inseparable α / β mixture.

To a stirred solution of the glycoside mixture in dry pyridine (0.9 ml) was added at 0 °C thioacetic acid (0.9 ml). The mixture was warmed to room temperature and stirred for 24 h at that temperature. The solution was co-evaporated with toluene (2x5 ml) and the residue was purified by flash chromatography (EtOAc/hexanes 1:2 to 2:1 to 5 6:1) to give acetamide **23*** (22 mg, 0.034 mmol, 29% over two steps) as a white solid, along with the corresponding β -isomer (21.6 mg, 0.034 mmol, 29%). HRMS (ESI) calcd for $C_{34}H_{46}N_2O_8S$ ($M+Na$)⁺ 665.2872 found 665.2865 *m/z*.


10 **Example 21: 6,6'-Dithiobis[2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 1)-1-hexanol] (24*):**



To a stirred solution of ester **23*** (10 mg, 0.016 mmol) in dry CH_2Cl_2 (1.0 ml) were added at room temperature first a mixture of pyridine (38 μ l, 0.467 mmol) and acetic acid (24.9 μ l, 0.436 mmol), and then hydrazine hydrate (1.0 μ l, 0.020 mmol). The 15 mixture was stirred for 2 h at that temperature, quenched with acetone (0.1 ml) and purified by size exclusion chromatography (Sephadex LH-20, $CH_2Cl_2/MeOH$ 2:1) to give the corresponding alcohol as a clear oil.

To a stirred solution of liquid ammonia (5 ml) was added at -78 °C a solution of the 20 intermediate alcohol in THF (1.2 ml). The mixture was treated with *t*BuOH (0.5 ml) and lumps of freshly cut sodium (80 mg) were added until a deeply blue color persisted. The reaction was stirred at -78 °C for 45 min and quenched by addition of 25 solid ammonium acetate (100 mg). The solution was warmed to room temperature under a stream of argon and co-evaporated with MeOH (2x10 ml) and water (2x5 ml). The residue was left under air for 16 h, purified by size exclusion chromatography (Sephadex G-25, 9:1 MeOH/5 mM aq. NH_4OAc) and lyophilized repeatedly to give disulfide **24*** (1.7 mg, 2.7 μ mol, 33% over two steps) as a white solid. HRMS (ESI) calcd. for $C_{28}H_{54}N_4O_8S_2$ ($M+Na$)⁺ 661.3281 found 661.3306 *m/z*.


30 **Example 22: 2-Acetamido-4-(benzyloxycarbonyl)amino-3-O-levulinoyl-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 1)-2-(benzylthio)ethanol (25*):**



2-(benzylthio)ethanol **11*** (71 mg, 0.421 mmol) and glycosyl phosphate **2*** (171 mg, 0.281 mmol) were co-evaproated with dry toluene (3x10 ml) and kept under high vacuum for 30 min. The mixture was dissolved in CH₂Cl₂ (1.8 ml) and stirred over

5 activated molecular sieves (4 Å-AW) for 1 h at room temperature. The solution was then cooled to -40 °C and treated with TMSOTf (56 µl, 0.309 mmol in 0.2 ml dry CH₂Cl₂). The mixture was slowly warmed to 0 °C (2 h), quenched with a 1:1 (v/v) mixture of MeOH and triethylamine (0.5 ml), diluted with CH₂Cl₂ (20 ml), filtered through Celite and concentrated. The residue was purified by flash chromatography
10 (EtOAc/hexanes 1:3 to 1:1) to give the corresponding α-glycoside (55 mg, 0.096 mmol, 34%) along with the corresponding β-glycoside (22 mg, 0.039 mmol, 14%).
To a stirred solution of the intermediate α-glycoside (40 mg, 0.070 mmol) in dry pyridine (0.4 ml) was added at 0 °C thioacetic acid (0.4 ml). The mixture was warmed to room temperature and stirred for 24 h at that temperature. The solution was co-
15 evaporated with toluene (2x5 ml) and the residue was purified by flash chromatography (EtOAc/hexanes 1:3 to acetone/hexanes 1:2 to 2:3) to give acetamide **25*** (31 mg, 0.053 mmol, 76%) as a white solid. HRMS (ESI) calcd for C₃₀H₃₈N₂O₈S (M+Na)⁺ 609.2246 found 609.2256 m/z.

20 **Example 23: 6,6'-Dithiobis[2-acetamido-4-amino-2,4,6-trideoxy-α-D-galactopyranosyl-(1→1)-1-ethanol] (26):**




To a stirred solution of ester **25*** (20.7 mg, 0.035 mmol) in dry CH₂Cl₂ (3.0 ml) were added at room temperature first a mixture of pyridine (86 µl, 1.058 mmol) and acetic acid (57 µl, 0.988 mmol), and then hydrazine hydrate (3.4 µl, 0.071 mmol). The mixture was stirred for 5 h at that temperature, diluted with EtOAc (2 ml), quenched

with acetone (0.1 ml) and poured into water (10 ml). The aqueous phase was extracted with EtOAc (4x5 ml), the combined organic fractions were dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (acetone/hexanes 1:1) to give the intermediate alcohol (17.5 mg) as a white solid.

- 5 To a stirred solution of liquid ammonia (6 ml) was added at -78 °C a solution of the intermediate alcohol in THF (1.5 ml). The mixture was treated with *t*BuOH (0.5 ml) and lumps of freshly cut sodium (45 mg) were added until a deeply blue color persisted. The reaction was stirred at -78 °C for 45 min and quenched by addition of solid ammonium acetate (100 mg). The solution was warmed to room temperature
- 10 under a stream of argon and co-evaporated with MeOH (2x10 ml) and water (2x5 ml). The residue was left under air for 16 h, purified by size exclusion chromatography (Sephadex G-25, 1:10 MeOH/5 mM aq. NH₄OAc) and lyophilized repeatedly to give disulfide **26*** as the diacetate salt (7.91 mg, 12.3 µmol, 70% over
- 15 two steps) as a white solid. HRMS (ESI) calcd. for C₂₀H₃₈N₄O₈S₂ (M+Na)⁺ 549.2029 found 549.2086 *m/z*.

Example 24: 2,2'-Dithiobis[α -D-galactopyranosyluronate-(1 \rightarrow 3)- α -D-galactopyranosyluronate-(1 \rightarrow 1)-1-ethanol] (27*):



20

- To a stirred solution of ester **13*** (8.6 mg, 9.5 µmol) in THF (0.6 ml) and MeOH (0.3 ml) was added at 0 °C a 1 M solution of NaOH in water (0.5 ml). The reaction was slowly warmed to room temperature and stirred for 16 h. The reaction was diluted with EtOAc (5 ml) and water (5 ml) and acidified to pH 4 with 0.5 M aq. NaHSO₄. After separation, the aqueous fraction was extracted with EtOAc (8x5 ml), the combined organic fractions were dried over Na₂SO₄ and concentrated to give the intermediate diacid as a white solid.

To a stirred solution of liquid ammonia (6 ml) was added at -78 °C a solution of the crude diacid in THF (1.5 ml). The mixture was treated with *t*BuOH (0.4 ml) and lumps of freshly cut sodium (75 mg) were added until a deeply blue color persisted. The reaction was stirred at -78 °C for 45 min and quenched by addition of solid 5 ammonium acetate (100 mg). The solution was warmed to room temperature under a stream of argon and co-evaporated with MeOH (2x10 ml) and water (2x5 ml). The residue was left under air for 16 h, purified by size exclusion chromatography (Sephadex G-25, 1:9 MeOH/5 mM aq. NH₄OAc) and lyophilized repeatedly to give disulfide **27*** (2.5 mg, 2.9 µmol, 61% over two steps) as a white solid. HRMS (MALDI) 10 calcd for C₂₈H₄₂O₂₆S₂ (M-H⁺) 901.0966 found 901.0981 *m/z*.

Example 25: Synthesis of glycoconjugates- Conjugation of saccharides of general formula (I) to CRM₁₉₇: To a stirred solution of CRM₁₉₇ (1 mg, 17.2 nmol) in 0.1 M sodium phosphate buffer (NaPi) pH 7.4 (1 ml) was added at room temperature 15 a solution of succinimidyl-3-(bromoacetamido)propionate (SBAP) (264 µg, 863 nmol) in DMF (20 µl). The mixture was stirred for 1 h at that temperature, and concentrated using membrane filtration (Amicon Ultra centrifuge membranes, 10 kDa cut-off). The protein solution was diluted with 0.1 M NaPi pH 7.4 and concentrated again. This process was repeated three times and the solution was diluted to 1 ml using 0.1 M 20 NaPi pH 7.4. The intermediate of general formula (II) (690 mmol) in 120 µl 0.1 M NaPi pH 7.4 was treated at room temperature with tris(2-carboxyethyl)phosphine (TCEP) (690 mmol), left for 1 h at that temperature under an argon atmosphere and added to the solution of the bromoacetamido-modified CRM₁₉₇ protein at room 25 temperature. The mixture was left at room temperature for 2 h and then at 4 °C for 16 h, and purified using membrane filtration (see above). The purified glycoconjugate in 0.1 M NaPi pH 7.4 (1 ml) was then treated at room temperature with L-cysteine (417 µg, 3.45 µmol) in 100 µl water. The mixture was left for 2 h at that temperature and purified by membrane filtration. Incorporation of the saccharide of general formula (I) 30 into the glycoconjugate was assessed by MALDI-TOF-MS, SDS-PAGE and size exclusion chromatography with right angle light scattering detection (SEC-RALS).

Example 26: Synthesis of glycoconjugates- In flow conjugation of the saccharides of general formula (I) to a glycosphingolipid with immunomodulatory properties

35 By using a photochemical flow reactor (*Chem. Eur. J.* **2013**, *19*, 3090) that was fitted with a loop of Teflon AF2400 tubing (566 µL), a solution of saccharide of general formula (I) (1.5 equiv.) in water (300 µL) was reacted with pentenyl modified (2S,3S,4R)-1-(α -D-galactopyranosyl)-2-hexacosanoylaminooctadecane-3,4-diol (1 equiv.) in water (300 µL) and AcOH (8 µL; residence time: 10 min, flow rate: 28.3

5 $\mu\text{L}/\text{min}^{-1}$ per syringe). The reactor output was lyophilized and the crude material was purified using size exclusion chromatography (Sephadex-G25, 5% EtOH in water, 10 mmx150 mm) to yield the glycoconjugate of saccharide of general formula (I) covalently linked to the modified (2S,3S,4R)-1-(α -D-galactopyranosyl)-2-hexacosanoylaminoctadecane-3,4-diol as white solid.

10 **Example 27: Synthesis of glycoconjugates- Conjugation of 2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(thio)ethanol to BSA**

15 To a stirred solution of BSA (0.5 mg, 7.6 nmol) in 0.1 M sodium phosphate buffer (NaPi) pH 7.4 (1 mL) was added at room temperature a solution of N-succinimidyl-3-(bromoacetamido)propionate (SBAP) (89 μg , 290 nmol) in DMF (20 μL). The mixture was stirred for 1 h at room temperature, and concentrated using membrane filtration (Amicon 0.5 mL Ultra centrifuge membranes, 10 kDa cut-off). The protein solution was diluted with 0.1 M NaPi pH 7.4 and concentrated again. This process was repeated three times and the solution was diluted to 0.5 mL using water. 20 μL were taken for analysis, and the protein solution was re-buffered to 0.1 M NaPi pH 7.4 using membrane filtration. Disulfide 18* (140 μg , 228 nmol resp. to the monomer) in 20 120 μL 0.1 M NaPi pH 7.4 was treated at room temperature with tris(2-carboxyethyl)phosphine (TCEP) (250 nmol), left for 1 h at that temperature under an argon atmosphere and added to the solution of the activated protein at room temperature. The mixture was left at 4 °C for 16 h, and purified using membrane filtration (see above). After washing with water and diluting to 0.5 mL, another 25 analytical sample (20 μL) was taken, and the solution was re-buffered. The purified glycoconjugate in 0.1 M NaPi pH 7.4 (0.5 mL) was then treated at room temperature with L-cysteine (417 μg , 3.45 μmol) in 100 μl water. The mixture was left for 2 h at that temperature and purified by membrane filtration. Incorporation of 2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-(α -D-galactopyranosyl)uronate-30 (1 \rightarrow 3)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(thio)ethanol into the glycoconjugate was assessed by MALDI-TOF-MS (positive mode):

30 Molecular weight measured:

BSA: 66341 m/z.

BSA-SBAP conjugate: 68316 m/z (incorporation of approximately 10 SBAP groups).

35 BSA-SBAP glycoconjugate: 69101 m/z (incorporation of approximately 1.3 molecules of 2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(thio)ethanol).

BSA-SBAP glycoconjugate after quenching with L-cysteine: 72074 m/z (incorporation of approximatively 24.5 L-cysteine molecules).

Example 28: Synthesis of glycoconjugates- Conjugation of 2-acetamido-4-

5 amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-(α -D-
galactopyranosyl)uronate-(1 \rightarrow 3)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-
(thio)ethanol to BSA

To a stirred solution of BSA (0.5 mg, 7.6 nmol) in 0.1 M sodium phosphate buffer (NaPi) pH 7.4 (1 mL), a solution of *N*-Succinimidyl-3-maleimidopropionate (101 μ g, 380 nmol) in DMF (20 μ L) was added at room temperature. The mixture was stirred

10 for 1 h at room temperature, and concentrated using membrane filtration (Amicon 0.5 mL Ultra centrifuge membranes, 10 kDa cut-off). The protein solution was diluted with 0.1 M NaPi pH 7.4 and concentrated again. This process was repeated three times and the solution was diluted to 0.5 mL using water. 20 μ L were taken for

15 analysis, and the protein solution was re-buffered to 0.1 M NaPi pH 7.4 using membrane filtration. Disulfide **18*** (140 μ g, 228 nmol resp. to the monomer) in 120 μ L 0.1 M NaPi pH 7.4 was treated at room temperature with tris(2-carboxyethyl)phosphine (TCEP) (250 nmol), left for 1 h at that temperature under an argon atmosphere and added to the solution of the activated protein at room

20 temperature. The mixture was left at 4 °C for 16 h, and purified using membrane filtration (see above). After washing with water and diluting to 0.5 mL, another analytical sample (20 μ L) was taken, and the solution was re-buffered. The purified glycoconjugate in 0.1 M NaPi pH 7.4 (0.5 mL) was then treated at room temperature with L-cysteine (417 μ g, 3.45 μ mol) in 100 μ L water. The mixture was left for 2 h at

25 that temperature and purified by membrane filtration. Incorporation of glycan into the glycoconjugate was assessed by MALDI-TOF-MS (positive mode):

Molecular weight measured:

BSA: 66341 m/z;

30 BSA-maleimide conjugate: 69254 m/z (incorporation of approximately 19 maleimide groups);

BSA-maleimide glycoconjugate: 71340 m/z (incorporation of approximately 3.4 molecules of 2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(thio)ethanol);

35 BSA-maleimide glycoconjugate after quenching with L-cysteine: 72106 m/z (incorporation of approx. 6.3 L-cysteine molecules).

Example 29: Conjugation to a solid support- Synthesis of microarrays using GAPSII slides

- Maleimide-functionalized microarrays were produced by submerging amine-coated slides (GAPS II slides, Corning) in 6-maleimidohexanoic acid *N*-hydroxysuccinimide ester (2 mM) in dry DMF with diisopropylethylamine (2.5% v/v) for 24 h at room temperature. Slides were washed three times with water and three times with
- 5 ethanol, centrifuged to dryness, and stored under argon until spotting. Diluted saccharides of general formula (I) were printed onto the modified microarray slides at 0.4 nL per spot by an automatic piezoelectric arraying robot (Scienion, Berlin, Germany). For completion of the immobilization reaction, printed slides were stored for 24 h in a humidified chamber.
- 10 Microarray slides were washed three times with water. Unreacted maleimide was quenched by submerging the slides in β -mercaptoethanol (0.1%, v/v) in PBS for 1 h at room temperature. Slides were washed three times with water and with ethanol, centrifuged to dryness, and blocked with BSA (1%, w/v) in PBS for 1 h at room temperature. Blocked slides were washed (2x PBS, 3 x water), centrifuged, and
- 15 incubated with the sera dilutions.

Example 30: Conjugation to a solid support- Synthesis of microarrays using CodeLink NHS slides

- CodeLink NHS slides were incubated for 24 h (1% w/v in PBS) at 4 °C. Slides were
- 20 incubated in blocking buffer (100 mM ethanolamine in 50 mM NaPi pH > 9) for 30 min at room temperature, washed three times each with water and ethanol, and dried. Slides were then subjected to maleimide functionalization and printing (see **Example 29**).

25 **Example 31: Binding experiments using the microarrays synthesized according to the procedure described at examples 29 and 30.**

- Binding experiments were performed by incubating microarray slides coated with the saccharides of general formula (I) with either a rabbit anti-SP1 typing serum or human pneumococcal reference serum 007sp (pooled sera of 287 humans
- 30 immunized with Pneumovax vaccine) in the dilutions indicated in the presence or absence of native SP1 polysaccharide, and using fluorescently labeled anti-rabbit or anti-human secondary antibodies.

35 **Example 32: Assessment of the immunogenicity of the linker A and of the interconnecting molecule**

To check the immunogenicity of the linker A and of the interconnecting molecule used in preparing the glycoconjugate 1 according to the present invention i.e. a glycoconjugate containing

a saccharide of general formula I presenting a linker A connected via an interconnecting molecule 1 to immunogenic carrier 1 such as:

5 H-(P)_{n3}-(N)_{n2}-(M)_{n3}-O-A-S-rest interconnecting molecule 1-immunogenic carrier 1,

three glycoconjugates need to be synthesized:

glycoconjugate 2:

10 H-(P)_{n3}-(N)_{n2}-(M)_{n3}-O-A-S-rest interconnecting molecule 2-immunogenic carrier 2;

glycoconjugate 3:

galactose-O-A-S-rest interconnecting molecule 1-immunogenic carrier 2;

15 **glycoconjugate 4:**

galactose-O-A-S-rest interconnecting molecule 2-immunogenic carrier 2;

The **immunogenic carrier 2** has to be non-related to immunogenic carrier 1 used in
20 immunization. For example, if CRM197 was used to prepare glycoconjugate 1, then BSA can be used as immunogenic carrier 2.

The galactose was chosen for preparing glycoconjugate 3, as it is non related to the saccharides according to the present invention i.e. to H-(P)_{n3}-(N)_{n2}-(M)_{n3}-OH.

25

The interconnecting molecule 2 used for the preparation of glycoconjugate 2 has to be non related to interconnecting molecule 1. For example, if Sulfo-GMBS was used as interconnecting molecule 1, then a non-related interconnecting molecule 2 would be Sulfo SIAB.

30

The choice of immunogenic carrier 2, interconnecting molecule 2 and of non related saccharide are obvious for the person skilled in the art of synthesis of glycoconjugates.

35 Protocol for ELISA:


A 96-well-plate is coated with 50 µl of the respective glycoconjugates (50 µg/ml) in PBS for 1 h at 37 °C. The plate is washed once with 100 µl washing buffer (PBS+0.1% (v/v) Tween-20) and blocked using 200 µl blocking solution (1% (w/v) BSA in PBS) for 1 h at 37 °C. The plate is washed 3x with washing buffer and then

incubated with dilutions of antiserum in blocking solution (50 μ l) for 16 h at 4 °C. The plate is washed 3x and incubated with 50 μ l of an appropriate secondary antibody (e.g. Goat Anti-Mouse IgG H&L (HRP), abcam ab6789) diluted in blocking solution for 1 h at 37 °C. The plate is washed 3x with washing buffer and incubated with 5 ELISA substrate (e.g. ABTS from Pierce, No. 37615) according to the manufacturer's manual.

Comparison of the optical density between the glycoconjugates applied will give a quantitative comparison between anti-linker-, anti-interconnecting molecule, and anti-10 saccharide immune responses.


Example 33: 2-Azido-4-(benzyloxycarbonyl)amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 1)-6-(benzylthio)ethanol (28*)

15



To a stirred solution of the intermediate Lev ester **25*** (17 mg, 0.03 mmol) in dry CH₂Cl₂ (1 mL) were added at room temperature first a mixture of pyridine (72 μ l, 0.894 mmol) and acetic acid (48 μ l, 0.834 mmol), and then hydrazine hydrate (3 μ l, 0.062 mmol). The mixture was stirred for 5 h at that temperature, diluted with EtOAc (2 ml), quenched with acetone (0.1 mL) and poured into water (10 mL). The aqueous phase was extracted with CH₂Cl₂ (4x5 ml), the combined organic fractions were dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 3:1) to give alcohol **28*** (13 mg, 0.028 mmol, 92%) as a clear oil. 25 HRMS (ESI) calcd. for C₂₃H₂₈N₄O₅S (M+Na)⁺ 495.1678 found 495.1679 m/z.

Example 34: Methyl (2,3-di-O-benzyl- α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-2-azido-4-(benzyloxycarbonyl)amino-2,4,6-trideoxy- α -D-galactopyranosyl -(1 \rightarrow 1)-2-(benzylthio)ethanol (29*)




Alcohol **28*** (13 mg, 0.028 mmol), TTBPy (45, 0.138 mmol) and thioglycoside **19*** (37 mg, 0.069 mmol) were co-evaporated with dry toluene (3x10 mL) and kept under high vacuum for 1 h. The mixture was dissolved in dry THF (1.5 mL) and stirred over

- 5 activated molecular sieves (3 Å) for 30 min at room temperature. The solution was cooled to 0 °C and treated dropwise with DMTST (17 mg, 0.069 mmol in 0.2 mL DCM) The mixture was warmed to room temperature and treated with an additional DMTST solution in DCM (2 equiv.) after 2 h. The reaction was stirred for 16 h and quenched with 1:1 (v/v) mixture of 10% aq. Na₂S₂O₃ and sat. aq. NaHCO₃ (5 mL).
 10 The mixture was extracted with CH₂Cl₂ (3x10 mL), dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (EtOAc/hexanes 1:2) to give the intermediate disaccharide as a clear oil.

15 To a stirred solution of the intermediate disaccharide in dry pyridine (0.2 ml) was added at 0 °C thioacetic acid (0.2 ml). The mixture was warmed to room temperature and stirred for 24 h at that temperature. The solution was co-evaporated with toluene (2x5 ml) and the residue was purified by flash chromatography (EtOAc/hexanes 1:3 to acetone/hexanes 1:2) to give the intermediate acetamide as a white foam.

- 20 To a stirred solution of the intermediate acetamide in dry CH₂Cl₂ (0.6 mL) and MeOH (60 µL) were added at room temperature first a mixture of pyridine (12 µL, 0.16 mmol) and acetic acid (8 µL, 0.15 mmol), and then hydrazine hydrate (1 µL, 0.021 mmol). The mixture was stirred for 3 h at that temperature, diluted with CH₂Cl₂ (2 mL), quenched with acetone (0.1 mL) and poured into water (5 mL). The aqueous phase was extracted with CH₂Cl₂ (4x5 mL), the combined organic fractions were dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (acetone/hexanes 0:1 to 1:1) to give acetamide **29*** (2.7 mg, 3.14 µmol, 21% over 3 steps based on recovered **28***) as a white foam. HRMS (ESI) calcd. for C₄₆H₅₄N₂O₁₂S (M+Na)⁺ 881.3295 found 881.3286 *m/z*.

Example 35: 2,2'-Dithiobis[α -D-galactopyranosyluronate-(1 \rightarrow 3)-2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl -(1 \rightarrow 1)-1-ethanol] (30*)




- 5 To a stirred solution of ester **29*** (2.7 mg, 3.14 μ mol) in THF (1 mL) and MeOH (0.25 mL) was added at 0 °C a 1 M solution of NaOH in water (0.4 mL). The reaction was slowly warmed to room temperature and stirred for 16 h. The reaction was diluted with EtOAc (5 ml) and water (5 ml) and acidified to pH 4 with 0.5 M aq. NaHSO₄. After separation, the aqueous fraction was extracted with EtOAc (8x5 ml), the combined organic fractions were dried over Na₂SO₄ and concentrated to give the intermediate diacid as a white solid.
- 10

- 15 To a stirred solution of liquid ammonia (10 ml) was added at -78 °C a solution of the crude diacid in THF (1.5 ml). The mixture was treated with *t*BuOH (0.4 ml) and lumps of freshly cut sodium (80 mg) were added until a deeply blue color persisted. The reaction was stirred at -78 °C for 45 min and quenched by addition of solid ammonium acetate (100 mg). The solution was warmed to room temperature under a stream of argon and co-evaporated with MeOH (2x10 ml) and water (2x5 ml). The residue was left under air for 16 h, purified by size exclusion chromatography (Sephadex G-25, 1:3 MeOH/5 mM aq. NH₄OAc) and lyophilized repeatedly to give disulfide **30*** (1.15 mg, 2.61 μ mol, 83% over two steps) as a white solid:
- 20

- 25 ¹H NMR (600 MHz, D₂O) δ 5.16 (d, *J* = 2.0 Hz, 1H), 5.02 (d, *J* = 3.2 Hz, 1H), 4.49 (d, *J* = 5.9 Hz, 1H), 4.44 – 4.35 (m, 1H), 4.34 – 4.20 (m, 2H), 4.15 – 4.04 (m, 1H), 4.00 – 3.78 (m, 5H), 3.06 (t, *J* = 5.6 Hz, 2H), 2.09 (s, 3H), 1.41 (d, *J* = 6.6 Hz, 3H). LRMS calcd. for C₃₂H₅₄N₄O₂₀S₂ (M+2H)²⁺ 440.4 found 440.2 *m/z*.

Example 36.1: General procedures for accessing the compound of general formula 3 according to the present invention:



wherein: R³ represents Me and R⁴ represents OH


To a mixture of aldehyde **A17** (1.0 mmol) and ketone **A16** (1.0 mmol) in i-PROH (100 μ L) were added propionic acid (0.1 mmol, 10 mol %) and pyrrolidine (0.1 mmol, 10 mol %). The reaction mixture was stirred at 45 °C for 1-25 h. NaHCO₃ was added, and the mixture was extracted with CH₂Cl₂ (3.5 mL). The combined extracts were washed with brine, dried (Na₂SO₄), and concentrated in vacuo. The residue was purified by flash chromatography to give intermediate unsaturated ketone **A15**.

A 9.5×10^{-3} M solution of intermediate unsaturated ketone **A15** (0.05 mmol) in THF and recently prepared Stryker's reagent (0.025 mmol), were mixed together forming a homogeneous solution that was stirred at room temperature for 2 h. The reaction was quenched with saturated aq. NH₄Cl. The mixture was stirred for 1 h. The reaction mixture was filtered, and the residue was washed with ethyl acetate. The organic phase was separated, and the aqueous phase was extracted with ethyl acetate. The combined organic phases were dried with MgSO₄ and the solvent was removed under vacuum. The residue was purified by column chromatography on silica gel to give ketone **A14**.


15

Neat tris-nonyloxy methyl titanium (10.1 mmol) was placed in a two-necked round bottom flask and subjected to Ar atmosphere. Ketone **A14** (4.65 mmol) in THF was added and the mixture was stirred at room temperature for 30 min. Oleic acid (17.8 mmol) was added and the mixture was heated to 110°C. The product was concentrated and purified by flash chromatography to give the intermediate tertiary alcohol.

20 To a solution of the PMB ether (0.1 mmol) in dichloroethane (5 mL), POCl₃ (0.5 mmol) was added and stirred at room temperature. After completion of the reaction, it was quenched in ice water and the organic layer was separated and the aqueous layer was extracted with dichloroethane (2 \times 5 mL). Combined organic layer was washed with brine solution, dried (Na₂SO₄), concentrated under reduced pressure and the residue was purified by column chromatography (silica gel, EtOAc : Hexanes) to afford alcohol **A13**.

Example 36.3


- 5 An oven-dried, 250-mL, round-bottomed flask was charged with 1,3-propanedithiol (20 mmol, 1.1 equiv) and tetrabutylammonium iodide (0.40 mmol, 2.2 mol %) in dry THF (100 mL). The mixture was stirred at room temperature and sodium hydride (60% suspension in mineral oil, 20 mmol, 1.1 equiv) was added by portions. The resulting mixture was stirred for 30 min, then benzyl bromide (18 mmol) was added dropwise. The solution was stirred for 1 h at room temperature, then filtered on a frit funnel and concentrated under vacuum. The resulting crude oil was distilled under vacuum to afford the title compound as colorless oil **A20**.
- 10 To a solution of **A19** (0.11 mmol) in anhydrous DMF (2 mL) was added thiol **A20** (0.22 mmol). The mixture was stirred at 25 °C for 18 h and then concentrated in vacuum. The residue was purified by flash chromatography to afford **A18**.
- A variety of dithiol derivatives of general formula **A20** are commercially available.

Example 36.4

- 5 Freshly cut sodium metal (4.67 mmol) was dissolved in isopropanol (10 mL) and benzyl mercaptan (6.23 mmol) was added. A solution of 4-(bromomethyl)cyclopent-1-ene (1.55 mmol) in isopropanol (5 mL) was added and the solution was heated under reflux for 4 days. The solution was allowed to cool to room temperature, diluted with water (50 mL) and extracted with diethyl ether (3X50 mL). The combined organic extracts were washed with 0.1 M aq potassium hydroxide (2X50 mL), dried and evaporated to yield crude **A25**. Column chromatography on silica gel eluting with ethyl acetate/hexane afforded the title compound **A25**.
- 10 A dry 50-mL flask equipped with a magnetic stirring bar, a septum inlet, an oil bubbler, and a reflux condenser was flushed with nitrogen. To the flask were added an alkene **A25** (5.5 mmol) and dry THF (2.5 mL) and then a solution of 9-BBN (0.5 M solution in THF, 5.5 mmol) at 0 °C. The mixture was warmed up slowly to room temperature and then stirred for 4-6 h to give a solution of B-alkyl-9-BBN **A24**.


To the above borane solution of **A24** were added DMF-THF (25 mL), $\text{PdCl}_2(\text{dppf})$ (0.15 mmol), haloalkene (5 mmol), and powdered K_3PO_4 (6 mmol). The mixture was stirred for 8 h at 50 °C and then poured into water. The product was extracted with benzene, washed with water four times, and dried over MgSO_4 . Column chromatography on silica gel eluting with ethyl acetate/hexane afforded the title compound **A22**.

Example 36.5



To a solution of 1.1 equiv aldehyde **A30** in dry methanol was dropwise added 1.3 equiv of amine **A28** in dry methanol at 0 °C under an argon atmosphere. The mixture was stirred for an additional 10 min at room temperature and followed by the sequential addition of 1.0 equiv acid **A27** in dry methanol and 1.1 equiv isocyanide **A29**. The reaction mixture was stirred for 24-48 h at room temperature. The resulting solution was diluted with dichloromethane and washed with 1 N HCl aqueous solution followed by saturated sodium bicarbonate aqueous solution and brine. The organic layer was dried over MgSO_4 , concentrated and purified by silica gel column chromatography (hexanes/ethyl acetate).

Example 36.6



Synthesis of A33:

- 5 To the solution of azide **A35** (0.03 mol) and ammonium chloride (0.07 mol) in ethyl alcohol (80 mL) and water (27 mL), zinc powder (0.04 mol) was added, the mixture was stirred vigorously at room temperature or at refluxing. After the reaction is over, ethyl acetate (200 mL) and aqueous ammonia (10 mL) was added. The mixture was filtered, and the filtrate was washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure.


10 Synthesis of **A31**:
 Acid **A32** (1 mmol) and amine **A33** (1 mmol) were coupled with 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) hydrochloride (1 mmol) in 3.5 mL methanol for 3 h at room temperature. The mixture was diluted with EtOAc (20 mL) and extracted with water (10

mL), 1 M aq. HCl (10 mL) and sat. aq. NaHCO₃ (10 mL). The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography to give the intermediate alcohol.

A suspension of NaH (0.83 mmol) in THF (2 mL) was cooled to 0 °C, and then to the suspension was added the intermediate alcohol (0.17 mmol) in THF (4 mL) slowly. The mixture was refluxed for 2 h and then allowed to cool to room temperature. To the mixture was added **A34** (0.11 mmol) in THF (2 mL) at that temperature dropwise. The reaction mixture was refluxed for 12 h and then allowed to cool to room temperature. After removal of the solvent by a rotary evaporator, the residue was dissolved in CHCl₃ and filtered. The filtrate was evaporated by a rotary evaporator and chromatographed.

To a solution of the PMB ether (0.1 mmol) in dichloroethane (5 mL), POCl₃ (0.5 mmol) was added and stirred at room temperature. After completion of the reaction, it was quenched in ice water and the organic layer was separated and the aqueous layer was extracted with dichloroethane (2 × 5 mL). Combined organic layer was washed with brine solution, dried (Na₂SO₄), concentrated under reduced pressure and the residue was purified by column chromatography (silica gel, EtOAc : Hexane) to afford the corresponding alcohol **A31**.

Example 36.7




15

In the test tube were introduced epoxide **A42** (5 mmol) and water (2 mL). Amine **A41** (6 mmol) was added in one portion and the test tube was kept at 0 °C and warmed to room temperature under vigorous stirring for 24 h. Water (2 mL) was added and the aqueous mixture was extracted with 10 mL of ethyl acetate and dried over anhydrous Na₂SO₄, and solvent was removed under reduced pressure to give the intermediate β-amino alcohol. The crude amine in EtOAc (2 mL) and sat. aq. NaHCO₃ (1 mL) was treated with CbzCl (6 mmol) at room temperature and stirred for 5 h at that temperature. The mixture was extracted with EtOAc (3 × 5 mL), the organic fraction was dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography to give alcohol **A40**.


10

20

Example 36.8

Bromide **A47** (8.26 mmol) was reacted with thiol **A48** in MeOH (5 mL) adjusted to pH 9 with NaOMe. The mixture was stirred for 16 h at room temperature and concentrated under reduced pressure to give the intermediate PMB ether.


To a solution of the PMB ether (0.1 mmol) in dichloroethane (5 mL), POCl_3 (0.5 mmol) was added and stirred at room temperature. After completion of the reaction, it was quenched in ice water and the organic layer was separated and the aqueous layer was extracted with dichloroethane (2×5 mL). Combined organic layer was washed with brine solution, dried (Na_2SO_4), concentrated under reduced pressure and the residue was purified by column chromatography (silica gel, EtOAc : Hexane) to afford the corresponding alcohol **A46**.

Example 36.9

A51

Acid **A51** (1 mmol) and amine **A50** (1 mmol) were coupled with 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) hydrochloride (1 mmol) in 3.5 mL methanol for 3 h at room temperature. The mixture was diluted with EtOAc (20 mL) and extracted with water (10 mL), 1 M aq. HCl (10 mL) and sat. aq. NaHCO₃ (10 mL). The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography to give alcohol **A49**.


Example 36.10



A52 **A53**

To a solution of **A53** (80 mmol) in pyridine (60 mL) was added portionwise MsCl (80 mmol) with efficient stirring during 30 min at 20 °C. After being stirred for 30 min, the reaction mixture was maintained for 20 h at room temperature, diluted with CH₂Cl₂ (100 mL), and washed with 2 N aq. HCl until the aqueous washings became acidic. The H₂O layer was extracted with CH₂Cl₂ (3 X 50 mL). The combined organic extract was dried (Na₂SO₄), concentrated, and concentrated to give the intermediate mesylate. Sodium benzylsulfide (4.1 mmol) was added portionwise to a stirred solution of the intermediate mesylate (2.7 mmol) in DMF (5 mL) at room temperature. After 3 h, the mixture was diluted with toluene (100 mL), washed with water (25 mL) and brine (25 mL), dried (MgSO₄), and concentrated in vacuo. The residue was purified by flash chromatography to give thioether **A52**.

Example 36.11




A54 **A55**

To a solution of the diol **A55** (8 mmol) in pyridine (6 mL) was added portionwise TsCl (8 mmol) with efficient stirring during 30 min at 20 °C. After being stirred for 30 min, the reaction mixture was maintained for 20 h at room temperature, diluted with CH_2Cl_2 (100 mL), and washed with 2 N aq. HCl until the aqueous washings became acidic. The H_2O layer was extracted with CH_2Cl_2 (3 X 50 mL). The combined organic extract was dried (Na_2SO_4) and concentrated to give the intermediate tosylate.

A solution of the intermediate tosylate (0.67 mmol) in 5 mL of dry HMPA was cooled in an ice bath under N_2 . This mixture was added to a cold solution of NaSBn (10 mmol) in 20 mL of dry HMPA (prepared from 400 mg of sodium and excess BnSH in dry ether, which was subsequently removed and replaced with HMPA). After the addition, the solution was stored in the freezer (-15 °C) for 14 h. It was then treated with 100 mL of water and extracted three times with ether. The ether extracts were washed four times with water and dried over MgSO_4 . The solvent was then removed under vacuum and the residue (140 mg) chromatographed over silica gel to give thioether **A54**.

10

Example 36.12




- 15 To freshly distilled CH_2Cl_2 (20 mL) was added Et_2Zn (1.0 M in hexanes) (20.0 mmol) under N_2 . The solution was cooled in an ice bath and a solution of trifluoroacetic acid (20.0 mmol) in CH_2Cl_2 (10 mL) was then dripped into the reaction mixture via syringe. Upon stirring for 20 min, a solution of CH_2I_2 (20.0 mmol) in CH_2Cl_2 (10 mL) was added. After an additional 20 min of stirring, a solution of diol **A58** (10.0 mmol) in CH_2Cl_2 (10 mL) was added, and the ice bath was removed. After an additional 30 min of stirring, the reaction mixture was quenched with 0.1 N HCl (50 mL) (alternatively with saturated aqueous NH_4Cl or Et_3N followed by saturated aqueous NaHCO_3) and hexanes (25 mL), and the layers were separated. The aqueous layer was extracted with hexanes. The combined organic layers were washed with saturated NaHCO_3 , H_2O , and brine and then dried (Na_2SO_4), filtered, concentrated, and purified by column chromatography (hexanes/ether = 50/1) to give diol **A57**.

To a solution of the diol **A57** (8 mmol) in pyridine (6 mL) was added portionwise TsCl (8 mmol) with efficient stirring during 30 min at 20 °C. After being stirred for 30 min, the reaction mixture was maintained for 20 h at room temperature, diluted with CH₂Cl₂ (100 mL), and washed with 2 N aq. HCl until the aqueous washings became acidic. The H₂O layer was extracted with CH₂Cl₂ (3 X 50 mL). The combined organic extract was dried (Na₂SO₄) and concentrated to give the intermediate tosylate.

A solution of the intermediate tosylate (0.67 mmol) in 5 mL of dry HMPA was cooled in an ice bath under N₂. This mixture was added to a cold solution of NaSBn (10 mmol) in 20 mL of dry HMPA (prepared from 400 mg of sodium and excess BnSH in dry ether, which was subsequently removed and replaced with HMPA). After the addition, the solution was stored in the freezer (-15 °C) for 14 h. It was then treated with 100 mL of water and extracted three times with ether. The ether extracts were washed four times with water and dried over MgSO₄. The solvent was then removed under vacuum and the residue (140 mg) chromatographed over silica gel to give thioether **A56**.

10

Example 36.13

**A59****A60****A61**

To a solution of diol **A61** (16.1 mmol), triethylamine (24.1 mmol), and DMAP (0.16 mmol) in CH₂Cl₂ (120 mL) was added TBDMSCl (19.3 mmol) in 5 portions over 1 h at 0 °C. The resulting heterogeneous reaction mixture was gradually warmed to rt. The mixture was stirred for 12 h before dilution with water and CH₂Cl₂. The organic layer was washed successively with solutions of saturated aq NaHCO₃, saturated aq NH₄Cl, water and brine, dried over MgSO₄, filtered, and concentrated in vacuo. The pale yellow oil was purified by vacuum chromatography to give the intermediate silyl ether.

A mixture of the intermediate silyl ether (1.0 mmol) and catalyst (Rh or Ru on activated carbon, N.E. Chemcat; 10 wt% of substrate) in iPrOH (1 mL) in a sealed tube was stirred at 60 °C at 5 atm H₂. After cooling to room temperature, the reaction mixture was diluted

15

20


with MeOH (20 mL) and the catalyst was removed by filtration through a membrane filter (Millipore, Millex-LH, 0.45 mm). The filtrate was concentrated in vacuo to give the corresponding decalin.

To a solution of the intermediate decalin alcohol (8 mmol) in pyridine (6 mL) was added portionwise TsCl (8 mmol) with efficient stirring during 30 min at 20 °C. After being stirred for 30 min, the reaction mixture was maintained for 20 h at room temperature, diluted with CH_2Cl_2 (100 mL), and washed with 2 N aq. HCl until the aqueous washings became acidic. The H_2O layer was extracted with CH_2Cl_2 (3 X 50 mL). The combined organic extract was dried (Na_2SO_4) and concentrated to give tosylate **A60**.

A solution of tosylate **A60** (0.67 mmol) in 5 mL of dry HMPA was cooled in an ice bath under N_2 . This mixture was added to a cold solution of NaSBn (10 mmol) in 20 mL of dry HMPA (prepared from 400 mg of sodium and excess BnSH in dry ether, which was subsequently removed and replaced with HMPA). After the addition, the solution was stored in the freezer (-15 °C) for 14 h. It was then treated with 100 mL of water and extracted three times with ether. The ether extracts were washed four times with water and dried over $MgSO_4$. The solvent was then removed under vacuum and the residue (140 mg) chromatographed over silica gel to give the intermediate silyl ether.

The intermediate silyl ether (0.235 mmol) was dissolved in THF (1 mL) at room temperature, followed by addition of 70% HF·pyridine (0.2 mL). After stirring for two days, the reaction mixture was carefully quenched with sat. aq. NaHCO₃ and the resulting solution was diluted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by silica gel column chromatography to give alcohol **A59**.

Example 36.14




A65 A66 A67 A68

A solution of squaric ethyl ester **A67** (0.5 mmol), **A68** (100 mg, 0.588 mmol) and Et₃N (15 drops) in CH₂Cl₂ (10 mL) was stirred at room temperature overnight. Then it was concentrated under reduced pressure. The resulting crude residue, **A66** (1.1 mmol) and Et₃N (15 drops) in CH₂Cl₂ (10 mL) was stirred at room temperature overnight. Then it was concentrated under reduced pressure and purified by column chromatography. PMB-ether (0.444 mmol) was dissolved in acetone (4.5 mL) and water (0.5 mL). CAN (0.845 mmol) was added as a solid, followed by the dropwise addition of a solution of CAN (0.845 mmol) in acetone (0.9 mL) and water (0.1 mL) over 70 minutes. After a further 15 minutes, the reaction was poured into aqueous sodium bicarbonate solution, and extracted with chloroform. The product was purified on chromatography to give **65**.

10


Example 36.15



15


Triethylamine (0.359 mL) was added to a solution of commercially available S-benzyl-(*R*)-cysteinol (2.56 mmol) in THF (7.50 mL). After 10 min of stirring, di-tert-butyl dicarbonate (2.56 mmol) was added at 0 °C. The reaction mixture was stirred for 2 h at room temperature. The solvent was then evaporated under reduced pressure; the residue was dissolved in ethyl acetate and washed with water. The organic layer was separated, dried with Na_2SO_4 , filtered, and evaporated under reduced pressure to yield **A69** as a colorless oil.

Example 37: Further examples of saccharides according to the present invention:




31*

Chemical formula: C₂₅H₄₁N₃O₁₇S; Molecular Weight = 687.68;




32* Chemical formula: $C_{23}H_{36}F_2N_2O_{16}S$; Molecular Weight = 666.61;




33*


Chemical formula: $C_{19}H_{32}N_2O_{10}S$, Molecular Weight = 480.54;



34*

Chemical formula: $C_{12}H_{22}O_6S_2$, Molecular Weight = 326.43;





Chemical formula: C₁₈H₃₂N₂O₁₁S, Molecular Weight = 484.53;

Example 38: Synthesis of glycoconjugates- Conjugation of 2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(thio)ethanol to CRM197

5

To a stirred solution of CRM197 (2 mg, 34.5 nmol) in 0.1 M sodium phosphate buffer (NaPi) pH 7.4 (1.33 mL) was added at room temperature a solution of *N*-Succinimidyl-3-(bromoacetamido)propionate (SBAP) (1.05 mg, 3.4 μ mol) in DMF (40 μ L). The mixture was stirred for 1 h at that temperature, and concentrated using membrane filtration (Amicon 4 mL Ultra centrifuge membranes, 10 kDa cut-off). The protein solution was diluted to 4 mL with sterile water and concentrated again. This process was repeated three times and the solution was diluted to 0.5 mL using sterile water. 20 μ L were taken for analysis, and the protein solution was re-buffered to 0.1 M NaPi pH 8.0 (0.5 mL) using membrane filtration. Disulfide **18*** (1.44 mg, 2.33 μ mol resp. to the monomer) in 0.1 M NaPi pH 8.0 (0.2 mL) was treated at room temperature with tris(2-carboxyethyl)phosphine (TCEP, 25 μ L of a 100 mM stock solution with pH 7.4), left for 1 h at that temperature under an argon atmosphere and added to the solution of the activated protein. The mixture was stirred at room

temperature for 16 h, and washed with sterile water using membrane filtration (see above). Another analytical sample was taken, and the solution was re-buffered to 0.1 M NaPi pH 7.4 (0.5 mL). The glycoconjugate was then treated at room temperature with L-cysteine (0.625 mg, 5.1 μ mol) in 100 μ l sterile water. The mixture was left for 2

5 h at that temperature and purified by membrane filtration. Incorporation of glycan into the glycoconjugate was assessed by MALDI-TOF-MS (positive mode):

Molecular weight measured:

CRM197: 58100 *m/z*

CRM197-SBAP conjugate: 61700 *m/z* (incorporation of approximatively 19 SBAP 10 groups)

CRM197-SBAP-glycoconjugate: 66000 *m/z* (incorporation of approximatively 5.9 molecules of 2-acetamido-4-amino-2,4,6-trideoxy- α -D-galactopyranosyl-(1 \rightarrow 4)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 3)-(α -D-galactopyranosyl)uronate-(1 \rightarrow 1)-2-(thio)ethanol).

15

Example 39: Immunization experiment


Mice (6-8 week old female NMRI mice, Charles River) were immunized s. c. with

20 CRM197-SBAP-glycoconjugate synthesized in **Example 38** (corresponding to 4 μ g synthetic glycan) formulated with or without Alum (Alhydrogel, Brenntag) at a total volume of 100 μ L at days 0, 14 and 28. Control groups comprised mice treated equally with Alum only or PBS. Blood was collected at days 0, 14, 28 and 35 and the immune response was assessed by glycan microarray and ELISA.

25 An immune response against the native Sp1 polysaccharide was found in a subset of mice immunized with glycoconjugate adjuvanted with Alum with an endpoint titer of 500 and 2500 at day 35 compared to day 0, respectively.

Claims


1. A saccharide of general formula (I):




wherein A represents $(\text{CH}_2)_{o1}$;

$o1$ is an integer selected from 1, 2, 3, 4, 5 and 6;


M, N and P represent independently of each other one of the following sugar fragments:



S1



S2




S3

wherein the sugar fragments S1, S2, S3 are connected to each other and to $-\text{O}-\text{A}-\text{SH}$ fragment via O-glycosidic bonds, each sugar fragment S1, S2, and S3 is not more than once present in the general formula (I), sugar fragment S1 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{SH}$ and sugar fragment S3, sugar fragment S3 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{SH}$ and sugar fragment S2, and sugar fragment S2 cannot be simultaneously connected to $-\text{O}-\text{A}-\text{SH}$ and sugar fragment S1, and $n1$, $n2$ and $n3$ are integers selected from 0 and 1, wherein at least one of the integers $n1$, $n2$ and $n3$ is 1

and pharmaceutically acceptable salts of these saccharides.


2. A synthesis of the saccharide of the general formula (I):



wherein A represents $(\text{CH}_2)_{o1}$;


$o1$ is an integer selected from 1, 2, 3, 4, 5 and 6;

M, N and P represent independently of each other one of the following sugar fragments:




wherein the sugar fragments S1, S2, S3 are connected to each other and to –O–A–SH fragment *via* O-glycosidic bonds, each sugar fragment S1, S2, and S3 is not more than once present in the general formula (I), sugar fragment S1 cannot be simultaneously connected to –O–A–SH and sugar fragment S3, sugar fragment S3 cannot be simultaneously connected to –O–A–SH and sugar fragment S2, and sugar fragment S2 cannot be simultaneously connected to –O–A–SH and sugar fragment S1, and n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1 and comprising the steps:




A1) Reacting the compound **2** of the formula:



wherein P¹ – P³ represent protecting groups, with the compound 3 of the formula




wherein P^4 represents a protecting group, in order to obtain compound **4** of general formula:

**4**wherein P¹-P⁴ and A are defined as above;**and**performing removal of protecting groups P¹-P⁴ on compound **4** to afford monosaccharide disulfide **5** of general formula:**5**wherein A is defined as above, and wherein monosaccharide disulfide **5** is further treated with a reducing agent to afford monosaccharide **6** of general formula:**6 (H-S2-O-A-SH)**

wherein A is defined as above;

orperforming selective deprotection on compound **4** to afford compound **7** of general formula




7

wherein P^5 is a protecting group and P^1 , P^3 , P^4 and A are defined as above.


or

A2) Reacting compound 8 of general formula



8

wherein P^6 and P^7 represent protecting groups, with compound 3 to afford compound 9 of general formula




9


wherein P^6 , P^7 and A are defined as above;

and

performing conversion of the azido group to acetamido group and removal of the protecting groups P^4 , P^6 and P^7 on compound 9 to afford monosaccharide disulfide 10 of general formula:




wherein A is defined as above, and wherein monosaccharide **10** is further treated with a reducing agent to afford monosaccharide **11** of general formula:



wherein A is defined as above;

or


performing selective deprotection on compound **9** to afford compound **12** of general formula:



wherein P⁴, P⁷ and A are defined as above.


or

A3) Reacting compound **13** of general formula



13

wherein $P^8 - P^{11}$ represent protecting groups, with compound **3** to afford compound **14** of general formula:




14

wherein $P^4, P^8 - P^{11}$ are defined as above

and

performing selective deprotection of compound **14** to afford compound **15** of general formula:




15

wherein P^4, P^8, P^9, P^{11} and A are defined as above.

and


B1) Reacting compound **7** with compound **13** to afford compound **16** of general formula:



wherein P^1 , P^3 - P^5 , P^8 - P^{11} and A are defined as above;


and

performing removal of protecting groups P^1 , P^3 - P^5 , P^8 - P^{11} on compound **16** to afford disaccharide disulfide **17** of general formula:



wherein A is defined as above and wherein disaccharide disulfide **17** is further treated with a reducing agent to afford disaccharide **18** of general formula:

2014289342 30 Apr 2018




18 (H-S3-S2-O-A-SH)

wherein A is defined as above;

or

performing selective removal of protecting group P^{10} on compound **16** to afford compound **19** of general formula:




19

wherein P^1 , P^3 - P^5 , P^8 , P^9 , P^{11} and A are defined as above.

or

B2) Reacting compound **15** with compound **8** to afford compound **20** of general formula:




20

wherein P^4 , P^6-P^9 , P^{11} and A are defined as above


and

performing conversion of the azido group to acetamido group and removal of the protecting groups P^4 , P^6-P^9 , P^{11} on compound **20** to afford disaccharide disulfide **21** of general formula:



21

wherein A is defined as above and wherein disaccharide disulfide **21** is treated with a reducing agent to afford disaccharide **22** of general formula:




22 (H-S1-S3-O-A-SH)

wherein A is defined as above;

or

performing selective removal of protecting group P⁶ on compound **20** to afford compound **23** of general formula:




23

wherein P⁴, P⁷-P⁹, P¹¹ and A are defined as above;

or

B3) Reacting compound **12** with compound **2** to afford compound **24** of general formula:


2014289342 30 Apr 2018

**24**


wherein P^1 - P^4 , P^7 and A are defined as above,

and

performing conversion of the azido group to acetamido group and removal of protecting groups P^1 - P^4 and P^7 on compound **24** to afford disaccharide disulfide **25** of general formula:

**25**


wherein A is defined as above, and wherein disaccharide disulfide **25** is further treated with a reducing agent to afford disaccharide **26** of general formula:

**26 (H-S2-S1-O-A-SH)**

wherein A is defined as above;

or

performing selective deprotection on compound **24** to afford compound **27** of general formula:




27

wherein P^{12} is a protecting group and P^1 , P^3 , P^4 , P^7 and A are defined as above.

and

C1) Reacting compound **19** with compound **8** to afford compound **28** of general formula:



28


wherein P^1 , P^3 - P^9 , P^{11} and A are defined as above;

30 Apr 2018

2014289342

and


wherein protecting group P^6 is replaced with protecting group P^{13} in order to obtain compound **29** of the following chemical formula:

**29**

wherein P^1 , P^3 - P^5 , P^7 - P^9 , P^{11} , P^{13} and A are defined as above;

and


conversion of compound **29** to trisaccharide disulfide **30** by conversion of the azido group in the acetamido group and cleavage of the protecting group P^1 , P^3 - P^5 , P^7 - P^9 , P^{11} , P^{13} , wherein compound **30** is of general formula:



and wherein A is defined as above;

and

conversion of trisaccharide disulfide **30** to trisaccharide **31** by treatment with a reducing agent, wherein compound **31** is of general formula:




31 (H-S1-S3-S2-O-A-SH)

and wherein A is defined as above.

or


C2) Reacting compound **23** with compound **2** to afford compound **32** of general formula:



32

wherein $P^1 - P^4$, $P^7 - P^9$, P^{11} and A are defined as above;
and

conversion of compound 32 to trisaccharide disulfide 33 by conversion of the azido group in the acetamido group and cleavage of the protecting group $P^1 - P^4$, $P^7 - P^9$, P^{11} , wherein compound 33 is of general formula:




33

wherein A is defined as above;

and

conversion of trisaccharide disulfide **33** to trisaccharide **34** by treatment with a reducing agent, wherein compound **34** is of general formula:




34 (H-S2-S1-S3-O-A-SH)

wherein A is defined as above;

or

C3) Reacting compound **27** with compound **13** to afford compound **35** of general formula:




35

wherein P¹, P³, P⁴, P⁷-P¹¹ and A are defined as above;

and

conversion of compound **35** to trisaccharide disulfide **36** by conversion of the azido group in the acetamido group and cleavage of the protecting group P¹, P³, P⁴, P⁷-P¹¹, wherein compound **36** is of general formula:




2

wherein A is defined as above;

and


conversion of trisaccharide disulfide **36** to trisaccharide **37** by treatment with a reducing agent, wherein compound **37** is of general formula:



37 (H-S3-S2-S1-O-A-SH)

wherein A is defined as above.

3. The synthesis according to claim 2 further comprising step D:
 - D) preparing a salt of the compound of general formula (I) or preparing a lyophilisate of the compound of general formula (I) or of the salt of the compound of general formula (I).
4. The synthesis according to claim 2, wherein the reactions between compounds **2** and **3**, compounds **2** and **12**, and compounds **2** and **23** are performed in presence of DMTST and TTBPY in a mixture of non-polar and polar aprotic solvents.
5. The synthesis according to claim 2, wherein the replacement of protecting group P^6 on compound **28** with protecting group P^{13} to obtain compound **29** is performed in two steps, first involving the reaction of compound **28** with hydrazine or a hydrazinium salt in a solvent or a mixture of solvents, and second by treatment of the product obtained after the first step with $BnOCH_2SCy$, DMTST and TTBPY in a non-polar solvent.
6. The synthesis according to claim 2, wherein the cleavage of the protecting groups involves first cleavage of the base-labile protecting groups by treatment with a base in a mixture of polar aprotic and polar protic solvents; and second cleavage of the protecting groups sensitive to hydrogenation by exposure to sodium and ammonia in a mixture of polar protic and polar aprotic solvents.
7. An intermediate of the general formula (II):




(II)


wherein A represents $(CH_2)_{o1}$;

$o1$ is an integer selected from 1, 2, 3, 4, 5 and 6;


M, N and P represent independently of each other one of the following fragments:



S1



S2



S3

wherein the sugar fragments S1, S2, S3 are connected to each other and to $-O-A-S-$ fragment via O-glycosidic bonds, each sugar fragment S1, S2, and S3 is not more than once present in the fragment $H-(P)_{n3}-(N)_{n2}-(M)_{n1}-O-A-S-$, sugar fragment S1 cannot be simultaneously connected to $-O-A-S-$ and sugar fragment S3, sugar fragment S3 cannot be simultaneously connected to $-O-A-S-$ and sugar fragment S2, sugar fragment S2 cannot be simultaneously connected to $-O-A-S-$ and sugar fragment S1, and n1, n2 and n3 are integers selected from 0 and 1, wherein at least one of the integers n1, n2 and n3 is 1 and pharmaceutically acceptable salts of these saccharides.

8. A glycoconjugate obtained by reacting a saccharide of general formula (I) according to claim 1 with an immunogenic carrier.
9. The glycoconjugate according to claim 8 for use as a vaccine in immunization against diseases associated with bacteria containing in their capsular polysaccharide a saccharide structure selected from:
 - α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp
 - α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp
 - α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp
 - α -D-GalAp
 - α -2,4,6-trideoxy-4-amino-D-GalNAc
 - α -D-GalAp-(1 \rightarrow 3)- α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc
 - α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc
 - α -D-GalAp-(1 \rightarrow 3)- α -2,4,6-trideoxy-4-amino-D-GalNAc-(1 \rightarrow 4)- α -D-GalAp.
10. A pharmaceutical composition comprising the glycoconjugate according to claim 8 and/or the saccharide according to claim 1 and/or the intermediate according to claim 7 together with at least one pharmaceutically acceptable cryoprotectant, lyoprotectant, excipient and/or diluent.
11. The pharmaceutical composition according to claim 10 for use in immunization against diseases associated with *Streptococcus pneumoniae* bacteria.
12. A saccharide according to claim 1 and/or a intermediate according to claim 7 for use as marker in immunological assays for diagnostics of diseases caused by bacteria containing in their capsular polysaccharide a saccharide structure selected from:

- α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp
α-D-GalAp-(1→3)-α-D-GalAp
α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp.
13. A method of vaccinating a subject against diseases associated with bacteria containing in their capsular polysaccharide a saccharide structure selected from:
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp
α-D-GalAp-(1→3)-α-D-GalAp
α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp,
comprising administering to said subject the glycoconjugate according to claim 8.
14. Use of the glycoconjugate according to claim 8 in the manufacture of a vaccine for immunization against diseases associated with bacteria containing in their capsular polysaccharide a saccharide structure selected from:
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp
α-D-GalAp-(1→3)-α-D-GalAp
α-D-GalAp
α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp.
15. The method of claim 13 or the use of claim 14, wherein the bacteria is *Streptococcus pneumoniae* type 1.

16. The method of claim 13 or the use of claim 14, wherein the diseases associated with bacteria include pneumonia, meningitis, otitis media, bacteremia and acute exacerbation of chronic bronchitis, sinusitis, arthritis and conjunctivitis.
17. A method of immunizing a subject against diseases associated with *Streptococcus pneumoniae* bacteria, comprising administering to said subject the pharmaceutical composition according to claim 10.
18. Use of the glycoconjugate according to claim 8 and/or the saccharide according to claim 1 and/or the intermediate according to claim 7 in the manufacture of a medicament for immunization against diseases associated with *Streptococcus pneumoniae* bacteria.
19. The method of claim 17 or the use of claim 18, wherein the *Streptococcus pneumoniae* bacteria is selected from the group containing or consisting of *Streptococcus pneumoniae* type 1, *Streptococcus pneumoniae* type 4, *Streptococcus pneumoniae* type 9V, *Streptococcus pneumoniae* type 2, *Streptococcus pneumoniae* type 19F, *Streptococcus pneumoniae* type 3, *Streptococcus pneumoniae* type 19A, *Streptococcus pneumoniae* type 12F, *Streptococcus pneumoniae* type 31, *Streptococcus pneumoniae* type 7F, *Streptococcus pneumoniae* type 5, *Streptococcus pneumoniae* type 14, *Streptococcus pneumoniae* type 6A, *Streptococcus pneumoniae* type 6B, *Streptococcus pneumoniae* type 18C and *Streptococcus pneumoniae* type 23F.
20. Use of a saccharide according to claim 1 and/or a intermediate according to claim 7 as a marker in immunological assays for diagnostics of diseases caused by bacteria containing in their capsular polysaccharide a saccharide structure selected from:
 - α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp-(1→3)-α-D-GalAp
 - α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp
 - α-D-GalAp-(1→3)-α-D-GalAp
 - α-D-GalAp
 - α-2,4,6-trideoxy-4-amino-D-GalNAc
 - α-D-GalAp-(1→3)-α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
 - α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc
 - α-D-GalAp-(1→3)-α-2,4,6-trideoxy-4-amino-D-GalNAc-(1→4)-α-D-GalAp.



Figure 1**Figure 3**


Figure 2


Figure 4

Figure 5





Figure 6

Serum dilution: 1:50 1:200 1:800 1:3200



Inhibition of binding
with native *Streptococcus*
pneumoniae type 1 polysaccharide

Figure 7




Figure 8

