

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199749301 B2
(10) Patent No. 708197

(54) Title
Improved rapidly disintegratable multiparticulate tablet

(51)⁶ International Patent Classification(s)
A61K 009/20 A61K 009/16

(21) Application No: **199749301** (22) Application Date: **1997.12.30**

(30) Priority Data

(31) Number	(32) Date	(33) Country
9709233	1997.07.21	FR

(43) Publication Date : **1999.02.04**

(43) Publication Journal Date : **1999.02.04**

(44) Accepted Journal Date : **1999.07.29**

(71) Applicant(s)
Laboratoires Prographarm

(72) Inventor(s)
Charles Chauveau; Edouard Gendrot; Alain Gilles Demichelis; Nouredine Nouri

(74) Agent/Attorney
**WATERMARK PATENT and TRADEMARK ATTORNEYS, Locked Bag 5, HAWTHORN
VIC 3122**



(12) PATENT ABSTRACT (11) Document No. AU-A-49301/97
(19) AUSTRALIAN PATENT OFFICE

- (54) Title
IMPROVED RAPIDLY DISINTEGRATABLE MULTIPARTICULATE TABLET
- International Patent Classification(s)
(51)⁶ **A61K 009/20 A61K 009/16**
- (21) Application No. : **49301/97** (22) Application Date : **30/12/97**
- (30) Priority Data
- (31) Number (32) Date (33) Country
97 09233 21/07/97 FR FRANCE
- (43) Publication Date : **04/02/99**
- (71) Applicant(s)
LABORATOIRES PROGRAPHARM
- (72) Inventor(s)
CHARLES CHAUVEAU; EDOUARD GENDROT; ALAIN GILLES DEMICHELIS; NOUREDDINE NOURI
- (74) Attorney or Agent
WATERMARK PATENT & TRADEMARK ATTORNEYS , Locked Bag 5, HAWTHORN VIC 3122
- (57) Claim

1. Improved multiparticulate tablet which disintegrates in the mouth in less than 40 seconds and comprises on the one hand an active ingredient in the form of coated microcrystals and on the other hand an excipient, characterized in that

- the excipient comprises at least one disintegration agent and at least one soluble diluent agent with binding properties consisting of a polyol having less than 13 carbon atoms either in the form of the directly compressible product the average particle diameter of which is from 100 to 500 micrometres or in the form of a powder the average particle diameter of which is less than 100 micrometres, this polyol preferably being selected from the group comprising mannitol, xylitol, sorbitol and maltitol, it being understood that sorbitol cannot be used alone and that, when there is a single soluble diluent agent with binding properties, it is used in the form of the directly compressible product whereas, when there are at least two soluble diluent agents with binding properties, one is in the directly compressible form and the other in powder form,

in which case the polyol is the same, the proportion of directly compressible polyol to powder polyol being from 99/1 to 50/50 and preferably from 80/20 to 50/50 and

- the coating of the microcrystals of active ingredient comprises at least one coating agent selected as a function of the physico-chemical characteristics of the active ingredient from the group comprising poly-methacrylates, cellulose polymers, in particular ethyl celluloses, hydroxypropyl-methyl celluloses, hydroxypropyl celluloses and cellulose acetophthalates and combinations of these polymers with each other, optionally combined with plastifiers or soluble agents, in particular polyols.

AUSTRALIA

Patents Act 1990

**ORIGINAL
COMPLETE SPECIFICATION
STANDARD PATENT**

Application Number:

Lodged:

Invention Title: IMPROVED RAPIDLY DISINTEGRATABLE MULTIPARTICULATE
 TABLET

The following statement is a full description of this invention, including the best method of performing it known to us :-

**IMPROVED RAPIDLY DISINTEGRATABLE
MULTIPARTICULATE TABLET**

The invention relates to an improved rapidly disintegratable multiparticulate tablet of the type which disintegrates in the mouth in less than 5 40 seconds and which comprises, on the one hand, an active ingredient in the form of microcrystals coated with a coating agent and, on the other hand, an excipient.



The active ingredient may be selected from the group comprising antalgics, antipyretics, antidiarrheals, antispasmodics, digestive motoricity 10 regulators and anti-inflammatories and more particularly from the group comprising paracetamol, ibuprofen, aspirin, ketoprofen and loperamide.

French Patent FR 91 09245 discloses multiparticulate tablets of the type in question which are generally satisfactory; however, the texture of some of these tablets leads to a gritty and pasty sensation on ingestion.

15 The object of the invention is above all to overcome this drawback and to supply a rapidly disintegratable tablet with a systematically pleasant texture, this tablet also leading to an optimum biological availability of the active ingredient.

The Applicants have had the merit of discovering, following extensive 20 research, that this object could be achieved on the one hand by incorporating into the excipient a disintegration agent and at least one particular soluble diluent agent with binding properties and, on the other hand, by selecting the coating agent as a function of the physico-chemical characteristics of the active ingredient.

As a result, the improved multiparticulate tablet according to the invention 25 is



- a multiparticulate tablet which disintegrates in the mouth in less than 40 seconds and including an excipient and an active ingredient in the form of microcrystals which have a taste-masking coating, wherein the excipient includes, with respect to the mass of the tablet, from 3 to 15% by weight of at least one disintegration agent and from 40 to 90% by weight of at least one soluble diluent agent with binding properties, said diluent agent consisting of a polyol having less than 13 carbon atoms, said polyol being either in the directly compressible form which is composed of particles whose average diameter is from 100 to 500 micrometers, or in the powder form which is composed of particles whose average diameter is less than 100 micrometres, said polyol being selected from the group consisting of mannitol, xylitol, sorbitol and maltitol, with the proviso that, when only one soluble diluent agent with binding properties is used, it is a polyol in the directly compressible form except sorbitol and, when at least two soluble diluent agents with binding properties are used, one is a polyol in the directly compressible form and the other is the same or another polyol in powder form, the proportion of directly compressibly polyol to powder polyol being from 99/1 to 50/50 and the coating of the microcrystals of active ingredient includes at least one coating agent selected as a function of the physico-chemical characteristics of the active ingredient from the group consisting of polymethacrylates and cellulose polymers and combinations thereof with each other.

The invention will be better understood using the additional description and non-limitative examples relating to advantageous methods of implementation which follow.

With the intention of preparing a multiparticulate tablet of the type in question, the procedure is as follows or in an equivalent fashion.



5 First the active ingredient is selected from the group comprising antalgics, antipyretics, antidiarrheals, antispasmodics, digestive motoricity regulators and anti-inflammatory and more particularly from the group comprising paracetamol, ibuprofen, aspirin, ketoprofen and loperamide.

10 The excipient is prepared by mixing at least one disintegration agent with at least one soluble diluent agent with binding properties consisting of a polyol having less than 13 carbon atoms either in the form of a directly compressible product the average particle diameter of which is from 100 to 500 micrometres or in the form of a powder the average particle diameter of which is less than 100 micrometres, this polyol preferably being selected from the group comprising mannitol, xylitol, sorbitol and maltitol, 15 it being understood that sorbitol cannot be used alone.

20 When there is a single soluble diluent agent with binding properties, therefore different from sorbitol, it is used in the form of a directly compressible product.

25 When at least two soluble diluent agents with binding properties are used, one is present in the form of the directly compressible product and the other, which can be the same polyol, is in the form of a powder the average component particle diameter of which is less than 100 micrometres, the proportion of directly compressible polyol to powder polyol being from 99/1 to 50/50 and preferably from 80/20 to 50/50.

30 The disintegration agent is preferably selected from the group comprising cross-linked polyvinylpyrrolidone, designated in the art by the term crospovidone and cross-linked sodium carboxymethylcellulose designated in the art by the term sodium croscarmellose.

35 The respective proportions of disintegration agent and binding agent or soluble diluent used for the constitution of the excipient are, with respect to the mass

of the tablet, from 3 to 15% by weight, preferably from 5 to 10% by weight for the first and from 40 to 90% by weight, preferably from 50 to 70% by weight for the second; the maximum proportion of sorbitol is 30% by weight.

5 The active ingredient which is a component of the composition of the tablet according to the invention is presented before coating in the form of microcrystals the average diameter of which is from approximately 1 to approximately 500 micrometres.

10 In the case of certain active ingredients, microcrystals with an average diameter of less than 100 micrometres are preferably used in order to increase the exchange surface between the active ingredient and the ambient environment; by proceeding in this way, the
15 solubilization speed and/or the intrinsic solubility of dissolution are optimized.

 The microcrystals are preferably coated using the technique known as the fluidized bed.

 Direct coating is applied in the case of
20 microcrystals having an average size of 100 to 500 micrometres; for microcrystals smaller than 100 micrometres, a prior treatment is carried out consisting either in the granulation of the microcrystals by a standard wet or dry granulation process, or by prior fixing of the microcrystals
25 on neutral supports known in themselves when the average size of said microcrystals is less than approximately 20 micrometres; the microcrystals are fixed to the neutral supports in a standard manner using a binding agent consisting for example of hydroxypropylmethyl cellulose.

30 The coating is obtained from at least one coating agent selected, according to the physico-chemical characteristics of the active ingredient, from the group comprising the polymethacrylates, in particular certain of those marketed under the EUDRAGIT trade mark and more
35 particularly the 30% dispersions of poly(ethylacrylate-

methyl-metacrylate) marketed under the trade mark EUDRAGIT NE 30 D, the type E aminoalkyl-methacrylate copolymers marketed under the trade mark EUDRAGIT E, the cellulose polymers, in particular the ethyl celluloses, hydroxypropyl-methyl celluloses, hydroxypropyl celluloses and the cellulose acetophthalate, the combinations of these polymers with each other and optionally in association with plastifiers, for example polyethyleneglycol 6000, or with soluble agents, in particular polyols, for example mannitol.

As an example, the coating can be constituted starting from

- EUDRAGIT NE 30 D alone or mixed with EUDRAGIT E in one or more organic solvents,
- ethyl cellulose alone or mixed with hydroxypropyl-methyl cellulose in association with a plastifier optionally in the presence of a hydroalcoholic solvent,
- a polymethacrylate, in particular EUDRAGIT NE 30 D mixed with a soluble cellulose derivative, in particular hydroxypropyl-methyl cellulose and a plastifier and/or a soluble diluent agent with binding properties,
- EUDRAGIT E 100 alone.

Due to the coating according to the invention of the microcrystals of active ingredient, said coating comprising in particular a in association or combination a soluble polymer and an insoluble polymer, the definitive tablet is preferably characterised in that

- on the one hand, in an acid medium with a pH lower than 5, the quantity of active ingredient which is dissolved, after disintegration of the tablet, in 5 to 20 minutes from the coated microcrystals is equal to at least 80% and preferably to at least 100% of the quantity of active ingredient which is dissolved in the same time period following disintegration from a tablet allowing the immediate release of the active ingredient consisting of microcrystals but in which said microcrystals are not coated



and,

- on the other hand, the active ingredient does not dissolve significantly after remaining for a period of less than 5 minutes in a medium the pH conditions of which are similar to those of saliva, namely 7.0 ± 0.5 , thus ensuring satisfactory masking of its taste.

When the quantity of active ingredient which is dissolved after disintegration of the tablet in 5 to 20 minutes from the coated microcrystals is equal to at least 100% of the quantity of active ingredient which is dissolved in the same time period after disintegration from a tablet allowing the immediate release of the active ingredient consisting of microcrystals but in which said microcrystals are not coated, the biological availability of the active ingredient is at least equivalent to that of the same active ingredient obtained from the above tablet in which said microcrystals are not coated.

For the production of the tablet, a mixture of the excipient and the coated microcrystals is first prepared and this mixture is then homogeneized in a dry mixer.

Preferably, a sweetener, a flavouring and a lubricant are incorporated to this mixture.

The sweetener can be selected from the group comprising Aspartame and sodium saccharinate, and the lubricant from the group comprising magnesium stearate, sodium stearyl fumarate, stearic acid and polyethyleneglycol 6000.

The mixture is then subjected to a compression force sufficient to confer on the resulting tablet a sufficient hardness for it to be handled and packaged industrially, then carried and handled by the patient without any particular precautions; for information only, it is indicated that the hardnesses meeting these conditions are generally comprised between 20 and 70 Newtons.

The tablets according to the invention present, with

respect to the tablets of the type in question which already exist, simultaneously an improvement in the speed at which the active ingredient is made available in the organism and an improvement in palatability.

5

EXAMPLE 1

Multiparticulate paracetamol tablet dosed at 500 mg.

The composition of the above tablet results from Table I below.

10

TABLE I

Constituents	Centesimal formula
Coated paracetamol	39.2
Mannitol for direct compression	36.7
Crystalline powder mannitol	12.3
Crospovidone	8.6
Aspartame	2.7
Blackcurrant flavouring	0.4
Magnesium stearate	0.1
Total	100.0%

15

20

This tablet is prepared as indicated below.

The paracetamol microcrystals are introduced into a fluidized bed installation and a dispersion in ethanol of EUDRAGIT E 100, EUDRAGIT NE 30 D and colloidal silica is sprayed onto the microcrystals in order to obtain microcrystals coated with 10% of polymer.

25

All the excipients are sieved and the mixture containing the coated paracetamol and the excipients is homogenized in a dry mixer.

30

The mixture is divided and moulded on a tableting machine equipped with 17 mm diameter punches.

The compression force is adjusted in order to obtain tablets having a hardness of 40 ± 10 Newtons.

35

The disintegration time in the mouth of the tablets thus obtained is less than 40 seconds.

This time corresponds to the time which separates,

on the one hand, the moment the tablet is placed in contact with the saliva in the mouth and, on the other hand, the moment the suspension, resulting from the disintegration of the tablet in contact with the saliva, is swallowed.

5 The disintegration time less than 40 seconds indicated above is the average of the values noted for a representative group of healthy subjects.

10 The pharmacokinetic study to which the above tablet was subjected shows that the biological availability of the active ingredient, namely paracetamol, is not significantly different from that which is observed after administration of a commercially available tablet based on paracetamol. The masking of the taste of the paracetamol by the coating present on the tablet according to the invention does not
15 therefore result in any delay in absorption, which is even slightly more rapid as the maximum is achieved in 0.5 hour on average whereas it is only achieved in 0.88 hour on average in the case of the commercially available reference tablet.

20 The values recorded with respect to the properties of the tablet according to the invention and those relating to a currently marketed paracetamol-based tablet with immediate release of the active ingredient and not previously coated are shown in Table II below.

25 **TABLE II**

Parameters	Tablet containing a 500 mg dose of paracetamol according to the invention	Commercially available tablet containing a 500 mg dose of paracetamol
Lag t (h)	0.17	0.17
t _{max} (h)	0.50	0.88
C _{max} (µg/ml)	6.28±1.61	6.26±2.37
AUC _{0-t} (µg.h/ml)	18.59±3.44	18.10±3.40
AUC _{inf} (µg.h/ml)	19.68±3.85	19.24±3.79

In this table,

- lag t designates the time in hours which elapsed between the administration of the medicament and the detection of the active ingredient in the subject's blood,
- 5 - tmax designates the time in hours after which the serous concentration of active ingredient reaches a maximum,
- Cmax ($\mu\text{g/ml}$) designates the value of the maximum concentration of active ingredient reached at the value tmax; this concentration is expressed in μg of active
10 ingredient per ml of serum,
- AUC0-t designates the surface area under the curve of the serous concentrations of active ingredient as a function of time until the last quantified sample is taken, and
- AUCinf designates the surface area under the curve of the
15 serous concentrations of active ingredient as a function of time extrapolated to infinity.

An examination of the results shown in Table II shows that the coating used to mask the taste of the active ingredient does not result in a modification of the
20 biological availability of the active ingredient or any delay in absorption, which is a determining result in the administration of certain active ingredients where rapid action is desired, in particular antalgics.

25 **EXAMPLE 2**

Multiparticulate loperamide tablet at a dose of 2 mg.

The composition of the above tablet results from Table III below.

30

TABLE III

Constituents	Centesimal formula
Coated loperamide	15.1
Mannitol for direct compression	56.6
Crystalline powder mannitol	18.8
Crospovidone	5.0
Aspartame	3.0
Mint flavouring	0.7
Magnesium stearate	0.8
Total	100.0%

This tablet is prepared in the manner indicated in Example 1.

The active ingredient is presented in the form of microcrystals the average size of which is less than 20 micrometres; consequently, fixing of these microcrystals on neutral supports consisting of a polyol and having an average size of the order of 60 micrometres is carried out using a hydroxypropyl-methyl cellulose solution.

The neutral supports comprising the loperamide microcrystals are then coated with 20% of a mixture of EUDRAGIT NE 30 D and EUDRAGIT E 100.

This tablet has a hardness of 30±5 Newtons.

It disintegrates in the mouth in 20 seconds.

EXAMPLE 3

Ketoprofen tablet at a dose of 12.5 mg.

The composition of the above tablet results from Table IV below.

TABLE IV

Constituents	Centesimal formula
Coated ketoprofen	6.7
Mannitol for direct compression	63.3
Crystalline powder mannitol	22.2
Crospovidone	5.0
Aspartame	1.3
Mint flavouring	0.7
Magnesium stearate	0.8
Total	100.0%

It is prepared in the manner indicated in Example 2.

The active ingredient is presented in the form of microcrystals the average size of which is less than 20 micrometres; consequently, fixing of these microcrystals on neutral supports consisting of a polyol and the size of which is of the order of 60 micrometres is carried out using a hydroxypropyl-methyl cellulose solution.

The neutral supports comprising the ketoprofen microcrystals constituting the above tablet are then coated with 20% of a mixture of EUDRAGIT NE 30 D, hydroxypropyl-methyl cellulose and polyethylene-glycol 6000.

It has a hardness of 35±5 Newtons.

It disintegrates in the mouth in 20 seconds.

EXAMPLE 4

Multiparticulate acetylsalicylic acid tablet at a dose of 325 mg.

The composition of the above tablet results from Table V below.

TABLE V

Constituents	Centesimal formula
Coated acetylsalicylic acid	37.4
Mannitol for direct compression	38.0
Crystalline powder mannitol	12.6
Crospovidone	8.6
Aspartame	2.9
Sodium stearylfumarate	0.5
Total	100.0%

This tablet is prepared in the manner indicated in Example 1.

The acetylsalicylic acid microcrystals constituting the above tablet are coated with 4.5% of ethyl cellulose N7 and polyethyleneglycol 6000.

The tablet has a hardness of 50±15 Newtons.

It disintegrates in the mouth in 20 seconds.

EXAMPLE 5

Multiparticulate ibuprofen tablet at a dose of 100 mg.

The composition of the above tablet results from Table VI below.

TABLE VI

Constituents	Centesimal formula
Coated ibuprofen	14.4
Mannitol for direct compression	52.5
Crystalline powder mannitol	17.5
Crospovidone	10.0
Aspartame	3.7
Cherry flavouring	0.9
Magnesium stearate	1.0
Total	100.0%

This tablet is prepared in the manner indicated in Example 1.

5 The ibuprofen microcrystals having an average size in the region of 30 micrometres are granulated by a standard wet route granulation process. The resulting particle is then coated with 20% of a mixture of ethyl cellulose N7 and polyethyleneglycol 6000.

The resulting coated ibuprofen crystals are mixed with the excipient, then subjected to compression.

10 The resulting tablet has a hardness of 40 ± 10 Newtons.

It disintegrates in the mouth in 30 seconds.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A multiparticulate tablet which disintegrates in the mouth in less than 40 seconds and including an excipient and an active ingredient in the form of microcrystals which have a taste-masking coating, wherein

- the excipient includes, with respect to the mass of the tablet, from 3 to 15% by weight of at least one disintegration agent and from 40 to 90% by weight of at least one soluble diluent agent with binding properties, said diluent agent consisting of a polyol having less than 13 carbon atoms, said polyol being either in the directly compressible form which is composed of particles whose average diameter is from 100 to 500 micrometers, or in the powder form which is composed of particles whose average diameter is less than 100 micrometres, said polyol being selected from the group consisting of mannitol, xylitol, sorbitol and maltitol, with the proviso that, when only one soluble diluent agent with binding properties is used, it is a polyol in the directly compressible form except sorbitol and, when at least two soluble diluent agents with binding properties are used, one is a polyol in the directly compressible form and the other is the same or another polyol in powder form, the proportion of directly compressible polyol to powder polyol being from 99/1 to 50/50 and

- the coating of the microcrystals of active ingredient includes at least one coating agent selected as a function of the physico-chemical characteristics of the active ingredient from the group consisting of polymethacrylates and cellulose polymers and combinations thereof with each other.

2. A multiparticulate tablet according to Claim 1, wherein the proportion of disintegrating agent and of soluble diluent agent with binding properties is respectively from 5 to 10% and from 50 to 70% by weight.

3. A multiparticulate tablet according to any one of Claims 1 and 2, wherein the maximum proportion of sorbitol is 30% by weight.



4. A multiparticulate tablet according to any one of Claims 1 to 3, wherein the proportion of directly compressible polyol to powder polyol is from 80/20 to 50/50.

5. A multiparticulate tablet according to any one of Claims 1 to 4, wherein the coating agent is selected from the group consisting of ethyl celluloses, hydroxypropyl-methyl celluloses, hydroxypropyl celluloses and cellulose acetophthalates.

6. A multiparticulate tablet according to any one of Claims 1 to 5, wherein the coating agent is combined with at least one of the products of the group consisting of plastifiers, soluble agents and polyols.

7. A multiparticulate tablet according to any one of Claims 1 to 6, wherein the biological availability of the active ingredient is at least equivalent to that of the active ingredient of the same tablet wherein the said active ingredient is not coated, allowing thus its immediate release.

8. A multiparticulate tablet according to Claim 7, wherein the quantity of active ingredient which is dissolved from the coated microcrystals, after disintegration of the tablet, in 5 to 20 minutes when placed in an acid medium having a pH lower than 5, is equal to at least 80% of the quantity of active ingredient which is dissolved in the same time after disintegration from a tablet which is identical except the fact that the microcrystals are not coated.

9. A multiparticulate tablet according to Claim 7, wherein the quantity of active ingredient which is dissolved from the coated microcrystals, after disintegration of the tablet, in 5 to 20 minutes when placed in an acid medium



having a pH lower than 5, is equal to at least 100% of the quantity of active ingredient which is dissolved in the same time after disintegration from a tablet which is identical except the fact that the microcrystals are not coated.

DATED this 26th day of May 1999

LABORATOIRES PROGRAPHARM

WATERMARK PATENT & TRADEMARK ATTORNEYS
290 BURWOOD ROAD
HAWTHORN VICTORIA 3122
AUSTRALIA

VAX DOC027 AU4930197.WPC : KJS/KMH/RES

1999

55

17



IMPROVED RAPIDLY DISINTEGRATABLE MULTIPARTICULATE TABLET

A B S T R A C T

The invention relates to an improved rapidly disintegratable multiparticulate tablet of the type which disintegrates in the mouth in less than 40 seconds and which comprises, on the one hand, an active ingredient in the form of microcrystals coated with a coating agent and, on the other hand, an excipient.

No figure.