



(22) Date de dépôt/Filing Date: 2016/07/22

(41) Mise à la disp. pub./Open to Public Insp.: 2017/01/23

(45) Date de délivrance/Issue Date: 2022/06/21

(30) Priorité/Priority: 2015/07/23 (US62/195.936)

(51) Cl.Int./Int.Cl. C12N 7/00 (2006.01),
C12N 5/071 (2010.01), C12N 7/02 (2006.01),
C12Q 1/70 (2006.01), G01N 33/48 (2006.01),
G01N 33/50 (2006.01)

(72) Inventeurs/Inventors:

BUNO, BRETT, US;
JOURNIGAN, TERRI, US;
HOTTA, JOANN, US;
BURDICK, MICHAEL, US



(73) Propriétaire/Owner:

GRIFOLS, S.A., ES

(74) Agent: ROBIC

(54) Titre : METHODE DE PRODUCTION DE STOCKS DE VIRUS DE L'HEPATITE E A HAUT DEGRE ET ESSAIS DE TITRAGE POUR LE VIRUS DE L'HEPATITE E

(54) Title: METHOD FOR GENERATING HIGH-TITER HEPATITIS E VIRUS STOCKS AND TITRATION ASSAY FOR HEPATITIS E VIRUS



(57) Abrégé/Abstract:

Provided herein are the use of polybrene™ as an additive in cell culture medium used in methods for the generation of high-titer hepatitis E virus stocks and assays for titration of hepatitis E virus. Also provided are a cell culture medium containing polybrene for high-titer HEV generation, a method for determining the presence and/or the level of HEV in a sample, and an HEV titration assay using polybrene.

ABSTRACT

Method for generating high-titer hepatitis E virus stocks and titration assay for hepatitis E virus

5

Provided herein are the use of polybrene™ as an additive in cell culture medium used in methods for the generation of high-titer hepatitis E virus stocks and assays for titration of hepatitis E virus. Also provided are a cell 10 culture medium containing polybrene for high-titer HEV generation, a method for determining the presence and/or the level of HEV in a sample, and an HEV titration assay using polybrene.

Method for generating high-titer hepatitis E virus stocks
and titration assay for hepatitis E virus

5

DESCRIPTION

PRIORITY AND CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of priority to U.S. Provisional Application 62/195,936, filed on July 23, 10 2015.

REFERENCE TO SEQUENCE LISTING

The present application is being filed along with a Sequence Listing in electronic format. The Sequence 15 Listing is provided as a file entitled DURC6_006AUS_SEQLIST.txt which is 1,093 bytes in size, created on May 31, 2016 and last modified on May 31, 2016.

BACKGROUND

20 Field

The present invention refers to the field of virology, more precisely to methods for generating Hepatitis E Virus (hereinafter HEV) stocks. Specifically, the present invention discloses methods for propagating and titrating 25 high titer HEV stocks.

Description of the Related Art

HEV (genus *Hepevirus*, family *Hepeviridae*) is a small non-enveloped/pseudo-enveloped virus, with a single-stranded 30 positive-sense, polyadenylated RNA genome of approximately

7.2 kb. There are four genotypes of HEV that have been identified, but only one serotype. Genotypes 1 and 2 are mainly responsible for waterborne infections in underdeveloped countries and cause disease primarily in 5 humans and higher primates. Infections are typically self-resolving and acute, lasting at most 2 to 7 weeks, but may be fatal especially in pregnant women. Genotypes 3 and 4 are associated with endemic (autochthonous) infections in industrialized countries. These two genotypes cause 10 disease largely in swine but humans can become accidental hosts as a result of food or zoonotic exposure. Clinical disease is typically asymptomatic and mild in young adults, but can become clinically apparent in older men. In addition genotype 3 and 4 infections can become chronic 15 in immune-suppressed persons, such as organ-transplant patients or AIDS patients.

Recently hepatitis E has been categorized as a transfusion transmittable infectious disease. Given the worldwide 20 spread of HEV in recent years, concerns have been raised regarding the safety of blood- and plasma-derived products. The virus safety profile of blood- and plasma-derived products can be assured by performing clearance studies that demonstrate the virus reduction and/or 25 clearance capacity of their manufacturing processes. During these clearance studies a known amount of virus is deliberately spiked into a blood or a plasma product intermediate and then the spiked material is processed using a bench scale model of the manufacturing process. 30 Virus reduction and/or clearance across a step is determined by comparing the amount of virus before and after treatment.

Virus clearance studies require large quantities of high titer virus and the lack of an efficient cell culture system for HEV has hampered the ability to perform such studies for HEV. Several HEV cell culture systems have 5 recently been developed to address this problem.

A genotype 3 and a genotype 4 strain were adapted by Okamoto and colleagues to grow in A549 human lung cells reaching HEV RNA titers of 3.9×10^8 copies/mL. In 10 addition, said genotype 4 strain was also cultured in PLC/PRF/5 cells (human hepatoma cells) but with lower titers.

A second genotype 3 (strain Kernow-C1) was adapted by 15 Emerson and colleagues to grow in HepG2/C3A human hepatoma cells obtaining a titer of 4.61×10^8 genomes/mL after 6 passages. Studies showed that the adaptation for growth in vitro resulted after the acquisition of 174 ribonucleotides of the S17 human ribosomal protein gene. 20 Because genomes with the same insertion were detected in the original virus inoculum, a fecal suspension from a chronically infected HIV-1 patient, the recombination/insertion event had occurred naturally and was not an artifact of cell culture. Attempts to grow said strain in 25 PLC/PRF/5 and A549 cells were unsuccessful or resulted in lower titers, but the virus infected and replicated in kidney cells from swine, the major zoonotic host for genotype 3 viruses.

30 Human hepatoma cells are difficult to grow and may require special cell plating methods including the use of coatings such as collagen, fibronectin, gelatin and/or poly-L-lysine to facilitate cell attachment and/or cell growth.

In addition, not all of these coatings work well for all cell types.

SUMMARY

5 In some embodiments, there is a method of producing a high titer of Hepatitis E Virus (HEV), the method comprising:
a) culturing a cell line *in vitro*, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741, in a medium
10 comprising polybrene in a concentration of about 1 μ g/mL - about 5 μ g/mL, and b) infecting the cell line with HEV.

In some embodiments, a method of producing a high titer of Hepatitis E Virus (HEV) is provided, the method
15 comprising:

a) culturing a cell line *in vitro*, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741, in a medium comprising
20 polybreneTM also known as 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide in a concentration of about 1 μ g/mL - about 5 μ g/mL,
b) infecting the cell line with HEV, and
25 c) culturing the infected cells for approximately one week,

wherein the high titer of HEV is about 10^7 RNA copies/mL - about 10^{10} RNA copies/mL.

30 In some embodiments, there is a method of determining a presence and/or a level of HEV in a sample, the method comprising the steps of: a) providing the sample to a

mixture comprising a cell line, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741, and a culture medium, wherein the culture medium comprises polybrene in 5 a concentration of about 1 μ g/mL - about 5 μ g/mL; b) incubating the mixture comprising the sample from step a), to allow for the propagation of HEV, if present in the sample; c) collecting a portion of step b), the portion comprising HEV, if present and propagated during step b); 10 and d) measuring the presence and/or the level of a biological substance associated with HEV in the collected portion.

In some embodiments, a method of determining a presence 15 and/or a concentration of HEV in a sample is provided, the method comprising the steps of:

- a) providing the sample to a mixture comprising a cell line, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741, and a culture medium, wherein the culture medium comprises polybreneTM also known as 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide in a concentration of about 1 μ g/mL - about 5 μ g/mL;
- b) incubating the mixture comprising the sample from step a) for approximately one week, to allow for the propagation of HEV, if present in 25 the sample;

- c) collecting a portion of step b), the portion comprising HEV, if present and propagated during step b); and
- d) measuring the presence and/or the concentration of a HEV polynucleotide or polypeptide in the collected portion.

5 In some embodiments, there is a culture medium for generating a high-titer HEV, the culture medium comprising polybrene in a range of about 1 μ g/mL - about 5 μ g/mL.

10 In some embodiments, there is a culture medium for generating a high-titer HEV, the culture medium comprising polybreneTM also known as 1,5-dimethyl-1,5-diazaundecamethylene polymetho-bromide in a range of about 1 μ g/mL - about 5 μ g/mL, wherein the high-titer HEV is about 10^7 RNA copies/mL - about 10^{10} RNA copies/mL.

15 In some embodiments, there is an HEV titration assay comprising the use of a culture medium comprising polybrene in the range of about 1 μ g/mL - about 5 μ g/mL and a cell line, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741.


20

In some embodiments, there is an HEV titration assay comprising the method of determining a presence and/or a concentration of HEV in a sample as defined herein.

25 In some embodiments, a method of producing a high- titer of Hepatitis E Virus is provided, the method comprising: culturing a cell line *in vitro* in a medium comprising a

concentration of polybrene, and infecting the cell line with HEV. In some embodiments of the method, the cell line used is HepG2 (ATCC number HB-8065) or HepG2/C3A (ATCC number CRL-10741). In some embodiments of the method, the 5 concentration of polybrene is about 1 μ g/mL - about 5 μ g/mL. In some embodiments of the method, the high titer is about 10^8 copies/mL - about 10^{10} copies/mL. In some embodiments of the method, the high titer is about 10^7 copies/mL - about 10^{10} copies/mL. In some embodiments, the 10 method further comprises the steps of adding a concentration of polybrene to the medium; a passaging the HEV-infected cell line in the medium comprising polybrene; and collecting the medium and/or infected cells. In some embodiments of the method, the concentration of polybrene 15 is about 1 μ g/mL - about 5 μ g/mL. In some embodiments of the method, the high titer of HEV obtained in the medium and/or infected cells is about 10^8 copies/mL - about 10^{10} copies/mL. In some embodiments of the method, the high titer of HEV obtained in the medium and/or infected cells 20 is about 10^7 copies/mL - about 10^{10} copies/mL.

In some embodiments, there is a method of determining a presence and/or a level of HEV in a sample, the method comprising the steps of: providing the sample to a mixture 25 comprising a cell line and a culture medium, wherein said



culture medium comprises polybrene; incubating the mixture comprising the sample from the previous step to allow for the propagation of the HEV, if present in the sample; collecting a portion of the previous step, said portion comprising HEV, if present and propagated during the previous step; and measuring the presence and/or the level of a biological substance associated with HEV in the collected portion. In some embodiments of the method, the cell line is selected from the group consisting of cell lines of HepG2 (ATCC number HB-8065) and HepG2/C3A (ATCC number CRL-10741). In some embodiments of the method, said biological substance comprises a polynucleotide and/or a polypeptide sequence of HEV. In some embodiments of the method, said measuring comprises the steps of: providing a first reaction mixture by mixing the collected portion with a first solution so as to expose a polynucleotide of HEV, if HEV is present in the collected portion, wherein the polynucleotide is an RNA of HEV; providing a second reaction mixture by adding, to the first reaction mixture, a first reagent so as to produce a complementary deoxyribonucleic acid (cDNA) that is at least partially complementary to the RNA of HEV; providing a third reaction mixture by adding, to the second reaction mixture, a second reagent comprising a pair of polynucleotides to amplify a sequence of the cDNA that is at least partially complementary to each of the pair of polynucleotides; providing a fourth reaction mixture by amplifying the sequence; and determining a concentration of the amplified sequence in the fourth reaction mixture. In some embodiments of the method, said determining the concentration comprises: providing the fourth reaction mixture; providing a one or more controls comprising a predetermined amount of the cDNA of HEV; adding an agent

to the fourth reaction mixture and the one or more controls, wherein said agent is at least partially specific to the amplified sequence in the fourth reaction mixture and the predetermined amount of the cDNA of HEV in 5 the one or more controls; and calculating a level of recognition of HEV by the agent in the fourth reaction mixture relative to the one or more controls. In some embodiments of the method, said pair of polynucleotides comprises:

10

- a) 5'-CGGCTATCGGCCAGAAGTT-3' (SEQ ID NO: 1)
- b) 5'-CCGTGGCTATAACTGTGGTCT-3' (SEQ ID NO: 2)

In some embodiments of the method, the agent comprises:

15

5'-FAMTM-TTTTACGC-ZENTM-AGGCTGCCAAGGCC-3IABkFQTM-3' (SEQ ID NO: 3)

In some embodiments, a culture medium for generating a 20 high-titer HEV comprising polybrene in the range of about 1 μ g/mL - about 5 μ g/mL is provided. In some embodiments, the high-titer HEV is about 10⁸ copies/mL - about 10¹⁰ copies/mL. In some embodiments, the high-titer HEV is about 10⁷ copies/mL - about 10¹⁰ copies/mL.

25

In some embodiments, an HEV titration assay comprising the use of polybrene is provided. In some embodiments of the assay, the concentration of polybrene used is in the range of about 1 μ g/mL - about 5 μ g/mL.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A - Figure 1C show images taken at 20x magnification of HepG2/C3A cells cultured for 1 day with different cell culture media according to some embodiments 5 of the disclosure herein.

Figure 1A shows HepG2/C3A cells cultured with DMEM + 10% FBS + Non-essential amino acids, fungizone, HEPES, gentamicin (NHG).

10

Figure 1B shows HepG2/C3A cells cultured with DMEM + 10% FBS + NHG + pyruvate.

Figure 1C shows HepG2/C3A cells cultured with DMEM + 10% 15 FBS + NHG + pyruvate + polybrene.

Figure 2A - Figure 2C show images taken at 10x magnification (left) or 20x magnification (right) of HepG2/C3A cells cultured for 3 days .

20

Figure 2A shows HepG2/C3A cells cultured with DMEM + 10% FBS + NHG.

Figure 2B shows HepG2/C3A cells cultured with DMEM + 10% 25 FBS + NHG + pyruvate.

Figure 2C shows HepG2/C3A cells cultured with DMEM + 10% FBS + NHG + pyruvate + polybrene.

30 **Figure 3** shows a comparison of HEV RNA titers in chronically infected HepG2 (human) to MPK (pig) cells assessed by HEV PCR assay according to some embodiments of the disclosure herein. Polybrene was not used for MPK

cells because MPK cell attachment to plating surfaces is very efficient.

Figure 4 shows the nucleotide sequences of primers and 5 probes used for HEV PCR assay and their location on the HEV genome according to some embodiments of the disclosure herein.

Figure 5 shows the standard curve for HEV PCR assay 10 according to some embodiments of the disclosure herein.

DETAILED DESCRIPTION

The invention relates to a process for generating high-titer stocks of hepatitis E virus (HEV) suitable for use 15 in virus clearance studies and a method for determining infectious HEV titers. The present invention comprises a simple method to propagate high titer HEV in cell culture for use in studies to evaluate the HEV clearance capacity of manufacturing processes of blood- and plasma-derived 20 products.

As previously discussed, HEV has been adapted to grow in cell culture by Okamoto and colleagues, using PLC/PRF/5 human hepatoma cells and A549 human lung cells, and by 25 Emerson and colleagues, using HepG2/C3A human hepatoma cells. Human hepatoma cells are difficult to grow and may require special cell plating methods. Emerson's laboratories coats plates with rat tail collagen I before seeding cells, while Okamoto's laboratories uses plates 30 purchased from IWAKI, some of which are collagen-coated. Table 1 compares HEV titers from various cell culture systems and lists the methods used by different laboratories to enhance cell plating. Collagen and other

coatings such as fibronectin, gelatin and poly-L-lysine facilitate cell attachment and/or cell growth so difficult-to-grow cells can reach higher densities and, thereby, yield high titer virus.

5

Table 1. Comparison of HEV Titers from Various Cell Culture Systems

Lab	Cell plating enhancement method	Cells	HEV strain (genotype)	HEV Pass #	Log ₁₀ RNA copies/mL	Reference
Okamoto	IWAKI plates	PLC/PR F/5 & A549	JE03-1760F (3) JE-JF5/15F (4)	45 25	8 - 9 9 - 10	Tanaka, et al. (2007) J Gen Virol 88: 903-911
Emerson	Rat Tail Collagen I coated plates	HepG2/ C3A	Kernow-C1 Passage 6 (3)	6	8.7	Patent Application US 2013 0302790 A1
Grifols	Polybrene in media	HepG2/ C3A	Kernow-C1 Passage 6 (3)	12 13 14 34 35 36	8.2 8.5 8.9 8.9 8.9 8.9	Experiment VM1893

In some embodiments of the present invention, instead of 10 using a coating such as collagen or other coatings to facilitate cell attachment and/or cell growth, a culture method for HEV involving the use of polybrene (also known as hexadimethrine bromide and 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide) is provided.

Polybrene is an inexpensive cationic polymer typically used in cell culture to increase the infection efficiency of retroviruses by decreasing the charge repulsion between 5 virus and cells. The capacity of polybrene to bring different objects closer together is used in our invention to facilitate the attachment of human hepatoma cells to plating surfaces and HEV to cells. Polybrene is simply added to the cell culture media that is used for cell and 10 virus propagation. The present invention obviates the need to purchase expensive plates pre-coated with collagen or other coating agents or to pre-coat plates with collagen or other coating agents before cell seeding and virus infection.

15

HEV usually replicates to low titers *in vivo* so growth *in vitro* is difficult. Methods for culturing HEV to yield high-titer stocks suitable for use in virus clearance studies were not previously available.

20

In one aspect of the invention disclosed herein, propagation of high titer HEV in cell culture is possible by using media supplemented with polybrene.

25 Polybrene is an inexpensive cationic polymer used in cell culture to increase the infection efficiency of retroviruses (enveloped viruses) by decreasing the charge repulsion between virus and cells. In some embodiments of the invention, polybrene is added to cell culture media to 30 facilitate the attachment of HEV, a non-enveloped/pseudo-enveloped virus to cells. In addition, the presence of polybrene improves cell attachment and proliferation,

thereby, enhancing the overall production of HEV in infected cells.

Therefore, in a first embodiment the present invention 5 refers to a method of producing high-titer HEV in *in vitro* cultures based on the addition of polybrene to the cell culture medium.

HEV does not produce cytopathic effects (CPE) in cell 10 culture so HEV detection is based on PCR. Thus, in some embodiments, detection of HEV at low concentrations is possible by a PCR assay of the present invention. The PCR assay can be used to develop, for example, an infectivity assay for HEV that is precise, accurate and reliable.

15

The PCR assay originally developed by the National Institutes of Health (NIH) was modified to increase sensitivity. The reverse primer was altered and a completely new probe was designed. Methods using the PCR 20 assay of the present invention consistently scored samples correctly as positive or negative, and are useful in titer calculations. Thus, in a further embodiment, the present invention discloses PCR-based HEV titration assays that comprise the use of polybrene or a culture medium 25 (preferably, cell culture medium) comprising said polybrene.

In one embodiment, the invention relates to a process that involves: propagation of high titer HEV in cell culture by 30 using media supplemented with polybrene; and the detection of HEV by a sensitive PCR assay.

The process can be specific to HEV but the methods for virus propagation can be applied to other non-enveloped viruses. For example, in some embodiments, the present invention was tested with HEV genotype 3 strain Kernow-C1

5 Passage 6 obtained from the Emerson lab.

It is contemplated that, in some embodiments, the method of producing high-titer HEV mentioned above is carried out in *in vitro* cultures, preferably in *in vitro* organ, tissue 10 or cell cultures. In the most preferable embodiment, the method of producing high-titer HEV of the present invention is carried out in an *in vitro* cell culture.

Both primary cell culture lines and established cell 15 culture lines can be used in the method of the present invention. Cell culture lines used can derive from any organism. For example, the use of insect cells or mammalian cells is contemplated. Preferably, cell culture lines used are derived from pigs (or minipigs) and humans.

20

Moreover, preferably, cell lines used are derived from liver or kidney. Said liver or kidney from which cell lines derive can be healthy, or diseased, or comprise a malignant or benign growth. In the most preferable 25 embodiment, established cell lines HepG2 (ATCC number HB-8065), or HepG2/C3A (ATCC number CRL-10741) are used.

In some embodiments, the concentration of polybrene is in the range of about 1 μ g/mL - about 5 μ g/mL. In some 30 embodiments, the concentration of polybrene is about 1, 2, 3, 4 or 5 μ g/mL. In a preferred embodiment, the concentration of polybrene is about 4 μ g/mL.

The method of the present invention, in some embodiments, yielded HEV titers in the range of about 10^8 copies/mL - about 10^{10} copies/mL. The method of the present invention, 5 in some embodiments, yielded HEV titers in the range of about 10^7 copies/mL - about 10^{10} copies/mL. In a preferred embodiment, the method of the present invention yielded HEV titers of about 10^9 copies/mL.

10 In a further embodiment, the present invention refers to a method of producing high-titer HEV comprising the steps of: a) plating cells in culture medium containing polybrene and b) infecting the *in vitro* culture with HEV. In some embodiments, the cell culture medium from the *in* 15 *vitro* cultures is collected to obtain high-titer HEV

After step b) above, cultured cells can be collected by any means known in the state of the art (preferably, by trypsinization) and an aliquot may be assayed for the 20 content of HEV. Said content of HEV is preferably determined by PCR, even more preferably with the PCR assay of the present invention and that is described below. See Figure 5. In alternative embodiments, the content of HEV can be determined by an assay based on immunofluorescence 25 (IF). See Figure 5.

Moreover, after said step b) cultured cells can be obtained by any means known in the state of the art (preferably, by trypsinization) and HEV stocks can be 30 obtained by any method known in the state of the art. In a preferred embodiment, HEV stock is generated by freeze-thawing cells a number of times, preferably 1 or 2 times.

Generated HEV stocks are preferably stored at -65°C or colder.

It is contemplated that in some embodiments, the method 5 described above also comprises the steps of: c) adding polybrene to the cell culture medium; and d) further passaging the cells infected with HEV.

Steps c) and d) can be repeated, for example, to increase 10 the number of cells producing HEV and to maximize virus spread and infection in the culture. A person skilled in the art can easily determine the number of times that steps c) and d) can be repeated, this is the number of 15 passages that the infected cell line can tolerate on the basis of, for example, the appearance of the cells and their growth curves.

After step d) of each passage, cultured cells can be obtained by any means known in the state of the art 20 (preferably, by trypsinization) and an aliquot may be assayed for the content of HEV. Said content of HEV is preferably determined by PCR, even more preferably with the PCR assay of the present invention and that is described below. See Figure 5.

25

Moreover, after said step d) of each passage cultured cells can be obtained by any means known in the state of the art (preferably, by trypsinization) and HEV stocks can be obtained by any method known in the state of the art. 30 In a preferred embodiment, HEV stock is generated by freeze-thawing cells a number of times, preferably 1 or 2 times. Generated HEV stocks are preferably stored at -65°C or colder.

Alternatively, medium can also be collected only after the desired or required number of passages have been performed with the infected cell line (this is, after steps c) and 5 d) have been repeated the desired or required number of times).

Step d) of passaging the cells infected with HEV, as a person skilled in the art knows, implies obtaining the 10 cells by any means known in the state of the art. In case cells grow attached to a surface (for example, from a flask or a plate), the cells may need to be detached by any means known in the state of the art, preferably by trypsinization. When enzymes (for example, trypsin) are 15 used to detach cells normally said enzymes need to be inactivated. Usually, in the state of the art, said inactivation is performed by dilution with a solution rich in proteins, preferably with cell culture medium with FBS.

20 Given the fact that polybrene and culture media comprising said reagent (preferably, cell culture media) allow efficient infection and production of HEV in *in vitro* cultures (preferably, *in vitro* cell cultures), they can also be used in HEV titration assays.

25

Therefore, in a further embodiment, the present invention discloses HEV titration assays are characterized in that they comprise the use of polybrene or a culture medium (preferably, cell culture medium) comprising said 30 polybrene.

The *in vitro* culture is preferably an *in vitro* organ, tissue or cell culture. In the most preferable embodiment, the *in vitro* culture is an *in vitro* cell culture.

5 Both primary cell culture lines and established cell culture lines can be used. Cell culture lines used can derive from any organism. For example, the use of insect cells or mammalian cells is contemplated. Preferably, cell culture lines used are derived from pigs (or minipigs) and
10 humans.

Moreover, preferably, cell lines used are derived from liver or kidney. Said liver or kidney from which cell lines derive can be healthy, or diseased, or comprise a
15 malignant or benign growth. In the most preferable embodiment, established cell lines HepG2 (ATCC number HB-8065 or HepG2/C3A (ATCC number CRL-10741) are used.

Preferably, the HEV titration assay of the present
20 invention is a 50% Tissue Culture Infective Dose (TCID₅₀) assay that is performed as a person skilled in the art knows, except the presence of virus is indicated by a positive PCR signal and not by viral cytopathology.

25 In said titration assays, polybrene or the cell culture medium comprising said polybrene are used for the culture of the cells in which HEV samples will be titered.

30 In another embodiment, a method for the detection of HEV in the HEV stocks is provided.

Said detection can be performed by any method or means known in the state of the art. Nevertheless, in a preferred embodiment, detection is performed by PCR.

5 In a preferred embodiment, PCR used for the detection of HEV is real time reverse transcriptase PCR that comprises a first step of reverse transcription of viral RNA to complementary DNA (cDNA), and a second step of real time PCR that is preferably performed using a double quenched
10 probe.

In the most preferred embodiment, in the second step, the following oligonucleotides are used as primers and probe (Figure 4):

15

- a) forward primer: 5'-CGGCTATCGGCCAGAAGTT-3' (SEQ ID NO: 1)
- b) reverse primer: 5'-CCGTGGCTATAACTGTGGTCT-3' (SEQ ID NO: 2)
- 20 c) probe: 5'-FAM-TTTTACGC-ZEN-AGGCTGCCAAGGCC-3IABkFQ-3' (SEQ ID NO: 3)

Data from studies to assess the performance of the HEV PCR-based infectivity assay is summarized in Table 2.

25

Table 2. Assessment of HEV PCR-based Infectivity Assay Performance

Assay Parameter	Method Description	Acceptance Criteria	Results
Precision - Repeatability	Multiple measurements of single sample by one analyst in one day	CV \leq 30%	High: Med: Low: Pass
Precision - Intermediate	Multiple measurements of single sample by multiple analysts over several days	p-value $>$ 0.01 or difference in mean titers $<$ 0.5 \log_{10}	High: P 0.4 Med: Diff mean titer p 0.4 Low: P 0.6 Pass
Accuracy	Measures closeness of results to expected titers	CV \leq 30% Nominal 50-150%	CV: 2-17% Nom: 105-136% Pass
Linearity	Measures relationship of assay results to actual virus concentration	R ² for regression line $>$ 0.95	R ² = 0.99 Pass
Limit of Quantitation	Lowest conc. of virus detected with acceptable precision, & accuracy	Lowest concentration with CV \leq 30%	10 ^{0.8} TCID ₅₀ /mL Pass

Assay Parameter	Method Description	Acceptance Criteria	Results
Range	Interval where virus is detected with acceptable precision, accuracy, & linearity	CV \leq 30%	$10^{0.8}$ to $10^{6.4}$ TCID ₅₀ /mL Pass
Detection Limit - Standard Titration	Lowest concentration of virus that can be detected	Lowest concentration with CV \leq 50%	$10^{0.8}$ TCID ₅₀ /mL Pass
Detection Limit - Large Volume Titration	Titrated 12.6 mL of $10^{6.4}$ TCID ₅₀ /mL stock after $10^{6.6}$ -fold dilution	Expected value: 0-2 positive/252 wells	2 positive/252 wells = $10^{-0.5}$ TCID ₅₀ /mL NA

In a preferred embodiment cells are titrated in 96 well-plate and HEV RNA extracted with magnetic Biocclone beads or magnetic Dynabeads, more preferably Dynabeads are used.

5

For a better understanding, certain embodiments of the present invention are described in more detail with reference to the accompanying figures, which are presented by way of example, and with reference to illustrative 10 examples which are not a limitation of the present invention.

EXAMPLES

Example 1. Cell growth patterns of HepG2/C3A cells in 15 different Growth Media

HepG2/C3A (10^6) cells were seeded in 25 cm² flasks and cultured in one of the following cell culture media:

- DMEM + 10% FBS + NHG (Non-essential amino acids, fungizone, HEPES, gentamicin);
- DMEM + 10% FBS + NHG + 1 mM pyruvate; or
- 5 - DMEM + 10% FBS + NHG + 1 mM pyruvate + 4 ug/mL polybrene.

Images, taken at 20x magnification after 1 day, are shown in Figure 1. HepG2/C3A cells cultured in medium DMEM + 10%
10 FBS + NHG (A) or DMEM + 10% FBS + NHG + pyruvate (B) grew in clumps, which would make infection with virus (e.g. HEV) very difficult. On the other hand, when HepG2/C3A cells were cultured in DMEM + 10% FBS + NHG + pyruvate + polybrene (C), cells attached and grew in a flat even
15 monolayer, which should be easily infected by a virus (e.g. HEV).

Images, taken at 10x and 20x magnification after 3 days, are shown in Figure 2. Similar to the 1 day cells,
20 HepG2/C3A cells cultured in DMEM + 10% FBS + NHG (A) or DMEM + 10% FBS + NHG + pyruvate (B) grew in clumps but cells grown in DMEM + 10% FBS + NHG + pyruvate + polybrene (C) grew in a flat even monolayer. Since cell clumps would impede uniform infection by viruses, the efficiency of HEV
25 infection would most likely be higher in cell grown in the presence of polybrene.

Example 2. Establishment of HEV-infected cells

Cell culture flasks were seeded for infection as follows:
30 HepG2 or HepG2/C3A cells were trypsinized according to protocols or procedures known in the state of the art. Ten mL of Growth Medium (base medium in accordance with the requirements of the cell line used plus polybrene at a

concentration of about 1 - about 5 μ g/mL) were added to neutralize the trypsin and the suspension was pipetted up and down to break up cell clumps. The cells were then seeded at a density of approximately 10^6 cells per 150 cm^2 flask and left overnight in a 37°C incubator.

Cells were infected with HEV by removing the medium from the flask and adding HEV Stock (clarified virus-infected cell lysate) at a Multiplicity of Infection (MOI) of 0.1 - 10 1.0 and in a total volume of 5 - 10 mL. The cells were incubated at 37°C for at least 1 hour, during which time, the flask was periodically rocked back and forth to prevent cells from drying out and to distribute the virus inoculum evenly across cells. Additional Growth Medium (10 15 - 20 mL per flask) was added and the flask was kept at 37°C until the cell monolayer was confluent (approximately 1 week).

Example 3. Propagation of HEV-infected cells

20 Infected cells (HepG2 or HepG2/C3A) in a 150 cm^2 flask were trypsinized in accordance with Example 1 when the cell monolayer became confluent. If multiple flasks were trypsinized, cell suspensions were pooled together before proceeding.

25

An aliquot, no less than 1 mL, of the trypsinized cell suspension was taken to analyze for HEV RNA by PCR. The remaining cell suspension was discarded or divided into other 150 cm^2 flasks at a density equivalent to the number 30 of cells in 1/6th of a confluent monolayer (1:6 split). Growth Medium was added to bring the final volume in each 150 cm^2 flask to 20 - 25 mL and the flasks were incubated until the cells reached confluency (approximately 1 week).

This procedure was repeated each week until the quantity of HEV RNA in the 1 mL sample, that had been removed for PCR analysis, reached high titers. In some embodiments, 5 high titer was in the range of about 10^8 copies/mL - about 10^{10} copies/mL. In some embodiments, high titer was in the range of about 10^7 copies/mL - about 10^{10} copies/mL. Some flasks were then trypsinized and split for continued passage of the HEV-infected cells while the remaining 10 flasks were processed as HEV stock.

Flasks of HEV stock were frozen and thawed 1 - 2 times to rupture infected cells and release virus. The infected lysates were then pooled, aliquoted into appropriate 15 containers and stored at no warmer than -65°C.

Example 4. HEV PCR-based infectivity assay

HEV titration in a 96-well plate format is described in this example but the assay could be easily adapted to 20 multiwell plates of other sizes.

Serial dilutions of a HEV stock were made and added to wells seeded with HepG2 or HepG2/C3A cells. Virus was allowed to adsorb for no less than 1 hr at 37°C and Growth 25 Media was added. Plates were incubated at 37°C for no less than 2 days before aspirating the media from the wells and washing/aspirating the cells no less than 2x with buffer (e.g. PBS). The plates were then extracted for PCR or were stored at no warmer than -65°C until ready for extraction.

30

The Dynabeads® mRNA DIRECT™ Micro Kit (Life Technologies) was used to extract polyadenylated RNA (e.g. HEV RNA) from cells in each well of the titration plate, following

manufacturer instructions. The resulting eluate (poly A RNA) was immediately processed for PCR amplification or stored at no warmer than -65°C until ready for PCR amplification.

5

One step RT-PCR was used to detect HEV RNA in samples, using the primers and probes as previously discussed. The assay conditions for each reaction were as follows:

10 a) Reagents; 5.0 μ l 4x TaqMan® Fast Virus 1-Step Master Mix (Life Technologies), 0.08 μ L 100 mM primer F+R, 0.04 μ L 100 mM probe, 0.4 μ L SUPERase In (Life Technologies), 4.4 μ L water and 10 μ L template (total 20 μ l)

15 b) Reaction: 52°C 10 minutes, 95°C 30 seconds, and 40 cycles of 95°C 15 seconds, 56°C 45 seconds,

PCR and cycle threshold (Ct) value determination were performed using an AB 7500 Real Time PCR System (Applied 20 Biosystems, Foster City, CA) and accompanying software according to manufacturer's instruction.

PCR was quantitative or qualitative. For quantitative PCR, a HEV cDNA plasmid obtained from the NIH was linearized 25 with MluI and transcribed using the mMESSAGE mMACHINE® Kit (Life Technologies) to generate 7.2 kb RNA transcripts with a 7-methyl guanosine cap and poly A tail. The transcripts were purified with the Ambion MegaClear kit (Life Technologies), quantified with the Quant-iT™ 30 RiboGreen® RNA Reagent and Kit (Invitrogen) and used as standards to construct RNA standard calibration curves.

RNA standard curves were generated by the AB7500 software system by plotting the Ct values against the logarithm of the calculated copy numbers for the standards.

5 Figure 5 (right panel) shows a typical standard curve. All curves had a wide dynamic range, ranging from 10^0 to 10^7 copies per reaction, and were linear with a correlation coefficient of $r^2 > 0.99$. The percent efficiency of amplifications was calculated as $\% E = [10^{(-1/\text{slope})-1}] * 100$. Based on $n = 22$ HEV qPCR standard curves, the efficiency of HEV quantitative PCR was 100.4% (data not shown).

15 For qualitative PCR, wells were scored positive or negative based on the presence of a positive PCR signal. Virus titers were calculated as $\text{TCID}_{50}/\text{mL}$ using the appropriate statistical methods: Spearman-Kärber, MPN or Poisson.

20 Example 5. Assessment of HEV PCR-based Infectivity Assay Performance

Assay qualification studies were performed to assess the operational characteristics of the HEV infectivity assay. The parameters evaluated were precision, accuracy, 25 linearity, limit of quantitation and detection, and dynamic range.

Acceptance criteria were the same as those typically used for other virus titration assays.

30

The results are summarized in Table 2 and show that the assay passed all tests.

DEFINITIONS

HepG2: Hepatocellular carcinoma cells obtained from the American Type Culture Collection (ATCC number HB-8065)

5 MPK: Minipig kidney cells obtained from the American Type Culture Collection (ATCC number CCL-66)

DMEM: Dulbecco's Modified Eagle Medium

FBS: Fetal Bovine Serum

10 HEPES: N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid

NEAA: Nonessential Amino Acids

NHG: A mixture of the following: Nonessential Amino Acids, Fungizone, HEPES, and Gentamicin

15 Titration: The process of serially diluting a sample to bracket an expected viral titer and transferring the diluted sample to a plate to determine the TCID₅₀/mL

20 Titer: Concentration of a substance (virus) in solution or the strength of such a substance determined by titration

25 SK: Spearman-Karber is a statistical method to calculate virus titers in samples with relatively high concentrations of virus. This method is used when the proportion of positive wells at any dilution is >25%

30 MPN: Most Probable Number is a statistical method to calculate virus titers in samples with relatively low concentrations of virus. MPN is used when the proportion of positives wells at all dilutions is <25%.

Poisson: Poisson is a statistical method to calculate virus titers in samples with extremely low

concentrations of virus. This method is used when no positive wells are observed.

ATCC: American Type Culture Collection

5 TCID₅₀: Corresponds to 50% Tissue Culture Infective Dose (Endpoint dilution assay). It is a measurement of infectious virus titer that quantifies the amount of virus required to kill 50% of infected hosts or to produce cytopathic effect in 50% of inoculated tissue culture cells.

10

In some aspects, embodiments of the present invention as described herein include the following items:

15 1. A method of producing a high titer of Hepatitis E Virus (HEV), the method comprising:

20 a) culturing a cell line *in vitro*, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741, in a medium comprising polybreneTM also known as 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide in a concentration of about 1 μ g/mL - about 5 μ g/mL,
25 b) infecting the cell line with HEV, and
c) culturing the infected cells for approximately one week,

wherein the high titer of HEV is about 10⁷ RNA copies/mL - about 10¹⁰ RNA copies/mL.

30 2. The method according to item 1, wherein the high titer of HEV is about 10⁸ RNA copies/mL - about 10¹⁰ RNA copies/mL.

3. The method according to item 1 or 2, wherein the method further comprises the steps of passaging the HEV-infected cell line of step b) in a culture medium comprising polybreneTM in a concentration 5 of about 1 μ g/mL - about 5 μ g/mL and culturing the infected cells for approximately one week.
4. The method according to item 3, wherein the high titer of HEV obtained in the medium and/or infected cells is 10 about 10⁸ RNA copies/mL - about 10¹⁰ RNA copies/mL.
5. The method according to item 3, wherein the high titer of HEV obtained in the medium and/or infected cells is about 10⁷ RNA copies/mL - about 10¹⁰ RNA copies/mL.
- 15 6. A method of determining a presence and/or a concentration of Hepatitis E Virus (HEV) in a sample, the method comprising the steps of:
 - 20 a) providing the sample to a mixture comprising a cell line, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741, and a culture medium, wherein the culture medium comprises polybreneTM also known as 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide in a concentration of about 1 μ g/mL - about 5 μ g/mL;
 - 25 b) incubating the mixture comprising the sample from step a) for approximately one week, to allow for the propagation of HEV, if present in the sample;
 - 30 c) collecting a portion of step b), the portion comprising HEV, if present and propagated during step b); and

d) measuring the presence and/or the concentration of a HEV polynucleotide or polypeptide in the collected portion.

5 7. The method according to item 6, wherein measuring the presence and/or the concentration of the HEV polynucleotide or polypeptide comprises the steps of:

10 e) providing a first reaction mixture by mixing the collected portion with a first solution so as to expose a polynucleotide of HEV, if HEV is present in the collected portion, wherein the polynucleotide is an RNA of HEV;

15 f) providing a second reaction mixture by adding, to the first reaction mixture, a first reagent so as to produce a complementary deoxyribonucleic acid (cDNA) that is at least partially complementary to the RNA of HEV;

20 g) providing a third reaction mixture by adding, to the second reaction mixture, a second reagent comprising a pair of polynucleotides to amplify a sequence of the cDNA that is at least partially complementary to each of the polynucleotides in the pair of polynucleotides;

25 h) providing a fourth reaction mixture by amplifying the sequence; and

25 i) determining a concentration of the amplified sequence in the fourth reaction mixture.

30 8. The method according to item 7, wherein said determining the concentration comprises:

j) providing the fourth reaction mixture;

30 k) providing one or more controls comprising a predetermined amount of the cDNA of HEV;

- 1) adding an oligonucleotide sequence to the fourth reaction mixture and the one or more controls, wherein said oligonucleotide sequence is at least partially specific to the amplified sequence in the fourth reaction mixture and the predetermined amount of the cDNA of HEV in the one or more controls; and
- 5 m) comparing the concentration of the amplified sequence in the fourth reaction mixture relative to the predetermined amount of the cDNA of HEV in the one or more controls.
- 10

9. The method according to item 7, wherein said pair of polynucleotides comprises:

15

- i. 5'-CGGCTATCGGCCAGAAGTT-3' (SEQ ID NO: 1) and
- ii. 5'-CCGTGGCTATAACTGTGGTCT-3' (SEQ ID NO: 2).

20 10. The method according to item 8 or 9, wherein the oligonucleotide sequence comprises:

5'- TTTTACGCAGGCTGCCAAGGCC-3' (SEQ ID NO: 3).

25 11. The method according to item 10, wherein probes for quantification of the amplification (qPCR) are attached to the oligonucleotide sequence.

30 12. The method according to item 11, wherein the probes for quantification of the amplification (qPCR) are 5(6)-carboxyfluorescein dye (5(6)-FAMTM), ZENTM double quencher and/or IOWA BLACK-FQTM quencher.

13. The method according to item 12, wherein FAM™, ZEN™ and IOWA BLACK-FQ™ are linked to the oligonucleotide sequence comprising SEQ ID NO:3 at positions 1, 9 and 23 respectively.

5

14. A culture medium for generating a high-titer Hepatitis E Virus (HEV), the culture medium comprising polybrene™ also known as 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide in a range of about 1 µg/mL - about 5 µg/mL, wherein the high-titer HEV is about 10⁷ RNA copies/mL - about 10¹⁰ RNA copies/mL.

15. The culture medium according to item 14, wherein the high-titer HEV is about 10⁸ RNA copies/mL - about 10¹⁰ RNA copies/mL.

20 16. An Hepatitis E Virus (HEV) titration assay comprising the method of determining a presence and/or a concentration of HEV in a sample as defined in any one of items 6 to 13.

REFERENCES

1. Okamoto H (2011) Hepatitis E virus cell culture models. Virus Research 161: 65- 77.
2. Tanaka T, et al (2007) Development and evaluation of an efficient cell-culture system for Hepatitis E virus. J Gen Virol 88: 903-911.
3. Tanaka T, et al (2009) Development and Characterization of a Genotype 4 Hepatitis E Virus Cell Culture System Using a HE-JF5/15F Strain Recovered from a Fulminant Hepatitis Patient. J Clin Microbiol 47: 1906-1910.

4. Shukla P, et al (2011) Cross-species infections of cultured cells by hepatitis E virus and discovery of an infectious virus-host recombinant. PNAS. 108 (6): 2438-2443.
5. Shukla P, et al (2012) Adaptation of a Genotype 3 Hepatitis E Virus to Efficient Growth in Cell Culture Depends on an Inserted Human Gene Segment Acquired by Recombination. J Virol 86 (10): 5697-5707
6. US Provisional Patent Application No 61/431,377
- 10 7. US Provisional Patent Application No 61/554,323
8. US Patent Application No. 13/978,839
9. International PCT Application No. PCT/US2012/020830

CLAIMS

1. A method of producing a high titer of Hepatitis E Virus (HEV), the method comprising:

5

- a) culturing a cell line *in vitro*, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741, in a medium comprising polybreneTM also known as 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide in a concentration of about 1 μ g/mL - about 5 μ g/mL,
- b) infecting the cell line with HEV, and
- c) culturing the infected cells for approximately one week,

10

wherein the high titer of HEV is about 10^7 RNA copies/mL - about 10^{10} RNA copies/mL.

15

2. The method according to claim 1, wherein the high titer of HEV is about 10^3 RNA copies/mL - about 10^{10} RNA copies/mL.

3. The method according to claim 1 or 2, wherein the method further comprises the steps of
25 passaging the HEV-infected cell line of step b) in a culture medium comprising polybreneTM in a concentration of about 1 μ g/mL - about 5 μ g/mL and culturing the infected cells for approximately one week.

30 4. The method according to claim 3, wherein the high titer of HEV obtained in the medium and/or infected cells is about 10^3 RNA copies/mL - about 10^{10} RNA copies/mL.

5. The method according to claim 3, wherein the high titer of HEV obtained in the medium and/or infected cells is about 10^7 RNA copies/mL – about 10^{10} RNA copies/mL.

5 6. A method of determining a presence and/or a concentration of Hepatitis E Virus (HEV) in a sample, the method comprising the steps of:

- 10 a) providing the sample to a mixture comprising a cell line, wherein the cell line is HepG2 deposited under ATCC number HB-8065 or HepG2/C3A deposited under ATCC number CRL-10741, and a culture medium, wherein the culture medium comprises polybreneTM also known as 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide in a concentration of about 1 μ g/mL – about 5 μ g/mL;
- 15 b) incubating the mixture comprising the sample from step a) for approximately one week, to allow for the propagation of HEV, if present in the sample;
- 20 c) collecting a portion of step b), the portion comprising HEV, if present and propagated during step b); and
- 25 d) measuring the presence and/or the concentration of a HEV polynucleotide or polypeptide in the collected portion.

7. The method according to claim 6, wherein measuring the presence and/or the concentration of the HEV polynucleotide or polypeptide comprises the steps of:

- e) providing a first reaction mixture by mixing the collected portion with a first solution so as to expose a polynucleotide of HEV, if HEV is present in the collected portion, wherein the polynucleotide is an RNA of HEV;
- f) providing a second reaction mixture by adding, to the first reaction mixture, a first reagent so as to produce a complementary deoxyribonucleic acid (cDNA) that is at least partially complementary to the RNA of HEV;
- g) providing a third reaction mixture by adding, to the second reaction mixture, a second reagent comprising a pair of polynucleotides to amplify a sequence of the cDNA that is at least partially complementary to each of the polynucleotides in the pair of polynucleotides;
- h) providing a fourth reaction mixture by amplifying the sequence; and
- i) determining a concentration of the amplified sequence in the fourth reaction mixture.

8. The method according to claim 7, wherein said determining the concentration comprises:

predetermined amount of the cDNA of HEV in the one or more controls; and

5 m) comparing the concentration of the amplified sequence in the fourth reaction mixture relative to the predetermined amount of the cDNA of HEV in the one or more controls.

9. The method according to claim 7, wherein said pair of polynucleotides comprises:

10

- i. 5'-CGGCTATCGGCCAGAAGTT-3' (SEQ ID NO: 1) and
- ii. 5'-CCGTGGCTATAACTGTGGTCT-3' (SEQ ID NO: 2) .

10. The method according to claim 8 or 9, wherein the 15 oligonucleotide sequence comprises:

5' - TTTTACGCAGGCTGCCAAGGCC-3' (SEQ ID NO: 3) .

11. The method according to claim 10, wherein probes for 20 quantification of the amplification (qPCR) are attached to the oligonucleotide sequence.

12. The method according to claim 11, wherein the probes for quantification of the amplification (qPCR) are 5(6)-25 carboxyfluorescein dye (5(6)-FAMTM), ZENTM double quencher and/or IOWA BLACK-FQTM quencher.

13. The method according to claim 12, wherein FAMTM, ZENTM and IOWA BLACK-FQTM are linked to the oligonucleotide 30 sequence comprising SEQ ID NO:3 at positions 1, 9 and 23 respectively.

14. A culture medium for generating a high-titer Hepatitis E Virus (HEV), the culture medium comprising polybreneTM also known as 1,5-dimethyl-1,5-diazaundeca-methylene polymetho-bromide in a range of about 1 μ g/mL - about 5 5 μ g/mL, wherein the high-titer HEV is about 10^7 RNA copies/mL - about 10^{10} RNA copies/mL.

15. The culture medium according to claim 14, wherein the high-titer HEV is about 10^3 RNA copies/mL - about 10^{10} RNA 10 copies/mL.

16. An Hepatitis E Virus (HEV) titration assay comprising the method of determining a presence and/or a concentration of HEV in a sample as defined in any one of 15 claims 6 to 13.




Figure 1A

2/9




Figure 1B

3/9




Figure 1C

4/9




Figure 2A

5/9




Figure 2B

6/9



Figure 2C

7/9







Figure 3



5'-CGGCTATCGGCCAGAAGTT-3' (SEQ ID NO: 1)
 5'-CCGTGGCTATAACTGTGGTCT-3' (SEQ ID NO: 2)
 5'-FAM-TTTTACGC-ZEN-AGGCTGCCAAGGCC-3' (SEQ ID NO: 3)

Figure 4



- Based on historical data, the threshold is manually set at 0.1 so it passes through the exponential phase of all standard curves and to ensure detection of 1 RNA copy per rxn

- Slope = -3.3
- $R^2 = 0.997$

Figure 5

