




(86) Date de dépôt PCT/PCT Filing Date: 1999/10/22
 (87) Date publication PCT/PCT Publication Date: 2000/05/04
 (45) Date de délivrance/Issue Date: 2009/12/08
 (85) Entrée phase nationale/National Entry: 2001/04/20
 (86) N° demande PCT/PCT Application No.: GB 1999/003430
 (87) N° publication PCT/PCT Publication No.: 2000/024253
 (30) Priorité/Priority: 1998/10/22 (GB9823010.5)

(51) Cl.Int./Int.Cl. *A01N 37/36* (2006.01),
A01N 35/06 (2006.01), *A01N 37/38* (2006.01),
A01N 43/24 (2006.01), *A01N 43/50* (2006.01),
A01N 43/54 (2006.01), *A01N 43/56* (2006.01),
A01N 43/653 (2006.01), *A01N 43/68* (2006.01),
A01N 43/713 (2006.01), *A01N 47/22* (2006.01),
A01N 47/36 (2006.01), *A01N 53/08* (2006.01)

(72) Inventeurs/Inventors:
 BARDSLEY, RICHARD ANDREW, GB;
 BICKERS, UDO MATTHIAS, DE;
 BRIGGS, GEOFFREY GOWER, GB;
 GREEN, SHIRLEY ANN, GB;

(73) Propriétaire/Owner:

(54) Titre : DERIVES D'ACIDE CITRIQUE
 (54) Title: CITRIC ACID DERIVATIVES



(57) Abrégé/Abstract:

The invention provides the use as a pesticidal adjuvant of at least one citric acid derivative and compositions containing the derivative, which has a log octanol-water coefficient (log P) of 2.6 to 11 and an equivalent hydrocarbon (EH) value of 29 to 47. The invention has been shown to enhance the efficacy of a range of pesticides.

(72) Inventeurs(suite)/Inventors(continued): PATE, ADRIENNE ELIZABETH, GB; SANWALD, ERICH FRIEDRICH, DE;
STOCK, DAVID, GB

(73) Propriétaires(suite)/Owners(continued):BAYER CROPSCIENCE SA, FR

(74) Agent: MACRAE & CO.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : A01N 37/36 // (A01N 37/36, 53:08, 47:36, 47:22, 43:713, 43:68, 43:653, 43:56, 43:54, 43:50, 43:24, 37:38, 35:06)</p>	A1	<p>(11) International Publication Number: WO 00/24253 (43) International Publication Date: 4 May 2000 (04.05.00)</p>
<p>(21) International Application Number: PCT/GB99/03430 (22) International Filing Date: 22 October 1999 (22.10.99) (30) Priority Data: 9823010.5 22 October 1998 (22.10.98) GB (71) Applicant (for all designated States except US): HOECHST SCHERING AGREVO GMBH [DE/DE]; Miraustrasse 54, D-13509 Berlin (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): BARDSLEY, Richard, Andrew [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). BICKERS, Udo, Matthias [DE/DE]; D-65926 Frankfurt am Main (DE). BRIGGS, Geoffrey, Gower [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). GREEN, Shirley, Ann [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). PATE, Adrienne, Elizabeth [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). SANWALD, Erich, Friedrich [DE/DE]; D-65926 Frankfurt am Main (DE). STOCK, David [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB).</p>	<p>(74) Agent: SEWELL, Richard, Charles; AgrEvo UK Limited, Patent Dept., Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). (81) Designated States: AU, BR, CA, CN, CZ, HU, IL, IN, JP, KP, KR, MX, PL, RU, TR, UA, US, ZA, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: USE OF CITRIC ACID DERIVATIVES AS PESTICIDAL ADJUVANTS</p>		
<p>(57) Abstract</p> <p>The invention provides the use as a pesticidal adjuvant of at least one citric acid derivative and compositions containing the derivative, which has a log octanol-water coefficient (log P) of 2.6 to 11 and an equivalent hydrocarbon (EH) value of 29 to 47. The invention has been shown to enhance the efficacy of a range of pesticides.</p>		

Citric Acid Derivatives

Field of the invention

This invention relates to the novel use of citric acid derivatives in association with pesticides and also pesticidal compositions containing citric acid derivatives. In particular, the invention relates to compositions having fungicidal, herbicidal, insecticidal and acaricidal activity.

Background to the invention

Pesticidal compounds are typically used in the form of compositions containing one or more co-formulants, for example surfactants. For example, WO96/22020 discloses the use in a pesticidal composition of at least one aliphatic mono-, di- or tri-ester, with no mention of citric acid derivatives. WO90/13222 discloses plant-protecting preparations comprising certain citric acid derivatives that are different from those of the present invention. EP 0 579 052 deals with plant-protecting compositions comprising a biocide and an accelerator, for example linear diacid and esters thereof. GB 2 002 635 and DE 27 38 878 disclose pyrethroid-containing insecticidal compositions comprising citric acid esters to reduce the volatility of the active ingredient. JP 52 141853 provides plastic films containing trialkyl acetylcitrate with good resistance to fungus growth. We have found a new group of compounds, not previously used in association with pesticidal compounds, that can be used with advantage in association with pesticidal compounds.

Description

In a first aspect, the invention provides the use as a pesticidal adjuvant of at least one citric acid derivative, which has a log octanol-water coefficient (log P) of 2.6 to 11 (preferably 2.7 to 11, particularly 4 to 10.7, especially 4 to 10.3) and an equivalent hydrocarbon (EH) value of 29 to 47 (preferably 32 to 44).

We have found that the use of citric acid derivatives according to the invention which have little or no pesticidal activity in their own right, surprisingly aid the penetration of the pesticidal compound into the plant or plant tissue, thereby enhancing the efficacy of the pesticide in the pesticidal composition. This enhanced efficacy enables lower application rates. Lower application rates coupled

31-10-2000

GB 009903430

1a

with low toxicity of citric acid derivatives, leads to a reduction in the environmental impact.

In the context of the invention, the term adjuvant is a compound which enhances
5 the bioactivity of a pesticide, having no bioactivity in its own right.

In accordance with its normal meaning, the term pesticide, pesticidal etc, includes plant growth regulators (PGRs).

10 Advantageously, we have found that citric acid derivatives used as part of the invention as well as improving penetration of the pesticidal compound into the

plant, are also effective solvents or co-solvents for use in pesticidal compositions. Citric acid derivatives are particularly effective in conjunction with those pesticides that are conventionally formulated as emulsifiable concentrates or emulsions in water, i.e. low to intermediate lipophilic pesticides.

5


Further advantages of citric acid derivatives include their low flash point and negligible odour compared with many other currently used adjuvants.

The log P value (octanol/water partition coefficient) of a compound is a measure of the solubility of the compound in water. The higher the log P the less soluble is the compound. The log P is calculated from the structure of the compound using the "clog P Program" (Pomona College Med Chem Release 3.54, January 1988, provided by Daylight Chemical Information Systems Inc., Claremont, California). The values generated by this program correlate well in general with those determined experimentally by methods well-known to those skilled in the art. Such experimental methods are difficult to apply accurately to compounds within the scope of this invention, and we therefore use the calculated values.

The equivalent hydrocarbon (EH) value of a compound is a measure of the volatility of a compound. The EH value is arrived at by adding the total number of carbon atoms in the molecule, adding five for any hydroxy group, one for any ether group, eight for any carboxyl group, and then adding three for each ester group. It has been found that EH values determined in this way correlate well with actual volatilities.

25

Preferred citric acid derivatives are those having the general formula (I)



wherein,

R¹, R² and R³, which may be the same or different, are C₁ to C₂₀ alkyl,

C₂ to C₂₀ alkenyl or C₂ to C₂₀ alkynyl, each of which may be substituted by -OR⁶, where R⁶ is hydrogen or alkyl; or hydrogen;

R⁴ is hydrogen or -C(=O)R⁵, where R⁵ is as defined for R¹; and

3

R^1 , R^2 , R^3 and R^4 are chosen such that the log P and EH value of the citric acid derivative is as defined hereinabove.

5 Any alkyl group may be straight or branched and is preferably of 1 to 10 carbon atoms, especially 1 to 7 and particularly 1 to 5 carbon atoms.

Any alkenyl or alkynyl group may be straight or branched and is preferably of 2 to 7 carbon atoms and may contain up to 3 double or triple bonds which may be conjugated, for example vinyl, allyl, butadienyl or propargyl.

10

Preferably, at least two R^1 , R^2 and R^3 groups are not hydrogen.

Preferably R^1 , R^2 and R^3 , which may be the same or different, are C_1 to C_{20} alkyl, especially C_3 to C_8 alkyl.

15

Particularly preferred compounds are tri-*n*-butyl citrate (log P = 4.3, EH = 32) and *O*-acetyl tri-*n*-butyl citrate (log P = 4.8, EH = 32).

20 The citric acid derivative of the invention may exist as a single compound, however more commonly it exists as a mixture of compounds. Often, citric acid derivatives are only commercially available as mixtures obtained by fractional distillation. In cases such as these, the log P and EH values are average values.

25 In a further aspect, the invention provides a pesticidal composition comprising at least one pesticidal compound and at least one citric acid derivative defined hereinabove.

30 In another aspect, the invention provides a method of combating pests at a locus infested or liable to be infested therewith which comprises applying to said locus an effective amount of a pesticidal composition which comprises a pesticidal compound and at least one citric acid derivative as hereinabove described.

Suitable active ingredients which can be combined with the citric acid derivatives are especially those disclosed in "The Pesticidal Manual", 11th Edition, published

by Crop Protection Publications, and may for example be a herbicide, fungicide, insecticide, acaricide or PGR.

A suitable fungicide is for example

- 5 (i) a conazole sterol Δ^{14} -demethylase inhibitor,
(ii) a sterol Δ^{14} -reductase/ $\Delta^{8,7}$ -isomerase reduction inhibitor based on a 1-[3-(4-tert-butylphenyl)-2-methylpropyl] group which is attached via the *N*-atom to piperidine or 2,6-dimethylmorpholine
(iii) a dithiocarbamate fungicide
10 (iv) a phthalimide fungicide in which a chloroalkylthio group is attached via the *N*-atom to the optionally hydrogenated phthalimide group.
(v) an anilide fungicide
(vi) an mbc fungicide.
(vii) a carbamate fungicide
15 (viii) a copper compound fungicide
(ix) a tin compound fungicide
(x) a strobilurin type fungicide,
(xi) a 2-anilinopyrimidine fungicide
(xii) a compound which causes systemic activated resistance, or
20 (xiii) a fungicide selected from the group consisting of chlorothalonil, dimethomorph, fenamidone, fencpiclonil, fluazinam, hymexazol, nuarimol, pencycuron, pyrifenox, thicyofen, probenazole, pyroquilon, tricyclazole, quaternary ammonium compounds, fludioxonil, quinoxifen, famoxadone, diclocymet, spiroxamine, flumetover,
25 fenhexamid, furametpyr, diflumetorim, fencaramid, carpropamid, sulfur and 5-chloro-6-(2-chloro-6-fluorophenyl)-7-*N*-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-*a*]pyrimidine.

30 Conazoles are defined in ISO standard 257 as compounds based on imidazole or 1,2,4-triazole and containing a halogenated phenyl group. Examples include prochloraz (and its metal complexes - especially the zinc, manganese or copper complex), propiconazole, flusilazole, hexaconazole, tebuconazole, difenoconazole, bromuconazole, cyproconazole, diniconazole, fenbuconazole, imibenconazole, furconazole, tetraconazole, myclobutanil, penconazole, fluquinconazole,

5

azaconazole, imazalil, triflumizole, epoxiconazole, triticonazole, metconazole and oxpoconazole.

Examples of type (ii) fungicides include fenpropimorph and fenpropidin.

5

Examples of type (iii) fungicides include mancozeb and thiram.

Examples of type (iv) fungicides include folpet, captafol and captan.

Examples of type (v) fungicides include

- 10 a) 3',5'-dichloroanilide fungicides in which the anilino nitrogen comprises a ring carrying two oxo substituents, in positions adjacent the nitrogen, e.g. iprodione, vinclozolin or procymidone, or
- b) acetanilide fungicides, e.g. metalaxyl or ofurace,
- c) sulfanilide fungicides, e.g. dichlofluanid,
- 15 d) benzanilide fungicides, e.g. flutolanil, and
- e) heteroarylanilide fungicides, e.g. thifluzamide.

Examples of type (vi) fungicides include carbendazim, benomyl and thiophanate-methyl.

20

Examples of type (vii) fungicides include diethofencarb and propamocarb.

Examples of type (viii) fungicides include Bordeaux mixture, oxine-copper, copper oxychloride and copper naphthenate.

25

Examples of type (ix) fungicides include tributyltin oxide and tributyltin naphthenate.

Strobilurin type fungicides (type (x) fungicides) are methyl esters or

30 N-methylamides of arylacetic acid in which the acetic acid also carries a methoxymethylene or methoxyimino substituent. The aryl group is usually a 2-substituted phenyl group and/or can be separated from the acetic acid by a linking group such as oxygen. Examples of such compounds are kresoxim-methyl, azoxystrobin, metominostrobin, trifloxystrobin and picoxystrobin.

35

Examples of type (xi) fungicides include pyrimethanil, mepanipyrim and cyprodinil.

An example of a type (xii) fungicide is that having the code number CGA 2425704, which is sold under the trade name "Bion"^{*} and whose proposed
5 common name is acibenzolar.

Examples of herbicides which can be used in conjunction with the citric acid derivatives include

- a) Acetolactate synthase inhibitors, e.g.
- 10 (i) sulfonylureas, such as chlorsulfuron, sulfometuron, metsulfuron, bensulfuron, chlorimuron, tribenuron, thifensulfuron, thiameturon, ethametsulfuron, nicosulfuron, rimsulfuron, azimsulfuron, cinosulfuron, prosulfuron, flazasulfuron, pyrazasulfuron, triasulfuron, primisulfuron, oxasulfuron, imazasulfuron, cyclosulfamuron,
- 15 amidosulfuron, ethoxysulfuron, iodosulfuron, halosulfuron, triflusulfuron, flurpysulfuron, sulfosulfuron, foramsulfuron, tritosulfuron, trifloxysulfuron and foramsulfuron,
- (ii) 4,6-dimethoxypyrimidinyloxy benzoic acid analogues, such as pyriothiobac, bispyribac, pyriminobac and pyribenzoxin;
- 20 (iii) arylsulfonanilides, such as cloransulam, diclosulam, flumetsulam, metosulam and florasulam, and
- (iv) benzenesulfonamides, such as flucarbazone and procarbazon
- b) choroacetanilides, such as alachlor, metolachlor, acetochlor and propachlor,
- c) dinitroanilines, such as trifluralin, pendimethalin and ethalfluralin,
- 25 d) HBNs, such as bromoxynil and ioxynil,
- e) benzoic acids, such as dicamba and propyzamide,
- f) phosphorus acid esters, such as glyphosate, glufosinate and bilanofos,
- g) quaternary ammonium compounds, such as paraquat, diquat and difenzoquat,
- 30 h) aryloxyalkanoic acids, such as 2,4-D, 2,4 DB, dichloprop, MCPA, mecoprop, diclofop, clomeprop, fluazifop, haloxyfop, fenoxaprop, quizalofop, propaquizafop, clodinafop and cyhalofop,
- i) anilides, such as propanil and mefenacet,

*Trade-mark

- j) protoporphorinogen oxidase inhibitors, e.g.
 (i) diphenyl ethers, such as bifenox, lactofen, acifluorfen, fluoroglycofen, fomesafen, oxyfluorfen, chlomethoxyfen and acloniphen, or
 5 (ii) 4-chlorophenylazoless, such as. pentoxazone, cinidon-ethyl, flumiclorac, pyraflufen, azafenidin, fluthiacet-methy, sulfentrazone, , carfentrazone, isopropazol, profluazol, and
 (iii) flumoxazin,
- k) ureas, such as chlortoluron, isoproturon, daimuron, linuron, monolinuron
 10 and thidiazuron,
- l) uracils, such as bromacil and lenacil,
- m) triazines, such as atrazine, simazine, cyanazine, symetryn, terbutryn, trietazine and triaziflam,
- n) carbamates, such as desmedipham, phenmedipham, triallate, molinate,
 15 dimepiperate, isopropilate, thiobencarb, esprocarm and asulam,
- o) pyridines, such as trichlopyr, picloram, diflufenican, fluroxypyr, thiazopyr and clopyralid,
- p) pyrazoles, such as pyrazolate, pyrazoxyfen and benzofenap,
- q) imidazolidinones, such as imazamethabenz, imazaquin, imazapyr,
 20 imazethapyr, imazamox and AC 263222,
- r) cyclohexanediones, such as alloxydim, sethoxydim, cycloxydim, tralkoxydim, clethodim and mesotrione,
- s) oxa- and thia-diazoles, such as oxadiazon, oxadiardyl and flufenacet, and
- t) triazinones, such as, met amitron and metribuzin,
- 25 u) miscellaneous compounds, such as isoxaflutole, cinmethylin, bentazon, ethofumesate, met amitron, metribuzin, fluorochloridinone and quinmerac.

Examples of insecticides and acaricides which can be used in conjunction with the citric acid derivates include

- 30 a) chlorinated hydrocarbons, such as endosulfan and lindane,
 b) nitroimines and cyanoimines, such as imidacloprid, thiamethoxam, thiacloprid and acetamiprid,
 c) pyrazoles, such as fipronil, vaniliprole, ethiprole, fenpyroximate, chlofenapyr, tebufenpyrad and tolfenpyrad,
 35 d) tin compounds, such as fenbutatin and azocyclotin,

- e) benzoylureas, such as chlorfluazuron, diflubenzuron, flucycloxuron, teflubenzuron, flufenoxuron, fluazuron, hexaflumuron, lufenuron, triflumuron and novaluron,
- f) other ureas, such as hexythiazox, triazamate, diafenthiuron and sulcofuron,
- 5 g) fermentation products, such as ivermectin, abamectin, spinosad, and, emamectin,
- h) benzoylhydrazines, such as tebufenozide, halofenozide and methoxyfenozide,
- i) carbamates, such as indoxacarb, bendiocarb, carbofuran, carbosulfan,
10 propoxur, methiocarb, aldicarb, methomyl, thiofanox, thiodicarb, pirimicarb and cartap,
- j) pyrethroids, e.g.
- (i) esters, such as permethrin, cypermethrin, deltamethrin, cyhalothrin, cyfluthrin, tralomethrin, flumethrin., acrinathrin,
15 fenvalerate, flucythrinate, fluvalinate, ZXI 8901, fenfluthrin, tefluthrin, biphenethrin, transfluthrin, tetramethrin, (including various stereo and/or geometric isomers of these compounds), and
- (ii) non-esters, such as etofenprox, silafluofen, flufenprox, halfenprox and protrifenbute
- 20 k) miscellaneous compounds, such as benzoximate, pyridaben, fenoxycarb, pyriproxyfen, hydamethylnon, buprofezin, chinomethionat, clofentezine, acequinocyl, chlofenapyrfenazaquin, pyrimidifen and cyromazine.

Examples of PGRs are chlormequat, paclobutrazol and ethephon.

25

The names quoted for these compounds are the non-proprietary common names and the chemical structure can be found for example by reference to "The Pesticide Manual" *ibid*. Structures of compounds not in the Pesticide Manual can be found in most compendia of pesticide common names.

30

We have found that the invention is particularly useful with pesticides of high melting point, i.e. greater than 100 °C (especially greater than 130 °C), and/or low solubility in organic solvents.

35

The citric acid derivatives of the invention are particularly efficacious with the following fungicides and groups of fungicides:

5 Conazoles (especially fluquinconazole or triticonazole), strobilurins (especially alkoxyacrylates e.g. azoxystrobin, or those containing an oximeamide moiety) or dimethomorph.

The citric acid derivatives of the invention are particularly efficacious with the following herbicides and groups of herbicides:

10

Alkyldiaminotriazines (especially triaziflam), aryloxyalkanoic acids and esters (especially fenoxaprop, quizalofop, fluazifop), pyridines (especially fluoroxyppy), carbamates (especially phenmedipham or desmedipham), protoporphorinogen oxidase inhibitors (especially carfentrazone), sulfonylureas (especially
15 nicosulfuron), HBNs, cyclohexanediones (especially mesotrione), imidazolidinones (especially imazapyr) and isoxaflutol.

The citric acid derivatives of the invention are particularly efficacious with the following insecticides:

20

Fipronil, indoxacarb, endosulfan, pirimicarb or clofentezine.

The citric acid derivative is preferably applied at a rate of 20 g to 2000 g, especially 50 g to 500 g, per 100 litres of the diluted formulation.

25

The citric acid derivative may be incorporated in conventional formulations (e.g. emulsions in water, suspoemulsions or solid formulation types after adsorption onto a suitable inert carrier, such as water dispersible granules) or may be added ("tank-mixed") to the pesticide just prior to use. It may be desirable also to add
30 small quantities of solvent and/or surfactant, especially a non-ionic surfactant, and other additives such as fatty acids to improve the emulsifiability of the citric acid derivative. Typically, the amount of emulsifier is 1 to 20% of the citric acid derivative. The choice of emulsifier is not critical to the performance of the invention, however alcohol ethoxylate surfactants form a preferred group.

35

The weight ratio of the pesticidal compound to citric acid derivative is preferably 1 to 50, especially 1 to 10.

Examples

The following Examples show how the pesticidal adjuvants of the invention enhance the efficacy of a variety of pesticidal active ingredients.

5

Examples 1 and 2 show enhancement of fungicidal efficacy against *Erysiphe graminis* (powdery mildew) and *Plasmopara viticola* respectively.

Example 3 shows enhancement of herbicidal efficacy against a number of
10 common weeds.

Example 4 shows enhancement of insecticidal efficacy against *Aphis fabae*.

Example 5 shows enhancement of insecticidal efficacy against the eggs of
15 *Tetranychus urticae* (red spider mite).

Example 1 – Fungicidal efficacy

In a glasshouse, wheat plants were inoculated with *Erysiphe graminis* (powdery
mildew). One day after inoculation, the plants were sprayed with various
20 fungicidal mixtures.

The spray mixtures were prepared by diluting with water a commercial
composition of the active ingredient and optionally a solution of the citric acid
derivative and emulsifier. When present, the citric acid derivative was either
25 tributyl citrate (TBC) or *O*-acetyl tributyl citrate (ATBC), and constituted 0.25%
w/v of the mixture (see Table A). The solution of citric acid derivative contained
20% emulsifier. The emulsifier was commercially available Synperonic A5^{*}
(supplied by Uniqema)

30 The concentration of active ingredient was selected to give an application rate
shown in Table A.

*Trade-mark

The commercial compositions of the active ingredients used in preparing the spray mixtures are given below.

- Fluquinconazole - Commercial composition sold as Castellan^{*}
 5 Azoxystrobin - Commercial composition sold as Amistar^{*}
 Triticonazole - Non-commercial composition formulated in-house to give a 10SC from technical grade triticonazole

10 One week after spraying, the wheat was assessed for control of disease. The results are shown in Table A. The degree of control, shown in the right hand side column, is the average of seven replicates.

Table A

Mix	Active ingredient (Rate g/ha)	Adjuvant	% Control
1a	Fluquinconazole (250)	nil	19.6
2a	Fluquinconazole (250)	TBC	97.1
3a	Fluquinconazole (250)	ATBC	96.1
4a	Azoxystrobin (250)	nil	19.6
5a	Azoxystrobin (250)	TBC	90.2
6a	Azoxystrobin (250)	ATBC	85.3
7a	Triticonazole (250)	nil	30.2
8a	Triticonazole (250)	TBC	65.1
9a	Triticonazole (250)	ATBC	73
10a	Triticonazole (125)	nil	24.8
11a	Triticonazole (125)	TBC	77.8
12a	Triticonazole (125)	ATBC	77.8
13a	Triticonazole (62.5)	nil	11.1
14a	Triticonazole (62.5)	TBC	63.5
15a	Triticonazole (62.5)	ATBC	65.1

*Trade-mark

Example 2 – Fungicidal efficacy

Using the same method as Example 1 vine seedlings were inoculated with *Plasmopara viticola* and sprayed with various mixtures shown in Table B. The spray mixtures were prepared in analogous fashion to Example 1 using dimethomorph as the active ingredient. The commercial composition of dimethomorph used was Forum.

The mixtures were sprayed on to the plants at a rate of 2.5 g of active ingredient per 100 litres of spray mixture.

One week after spraying, the wheat was assessed for control of disease. The results are shown in Table B. The degree of control, shown in the right hand side column, is the average of five replicates.

15

Table B

Mix	Active ingredient	Adjuvant	% Control
1b	Dimethomorph	nil	65.3
2b	Dimethomorph	TBC	96.4
3b	Dimethomorph	ATBC	72.02

Example 3 – Herbicidal efficacy

Seeds of the test species shown in Table C were sown in sandy loam soil and were grown in the glasshouse to the growth stages shown in Table C. Each species was then sprayed with various spray mixtures.

Table C

BROAD-LEAVED			GRASSES		
Scientific name	Code	Growth Stage	Scientific name	Code	Growth Stage
<i>Amaranthus retroflexus</i>	AMARE	11	<i>Lolium multiflorum</i>	LOLMU	12
<i>Chenopodium album</i>	CHEAL	11	<i>Alopecurus myosuroides</i>	ALOMY	12
<i>Stellaria media</i>	STEME	11-12	<i>Echinochloa crus-galli</i>	ECHCG	12
<i>Galium aparine</i>	GALAP	10-11	<i>Setaria viridis</i>	SETVI	11
<i>Pharbitis purpurea</i>	PHBPU	11-12	<i>Digitaria adscendens</i>	DIGSA	11-12

The spray mixtures were prepared in analogous fashion to Example 1. The active ingredient and citric acid derivative, and the concentration thereof, was selected to give an application rate shown in Table D.

- 5 The commercial compositions of the active ingredients used in preparing the spray mixtures are given below.

	Carfentrazone	- Commercial composition (WG50) sold as Affinity*
	Triaziflam	- Commercial composition (EW30)
10	Imazethapyr	- Commercial composition (EC24) sold as Pursuit.*
	Nicosulfuron	- Commercial composition (SC04) sold as Milagro*.
	Mesotrione	- Non-commercial composition (WP20) made in-house.

- 15 Mixtures were sprayed at a rate of 300 litre of spray mixture/hectare. In all cases two replicates were done. After spraying, the sprayed mixture was allowed to dry and the pots were returned to the glasshouse.

- 20 The herbicidal effects were assessed visually over a period of 3 weeks. Visual assessment were on a 100% basis (0 = no effect compared to untreated plants; 100 = complete kill of test species). The results are shown in Table D. Tributyl citrate and *O*-acetyl tributyl citrate in the absence of any active ingredient caused no herbicidal effects (see experiments 26d to 29d).

Table D

Mix	Active Ingredient (rate kg/ha)	Adjuvant (rate kg/ha)	LOLMU	ALOMY	ECHCG	SETVI	DIGSA	AMARE	CHEAL	STEME	GALAP	PHBPU	MEAN
1d	Carfentrazone (0.02)	nil	0	0	0	5	0	80	85	30	70	10 0	37
2d	Carfentrazone (0.02)	ATBC (0.3)	0	0	40	50	0	95	10 0	10	75	10 0	47
3d	Carfentrazone (0.02)	ATBC (0.9)	0	0	30	45	0	90	90	10	10 0	10 0	47
4d	Carfentrazone (0.02)	TBC (0.3)	0	0	40	55	10	80	95	0	98	10 0	48
5d	Carfentrazone (0.02)	TBC (0.9)	0	0	10	50	0	95	90	10	10 0	10 0	46
6d	Triaziflam (0.08)	nil	0	10	10	10	0	70	70	10	70	95	31

*Trade-mark

Mix	Active Ingredient (rate kg/ha)	Adjuvant (rate kg/ha)	LOLMU	ALOMY	ECHCG	SETVI	DIGSA	AMARE	CHEAL	STEME	GALAP	PHBPU	MEAN
7d	Triaziflam (0.08)	ATBC (0.3)	40	40	40	20	10	90	95	10	75	70	49
8d	Triaziflam (0.08)	ATBC (0.9)	0	30	50	50	0	95	95	20	80	90	51
9d	Triaziflam (0.08)	TBC (0.3)	50	30	50	10	10	80	90	10	60	95	49
10d	Triaziflam (0.08)	TBC (0.9)	10	10	90	20	30	85	95	20	60	98	52
11d	Imazethapyr (0.02)	nil	0	0	10	40	0	80	30	75	40	15	29
12d	Imazethapyr (0.02)	ATBC (0.3)	10	0	80	55	55	75	55	75	40	20	47
13d	Imazethapyr (0.02)	ATBC (0.9)	30	20	80	55	50	75	40	70	85	15	52
14d	Imazethapyr (0.02)	TBC (0.3)	0	10	80	60	55	70	60	80	70	20	51
15d	Imazethapyr (0.02)	TBC (0.9)	10	0	80	65	50	85	60	80	65	20	52
16d	Nicosulfuron (0.02)	nil	0	10	0	10	0	50	5	5	25	0	11
17d	Nicosulfuron (0.02)	ATBC (0.3)	0	10	20	40	0	60	10	0	25	10	18
18d	Nicosulfuron (0.02)	ATBC (0.9)	10	30	40	70	10	60	10	0	40	10	28
19d	Nicosulfuron (0.02)	TBC (0.3)	0	10	30	50	0	40	10	0	20	0	16
20d	Nicosulfuron (0.02)	TBC (0.9)	0	10	45	50	15	55	10	0	0	15	20
21d	Mesotrione (0.08)	nil	0	0	40	0	30	80	85	85	65	65	45
22d	Mesotrione (0.08)	ATBC (0.3)	0	5	80	0	75	70	90	70	65	65	52
23d	Mesotrione (0.08)	ATBC (0.9)	0	25	85	20	80	70	95	73	68	75	59
24d	Mesotrione (0.08)	TBC (0.3)	0	10	85	10	85	70	90	75	63	65	55
25d	Mesotrione (0.08)	TBC (0.9)	0	40	85	20	83	73	90	58	80	70	60
26d	-	ATBC (0.3)	0	0	0	0	0	0	0	0	0	0	0
27d	-	ATBC (0.9)	0	0	0	0	0	0	0	0	0	0	0
28d	-	TBC (0.3)	0	0	0	0	0	0	0	0	0	0	0
29d	-	TBC (0.9)	0	0	0	0	0	0	0	0	0	0	0

Example 4 – Insecticidal efficacy

Three-week old bean plants (*Vicia faba*) were sprayed three times to runoff with various mixtures.

5

The spray mixtures were prepared in analogous fashion to Example 1. The active ingredient and citric acid derivative, and the concentration thereof, was selected

SUBSTITUTE SHEET (RULE 26)

to give an application rate shown in Table E. The citric acid derivative was tributyl citrate (TBC).

The commercial compositions of the active ingredients used in preparing the spray mixtures are given below.

Endosulfan	- Commercial composition (EC35) sold as Thiodan.*
Deltamethrin	- Commercial composition (EC2.5) sold as Decis.*
Pirimicarb	- Commercial composition (SG 50) sold as Pirimor.*
10 Fipronil	- Commercial composition (WG80) sold as Regent.*

The plants were infested with *Aphis fabae* and the degree of mortality was measured 3 days and 7 days after infestation. The results are shown in columns 4 and 5 of Table E.

15

In order to test persistency, some of the plants were re-infested 3 days after the first infestation and the degree of mortality was measured 3 and 7 days after re-infestation. The results are shown in columns 6 and 7 of Table E. To test the persistency further, some of the plants were re-infested 7 days after the first infestation and the degree of mortality was measured 3 and 7 days after the second re-infestation. The results are shown in columns 8 and 9 of Table E.

20

The experiments were conducted at 25 °C and 60% humidity. Each result was the average of two replications.

25

Table E

Mix	Active ingredient (rate g/ha)	Adj.	Infested		Reinfested		Reinfested	
			@ 0 days		@ 3 days		@ 7 days	
			3 days	7 days	3 days	7 days	3 days	7 days
1e	Endosulfan (10)	nil	75	40	15	10	0	0
2e	Endosulfan (10)	TBC	98	96	80	63	23	10
3e	Endosulfan (3)	nil	23	8	0	0	0	0
4e	Endosulfan (3)	TBC	88	83	15	0	0	0
5e	Endosulfan (1)	nil	0	0	0	0	0	0

*Trade-mark

Mix	Active ingredient (rate g/ha)	Adj.	Infested @ 0 days		Reinfested @ 3 days		Reinfested @ 7 days	
			3 days	7 days	3 days	7 days	3 days	7 days
6e	Endosulfan (1)	TBC	20	15	0	0	0	0
7e	Deltamethrin (0.03)	nil	99	99	98	97	90	80
8e	Deltamethrin (0.03)	TBC	100	98	93	92	90	85
9e	Deltamethrin (0.01)	nil	96	93	80	68	65	30
10e	Deltamethrin (0.01)	TBC	97	94	93	88	65	45
11e	Deltamethrin (0.003)	nil	80	75	15	8	10	0
12e	Deltamethrin (0.003)	TBC	78	68	75	58	20	10
13e	Pirimicarb (0.3)	nil	20	20	0	0	0	0
14e	Pirimicarb (0.3)	TBC	83	65	30	15	15	0
15e	Pirimicarb (0.1)	nil	0	0	0	0	0	0
16e	Pirimicarb (0.1)	TBC	25	10	18	0	0	0
17e	Fipronil (0.3)	nil	88	85	35	30	0	0
18e	Fipronil (0.3)	TBC	98	98	95	93	85	80
19e	Fipronil (0.1)	nil	65	55	15	10	0	0
20e	Fipronil (0.1)	TBC	63	63	65	65	45	40

Example 5 – Insecticidal (ovicide) efficacy

Four week old bean plants were infested with a mixed population of the eggs of *Tetranychus urticae* (red spider mites). After 24 and 48 hours the motile stages were removed with pressurised air. The plants were sprayed to runoff with various spray mixtures.

The spray mixtures were prepared in analogous fashion to Example 1 using clofentezine at rates given in Table F. The citric acid derivative was tributyl citrate (TBC).

The compositions were prepared from commercially available Clofentezine* SC 42 (Apollo). The plants were placed in the glass house at 23 °C and 60% humidity.

The mortality of the eggs was assessed after 10 and 14 days. The results are shown in Table F.


*Trade-mark

Table F

Mix	Active ingredient (Rate g/ha)	Ajuvant	10 days	14 days
1f	Clofentezine (10)	nil	99	97
2f	Clofentezine (10)	TBC	100	100
3f	Clofentezine (3)	nil	98	95
4f	Clofentezine (3)	TBC	100	100
5f	Clofentezine (1)	nil	70	50
6f	Clofentezine (1)	TBC	100	100
7f	Clofentezine (0.3)	nil	40	25
8f	Clofentezine (0.3)	TBC	70	93

CLAIMS

1. The use of at least one citric acid derivative, which has a log octanol-water coefficient (log P) of 2.6 to 11 and an equivalent hydrocarbon (EH) value of 29 to 47, where the citric acid derivative has the general formula (I)



wherein:

R^1 , R^2 and R^3 are independently selected from the group consisting of:

- a) hydrogen,
- b) C_1 to C_{20} alkyl which may be substituted by $-\text{OR}^6$, where R^6 is hydrogen or alkyl,
- c) C_2 to C_{20} alkenyl which may be substituted by $-\text{OR}^6$, where R^6 is hydrogen or alkyl,
and
- d) C_2 to C_{20} alkynyl which may be substituted by $-\text{OR}^6$, where R^6 is hydrogen or alkyl;


R^4 is hydrogen or $-\text{C}(=\text{O})\text{R}^5$, where R^5 is as defined for R^1 ; and

R^1 , R^2 , R^3 and R^4 are chosen such that the log P and EH value of the citric acid derivative are as defined above;

as a fungicidal or herbicidal adjuvant.

2. The use according to claim 1 where in the log P is 2.7 to 11.
3. The use according to claim 2 where in the log P is 4 to 10.7,
4. The use according to claim 3 where in the log P is 4 to 10.3.
5. The use according to any one of claims 1 to 4 where the EH is 32 to 44.

6. A fungicidal or herbicidal composition comprising at least one fungicidal or herbicidal compound and at least one citric acid derivative defined in any one of claims 1 to 5.
7. A composition according to claim 6 wherein the fungicidal or herbicidal compound has a melting point greater than 100°C.
8. A composition according to claim 7 wherein the fungicidal or herbicidal compound has a melting point greater than 130°C.
9. Use of a fungicidal or herbicidal composition as defined in claim 6, 7 or 8 to combat a fungal or a herbal pest at a locus infested with said fungal or a herbal pest.
10. A method of combatting a fungal or herbal pest at a locus other than a human or animal body, wherein said locus is infested with or liable to be infested with said fungal or herbal pest, which comprises applying to said locus an effective amount of a fungicidal or herbicidal composition as defined in claim 6, 7 or 8.



(1)