



(51) International Patent Classification:

C07D 495/04 (2006.01) A61P 35/00 (2006.01)
A61K 31/407 (2006.01) A61P 3/04 (2006.01)

MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/EP2017/062009

Published:

— with international search report (Art. 21(3))

(22) International Filing Date:

18 May 2017 (18.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

16170238.6 18 May 2016 (18.05.2016) EP

(71) Applicant: ISTITUTO EUROPEO DI ONCOLOGIA
S.R.L. [IT/IT]; Via Filodrammatici 10, I-20121 Milano
(IT).


(72) Inventors: VIANELLO, Paola; Corso Sempione 63,
20149 Milano (IT). ROMUSSI, Alessia; Via Bozzano
11/38, 16143 Genova (IT). CAPPÀ, Anna; Via Villa S. An-
tonio 17, I-62039 Visso (IT). TRIFIRO', Paolo; Via Stra-
bone 35, I-22100 Como (IT). VARASI, Mario; Via Mon-
cucco 24/A, I-20142 Milano (IT). SARTORI, Luca; Via
per Caviglio 57, 22034 Brunate (IT). MERCURIO, Ciro;
Via Pietro Micca 23, I-20025 Legnano (IT).

(74) Agent: COOLEY (UK) LLP; Dashwood, 69 Old Broad
Street, London Greater London EC2M 1QS (GB).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR,
KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

(54) Title: IMIDAZOLES AS HISTONE DEMETHYLASE INHIBITORS



(I)

(57) Abstract: The present invention relates to imidazole derivatives, wherein A, R, R¹, and R² are as defined in the specification, pharmaceutical compositions containing such compounds and to their use in therapy.



Imidazoles as Histone Demethylase Inhibitors

Field of the Invention

The present invention relates to imidazole derivatives, pharmaceutical compositions containing such compounds and to their use in therapy.

Background of the invention

Alterations in the structural and functional states of chromatin, mainly determined by post-translational modifications of histone components, are involved in the pathogenesis of a variety of diseases. The enzymatic processes, which govern these post-translational modifications on the nucleosomes, have become potential targets for the so-called epigenetic therapies (Portela, A. *et al.* Nat. Biotechnol. 2010, 28, 1057-1068).

The discovery of an increasing number of histone lysine demethylases has highlighted the dynamic nature of the regulation of histone methylation, a key chromatin modification that is involved in eukaryotic genome and gene regulation. Histone lysine demethylases represent attractive targets for epigenetic drugs, since their expression and/or activities are often misregulated in cancer (Varier, R. A. *et al.* Biochim. Biophys. Acta. 2011, 1815, 75-89). A lysine can be mono-, di-, and tri-methylated and each modification, even on the same amino acid, can have different biological effects.

Histone lysine demethylases exert their activity through two different type of mechanism (Anand, R. *et al.* J. Biol. Chem. 2007, 282, 35425-35429; Metzger, E. *et al.* Nat. Struct. Mol. Biol. 2007, 14, 252-254). While the Jumonji domain-containing histone demethylases, which are iron and 2-oxoglutarate dependent oxygenases, act on mono-, di- and trimethylated lysines, the flavin-dependent (FAD) histone demethylases catalyse the cleavage of mono and dimethylated lysine residues. Currently, two FAD dependent demethylases have been identified: LSD1, also known as KDM1A, and LSD2, also known as KDM1B. (Culhane, J. C. *et al.* Curr. Opin. Chem. Biol. 2007, 11, 561-568, Ciccone, D. N. *et al.* Nature 2009, 461, 415-418).

KDM1A is a constituent in several chromatin-remodeling complexes and is often associated with the co-repressor protein CoREST. KDM1A specifically removes the methyl groups from mono- and di-methyl Lys4 of histone H3, which is a well-characterized

gene activation mark. KDM1A represents an interesting target for epigenetic drugs due to its over-expression in solid and hematological tumors (Schulte, J. H. *et al.* *Cancer Res.* 2009, 69, 2065-2071; Lim, S. *et al.* *Carcinogenesis* 2010, 31, 512–520; Hayami, S. *et al.* *Int. J. Cancer* 2011, 128, 574–586; Schildhaus, H. U. *et al.* *Hum. Pathol.* 2011, 42, 1667-1675; Bennani-Baiti, I. M. *et al.* *Hum. Pathol.* 2012, 43, 1300-1307). Its over-expression correlates to tumor recurrence in prostate cancer (Kahl, P. *et al.* *Cancer Res.* 2006, 66, 11341-11347), and KDM1A has a role in various differentiation processes, such as adipogenesis (Musri, M. M. *et al.* *J. Biol. Chem.* 2010, 285, 30034-30041), muscle skeletal differentiation (Choi, J. *et al.* *Biochem. Biophys. Res. Commun.* 2010, 401, 327-332), and hematopoiesis (Hu, X. *et al.* *Proc. Natl. Acad. Sci. USA* 2009, 106, 10141-10146; Li, Y. *et al.* *Oncogene.* 2012, 31, 5007-5018). KDM1A is further involved in the regulation of cellular energy expenditure (Hino S. *Et al.* *Nat Commun.* 2012, doi: 10.1038/ncomms1755), in the regulation of thermogenesis and oxidative metabolism in adipose tissue (Duteil *et al.* *Nat Commun.* 2014 Jun 10; 5:4093. doi: 10.1038/ncomms5093.), in the control of checkpoints of viral gene expression in productive and latent infections (Roizman, B. J. *Viol.* 2011, 85, 7474-7482), and more specifically in the control of herpes virus infection (Gu, H. J. *Viol.* 2009, 83, 4376-4385) and HIV transcription (Sakane, N. *et al.* *PLoS Pathog.* 2011, 7(8):e1002184). The role of KDM1A in the regulation of neural stem cell proliferation (Sun, G. *et al.* *Mol. Cell Biol.* 2010, 30, 1997-2005) and in the control of neuritis morphogenesis (Zibetti, C. *et al.* *J. Neurosci.* 2010, 30, 2521-2532) suggests its possible involvement in neurodegenerative diseases.

Furthermore, KDM1A has been found to be relevant in the control of other important cellular processes, such as DNA methylation (Wang, J. *et al.* *Nat. Genet.* 2009, 41(1):125-129), cell proliferation (Scoumanne, A. *et al.* *J. Biol. Chem.* 2007, 282, 15471-15475; Cho, H. S. *et al.* *Cancer Res.* 2011, 71, 655-660), epithelial mesenchimal transition (Lin, T. *et al.* *Oncogene.* 2010, 29, 4896-4904) and chromosome segregation (Lv, S. *et al.* *Eur. J. Cell Biol.* 2010, 89, 557-563). Moreover, KDM1A inhibitors were able to reactivate silenced tumor suppressor genes (Huang, Y. *et al.* *Proc. Natl. Acad. Sci. U S A.* 2007, 104, 8023-8028; Huang, Y. *et al.* *Clin. Cancer Res.* 2009, 15, 7217-7228), to target selectively cancer cells with pluripotent stem cell properties (Wang, J. *et al.* *Cancer Res.*

2011, 71, 7238-7249), as well as to reactivate the all-*trans*-retinoic acid differentiation pathway in acute myeloid leukemia (Schenk, T. *et al.* Nat Med. 2012, 18, 605-611). Moreover, KDM1A has a clear role in sustaining the oncogenic potential of MLL-AF9 translocation in leukaemia stem cells (Harris *et al.* Cancer Cell, 21 (2012), 473–487), as well as in the stem-like tumor propagating cells of human glioblastoma (Suvà *et al.* Cell 2014, 157, 580-594).

The more recently discovered demethylase KDM1B (LSD2) displays similarly to KDM1A specificity for mono- and di-methylated Lys4 of histone H3. KDM1B, differently from KDM1A, does not bind CoREST and it has not been found up to now in any of KDM1A-containing protein complexes (Karytinis, A. *et al.* J. Biol. Chem. 2009, 284, 17775-17782). On the contrary, KDM1B forms active complexes with euchromatic histone methyltransferases G9a and NSD3 as well as with cellular factors involved in transcription elongation. KDM1B has been reported to have a role as regulator of transcription elongation rather than that of a transcriptional repressor as proposed for KDM1A (Fang, R. *et al.* Mol. Cell 2010, 39, 222-233).

KDM1A and KDM1B are both flavo amino oxidase dependent proteins sharing a FAD coenzyme-binding motif, a SWIRM domain and an amine oxidase domain, all of which are integral to the enzymatic activity of KDM1 family members. Moreover, both KDM1A and KDM1B show a structural similarity with the monoamine oxidases MAO-A and MAO-B. Indeed, tranylcypromine, a MAO inhibitor used as antidepressant agent, was found to be also able to inhibit KDM1A. The compound acts as an irreversible inhibitor forming a covalent adduct with the FAD cofactor. (Lee, M. G. *et al.* Chem. Biol. 2006, 13, 563; Schmidt, D. M. Z. *et al.* Biochemistry 2007, 46, 4408). Tranylcypromine analogs and their KDM1A inhibitory activity have been described in Bioorg. Med. Chem. Lett. 2008, 18, 3047-3051, in Bioorg. Med. Chem. 2011, 19, 3709-3716, and in J. Am. Chem. Soc 2011, 132, 6827-6833. Further arylcyclopropylamine and heteroaryl cyclopropylamine derivatives as KDM1A, MAO-A and/or MAO-B enzyme inhibitors are disclosed in US2010/324147, in WO2012/045883, in WO2013/022047 and in WO2011/131576.


Reversible KDM1A inhibitors are not so common and no clinical data for them are so far available. Examples of reversible inhibitors are aminothiazoles as described in Med. Chem. Commun. 2013, 4, 1513–1522, a *N'*-(1-phenylethylidene)-benzohydrazide series

(J. Med. Chem. 2013, 56, 9496–9508, WO2013025805), or thienopyrrole derivatives (WO2016/034946). Thus, there is still a need for further reversible inhibitors having useful antitumor properties, adequate selectivity and stability of action, and possibly showing a higher activity with respect to specific subclasses thereof.

Description of the Invention


The present invention relates to substituted imidazole derivatives having highly potent inhibitory activities of the KDM1A enzyme and selective over monoamine oxidases (MAOs), useful in the prevention or therapy of diseases and conditions associated with the activity of the histone demethylases.

According to the present invention there are provided compounds of general formula (I):



(I)

wherein



R is hydrogen or **L¹-R⁵**;

R¹, R² are independently, hydrogen, C₁-C₄-alkyl, C₃-C₇ cycloalkyl, aryl, heteroaryl, -(CH₂)_zCF₃, or **R¹** and **R²**, taken together with the carbon atoms they are bound to, form with the imidazole a fused heterocycle or a fused heteroaryl;

z is an integer from zero to 6

R³, R⁴ are hydrogen or C₁-C₄-alkyl;

L¹ is -(CH₂)_j-Y-, or -CH₂-

j is an integer from 2 to 6;

Y is a bond, oxygen, or CH₂;

- R⁵** is C₁-C₄-alkyl or aryl, wherein the aryl is optionally substituted by halogen, C₁-C₆-alkyl or **L²-R⁶**;
- L²** is -(CH₂)_m-; -(CH₂)_n-**W**-(CH₂)_p-;
- R⁶** is heterocyclyl, wherein the heterocyclyl is optionally substituted by C₁-C₆-alkyl;
- m, n, p** are, independently, zero or an integer from 1 to 6;
- W** is oxygen, sulphur, NH, or CH₂;

or stereoisomers or pharmaceutically acceptable salts thereof.

In one embodiment, **R¹** and **R²** are independently hydrogen, C₁-C₄-alkyl, C₃-C₇ cycloalkyl, aryl, heteroaryl, or **R¹** and **R²**, taken together with the carbon atoms they are bound to, form with the imidazole a fused heterocycle or a fused heteroaryl.

Preferably, **R³** or **R⁴** are methyl or ethyl.

In a more preferred embodiment:

- R** is hydrogen or **L¹-R⁵**;
- R¹, R²** are independently, hydrogen, C₁-C₄-alkyl, cyclopropyl, cyclobutyl, phenyl, pyridyl, or **R¹** and **R²**, taken together with the carbon atoms they are bound to, form with the imidazole a fused heterocycle or a fused heteroaryl;
- R³, R⁴** are methyl or ethyl;
- L¹** is -(CH₂)₂-**Y**- or -CH₂-;
- Y** is a bond, oxygen, or CH₂;
- R⁵** is methyl or phenyl optionally substituted by **L²-R⁶**;
- L²** is -**W**-(CH₂)_p-;
- R⁶** is heterocyclyl, wherein the heterocyclyl is optionally substituted by C₁-C₆-alkyl;
- p** is zero or an integer from 1 to 6;
- W** is oxygen;

or stereoisomers or pharmaceutically acceptable salts thereof.

Particularly preferred compounds of general formula (I) include:

4-methyl-5-(4-phenyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(4-ethyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-(4-methyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(4-isopropyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
5-(4-cyclopropyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-(4-propyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(4-cyclobutyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-(5-methyl-4-phenyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(5-ethyl-4-phenyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
5-(5-isopropyl-4-phenyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-ethyl-5-(4-ethyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(4-cyclobutyl-1H-imidazol-2-yl)-4-ethyl-thieno[3,2-b]pyrrole;
4-methyl-5-(4-phenyl-5-propyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(4-ethyl-1H-imidazol-2-yl)-6-methyl-thieno[2,3-b]pyrrole;
5-(5-ethyl-4-phenyl-1H-imidazol-2-yl)-6-methyl-thieno[2,3-b]pyrrole;
4-methyl-5-(4,5,6,7-tetrahydro-1H-benzimidazol-2-yl)thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[4-(pyrrolidin-3-ylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-ethyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-cyclobutyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-cyclobutyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-ethyl-4-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-ethyl-4-phenyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-5-phenyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[3-(azepan-4-yloxy)phenoxy]ethyl]-4-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[3-(azepan-4-yloxy)phenoxy]ethyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole;

4-methyl-5-[5-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-4-propyl-imidazol-2-

yl]thieno[3,2-b]pyrrole;
4-methyl-5-[4-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-4-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-4-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-ethyl-5-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-isopropyl-5-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-[4-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole;
5-[1-[3-[3,4-bis(4-piperidylmethoxy)phenyl]propyl]-4-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;
4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;
5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-4-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[3-[3,4-bis(4-piperidylmethoxy)phenyl]propyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenyl]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-cyclobutyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-cyclobutyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-4-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;

4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;

5-[4-cyclobutyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-cyclobutyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

4-methyl-5-[4-propyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole;

4-methyl-5-[5-propyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole;

5-[4-cyclobutyl-1-[2-[4-(4-piperidyloxy)phenyl]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-ethyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-

b]pyrrole;
4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenyl]ethyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;
4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenyl]ethyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;
4-methyl-5-[1-[3-[4-(4-piperidyloxymethyl)phenyl]propyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;
4-methyl-5-[1-[3-[4-(4-piperidyloxymethyl)phenyl]propyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;
5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-1-(2-methoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-ethyl-1-(2-methoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-1-(2-phenoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-ethyl-1-(2-phenoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-(1-benzyl-5-ethyl-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-[4-(4-pyridyl)-1H-imidazol-2-yl]thieno[3,2-b]pyrrole;
5-(1H-benzimidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]benzimidazol-2-yl]thieno[3,2-b]pyrrole;
5-[4-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole
5-[5-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole
5-[4-cyclobutyl-1-[3-[4-[[[(3R)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole
5-[5-cyclobutyl-1-[3-[4-[[[(3R)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole
5-[4-cyclobutyl-1-[3-[4-[[[(3S)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

5-[5-cyclobutyl-1-[3-[4-[[[(3S)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

5-[4-cyclobutyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

5-[4-cyclobutyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

5-[5-cyclobutyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

5-[4-cyclobutyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

5-[5-cyclobutyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

4-methyl-5-[4-methyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole

4-methyl-5-[5-methyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole

5-[4-cyclobutyl-1-[2-[4-(4-piperidylmethoxy)phenyl]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

4-methyl-5-[1-[3-[3-(4-piperidyloxy)phenyl]propyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole

4-methyl-5-[1-[3-[3-(4-piperidyloxy)phenyl]propyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole

4-methyl-5-[1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole

4-methyl-5-[1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole

4-methyl-5-[1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole

4-methyl-5-[1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole

5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-

b]pyrrole

5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole

4-methyl-5-[4-(3,3,3-trifluoropropyl)-1H-imidazol-2-yl]thieno[3,2-b]pyrrole

4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-4-(3,3,3-trifluoropropyl)imidazol-2-yl]thieno[3,2-b]pyrrole or stereoisomers or pharmaceutically acceptable salts thereof.

In one embodiment, the compound of general formula (I) is a reversible KDM1A (LSD1) inhibitor.

In another embodiment, the invention provides the compounds of general formula (I) for use as medicament.

In another embodiment, the invention provides the compounds of general formula (I) for the use in the treatment and/or prevention of cancer, infectious diseases or a disease characterized by aberration of cellular energy metabolism, such as obesity. Preferably, the compounds of general formula (I) are for the use in the treatment and/or prevention of leukemia, non-small cell lung cancer, hepatocellular carcinoma, or glioblastomas. Still preferably the glioblastomas are giant cell glioblastoma or gliosarcoma.

In another embodiment, the invention provides a method for treating or preventing cancer, such as leukemia, non-small cell lung cancer, hepatocellular carcinoma, or glioblastomas, infectious diseases or a disease characterized by aberration of cellular energy metabolism, such as obesity, wherein the method comprises the step of administering a compound of general formula (I) to a subject in an effective amount.

In a further embodiment, the invention provides the use of a compound of general formula (I) for the manufacture of a medicament for treating or preventing cancer, such as leukemia, non-small cell lung cancer, hepatocellular carcinoma, or glioblastomas, infectious diseases or a disease characterized by aberration of cellular energy metabolism, such as obesity.

A further embodiment of the invention is a pharmaceutical composition comprising a compound of general formula (I), together with a pharmaceutically acceptable excipient and/or diluent. The pharmaceutical composition may further comprise at least one further therapeutic agent, preferably selected from the group consisting of histone deacetylase inhibitors, retinoid receptor modulators, anti-proliferative/antineoplastic agents, cytostatic

agents, agents which inhibit cancer cell invasion, inhibitors of growth factor function, anti-angiogenic agents, cell cycle inhibitors, proteasome inhibitors, HSP90 inhibitors, selective COX-2 inhibitors or a chemotherapeutic agent.

In the present invention, “aryl” represents a mono or bicyclic aromatic ring system of, respectively, 6, 9 or 10 atoms, examples of such an aryl are phenyl, indenyl, indanyl and naphthyl and tetrahydronaphthalenyl. Substituted aryl means that the hydrogen atoms on independently each carbon atom may be independently replaced by a substituent as defined herein above.

“Heteroaryl” represents a mono or bicyclic heteroaromatic ring system of, respectively, 5 to 10 members, which contains one, two, three or four heteroatoms selected from nitrogen, oxygen or sulphur and one to nine carbon atoms. Examples of said heteroaryls include, but are not limited to: pyrrolyl, furyl, thienyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, 1,2,3-oxadiazolyl, 1,3,4-thiadiazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzo[*b*]furanyl, benzo[*b*]thienyl, benzopyranyl, indazolyl, benzimidazolyl, purinyl, quinolyl, isoquinolyl, quinazolinyl, and quinoxalinyl.

“Heterocyclyl” represents a mono, bicyclic or a spirocyclic saturated or partially saturated non-aromatic ring system of, respectively, 4 to 12 members, which contains one, two, or three heteroatoms selected from nitrogen, oxygen, and sulphur and three to eleven carbon atoms. Examples of such heterocycles include, but are not limited to: pyrrolidyl, pyrrolidinyl, piperidyl, piperidinyl, piperazinyl, tetrahydrofuranyl, tetrahydropyranyl, morpholyl, thiomorpholyl, tetrahydroquinolyl, tetrahydroisoquinolyl, tetrahydroquinoxalinyl, benzodioxolyl, 2,3-dihydro-benzodioxinyl, benzoxazolyl, azetidyl, azepinyl, and diazapinyl. Examples of bicyclic ring systems include, but are not limited to, 2-aza-bicyclo[2.2.1]heptanyl, 2,5-diaza-bicyclo[2.2.1]hept-2-yl or 8-azabicyclo[3.2.1]octanyl. Examples of spirocyclic ring systems include, but are not limited to, 3,8-diazaspiro[4.5]decane. Substituted heterocyclyl means that one or two hydrogen atoms on independently each carbon atom or heteroatom may be independently replaced by a substituent as defined herein above.

“Fused heterocycle” represents a bicyclic partially saturated non-aromatic ring system of, respectively, 8 to 12 members. Examples of such heterocycles include, but are not limited to: tetrahydrocyclopenta[*d*]imidazole, tetrahydrobenzimidazol or

hexahydrocyclohepta[d]imidazole.

"Fused heteroaryl" represents a bicyclic aromatic ring system. In accordance with the invention, one ring of the fused heteroaryl is formed from the imidazole ring of the compound of formula I. An example of such heteroaryl includes but is not limited to benzimidazol.

The term "C₁-C₆ alkyl" refers to a straight or branched hydrocarbon chain radical, consisting solely of carbon and hydrogen atoms, having from one to six carbon atoms. The "C₁-C₆ alkyl" group is preferably a linear or branched C₁-C₄ alkyl group, more preferably a C₁-C₂ alkyl group. Suitable examples of C₁-C₆ alkyl include methyl, ethyl, n-propyl, isopropyl, butyl, *tert*-butyl, pentyl, and hexyl. The term "C₁-C₄ alkyl" refers to a straight or branched hydrocarbon chain radical, consisting solely of carbon and hydrogen atoms, having from one to four carbon atoms.

The term "C₃₋₇ cycloalkyl" refers to a saturated monocyclic hydrocarbon ring system having three to seven carbon atoms. Suitable examples of C₃₋₆-cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

The term "halogen" refers to fluoro, chloro, bromo, or iodo. "Halogens" are preferably fluorine, chlorine or bromine, being in particular fluorine or chlorine.

"Leaving group" refers to halogen, preferably to chloride, bromide or iodide.

"Reversible inhibitor" is an inhibiting molecular entity that interacts with an enzyme by non-covalent interactions and is able to associate/dissociate to the enzyme.

Pharmaceutically acceptable salts comprise conventional non-toxic salts obtained by salification of a compound of formula (I) with inorganic acids (e.g. hydrochloric, hydrobromic, sulphuric, or phosphoric acids), or with organic acids (e.g. acetic, propionic, succinic, benzoic, sulfanilic, 2-acetoxy-benzoic, cinnamic, mandelic, salicylic, glycolic, lactic, oxalic, malic, maleic, malonic, fumaric, tartaric, citric, *p*-toluenesulfonic, methanesulfonic, ethanesulfonic, or naphthalensulfonic acids). For reviews on suitable pharmaceutical salts see Berge S. M. *et al.*, J. Pharm. Sci. 1977, 66, 1-19; Gould P. L. Int. J. Pharm 1986, 33, 201-217; and Bighley *et al.* Encyclopedia of Pharmaceutical Technology, Marcel Dekker Inc, New York 1996, Volume 13, page 453-497. Other salts, which are not pharmaceutically acceptable, for example the trifluoroacetate salt, may be useful in the preparation of compounds of this invention and these form a further aspect


of the invention. The invention includes within its scope all possible stoichiometric and non-stoichiometric forms of the salts of the compounds of formula (I).

In addition, the compounds of formula (I) may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, EtOH and the like.

Certain compounds of formula (I) may exist in stereoisomeric forms (e.g. they may contain one or more asymmetric carbon atoms). The individual stereoisomers (enantiomers and diastereomers) and mixtures of these are included within the scope of the present invention. The present invention also covers the individual isomers of the compounds represented by formula (I) as mixtures with isomers thereof in which one or more chiral centres are inverted.


The invention also includes all suitable isotopic variations of a compound of the invention. Examples of isotopes that can be incorporated into compounds of the invention include isotopes such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl , respectively. Certain isotopic variations of the invention, for example, those in which a radioactive isotope such as ^3H or ^{14}C is incorporated, are useful in drug and/or substrate tissue distribution studies. Further, substitution with isotopes such as deuterium ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability. Isotopic variations of the compounds of the invention can generally be prepared by conventional procedures such as by the illustrative methods or by the preparations described in the examples hereafter using appropriate isotopic variations of suitable reagents.

Imidazoles of formula (I), wherein **R** is hydrogen, can be synthesized from amidine of formula **A1** by condensation with α -halogeno ketones of formula **A2**, which are commercially available or can be prepared according conventional methods well known to a person skilled in art or following the procedures described in WO 2012/049277, and where **X** is chlorine, bromine or iodine (Li, B. *et al.* Org. Process Res. Dev. 2002, 6, 682-683) in a suitable solvent, for instance THF or a THF/water mixture, at temperatures ranging from room temperature to reflux, as represented in **Scheme A-I** below.

**Scheme A-I**


wherein **A**, **R**¹, and **R**² are as defined above and **X** is a chlorine, bromine or iodine.

Imidazoles of formula (I), wherein **R** is **L**¹-**R**⁵ and **L**¹ and **R**⁵ are as defined above, can be obtained by reaction of compounds of formula (I), wherein **R** is hydrogen, with derivatives of formula **A3**, wherein **LG** is a leaving group, for example bromine, in a suitable solvent, for instance dimethylacetamide, and in presence of a base as represented in **Scheme A2** below:

**Scheme A-II**

wherein **A**, **R**¹, and **R**² are as defined above, **R** is **L**¹-**R**⁵ (with **L**¹ and **R**⁵ as defined above) and **LG** is a leaving group.

Amidines of formula **A1** can be prepared starting from compounds of formula **B1** and **B2** as described in **Scheme B** below:

**Scheme B**

wherein R^3 , R^4 and **A** are as defined above.

Ethyl 4H-thieno[3,2-b]pyrrole-5-carboxylate (compound **B1**, Fluorochem, Cat No. 067104) and ethyl 6H-thieno[2,3-b]pyrrole-5-carboxylate (compound **B2**, Sigma Aldrich, Cat. No. PH011284) are commercially available. Compound **B1** or **B2** can be alkylated with a suitable alkyl halide R^3-LG^1 or R^4-LG^1 , where R^3 and R^4 are as defined above and LG^1 is a leaving group, for example bromide or iodide, in presence of a suitable base, such as NaH, in a suitable solvent, such as DMF, at a temperature ranging from about 0 °C to about 50 °C.

The ethyl ester group of a compound of formula **B3** or **B4** can be hydrolysed into a carboxylic acid of formula **B5** according to known methods, e.g. by treatment with a base, such as LiOH or NaOH, in a suitable solvent, for example in ethanol, in THF or in a ethanol/water mixture. The hydrolysis of the ethyl ester can be carried out at a temperature ranging from 0 °C to the boiling point of the solvent.


The carboxylic acid of formula **B5** is converted into the primary amide of formula **B6** according to conventional methods well known to a person skilled in art. For example, the carboxylic acid of formula **B5** is converted into a suitable acylating agent, such as an acyl chloride, which is then treated with NH₃ in a suitable solvent such as water at 0°C.

Compound of formula **B6** can be transformed into nitrile of formula **B7** with dehydrating agents such as phosphorus pentoxide, SOCl₂, DBU/ethyl dichlorophosphate in a suitable solvent, for example CH₂Cl₂, toluene, acetonitrile at a temperature ranging from about 0 °C to reflux.

Amidine of formula **A1** can be prepared from a compound of formula **B7** by amidation of imidates (Pinner reaction), hydrolysis of *N*-silylated arylamidines (A.R. Sanger, *Inorg. Nucl. Chem. Lett.* 1973,9, 351), addition of aluminium amide to nitrile (Tetrahedron Letters 1995, 48, 8761-8764), or by reduction of acylated amidoxime of formula **B9**, that can be obtained by reaction of nitrile of formula **B7** with hydroxyl amine, followed by reaction with acetic anhydride. Reduction of a compound of formula **B9** can be carried out for instance by catalytic hydrogenation with palladium on charcoal using reducing agents such as H₂ (Ismail, M.A. *et al.* *J. Med. Chem.* 2003, 46, 4761-4769, Lange, U. E. W *et al.* *Bioorg. Med. Chem. Lett.* 2003, 13, 2029-2033), potassium or ammonium formate (Nadrah, K. *et al.* *Synlett*, 2007, 1257-125, Jnaneshwara, G. K. *et al.* *J. Chem. Research* 1998, 160-161), triethyl silane (Mahajan, U. S. *et al.* *Synthetic Communications* 2011, 41, 2195-2199) or with Zn in acetic acid (Goswami, R. *et al.* *Bioorg. Med. Chem.* 2014, 22, 3187-3203).

Compounds of formula **A3** can be prepared by known methods. For instance a compound of formula **C5**, where **LG** and **R⁶** are as defined above, **Y** and **L²** are oxygen can be obtained by the reaction of a diol derivative of formula **C1** with an alcohol of formula **C2**, that is carried out under the standard conditions of the Mitsunobu reaction, for instance by reaction with triphenylphosphine and diethylazodicarboxylate, at a temperature

ranging from about 0 °C to 80 °C, in a suitable solvent, such as tetrahydrofuran or toluene, for a time varying from about 30 min up to 72 h, to give a compound of formula **C3**. Reaction of a compound of formula **C3** with an alcohol of formula **C4** under the standard condition of the Mitsunobu reaction provides the intermediate **C5** as represented in **Scheme C** below:




Scheme C

wherein **p**, **R⁶** and **LG** are as defined above.

Compounds of formula **C1**, **C2**, and **C4** are known compounds or can be prepared by known methods.

Alternatively, reaction of a compound of formula **D1** with an alcohol of formula **D2** under the standard condition of the Mitsunobu reaction provides the intermediate **D3** as represented in **Scheme D** below:



Scheme D

wherein **p**, **R⁶** and **LG** are as defined above.

Compounds of formula **D1** and **D2** are known compounds or can be prepared by known methods.

In the case it is necessary to protect a chemical group of a compound of the present invention and/or an intermediate thereof, before carrying out one of the before described reactions, said chemical group can be protected and deprotected according to known methods. A thorough discussion for protection/deprotection steps is provided for example in Greene and Wuts (Greene, T.W.; Wuts, P.G.M. "Protective Groups in Organic Synthesis", John Wiley & Sons Inc., 2006) or in Kocienski (Kocienski, P.J. "Protecting Groups", George Thieme Verlag, 2005).

Salification of the compounds of formula (I), and preparation of compounds of formula (I), free of their salts, can be carried out by known conventional methods.

In view of the above described mechanisms of action, the compounds of the present invention are useful in the prevention or treatment of tumor type diseases, including but not limited to: acute and chronic myeloid leukaemia, acute and chronic lymphoblastic leukaemia, myelodysplastic syndromes, multiple myeloma, Hodgkin's disease, non-Hodgkin's lymphomas, cutaneous and peripheral T-cell lymphoma, adult T-cell leukemia, large B-cell lymphoma; mammary tumors; pulmonary tumors and pleural mesotheliomas, adenocarcinoma, non-small lung cancer, small-cell lung cancer; skin tumors including basal cell carcinomas (basaliomas), melanomas, squamous cell carcinoma, Kaposi's sarcoma, keratocanthomas, osteosarcomas, fibrosarcomas, rhabdomyosarcomas, neuroblastomas, glioblastomas, cerebral tumors, head and neck cancer, testicular and ovarian tumors, cervical carcinoma, endometrial and prostate tumors (for example advanced prostate cancer), thyroid carcinomas (for example thyroid follicular cancer), colon cancers (for example colon adenocarcinoma, colon adenoma), gastric tumors and gastrointestinal adenocarcinomas, hepatocellular carcinomas, pancreatic carcinomas (for example exocrine pancreatic carcinoma), renal tumors, teratocarcinomas and embryonic carcinomas.

The compounds of the invention are also useful in the prevention or treatment of infections, including, but not limited to, infections caused by protozoa, fungi, phytotoxic agents, viruses and parasites, for example HIV or herpes virus infections.

Furthermore, the compounds of the invention are also useful in the prevention or treatment of obesity.

The compounds of formula (I), can also be used in combination with additional agents, in particular anti-tumor and differentiating agents, either by separate administrations, or by including the two active principles in the same pharmaceutical formulation. Non-exhaustive examples of suitable additional agents include:

- a) histone deacetylase inhibitors (for example, but not limited to SAHA, PXD101, JNJ-26481585, SB939, ITF-2357, LBH589, PCI-24781, valproic acid, butyric acid, MS-275, MGCD0103 and FK-228);
- b) retinoid receptor modulators such as 13-*cis*-retinoic acid, 9-*cis*-retinoic acid, bexarotene, alitretinoin, or tretinoin; vitamin D;
- c) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical

oncology, such as alkylating agents (for example platin derivatives like *cis*-platin, carboplatin, oxaliplatin, lobaplatin, satraplatin, nedaplatin, heptaplatin; nitrogen mustard such as chlorambucil, melphalan, chlormethine, cyclophosphamide, ifosfamide, trofosfamide, uramustine, bendamustine, estramustine; busulphan, temozolomide or nitrosoureas); antimetabolites (for example antifolates such as aminopterin, methotrexate, pemetrexed, raltitrexed); purines such as cladribine, clofarabine, fludarabine, mercaptopurine, pentostatin, thioguanine; pyrimidines like capecitabine, cytarabine, fluorouracil, floxuridine, gemcitabine; azacitidine, decitabine; cytosine arabinoside or hydroxyurea; antitumour antibiotics (for example anthracyclines like aclarubicin, amrubicin, daunomycin, doxorubicin, epirubicin, idarubicin, valrubicin, zorubicine; mitoxantrone; or antibiotics from streptomyces like actinomycin, bleomycin, mitomycin, or plicamycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine or vinorelbine; taxoids like docetaxel, paclitaxel or tesetaxel; epothilones like ixabepilone) and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide; amsacrine, camptothecin, irinotecan, rubitecan, and topotecan);

d) cytostatic agents such as antioestrogens (for example but not limited to tamoxifen, toremifene, raloxifene, droloxifene and idoxifene), oestrogen receptor down regulators (for example but not limited to fulvestrant), antiandrogens (for example but not limited to bicalutamide, flutamide, nilutamide, liarozole or cyproterone acetate), LHRH antagonists or LHRH agonists (for example but not limited to goserelin, leuprorelin or buserelin), progestogens (for example but not limited to megestrol acetate), aromatase inhibitors (for example but not limited to anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5-alpha-reductase such as finasteride;

e) agents which inhibit cancer cell invasion (for example metalloproteinase inhibitors and inhibitors of urokinase plasminogen activator receptor function);

f) inhibitors of growth factor function, for example growth factor antibodies, growth factor receptor antibodies (for example but not limited to the anti-erbb2 antibody trastuzumab, the anti-erbbl antibody cetuximab and panitumumab, the anti IGF1R antibody figitumumab), farnesyl transferase inhibitors, MEK inhibitors, tyrosine kinase inhibitors and serine/threonine kinase inhibitors, for example enzastaurin, dasatinib,

erlotinib, gefitinib, imatinib, lapatinib, nilotinib, sorafenib, sunitinib, everolimus, sirolimus or temsirolimus;

g) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth factor, for example the anti-vascular endothelial cell growth factor antibody bevacizumab, lenalidomide or thalidomide;

h) cell cycle inhibitors including for example CDK inhibitors (for example but not limited to flavopiridol, roscovitine) and other inhibitors of cell cycle checkpoints; inhibitors of aurora kinase and other kinases involved in mitosis and cytokinesis regulation;

i) proteasome inhibitors (for example but not limited to lactacystin, bortezomib, epoxomicin);

j) HSP90 inhibitors (for example but not limited to AT-13387, KOS-953, KOS-1022, CNF-1010, CNF-2024, SNX 5422, STA-9090, NVP-HSP990, NVP-AUY922, PU-H17 and XL-888)

k) Selective COX-2 inhibitors (for example but not limited to celecoxib), or non selective NSAIDs (for example but not limited to diclofenac, flurbiprofen, ibuprofen, ketoprofen, or naproxen).

In another aspect, a compound of general formula (I) can be used in combination with radiation therapy. In yet another aspect, a compound of general formula (I) may be administered in combination with standard chemotherapy combinations such as, but not restricted to, CMF (cyclophosphamide, methotrexate and 5-fluorouracil), CAF (cyclophosphamide, doxorubicin and 5-fluorouracil), AC (doxorubicin and cyclophosphamide), FEC (5-fluorouracil, epirubicin, and cyclophosphamide), ACT or ATC (doxorubicin, cyclophosphamide, and paclitaxel), or CMFP (cyclophosphamide, methotrexate, 5-fluorouracil and prednisone).

The invention also provides pharmaceutical compositions comprising one or more compounds of formula (I), and one or more pharmaceutically acceptable excipient and/or diluent. The pharmaceutical compositions can be chosen on the basis of the treatment requirements. Such compositions are prepared by blending and are suitably adapted to oral or parenteral administration, and as such can be administered in the form of tablets, capsules, oral preparations, powders, granules, pills, injectable or infusible liquid, solutions, suspensions, emulsions, suppositories, ointments, creams, lotions, gels,

pastes, transdermal delivery devices.

Tablets and capsules for oral administration are normally presented in unit dose form and contain conventional excipients such as binders, fillers (including cellulose, mannitol, lactose), diluents, tableting agents, lubricants (including magnesium stearate), detergents, disintegrants (e.g. polyvinylpyrrolidone and starch derivatives such as sodium glycolate starch), coloring agents, flavoring agents, and wetting agents (for example sodium lauryl sulfate).

The oral solid compositions can be prepared by conventional methods of blending, filling or tableting. The blending operation can be repeated to distribute the active principle throughout compositions containing large quantities of fillers. Such operations are conventional.

Oral liquid preparations can be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or can be presented as a dry product for reconstitution with water or with a suitable vehicle before use. Such liquid preparations can contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethyl cellulose, carboxymethyl cellulose, aluminium stearate gel, or hydrogenated edible fats; emulsifying agents, such as lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which can include edible oils), such as almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, such as methyl or propyl *p*-hydroxybenzoate or sorbic acid, and if desired, conventional flavoring or coloring agents. Oral formulations also include conventional slow-release formulations such as enterically coated tablets or granules.

Pharmaceutical preparation for administration by inhalation can be delivered from an insufflator or a nebulizer pressurized pack.

For parenteral administration fluid unit dosages can be prepared, containing the compound and a sterile vehicle. The compound can be either suspended or dissolved, depending on the vehicle and concentration. The parenteral solutions are normally prepared by dissolving the compound in a vehicle, sterilising by filtration, filling suitable vials and sealing. Advantageously, adjuvants such as local anaesthetics, preservatives and buffering agents can also be dissolved in the vehicle. To increase the stability, the

composition can be frozen after having filled the vials and removed the water under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound can be suspended in the vehicle instead of being dissolved, and sterilized by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent can be included in the composition to facilitate uniform distribution of the compound of the invention.

For buccal or sublingual administration the compositions may be tablets, lozenges, pastilles, or gel.

The compounds can be pharmaceutically formulated as suppositories or retention enemas, e.g. containing conventional suppositories bases such as cocoa butter, polyethylene glycol, or other glycerides, for a rectal administration.

Another means of administering the compounds of the invention regards topical treatment. Topical formulations can contain for example ointments, creams, lotions, gels, solutions, pastes and/or can contain liposomes, micelles and/or microspheres. Examples of ointments include oleaginous ointments such as vegetable oils, animal fats, semisolid hydrocarbons, emulsifiable ointments such as hydroxystearin sulfate, anhydrous lanolin, hydrophilic petrolatum, cetyl alcohol, glycerol monostearate, stearic acid, water soluble ointments containing polyethylene glycols of various molecular weights. Creams, as known to formulation experts, are viscous liquids or semisolid emulsions, and contain an oil phase, an emulsifier and an aqueous phase. The oil phase generally contains petrolatum and an alcohol such as cetyl or stearic alcohol. Formulations suitable for topical administration to the eye also include eye drops, wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent for the active ingredient.

A further method of administering the compounds of the invention regards transdermal delivery. Typical transdermal formulations comprise conventional aqueous and non-aqueous vectors, such as creams, oils, lotions or pastes or can be in the form of membranes or medicated patches.

A reference for the formulations is the book by Remington ("Remington: The Science and Practice of Pharmacy", Lippincott Williams & Wilkins, 2000).

The compounds of the present invention may be employed alone as a sole therapy or in

combination with other therapeutic agents for the treatment of the above-mentioned conditions. The combination can be administered as separate compositions (simultaneous, sequential) of the individual components of the treatment or as a single dosage form containing both agents. When the compounds of this invention are in combination with others active ingredients, the active ingredients may be separately formulated into single-ingredient preparations of one of the above-described forms and then provided as combined preparations, which are given at the same time or different times, or may be formulated together into a two- or more- ingredient preparation.

Compounds of general formula (I) may be administered to a patient in a total daily dose of, for example, from 0.001 to 1000 mg/kg body weight daily. Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose. The determination of optimum dosages for a particular patient is well known to one skilled in the art.

As is common practice, the compositions are normally accompanied by written or printed instructions for use in the treatment in question.

The following examples and biological data are presented in order to further illustrate the invention.

1. CHEMICAL SYNTHESIS

Unless otherwise indicated, commercially available reagents and solvents (HPLC grade) were used without further purification. Specifically, the following abbreviations may have been used in the descriptions of the experimental methods:

NMR (Nuclear Magnetic Resonance)	¹ H (proton)
MHz (Megahertz)	Hz (Hertz)
HPLC (High Performance Liquid Chromatography)	LC-MS (Liquid Chromatography Mass Spectrum)
s (seconds)	min (minutes)
h (hours)	mg (milligrams)
g (grams)	μL (microlitres)
mL (millilitres)	mmol (millimoles)

nm (nanometers)	μ M (micromolar)
M (molarity)	RT (room temperature)
AcOH (acetic acid)	
BOC or boc (<i>tert</i> -butoxycarbonyl)	(BOC) ₂ O (di- <i>tert</i> -butyl dicarbonate)
CBr ₄ (carbon tetrabromide)	CDCl ₃ (euterated chloroform)
CH ₂ Cl ₂ (dichloromethane)	DBU (1,8-diazabicyclo[5.4.0]undec-7-ene)
CH ₃ CN (acetonitrile)	DIAD (diisopropyl azodicarboxylate)
DIPEA (<i>N,N</i> -diisopropylethylamine)	
DMA (dimethylacetamide)	DMAP (dimethylaminopyridine)
DMF (dimethylformamide)	DMSO (dimethyl sulfoxide)
DMSO-d ₆ (deuterated dimethyl sulfoxide)	DTT (dithiothreitol)
Et ₂ O (diethyl ether)	EtOAc (ethyl acetate)
EtOH (ethanol)	HCl (hydrochloric acid)
LiHMDS ([bis(trimethylsilyl)amino]lithium)	LiOH (lithium hydroxide)
K ₂ CO ₃ (potassium carbonate)	MAO A (monoamine oxidase A)
MAO B (monoamine oxidase B)	MeOH (methanol)
MeOH-d ₄ (deuterated methanol)	NaBH ₄ (sodium borohydride)
NaBH(OAc) ₃ (sodium triacetoxyborohydride)	NaH (sodium hydride)
NaCl (sodium chloride)	NaHCO ₃ (sodium bicarbonate)
Na ₂ SO ₄ (sodium sulphate)	NBS (1-bromopyrrolidine-2,5-dione)
NH ₃ (ammonia)	NH ₄ Cl (ammonium chloride)
PPh ₃ (triphenylphosphine)	SOCl ₂ (thionyl chloride)
TEA (triethylamine)	THF (tetrahydrofuran)
Tris (tris(hydroxymethyl)aminomethane)	Pd(dba) ₂ (<i>bis</i> (dibenzylideneacetone)palladium(0))

Except where indicated otherwise, all temperatures are expressed in °C (degrees centigrade) or K (Kelvin).

The ^1H -NMR spectra were acquired with a Varian 500 MHz instrument. The chemical shifts are expressed in parts per million (ppm, δ units). The coupling constants are expressed in Hertz (Hz) and the splitting patterns are described as s (singlet), bs (broad signal), d (doublet), t (triplet), q (quartet), quint (quintet), m (multiplet).

The LC-MS analyses were carried out on a Waters Acquity UPLC or Waters Acquity UPLC H-Class linked to with a SQD Single quadrupole (Waters) using an Acquity UPLC BEH C18 (50 x 2.1 mm, 1.7 μm) or Acquity UPLC HSS T3 (50 x 2.1 mm, 1.8 μm) column. Phase A was composed by either Milli-Q water/ CH_3CN 95/5 + 0.07% formic acid or Milli-Q water + 0.07% formic acid; Phase B by CH_3CN + 0.05% formic acid; flow rate: 0.6 mL/min; UV detection (DIODE array) from 210 to 400 nm; ESI+ detection in the 100-2000 m/z range. The yields were calculated assuming that products were 100% pure if not stated otherwise.


Intermediate 1: 4-Methylthieno[3,2-b]pyrrole-5-carboxylic acid

4-Methylthieno[3,2-b]pyrrole-5-carboxylate was prepared as described in EP application 14183755.9.

Intermediate 2: 4-Ethylthieno[3,2-b]pyrrole-5-carboxylic acid

4-Ethylthieno[3,2-b]pyrrole-5-carboxylate was prepared as described in EP application 14183755.9.


Intermediate 3: 4-Methylthieno[3,2-b]pyrrole-5-carboxamide



1.7 g (14 mmol, 1 mL) of SOCl_2 and 3 drops of dry DMF were added at RT to a solution of 2.12 g (11.7 mmol) of 4-methylthieno[3,2-b]pyrrole-5-carboxylic acid (Intermediate 1) dissolved in 15 mL of dry CH_2Cl_2 /THF (3:1, v:v). The mixture was stirred for 2 h at reflux, then cooled down to RT and added dropwise to 8.2 mL (58.5 mmol) of a 28% solution in


water of ammonium hydroxide at 0°C. After 5 min the reaction mixture was diluted with water and extracted with EtOAc. The organic phases were dried over Na₂SO₄, filtered and evaporated to dryness to give 2.1 g of 4-methylthieno[3,2-b]pyrrole-5-carboxamide (99%) as brown solid. ¹H NMR (DMSO) δ (ppm): 7.70-7.65 (bs, 1H), 7.44 (d, *J*=5.4 Hz, 1 H), 7.18 (d, *J*=5.4 Hz, 1 H), 6.87–6.84 (bs, 1 H), 7.09 (s, 1H), 3.97 (s, 3 H); MS (ESI): *m/z*: 181 [M+H]⁺.

Intermediate 4: 4-Ethylthieno[3,2-b]pyrrole-5-carboxamide



4-Ethylthieno[3,2-b]pyrrole-5-carboxamide was prepared according to the procedure for Intermediate 3, starting from 4-ethylthieno[3,2-b]pyrrole-5-carboxylic acid (Intermediate 2), providing 4-ethylthieno[3,2-b]pyrrole-5-carboxamide (99%) as a brown solid. ¹H NMR (CDCl₃) δ (ppm): 7.31 (d, *J*=5.4 Hz, 1 H), 6.98 (d, *J*=5.4 Hz, 1 H), 6.85 (s, 1 H), 5.57 (bs, 2 H), 4.59 (q, *J*=6.8 Hz, 2 H), 1.44 (t, *J*=6.8 Hz, 3 H); MS (ESI): *m/z*: 195 [M+H]⁺.


Intermediate 5: 4-Methylthieno[3,2-b]pyrrole-5-carbonitrile



0.62 g (3.84 mmol, 0.39 mL) of ethyl dichlorophosphate and 0.93 g (5.76 mmol, 0.58 mL) of DBU were added to a solution of 0.35 g (1.92 mmol) of 4-ethylthieno[3,2-b]pyrrole-5-carboxamide (Intermediate 3) in dry CH₂Cl₂ (10 mL) at 0°C. After stirring at RT for 12 h further 0.62 g (3.84 mmol) of ethyl dichlorophosphate and 0.93 g (5.76 mmol, 0.58 mL) of DBU were added and the reaction was carried out for additional 2 h. The mixture was then diluted with water and brought to acidic pH with 2 M HCl; the aqueous phase was extracted with CH₂Cl₂ and the combined organic layers were dried over Na₂SO₄, filtered and evaporated to dryness. The crude product was purified by flash chromatography on silica gel (hexane/EtOAc, 1% of EtOAc to 10% of EtOAc) to give 0.234 g of 4-


methylthieno[3,2-b]pyrrole-5-carbonitrile (75%) as pale yellow solid. ^1H NMR (CDCl_3) δ (ppm): 7.41 (d, $J=5.4$ Hz, 1 H), 7.02 (s, 1 H), 6.97 (d, $J=5.4$ Hz, 1 H), 3.90 (s, 3 H).

Intermediate 6: 4-Ethylthieno[3,2-b]pyrrole-5-carbonitrile




4-Ethylthieno[3,2-b]pyrrole-5-carbonitrile was prepared according to the procedure for Intermediate 5, starting from 4-ethylthieno[3,2-b]pyrrole-5-carboxamide (Intermediate 4), providing 4-ethylthieno[3,2-b]pyrrole-5-carbonitrile intermediate (73%) as a pale yellow solid. ^1H NMR (CDCl_3) δ (ppm): 7.43 (d, $J=5.4$ Hz, 1 H), 7.01 (s, 1 H), 6.95 (d, $J=5.4$ Hz, 1 H), 4.28 (q, $J=7.3$ Hz, 2 H), 1.51 (t, $J=7.3$ Hz, 3 H).

Intermediate 7: *N'*-Hydroxy-4-methyl-thieno[3,2-b]pyrrole-5-carboxamidine




0.58 g (8.3 mmol) of hydroxylamine hydrochloride and 1.07 g (8.34 mmol, 1.45 mL) of DIPEA were added to a solution of 1.23 g (7.58 mmol) of 4-methylthieno[3,2-b]pyrrole-5-carbonitrile (Intermediate 5) in 16 mL of dry EtOH. The mixture was heated at 55°C for 2 h, then further 0.2 g (2 mmol) of hydroxylamine hydrochloride and 0.3 g (2 mmol, 0.4 mL) of DIPEA were added and the mixture was stirred for additional 2 h. The solution was concentrated and the residue was portioned between CH_2Cl_2 /water. The aqueous layer was extracted with CH_2Cl_2 and the combined organic layers were dried over Na_2SO_4 , filtered and evaporated to dryness to give 1.47 g of *N'*-hydroxy-4-methyl-thieno[3,2-b]pyrrole-5-carboxamidine (99%) as pale yellow solid. ^1H NMR ($\text{DMSO}-d_6$) δ (ppm): 9.62 (s, 1 H), 7.28 (d, $J=5.4$ Hz, 1 H), 7.13 (d, $J=5.4$ Hz, 1 H), 6.71 (s, 1 H), 5.72 (bs, 2 H), 3.86 (s, 3 H); MS (ESI): m/z : 196 $[\text{M}+\text{H}]^+$.

Intermediate 8: 4-Ethyl-*N'*-hydroxy-thieno[3,2-b]pyrrole-5-carboxamidine




4-Ethyl-*N'*-hydroxy-thieno[3,2-b]pyrrole-5-carboxamidine was prepared according to the procedure for Intermediate 7 starting from 4-ethylthieno[3,2-b]pyrrole-5-carbonitrile (Intermediate 6) providing 4-ethyl-*N'*-hydroxy-thieno[3,2-b]pyrrole-5-carboxamidine (99%) as a pale yellow solid. ^1H NMR (CDCl_3) δ (ppm): 7.19 (d, $J=5.4$ Hz, 1 H), 6.95 (d, $J=45.4$ Hz, 1 H), 6.65 (s, 1 H), 4.86 (bs, 2 H), 4.39 (q, $J=7.3$ Hz, 2 H), 1.38 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 210 $[\text{M}+\text{H}]^+$.

Intermediate 9: Amino-(4-methylthieno[3,2-b]pyrrol-5-yl)methylene]amino] acetate




0.95 g (9.3 mmol, 0.88 mL) of acetic anhydride were added to a solution of 1.52 g (7.79 mmol) of *N'*-hydroxy-4-methyl-thieno[3,2-b]pyrrole-5-carboxamidine (Intermediate 7) in 20 mL of AcOH at RT and the mixture was stirred at RT for 15 min. Then the mixture was diluted with water/ CH_2Cl_2 and neutralized with a saturated solution of NaHCO_3 . The aqueous phase was extracted with CH_2Cl_2 and the organic layers were dried over Na_2SO_4 , filtered and evaporated to dryness. The obtained dark solid was purified by column chromatography on silica gel (EtOAc/hexane 1:10, v:v to 8:2, v:v) to provide 1.57 g of amino-(4-methylthieno[3,2-b]pyrrol-5-yl)methylene]amino] acetate (85%) as a dark solid. ^1H NMR ($\text{DMSO}-d_6$) δ (ppm): 7.39 (d, $J=5.4$ Hz, 1 H), 7.17 (d, $J=5.4$ Hz, 1 H), 6.89 (s, 1 H), 6.70 (s, 2 H), 3.92 (s, 3 H), 2.15 (s, 3 H); MS (ESI): m/z : 497 $[2\text{M}+\text{Na}]^+$.

Intermediate 10: Amino-(4-ethylthieno[3,2-b]pyrrol-5-yl)methylene]amino] acetate



Amino-(4-ethylthieno[3,2-b]pyrrol-5-yl)methylene]amino] acetate was obtained as a dark solid following the procedure for Intermediate **9** starting from 4-ethyl-*N'*-hydroxy-thieno[3,2-b]pyrrole-5-carboxamidine (Intermediate **8**) (80%). ¹H NMR (CDCl₃) δ (ppm): 7.23 (d, *J*=5.4 Hz, 1H), 6.96 (d, *J*=5.4 Hz, 1 H), 6.75 (s, 1 H), 5.00 (bs, 2 H), 4.50 (q, *J*=7.2 Hz, 2 H), 2.28 (s, 3 H), 1.42 (t, *J*=7.2 Hz, 3 H); MS (ESI): *m/z*: 274 [M+Na]⁺.


Intermediate 11: 4-Methylthieno[3,2-b]pyrrole-5-carboxamidine acetate



1.56 g (6.57 mmol) of amino-(4-methylthieno[3,2-b]pyrrol-5-yl)methylene]amino] acetate (Intermediate **9**) were dissolved in 250 mL of EtOH and 3 mL of AcOH were added. The mixture was hydrogenated in an H-cube apparatus using a 10% Pd/C cartridge at 35°C and at atmospheric pressure with a flow of 0.5 mL/min. Then the solution was concentrated and the residue was precipitated with Et₂O. The solid was filtered off and dried to afford 1.07 g of 4-methylthieno[3,2-b]pyrrole-5-carboxamidine acetate (70%) as a grey powder. ¹H NMR (DMSO-*d*₆) δ (ppm): 10.99–8.34 (bs, 4 H), 7.52 (d, *J*=5.4 Hz, 1 H), 7.24 (d, *J*=5.4 Hz, 1 H), 7.00 (s, 1 H), 3.91 (s, 3 H), 1.74 (s, 3 H); MS (ESI): *m/z*: 180 [M+H]⁺.


Alternatively intermediate **11** can be obtained directly from 4-methylthieno[3,2-b]pyrrole-5-carbonitrile intermediate **5**. 2.09 g (12.88 mmol) of intermediate **5** was dissolved in dry THF (7 mL) and added to a 1M solution of lithium bis(trimethylsilyl)amide (14.17 mmol, 14.17 mL). After stirring at rt for 4.5h the reaction mixture was quenched at 0°C with a 2N solution of HCl in ether (57.96 mmol, 30 mL) and stirred at 0°C for 15 minutes. After overnight standing in the fridge the mixture was filtered and the solid was washed with ether and dried to give 4-methylthieno[3,2-b]pyrrole-5-carboxamidine hydrochloride intermediate **11** (96%) as beige powder.

Intermediate 12: 4-Ethylthieno[3,2-b]pyrrole-5-carboxamidine acetate



0.097 g (0.83 mmol, 0.13 mL) of triethylsilane were added dropwise at 0°C 0.06 g (0.06 mmol) to a suspension of 10% Pd/C and of 0.07 g (0.28 mmol) of amino-(4-ethylthieno[3,2-b]pyrrol-5-yl)methylene]amino] acetate (Intermediate **10**) in 0.5 mL of dry MeOH. The mixture was allowed to reach RT within 1 h and was stirred at RT for further 30 min. The mixture was filtered on a celite bed, concentrated and the residue was triturated with CH₂Cl₂/Et₂O to afford 49 mg of 4-ethylthieno[3,2-b]pyrrole-5-carboxamidine acetate (69%) as a grey powder. ¹H NMR (DMSO-d₆) δ (ppm): 10.60–8.43 (bs, 4 H), 7.48 (d, *J*=5.4 Hz, 1 H), 7.25 (d, *J*=45.4 Hz, 1 H), 6.96 (s, 1 H), 4.45 (q, *J*=7.1 Hz, 2 H), 1.74 (s, 3 H), 1.22 (t, *J*=7.1 Hz, 3 H); MS (ESI): *m/z*: 194 [M+H]⁺.

Intermediate 13: 6-methylthieno[2,3-b]pyrrole-5-carboxamidine hydrochloride



6-Methylthieno[2,3-b]pyrrole-5-carboxylic acid

6-methylthieno[2,3-b]pyrrole-5-carboxylic acid was prepared as described in EP application 14183755.9.

6-methylthieno[2,3-b]pyrrole-5-carboxamide

6-methylthieno[2,3-b]pyrrole-5-carboxamide was prepared according to the procedure for Intermediate **3**, starting from 6-methylthieno[2,3-b]pyrrole-5-carboxylic acid providing 6-methylthieno[2,3-b]pyrrole-5-carboxamide (99%) as a pale yellow solid. ¹H NMR (DMSO-d₆) δ (ppm): 7.65 (bs, 1 H), 7.11 (d, *J*=4.9 Hz, 1 H), 7.06 (s, 1 H), 7.05–7.02 (m, 2 H), 7.05 (3.94 (s, 3 H). MS (ESI): *m/z*: 181 [M+H]⁺.

6-methylthieno[2,3-b]pyrrole-5-carbonitrile


6-methylthieno[2,3-b]pyrrole-5-carbonitrile was prepared according to the procedure for Intermediate **5**, starting from 6-methylthieno[2,3-b]pyrrole-5-carboxamide providing 6-methylthieno[2,3-b]pyrrole-5-carbonitrile (64%) as a pale yellow solid. ¹H NMR (CDCl₃) δ

(ppm): 7.03–7.02 (d, $J=5.4$ Hz, 1 H), 7.00 (s, 1 H), 6.99 (d, $J=5.4$ Hz, 1 H), 3.87 (s, 3 H).

6-methylthieno[2,3-b]pyrrole-5-carboxamidinium hydrochloride

0.391 g (2.41 mmol) of 6-methylthieno[2,3-b]pyrrole-5-carbonitrile was dissolved in 2.5 mL of dry THF and was added at RT dropwise to 3.1 mL (3.1 mmol) of a 1 M solution of lithium hexamethyldisilazide in THF. The mixture was stirred at RT overnight. Then further 0.2 mL (0.2 mmol) of a 1 M solution of lithium hexamethyldisilazide were added. After stirring at RT for further 3 h, 1.2 mL (2.4 mmol) of a 2 M solution of HCl in Et₂O were added dropwise at 0°C and the mixture was allowed to reach RT in 15 min. Some precipitate was observed and the mixture was diluted with Et₂O, filtered and the residue was washed with Et₂O providing 0.444 g of 6-methylthieno[2,3-b]pyrrole-5-carboxamidinium hydrochloride (86%) as a beige powder. ¹H NMR (DMSO-*d*₆) δ (ppm): 8.61 (bs, 4 H), 7.23 (d, $J=4.9$ Hz, 1 H), 7.10 (d, $J=4.9$ Hz, 1 H), 7.04 (s, 1 H), 3.88 (s, 3 H). MS (ESI): m/z : 180 [M+H]⁺.




Example 1: 4-Methyl-5-(4-phenyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole
















A solution of 0.018 g (0.092 mmol) of 2-bromo-1-phenyl-ethanone (Sigma-Aldrich, Cat. No. 115835) in THF (0.05 mL) was added dropwise to 0.016 g (0.092 mmol) of 4-methylthieno[3,2-b]pyrrole-5-carboxamidinium acetate (Intermediate 11) suspended in 0.55 mL THF/water (4:1, v:v). The mixture was heated to reflux for 30 min. Then the solution was concentrated and the residue was diluted with water and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The obtained crude product was purified by flash chromatography on silica gel (hexane/EtOAc, 97:3, v:v to 7:3, v:v) and then triturated with Et₂O to give 0.013 g of *N'*-hydroxy-4-methylthieno[3,2-b]pyrrole-5-carboxamidinium (51%) as a white powder. ¹H NMR (CDCl₃) δ (ppm): 7.83–7.73 (m, 2 H), 7.46–7.38 (m, 2 H), 7.34 (s, 1 H), 7.32–7.28 (m, 1 H), 7.17 (d, $J=5.4$ Hz, 1 H), 6.96 (d, $J=5.4$ Hz, 1 H), 6.73 (s, 1 H), 4.18 (s, 3 H); MS (ESI): m/z : 280 [M+H]⁺.







The following compounds (see Table 1) were prepared starting from 4-methylthieno[3,2-b]pyrrole-5-carboxamide acetate (Intermediate 11: Examples 2-10, 13 and 16-17) 4-ethylthieno[3,2-b]pyrrole-5-carboxamide acetate (Intermediate 12: Examples 11-12) or 6-methylthieno[2,3-b]pyrrole-5-carboxamide hydrochloride (Intermediate 13: Examples 14-15) and the appropriate α -bromo ketone or α -chloro ketone according to the procedure described for Example 1.









Table 1

Ex.	Name	α -halogeno ketone	Structure	Analytical Data
2	5-(4-ethyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole	 (Sigma-Aldrich, Cat. No. 243299)		^1H NMR (DMSO- d_6) δ (ppm): 12.19–11.77 (bs, 1 H), 7.26 (d, $J=5.4$ Hz, 1 H), 7.15 (d, $J=5.4$ Hz, 1 H), 6.82 (bs, 1 H), 6.77 (s, 1 H), 4.08 (s, 3 H), 2.63–2.52 (m, 2 H), 1.21 (t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 232 $[\text{M}+\text{H}]^+$.
3	4-methyl-5-(4-methyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole	 (Sigma-Aldrich, Cat. No. 167479) http://www.sigmaaldrich.com/catalog/product/aldrich/167479?lang=it&region=IT		^1H NMR (DMSO- d_6) δ (ppm): 12.30 (bs, 1 H), 7.28 (d, $J=5.4$ Hz, 1 H), 7.16 (d, $J=4.9$ Hz, 1 H), 6.84 (bs, 1 H), 6.78 (s, 1 H), 4.06 (s, 3 H), 2.20 (s, 3 H); MS (ESI): m/z : 218 $[\text{M}+\text{H}]^+$.

4	5-(4-isopropyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole	 (WO 2012/049277)		¹ H NMR (CDCl ₃) δ (ppm): 7.14 (d, J=5.4 Hz, 1 H), 6.92 (d, J=5.4 Hz, 1 H), 6.81 (s, 1 H), 6.64 (s, 1 H), 4.06 (s, 3 H), 3.05-2.94 (m, 1 H), 1.35-1.29 (m, 6 H); MS (ESI): m/z: 246 [M+H] ⁺ .
5	5-(4-cyclopropyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole	 (Enamine, Cat. No. EN300-57214)		¹ H NMR (CDCl ₃) δ (ppm): 7.16-7.14 (d, J=5.4 Hz, 1 H), 6.95 (d, J=5.4 Hz, 1 H), 6.78 (s, 1 H), 6.59 (s, 1 H), 4.08 (s, 3 H), 1.92-1.82 (m, 1 H), 0.96-0.70 (m, 4 H); MS (ESI): m/z: 244 [M+H] ⁺ .
6	4-methyl-5-(4-propyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole	 (WO2014/86701)		¹ H NMR (CDCl ₃) δ (ppm): 7.14 (d, J=5.4 Hz, 1 H), 6.92 (d, J=5.4 Hz, 1 H), 6.82 (s, 1 H), 6.65 (s, 1 H), 4.07 (s, 3 H), 2.63 (t, J=7.4 Hz, 2 H), 1.70 (qd, J=7.4, 15.0 Hz, 2 H), 1.00 (t, J=7.4 Hz, 3 H); MS (ESI): m/z: 246 [M+H] ⁺ .
7	5-(4-cyclobutyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole	 (WO2014/86701)		¹ H NMR (CDCl ₃) δ (ppm): 7.14 (d, J=5.4 Hz, 1 H), 6.91 (d, J=5.4 Hz, 1 H), 6.86 (s, 1 H), 6.66 (s, 1 H), 4.06 (s, 3 H)


	methyl-thieno[3,2-b]pyrrole			H), 3.61–3.51 (m, 1 H), 2.43–2.33 (m, 2 H), 2.25–2.14 (m, 2 H), 2.10–1.88 (m, 2 H); MS (ESI): m/z: 258 [M+H] ⁺ .
8	4-methyl-5-(5-methyl-4-phenyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole	 (Sigma-Aldrich, Cat. No. 471607)		¹ H NMR (CDCl ₃) δ (ppm): 7.67–7.60 (m, 2 H), 7.46–7.39 (m, 2 H), 7.33–7.28 (m, 1 H), 7.15 (d, J=5.4 Hz, 1 H), 6.92 (d, J=5.4 Hz, 1 H), 6.71 (s, 1 H), 4.13 (s, 3 H), 2.47 (s, 3 H); MS (ESI): m/z: 294 [M+H] ⁺ .
9	5-(5-ethyl-4-phenyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole	 (Enamine, Cat. No. EN300-21953)		¹ H NMR (CDCl ₃) δ (ppm): 7.69–7.58 (m, 2 H), 7.47–7.40 (m, 2 H), 7.33–7.28 (m, 1 H), 7.15 (d, J=5.4 Hz, 1 H), 6.95 (d, J=5.4 Hz, 1 H), 6.68 (s, 1 H), 4.15 (s, 3 H), 2.88 (q, J=7.7 Hz, 2 H), 1.36 (t, J=7.6 Hz, 3 H); MS (ESI): m/z: 308 [M+H] ⁺ .
10	5-(5-isopropyl-4-phenyl-1H-imidazol-2-yl)-4-methyl-	 (Enamine, Cat. No. EN300-42834)		¹ H NMR (CDCl ₃) δ (ppm): 7.68–7.52 (m, 2 H), 7.47–7.39 (m, 2H), 7.35–7.29 (m, 1 H), 7.14 (d, J=5.4 Hz, 2 H), 6.94 (d, J=5.4 Hz, 2 H), 6.70 (s, 1 H), 4.13 (s, 3 H), 3.42–3.27

	thieno[3,2-b]pyrrole			(m, 1 H), 1.38 (d, $J=6.8$ Hz, 6 H)MS (ESI): m/z : 322 $[M+H]^+$.
11	4-ethyl-5-(4-ethyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole	 (Sigma-Aldrich, Cat. No. 243299)		^1H NMR (CDCl_3) δ (ppm): 7.15 (d, $J=5.4$ Hz, 1 H), 6.97 (d, $J=5.4$ Hz, 1 H), 6.82 (s, 1 H), 6.62 (s, 1 H), 4.63 (q, $J=7.2$ Hz, 2 H), 2.69 (q, $J=7.3$ Hz, 2 H), 1.38 (t, $J=7.2$ Hz, 3 H), 1.29 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 246 $[M+H]^+$.
12	5-(4-cyclobutyl-1H-imidazol-2-yl)-4-ethyl-thieno[3,2-b]pyrrole	 (WO2014/86701)		^1H NMR (CDCl_3) δ (ppm): 7.14 (d, $J=5.4$ Hz, 1 H), 6.96 (d, $J=5.4$ Hz, 1 H), 6.86 (s, 1 H), 6.62 (s, 1 H), 4.63 (q, $J=7.2$ Hz, 2 H), 3.60-3.50 (m, 1 H), 2.42-2.32 (m, 2 H), 2.25-2.11 (m, 2 H), 2.09-1.86 (m, 2 H), 1.38 (t, $J=7.2$ Hz, 3 H); MS (ESI): m/z : 272 $[M+H]^+$.
13	4-methyl-5-(4-phenyl-5-propyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole	 (Hyde, J. F. <i>et al.</i> J. Am. Chem. Soc, 1928, 50, 2287)		^1H NMR (CDCl_3) δ (ppm): = 7.70-7.58 (m, 2H), 7.46-7.40 (m, 2H), 7.34-7.28 (m, 1 H)7.15 (d, $J=5.4$ Hz, 1 H), 6.95 (d, $J=5.4$ Hz, 1 H), 6.66 (s, 1 H), 4.15 (s, 3 H), 2.82 (t, $J=7.3$ Hz, 2 H), 1.78 (qd, $J=7.3, 15.0$ Hz,

				2 H), 1.03 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 322 $[M+H]^+$.
14	5-(4-ethyl-1H-imidazol-2-yl)-6-methyl-thieno[2,3-b]pyrrole	 (Sigma-Aldrich, Cat. No. 243299)		^1H NMR (CDCl_3) δ (ppm): 6.97 (d, $J=5.4$ Hz, 1 H), 6.87 (d, $J=5.4$ Hz, 1 H), 6.80 (s, 1 H), 6.59 (s, 1 H), 4.05 (s, 3 H), 2.69 (q, $J=7.6$ Hz, 2 H), 1.30 (t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 232 $[M+H]^+$.
15	5-(5-ethyl-4-phenyl-1H-imidazol-2-yl)-6-methyl-thieno[2,3-b]pyrrole	 (Enamine, Cat. No. EN300-21953)		^1H NMR (CDCl_3) δ (ppm): = 7.67–7.28 (m, 5 H), 6.97 (d, $J=4.9$ Hz, 1 H), 6.87 (d, $J=4.9$ Hz, 1 H), 6.66 (s, 1 H), 4.13 (s, 3 H), 2.87 (q, $J=7.3$ Hz, 2 H), 1.36 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 308 $[M+H]^+$.
16	4-methyl-5-(4,5,6,7-tetrahydro-1H-benzimidazol-2-yl)thieno[3,2-b]pyrrole	 (Sigma-Aldrich, Cat. No. 690422)		^1H NMR (CDCl_3) δ (ppm): = 7.14 (d, $J=5.4$ Hz, 1 H), 6.93 (d, $J=5.4$ Hz, 1 H), 6.61 (s, 1 H), 4.08 (s, 3 H), 2.66–2.61 (m, 4 H), 1.86–1.81 (m, 4 H); MS (ESI): m/z : 258 $[M+H]^+$.
17	4-methyl-5-[4-(3,3,3-trifluoropropyl)-1H-imidazol-2-yl]thieno[3,2-b]pyrrole	 (FCH group,		^1H NMR (CDCl_3) δ (ppm): = 7.16 (d, $J = 5.4$ Hz, 1 H), 6.95 (d, $J = 5.4$ Hz, 1 H), 6.84 (s, 1 H), 6.64 (s, 1 H), 4.09 (s, 3

	imidazol-2-yl]thieno[3,2-b]pyrrole	Cat. No FCH2646621-1		H), 2.95 - 2.86 (m, 2 H), 2.62 - 2.46 (m, 2 H)
--	------------------------------------	-------------------------	--	---

Intermediate 14: *tert*-Butyl 4-[3-(2-bromoethoxy)phenoxy]piperidine-1-carboxylate



tert-Butyl 4-(3-hydroxyphenoxy)piperidine-1-carboxylate


0.64 g (3.0 mmol, 0.62 mL) of DIAD (Sigma-Aldrich, Cat. No. 225541) were added dropwise to a solution of 0.33 g (3.0 mmol) of resorcinol (Sigma-Aldrich, Cat. No. 398047), 0.42 g (2.0 mmol) of *tert*-butyl 4-hydroxypiperidine-1-carboxylate (Apollo, Cat. No. OR5404) and 0.80 g (3.0 mmol) of PPh₃ (Sigma-Aldrich, Cat. No. T84409) in 20 mL of dry THF at 0°C. The reaction mixture was allowed to reach RT and was stirred overnight. The solvent was then removed and the crude mixture was purified by flash chromatography on silica gel (hexane/acetone, 0% to 15% of acetone) to give 0.217 g of *tert*-butyl 4-(3-hydroxyphenoxy)piperidine-1-carboxylate (37%) as white solid. ¹H NMR (CDCl₃) δ (ppm): 7.16–7.11 (m, 1 H), 6.52–6.48 (m, 1 H), 6.46–6.41 (m, 2 H), 4.96 (bs, 1 H), 4.50–4.38 (m, 1 H), 3.75–3.65 (m, 2 H), 3.39–3.29 (m, 2 H), 1.97–1.86 (m, 2 H), 1.81–1.69 (m, 2 H), 1.48 (s, 9 H); MS (ESI): m/z: 238 [M-56+H]⁺.

tert-Butyl 4-[3-(2-bromoethoxy)phenoxy]piperidine-1-carboxylate

0.16 g (0.76 mmol, 0.158 mL) of DIAD (Sigma-Aldrich, Cat. No. 225541) were added dropwise to a solution comprising 0.150 g (0.51 mmol) of *tert*-butyl 4-(3-hydroxyphenoxy)piperidine-1-carboxylate, 0.100 g (0.76 mmol, 0.057 mL) of 2-bromoethanol (Sigma-Aldrich, Cat. No. B65586) and 0.203 g (0.76 mmol) of PPh₃ (Sigma-Aldrich, Cat. No. T84409) in 5 mL of dry THF at 0°C. The solution was allowed to reach RT and was stirred overnight. Then further 0.028 mL of 2-bromoethanol, 0.101 mg of PPh₃ and 0.079 mL of DIAD were added to the reaction mixture cooled down to 0°C

and the mixture was stirred at RT for further 24 h. The solvent was removed and the crude product was purified by flash chromatography on silica gel (hexane/acetone, 0% to 5% of acetone) to provide 0.105 g of *tert*-butyl 4-[3-(2-bromoethoxy)phenoxy]piperidine-1-carboxylate (51%) as colourless oil. ^1H NMR (CDCl_3) δ (ppm): 7.23–7.15 (m, 1 H), 6.58–6.47 (m, 3 H), 4.50–4.43 (m, 1 H), 4.28 (t, $J=6.4$ Hz, 2 H), 3.75–3.67 (m, 2 H), 3.64 (t, $J=6.4$ Hz, 2 H), 3.38–3.28 (m, 2 H), 1.96–1.86 (m, 2 H), 1.81–1.69 (m, 2 H), 1.48 (s, 9 H); MS (ESI): m/z : 344 $[\text{M}-56+\text{H}]^+$.

Intermediate 15: *tert*-Butyl 3-[[4-(2-bromoethoxy)phenoxy]methyl]pyrrolidine-1-carboxylate




tert-Butyl 3-[(4-hydroxyphenoxy)methyl]pyrrolidine-1-carboxylate

0.37 g of *tert*-butyl 3-[(4-hydroxyphenoxy)methyl]pyrrolidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 1, starting from 0.23 g (2.1 mmol) of hydroquinone (Sigma-Aldrich, Cat. No. H9003) and 0.22 g (1.0 mmol) of *tert*-butyl 3-(hydroxymethyl)pyrrolidine-1-carboxylate (Fluorochem, Cat. No. 048620) (yield: 24%). ^1H NMR (CDCl_3) δ (ppm): 6.82–6.73 (m, 4 H), 4.59 (s, 1 H), 3.94–3.80 (m, 2 H), 3.63–3.55 (m, 1 H), 3.53–3.44 (m, 1 H), 3.42–3.32 (m, 1 H), 3.24–3.14 (m, 1 H), 2.70–2.60 (m, 1 H), 2.12–2.00 (m, 1 H), 1.87–1.73 (m, 1 H), 1.47 (s, 9 H); MS (ESI): m/z : 238 $[\text{M}-56+\text{H}]^+$.

tert-Butyl 3-[[4-(2-bromoethoxy)phenoxy]methyl]pyrrolidine-1-carboxylate

0.053 g of *tert*-butyl 3-[[4-(2-bromoethoxy)phenoxy]methyl]pyrrolidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 2, starting from 0.070 g (0.25 mmol) *tert*-butyl 3-[(4-hydroxyphenoxy)methyl]pyrrolidine-1-carboxylate and 2-bromoethanol (yield: 54%). ^1H NMR (CDCl_3) δ (ppm): 6.92–6.79 (m, 4 H), 4.25 (t, $J=6.4$ Hz, 2 H), 3.97–3.81 (m, 2 H), 3.62 (t, $J=6.1$ Hz, 2 H), 3.60–3.56 (m, 1 H), 3.51–3.44 (m, 1 H), 3.41–3.32 (m, 1 H), 3.23–3.16 (m, 1 H), 2.70–2.61 (m, 1 H), 2.11–2.02 (m, 1 H), 1.85–1.74 (m, 1 H), 1.47 (s, 9 H); MS (ESI): m/z : 344 $[\text{M}-56+\text{H}]^+$.

Intermediate 16: *tert*-Butyl 4-[[4-(2-bromoethoxy)phenoxy]methyl]piperidine-1-carboxylate




tert-Butyl 4-[(4-hydroxyphenoxy)methyl]piperidine-1-carboxylate

0.28 g of *tert*-butyl 4-[(4-hydroxyphenoxy)methyl]pyrrolidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 1, starting from 0.50 g (4.5 mmol) of hydroquinone (Sigma-Aldrich, Cat. No. H9003) and 0.65 g (3.0 mmol) of *tert*-butyl 4-(hydroxymethyl)piperidine-1-carboxylate (Sigma-Aldrich, Cat. No. 556017) (yield: 31%). ¹H NMR (CDCl₃) δ (ppm): 6.83–6.70 (m, 4 H), 4.58 (bs, 1 H), 4.22–4.10 (m, 2 H), 3.75 (d, *J*=6.4 Hz, 2 H), 2.81–2.69 (m, 2 H), 2.00–1.88 (m, 1 H), 1.86–1.77 (m, 2 H), 1.48 (s, 9 H), 1.31–1.23 (m, 2 H); MS (ESI): *m/z*: 252 [M-56+H]⁺.

tert-Butyl 4-[[4-(2-bromoethoxy)phenoxy]methyl]piperidine-1-carboxylate

0.11 g of *tert*-butyl 4-[[4-(2-bromoethoxy)phenoxy]methyl]piperidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 2, starting from 0.28 g (0.90 mmol) of *tert*-butyl 4-[(4-hydroxyphenoxy)methyl]pyrrolidine-1-carboxylate and 2-bromoethanol (yield: 29%). ¹H NMR (CDCl₃) δ (ppm): 6.89–6.79 (m, 4 H), 4.25 (t, *J*=6.4 Hz, 2 H), 4.20–4.11 (m, 2 H), 3.76 (d, *J*=6.4 Hz, 2 H), 3.62 (t, *J*=6.4 Hz, 2 H), 2.81–2.70 (m, 2 H), 2.00–1.88 (m, 1 H), 1.87–1.77 (m, 2 H), 1.47 (s, 9 H), 1.32–1.23 (m, 2 H); MS (ESI): *m/z*: 358 [M-56+H]⁺.

Intermediate 17: *tert*-Butyl 4-[4-(2-bromoethoxy)phenoxy]piperidine-1-carboxylate



tert-Butyl 4-(4-benzyloxyphenoxy)piperidine-1-carboxylate

0.8 g of *tert*-butyl 4-(4-benzyloxyphenoxy)piperidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 1 starting from 0.60 g (3.0 mmol) of

4-benzyloxyphenol (Sigma-Aldrich, Cat. No. 158348) and 0.93 g (4.5 mmol) of *tert*-butyl 4-hydroxypiperidine-1-carboxylate (Apollo, Cat. No. OR5404) (Yield: 70%). ^1H NMR (CDCl_3) δ (ppm): 7.46–7.42 (m, 2 H), 7.42–7.37 (m, 2 H), 7.36–7.31 (m, 1 H), 6.93–6.84 (m, 4 H), 5.03 (s, 2 H), 4.38–4.28 (m, 1 H), 3.76–3.68 (m, 2 H), 3.35–3.24 (m, 2 H), 1.95–1.84 (m, 2 H), 1.78–1.67 (m, 2 H), 1.48 (s, 9 H); MS (ESI): m/z : 328 $[\text{M}-56+\text{H}]^+$.


tert-Butyl 4-(4-hydroxyphenoxy)piperidine-1-carboxylate

0.454 g (1.18 mmol) of *tert*-butyl 4-(4-benzyloxyphenoxy)piperidine-1-carboxylate dissolved in 47 mL of dry EtOH were hydrogenated in an H-cube apparatus using a 10% Pd/C cartridge at 25°C, at atmospheric pressure and with a flow of 0.5 mL/min for 5 h. The solution was then concentrated to provide 0.340 g of *tert*-butyl 4-(4-hydroxyphenoxy)piperidine-1-carboxylate (98%) as a white solid. ^1H NMR (CDCl_3) δ (ppm): 6.87–6.71 (m, 4 H), 4.58 (bs, 1 H), 4.36–4.27 (m, 1 H), 3.76–3.67 (m, 2 H), 3.37–3.24 (m, 2 H), 1.95–1.82 (m, 2 H), 1.78–1.66 (m, 2 H), 1.48 (s, 9 H); MS (ESI): m/z : 238 $[\text{M}-56+\text{H}]^+$.

tert-Butyl 4-[4-(2-bromoethoxy)phenoxy]piperidine-1-carboxylate

0.31 g of *tert*-Butyl 4-[4-(2-bromoethoxy)phenoxy]piperidine-1-carboxylate was obtained according to the procedure for Intermediate **14**, step 2, starting from 0.340 g (1.16) of *tert*-butyl 4-(4-hydroxyphenoxy)piperidine-1-carboxylate and 2-bromoethanol (yield: 66%). ^1H NMR (CDCl_3) δ (ppm): 6.92–6.80 (m, 4 H), 4.37–4.31 (m, 1 H), 4.25 (t, $J=6.1$ Hz, 2 H), 3.76–3.67 (m, 2 H), 3.63 (t, $J=6.4$ Hz, 2 H), 3.35–3.26 (m, 2 H), 1.95–1.85 (m, 2 H), 1.77–1.67 (m, 2 H), 1.48 (s, 9 H); MS (ESI): m/z : 344 $[\text{M}-56+\text{H}]^+$.

Intermediate 18: *tert*-Butyl 4-[4-(2-bromoethoxy)phenoxy]azepane-1-carboxylate



tert-Butyl 4-(4-benzyloxyphenoxy)azepane-1-carboxylate

0.74 g of *tert*-butyl 4-(4-benzyloxyphenoxy)azepane-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 1, starting from 0.60 g (3.0 mmol) of 4-benzyloxyphenol and 0.97 g (4.5 mmol) of *tert*-butyl 4-hydroxyazepane-1-carboxylate (Sigma-Aldrich, Cat. No. CDS009029) (yield: 62%). ^1H NMR (CDCl_3) δ (ppm): 7.46–7.42

(m, 2 H), 7.42–7.37 (m, 2 H), 7.36–7.31 (m, 1 H), 6.92–6.80 (m, 4 H), 5.02 (s, 2 H), 4.37–4.26 (m, 1 H), 3.65–3.23 (m, 4 H), 2.09–1.84 (m, 5 H), 1.70–1.59 (m, 1 H), 1.48 (s, 9 H); MS (ESI): m/z: 342 [M-56+H]⁺.


tert-Butyl 4-(4-hydroxyphenoxy)azepane-1-carboxylate

0.51 g of *tert*-butyl 4-(4-hydroxyphenoxy)azepane-1-carboxylate was prepared according to the procedure for Intermediate 17, step 2, starting from 0.71 g (1.8 mmol) of *tert*-butyl 4-(4-benzyloxyphenoxy)azepane-1-carboxylate (yield: 92%). ¹H NMR (CDCl₃) δ (ppm): 6.82–6.72 (m, 4 H), 4.58 (bs, 1 H), 4.36–4.25 (m, 1 H), 3.65–3.22 (m, 4 H), 2.12–1.83 (m, 5 H), 1.70–1.59 (m, 1 H), 1.48 (s, 9 H); MS (ESI): m/z: 252 [M-56+H]⁺.

tert-Butyl 4-[4-(2-bromoethoxy)phenoxy]azepane-1-carboxylate


0.22 g of *tert*-butyl 4-[4-(2-bromoethoxy)phenoxy]azepane-1-carboxylate was prepared according to the procedure for Intermediate 14, step 2, starting from 0.36 g (1.2 mmol) of *tert*-butyl 4-(4-hydroxyphenoxy)azepane-1-carboxylate and 2-bromoethanol (yield: 47%). ¹H NMR (CDCl₃) δ (ppm): 6.92–6.76 (m, 4 H), 4.38–4.29 (m, 1 H), 4.25 (t, *J*=6.4 Hz, 2 H), 3.67–3.22 (m, 6 H), 2.11–1.84 (m, 5 H), 1.70–1.59 (m, 1 H), 1.48 (s, 9 H); MS (ESI): m/z: 358 [M-56+H]⁺.

Intermediate 19: *tert*-Butyl 4-[4-(3-bromopropyl)phenoxy]piperidine-1-carboxylate




0.33 g of *tert*-butyl 4-[4-(3-bromopropyl)phenoxy]piperidine-1-carboxylate was prepared according to the procedure for Intermediate 14, step 1, starting from 0.22 g (1.0 mmol) of 4-(3-bromopropyl)phenol (US5204018) and 0.30 g (1.5 mmol) of *tert*-butyl 4-hydroxypiperidine-1-carboxylate (Apollo, Cat. No. OR5404) (yield: 82%). ¹H NMR (CDCl₃) δ (ppm): 7.17–7.08 (m, 2 H), 6.89–6.81 (m, 2 H), 4.49–4.38 (m, 1 H), 3.78–3.65 (m, 2 H), 3.40 (t, *J*=6.6 Hz, 2 H), 3.37–3.30 (m, 2 H), 2.73 (t, *J*=7.3 Hz, 2 H), 2.18–2.10 (m, 2 H), 1.95–1.87 (m, 2 H), 1.80–1.69 (m, 2 H), 1.47 (s, 9 H); MS (ESI): m/z: 342 [M-56+H]⁺.

Intermediate 19A: *tert*-Butyl 4-[4-(3-bromopropyl)phenoxy]piperidine-1-carboxylate




0.41 g of *tert*-butyl 4-[3-(3-bromopropyl)phenoxy]piperidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 1, starting from 0.28 g (1.3 mmol) of 3-(3-bromopropyl)phenol (Murphy, W. *et al.* J. Chem. Soc., Perkin Trans. 1: Org. Bio-Org. Chem. 1980, 7, 1567-1577) and 0.40 g (1.95 mmol) of *tert*-butyl 4-hydroxypiperidine-1-carboxylate (Apollo, Cat. No. OR5404) (yield: 79%). ^1H NMR (CDCl_3) δ (ppm): 7.25–7.18 (m, 1 H), 6.83–6.74 (m, 3 H), 4.51–4.42 (m, 1 H), 3.76–3.67 (m, 2 H), 3.41 (t, $J=6.4$ Hz, 2 H), 3.38–3.31 (m, 2 H), 2.75 (t, $J=7.3$ Hz, 2 H), 2.21–2.13 (m, 2 H), 1.97–1.87 (m, 2 H), 1.81–1.69 (m, 2 H), 1.48 (s, 9 H); MS (ESI): m/z : 342 $[\text{M}-56+\text{H}]^+$.

Intermediate 20: *tert*-Butyl 3-[[4-(3-bromopropyl)phenoxy]methyl]pyrrolidine-1-carboxylate



0.37 g of *tert*-butyl 3-[[4-(3-bromopropyl)phenoxy]methyl]pyrrolidine-1-carboxylate was prepared according to the procedure for Intermediate **15**, step 1, starting from 0.39 g (1.8 mmol) 4-(3-bromopropyl)phenol (US5204018) and 0.57 g (2.7 mmol) of *tert*-butyl 3-(hydroxymethyl)pyrrolidine-1-carboxylate (Fluorochem, Cat. No. 048620) (yield: 52%). ^1H NMR (CDCl_3) δ (ppm): 7.16–7.07 (m, 2 H), 6.87–6.78 (m, 2 H), 3.95–3.85 (m, 2 H), Part AB of ABX System: $V_A=3.6$, $V_B=3.21$, $J_{AB}=10.9$ Hz, $J_{AX}=7.6$ Hz, $J_{BX}=6.8$ Hz, 3.52–3.44 (m, 1 H), 3.42–3.33 (m, 3 H), 2.76–2.62 (m, 3 H), 2.17–2.03 (m, 3 H), 1.85–1.75 (m, 1 H), 1.47 (s, 9 H); MS (ESI): m/z : 342 $[\text{M}-56+\text{H}]^+$.

Intermediate 21: *tert*-Butyl 4-[3-(2-bromoethoxy)phenoxy]azepane-1-carboxylate




tert-Butyl 4-(3-hydroxyphenoxy)azepane-1-carboxylate

0.43 g of *tert*-butyl 4-(3-hydroxyphenoxy)azepane-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 1, starting from 0.50 g (4.5 mmol) of resorcinol (Sigma-Aldrich, Cat. No. 398047) and 0.65 g (3.0 mmol) of *tert*-butyl 4-hydroxyazepane-1-carboxylate (Sigma-Aldrich, Cat. No. CDS009029) (yield: 46%). ¹H NMR (CDCl₃+D₂O) δ (ppm): 7.19–7.05 (m, 1 H), 6.52–6.33 (m, 3 H), 4.48–4.32 (m, 1 H), 3.68–3.15 (m, 4 H), 2.16–1.81 (m, 5 H), 1.72–1.61 (m, 1 H), 1.48 (s, 9 H); MS (ESI): m/z: 252 [M-56+H]⁺.

tert-Butyl 4-[3-(2-bromoethoxy)phenoxy]azepane-1-carboxylate

0.13 g of *tert*-butyl 3-[[4-(2-bromoethoxy)phenoxy]methyl]pyrrolidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 2, starting from 0.19 g (0.61 mmol) of *tert*-butyl 4-(3-hydroxyphenoxy)azepane-1-carboxylate and 2-bromoethanol (yield: 51%). ¹H NMR (CDCl₃) δ (ppm): 7.23–7.14 (m, 1 H), 6.55–6.48 (m, 2 H), 6.48–6.44 (m, 1 H), 4.48–4.38 (m, 1 H), 4.28 (t, *J*=6.4 Hz, 2 H), 3.64 (t, *J*=6.1 Hz, 2 H), 3.60–3.20 (m, 4 H), 2.13–1.86 (m, 5 H), 1.71–1.60 (m, 1 H), 1.49 (s, 9 H); MS (ESI): m/z: 358 [M-56+H]⁺.


Intermediate 22: *tert*-butyl 4-[[4-(3-bromopropyl)phenoxy]methyl]piperidine-1-carboxylate



0.28 g of *tert*-butyl 4-[[4-(3-bromopropyl)phenoxy]methyl]piperidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 1, starting from 0.20 g (0.91 mmol) of 4-(3-bromopropyl)phenol (US5204018) and 0.29 g (1.4 mmol) of *tert*-butyl 4-


(hydroxymethyl)piperidine-1-carboxylate (Sigma-Aldrich, Cat. No. 556017) (yield: 76%).
 ^1H NMR (CDCl_3) δ (ppm): 7.17–7.07 (m, 2 H), 6.87–6.76 (m, 2 H), 4.21–4.11 (m, 2 H), 3.79 (d, $J=6.4$ Hz, 2 H), 3.39 (t, $J=6.6$ Hz, 2 H), 2.83–2.69 (m, 4 H), 2.19–2.09 (m, 2 H), 2.01–1.90 (m, 1 H), 1.88–1.76 (m, 2 H), 1.47 (s, 9 H), 1.34–1.19 (m, 2 H); MS (ESI): m/z : 356 $[\text{M}-56+\text{H}]^+$.

Intermediate 23: *tert*-butyl 4-[4-(3-bromopropyl)phenoxy]azepane-1-carboxylate



0.28 g of *tert*-butyl 4-[4-(3-bromopropyl)phenoxy]azepane-1-carboxylate was prepared according to the procedure for Intermediate 14, step 1, starting from 0.20 g (0.91 mmol) of 4-(3-bromopropyl)phenol (US5204018) and 0.29 g (1.4 mmol) of *tert*-butyl 4-hydroxyazepane-1-carboxylate (Intermediate 18, Step 2) (yield: 76%). ^1H NMR (CDCl_3) δ (ppm): 7.14–7.07 (m, 2 H), 6.85–6.77 (m, 2 H), 4.45–4.38 (m, 1 H), 3.65–3.24 (m, 6 H), 2.72 (t, $J=7.3$ Hz, 2 H), 2.18–2.10 (m, 2 H), 2.10–2.02 (m, 1 H), 1.99–1.85 (m, 4 H), 1.70–1.59 (m, 1 H), 1.48 (s, 9 H); MS (ESI): m/z : 356 $[\text{M}-56+\text{H}]^+$.

Intermediate 24: *tert*-butyl 4-[[4-(3-bromopropyl)-2-[(1-*tert*-butoxycarbonyl-4-piperidyl)methoxy]phenoxy]methyl]piperidine-1-carboxylate




tert-butyl 4-[[2-[(1-*tert*-butoxycarbonyl-4-piperidyl)methoxy]-4-(3-hydroxypropyl)phenoxy]methyl]piperidine-1-carboxylate


0.079 g (0.47 mmol) of 4-(3-hydroxypropyl)benzene-1,2-diol (Yang, J. *et al.* Biorg. Med. Chem Lett. 2014, 24, 2680-2684), 0.03 g (0.2 mmol) of NaI and 0.61 g (1.9 mmol) of Cs₂CO₃ were suspended in 1.5 mL dry DMF under nitrogen atmosphere. 0.46 g (1.6 mmol) of *tert*-butyl 4-(bromomethyl)piperidine-1-carboxylate (Sigma-Aldrich, Cat. No. 796719) in 0.8 mL dry DMF was added and the suspension was stirred at 80 °C for 7 h. The mixture was cooled down to RT and a further portion of NaI (0.014 g, 0.09 mmol), Cs₂CO₃ (0.23 g, 0.72 mmol) and *tert*-butyl 4-(bromomethyl)piperidine-1-carboxylate (0.2 g, 0.72 mmol) were added. The resulting mixture was heated at 80 °C overnight, then cooled down to RT and the product was extracted with EtOAc, dried over Na₂SO₄ and purified by column chromatography over silica gel (eluent: hexane/acetone, 0% to 18% of acetone) providing 148 mg (56%) of *tert*-butyl 4-[[2-[(1-*tert*-butoxycarbonyl-4-piperidyl)methoxy]-4-(3-hydroxypropyl)phenoxy]methyl]piperidine-1-carboxylate as a pale yellow oil. ¹H NMR (CDCl₃) δ (ppm): 6.85–6.69 (m, 3 H), 4.22–4.08 (m, 4 H), 3.86–3.78 (m, 4 H), 3.69 (t, *J*=7.3 Hz, 2 H), 2.82–2.71 (m, 4 H), 2.65 (t, *J*=7.3 Hz, 2 H), 2.06–1.94 (m, 2 H), 1.92–1.78 (m, 6 H), 1.48 (s, 9 H), 1.47 (s, 9 H), 1.33–1.19 (m, 4 H); MS (ESI): *m/z*: 585 [M+Na]⁺.

tert-butyl 4-[[4-(3-bromopropyl)-2-[(1-*tert*-butoxycarbonyl-4-piperidyl)methoxy]phenoxy]methyl]piperidine-1-carboxylate

A solution of 0.10 g (0.31 mmol) CBr₄ in 0.75 mL CH₂Cl₂ was added dropwise at -18 °C to a solution of 0.144 g (0.256 mmol) of *tert*-butyl 4-[[2-[(1-*tert*-butoxycarbonyl-4-piperidyl)methoxy]-4-(3-hydroxypropyl)phenoxy]methyl]piperidine-1-carboxylate and 0.081 g (0.31 mmol) of PPh₃ in 1.75 mL CH₂Cl₂. The reaction mixture was stirred at RT for 4 h. Then, the solution was concentrated and the residue was purified by column chromatography eluent: hexane/acetone, 0% to 7% of acetone) affording 118 mg (74%) of *tert*-butyl 4-[[4-(3-bromopropyl)-2-[(1-*tert*-butoxycarbonyl-4-piperidyl)methoxy]phenoxy]methyl]piperidine-1-carboxylate as a pale yellow oil. ¹H NMR (CDCl₃) δ (ppm): 6.85–6.67 (m, 3 H), 4.24–4.05 (m, 4 H), 3.87–3.75 (m, 4 H), 3.39 (t, *J*=6.6 Hz, 2 H), 2.83–2.63 (m, 6 H), 2.19–2.10 (m, 2 H), 2.05–1.93 (m, 2 H), 1.88–1.79 (m, 4 H), 1.48 (s, 9 H), 1.47 (s, 9 H), 1.32–1.18 (m, 4 H); MS (ESI): *m/z*: 647 [M+Na]⁺.

Intermediate 25: *tert*-Butyl 4-[4-(2-bromoethyl)phenoxy]piperidine-1-carboxylate


0.56 g of *tert*-butyl 4-[4-(2-bromoethyl)phenoxy]piperidine-1-carboxylate was prepared according to the procedure for Intermediate **14**, step 1, starting from 0.40 g (2.0 mmol) of 4-(2-bromoethyl)phenol (Fluorochem, Cat. No. 233801) and 0.61 g (3.0 mmol) of *tert*-butyl 4-hydroxypiperidine-1-carboxylate (Apollo, Cat. No. OR5404) (yield: 73%). ¹H NMR (CDCl₃) δ (ppm): 7.18–7.08 (m, 2 H), 6.91–6.81 (m, 2 H), 4.50–4.37 (m, 1 H), 3.76–3.65 (m, 2 H), 3.54 (t, *J*=7.6 Hz, 2 H), 3.39–3.28 (m, 2 H), 3.11 (t, *J*=7.6 Hz, 2 H), 1.99–1.85 (m, 2 H), 1.80–1.68 (m, 2 H), 1.48 (s, 9 H); MS (ESI): *m/z*: 328 [M-56+H]⁺.

Intermediate 26: *tert*-Butyl 4-[3-[2-[4-ethyl-2-(4-methylthieno[3,2-*b*]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate

0.03 g (0.13 mmol) of 5-(4-ethyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-*b*]pyrrole (Example 2) were added to a suspension of 0.0060 g (0.16 mmol) of NaH (60% dispersion in mineral oil) in dry DMA (0.2 mL) cooled down to -15°C. The mixture was stirred at a temperature ranging from -15°C to 0°C for 1 h, then a solution of 0.065 g of *tert*-butyl 4-[3-(2-bromoethoxy)phenoxy]piperidine-1-carboxylate (Intermediate **14**) in 0.5 mL of dry DMA was added at 0°C and the mixture was stirred overnight at RT. The reaction was stopped by adding water and the product was extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered and evaporated to dryness. The crude product was purified by flash chromatography on silica gel (hexane/acetone, 94:6% →

60:40, v:v) to afford 0.043 g of *tert*-butyl 4-[3-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate (Intermediate **26**, yield: 60%) as white foam. ¹H NMR (DMSO-*d*₆) δ (ppm): 7.32 (d, *J*=5.4 Hz, 1 H), 7.18 (d, *J*=5.4 Hz, 1 H), 7.16–7.07 (m, 2 H), 6.67 (s, 1 H), 6.57–6.39 (m, 3 H), 4.54–4.45 (m, 1 H), 4.37–4.20 (m, 4 H), 3.76 (s, 3 H), 3.68–3.55 (m, 2 H), 3.21–3.09 (m, 2 H), 2.51 (q, *J*=7.3 Hz, 2H), 1.91–1.78 (m, 2 H), 1.52–1.42 (m, 2 H), 1.39 (s, 9 H), 1.18 (t, *J*=7.3 Hz, 3 H); MS (ESI): *m/z*: 551 [M+H]⁺.



Intermediate 27: *tert*-Butyl 4-[3-[2-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate





Purification of the crude Intermediate **26** by flash chromatography (hexane/acetone, from 94:6% to 60:40, v:v) provided also 0.0092 g of *tert*-butyl 4-[3-[2-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate (Intermediate **27**, yield: 13%) as pale yellow foam. ¹H NMR (DMSO-*d*₆) δ (ppm): 7.32 (d, *J*=5.4 Hz, 1 H), 7.19 (d, *J*=5.4 Hz, 1 H), 6.86 (s, 1 H), 6.72 (s, 1 H), 6.55–6.30 (m, 3 H), 4.52–4.43 (m, 1 H), 4.39–4.33 (m, 2 H), 4.12–4.07 (m, 2 H), 3.69 (s, 3 H), 3.64–3.56 (m, 2 H), 3.21–3.05 (m, 2 H), 2.75–2.67 (m, 2 H), 1.87–1.77 (m, 2 H), 1.52–1.43 (m, 2 H), 1.39 (s, 9 H), 1.30–1.24 (m, 3 H); MS (ESI): *m/z*: 551 [M+H]⁺.



The following intermediates (see Table 2) were prepared starting from the appropriate mono or bi-substituted 5-(1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrroles (Examples **1**, **2**, **6**, **7**, **9**, **10**, **13** for Intermediates **28-46** and **48-85**) or from 5-(4-ethyl-1H-imidazol-2-yl)-6-methyl-thieno[2,3-b]pyrrole (Example **14** for Intermediate **47**) and the appropriate alkyl-bromides (Intermediates **14-24**) according to the procedure for Intermediates **26** and **27**.



Table 2: BOC Intermediates

Int.	Name	Structure	Analytical Data
28	<i>tert</i> -butyl 4-[4-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		^1H NMR (CDCl_3) δ (ppm): 7.17 (d, $J=5.4$ Hz, 1 H), 7.97(d, $J=5.4$ Hz, 1 H), 6.94 (s, 1H), 6.86–6.72 (m, 4 H), 6.59 (s, 1 H), 4.43–4.27 (m, 3 H), 4.20-4.13 (m, 2 H), 3.85 (s, 3 H), 3.76–3.25 (m, 4 H), 2.79–2.64 (m, 2 H), 1.94–1.65 (m, 4 H), 1.48 (s, 9 H), 1.34–1.22 (m, 3 H); MS (ESI): m/z : 551 $[\text{M}+\text{H}]^+$.
29	<i>tert</i> -butyl 4-[4-[2-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		^1H NMR (CDCl_3) δ (ppm): 7.20 (d, $J=5.4$ Hz, 1H), 7.05 (s, 1 H), 6.98 (d, $J=5.4$ Hz, 1H), 6.81–6.58 (m, 5 H), 4.41 (t, $J=5.7$, 2H), 4.35-4.27 (m, 1 H), 4.05 (t, $J=5.7$, 2H), 3.80 (s, 3 H), 3.74–3.25 (m, 4 H), 2.76 (q, $J=7.6$, 2 H), 1.92–1.65 (m, 4 H), 1.47 (s, 9 H), 1.40 (t, $J=7.6$, 3 H); MS (ESI): m/z : 551 $[\text{M}+\text{H}]^+$.



30	<i>tert</i>-butyl 3-[[4-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.17 (d, <i>J</i>=5.4 Hz, 1H), 6.97 (d, <i>J</i>=5.4 Hz, 1H), 6.94 (s, 1 H), 6.84–6.69 (m, 4 H), 6.58 (s, 1 H), 4.37 (t, <i>J</i>=5.4 Hz, 2H), 4.15 (t, <i>J</i>=5.4 Hz, 2H), 3.83 (s, 3 H), 3.94–3.78 (m, 2 H), 3.66–3.09 (m, 4 H), 2.75–2.60 (m, 3 H), 2.12–1.69 (m, 2 H), 1.47 (s, 9 H), 1.29 (t, <i>J</i>=7.6, 3H); MS (ESI): <i>m/z</i>: 551 [M+H]⁺.
31	<i>tert</i>-butyl 4-[[4-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]methyl]piperidine-1-carboxylate		¹H NMR (DMSO-<i>d</i>₆) δ (ppm): 7.32 (d, <i>J</i>=4.9 Hz, 1 H), 7.18 (d, <i>J</i>=5.4 Hz, 1 H), 7.14 (s, 1 H), 6.82–6.73 (m, 4 H), 6.67 (s, 1 H), 4.31 (t, <i>J</i>=5.4 Hz, 2 H), 4.18 (t, <i>J</i>=5.4 Hz, 2 H), 4.03–3.88 (m, 2 H), 3.77 (s, 3 H), 3.73 (d, <i>J</i>=6.4 Hz, 2 H), 2.71 (bs, 2 H), 2.52 (q, <i>J</i>=7.6 Hz, 2 H), 1.93–1.80 (m, 1 H), 1.76–1.65 (m, 2 H), 1.39 (s, 9 H), 1.18 (t, <i>J</i>=7.6 Hz, 3 H), 1.16–1.06 (m, 2 H); MS (ESI): <i>m/z</i>: 565 [M+H]⁺.



32	<i>tert</i>-butyl 4-[[4-[2-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]methyl]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.37 (d, <i>J</i>=5.4 Hz, 1 H), 7.21 (d, <i>J</i>=5.4 Hz, 1 H), 7.03 (s, 1 H), 6.80 (s, 1 H), 6.76–6.62 (m, 4 H), 4.39 (t, <i>J</i>=5.4 Hz, 2 H), 4.06 (t, <i>J</i>=5.4 Hz, 2 H), 3.94 (bs, 2 H), 3.73–3.66 (m, 5 H), 2.77–2.64 (m, 4 H), 1.89–1.78 (m, 1 H), 1.74–1.66 (m, 2 H), 1.39 (s, 9 H), 1.28 (t, <i>J</i>=7.3 Hz, 3 H), 1.16–1.04 (m, 2 H); MS (ESI): <i>m/z</i>: 565 [M+H]⁺.
33	<i>tert</i>-butyl 4-[4-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.18 (d, <i>J</i>=5.4 Hz, 1 H), 7.14 (s, 1 H), 6.82–6.74 (m, 4 H), 6.67 (s, 1 H), 4.38–4.27 (m, 3 H), 4.18 (t, <i>J</i>=5.4 Hz, 2 H), 3.77 (s, 3 H), 3.42–3.20 (m, 4 H), 2.53 (q, <i>J</i>=7.4 Hz, 2 H), 1.99–1.87 (m, 1 H), 1.84–1.63 (m, 4 H), 1.60–1.50 (m, 1 H), 1.40 (s, 9 H), 1.18 (t, <i>J</i>=7.6 Hz, 3 H); MS (ESI): <i>m/z</i>: 566 [M+H]⁺.



34	<i>tert</i>-butyl 4-[4-[2-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.18 (d, <i>J</i>=5.4 Hz, 1 H), 6.86 (s, 1 H), 6.77–6.63 (m, 5 H), 4.38–4.26 (m, 3 H), 4.05 (t, <i>J</i>=5.4 Hz, 2 H), 3.69 (s, 3 H), 3.30–3.19 (m, 4 H), 2.71 (q, <i>J</i>=7.4 Hz, 2 H), 1.98–1.84 (m, 1 H), 1.83–1.61 (m, 4 H), 1.60–1.48 (m, 1 H), 1.40 (s, 9 H), 1.27 (t, <i>J</i>=7.4 Hz, 3 H); MS (ESI): <i>m/z</i>: 566 [M+H]⁺.
35	<i>tert</i>-butyl 4-[4-[3-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.07 (s, 1 H), 7.01–6.97 (m, 2 H), 6.81–6.75 (m, 2 H), 6.41 (s, 1 H), 4.48–4.39 (m, 1 H), 3.94 (t, <i>J</i>=7.3 Hz, 2 H), 3.78 (s, 3 H), 3.69–3.59 (m, 2 H), 3.22–3.07 (m, 2 H), 2.53 (q, <i>J</i>=7.3 Hz, 2 H), 2.44 (t, <i>J</i>=7.3 Hz, 2 H), 2.00–1.92 (m, 2 H), 1.89–1.80 (m, 2 H), 1.51–1.42 (m, 2 H), 1.40 (s, 9 H), 1.18 (t, <i>J</i>=7.6 Hz, 3 H); MS (ESI): <i>m/z</i>: 549 [M+H]⁺.



36	<i>tert</i>-butyl 4-[4-[3-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.02–6.95 (m, 2 H), 6.84 (s, 1 H), 6.81–6.75 (m, 2 H), 6.46 (s, 1 H), 4.49–4.40 (m, 1 H), 3.92 (t, <i>J</i>=7.8 Hz, 2 H), 3.72 (s, 3 H), 3.68–3.59 (m, 2 H), 3.21–3.08 (m, 2 H), 2.57 (q, <i>J</i>=7.3 Hz, 2 H), 2.43 (t, <i>J</i>=7.3 Hz, 2 H), 1.90–1.78 (m, 4 H), 1.52–1.42 (m, 2 H), 1.40 (s, 9 H), 1.23 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 549 [M+H]⁺.
37	<i>tert</i>-butyl 4-[4-[2-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.33 (d, <i>J</i>=5.4 Hz, 1 H), 7.21–7.16 (m, 2 H), 6.88–6.74 (m, 4 H), 6.68 (s, 1 H), 4.42–4.34 (m, 1 H), 4.31 (t, <i>J</i>=5.3 Hz, 2 H), 4.19 (t, <i>J</i>=5.3 Hz, 2 H), 3.78 (s, 3 H), 3.66–3.57 (m, 2 H), 3.47–3.37 (m, 1 H), 3.20–3.05 (m, 2 H), 2.25–2.09 (m, 4 H), 1.99–1.87 (m, 1 H), 1.86–1.76 (m, 3 H), 1.50–1.42 (m, 2 H), 1.39 (s, 9 H); MS (ESI): <i>m/z</i>: 577 [M+H]⁺.



38	<i>tert</i>-butyl 4-[4-[2-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.33 (d, <i>J</i>=5.4 Hz, 1 H), 7.19 (d, <i>J</i>=5.4 Hz, 1 H), 6.98 (s, 1 H), 6.82–6.76 (m, 2 H), 6.72 (s, 1 H), 6.68–6.61 (m, 2 H), 4.38–4.32 (m, 1 H), 4.28 (t, <i>J</i>=5.4 Hz, 2 H), 3.99 (t, <i>J</i>=5.8 Hz, 2 H), 3.71–3.57 (m, 6 H), 3.19–3.08 (m, 2 H), 2.39–2.30 (m, 2 H), 2.19–2.09 (m, 2 H), 2.05–1.94 (m, 1 H), 1.90–1.76 (m, 3 H), 1.50–1.36 (m, 11 H); MS (ESI): <i>m/z</i>: 577 [M+H]⁺.
39	<i>tert</i>-butyl 4-[4-[2-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-phenyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.72–7.66 (m, 2 H), 7.46–7.38 (m, 2 H), 7.35 (d, <i>J</i>=5.3 Hz, 1 H), 7.28–7.23 (m, 1 H), 7.21 (d, <i>J</i>=4.9 Hz, 1 H), 6.82–6.76 (m, 3 H), 6.69–6.64 (m, 2 H), 4.46 (t, <i>J</i>=5.4 Hz, 2 H), 4.37–4.29 (m, 1 H), 4.08 (t, <i>J</i>=5.4 Hz, 2 H), 3.77 (s, 3 H), 3.66–3.56 (m, 2 H), 3.19–3.05 (m, 2 H), 2.97 (q, <i>J</i>=7.3 Hz, 2 H), 1.85–1.74 (m, 2 H), 1.49–1.37 (m, 11 H), 1.26 (t, <i>J</i>=7.3



			Hz, 3 H); MS (ESI): m/z: 627 [M+H] ⁺ .
40	<i>tert</i> -butyl 4-[3-[2-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-phenyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹ H NMR (DMSO-d ₆) δ (ppm): 7.73–7.65 (m, 2 H), 7.45–7.39 (m, 2 H), 7.35 (d, <i>J</i> =5.4 Hz, 1 H), 7.29–7.23 (m, 1 H), 7.21 (d, <i>J</i> =5.4 Hz, 1 H), 7.09–7.04 (m, 1 H), 6.81 (s, 1 H), 6.56–6.49 (m, 1 H), 6.35–6.30 (m, 2 H), 4.51–4.41 (m, 3 H), 4.13 (t, <i>J</i> =5.4 Hz, 2 H), 3.77 (s, 3 H), 3.63–3.54 (m, 2 H), 3.20–3.08 (m, 2 H), 2.96 (q, <i>J</i> =7.3 Hz, 2 H), 1.86–1.76 (m, 2 H), 1.51–1.36 (m, 11 H), 1.26 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): m/z: 627 [M+H] ⁺ .
41	<i>tert</i> -butyl 4-[3-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-phenyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹ H NMR (DMSO-d ₆) δ (ppm): 7.55–7.41 (m, 5 H), 7.34 (d, <i>J</i> =5.4 Hz, 1 H), 7.20 (d, <i>J</i> =5.4 Hz, 1 H), 7.02–6.95 (m, 1 H), 6.77 (s, 1 H), 6.49–6.43 (m, 1 H), 6.15–6.12 (m, 1 H), 6.11–6.07 (m, 1 H), 4.47–4.35 (m, 3 H), 3.79 (s, 3 H), 3.74 (t, <i>J</i> =5.8 Hz, 2 H), 3.64–3.54 (m, 2 H), 3.20–3.08 (m, 2 H), 2.46 (q,



			$J=7.3$ Hz, 2 H), 1.86–1.76 (m, 2 H), 1.49–1.37 (m, 11 H), 1.13 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 627 $[M+H]^+$.
42	<i>tert</i> -butyl 4-[3-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.32 (d, $J=5.4$ Hz, 1 H), 7.19 (d, $J=5.4$ Hz, 1 H), 7.15–7.09 (m, 2 H), 6.67 (s, 1 H), 6.50–6.46 (m, 1 H), 6.45–6.41 (m, 1 H), 6.41–6.37 (m, 1 H), 4.50–4.40 (m, 1 H), 4.33 (t, $J=5.4$ Hz, 2 H), 4.22 (t, $J=5.8$ Hz, 2 H), 3.77 (s, 3 H), 3.42–3.34 (m, 4 H), 2.53 (q, $J=7.3$ Hz, 2 H), 2.00–1.90 (m, 1 H), 1.83–1.65 (m, 4 H), 1.63–1.50 (m, 1 H), 1.40 (s, 9 H), 1.18 (t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 565 $[M+H]^+$.
43	<i>tert</i> -butyl 4-[3-[2-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.32 (d, $J=4.9$ Hz, 1 H), 7.18 (d, $J=5.4$ Hz, 1 H), 7.11–7.03 (m, 1 H), 6.86 (s, 1 H), 6.71 (s, 1 H), 6.49–6.42 (m, 1 H), 6.35–6.31 (m, 1 H), 6.31–6.29 (m, 1 H), 4.46–4.40 (m, 1 H), 4.37 (t, $J=5.4$ Hz, 2 H), 4.08 (t, $J=5.3$ Hz, 2 H),



			3.69 (s, 3 H), 3.42–3.34 (m, 4 H), 2.71 (q, $J=7.3$ Hz, 2 H), 2.02–1.88 (m, 1 H), 1.83–1.64 (m, 4 H), 1.61–1.50 (m, 1 H), 1.39 (s, 9 H), 1.27 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 565 $[M+H]^+$.
44	<i>tert</i> -butyl 4-[4-[2-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.33 (d, $J=4.9$ Hz, 1 H), 7.22–7.12 (m, 2 H), 6.83–6.73 (m, 4 H), 6.68 (s, 1 H), 4.38–4.26 (m, 3 H), 4.19 (t, $J=5.4$ Hz, 2 H), 3.78 (s, 3 H), 3.46–3.35 (m, 3 H), 2.27–2.09 (m, 4 H), 1.99–1.49 (m, 10 H), 1.40 (s, 9 H); MS (ESI): m/z : 591 $[M+H]^+$.
45	<i>tert</i> -butyl 4-[4-[2-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.32 (d, $J=5.4$ Hz, 1 H), 7.18 (d, $J=5.4$ Hz, 1 H), 6.98 (s, 1 H), 6.77–6.60 (m, 5 H), 4.37–4.23 (m, 3 H), 4.00 (t, $J=5.3$ Hz, 2 H), 3.75–3.59 (m, 4 H), 3.42–3.36 (m, 2 H), 2.41–2.29 (m, 2 H), 2.20–1.49 (m, 12 H), 1.40 (s, 9 H); MS (ESI): m/z : 591 $[M+H]^+$.



46	<i>tert</i>-butyl 3-[[4-[3-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.29 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=4.9 Hz, 1 H), 7.06 (s, 1 H), 7.03–6.98 (m, 2 H), 6.78–6.73 (m, 2 H), 6.34 (s, 1 H), 3.95–3.81 (m, 4 H), 3.78 (s, 3 H), 3.48–3.35 (m, 2 H), 3.29–3.18 (m, 1 H), 3.11–3.03 (m, 1 H), 2.62–2.52 (m, 3 H), 2.48–2.43 (m, 2 H), 2.03–1.91 (m, 3 H), 1.76–1.61 (m, 1 H), 1.39 (s, 9 H), 1.18 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 549 [M+H]⁺.
47	<i>tert</i>-butyl 4-[4-[2-[4-ethyl-2-(6-methylthieno[2,3-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.14 (s, 1 H), 7.082 (d, <i>J</i>=7.04 Hz, 1 H), 7.45 (d, <i>J</i>=7.04 Hz, 1 H), 6.89–6.74 (m, 4 H), 6.63 (s, 1 H), 4.41–4.35 (m, 1 H), 4.31 (t, <i>J</i>=5.4 Hz, 2 H), 4.19 (t, <i>J</i>=5.4 Hz, 2 H), 3.72 (s, 3 H), 3.66–3.56 (m, 2 H), 3.21–3.05 (m, 2 H), 2.49 (q, <i>J</i>=7.3 Hz, 2 H), 1.88–1.77 (m, 2 H), 1.51–1.42 (m, 2 H), 1.39 (s, 9 H), 1.18 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 551 [M+H]⁺.



48	<i>tert</i>-butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-phenyl-4-propyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.55–7.41 (m, 5 H), 7.34 (d, <i>J</i>=5.4 Hz, 1 H), 7.20 (d, <i>J</i>=5.4 Hz, 1 H), 6.76 (s, 1 H), 6.71 (d, <i>J</i>=9.3 Hz, 2 H), 6.43 (d, <i>J</i>=9.3 Hz, 2 H), 4.35 (t, <i>J</i>=5.9 Hz, 2 H), 4.33–4.27 (m, 1 H), 3.78 (s, 3 H), 3.69 (t, <i>J</i>=5.9 Hz, 2 H), 3.64–3.55 (m, 2 H), 3.21–3.04 (m, 2 H), 2.42 (t, <i>J</i>=7.6 Hz, 2 H), 1.85–1.74 (m, 2 H), 1.64–1.53 (m, 2 H), 1.47–1.35 (m, 11 H), 0.84 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 641 [M+H]⁺.
49	<i>tert</i>-butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-phenyl-5-propyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.71–7.63 (m, 2 H), 7.45–7.38 (m, 2 H), 7.35 (d, <i>J</i>=5.4 Hz, 1 H), 7.27–7.23 (m, 1 H), 7.21 (d, <i>J</i>=5.4 Hz, 1 H), 6.83–6.74 (m, 3 H), 6.70–6.63 (m, 2 H), 4.45 (t, <i>J</i>=5.4 Hz, 2 H), 4.37–4.30 (m, 1 H), 4.07 (t, <i>J</i>=5.4 Hz, 2 H), 3.77 (s, 3 H), 3.65–3.55 (m, 2 H), 3.19–3.05 (m, 2 H), 2.93–2.84 (m, 2 H), 1.84–1.75 (m, 2 H), 1.67–1.58 (m, 2 H), 1.48–



			1.40 (m, 2 H), 1.38 (s, 9 H), 0.98 (t, $J=7.1$ Hz, 3 H); MS (ESI): m/z : 641 $[M+H]^+$.
50	<i>tert</i> -butyl 4-[[4-[3-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]piperidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.31 (d, $J=5.4$ Hz, 1 H), 7.17 (d, $J=5.4$ Hz, 1 H), 7.07 (s, 1 H), 7.03–6.98 (m, 2 H), 6.79–6.71 (m, 2 H), 6.38 (s, 1 H), 4.02–3.89 (m, 4 H), 3.78 (s, 3 H), 3.75 (d, $J=6.4$ Hz, 2 H), 2.84–2.64 (m, 2 H), 2.53 (q, $J=7.3$ Hz, 2 H), 2.45 (t, $J=7.3$ Hz, 2 H), 2.01–1.92 (m, 2 H), 1.92–1.81 (m, 1 H), 1.77–1.69 (m, 2 H), 1.40 (s, 9 H), 1.19 (t, $J=7.3$ Hz, 3 H), 1.16–1.11 (m, 2 H); MS (ESI): m/z : 563 $[M+H]^+$.
51	<i>tert</i> -butyl 4-[[4-[3-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]piperidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.31 (d, $J=5.4$ Hz, 1 H), 7.16 (d, $J=5.4$ Hz, 1 H), 7.03–6.97 (m, 2 H), 6.83 (s, 1 H), 6.77–6.71 (m, 2 H), 6.40 (s, 1 H), 4.03–3.89 (m, 4 H), 3.76 (d, $J=6.4$ Hz, 2 H), 3.72 (s, 3 H), 2.83–2.65 (m, 2 H), 2.58 (q, $J=7.3$ Hz, 2 H), 2.45 (t,




			$J=7.3$ Hz, 2 H), 1.93–1.79 (m, 3 H), 1.77–1.70 (m, 2 H), 1.40 (s, 9 H), 1.27–1.07 (m, 5 H); MS (ESI): m/z : 563 $[M+H]^+$.
52	<i>tert</i> -butyl 4-[4-[3-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]azepane-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.31 (d, $J=5.4$ Hz, 1 H), 7.17 (d, $J=5.4$ Hz, 1 H), 7.07 (s, 1 H), 7.01–6.96 (m, 2 H), 6.75–6.69 (m, 2 H), 6.41 (s, 1 H), 4.43–4.35 (m, 1 H), 3.94 (t, $J=7.3$ Hz, 2 H), 3.78 (s, 3 H), 3.43–3.23 (m, 4 H), 2.51 (q, $J=7.3$ Hz, 2 H), 2.44 (t, $J=7.3$ Hz, 2 H), 1.99–1.90 (m, 2 H), 1.83–1.50 (m, 6 H), 1.40 (s, 9 H), 1.18 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 563 $[M+H]^+$.
53	<i>tert</i> -butyl 4-[4-[3-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]azepane-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.32 (d, $J=5.4$ Hz, 1 H), 7.17 (d, $J=5.4$ Hz, 1 H), 7.02–6.96 (m, 2 H), 6.84 (s, 1 H), 6.75–6.69 (m, 2 H), 6.45 (s, 1 H), 4.45–4.36 (m, 1 H), 3.92 (t, $J=7.3$ Hz, 2 H), 3.73 (s, 3 H), 3.44–3.22 (m, 4 H), 2.57 (q, $J=7.3$ Hz, 2 H), 2.44 (t, $J=7.3$ Hz, 2 H), 2.03–1.48


			(m, 8 H), 1.40 (s, 9 H), 1.23 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 563 $[M+H]^+$.
54	<i>tert</i> -butyl 4-[4-[3-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-phenyl-imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.74–7.63 (m, 2 H), 7.44–7.38 (m, 2 H), 7.35 (d, $J=5.2$ Hz, 1 H), 7.28–7.22 (m, 1 H), 7.20 (d, $J=5.2$ Hz, 1 H), 7.04–6.75 (m, 4 H), 6.56 (s, 1 H), 4.50–4.42 (m, 1 H), 4.05–3.98 (m, 2 H), 3.81 (s, 3 H), 3.68–3.59 (m, 2 H), 3.22–3.05 (m, 2 H), 2.81 (q, $J=7.5$ Hz, 2 H), 2.54–2.44 (m, 2 H), 1.92–1.81 (m, 4 H), 1.53–1.43 (m, 2 H), 1.40 (s, 9 H), 1.18 (t, $J=7.5$ Hz, 3 H); MS (ESI): m/z : 625 $[M+H]^+$.
55	<i>tert</i> -butyl 4-[4-[3-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-phenyl-imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.51–7.46 (m, 2 H), 7.45–7.38 (m, 3 H), 7.34 (d, $J=5.4$ Hz, 1 H), 7.20 (d, $J=5.4$ Hz, 1 H), 6.76–6.65 (m, 4 H), 6.54 (s, 1 H), 4.46–4.33 (m, 1 H), 3.97 (t, $J=7.3$ Hz, 2 H), 3.83 (s, 3 H), 3.68–3.57 (m, 2 H), 3.22–3.07 (m, 2 H), 2.45 (q, $J=7.6$ Hz, 2 H), 2.14 (t,



			$J=7.3$ Hz, 2 H), 1.88–1.78 (m, 2 H), 1.56–1.37 (m, 13 H), 1.13 (t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 625 $[M+H]^+$.
56	<i>tert</i> -butyl 4-[4-[2-[4-isopropyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-phenyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		^1H NMR (CDCl_3) δ (ppm): 7.52–7.37 (m, 5 H), 7.16 (d, $J=5.4$ Hz, 1 H), 6.98 (d, $J=5.4$ Hz, 1 H), 6.70–6.65 (m, 3 H), 6.44–6.39 (m, 2 H), 4.35 (t, $J=6.4$ Hz, 2 H), 4.30–4.23 (m, 1 H), 3.83 (s, 3 H), 3.74 (t, $J=6.4$ Hz, 2 H), 3.72–3.64 (m, 2 H), 3.34–3.23 (m, 2 H), 2.90 (sep, $J=6.7$ Hz, 1 H), 1.90–1.62 (m, 4 H), 1.47 (s, 9 H), 1.27 (d, $J=6.7$ Hz, 6 H); MS (ESI): m/z : 641 $[M+H]^+$.
57	<i>tert</i> -butyl 4-[4-[3-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		^1H NMR (CDCl_3) δ (ppm): 7.15 (d, $J=5.4$ Hz, 1 H), 7.01–6.93 (m, 3 H), 6.81–6.75 (m, 3 H), 6.40 (s, 1 H), 4.44–4.36 (m, 1 H), 3.98 (t, $J=7.6$ Hz, 2 H), 3.83 (s, 3 H), 3.75–3.66 (m, 2 H), 3.59–3.47 (m, 1 H), 3.37–3.28 (m, 2 H), 2.54 (t, $J=7.6$ Hz, 2 H), 2.38–2.20 (m, 4 H), 2.09–1.85 (m, 6 H), 1.78–1.70 (m, 2 H), 1.48

			(s, 9 H); MS (ESI): m/z: 575 [M+H] ⁺ .
58	<i>tert</i> -butyl 4-[4-[3-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹ H NMR (CDCl ₃) δ (ppm): 7.16 (d, <i>J</i> =5.4 Hz, 1 H), 7.01 (s, 1 H), 7.00–6.95 (m, 3 H), 6.83–6.77 (m, 2 H), 6.42 (s, 1 H), 4.46–4.37 (m, 1 H), 3.91–3.83 (m, 2 H), 3.81 (s, 3 H), 3.76–3.63 (m, 2 H), 3.41–3.27 (m, 3 H), 2.51 (t, <i>J</i> =7.3 Hz, 2 H), 2.34–2.13 (m, 4 H), 2.09–1.85 (m, 6 H), 1.80–1.67 (m, 2 H), 1.48 (s, 9 H); MS (ESI): m/z: 575 [M+H] ⁺ .
59	<i>tert</i> -butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-phenyl-imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹ H NMR (DMSO-d ₆) δ (ppm): 7.90 (s, 1 H), 7.85–7.79 (m, 2 H), 7.41–7.33 (m, 3 H), 7.25–7.18 (m, 2 H), 7.06–6.98 (m, 2 H), 6.82–6.76 (m, 2 H), 6.53 (s, 1 H), 4.49–4.38 (m, 1 H), 4.06 (t, <i>J</i> =7.3 Hz, 2 H), 3.88 (s, 3 H), 3.68–3.59 (m, 2 H), 3.22–3.06 (m, 2 H), 2.52 (t, <i>J</i> =7.3 Hz, 2 H), 2.10–2.01 (m, 2 H), 1.89–1.80 (m, 2 H), 1.51–1.42 (m, 2 H), 1.40 (s, 9 H); MS (ESI): m/z: 597 [M+H] ⁺ .



60			¹ H NMR (DMSO-d ₆) δ (ppm): 7.29 (d, <i>J</i> =5.4 Hz, 1 H), 7.16 (d, <i>J</i> =5.4 Hz, 1 H), 7.06 (s, 1 H), 6.78–6.71 (m, 2 H), 6.60–6.55 (m, 1 H), 6.39 (s, 1 H), 4.02–3.87 (m, 6 H), 3.78 (s, 3 H), 3.76–3.71 (m, 4 H), 2.83–2.58 (m, 4 H), 2.53 (q, <i>J</i> =7.3 Hz, 2 H), 2.44 (t, <i>J</i> =7.3 Hz, 2 H), 2.01–1.93 (m, 2 H), 1.91–1.80 (m, 2 H), 1.75–1.65 (m, 4 H), 1.39 (s, 9 H), 1.38 (s, 9 H), 1.21–1.08 (m, 7 H); MS (ESI): <i>m/z</i> : 777 [M+H] ⁺ .
61	<i>tert</i> -butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹ H NMR (CDCl ₃) δ (ppm): 7.16 (d, <i>J</i> =5.4 Hz, 1 H), 6.97 (d, <i>J</i> =5.4 Hz, 1 H), 6.93 (s, 1 H), 6.85–6.71 (m, 4 H), 6.57 (bs, 1 H), 4.42–4.27 (m, 3 H), 4.15 (t, <i>J</i> =5.1 Hz, 2 H), 3.83 (bs, 3 H), 3.75–3.66 (m, 2 H), 3.35–3.25 (m, 2 H), 2.66–2.55 (m, 2 H), 1.94–1.80 (m, 2 H), 1.77–1.65 (m, 4 H), 1.48 (s, 9 H), 1.00 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 565 [M+H] ⁺ .



62	<i>tert</i>-butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.18 (d, <i>J</i>=5.4 Hz, 1 H), 7.00 (s, 1 H), 6.98 (d, <i>J</i>=5.4 Hz, 1 H), 6.79–6.73 (m, 2 H), 6.67–6.59 (m, 3 H), 4.39 (t, <i>J</i>=5.9 Hz, 2 H), 4.33–4.24 (m, 1 H), 4.03 (t, <i>J</i>=5.9 Hz, 2 H), 3.78 (s, 3 H), 3.73–3.66 (m, 2 H), 3.34–3.26 (m, 2 H), 2.68 (t, <i>J</i>=7.3 Hz, 2 H), 1.92–1.65 (m, 6 H), 1.47 (s, 9 H), 1.09 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 565 [M+H]⁺.
63	<i>tert</i>-butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]propyl]phenoxy]azepane-1-carboxylate		¹H NMR (DMSO-<i>d</i>₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.07 (s, 1 H), 7.01–6.94 (m, 2 H), 6.76–6.66 (m, 2 H), 6.40 (s, 1 H), 4.43–4.32 (m, 1 H), 3.94 (t, <i>J</i>=7.3 Hz, 2 H), 3.78 (s, 3 H), 3.43–3.21 (m, 4H), 2.51–2.38 (m, 4 H), 2.02–1.49 (m, 10 H), 1.40 (s, 9 H), 0.93 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 577 [M+H]⁺.
64	<i>tert</i>-butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-		¹H NMR (DMSO-<i>d</i>₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.02–6.95 (m, 2 H), 6.82 (s,



	propyl-imidazol-1-yl]propyl]phenoxy]azepane-1-carboxylate		1 H), 6.76-6.70 (m, 2 H), 6.46 (s, 1 H), 4.46-4.34 (m, 1 H), 3.91 (t, $J=8.3$ Hz, 2 H), 3.72 (s, 3 H), 3.44-3.23 (m, 4 H), 2.50-2.40 (m, 4 H), 2.05-1.50 (m, 10 H), 1.40 (s, 9 H), 0.96 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 577 $[M+H]^+$.
65	<i>tert</i> -butyl 4-[[2-[(1- <i>tert</i> -butoxycarbonyl-4-piperidyl)methoxy]-4-[3-[5-ethyl-2-(4-methylthieno[3,2- <i>b</i>]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]piperidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.29 (d, $J=5.4$ Hz, 1 H), 7.15 (d, $J=5.4$ Hz, 1 H), 6.82 (s, 1 H), 6.78-6.72 (m, 2 H), 6.61-6.55 (m, 1 H), 6.43 (s, 1 H), 4.03-3.86 (m, 6 H), 3.78-3.69 (m, 7 H), 2.83-2.63 (m, 4 H), 2.57 (q, $J=7.3$ Hz, 2 H), 2.43 (t, $J=7.3$ Hz, 2 H), 1.92-1.80 (m, 4 H), 1.77-1.65 (m, 4 H), 1.39 (s, 9 H), 1.38 (s, 9 H), 1.27-1.08 (m, 7 H) MS (ESI): m/z : 777 $[M+H]^+$.



66	<i>tert</i>-butyl 4-[4-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.14 (d, <i>J</i>=5.4 Hz, 1 H), 7.06 (s, 1 H), 6.98–6.92 (m, 2 H), 6.84–6.79 (m, 2 H), 6.36 (s, 1 H), 4.51–4.42 (m, 1 H), 4.14 (t, <i>J</i>=7.3 Hz, 2 H), 3.68–3.58 (m, 5 H), 3.21–3.08 (m, 2 H), 2.91 (t, <i>J</i>=7.3 Hz, 2 H), 2.52 (q, <i>J</i>=7.6 Hz, 2 H), 1.90–1.79 (m, 2 H), 1.51–1.42 (m, 2 H), 1.39 (s, 9 H), 1.17 (t, <i>J</i>=7.6 Hz, 3 H) MS (ESI): <i>m/z</i>: 535 [M+H]⁺.
67	<i>tert</i>-butyl 3-[[4-[3-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.11 (s, 1 H), 7.04–6.97 (m, 2 H), 6.79–6.73 (m, 2 H), 6.35 (s, 1 H), 3.95–3.91 (m, 2 H), 3.90–3.81 (m, 2 H), 3.79 (s, 3 H), 3.48–3.33 (m, 3 H), 3.29–3.18 (m, 1 H), 3.11–3.04 (m, 1 H), 2.64–2.51 (m, 1 H), 2.48–2.42 (m, 2 H), 2.26–2.09 (m, 4 H), 2.06–1.58 (m, 6 H), 1.39 (s, 9 H); MS (ESI): <i>m/z</i>: 575 [M+H]⁺.



68	<i>tert</i>-butyl 3-[[4-[3-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.04–6.99 (m, 2 H), 6.93 (s, 1 H), 6.81–6.76 (m, 2 H), 6.39 (s, 1 H), 3.91–3.81 (m, 4 H), 3.72 (s, 3 H), 3.49–3.35 (m, 3 H), 3.28–3.17 (m, 1 H), 3.10–3.04 (m, 1 H), 2.61–2.53 (m, 1 H), 2.45 (t, <i>J</i>=7.3 Hz, 2 H), 2.30–1.62 (m, 10 H), 1.39 (s, 9 H); MS (ESI): <i>m/z</i>: 575 [M+H]⁺.
69	<i>tert</i>-butyl 4-[4-[3-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]azepane-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.12 (s, 1 H), 7.02–6.96 (m, 2 H), 6.76–6.68 (m, 2 H), 6.41 (s, 1 H), 4.45–4.31 (m, 1 H), 3.94 (t, <i>J</i>=7.3 Hz, 2 H), 3.79 (s, 3 H), 3.45–3.35 (m, 3 H), 3.31–3.24 (m, 2 H), 2.44 (t, <i>J</i>=7.3 Hz, 2 H), 2.26–2.08 (m, 4 H), 2.01–1.50 (m, 10 H), 1.40 (s, 9 H); MS (ESI): <i>m/z</i>: 589 [M+H]⁺.


70	<i>tert</i>-butyl 4-[4-[3-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]azepane-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.03–6.97 (m, 2 H), 6.92 (s, 1 H), 6.78–6.70 (m, 2 H), 6.47 (s, 1 H), 4.47–4.36 (m, 2 H), 3.84 (t, <i>J</i>=6.8 Hz, 1 H), 3.72 (s, 3 H), 3.46–3.35 (m, 5 H), 2.43 (t, <i>J</i>=6.8 Hz, 2 H), 2.27–1.50 (m, 14 H), 1.40 (s, 9 H); MS (ESI): <i>m/z</i>: 589 [M+H]⁺.
71	<i>tert</i>-butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.07 (s, 1 H), 7.01–6.95 (m, 2 H), 6.80–6.74 (m, 2 H), 6.42 (s, 1 H), 4.48–4.39 (m, 1 H), 3.95 (t, <i>J</i>=7.3 Hz, 2 H), 3.78 (s, 3 H), 3.68–3.58 (m, 2 H), 3.21–3.06 (m, 2 H), 2.48–2.39 (m, 4 H), 2.00–1.91 (m, 2 H), 1.89–1.80 (m, 2 H), 1.67–1.56 (m, 2 H), 1.51–1.42 (m, 2 H), 1.40 (s, 9 H), 0.93 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 563 [M+H]⁺.

72	<i>tert</i>-butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.03–6.97 (m, 2 H), 6.82 (s, 1 H), 6.81–6.76 (m, 2 H), 6.47 (s, 1 H), 4.50–4.39 (m, 1 H), 3.92 (t, <i>J</i>=7.8 Hz, 2 H), 3.72 (s, 3 H), 3.68–3.59 (m, 2 H), 3.21–3.05 (m, 2 H), 2.48–2.41 (m, 4 H), 1.90–1.76 (m, 4 H), 1.66–1.56 (m, 2 H), 1.52–1.43 (m, 2 H), 1.40 (s, 9 H), 0.96 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 563 [M+H]⁺.
73	<i>tert</i>-butyl 4-[[4-[3-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.11 (s, 1 H), 7.03–6.97 (m, 2 H), 6.77–6.71 (m, 2 H), 6.38 (s, 1 H), 4.02–3.89 (m, 4 H), 3.79 (s, 3 H), 3.75 (d, <i>J</i>=6.4 Hz, 2 H), 3.46–3.36 (m, 1 H), 2.83–2.64 (m, 2 H), 2.45 (t, <i>J</i>=7.3 Hz, 2 H), 2.26–2.09 (m, 4 H), 2.01–1.79 (m, 5 H), 1.77–1.68 (m, 2 H), 1.39 (s, 9 H), 1.19–1.06 (m, 2 H); MS (ESI): <i>m/z</i>: 589 [M+H]⁺.

74	<i>tert</i>-butyl 4-[[4-[3-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.03–6.97 (m, 2 H), 6.93 (s, 1 H), 6.79–6.73 (m, 2 H), 6.42 (s, 1 H), 4.02–3.90 (m, 2 H), 3.84 (t, <i>J</i>=7.8 Hz, 2 H), 3.77 (d, <i>J</i>=6.4 Hz, 2 H), 3.72 (s, 3 H), 3.48–3.39 (m, 1 H), 2.82–2.64 (m, 2 H), 2.44 (t, <i>J</i>=7.3 Hz, 2 H), 2.30–2.21 (m, 2 H), 2.14–2.03 (m, 2 H), 2.02–1.68 (m, 7 H), 1.39 (s, 9 H), 1.20–1.07 (m, 2 H); MS (ESI): <i>m/z</i>: 589 [M+H]⁺.
75	<i>tert</i>-butyl 3-[[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.29 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.06 (s, 1 H), 7.03–6.96 (m, 2 H), 6.80–6.72 (m, 2 H), 6.35 (s, 1 H), 3.93 (t, <i>J</i>=7.3 Hz, 2 H), 3.89–3.81 (m, 2 H), 3.78 (s, 3 H), 3.49–3.39 (m, 1 H), 3.38–3.34 (m, 1 H), 3.29–3.17 (m, 1 H), 3.11–3.04 (m, 1 H), 2.61–2.53 (m, 1 H), 2.48–2.39 (m, 4 H), 2.05–1.90 (m, 3 H), 1.76–1.56 (m, 3 H), 1.39 (s, 9 H), 0.93



			(t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 563 $[M+H]^+$.
76	<i>tert</i> -butyl 3-[[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.30 (d, $J=5.4$ Hz, 1 H), 7.16 (d, $J=5.4$ Hz, 1 H), 7.04–6.97 (m, 2 H), 6.82 (s, 1 H), 6.79–6.74 (m, 2 H), 6.39 (s, 1 H), 3.95–3.80 (m, 4 H), 3.71 (s, 3 H), 3.50–3.40 (m, 1 H), 3.38–3.34 (m, 1 H), 3.27–3.17 (m, 1 H), 3.11–3.04 (m, 1 H), 2.62–2.52 (m, 1 H), 2.48–2.42 (m, 4 H), 2.03–1.96 (m, 1 H), 1.86–1.78 (m, 2 H), 1.75–1.57 (m, 3 H), 1.39 (s, 9 H), 0.96 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 563 $[M+H]^+$.
77	<i>tert</i> -butyl 4-[4-[2-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethyl]phenoxy]piperidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.30 (d, $J=5.4$ Hz, 1 H), 7.15 (d, $J=5.4$ Hz, 1 H), 7.10 (s, 1 H), 6.98–6.93 (m, 2 H), 6.84–6.79 (m, 2 H), 6.38 (s, 1 H), 4.51–4.42 (m, 1 H), 4.14 (t, $J=7.3$ Hz, 2 H), 3.68 (s, 3 H), 3.65–3.58 (m, 2 H), 3.44–3.35 (m, 1 H), 3.19–3.09 (m, 2 H), 2.91 (t, $J=7.3$ Hz, 2 H), 2.27–2.07 (m, 4 H), 1.98–



			1.81 (m, 4 H), 1.51–1.42 (m, 2 H), 1.39 (s, 9 H); MS (ESI): m/z: 561 [M+H] ⁺ .
78	<i>tert</i> -butyl 4-[3-[3-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹ H NMR (DMSO-d ₆) δ (ppm): 7.30 (d, <i>J</i> =5.4 Hz, 1 H), 7.15 (d, <i>J</i> =5.4 Hz, 1 H), 7.14–7.09 (m, 1 H), 7.07 (s, 1 H), 6.79–6.74 (m, 2 H), 6.71–6.66 (m, 1 H), 6.37 (s, 1 H), 4.55–4.40 (m, 1 H), 3.96 (t, <i>J</i> =7.3 Hz, 2 H), 3.78 (s, 3 H), 3.67–3.54 (m, 2 H), 3.21–3.07 (m, 2 H), 2.56–2.52 (m, 4 H), 2.04–1.94 (m, 2 H), 1.89–1.79 (m, 2 H), 1.50–1.42 (m, 2 H), 1.39 (s, 9 H), 1.18 (t, <i>J</i> =7.6 Hz, 3 H); MS (ESI): m/z: 549 [M+H] ⁺ .
79	<i>tert</i> -butyl 4-[3-[3-[5-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹ H NMR (DMSO-d ₆) δ (ppm): 7.31 (d, <i>J</i> =5.4 Hz, 1 H), 7.16 (d, <i>J</i> =5.4 Hz, 1 H), 7.14–7.08 (m, 1 H), 6.83 (s, 1 H), 6.79–6.74 (m, 2 H), 6.70–6.66 (m, 1 H), 6.41 (s, 1 H), 4.56–4.42 (m, 1 H), 3.94 (t, <i>J</i> =7.8 Hz,



			2 H), 3.73 (s, 3 H), 3.66–3.56 (m, 2 H), 3.20–3.05 (m, 2 H), 2.57 (q, $J=7.3$ Hz, 2 H), 2.48–2.43 (m, 2 H), 1.92–1.78 (m, 4 H), 1.51–1.41 (m, 2 H), 1.39 (s, 9 H), 1.23 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 549 $[M+H]^+$.
80	<i>tert</i> -butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]ethyl]phenoxy]piperidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.30 (d, $J=5.4$ Hz, 1 H), 7.14 (d, $J=5.4$ Hz, 1 H), 7.04 (s, 1 H), 6.96–6.90 (m, 2 H), 6.83–6.77 (m, 2 H), 6.35 (s, 1 H), 4.50–4.42 (m, 1 H), 4.15 (t, $J=7.3$ Hz, 2 H), 3.68–3.57 (m, 5 H), 3.20–3.08 (m, 2 H), 2.90 (t, $J=7.1$ Hz, 2 H), 2.45 (t, $J=7.3$ Hz, 2 H), 1.89–1.78 (m, 2 H), 1.65–1.55 (m, 2 H), 1.50–1.42 (m, 2 H), 1.39 (s, 9 H), 0.92 (t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 549 $[M+H]^+$.


81	<i>tert</i>-butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]ethyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 6.88–6.83 (m, 2 H), 6.81–6.74 (m, 3 H), 6.57 (s, 1 H), 4.51–4.40 (m, 1 H), 4.16 (t, <i>J</i>=7.3 Hz, 2 H), 3.68–3.56 (m, 5 H), 3.22–3.07 (m, 2 H), 2.74 (t, <i>J</i>=7.1 Hz, 2 H), 2.44 (t, <i>J</i>=7.3 Hz, 2 H), 1.91–1.78 (m, 2 H), 1.68–1.55 (m, 2 H), 1.51–1.42 (m, 2 H), 1.39 (s, 9 H), 0.97 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 549 [M+H]⁺.
82	<i>tert</i>-butyl 4-[[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]propyl]phenyl]methoxy]piperidine-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.15 (d, <i>J</i>=5.4 Hz, 1 H), 7.01–6.93 (m, 3 H), 6.79–6.71 (m, 3 H), 6.37 (s, 1 H), 4.30–4.05 (m, 2 H), 3.98 (t, <i>J</i>=7.3 Hz, 2 H), 3.83 (s, 3 H), 3.79–3.74 (m, 2 H), 2.82–2.69 (m, 2 H), 2.61 (t, <i>J</i>=7.3 Hz, 2 H), 2.53 (t, <i>J</i>=7.3 Hz, 2 H), 2.01–1.99 (m, 2 H), 1.98–1.89 (m, 1 H), 1.87–1.79 (m, 2 H), 1.77–1.68 (m, 2 H), 1.48 (s, 9 H), 1.32–1.22 (m, 2 H), 1.00 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 577 [M+H]⁺.



83	<i>tert</i>-butyl 4-[[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]propyl]phenyl]methoxy]piperidine-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.01–6.91 (m, 4 H), 6.79–6.73 (m, 2 H), 6.39 (s, 1 H), 4.31–4.02 (m, 2 H), 3.94 (t, <i>J</i>=7.3 Hz, 2 H), 3.83–3.74 (m, 5 H), 2.85–2.65 (m, 2 H), 2.57–2.45 (m, 4 H), 2.01–1.88 (m, 3 H), 1.87–1.80 (m, 2 H), 1.77–1.66 (m, 2 H), 1.48 (s, 9 H), 1.34–1.20 (m, 2 H), 1.04 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 577 [M+H]⁺.
84	<i>tert</i>-butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 6.97 (d, <i>J</i>=5.4 Hz, 1 H), 6.93 (s, 1 H), 6.83–6.69 (m, 4 H), 6.57 (s, 1 H), 4.37 (t, <i>J</i>=5.1 Hz, 2 H), 4.33–4.27 (m, 1 H), 4.15 (t, <i>J</i>=5.1 Hz, 2 H), 3.82 (s, 3 H), 3.64–3.36 (m, 4 H), 2.61 (m, 2 H), 2.12–1.82 (m, 4 H), 1.79–1.68 (m, 2 H), 1.60 (m, 2 H), 1.48 (s, 9 H), 1.00 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 579 [M+H]⁺.



85	<i>tert</i>-butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.00–6.95 (m, 2 H), 6.77–6.59 (m, 5 H), 4.37 (t, <i>J</i>=5.9 Hz, 2 H), 4.32–4.23 (m, 1 H), 4.02 (t, <i>J</i>=5.9 Hz, 2 H), 3.76 (s, 3 H), 3.65–3.36 (m, 4 H), 2.68 (t, <i>J</i>=7.3 Hz, 2 H), 2.09–1.82 (m, 4 H), 1.84–1.74 (m, 2 H), 1.68–1.56 (m, 2 H), 1.48 (s, 9 H), 1.09 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 579 [M+H]⁺.
86	<i>tert</i>-butyl 4-[4-[3-[4-cyclobutyl-2-(6-methylthieno[2,3-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.11 (s, 1 H), 7.07 (d, <i>J</i>=5.4 Hz, 1 H), 7.03 – 6.98 (m, 3 H), 6.81 – 6.76 (m, 2 H), 6.34 (s, 1 H), 4.48 – 4.35 (m, 1 H), 3.93 (t, <i>J</i>=7.3 Hz, 2 H), 3.75 (s, 3 H), 3.68 – 3.59 (m, 2 H), 3.46 – 3.36 (m, 1 H), 3.14 (bs, 2 H), 2.45 (t, <i>J</i>=7.3 Hz, 2 H), 2.26 – 2.09 (m, 4 H), 2.02 – 1.79 (m, 6 H), 1.50 – 1.41 (m, 2 H), 1.40 (s, 9 H).; MS (ESI): <i>m/z</i>: 575 [M+H]⁺.


87	<i>tert</i>-butyl 4-[4-[3-[5-cyclobutyl-2-(6-methylthieno[2,3-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.07 (d, <i>J</i>=5.4 Hz, 1 H), 7.04 - 6.99 (m, 3 H), 6.92 (s, 1 H), 6.84 - 6.79 (m, 2 H), 6.39 (s, 1 H), 4.50 - 4.39 (m, 1 H), 3.83 (t, <i>J</i>=7.3 Hz, 2 H), 3.71 - 3.58 (m, 5 H), 3.46 - 3.36 (m, 1 H), 3.14 (br. s, 2 H), 2.44 (t, <i>J</i>=7.3 Hz, 2 H), 2.27 - 2.17 (m, 2 H), 2.14 - 2.03 (m, 2 H), 2.00 - 1.74 (m, 6 H), 1.54 - 1.43 (m, 2 H), 1.40 (s, 9 H); MS (ESI): <i>m/z</i>: 575 [M+H]⁺.
88	<i>tert</i>-butyl (3<i>R</i>)-3-[[4-[3-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.11 (s, 1 H), 7.04 - 6.98 (m, 2 H), 6.79 - 6.73 (m, 2 H), 6.35 (s, 1 H), 3.97 - 3.81 (m, 4 H), 3.79 (s, 3 H), 3.47 - 3.33 (m, 3 H), 3.28 - 3.17 (m, 1 H), 3.11 - 3.01 (m, 1 H), 2.62 - 2.53 (m, 1 H), 2.46 (t, <i>J</i>=7.3 Hz, 2 H), 2.26 - 2.10 (m, 4 H), 2.03 - 1.89 (m, 4 H), 1.88 - 1.77 (m, 1 H), 1.74 - 1.63 (m, 1 H), 1.39 (s, 9 H); MS (ESI): <i>m/z</i>: 575 [M+H]⁺.



89	<i>tert</i>-butyl (3R)-3-[[4-[3-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.15 (d, <i>J</i>=5.4 Hz, 1 H), 7.04 - 6.99 (m, 2 H), 6.93 (s, 1 H), 6.81 - 6.74 (m, 2 H), 6.39 (s, 1 H), 3.93 - 3.80 (m, 4 H), 3.72 (s, 3 H), 3.49 - 3.36 (m, 3 H), 3.26 - 3.15 (m, 1 H), 3.11 - 3.01 (m, 1 H), 2.62 - 2.53 (m, 1 H), 2.45 (t, <i>J</i>=7.3 Hz, 2 H), 2.30 - 2.20 (m, 2 H), 2.14 - 2.03 (m, 2 H), 2.03 - 1.90 (m, 2 H), 1.88 - 1.63 (m, 4 H), 1.39 (s, 9 H); MS (ESI): <i>m/z</i>: 575 [M+H]⁺.
90	<i>tert</i>-butyl (3S)-3-[[4-[3-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.11 (s, 1 H), 7.04 - 6.99 (m, 2 H), 6.79 - 6.73 (m, 2 H), 6.35 (s, 1 H), 3.93 (t, <i>J</i>=7.3 Hz, 2 H), 3.89 - 3.81 (m, 2 H), 3.79 (s, 3 H), 3.47 - 3.35 (m, 3 H), 3.29 - 3.17 (m, 1 H), 3.11 - 3.03 (m, 1 H), 2.61 - 2.53 (m, 1 H), 2.45 (t, <i>J</i>=7.3 Hz, 2 H), 2.26 - 2.09 (m, 4 H), 2.02 - 1.60 (m, 6 H), 1.39 (s, 9



			H).; MS (ESI): m/z: 575 [M+H] ⁺ .
91	<i>tert</i>-butyl (3<i>S</i>)-3-[[4-[3-[5-cyclobutyl-2-(4-methylthieno[3,2-<i>b</i>]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]methyl]pyrrolidine-1-carboxylate		¹H NMR (DMSO-<i>d</i>₆) δ (ppm): 7.29 (d, <i>J</i>=5.4 Hz, 1 H), 7.15 (d, <i>J</i>=5.4 Hz, 1 H), 7.04 - 6.97 (m, 2 H), 6.92 (s, 1 H), 6.80 - 6.75 (m, 2 H), 6.39 (s, 1 H), 3.93 - 3.80 (m, 4 H), 3.72 (s, 3 H), 3.48 - 3.36 (m, 3 H), 3.27 - 3.16 (m, 1 H), 3.13 - 3.04 (m, 1 H), 2.62 - 2.55 (m, 1 H), 2.44 (t, <i>J</i>=7.3 Hz, 2 H), 2.31 - 2.20 (m, 2 H), 2.14 - 1.61 (m, 8 H), 1.39 (s, 9 H); MS (ESI): m/z: 575 [M+H]⁺.


92	<i>tert</i>-butyl 4-[[4-[2-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]methyl]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.33 (d, <i>J</i>=5.4 Hz, 1 H), 7.21 - 7.15 (m, 2 H), 6.83 - 6.73 (m, 4 H), 6.68 (s, 1 H), 4.31 (t, <i>J</i>=5.3 Hz, 2 H), 4.18 (t, <i>J</i>=5.3 Hz, 2 H), 3.95 (bs, 2 H), 3.78 (s, 3 H), 3.73 (d, <i>J</i>=6.4 Hz, 2 H), 3.47 - 3.37 (m, 1 H), 2.73 (bs, 2 H), 2.25 - 2.09 (m, 4 H), 1.98 - 1.78 (m, 3 H), 1.75 - 1.67 (m, 2 H), 1.38 (s, 9 H), 1.17 - 1.05 (m, 2 H); MS (ESI): <i>m/z</i>: 591 [M+H]⁺.
93	<i>tert</i>-butyl 4-[3-[3-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.29 (d, <i>J</i>= 5.4 Hz, 1 H), 7.16 - 7.06 (m, 3 H), 6.78 - 6.71 (m, 2 H), 6.70 - 6.65 (m, 1 H), 6.33 (s, 1 H), 4.53 - 4.43 (m, 1 H), 3.92 (t, <i>J</i>=7.3 Hz, 2 H), 3.75 (s, 3 H), 3.63 - 3.57 (m, 2 H), 3.45 - 3.36 (m, 1 H), 3.12 (bs, 2 H), 2.46 (t, <i>J</i>=7.3 Hz, 2 H), 2.24 - 2.08 (m, 4 H), 2.02 - 1.76 (m, 6 H), 1.48 - 1.39 (m, 2 H), 1.37 (s, 9 H); MS (ESI): <i>m/z</i>: 575 [M+H]⁺.



94	<i>tert</i>-butyl 4-[3-[3-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.18 - 7.09 (m, 2 H), 6.93 (s, 1 H), 6.81 - 6.75 (m, 2 H), 6.72 - 6.65 (m, 1 H), 6.44 (s, 1 H), 4.55 - 4.46 (m, 1 H), 3.86 (t, <i>J</i>=7.3 Hz, 2 H), 3.73 (s, 3 H), 3.66 - 3.56 (m, 2 H), 3.47 - 3.37 (m, 1 H), 3.14 (bs, 2 H), 2.46 (t, <i>J</i>=7.3 Hz, 2 H), 2.27 - 2.18 (m, 2 H), 2.13 - 2.03 (m, 2 H), 2.00 - 1.78 (m, 6 H), 1.51 - 1.43 (m, 2 H), 1.39 (s, 9 H); MS (ESI): <i>m/z</i>: 575 [M+H]⁺.
95	<i>tert</i>-butyl 4-[3-[2-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.21 - 7.16 (m, 2 H), 7.15 - 7.08 (m, 1 H), 6.68 (s, 1 H), 6.57 - 6.52 (m, 1 H), 6.47 - 6.41 (m, 2 H), 4.55 - 4.47 (m, 1 H), 4.32 (t, <i>J</i>=5.4 Hz, 2 H), 4.24 (t, <i>J</i>=5.4 Hz, 2 H), 3.78 (s, 3 H), 3.66 - 3.57 (m, 2 H), 3.45 - 3.37 (m, 1 H), 3.15 (bs, 2 H), 2.26 - 2.10 (m, 4 H), 1.99 - 1.79 (m, 4 H), 1.53 - 1.43 (m, 2 H), 1.39


			(s, 9 H); MS (ESI): m/z: 577 [M+H] ⁺ .
96	<i>tert</i>-butyl 4-[3-[2-[5-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.18 (d, <i>J</i>=5.4 Hz, 1 H), 7.10 - 7.04 (m, 1 H), 6.98 (s, 1 H), 6.71 (s, 1 H), 6.55 - 6.49 (m, 1 H), 6.36 - 6.28 (m, 2 H), 4.52 - 4.44 (m, 1 H), 4.31 (t, <i>J</i>=5.4 Hz, 2 H), 4.04 (t, <i>J</i>=5.4 Hz, 2 H), 3.73 - 3.55 (m, 6 H), 3.17 (bs, 2 H), 2.41 - 2.32 (m, 2 H), 2.20 - 1.77 (m, 6 H), 1.51 - 1.42 (m, 2 H), 1.39 (s, 9 H); MS (ESI): m/z: 577 [M+H]⁺.



97	<i>tert</i>-butyl 4-[4-[3-[4-methyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.07 (s, 1 H), 7.01 - 6.95 (m, 2 H), 6.81 - 6.73 (m, 2 H), 6.40 (s, 1 H), 4.47 - 4.38 (m, 1 H), 3.93 (t, <i>J</i>=7.3 Hz, 2 H), 3.78 (s, 3 H), 3.68 - 3.59 (m, 2 H), 3.14 (bs, 2 H), 2.43 (t, <i>J</i>=7.3 Hz, 2 H), 2.15 (s, 3 H), 1.98 - 1.89 (m, 2 H), 1.88 - 1.80 (m, 2 H), 1.50 - 1.42 (m, 2 H), 1.39 (s, 9 H); MS (ESI): <i>m/z</i>:535 [M+H]⁺.
98	<i>tert</i>-butyl 4-[4-[3-[5-methyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.04 - 6.95 (m, 2 H), 6.82 (s, 1 H), 6.80 - 6.76 (m, 2 H), 6.43 (s, 1 H), 4.49 - 4.40 (m, 1 H), 3.92 (t, <i>J</i>=7.3 Hz, 2 H), 3.72 (s, 3 H), 3.68 - 3.59 (m, 2 H), 3.14 (bs, 2 H), 2.45 (t, <i>J</i>=7.3 Hz, 2 H), 2.21 (s, 3 H), 1.89 - 1.80 (m, 4 H), 1.51 - 1.42 (m, 2 H), 1.39 (s, 9 H); MS (ESI): <i>m/z</i>: 535 [M+H]⁺.



99	<i>tert</i>-butyl 4-[[4-[2-[4-cyclobutyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethyl]phenoxy]methyl]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.31 (d, <i>J</i>=5.4 Hz, 1 H), 7.15 (d, <i>J</i>=5.4 Hz, 1 H), 7.10 (s, 1 H), 7.00 - 6.93 (m, 2 H), 6.79 - 6.72 (m, 2 H), 6.46 (s, 1 H), 4.14 (t, <i>J</i>=7.3 Hz, 2 H), 3.94 (bs, 2 H), 3.75 (d, <i>J</i>=6.4 Hz, 2 H), 3.67 (s, 3 H), 3.45 - 3.35 (m, 1 H), 2.90 (t, <i>J</i>=7.3 Hz, 2 H), 2.72 (bs, 2 H), 2.27 - 2.06 (m, 4 H), 2.00 - 1.78 (m, 3 H), 1.76 - 1.69 (m, 2 H), 1.38 (s, 9 H), 1.17 - 1.04 (m, 2 H); MS (ESI): <i>m/z</i>: 575 [M+H]⁺.
100	<i>tert</i>-butyl 4-[3-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.30 (d, <i>J</i>=5.4 Hz, 1 H), 7.15 (d, <i>J</i>=5.4 Hz, 1 H), 7.13 - 7.09 (m, 1 H), 7.07 (s, 1 H), 6.79 - 6.74 (m, 2 H), 6.70 - 6.66 (m, 1 H), 6.38 (s, 1 H), 4.54 - 4.45 (m, 1 H), 3.96 (t, <i>J</i>=7.3 Hz, 2 H), 3.77 (s, 3 H), 3.66 - 3.57 (m, 2 H), 3.20 - 3.07 (m, 2 H), 2.49 - 2.43 (m, 4 H), 2.04 - 1.95 (m, 2 H), 1.88 - 1.80 (m, 2 H), 1.67 - 1.56 (m, 2 H), 1.50 - 1.42 (m, 2 H), 1.39 (s, 9 H), 0.93

			(t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 563 $[M+H]^+$.
101	<i>tert</i> -butyl 4-[3-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		^1H NMR (DMSO- d_6) δ (ppm): 7.30 (d, $J=5.4$ Hz, 1 H), 7.16 (d, $J=5.4$ Hz, 1 H), 7.14 - 7.09 (m, 1 H), 6.82 (s, 1 H), 6.79 - 6.75 (m, 2 H), 6.70 - 6.66 (m, 1 H), 6.43 (s, 1 H), 4.56 - 4.45 (m, 1 H), 3.98 - 3.89 (m, 2 H), 3.72 (s, 3 H), 3.66 - 3.56 (m, 2 H), 3.14 (bs, 2 H), 2.49 - 2.44 (m, 4 H), 1.91 - 1.80 (m, 4 H), 1.66 - 1.56 (m, 2 H), 1.51 - 1.42 (m, 2 H), 1.39 (s, 9 H), 0.96 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 563 $[M+H]^+$.


102	<i>tert</i>-butyl 4-[3-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-d₆) δ (ppm): 7.32 (d, <i>J</i>=5.4 Hz, 1 H), 7.18 (d, <i>J</i>=5.4 Hz, 1 H), 7.15 - 7.09 (m, 2 H), 6.68 (s, 1 H), 6.57 - 6.53 (m, 1 H), 6.47 - 6.42 (m, 2 H), 4.54 - 4.46 (m, 1 H), 4.32 (t, <i>J</i>=5.1 Hz, 2 H), 4.24 (t, <i>J</i>=5.1 Hz, 2 H), 3.76 (s, 3 H), 3.66 - 3.58 (m, 2 H), 3.15 (bs, 2 H), 2.46 (t, <i>J</i>=7.3 Hz, 2 H), 1.89 - 1.81 (m, 2 H), 1.67 - 1.56 (m, 2 H), 1.52 - 1.42 (m, 2 H), 1.40 (s, 9 H), 0.93 (t, <i>J</i>=7.6 Hz, 3 H); MS (ESI): <i>m/z</i>: 565 [M+H]⁺.
103	<i>tert</i>-butyl 4-[3-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.17 (d, <i>J</i>=4.9 Hz, 1 H), 7.14 - 7.07 (m, 1 H), 7.00 - 6.94 (m, 2 H), 6.61 (s, 1 H), 6.54 - 6.49 (m, 1 H), 6.35 - 6.29 (m, 2 H), 4.46 - 4.36 (m, 3 H), 4.05 (t, <i>J</i>=5.9 Hz, 2 H), 3.76 (s, 3 H), 3.72 - 3.64 (m, 2 H), 3.37 - 3.29 (m, 2 H), 2.68 (t, <i>J</i>=7.8 Hz, 2 H), 1.94 - 1.85 (m, 2 H), 1.84 - 1.68 (m, 4 H), 1.48 (s, 9 H), 1.10 (t, <i>J</i>=7.3 Hz,

			3 H); MS (ESI): m/z: 565 [M+H] ⁺ .
104	<i>tert</i> -butyl 4-[[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]propyl]phenoxy]methyl]piperidine-1-carboxylate		¹ H NMR (CDCl ₃) δ (ppm): 7.15 (d, <i>J</i> =5.4 Hz, 1 H), 7.01 - 6.93 (m, 3 H), 6.79 - 6.71 (m, 3 H), 6.37 (s, 1 H), 4.30 - 4.05 (m, 2 H), 3.99 (t, <i>J</i> =7.3 Hz, 2 H), 3.83 (s, 3 H), 3.77 (d, <i>J</i> =6.4 Hz, 2 H), 2.82 - 2.69 (m, 2 H), 2.62 (t, <i>J</i> =7.1 Hz, 2 H), 2.53 (t, <i>J</i> =7.3 Hz, 2 H), 2.04 (t, <i>J</i> =7.3 Hz, 2 H), 1.98 - 1.89 (m, 1 H), 1.87 - 1.79 (m, 2 H), 1.77 - 1.68 (m, 2 H), 1.48 (s, 9 H), 1.34 - 1.21 (m, 2 H), 1.00 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): m/z: 577 [M+H] ⁺ .

105	<i>tert</i>-butyl 4-[[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]propyl]phenoxy]methyl]piperidine-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 7.01 - 6.91 (m, 4 H), 6.82 - 6.72 (m, 2 H), 6.39 (s, 1 H), 4.31 - 4.02 (m, 2 H), 3.98 - 3.90 (m, 2 H), 3.83 - 3.74 (m, 5 H), 2.85 - 2.65 (m, 2 H), 2.56 - 2.44 (m, 4 H), 2.01 - 1.88 (m, 3 H), 1.87 - 1.78 (m, 2 H), 1.76 - 1.65 (m, 2 H), 1.48 (s, 9 H), 1.34 - 1.20 (m, 2 H), 1.04 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 577 [M+H]⁺.
106	<i>tert</i>-butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-propyl-imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.16 (d, <i>J</i>=5.4 Hz, 1 H), 6.97 (d, <i>J</i>=5.4 Hz, 1 H), 6.93 (s, 1 H), 6.83 - 6.69 (m, 4 H), 6.57 (s, 1 H), 4.37 (t, <i>J</i>=5.1 Hz, 2 H), 4.33 - 4.27 (m, 1 H), 4.15 (t, <i>J</i>=5.1 Hz, 2 H), 3.82 (s, 3 H), 3.64 - 3.36 (m, 4 H), 2.61 (bs, 2 H), 2.10 - 1.81 (m, 4 H), 1.78 - 1.68 (m, 2 H), 1.60 (bs, 2 H), 1.48 (s, 9 H), 1.00 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 579 [M+H]⁺.

107	<i>tert</i>-butyl 4-[4-[2-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-5-propyl-imidazol-1-yl]ethoxy]phenoxy]azepane-1-carboxylate		¹H NMR (CDCl₃) δ (ppm): 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.01 - 6.94 (m, 2 H), 6.77 - 6.71 (m, 2 H), 6.66 - 6.56 (m, 3 H), 4.37 (t, <i>J</i>=5.9 Hz, 2 H), 4.31 - 4.24 (m, 1 H), 4.02 (t, <i>J</i>=5.9 Hz, 2 H), 3.76 (s, 3 H), 3.64 - 3.36 (m, 4 H), 2.68 (t, <i>J</i>=7.8 Hz, 2 H), 2.08 - 1.57 (m, 8 H), 1.48 (s, 9 H), 1.09 (t, <i>J</i>=7.3 Hz, 3 H); MS (ESI): <i>m/z</i>: 579 [M+H]⁺.
108	<i>tert</i>-butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)-4-(3,3,3-trifluoropropyl)imidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate		¹H NMR (DMSO-<i>d</i>₆) δ (ppm): 7.17 (d, <i>J</i>=5.4 Hz, 1 H), 7.01 - 6.93 (m, 3 H), 6.83 - 6.72 (m, 3 H), 6.40 (s, 1 H), 4.45 - 4.35 (m, 1 H), 3.99 (t, <i>J</i>=7.3 Hz, 2 H), 3.81 (s, 3 H), 3.75 - 3.64 (m, 2 H), 3.37 - 3.27 (m, 2 H), 2.92 - 2.84 (m, 2 H), 2.62 - 2.47 (m, 4 H), 2.04 (t, <i>J</i>=7.6 Hz, 2 H), 1.95 - 1.84 (m, 2 H), 1.78 - 1.67 (m, 2 H), 1.48 (s, 9 H); MS (ESI): <i>m/z</i>: 617 [M+H]⁺.

Example 18: 5-[4-ethyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride








0.042 g (0.076 mmol) of *tert*-butyl 4-[3-[2-[4-ethyl-2-(4-methylthieno[3,2-b]pyrrol-5-yl)imidazol-1-yl]ethoxy]phenoxy]piperidine-1-carboxylate (Intermediate **26**) was dissolved at 0°C in 0.5 mL of 4 M HCl in dioxane. The mixture was stirred at 0°C for 30 min, then the solvent was removed, the residue was triturated with Et₂O and dried to afford 0.036 g of 5-[4-ethyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride (98%) as a white powder. ¹H NMR (DMSO-d₆) δ (ppm): 8.87–8.71 (m, 2 H), 7.76 (s, 1 H), 7.55 (d, *J*=5.4 Hz, 1 H), 7.28 (d, *J*=5.4 Hz, 1 H), 7.20–7.11 (m, 1 H), 7.04 (s, 1 H), 6.63–6.45 (m, 3 H), 4.65–4.56 (m, 1 H), 4.52–4.44 (m, 2 H), 4.39–4.28 (m, 2 H), 3.77 (s, 3 H), 3.25–3.14 (m, 2 H), 3.15–2.99 (m, 2 H), 2.71 (q, *J*=7.6 Hz, 2 H), 2.14–1.70 (m, 4 H), 1.26 (t, *J*=7.6 Hz, 3 H); MS (ESI): *m/z*: 451 [M+H]⁺.

The following compounds were obtained starting from the corresponding Boc-intermediates according to the procedure described Example **18**.



In case of the examples **19**, **21**, **24**, **26**, **31-33**, **37-39** the compounds were purified by preparative HPLC providing the products as trifluoroacetate salts (Examples **24** and **26**) or as free bases after filtration on a bicarbonate SPE cartridge (200 mg) and eluting with MeOH (Examples **19**, **21**, **31**, **32**, **33**, **39**). In case of the examples **37** and **38** the compounds were filtered on a bicarbonate SPE cartridge (200 mg) and purified by flash chromatography on silica gel. Compounds **41**, **43**, **48**, **49**, **58** and **60** were purified by column chromatography on silica gel (CH₂Cl₂/MeOH/NH₃).




Table 3: Final Products




Ex.	Name	Structure	Analytical Data
19	5-[5-ethyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		^1H NMR (CDCl_3) δ (ppm): 7.19–7.15 (m, 1 H), 7.13–7.07 (m, 1 H), 7.01–6.95 (m, 2 H), 6.61 (s, 1 H), 6.54–6.27 (m, 3 H), 4.42–4.37 (m, 2 H), 4.36–4.30 (m, 1 H), 4.09–4.03 (m, 2 H), 3.78 (s, 3 H), 3.19–3.10 (m, 2 H), 2.81–2.71 (m, 4 H), 2.06–1.94 (m, 2 H), 1.73–1.59 (m, 2 H), 1.41–1.35 (m, 3 H); MS (ESI): m/z : 451 $[\text{M}+\text{H}]^+$.
20	5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR ($\text{DMSO}-d_6$) δ (ppm): 8.05–8.79 (m, 2 H), 7.79 (s, 1 H), 7.60–7.52 (m, 1 H), 7.32–7.27 (m, 1 H), 7.07 (s, 1 H), 6.94–6.72 (m, 4 H), 4.54–4.41 (m, 3 H), 4.35–4.24 (m, 2 H), 3.78 (s, 3 H), 3.24–2.95 (m, 4 H), 2.80–2.67 (m, 2 H), 2.07–1.69 (m, 4 H), 1.30–1.21 (m, 3 H); MS (ESI): m/z : 451 $[\text{M}+\text{H}]^+$.
21	5-[5-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		^1H NMR (CDCl_3) δ (ppm): 7.19–7.15 (m, 1 H), 7.00–6.95 (m, 2 H), 6.80–6.56 (m, 5 H), 4.40–4.34 (m, 2 H), 4.27–4.19 (m, 1 H), 4.05–4.00 (m, 2 H), 3.77 (s, 3 H), 3.21–3.09 (m, 2 H), 2.81–2.70 (m, 4 H), 2.02–1.60 (m, 4 H), 1.38 (m, 3 H); MS (ESI): m/z : 451 $[\text{M}+\text{H}]^+$.



22	5-[4-ethyl-1-[2-[4-(pyrrolidin-3-ylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 9.18–8.89 (m, 2 H), 7.78 (s, 1 H), 7.63–7.46 (m, 1 H), 7.32–7.27 (m, 1 H), 7.06 (s, 1 H), 6.90–6.72 (m, 4 H), 4.53–4.41 (m, 2 H), 4.54–4.41 (m, 2 H), 3.95–3.92 (m, 2 H), 3.77 (s, 3 H), 3.32–2.91 (m, 4 H), 2.80–2.56 (m, 3 H), 2.14–1.65 (m, 2 H), 1.31–1.23 (m, 3 H); MS (ESI): m/z: 451 [M+H] ⁺ .
23	5-[4-ethyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.84 (bs, 1 H), 8.57 (bs, 1 H), 7.78 (bs, 1 H), 7.56 (d, J=5.4 Hz, 1 H), 7.29 (d, J=5.4 Hz, 1 H), 7.06 (bs, 1 H), 6.85–6.71 (m, 4 H), 4.47 (t, J=4.6 Hz, 2 H), 4.29 (t, J=4.6 Hz, 2 H), 3.80–3.71 (m, 5 H), 3.32–3.21 (m, 2 H), 2.94–2.80 (m, 2 H), 2.73 (q, J=7.5 Hz, 2 H), 2.06–1.94 (m, 1 H), 1.92–1.83 (m, 2 H), 1.53–1.39 (m, 2 H), 1.27 (t, J=7.6 Hz, 3 H); MS (ESI): m/z: 465 [M+H] ⁺ .




24	5-[5-ethyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole trifluoroacetate		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.74–7.70 (m, 1 H), 7.54 (d, <i>J</i> =5.4 Hz, 1 H), 7.28 (d, <i>J</i> =5.4 Hz, 1 H), 7.09 (s, 1 H), 6.79–6.64 (m, 4 H), 4.56 (t, <i>J</i> =4.9 Hz, 2 H), 4.15 (t, <i>J</i> =4.9 Hz, 2 H), 3.72 (d, <i>J</i> =6.4 Hz, 2 H), 3.67 (s, 3 H), 3.32–3.24 (m, 2 H), 2.93–2.80 (m, 4 H), 2.03–1.92 (m, 1 H), 1.91–1.83 (m, 2 H), 1.47–1.35 (m, 2 H), 1.32 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 465 [M+H] ⁺ .
25	5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.91 (bs, 2 H), 7.79 (s, 1 H), 7.56 (d, <i>J</i> =4.9 Hz, 1 H), 7.30 (d, <i>J</i> =5.4 Hz, 1 H), 7.06 (bs, 1 H), 6.90–6.74 (m, 4 H), 4.60–4.41 (m, 3 H), 4.29 (t, <i>J</i> =4.6 Hz, 2 H), 3.77 (s, 3 H), 3.25–3.01 (m, 4 H), 2.72 (q, <i>J</i> =7.3 Hz, 2 H), 2.18–2.04 (m, 1 H), 2.03–1.75 (m, 4 H), 1.74–1.61 (m, 1 H), 1.26 (t, <i>J</i> =7.6 Hz, 3 H); MS (ESI): <i>m/z</i> : 465 [M+H] ⁺ .
26	5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole trifluoroacetate		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.67 (s, 1 H), 7.53 (d, <i>J</i> =5.3 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.11–7.03 (m, 1 H), 6.84–6.64 (m, 4 H), 4.60–4.41 (m, 3 H), 4.14 (t, <i>J</i> =4.9 Hz, 2 H), 3.67 (s, 3 H), 3.26–3.00 (m, 4 H), 2.83 (q, <i>J</i> =7.3 Hz, 2 H),



			2.13–2.01 (m, 1 H), 1.99–1.72 (m, 4 H), 1.72–1.61 (m, 1 H), 1.31 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 465 $[M+H]^+$.
27	5-[4-ethyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 +D $_2$ O): 7.66 (s, 1 H), 7.51 (d, $J=5.4$ Hz, 1 H), 7.25 (d, $J=5.4$ Hz, 1 H), 7.05–6.97 (m, 2 H), 6.83–6.75 (m, 3 H), 4.56–4.40 (m, 1 H), 4.03 (t, $J=7.5$ Hz, 2 H), 3.71 (s, 3 H), 3.26–3.15 (m, 2 H), 3.09–2.99 (m, 2 H), 2.68 (q, $J=7.3$ Hz, 2 H), 2.45 (t, $J=7.8$ Hz, 2 H), 2.08–1.97 (m, 4 H), 1.82–1.67 (m, 2 H), 1.24 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 449 $[M+H]^+$.
28	5-[5-ethyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 +D $_2$ O) δ (ppm): 7.61 (s, 1 H), 7.53 (d, $J=5.4$ Hz, 1 H), 7.25 (d, $J=5.4$ Hz, 1 H), 7.02–6.95 (m, 2 H), 6.85 (s, 1 H), 6.81–6.73 (m, 2 H), 4.55–4.47 (m, 1 H), 4.03 (t, $J=7.8$ Hz, 2 H), 3.67 (s, 3 H), 3.26–3.16 (m, 2 H), 3.09–3.00 (m, 2 H), 2.69 (q, $J=7.3$ Hz, 2 H), 2.45 (t, $J=6.9$ Hz, 2 H), 2.10–1.98 (m, 2 H), 1.96–1.85 (m, 2 H), 1.81–1.69 (m, 2 H), 1.27 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 449 $[M+H]^+$.




29	5-[4-cyclobutyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.81 (s, 1 H), 7.54 (d, J=5.4 Hz, 1 H), 7.27 (d, J=4.9 Hz, 1 H), 7.04 (s, 1 H), 6.91–6.84 (m, 2 H), 6.81–6.73 (m, 2 H), 4.51–4.39 (m, 3 H), 4.32–4.25 (m, 2 H), 3.73 (s, 3 H), 3.67–3.58 (m, 1 H), 3.24–2.97 (m, 4 H), 2.39–1.68 (m, 10 H); MS (ESI): m/z: 477 [M+H] ⁺ .
30	5-[5-cyclobutyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.82 (s, 1 H), 7.52 (d, J=5.4 Hz, 1 H), 7.25 (d, J=5.4 Hz, 1 H), 7.06 (s, 1 H), 6.85–6.79 (m, 2 H), 6.68–6.61 (m, 2 H), 4.50–4.38 (m, 3 H), 4.11–4.05 (m, 2 H), 3.85–3.73 (m, 1 H), 3.66 (s, 3 H), 3.25–2.93 (m, 4 H), 2.46–1.63 (m, 10 H) MS (ESI): m/z: 477 [M+H] ⁺ .
31	5-[5-ethyl-4-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.79–7.70 (m, 2 H), 7.47–7.40 (m, 2 H), 7.32–7.28 (m, 1 H), 7.19 (d, J=5.4 Hz, 1 H), 6.99 (d, J=5.4 Hz, 1 H), 6.75–6.69 (m, 2 H), 6.67–6.59 (m, 3 H), 4.53–4.39 (m, 3 H), 4.06 (t, J=5.9 Hz, 2 H), 3.83 (s, 3 H), 3.39–3.19 (m, 4 H), 3.00 (q, J=7.8 Hz, 2 H), 2.27–2.15 (m, 2 H), 2.13–2.02 (m, 2 H), 1.37 (t, J=7.6 Hz, 3 H); MS (ESI): m/z: 527 [M+H] ⁺ .




32	5-[5-ethyl-4-phenyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.77–7.71 (m, 2 H), 7.46–7.39 (m, 2 H), 7.32–7.27 (m, 1 H), 7.18 (d, <i>J</i> =5.4 Hz, 1 H), 7.13–7.07 (m, 1 H), 6.99 (d, <i>J</i> =5.4 Hz, 1 H), 6.66 (s, 1 H), 6.50–6.44 (m, 1 H), 6.36–6.31 (m, 1 H), 6.30–6.26 (m, 1 H), 4.54–4.42 (m, 3 H), 4.09 (t, <i>J</i> =5.9 Hz, 2 H), 3.82 (s, 3 H), 3.33–3.21 (m, 2 H), 3.17–3.07 (m, 2 H), 3.00 (q, <i>J</i> =7.5 Hz, 2 H), 2.23–2.11 (m, 2 H), 2.05–1.91 (m, 2 H), 1.37 (t, <i>J</i> =7.6 Hz, 3 H); MS (ESI): <i>m/z</i> : 527 [M+H] ⁺ .
33	5-[4-ethyl-5-phenyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.53–7.48 (m, 2 H), 7.47–7.38 (m, 3 H), 7.18 (d, <i>J</i> =4.9 Hz, 1 H), 7.05–6.96 (m, 2 H), 6.66 (s, 1 H), 6.47–6.39 (m, 1 H), 6.14–6.06 (m, 2 H), 4.52–4.35 (m, 3 H), 3.85 (s, 3 H), 3.75 (t, <i>J</i> =6.1 Hz, 2 H), 3.31–3.20 (m, 2 H), 3.12 (bs, 2 H), 2.58 (q, <i>J</i> =7.2 Hz, 2 H), 2.15 (bs, 2 H), 1.95 (bs, 2 H), 1.23 (t, <i>J</i> =7.6 Hz, 3 H); MS (ESI): <i>m/z</i> : 527 [M+H] ⁺ .
34	5-[1-[2-[3-(azepan-4-yloxy)phenoxy]ethyl]-4-ethylimidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (DMSO- <i>d</i> ₆ +D ₂ O) δ (ppm): 7.75 (s, 1 H), 7.54 (d, <i>J</i> =5.4 Hz, 1 H), 7.28 (d, <i>J</i> =4.9 Hz, 1 H), 7.17–7.11 (m, 1 H), 7.04 (s, 1 H), 6.56–6.51 (m, 1 H), 6.47–6.40 (m, 2 H), 4.67–



	b]pyrrole hydrochloride		4.59 (m, 1 H), 4.47 (t, $J=4.9$ Hz, 2 H), 4.32 (t, $J=4.9$ Hz, 2 H), 3.74 (s, 3 H), 3.26–3.04 (m, 4 H), 2.71 (q, $J=7.3$ Hz, 2 H), 2.16–2.06 (m, 1 H), 2.02–1.91 (m, 2 H), 1.91–1.75 (m, 2 H), 1.74–1.64 (m, 1 H), 1.25 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 465 $[M+H]^+$.
35	5-[1-[2-[3-(azepan-4-yloxy)phenoxy]ethyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 +D $_2$ O) δ (ppm): 7.64 (bs, 1 H), 7.53 (d, $J=5.4$ Hz, 1 H), 7.28 (d, $J=5.4$ Hz, 1 H), 7.14–7.05 (m, 2 H), 6.53–6.47 (m, 1 H), 6.36–6.27 (m, 2 H), 4.64–4.51 (m, 3 H), 4.18 (t, $J=4.9$ Hz, 2 H), 3.67 (s, 3 H), 3.24–3.02 (m, 4 H), 2.83 (q, $J=7.3$ Hz, 2 H), 2.15–2.05 (m, 1 H), 2.01–1.73 (m, 4 H), 1.72–1.64 (m, 1 H), 1.31 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 465 $[M+H]^+$.
36	5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 +D $_2$ O) δ (ppm): 7.83 (bs, 1 H), 7.53 (d, $J=5.4$ Hz, 1 H), 7.28 (d, $J=5.4$ Hz, 1 H), 7.05 (s, 1 H), 6.86–6.72 (m, 4 H), 4.55–4.40 (m, 3 H), 4.26 (t, $J=4.9$ Hz, 2 H), 3.73 (s, 3 H), 3.67–3.58 (m, 1 H), 3.24–3.16 (m, 1 H), 3.13–3.01 (m, 3 H), 2.38–2.28 (m, 2 H), 2.25–2.14 (m, 2 H), 2.13–1.61




			(m, 8 H); MS (ESI): m/z: 491 [M+H] ⁺ .
37	5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.17 (d, <i>J</i> =5.4 Hz, 1 H), 7.04 (s, 1 H), 6.98 (d, <i>J</i> =4.9 Hz, 1 H), 6.77–6.58 (m, 5 H), 4.57–4.49 (m, 1 H), 4.31 (t, <i>J</i> =5.9 Hz, 2 H), 3.99 (t, <i>J</i> =5.9 Hz, 2 H), 3.77 (s, 3 H), 3.64–3.56 (m, 1 H), 3.35–3.12 (m, 4 H), 2.49–2.38 (m, 2 H), 2.32–1.76 (m, 10 H); MS (ESI): m/z: 491 [M+H] ⁺ .
38	5-[4-ethyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.15 (d, <i>J</i> =5.4 Hz, 1 H), 7.02–6.94 (m, 3 H), 6.81–6.73 (m, 3 H), 6.35 (s, 1 H), 3.97 (t, <i>J</i> =7.4 Hz, 2 H), 3.94–3.81 (m, 5 H), Part AB of ABX System: VA=3.28, VB=2.98, JAB= 11.7 Hz, JAX= 7.8 Hz, JBX= 6.3 Hz, 3.22–3.15 (m, 1 H), 3.13–3.05 (m, 1 H), 2.72–2.63 (m, 3 H), 2.54 (t, <i>J</i> =7.3 Hz, 2 H), 2.14–1.99 (m, 3 H), 1.77–1.66 (m, 1 H), 1.28 (t, <i>J</i> =7.6 Hz, 3 H); MS (ESI): m/z: 449 [M+H] ⁺ .
39	5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole		¹ H NMR (DMSO-d ₆) δ (ppm): 7.13 (s, 1 H), 7.08 (d, <i>J</i> =5.4 Hz, 1 H), 7.04 (d, <i>J</i> =5.4 Hz, 1 H), 6.86–6.72 (m, 4 H), 6.63 (s, 1 H), 4.34–4.27 (m, 2 H), 4.25–4.12 (m, 3 H), 3.72 (s, 3 H), 2.98–2.89 (m, 2 H), 2.60–2.50



			(m, 4 H), 1.89–1.76 (m, 2 H), 1.45–1.34 (m, 2 H), 1.18 (t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 451 $[M+H]^+$.
40	4-methyl-5-[5-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-4-propylimidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		^1H NMR (D_2O) δ (ppm): 7.51–7.36 (m, 5 H), 7.26 (d, $J=5.4$ Hz, 1 H), 6.93 (d, $J=5.4$ Hz, 1 H), 6.77 (s, 1 H), 6.62–6.55 (m, 2 H), 6.40–6.34 (m, 2 H), 4.47 (t, $J=4.9$ Hz, 2 H), 4.36–4.27 (m, 1 H), 3.75 (t, $J=4.9$ Hz, 2 H), 3.53 (s, 3 H), 3.29–3.20 (m, 2 H), 3.09–3.00 (m, 2 H), 2.49 (t, $J=7.3$ Hz, 2 H), 2.01–1.89 (m, 2 H), 1.83–1.73 (m, 2 H), 1.54–1.43 (m, 2 H), 0.71 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 541 $[M+H]^+$.
41	4-methyl-5-[4-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-5-propylimidazol-2-yl]thieno[3,2-b]pyrrole		^1H NMR ($\text{DMSO}-d_6$) δ (ppm): 7.71–7.66 (m, 2 H), 7.45–7.38 (m, 2 H), 7.35 (d, $J=5.4$ Hz, 1 H), 7.28–7.23 (m, 1 H), 7.22 (d, $J=5.4$ Hz, 1 H), 6.80 (s, 1 H), 6.78–6.73 (m, 2 H), 6.68–6.63 (m, 2 H), 4.45 (t, $J=5.6$ Hz, 2 H), 4.25–4.16 (m, 1 H), 4.07 (t, $J=5.6$ Hz, 2 H), 3.77 (s, 3 H), 2.98–2.83 (m, 4 H), 2.61–2.52 (m, 2 H), 1.88–1.78 (m, 2 H), 1.69–1.57 (m, 2 H), 1.47–1.34 (m, 2 H), 0.98 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 541 $[M+H]^+$.



42	5-[4-ethyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (D ₂ O) δ (ppm): 7.33 (d, <i>J</i> =5.4 Hz, 1 H), 7.24 (s, 1 H), 7.01 (d, <i>J</i> =5.4 Hz, 1 H), 6.83–6.77 (m, 2 H), 6.58–6.52 (m, 3 H), 3.93 (t, <i>J</i> =7.3 Hz, 2 H), 3.70 (d, <i>J</i> =6.4 Hz, 2 H), 3.52 (s, 3 H), 3.40–3.33 (m, 2 H), 2.96–2.86 (m, 2 H), 2.59 (q, <i>J</i> =7.8 Hz, 2 H), 2.40 (t, <i>J</i> =7.3 Hz, 2 H), 2.05–1.89 (m, 5 H), 1.49–1.36 (m, 2 H), 1.14 (t, <i>J</i> =7.8 Hz, 3 H); MS (ESI): <i>m/z</i> : 463 [M+H] ⁺ .
43	5-[5-ethyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.15 (d, <i>J</i> =5.4 Hz, 1 H), 7.00–6.95 (m, 3 H), 6.94 (s, 1 H), 6.79–6.73 (m, 2 H), 6.39 (s, 1 H), 3.94 (t, <i>J</i> =7.3 Hz, 2 H), 3.80 (s, 3 H), 3.77 (d, <i>J</i> =6.4 Hz, 2 H), 3.19–3.11 (m, 2 H), 2.72–2.64 (m, 2 H), 2.57 (q, <i>J</i> =7.5 Hz, 2 H), 2.52 (t, <i>J</i> =7.3 Hz, 2 H), 1.98–1.90 (m, 3 H), 1.88–1.82 (m, 2 H), 1.36–1.25 (m, 5 H); MS (ESI): <i>m/z</i> : 463 [M+H] ⁺ .
44	5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-4-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO- <i>d</i> ₆) δ (ppm): 9.11–8.86 (m, 2 H), 7.76 (s, 1 H), 7.56 (d, <i>J</i> =5.4 Hz, 1 H), 7.29 (d, <i>J</i> =5.4 Hz, 1 H), 7.05–6.99 (m, 2 H), 6.89 (bs, 1 H), 6.80–6.64 (m, 2 H), 4.61–4.53 (m, 1 H), 4.07 (t, <i>J</i> =7.3 Hz, 2 H), 3.77 (s, 3 H), 3.24–3.02 (m, 4 H), 2.71 (q, <i>J</i> =7.3 Hz, 2 H), 2.50–



			2.44 (m, 2 H), 2.19–1.63 (m, 8 H), 1.27 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 463 [M+H] ⁺ .
45	5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.99–8.78 (m, 2 H), 7.71–7.61 (bs, 1 H), 7.55 (d, $J=5.4$ Hz, 1 H), 7.27 (d, $J=5.4$ Hz, 1 H), 7.03–7.96 (m, 2 H), 6.90 (s, 1 H), 6.78–6.72 (m, 2 H), 4.64–4.53 (m, 1 H), 4.09–4.00 (m, 2 H), 3.71 (s, 3 H), 3.24–3.03 (m, 4 H), 2.76–2.65 (m, 2 H), 2.49–2.42 (m, 2 H), 2.18–1.63 (m, 8 H), 1.28 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 463 [M+H] ⁺ .
46	5-[5-ethyl-4-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.67–7.45 (m, 6 H), 7.28 (d, $J=4.9$ Hz, 1 H), 7.03–6.98 (m, 2 H), 6.96 (bs, 1 H), 6.81–6.76 (m, 2 H), 4.57–4.49 (m, 1 H), 4.18–4.09 (m, 2 H), 3.77 (s, 3 H), 3.25–3.01 (m, 4 H), 2.86 (q, $J=7.7$ Hz, 2 H), 2.54–2.46 (m, 2 H), 2.09–1.99 (m, 2 H), 1.97–1.88 (m, 2 H), 1.82–1.71 (m, 2 H), 1.18 (t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 525 [M+H] ⁺ .
47	5-[4-ethyl-5-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.60–7.50 (m, 6 H), 7.29 (d, $J=5.4$ Hz, 1 H), 6.95 (s, 1 H), 6.76–6.67 (m, 4 H), 4.53–4.46 (m, 1 H), 4.02 (t, $J=7.3$ Hz, 2 H), 3.79 (s, 3 H), 3.23–3.01 (m, 4



	b]pyrrole hydrochloride		H), 2.59 (q, $J=7.3$ Hz, 2 H), 2.18 (t, $J=7.3$ Hz, 2 H), 2.06–1.69 (m, 4 H), 1.60–1.52 (m, 2 H), 1.17 (t, $J=7.6$ Hz, 3 H); MS (ESI): m/z : 525 $[M+H]^+$.
48	5-[4-isopropyl-5-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		^1H NMR (CDCl_3) δ (ppm): 7.53–7.36 (m, 5 H), 7.16 (d, $J=5.4$ Hz, 1 H), 6.98 (d, $J=5.4$ Hz, 1 H), 6.70–6.62 (m, 3 H), 6.45–6.36 (m, 2 H), 4.35 (t, $J=6.4$ Hz, 2 H), 4.30–4.23 (m, 1 H), 3.83 (s, 3 H), 3.74 (t, $J=6.4$, 2 H), 3.24–3.14 (m, 2 H), 2.95–2.85 (m, 3 H), 2.09–2.00 (m, 2 H), 1.82–1.72 (m, 2 H), 1.27 (d, $J=6.8$ Hz, 6 H); MS (ESI): m/z : 541 $[M+H]^+$.
49	5-[4-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR ($\text{DMSO}-d_6$) δ (ppm): 8.84–8.64 (m, 2 H), 7.87 (s, 1 H), 7.60–7.51 (m, 1 H), 7.27 (d, $J=4.9$ Hz, 1 H), 7.07–7.01 (m, 2 H), 6.90 (s, 1 H), 6.85–6.79 (m, 2 H), 4.57–4.49 (m, 1 H), 4.11–4.01 (m, 2 H), 3.76 (s, 3 H), 3.66–3.57 (m, 1 H), 3.25–3.15 (m, 2 H), 3.09–2.96 (m, 2 H), 2.51–2.45 (m, 2 H), 2.40–2.29 (m, 2 H), 2.27–2.18 (m, 2 H), 2.13–1.70 (m, 8 H); MS (ESI): m/z : 475 $[M+H]^+$.



50	5-[5-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.16 (d, <i>J</i> =5.4 Hz, 1 H), 7.02–6.93 (m, 4 H), 6.83–6.77 (m, 2 H), 6.42 (s, 1 H), 4.39–4.30 (m, 1 H), 3.91–3.84 (m, 2 H), 3.80 (s, 3 H), 3.42–3.31 (m, 1 H), 3.20–3.13 (m, 2 H), 2.82–2.73 (m, 2 H), 2.51 (t, <i>J</i> =7.3 Hz, 2 H), 2.34–2.26 (m, 2 H), 2.25–2.13 (m, 2 H), 2.08–1.84 (m, 6 H), 1.70 (m, 2 H); MS (ESI): <i>m/z</i> : 475 [M+H] ⁺ .
51	4-methyl-5-[4-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 8.25 (s, 1 H), 7.90–7.72 (m, 2 H), 7.54–7.45 (m, 3 H), 7.43–7.36 (m, 1 H), 7.26 (d, <i>J</i> =5.4 Hz, 1 H), 7.07–7.00 (m, 2 H), 6.85–6.76 (m, 3 H), 4.55–4.46 (m, 1 H), 4.12 (t, <i>J</i> =7.3 Hz, 2 H), 3.81 (s, 3 H), 3.24–3.15 (m, 2 H), 3.07–2.99 (m, 2 H), 2.53 (t, <i>J</i> =7.3 Hz, 2 H), 2.15–2.06 (m, 2 H), 2.05–1.96 (m, 2 H), 1.80–1.68 (m, 2 H); MS (ESI): <i>m/z</i> : 497 [M+H] ⁺ .
52	5-[1-[3-[3,4-bis(4-piperidylmethoxy)phenyl]propyl]-4-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole dihydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.66 (s, 1 H), 7.53 (d, <i>J</i> =5.4 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 6.83–6.75 (m, 2 H), 6.75–6.70 (m, 1 H), 6.60–6.55 (m, 1 H), 4.04 (t, <i>J</i> =7.3 Hz, 2 H), 3.77–3.69 (m, 7 H), 3.32–3.23 (m, 4 H), 2.93–2.81 (m, 4 H),

			2.68 (q, $J=7.3$ Hz, 2 H), 2.44 (t, $J=7.1$ Hz, 2 H), 2.10–1.94 (m, 4 H), 1.93–1.82 (m, 4 H), 1.51–1.37 (m, 4 H), 1.24 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 576 $[M+H]^+$.
53	4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-4-propylimidazol-2-yl]thieno[3,2-b]pyrrole		^1H NMR (DMSO- d_6) δ (ppm): 8.93–8.65 (m, 2 H), 7.78 (bs, 1 H), 7.55 (d, $J=5.4$ Hz, 1 H), 7.29 (d, $J=5.4$ Hz, 1 H), 7.06 (bs, 1 H), 6.94–6.73 (m, 4 H), 4.53–4.43 (m, 3 H), 4.29 (t, $J=4.6$ Hz, 2 H), 3.77 (s, 3 H), 3.25–3.14 (m, 2 H), 3.12–2.94 (m, 2 H), 2.68 (t, $J=7.3$ Hz, 2 H), 2.09–1.96 (m, 2 H), 1.83–1.60 (m, 4 H), 0.95 (t, $J=7.3$ Hz, 3 H) MS (ESI): m/z : 465 $[M+H]^+$.
54	4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-5-propylimidazol-2-yl]thieno[3,2-b]pyrrole		^1H NMR (DMSO- d_6) δ (ppm): 7.17 (d, $J=5.4$ Hz, 1 H), 7.01–6.93 (m, 2 H), 6.79–6.73 (m, 2 H), 6.67–6.55 (m, 3 H), 4.37 (t, $J=5.9$ Hz, 2 H), 4.27–4.17 (m, 1 H), 4.02 (t, $J=5.9$ Hz, 2 H), 3.76 (s, 3 H), 3.22–3.10 (m, 2 H), 3.79–2.71 (m, 2 H), 2.68 (t, $J=7.3$ Hz, 2 H), 2.05–1.94 (m, 2 H), 1.79 (sxt, $J=7.3$ Hz, 2 H), 1.71–1.61 (m, 2 H), 1.09 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 465 $[M+H]^+$.

55	5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.99–8.76 (m, 2 H), 7.77 (s, 1 H), 7.56 (d, J=5.4 Hz, 1 H), 7.29 (d, J=5.4 Hz, 1 H), 7.06–6.98 (m, 2 H), 6.90 (s, 1 H), 6.81–6.73 (m, 2 H), 4.62–4.51 (m, 1 H), 4.07 (t, J=7.1 Hz, 2 H), 3.76 (s, 3 H), 3.25–3.02 (m, 4 H), 2.66 (t, J=7.1 Hz, 2 H), 2.50–2.41 (m, 2 H), 2.17–1.59 (m, 10 H), 0.96 (t, J=7.6 Hz, 3 H); MS (ESI): m/z: 477 [M+H] ⁺ .
56	5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.94–8.71 (m, 2 H), 7.62 (s, 1 H), 7.54 (d, J=5.4 Hz, 1 H), 7.27 (d, J=5.4 Hz, 1 H), 7.04–6.97 (m, 2 H), 6.89 (s, 1 H), 6.79–6.71 (m, 2 H), 4.63–4.54 (m, 1 H), 4.10–4.00 (m, 2 H), 3.71 (s, 3 H), 3.25–3.03 (m, 4 H), 2.67–2.58 (m, 2 H), 2.50–2.44 (m, 2 H), 2.18–1.61 (m, 10 H), 1.00 (s, 3 H); MS (ESI): m/z: 477 [M+H] ⁺ .



57	5-[1-[3-[3,4-bis(4-piperidylmethoxy)phenyl]propyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole dihydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 7.59–7.43 (m, 2 H), 7.21 (d, <i>J</i> =5.4 Hz, 1 H), 6.80–6.69 (m, 3 H), 6.60–6.52 (m, 1 H), 4.06–3.97 (m, 2 H), 3.76–3.65 (m, 7 H), 3.32–3.24 (m, 4 H), 2.93–2.83 (m, 4 H), 2.68 (q, <i>J</i> =7.3 Hz, 2 H), 2.44 (t, <i>J</i> =1.0 Hz, 2 H), 2.05–1.84 (m, 8 H), 1.51–1.37 (m, 4 H), 1.26 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 576 [M+H] ⁺ .
58	5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenyl]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.13 (d, <i>J</i> =5.4 Hz, 1 H), 6.98–6.88 (m, 3 H), 6.82–6.74 (m, 2 H), 6.71 (s, 1 H), 6.27 (s, 1 H), 4.36–4.28 (m, 1 H), 4.18 (t, <i>J</i> =7.3 Hz, 2 H), 3.71 (s, 3 H), 3.19–3.09 (m, 2 H), 2.95 (t, <i>J</i> =7.3 Hz, 2 H), 2.79–2.70 (m, 2 H), 2.65 (q, <i>J</i> =7.6 Hz, 2 H), 2.07–2.00 (m, 2 H), 1.72–1.59 (m, 2 H), 1.27 (t, <i>J</i> =7.6 Hz, 3 H); MS (ESI): <i>m/z</i> : 435 [M+H] ⁺ .



59	5-[4-cyclobutyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.75 (s, 1 H), 7.52 (d, <i>J</i> =5.4 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.06–6.97 (m, 2 H), 6.81 (s, 1 H), 6.77–6.71 (m, 2 H), 4.03 (t, <i>J</i> =7.3 Hz, 2 H), 3.93–3.81 (m, 2 H), 3.71 (s, 3 H), 3.58–3.54 (m, 1 H), Part AB of ABX System: VA=3.34, VB=2.97, JAB= 11 Hz, JAX= 6.8 Hz, JBX= 7.4 Hz, 3.29–3.21 (m, 1 H), 3.20–3.12 (m, 1 H), 2.73–2.64 (m, 1 H), 2.46 (t, <i>J</i> =7.3 Hz, 2 H), 2.37–2.28 (m, 2 H), 2.24–2.14 (m, 2 H), 2.13–1.95 (m, 4 H), 1.92–1.82 (m, 1 H), 1.78–1.67 (m, 1 H); MS (ESI): <i>m/z</i> : 475 [M+H] ⁺ .
60	5-[5-cyclobutyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.15 (d, <i>J</i> =5.4 Hz, 1 H), 7.02–6.91 (m, 4 H), 6.82–6.74 (m, 2 H), 6.38 (s, 1 H), 3.93–3.76 (m, 7 H), 3.42–3.33 (m, 1 H), Part AB of ABX System: VA=3.15, VB=2.84, JAB= 11.2 Hz, JAX= 7.9 Hz, JBX= 5.8 Hz, 3.08–3.01 (m, 1 H), 2.99–2.91 (m, 1 H), 2.63–2.55 (m,



			1 H), 2.51 (t, $J=7.3$ Hz, 2 H), 2.36–1.84 (m, 9 H), 1.65–1.53 (m, 1 H); MS (ESI): m/z : 475 $[M+H]^+$.
61	5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-4-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 +D $_2$ O) δ (ppm): 7.82 (bs, 1 H), 7.54 (d, $J=5.4$ Hz, 1 H), 7.27 (d, $J=5.4$ Hz, 1 H), 7.05–6.98 (m, 2 H), 6.87 (s, 1 H), 6.79–6.69 (m, 2 H), 4.62–4.50 (m, 1 H), 4.05 (t, $J=7.3$ Hz, 2 H), 3.73 (s, 3 H), 3.64–3.54 (m, 1 H), 3.25–3.03 (m, 4 H), 2.46 (t, $J=7.3$ Hz, 2 H), 2.39–2.29 (m, 2 H), 2.26–2.16 (m, 2 H), 2.15–1.64 (m, 10 H); MS (ESI): m/z : 489 $[M+H]^+$.
62	5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 +D $_2$ O) δ (ppm): 7.68 (bs, 1 H), 7.50 (d, $J=5.4$ Hz, 1 H), 7.24 (d, $J=5.4$ Hz, 1 H), 7.03–6.95 (m, 2 H), 6.84 (s, 1 H), 6.78–6.72 (m, 2 H), 4.62–4.53 (m, 1 H), 3.93 (t, $J=7.3$ Hz, 2 H), 3.68 (s, 3 H), 3.26–3.04 (m, 5 H), 2.44 (t, $J=7.3$ Hz, 2 H), 2.32–1.62 (m, 14 H); MS (ESI): m/z : 489 $[M+H]^+$.



63	4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-4-propylimidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.73 (s, 1 H), 7.55 (d, <i>J</i> =5.4 Hz, 1 H), 7.26 (d, <i>J</i> =5.4 Hz, 1 H), 7.05–6.98 (m, 2 H), 6.88 (s, 1 H), 6.84–6.73 (m, 2 H), 4.59–4.48 (m, 1 H), 4.07 (t, <i>J</i> =7.3 Hz, 2 H), 3.73 (s, 3 H), 3.26–3.16 (m, 2 H), 3.11–3.00 (m, 2 H), 2.65 (t, <i>J</i> =7.3 Hz, 2 H), 2.45 (t, <i>J</i> =7.3 Hz, 2 H), 2.10–1.99 (m, 4 H), 1.80–1.63 (m, 4 H), 0.95 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 463 [M+H] ⁺ .
64	4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-5-propylimidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.61–7.46 (m, 2 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.04–6.95 (m, 2 H), 6.85 (s, 1 H), 6.81–6.75 (m, 2 H), 4.57–4.46 (m, 1 H), 4.00 (t, <i>J</i> =7.3 Hz, 2 H), 3.67 (s, 3 H), 3.26–3.14 (m, 2 H), 3.10–2.97 (m, 2 H), 2.59 (t, <i>J</i> =7.3 Hz, 2 H), 2.45 (t, <i>J</i> =7.3 Hz, 2 H), 2.09–1.97 (m, 2 H), 1.94–1.83 (m, 2 H), 1.80–1.71 (m, 2 H), 1.69–1.58 (m, 2 H), 0.98 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 463 [M+H] ⁺ .



65	5-[4-cyclobutyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.77 (s, 1 H), 7.54 (d, <i>J</i> =5.4 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.04–6.95 (m, 2 H), 6.81 (s, 1 H), 6.75–6.69 (m, 2 H), 4.03 (t, <i>J</i> =7.3 Hz, 2 H), 3.76–3.68 (m, 5 H), 3.57–3.54 (m, 1 H), 3.32–3.25 (m, 2 H), 2.93–2.83 (m, 2 H), 2.45 (t, <i>J</i> =7.3 Hz, 2 H), 2.37–2.28 (m, 2 H), 2.25–2.14 (m, 2 H), 2.10–1.95 (m, 4 H), 1.92–1.84 (m, 3 H), 1.50–1.38 (m, 2 H); MS (ESI): <i>m/z</i> : 489 [M+H] ⁺ .
66	5-[5-cyclobutyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.63 (bs, 1 H), 7.48 (d, <i>J</i> =5.4 Hz, 1 H), 7.23 (d, <i>J</i> =5.4 Hz, 1 H), 7.01–6.93 (m, 2 H), 6.79–6.68 (m, 3 H), 3.92 (t, <i>J</i> =7.8 Hz, 2 H), 3.76 (d, <i>J</i> =6.4 Hz, 2 H), 3.67 (s, 3 H), 3.57–3.51 (m, 1 H), 3.33–3.23 (m, 2 H), 2.93–2.82 (m, 2 H), 2.44 (t, <i>J</i> =7.3 Hz, 2 H), 2.34–2.25 (m, 2 H), 2.21–2.11 (m, 2 H), 2.07–1.96 (m, 2 H), 1.94–1.80 (m, 5 H), 1.50–1.36 (m, 2 H); MS (ESI): <i>m/z</i> : 489 [M+H] ⁺ .



67	4-methyl-5-[4-propyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ +D ₂ O) δ (ppm): 7.69 (bs, 1 H), 7.53 (d, <i>J</i> =5.4 Hz, 1 H), 7.26 (d, <i>J</i> =5.4 Hz, 1 H), 7.04–6.97 (m, 2 H), 6.81 (bs, 1 H), 6.78–6.72 (m, 2 H), 4.05 (t, <i>J</i> =7.3 Hz, 2 H), 3.94–3.81 (m, 2 H), 3.72 (s, 3 H), Part AB of ABX System: VA=3.34, VB=2.97, JAB=11.6 Hz, JAX=8.1 Hz, JBX=7.3 Hz, 3.29–3.21 (m, 1 H), 3.20–3.12 (m, 1 H), 2.73–2.61 (m, 3 H), 2.46 (t, <i>J</i> =7.3 Hz, 2 H), 2.14–1.99 (m, 3 H), 1.77–1.62 (m, 3 H), 0.94 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 463 [M+H] ⁺ .
68	4-methyl-5-[5-propyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.16 (d, <i>J</i> =4.9 Hz, 1 H), 7.00–6.95 (m, 3 H), 6.93 (s, 1 H), 6.81–6.73 (m, 2 H), 6.38 (s, 1 H), 3.97–3.81 (m, 4 H), 3.79 (s, 3 H), Part AB of ABX System: VA=3.17, VB=2.86, JAB=11.3 Hz, JAX=7.9 Hz, JBX=5.9 Hz, 3.10–3.02 (m, 1 H), 3.01–2.94 (m, 1 H), 2.64–2.57 (m, 1 H), 2.56–2.45 (m, 4 H), 2.06–1.98 (m, 1 H), 1.97–



			1.89 (m, 2 H), 1.76–1.67 (m, 2 H), 1.65–1.55 (m, 1 H), 1.04 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 463 $[M+H]^+$.
69	5-[4-cyclobutyl-1-[2-[4-(4-piperidyl)oxy]phenyl]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		^1H NMR (CDCl_3) δ (ppm): 7.13 (d, $J=5.4$ Hz, 1 H), 6.96–6.90 (m, 3 H), 6.82–6.76 (m, 2 H), 6.73 (s, 1 H), 6.27 (s, 1 H), 4.42–4.33 (m, 1 H), 4.18 (t, $J=7.3$ Hz, 2 H), 3.73 (s, 3 H), 3.56–3.46 (m, 1 H), 3.24–3.13 (m, 2 H), 2.96 (t, $J=7.3$ Hz, 2 H), 2.90–2.79 (m, 2 H), 2.38–2.17 (m, 4 H), 2.11–1.71 (m, 6 H); MS (ESI): m/z : 461 $[M+H]^+$.
70	5-[4-ethyl-1-[3-[3-(4-piperidyl)oxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR ($\text{DMSO}-d_6$) δ (ppm): 8.80 (bs, 2 H), 7.79 (bs, 1 H), 7.56 (d, $J=5.4$ Hz, 1 H), 7.28 (d, $J=5.4$ Hz, 1 H), 7.16–7.06 (m, 1 H), 6.88 (bs, 1 H), 6.82–6.77 (m, 2 H), 6.74–6.67 (m, 1 H), 4.63–4.52 (m, 1 H), 4.09 (t, $J=7.3$ Hz, 2 H), 3.77 (s, 3 H), 3.23–3.15 (m, 2 H), 3.08–2.99 (m, 2 H), 2.72 (q, $J=7.3$ Hz, 2 H), 2.54 (t, $J=7.3$ Hz, 2 H), 2.16–1.99


			(m, 4 H), 1.82–1.71 (m, 2 H), 1.27 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 449 $[M+H]^+$.
71	5-[5-ethyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 +D $_2$ O) δ (ppm): 7.54–7.41 (m, 2 H), 7.22 (d, $J=5.4$ Hz, 1 H), 7.13–7.03 (m, 1 H), 6.82–6.63 (m, 4 H), 4.61–4.51 (m, 1 H), 4.03 (t, $J=7.3$ Hz, 2 H), 3.67 (s, 3 H), 3.26–3.14 (m, 2 H), 3.09–2.98 (m, 2 H), 2.65 (q, $J=7.3$ Hz, 2 H), 2.48–2.45 (m, 2 H), 2.08–1.98 (m, 2 H), 1.97–1.88 (m, 2 H), 1.80–1.68 (m, 2 H), 1.25 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 449 $[M+H]^+$.
72	4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenyl]ethyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 +D $_2$ O) δ (ppm): 7.72 (s, 1 H), 7.53 (d, $J=5.4$ Hz, 1 H), 7.23 (d, $J=5.4$ Hz, 1 H), 6.97–6.90 (m, 2 H), 6.86–6.78 (m, 2 H), 6.66 (s, 1 H), 4.59–4.50 (m, 1 H), 4.31 (t, $J=6.8$ Hz, 2 H), 3.50 (s, 3 H), 3.24–3.16 (m, 2 H), 3.08–2.97 (m, 4 H), 2.63 (t, $J=7.3$ Hz, 2 H), 2.08–1.98 (m, 2 H), 1.80–1.71 (m,


			2 H), 1.69–1.60 (m, 2 H), 0.93 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 449 $[M+H]^+$.
73	4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenyl]ethyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole		^1H NMR ($\text{CDCl}_3+\text{D}_2\text{O}$) δ (ppm): 7.16 (d, $J=5.4$ Hz, 1 H), 6.96 (d, $J=5.4$ Hz, 1 H), 6.93 (s, 1 H), 6.88–6.83 (m, 2 H), 6.77–6.72 (m, 2 H), 6.46 (s, 1 H), 4.36–4.26 (m, 1 H), 4.18 (t, $J=7.3$ Hz, 2 H), 3.70 (s, 3 H), 3.19–3.07 (m, 2 H), 2.81 (t, $J=7.3$ Hz, 2 H), 2.77–2.67 (m, 2 H), 2.48 (t, $J=7.3$ Hz, 2 H), 2.04–1.96 (m, 2 H), 1.78–1.62 (m, 4 H), 1.05 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 449 $[M+H]^+$.
74	4-methyl-5-[1-[3-[4-(4-piperidyloxymethyl)phenyl]propyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		^1H NMR ($\text{DMSO}-d_6$) δ (ppm): 8.87–8.72 (m, 1 H), 8.62–8.42 (m, 1 H), 7.84–7.64 (bs, 1 H), 7.56 (d, $J=5.4$ Hz, 1 H), 7.28 (d, $J=5.4$ Hz, 1 H), 7.30–7.24 (m, 2 H), 6.91–6.81 (bs, 1 H), 6.78–6.72 (m, 2 H), 4.06 (t, $J=7.3$ Hz, 2 H), 3.80–3.71 (m, 5 H), 3.34–3.23 (m, 2 H), 2.97–2.82 (m, 2 H), 2.66 (t, $J=7.3$ Hz, 2 H), 2.49–2.41 (m, 2 H), 2.13–1.96 (m, 3 H), 1.93–1.83 (m, 2 H), 1.75–1.63 (m, 2 H), 1.52–



			1.39 (m, 2 H), 0.96 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 477 $[M+H]^+$.
75	4-methyl-5-[1-[3-[4-(4-piperidyloxymethyl)phenyl]propyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6) δ (ppm): 8.85–8.68 (m, 1 H), 8.58–8.40 (m, 1 H), 7.71–7.62 (bs, 1 H), 7.56 (d, $J=5.4$ Hz, 1 H), 7.27 (d, $J=5.4$ Hz, 1 H), 7.03–6.97 (m, 2 H), 6.86 (bs, 1 H), 6.76–6.70 (m, 2 H), 4.04 (t, $J=7.3$ Hz, 2 H), 3.80–3.74 (m, 2 H), 3.71 (s, 3 H), 3.34–3.25 (m, 2 H), 2.96–2.80 (m, 2 H), 2.63 (t, $J=7.3$ Hz, 2 H), 2.49–2.43 (m, 2 H), 2.06–1.98 (m, 1 H), 1.95–1.84 (m, 4 H), 1.76–1.62 (m, 2 H), 1.54–1.39 (m, 2 H), 1.01 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 477 $[M+H]^+$.
76	5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		^1H NMR (CDCl_3) δ (ppm): 7.15 (d, $J=4.9$ Hz, 1 H), 6.97 (d, $J=4.9$ Hz, 1 H), 6.92 (s, 1 H), 6.83–6.70 (m, 4 H), 6.56 (s, 1 H), 4.53–4.46 (m, 1 H), 4.36 (t, $J=5.1$ Hz, 2 H), 4.14 (t, $J=5.1$ Hz, 2 H), 3.82 (s, 3 H), 3.21–2.98 (m, 2 H), 2.90–2.65 (m, 2 H), 2.60 (t, $J=7.3$ Hz, 2 H), 2.23–1.81 (m, 6 H), 1.78–1.66 (m, 2 H),



			1.00 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 479 $[M+H]^+$.
77	5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-propylimidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		^1H NMR (DMSO- d_6) δ (ppm): 7.32 (d, $J=5.4$ Hz, 1 H), 7.19 (d, $J=5.4$ Hz, 1 H), 6.86 (s, 1 H), 6.78–6.61 (m, 5 H), 4.46–4.39 (m, 1 H), 4.35 (t, $J=5.4$ Hz, 2 H), 4.03 (t, $J=5.4$ Hz, 2 H), 3.68 (s, 3 H), 3.07–2.79 (m, 4 H), 2.66 (t, $J=7.3$ Hz, 2 H), 2.10–1.45 (m, 8 H), 1.01 (t, $J=7.3$ Hz, 3 H); MS (ESI): m/z : 479 $[M+H]^+$.
78	5-[4-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole hydrochloride		^1H NMR (DMSO- d_6 + D_2O) δ (ppm): 7.74 (bs, 1 H), 7.20 (d, $J=5.4$ Hz, 1 H), 7.07 (d, $J=5.4$ Hz, 1 H), 7.04 - 6.99 (m, 2 H), 6.83 - 6.73 (m, 3 H), 4.57 - 4.46 (m, 1 H), 4.02 (t, $J=7.3$ Hz, 2 H), 3.67 (s, 3 H), 3.65 - 3.56 (m, 1 H), 3.25 - 3.16 (m, 2 H), 3.08 - 3.00 (m, 2 H), 2.45 (t, $J=7.3$ Hz, 2 H), 2.38 - 2.27 (m, 2 H), 2.24 - 2.13 (m, 2 H), 2.10 - 1.96 (m, 5 H), 1.91 - 1.82 (m, 1 H), 1.80 - 1.70 (m, 2 H); MS (ESI): m/z : 475 $[M+H]^+$.


79	5-[5-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole		¹ H NMR (CDCl ₃ + D ₂ O) δ (ppm): 7.04 - 6.94 (m, 4 H), 6.89 (d, <i>J</i> =5.4 Hz, 1 H), 6.83 - 6.76 (m, 2 H), 6.40 (s, 1 H), 4.39 - 4.27 (m, 1 H), 3.87 (t, <i>J</i> =7.8 Hz, 2 H), 3.75 (s, 3 H), 3.42 - 3.31 (m, 1 H), 3.19 - 3.09 (m, 2 H), 2.79 - 2.68 (m, 2 H), 2.51 (t, <i>J</i> =7.3 Hz, 2 H), 2.34 - 2.24 (m, 2 H), 2.23 - 2.12 (m, 2 H), 2.06 - 1.85 (m, 6 H), 1.73 - 1.62 (m, 2 H).; MS (ESI): <i>m/z</i> : 475 [M+H] ⁺ .
80	5-[4-cyclobutyl-1-[3-[4-[(3R)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.75 (s, 1 H), 7.53 (d, <i>J</i> =5.4 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.04 - 6.98 (m, 2 H), 6.81 (s, 1 H), 6.76 - 6.70 (m, 2 H), 4.03 (t, <i>J</i> =7.3 Hz, 2 H), Part AB of ABX System: VA=3.9, VB=3.85, <i>J</i> AB=9.5 Hz, <i>J</i> AX=6.1 Hz, <i>J</i> BX=7.1 Hz, 3.71 (s, 3 H), 3.58 - 3.53 (m, 1 H), Part AB of ABX System: VA=3.34, VB=2.97, <i>J</i> AB=11.6 Hz, <i>J</i> AX=8.1 Hz, <i>J</i> BX=7.3 Hz), 3.28 - 3.21 (m, 1 H), 3.20 - 3.12 (m, 1 H), 2.72 - 2.64 (m, 1 H), 2.45 (t, <i>J</i> =7.3 Hz, 2 H), 2.37 - 2.28 (m, 2 H), 2.24 -



			1.96 (m, 6 H), 1.93 - 1.83 (m, 1 H), 1.76 - 1.67 (m, 1 H); MS (ESI): m/z: 475 [M+H] ⁺ .
81	5-[5-cyclobutyl-1-[3-[4-[(3R)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.72 (s, 1 H), 7.50 (d, J=5.4 Hz, 1 H), 7.23 (d, J=5.4 Hz, 1 H), 7.02 - 6.96 (m, 2 H), 6.81 (s, 1 H), 6.77 - 6.71 (m, 2 H), 3.98 - 3.82 (m, 4 H), 3.67 (s, 3 H), 3.58 - 3.52 (m, 1 H), Part AB of ABX System: VA=3.34, VB=2.97, JAB=11.6 Hz, JAX=8.1 Hz, JBX=7.8 Hz, 3.29 - 3.21 (m, 1 H), 3.19 - 3.12 (m, 1 H), 2.73 - 2.65 (m, 1 H), 2.44 (t, J=7.3 Hz, 2 H), 2.34 - 2.25 (m, 2 H), 2.22 - 1.96 (m, 4 H), 1.91 - 1.80 (m, 3 H), 1.78 - 1.67 (m, 1 H); MS (ESI): m/z: 475 [M+H] ⁺ .



82	5-[4-cyclobutyl-1-[3-[4-[[[(3S)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.77 (bs, 1 H), 7.54 (d, <i>J</i> =5.4 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.05 - 6.99 (m, 2 H), 6.84 (bs, 1 H), 6.79 - 6.72 (m, 2 H), 4.03 (t, <i>J</i> =7.3 Hz, 2 H), Part AB of ABX System: VA=3.91, VB=3.86, <i>J</i> AB=9.4 Hz, <i>J</i> AX=6.1 Hz, <i>J</i> BX=7.4 Hz, 3.72 (s, 3 H), 3.65 - 3.55 (m, 1 H), Part AB of ABX System: VA=3.34, VB=2.97, <i>J</i> AB=11.5 Hz, <i>J</i> AX=7.9 Hz, <i>J</i> BX=7.3 Hz, 3.28 - 3.21 (m, 1 H), 3.20 - 3.12 (m, 1 H), 2.72 - 2.65 (m, 1 H), 2.46 (t, <i>J</i> =7.3 Hz, 2 H), 2.38 - 2.28 (m, 2 H), 2.25 - 2.15 (m, 2 H), 2.14 - 1.96 (m, 4 H), 1.92 - 1.84 (m, 1 H), 1.78 - 1.68 (m, 1 H); MS (ESI): <i>m/z</i> : 475 [M+H] ⁺ .
----	--	--	--



83	5-[5-cyclobutyl-1-[3-[4-[(3S)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃ + D ₂ O) δ (ppm): 7.15 (d, <i>J</i> =5.4 Hz, 1 H), 7.02 - 6.92 (m, 4 H), 6.81 - 6.73 (m, 2 H), 6.36 (s, 1 H), 3.94 - 3.81 (m, 4 H), 3.78 (s, 3 H), 3.42 - 3.32 (m, 1 H), Part AB of ABX System: VA=3.14, VB=2.84, JAB=11.4 Hz, JAX=7.4 Hz, JBX=5.6 Hz, 3.08 - 3.01 (m, 1 H), 2.98 - 2.91 (m, 1 H), 2.64 - 2.55 (m, 1 H), 2.50 (t, <i>J</i> =7.3 Hz, 2 H), 2.37 - 2.27 (m, 2 H), 2.24 - 2.14 (m, 2 H), 2.09 - 1.84 (m, 5 H), 1.66 - 1.55 (m, 1 H); MS (ESI): <i>m/z</i> : 475 [M+H] ⁺ .
84	5-[4-cyclobutyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.76 (s, 1 H), 7.51 (d, <i>J</i> =5.4 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.01 (s, 1 H), 6.83 - 6.70 (m, 4 H), 4.44 (t, <i>J</i> =4.9 Hz, 2 H), 4.25 (t, <i>J</i> =4.9 Hz, 2 H), 3.75 - 3.69 (m, 5 H), 3.62 - 3.54 (m, 1 H), 3.31 - 3.22 (m, 2 H), 2.91 - 2.81 (m, 2 H), 2.35 - 2.27 (m, 2 H), 2.22 - 2.12 (m, 2 H), 2.05 - 1.93 (m, 2 H), 1.91 - 1.82 (m, 3 H), 1.47 - 1.35 (m, 2 H); MS (ESI): <i>m/z</i> : 491 [M+H] ⁺ .



85	5-[4-cyclobutyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.86 (bs, 2 H), 7.87 (bs, 1 H), 7.56 (d, <i>J</i> =5.4 Hz, 1 H), 7.28 (d, <i>J</i> =5.4 Hz, 1 H), 7.17 - 7.08 (m, 1 H), 6.87 (bs, 1 H), 6.82 - 6.77 (m, 2 H), 6.74 - 6.68 (m, 1 H), 4.63 - 4.55 (m, 1 H), 4.08 (t, <i>J</i> =7.3 Hz, 2 H), 3.76 (s, 3 H), 3.65 - 3.57 (m, 1 H), 3.23 - 3.14 (m, 2 H), 3.08 - 2.98 (m, 2 H), 2.53 (t, <i>J</i> =7.6 Hz, 2 H), 2.39 - 2.30 (m, 2 H), 2.29 - 2.18 (m, 2 H), 2.16 - 1.98 (m, 5 H), 1.93 - 1.84 (m, 1 H), 1.82 - 1.72 (m, 2 H); MS (ESI): <i>m/z</i> : 475 [M+H] ⁺ .
86	5-[5-cyclobutyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.66 (bs, 1 H), 7.49 (d, <i>J</i> =5.4 Hz, 1 H), 7.22 (d, <i>J</i> =5.4 Hz, 1 H), 7.15 - 7.06 (m, 1 H), 6.83 - 6.74 (m, 3 H), 6.71 - 6.63 (m, 1 H), 4.62 - 4.52 (m, 1 H), 3.94 (t, <i>J</i> =7.6 Hz, 2 H), 3.67 (s, 3 H), 3.58 - 3.51 (m, 1 H), 3.24 - 3.16 (m, 2 H), 3.09 - 2.98 (m, 2 H), 2.47 (t, <i>J</i> =7.3 Hz, 2 H), 2.31 - 2.23 (m, 2 H), 2.20 - 2.11 (m, 2 H), 2.08 - 1.93 (m, 3 H), 1.92 - 1.82 (m, 3 H), 1.80 -



			1.69 (m, 2 H); MS (ESI): m/z: 475 [M+H] ⁺ .
87	5-[4-cyclobutyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.78 (bs, 1 H), 7.53 (d, J=5.4 Hz, 1 H), 7.27 (d, J=5.4 Hz, 1 H), 7.18 - 7.11 (m, 1 H), 7.02 (s, 1 H), 6.62 - 6.55 (m, 1 H), 6.48 - 6.40 (m, 2 H), 4.62 - 4.54 (m, 1 H), 4.46 (t, J=4.9 Hz, 2 H), 4.32 (t, J=4.9 Hz, 2 H), 3.72 (s, 3 H), 3.60 - 3.55 (m, 1 H), 3.24 - 3.15 (m, 2 H), 3.09 - 2.99 (m, 2 H), 2.37 - 2.28 (m, 2 H), 2.24 - 2.13 (m, 2 H), 2.08 - 1.69 (m, 6 H); MS (ESI): m/z: 477 [M+H] ⁺ .



88	5-[5-cyclobutyl-1-[2-[3-(4-piperidyloxy)phenyl]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.76 (bs, 1 H), 7.50 (d, <i>J</i> =5.4 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.13 - 7.07 (m, 1 H), 7.03 (s, 1 H), 6.57 - 6.51 (m, 1 H), 6.35 - 6.27 (m, 2 H), 4.58 - 4.50 (m, 1 H), 4.45 (t, <i>J</i> =4.9 Hz, 2 H), 4.12 (t, <i>J</i> =4.9 Hz, 2 H), 3.81 - 3.72 (m, 1 H), 3.66 (s, 3 H), 3.24 - 3.13 (m, 2 H), 3.09 - 2.98 (m, 2 H), 2.45 - 2.37 (m, 2 H), 2.27 - 2.16 (m, 2 H), 2.09 - 1.68 (m, 6 H); MS (ESI): <i>m/z</i> : 477 [M+H] ⁺ .
89	4-methyl-5-[4-methyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.66 (s, 1 H), 7.53 (d, <i>J</i> =5.4 Hz, 1 H), 7.26 (d, <i>J</i> =5.4 Hz, 1 H), 7.04 - 6.97 (m, 2 H), 6.85 (s, 1 H), 6.81 - 6.76 (m, 2 H), 4.55 - 4.45 (m, 1 H), 4.05 (t, <i>J</i> =7.3 Hz, 2 H), 3.72 (s, 3 H), 3.25 - 3.16 (m, 2 H), 3.09 - 2.98 (m, 2 H), 2.44 (t, <i>J</i> =7.3 Hz, 2 H), 2.33 (s, 3 H), 2.10 - 1.97 (m, 4 H), 1.79 - 1.66 (m, 2 H); MS (ESI): <i>m/z</i> : 435 [M+H] ⁺ .


90	4-methyl-5-[5-methyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆ + D ₂ O) δ (ppm): 7.60 - 7.47 (m, 2 H), 7.24 (d, J=5.4 Hz, 1 H), 7.04 - 6.95 (m, 2 H), 6.84 (s, 1 H), 6.80 - 6.72 (m, 2 H), 4.56 - 4.45 (m, 1 H), 4.03 (t, J=7.3 Hz, 2 H), 3.67 (s, 3 H), 3.26 - 3.15 (m, 2 H), 3.10 - 2.98 (m, 2 H), 2.46 (t, J=7.3 Hz, 2 H), 2.34 (s, 3 H), 2.06 - 1.98 (m, 2 H), 1.96 - 1.87 (m, 2 H), 1.80 - 1.69 (m, 2 H); MS (ESI): m/z: 435 [M+H] ⁺ .
91	5-[4-cyclobutyl-1-[2-[4-(4-piperidylmethoxy)phenyl]ethyl]imidazol-2-yl]-4-methylthieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃ + D ₂ O) δ (ppm): 7.14 (d, J=5.4 Hz, 1 H), 6.98 - 6.88 (m, 3 H), 6.79 - 6.69 (m, 3 H), 6.35 (s, 1 H), 4.18 (t, J=7.3 Hz, 2 H), 3.75 (d, J=6.4 Hz, 2 H), 3.70 (s, 3 H), 3.55 - 3.46 (m, 1 H), 3.19 - 3.10 (m, 2 H), 2.94 (t, J=7.3 Hz, 2 H), 2.70 - 2.60 (m, 2 H), 2.37 - 2.17 (m, 4 H), 2.06 - 1.80 (m, 5 H), 1.35 - 1.26 (m, 2 H); MS (ESI): m/z: 475 [M+H] ⁺ .

92	4-methyl-5-[1-[3-[3-(4-piperidyloxy)phenyl]propyl]-4-propylimidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.84 (bs, 2 H), 7.78 (bs, 1 H), 7.56 (d, <i>J</i> =5.4 Hz, 1 H), 7.29 (d, <i>J</i> =5.4 Hz, 1 H), 7.16 - 7.09 (m, 1 H), 6.88 (bs, 1 H), 6.83 - 6.78 (m, 2 H), 6.74 - 6.68 (m, 1 H), 4.65 - 4.54 (m, 1 H), 4.09 (t, <i>J</i> =7.3 Hz, 2 H), 3.76 (s, 3 H), 3.24 - 3.14 (m, 2 H), 3.08 - 2.94 (m, 2 H), 2.67 (t, <i>J</i> =7.3 Hz, 2 H), 2.53 (t, <i>J</i> =7.3 Hz, 2 H), 2.15 - 1.99 (m, 4 H), 1.83 - 1.64 (m, 4 H), 0.96 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 463 [M+H] ⁺ .
93	4-methyl-5-[1-[3-[3-(4-piperidyloxy)phenyl]propyl]-5-propylimidazol-2-yl]thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃ + D ₂ O) δ (ppm): 7.19 - 7.13 (m, 2 H), 6.97 (d, <i>J</i> =5.4 Hz, 1 H), 6.92 (s, 1 H), 6.78 - 6.73 (m, 1 H), 6.70 - 6.63 (m, 2 H), 6.40 (s, 1 H), 4.41 - 4.34 (m, 1 H), 3.96 (t, <i>J</i> =7.8 Hz, 2 H), 3.79 (s, 3 H), 3.19 - 3.10 (m, 2 H), 2.80 - 2.71 (m, 2 H), 2.58 - 2.42 (m, 4 H), 2.06 - 1.91 (m, 4 H), 1.75 - 1.62 (m, 4 H), 1.03 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 463 [M+H] ⁺ .


94	4-methyl-5-[1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]-4-propylimidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 7.68 (bs, 1 H), 7.50 (d, <i>J</i> =5.4 Hz, 1 H), 7.25 (d, <i>J</i> =5.4 Hz, 1 H), 7.17 - 7.10 (m, 1 H), 7.01 (s, 1 H), 6.61 - 6.55 (m, 1 H), 6.46 - 6.42 (m, 2 H), 4.62 - 4.53 (m, 1 H), 4.46 (t, <i>J</i> =4.9 Hz, 2 H), 4.30 (t, <i>J</i> =4.9 Hz, 2 H), 3.71 (s, 3 H), 3.26 - 3.13 (m, 2 H), 3.11 - 2.97 (m, 2 H), 2.64 (t, <i>J</i> =7.3 Hz, 2 H), 2.11 - 1.96 (m, 2 H), 1.81 - 1.71 (m, 2 H), 1.70 - 1.57 (m, 2 H), 0.91 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 465 [M+H] ⁺ .
95	4-methyl-5-[1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]-5-propylimidazol-2-yl]thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃ + D ₂ O) δ (ppm): 7.17 (d, <i>J</i> =5.4 Hz, 1 H), 7.13 - 7.05 (m, 1 H), 7.02 - 6.90 (m, 2 H), 6.60 (s, 1 H), 6.55 - 6.46 (m, 1 H), 6.38 - 6.26 (m, 2 H), 4.39 (t, <i>J</i> =5.9 Hz, 2 H), 4.35 - 4.28 (m, 1 H), 4.05 (t, <i>J</i> =5.9 Hz, 2 H), 3.76 (s, 3 H), 3.18 - 3.05 (m, 2 H), 2.76 - 2.63 (m, 4 H), 2.04 - 1.93 (m, 2 H), 1.84 - 1.73 (m, 2 H), 1.70 - 1.56 (m, 2 H), 1.09 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): <i>m/z</i> : 465 [M+H] ⁺ .




96	4-methyl-5-[1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.79 (bs, 1 H), 8.52 (bs, 1 H), 7.74 (bs, 1 H), 7.54 (d, J=5.4 Hz, 1 H), 7.28 (d, J=5.4 Hz, 1 H), 7.05 - 6.98 (m, 2 H), 6.86 (s, 1 H), 6.78 - 6.69 (m, 2 H), 4.06 (t, J=7.3 Hz, 2 H), 3.80 - 3.71 (m, 5 H), 3.33 - 3.21 (m, 2 H), 2.96 - 2.81 (m, 2 H), 2.66 (t, J=7.1 Hz, 2 H), 2.48 (t, J=7.3 Hz, 2 H), 2.13 - 1.96 (m, 3 H), 1.93 - 1.82 (m, 2 H), 1.75 - 1.61 (m, 2 H), 1.52 - 1.39 (m, 2 H), 0.96 (t, J=7.3 Hz, 3 H); MS (ESI): m/z: 477 [M+H] ⁺ .
97	4-methyl-5-[1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride		¹ H NMR (DMSO-d ₆) δ (ppm): 8.77 (bs, 1 H), 8.49 (bs, 1 H), 7.66 (bs, 1 H), 7.56 (d, J=5.4 Hz, 1 H), 7.27 (d, J=5.4 Hz, 1 H), 7.03 - 6.95 (m, 2 H), 6.86 (bs, 1 H), 6.76 - 6.70 (m, 2 H), 4.04 (d, J=7.3 Hz, 2 H), 3.76 (d, J=6.4 Hz, 2 H), 3.71 (s, 3 H), 3.34 - 3.21 (m, 2 H), 2.96 - 2.80 (m, 2 H), 2.63 (t, J=7.6 Hz, 2 H), 2.49 - 2.43 (m, 2 H), 2.06 - 1.97 (m, 1 H), 1.95 - 1.84 (m, 4 H), 1.75 - 1.62 (m, 2 H), 1.54 - 1.39 (m, 2 H), 1.01 (t, J=7.3


			Hz, 3 H); MS (ESI): m/z: 477 [M+H] ⁺ .
98	5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.16 (d, <i>J</i> =5.4 Hz, 1 H), 6.97 (d, <i>J</i> =5.4 Hz, 1 H), 6.92 (s, 1 H), 6.83 - 6.70 (m, 4 H), 6.56 (s, 1 H), 4.53 - 4.46 (m, 1 H), 4.36 (t, <i>J</i> =5.4 Hz, 2 H), 4.14 (t, <i>J</i> =5.4 Hz, 2 H), 3.82 (s, 3 H), 3.21 - 2.98 (m, 4 H), 2.60 (t, <i>J</i> =7.6 Hz, 2 H), 2.23 - 1.81 (m, 6 H), 1.77 - 1.65 (m, 2 H), 1.00 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): m/z: 479 [M+H] ⁺ .
99	5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (DMSO- <i>d</i> ₆) δ (ppm): 7.32 (d, <i>J</i> =5.4 Hz, 1 H), 7.19 (d, <i>J</i> =5.4 Hz, 1 H), 6.86 (s, 1 H), 6.78 - 6.61 (m, 5 H), 4.46 - 4.39 (m, 1 H), 4.35 (t, <i>J</i> =5.4 Hz, 2 H), 4.03 (t, <i>J</i> =5.4 Hz, 2 H), 3.68 (s, 3 H), 3.06 - 2.78 (m, 4 H), 2.66 (t, <i>J</i> =7.8 Hz, 2 H), 2.05 - 1.48 (m, 8 H), 1.01 (t, <i>J</i> =7.3 Hz, 3 H); MS (ESI): m/z: 479 [M+H] ⁺ .

100	4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-4-(3,3,3-trifluoropropyl)imidazol-2-yl]thieno[3,2-b]pyrrole		¹ H NMR (DMSO-d ₆) δ (ppm): 7.32 (d, <i>J</i> =5.4 Hz, 1 H), 7.21 (s, 1 H), 7.17 (d, <i>J</i> =5.4 Hz, 1 H), 7.00 - 6.95 (m, 2 H), 6.79 - 6.72 (m, 2 H), 6.44 (s, 1 H), 4.35 - 4.24 (m, 1 H), 3.95 (t, <i>J</i> =7.3 Hz, 2 H), 3.78 (s, 3 H), 3.01 - 2.90 (m, 2 H), 2.79 - 2.71 (m, 2 H), 2.67 - 2.54 (m, 4 H), 2.43 (t, <i>J</i> =7.3 Hz, 2 H), 2.01 - 1.82 (m, 4 H), 1.50 - 1.38 (m, 2 H); MS (ESI): <i>m/z</i> : 517 [M+H] ⁺ .
-----	--	--	---


Compounds **101** and **102** were obtained starting from compound **2** and 1-bromo-2-methoxy-ethane (Sigma-Aldrich, Cat. No. 238155) according to the procedure described for Intermediate **26** and separated by column chromatography on silica gel (hexane/acetone, 95:5 → 60:40, v:v). Compounds **103** and **104** were obtained starting from compound **2** and 2-bromoethoxybenzene (Sigma-Aldrich, Cat. No. B75506-25G) and compound **105** from compound **2** and benzylbromide (Sigma-Aldrich, Cat. No. B17905) according to the procedure described for Intermediate **26**.

101	5-[5-ethyl-1-(2-methoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.18-7.15 (m, 1 H), 7.00-6.93 (m, 2 H), 6.60 (s, 1 H), 4.21-4.15 (m, 2 H), 3.81 (s, 3 H), 3.56-3.50 (m, 2 H), 3.25 (s, 3 H), 2.74-2.60 (m, 2 H), 1.39-1.33 (m, 3 H); MS (ESI): <i>m/z</i> : 290 [M+H] ⁺ .
-----	---	---	---

102	5-[4-ethyl-1-(2-methoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.18-7.14 (m, 1 H), 6.98-6.95 (m, 1 H), 6.88 (s, 1 H), 6.55 (s, 1 H), 4.19-4.14 (m, 2 H), 3.85 (s, 3 H), 3.66-3.61 (m, 2 H), 3.34 (s, 3 H), 2.80-2.62 (m, 2 H), 1.34-1.27 (m, 3 H); MS (ESI): m/z: 290 [M+H] ⁺ .
103	5-[5-ethyl-1-(2-phenoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.32-7.25 (m, 2 H), 7.20-7.15 (m, 1 H), 7.02-6.93 (m, 3 H), 6.88-6.80 (m, 2 H), 6.60 (s, 1 H), 4.44-4.37 (m, 2 H), 4.23-4.18 (m, 2 H), 3.84 (s, 3 H), 2.76-2.62 (m, 2 H), 1.34-1.24 (m, 3 H); MS (ESI): m/z: 352 [M+H] ⁺ .
104	5-[4-ethyl-1-(2-phenoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.32-7.25 (m, 2 H), 7.20-7.15 (m, 1 H), 7.02-6.93 (m, 3 H), 6.88-6.80 (m, 2 H), 6.60 (s, 1 H), 4.44-4.37 (m, 2 H), 4.23-4.18 (m, 2 H), 3.84 (s, 3 H), 2.76-2.62 (m, 2 H), 1.34-1.24 (m, 3 H); MS (ESI): m/z: 352 [M+H] ⁺ .


105	5-(1-benzyl-5-ethyl-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole		¹ H NMR (CDCl ₃) δ (ppm): 7.40–7.29 (m, 3 H), 7.17–7.12 (m, 1 H), 7.10–7.04 (m, 2 H), 6.98–6.93 (m, 1 H), 6.73 (s, 1 H), 6.41 (s, 1 H), 5.22 (s, 2 H), 3.85 (s, 3 H), 2.78–2.61 (m, 2 H), 1.33–1.26 (m, 3 H); MS (ESI): m/z: 322 [M+H] ⁺ .
-----	--	---	--

Example 106: 4-Methyl-5-[4-(4-pyridyl)-1H-imidazol-2-yl]thieno[3,2-b]pyrrole



4-Methyl-5-[4-(4-pyridyl)-1H-imidazol-2-yl]thieno[3,2-b]pyrrole was prepared starting from 4-methylthieno[3,2-b]pyrrole-5-carboxamide acetate (Intermediate 11) and 2-bromo-1-(4-pyridyl)ethanone (Fluorochem, Cat No. 017122) according to the procedure described for Example 1 and purified by flash chromatography on silica gel (CH₂Cl₂/MeOH/NH₃, 97:3:0.5, v:v:v). ¹H NMR (MeOH-d₄) δ (ppm): 8.53–8.46 (m, 2 H), 7.88–7.79 (m, 3 H), 7.25 (d, *J*=5.4 Hz, 1 H), 7.09 (d, *J*=5.4 Hz, 1 H), 6.83 (s, 1 H), 4.14 (s, 3 H); MS (ESI): m/z: 281 [M+H]⁺.


Example 107: 5-(1H-Benzimidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole



0.050 g (0.30 mmol) of 4-methylthieno[3,2-b]pyrrole-5-carbaldehyde (Aldrich, Cat. N.

CDS015804), 0.033 g (0.30 mmol) of *o*-phenyldiamine (Aldrich, Cat. N. P23938-5G) and iodobenzene diacetate (Aldrich, Cat. N. 178721-25G) were stirred at -15°C in solvent free conditions. The mixture was then warmed to RT becoming a sticky brown oil. After stirring at RT for 1.5 h a further portion of 0.018g (0.18 mmol) of *o*-phenyldiamine was added. The obtained dark oil was purified by column chromatography on silica gel (EtOAc/hexane 3:97, v:v to 30:70 v:v) to provide 0.025 g of 5-(1H-benzimidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole (33%). ¹H NMR (DMSO-d₆) δ (ppm): 12.74–12.57 (bs, 1 H), 7.68–7.46 (m, 2 H), 7.44–7.39 (m, 1 H), 7.28–7.22 (m, 1 H), 7.21–7.14 (m, 3 H), 4.26 (s, 3 H); MS (ESI): m/z: 254 [M+H]⁺.

Example 108: 4-Methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]benzimidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride



tert-Butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)benzimidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate

tert-Butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)benzimidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate was prepared starting from 5-(1H-benzimidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole (Example 107) and *tert*-butyl 4-[4-(3-bromopropyl)phenoxy]piperidine-1-carboxylate (Intermediate 19) according to the procedure for Intermediate 26. ¹H NMR (CDCl₃) δ (ppm): 7.95–7.74 (m, 1 H), 7.39–7.29 (m, 3 H), 7.26 (d, *J*=5.4 Hz, 1 H), 7.06–6.99 (m, 3 H), 6.85–6.80 (m, 2 H), 6.59 (s, 1 H), 4.48–4.39 (m, 1 H), 4.38–4.31 (t, *J*=7.1 Hz, 2 H), 4.02 (s, 3 H), 3.76–3.66 (m, 2 H), 3.38–3.26 (m, 2 H), 2.64 (t, *J*=7.1 Hz, 2 H), 2.25–2.15 (m, 2 H), 1.97–1.85 (m, 2 H), 1.80–1.69 (m, 2 H), 1.48 (s, 9 H); MS (ESI): m/z: 571 [M+H]⁺.

4-Methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]benzimidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride

4-Methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]benzimidazol-2-yl]thieno[3,2-b]pyrrole hydrochloride was obtained starting from *tert*-butyl 4-[4-[3-[2-(4-methylthieno[3,2-b]pyrrol-5-yl)benzimidazol-1-yl]propyl]phenoxy]piperidine-1-carboxylate according to the procedure described Example 18. ¹H NMR (DMSO-d₆) δ (ppm): 8.80–8.57 (m, 2 H), 7.86–7.75 (m, 2 H), 7.54 (d, *J*=5.4 Hz, 1 H), 7.49–7.38 (m, 2 H), 7.29 (d, *J*=5.4 Hz, 1 H), 7.12–7.05 (m, 2 H), 6.89–6.81 (m, 3 H), 4.59–4.52 (m, 1 H), 4.40 (t, *J*=7.3 Hz, 2 H), 3.94 (s, 3 H), 3.27–3.14 (m, 2 H), 3.10–3.00 (m, 2H),, 2.58 (t, *J*=7.3 Hz, 2 H), 2.16–1.99 (m, 4 H), 1.83–1.71 (m, 2 H); MS (ESI): *m/z*: 471 [M+H]⁺.

2. BIOLOGICAL TESTING

2.1 Assay of enzyme inhibition of KDM1A (LSD1)

The KDM1A inhibiting activity was determined using a TR-FRET assay (time resolved fluorescence resonance energy transfer, Lance® Ultra Demethylase technology (Perkin Elmer, Waltham, MA, USA)), which comprises a Europium chelate donor dye (TRF0404, Perkin Elmer, Waltham, MA, USA) together with *ULight*TM (TR0102, Perkin Elmer, Waltham, MA, USA), a small molecular weight acceptor dye with a red-shifted fluorescent emission, and a biotinylated 21 aminoacids histone H3-derived monomethylated peptide (H3K4me) [Lys(Me1)4]-Histone H3 (1-21)-GGK(biotin), (64355, Anaspec, Fremont, CA, USA) as substrate. The intensity of the light emission is proportional to the level of biotinylated reaction product. The complex of human recombinant KDM1A/CoREST protein was produced in *E. coli* as separate proteins and co-purified as previously described.(Forneris, F. *et al.* Trends Biochem, Sci, 2008, 33, 181-189) (Forneris, F. *et al.* J. Biol. Chem. 2007, 282, 20070-20074).

Demethylase Assay conditions: 0.25 nM KDM1A/CoREST protein and compound in 100% DMSO were added in a final volume of 48 µL assay buffer (Tris HCl 50 mM pH 8.8, NaCl 50 mM, DTT 1 mM, Tween-20 0.01%) to each well of a 96 well half area flat bottom white plate (3693 Costar, Sigma-Aldrich, St. Louis, M, USA).

Demethylase reaction was started by the addition of 50nM histone H3K4 monomethylated. After 20 min at RT, 300 µM tranilcypromine (P8511-1G, Sigma-Aldrich, St. Louis, MO 63103) was added to stop the reaction.

Detection step conditions: 10 µL of the assay mixture was transferred from the original

plate into a 384 well white plate (6007290 OptiPlate™, Perkin Elmer, Waltham, MA, USA) and 10 µL of the detection Mix containing 2 nM Eu-antibody and 10 nM U-*Light*-Streptavidin in 1X Lance Detection Buffer (TRF0404, TR0102, CR97100, Perkin Elmer, Waltham, MA, USA). The resulting mixture was incubated in the dark for 1 h at RT. Then, TR-FRET signal was read by a fluorimeter (Infinite® F200, Tecan, Männedorf, Switzerland) (Excitation 320nm, Emission 665 nm and 620 nm, delay time 50µs, window time 100µs).

IC₅₀ determination: The inhibitor concentrations ranged from 0.025 to 500 µM (serial 1:3 dilutions). The IC₅₀ was calculated using GraphPad Software.

Compounds **2**, **4-8**, **11**, **13**, **15**, **16** and **70** exhibit IC₅₀ values of less than 10 µM, examples **9**, **10** and **12** exhibit IC₅₀ values of less than 1 µM, examples **18-54** and **57-62** exhibit an IC₅₀ value of less than 0.1 µM.

2.2 Cell growth

CellTiter-Fluor® (Promega) is as a nonlytic, single-reagent-addition fluorescence assay that measures the relative number of living cells in a culture population after experimental manipulation. The CellTiter-Fluor™ Cell Viability Assay measures the conserved and constitutive protease activity within live cells and therefore acts as a marker for cell viability.

Human leukemia MV4-11 cells, (obtained from the Deutsche Sammlung von Mikroorganismen und Zellkulturen, **ACC 102**) or NB4 cells, (obtained from the Deutsche Sammlung von Mikroorganismen und Zellkulturen) in exponential growth, were incubated for 48 h with different concentrations of the inhibitors. After 48 h a volume of CellTiter-Fluor® Reagent equal to one fifth of volume of cell culture medium was added. The content was mixed and incubates for at least 90 min at 37 °C degree to obtain a stable signal. The fluorescence was recorded using an excitation wavelength of 360 nm and an emission at 535 nm. The IC₅₀ was calculated using GraphPad Software.

Compounds **18-33** exhibit IC₅₀ values of less than 10 µM against human leukemia MV4-11 cells and IC₅₀ values of less than 20 µM against human leukemia NB4 cells, compounds **39-50** and **57** IC₅₀ values of less than 10 µM against human leukemia NB4


cells.

2.3 Bioluminescent-Coupled Assay for Monoamine oxidases (MAO-Glo Assay)

The MAO Glo Assay from Promega (cat. V1402, Promega, Madison, WI) was used to measure the effect of inhibitors on MAO A and MAO B activity. Human recombinant MAO A and MAO B were expressed in *Pichia pastoris* and purified as published (Binda C. *et al.* Proc. Natl. Acad. Sci. USA, 2003, 9750-9755). The assay was performed at RT in 50 μ L (25 μ L reaction solution + 25 μ L detection reagent) in 96 well half area white plates (cat. 3693, Corning, Corning, NY). Luminescence was measured after 20 min incubation in the dark using a microplate reader (Infinite F200, Tecan Group, Switzerland) with an integration time of 0.25 s per well. 50 nM MAO A or 125 nM MAO B were incubated with five different inhibitor concentrations (from 0.004 μ M to 100 μ M) for 15 min at RT in Promega MAO Buffer or Promega MAO B Buffer (MAO Glo Assay kit, catalogue number V1402, Promega, Madison, WI). After 30 min of incubation the reaction was stopped with the Promega detection reagent. All compounds were tested twice and IC₅₀ values were calculated using GraphPad Prism version 4.0 (GraphPad Software, San Diego, CA). Compounds **7**, **10**, **19-20**, and **23-26** were at least 10 times more active against **KDM1A (LSD1)** compared to both MAO A and MAO B.

CLAIMS


1. A compound of formula (I)




(I)

wherein

A



or



R

is hydrogen or **L¹-R⁵**;

R¹, R²

are independently, hydrogen, C₁-C₄-alkyl, C₃-C₇ cycloalkyl, aryl, heteroaryl, -(CH₂)_zCF₃, or **R¹** and **R²**, taken together with the carbon atoms they are bound to, form with the imidazole a fused heterocycle or a fused heteroaryl;

z

is an integer from zero to 6

R³, R⁴

are hydrogen or C₁-C₄-alkyl;

L¹

is -(CH₂)_j-Y- or -CH₂-;

j

is an integer from 2 to 6;

Y

is a bond, oxygen, or CH₂;

R⁵

is C₁-C₄-alkyl or aryl, wherein the aryl is optionally substituted by halogen, C₁-C₆-alkyl or **L²-R⁶**;

L²

is -(CH₂)_m-; -(CH₂)_n-**W**-(CH₂)_p-;

R⁶

is heterocyclyl, wherein the heterocyclyl is optionally substituted by C₁-C₆-alkyl;

m, n, p

are, independently, zero or an integer from 1 to 6;

W

is oxygen, sulphur, NH, or CH₂;

or stereoisomers or pharmaceutically acceptable salts thereof.

2. A compound of formula (I) according to claim 1, wherein **R**¹ and **R**² are independently hydrogen, C₁-C₄-alkyl, C₃-C₇ cycloalkyl, aryl, heteroaryl, or **R**¹ and **R**², taken together with the carbon atoms they are bound to, form with the imidazole a fused heterocycle or a fused heteroaryl.

3. A compound of formula (I) according to claim 1 or claim 2, wherein **R**² or **R**³ are methyl or ethyl.

4. A compound of formula (I) according to any one of claims 1 to 3, wherein

R is hydrogen or **L**¹-**R**⁵;

R¹, **R**² are independently, hydrogen, C₁-C₄-alkyl, cyclopropyl, cyclobutyl, phenyl, pyridyl, or **R**¹ and **R**², taken together with the carbon atoms they are bound to, form with the imidazole a fused heterocycle or a fused heteroaryl;

R³, **R**⁴ are methyl or ethyl;

L¹ is -(CH₂)₂-**Y**- or -CH₂-;

Y is a bond, oxygen, or CH₂;

R⁵ is methyl or phenyl optionally substituted by **L**²-**R**⁶;

L² is -**W**-(CH₂)_p-;

R⁶ is heterocyclyl, wherein the heterocyclyl is optionally substituted by C₁-C₆-alkyl;

p is zero or an integer from 1 to 6;

W is oxygen;

or stereoisomers or pharmaceutically acceptable salts thereof.

5. A compound according to claim 1 selected from:

4-methyl-5-(4-phenyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;

5-(4-ethyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;

4-methyl-5-(4-methyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;

5-(4-isopropyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;

5-(4-cyclopropyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;

4-methyl-5-(4-propyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(4-cyclobutyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-(5-methyl-4-phenyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(5-ethyl-4-phenyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
5-(5-isopropyl-4-phenyl-1H-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-ethyl-5-(4-ethyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(4-cyclobutyl-1H-imidazol-2-yl)-4-ethyl-thieno[3,2-b]pyrrole;
4-methyl-5-(4-phenyl-5-propyl-1H-imidazol-2-yl)thieno[3,2-b]pyrrole;
5-(4-ethyl-1H-imidazol-2-yl)-6-methyl-thieno[2,3-b]pyrrole;
5-(5-ethyl-4-phenyl-1H-imidazol-2-yl)-6-methyl-thieno[2,3-b]pyrrole;
4-methyl-5-(4,5,6,7-tetrahydro-1H-benzimidazol-2-yl)thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[5-ethyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[5-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[4-(pyrrolidin-3-ylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[5-ethyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-ethyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-ethyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-
b]pyrrole;

5-[5-ethyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-cyclobutyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-cyclobutyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-ethyl-4-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-ethyl-4-phenyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-5-phenyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[3-(azepan-4-yloxy)phenoxy]ethyl]-4-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[3-(azepan-4-yloxy)phenoxy]ethyl]-5-ethyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole;

4-methyl-5-[5-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;

4-methyl-5-[4-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-ethyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-

thieno[3,2-b]pyrrole;
5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-4-ethyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-ethyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[5-ethyl-4-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[4-ethyl-5-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[4-isopropyl-5-phenyl-1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-
methyl-thieno[3,2-b]pyrrole;
5-[4-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[5-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
4-methyl-5-[4-phenyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-
yl]thieno[3,2-b]pyrrole;
5-[1-[3-[3,4-bis(4-piperidylmethoxy)phenyl]propyl]-4-ethyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-4-propyl-imidazol-2-
yl]thieno[3,2-b]pyrrole;
4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenoxy]ethyl]-5-propyl-imidazol-2-
yl]thieno[3,2-b]pyrrole;
5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-4-propyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-propyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[1-[3-[3,4-bis(4-piperidylmethoxy)phenyl]propyl]-5-ethyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole;
5-[4-ethyl-1-[2-[4-(4-piperidyloxy)phenyl]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-
b]pyrrole;

5-[4-cyclobutyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-cyclobutyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-4-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[1-[3-[4-(azepan-4-yloxy)phenyl]propyl]-5-cyclobutyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;

4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;

5-[4-cyclobutyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-cyclobutyl-1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

4-methyl-5-[4-propyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole;

4-methyl-5-[5-propyl-1-[3-[4-(pyrrolidin-3-ylmethoxy)phenyl]propyl]imidazol-2-yl]thieno[3,2-b]pyrrole;

5-[4-cyclobutyl-1-[2-[4-(4-piperidyloxy)phenyl]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[4-ethyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

5-[5-ethyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;

4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenyl]ethyl]-4-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;

4-methyl-5-[1-[2-[4-(4-piperidyloxy)phenyl]ethyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;

4-methyl-5-[1-[3-[4-(4-piperidyloxymethyl)phenyl]propyl]-4-propyl-imidazol-2-

yl]thieno[3,2-b]pyrrole;
4-methyl-5-[1-[3-[4-(4-piperidyloxymethyl)phenyl]propyl]-5-propyl-imidazol-2-yl]thieno[3,2-b]pyrrole;
5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-1-(2-methoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-ethyl-1-(2-methoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[5-ethyl-1-(2-phenoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-[4-ethyl-1-(2-phenoxyethyl)imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole;
5-(1-benzyl-5-ethyl-imidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-[4-(4-pyridyl)-1H-imidazol-2-yl]thieno[3,2-b]pyrrole;
5-(1H-benzimidazol-2-yl)-4-methyl-thieno[3,2-b]pyrrole;
4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]benzimidazol-2-yl]thieno[3,2-b]pyrrole;
5-[4-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole
5-[5-cyclobutyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-6-methyl-thieno[2,3-b]pyrrole
5-[4-cyclobutyl-1-[3-[4-[[[(3R)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole
5-[5-cyclobutyl-1-[3-[4-[[[(3R)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole
5-[4-cyclobutyl-1-[3-[4-[[[(3S)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole
5-[5-cyclobutyl-1-[3-[4-[[[(3S)-pyrrolidin-3-yl]methoxy]phenyl]propyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole
5-[4-cyclobutyl-1-[2-[4-(4-piperidylmethoxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole
5-[4-cyclobutyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-

thieno[3,2-b]pyrrole
5-[5-cyclobutyl-1-[3-[3-(4-piperidyloxy)phenyl]propyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole
5-[4-cyclobutyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole
5-[5-cyclobutyl-1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole
4-methyl-5-[4-methyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-
yl]thieno[3,2-b]pyrrole
4-methyl-5-[5-methyl-1-[3-[4-(4-piperidyloxy)phenyl]propyl]imidazol-2-
yl]thieno[3,2-b]pyrrole
5-[4-cyclobutyl-1-[2-[4-(4-piperidylmethoxy)phenyl]ethyl]imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole
4-methyl-5-[1-[3-[3-(4-piperidyloxy)phenyl]propyl]-4-propyl-imidazol-2-
yl]thieno[3,2-b]pyrrole
4-methyl-5-[1-[3-[3-(4-piperidyloxy)phenyl]propyl]-5-propyl-imidazol-2-
yl]thieno[3,2-b]pyrrole
4-methyl-5-[1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]-4-propyl-imidazol-2-
yl]thieno[3,2-b]pyrrole
4-methyl-5-[1-[2-[3-(4-piperidyloxy)phenoxy]ethyl]-5-propyl-imidazol-2-
yl]thieno[3,2-b]pyrrole
4-methyl-5-[1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]-4-propyl-imidazol-2-
yl]thieno[3,2-b]pyrrole
4-methyl-5-[1-[3-[4-(4-piperidylmethoxy)phenyl]propyl]-5-propyl-imidazol-2-
yl]thieno[3,2-b]pyrrole
5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-4-propyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole
5-[1-[2-[4-(azepan-4-yloxy)phenoxy]ethyl]-5-propyl-imidazol-2-yl]-4-methyl-
thieno[3,2-b]pyrrole
4-methyl-5-[4-(3,3,3-trifluoropropyl)-1H-imidazol-2-yl]thieno[3,2-b]pyrrole
4-methyl-5-[1-[3-[4-(4-piperidyloxy)phenyl]propyl]-4-(3,3,3-

trifluoropropyl)imidazol-2-yl]thieno[3,2-b]pyrrole

or stereoisomers or pharmaceutically acceptable salts thereof.

6. A pharmaceutical composition comprising a compound as defined in any one of claims 1 to 5.

7. The pharmaceutical composition according to claim 6 further comprising at least one therapeutic agent, selected from the group consisting of histone deacetylase inhibitors, retinoid receptor modulators, antiproliferative/antineoplastic agents, cytostatic agents, agents which inhibit cancer cell invasion, inhibitors of growth factor function, antiangiogenic agents, cell cycle inhibitors, proteasome inhibitors, HSP90 inhibitors, Selective COX-2 inhibitors or a chemotherapeutic agent.

8. The pharmaceutical composition according to claim 6 or 7 in the form of tablets, capsules, oral preparations, powders, granules, pills, injectable or infusible liquid, solutions, suspensions, emulsions, suppositories, ointments, creams, lotions, gels, pastes, transdermal delivery devices.


9. The compound according to any one of claims 1 to 5 for use as medicament.

10. The compound according to any one of claims 1 to 5 for use in the treatment and/or prevention of cancer, infectious diseases or a disease characterized by aberration of cellular energy metabolism, such as obesity.

11. The compound according to claim 10 for use in the treatment and/or prevention of leukemia, non-small cell lung cancer, hepatocellular carcinoma, or glioblastomas.

12. A process for obtaining a compound of formula (I) according any one of claims 1 to 5, wherein **R** is hydrogen, the process comprising the preparation of compounds of formula (I) by reaction of a compound of formula **A1** with the suitable with α -halogeno


ketones of formula **A2** as represented in **Scheme A-I** below:



Scheme A-I

wherein **A**, **R¹**, and **R²** are as defined in claim 1 and **X** is a chlorine, bromine or iodine.

13. A process for obtaining a compound of formula **(I)** according to any one of claims 1 to 5, wherein **R** is **L¹-R⁵**, the process comprising the reaction of a compound of formula **(I)** according to claim 1, wherein **R** is hydrogen, with derivatives of formula **A3** in presence of a base, as represented in **Scheme A-II** below:



Scheme A-II

wherein **A**, **R¹**, and **R²** are as defined in claim 1, **R** is **L¹-R⁵**, wherein **L¹** and **R⁵** are as defined in claim 1, and **LG** is a leaving group.

14. The compound according to any one of claims 1 to 5, wherein the compound of formula **(I)** is a reversible KDM1A (LSD1) inhibitor.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/062009

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D495/04 A61K31/407 A61P35/00 A61P3/04 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, WPI Data, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 2 993 175 A1 (IEO ISTITUTO EUROP DI ONCOLOGIA SRL [IT]) 9 March 2016 (2016-03-09) cited in the application abstract examples claims ----- -/--	1-14
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 20 June 2017		Date of mailing of the international search report 17/07/2017
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Stix-Malaun, Elke

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2017/062009

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VARIER R A ET AL: "Histone lysine methylation and demethylation pathways in cancer", BBA - REVIEWS ON CANCER, ELSEVIER SCIENCE BV, AMSTERDAM, NL, vol. 1815, no. 1, 1 January 2011 (2011-01-01), pages 75-89, XP027577984, ISSN: 0304-419X [retrieved on 2010-10-15] Chapter 3.1 the whole document -----	1-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2017/062009

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
EP 2993175	A1	09-03-2016	EP 2993175 A1	09-03-2016
			EP 3189059 A2	12-07-2017
			WO 2016034946 A2	10-03-2016
